
a widely expressed cytosolic chaperone (presenter protein) in the cell (e.g., cyclophilin A). More specifically, in some examples, the inhibitors of Ras described herein induce a new binding pocket in Ras by driving formation of a high affinity tri-complex, or conjugate, between the Ras protein and the widely expressed cytosolic chaperone, cyclophilin A (CYPA). Without being bound by theory, the inventors believe that one way the inhibitory effect on Ras is effected by compounds of the disclosure and the complexes, or conjugates, they form is by steric occlusion of the interaction site between Ras and downstream effector molecules, such as RAF and PI3K, which are required for propagating the oncogenic signal.

As such, in an example, the disclosure features a compound, or pharmaceutically acceptable salt thereof, of structural Formula I:



Formula I

wherein the dotted lines represent zero, one, two, three, or four non-adjacent double bonds;

A is -N(H or CH_3)C(O)-(CH_2)- where the amino nitrogen is bound to the carbon atom of -CH(R^{10})-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or optionally substituted 5 to 10-membered heteroarylene;

B is absent, -CH(R^9)-, >C=C$R^9R^{9'}$, or >C$R^9R^{9'}$ where the carbon is bound to the carbonyl carbon of -N(R^{11})C(O)-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or 5 to 6-membered heteroarylene;

G is optionally substituted C_1-C_4 alkylene, optionally substituted C_1-C_4 alkenylene, optionally substituted C_1-C_4 heteroalkylene, -C(O)O-**C**H(R^6)- where **C** is bound to -C(R^7R^8)-, -C(O)NH-**C**H(R^6)- where **C** is bound to -C(R^7R^8)-, optionally substituted C_1-C_4 heteroalkylene, or 3 to 8-membered heteroarylene;

L is absent or a linker;

W is a cross-linking group comprising a vinyl ketone, a vinyl sulfone, an ynone, a haloacetal, or an alkynyl sulfone;

X^1 is optionally substituted C_1-C_2 alkylene, NR, O, or $S(O)_n$;

substituted 3 to 14-membered heterocycloalkyl, optionally substituted 5 to 10-membered heteroaryl, or optionally substituted 6 to 10-membered aryl, or

$R^{7'}$ and $R^{8'}$ combine with the carbon atom to which they are attached to form optionally substituted 3 to 6-membered cycloalkyl or optionally substituted 3 to 7-membered heterocycloalkyl;

R^9 is H, F, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl; or

R^9 and L combine with the atoms to which they are attached to form an optionally substituted 3 to 14-membered heterocycloalkyl;

$R^{9'}$ is hydrogen or optionally substituted C_1-C_6 alkyl; or

R^9 and $R^{9'}$, combined with the atoms to which they are attached, form a 3 to 6-membered cycloalkyl or a 3 to 6-membered heterocycloalkyl;


R^{10} is hydrogen, halo, hydroxy, C_1-C_3 alkoxy, or C_1-C_3 alkyl;

R^{10a} is hydrogen or halo;

R^{11} is hydrogen or C_1-C_3 alkyl; and

R^{21} is hydrogen or C_1-C_3 alkyl (e.g., methyl).

In an aspect, there is provided a compound having the structure of Formula If:



Formula If


or pharmaceutically acceptable salt thereof,

wherein A is -N(H or CH_3)C(O)-(CH_2)- where the amino nitrogen is bound to the carbon atom of -CH(R^{10})-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or optionally substituted 5 to 6-membered heteroarylene;


B is -CH(R^9)- where the carbon is bound to the carbonyl carbon of -NHC(O)-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or 5 to 6-membered heteroarylene;


L is absent or a linker;


W is a cross-linking group comprising


a ketone of formula: 

,

a vinyl ketone of formula: ,

a vinyl sulfone of formula: ,

an ynone of formula: , or

an alkynyl sulfone of formula: ;

R^1 is cyano, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 6-membered cycloalkenyl, optionally substituted 3 to 6-membered heterocycloalkyl, optionally substituted 6 to 10-membered aryl, or optionally substituted 5 to 10-membered heteroaryl;

R^2 is C_1-C_6 alkyl, C_1-C_6 fluoroalkyl, or 3 to 6-membered cycloalkyl;

R^7 is C_1-C_3 alkyl;

R^8 is C_1-C_3 alkyl; and

R^9 is optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl;

R^{16a}, R^{16b}, and R^{16c} are, independently, hydrogen, -CN, halogen, or -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl;

R^{17} is hydrogen, -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), or a 4 to 7-membered saturated cycloalkyl, or a 4 to 7-membered saturated heterocycloalkyl;


R^{18a}, R^{18b}, and R^{18c} are, independently, hydrogen, -CN, or -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl;


R^{19} is hydrogen, -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl, or a 4 to 7-membered saturated heterocycloalkyl;


X^e is a halogen; and


R^{20} is hydrogen, -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl.


In another aspect, there is provided a compound selected from:












































































































































































































































assumed





assumed





4h









































































4j






assumed






assumed





































































































































































































































































assumed








assumed














































4s





































4t









































































OMe





































4w





































4x





































4y









































































4aa





































4ab





































4ac





































4ad





































4ae





































4af






































































4ah









































































4aj









































































4al





































4am





































4an





















































































































































































4as




























































































































































assumed






assumed









OH














































4ay


































assumed








































4ba









































































4bc


















































































































































































4bh





































4bi





















assumed






































or a pharmaceutically acceptable salt thereof.

Also provided are pharmaceutical compositions comprising a compound of Formula I, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.

In another aspect, there is provided a pharmaceutical composition comprising a compound, or a pharmaceutically acceptable salt thereof, as defined above, and a pharmaceutically acceptable excipient.

In another aspect, there is further provided a conjugate, or salt thereof, comprising the structure of Formula IV:


M-L-P

Formula IV

wherein L is a linker;

P is a monovalent organic moiety; and

M has the structure of Formula V:



Formula V

wherein the dotted lines represent zero, one, two, three, or four non-adjacent double bonds;

A is -N(H or CH_3)C(O)-(CH_2)- where the amino nitrogen is bound to the carbon atom of -CH(R^{10})-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or optionally substituted 5 to 6-membered heteroarylene;

B is absent, -CH(R^9)-, >C=C$R^9R^{9'}$, or >C$R^9R^{9'}$ where the carbon is bound to the carbonyl carbon of -N(R^{11})C(O)-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-

R^7 and R^8 combine with the carbon atom to which they are attached to form $C=CR^{7'}R^{8'}$; C=N(OH), C=N(O-C_1-C_3 alkyl), C=O, C=S, C=NH, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl;

R^{7a} and R^{8a} are, independently, hydrogen, halo, optionally substituted C_1-C_3 alkyl, or combine with the carbon to which they are attached to form a carbonyl;

$R^{7'}$ is hydrogen, halogen, or optionally substituted C_1-C_3 alkyl; $R^{8'}$ is hydrogen, halogen, hydroxy, cyano, optionally substituted C_1-C_3 alkoxy, optionally substituted C_1-C_3 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 8-membered cycloalkyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 5 to 10-membered heteroaryl, or optionally substituted 6 to 10-membered aryl, or

$R^{7'}$ and $R^{8'}$ combine with the carbon atom to which they are attached to form optionally substituted 3 to 6-membered cycloalkyl or optionally substituted 3 to 7-membered heterocycloalkyl;

R^9 is H, F, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl, or

R^9 and L combine with the atoms to which they are attached to form an optionally substituted 3 to 14-membered heterocycloalkyl;

$R^{9'}$ is hydrogen or optionally substituted C_1-C_6 alkyl; or

R^9 and $R^{9'}$, combined with the atoms to which they are attached, form a 3 to 6-membered cycloalkyl or a 3 to 6-membered heterocycloalkyl;

R^{10} is hydrogen, halo, hydroxy, C_1-C_3 alkoxy, or C_1-C_3 alkyl;

R^{10a} is hydrogen or halo;

R^{11} is hydrogen or C_1-C_3 alkyl; and

R^{21} is H or C_1-C_3 alkyl.

In another aspect, there is provided a compound, or a pharmaceutically acceptable salt thereof, as defined above or a pharmaceutical composition as defined above for use as a medicament.

Also provided is a method of treating cancer in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of the present disclosure, or a pharmaceutically acceptable salt thereof.


In another aspect, there is provided the use of a compound, or a pharmaceutically acceptable salt thereof, as defined above or a pharmaceutical composition as defined above in the manufacture of a medicament for the treatment of cancer.

In some examples, a method is provided of treating a Ras protein-related disorder in a subject in need thereof, the method comprising administering to the subject a therapeutically effective amount of a compound of the present disclosure, or a pharmaceutically acceptable salt thereof.

Further provided is a method of inhibiting a Ras protein in a cell, the method comprising contacting the cell with an effective amount of a compound of the present disclosure, or a pharmaceutically acceptable salt thereof.

Claims

1. A compound having the structure of Formula If:



Formula If


or pharmaceutically acceptable salt thereof,


wherein A is -N(H or CH_3)C(O)-(CH_2)- where the amino nitrogen is bound to the carbon atom of -CH(R^{10})-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or optionally substituted 5 to 6-membered heteroarylene;


B is -CH(R^9)- where the carbon is bound to the carbonyl carbon of -NHC(O)-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or 5 to 6-membered heteroarylene;


L is absent or a linker;


W is a cross-linking group comprising

a ketone of formula: ,

a vinyl ketone of formula: ,

a vinyl sulfone of formula: ,

an ynone of formula: , or

an alkynyl sulfone of formula: ;

R^1 is cyano, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 6-membered cycloalkenyl, optionally

substituted 3 to 6-membered heterocycloalkyl, optionally substituted 6 to 10-membered aryl, or optionally substituted 5 to 10-membered heteroaryl;

R^2 is C_1-C_6 alkyl, C_1-C_6 fluoroalkyl, or 3 to 6-membered cycloalkyl;

R^7 is C_1-C_3 alkyl;

R^8 is C_1-C_3 alkyl; and

R^9 is optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl;

R^{16a}, R^{16b}, and R^{16c} are, independently, hydrogen, -CN, halogen, or -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl;

R^{17} is hydrogen, -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), or a 4 to 7-membered saturated cycloalkyl, or a 4 to 7-membered saturated heterocycloalkyl;

R^{18a}, R^{18b}, and R^{18c} are, independently, hydrogen, -CN, or -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl;


R^{19} is hydrogen, -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl, or a 4 to 7-membered saturated heterocycloalkyl;

X^e is a halogen; and

R^{20} is hydrogen, -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl.

2. The compound, or pharmaceutically acceptable salt thereof, of claim 1, wherein R^1 is optionally substituted 6 to 10-membered aryl, optionally substituted 3 to 6-membered cycloalkenyl, or optionally substituted 5 to 10-membered heteroaryl.

3. The compound of claim 1, wherein the compound has the structure of Formula Ig:




or pharmaceutically acceptable salt thereof;


wherein A is -N(H or CH_3)C(O)-(CH_2)- where the amino nitrogen is bound to the carbon atom of -CH(R^{10})-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or optionally substituted 5 to 6-membered heteroarylene;


B is -CH(R^9)- where the carbon is bound to the carbonyl carbon of -NHC(O)-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or 5 to 6-membered heteroarylene;


L is absent or a linker;


W is a cross-linking group comprising

a ketone of formula ;

a vinyl ketone of formula: ,

a vinyl sulfone of formula: ,

an ynone of formula: , or

an alkynyl sulfone of formula: ;

R^2 is C_1-C_6 alkyl, C_1-C_6 fluoroalkyl, or 3 to 6-membered cycloalkyl;

R^7 is C_1-C_3 alkyl;

R^8 is C_1-C_3 alkyl; and

R^9 is optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl

X^e and X^f are, independently, N or CH; and

R^{12} is optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl;

R^{16a}, R^{16b}, and R^{16c} are, independently, hydrogen, -CN, halogen, or -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl;

R^{17} is hydrogen, -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), or a 4 to 7-membered saturated cycloalkyl, or a 4 to 7-membered saturated heterocycloalkyl;

R^{18a}, R^{18b}, and R^{18c} are, independently, hydrogen, -CN, or -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl;

R^{19} is hydrogen, -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl, or a 4 to 7-membered saturated heterocycloalkyl;

X^e is a halogen; and

R^{20} is hydrogen, -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl.

4. The compound, or pharmaceutically acceptable salt thereof, of claim 1, wherein A is optionally substituted 6-membered arylene.

5. The compound, or pharmaceutically acceptable salt thereof, of claim 1, wherein A is optionally substituted 5 to 6-membered heteroarylene.

6. The compound, or pharmaceutically acceptable salt thereof, of claim 1, wherein A is optionally substituted C_1-C_4 heteroalkylene.

7. The compound, or pharmaceutically acceptable salt thereof, of claim 1, wherein A is optionally substituted 3 to 6-membered heterocycloalkylene.

8. The compound, or pharmaceutically acceptable salt thereof, of any one of claims 1 to 5, wherein B is -CHR^9-.

9. The compound, or pharmaceutically acceptable salt thereof, of claim 8, wherein R^9 is optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl.

10. The compound, or pharmaceutically acceptable salt thereof, of any one of claims 1 to 7, wherein B is optionally substituted 6-membered arylene.

11. The compound, or pharmaceutically acceptable salt thereof, of claim 10, wherein B is 6-membered arylene.

12. The compound, or pharmaceutically acceptable salt thereof, of any one of claims 1 to 11, wherein the linker is the structure of Formula II:


A^1-$(B^1)_f$-$(C^1)_g$-$(B^2)_h$-(D^1)-$(B^3)_i$-$(C^2)_j$-$(B^4)_k$–A^2

Formula II

where A^1 is a bond between the linker and B; A^2 is a bond between W and the linker; B^1, B^2, B^3, and B^4 each, independently, is selected from optionally substituted C_1-C_2 alkylene, optionally substituted C_1-C_3 heteroalkylene, O, S, and NR^N; R^N is hydrogen, optionally substituted C_1–C_4 alkyl, optionally substituted C_2-C_4 alkenyl, optionally substituted C_2-C_4 alkynyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 6 to 10-membered aryl, or optionally substituted C_1-C_7 heteroalkyl; C^1 and C^2 are each, independently, selected from carbonyl, thiocarbonyl, sulphonyl, or phosphoryl; f, g, h, i, j, and k are each, independently, 0 or 1; and D^1 is optionally substituted C_1-C_{10} alkylene, optionally substituted C_2-C_{10} alkenylene, optionally substituted C_2-C_{10} alkynylene, optionally substituted 3 to 14-membered heterocycloalkylene, optionally substituted 5 to 10-membered heteroarylene, optionally substituted 3 to 8-membered cycloalkylene, optionally substituted 6 to 10-membered arylene, optionally substituted C_2-C_{10} polyethylene glycolene, or optionally substituted C_1-C_{10} heteroalkylene, or a chemical bond linking A^1-$(B^1)_f$-$(C^1)_g$-$(B^2)_h$- to -$(B^3)_i$-$(C^2)_j$-$(B^4)_k$–A^2.

13. The compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 12, wherein the linker is acyclic.

14. The compound, or a pharmaceutically acceptable salt thereof, of claim 13, wherein the linker has the structure of Formula IIa:



Formula IIa

wherein X^a is absent or N;


R^{14} is absent, hydrogen or optionally substituted C_1-C_6 alkyl; and

L^2 is absent, -SO_2-, optionally substituted C_1-C_4 alkylene or optionally substituted C_1-C_4 heteroalkylene,

wherein at least one of X^a, R^{14}, or L^2 is present.

15. The compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 12, wherein the linker is or comprises a cyclic moiety.

16. The compound, or a pharmaceutically acceptable salt thereof, of claim 15, wherein the linker has the structure of Formula IIb:



Formula IIb

wherein o is 0 or 1;


R^{15} is hydrogen, optionally substituted C_1-C_6 alkyl, optionally substituted 3 to 8-membered cycloalkylene, or optionally substituted 3 to 8-membered heterocycloalkylene;

X^4 is absent, optionally substituted C_1-C_4 alkylene, O, NCH_3, or optionally substituted C_1-C_4 heteroalkylene;

Cy is optionally substituted 3 to 8-membered cycloalkylene, optionally substituted 3 to 8-membered heterocycloalkylene, optionally substituted 6-10 membered arylene, or optionally substituted 5 to 10-membered heteroarylene; and

L^3 is absent, -SO_2-, optionally substituted C_1-C_4 alkylene or optionally substituted C_1-C_4 heteroalkylene.


17. The compound, or a pharmaceutically acceptable salt thereof, of claim 1, wherein W has the structure of Formula IIIa:



Formula IIIa

wherein R^{16a}, R^{16b}, and R^{16c} are, independently, hydrogen, -CN, halogen, or -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl.


18. The compound, or a pharmaceutically acceptable salt thereof, of claim 1, wherein W has the structure of Formula IIIb:



Formula IIIb

wherein R^{17} is hydrogen, -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), or a 4 to 7-membered saturated cycloalkyl, or a 4 to 7-membered saturated heterocycloalkyl.


19. The compound, or a pharmaceutically acceptable salt thereof, of claim 1, wherein W has the structure of Formula IIIc:



Formula IIIc

wherein R^{18a}, R^{18b}, and R^{18c} are, independently, hydrogen, -CN, or -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl.


20. The compound, or a pharmaceutically acceptable salt thereof, of claim 1, wherein W has the structure of Formula IIId:



Formula IIId

wherein R^{19} is hydrogen, -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl, or a 4 to 7-membered saturated heterocycloalkyl.

21. The compound, or a pharmaceutically acceptable salt thereof, of claim 1, wherein W has the structure of Formula IIIe:





Formula IIIe


wherein X^e is a halogen; and


R^{20} is hydrogen, -C_1-C_3 alkyl optionally substituted with one or more substituents independently selected from -OH, -O-C_1-C_3 alkyl, -NH_2, -NH(C_1-C_3 alkyl), -N(C_1-C_3 alkyl)$_2$, or a 4 to 7-membered saturated heterocycloalkyl.


22. A compound selected from:
















































































































































































































































assumed






assumed














































































assumed






assumed








































































































































































































































































assumed






assumed







































































































































































































































































































































































































































assumed






assumed




























































































































































































































































































































































































































































































































































































































































































































assumed



assumed





































































































































































































































































































































































assumed






assumed




























































assumed


















assumed













MeO







or a pharmaceutically acceptable salt thereof.

23. A pharmaceutical composition comprising a compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 22, and a pharmaceutically acceptable excipient.

24. A conjugate, or salt thereof, comprising the structure of Formula IV:


M-L-P

Formula IV

wherein L is a linker;

P is a monovalent organic moiety; and

M has the structure of Formula V:



Formula V

wherein the dotted lines represent zero, one, two, three, or four non-adjacent double bonds;

A is -N(H or CH_3)C(O)-(CH_2)- where the amino nitrogen is bound to the carbon atom of -CH(R^{10})-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-membered heterocycloalkylene, optionally substituted 6-membered arylene, or optionally substituted 5 to 6-membered heteroarylene;

B is absent, -CH(R^9)-, >C=C$R^9R^{9'}$, or >C$R^9R^{9'}$ where the carbon is bound to the carbonyl carbon of -N(R^{11})C(O)-, optionally substituted 3 to 6-membered cycloalkylene, optionally substituted 3 to 6-

membered heterocycloalkylene, optionally substituted 6-membered arylene, or 5 to 6-membered heteroarylene;

G is optionally substituted C_1-C_4 alkylene, optionally substituted C_1-C_4 alkenylene, optionally substituted C_1-C_4 heteroalkylene, -C(O)O-**C**H(R^6)- where **C** is bound to -C(R^7R^8)-, -C(O)NH-**C**H(R^6)- where **C** is bound to -C(R^7R^8)-, optionally substituted C_1-C_4 heteroalkylene, or 3 to 8-membered heteroarylene;

X^1 is optionally substituted C_1-C_2 alkylene, NR, O, or $S(O)_n$;

X^2 is O or NH;

X^3 is N or CH;

n is 0, 1, or 2;

R is hydrogen, cyano, optionally substituted C_1-C_4 alkyl, optionally substituted C_2-C_4 alkenyl, optionally substituted C_2-C_4 alkynyl, C(O)R', C(O)OR', C(O)N(R')$_2$, S(O)R', $S(O)_2$R', or $S(O)_2$N(R')$_2$;

each R' is, independently, H or optionally substituted C_1-C_4 alkyl;

Y^1 is C, CH, or N;

Y^2, Y^3, Y^4, and Y^7 are, independently, C or N;

Y^5 is CH, CH_2, or N;

Y^6 is C(O), CH, CH_2, or N;

R^1 is cyano, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 6-membered cycloalkenyl, optionally substituted 3 to 6-membered heterocycloalkyl, optionally substituted 6 to 10-membered aryl, or optionally substituted 5 to 10-membered heteroaryl, or

R^1 and R^2 combine with the atoms to which they are attached to form an optionally substituted 3 to 14-membered heterocycloalkyl;

R^2 is absent, hydrogen, optionally substituted C_1-C_6 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 6-membered cycloalkyl, optionally substituted 3 to 7-membered heterocycloalkyl, optionally substituted 6-membered aryl, optionally substituted 5 or 6-membered heteroaryl; R^3 is absent, or

R^2 and R^3 combine with the atom to which they are attached to form an optionally substituted 3 to 8-membered cycloalkyl or optionally substituted 3 to 14-membered heterocycloalkyl;

R^4 is absent, hydrogen, halogen, cyano, or methyl optionally substituted with 1 to 3 halogens;

R^5 is hydrogen, C_1-C_4 alkyl optionally substituted with halogen, cyano, hydroxy, or C_1-C_4 alkoxy, cyclopropyl, or cyclobutyl;

R^6 is hydrogen or methyl; R^7 is hydrogen, halogen, or optionally substituted C_1-C_3 alkyl, or

R^6 and R^7 combine with the carbon atoms to which they are attached to form an optionally substituted 3 to 6-membered cycloalkyl or optionally substituted 3 to 7-membered heterocycloalkyl;

R^8 is hydrogen, halogen, hydroxy, cyano, optionally substituted C_1-C_3 alkoxy, optionally substituted C_1-C_3 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 8-membered cycloalkyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 5 to 10-membered heteroaryl, or optionally substituted 6 to 10-membered aryl, or

R^7 and R^8 combine with the carbon atom to which they are attached to form C=C$R^{7'}R^{8'}$; C=N(OH), C=N(O-C_1-C_3 alkyl), C=O, C=S, C=NH, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl;

R^{7a} and R^{8a} are, independently, hydrogen, halo, optionally substituted C_1-C_3 alkyl, or combine with the carbon to which they are attached to form a carbonyl;

$R^{7'}$ is hydrogen, halogen, or optionally substituted C_1-C_3 alkyl; $R^{8'}$ is hydrogen, halogen, hydroxy, cyano, optionally substituted C_1-C_3 alkoxy, optionally substituted C_1-C_3 alkyl, optionally substituted C_2-C_6 alkenyl, optionally substituted C_2-C_6 alkynyl, optionally substituted 3 to 8-membered cycloalkyl, optionally substituted 3 to 14-membered heterocycloalkyl, optionally substituted 5 to 10-membered heteroaryl, or optionally substituted 6 to 10-membered aryl, or

$R^{7'}$ and $R^{8'}$ combine with the carbon atom to which they are attached to form optionally substituted 3 to 6-membered cycloalkyl or optionally substituted 3 to 7-membered heterocycloalkyl;

R^9 is H, F, optionally substituted C_1-C_6 alkyl, optionally substituted C_1-C_6 heteroalkyl, optionally substituted 3 to 6-membered cycloalkyl, or optionally substituted 3 to 7-membered heterocycloalkyl, or

R^9 and L combine with the atoms to which they are attached to form an optionally substituted 3 to 14-membered heterocycloalkyl;

$R^{9'}$ is hydrogen or optionally substituted C_1-C_6 alkyl; or

R^9 and $R^{9'}$, combined with the atoms to which they are attached, form a 3 to 6-membered cycloalkyl or a 3 to 6-membered heterocycloalkyl;

R^{10} is hydrogen, halo, hydroxy, C_1-C_3 alkoxy, or C_1-C_3 alkyl;

R^{10a} is hydrogen or halo;

R^{11} is hydrogen or C_1-C_3 alkyl; and

R^{21} is H or C_1-C_3 alkyl.

25. A compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 22 or a pharmaceutical composition of claim 23 for use as a medicament.

26. Use of a compound, or a pharmaceutically acceptable salt thereof, of any one of claims 1 to 22 or a pharmaceutical composition of claim 23 in the manufacture of a medicament for the treatment of cancer.

27. The use of claim 26, wherein the cancer is pancreatic cancer, colorectal cancer, non-small cell lung cancer, or endometrial cancer.

28. The use of claim 26 or 27, wherein the cancer comprises a Ras mutation.

29. The use of claim 28, wherein the Ras mutation is K-Ras G12C, K-Ras G13C, H-Ras G12C, H-Ras G13C, N-Ras G12C, or N-Ras G13C.

30. The use of any one of claims 26 to 28, wherein an additional anti-cancer therapy is to be administered.

31. The use of claim 30, wherein the additional anti-cancer therapy is an EGFR inhibitor, a second Ras inhibitor, a SHP2 inhibitor, a SOS1 inhibitor, a Raf inhibitor, a MEK inhibitor, an ERK

inhibitor, a PI3K inhibitor, a PTEN inhibitor, an AKT inhibitor, an mTORC1 inhibitor, a BRAF inhibitor, a PD-L1 inhibitor, a PD-1 inhibitor, a CDK4/6 inhibitor, a HER2 inhibitor, or a combination thereof.

32. The use of claim 30 or 31, wherein the additional anti-cancer therapy is a SHP2 inhibitor.