DESCRIPTION

Title of the Invention: CULTURE METHOD FOR STABLE UNDIFFERENTIATED PROLIFERATION OF PLURIPOTENT STEM CELLS

Technical Field

[0001]

The present invention relates to a culture method for proliferation of pluripotent stem cells while maintaining an undifferentiated state, and particularly relates to a culture method for stable proliferation of human pluripotent stem cells under serum-free, feeder-free conditions and under single seeding conditions while maintaining an undifferentiated state and the like.

Background Art

[0002]

Pluripotent stem cells such as ES (Embryonic stem) cell, iPS (induced pluripotent stem) cell and the like are expected to be usable for the regenerative medicine and the like in view of the superior proliferativity and pluripotency. In particular, iPS cell is considered a highly superior material for the regenerative medicine, since production and obtainment are relatively easy, there are few ethical restrictions for production, and further, from the aspect of rejection in transplantation.

[0003]

These pluripotent stem cells have conventionally been cultured by co-culture with a carrying cell (hereinafter feeder cell) such as fibroblast and the like in a medium containing serum. For example, in non-patent document 1, which is the world-first report by Yamanaka et al. on the production of iPS cell, iPS cell was established and maintained and proliferated under conditions using a feeder cell and serum. These pluripotent stem cells proliferate while forming a colony where individual cells are clustered. When a colony is dissociated into single cells and used for seeding (hereinafter single

cell-seeding), the cells become unstable. It is therefore a general practice to seed a colony maintained to have a certain size (hereinafter colony seeding). For example, non-patent document 2 discloses an example showing that the proliferation conditions of cells are prone to influence from culture environment in the case of single cell-seeding compared to colony seeding. In other words, single cell-seeding poses higher culture difficulty than colony seeding.

[0004]

To perform feeder-free culture, it is necessary to coat the bottom of a culture vessel with a substrate or scaffolding material replacing a feeder cell. As the substrate, an extracellular matrix component is often used. Patent document 1 discloses that use of an active fragment of laminin 511 as a substrate is preferable for the proliferation of human ES/iPS cell, and single cell-seeding is also possible.

[0005]

Patent document 2 and non-patent document 3 disclose the composition of a serum-free medium for human pluripotent stem cells. This composition called E8 contains DMEM/F12 as a basal medium, and further contains some factors such as bFGF, insulin and the like. At present, it is considered the minimum composition for cultivating human pluripotent stem cells.

[0006]

Ethanolamine is known to contribute to the promoted proliferation of mesenchymal stem cells when used as an additive in a medium. For example, patent document 3 discloses an example suggesting that ethanolamine promotes proliferation of mesenchymal stem cells.

[0007]

In addition, patent document 4 describes a method of maintaining primate embryonic stem cells in a medium containing ethanolamine, 2-mercaptoethanol, a complex of oleic acid with fatty acid-free bovine albumin, heparin and the like, and the like, and patent document 5 describes a medium for culturing ES

cell, which contains 2-mercaptoethanol, 2-ethanolamine, a complex of oleic acid with fatty acid-free bovine serum albumin and the like. Patent document 6 discloses a medium for ES cell, which contains human albumin, ethanolamine, β-mercaptoethanol and the like.

[0008]

The above-mentioned patent documents 4, 5 and 6 describe given amounts (10 μM, 10 μM, 100 μM, respectively) of 2-mercaptoethanol (β-mercaptoethanol) as an essential component of medium. In patent document 4, oleic acid is added to bovine albumin after removal of fatty acid, such that the bovine albumin carries oleic acid, and 9.4 mg/g of oleic acid is added to albumin.

[0009]

In the meantime, it has been reported heretofore that sulfated polysaccharides have an effect to protect growth factors from degradation, denaturation, inactivation and the like. For example, patent document 7 discloses that carrageenan stabilizes bFGF, and describes in the Examples that a protecting agent containing sulfated polysaccharides such as heparin, dextran sulfate, carrageenan and the like protects bFGF from hydrolysis and heat denaturation. However, an effect provided by a combination of ethanolamine and sulfated polysaccharides is not disclosed. Moreover, the above-mentioned patent document 4 discloses a medium integrally containing ethanolamine and heparin; however, a detailed effect of each of them has not been known to date.

[Document List]

[patent documents]

[0010]

patent document 1: JP-A-2011-78370
patent document 2: WO 2012/019122
patent document 3: JP-A-2006-325445
patent document 4: JP-A- 2009-542247
patent document 5: WO 2005/063968

patent document 6: US-B-8569061

patent document 7: WO 92/13526

[non-patent documents]

[0011]

non-patent document 1: Cell, 2006, 126, 663-76

non-patent document 2: Nature Communications, 2012, 3:1236

non-patent document 3: Nature Methods, 2011, 8, 424-429

SUMMARY OF THE INVENTION

Problems to be Solved by the Invention

[0012]

The above-mentioned conventional culture methods pose various problems when practicing a regenerative medicine using a pluripotent stem cell, particularly at an industrial level. As a feeder cell, xenogenic cells such as mouse embryo-derived fibroblast and the like are generally used, and the problem of safety after transplantation has been pointed out. Also, from the aspects of cost and cell quality management, culture without using a feeder cell (hereinafter feeder-free culture) is preferable. As for serum, the problem of infection source, and concern about property difference between lots which causes inconsistent culture results have been pointed out. As for the colony seeding, the number of seeded cells cannot be adjusted rigorously, which makes it difficult to manage culture schedule, and produces personal variation in the culture results. To produce pluripotent stem cells for a regenerative medicine at an industrial level, the work needs to be performed by plural workers under the conditions where procedures and schedule are rigorously managed. Moreover, since β-mercaptoethanol, which is generally added to a medium for pluripotent stem cell, is designated as toxin, it is preferable to not add same to a medium or reduce the amount thereof to be added as much as possible, considering the complicated handling and the like. The influence of its amount to be added on the culture of pluripotent stem cells has not been clarified in detail yet. From the foregoing, there is a demand for the development of a

culture method, which adopts serum-free, feeder-free and single cell-seeding of pluripotent stem cells, and is different from conventional culture methods. In addition, cost is a major issue at an industrial level. Therefore, a mere success in the performance of serum-free, feeder-free and single cell-seeding culture is not sufficient, and such culture affording the largest possible number of cells per unit time, namely, culture with superior proliferation efficiency, is desirable.

[0013]

The present inventors tried serum-free, feeder-free and single cell-seeding culture by a conventional culture method using the active fragment of laminin 511 described in patent document 1 as a substrate and a medium having the E8 composition. However, they could not perform stable maintenance culture for a long term. It is necessary to elucidate and overcome the cause of such instability in culture, and further improve the proliferative ability.

[0014]

It is therefore an object of the present invention to provide a means for proliferating pluripotent stem cells while maintaining an undifferentiated state. A further object of the present invention is to provide a means for stably and highly efficiently proliferating pluripotent stem cells while maintaining an undifferentiated state, in serum-free, feeder-free and single cell-seeding culture.

Means of Solving the Problems

[0015]

The present inventors have conducted intensive studies in an attempt to solve the aforementioned problems and found that a feeder cell releases ethanolamine, and ethanolamine promotes the proliferation of pluripotent stem cells. They have also found that E8 causes remarkable property degradation after thawing and preparation, which may possibly cause instability of culture using E8, and further found that albumin can suppress such property degradation. They have also found that

albumin not only has a medium stabilizing effect but also has an action to potentiate the above-mentioned cell proliferation effect of ethanolamine. They have also found that sulfated saccharides have a proliferation promoting and medium stabilizing effect in the presence of ethanolamine. The present inventors further showed that the amounts of β-mercaptoethanol, and fatty acid carried by albumin influence proliferation of pluripotent stem cells while maintaining an undifferentiated state. The present inventors have conducted further studies based on these findings and completed the present invention.

[0016]

Accordingly, the present invention is as described below.

[1] A culture method for a proliferation of a pluripotent stem cell while maintaining an undifferentiated state, comprising a step of cultivating the pluripotent stem cell in a medium added with at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof, and substantially free of β-mercaptoethanol or containing β-mercaptoethanol at a concentration of not more than 9 μM.

[2] The method of [1], wherein the concentration of at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof in the medium is 1 μM - 1000 μM.

[3] The method of [1], wherein the concentration of at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof in the medium is 5 μM - 200 μM.

[4] The method of any of [1] - [3], wherein the ethanolamine analog is a compound represented by the following formula

[0017]

$X-CH_2-CH_2-O-Y$

[0018]

wherein

X is R^1-N(R^2)- [R^1 and R^2 are the same or different and each is a hydrogen atom or an amino-protecting group] or R^3-CH=N- [R^3-CH is H-CH or a Shiff base type amino-protecting group]; and Y is -P(=O)(OH)-O-R^4 [R^4 is -CH_2-CH(O-R^5)-CH_2-O-R^6 (R^5 and R^6 are the same or different and each is an acyl group having 2 - 30 carbon atoms or a hydrogen atom) or a hydrogen atom], a hydrogen atom or a hydroxy-protecting group.

[5] The method of [4], wherein R^1 and R^2 are the same or different and each is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms, and R^3 is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms.

[6] The method of any of [1] - [3], wherein the ethanolamine analog is one or plural selected from the group consisting of phosphoethanolamine, monomethylethanolamine, dimethylethanolamine, N-acylphosphatidylethanolamine, phosphatidylethanolamine, and lysophosphatidylethanolamine.

[7] The method of any of [1] - [6], wherein the medium is further added with albumin.

[8] The method of [7], wherein the concentration of the albumin in the medium is 0.1 g/l - 20 g/l.

[9] The method of [7], wherein the concentration of the albumin in the medium is 1 g/l - 8 g/l.

[10] The method of any of [7] - [9], wherein the albumin is obtained from plasma of an animal (including human) or by gene

recombination technology.

[11] The method of any of [7] - [10], wherein the amount of fatty acid carried by the albumin in the medium is not more than 9 mg/g.

[12] The method of any of [7] - [10], wherein the amount of fatty acid carried by the albumin in the medium is not more than 2.2 mg/g.

[13] The method of any of [1] - [12], wherein the medium is further added with sulfated saccharide and/or a pharmaceutically acceptable salt thereof.

[14] The method of [13], wherein the concentration of the sulfated saccharide and/or a pharmaceutically acceptable salt thereof in the medium is 1 - 1000 ng/ml.

[15] The method of [13] or [14], wherein the sulfated saccharide or a pharmaceutically acceptable salt thereof is dextran sulfate Na having an average molecular weight of 2,500 - 7,500.

[16] A culture method for a proliferation of a pluripotent stem cell while maintaining an undifferentiated state, comprising a step of cultivating the pluripotent stem cell in a medium added with at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof, and added with sulfated saccharide and/or a pharmaceutically acceptable salt thereof.

[17] The method of [16], wherein the concentration of the sulfated saccharide and/or a pharmaceutically acceptable salt thereof in the medium is 1 - 1000 ng/ml.

[18] The method of [16] or [17], wherein the sulfated saccharide or a pharmaceutically acceptable salt thereof is dextran sulfate Na having an average molecular weight of 2,500 - 7,500.

[19] The method of any of [16] - [18], wherein the medium is further added with albumin.

[20] The method of [19], wherein the concentration of the albumin in the medium is 0.1 g/l - 20 g/l.

[21] The method of [19], wherein the concentration of the albumin in the medium is 1 g/l - 8 g/l.

[22] The method of any of [19] - [21], wherein the albumin is obtained from plasma of an animal (including human) or by gene recombination technology.

[23] The method of any of [1] - [22], wherein the culture is performed in the absence of a feeder cell.

[24] The method of [23], wherein the culture is performed by using an extracellular matrix or an active fragment thereof, or an artificial product mimicking the function thereof.

[25] The method of [23], wherein the culture is performed by using laminin 511 or an active fragment thereof or matrigel.

[26] The method of any of [1] - [25], wherein the culture is performed by single cell-seeding.

[27] The method of any of [1] - [26], wherein the culture is performed under serum-free conditions.

[28] The method of any of [1] - [27], wherein the medium is substantially free of a component derived from an animal other than human.

[29] The method of any of [1] - [28], wherein the pluripotent stem cell is an embryonic stem cell (ES cell) or induced pluripotent stem cell (iPS cell).

[30] The method of any of [1] - [29], wherein the pluripotent stem cell is derived from primates.

[31] The method of any of [1] - [30], wherein the pluripotent stem cell is human iPS cell.

[32] A preservation stabilizing method for a medium for a proliferation of a pluripotent stem cell, comprising adding at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof, and adding sulfated saccharide and/or a pharmaceutically acceptable salt thereof.

[33] The method of [32], wherein the final concentration of the sulfated saccharide and/or a pharmaceutically acceptable salt thereof in use is 1 - 1000 ng/ml.

[34] The method of [32] or [33], further comprising adding albumin.

[35] The method of [34], wherein the final concentration of the albumin in use is 0.1 g/l - 20 g/l.

[36] The method of [34], wherein the final concentration of the albumin in use is 1 g/l - 8 g/l.

[37] The method of any of [34] - [36], wherein the albumin is obtained from plasma of an animal (including human) or by gene recombination technology.

[38] The method of any of [34] - [37], wherein the amount of fatty acid carried by the albumin is not more than 9 mg/g.

[39] The method of any of [34] - [37], wherein the amount of fatty acid carried by the albumin is not more than 2.2 mg/g.

[40] The method of any of [32] - [39], wherein the medium is substantially free of β-mercaptoethanol or contains β-mercaptoethanol at a final concentration of not more than 9 μM.

[41] The method of any of [32] - [40], wherein the medium is for a proliferation of a pluripotent stem cell while maintaining an undifferentiated state.

[42] A medium additive for a proliferation of a pluripotent stem cell while maintaining an undifferentiated state, comprising at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof, and substantially free of β-mercaptoethanol or containing β-mercaptoethanol at a concentration of not more than 9 μM when in use.

[43] The medium additive of [42], wherein the ethanolamine analog is a compound represented by the following formula

[0019]

$X\text{-}CH_2\text{-}CH_2\text{-}O\text{-}Y$

[0020]

wherein

X is $R^1\text{-}N(R^2)\text{-}$ [R^1 and R^2 are the same or different and each is a hydrogen atom or an amino-protecting group] or $R^3\text{-}CH{=}N\text{-}$ [R^3-CH is H-CH or a Shiff base type amino-protecting group]; and

Y is $-P(=O)(OH)-O-R^4$ [R^4 is $-CH_2-CH(O-R^5)-CH_2-O-R^6$ (R^5 and R^6 are the same or different and each is an acyl group having 2 - 30 carbon atoms or a hydrogen atom) or a hydrogen atom], a hydrogen atom or a hydroxy-protecting group.

[44] The medium additive of [43], wherein R^1 and R^2 are the same or different and each is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms, and R^3 is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms.

[45] The medium additive of any of [42] - [44], wherein the ethanolamine analog is one or plural selected from the group consisting of phosphoethanolamine, monomethylethanolamine, dimethylethanolamine, N-acylphosphatidylethanolamine, phosphatidylethanolamine, and lysophosphatidylethanolamine.

[46] The medium additive of any of [42] - [45], further comprising albumin.

[47] The medium additive of [46], wherein the final concentration of the albumin in use is 0.1 g/l - 20 g/l.

[48] The medium additive of [46], wherein the final concentration of the albumin in use is 1 g/l - 8 g/l.

[49] The medium additive of any of [46] - [48], wherein the amount of fatty acid carried by the albumin is not more than 9 mg/g.

[50] The medium additive of any of [46] - [48], wherein the amount of fatty acid carried by the albumin is not more than

2.2 mg/g.

[51] The medium additive of any of [46] - [50], wherein the albumin is obtained from plasma of an animal (including human) or by gene recombination technology.

[52] The medium additive of any of [42] - [51], further comprising sulfated saccharide and/or a pharmaceutically acceptable salt thereof.

[53] The medium additive of [52], wherein the final concentration of the sulfated saccharide and/or a pharmaceutically acceptable salt thereof in use is 1 - 1000 ng/ml.

[54] The medium additive of [52] or [53], wherein the sulfated saccharide is at least one selected from the group consisting of sulfated monosaccharide, sulfated disaccharide, sulfated polysaccharide, sulfated sugar alcohol and sulfated cyclitol.

[55] The medium additive of any of [52] - [54], wherein the aforementioned sulfated saccharide or a pharmaceutically acceptable salt thereof is at least one selected from the group consisting of dextran sulfate Na, cellulose SO_3Na, xanthan gum SO_3Na, fucoidan, alginate SO_3Na, inulin SO_3Na, maltoheptaose SO_3Na, stachyose SO_3Na, maltotriose SO_3Na, multitol SO_3Na, sucrose8SO_3K, glucose SO_3Na, myo-6 inositol SO_3K, α-cyclodextrin SO_3Na, mannitol SO_3Na, xylitol SO_3Na and erythritol SO_3Na.

[56] The medium additive of [55], wherein the aforementioned sulfated saccharide or a pharmaceutically acceptable salt thereof is at least one selected from the group consisting of dextran sulfate Na, fucoidan, maltoheptaose SO_3Na, maltotriose SO_3Na, multitol SO_3Na and sucrose8SO_3K.

[57] The medium additive of any of [52] - [56], wherein the aforementioned sulfated saccharide or a pharmaceutically acceptable salt thereof is dextran sulfate Na having an average molecular weight of 2,500 - 7,500.

[58] The medium additive of any of [42] - [57], wherein the proliferation of a pluripotent stem cell while maintaining an undifferentiated state is performed under the conditions

without using a feeder cell.

[59] The medium additive of [58], wherein the conditions without using a feeder cell do not include use of a feeder cell but include use of an extracellular matrix or an active fragment thereof, or an artificial product mimicking the function thereof.

[60] The medium additive of [58], wherein the conditions without using a feeder cell do not include use of a feeder cell but include use of laminin 511 or an active fragment thereof or matrigel.

[61] The medium additive of any of [42] - [60], wherein the proliferation of a pluripotent stem cell while maintaining an undifferentiated state is performed by single cell-seeding.

[62] The medium additive of any of [42] - [61], wherein the proliferation of a pluripotent stem cell while maintaining an undifferentiated state is performed under serum-free conditions.

[63] The medium additive of any of [42] - [62], comprising substantially free of a component derived from an animal other than human.

[64] The medium additive of any of [42] - [63], wherein the pluripotent stem cell is an embryonic stem cell (ES cell) or induced pluripotent stem cell (iPS cell).

[65] The medium additive of any of [42] - [64], wherein the pluripotent stem cell is derived from primates.

[66] The medium additive of any of [42] - [65], wherein the pluripotent stem cell is human iPS cell.

[67] A medium for proliferation of a pluripotent stem cell while maintaining an undifferentiated state, which comprises the medium additive of any of [42] - [66].

[68] A culture method for a proliferation of an induced pluripotent stem cell (iPS cell) while maintaining an undifferentiated state, comprising a step of cultivating the induced pluripotent stem cell (iPS cell) in a medium added with at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts

thereof.

[69] The method of [68], wherein the concentration of at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof in the medium is 1 μM - 1000 μM.

[70] The method of [68], wherein the concentration of at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof in the medium is 5 μM - 200 μM.

[71] The method of any of [68] - [70], wherein the ethanolamine analog is a compound represented by the following formula

[0021]

$X-CH_2-CH_2-O-Y$

[0022]

wherein

X is $R^1-N(R^2)-$ [R^1 and R^2 are the same or different and each is a hydrogen atom or an amino-protecting group] or $R^3-CH=N-$ [R^3-CH is H-CH or a Shiff base type amino-protecting group]; and

Y is $-P(=O)(OH)-O-R^4$ [R^4 is $-CH_2-CH(O-R^5)-CH_2-O-R^6$ (R^5 and R^6 are the same or different and each is an acyl group having 2 - 30 carbon atoms or a hydrogen atom) or a hydrogen atom], a hydrogen atom or a hydroxy-protecting group.

[72] The method of [71], wherein R^1 and R^2 are the same or different and each is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms, and

R^3 is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group

having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms.

[73] The method of any of [68] - [72], wherein the ethanolamine analog is one or plural selected from the group consisting of phosphoethanolamine, monomethylethanolamine, dimethylethanolamine, N-acylphosphatidylethanolamine, phosphatidylethanolamine, and lysophosphatidylethanolamine.

[74] The method of any of [68] - [73], wherein the medium is further added with albumin.

[75] The method of [74], wherein the concentration of the albumin in the medium is 0.1 g/l - 20 g/l.

[76] The method of [74], wherein the concentration of the albumin in the medium is 1 g/l - 8 g/l.

[77] The method of any of [74] - [76], wherein the albumin is obtained from plasma of an animal (including human) or by gene recombination technology.

[78] The method of any of [74] - [77], wherein the amount of fatty acid carried by the albumin in the medium is not more than 9 mg/g.

[79] The method of any of [74] - [77], wherein the amount of fatty acid carried by the albumin in the medium is not more than 2.2 mg/g.

[80] The method of any of [68] - [79], wherein the culture is performed in the absence of a feeder cell.

[81] The method of [80], wherein the culture is performed by using an extracellular matrix or an active fragment thereof, or an artificial product mimicking the function thereof.

[82] The method of [80], wherein the culture is performed by using laminin 511 or an active fragment thereof or matrigel.

[83] The method of any of [68] - [82], wherein the culture is performed by single cell-seeding.

[84] The method of any of [68] - [83], wherein the culture is performed under serum-free conditions.

[85] The method of any of [68] - [84], wherein the medium is substantially free of a component derived from an animal other

than human.

[86] The method of any of [68] - [85], wherein the induced pluripotent stem cell is derived from primates.

[87] The method of any of [68] - [86], wherein the induced pluripotent stem cell is human iPS cell.

[88] A preservation stabilizing method for a medium for a proliferation of a pluripotent stem cell, comprising adding at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof.

[89] The method of [88], further comprising adding sulfated saccharide and/or a pharmaceutically acceptable salt thereof.

[90] The method of [89], wherein the final concentration of the sulfated saccharide and/or a pharmaceutically acceptable salt thereof in use is 1 - 1000 ng/ml.

[91] The method of any of [88] - [90], further comprising adding albumin.

[92] The method of [91], wherein the final concentration of the albumin in use is 0.1 g/l - 20 g/l.

[93] The method of [91], wherein the final concentration of the albumin in use is 1 g/l - 8 g/l.

[94] The method of any of [91] - [93], wherein the albumin is obtained from plasma of an animal (including human) or by gene recombination technology.

[95] The method of any of [91] - [94], wherein the amount of fatty acid carried by the albumin is not more than 9 mg/g.

[96] The method of any of [91] - [94], wherein the amount of fatty acid carried by the albumin is not more than 2.2 mg/g.

[97] The method of any of [88] - [96], wherein the medium is substantially free of β-mercaptoethanol or contains β-mercaptoethanol at a final concentration of not more than 9 μM.

[98] The method of any of [88] - [97], wherein the medium is for a proliferation of a pluripotent stem cell while maintaining an undifferentiated state.

[99] A medium additive for a proliferation of a pluripotent

stem cell while maintaining an undifferentiated state, comprising at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof.

[100] The medium additive of [99], wherein the ethanolamine analog is a compound represented by the following formula

[0023]

$X-CH_2-CH_2-O-Y$

[0024]

wherein

X is $R^1-N(R^2)-$ [R^1 and R^2 are the same or different and each is a hydrogen atom or an amino-protecting group] or $R^3-CH=N-$ [R^3-CH is H-CH or a Shiff base type amino-protecting group]; and

Y is $-P(=O)(OH)-O-R^4$ [R^4 is $-CH_2-CH(O-R^5)-CH_2-O-R^6$ (R^5 and R^6 are the same or different and each is an acyl group having 2 - 30 carbon atoms or a hydrogen atom) or a hydrogen atom], a hydrogen atom or a hydroxy-protecting group.

[101] The medium additive of [100], wherein R^1 and R^2 are the same or different and each is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms, and

R^3 is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms.

[102] The medium additive of any of [99] - [101], wherein the ethanolamine analog is one or plural selected from the group consisting of phosphoethanolamine, monomethylethanolamine,

dimethylethanolamine, N-acylphosphatidylethanolamine, phosphatidylethanolamine, and lysophosphatidylethanolamine.

[103] The medium additive of any of [99] - [102], further comprising albumin.

[104] The medium additive of [103], wherein the albumin is obtained from plasma of an animal (including human) or by gene recombination technology.

[105] The medium additive of any of [99] - [104], further comprising sulfated saccharide and/or a pharmaceutically acceptable salt thereof.

[106] The medium additive of [105], wherein the final concentration of the sulfated saccharide and/or a pharmaceutically acceptable salt thereof in use is 1 - 1000 ng/ml.

[107] The medium additive of [105] or [106], wherein the sulfated saccharide is at least one selected from the group consisting of sulfated monosaccharide, sulfated disaccharide, sulfated polysaccharide, sulfated sugar alcohol and sulfated cyclitol.

[108] The medium additive of any of [105] - [107], wherein the aforementioned sulfated saccharide or a pharmaceutically acceptable salt thereof is at least one selected from the group consisting of dextran sulfate Na, cellulose SO_3Na, xanthan gum SO_3Na, fucoidan, alginate SO_3Na, inulin SO_3Na, Maltoheptaose SO_3Na, stachyose SO_3Na, maltotriose SO_3Na, multitol SO_3Na, sucrose8SO_3K, glucose SO_3Na, myo-6 inositol SO_3K, α-cyclodextrin SO_3Na, mannitol SO_3Na, xylitol SO_3Na and erythritol SO_3Na.

[109] The medium additive of [108], wherein the aforementioned sulfated saccharide or a pharmaceutically acceptable salt thereof is at least one selected from the group consisting of dextran sulfate Na, fucoidan, Maltoheptaose SO_3Na, maltotriose SO_3Na, multitol SO_3Na and sucrose8SO_3K.

[110] The medium additive of any of [105] - [109], wherein the aforementioned sulfated saccharide or a pharmaceutically acceptable salt thereof is dextran sulfate Na having an average

molecular weight of 2,500 - 7,500.

[111] The medium additive of any of [99] - [110], wherein the proliferation of a pluripotent stem cell while maintaining an undifferentiated state is performed under the conditions without using a feeder cell.

[112] The medium additive of [111], wherein the conditions without using a feeder cell do not include use of a feeder cell but include use of an extracellular matrix or an active fragment thereof, or an artificial product mimicking the function thereof.

[113] The medium additive of [111], wherein the conditions without using a feeder cell do not include use of a feeder cell but include use of laminin 511 or an active fragment thereof or matrigel.

[114] The medium additive of any of [99] - [113], wherein the proliferation of a pluripotent stem cell while maintaining an undifferentiated state is performed by single cell-seeding.

[115] The medium additive of any of [99] - [114], wherein the proliferation of a pluripotent stem cell while maintaining an undifferentiated state is performed under serum-free conditions.

[116] The medium additive of any of [99] - [115], comprising substantially free of a component derived from an animal other than human.

[117] The medium additive of any of [99] - [116], wherein the pluripotent stem cell is an embryonic stem cell (ES cell) or induced pluripotent stem cell (iPS cell).

[118] The medium additive of any of [99] - [117], wherein the pluripotent stem cell is derived from primates.

[119] The medium additive of any of [99] - [118], wherein the pluripotent stem cell is human iPS cell.

[120] A medium for proliferation of a pluripotent stem cell while maintaining an undifferentiated state, which comprises the medium additive of any of [99] - [119].

Effect of the Invention

[0025]

According to the present invention, pluripotent stem cells can be stably and efficiently proliferated, and can be stably proliferated for a long term while maintaining an undifferentiated state even in serum-free, feeder-free and single cell-seeding culture.

Brief Description of the Drawings

[0026]

Fig. 1 shows a medium stabilizing effect by albumin under preservation conditions at 4°C.

Fig. 2 shows a cell proliferation promoting effect of ethanolamine and an effect of combination with albumin.

Fig. 3 shows a cell proliferation promoting effect by a combination of ethanolamine and dextran sulfate sodium, which is expressed by a cumulative viable cell increase rate (fold).

Fig. 4 shows the results of alkaline phosphatase staining of iPS cell colonies after long-term culture in an E8 minimum composition medium added with albumin, ethanolamine and dextran sulfate sodium.

Fig. 5 shows a cell proliferation promoting effect of ethanolamine in culture using matrigel as a basal lamina matrix.

Fig. 6 shows that medium change can be omitted in culture of iPS cells, by adding dextran sulfate sodium to a medium containing albumin and ethanolamine.

Fig. 7 shows a medium stabilizing effect of a combination of ethanolamine and dextran sulfate sodium under room temperature preservation conditions.

Fig. 8 shows a cell proliferation promoting effect of O-phosphoryl ethanolamine.

Fig. 9 shows a cell proliferation promoting effect of 2-(methylamino)ethanol.

Fig. 10 shows a cell proliferation promoting effect of 2-dimethylaminoethanol.

Fig. 11 shows a cell proliferation promoting effect of ethanolamine hydrochloride.

Fig. 12 shows a cell proliferation suppressive effect

along with an increase in the amount of oleic acid carried by albumin.

Description of Embodiments

[0027]

(culture method for proliferation of pluripotent stem cells while maintaining an undifferentiated state)

The present invention provides a culture method for the proliferation of a pluripotent stem cell while maintaining an undifferentiated state, comprising a step of cultivating a pluripotent stem cell in a medium added with at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof, and substantially free of β-mercaptoethanol or containing β-mercaptoethanol at a concentration of not more than 9 μM (hereinafter to be also referred to as the culture method of the present invention).

The present invention is based on the new finding that a feeder cell releases ethanolamine, and addition of ethanolamine can promote proliferation of pluripotent stem cells while maintaining an undifferentiated state even in feeder-free culture. As a feeder cell, xenogenic cells are generally used, and incidence of infection with a feeder cell-derived virus is also known. The present invention can solve these problems, and is extremely useful in the field of regenerative medicine.

Also, the present invention is based on the finding that pluripotent stem cells can be proliferated well while maintaining an undifferentiated state even when the concentration of β-mercaptoethanol in the medium is reduced. In the culture method of the present invention, the medium is preferably substantially or completely free of β-mercaptoethanol. Even when the medium contains β-mercaptoethanol, its concentration is preferably not more than 9 μM.

[0028]

In the present specification, being "substantially free

of β-mercaptoethanol" means that the content concentration of β-mercaptoethanol is below detection limit.

[0029]

In the present invention, the "pluripotent stem cell" means an immature cell having self-renewal capacity and differentiation/proliferation capacity, which is capable of differentiating into any tissue or cell constituting living organisms. Examples of the pluripotent stem cell include embryonic stem cells (ES cell), embryonic germ cell (EG cell), induced pluripotent stem cell (iPS cell) and the like. A stem cell established by cultivating an early embryo generated by nuclear transplantation of the nucleus of a somatic cell is also included in the pluripotent stem cell (Nature, 385, 810 (1997); Science, 280, 1256 (1998); Nature Biotechnology, 17, 456 (1999); Nature, 394, 369 (1998); Nature Genetics, 22, 127 (1999); Proc. Natl. Acad. Sci. USA, 96, 14984 (1999); Nature Genetics, 24, 109 (2000)). The pluripotent stem cell in the present invention does not include a multipotent stem cell. The multipotent stem cell means a cell capable of differentiating into plural, though not all, types of tissues and cells and includes somatic stem cells such as mesenchymal stem cell and the like.

[0030]

While the culture method of the present invention can be preferably used for any pluripotent stem cells, it is preferably used for proliferation of embryonic stem cell (ES cell) or induced pluripotent stem cell (iPS cell) while maintaining an undifferentiated state.

[0031]

Also, the culture method of the present invention can be preferably used for pluripotent stem cells derived from any animals. The pluripotent stem cells cultured by using the medium of the present invention are, for example, pluripotent stem cells derived from rodents such as mouse, rat, hamster, guinea pig and the like, Lagomorpha such as rabbit and the like,

Ungulata such as swine, bovine, goat, horse, sheep and the like, Carnivora such as dog, cat and the like, primates such as human, monkey, Macaca mulatta, marmoset, orangutan, chimpanzee and the like. Preferred are pluripotent stem cells derived from primates. When it is used for regenerative medicine, human iPS cells are preferable.

[0032]

In the present invention, the "proliferation while maintaining an undifferentiated state" of pluripotent stem cells means that pluripotent stem cells can proliferate in an undifferentiated state while maintaining the pluripotency. That is, the culture method of the present invention can also be said a method of proliferating pluripotent stem cells in an undifferentiated state while maintaining pluripotency. Whether a pluripotent stem cell is maintained in an undifferentiated state is confirmed by alkaline phosphatase staining as shown in the below-mentioned Examples. Stained cells are evaluated as being maintained in an undifferentiated state.

[0033]

The "ethanolamine" (also called 2-aminoethanol, monoethanolamine) used in the present invention may be isolated and purified from a natural product or a processed product thereof, or a synthesized product. Ethanolamine can be produced by reacting ethylene oxide and ammonia. Ethanolamine can also be isolated and purified from a natural product or a processed product thereof by known techniques such as solvent extraction, various chromatographys and the like. Ethanolamine may be a commercially available product and can be obtained from, for example, Sigma-Aldrich Co., Ltd. and the like.

[0034]

Examples of the "ethanolamine analog" to be used in the present invention include a compound represented by the following formula

[0035]

$X-CH_2-CH_2-O-Y$

[0036]

wherein

X is R^1-N(R^2)- [R^1 and R^2 are the same or different and each is a hydrogen atom or an amino-protecting group] or R^3-CH=N- [R^3-CH is H-CH or a Shiff base type amino-protecting group]; and

Y is -P(=O)(OH)-O-R^4 [R^4 is -CH_2-CH(O-R^5)-CH_2-O-R^6 (R^5 and R^6 are the same or different and each is an acyl group having 2 - 30 carbon atoms or a hydrogen atom) or a hydrogen atom], a hydrogen atom or a hydroxy-protecting group].

[0037]

As for the "amino-protecting group", for example, books such as Green et al., Protective Groups in Organic Synthesis, 3rd Edition, 1999, John Wiley & Sons, Inc. and the like can be referred to, and an appropriate protecting group can be selected, introduced and removed. Examples of the "amino-protecting group" include a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms. Furthermore, a leaving group that can be bound to an amino group for forming a prodrug can also be mentioned.

[0038]

The "halogen atom" means a fluorine atom, a chlorine atom, a bromine atom, or an iodine atom.

[0039]

Examples of the "aryl group" include phenyl, 1-naphthyl, 2-naphthyl and the like.

[0040]

Examples of the "acyl group having 2 - 30 carbon atoms" include a saturated carboxylic acid acyl group and an unsaturated carboxylic acid acyl group. Examples of the saturated carboxylic acid acyl group include acetyl(ethanoyl),

propanoyl, butanoyl, pentanoyl, hexanoyl, heptanoyl, octanoyl, nonanoyl, decanoyl, undecanoyl, dodecanoyl, tridecanoyl, tetradecanoyl, pentadecanoyl, hexadecanoyl, heptadecanoyl, octadecanoyl, nonadecanoyl, icosanoyl, eicosanoyl, Henicosanoyl, Heneicosanoyl, docosanoyl , tricosanoyl, tetracosanoyl, pentacosanoyl, hexacosanoyl, heptacosanoyl, octacosanoyl, nonacosanoyl, triacontanoyl and the like. Examples of the unsaturated carboxylic acid acyl group include acryloyl, methacryloyl, crotonoyl, isocrotonoyl, butenoyl, butadienoyl, pentenoyl, hexenoyl, heptenoyl, octenoyl, nonenoyl, decenoyl, undecenoyl, dodecenoyl, tetradecenoyl, oleloyl, elaidinoyl, cyclopentanoyl, cyclohexanoyl, cycloheptanoyl, methylcyclopentanoyl, methylcyclohexanoyl, methylcycloheptanoyl, cyclopentenoyl, 2,4-cyclopentadienoyl, cyclohexenoyl, 2,4-cyclohexadienoyl, cycloheptenoyl, methylcyclopentenoyl, methylcyclohexenoyl, methylcycloheptenoyl and the like.

[0041]

The "alkyl group having 1 - 6 carbon atoms" means a linear or branched alkyl group having 1 - 6 carbon atoms. Specific examples thereof include methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, tert-pentyl, neopentyl, 2-pentyl, 3-pentyl, n-hexyl, 2-hexyl and the like.

[0042]

The "alkoxyl group having 1 - 6 carbon atoms" means a linear or branched alkoxyl group having 1 - 6 carbon atoms. Specific examples thereof include a methoxy group, an ethoxy group, a propoxy group, a butoxy group, a pentyloxy group, a hexyloxy group, a methoxymethoxy group, a methoxyethoxy group, a methoxypropoxy group, an ethoxyethoxy group, an ethoxypropoxy group and the like.

[0043]

The "hydroxyalkyl group having 1 - 6 carbon atoms" means a linear or branched hydroxyalkyl group having 1 - 6 carbon atoms, wherein a part of the hydrogen atom in the alkyl group

is substituted by a hydroxyl group. Specific examples thereof include a hydroxymethyl group, a 2-hydroxyethyl group, a 3-hydroxypropyl group, a 2-hydroxypropyl group, a 2-hydroxy-2-methylpropyl group, a 4-hydroxybutyl group, a 3-hydroxybutyl group, a 2-hydroxybutyl group and the like.

[0044]

The "haloalkyl group having 1 - 6 carbon atoms" means a linear or branched haloalkyl group having 1 - 6 carbon atoms, wherein a part of the hydrogen atom in the alkyl group is substituted by a halogen atom, and the halogen atom is a fluorine atom, a chlorine atom, a bromine atom or an iodine atom. Specific examples thereof include a trifluoromethyl group, a chloromethyl group, a bromomethyl group, a dichloromethyl group, a difluoromethyl group, a trichloromethyl group, a 2-fluoroethyl group, a 2-chloroethyl group, a 2-bromoethyl group, a 1,1-difluoroethyl group, a 2,2,2-trifluoroethyl group, a 3-chloropropyl group, a 3-iodopropyl group and the like.

[0045]

The "haloalkoxyl group having 1 - 6 carbon atoms" means a linear or branched haloalkoxyl group having 1 - 6 carbon atoms, wherein a part of the hydrogen atom in the alkoxy group is substituted by a halogen atom. Specific examples thereof include a trifluoromethoxy group, a pentafluoroethoxy group, a 2-chloroethoxy group, a 2,2,2-trifluoroethoxy group, a heptafluoro-n-propoxy group, a heptafluoro-i-propoxy group, a 1,1,1,3,3,3-hexafluoro-2-propoxy group, a 3-fluoro-n-propoxy group, a 1-chlorocyclopropoxy group, a 2-bromocyclopropoxy group, a 3,3,4,4,4-pentafluoro-2-butoxy group, a noanfluoro-n-butoxy group, a nonafluoro-2-butoxy group, a 5,5,5-trifluoro-n-pentyloxy group, a 4,4,5,5,5-pentafluoro-2-pentyloxy group, a 3-chloro-n-pentyloxy group, a 4-bromo-2-pentyloxy group, a 4-chlorobutyloxy group, a 2-iodo-n-propyloxy group and the like.

[0046]

The "halohydroxyalkyl group having 1 - 6 carbon atoms"

means a linear or branched halohydroxyalkyl group having 1 - 6 carbon atoms, wherein a part of the hydrogen atom in the hydroxylalkyl group is substituted by a halogen atom. Specific examples thereof include difluorohydroxymethyl, 1,1-difluoro-2-hydroxyethyl, 2,2-difluoro-2-hydroxyethyl, 1,1,2,2-tetrafluoro-2-hydroxyethyl group and the like.

[0047]

The "forming a prodrug" in the "leaving group that can be bound to an amino group for forming a prodrug" means conversion of the target compound such that the compound which shows a small or no effect of the present invention shows the effect of the present invention upon removal of the leaving group in a medium and/or during culture. It means formation of a temporary bond between the amino group and the leaving group in the compound showing the effect of the present invention, which is removed in a medium and/or during culture.

The "leaving group that can be bound to an amino group for forming a prodrug" is not particularly limited as long as it is used in the field of synthetic organic chemistry. Examples thereof include a halogen atom (e.g., chlorine atom, bromine atom, iodine atom etc.), a sulfonyloxy group (e.g., methanesulfonyloxy group, trifluoromethanesulfonyloxy group, benzenesulfonyloxy group, p-toluenesulfonyloxy group etc.) and the like.

[0048]

When R^3-CH is a Shiff base type amino-protecting group, R^3 is a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms.

[0049]

While the "hydroxyl-protecting group" is not particularly

limited as long as it is a hydroxyl-protecting group to be used in the field of synthetic organic chemistry, for example, an alkyl group having 1 - 6 carbon atoms (e.g., methyl, ethyl, propyl, isopropyl, butyl, tert-butyl), a phenyl group, a trityl group, an aralkyl group having 7 - 10 carbon atoms (e.g., benzyl, p-methoxybenzyl), a formyl group, an alkyl-carbonyl group having 1 - 6 carbon atoms (e.g., acetyl, propionyl), a benzoyl group, an aralkyl-carbonyl group having 7 - 10 carbon atoms (e.g., benzylcarbonyl), methoxymethyl, ethoxyethyl, a 2-tetrahydropyranyl group, a 2-tetrahydrofuranyl group, substituted silyl group (e.g., trimethylsilyl, triethylsilyl, dimethylphenylsilyl, tert-butyldimethylsilyl, tert-butyldiethylsilyl), an alkenyl group having 2 - 6 carbon atoms (e.g., 1-allyl) and the like can be mentioned. Furthermore, a leaving group that can be bound to a hydroxy group for forming a prodrug can also be mentioned.

[0050]

The "forming a prodrug" in the "leaving group that can be bound to a hydroxy group for forming a prodrug" means conversion of the target compound such that the compound which shows a small or no effect of the present invention shows the effect of the present invention upon removal of the leaving group in a medium and/or during culture. It means formation of a temporary bond between the hydroxy group and the leaving group in the compound showing the effect of the present invention, which is removed in a medium and/or during culture.

The "leaving group that can be bound to a hydroxy group for forming a prodrug" is not particularly limited as long as it is used in the field of synthetic organic chemistry. Examples thereof include a halogen atom (e.g., chlorine atom, bromine atom, iodine atom etc.), a sulfonyloxy group (e.g., methanesulfonyloxy group, trifluoromethanesulfonyloxy group, benzenesulfonyloxy group, p-toluenesulfonyloxy group etc.) and the like.

[0051]

The ethanolamine analog is preferably one or plural selected from the group consisting of phosphoethanolamine (aka phosphoryl ethanolamine), monomethylethanolamine, dimethylethanolamine, N-acylphosphatidylethanolamine, phosphatidylethanolamine, and lysophosphatidylethanolamine.

[0052]

The ethanolamine and/or ethanolamine analog to be used in the present invention may be in the form of a pharmaceutically acceptable salt. Examples of such salt when an acidic group such as a carboxyl group and the like is present in the compound include salts with alkali metals such as ammonium salt, sodium, potassium and the like, salts with alkaline earth metals such as calcium, magnesium and the like, salts with organic amines such as aluminum salt, zinc salt, triethylamine, morpholine, piperidine, dicyclohexylamine and the like, and salts with basic amino acids such as arginine, lysine and the like. Examples thereof when a basic group is present in the compound include salts with inorganic acids such as hydrochloric acid, sulfuric acid, phosphoric acid, nitric acid, hydrobromic acid and the like, salts with organic carboxylic acids such as acetic acid, trifluoroacetic acid, citric acid, benzoic acid, maleic acid, fumaric acid, tartaric acid, succinic acid, tannic acid, butyric acid, hibenzoic acid, pamoic acid, enanthic acid, decanoic acid, teoclic acid, salicylic acid, lactic acid, oxalic acid, mandelic acid, malic acid and the like, and salts with organic sulfonic acids such as methanesulfonic acid, benzenesulfonic acid, p-toluenesulfonic acid and the like. Particularly preferred is hydrochloride.

[0053]

The final concentration (concentration when in use) of ethanolamine and/or ethanolamine analogs and pharmaceutically acceptable salts thereof in the medium can be determined to fall within any range as long as it can promote proliferation of pluripotent stem cells while maintaining an undifferentiated

state under feeder-free conditions. While the final concentration can vary depending on the kind thereof, it is generally 1 μM - 1000 μM, preferably 5 μM - 200 μM, or 11 μM - 200 μM. When it is less than 1 μM, the effect of promoting proliferation of pluripotent stem cells while maintaining an undifferentiated state tends to be weak. When it exceeds 1000 μM, proliferation of pluripotent stem cells may sometimes be suppressed. The concentration of at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof in the medium is generally 1 μM - 1000 μM, preferably 5 μM - 200 μM, or 11 μM - 200 μM. When plural kinds are used, the total amount thereof is set to fall within the above-mentioned range. However, it can be increased or decreased as appropriate depending on the number of the kinds thereof.

[0054]

In the present invention, that at least one selected from the group consisting of ethanolamine and/or ethanolamine analogs and pharmaceutically acceptable salts thereof "promotes proliferation of pluripotent stem cells while maintaining an undifferentiated state under feeder-free conditions" means that the number of cells exceeding 100% can be obtained by adding at least one selected from the group consisting of ethanolamine and/or ethanolamine analogs and pharmaceutically acceptable salts thereof to the medium, wherein the number of pluripotent stem cells cultured under the same conditions except that ethanolamine and/or ethanolamine analogs and pharmaceutically acceptable salts thereof are not added and in the absence of a feeder cell is the standard (100%). Whether it promotes proliferation of pluripotent stem cells while maintaining an undifferentiated state under feeder-free conditions can be evaluated by a method using a known cell proliferation system, such as the method described in Examples and the like.

[0055]

High concentration of β-mercaptoethanol is feared for its

toxicity. Therefore, the concentration of β-mercaptoethanol used in the present invention is not more than 9 μM, more preferably not more than 7 μM, further preferably not more than 5 μM, as the final concentration when in use. Furthermore, substantial or complete absence of β-mercaptoethanol is preferable. In the present invention, pluripotent stem cell can be stably proliferated while maintaining an undifferentiated state even when β-mercaptoethanol is not contained substantially.

[0056]

The definition of the "substantial absence of β-mercaptoethanol" is as mentioned above.

[0057]

The medium used in the present invention may be further added with albumin. Addition of albumin affords a preservation stabilizing effect on the medium, as well as enhances the effect of ethanolamine to promote proliferation of pluripotent stem cells while maintaining an undifferentiated state.

The "preservation stabilization" of the medium means alleviation of the time-dependent deterioration of the medium during preservation of the medium (generally, about -80°C - about 40°C) and when in use. The "deterioration of the medium" in the present invention means deterioration of the function to proliferate pluripotent stem cells while maintaining an undifferentiated state, and the level thereof can be evaluated by culturing pluripotent stem cells in said medium for a given period, and counting the cell number, as described in the below-mentioned Examples. Use of a medium immediately after preparation being the standard, a smaller cell number after culture is evaluated as further deterioration of the medium.

[0058]

The albumin to be used in the present invention is an animal-derived serum albumin. Examples of the animal include, but are not particularly limited to, rodents such as mouse, rat, hamster, guinea pig and the like, experiment animals such as

rabbit and the like, pets such as dog, cat and the like, domestic animals such as bovine, swine, goat, horse, sheep and the like, primates such as human, monkey, orangutan, chimpanzee and the like, and the like. When cells to be used for regenerative medicine are cultured, the albumin to be used in the present invention is preferably human albumin. Being "animal-derived" means that the amino acid sequence of the albumin is that of an animal.

[0059]

The albumin used in the present invention may be isolated and purified from a biological sample of an animal (e.g., blood, plasma, serum etc.), or isolated and purified after production by gene recombination technology. The preparation method of albumin is known. In addition, albumin may be a commercially available product and can be obtained from, for example, Sigma-Aldrich Co. LLC. and the like.

[0060]

The albumin used in the present invention is preferably obtained from plasma of an animal (including human) or by gene recombination technology.

[0061]

The albumin used in the present invention carries fatty acid in an amount of preferably not more than 9 mg/g, more preferably not more than 7 mg/g, further preferably not more than 2.2 mg/g.

[0062]

When a medium added with albumin is used in the present invention, the final concentration (concentration when in use) of albumin in the medium is not particularly limited as long as it affords a medium-stabilizing effect, and enhances a promoting effect of at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof on the proliferation of pluripotent stem cells while maintaining an undifferentiated state. It is generally 0.1 g/l - 20 g/l, preferably 1 g/l - 8

g/l.

[0063]

The medium to be used in the present invention may be further added with sulfated saccharides and/or a pharmaceutically acceptable salt thereof. Addition of sulfated saccharides in combination with ethanolamine to the medium affords a stabilizing effect on the medium, as well as enhances the effect of ethanolamine to promote proliferation of pluripotent stem cells while maintaining an undifferentiated state. When desired, plural kinds of sulfated saccharides and/or pharmaceutically acceptable salts thereof may be used.

[0064]

In the present invention, the "sulfated saccharide" is sulfated substance of saccharides. The "saccharide" is not particularly limited as long as it is known in the technical field, or may be novel. The saccharide may be a natural product or synthesized product. The sulfated saccharides to be added to the medium of the present invention preferably include sulfated monosaccharide, sulfated disaccharide, sulfated polysaccharide, sulfated sugar alcohol and sulfated cyclitol.

[0065]

The "monosaccharide" may be known in the technical field or novel. The number of carbons constituting carbohydrate is not limited and may be any of, for example, tetrose, pentose, hexose, heptose and the like. Specific examples of the monosaccharide include glucose, galactose, mannose, talose, idose, altrose, allose, gulose, xylose, arabinose, rhamnose, fucose, fructose, ribose, deoxyribose, glucosamine, galactosamine, glucuronic acid, galacturonic acid and the like. The sulfated monosaccharide is sulfated substance of these monosaccharides.

[0066]

The "disaccharide" is a carbohydrate wherein two molecules of the aforementioned monosaccharide are bonded by a glycosidic bond to become one molecule, and may be known in the

technical field or novel. The manner of glycosidic bond is not particularly limited, and may be any of α-1,2 bond, β-1,2 bond, α-1,3 bond, β-1,3 bond, α-1,4 bond, β-1,4 bond, α-1,5 bond, β-1,5 bond, α-1,6 bond, β-1,6 bond, α-1,α-1 bond, α-1,β-1 bond, α-1,β-2 bond and the like. Specific examples of the disaccharide include sucrose, lactose, maltose, trehalose, cellobiose, maltitol and the like. The sulfated disaccharide is sulfated substance of these disaccharides.

[0067]

The polysaccharide is a carbohydrate wherein three or more molecules of the aforementioned monosaccharide are bonded by a glycosidic bond to become one molecule, and may be known in the technical field or novel. Polysaccharide may consist of only one kind of the aforementioned saccharides, or two or more kinds thereof may be combined. Polysaccharide may be any of linear, branched and cyclic. Examples of the polysaccharide include amylose, amylopectin, glycogen, dextrin, α-cyclodextrin, β-cyclodextrin, γ-cyclodextrin, dextran, maltoheptaose, stachyose, maltotriose, pullulan, cellulose and a derivative thereof (e.g., hydroxyethylcellulose, hydroxypropylcellulose etc.), laminaran, curdlan, callose, mannan, glucomannan, galactomannan, xylan, glucuronoxylan, arabinoxylan, araban, galactan, galacturonan, chitin, chitosan, xyloglucan, pectic acid and pectin, alginic acid, arabinogalactan, glycosaminoglycan (e.g., dextran sulfate, heparan sulfate, heparin, hyaluronic acid, chondroitin 4-sulfate, chondroitin 6-sulfate, dermatan sulfate, ketaran sulfate etc.), guar gum, xanthan gum, fucoidan, inulin and the like. The sulfated polysaccharide is sulfated substance of these polysaccharides. Among the above-mentioned saccharides, those already sulfated (e.g., dextran sulfate, heparan sulfate, heparin, chondroitin 4-sulfate, chondroitin 6-sulfate, dermatan sulfate, ketaran sulfate, fucoidan etc.) include the saccharides themselves. As the sulfated polysaccharide, dextran sulfate, sulfated substance of cellulose (i.e., cellulose SO_3H), sulfated

substance of xanthan gum (i.e., xanthan gum SO_3H), fucoidan, sulfated substance of alginic acid (i.e., alginate SO_3H), sulfated substance of inulin (i.e., inulin SO_3H), sulfated substance of α-cyclodextrin (i.e., α-cyclodextrin SO_3H), sulfated substance of maltoheptaose (i.e., maltoheptaose SO_3H), sulfated substance of stachyose (i.e., stachyose SO_3H) and sulfated substance of maltotriose (i.e., maltotriose SO_3H) are preferable, and dextran sulfate is particularly preferable.

[0068]

The "sugar alcohol" is a compound produced by reducing the carbonyl group of the aforementioned monosaccharide, and may be known in the technical field or novel. Examples of the sugar alcohol include glycerol, erythritol, threitol, arabinitol, xylitol, sorbitol, mannitol, volemitol, perseitol and the like, and erythritol, xylitol and mannitol are preferable. The sulfated sugar alcohol is a sulfated substance of these sugar alcohols.

[0069]

The "cyclitol" is polyhydroxycycloalkane, and also called cyclic sugar alcohol or cyclit. The cyclitol may be known in the technical field or novel. While cyclitol is known to include many isomers, any isomer may be used. While the number of carbons constituting the ring is not particularly limited, a 6-membered ring is preferable. Examples of the cyclitol include inositol (1,2,3,4,5,6-cyclohexanehexaol), a derivative of inositol (derivative wherein hydroxy group is substituted by amino group, ketone group, carboxyl group etc.) and the like. The sulfated cyclitol is a sulfated substance of these cyclitols.

[0070]

The sulfated saccharides to be added to the medium in the present invention may be in the form of a pharmaceutically acceptable salt. Examples of such salt include salts of a sulfate group etc. present in the sulfated saccharides and a base. Specific examples thereof include alkali metal salts

such as sodium salt, potassium salt and the like; alkaline earth metal salts such as calcium salt, magnesium salt and the like; salts with inorganic base such as aluminum salt, ammonium salt and the like; salts with organic base such as trimethylamine, triethylamine, pyridine, picoline, 2,6-lutidine, ethanolamine, diethanolamine, triethanolamine, cyclohexylamine, dicyclohexylamine, N,N'-dibenzylethylenediamine and the like, which can be prepared from a free form by a conventional method. As the pharmaceutically acceptable salt of the sulfated saccharides, a sodium salt or potassium salt of a sulfate group is preferable. Examples thereof include sucrose $8SO_3K$, dextran sulfate Na (molecular weight 5,000, 25,000, 500,000 etc.), cellulose SO_3Na, xanthan gum SO_3Na, alginic acid SO_3Na, inulin SO_3Na, α-cyclodextrin SO_3Na, erythritol SO_3Na, mannitol SO_3Na, myo-inositol $6SO_3K$ and the like, and dextran sulfate Na is particularly preferable.

[0071]

When a medium added with sulfated saccharides and/or a pharmaceutically acceptable salt thereof is used in the present invention, the final concentration (concentration when in use) of sulfated saccharides and/or a pharmaceutically acceptable salt thereof in the medium is not particularly limited as long as the medium stabilizing effect is afforded and a promoting effect of at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof on the proliferation of pluripotent stem cells while maintaining an undifferentiated state is enhanced. It is generally 1 - 1000 ng/ml, preferably 10 - 250 ng/ml. When plural kinds are used, the total amount thereof is set to fall within the above-mentioned range. However, it can be increased or decreased as appropriate depending on the kind.

[0072]

The average molecular weight of sulfated saccharides or a pharmaceutically acceptable salt thereof is not particularly limited, and varies depending on the kind of the sulfated

saccharides to be employed and the kind of the salt. It is generally 50 - 1,000,000, preferably 100 - 700,000, more preferably 300 - 500,000, most preferably 500 - 100,000. When the average molecular weight exceeds 1,000,000, addition thereof at a concentration not less than a given level tends to cause toxicity or suppression of cell proliferation seemingly due to the inhibition of cell adhesion and the like. The average molecular weight can be measured by gel permeation chromatography and the like.

For example, the average molecular weight of the dextran sulfate Na is generally 1000 - 700,000, preferably 1000 - 300,000, more preferably 1000 - 100,000, most preferably 2,500 - 7,500.

[0073]

The medium to be used in the present invention may or may not contain serum. When cells to be used for regenerative medicine are cultivated, in view of the possibility of serum being a virus infection source, and concern about property difference between lots which causes inconsistent culture results and the like, culture is preferably performed under serum-free conditions.

[0074]

The medium to be used in the present invention may or may not contain a component derived from a species different from the cell to be cultured. When human cells to be used for regenerative medicine are cultivated, a component derived from an animal other than human is preferably absent from the aspects of safety after transplantation.

[0075]

As the basal medium to be used in the present invention, one known per se can be used depending on the kind of the pluripotent stem cells, and is not particularly limited as long as it does not inhibit proliferation of the pluripotent stem cells while maintaining an undifferentiated state. Examples thereof include DMEM, EMEM, IMDM (Iscove's Modified Dulbecco's

Medium), GMEM (Glasgow's MEM), RPMI-1640, α-MEM, Ham's Medium F-12, Ham's Medium F-10, Ham's Medium F12K, Medium 199, ATCC-CRCM30, DM-160, DM-201, BME, Fischer, McCoy's 5A, Leibovitz's L-15, RITC80-7, MCDB105, MCDB107, MCDB131, MCDB153, MCDB201, NCTC109, NCTC135, Waymouth's MB752/1, CMRL-1066, Williams' medium E, Brinster's BMOC-3 Medium, E8 medium (Nature Methods, 2011, 8, 424-429), a mixed medium thereof and the like. In addition, a medium altered for culture of pluripotent stem cells, a mixture of the above-mentioned basal medium and other medium, and the like may also be used.

[0076]

The medium to be used in the present invention can further contain an additive known per se. The additive is not particularly limited as long as it does not inhibit proliferation of pluripotent stem cells while maintaining an undifferentiated state. Examples thereof include growth factor (e.g., insulin etc.), iron source (e.g., transferrin etc.), polyamines (e.g., putrescine etc.), mineral (e.g., sodium selenate etc.), saccharides (e.g., glucose etc.), organic acid (e.g., pyruvic acid, lactic acid etc.), amino acid (e.g., L-glutamine etc.), reducing agent (e.g., sodium thioglycolate), vitamins (e.g., ascorbic acid, d-biotin etc.), steroid (e.g., β-estradiol, progesterone etc.), antibiotic (e.g., streptomycin, penicillin, gentamicin etc.), buffering agent (e.g., HEPES etc.) and the like. In addition, additives that have been conventionally used for culturing pluripotent stem cells can be contained as appropriate. The additive is preferably contained within a concentration range known per se.

[0077]

In the culture method of the present invention, a feeder cell may or may not be used. When cells to be used for regenerative medicine are cultivated, from the aspect of safety after transplantation, a feeder cell is preferably absent (feeder-free). Although not bound by any theory, in the culture method of the present invention, proliferation of

pluripotent stem cells while maintaining an undifferentiated state can be promoted even in the absence of a feeder cell, since ethanolamine generally secreted from a feeder cell is added to the medium.

[0078]

When a feeder cell is not used, culture is preferably performed by using an extracellular matrix or an active fragment thereof or an artificial product mimicking the functions thereof.

The extracellular matrix is not particularly limited as long as it is generally used for cell culture with the aim to improve adhesion between the surface of a culture vessel and the cell. For example, known ones such as laminin (laminin 511, laminin 332 etc.), fibronectin, vitronectin, collagen, elastin, adhesamine and the like can be used. The active fragment of an extracellular matrix only needs to be a fragment thereof having a cell adhesion activity equivalent to that of the extracellular matrix, and known ones can be used. For example, E8 fragment of laminin 511, E8 fragment of laminin 332 and the like disclosed in JP-A-2011-78370 can be mentioned. The extracellular matrix and an active fragment thereof may be commercially available products and available from, for example, (Life Technologies, BD Falcon, BioLamina) and the like. Two or more kinds of these extracellular matrices and active fragments thereof may be used in combination. Also, a matrigel (trade name) which is a mixture of complicated basal lamina components containing protein and polysaccharides, that are extracted and purified from EHS sarcoma of mouse overproducing the basal lamina, may also be used. The extracellular matrix and an active fragment thereof may be suspended in a suitable solution, and applied to a container suitable for cultivating cells.

An artificial product mimicking the function of extracellular matrix is not particularly limited as long as it is generally used for culturing cells and, for example, known ones such as Synthemax (registered trade mark)and Ultra-Web

(registered trade mark) of Corning Incorporated, Hy-STEM series , polylysine and polyornithine of Sigma Aldrich Co., Ltd. and the like can be used.

The extracellular matrix or an active fragment thereof or an artificial product mimicking the functions thereof to be used in the present invention are preferably matrigel or laminin 511 or an active fragment of laminin 511, more preferably an active fragment of laminin 511 (i.e., E8 fragment of laminin 511).

[0079]

In the culture method of the present invention, the cell seeding method is not particularly limited, and may be colony seeding or single cell-seeding. To produce pluripotent stem cells for regenerative medicine at an industrial level, the work needs to be performed by plural workers under the conditions where procedures and schedule are rigorously managed. Therefore, single cell-seeding permitting rigorous adjustment of the seeding cell number is preferable.

For single cell-seeding, colonies of pluripotent stem cells are dissociated to single cells, and seeded in the medium. Single cell seeding can be performed by a method known per se. For example, cell-cell adhesion and cell-matrix adhesion are weakened with a cell detaching solution (trypsin solution etc.), and the cells are detached from the matrix with a scraper (IWAKI, 9000-220 etc.) and the like (in this state, the cells forming cell clusters are suspended in a solution, not complete single cells). The cells are thereafter dissociated by pipetting into single cells, and seeded in the medium. When seeding, ROCK inhibitor such as Y-27632 (Nacalai Tesque: 08945-84) and the like is preferably added to the medium to ensure survival of the pluripotent stem cells. Since ROCK inhibitor is not necessary for the proliferation of pluripotent stem cells from the following day of the seeding, it is preferably excluded from the medium.

[0080]

Other culture conditions can be appropriately determined. For example, while the culture temperature is not particularly limited, it can be about 30 - 40°C, preferably about 37°C. The CO_2 concentration can be about 1 - 10%, preferably about 2 - 5%. The oxygen partial pressure can be 1 - 10%.

In cell culture, medium change is sometimes necessary during the culture due to the deterioration of medium components, accumulation of waste products discharged from the cell and the like. In the culture method of the present invention, it is also possible to omit the medium change as shown in the Examples. For example, pluripotent stem cells can be cultivated for 4 or more consecutive days (4 days, 5 days, 6 days etc.) without medium change.

[0081]

In one embodiment, the present invention provides a culture method for the proliferation of a pluripotent stem cell while maintaining an undifferentiated state, comprising a step of cultivating a pluripotent stem cell in a medium added with at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof, and added with sulfated saccharides and/or a pharmaceutically acceptable salt thereof. The definitions of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof, and sulfated saccharides and a pharmaceutically acceptable salt thereof in the culture method are the same as those mentioned above, and use concentrations thereof are also as mentioned above. In addition, other conditions relating to the culture method are also as mentioned above.

[0082]

(Preservation stabilizing method for medium for proliferation of pluripotent stem cells)

The present invention provides a preservation stabilizing method for a medium for the proliferation of a pluripotent stem cell, comprising adding at least one selected from the group

consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof, and adding sulfated saccharides and/or a pharmaceutically acceptable salt thereof (hereinafter to be also referred to as the stabilizing method of the present invention).

[0083]

The "preservation stabilization" of the medium in the present invention means alleviation of the time-dependent deterioration of the medium during preservation of the medium (generally, about -80°C - 40°C). The "deterioration of the medium" here means deterioration of the function to proliferate pluripotent stem cells while maintaining an undifferentiated state, and the level thereof can be evaluated by culturing pluripotent stem cells in said medium for a given period, and counting the cell number, as described in the below-mentioned Examples. Use of a medium immediately after preparation being the standard, a smaller cell number after culture is evaluated as further deterioration of the medium.

[0084]

The medium for the proliferation of pluripotent stem cells in the present invention is the same as the medium used in the above-mentioned culture method of the present invention, and preferably a medium for the proliferation of pluripotent stem cells while maintaining an undifferentiated state.

[0085]

Ethanolamine and ethanolamine analogs and pharmaceutically acceptable salts thereof to be added to the medium for preservation stability of the medium are the same as those used for the above-mentioned culture method of the present invention. The concentration of addition to the medium is the same as those in the above-mentioned culture method of the present invention. By adding ethanolamine and ethanolamine analogs and pharmaceutically acceptable salts thereof to the medium, the medium after addition can be stabilized.

[0086]

The stabilizing method of the present invention further comprises a step of adding sulfated saccharides and/or a pharmaceutically acceptable salt thereof. The sulfated saccharides and/or a pharmaceutically acceptable salt thereof are the same as those used for the above-mentioned culture method of the present invention. The concentration of addition to the medium is the same as that in the above-mentioned culture method of the present invention. By adding at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof, and sulfated saccharides and/or a pharmaceutically acceptable salt thereof in combination to the medium, the stability of the medium can be further enhanced.

[0087]

According to the present invention, for example, when the medium is preserved at room temperature (generally about 15 - 25°C) for up to about 8 days, after preparation of the medium, deterioration of the medium can be alleviated. Hence, the medium can be preserved for a longer term than before, and the present invention is useful for the production of pluripotent stem cells and the like.

[0088]

(Medium additive for proliferation of pluripotent stem cells while maintaining an undifferentiated state)

The component to be added to the basal medium in the above-mentioned culture method of the present invention can be a medium additive. That is, the present invention provides a medium additive for the proliferation of a pluripotent stem cell while maintaining an undifferentiated state, comprising at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof, and substantially free of β-mercaptoethanol or containing β-mercaptoethanol at a concentration of not more than 9 μM when in use (hereinafter to be also referred to as the medium additive of the present invention).

[0089]

Ethanolamine and ethanolamine analogs and pharmaceutically acceptable salts thereof are the same as those used for the above-mentioned culture method of the present invention.

[0090]

The amount of the medium additive of the present invention to be added to a medium can be determined to fall within any range as long as it can promote proliferation of pluripotent stem cells while maintaining an undifferentiated state when cultured under feeder-free conditions using the medium after preparation. It can be added such that the final concentration of ethanolamine and ethanolamine analogs and pharmaceutically acceptable salts thereof in the medium is generally 1 μM - 1000 μM, preferably 5 μM - 200 μM, or 11 μM - 200 μM.

[0091]

High concentration β-mercaptoethanol is feared for its toxicity. Therefore, the concentration of β-mercaptoethanol used in the present invention is not more than 9 μM, more preferably not more than 7 μM, further preferably not more than 5 μM, as the final concentration when in use. Furthermore, substantial or complete absence of β-mercaptoethanol is preferable. In the present invention, pluripotent stem cell can be stably proliferated while maintaining an undifferentiated state even when β-mercaptoethanol is not contained substantially.

[0092]

The definition of the "substantially free of β-mercaptoethanol" is as mentioned above.

[0093]

The medium additive of the present invention may further contain albumin, sulfated saccharides and/or a pharmaceutically acceptable salt thereof. These are the same as those used for the above-mentioned culture method of the present invention.

[0094]

Where necessary, the medium additive of the present invention may contain other components other than the above-mentioned components. Examples of other components include, but are not particularly limited to, additives generally used for the preparation of a medium, for example, additives contained in the medium used in the above-mentioned production method of the present invention.

[0095]

While the medium additive of the present invention may or may not contain a component derived from a species different from the cell to be cultured. When human cells to be used for regenerative medicine are cultivated, it is preferable that a component derived from an animal other than human not be contained.

[0096]

Using the medium additive of the present invention, a medium capable of stable proliferation for a long term while maintaining an undifferentiated state even in serum-free, feeder-free and single cell-seeding culture can be prepared.

[0097]

The present invention is explained in more detail in the following by referring to Examples, which are not to be construed as limitative.

Examples

[0098]

In the following Examples, proliferation effects of various test compounds on human pluripotent stem cells were evaluated. As human pluripotent stem cells, induced pluripotent stem cells (iPS cells, 201B7 strain) purchased from iPS Academia Japan, Inc. were used, unless particularly specified. Cell culture was performed using a culture vessel (Becton, Dickinson and Company, Falcon culture petri dish or Falcon culture plate) coated with a basal lamina matrix under the conditions of 5% CO_2/37°C.

Various test compounds were added to a medium having the "E8" composition (disclosed in Nature Methods, 2011, 8, 424-429) considered at present to be the minimum composition for cultivating human pluripotent stem cells at given concentrations and used for culture, and the effects thereof were studied. The medium was prepared by using Essential 8 (Life Technologies: A14666SA) considered to have the "E8" composition or a medium formulated to have an equivalent composition.

[0099]

Reference Example 1 medium stabilizing effect of albumin

Human serum-derived albumin (Sigma-Aldrich Co. LLC: A1887) was added at a final concentration of 2.6 g/l, and the effect of albumin was examined by studying the cell numbers after culture when used immediately after preparation for culture or when used after preservation at 4°C for 3 weeks after the preparation. The culture period was 1 week. 13,000 viable cells per well were used for single cell seeding. As a basal lamina matrix, a fragment containing an active domain of laminin 511, which was purchased from Osaka University, was applied at 5 μg/well. Y-27632 (final concentration 10 μM, Nacalai Tesque: 08945-84) was added to the medium used for seeding. The cells were cultured in a medium free of Y-27632 from the following day. Viable cell number was measured by Trypan Blue (Life Technologies: 15250-061) staining using a hemocytometer.

The average results of 3 series of experiments for each medium are shown in Fig. 1. When used immediately after preparation, equivalent cell proliferation was observed with or without albumin addition. When a medium after 3 weeks from preparation was used, cell proliferation was scarcely found in an albumin-free medium, whereas obvious cell proliferation was observed in an albumin addition medium. The above results reveal that E8 minuimun composition medium shows remarkable degradation of the property when kept under general use

(preservation) conditions for 3 weeks, but addition of albumin can improve such deteriorative phenomenon. To conclude, albumin was found to contribute to the stabilization of a medium stored at 4°C.

[0100]

Example 1 Proliferation promoting effect of ethanolamine and combination effect with albumin

Ethanolamine (Sigma-Aldrich Co. LLC: E0135) was added at a final concentration of 6, 30, 150, 750 or 3,750 μM, and the medium was used for culture immediately after preparation, and the effect of ethanolamine was examined by studying the cell numbers after culture. The culture period was 1 week. To study the effect of combination with albumin, human serum-derived albumin (Sigma-Aldrich Co. LLC: A1887) was further added at a final concentration of 2.6 g/l to the above-mentioned ethanolamine addition medium, and a similar examination was performed. 13,000 viable cells per 1 well were used for single cell-seeding. As a basal lamina matrix, a fragment containing the active domain of laminin 511, which was purchased from Osaka University, was applied at 5 μg/well. Y-27632 was added (final concentration 10 μM, Nacalai Tesque: 08945-84) to the medium used for seeding. The cells were cultured in a medium free of Y-27632 from the following day. The number of viable cells was measured by Trypan Blue (Life Technologies: 15250-061) staining using a hemocytometer.

The experiment was performed in 3 series for each medium, and the average results are shown in Fig. 2. The values are shown as relative values to the ethanolamine non-addition group (0 μM). Ethanolamine was found to show a proliferation-promoting effect in a relatively wide concentration range. It was found that a proliferation suppressive effect conversely appears in a high concentration range. It was found that combination with albumin further enhances the proliferation promoting effect of ethanolamine, resists a proliferation suppressive effect in a high concentration range, and shows a

proliferation promoting effect.

[0101]

Example 2 Effect of dextran sulfate

Human serum derived albumin (Sigma-Aldrich Co. LLC: A1887) was added to a medium (final concentration 2.6 g/l), ethanolamine (final concentration 30 μM) was added singly thereto, dextran sulfate sodium (Wako Pure Chemical Industries, Ltd., final concentration 50 ng/ml) was further added to the medium, and a further proliferation promoting effect of a combination of ethanolamine and dextran sulfate sodium was verified. Culture was performed for about 3 weeks, which included two passages, and the cumulative viable cell increase rate for the total culture period was calculated. 13,000 viable cells per 1 well were used for single cell-seeding. As a basal lamina matrix, a fragment containing the active domain of laminin 511, which was purchased from Osaka University, was applied at 5 μg/well. Y-27632 was added (final concentration 10 μM, Nacalai Tesque: 08945-84) to the medium used for seeding. The cells were cultured in a medium free of Y-27632 from the following day. When passaged, cells were detached by TrypLE™ Select (Life Technologies: 12563-011), 13,000 viable cells were seeded again in Y-27632 added medium, and cultured in a medium free of Y-27632 from the following day. The number of viable cells was measured by Trypan Blue (Life Technologies: 15250-061) staining using a hemocytometer.

[0102]

The experiment was performed in 3 series for each medium, and the average results are shown in Fig. 3. It was found that the cell increase rate was higher when dextran sulfate sodium was combined than the single addition of ethanolamine.

Both ethanolamine and dextran sulfate sodium, which showed the highest increase rate, were added to a medium, and culture was performed for about one month. Alkaline phosphatase staining was performed to confirm maintenance of undifferentiated state. The results of staining with an

alkaline phosphatase staining kit (Sigma-Aldrich Co. LLC: 86-R) are shown in Fig. 4. iPS cell colonies in the whole well were stained, which confirmed that long-term culture in an E8 minimum composition medium added with albumin, ethanolamine, dextran sulfate sodium resulted in the proliferation of iPS cells while maintaining an undifferentiated state.

[0103]

Example 3 Proliferation promoting effect of ethanolamine - results of culture using matrigel

Ethanolamine (Sigma-Aldrich Co. LLC: E0135) was added at a final concentration of 6, 30, 150, 750 or 3,750 μM, and the medium was used for culture immediately after preparation, and the effect of ethanolamine was examined by studying the cell numbers after culture. The culture period was 8 days. 100,000 cells per 1 well were used for single cell-seeding. As a basal lamina matrix, matrigel (Japan Becton Dickinson) was applied. Y-27632 was added (final concentration 10 μM, Nacalai Tesque: 08945-84) to the medium used for seeding. The cells were cultured in a medium free of Y-27632 from the following day.

The experiment was performed in 3 series for each medium, and the average results are shown in Fig. 5. In the same manner as in Example 1, the results showing that ethanolamine affords a proliferation promoting effect in a wide concentration range were obtained. Therefore, it was found that the proliferation promoting effect of ethanolamine is not limited to culture using laminin 511.

[0104]

Example 4 Effect of combination of ethanolamine and dextran sulfate sodium - results of culture using matrigel

(1) Effect in medium containing albumin

Essential 8 medium added with human serum derived albumin (final concentration 2.6 g/l) and ethanolamine (final concentration 30 μM) (control), and the medium added with dextran sulfate sodium at a given concentration were prepared, and the effect of combination of ethanolamine and dextran

sulfate sodium was verified. The culture period was 6-12 days. For single cell-seeding, 40,000 cells per 1 well were seeded on a 12-well culture plate coated with matrigel. Y-27632 was added (final concentration 10 μM, Nacalai Tesque: 08945-84) to the medium used for seeding. The cells were cultured in a medium free of Y-27632 from the following day. For seeding as colony, the cells diluted 2.5- to 3.5-fold of the original culture per well were plated on a 6 well culture plate coated with matrigel. In this case, Y-27632 was not added to the medium to be used for seeding. The medium change was performed every 2 - 3 days.

The evaluation criteria of cell proliferation were as follows.

⊙: cell number is not less than 120% of that of control

O: cell number is not less than 100% and less than 120% of that of control

-: cell number is not less than 50% and less than 100% of that of control

x: cell number is not more than 50% of that of control

The experiment was performed in 3 series for each medium, and the results are shown in Table 1. Higher cell proliferation was observed when dextran sulfate sodium was added. Therefore, it was found that the cell proliferation promoting effect afforded by a combination of ethanolamine and dextran sulfate sodium is not limited to culture using laminin 511. It was also found that the effect was exhibited even when seeded as colony, and the effect is not limited to single cell-seeding.

[0105]

Table 1

	final concentration (ng/ml) of dextran sulfate sodium			
	1	10	50	100
single cell-seeding	○	○	◎	not evaluated
colony seeding	not evaluated	◎	not evaluated	◎

[0106]

(2) Effect of omission of medium change

In the evaluation of the above-mentioned (1), culture was performed in a medium added with dextran sulfate sodium (final concentration 10 ng/ml) without medium change, and whether medium change can be omitted by adding ethanolamine and dextran sulfate sodium in combination to the medium was verified. The culture period was 6 days.

The experiment was performed in 3 series for each medium, and the results are shown in Fig. 6. When medium change was not performed, high cell proliferation was found by the addition of dextran sulfate sodium. The effect thereof was close to that afforded by medium change with a medium free of dextran sulfate sodium. Therefore, it was found that a combination of ethanolamine and dextran sulfate sodium provides an effect of omitting medium change.

[0107]

(3) Stabilizing effect in medium without albumin

Essential 8 medium added with ethanolamine (final concentration 30 μM) alone, the medium added with ethanolamine (final concentration 30 μM) and dextran sulfate sodium (Wako Pure Chemical Industries, Ltd., final concentration 50 ng/ml), and the medium added with human serum derived albumin (final concentration 2.6 g/l) alone were prepared, and a medium stabilizing effect of a combination of ethanolamine and dextran sulfate sodium was verified. Each medium was left standing at room temperature for 8 days after preparation and used for

culture. The culture period was 8 days. 100,000 cells per 1 well were used for single cell-seeding on a 6-well culture plate coated with matrigel. Y-27632 was added (final concentration 10 μM, Nacalai Tesque: 08945-84) to the medium used for seeding. The cells were cultured in a medium free of Y-27632 from the following day.

The experiment was performed in 3 series for each medium, and the average results are shown in Fig. 7. The effect afforded by adding both ethanolamine and dextran sulfate sodium was equal to that afforded by adding albumin. Therefore, it was found that a combination of ethanolamine and dextran sulfate sodium provides a medium stabilizing effect at room temperature, and has a possibility of replacing or reducing albumin widely used for medium.

[0108]

Example 5 Effect of ethanolamine analogs

(1) Effect of O-phosphoryl ethanolamine (aka phosphoethanolamine)

To a medium added with human serum-derived albumin (Sigma-Aldrich Co. LLC.: A1887) at a final concentration of 2.6 g/l was added O-phosphoryl ethanolamine (Sigma-Aldrich Co. LLC.: P0503-25G) at a final concentration of 6, 30, 150 or 750 μM, and used for the culture from the next day of preparation. After the culture, the cell number was counted to examine the effect of O-phosphoryl ethanolamine. The culture period was set to 1 week. 13,000 viable cells per well were used for single cell-seeding. As a basal lamina matrix, a fragment containing the active domain of laminin 511, which was purchased from Osaka University, was applied at 4.8 μg/well. Y-27632 was added (final concentration 10 μM, Nacalai Tesque: 08945-84) to the medium used for seeding. The cells were cultured in a medium free of Y-27632 from the following day. The number of viable cells was measured by Cell Viability autoanalyzer ViCELL™ XR (BECKMAN COULTER).

The experiment was performed in 3 series for each medium,

and the average results are shown in Fig. 8. The values are shown as relative values to the O-phosphoryl ethanolamine non-addition group (0 μM). O-phosphoryl ethanolamine was found to show a proliferation-promoting effect in a relatively wide concentration range.

[0109]

(2) Effect of 2-(methylamino)ethanol

To a medium added with human serum-derived albumin (Sigma-Aldrich Co. LLC.: A1887) at a final concentration of 2.6 g/l was added 2-(methylamino)ethanol (Sigma-Aldrich Co. LLC.: 471445-25ML) at a final concentration of 6, 30, 150 or 750 μM, and used for the culture from the next day of preparation. After the culture, the cell number was counted to examine the effect of 2-(methylamino)ethanol. The culture period was set to 1 week. 13,000 viable cells per well were used for single cell-seeding. As a basal lamina matrix, a fragment containing the active domain of laminin 511, which was purchased from Osaka University, was applied at 4.8 μg/well. Y-27632 was added (final concentration 10 μM, Nacalai Tesque: 08945-84) to the medium used for seeding. The cells were cultured in a medium free of Y-27632 from the following day. The number of viable cells was measured by Cell Viability autoanalyzer ViCELL™ XR (BECKMAN COULTER).

The experiment was performed in 3 series for each medium, and the average results are shown in Fig. 9. The values are shown as relative values to the 2-(methylamino)ethanol non-addition group (0 μM). 2-(Methylamino)ethanol was found to show a proliferation-promoting effect in a relatively wide concentration range.

[0110]

(3) Effect of 2-dimethylaminoethanol

To a medium added with human serum-derived albumin (Sigma-Aldrich Co. LLC.: A1887) at a final concentration of 2.6 g/l was added 2-dimethylaminoethanol (Sigma-Aldrich Co. LLC.: 471453-100 ml) at a final concentration of 6, 30, 150 or 750 μM,

and used for the culture from the next day of preparation. After the culture, the cell number was counted to examine the effect of 2-dimethylaminoethanol. The culture period was set to 1 week. 13,000 viable cells per well were used for single cell-seeding. As a basal lamina matrix, a fragment containing the active domain of laminin 511, which was purchased from Osaka University, was applied at 4.8 μg/well. Y-27632 was added (final concentration 10 μM, Nacalai Tesque: 08945-84) to the medium used for seeding. The cells were cultured in a medium free of Y-27632 from the following day. The number of viable cells was measured by Cell Viability autoanalyzer ViCELL™ XR (BECKMAN COULTER).

The experiment was performed in 3 series for each medium, and the average results are shown in Fig. 10. The values are shown as relative values to the 2-dimethylaminoethanol non-addition group (0 μM). It was found that 2-dimethylaminoethanol shows a proliferation promoting effect in a relatively wide concentration range, but a proliferation suppressive effect appears in a high concentration range.

[0111]

(4) Effect of ethanolamine hydrochloride

To a medium added with human serum-derived albumin (Sigma-Aldrich Co. LLC.: A1887) at a final concentration of 2.6 g/l was added ethanolamine hydrochloride (Tokyo Chemical Industry Co., Ltd.: A0298) at a final concentration of 6, 30, 150 or 750 μM, and used for the culture from the next day of preparation. After the culture, the cell number was counted to examine the effect of ethanolamine hydrochloride. The culture period was set to 1 week. 13,000 viable cells per well were used for single cell-seeding. As a basal lamina matrix, a fragment containing the active domain of laminin 511, which was purchased from Osaka University, was applied at 4.8 μg/well. Y-27632 was added (final concentration 10 μM, Nacalai Tesque: 08945-84) to the medium used for seeding. The cells were cultured in a medium free of Y-27632 from the following day.

The number of viable cells was measured by Cell Viability autoanalyzer ViCELL™ XR (BECKMAN COULTER).

The experiment was performed in 3 series for each medium, and the average results are shown in Fig. 11. The values are shown as relative values to the ethanolamine hydrochloride non-addition group (0 μM). Ethanolamine hydrochloride was found to show a proliferation-promoting effect in a relatively wide concentration range.

[0112]

Example 6 Influence of oleic acid

To a solution (25%, 40 ml) of human serum albumin (Nova Biologics) in saline was added phosphate buffer (pH 7.2, 40 ml). Furthermore, a suspension of activated carbon (5 g, manufactured by Wako Pure Chemical Industries, Ltd.) heated at 200°C for 30 min in advance in phosphate buffer (20 ml) was added to 100 ml. After stirring at 4°C for 3 hr, the mixture was centrifuged at 4°C, 11,900 rpm for 20 min. Sedimented activated carbon was removed by decantation, and the reaction mixture was filtered with a 0.22 μm syringe filter, whereby human serum albumin with fatty acid removal treatment was obtained. Then, oleic acid was added to the obtained human serum albumin with fatty acid removal treatment to prepare human serum-derived albumin having a fatty acid carried amount of 2.2, 6.5, 21.7 mg/g, and each albumin was added to the medium to a final concentration of 2.6 g/l (final concentration of oleic acid in the medium 20, 60, 200 μM, respectively). Ethanolamine (final concentration 30 μM) was added to each medium. Using each medium, the cells were cultured for one week, and the cell number after culture was confirmed to study the influence of oleic acid. 13,000 viable cells per well were used for single cell-seeding. As a basal lamina matrix, a fragment containing the active domain of laminin 511, which was purchased from Osaka University, was applied at 4.8 μg/well. Y-27632 was added (final concentration 10 μM, Nacalai Tesque: 08945-84) to the medium used for seeding. The cells were

cultured in a medium free of Y-27632 from the following day. The number of viable cells was measured by Cell Viability autoanalyzer ViCELL™ XR (BECKMAN COULTER).

The results of culture using each medium are shown in Fig. 12. It was found that cell proliferation was suppressed along with an increasing amount of oleic acid carried by albumin.

Industrial Applicability

[0113]

According to the present invention, pluripotent stem cells can be stably and efficiently proliferated, and can be stably proliferated for a long term while maintaining an undifferentiated state even in serum-free, feeder-free and single cell-seeding culture, and therefore, it is useful in the field of regenerative medicine and the like.

[0114]

This application is based on a patent application No. 2013-016592 filed in Japan (filing date: January 31, 2013), the contents of which are incorporated in full herein.

Claims

1. A culture method for a proliferation of a pluripotent stem cell while maintaining an undifferentiated state, comprising a step of cultivating the pluripotent stem cell in a medium added with at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof, and substantially free of β-mercaptoethanol or containing β-mercaptoethanol at a concentration of not more than 9 μM.

2. The method according to claim 1, wherein the concentration of at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof in the medium is 1 μM - 1000 μM.

3. The method according to claim 1, wherein the concentration of at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof in the medium is 5 μM - 200 μM.

4. The method according to any one of claims 1 to 3, wherein the ethanolamine analog is a compound represented by the following formula

$X-CH_2-CH_2-O-Y$

wherein

X is $R^1-N(R^2)-$ [R^1 and R^2 are the same or different and each is a hydrogen atom or an amino-protecting group] or $R^3-CH=N-$ [R^3-CH is H-CH or a Shiff base type amino-protecting group]; and

Y is $-P(=O)(OH)-O-R^4$ [R^4 is $-CH_2-CH(O-R^5)-CH_2-O-R^6$ (R^5 and R^6 are the same or different and each is an acyl group having 2 - 30 carbon atoms or a hydrogen atom) or a hydrogen atom], a hydrogen atom or a hydroxy-protecting group.

5. The method according to claim 4, wherein R^1 and R^2 are the

same or different and each is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms, and R^3 is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms.

6. The method according to any one of claims 1 to 3, wherein the ethanolamine analog is one or plural selected from the group consisting of phosphoethanolamine, monomethylethanolamine, dimethylethanolamine, N-acylphosphatidylethanolamine, phosphatidylethanolamine, and lysophosphatidylethanolamine.

7. The method according to any one of claims 1 to 6, wherein the medium is further added with albumin.

8. The method according to claim 7, wherein the concentration of the albumin in the medium is 0.1 g/l - 20 g/l.

9. The method according to claim 7, wherein the concentration of the albumin in the medium is 1 g/l - 8 g/l.

10. The method according to any one of claims 7 to 9, wherein the albumin is obtained from plasma of an animal (including human) or by gene recombination technology.

11. The method according to any one of claims 7 to 10, wherein

the amount of fatty acid carried by the albumin in the medium is not more than 9 mg/g.

12. The method according to any one of claims 7 to 10, wherein the amount of fatty acid carried by the albumin in the medium is not more than 2.2 mg/g.

13. The method according to any one of claims 1 to 12, wherein the medium is further added with sulfated saccharide and/or a pharmaceutically acceptable salt thereof.

14. The method according to claim 13, wherein the concentration of the sulfated saccharide and/or a pharmaceutically acceptable salt thereof in the medium is 1 - 1000 ng/ml.

15. The method according to claim 13 or 14, wherein the sulfated saccharide or a pharmaceutically acceptable salt thereof is dextran sulfate Na having an average molecular weight of 2,500 - 7,500.

16. A culture method for a proliferation of a pluripotent stem cell while maintaining an undifferentiated state, comprising a step of cultivating the pluripotent stem cell in a medium added with at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof, and added with sulfated saccharide and/or a pharmaceutically acceptable salt thereof.

17. The method according to claim 16, wherein the concentration of the sulfated saccharide and/or a pharmaceutically acceptable salt thereof in the medium is 1 - 1000 ng/ml.

18. The method according to claim 16 or 17, wherein the sulfated saccharide or a pharmaceutically acceptable salt thereof is dextran sulfate Na having an average molecular

weight of 2,500 - 7,500.

19. The method according to any one of claims 16 to 18, wherein the medium is further added with albumin.

20. The method according to claim 19, wherein the concentration of the albumin in the medium is 0.1 g/l - 20 g/l.

21. The method according to claim 19, wherein the concentration of the albumin in the medium is 1 g/l - 8 g/l.

22. The method according to any one of claims 19 to 21, wherein the albumin is obtained from plasma of an animal (including human) or by gene recombination technology.

23. The method according to any one of claims 1 to 22, wherein the culture is performed in the absence of a feeder cell.

24. The method according to claim 23, wherein the culture is performed by using an extracellular matrix or an active fragment thereof, or an artificial product mimicking the function thereof.

25. The method according to claim 23, wherein the culture is performed by using laminin 511 or an active fragment thereof or matrigel.

26. The method according to any one of claims 1 to 25, wherein the culture is performed by single cell-seeding.

27. The method according to any one of claims 1 to 26, wherein the culture is performed under serum-free conditions.

28. The method according to any one of claims 1 to 27, wherein the medium is substantially free of a component derived from an

animal other than human.

29. The method according to any one of claims 1 to 28, wherein the pluripotent stem cell is an embryonic stem cell (ES cell) or induced pluripotent stem cell (iPS cell).

30. The method according to any one of claims 1 to 29, wherein the pluripotent stem cell is derived from primates.

31. The method according to any one of claims 1 to 30, wherein the pluripotent stem cell is human iPS cell.

32. A preservation stabilizing method for a medium for a proliferation of a pluripotent stem cell, comprising adding at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof, and adding sulfated saccharide and/or a pharmaceutically acceptable salt thereof.

33. The method according to claim 32, wherein the final concentration of the sulfated saccharide and/or a pharmaceutically acceptable salt thereof in use is 1 - 1000 ng/ml.

34. The method according to claim 32 or 33, further comprising adding albumin.

35. The method according to claim 34, wherein the final concentration of the albumin in use is 0.1 g/l - 20 g/l.

36. The method according to claim 34, wherein the final concentration of the albumin in use is 1 g/l - 8 g/l.

37. The method according to any one of claims 34 to 36, wherein the albumin is obtained from plasma of an animal (including

human) or by gene recombination technology.

38. The method according to any one of claims 34 to 37, wherein the amount of fatty acid carried by the albumin is not more than 9 mg/g.

39. The method according to any one of claims 34 to 37, wherein the amount of fatty acid carried by the albumin is not more than 2.2 mg/g.

40. The method according to any one of claims 32 to 39, wherein the medium is substantially free of β-mercaptoethanol or contains β-mercaptoethanol at a final concentration of not more than 9 μM.

41. The method according to any one of claims 32 to 40, wherein the medium is for a proliferation of a pluripotent stem cell while maintaining an undifferentiated state.

42. A medium additive for a proliferation of a pluripotent stem cell while maintaining an undifferentiated state, comprising at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof, and substantially free of β-mercaptoethanol or containing β-mercaptoethanol at a concentration of not more than 9 μM when in use.

43. The medium additive according to claim 42, wherein the ethanolamine analog is a compound represented by the following formula

$X-CH_2-CH_2-O-Y$

wherein

X is $R^1-N(R^2)-$ [R^1 and R^2 are the same or different and each is a hydrogen atom or an amino-protecting group] or $R^3-CH=N-$ [R^3-CH is H-CH or a Shiff base type amino-protecting group]; and

Y is $-P(=O)(OH)-O-R^4$ [R^4 is $-CH_2-CH(O-R^5)-CH_2-O-R^6$ (R^5 and R^6 are the same or different and each is an acyl group having 2 - 30 carbon atoms or a hydrogen atom) or a hydrogen atom], a hydrogen atom or a hydroxy-protecting group.

44. The medium additive according to claim 43, wherein R^1 and R^2 are the same or different and each is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms, and
R^3 is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms.

45. The medium additive according to any one of claims 42 to 44, wherein the ethanolamine analog is one or plural selected from the group consisting of phosphoethanolamine, monomethylethanolamine, dimethylethanolamine, N-acylphosphatidylethanolamine, phosphatidylethanolamine, and lysophosphatidylethanolamine.

46. The medium additive according to any one of claims 42 to 45, further comprising albumin.

47. The medium additive according to claim 46, wherein the final concentration of the albumin in use is 0.1 g/l - 20 g/l.

48. The medium additive according to claim 46, wherein the final concentration of the albumin in use is 1 g/l - 8 g/l.

49. The medium additive according to any one of claims 46 to 48, wherein the amount of fatty acid carried by the albumin is not more than 9 mg/g.

50. The medium additive according to any one of claims 46 to 48, wherein the amount of fatty acid carried by the albumin is not more than 2.2 mg/g.

51. The medium additive according to any one of claims 46 to 50, wherein the albumin is obtained from plasma of an animal (including human) or by gene recombination technology.

52. The medium additive according to any one of claims 42 to 51, further comprising sulfated saccharide and/or a pharmaceutically acceptable salt thereof.

53. The medium additive according to claim 52, wherein the final concentration of the sulfated saccharide and/or a pharmaceutically acceptable salt thereof in use is 1 - 1000 ng/ml.

54. The medium additive according to claim 52 or 53, wherein the sulfated saccharide is at least one selected from the group consisting of sulfated monosaccharide, sulfated disaccharide, sulfated polysaccharide, sulfated sugar alcohol and sulfated cyclitol.

55. The medium additive according to any one of claims 52 to 54, wherein the aforementioned sulfated saccharide or a pharmaceutically acceptable salt thereof is at least one selected from the group consisting of dextran sulfate Na, cellulose SO_3Na, xanthan gum SO_3Na, fucoidan, alginate SO_3Na,

inulin SO_3Na, maltoheptaose SO_3Na, stachyose SO_3Na, maltotriose SO_3Na, multitol SO_3Na, sucrose8SO_3K, glucose SO_3Na, myo-6 inositol SO_3K, α-cyclodextrin SO_3Na, mannitol SO_3Na, xylitol SO_3Na and erythritol SO_3Na.

56. The medium additive according to claim 55, wherein the aforementioned sulfated saccharide or a pharmaceutically acceptable salt thereof is at least one selected from the group consisting of dextran sulfate Na, fucoidan, maltoheptaose SO_3Na, maltotriose SO_3Na, multitol SO_3Na and sucrose8SO_3K.

57. The medium additive according to any one of claims 52 to 56, wherein the aforementioned sulfated saccharide or a pharmaceutically acceptable salt thereof is dextran sulfate Na having an average molecular weight of 2,500 - 7,500.

58. The medium additive according to any one of claims 42 to 57, wherein the proliferation of a pluripotent stem cell while maintaining an undifferentiated state is performed under the conditions without using a feeder cell.

59. The medium additive according to claim 58, wherein the conditions without using a feeder cell do not include use of a feeder cell but include use of an extracellular matrix or an active fragment thereof, or an artificial product mimicking the function thereof.

60. The medium additive according to claim 58, wherein the conditions without using a feeder cell do not include use of a feeder cell but include use of laminin 511 or an active fragment thereof or matrigel.

61. The medium additive according to any one of claims 42 to 60, wherein the proliferation of a pluripotent stem cell while maintaining an undifferentiated state is performed by single

cell-seeding.

62. The medium additive according to any one of claims 42 to 61, wherein the proliferation of a pluripotent stem cell while maintaining an undifferentiated state is performed under serum-free conditions.

63. The medium additive according to any one of claims 42 to 62, comprising substantially free of a component derived from an animal other than human.

64. The medium additive according to any one of claims 42 to 63, wherein the pluripotent stem cell is an embryonic stem cell (ES cell) or induced pluripotent stem cell (iPS cell).

65. The medium additive according to any one of claims 42 to 64, wherein the pluripotent stem cell is derived from primates.

66. The medium additive according to any one of claims 42 to 65, wherein the pluripotent stem cell is human iPS cell.

67. A medium for proliferation of a pluripotent stem cell while maintaining an undifferentiated state, which comprises the medium additive according to any one of claims 42 to 66.

68. A culture method for a proliferation of an induced pluripotent stem cell (iPS cell) while maintaining an undifferentiated state, comprising a step of cultivating the induced pluripotent stem cell (iPS cell) in a medium added with at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof.

69. The method according to claim 68, wherein the concentration of at least one selected from the group consisting of

ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof in the medium is 1 μM - 1000 μM.

70. The method according to claim 68, wherein the concentration of at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof in the medium is 5 μM - 200 μM.

71. The method according to any one of claims 68 to 70, wherein the ethanolamine analog is a compound represented by the following formula

$X-CH_2-CH_2-O-Y$

wherein

X is $R^1-N(R^2)-$ [R^1 and R^2 are the same or different and each is a hydrogen atom or an amino-protecting group] or $R^3-CH=N-$ [R^3-CH is H-CH or a Shiff base type amino-protecting group]; and

Y is $-P(=O)(OH)-O-R^4$ [R^4 is $-CH_2-CH(O-R^5)-CH_2-O-R^6$ (R^5 and R^6 are the same or different and each is an acyl group having 2 - 30 carbon atoms or a hydrogen atom) or a hydrogen atom], a hydrogen atom or a hydroxy-protecting group.

72. The method according to claim 71, wherein R^1 and R^2 are the same or different and each is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms, and R^3 is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1

- 6 carbon atoms.

73. The method according to any one of claims 68 to 72, wherein the ethanolamine analog is one or plural selected from the group consisting of phosphoethanolamine, monomethylethanolamine, dimethylethanolamine, N-acylphosphatidylethanolamine, phosphatidylethanolamine, and lysophosphatidylethanolamine.

74. The method according to any one of claims 68 to 73, wherein the medium is further added with albumin.

75. The method according to claim 74, wherein the concentration of the albumin in the medium is 0.1 g/l - 20 g/l.

76. The method according to claim 74, wherein the concentration of the albumin in the medium is 1 g/l - 8 g/l.

77. The method according to any one of claims 74 to 76, wherein the albumin is obtained from plasma of an animal (including human) or by gene recombination technology.

78. The method according to any one of claims 74 to 77, wherein the amount of fatty acid carried by the albumin in the medium is not more than 9 mg/g.

79. The method according to any one of claims 74 to 77, wherein the amount of fatty acid carried by the albumin in the medium is not more than 2.2 mg/g.

80. The method according to any one of claims 68 to 79, wherein the culture is performed in the absence of a feeder cell.

81. The method according to claim 80, wherein the culture is performed by using an extracellular matrix or an active fragment thereof, or an artificial product mimicking the

function thereof.

82. The method according to claim 80, wherein the culture is performed by using laminin 511 or an active fragment thereof or matrigel.

83. The method according to any one of claims 68 to 82, wherein the culture is performed by single cell-seeding.

84. The method according to any one of claims 68 to 83, wherein the culture is performed under serum-free conditions.

85. The method according to any one of claims 68 to 84, wherein the medium is substantially free of a component derived from an animal other than human.

86. The method according to any one of claims 68 to 85, wherein the induced pluripotent stem cell is derived from primates.

87. The method according to any one of claims 68 to 86, wherein the induced pluripotent stem cell is human iPS cell.

88. A preservation stabilizing method for a medium for a proliferation of a pluripotent stem cell, comprising adding at least one selected from the group consisting of ethanolamine, ethanolamine analogs and pharmaceutically acceptable salts thereof.

89. The method according to claim 88, further comprising adding sulfated saccharide and/or a pharmaceutically acceptable salt thereof.

90. The method according to claim 89, wherein the final concentration of the sulfated saccharide and/or a pharmaceutically acceptable salt thereof in use is 1 - 1000

ng/ml.

91. The method according to any one of claims 88 to 90, further comprising adding albumin.

92. The method according to claim 91, wherein the final concentration of the albumin in use is 0.1 g/l - 20 g/l.

93. The method according to claim 91, wherein the final concentration of the albumin in use is 1 g/l - 8 g/l.

94. The method according to any one of claims 91 to 93, wherein the albumin is obtained from plasma of an animal (including human) or by gene recombination technology.

95. The method according to any one of claims 91 to 94, wherein the amount of fatty acid carried by the albumin is not more than 9 mg/g.

96. The method according to any one of claims 91 to 94, wherein the amount of fatty acid carried by the albumin is not more than 2.2 mg/g.

97. The method according to any one of claims 88 to 96, wherein the medium is substantially free of β-mercaptoethanol or contains β-mercaptoethanol at a final concentration of not more than 9 μM.

98. The method according to any one of claims 88 to 97, wherein the medium is for a proliferation of a pluripotent stem cell while maintaining an undifferentiated state.

99. A medium additive for a proliferation of a pluripotent stem cell while maintaining an undifferentiated state, comprising at least one selected from the group consisting of ethanolamine,

ethanolamine analogs and pharmaceutically acceptable salts thereof.

100. The medium additive according to claim 99, wherein the ethanolamine analog is a compound represented by the following formula

$X-CH_2-CH_2-O-Y$

wherein

X is $R^1-N(R^2)-$ [R^1 and R^2 are the same or different and each is a hydrogen atom or an amino-protecting group] or $R^3-CH=N-$ [R^3-CH is H-CH or a Shiff base type amino-protecting group]; and

Y is $-P(=O)(OH)-O-R^4$ [R^4 is $-CH_2-CH(O-R^5)-CH_2-O-R^6$ (R^5 and R^6 are the same or different and each is an acyl group having 2 - 30 carbon atoms or a hydrogen atom) or a hydrogen atom], a hydrogen atom or a hydroxy-protecting group.

101. The medium additive according to claim 100, wherein R^1 and R^2 are the same or different and each is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms, and

R^3 is a hydrogen atom, a halogen atom, a hydroxy group, an aryl group, an acyl group having 2 - 30 carbon atoms, an alkyl group having 1 - 6 carbon atoms, an alkoxyl group having 1 - 6 carbon atoms, a hydroxyalkyl group having 1 - 6 carbon atoms, a haloalkyl group having 1 - 6 carbon atoms, a haloalkoxyl group having 1 - 6 carbon atoms or a halohydroxyalkyl group having 1 - 6 carbon atoms.

102. The medium additive according to any one of claims 99 to 101, wherein the ethanolamine analog is one or plural selected

from the group consisting of phosphoethanolamine, monomethylethanolamine, dimethylethanolamine, N-acylphosphatidylethanolamine, phosphatidylethanolamine, and lysophosphatidylethanolamine.

103. The medium additive according to any one of claims 99 to 102, further comprising albumin.

104. The medium additive according to claim 103, wherein the albumin is obtained from plasma of an animal (including human) or by gene recombination technology.

105. The medium additive according to any one of claims 99 to 104, further comprising sulfated saccharide and/or a pharmaceutically acceptable salt thereof.

106. The medium additive according to claim 105, wherein the final concentration of the sulfated saccharide and/or a pharmaceutically acceptable salt thereof in use is 1 - 1000 ng/ml.

107. The medium additive according to claim 105 or 106, wherein the sulfated saccharide is at least one selected from the group consisting of sulfated monosaccharide, sulfated disaccharide, sulfated polysaccharide, sulfated sugar alcohol and sulfated cyclitol.

108. The medium additive according to any one of claims 105 to 107, wherein the aforementioned sulfated saccharide or a pharmaceutically acceptable salt thereof is at least one selected from the group consisting of dextran sulfate Na, cellulose SO_3Na, xanthan gum SO_3Na, fucoidan, alginate SO_3Na, inulin SO_3Na, Maltoheptaose SO_3Na, stachyose SO_3Na, maltotriose SO_3Na, multitol SO_3Na, sucrose8SO_3K, glucose SO_3Na, myo-6 inositol SO_3K, α-cyclodextrin SO_3Na, mannitol SO_3Na, xylitol

SO_3Na and erythritol SO_3Na.

109. The medium additive according to claim 108, wherein the aforementioned sulfated saccharide or a pharmaceutically acceptable salt thereof is at least one selected from the group consisting of dextran sulfate Na, fucoidan, Maltoheptaose SO_3Na, maltotriose SO_3Na, multitol SO_3Na and sucrose8SO_3K.

110. The medium additive according to any one of claims 105 to 109, wherein the aforementioned sulfated saccharide or a pharmaceutically acceptable salt thereof is dextran sulfate Na having an average molecular weight of 2,500 - 7,500.

111. The medium additive according to any one of claims 99 to 110, wherein the proliferation of a pluripotent stem cell while maintaining an undifferentiated state is performed under the conditions without using a feeder cell.

112. The medium additive according to claim 111, wherein the conditions without using a feeder cell do not include use of a feeder cell but include use of an extracellular matrix or an active fragment thereof, or an artificial product mimicking the function thereof.

113. The medium additive according to claim 111, wherein the conditions without using a feeder cell do not include use of a feeder cell but include use of laminin 511 or an active fragment thereof or matrigel.

114. The medium additive according to any one of claims 99 to 113, wherein the proliferation of a pluripotent stem cell while maintaining an undifferentiated state is performed by single cell-seeding.

115. The medium additive according to any one of claims 99 to

114, wherein the proliferation of a pluripotent stem cell while maintaining an undifferentiated state is performed under serum-free conditions.

116. The medium additive according to any one of claims 99 to 115, comprising substantially free of a component derived from an animal other than human.

117. The medium additive according to any one of claims 99 to 116, wherein the pluripotent stem cell is an embryonic stem cell (ES cell) or induced pluripotent stem cell (iPS cell).

118. The medium additive according to any one of claims 99 to 117, wherein the pluripotent stem cell is derived from primates.

119. The medium additive according to any one of claims 99 to 118, wherein the pluripotent stem cell is human iPS cell.

120. A medium for proliferation of a pluripotent stem cell while maintaining an undifferentiated state, which comprises the medium additive according to any one of claims 99 to 119.