CLAIMS

1. A composition for treating and/or preventing Hepatitis B virus infection in a human or non-human primate, comprising polyriboinosinic acid-polyribocytidylic acid (PIC); at least one antibiotic or polyamide compound selected from spermidine salt, spermidine, N-(3-aminopropyl), N-(3-aminopropyl)-1,4-butandiamine, spermine BR, spermine, OS-dimethylphosphoramidothioate, polylysine, and aminoglycoside; at least one positive ion selected from calcium, cadmium, lithium, magnesium, cerium, cesium, chromium, cobalt, deuterium, gallium, iodine, iron, and zinc; and HBsAg, wherein the concentration of the HBsAg in the composition ranges from 20 μg/unit dose to 60 μg/unit dose.

2. The composition of claim 1, further comprising HBcAg.

3. The composition of claim 1 or 2, wherein the HBsAg is produced *by Hansenula polymorpha*, CHO cells, insect expression system, *Saccharomyces cerevisiae*, or *Pichia pastoris*.

4. The composition of claim 2, wherein the HBcAg is produced by *Escherichia coli*, *Hansenula polymorpha*, insect expression system, *Saccharomyces cerevisiae*, or *Pichia pastoris*.

5. The composition of claim 1 or 2, wherein the composition is for administration to a Hepatitis B virus uninfected subject, a chronic Hepatitis B virus subject, or an acute Hepatitis B virus subject.

6. The composition of claim 1 or 2, wherein the composition is for administration to a subject by one way selected from a group including intramuscular injection, intraperitoneal injection, intravenous injection, subcutaneous injection, transdermal

delivery, intradermal delivery, nasal delivery, opthamalical delivery, oral delivery, sublingual delivery, peritumoral delivery, and intratumor delivery.

7. The composition of claim 1 or 2, wherein the antibiotic is selected from a group including tobramycin, anthracycline, butyrosin sulfate, gentamicin, hygromycin, amikacin, dibekacin, nebramycin, beta-lactam, neomycin, puromycin, streptomycin, streptozotocin, and kanamycin.

8. The composition of any one of claims 1 to 7, wherein the ratio of the HBsAg to PIC is 1:30, 1:25, or 1:20.

9. The composition of claim 2 or 4, wherein the ratio of the HBcAg to PIC is 1:30, 1:25, 1:20.

10. The composition of any one of the preceding claims, wherein the concentration of PIC in the composition ranges from 250 μg/unit dose to 6000 μg/unit dose.

11. The composition of claim 10, wherein the concentration of PIC in the composition ranges from 500 μg/unit dose to 1500 μg/unit dose.

12. The composition of any one of the preceding claims, wherein the concentration of the HBsAg in the composition is 20 μg/unit dose, 30 μg/unit dose, 40 μg/unit dose, 50 μg/unit dose, or 60 μg/unit dose, .

13. The composition of claim 2, wherein the concentration of the HBcAg in the composition ranges from 10 μg/unit dose to 100 μg/unit dose.

14. The composition of claim 13, wherein the concentration of the HBcAg in the composition is 10 μg/unit dose, 20 μg/unit dose, 30 μg/unit dose, 40 μg/unit dose, 50 μg/unit dose, 60 μg/unit dose, 70 μg/unit dose, 80 μg/unit dose, 90 μg/unit dose,

or 100 μg/unit dose.

15. The composition of any of the preceding claims, wherein one unit dose ranges from 0.1 ml to 250.0 ml.

16. The composition of claim 15, wherein said unit dose is 0.1 ml, 0.15 ml, 0.2 ml, 0.5 ml, 1.0 ml, 1.5 ml, 2.0 ml, 2.5 ml, 3.0 ml, 4.0 ml, 5.0 ml, 10.0 ml, 20.0 ml, 30.0 ml, 40.0 ml, 50.0 ml, 60.0 ml, 70.0 ml, 80.0 ml, 90.0 ml, 100.0 ml, 150.0 ml, 200.0 ml, or 250.0 ml.

17. The composition of claim 1 or 2, wherein the composition further comprises at least one substance selected from a group including gelatin, saccharose, granulated sugar, lactose, maltose, trehalose, glucose, low molecular dextran, sorbitol, polysorbate 20, mannitol PEG, human blood albumin, recombinant human serum albumin, sodium caprylate, urea, aluminum hydroxide, phenol red, magnesium chloride, potassium chloride, sodium chloride, sodium thiosulphate, potassium dihydrogen, ascorbic acid, trichloromethane, phenol, and thiomersal.

18. The composition of claim 1 or 2, the composition further comprising at least one physiologically acceptable buffer, wherein the physiological acceptable buffer is selected from a group including acetate buffer, trisamine buffer, bicarbonate buffer, carbonate buffer, and phosphate buffered saline.

19. The composition of claim 1 or 2, wherein the pH of said composition ranges from 6.0 to 8.0.

20. The composition of claim 1 or 2, wherein the composition is in a liquid or solid form, wherein said liquid form is selected from a group including parenteral

solution, suspension, ointment, emulsion, drop, syrup, and gel; wherein said solid form is selected from a group including dry powder, freeze-dried powder, tablet, capsule, suppository, granule, and sugar-coated tablet.

21. The composition of any one of claims 1 to 20, for use in medicine.

22. Use of PIC, at least one antibiotic or polyamide compound selected from spermidine salt, spermidine, N-(3-aminopropyl), N-(3-aminopropyl)-1,4-butandiamine, spermine BR, spermine, OS-dimethylphosphoramidothioate, polylysine, and aminoglycoside, at least one positive ion selected from calcium, cadmium, lithium, magnesium, cerium, cesium, chromium, cobalt, deuterium, gallium, iodine, iron, and zinc, and HBsAg, in the preparation of a pharmaceutical composition for treating and/or preventing Hepatitis B virus infection in a human or non-human primate, wherein the concentration of the HBsAg in the composition ranges from 20 μg/unit dose to 60 μg/unit dose.

23. The use of claim 22, wherein the composition further comprises HBcAg.

24. The use of claim 22 or 23, wherein the composition is to be administered to a subject once every two months, twice every two months, three times every two months, four times every two months, five times every two months, once every month, twice every month, once every three weeks, twice every three weeks, twice every two weeks, once every two weeks, or once every week.

25. The use of any one of claims 22-24, wherein said subject is a HBV uninfected subject, a chronic HBV subject, or an acute HBV subject.