In certain embodiments, one or more glutamine (Q) in SEQ ID NO: 16 deamidates to glutamic acid (Glu), i.e., α-glutamic acid, γ-glutamic acid (Glu), or a blend of α- and γ-glutamic acid, which may interconvert through a common glutarinimide intermediate. Any suitable ratio of α- and γ-glutamic acid may be present. For example, in certain embodiments, the ratio may be from 10:1 to 1:10 α to γ, about 50:50 α: γ, or about 1:3 α : γ, or another selected ratio.



Thus, an rAAVhu68 includes subpopulations within the rAAVhu68 capsid of vp1, vp2 and/or vp3 proteins with deamidated amino acids, including at a minimum, at least one subpopulation comprising at least one highly deamidated asparagine. In addition, other modifications may include isomerization, particularly at selected aspartic acid (D or Asp) residue positions. In still other embodiments, modifications may include an amidation at an Asp position.

In certain embodiments, an AAVhu68 capsid contains subpopulations of vp1, vp2 and vp3 having at least 4 to at least about 25 deamidated amino acid residue positions, of which at least 1 to 10% are deamidated as compared to the encoded amino acid sequence of SEQ ID NO: 16. The majority of these may be N residues. However, Q residues may also be deamidated.

In certain embodiments, an AAV68 capsid is further characterized by one or more of the following. AAV hu68 capsid proteins comprise: AAVhu68 vp1 proteins produced by expression from a nucleic acid sequence which encodes the predicted amino acid sequence

non-enzymatic in nature, being caused by functional groups within the capsid protein which deamidate selected asparagines, and to a lesser extent, glutamine residues. Efficient capsid assembly of the majority of deamidation vp1 proteins indicates that either these events occur following capsid assembly or that deamidation in individual monomers (vp1, vp2 or vp3) is well-tolerated structurally and largely does not affect assembly dynamics. Extensive deamidation in the VP1-unique (VP1-u) region (~aa 1-137), generally considered to be located internally prior to cellular entry, suggests that VP deamidation may occur prior to capsid assembly.

Without wishing to be bound by theory, the deamidation of N may occur through its C-terminus residue's backbone nitrogen atom conducts a nucleophilic attack to the Asn's side chain amide group carbon atom. An intermediate ring-closed succinimide residue is believed to form. The succinimide residue then conducts fast hydrolysis to lead to the final product aspartic acid (Asp) or iso aspartic acid (IsoAsp). Therefore, in certain embodiments, the deamidation of asparagine (N or Asn) leads to an Asp or IsoAsp, which may interconvert through the succinimide intermediate e.g., as illustrated below.



As provided herein, each deamidated N of SEQ ID NO: 16 may independently be aspartic acid (Asp), isoaspartic acid (isoAsp), aspartate, and/or an interconverting blend of Asp and isoAsp, or combinations thereof. Any suitable ratio of α- and isoaspartic acid may be present. For example, in certain embodiments, the ratio may be from 10:1 to 1:10 aspartic to isoaspartic, about 50:50 aspartic: isoaspartic, or about 1:3 aspartic: isoaspartic, or another selected ratio.