


R_{11} and R_{12} are independently hydrogen, halogen, $-C_{1-10}$ alkyl, $-C_{2-10}$ alkenyl, $-C_{2-10}$ alkynyl, $-C_{1-10}$ heteroalkyl, $-C_{3-10}$ aryl, $-C_{1-10}$ hetaryl, $-C_{3-10}$ cycloalkyl, $-C_{1-10}$ heterocyclyl, $-OH$, $-CF_3$, $-OCF_3$, $-OR^{31}$, $-NR^{31}R^{32}$, $-C(O)R^{31}$, $-CO_2R^{31}$, $-C(=O)NR^{31}$, $-NO_2$, $-CN$, $-S(O)_{0-2}R^{31}$, $-SO_2NR^{31}R^{32}$, $-NR^{31}C(=O)R^{32}$, $-NR^{31}C(=O)OR^{32}$, $-NR^{31}C(=O)NR^{32}R^{33}$, $-NR^{31}S(O)_{0-2}R^{32}$, $-C(=S)OR^{31}$, $-C(=O)SR^{31}$, $-NR^{31}C(=NR^{32})NR^{32}R^{33}$, $-NR^{31}C(=NR^{32})OR^{33}$, $-NR^{31}C(=NR^{32})SR^{33}$, $-OC(=O)OR^{33}$, $-OC(=O)NR^{31}R^{32}$, $-OC(=O)SR^{31}$, $-SC(=O)SR^{31}$, $-P(O)OR^{31}OR^{32}$, or $-SC(=O)NR^{31}NR^{32}$; and each of R^{31} , R^{32} , and R^{33} is independently hydrogen, halogen, $-C_{1-10}$ alkyl, $-C_{2-10}$ alkenyl, $-C_{2-10}$ alkynyl, $-C_{1-10}$ heteroalkyl, $-C_{3-10}$ aryl, $-C_{1-10}$ hetaryl, $-C_{3-10}$ cycloalkyl, or $-C_{1-10}$ heterocyclyl, or R^{31} together with R^{32} form a heterocyclic ring.

[0006b] In one embodiment, there is provided use of the compound as described herein in the manufacture of a medicament for treatment of a disorder mediated by ERK1 and/or ERK2.


[0006c] In one embodiment, there is provided use of the compound as described herein in the manufacture of a medicament for treatment of cancer, optionally wherein the cancer is selected from the group consisting of breast cancer, pancreatic cancer, non-small cell lung cancer (NSCLC), thyroid cancer, seminomas, melanoma, bladder cancer, liver cancer, kidney cancer, myelodysplastic syndrome (MDS), acute myelogenous leukemia (AML), and colorectal cancer.

[0007] In some embodiments, the invention provides a compound of Formula I-A:




or a pharmaceutically acceptable salt or prodrug thereof.

[0008] In some embodiments, the invention provides a compound of Formula I-B:



or a pharmaceutically acceptable salt or prodrug thereof.

[0009] In some cases, the invention provides a compound of Formula II:




or a pharmaceutically acceptable salt or prodrug thereof.

In some embodiments, X_1 is $C=O$. In some embodiments, X_2 is NR_1 . In some embodiments, X_5 is N and X_6 is C . In other embodiments, X_5 is C and X_6 is C . In some embodiments, X_9 is CR_{21} .

Claims

1. A compound of Formula II-E:



or a pharmaceutically acceptable salt thereof; wherein:

R_1 is $-C_{1-10}$ heterocyclyl, $-C_{1-10}$ heterocyclyl- C_{1-10} alkyl, or $-C_{1-10}$ heterocyclyl- C_{3-10} aryl, each of which is unsubstituted or substituted by one or more independent R_{10} or R_{11} substituents;

R_{21} is $-L-C_{1-10}$ heteroalkyl, $-L-C_{3-10}$ aryl, $-L-C_{1-10}$ hetaryl, $-L-C_{3-10}$ cycloalkyl, or $-L-C_{1-10}$ heterocyclyl, each of which is unsubstituted or substituted by one or more independent R_{12} substituents;

L is a bond, $-O-$, $-N(R^{31})-$, $-S(O)_{0-2}-$, $-C(=O)-$, $-C(=O)O-$, $-OC(=O)-$, $-C(=O)N(R^{31})-$, $-N(R^{31})C(=O)-$, $-NR^{31}C(=O)O-$, $-NR^{31}C(=O)NR^{32}-$, $-NR^{31}S(O)_{0-2}-$, $-S(O)_{0-2}N(R^{31})-$, $-C(=S)O-$, $-C(=O)S-$, $-NR^{31}C(=NR^{32})NR^{32}-$, $-NR^{31}C(=NR^{32})O-$, $-NR^{31}C(=NR^{32})S-$, $-OC(=O)O-$, $-OC(=O)NR^{31}-$, $-OC(=O)S-$, $-SC(=O)S-$, $-P(O)OR^{31}O-$, or $-SC(=O)NR^{31}-$;

R_{72} is hydrogen;

R_{10} is $-C_{1-10}$ alkyl, $-C_{2-10}$ alkenyl, $-C_{2-10}$ alkynyl, $-C_{1-10}$ heteroalkyl, $-C_{3-10}$ aryl, $-C_{1-10}$ hetaryl, $-C_{3-10}$ cycloalkyl, or $-C_{1-10}$ heterocyclyl, each of which is unsubstituted or substituted by one or more independent R_{11} substituents;

R_{11} and R_{12} are independently hydrogen, halogen, $-C_{1-10}$ alkyl, $-C_{2-10}$ alkenyl, $-C_{2-10}$ alkynyl, $-C_{1-10}$ heteroalkyl, $-C_{3-10}$ aryl, $-C_{1-10}$ hetaryl, $-C_{3-10}$ cycloalkyl, $-C_{1-10}$ heterocyclyl, $-OH$, $-CF_3$, $-OCF_3$, $-OR^{31}$, $-NR^{31}R^{32}$, $-C(O)R^{31}$, $-CO_2R^{31}$, $-C(=O)NR^{31}$, $-NO_2$, $-CN$, $-S(O)_{0-2}R^{31}$, $-SO_2NR^{31}R^{32}$, $-NR^{31}C(=O)R^{32}$, $-NR^{31}C(=O)OR^{32}$, $-NR^{31}C(=O)NR^{32}R^{33}$, $-NR^{31}S(O)_{0-2}R^{32}$, $-C(=S)OR^{31}$, $-C(=O)SR^{31}$, $-NR^{31}C(=NR^{32})NR^{32}R^{33}$, $-NR^{31}C(=NR^{32})OR^{33}$, $-NR^{31}C(=NR^{32})SR^{33}$, $-OC(=O)OR^{33}$, $-OC(=O)NR^{31}R^{32}$, $-OC(=O)SR^{31}$, $-SC(=O)SR^{31}$, $-P(O)OR^{31}OR^{32}$, or $-SC(=O)NR^{31}NR^{32}$; and

each of R^{31} , R^{32} , and R^{33} is independently hydrogen, halogen, $-C_{1-10}$ alkyl, $-C_{2-10}$ alkenyl, $-C_{2-10}$ alkynyl, $-C_{1-10}$ heteroalkyl, $-C_{3-10}$ aryl, $-C_{1-10}$ hetaryl, $-C_{3-10}$ cycloalkyl, or $-C_{1-10}$ heterocyclyl, or R^{31} together with R^{32} form a heterocyclic ring.

2. The compound of claim 1, wherein R_{21} is selected from the group consisting of $-L-C_{1-10}$ heteroalkyl, $-L-C_{3-10}$ aryl, $-L-C_{1-10}$ hetaryl, $-L-C_{3-10}$ cycloalkyl, and $-L-C_{1-10}$ heterocyclyl, each of which is unsubstituted or substituted by one or more independent R_{12} substituents; and wherein L is a bond.

3. The compound of claim 2, wherein R_{21} is $-L-C_{1-10}$ hetaryl unsubstituted or substituted by one or more independent R_{12} substituents; and wherein L is a bond.

4. The compound of claim 3, wherein the C_{1-10} hetaryl of R_{21} comprises one or more nitrogen atoms.


5. The compound of claim 4, wherein the C_{1-10} hetaryl of R_{21} is selected from the group consisting of pyrazolyl, pyridinyl, pyrazinyl, pyrimidinyl, and pyridazinyl.

6. The compound of claim 5, wherein the C_{1-10} hetaryl of R_{21} is unsubstituted.

7. The compound of claim 5, wherein the C_{1-10} hetaryl or R_{21} is substituted with one, two, or three independent R_{12} substituents.

8. The compound of any one of claims 2-5 or 7, wherein each R_{12} substituent is independently selected from the group consisting of $-C_{1-10}$ alkyl, $-C_{2-10}$ alkenyl, $-C_{2-10}$ alkynyl, $-C_{1-10}$ heteroalkyl, $-C_{3-10}$ aryl, $-C_{1-10}$ hetaryl, $-C_{3-10}$ cycloalkyl, $-C_{1-10}$ heterocyclyl, $-OH$, $-CF_3$, $-OCF_3$, and $-OR^{31}$; wherein each R^{31} is independently hydrogen or $-C_{1-10}$ alkyl.

9. The compound of claim 8, wherein each R_{12} substituent is independently selected from the group consisting $-Me$, $-Et$, $-i-Pr$, $-n-Pr$, $-OH$, $-OMe$, $-OEt$, and $-OPr$.



11. Use of the compound of any one of claims 1-10 for the manufacture of a medicament for treatment of a disorder mediated by ERK1 and/or ERK2.

12. The use of claim 11, wherein the disorder is cancer.

13. Use of the compound of any one of claims 1-10 for the manufacture of a medicament for treatment of cancer, optionally wherein the cancer is selected from the group consisting of breast cancer, pancreatic cancer, non-small cell lung cancer (NSCLC), thyroid cancer, seminomas, melanoma, bladder cancer, liver cancer, kidney cancer, myelodysplastic syndrome (MDS), acute myelogenous leukemia (AML), and colorectal cancer.

14. A pharmaceutical composition comprising the compound of any one of claims 1-10 and a pharmaceutically acceptable excipient, optionally wherein the composition is formulated in an oral dosage.

15. The compound of claim 1, wherein the compound is selected from:

