

DESCRIPTION

METHOTREXATE DERIVATIVE


[Technical Field]

This invention relates to a novel methotrexate
5 derivative, and, more specifically, to a novel methotrexate
derivative useful as an antirheumatic agent.

[Technical Background]

Hitherto, methotrexate has been used for many years
as a treating agent for leukemia, but, since Gubner et al
10 reported in 1951 the effectiveness by using aminopterin
for rheumatoid arthritis (RA) and psoriasis, an attention
has been drawn to aminopterin as a treating agent for RA and
it has mainly been used in Europe and USA. In relatively
recent years, the method of use and the dosage thereof have
15 been studied in more detail, and it becomes apparent that
the treatment at a low dose of methotrexate exhibits low
side effects and excellent effectiveness. However, since
side effects such as hepatic disorders and lung fibrosis
caused by administration of methotrexate is not negligible,
20 development of a drug having less side effects and yet
having excellent effects is desired.


Hitherto, methotrexate derivatives in which an alkyl
group other than a methyl group has been introduced into N¹⁰,
for example, having the following formula:



This Sheet Has Been Replaced

THIS SHEET HAS BEEN
REPLACED

(J. Med. Chem., 22, 862 (1979) and having the following formula:




(J. Med. Chem., 25, 877 (1982) have been known, but none of these compounds showed a satisfactory activity.

5 The present inventors made extensively studies for developing compounds having a good balance between a pharmacological effect as an antirheumatic agent and side effects in the above type of methotrexate derivatives, and as a result completed the present invention.

10 [Disclosure of the Invention]

The present invention relates to a methotrexate derivative represented by the following general formula (I):



wherein R_1 and R_2 may be the same or different and each represents a hydrogen atom or a lower alkyl group having
 15 from 1 to 4 carbon atoms, and a salt thereof. Further, the present invention provides an antirheumatic agent containing at least one of these compounds as an effective ingredient.

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

[Brief Description of Drawing]

Fig. 1 shows a percent cell propagation at each of the concentrations of the test drugs.






[Best Mode for Working the Invention]

5 The compounds according to the present invention are described as a general formula in broad concept in WO 92/9436, but are novel compounds, none of which has been actually produced and have not been disclosed in literature references. These compounds are synthesized, for example,
10 in the following manner:

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED



wherein R_1 and R_2 may be the same or different and each represents a hydrogen atom or a lower alkyl group having from 1 to 4 carbon atom, A represents a protective group,

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED



and X represents a halogen atom.

The reaction for obtaining a compound of the general formula (3) from a compound of the general formula (1) and a compound of the general formula (2) is performed by an amido-bond forming reaction generally used. For example, the compound of the general formula (1) is suspended in an acid halogenating agent such as thionyl chloride and oxalyl chloride and stirred in the presence of a catalytic amount of dimethylformamide, etc. at room temperature to obtain an acid halide which is then dissolved in a solvent such as dichloromethane and added to an aqueous solution of the compound of the general formula (2) under ice cooling or water cooling, followed by stirring at room temperature in the presence of an inorganic salt such as potassium carbonate, sodium hydroxide and sodium bicarbonate.

Examples of the protective group represented by A in the formula include a protective group for an amino group such as an acyl group, and preferably, a carbobenzoxy group, a tosyl group and an acetyl group. The compound of the general formula (1) can be obtained, for example, by the method described in International Publication WO 92/9436.

The compound of the general formula (3) is de-protected by the usual method thereby converting into the compound of the general formula (4). For example, when the protective group is a carbobenzoxy group or a tosyl group, removal of the protective group is achieved by adding the compound of the general formula (3) to a solution of anisole or phenol dissolved in a hydrogen bromide-acetic

acid solution, followed by stirring at from 10°C to 60°C, preferably at room temperature. Also, when the protective group is a carbobenzoxy group, removal of the protective group may be conducted by dissolving the compound of the
5 general formula (3) in a solvent such as methanol, ethanol or acetic acid, and, after adding palladium-carbon thereto, the mixture is stirred in a hydrogen atmosphere at room temperature.





The reaction for obtaining the compound of the
10 general formula (6) from the compound of the general formula (4) and the compound of the general formula (5) is carried out by stirring the compound of the general formula (4) and the compound of the general formula (5) in a solvent such as dimethylacetamide or dimethylformamide at from 0°C to 100°C,
15 preferably from 50°C to 70°C. In particular, when R₁ and R₂ are hydrogen atoms, an aqueous alkali solution such as 1N aqueous solution of sodium hydroxide is added to a solvent such as methanol or ethanol, and the stirring is conducted at from 15°C to 35°C to obtain the object compound.
20 Examples of the halogen atom represented by X in the formula include a bromine atom and a chlorine atom.

The compound of the present invention can also be synthesized by the following process:

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED



wherein R_1 , R_2 and R_3 may be the same or different and each represents a hydrogen atom or a lower alkyl group having from 1 to 4 carbon atoms, and X represents a halogen atom.




The reaction for obtaining the compound of the general formula (8) from the compound of the general formula (5) and the compound of the general formula (7) is carried out in the same manner as described for obtaining
5 the compound of the general formula (6) from the compound of the general formula (4) and the compound of the general formula (5). The compound of the general formula (7) can be obtained, for example, by the process described in International Publication WO 92/9436.

10 The reaction for obtaining the compound of the general formula (6) from the compound of the general formula (8) and the compound of the general formula (2) is carried out by the amido-bond forming reaction generally used.

15 The compound of the present invention can also be obtained as a salt thereof by the usual method. The salt which can be used include, for example, an inorganic acid salt such as hydrochloride, hydrobromide, hydroiodide, sulfate and phosphate, an organic acid salt such as
20 succinate, malonate, acetate, maleate, fumarate, oxalate, gluconate, mandelate, benzoate, salicylate, methane-sulfonate, benzenesulfonate and p-toluenesulfonate, and a metal salt such as a sodium salt, a potassium salt and a magnesium salt, and preferably an inorganic salt and
25 an organic salt, and more preferably hydrobromide or methanesulfonate.

The pharmaceutical agent containing the compound of the present invention can be administered orally or



parenterally, but the oral administration or a topical administration such as intra-articular is preferred. The dose to be administered varies depending upon the body weight and the conditions of disease of patients, but is
5 generally preferably at from 0.01 to 100 mg/day/person.

The form of the pharmaceutical agent containing the compound of the present invention include a liquid preparation such as injection, tablet, capsule, powder, etc.

The compounds represented by the general formula (I)
10 according to the present invention are excellent in the balance between the antirheumatic activity and the toxicity as compared with methotrexate and the existing methotrexate derivatives and, therefore, are useful as antirheumatic agents.

15 [Examples]

The compounds of the present invention are further illustrated in more detail with reference to examples thereof, but the present invention is not limited by these examples.

20 [Reference Example 1]

Synthesis of dimethyl N-(4-carbobenzoxy-3,4-dihydro-2H-1,4-benzothiazine-7-carbonyl)-L- α -aminoadipate

Thionyl chloride (10 ml) was added to 4-carbobenzoxy-3,4-dihydro-2H-1,4-benzothiazine-7-carboxylic acid (2.0 g),
25 followed by stirring at room temperature for 2 hours. Then, the reaction solution was concentrated under reduced pressure to dryness. The resulting solid material was dissolved in dichloromethane (25 ml), and to the resulting

solution were added subsequently an aqueous solution (25 ml) of dimethyl L- α -aminoadipate hydrochloride (1.4 g) and then potassium carbonate (3.4 g), followed by stirring overnight. The reaction mixture was poured into a saturated aqueous solution of sodium bicarbonate, and an organic substance was extracted with chloroform. Further, the chloroform layer was washed with 1N hydrochloric acid, dried over sodium sulfate, and the solvent was distilled off under reduced pressure. The resulting residue was subjected to silica gel column chromatography using chloroform:methanol = 100:1 as an eluting solvent to obtain the titled compound (2.35 g).

^1H NMR (CDCl_3 , δ values):

1.6-2.1 (4H,m), 2.37 (2H,t,J=7.1Hz), 3.20 (2H,m),
3.67 (3H,s), 3.78 (3H,s), 3.98 (2H,m), 4.77
(1H,m), 5.23 (2H,s), 6.75 (1H,d,J=7.3Hz),
7.36 (5H,m), 7.48 (2H,m), 7.63 (1H,d,J=1.5Hz)

[Reference Example 2]

Synthesis of dimethyl N-(3,4-dihydro-2H-1,4-benzothiazine-7-carbonyl)-L- α -aminoadipate

The compound (14.8 g) of Reference Example 1 was added to a 30% hydrogen bromide-acetic acid solution (200 ml) of amisole (15 g), followed by stirring at room temperature for 4 hours. Then, a large amount of ether was added to the reaction mixture whereby a red brown oily substance was precipitated. Most of the ethereal layer was removed and the oily substance was suspended in chloroform. The resulting suspension was washed with a saturated aqueous solution of sodium bicarbonate and extracted with

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

chloroform. The chloroform layer was dried over sodium sulfate, and the solvent was distilled off under reduced pressure to obtain the titled compound (7.6 g).

¹H NMR (CDCl₃-CD₃OD, δ values):

5 1.6-2.0 (4H,m), 2.40 (2H,t,J=6.8Hz), 3.01 (2H,m),
 3.68 (5H,m), 3.78 (3H,s), 4.65 (1H,m), 6.50
 (1H,d,J=8.8Hz), 7.41 (2H,m), 7.51 (1H,d,J=2.0Hz)

[Example 1]

10 Synthesis of dimethyl N-(1-((2,4-diamino-6-pteridinyl)methyl)-3,4-dihydro-2H-1,4-benzothiazine-7-carbonyl)-L-α-aminoadipate

 The compound (7.6 g) of Reference Example 2 and a 6-bromomethyl-2,4-diaminopteridine hydrobromide salt isopropanol adduct (8.2 g) were suspended in dimethyl-
15 acetamide (120 ml), followed by stirring at 60°C for 13 hours. After cooling, the reaction mixture was poured into a saturated aqueous solution of sodium bicarbonate, and the mixture was extracted with a mixed solvent of chloroform:methanol = 10:1. The organic layer was dried
20 over sodium sulfate, and then the solvent was distilled off under reduced pressure. The resulting residue was subjected to silica gel column chromatography using chloroform:methanol = 10:1 as an eluting solvent to obtain the titled compound (5.8 g).


25 ¹H NMR (CDCl₃-CD₃OD, δ values):

 1.6-2.0 (4H,m), 2.39 (2H,t,J=7.1Hz), 3.17 (2H,m),
 3.68 (3H,s), 3.76 (3H,s), 3.94 (2H,m), 4.66 (1H,m),
 4.79 (2H,s), 6.70 (1H,d,J=8.8Hz), 7.42 (2H,m),

This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED



7.62 (1H,d,J=2.4Hz), 8.67 (1H,s)

[Example 2]

Synthesis of N-(1-((2,4-diamino-6-pteridinyl)methyl)-
3,4-dihydro-2H-1,4-benzothiazine-7-carbonyl)-L- α -
5 aminoadipic acid

The compound (5.8 g) of Example 1 was dissolved
in ethanol (300 ml), and 1N aqueous solution of sodium
hydroxide (32.2 ml) was added thereto at 35°C, followed by
stirring at room temperature overnight. Then, the reaction
10 solution was evaporated under reduced pressure to dryness.
The resulting solid substance was dissolved in water
(100 ml), and the solution was adjusted to a pH of 3.7
with 1N hydrochloric acid, followed by allowing to stand
in a cool place for 2 hours. The precipitate formed was
15 separated by filtration to obtain the titled compound
(4.5 g).

¹H NMR (DMSO-d₆ δ values):

1.5-1.9 (4H,m), 2.21 (2H,t,J=7.1Hz), 3.15 (2H,m),
3.96 (2H,m), 4.30 (1H,m), 4.76 (2H,s), 6.78
20 (1H,d,J=8.8Hz), 7.44 (1H,m), 7.60 (1H,s),
8.17 (1H,d,J=7.8Hz), 8.65 (1H,s)

[Example 3]

Synthesis of N-(1-((2,4-diamino-6-pteridinyl)methyl)-
3,4-dihydro-2H-1,4-benzothiazine-7-carbonyl)-L- α -
25 aminoadipic acid hydrobromide (2/3 hydrate)

The compound (100 mg) obtained in Example 2 was
suspended in hydrobromic acid (10 ml) and dissolved by
warming up to 60°C in a water bath. Then, the solution



was filtered through a cotton plug, and the filtrate was cooled to room temperature, followed by allowing to stand overnight. The precipitated needle crystals were separated by filtration and dried while heating in vacuum to obtain the titled compound (92 mg).

^1H NMR ($\text{DMSO}-d_6$, δ values):

1.42-1.90 (4H,m), 2.21 (2H,t,J=7.3Hz), 3.19 (2H,m), 3.97 (2H,m), 4.26 (1H,m), 4.84 (2H,s), 6.76 (1H,d,J=8.9Hz), 7.44 (1H,dd,J=2.2Hz,8.8Hz), 7.62 (1H,d,J=1.97Hz), 8.23 (1H,d,J=8.2Hz), 8.80 (1H,s)

Elementary Analysis

C:43.76, H:4.31, N:18.73, S:5.12, Br:13.43

Calc'd Values

C:43.73, H:4.37, N:18.54, S:5.31, Br:13.22


Empirical Formula

$\text{C}_{22}\text{H}_{24}\text{N}_8\text{O}_5\text{S} \cdot \text{HBr} \cdot 2/3\text{H}_2\text{O}$


[Test Example]

2.5×10^3 synovial cells from RA patient were cultured (in 96-well culture plate) with various concentrations of a test compound. Three days later, 1 μCi ^3H -UdR was added to each well and cultured for further 2 days. After culture, intracellular uptake of ^3H -UdR was measured by a scintillation counter.

The drugs used for the test were as follows:



THIS SHEET HAS BEEN
REPLACED



Control Compound

The results are shown in Fig. 1. As is apparent from Fig. 1, the compound of the present invention was confirmed to have a more excellent inhibitory activity to synovial cell proliferation than the control compound.

[Possibility of Utilization in Industry]

The compounds represented by the general formula (I) according to the present invention have excellent antirheumatic activity and are useful as antirheumatic agents in view of their low toxicity as compared to methotrexate.


This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED

CLAIMS

1. A compound represented by the general formula (I):



wherein R_1 and R_2 , which may be the same or different, each represents a hydrogen atom or a lower alkyl group having from 1 to 4 carbon atoms, or a salt thereof.

2. Antirheumatic agent containing, as an active ingredient, at least one of the compounds represented by the general formula (I) as defined in Claim 1 and the salts thereof.



This Sheet Has Been Replaced



THIS SHEET HAS BEEN
REPLACED