

Description

BACKGROUND OF THE INVENTION

5 Field of the Invention:

This invention relates to a mutant which is resistant to a cell membrane synthesis inhibitor, and, more particularly, to a cell membrane synthesis inhibitor-resistant mutant of an extracellular enzyme-producing microorganism belonging to genus Bacillus, such as cellulase-, protease-, or amylase-producing microor-
 10 ganism, belonging to genus Bacillus. The invention also relates to a process for preparing such a mutant.

Description of the Background:

Cellulase is realized as a group of complicated enzymes catalyzing enzymatic reactions decomposing
 15 cellulose into glucose, cellobiose, or even into a cello-oligosaccharide. Depending on the functions, it encompasses enzymes such as C_x cellulase (CMC-ase or endo- β -1,4-glucanase), C_1 cellulase (cellobiohydrolase or exo- β -1,4-glucanase), and β -glucosidase (cellobiose β -1,4-glucosidase), and the like.

Conventionally, studies on cellulase have been directed, for the most part, to the effective utilization of biomass resources. For this reason major supply sources of cellulase have been fungus belonging to genera
 20 such as Trichoderma, Aspergillus, Acremonium, Humichola, and the like.

Recent development of a new application of cellulase as an additive to detergents for washing clothes is requiring supply of alkaline cellulase having a better stability and exhibiting maximum activity at a pH of alkaline side at which cloth washing is performed with a better efficiency.

There have been known several processes for producing alkaline cellulase. Until recently, such
 25 processes have been typified by a process for collecting cellulase A from a culture broth of an alkalophilic microorganism belonging to genus Bacillus (Japanese Patent Laid-open No. 28515/1985), a process for producing alkaline cellulase 301-A through culturing of a microorganism belonging to genus of Cel- lulomonas (Japanese Patent Laid-open No. 224686/1983), a process for obtaining carboxymethyl cellulase through culturing of an alkalophilic strain, Bacillus sp. no. 1139 [Fukumori F., Kudo T., and Horikoshi K.; J.
 30 Gen. Microbiol., 131, 3339 (1985)], a process for producing alkaline cellulase with the use of a microorganism belonging to Streptomyces (Japanese Patent Laid-open No. 19483/1986). Microorganism used in these processes had only a low alkaline cellulase productivity and thus the processes have not been suitable for industrial application.

In view of this situation the present inventors have undertaken extensive studies in order to obtain a
 35 microorganism having a better alkaline cellulase productivity. As a result, inventors have previously discovered several novel microorganisms having superior alkaline cellulase producing capability, including Bacillus sp. KSM-635 (FERM 8872), and filed patent applications on these microorganisms (Japanese Patent Application No. 257775/1986 and others).

Formulating protease into detergents have long been practiced. Earlier practice was to formulate into
 40 detergents protease having an optimum pH close to the neutral region such as papain, pancreas trypsin, or the like. Like cellulase, however, protease having an optimum pH in an alkaline side and stable in surface active agents have been desired for the detergent use. Development of such protease has been undertaken.

Typical alkaline proteases used for detergent purpose are alkalase, savinase, esperase (manufactured
 45 by Novo Industries Co.), maxatase (produced by Gist Brocades Co.), and the like. The optimum reaction temperature for these enzymes is in the neighborhood of 60°C and is in conformity with the washing practice of some region, for example, European countries and U.S.A.

However, in other regions where washing is performed near room temperature, supply of protease are
 50 desired which has a strong activity at a temperature lower than that at which conventional protease are active. Furthermore, because of common knowledge that a protease with a superior capability of decomposing insoluble proteins such as keratin can contribute to detergency [Minagawa M.; Sensho-shi (J. Jpn. Res. Assn. Text. End-Uses, 26, 322 (1985))], the desire has existed for a protease having a superior insoluble protein decomposing capability.

The present inventors have carried out extensive studies in order to obtain an alkaline protease having
 55 keratin-decomposing capability, and a lower optimum temperature, i.e., exhibiting high activity at a low temperature, and discovered a group of microorganisms, including Bacillus sp. KSM-2001 (FERM P-9449), satisfying these requirements. The inventors previously filed a patent application (Japanese Patent Application No. 261487/1987) based on these findings.

Attention has been given also to amylase capable of hydrolyzing starch as a component of detergents, since starch are abundantly contained in the dirt of foods and the like. Examples of known amylases are neutral amylase produced, for example, by Bacillus amyloliquefaciens [N. E. Welker and L. L. Campbell; J. Bacteriol., **94**, 1124, (1967)], heat resistant amylase produced, for example, by Bacillus licheniformis - (Japanese Patent Laid-open No. 37094/1986), alkaline amylase produced, for example, by Bacillus ohbensis (Japanese Patent Publication No. 31949/1977), and the like.

Development of the above-mentioned new microorganisms has made it possible to produce extracellular enzymes for practical application. For the purpose of industrial production, however, a need has continued to exist for the further development of extracellular enzyme-producing microorganisms other than those mentioned above, as well as improvement in productivity of extracellular enzyme by microorganisms.

The production of extracellular α -Amylase in Bacillus subtilis has been studied by Hltotsyanagi, K. et al, (1979) Agric. Biol. Chem. **43**(11) pp.2343-2349. The document discloses mutants resistant to D-cycloserine and ampicillin strains.

In this situation the inventors have undertaken continued studies for the improvement of productivity of extracellular enzymes, especially by means of mutation, and found that a remarkable productivity improvement was attained by providing extracellular enzyme-producing microorganisms belonging to genus Bacillus with resistance to antibiotics cell membrane synthesis inhibitors. This finding has led to the completion of this invention.

SUMMARY OF THE INVENTION

Accordingly, an object of this invention is to provide a cell membrane synthesis inhibitor-resistant mutant of an extracellular enzyme-producing microorganism belonging to genus Bacillus, as defined in claim 1.

Another object of this invention is to provide a process for producing a cell membrane synthesis inhibitor-resistant mutant of an extracellular enzyme-producing microorganism belonging to genus Bacillus as defined in claim 1, comprising:

subjecting an extracellular enzyme-producing microorganism belonging to genus Bacillus to a mutation treatment, and culturing said microorganism in a culture medium containing vancomycin or ristocetin as a cell membrane synthesis inhibitor.

DETAILED DESCRIPTION OF THE INVENTION AND PREFERRED EMBODIMENTS

A cell membrane synthesis inhibitor used in this invention is defined as a substance inhibiting the lipid cycle involved in the cell membrane synthesis in a microorganism. Following antibiotics are given as examples of such substances:

Vancomycin, Ristocetin: These substances inhibit the polymerization reaction of undecaprenol-PP-MurNAc(-GlcNAc)-peptide, a lipid intermediate, to produce a linear glycopeptide [Anderson, J. S. et al, Proc. Natl. Acad. Sci. U.S.A., **53**, 881 (1965); Lugtenberg, E. J. J. et al, J. Bacteriol., **108**, 20 (1971)].

The cell membrane synthesis inhibitor-resistant mutant of an enzyme-producing microorganism belonging to genus of Bacillus of this invention (such mutant being hereinafter referred to as "inhibitor-resistant mutant") can be produced by inducing mutation in an extracellular enzyme-producing microorganism belonging to genus Bacillus (hereinafter referred to as "parent strain") through conventional method such as reaction of a mutagenic agent on the parent strain or irradiation of UV rays or radioactive rays, culturing the mutant thus produced in a culture medium containing cell membrane synthesis inhibitor vancomycin or ristocetin, at a high concentration, and collecting cells which are resistant to the cell membrane synthesis inhibitor.

Any microorganisms belonging to genus Bacillus and possessing extracellular enzyme-producing capability can be used as a parent strain. Preferable parent strains are cellulase-, protease-, or amylase-producing microorganisms, and specifically include, as cellulase-producing parent strains, Bacillus sp. KSM-635 (Japanese Patent Application No. 257775/1986), Bacillus sp. KSM-425 (Japanese Patent Application No. 283742/1986), Bacillus sp. KSM-521 (Japanese Patent Application No. 57644/1987), and Bacillus sp. KSM-580 (Japanese Patent Application No. 68597/1987), all previously discovered by the present inventors in the soils in Haga-gun, Tochigi-ken, Japan; as protease-producing parent strains, Bacillus sp. KSM-2001, Bacillus sp. KSM-2002, Bacillus sp. KSM-2003, and Bacillus sp. KSM-2005, all of which were discovered by the present inventors in the soils in Haga-gun, Tochigi-ken, Japan and filed for patent (Japanese Patent Application No. 261487/1987); and as amylase-producing microorganisms, Bacillus amyloliquefaciens KSM-21 (ATCC 23844), Bacillus amyloliquefaciens KSM-22 (ATCC 23845), and Bacillus licheniformis KSM-23

(ATCC 27811).

Given as examples of chemicals inducing mutation are bases or base-like substances, including 5-bromouracil, bromodeoxyuridine, and the like, acridine, nitrous acid, hydroxylamines, alkylating reagents such as ethylmethane sulfonate (EMS), N-methyl-N'-nitrosoguanidine (NTG), mustard gas, and the like.

Ionizing radiation and ultraviolet radiation are given as examples of radiation.

The term "inhibitor-resistant" as used in this invention means the state of a microorganism capable of growing in the conditions where a cell membrane synthesis inhibitor is present at a concentration of at least the minimum inhibitory concentration (MIC) of the parent strain. Therefore, in order to select inhibitor-resistant strains among mutants, MIC of the parent strain is first measured, a mutant is cultured in a medium containing the cell membrane synthesis inhibitor of a concentration above the MIC, and cells forming colony is collected.

Typical examples of inhibitor-resistant strains of cellulase-producing microorganisms are vancomycin-resistant *Bacillus* sp. KSM-635, and ristocetin-resistant *Bacillus* sp. KSM-635, both having been induced from *Bacillus* sp. KSM-635 as the parent strain. Mycological characteristics of these two antibiotic-resistant strains are now discussed.

(1) Observation under microscope

Cells are gram-positive rods of a size of 0.5 - 1.2 μm x 1.5 - 4.0 μm , with an endospore (0.7 - 1.2 μm x 1.0 - 2.0 μm) forming at one of their ends. They have flagella and are motile.

(2) Growth in various culture media

Cells grow in a pH range of 8 to 11. Growth is most proliferous in a medium to which 0.5 to 1.0% of sodium carbonate (Na_2CO_3) is added. The mycological characteristics of the cells described below are those in the presence of 1.0% by weight of Na_2CO_3 .

(a) Nutrient agar plate

Colony has a circular shape, with its surface being flat. The color of the colony is white - yellow, semi-transparent, and glossy.

(b) Nutrient broth

Cells can grow and the broth become turbid. Cells do not grow in a neutral pH.

(c) Cells can grow in nutrient broth containing 7% NaCl.

(d) Nutrient agar containing gelatin

Cells do not grow.

(e) Litmus milk medium

Milk is not coagulated.

Peptonization: negative

(3) Physiological characteristics

(a) Nitric acid reduced to nitrite: positive

Denitrification: negative

(b) MR test: undeterminable

VP test: positive

(c) Do not produce indole

(d) Production of hydrogen sulfide: negative

(e) Do not hydrolyze or utilize starch

(f) Utilize citric acid

(g) Utilize nitric acid and nitrous acid. Utilize ammonium chloride only slightly, but utilize ammonium phosphate efficiently.

(h) Turn light yellow, but are not fluorescent on King B agar plate

(i) Urease: negative

(j) Oxidase: unclear

(k) Catalase: positive

(l) Growth temperature: 20 - 45 °C

Optimum growth temperature: 29 - 37 °C

(m) Growth pH range: 8 - 11

Optimum pH range: 9.5 - 10.2

(n) Behavior on oxygen: aerobic

(o) Sugar utilization

Utilize D-ribose, L-arabinose, D-xylose, D-glucose, D-mannose, D-fructose, maltose, sucrose, trehalose, D-mannitol, inositol, and glycerol, and do not utilize D-galactose, lactose, D-sorbitol, starch, dextrin, and raffinose.

(p) Hydrolysis of casein and gelatin: negative

(q) Require biotin (or desthiobiotin) for the growth

The above-mentioned characteristics of inhibitor-resistant mutant strains, *Bacillus* sp. KSM-635_v and *Bacillus* sp. KSM-635_r, are the same as those of the parent strain, *Bacillus* sp. KSM-635. However, different from the parent strain these two inhibitor-resistant mutants can grow in the presence of vancomycin or ristocetin, and possess alkaline cellulase-producing capability 2 to 4 times greater than the parent strain.

Based on the above-mentioned similarity of the inhibitor-resistant mutants to the parent strain, the mutants are judged as belonging to the same group as the parent strain, an alkalophilic *Bacillus*, which was defined by Horikoshi and Akiba as the *Bacillus* "growing in an alkaline medium of the pH of at least 8 but not below pH 8 [Alkalophilic Microorganisms, Japan Scientific Society Press (Tokyo), 1982].

Typical examples of inhibitor-resistant strains of protease-producing microorganisms are vancomycin-resistant *Bacillus* sp. KSM-2002_v and ristocetin-resistant *Bacillus* sp. KSM-2002_r, both having been induced from *Bacillus* sp. KSM-2002 as the parent strain. Mycological characteristics of these two inhibitor-resistant strains are as follows:

(1) Observation under microscope

Cells are gram-positive rods of a size of 0.5 - 0.8 μm x 1.5 - 3.0 μm , with a circular or oval endospore (0.4 - 0.8 μm x 0.4 - 0.8 μm) forming at central-subterminal of the cell. They have peripheral flagella and are motile.

(2) Growth in various culture media

(a-i) Nutrient agar plate

Cells form circular, convex, or smooth wave-like colonies with a diameter of 1.0 to 5.0 mm. The colony has a dewdrop shape with a smooth, fatty, semi-transparent surface colored pale yellow - milky white.

(a-ii) Nutrient agar slant

Grow only weakly forming a band in a broth at pH 7.0. Colony is milky white and semi-transparent. The band-like growth expands at pH 10.0.

(b) Nutrient broth

Cells can slightly grow at pH 7.0 without forming cell membrane.

(c) Cells can grow in the presence of a 10% sodium chloride.

(d) Stab culturing in nutrient agar

Liquefy gelatin at pH 7.0.

(e) Litmus milk medium

Slightly grow. Milk coagulation and litmus discoloration are not observed.

(3) Physiological characteristics

(a) Nitric acid reduced to nitrite: negative Denitrification reaction: negative

(b) MR test: negative VP test: negative

(c) Do not produce indole

(d) Production of hydrogen sulfide: negative

(e) Hydrolysis of starch: negative

(f) Utilization of citric acid: negative (in Christensen-Simons medium)

(g) Utilization of nitric acid and ammonium sulfate: negative

(h) Do not produce pigment.

(i) Urease: negative

(j) Oxidase: positive

(k) Catalase: positive

(l) Growth temperature: 10 - 37°C, growth is good at 20 - 30°C.

(m) Growth pH range: 6.0 - 11.0

(n) Grow in either aerobic or anaerobic conditions.

(o) Sugar utilization

Utilize L-arabinose, D-xylose, D-glucose, D-fructose, D-galactose, maltose, lactose, D-sorbitol, D-mannitol, and starch, and not utilize D-mannose, sucrose, trehalose, inositol, and glycerol.

(p) Production of acids and gases from sugars

Both inhibitor-resistant strains do not produce acids and gases from following sugars.

L-arabinose, D-xylose, D-glucose, D-mannose, D-fructose, D-galactose, maltose, sucrose, lactose, trehalose, D-sorbitol, D-mannitol, inositol, glycerol, and starch.

The above-mentioned characteristics of inhibitor-resistant mutant strains, *Bacillus* sp. KSM-2002, and *Bacillus* sp. KSM-2002, are the same as those of the parent strain, *Bacillus* sp. KSM-2002. However, different from the parent strain these two inhibitor-resistant mutants can grow in the presence of vancomycin or ristocetin at a concentration above MIC, and possess alkaline protease-producing capability 2 to 3 times greater than the parent strain.

Further, given as typical examples of inhibitor-resistant strains of amylase-producing microorganisms are vancomycin-resistant *Bacillus amyloliquefaciens* KSM-22, and ristocetin-resistant *Bacillus amyloliquefaciens* KSM-22, both having been induced from a known microorganism, *Bacillus amyloliquefaciens* KSM-22 [ATCC 23845; N. E. Welker and L. L. Campbell, *J. Bacteriol.*, 94, 1124 (1967)], as the parent strain. These inhibitor-resistant mutant strains have the same mycological characteristics as the parent strain, *Bacillus amyloliquefaciens* KSM-22. However, different from the parent strain the inhibitor-resistant mutants can grow in the presence of vancomycin or ristocetin, and possess amylase-producing capability 1.3 to 4 times greater than the parent strain.

Culturing of the inhibitor-resistant mutant of this invention may be carried out by inoculating the cells into a normal liquid medium and culturing the cells according to a conventional aerobic culture method. Inclusion of a suitable amount of carbon and nitrogen sources which the microorganism can utilize in the medium is desirable. There are no specific limitations as to the carbon and nitrogen sources. Given as examples of carbon sources are fibrous substances such as chaff, hull, filter paper, general paper, sawdust; molasses, invert sugar, CMC, Avicel, cellulose powder, xylan, pectin, soluble starch, and the like. In addition, other utilizable carbon sources are included, such as ribose, arabinose, xylose, glucose, mannose, fructose, galactose, lactose, maltose, sucrose, trehalose, sorbitol, mannitol, inositol, glycerol, as well as organic acid such as citric acid, acetic acid, and the like. Enumerated as nitrogen sources are inorganic nitrogen-containing compounds such as ammonium nitrate, ammonium sulfate, ammonium chloride, ammonium phosphate, sodium nitrate, and the like; corn gluten meal, soybean flour, soybean meal, corn steep liquor, casamino acid, yeast extract, pharma media, sardine meal, meat extract, peptone, hypro, ajipower, corn-soybean meal, coffee meal, cotton seed meal, cultivator, amiflex, ajipron, zest, ajix, and the like. In addition to these carbon and nitrogen sources, phosphoric acid, salts of metal such as Mg^{+2} , Ca^{+2} , Mn^{+2} , Zn^{+2} , Co^{+2} , Na^{+} , K^{+} , and the like, as required, other micro-nutritious organic or inorganic substances can be added into the culture medium.

Enzymes such as cellulase, protease, amylase produced by the inhibitor-resistant mutant outside the cells can be collected and purified by a conventional method. The culture liquid containing such an enzyme can be used as is as an enzyme source. Such an enzyme liquid may be obtained by separating cells from the culture broth by means of centrifugation, filtration, or the like, and by precipitating proteins by a conventional method, including salting out, isoelectric precipitation, precipitation from solvent, or the like, from the cells or the culture liquid filtrate, or by condensing said filtrate by means of ultrafiltration. Desalting can be performed by a conventional method, including dialysis, gel filtration, or the like. Although it is possible to use the enzyme liquid thus obtained as is, they can be served for use after purifying by means of a suitable combination of hydroxyapatite chromatography, ion-exchange chromatography such as DEAE-Sephadex or DEAE-cellulose, molecular sieving gel chromatography, and the like.

Since the inhibitor-resistant mutants of this invention have extracellular enzyme-producing capability of 2 to 4 times of the parent strains, they can be advantageously used for the industrial production of cellulase, protease, amylase, and the like.

Other features of the invention will become apparent in the course of the following description of the exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

EXAMPLES

In the examples below enzymatic activities were measured according to the following method.

(1) CMC-ase activity

To the solution comprised of 0.2 ml of 25% CMC, 0.1 ml of 0.5 M glycine buffer (pH 9.0), and 0.1 ml of deionized water was added 0.1 ml of enzyme solution diluted to an appropriate concentration, and the mixture was reacted for 20 minutes at 40 °C. After the reaction, reducing sugar was quantitatively analyzed by 3,5-dinitrosalicylic acid (DNS) method. Specifically, in order to determine the amount of produced reducing sugar 1 ml of DNS reagent was added to 0.5 ml of reaction liquid, and mixture was caused to color by heating at 100 °C for 5 minutes. After cooling, the colored mixture was diluted with 4.5 ml of deionized water and subjected to colorimetric quantitative analysis at 535 nm.

One (1) unit of enzyme titer is defined as the amount of the enzyme producing reducing sugar corresponding to 1 μmol of glucose per minute. Fermentation productivity is expressed by the enzyme unit per 1 liter of the culture liquid.

(2) Protease activity

Following the method proposed by Anson [Anson, M. L., *J. Ger. Physiol.*, **22**, 79 (1938)] bovine serum hemoglobin was denatured by urea, and adjusted to pH 10.5 with sodium hydroxide. Into 0.5 ml of this substrate solution (2.2% as hemoglobin) 0.1 ml [1.0×10^{-5} - 1.0×10^{-3} $\mu\text{A.U}$ (A.U.: Anson Unit)] of an enzyme solution was added, and mixture was reacted at 25 °C for 10 minutes, upon which 1.0 ml of 4.9% trichloroacetic acid was added to terminate the reaction. After termination of the reaction, the reaction mixture was centrifuged at 3,000 rpm for 10 minutes. Proteolysates contained in the supernatant was quantitatively analyzed by Folin-Lowry method [O. H. Lowry et al; *J. Biol. Chem.*, **193**, 265 (1951)]. 1 A.U. is defined as the amount of enzyme releasing 1 mmol of tyrosine in 1 minutes under the above reaction conditions.

(3) Amylase activity

To the solution comprised of 0.4 ml of 1.25% soluble starch, 0.025 ml of 1 M Tris-HCl buffer (pH 7.0), 0.05 ml of 0.2 M calcium chloride, and 0.425 ml of deionized water was added 0.1 ml of an enzyme solution diluted to an appropriate concentration, and the mixture was reacted for 5 minutes at 30 °C. After the reaction, reducing sugar was quantitatively analyzed by 3,5-dinitrosalicylic acid (DNS) method. Specifically, in order to determine the amount of produced reducing sugar 1 ml of DNS reagent was added to 1 ml of the reaction liquid, and the mixture was caused to color by heating at 100 °C for 5 minutes. After cooling, the colored mixture was diluted with 4 ml of deionized water and subjected to colorimetric quantitative analysis at 535 nm.

One (1) unit of enzyme titer is defined as the amount of the enzyme producing reducing sugar corresponding to 1 μmol of maltose per minute.

Reference Example 1

Bacillus sp. KSM-635 (FERM P-8872) was spread on the medium A described below, each medium containing vancomycin at a concentration shown in Table 1 below, to determine minimum inhibitory concentration (MIC). The media A containing vancomycin to which *Bacillus* sp. KSM-635 (FERM P-8872) was spread was left for 3 days at 30 °C, and the cell growth conditions on the medium surface was observed by naked eyes. The results are shown in Table 1.

Medium A:	
Sucrose	2%
Mixed amino acid solution	6%
KH ₂ PO ₄	0.1%
Tris-HCl buffer	0.1 M (Final conc. pH 8)
Bacto agar-agar	2.0%
Vancomycin	Concentration in Table 1

TABLE 1

Concentration of vancomycin	Cell growth on medium surface
0 (Control)	+++
0.1 γ	++
0.2 γ	±
0.3 γ	-
0.5 γ	-
1.0 γ	-
1.5 γ	-

As evident from the table MIC of vancomycin for Bacillus sp. KSM-635 was about 0.2 γ .

Reference Example 2

Bacillus sp. KSM-635 (FERM P-8872) was spread on the medium B described below, each medium containing ristocetin at a prescribed concentration, to determine minimum inhibitory concentration (MIC) of the antibiotic. As a result, the MIC of ristocetin for Bacillus sp. KSM-635 (FERM P-8872) was found to be about 0.1 γ .

Medium B:	
Sucrose	2%
Mixed amino acid solution	6%
KH ₂ PO ₄	0.1%
Tris-HCl buffer	0.1 M (Final conc. pH 8)
Bacto agar-agar	2.0%
Ristocetin	Suitable concentration

Example 1

Bacillus sp. KSM-635 (FERM P-8872) was inoculated into the liquid medium C and cultured at 30°C for 20 hours. EMS was added to the culture broth to a concentration of 0.5 to 3% (normally 1%) and the broth was left for 20 minutes at 30°C. Then, a 3% treated cultured liquid was inoculated into the liquid medium and the culturing was continued at 30°C overnight.

The culture liquid was spread on medium A containing not less than 0.2 γ of vancomycin and cultured at 30°C for 2 days. Colonies grown on the surface of the medium were collected. The above culturing procedure was repeated 5 times to obtain 18 vancomycin-resistant strains.

Each of these vancomycin-resistant strains was inoculated into 100 ml of liquid medium D and cultured at 30°C for 3 days. The alkaline cellulase activity was determined on each broth. As a result, the production of vancomycin-resistant strains with a high alkaline cellulase activity as shown in Table 2 was confirmed. The strain having the highest titer (No. 3215) was named Bacillus sp. KSM-635_v. The strain was deposited with Fermentation Research Institute, Agency of Industrial Science and Technology (FERM P-9084).

Medium C:	
Meat extract	1%
Yeast extract	0.5%
NaCl	0.5%
Glucose	2.0%
KH ₂ PO ₄	1.0%
Na ₂ CO ₃	0.75%

Medium D:	
Sugar	2%
Mixed amino acid solution	6%
Yeast extract	0.2%
Na ₂ CO ₃	0.5%

TABLE 2

Mutant No.	Vancomycin Resistance Acquired	Alkaline Cellulase Production (unit/l)
3215	1.0 γ	16,300
3422	1.0 γ	14,200
3424	0.5 γ	9,500
3681	0.5 γ	11,230
3691	0.5 γ	14,100
3693	0.5 γ	10,990
<u>Bacillus</u> sp. KSM-635 (Parent strain: Control)	0.2 γ	4,280

Example 2

Bacillus sp. KSM-635 (FERM P-8872) was inoculated into the liquid medium C and cultured at 30°C for 20 hours. EMS was added to the culture broth to a concentration of 0.5 to 3% (normally 1%) and the broth was left for 20 minutes at 30°C. Then, a 3% treated cultured liquid was inoculated into the liquid medium and the culturing was continued under shaking at 30°C overnight.

The culture liquid was spread on medium B containing not less than 0.1 γ of ristocetin and cultured at 30°C for 2 days. Colonies grown on the surface of the medium were collected. The above culturing procedure was repeated 5 times to obtain 14 ristocetin-resistant strains.

Each of these ristocetin-resistant strains was inoculated into 100 ml of liquid medium D and cultured at 30°C for 3 days. The alkaline cellulase titer was determined on each broth. As a result, the production of ristocetin-resistant strains with a high alkaline cellulase titer as shown in Table 3 was confirmed. The strain having the highest titer (No. 4021) was named Bacillus sp. KSM-635. The strain was deposited with Fermentation Research Institute, Agency of Industrial Science and Technology (FERM P-9085).

TABLE 2

TABLE 2		
Mutant No.	Ristocetin Resistance	Alkaline Cellulase Production (unit/l)
4021	1.5 γ	20,050
4033	1.5 γ	15,600
4091	1.0 γ	18,200
4180	0.5 γ	16,800
<u>Bacillus</u> sp. KSM-635 (Parent strain: Control)	0.1 γ	4,590

Example 3

Bacillus sp. KSM-635, (FERM P-9084) was cultured in a medium (30 ml in a 500 ml Sakaguchi flask), containing 2% of sucrose, 12% of amino acid mixture, 0.05% yeast extract, 0.1% of KH_2PO_4 , and 0.5% of Na_2CO_3 , under shaking at 30 C for 3 days. The amount of alkaline cellulase produced was measured to obtain the results of 21,520 unit/liter.

Example 4

The amount of alkaline cellulase production by Bacillus sp. KSM-635, (FERM P-9085) was measured following the manner as described in Example 3. The results obtained was 20,290 unit/liter.

Reference Example 3

Bacillus sp. KSM-2002 (FERM P-9450) was spread on the media E described below, each medium containing vancomycin or ristocetin at a concentration shown in Table 4 below, to determine minimum inhibitory concentration (MIC) of vancomycin or ristocetin. The medium E containing vancomycin or ristocetin to which Bacillus sp. KSM-2002 was spread and was left for 3 days at 30 °C, and then the cell growth conditions on the medium surface was observed by naked eyes. The results are shown in Table 4.

Medium E:	
Bacto tryptone	1.5%
Bacto soytone	0.5%
NaCl	0.5%
Bacto agar-agar	2.0%
Vancomycin or Ristocetin	Concentration in Table 4
(pH 7.0)	

TABLE 4

Concentration of vancomycin	Cell growth on medium surface	Concentration of ristocetin	Cell growth on medium surface
0 (Control)	+++	0 (Control)	+++
0.008 γ	++	0.008 γ	++
0.01 γ	+	0.01 γ	+
0.02 γ	$\pm$	0.02 γ	+
0.03 γ	-	0.03 γ	$\pm$
0.04 γ	-	0.04 γ	-
0.05 γ	-	0.05 γ	-
0.1 γ	-	0.1 γ	-

As evident from the table, vancomycin and ristocetin MIC for *Bacillus* sp. KSM-2002 were about 0.02 γ and 0.03 γ , respectively.

Example 5

Bacillus sp. KSM-2002 (FERM P-9450) was inoculated into the liquid medium F and cultured at 30°C for 18 hours. NTG was added to the culture broth to the concentration of 30 to 100 γ , and the broth was left for 30 - 60 minutes at 20°C. Then, a 1% treated culture liquid was inoculated into the liquid medium and the culturing was continued at 30°C overnight.

The culture liquid was spread on medium E containing not less than 0.02 γ of vancomycin or not less than 0.03 γ of ristocetin and cultured at 30°C for 3 days. Colonies grown on the surface of the medium were collected. The above culturing procedure was repeated 5 times to obtain 7 vancomycin-resistant strains and 4 ristocetin-resistant strains.

Each of these strains was inoculated into 5 ml of liquid medium G in a test tube and cultured at 30°C for 3 days. The alkaline protease titer was determined on each broth. As a result, high titer strains as shown in Table 5 were obtained. Among these, a vancomycin-resistant strain having the highest titer (No. 2016) and a ristocetin-resistant strain having the highest titer (No. 2014) were named *Bacillus* sp. KSM-2002, and *Bacillus* sp. KSM-2002. The strains were deposited with Fermentation Research Institute, Agency of Industrial Science and Technology as FERM P-9939 and FERM P-9938, respectively.

Medium F:	
Bacto tryptone	1.5%
Bacto sottonne	0.5%
NaCl	0.5%
(pH 7.0)	

Medium G:	
Glucose	0.2%
Wool keratin	0.5%
Yeast extract	0.05%
KH ₂ PO ₄	0.1%
MgSO ₄ ·7H ₂ O	0.02%
Na ₂ CO ₃	0.4%
(pH 9.0)	

TABLE 5

Mutant No.	Vancomycin Resistance	Alkaline protease Production	Mutant No.	Ristocetin Resistance	Alkaline protease Production
2003	0.05 γ	2.23 A.U./l	2012	0.05 γ	1.45 A.U./l
2016	0.05 γ	2.78 A.U./l	2013	0.05 γ	1.86 A.U./l
2006	0.06 γ	1.43 A.U./l	2014	0.05 γ	2.53 A.U./l
<i>Bacillus</i> sp. KSM-2002 (Parent strain: Control)	0.02 γ	0.71 A.U./l	<i>Bacillus</i> sp. KSM-2002 (Parent strain: Control)	0.03 γ	0.71 A.U./l

15 Reference Example 5

Bacillus amyloliquefaciens KSM (ATCC 23845) was spread on the medium E, each medium containing vancomycin or ristocetin at a prescribed concentration, to determine minimum inhibitory concentration (MIC) of vancomycin or ristocetin in the same manner as in Reference Example 3. The results are shown in Table

20 6.

TABLE 6

Concentration of vancomycin	Cell growth on medium surface	Concentration of ristocetin	Cell growth on medium surface
0 (Control)	+++	0 (Control)	+++
0.09 γ	++	0.7 γ	++
0.1 γ	++	0.8 γ	+
0.2 γ	+	0.9 γ	$\pm$
0.3 γ	$\pm$	1.0 γ	-
0.4 γ	-	1.3 γ	-
0.5 γ	-	1.5 γ	-
1.0 γ	-		

35 Example 6

Bacillus amyloliquefaciens KSM-22 (ATCC 23845) was inoculated into the liquid medium F and cultured at 30 °C for 16 hours. NTG was added to the culture broth to a concentration of 30 to 100 γ and the broth was left for 30 -60 minutes at 20 °C. Then, a 1% treated culture liquid was inoculated into the liquid medium and the culturing was continued at 30 °C overnight.

The culture liquid was spread on medium E containing not less than 0.3 γ of vancomycin or 0.9 γ of ristocetin and cultured at 30 C for 3 days. Colonies grown on the surface of the medium were collected. The above culturing procedure was repeated 5 times to obtain 5 vancomycin-resistant strains and 4 ristocetin-resistant strains.

Each of these strains was inoculated into 5 ml of liquid medium F and cultured at 30 °C for 2 days. The amylase-titer was determined on each broth. As a result, high titer strains as shown in Table 7 were obtained. Among these, a vancomycin-resistant strain having the highest titer (No. 4591) and a ristocetin-resistant strain having the highest titer (No. 4551) were named *Bacillus amyloliquefaciens* KSM-22, and *Bacillus amyloliquefaciens* KSM-22. The strains were deposited with Fermentation Research Institute, Agency of Industrial Science and Technology as FERM P-9937 and FERM P-9936, respectively.

TABLE 7

Mutant No.	Vancomycin Resistance	Amylase Production	Mutant No.	Ristocetin Resistance	Amylase Production
4504	0.5 γ	20900 unit/l	4551	2.0 γ	49400 unit/l
4591	0.5 γ	46900 unit/l	4554	2.0 γ	13050 unit/l
<u>Bacillus</u> amyloliquefaciens KSM-22 (ATCC 23845) (Parent strain: Control)	0.3 γ	10700 unit/l	<u>Bacillus</u> amyloliquefaciens KSM-22 (ATCC 23845) (Parent strain: Control)	0.9 γ	10700 unit/l

Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings.

Particulars with respect to microorganisms

Strain identification reference	Accession number by FERM	Converted from	Date of original deposit
KSM-635v	FERM BP 2035	FERM P-9084	December 15, 1986
KSM-635r	FERM BP-2036	FERM P-9085	December 15, 1986
KSM-2002	FERM BP-2037	FERM P-9450	July 2, 1987
KSM-22r	FERM BP-2038	FERM P-9936	March 15, 1988
KSM-22v	FERM BP-2039	FERM P-9937	March 15, 1988
KSM-2002v	FERM BP-2041	FERM P-9939	March 15, 1988
KSM-2002r	FERM BP-2040	FERM P-9938	March 15, 1988

Claims

55 Claims for the following Contracting States : CH, DE, FR, GB, IT, LI, NL, SE

1. A cell membrane synthesis inhibitor-resistant mutant of an extra-cellular enzyme-producing microorganism belonging to the genus Bacillus, said mutant being resistant to vancomycin or ristocetin as a



Particulars with respect to microorganisms

Strain identification reference	Accession number by FERM	Converted from	Date of original deposit
KSM-635v	FERM BP-2035	FERM P-9084	December 15, 1986
KSM-635r	FERM BP-2036	FERM P-9085	December 15, 1986
KSM-2002	FERM BP-2037	FERM P-9450	July 2, 1987
KSM-22r	FERM BP-2038	FERM P-9936	March 15, 1988
KSM-22v	FERM BP-2039	FERM P-9937	March 15, 1988
KSM-2002v	FERM BP-2041	FERM P-9939	March 15, 1988
KSM-2002r	FERM BP-2040	FERM P-9938	March 15, 1988

Claims

Claims for the following Contracting States : CH, DE, FR, GB, IT, LI, NL, SE

1. A cell membrane synthesis inhibitor-resistant mutant of an extra-cellular enzyme-producing microorganism belonging to the genus Bacillus, said mutant being resistant to vancomycin or ristocetin as a



substance inhibiting the lipid cycle involved in the cell membrane synthesis in a microorganism.

2. A process for producing a cell membrane synthesis inhibitor-resistant mutant as defined in Claim 1 comprising: subjecting an extracellular enzyme-producing microorganism belonging to genus Bacillus to a mutation treatment, and culturing said microorganism in a culture medium containing vancomycin or ristocetin as a cell membrane synthesis inhibitor.

Claim for the following Contracting State : ES

1. A process for producing a cell membrane synthesis inhibitor-resistant mutant of an extra-cellular enzyme-producing microorganism belonging to the genus Bacillus, said mutant being resistant to vancomycin or ristocetin as a substance inhibiting the lipid cycle involved in the cell membrane synthesis in a microorganism, said process being characterized by subjecting an extracellular enzyme-producing microorganism belonging to genus Bacillus to a mutation treatment, and culturing said microorganism in a culture medium containing vancomycin or ristocetin as a cell membrane synthesis inhibitor, and collecting cells which are resistant to the cell membrane synthesis inhibitor.

Patentansprüche

Patentansprüche für folgenden Vertragsstaaten : CH, DE, FR, GB, IT, LI, NL, SE

1. Eine gegen den Zellmembransynthese-Inhibitor resistente Mutante eines ein extrazelluläres Enzym erzeugenden Mikroorganismus der Gattung Bacillus, wobei die Mutante resistent ist gegenüber Vancomycin oder Ristocetin als eine Substanz, welche den Lipidzyklus inhibiert, der bei der Zellmembransynthese in einem Mikroorganismus involviert ist.
2. Verfahren zur Erzeugung einer gegen den Zellmembransynthese-Inhibitor resistenten Mutante, wie in Anspruch 1 definiert, umfassend: Durchführung einer Mutationsbehandlung bei einem extrazelluläres Enzym erzeugenden Mikroorganismus der Gattung Bacillus und Kultivierung des Mikroorganismus in einem Kulturmedium, welches Vancomycin oder Ristocetin als einen Zellmembransynthese-Inhibitor enthält.

Patentanspruch für folgenden Vertragsstaat : ES

1. Verfahren zur Erzeugung einer gegen den Zellmembransynthese-Inhibitor resistenten Mutante, wobei die Mutante resistent ist gegenüber Vancomycin oder Ristocetin als eine Substanz, welche den Lipidzyklus inhibiert, der bei der Zellmembransynthese in einem Mikroorganismus involviert ist, wobei das Verfahren gekennzeichnet ist durch Durchführung einer Mutationsbehandlung bei einem extrazelluläres Enzym erzeugenden Mikroorganismus der Gattung Bacillus und Kultivierung des Mikroorganismus in einem Kulturmedium, welches Vancomycin oder Ristocetin als einen Zellmembransynthese-Inhibitor enthält, und Sammlung von Zellen, welche gegenüber dem Zellmembransynthese-Inhibitor resistent sind.

Revendications

Revendications pour les Etats contractants suivants : CH, DE, FR, GB, IT, LI, NL, SE

1. Mutant résistant à un inhibiteur de synthèse de la membrane cellulaire issu d'un micro-organisme producteur d'enzyme extracellulaire et appartenant au genre Bacillus, ledit mutant étant résistant à la vancomycine ou à la ristocétine à titre de substance inhibant le cycle lipidique impliqué dans la synthèse de la membrane cellulaire chez un micro-organisme.
2. Procédé de production d'un mutant résistant à un inhibiteur de synthèse de la membrane cellulaire tel que défini dans la revendication 1 comprenant : les étapes consistant à soumettre un micro-organisme producteur d'enzyme extracellulaire et appartenant au genre Bacillus à un traitement de mutation et la culture dudit micro-organisme dans un milieu de culture contenant de la vancomycine ou de la ristocétine à titre d'inhibiteur de synthèse de la membrane cellulaire.



Revendication pour l'Etat contractant suivant : ES

1. Procédé de production d'un mutant résistant à un inhibiteur de synthèse de la membrane cellulaire issu d'un micro-organisme producteur d'enzyme extracellulaire et appartenant au genre Bacillus, ledit mutant étant résistant à la vancomycine ou à la ristocétine à titre de substance capable d'inhiber le cycle lipidique impliqué dans la synthèse de la membrane cellulaire chez un micro-organisme, ledit procédé étant caractérisé en ce que l'on soumet un micro-organisme producteur d'enzyme extracellulaire et appartenant au genre Bacillus à un traitement de mutation, cultive ledit micro-organisme dans un milieu de culture contenant de la vancomycine ou de la ristocétine à titre de substance capable d'inhiber la synthèse de la membrane cellulaire, et recueille les cellules résistantes à l'inhibiteur de synthèse de la membrane cellulaire.