

PROCESS FOR THE PREPARATION OF BENZO-FUSED HETEROARYL SULFAMATES AND
CRYSTALLINE FORM OF
N-((2S)-6-CHLORO-2,3-DIHYDRO-1,4-BENZODIOXIN-2-YL) METHYL-SULFAMIDE

FIELD OF THE INVENTION

5 The present invention is directed to a process for the preparation of benzo-fused heteroaryl sulfamates, useful for the treatment of epilepsy and related disorders.

BACKGROUND OF THE INVENTION

10 Epilepsy describes a condition in which a person has recurrent seizures due to a chronic, underlying process. Epilepsy refers to a clinical phenomenon rather than a single disease entity, since there are many forms and causes of epilepsy. Using a definition of epilepsy as two or more unprovoked seizures, the incidence of epilepsy is estimated at approximately 0.3 to 0.5 percent in
15 different populations throughout the world, with the prevalence of epilepsy estimated at 5 to 10 people per 1000.

 An essential step in the evaluation and management of a patient with a seizure is to determine the type of seizure that has occurred. The main characteristic that distinguishes the different categories of seizures is whether
20 the seizure activity is partial (synonymous with focal) or generalized.

 Partial seizures are those in which the seizure activity is restricted to discrete areas of the cerebral cortex. If consciousness is fully preserved during the seizure, the clinical manifestations are considered relatively simple and the seizure is termed a simple-partial seizure. If consciousness is impaired, the
25 seizure is termed a complex-partial seizure. An important additional subgroup comprises those seizures that begin as partial seizures and then spread diffusely throughout the cortex, which are known as partial seizures with secondary generalization.

 Generalized seizures involve diffuse regions of the brain simultaneously
30 in a bilaterally symmetric fashion. Absence or petit mal seizures are characterized by sudden, brief lapses of consciousness without loss of postural control. Atypical absence seizures typically include a longer duration in the lapse of consciousness, less abrupt onset and cessation, and more obvious


motor signs that may include focal or lateralizing features. Generalized Tonic-clonic or grand mal seizures, the main type of generalized seizures, are characterized by abrupt onset, without warning. The initial phase of the seizure is usually tonic contraction of muscles, impaired respiration, a marked
5 enhancement of sympathetic tone leading to increased heart rate, blood pressure, and pupillary size. After 10-20 s, the tonic phase of the seizure typically evolves into the clonic phase, produced by the superimposition of periods of muscle relaxation on the tonic muscle contraction. The periods of relaxation progressively increase until the end of the ictal phase, which usually
10 lasts no more than 1 min. The postictal phase is characterized by unresponsiveness, muscular flaccidity, and excessive salivation that can cause stridorous breathing and partial airway obstruction. Atonic seizures are characterized by sudden loss of postural muscle tone lasting 1-2 s. Consciousness is briefly impaired, but there is usually no postictal confusion.
15 Myoclonic seizures are characterized by a sudden and brief muscle contraction that may involve one part of the body or the entire body. (Harrison's Online, March 29, 2001)

McComsey, D., et al. in US Patent Publication US 2006/0041008 A1,
20 published February 23, 2006 and McComsey, D., et al. in US Patent Publication US 2005/0282887 A1, published December 22, 2005 disclose compounds of formula (I) and their use in the treatment of epilepsy and related disorders. McComsey, D., et al. in US Patent Publication US 2006/0041008 A1 and McComsey, D., et al. in US Patent Publication US 2005/0282887 A1
25 further disclose a process for the preparation of the compounds of formula (I) comprising reacting a suitable substituted amine with sulfamide.

In the process(es) as disclosed in McComsey D., et al. compounds of formula (I) wherein R^1 and R^2 are each hydrogen describes the use of sulfamoyl chloride ($Cl-SO_2-NH_2$) as a reagent, which reagent is unsuitable for
30 large scale / commercial preparation. There remains, however, a need for a process suitable for the preparation of large scale material and / or for commercial preparation of the compounds of formula (I).

SUMMARY OF THE INVENTION

The present invention is directed to processes for the preparation of compounds of formula (I)



5

wherein


R^1 is selected from the group consisting of hydrogen and lower alkyl;

R^4 is selected from the group consisting of hydrogen and lower alkyl;

a is an integer from 1 to 2;




is selected from the group consisting of




15

2;

each R^5 is independently selected from the group consisting of halogen, lower alkyl and nitro;



- 5 The present invention is directed to a process for the preparation of compounds of formula (IA)



wherein


R¹ is hydrogen;


- 10 R⁴ is selected from the group consisting of hydrogen and lower alkyl;

a is an integer from 1 to 2;




is selected from the group consisting of






wherein b is an integer from 0 to 4; and wherein c is an integer from 0 to 2;

- 5 each R^5 is independently selected from the group consisting of halogen, lower alkyl and nitro;




or a pharmaceutically acceptable salt thereof;

- 10 comprising




- reacting a compound of formula (X) with a compound of formula (XI)
 wherein $-\text{C}(\text{O})\text{OR}^0$ is a nitrogen protecting group; in the presence of an organic
 or inorganic base, wherein the organic or inorganic base is not reactive with the
 15 chloro group on the compound of formula (XI); in an aprotic organic solvent; to
 yield the corresponding compound of formula (XII);



de-protecting the compound of formula (XII); to yield the corresponding compound of formula (IA).

- 5 The present invention is further directed to a process for the preparation of a compound of formula (IB)



wherein





R¹ is selected from the group consisting of lower alkyl;


10 R⁴ is selected from the group consisting of hydrogen and lower alkyl;

a is an integer from 1 to 2;




is selected from the group consisting of






wherein b is an integer from 0 to 4; and wherein c is an integer from 0 to 2;

- 5 each R^5 is independently selected from the group consisting of halogen, lower alkyl and nitro;




or a pharmaceutically acceptable salt thereof;


- 10 comprising



- reacting a compound of formula (X) with a compound of formula (XI)
- wherein $-C(O)OR^0$ is a nitrogen protecting group; in the presence of an organic or inorganic base, wherein the organic or inorganic base is not reactive with the
- 15 chloro group on the compound of formula (XI); in an aprotic organic solvent; to yield the corresponding compound of formula (XII);



reacting the compound of formula (XII) with a compound of formula (XV), wherein Q is a leaving group; in an organic solvent; to yield the corresponding compound of formula (XVI)




5


de-protecting the compound of formula (XVI); to yield the corresponding compound of formula (IB).

In an embodiment, the present invention is directed to a process for the preparation of a compound of formula (I-S)

10




or a pharmaceutically acceptable salt thereof (also known as *N*-[[(2*S*)-6-chloro-2,3-dihydro-1,4-benzodioxin-2-yl]methyl]-sulfamide) comprising



15

reacting a compound of formula (X-S) with a compound of formula (XI), wherein —C(O)OR^0 is a nitrogen protecting group; in the presence of an organic or inorganic base, wherein the organic or inorganic base is not reactive with the


chloro group on the compound of formula (XI); in an aprotic organic solvent; to yield the corresponding compound of formula (XII-S);



- de-protecting the compound of formula (XII-S); to yield the
5 corresponding compound of formula (I-S).

The present invention is further directed to a crystalline form of the compound of formula (I-S).

- 10 The present invention is further directed to compounds of formula (XII)



- wherein $-C(O)OR^0$ is a nitrogen protecting group and wherein ,
a, R^4 and R^1 are as herein defined. Preferably, $-C(O)OR^0$ is a nitrogen
protecting group wherein R^0 is selected from the group consisting of C_{1-4} alkyl
15 (preferably t-butyl), benzyl, p-methoxybenzyl and 9-fluorenylmethyl. The
compounds of formula (XII) are useful as intermediates in the synthesis of the
compounds of formula (I).

- The present invention is further directed to a product prepared according
20 to the process described herein.

Illustrative of the invention is a pharmaceutical composition comprising a
pharmaceutically acceptable carrier and the product prepared according to the
process described herein. An illustration of the invention is a pharmaceutical
composition made by mixing the product prepared according to the process

described herein and a pharmaceutically acceptable carrier. Illustrating the invention is a process for making a pharmaceutical composition comprising mixing the product prepared according to the process described herein and a pharmaceutically acceptable carrier.


- 5 Exemplifying the invention are methods of treating epilepsy or a related disorder comprising administering to a subject in need thereof, a therapeutically effective amount of any of the compounds or pharmaceutical compositions described above.

10 BRIEF DESCRIPTION OF THE FIGURES

Figure 1 illustrates a representative XRD Spectra for Crystalline Form of the Compound of Formula (I-S)

DETAILED DESCRIPTION OF THE INVENTION


- 15 The present invention is directed to processes for the preparation of compound of formula (I)



wherein R^1 , R^4 , a and R are as herein defined. The compounds of formula (I) are useful in the treatment of epilepsy and related disorders.


20

In an embodiment, the present invention is directed to a process for the synthesis of compounds of formula (IA)



- 25 wherein R^1 is hydrogen, and wherein R^4 , a and R are as herein defined; and pharmaceutically acceptable salts thereof.


In another embodiment, the present invention is directed to a process for the preparation of compounds of formula (IB)



wherein R^1 is selected from the group consisting of lower alkyl, and

- 5 wherein R^4 , a and  are as herein defined; and pharmaceutically acceptable salts thereof.

In another embodiment, the present invention is directed to a process for the synthesis of compounds of formula (IC)



- 10 and pharmaceutically acceptable salts thereof, wherein b and R^5 are as herein defined. Preferably, b is an integer from 0 to 2; more preferably, b is an integer from 0 to 1. Preferably R^5 is halogen, more preferably chloro.

- In an embodiment of the present invention R^1 is selected from the group consisting of hydrogen and methyl. In another embodiment of the present
15 invention, R^1 is hydrogen.

In an embodiment of the present invention $-(\text{CH}_2)_a-$ is selected from the group consisting of $-\text{CH}_2-$ and $-\text{CH}_2\text{CH}_2-$. In another embodiment of the present invention $-(\text{CH}_2)_a-$ is $-\text{CH}_2-$.

- In an embodiment of the present R^4 is selected from the group
20 consisting of hydrogen and methyl, preferably, R^4 is hydrogen.

In an embodiment of the present invention a is 1.


- In an embodiment of the present invention b is an integer from 0 to 2. In another embodiment of the present invention c is an integer from 0 to 2. In another embodiment of the present invention b is an integer from 0 to 1. In
25 another embodiment of the present invention c is an integer from 0 to 1. In yet another embodiment of the present invention the sum of b and c is an integer

form 0 to 2, preferably an integer from 0 to 1. In yet another embodiment of the present invention b is an integer from 0 to 2 and c is 0.



In an embodiment of the present invention,  is a ring structure



selected from the group consisting of




5



. In another embodiment of the present

invention, is a ring structure selected from the group consisting of



In an embodiment of the present invention,  is a ring structure





10 selected from the group consisting of 2-(chromanyl), 2-(6-chloro-2,3-dihydro-benzo[1,4]dioxinyl), 2-(benzo[1,3]dioxolyl), 2-(5-chloro-2,3-dihydro-benzo[1,4]dioxinyl), 2-(7-nitro-2,3-dihydro-benzo[1,4]dioxinyl), 2-(6,7-dichloro-2,3-dihydro-benzo[1,4]dioxinyl), 2-(2,3-dihydro-naphtho[2,3-b][1,4]dioxinyl) and 2-(7-chloro-benzo[1,3]dioxolyl). In another embodiment of the present



15 invention,  is a ring structure selected from the group consisting of 2-(5-chloro-2,3-dihydro-benzo[1,4]dioxinyl), 2-(7-nitro-2,3-dihydro-




benzo[1,4]dioxinyl), 2-(6,7-dichloro-2,3-dihydro-benzo[1,4]dioxinyl) and 2-(2,3-dihydro-naphtho[2,3-b][1,4]dioxinyl).

In an embodiment of the present invention,  is selected from the group consisting of


5 , ,



,  and

.

In another embodiment of the present invention,

 is selected from the group consisting of

,

 and .

In an embodiment of the present invention,  is selected from the group consisting of 2-(2,3-dihydro-benzo[1,4]dioxinyl), 2-(benzo[1,3]dioxolyl), 2-(3,4-dihydro-benzo[1,4]dioxepinyl), 2-(6-chloro-2,3-dihydro-benzo[1,4]dioxinyl), 2-(6-fluoro-2,3-dihydro-benzo[1,4]dioxinyl), 2-(chromanyl), 2-(5-fluoro-2,3-dihydro-benzo[1,4]dioxinyl), 2-(7-chloro-2,3-dihydro-benzo[1,4]dioxinyl), 2-(6-chloro-benzo[1,3]dioxolyl), 2-(7-nitro-2,3-dihydro-benzo[1,4]dioxinyl), 2-(7-methyl-2,3-dihydro-benzo[1,4]dioxinyl), 2-(5-chloro-2,3-dihydro-benzo[1,4]dioxinyl), 2-(6-bromo-2,3-dihydro-benzo[1,4]dioxinyl), 2-(6,7-dichloro-

2,3-dihydro-benzo[1,4]dioxinyl), 2-(8-chloro-2,3-dihydro-benzo[1,4]dioxinyl), 2-(2,3-dihydro-naphtho[2,3-b][1,4]dioxinyl) and 2-(4-methyl-benzo[1,3]dioxolyl).



In another embodiment of the present invention,  is selected from the group consisting 2-(benzo[1,3]dioxolyl), 2-(2,3-dihydro-
5 benzo[1,4]dioxinyl), 2-(6-chloro-2,3-dihydro-benzo[1,4]dioxinyl), 2-(7-chloro-2,3-dihydro-benzo[1,4]dioxinyl), 2-(7-methyl-2,3-dihydro-benzo[1,4]dioxinyl), 2-(6-bromo-2,3-dihydro-benzo[1,4]dioxinyl) and 2-(6,7-dichloro-2,3-dihydro-





benzo[1,4]dioxinyl). In another embodiment of the present invention,  is selected from the group consisting of 2-(2,3-dihydro-benzo[1,4]dioxinyl), 2-(7-methyl-2,3-dihydro-benzo[1,4]dioxinyl) and 2-(6-bromo-2,3-dihydro-benzo[1,4]dioxinyl).
10

In an embodiment of the present invention R^3 is selected from the group consisting of halogen, lower alkyl, hydroxy substituted lower alkyl, -O-(lower alkyl), nitro, cyano, amino, lower alkylamino and di(lower alkyl)amino. In
15 another embodiment of the present invention R^3 is selected from the group consisting of halogen and nitro. In another embodiment of the present invention R^3 is selected from the group consisting of chloro and nitro.

In an embodiment of the present invention R^5 is selected from the group consisting of (II) halogen and lower alkyl. In another embodiment of the
20 present invention R^5 is selected from chloro, fluoro, bromo and methyl.

In an embodiment of the present invention, in the compound of formula






In an embodiment of the present invention, the stereo-center on the
 5 compound of formula (I) is in the S-configuration. In another embodiment of
 the present invention, the stereo-center on the compound of formula (I) is in the
 R-configuration.

In an embodiment of the present invention the compound of formula (I)
 is present as an enantiomerically enriched mixture, wherein the % enantiomeric
 10 enrichment (%ee) is greater than about 75%, preferably greater than about
 85%, more preferably greater than about 90%, more preferably greater than
 about 95%, most preferably greater than about 98%.

In another embodiment, the present invention is directed to one or more
 of the representative compounds of formula (I) as listed in Table 1, below. In
 15 Table 1 below, the column headed "stereo" defines the stereo-configuration at
 the carbon atom of the heterocycle attached at the starred bond. Where no
 designation is listed, the compound was prepared as a mixture of stereo-
 configurations. Where an "R" or "S" designation is listed, the stereo-
 configuration was based on the enantiomerically enriched starting material.

20

Table 1: Representative Compounds of Formula (I)

						
ID No.		Stereo	(CH ₂) _a	NR ⁴	R ¹	R ²
1	2-(2,3-dihydro- benzo[1,4]dioxinyl)		CH ₂	NH	H	H


2	2-(benzo[1,3]dioxolyl)		CH ₂	NH	H	H
3	2-(3,4-dihydro-2H-benzo[1,4]dioxepinyl)		CH ₂	NH	H	H
4	2-(2,3-dihydro-benzo[1,4]dioxinyl)	S	CH ₂	NH	H	H
5	2-(2,3-dihydro-benzo[1,4]dioxinyl)	R	CH ₂	NH	H	H
7	2-(2,3-dihydro-benzo[1,4]dioxinyl)		CH ₂	N(CH ₃)	H	H
8	2-(6-chloro-2,3-dihydro-benzo[1,4]dioxinyl)	S	CH ₂	NH	H	H
9	2-(6-fluoro-2,3-dihydro-benzo[1,4]dioxinyl)	S	CH ₂	NH	H	H
10	2-(chromanyl)		CH ₂	NH	H	H
13	2-(5-fluoro-2,3-dihydro-benzo[1,4]dioxinyl)	S	CH ₂	NH	H	H
14	2-(7-chloro-2,3-dihydro-benzo[1,4]dioxinyl)	S	CH ₂	NH	H	H
15	2-(6-chloro-benzo[1,3]dioxolyl)		CH ₂	NH	H	H
16	2-(2,3-dihydro-benzo[1,4]dioxinyl)		CH ₂ CH ₂	NH	H	H
18	2-(7-nitro-2,3-dihydro-benzo[1,4]dioxinyl)	S	CH ₂	NH	H	H
19	2-(7-methyl-2,3-dihydro-benzo[1,4]dioxinyl)	S	CH ₂	NH	H	H
20	2-(5-chloro-2,3-dihydro-benzo[1,4]dioxinyl)	S	CH ₂	NH	H	H
22	2-(8-methoxy-2,3-dihydro-benzo[1,4]dioxinyl)	S	CH ₂	NH	H	H

24	2-(6-bromo-2,3-dihydro-benzo[1,4]dioxinyl)	S	CH ₂	NH	H	H
29	2-(6,7-dichloro-2,3-dihydro-benzo[1,4]dioxinyl)	S	CH ₂	NH	H	H
30	2-(8-chloro-2,3-dihydro-benzo[1,4]dioxinyl)	S	CH ₂	NH	H	H
33	2-(2,3-dihydro-naphtho[2,3-b][1,4]dioxinyl)	S	CH ₂	NH	H	H
35	2-(4-methyl-benzo[1,3]dioxolyl)		CH ₂	NH	H	H


Additional embodiments of the present invention, include those wherein the substituents selected for one or more of the variables defined herein (e.g. a,

5 R^1 , R^4 , R^5 , , etc.) are independently selected to be any individual substituent or any subset of substituents selected from the complete list as defined herein.

The present invention is further directed to compounds of formula (XII)



10 wherein $-C(O)OR^0$ is a nitrogen protecting group and wherein , a, R^4 and R^1 are as herein defined. In an embodiment, the present invention is directed to a compound of formula (XII-S)



wherein $-C(O)OR^0$ is a nitrogen protecting group. Preferably, $-C(O)OR^0$ is a nitrogen protecting group wherein R^0 is selected from the group consisting of alkoxycarbonyl, aryloxycarbonyl, aralkyloxycarbonyl. More preferably, R^0 is selected from the group consisting of C_{1-4} alkyl (preferably t-butyl), benzyl, p-methoxybenzyl, phenylethyl, phenyl, naphthyl, cycloalkyl and 9-fluorenylmethyl. More preferably, R^0 is selected from the group consisting of C_{1-4} alkyl (preferably t-butyl), benzyl, p-methoxybenzyl and 9-fluorenylmethyl. More preferably still, R^0 is selected from the group consisting of t-butyl, benzyl, p-methoxybenzyl and 9-fluorenylmethyl.

As used herein, unless otherwise noted, “**halogen**” shall mean chlorine, bromine, fluorine and iodine.

As used herein, unless otherwise noted, the term “**alkyl**” whether used
15 alone or as part of a substituent group, includes straight and branched chains.
For example, alkyl radicals include methyl, ethyl, propyl, isopropyl, butyl,
isobutyl, sec-butyl, t-butyl, pentyl and the like. Unless otherwise noted, “**lower**”
when used with alkyl means a carbon chain composition of 1-4 carbon atoms.

As used herein, unless otherwise noted, “**alkoxy**” shall denote an oxygen
20 ether radical of the above described straight or branched chain alkyl groups. For
example, methoxy, ethoxy, n-propoxy, sec-butoxy, t-butoxy, n-hexyloxy and the
like.

As used herein, unless otherwise noted, “**aryl**” shall refer to unsubstituted
25 carbocyclic aromatic groups such as phenyl, naphthyl, and the like, preferably
phenyl.

As used herein, unless otherwise noted, "**aralkyl**" shall mean any lower alkyl group substituted with an aryl group such as phenyl, naphthyl and the like. For example, benzyl, phenylethyl, phenylpropyl, naphthylmethyl, and the like, preferably benzyl.

As used herein, unless otherwise noted, the term “**cycloalkyl**” shall mean any stable monocyclic, bicyclic or polycyclic, saturated ring system, preferably a monocyclic or bicyclic saturated ring system, more preferably a monocyclic saturated ring system. Suitable examples include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, adamantyl, and the like.

As used herein, the notation “*” shall denote the presence of a stereogenic center.

As used herein, unless otherwise noted, the term “**enantiomerically enriched**” when used to describe a compound with one stereogenic center, shall mean that one stereo-configuration of the compound is present in a greater amount than the opposite stereo-configuration of said compound. Preferably, when the compound is said to be enantiomerically enriched, the desired enantiomer of said compound is present in an enantiomeric excess of at least about 75 percent ee, more preferably at least 85 percent ee, more preferably at least 90 percent ee, more preferably at least 95 percent ee, more preferably at least 98 percent ee, most preferably at least 99 percent ee.

As used herein, unless otherwise noted, the term “**nitrogen protecting group**” shall mean any ester group which can act as a protecting group for a amine, amide, sulfamide or sulfanamide nitrogen. Suitable examples include, but are not limited to alkoxycarbonyl, aryloxycarbonyl, aralkyloxycarbonyl, and the like. For example, the nitrogen protecting group may be of the formula – C(O)OR⁰, wherein R⁰ is C₁₋₄alkyl (preferably t-butyl), benzyl, p-methoxybenzyl, phenylethyl, phenyl, naphthyl, cycloalkyl, 9-fluorenylmethyl, and the like, and wherein any of the R⁰ groups may be further substituted.

Abbreviations used in the specification, particularly the Schemes and Examples, are as follows:

BOC or Boc	=	t-Butoxycarbonyl (-C(O)O-C(CH ₃) ₃)
DIPEA or DIEA	=	Diisopropylethylamine
DMAP	=	4-(N,N-Dimethylamino)pyridine

DMF	=	N,N-Dimethylformamide
EtOAc	=	Ethyl acetate
Fmoc	=	9-Fluorenylmethoxycarbonyl
MTBE	=	Methyl t-butyl ether
Pd/C	=	Palladium on Carbon Catalyst
RBF	=	Round bottom flask
RT or rt	=	Room temperature
TEA	=	Triethylamine
TFA	=	Trifluoroacetic Acid
THF	=	Tetrahydrofuran
XRD	=	X-ray Diffraction

As used herein, unless otherwise noted, the terms “**epilepsy and related disorders**” or “**epilepsy or related disorder**” shall mean any disorder in which a subject (preferably a human adult, child or infant) experiences one or more

5 seizures and / or tremors. Suitable examples include, but are not limited to, epilepsy (including, but not limited to, localization-related epilepsies, generalized epilepsies, epilepsies with both generalized and local seizures, and the like), seizures as a complication of a disease or condition (such as seizures associated with encephalopathy, phenylketonuria, juvenile Gaucher’s disease, Lundborg’s

10 progressive myoclonic epilepsy, stroke, head trauma, stress, hormonal changes, drug use or withdrawal, alcohol use or withdrawal, sleep deprivation, and the like), essential tremor, restless limb syndrome, and the like. Preferably, the disorder is selected from epilepsy (regardless of type, underlying cause or origin), essential tremor or restless limb syndrome, more preferably, the disorder is epilepsy

15 (regardless of type, underlying cause or origin) or essential tremor.

The term “**subject**” as used herein, refers to an animal, preferably a mammal, most preferably a human, who is or has been the object of treatment, observation or experiment.

20 The term “**therapeutically effective amount**” as used herein, means that amount of active compound or pharmaceutical agent that elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a

researcher, veterinarian, medical doctor or other clinician, which includes alleviation of the symptoms of the disease or disorder being treated.

As used herein, the term “**composition**” is intended to encompass a product comprising the specified ingredients in the specified amounts, as well
5 as any product which results, directly or indirectly, from combinations of the specified ingredients in the specified amounts.

One skilled in the art will recognize that, where not otherwise specified, the reaction step(s) in the specification and claims are performed under suitable conditions (e.g. temperature, pressure, with appropriate solvents
10 and/or reactants), according to known methods, to provide the desired product. The term “**suitable conditions**” shall mean a reaction step is performed under appropriate conditions (e.g. temperature, pressure, with appropriate solvents and/or reactants) according to known methods to provide the desired product.

One skilled in the art will also recognize that, in the specification and
15 claims as presented herein, wherein a reagent or reagent class/type/ (e.g. base, solvent, etc.) is recited in more than one step of a process, the individual reagents are independently selected for each reaction step and may be the same or different from each other. For example wherein two steps of a process recite an organic or inorganic base as a reagent, the organic or inorganic base
20 selected for the first step may be the same or different than the organic or inorganic base of the second step.

To provide a more concise description, some of the quantitative expressions given herein are not qualified with the term “**about**”. It is understood that whether the term “about” is used explicitly or not, every
25 quantity given herein is meant to refer to the actual given value, and it is also meant to refer to the approximation to such given value that would reasonably be inferred based on the ordinary skill in the art, including approximations due to the experimental and/or measurement conditions for such given value.

As used herein, unless otherwise noted, the term “**aprotic solvent**” shall
30 mean any solvent that does not yield a proton. Suitable examples include, but are not limited to DMF, dioxane, THF, acetonitrile, pyridine, dichloroethane, dichloromethane, MTBE, toluene, and the like.

As used herein, unless otherwise noted, the term “**leaving group**” shall mean a charged or uncharged atom or group which departs during a substitution or displacement reaction. Suitable examples include, but are not limited to, Br, Cl, I, mesylate, tosylate, and the like.

5 As used herein, unless otherwise noted, the term “**nitrogen protecting group**” shall mean a group which may be attached to a nitrogen atom to protect said nitrogen atom from participating in a reaction and which may be readily removed following the reaction. Suitable nitrogen protecting groups include, but are not limited to carbamates – groups of the formula $-C(O)O-R$
10 wherein R is for example methyl, ethyl, t-butyl, benzyl, phenylethyl, p-methoxybenzyl, 9-fluorenylmethyl, and the like, $CH_2=CH-CH_2-$, and the like; amides – groups of the formula $-C(O)-R'$ wherein R' is for example methyl, phenyl, trifluoromethyl, and the like; N-sulfonyl derivatives - groups of the formula $-SO_2-R''$ wherein R'' is for example tolyl, phenyl, trifluoromethyl,
15 2,2,5,7,8-pentamethylchroman-6-yl-, 2,3,6-trimethyl-4-methoxybenzene, and the like. Other suitable nitrogen protecting groups may be found in texts such as T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John Wiley & Sons, 1991.

20 Where the compounds according to this invention have at least one **chiral center**, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral centers, they may additionally exist as diastereomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention. Preferably,
25 wherein the compound is present as an enantiomer, the enantiomer is present at an enantiomeric excess of greater than or equal to about 80%, more preferably, at an enantiomeric excess of greater than or equal to about 90%, more preferably still, at an enantiomeric excess of greater than or equal to about 95%, more preferably still, at an enantiomeric excess of greater than or
30 equal to about 98%, most preferably, at an enantiomeric excess of greater than or equal to about 99%. Similarly, wherein the compound is present as a diastereomer, the diastereomer is present at an diastereomeric excess of greater than or equal to about 80%, more preferably, at an diastereomeric

excess of greater than or equal to about 90%, more preferably still, at an diastereomeric excess of greater than or equal to about 95%, more preferably still, at an diastereomeric excess of greater than or equal to about 98%, most preferably, at an diastereomeric excess of greater than or equal to about 99%.

5 Furthermore, some of the crystalline forms for the compounds of the present invention may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds of the present invention may form solvates with water (i.e., hydrates) or common organic solvents, and such solvates are also intended to be encompassed
10 within the scope of this invention.

One skilled in the art will recognize that wherein a reaction step of the present invention may be carried out in a variety of solvents or solvent systems, said reaction step may also be carried out in a mixture of the suitable solvents or solvent systems.


15 Where the processes for the preparation of the compounds according to the invention give rise to mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The
20 compounds may, for example, be resolved into their component enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-D-tartaric acid and/or (+)-di-p-toluoyl-L-tartaric acid followed by fractional crystallization and regeneration of the free base. The compounds may also be resolved by
25 formation of diastereomeric esters or amides, followed by chromatographic separation and removal of the chiral auxiliary. Alternatively, the compounds may be resolved using a chiral HPLC column.

During any of the processes for preparation of the compounds of the present invention, it may be necessary and/or desirable to protect sensitive or
30 reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in Protective Groups in Organic Chemistry, ed. J.F.W. McOmie, Plenum Press, 1973; and T.W. Greene & P.G.M. Wuts, Protective Groups in Organic Synthesis, John

Wiley & Sons, 1991. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

For use in medicine, the salts of the compounds of this invention refer to
 5 non-toxic "**pharmaceutically acceptable salts**." Other salts may, however, be useful in the preparation of compounds according to this invention or of their pharmaceutically acceptable salts. Suitable pharmaceutically acceptable salts of the compounds include acid addition salts which may, for example, be formed by mixing a solution of the compound with a solution of a
 10 pharmaceutically acceptable base (preferably a strong base) such as NaOH, KOH, NaH, choline hydroxide, and the like.

The present invention is directed to a process for the preparation of compounds of formula (I). Compounds of formula (IA) (compounds of formula
 15 (I) wherein R¹ is hydrogen) may be prepared as outlined in more detail in Scheme 1, below.



Scheme 1

20 Accordingly, a suitably substituted compound of formula (X), a known compound or compound prepared by known methods, is reacted with a suitably substituted compound of formula (XI), wherein $-C(O)OR^0$ is a suitably selected nitrogen protecting group, for example, an alkoxycarbonyl, aryloxycarbonyl, aralkyloxycarbonyl, and the like, for example, wherein R⁰ is C₁₋₄alkyl (preferably
 25 t-butyl), benzyl, p-methoxybenzyl, phenylethyl, phenyl, naphthyl, cycloalkyl, 9-


fluorenylmethyl, and the like, and wherein any of the R⁰ groups may be further substituted, a known compound or compound prepared by known methods;

in the presence of an organic or inorganic base, wherein the organic or inorganic base is not reactive with the chloro group on the compound of

5 formula (XI), preferably, an organic base, more preferably a tertiary amine base such as DIPEA, TEA, pyridine, N-methylmorpholine, N-methylpiperidine, and the like, preferably pyridine; wherein the base is preferably present in an amount greater than about 1 molar equivalent, more preferably, the base is present in an amount in the range of from about 1.1 to about 3.0 molar
10 equivalents; most preferably in an amount of about 2.0 molar equivalents; in an aprotic organic solvent such as DMF, THF, acetonitrile, and the like, preferably acetonitrile; preferably at a temperature in the range of from about 0°C to about 50°C; to yield the corresponding compound of formula (XII).

15 The compound of formula (XII) is de-protected according to known methods, to yield the corresponding compound of formula (IA). For example, by reacting the compound of formula (XII) with an acid or reacting the compound of formula (XII) with hydrogen or a source of hydrogen. Preferably, wherein the nitrogen protecting group (–C(O)OR⁰) is BOC, the compound of
20 formula (XII) is de-protected by reacting with an acid such as TFA, HCl, and the like, preferably HCl; in an organic solvent such as THF, ethyl acetate, and the like. Alternatively, wherein the nitrogen protecting group (–C(O)OR⁰) is benzyl, the compound of formula (XII) is de-protected by reacting with hydrogen or a source of hydrogen, such as hydrogen gas, in the presence of a catalyst such as
25 Pd/C, at a pressure in the range of from about 10 psi to about 60 psi, in an organic solvent such as ethanol, methanol, toluene acetic acid, and the like. Alternatively, wherein the nitrogen protecting group (–C(O)OR⁰) is 9-fluorenylmethyl, the compound of formula (XII) is de-protected by reacting with a base such as an amine base such as piperidine, morpholine, and the like, in
30 an organic solvent DMF, and the like.

Compounds of formula (IB) (compounds of formula (I) wherein R^1 is selected from the group consisting of lower alkyl) may be prepared according to the process as outlined in Scheme 2, below.



Scheme 2


Accordingly, a suitably substituted compound of formula (XII), prepared as in Scheme 1 above, is reacting with a suitably selected compound of formula (XV), an alkyl halide or alkyl sulfonate, wherein Q is a suitable leaving group such as Cl, Br, I, $-O-SO_2-CH_3$ (mesylate) $-O-SO_2-CF_3$ (triflate) $-O-SO_2-$ tolyl (tosylate), and the like; in the presence of an organic or inorganic base such as K_2CO_3 , Na_2CO_3 , NaOH, KOH, pyridine, DIPEA, TEA, and the like, preferably a tertiary amine base, more preferably pyridine; in an organic solvent such as THF, acetonitrile, DMF, and the like; preferably at a temperature in the range of from about $0^\circ C$ to about $50^\circ C$; to yield the corresponding compound of formula (XVI).

The compound of formula (XVI) is de-protected according to known methods, to yield the corresponding compound of formula (IB). For example, by reacting the compound of formula (XVI) with an acid or reacting the compound of formula (XVI) with hydrogen or a source of hydrogen. Preferably, wherein the nitrogen protecting group $(-C(O)OR^0)$ is BOC, the compound of formula (XVI) is de-protected by reacting with an acid such as TFA, HCl, and the like, preferably HCl; in an organic solvent such as THF, ethyl acetate, and the like. Alternatively, wherein the nitrogen protecting group $(-C(O)OR^0)$ is benzyl, the compound of formula (XVI) is de-protected by reacting with hydrogen or a source of hydrogen, such as hydrogen gas, in the present of a catalyst such as Pd/C, at a pressure in the range of from about 10 psi to about

60 psi, in an organic solvent such as ethanol, methanol, toluene acetic acid, and the like. Alternatively, wherein the nitrogen protecting group ($-\text{C}(\text{O})\text{OR}^0$) is 9-fluorenylmethyl, the compound of formula (XVI) is de-protected by reacting with a base such as an amine base such as piperidine, morpholine, and the like, in an organic solvent DMF, and the like.

The compound of formula (I) is isolated according to known methods, for example by filtration, solvent evaporation, and the like. Preferably, the compound of formula (I) is further purified according to known methods, for example by recrystallization from a suitable solvent such as water, toluene, and the like, preferably toluene.

In an embodiment, the present invention is directed to a process for the preparation of the compound of formula (I-S), as outlined in Scheme 3, below.



Scheme 3

Accordingly, a suitably substituted compound of formula (X-S), also known as C-(6-chloro-2,3-dihydro-benzo[1,4]dioxin-2-yl)-methanamine, a known compound, wherein $-\text{C}(\text{O})\text{OR}^0$ is a suitably selected nitrogen protecting group, for example, an alkoxycarbonyl, aryloxycarbonyl, aralkyloxycarbonyl, and the like, for example, wherein R^0 is C_{1-4} alkyl (preferably t-butyl), benzyl, p-methoxybenzyl, phenylethyl, phenyl, naphthyl, cycloalkyl, 9-fluorenylmethyl,

and the like, and wherein any of the R⁰ groups may be further substituted, a known compound or compound prepared by known methods;


in the presence of an organic or inorganic base, wherein the organic or inorganic base is not reactive with the chloro group on the compound of formula (XI), preferably, an organic base, more preferably a tertiary amine base such as DIPEA, TEA, pyridine, N-methylmorpholine, N-methylpiperidine, and the like, preferably pyridine; wherein the base is preferably present in an amount greater than about 1 molar equivalent, more preferably, the base is present in an amount in the range of from about 1.1 to about 3.0 molar equivalents; most preferably in an amount of about 2.0 molar equivalents; in an aprotic organic solvent such as DMF, THF, acetonitrile, and the like, preferably acetonitrile; preferably at a temperature in the range of from about 0°C to about 50°C; to yield the corresponding compound of formula (XII-S).

The compound of formula (XII-S) is de-protected according to known methods, to yield the corresponding compound of formula (I-S). For example, by reacting the compound of formula (XII-S) with an acid or reacting the compound of formula (XII-S) with hydrogen or a source of hydrogen. Preferably, wherein the nitrogen protecting group (–C(O)OR⁰) is BOC, the compound of formula (XII-S) is de-protected by reacting with an acid such as TFA, HCl, and the like, preferably HCl; in an organic solvent such as THF, ethyl acetate, and the like. Alternatively, wherein the nitrogen protecting group (–C(O)OR⁰) is benzyl, the compound of formula (XII-S) is de-protected by reacting with hydrogen or a source of hydrogen, such as hydrogen gas, in the present of a catalyst such as Pd/C, at a pressure in the range of from about 10 psi to about 60 psi, in an organic solvent such as ethanol, methanol, toluene acetic acid, and the like. Alternatively, wherein the nitrogen protecting group (–C(O)OR⁰) is 9-fluorenylmethyl, the compound of formula (XII-S) is de-protected by reacting with a base such as an amine base such as piperidine, morpholine, and the like, in an organic solvent DMF, and the like.

The compound of formula (I-S) is isolated according to known methods, for example by filtration. Preferably, the compound of formula (I-S) is further

purified according to known methods, for example by recrystallization from a suitable solvent such as water, toluene, and the like, preferably toluene.


- Compounds of formula (XI) are known compounds or compounds which may be prepared according to known methods. For example, the compounds of formula (XI) may be prepared according to the process as outlined in Scheme 4, below.



Scheme 4

- Accordingly, a suitably substituted compound of formula (XIII), a known compound or compound prepared by known methods, is reacted with a suitably selected alcohol, a compound of formula (XIV); wherein the compound of formula (XIV) is preferably present in about 1 molar equivalent; neat or in an aprotic organic solvent such as acetonitrile, ethyl acetate, toluene, and the like, provided that the compound of formula (XIII) and the compound of formula (XIV) are at least partially soluble in the solvent and unreactive to the solvent; preferably, at a temperature in the range of from about 0° to about room temperature; to yield the corresponding compound of formula (XI).

- Compounds of formula (XI-S) wherein R⁰ is t-butyl may be prepared according to the process outlined in Scheme 5.



Scheme 5

- Accordingly, isocyanatidosulfonyl chloride, a known compound, is reacted t-butanol, a known compound, wherein the t-butanol is preferably present in about 1 molar equivalent; neat or in an aprotic organic solvent such

- as, acetonitrile, ethyl acetate, toluene, and the like, provided that the compound of formula (XIII) and the compound of formula (XIV) are at least partially soluble in the solvent and unreactive to the solvent, preferably neat or in acetonitrile; preferably, at a temperature in the range of from about 0° to about room
- 5 temperature, morepreferably, at a temperature of about 0°C; to yield the corresponding compound of formula (XI-S), also known as [(1,1-dimethylethoxy)carbonyl]-sulfamoyl chloride.

The present invention is further directed to a crystalline form of the compound of formula (I-S). The crystalline form of the compound of formula (I-S) may be characterized by their corresponding Powder X-ray Diffraction (PXRD) spectra.

In an embodiment, the crystalline form of the compound of formula (I-S) may be characterized by its corresponding PXRD peaks, wherein the peaks

15 have a relative intensity of greater than or equal to about 10% relative intensity; preferably, wherein the peaks have a relative intensity of greater than or equal to about 25% relative intensity.

In an embodiment, the crystalline form of the compound of formula (I-S) may be characterized by its corresponding PXRD peaks, wherein the peaks are

20 defined by their position ($^{\circ}2\theta$), d-spacing (Å) and relative intensity (%). In another embodiment, the crystalline form of the compound of formula (I-S) may be characterized by its corresponding PXRD peaks, wherein the peaks are defined by their position ($^{\circ}2\theta$) and d-spacing (Å).

25 A powder XRD spectra was measured for a representative sample of the crystalline form (I-SA) of the compound of formula (I-S), as shown in Figure 1, with characteristic peaks as listed in Table 2 below. The PXRD spectra were measured using an X-Celerator detector, scanning from 3 to 35 $^{\circ}2\theta$, at a step size of 0.0165 $^{\circ}2\theta$, a time per step of 10.16 sec, an effective scan speed of

30 0.2067 $^{\circ}$ /sec, instrument voltage of 45 kV and a current setting of 40 mA.

Table 2: Form (I-S)

Position ($^{\circ}2\theta$)	d-spacing (Å)	Relative Intensity (%)
4.44	19.92	33
15.50	5.72	14
17.32	5.12	48
18.57	4.78	100
19.39	4.58	10
19.86	4.47	30
20.03	4.43	20
20.88	4.26	51
21.57	4.12	23
21.93	4.05	24
22.71	3.92	13
23.19	3.84	14
23.90	3.72	29
24.53	3.63	16
25.02	3.56	25
26.04	3.42	19
26.71	3.34	16
26.84	3.32	13
28.28	3.16	26
29.96	2.98	12
30.70	2.91	21

The present invention further comprises pharmaceutical compositions containing one or more of the compounds prepared according to any of the processes described herein with a pharmaceutically acceptable carrier.

- 5 Pharmaceutical compositions containing one or more of the compounds of the invention described herein as the active ingredient can be prepared by intimately mixing the compound or compounds with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques. The carrier may take a wide variety of forms depending upon the desired route of
- 10 administration (e.g., oral, parenteral). Thus for liquid oral preparations such as suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, stabilizers, coloring agents and the like; for solid oral preparations, such as powders, capsules and tablets, suitable carriers and additives include starches, sugars, diluents,
- 15 granulating agents, lubricants, binders, disintegrating agents and the like. Solid oral preparations may also be coated with substances such as sugars or be enteric-coated so as to modulate major site of absorption. For parenteral administration, the carrier will usually consist of sterile water and other

ingredients may be added to increase solubility or preservation. Injectable suspensions or solutions may also be prepared utilizing aqueous carriers along with appropriate additives.

To prepare the pharmaceutical compositions of this invention, one or more compounds of the present invention as the active ingredient is intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending of the form of preparation desired for administration, e.g., oral or parenteral such as intramuscular. In preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed. Thus, for liquid oral preparations, such as for example, suspensions, elixirs and solutions, suitable carriers and additives include water, glycols, oils, alcohols, flavoring agents, preservatives, coloring agents and the like; for solid oral preparations such as, for example, powders, capsules, caplets, gelcaps and tablets, suitable carriers and additives include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit form, in which case solid pharmaceutical carriers are obviously employed. If desired, tablets may be sugar coated or enteric coated by standard techniques. For parenterals, the carrier will usually comprise sterile water, through other ingredients, for example, for purposes such as aiding solubility or for preservation, may be included. Injectable suspensions may also be prepared, in which case appropriate liquid carriers, suspending agents and the like may be employed. The pharmaceutical compositions herein will contain, per dosage unit, e.g., tablet, capsule, powder, injection, teaspoonful and the like, an amount of the active ingredient necessary to deliver an effective dose as described above. The pharmaceutical compositions herein will contain, per unit dosage unit, e.g., tablet, capsule, powder, injection, suppository, teaspoonful and the like, of from about 1-1000 mg and may be given at a dosage of from about 0.01-300 mg/kg/day, or any range therein, preferably from about 0.5-100 mg/kg/day, or any range therein, more preferably from about 1.0-25.0 mg/kg/day, or any range therein. The dosages, however, may be varied depending upon the

requirement of the patients, the severity of the condition being treated and the compound being employed. The use of either daily administration or post-periodic dosing may be employed.

Preferably these compositions are in unit dosage forms from such as
5 tablets, pills, capsules, powders, granules, sterile parenteral solutions or suspensions, metered aerosol or liquid sprays, drops, ampoules, autoinjector devices or suppositories; for oral parenteral, intranasal, sublingual or rectal administration, or for administration by inhalation or insufflation. Alternatively, the composition may be presented in a form suitable for once-weekly or once-
10 monthly administration; for example, an insoluble salt of the active compound, such as the decanoate salt, may be adapted to provide a depot preparation for intramuscular injection. For preparing solid compositions such as tablets, the principal active ingredient is mixed with a pharmaceutical carrier, e.g. conventional tableting ingredients such as corn starch, lactose, sucrose,
15 sorbitol, talc, stearic acid, magnesium stearate, dicalcium phosphate or gums, and other pharmaceutical diluents, e.g. water, to form a solid preformulation composition containing a homogeneous mixture of a compound of the present invention, or a pharmaceutically acceptable salt thereof. When referring to these preformulation compositions as homogeneous, it is meant that the active
20 ingredient is dispersed evenly throughout the composition so that the composition may be readily subdivided into equally effective dosage forms such as tablets, pills and capsules. This solid preformulation composition is then subdivided into unit dosage forms of the type described above containing from 0.1 to about 500 mg of the active ingredient of the present invention. The
25 tablets or pills of the novel composition can be coated or otherwise compounded to provide a dosage form affording the advantage of prolonged action. For example, the tablet or pill can comprise an inner dosage and an outer dosage component, the latter being in the form of an envelope over the former. The two components can be separated by an enteric layer which
30 serves to resist disintegration in the stomach and permits the inner component to pass intact into the duodenum or to be delayed in release. A variety of material can be used for such enteric layers or coatings, such materials

including a number of polymeric acids with such materials as shellac, cetyl alcohol and cellulose acetate.

The liquid forms in which the novel compositions of the present invention may be incorporated for administration orally or by injection include, aqueous solutions, suitably flavoured syrups, aqueous or oil suspensions, and flavoured emulsions with edible oils such as cottonseed oil, sesame oil, coconut oil or peanut oil, as well as elixirs and similar pharmaceutical vehicles. Suitable dispersing or suspending agents for aqueous suspensions, include synthetic and natural gums such as tragacanth, acacia, alginate, dextran, sodium carboxymethylcellulose, methylcellulose, polyvinyl-pyrrolidone or gelatin.

The method of treating epilepsy or a related disorder described in the present invention may also be carried out using a pharmaceutical composition comprising any of the compounds as defined herein and a pharmaceutically acceptable carrier. The pharmaceutical composition may contain between about 0.1 mg and 1000 mg, preferably about 50 to 500 mg, of the compound, or any range therein, and may be constituted into any form suitable for the mode of administration selected. Carriers include necessary and inert pharmaceutical excipients, including, but not limited to, binders, suspending agents, lubricants, flavorants, sweeteners, preservatives, dyes, and coatings. Compositions suitable for oral administration include solid forms, such as pills, tablets, caplets, capsules (each including immediate release, timed release and sustained release formulations), granules, and powders, and liquid forms, such as solutions, syrups, elixirs, emulsions, and suspensions. Forms useful for parenteral administration include sterile solutions, emulsions and suspensions.

Advantageously, compounds of the present invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three or four times daily. Furthermore, compounds for the present invention can be administered in intranasal form via topical use of suitable intranasal vehicles, or via transdermal skin patches well known to those of ordinary skill in that art. To be administered in the form of a transdermal delivery system, the dosage administration will, of course, be continuous rather than intermittent throughout the dosage regimen.

For instance, for oral administration in the form of a tablet or capsule, the active drug component can be combined with an oral, non-toxic pharmaceutically acceptable inert carrier such as ethanol, glycerol, water and the like. Moreover, when desired or necessary, suitable binders; lubricants, disintegrating agents and coloring agents can also be incorporated into the mixture. Suitable binders include, without limitation, starch, gelatin, natural sugars such as glucose or beta-lactose, corn sweeteners, natural and synthetic gums such as acacia, tragacanth or sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride and the like. Disintegrators include, without limitation, starch, methyl cellulose, agar, bentonite, xanthan gum and the like.

The liquid forms in suitably flavored suspending or dispersing agents such as the synthetic and natural gums, for example, tragacanth, acacia, methyl-cellulose and the like. For parenteral administration, sterile suspensions and solutions are desired. Isotonic preparations which generally contain suitable preservatives are employed when intravenous administration is desired.

To prepare a pharmaceutical composition of the present invention, a compound of formula (I) as the active ingredient is intimately admixed with a pharmaceutical carrier according to conventional pharmaceutical compounding techniques, which carrier may take a wide variety of forms depending of the form of preparation desired for administration (e.g. oral or parenteral). Suitable pharmaceutically acceptable carriers are well known in the art. Descriptions of some of these pharmaceutically acceptable carriers may be found in The Handbook of Pharmaceutical Excipients, published by the American Pharmaceutical Association and the Pharmaceutical Society of Great Britain.

Methods of formulating pharmaceutical compositions have been described in numerous publications such as Pharmaceutical Dosage Forms: Tablets, Second Edition, Revised and Expanded, Volumes 1-3, edited by Lieberman et al; Pharmaceutical Dosage Forms: Parenteral Medications, Volumes 1-2, edited by Avis et al; and Pharmaceutical Dosage Forms: Disperse Systems, Volumes 1-2, edited by Lieberman et al; published by Marcel Dekker, Inc.

Compounds of this invention may be administered in any of the foregoing compositions and according to dosage regimens established in the art whenever treatment of epilepsy or related disorders is required.

The daily dosage of the products may be varied over a wide range from
5 0.01 to 1,000 mg per adult human per day. For oral administration, the compositions are preferably provided in the form of tablets containing, 0.01, 0.05, 0.1, 0.5, 1.0, 2.5, 5.0, 10.0, 15.0, 25.0, 50.0, 100, 150, 200, 250 and 500 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the patient to be treated. An effective amount of the drug is ordinarily supplied
10 at a dosage level of from about 0.01 mg/kg to about 300.0 mg/kg of body weight per day, or any range therein. Preferably, the range is from about 0.5 to about 100.0 mg/kg of body weight per day, or any range therein, more preferably, from about 1.0 to about 25.0 mg/kg of body weight per day, or any range therein. The compounds may be administered on a regimen of 1 to 4 times per day.

15 Optimal dosages to be administered may be readily determined by those skilled in the art, and will vary with the particular compound used, the mode of administration, the strength of the preparation, the mode of administration, and the advancement of the disease condition. In addition, factors associated with the particular patient being treated, including patient age, weight, diet and time of
20 administration, will result in the need to adjust dosages.

One skilled in the art will recognize that, both *in vivo* and *in vitro* trials using suitable, known and generally accepted cell and / or animal models are predictive of the ability of a test compound to treat or prevent a given disorder.


One skilled in the art will further recognize that human clinical trails
25 including first-in-human, dose ranging and efficacy trials, in healthy patients and / or those suffering from a given disorder, may be completed according to methods well known in the clinical and medical arts.

The following Examples are set forth to aid in the understanding of the
30 invention, and are not intended and should not be construed to limit in any way the invention set forth in the claims which follow thereafter.


In the Examples which follow, some synthesis products are listed as having been isolated as a residue. It will be understood by one of ordinary skill

in the art that the term “**residue**” does not limit the physical state in which the product was isolated and may include, for example, a solid, an oil, a foam, a gum, a syrup, and the like.

5

Example 1**Toluene-4-sulfonic acid 6-chloro-2,3-dihydro-benzo[1,4]dioxin-2-ylmethyl ester**

A 5L three-neck-round bottom flask equipped with a reflux condenser,
10 nitrogen outlet, overhead stirrer, heating mantle and temperature control unit
was charged with 6-chloro-2,3-dihydro-benzo[1,4]dioxin-2-yl)-methanol (176 g
877.28 mmoles), tetrahydrofuran (2 L), 4-pyridinamine, DMAP (25 g; 204.63
mmoles) and p-toluenesulfonyl chloride (1.05 moles; 200.70 g;). The reaction
mixture was then warmed to 40°C. An additional portion of DMAP (5g) and p-
15 toluenesulfonyl chloride (10 g; 52.45 mmoles) were added and the reaction was
continued overnight. Upon consumption of starting material the reaction was
quenched with water (500mL) and the resulting mixture extract with MTBE
(1.2L), washed with water (300mL, 200mL), then with 1N HCl wash (400mL,
100mL), and an additional water wash (100mL), followed by a bicarbonate
20 wash (100mL). The organic layer was dried over Na₂SO₄, (50g) and silica
(60g). The resulting solution was filtered and concentrated to the title
compound as a thick oil.


Example 2**2-(6-Chloro-2,3-dihydro-benzo[1,4]dioxin-2-ylmethyl)-isoindole-1,3-dione**

A three-necked-round bottom flask equipped with a nitrogen outlet, magnetic stir bar, heating mantle and temperature control unit was charged with toluene-4-sulfonic acid 6-chloro-2,3-dihydro-benzo[1,4]dioxin-2-ylmethyl ester (300 g; 845.52 mmoles), dimethylformamide (330.00 mL) and potassium phthalimide (1.10 moles; 203.59 g). The resulting slurry was warmed to 100°C and allowed to heat for 2h. The reaction mixture was then cooled to room temperature in an ice bath and quenched into a stirred ice / water mixture (1L). A white solid developed upon stirring. After 2h at room temperature, the solid was filtered and air dried to yield the title compound as an off-white solid.

10

Example 3

C-(6-Chloro-2,3-dihydro-benzo[1,4]dioxin-2-yl)-methylamine




A 5L four-neck round bottom flask equipped with a nitrogen outlet, reflux condenser, overhead stirrer, heating mantle and temperature control unit was charged with (S)-2-((6-chloro-2,3-dihydrobenzo[b][1,4]dioxin-2-yl)methyl)isoindoline-1,3-dione (242 g; 733.91 mmoles), ethanol (2.6 L) and hydrazine (48 g; 1.50 moles). The resulting mixture was heated to reflux. After 7h the reaction mixture was cooled to room temperature and treated with HCl (2N) until acidic; and the resulting solids were filtered off. The filtrate was concentrated, then treated with MTBE (1L) and sodium hydroxide (3N, 200mL). The organic layer was separated and washed with water (200mL), followed by a brine (200mL) wash. The organic layer was dried (Na₂SO₄), filtered and concentrated to yield the title compound.

25

Example 4


N-[[[(2S)-6-chloro-2,3-dihydro-1,4-benzodioxin-2-yl]methyl]-N'-[(1,1-dimethylethoxy)carbonyl]-sulfamide



A 5L three-necked-RBF equipped with a nitrogen outlet, magnetic stir bar, addition funnel, internal thermometer and cooling bath was treated with C-(6-chloro-2,3-dihydro-benzo[1,4]dioxin-2-yl)-methylamine (135 g; 676.23
 5 mmoles), acetonitrile (1000 mL), 4-pyridinamine, DMAP (16.91 mmoles; 2.07 g) and pyridine (82.02 mL; 80.24 g). Upon cooling the reaction mixture to 0°C an acetonitrile solution (50mL) of Cl-SO₂-NH-Boc (168 g; 779.01 mmoles) was added via addition funnel. The resulting mixture was then allowed to warm to room temperature with subsequent heating to 45°C (5h). The reaction was
 10 then quenched by addition of 1N HCl until acidic. The product was extracted with ethyl acetate (500mL). After phase separation, the organic layer was treated with 3N NaOH (300mL), the basic aqueous layer was then cooled in an ice bath and treated with water (20mL) and 2N HCl. The product was extracted with ethyl acetate (500 mL), dried (Na₂SO₄), filtered and concentrated to yield
 15 the title compound as a light yellow solid.

Example 5

N-[[[(2S)-6-chloro-2,3-dihydro-1,4-benzodioxin-2-yl]methyl]-sulfamide




20 A 2L three-necked round bottom flask equipped with and 250mL addition funnel, nitrogen outlet, overhead stirrer, heating mantle and temperature control unit was charged with (S)-tert-butyl N-((6-chloro-2,3-dihydrobenzo[b][1,4]dioxin-2-yl)methyl)sulfamoylcarbamate (256 g; 540.61 mmoles), ethyl Acetate (409.60 mL) followed by addition of hydrogen chloride (2.16 moles; 177.54 mL). The
 25 resulting mixture was warmed to 50°C for 1h, then cooled to room temperature then neutralized by addition of NaOH (2N, 220mL) and the product extracted using ethyl acetate (2 x 200mL), the organic layer was washed with brine

(100mL), phase separated and dried with Na₂SO₄ (20g), filtered and concentrated to yield a thick oil (100 g crude).

The oil was taken up in hot water (750mL + 200mL) hot filtered to remove insoluble oil and other impurities. The resulting mixture was cooled to 25°C and the resulting solids were filtered and dried to yield the title compound as a white solid.

Example 6

[(1,1-dimethylethoxy)carbonyl]- sulfamoyl chloride




10

A 100mL three-necked-RBF equipped with a nitrogen outlet, addition funnel, temperature sensor and ice bath was treated with chlorosulfonyl isocyanate (811.47 mmols; 70.76 mL) and acetonitrile (50mL). Upon cooling to 0°C t-butyl alcohol (811.47 mmols; 77.08 mL; 60.15 g) was added as a solution in acetonitrile (50mL) via addition funnel, while the internal temperature of the reaction mixture was maintained at less than 10°C. After addition the reaction was determined to be complete. The reaction mixture was concentrated to a white foam (168g), which was washed with heptane on the filter pad to yield the title compound.

20

Example 7

Toluene Recrystallization of N-[(2S)-6-chloro-2,3-dihydro-1,4-benzodioxin-2-yl]methyl]-sulfamide



25

N-[(2S)-6-chloro-2,3-dihydro-1,4-benzodioxin-2-yl]methyl]-sulfamide (0.5g) was dissolved in hot toluene (5mL), hot-filtered to remove insoluble material and then allowed to slowly cool to room temperature. The resulting

solid was collected by filtration, washed with heptane and air dried to yield the title compound as a crystalline, white solid.

m.p. 98°C.

5

Example 8

Liquid Formulation

The compound of formula (I-S), prepared for example, as in Example 7 above, was formulated according to known methods into liquid formulations of 25mg and 100 mg, respectively, with components as listed in Table 3 below.

10

Table 3: Liquid Formulations

Component	Role	25 mg/mL Suspension	100 mg/mL Suspension
Compound of formula (I-S)	Active	25 mg	100 mg
Hypromellose (also known as HPMC or hydroxypropylmethylcellulose)	Suspending agent	5 mg	5 mg
Purified water	Solvent	q.s. ad. 1 mL	q.s. ad. 1 mL

Example 9 – Prophetic Example


As a specific embodiment of an oral composition, 100 mg of the compound of formula (I-S), prepared as in Example 7 is formulated with sufficient finely divided lactose to provide a total amount of 580 to 590 mg to fill a size O hard gel capsule.

While the foregoing specification teaches the principles of the present invention, with examples provided for the purpose of illustration, it will be understood that the practice of the invention encompasses all of the usual variations, adaptations and/or modifications as come within the scope of the following claims and their equivalents.

25

We Claim:

1. A process for the preparation of a compound of formula (IA)



wherein


- 5 R^1 is hydrogen;

R^4 is selected from the group consisting of hydrogen and lower alkyl;

a is an integer from 1 to 2;





is selected from the group consisting of




wherein b is an integer from 0 to 4; and wherein c is an integer from 0 to

2;

- 15 each R^5 is independently selected from the group consisting of halogen, lower alkyl and nitro;



reacting a compound of formula (X) with a compound of formula (XI)
wherein $-\text{C}(\text{O})\text{OR}^0$ is a nitrogen protecting group; in the presence of an organic
or inorganic base, wherein the organic or inorganic base is not reactive with the
chloro group on the compound of formula (XI); in an aprotic organic solvent; to
10 yield the corresponding compound of formula (XII);




de-protecting the compound of formula (XII); to yield the corresponding
compound of formula (IA).

- 15 2. A process as in Claim 1, wherein a is 1; R^4 is hydrogen and  is 2-(6-chloro-2,3-dihydro-benzo[1,4]dioxinyl).

3. A process as in Claim 1, wherein the organic or inorganic base is a tertiary amine base selected from the group consisting of DIPEA, TEA, pyridine, N-methylmorpholine and N-methylpiperidine.
- 5 4. A process as in Claim 1, wherein the organic or inorganic base is pyridine.
5. A process as in Claim 3, wherein the tertiary amine base is present in an amount in the range of from about 1.1 to about 3.0 molar equivalents.
- 10 6. A process as in Claim 5, wherein the tertiary amine base is present in an amount of about 2.0 molar equivalents.
7. A process as in Claim 1, wherein the aprotic organic solvent is selected
15 from the group consisting of DMF, THF and acetonitrile.
8. A process as in Claim 7, wherein the aprotic organic solvent is acetonitrile.
- 20 9. A process as in Claim 1, wherein $-C(O)OR^0$ is selected from the group consisting of C_{1-4} alkoxycarbonyl, aryloxycarbonyl and aralkyloxycarbonyl.
10. A process as in Claim 9, wherein $-C(O)OR^0$ is selected from the group consisting of lower alkyl, benzyl, p-methoxybenzyl and 9-fluorenylmethyl.
- 25 11. A process as in Claim 1, wherein $-C(O)OR^0$ is $-C(O)O$ -t-butyl.
12. A process as in Claim 1, wherein the compound of formula (XII) is de-protected by reacting the compound of formula (XII) with an acid.
- 30 13. A process as in Claim 12, wherein the compound of formula (XII) is de-protected by reacting the compound of formula (XII) with hydrochloric acid.

14. A process for the preparation of a compound of formula (IB)



wherein


R^1 is selected from the group consisting of lower alkyl;

- 5 R^4 is selected from the group consisting of hydrogen and lower alkyl;

a is an integer from 1 to 2;



is selected from the group consisting of





10

wherein b is an integer from 0 to 4; and wherein c is an integer from 0 to


2;

each R^5 is independently selected from the group consisting of halogen,


- 15 lower alkyl and nitro;



reacting a compound of formula (X) with a compound of formula (XI)
 wherein $-\text{C}(\text{O})\text{OR}^0$ is a nitrogen protecting group; in the presence of an organic
 or inorganic base, wherein the organic or inorganic base is not reactive with the
 chloro group on the compound of formula (XI); in an aprotic organic solvent; to
 10 yield the corresponding compound of formula (XII);



reacting the compound of formula (XII) with a compound of formula (XV),
 wherein Q is a leaving group; in an organic solvent; to yield the corresponding
 compound of formula (XVI)



de-protecting the compound of formula (XVI); to yield the corresponding compound of formula (IB).

15. A process as in Claim 14, wherein the organic or inorganic base is a
5 tertiary amine base selected from the group consisting of DIPEA, TEA, pyridine, N-methylmorpholine and N-methylpiperidine.

16. A process as in Claim 15, wherein the organic or inorganic base is pyridine.

10

17. A process as in Claim 14, wherein the tertiary amine base is present in an amount in the range of from about 1.1 to about 3.0 molar equivalents.

18. A process as in Claim 17, wherein the tertiary amine base is present in
15 an amount of about 2.0 molar equivalents.

19. A process as in Claim 14, wherein the aprotic organic solvent is selected from the group consisting of DMF, THF and acetonitrile.

20. A process as in Claim 19, wherein the aprotic organic solvent is acetonitrile.

21. A process as in Claim 14, wherein $-C(O)OR^0$ is selected from the group consisting of C_{1-4} alkoxycarbonyl, aryloxycarbonyl and aralkyloxycarbonyl.

25

22. A process as in Claim 21, wherein $-C(O)OR^0$ is selected from the group consisting of lower alkyl, benzyl, p-methoxybenzyl and 9-fluorenylmethyl.

23. A process as in Claim 14, wherein $-C(O)OR^0$ is $-C(O)O$ -t-butyl.


30

24. A process as in Claim 14, wherein Q is selected from the group consisting of Cl, Br, I, $-O-SO_2-CH_3$, $-O-SO_2-CF_3$, $-O-SO_2$ -tolyl.


25. A process as in Claim 14, wherein the compound of formula (XII) is de-protected by reacting the compound of formula (XII) with an acid.

26. A process as in Claim 25, wherein the compound of formula (XII) is de-protected by reacting the compound of formula (XII) with hydrochloric acid.


27. A process for the preparation of a compound of formula (I-S)



or a pharmaceutically acceptable salt thereof; comprising



reacting a compound of formula (X-S) with a compound of formula (XI), wherein $-C(O)OR^0$ is a nitrogen protecting group; in the presence of an organic or inorganic base, wherein the organic or inorganic base is not reactive with the chloro group on the compound of formula (XI); in an aprotic organic solvent; to yield the corresponding compound of formula (XII-S);



de-protecting the compound of formula (XII-S); to yield the corresponding compound of formula (I-S).


28. A process as in Claim 27, wherein the organic or inorganic base is a tertiary amine base selected from the group consisting of DIPEA, TEA, pyridine, N-methylmorpholine and N-methylpiperidine.

29. A process as in Claim 27, wherein the organic or inorganic base is pyridine.
- 5 30. A process as in Claim 28, wherein the tertiary amine base is present in an amount in the range of from about 1.1 to about 3.0 molar equivalents.
31. A process as in Claim 30, wherein the tertiary amine base is present in an amount of about 2.0 molar equivalents.
- 10 32. A process as in Claim 27, wherein the aprotic organic solvent is selected from the group consisting of DMF, THF and acetonitrile.
33. A process as in Claim 32, wherein the aprotic organic solvent is
15 acetonitrile.
34. A process as in Claim 27, wherein $-C(O)OR^0$ is selected from the group consisting of C_{1-4} alkoxycarbonyl, aryloxycarbonyl and aralkyloxycarbonyl.
- 20 35. A process as in Claim 34, wherein $-C(O)OR^0$ is selected from the group consisting of lower alkyl, benzyl, p-methoxybenzyl and 9-fluorenylmethyl.
36. A process as in Claim 27, wherein $-C(O)OR^0$ is $-C(O)O$ -t-butyl.
- 25 37. A process as in Claim 27, wherein the compound of formula (XII-S) is de-protected by reacting the compound of formula (XII-S) with an acid.
38. A process as in Claim 37, wherein the compound of formula (XII-S) is de-protected by reacting the compound of formula (XII-S) with hydrochloric
30 acid.
39. A process as in Claim 27, wherein the compound of formula (I-S) is further recrystallized.

40. A process as in Claim 39, wherein the compound of formula (I-S) is recrystallized from a solvent selected from the group consisting of water and toluene.

5

41. A compound of formula (XII)



wherein

$-\text{C}(\text{O})\text{OR}^0$ is a nitrogen protecting group;





10 R^1 is selected from the group consisting of hydrogen and lower alkyl;

R^4 is selected from the group consisting of hydrogen and lower alkyl;


a is an integer from 1 to 2;



is selected from the group consisting of




15




wherein b is an integer from 0 to 4; and wherein c is an integer from 0 to 2;

- 5 each R^5 is independently selected from the group consisting of halogen, lower alkyl and nitro;



- 10 42. A compound as in Claim 41, wherein a is 1; R^4 is hydrogen and



43. A compound as in Claim 41, wherein $-C(O)OR^0$ is selected from the group consisting of C_{1-4} alkoxycarbonyl, aryloxycarbonyl and aralkyloxycarbonyl.


15

44. A process as in Claim 43, wherein $-C(O)OR^0$ is selected from the group consisting of lower alkyl, benzyl, p-methoxybenzyl and 9-fluorenylmethyl.

45. A process as in Claim 41, wherein $-C(O)OR^0$ is $-C(O)O$ -t-butyl.

46. A process as in Claim 39, wherein the compound of formula (I-S) is recrystallized from toluene.

5 47. A compound of formula (XII-S)



wherein $-C(O)OR^0$ is a nitrogen protecting group.

48. A compound as in Claim 47, wherein $-C(O)OR^0$ is selected from the
10 group consisting of C_{1-4} alkoxycarbonyl, aryloxycarbonyl and aralkyloxycarbonyl.

49. A process as in Claim 47, wherein $-C(O)OR^0$ is selected from the group consisting of lower alkyl, benzyl, p-methoxybenzyl and 9-fluorenylmethyl.

15 50. A process as in Claim 47, wherein $-C(O)OR^0$ is $-C(O)O$ -t-butyl.

51. A product prepared according to the process of Claim 1.

52. A product prepared according to the process of Claim 27.

20

53. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and the product of Claim 52.


54. A pharmaceutical composition made by mixing the product of Claim 52
25 and a pharmaceutically acceptable carrier.

55. A process for making a pharmaceutical composition comprising mixing the product of Claim 52 and a pharmaceutically acceptable carrier.

56. A method of treating epilepsy or a related disorder comprising administering to a subject in need thereof a therapeutically effective amount of the product of Claim 52.

5 57. A method as in Claim 56, wherein the disorder is epilepsy.

58. Crystalline form I-SA of the compound of formula (I-S)



59. The crystalline form I-SA of claim 58 wherein crystalline form I-SA has
10 the following powder X-ray diffraction peaks, as defined to position and d-spacing:

Position ($^{\circ}2\theta$)	d-spacing ($\text{\AA}$)
4.44	19.92
15.50	5.72
17.32	5.12
18.57	4.78
19.39	4.58
19.86	4.47
20.03	4.43
20.88	4.26
21.57	4.12
21.93	4.05
22.71	3.92
23.19	3.84
23.90	3.72
24.53	3.63
25.02	3.56
26.04	3.42
26.71	3.34
26.84	3.32
28.28	3.16
29.96	2.98
30.70	2.91