


WHAT IS CLAIMED IS:

1. A compound having the following formula (I):



I

or a stereoisomer or pharmaceutically-acceptable salt thereof, wherein

R^1 is C_{1-3} alkyl substituted by 0-7 deuterium atoms;

R^2 is methyl, ethyl, propyl, furyl, pyranyl, cyclopropyl, cyclobutyl or cyclohexyl, cyclopentyl, pyridyl, pyrimidinyl, pyridazinyl, pyrazinyl, quinolinyl or pyrrolopyridinyl,

- 10 each group substituted as valence allows by 0-4 groups selected from R^{2a} ;

R^{2a} at each occurrence is independently hydrogen, =O, halo, OCF_3 , CN, NO_2 ,


$-(CH_2)_rOR^b$, $-(CH_2)_rSR^b$, $-(CH_2)_rC(O)R^b$, $-(CH_2)_rC(O)OR^b$, $-(CH_2)_rOC(O)R^b$,

$(CH_2)_rNR^{11}R^{11}$, $-(CH_2)_rC(O)NR^{11}R^{11}$, $-(CH_2)_rNR^bC(O)R^c$, $-(CH_2)_rNR^bC(O)OR^c$,

$-NR^bC(O)NR^{11}R^{11}$, $-S(O)_pNR^{11}R^{11}$, $-NR^bS(O)_pR^c$, $-S(O)_pR^c$, C_{1-6} alkyl substituted with

- 15 0-3 R^a , C_{1-6} haloalkyl, $-(CH_2)_r$ -3-14 membered carbocycle substituted with 0-1 R^a or

$a-(CH_2)_r$ -5-7 membered heterocycle comprising carbon atoms or 1-4 heteroatoms selected from N, O, and $S(O)_p$ substituted with 0-2 R^a ;



R^3 has the formula,

R^{3aa} is C_{1-6} alkyl substituted with 0-3 R^{a2} , $S(O)_pR^{c2}$ or OR^{b2} ;

- 20 R^{3ab} , R^{3ac} , or R^{3ad} are independently hydrogen, pyrazolyl, thiazolyl, pyrimidinyl or oxadiazolyl, each group substituted with 0-3 R^{a2} ;

R^{a2} is at each occurrence independently halo, OH or C_{1-6} alkyl substituted with 0-3 R^{12} ;

R^{b2} is hydrogen or C_{1-6} alkyl substituted with 0-2 R^{d2}

- 25 R^{c2} is C_{1-6} alkyl substituted with 0-3 R^{12} ;

R^{d2} independently at each occurrence is F or OH;

R^{12} is halo, CN or OH;

- R^4 and R^5 are independently hydrogen;
 R^{11} at each occurrence independently is hydrogen;
 R^a at each occurrence is hydrogen, F, Cl, Br, OCF_3 , CF_3 , CHF_2 , CN, NO_2 ,
 $-(CH_2)_rOR^b$, $-(CH_2)_rSR^b$, $-(CH_2)_rC(O)R^b$, $-(CH_2)_rC(O)OR^b$, $-(CH_2)_rOC(O)R^b$,
5 $-(CH_2)_rNR^{11}R^{11}$, $-(CH_2)_rC(O)NR^{11}R^{11}$, $-(CH_2)_rNR^bC(O)R^c$, $-(CH_2)_rNR^bC(O)OR^c$,
 $NR^bC(O)NR^{11}R^{11}$, $-S(O)_pNR^{11}R^{11}$, $NR^bS(O)_pR^c$, $-S(O)R^c$, $-S(O)_2R^c$, C_{1-6} alkyl
substituted with 0-3 R^f , C_{1-6} haloalkyl, $-(CH_2)_r$ -3-14 membered carbocycle, or $-(CH_2)_r$ -5-7
membered heterocycle comprising carbon atoms and 1-4 heteroatoms selected from N, O,
and $S(O)_p$ substituted with 0-3 R^f ;
10 R^b at each occurrence is hydrogen, C_{1-6} alkyl substituted with 0-3 R^d , C_{1-6}
haloalkyl, C_{3-6} cycloalkyl substituted with 0-2 R^d , or $-(CH_2)_r$ -5-7 membered heterocycle
comprising carbon atoms and 1-4 heteroatoms selected from N, O, and $S(O)_p$ substituted
with 0-3 R^f , or $(CH_2)_r$ -phenyl substituted with 0-3 R^d ;
 R^c is C_{1-6} alkyl substituted with 0-3 R^f , $(CH_2)_r$ - C_{3-6} cycloalkyl substituted with 0-3
15 R^f or $(CH_2)_r$ -phenyl substituted with 0-3 R^f ;
 R^d at each occurrence is independently hydrogen, F, Cl, Br, OCF_3 , CF_3 , CN, NO_2 ,
 $-OR^c$, $-(CH_2)_rC(O)R^c$, $-NR^cR^c$, $-NR^cC(O)OR^c$, C_{1-6} alkyl or $(CH_2)_r$ -phenyl substituted
with 0-3 R^f ;
 R^e at each occurrence is independently selected from hydrogen, C_{1-6} alkyl,
20 C_{3-6} cycloalkyl and $(CH_2)_r$ -phenyl substituted with 0-3 R^f ;
 R^f independently at each occurrence is hydrogen, halo, CN, NH_2 , OH, C_{3-6}
cycloalkyl, CF_3 , $O(C_{1-6}alkyl)$ or a $-(CH_2)_r$ -5-7 membered heteroaryl comprising carbon
atoms and 1-4 heteroatoms selected from N, O, and $S(O)_p$;
p is 0, 1, or 2; and
25 r is 0, 1, 2, 3, or 4.


2. The compound according to claim 1, or a stereoisomer or
pharmaceutically-acceptable salt thereof, wherein R^1 is CH_3 , C_2H_5 , CD_3 or CD_2CD_3 .
30 3. A compound according to any one of claims 1-2, or a stereoisomer or
pharmaceutically-acceptable salt thereof, wherein R^2 is cyclopropyl substituted by 0-4
groups selected from R^{2a} .

4. The compound according to any one of claims 1-3, or a stereoisomer or pharmaceutically-acceptable salt thereof, wherein R^{3aa} is $S(O)_2CH_3$ or OCH_3 .

5. The compound according to any one of claims 1-4, or a stereoisomer or pharmaceutically-acceptable salt thereof, wherein both R^4 and R^5 are hydrogen.

6. The compound of any one of claims 1-5, or a stereoisomer or pharmaceutically-acceptable salt thereof, having the formula:

10



7. A pharmaceutical composition comprising one or more compounds according to any one of claims 1-6 and a pharmaceutically acceptable carrier or diluent.