

COMPOUNDS POSSESSING NEURONAL ACTIVITY

TECHNICAL FIELD OF THE INVENTION

The present invention relates to compounds, methods and pharmaceutical compositions which possess neuronal activity. The compounds and compositions that can be used either alone or in conjunction with a neurotrophic factor, such as nerve growth factor, in methods for treating or preventing neuronal damage caused by disease or physical trauma.

BACKGROUND OF THE INVENTION

Neurological diseases are associated with the death of or injury to neuronal cells. For example, the loss of dopaminergic neurons in the substantia nigra is the basis for Parkinson's disease. Although the molecular mechanism of neurodegeneration in Alzheimer's disease is yet to be established, inflammation and deposition of beta-amyloid protein and other such agents may compromise neuronal function or survival. In patients suffering from brain ischemia or spinal cord injuries, extensive neuronal cell death is observed. Currently, there are no satisfactory treatments for these diseases.

One approach to treating neurological diseases involves the use of drugs capable of inhibiting neuronal cell death. A more recent approach involves the promotion of nerve regeneration by drugs which stimulate neurite
5 outgrowth.

Neurite outgrowth may be stimulated *in vitro* by nerve growth factors, such as NGF. For example, Glial Cell Line-Derived Neurotrophic Factor (GDNF) demonstrates neurotrophic activity both, *in vivo* and *in vitro*, and is
10 currently being investigated for the treatment of Parkinson's disease. Insulin and Insulin-like growth factors have been shown to stimulate growth of neurites in rat pheochromocytoma PC12 cells and in cultured sympathetic and sensory neurons [Recio-Pinto et al., J.
15 Neurosci., 6, pp. 1211-1219 (1986)]. Insulin and Insulin-like growth factors also stimulate the regeneration of injured motor nerves *in vivo* and *in vitro* [Near et al., PNAS, pp. 89, 11716-11720 (1992); and Edbladh et al., Brain Res., 641, pp. 76-82 (1994)].
20 Similarly, fibroblast growth factor (FGF) stimulates neural proliferation [D. Gospodarowicz et al., Cell Differ., 19, p. 1 (1986)] and growth [M. A. Walter et al., Lymphokine Cytokine Res., 12, p. 135 (1993)].

There are, however, several disadvantages
25 associated with the use of nerve growth factors for treating neurological diseases. They do not readily cross the blood-brain barrier. They are unstable in plasma. And they have poor drug delivery properties.

Recently, small molecules have been shown to
30 stimulate axonal outgrowth *in vivo*. In individuals suffering from a neurological disease, this stimulation of neurite outgrowth may protect neurons from further degeneration, and accelerate the regeneration of nerve cells. For example, estrogen has been shown to promote

the growth of axons and dendrites, which are neurites sent out by nerve cells to communicate with each other in a developing or injured adult brain [(C. Dominique Toran-Allerand et al., J. Steroid Biochem. Mol. Biol., 56, pp. 169-78 (1996); and B. S. McEwen et al., Brain Res. Dev. Brain. Res., 87, pp. 91-95 (1995)]. The progress of Alzheimer's disease may be slowed in women who take estrogen. Estrogen is hypothesized to complement NGF and other neurotrophins and thereby help neurons differentiate and survive.

It has been reported that compounds with an affinity for the FK506 binding protein (FKBP) that inhibit that protein's rotamase activity also possess nerve growth stimulatory activity. [Lyons et al., PNAS, 91, pp. 3191-3195 (1994)]. Many of these such compounds also have immunosuppressive activity.

FK506 (Tacrolimus), an immunosuppressive drug, has been demonstrated to act synergistically with NGF in stimulating neurite outgrowth in PC12 cells as well as sensory ganglia [Lyons et al. (1994)]. This compound has also been shown to be neuroprotective in focal cerebral ischemia [J. Sharkey and S. P. Butcher, Nature, 371, pp. 336-339 (1994)] and to increase the rate of axonal regeneration in injured sciatic nerve [B. Gold et al., J. Neurosci., 15, pp. 7509-16 (1995)].

The use of immunosuppressive compounds, however, has obvious drawbacks. In addition to compromising immune function, prolonged treatment with these compounds can cause nephrotoxicity [Kopp et al., J. Am. Soc. Nephrol., 1, p. 162 (1991)], neurological deficits [P.C. DeGroen et al., N. Eng. J. Med., 317, p. 861 (1987)] and vascular hypertension [Kahan et al., N. Eng. J. Med., 321, p. 1725 (1989)].

More recently, sub-classes of FKBP binding compounds which inhibit rotamase activity, but which purportedly lack immunosuppressive activity have been disclosed for use in stimulating nerve growth [see United States patents 5,614,547; 5,696,135; WO 96/40633; WO 96/40140; WO 97/16190; J. P. Steiner et al., Proc. Natl. Acad. Sci. USA , 94, pp. 2019-23 (1997); and G.S. Hamilton et al., Bioorg. Med. Chem. Lett., 7, pp. 1785-90 (1997)]. While these compounds supposedly avoid certain unwanted side effects of immunosuppressive FKBP binding compounds, they still bind to FKBP and inhibit its rotamase activity. This latter property may still lead to undesirable side effects due to other roles FKBP may play in mammals.

Surprisingly, it is now known that binding to FKBP is not necessary for neuronal activity. Co-pending United States patent application Serial Nos. 08/748,447, 08/748,448 and 08/749,114 each describe the use of non-FKBP binding, non-immunosuppressive compounds for stimulating nerve growth and preventing neurodegeneration. Due to their lack of affinity for FKBP, these compounds advantageously avoid any potential interference with FKBP-associated biochemical pathways. These compounds do, however, inhibit multi-drug resistance ("MDR") through inhibition of the p-glycoprotein and MRP. While it appears that the dosages of those compounds necessary to stimulate nerve growth and prevent neurodegeneration are lower than those that effect MDR, it would still be desirable to obtain compounds which are specific for neuronal activity, without other significant mechanisms of action.

Though a wide variety of neurological degenerative disorders may be treated by stimulating neurite outgrowth, there are relatively few agents known

to possess these properties. Moreover, the newer non-immunosuppressive compounds have only recently begun being tested in living organisms. Thus, there remains a need for new pharmaceutically acceptable compounds and compositions
5 that have the ability to stimulate neurite outgrowth and prevent neurodegeneration in patients without causing immunosuppression, without interfering with FKBP and without affecting cellular pumps, such as p-glycoprotein or MRP.


International Publication WO 92/21313 discloses a
10 genus of compounds that have an affinity for FK-506 binding protein and immunosuppressive activity. International Publications WO 94/07858 and WO 95/26337 each disclose a genera of compounds possessing multi-drug reversal activity. J. Med. Chem. 12 (1969), pp. 677-680 discloses synthesized
15 compounds and studies on the inhibition of sheep red blood cells lysis by hemolysin and complement from the guinea pig. Anal. Biochem. 129 (1983), pp. 502-512 discloses the synthesis of a fluorescent substrate for porcine pepsin.

SUMMARY OF THE INVENTION


20 Applicants have identified several subclasses of compounds that do not bind FKBP, do not inhibit MDR, but have potent neuronal activity.

The term "neuronal activity", as used herein, includes stimulation of damaged neurons, promotion of
25 neuronal regeneration, prevention of neurodegeneration and treatment of a neurological disorder. The compounds of this invention have activity in both peripheral nerves and the central nervous system.

Two of these subgenera fall within the genus of
30 compounds described in WO 92/21313. These compounds are characterized by the formulae:



Two other subgenera of compounds of this invention have the formulae:



The invention also includes pharmaceutically acceptable derivatives of any of compounds (I) through (IV).

In each of these compounds, A and B are independently selected from hydrogen, Ar, (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, (C₅-C₇)-cycloalkyl substituted-(C₁-C₆)-straight or branched alkyl, (C₅-C₇)-cycloalkyl substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl, (C₅-C₇)-cycloalkenyl substituted-(C₁-C₆)-straight or branched alkyl, (C₅-C₇)-cycloalkenyl substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl, Ar-substituted-(C₁-C₆)-straight or branched alkyl, or Ar-substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl; wherein any one of the CH₂ groups of said alkynyl, alkenyl or alkyl chains in A or B is optionally replaced by O, S, S(O), S(O)₂ or N(R); wherein

R is selected from hydrogen, (C₁-C₆)-straight or branched alkyl, or (C₂-C₆)-straight or branched alkenyl or alkynyl;

Ar is selected from phenyl, 1-naphthyl, 2-naphthyl, indenyl, azulenyl, fluorenyl, anthracenyl, 2-furyl, 3-furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, 2-pyrazolinyl, pyrazolidinyl, isoxazolyl, isothiazolyl, 1,2,3-oxadiazolyl, 1,2,3-triazolyl, 1,2,3-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,4-triazolyl, 1,2,4-oxadiazolyl, 1,2,4-thiadiazolyl, benzoxazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, 1,3,5-triazinyl, 1,3,5-trithianyl, indolizinyl, indolyl, isoindolyl, 3H-indolyl, indolinyl, benzo[b]furanyl, benzo[b]thiophenyl, 1H-indazolyl, benzimidazolyl, benzthiazolyl, purinyl, 4H-quinolizinyl, quinolinyl, 1,2,3,4-tetrahydro-isoquinolinyl, isoquinolinyl, 1,2,3,4-tetrahydro-isoquinolinyl, cinnolinyl, phthalazinyl, quinazolinyl, quinoxalinyl, 1,8-naphthyridinyl, peridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl or phenoxazinyl or other chemically feasible monocyclic, bicyclic or tricyclic ring systems, wherein each ring consists of 5 to 7 ring atoms and wherein each ring comprises 0 to 3 heteroatoms independently selected from N, N(R), O, S, S(O), or S(O)₂ and wherein

each Ar is optionally substituted with one to three substituents independently selected from halogen, hydroxyl, nitro, -SO₃H, trifluoromethyl, trifluoromethoxy, (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl, O-[(C₁-C₆)-straight or branched alkyl], O-[(C₂-C₆)-straight or branched alkenyl], O-benzyl, O-phenyl, 1,2-

methylenedioxy, $-N(R^1)(R^2)$, carboxyl, $N-(C_1-C_5-$
straight or branched alkyl or C_2-C_5 -straight or
branched alkenyl) carboxamides, N,N -di- $(C_1-C_5-$
5 straight or branched alkyl or C_2-C_5 -straight or
branched alkenyl) carboxamides, $N-(C_1-C_5$ -straight or
branched alkyl or C_2-C_5 -straight or branched alkenyl)
sulfonamides, N,N -di- $(C_1-C_5$ -straight or branched
alkyl or C_2-C_5 -straight or branched alkenyl)
sulfonamides, morpholinyl, piperidinyl, $O-Z$, CH_2-
10 $(CH_2)_q-Z$, $O-(CH_2)_q-Z$, $(CH_2)_q-Z-O-Z$, or $CH=CH-Z$;

wherein R^1 and R^2 are independently
selected from (C_1-C_6) -straight or branched
alkyl, (C_2-C_6) -straight or branched alkenyl or
alkynyl, hydrogen or benzyl; or wherein R_1 and
15 R_2 are taken together with the nitrogen atom to
which they are bound to form a 5-7 membered
heterocyclic ring;

Z is selected from 4-methoxyphenyl, 2-
pyridyl, 3-pyridyl, 4-pyridyl, pyrazyl,
20 quinolyl, 3,5-dimethylisoxazolyl, isoxazolyl, 2-
methylthiazolyl, thiazolyl, 2-thienyl, 3-thienyl,
or pyrimidyl; and

q is 0, 1 or 2;

X is N, O or CH;

25 when X is N or CH, Y is selected from hydrogen, Ar,
 (C_1-C_6) -straight or branched alkyl, (C_2-C_6) -straight or
branched alkenyl or alkynyl, (C_5-C_7) -cycloalkyl-
substituted- (C_1-C_6) -straight or branched alkyl, (C_5-C_7) -
cycloalkyl-substituted- (C_2-C_6) -straight or branched
30 alkenyl or alkynyl, (C_5-C_7) -cycloalkenyl-substituted- $(C_1-$
 $C_6)$ -straight or branched alkyl, (C_5-C_7) -cycloalkenyl-
substituted- (C_2-C_6) -straight or branched alkenyl of
alkynyl, Ar-substituted- (C_1-C_6) -straight or branched

alkyl, or Ar-substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl;

when X is O, Y is a lone pair of electrons;

n is 0, 1 or 2;

5 m is 0, 1 or 2;

n + m is less than 4 and greater than 0;

the indicated ring in formulae II and IV is saturated, partially unsaturated or unsaturated;

1 to 2 carbon atoms in the ring in formulae II and
10 IV are optionally replaced with a heteroatom independently selected from O, S, S(O), S(O)₂ or NR; and

said ring in formula II and IV is optionally benzofused.

J is selected from hydrogen, (C₁-C₆)-straight or
15 branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, Ar-substituted-(C₁-C₆)-straight or branched alkyl, Ar-substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl, or cyclohexylmethyl;

K is selected from (C₁-C₆)-straight or branched
20 alkyl, Ar-substituted-(C₁-C₆)-straight or branched alkyl, (C₁-C₆)-straight or branched alkenyl or alkynyl, Ar-substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl, or cyclohexylmethyl; wherein any one of the CH₂ groups of said alkyl, alkenyl or alkynyl chains in K is
25 optionally replaced by O, S, S(O), S(O)₂ or N(R);

D is selected from is selected from Ar, (C₁-C₆) straight or branched alkyl, (C₂-C₆) straight or branched alkenyl or alkynyl, (C₅-C₇) cycloalkyl substituted (C₁-C₆) straight or branched alkyl, (C₅-C₇) cycloalkyl substituted
30 (C₂-C₆) straight or branched alkenyl or alkynyl, (C₅-C₇) cycloalkenyl substituted (C₁-C₆) straight or branched alkyl, (C₅-C₇) cycloalkenyl substituted (C₂-C₆) straight or branched alkenyl of alkynyl, Ar-substituted (C₁-C₆)

straight or branched alkyl, or Ar-substituted (C₂-C₆)
straight or branched alkenyl or alkynyl;

wherein any one of the CH₂ groups of said alkyl
chains in D other than the one that is directly bound to
5 SO₂ in the compound, is optionally replaced by O, S, SO,
SO₂ or NR.

R³ is (C₁-C₆)-straight or branched alkyl, Ar-
substituted-(C₁-C₆)-straight or branched alkyl, (C₂-C₆)-
straight or branched alkenyl or alkynyl, or Ar-
10 substituted-(C₂-C₆)-straight or branched alkenyl or
alkynyl; wherein any one of the CH₂ groups of said alkyl,
alkenyl or alkynyl chains in R³ is optionally replaced by
O, S, S(O), S(O)₂, or N(R); and wherein any one of the CH₂
groups of said alkyl, alkenyl or alkynyl chains in R³
15 except the CH₂ group bound to nitrogen, is optionally
replaced with C(O), with the proviso that in formula II,
when n is 0 and m is 1, the second CH₂ group in the alkyl
chain of R³ is not replaced with C(O).

The compositions disclosed herein comprise a
20 compound of this invention, a carrier, and optionally, a
neuronal growth factor.

The methods of stimulating nerve growth and
preventing neurodegeneration disclosed herein employ the
above compounds either alone or in combination with a
25 neuronal growth factor. The methods are useful in
treating or preventing nerve damage caused by various
neurological diseases and physical traumas and also in ex
vivo nerve regeneration.

DETAILED DESCRIPTION OF THE INVENTION

30 Applicants have discovered diverse genera of
compounds with neuronal activity which do not bind to
FKBP, and which do not have multi-drug resistance
reversal activity. Without being bound by theory,

applicants believe that the compounds disclosed in this application exert their neuronal activity by increasing cytoplasmic Ca^{2+} concentrations. This is likely achieved by interaction, either direct or indirect, with a calcium release channel, such as the ryanodine receptor or the inositol 1,4,5-trisphosphate receptor, in the endoplasmic reticulum of the nerve cell.

Thus, according to one embodiment, the invention provides a method of stimulating nerve growth or preventing neurodegeneration by contacting nerve cells with a compound that:

- a. increases cytoplasmic Ca^{2+} concentration or binds to the ryanodine receptor;
- b. does not bind to FKBP; and
- c. does not possess MDR reversal activity.

According to a related embodiment, the present invention provides a compound that:

- a. has neuronal activity;
- b. increases cytoplasmic Ca^{2+} concentration or binds to the ryanodine receptor;
- c. does not bind to FKBP; and
- d. does not possess MDR reversal activity.

The term "increases cytoplasmic Ca^{2+} concentration", as used herein means a detectable increase in channel current recorded in the single channel recording assay described below in the presence of such a compound as compared to an appropriate control. Alternatively, the term "increases cytoplasmic Ca^{2+} concentration", as used herein means a detectable shift in the fluorescence spectrum in the cell assay described herein.

The term "binds to the ryanodine receptor", as used herein, means that the compound specifically

competes with ryanodine for binding to microsomes in the assay described below.

The term "does not bind FKBP", as used herein means that the compound demonstrates a K_i of 10 μM or greater in at least one of the rotamase inhibitory assays described below.

The term "does not possess MDR reversal activity", as used herein means that at a concentration of 2.5 μM , the compound has an MDR ratio of less than 7.0, and preferably less than 3.0 in at least one of the MDR assays described below.

Single-channel recording experiments are useful to determine if the compounds of this invention cause the requisite increase in cytoplasmic Ca^{2+} concentration. These experiments are conducted as described in E. Kaftan et al., Circulation Research, 78, pp. 990-997 (1996), the disclosure of which is herein incorporated by reference. Single channel recordings are conducted under voltage clamp conditions with a pair of Ag/AgCl electrodes contacting the solutions via CsCl junctions. Vesicles are added to the *cis* chamber and fused with planar lipid bilayers composed of phosphatidylethanolamine/phosphatidylcholine (3:1, 30 mg/ml in decane, Avanti Polar Lipids). The *trans* chamber contains 250 mM HEPES and 53 mM $\text{Ba}(\text{OH})_2$, pH 7.35; the *cis* chamber contains 250 mM HEPES-Tris pH 7.35. Compounds dissolved in methanol are added to the *cis* chamber. Channel currents are

amplified using a bilayer clamp amplifier (BC-525A, Warner Instruments) and recorded on VHS tape (Dagen Corp.). Data are filtered to an eight-pole Bessel filter (Frequency Devices) to 500 Hz, digitized at 2 kHz, transferred to a personal computer, and analyzed with pClamp version 6.0 (Axon Instruments). Single channel recordings are done at least 3 times for each compound condition.

Ryanodine binding may be measured by incubating microsomal protein with ^3H -ryanodine in buffer containing 36 mM Tris pH 7.2 and 50 mM KCl in the absence or presence of test compounds. Controls for maximum binding were done in the presence of 0.72 mM ATP and 36 μM CaCl_2 . Nonspecific binding was measured in the presence of 25 μM unlabelled ryanodine. Binding reactions were incubated for 2 hours at room temperature, and then centrifuged for 15 minutes at 30,000 x g. The pellets were solubilized and the radioactivity was measured by scintillation counting.

Alternatively, the flux of cytoplasmic Ca^{2+} into the cell can be followed fluorescently. For example, neuronal cells can be incubated with NGF and a calcium binding fluorescent dye, such as Fura-2, in a calcium-containing buffer. Cells are imaged continuously both before and after the addition of a test compound of this invention. The difference in fluorescent intensity before and after the addition of compounds is then plotted as a ration of fluorescence units at 340 nm and 380 nm.

Testing a compound of this invention to confirm that it binds to FKBP12 with a K_i of 10 μM or higher may be achieved using several assays known in the art. In particular, those compounds may be assayed for their ability (or lack thereof) to inhibit rotamase. Examples


of assays that measure inhibition of FKBP12 rotamase activity are those in which the isomerization of an artificial substrate -- N-succinyl-Ala-Ala-Pro-Phe-p-nitroanilide -- is followed spectrophotometrically [M. W. 5 Harding et al., Nature, 341, pp. 758-60 (1989); by J. J. Siekierka et al., Nature, 341, pp. 755-57 (1989); and S. T. Park et al., J. Biol. Chem., 267, pp. 3316-24 (1992)]. The assay includes the cis form of the substrate, FKBP12, the compound to be tested and chymotrypsin. Chymotrypsin 10 is able to cleave p-nitroanilide from the trans form of the substrate, but not the cis form. Release of p-nitroanilide is measured.

Other FKBP binding assays include a competitive LH20 binding assay using labeled FK-506 as a reporting 15 ligand. These have been described by M. W. Harding et al., Nature, 341, pp. 758-60 (1989) and by J. J. Siekierka et al., Nature, 341, pp. 755-57 (1989).

To determine whether a compound according to this invention has the requisite MDR ratio below 7.0, any 20 of the assays described in United States patents 5,543,423, 5,717,092, 5,726,184 or 5,744,485, the disclosures of which are herein incorporated by reference, may be utilized.

In particular, cell lines which are known to be 25 resistant to a particular drug are employed. These cell lines include, but are not limited to, the L1210, P388D, HL60 and MCF7 cell lines. Alternatively, resistant cell lines may be developed. The cell line is exposed to the drug to which it is resistant, or to the test compound; 30 cell viability is then measured and compared to the viability of cells which are exposed to the drug in the presence of the test compound ("MDR ratio").

According to one embodiment, the invention provides a compound of the formula:



and pharmaceutically acceptable derivatives thereof
wherein:

A and B are independently selected from hydrogen,
5 Ar, (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or
branched alkenyl or alkynyl, (C₅-C₇)-cycloalkyl
substituted-(C₁-C₆)-straight or branched alkyl, (C₅-C₇)-
cycloalkyl substituted-(C₂-C₆)-straight or branched
10 alkenyl or alkynyl, (C₅-C₇)-cycloalkenyl substituted-(C₁-
C₆)-straight or branched alkyl, (C₅-C₇)-cycloalkenyl
substituted-(C₂-C₆)-straight or branched alkenyl or
alkynyl, Ar-substituted-(C₁-C₆)-straight or branched
alkyl, or Ar-substituted-(C₂-C₆)-straight or branched
15 alkenyl or alkynyl; wherein any one of the CH₂ groups of
said alkyl, alkenyl or alkynyl chains in A or B is
optionally replaced by O, S, S(O), S(O)₂, or N(R); wherein

R is selected from hydrogen, (C₁-C₆)-straight or
branched alkyl, or (C₂-C₆)-straight or branched alkenyl or
alkynyl;

20 Ar is selected from phenyl, 1-naphthyl, 2-naphthyl,
indenyl, azulenyl, fluorenyl, anthracenyl, 2-furyl, 3-
furyl, 2-thienyl, 3-thienyl, 2-pyridyl, 3-pyridyl, 4-
pyridyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl,
pyrazolyl, 2-pyrazolyl, pyrazolidinyl, isoxazolyl,
25 isothiazolyl, 1,2,3-oxadiazolyl, 1,2,3-triazolyl, 1,3,4-
thiadiazolyl, 1,2,4-triazolyl, 1,2,4-oxadiazolyl, 1,2,4-
thiadiazolyl, 1,2,3-thiadiazolyl, benzoxazolyl,
pyridazinyl, pyrimidinyl, pyrazinyl, 1,3,5-triazinyl,
1,3,5-trithianyl, indolizinyl, indolyl, isoindolyl, 3H-
30 indolyl, indolinyl, benzo[b]furanyl, benzo[b]thiophenyl,

1H-indazolyl, benzimidazolyl, benzthiazolyl, purinyl, 4H-quinoliziny, quinolinyl, 1,2,3,4-tetrahydro-isoquinolinyl, isoquinolinyl, 1,2,3,4-tetrahydro-isoquinolinyl, cinnolinyl, phthalazinyl, quinazolinyl, 5 quinoxalinyl, 1,8-naphthyridinyl, peridinyl, carbazolyl, acridinyl, phenazinyl, phenothiazinyl or phenoxazinyl or other chemically feasible monocyclic, bicyclic or tricyclic ring systems, wherein each ring consists of 5 to 7 ring atoms and wherein each ring comprises 0 to 3 10 heteroatoms independently selected from N, N(R), O, S, S(O), or S(O)₂ and wherein

each Ar is optionally substituted with one to three substituents independently selected from halogen, hydroxyl, nitro, -SO₃H, trifluoromethyl, trifluoromethoxy, 15 (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl, O-[(C₁-C₆)-straight or branched alkyl], O-[(C₂-C₆)-straight or branched alkenyl], O-benzyl, O-phenyl, 1,2-methylenedioxy, -N(R¹)(R²), carboxyl, N-(C₁-C₅-straight or branched alkyl or C₂-C₅-straight or branched 20 alkenyl) carboxamides, N,N-di-(C₁-C₅-straight or branched alkyl or C₂-C₅-straight or branched alkenyl) carboxamides, N-(C₁-C₅-straight or branched alkyl or C₂-C₅-straight or branched alkenyl) sulfonamides, N,N-di-(C₁-C₅-straight or branched alkyl or C₂-C₅-straight or branched alkenyl) 25 sulfonamides, morpholinyl, piperidinyl, O-Z, CH₂-(CH₂)_q-Z, O-(CH₂)_q-Z, (CH₂)_q-Z-O-Z, or CH=CH-Z;

wherein R¹ and R² are independently selected from (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, hydrogen or 30 benzyl; or wherein R₁ and R₂ are taken together with the nitrogen atom to which they are bound to form a 5-7 membered heterocyclic ring;

Z is selected from 4-methoxyphenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, pyrazyl, quinolyl, 3,5-

dimethylisoxazoyl, isoxazoyl, 2-methylthiazoyl, thiazoyl, 2-thienyl, 3-thienyl, or pyrimidyl; and

q is 0, 1 or 2;

X is N, O or C(R);

- 5 wherein when X is N or C(R), Y is selected from hydrogen, Ar, (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, (C₅-C₇)-cycloalkyl-substituted-(C₁-C₆)-straight or branched alkyl, (C₅-C₇)-cycloalkyl-substituted-(C₂-C₆)-straight or branched
10 alkenyl or alkynyl, (C₅-C₇)-cycloalkenyl-substituted-(C₁-C₆)-straight or branched alkyl, (C₅-C₇)-cycloalkenyl-substituted-(C₂-C₆)-straight or branched alkenyl of alkynyl, Ar-substituted-(C₁-C₆)-straight or branched alkyl, or Ar-substituted-(C₂-C₆)-straight or branched
15 alkenyl or alkynyl;

when X is O, Y is a lone pair of electrons;

- K is selected from (C₁-C₆)-straight or branched alkyl, Ar-substituted-(C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, Ar-substituted-(C₂-C₆)-straight or branched alkenyl or
20 alkynyl, or cyclohexylmethyl; wherein any one of the CH₂ groups of said alkyl, alkenyl or alkynyl chains in K is optionally replaced by O, S, S(O), S(O)₂ or N(R);

n is 0, 1 or 2;

- 25 J is selected from hydrogen, (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, Ar-substituted-(C₁-C₆)-straight or branched alkyl, Ar-substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl, or cyclohexylmethyl; and

- 30 D is selected from is selected from Ar, (C₁-C₆) straight or branched alkyl, (C₂-C₆) straight or branched alkenyl or alkynyl, (C₅-C₇) cycloalkyl substituted (C₁-C₆) straight or branched alkyl, (C₅-C₇) cycloalkyl substituted (C₂-C₆) straight or branched alkenyl or alkynyl, (C₅-C₇)

cycloalkenyl substituted (C_1-C_6) straight or branched alkyl, (C_5-C_7) cycloalkenyl substituted (C_2-C_6) straight or branched alkenyl or alkynyl, Ar-substituted (C_1-C_6) straight or branched alkyl, or Ar-substituted (C_2-C_6) straight or branched alkenyl or alkynyl; wherein any one of the CH_2 groups of said alkyl chains in D other than the one that is directly bound to SO_2 in the compound, is optionally replaced by O, S, SO, SO_2 or NR.

In one preferred embodiment of formula I, A and B are not simultaneously hydrogen. Even more preferred is when at least one of A or B is a (C_1-C_6)-straight alkyl terminally substituted with Ar (which itself is substituted or unsubstituted). More preferably, at least one of A or B is a (C_1-C_6)-straight alkyl terminally substituted with pyridine (which itself substituted or unsubstituted).

According to another preferred embodiment of formula I, X is nitrogen or oxygen.

In another preferred embodiment of formula I, J is a (C_1-C_3)-straight alkyl.

In yet another preferred embodiment of formula I, K is an Ar-substituted (C_1-C_3)-straight alkyl. More preferred is when K is a (C_1-C_3)-straight alkyl terminally substituted with an unsubstituted phenyl.

According to another preferred embodiment of formula I, D is selected from (C_1-C_6) straight or branched alkyl, Ar, Ar-substituted (C_1-C_6) straight or branched alkyl, Ar-substituted (C_2-C_6) straight or branched alkenyl or alkynyl, or $-CH_2-S(O)_2-(C_1-C_4)$ straight or branched alkyl.

More preferably, D is selected from aminophenyl, nitrophenyl, isopropyl, benzyl, fluorophenyl, cyanophenyl, methoxyphenyl,


dimethoxyphenyl, methylsulfonylmethyl, ethylenephenyl, dinitroanilinophenyl, N,N-dimethylaminophenylazophenyl N,N-dimethylaminonaphthyl or acetamidophenyl.

Most preferably, D is selected from 4-aminophenyl, 2-nitrophenyl, methylsulfonylmethyl, benzyl, ethylenephenyl, 4-fluorophenyl, 4-cyanophenyl, 3,4-dimethoxyphenyl, 4-methoxyphenyl, 4-(2,4-dinitroanilino)phenyl, 4-((4-(N,N-dimethylamino)phenyl)azo)phenyl, 5-(N,N-dimethylamino)naphthyl or 4-acetamidophenyl.

According to another preferred embodiment, Y is methyl.

In another preferred embodiment, n is 0.

According to another embodiment, the invention provides a compound of the formula:.



and pharmaceutically acceptable derivatives thereof, wherein A, B, Y, D, n and subcomponents thereof are as defined above;

m is 0, 1 or 2;

n + m is less than 4 and greater than 0;
the indicated ring in formula II is saturated,
partially unsaturated or unsaturated;

1 to 2 carbon atoms in the ring in formula II are
5 optionally replaced with a heteroatom independently
selected from O, S, S(O), S(O)₂ or NR; and

said ring in formula II is optionally benzofused.

Preferably, in formula II, n is 0, m is 2 and
the ring is fully saturated.

10 According to another preferred embodiment of
formula II one carbon in said ring is optionally replaced
with a heteroatom selected from O, S, SO or SO₂.

Preferred embodiments for the individual
components in the compounds of formula II are the same as
15 those set forth above for compounds of formula I.

The most preferred compounds of formula I and II are set
forth in Tables 1 and 2, below:

Table 1. Compounds of formula I.

Cmpd #	Structure	Cmpd #	Structure
2		4	
5		6	
7		8	












Cmpd #	Structure	Cmpd #	Structure
11		12	
13		14	
15		16	
17		18	

Table 2. Compounds of formula II

Cmpd #	Structure	Cmpd #	Structure
1		3	

According to another embodiment of this invention is provided a compound of the formula:




and pharmaceutically acceptable derivatives thereof, wherein:

A, B, X, Y, K, J, n and subcomponents thereof are as defined above for compounds of formula I; and

10 R^3 is $(\text{C}_1\text{-C}_6)$ -straight or branched alkyl, Ar-substituted- $(\text{C}_1\text{-C}_6)$ -straight or branched alkyl, $(\text{C}_2\text{-C}_6)$ -straight or branched alkenyl or alkynyl, or Ar-substituted- $(\text{C}_2\text{-C}_6)$ -straight or branched alkenyl or alkynyl; wherein any one of the CH_2 groups of said alkyl, alkenyl or alkynyl chains in R^3 is optionally replaced by
 15 O, S, S(O), S(O)₂ or N(R); and wherein any one of the CH_2 groups of said alkyl, alkenyl or alkynyl in R^3 except the CH_2 group bound to nitrogen, is optionally replaced with C(O).

20 According to another embodiment, the invention provides a compound of the formula:



10


and pharmaceutically acceptable derivatives thereof;
wherein:

15 A, B, X, Y, n, m and subcomponents thereof are as
defined for compounds of formula (II); and

 R³ is as defined for compounds of formula (III),
with the proviso that in formula IV, when n is 0 and m is
1, the second CH₂ group in the alkyl, alkenyl or alkynyl
20 chain of R³ is not replaced with C(O).

 The term "second CH₂ group in the alkyl chain of
R³" refers to the CH₂ group bound immediately to the CH₂
group that is bound to nitrogen (indicated in bold in the
formula below:

25



30

 Preferably, in the compounds of formulae III
and IV, at least one of A or B is Ar-substituted (C₁-C₆)-
alkyl chain. Even more preferred is when at least one of
A or B is is a (C₁-C₆)-alkyl chain terminally substituted
with phenyl or pyridinyl.

35

 According to another preferred embodiment in
the compounds of formulae III or IV, X is N or O.

 According to yet another preferred embodiment,
in the compounds of formula III, K is an Ar-substituted
alkyl, alkenyl or alkynyl. More preferably, K is benzyl.

According to another preferred embodiment, in compounds of formula III, J is hydrogen or alkyl, preferably methyl.

In compounds of formulae III and IV, R³ is preferably hydrogen, (C₁-C₆)-alkyl, (C₁-C₆)-alkyl terminally substituted with pyridyl, or 3,4,5-trimethoxybenzoylmethyl.





In a preferred embodiment of formula III, n is 0.










In a preferred embodiment the indicated ring in formula IV is fully saturated.

In another preferred embodiment of the compounds of formula IV, m + n is 1 or 2. Even more preferred is when n is 0 and m is 1 or 2. Most preferably, n is 0 and m is 2.

The most preferred compounds of formulae III and IV are listed in the table below, as well as being described in the examples:

Table 3. Compounds of formulae III and IV.

Cmpd #	Structure	Cmpd#	Structure
101		107	
102		108	

Cmpd #	Structure	Cmpd #	Structure
103		109	
104		110	
105		111	
106		112	
113			

This invention includes all optical and racemic isomers of the compounds of formulae I through IV, as well as pharmaceutically acceptable derivatives thereof.

A "pharmaceutically acceptable derivative," as used herein denotes any pharmaceutically acceptable salt, ester, prodrug, or salt of such ester or prodrug, of a compound of this invention or any other compound which, upon administration to a patient, is capable of providing (directly or indirectly) a compound of this invention, or a metabolite or residue thereof, characterized by neuronal activity.

Surprisingly and unexpectedly, the compounds of formulae (I) through (IV) of this invention do not bind FKBP, do not inhibit its rotamase activity and are not immunosuppressive. Moreover, although the compounds of this invention bear some structural resemblance to compounds which are known to reverse multi-drug resistance (see United States patent 5,543,423, WO 95/26337 and WO 94/07858), the present compounds appear to have no activity against MDR. Thus, the presently disclosed compounds advantageously possess neuronal activity, without interfering with other pathways known to be affected by structurally similar compounds.

The nerve growth activity of the compounds of this invention may be initially assayed using several cell culture assays known in the art. For example, the compounds of this invention may be tested in a neurite outgrowth assay using pheochromocytoma PC12 cells as described by Lyons et al., PNAS, 91, pp. 3191-3195 (1994). A similar assay may be carried out in SH-SY5Y human neuroblastoma cells. Alternatively, the chick dorsal root ganglia assay described in United States patent 5,614,547 or in G. S. Hamilton et al., Bioorg. Med. Chem. Lett., (1997) and references cited therein, may be utilized.

The compounds of this invention may also be assayed for nerve growth activity *in vivo* using a mouse

model of Parkinson's disease [J. P. Steiner et al., Proc. Natl. Acad. Sci. USA, 94, pp. 2019-23 (1997), United States patent 5,721,256] or following surgical sciatic nerve crush in rats.

5 According to another embodiment, the invention provides compositions comprising a compound of any of formula (I) through (IV) and a pharmaceutically acceptable carrier. Preferably, the compositions of this invention are formulated for administration to a mammal
10 (i.e., pharmaceutical compositions).

(If pharmaceutically acceptable salts of the compounds of this invention are utilized in these compositions, those salts are preferably derived from inorganic or organic acids and bases. Included among
15 such acid salts are the following: acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate,
20 glucoheptanoate, glycerophosphate, glycolate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oxalate, palmoate,
25 pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, salicylate, succinate, sulfate, tartrate, thiocyanate, tosylate and undecanoate. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in the
30 preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable acid addition salts.

Base salts include ammonium salts, alkali metal salts, such as sodium and potassium salts, alkaline

earth metal salts, such as calcium and magnesium salts, salts with organic bases, such as dicyclohexylamine salts, N-methyl-D-glucamine, and salts with amino acids such as arginine, lysine, and N-(C₁₋₄ alkyl)₄⁺ salts.

5 Also, the basic nitrogen-containing groups can be quaternized with such agents as lower alkyl halides, such as methyl, ethyl, propyl, and butyl chloride, bromides and iodides; dialkyl sulfates, such as dimethyl, diethyl, dibutyl and diamyl sulfates, long chain halides
10 such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides, aralkyl halides, such as benzyl and phenethyl bromides and others. Water or oil-soluble or dispersible products are thereby obtained.

 The compounds utilized in the compositions and
15 methods of this invention may also be modified by appending appropriate functionalities to enhance selective biological properties. Such modifications are known in the art and include those which increase biological penetration into a given biological system
20 (e.g., blood, lymphatic system, central nervous system), increase oral availability, increase solubility to allow administration by injection, alter metabolism and alter rate of excretion.

 Pharmaceutically acceptable carriers that may
25 be used in these pharmaceutical compositions include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, self-emulsifying drug delivery systems (SEDDS) such as α -tocopherol, polyethyleneglycol 1000 succinate, or TPGS, surfactants used in
30 pharmaceutical dosage forms such as Tweens or other similar polymeric delivery matrices, serum proteins, such as human serum albumin, gelatin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty

acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, polylactic acid, polyaceticpolyglycollic acid, citric acid, cellulose-based substances, such as HPC and HPMC, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol, wool fat. Cyclodextrins such as α -, β -, and γ -cyclodextrin, or chemically modified derivatives such as hydroxyalkylcyclodextrins, including 2- and 3-hydroxypropyl- β -cyclodextrins, or other solubilized derivatives may also be advantageously used to enhance delivery of compounds of the compounds of this invention.

According to another embodiment, the pharmaceutical compositions of this invention additionally comprise a neurotrophic factor. The term "neurotrophic factor", as used herein, refers to compounds which are capable of stimulating growth or proliferation of nervous tissue. As used in this application, the term "neurotrophic factor" excludes the compounds described herein.

Numerous neurotrophic factors have been identified in the art and any of those factors may be utilized in the compositions of this invention. These

neurotrophic factors include, but are not limited to, nerve growth factor (NGF), insulin growth factor (IGF-1) and its active truncated derivatives such as gIGF-1, acidic and basic fibroblast growth factor (aFGF and bFGF, respectively), platelet-derived growth factors (PDGF), brain-derived neurotrophic factor (BDNF), ciliary neurotrophic factors (CNTF), glial cell line-derived neurotrophic factor (GDNF), neurotrophin-3 (NT-3), neurotrophin 4/5 (NT-4/5), or any of the compounds described in WO 97/36869, WO 96/41609, WO 97/16190, WO 96/40633, WO 97/18828, WO 96/40140 or WO 98/13355. The most preferred neurotrophic factor in the compositions of this invention is NGF.

The compositions of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intrahepatic, intralesional and intracranial injection or infusion techniques. Preferably, the compositions are administered orally, intraperitoneally or intravenously.

The pharmaceutical compositions of this invention may contain any conventional non-toxic pharmaceutically-acceptable carriers, adjuvants or vehicles. In some cases, the pH of the formulation may be adjusted with pharmaceutically acceptable acids, bases or buffers to enhance the stability of the formulated compound or its delivery form.

Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or

wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution
5 in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any
10 bland fixed oil may be employed including synthetic mono- or di-glycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil,
15 especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as Ph. Helv or similar alcohol.

The pharmaceutical compositions of this
20 invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers which are commonly used include lactose, corn starch, dicalcium
25 phosphate and microcrystalline cellulose (Avicel). Lubricating agents, such as magnesium stearate and talc, are also typically added. For oral administration in a capsule form, useful diluents include lactose, dried corn starch and TPGS, as well as the other diluents used in
30 tablets. For oral administration in a soft gelatin capsule form (filled with either a suspension or a solution of a compound of this invention), useful diluents include PEG400, TPGS, propylene glycol, Labrasol, Gelucire, Transcutol, PVP and potassium

acetate. When aqueous suspensions are administered orally, the active ingredient is combined with emulsifying and suspending agents, such as sodium CMC, methyl cellulose, pectin and gelatin. If desired, certain
5 sweetening and/or flavoring and/or coloring agents may be added.

Alternatively, the pharmaceutical compositions of this invention may be administered in the form of suppositories for rectal administration. These can be
10 prepared by mixing the agent with a suitable non-irritating excipient which is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax, gelatin, glycerin and
15 polyethylene glycols.

The pharmaceutical compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including
20 diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

Topical application for the lower intestinal tract can be effected in a rectal suppository formulation
25 (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

For topical applications, the pharmaceutical compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in
30 one or more carriers. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying

wax, stearic acid, cetyl stearate, cetyl alcohol, lanolin, magnesium hydroxide, kaolin and water.

Alternatively, the pharmaceutical compositions can be formulated in a suitable lotion or cream containing the
5 active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters, wax, cetyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

10 For ophthalmic use, the pharmaceutical compositions may be formulated as micronized suspensions in isotonic, pH adjusted sterile saline, or, preferably, as solutions in isotonic, pH adjusted sterile saline, either with or without a preservative such as
15 benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutical compositions may be formulated in an ointment such as petrolatum.

The pharmaceutical compositions of this invention may also be administered by nasal aerosol or
20 inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability,
(
25 fluorocarbons, and/or other conventional solubilizing or dispersing agents.

The amount of both, the compound and the neurotrophic factor (in those compositions which comprise a neurotrophic factor) that may be combined with the
30 carrier materials to produce a single dosage form will vary depending upon the host treated, the particular mode of administration. Preferably, the compositions of this invention should be formulated so that a dosage of between 0.01 - 10 mg/kg body weight/day of a compound of

this invention can be administered. More preferably, the dosage is between 0.1 - 10 mg/kg body weight/day.

In those compositions which comprise a neurotrophic factor, that factor and the compounds of this invention act synergistically to stimulate neurite outgrowth or prevent neurodegeneration. Therefore, the amount of neurotrophic factor in such compositions will be less than that required in a monotherapy utilizing only that factor. In such compositions a dosage of between 0.01 - 10 mg/kg body weight/day of the neurotrophic factor can be administered.

It should also be understood that a specific dosage and treatment regimen for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health, sex, diet, time of administration, rate of excretion, drug combination, and the judgment of the treating physician and the severity of the particular disease being treated. The amount of active ingredients will also depend upon the particular compound and neurotrophic factor in the composition.

According to another embodiment, this invention provides methods for stimulating neurite outgrowth and nerve growth and for preventing neurodegeneration. In one aspect of this embodiment, the method is used to stimulate neurite outgrowth and nerve growth and prevent neurodegeneration in a patient and is achieved by administering to the patient a pharmaceutically acceptable composition comprising any of the compounds of this invention and a pharmaceutically acceptable carrier. The amount of compound utilized in these methods is between about 0.01 and 10 mg/kg body weight/day.

This method may be used to treat nerve damage and prevent neurodegeneration caused by a wide variety of

diseases or physical traumas. These include, but are not limited to, trigeminal neuralgia, glossopharyngeal neuralgia, Bell's Palsy, myasthenia gravis, muscular dystrophy, muscle injury, progressive muscular atrophy, progressive bulbar inherited muscular atrophy, herniated, ruptured or prolapsed intervertebrae disk syndrome's, cervical spondylosis, plexus disorders, thoracic outlet destruction syndromes, peripheral neuropathies, such as those caused by lead, dapson, ticks, or porphyria, other peripheral myelin disorders, Alzheimer's disease, Gullain-Barre syndrome, Parkinson's disease and other Parkinsonian disorders, ALS, multiple sclerosis, other central myelin disorders, stroke and ischemia associated with stroke, neural paropathy, other neural degenerative diseases, motor neuron diseases, sciatic crush, neuropathy associated with diabetes, spinal cord injuries, facial nerve crush and other trauma, chemotherapy- and other medication-induced neuropathies and Huntington's disease.

In another aspect of this embodiment, the method is used to stimulate nerve growth ex vivo. For this aspect, the compounds or compositions described above can be applied directly to the nerve cells in culture. This aspect of the invention is useful for ex vivo nerve regeneration.

According to an alternate embodiment, the method of stimulating neurite outgrowth or preventing neurodegeneration comprises the additional step of treating a patient or ex vivo nerve cells in culture with a neurotrophic factor, such as those contained in the compositions of this invention described above. This embodiment includes administering the compound and the neurotrophic agent in a single dosage form or in separate, multiple dosage forms when they are to be

administered to a patient. If separate dosage forms are utilized, they may be administered concurrently, consecutively or within less than about 5 hours of one another.

5 In order that the invention described herein may be more fully understood, the following examples are set forth. It should be understood that these examples are for illustrative purposes only and are not to be construed as limiting this invention in any manner.

10 General Methods


Proton nuclear magnetic resonance (^1H NMR) spectra were recorded at 500 MHz on a Bruker AMX 500. Chemical shifts are reported in parts per million relative to Me_4Si . Analytical high performance liquid
15 chromatography was performed on a Hewlett-Packard 1050 liquid chromatograph.

EXAMPLE 1

N-(4-Aminobenzenesulfonamido)-(S)-Piperidine-2-Carboxylic
Acid-2-((N Methyl)-2-Pyridylethyl) Amide

20 The synthesis of N-(4-aminobenzenesulfonamido)-(S)-piperidine-2-carboxylic acid-2-((N-methyl)-2-pyridylethyl) amide (compound 1) is set forth below.

A. (S)-Piperidine-1,2-Dicarboxylic Acid-1-(Tert-Butyl ester)-2-((N Methyl)-2-Pyridinylethyl)
25 Amide.




To a solution of (S)-piperidine-1,2 dicarboxylic acid 1-tert-butylester (5.0 g, 21.8 mmol) in methylene chloride (50 ml) was added EDC (6.0 g, 191.71,

31.2 mmol) followed by the addition of 2-(2-methylaminoethyl)pyridine (3.0 g, 22 mmol). The mixture was allowed to stir at ambient temperature for 24 hours.

The solution was diluted with 200 ml ethyl acetate and water (50 ml). The aqueous layer made basic by the addition of 2N NaOH until pH 12-13 was achieved. The separated organics were dried over anhydrous MgSO_4 and concentrated under reduced pressure. The crude product was purified via medium pressure liquid chromatography using a gradient solvent system of 1:99 methanol/ methylene chloride followed by 2:98 methanol/ methylene chloride solution to give 3.8 g (50% yield) of the title compound as a colorless oil. TLC: $R_f=0.49$ (5:95 methanol/ methylene chloride), ^1H -NMR (CDCl_3) consistent with structure.

B. (S)-Piperidine-2-Carboxylic Acid-2-((N Methyl)-2-Pyridylethyl) Amide.




106

To the compound of part A (3.8 g, 10.9 mmol) in methylene chloride (25 ml) was added trifluoroacetic acid

(10 ml, 130 mmol). The mixture was allowed to stir 2 hours at ambient temperature. The solution was concentrated under reduced pressure to dryness. The residue was taken up in ethyl acetate (200 ml) and water (50 ml). The aqueous layer was made basic by the addition of 2N NaOH until pH 14-15 was achieved.

The separated organics were dried over anhydrous MgSO_4 and concentrated under reduced pressure. The crude product was purified via medium pressure liquid chromatography using a gradient solvent system of 1:99 methanol/methylene chloride followed by 1:10:90 NH_4OH /methanol/methylene chloride to give 2.2 g (81% yield) of the title compound as a colorless oil. TLC: $R_f=0.11$ (1:10:90 NH_4OH /methanol/methylene chloride), HPLC: $R_t=5.22$ min, ^1H -NMR (CDCl_3) consistent with structure.

C. N-(4-Nitrobenzenesulfonamido)-(S)-Piperidine-2-Carboxylic Acid-2-((N Methyl)-2-Pyridylethyl) Amide.




To a solution of the compound of part B (200 mg, 0.81 mmol) in methylene chloride (5 ml) was added triethylamine (2.0 ml, 101.19, 19.8 mmol) followed by the addition of 4-nitrobenzenesulfonylchloride (260 mg, 1.22 mmol). The mixture was allowed to stir at ambient temperature for 24 hours. The solution was diluted with 100 ml ethyl acetate and a saturated solution of sodium

bicarbonate (50 ml). The separated organics were dried over anhydrous MgSO_4 and concentrated under reduced pressure. The crude product was purified via medium pressure liquid chromatography using a gradient solvent system of methylene chloride followed by 1:99 methanol/methylene chloride solution to give 273 mg (78% yield) of the title compound as a yellowish oil. TLC: $R_f=0.57$ (5:95 methanol/ methylene chloride), ^1H -NMR (CDCl_3) consistent with structure.

10

D. N-(4-aminobenzenesulfonamido)-(S)-piperidine-2-carboxylic acid-2-((N-methyl)-2-pyridylethyl) amide (compound 1).




15 A solution of the compound from step C (273 mg, 0.63 mmol) in ethyl acetate (10 ml) and ethanol (10 ml) was treated under ambient temperature with 150 mg of 10% palladium on carbon and hydrogenated for 24 hours under a slight positive pressure of hydrogen. The mixture was
20 filtered and concentrated in vacuo and the crude product purified via medium pressure liquid chromatography using methylene chloride followed by 2:98 methanol/ methylene chloride followed by 0.5:5:95 NH_4OH /methanol/ methylene chloride solution as the solvent system to give 102 mg
25 (40% yield) of the title compound as a yellowish oil. TLC: $R_f=0.36$ (1:10:90 NH_4OH /methanol/methylene chloride), HPLC: $R_t= 6.86$ min, ^1H -NMR (CDCl_3) consistent with structure.

EXAMPLE 2

(S)-(N-Methyl)-2-(Methyl-(4-Amino-Benzenesulfanilamido))-
3-Phenyl-N-(2-(Pyridin-2yl) Ethyl) Propionamide

5 The synthesis of (S)-(N-methyl)-2-(methyl-(4-amino-benzenesulfanilamido))-3-phenyl-N-(2-(pyridin-2yl) ethyl) propionamide (compound 2) is set forth below.

A. (N-Methyl)-2-(Methyl-2-(Tertbutoxycarbonyl) Amino-3-Phenyl-N-(3-Pyridin-4-yl)-Propylbutyl Propionamide.



10

To a solution of Boc-(N-Methyl)phenylalanine (5.0 g, 17.8 mmol) in methylene chloride (50 ml) was added EDC (6.0 g, 191.71, 31.2 mmol) followed by the addition of 2-(2-methylaminoethyl)pyridine (2.5 g, 18.4 mmol). The mixture was allowed to stir at ambient temperature for 24 hours. The solution was diluted with 200 ml ethyl acetate and water (50 ml). The aqueous layer was made basic by the addition of 2N NaOH until pH 12 was achieved. The separated organics were dried over anhydrous MgSO₄ and concentrated under reduced pressure.


20

The crude product was purified via medium pressure liquid chromatography using a gradient solvent system of 1:99 methanol/methylene chloride followed by 2:98 methanol/methylene chloride solution to give 2.83 g (40%

- 42 -

yield) of the title compound as a colorless oil. TLC:
R_f=0.56 (5:95 methanol/ methylene chloride), [¹H]-NMR
(CDCl₃) consistent with structure.

B. (N-methyl)-2-(methylamino)-3-phenyl-N-(2-(pyridin-2yl)ethyl) propionamide.




107

5 To the compound of step A (2.83 g, 7.1 mmol) in methylene chloride (25 ml) was added trifluoroacetic acid (10 ml, 130 mmol). The mixture was allowed to stir 2 hours at ambient temperature. The solution was concentrated under reduced pressure to dryness. The residue was taken up in ethyl acetate (200 ml) and water (50 ml). The aqueous layer was made basic by the addition of 2N NaOH until pH 14-15 was achieved.

The separated organics were dried over anhydrous MgSO₄ and concentrated under reduced pressure.

15 The crude product was purified via medium pressure liquid chromatography using a gradient solvent system of 1:99 methanol/methylene chloride followed by 1:10:90 NH₄OH/methanol/methylene chloride) to give 1.53 g (75% yield) of the title compound as a colorless oil. TLC: 20 R_f=0.70 (1:10:90 NH₄OH/methanol/methylene chloride). HPLC: Rt= 5.54 min, [¹H]-NMR (CDCl₃) consistent with structure.


- C. (S)-(N-methyl)-2-(methyl-(4-nitro-benzenesulfanilamido))-3-phenyl-N-(2-(pyridin-2-yl)Ethyl) propionamide.



- 5 To a solution of the compound of step B (300 mg, 1.05 mmol) in methylene chloride (15 ml) was added triethylamine (2.0 ml, 101.19, 19.8 mmol) followed by the addition of 4-nitrobenzenesulfonylchloride (330 mg, 1.56 mmol). The mixture was allowed to stir at ambient
10 temperature for 24 hours. The solution was diluted with 100 ml ethyl acetate and a saturated solution of sodium bicarbonate (50 ml). The separated organics were dried over anhydrous MgSO₄ and concentrated under reduced pressure. The crude product was purified via medium
15 pressure liquid chromatography using a gradient solvent system of methylene chloride followed by 1:99 methanol/methylene chloride solution to give 453 mg (89% yield) of the title compound as a yellowish oil. TLC: R_f=0.63 (5:95 methanol/ methylene chloride), [¹H]-NMR (CDCl₃)
20 consistent with structure.

- D. (S)-(N-methyl)-2-(methyl-(4-amino-benzenesulfanilamido))-3-phenyl-N-(2-(pyridin-2-yl)ethyl) propionamide (compound 2).

1




- 5 A solution of the compound from step C (453 mg, 0.94 mmol) in ethyl acetate (20 ml) was treated under ambient temperature with 150 mg of 10% palladium on carbon and hydrogenated for 24 hours under a slight positive pressure of hydrogen. The mixture was filtered and
10 concentrated in vacuo and the crude product purified via medium pressure liquid chromatography using 2:98 methanol/methylene chloride followed by 0.5:5:95 NH₄OH/methanol/methylene chloride solution as the solvent system to give 194 mg (46% yield) of the title compound
15 as a yellowish oil. TLC: R_f=0.46 (1:10:90 NH₄OH/methanol/methylene chloride), HPLC: R_t= 9.04 min, [¹H]-NMR (CDCl₃) consistent with structure.

EXAMPLE 3

- 20 N-(4-Nitrobenzenesulfonamido)-(S)-Piperidine-2-Carboxylic Acid-((N Methyl)-3-(Pyridin-3-Yl)Propyl) Amide

The synthesis of N-(4-nitrobenzenesulfonamido)-(S)-piperidine-2-carboxylic acid-((N-methyl)-3-(pyridin-3-yl)propyl) amide (compound 3) is described below.


A. 3-(3-Bromopropyl)-Pyridine.



To a solution of 3-pyridinepropanol (9.50 g, 69.3 mmol) in DMF (50 ml) was added triphenylphosphine (20.0 g, 76.2 mmol) followed by the addition of bromine (5.4 ml, 104 mmol). The mixture was allowed to stir at ambient temperature for 48 hours. The solution was diluted with 200 ml water and the aqueous layer made basic by the addition of 2N NaOH until pH 12-13 was achieved. The organics were taken up in ethyl acetate (200 ml). The separated organics were dried over anhydrous MgSO_4 and concentrated under reduced pressure.

The crude product was purified via medium pressure liquid chromatography using a gradient solvent system of methylene chloride followed by a 1:99 methanol/ methylene chloride solution to give 11.8 g (85% yield) of the title compound as a yellowish orange oil. TLC: R_f =0.8 (3:97 methanol/ methylene chloride), ^1H -NMR (CDCl_3) consistent with structure.


B. N-Methyl-3-(3-Aminopropyl)-Pyridine.



A solution of the compound from step A (11.8 g, 59 mmol) in methanol (50 ml) was bubbled N-methylamine gas for a period of 10 to 15 min until saturated. The flask was sealed and the mixture allowed to stir overnight at ambient temperature. The mixture was concentrated in vacuo and the residue taken up in ethyl

acetate and water. The aqueous layer was made basic by the addition of 2N NaOH until pH 14-15 was achieved. The organics were taken up in ethyl acetate (250 ml). The separated organics were dried over anhydrous MgSO₄ and concentrated under reduced pressure. The crude product purified via medium pressure liquid chromatography using methylene chloride followed by 5:95 methanol/ methylene chloride followed by a 1:10:90 NH₄OH/methanol/ methylene chloride solution as the solvent system to give 3.2 g (36% yield) of the title compound as colorless oil. TLC: R_f=0.14 (1:10:90 NH₄OH/methanol/methylene chloride), [¹H]-NMR (CDCl₃) consistent with structure.

C. N-methyl-(S)-piperidine-1,2-dicarboxylic acid-1-(tert-butyl ester)-3-((pyridin-3-yl)propyl) amide.




To a solution of (S)-piperidine-1,2 dicarboxylic acid 1-tert-butylester (2.28 g, 9.9 mmol) in methylene chloride (30 ml) was added EDC (2.0 g, 191.71, 10.4 mmol) followed by the compound from step B (11.8 g, 59 mmol). The mixture was allowed to stir at ambient temperature for 24 hours. The solution was diluted with 200 ml ethyl acetate and a saturated solution of sodium bicarbonate (50 ml). The separated organics were dried over anhydrous MgSO₄ and concentrated under reduced pressure.

The crude product was purified via medium pressure liquid chromatography using a gradient solvent

system of 1:99 methanol/methylene chloride followed by 5:95 methanol/ methylene chloride solution to give 1.2 g (33% yield) of the title compound as a colorless oil. TLC: R_f =0.28 (5:95 methanol/ methylene chloride), ^1H -NMR (CDCl_3) consistent with structure.

D. (S)-piperidine-2-carboxylic acid-((N-methyl)-3-(pyridin-3-yl)propyl) amide.




110

To the compound from step C (1.2 g, 3.3 mmol) in methylene chloride (20mL) was added trifluoroacetic acid (10 ml, 130 mmol). The mixture was allowed to stir 3 hours at ambient temperature. The solution was concentrated under reduced pressure to dryness. The residue was taken up in ethyl acetate (200 ml) and 2N NaOH (50 ml). The separated organics were dried over anhydrous MgSO_4 and concentrated under reduced pressure.


The crude product was purified via medium pressure liquid chromatography using a gradient solvent system of 5:95 methanol/methylene chloride followed by 1:10:90 NH_4OH /methanol/methylene chloride) to give 850 mg (98% yield) of the title compound as a colorless oil. TLC: R_f =0.17 (1:10:90 NH_4OH /methanol/methylene chloride), HPLC: R_t = 6.67 min, ^1H -NMR (CDCl_3) consistent with structure.

- E. N-(4-nitrobenzenesulfonamido)-(S)-piperidine-2-carboxylic acid-((N-methyl)-3-(pyridin-3-yl)propyl) amide.



To a solution of the compound from step D (250 mg, 0.96 mmol) in methylene chloride (15 ml) was added triethylamine (2.0 ml, 101.19, 19.8 mmol) followed by the addition of 4-nitrobenzenesulfonylchloride (300 mg, 1.42 mmol). The mixture was allowed to stir at ambient temperature for 24 hours. The solution was diluted with 200 ml ethyl acetate and a saturated solution of sodium bicarbonate (50 ml). The separated organics were dried over anhydrous MgSO_4 and concentrated under reduced pressure. The crude product was purified via medium pressure liquid chromatography using a gradient solvent system of methylene chloride followed by 1:99 methanol/methylene chloride solution to give 165 mg (37% yield) of the title compound as a yellowish oil. TLC: $R_f=0.52$ (5:95 methanol/ methylene chloride), ^1H -NMR (CDCl_3) consistent with structure.

F. N-(4-nitrobenzenesulfonamido)-(S)-piperidine-2-carboxylic acid-((N-methyl)-3-(pyridin-3-yl)propyl) amide (compound 3).



5 A solution of the compound from step E (165 mg, 0.35 mmol) in ethyl acetate (20 ml) was treated under ambient temperature with 150 mg of 10% palladium on carbon and hydrogenated for 24 hours under a slight positive pressure of hydrogen. The mixture was filtered and
10 concentrated in vacuo and the crude product purified via medium pressure liquid chromatography using methylene chloride followed by 2:98 methanol/
methylene chloride followed by 0.5:5:95 NH₄OH/methanol/
methylene chloride solution as the solvent system to give
15 60 mg (41% yield) of the title compound as a yellowish oil. TLC: R_f=0.20 (5:95 methanol/methylene chloride), HPLC: R_t= 7.25 min, [¹H]-NMR (CDCl₃) consistent with structure.


EXAMPLE 4

20 (S)-(N-Methyl)-2-(Methyl-(4-Amino-Benzenesulfanilamido))-3-Phenyl-N-(4-(Pyridin-3-Yl)-1-(3-(Pyridin-3yl)Propyl) Butyl) propionamide

The synthesis of (S)-(N-methyl)-2-(methyl-(4-amino-benzenesulfanilamido))-3-phenyl-N-(4-(pyridin-3-yl)-1-(3-(pyridin-3yl)Propyl) butyl) propionamide is set
25 forth below.


A. (S)-(N-methyl)-2-(methyl-2-tertbutyloxycarbonyl)amino)-3-phenyl-N-(4-(pyridin-3-yl)-1-(3-(pyridin-3-yl)propyl)butyl)propionamide:

5




To a solution of Boc-(N-Methyl)phenylalanine (1.42 g, 5.1 mmol) in methylene chloride (10 ml) was added EDC (0.98 g, 191.71, 5.1 mmol) followed by the addition of N-methyl-1,7 bis(3-pyridyl)-4-heptylamine (1.2 g, 18.4 mmol). The mixture was allowed to stir at ambient temperature for 24 hours. The solution was diluted with 100 ml ethyl acetate and a saturated solution of sodium bicarbonate (50 ml). The separated organics were dried over anhydrous MgSO₄ and concentrated under reduced pressure. The crude product was purified via medium pressure liquid chromatography using 2:98 methanol/methylene chloride solution to give 970 mg (42% yield) of the title compound as a colorless oil. TLC: R_f=0.51 (5:95 methanol/ methylene chloride), [¹H]-NMR (CDCl₃) consistent with structure.

B. (2S)-N-Methyl-2-methylamino-3-phenyl-N-[4-pyridin-3-yl-1-(3-pyridin-3-yl-propyl)-butyl]-propionamide




To the compound of step A (5.0 g, 9.2 mmol) in
5 methylene chloride (20 ml) was added trifluoroacetic acid (20
ml, 260 mmol). The mixture was allowed to stir 3 hours at
ambient temperature. The solution was concentrated under
reduced pressure to dryness. The residue was taken up in
ethyl acetate (200 ml) and a saturated solution of sodium
10 bicarbonate (100 ml). The separated organics were dried over
anhydrous MgSO_4 and concentrated under reduced pressure. The
crude product was purified via medium pressure liquid
chromatography using a gradient solvent system of 1:99
methanol/ methylene chloride followed by 5:95 methanol/
15 methylene chloride followed by 1:10:90
 NH_4OH /methanol/methylene chloride) to give 3.12 g (76% yield)
of the title compound as a colorless oil. TLC: R_f =0.38
(1:10:90 NH_4OH /methanol/methylene chloride). HPLC: R_t = 9.52
min, ^1H -NMR (CDCl_3) consistent with structure.

C. (S)-(N-methyl)-2-(methyl-(4-nitrobenzenesulfanilamido))-3-phenyl-N-(4-(pyridin-3-yl)-1-(3-(pyridin-3-yl)propyl)Butyl)propionamide.



5 To a solution of the compound of step B (1.5 g, 3.4 mmol) in methylene chloride (20 ml) was added triethylamine (5.0 ml, 101.19, 35.8 mmol) followed by the addition of 4-nitrobenzenesulfonylchloride (1.0 g, 4.7
10 mmol). The mixture was allowed to stir at ambient temperature for 24 hours. The solution was diluted with 150 ml ethyl acetate and washed with water (50 ml). The separated organics were dried over anhydrous MgSO₄ and concentrated under reduced pressure. The crude product
15 was purified via medium pressure liquid chromatography using a gradient solvent system of methylene chloride followed by 1:99 methanol/methylene chloride solution to give 1.75 g (83% yield) of the title compound as a yellowish oil. TLC: R_f=0.67 (5:95 methanol/methylene
20 chloride), [¹H]-NMR (CDCl₃) consistent with structure.

- D. (S)-(N-methyl)-2-(methyl-(4-amino-benzenesulfanilamido))-3-phenyl-N-(4-(pyridin-3-yl)-1-(3-(pyridin-3-yl)propyl)butyl)propionamide (compound 4).



5 A solution of the compound of step C (1.75 g, 2.78 mmol) in ethyl acetate (50 ml) was treated under ambient temperature with 1.0 g of 10% palladium on carbon and hydrogenated for 24 hours under a slight positive
10 pressure of hydrogen. The mixture was filtered and concentrated in vacuo and the crude product purified via medium pressure liquid chromatography using methylene chloride followed by 1:99 methanol/methylene chloride followed by 3:97 methanol/methylene chloride solution as
15 the solvent system to give 0.79 mg (47% yield) of the title compound as a yellowish oil. TLC: R_f=0.36 (1:10:90 NH₄OH/methanol/ methylene chloride). HPLC: R_t= 7.97 min, [¹H]-NMR (CDCl₃) consistent with structure.


20 The synthesis of other compounds of this invention, including those listed in Table 1 above, may be achieved by modifying the synthesis schemes set forth in Examples 1-4 using appropriate reagents that are well known in the art.

EXAMPLE 5

(S)-Piperidine-2-Carboxylic Acid,
(4-Pyridylmethyl) Amide, Citrate Salt


The synthesis of (S)-piperidine-2-carboxylic
5 acid-1-(tert-butyl ester)-2-(4-pyridylmethyl) amide,
citrate salt (compound 1) is set forth below.

A. (S)-piperidine-1,2-dicarboxylic acid-1-(tert-
butyl ester)-2-(4-pyridylmethyl) amide




10 Following the method described in Example 1,
part A, (S)-piperidine-1,2 dicarboxylic acid-1-(tert-
butyl ester) (2.0 g, 8.72 mmol) and 4-(aminomethyl)
pyridine (3.18 g, 29.41 mmol) was converted to 0.75 g
(27% yield) of product. ¹H-NMR (CDCl₃) consistent with
15 structure.

B. (S)-Piperidine-2-Carboxylic acid,
(4-Pyridylmethyl) Amide



20 Following the method of Example 1, part B, (S)-
piperidine-1-carboxylic acid-1-(tert-butyl ester)-2-(4-
pyridylmethyl) amide (0.75 g, 2.35 mmol) gave 0.49 g (95%
yield) of the title compound. ¹H-NMR (CDCl₃)
consistent with structure.

C. (S)-Piperidine-2-Carboxylic acid, (4-Pyridylmethyl) Amide, Citrate Salt



A solution of the amine (107 mg, 0.48 mmol) from part B and citric acid (94 mg, 0.48 mmol) in absolute ethanol was warmed to 60°C until dissolved. The solution was concentrated in vacuo and the residue was dissolved in absolute ethanol and concentrated in vacuo to give a foam. ^1H -NMR (CDCl_3) consistent with structure.

10


EXAMPLE 6

(S)-Piperidine-2-Carboxylic Acid, (4-Pyridylmethyl) Amide, Citrate Salt

The synthesis of (S)-piperidine-1,2-dicarboxylic acid-1-(tert-butyl ester)-2-(3-pyridylmethyl) amide, citrate salt is set forth below.


15

A. (S)-piperidine-1,2-dicarboxylic acid-1-(tert-butyl ester)-2-(3-pyridylmethyl) amide




Following the method described in Example 1, part A, (S)-piperidine-1,2-dicarboxylic acid-1-(tert-butyl ester) (2.0 g, 8.72 mmol) and 3-(aminomethyl) pyridine (3.18 g, 29.41 mmol) was converted to 1.0 g (36% yield) of product. ¹H-NMR (CDCl₃) consistent with structure.

B. (S)-Piperidine-2-Carboxylic acid, (3-Pyridylmethyl) Amide



Following the method of Example 1, part B, (S)-piperidine-1,2-dicarboxylic acid-1-(tert-butyl ester)-2-(3-pyridylmethyl) amide (1.0 g, 3.13 mmol) gave 0.56 g (82% yield) of the title compound. ¹H-NMR (CDCl₃) consistent with structure.

C. (S)-Piperidine-2-Carboxylic acid, (3-Pyridylmethyl) Amide, Citrate Salt




103

A solution of the amine (111 mg, 0.51 mmol) in part B and citric acid (97 mg, 0.51 mmol) was warmed to 60°C until dissolved. The solution was concentrated in vacuo and the residue was dissolved in absolute ethanol and concentrated in vacuo to give a foam. ¹H-NMR (CDCl₃)

consistent with structure.

EXAMPLE 7


(S)-Piperidine-2-Carboxylic Acid-2-((N Methyl)-2-Pyridylethyl) Amide, Citrate Salt



Following the method described in Example 5, part C, (S)-piperidine-2-carboxylic acid-2-((N methyl)-2-pyridylethyl) amide (product in example 1, part B) was converted to a citrate salt. $[^1\text{H}]$ -NMR (CDCl_3) consistent with structure.

EXAMPLE 8

(S)-Piperidine-2-Carboxylic acid-((N-Methyl)-3-(Pyridin-3-yl)Propyl) Amide, Citrate Salt




Following the procedure described in Example 6, part C, S)-piperidine-2-carboxylic acid-((N-methyl)-3-(pyridin-3-yl)propyl) amide (product from Example 3, part D) was converted to a citrate salt. $[^1\text{H}]$ -NMR (CDCl_3) consistent with structure.

EXAMPLE 9

(S)-Piperidine-2-Carboxylic Acid,
(1,7-di-pyridin-3-yl)heptan-4-yl ester, Fumarate Salt

5

- A. (S)-Piperidine-1,2-Dicarboxylic Acid-1-(Tert-Butyl ester)-2-(1,7-di-pyridin-3-yl)heptan-4-yl ester




10

A solution of (1,7-di-pyridin-3-yl)heptan-4-ol (6.6 g, 24.41 mmol) in THF (30 ml) was added to (S)-piperidine-1,2 dicarboxylic acid-1-(tert-butyl ester) (5.0 g, 21.81 mmol) and EDC (4.7 g, 24.52 mmol) and allowed to stir for 18 h at room temperature. The reaction was diluted with ethyl acetate (200 ml) and washed with water. The organic layer was dried over MgSO₄, concentrated in vacuo and purified by medium pressure liquid chromatography using 1:100 methanol/methylene chloride as the solvent system to give 2.0 g (20% yield) of the title compound. [¹H]-NMR (CDCl₃) consistent with structure.


20

B. (S)-Piperidine-2-Carboxylic Acid, (1,7-di-pyridin-3-yl)heptan-4-yl) ester



Following the method of Example 1, part B, the compound from Example 9, part A, (1.0 g, 4.30 mmol) gave 1.45 g (88% yield) of the title compound. $[^1\text{H}]\text{-NMR}$ (CDCl_3) consistent with structure.


C. (S)-Piperidine-2-Carboxylic Acid, (1,7-di-pyridin-3-yl)heptan-4-yl) ester, Bis-fumarate Salt



Following the procedure in Example 6, part C, the amine in example 9, part B may be converted to the title compound using 1 equivalent of the amine and two equivalents of fumaric acid. $[^1\text{H}]\text{-NMR}$ (CDCl_3) consistent with structure.

EXAMPLE 10

(S)-1-Methyl-Piperidine-2-Carboxylic Acid, (1,7-di-pyridin-3-yl)heptan-4-yl) ester




111

5 A mixture of the amine (300 mg, 0.79 mmol) from
Example 9, part B and paraformaldehyde (500 mg) in methanol
(15 ml) was added to Na(CN)BH₃ (500 mg). The mixture was
stirred for 65 h at room temperature. The reaction was
concentrated in vacuo and taken up in 2N aqueous NaOH and
10 extracted with ethyl acetate (150ml). The organic layer was
dried over MgSO₄ concentrated in vacuo and purified by medium
pressure liquid chromatography using a gradient solvent
system of 2:98 methanol/methylene chloride followed by
0.5:5:95 NH₄OH/methanol/methylene chloride to give 230 mg
15 (74% yield) of the title compound as a clear oil. [¹H]-NMR
(CDCl₃) consistent with structure.

EXAMPLE 11


(S)-1-(2-Methylpropyl)-Piperidine-2-Carboxylic Acid,
1,7-di-pyridin-3-yl)heptan-4-yl) ester



5 Following the method described in Example 10,
mixture of the amine (300 mg, 0.79 mmol) from Example 9, part
B and 2-methylpropionaldehyde (1.6 g, 22.0 mmol) in methanol
(15 ml) was added to Na(CN)BH₃ (500 mg). The mixture was
stirred for 65 h at room temperature. The reaction was
10 concentrated in vacuo and taken up in 2N aqueous NaOH (20 ml)
and extracted with ethyl acetate (150ml). The organic layer
was dried over MgSO₄, concentrated in vacuo and purified by
medium pressure liquid chromatography using a gradient
solvent system of 2:98 methanol/methylene chloride followed
15 by 0.5:5:95 NH₄OH/methanol/methylene chloride to give 170 mg
(49% yield) of the title compound as a clear oil. [¹H]-NMR
(CDCl₃) consistent with structure.

EXAMPLE 12

(S)-1-(Pyridin-4-ylmethyl)-Piperidine-2-Carboxylic Acid
(1,7-di-pyridin-3-yl)heptan-4-yl ester




113

5

Following the method described in Example 10,
mixture of the amine (300 mg, 0.79 mmol) from Example 9, part
B and 4-pyridinecarboxaldehyde (0.5 g, 4.67 mmol) in methanol
(15 ml) was added to Na(CN)BH₃ (500 mg). The mixture was
10 stirred for 65 h at room temperature. The reaction was
concentrated in vacuo and taken up in 2N aqueous NaOH (20 ml)
and extracted with ethyl acetate (150ml). The organic layer
was dried over MgSO₄, concentrated in vacuo and purified by
medium pressure liquid chromatography using a gradient
15 solvent system of 2:98 methanol/methylene chloride followed
by 0.5:5:95 NH₄OH/methanol/methylene chloride to give 70 mg
(19% yield) of the title compound as a clear oil. [¹H]-NMR
(CDCl₃) consistent with structure.

EXAMPLE 13

(S) - (N-Methyl) - 2 - (Methylamino) - 3 - Phenyl - N - (2 - (Pyridin-2-yl)ethyl) Propionamide, Citrate Salt




5 Following the method described in Example 6, part C, (N-methyl)-2-(methylamino)-3-phenyl-N-(2-(pyridin-2-yl)ethyl) propionamide (product in Example 2, part B) was converted to a citrate salt. ^1H -NMR (CDCl_3) consistent with structure.

10

EXAMPLE 14

(2S) - N-Methyl-2-methylamino-3-phenyl-N - [4-pyridin-3-yl-1-(3-pyridin-3-yl-propyl)-butyl]-propionamide, Citrate Salt



15 Following the method described in Example 6, part C, (2S)-N-Methyl-2-methylamino-3-phenyl-N-[4-pyridin-3-yl-1-(3-pyridin-3-yl-propyl)-butyl]-propionamide (product in

Example 4, part B) was converted to a citrate salt. [^1H]-NMR (CDCl_3) consistent with structure.

EXAMPLE 15

Rotamase Inhibition Assay

5 Inhibition of FKBP enzyme activity was determined in the PPIase assay described in S. T. Park et al., J. Biol. Chem., 267, pp. 3316-24 (1992), the disclosure of which is herein incorporated by reference. This is a chymotrypsin-coupled assay in which FKBP catalyzes the cis to trans
(
10 isomerization of the Leu-Pro bond in the peptide substrate Suc-Ala-Leu-Pro-Phe-pNA. The trans form, but not the cis form, is cleavable by chymotrypsin. The release Phe-pNA was monitored by absorbance at 400 nm. Chymotrypsin cleavage is very rapid and therefore cis-to-trans isomerization is rate-
15 limiting.

Each reaction mix consisted of 0.1 M Tris buffer, pH 7.8, 15 nM FKBP, 30 μM substrate and 0.5 nM to 10 μM of compound to be tested diluted in Me_2SO . The mixture was incubated at 15°C for 5 minutes and then reaction was
20 initiated by addition of chymotrypsin (100 $\mu\text{g}/\text{ml}$ final concentration) and followed spectrophotometrically for 5 minutes. Total reaction volume was 1 ml. None of the compounds of this invention demonstrated a K_i of less than 10 μM , as demonstrated in the table below.

Table 4. Rotamase Inhibition.

Compound #	Rotamase Inhibition (KI) (nM)	Compound #	Rotamase Inhibition (KI) (nM)
2	>50,000	106	>50,000
3	>50,000	107	>25,000
4	>25,000	108	>25,000
8	>50,000	110	>25,000
18	>50,000	111	>50,000
101	>50,000	112	>50,000
103	>50,000	113	>50,000
104	>50,000		

EXAMPLE 16

MDR Sensitization Assay

5 To demonstrate that the compounds according to this invention do not possess MDR reversal activity, cell lines which are known to be resistant to a particular drug were used.

We carried out multidrug resistance reversal (MDR) assays using L1210vMDRC.06 or HL60/Vinc cell lines.

10 L1210vMDRC.06 are L1210 mouse leukemia cells transduced with the pHaMDR1/A retrovirus carrying a MDR1 cDNA, as described by Pastan et al., Proc. Natl. Acad. Sci. USA, 85, pp. 4486-4490 (1988). The L1210vMDRC.06 multidrug resistant line is a cell line that has been drug-selected by culturing transfected cells in 0.06 µg/ml colchicine.

15 The HL60/Vinc human promyleocytic leukemia cell line is a multidrug resistant cell line derived from HL60 drug-sensitive parental cells by selection in increasing concentrations of vincristine.

20 Using L1210vMDRC.06, multidrug resistance reversal assays were conducted by plating 1×10^4 cells/well in 96 well microtiter plates and exposing them

to a concentration range of doxorubicin (50 nM - 10 μ M) in the presence or absence of compounds of this invention (0.1, 0.25, 0.5, 1.0, or 2.5 μ M) as described in U. Germann et al., Anti-Cancer Drugs, 8, pp. 125-140 (1997).

5 After culture for 3 days, the viability of cells was quantitated using XTT dye to assess mitochondrial function [Roehm et al., J. Immunol. Methods, 142, pp. 257-265, (1991)]. All determinations were made in replicates of at least 4. Results were determined by
10 comparison of the IC50 for doxorubicin alone to the IC50 for doxorubicin + compound. An MDR ratio was calculated (IC50 Dox/ IC50 Dox + Inhibitor) and the integer value used for comparison of compound potencies.

Assays using HL60/Vinc cells were performed by
15 plating cells in 96 well microtiter plates at a concentration of 4×10^4 cells/well. The cells were then exposed to various concentrations of doxorubicin (9 nM to 6.7 μ M) in the presence or absence of various compounds of this invention at various concentrations (0.5, 1.0,
20 2.5, 5.0 or 10 μ M) as described in U. Germann et al., Anti-Cancer Drugs, 8, pp. 141-155 (1997). After culturing the cells for 3 days, their viability was quantitated using the XTT dye method to assess
25 mitochondrial function (Roehm et al., supra). Results were expressed as a ratio of the IC50 for doxorubicin alone to the IC50 for doxorubicin plus compound. In all assays the intrinsic antiproliferative or cytotoxicity activity of the MDR inhibitors was also determined for HL60/Vinc cells.

30 The results of these assays on several compounds of this invention are depicted in the table below.

35 Table 5. MDR Reversal Assay.

Compound #	MDR Ratio*	Cell line	Compound #	MDR Ratio*	Cell line
1	0.6	HL60/Vinc	16	19.9	HL60/Vinc
2	0.7	HL60/Vinc	17	14.6	HL60/Vinc
3	0.6	HL60/Vinc	18	10.8	HL60/Vinc
4	6.2	HL60/Vinc	102	1.2	HL60/Vinc
5	30.4	HL60/Vinc	103	0.7	HL60/Vinc
6	17.7	HL60/Vinc	104	0.6	HL60/Vinc
7	21.9	HL60/Vinc	105	10	L1210vMDR C.D6
8	28.5	HL60/Vinc	106	1.2	HL60/Vinc
9	24.3	HL60/Vinc	107	0.9	HL60/Vinc
10	9.1	HL60/Vinc	108	0.4	HL60/Vinc
11	1.7	HL60/Vinc	110	1.1	HL60/Vinc
12	22.6	HL60/Vinc	111	1.2	HL60/Vinc
13	19.3	HL60/Vinc	112	2.4	HL60/Vinc
14	19.9	HL60/Vinc	113	2.3	HL60/Vinc
15	2.4	HL60/Vinc			

* MDR ratios tested at compound concentrations of 2.5 μ M, except for compounds 5, 6 and 7, which were tested at 10 μ M.

5 As can be seen from these results, there does not seem to be a strong correlation between structure and MDR ratio. Factors which could be responsible for this are the ability of the compound to get into the cell, the toxicity of the compound and the metabolism of the compound inside the cell. As such, compounds with an MDR ratio higher than 7 would not be included as compounds which lacks MDR reversal activity according to the parameters of this invention.

EXAMPLE 17

15 Neurite Outgrowth Quantitation In PC12 Cell System

In order to directly determine the neurotrophic activity of compounds described in this invention, the neurite outgrowth assay was carried out with

pheochromocytoma PC12 cells as described by Lyons et al. (1994).

PC12 cells were maintained at 37°C and 5% CO₂ in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% heat-inactivated horse serum, 5% heat-inactivated fetal bovine serum (FBS), and 1% glutamate.

The cells were then plated at 105 per well in 96 well plates coated with 5 µg/cm² rat tail collagen and allowed to attach overnight. The medium was then replaced with DMEM, 2% heat-inactivated horse serum, 1% glutamate, 1-5 ng/ml of NGF (Sigma) and varying concentrations of the compounds of this invention (0.01 nM - 10000 nM). The background control culture was administered with 105 ng/ml of NGF alone without compound. Positive control cultures were administered with high concentration of NGF (50 ng/ml). The cells were then incubated at 37°C at 5% CO₂ for 72 hours, fixed with 3% formaldehyde and nerve outgrowth was determined visually on a scale of 0 to 4.

The results are shown in the table below.

Table 6: Neurite Outgrowth Activity of Compounds of this Invention.

Compound	0.01 nM	0.1 nM	1 nM	10 nM	100 nM	1000 nM	10000 nM
1	ND	ND	ND	3 ^{***}	1 ^{***}	3 ^{***}	ND
2	ND	ND	ND	3 ^{***}	3 ^{***}	4 ^{***}	ND
3	ND	ND	ND	2 ^{***}	4 ^{***}	4 ^{***}	ND
4	ND	ND	ND	3 [*]	3 [*]	4 [*]	ND
5	ND	ND	ND	2 [*]	3 [*]	3 [*]	ND
6	ND	ND	ND	4 [*]	2 [*]	3 [*]	ND
7	ND	ND	ND	2 [*]	3 [*]	3 [*]	ND
8	ND	ND	ND	4 [*]	2 [*]	3 [*]	ND
9	ND	ND	ND	4 [*]	2 [*]	3 [*]	ND
10	ND	ND	ND	3 [*]	4 [*]	2 [*]	ND
11	ND	ND	ND	1 [*]	3 [*]	3 [*]	ND

Compound	0.01 nM	0.1 nM	1 nM	10 nM	100 nM	1000 nM	10000 nM
12	ND	ND	ND	3*	3*	4*	ND
13	ND	ND	ND	3*	3*	4*	ND
14	ND	ND	ND	3*	2*	4*	ND
15	ND	ND	ND	3*	3*	3*	ND
16	ND	ND	ND	3*	4*	3*	ND
17	ND	ND	ND	3*	3*	4*	ND
18	ND	ND	ND	3*	4*	4*	ND
101	ND	ND	ND	ND	ND	ND	ND
102	3*	3*	4*	4**	2**	3**	4*
103	ND	ND	ND	0	0	0	ND
104	3*	3*	2*	2*	4*	3*	4*
105	ND	ND	ND	4	4	0	ND
106	ND	ND	ND	ND	1*	2*	ND
107	ND	ND	ND	ND	3*	0*	ND
108	ND	ND	ND	0	3	0	ND
109	3*	4*	4*	3*	2*	4*	3*
110	ND	ND	ND	ND	3*	0*	ND
111	4*	4*	3*	2**	4**	4**	4*
112	ND	ND	ND	3*	4*	3*	ND
113	ND	ND	ND	3*	4*	4*	ND

* Assay done in triplicate for each concentration tested. Indicated result is average of the three samples.

** Assay repeated twice, each time in triplicate.
Indicated result is average of the six samples.

*** Assay done in quadruplicate for each
concentration tested. Indicated result is average of the
5 three samples.

The compounds of this invention as exemplified
by compounds 1 - 18, 101 and 103-113 caused a significant
increase in neurite outgrowth over background control
10 cultures. Lack of nerve growth stimulatory activity (as
indicated by the "0") at high compound concentrations of
certain compounds is attributed to the toxic effects of
those compounds on the cells at higher concentrations.
Negative results for compound 103 were the result of only
15 a single experiment with one sample assayed at each of 3
concentrations. We believe this compound will display
some nerve growth stimulatory activity in the PC12 assay
when this experiment is repeated.


Other compounds of this invention will also
20 demonstrate significant nerve growth stimulatory
activity.

While we have hereinbefore presented a number
of embodiments of this invention, it is apparent that my
basic construction can be altered to provide other
(25 embodiments which utilize the methods of this invention.
Therefore, it will be appreciated that the scope of this
invention is to be defined by the claims appended hereto
rather than the specific embodiments which have been
presented hereinbefore by way of example.

We claim:

1. A compound of the formula:

(III)



and pharmaceutically acceptable derivatives thereof, wherein:

A and B are independently selected from Ar-substituted-(C₁-C₆)-straight or branched alkyl, or Ar-substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl; Ar is selected from 2-pyridyl, 3-pyridyl, or 4-pyridyl; and wherein

each Ar is optionally substituted with one to three substituents independently selected from halogen, hydroxyl, nitro, --SO₃H, trifluoromethyl, trifluoromethoxy, (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl, O-((C₁-C₆)-straight or branched alkyl), O-((C₂-C₆)-straight or branched alkenyl), O-benzyl, O-phenyl, 1,2-methylenedioxy, --N(R¹)(R²), carboxyl, N-(C₁-C₅ -straight or branched alkyl or C₂-C₅ -straight or branched alkenyl) carboxamide, N,N-di-(C₂-C₅ -straight or branched alkyl or C₂-C₅ -straight or branched alkenyl) carboxamide, N-(C₁-C₅ straight or branched alkyl or C₂-C₅ -straight or branched alkenyl) sulfonamide, N,N-di-(C₁-C₅ -straight or branched alkyl or C₂-C₅ -straight or branched alkenyl) sulfonamide, morpholinyl, piperidinyl, O--Z, CH₂ --(CH₂)_q --Z, O--(CH₂)_q --Z, (CH₂)_q --Z--O--Z, or CH=CH--Z;

wherein R¹ and R² are independently selected from (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, hydrogen or benzyl; or wherein R₁ and R₂ are taken together with the nitrogen atom to which they are bound to form a 5-7 membered heterocyclic ring;

Z is selected from 2-pyridyl, 3-pyridyl, or 4-pyridyl; and

q is 0, 1 or 2;

X is N, O or C(R);

wherein R is selected from hydrogen, (C₁-C₆)-straight or branched alkyl, or (C₂-C₆)-straight or branched alkenyl or alkynyl;

wherein when X is N or C(R), Y is selected from hydrogen, Ar, (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, (C₅-C₇)-cycloalkyl-substituted-(C₁-C₆)-straight or branched alkyl, (C₅-C₇)-cycloalkyl-substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl, (C₅-C₇)-cycloalkenyl-substituted-(C₁-C₆)-straight or branched alkyl, (C₅-C₇)-cycloalkenyl-substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl, Ar-substituted-(C₁-C₆)-straight or branched alkyl, Ar-substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl, phenyl substituted-(C₁-C₆)-straight or branched alkyl, or phenyl substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl;

when X is O, Y is a lone pair of electrons;

K is selected from (C₁-C₆)-straight or branched alkyl, Ar-substituted-(C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, Ar-substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl, phenyl substituted-(C₁-C₆)-straight or branched alkyl, phenyl substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl, or cyclohexylmethyl; wherein any one of the CH₂ groups of said alkyl, alkenyl or alkynyl chains in K is optionally replaced by O, S, S(O), S(O)₂ or N(R);

n is 0, 1 or 2;

J is selected from hydrogen, (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, Ar-substituted-(C₁-C₆)-straight or branched alkyl, Ar-substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl, phenyl substituted-(C₁-C₆)-straight or branched alkyl, phenyl substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl, or cyclohexylmethyl;

R³ is selected from (C₁-C₆)-straight or branched alkyl, Ar-substituted-(C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, Ar-substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl, phenyl substituted-(C₁-C₆)-straight or branched alkyl, or phenyl substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl;

wherein any one of the CH₂ groups of said alkyl, alkenyl or alkynyl chains in R³ is optionally replaced by O, S, S(O), or N(R); and

wherein any one of the CH₂ groups of said alkyl, alkenyl or alkynyl in R³, except the CH₂ group bound to nitrogen, is optionally replaced with C(O) or S(O)₂.

2. The compound according to claim 1, wherein at least one of A or B is an Ar-substituted (C₁-C₆)-alkyl chain.

3. The compound according to claim 2, wherein at least one of A or B is a (C₁-C₆)-alkyl chain terminally substituted with pyridinyl.

4. The compound according to claim 1, wherein X is N or O.

5. The compound according to claim 1, wherein K is benzyl.

6. The compound according to claim 1, wherein J is hydrogen or methyl.

7. The compound according to claim 1, wherein R³ is selected from hydrogen, (C₁-C₆)-alkyl, (C₁-C₆)-alkyl terminally substituted with pyridyl, or 3,4,5-trimethoxybenzoylmethyl.


8. The compound according to claim 1, wherein n is 0.

9. The compound according to claim 1, wherein said compound is (2S)-N-Methyl-2-methylamino-3-phenyl-N-[4-pyridin-3-yl-1-(3-pyridin-3-yl-propyl)-butyl]-propionamide.

10. A composition comprising:

- a) a neurotrophic amount of a compound according to claim 1; and
- b) a pharmaceutically suitable carrier.

11. A compound of the formula:



and pharmaceutically acceptable derivatives thereof; wherein:

at least one of A or B is a (C₁-C₆)-alkyl chain terminally substituted with phenyl or pyridinyl;

the other of A or B is independently selected from hydrogen, Ar, (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, Ar-substituted-(C₁-C₆)-straight or branched alkyl, or Ar-substituted-(C₂-C₆)-straight or branched alkenyl or alkynyl; wherein any one of the CH₂ groups of said alkyl, alkenyl or alkynyl chains in A or B is optionally replaced by O;

Ar is selected from phenyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, or thiazolyl, and wherein

each Ar is optionally substituted with one to three substituents independently selected from halogen, hydroxyl, nitro, --SO₃H, trifluoromethyl, trifluoromethoxy, (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl, O--[(C₁-C₆)-straight or branched alkyl], O--[(C₂-C₆)-straight or branched alkenyl], O-benzyl, O-phenyl, 1,2-methylenedioxy, --N(R¹)(R²), carboxyl, N--(C₁-C₅-straight or branched alkyl or C₂-C₅-straight or branched alkenyl) carboxamide, N,N-di-(C₁-C₅-straight or branched alkyl or C₂-C₅-straight or branched alkenyl) carboxamide, N--(C₁-C₅-straight or branched alkyl or C₂-C₅-straight or branched alkenyl) sulfonamide, N,N-di-(C₁-C₅-straight or branched alkyl or C₂-C₅-straight or branched alkenyl) sulfonamide, morpholinyl, piperidinyl, O-Z, CH₂--(CH₂)_q-Z, O--(CH₂)_q-Z, (CH₂)_q-Z-O-Z, or CH=CH-Z;

wherein R₁ and R₂ are independently selected from (C₁-C₆)-straight or branched alkyl, (C₂-C₆)-straight or branched alkenyl or alkynyl, hydrogen or

benzyl; or wherein R_1 and R_2 are taken together with the nitrogen atom to which they are bound to form a 5-7 membered heterocyclic ring;

Z is selected from 2-pyridyl, 3-pyridyl, or 4-pyridyl; and

q is 0, 1 or 2;

R_3 is selected from (C_1C_6) -alkyl, (C_1C_6) alkyl terminally substituted with pyridyl, or 3,4,5-trimethoxybenzoylmethyl;

X is N or O;

wherein when X is N, Y is selected from hydrogen, Ar, (C_1-C_6) -straight or branched alkyl, (C_2-C_6) -straight or branched alkenyl or alkynyl, Ar-substituted- (C_1-C_6) -straight or branched alkyl, or Ar-substituted- (C_1-C_6) -straight or branched alkenyl or alkynyl;

when X is O, Y is a lone pair of electrons;

n is 0;

m is 2; and

the indicated ring is fully saturated.

12. The compound of claim 11, wherein X is nitrogen.

13. The compound of claim 11 or 12, wherein Y is selected from hydrogen, (C_1-C_6) -straight or branched alkyl or Ar-substituted- (C_1-C_6) -straight or branched alkyl.

14. A composition comprising:

- a) a therapeutically effective amount of a compound according to claim 1;
- and
- b) a pharmaceutically suitable carrier.