

DESCRIPTION

SUGAR CHAIN-RELATED GENE AND USE THEREOF

5 Technical Field

The present invention relates to inhibitors of fibrogenesis at the physiological tissue level by inhibiting sugar chain-related genes.

Background Art

10 Intensive studies have been conducted on nucleic acids and proteins, revealing many findings. However, these studies also showed that there are only about 22,000 human genes and also that post-translational modification of proteins plays an important role *in vivo*. They also suggested limitations of conventional study approaches. In recent years, the importance of sugar chains has been rediscovered with the post-genome and post-proteomics trends (the journal
15 “Nature” extensively featured sugar chains in Vol. 446 published in April 26, 2007 (Non-patent Documents 1 to 7). Sugar chains have not been analyzed intensively because of the difficulty to perform structural analysis, etc. Although they are assumed to be involved in cancer, inflammation, immunity, viral infection, etc., at present, little is known about their roles and such, and therefore, elucidation is being awaited.

20 There are various known sugars (monosaccharides) that constitute sugar chains. Such known sugars include, for example, glucose (Glc), galactose (Gal), mannose (Man), glucuronic acid (GlcUA), iduronic acid (IdoA), fucose (Fuc), glucosamine (GlcN), N-acetylglucosamine (GlcNAc), N-acetylgalactosamine (GalNAc), xylose (Xyl), and sialic acid (SA).

 Furthermore, it has been reported that sugars constituting sugar chains are subject to a
25 variety of chemical modifications. Such chemical modifications include, for example, methylation, acetylation, formylation, myristoylation, amidation, ubiquitination, acylation, phosphorylation, epimerization, and sulfation. Examples of chemical modifications also include sialylation, asialylation, fucosylation, glycosylation, galactosylation, lactosylation, and mannosylation.

30 Sugars have a number of sites for such chemical modifications. For example, it is known that GlcNAc can be chemically-modified in any of carbons at positions 1 to 6. It is reported that other sugars are also chemically-modified at various sites.

[Non-patent Document 1] Danica P. Galonic and David Y. Gin, Nature 446: 1000-1007
(2007)

35 [Non-patent Document 2] Christopher J. Thibodeaux *et al.*, Nature 446: 1008-1016
(2007)

[Non-patent Document 3] Gerald W. Hart *et al.*, Nature 446: 1017-1022 (2007)

[Non-patent Document 4] Ajit Varki, Nature 446: 1023-1029 (2007)

[Non-patent Document 5] Joseph R. Bishop *et al.*, Nature 446: 1030-1037 (2007)

[Non-patent Document 6] Christopher N. *et al.*, Nature 446: 1038-1045 (2007)

5 [Non-patent Document 7] Peter H. Seeberger and Daniel B. Werz, Nature 446: 1046-1051 (2007)

Disclosure of the Invention

[Problems to be Solved by the Invention]

10 The present invention is based on a new finding obtained through studies on sugar chain-related genes. An objective of the present invention is to provide novel uses of sugar chain-related genes. Specifically, the objective is to provide agents that suppress fibrogenesis at the physiological tissue level by inhibiting the function of sugar chain-related genes, and methods of screening for the agents.

15

[Means for Solving the Problems]

Sugar chains have been suggested to play a very important role *in vivo*. However, little is known about the *in vivo* functions of sugar chains.

As described above, several types of sugars constituting sugar chains are known.

20 Sugars are variously chemically-modified at multiple sites, and such modifications are considered to assume important physiological effects *in vivo*.

As described above, there are various types of sugars that constitute sugar chains, many types of chemical modifications that target sugars, and many chemical modification sites in the sugars. This suggests that possible sugar chain variations are innumerable. Thus, it is very
25 difficult to determine a sugar chain structure that plays a certain role, and it is also extremely difficult to reveal the relationship between a pathological condition caused by a disease and a specific action of a sugar chain.

The present inventors conducted dedicated studies, and as a result succeeded for the first time in discovering that tissue fibrogenesis can be suppressed at the physiological level by
30 inhibiting sulfation at position 4 or 6 of GalNAc, a sugar that constitutes sugar chains. Furthermore, by studies using various animal disease models, the present inventors demonstrated that inhibitors of sulfation at position 4 or 6 of GalNAc produce therapeutic effects against diseases caused by tissue fibrogenesis (tissue fibrogenic disorders).

35 The present invention relates to agents that suppress fibrogenesis at the physiological tissue level by inhibiting the functions of sugar chain-related genes, and methods of screening for the agents. Specifically, the present invention provides:

- [1] a tissue fibrogenesis suppressing-agent, which comprises as an ingredient an inhibitor of sulfation at position 4 or 6 of N-acetylgalactosamine;
- [2] the agent of [1], which has an effect of suppressing fibrogenesis of a physiological tissue;
- [3] the agent of [1] or [2], wherein the inhibitor has the activity of inhibiting the function of a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine;
- [4] the agent of [3], wherein the inhibitor is an siRNA that suppresses the expression of the sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine;
- [5] the agent of [1] or [2], wherein the inhibitor is a desulfating enzyme that desulfates at position 4 or 6 of N-acetylgalactosamine;
- [6] the agent of any one of [1] to [5] for treating or preventing a fibrogenic disorder;
- [7] a method of screening for a tissue fibrogenesis suppressing-agent, which comprises the step of selecting a compound that inhibits sulfation at position 4 or 6 of N-acetylgalactosamine that constitutes a sugar chain;
- [8] a method of screening for a tissue fibrogenesis suppressing-agent, which comprises the steps of:
 - (a) contacting a test compound with N-acetylgalactosamine or a sugar chain comprising N-acetylgalactosamine;
 - (b) determining the degree of sulfation at position 4 or 6 of N-acetylgalactosamine; and
 - (c) selecting a compound that reduces the degree of sulfation as compared to when the test compound is not contacted;
- [9] a method of screening for a tissue fibrogenesis suppressing-agent, which comprises the steps of:
 - (a) contacting a test compound with a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine;
 - (b) determining the sulfotransferase activity of the enzyme; and
 - (c) selecting a compound that reduces the activity as compared to when the test compound is not contacted;
- [10] a method of screening for a tissue fibrogenesis suppressing-agent, which comprises the steps of:
 - (a) contacting a test compound with a cell expressing a gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine;
 - (b) determining the expression level of the gene in the cell; and
 - (c) selecting a compound that reduces the expression level of the gene as compared to when the test compound is not contacted;
- [11] a method of screening for a tissue fibrogenesis suppressing-agent, which comprises the steps of:

(a) contacting a test compound with a cell or cell extract containing a DNA wherein a reporter gene is operably linked to the transcriptional regulatory region of the gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine;

(b) determining the expression level of the reporter gene; and

5 (c) selecting a compound that reduces the expression level of the reporter gene as compared to when the test compound is not contacted; and

[12] a method of producing a pharmaceutical composition for treating or preventing a fibrogenic disorder, which comprises the steps of:

10 (a) selecting a tissue fibrogenesis suppressing-agent from test compounds by the method of any one of claims 7 to 11; and

(b) combining the agent with a pharmaceutically acceptable carrier.

The present invention also provides:

[13] a method of suppressing tissue fibrogenesis, which comprises the step of administering an inhibitor of sulfation at position 4 or 6 of N-acetylgalactosamine to an individual;

15 [14] use of an inhibitor of sulfation at position 4 or 6 of N-acetylgalactosamine in the manufacture of a tissue fibrogenesis suppressing-agent; and

[15] an inhibitor of sulfation at position 4 or 6 of N-acetylgalactosamine for use in suppressing tissue fibrogenesis.

20 Brief Description of the Drawings

Fig. 1 shows the result of quantitative reverse transcription PCR method for gene expressions of cardiomyopathy model mice in which cardiomyopathy was induced by intratracheal administration of DOX. The examined items were GalNAc4S-6ST as an siRNA target gene, and α -SMA and type I collagen, both of which are fibrogenesis markers. The graph indicates

25 relative ratios between a target gene and a house keeping gene (ribosome 18S). * $P < 0.001$ (t-test)

Fig. 2 shows the result analyzing the heart weight/body weight ratio of cardiomyopathy model mice in which cardiomyopathy was induced by intraperitoneal administration of Doxorubicin hydrochloride (DOX; Kyowa Hakko). * $P < 0.05$ (t-test)

30 Fig. 3 depicts photographs showing histological observations indicating the suppressive effect on type I collagen deposition in an siRNA-treated group of cardiomyopathy model mice in which cardiomyopathy was induced by intratracheal administration of DOX. The magnification is 100 fold.

35 Fig. 4 depicts photographs showing histological observations indicating the suppressive effect on type III collagen deposition in an siRNA-treated group of cardiomyopathy model mice in which cardiomyopathy was induced by intratracheal administration of DOX. The magnification is 100 fold.

Fig. 5 depicts photographs showing the suppressive effect on fibroblast accumulation in an siRNA-treated group of cardiomyopathy model mice in which cardiomyopathy was induced by intratracheal administration of DOX. The magnification is 50 fold.

Fig. 6 depicts graphs showing clinical features of a mouse intestinal fibrogenesis model, in which intestinal fibrogenesis was induced by DSS. The administration of GalNAc4S-6ST siRNA resulted in significant reduction of clinical score (left) and colon shortening (right).

Fig. 7 depicts graphs showing the expression of fibrogenesis-related genes in a mouse intestinal fibrogenesis model. GalNAc4S-6ST, and α -SMA and type I collagen, both of which are fibrogenesis markers, were assessed for the expression in colonic tissue. The graphs indicate relative ratios between a target gene and a house keeping gene (ribosome 18S). The enhanced expression of type I collagen (left, bottom) and α -SMA (right, bottom) is significantly suppressed due to the silencing effect of GalNAc4S-6ST siRNA (upper).

Fig. 8 depicts photographs showing collagen deposition in a mouse intestinal fibrogenesis model. Images of Masson-stained colonic tissues (blue). The magnification is 100 fold. GalNAc4S-6ST siRNA reduces the collagen deposition in tissues.

Fig. 9 depicts photographs showing fibroblast infiltration in a mouse model for intestinal fibrogenesis. Images of stained fibroblasts (brown) in colonic tissues. The magnification is 100 fold. GalNAc 4S-6ST siRNA suppresses the full-thickness infiltration of fibroblasts.

Fig. 10 depicts photographs showing images of macrophage infiltration in a mouse intestinal fibrogenesis model. Images of stained macrophages (brown) in colonic tissues. Magnification: 100x. GalNAc4S-6ST siRNA suppresses the full-thickness infiltration of macrophages.

Fig. 11 depicts a graph showing colon lengths in a mouse intestinal fibrogenesis model. GalNAc4ST siRNA significantly suppresses colon shortening.

Fig. 12 depicts photographs showing clinical images of a mouse emphysema model, in which emphysema was induced by PPE. C6ST-1 siRNA administration significantly suppresses the histological disruption of the pulmonary alveolar wall. Magnification: 100x.

Fig. 13 depicts graphs showing the expression of fibrogenesis-related genes in a mouse emphysema model. GalNAc4S-6ST, and α -SMA and type I collagen, both of which are fibrogenesis markers, were assessed for the expression in pulmonary tissues. The graphs indicate relative ratios between a target gene and a house keeping gene (ribosome 18S). The enhanced expression of type I collagen (middle) and α -SMA (right) is significantly suppressed due to the silencing effect of C6ST-1 siRNA (left).

Fig. 14 depicts photographs showing fibroblast infiltration in a mouse emphysema model. Images of stained fibroblasts (brown) in pulmonary tissues. Magnification: 200x. C6ST-1 siRNA suppresses fibroblast infiltration into the pulmonary alveolar interstitium.

Fig. 15 depicts photographs showing images of macrophage infiltration in a mouse emphysema

model. Images of stained macrophages (brown) in pulmonary tissues. Magnification: 200x. C6ST-1 siRNA suppresses macrophage infiltration into the pulmonary alveolar interstitium. Fig. 16 depicts a graph showing static lung compliance (Cst) in a mouse emphysema model. C6ST-1 siRNA significantly reduces Cst.

5 Fig. 17 depicts a graph showing the right lung volume (μ l) in a mouse emphysema model. C6ST-1 siRNA significantly reduces lung volume.

Fig. 18 depicts graphs showing the expression of fibrogenesis-related genes in a mouse emphysema model. α -SMA, type I collagen, and TGF- β , which are fibrogenesis markers, were assessed for the expression in pulmonary tissues. The graphs indicate relative ratios between a target gene and a house keeping gene (ribosome 18S). The enhanced expression of each

10 fibrogenesis marker is significantly suppressed due to the silencing effect of GalNAcST siRNA. Fig. 19 depicts a graph showing static lung compliance (Cst) in a mouse emphysema model. GalNAcST siRNA significantly reduces Cst.

Fig. 20 depicts a graph showing the right lung volume (μ l) in a mouse emphysema model. GalNAcST siRNA significantly reduces lung volume.

15 Fig. 21 depicts a graph showing obesity changes in a mouse type 2 diabetes model. C4ST-1, C4ST-2, and C4ST-3 siRNAs tends to suppress obesity. C4ST-2 and C4ST-3 siRNAs produce a significant anti-obesity effect.

Fig. 22 depicts a graph showing insulin resistance in a mouse type 2 diabetes model. C4ST-1, C4ST-2, and C4ST-3 siRNAs significantly improve insulin resistance.

20 Fig. 23 depicts photographs showing gene expressions in pancreatic tissues in a mouse type 2 diabetes model. C4ST-1, C4ST-2, and C4ST-3 siRNAs suppress the expression of C4ST-1, C4ST-2, and C4ST-3 genes in pancreatic tissues.

Fig. 24 depicts photographs showing APP deposition in pancreatic islets in a mouse type 2 diabetes model. C4ST-2 siRNA suppresses the deposition of APP (green) in islets. Cell nucleus (red); magnification: 400x.

Fig. 25 depicts photographs showing fibroblast infiltration in pancreatic islets in a mouse type 2 diabetes model. C4ST-1, C4ST-2, C4ST-3 siRNAs suppress fibroblast infiltration (brown) into islets. Magnification: 200x.

30 Fig. 26 depicts photographs showing macrophage infiltration into pancreatic islets in a mouse type 2 diabetes model. C4ST-1, C4ST-2, and C4ST-3 siRNAs suppress macrophage infiltration (brown) into islets. Magnification: 200x.

Fig. 27 depicts a graph showing insulin resistance in a mouse type 2 diabetes model. GalNAcST siRNA administration decreases the blood glucose level well after insulin loading. Specifically,

35 this shows the improvement of insulin resistance. Fig. 28 depicts a graph and photographs showing the accumulation of fibroblasts in the

interstitium of kidney tissue in a mouse diabetic nephropathy model. C4ST-1 siRNA significantly suppresses the accumulation of fibroblasts (brown). Magnification: 200x.

Fig. 29 depicts a graph and photographs showing the accumulation of macrophages in the interstitium of kidney tissue in a mouse model for diabetic nephropathy. C4ST-1 siRNA

5 significantly suppresses the accumulation of macrophages (brown). Magnification: 200x.

Fig. 30 depicts photographs showing the accumulation of α SMA-positive cells in the interstitium of kidney tissue in a mouse diabetic nephropathy model. C4ST-1 siRNA significantly suppresses the accumulation of α SMA-positive cells. Magnification: 200x.

Fig. 31 depicts graphs showing an anti-fibrogenic effect in a mouse diabetic nephropathy model.

10 GalNAc4S-6ST (G#1) siRNA administration significantly suppresses the enhanced expression of GalNAc4S-6ST (G#1), α SMA, and TGF β in kidney tissues. ARB; angiotensin receptor antagonist (Valsartan).

Fig. 32 depicts a graph and photographs showing fibroblast accumulation in the interstitium of kidney tissue in a mouse diabetic nephropathy model. GalNAc4S-6ST (G#1) siRNA

15 significantly suppresses the accumulation of fibroblasts (brown). Magnification: 200x.

Fig. 33 depicts a graph and photographs showing macrophage accumulation in the interstitium of kidney tissue in a mouse diabetic nephropathy model. GalNAc4S-6ST (G#1) siRNA significantly suppresses the accumulation of macrophages (brown). Magnification: 200x.

Fig. 34 depicts a graph and photographs showing the accumulation of type IV collagen in a

20 mouse diabetic nephropathy model. GalNAc4S-6ST (G#1) siRNA significantly suppresses the thickening of glomerular basement membrane, which can be confirmed by the positivity for type IV collagen (brown). Magnification: 400x.

Fig. 35 depicts graphs showing the renal protective effect in a mouse diabetic nephropathy model.

25 GalNAc4S-6ST (G#1) siRNA significantly suppresses the enhanced expression of angiotensinogen and ACE in kidney tissues.

Fig. 36 depicts a graph showing the renal function protective effect in a mouse diabetic nephropathy model. GalNAc4S-6ST (G#1) siRNA significantly suppresses the increase of serum creatinine, i.e., suppresses decline of renal function.

Fig. 37 depicts graphs showing gene expression in a mouse model for diabetic nephropathy.

30 GalNAcST siRNA significantly suppresses the enhanced expression of GalNAc4ST-1, GalNAc4ST-2, and GalNAc4S-6ST in kidney tissues.

Fig. 38 depicts graphs showing an anti-fibrogenic effect in a mouse diabetic nephropathy model. GalNAcST siRNA significantly suppresses the enhanced expression of CTGF, α SMA, type I collagen, and ACE in kidney tissues.

35 Fig. 39 depicts a graph showing gene expression in mice with drug-induced interstitial nephritis. GalNAc4S-6ST (G#1) siRNA significantly suppresses the enhanced expression of GalNAc4S-

6ST (G#1) in kidney tissues.

Fig. 40 depicts photographs showing collagen deposition in mice with drug-induced interstitial nephritis. GalNac4S-6ST (G#1) siRNA significantly decreases the deposition of type I collagen (brown) in renal interstitium. Magnification: 200x.

- 5 Fig. 41 depicts graphs showing an anti-fibrogenic effect in a mouse UUO fibrogenesis model. C6ST siRNA significantly suppresses the enhanced expression of C6ST-2 (G#10), TGF β , α SMA, type I collagen, and CTGF in kidney tissues.

Fig. 42 depicts a graph and photographs showing fibroblast accumulation in the interstitium in a mouse UUO fibrogenesis model. C6ST siRNA significantly suppresses fibroblast accumulation (brown) in juxtaglomerular and interstitial area. Magnification: 200x.

Fig. 43 depicts a graph and photographs showing macrophage accumulation in the interstitium in a mouse UUO fibrogenesis model. C6ST siRNA significantly suppresses the accumulation of macrophages (brown) in juxtaglomerular and interstitial area. Magnification: 200x.

- Fig. 44 depicts a graph and photographs showing collagen deposition in a mouse UUO fibrogenesis model. C6ST siRNA significantly suppresses the thickening of glomerular basement membrane, which can be confirmed by the positivity of type IV collagen (brown). Magnification: 400x.

Fig. 45 depicts photographs showing the activation of fibroblasts accumulated in tissues in a mouse UUO fibrogenesis model. C6ST siRNA significantly suppresses the accumulation of α SMA-positive cells (brown) in juxtaglomerular and interstitial area. Magnification: 400x.

Fig. 46 depicts photographs showing the accumulation of ACE-producing cells in the interstitium in a mouse UUO fibrogenesis model. C6ST siRNA significantly suppresses the accumulation of ACE-producing cells (brown) in juxtaglomerular and interstitial area. Magnification: 400x.

- 25 Fig. 47 depicts photographs showing the tissue accumulation of type IV collagen in a mouse diabetic retinopathy model. GalNac4S-6ST (G#1) siRNA significantly suppresses the accumulation of type IV collagen. Magnification: 200x. GCL, ganglion cell layer; INL, inner nuclear layer; ONL, external granular layer.

Fig. 48 depicts photographs showing the tissue accumulation of CSPG in a mouse diabetic retinopathy model. GalNac4S-6ST (G#1) siRNA significantly suppresses CSPG accumulation (brown). In particular, the suppressing effect is prominent in GCL, ONL, and pigment epithelial cell layer. Magnification: 200x.

Fig. 49 depicts photographs showing GFAP-positive cells in a mouse diabetic retinopathy model. GalNac4S-6ST (G#1) siRNA significantly increases GFAP-positive cells (brown) in the region from INL to GCL. Magnification: 200x.

Fig. 50 depicts a graph showing the number of gangliocytes in a mouse diabetic retinopathy

model. Counts show the number of gangliocytes in GGL. GalNac4S-6ST (G#1) siRNA significantly suppresses the reduction in gangliocyte number.

Fig. 51 depicts graphs showing the optic nerve regeneration effect in a mouse diabetic retinopathy model. GalNac4S-6ST (G#1) siRNA significantly increases the expression of GS in ocular tissues.

Fig. 52 depicts a graph showing gene expression in a mouse fatty liver injury model. The enhanced expression of GalNac4S-6ST in liver tissues, and significant suppression of the expression by GalNacST siRNA are shown.

Fig. 53 depicts graphs showing the expression of fibrogenesis-related genes in a mouse fatty liver injury model. The enhanced expression of type I collagen and α SAM in liver tissues, and significant suppression of the expression by GalNacST siRNA are shown.

Fig. 54 depicts photographs showing the infiltration of fibrogenic cells in a mouse fatty liver injury model. GalNacST siRNA suppresses the bridge-like accumulation of fibroblasts (brown) in liver tissues. Magnification: 100x.

Fig. 55 depicts a graph showing clinical scores for fibrogenesis in a mouse fatty liver injury model. GalNacST siRNA significantly suppresses the increase in the fibrogenesis score in liver tissues.

Fig. 56 depicts photographs showing macrophage infiltration in a mouse fatty liver injury model. GalNacST siRNA significantly suppresses the accumulation of macrophages (brown) in liver tissues. Magnification: 100x.

Fig. 57 depicts graphs showing the expression of lipid metabolism-related genes in a mouse fatty liver injury model. GalNacST siRNA significantly suppresses the increased expression of ChREBP and ACC2 in liver tissues.

Fig. 58 depicts photographs showing the infiltration of fibrogenic cells in a mouse fatty liver injury model. C4ST-1, C4ST-2, and C4ST-3 siRNAs suppress the bridge-like accumulation of fibroblasts (brown) in liver tissues. Magnification: 100x.

Fig. 59 depicts a graph showing clinical scores for fibrogenesis in a mouse fatty liver injury model. C4ST-1, C4ST-2, and C4ST-3 siRNAs significantly suppress the increase in the fibrogenesis score in liver tissues.

Fig. 60 depicts a graph showing clinical hepatic disorder in a mouse fatty liver injury model. C4ST-1, C4ST-2, and C4ST-3 siRNAs suppress the increase in ALT, which is an indicator of hepatocyte disorder.

Fig. 61 depicts photographs showing the infiltration of fibroblasts in a mouse hepatic fibrosis model. C6ST siRNA significantly suppresses the accumulation of fibroblasts (brown) in liver tissues. Magnification: 50x.

Fig. 62 depicts a graph showing clinical scores for fibrogenesis in a mouse hepatic fibrosis

model. C6ST siRNA significantly suppresses the increase in the clinical fibrogenesis score in liver tissues.

Fig. 63 depicts graphs showing the anti-fibrogenic effect in a mouse hepatic fibrosis model.

C6ST siRNA significantly suppresses the expression of α SMA, type I collagen, CTGF, and TGF β .

Fig. 64 depicts graphs showing the anti-fibrogenic effect in a mouse Parkinson's disease model.

GalNAc4S-6ST siRNA significantly suppresses the expression of GalNAc4S-6ST, TGF β , type I collagen, and α SMA in brain tissues.

Fig. 65 depicts photographs showing the accumulation of fibroblasts in a mouse Parkinson's

disease model. GalNAc4S-6ST siRNA drastically decreases the accumulation of fibroblasts (brown) in brain tissues. Magnification: 200x.

Fig. 66 depicts graphs showing the nerve protective effect in a mouse Parkinson's disease model.

GalNAc4S-6ST siRNA significantly enhances the expression of GDNF and Nurr1.

Fig. 67 depicts photographs showing the dopamine neuron regeneration effect in a mouse

Parkinson's disease model. GalNAc4S-6ST siRNA suppresses the degeneration of TH-positive dopamine neurons (green). Magnification: 200x.

Fig. 68 depicts photographs showing the dopamine neuron regeneration effect in a mouse Parkinson's disease model. GalNAc4ST siRNA suppresses the degeneration of TH-positive dopamine neurons (green). Magnification: 200x.

Fig. 69 depicts photographs showing the CSPG-reducing effect of C4-sulfatase in a mouse type 2 diabetic retinopathy model. The photographs show images of stained CSPG (CS56) (brown, arrow) in the retina of type 2 diabetic retinopathy model mice.

Fig. 70 depicts photographs showing the angiogenesis-suppressing effect of C4-sulfatase in a mouse type 2 diabetic retinopathy model. The photographs show images of stained vascular endothelial cells (CD31) (brown, arrow) in the retina of type 2 diabetic retinopathy model mice.

Fig. 71 depicts photographs showing the collagen augmentation-suppressing effect of C4-sulfatase in a mouse type 2 diabetic retinopathy model. The photographs show images of stained type IV collagen (brown, arrow) in the retina of type 2 diabetic retinopathy model mice.

Fig. 72 depicts photographs showing the suppressive effect of C4-sulfatase on fibroblast accumulation in the liver of a mouse type 2 diabetes model. The magnification is 50 or 100 fold.

Fig. 73 depicts photographs showing the macrophage infiltration-suppressing effect in the liver of a mouse type 2 diabetes model. The magnification is 50 or 100 fold.

Fig. 74 depicts graphs showing the result of serum biochemical tests (AST, ALT, and TG) in a mouse type 2 diabetes model. In the graphs: unt, untreated group; nor, control group; C4sul, C4-sulfatase.

Fig. 75 depicts photographs showing the localization of CSPG in brain tissues. The photographs

show the result of analyzing the dynamics of CSPG expression in brain tissues using an enzyme-antibody immunostaining method. Assay was carried out using CS-56 (Seikagaku Co.) for primary antibody and Mouse Stain Kit for staining.

Fig. 76 depicts photographs showing the localization of dopaminergic neurons in brain tissues.

5 The photographs show an analysis result obtained by a fluorescence immunostaining method.

Fig. 77 depicts a graph showing the result of TNF- α gene expression analysis, which was obtained by Real-time PCR method. The graph shows the result of real-time PCR for the expression of TGF- β as a fibrosis marker, and TNF- α as an indicator for inflammation associated with macrophage infiltration.

10 Fig. 78 depicts a graph showing the result of Nurr1 gene expression analysis by real-time PCR method. The graph shows the result for Nurr1 gene expression in brain tissues, which was obtained using Cyber premix kit (Takara Bio) and Real-time PCR thermal cycler DICE (Takara Bio). The graph indicates relative ratios between Nurr1 and a house keeping gene (β -actin).

15 Mode for Carrying Out the Invention

The present invention will be specifically described below.

The present invention relates to tissue fibrogenesis inhibitors based on the mechanism of inhibiting sulfation at position 4 or 6 of N-acetylgalactosamine, a sugar constituting sugar chains.


20 Specifically, the present invention provides tissue fibrogenesis inhibitors comprising as an ingredient an inhibitor of sulfation at position 4 or 6 of N-acetylgalactosamine (herein sometimes also referred to as “inhibitors of the present invention” or simply as “inhibitors”).

Herein, “N-acetylgalactosamine (GalNAc)” refers to the N-acetylated form of galactosamine, which is a hexosamine.

25 Furthermore, it is known that N-acetylgalactosamine can be chemically-modified at positions 1 to 6.

The present invention is characterized by inhibiting sulfation at position 4 or 6 of N-acetylgalactosamine.

30 Specifically, the sites where the sulfation is inhibited by the inhibitors of the present invention are indicated by arrow in the following formula of GalNAc.



The sites where the sulfation is inhibited by the inhibitors of the present invention are

positions 4 or 6 of GalNAc. In the present invention, the sulfation of GalNAc may be inhibited at both positions 4 and 6. In the present invention, preferred GalNAc is a sugar in chondroitin sulfate proteoglycan (CSPG).

Herein, the inhibition of sulfation refers to inhibition of transfer of a sulfate group to position 4 or 6 in GalNAc, elimination of a sulfate group from a site where GalNAc has been already sulfated, or substitution of a sulfate group with other chemically-modified group.

The agents for suppressing tissue fibrogenesis of the present invention (herein sometimes referred to as “agents of the present invention”) preferably have an *in vivo* fibrogenesis-suppressing effect.

Tissues where fibrogenesis is suppressed by the agents of the present invention are not particularly limited. Such tissues include, for example, cardiac tissues, gastrointestinal tissues, lung tissues, pancreatic tissues, kidney tissues, ocular tissues, liver tissues, cranial nerve tissues, and skin tissues.

Herein, “fibrogenesis” may be referred to as “fibrosis”. Alternatively, “fibrogenesis” may be synonymous with other phrases such as “fibrogenic lesion in tissues”, “fibrogenic tissue alteration”, and “neofibrogenesis”.

The inhibitors of the present invention are not particularly limited as long as they are substances having the activity of inhibiting the sulfation at position 4 or 6 of N-acetylgalactosamine.

A preferred embodiment of inhibitors of the present invention includes, for example, substances having the activity of inhibiting the function of sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine.

Preferred embodiments of the above-described substances include, for example, compounds (nucleic acids) selected from the group consisting of:

- (a) antisense nucleic acids against transcripts of the genes encoding the sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine, or portions thereof;
- (b) nucleic acids with the ribozyme activity of specifically cleaving transcripts of genes encoding the sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine; and
- (c) nucleic acids with the activity of using RNAi effect to inhibit the expression of genes encoding the sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine (siRNAs that suppress the expression of sulfotransferase genes).

The “substances with the activity of inhibiting sulfation” also include, for example, compounds selected from the group consisting of:

- (a) antibodies that bind to sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine;
- (b) sulfotransferase variants for sulfotransferases that transfer a sulfate to position 4 or 6 of N-

acetylgalactosamine; and

(c) low-molecular-weight compounds that bind to sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine.

Another embodiment of inhibitors of the present invention includes, for example, substances having the activity of desulfating a sulfate group at position 4 or 6 of N-acetylgalactosamine. Such substances include, for example, enzymes that desulfate the sulfate group (desulfating enzymes) at position 4 or 6 of N-acetylgalactosamine.

The “desulfating” sulfate group at position 4 or 6 of N-acetylgalactosamine means that a sulfate group at position 4 or 6 is eliminated from N-acetylgalactosamine.

Such desulfating enzymes include, for example, chondroitin-4-sulfatase (C4-sulfatase) and chondroitin-6-sulfatase.

Sulfotransferases of the present invention are not particularly limited as long as enzymes have an activity of transferring a sulfate to position 4 or 6 of GalNAc, but include, for example:

1) GalNAc4ST-1: N-acetylgalactosamine 4-sulfotransferase-1

Alias CHST8: Carbohydrate (N-acetylgalactosamine 4-O) sulfotransferase 8

2) GalNAc4ST-2

Alias CHST9: Carbohydrate (N-acetylgalactosamine 4-O) sulfotransferase 9

3) C4ST-1: chondroitin-4-O-sulfotransferase-1

Alias CHST11: Carbohydrate (chondroitin 4) sulfotransferase 11

4) C4ST-2

Alias CHST12

5) C4ST-3

Alias CHST13

6) C6ST-1: chondroitin-6-O-sulfotransferase-1

Alias CHST3: Carbohydrate (chondroitin 6) sulfotransferase 3

7) GalNAc4S-6ST: N-acetylgalactosamine 4-sulfate 6-O sulfotransferase

8) D4ST-1: dermatan 4 sulfotransferase 1

9) C6ST-2: chondroitin-6-O-sulfotransferase-2

Alias CHST7: Carbohydrate (chondroitin 6) sulfotransferase 7

Further, on a genomic DNA level, such groups of enzymes sharing features do not necessarily correspond to single genes. For example, both chondroitin-4-sulfatase and chondroitin-6-sulfatase can be retrieved from the public gene database GenBank as sequences referred to by multiple accession numbers (for example, GenBank accession Nos: NT_039500 (a portion thereof is shown under accession No: CAAA01098429 (SEQ ID NO: 1)), NT_078575, NT_039353, NW_001030904, NW_001030811, NW_001030796, and NW_000349).

Specifically, below are examples of sulfotransferases of the present invention with accession numbers in the public gene database GenBank, nucleotide sequences, and amino acid sequences:

- GalNAc4ST-1 (Accession number NM_175140; nucleotide sequence: SEQ ID NO: 2; amino acid sequence, SEQ ID NO: 3)
- GalNAc4ST-2 (Accession number NM_199055; nucleotide sequence: SEQ ID NO: 4; amino acid sequence, SEQ ID NO: 5)
- C4ST-1 (Accession number NM_021439; nucleotide sequence: SEQ ID NO: 6, amino acid sequence, SEQ ID NO: 7)
- C4ST-2 (Accession number NM_021528; nucleotide sequence: SEQ ID NO: 8; amino acid sequence, SEQ ID NO: 9)
- C4ST-3 (Accession number XM_355798; nucleotide sequence: SEQ ID NO: 10; amino acid sequence, SEQ ID NO: 11)
- D4ST (Accession number NM_028117; nucleotide sequence: SEQ ID NO: 12; amino acid sequence, SEQ ID NO: 13)
- C6ST-1 (Accession number NM_016803; nucleotide sequence: SEQ ID NO: 14; amino acid sequence, SEQ ID NO: 15)
- C6ST-2 (Accession number AB046929; nucleotide sequence: SEQ ID NO: 16; amino acid sequence, SEQ ID NO: 17)
- GalNAc4S-6ST (Accession number NM_015892; nucleotide sequence: SEQ ID NO: 18; amino acid sequence, SEQ ID NO: 19)

In addition to the proteins listed above, the proteins of the present invention include those exhibiting high homology (typically 70% or higher, preferably 80% or higher, more preferably 90% or higher, and most preferably 95% or higher) to sequences shown in the Sequence Listing and having a function of the proteins listed above (for example, the function of binding to intracellular components). The proteins listed above are, for example, proteins comprising an amino acid sequence with an addition, deletion, substitution, or insertion of one or more amino acids in any of the amino acid sequences of SEQ ID NOs: 3, 5, 7, 9, 11, 13, 15, and 17, in which the number of altered amino acids is typically 30 amino acids or less, preferably ten amino acids or less, more preferably five amino acids or less, and most preferably three amino acids or less.

The above-described genes of the present invention include, for example, endogenous genes of other organisms which correspond to DNAs comprising any of the nucleotide sequences of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, and 16 (homologues to the human genes described above, or the like).

Each of the endogenous DNAs of other organisms which correspond to DNAs

comprising any of the nucleotide sequences of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, and 16 are generally highly homologous to a DNA of any of SEQ ID NOs: 2, 4, 6, 8, 10, 12, 14, and 16. High homology means 50% or higher homology, preferably 70% or higher homology, more preferably 80% or higher homology, and still more preferably 90% or higher homology (for example, 95% or higher, or 96%, 97%, 98%, or 99% or higher). Homology can be determined using the mBLAST algorithm (Altschul, *et al.* Proc. Natl. Acad. Sci. USA, 1990, 87, 2264-8; Karlin and Altschul, Proc. Natl. Acad. Sci. USA, 1993, 90, 5873-7). When the DNAs have been isolated from the body, each of them may hybridize under stringent conditions to a DNA of SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, and 16. Herein, stringent conditions include, for example, "2x SSC, 0.1% SDS, 50°C", "2x SSC, 0.1% SDS, 42°C", and "1x SSC, 0.1% SDS, 37°C"; more stringent conditions include "2x SSC, 0.1% SDS, 65°C", "0.5x SSC, 0.1% SDS, 42°C", and "0.2x SSC, 0.1% SDS, 65°C".

Those skilled in the art can appropriately obtain proteins functionally equivalent to the above-described proteins from the above-described highly homologous proteins by using methods for assaying the activity of desulfating or inhibiting the sulfation at position 4 or 6 of N-acetylgalactosamine.

Further, based on the nucleotide sequences of the above-described genes, those skilled in the art can appropriately obtain endogenous genes of other organisms that correspond to the above-described genes. In the present invention, the above-described proteins and genes in non-human organisms, which correspond to the above-described proteins and genes, or the above-described proteins and genes that are functionally equivalent to the above-described proteins and genes, may simply be referred to using the above-described names.

The proteins of the present invention can be prepared not only as natural proteins but also as recombinant proteins using genetic recombination techniques. The natural proteins can be prepared by, for example, methods of subjecting cell extracts (tissue extracts) that may express the above-described proteins to affinity chromatography using antibodies against the above-described proteins. On the other hand, the recombinant proteins can be prepared, for example, by culturing cells transformed with DNAs encoding the proteins described above. The above-described proteins of the present invention can be suitably used, for example, in the screening methods described herein below.

In the present invention, "nucleic acids" refer to both RNAs and DNAs. Chemically synthesized nucleic acid analogs, such as so-called "PNAs" (peptide nucleic acids), are also included in the nucleic acids of the present invention. PNAs are nucleic acids in which the fundamental backbone structure of nucleic acids, the pentose-phosphate backbone, is replaced by a polyamide backbone with glycine units. PNAs have a three-dimensional structure quite similar to that of nucleic acids.

Methods for inhibiting the expression of specific endogenous genes using antisense technology are well known to those skilled in the art. There are a number of causes for the action of antisense nucleic acids in inhibiting target gene expression, including:

inhibition of transcription initiation by triplex formation;

5 transcription inhibition by hybrid formation at a site with a local open loop structure generated by an RNA polymerase;

transcription inhibition by hybrid formation with the RNA being synthesized;

splicing inhibition by hybrid formation at an intron-exon junction;

splicing inhibition by hybrid formation at the site of spliceosome formation;

10 inhibition of transport from the nucleus to the cytoplasm by hybrid formation with mRNA;

splicing inhibition by hybrid formation at the capping site or poly(A) addition site;

inhibition of translation initiation by hybrid formation at the translation initiation factor binding site;

15 inhibition of translation by hybrid formation at the ribosome binding site adjacent to the start codon;

inhibition of peptide chain elongation by hybrid formation in the translational region of mRNA or at the polysome binding site of mRNA; and

inhibition of gene expression by hybrid formation at the protein-nucleic acid interaction sites.

Thus, antisense nucleic acids inhibit the expression of target genes by inhibiting various

20 processes, such as transcription, splicing, and translation (Hirashima and Inoue, Shin Seikagaku

Jikken Koza 2 (New Courses in Experimental Biochemistry 2), Kakusan (Nucleic Acids) IV:

“Idenshi no Fukusei to Hatsugen (Gene replication and expression)”. Ed. The Japanese Biochemical Society, Tokyo Kagakudojin, 1993, pp. 319-347).

25 The antisense nucleic acids used in the present invention may inhibit the expression and/or function of genes encoding any of the sulfotransferases described above, based on any of the actions described above.

In one embodiment, antisense sequences designed to be complementary to an untranslated region adjacent to the 5' end of an mRNA for a gene encoding an above-described sulfotransferase may be effective for inhibiting translation of the gene. Sequences

30 complementary to a coding region or 3'-untranslated region can also be used. Thus, the antisense nucleic acids to be used in the present invention include not only nucleic acids comprising sequences antisense to the coding regions, but also nucleic acids comprising

sequences antisense to untranslated regions of genes encoding the above-described

sulfotransferases. Such antisense nucleic acids to be used are linked downstream of adequate

35 promoters and are preferably linked with transcription termination signals on the 3' side.

Nucleic acids thus prepared can be introduced into desired animals (cells) using known methods.

The sequences of the antisense nucleic acids are preferably complementary to a gene or portion thereof encoding a sulfotransferase that is endogenous to the animals (cells) to be transformed with them. However, the sequences need not be perfectly complementary, as long as the antisense nucleic acids can effectively suppress expression of a gene. The transcribed RNAs preferably have 90% or higher, and most preferably 95% or higher complementarity to target gene transcripts. To effectively inhibit target gene expression using antisense nucleic acids, the antisense nucleic acids are preferably at least 15 nucleotides long, and less than 25 nucleotides long. However, the lengths of the antisense nucleic acids of the present invention are not limited to the lengths mentioned above, and they may be 100 nucleotides or more, or 500 nucleotides or more.

The antisense nucleic acids of the preset invention are not particularly limited, and can be prepared, for example, based on the nucleotide sequence of C4ST-1 (GenBank Accession No: NM_021439; SEQ ID NO: 6), C4ST-2 (GenBank Accession No: NM_021528; SEQ ID NO: 8), C4ST-3 (GenBank Accession No: XM_355798; SEQ ID NO: 10), or such.

Expression of the above-mentioned genes encoding sulfotransferases can also be inhibited using ribozymes or ribozyme-encoding DNAs. Ribozymes refer to RNA molecules with catalytic activity. There are various ribozymes with different activities. Among others, studies that focused on ribozymes functioning as RNA-cleaving enzymes have enabled the design of ribozymes that cleave RNAs in a site-specific manner. Some ribozymes have 400 or more nucleotides, such as group I intron type ribozymes and M1 RNA, which is comprised by RNase P, but others, called hammerhead and hairpin ribozymes, have a catalytic domain of about 40 nucleotides (Koizumi, M. and Otsuka E., Tanpakushitsu Kakusan Koso (Protein, Nucleic Acid, and Enzyme), 1990, 35, 2191).

For example, the autocatalytic domain of a hammerhead ribozyme cleaves the sequence G13U14C15 at the 3' side of C15. Base pairing between U14 and A9 has been shown to be essential for this activity, and the sequence can be cleaved when C15 is substituted with A15 or U15 (Koizumi, M. *et al.*, FEBS Lett., 1988, 228, 228). Restriction enzyme-like RNA-cleaving ribozymes that recognize the sequence UC, UU, or UA in target RNAs can be created by designing their substrate-binding sites to be complementary to an RNA sequence adjacent to a target site (Koizumi, M. *et al.*, FEBS Lett., 1988, 239, 285; Koizumi, M. and Otsuka, E., Tanpakushitsu Kakusan Koso (Protein, Nucleic Acid, and Enzyme), 1990, 35, 2191; and Koizumi, M. *et al.*, Nucl Acids Res., 1989, 17, 7059).

In addition, hairpin ribozymes are also useful for the purposes of the present invention. Such ribozymes are found in, for example, the minus strand of satellite RNAs of tobacco ring spot viruses (Buzayan, J.M., Nature, 1986, 323, 349). It has been shown that target-specific RNA-cleaving ribozymes can also be created from hairpin ribozymes (Kikuchi, Y. and Sasaki,

N., Nucl Acids Res., 1991, 19, 6751; and Kikuchi, Y. Kagaku to Seibutsu (Chemistry and Biology), 1992, 30, 112). Thus, the expression of the above-described genes encoding sulfotransferases can be inhibited by using ribozymes to specifically cleave the gene transcripts.

The expression of endogenous genes can also be suppressed by RNA interference (hereinafter abbreviated as “RNAi”), using double-stranded RNAs comprising a sequence the same as or similar to a target gene sequence.

A great many disease-related genes have been rapidly identified since the entire human nucleotide sequence was revealed upon the recent completion of the genome project, and currently specific gene-targeted therapies and drugs are being actively developed. Of these, the application to gene therapy of small interfering RNAs (siRNAs), which produce the effect of specific post-transcriptional suppression, has been drawing attention.

RNAi is a phenomenon discovered by Fire *et al.* in 1998 (Fire, A., Nature (1998) 391: 806-811), where double strand RNA strongly suppresses expression of homologous target genes. RNAi has been drawing attention recently as a method applicable in gene therapy, because it is simpler than conventional gene transfer methods using vectors or such, and its target specificity is high. Furthermore, in mammalian cells, RNAi can be induced using short dsRNAs (siRNAs) and has many advantages: compared to knockout mice, RNAi has a stable effect, is easy to experiment with, has a low cost, and so on.

Nucleic acids with inhibitory activity based on the RNAi effect are generally referred to as siRNAs or shRNAs. RNAi is a phenomenon in which, when cells or such are introduced with short double-stranded RNAs (hereinafter abbreviated as “dsRNAs”) comprising sense RNAs that have sequences homologous to the mRNAs of a target gene, and antisense RNAs that comprise sequences homologous a sequence complementary thereto, the dsRNAs bind specifically and selectively to the target gene mRNAs, induce their disruption, and cleave the target gene, thereby effectively inhibiting (suppressing) target gene expression. For example, when dsRNAs are introduced into cells, the expression of genes with sequences homologous to the RNAs is suppressed (the genes are knocked down). As described above, RNAi can suppress the expression of target genes, and is thus drawing attention as a method applicable to gene therapy, or as a simple gene knockout method replacing conventional methods of gene disruption, which are based on complicated and inefficient homologous recombination.

In the present invention, the RNAs to be used in RNAi are not necessarily perfectly identical to the genes or portions thereof that encode an above-described sulfotransferase; however, the RNAs are preferably perfectly homologous to the genes or portions thereof. Furthermore, the terminal portion may include an overhang of about two bases.

The targets of the siRNAs to be designed are not particularly limited, as long as they are genes encoding an above-described sulfotransferase. Any region of the gene can be a candidate

for a target.

For example, siRNAs may be prepared based on a nucleotide sequence of C4ST-1 gene (SEQ ID NO: 6), C4ST-2 gene (SEQ ID NO: 8), C4ST-3 gene (SEQ ID NO: 10), and such.

More specifically, partial regions of such sequences may be used as candidates for the targets.

5 For example, siRNAs may be prepared based on portions of the nucleotide sequences of C4ST-1 gene (SEQ ID NO: 20), C4ST-2 gene (SEQ ID NO: 21), C4ST-3 gene (SEQ ID NO: 22), C6ST-1 gene (SEQ ID NO: 23), C6ST-2 gene (SEQ ID NO: 24), or such. More specifically, examples of the siRNAs also include those targeted to the DNA sequences (SEQ ID NOs: 25, 26, 35 to 50, 55 to 65, and 82 to 88) specifically shown herein.

10 The siRNAs can be introduced into cells by adopting methods of introducing cells with plasmid DNAs linked with siRNAs synthesized *in vitro* or methods that comprise annealing two RNA strands.

The two RNA molecules described above may be closed at one end or, for example, may be siRNAs with hairpin structures (shRNAs). shRNAs refer to short hairpin RNAs, which
15 are RNA molecules with a stem-loop structure, since a portion of the single strand constitutes a strand complementary to another portion. Thus, molecules capable of forming an intramolecular RNA duplex structure are also included in the siRNAs of the present invention.

In a preferred embodiment of the present invention, the siRNAs of the present invention also include, for example, double-stranded RNAs with additions or deletions of one or a few
20 RNAs in an siRNA which targets a specific DNA sequence (SEQ ID NOs: 25, 26, 35 to 50, 55 to 65, and 82 to 88) shown herein and which can suppress the expression of C4ST-1, C4ST-2, C4ST-3, or such *via* RNAi effect, as long as the double-stranded RNAs have the function of suppressing the expression of a gene encoding an above-described sulfotransferase.

The RNAs used in RNAi (siRNAs) do not need to be perfectly identical (homologous)
25 to the genes encoding the above proteins or portions thereof; however, the RNAs are preferably perfectly identical (homologous).

Some details of the RNAi mechanism still remain unclear, but it is understood that an enzyme called "DICER" (a member of the RNase III nuclease family) is contacted with a double-stranded RNA and degrades it into small fragments, called "small interfering RNAs" or
30 "siRNAs". The double-stranded RNAs of the present invention that have RNAi effect include such double-stranded RNAs prior to being degraded by DICER. Specifically, since even long RNAs that have no RNAi effect when intact can be degraded into siRNAs which have RNAi effect in cells, the length of the double-stranded RNAs of the present invention is not particularly limited.

35 For example, long double-stranded RNAs covering the full-length or near full-length mRNA of a gene encoding an above-described sulfotransferase can be pre-digested, for example,

by DICER, and then the degradation products can be used as agents of the present invention. These degradation products are expected to contain double-stranded RNA (siRNA) molecules with an RNAi effect. With this method, it is not necessary to specifically select the mRNA regions expected to have RNAi effect. In other words, it is not necessary to accurately determine regions with RNAi effect in the mRNAs of the genes described above.

The above-described “double-stranded RNAs capable of suppression *via* RNAi effect” can be suitably prepared by those skilled in the art based on nucleotide sequences of the above-described sulfotransferases, which are targeted by the double-stranded RNAs. For example, the double-stranded RNAs of the present invention can be prepared based on the nucleotide sequence of SEQ ID NO: 25. In other words, it is within the range of ordinary experimentation for those skilled in the art to select an arbitrary consecutive RNA region in an mRNA that is a transcript of the nucleotide sequence of SEQ ID NO: 25, and prepare double-stranded RNA corresponding to the region. Those skilled in the art can also use known methods to properly select siRNA sequences with stronger RNAi effect from the mRNA sequence, which is the transcript of the nucleotide sequence of SEQ ID NO: 25. When one of the strands is already identified, those skilled in the art can readily determine the nucleotide sequence of the other strand (complementary strand). Those skilled in the art can appropriately prepare siRNAs using a commercially available nucleic acid synthesizer. Alternatively, general custom synthesis services may be used to synthesize desired RNAs.

The siRNAs of the present invention are not necessarily single pairs of double-stranded RNAs directed to target sequences, but may be mixtures of multiple double-stranded RNAs directed to regions that cover the target sequence. Herein, those skilled in the art can appropriately prepare the siRNAs as nucleic acid mixtures matched to a target sequence by using a commercially available nucleic acid synthesizer or DICER enzyme. Meanwhile, general custom synthesis services may be used to synthesize desired RNAs. The siRNAs of the present invention include so-called “siRNA cocktails”.

All nucleotides in the siRNAs of the present invention do not necessarily need to be ribonucleotides (RNAs). Specifically, one or more of the ribonucleotides constituting the siRNAs of the present invention may be replaced with corresponding deoxyribonucleotides. The term “corresponding” means that although the sugar moieties are structurally differently, the nucleotide residues (adenine, guanine, cytosine, or thymine (uracil)) are the same. For example, deoxyribonucleotides corresponding to ribonucleotides with adenine refer to deoxyribonucleotides with adenine. The term “or more” described above is not particularly limited, but preferably refers to a small number of about two to five ribonucleotides.

Furthermore, DNAs (vectors) capable of expressing the RNAs of the present invention are also included in the preferred embodiments of compounds capable of suppressing the

expression of the genes encoding the above-described proteins of the present invention. The DNAs (vectors) capable of expressing the double-stranded RNAs of the present invention are, for example, DNAs structured such that a DNA encoding one strand of a double-stranded RNA and a DNA encoding the other strand of the double-stranded RNA are linked with promoters so that each DNA can be expressed. The above DNAs of the present invention can be appropriately prepared by those skilled in the art using standard genetic engineering techniques. More specifically, the expression vectors of the present invention can be prepared by adequately inserting DNAs encoding the RNAs of the present invention into various known expression vectors.

Furthermore, the expression-inhibiting substances of the present invention also include compounds that inhibit the expression of the above-described sulfotransferases by binding to an expression regulatory region of a gene encoding the above-described sulfotransferases (for example, a promoter region). Such compounds can be obtained, for example, using a fragment of a promoter DNA of the gene encoding an above-described sulfotransferase to perform screening methods using as an indicator the activity of binding to the DNA fragment. Those skilled in the art can appropriately determine whether compounds of interest inhibit the expression of the above-described genes encoding sulfotransferases by using known methods, for example, reporter assays and such.

Furthermore, DNAs (vectors) capable of expressing the above-described RNAs of the present invention are also included in preferred embodiments of the compounds capable of inhibiting the expression of a gene encoding an above-described sulfotransferase of the present invention. For example, DNAs (vectors) capable of expressing the above-described double-stranded RNAs of the present invention are structured such that a DNA encoding one strand of a double-stranded RNA and a DNA encoding the other strand of the double-stranded RNA are linked to promoters so that both can be expressed. Those skilled in the art can appropriately prepare the above-described DNAs of the present invention using standard genetic engineering techniques. More specifically, the expression vectors of the present invention can be prepared by appropriately inserting DNAs encoding the RNAs of the present invention into various known expression vectors.

Preferred embodiments of the above-described vector of the present invention include vectors expressing RNAs (siRNAs) that can suppress the expression of C4ST-1, C4ST-2, C4ST-3, or the like by the RNAi effect.

Antibodies that bind to the above-described sulfotransferases can be prepared by methods known to those skilled in the art. Polyclonal antibodies can be obtained, for example, by the following procedure: small animals such as rabbits are immunized with an above-described natural protein or a recombinant protein expressed in microorganisms as a fusion

protein with GST, or a partial peptide thereof. Sera are obtained from these animals and purified by, for example, ammonium sulfate precipitation, Protein A or G column, DEAE ion exchange chromatography, affinity column coupled with the sulfotransferase described above, synthetic peptide, or such, to prepare antibodies. Monoclonal antibodies can be obtained by the following procedure: small animals such as mice are immunized with an above-described sulfotransferase, or a partial peptide thereof. Spleens are removed from the mice and crushed to isolate cells. The cells are fused with mouse myeloma cells using a reagent such as polyethylene glycol. Clones producing antibodies that bind to an above-described sulfotransferase are selected from among the resulting fused cells (hybridomas). The obtained hybridomas are then transplanted into the peritoneal cavities of mice, and ascites collected. The obtained monoclonal antibodies can be purified by, for example, ammonium sulfate precipitation, Protein A or G columns, DEAE ion exchange chromatography, affinity columns coupled with an above-described sulfotransferase, synthetic peptides, or such.

The antibodies of the present invention are not particularly limited as long as they bind to an above-described sulfotransferase of the present invention. The antibodies of the present invention may be human antibodies, humanized antibodies created by gene recombination, fragments or modified products of such antibodies, in addition to the polyclonal and monoclonal antibodies described above.

The proteins of the present invention used as sensitizing antigens to prepare antibodies are not limited in terms of the animal species from which the proteins are derived. However, the proteins are preferably derived from mammals, for example, mice and humans. Human-derived proteins are particularly preferred. The human-derived proteins can be appropriately obtained by those skilled in the art using the gene or amino acid sequences disclosed herein.

In the present invention, the proteins to be used as sensitizing antigens may be whole proteins or partial peptides thereof. Such partial peptides of the proteins include, for example, amino-terminal (N) fragments and carboxyl-terminal (C) fragments of the proteins. Herein, "antibodies" refer to antibodies that react with a full-length protein or fragment thereof.

In addition to immunizing nonhuman animals with antigens to obtain the above hybridomas, human lymphocytes, for example, EB virus-infected human lymphocytes, can be sensitized *in vitro* with the proteins or with cells expressing the proteins, or with lysates thereof, and the sensitized lymphocytes can be fused with human-derived myeloma cells with the ability to divide permanently, for example, U266, to obtain hybridomas that produce desired human antibodies with binding activity to the proteins.

It is expected that antibodies against the above-described sulfotransferases of the present invention exhibit the effect of inhibiting protein expression or function by binding to the proteins. When using the prepared antibodies for human administration (antibody therapy), the antibodies

are preferably human or humanized antibodies in order to reduce immunogenicity.

Furthermore, in the present invention, low-molecular-weight substances (low-molecular-weight compounds) that bind to the above-described sulfotransferases are also included in the substances capable of inhibiting the function of the above-described sulfotransferases. Such low-molecular-weight substances may be natural or artificial compounds. In general, the compounds can be produced or obtained by methods known to those skilled in the art. The compounds of the present invention can also be obtained by the screening methods described below.

In addition, the substances of the present invention capable of inhibiting the expression or function of the above-described sulfotransferases include dominant-negative mutants (dominant-negative proteins) for the above-described sulfotransferases. The “dominant-negative protein mutants for the above sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine” refer to proteins with the function of reducing or abolishing the activity of endogenous wild-type proteins by expressing the genes encoding the sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine.

The inhibitors of the present invention that inhibit sulfation at position 4 or 6 of N-acetylgalactosamine have therapeutic or preventive effect for fibrogenic disorders. Therefore, in a preferred embodiment, the agents of the present invention are therapeutic or preventive agents for fibrogenic disorders.

Herein, “therapeutic or preventive” does not necessarily refer to a perfect therapeutic or preventive effect on organs or tissues with tissue fibrogenesis, and may refer to a partial effect.

The tissue fibrogenesis-suppressing agents of the present invention have the activity of suppressing fibrogenesis through inhibiting the sulfation at position 4 or 6 of N-acetylgalactosamine, which is a cause of fibrogenesis. Thus, preferred embodiments of the present invention provide, for example, therapeutic or preventive agents for tissue fibrogenic disorders which comprise as an active ingredient a tissue fibrogenesis-suppressing agent of the present invention.

The “therapeutic agents for tissue fibrogenic disorders” of the present invention can also be referred to as “improving agents for tissue fibrogenic disorders”, “anti-tissue fibrogenesis agents”, or the like. Meanwhile, the agents of the present invention can also be referred to as “pharmaceutical agents”, “pharmaceutical compositions”, “therapeutic medicines”, or the like.

The “treatments” of the present invention also comprise preventive effects that can suppress the onset of fibrogenesis in advance. The treatments are not limited to those producing a complete therapeutic effect on fibrogenic organs (tissues), and the effects may be partial.

The agents of the present invention can be combined with physiologically acceptable carriers, excipients, diluents and such, and orally or parenterally administered as pharmaceutical

compositions. Oral agents may be in the form of granules, powders, tablets, capsules, solutions, emulsions, suspensions, or the like. The dosage forms of parenteral agents can be selected from injections, infusions, external preparations, inhalants (nebulizers), suppositories, and the like. Injections include preparations for subcutaneous, intramuscular, intraperitoneal, intracranial, and intranasal injections, and the like. The external preparations include nasal preparations, ointments, and such. Techniques for formulating the above-described dosage forms that contain the agents of the present invention as primary ingredients are known.

For example, tablets for oral administration can be produced by compressing and shaping the agents of the present invention in combination with excipients, disintegrants, binders, lubricants, and the like. Excipients commonly used include lactose, starch, mannitol, and the like. Commonly used disintegrants include calcium carbonate, carboxymethylcellulose calcium, and the like. Binders include gum arabic, carboxymethylcellulose, and polyvinylpyrrolidone. Known lubricants include talc, magnesium stearate, and such.

Known coatings can be applied to tablets comprising the agents of the present invention to prepare enteric coated formulations or for masking. Ethylcellulose, polyoxyethylene glycol, or such can be used as a coating agent.

Meanwhile, injections can be prepared by dissolving the agents of the present invention, which are chief ingredients, together with an appropriate dispersing agent, or dissolving or dispersing the agents in a dispersion medium. Both water-based and oil-based injections can be prepared, depending on the selection of dispersion medium. When preparing water-based injections, the dispersing agent is distilled water, physiological saline, Ringer's solution or such. For oil-based injections, any of the various vegetable oils, propylene glycols, or such is used as a dispersing agent. If required, a preservative such as paraben may be added at this time. Known isotonicizing agents such as sodium chloride and glucose can also be added to the injections. In addition, soothing agents such as benzalkonium chloride and procaine hydrochloride can be added.

Alternatively, the agents of the present invention can be formed into solid, liquid, or semi-solid compositions to prepare external preparations. Such solid or liquid compositions can be prepared as the same compositions as described above and then used as external preparations. The semi-solid compositions can be prepared using an appropriate solvent, to which a thickener is added if required. Water, ethyl alcohol, polyethylene glycol, and the like can be used as the solvent. Commonly used thickeners are bentonite, polyvinyl alcohol, acrylic acid, methacrylic acid, polyvinylpyrrolidone, and the like. Preservatives such as benzalkonium chloride can be added to these compositions. Alternatively, suppositories can be prepared by combining the compositions with carriers, like oil bases such as cacao butter, or aqueous gel bases such as cellulose derivatives.

When the agents of the present invention are used as gene therapy agents, the agents may be directly administered by injection, or vectors carrying the nucleic acid may be administered. Such vectors include adenovirus vectors, adeno-associated virus vectors, herpes virus vectors, vaccinia virus vectors, retroviral vectors, and lentivirus vectors. These vectors allow efficient administration.

Alternatively, the agents of the present invention can be encapsulated into phospholipid vesicles such as liposomes, and then the vesicles can be administered. Vesicles carrying siRNAs or shRNAs are introduced into given cells by lipofection. The resulting cells are then systemically administered, for example, intravenously or intra-arterially. The cells can also be locally administered into tissues or such with fibrogenesis. siRNAs exhibit a quite superior and specific post-transcriptional suppression effect *in vitro*; however, *in vivo* they are rapidly degraded due to serum nuclease activity, and thus, their time was limited. There is therefore demand for the development of optimized and effective delivery systems. As one example, Ochiya *et al.* have reported that atelocollagen, a bio-affinity material, is a highly suitable siRNA carrier because it has the activity of protecting nucleic acids from nucleases in the body when mixed with the nucleic acids to form a complex (Ochiya, T. *et al.*, Nat. Med., 1999, 5, 707-710; Ochiya, T. *et al.*, Curr. Gene Ther., 2001, 1, 31-52); however, the methods for introducing drugs of the present invention are not limited thereto.

The agents of the present invention are administered to mammals including humans at required (effective) doses, within a dose range considered to be safe. Ultimately, the doses of the agents of the present invention can be appropriately determined by medical practitioners or veterinarians after considering the dosage form and administration method, and the patient's age and weight, symptoms, and the like. For example, adenoviruses are administered once a day at a dose of about 10^6 to 10^{13} viruses every one to eight weeks, although the doses vary depending on the age, sex, symptoms, administration route, administration frequency, and dosage form.

Commercially available gene transfer kits (for example: AdenoExpressTM, Clontech) may be used to introduce siRNAs or shRNAs into target tissues or organs.

Diseases to be treated or prevented by the agents of the present invention is not particularly limited as long as they are caused by tissue fibrogenesis, but preferably include, cardiac disorders, intestinal diseases, liver diseases, hepatic disorders, kidney disorders, cranial nerve diseases, eye disorders, pancreas disorders.

The "diseases caused by fibrogenesis" in the present invention is not particularly limited, and specifically include, for example, elastosis, scleroderma, chronic peritonitis, and retroperitoneal fibrosis in integumentary and epithelial tissues such as skin; polymyositis, dermatomyositis, polyarteritis nodosa, soft tissue fibrosis, chronic rheumatoid arthritis, palmar fibromatosis, tendinitis, tenovaginitis, Achilles tendinitis, mycetoma pedis, and

such in supportive tissues such as connective tissues and muscles;

myelofibrosis, hypersplenism, vasculitis, bradyarrhythmia, arteriosclerosis, obstructive thrombotic angiitis, nodular fibrosis, angina pectoris, dilated congestive cardiomyopathy, heart failure, restrictive cardiomyopathy, diffuse nonobstructive cardiomyopathy, obstructive
5 cardiomyopathy, cor pulmonale, mitral stenosis, aortic valve stenosis, chronic pericarditis, endocardial fibrosis, endomyocardial fibrosis, and such in blood tissues and vascular system such as bone marrow and heart;

chronic pancreatitis, Crohn's disease, ulcerative colitis, alcoholic hepatitis, chronic hepatitis B, chronic hepatitis C, Wilson's disease, cirrhosis, viral hepatitis, Gaucher's disease, glycogen
10 storage disease, alpha 1-antitrypsin deficiency, hemochromatosis, tyrosinemia, levulosemia, galactosemia, Zellweger syndrome, congenital hepatic fibrosis, portal hypertension, hepatic granulomatosis, Budd-Chiari syndrome, primary sclerosing cholangitis, fatty liver, nonalcoholic hepatitis, hepatic fibrosis, congenital hepatic fibrosis, alcoholic cirrhosis, viral cirrhosis, parasitic cirrhosis, toxic cirrhosis, trophopathic cirrhosis, congestive cirrhosis, hepatic sclerosis, Charcot's
15 cirrhosis, Todd's cirrhosis, secondary biliary cirrhosis, unilobar cirrhosis, cirrhosis resulting from chronic nonsuppurative destructive cholangitis, obstructive cirrhosis, cholangiolitic cirrhosis, biliary cirrhosis, atrophic cirrhosis, postnecrotic cirrhosis, posthepatic cirrhosis, nodular cirrhosis of the liver, mixed cirrhosis, micronodular cirrhosis, compensatory cirrhosis, decompensated cirrhosis, macronodular cirrhosis, septal cirrhosis, cryptogenic cirrhosis,
20 periportal cirrhosis, portal cirrhosis, primary biliary cirrhosis, and such in the gastrointestinal system such as liver;

coccidioidomycosis, blastomycosis, allergic bronchopulmonary aspergillosis, Goodpasture's syndrome, pulmonary fibrosis associated with adult respiratory distress syndrome, chronic obstructive pulmonary disease, pulmonary atelectasis, pneumonia, chalicosis, asbestosis,
25 hypersensitivity pneumonitis, idiopathic pulmonary fibrosis, lymphocytic interstitial pneumonia, Langerhans-cell granulomatosis, cystic fibrosis, pustular fibrosis, pulmonary fibrosis, fibrosing pulmonary alveolitis, interstitial fibrosis, diffuse pulmonary fibrosis, chronic interstitial pneumonia, bronchiectasis, bronchiolar fibrosis, peribronchial fibrosis, pleural fibrosis, and such in the respiratory system such as lung;

male hypogonadism, myotonic dystrophy, fibrosis such as associated with Peyronie's disease, chronic tubulointerstitial nephritis, autosomal recessive cystic kidney, myeloma kidney, hydronephrosis, rapidly progressive glomerulonephritis, nephrotoxic diseases,
xanthogranulomatous pyelonephritis, sickle cell nephropathy, nephrogenic diabetes insipidus, autosomal dominant polycystic kidney disease, chronic glomerular nephritis, IgA nephropathy,
35 renal sclerosis, focal glomerulosclerosis, membranous nephritis, membranoproliferative glomerulonephritis, chronic pyelonephritis, renal amyloidosis, polycystic kidney disease,

retroperitoneal fibrosis, pathology in the kidney associated with a connective tissue disease such as lupus nephritis, diabetic nephropathy, chronic prostatitis, and urocystitis associated with schistosomiasis in the urogenital system such as kidney;

fibrotic breast disease, mammary fibroadenoma, and such;

- 5 congenital torticollis, ankylosing spondylitis, spinal cord disorders such as neurofibroma and neurological dysfunction after spinal cord injury, and cranial nerve diseases such as Parkinson's disease and Alzheimer's disease in the nervous system such as spinal cord;

retrolental fibrosis and proliferative retinopathy in the eyeball; and

- 10 sarcoidosis that develops systemic involvement, fibrosis and systemic scleroderma associated with systemic lupus erythematosus, polymyositis, dermatomyositis, and such. However, in the present invention, the "disease caused by fibrogenesis" is not limited thereto, and includes diseases caused by fibrosis in each body tissue such as skin and organs.

- The present invention also relates to methods of screening for agents for suppressing tissue fibrogenesis (herein sometimes referred to as "methods of the present invention"), which
15 use as an indicator the degree of sulfation at position 4 or 6 of N-acetylgalactosamine.

A preferred embodiment of methods of the present invention is the methods comprising the step of selecting compounds that inhibit the sulfation at position 4 or 6 of N-acetylgalactosamine that constitute sugar chains.

- Using the screening methods of the present invention, tissue fibrogenesis-suppressing
20 agents or candidate compounds for agents for treating or preventing fibrogenic disorders can be efficiently acquired.

Preferred embodiments of the screening methods of the present invention are methods of screening for tissue fibrogenesis-suppressing agents that comprise the steps of (a) to (c):

- (a) contacting test compounds with N-acetylgalactosamines or sugar chain having N-
25 acetylgalactosamines;

(b) measuring a degree of sulfation at position 4 or 6 of N-acetylgalactosamines; and

(c) selecting compounds that reduce the degree of sulfation as compared with those without contacting with the test compounds.

Embodiments of the screening methods of the present invention are exemplified below.

- 30 In the embodiments described below, N-acetylgalactosamines, sulfotransferases, desulfatases, or such to be used include those derived from humans, mice, rats, and others, but are not particularly limited thereto.

- The test compounds to be used in the embodiments described below are not particularly limited, but include, for example, single compounds, such as natural compounds, organic
35 compounds, inorganic compounds, proteins, and peptides, as well as compound libraries, expression products of gene libraries, cell extracts, cell culture supernatants, products of

fermenting microorganisms, extracts of marine organisms, and plant extracts.

In the methods of the present invention, the “contact” with test compounds is typically achieved by mixing the test compounds with N-acetylgalactosamines, sulfotransferases, , or desulfatases, but the “contact” is not limited to this methods. For example, the “contact” can
 5 also be achieved by contacting test compounds with cells expressing these proteins or portions thereof.

In the embodiments described below, the “cells” include those derived from humans, mice, rats, and such, but are not limited thereto. Cells of microorganisms, such as *Escherichia coli* and yeasts, which are transformed to express the proteins used in each embodiment, can also
 10 be used. For example, the “cells that express genes encoding sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine” include cells that express endogenous genes encoding sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine, or cells that express introduced foreign genes encoding sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine. Such cells that express foreign genes encoding
 15 sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine can typically be prepared by introducing host cells with expression vectors carrying a gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine as an insert. The expression vectors can be prepared using standard genetic engineering techniques.

In addition, the degree of sulfation in the methods of the present invention can be
 20 determined by methods known to those skilled in the art. For example, the degree of sulfation can be determined by measuring the amount of label using a labeled compound, antibody, or such that binds to a sulfated structure at position 4 or 6 of N-acetylgalactosamine, or a portion thereof. Alternatively, the degree of sulfation can be detected by chromatography, mass spectrometry, or the like.

Those skilled in the art can appropriately evaluate the degree of sulfation at position 4 or
 25 6 of N-acetylgalactosamine, for example, by the following known methods:

(1) method based on quantitative dye binding using a labeling dye (1-9-dimethylene blue)
 (Nature. 1998 Feb 26; 391 (6670): 908-11)

(2) method based on photo-affinity labeling using [³²P]3',5'-ABP (Mandon, E.C., Milla, M.E.,
 30 Kempner, E., and Hirschberg, C.B. (1994) Proc. Natl. Acad. Sci. USA, 91, 10707-10711)

(3) method based on photo-affinity labeling using 3'-[³²P]-β methyleneb PAPS (Ozeran, J.D., Wesley, J., and Schwarz, N.B. (1996) Biochemistry, 35, 3695-3703)

(4) method using anion exchange resins (method for isolating sulfated glycoproteins) (Vol. 16,
 No. 2 (19860430) pp. 69-72, Kitasato University, ISSN: 03855449)

(5) colorimetric staining of sGAG with Alcian blue (Anal Biochem. 1998 Feb 15; 256(2): 229-
 35 37)

Furthermore, those skilled in the art can readily evaluate by the above-listed methods or the like whether a substance is an inhibitor of sulfation at position 4 or 6 of N-acetylgalactosamine of the present invention.

Another embodiment of screening methods of the present invention includes the methods comprising the step of selecting substances (compounds) which reduce the activity of sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine.

The above-described methods of the present invention comprise, for example, the steps of:

- (a) contacting a test compound with sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine;
- (b) measuring the sulfotransferase activity of the enzymes; and
- (c) selecting a compound that reduces the activity as compared to when the test compound is not contacted.

In the above-described methods, first, a test compound is contacted with a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine.

Then, the sulfotransferase activity of the enzyme is measured. Next, a compound that reduces the activity as compared to without contact with the test compound is selected. Such a compound that reduces the activity can be used as a fibrogenesis inhibitor or a therapeutic agent for fibrogenic disorders.

Methods that enable evaluation (determination) of whether a test compound has the above-described sulfotransferase activity include, for example, the methods described below.

Various test compounds are mixed during a set period of culture of cells or cell lines that promote the sulfation at position 4 or 6 of N-acetylgalactosamine, and the degree of sulfation before and after the culture can be easily determined by, for example, using an antibody that recognizes sulfation at position 4 (clone: LY111, 2H6) or an antibody that recognizes sulfation at position 6 (clone: MC21C, MO225, and CS-56) (all from Seikagaku Co.). Fluorescence values may be compared between before and after the culture by using fluorescently labeled antibodies. Alternatively, the same detection method can be conducted using 2-B-6 or 3-B-3 antibodies before and after culture. Compounds that suppress an increase in the sulfation after cell culture (an increase in the fluorescence value for LY111 or MC21C), or compounds that promote the progression of desulfation after cell culture (an increase in the fluorescence value for 2-B-6 or 3-B-3) can be selected as a desired candidate compound in the methods of the present invention.

As a further option, cell lines that constitutively express sulfotransferase genes such as C4ST-1 and C6ST-1 can be prepared by introducing the genes into CHO cells, L cells, or such by well-known methods. The use of such cell lines that constitutively add sulfate groups allows

a more clear determination of candidates for therapeutic compounds.

Another preferred embodiment of the screening methods for a tissue fibrogenesis inhibitor of present invention includes methods comprising the step of selecting compounds that reduce the expression of N-acetylgalactosamine sulfotransferase genes of the present invention.

- 5 The above methods of the present invention comprise, for example, the steps of:
- (a) contacting a test compound with cells expressing genes encoding sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine;
 - (b) measuring the gene expression level of the cells; and
 - (c) selecting a compound that reduces the gene expression level as compared to when the test
- 10 compound is not contacted.

In the above methods, test compounds are first contacted with cells expressing a gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine.

- Next, the expression level of the gene encoding the sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine is measured. Herein, "expression of the
- 15 gene" includes both transcription and translation. Gene expression level can be measured by methods known to those skilled in the art.

- For example, mRNAs are extracted from cells expressing any one of the above-described proteins by conventional methods, and these mRNAs can be used as templates in Northern hybridization, RT-PCR, DNA arrays, or such to measure the transcription level of the
- 20 gene. Alternatively, protein fractions are collected from cells expressing a gene encoding any of the above-described proteins, and expression of the protein can be detected by electrophoresis such as SDS-PAGE to measure the level of gene translation. Alternatively, the level of gene translation can be measured by detecting the expression of any of the above-described proteins by Western blotting using an antibody against the proteins. Such antibodies for use in detecting
- 25 the proteins are not particularly limited, as long as they are detectable. For example, both monoclonal and polyclonal antibodies can be used.

Next, the expression level is compared with that in the absence of the test compounds (the control).

- Then, compounds that reduce (suppress) the expression level of the gene as compared to
- 30 when the test compounds are absent are selected. The compounds resulting in a reduction (suppression) can be agents for suppressing tissue fibrogenesis or candidate compounds for treating fibrogenic disorders.

- Furthermore, an embodiment of the screening methods of the present invention includes methods of selecting the present invention's compounds that reduce the expression level of a
- 35 gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine, using as an indicator, the amount (level) of reporter gene expression. The

above-described methods comprise, for example, the steps of:

- (a) contacting a test compound with cells or cell extracts containing a DNA structured such that a reporter gene is operably linked to a transcriptional regulatory region of a gene encoding a sulfotransferase that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine;
- 5 (b) measuring the expression amount (level) of the reporter gene; and
- (c) selecting a compound that reduces the expression amount (level) of the reporter gene as compared to when the test compound is not contacted.

In the above methods, test compounds are first contacted with cells or cell extracts containing DNAs structured such that a reporter gene is operably linked with a transcriptional
10 regulatory region of a gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine.

Herein, “operably linked” means that a reporter gene is linked with a transcriptional regulatory region of a gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine, such that expression of the reporter gene is induced upon binding of
15 transcriptional factors to the transcriptional regulatory region. Therefore, the meaning of “operably linked” also includes cases where a reporter gene is linked with a different gene and produces a fusion protein with a different gene product, as long as expression of the fusion protein is induced upon the binding of transcriptional factors to the transcriptional regulatory region of the gene encoding the sulfotransferase that transfers a sulfate to position 4 or 6 of N-
20 acetylgalactosamine. Those skilled in the art can obtain the transcriptional regulatory regions of genes encoding sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine that are present in the genome, based on the cDNA nucleotide sequences of the genes encoding the sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine.

The reporter genes for use in these methods are not particularly limited, as long as their expression is detectable. The reporter genes include, for example, the CAT gene, the lacZ gene, the luciferase gene, and the GFP gene. The “cells containing a DNA structured such that a reporter gene is operably linked with a transcriptional regulatory region of a gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine” include, for
30 example, cells introduced with vectors carrying such structures as inserts. Such vectors can be prepared by methods well known to those skilled in the art. The vectors can be introduced into cells by standard methods, for example, calcium phosphate precipitation, electroporation, lipofection, and microinjection. The “cells containing a DNA structured such that a reporter gene is operably linked with a transcriptional regulatory region of a gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine” include cells
35 in which the structure has been integrated into the chromosomes. A DNA structure can be

integrated into chromosomes by methods generally used by those skilled in the art, for example, gene transfer methods using homologous recombination.

The “cell extracts containing a DNA structured such that a reporter gene is operably linked with a transcriptional regulatory region of a gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine” include, for example, mixtures of cell extracts included in commercially available *in vitro* transcription-translation kits and DNAs structured such that a reporter gene is operably linked with the transcriptional regulatory region of the gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine.

“Contact” can be achieved by adding test compounds to a culture medium of “cells containing a DNA structured such that a reporter gene is operably linked with a transcriptional regulatory region of a gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine”, or by adding test compounds to the above-described commercially available cell extracts containing the DNAs. When the test compound is a protein, contact may also be achieved, for example, by introducing a DNA vector expressing the protein into the cells.

In the above methods, the expression level of the reporter gene is then measured. The expression level of the reporter gene can be measured by methods known to those skilled in the art, depending on the type of the reporter gene. When the reporter gene is the CAT gene, its expression can be determined, for example, by detecting the acetylation of chloramphenicol by the gene product. When the reporter gene is the lacZ gene, its expression level can be determined by detecting the color development of chromogenic compounds due to the catalytic action of the gene expression product. Alternatively, when the reporter gene is the luciferase gene, its expression level can be determined by detecting the fluorescence of fluorogenic compounds due to the catalytic action of the gene expression product. Furthermore, when the reporter gene is the GFP gene, its expression level can be determined by detecting the fluorescence of the GFP protein.

In the above methods, the expression amount (level) of the reporter gene is then compared with that in the absence of the test compounds (the control). Compounds that reduce (suppress) the expression level of the reporter gene as compared with a control are then selected, where the reporter gene is operably linked with a gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine. Compounds resulting in a reduction (suppression) can be agents for suppressing tissue fibrogenesis or candidate compounds for treating fibrogenic disorders.

The tissue fibrogenesis inhibitors that are found by the screening methods of the present invention are preferably therapeutic or preventive agents for fibrogenic disorders.

The present invention also provides methods of producing pharmaceutical compositions

for treating or preventing fibrogenic disorders. The above-described production methods of the present invention comprise, for example, the steps of:

(a) selecting a tissue fibrogenesis inhibitor from test samples by the above-described methods of screening for tissue fibrogenesis inhibitors; and

5 (b) combining the agent with a pharmaceutically acceptable carrier.

In these methods, first, a tissue fibrogenesis inhibitor is selected from test samples by the above-described methods of screening for tissue fibrogenesis inhibitors.

Then, the selected agent is combined with a pharmaceutically acceptable carrier. The pharmaceutically acceptable carrier includes, for example, those described above.

10 The present invention also provides kits comprising various agents, reagents, and the like, which are used to conduct the screening methods of the present invention.

The kits of the present invention can be prepared, for example, by selecting adequate reagents from the above-described various reagents, depending on the screening method to be conducted. The kits of the present invention may contain, for example, the sulfotransferases that transfer a sulfate to position 4 or 6 of N-acetylgalactosamine of the present invention. The kits of the present invention may further contain various reagents, vessels, and the like to be used in the methods of the present invention. The kits may appropriately contain, for example, antibodies, probes, various reaction reagents, cells, culture media, control samples, buffers, and instruction manuals containing a description of how to use the kits.

20 The present invention also provides therapeutic or preventive methods for fibrogenic disorders, which comprise the step of administering the agents of the present invention to individuals (for example, to patients and such).

The individuals subjected to the therapeutic or preventive methods of the present invention are not particularly limited, as long as they are organisms that can develop a fibrogenic disorder; however, humans are preferred.

In general, administration to individuals can be achieved, for example, by methods known to those skilled in the art, such as intraarterial injections, intravenous injections, and subcutaneous injections. The administered dose varies depending on the patient's weight and age, and the administration method or such; however, those skilled in the art (medical practitioners, veterinarians, pharmacists, and the like) can appropriately select a suitable dose.

30 The present invention also relates to the uses of agents of the present invention in producing tissue fibrogenesis inhibitors.

All prior art documents cited in this specification are incorporated herein by reference.

35 [Examples]

Herein below, the present invention will be specifically described with reference to

Examples, but the technical scope of the present invention is not to be construed as being limited thereto.

Herein, occasionally, an siRNA structure (sequence) is presented by showing a DNA region of a target gene. Those skilled in the art can readily understand the structure of an siRNA comprising double-stranded RNA corresponding to the DNA sequence based on the information of DNA sequence described as a target sequence.

[Cardiac tissue]

[Example 1] Assessment of the target sugar chain-related gene knockdown effect of siRNAs and anti-fibrogenic effect at the gene level in a mouse cardiomyopathy model

A model prepared by intraperitoneal administration of Doxorubicin hydrochloride (DOX; Kyowa Hakko), as a standard mouse cardiomyopathy model, was used in this Example and those below. This mouse model is classical, but highly reproducible and simple. Thus, the model has been widely used as a cardiomyopathy model for elucidating pathological conditions, experimenting new therapeutics, or such (Longhu Li, Circulation (2006) 113: 535-543; Xiaoming Yi, Am J Physiol Heart Circ Physiol (2006) 290: H1098-H1102; Kang YJ, J Biol Chem. 2000 May 5; 275(18): 13690-8; Nozaki N, Circulation (2004) 110: 2869-2874; Fisher PW, Circulation (2005) 111: 1601-1610).

The model mouse histologically develops fibrogenesis of the myocardial interstitium. This pathological findings is commonly observed in dilated cardiomyopathy, restrictive cardiomyopathy, hypertrophic cardiomyopathy, and arrhythmogenic right ventricular cardiomyopathy (ARVC), as well as left ventricular remodeling after acute myocardial infarction, stable angina pectoris, unstable angina pectoris, myocarditis, valvular heart disease, arrhythmia, or hypertension. The fibrogenesis is a pathological feature responsible for the myocardial dysfunction in chronic heart failure caused by the above-listed diseases (Jugdutt BI, Circulation. 108: 1395-1403, 2003).

First, the method of preparing the mouse model is described below. DOX (15 mg/kg; Kyowa Hakko) is administered to the peritoneal cavities of C57BL6/J mice (male, eight weeks old, CLEA Japan Inc.). The mice were reared for one week after administration, and then heart tissues were collected from them. As a control group, similar mice were also purchased and reared around the same time without DOX administration.

The GalNac4S-6ST siRNA agent was administered by the following procedure: 1 µg of GalNac4S-6ST siRNA (Hokkaido System Science, Co., Ltd.) was combined with 200 µl of 1% atelocollagen (Koken Co.) as a vehicle, and the mixture was intraperitoneally administered to each mouse 24 hours before DOX administration. The nucleotide sequence of the GalNac4S-6ST siRNA agent used in this Example is shown below, but the sequence is not limited to this

Example.

[human GalNac4S-6ST siRNA] (Gene Bank accession number NM_015892)
(Hokkaido System Science, Co., Ltd.)

5'-ggagcagagcaagaugaauacaauc-ag -3' (SEQ ID NO: 25)

5 3'-ua-ccucgucucguucuacuuauguuag -5' (SEQ ID NO: 26)

1 ml of RNA iso (TAKARA BIO INC.) was added to 50 mg each of organs (heart) excised from cardiomyopathy model mouse. The organs were crushed using an electrical homogenizer (DIGITAL HOMOGENIZER; AS ONE), then, 200 µl of chloroform (Sigma-Aldrich Japan) was added to the resulting suspension. The mixture was gently mixed and then cooled on ice for about five minutes, and centrifuged in a centrifuge (Centrifuge 5417R; Eppendorf) at 12,000 rpm and 4°C for 15 minutes. After centrifugation, 500 µl of the supernatant was transferred to a fresh Eppendorf tube, and an equal volume of isopropanol (500 µl; Sigma-Aldrich Japan) was added thereto. The solution was mixed, and then 1 µl of glycogen (Invitrogen) was added thereto. The mixture was cooled on ice for 15 minutes, and then centrifuged at 12,000 rpm and 4°C for 15 minutes. Next, RNA precipitate obtained after washing three times with 1,000 µl of 75% ethanol (Sigma-Aldrich Japan) was air-dried for 30 minutes to one hour, and then dissolved in Otsuka distilled water (Otsuka Pharmaceutical Co., Ltd). The solution was 100 times diluted with Otsuka distilled water. The RNA concentrations of extracted samples in UV plates (Corning Costar) were determined using a plate reader (POWER Wave XS; BIO-TEK).

Next, reverse transcription reaction (cDNA synthesis) is conducted by the following procedure. The concentrations of the obtained RNA samples were adjusted to 500 ng/20 µl. The samples were heated at 68°C for three minutes in a BLOCK INCUBATOR (ASTEC), and cooled on ice for ten minutes. After cooling on ice, 80 µl of RT PreMix solution (composition: 18.64 µl of 25 mM MgCl₂ (Invitrogen), 20 µl of 5x Buffer (Invitrogen), 6.6 µl of 0.1 M DTT (Invitrogen), 10 µl of 10 mM dNTP mix (Invitrogen), 2 µl of RNase Inhibitor (Invitrogen), 1.2 µl of MMLV Reverse Transcriptase (Invitrogen), 2 µl of Random primer (Invitrogen), and 19.56 µl of sterile distilled water (Otsuka distilled water; Otsuka Pharmaceutical Co., Ltd.)), which had been prepared in advance, was added to the samples. The mixtures were heated in a BLOCK INCUBATOR (ASTEC) at 42°C for one hour and at 99°C for five minutes, and then cooled on ice. 100 µl of desired cDNAs were prepared and quantitative PCR reaction was carried out using the prepared cDNAs in the following composition. For quantitative PCR, SYBR Premix Kit (TAKARA BIO INC.) and Real-time PCR thermal cycler DICE (TAKARA BIO INC.) were used. Conditions of PCR reaction was: 95°C for 10 seconds, 40 cycles of 95°C for 5 seconds and 60°C for 30 seconds, finally, melting curve analysis was conducted. Nucleotide sequences of primers used in the quantitative PCR were described below.

[Quantitative PCR Primer sequences]

*mouse GalNAc4S-6ST (TAKARA BIO INC.)

Forward: 5'-GTGAGTTCTGCTGCGGTCCA-3' (SEQ ID NO: 27)

Reverse: 5'-AGTCCATGCTGATGCCAGAG-3' (SEQ ID NO: 28)

5 *mouse procollagen Type 1 alpha 2 (TAKARA BIO INC.)

Forward: 5'-ACCCGATGGCAACAATGGA-3' (SEQ ID NO: 29)

Reverse: 5'-ACCAGCAGGGCCTTGTTTAC-3' (SEQ ID NO: 30)

*mouse α -SMA (TAKARA BIO INC.)

Forward: 5'-CATCCGTAAAGACCTCTATGCCAAC-3' (SEQ ID NO: 31)

10 Reverse: 5'-ATGGAGCCACCGATCCACA-3' (SEQ ID NO: 32)

*mouse rRibosome 18S (TAKARA BIO INC.)

Forward: 5'-TTCTGGCCAACGGTCTAGACAAC-3' (SEQ ID NO: 33)

Reverse: 5'-CCAGTGGTCTTGGTGTGCTGA-3' (SEQ ID NO: 34)

As shown in Fig. 2, the expressions of GalNAc4S-6ST, type I collagen, and α -SMA
 15 genes were determined, and the result showed that the expression of GalNAc4S-6ST was
 significantly suppressed in the GalNAc4S-6ST siRNA-treated group as compared to the
 untreated group ($P < 0.001$; when compared to the untreated group). Furthermore, the expressions
 of α -SMA and type I collagen genes were measured as indicators for fibrogenesis, which is an
 important pathological condition of cardiomyopathy. As a result, the significant reduction of
 20 expression were confirmed in the GalNAc4S-6ST siRNA-treated group as compared to the
 untreated group ($P < 0.001$; when compared to the untreated group). This result demonstrates that
 the target knockdown effect of the GalNAc4S-6ST siRNA results in suppression of the
 progression of myocardial fibrogenesis at the gene expression level.

The agents of the present invention are thus useful, for example, as myocardial
 25 fibrogenesis inhibitors.

[Example 2] Cardiac hypertrophy-suppressing effect of GalNAc4S-6ST siRNA in a mouse
 cardiomyopathy model

In this Example, the heart weights (mg) and body weights (g) of cardiomyopathy model
 30 mice were measured to calculate the heart/body weight ratio which is an indicator for cardiac
 hypertrophy. The cardiac hypertrophy-suppressing effect of the GalNAc4S-6ST (GalNac)
 siRNA was evaluated. Cardiac hypertrophy also serves as an indicator for tissue fibrotic change.

Fig. 1 shows the result of calculating the heart weight (mg)/body weight (g) ratios in the
 siRNA-treated group ($n=4$) and untreated group ($n=4$). The result showed that the ratio was
 35 6.376 ± 0.484 and 5.442 ± 0.203 in the untreated and siRNA-treated groups, respectively. Thus,
 the significant reduction of the ratio was found in the siRNA-treated group as compared to the

untreated group ($p < 0.05$; t-test). This suggests that GalNac4S-6ST siRNA has the effect of suppressing pathological cardiac hypertrophy.

The agents of the present invention are thus useful, for example, as cardiac hypertrophy-suppressing agents (therapeutic agents for cardiac hypertrophy).

5
[Example 3] Assessment of type I collagen deposition-suppressing effect of GalNac4S-6ST siRNA in a mouse cardiomyopathy model

In this Example, the type I collagen deposition (an indicator of fibrogenesis)-suppressing effect of GalNac4S-6ST siRNA was assessed using heart samples of
10 cardiomyopathy model mice. Cardiac tissue samples were collected from the same mice as described in Example 1, and embedded in OCT compound (Miles), an embedding medium for cryosectioning. The samples were sliced into thin sections using Cryostat (Carl Zeiss). The resulting sections were fixed with acetone (Sigma Aldrich Japan) for ten minutes, and then washed with phosphate buffer. A rabbit antiserum anti-type I collagen (rabbit polyclonal
15 antibody, 1:2,000 dilution; LSL) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, a peroxidase-labeled goat anti-rabbit IgG antibody (1:200 dilution; Cappel) was added as the secondary antibody, and the sections were incubated at room temperature for 30 minutes. After incubation, DAB substrate (Nichirei) was added to the samples. The samples were observed under a light microscope (Leica
20 Microsystems).

The histological findings were shown in Fig. 3. Very intense positive signals for type I collagen were observed between myocardial fibers in the untreated group. Meanwhile, in the siRNA-treated group, the type I collagen-positive signals were considerably weaker than those of the untreated group. The above-described type I collagen immunostaining result demonstrates
25 that the GalNac4S-6ST siRNA has the effect of suppressing the excessive deposition of type I collagen in myocardial tissues. This result correlates with the result of quantitative PCR described in Example 1.

The agents of the present invention are thus useful, for example, as agents for suppressing type I collagen deposition in myocardial tissues.

30
[Example 4] Assessment of type III collagen deposition-suppressing effect of GalNac4S-6ST siRNA in a mouse cardiomyopathy model

In this Example, the type III collagen deposition (an indicator of fibrogenesis activity) - suppressing effect of GalNac4S-6ST siRNA was assessed using heart samples of
35 cardiomyopathy model mice. Tissue sections obtained by the same method as described in Example 3 were fixed with acetone (Sigma Aldrich Japan) for ten minutes, and then washed with

phosphate buffer. A rabbit antiserum anti-type III collagen (rabbit polyclonal antibody, 1:2000 dilution; LSL) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, a peroxidase-labeled goat anti-rabbit IgG antibody (1:200 dilution; Cappel) was added as the secondary antibody, and the sections were incubated at room temperature for 30 minutes. After incubation, DAB substrate (Nichirei) was added to the samples. The samples were observed under a light microscope (Leica Microsystems).

The histological findings are shown in Fig. 4. Moderately strong positive signals for type III collagen were observed between myocardial fibers in the untreated group. Meanwhile, in the siRNA-treated group, the type III collagen-positive signals were comparable to those of the control group. The above-described result of type III collagen immunostaining demonstrates that the GalNac4S-6ST siRNA has the effect of suppressing the type III collagen deposition in heart tissues, implying that the siRNA is also effective in suppressing active collagen deposition.

The agents of the present invention are thus useful, for example, as agents for suppressing type III collagen deposition in myocardial tissues.

[Example 5] Assessment of fibroblast infiltration-suppressing effect of GalNac4S-6ST siRNA in a mouse cardiomyopathy model

This Example assesses the pharmacological effect of GalNac4S-6ST siRNA on the kinetics of fibroblasts that infiltrate into cardiac tissues of cardiomyopathy model mice due to DOX administration. Tissue sections obtained by the same method as described in Example 3 were fixed with acetone (Sigma Aldrich Japan) for ten minutes, and then washed with phosphate buffer. An anti-mouse fibroblast antibody (ER-TR7, rat monoclonal antibody, 1:400 dilution; BMA Biomedicals Ltd.) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, a peroxidase-labeled goat anti-rat immunoglobulin antibody (1:200 dilution; Biosource International, Inc.) was added as the secondary antibody, and the sections were incubated at room temperature for 30 minutes. After incubation, DAB substrate (Nichirei Biosciences) was added to the samples. The samples were observed under a light microscope (Leica Microsystems).

The histological findings were shown in Fig. 5. The photograph focuses on the ventricular septum. Infiltration of numerous fibroblasts was observed in the untreated group as compared to the control group. In contrast, the degree of fibroblast infiltration in the siRNA-treated group was less as compared to the untreated group. The above-described result shows that GalNac4S-6ST siRNA has the pharmacological effect of suppressing the fibroblast infiltration into myocardial tissues and this activity contributes to the anti-fibrogenic effect.

The agents of the present invention are thus useful, for example, as agents for suppressing fibroblast infiltration into myocardial tissues.


[Gastrointestinal tissue]

[Example 6] Clinical fibrogenesis-suppressing effect of GalNAc4S-6ST in a mouse intestinal fibrosis model

The colitis model mice were prepared by allowing C57BL/6J mice (female, six weeks old; CLEA Japan Inc.) to freely drink high-concentration chlorine water containing 3% dextran sulfate sodium (DSS; Wako Pure Chemical Industries Ltd.) for eight days. The DSS-induced colitis model has excellent reproducibility, and is thus widely used as a typical experimental mouse model for inflammatory bowel diseases such as ulcerative colitis or Crohn's disease, as well as a model with full-thickness inflammation and fibrotic changes, and muscle layer thickening, which are histological characteristics of the narrowing of colon lumen (Sasaki N, J Inflamm. 2005 2: 13, Review: Pucilowska JB *et al.* Am J Physiol Gastroenterol Liver Physiol. 279: G653-G659, 2000). Therefore, the histological findings are commonly and widely observed in inflammatory bowel diseases as well as pathological conditions with the histological narrowing of the intestinal lumen, specifically diseases such as intestinal Behcet's disease (simple ulcer), irritable bowel syndrome, ischemic enteritis, drug-induced enteritis, radiation enteritis, esophagus achalasia, esophagostenosis associated with scleroderma, narrowing of the colon lumen associated with systemic lupus erythematosus (SLE), Hirschsprung's disease, stenosis after removal of intestine (postoperative stricture), narrowing of the intestinal lumen after endoscopic mucosal resection for gastrointestinal cancer (tongue cancer, epipharynx carcinoma, pharyngeal cancer, esophageal cancer, stomach cancer, small intestinal cancer, colon cancer, and rectal cancer), and ileus.

Simultaneously to feeding mice with 3% DSS water, the same GalNAc4S-6ST siRNA (1 µg/head) as described in Example 1 was combined with atelocollagen (Koken Co.) prediluted 10-fold with PBS and 200 µl of the mixture was injected to the peritoneal cavities of the mice. The group of mice treated as described above was named "GalNAc4S-6ST siRNA group", while a group treated with atelocollagen alone without combining GalNAc4S-6ST siRNA was named the "control group". The body weight and the disease activity index (DAI) score were recorded during seven days of 3% DSS water feeding (Kihara M., Gut. 2003, 52, 713-9). The evaluation criteria for DAI are shown below.

[Table 1]



The result of scoring the DAI of each mouse, setting the score on the first day of DSS water feeding (day 0) as 1, is shown in Fig. 6. On the third day, the GalNAc4S-6ST siRNA-administered group exhibited a significantly lower score as compared to the control group (p<0.001; t-test). This result suggests that suppression of GalNAc4S-6ST gene expression produces the effect of suppressing inflammatory activity at relatively earlier stages.

Furthermore, the mice were sacrificed and their colon lengths were measured on the fifth day. The colon shortening was significantly suppressed in the GalNAc4S-6ST siRNA-administered group (p<0.005; t-test) (Fig. 6). The colon length is a definitive indicator that reflects intestinal fibrogenesis or stenosis. Thus, it was also clinically demonstrated that the fibrotic change of intestine was suppressed in the GalNAc4S-6ST siRNA-administered group.

The agents of the present invention are thus useful, for example, as agents for suppressing fibrotic changes of the intestine.

[Example 7] Intestinal fibrogenesis-suppressing effect of GalNAc4S-6ST siRNA in a mouse intestinal fibrosis model

In this Example, the expression of fibrogenesis-related genes in the colon after GalNAc4S-6ST siRNA administration was assessed by the quantitative real-time PCR method.

The intestinal fibrogenesis model was prepared by the same method as described in Example 6. The mice were sacrificed on day 7. A part of the collected colon was placed in 1.5-ml tubes, and frozen with liquid nitrogen. cDNA was synthesized by the same method as that described in Example 1, and quantitative PCR was carried out. The primer sequences and the number of PCR cycle conditions were the same as those described in Example 1.

The result is shown in Fig. 7. The expression of GalNAc4S-6ST gene was enhanced in this model. The significant knockdown of the gene was confirmed by GalNAc4S-6ST siRNA treatment (p<0.001; t-test). Furthermore, the enhanced expression of type I collagen and α -SMA as indicators of fibrogenesis were significantly suppressed by GalNAc4S-6ST siRNA (for both genes, p<0.001; t-test). The result suggests that the enhanced fibrotic changes of the colon can be effectively suppressed by suppressing the expression of GalNAc4S-6ST.

The agents of the present invention are thus useful, for example, as agents for suppressing the fibrotic changes of the colon.

[Example 8] Tissue fibrogenesis-suppressing effect of GalNAc4S-6ST siRNA in a mouse intestinal fibrosis model

The intestinal fibrogenesis model was prepared by the same method as described in Example 6. The mice were sacrificed on day 7. Cryoblocks and tissue sections were prepared from the collected colon by the same method as described in Example 3. Masson-stained images of colonic tissue sections are shown in Fig. 8. Masson staining serves as an indicator to assess fibrotic change of tissues by visualizing collagen fibers. In the GalNAc4S-6ST siRNA-administered group, the full-thickness (lamina propria mucosae, submucosa, and muscle layer) collagen fiber deposition was significantly suppressed as compared to the control group.

The agents of the present invention are thus useful, for example, as agents for suppressing the full-thickness (lamina propria mucosae, submucosa, and muscle layer) collagen fiber deposition.

[Example 9] Histological fibroblast infiltration-suppressing effect of GalNAc4S-6ST siRNA in a mouse intestinal fibrosis model

The intestinal fibrogenesis model was prepared by the same method as described in Example 6. The mice were sacrificed on day 7. Cryoblocks and tissue sections were prepared from the collected colon by the same method as described in Example 3. The resulting sections were fixed with acetone (Wako Pure Chemical Industries) for ten minutes, and then washed with phosphate buffer. An anti-ER-TR7 antibody (rat monoclonal antibody, 1 µg/ml; BMA) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, the secondary antibody reaction was carried out by adding a peroxidase-labeled anti-rat IgG antibody (1:200 dilution), which was followed by color development by adding DAB substrate (Nichirei Biosciences). Then, the nucleus was stained by Lillie-Mayer hematoxylin (Muto Pure Chemicals Co.). The samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

As a result, the full-thickness infiltration of fibroblasts was significantly suppressed in the GalNAc4S-6ST siRNA-treated group as compared to the control group (Fig. 9). This result demonstrates that the inhibition of GalNAc4S-6ST gene expression results in suppression of the infiltration and retention of fibroblasts in focal tissue lesion and thereby reduces the enhanced fibrotic change.

The agents of the present invention are thus useful, for example, as agents for suppressing the infiltration or retention of fibroblasts.

[Example 10] Histological macrophage infiltration-suppressing effect of GalNAc4S-6ST siRNA in a mouse intestinal fibrosis model

The intestinal fibrogenesis model was prepared by the same method as described in Example 6. The mice were sacrificed on day 7. Cryoblocks and tissue sections were prepared from the collected colon by the same method as described in Example 3. The resulting sections were fixed with acetone (Wako Pure Chemical Industries) for ten minutes, and then washed with phosphate buffer. An anti-F4/80 antibody (clone A3-1, rat monoclonal antibody, 2 µg/ml: CALTAG LABORATORIES) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, the secondary antibody reaction was carried out by adding a peroxidase-labeled anti-rat IgG antibody (1:200 dilution), which was followed by color development by adding DAB substrate (Nichirei Biosciences). Then, the nucleus was stained by Lillie-Mayer hematoxylin (Muto Pure Chemicals Co.). The samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

The result showed that the full-thickness macrophage infiltration was significantly suppressed in the GalNAc4S-6ST siRNA-treated group as compared to the control group (Fig. 10). This result demonstrates that the inhibition of GalNAc4S-6ST gene expression resulted in suppression of the infiltration of macrophages and fibroblasts, which are cell groups responsible for the persistent or enhanced fibrotic changes, and thereby comprehensively suppressed the tissue fibrotic changes.

The agents of the present invention are thus useful, for example, as agents for suppressing the infiltration of macrophages or fibroblasts.

[Example 11] Histological macrophage infiltration-suppressing effect of GalNAcST siRNA in a mouse intestinal fibrosis model

GalNAc4S-6ST is an enzyme that transfers a sulfate group to position 6 in N-acetylgalactosamine sulfated at position 4. GalNAc-4ST1 and GalNAc-4ST2, which belong to the 4-O-sulfotransferase family, were assessed in this Example. The intestinal fibrogenesis model was prepared by the same method as described in Example 6. The mice were sacrificed on day 7. In this Example, GalNAc4S-6ST siRNA, GalNAc4ST-1, and GalNAc4ST-2 (GeneWorld) were combined together; 1 µg of the mixture was combined with 200 µl of 1% atelocollagen (Koken Co.), which is a vehicle, and administered intraperitoneally to each mouse. A group administered with the siRNA is referred to as "GalNAc ST siRNA-administered group". The control group is the same as described in Example 6. The siRNA nucleotide sequences of GalNAc4S-6ST, GalNAc4ST-1, and GalNAc4ST-2 used in this Example are shown below, but

the sequence are not limited the Example.

[GalNAc4ST-1 siRNA cocktail sequences] (GenBank accession number NM_175140)
(GeneWorld)

5'-ACCCCCAACTCGGAACGATGCGGCT-3' (SEQ ID NO: 35)

5 5'-TGCATGTTCTCGTCCATCCTGCTG-3' (SEQ ID NO: 36)

5'-CGCCACCGTGTACTGTACTGTGAAGT-3' (SEQ ID NO: 37)

5'-AGGCT GCTCCAACCTG GAAGAGGGTG-3' (SEQ ID NO: 38)

[GalNAc4ST-2 siRNA cocktail sequences] (GenBank accession number NM_199055)
(GeneWorld)

10 5'-ATATAGTATCTAGGATATATGTAG-3' (SEQ ID NO: 39)

5'-GAAGTACCAAAAGCTGGCTGCTCTA-3' (SEQ ID NO: 40)

5'-TTCTATCACTTGGACTATTTGATGTT-3' (SEQ ID NO: 41)

5'-TACACAACCTCCACATTTGTAATTTG-3' (SEQ ID NO: 42)

[GALNAc4S-6ST siRNA cocktail sequences] (GenBank accession number NM_029935)

15 (GeneWorld)

5'-CCAGAAGCCAAGCTCATTGTTATG-3' (SEQ ID NO: 43)

5'-CTGTGGAGAGGTTGTACTCAGACTA-3' (SEQ ID NO: 44)

5'-ATTTGCCTGGAAGACAACGTGAGAGC-3' (SEQ ID NO: 45)

5'-GTCCCTTCTGCAGAAGCTGGGCCCCACT-3' (SEQ ID NO: 46)

20 As investigated in detail in Examples 6 to 10, the tissue fibrotic changes in the mouse intestinal fibrogenesis model can be assessed representatively by the colon length. Accordingly, the colon length on day 7 is shown as an essential evaluation item in this Example. In the GalNAc ST siRNA-administered group, the shortening of colon was significantly suppressed as compared to the control group ($p < 0.01$; t-test) (Fig. 11). It was thus demonstrated that the
25 intestinal fibrogenesis was inhibited by suppressing the expression of the GalNAc4ST-1 and GalNAc4ST-2 genes.

The agents of the present invention are thus useful, for example, as intestinal fibrogenesis inhibitors.

30 [Lung tissue]

[Example 12] Effect of C6ST-1 siRNA on pulmonary alveolar interstitium in a mouse pulmonary emphysema model

A basic mouse pulmonary emphysema model, which is prepared by intratracheal administration of porcine pancreatic elastase (PPE), is used in this Example. This mouse model
35 is classical, but highly reproducible and simple. Thus, this mouse model has been used commonly as a pulmonary emphysema model. Inflammatory cell infiltration to the pulmonary

alveolar interstitium is observed as a histological feature. This histological finding is commonly detected in chronic obstructive pulmonary disease (COPD) such as emphysema and chronic bronchitis as well as diseases causing chronic respiratory failure, such as idiopathic interstitial pneumonias (IIPs), coniosis, and pulmonary tuberculosis sequelae (Karlinsky JB *et al.*, *Am Rev Respir Dis* 1978; 117: 1109-1133; Otto-Verberne CJ *et al.*, Protective effect of pulmonary surfactant on elastase-induced emphysema in mice. *Eur Respir J* 1992; 5: 1223-1230; Janoff A *et al.*, Prevention of elastase-induced experimental emphysema by oral administration of a synthetic elastase inhibitor. *Am Rev Respir Dis* 1980; 121: 1025-1029; Christensen TG, *et al.*, Irreversible bronchial goblet cell metaplasia in hamsters with elastase-induced panacinar emphysema. *J Clin Invest* 1977; 59: 397-404; Lucey EC, *et al.*, Remodeling of alveolar walls after elastase treatment of hamsters: results of elastin and collagen mRNA in situ hybridization. *Am J Respir Crit Care Med* 1998; 158: 555-564; Snider GL, Lucey EC, Stone PJ. Animal models of emphysema. *Am Rev Respir Dis* 1986; 133: 149-169).

In this Example, the effect of C6ST-1 siRNA in suppressing emphysematous lesions was examined by hematoxylin-eosin staining (HE staining) of lung tissue samples from pulmonary emphysema model mice.

First, the model mice were prepared. PPE (4 units; Calbiochem-Novabiochem) was administered intratracheally to C57BL6/J mice (female, 5- to 6-weeks old; CLEA Japan). The mice were grown for three weeks after administration, and then lung tissues were collected from them. Mice that did not have PPE administration were used as the control group.

The C6ST-1 siRNA was administered by the same procedure as shown in Example 1: 1 µg of C6ST-1 siRNA (Ambion) was combined with 1% atelocollagen (Koken Co.), which is an siRNA vehicle, and administered to the peritoneal cavities once a week after PPE administration. The dose was 200 µl/head.

*[C6ST-1 siRNA cocktail sequences]

(GenBank accession number NM_016803)

5'-gcgccccctctcccatggagaaag-3' (SEQ ID NO: 47)

5'-gctttgcctcaggatttcgggacc-3' (SEQ ID NO: 48)

5'-gggtcagccttggtctaccgtgatgtc-3' (SEQ ID NO: 49)

5'-gcagttgttgctatgcgacctgtat-3' (SEQ ID NO: 50)

The collected right lung tissues were embedded in the OCT compound (Miles), an embedding medium for cryosectioning, and cryoblocks were prepared using liquid nitrogen. The cryoblocks were sliced into 6-µm sections using cryostat (Microm).

The resulting sections were fixed with 1% glutaraldehyde (Nacalai Tesque) for 10 minutes, and further fixed with formol-calcium solution for 10 minutes. The sections were washed with phosphate buffer, and then stained with Lillie-Mayer's hematoxylin solution

(Sigma Aldrich Japan) at room temperature for 5 minutes. The sections were washed gently with a decolorizing solution (70% ethanol containing 0.5% HCl; prepared using reagents from Nacalai Tesque), and then washed with water for 10 minutes. The sections were stained with eosin-alcohol at room temperature for 5 minutes, and then washed with water for 10 minutes.

The sections were washed gently with 100% ethanol, and then allowed to stand for 3 minutes. The sections were further washed gently with xylene (Nacalai Tesque), and allowed to stand for 10 minutes. This sample was histologically observed using a light microscope (Leica Microsystems).

The obtained histological images are shown in Fig. 12. The histological features of a normal lung parenchyma with characteristic faveolate alveolar septal walls are found in the control group (PPE non-administered mice). Meanwhile, emphysematous lesions due to destruction and abolishment of alveolar septal walls and enlargement of air space (characteristics of pulmonary emphysema) can be observed in the untreated group (administered with PPE but not with C6ST-1 siRNA) shown in the middle photograph. On the other hand, in the enzyme-treated group (administered with PPE and C6ST-1 siRNA), slight abolishment of alveolar septal walls and emphysematous lesions can be observed; however, their levels are significantly improved.

[Example 13] Pulmonary interstitial fibrosis-suppressing effect of C6ST-1 siRNA in a mouse pulmonary emphysema model

In this Example, the expression of fibrosis-related genes in the pulmonary alveolar interstitium after C6ST-1 siRNA administration is assessed by quantitative real-time PCR method.

The COPD model was prepared by the same method as described in Example 12. A part of the collected lung tissue were placed in 1.5-ml tubes, and frozen with liquid nitrogen. cDNA synthesis was carried out by the same method as described in Example 1, and quantitative PCR was conducted. The primer sequences for type I collagen and α -SMA and PCR conditions were also the same as described in Example 1. The sequences of C6ST-1 primers are shown below.

[Quantitative PCR primer sequences]

*mouse C6ST-1 (Takara Bio)

Forward: 5'-TGTTCTGGCATTGTGGTCATA -3' (SEQ ID NO: 51)

Reverse: 5'-CCAACTC GCTCAGGGACAAGA-3' (SEQ ID NO: 52)

The result is shown in Fig. 13. The expression of C6ST-1 gene is enhanced in this model. Significant knockdown of the C6ST-1 gene was confirmed by C6ST-1 siRNA treatment ($p < 0.01$; t-test). Furthermore, the enhanced expression of type I collagen and α -SMA as

indicators for fibrogenesis were significantly suppressed by C6ST-1 siRNA (for both genes, $p < 0.01$; t-test). This result suggests that the enhanced fibrotic changes of pulmonary interstitium can be effectively suppressed by inhibiting the expression of C6ST-1.

The agents of the present invention are thus useful, for example, as agents for suppressing fibrotic changes of the pulmonary interstitium.

[Example 14] Histological fibroblast cell infiltration-suppressing effect of C6ST-1 siRNA in a mouse pulmonary emphysema model

The intestinal fibrogenesis model was prepared by the same method as described in Example 12. Cryoblocks and tissue sections were prepared from the collected lung tissues by the same method as described in Example 3. The resulting sections were fixed with acetone (Wako Pure Chemical Industries) for ten minutes, and then washed with phosphate buffer. An anti-ER-TR7 antibody (rat monoclonal antibody, 1 $\mu\text{g/ml}$; BMA) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, the secondary antibody reaction was carried out by adding a peroxidase-labeled anti-rat IgG antibody (1:200 dilution), which was followed by color development by adding DAB substrate (Nichirei Biosciences). Then, the nucleus was stained by Lillie-Mayer hematoxylin (Muto Pure Chemicals Co.). The samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

As a result, a strong accumulation of fibroblasts was observed in the interstitium with damaged alveolar septa in the control group (Fig. 14; left panel for the control group). Furthermore, accumulation of many fibroblasts was also observed in the interstitium where the alveolar septa were being damaged (Fig. 14; right panel for the control group). Thus, the present inventors obtained the unexpected result that excessive accumulation of fibroblasts lead to the alveolar wall damaging process in pathological conditions such as COPD. Meanwhile, fibroblast infiltration to the interstitium of alveolar tissue was obviously suppressed in the C6ST-1 siRNA-treated group (Fig. 14). It was thus demonstrated that the suppression of C6ST-1 gene expression resulted in inhibition of fibroblast infiltration and retention in the interstitium of alveolar tissue and thereby reduced the enhanced fibrotic changes.

The agents of the present invention are thus useful, for example, as agents for suppressing fibroblast infiltration and retention in the interstitium of alveolar tissue.

[Example 15] Histological macrophage infiltration-suppressing effect of C6ST-1 siRNA in a mouse pulmonary emphysema model

The COPD model was prepared by the same method as described in Example 12. The collected lung tissues were processed by the same method as described in Example 3 to prepare

tissue cryoblocks and sections. The resulting sections were fixed with acetone (Wako Pure Chemical Industries) for ten minutes, and then washed with phosphate buffer. An anti-F4/80 antibody (clone A3-1, rat monoclonal antibody, 2 µg/ml; CALTAG LABORATORIES) was added as the primary antibody, and the sections were incubated at room temperature for one hour.

Then, the secondary antibody reaction was carried out by adding a peroxidase-labeled anti-rat IgG antibody (1:200 dilution), which was followed by color development by adding DAB substrate (Nichirei Biosciences). Then, the nucleus was stained by Lillie-Mayer hematoxylin (Muto Pure Chemicals Co.). The samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

The result showed that the macrophage infiltration to the pulmonary alveolar interstitium was clearly suppressed in the C6ST-1 siRNA-treated group as compared to the control group (Fig. 15). It was thus demonstrated that the suppression of C6ST-1 gene expression resulted in suppression of the infiltration of macrophages and fibroblasts, which are cell groups responsible for the persistent or enhanced fibrotic changes, and thereby comprehensively suppressed the tissue fibrotic changes.

The agents of the present invention are thus useful, for example, as agents for suppressing the infiltration of macrophages to the pulmonary alveolar interstitium.

[Example 16] Respiratory function-preserving effect of C6ST-1 siRNA in a mouse pulmonary emphysema model

In this Example, C6ST-1 siRNA was assessed for its influence on respiratory function using static lung compliance (static compliance (Cst)) as an indicator to evaluate the clinical effect of C6ST-1 siRNA in pulmonary emphysema model mice. Cst represents a measure of lung tissue flexibility. Cst is increased in pulmonary emphysema, which is a disease with alveolar tissue damage.

Pulmonary emphysema model mice were prepared by the same procedure described in Example 12, and then treated with C6ST-1 siRNA. An anesthetic agent was given to the mice to stop spontaneous respiration, and then their Cst was monitored using FlexiVent (SCIREQ) respiratory function analyzer in the PV loop mode. Mice were connected to the FlexiVent by the following procedure: a median incision was performed after stopping spontaneous respiration, and then a special cannula was inserted into the trachea, which was followed by peribronchial ligation.

The result of this Example is shown in Fig. 16. Cst was 42.62 ± 2.25 µl/cm H₂O in the control group, while it was 51.22 ± 5.2 µl/cm H₂O in the untreated group (when compared to the control group, $P=0.03$; t-test). Thus, a statistically significant increase was observed in the untreated group. In contrast, Cst was 42.92 ± 1.82 µl/cm H₂O (when compared to the untreated

group, $P=0.03$; t-test) and thus significantly decreased in the C6ST-1 siRNA-administered group when compared to the untreated group.

It was thus demonstrated that the C6ST-1 siRNA-administered group significantly suppressed the increase in Cst caused by pulmonary emphysema induced by intratracheal administration of PPE. Furthermore, the result described in this Example suggests that the suppression of C6ST-1 gene expression not only suppresses the alveolar damage caused by fibrotic changes of the interstitium of lung tissue but also improves the actual clinical symptoms (respiratory conditions).

The agents of the present invention are thus useful, for example, as agents for suppressing alveolar damage.

[Example 17] Tissue-preserving effect of C6ST-1 siRNA in a mouse pulmonary emphysema model

In pulmonary emphysema, whose characteristic pathological feature is enlarged alveolar air space, the lung capacity is increased with progression of emphysematous lesions. The purpose of this Example is to prove that the therapeutic effect of C6ST-1 siRNA is not only suppression of damage at the cell level but also the effect of morphological maintenance and preservation of the organ.

The pulmonary tissues used in this Example were the right lungs of the same lung tissues as used in Example 12. The lung tissues isolated from mice were gently washed with phosphate buffer and then immersed into phosphate buffer saturated in a glass container. The glass container filled with phosphate buffer was weighed in advance. After the lung tissues were added to the container, lung capacity was calculated by converting the increased weight into a liquid volume.

The result of this Example is shown in Fig. 17. The lung capacity was $277.5 \pm 61.85 \mu\text{l}$ in the control group, while it was $413.33 \pm 77.67 \mu\text{l}$ in the untreated group (when compared to the control group, $P=0.024$; t-test). Thus, a statistically significant increase of lung capacity was observed in the untreated group. In contrast, the lung capacity was $292.5 \pm 51.23 \mu\text{l}$ and thus significantly decreased in the C6ST-1 siRNA-treated group (when compared to the untreated group $P=0.027$; t-test) when compared to the untreated group.

These results revealed that the inhibition of C6ST-1 gene expression effectively suppresses the increase in the lung capacity associated with pulmonary emphysema induced by intratracheal administration of PPE. This suggests that the effect is not only the suppression of the damage at the cell level but also the effect of morphological maintenance and preservation of the organ or effect of repairing damaged tissues.

The agents of the present invention are thus useful, for example, as agents for

suppressing the increase of lung capacity caused by pulmonary emphysema.

[Example 18] Pulmonary interstitial fibrosis-suppressing effect of GalNAcST siRNA in a mouse pulmonary emphysema model

The importance of sulfation at position 4 and 6 is demonstrated with an additional Example. In this Example, the expression of fibrosis-related genes in the pulmonary alveolar interstitium after GalNAcST siRNA administration was assessed by quantitative real-time PCR method using the same method as described in Example 11. The siRNA sequences are the same as shown in Example 11.

The pulmonary emphysema model was prepared by the same method as described in Example 12. A part of the collected lung tissues were placed into 1.5-ml tubes, and frozen with liquid nitrogen. cDNA synthesis was carried out by the same method as described in Example 1, and assessed by quantitative PCR method. The primer sequences and of PCR conditions were the same as described in Examples 1 and 13. The primer sequences for TGF- β are shown below.

[Primer sequences used in quantitative PCR]

*mouse TGF- β (Takara Bio Inc.)

Forward: 5'-GTGTGGAGCAACATGTGGAAGCTCTA -3' (SEQ ID NO: 53)

Reverse: 5'-TTGGTTCAGCCACTGCCGTA -3' (SEQ ID NO: 54)

The result is shown in Fig. 18. The enhanced expression of type I collagen, α -SMA, and TGF- β as indicators for fibrogenesis were also significantly suppressed by GalNAcST siRNA (for all genes, $p < 0.01$; t-test) in this Example. This result demonstrates that the suppression of GAINAc4ST-1, GAINAc4ST-2, and GAINAc4S-6ST expression can effectively inhibit the enhanced fibrotic changes of pulmonary interstitium.

The agents of the present invention are thus useful, for example, as agents for suppressing fibrotic changes in lung interstitium.

[Example 19] Respiratory function-preserving effect of GalNAcST siRNA in a mouse pulmonary emphysema model

In this Example, GAINAcST siRNA was assessed for its influence on respiratory function using static lung compliance (static compliance (Cst)) as an indicator to evaluate the clinical effect of GAINAcST siRNA in pulmonary emphysema model mice.

Pulmonary emphysema model mice were prepared by the same procedure described in Example 12, and then treated with GAINAcST siRNA. The spontaneous respiration of mice was ceased by an anesthetic agent, and then their Cst was monitored using FlexiVent (SCIREQ) respiratory function analyzer in the PV loop mode. Mice were connected to the FlexiVent by the following procedure: a median incision was performed after stopping spontaneous respiration,

and then a special cannula was inserted into the trachea, which was followed by peribronchial ligation.

The result of this Example is shown in Fig. 18. Cst was 42.62 ± 2.25 $\mu\text{l}/\text{cm H}_2\text{O}$ in the control group, while it was 51.22 ± 5.2 $\mu\text{l}/\text{cm H}_2\text{O}$ in the untreated group (when compared to the control group, $P=0.03$; t test). Thus, a statistically significant increase was observed in the untreated group. In contrast, Cst was 44.15 ± 2.29 $\mu\text{l}/\text{cm H}_2\text{O}$ (when compared to the untreated group, $P=0.0018$; t-test) and thus significantly decreased in the C6ST-1 siRNA-administered group when compared to the untreated group.

It was thus demonstrated that the increase in Cst caused by pulmonary emphysema induced by intratracheal administration of PPE was significantly suppressed in the GalNAcST siRNA-administered group. Furthermore, the result described in this Example suggests that the inhibition of GAINAc4ST-1, GAINAc4ST-2, and GAINAc4S-6ST gene expression not only suppresses the alveolar damage caused by the fibrotic change of interstitium of lung tissue but also improves the actual clinical symptom (respiratory condition).

The agents of the present invention are thus useful, for example, as agents for suppressing alveolar damage or as an agent for improving respiratory conditions.

[Example 20] Tissue-preserving effect of GalNAcST siRNA in a mouse pulmonary emphysema model

The objective of this Example is to prove that the therapeutic effect of GalNAcST siRNA is not only the suppression at the cell level but also the effect of morphological maintenance and preservation of the organ.

The pulmonary tissues used in this Example were the right lungs of the same lung tissues as used in Example 12. The lung tissues isolated from mice were gently washed with phosphate buffer and then immersed in phosphate buffer saturated in a glass container. The glass container filled with phosphate buffer was weighed in advance. After the lung tissues were added to the container, lung capacity was calculated by converting the increased weight into a liquid volume.

The result of this Example is shown in Fig. 20. The lung capacity was 277.5 ± 61.85 μl in the control group, while it was 413.33 ± 77.67 μl in the untreated group (when compared to the control group, $P=0.024$; t-test). Thus, a statistically significant increase of lung capacity was observed in the untreated group. In contrast, the lung capacity was 315 ± 51.96 μl and thus significantly decreased in the GalNAcST siRNA-treated group (when compared to the untreated group $P=0.049$; t-test).

When taken together, these results reveal that the inhibition of GAINAc4ST-1, GAINAc4ST-2, and GAINAc4S-6ST gene expressions effectively suppresses the increase in the

lung capacity associated with pulmonary emphysema induced by intratracheal administration of PPE. This suggests that the effect is not only the suppression of the damage at the cell level but also the effect of maintaining and preserving the morphology of the organ or the effect of repairing damaged tissues.

5 The agents of the present invention are thus useful, for example, as agents for maintaining or preserving lung morphology.

[Pancreatic tissue]

10 The Examples below describe preparation of a type 2 diabetes model by administration of Streptozotocin to C57BL/6JcL mice (female; CLEA Japan Inc.) on day 2 after birth and assessment of the mice for weight and blood glucose changes, gene expression, and fibrotic change of pancreatic tissues caused by treatment with chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 1 (C4ST-1) siRNA, chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 2 (C4ST-2) siRNA, and chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 3 (C4ST-3) siRNA. Each siRNA was administered by the same method as described in Example 1. Sequences are shown below.

*[C4ST-1 siRNA cocktail sequences]

C4ST1 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 1)

(GenBank accession number NM_021439)

20 5'-ACAAAGCCATGAAGCCGCGCTGCTGGAAGTGATGAGGATGAACAGAATT-3'
(SEQ ID NO: 55)

5'-CAACCTGAAGACCCTTAACCAGTACA-3' (SEQ ID NO: 56)

5'-GCATCCCAGAGATCAACCACCGCTTG-3' (SEQ ID NO: 57)

*[C4ST-2 siRNA cocktail sequences]

25 C4ST2 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 2)

(GenBank accession number NM_021528)

5'-GCCAGGAGTGGGCCCAGCCCAGGGC -3' (SEQ ID NO: 58)

5'-ATGACCAAGCCGCGGCTCTTCCGGCTG -3' (SEQ ID NO: 59)

5'-AGAGCCTGCTGGACCGGGGCAGCCCCTA -3' (SEQ ID NO: 60)

30 5'-GAGACCCCCTGGACATCCCCCGGGAACA-3' (SEQ ID NO: 61)

*[C4ST-3 siRNA cocktail sequences]

C4ST3 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 3)

(GenBank accession number XM_355798)

5'-ATGACTGTGCGCTGCCACGCGTGCCA -3' (SEQ ID NO: 62)

35 5'-CAGCATGGGAAGACGCTCCTGTTGCA -3' (SEQ ID NO: 63)

5'-TCCAAGCGCAATCCCTGCGCACGAGGCG -3' (SEQ ID NO: 64)

5'-GCCTGGCCTGCTGCCCTCGCTGGCC -3' (SEQ ID NO: 65)

First, sample preparation was conducted as described below.

[Example 21] Assessment of anti-obesity effect of C4ST-1, C4ST-2, and C4ST-3 siRNA treatment in Streptozocin-induced C57BL/6JcL type 2 diabetes model mice

Gestational Day 14 C57BL/6JcL mice (CLEA Japan Inc.) were reared and allowed to deliver. 10 mg/ml Streptozocin (SIGMA) was subcutaneously administered at 20 µl/head to Day 2 postnatal female C57BL/6JcL mice. The mice were reared with sterile water and CE-2 Diet (CLEA Japan Inc.) until they were four weeks old, and then on sterile water and a High Fat Diet (CLEA Japan Inc.) for the next two weeks. On the second week, 1 µg of a mixture of chondroitin D-N-acaetylgalactosamine-4-*O*-sulfotransferase 1 (C4ST-1), chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 2 (C4ST-2), and chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 3 (C4ST-3) siRNAs (GeneWorld) were combined with 1% atelocollagen (Koken Co.), which is a vehicle for siRNA, and 200 µl of the mixture was administered into peritoneal cavities once a week, twice in total (for two weeks). On Day 14 of this experiment, 100 µl of 5 mg/ml BrdU (ZyMED Laboratory Inc.) was administered into the tail vein, and one hour after administration the mice were dissected and the liver, pancreas, kidney, testis, ovary, and muscle were isolated to obtain samples for immunostaining and gene expression analysis. The body weight change was monitored over time for 14 days of this

Example.

The result is shown in Fig. 21. The vertical axis indicates the body weight (g), while the horizontal axis indicates the monitoring days. Fig. 21 shows that the weight increase tended to be suppressed 10 days after treatment in the C4ST-1 siRNA-treated group, C4ST-2 siRNA-treated group, and C4ST-3 siRNA-treated group as compared to the control group. The weight increase was significantly suppressed on day 18 in the C4ST-2 siRNA-treated group and C4ST-3 siRNA-treated group (both groups; $p < 0.05$). This result suggests that obesity associated with type 2 diabetes can be suppressed by inhibiting the expression of C4ST-1, C4ST-2, and C4ST-3.

The agents of the present invention are thus useful, for example, as agents for suppressing body weight increase, or as agents for suppressing obesity associated with type 2 diabetes.

[Example 22] Assessment of insulin resistance after treatment with C4ST-1, C4ST-2, and C4ST-3 siRNAs in Streptozocin-induced C57BL/6JcL type 2 diabetes model mice

As an insulin-tolerance test, human crystalline insulin (0.75 U/kg) was administered into peritoneal cavities the day before dissection, and 0, 15, and 60 minutes after administration, the blood glucose levels were measured and evaluated using a blood glucose monitor, Glu-Test Ace

(BOMBYX Medicine co.). The blood glucose level changes monitored at 0, 15, and 60 minutes after siRNA treatment are shown in Fig. 22. The vertical axis indicates the blood glucose level (mg/dl), while the horizontal axis indicates 0, 15, and 60 minutes after insulin-tolerance test.

Fig. 22 shows that the blood glucose level was not significantly decreased 0 and 15 minutes after treatment in any of the C4ST-1 siRNA-treated group, C4ST-2 siRNA-treated group, and C4ST-3 siRNA-treated group as compared to the control group, while it significantly decreased 60 minutes after treatment in each of the C4ST-1 siRNA-treated group, C4ST-2 siRNA-treated group, and C4ST-3 siRNA-treated group. This result suggests that insulin resistance, which is an essential functional disorder in type 2 diabetes, can be effectively improved by suppressing the expression of C4ST-1, C4ST-2, and C4ST-3.

The agents of the present invention are thus useful, for example, as agents for improving insulin resistance in type 2 diabetes.

[Example 23] Assessment of gene expression in pancreatic tissues after treatment with C4ST-1, C4ST-2, and C4ST-3 siRNAs in Streptozocin-induced C57BL/6JcL type 2 diabetes model mice

cDNA was prepared by the same method as described in Example 1 from 50 mg of each organ (pancreas) isolated from Streptozocin-induced female C57BL/6JcL mice. PCR was conducted in the following composition.

2 µl of PCR Buffer [composition: 166 mM (NH₄)₂SO₄ (Sigma Aldrich Japan), 670 mM Tris pH8.8 (Invitrogen), 67 mM MgCl₂·6H₂O (Sigma Aldrich Japan), 100 mM 2-mercaptoethanol (WAKO)], 0.8 µl of 25 mM dNTP mix (Invitrogen), 0.6 µl of DMSO (Sigma Aldrich Japan), 0.2 µl of Primer Forward (GeneWorld), 0.2 µl of Primer Reverse (GeneWorld), 15.7 µl of Otsuka distilled water (Otsuka Pharmaceuticals, Inc.), 0.1 µl of Taq polymerase (Perkin Elmer), and 1 µl of cDNA obtained as described above were combined, and reacted using Authorized Thermal Cycler (eppendorf) at 35 cycles of 94°C for 45 seconds, 56°C for 45 seconds, and 72°C for 60 seconds. After the reaction, the obtained PCR products were combined with 2 µl of Loading Dye (Invitrogen). 1.5% agarose gel was prepared using UltraPure Agarose (Invitrogen), and the samples were electrophoresed in a Mupid-2 plus (ADVANCE) at 100 V for 20 minutes. After electrophoresis, the gel was shaken for 20-30 minutes in a stain solution prepared by 10,000 times diluting Ethidium Bromide (Invitrogen) with 1x LoTE (composition: 3 mM Tris-HCl (pH 7.5) (Invitrogen), 0.2 mM EDTA (pH 7.5) (Sigma Aldrich Japan)). The gel was photographed with EXILIM (CASIO) positioned on I-Scope WD (ADVANCE) and the gene expression was confirmed.

Primers (Forward and Reverse) (GeneWorld) used are described in the following.

[Primer sequences]

*GAPDH

Forward: 5'-CTGCCAAGTATGACATCA -3' (SEQ ID NO: 66)

Reverse: 5'-TACTCCTTGGAGGCCATGTAG -3' (SEQ ID NO: 67)

*C4ST1 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 1)

Forward: 5'-gtggatgaggaccacgaact-3' (SEQ ID NO: 68)

5 Reverse: 5'-ctttcaagcgggtggtgat-3' (SEQ ID NO: 69)

*C4ST2 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 2)

Forward: 5'-acctcctagaccacacacg-3' (SEQ ID NO: 70)

Reverse: 5'-ggatgttgcaaaccagtct-3' (SEQ ID NO: 71)

*C4ST3 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 3)

10 Forward: 5'-atgagccctcaacgaacac-3' (SEQ ID NO: 72)

Reverse: 5'-tggtagaaggggctgatgac-3' (SEQ ID NO: 73)

Result is shown in Fig. 23. GAPDH expression, which is used as a positive control, was confirmed by RT-PCR in the control group, C4ST1 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 1), C4ST2 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 2), and C4ST3 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 3). Compared to the control group, expression was reduced in C4ST1 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 1) siRNA-, C4ST2 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 2) siRNA-, C4ST3 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 3) siRNA-treated groups, and C4ST1 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 1), C4ST2 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 2), C4ST3 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 3) gene knockdown was confirmed by administration of Atellocollagen-mediated C4ST1 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 1) siRNA, C4ST2 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 2) siRNA, C4ST3 (Chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 3) siRNA.

25

[Example 24] Assessment of accumulation of amyloid precursor protein in pancreas after treatment with C4ST-1, C4ST-2, and C4ST-3 siRNAs in Streptozocin-induced C57BL/6JcL type 2 diabetes model mice

In this Example, the amyloid precursor protein (APP) deposition-suppressing effect of C4ST-1, C4ST-2, and C4ST-3 siRNAs was assessed using pancreatic tissue samples of type 2 diabetes model mice. The deposition of APP and amyloid fibers in islets including β cells has long been known to be an important histopathological feature of type 2 diabetes. In recent years, APP has been demonstrated to be involved in Alzheimer's disease (Johnson KH *et al.* N Eng J Med 321: 513, 1989; Rhodes CJ. Science 307: 380, 2005; Haan MN. Nat Clin Pract Neurol 3: 159, 2006; Prentki M *et al.* J Clin Invest 116: 1802, 2006).

35 The prepared sections of tissue samples were stained with a goat anti-amyloid precursor

protein antibody (Calbiochem) by the same method as described in Example 3 to assess its expression at the tissue level. Fig. 24 shows histological images of the normal mouse group, control group, and C4ST-2 siRNA-treated group. The APP deposition is enhanced in the islets of type 2 diabetes model mice. The deposition was demonstrated to be clearly suppressed in the C4ST-2 siRNA-treated group as compared to the control group.

The agents of the present invention are thus useful, for example, as agents for suppressing amyloid fiber deposition.

[Example 25] Histological fibroblast infiltration-suppressing effect of C4ST-1, C4ST-2, and C4ST-3 siRNA treatment in Streptozocin-induced C57BL/6JcL type 2 diabetes model mice

Cryoblocks and tissue sections were prepared from the collected pancreatic tissues by the same method as described in Example 3. The resulting sections were fixed with acetone (Wako Pure Chemical Industries) for ten minutes, and then washed with phosphate buffer. An anti-ER-TR7 antibody (rat monoclonal antibody, 1 μ g/ml; BMA) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, the secondary antibody reaction was carried out by adding a peroxidase-labeled anti-rat IgG antibody (1:200 dilution), and color development was performed by adding DAB substrate (Nichirei Biosciences). Then, the nucleus was stained by Lillie-Mayer hematoxylin (Muto Pure Chemicals Co.). The samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

The result showed that the fibroblast infiltration to pancreatic islets was significantly suppressed in each of the C4ST-1 siRNA-, C4ST-2 siRNA-, and C4ST-3 siRNA-treated groups as compared to the control group (Fig. 25). This result demonstrates that the inhibition of C4ST-1, C4ST-2, and C4ST-3 gene expression reduces the enhanced tissue fibrotic change by suppressing the fibroblast infiltration and retention in islets containing β cells.

The agents of the present invention are thus useful, for example, as agents for suppressing fibroblast infiltration to islets.

[Example 26] Histological macrophage infiltration-suppressing effect of treatment with C4ST-1, C4ST-2, and C4ST-3 siRNAs in Streptozocin-induced C57BL/6JcL type 2 diabetes model mice

Cryoblocks and tissue sections were prepared from the collected pancreatic tissues by the same method as described in Example 3. The resulting sections were fixed with acetone (Wako Pure Chemical Industries) for ten minutes, and then washed with phosphate buffer. An anti-F4/80 antibody (clone A3-1, rat monoclonal antibody, 2 μ g/ml; CALTAG LABORATORIES) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, the secondary antibody reaction was carried out by adding a

peroxidase-labeled anti-rat IgG antibody (1:200 dilution), and color development was performed by adding DAB substrate (Nichirei). Then, the nucleus was stained by Lillie-Mayer hematoxylin (Muto Pure Chemicals Co.). The samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

The result showed that the macrophage infiltration to the pancreatic islets was significantly suppressed in each of the C4ST-1 siRNA-, C4ST-2 siRNA-, and C4ST-3 siRNA-treated groups as compared to the control group (Fig. 26). This finding demonstrates that the suppression of C4ST-1, C4ST-2, and C4ST-3 gene expressions resulted in general inhibition of the tissue fibrotic changes *via* suppression of the infiltration of macrophages and fibroblasts, which are cell groups responsible for the persistent or enhanced fibrotic changes.

The agents of the present invention are thus useful, for example, as agents for suppressing the infiltration of macrophages to the pancreatic islets.

[Example 27] Assessment of insulin resistance after GalNAcST siRNA treatment in Streptozocin-induced C57BL/6JcL type 2 diabetes model mice

The importance of sulfation at position 4 and 6 is demonstrated with an additional Example. In this Example, the improvement of insulin resistance by GalNAcST siRNA administration is assessed using the same methods as described in Examples 11 and 18. The type 2 diabetes model was prepared by the same method as described in Example 21, and the insulin resistance was tested by the same method as described in Example 22. The result is shown in Fig. 27.

GalNAcST siRNA administration exhibited a good antihyperglycemic effect after insulin loading. This result suggests that the insulin resistance arising from fibrotic changes in islets of pancreatic tissues can be effectively improved by suppressing the expression of GalNAc4ST-1, GalNAc4ST-2, and GalNAc4S-6ST.

The agents of the present invention are thus useful, for example, as hypoglycemic agents or as agents for improving the insulin resistance of pancreatic tissue.

[Kidney tissue]

Fibrotic changes of kidney tissue are thought to be the endpoints of various kidney diseases. (1) From a classical point of view, tubulointerstitial disease is understood as a representative disease caused by renal fibrogenesis. This disease includes Sjogren's syndrome, transplant rejection (chronic allograft nephropathy, etc.), and graft-versus-host reaction (graft-versus-host disease, etc.), in addition to drug-induced, infective, radiation-induced, and heavy metal-induced interstitial renal disorders. (2) Renal fibrogenesis also includes renovascular disease. Renovascular disease includes nephrosclerosis associated with hypertension. In recent

years, this disease also includes fibrogenesis associated with arteriosclerosis and metabolic syndrome. Furthermore, interstitial fibrogenesis also occurs as a result of proteinuria or the like in primary glomerular disease, which leads to renal failure. Thus, renal fibrotic changes can also be developed in: (3) primary glomerular disease and (4) secondary glomerular diseases. The diseases of (3) include IgA nephropathy, minimal lesion nephrotic syndrome, membranous nephropathy, membranoproliferative glomerulonephritis, and focal segmental glomerulosclerosis (FGFS). The diseases of (4) include diabetic nephropathy, lupus nephritis associated with systemic lupus erythematosus (SLE), nephropathy associated with chronic rheumatoid arthritis, amyloid nephropathy, and nephropathy associated with type B or C hepatitis. (5) Finally, renal fibrogenesis also includes interstitial kidney diseases caused by urinary obstruction, including urinary lithiasis, tumors, and neurogenic bladder dysfunctions.

Recently, the group of diseases listed above has been collectively named “chronic kidney disease (CKD)” as a clinical concept which is classified and diagnosed based on the degree of kidney function. CKD gradually progresses through interstitial fibrotic changes, leading to chronic renal failure, and then end-stage renal disease (ESRD). Conventionally, only dialysis has been a definitive treatment for ESRD. Although the possibility of slowing down progression to ESRD has been suggested by using an antihypertensive agent that act on the angiotensin system, therapeutic methods that target renal fibrogenesis itself have not been established. There is a desperate need for new CKD treatment. In terms of the progression process of pathological conditions, treatment targeting fibrotic changes is of the greatest significance as a common therapeutic strategy regardless of the type of primary disease. It has been reported that there are 500 million CKD patients worldwide. It is predicted that the number of patients will continue to increase in the future due to altered lifestyle habits. In addition, a large-scale trial revealed the very high risk of death from a cardiovascular disorder before receiving dialysis. Thus, considering CKD as a disease of the 21st century, active treatment of CKD is a significant challenge in the medical community (Sergio A *et al.*, Hypertension 38: 635, 2001; Weiner DE *et al.*, JASN 15: 1307, 2004; Anavekar NS *et al.*, N Eng J Med 351: 1285, 2004; Remuzzi G *et al.*, J Clin Invest 116: 288, 2006; Tonelli M *et al.*, BMJ 332: 1426, 2006; Khwaja A *et al.*, Kidney International doi:10.1038/sj.ki.5002489).

In the next Examples, the inhibition of sugar chain-related genes that modify the sulfate group at position 4 or 6 was assessed for the effect on histological fibrotic changes in the renal interstitium.

[Example 28] Assessment of anti-fibrogenic effect of C4ST-1 siRNA in a mouse diabetic nephropathy model

A type 2 diabetes model was prepared by the same method as described in Example 21.

Gestational Day 14 of C57BL/6JcL mice (CLEA Japan Inc.) were reared and allowed to deliver. Streptozotocin (STZ; Sigma) was administered to Day 2 postnatal C57BL/6JcL mice to prepare a STZ-induced diabetes model. The mice were subcutaneously injected with 20 μ l of STZ (10 mg/ml) three times for two days. Thus, a total of 60 μ l was administered to the mice. Together
 5 with their mothers, the mice were fed with a normal diet until they are four weeks old. After weanling at the age of four completed weeks, the mice were fed with a High Fat Diet (CLEA Japan Inc.) for two weeks. In the second week, 1 μ g of chondroitin D-N-acetylgalactosamine-4-*O*-sulfotransferase 1 (C4ST-1) siRNA (GeneWorld) was combined with 0.1% atelocollagen (Koken Co.), which is a vehicle for siRNA, and 200 μ l of the resulting mixture was administered
 10 into peritoneal cavities once a week (one shot/week) twice in total (for two weeks). On day 14 of this experiment, the mice were dissected and their kidneys were isolated to obtain samples for immunostaining. Gene expression, body weight, and effect on insulin resistance are described in Examples 21, 22, and 23.

The prepared sections of kidney tissue samples were fixed with acetone (Wako Pure Chemical Industries) for ten minutes, and then washed with phosphate buffer. An anti-ER-TR7 antibody (rat monoclonal antibody, 1 μ g/ml; BMA) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, the secondary antibody reaction was carried out by adding a peroxidase-labeled anti-rat IgG antibody (1:200 dilution), and color development was performed by adding DAB substrate (Nichirei Biosciences). Then,
 15 the nucleus was stained by Lillie-Mayer hematoxylin (Muto Pure Chemicals Co.). The samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

The result showed that the infiltration of fibroblasts (ER-TR7-positive cells) in the renal cortex and medulla was less in the C4ST-1 siRNA-treated group as compared to that in the control group (Fig. 28; the original images are in color). The ER-TR7-positive cells were
 25 counted to quantify the fibrogenesis in inflammatory cells. Each sample was observed with ten microscopic optical fields under a microscope at a magnification of 400 fold, and the number of positive cell were counted and compared between the control group and C4ST-1 siRNA-treated group. The result showed that the ER-TR7 positivity ratio was significantly reduced in the
 30 C4ST-1 siRNA-treated group as compared to the control group ($p < 0.001$).

[Example 29] Assessment of C4ST-1 siRNA for its effect on macrophage infiltration in a mouse diabetic nephropathy model

The prepared sections of kidney tissue samples were fixed with acetone (Wako Pure Chemical Industries) for ten minutes, and then washed with phosphate buffer. An anti-F4/80 antibody (clone A3-1, rat monoclonal antibody, 2 μ g/ml; CALTAG LABORATORIES) was
 35 added as the primary antibody, and the

sections were incubated at room temperature for one hour. Then, the secondary antibody reaction was carried out by adding a peroxidase-labeled anti-rat IgG antibody (1:200 dilution), and color development was performed by adding DAB substrate (Nichirei Biosciences). Then, the nucleus was stained by Lillie-Mayer hematoxylin (Muto Pure Chemicals Co.). The samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

The result showed that the infiltration of macrophages (F4/80-positive cells) in the renal cortex and medulla was less in the C4ST-1 siRNA-treated group as compared to that in the control group (Fig. 29; the original images are in color). Furthermore, the tissue lesions were quantified by the same method as described in Example 28. The result showed that the macrophage positivity ratio was significantly reduced in the C4ST-1 siRNA-treated group as compared to the control group ($p < 0.001$).

[Example 30] Assessment of C4ST-1 siRNA for fibroblast activation in tissues in a mouse diabetic nephropathy model

The sections of kidney tissue samples were fixed with acetone (Wako Pure Chemical Industries) for ten minutes, and washed with phosphate buffer. An anti-human smooth muscle actin antibody (α SMA: mouse monoclonal antibody, 1:100; DACO) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, the secondary antibody reaction was carried out using Histofine Mouse Stain kit (Nichirei), and color development was performed by adding DAB substrate (Nichirei Biosciences). Then, the nucleus was stained by Lillie-Mayer hematoxylin (Muto Pure Chemicals Co.). The samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

The result showed that α SMA-positive cells retained in juxtaglomerular and interstitial areas were significantly reduced in the C4ST-1 siRNA-treated group as compared to the control group (Fig. 30; the original images are in color). α SMA serves as a functional marker for fibroblast activation. This result demonstrates that inhibition of C4ST-1 gene expression suppress activation of fibroblasts infiltrating in the tissue.

The agents of the present invention are thus useful, for example, as fibroblast activation inhibitors.

[Example 31] Assessment of GalNAc4S-6ST siRNA for tissue fibrogenic alteration in a mouse diabetic nephropathy model

The diabetic nephropathy model was prepared by the same method as described in Example 28 to assess the effect of GalNAc4S-6ST siRNA. In this Example, the follow-up

examination was carried out over a longer period of time. The sequence of GalNAc4S-6ST siRNAs was the same as shown in Example 1. The siRNAs were administered into peritoneal cavities once at the age of eight weeks and once again at age of nine weeks. Furthermore, as a control for comparison, Valsartan, an angiotensin II receptor blocker (ARB), was orally administered at a dose of 30 mg/kg on the same schedule. Kidney tissues were collected at the age of ten weeks to conduct immunohistochemical and gene expression tests. To assess the gene expression in the kidney tissues, quantitative PCR was carried out by the same method as described in Example 1. In this Example, 36B4 was used as an internal control. The sequence of 36B4 is shown below.

[Primer sequences used in quantitative PCR]

*mouse 36B4 (Takara Bio)

Forward: 5'-TTCCAGGCTTTGGGCATCA -3' (SEQ ID NO: 74)

Reverse: 5'-ATGTTTCAGCATGTTTCAGCAGTGTG -3' (SEQ ID NO: 75)

The result is shown in Fig. 31 (GalNAc4S-6ST is abbreviated as G#1 in this figure). In the diabetic nephropathy model, the expression of GalNAc4S-6ST in kidney tissues is enhanced. The administration of GalNAc4S-6ST siRNA significantly suppressed not only the expression of GalNAc4S-6ST gene in kidney tissues but also the enhanced expression of α SMA and TGF β , which are fibrosis markers. The therapeutic effect was evaluated by comparing with that of ARB, and the result showed that the TGF β -suppressing effect was observed in both groups and there was no significant difference between the two while the α SMA-suppressing effect was significant in the GalNAc4S-6ST siRNA-administered group as compared to the ARB-administered group. This result suggests that the markers for fibrotic changes in kidney tissues can be suppressed by inhibiting the expression of GalNAc4S-6ST gene. The result also demonstrates that GalNAc4S-6ST siRNA showed a markedly superior effect in terms of fibroblast activation (enhancement of α SMA).

[Example 32] Assessment of GalNAc4S-6ST siRNA for fibroblast infiltration in a mouse diabetic nephropathy model

The degree of fibroblast infiltration into kidney tissues was quantitatively evaluated by conducting an immunohistochemical study using the same method as described in Example 28. The result is shown in Fig. 32. Fibroblast infiltration to kidney tissues was significantly suppressed by GalNAc4S-6ST siRNA administration. A quantitative evaluation over the entire interstitium did not show any significant difference between GalNAc4S-6ST siRNA administration and ARB treatment. Meanwhile, when the evaluation was restricted to fibroblasts infiltrating in juxtaglomerular areas, a significant suppressing effect was observed in the GalNAc4S-6ST siRNA-administered group, as shown in the figure.

The agents of the present invention are thus useful, for example, as agents for suppressing fibroblast infiltration into renal interstitium.

[Example 33] Assessment of GalNAc4S-6ST siRNA in macrophage infiltration in a mouse diabetic nephropathy model

The degree of fibroblast infiltration into kidney tissues was quantitatively evaluated by conducting an immunohistochemical study using the same method as described in Example 29. The result is shown in Fig. 33. Macrophage infiltration into kidney tissues was significantly suppressed by GalNAc4S-6ST administration. A quantitative evaluation over the entire interstitium did not show any significant difference between GalNAc4S-6ST siRNA administration and ARB treatment. Meanwhile, when the evaluation was restricted to macrophages infiltrating in juxtaglomerular areas, a significant suppressing effect was observed in the GalNAc4S-6ST siRNA-administered group, as shown in the figure.

The agents of the present invention are thus useful, for example, as agents for suppressing macrophage infiltration to renal interstitium.

[Example 34] Assessment of GalNAc4S-6ST siRNA in glomerular basement membrane thickening in a mouse diabetic nephropathy model

Type IV collagen was immunostained by the same method as described in Examples 32 and 33. A rabbit anti-mouse type IV collagen antiserum (LSL) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, the secondary antibody reaction was carried out by adding a peroxidase-labeled anti-rabbit IgG antibody (1:200 dilution), and color development was performed by adding DAB substrate (Nichirei Biosciences). Then, the nucleus was stained by Lillie-Mayer hematoxylin (Muto Pure Chemicals Co.). The samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

The thickness of collagen visualized as brown signals surrounding glomeruli was measured to quantify the accumulation of type IV collagen. 15 to 20 glomeruli were assessed for each sample. The thickness of the thickest portion around a glomerulus was measured on a display monitor using vernier calipers. The result showed that glomerular basement membrane (GBM) thickening was significantly suppressed in the GalNAc4S-6ST siRNA-administered group as compared to the control group (Fig. 34). The thickening tended to be suppressed more markedly in the GalNAc4S-6ST siRNA-administered group as compared to ARB.

The agents of the present invention are thus useful, for example, as agents for suppressing glomerular basement membrane thickening.

[Example 35] Assessment of GalNAc4S-6ST siRNA in the angiotensin pathway in a mouse diabetic nephropathy model

Angiotensin II has been reported to be involved in fibrogenesis in diabetic nephropathy. In this Example, the angiotensin pathway in kidney tissues was assessed by quantitative PCR.

5 The expression of angiotensinogen and angiotensin converting enzyme (ACE) is enhanced in this model (Fig. 35). The enhanced expression was speculated to be a factor responsible for the renal enhancement of angiotensin II. The effect on suppressing angiotensinogen and ACE was observed in the GalNAc4S-6ST siRNA-administered group (Fig. 35). The result demonstrates that GalNAc4S-6ST siRNA also produces an improving effect on the angiotensin pathway
10 through suppression of fibroblast infiltration and activation by suppressing the GalNAc4S-6ST gene.

The agents of the present invention are thus useful, for example, as agents for suppressing the expression of angiotensinogen or as an angiotensin converting enzyme inhibitor.

15 [Example 36] Assessment of GalNAc4S-6ST siRNA in serum creatinine concentration in a mouse diabetic nephropathy model

Serum creatinine is a most commonly used clinical marker for renal function. The serum creatinine concentration is elevated due to impaired renal function during the process from diabetic nephropathy to ESRD. However, it has been revealed that functional nephrons are
20 already functionally impaired by 50% when the serum creatinine level is elevated. Protecting renal function before the creatinine level increases is a clinically important challenge.

In this model as well, elevation of serum creatinine level was observed eventually at the age of 18 weeks when histological fibrogenesis had already progressed (Fig. 36). The elevation of serum creatinine was suppressed in the GalNAc4S-6ST siRNA-administered group as
25 compared to the control group (Fig. 36). This result suggests that inhibition of the GalNAc4S-6ST gene results in suppression of fibrotic changes of renal tissue and thereby suppresses the elevation of serum creatinine, i.e., the deterioration of renal function. Thus, suppression of the GalNAc4S-6ST gene produces a very beneficial renal protective effect.

30 The agents of the present invention are thus useful, for example, as agents for suppressing the deterioration of renal function, or as a renal protective agent.

[Example 37] Assessment of anti-fibrogenic effect of GalNAcST siRNA in a mouse diabetic nephropathy model

The importance of sulfation at position 4 and 6 is shown with an additional Example.

35 The effect of GalNAcST siRNA administration on fibrotic changes in the renal interstitium was assessed by the same method as described in Example 11. The schedule of GalNAcST siRNA

administration is the same as described in Example 28. Quantitative PCR was carried out using kidney tissues by the same method as described Example 31. In this Example, β -actin was used as an internal control. The sequence of β -actin is shown below.

[Quantitative PCT primer sequences]

5 *mouse β actin (Takara Bio)

Forward: 5'-CATCCGTAAAGACCTCTATGCCAAC -3' (SEQ ID NO: 76)

Reverse: 5'-ATGGAGCCACCGATCCACA -3' (SEQ ID NO: 77)

As shown in Fig. 37, the expression of GalNac4ST-1, GalNac4ST-2, and GalNac4S-6ST was enhanced in kidney tissues in the diabetic nephropathy model. The expression of all the genes was significantly suppressed by administering GalNacST siRNA.

[Example 38] Assessment of anti-fibrogenic and kidney-protecting effects of GalNacST siRNA in a mouse diabetic nephropathy model

The enhanced expression of α SMA, TGF β , and CTGF, which are fibrogenesis markers for kidney tissue fibrogenesis, was significantly suppressed in the GalNacST siRNA-administered group as compared to the control group (Fig. 38). The enhanced expression of ACE was also significantly suppressed in the GalNacST siRNA-administered group. This result demonstrates that inhibition of GalNac4ST-1, GalNac4ST-2, and GalNac4S-6ST expression results in suppression of kidney tissue fibrogenesis and renal protection. Together with the result of Example 35, the fact that ACE was markedly reduced by suppressing any of the genes demonstrates that they have a hypotensive effect.

The agents of the present invention are thus useful, for example, as antihypertensive agents.

25 [Example 39] Assessment of GalNac4S-6ST siRNA for gene expression in a mouse drug-induced interstitial nephritis model

This Example assesses the effect of GalNac4S-6ST siRNA on a typical drug-induced interstitial nephritis. First, the mouse model is prepared as described below. Adriamycin (15 mg/kg; Kyowa Hakko) was administered to the peritoneal cavities of C57BL6/J mice (male, eight weeks old, CLEA Japan Inc.). The mice were reared for one week after administration, and then kidney tissues were collected from them. As a control group, mice of the same lineage and age were also purchased and reared in the same period, but Adriamycin was not given to them.

GalNac 4S-6ST siRNA was administered by the same method as described in Example 1: 1 μ g of GalNac 4S-6ST siRNA (Hokkaido System Science Co.) was combined with 200 μ l of 1% atelocollagen (Koken Co.), which is a vehicle, and the resulting mixture was administered intraperitoneally to each mouse 24 hours before Adriamycin administration. The expression of

GalNac4S-6ST is also enhanced in kidney tissues in the typical drug-induced interstitial nephritis model described in this Example (Fig. 39). The expression was significantly suppressed by administering GalNac4S-6ST siRNA.

5 [Example 40] Assessment of GalNac4S-6ST siRNA in gene expression in a mouse drug-induced interstitial nephritis model

In this Example, immunostaining of type I collagen was carried out by the same method as described in Example 34. A rabbit anti-rat type I collagen antiserum (LSL) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, the secondary antibody reaction was carried out by adding a peroxidase-labeled anti-rabbit IgG antibody (1:200 dilution), and color development was performed by adding DAB substrate (Nichirei Biosciences). Then, the nucleus was stained by Lillie-Mayer hematoxylin (Muto Pure Chemicals Co.). The samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

15 The result is shown in Fig. 40. Diffuse deposition of type I collagen was observed over juxtaglomerular and interstitial areas in the control group, while the deposition was markedly suppressed in the GalNac4S-6ST-administered group. This result suggests that fibrotic changes in kidney tissues can also be suppressed by suppressing the expression of GalNac4S-6ST in the drug-induced interstitial nephritis.

20 The agents of the present invention are thus useful, for example, as agents for suppressing fibrotic changes in kidney tissues.

[Example 41] Assessment of anti-fibrosis effect of C6ST siRNA in renal fibrosis model mice induced by unilateral ureteral obstruction (UUO)

25 First, a mouse model for renal fibrosis is prepared as described below. Renal fibrosis model mouse was prepared by conducting unilateral ureteral obstruction (UUO) to C57BL/6JcL mice (female, eight weeks old; CLEA Japan Inc.). This model has excellent reproducibility, and is thus widely used as an experimental mouse renal fibrosis model (American Journal of Pathology 2003 163 (4): 1261-1273). Mice were subjected to laparotomy under Ketalar/xylazine anesthesia. The ureters were exposed and the right ureter was ligated at two sites with 4-0 surgical suture. The peritoneum and skin were closed with 1-0 surgical suture.

30 The effect of inhibiting C6ST expression by C6ST siRNA administration was checked by PCR method using renal fibrosis model mice prepared by unilateral ureteral obstruction (UUO), as a typical example of renal fibrosis model mice. The renal fibrosis model was prepared by conducting UUO to C57BL/6JcL mice (female, eight weeks old; CLEA Japan Inc.). A mixture of C6ST-1 and C6ST-2 siRNAs (1 µg/head; GeneWorld) or PBS was combined with

0.1% atelocollagen (Koken Co.), which is a vehicle for siRNA, and 200 µl of the resulting mixture was injected into the peritoneal cavity of each mouse. Groups of mice treated as described above were named C6ST siRNA group and control group. On day 8 of the experiment, the mice were dissected to excise the UUO-treated kidney. Thus, samples for immunostaining and gene expression analysis were obtained from the mice. Quantitative PCR was carried out by the same method as described in Example 1.

C6ST-1 siRNA, C6ST-2 primers (Forward, Reverse) (GeneWorld) used herein are shown below.

[Primer sequences]

*C6ST1 (Chondroitin 6-sulfotransferase-1)

Forward: 5'-tgtgtggacacacctcccta-3' (SEQ ID NO: 78)

Reverse: 5'-cttcaaaggtccccttctc-3' (SEQ ID NO: 79)

*C6ST2 (Chondroitin 6-sulfotransferase-2)

Forward: 5'-cagcttgagccattcaaca-3' (SEQ ID NO: 80)

Reverse: 5'-gggtgaggccttaggaac-3' (SEQ ID NO: 81)

[C6ST-1 cocktail sequences] (Gene Bank accession number NM_016803)

(GeneWorld)

5'-gcgccccctctcccatggagaaag-3' (SEQ ID NO: 82)

5'-gctttgcctcaggatttcgggacc-3' (SEQ ID NO: 83)

5'-ggttcagccttggtctaccgtgatgtc-3' (SEQ ID NO: 84)

5'-gcagttgttgctatgcgacctgtat-3' (SEQ ID NO: 85)

[C6ST-2 cocktail sequences] (Gene Bank accession number NM_021715)

(GeneWorld)

5'-tgaggagagtgaggattcggtgaa-3' (SEQ ID NO: 86)

5'-cggacgtgggactcgtcgaggacaaag-3' (SEQ ID NO: 87)

5'-cgaaagtacctgcccgccttctgc-3' (SEQ ID NO: 88)

The result showed that the expression of C6ST-2 (G#10) was enhanced in the kidney (Fig. 41; C6ST-2 is abbreviated as G#10). The expression level was decreased in the C6ST siRNA-treated group. The C6ST-2 gene knockdown was confirmed in the atelocollagen-mediated C6ST siRNA administration (Fig. 41). The C6ST siRNA administration also significantly suppressed the enhanced expression of fibrogenesis markers: TGFβ, αSMA, type I collagen, and CTGF.

[Example 42] Assessment of C6ST siRNA for fibroblast infiltration in renal fibrosis model mice induced by unilateral ureteral obstruction (UUO)

The isolated tissues were embedded in OCT compound (Sakura Finetechnical Co.), an

embedding medium for cryosectioning. Cryoblocks were prepared using liquid nitrogen, and sliced into 6- μ m thick sections using Cryostat (Micro-edge Instruments Co.). The resulting sections were immunostained by the same method as described in Example 28 using an anti-ER-TR7 antibody. The samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals and quantified.

The result showed that the accumulation of fibroblasts in the renal interstitium was significantly suppressed in the UUO-treated kidney in the C6ST siRNA-treated group as compared to the control group (Fig. 42).

The agents of the present invention are thus useful, for example, as agents for suppressing fibroblast accumulation in kidney interstitium.

[Example 43] Assessment of C6ST siRNA in macrophage infiltration in renal fibrosis model mice induced by unilateral ureteral obstruction (UUO)

Immunostaining and quantitation was carried out by the same method as described in Example 42 using an anti-F4/80 antibody. The result showed that the accumulation of macrophages in the renal interstitium was significantly suppressed in the UUO-treated kidney in the C6ST siRNA-treated group as compared to the control group (Fig. 43).

The agents of the present invention are thus useful, for example, as agents for suppressing macrophage accumulation in kidney interstitium.

[Example 44] Assessment of C6ST siRNA in collagen accumulation in a renal fibrosis mouse model included by unilateral ureteral obstruction (UUO)

Immunostaining was carried out by the same method as described in Example 42 using an anti-type IV collagen antibody, and quantitation was achieved by the same method as described in Example 34. The result showed that the fibrous thickening of glomerular basement membrane in the UUO-treated kidney was significantly suppressed in the C6ST siRNA-treated group as compared to the control group (Fig. 44). This suggests that the infiltration and fibrogenesis of inflammatory cells can be suppressed by inhibiting the expression of C6ST-2 gene.

The agents of the present invention are thus useful, for example, as agents for suppressing the infiltration of inflammatory cells.

[Example 45] Assessment of C6ST siRNA for fibroblast activation in a renal fibrosis mouse model induced by unilateral ureteral obstruction (UUO)

Immunostaining was carried out by the same method as described in Example 30 using an α SMA antibody. The result showed that α SMA-positive cells were clearly decreased in the

interstitium, juxtaglomerular areas in particular, of the UUO-treated kidneys in the C6ST siRNA-treated group as compared to the control group (Fig. 45). This result suggests that the activation of fibroblasts accumulating in the renal interstitium is suppressed by inhibiting the expression of C6ST-1 and C6ST-2 genes.

The agents of the present invention are thus useful, for example, as agents for suppressing the activation of fibroblasts in the renal interstitium.

[Example 46] Assessment of C6ST siRNA in ACE expression in a renal fibrosis mouse model induced by unilateral ureteral obstruction (UUO)

Immunostaining was carried out by the same method using a rabbit anti-human ACE antibody (Santa Cruz). The result showed that ACE-positive cells were clearly decreased in the interstitium, juxtaglomerular areas in particular, of the UUO-treated kidneys in the C6ST siRNA-treated group as compared to the control group (Fig. 46). This result demonstrates that inhibition of C6ST-2 gene expression results in the suppression of ACE expression with the suppression of the activation of fibroblasts accumulated in the renal interstitium. Suppression of ACE produces a hypotensive effect. Thus, the result described above strongly suggests that C6ST siRNA has antihypertensive activity or arteriosclerosis-suppressing effect.

The agents of the present invention are thus useful, for example, as arteriosclerosis-suppressing agents.

[Ocular tissue]

Like other organs, fibrotic changes in ocular tissues occur as a result of invasion due to various causes. This leads to impairment and/or loss of vision. Such major diseases include diabetic retinopathy, retinal vein occlusion, retinopathy of prematurity, age-related macular degeneration, and retinitis pigmentosa. The diseases also include fibrogenesis associated with corneal inflammation, glaucoma, or cataract (reviews: Fiedlander M. J Clin Invest 117: 576-586, 2007; Harada T *et al.* Genes and Dev. 21: 367-378, 2007). From the histopathological viewpoint, damage/decrease of photoreceptor cells caused by fibrogenesis is the major cause of visual loss. Thus, suppressing fibrogenesis in ocular tissues has been expected as a novel therapeutic strategy to prevent visual loss in all ocular diseases.

Next Examples focus on retinal fibrogenesis and the resulting loss of photoreceptor cells in a diabetic retinopathy model.

[Example 47] Assessment of GalNAc4S-6ST (G#1) siRNA in collagen accumulation in a mouse diabetic retinopathy model

Gestational Day 14 C57BL6J/JcL mice (CLEA Japan Inc.) were reared and allowed to

deliver. 10 mg/ml Streptozocin (SIGMA) was subcutaneously administered at 20 µl/head to Day 2 postnatal female C57BL6J/JcL mice. The mice were reared with sterile water and CE-2 Diet (CLEA Japan Inc.) until they were four weeks old, and then with sterile water and a High Fat Diet (CLEA Japan Inc.) for subsequent two weeks. 1 µg of GalNac4S-6ST (G#1) siRNA (Hokkaido System Science Co.) was combined with 200 µl of 1% atelocollagen (Koken Co.), which is a vehicle, and the resulting mixture was administered into the peritoneal cavity of each mouse in the eighth and ninth week by the same method as described in Example 11. In the 18th week, eye balls were excised from mice of the two groups and immunohistochemically examined to assess the effect of GalNac4S-6ST (G#1) siRNA.

Cryoblocks and sections were prepared from the excised eye balls. The sections were fixed with acetone (Sigma Aldrich Japan) for ten minutes, and then washed with phosphate buffer. A rabbit anti-type IV collagen antiserum (1:2000 dilution; LSL) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, a peroxidase-labeled anti-rabbit IgG antibody (1:25 dilution; Cappel) was added as the secondary antibody, and the samples were incubated at room temperature for 30 minutes. After incubation, an enzyme-mediated chromogenic reaction was conducted by adding DAB substrate (Nichirei Biosciences). The samples were observed under a light microscope (Leica Microsystems).

Obtained histology of the retina is shown in Fig. 47. In the control group, the deposition of type IV collagen was increased over the region from ganglion cell layer (GCL) to inner nuclear layer (INL), which is essential for vision. By contrast, the increase of collagen in GCL in particular, was significantly suppressed in the GalNac4S-6ST (G#1) siRNA-administered group.

The agents of the present invention are thus useful, for example, as agents for suppressing collagen increase in the ganglion cell layer.

[Example 48] Assessment of GalNac4S-6ST (G#1) siRNA in the accumulation of sodium chondroitin sulfate proteoglycan in a mouse diabetic retinopathy model

The retina was immunohistochemically assessed using the same method as described in Example 47. An anti-chondroitin sulfate proteoglycan (CSPG) antibody (clone CS56, mouse monoclonal antibody, 1:100; Seikagaku Co.) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, the secondary antibody reaction was carried out using Histofine Mouse Stain kit (Nichirei; used for mouse monoclonal antibody).

As shown in Fig. 48, a significant enhancement of CS56-positive signals were observed in GCL, and the segment from outer nuclear layer (ONL) to pigmented cell layer of retina in the control group. In contrast, the CS56 signal intensity was markedly reduced in the GalNac4S-6ST (G#1) siRNA-administered group. Thus, gangliocytes and retinal pigment epithelial cells

were morphologically well preserved in GCL.

The result described above demonstrates that *in vivo* administration of GalNac4S-6ST (G#1) siRNA significantly suppresses the induced CSPG deposition in retinal tissues in this model mouse. Together with collagen, CSPG is considered to be essential for the formation of fibrogenic lesions. Furthermore, CSPG has been reported to inhibit the process of axon extension of gangliocyte (Brittis PA *et al.*, Science 255: 733, 1992). Previously published reports only describe results of *in vitro* experiments and developmental process. Thus, the role in *in vivo* pathological lesions still remains unknown. However, the result described herein for the first time suggests the role of CSPG in lesional tissues.

[Example 49] Assessment of GalNac4S-6ST (G#1) siRNA in the accumulation of glial cells in a mouse diabetic retinopathy model

Optic nerve regeneration has been reported to serve as a biological defense mechanism after retinal damage. Meanwhile, it is reported that the optic nerve progenitor cells responsible for such regeneration after injury are glial cells (Fischer AJ *et al.*, Nature neuroscience 4: 247, 2001; Ooto S *et al.*, PNAS 101: 13645, 2004). In this Example, immunostaining was carried out using a goat anti-GFAP antibody (Santa Cruz) as a glial cell marker by the same method as described in Example 47.

The result is shown in Fig. 49. The number of GFAP-positive glial cells was not altered in both normal and control groups. In contrast, the cell count was markedly increased in the area from INL to GCL in the GalNac4S-6ST (G#1) siRNA-administered group. Optic nerve regeneration has been reported to occur from INL toward GCL. Thus, the result described above suggests the process of active optic nerve regeneration induced by GalNac4S-6ST (G#1) siRNA.

The agents of the present invention are thus useful, for example, as agents for regenerating the optic nerve.

[Example 50] Assessment of GalNac4S-6ST (G#1) siRNA in gangliocytes in a mouse diabetic retinopathy model

Gangliocytes were quantified using samples prepared in Example 47. The result revealed that GCL gangliocytes were reduced in the diabetic retinopathy model but the loss was significantly recovered by GalNac4S-6ST (G#1) siRNA administration (Fig. 50).

[Example 51] Assessment of GalNac4S-6ST (G#1) siRNA in gene expression in a mouse diabetic retinopathy model

RNA was extracted from ocular tissues and quantitative PCR was carried out by the same method as described in Example 1. The ability to regenerate the optic nerve was assessed

by analyzing changes in the expression of glutamate synthetase (GS), which is a Muller cell marker. The sequences of PCR primers are shown below.

[Quantitative PCR primer sequences]

*mouse GS (Takara Bio Inc.)

5 Forward: 5'-CTGTGAGCCCAAGTGTGTGGA -3' (SEQ ID NO: 89)

Reverse: 5'-GTCTCGAAACATGGCAACAGGA -3' (SEQ ID NO: 90)

The result is shown in Fig. 51. Administration of GalNac4S-6ST (G#1) siRNA could significantly inhibit the enhanced expression of GalNac4S-6ST in ocular tissues. The expression inhibition resulted in a significant increase in the expression of GS in ocular tissues in the GalNac4S-6ST (G#1) siRNA-administered group. The result described above shows that GalNac4S-6ST (G#1) siRNA administration results in Muller cell regeneration, i.e., optic nerve restoration.

This Example revealed that fibrotic changes of retinal tissues could be suppressed by inhibiting the expression of GalNac4S-6ST gene, which led to the prevention of photoreceptor cell loss through the regeneration of gangliocytes. This histological feature is commonly observed in a wide variety of ocular diseases with fibrosis including glaucoma and diabetic retinopathy.

The agents of the present invention are thus useful, for example, as agents for regenerating Muller cells or gangliocytes.

20

[Liver tissue]

Fibrotic changes in liver tissues are the progressive or terminal stage of various liver diseases. Liver fibrogenesis results from various liver diseases, including viral hepatitis (type A, B, C, D, E, and G viral hepatitis), alcohol liver disease, nonalcoholic fatty liver diseases (NAFLD and NASH), metabolic liver disease, drug-induced liver disease, idiopathic portal hypertension, Budd-Chiari's syndrome, autoimmune hepatitis, primary biliary cirrhosis, primary sclerosing cholangitis, biliary disorders (including biliary atresia and biliary dilation), biliary atresia caused by pancreatic diseases such as tumor, graft-versus-host reaction, and chronic rejection (Bataller R *et al.*, J Clin Invest 115: 209, 2005; Iredale JP. J Clin Invest 117: 539, 2007).

25 Fibrotic changes of liver at the tissue level were assessed in the next Examples.

30

[Example 52] Assessment of GalNacST siRNA in gene expression in a mouse fatty liver disease model

Gestational day 14 C57BL/6JcL mice (CLEA Japan Inc.) were reared and allowed to deliver. Streptozocin (STZ; SIGMA) was administered to Day 2 postnatal C57BL/6JcL mice. STZ (10 mg/ml) was subcutaneously administered at 20 µl/head three times for two days. Thus,

35

a total of 60 μ l was administered to the mice. Together with their mothers, the mice were fed with a normal diet until they were four weeks old. After weanling at the age of four weeks, the mice were fed with a High Fat Diet (CLEA Japan Inc.) for two weeks. In the second week, 200 μ l of GalNAcST siRNA described in Example 11 was administered into peritoneal cavities once a week (one shot/week) twice in total (for two weeks). On day 14 of the experiment, the mice were dissected and their livers were isolated to prepare samples for gene expression analysis and immunostaining.

Quantitative PCR was carried out by the same method as described in Example 1 using liver tissues. The result is shown in Fig. 52. The expression of GalNAc4S-6ST is enhanced in liver tissues in this model. GalNAcST siRNA administration resulted in significant suppression of the expression.

[Example 53] Assessment of anti-fibrogenic effect of GalNAcST siRNA in a mouse fatty liver disease model

The expression of fibrogenesis markers in liver tissues was assessed by the same method as described in Example 52. The expression of type I collagen and α SMA was significantly enhanced in this model (Fig. 53), suggesting enhanced fibrotic changes in liver tissues. Meanwhile, the enhanced expression of the fibrogenesis markers was significantly suppressed by administering GalNAcST siRNA.

[Example 54] Assessment of GalNAcST siRNA in fibroblast infiltration in a mouse fatty liver disease model

Next, liver tissues were immunostained by the same method as described in Example 3. A rat anti-mouse fibroblast antibody (clone ER-TR7, 1:500 dilution; BMA) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, a peroxidase-labeled anti-rat IgG antibody (1:200 dilution; Biosource International, Inc.) was added as the secondary antibody, and the sections were incubated at room temperature for 30 minutes. After incubation, DAB substrate (Nichirei Biosciences) was added, and the samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

Examples of obtained images of immunostained liver tissues are shown in Fig. 53. The accumulation of fibroblasts was clearly observed and a bridge formation was confirmed in the histological picture in the control group. In contrast, there was almost no accumulation of fibroblasts in the GalNAcST siRNA-administered group.

[Example 55] Assessment of GalNAcST siRNA in fibrogenic score in a mouse fatty liver disease

model

Each of the samples immunohistochemically stained in Example 54 was assessed for the degree of live fibrogenesis using fibrogenesis scores based on previous reports (Dai K, *et al.*, World J Gastroenterol. 31: 4822-4826, 2005; Hillebrandt S, *et al.*, Nature Genetics 37: 835-843, 2005). The fibrogenesis scores were defined according to the following criteria: 0, normal; 1, few collagen fibrils extend from the central vein; 2, extension of collagen fibrils is apparent but collagen fibrils have not yet encompassed the whole liver; 3, extension of collagen fibrils is apparent and collagen fibrils have encompassed the whole liver; 4, diffuse extension of collagen fibrils is observed in the whole liver and pseudolobules are formed.

The result is shown as a graph in Fig. 55. Each bar indicates mean $\pm$ standard deviation of the fibrogenesis score in each group. The fibrogenesis was statistically significantly reduced in the GalNAcST siRNA-administered group as compared to the control group ($p < 0.01$; t-test). This finding suggests that GalNAcST siRNA administration also clinically produces a superior liver fibrogenesis-suppressing effect.

[Example 56] Assessment of GalNAcST siRNA in macrophage infiltration in a mouse fatty liver disease model

Next, liver tissues were immunostained by the same method as described in Example 54. A rat anti-mouse F4/80 antibody (clone A3-1, 2 μ g/ml; CALTAG LABORATORIES) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, a peroxidase-labeled anti-rat IgG antibody (1:200 dilution; Biosource International, Inc.) was added as the secondary antibody, and the sections were incubated at room temperature for 30 minutes. After incubation, DAB substrate (Nichirei Biosciences) was added, and the samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

Examples of obtained images of immunostained liver tissues are shown in Fig. 56. The accumulation of macrophages was clearly observed in the control group. Formation of inflammatory accumulation lesions were seen in the histological pictures. In contrast, there was no excessive accumulation of macrophages in the GalNAcST siRNA-administered group.

[Example 57] Assessment of GalNAcST siRNA in the hepatic lipid metabolism in a mouse fatty liver disease model

The expression of lipid metabolism-related genes in the liver was assessed by the same method as described in Example 52. The enhanced expression of carbohydrate response element-binding protein (ChREBP) and acetyl-CoA carboxylase-2 (ACC2) was observed. Meanwhile, the enhanced expression was significantly suppressed in the GalNAcST siRNA-

administered group (Fig. 57). This result shows that the glycolipid metabolism can be improved *via* suppression of fibrotic changes in liver tissues by inhibiting the expression of GalNAc4ST-1, GalNAc4ST-2, and GalNAc4S-6ST genes.

The agents of the present invention are thus useful, for example, as agents for improving glycolipid metabolism.

[Example 58] Assessment of C4ST-1 siRNA, C4ST-2 siRNA, and C4ST-3 siRNA for fibroblast accumulation in a mouse fatty liver disease model

A fatty liver disease model was prepared by the same method as described in Example 52. C4ST-1 siRNA, C4ST-2 siRNA, and C4ST-3 siRNA described in Example 21 were administered according to the same administration protocol. Liver tissues were collected from the mice.

Next, liver tissues were immunostained by the same method as described in Example 3. A rat anti-mouse fibroblast antibody (clone ER-TR7, 1:500 dilution; BMA) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, a peroxidase-labeled anti-rat IgG antibody (1:200 dilution; Biosource International, Inc.) was added as the secondary antibody, and the sections were incubated at room temperature for 30 minutes. After incubation, DAB substrate (Nichirei Biosciences) was added, and the samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

Examples of obtained images of immunostained liver tissues are shown in Fig. 58. The accumulation of fibroblasts was clearly observed in the control group. A bridge formation was seen in the histological pictures. In contrast, fibroblast accumulation was not observed in any of the C4ST-1 siRNA-, C4ST-2 siRNA-, and C4ST-3 siRNA-administered group.

[Example 59] Assessment of C4ST-1 siRNA, C4ST-2 siRNA, and C4ST-3 siRNA in the fibrogenic score in a mouse fatty liver disease model

Each sample was assessed for the degree of liver fibrogenesis using the fibrogenesis scores based on the immunohistochemical staining carried out in Example 58. The result is shown in Fig. 59. Each bar indicates mean $\pm$ standard deviation of the fibrogenesis score in each group. The fibrogenesis was statistically significantly reduced in all of the C4ST-1 siRNA-, C4ST-2 siRNA-, C4ST-3 siRNA-administered groups as compared to the control group ($p < 0.001$; t-test). This finding suggests that the administration of siRNAs against C4ST-1, C4ST-2, or C4ST-3 also clinically produces a superior liver fibrogenesis-suppressing effect.

[Example 60] Assessment of C4ST-1 siRNA, C4ST-2 siRNA, and C4ST-3 siRNA in hepatocyte

disorders in a mouse fatty liver disease model

When mice were sacrificed, blood was collected from them according to the protocol as described in Example 58. The blood samples were custom-assayed for the serum alanine transferase (ALT) levels through SRL Inc. The result is shown in Fig. 60. The serum ALT level is the most widely used clinical indicator for hepatocyte destruction. The serum ALT level was elevated in the control group. Meanwhile, the mean value was decreased to 50% or less in each of the C4ST-1 siRNA-, C4ST-2 siRNA-, and C4ST-3 siRNA-administered groups as compared to the control group (Fig. 60). This result demonstrates that hepatocyte damages can be reduced *via* suppression of fibrotic changes by inhibiting the expression of C4ST-1, C4ST-2, or C4ST-3 gene in liver tissues.

The agents of the present invention are thus useful, for example, as agents for reducing hepatocyte damage.

[Example 61] Assessment of the anti-fibrogenic effect of C6ST siRNA in a mouse hepatic fibrosis model

In this Example, experiments were carried out using a mouse cirrhosis model induced by carbon tetrachloride, which is the most widely used cirrhosis model. First, the mouse model was prepared as described below. Carbon tetrachloride (25 μ l/100 g body weight; Sigma-Aldrich) was injected into peritoneal cavities of C57BL6/J mice (female, five or six weeks old; CLEA Japan Inc.) twice a week for four weeks (eight times) to induce hepatic fibrosis. Then, carbon tetrachloride was additionally administered twice a week for two weeks (a total of 12 times) to induce cirrhosis. Mice with induced cirrhosis were sacrificed, and their livers were collected (cirrhotic livers). Meanwhile, in the control experiment, livers were collected from C57BL6/J mice (female, CLEA Japan Inc.) of the same age without carbon tetrachloride administration (normal livers).

The same C6ST siRNA as described in Example 41 and a control were administered into peritoneal cavities four times at the same time of additional carbon tetrachloride administration (a total four times from the ninth to twelfth administration). After the additional administration, mice in the both groups were sacrificed. Liver tissue sections were prepared from the mice and immunohistochemically assessed. A rat anti-mouse fibroblast antibody (clone ER-TR7, 1:500 dilution; BMA) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, a peroxidase-labeled anti-rat IgG antibody (1:200 dilution; Biosource International, Inc.) was added as the secondary antibody, and the sections were incubated at room temperature for 30 minutes. After incubation, DAB substrate (Nichirei Biosciences) was added, and the samples were observed under a light microscope (Leica Microsystems). The antibody binding was detected by visualizing it as brown signals.

Examples of obtained images of immunostained liver tissues are shown in Fig. 61. The accumulation of fibroblasts was clearly observed in the control group. A bridge formation was shown in the histological pictures. In contrast, there was no accumulation of fibroblasts in any of the C6ST siRNA-administered groups.

5

[Example 62] Assessment of C6ST siRNA in the fibrogenesis score in a mouse hepatic fibrosis model

The degree of liver fibrogenesis in each sample was assessed using the fibrogenesis scores based on the immunohistochemical staining carried out in Example 61. The result is shown in Fig. 62. Each bar indicates mean $\pm$ standard deviation of the fibrogenesis score in each group. The fibrogenesis was statistically significantly reduced in the C6ST siRNA-administered group as compared to the control group ($p < 0.05$; t-test). This finding suggests that the administration of C4ST-1 siRNA, C4ST-2 siRNA, or C4ST-3 siRNA also produces a clinically beneficial effect of suppressing liver fibrogenesis.

15

[Example 63] Assessment of the anti-fibrogenic effect of C6ST siRNA in a mouse hepatic fibrosis model

RNA was extracted from liver tissues and quantitative PCR was carried out by the same method as described in Example 1. The result is shown in Fig. 63. The expression of α SMA, type I collagen, CTGF, and TGF β as fibrogenesis markers was enhanced in the control group. Meanwhile, the expression was significantly suppressed in the C6ST siRNA-administered group. This result suggests that fibrotic changes in liver tissues can be suppressed by inhibiting the expression of C6ST-1 and C6ST-2 genes.

25 [Cranial nerve tissue]

In these Examples, an MPTP-induced Parkinson's disease model was used as a fundamental mouse model for Parkinson's disease. This model is a classical but highly reproducible and simple model, and thus has been widely used as a Parkinson's disease model. The histological features are: infiltration of inflammatory cells into brain parenchymal tissue and reduction in the number of dopamine-producing neurons. Classically, neurofibrillary tangles were thought to be a cause of the reduction in the number of neurons in such pathological conditions.

Thus, such diseases include pathological conditions with neuronal disorders, specifically, not only representative neurodegenerative diseases such as Parkinson's disease, progressive supranuclear palsy, corticobasal degeneration, Alzheimer's disease, polyglutamine disease, amyotrophic lateral sclerosis (ALS), spinal progressive muscular atrophy, spinobulbar muscular

35

atrophy, Huntington's disease, and multiple sclerosis, but also other diseases such as multiple system atrophy (striatonigral degeneration, olivopontocerebellar atrophy, and Shy-Drager syndrome), adrenoleukodystrophy, Guillain-Barre syndrome, myasthenia gravis, Fisher syndrome, chronic inflammatory demyelinating polyneuropathy, Lewis-Sumner syndrome, Crow-Fukase syndrome, normal pressure hydrocephalus, syringomyelia, prion disease (Creutzfeldt-Jakob disease, Gerstmann-Straussler-Scheinker disease, and fatal familial insomnia), subacute sclerosing panencephalitis (SSPE), progressive multifocal leukoencephalopathy (PML), and spinocerebellar degeneration. Such diseases also include posttraumatic cerebral sequelae, sequelae of cerebrovascular disease (cerebral infarction and hemorrhage), sequelae of viral encephalitis, sequelae of bacterial meningitis, sequelae of spinal cord injury, and neurofibrillary tangle in the spinal nerve, peripheral nerve, auditory nerve, and optical nerve, etc. In particular, the above-listed sequelae have long been speculated as a basis for psychiatric disorders such as depression, and thus neurofibrillary tangle is important as a cause of psychiatric symptoms.

[Example 64] Assessment of the anti-fibrogenic effect of GalNAc4S-6ST siRNA treatment in a C57BL/6JcL mouse Parkinson's disease model induced by MPTP

In this Example, a mouse Parkinson's disease model was prepared by selectively degenerating dopamine neurons using 1-methyl-4-phenyl-1,2,3,6 tetrahydropyridine (MPTP) (Amende *et al.*, (2005) Journal of NeuroEngineering and Rehabilitation 2(20): 1-13). The mice were administered with GalNAc4S-6ST siRNA, and the gene expression and histological features after treatment was assessed.

Gestational day 14 C57BL/6JcL mice (CLEA Japan Inc.) were reared and allowed to deliver. 1 µg of the same GalNAc4S-6ST siRNA (GeneWorld) as described in Example 1 was combined with 1% atelocollagen (Koken Co.), which is a vehicle for siRNA, and the mixture was administered at 200 µl/head to the peritoneal cavities of C57BL/6JcL mice (female, eight weeks old; CLEA Japan Inc.). Two, three, and four days after administration, MPTP (Sigma Aldrich Japan), which selectively destroys dopamine neurons, was administered to the mice at 30 mg/kg (three times in total). The mice were reared, and on day 8 of the experiment 100 µl of 5 mg/ml BrdU (ZyMED Laboratory Inc.) was administered into the tail vein. After one hour, the mice were dissected and their brains were isolated to prepare samples for immunostaining and gene expression analysis.

The gene expression was assessed quantitatively by the same method as described in Example 1. The result is shown in Fig. 64. The expressions of GalNAc4S-6ST, and TGFβ, type I collagen, and αSMA, which are fibrogenesis markers, are enhanced in brain tissues in the Parkinson's disease model. Meanwhile, the expression was significantly suppressed by GalNAc4S-6ST siRNA administration. This demonstrates that fibrotic changes in brain tissues

can be suppressed by inhibiting the expression of GalNAc4S-6ST.

The agents of the present invention are thus useful, for example, as agents for suppressing fibrotic changes in brain tissues.

5 [Example 65] Assessment of GalNAc4S-6ST siRNA treatment for fibroblast accumulation in a C57BL/6JcL mouse Parkinson's disease model induced by MPTP

Brain tissue samples were treated by the same method as described in Example 3 to compare histological features after *in vivo* administration of GalNAc4S-6ST against fibrogenesis of neurons in the brain. The resulting sections were fixed with 4% PFA-phosphate buffer
10 (Nacalai Tesque) for ten minutes, and then washed with deionized water. An anti-fibroblast antibody (ER-TR7, 1:100 dilution; BMA) was added as the primary antibody, and the sections were incubated at 4°C overnight. Then, an Alexa488-labeled goat anti-rat IgG antibody (1:200 dilution; Invitrogen) was added as the secondary antibody, and the sections were incubated at room temperature for 30 minutes.

15 Images of tissues obtained by the method described above are shown in Fig. 65. The strong positive signals in the untreated group suggest intracranial infiltration of fibroblasts around the granular cortex in the splenium of posterior corpus callosum as compared to the control group. The positive signals of fibroblasts were drastically decreased in the GalNAc4S-6ST siRNA-treated group. The result described above demonstrates that the feature represented
20 by ER-TR7-positive signals in brain tissues induced in the mouse Parkinson's disease model was significantly suppressed by *in vivo* administration of GalNAc4S-6ST siRNA.

[Example 66] Assessment of the neuroprotective effect of GalNAc4S-6ST siRNA treatment in a C57BL/6JcL mouse Parkinson's disease model induced by MPTP

25 Next, to assess whether the above-described fibrogenesis was associated with the decrease of neurons, the expression of nerve regeneration-related genes in brain tissues was quantified by the same method as described in Example 64. The administration of GalNAc4S-6ST siRNA resulted in enhanced expression of GDNF, which is a factor that regulates the survival and differentiation of dopamine neurons as well as enhances the regeneration of the
30 neurons, and Nurr1, which is a factor for forming dopamine neurons (Fig. 66). This result suggests that the regeneration of dopamine neurons in brain tissues can be stimulated by inhibiting the expression of GalNAc4S-6ST.

The agents of the present invention are thus useful, for example, as agents for stimulating the regeneration of dopamine neurons in brain tissues.

35 [Example 67] Assessment of the neuroprotective effect of GalNAc4S-6ST siRNA treatment in a

C57BL/6JcL mouse Parkinson's disease model induced by MPTP

To finally verify the results described above in the Examples, histological features of the prepared tissue sample sections were weighed by staining dopamine neurons of the sections with an antibody against tyrosine hydroxylase, which is a marker for dopamine neuron. Tyrosine hydroxylase (TH) is an enzyme that converts the dopamine precursor into dopamine. The sections prepared by the same method as described in Example 64 were fixed with 4% PFA-phosphate buffer (Nacalai Tesque) for ten minutes, and then washed with deionized water. A rabbit anti-tyrosine hydroxylase polyclonal antibody (1:50 dilution; Calbiochem) was added as the primary antibody, and the sections were incubated at room temperature for one hour. An Alexa488-labeled donkey anti-rabbit antibody (1:200 dilution; Invitrogen) was added as the secondary antibody, and the sections were incubated at room temperature for 30 minutes.

Histological images for the control, untreated, and GalNAc4S-6ST siRNA-treated groups are shown in Fig. 67 (the original images are in color). Normal expression of tyrosine hydroxylase was confirmed in the superior colliculus of midbrain in the control group. Meanwhile, the signal for the expression was negative in the untreated group. This finding suggests that MPTP selectively destroyed dopamine neurons. On the other hand, a stronger signal was confirmed in the GalNAc4S-6ST siRNA-treated group as compared to the untreated group. In sum, it was concluded that functional recovery of dopamine neurons can be achieved *via* suppression of fibrotic changes by *in vivo* administration of GalNAc4S-6ST siRNA.

The agents of the present invention are thus useful, for example, as agents for recovering the function of dopamine neuron.

[Example 68] Assessment of neuroprotective effect of GalNAcST siRNA treatment in a C57BL/6JcL mouse Parkinson's disease model induced by MPTP

By the same procedure as described in Example 64, gestational day 14 of C57BL/6JcL mice (CLEA Japan Inc.) were reared and allowed to deliver. 1 µg of GalNAcST (a mixture of GalNAc4ST-1, GalNAc4ST-2, and GALNAC4S-6ST cocktail sequences) siRNAs (GeneWorld) was combined with 1% atelocollagen (Koken Co.), which is a vehicle for siRNA, and the resulting mixture was administered at 200 µl/head to the peritoneal cavities of C57BL/6JcL mice (female, eight weeks old; CLEA Japan Inc.). Two, three, and four days after administration, MPTP (Sigma Aldrich Japan), which selectively destroys dopamine neurons, was administered to the mice at 30 mg/kg (three times in total). The mice were reared, and on day 8 of the experiment 100 µl of 5 mg/ml BrdU (ZyMED Laboratory Inc.) was administered into the tail vein. After one hour, the mice were dissected and their brains were isolated to prepare samples for immunostaining and gene expression analysis.

The resulting sections were immunostained with an anti-TH antibody by the same

method as described in Example 67. The result is shown in Fig. 68. The result demonstrated that the reduction of TH-positive dopamine neurons was suppressed in the GalNAcST siRNA-treated group. Specifically, like Example 67, it was concluded that function recovery of dopamine neurons can be achieved *via* suppression of neurofibrillary tangle by inhibiting the expression of GalNAc4ST-1, GalNAc4ST-2, and GalNAc4S-6ST.

The agents of the present invention are thus useful, for example, as agents for suppressing neurofibrillary tangle.

[Example 69] Assessment of the effect of C4-sulfatase in a mouse type 2 diabetic retinopathy model induced by Streptozotocin: reduction of sulfated CSPG

Gestational day 14 C57BL/6JcL mice (CLEA Japan Inc.) were reared and allowed to deliver. 10 mg/ml Streptozocin (SIGMA) was subcutaneously administered at 20 µl/head to postnatal Day 2 female C57BL/6JcL mice. The mice were reared with sterile water and CE-2 Diet (CLEA Japan Inc.) until they were four weeks old, and then with sterile water and a High Fat Diet (CLEA Japan Inc.) for subsequent two weeks. In the second week, chondro-4-desulfating enzyme (C4-sulfatase) (20 units/ml; Seikagaku Co.) was administered at 4 units/head or medium (phosphate buffer) was administered into the peritoneal cavities twice a week four times (two weeks). On day 14, eye balls were isolated from mice of both groups and immunohistochemically assayed to assess the effect of C4-sulfatase.

Cryoblocks and sections were prepared from the isolated eye balls. The sections were fixed with acetone (Sigma Aldrich Japan) for ten minutes, and then washed with phosphate buffer. Then, an anti-chondroitin sulfate proteoglycan (CSPG) antibody (clone CS56, mouse monoclonal antibody, 10 µg/ml; Seikagaku Co.) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Next, the secondary antibody reaction was carried out using a Histofine Mouse Stain kit (Nichirei Biosciences; used for mouse monoclonal antibody), and color development was performed by adding DAB substrate (Nichirei Biosciences). The samples were observed under a light microscope (Leica Microsystems).

The obtained histological pictures are shown in Fig. 69 (the original images are in color). In the untreated group, CS56-positive signals are newly found in the vitreous side of the retina. In contrast, the intensity of CS56 is reduced in the enzyme-treated group. CS56 is an antibody that recognizes a sulfate group, and the reduction is suggested to reflect a decrease of sulfate groups at position 4. The result described above demonstrates that the induced deposition of CSPG in retinal tissues is suppressed *via* modification by *in vivo* administration of C4-sulfatase.

C4-sulfatase is an enzyme that desulfates GalNAc at position 4. Thus, it was demonstrated that tissue fibrogenesis at the biological level could be suppressed by inhibiting the sulfation of GalNAc at position 4. Specifically, desulfating enzymes for the sulfate group of

GalNAc at position 4 are useful as tissue fibrogenesis inhibitors.

[Example 70] Assessment of the effect of C4-sulfatase in a mouse type 2 diabetic retinopathy model induced by Streptozotocin: suppression of vascular endothelial cell proliferation

The sections prepared by the same method as described in Example 69 were fixed with acetone (Sigma Aldrich Japan) for ten minutes, and then washed with phosphate buffer. A rat anti-vascular endothelial cell antibody (CD31; 1:200 dilution; Phamingen) was added as the primary antibody, and the sections were incubated at room temperature for one hour. Then, a donkey peroxidase-labeled anti-rat IgG antibody (1:200 dilution; Biosource International, Inc.) was added as the secondary antibody, and the sections were incubated at room temperature for 30 minutes. After incubation, DAB substrate (Nichirei Biosciences) was added to the samples. The samples were observed under a light microscope (Leica Microsystems). The obtained histological pictures are shown in Fig. 70 (the original images are in color). The images show that the number of CD31-positive cells was increased in the vitreous side of the retina and some of them were protruded into the vitreum in the untreated group. This indicates that the model reflects the stage of preproliferative retinopathy in diabetic retinopathy and shows the effectiveness of this model. Meanwhile, the number of CD31-positive cells was significantly reduced in the corresponding region in the enzyme-treated group. In sum, it is suggested that the number of vascular endothelial cells is increased in the vitreous side of the retina in type 2 diabetes model, and that such proliferation of blood vessels can be suppressed by administering C4-sulfatase.

Specifically, desulfating enzymes for the sulfate group at position 4 of GalNAc are useful as an agent for suppressing proliferation of blood vessels.

[Example 71] Assessment of the effect of C4-sulfatase in a mouse type 2 diabetic retinopathy model induced by Streptozotocin: suppression of collagen-proliferative alteration

The sections prepared by the same method as described in Example 69 were fixed with acetone (Sigma Aldrich Japan) for ten minutes, and then washed with phosphate buffer. A rabbit anti-type IV collagen antibody (1:250 dilution; Sigma) was added as the primary antibody, and the sections were incubated at room temperature for one hour. A peroxidase-labeled anti-rabbit IgG antibody (1:200 dilution; Jackson ImmunoResearch) was added as the secondary antibody, and the sections were incubated at room temperature for 30 minutes. After incubation, DAB substrate (Nichirei) was added to the samples. The samples were observed under a light microscope (Leica Microsystems).

The obtained histological pictures are shown in Fig. 71 (the original images are in color). In the untreated group, type IV collagen-positive signals were increased in the vitreous side of

the retina and arranged parallel to the internal limiting membrane of the retina. This suggests morphological aberration of vein and collagen proliferation, i.e., fibrotic changes. In contrast, type IV collagen proliferation was markedly suppressed in the corresponding region in the enzyme-treated group. This demonstrated that retinal collagen proliferation is observed in the type 2 diabetes model but the collagen proliferation can be suppressed by administering C4-sulfatase.

Specifically, desulfating enzymes for the sulfate group at position 4 of GalNAc are useful as agents for treating type 2 diabetic retinopathy.

[Example 72] Localization of fibroblasts in the liver of type 2 diabetes model mouse

The anti-fibrogenic effect (using tissue infiltration of fibroblasts as an indicator) of C4-sulfatase on the liver was assessed in Examples 72 using the livers collected from the above-described type 2 diabetes model mice (Examples 69-71). Cryoblock preparation, immunostaining, and such were all achieved by the methods described above. As shown in Fig. 72, infiltration of many fibroblasts was observed in the untreated group. Meanwhile, the degree of fibroblast infiltration was reduced in the C4-sulfatase-treated group as compared to the untreated group. This suggests that C4-sulfatase has the pharmacological effect of suppressing fibroblast infiltration and this effect contributes to the anti-fibrogenic effect.

Specifically, desulfating enzymes for the sulfate group at position 4 of GalNAc are useful as agents for suppressing fibroblast infiltration.

[Example 73] Localization of macrophages in the liver of type 2 diabetes model mouse

The anti-inflammatory effect (using tissue infiltration of macrophages as an indicator) of C4-sulfatase on the liver was assessed using the livers collected from the above-described type 2 diabetes model mice (Examples 69-71). Cryoblock preparation, immunostaining, and such were all achieved by the methods described above. As shown in Fig. 73, infiltration of many macrophages accompanying spot formation was observed in the untreated group. Meanwhile, the degree of macrophage infiltration was reduced in the C4-sulfatase-treated group as compared to the untreated group. This result suggests that C4-sulfatase has the pharmacological effect of suppressing macrophage infiltration and this effect contributes to the anti-inflammatory effect.

Specifically, desulfating enzymes for the sulfate group at position 4 of GalNAc are useful as agents for suppressing macrophage infiltration or as anti-inflammatory agents.

[Example 74] Serum biochemical test findings on type 2 diabetes model mice

An additional analysis was carried out in Example 74 to supplement the results of

Examples 72 and 73. This Example assayed aspartate aminotransferase (AST) and alanine aminotransferase (ALT) as indicators for liver function, and triacylglycerol (TG) as an indicator for lipid metabolism, using sera collected from the above-described type 2 diabetes model mice (Example 69-71). The result is shown in Fig. 74. Biochemical tests were outsourced. AST, ALT, and TG values all tended to be increased in the untreated group (unt) as compared to the control group (nor). Meanwhile, the increases in the AST, ALT, and TG values tended to be suppressed in the C4-sulfatase-treated group (C4sul) as compared to the untreated group (unt). The result described above supports the results of Examples 72 and 73, suggesting that the liver function is preserved due to the anti-fibrogenic and anti-inflammatory effects of C4-sulfatase.

Specifically, desulfating enzymes for the sulfate group at position 4 of GalNAc are useful as agents for treating liver function disorders.

[Example 75] Comparison of CSPG expression in brain tissues in a C57BL/6JcL mouse model for Parkinson's disease induced by MPDP

The experiments in this Example was performed by preparing a model using MPDP, which is a metabolite of 1-methyl-4-phenyl-1,2,3,6 tetrahydropyridine (MPTP).

On Day 0 and 2, 100 μ l of 4 U/ml C4-sulfatase (Seikagaku Co.) was administered into the peritoneal cavity of each of C57BL/6JcL mice (female, eight weeks old; CLEA Japan Inc.). Alternatively, 200 μ l of a mixture of GalNAc4S-6ST siRNA (1 μ g; Hokkaido System Science Co.) and 1% atelocollagen was pre-administered on Day 0. On Day 2 to 4, MPDP (Sigma Aldrich Japan) was administered at 30 mg/kg every day.

The nucleotide sequence of mouse GalNAc4S-6ST siRNA agent used in this Example is shown below, but the sequences are not limited to these examples.

[mouse GalNAc4-6ST siRNAs] (Gene Bank accession number NM_029935)

(Hokkaido System Science, Co., Ltd.)

5'-gcagcccagcaagaugaauaagauc-ag-3' (SEQ ID NO: 91)

3'-ua-cgucgggucguucuacuauuucag-5' (SEQ ID NO: 92)

On day 8, 100 μ l of BrdU 5 μ g/mL (ZyMED Laboratory, Inc) was administered into mouse tail vein. One hour after administration, mice were dissected and their brains were isolated to prepare samples for immunostaining and gene expression analysis. 1 ml of RNA-Bee (TEL-TEST) was added to 50 mg each of excised organs (brains). The organs were crushed using an electrical homogenizer (DIGITAL HOMOGENIZER; AS ONE), then, 200 μ l of chloroform (Sigma-Aldrich Japan) was added to the resulting suspension. The mixture was gently mixed and then cooled on ice for about five minutes, and centrifuged in a centrifuge (Centrifuge 5417R; Eppendorf) at 12,000 rpm and 4°C for 15 minutes. After centrifugation, 500 μ l of the supernatant was transferred to a fresh eppendorf tube, and an equal volume of

isopropanol (500 μ l; Sigma-Aldrich Japan) was added thereto. The solution was mixed, and then 1 μ l of glycogen (Invitrogen) was added thereto. The mixture was cooled on ice for 15 minutes, and then centrifuged at 12,000 rpm and 4°C for 15 minutes. Next, RNA precipitate obtained after washing three times with 1,000 μ l of 75% ethanol (Sigma-Aldrich Japan) was air-dried for 30 minutes to one hour, and then dissolved in Otsuka distilled water (Otsuka Pharmaceutical Co., Ltd). The solution was 100 times diluted with Otsuka distilled water. The RNA concentrations of extracted samples in UV plates (Corning Costar) were determined using a plate reader (POWER Wave XS; BIO-TEK).

Next, an RT reaction (cDNA synthesis) was conducted by the following procedure. The concentrations of the obtained RNA samples were adjusted to 500 ng/20 μ l. The samples were heated at 68°C for three minutes in a BLOCK INCUBATOR (ASTEC), and cooled on ice for ten minutes. After cooling on ice, 80 μ l of RT PreMix solution (composition: 18.64 μ l of 25 mM MgCl₂ (Invitrogen), 20 μ l of 5x Buffer (Invitrogen), 6.6 μ l of 0.1 M DTT (Invitrogen), 10 μ l of 10 mM dNTP mix (Invitrogen), 2 μ l of RNase Inhibitor (Invitrogen), 1.2 μ l of MMLV Reverse Transcriptase (Invitrogen), 2 μ l of Random primer (Invitrogen), and 19.56 μ l of sterile distilled water (Otsuka distilled water; Otsuka Pharmaceutical Co., Ltd.)), which had been prepared in advance, was added to the samples. The mixtures were heated in a BLOCK INCUBATOR (ASTEC) at 42°C for one hour and at 99°C for five minutes, and then cooled on ice. 100 μ l of desired cDNAs were prepared and PCR reaction was carried out using the prepared cDNAs in the following composition.

2 μ l of PCR Buffer [composition: 166 mM (NH₄)₂SO₄ (Sigma Aldrich Japan), 670 mM Tris pH8.8 (Invitrogen), 67 mM MgCl₂·6H₂O (Sigma Aldrich Japan), 100 mM 2-mercaptoethanol (WAKO)], 0.8 μ l of 25 mM dNTP mix (Invitrogen), 0.6 μ l of DMSO (Sigma Aldrich Japan), 0.2 μ l of Primer Forward (GeneWorld), 0.2 μ l of Primer Reverse (GeneWorld), 15.7 μ l of Otsuka distilled water (Otsuka Pharmaceuticals, Inc.), 0.1 μ l of Taq polymerase (Perkin Elmer), and 1 μ l of cDNA obtained as described above were combined, and reacted using Authorized Thermal Cycler (eppendorf) at 30 cycles of 94°C for 45 seconds, 55°C for 45 seconds, 72°C for 60 seconds. After the reaction, the obtained PCR products were combined with 2 μ l of Loading Dye (Invitrogen). 1.5% agarose gel was prepared using UltraPure Agarose (Invitrogen), and the samples were electrophoresed in a Mupid-2 plus (ADVANCE) at 100 V for 20 minutes. After electrophoresis, the gel was shaken for 20 minutes in a stain solution prepared by 10,000 times diluting Ethidium Bromide (Invitrogen) with 1x LoTE (composition: 3 mM Tris-HCl (pH 7.5) (Invitrogen), 0.2 mM EDTA (pH 7.5) (Sigma Aldrich Japan)). The gel was photographed with EXILIM (CASIO) positioned on I-Scope WD (ADVANCE) and confirmed the gene expression. Cryoblock preparation, immunostaining, and such were all achieved by the methods described above.

The expression of CSPG in the MPDP-induced Parkinson's disease model was assessed by immunostaining using antibody against CS-56 (an anti-CSPG antibody, 1:100 dilution; Seikagaku Co.). The result is shown in Fig. 75. Strong CSPG-positive signals were observed in the untreated group, as shown in Fig. 75. Meanwhile, the positive signals were reduced in the C4-sulfatase-treated group and gene therapy group.

[Example 76] Localization of dopaminergic neurons in brain tissues in a C57BL/6JcL mouse model for Parkinson's disease induced by MPDP

To assess the pharmacological effects of C4-sulfatase and GalNAc4S-6ST siRNA on dopaminergic neurons, localization of dopaminergic neurons in the MPDP-induced Parkinson's disease model was analyzed by fluorescent immunostaining using an anti-tyrosine hydroxylase (TH) antibody (1:20 dilution) and an Alexa-488-labeled anti-rabbit IgG antibody (1:200 dilution; Invitrogen) for the secondary antibody. As shown in Fig. 76, the positive signals for dopaminergic neurons localized in brain tissues were weaker in the untreated group as compared to the other groups. Meanwhile, there was no great difference of the positive signal intensity in the C4-sulfatase-treated group and gene therapy group compared to the control group. The result suggests that the protective effect or regeneration/repair-promoting effect on dopaminergic neuron is produced by the administration of C4-sulfatase or the siRNA.

Specifically, inhibitors that act on GalNAc at position 4 or 6 (desulfating enzymes for the sulfate group at position 4 of GalNAc, and GalNAc4S-6ST siRNA) are useful as agents for protecting dopaminergic neurons or as agents for promoting the regeneration/repair of dopaminergic neurons.

[Example 77] Analysis of inflammation-related gene expression in brain tissues in a C57BL/6JcL mouse Parkinson's disease model induced by MPDP

To compare the anti-inflammatory effects of C4-sulfatase and GalNAc siRNA, total RNAs were extracted by the method described above from the same samples used to prepare tissue sections described in Examples 75 and 76, and the expression of TNF- α was analyzed by quantitative PCR. For quantitative PCR, SYBR Premix Kit (TAKARA BIO INC.) and Real-time PCR thermal cycler DICE (TAKARA BIO INC.) were used. Conditions of PCR reaction was: 95°C for 10 seconds, 40 cycles of 95°C for 5 seconds and 60°C for 30 seconds, finally, melting curve analysis was conducted. Nucleotide sequences of primers used in the quantitative PCR were described below.

mouse β actin (TAKARA BIO INC.)

Forward: 5'-CATCCGTAAAGACCTCTATGCCAAC -3' (SEQ ID NO: 93)

Reverse: 5'-ATGGAGCCACCGATCCACA -3' (SEQ ID NO: 94)

Tumor Necrosis Factor (TNF- α) (TAKARA BIO INC.)

Forward: 5'-CAGGAGGGAGAACAGAACTCCA-3 (SEQ ID NO: 95)

Reverse: 5'-CCTGGTTGGCTGCTTGCTT-3' (SEQ ID NO: 96)

As shown in Fig. 77, the expression of TNF- α was suppressed in the C4-sulfatase-treated group as compared to the untreated group. Meanwhile, the gene therapy group only showed a tendency of suppressing the expression. The result described above demonstrates that C4-sulfatase has the activity of suppressing inflammation.

Specifically, desulfating enzymes for the sulfate group at position 4 of GalNAc are useful as anti-inflammatory agents.

[Example 78] Analysis of inflammation-related genes for their expression in brain tissues in a C57BL/6JcL mouse Parkinson's disease model induced by MPDP

To complement the results described in Example 77, Examination was conducted using Nurr1, which is a gene involved in the generation of dopaminergic neurons, as a marker in this Example. The effects of C4-sulfatase and GalNAc4S-6ST were evaluated by the above-described quantitative PCR. The result is described below. The nucleotide sequences of primers used are as follows:

Nuclear receptor subfamily 4 Group A member 2 (Nurr1): (TAKARA BIO INC.)

Forward: 5'-CTGCCCTGGCTATGGTCACA-3' (SEQ ID NO: 97)

Reverse: 5'-AGACAGGTAGTTGGGTCGGTTCA-3' (SEQ ID NO: 98)

As shown in Fig. 78, the expression level of Nurr1 was significantly elevated in the C4-sulfatase-treated group and gene therapy group as compared to the untreated group ($P < 0.001$). This result supports the result described in Example 76 and shows the dopaminergic neuron-protecting effect or dopaminergic neuron regeneration/repair-promoting effect by administration of C4-sulfatase or the siRNA.

Specifically, inhibitors that act on GalNAc at position 4 or 6 (desulfating enzymes for the sulfate group at position 4 of GalNAc, and GalNAc4S-6ST siRNA) are useful as agents for protecting dopaminergic neurons or as agents for promoting the regeneration/repair of dopaminergic neurons.

Industrial Applicability

The present invention provides agents for suppressing fibrogenesis at the physiological tissue level through inhibiting the functions of sugar chain-related genes. The agents are useful for treating or preventing tissue fibrogenic disorders.

Chronic tissue fibrogenesis (fibrogenic tissue alterations) can develop in any organ in the body, and is a general term for a group of diseases that cause organ dysfunctions, leading to

death (review: Wynn TA, J. Clin. Invest. 117: 524-529, 2007).

Fibrogenic disorders are thought to be a terminal stage of chronic inflammation and can develop in any organ of the body. "Fibrogenic disorders" is a general term for a group of diseases that cause organ dysfunctions, leading to death. Fibrogenesis and the resulting dysfunctions are suspected to be the root of diseases with a high mortality rate, such as cardiovascular and cerebrovascular diseases. There is a view that 45% of deaths are caused by fibrogenesis in Western countries. Under this view, causes of individual death from disease are summarized into three groups: cancer, infections, and fibrogenesis. In addition to the conventional definitions of fibrogenic disorders such as cirrhosis, pulmonary fibrosis, and nephrosclerosis, a wide variety of diseases (mainly chronic diseases and excluding cancer and infections), can be defined as "fibrogenic disorders". Particularly, recent changes in lifestyle habits (commonly called "Westernization") have led to a rapid increase of new life-threatening diseases (disease concepts) such as non-alcoholic steatohepatitis (NASH) and chronic kidney disease (CKD). The finding that the diseases are caused by tissue fibrogenesis suggests the urgency of establishing therapeutic agents for "fibrogenic disorders". To date, however, only transplantation and artificial organs are available and there is no fundamental therapeutic method. This is a very urgent problem to be solved.

The present invention provides techniques for suppressing fibrogenic disorders (fibrogenic tissue lesions and resulting dysfunctions) based on a completely new method that targets the recently-identified sugar chain-related genes, functions of which were unknown.

Fibrogenic disorders that lead to clinically intractable dysfunctions are not only a deciding factor of death, but are also disorders that severely impair the daily quality of life (QOL). Thus, it is very important to establish tissue fibrogenesis-suppressing agents. However, no such agents are commercially available to date. Dedicated experimental studies are being conducted to assess TGF- β inhibitors, angiotensin inhibitors, inflammatory cytokine inhibitors, TLR inhibitors, MMP inhibitors, and the like. Still, however, their efficacy has not yet been established (review: Wynn TA, J. Clin. Invest. 117: 524-529, 2007).

The present inventors administered inhibitors that target sugar chain-related genes to subjects, which were model animals for fibrogenic disorders of various organs, in order to relieve fibrogenic lesions and to restore organ function.

The fibrogenic disorders were definitively confirmed based on histopathological features (including immunostaining for fibroblasts or collagen, and Masson staining). The therapeutic effect (fibrogenesis inhibitory effect) was determined based on clinical symptoms and the expression level of collagen in each organ, in addition to the histopathological features.

The therapeutic effect was proven by using as a technical method direct gene knockdown with nucleic acid pharmaceuticals (siRNA) against sugar chain-related genes.

CLAIMS

1. A tissue fibrogenesis suppressing-agent, which comprises as an ingredient an inhibitor of sulfation at position 4 or 6 of N-acetylgalactosamine.
5
2. The agent of claim 1, which has an effect of suppressing fibrogenesis of a physiological tissue.
3. The agent of claim 1 or 2, wherein the inhibitor has the activity of inhibiting the function of a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine.
10
4. The agent of claim 3, wherein the inhibitor is an siRNA that suppresses the expression of the sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine.
5. The agent of claim 1 or 2, wherein the inhibitor is a desulfating enzyme that desulfates at
15 position 4 or 6 of N-acetylgalactosamine.
6. The agent of any one of claims 1 to 5 for treating or preventing a fibrogenic disorder.
7. A method of screening for a tissue fibrogenesis suppressing-agent, which comprises the step
20 of selecting a compound that inhibits sulfation at position 4 or 6 of N-acetylgalactosamine that constitutes a sugar chain.
8. A method of screening for a tissue fibrogenesis suppressing-agent, which comprises the steps of:
25 (a) contacting a test compound with N-acetylgalactosamine or a sugar chain comprising N-acetylgalactosamine;
(b) determining the degree of sulfation at position 4 or 6 of N-acetylgalactosamine; and
(c) selecting a compound that reduces the degree of sulfation as compared to when the test compound is not contacted.
30
9. A method of screening for a tissue fibrogenesis suppressing-agent, which comprises the steps of:
(a) contacting a test compound with a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine;
35 (b) determining the sulfotransferase activity of the enzyme; and

(c) selecting a compound that reduces the activity as compared to when the test compound is not contacted.

10. A method of screening for a tissue fibrogenesis suppressing-agent, which comprises the steps of:

(a) contacting a test compound with a cell expressing a gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine;

(b) determining the expression level of the gene in the cell; and

(c) selecting a compound that reduces the expression level of the gene as compared to when the test compound is not contacted.

11. A method of screening for a tissue fibrogenesis suppressing-agent, which comprises the steps of:

(a) contacting a test compound with a cell or cell extract containing a DNA wherein a reporter gene is operably linked to the transcriptional regulatory region of the gene encoding a sulfotransferase that transfers a sulfate to position 4 or 6 of N-acetylgalactosamine;

(b) determining the expression level of the reporter gene; and

(c) selecting a compound that reduces the expression level of the reporter gene as compared to when the test compound is not contacted.

12. A method of producing a pharmaceutical composition for treating or preventing a fibrogenic disorder, which comprises the steps of:

(a) selecting a tissue fibrogenesis suppressing-agent from test compounds by the method of any one of claims 7 to 11; and

(b) combining the agent with a pharmaceutically acceptable carrier.

SEQUENCE LISTING

<110> STELIC INSTITUTE OF REGENERATIVE MEDICINE
STELIC INSTITUTE & CO.

<120> Glycochain related genes, and use thereof

<130> S10-A0703P

<150> JP 2007-336518

<151> 2007-12-27

<160> 98

<170> PatentIn version 3.3

<210> 1

<211> 9120

<212> DNA

<213> Mus musculus

<400> 1

```
tgtatatctg agtatctgtg tatgtgcata tatgtgtgtc tatgtgtatg tgtggggttg      60
tgcattgtgtg tctgtgtgta catgtctgtg tgtctctgtg tgtgtctgtg tatgtgtgtt    120
tacattgtgtg tgtgtgtttc tgtgtggacg catgtgtatg tgtctccgtg tgcctgcatg      180
tgtgtttgtg tatgtgtgta ggcatgagta catcatggtg tacattgtaga aattgagaga      240
gactcttggg tgtggttctc atctcccacc atatctgaag taaatgtctt attcacacca      300
ctgtgcacat cagaaagcta gccttgacct ccaaggaact ctgtctctgt ctctgtctt      360
attgtaggta cctgggggtt ataagcctat gctgctgcat atggcttcac ttgggttctg      420
gggatttgaa cttagggtcct cacagtttta gggcaagcat ttatttact gaacaatctc      480
ccatgatccc tagagtttat ttagaggcaac ttaaccttta gtcgtcattg agatttagca      540
ttgatttcag ttccagttgt attctgatgt tgagcgctac agcagaggta cccagggact      600
gctgattttc tgtggccatg aatcctctag tgattaataa atttcatcat aatgttctgt      660
ttatttctaa atgtccagac aactctaatt tcaaattaat ctcatcctaa acagcacagg      720
ataaaagtta cagattgtcc tatcttcatt catgtgatgc ctcaaatac agtgttaatc      780
cttttagtcag ttatttgtga aacggaacct aattacacac actcgcatgc gcgtgcgcgc      840
acacacacac acacacacac acacacacac acacacacac acttgtagat ttgccacac      900
agctgcttaa gcaatttgga gcaaatgtag aaatattatg aaatttcaca taaaaatggg      960
ttttcctctg tcttcttgag aaatgtgtga gtttcaatat gacagtaatc ttctggtgtt    1020
ggactataac cacttgaacc cctgaggtag taagtgtgta cctactacgt gcctggattc     1080
catcatttcc ctacttatat aagcttctcc atcctgtaga tcagcactgt gtaatggagt     1140
aacacaagtc attgaatcat gcaatcattt aatcaattta attggcactt ctgataagag     1200
aagtgggcac atatttagtt gtgtttgcag cctgtgctat gtcatttttc tctatatata     1260
ctttaagatt tttttctttt ttccatttac gggggggggg ttattccatt taatttacat     1320
gcttaatgac atataatttg catatcaggt gttcctttct ttagagttaa cttcttttag     1380
gaccattgtt tataatggat gtatatcatc cctaaattaa atcagccatt gtctacttaa     1440
cacaccagtt tctctgatgt ccctctggct cttcctcctc tgaggttcta gttataggaa     1500
```

agtctggcat	tttgatgttt	ttccatggct	tcgtgggctc	tgatcatcca	cttacagtgt	1560
actttcatct	cttgtctaca	aactgcatat	catctattga	tctgcgttca	aggctactac	1620
attttcatgc	aattccactt	tgctgttaaa	ccttttctat	gatccgtttt	tatacttata	1680
aacaatccaa	gtgttgact	ctttgggtat	aaatgttctg	tatttcacag	gtcttaactt	1740
cttttcagac	tatacaaaga	aagattttct	atttttttta	tttgttgtat	gcattttatg	1800
catttagcac	tgggattttcc	ataacaggca	gttaagatca	ttgaatacat	actttgtgtg	1860
gtaaatataa	tacggcttat	ttgattctcc	cccttaaaat	gctcccatct	tctcagtcac	1920
gtcaggttgc	tatggagtcc	accctgagca	ccatgagtgt	gatgctgtac	agacactcct	1980
gatatgcctt	ctgttatttg	aactccacct	gcaagttttc	attcaagcat	agctcctaata	2040
cccatcagc	gtgcactttt	cttaattgat	ttgctggtag	agagctatat	gtaagcatgg	2100
cttagcatca	aattgaactt	tctcctctcc	ctagcttctt	tccagtatat	tcttcctcac	2160
cttcatcaga	ttttgaggtc	ttgagatcta	aaccttcatg	ttacaatttg	cagagataacc	2220
aggatctcta	tcagacttga	tgtgtctcca	tgtaattcta	gtatcagcct	ttcataagat	2280
aattgcaaaa	tatagaacct	cccctctctc	cttttgacta	ttacagcttt	gtttctctcc	2340
caagttaata	cagatgtatg	caaatcaact	aagaaaagca	aacactgaat	gggataagga	2400
gctatactca	aaagaaagct	tgtagttgga	gttcaaataa	cacaagtgga	accaagagta	2460
tctgcacagc	atcagactca	ctgtgctcag	gatggaatcc	atagcaacct	cccataactg	2520
agaggatgtg	agcccccttt	aaagggaaaa	tcaaataagc	agtattatat	taaccacatg	2580
atgtgtatat	tcaatgtaga	ttgggctgca	ccgaagttat	gatttttagta	atgagtattt	2640
ctccttggg	tttcttctgg	accaccctgg	ttttcctttt	aaaaatagta	aaatagcagc	2700
aatgactgct	tctcttaaaa	tctgaacaac	acagagcact	tcagcatcct	ttcaacctat	2760
ctaaatccat	tgtagacctg	cactgacttg	ctgtgagcta	ttgtgtcaca	actattattg	2820
tagctaaaca	ccttgaattt	tcaaagctgc	tctggactgt	gtcttctttt	agaattaaac	2880
ttgctaaaga	tattggactc	tgagatgctt	attcaccact	ttctataaaa	gagtaaggat	2940
gacaatgctt	tgagtagaag	aagattgcat	acgagggtag	cagcatgtca	ttatgtcccc	3000
agcagtcagt	aaagtaagga	tccagtgatg	gtagagttat	tatgtccatt	atggaactaa	3060
tgtaaattgt	tatcaatgac	tagaatggct	gtaatcacca	atccttagac	aagtgtgagt	3120
aacatcatgc	tcttttctta	taataataga	tagtcacaca	caccatgggtc	ttaaactatta	3180
cattttgatt	tttctctttg	tgtctgtctt	ttcgtacctc	tactccactt	acctgagcct	3240
tcagctaggg	ctcctccaag	acggcagtca	cagtactagt	gaggataatt	ctcatatgga	3300
gacctgaagc	agaataacct	tcttccaagg	aaacaaaaat	cctagtcaac	atcttctcca	3360
ctctttgggt	atttattttt	atttctattt	ctattttgcc	atttaagaat	ttttaaactc	3420
ttttctgttt	tcttatatat	attctttttt	ttacaacttt	tgatattctt	tatcaaatga	3480
tttatcccat	tgcttttctt	cctgcagcta	tctgtgagtt	ttagcatgga	tatttagaat	3540
acaattggaa	attatgaagg	tcgaataaaa	gttctctaag	agccaggacc	tcactagcct	3600

agttagttgc	ttagtttcta	atataaggca	tggtttatct	cctactgtat	ggcccttaaa	3660
tccactagac	atctgttggc	ttccaacagc	atacaattgc	tgctattgta	tgtttaggta	3720
tatatTTTTA	tgaaacacta	tattctgatt	ggtgttcatt	cttcctatga	aggggaagat	3780
tttattgcct	gctcaaattc	aagtcagtc	tttttgagta	gacaaagggt	gtccaatcct	3840
gctaaaccct	tttctTTTTT	gcttggggaa	tttcttttagg	gaaaattctc	aaccatgaac	3900
caagcctaga	gattgataat	agctgaaggc	agagtagcca	gtcttagtaa	gggatgagag	3960
aaagactgac	caaagactga	ataggaaatg	gttgatagca	ttcattggga	ctagacaata	4020
aagttagcta	ccccaacatg	ggcaaattga	gaaaatgggg	agagtcatat	ggatatgtag	4080
ggaggagaag	aaaggaaggg	gtgagtttga	agtatggaag	agcctttaaa	gcatttgtaa	4140
gatttgtaag	aaaaagcttt	tgagagacag	agtattacct	gccttcaa	cccttcaatt	4200
actgtcccta	caagacatac	ctgtgctcag	ctgctcttgg	accacaagg	ggcctaagag	4260
catgtttgtg	tccctgtctt	tatacaggct	tctcatctgt	atttctcaa	cgtaaataca	4320
aatgggggag	ataacaaatg	atatttctaa	gtggtacaca	ggtatgagaa	ctgcacacaa	4380
tctatttata	tgtataatgg	ccagagctct	cagcagggaa	cagaggaagc	ctaggtaact	4440
cctgctccag	agcatgtctg	tatggagacc	tcagttcagc	aactatgaaa	tctgtgttca	4500
aagggtcatc	actgggaaca	gttcttgga	tagcagacat	attcattctc	tctctctctc	4560
tctctctctc	tctctctctc	tctctctctc	tctctctctc	tctctgtctg	tctctgtctc	4620
tctctgtctg	tctgtctgtc	tctcccttct	gtcaggggta	tgcatatcatg	ggctgggaat	4680
tctatcttcc	ctgcctgctg	tctttgggtg	ctccctaattg	tggtgccatg	tttacttaac	4740
ctgtctgaaa	tggaggataa	cctctgggtct	gataactgtc	aagttactga	aatgtcaggg	4800
aattgtgaat	gaaccctctc	gaatgctggg	cagatgtttt	gagcacgtgg	gtatccatag	4860
attttttttt	cagcaggatt	tccttcttta	atataattgtg	ggaacgaaac	attgaagttt	4920
ccaacactaa	gaaaagcaat	gctctggggt	tgcaaagaag	tctccaaagc	tgtgggagag	4980
gcaatggctt	tgccctacaa	aggtgatcct	gatttggtg	gggtgttcct	cagtctccct	5040
catcctttgt	ttgtggacca	aacctctgat	caaataatca	gtatacagac	aagtactgac	5100
tgactcctgt	cactgtttcc	catgtaaatt	ctcaaccatg	tgtgtcagaa	ttgtttgagt	5160
catcaaagtt	ccgtgatgct	caggatgggtg	aataccagaa	actgaagtct	gaaatccaat	5220
ttggcctttc	tcgggcactg	taaactgagt	gcaaatgtgc	ccagttccct	cgtagggccat	5280
ttatagcaca	ataacacagt	gcttgcttac	aaaatctagt	cccgattaaa	gtattttattt	5340
aaagttgctt	tagacaaata	tatttctaaa	tggaaggatg	aaacagaaag	aatattccaa	5400
ctgtacagtc	tatatTTTTG	ttctaaaaat	ttttattaga	attctgactt	ttagaaaaca	5460
aatagatatt	caaaaactac	ttctttaatt	aaataataac	atcattatag	acattctcca	5520
gccatttaga	agtagcatga	ttacaaagca	ttcatgttcc	ctttgttttt	tgaacagaa	5580
gtattttatat	gacaaataaa	aagcaaaaat	ctcatatatg	tgatagctat	agatgcattg	5640
ataggaatta	aatttaaatca	aaaagaactt	ttaatggcaa	taatgtggct	actaacttca	5700
ttgtttgtaa	taatgctgca	aatgtttaat	gaaaactgtt	actaacaata	caggagaaga	5760

ttacagatat	gattaggtcg	tgttgataga	aatggaaaga	gaaagggtcac	tcactacctc	5820
agccacagaa	cttgaggaat	gtgtgagggg	ttttgaggaa	aaaagagaac	aatgtaacat	5880
cacagcaaca	tcttaatctg	aaagtagacg	acaaagggtt	cagagtaggg	gatcagaaat	5940
ttaagtatgc	agagaagcaa	ggcatatacg	gcttagacaa	tgtatgaaga	acagcagcag	6000
acagcaagtg	aaagaaatca	cagtccacat	ggaaatcaca	cagaccacag	tgtgcacaca	6060
gagacagaaa	tcagcaaggg	acacacggct	tccagttatg	aaagttagac	agcaaaattt	6120
aagttgtccc	acctaaaaca	tgatgtatcg	tctctcaa	ggggaagggg	aaggagcagc	6180
aaacatctgg	atatgtctaa	aacctcacag	agaacatctg	ggatctttgc	atttctccct	6240
ctgactttct	caatatgtta	gctacttcta	gatagtttca	ataatgggtct	ctaattgact	6300
atggagagac	aggtctcatg	tatgtgaaca	tccttaacca	ggagatggaa	tgtgtttcct	6360
tgtcactcat	tttctaagga	agtgaattcc	ttctgcattg	cccaggattg	gatgttacgt	6420
gaaactaaaa	atcattgatt	aaaaagaaat	gtcattacca	cagttggggt	gaatcaatca	6480
agacttatcc	tagttacagg	catcacctcc	ccttaagcac	aaatactgag	aacagctaag	6540
ctcactttga	aatctgggat	gaggaggctt	taggtcaagc	atctgacaat	gtgtgtgtgt	6600
atgtgtgtgt	gtgtgtgtgt	gtgtagtatg	attatatgta	tatatgtaga	tgtgtgtata	6660
tatttatatg	tatatatgta	tatgtgagta	tgtatatatg	tatatgtgtg	tatgtgtatg	6720
agtgtgtgta	tgtatgacta	tgagtgtgta	tgtattgtgc	aagtgtgtgt	atatatgtgt	6780
gtatatgttt	atgtatatgt	gtattgtatg	ggtgtgaatg	tattatatat	gtgtgtaggt	6840
acatatgtgt	atatgtttat	gcgtgtatat	gtgtatgcat	tatatgtatg	tgtgtatgtg	6900
tgtatgtttt	catgtgtatg	tagcatatgt	atatgcatgt	gtgtgtctgt	gtgtgtaagc	6960
atgtattttac	taagaaaaaa	tcagtttctc	tagctaata	gtcattgaag	tgaagaaacc	7020
ttacgctatc	acattactat	gtttccagca	taaattgcaa	tatcagtaag	acattcccca	7080
tcataactta	atggacactt	agattaggtc	atgcagcagt	aaattatgaa	agcgtctatg	7140
ctgagctgag	gtgggaaatc	gtctttcctg	cctccttctc	cctgttcttt	ctcctaaagg	7200
ccctcatcct	ctttctcccc	ttcctttctt	ttaaaatatt	ctatcattga	tccaagctaa	7260
ctaacccttc	ccttgctcac	atgtgacctg	cccagaggaa	tctatttata	gtcaacaata	7320
acaggccaag	gacatgcaaa	caaagtactt	tacttagccc	cacttgggtga	atctattggg	7380
tgaataggta	aggggttatt	tataggagct	tgactgactg	gcacagctgt	atctctgaaa	7440
agtcccccaa	agcatcatgg	gtgggaacta	aggaaagccg	gaaatgctgg	agctggaatc	7500
ccaacctgca	gtggctacat	gatgaatgtg	tctcctttcc	tactaaatat	ttaagactct	7560
gggcaggctg	gttagtcttg	taaaagtctt	gtgggtttct	ggagtcctgg	gaggctcact	7620
gtccctgcag	aagcttggtt	ttattgtata	tgaaagaaag	tgctcaacac	tgagtcacaa	7680
aagaagcccc	cgagaatcgg	tggcagaggt	ggaaatagaa	gaatccgatg	cagttccccga	7740
ggcaagagtt	gatcctctcc	atggtgacaa	gaaaacaaga	gccacagaag	cctgaaggct	7800
aagactatga	cctgtgaggt	tctcctctcc	taggacttga	atgcacttaa	ctctgagggc	7860

tagcagggca	gagattcagg	agcgttctca	gaggttgtcc	cttctccttg	tatgattcta	7920
ggagaagctc	agtgcatagg	cagaagttgc	aaactctaac	acattaaccc	acatcctgaa	7980
ttgttttata	tcaattgttt	tataactaca	tgttttaact	tttgttttgg	tttttgaatt	8040
ttgtttgttt	gtttgttttc	tatttttcat	atgtgtgcat	gtgagccaga	agacaggcta	8100
agggactctg	ttctctcttt	ccagcatgtg	ggtttcaggg	atcaaagtta	ggccccagg	8160
cttcaactggg	caacgggacc	cttagcctct	gagtcactct	cttaactctt	ttctttgaaa	8220
ctttttcatt	acattgtgtc	agtgcacaca	cacacacaca	cacacatgca	tatacatcca	8280
catgcatata	catccacatg	ccatatgtca	tggcacatgt	gtggaaagaa	atgggtgcct	8340
tgaggaaatc	agttttcttc	ttccatcacg	tggatcttat	gacttcaggt	aacccatctc	8400
tgctcccca	gcatggctac	aagcatttac	ctgccaaagtc	atctccctgg	ctccacatt	8460
aatttataat	aatacaccct	ttcagcctag	agagatgact	cagtggtta	gaactctggc	8520
tgtgcttcag	aaggttctga	gttcaattcc	cagcagctat	aagggtggctc	acaactatct	8580
ctaattgtgt	tttctgtcct	cttctggcat	gcacttgtat	gtgcagatag	agtacttata	8640
tgcataaata	aataaata	cattaataat	cttttaaaaa	gaaggaaaga	ttaaaagaac	8700
aatacaccct	ttcttttaag	taagaaaaca	tgcttagtat	atagaataaa	tgctggtggt	8760
gaaaggtgaa	attcagaaga	aattataggt	tggaaatatt	tgaactcttt	tgtgtcaaac	8820
caaagtgtaa	agatgatgtg	ataataaggt	aagggttttct	tcctacacat	atacttgttt	8880
aataaccagt	ataagactat	gagtgaacat	aactaaaaat	acatgaaaaa	aaactatttt	8940
aaaaaaacat	cactttcaga	tgactctggt	gaaactcact	gataaccagc	aatctctttg	9000
atagaccaga	gtgtccagtc	tgaagttaat	tcttttgctt	tattctcaaa	caatgttgca	9060
gaagtatgaa	ccatagtatc	aagatacact	gtccacatat	gcacctggcc	tgacttgtgt	9120

<210> 2
 <211> 2105
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (418)..(1671)

<400> 2	
gagccttccc	ggcgcgtgag ccggatccgg tggcaccgcg gggaagagac aggaccgggc 60
gggtggcggca	gagacagggg gacgcacccg gtgcagaaga tccaatagga gcacgccgcc 120
gcaacctctc	ccgcgcgctc cggtcgccga ctctacgccg atcgccact ccccgcacct 180
tggactacac	cgggaaaaag acgcagcctg gcccgaaactg gggcggagat cttcttgct 240
tccaagagc	tcgggaggga agatttctcc gccgcgaga acccgccgg actggaggaa 300
cccgtggcct	ggagagcgct ggctgtgcca gacccagcc tgatggatgt ctggtgtgga 360
taatgaggga	agaacgtgcc ttttacaccc aagaggtgac cccggagcgt gccccgg 417
atg acc cca caa ctc gga acg atg cgg cta gcc tgc atg ttc tcg tcc	465
Met Thr Pro Gln Leu Gly Thr Met Arg Leu Ala Cys Met Phe Ser Ser	
1 5 10 15	

atc Ile	ctg Leu	ctg Leu	ttt Phe 20	gga Gly	gct Ala	gcg Ala	ggc Gly	ctg Leu 25	ctc Leu	ctc Leu	ttc Phe	atc Ile	agc Ser 30	ctc Leu	cag Gln	513
gac Asp	cct Pro	ata Ile 35	gag Glu	ctc Leu	agc Ser	ccc Pro	cag Gln 40	caa Gln	gtt Val	cca Pro	ggg Gly	ata Ile 45	aag Lys	ttc Phe	agc Ser	561
atc Ile	agg Arg 50	ccc Pro	cag Gln	caa Gln	ccc Pro	cag Gln 55	cat His	gat Asp	agc Ser	cac His	ttg Leu 60	agg Arg	ata Ile	tcc Ser	aca Thr	609
gaa Glu 65	aag Lys	ggc Gly	aca Thr	cga Arg	gat Asp 70	tca Ser	ccc Pro	agc Ser	ggg Gly	tcg Ser 75	cca Pro	aga Arg	ggc Gly	ctc Leu	cag Gln 80	657
ctg Leu	caa Gln	gcg Ala	cct Pro	gac Asp 85	caa Gln	cct Pro	cga Arg	cct Pro	cac His 90	ccg Pro	aag Lys	gca Ala	gcg Ala	gga Gly 95	tct Ser	705
cct Pro	ttg Leu	cgc Arg	ctc Leu 100	cgg Arg	cag Gln	cgc Arg	agg Arg	cgg Arg 105	aga Arg	ctg Leu	ctc Leu	atc Ile	aaa Lys 110	aag Lys	atg Met	753
cca Pro	gcc Ala	gca Ala 115	ggg Gly	act Thr	aac Asn	caa Gln	ggc Gly 120	aac Asn	aac Asn	tcg Ser	tcc Ser	gaa Glu 125	acc Thr	ttt Phe	atc Ile	801
cag Gln 130	ccg Pro	aga Arg	ccc Pro	cgc Arg	acc Thr	atg Met 135	gac Asp	agt Ser	cgt Arg	tgg Trp	gtc Val 140	agc Ser	ctg Leu	cac His	cag Gln	849
acc Thr 145	caa Gln	cag Gln	gag Glu	cgc Arg	aag Lys 150	cgt Arg	gtg Val	atg Met	cga Arg	gaa Glu 155	gcc Ala	tgc Cys	gct Ala	aaa Lys	tac Tyr 160	897
agg Arg	gcc Ala	agc Ser	agc Ser	agc Ser 165	cgc Arg	aga Arg	gct Ala	gtc Val	act Thr 170	ccc Pro	cgc Arg	cac His	gtc Val	tcc Ser 175	cgc Arg	945
atc Ile	ttc Phe	gtg Val 180	gag Glu	gac Asp	cgc Arg	cac His	cgt Arg	gta Val 185	ctg Leu	tac Tyr	tgt Cys	gaa Glu	gta Val 190	ccc Pro	aag Lys	993
gca Ala	ggc Gly	tgc Cys 195	tcc Ser	aac Asn	tgg Trp	aag Lys	agg Arg 200	gtg Val	ctc Leu	atg Met	gtg Val	ctg Leu 205	gca Ala	ggg Gly	tta Leu	1041
gcc Ala 210	tca Ser	tcc Ser	acg Thr	gca Ala	gat Asp	atc Ile 215	caa Gln	cac His	aac Asn	acc Thr	gtc Val 220	cac His	tat Tyr	ggc Gly	agc Ser	1089
gcc Ala 225	ctt Leu	aag Lys	cgc Arg	ctg Leu	gat Asp 230	act Thr	ttt Phe	gac Asp	cgg Arg	cag Gln 235	ggc Gly	ata Ile	gtg Val	cac His	cgc Arg 240	1137
ctc Leu	agt Ser	acc Thr	tac Tyr	acc Thr 245	aag Lys	atg Met	ctc Leu	ttt Phe	gtc Val 250	cgg Arg	gaa Glu	ccc Pro	ttt Phe 255	gag Glu 255	cgg Arg	1185
ctg Leu	gtc Val	tct Ser	gct Ala 260	ttc Phe	cga Arg	gac Asp	aag Lys	ttt Phe 265	gag Glu	cat His	cct Pro	aac Asn	agc Ser 270	tac Tyr	tat Tyr	1233
cat His	cct Pro	gtc Val 275	ttt Phe	ggc Gly	aag Lys	gct Ala	atc Ile 280	ctg Leu	gcc Ala	cgg Arg	tac Tyr	cgc Arg 285	gcc Ala	aac Asn	gcc Ala	1281
tcg Ser	cgg Arg	gag Glu	gca Ala	ctg Leu	cgg Arg	act Thr	ggc Gly	tcc Ser	ggg Gly	gtg Val	cag Gln	ttc Phe	ccc Pro	gag Glu	ttc Phe	1329

290	295	300	
gtc cag tac ctg ttg gat gtc cac cgg ccc gtg ggc atg gac atc cac Val Gln Tyr Leu Leu Asp Val His Arg Pro Val Gly Met Asp Ile His 305 310 315 320			1377
tgg gac cat gtt agc cgg ctg tgc agc ccc tgc ctc atc gac tat gac Trp Asp His Val Ser Arg Leu Cys Ser Pro Cys Leu Ile Asp Tyr Asp 325 330 335			1425
ttt gtg ggc aag ttc gag agc atg gaa gac gat gcc aac ttc ttc ctg Phe Val Gly Lys Phe Glu Ser Met Glu Asp Asp Ala Asn Phe Phe Leu 340 345 350			1473
cgt ctc atc cat gcg ccc ggg aac ctg act ttc ccg agg ttc aag gac Arg Leu Ile His Ala Pro Gly Asn Leu Thr Phe Pro Arg Phe Lys Asp 355 360 365			1521
agg cac tcc gag gag gcg cgg acc aca tcg aga atc acc cat cag tac Arg His Ser Glu Glu Ala Arg Thr Thr Ser Arg Ile Thr His Gln Tyr 370 375 380			1569
ttc gct cag ctc tcc tcg ctg cag cga cag cga acc tac gac ttc tac Phe Ala Gln Leu Ser Ser Leu Gln Arg Gln Arg Thr Tyr Asp Phe Tyr 385 390 395 400			1617
tac atg gat tac ctg atg ttc aac tac tcc aaa cct ttc tcg gac ctg Tyr Met Asp Tyr Leu Met Phe Asn Tyr Ser Lys Pro Phe Ser Asp Leu 405 410 415			1665
tac tga gggcggggcc tgctgggtcag gggcgggggtc tgccgggtcat gcccactcac Tyr			1721
ctgcgcatag gcggcctccg gggactgagc tctgaggatg tgaggccttg tggtgtggc			1781
cctaggggtgg gccacagagg ccagacaat ggaccttgac cttgtccca caccatttc			1841
ctcattgggt tggctgagtt tgagacggag cagcactcgg atggatgctt taagaactca			1901
gctgagctat gctgagctat gctgtccagg gaagcctgag acccagaaga gggccccagc			1961
gtcaggggat gtcctacatc cccttatcct ttgccttgta ccaaaccacg tggtttgctg			2021
cttttctatg acccaggggtc atctgaataa agcacatggt tttcagagca aaaaaaaaaa			2081
aaaaaaaaa aaaaaaaaaa aaaa			2105
 <210> 3 <211> 417 <212> PRT <213> Mus musculus <400> 3			
Met Thr Pro Gln Leu Gly Thr Met Arg Leu Ala Cys Met Phe Ser Ser 1 5 10 15			
Ile Leu Leu Phe Gly Ala Ala Gly Leu Leu Leu Phe Ile Ser Leu Gln 20 25 30			
Asp Pro Ile Glu Leu Ser Pro Gln Gln Val Pro Gly Ile Lys Phe Ser 35 40 45			
Ile Arg Pro Gln Gln Pro Gln His Asp Ser His Leu Arg Ile Ser Thr 50 55 60			

Glu Lys Gly Thr Arg Asp Ser Pro Ser Gly Ser Pro Arg Gly Leu Gln
 65 70 75 80
 Leu Gln Ala Pro Asp Gln Pro Arg Pro His Pro Lys Ala Ala Gly Ser
 85 90 95
 Pro Leu Arg Leu Arg Gln Arg Arg Arg Leu Leu Ile Lys Lys Met
 100 105 110
 Pro Ala Ala Gly Thr Asn Gln Gly Asn Asn Ser Ser Glu Thr Phe Ile
 115 120 125
 Gln Pro Arg Pro Arg Thr Met Asp Ser Arg Trp Val Ser Leu His Gln
 130 135 140
 Thr Gln Gln Glu Arg Lys Arg Val Met Arg Glu Ala Cys Ala Lys Tyr
 145 150 155 160
 Arg Ala Ser Ser Ser Arg Arg Ala Val Thr Pro Arg His Val Ser Arg
 165 170 175
 Ile Phe Val Glu Asp Arg His Arg Val Leu Tyr Cys Glu Val Pro Lys
 180 185 190
 Ala Gly Cys Ser Asn Trp Lys Arg Val Leu Met Val Leu Ala Gly Leu
 195 200 205
 Ala Ser Ser Thr Ala Asp Ile Gln His Asn Thr Val His Tyr Gly Ser
 210 215 220
 Ala Leu Lys Arg Leu Asp Thr Phe Asp Arg Gln Gly Ile Val His Arg
 225 230 235 240
 Leu Ser Thr Tyr Thr Lys Met Leu Phe Val Arg Glu Pro Phe Glu Arg
 245 250 255
 Leu Val Ser Ala Phe Arg Asp Lys Phe Glu His Pro Asn Ser Tyr Tyr
 260 265 270
 His Pro Val Phe Gly Lys Ala Ile Leu Ala Arg Tyr Arg Ala Asn Ala
 275 280 285
 Ser Arg Glu Ala Leu Arg Thr Gly Ser Gly Val Gln Phe Pro Glu Phe
 290 295 300
 Val Gln Tyr Leu Leu Asp Val His Arg Pro Val Gly Met Asp Ile His
 305 310 315 320
 Trp Asp His Val Ser Arg Leu Cys Ser Pro Cys Leu Ile Asp Tyr Asp
 325 330 335
 Phe Val Gly Lys Phe Glu Ser Met Glu Asp Asp Ala Asn Phe Phe Leu
 340 345 350

Arg Leu Ile His Ala Pro Gly Asn Leu Thr Phe Pro Arg Phe Lys Asp
 355 360 365

Arg His Ser Glu Glu Ala Arg Thr Thr Ser Arg Ile Thr His Gln Tyr
 370 375 380

Phe Ala Gln Leu Ser Ser Leu Gln Arg Gln Arg Thr Tyr Asp Phe Tyr
 385 390 395 400

Tyr Met Asp Tyr Leu Met Phe Asn Tyr Ser Lys Pro Phe Ser Asp Leu
 405 410 415

Tyr

<210> 4
 <211> 1369
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (40)..(1281)

<400> 4
 attcagacat gaagagagac acagtggctt gaagtggctc atg aaa gcc aaa caa 54
 Met Lys Ala Lys Gln
 1 5
 gtc ttc ttt tct gtc ctg ctg ttt ggg aca gca ggg ctt ctg ctc ttc 102
 Val Phe Phe Ser Val Leu Leu Phe Gly Thr Ala Gly Leu Leu Leu Phe
 10 15 20
 atg tac ttg caa gca tgg att gaa gaa cat cat aca ggg aaa ata gag 150
 Met Tyr Leu Gln Ala Trp Ile Glu Glu His His Thr Gly Lys Ile Glu
 25 30 35
 aag aaa agg gat cag aaa gga gta tcg gtg act acg gga aaa atc cag 198
 Lys Lys Arg Asp Gln Lys Gly Val Ser Val Thr Thr Gly Lys Ile Gln
 40 45 50
 aaa cag atc acg aat cag aac tct gag gtt cac atg cct gaa gat ctg 246
 Lys Gln Ile Thr Asn Gln Asn Ser Glu Val His Met Pro Glu Asp Leu
 55 60 65
 aag aag aaa ggg gga gat ctg ctc aac cta ggg agt cca aca agg gtt 294
 Lys Lys Lys Gly Gly Asp Leu Leu Asn Leu Gly Ser Pro Thr Arg Val
 70 75 80 85
 tta agg aag atc agc cat tca caa agg gag aac gga gct tac aga tca 342
 Leu Arg Lys Ile Ser His Ser Gln Arg Glu Asn Gly Ala Tyr Arg Ser
 90 95 100
 act gaa gca cat caa gga gct aaa att gaa gtt ttt cag aaa ccc atc 390
 Thr Glu Ala His Gln Gly Ala Lys Ile Glu Val Phe Gln Lys Pro Ile
 105 110 115
 cag atg gac tgg cca ctg gtc act cag ccc tta aac aaa agt ttg gtc 438
 Gln Met Asp Trp Pro Leu Val Thr Gln Pro Leu Asn Lys Ser Leu Val
 120 125 130
 caa ggc aac aaa tgg aag aaa gca gat gca acc caa gag aag cgt cgg 486
 Gln Gly Asn Lys Trp Lys Lys Ala Asp Ala Thr Gln Glu Lys Arg Arg

135	140	145	
tca ttc ctt cat gag ttt tgc aag aaa tat ggt aga gta aat gat ccc Ser Phe Leu His Glu Phe Cys Lys Lys Tyr Gly Arg Val Asn Asp Pro 150 155 160 165			534
aag ttc aac ctt ttt cat ata gta tct agg ata tat gta gaa gac aaa Lys Phe Asn Leu Phe His Ile Val Ser Arg Ile Tyr Val Glu Asp Lys 170 175 180			582
cac aaa atc ctg tac tgt gaa gta cca aaa gct ggc tgc tct aat tgg His Lys Ile Leu Tyr Cys Glu Val Pro Lys Ala Gly Cys Ser Asn Trp 185 190 195			630
aaa aga att ctg atg gtc cta aat gga ttg gct tcc tct gca tac aat Lys Arg Ile Leu Met Val Asn Gly Leu Ala Ser Ser Ala Tyr Asn 200 205 210			678
atc tcc cat gat act gtg cac tat ggg aag cat ctg aaa aca ctg gat Ile Ser His Asp Thr Val His Tyr Gly Lys His Leu Lys Thr Leu Asp 215 220 225			726
agt ttt gac tta aaa gga gta cac atg cgt ttg aat aca tat acc aaa Ser Phe Asp Leu Lys Gly Val His Met Arg Leu Asn Thr Tyr Thr Lys 230 235 240 245			774
gct gtg ttt gtt aga gat ccc atg gaa aga tta gtc tcc gca ttt agg Ala Val Phe Val Arg Asp Pro Met Glu Arg Leu Val Ser Ala Phe Arg 250 255 260			822
gat aaa ttt gag cat ccc aat agt tac tac cat ccg gtg ttt gga aag Asp Lys Phe Glu His Pro Asn Ser Tyr Tyr His Pro Val Phe Gly Lys 265 270 275			870
gca att atc aag aaa tat cga cca aat gcc tct gca gaa gca tta aat Ala Ile Ile Lys Lys Tyr Arg Pro Asn Ala Ser Ala Glu Ala Leu Asn 280 285 290			918
aat gga tct gga gtc aaa ttc aaa gaa ttc gcc tac tat ttg ctg gat Asn Gly Ser Gly Val Lys Phe Lys Glu Phe Ala Tyr Tyr Leu Leu Asp 295 300 305			966
gct cac cgt cca gta gga atg gat att cac tgg gaa aga gtc agc aaa Ala His Arg Pro Val Gly Met Asp Ile His Trp Glu Arg Val Ser Lys 310 315 320 325			1014
ctg tgt tat ccg tgt ttg atc aac tat gac ttt gta ggg aag ttt gag Leu Cys Tyr Pro Cys Leu Ile Asn Tyr Asp Phe Val Gly Lys Phe Glu 330 335 340			1062
acc tta gga gag gat gcc aat tac ttt cta cag ttg att ggt gct cca Thr Leu Gly Glu Asp Ala Asn Tyr Phe Leu Gln Leu Ile Gly Ala Pro 345 350 355			1110
aaa gag ttg aca ttt cca aac ttt aag gat agg cac tcc tct gat gaa Lys Glu Leu Thr Phe Pro Asn Phe Lys Asp Arg His Ser Ser Asp Glu 360 365 370			1158
aga acc aat gcc cac gtg gta agg cag tat tta aag gac ctg agc aca Arg Thr Asn Ala His Val Val Arg Gln Tyr Leu Lys Asp Leu Ser Thr 375 380 385			1206
gcc gaa aga cag ctc atc tat gac ttc tat cac ttg gac tat ttg atg Ala Glu Arg Gln Leu Ile Tyr Asp Phe Tyr His Leu Asp Tyr Leu Met 390 395 400 405			1254
ttt aat tac aca act cca cat ttg taa tttgcattca tttttctaaa Phe Asn Tyr Thr Thr Pro His Leu 410			1301
gccctacata gatttaaatga tgatggcctc aaataagcta ctgtaattgt cctacaattc			1361

tctgtatg

1369

<210> 5

<211> 413

<212> PRT

<213> Mus musculus

<400> 5

Met Lys Ala Lys Gln Val Phe Phe Ser Val Leu Leu Phe Gly Thr Ala
1 5 10 15

Gly Leu Leu Leu Phe Met Tyr Leu Gln Ala Trp Ile Glu Glu His His
20 25 30

Thr Gly Lys Ile Glu Lys Lys Arg Asp Gln Lys Gly Val Ser Val Thr
35 40 45

Thr Gly Lys Ile Gln Lys Gln Ile Thr Asn Gln Asn Ser Glu Val His
50 55 60

Met Pro Glu Asp Leu Lys Lys Lys Gly Gly Asp Leu Leu Asn Leu Gly
65 70 75 80

Ser Pro Thr Arg Val Leu Arg Lys Ile Ser His Ser Gln Arg Glu Asn
85 90 95

Gly Ala Tyr Arg Ser Thr Glu Ala His Gln Gly Ala Lys Ile Glu Val
100 105 110

Phe Gln Lys Pro Ile Gln Met Asp Trp Pro Leu Val Thr Gln Pro Leu
115 120 125

Asn Lys Ser Leu Val Gln Gly Asn Lys Trp Lys Lys Ala Asp Ala Thr
130 135 140

Gln Glu Lys Arg Arg Ser Phe Leu His Glu Phe Cys Lys Lys Tyr Gly
145 150 155 160

Arg Val Asn Asp Pro Lys Phe Asn Leu Phe His Ile Val Ser Arg Ile
165 170 175

Tyr Val Glu Asp Lys His Lys Ile Leu Tyr Cys Glu Val Pro Lys Ala
180 185 190

Gly Cys Ser Asn Trp Lys Arg Ile Leu Met Val Leu Asn Gly Leu Ala
195 200 205

Ser Ser Ala Tyr Asn Ile Ser His Asp Thr Val His Tyr Gly Lys His
210 215 220

Leu Lys Thr Leu Asp Ser Phe Asp Leu Lys Gly Val His Met Arg Leu
225 230 235 240

tgc Cys 15	cgg Arg	atg Met	gtg Val	ctg Leu	gcc Ala 20	act Thr	tgc Cys	ttc Phe	gga Gly 25	tcc Ser	ttt Phe	atc Ile	ttg Leu	gtc Val	atc Ile 30	458
ttc Phe	tat Tyr	ttc Phe	caa Gln	agt Ser 35	atg Met	ttg Leu	cac His	cca Pro	gtc Val 40	atg Met	cgg Arg	agg Arg	aac Asn	ccc Pro 45	ttc Phe	506
ggt Gly	gtg Val	gac Asp	atc Ile 50	tgc Cys	tgc Cys	cgg Arg	aag Lys	gga Gly 55	tcg Ser	aga Arg	agt Ser	ccc Pro	ctg Leu 60	cag Gln	gag Glu	554
ctc Leu	tac Tyr	aat Asn 65	ccc Pro	atc Ile	cag Gln	ctg Leu	gag Glu 70	cta Leu	tcc Ser	aac Asn	act Thr	gcc Ala 75	atc Ile	ctg Leu	cac His	602
cag Gln 80	atg Met	aga Arg	cgg Arg	gac Asp	cag Gln	gtg Val 85	aca Thr	gac Asp	acc Thr	tgc Cys	cgg Arg	gcc Ala	aac Asn	agt Ser	gcc Ala	650
atg Met 95	agc Ser	cgc Arg	aag Lys	cgc Arg	agg Arg 100	gtg Val	ctg Leu	acc Thr	ccc Pro	aac Asn 105	gac Asp	ctg Leu	aag Lys	cac His	ctg Leu 110	698
gtg Val	gtg Val	gat Asp	gag Glu	gac Asp 115	cac His	gaa Glu	ctc Leu	atc Ile	tac Tyr 120	tgc Cys	tat Tyr	gtg Val	ccc Pro	aag Lys 125	gta Val	746
gcg Ala	tgc Cys	acc Thr	aac Asn 130	tgg Trp	aag Lys	agg Arg	ctc Leu	atg Met 135	atg Met	gtc Val	ctg Leu	agt Ser	ggc Gly 140	cgg Arg	ggc Gly	794
aag Lys	tac Tyr	agc Ser 145	gat Asp	ccc Pro	atg Met	gag Glu	atc Ile 150	cca Pro	gcc Ala	aac Asn	gaa Glu	gcc Ala 155	cac His	gtg Val	tcg Ser	842
gcc Ala 160	aac Asn	ctg Leu	aag Lys	acc Thr	ctt Leu	aac Asn 165	cag Gln	tac Tyr	agc Ser	atc Ile	cca Pro 170	gag Glu	atc Ile	aac Asn	cac His	890
cgc Arg 175	ttg Leu	aaa Lys	agc Ser	tac Tyr	atg Met 180	aag Lys	ttc Phe	ctg Leu	ttc Phe 185	gtg Val	cgg Arg	gaa Glu	ccc Pro	ttc Phe	gag Glu 190	938
agg Arg	ctg Leu	gtg Val	tct Ser	gcc Ala 195	tac Tyr	cgc Arg	aac Asn	aag Lys	ttc Phe 200	acg Thr	cag Gln	aag Lys	tac Tyr	aac Asn 205	acc Thr	986
tcc Ser	ttc Phe	cac His	aag Lys 210	cgc Arg	tac Tyr	ggc Gly	acc Thr	aag Lys 215	atc Ile	atc Ile	cga Arg	cgc Arg	cag Gln 220	cgg Arg	aag Lys	1034
aac Asn	gcc Ala	acg Thr 225	cag Gln	gag Glu	gcc Ala	ctg Leu	cgc Arg 230	aag Lys	ggg Gly	gac Asp	gat Asp	gtc Val 235	aag Lys	ttc Phe	gag Glu	1082
gag Glu 240	ttc Phe	gtg Val	gcc Ala	tac Tyr	ctc Leu	atc Ile 245	gac Asp	ccc Pro	cac His	acc Thr	cag Gln 250	cgg Arg	gag Glu	gag Glu	ccc Pro	1130
ttc Phe 255	aac Asn	gag Glu	cac His	tgg Trp	cag Gln 260	acg Thr	gtc Val	tac Tyr	tct Ser	ctc Leu 265	tgc Cys	cac His	ccg Pro	tgc Cys	cac His 270	1178
atc Ile	cac His	tac Tyr	gac Asp	ctc Leu 275	gtg Val	ggc Gly	aag Lys	tat Tyr	gag Glu 280	aca Thr	ctg Leu	gag Glu	gag Glu	gac Asp 285	tcc Ser	1226
aat Asn	tac Tyr	gta Val	ctg Leu 290	cag Gln	ctg Leu	gcc Ala	gga Gly	gtg Val 295	agc Ser	ggc Gly	tac Tyr	ctg Leu	aag Lys 300	ttc Phe	ccc Pro	1274

acc tat gca aag tcc acc cga act acc gac gag atg acc acg gag ttc	1322
Thr Tyr Ala Lys Ser Thr Arg Thr Thr Asp Glu Met Thr Thr Glu Phe	
305 310 315	
ttc cag aac atc agc gcc gag cac cag aca cag ctg tac gaa gtc tac	1370
Phe Gln Asn Ile Ser Ala Glu His Gln Thr Gln Leu Tyr Glu Val Tyr	
320 325 330	
aaa ctg gac ttt tta atg ttc aac tac tca gtg cca aac tac ctg aag	1418
Lys Leu Asp Phe Leu Met Phe Asn Tyr Ser Val Pro Asn Tyr Leu Lys	
335 340 345 350	
ttg gat tag agggctgcgg aggggagggg gaggggtggt tgggggagag	1467
Leu Asp	
ggagagaatc ctgcttttta atttaagatt tttatttgtc aaaagaattc tatggagact	1527
gggttatattt gtaagttaat atgtcttcgg gggagatgct gcgagcggca tggttaagaa	1587
ttatttaaaa attctccacg gggaaggaca gctgtctttg caggggagcg ggggtgaata	1647
tccctgtttt tagaagtgga tactgcaaca ctgtctccaa ggtgtccttg tgttctggtg	1707
aagtccacaa actgcattcc ataaagtcta atgaatctta tttatagtta tttaaacgtg	1767
gtctgtggca gcagcctgct cctgtctgtg cagaggagag tacagtctgt gcttcgctgg	1827
ctatggctca ttgggggagg ggaggggggtt atctcccca ctgaccgtgt gtagaatccc	1887
atggtgacag ctgcgggatg tgtgcctgtg gccattcaag aaaccctgct agaaagaagt	1947
caattgtcct tgcactaaga aacagggatg agactgggtc aaaccgattg ctagcaaccc	2007
caaagtccct tgtagtaaag aaatgcaa atttttaacc caagaagaga gcagatgata	2067
gcgatttctc tttaccgaaa ttagcaggct ataagaatag gctttctgtg gaaatcttta	2127
agtggcactt tccagtgcct gcgaaggcag aataatgggt gttttctagt caataagtat	2187
tgagcatcta ctattctcca ggcttgactg cagatagcac attgagtga accaggagcc	2247
ttggccttca gcaagcttta tgtgaacaca gggcatccag ctcaacagtg aaccaacacc	2307
cccatccctt tggctctcac acagacttta gactattctc aagtgggtcca gaggggtccag	2367
gaccctagca aatttgcgta tttctacagt tcaaatttga aagccacaca tttggatcag	2427
gccacttaac tagctaagta gttcattgcc caacacaaga aatgacagaa acccttttga	2487
tcatgtatat ctagtgcagt gtcttcatag tagaagccta taagccacaa tattctccaa	2547
gcgcaaggat ggaggactgc tgtgattgat tagtgatgtc tgccttgact aaggaaaggg	2607
aggaacattt ctcaagggcc tcctactgtg atctccagtc cacatcttgt cctgcctgcc	2667
acttaaaaca gtgcactgca tacatttgga gactggcgga gttcaagtgt gaaggtcaaa	2727
ggtggagtgt tggtaaccac aggcacatgc ccaccctcca tgtcctgcat tggcagagca	2787
tccaggacct cctctctctc cagaagggtg gaaatttgct ctttctgtt ctttgctgat	2847
caatgccccaa acctattcct ctttctagt tgtttctgat actttgcggt tgacatttga	2907
gccacattcc catgacagcc agtgtccggg tacgtagcag atgctgtaca tagttctgat	2967
gataagggag ctggtaggct gttaaatttg cttttgtgt ttttggtggg ttttgtttgt	3027
tttgatgcag tatagagtga agggggtaag aatattctga aaaacataaa attgagtaat	3087

ttattcacag	aaactattaa	aatggtattg	ttaggctcac	aaggaggctg	ggggcagcgg	3147
taactgctac	actgtctcgc	agaggaaatt	attttattta	atgtgagaga	aatgtgagcc	3207
agagtggctg	cctgaggcta	tttataaaac	gcaggtagct	tccccaattt	ctctccccctc	3267
gcctttaata	aatagattta	tgtcatgtgt	gtgaccaaga	acagtgcacg	atcttggtg	3327
tccctggaat	agagaggcag	tcatggtgct	caggccctgg	gcgtggtgta	gctgatgcat	3387
gtcccaccga	gcccaggctc	tccaaggcat	cagtttgtgg	tcacatatcc	ttagaaaaca	3447
aaattaagtc	aggttcctgt	ctgcaggagt	tttcagagcc	ttgaatacat	agtctgcctt	3507
gtgaaactgg	gagaatgtag	catttctggg	ctgcatgacc	taggtatcct	gtatggcaga	3567
ccatgtctta	gctggttttt	cacacagtag	ccttcagtga	gggtgatcta	gcctgtctga	3627
ggctctgcaa	gattgacagc	tgcttcctag	atccttcac	aaatcagcag	aaacagtgtc	3687
catctgggaa	ctttatgaga	gtgttgcttt	acctagttag	gggtgtctca	tctggatact	3747
gccccagaga	gtgatcccg	ccctccaagc	acagcggagg	gaagcctagg	gtgttatgga	3807
gtggggcagg	tggcttaagg	acaccttaaa	ggccacttgg	tggggataat	ggcctcgtgg	3867
ggatactcct	cacttccacc	cacagttagg	aactgttcac	tgtgtgcttc	aggagtttta	3927
aatggagatt	cgtgttgatt	tccatggtcc	gcatctgttt	gtcttccgtc	tgtggtccag	3987
tgtgtgtgc	agatggtact	tactcccttt	ctctctctta	caggagaaa	gactgtcctt	4047
ctggggagag	tgtatcaata	cgcttcacga	atgaggaggt	ctttctcctg	tctgtgttc	4107
acacttccta	gtgacaattt	tcaaaatatg	tatcggttg	actgtttatt	gtagagtgtg	4167
gaaaggtttc	tgtgttcctg	gctaaagaca	tccattggtg	aaatgtgctt	ggaaatccaa	4227
gtgttttcta	ctgaggaaaa	aaaaatgttg	gaaaagcttt	ctcctgcatg	cctctgtgtc	4287
ttagctagaa	aggggagctg	ccagcgtccc	agctctgagc	cactgttcaa	aggtgcagct	4347
gtgttttagg	actaggtaca	ggcagagtga	taaggacatg	cctgcttagt	gtactcat	4407
atcgagaaag	tttagactat	gagccacagg	ggtccctgcc	tgcagtcctt	atagcctccc	4467
cctgccccag	tgcttttctc	agctggtctg	ggaatcaggg	cagcccattg	aggggggctc	4527
agacaaccct	ttcttcctgg	ttctgggagt	aaggctttct	ctgcagccac	aaggagaaca	4587
tgggaaccag	aaacttctct	ttggtcagcc	cttctgagca	cacaggttta	gcccagctc	4647
caagaatgct	cctgagcatt	gtataagccc	ctaggactct	tagctcatgt	gataatgatt	4707
gcattactat	cctgctttta	ctgacaaaac	cacagagaat	tttgcttgct	ttcctactct	4767
cttctctg	ccaggcctcc	tccatacgta	tacatacaca	catgggcatg	tatacacaca	4827
catgcacaat	agacatgcac	acacacatgc	acacatgggt	atacatatac	acacatgcac	4887
aatagacatg	catatataca	tatacatgca	cacattgata	ggcatacaca	catgcacagt	4947
agacatgcac	acatgggtat	acatagacac	acatgcacag	acatgcatac	acacacatgc	5007
acacataaat	gtgcacacac	acatacacia	attgcccag	cctgtgcact	cctggtaccc	5067
atgtcccatc	ctggcccacc	tctttgtcag	caggactgac	tccatcactc	cctcaggttt	5127
tgcccaccag	cacccccag	ctttcccca	gcagagctaa	acagcccctc	tgaatagaca	5187
acaggacctt	ctagaaacag	ctgaaccctt	ggacagcagc	aggcagattt	tgatctgctc	5247

tcatacaggta caaaagggtgg tatccgtatt ctcttctgag catgctcagt aagcatatgg 5307
 acatagaatt taacatctct gtggagtgtg tgttttttac atatttgtat gcagtcgagg 5367
 agggcctgtt gtagaattct ctccctgtat cttactatac tgttaaagaa gctgaattct 5427
 atgttgccaa cagatgctg aaatgttcct ccaggaaaag ccattcaagc ctgattatct 5487
 ttctaagtaa cttcaattaa attgaagaag aaaaaaaaaa aaaaa 5532

<210> 7
 <211> 352
 <212> PRT
 <213> Mus musculus

<400> 7

Met Lys Pro Ala Leu Leu Glu Val Met Arg Met Asn Arg Ile Cys Arg
 1 5 10 15

Met Val Leu Ala Thr Cys Phe Gly Ser Phe Ile Leu Val Ile Phe Tyr
 20 25 30

Phe Gln Ser Met Leu His Pro Val Met Arg Arg Asn Pro Phe Gly Val
 35 40 45

Asp Ile Cys Cys Arg Lys Gly Ser Arg Ser Pro Leu Gln Glu Leu Tyr
 50 55 60

Asn Pro Ile Gln Leu Glu Leu Ser Asn Thr Ala Ile Leu His Gln Met
 65 70 75 80

Arg Arg Asp Gln Val Thr Asp Thr Cys Arg Ala Asn Ser Ala Met Ser
 85 90 95

Arg Lys Arg Arg Val Leu Thr Pro Asn Asp Leu Lys His Leu Val Val
 100 105 110

Asp Glu Asp His Glu Leu Ile Tyr Cys Tyr Val Pro Lys Val Ala Cys
 115 120 125

Thr Asn Trp Lys Arg Leu Met Met Val Leu Ser Gly Arg Gly Lys Tyr
 130 135 140

Ser Asp Pro Met Glu Ile Pro Ala Asn Glu Ala His Val Ser Ala Asn
 145 150 155 160

Leu Lys Thr Leu Asn Gln Tyr Ser Ile Pro Glu Ile Asn His Arg Leu
 165 170 175

Lys Ser Tyr Met Lys Phe Leu Phe Val Arg Glu Pro Phe Glu Arg Leu
 180 185 190

Val Ser Ala Tyr Arg Asn Lys Phe Thr Gln Lys Tyr Asn Thr Ser Phe
 195 200 205

His Lys Arg Tyr Gly Thr Lys Ile Ile Arg Arg Gln Arg Lys Asn Ala
 210 215 220
 Thr Gln Glu Ala Leu Arg Lys Gly Asp Asp Val Lys Phe Glu Glu Phe
 225 230 235 240
 Val Ala Tyr Leu Ile Asp Pro His Thr Gln Arg Glu Glu Pro Phe Asn
 245 250 255
 Glu His Trp Gln Thr Val Tyr Ser Leu Cys His Pro Cys His Ile His
 260 265 270
 Tyr Asp Leu Val Gly Lys Tyr Glu Thr Leu Glu Glu Asp Ser Asn Tyr
 275 280 285
 Val Leu Gln Leu Ala Gly Val Ser Gly Tyr Leu Lys Phe Pro Thr Tyr
 290 295 300
 Ala Lys Ser Thr Arg Thr Thr Asp Glu Met Thr Thr Glu Phe Phe Gln
 305 310 315 320
 Asn Ile Ser Ala Glu His Gln Thr Gln Leu Tyr Glu Val Tyr Lys Leu
 325 330 335
 Asp Phe Leu Met Phe Asn Tyr Ser Val Pro Asn Tyr Leu Lys Leu Asp
 340 345 350

<210> 8
 <211> 1785
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (149)..(1408)

<400> 8
 ggcgatttcg gctgcagaat cagcatcacc agcaacagca gcagcggcgg tgactgtggc 60
 gggcgctagg tccgtctcct agggaccatg tcccagctgt gcacaaggct gaagtgaagg 120
 gccaggagtg ggcccagccc agggcagc atg acc aag ccg cgg ctc ttc cgg 172
 Met Thr Lys Pro Arg Leu Phe Arg
 1 5
 ctg tgg ctg gta cta ggg tcg gct ctc atg atc ctt ttg atc att gta 220
 Leu Trp Leu Val Leu Gly Ser Ala Leu Met Ile Leu Leu Ile Ile Val
 10 15 20
 tat tgg gac aac gtg gga acc gcc cac ttc tat ctg cac acg tct ctc 268
 Tyr Trp Asp Asn Val Gly Thr Ala His Phe Tyr Leu His Thr Ser Leu
 25 30 35 40
 tcc agg cca cac atc cta gaa ccc ctt ccc acc cag gga ttg gtg gag 316
 Ser Arg Pro His Ile Leu Glu Pro Leu Pro Thr Gln Gly Leu Val Glu
 45 50 55
 gag aac gtg ttc aca tct gac gtg gat gag ttt ttg gat act ctc ctt 364
 Glu Asn Val Phe Thr Ser Asp Val Asp Glu Phe Leu Asp Thr Leu Leu
 60 65 70

agt tct gac gcg aag cac aac gac ctt tcc agg aga aaa act gag cag Ser Ser Asp Ala Lys His Asn Asp Leu Ser Arg Arg Lys Thr Glu Gln 75 80 85	412
ccc ccg gcg ccc gcc ccc agc aag cca gtc ttg agc cac atg gag gag Pro Pro Ala Pro Ala Pro Ser Lys Pro Val Leu Ser His Met Glu Glu 90 95 100	460
aac gtg aga ggc tac gac tgg tcc act cat gat gcc cat cag aac cct Asn Val Arg Gly Tyr Asp Trp Ser Thr His Asp Ala His Gln Asn Pro 105 110 115 120	508
gac cgg gac agg cag cag gcc gag agg agg agc ctg ctg aga gac ttc Asp Arg Asp Arg Gln Gln Ala Glu Arg Arg Ser Leu Leu Arg Asp Phe 125 130 135	556
tgt gcc aac gcc agc ctg gca ttc ccc acc aag gac cgc tct ttt gac Cys Ala Asn Ala Ser Leu Ala Phe Pro Thr Lys Asp Arg Ser Phe Asp 140 145 150	604
gac atc ccc aac tac gaa ctg aac cac ctg atc gtg gac gac cgc cac Asp Ile Pro Asn Tyr Glu Leu Asn His Leu Ile Val Asp Asp Arg His 155 160 165	652
ggg gtc atc tac tgc tac gtg ccc aag gtg gcc tgc acc aac tgg aag Gly Val Ile Tyr Cys Tyr Val Pro Lys Val Ala Cys Thr Asn Trp Lys 170 175 180	700
cga gtg atg atc gtg ctg agc gag agc ctg ctg gac cgg ggc agc ccc Arg Val Met Ile Val Leu Ser Glu Ser Leu Leu Asp Arg Gly Ser Pro 185 190 200	748
tac cga gac ccc ctg gac atc ccc cgg gaa cac gtg cac aac acc agc Tyr Arg Asp Pro Leu Asp Ile Pro Arg Glu His Val His Asn Thr Ser 205 210 215	796
acg cac ctc acc ttc aac aag ttc tgg cgc cgc tac gga aag ttc tcc Thr His Leu Thr Phe Asn Lys Phe Trp Arg Arg Tyr Gly Lys Phe Ser 220 225 230	844
cgt cac ctc atg aag gtg aag ctg aag aag tac acc aag ttc ctg ttc Arg His Leu Met Lys Val Lys Leu Lys Lys Tyr Thr Lys Phe Leu Phe 235 240 245	892
gtg cgc gac ccc ttt gtg cgc ctc atc tca gcc ttc cgc agc aag ttc Val Arg Asp Pro Phe Val Arg Leu Ile Ser Ala Phe Arg Ser Lys Phe 250 255 260	940
gag ctg gag aac gaa gag ttt tac cgc aag ttc gcg gtg ccc atg ctc Glu Leu Glu Asn Glu Glu Phe Tyr Arg Lys Phe Ala Val Pro Met Leu 265 270 275 280	988
cga ctg tac gcc aac cac acc agc ctg ccc gcc tcg gtg agt gag gct Arg Leu Tyr Ala Asn His Thr Ser Leu Pro Ala Ser Val Ser Glu Ala 285 290 295	1036
ttc agc gcc ggg ctc aag gtc tcc ttc gcc aac ttc atc cag tac ctc Phe Ser Ala Gly Leu Lys Val Ser Phe Ala Asn Phe Ile Gln Tyr Leu 300 305 310	1084
cta gac cca cac acg gag aag ctg gcg cct ttc aac gag cac tgg cga Leu Asp Pro His Thr Glu Lys Leu Ala Pro Phe Asn Glu His Trp Arg 315 320 325	1132
cag gtg tac cgc ctc tgc cac ccg tgc cag ata gac tat gac ttc gtg Gln Val Tyr Arg Leu Cys His Pro Cys Gln Ile Asp Tyr Asp Phe Val 330 335 340	1180
ggg aag ctg gag acg ctc gat gag gac gct gcc cag ctc ctg agg ttc Gly Lys Leu Glu Thr Asp Glu Asp Ala Ala Gln Leu Leu Arg Phe 345 350 355 360	1228

ctc aag gta gac tcc cag ctc cac ttc ccc ccc agt tat cgg aac agg 1276
 Leu Lys Val Asp Ser Gln Leu His Phe Pro Pro Ser Tyr Arg Asn Arg 375
 365
 acg gcc agc agc tgg gag gaa gac tgg ttt gcc aac atc ccc ctg gca 1324
 Thr Ala Ser Ser Trp Glu Glu Asp Trp Phe Ala Asn Ile Pro Leu Ala 390
 380
 tgg agg caa cag ctc tat aaa ctc tac gag gcc gac ttt gtt ctc ttt 1372
 Trp Arg Gln Gln Leu Tyr Lys Leu Tyr Glu Ala Asp Phe Val Leu Phe 405
 395
 ggc tac ccc aag cca gaa aac ctg ctc agg gac tga gccccagaa 1418
 Gly Tyr Pro Lys Pro Glu Asn Leu Leu Arg Asp 415
 410
 gccctcacgc tgcccccaac aaattgaatg gctgtcccat gaggccgtcc tttgaggatg 1478
 ggaccctgtg gcctcctggg ttctctcctg gcttcctttg cttcctggtg tgacaggcag 1538
 aggattccac gcccccccct cgcattctgga gaccgtggta cagccaagac cgaagcacct 1598
 cactctccag agttttgctg tccccacccc cgcccttttg caatctggat ttgtttactc 1658
 cacagcctgt attcatggaa cactgtgtta aatactgttt tctaagatta atatatttca 1718
 gatatatatta atacgaaagt gggaggaagc tggagtaaag tgtggcgccc gcaaaaaaaaa 1778
 aaaaaaaaa 1785

<210> 9
 <211> 419
 <212> PRT
 <213> Mus musculus

<400> 9

Met Thr Lys Pro Arg Leu Phe Arg Leu Trp Leu Val Leu Gly Ser Ala
1 5 10 15

Leu Met Ile Leu Leu Ile Ile Val Tyr Trp Asp Asn Val Gly Thr Ala
20 25 30

His Phe Tyr Leu His Thr Ser Leu Ser Arg Pro His Ile Leu Glu Pro
35 40 45

Leu Pro Thr Gln Gly Leu Val Glu Glu Asn Val Phe Thr Ser Asp Val
50 55 60

Asp Glu Phe Leu Asp Thr Leu Leu Ser Ser Asp Ala Lys His Asn Asp
65 70 75 80

Leu Ser Arg Arg Lys Thr Glu Gln Pro Pro Ala Pro Ala Pro Ser Lys
85 90 95

Pro Val Leu Ser His Met Glu Glu Asn Val Arg Gly Tyr Asp Trp Ser
100 105 110

Thr His Asp Ala His Gln Asn Pro Asp Arg Asp Arg Gln Gln Ala Glu
115 120 125

Arg Arg Ser Leu Leu Arg Asp Phe Cys Ala Asn Ala Ser Leu Ala Phe
 130 135 140

Pro Thr Lys Asp Arg Ser Phe Asp Asp Ile Pro Asn Tyr Glu Leu Asn
 145 150 155 160

His Leu Ile Val Asp Asp Arg His Gly Val Ile Tyr Cys Tyr Val Pro
 165 170 175

Lys Val Ala Cys Thr Asn Trp Lys Arg Val Met Ile Val Leu Ser Glu
 180 185 190

Ser Leu Leu Asp Arg Gly Ser Pro Tyr Arg Asp Pro Leu Asp Ile Pro
 195 200 205

Arg Glu His Val His Asn Thr Ser Thr His Leu Thr Phe Asn Lys Phe
 210 215 220

Trp Arg Arg Tyr Gly Lys Phe Ser Arg His Leu Met Lys Val Lys Leu
 225 230 235 240

Lys Lys Tyr Thr Lys Phe Leu Phe Val Arg Asp Pro Phe Val Arg Leu
 245 250 255

Ile Ser Ala Phe Arg Ser Lys Phe Glu Leu Glu Asn Glu Glu Phe Tyr
 260 265 270

Arg Lys Phe Ala Val Pro Met Leu Arg Leu Tyr Ala Asn His Thr Ser
 275 280 285

Leu Pro Ala Ser Val Ser Glu Ala Phe Ser Ala Gly Leu Lys Val Ser
 290 295 300

Phe Ala Asn Phe Ile Gln Tyr Leu Leu Asp Pro His Thr Glu Lys Leu
 305 310 315 320

Ala Pro Phe Asn Glu His Trp Arg Gln Val Tyr Arg Leu Cys His Pro
 325 330 335

Cys Gln Ile Asp Tyr Asp Phe Val Gly Lys Leu Glu Thr Leu Asp Glu
 340 345 350

Asp Ala Ala Gln Leu Leu Arg Phe Leu Lys Val Asp Ser Gln Leu His
 355 360 365

Phe Pro Pro Ser Tyr Arg Asn Arg Thr Ala Ser Ser Trp Glu Glu Asp
 370 375 380

Trp Phe Ala Asn Ile Pro Leu Ala Trp Arg Gln Gln Leu Tyr Lys Leu
 385 390 395 400

Tyr Glu Ala Asp Phe Val Leu Phe Gly Tyr Pro Lys Pro Glu Asn Leu
 405 410 415

Leu Arg Asp

<210> 10
 <211> 1728
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (1)..(1116)

<400> 10
 atg act gtc gcc tgc cac gcg tgc cag gca cag cat ggg aag acg ctc 48
 Met Thr Val Ala Cys His Ala Cys Gln Ala Gln His Gly Lys Thr Leu
 1 5 10 15
 ctg ttg cag gcg gcc ctt gcc ggt ggt ggc aag tct ggg tgc tgc act 96
 Leu Leu Gln Ala Ala Leu Ala Gly Gly Gly Lys Ser Gly Cys Cys Thr
 20 25 30
 cct gct cct gtg cgc cct gcg tcc cgg gta acc aca gga aag gat gcc 144
 Pro Ala Pro Val Arg Pro Ala Ser Arg Val Thr Thr Gly Lys Asp Ala
 35 40 45
 cag gac act gaa tgg cag ggc tcc cca aaa gcc ctt ttg ggg gtt ccg 192
 Gln Asp Thr Glu Trp Gln Gly Ser Pro Lys Ala Leu Leu Gly Val Pro
 50 55 60
 aca ttt gaa aat aaa gct ctg ggc tcc agc tgg ttc ggt gga gtg agg 240
 Thr Phe Glu Asn Lys Ala Leu Gly Ser Ser Trp Phe Gly Gly Val Arg
 65 70 75 80
 aag agt ccc cta cag ctg ttg cgt gac ctg gac cag ggt cca cgc tcc 288
 Lys Ser Pro Leu Gln Leu Leu Arg Asp Leu Asp Gln Gly Pro Arg Ser
 85 90 95
 gcg atg gcc gag gtg cac cag cag cgg cgt gag ctg ctg cgc cgg gcc 336
 Ala Met Ala Glu Val His Gln Gln Arg Arg Glu Leu Leu Arg Arg Ala
 100 105 110
 tgc agc cgc cac acg cga cgc caa cgc ctg ctg cag ccg gag gac ctg 384
 Cys Ser Arg His Thr Arg Arg Gln Arg Leu Leu Gln Pro Glu Asp Leu
 115 120 125
 cgt cac gtg ctg gtg gac gac gcg cac cgg ctg ctg tac tgc tac gtg 432
 Arg His Val Leu Val Asp Asp Ala His Arg Leu Tyr Cys Tyr Val
 130 135 140
 cct aag gtg gcc tgc acc aac tgg aag cgt gtg atg ctg gcg ttg cgc 480
 Pro Lys Val Ala Cys Thr Asn Trp Lys Arg Val Met Leu Ala Leu Arg
 145 150 155 160
 ggc cgt ggg gat cca agc gca atc cct gcg cac gag gcg cat gcg cct 528
 Gly Arg Gly Asp Pro Ser Ala Ile Pro Ala His Glu Ala His Ala Pro
 165 170 175
 ggc ctg ctg ccc tcg ctg gcc gac ttt gcg ccg gct gag gtc aac tgg 576
 Gly Leu Leu Pro Ser Leu Ala Asp Phe Ala Pro Ala Glu Val Asn Trp
 180 185 190
 cgg ctg cgc gac tac ctg acc ttt ctc ttc gtg cgg gag ccc ttc gag 624
 Arg Leu Arg Asp Tyr Leu Thr Phe Leu Phe Val Arg Glu Pro Phe Glu
 195 200 205
 cgc ctg gcg tca gcc tac cgc aac aag ctg gcg cgg cca cac agc gcg 672
 Arg Leu Ala Ser Ala Tyr Arg Asn Lys Leu Ala Arg Pro His Ser Ala
 210 215 220

gcc ttc cag cgg cgc tat ggc aca cgc atc gtg cgt cgc cta cga cca Ala Phe Gln Arg Arg Tyr Gly Thr Arg Ile Val Arg Arg Leu Arg Pro 225 230 235 240	720
cac gcg cag ccc gat gcg ctg gcc cgc ggc cac gac gtg cgc ttc gcc His Ala Gln Pro Asp Ala Leu Ala Arg Gly His Asp Val Arg Phe Ala 245 250 255	768
gag ttc ctg gcc tac ctg ctc gac ccg cgc acg cgc cgt cat gag ccc Glu Phe Leu Ala Tyr Leu Leu Asp Pro Arg Thr Arg Arg His Glu Pro 260 265 270	816
ttc aac gaa cac tgg gag cgc gca cac gcg ctg tgc cat ccg tgc cta Phe Asn Glu His Trp Glu Arg Ala His Ala Leu Cys His Pro Cys Leu 275 280 285	864
gtg cgc tat gat gta gtg ggc aag ttt gag acg ata gca gat gat gct Val Arg Tyr Asp Val Val Gly Lys Phe Glu Thr Ile Ala Asp Asp Ala 290 295 300	912
gcc ttc gtg ctg gac ctg gtg ggt gag cct ggg cta cgt ttc cct gct Ala Phe Val Leu Asp Leu Val Gly Glu Pro Gly Leu Arg Phe Pro Ala 305 310 315 320	960
cca ccg ctg agg cca gag aag gac ctt acg cgt gag cag gcc cgg cgc Pro Pro Leu Arg Pro Glu Lys Asp Leu Thr Arg Glu Gln Ala Arg Arg 325 330 335	1008
ctt ttc cag gac atc agc ccc ttc tac cag cgt cgc ctc ttt aac ctc Leu Phe Gln Asp Ile Ser Pro Phe Tyr Gln Arg Arg Leu Phe Asn Leu 340 345 350	1056
tat aag atg gac ttt ctg ctc ttc aat tac tct gcc cct tcc tac ctg Tyr Lys Met Asp Phe Leu Leu Phe Asn Tyr Ser Ala Pro Ser Tyr Leu 355 360 365	1104
cga ctg caa taa ggggtgttggg tgcaatagag ccagtggctg ctgtgaccag Arg Leu Gln 370	1156
gaggccacca ggaggctctgg aacgaacctg gttgtgtgga ttggagacct tatccagtgg	1216
gcctgaccaa gagtctggcc actggtcaca ctcatccga ctgggtaggg tacaggttgc	1276
tttaggtgac cataaccttg tcaggccggt tctgctgtta gtttgatgtg tgtctcttcc	1336
tcccactctg cagatgtcag gcttcttcct aggactccag gtttgtagtt ctttggtttg	1396
gtttgagagg ccatttctca gtcttgtctg tgtagaacct gtccgtggca tgggtctaca	1456
agacagaatg tcatggcttg gttcagtgtg gcctaaggctc ttgtcagcat ttactgctta	1516
ggagttaaca tgagctgcct gccacccag cagtcacagg atggtgagca ctaccactcc	1576
acatctacct ggctgatcta cctttgatct cagccttgca agaggctgga tcttcccctg	1636
tgtcagcaaa ggccaagatg caatactgtg gcagcttttc cagctcactt ttattttttt	1696
tgttggtttt ttaaataaat atgttttggt ac	1728

<210> 11
 <211> 371
 <212> PRT
 <213> Mus musculus

<400> 11

Met Thr val Ala Cys His Ala Cys Gln Ala Gln His Gly Lys Thr Leu
1 5 10 15

Leu Leu Gln Ala Ala Leu Ala Gly Gly Gly Lys Ser Gly Cys Cys Thr
 20 25 30
 Pro Ala Pro Val Arg Pro Ala Ser Arg Val Thr Thr Gly Lys Asp Ala
 35 40 45
 Gln Asp Thr Glu Trp Gln Gly Ser Pro Lys Ala Leu Leu Gly Val Pro
 50 55 60
 Thr Phe Glu Asn Lys Ala Leu Gly Ser Ser Trp Phe Gly Gly Val Arg
 65 70 75 80
 Lys Ser Pro Leu Gln Leu Leu Arg Asp Leu Asp Gln Gly Pro Arg Ser
 85 90 95
 Ala Met Ala Glu Val His Gln Gln Arg Arg Glu Leu Leu Arg Arg Ala
 100 105 110
 Cys Ser Arg His Thr Arg Arg Gln Arg Leu Leu Gln Pro Glu Asp Leu
 115 120 125
 Arg His Val Leu Val Asp Asp Ala His Arg Leu Leu Tyr Cys Tyr Val
 130 135 140
 Pro Lys Val Ala Cys Thr Asn Trp Lys Arg Val Met Leu Ala Leu Arg
 145 150 155 160
 Gly Arg Gly Asp Pro Ser Ala Ile Pro Ala His Glu Ala His Ala Pro
 165 170 175
 Gly Leu Leu Pro Ser Leu Ala Asp Phe Ala Pro Ala Glu Val Asn Trp
 180 185 190
 Arg Leu Arg Asp Tyr Leu Thr Phe Leu Phe Val Arg Glu Pro Phe Glu
 195 200 205
 Arg Leu Ala Ser Ala Tyr Arg Asn Lys Leu Ala Arg Pro His Ser Ala
 210 215 220
 Ala Phe Gln Arg Arg Tyr Gly Thr Arg Ile Val Arg Arg Leu Arg Pro
 225 230 235 240
 His Ala Gln Pro Asp Ala Leu Ala Arg Gly His Asp Val Arg Phe Ala
 245 250 255
 Glu Phe Leu Ala Tyr Leu Leu Asp Pro Arg Thr Arg Arg His Glu Pro
 260 265 270
 Phe Asn Glu His Trp Glu Arg Ala His Ala Leu Cys His Pro Cys Leu
 275 280 285
 Val Arg Tyr Asp Val Val Gly Lys Phe Glu Thr Ile Ala Asp Asp Ala

290 295 300

Ala Phe Val Leu Asp Leu Val Gly Glu Pro Gly Leu Arg Phe Pro Ala
 305 310 315 320

Pro Pro Leu Arg Pro Glu Lys Asp Leu Thr Arg Glu Gln Ala Arg Arg
 325 330 335

Leu Phe Gln Asp Ile Ser Pro Phe Tyr Gln Arg Arg Leu Phe Asn Leu
 340 345 350

Tyr Lys Met Asp Phe Leu Leu Phe Asn Tyr Ser Ala Pro Ser Tyr Leu
 355 360 365

Arg Leu Gln
 370

<210> 12
 <211> 2073
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (129)..(1259)

<400> 12
 ccggcccttg tcctgctg caccgccgga gctggccacc cactatccct cccctccga 60
 gacctccagc ccctgctgca gtcacctccc ctgcagcctc gaggtcggcg aggtctggcc 120
 gcagcacc atg ttt ccc cgc cct ctg acc cca ctg gct gcc ccg aaa agc 170
 Met Phe Pro Arg Pro Leu Thr Pro Leu Ala Ala Pro Lys Ser
 1 5 10
 gcg gag acc ctg ggc cgc acg cca agg cgg gcc cca ttg ggc cgg gcc 218
 Ala Glu Thr Leu Gly Arg Thr Pro Arg Arg Ala Pro Leu Gly Arg Ala
 15 20 25 30
 cgg gct ggg ctc ggg ggg ccg ccc ctg ctg ctg ccg tcc atg ctg atg 266
 Arg Ala Gly Leu Gly Gly Pro Pro Leu Leu Leu Pro Ser Met Leu Met
 35 40 45
 ttc gct gta atc gtg gcc tcc agc gga ctg ctg ctc atg atc gag cga 314
 Phe Ala Val Ile Val Ala Ser Ser Gly Leu Leu Leu Met Ile Glu Arg
 50 55 60
 ggc atc cta tcg gag atg aaa ccc ctt ccc ctg cac cct ccc agc cac 362
 Gly Ile Leu Ser Glu Met Lys Pro Leu Pro Leu His Pro Pro Ser His
 65 70 75
 aaa ggc gcg gcc tgg agc ggg aca gat cct aag cct aga ggc cta tcc 410
 Lys Gly Ala Ala Trp Ser Gly Thr Asp Pro Lys Pro Arg Gly Leu Ser
 80 85 90
 ttg gat gct ggg gac tcg gac ttg caa gtg agg gag gac atc cga aac 458
 Leu Asp Ala Gly Asp Ser Asp Leu Gln Val Arg Glu Asp Ile Arg Asn
 95 100 105 110
 cgg acc ttg agg gcc gtg tgc gga caa cca ggc atg ccc cgg gac ccc 506
 Arg Thr Leu Arg Ala Val Cys Gly Gln Pro Gly Met Pro Arg Asp Pro
 115 120 125
 tgg gac ttg ccg gtg gga cag cgg cgc acc ctg ctg cgc cac att ctc 554

Trp	Asp	Leu	Pro	Val	Gly	Gln	Arg	Arg	Thr	Leu	Leu	Arg	His	Ile	Leu		
			130					135					140				
gta	agt	gac	cgc	tac	cgc	ttc	ctc	tac	tgc	tat	gtc	ccc	aaa	gtg	gcc	602	
Val	Ser	Asp	Arg	Tyr	Arg	Phe	Leu	Tyr	Cys	Tyr	Val	Pro	Lys	Val	Ala		
		145					150					155					
tgc	tct	aac	tgg	aaa	cgt	gtg	ctg	aag	gtg	ctg	gct	ggc	gtc	ctg	aac	650	
Cys	Ser	Asn	Trp	Lys	Arg	Val	Leu	Lys	Val	Leu	Ala	Gly	Val	Leu	Asn		
	160					165					170						
aac	gtg	gat	gtc	cgc	ctc	aag	atg	gac	cac	ccc	agt	gac	ttg	gtg	ttt	698	
Asn	Val	Asp	Val	Arg	Leu	Lys	Met	Asp	His	Pro	Ser	Asp	Leu	Val	Phe		
175					180					185					190		
ctg	gca	gac	ctg	cgg	cct	gag	gag	att	cgc	tac	cgt	ctg	cag	cac	tac	746	
Leu	Ala	Asp	Leu	Arg	Pro	Glu	Glu	Ile	Arg	Tyr	Arg	Leu	Gln	His	Tyr		
				195					200					205			
ttc	aag	ttc	ctg	ttt	gtg	cga	gac	ccc	ttg	gaa	cgc	ctc	ctg	tct	gct	794	
Phe	Lys	Phe	Leu	Phe	Val	Arg	Asp	Pro	Leu	Glu	Arg	Leu	Leu	Ser	Ala		
			210					215					220				
tac	cgt	aac	aag	ttt	gga	gag	atc	cga	gag	tac	cag	cag	cga	tat	ggg	842	
Tyr	Arg	Asn	Lys	Phe	Gly	Glu	Ile	Arg	Glu	Tyr	Gln	Gln	Arg	Tyr	Gly		
		225					230					235					
gcc	gaa	att	gtc	agg	cgc	tac	agg	gct	gga	gct	ggt	ccc	agc	cct	gca	890	
Ala	Glu	Ile	Val	Arg	Arg	Tyr	Arg	Ala	Gly	Ala	Gly	Pro	Ser	Pro	Ala		
	240					245					250						
ggg	gac	gat	gtc	acc	ttc	cca	gag	ttc	ctg	aga	tac	ctg	gtg	gat	gag	938	
Gly	Asp	Asp	Val	Thr	Phe	Pro	Glu	Phe	Leu	Arg	Tyr	Leu	Val	Asp	Glu		
255					260				265						270		
gat	cct	gaa	cat	atg	aat	gag	cat	tgg	atg	cct	gtg	tac	cac	ctg	tgc	986	
Asp	Pro	Glu	His	Met	Asn	Glu	His	Trp	Met	Pro	Val	Tyr	His	Leu	Cys		
				275					280					285			
caa	cca	tgt	gct	gtg	cac	tac	gac	ttc	gtg	ggt	tcc	tat	gag	agg	ctg	1034	
Gln	Pro	Cys	Ala	Val	His	Tyr	Asp	Phe	Val	Gly	Ser	Tyr	Glu	Arg	Leu		
			290					295					300				
gag	gct	gat	gcc	aac	cag	gtg	ctg	gag	tgg	gtg	cgg	gcc	cca	ccc	cat	1082	
Glu	Ala	Asp	Ala	Asn	Gln	Val	Leu	Glu	Trp	Val	Arg	Ala	Pro	Pro	His		
		305					310					315					
gtc	cgg	ttc	cca	gct	cgc	cag	gcc	tgg	tac	cgg	cca	gcc	agc	cca	gaa	1130	
Val	Arg	Phe	Pro	Ala	Arg	Gln	Ala	Trp	Tyr	Arg	Pro	Ala	Ser	Pro	Glu		
	320					325					330						
agt	ctg	cat	tac	cac	ttg	tgc	aat	gtt	cca	cgg	gcc	ctg	ctt	caa	gat	1178	
Ser	Leu	His	Tyr	His	Leu	Cys	Asn	Val	Pro	Arg	Ala	Leu	Leu	Gln	Asp		
335					340					345					350		
gtg	cta	cct	aag	tat	atc	ctg	gac	ttc	tcc	ctc	ttt	gct	tac	cca	ctg	1226	
Val	Leu	Pro	Lys	Tyr	Ile	Leu	Asp	Phe	Ser	Leu	Phe	Ala	Tyr	Pro	Leu		
				355					360					365			
ccc	aat	gtc	acc	aag	gaa	gcc	tgt	cac	caa	tga	cagtaggcca	gcaccttttg				1279	
Pro	Asn	Val	Thr	Lys	Glu	Ala	Cys	His	Gln								
			370					375									
gagtttgggt	ttaatgat	at	cagctttggg	atgtctttca	gagaaactcc	tggctctggg										1339	
tggcttcctg	gtttctctag	gtgtctccat	atctcagtg	taaggactgt	ccttgagggt											1399	
ccttggtccac	agtggtcag	aggacagagc	tagaaaggag	gcctgctgct	ttcactgggtg											1459	
aactgcctct	cttaggggcc	tgtggtatcc	gtgtctgcag	ggcaccagtg	gttattaaag											1519	

ccatatgttt gatcgaaaga ctgacttcag cccctggct gctgggtcta tgcagtccac 1579
ctggtctgtt gtaatttaac ctgtggccaa atcccaaata tgacactagc caagcacatg 1639
atcatgccta ggaccaatgg ctgtgacccc ctattcaccc atcccatgga cctcaggact 1699
ggagtgagct gtggtgcctt agaaatgaaa tgtgtgcaat tctactccag acttttacat 1759
ttcctcctct tgctaggtct gaatcathtt tctaaggaaa gagaaacgga agtggggccc 1819
ttacctcgaa gctctaaagc ccagcccctc aagcatccaa agacgcctgt gcctgacctc 1879
ttcctagggc tcttgagca tcttcaataa gcctcccttc cctacaaacc tttggagact 1939
atgtgagact gtatggccca tatatctggc tgtcacttgt ctaatgcatt tattttaaat 1999
gtgtatatatt taataggatc cttgtaaggg ctgactttta ataaagcttt ttcatatata 2059
aaaaaaaaa aaaa 2073

<210> 13
<211> 376
<212> PRT
<213> Mus musculus

<400> 13

Met Phe Pro Arg Pro Leu Thr Pro Leu Ala Ala Pro Lys Ser Ala Glu
1 5 10 15

Thr Leu Gly Arg Thr Pro Arg Arg Ala Pro Leu Gly Arg Ala Arg Ala
20 25 30

Gly Leu Gly Gly Pro Pro Leu Leu Leu Pro Ser Met Leu Met Phe Ala
35 40 45

Val Ile Val Ala Ser Ser Gly Leu Leu Leu Met Ile Glu Arg Gly Ile
50 55 60

Leu Ser Glu Met Lys Pro Leu Pro Leu His Pro Pro Ser His Lys Gly
65 70 75 80

Ala Ala Trp Ser Gly Thr Asp Pro Lys Pro Arg Gly Leu Ser Leu Asp
85 90 95

Ala Gly Asp Ser Asp Leu Gln Val Arg Glu Asp Ile Arg Asn Arg Thr
100 105 110

Leu Arg Ala Val Cys Gly Gln Pro Gly Met Pro Arg Asp Pro Trp Asp
115 120 125

Leu Pro Val Gly Gln Arg Arg Thr Leu Leu Arg His Ile Leu Val Ser
130 135 140

Asp Arg Tyr Arg Phe Leu Tyr Cys Tyr Val Pro Lys Val Ala Cys Ser
145 150 155 160

Asn Trp Lys Arg Val Leu Lys Val Leu Ala Gly Val Leu Asn Asn Val
165 170 175

Asp Val Arg Leu Lys Met Asp His Pro Ser Asp Leu Val Phe Leu Ala
 180 185 190
 Asp Leu Arg Pro Glu Glu Ile Arg Tyr Arg Leu Gln His Tyr Phe Lys
 195 200 205
 Phe Leu Phe Val Arg Asp Pro Leu Glu Arg Leu Leu Ser Ala Tyr Arg
 210 215 220
 Asn Lys Phe Gly Glu Ile Arg Glu Tyr Gln Gln Arg Tyr Gly Ala Glu
 225 230 235 240
 Ile Val Arg Arg Tyr Arg Ala Gly Ala Gly Pro Ser Pro Ala Gly Asp
 245 250 255
 Asp Val Thr Phe Pro Glu Phe Leu Arg Tyr Leu Val Asp Glu Asp Pro
 260 265 270
 Glu His Met Asn Glu His Trp Met Pro Val Tyr His Leu Cys Gln Pro
 275 280 285
 Cys Ala Val His Tyr Asp Phe Val Gly Ser Tyr Glu Arg Leu Glu Ala
 290 295 300
 Asp Ala Asn Gln Val Leu Glu Trp Val Arg Ala Pro Pro His Val Arg
 305 310 315 320
 Phe Pro Ala Arg Gln Ala Trp Tyr Arg Pro Ala Ser Pro Glu Ser Leu
 325 330 335
 His Tyr His Leu Cys Asn Val Pro Arg Ala Leu Leu Gln Asp Val Leu
 340 345 350
 Pro Lys Tyr Ile Leu Asp Phe Ser Leu Phe Ala Tyr Pro Leu Pro Asn
 355 360 365
 Val Thr Lys Glu Ala Cys His Gln
 370 375

<210> 14
 <211> 6000
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (496)..(1914)

<400> 14
 cttagccact gaaccatctc tccagcctct gttttgggggt gtttgtttga gacagaatca 60
 aaaggctgggt cttaaagtaca tattgtagct aaagatggct tccaactcct gccttgccct 120
 cagagtgtta ggggtgtgacc tttaaagtac ctattaaaga aacccaaca ggtgtagagt 180
 acacctgtga tcccagctct taggaggttg gacagatgga taaagagggtc cagggtgtccc 240

ggactacata gagagaccct atctcaaaag agaggcgaag acctaggttt tgagagtggg	300
cctgaggtct ctcttcaaaa gttttataga aagatgtctc tgttccctgt ctatgaacac	360
ggatggcctg agcacctgtc tcttcacagg atcagagtgt ccccccacctg aagagggctg	420
attgggtccc aagctatgct cctgagctga gtgcctgcag ccagtctgag gaactccatg	480
gcgccccctc tcccc atg gag aaa gga ctc gct ttg cct cag gat ttc cgg	531
Met Glu Lys Gly Leu Ala Leu Pro Gln Asp Phe Arg	
1 5 10	
gac ctt gta cac agc cta aag att cga ggc aga tac gtc ttg ttc ctg	579
Asp Leu Val His Ser Leu Lys Ile Arg Gly Arg Tyr Val Leu Phe Leu	
15 20 25	
gca ttt gtg gtc ata gtt ttt atc ttc att gaa aag gaa aat aaa atc	627
Ala Phe Val Val Ile Val Phe Ile Phe Ile Glu Lys Glu Asn Lys Ile	
30 35 40	
ata tcc agg gtc tcc gac aag ctg aag cag atc cct cat ttt gtg gca	675
Ile Ser Arg Val Ser Asp Lys Leu Lys Gln Ile Pro His Phe Val Ala	
45 50 55 60	
gat gcc aac agc act gac cca gcc ctg ctc tta tcg gag aat gca tct	723
Asp Ala Asn Ser Thr Asp Pro Ala Leu Leu Leu Ser Glu Asn Ala Ser	
65 70 75	
ctc ttg tcc ctg agc gag ttg gat tcc acc ttt tcc cat ctg cgg agc	771
Leu Leu Ser Leu Ser Glu Leu Asp Ser Thr Phe Ser His Leu Arg Ser	
80 85 90	
cgc ctg cac aac ctg agc ctg cag ctg ggc gtg gag cca gca atg gag	819
Arg Leu His Asn Leu Ser Leu Gln Leu Gly Val Glu Pro Ala Met Glu	
95 100 105	
agc cag gag gct ggg gca gag aag cca tcc cag cag gct gga gca ggg	867
Ser Gln Glu Ala Gly Ala Glu Lys Pro Ser Gln Gln Ala Gly Ala Gly	
110 115 120	
acc cgg cgc cac gtg ctt ctc atg gcc acc acc cgc acg ggt tcc tcg	915
Thr Arg Arg His Val Leu Leu Met Ala Thr Thr Arg Thr Gly Ser Ser	
125 130 135 140	
ttc gtg ggc gag ttc ttc aac cag cag ggc aat atc ttc tac ctc ttc	963
Phe Val Gly Glu Phe Phe Asn Gln Gln Gly Asn Ile Phe Tyr Leu Phe	
145 150 155	
gag cca ctg tgg cac atc gag cgc acc gtg ttc ttc cag cag cga ggc	1011
Glu Pro Leu Trp His Ile Glu Arg Thr Val Phe Phe Gln Gln Arg Gly	
160 165 170	
gcc agc gcg gct ggt tca gcc ttg gtc tac cgt gat gtc ctc aag cag	1059
Ala Ser Ala Ala Gly Ser Ala Leu Val Tyr Arg Asp Val Leu Lys Gln	
175 180 185	
ttg ttg cta tgc gac ctg tat gtg ctg gag ccc ttc atc agc cct ccg	1107
Leu Leu Leu Cys Asp Leu Tyr Val Leu Glu Pro Phe Ile Ser Pro Pro	
190 195 200	
ccc gag gac cac ttg act cag ttc ctg ttc cgc cgg gga tcc agc cgt	1155
Pro Glu Asp His Leu Thr Gln Phe Leu Phe Arg Arg Gly Ser Ser Arg	
205 210 215 220	
tca ctc tgc gag gat ccg gtg tgc aca ccc ttc gtc aag aag gtc ttt	1203
Ser Leu Cys Glu Asp Pro Val Cys Thr Pro Phe Val Lys Lys Val Phe	
225 230 235	
gag aag tac cac tgc agg aac cgt cgc tgc ggg cca ctc aac gtg acc	1251
Glu Lys Tyr His Cys Arg Asn Arg Arg Cys Gly Pro Leu Asn Val Thr	

240	245	250			
ttg gcg ggc gag gcc tgc cgc cgc aag gac cac gtg gcc ctc aag gct Leu Ala Gly Glu Ala Cys Arg Arg Lys Asp His Val Ala Leu Lys Ala	255	260	265	1299	
gtg cgc atc cgt cag ctg gag ttc ctg cag ccg cta gtt gag gac ccg Val Arg Ile Arg Gln Leu Glu Phe Leu Gln Pro Leu Val Glu Asp Pro	270	275	280	1347	
agg ttg gat cta cga gtc att cag ctg gtg cgc gac ccc cgg gcc gtg Arg Leu Asp Leu Arg Val Ile Gln Leu Val Arg Asp Pro Arg Ala Val	285	290	295	300	1395
ctg gct tca cgc ata gtg gcc ttt gcg ggc aag tat gag aac tgg aag Leu Ala Ser Arg Ile Val Ala Phe Ala Gly Lys Tyr Glu Asn Trp Lys	305	310	315	1443	
aag tgg ctg tcc gag ggg cag gac cag ctg agc gag gat gag gtg cag Lys Trp Leu Ser Glu Gly Gln Asp Gln Leu Ser Glu Asp Glu Val Gln	320	325	330	1491	
cga ttg cgg ggc aac tgt gag agc atc cgc ctg tct gca gag ctg ggc Arg Leu Arg Gly Asn Cys Glu Ser Ile Arg Leu Ser Ala Glu Leu Gly	335	340	345	1539	
ttg cgg cag cca gcc tgg ctg cgc ggt cgt tac atg ctg gtg cgc tat Leu Arg Gln Pro Ala Trp Leu Arg Gly Arg Tyr Met Leu Val Arg Tyr	350	355	360	1587	
gag gat gtg gca cgc agg cca ctg cag aag gcc cga gag atg tac agc Glu Asp Val Ala Arg Arg Pro Leu Gln Lys Ala Arg Glu Met Tyr Ser	365	370	375	380	1635
ttt gcg ggc atc ccc ttg acc ccg cag gtg gag gac tgg atc cag aag Phe Ala Gly Ile Pro Leu Thr Pro Gln Val Glu Asp Trp Ile Gln Lys	385	390	395	1683	
aac acg cag gcg aca cgc gac agc agc gat gtc tac tcc act cag aaa Asn Thr Gln Ala Thr Arg Asp Ser Ser Asp Val Tyr Ser Thr Gln Lys	400	405	410	1731	
aac tct tct gag cag ttt gag aag tgg cgc ttc agc atg cct ttc aag Asn Ser Ser Glu Gln Phe Glu Lys Trp Arg Phe Ser Met Pro Phe Lys	415	420	425	1779	
ctg gca cag gtg gta cag gct gcc tgt ggc ccg acc atg cac ctc ttt Leu Ala Gln Val Val Gln Ala Ala Cys Gly Pro Thr Met His Leu Phe	430	435	440	1827	
ggc tac aag ttg gcc agg gat gcc gcc tca ctc acc aac cgc tcc atc Gly Tyr Lys Leu Ala Arg Asp Ala Ala Ser Leu Thr Asn Arg Ser Ile	445	450	455	460	1875
agc ctg ctg gag gag cgg ggc acc ttc tgg gtc acg tag tgggggatgt Ser Leu Leu Glu Glu Arg Gly Thr Phe Trp Val Thr	465	470		1924	
ctgggaccct tggaactcct tcttgtgaaa ggctggccct gcttcctca caccagcct				1984	
ggcagtgaga catagccctg gcagaaagtc aaaatgggga gcgatgatgg gaacatagcc				2044	
cctggctgta gcggtagggc cccctgccag ctagactccc cagagcagct acagctaggg				2104	
ccttgggctc ctctgaggac accctgtctc cttagtggta ctggtcataa ggtgtcctca				2164	
ccctatgacc tgcagtgtcc cgagcaggtc aaggtagggt cctgtgtgtg gacacacctc				2224	
cctagctctt tctcacacag tctacacgaa gcctgtaaag gcctgtaggt tgtgtgggct				2284	
ggagtccaag tattaacaa ccagaagggg tgtagaccct ctgcagcctc tcagacctcc				2344	

tactgtatta	agtgttaacc	tcttccgctc	tgagtcagag	aagctgggat	ctggctgagt	2404
cctgggaagg	aggaggacag	cctaggggtga	ggaagggggac	ctttgaagct	cctcagggaa	2464
cagtggctgt	gtaccagctc	atagaaaatg	gtttgatcag	ctgttctagt	cactcatgag	2524
tatcaatcag	cctgtgtaga	gcaagacaca	caaagtgcag	tgaaaacaga	caaggccagg	2584
ggtgtggctc	agcggtaaag	cccttgccctc	tctcatgccc	agggccctag	gtgtgattct	2644
tagcactata	aataataagg	aaaataagcc	atacatagct	gtgcagtaca	gagagagaca	2704
gaaaagcgta	ataaacattc	ccccctctca	taagcacacg	ctctgcctct	gtctctgtct	2764
ctgtctctct	ctgtctctgt	ctctctctgt	ctctctctct	cactctctct	ctctctctct	2824
ctctctcaca	cacacacaca	cacacacaca	cacaaacaca	cacacacaca	ctggtacaga	2884
aaagcttccc	ttctccactc	tttatgctca	ccatgttttt	aggatttttg	tttgtttgtt	2944
ttgttttggg	tttctatata	tctaaacaag	gagccttaaa	ttacatcttt	ttgggattac	3004
gtggcgaggg	ggagcagcag	ggagcatttg	cttctggtaa	aaggatagac	cagaaaggac	3064
ttcccccttg	atgatcggag	tcatgcagag	cccccagggt	cccacatgtt	ctgtgtgaac	3124
ttcatgtgga	atgactcaca	gaagtgacta	ccttagggcg	tcgctctgta	gctcaatgtg	3184
cgtagagttt	gtctattgta	taggaagccc	tgggttctgt	gccctgcccc	acaccatata	3244
ggtatgatgg	cgtgtaccca	tatcccagca	ttcaggaaat	caaaggtttg	cggtcaaaag	3304
ttcaagggtg	agctatgtat	cttttgagat	gtccgagtct	ataagctccc	tctactgaga	3364
gctgctagta	ttccttctca	agaacagggt	atttgaaaat	ctcttcttgt	aagccccaac	3424
atttggggga	ggggaggatg	ggggaaactc	tgactgagca	gctaggtttc	cccaagcaga	3484
agttcttctc	tcccagtgca	agctgagggt	tagtgccccg	gctagcctcg	gaactggctc	3544
agtgggggtt	ggcacgctga	cttcctggac	aattgctgca	tctgacactg	tgagcgccca	3604
ggacaccgga	tgtgtagcat	ggatgtgggc	ctagcacagg	tgtggaagag	gacagtcagg	3664
acactcactg	ttggtgtttg	ttatccacct	gttcagccgc	cgggtcccac	acatagggtca	3724
tctttcgtca	ctcaagccgg	cctctgtttt	tgattttaac	aatccaagag	caagtgtgta	3784
ggggacaaaag	agacaccagc	tcttcatact	tgatggagac	cgggacagga	tgctgcaggc	3844
aggctcggga	gtgcattatc	gtttcttctg	agctcttcat	ccaggccata	ttcccttttc	3904
tgcatcgcag	ctgggggtgg	aggggctgag	ctgggagttt	ccgtccttct	tccctgtggg	3964
ggtgggagat	ggggctcaag	gtctccatct	ctcggtccc	tgaggacaga	acccccacag	4024
tggacctttg	ggccttctca	ggacattgac	aatgttgtgt	gcactgcaag	ttgactttat	4084
ttattttgta	ggaaaaagag	atggtattct	ttagcaggat	ctgaaactgt	accctagtca	4144
agaagtacaa	ttaataacat	tattattaat	aacaattggg	atgattcaaa	ggtcacacac	4204
acaccagacc	acccaagaac	catgtaggag	atgaggcgga	gccagggtcat	ggaagcccag	4264
aatgctacag	agattctgtg	ggtctattat	tgatcagaaa	atacaacatg	ggggggcctt	4324
ttctatggca	taactcaggt	gtctgcaaag	acagagcagc	tcagggtgggt	ggataggcag	4384
gcagatggag	ccttgaatca	catcccctga	ggctcagcca	gccagtgcga	atcctcaagg	4444

```

ttcggaaagg gacagtgtag cggcaaacta ctgggcatct tctctgatgc ttagaggctc 4504
ttaccaagga gcttgatgga gaaagtggcc atgttggtgg catgggacaa ctgctcaccc 4564
aacttcaaag atcaaagggc acccaggtgg cctatgacag tgacactctt tcctaccatt 4624
aggggtgtct gccccactg aaagccatga aatttctggc cctaaggggg taggaaggac 4684
ttaggagtag cagatggttt gatcccatcc cccccacaca cacacacaca tccctcagct 4744
gtcttccaca ttagagccac ttcagttgtc catggacttg tccctttgag acatcctgga 4804
ttttgaagga tagaaacatc cccaagatgg tctcgtgtta atcccacaac agaggcagaa 4864
gggtcatact gagagagaga gagagagaga gagagagaga gagaggagtg ggggccagac 4924
cctcaagaag ccaagtgggt ctggtcaacc tgtgcacatg agaaggaggg aacatcacta 4984
aaatcagggc ctgggctggt gtgttggtgg tgagatcccg tggagtggct ggctagatat 5044
ggatgagttt tctgagcatg ctcacacacc cccaacttca cattcttaga aatagcacia 5104
ccataatgcc ttacctcaaa aggatgaggc aaagcttgca attaatccc agtgtctgaa 5164
gacctccggg atcctttgag ctgtgtgtcc ttgtcagtag atggggacag gtccttagg 5224
tattcctgac atagaggtaa ggtgctgccc ttgctgcctt gcataagctg tgacaagctt 5284
cttactgggc acatagaacg gctctgtcat ctgcttccac taaatttggg cttggacttt 5344
gcctcctgcc aaatctccat ctctgctgga tagtctagtc cctagggctt aaccaccacc 5404
ctccacttcc tgggtgggtcc tacaagcgtc gtctttgtgc cagtaccggg atgggtgcct 5464
gcctcatggc taaatggtac aggacatctt cccagactga gtggtcatgg catgtgcatg 5524
tatgttcatg agtacatgca tgggtgtgtg ggggtggatg gtgcattctg ttctgcttct 5584
ggtgcagcta cacagccaca acctcttctt gtcattgacc ttcttgggtt tctttagacc 5644
acagataatc ttccaatgcc cgattctgct gttctcatct gagagctgac aaccagcgc 5704
tcagagtaga gttcatgacc taggaaaccc ttctcctggt agcttactga acttatataa 5764
ttaaaaacga acatataggg gttggagaga tggttcattg gttaagagca ccggctgctt 5824
ttgcaaagga cctgagttca aatcccagca accatatggt ggcttacaac catctgtaat 5884
gagttctggt gccctcttct ggcatgcagg tgtacatgca gatagagcac tcctatgcat 5944
aaaataaata aatcttttaa ataaataaat aaatgaacat gaaaaaaaaa aaaaaa 6000

```

<210> 15

<211> 472

<212> PRT

<213> Mus musculus

<400> 15

Met Glu Lys Gly Leu Ala Leu Pro Gln Asp Phe Arg Asp Leu Val His
1 5 10 15

Ser Leu Lys Ile Arg Gly Arg Tyr Val Leu Phe Leu Ala Phe Val Val
20 25 30

Ile Val Phe Ile Phe Ile Glu Lys Glu Asn Lys Ile Ile Ser Arg Val
35 40 45

Ser Asp Lys Leu Lys Gln Ile Pro His Phe Val Ala Asp Ala Asn Ser
 50 55 60

Thr Asp Pro Ala Leu Leu Leu Ser Glu Asn Ala Ser Leu Leu Ser Leu
 65 70 75 80

Ser Glu Leu Asp Ser Thr Phe Ser His Leu Arg Ser Arg Leu His Asn
 85 90 95

Leu Ser Leu Gln Leu Gly Val Glu Pro Ala Met Glu Ser Gln Glu Ala
 100 105 110

Gly Ala Glu Lys Pro Ser Gln Gln Ala Gly Ala Gly Thr Arg Arg His
 115 120 125

Val Leu Leu Met Ala Thr Thr Arg Thr Gly Ser Ser Phe Val Gly Glu
 130 135 140

Phe Phe Asn Gln Gln Gly Asn Ile Phe Tyr Leu Phe Glu Pro Leu Trp
 145 150 155 160

His Ile Glu Arg Thr Val Phe Phe Gln Gln Arg Gly Ala Ser Ala Ala
 165 170 175

Gly Ser Ala Leu Val Tyr Arg Asp Val Leu Lys Gln Leu Leu Cys
 180 185 190

Asp Leu Tyr Val Leu Glu Pro Phe Ile Ser Pro Pro Pro Glu Asp His
 195 200 205

Leu Thr Gln Phe Leu Phe Arg Arg Gly Ser Ser Arg Ser Leu Cys Glu
 210 215 220

Asp Pro Val Cys Thr Pro Phe Val Lys Lys Val Phe Glu Lys Tyr His
 225 230 235 240

Cys Arg Asn Arg Arg Cys Gly Pro Leu Asn Val Thr Leu Ala Gly Glu
 245 250 255

Ala Cys Arg Arg Lys Asp His Val Ala Leu Lys Ala Val Arg Ile Arg
 260 265 270

Gln Leu Glu Phe Leu Gln Pro Leu Val Glu Asp Pro Arg Leu Asp Leu
 275 280 285

Arg Val Ile Gln Leu Val Arg Asp Pro Arg Ala Val Leu Ala Ser Arg
 290 295 300

Ile Val Ala Phe Ala Gly Lys Tyr Glu Asn Trp Lys Lys Trp Leu Ser
 305 310 315 320

Glu Gly Gln Asp Gln Leu Ser Glu Asp Glu Val Gln Arg Leu Arg Gly
 325 330 335

Asn Cys Glu Ser Ile Arg Leu Ser Ala Glu Leu Gly Leu Arg Gln Pro
 340 345 350
 Ala Trp Leu Arg Gly Arg Tyr Met Leu Val Arg Tyr Glu Asp Val Ala
 355 360 365
 Arg Arg Pro Leu Gln Lys Ala Arg Glu Met Tyr Ser Phe Ala Gly Ile
 370 375 380
 Pro Leu Thr Pro Gln Val Glu Asp Trp Ile Gln Lys Asn Thr Gln Ala
 385 390 395 400
 Thr Arg Asp Ser Ser Asp Val Tyr Ser Thr Gln Lys Asn Ser Ser Glu
 405 410 415
 Gln Phe Glu Lys Trp Arg Phe Ser Met Pro Phe Lys Leu Ala Gln Val
 420 425 430
 Val Gln Ala Ala Cys Gly Pro Thr Met His Leu Phe Gly Tyr Lys Leu
 435 440 445
 Ala Arg Asp Ala Ala Ser Leu Thr Asn Arg Ser Ile Ser Leu Leu Glu
 450 455 460
 Glu Arg Gly Thr Phe Trp Val Thr
 465 470

<210> 16
 <211> 2080
 <212> DNA
 <213> Mus musculus

<220>
 <221> CDS
 <222> (147)..(1601)

<400> 16
 caccgcccgg ctgctccact ccgctccacc caacattgag ggagacccga agaggccgga 60
 gccgaggact ttgggctggg ttttctggac agaaccagca ggcgcctact ctgctctggg 120
 tggggagagt gaggattcgg tgaact atg aag ggc cgg cgg cgg cgc cga 173
 Met Lys Gly Arg Arg Arg Arg Arg
 1 5
 aag tat tgc aag ttc acg ctg ctc ttg gcg ctg tac acg ctt ttg cta 221
 Lys Tyr Cys Lys Phe Thr Leu Leu Leu Ala Leu Tyr Thr Leu Leu Leu
 10 15 20 25
 ctt ctt gtc ccc tct gta ctg gac agc cac agc gag cag gac aag ggc 269
 Leu Leu Val Pro Ser Val Leu Asp Ser His Ser Glu Gln Asp Lys Gly
 30 35 40
 agg aac tgc ccc ggc ctg cag cgc agc ttg ggt gtg tgg agc ctg gag 317
 Arg Asn Cys Pro Gly Leu Gln Arg Ser Leu Gly Val Trp Ser Leu Glu
 45 50 55
 gcg gcg gcg gcc ggg gaa cgt gag cag ggc gct gag gtg cgg tcc ctg 365
 Ala Ala Ala Ala Gly Glu Arg Glu Gln Gly Ala Glu Val Arg Ser Leu

60	65	70	
gcc gaa gga aac ccg gat cga tcc ccc ggg tcc ccc ggc aac ctc agc Ala Glu Gly Asn Pro Asp Arg Ser Pro Gly Ser Pro Gly Asn Leu Ser 75 80 85			413
gcc gtc ggt gag gcg gtg acc cag gaa aag caa cac atc tat gtg cat Ala Val Gly Glu Ala Val Thr Gln Glu Lys Gln His Ile Tyr Val His 90 95 100			461
gcc acc tgg cgc acc ggc tgc tcc ttc ttg ggc gaa ctc ttc aac cag Ala Thr Trp Arg Thr Gly Ser Ser Phe Leu Gly Glu Leu Phe Asn Gln 110 115 120			509
cac ccg gac gtt ttc tac ttg tac gac ccc atg tgg cat ctg tgg cag His Pro Asp Val Phe Tyr Leu Tyr Asp Pro Met Trp His Leu Trp Gln 125 130 135			557
gca ctg tat ccg ggc gac gcg gag agc ctg cag ggc gca cta aga gac Ala Leu Tyr Pro Gly Asp Ala Glu Ser Leu Gln Gly Ala Leu Arg Asp 140 145 150			605
atg ctg cgc tcc ctc ttc cgc tgt gat ttc tct gtg ctg cgc ctg tac Met Leu Arg Ser Leu Phe Arg Cys Asp Phe Ser Val Leu Arg Leu Tyr 155 160 165			653
gcg cag cct ggg gac cct ggg gag cga gca ccg gac tcg gcc aac ctc Ala Gln Pro Gly Asp Pro Gly Glu Arg Ala Pro Asp Ser Ala Asn Leu 170 175 180 185			701
acc acg gcc atg ctt ttc cgc tgg cgg acc aac aag gtc atc tgc tcg Thr Thr Ala Met Leu Phe Arg Trp Arg Thr Asn Lys Val Ile Cys Ser 190 195 200			749
ccg cct ctg tgc ccc gcc gcg ccc cgg gca cgc gcg gac gtg gga ctc Pro Pro Leu Cys Pro Ala Ala Pro Arg Ala Arg Ala Asp Val Gly Leu 205 210 215			797
gtc gag gac aaa gcc tgc gaa agt acc tgc ccg ccc gtt tcg ctc cgc Val Glu Asp Lys Ala Cys Glu Ser Thr Cys Pro Pro Val Ser Leu Arg 220 225 230			845
gcc ctg gag gcc gag tgc cgc aag tac ccg gtg gtg gtc atc aaa gac Ala Leu Glu Ala Glu Cys Arg Lys Tyr Pro Val Val Val Ile Lys Asp 235 240 245			893
gtg cgg cta ctg gac ctg gga gtg ctg gtc cct ctg ctg cgt gac cca Val Arg Leu Leu Asp Leu Gly Val Leu Val Pro Leu Leu Arg Asp Pro 250 255 260 265			941
ggc ctc aac cta aag gtg gtg caa ctc ttc cga gac cct cgg gcc gtg Gly Leu Asn Leu Lys Val Val Gln Leu Phe Arg Asp Pro Arg Ala Val 270 280			989
cac aac tcg cgc ctc aag tcg agg cag gga ctg ctg cgc gaa agc atc His Asn Ser Arg Leu Lys Ser Arg Gln Gly Leu Leu Arg Glu Ser Ile 285 290 295			1037
cag gtg ctg cgc acg cgc cag agg ggc gac cac ttc cac cgg gtg ctg Gln Val Leu Arg Thr Arg Gln Arg Gly Asp His Phe His Arg Val Leu 300 305 310			1085
ctg gcg cat gga gtg gat gcc cgt ccg gga ggc cag gcc cgg gct ctg Leu Ala His Gly Val Asp Ala Arg Pro Gly Gly Gln Ala Arg Ala Leu 315 320 325			1133
ccc tcg gcg cca cgc gct gat ttc ttc tta acc agc gcg ctt gag gtg Pro Ser Ala Pro Arg Ala Asp Phe Phe Leu Thr Ser Ala Leu Glu Val 330 335 340 345			1181
atc tgt gaa gcg tgg ctt cgc gac ctg cta ttc acc cgc ggc gcg ccc			1229

Ile	Cys	Glu	Ala	Trp	Leu	Arg	Asp	Leu	Leu	Phe	Thr	Arg	Gly	Ala	Pro		
				350					355					360			
gcc	tgg	ctg	agg	cgt	cgc	tac	ctg	cgg	ctg	cgt	tat	gag	gac	ctg	gtg	1277	
Ala	Trp	Leu	Arg	Arg	Arg	Tyr	Leu	Arg	Leu	Arg	Tyr	Glu	Asp	Leu	Val		
			365					370					375				
tgg	cag	ccc	caa	gcc	cag	ctg	cgc	cgc	ctg	ctg	cgc	ttc	tct	ggg	ttg	1325	
Trp	Gln	Pro	Gln	Ala	Gln	Leu	Arg	Arg	Leu	Leu	Arg	Phe	Ser	Gly	Leu		
		380					385					390					
cgg	aca	ctc	gcc	gcg	ctt	gat	gcc	ttc	gca	ttc	aat	atg	acg	cgg	ggc	1373	
Arg	Thr	Leu	Ala	Ala	Leu	Asp	Ala	Phe	Ala	Phe	Asn	Met	Thr	Arg	Gly		
	395					400					405						
tcg	gcc	tac	ggc	gcc	gat	cgt	ccc	ttc	cac	ttg	tct	gcg	cgg	gac	gcc	1421	
Ser	Ala	Tyr	Gly	Ala	Asp	Arg	Pro	Phe	His	Leu	Ser	Ala	Arg	Asp	Ala		
410					415					420					425		
cga	gag	gct	gtg	cac	gcc	tgg	cgc	gaa	cgt	ctg	agc	caa	gag	cag	gtg	1469	
Arg	Glu	Ala	Val	His	Ala	Trp	Arg	Glu	Arg	Leu	Ser	Gln	Glu	Gln	Val		
				430				435						440			
cgc	caa	gtg	gaa	acc	gcc	tgc	gcc	cct	gcc	atg	cgt	ctg	ctt	gcc	tac	1517	
Arg	Gln	Val	Glu	Thr	Ala	Cys	Ala	Pro	Ala	Met	Arg	Leu	Leu	Ala	Tyr		
			445				450						455				
cct	cga	agt	ggg	gac	gaa	cgc	gac	agg	aag	acc	gtc	agg	gaa	ggg	gag	1565	
Pro	Arg	Ser	Gly	Asp	Glu	Arg	Asp	Arg	Lys	Thr	Val	Arg	Glu	Gly	Glu		
		460					465				470						
aca	cca	ctg	gag	acc	aag	gcc	aat	tgg	gct	gtg	taa	taccctgac				1611	
Thr	Pro	Leu	Glu	Thr	Lys	Ala	Asn	Trp	Ala	Val							
	475					480											
cctgaaccct	gccccggggc	gtattcaggt	cagtggccat	aaaaagggtga	actcagcatg											1671	
ctgcccccg	actggagagg	ctgcacgggtg	gaggcgatct	atcacactgt	gagacactgg											1731	
gactgatttg	gtatcaactg	ctgtgccatt	ctcctgggtca	ggagcatcac	aagctgttaa											1791	
gtaatgacag	acaccttggc	tgagatgaag	tttccagaaa	ggaagtaaca	gtgcaatgtg											1851	
gatattttgtg	accacaacat	aggaaaagct	gtacttccca	ggctgaactt	ggctcagctt											1911	
gagccatttc	aacaaggcat	cctcacaata	atgaagagat	gtgatctggt	ttcctttcac											1971	
atcagccaag	atgtctggac	aaaaccatca	atgtgaataa	gggccaaagtg	cagttgtgtc											2031	
tctcttgatt	aaattacttc	atattaaata	aaaaaaaaaa	aaaaaaaaaa												2080	

<210> 17
 <211> 484
 <212> PRT
 <213> Mus musculus

<400> 17

Met	Lys	Gly	Arg	Arg	Arg	Arg	Arg	Arg	Lys	Tyr	Cys	Lys	Phe	Thr	Leu		
1			5						10					15			
Leu	Leu	Ala	Leu	Tyr	Thr	Leu	Leu	Leu	Leu	Val	Pro	Ser	Val	Leu			
		20					25				30						
Asp	Ser	His	Ser	Glu	Gln	Asp	Lys	Gly	Arg	Asn	Cys	Pro	Gly	Leu	Gln		
		35					40					45					

Arg Ser Leu Gly Val Trp Ser Leu Glu Ala Ala Ala Ala Gly Glu Arg
 50 55 60
 Glu Gln Gly Ala Glu Val Arg Ser Leu Ala Glu Gly Asn Pro Asp Arg
 65 70 75 80
 Ser Pro Gly Ser Pro Gly Asn Leu Ser Ala Val Gly Glu Ala Val Thr
 85 90 95
 Gln Glu Lys Gln His Ile Tyr Val His Ala Thr Trp Arg Thr Gly Ser
 100 105 110
 Ser Phe Leu Gly Glu Leu Phe Asn Gln His Pro Asp Val Phe Tyr Leu
 115 120 125
 Tyr Asp Pro Met Trp His Leu Trp Gln Ala Leu Tyr Pro Gly Asp Ala
 130 135 140
 Glu Ser Leu Gln Gly Ala Leu Arg Asp Met Leu Arg Ser Leu Phe Arg
 145 150 155 160
 Cys Asp Phe Ser Val Leu Arg Leu Tyr Ala Gln Pro Gly Asp Pro Gly
 165 170 175
 Glu Arg Ala Pro Asp Ser Ala Asn Leu Thr Thr Ala Met Leu Phe Arg
 180 185 190
 Trp Arg Thr Asn Lys Val Ile Cys Ser Pro Pro Leu Cys Pro Ala Ala
 195 200 205
 Pro Arg Ala Arg Ala Asp Val Gly Leu Val Glu Asp Lys Ala Cys Glu
 210 215 220
 Ser Thr Cys Pro Pro Val Ser Leu Arg Ala Leu Glu Ala Glu Cys Arg
 225 230 235 240
 Lys Tyr Pro Val Val Val Ile Lys Asp Val Arg Leu Leu Asp Leu Gly
 245 250 255
 Val Leu Val Pro Leu Leu Arg Asp Pro Gly Leu Asn Leu Lys Val Val
 260 265 270
 Gln Leu Phe Arg Asp Pro Arg Ala Val His Asn Ser Arg Leu Lys Ser
 275 280 285
 Arg Gln Gly Leu Leu Arg Glu Ser Ile Gln Val Leu Arg Thr Arg Gln
 290 295 300
 Arg Gly Asp His Phe His Arg Val Leu Leu Ala His Gly Val Asp Ala
 305 310 315 320
 Arg Pro Gly Gly Gln Ala Arg Ala Leu Pro Ser Ala Pro Arg Ala Asp
 325 330 335

Phe Phe Leu Thr Ser Ala Leu Glu Val Ile Cys Glu Ala Trp Leu Arg
 340 345 350
 Asp Leu Leu Phe Thr Arg Gly Ala Pro Ala Trp Leu Arg Arg Arg Tyr
 355 360 365
 Leu Arg Leu Arg Tyr Glu Asp Leu Val Trp Gln Pro Gln Ala Gln Leu
 370 375 380
 Arg Arg Leu Leu Arg Phe Ser Gly Leu Arg Thr Leu Ala Ala Leu Asp
 385 390 395 400
 Ala Phe Ala Phe Asn Met Thr Arg Gly Ser Ala Tyr Gly Ala Asp Arg
 405 410 415
 Pro Phe His Leu Ser Ala Arg Asp Ala Arg Glu Ala Val His Ala Trp
 420 425 430
 Arg Glu Arg Leu Ser Gln Glu Gln Val Arg Gln Val Glu Thr Ala Cys
 435 440 445
 Ala Pro Ala Met Arg Leu Leu Ala Tyr Pro Arg Ser Gly Asp Glu Arg
 450 455 460
 Asp Arg Lys Thr Val Arg Glu Gly Glu Thr Pro Leu Glu Thr Lys Ala
 465 470 475 480
 Asn Trp Ala Val

<210> 18
 <211> 4713
 <212> DNA
 <213> Homo sapiens

<220>
 <221> CDS
 <222> (514)..(2199)

<400> 18
 ggaaatctgg cattttttaaa agtttgcgcc ccacaaagag gaaatattcc aaagggtactc 60
 aggatgtaaa aggggagatc ttcacagatg cctccgtgga tggcatggca atccatccat 120
 caatgagaag accatgattt cttttaattt tctgtgtgtt tccacattcc ccagtggagaa 180
 ttcttccacc tttttttgtg ccatgggaaa aacctgaagg gcaggcagag ctgctcccga 240
 acttgtgacc ttctctgagg ttgcagcggc tcttgtagaa catgactctg ggacatcact 300
 tccttttgtt ttctttcgga gctgaaccaa agaatgtgca ccctctttct ctagtgtgtg 360
 ggtgtctgct tatttttgta tttgtgcttt ccatccatct tctgtgatca caaggcattc 420
 ttaagggttt ctagcacgac ttgcggacat ccagactcgt gggggggccca cccatggctc 480
 ggtaagccag cagcccaggg cactggcact acc atg agg cac tgc att aat tgc 534
 Met Arg His Cys Ile Asn Cys
 1 5

tgc Cys	ata Ile	cag Gln 10	ctg Leu	tta Leu	ccc Pro	gac Asp	ggc Gly 15	gca Ala	cac His	aag Lys	cag Gln 20	cag Gln	gtc Val	aac Asn	tgc Cys	582
caa Gln	ggg Gly 25	ggc Gly	ccc Pro	cat His	cac His	ggt Gly 30	cac His	cag Gln	gcg Ala	tgc Cys	ccc Pro 35	acg Thr	tgc Cys	aaa Lys	gga Gly	630
gaa Glu 40	aac Asn	aaa Lys	att Ile	ctg Leu	ttt Phe 45	cgt Arg	gtg Val	gac Asp	agt Ser	aag Lys 50	cag Gln	atg Met	aac Asn	ttg Leu	ctt Leu 55	678
gct Ala	gtt Val	ctc Leu	gaa Glu	gtg Val 60	agg Arg	act Thr	gaa Glu	ggg Gly 65	aac Asn	gaa Glu	aac Asn	tgg Trp	ggt Gly 70	ggg Gly	ttt Phe	726
ttg Leu	cgc Arg	ttc Phe	aaa Lys 75	aag Lys	ggg Gly	aag Lys	cga Arg	tgt Cys 80	agc Ser	ctc Leu	gtt Val	ttt Phe	gga Gly 85	ctg Leu	ata Ile	774
ata Ile	atg Met	acc Thr 90	ttg Leu	gta Val	atg Met	gct Ala	tct Ser 95	tac Tyr	atc Ile	ctt Leu	tct Ser	ggg Gly 100	gcc Ala	cac His	caa Gln	822
gag Glu	ctt Leu 105	ctg Leu	atc Ile	tca Ser	tca Ser	cct Pro 110	ttc Phe	cat His	tac Tyr	gga Gly	ggc Gly 115	ttc Phe	ccc Pro	agc Ser	aac Asn	870
ccc Pro 120	agc Ser	ttg Leu	atg Met	gac Asp	agc Ser 125	gaa Glu	aac Asn	cca Pro	agt Ser	gac Asp 130	aca Thr	aag Lys	gag Glu	cat His	cac His 135	918
cac His	caa Gln	tcc Ser	tct Ser	gta Val 140	aat Asn	aat Asn	att Ile	tca Ser	tac Tyr 145	atg Met	aag Lys	gac Asp	tat Tyr	cca Pro 150	agc Ser	966
att Ile	aaa Lys	tta Leu	att Ile 155	atc Ile	aac Asn	agc Ser	atc Ile	aca Thr 160	act Thr	agg Arg	att Ile	gag Glu	ttc Phe 165	acg Thr	acc Thr	1014
aga Arg	cag Gln 170	ctc Leu	cca Pro	gac Asp	tta Leu	gaa Glu	gac Asp 175	ctt Leu	aag Lys	aag Lys	cag Gln	gag Glu 180	ttg Leu	cat His	atg Met	1062
ttt Phe 185	tca Ser	gtc Val	atc Ile	ccc Pro	aac Asn	aaa Lys 190	ttc Phe	ctt Leu	cca Pro	aac Asn	agt Ser 195	aag Lys	agc Ser	ccc Pro	tgt Cys	1110
tgg Trp 200	tac Tyr	gag Glu	gag Glu	ttc Phe	tgc Ser 205	ggg Gly	cag Gln	aac Asn	acc Thr	acc Thr 210	gac Asp	ccc Pro	tac Tyr	ctc Leu	acc Thr 215	1158
aac Asn	tcc Ser	tac Tyr	gtg Val 220	ctc Leu	tac Tyr	tcc Ser	aag Lys	cgc Arg	ttc Phe 225	cgc Arg	tcc Ser	acc Thr	ttc Phe	gac Asp 230	gcc Ala	1206
ctg Leu	cgc Arg	aag Lys	gcc Ala 235	ttc Phe	tgg Trp	ggc Gly	cac His	ctg Leu 240	gcg Ala	cac His	gcg Ala	cac His	ggg Gly 245	aag Lys	cac His	1254
ttc Phe	cgc Arg	ctg Leu 250	cgc Arg	tgc Cys	ctg Leu	ccg Pro	cac His 255	ttc Phe	tac Tyr	atc Ile	ata Ile	ggg Gly 260	cag Gln	ccc Pro	aag Lys	1302
tgc Cys	ggg Gly 265	acc Thr	aca Thr	gac Asp	ctc Leu	tat Tyr 270	gac Asp	cgc Arg	ctg Leu	cgg Arg	ctg Leu 275	cac His	cct Pro	gag Glu	gtc Val	1350
aag Lys	ttc Phe	tcc Ser	gcc Ala	atc Ile	aag Lys	gag Glu	cca Pro	cac His	tgg Trp	tgg Trp	acc Thr	cgg Arg	aag Lys	cgc Arg	ttt Phe	1398

280		285		290		295	
gga atc gtc cgc cta aga gat ggg ctg cga gac cgc tat ccc gtg gaa	Gly Ile Val Arg Leu Arg Asp Gly Leu Arg Asp Arg Tyr Pro Val Glu	300	305	310	1446		
gat tat ctg gac ctc ttt gac ctg gcc gca cac cag atc cat caa gga	Asp Tyr Leu Asp Leu Phe Asp Leu Ala Ala His Gln Ile His Gln Gly	315	320	325	1494		
ctg cag gcc agc tct gca aag gag cag agc aag atg aat aca atc att	Leu Gln Ala Ser Ser Ala Lys Glu Gln Ser Lys Met Asn Thr Ile Ile	330	335	340	1542		
atc ggg gag gcc agt gcc tcc acg atg tgg gat aat aat gcc tgg acg	Ile Gly Glu Ala Ser Ala Ser Thr Met Trp Asp Asn Asn Ala Trp Thr	345	350	355	1590		
ttc ttc tac gac aac agc acg gat ggc gag cca ccg ttt ctg acg cag	Phe Phe Tyr Asp Asn Ser Thr Asp Gly Glu Pro Pro Phe Leu Thr Gln	360	365	370	1638		
gac ttc atc cac gcc ttt cag cca aat gcc aga ctg att gtc atg ctc	Asp Phe Ile His Ala Phe Gln Pro Asn Ala Arg Leu Ile Val Met Leu	380	385	390	1686		
agg gac cct gtg gag agg ttg tac tca gac tat ctc tac ttt gca agt	Arg Asp Pro Val Glu Arg Leu Tyr Ser Asp Tyr Leu Tyr Phe Ala Ser	395	400	405	1734		
tcg aat aaa tcc gcg gac gac ttc cat gag aaa gtg aca gaa gca ctg	Ser Asn Lys Ser Ala Asp Asp Phe His Glu Lys Val Thr Glu Ala Leu	410	415	420	1782		
cag ctg ttt gaa aat tgc atg ctt gat tat tca ctg cgc gcc tgc gtc	Gln Leu Phe Glu Asn Cys Met Leu Asp Tyr Ser Leu Arg Ala Cys Val	425	430	435	1830		
tac aac aac acc ctc aac aac gcc atg cct gtg agg ctc cag gtt ggg	Tyr Asn Asn Thr Leu Asn Asn Ala Met Pro Val Arg Leu Gln Val Gly	440	445	450	1878		
ctc tat gct gtg tac ctt ctg gac tgg ctc agc gtt ttt gac aag caa	Leu Tyr Ala Val Tyr Leu Leu Asp Trp Leu Ser Val Phe Asp Lys Gln	460	465	470	1926		
cag ttt ctc att ctt cgc ctg gaa gat cat gca tcc aac gtc aag tac	Gln Phe Leu Ile Leu Arg Leu Glu Asp His Ala Ser Asn Val Lys Tyr	475	480	485	1974		
acc atg cac aag gtc ttc cag ttt ctg aac cta ggg ccc tta agt gag	Thr Met His Lys Val Phe Gln Phe Leu Asn Leu Gly Pro Leu Ser Glu	490	495	500	2022		
aag cag gag gct ttg atg acc aag agc ccc gca tcc aat gca cgg cgt	Lys Gln Glu Ala Leu Met Thr Lys Ser Pro Ala Ser Asn Ala Arg Arg	505	510	515	2070		
ccc gag gac cgg aac ctg ggg ccc atg tgg ccc atc aca cag aag att	Pro Glu Asp Arg Asn Leu Gly Pro Met Trp Pro Ile Thr Gln Lys Ile	520	525	530	2118		
ctg cgg gat ttc tac agg ccc ttc aac gct agg ctg gcg cag gtc ctc	Leu Arg Asp Phe Tyr Arg Pro Phe Asn Ala Arg Leu Ala Gln Val Leu	540	545	550	2166		
gcg gat gag gcg ttt gcg tgg aag acg acg tga gagctgaatt gttgctgcac	Ala Asp Glu Ala Phe Ala Trp Lys Thr Thr	555	560		2219		
gtgctggggc cgccaatgcc gtcacatca ggattttaca aatctctttg cggggaactg					2279		

tttcactcat	ggtatgaaa	accccaggac	tctgccactc	taggcacaca	tgaattataa	2339
ccatttttga	atttccttcg	tgatgttcga	gagctcagca	atggaccctt	cacagagctc	2399
ctctatccga	ggccattgga	gaccccagtt	tctcaagaat	tcagctctgc	tctgagcgtc	2459
ctggagcttg	gggatgcagc	cagctggcct	gcactgggtg	tggagagAAC	acctagggaa	2519
ggcagcctgg	ccctgcccgc	ctccgccttc	tggagagcct	ctgggttctg	agtcagcaag	2579
ccagaggcca	tgccacaggc	ctggctggaa	cttacacttc	acgttccctt	tttttcccc	2639
tagagatggg	gtctcgccgt	gttgacacaga	ctgtctgtat	tcaatggcta	tcttcacagg	2699
tgtgatcata	ccacattcac	ttctgaaaca	ctcttggtgc	gatcgctaac	ctcactggga	2759
cagagaaccg	cagtctttcg	agaatggagg	ctcttcattt	tttttttctc	ctttactcca	2819
aactcagccc	tccagtttct	tcagatgtaa	accctgttaa	cgctactgtt	tccaaaagga	2879
aaaaaataag	tcagtttttg	gcagcacctt	catctttctg	acctcctcct	attctgtcct	2939
tgtggactta	tgtttaacat	agaaaatgaa	tgcgttttaa	acaaaaccac	tttctgcatt	2999
taaccagtcc	tggtctcttc	tctgctgcct	cttcatacgt	tttctcaaga	acttcagttt	3059
ataattggaa	gagaaatttt	tgctgttaat	gccagaatga	gcaacctcaa	ggaattgaac	3119
acttcttggg	aaatctaggt	aattcaagcc	ctcatcaggt	ttacaagatc	atcagagaaa	3179
cagaggattt	taatttttag	ttctggccgg	ctacaggctc	catttctctg	ccttccccatt	3239
ggaaatagtt	tatttccaca	ttctccactg	cgtgtgggtc	aagtccctca	cccagcaagg	3299
gactatagat	actcgtgtcc	caattccaaa	acacaatgca	caagctgaac	ttgggctgaa	3359
cgtggcgtgt	tgagatttgg	aatgagggtt	ctaagagccg	tggtcttcat	ggaattttcc	3419
aggccacttg	gcagcttggt	ttaccgatgg	atgggctaga	gatcttgtcg	tttcttggaa	3479
gtcacaggga	agattgaaga	gaacgcctga	gcaccccttg	caacagccca	ggtgggacct	3539
ggatgaagct	ttgcactcaa	gtattgtcaa	gggaagcttc	ctgtgaacca	aagtctctcag	3599
gccaagggtct	cgcccaccaa	agccagaaag	tgcaagcacc	cgtctacca	gctctaactt	3659
gtatgtgtga	gacagaccag	gcttcggggg	taggaggatc	tgcagttgtt	cagccgtctt	3719
tctgctggtg	ttgtctttct	gccatcagag	aagggaacaca	cagcccgttc	gaagggtgtc	3779
agagggctct	gagcgccagg	atggccaggg	ctgtttttgc	tactgaagga	gcgtgtgtcc	3839
tgaactccca	cttgcaggga	cagtccccac	cttctctata	gccggcactg	ggagcagccg	3899
ccagcaggga	aatctggcct	gagcacaagg	atgcttttagg	gagagatcac	ttcagtgtgt	3959
gtgtatatatt	atttgcagta	cagtgcgcgc	gtgtgtgtgt	gtgtacgcgc	acgtgtgggt	4019
gagtgcgtct	tctgagtggg	ttctgttcag	ttgctaataa	ggctcctccg	ctctggacac	4079
aaccctttta	tagattaatt	tctctgccaa	ttaacttgtc	attttcagta	catattttac	4139
tattccacac	caaccataat	tacaacaagg	gatttttctt	atgcactcct	atgcatgtga	4199
ataacatgtg	gtgtaattct	gcttcttaca	gaagtattac	tgaaggattt	atttccaata	4259
ttatttggtt	tattatgcgg	atctttttta	tatatgcagt	cccatccctt	ctgtgccact	4319
caatgccatc	cagacatggt	ttttccctcc	aggggccttt	ctctccagag	ggcacttcgg	4379

ctgcctctgc ttctctcat tcgaggcccg gctcttgctg acagaatagg ttccgttctg 4439
ggcgggtggtt ctcgagcctg ccattcaaaa ccaaagcaaa ttggagcatt tctcacaaca 4499
tggatttgaa gttccttttt gttctcaaaa gttgtgaccg tgttaaattg tactccctta 4559
gtcctgtaag gtatgttaag tgaatcgagc ttacgctgta cttttattaa tatttaacat 4619
aattaaagat ggaccataa gagtgacgcc tgtggagcgc gtgctcttcc tctgcagcca 4679
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaa 4713

<210> 19
<211> 561
<212> PRT
<213> Homo sapiens

<400> 19

Met Arg His Cys Ile Asn Cys Cys Ile Gln Leu Leu Pro Asp Gly Ala
1 5 10 15

His Lys Gln Gln Val Asn Cys Gln Gly Gly Pro His His Gly His Gln
20 25 30

Ala Cys Pro Thr Cys Lys Gly Glu Asn Lys Ile Leu Phe Arg Val Asp
35 40 45

Ser Lys Gln Met Asn Leu Leu Ala Val Leu Glu Val Arg Thr Glu Gly
50 55 60

Asn Glu Asn Trp Gly Gly Phe Leu Arg Phe Lys Lys Gly Lys Arg Cys
65 70 75 80

Ser Leu Val Phe Gly Leu Ile Ile Met Thr Leu Val Met Ala Ser Tyr
85 90 95

Ile Leu Ser Gly Ala His Gln Glu Leu Leu Ile Ser Ser Pro Phe His
100 105 110

Tyr Gly Gly Phe Pro Ser Asn Pro Ser Leu Met Asp Ser Glu Asn Pro
115 120 125

Ser Asp Thr Lys Glu His His His Gln Ser Ser Val Asn Asn Ile Ser
130 135 140

Tyr Met Lys Asp Tyr Pro Ser Ile Lys Leu Ile Ile Asn Ser Ile Thr
145 150 155 160

Thr Arg Ile Glu Phe Thr Thr Arg Gln Leu Pro Asp Leu Glu Asp Leu
165 170 175

Lys Lys Gln Glu Leu His Met Phe Ser Val Ile Pro Asn Lys Phe Leu
180 185 190

Pro Asn Ser Lys Ser Pro Cys Trp Tyr Glu Glu Phe Ser Gly Gln Asn
195 200 205

Thr Thr Asp Pro Tyr Leu Thr Asn Ser Tyr Val Leu Tyr Ser Lys Arg
 210 215 220
 Phe Arg Ser Thr Phe Asp Ala Leu Arg Lys Ala Phe Trp Gly His Leu
 225 230 235 240
 Ala His Ala His Gly Lys His Phe Arg Leu Arg Cys Leu Pro His Phe
 245 250 255
 Tyr Ile Ile Gly Gln Pro Lys Cys Gly Thr Thr Asp Leu Tyr Asp Arg
 260 265 270
 Leu Arg Leu His Pro Glu Val Lys Phe Ser Ala Ile Lys Glu Pro His
 275 280 285
 Trp Trp Thr Arg Lys Arg Phe Gly Ile Val Arg Leu Arg Asp Gly Leu
 290 295 300
 Arg Asp Arg Tyr Pro Val Glu Asp Tyr Leu Asp Leu Phe Asp Leu Ala
 305 310 315 320
 Ala His Gln Ile His Gln Gly Leu Gln Ala Ser Ser Ala Lys Glu Gln
 325 330 335
 Ser Lys Met Asn Thr Ile Ile Ile Gly Glu Ala Ser Ala Ser Thr Met
 340 345 350
 Trp Asp Asn Asn Ala Trp Thr Phe Phe Tyr Asp Asn Ser Thr Asp Gly
 355 360 365
 Glu Pro Pro Phe Leu Thr Gln Asp Phe Ile His Ala Phe Gln Pro Asn
 370 375 380
 Ala Arg Leu Ile Val Met Leu Arg Asp Pro Val Glu Arg Leu Tyr Ser
 385 390 395 400
 Asp Tyr Leu Tyr Phe Ala Ser Ser Asn Lys Ser Ala Asp Asp Phe His
 405 410 415
 Glu Lys Val Thr Glu Ala Leu Gln Leu Phe Glu Asn Cys Met Leu Asp
 420 425 430
 Tyr Ser Leu Arg Ala Cys Val Tyr Asn Asn Thr Leu Asn Asn Ala Met
 435 440 445
 Pro Val Arg Leu Gln Val Gly Leu Tyr Ala Val Tyr Leu Leu Asp Trp
 450 455 460
 Leu Ser Val Phe Asp Lys Gln Gln Phe Leu Ile Leu Arg Leu Glu Asp
 465 470 475 480
 His Ala Ser Asn Val Lys Tyr Thr Met His Lys Val Phe Gln Phe Leu
 485 490 495

Asn Leu Gly Pro Leu Ser Glu Lys Gln Glu Ala Leu Met Thr Lys Ser
500 505 510

Pro Ala Ser Asn Ala Arg Arg Pro Glu Asp Arg Asn Leu Gly Pro Met
515 520 525

Trp Pro Ile Thr Gln Lys Ile Leu Arg Asp Phe Tyr Arg Pro Phe Asn
530 535 540

Ala Arg Leu Ala Gln Val Leu Ala Asp Glu Ala Phe Ala Trp Lys Thr
545 550 555 560

Thr

<210> 20
<211> 536
<212> DNA
<213> Mus musculus

<400> 20
acaaagccat gaagccggcg ctgctggaag tgatgaggat gaacagaatt tgccggatgg 60
tgctggccac ttgcttcgga tcctttatct tggatcatctt ctattttcaa agtatgttgc 120
accagtcacat gcgagggaac cccttcggtg tggacatctg ctgccggaag ggatcgagaa 180
gtccccctgca ggagctctac aatcccatcc agctggagct atccaacact gccatcctgc 240
accagatgag acgggaccag gtgacagaca cctgccgggc caacagtgcc atgagccgca 300
agcgcaggggt gctgaccccc aacgacctga agcacctggt ggtggatgag gaccacgaac 360
tcacttactg ctatgtgccc aaggtagcgt gcaccaactg gaagaggctc atgatgggtcc 420
tgagtggccg gggcaagtac agcgatccca tggagatccc agccaacgaa gcccacgtgt 480
cggccaacct gaagaccctt aaccagtaca gcatcccaga gatcaaccac cgcttg 536

<210> 21
<211> 660
<212> DNA
<213> Mus musculus

<400> 21
gccaggagtg ggcccagccc agggcagcat gaccaagccg cggtcttcc ggctgtggct 60
ggtactaggg tcggctctca tgatcctttt gatcattgta tattgggaca acgtgggaac 120
cgcccacttc tatctgcaca cgtctctctc caggccacac atcctagaac cccttcccac 180
ccagggattg gtggaggaga acgtgttcac atctgacgtg gatgagtgtt tggatactct 240
ccttagttct gacgcgaagc acaacgacct ttccaggaga aaaactgagc agcccccggc 300
gcccccccc agcaagccag tcttgagcca catggaggag aacgtgagag gctacgactg 360
gtccactcat gatgcccac agaaccctga ccgggacagg cagcaggccg agaggaggag 420
cctgctgaga gacttctgtg ccaacgccag cctggcattc cccaccaagg accgctcttt 480
tgacgacatc cccaactacg aactgaacca cctgatcgtg gacgaccgcc acgggggtcat 540

ctactgctac	gtgccaag	tggcctgcac	caactggaag	cgagtgatga	tcgtgctgag	600
cgagagcctg	ctggaccggg	gcagccccta	ccgagacccc	ctggacatcc	cccgggaaca	660

<210> 22
 <211> 600
 <212> DNA
 <213> Mus musculus

<400> 22						
atgactgtcg	cctgccacgc	gtgccaggca	cagcatggga	agacgctcct	gttgcaggcg	60
gcccttgccg	gtggtggcaa	gtctgggtgc	tgcactcctg	ctcctgtgcg	ccctgctgcc	120
cgggtaacca	caggaaagga	tgccaggac	actgaatggc	agggctcccc	aaaagccctt	180
ttgggggttc	cgacatttga	aaataaagct	ctgggctcca	gctggttcgg	tggagtgagg	240
aagagtcccc	tacagctggt	gcgtgacctg	gaccagatgt	ttggcagctg	tgagctatgg	300
gtagtcagtg	gggagcggca	gaaagtgggt	ccacgctccg	cgatggccga	gggtgcaccag	360
cagcggcgtg	agctgctgcg	ccgggcctgc	agccgccaca	cgcgacgcca	acgcctgctg	420
cagccggagg	acctgcgtca	cgtgctggtg	gacgacgcgc	accggctgct	gtactgctac	480
gtgcctaagg	tggcctgcac	caactggaag	cgtgtgatgc	tggcgttgcg	cggccgtggg	540
gatccaagcg	caatccctgc	gcacgaggcg	catgcgcctg	gcctgctgcc	ctcgctggcc	600

<210> 23
 <211> 600
 <212> DNA
 <213> Mus musculus

<400> 23						
gcgccccctc	tccccatgga	gaaaggactc	gctttgcctc	aggatttccg	ggaccttgta	60
cacagcctaa	agattcgagg	cagatacgtc	ttgttcctgg	catttggtgt	catagttttt	120
atcttcattg	aaaaggaaaa	taaaatcata	tccagggtct	ccgacaagct	gaagcagatc	180
cctcattttg	tggcagatgc	caacagcact	gacccagccc	tgctcttata	ggagaatgca	240
tctctcttgt	ccctgagcga	gttggattcc	accttttccc	atctgcggag	ccgcctgcac	300
aacctgagcc	tgcagctggg	cgtggagcca	gcaatggaga	gccaggaggc	tggggcagag	360
aagccatccc	agcaggctgg	agcagggacc	cggcgccacg	tgcttctcat	ggccaccacc	420
cgcacgggtt	cctcgttcgt	gggcgagttc	ttcaaccagc	agggcaatat	cttctacctc	480
ttcgagccac	tgtggcacat	cgagcgcacc	gtgttcttcc	agcagcgagg	cgccagcgcg	540
gctggttcag	ccttgggtcta	ccgtgatgtc	ctcaagcagt	tgttgctatg	cgacctgtat	600

<210> 24
 <211> 720
 <212> DNA
 <213> Mus musculus

<400> 24						
tggggagagt	gaggattcgg	tgaactatga	agggccggcg	gcggcggcgc	cgagagtatt	60
gcaagttcac	gctgctcttg	gcgctgtaca	cgcttttgct	acttcttgct	ccctctgtac	120
tggacagcca	cagcgagcag	gacaagggca	ggaactgccc	cggcctgcag	cgcagcttgg	180

gtgtgtggag cctggaggcg gcggcgccg gggaacgtga gcagggcgct gaggtgcggt 240
ccctggccga aggaaacccg gatcgatccc ccgggtcccc cggcaacctc agcgccgtcg 300
gtgaggcggt gaccagga aagcaacaca tctatgtgca tgccacctgg cgcaccggct 360
cgtccttctt gggcgaactc ttcaaccagc acccggaagt tttctacttg tacgagccca 420
tgtggcatct gtggcaggca ctgtatccgg gcgacgcgga gagcctgcag ggcgcactaa 480
gagacatgct gcgctccctc ttccgctgtg atttctctgt gctgcgcctg tacgcgcagc 540
ctggggaccc tggggagcga gcaccggact cggccaacct caccacggcc atgcttttcc 600
gctggcggac caacaaggtc atctgctcgc cgctctgtg ccccgccg cgccggggcac 660
gcgcggacgt gggactcgtc gaggacaaag cctgcgaaag tacctgcccg cccgtttcgc 720

<210> 25
<211> 27
<212> RNA
<213> Artificial

<220>
<223> an artificially synthesized sequence

<400> 25
ggagcagagc aaugaaua caucag 27

<210> 26
<211> 27
<212> RNA
<213> Artificial

<220>
<223> an artificially synthesized sequence

<400> 26
gauugauuc aucuugcucu gcuccau 27

<210> 27
<211> 20
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized primer sequence

<400> 27
gtgagttctg ctgcggtcca 20

<210> 28
<211> 21
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized primer sequence

<400> 28
agtccatgct gatgccaga g 21

<210> 29
<211> 19
<212> DNA
<213> Artificial

<220>
 <223> an artificially synthesized primer sequence
 <400> 29
 acccgatggc aacaatgga 19

<210> 30
 <211> 20
 <212> DNA
 <213> Artificial

<220>
 <223> an artificially synthesized primer sequence
 <400> 30
 accagcaggg ccttggtcac 20

<210> 31
 <211> 25
 <212> DNA
 <213> Artificial

<220>
 <223> an artificially synthesized primer sequence
 <400> 31
 catccgtaaa gacctctatg ccaac 25

<210> 32
 <211> 19
 <212> DNA
 <213> Artificial

<220>
 <223> an artificially synthesized primer sequence
 <400> 32
 atggagccac cgatccaca 19

<210> 33
 <211> 23
 <212> DNA
 <213> Artificial

<220>
 <223> an artificially synthesized primer sequence
 <400> 33
 ttctggccaa cgttctagac aac 23

<210> 34
 <211> 21
 <212> DNA
 <213> Artificial

<220>
 <223> an artificially synthesized primer sequence
 <400> 34
 ccagtgggtct tgggtgtgctg a 21

<210> 35
 <211> 25
 <212> DNA

<213> Artificial
 <220>
 <223> an artificially synthesized sequence
 <400> 35
 accccaact cggaacgatg cggct 25

<210> 36
 <211> 24
 <212> DNA
 <213> Artificial
 <220>
 <223> an artificially synthesized sequence
 <400> 36
 tgcattgttct cgtccatcct gctg 24

<210> 37
 <211> 26
 <212> DNA
 <213> Artificial
 <220>
 <223> an artificially synthesized sequence
 <400> 37
 cgccaccgtg tactgtactg tgaagt 26

<210> 38
 <211> 25
 <212> DNA
 <213> Artificial
 <220>
 <223> an artificially synthesized sequence
 <400> 38
 aggctgctcc aactggaaga gggtg 25

<210> 39
 <211> 24
 <212> DNA
 <213> Artificial
 <220>
 <223> an artificially synthesized sequence
 <400> 39
 atatagtatc taggatatat gtag 24

<210> 40
 <211> 25
 <212> DNA
 <213> Artificial
 <220>
 <223> an artificially synthesized sequence
 <400> 40
 gaagtaccaa aagctggctg ctcta 25

<210> 41
 <211> 26

<212> DNA
 <213> Artificial
 <220>
 <223> an artificially synthesized sequence
 <400> 41
 ttctatcact tggactatgt gatgtt 26

<210> 42
 <211> 25
 <212> DNA
 <213> Artificial
 <220>
 <223> an artificially synthesized sequence
 <400> 42
 tacacaactc cacatttgta atttg 25

<210> 43
 <211> 24
 <212> DNA
 <213> Artificial
 <220>
 <223> an artificially synthesized sequence
 <400> 43
 ccagaagcca agctcattgt tatg 24

<210> 44
 <211> 25
 <212> DNA
 <213> Artificial
 <220>
 <223> an artificially synthesized sequence
 <400> 44
 ctgtggagag gttgtactca gacta 25

<210> 45
 <211> 26
 <212> DNA
 <213> Artificial
 <220>
 <223> an artificially synthesized sequence
 <400> 45
 atttgcctgg aagacaacgt gagagc 26

<210> 46
 <211> 27
 <212> DNA
 <213> Artificial
 <220>
 <223> an artificially synthesized sequence
 <400> 46
 gtcccttctg cagaagctgg gccact 27

<210> 47

<211> 25
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 47
 gcgccccctc tccccatgga gaaag 25

<210> 48
 <211> 25
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 48
 gctttgcctc aggatttccg ggacc 25

<210> 49
 <211> 27
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 49
 ggttcagcct tggctaccg tgatgtc 27

<210> 50
 <211> 25
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 50
 gcagttgttg ctatgcgacc tgtat 25

<210> 51
 <211> 23
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized primer sequence

 <400> 51
 tgttcctggc atttgtggtc ata 23

<210> 52
 <211> 21
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized primer sequence

 <400> 52
 ccaactcgct cagggacaag a 21

<210> 53
 <211> 25
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized primer sequence

 <400> 53
 gtgtggagca acatgtggaa ctcta 25

<210> 54
 <211> 20
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized primer sequence

 <400> 54
 ttggttcagc cactgccgta 20

<210> 55
 <211> 50
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 55
 acaaagccat gaagccggcg ctgctggaag tgatgaggat gaacagaatt 50

<210> 56
 <211> 26
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 56
 caacctgaag acccttaacc agtaca 26

<210> 57
 <211> 26
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 57
 gcatcccaga gatcaaccac cgcttg 26

<210> 58
 <211> 25
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 58
 gccaggagtg ggcccagccc agggc 25

<210> 59
 <211> 27
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 59
 atgaccaagc cgcggtctt ccggctg 27

<210> 60
 <211> 28
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 60
 agagcctgct ggaccggggc agccccta 28

<210> 61
 <211> 28
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 61
 gagaccccct ggacatcccc cggaaca 28

<210> 62
 <211> 26
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 62
 atgactgtcg cctgccacgc gtgcca 26

<210> 63
 <211> 26
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 63
 cagcatggga agacgctcct gttgca 26

<210> 64
 <211> 28
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 64
 tccaagcgca atccctgcgc acgaggcg 28

<210> 65
 <211> 25
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized sequence

 <400> 65
 gcctggcctg ctgccctcgc tggcc 25

<210> 66
 <211> 18
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized primer sequence

 <400> 66
 ctgccaagta tgacatca 18

<210> 67
 <211> 21
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized primer sequence

 <400> 67
 tactccttgg aggccatgta g 21

<210> 68
 <211> 20
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized primer sequence

 <400> 68
 gtggatgagg accacgaact 20

<210> 69
 <211> 20
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized primer sequence

 <400> 69
 cttttcaagc ggtggttgat 20

<210> 70
 <211> 20
 <212> DNA
 <213> Artificial

 <220>
 <223> an artificially synthesized primer sequence

 <400> 70

acctcctaga cccacacacg 20

<210> 71
<211> 20
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized primer sequence

<400> 71
ggatgttggc aaaccagtct 20

<210> 72
<211> 20
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized primer sequence

<400> 72
atgagccctt caacgaacac 20

<210> 73
<211> 20
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized primer sequence

<400> 73
tggtagaagg ggctgatgtc 20

<210> 74
<211> 19
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized primer sequence

<400> 74
ttccaggctt tgggcatca 19

<210> 75
<211> 24
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized primer sequence

<400> 75
atgttcagca tggtcagcag tgtg 24

<210> 76
<211> 25
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized primer sequence

<400> 76
catccgtaaa gacctctatg ccaac 25

<210> 77
<211> 19
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized primer sequence

<400> 77
atggagccac cgatccaca 19

<210> 78
<211> 20
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized primer sequence

<400> 78
tgtgtggaca cacctcccta 20

<210> 79
<211> 20
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized primer sequence

<400> 79
cttcaaaggt ccccttcctc 20

<210> 80
<211> 20
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized primer sequence

<400> 80
cagcttgagc catttcaaca 20

<210> 81
<211> 20
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized primer sequence

<400> 81
gggtgaggcc ttaggaaac 20

<210> 82
<211> 25
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized sequence

<400> 82
gcgccccctc tccccatgga gaaag 25

<210> 83
<211> 25
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized sequence

<400> 83
gctttgcctc aggatttcg ggacc 25

<210> 84
<211> 27
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized sequence

<400> 84
ggttcagcct tggctaccg tgatgtc 27

<210> 85
<211> 25
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized sequence

<400> 85
gcagttgttg ctatgcgacc tgtat 25

<210> 86
<211> 24
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized sequence

<400> 86
tggggagagt gaggattcgg tgaa 24

<210> 87
<211> 27
<212> DNA
<213> Artificial

<220>
<223> an artificially synthesized sequence

<400> 87
cggacgtggg actcgtcgag gacaaag 27

<210> 88
<211> 26
<212> DNA
<213> Artificial

<220>

<223> an artificially synthesized sequence
 <400> 88
 cgaaagtacc tgcccgccg tttcgc 26

<210> 89
 <211> 21
 <212> DNA
 <213> Artificial

<220>
 <223> an artificially synthesized primer sequence
 <400> 89
 ctgtgagccc aagtgtgtgg a 21

<210> 90
 <211> 22
 <212> DNA
 <213> Artificial

<220>
 <223> an artificially synthesized primer sequence
 <400> 90
 gtctcgaaac atggcaacag ga 22

<210> 91
 <211> 27
 <212> RNA
 <213> Artificial

<220>
 <223> An artificially synthesized siRNA sequence
 <400> 91
 gcagcccagc aagaugaaua agaucag 27

<210> 92
 <211> 27
 <212> RNA
 <213> Artificial

<220>
 <223> An artificially synthesized siRNA sequence
 <400> 92
 gaucuauuc aucuugcugg gcugcau 27

<210> 93
 <211> 25
 <212> DNA
 <213> Artificial

<220>
 <223> An artificially synthesized primer sequence
 <400> 93
 catccgtaaa gacctctatg ccaac 25

<210> 94
 <211> 19
 <212> DNA
 <213> Artificial

<220>
 <223> An artificially synthesized primer sequence
 <400> 94
 atggagccac cgatccaca 19

<210> 95
 <211> 23
 <212> DNA
 <213> Artificial

<220>
 <223> An artificially synthesized primer sequence
 <400> 95
 caggagggag aacagaaact cca 23

<210> 96
 <211> 19
 <212> DNA
 <213> Artificial

<220>
 <223> An artificially synthesized primer sequence
 <400> 96
 cctggttggc tgcttgctt 19

<210> 97
 <211> 20
 <212> DNA
 <213> Artificial

<220>
 <223> An artificially synthesized primer sequence
 <400> 97
 ctgccctggc tatggtcaca 20

<210> 98
 <211> 23
 <212> DNA
 <213> Artificial

<220>
 <223> An artificially synthesized primer sequence
 <400> 98
 agacaggtag ttgggtcggt tca 23