

METHODS OF TREATING CANCER WITH ANTI-TIM-3 ANTIBODIES

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] The present application claims benefit of U.S. Provisional Application No. 62/444,354, filed January 9, 2017, and U.S. Provisional Application No. 62/582,272, filed November 6, 2017, each of which is incorporated by reference in its entirety.

SEQUENCE LISTING

[0002] The present specification makes reference to a Sequence Listing provided in electronic form as an ASCII.txt file named “TSR-002 SEQ LIST_ST25” that was generated on January 8, 2018, and is 39,647 bytes in size.

BACKGROUND

[0003] Cancer is a serious public health problem, with about 600,920 people in the United States of America expected to die of cancer in 2017 alone according to the American Cancer Society, Cancer Facts & Figures 2017 (<https://www.cancer.org/research/cancer-facts-statistics/all-cancer-facts-figures/cancer-facts-figures-2017.html>). Accordingly, there continues to be a need for effective therapies to treat cancer patients.

SUMMARY

[0004] The present invention encompasses a recognition that certain dosage regimens for agents that are capable of inhibiting T Cell Immunoglobulin and Mucin Domain-3 (TIM-3) signaling (e.g., anti-TIM-3 antibody agents) are useful for treating disorders such as cancer.

[0005] In some embodiments, the present disclosure provides methods of treating disorders such as cancer that include administering compositions that deliver particular TIM-3 inhibitors (e.g., anti-TIM-3 antibody agents) according to dosing regimens that may achieve clinical benefit in at least some patients.

[0006] In embodiments, a TIM-3 inhibitor is a TIM-3 binding agent. In embodiments, a TIM-3 binding agent is an antibody, an antibody conjugate, or an antigen-binding fragment thereof. In embodiments, a TIM-3 binding agent is an antibody agent (i.e., an anti-TIM-3 antibody agent).

[0007] In embodiments, an anti-TIM-3 antibody agent comprises a heavy chain variable region with one or more CDR sequences selected from SEQ ID NOs: 21, 22, and 23 and/or a light chain variable region with one or more CDR sequences selected from SEQ ID NOs: 24, 25, and 26. In embodiments, an anti-TIM-3 antibody agent comprises a heavy chain variable region with two or three CDR sequences selected from SEQ ID NOs: 21, 22, and 23 and/or a light chain variable region with two or three CDR sequences selected from SEQ ID NOs: 24, 25, and 26. In embodiments, an anti-TIM-3 antibody agent comprises a heavy chain variable region with three CDR sequences of SEQ ID NOs: 21, 22, and 23 and/or a light chain variable region with three CDR sequences of SEQ ID NOs: 24, 25, and 26. In embodiments, an anti-TIM-3 antibody agent comprises a heavy chain variable region with three CDR sequences of SEQ ID NOs: 21, 22, and 23 and a light chain variable region with three CDR sequences of SEQ ID NOs: 24, 25, and 26.

[0008] In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain variable domain whose amino acid sequence having at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NO:1 or SEQ ID NO:7. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin light chain variable domain whose amino acid sequence having at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NO:2 or SEQ ID NO:8. In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain variable domain whose amino acid sequence comprises SEQ ID NO: 1 or SEQ ID NO: 7 and/or an immunoglobulin light chain variable domain whose amino acid sequence comprises SEQ ID NO: 2 or SEQ ID NO: 8. In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain variable domain whose amino acid sequence comprises SEQ ID NO: 1 or SEQ ID NO: 7 and an immunoglobulin light chain variable domain whose amino acid sequence comprises SEQ ID NO: 2 or SEQ ID NO: 8.

[0009] In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain polypeptide comprising an amino acid sequence having about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NO:3. In

some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin light chain polypeptide comprising an amino acid sequence having about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NO:4. In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain whose amino acid sequence comprises SEQ ID NO: 3 and/or an immunoglobulin light chain whose amino acid sequence comprises SEQ ID NO: 4. In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain whose amino acid sequence comprises SEQ ID NO: 3 and an immunoglobulin light chain whose amino acid sequence comprises SEQ ID NO: 4.

[0010] The present disclosure provides, in some embodiments, methods of treating a disorder in a subject that is responsive to T Cell Immunoglobulin and Mucin Protein 3 (TIM-3) inhibition comprising administering a therapeutically effective dose of an agent that is capable of inhibiting TIM-3 signaling. In embodiments, a therapeutically effective dose is about 1, 3 or 10 mg/kg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is about 100 - 1500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg; a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; a flat dose about 900 mg; a flat dose about 1000 mg; a flat dose about 1100 mg; a flat dose about 1200 mg; a flat dose about 1300 mg; a flat dose about 1400 mg; or a flat dose about 1500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is about 1 mg/kg. In embodiments, a therapeutically effective dose is about 3 mg/kg. In embodiments, a therapeutically effective dose is about 10 mg/kg. In embodiments, a therapeutically effective dose is a flat dose about 100 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 300 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 1200 mg of a TIM-3 inhibitor. In embodiments, a TIM-3 inhibitor is any anti-TIM-3 antibody agent described herein.

[0011] The present disclosure provides, in some embodiments, methods of increasing T cell activation or T cell effector function in a subject that is responsive to T Cell Immunoglobulin and Mucin Protein 3 (TIM-3) inhibition comprising administering a therapeutically effective dose of an agent that is capable of inhibiting TIM-3 signaling. In embodiments, a therapeutically effective dose is about 1, 3 or 10 mg/kg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is about 100 - 1500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically

effective dose is a flat dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg; a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; a flat dose about 900 mg; a flat dose about 1000 mg; a flat dose about 1100 mg; a flat dose about 1200 mg; a flat dose about 1300 mg; a flat dose about 1400 mg; or a flat dose about 1500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is about 1 mg/kg. In embodiments, a therapeutically effective dose is about 3 mg/kg. In embodiments, a therapeutically effective dose is about 10 mg/kg. In embodiments, a therapeutically effective dose is a flat dose about 100 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 300 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 900 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 1200 mg of a TIM-3 inhibitor. In embodiments, a TIM-3 inhibitor is any anti-TIM-3 antibody agent described herein.

[0012] The present disclosure provides, in some embodiments, methods of reducing tumors or inhibiting the growth of tumor cells in a subject that is responsive to T Cell Immunoglobulin and Mucin Protein 3 (TIM-3) inhibition comprising administering a therapeutically effective dose of an agent that is capable of inhibiting TIM-3 signaling. In embodiments, a therapeutically effective dose is about 1, 3 or 10 mg/kg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is about 100 - 1500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg; a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; a flat dose about 900 mg; a flat dose about 1000 mg; a flat dose about 1100 mg; a flat dose about 1200 mg; a flat dose about 1300 mg; a flat dose about 1400 mg; or a flat dose about 1500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is about 1 mg/kg. In embodiments, a therapeutically effective dose is about 3 mg/kg. In embodiments, a therapeutically effective dose is about 10 mg/kg. In embodiments, a therapeutically effective dose is a flat dose about 100 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 300 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 900 mg of a TIM-3

inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 1200 mg of a TIM-3 inhibitor. In embodiments, a TIM-3 inhibitor is any anti-TIM-3 antibody agent described herein.

[0013] The present disclosure provides, in some embodiments, methods of inducing an immune response in a subject that is responsive to T Cell Immunoglobulin and Mucin Protein 3 (TIM-3) inhibition comprising administering a therapeutically effective dose of an agent that is capable of inhibiting TIM-3 signaling. In embodiments, a therapeutically effective dose is about 1, 3 or 10 mg/kg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is about 100 - 1500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg; a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; a flat dose about 900 mg; a flat dose about 1000 mg; a flat dose about 1100 mg; a flat dose about 1200 mg; a flat dose about 1300 mg; a flat dose about 1400 mg; or a flat dose about 1500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is about 1 mg/kg. In embodiments, a therapeutically effective dose is about 3 mg/kg. In embodiments, a therapeutically effective dose is about 10 mg/kg. In embodiments, a therapeutically effective dose is a flat dose about 100 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 300 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 900 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 1200 mg of a TIM-3 inhibitor. In embodiments, a TIM-3 inhibitor is any anti-TIM-3 antibody agent described herein.

[0014] The present disclosure provides, in some embodiments, methods of enhancing an immune response or increasing the activity of an immune cell in a subject that is responsive to T Cell Immunoglobulin and Mucin Protein 3 (TIM-3) inhibition comprising administering a therapeutically effective dose of an agent that is capable of inhibiting TIM-3 signaling. In embodiments, an immune response is a humoral or cell mediated immune response. In embodiments, an immune response is a CD4 or CD8 T cell response. In embodiments, an immune response is a B cell response. In embodiments, a therapeutically effective dose is about 1, 3 or 10 mg/kg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is about 100 - 1500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat

dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg; a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; a flat dose about 900 mg; a flat dose about 1000 mg; a flat dose about 1100 mg; a flat dose about 1200 mg; a flat dose about 1300 mg; a flat dose about 1400 mg; or a flat dose about 1500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is about 1 mg/kg. In embodiments, a therapeutically effective dose is about 3 mg/kg. In embodiments, a therapeutically effective dose is about 10 mg/kg. In embodiments, a therapeutically effective dose is a flat dose about 100 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 300 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 900 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 1200 mg of a TIM-3 inhibitor. In embodiments, a TIM-3 inhibitor is any anti-TIM-3 antibody agent described herein.

[0015] The present disclosure provides, in some embodiments, methods of treating cancer that comprises administering to a patient in need of treatment an anti-T Cell Immunoglobulin and Mucin Domain-3 (TIM-3) at a therapeutically effective dose. In embodiments, a therapeutically effective dose is about 1, 3 or 10 mg/kg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is about 100 - 1500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg; a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; a flat dose about 900 mg; a flat dose about 1000 mg; a flat dose about 1100 mg; a flat dose about 1200 mg; a flat dose about 1300 mg; a flat dose about 1400 mg; or a flat dose about 1500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is about 1 mg/kg. In embodiments, a therapeutically effective dose is about 3 mg/kg. In embodiments, a therapeutically effective dose is about 10 mg/kg. In embodiments, a therapeutically effective dose is a flat dose about 100 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 300 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 900 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 1200 mg of a

TIM-3 inhibitor. In embodiments, a TIM-3 inhibitor is any anti-TIM-3 antibody agent described herein.

[0016] The present disclosure provides, in some embodiments, methods of treating cancer comprising administering to a patient in need of treatment an anti-T Cell Immunoglobulin and Mucin Domain-3 (TIM-3) antibody at a therapeutically effective dose at an administration interval for a period sufficient to achieve clinical benefit. In embodiments, an anti-TIM-3 antibody comprises a heavy chain comprising three CDRs that have sequences of SEQ ID NOs: 21, 22, or 23; and/or a light chain comprising three CDRs that have sequences of SEQ ID NOs: 24, 25, or 26. In embodiments, an anti-TIM-3 antibody comprises an immunoglobulin heavy chain variable domain comprising SEQ ID NO: 1 or SEQ ID NO: 7 and an immunoglobulin light chain variable domain comprising SEQ ID NO: 2 or SEQ ID NO: 8. In embodiments, an anti-TIM-3 antibody comprises a heavy chain polypeptide comprising SEQ ID NO: 3 and a light chain polypeptide comprising SEQ ID NO: 4. In embodiments, a therapeutically effective dose is about 1, 3 or 10 mg/kg of an anti-TIM-3 antibody. In embodiments, a therapeutically effective dose is about 100 - 1500 mg of an anti-TIM-3 antibody. In embodiments, a therapeutically effective dose is a flat dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg; a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; a flat dose about 900 mg; a flat dose about 1000 mg; a flat dose about 1100 mg; a flat dose about 1200 mg; a flat dose about 1300 mg; a flat dose about 1400 mg; or a flat dose about 1500 mg of an anti-TIM-3 antibody. In embodiments, a therapeutically effective dose is about 1 mg/kg of an anti-TIM-3 antibody. In embodiments, a therapeutically effective dose is about 3 mg/kg of an anti-TIM-3 antibody. In embodiments, a therapeutically effective dose is about 10 mg/kg of an anti-TIM-3 antibody. In embodiments, a therapeutically effective dose is a flat dose about 100 mg of an anti-TIM-3 antibody. In embodiments, a therapeutically effective dose is a flat dose about 300 mg of an anti-TIM-3 antibody. In embodiments, a therapeutically effective dose is a flat dose about 500 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 900 mg of a TIM-3 inhibitor. In embodiments, a therapeutically effective dose is a flat dose about 1200 mg of an anti-TIM-3 antibody.

[0017] In any of the methods described herein, a therapeutically effective dose is about 1 mg/kg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective

dose is about 3 mg/kg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 5 mg/kg of a TIM-3 inhibitor. In embodiments, a TIM-3 inhibitor is any anti-TIM-3 antibody agent described herein.

[0018] In any of the methods described herein, a therapeutically effective dose is about 100 mg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 200 mg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 300 mg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 400 mg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 500 mg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 600 mg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 700 mg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 800 mg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 900 mg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 1000 mg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 1100 mg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 1200 mg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 1300 mg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 1400 mg of a TIM-3 inhibitor. In any of the methods described herein, a therapeutically effective dose is about 1500 mg of a TIM-3 inhibitor. In embodiments, a TIM-3 inhibitor is any anti-TIM-3 antibody agent described herein.

[0019] In embodiments, a TIM-3 inhibitor is administered at an administration interval (or treatment cycle) of once a week (Q1W), once every 2 weeks (Q2W), once every 3 weeks (Q3W), once every 4 weeks (Q4W), once every 5 weeks (Q5W), or once every 6 weeks (Q6W). In embodiments, a TIM-3 inhibitor is administered for a period of at least about 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 weeks, or more. In embodiments, a TIM-3 inhibitor is administered at an administration interval (or treatment cycle) of once a week (Q1W). In embodiments, a TIM-3 inhibitor is administered at an administration interval (or treatment cycle) of once every 2 weeks (Q2W). In embodiments, a TIM-3 inhibitor is administered at an administration interval (or treatment cycle) of once every three weeks (Q3W). In embodiments,

a TIM-3 inhibitor is administered at an administration interval (or treatment cycle) of once every 4 weeks (Q4W). In embodiments, a TIM-3 inhibitor is administered at an administration interval (or treatment cycle) of once every 5 weeks (Q5W). In embodiments, a TIM-3 inhibitor is administered at an administration interval (or treatment cycle) of once every 6 weeks (Q6W).

[0020] In embodiments, a TIM-3 inhibitor (e.g., a therapeutically effective dose of about 100 mg, 300 mg, 500 mg, or 900 mg) is administered at an administration interval (or treatment cycle) of once every 3 weeks. In embodiments, a therapeutically effective dose of about 100 mg is administered at an administration interval (or treatment cycle) of once every 3 weeks. In embodiments, a therapeutically effective dose of about 300 mg is administered at an administration interval (or treatment cycle) of once every 3 weeks. In embodiments, a therapeutically effective dose of about 500 mg is administered at an administration interval (or treatment cycle) of once every 3 weeks. In embodiments, a therapeutically effective dose of about 900 mg is administered at an administration interval (or treatment cycle) of once every 3 weeks.

[0021] In embodiments, a TIM-3 inhibitor is administered on the first day of a treatment cycle or within 1, 2, or 3 days of the first day of a treatment cycle. In embodiments, a TIM-3 inhibitor is any anti-TIM-3 antibody agent described herein.

[0022] In embodiments, a TIM-3 inhibitor described herein is administered according to dosing regimens demonstrated to achieve a clinical benefit in some patients (for example, according to a regimen as determined by a physician, including dosing modifications). In embodiments, a TIM-3 inhibitor described herein is administered until treatment is discontinued due to, e.g., disease progression or an adverse reaction or as determined by a physician. In embodiments, a clinical benefit is stable disease (“SD”), a partial response (“PR”) and/or a complete response (“CR”). In embodiments, a clinical benefit is stable disease (“SD”). In embodiments, a clinical benefit is a partial response (“PR”). In embodiments, a clinical benefit is a complete response (“CR”). In embodiments, PR or CR is determined in accordance with Response Evaluation Criteria in Solid Tumors (RECIST). In embodiments, a TIM-3 inhibitor is administered for a longer period to maintain clinical benefit. In embodiments, a TIM-3 inhibitor is any anti-TIM-3 antibody agent described herein.

[0023] In embodiments, a TIM-3 inhibitor is administered periodically to a subject at a dose of about 100 mg, about 300 mg, or about 1200 mg. In embodiments, a TIM-3 inhibitor is

administered periodically to a subject at a dose of about 100 mg (e.g., once every three weeks (Q3W) and/or for 2, 3, 4, 5, 6, or more cycles). In embodiments, a TIM-3 inhibitor is administered periodically to a subject at a dose of about 300 mg (e.g., once every three weeks (Q3W) and/or for 2, 3, 4, 5, 6, or more cycles). In embodiments, a TIM-3 inhibitor is administered periodically to a subject at a dose of about 1200 mg (e.g., once every three weeks (Q3W) and/or for 2, 3, 4, 5, 6, or more cycles). In embodiments, a TIM-3 inhibitor is any anti-TIM-3 antibody agent described herein.

[0024] In embodiments, a subject has been further administered or will be administered a further therapeutic agent, such that the subject receives a TIM-3 inhibitor (e.g., any anti-TIM-3 antibody agent described herein) and a further therapeutic agent (e.g., one, two, three, four, or more further therapeutic agents).

[0025] In embodiments, a subject has been further administered or will be administered an immune checkpoint inhibitor, such that the subject receives a TIM-3 inhibitor (e.g., any anti-TIM-3 antibody agent described herein) and an immune checkpoint inhibitor. That is, a subject can be administered a TIM-3 inhibitor in combination with at least one immune checkpoint inhibitor.

[0026] In embodiments, a checkpoint inhibitor is an agent capable of inhibiting any of the following: PD-1 (e.g., inhibition via anti-PD-1, anti-PD-L1, or anti-PD-L2 therapies), CTLA-4, TIM-3, TIGIT, LAGs (e.g., LAG-3), CEACAM (e.g., CEACAM-1, -3 and/or -5), VISTA, BTLA, LAIR1, CD160, 2B4, CD80, CD86, B7-H3 (CD276), B7-H4 (VTCN1), HVEM (TNFRSF14 or CD270), KIR, A2aR, MHC class I, MHC class II, GALS, adenosine, TGFR (e.g., TGFR beta), B7-H1, B7-H4 (VTCN1), OX-40, CD137, CD40, IDO, or CSF-1R. In embodiments, a checkpoint inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, or a toxin. In embodiments, a checkpoint inhibitor is an antibody, an antibody conjugate, or an antigen-binding fragment thereof.

[0027] In embodiments, an immune checkpoint inhibitor is an agent that inhibits programmed death-1 protein (PD-1) signaling, cytotoxic T-lymphocyte-associated protein 4 (CTLA-4), lymphocyte activation gene-3 (LAG-3), T cell immunoglobulin and ITIM domain (TIGIT), indoleamine 2,3-dioxygenase (IDO), or colony stimulating factor 1 receptor (CSF1R).

[0028] In embodiments, an immune checkpoint inhibitor is a PD-1 inhibitor. In embodiments, a PD-1 inhibitor is a PD-1 binding agent (e.g., an antibody, an antibody conjugate,

or an antigen-binding fragment thereof). In embodiments, a PD-1 binding agent is nivolumab, pembrolizumab, TSR-042, PDR-001, tislelizumab (BGB-A317), cemiplimab (REGN2810), LY-3300054, JNJ-63723283, MGA012, BI-754091, IBI-308, camrelizumab (HR-301210), BCD-100, JS-001, CX-072, AMP-514 / MEDI-0680, AGEN-2034, CS1001, TSR-042, Sym-021, PF-06801591, LZM009, KN-035, AB122, genolimzumab (CBT-501), AK 104, or GLS-010, or derivatives thereof. In embodiments, a PD-1 inhibitor is a PD-L1 or PD-L2 binding agent (e.g., an antibody, an antibody conjugate, or an antigen-binding fragment thereof). In embodiments, a PD-1 inhibitor is a PD-L1 or PD-L2 binding agent is durvalumab, atezolizumab, avelumab, BGB-A333, SHR-1316, FAZ-053, CK-301, or, PD-L1 millamolecule, or derivatives thereof. In embodiments, PD-1 inhibitor (e.g., TSR-042) is administered to the subject periodically at a dose of about 500 mg or 1000 mg. In embodiments, a PD-1 inhibitor (e.g., TSR-042) is administered to the subject periodically at a dose of about 500 mg or 1000 mg. In embodiments, a PD-1 inhibitor (e.g., TSR-042) is administered to the subject periodically at a dose of about 500 mg. In embodiments, a PD-1 inhibitor (e.g., TSR-042) is administered to the subject once every 3 weeks. In embodiments, a PD-1 inhibitor (e.g., TSR-042) is administered for 2, 3, 4, 5, 6, or more cycles. In embodiments, a PD-1 inhibitor (e.g., TSR-042) is administered for 4 cycles. In embodiments, a PD-1 inhibitor (e.g., TSR-042) is administered to the subject periodically at a dose of about 1000 mg. In embodiments, a PD-1 inhibitor (e.g., TSR-042) is administered to the subject once every 6 weeks. In embodiments, a PD-1 inhibitor (e.g., TSR-042) is administered at a first dose of about 500 mg once every 3 weeks for 4 cycles followed by a second dose of about 1000 mg once every 6 weeks (e.g., until treatment is discontinued).

[0029] In embodiments, an immune checkpoint inhibitor is a CTLA-4 inhibitor (e.g., an antibody, an antibody conjugate, or an antigen-binding fragment thereof). In embodiments, a CTLA-4 inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, or a toxin. In embodiments, a CTLA-4 inhibitor is a small molecule. In embodiments, a CTLA-4 inhibitor is a CTLA-4 binding agent. In embodiments, a CTLA-4 inhibitor is an antibody, an antibody conjugate, or an antigen-binding fragment thereof. In embodiments, a CTLA-4 inhibitor is ipilimumab (Yervoy), AGEN1884, or tremelimumab.

[0030] In embodiments, an immune checkpoint inhibitor is a LAG-3 inhibitor (e.g., an antibody, an antibody conjugate, or an antigen-binding fragment thereof). In embodiments, a LAG-3 inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a

carbohydrate, a lipid, a metal, or a toxin. In embodiments, a LAG-3 inhibitor is a small molecule. In embodiments, a LAG-3 inhibitor is a LAG-3 binding agent. In embodiments, a LAG-3 inhibitor is an antibody, an antibody conjugate, or an antigen-binding fragment thereof. In embodiments, a LAG-3 inhibitor is a IMP321, BMS-986016, GSK2831781, Novartis LAG525, or a LAG-3 inhibitor described in WO 2016/126858, WO 2017/019894, or WO 2015/138920, each of which is hereby incorporated by reference in its entirety.

[0031] In embodiments, an immune checkpoint inhibitor is a TIGIT inhibitor (e.g., an antibody, an antibody conjugate, or an antigen-binding fragment thereof). In embodiments, a TIGIT inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, or a toxin. In embodiments, a TIGIT inhibitor is small molecule. In embodiments, a TIGIT inhibitor is a TIGIT binding agent. In embodiments, a TIGIT inhibitor is an antibody, an antibody conjugate, or an antigen-binding fragment thereof. In embodiments, a TIGIT inhibitor is MTIG7192A, BMS-986207, or OMP-31M32.

[0032] In embodiments, an immune checkpoint inhibitor is an IDO inhibitor. In embodiments, an IDO inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, or a toxin. In embodiments, an IDO inhibitor is small molecule. In embodiments, an IDO inhibitor is an IDO binding agent. In embodiments, an IDO inhibitor is an antibody, an antibody conjugate, or an antigen-binding fragment thereof.

[0033] In embodiments, an immune checkpoint inhibitor is a CSF1R inhibitor. In embodiments, a CSF1R inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, or a toxin. In embodiments, a CSF1R inhibitor is small molecule. In embodiments, a CSF1R inhibitor is a CSF1R binding agent. In embodiments, a CSF1R inhibitor is an antibody, an antibody conjugate, or an antigen-binding fragment thereof.

[0034] In embodiments, a method comprises administering a TIM-3 inhibitor (e.g., any anti-TIM-3 antibody agent described herein) with at least two of the immune checkpoint inhibitors. In embodiments, a method comprises administering a third checkpoint inhibitor. In embodiments, a method comprises administering a TIM-3 inhibitor with a PD-1 inhibitor, and a LAG-3 inhibitor, such that the subject receives all three. In embodiments, a method comprises administering a TIM-3 inhibitor with a PD-1 inhibitor, a LAG-3 inhibitor, and a CTLA-4 inhibitor, such that the subject receives all four.

[0035] In embodiments, a subject has been further administered or will be administered an agent that inhibits poly (ADP-ribose) polymerase (PARP), such that the subject receives treatment with a TIM-3 inhibitor and a PARP inhibitor.

[0036] In embodiments, a PARP inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, or a toxin. In embodiments, a PARP inhibitor is selected from the group consisting of: ABT-767, AZD 2461, BGB-290, BGP 15, CEP 8983, CEP 9722, DR 2313, E7016, E7449, fluzoparib, IMP 4297, INO1001, JPI 289, JPI 547, monoclonal antibody B3-LysPE40 conjugate, MP 124, niraparib, NU 1025, NU 1064, NU 1076, NU1085, olaparib, ONO2231, PD 128763, R 503, R554, rucaparib, SBP 101, SC 101914, simmiparib, talazoparib, veliparib, WW 46, 2-(4-(trifluoromethyl)phenyl)-7,8-dihydro-5H-thiopyrano[4,3-d]pyrimidin-4-ol, and salts or derivatives thereof. In embodiments, a PARP inhibitor is niraparib, olaparib, rucaparib, talazoparib, or veliparib. In embodiments, a PARP inhibitor is niraparib (e.g., niraparib free base, niraparib tosylate, or niraparib tosylate monohydrate, or any combination thereof).

[0037] In embodiments, a subject is further administered or will be administered one or more immune checkpoint inhibitors (e.g., a PD-1 inhibitor and/or a LAG-3 inhibitor) such that the subject receives treatment with a TIM-3 inhibitor, a PARP inhibitor (e.g., niraparib), and the one or more immune checkpoint inhibitors. In embodiments, a subject is administered a TIM-3 inhibitor, a PD-1 inhibitor (e.g., TSR-042) and a PARP inhibitor (e.g., niraparib). In embodiments, a subject is administered a TIM-3 inhibitor, a PD-1 inhibitor (e.g., TSR-042), a LAG-3 inhibitor, and a PARP inhibitor (e.g., niraparib).

[0038] In embodiments, a patient has a disorder that is a T-cell dysfunctional disorder.

[0039] In embodiments, a patient has a disorder that is cancer.

[0040] In embodiments, a cancer is associated with a high tumor mutation burden (TMB).

[0041] In embodiments, a cancer is microsatellite stable (MSS).

[0042] In embodiments a cancer is characterized by microsatellite instability.

[0043] In embodiments, a cancer has a high microsatellite instability status (MSI-H).

[0044] In embodiments, a cancer has a low microsatellite instability status (MSI-L).

[0045] In embodiments, a cancer is associated with high TMB and MSI-H.

[0046] In embodiments, a cancer is associated with high TMB and MSI-L or MSS. In embodiments, a cancer is associated with high TMB and MSI-L. In embodiments, a cancer is associated with high TMB and MSS.

[0047] In embodiments, a cancer has a defective DNA mismatch repair system.

[0048] In embodiments, a cancer has a defect in a DNA mismatch repair gene.

[0049] In embodiments, a cancer is a hypermutated cancer.

[0050] In embodiments, a cancer has homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0051] In embodiments, a cancer comprises a mutation in polymerase delta (POLD).

[0052] In embodiments, a cancer comprises a mutation in polymerase epsilon (POLE).

[0053] In embodiments, a cancer is adenocarcinoma, endometrial cancer, breast cancer, ovarian cancer, cervical cancer, fallopian tube cancer, testicular cancer, primary peritoneal cancer, colon cancer, colorectal cancer, stomach cancer, small intestine cancer, squamous cell carcinoma of the anogenital region (e.g., squamous cell carcinoma of the anus, penis, cervix, vagina, or vulva), soft tissue sarcoma (e.g., leiomyosarcoma), melanoma, renal cell carcinoma, lung cancer, non-small cell lung cancer, adenocarcinoma of the lung, squamous cell carcinoma of the lung, stomach cancer, bladder cancer, gall bladder cancer, liver cancer, thyroid cancer, laryngeal cancer, salivary gland cancer, esophageal cancer, head and neck cancer, squamous cell carcinoma of the head and neck, prostate cancer, pancreatic cancer, mesothelioma, Merkel cell carcinoma, sarcoma, glioblastoma, a hematological cancer, multiple myeloma, B-cell lymphoma, T-cell lymphoma, Hodgkin’s lymphoma (HL)/primary mediastinal B-cell lymphoma, chronic myelogenous leukemia, acute myeloid leukemia, acute lymphoblastic leukemia, non-Hodgkin’s lymphoma, neuroblastoma, a CNS tumor, diffuse intrinsic pontine glioma (DIPG), Ewing’s sarcoma, embryonal rhabdomyosarcoma, osteosarcoma, or Wilms tumor. In embodiments, a cancer is MSS or MSI-L, is characterized by microsatellite instability, is MSI-H, has high TMB, has high TMB and is MSS or MSI-L, has high TMB and is MSI-H, has a defective DNA mismatch repair system, has a defect in a DNA mismatch repair gene, is a hypermutated cancer, is an HRD cancer, comprises a mutation in polymerase delta (POLD) or comprises a mutation in polymerase epsilon (POLE).

[0054] In embodiments, a cancer is endometrial cancer (e.g., MSI-H or MSS/MSI-L endometrial cancer). In embodiments, a cancer is a MSI-H cancer comprising a mutation in

POLE or POLD (e.g., a MSI-H non-endometrial cancer comprising a mutation in POLE or POLD). In embodiments, a cancer is breast cancer (triple negative breast cancer (TNBC)). In embodiments, a cancer is lung cancer (e.g., non-small cell lung cancer). In embodiments, a cancer is melanoma. In embodiments, a cancer is colorectal cancer. In embodiments, a cancer is squamous cell carcinoma of the anus, squamous cell carcinoma of the penis, squamous cell carcinoma of the cervix, squamous cell carcinoma of the vagina, or squamous cell carcinoma of the vulva.

[0055] In embodiments, a cancer has homologous recombination repair deficiency/homologous repair deficiency (“HRD”). In embodiments, a cancer is acute myeloid leukemia. In embodiments, a cancer is acute lymphoblastic leukemia. In embodiments, a cancer is non-Hodgkin’s lymphoma. In embodiments, a cancer is Hodgkin’s lymphoma. In embodiments, a cancer is neuroblastoma. In embodiments, a cancer is a CNS tumor. In embodiments, a cancer is diffuse intrinsic pontine glioma (DIPG). In embodiments, a cancer is Ewing’s sarcoma. In embodiments, a cancer is embryonal rhabdomyosarcoma. In embodiments, a cancer is osteosarcoma. In embodiments, a cancer is Wilms tumor. In embodiments, a cancer is a soft tissue sarcoma (e.g., leiomyosarcoma).

[0056] In some embodiments, a patient has cancer, such as: a non-small cell lung cancer (NSCLC), a hepatocellular cancer, a renal cancer, a melanoma, a cervical cancer, a colorectal cancer, a squamous cell carcinoma of the anogenital region, a head and neck cancer, a triple negative breast cancer, an ovarian cancer or an endometrial cancer. In some embodiments, a patient has a cancer with microsatellite instability. In some embodiments, the microsatellite instability is considered high, wherein the instability is significantly higher than that observed in a control cell (e.g., MSI-H status). In some embodiments, the patient has a solid tumor. In some embodiments, the patient has an advanced stage solid tumor. In some embodiments, a patient has an advanced stage solid tumor, such as a non-small cell lung cancer (NSCLC), a hepatocellular cancer, a renal cancer, a melanoma, a cervical cancer, a colorectal cancer, a squamous cell carcinoma of the anogenital region, a head and neck cancer, a triple negative breast cancer, an ovarian cancer or a endometrial cancer. In some embodiments, a patient has an advanced stage solid tumor with microsatellite instability.

[0057] In some embodiments, the patient has a hematological cancer. In some embodiments, the patient has a hematological cancer such as Diffuse large B cell lymphoma (“DLBCL”),

Hodgkin's lymphoma ("HL"), Non-Hodgkin's lymphoma ("NHL"), Follicular lymphoma ("FL"), acute myeloid leukemia ("AML"), acute lymphoblastic leukemia ("ALL"), or Multiple myeloma ("MM"). In some embodiments, a patient has a hematological cancer with microsatellite instability.

[0058] In some embodiments, a patient has a cancer characterized by PD-1 and/or PD-L1 expression. In some embodiments, a cancer has high PD-1 and/or PD-L1 expression (e.g., by high PD-1 and/or high PD-L1 expression). In some embodiment, a cancer characterized by PD-1 and/or PD-L1 expression is a head and neck cancer, a lung cancer (e.g., a non-small cell lung cancer (NSCLC)), a renal cancer, a bladder cancer, a melanoma, Merkel cell carcinoma, a cervical cancer, a vaginal cancer, a vulvar cancer, a uterine cancer, an endometrial cancer, an ovarian cancer, a fallopian tube cancer, a breast cancer, a prostate cancer, a salivary gland tumor, a thymoma, an adrenocortical carcinoma, an esophageal cancer, a gastric cancer, a colorectal cancer, an appendiceal cancer, a urothelial cell carcinoma, or a squamous cell carcinoma (e.g., of the lung; of the anogenital region including anus, penis, cervix, vagina, or vulva; or of the esophagus). In some certain embodiments, a cancer characterized by PD-1 and/or PD-L1 expression is an anal cancer, a fallopian tube cancer, an ovarian cancer, or a lung cancer.

[0059] In some embodiments, the patient has a head and neck cancer, a lung cancer (e.g., a non-small cell lung cancer (NSCLC)), a renal cancer, a bladder cancer, a melanoma, Merkel cell carcinoma, a cervical cancer, a vaginal cancer, a vulvar cancer, a uterine cancer, an endometrial cancer, an ovarian cancer, a fallopian tube cancer, a breast cancer, a prostate cancer, a salivary gland tumor, a thymoma, an adrenocortical carcinoma, an esophageal cancer, a gastric cancer, a colorectal cancer, an appendiceal cancer, a urothelial cell carcinoma, or a squamous cell carcinoma.

[0060] In embodiments, a cancer is an advanced cancer. In embodiments, a cancer is a metastatic cancer. In embodiments, a cancer is a MSI-H cancer. In embodiments, a cancer is a MSS cancer. In embodiments, a cancer is a POLE-mutant cancer. In embodiments, a cancer is a POLD-mutant cancer. In embodiments, a cancer is a high TMB cancer. In embodiments, a cancer is associated with homologous recombination repair deficiency/homologous repair deficiency ("HRD").

[0061] In embodiments, a cancer is a solid tumor. In embodiments, a solid tumor is advanced. In embodiments, a solid tumor is a metastatic solid tumor. In embodiments, a solid

tumor is a MSI-H solid tumor. In embodiments, a solid tumor is a MSS solid tumor. In embodiments, a solid tumor is a POLE-mutant solid tumor. In embodiments, a solid tumor is a POLD-mutant solid tumor. In embodiments, a solid tumor is a high TMB solid tumor. . In embodiments, a solid tumor is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0062] In embodiments, a cancer is a non-endometrial cancer (e.g., a non-endometrial solid tumor). In embodiments, a non-endometrial cancer is an advanced cancer. In embodiments, a non-endometrial cancer is a metastatic cancer. In embodiments, a non-endometrial cancer is a MSI-H cancer. In embodiments, a non-endometrial cancer is a MSS cancer. In embodiments, a non-endometrial cancer is a POLE-mutant cancer. In embodiments, a non-endometrial cancer is a solid tumor (e.g., a MSS solid tumor, a MSI-H solid tumor, a POLD mutant solid tumor, or a POLE-mutant solid tumor). In embodiments, a non-endometrial cancer is a high TMB cancer. In embodiments, a non-endometrial cancer is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0063] In embodiments, a cancer is endometrial cancer (e.g., a solid tumor). In embodiments, an endometrial cancer is an advanced cancer. In embodiments, an endometrial cancer is a metastatic cancer. In embodiments, an endometrial cancer is a MSI-H endometrial cancer. In embodiments, an endometrial cancer is a MSS endometrial cancer. In embodiments, an endometrial cancer is a POLE-mutant endometrial cancer. In embodiments, an endometrial cancer is a POLD-mutant endometrial cancer. In embodiments, an endometrial cancer is a high TMB endometrial cancer. In embodiments, an endometrial cancer is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0064] In embodiments, a cancer is a lung cancer (e.g., a solid tumor). In embodiments, a lung cancer is an advanced lung cancer. In embodiments, a lung cancer is a metastatic lung cancer. In embodiments, a lung cancer is squamous cell carcinoma of the lung. In embodiments, a lung cancer is small cell lung cancer (SCLC). In embodiments, a lung cancer is non-small cell lung cancer (NSCLC). In embodiments, a lung cancer is an ALK-translocated lung cancer (e.g., a lung cancer with a known ALK-translocation). In embodiments, a lung cancer is an EGFR-mutant lung cancer (e.g., a lung cancer with a known EGFR mutation). In embodiments, a lung cancer is a MSI-H lung cancer. In embodiments, a lung cancer is a MSS lung cancer. In embodiments, a lung cancer is a POLE-mutant lung cancer. In embodiments, a lung cancer is a

POLD-mutant lung cancer. In embodiments, a lung cancer is a high TMB lung cancer. In embodiments, a lung cancer is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0065] In embodiments, a cancer is a colorectal (CRC) cancer (e.g., a solid tumor). In embodiments, a colorectal cancer is an advanced colorectal cancer. In embodiments, a colorectal cancer is a metastatic colorectal cancer. In embodiments, a colorectal cancer is a MSI-H colorectal cancer. In embodiments, a colorectal cancer is a MSS colorectal cancer. In embodiments, a colorectal cancer is a POLE-mutant colorectal cancer. In embodiments, a colorectal cancer is a POLD-mutant colorectal cancer. In embodiments, a colorectal cancer is a high TMB colorectal cancer. In embodiments, a colorectal cancer is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0066] In embodiments, a cancer is a melanoma. In embodiments, a melanoma is an advanced melanoma. In embodiments, a melanoma is a metastatic melanoma. In embodiments, a melanoma is a MSI-H melanoma. In embodiments, a melanoma is a MSS melanoma. In embodiments, a melanoma is a POLE-mutant melanoma. In embodiments, a melanoma is a POLD-mutant melanoma. In embodiments, a melanoma is a high TMB melanoma. In embodiments, a melanoma is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0067] In embodiments, a cancer is squamous cell carcinoma of the anogenital region (e.g., of the anus, penis, cervix, vagina, or vulva). In embodiments, a squamous cell carcinoma of the anogenital region (e.g., of the anus, penis, cervix, vagina, or vulva) is an advanced cancer. In embodiments, a squamous cell carcinoma of the anogenital region (e.g., of the anus, penis, cervix, vagina, or vulva) is a metastatic cancer. In embodiments, a squamous cell carcinoma of the anogenital region (e.g., of the anus, penis, cervix, vagina, or vulva) is MSI-H. In embodiments, a squamous cell carcinoma of the anogenital region (e.g., of the anus, penis, cervix, vagina, or vulva) is MSS. In embodiments, a lung cancer is a POLE-mutant cancer. In embodiments, a squamous cell carcinoma of the anogenital region (e.g., of the anus, penis, cervix, vagina, or vulva) is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0068] In embodiments, a cancer is an ovarian cancer. In embodiments, an ovarian cancer is an advanced ovarian cancer. In embodiments, an ovarian cancer is a metastatic ovarian cancer.

In embodiments, an ovarian cancer is a MSI-H ovarian cancer. In embodiments, an ovarian cancer is a MSS ovarian cancer. In embodiments, an ovarian cancer is a POLE-mutant ovarian cancer. In embodiments, an ovarian cancer is a POLD-mutant ovarian cancer. In embodiments, an ovarian cancer is a high TMB ovarian cancer. In embodiments, an ovarian cancer is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”). In embodiments, an ovarian cancer is a serous cell ovarian cancer. In embodiments, an ovarian cancer is a clear cell ovarian cancer.

[0069] In embodiments, a cancer is a fallopian cancer. In embodiments, a fallopian cancer is an advanced fallopian cancer. In embodiments, a fallopian cancer is a metastatic fallopian cancer. In embodiments, a fallopian cancer is a MSI-H fallopian cancer. In embodiments, a fallopian cancer is a MSS fallopian cancer. In embodiments, a fallopian cancer is a POLE-mutant fallopian cancer. In embodiments, a fallopian cancer is a POLD-mutant fallopian cancer. In embodiments, a fallopian cancer is a high TMB fallopian cancer. In embodiments, a fallopian cancer is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”). In embodiments, a fallopian cancer is a serous cell fallopian cancer. In embodiments, a fallopian cancer is a clear cell fallopian cancer.

[0070] In embodiments, a cancer is a primary peritoneal cancer. In embodiments, a primary peritoneal cancer is an advanced primary peritoneal cancer. In embodiments, a primary peritoneal cancer is a metastatic primary peritoneal cancer. In embodiments, a primary peritoneal cancer is a MSI-H primary peritoneal cancer. In embodiments, a primary peritoneal cancer is a MSS primary peritoneal cancer. In embodiments, a primary peritoneal cancer is a POLE-mutant primary peritoneal cancer. In embodiments, a primary peritoneal cancer is a POLD-mutant primary peritoneal cancer. In embodiments, a primary peritoneal cancer is a high TMB primary peritoneal cancer. In embodiments, a primary peritoneal cancer is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”). In embodiments, a primary peritoneal cancer is a serous cell primary peritoneal cancer. In embodiments, a primary peritoneal cancer is a clear cell primary peritoneal cancer.

[0071] In embodiments, a cancer is acute lymphoblastic leukemia (“ALL”). In embodiments, acute lymphoblastic leukemia is advanced acute lymphoblastic leukemia. In embodiments, acute lymphoblastic leukemia is metastatic acute lymphoblastic leukemia. In embodiments, acute lymphoblastic leukemia is MSI-H acute lymphoblastic leukemia. In

embodiments, acute lymphoblastic leukemia is MSS acute lymphoblastic leukemia. In embodiments, acute lymphoblastic leukemia is POLE-mutant acute lymphoblastic leukemia. In embodiments, acute lymphoblastic leukemia is POLD-mutant acute lymphoblastic leukemia. In embodiments, an acute lymphoblastic leukemia is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0072] In embodiments, a cancer is acute myeloid leukemia (“AML”). In embodiments, acute myeloid leukemia is advanced acute myeloid leukemia. In embodiments, acute myeloid leukemia is metastatic acute myeloid leukemia. In embodiments, acute myeloid leukemia is MSI-H acute myeloid leukemia. In embodiments, acute myeloid leukemia is MSS acute myeloid leukemia. In embodiments, acute myeloid leukemia is POLE-mutant acute myeloid leukemia. In embodiments, acute myeloid leukemia is POLD-mutant acute myeloid leukemia. In embodiments, an acute myeloid leukemia is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0073] In embodiments, a cancer is non-Hodgkin’s lymphoma (NHL). In embodiments, non-Hodgkin’s lymphoma is advanced non-Hodgkin’s lymphoma. In embodiments, non-Hodgkin’s lymphoma is metastatic non-Hodgkin’s lymphoma. In embodiments, non-Hodgkin’s lymphoma is MSI-H non-Hodgkin’s lymphoma. In embodiments, non-Hodgkin’s lymphoma is MSS non-Hodgkin’s lymphoma. In embodiments, non-Hodgkin’s lymphoma is POLE-mutant non-Hodgkin’s lymphoma. In embodiments, non-Hodgkin’s lymphoma is POLD-mutant non-Hodgkin’s lymphoma. In embodiments, non-Hodgkin’s lymphoma is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0074] In embodiments, a cancer is Hodgkin’s lymphoma (HL). In embodiments, Hodgkin’s lymphoma is advanced Hodgkin’s lymphoma. In embodiments, Hodgkin’s lymphoma is metastatic Hodgkin’s lymphoma. In embodiments, Hodgkin’s lymphoma is MSI-H Hodgkin’s lymphoma. In embodiments, Hodgkin’s lymphoma is MSS Hodgkin’s lymphoma. In embodiments, Hodgkin’s lymphoma is POLE-mutant Hodgkin’s lymphoma. In embodiments, Hodgkin’s lymphoma is POLD-mutant Hodgkin’s lymphoma. In embodiments, Hodgkin’s lymphoma is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0075] In embodiments, a cancer is a neuroblastoma (NB). In embodiments, a neuroblastoma is an advanced neuroblastoma. In embodiments, a neuroblastoma is a metastatic

neuroblastoma. In embodiments, neuroblastoma is a MSI-H neuroblastoma. In embodiments, a neuroblastoma is a MSS neuroblastoma. In embodiments, a neuroblastoma is a POLE-mutant neuroblastoma. In embodiments, a neuroblastoma is a POLD-mutant neuroblastoma. In embodiments, a neuroblastoma is a high TMB neuroblastoma. In embodiments, a neuroblastoma is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0076] In embodiments, a cancer is a CNS tumor. In embodiments, a CNS tumor is advanced. In embodiments, a CNS tumor is a metastatic CNS tumor. In embodiments, a CNS tumor is a MSI-H CNS tumor. In embodiments, a CNS tumor is a MSS CNS tumor. In embodiments, a CNS tumor is a POLE-mutant CNS tumor. In embodiments, a CNS tumor is a POLD-mutant CNS tumor. In embodiments, a CNS tumor is a high TMB CNS tumor. . In embodiments, a CNS tumor is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0077] In embodiments, a cancer is diffuse intrinsic pontine glioma (DIPG). In embodiments, a DIPG is an advanced DIPG. In embodiments, a DIPG is a metastatic DIPG. In embodiments, DIPG is a MSI-H DIPG. In embodiments, a DIPG is a MSS DIPG. In embodiments, a DIPG is a POLE-mutant DIPG. In embodiments, a DIPG is a POLD-mutant DIPG. In embodiments, a DIPG is a high TMB DIPG. In embodiments, a DIPG is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0078] In embodiments, a cancer is Ewing’s sarcoma. In embodiments, Ewing’s sarcoma is an advanced Ewing’s sarcoma. In embodiments, Ewing’s sarcoma is a metastatic Ewing’s sarcoma. In embodiments, Ewing’s sarcoma is a MSI-H Ewing’s sarcoma. In embodiments, Ewing’s sarcoma is a MSS Ewing’s sarcoma. In embodiments, Ewing’s sarcoma is a POLE-mutant Ewing’s sarcoma. In embodiments, Ewing’s sarcoma is a POLD-mutant Ewing’s sarcoma. In embodiments, Ewing’s sarcoma is a high TMB Ewing’s sarcoma. In embodiments, Ewing’s sarcoma is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0079] In embodiments, a cancer is an embryonal rhabdomyosarcoma (ERS). In embodiments, an embryonal rhabdomyosarcoma is an advanced embryonal rhabdomyosarcoma. In embodiments, an embryonal rhabdomyosarcoma is a metastatic embryonal rhabdomyosarcoma. In embodiments, an embryonal rhabdomyosarcoma is a MSI-H embryonal

rhabdomyosarcoma. In embodiments, an embryonal rhabdomyosarcoma is a MSS embryonal rhabdomyosarcoma. In embodiments, an embryonal rhabdomyosarcoma is a POLE-mutant embryonal rhabdomyosarcoma. In embodiments, an embryonal rhabdomyosarcoma is a POLD-mutant embryonal rhabdomyosarcoma. In embodiments, an embryonal rhabdomyosarcoma is a high TMB embryonal rhabdomyosarcoma. In embodiments, an embryonal rhabdomyosarcoma is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0080] In embodiments, a cancer is an osteosarcoma (OS). In embodiments, an osteosarcoma is an advanced osteosarcoma. In embodiments, an osteosarcoma is a metastatic osteosarcoma. In embodiments, an osteosarcoma is a MSI-H osteosarcoma. In embodiments, an osteosarcoma is a MSS osteosarcoma. In embodiments, an osteosarcoma is a POLE-mutant osteosarcoma. In embodiments, an osteosarcoma is a POLD-mutant osteosarcoma. In embodiments, an osteosarcoma is a high TMB osteosarcoma. In embodiments, an osteosarcoma is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0081] In embodiments, a cancer is a soft tissue sarcoma. In embodiments, a soft tissue sarcoma is an advanced soft tissue sarcoma. In embodiments, a soft tissue sarcoma is a metastatic soft tissue sarcoma. In embodiments, a soft tissue sarcoma is a MSI-H soft tissue sarcoma. In embodiments, a soft tissue sarcoma is a MSS soft tissue sarcoma. In embodiments, a soft tissue sarcoma is a POLE-mutant soft tissue sarcoma. In embodiments, a soft tissue sarcoma is a POLD-mutant soft tissue sarcoma. In embodiments, a soft tissue sarcoma is a high TMB soft tissue sarcoma. In embodiments, a soft tissue sarcoma is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”). In embodiments, a soft tissue sarcoma is leiomyosarcoma.

[0082] In embodiments, a cancer is Wilms tumor. In embodiments, Wilms tumor is an advanced Wilms tumor. In embodiments, Wilms tumor is a metastatic Wilms tumor. In embodiments, Wilms tumor is a MSI-H Wilms tumor. In embodiments, Wilms tumor is a MSS Wilms tumor. In embodiments, Wilms tumor is a POLE-mutant Wilms tumor. In embodiments, Wilms tumor is a POLD-mutant Wilms tumor. In embodiments, Wilms tumor is a high TMB Wilms tumor. In embodiments, Wilms tumor is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

[0083] In embodiments, a subject has previously been treated with one or more different cancer treatment modalities (e.g., one or more of surgery, radiotherapy, chemotherapy, or immunotherapy).

[0084] In embodiments, a subject has previously been treated with one different cancer treatment modalities (e.g., one or more of surgery, radiotherapy, chemotherapy, or immunotherapy). In embodiments, a subject has previously been treated with two or more different cancer treatment modalities (e.g., one or more of surgery, radiotherapy, chemotherapy, or immunotherapy). In embodiments, a subject has been previously treated with a cytotoxic therapy. In embodiments, a subject has been previously treated with chemotherapy. In embodiments, a subject has previously been treated with two different cancer treatment modalities (e.g., one or more of surgery, radiotherapy, chemotherapy, or immunotherapy). In embodiments, a subject has previously been treated with three different cancer treatment modalities (e.g., one or more of surgery, radiotherapy, chemotherapy, or immunotherapy).

[0085] In embodiments of methods described herein, a method further comprises administering one or more of surgery, a radiotherapy, a chemotherapy, an immunotherapy, an anti-angiogenic agent, or an anti-inflammatory. In embodiments, a method further comprises administering a chemotherapy.

[0086] In some embodiments, at least some of the patients in the cancer patient population have previously been treated with chemotherapy (e.g., platinum-based chemotherapy). For example, a patient who has received two lines of cancer treatment can be identified as a 2L cancer patient (e.g., a 2L NSCLC patient). In embodiments, a patient has received two lines or more lines of cancer treatment (e.g., a 2L+ cancer patient such as a 2L+ endometrial cancer patient). In embodiments, a patient has not been previously treated with an anti-PD-1 therapy. In embodiments, a patient previously received at least one line of cancer treatment (e.g., a patient previously received at least one line or at least two lines of cancer treatment). In embodiments, a patient previously received at least one line of treatment for metastatic cancer (e.g., a patient previously received one or two lines of treatment for metastatic cancer).

[0087] In embodiments, a subject is resistant to treatment with an agent that inhibits PD-1.

[0088] In embodiments, a subject is refractory to treatment with an agent that inhibits PD-1.

[0089] In embodiments, a method described herein sensitizes a subject to treatment with an agent that inhibits PD-1.

[0090] In embodiments, a subject comprises an exhausted immune cell (e.g., an exhausted immune cell that is an exhausted T cell).

[0091] In embodiments of methods described herein, a subject is an animal (e.g., a mammal). In embodiments, a subject is a human. In embodiments, a subject is a non-human mammal (e.g., mice, rats, rabbits, or non-human primates). Accordingly, methods described herein can be useful in both treatment of humans and in veterinary medicine.

[0092] In embodiments, a TIM-3 inhibitor is administered intravenously (e.g., by intravenous infusion).

[0093] The present disclosure provides, in some embodiments, methods of treating cancer in a patient in need thereof, the method comprising administering a composition that delivers an anti-TIM-3 antibody agent.

[0094] In some embodiments, the patient has not previously been treated with a cancer treatment modality.

[0095] In some embodiments, the patient has previously been treated with one or more different cancer treatment modalities. In some embodiments, the patient has previously been treated with one or more of radiotherapy, chemotherapy or immunotherapy. In some embodiments, the patient has previously been treated with chemotherapy (e.g., platinum-based chemotherapy).

[0096] In some embodiments, a composition that delivers a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is administered at a dose of 1, 3 or 10 mg/kg. In some embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of 1, 3 or 10 mg/kg every two weeks. In some embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of 1, 3 or 10 mg/kg every three weeks. In some embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of 1, 3 or 10 mg/kg every four weeks.

[0097] In some embodiments, a composition that delivers an anti-TIM-3 antibody agent at a fixed dose within a range of about 100 mg to about 1500 mg (e.g., about 100 mg to 1500 mg). In some embodiments, a composition that delivers an anti-TIM-3 antibody agent at a fixed dose within a range of about 100 mg to about 300 mg, about 300 mg to about 1,000 mg, or about 1000 mg to about 1200 mg. In embodiments, the fixed dose is about 100 mg. In embodiments, the

fixed dose is about 300 mg. In embodiments, the fixed dose is about 500 mg. In embodiments, the fixed dose is about 900 mg. In embodiments, the fixed dose is about 1200 mg. In some embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a fixed dose every two weeks (Q2W). In some embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a fixed dose every three weeks (Q3W). In some embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a fixed dose every four weeks (Q4W). In embodiments, the anti-TIM-3 antibody is administered at the administration interval of once a week (Q1W), once every 2 weeks (Q2W), once every 3 weeks (S3W), once every 4 weeks (Q4W), once every 5 weeks (Q5W), or once every 6 weeks (Q6W). In embodiments, the anti-TIM-3 antibody is administered for the period of at least 2, 4, 6, 8, 10, 12, 14, 16, 18, or 20 weeks. In embodiments, a dose is administered as a monotherapy (e.g., a therapeutically effective amount of 1200 mg of an anti-TIM-3 antibody administered Q2W or Q3W) or a dose is administered in combination with one or more other therapies. For example, 100 mg, 300 mg, 500 mg, or 900 mg (e.g., 100 mg or 300 mg) of an anti-TIM-3 antibody can be administered in combination with an anti-PD-1 antibody according to regimens described herein (e.g., 500 mg of an anti-PD-1 antibody administered Q3W for four treatment cycles followed by administration of 1000 mg of the anti-PD-1 antibody Q6W until treatment is discontinued (e.g., due to disease progression)).

[0098] In some embodiments, the composition is administered intravenously. In some embodiments, the composition is administered by intravenous infusion.

[0099] In some embodiments, a clinical benefit is a complete response (“CR”), a partial response (“PR”) or a stable disease (“SD”). In some embodiments, a clinical benefit corresponds to at least SD. In some embodiments, a clinical benefit corresponds to at least a PR. In some embodiments, a clinical benefit corresponds to a CR. In some embodiments, at least 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90% or 95% of patients achieve a clinical benefit. In some embodiments, at least 5% of patients achieve a clinical benefit. In some embodiments, at least 5% of patients achieve SD. In some embodiments, at least 5% of patients achieve at least a PR. In some embodiments, at least 5% of patients achieve CR. In some embodiments, at least 20% of patients achieve a clinical benefit. In some embodiments, at least 20% of patients achieve SD.

[00100] In some embodiments, the clinical benefit (e.g., SD, PR and/or CR) is determined in accordance with Response Evaluation Criteria in Solid Tumors (RECIST). In some embodiments, the clinical benefit (e.g., SD, PR and/or CR) is determined in accordance RECIST guidelines. In some embodiments, the clinical benefit (e.g., SD, PR and/or CR) is determined in accordance RECIST guidelines (version 1.1). In some embodiments, the clinical benefit (e.g., SD, PR and/or CR) is determined in accordance immune-related RECIST (irRECIST) guidelines. In some embodiments, tumor response can be assessed by either irRECIST or RECIST version 1.1. In some embodiments, tumor response can be assessed by both irRECIST and RECIST version 1.1. When used herein, the term “RECIST guidelines” can refer to RECIST 1.0, RECIST 1.1 or ir RECIST interchangeably.

[00101] In some embodiments, the patient is receiving or will receive an additional therapy in combination with the anti-TIM-3 antibody agent. In some embodiments, the additional therapy is radiotherapy, chemotherapy or immunotherapy. In some embodiments, the additional therapy includes treatment with a composition that delivers a PD-1 inhibitor (e.g., a PD-1-binding agent) and/or a LAG-3-binding agent. In some embodiments, the additional PD-1 inhibitor is nivolumab, pembrolizumab, atezolizumab, durvalumab, avelumab, TSR-042, PDR-001, tislelizumab (BGB-A317), cemiplimab (REGN2810), LY-3300054, JNJ-63723283, MGA012, BI-754091, IBI-308, camrelizumab (HR-301210), BCD-100, JS-001, CX-072, BGB-A333, AMP-514 (MEDI-0680), AGEN-2034, CS1001, Sym-021, SHR-1316, PF-06801591, LZM009, KN-035, AB122, genolimzumab (CBT-501), FAZ-053, CK-301, AK 104, GLS-010, PD-1VR or PD-1FL, or any of the PD-1 antibodies disclosed in WO2014/179664. In some embodiments, the additional therapy is a PARP inhibitor. In some embodiments, the PARP inhibitor is niraparib, olaparib, rucaparib, talazoparib, and veliparib. . In some embodiments, the PARP inhibitor is niraparib.

[00102] The present disclosure provides, in some embodiments, methods of treating cancer in a patient in need thereof, the method comprising administering a one or more compositions that deliver an anti-TIM-3 antibody agent in combination with a PD-1-binding agent. In some embodiments, a patient or patient population is receiving a combination therapy that comprises administration of an anti-TIM-3 antibody agent and a PD-1-binding agent (e.g., an anti-PD-1 antibody).

[00103] In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain variable domain whose amino acid sequence comprises SEQ ID NO: 1 or SEQ ID NO: 7 and an immunoglobulin light chain variable domain whose amino acid sequence comprises SEQ ID NO: 2 or SEQ ID NO: 8. In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain whose amino acid sequence comprises SEQ ID NO: 3 and an immunoglobulin light chain whose amino acid sequence comprises SEQ ID NO: 4.

[00104] In some embodiments, a PD-1-binding agent comprises an immunoglobulin heavy chain variable domain whose amino acid sequence comprises SEQ ID NO: 11 or SEQ ID NO: 17 and an immunoglobulin light chain variable domain whose amino acid sequence comprises SEQ ID NO: 12 or SEQ ID NO: 18 (“PD-1VR”). In some embodiments, a PD-1-binding agent comprises an immunoglobulin heavy chain whose amino acid sequence comprises SEQ ID NO: 13 and an immunoglobulin light chain whose amino acid sequence comprises SEQ ID NO: 14 (“PD1-FL”).

[00105] In some embodiments, an anti-TIM-3 antibody agent (e.g., an anti-TIM-3 antibody) is administered at a dose of about 1, 3 or 10 mg/kg. In some embodiments, an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of 1, 3 or 10 mg/kg every two weeks. In some embodiments, an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of about 1, 3 or 10 mg/kg every three weeks. In some embodiments, an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of 1, 3 or 10 mg/kg every four weeks. In some embodiments, an anti-TIM-3 antibody agent is administered at a fixed dose within a range of about 100 mg to 1,500 mg. In some embodiments, an anti-TIM-3 antibody agent is administered at a fixed dose within a range of about 300 mg to 1,000 mg, about 100 mg to about 500 mg, about 100 mg to about 1200 mg, or about 1000 mg to about 1500 mg. In some embodiments, an anti-TIM-3 antibody agent is administered at a fixed dose of about 100, 200 mg, 300 mg, 400 mg, 500 mg, 600 mg, 700 mg, 800 mg, 900 mg, 1000 mg, 1100 mg, 1200 mg, 1300 mg, 1400 mg, or 1500 mg. In some embodiments, an anti-TIM-3 antibody agent is administered at a fixed dose of about 100 mg. In some embodiments, an anti-TIM-3 antibody agent is administered at a fixed dose of about 300 mg. In some embodiments, an anti-TIM-3 antibody agent is administered at a fixed dose of about 500 mg. In some embodiments, an anti-TIM-3 antibody agent is administered at a fixed dose of about 900 mg. In some embodiments, an anti-TIM-3 antibody agent is administered at a

fixed dose of about 1200 mg. In some embodiments, an anti-TIM-3 antibody agent is administered according to a regimen that includes a fixed dose every two weeks (Q2W). In some embodiments, an anti-TIM-3 antibody agent is administered according to a regimen that includes a fixed dose every three weeks (Q3W). In some embodiments, an anti-TIM-3 antibody agent is administered according to a regimen that includes a fixed dose every four weeks (Q4W).

[00106] In some embodiments, a therapeutically effective dose is a dose of an anti-TIM-3 antibody of about 100 mg to about 1500 mg such as a dose that is about 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200 mg, 1300 mg, 1400 mg, or 1500 mg. In embodiments, a therapeutically effective dose is about 100 mg (e.g., administered Q2W or Q3W). In embodiments, a therapeutically effective dose is about 300 mg (e.g., administered Q2W or Q3W). In embodiments, a therapeutically effective dose is about 500 mg (e.g., administered Q2W or Q3W). In embodiments, a therapeutically effective dose is about 900 mg (e.g., administered Q2W or Q3W). In embodiments, a therapeutically effective dose is about 1200 mg (e.g., administered Q2W or Q3W). In embodiments, the anti-TIM-3 antibody is administered at the administration interval of once a week (Q1W), once every 2 weeks (Q2W), once every 3 weeks (S3W), once every 4 weeks (Q4W), once every 5 weeks (Q5W), or once every 6 weeks (Q6W). In embodiments, the anti-TIM-3 antibody is administered for the period of at least 2, 4, 6, 8, 10, 12, 14, 16, 18, or 20 weeks.

[00107] In embodiments, a therapeutically effective dose is about 100 mg, and the agent is administered at an administration interval of once every 2 weeks. In embodiments, a therapeutically effective dose is about 100 mg, and the agent is administered at an administration interval of once every 3 weeks.

[00108] In embodiments, a therapeutically effective dose is about 300 mg, and the agent is administered at an administration interval of once every 2 weeks. In embodiments, a therapeutically effective dose is about 300 mg, and the agent is administered at an administration interval of once every 3 weeks.

[00109] In embodiments, a therapeutically effective dose is about 500 mg, and the agent is administered at an administration interval of once every 2 weeks. In embodiments, a therapeutically effective dose is about 500 mg, and the agent is administered at an administration interval of once every 3 weeks.

[00110] In embodiments, a therapeutically effective dose is about 900 mg, and the agent is administered at an administration interval of once every 2 weeks. In embodiments, a therapeutically effective dose is about 900 mg, and the agent is administered at an administration interval of once every 3 weeks.

[00111] In embodiments, a therapeutically effective dose is about 1200 mg, and the agent is administered at an administration interval of once every 2 weeks. In embodiments, a therapeutically effective dose is about 1200 mg, and the agent is administered at an administration interval of once every 3 weeks.

[00112] In embodiments, a dose is administered as a monotherapy (e.g., a therapeutically effective amount of 1200 mg of an anti-TIM-3 antibody administered Q2W or Q3W) or a dose is administered in combination with one or more other therapies. For example, 100 mg or 300 mg of an anti-TIM-3 antibody can be administered in combination with an anti-PD-1 antibody according to regimens described herein (e.g., 500 mg of an anti-PD-1 antibody administered Q3W for four treatment cycles followed by administration of 1000 mg of the anti-PD-1 antibody Q6W until treatment is discontinued (e.g., due to disease progression)).

[00113] In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) is administered at a dose of 0.1, 1, 3, 10, or 20 mg/kg. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) is administered according to a regimen that includes a dose of 1, 3 or 10 mg/kg every two weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) is administered according to a regimen that includes a dose of 1, 3 or 10 mg/kg every three weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) is administered according to a regimen that includes a dose of 1, 3 or 10 mg/kg every four weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) at a dose of 500 mg. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) is administered according to a regimen that includes a dose of 500 mg every two weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) is administered according to a regimen that includes a dose of 500 mg every three weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) is administered according to a regimen that includes a dose of 500 mg every four weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) is administered according to a regimen that includes a dose of 500 mg every three weeks for at least one treatment cycle (e.g., for at least one, two, three, or four treatment cycles), followed by

administration. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) is administered according to a regimen that includes a dose of 1000 mg every three weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) is administered according to a regimen that includes a dose of 1000 mg every four weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) is administered according to a regimen that includes a dose of 1000 mg every five weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) is administered according to a regimen that includes a dose of 1000 mg every six weeks. In embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) is administered according to a regimen that includes a dose of 500 mg every three weeks for four treatment cycles, followed by administration of a dose of 1000 mg every six weeks until treatment is discontinued (e.g., due to disease progression). In embodiments, an anti-TIM-3 antibody agent is administered Q2W or Q3W in a dose that is about 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, or 1200 mg or in a dose that is about 1-10 mg/kg (e.g., a dose that is about 1, about 3, or about 10 mg/kg). In embodiments, an anti-TIM-3 antibody agent is administered Q2W or Q3W in a dose that is about 100 mg. In embodiments, an anti-TIM-3 antibody agent is administered Q2W or Q3W in a dose that is about 300 mg.

[00114] In embodiments, a PD-1 inhibitor (e.g., any anti-PD-1 antibody described herein) is administered at a first dose of about 500 mg once every 3 weeks for 3, 4, or 5 cycles followed by a second dose of about 1000 mg once every 6 weeks or more (e.g., a second dose of about 1000 mg once every 6 weeks). In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 100 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 300 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 500 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 900 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is administered once every three weeks.

[00115] In embodiments, a PD-1 inhibitor (e.g., any anti-PD-1 antibody described herein) is administered at a first dose of about 500 mg once every 3 weeks for 3 cycles followed by a second dose of about 1000 mg once every 6 weeks or more (e.g., a second dose of about 1000 mg once every 6 weeks). In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 100 mg. In embodiments, a therapeutically effective

dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 300 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 500 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 900 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is administered once every three weeks.

[00116] In embodiments, a PD-1 inhibitor (e.g., any anti-PD-1 antibody described herein) is administered at a first dose of about 500 mg once every 3 weeks for 4 cycles followed by a second dose of about 1000 mg once every 6 weeks or more (e.g., a second dose of about 1000 mg once every 6 weeks). In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 100 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 300 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 500 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 900 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is administered once every three weeks.

[00117] In embodiments, a PD-1 inhibitor (e.g., any anti-PD-1 antibody described herein) is administered at a first dose of about 500 mg once every 3 weeks for 5 cycles followed by a second dose of about 1000 mg once every 6 weeks or more (e.g., a second dose of about 1000 mg once every 6 weeks). In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 100 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 300 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 500 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is a flat dose about 900 mg. In embodiments, a therapeutically effective dose of the agent (e.g., an anti-TIM-3 antibody) is administered once every three weeks.

[00118] The present disclosure provides, in some embodiments, compositions comprising an anti-TIM-3 antibody agent for use in treatment of cancer in a selected cancer patient population. In some embodiments, an anti-TIM-3 antibody agent comprises a heavy chain comprising three CDRs that have sequences of SEQ ID NOs: 21, 22, or 23. In some embodiments, an anti-TIM-3 antibody agent comprises a light chain comprising three CDRs that have sequences of SED ID NOs: 24, 25, or 26. In some embodiments, an anti-TIM-3 antibody agent comprises a heavy

chain comprising three CDRs that have sequences of SEQ ID NOs: 21, 22, or 23; and a light chain comprising three CDRs that have sequences of SEQ ID NOs: 24, 25, or 26. In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain variable domain whose amino acid sequence comprises SEQ ID NO: 1 or SEQ ID NO:7. In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin light chain variable domain whose amino acid sequence comprises SEQ ID NO: 2 or SEQ ID NO: 8. In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain variable domain whose amino acid sequence comprises SEQ ID NO: 1 or SEQ ID NO:7 and an immunoglobulin light chain variable domain whose amino acid sequence comprises SEQ ID NO: 2 or SEQ ID NO: 8. In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain whose amino acid sequence comprises SEQ ID NO: 3. In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin light chain whose amino acid sequence comprises SEQ ID NO: 4. In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain whose amino acid sequence comprises SEQ ID NO: 3 and an immunoglobulin light chain whose amino acid sequence comprises SEQ ID NO: 4.

[00119] In some embodiments, the patients in the cancer patient population each have a tumor. In some embodiments, the patients in the cancer patient population each have a solid tumor. In some embodiments, at least some of the patients in the cancer patient population have an advanced stage solid tumor. In some embodiments, at least some of the patients in the cancer patient population have a metastatic solid tumor. In some embodiments, the patient has a MSI-H solid tumor. In some embodiments, the patients in the cancer patient population each have a cancer such as a non-small cell lung cancer (NSCLC), a hepatocellular cancer, a renal cancer, a melanoma, a cervical cancer, a colorectal cancer, a squamous cell carcinoma of the anogenital region (e.g., squamous cell carcinoma of the anus, penis, cervix, vagina, or vulva), a head and neck cancer, a triple negative breast cancer, an ovarian cancer or an endometrial cancer.

[00120] In some embodiments, the patient has a lung cancer (e.g., non-small cell lung cancer (NSCLC)). In some embodiments, the patient has a melanoma.

[00121] In some embodiments, the patient has a cancer associated with a POLE (DNA polymerase epsilon) or a POLD (DNA polymerase delta) mutation. In some embodiments, the POLE or POLD mutation is in an exonuclease domain. In some embodiments, the POLE or POLD mutation is a germline mutation. In some embodiments, the POLE or POLD mutation is

a sporadic mutation. In some embodiments, a method described herein further comprises a step of first identifying the patient having the cancer with the POLE or POLD mutation. In some embodiments, a POLE or POLD mutation is identified using sequencing.

[00122] In some embodiments, the patients in the cancer patient population each have a cancer with microsatellite instability. In some embodiments, the microsatellite instability is considered high, wherein the instability is significantly higher than that observed in a control cell (e.g., MSI-H status). In some embodiments, the microsatellite instability is MSI-Low (MSI-L). In some embodiments, the microsatellite instability is microsatellite stable (e.g., MSS status). In some embodiments, a patient has an advanced stage solid tumor with microsatellite instability.

[00123] In some embodiments, the patients in the cancer patient population each have a hematological cancer. In some embodiments, the patients in the cancer patient population each have a hematological cancer such as Diffuse large B cell lymphoma (“DLBCL”), Hodgkin’s lymphoma (“HL”), Non-Hodgkin’s lymphoma (“NHL”), Follicular lymphoma (“FL”), acute myeloid leukemia (“AML”), acute lymphoblastic leukemia (“ALL”), or Multiple myeloma (“MM”). In some embodiments, the patients in the cancer patient population each have a hematological cancer with microsatellite instability.

[00124] In some embodiments, at least some of the patients in the cancer patient population have previously been treated with one or more different cancer treatment modalities. In some embodiments, at least some of the patients in the cancer patient population have previously been treated with one or more of radiotherapy, chemotherapy or immunotherapy. In some embodiments, at least some of the patients in the cancer patient population have previously been treated with chemotherapy (e.g., platinum-based chemotherapy).

[00125] In some embodiments, at least some of the patients in the cancer patient population have not previously been treated with one or more different cancer treatment modalities.

[00126] The present disclosure provides, in some embodiments, a combination therapy for use in treatment of cancer in a selected cancer patient population, wherein the combination therapy comprises administering an anti-TIM-3 antibody agent and a PD-1-binding agent.

BRIEF DESCRIPTION OF THE DRAWINGS

[00127] **Figure 1** depicts a schematic illustration, not to scale, of enhancement of immune cell activation by anti-TIM-3 and anti-PD1.

[00128] **Figures 2A-2B** depict results from an exemplary T cell exhaustion model. (A) Target expression of PD-1 and TIM-3 in responsive (pre-stimulated) cells and exhausted (post-stimulated cells). (B) Quantification of IFN γ production in exhausted (post-stimulated) cells treated with a combination of an anti-PD-1 antibody agent and an anti-TIM-3 antibody agent, an anti-PD-1 antibody agent, an anti-TIM-3 antibody agent, and isotype control.

[00129] **Figure 3** depicts results from an *in vivo* efficacy study of a combination of an exemplary anti-PD-1 antibody (TSR-042) and an exemplary anti-TIM-3 antibody (TSR-022) antibodies. In the study, huNOG-EXL mice transplanted neonatally with CD34+ hematopoietic stem cells were implanted with A549 NSCLC cells and treated with single agents and combinations of anti-PD-1 and anti-TIM-3 antibodies.

[00130] **Figure 4A** relates to a dose escalation study for an exemplary anti-TIM-3 antibody (TSR-022) as monotherapy or in combination with an anti-PD-1 antibody.

[00131] **Figure 4B** relates to an expansion cohort in order to evaluate the anti-tumor activity of an exemplary anti-TIM-3 antibody (TSR-022) as a monotherapy and in combination with an anti-PD-1 antibody in patients with specific tumor types.

[00132] **Figure 5** describes patient demographics and baseline characteristics for participants in the dose-escalation study.

[00133] **Figure 6A** depicts mean PK profiles of an exemplary anti-TIM-3 antibody (TSR-022) after the first dose.

[00134] **Figure 6B** depicts TIM-3 occupancy on circulating monocytes as measured by flow cytometry from whole blood samples collected from patients treated with an exemplary anti-TIM-3 antibody (TSR-022).

[00135] **Figures 7A-7C** depict TIM-3 receptor occupancy for an exemplary anti-TIM-3 antibody (TSR-022) at doses of 1 mg/kg (**FIG. 7A**), 3 mg/kg (**FIG. 7B**), and 10 mg/kg (**FIG. 7C**) administered once every two weeks (Q2W).

[00136] **Figure 8** depicts the effects of treatment with an exemplary anti-TIM-3 antibody (TSR-022). Duration of treatment associated with specific dosages are shown, and a partial response is noted by black squares and stable disease is noted by black triangles. .

[00137] **Figure 9** depicts the brain scan of a patient with leiomyosarcoma, metastatic to the lung and kidney and who received three doses of an exemplary anti-TIM-3 antibody (TSR-022) at 10 mg/kg prior to re-staging imaging.

[00138] **Figure 10** depicts a receptor occupancy study of a 100 mg flat dose of an exemplary anti-TIM-3 antibody (TSR-022) administered once every three weeks (Q3W).

[00139] **Figure 11** depicts a receptor occupancy study of a 300 mg flat dose of an exemplary anti-TIM-3 antibody (TSR-022) administered in combination with a 500 mg flat dose of an exemplary anti-PD-1 antibody (TSR-042). A second dose of TSR-022 was administered on day 22, with the RO sample collected prior to the second dose.

[00140] **Figure 12** is a composite of mean receptor occupancy data for doses of an exemplary anti-TIM-3 antibody (TSR-022) of 1 mg/kg, 3 mg/kg, 10 mg/kg and of flat doses of 100 mg, 300 mg, and 1200 mg. The figure shows the occupancy ratio (free TIM-3:total TIM-3) as measured over a range of days.

DETAILED DESCRIPTION OF CERTAIN EMBODIMENTS

Definitions

[00141] **About:** The term “about”, when used herein in reference to a value, refers to a value that is similar, in context to the referenced value. In general, those skilled in the art, familiar with the context, will appreciate the relevant degree of variance encompassed by “about” in that context. For example, in some embodiments, the term “about” may encompass a range of values that within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction of the referred value.

[00142] **Administration:** As used herein, the term “administration” typically refers to the administration of a composition to a subject or system to achieve delivery of an agent that is, or is included in, the composition. Those of ordinary skill in the art will be aware of a variety of routes that may, in appropriate circumstances, be utilized for administration to a subject, for example a human. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (i.e., topical), transmucosal, and rectal administration. For example, in some embodiments, administration may be ocular, oral, parenteral, topical, etc. In embodiments, administration is parenteral (e.g., intravenous administration). In embodiments, intravenous administration is intravenous infusion. In some particular embodiments, administration may be bronchial (e.g., by bronchial instillation), buccal, dermal (which may be or comprise, for example, one or more of topical to the dermis,

intradermal, interdermal, transdermal, etc), enteral, intra-arterial, intradermal, intragastric, intramedullary, intramuscular, intranasal, intraperitoneal, intrathecal, intravenous, intraventricular, within a specific organ (e. g. intrahepatic), mucosal, nasal, oral, rectal, subcutaneous, sublingual, topical, tracheal (e.g., by intratracheal instillation), vaginal, vitreal, etc. In some embodiments, administration may involve only a single dose. In some embodiments, administration may involve application of a fixed number of doses. In some embodiments, administration may involve dosing that is intermittent (e.g., a plurality of doses separated in time) and/or periodic (e.g., individual doses separated by a common period of time) dosing. In some embodiments, administration may involve continuous dosing (e.g., perfusion) for at least a selected period of time.

[00143] Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid (EDTA); buffers such as acetates, citrates or phosphates, and agents for the adjustment of tonicity such as sodium chloride or dextrose. The pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

[00144] For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

[00145] Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

[00146] The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

[00147] *Affinity*: As is known in the art, “affinity” is a measure of the tightness with a particular ligand binds to its partner. Affinities can be measured in different ways. In some embodiments, affinity is measured by a quantitative assay. In some such embodiments, binding partner concentration may be fixed to be in excess of ligand concentration so as to mimic physiological conditions. Alternatively or additionally, in some embodiments, binding partner concentration and/or ligand concentration may be varied. In some such embodiments, affinity may be compared to a reference under comparable conditions (e.g., concentrations).

[00148] *Antibody*: As used herein, the term “antibody” refers to a polypeptide that includes canonical immunoglobulin sequence elements sufficient to confer specific binding to a particular target antigen. As is known in the art, intact antibodies as produced in nature are approximately 150 kD tetrameric agents comprised of two identical heavy chain polypeptides (about 50 kD each) and two identical light chain polypeptides (about 25 kD each) that associate with each other into what is commonly referred to as a “Y-shaped” structure. Each heavy chain is comprised of at least four domains (each about 110 amino acids long)— an amino-terminal variable (VH) domain (located at the tips of the Y structure), followed by three constant domains: CH1, CH2, and the carboxy-terminal CH3 (located at the base of the Y’s stem). A short region, known as the “switch”, connects the heavy chain variable and constant regions. The “hinge” connects CH2 and CH3 domains to the rest of the antibody. Two disulfide bonds in this hinge region connect the two heavy chain polypeptides to one another in an intact antibody. Each light chain is comprised of two domains – an amino-terminal variable (VL) domain, followed by a carboxy-terminal constant (CL) domain, separated from one another by another “switch”. Those skilled in the art are well familiar with antibody structure and sequence elements, recognize “variable” and “constant” regions in provided sequences, and understand that there may be some flexibility in definition of a “boundary” between such domains such that different presentations of the same antibody chain sequence may, for example, indicate such a boundary at a location that is shifted one or a few residues relative to a different presentation of the same antibody chain sequence. Intact antibody tetramers are comprised of two heavy chain-light chain dimers in which the heavy and light chains are linked to one another by a single

disulfide bond; two other disulfide bonds connect the heavy chain hinge regions to one another, so that the dimers are connected to one another and the tetramer is formed. Naturally-produced antibodies are also glycosylated, typically on the CH2 domain. Each domain in a natural antibody has a structure characterized by an “immunoglobulin fold” formed from two beta sheets (e.g., 3-, 4-, or 5-stranded sheets) packed against each other in a compressed antiparallel beta barrel. Each variable domain contains three hypervariable loops known as “complementarity determining regions” (CDR1, CDR2, and CDR3) and four somewhat invariant “framework” regions (FR1, FR2, FR3, and FR4). When natural antibodies fold, the FR regions form the beta sheets that provide the structural framework for the domains, and the CDR loop regions from both the heavy and light chains are brought together in three-dimensional space so that they create a single hypervariable antigen binding site located at the tip of the Y structure. The Fc region of naturally-occurring antibodies binds to elements of the complement system, and also to receptors on effector cells, including for example effector cells that mediate cytotoxicity. As is known in the art, affinity and/or other binding attributes of Fc regions for Fc receptors can be modulated through glycosylation or other modification. In some embodiments, antibodies produced and/or utilized in accordance with the present invention include glycosylated Fc domains, including Fc domains with modified or engineered such glycosylation. For purposes of the present invention, in certain embodiments, any polypeptide or complex of polypeptides that includes sufficient immunoglobulin domain sequences as found in natural antibodies can be referred to and/or used as an “antibody”, whether such polypeptide is naturally produced (e.g., generated by an organism reacting to an antigen), or produced by recombinant engineering, chemical synthesis, or other artificial system or methodology. In some embodiments, an antibody is polyclonal; in some embodiments, an antibody is monoclonal. In some embodiments, an antibody has constant region sequences that are characteristic of mouse, rabbit, primate, or human antibodies. In some embodiments, antibody sequence elements are humanized, primatized, chimeric, etc, as is known in the art. Moreover, the term “antibody” as used herein, can refer in appropriate embodiments (unless otherwise stated or clear from context) to any of the art-known or developed constructs or formats for utilizing antibody structural and functional features in alternative presentation. For example, in some embodiments, an antibody utilized in accordance with the present invention is in a format selected from, but not limited to, intact IgA, IgG, IgE or IgM antibodies; bi- or multi- specific antibodies (e.g., Zybodies®, etc);

antibody fragments such as Fab fragments, Fab' fragments, F(ab')₂ fragments, Fd' fragments, Fd fragments, and isolated CDRs or sets thereof; single chain Fvs; polypeptide-Fc fusions; single domain antibodies (e.g., shark single domain antibodies such as IgNAR or fragments thereof); cameloid antibodies; masked antibodies (e.g., Probodies®); Small Modular ImmunoPharmaceuticals (“SMIPsTM”); single chain or Tandem diabodies (TandAb®); VHHs; Anticalins®; Nanobodies® minibodies; BiTE®s; ankyrin repeat proteins or DARPINs®; Avimers®; DARTs; TCR-like antibodies; Adnectins®; Affilins®; Trans-bodies®; Affibodies®; TrimerX®; MicroProteins; Fynomers®, Centyrins®, and KALBITOR®s. In some embodiments, an antibody may lack a covalent modification (e.g., attachment of a glycan) that it would have if produced naturally. In some embodiments, an antibody may contain a covalent modification (e.g., attachment of a glycan, a payload [e.g., a detectable moiety, a therapeutic moiety, a catalytic moiety, etc], or other pendant group [e.g., poly-ethylene glycol, etc].

[00149] Antibodies include antibody fragments. Antibodies also include, but are not limited to, polyclonal monoclonal, chimeric dAb (domain antibody), single chain, F_{ab}, F_{ab}', F_(ab')₂ fragments, scFvs, and F_{ab} expression libraries. An antibody may be a whole antibody, or immunoglobulin, or an antibody fragment.

[00150] ***Antibody agent:*** As used herein, the term “antibody agent” refers to an agent that specifically binds to a particular antigen. In some embodiments, the term encompasses any polypeptide or polypeptide complex that includes immunoglobulin structural elements sufficient to confer specific binding. Exemplary antibody agents include, but are not limited to monoclonal antibodies or polyclonal antibodies. In some embodiments, an antibody agent may include one or more constant region sequences that are characteristic of mouse, rabbit, primate, or human antibodies. In some embodiments, an antibody agent may include one or more sequence elements are humanized, primatized, chimeric, etc, as is known in the art. In many embodiments, the term “antibody agent” is used to refer to one or more of the art-known or developed constructs or formats for utilizing antibody structural and functional features in alternative presentation. For example, embodiments, an antibody agent utilized in accordance with the present invention is in a format selected from, but not limited to, intact IgA, IgG, IgE or IgM antibodies; bi- or multi- specific antibodies (e.g., Zybodies®, etc); antibody fragments such as Fab fragments, Fab' fragments, F(ab')₂ fragments, Fd' fragments, Fd fragments, and isolated CDRs or sets thereof; single chain Fvs; polypeptide-Fc fusions; single domain antibodies (e.g.,

shark single domain antibodies such as IgNAR or fragments thereof); cameloid antibodies; masked antibodies (e.g., Probodies®); Small Modular ImmunoPharmaceuticals (“SMIPsTM”); single chain or Tandem diabodies (TandAb®); VHHs; Anticalins®; Nanobodies® minibodies; BiTE®s; ankyrin repeat proteins or DARPINs®; Avimers®; DARTs; TCR-like antibodies; Adnectins®; Affilins®; Trans-bodies®; Affibodies®; TrimerX®; MicroProteins; Fynomers®, Centyrins®, and KALBITOR®s. In some embodiments, an antibody may lack a covalent modification (e.g., attachment of a glycan) that it would have if produced naturally. In some embodiments, an antibody may contain a covalent modification (e.g., attachment of a glycan, a payload [e.g., a detectable moiety, a therapeutic moiety, a catalytic moiety, etc], or other pendant group [e.g., poly-ethylene glycol, etc.]. In many embodiments, an antibody agent is or comprises a polypeptide whose amino acid sequence includes one or more structural elements recognized by those skilled in the art as a complementarity determining region (CDR); in some embodiments, an antibody agent is or comprises a polypeptide whose amino acid sequence includes at least one CDR (e.g., at least one heavy chain CDR and/or at least one light chain CDR) that is substantially identical to one found in a reference antibody. In some embodiments, an included CDR is substantially identical to a reference CDR in that it is either identical in sequence or contains between 1-5 amino acid substitutions as compared with the reference CDR. In some embodiments, an included CDR is substantially identical to a reference CDR in that it shows at least 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 100% sequence identity with the reference CDR. In some embodiments, an included CDR is substantially identical to a reference CDR in that it shows at least 96%, 96%, 97%, 98%, 99%, or 100% sequence identity with the reference CDR. In some embodiments, an included CDR is substantially identical to a reference CDR in that at least one amino acid within the included CDR is deleted, added, or substituted as compared with the reference CDR but the included CDR has an amino acid sequence that is otherwise identical with that of the reference CDR. In some embodiments, an included CDR is substantially identical to a reference CDR in that 1-5 amino acids within the included CDR are deleted, added, or substituted as compared with the reference CDR but the included CDR has an amino acid sequence that is otherwise identical to the reference CDR. In some embodiments, an included CDR is substantially identical to a reference CDR in that at least one amino acid within the included CDR is substituted as compared with the reference CDR but the included CDR has an amino acid

sequence that is otherwise identical with that of the reference CDR. In some embodiments, an included CDR is substantially identical to a reference CDR in that 1-5 amino acids within the included CDR are deleted, added, or substituted as compared with the reference CDR but the included CDR has an amino acid sequence that is otherwise identical to the reference CDR. In some embodiments, an antibody agent is or comprises a polypeptide whose amino acid sequence includes structural elements recognized by those skilled in the art as an immunoglobulin variable domain. In some embodiments, an antibody agent is a polypeptide protein having a binding domain which is homologous or largely homologous to an immunoglobulin-binding domain.

[00151] Binding: It will be understood that the term “binding”, as used herein, typically refers to a non-covalent association between or among two or more entities. “Direct” binding involves physical contact between entities or moieties; indirect binding involves physical interaction by way of physical contact with one or more intermediate entities. Binding between two or more entities can typically be assessed in any of a variety of contexts – including where interacting entities or moieties are studied in isolation or in the context of more complex systems (e.g., while covalently or otherwise associated with a carrier entity and/or in a biological system or cell). In some embodiments, “binding” refers to the non-covalent interactions of the type which occur between an immunoglobulin molecule and an antigen for which the immunoglobulin is specific. The strength, or affinity of immunological binding interactions can be expressed in terms of the dissociation constant (K_d) of the interaction, wherein a smaller K_d represents a greater affinity. Immunological binding properties of selected polypeptides can be quantified using methods well known in the art. One such method entails measuring the rates of antigen-binding site/antigen complex formation and dissociation, wherein those rates depend on the concentrations of the complex partners, the affinity of the interaction, and geometric parameters that equally influence the rate in both directions. Thus, both the “on rate constant” (K_{on}) and the “off rate constant” (K_{off}) can be determined by calculation of the concentrations and the actual rates of association and dissociation. (*See* Nature 361:186-87 (1993)). The ratio of K_{off}/K_{on} enables the cancellation of all parameters not related to affinity, and is equal to the dissociation constant K_d . (*See, generally,* Davies et al. (1990) Annual Rev Biochem 59:439-473).

[00152] Binding agent: In general, the term “binding agent” is used herein to refer to any entity that binds to a target of interest as described herein. In many embodiments, a binding

agent of interest is one that binds specifically with its target in that it discriminates its target from other potential binding partners in a particular interaction context. In general, a binding agent may be or comprise an entity of any chemical class (e.g., polymer, non-polymer, small molecule, polypeptide, carbohydrate, lipid, nucleic acid, etc). In some embodiments, a binding agent is a single chemical entity. In some embodiments, a binding agent is a complex of two or more discrete chemical entities associated with one another under relevant conditions by non-covalent interactions. For example, those skilled in the art will appreciate that in some embodiments, a binding agent may comprise a “generic” binding moiety (e.g., one of biotin/avidin/streptavidin and/or a class-specific antibody) and a “specific” binding moiety (e.g., an antibody or aptamers with a particular molecular target) that is linked to the partner of the generic binding moiety. In some embodiments, such an approach can permit modular assembly of multiple binding agents through linkage of different specific binding moieties with the same generic binding moiety partner. In some embodiments, binding agents are or comprise polypeptides (including, e.g., antibodies or antibody fragments). In some embodiments, binding agents are or comprise small molecules. In some embodiments, binding agents are or comprise nucleic acids. In some embodiments, binding agents are aptamers. In some embodiments, binding agents are polymers; in some embodiments, binding agents are not polymers. In some embodiments, binding agents are non-polymeric in that they lack polymeric moieties. In some embodiments, binding agents are or comprise carbohydrates. In some embodiments, binding agents are or comprise lectins. In some embodiments, binding agents are or comprise peptidomimetics. In some embodiments, binding agents are or comprise scaffold proteins. In some embodiments, binding agents are or comprise mimeotopes. In some embodiments, binding agents are or comprise nucleic acids, such as DNA or RNA.

[00153] *Cancer*: The terms “cancer”, “malignancy”, “neoplasm”, “tumor”, and “carcinoma”, are used herein to refer to cells that exhibit relatively abnormal, uncontrolled, and/or autonomous growth, so that they exhibit an aberrant growth phenotype characterized by a significant loss of control of cell proliferation. In some embodiments, a tumor may be or comprise cells that are precancerous (e.g., benign), malignant, pre-metastatic, metastatic, and/or non-metastatic. The present disclosure specifically identifies certain cancers to which its teachings may be particularly relevant. In some embodiments, a relevant cancer may be characterized by a solid tumor. In some embodiments, a relevant cancer may be characterized by a hematologic tumor.

In general, examples of different types of cancers known in the art include, for example, hematopoietic cancers including leukemias, lymphomas (Hodgkin's and non-Hodgkin's), myelomas and myeloproliferative disorders; sarcomas, melanomas, adenomas, carcinomas of solid tissue, squamous cell carcinomas of the mouth, throat, larynx, and lung, liver cancer, genitourinary cancers such as prostate, cervical, bladder, uterine, and endometrial cancer and renal cell carcinomas, bone cancer, pancreatic cancer, skin cancer, cutaneous or intraocular melanoma, cancer of the endocrine system, cancer of the thyroid gland, cancer of the parathyroid gland, head and neck cancers, breast cancer, gastro-intestinal cancers and nervous system cancers, benign lesions such as papillomas, and the like.

[00154] *Carrier:* as used herein, refers to a diluent, adjuvant, excipient, or vehicle with which a composition is administered. In some exemplary embodiments, carriers can include sterile liquids, such as, for example, water and oils, including oils of petroleum, animal, vegetable or synthetic origin, such as, for example, peanut oil, soybean oil, mineral oil, sesame oil and the like. In some embodiments, carriers are or include one or more solid components. In some embodiments, the carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

[00155] *Combination therapy:* As used herein, the term "combination therapy" refers to a clinical intervention in which a subject is simultaneously exposed to two or more therapeutic regimens (e.g., two or more therapeutic agents). In some embodiments, the two or more therapeutic regimens may be administered simultaneously. In some embodiments, the two or more therapeutic regimens may be administered sequentially (e.g., a first regimen administered prior to administration of any doses of a second regimen). In some embodiments, the two or

more therapeutic regimens are administered in overlapping dosing regimens. In some embodiments, administration of combination therapy may involve administration of one or more therapeutic agents or modalities to a subject receiving the other agent(s) or modality. In some embodiments, combination therapy does not necessarily require that individual agents be administered together in a single composition (or even necessarily at the same time). In some embodiments, two or more therapeutic agents or modalities of a combination therapy are administered to a subject separately, *e.g.*, in separate compositions, via separate administration routes (*e.g.*, one agent orally and another agent intravenously), and/or at different time points. In some embodiments, two or more therapeutic agents may be administered together in a combination composition, or even in a combination compound (*e.g.*, as part of a single chemical complex or covalent entity), via the same administration route, and/or at the same time.

[00156] Complete Response: As used herein, the term “complete response” or “CR” is used to mean the disappearance of all or substantially all target lesions. In some embodiments, CR refers to an about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% decrease in the sum of the diameters of the target lesions (*i.e.*, loss of lesions), taking as reference the baseline sum diameters. In some embodiments, CR indicates that less than about 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1% or less of the total lesion diameter remains after treatment. Exemplary methods for evaluating complete response are identified by RECIST guidelines. See, *e.g.*, E.A. Eisenhauer, *et al.*, “New response evaluation criteria in solid tumors: Revised RECIST guideline (version 1.1.),” *Eur. J. of Cancer*, 45: 228-247 (2009).

[00157] Dosage form or unit dosage form: Those skilled in the art will appreciate that the term “dosage form” may be used to refer to a physically discrete unit of an active agent (*e.g.*, a therapeutic or diagnostic agent) for administration to a subject. Typically, each such unit contains a predetermined quantity of active agent. In some embodiments, such quantity is a unit dosage amount (or a whole fraction thereof) appropriate for administration in accordance with a dosing regimen that has been determined to correlate with a desired or beneficial outcome when administered to a relevant population (*i.e.*, with a therapeutic dosing regimen). Those of ordinary skill in the art appreciate that the total amount of a therapeutic composition or agent administered to a particular subject is determined by one or more attending physicians and may involve administration of multiple dosage forms.

[00158] *Dosing regimen or regimen:* Those skilled in the art will appreciate that the term “regimen” may be used to refer to a set of unit doses (typically more than one) that are administered individually to a subject, typically separated by periods of time. In some embodiments, a given therapeutic agent has a recommended dosing regimen, which may involve one or more doses. In some embodiments, a dosing regimen comprises a plurality of doses each of which is separated in time from other doses. In some embodiments, individual doses are separated from one another by a time period of the same length; in some embodiments, a dosing regimen comprises a plurality of doses and at least two different time periods separating individual doses. In some embodiments, all doses within a dosing regimen are of the same unit dose amount. In some embodiments, different doses within a dosing regimen are of different amounts. In some embodiments, a dosing regimen comprises a first dose in a first dose amount, followed by one or more additional doses in a second dose amount different from the first dose amount. In some embodiments, a dosing regimen comprises a first dose in a first dose amount, followed by one or more additional doses in a second dose amount same as the first dose amount. In some embodiments, a dosing regimen is correlated with a desired or beneficial outcome when administered across a relevant population (i.e., is a therapeutic dosing regimen). In some embodiments, a regimen comprises at least one dose, wherein the dose comprises one unit dose of a therapeutic agent (e.g., an anti-TIM-3 antibody agent). In some embodiments, a regimen comprises at least one dose, wherein the dose comprises two or more unit doses of a therapeutic agent. For example, a dose of 500 mg can be administered as a single 500 mg unit dose or as two 250 mg unit doses. In some embodiments, a regimen is correlated with or result in a desired or beneficial outcome when administered across a relevant population (i.e., is a therapeutic regimen).

[00159] *Hazard Ratio:* As used herein, a “hazard ratio” is the expression of the hazard or chance of events occurring in the treatment arm as a ratio of the events occurring in the control arm. Hazard ratios may be determined by the Cox model, a regression method for survival data, which provides an estimate of the hazard ratio and its confidence interval. The hazard ratio is an estimate of the ratio of the hazard rate in the treated versus the control group. The hazard rate is the probability that if the event in question has not already occurred, it will occur in the next time interval, divided by the length of that interval. An assumption of proportional hazards regression is that the hazard ratio is constant over time.

[00160] *Homology*: As used herein, the term “homology” refers to the overall relatedness between polymeric molecules, e.g., between nucleic acid molecules (e.g., DNA molecules and/or RNA molecules) and/or between polypeptide molecules. In some embodiments, polymeric molecules are considered to be “homologous” to one another if their sequences are at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% identical. In some embodiments, polymeric molecules are considered to be “homologous” to one another if their sequences are at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 99% similar (e.g., containing residues with related chemical properties at corresponding positions). For example, as is well known by those of ordinary skill in the art, certain amino acids are typically classified as similar to one another as “hydrophobic” or “hydrophilic” amino acids, and/or as having “polar” or “non-polar” side chains. Substitution of one amino acid for another of the same type may often be considered a “homologous” substitution.

[00161] As will be understood by those skilled in the art, a variety of algorithms are available that permit comparison of sequences in order to determine their degree of homology, including by permitting gaps of designated length in one sequence relative to another when considering which residues “correspond” to one another in different sequences. Calculation of the percent homology between two nucleic acid sequences, for example, can be performed by aligning the two sequences for optimal comparison purposes (e.g., gaps can be introduced in one or both of a first and a second nucleic acid sequences for optimal alignment and non-corresponding sequences can be disregarded for comparison purposes). In certain embodiments, the length of a sequence aligned for comparison purposes is at least 30%, at least 40%, at least 50%, at least 60%, at least 70%, at least 80%, at least 90%, at least 95%, or substantially 100% of the length of the reference sequence. The nucleotides at corresponding nucleotide positions are then compared. When a position in the first sequence is occupied by the same nucleotide as the corresponding position in the second sequence, then the molecules are identical at that position; when a position in the first sequence is occupied by a similar nucleotide as the corresponding position in the second sequence, then the molecules are similar at that position. The percent homology between the two sequences is a function of the number of identical and similar positions shared by the sequences, taking into account the number of gaps, and the length of each gap, which needs to be introduced for optimal alignment of the two sequences. Representative

algorithms and computer programs useful in determining the percent homology between two nucleotide sequences include, for example, the algorithm of Meyers and Miller (CABIOS, 1989, 4: 11-17), which has been incorporated into the ALIGN program (version 2.0) using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4. The percent homology between two nucleotide sequences can, alternatively, be determined for example using the GAP program in the GCG software package using an NWSgapdna.CMP matrix

[00162] K_D : as used herein, refers to the dissociation constant of a binding agent (e.g., an antibody or binding component thereof) from a complex with its partner (e.g., the epitope to which the antibody or binding component thereof binds).

[00163] K_{off} : as used herein, refers to the off rate constant for dissociation of a binding agent (e.g., an antibody or binding component thereof) from a complex with its partner (e.g., the epitope to which the antibody or binding component thereof binds).

[00164] K_{on} : as used herein, refers to the on rate constant for association of a binding agent (e.g., an antibody or binding component thereof) with its partner (e.g., the epitope to which the antibody or binding component thereof binds).

[00165] *Patient or subject*: As used herein, the term “patient” or “subject” refers to any organism to which provided compound or compounds described herein are administered in accordance with the present invention e.g., for experimental, diagnostic, prophylactic, and/or therapeutic purposes. Typical subjects include animals. The term “animal” refers to any member of the animal kingdom. In some embodiments, “animal” refers to humans, at any stage of development. In some embodiments, “animal” refers to non-human animals, at any stage of development. In certain embodiments, the non-human animal is a mammal (e.g., a rodent, a mouse, a rat, a rabbit, a monkey, a dog, a cat, a sheep, cattle, a primate, and/or a pig). In some embodiments, animals include, but are not limited to, mammals, birds, reptiles, amphibians, fish, insects, and/or worms. In some embodiments, an animal may be a transgenic animal, genetically-engineered animal, and/or a clone. In embodiments, animals are mammals such as mice, rats, rabbits, non-human primates, and humans; insects; worms; etc. In a preferred embodiment, a subject is a human. In some embodiments, a subject may be suffering from, and/or susceptible to a disease, disorder, and/or condition (e.g., cancer). As used herein, a “patient population” or “population of subjects” refers to a plurality of patients or subjects.

[00166] *Partial Response:* As used herein, the term “partial response” (“PR”) refers to a decrease in tumor progression in a subject as indicated by a decrease in the sum of the diameters of the target lesions, taking as reference the baseline sum diameters. In some embodiments, PR refers to at least a 30% decrease in the sum of diameters or target lesions, taking as reference the baseline sum diameters. Exemplary methods for evaluating partial response are identified by RECIST guidelines. See e.g., E.A. Eisenhauer, *et al.*, “New response evaluation criteria in solid tumors: Revised RECIST guideline (version 1.1.),” *Eur. J. of Cancer*, 45: 228-247 (2009).

[00167] *Pharmaceutical composition:* As used herein, the term “pharmaceutical composition” refers to a composition in which an active agent (e.g., an anti-TIM-3 antibody agent and/or a PD-1-binding agent) is formulated together with one or more pharmaceutically acceptable carriers. In some embodiments, the active agent is present in unit dose amount appropriate for administration in a therapeutic regimen that shows a statistically significant probability of achieving a predetermined therapeutic effect when administered to a relevant population. In some embodiments, a pharmaceutical composition may be specially formulated for administration in solid or liquid form, including those adapted for the following: oral administration, for example, drenches (aqueous or non-aqueous solutions or suspensions), tablets, e.g., those targeted for buccal, sublingual, and systemic absorption, boluses, powders, granules, pastes for application to the tongue; parenteral administration, for example, by subcutaneous, intramuscular, intravenous or epidural injection as, for example, a sterile solution or suspension, or sustained-release formulation; topical application, for example, as a cream, ointment, or a controlled-release patch or spray applied to the skin, lungs, or oral cavity; intravaginally or intrarectally, for example, as a pessary, cream, or foam; sublingually; ocularly; transdermally; or nasally, pulmonary, and to other mucosal surfaces. In some preferred embodiments, an active agent (e.g., an anti-TIM-3 antibody agent and/or a PD-1-binding agent) is formulated for parenteral administration.

[00168] *Pharmaceutically acceptable:* As used herein, the term “pharmaceutically acceptable” applied to the carrier, diluent, or excipient used to formulate a composition as disclosed herein means that the carrier, diluent, or excipient must be compatible with the other ingredients of the composition and not deleterious to the recipient thereof.

[00169] *Progression Free Survival:* As used herein, the term “progression free survival” means the time period for which a subject having a disease (e.g., cancer) survives, without a

significant worsening of the disease state. Progression free survival may be assessed as a period of time in which there is no progression of tumor growth and/or wherein the disease status of a patient is not determined to be a progressive disease. In some embodiments, progression free survival of a subject having cancer is assessed by evaluating tumor (lesion) size, tumor (lesion) number, and/or metastasis.

[00170] *Progression or Progressive Disease:* The term “progression” of tumor growth or a “progressive disease” (“PD”) as used herein in reference to cancer status indicates an increase in the sum of the diameters of the target lesions (tumors). In some embodiments, progression of tumor growth refers to at least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In some embodiments, in addition to a relative increase of 20%, the sum of diameters of target lesions must also demonstrate an absolute increase of at least 5 mm. An appearance of one or more new lesions may also be factored into the determination of progression of tumor growth. Progression for the purposes of determining progression free survival may also be determined if at least one of the following criteria is met: 1) tumor assessment by CT/MRI unequivocally shows progressive disease according to RECIST 1.1 or irRECIST criteria; or 2) additional diagnostic tests (e.g., histology/cytology, ultrasound techniques, endoscopy, positron emission tomography) identify new lesions or determine existing lesions qualify for unequivocal progressive disease AND CA-125- progression according to Gynecologic Cancer Intergroup (GCIG)-criteria (see Rustin et al., Int J Gynecol Cancer 2011;21: 419-423 which is incorporated herein in its entirety); 3) definitive clinical signs and symptoms of PD unrelated to non-malignant or iatrogenic causes ([i] intractable cancer-related pain; [ii] malignant bowel obstruction/worsening dysfunction; or [iii] unequivocal symptomatic worsening of ascites or pleural effusion) AND CA-125-progression according to GCIG-criteria.

[00171] *Solid Tumor:* As used herein, the term “solid tumor” refers to an abnormal mass of tissue that usually does not contain cysts or liquid areas. In some embodiments, a solid tumor may be benign; in some embodiments, a solid tumor may be malignant. Those skilled in the art will appreciate that different types of solid tumors are typically named for the type of cells that form them. Examples of solid tumors are carcinomas, lymphomas, and sarcomas. In some embodiments, solid tumors may be or comprise adrenal, bile duct, bladder, bone, brain, breast, cervix, colon, endometrium, esophagus, eye, gall bladder, gastrointestinal tract, kidney, larynx,

liver, lung, nasal cavity, nasopharynx, oral cavity, ovary, penis, pituitary, prostate, retina, salivary gland, skin, small intestine, stomach, testis, thymus, thyroid, uterine, vaginal, and/or vulval tumors.

[00172] *Stabilization or Stable Disease:* As used herein, “stabilization” of tumor growth or a “stable disease” (“SD”) refers to neither sufficient shrinkage to qualify for PR nor sufficient increase to qualify for PD. In some embodiments, stabilization refers to a less than 30%, 25%, 20%, 15%, 10% or 5% change (increase or decrease) in the sum of the diameters of the target lesions, taking as reference the baseline sum diameters. Exemplary methods for evaluating stabilization of tumor growth or a stable disease are identified by RECIST guidelines. See e.g., E.A. Eisenhauer, *et al.*, “New response evaluation criteria in solid tumors: Revised RECIST guideline (version 1.1.),” *Eur. J. of Cancer*, 45: 228-247 (2009).

[00173] *Therapeutically Effective Amount:* As used herein, is meant an amount that produces the desired effect for which it is administered. In some embodiments, the term refers to an amount that is sufficient, when administered to a population suffering from or susceptible to a disease, disorder, and/or condition in accordance with a therapeutic dosing regimen, to treat the disease, disorder, and/or condition. In some embodiments, a therapeutically effective amount is one that reduces the incidence and/or severity of, and/or delays onset of, one or more symptoms of the disease, disorder, and/or condition. Those of ordinary skill in the art will appreciate that the term “*therapeutically effective amount*” does not in fact require successful treatment be achieved in a particular individual. Rather, a therapeutically effective amount may be that amount that provides a particular desired pharmacological response in a significant number of subjects when administered to patients in need of such treatment. In some embodiments, reference to a therapeutically effective amount may be a reference to an amount as measured in one or more specific tissues (e.g., a tissue affected by the disease, disorder or condition) or fluids (e.g., blood, saliva, serum, sweat, tears, urine, etc.). Those of ordinary skill in the art will appreciate that, in some embodiments, a therapeutically effective amount of a particular agent or therapy may be formulated and/or administered in a single dose. In some embodiments, a therapeutically effective agent may be formulated and/or administered in a plurality of doses, for example, as part of a dosing regimen.

[00174] *Treatment:* As used herein, the term “treatment” (also “treat” or “treating”) refers to any administration of a therapy that partially or completely alleviates, ameliorates, relieves,

inhibits, delays onset of, reduces severity of, and/or reduces incidence of one or more symptoms, features, and/or causes of a particular disease, disorder, and/or condition. In some embodiments, such treatment may be of a subject who does not exhibit signs of the relevant disease, disorder and/or condition and/or of a subject who exhibits only early signs of the disease, disorder, and/or condition. Alternatively or additionally, such treatment may be of a subject who exhibits one or more established signs of the relevant disease, disorder and/or condition. In some embodiments, treatment may be of a subject who has been diagnosed as suffering from the relevant disease, disorder, and/or condition. In some embodiments, treatment may be of a subject known to have one or more susceptibility factors that are statistically correlated with increased risk of development of the relevant disease, disorder, and/or condition.

Methods of Treatment, Including Methods of Treating Cancer

[00175] Described herein are methods of treating disorders in a subject (e.g., disorders that benefit from administration of an anti-TIM-3 therapy). For example, an anti-TIM-3 therapy described herein can agent is administered e.g., as a monotherapy or in combination therapy, for a period sufficient to achieve clinical benefit or according to a regimen as determined by a physician (e.g., an anti-TIM-3 therapy is administered in dosage amounts and number of treatment cycles as determined by a physician).

[00176] In embodiments, methods described herein are useful for increasing T cell activation or T cell effector function in a subject.

[00177] In embodiments, methods described herein are useful for inducing an immune response in a subject.

[00178] In embodiments, methods described herein are useful for enhancing an immune response or increasing the activity of an immune cell in a subject.

[00179] The inventive methods can be used to treat any type of infectious disease (i.e., a disease or disorder caused by a bacterium, a virus, a fungus, or a parasite). Examples of infectious diseases that can be treated by the inventive method include, but are not limited to, diseases caused by a human immunodeficiency virus (HIV), a respiratory syncytial virus (RSV), an influenza virus, a dengue virus, a hepatitis B virus (HBV, or a hepatitis C virus (HCV)). When the inventive method treats an infectious disease, an anti-TIM-3 antibody agent can be administered in combination with at least one anti-bacterial agent or at least one anti-viral agent.

In this respect, the anti-bacterial agent can be any suitable antibiotic known in the art. The anti-viral agent can be any vaccine of any suitable type that specifically targets a particular virus (e.g., live-attenuated vaccines, subunit vaccines, recombinant vector vaccines, and small molecule anti-viral therapies (e.g., viral replication inhibitors and nucleoside analogs).

[00180] The inventive methods can be used to treat any type of autoimmune disease (i.e., as disease or disorder caused by immune system over-activity in which the body attacks and damages its own tissues), such as those described in, for example, MacKay I.R. and Rose N.R., eds., *The Autoimmune Diseases, Fifth Edition*, Academic Press, Waltham, MA (2014). Examples of autoimmune diseases that can be treated by the inventive method include, but are not limited to, multiple sclerosis, type 1 diabetes mellitus, rheumatoid arthritis, scleroderma, Crohn's disease, psoriasis, systemic lupus erythematosus (SLE), and ulcerative colitis. When the inventive method treats an autoimmune disease, an anti-TIM-3 antibody agent can be used in combination with an anti-inflammatory agent including, for example, corticosteroids (e.g., prednisone and fluticasone) and non-steroidal anti-inflammatory drugs (NSAIDs) (e.g., aspirin, ibuprofen, and naproxen).

[00181] In embodiments, methods described herein are useful for treating T-cell dysfunctional disorders (e.g., cancer).

[00182] In embodiments, methods described herein are useful for reducing tumors or inhibiting the growth of tumor cells in a subject.

[00183] The inventive methods can be used to treat any type of cancer known in the art.

[00184] In embodiments, a cancer is adenocarcinoma, adenocarcinoma of the lung, acute myeloid leukemia ("AML"), acute lymphoblastic leukemia ("ALL"), adrenocortical carcinoma, anal cancer, appendiceal cancer, B-cell derived leukemia, B-cell derived lymphoma, bladder cancer, brain cancer, breast cancer (e.g., triple negative breast cancer (TNBC)), cancer of the fallopian tube(s), cancer of the testes, cerebral cancer, cervical cancer, choriocarcinoma, chronic myelogenous leukemia, a CNS tumor, colon adenocarcinoma, colon cancer, colorectal cancer, diffuse intrinsic pontine glioma (DIPG), diffuse large B cell lymphoma ("DLBCL"), embryonal rhabdomyosarcoma (ERMS), endometrial cancer, epithelial cancer, esophageal cancer, Ewing's sarcoma, follicular lymphoma ("FL"), gall bladder cancer, gastric cancer, gastrointestinal cancer, glioma, head and neck cancer, a hematological cancer, hepatocellular cancer, Hodgkin's lymphoma (HL)/primary mediastinal B-cell lymphoma, kidney cancer, kidney clear cell cancer,

laryngeal cancer, leukemia, liver cancer, lung cancer, lymphoma, melanoma, Merkel cell carcinoma, mesothelioma, monocytic leukemia, multiple myeloma, myeloma, a neuroblastic-derived CNS tumor (e.g., neuroblastoma (NB)), non-Hodgkin's lymphoma (NHL), non-small cell lung cancer (NSCLC), oral cancer, osteosarcoma, ovarian cancer, ovarian carcinoma, pancreatic cancer, peritoneal cancer, primary peritoneal cancer, prostate cancer, relapsed or refractory classic Hodgkin's Lymphoma (cHL), renal cell carcinoma, rectal cancer, salivary gland cancer (e.g., a salivary gland tumor), sarcoma, skin cancer, small cell lung cancer, small intestine cancer, squamous cell carcinoma of the anogenital region (e.g., squamous cell carcinoma of the anus, penis, cervix, vagina, or vulva), squamous cell carcinoma of the esophagus, squamous cell carcinoma of the head and neck (SCHNC), squamous cell carcinoma of the lung, stomach cancer, T-cell derived leukemia, T-cell derived lymphoma, thymic cancer, a thymoma, thyroid cancer, uveal melanoma, urothelial cell carcinoma, uterine cancer, uterine endometrial cancer, uterine sarcoma, vaginal cancer, vulvar cancer, or Wilms tumor.

[00185] In other embodiments, a cancer is a head and neck cancer, a lung cancer (e.g., a non-small cell lung cancer (NSCLC)), a renal cancer, a bladder cancer, a melanoma, Merkel cell carcinoma (see, e.g., Bhatia et al., Curr. Oncol. Rep., 13(6): 488-497 (2011), a cervical cancer, a vaginal cancer, a vulvar cancer, a uterine cancer, an endometrial cancer, an ovarian cancer, a fallopian tube cancer, a breast cancer, a prostate cancer, a salivary gland tumor, a thymoma, an adrenocortical carcinoma, an esophageal cancer, a gastric cancer, a colorectal cancer, an appendiceal cancer, a urothelial cell carcinoma, or a squamous cell carcinoma (e.g., of the lung; of the anogenital region including anus, penis, cervix, vagina, or vulva; or of the esophagus). In some embodiments, a cancer for treatment in the context of the present disclosure is a melanoma, renal cell carcinoma, lung cancer, bladder cancer, breast cancer, cervical cancer, colon cancer, gall bladder cancer, laryngeal cancer, liver cancer, thyroid cancer, stomach cancer, salivary gland cancer, prostate cancer, pancreatic cancer, or Merkel cell carcinoma.

[00186] In embodiments a cancer is a lymphoma such as Hodgkin's disease, non-Hodgkin's Lymphoma, Multiple myeloma, Waldenstrom's macroglobulinemia, Heavy chain disease and Polycythemia vera.

[00187] In embodiments, a cancer is a squamous cell carcinoma. In embodiments, a cancer is squamous cell carcinoma of the lung. In embodiments, a cancer is squamous cell carcinoma of the esophagus. In embodiments, a cancer is squamous cell carcinoma of the anogenital region

(e.g., of the anus, penis, cervix, vagina, or vulva). In embodiments, a cancer is head and neck squamous cell carcinoma (HNSCC).

[00188] In embodiments, a cancer is bladder cancer, breast cancer (e.g., triple negative breast cancer (TNBC)), cancer of the fallopian tube(s), cholangiocarcinoma, colon adenocarcinoma, endometrial cancer, esophageal cancer, Ewing's sarcoma, gastric cancer, kidney clear cell cancer, lung cancer (e.g., lung adenocarcinoma or lung squamous cell cancer), mesothelioma, ovarian cancer, pancreatic cancer, peritoneal cancer, prostate cancer, uterine endometrial cancer, or uveal melanoma. In embodiments, a cancer is ovarian cancer, cancer of the fallopian tube(s), or peritoneal cancer. In embodiments, a cancer is breast cancer (e.g., TNBC). In embodiments, a cancer is lung cancer (e.g., non-small cell lung cancer). In embodiments, a cancer is prostate cancer.

[00189] In embodiments, a cancer is a CNS or brain cancer such as neuroblastoma (NB), glioma, diffuse intrinsic pontine glioma (DIPG), pilocytic astrocytoma, astrocytoma, anaplastic astrocytoma, glioblastoma multiforme, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, vestibular schwannoma, adenoma, metastatic brain tumor, meningioma, spinal tumor, or medulloblastoma. In embodiments, a cancer is a CNS tumor.

[00190] In other embodiments, a cancer is melanoma, renal cell carcinoma, lung cancer, bladder cancer, breast cancer, cervical cancer, colon cancer, gall bladder cancer, laryngeal cancer, liver cancer, thyroid cancer, stomach cancer, salivary gland cancer, prostate cancer, pancreatic cancer, or Merkel cell carcinoma (see, e.g., Bhatia et al., *Curr. Oncol. Rep.*, 13(6): 488-497 (2011)).

[00191] In some embodiments, a patient or population of patients have a hematological cancer. In some embodiments, the patient has a hematological cancer such as Diffuse large B cell lymphoma ("DLBCL"), Hodgkin's lymphoma ("HL"), Non-Hodgkin's lymphoma ("NHL"), Follicular lymphoma ("FL"), acute myeloid leukemia ("AML"), acute lymphoblastic leukemia ("ALL"), or Multiple myeloma ("MM"). In embodiments, a cancer is a blood-borne cancer such as acute lymphoblastic leukemia ("ALL"), acute lymphoblastic B-cell leukemia, acute lymphoblastic T-cell leukemia, acute myeloblastic leukemia ("AML"), acute lymphoblastic leukemia ("ALL"), acute promyelocytic leukemia ("APL"), acute monoblastic leukemia, acute erythroleukemic leukemia, acute megakaryoblastic leukemia, acute myelomonocytic leukemia,

acute nonlymphocytic leukemia, acute undifferentiated leukemia, chronic myelocytic leukemia("CML"), chronic lymphocytic leukemia("CLL"), hairy cell leukemia and multiple myeloma; acute and chronic leukemias such as lymphoblastic, myelogenous, lymphocytic, and myelocytic leukemias.

[00192] In some embodiments, a patient or population of patients have a solid tumor. In embodiments, a cancer is a solid tumor such as fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, osteosarcoma, colon cancer, colorectal cancer, kidney cancer, pancreatic cancer, bone cancer, breast cancer, ovarian cancer, prostate cancer, esophageal cancer, stomach cancer, oral cancer, nasal cancer, throat cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms tumor, cervical cancer, uterine cancer, testicular cancer, non small cell lung cancer (NSCLC), small cell lung carcinoma, bladder carcinoma, lung cancer, epithelial carcinoma, skin cancer, melanoma, neuroblastoma (NB), or retinoblastoma. In some embodiments, the tumor is an advanced stage solid tumor. In some embodiments, the tumor is a metastatic solid tumor. In some embodiments, the patient has a MSI-H solid tumor.

[00193] In some embodiments, a patient or population of patients to be treated by the methods of the present invention have or are susceptible to cancer, such as a head and neck cancer, a lung cancer (e.g., a non-small cell lung cancer (NSCLC)), a renal cancer, a bladder cancer, a melanoma, Merkel cell carcinoma, a cervical cancer, a vaginal cancer, a vulvar cancer, a uterine cancer, a endometrial cancer, an ovarian cancer, a fallopian tube cancer, a breast cancer, a prostate cancer, a salivary gland tumor, a thymoma, a adrenocortical carcinoma, a esophageal cancer, a gastric cancer, a colorectal cancer, an appendiceal cancer, a urothelial cell carcinoma, or a squamous cell carcinoma (e.g., of the lung; of the anogenital region including anus, penis, cervix, vagina, or vulva; or of the esophagus). In some embodiments, a patient or population of patients to be treated by the methods of the present invention have or are susceptible to lung cancer (e.g., NSCLC), renal cancer, melanoma, cervical cancer, colorectal cancer, or endometrial cancer (e.g., MSS endometrial cancer or MSI-H endometrial cancer).

[00194] In some embodiments, a patient or population of patients to be treated by the methods of the present invention have or are susceptible to non-small cell lung cancer (NSCLC), a hepatocellular cancer, a renal cancer, a melanoma, a cervical cancer, a colorectal cancer, a squamous cell carcinoma of the anogenital region (e.g., squamous cell carcinoma of the anus, penis, cervix, vagina, or vulva), a head and neck cancer, a triple negative breast cancer, an ovarian cancer or a endometrial cancer. In some embodiments, a patient has an advanced stage solid tumor, such as a non-small cell lung cancer (NSCLC), a hepatocellular cancer, a renal cancer, a melanoma, a cervical cancer, a colorectal cancer, a squamous cell carcinoma of the anogenital region (e.g., squamous cell carcinoma of the anus, penis, cervix, vagina, or vulva), a head and neck cancer, a triple negative breast cancer, an ovarian cancer or a endometrial cancer. In some embodiments, a patient has an advanced stage solid tumor with microsatellite instability.

[00195] In some embodiments, a cancer is a gynecologic cancer (i.e., a cancer of the female reproductive system such as ovarian cancer, fallopian tube cancer, cervical cancer, vaginal cancer, vulvar cancer, uterine cancer, or primary peritoneal cancer, or breast cancer). In some embodiments, cancers of the female reproductive system include, but are not limited to, ovarian cancer, cancer of the fallopian tube(s), peritoneal cancer, and breast cancer.

[00196] In embodiments, a cancer is ovarian cancer (e.g., serous or clear cell ovarian cancer). In embodiments, a cancer is fallopian tube cancer (e.g., serous or clear cell fallopian tube cancer). In embodiments, a cancer is primary peritoneal cancer (e.g., serous or clear cell primary peritoneal cancer).

[00197] In some embodiments, an ovarian cancer is an epithelial carcinoma. Epithelial carcinomas make up 85% to 90% of ovarian cancers. While historically considered to start on the surface of the ovary, new evidence suggests at least some ovarian cancer begins in special cells in a part of the fallopian tube. The fallopian tubes are small ducts that link a woman's ovaries to her uterus that are a part of a woman's reproductive system. In a normal female reproductive system, there are two fallopian tubes, one located on each side of the uterus. Cancer cells that begin in the fallopian tube may go to the surface of the ovary early on. The term 'ovarian cancer' is often used to describe epithelial cancers that begin in the ovary, in the fallopian tube, and from the lining of the abdominal cavity, call the peritoneum. In some embodiments, the cancer is or comprises a germ cell tumor. Germ cell tumors are a type of ovarian cancer develops in the egg-producing cells of the ovaries. In some embodiments, a cancer is or comprises a stromal tumor.

Stromal tumors develop in the connective tissue cells that hold the ovaries together, which sometimes is the tissue that makes female hormones called estrogen. In some embodiments, a cancer is or comprises a granulosa cell tumor. Granulosa cell tumors may secrete estrogen resulting in unusual vaginal bleeding at the time of diagnosis. In some embodiments, a gynecologic cancer is associated with homologous recombination repair deficiency/homologous repair deficiency (“HRD”) and/or BRCA1/2 mutation(s). In some embodiments, a gynecologic cancer is platinum-sensitive. In some embodiments, a gynecologic cancer has responded to a platinum-based therapy. In some embodiments, a gynecologic cancer has developed resistance to a platinum-based therapy. In some embodiments, a gynecologic cancer has at one time shown a partial or complete response to platinum-based therapy (e.g., a partial or complete response to the last platinum-based therapy or to the penultimate platinum-based therapy). In some embodiments, a gynecologic cancer is now resistant to platinum-based therapy.

[00198] In embodiments, a cancer is a breast cancer. Usually breast cancer either begins in the cells of the milk producing glands, known as the lobules, or in the ducts. Less commonly breast cancer can begin in the stromal tissues. These include the fatty and fibrous connective tissues of the breast. Over time the breast cancer cells can invade nearby tissues such the underarm lymph nodes or the lungs in a process known as metastasis. The stage of a breast cancer, the size of the tumor and its rate of growth are all factors which determine the type of treatment that is offered. Treatment options include surgery to remove the tumor, drug treatment which includes chemotherapy and hormonal therapy, radiation therapy and immunotherapy. The prognosis and survival rate varies widely; the five year relative survival rates vary from 98% to 23% depending on the type of breast cancer that occurs. Breast cancer is the second most common cancer in the world with approximately 1.7 million new cases in 2012 and the fifth most common cause of death from cancer, with approximately 521,000 deaths. Of these cases, approximately 15% are triple-negative, which do not express the estrogen receptor, progesterone receptor (PR) or HER2. In some embodiments, triple negative breast cancer (TNBC) is characterized as breast cancer cells that are estrogen receptor expression negative (<1% of cells), progesterone receptor expression negative (<1% of cells), and HER2-negative.

[00199] In embodiments, a cancer is ER-positive breast cancer, ER-negative breast cancer, PR-positive breast cancer, PR-negative breast cancer, HER2-positive breast cancer, HER2-negative breast cancer, BRCA1/2-positive breast cancer, BRCA1/2-negative cancer, or triple

negative breast cancer (TNBC). In embodiments, a cancer is triple negative breast cancer (TNBC). In some embodiments, a breast cancer is a metastatic breast cancer. In some embodiments, a breast cancer is an advanced breast cancer. In some embodiments, a cancer is a stage II, stage III or stage IV breast cancer. In some embodiments, a cancer is a stage IV breast cancer. In some embodiments, a breast cancer is a triple negative breast cancer.

[00200] In some embodiments, a patient or a population of patients to be treated by the methods of the present disclosure have or are susceptible to endometrial cancer (“EC”). Endometrial carcinoma is the most common cancer of the female genital, tract accounting for 10–20 per 100,000 person-years. The annual number of new cases of endometrial cancer (EC) is estimated at about 325 thousand worldwide. Further, EC is the most commonly occurring cancer in post-menopausal women. About 53% of endometrial cancer cases occur in developed countries. In 2015, approximately 55,000 cases of EC were diagnosed in the U.S. and no targeted therapies are currently approved for use in EC. There is a need for agents and regimens that improve survival for advanced and recurrent EC in 1L and 2L settings. Approximately 10,170 people are predicted to die from EC in the U.S. in 2016. The most common histologic form is endometrioid adenocarcinoma, representing about 75-80% of diagnosed cases. Other histologic forms include uterine papillary serous (less than 10%), clear cell 4%, mucinous 1%, squamous less than 1% and mixed about 10%.

[00201] From the pathogenetic point of view, EC falls into two different types, so-called types I and II. Type I tumors are low-grade and estrogen-related endometrioid carcinomas (EEC) while type II are non-endometrioid (NEEC) (mainly serous and clear cell) carcinomas. The World Health Organization has recently updated the pathologic classification of EC, recognizing nine different subtypes of EC, but EEC and serous carcinoma (SC) account for the vast majority of cases. EECs are estrogen-related carcinomas, which occur in perimenopausal patients, and are preceded by precursor lesions (endometrial hyperplasia/endometrioid intraepithelial neoplasia). Microscopically, lowgrade EEC (EEC 1–2) contains tubular glands, somewhat resembling the proliferative endometrium, with architectural complexity with fusion of the glands and cribriform pattern. High-grade EEC shows solid pattern of growth. In contrast, SC occurs in postmenopausal patients in absence of hyperestrogenism. At the microscope, SC shows thick, fibrotic or edematous papillae with prominent stratification of tumor cells, cellular budding, and anaplastic cells with large, eosinophilic cytoplasm. The vast majority of EEC are low grade

tumors (grades 1 and 2), and are associated with good prognosis when they are restricted to the uterus. Grade 3 EEC (EEC3) is an aggressive tumor, with increased frequency of lymph node metastasis. SCs are very aggressive, unrelated to estrogen stimulation, mainly occurring in older women. EEC 3 and SC are considered high-grade tumors. SC and EEC3 have been compared using the surveillance, epidemiology and End Results (SEER) program data from 1988 to 2001. They represented 10% and 15% of EC respectively, but accounted for 39% and 27% of cancer death respectively. Endometrial cancers can also be classified into four molecular subgroups: (1) ultramutated/POLE-mutant; (2) hypermutated MSI+ (e.g., MSI-H or MSI-L); (3) copy number low/microsatellite stable (MSS); and (4) copy number high/serous-like. Approximately 28% of cases are MSI-high. (Murali, *Lancet Oncol.* (2014). In some embodiments, a patient has a mismatch repair deficient subset of 2L endometrial cancer. In embodiments, an endometrial cancer is metastatic endometrial cancer. In embodiments, a patient has a MSS endometrial cancer. In embodiments, a patient has a MSI-H endometrial cancer.

[00202] In embodiments, a cancer is a lung cancer. In embodiments, a lung cancer is a squamous cell carcinoma of the lung. In embodiments, a lung cancer is small cell lung cancer (SCLC). In embodiments, a lung cancer is non-small cell lung cancer (NSCLC) such as squamous NSCLC. In embodiments, a lung cancer is an ALK-translocated lung cancer (e.g., ALK-translocated NSCLC). In embodiments, a cancer is NSCLC with an identified ALK translocation. In embodiments, a lung cancer is an EGFR-mutant lung cancer (e.g., EGFR-mutant NSCLC). In embodiments, a cancer is NSCLC with an identified EGFR mutation.

[00203] In embodiments, a cancer is a colorectal (CRC) cancer (e.g., a solid tumor). In embodiments, a colorectal cancer is an advanced colorectal cancer. In embodiments, a colorectal cancer is a metastatic colorectal cancer. In embodiments, a colorectal cancer is a MSI-H colorectal cancer. In embodiments, a colorectal cancer is a MSS colorectal cancer. In embodiments, a colorectal cancer is a POLE-mutant colorectal cancer. In embodiments, a colorectal cancer is a POLD-mutant colorectal cancer. In embodiments, a colorectal cancer is a high TMB colorectal cancer.

[00204] In embodiments, a cancer is a melanoma. In embodiments, a melanoma is an advanced melanoma. In embodiments, a melanoma is a metastatic melanoma. In embodiments, a melanoma is a MSI-H melanoma. In embodiments, a melanoma is a MSS melanoma. In

embodiments, a melanoma is a POLE-mutant melanoma. In embodiments, a melanoma is a POLD-mutant melanoma. In embodiments, a melanoma is a high TMB melanoma.

[00205] In embodiments, a cancer is an advanced cancer.

[00206] In embodiments, a cancer is a metastatic cancer.

[00207] In embodiments, a cancer is a recurrent cancer (e.g., a recurrent gynecological cancer such as recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, recurrent primary peritoneal cancer, or recurrent endometrial cancer).

[00208] Cancers that can be treated with methods described herein include cancers associated with a high tumor mutation burden (TMB), cancers that microsatellite stable (MSS), cancers that are characterized by microsatellite instability, cancers that have a high microsatellite instability status (MSI-H), cancers that have low microsatellite instability status (MSI-L), cancers associated with high TMB and MSI-H (e.g., cancers associated with high TMB and MSI-L or MSS), cancers having a defective DNA mismatch repair system, cancers having a defect in a DNA mismatch repair gene, hypermutated cancers, cancers having homologous recombination repair deficiency/homologous repair deficiency (“HRD”), cancers comprising a mutation in polymerase delta (POLD), and cancers comprising a mutation in polymerase epsilon (POLE).

[00209] In some embodiments, a tumor to be treated is characterized by microsatellite instability. In some embodiments, a tumor is characterized by microsatellite instability high status (MSI-H). Microsatellite instability (“MSI”) is or comprises a change that in the DNA of certain cells (such as tumor cells) in which the number of repeats of microsatellites (short, repeated sequences of DNA) is different than the number of repeats that was contained in the DNA from which it was inherited. About 15% of sporadic colorectal cancers (CRC) harbor widespread alterations in the length of microsatellite (MS) sequences, known as microsatellite instability (MSI) (Boland and Goel, 2010). Sporadic MSI CRC tumors display unique clinicopathological features including near-diploid karyotype, higher frequency in older populations and in females, and a better prognosis (de la Chapelle and Hampel, 2010; Popat et al., 2005). MSI is also present in other tumors, such as in endometrial cancer (EC) of the uterus, the most common gynecological malignancy (Duggan et al., 1994). The same reference Bethesda panel originally developed to screen an inherited genetic disorder (Lynch syndrome) (Umar et al., 2004) is currently applied to test MSI for CRCs and ECs. However, the genes frequently

targeted by MSI in CRC genomes rarely harbor DNA slippage events in EC genomes (Gurin et al., 1999).

[00210] Microsatellite instability arises from a failure to repair replication-associated errors due to a defective DNA mismatch repair (MMR) system. This failure allows persistence of mismatch mutations all over the genome, but especially in regions of repetitive DNA known as microsatellites, leading to increased mutational load. It has been demonstrated that at least some tumors characterized by MSI-H have improved responses to certain anti-PD-1 agents (Le et al., (2015) *N. Engl. J. Med.* 372(26):2509-2520; Westdorp et al., (2016) *Cancer Immunol. Immunother.* 65(10):1249-1259). In some embodiments, a cancer has a microsatellite instability of high microsatellite instability (e.g., MSI-H status). In some embodiments, a cancer has a microsatellite instability status of low microsatellite instability (e.g., MSI-Low). In some embodiments, a cancer has a microsatellite instability status of microsatellite stable (e.g., MSS status). In some embodiments microsatellite instability status is assessed by a next generation sequencing (NGS)-based assay, an immunohistochemistry (IHC)-based assay, and/or a PCR-based assay. In some embodiments, microsatellite instability is detected by NGS. In some embodiments, microsatellite instability is detected by IHC. In some embodiments, microsatellite instability is detected by PCR.

[00211] In embodiments, a patient has a MSI-L cancer.

[00212] In embodiments, a patient has a MSI-H cancer. In some embodiments, a patient has a MSI-H solid tumor. In embodiments, a MSI-H cancer is MSI-H endometrial cancer. In embodiments, a MSI—H cancer is a solid tumor. In embodiments, a MSI-H cancer is a metastatic tumor. In embodiments, a MSI-H cancer is endometrial cancer. In embodiments, a MSI-H cancer is a non-endometrial cancer. In embodiments, a MSI-H cancer is colorectal cancer.

[00213] In embodiments, a patient has a MSS cancer. In embodiments, a MSS cancer is MSS endometrial cancer.

[00214] In embodiments, a cancer is associated with a POLE (DNA polymerase epsilon) mutation (i.e., a cancer is a POLE-mutant cancer). In embodiments, a POLE mutation is a mutation in the exonuclease domain. In embodiments, a POLE mutation is a germline mutation. In embodiments, a POLE mutation is a sporadic mutation. In embodiments, a MSI cancer also is associated with a POLE mutation. In embodiments, a MSS cancer also is associated with a

POLE mutation. In embodiments, a POLE mutation is identified using sequencing. In embodiments, a POLE-mutant cancer is endometrial cancer. In embodiments, a POLE-mutant cancer is colon cancer. In embodiments, a POLE-mutant cancer is pancreatic cancer, ovarian cancer, or cancer of the small intestine.

[00215] In embodiments, a cancer is associated with a POLD (DNA polymerase delta) mutation (i.e., a cancer is a POLD-mutant cancer). In embodiments, a POLD mutation is a mutation in the exonuclease domain. In embodiments, a POLD mutation is a somatic mutation. In embodiments, a POLD mutation is a germline mutation. In embodiments, a POLD-mutant cancer is identified using sequencing. In embodiments, a POLD-mutant cancer is endometrial cancer. In embodiments, a POLD-mutant cancer is colorectal cancer. In embodiments, a POLD-mutant cancer is brain cancer.

[00216] In some embodiments, a patient has a mismatch repair deficient cancer.

[00217] In embodiments, a MMRd cancer is colorectal cancer.

[00218] Microsatellite instability may arise from a failure to repair replication-associated errors due to a defective DNA mismatch repair (MMR) system. This failure allows persistence of mismatch mutations all over the genome, but especially in regions of repetitive DNA known as microsatellites, leading to increased mutational load that may improve responses to certain anti-PD-1 agents. *Id.* In some embodiments, MSI-H status is assessed by a NGS-based assay and/or a PCR-based MSI assay. In some embodiments, microsatellite instability is detected by next generation sequencing. In embodiments, microsatellite instability is detected using immunohistochemistry (IHC) testing.

[00219] In embodiments, a cancer (e.g., a MMRd cancer) is characterized by a high tumor mutation burden (i.e., a cancer is a high TMB cancer). In some embodiments, the cancer is associated with high TMB and MSI-H. In some embodiments, the cancer is associated with high TMB and MSI-L or MSS. In some embodiments, the cancer is endometrial cancer associated with high TMB. In some related embodiments, the endometrial cancer is associated with high TMB and MSI-H. In some related embodiments, the endometrial cancer is associated with high TMB and MSI-L or MSS. In embodiments, a high TMB cancer is colorectal cancer. In embodiments, a high TMB cancer is lung cancer (e.g., small cell lung cancer (SCLC) or non-small cell lung cancer (NSCLC) such as squamous NSCLC or non-squamous NSCLC). In

embodiments, a high TMB cancer is melanoma. In embodiments, a high TMB cancer is urothelial cancer.

[00220] In embodiments, a patient has a cancer with elevated expression of tumor-infiltrating lymphocytes (TILs), i.e., a patient has a high-TIL cancer. In embodiments, a high-TIL cancer is breast cancer (e.g., triple negative breast cancer (TNBC) or HER2-positive breast cancer). In embodiments, a high-TIL cancer is a metastatic cancer (e.g., a metastatic breast cancer).

[00221] In embodiments, immune-related gene expression signatures can be predictive of a response to an anti-PD-1 therapy for cancer as described herein. For example, a gene panel that includes genes associated with IFN- γ signaling can be useful in identifying cancer patients who would benefit from anti-PD-1 therapy. Exemplary gene panels are described in Ayers et al., *J. Clin. Invest.*, 127(8):2930-2940, 2017. In embodiments, a cancer patient has a cancer that is breast cancer (e.g., TNBC) or ovarian cancer. In embodiments, a cancer patient has a cancer that is bladder cancer, gastric cancer, biliary cancer, esophageal cancer, or head and neck squamous cell carcinoma (HNSCC). In embodiments, a cancer patient has a cancer that is anal cancer or colorectal cancer.

[00222] In some embodiments, a patient has a tumor that expresses PD-L1. In some embodiments, PD-L1 status is evaluated in a patient or patient population. In some embodiments, mutational load and baseline gene expression profiles in archival or fresh pre-treatment biopsies are evaluated before, during and/or after treatment with an anti-PD-1 antibody agent. In some embodiments, the status and/or expression of TIM-3 and/or LAG-3 are evaluated in patients.

[00223] In some embodiments, a patient has previously been treated with one or more different cancer treatment modalities. In some embodiments, at least some of the patients in the cancer patient population have previously been treated with one or more of surgery, radiotherapy, chemotherapy or immunotherapy. In some embodiments, at least some of the patients in the cancer patient population have previously been treated with chemotherapy (e.g., platinum-based chemotherapy). For example, a patient who has received two lines of cancer treatment can be identified as a 2L cancer patient (e.g., a 2L NSCLC patient). In embodiments, a patient has received two lines or more lines of cancer treatment (e.g., a 2L+ cancer patient such as a 2L+ endometrial cancer patient). In embodiments, a patient has not been previously treated with an anti-PD-1 therapy. In embodiments, a patient previously received at least one line of cancer treatment (e.g., a patient previously received at least one line or at least two lines of

cancer treatment). In embodiments, a patient previously received at least one line of treatment for metastatic cancer (e.g., a patient previously received one or two lines of treatment for metastatic cancer). In embodiments, a subject is resistant to treatment with an agent that inhibits PD-1. In embodiments, a subject is refractory to treatment with an agent that inhibits PD-1. In embodiments, a method described herein sensitizes the subject to treatment with an agent that inhibits PD-1.

T Cell Immunoglobulin and Mucin Domain-3 (TIM-3)

[00224] The protein T Cell Immunoglobulin and Mucin Domain-3 (TIM-3), also known as Hepatitis A Virus Cellular Receptor 2 (HAVCR2), is a Th1-specific cell surface protein that regulates macrophage activation and enhances the severity of experimental autoimmune encephalomyelitis in mice. TIM-3 is highly expressed on the surface of multiple immune cell types, including, for example, Th1 IFN- γ + cells, Th17 cells, natural killer (NK) cells, monocytes, and tumor-associated dendritic cells (DCs) (see, e.g., Clayton et al., *J. Immunol.*, 192(2): 782-791 (2014); Jones et al., *J. Exp. Med.*, 205: 2763-2779 (2008); Monney et al., *Nature*, 415: 536-541 (2002); Hastings et al., *Eur. J. Immunol.*, 39: 2492-2501 (2009); Seki et al., *Clin. Immunol.*, 127: 78-88 (2008); Ju et al., *B. J. Hepatol.*, 52: 322-329 (2010); Anderson et al., *Science*, 318: 1141-1143 (2007); Baitsch et al., *PLoS ONE*, 7: e30852 (2012); Ndhlovu et al., *Blood*, 119: 3734-3743 (2012). TIM-3 also is highly expressed on “exhausted” or impaired CD8+ T-cells in a variety of chronic viral infections (e.g., HIV, HCV, and HBV) and in certain cancers (see, e.g., McMahan et al., *J. Clin. Invest.*, 120(12): 4546-4557 (2010); Jin et al., *Proc Natl Acad Sci USA*, 107(33): 14733-14738 (2010); Golden-Mason et al., *J. Virol.*, 83(18): 9122-9130 (2009); Jones et al., *supra*; Fourcade et al., *J. Exp. Med.*, 207(10): 2175-2186 (2010); Sakuishi et al., *J. Exp. Med.*, 207(10): 2187-2194 (2010); Zhou et al., *Blood*, 117(17): 4501-4510 (2011); Ngiow et al., *Cancer Res.*, 71(10): 3540-3551 (2011)).

[00225] Putative ligands for TIM-3 include phosphatidylserine (Nakayama et al., *Blood*, 113: 3821-3830 (2009)), galectin-9 (Zhu et al., *Nat. Immunol.*, 6: 1245-1252 (2005)), high-mobility group protein 1 (HMGB1) (Chiba et al., *Nature Immunology*, 13: 832-842 (2012)), and carcinoembryonic antigen cell adhesion molecule 1 (CEACAM1) (Huang et al., *Nature*, 517(7534): 386-90 (2015)).

[00226] TIM-3 functions to regulate various aspects of the immune response. The interaction of TIM-3 and galectin-9 (Gal-9) induces cell death and *in vivo* blockade of this interaction exacerbates autoimmunity and abrogates tolerance in experimental models, strongly suggesting that TIM-3 is a negative regulatory molecule. In contrast to its effect on T-cells, the TIM-3-Gal-9 interaction exhibits antimicrobial effects by promoting macrophage clearance of intracellular pathogens (see, e.g., Sakuishi et al., *Trends in Immunology*, 32(8): 345-349 (2011)). *In vivo*, suppression of TIM-3 has been shown to enhance the pathological severity of experimental autoimmune encephalomyelitis (Monney et al., *supra*; and Anderson, A. C. and Anderson, D. E., *Curr. Opin. Immunol.*, 18: 665-669 (2006)). Studies also suggest that dysregulation of the TIM-3-galectin-9 pathway could play a role in chronic autoimmune diseases, such as multiple sclerosis (Anderson and Anderson, *supra*). TIM-3 promotes clearance of apoptotic cells by binding phosphatidyl serine through its unique binding cleft (see, e.g., DeKruyff et al., *J. Immunol.*, 184(4):1918-1930 (2010)).

[00227] Inhibition of TIM-3 activity, such as through use of monoclonal antibodies, is currently under investigation as an immunotherapy for tumors based on preclinical studies (see, e.g., Ngiew et al., *Cancer Res.*, 71(21): 1-5 (2011); Guo et al., *Journal of Translational Medicine*, 11: 215 (2013); and Ngiew et al., *Cancer Res.*, 71(21): 6567-6571 (2011)).

[00228] The present disclosure provides particular antibody agents and methods relating thereto for the treatment of cancer.

Anti-TIM-3 antibody agents

[00229] The present disclosure provides methods of treating cancer that include administering compositions that deliver particular anti-TIM-3 antibody agents according to regimens that may achieve clinical benefit(s). The present disclosure describes, at least in part, anti-TIM-3 antibody agents and various compositions and methods relating thereto.

[00230] In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain variable domain whose amino acid sequence comprises SEQ ID NO: 1 or SEQ ID NO: 7.

SEQ ID NO: 1

EVQLLES GGGLVQPGGSLRLSCAAASGFTFSSYDMSWVRQAPGKGLDWVSTISG
GGTYTYYQDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCASMDYWGQG
TTVTVSSA

SEQ ID NO: 7

EVQLLES GGG LVQP GGS LRLSCAAASGFTFSSYDMSWVRQAPGKGLDWVSTISG
GGTYTTYQDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCASMDYWGQG
TTVTVSS

[00231] In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin light chain variable domain whose amino acid sequence comprises SEQ ID NO: 2 or SEQ ID NO: 8.

SEQ ID NO: 2

DIQMTQSPSSLSASVGDRVTITCRASQSIRRYLNWYHQKPGKAPKLLIYGASTLQ
SGVPSRFS GSGSGTDFTLTIS SLQPEDFAVYYCQQSHSAPLTFGGGTKVEIKR

SEQ ID NO: 8

DIQMTQSPSSLSASVGDRVTITCRASQSIRRYLNWYHQKPGKAPKLLIYGASTLQ
SGVPSRFS GSGSGTDFTLTIS SLQPEDFAVYYCQQSHSAPLTFGGGTKVEIK

[00232] In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain variable domain whose amino acid sequence comprises SEQ ID NO: 1 or SEQ ID NO: 7 and/or an immunoglobulin light chain variable domain whose amino acid sequence comprises SEQ ID NO: 2 or SEQ ID NO: 8.

[00233] In some embodiments, an anti-TIM-3 antibody agent comprises a variable heavy chain complementarity determining region 1, 2, and/or 3 (VH-CDR) comprising the amino acid sequences GFTFSSYDMS (SEQ ID NO: 21), TISGGGTYTTYQDSVK (SEQ ID NO: 22), and/or MDY (SEQ ID NO: 23), respectively. In some embodiments an anti-TIM-3 antibody agent comprises a variable light chain complementary determining region 1, 2, and/or 3 (VL-CDR) comprising the amino acid sequences RASQSIRRYLN (SEQ ID NO: 24), GASTLQS (SEQ ID NO: 25), and/or QQSHSAPLT (SEQ ID NO: 26), respectively. In some embodiments, an anti-TIM-3 antibody comprises VH-CDR sequences of SEQ ID NOs: 21, 22, and 23, and VL-CDR sequences of SEQ ID NOs: 24, 25, and 26. See **Table 1**.

Table 1. Amino Acid Sequences of Heavy and Light Chain Complementarity Determining Regions (CDRs)

Antibody	Variable region	CDR1	CDR2	CDR3
Anti-TIM-3	VH	GFTFSSYDMS (SEQ ID NO: 21)	TISGGGTYTTYQDSVK (SEQ ID NO: 22)	MDY (SEQ ID NO: 23)
Anti-TIM-3	VL	RASQSIRRYLN (SEQ ID NO: 24)	GASTLQS (SEQ ID NO: 25)	QQSHSAPLT (SEQ ID NO: 26)

[00234] In some embodiments, an anti-TIM-3 antibody agent is a monoclonal antibody. Particular antibodies of the present invention bind to TIM-3 with high affinity and effectively neutralize TIM-3 activity. Particular antibody heavy chain polypeptide (SEQ ID NO:3) and light chain polypeptide (SEQ ID NO:4) sequences are explicitly provided.

An anti-TIM-3 antibody heavy chain polypeptide (**SEQ ID NO:3**)

EVQLLESGGGLVQPGGSLRLSCAAASGFTFSSYDMSWVRQAPGKGLDW
VSTISGGGTYTTYQDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYC
ASMDYWGQGTTVTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFP
EPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTKTYTCN
VDHKPSNTKVKDKRVESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISR
TPEVTCVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVV
SVLTVLHQDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPP
SQEEMTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPVLDSDGS
FFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYTQKSLSLSLGK

An anti-TIM-3 antibody light chain polypeptide (**SEQ ID NO:4**)

DIQMTQSPSSLSASVGDRVTITCRASQSIRRYLNWYHQKPGKAPKLLIYGA
STLQSGVPSRFSGSGSGTDFTLTISLQPEDFAVYYCQQSHSAPLTFGGGT
KVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDN
ALQSGNSQESVTEQDSKDSSTLSLTLSKADYEKHKVYACEVTHQGLS
SPVTKSFNRGEC

[00235] The present disclosure provides an isolated immunoglobulin heavy chain polypeptide having an amino acid sequence as set forth in SEQ ID NO:3. The present disclosure further provides an isolated immunoglobulin heavy chain polypeptide having an amino acid sequence that shares at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with that set forth in SEQ ID NO:3. In some embodiments, sequence differences relative to the sequence set forth in SEQ ID NO:3 are not within the CDRs. In some

embodiments, an isolated immunoglobulin heavy chain polypeptide includes all three CDRs of SEQ ID NO:3. In some embodiments, an immunoglobulin heavy chain polypeptide includes a signal peptide. In some embodiments, an immunoglobulin heavy chain polypeptide which includes a signal peptide has an amino acid sequence as set forth in SEQ ID NO:5.

An anti-TIM-3 antibody heavy chain polypeptide with a signal sequence (**SEQ ID NO:5**)

MEFGLSWLFLVAILKGVQCEVQLLESGGGLVQPGGSLRLSCAAASGFTFSSYDM
SWVRQAPGKGLDWVSTISGGGTYTYQDSVKGRFTISRDNSKNTLYLQMNSLR
AEDTAVYYCASMDYWGQGTITVSSASTKGPSVFPLAPCSRSTSESTAALGCLV
KDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGKTKTYTCN
VDHKPSNTKVDKRVESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVT
CVVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQD
WLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTC
LVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSDGSFFLYSRLTVDKSRWQEGN
VFSCSVMHREALHNHYTQKSLSLGLGK

[00236] In some embodiments, a provided immunoglobulin heavy chain polypeptide is or comprises an IgG4 polypeptide. In some embodiments, a provided immunoglobulin heavy chain polypeptide comprises a human IGHG4*01 polypeptide. In some embodiments, a provided immunoglobulin heavy chain polypeptide comprises one or more mutations within the IgG heavy chain region. In some embodiments, a provided immunoglobulin heavy chain polypeptide comprises an IgG4 heavy chain constant region having one or more mutations in the heavy chain constant region. In some embodiments, a provided immunoglobulin heavy chain polypeptide comprises an IgG4 heavy chain constant region having one or more mutations in hinge region. It is envisioned that in some embodiments, a mutation in the IgG4 hinge region may prevent half molecule exchange with other IgG4 molecules. In some embodiments, the one or more mutations in hinge region of IgG4 may include a serine to proline stabilizing mutation that prevents half molecule exchange with other IgG4 molecules. In some embodiments, the one or more mutations in hinge region of IgG4 may include an S228P mutation. See, e.g., J. Biol. Chem. 2015; 290(9):5462-5469.

[00237] The present disclosure provides an isolated immunoglobulin light chain polypeptide having an amino acid sequence as set forth in SEQ ID NO:4. The present disclosure further provides an isolated immunoglobulin light chain polypeptide having an amino acid sequence that shares at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with that set forth in SEQ ID NO:4. In some embodiments, sequence differences relative to the sequence set forth in SEQ ID NO:4 are not within the CDRs. In some

embodiments, an isolated immunoglobulin light chain polypeptide includes all three CDRs of SEQ ID NO:4. In some embodiments, a provided immunoglobulin light chain polypeptide is a kappa light chain. In some embodiments, a provided immunoglobulin light chain polypeptide comprises a human IGKC*01 polypeptide. In some embodiments, the immunoglobulin light chain polypeptide includes a signal peptide. In some embodiments, the immunoglobulin light chain polypeptide which includes a signal peptide has an amino acid sequence as set forth in SEQ ID NO:6.

An anti-TIM-3 antibody light chain polypeptide with a signal sequence (**SEQ ID NO:6**)
MDMRVPAQLLGLLLLWLRGARCDIQMTQSPSSLSASVGDRVITTCRASQSIRRYL
NWKYHQKPGKAPKLLIYGASTLQSGVPSRFSGSGSGTDFTLTISSLQPEDFAVYYC
QQSHSAPLTFGGGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREA
KVQWKVDNALQSGNSQESVTEQDSKDSSTYSLSSTLTLSKADYEEKHKVYACEVT
HQGLSSPVTKSFNRGEC

[00238] In some embodiments, the present disclosure provides an anti-TIM-3 antibody agent comprising at least one immunoglobulin heavy chain having an amino acid sequence as set forth in SEQ ID NO:3 and at least one immunoglobulin light chain having an amino acid sequence as set forth in SEQ ID NO:4. In some embodiments an anti-TIM-3 antibody agent comprises two immunoglobulin heavy chains, each having an amino acid sequence as set forth in SEQ ID NO:3. Alternatively or additionally, in some embodiments an anti-TIM-3 antibody agent comprises two immunoglobulin light chains, each having an amino acid sequence as set forth in SEQ ID NO:4. In some embodiments, an anti-TIM-3 antibody agent has a canonical antibody format.

[00239] In some embodiments, a provided heavy chain, light chain and/or antibody agent is glycosylated and one or more sites. In some embodiments, a glycan is N-linked to an Fc region. In some embodiments, an antibody agent is glycosylated at Asn297 (Kabat numbering). In some embodiments, present disclosure provides a composition comprising one or more glycoforms of an heavy chain, light chain, and/or antibody agent as described herein. In some embodiments, a provided composition comprises plurality of such glycoforms, present in specified absolute and/or relative amounts. In some embodiments, the present disclosure provides compositions that may be substantially free of one or more particular glycoforms of an heavy chain, light chain, and/or antibody agent as described herein.

[00240] In embodiments, a TIM-3 binding agent is an anti-TIM-3 antibody that is TSR-022, which comprises a humanized monoclonal anti-TIM-3 antibody comprising a heavy chain whose amino acid sequence comprises SEQ ID NO: 3 and a light chain whose amino acid sequence comprises SEQ ID NO:4. This anti-TIM-3 antibody utilizes a human IGHG4*01 heavy chain gene, and a human IGKC*01 kappa light chain gene, as scaffolds. Further, there is a single Ser to Pro point mutation in the hinge region of the IgG4 heavy chain at the canonical S228 position, corresponding to residue 240 in SEQ ID NO: 5, which includes the signal sequence. Without wishing to be bound by theory, it is envisioned that this point mutation serves to stabilize the hinge of the antibody heavy chain.

[00241] Additional biophysical and biochemical characterization of this exemplary humanized monoclonal anti-TIM-3 antibody is also provided regarding observed disulfide linkages and glycosylation. Lys-C and trypsin digested peptides were well separated and detected by on-line LC-MS analysis. The disulfide bond linkages were confirmed by comparison of total ion chromatograms in the non-reduced (NR) condition with the reduced condition. Disulfide linkages are consistent with the expected disulfide linkage pattern for an IgG4 molecule. The residues involved in the expected inter- and intrachain disulfide linkages are tabulated below (**Tables 2, 3, and 4**).

Table 2 – Expected residues involved in disulfide linkages of an exemplary anti-TIM-3 antibody agent heavy chain having an amino acid sequence as set forth in SEQ ID NO: 1.

Cysteine residue ID	anti-TIM-3 mAb HC Residue (position in SEQ ID NO: 1)
I	22
II	96
III	127
IV	140
V	196
VI	219
VII	222
VIII	254
IX	314
X	360
XI	418

Table 3 – Expected residues involved in disulfide linkages of an exemplary anti-TIM-3 antibody agent light chain having an amino acid sequence as set forth in SEQ ID NO: 2.

Cysteine residue ID	anti-TIM-3 mAb LC Residue (position in SEQ ID NO: 2)
I	23
II	88
III	134
IV	194
V	214

Table 4. Exemplary disulfide bond assignments for an anti-TIM-3 antibody

Disulfide bond NO.	Disulfide-containing peptides	Linkage site on HC (position in SEQ ID NO:1)	Linkage site on LC (position in SEQ ID NO:2)
DS1	VTITCR=FSGSGSGTDFLTISLQPEDF AVYYCQQSHSAPLTFGGGTK		23 88
DS2	SGTASVVCLLNNFYPR=VYACEVTHQGLS SPVTK		134 194
DS3	SFNRGEC=GPSVFPLAPCSR GEC=GPSVFPLAPCSR	127	214
DS4	LSCAAASGFTFSSYDMSWVR=AEDTA VYYCASMDYWGQGTTVTVSSASTK	22 97	
DS5	STSESTAALGCLVK=TYTCNVDHK STSESTAALGCLVK=TYTCNVDHKPSNTK	140 196	
DS6	YGPPCPPCPAPEFLGGPSVFLFPPK=YGPPC PPCPAPEFLGGPSVFLFPPK YGPPCPPCPAPEFLGGPSVFLFPPK=YGPPC PPCPAPEFLGGPSVFLFPPKPK	219 222	
DS7	TPEVTCVVVDVSQEDPEVQFNWYVDGVE VHNAK=CK	254 314	

Disulfide bond NO.	Disulfide-containing peptides	Linkage site on HC (position in SEQ ID NO:1)	Linkage site on LC (position in SEQ ID NO:2)
DS8	NQVSLTCLVK=WQEGNVFSCSVMHEALH NHYTQK	360 418	

LC: light chain; HC: heavy chain

[00242] This exemplary anti-TIM-3 antibody exhibits an occupied N-glycosylation site at asparagine residue 290 in the CH2 domain of each heavy chain in the mature protein sequence (SEQ ID NO:1). The expressed N-glycosylation at this site is a mixture of oligosaccharide species typically observed on IgGs expressed in mammalian cell culture, for example, shown below is the relative abundance of glycan species from a preparation of this exemplary anti-TIM-3 antibody cultured in Chinese Hamster Ovary (CHO) cells (**Table 5**).

Table 5 – Glycan Analysis of an anti- TIM-3 antibody binding agent

Species	Abundance (% of total oligosaccharide)	Description of Glycan
G0F	20.1%	Core fucosylated agalactobiantennary complex-type oligosaccharide
G1F	41.9%	Core fucosylated monogalactosylated biantennary complex type oligosaccharide
G2F	29.0%	Core-fucosylated galactosylated biantennary complex type oligosaccharide
G2FS1	3.2%	Monosialylated core fucosylated galactosylated biantennary complex type oligosaccharide
G2FS2	1.2%	Disialylated core fucosylated galactosylated biantennary complex type oligosaccharide
M5	0.4%	Oligomannosidic N-linked oligosaccharide, Man ₅ GlcNAc ₂

PD-1-binding agents

[00243] The present disclosure provides methods of treating cancer that further include administering compositions that deliver particular programmed death-1 protein (PD-1)-binding agents according to regimens that may achieve clinical benefit(s). The present disclosure describes, at least in part, PD-1-binding agents (e.g., anti-PD-1 antibody agents) and various compositions and methods relating thereto. In some embodiments, a PD-1-binding agent is a monoclonal antibody.

[00244] In some embodiments, a PD-1-binding agent comprises an immunoglobulin heavy chain variable domain whose amino acid sequence comprises SEQ ID NO: 11 or SEQ ID NO: 17.

SEQ ID NO: 11

EVQLLES GGGLVQP GGSLRLSCAASGFTFSSYDMSWVRQAPGKGLEWVSTISGG
GSYTTYQDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCASPYAMDYW
GQGTTVTVSSA

SEQ ID NO: 17

EVQLLES GGGLVQP GGSLRLSCAASGFTFSSYDMSWVRQAPGKGLEWVSTISGG
GSYTTYQDSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCASPYAMDYW
GQGTTVTVSS

[00245] In some embodiments, a PD-1-binding agent comprises an immunoglobulin light chain variable domain whose amino acid sequence comprises SEQ ID NO: 12 or SEQ ID NO: 18.

SEQ ID NO: 12

DIQLTQSPSFLSAYVGDRVTITCKASQDVGTAVAWYQQKPGKAPKLLIYWASTL
HTGVPSRFS GSGSGTEFTLTIS SLQPEDFATYYCQHYSSYPWTFGQGTKLEIKR

SEQ ID NO: 18

DIQLTQSPSFLSAYVGDRVTITCKASQDVGTAVAWYQQKPGKAPKLLIYWASTL
HTGVPSRFS GSGSGTEFTLTIS SLQPEDFATYYCQHYSSYPWTFGQGTKLEIK

[00246] In some embodiments, a PD-1-binding agent comprises an immunoglobulin heavy chain variable domain whose amino acid sequence comprises SEQ ID NO: 11 or SEQ ID NO: 17 and/or an immunoglobulin light chain variable domain whose amino acid sequence comprises SEQ ID NO: 12 or SEQ ID NO: 18. In some embodiments a PD-1-binding agent is or comprises

an immunoglobulin G4 (IgG4) humanized monoclonal antibody (mAb). In some embodiments, a PD-1-binding agent comprises a human IGHG4*01 polypeptide. In some embodiments, a PD-1-binding agent comprises one or more mutations within the IgG heavy chain region. In some embodiments, a PD-1-binding agent comprises an IgG4 heavy chain constant region having one or more mutations in the heavy chain constant region. In some embodiments, a PD-1-binding agent comprises an IgG4 heavy chain constant region having one or more mutations in hinge region. It is envisioned that in some embodiments, a mutation in the IgG4 hinge region may prevent half molecule exchange with other IgG4 molecules. In some embodiments, the one or more mutations in hinge region of IgG4 may include a serine to proline stabilizing mutation that prevents half molecule exchange with other IgG4 molecules. In some embodiments, the one or more mutations in hinge region of IgG4 may include an S228P mutation. See, e.g., J. Biol. Chem. 2015; 290(9):5462-5469.

[00247] In some embodiments, a PD-1-binding agent comprises an immunoglobulin heavy chain polypeptide whose amino acid sequence comprises SEQ ID NO: 13.

SEQ ID NO: 13 – An anti-PD-1 antibody heavy chain polypeptide (CDR sequences)

EVQLLESGGGLVQPGGSLRLSCAASGFTFSSYDMSWVRQAPGKGLEWVSTISGG
GSYTTYQDSVKGRFTISRDNSKNTLYLQMNSLRAEDTAVYYCASPYYAMDYW
 GQGTTVTVSSASTKGPSVFPLAPCSRSTSESTAALGCLVKDYFPEPVTVSWNSGA
 LTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTKTYTCNVDHKPSNTKVDKRVES
 KYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSQEDPEVQFN
 WYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKGL
 PSSIEKTISKAKGQPREPQVYTLPPSQEEMTKNQVSLTCLVKGFYPSDIAVEWESN
 GQPENNYKTTPPVLDSDGSFFLYSRLTVDKSRWQEGNVFSCSVMHEALHNHYT
 QKSLSLSLGK

[00248] In some embodiments, a PD-1-binding agent comprises an immunoglobulin light chain polypeptide whose amino acid sequence comprises SEQ ID NO: 14.

SEQ ID NO: 14 – An anti-PD-1 antibody light chain polypeptide (CDR sequences)

DIQLTQSPSFLSAYVGDRVTITCKASQDVGTAVAWYQQKPGKAPKLLIYWASTL
HTGVPSRFGSGSGTEFTLTISSLQPEDFATYYCQHYSSYPWTFGQGTKLEIKRTV
 AAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQSGNSQESVT
 EQDSKDSTYSLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC

[00249] The present disclosure provides an isolated immunoglobulin heavy chain polypeptide having an amino acid sequence as set forth in SEQ ID NO:13. The present disclosure further provides an isolated immunoglobulin heavy chain polypeptide having an amino acid sequence

that shares at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with that set forth in SEQ ID NO:13. In some embodiments, sequence differences relative to the sequence set forth in SEQ ID NO:13 are not within the CDRs. In some embodiments, an isolated immunoglobulin heavy chain polypeptide includes all three CDRs of SEQ ID NO:13. In some embodiments, an immunoglobulin heavy chain polypeptide includes a signal peptide. In some embodiments, an immunoglobulin heavy chain polypeptide which includes a signal peptide has an amino acid sequence as set forth in SEQ ID NO:15.

An anti-PD-1 antibody heavy chain polypeptide with a signal sequence (**SEQ ID NO:15**)

MEFGLSWLFLVAILKGVQCEVQLLESGGGLVQPGGSLRLSCAASGFTFSSYDMS
WVRQAPGKGLEWVSTISGGGSYTTYQDSVKGRFTISRDN SKNTLYLQMN SLRA
EDTAVYYCASPYAMDYWGQGT TTVSSASTKGPSVFPLAPCSRSTSESTAALG
CLVKDYFPEPVT VSWNSGALTSGVHTFPAVLQSSGLYSLSSV VTPSSSLG TKTY
TCNV DHKPSNTKVDKR VESKYGPPCPPCPAPEFLGGPSVFLFPPKPKDTLMIS RTP
EVT CVVDVSQEDPEVQFNWYVDGVEVHNAKTKPREEQFNSTYRVVSVLTVLH
QDWLNGKEYKCKVSNKGLPSSIEKTISKAKGQPREPQVYTLPPS QEEMTKNQVS
LTCLVKGFYPSDIAVEWESNGQPENNYK TTPPVLDSDGSFFLYSRLTVDKSRWQ
EGNVFSCSVMHEALHNHYTQKSLSL SLGK

[00250] The present disclosure provides an isolated immunoglobulin light chain polypeptide having an amino acid sequence as set forth in SEQ ID NO:14. The present disclosure further provides an isolated immunoglobulin light chain polypeptide having an amino acid sequence that shares at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with that set forth in SEQ ID NO:14. In some embodiments, sequence differences relative to the sequence set forth in SEQ ID NO:14 are not within the CDRs. In some embodiments, an isolated immunoglobulin light chain polypeptide includes all three CDRs of SEQ ID NO:14. In some embodiments, a provided immunoglobulin light chain polypeptide is a kappa light chain. In some embodiments, a provided immunoglobulin light chain polypeptide comprises a human IGKC*01 polypeptide. In some embodiments, the immunoglobulin light chain polypeptide includes a signal peptide. In some embodiments, the immunoglobulin light chain polypeptide which includes a signal peptide has an amino acid sequence as set forth in SEQ ID NO:16.

An anti-PD-1 antibody light chain polypeptide with a signal sequence (**SEQ ID NO:16**)

MDMRVPAQLLGLLLWLPGARCDIQLTQSPSFLSAYVGDRVTITCKASQDVGTA
VAWYQQKPGKAPKLLIYWASTLHTGVPSRFSGSGSGTEFTLTIS SLQPEDFATYY
CQHYSSYPWTFGQGTKLEIKRTVAAPS VFIFPPSDEQLKSGTASVVCLLNNFYPRE
AKVQWKVDNALQSGNSQESVTEQDSKDSTYLSSTLTLSKADYEKHKVYACEV
THQGLSSPVTKSFNRGEC

[00251] SEQ ID NOs: 13 and 14 describe an exemplary humanized monoclonal anti-PD-1 antibody (TSR-042) utilizing a human IGHG4*01 heavy chain gene, and a human IGKC*01 kappa light chain gene, as scaffolds. There is a single Ser to Pro point mutation in the hinge region of the IgG4 heavy chain. This mutation is at the canonical S228 position. Without wishing to be bound by theory, it is envisioned that this point mutation serves to stabilize the hinge of the antibody heavy chain.

[00252] Additional biophysical and biochemical characterization of this exemplary humanized monoclonal anti-PD-1 antibody is provided herein, which is consistent with the expected disulfide linkage pattern for an IgG4 molecule. The residues involved in the expected inter- and intrachain disulfide linkages are tabulated below (**Tables 6 and 7**).

Table 6– Expected residues involved in disulfide linkages of an exemplary anti-PD-1 antibody agent heavy chain having an amino acid sequence as set forth in SEQ ID NO: 13.

Cysteine residue ID after Edelman ^a	anti-PD-1 mAb HC Residue (position in SEQ ID NO: 13)
I	22
II	96
III	130
IV	143
V	199
VI	222
VII	225
VIII	257
IX	317
X	363
XI	421

Table 7 – Expected residues involved in disulfide linkages of an exemplary anti-PD-1 antibody agent light chain having an amino acid sequence as set forth in SEQ ID NO: 14.

Cysteine residue ID after Edelman ^a	anti-PD-1 mAb LC Residue (position in SEQ ID NO: 14)
I	23
II	88
III	134

Cysteine residue ID after Edelman ^a	anti-PD-1 mAb LC Residue (position in SEQ ID NO: 14)
IV	194
V	214

[00253] This exemplary anti-PD-1 antibody exhibits an occupied N-glycosylation site at asparagine residue 293 in the CH2 domain of each heavy chain in the mature protein sequence (SEQ ID NO:13). The expressed N-glycosylation at this site is a mixture of oligosaccharide species typically observed on IgGs expressed in mammalian cell culture, for example, shown below is the relative abundance of glycan species from a preparation of this exemplary anti-PD-1 antibody cultured in Chinese Hamster Ovary (CHO) cells (**Table 8**).

Table 8 – Glycan Analysis of an anti-PD-1 antibody binding agent

Species	Abundance (% of total oligosaccharide)	Description of Glycan
G0	<0.1%	Nonfucosylated agalactobiantennary complex-type oligosaccharide
G0F	19.5%	Core fucosylated agalactobiantennary complex type oligosaccharide
G1	0.1%	Nonfucosylated monogalactosylated biantennary complex type oligosaccharide
G1F	45.6%	Core fucosylated monogalactosylated biantennary complex type oligosaccharide
G2F	27.4%	Core fucosylated galactosylated biantennary complex type oligosaccharide
M5	0.5%	Oligomannosidic N-glycan, Man ₅ GlcNAc ₂

[00254] In some embodiments, the present disclosure provides an anti-PD-1 antibody agent comprising at least one immunoglobulin heavy chain having an amino acid sequence as set forth in SEQ ID NO: 13 and at least one immunoglobulin light chain having an amino acid sequence as set forth in SEQ ID NO: 14. In some embodiments an anti-PD-1 antibody agent comprises two immunoglobulin heavy chains, each having an amino acid sequence as set forth in SEQ ID NO: 13. Alternatively or additionally, in some embodiments an anti-PD-1 antibody agent comprises two immunoglobulin light chains, each having an amino acid sequence as set forth in SEQ ID NO: 14. In some embodiments, an anti-PD-1 antibody agent has a canonical antibody format.

[00255] In some embodiments, a provided heavy chain, light chain and/or antibody agent has a structure that includes one or more disulfide bonds. In some embodiments, the one or more disulfide bonds are or include a disulfide bond at the expected position for an IgG4 immunoglobulin.

[00256] In some embodiments, a PD-1-binding agent is glycosylated and one or more sites. As used herein, “glycan” is a sugar polymer (moiety) component of a glycoprotein. The term “glycan” encompasses free glycans, including glycans that have been cleaved or otherwise released from a glycoprotein. In some embodiments, present disclosure provides a composition comprising one or more glycoforms of a heavy chain, light chain, and/or antibody agent as described herein. In some embodiments, a glycan is N-linked to an Fc region. In some embodiments, a PD-1-binding agent is glycosylated at Asn297 (Kabat numbering).

[00257] The term “glycoform” is used herein to refer to a particular form of a glycoprotein. That is, when a glycoprotein includes a particular polypeptide that has the potential to be linked to different glycans or sets of glycans, then each different version of the glycoprotein (i.e., where the polypeptide is linked to a particular glycan or set of glycans) is referred to as a “glycoform.” In some embodiments, a provided composition comprises a plurality of glycoforms of one or more of an heavy chain, light chain, and/or antibody agent as described herein.

[00258] In some embodiments, antagonist activity of a PD-1-binding agent in blocking the PD-1/PD-L1 or PD-L2 interaction may be confirmed or determined using a flow cytometry-based assay that measured binding of labeled PD-L1 and PD-L2 expressed as a mouse IgG1 Fc fusion proteins (PD-L1 mFc or PD-L2 mFc) to PD-1-expressing cells. In some embodiments, a PD-1-binding agent can efficiently block PD-1/PD-L1 and PD-1/PD-L2 binding compared to an IgG4 isotype control.

[00259] In some embodiments, a PD-1-binding agent can effectively neutralize PD-1 activity (e.g., can inhibit binding of PD-1 to PD-L1 and PD-L2). In some embodiments, functional antagonist activity of a PD-1-binding agent may be confirmed or determined in a mixed lymphocyte reaction (MLR) demonstrating enhanced interleukin (IL)-2 production upon addition of a PD-1-binding agent. In some embodiments, a MLR assay may be carried out using primary human CD4+ T cells as responders and human dendritic cells as stimulators.

Expression and Formulation

[00260] In some embodiments, an anti-TIM-3 antibody agent and/or a PD-1-binding agent is expressed from a vector comprising one or more nucleic acid sequences.

[00261] In some embodiments, anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain polypeptide that is encoded by a nucleotide sequence which comprises SEQ ID NO: 9.

Nucleotide sequence encoding anti-TIM-3 antibody heavy chain polypeptide (**SEQ ID NO: 9**)

```
GAG GTG CAG CTG TTG GAG TCT GGG GGA GGC TTG GTA CAG CCT GGG
GGG TCC CTG AGA CTC TCC TGT GCA GCA GCC TCT GGA TTC ACT TTC
AGT AGC TAT GAC ATG TCT TGG GTC CGC CAG GCT CCA GGG AAG GGG
CTG GAC TGG GTC TCA ACC ATT AGT GGT GGT GGT ACT TAC ACC TAC
TAT CAA GAC AGT GTG AAG GGG CGG TTC ACC ATC TCC AGA GAC AAT
TCC AAG AAC ACG CTG TAT CTG CAA ATG AAC AGC CTG AGA GCC GAG
GAC ACG GCC GTA TAT TAC TGT GCG TCC ATG GAC TAC TGG GGG CAA
GGG ACC ACG GTC ACC GTC TCC TCA GCA TCC ACC AAG GGC CCA TCG
GTC TTC CCG CTA GCA CCC TGC TCC AGG AGC ACC TCC GAG AGC ACA
GCC GCC CTG GGC TGC CTG GTC AAG GAC TAC TTC CCC GAA CCA GTG
ACG GTG TCG TGG AAC TCA GGC GCC CTG ACC AGC GGC GTG CAC ACC
TTC CCG GCT GTC CTA CAG TCC TCA GGA CTC TAC TCC CTC AGC AGC
GTG GTG ACC GTG CCC TCC AGC AGC TTG GGC ACG AAG ACC TAC ACC
TGC AAC GTA GAT CAC AAG CCC AGC AAC ACC AAG GTG GAC AAG AGA
GTT GAG TCC AAA TAT GGT CCC CCA TGC CCA CCA TGC CCA GCA CCT
GAG TTC CTG GGG GGA CCA TCA GTC TTC CTG TTC CCC CCA AAA CCC
AAG GAC ACT CTC ATG ATC TCC CGG ACC CCT GAG GTC ACG TGC GTG
GTG GTG GAC GTG AGC CAG GAA GAC CCC GAG GTC CAG TTC AAC TGG
TAC GTG GAT GGC GTG GAG GTG CAT AAT GCC AAG ACA AAG CCG CGG
GAG GAG CAG TTC AAC AGC ACG TAC CGT GTG GTC AGC GTC CTC ACC
GTC CTG CAC CAG GAC TGG CTG AAC GGC AAG GAG TAC AAG TGC AAG
GTC TCC AAC AAA GGC CTC CCG TCC TCC ATC GAG AAA ACC ATC TCC
AAA GCC AAA GGG CAG CCC CGA GAG CCA CAG GTG TAC ACC CTG CCC
CCA TCC CAG GAG GAG ATG ACC AAG AAC CAG GTC AGC CTG ACC TGC
CTG GTC AAA GGC TTC TAC CCC AGC GAC ATC GCC GTG GAG TGG GAG
AGC AAT GGG CAG CCG GAG AAC AAC TAC AAG ACC ACG CCT CCC GTG
CTG GAC TCC GAC GGC TCC TTC TTC CTC TAC AGC AGG CTA ACC GTG
GAC AAG AGC AGG TGG CAG GAG GGG AAT GTC TTC TCA TGC TCC GTG
ATG CAT GAG GCT CTG CAC AAC CAC TAC ACA CAG AAG AGC CTC TCC
CTG TCT CTG GGT AAA
```

[00262] In some embodiments, anti-TIM-3 antibody agent comprises an immunoglobulin light chain polypeptide that is encoded by a nucleotide sequence which comprises SEQ ID NO: 10.

Nucleotide sequence encoding anti-TIM-3 antibody light chain polypeptide (**SEQ ID NO: 10**)

GAC ATC CAG ATG ACC CAG TCT CCA TCC TCC CTG TCT GCA TCT GTA
GGA GAC AGA GTC ACC ATC ACT TGC CGG GCA AGT CAG AGC ATT AGG
AGG TAT TTA AAT TGG TAT CAC CAG AAA CCA GGG AAA GCC CCT AAG
CTC CTG ATC TAT GGT GCA TCC ACC TTG CAA AGT GGG GTC CCA TCA
AGG TTC AGT GGT AGT GGA TCT GGG ACA GAT TTC ACT CTC ACC ATC
AGC AGT CTG CAA CCT GAA GAT TTT GCA GTG TAT TAC TGT CAA CAG
AGT CAC AGT GCC CCC CTC ACT TTC GGC GGA GGG ACC AAG GTG GAG
ATC AAA CGA ACT GTG GCT GCA CCA TCT GTC TTC ATC TTC CCG CCA TCT
GAT GAG CAA TTG AAA TCT GGA ACT GCC TCT GTT GTG TGC CTG CTG
AAT AAC TTC TAT CCC AGA GAG GCC AAA GTA CAG TGG AAG GTG GAT
AAC GCC CTC CAA TCG GGT AAC TCC CAG GAG AGT GTC ACA GAG CAG
GAC AGC AAG GAC AGC ACC TAC AGC CTC AGC AGC ACC CTG ACG CTG
AGC AAA GCA GAC TAC GAG AAA CAC AAA GTC TAC GCC TGC GAA GTC
ACC CAT CAG GGC CTC AGC TCG CCC GTC ACA AAG AGC TTC AAC AGG
GGA GAG TGT

[00263] In some embodiments, PD-1 binding agent comprises an immunoglobulin heavy chain polypeptide that is encoded by a nucleotide sequence which comprises SEQ ID NO: 19.

SEQ ID NO: 19 – Nucleotide sequence encoding a immunoglobulin heavy chain polypeptide of a PD-1 binding agent

GAG GTG CAG CTG TTG GAG TCT GGG GGA GGC TTG GTA CAG CCT GGG
GGG TCC CTG AGA CTC TCC TGT GCA GCC TCT GGA TTC ACT TTC AGT
AGC TAT GAC ATG TCT TGG GTC CGC CAG GCT CCA GGG AAG GGG CTG
GAG TGG GTC TCA ACC ATT AGT GGT GGT GGT AGT TAC ACC TAC TAT
CAA GAC AGT GTG AAG GGG CGG TTC ACC ATC TCC AGA GAC AAT TCC
AAG AAC ACG CTG TAT CTG CAA ATG AAC AGC CTG AGA GCC GAG GAC
ACG GCC GTA TAT TAC TGT GCG TCC CCT TAC TAT GCT ATG GAC TAC
TGG GGG CAA GGG ACC ACG GTC ACC GTC TCC TCA GCA TCC ACC AAG
GGC CCA TCG GTC TTC CCG CTA GCA CCC TGC TCC AGG AGC ACC TCC
GAG AGC ACA GCC GCC CTG GGC TGC CTG GTC AAG GAC TAC TTC CCC
GAA CCA GTG ACG GTG TCG TGG AAC TCA GGC GCC CTG ACC AGC GGC
GTG CAC ACC TTC CCG GCT GTC CTA CAG TCC TCA GGA CTC TAC TCC CTC
AGC AGC GTG GTG ACC GTG CCC TCC AGC AGC TTG GGC ACG AAG ACC
TAC ACC TGC AAC GTA GAT CAC AAG CCC AGC AAC ACC AAG GTG GAC
AAG AGA GTT GAG TCC AAA TAT GGT CCC CCA TGC CCA CCA TGC CCA
GCA CCT GAG TTC CTG GGG GGA CCA TCA GTC TTC CTG TTC CCC CCA
AAA CCC AAG GAC ACT CTC ATG ATC TCC CGG ACC CCT GAG GTC ACG
TGC GTG GTG GTG GAC GTG AGC CAG GAA GAC CCC GAG GTC CAG TTC
AAC TGG TAC GTG GAT GGC GTG GAG GTG CAT AAT GCC AAG ACA AAG

CCG CGG GAG GAG CAG TTC AAC AGC ACG TAC CGT GTG GTC AGC GTC
 CTC ACC GTC CTG CAC CAG GAC TGG CTG AAC GGC AAG GAG TAC AAG
 TGC AAG GTC TCC AAC AAA GGC CTC CCG TCC TCC ATC GAG AAA ACC
 ATC TCC AAA GCC AAA GGG CAG CCC CGA GAG CCA CAG GTG TAC ACC
 CTG CCC CCA TCC CAG GAG GAG ATG ACC AAG AAC CAG GTC AGC CTG
 ACC TGC CTG GTC AAA GGC TTC TAC CCC AGC GAC ATC GCC GTG GAG
 TGG GAG AGC AAT GGG CAG CCG GAG AAC AAC TAC AAG ACC ACG CCT
 CCC GTG CTG GAC TCC GAC GGC TCC TTC TTC CTC TAC AGC AGG CTA
 ACC GTG GAC AAG AGC AGG TGG CAG GAG GGG AAT GTC TTC TCA TGC
 TCC GTG ATG CAT GAG GCT CTG CAC AAC CAC TAC ACA CAG AAG AGC
 CTC TCC CTG TCT CTG GGT AAA

[00264] In some embodiments, a PD-1-binding agent comprises an immunoglobulin light chain polypeptide that is encoded by a nucleotide sequence which comprises SEQ ID NO: 20.

SEQ ID NO: 20 – Nucleotide sequence encoding a immunoglobulin light chain polypeptide of a PD-1 binding agent

GAC ATC CAG TTG ACC CAG TCT CCA TCC TTC CTG TCT GCA TAT GTA
 GGA GAC AGA GTC ACC ATC ACT TGC AAG GCC AGT CAG GAT GTG GGT
 ACT GCT GTA GCC TGG TAT CAG CAA AAA CCA GGG AAA GCC CCT AAG
 CTC CTG ATC TAT TGG GCA TCC ACC CTG CAC ACT GGG GTC CCA TCA
 AGG TTC AGC GGC AGT GGA TCT GGG ACA GAA TTC ACT CTC ACA ATC
 AGC AGC CTG CAG CCT GAA GAT TTT GCA ACT TAT TAC TGT CAG CAT
 TAT AGC AGC TAT CCG TGG ACG TTT GGC CAG GGG ACC AAG CTG GAG
 ATC AAA CGG ACT GTG GCT GCA CCA TCT GTC TTC ATC TTC CCG CCA TCT
 GAT GAG CAA TTG AAA TCT GGA ACT GCC TCT GTT GTG TGC CTG CTG
 AAT AAC TTC TAT CCC AGA GAG GCC AAA GTA CAG TGG AAG GTG GAT
 AAC GCC CTC CAA TCG GGT AAC TCC CAG GAG AGT GTC ACA GAG CAG
 GAC AGC AAG GAC AGC ACC TAC AGC CTC AGC AGC ACC CTG ACG CTG
 AGC AAA GCA GAC TAC GAG AAA CAC AAA GTC TAC GCC TGC GAA GTC
 ACC CAT CAG GGC CTC AGC TCG CCC GTC ACA AAG AGC TTC AAC AGG
 GGA GAG TGT

[00265] In some embodiments, an anti-TIM-3 antibody agent and/or a PD-1-binding agent is expressed from a vector comprising one or more nucleic acid sequences encoding an immunoglobulin heavy chain variable domain polypeptide and/or an immunoglobulin light chain variable domain polypeptide. In some embodiments, an anti-TIM-3 antibody agent and/or a PD-1-binding agent is expressed from a vector comprising one or more nucleic acid sequences encoding an immunoglobulin heavy chain polypeptide and/or an immunoglobulin light chain polypeptide. The vector can be, for example, a plasmid, episome, cosmid, viral vector (e.g., retroviral or adenoviral), or phage. Suitable vectors and methods of vector preparation are well

known in the art (see, e.g., Sambrook et al., *Molecular Cloning, a Laboratory Manual*, 3rd edition, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (2001), and Ausubel et al, *Current Protocols in Molecular Biology*, Greene Publishing Associates and John Wiley & Sons, New York, N.Y. (1994)).

[00266] In some embodiments, vector(s) for expression of an anti-TIM-3 antibody agent and/or a PD-1-binding agent further comprises expression control sequences, such as promoters, enhancers, polyadenylation signals, transcription terminators, internal ribosome entry sites (IRES), and the like, that provide for the expression of the coding sequence in a host cell. Exemplary expression control sequences are known in the art and described in, for example, Goeddel, *Gene Expression Technology: Methods in Enzymology*, Vol. 185, Academic Press, San Diego, Calif. (1990).

[00267] The vector(s) comprising the nucleic acid(s) encoding an anti-TIM-3 antibody agent and/or a PD-1-binding agent of the present disclosure can be introduced into a host cell that is capable of expressing the polypeptides encoded thereby, including any suitable prokaryotic or eukaryotic cell. Some preferable qualities of host cells include easy and reliable growth, a reasonably fast growth rate, having well-characterized expression systems, and/or ease/efficient transformation or transfection.

[00268] In some embodiments, mammalian cells are utilized. A number of suitable mammalian host cells are known in the art, and many are available from the American Type Culture Collection (ATCC, Manassas, VA). Examples of suitable mammalian cells include, but are not limited to, Chinese hamster ovary cells (CHO) (ATCC No. CCL61), CHO DHFR-cells (Urlaub et al, *Proc. Natl. Acad. Sci. USA*, 97: 4216-4220 (1980)), human embryonic kidney (HEK) 293 or 293T cells (ATCC No. CRL1573), and 3T3 cells (ATCC No. CCL92). Other suitable mammalian cell lines are the monkey COS-1 (ATCC No. CRL1650) and COS-7 cell lines (ATCC No. CRL1651), as well as the CV-1 cell line (ATCC No. CCL70).

[00269] Further exemplary mammalian host cells include primate cell lines and rodent cell lines, including transformed cell lines. Normal diploid cells, cell strains derived from in vitro culture of primary tissue, as well as primary explants, are also suitable. Other suitable mammalian cell lines include, but are not limited to, mouse neuroblastoma N2A cells, HeLa, mouse L-929 cells, and BHK or HaK hamster cell lines, all of which are available from the

ATCC. Methods for selecting suitable mammalian host cells and methods for transformation, culture, amplification, screening, and purification of cells are known in the art.

[00270] In some embodiments, the mammalian cell is a human cell. For example, the mammalian cell can be a human lymphoid or lymphoid derived cell line, such as a cell line of pre-B lymphocyte origin. Examples of human lymphoid cells lines include, without limitation, RAMOS (CRL-1596), Daudi (CCL-213), EB-3 (CCL-85), DT40 (CRL-2111), 18-81 (Jack et al, Proc. Natl. Acad. Sci. USA, 85: 1581-1585 (1988)), Raji cells (CCL-86), and derivatives thereof.

[00271] In some embodiments, an anti-TIM-3 antibody agent is formulated as a pharmaceutical composition, containing one or a combination of monoclonal antibodies, or antigen-binding portion(s) thereof, formulated with a pharmaceutically acceptable carrier. An anti-TIM-3 antibody agent may be formulated alone or in combination with other drugs (e.g., as an adjuvant). For example, an anti-TIM-3 antibody agent can be administered in combination with other agents for the treatment or prevention of the diseases disclosed herein (e.g., cancer).

[00272] Therapeutic compositions typically must be sterile and stable under the conditions of manufacture and storage. The composition can be formulated as a solution, microemulsion, liposome, or other ordered structure suitable to high drug concentration. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as mannitol, sorbitol, or sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent that delays absorption, for example, monostearate salts and gelatin.

[00273] Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by sterilization microfiltration. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle that contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders is the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying (lyophilization) that yield a powder of the

active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

[00274] In some embodiments, a therapeutic composition is formulated as a sterile liquid. In some embodiments, the composition is free from visible particles. In some embodiments, the composition is formulated in a buffer. In some embodiments, an anti-TIM-3 antibody agent is stored at 2-8 °C. In some embodiments, a drug product of the present disclosure is free of preservatives.

General Protocol

[00275] As described herein, provided methods comprise administering an anti-TIM-3 antibody agent to a patient, a subject, or a population of subjects (e.g., according to a regimen that achieves clinical benefit).

[00276] Provided methods can provide various benefits (e.g., a clinical benefit). In embodiments, a method described herein achieves a clinical benefit. In embodiments, a clinical benefit is stable disease (SD). In embodiments, a clinical benefit is a partial response (PR). In embodiments, a clinical benefit is a complete response (CR).

[00277] In embodiments, a combination therapy achieves a clinical benefit for each therapy administered to a patient. For example, a subject may be resistant to treatment with an agent that inhibits PD-1 or a subject may be refractory to treatment with an agent that inhibits PD-1. In embodiments, a method described herein sensitizes the subject to treatment with an agent that inhibits PD-1. Therefore, in embodiments, a benefit of a combination therapy comprising administration of a TIM-3 inhibitor (e.g., any anti-TIM-3 antibody described herein) and a PD-1 inhibitor (e.g., any anti-PD-1 antibody described herein) is to achieve a clinical benefit with a PD-1 inhibitor (e.g., any anti-PD-1 antibody described herein) or to improve a clinical benefit.

[00278] In embodiments, a patient or subject is an animal. In embodiments, a patient or subject is a human.

[00279] In embodiments, administration of an anti-TIM-3 antibody agent is parenteral administration. In embodiments, parenteral administration is intravenous administration. In embodiments, intravenous administration is intravenous infusion.

[00280] In some embodiments, the regimen comprises at least one parenteral dose of an anti-TIM-3 antibody agent. In some embodiments, the regimen comprises a plurality of parenteral doses.

[00281] In some embodiments, the parenteral dose is an amount of an anti-TIM-3 antibody agent is within a range of about 5 to about 5000 mg (e.g., about 5 mg, about 10 mg, about 50 mg, about 100 mg, about 500 mg, about 600 mg, about 700 mg, about 800 mg, about 900 mg, about 1000 mg, about 1100 mg, about 1200 mg, about 1300 mg, about 1400 mg, about 1500 mg, about 2000 mg, about 3000 mg, about 4000 mg, about 5000 mg, or a range defined by any two of the foregoing values). In some embodiments, the parenteral dose of an anti-TIM-3 antibody agent is 500 mg or 1000 mg. In some embodiments, the parenteral dose of an anti-TIM-3 antibody agent is about 100 mg, about 300 mg, or about 1200 mg.

[00282] In some embodiments, the dose is in an amount relative to body weight. In some embodiments, the parenteral dose of an anti-TIM-3 antibody agent is within a range of about 0.01 mg/kg to 100 mg/kg of animal or human body weight; however, doses below or above this exemplary range are within the scope of the invention. The dose (e.g., a daily parenteral dose) can be about 0.01 mg/kg to about 50 mg/kg of total body weight (e.g., about 0.1 mg/kg, about 0.5 mg/kg, about 1 mg/kg, about 2 mg/kg, about 3 mg/kg, about 4 mg/kg, about 5 mg/kg, about 6 mg/kg, about 7 mg/kg, about 8 mg/kg, about 9 mg/kg, about 10 mg/kg, about 12 mg/kg, about 15 mg/kg, about 20 mg/kg, or a range defined by any two of the foregoing values).

[00283] In some embodiments, a composition that delivers an anti-TIM-3 antibody agent is administered to a patient at a dose of 0.1, 1, 3 or 10 mg/kg. In some embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of 0.1, 1, 3 or 10 mg/kg every two weeks. In some embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of 0.1, 1, 3 or 10 mg/kg every three weeks. In some embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of 0.1, 1, 3 or 10 mg/kg every four weeks.

[00284] In some embodiments, a composition that delivers an anti-TIM-3 antibody agent is administered to a patient at a dose of about 100 to about 500 mg (e.g., 200 to 500 mg). In some embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of about 100 to about 500 mg (e.g., 200 mg, 300 mg, 400 mg, 500 mg) every two weeks. In some embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of about 100 to about 500 mg (e.g., 200 mg, 300 mg, 400 mg, 500 mg) every three weeks. In some

embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of about 100 to about 500 mg (e.g., 200 mg, 300 mg, 400 mg, 500 mg) mg every four weeks.

[00285] In some embodiments, a composition that delivers an anti-TIM-3 antibody agent is administered to a patient at a dose of about 800 to about 1500 mg (e.g., 800 mg, 900 mg, 1000 mg, 1100 mg, 1200 mg, 1300 mg, 1400 mg, 1500 mg). In some embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of about 800 to about 1500 mg (e.g., 800 mg, 900 mg, 1000 mg, 1100 mg, 1200 mg, 1300 mg, 1400 mg, 1500 mg) mg every four weeks. In some embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of about 800 to about 1500 mg (e.g., 800 mg, 900 mg, 1000 mg, 1100 mg, 1200 mg, 1300 mg, 1400 mg, 1500 mg) every six weeks. In some embodiments, the composition that delivers an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of about 800 to about 1500 mg (e.g., 800 mg, 900 mg, 1000 mg, 1100 mg, 1200 mg, 1300 mg, 1400 mg, 1500 mg) every eight weeks.

[00286] In embodiments, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 1 mg/kg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody

such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, an anti-TIM-3 antibody agent is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00287] In embodiments, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 3 mg/kg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00288] In embodiments, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 10 mg/kg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00289] In embodiments, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 100 - 1500 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every week (Q1W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such

as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00290] In embodiments, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 100 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in

combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00291] In any of the methods described herein, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 200 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00292] In any of the methods described herein, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 300 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00293] In any of the methods described herein, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 400 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g.,

intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00294] In any of the methods described herein, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 500 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-

PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00295] In any of the methods described herein, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 600 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00296] In any of the methods described herein, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 700 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00297] In any of the methods described herein, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 800 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g.,

intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00298] In any of the methods described herein, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 900 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-

PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00299] In any of the methods described herein, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 1000 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00300] In any of the methods described herein, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 1100 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00301] In any of the methods described herein, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 1200 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g.,

intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00302] In any of the methods described herein, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 1300 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-

PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00303] In any of the methods described herein, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 1400 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00304] In any of the methods described herein, a dose (e.g., a therapeutically effective dose, a dose administered by a composition that delivers an anti-TIM-3 antibody agent, or a parenteral dose) is about 1500 mg of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every two weeks (Q2W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every three weeks (Q3W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every four weeks (Q4W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every five weeks (Q5W). In embodiments, administration of a dose (e.g., intravenous administration such as intravenous infusion) is once every six weeks (Q6W). In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is intravenous infusion. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for monotherapy. In embodiments, administration of a TIM-3 inhibitor (e.g., an anti-TIM-3 antibody agent) is for combination therapy (e.g., in combination with an immune checkpoint inhibitor (e.g., an anti-PD-1 antibody such as TSR-042). In embodiments, a TIM-3 inhibitor is an anti-TIM-3 antibody agent. In embodiments, a TIM-3 inhibitor is TSR-022. In embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:3 and an immunoglobulin light chain comprising an amino acid sequence having at least about 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% overall identity with SEQ ID NO:4.

[00305] Therapeutic or prophylactic efficacy can be monitored by periodic assessment of treated patients. For repeated administrations over several days or longer, depending on the condition, the treatment can be repeated until a desired suppression of disease symptoms occurs. However, other dosage regimens may be useful and are within the scope of the invention. The desired dosage can be delivered by a single bolus administration of the composition, by multiple bolus administrations of the composition, or by continuous infusion administration of the composition.

[00306] In some embodiments, an anti-TIM-3 antibody agent is administered to a patient or population of subjects who has exhibited response to prior therapy. In some embodiments, the patient or population of subjects has exhibited response to a prior cancer therapy.

[00307] In some embodiments, an anti-TIM-3 antibody agent is administered to a patient or population of subjects who has not exhibited response to prior therapy. In some embodiments, the patient or population of subjects has not received or exhibited response to a prior cancer therapy.

[00308] In embodiments, a subject is resistant to treatment with an agent that inhibits PD-1. In embodiments, a subject is refractory to treatment with an agent that inhibits PD-1. In embodiments, a method described herein sensitizes the subject to treatment with an agent that inhibits PD-1.

[00309] In embodiments, an anti-TIM-3 antibody agent as described herein is administered in combination with one or more additional therapies (e.g., therapies as described herein). That is, a subject is treated with an anti-TIM-3 antibody agent and one or more additional therapies is administered to a subject such that the subject receives each therapy.

[00310] In embodiments, an additional therapy is surgery. In embodiments, an additional therapy is radiotherapy. In embodiments, an additional therapy is chemotherapy. In embodiments, an additional therapy is immunotherapy.

[00311] In some embodiments, an anti-TIM-3 antibody agent is administered as a monotherapy.

[00312] In some embodiments, an anti-TIM-3 antibody agent is administered in combination therapy. In embodiments, an anti-TIM-3 antibody agent is administered in combination with another treatment modality (e.g., with one or more of surgery, radiotherapy, chemotherapy, or immunotherapy). In embodiments, an anti-TIM-3 antibody agent is administered in combination with surgery. In embodiments, an anti-TIM-3 antibody agent is administered in combination with radiotherapy. In embodiments, an anti-TIM-3 antibody agent is administered in combination with chemotherapy. In embodiments, an anti-TIM-3 antibody agent is administered in combination with immunotherapy.

[00313] In some embodiments, an anti-TIM-3 antibody agent is administered simultaneously or sequentially with an additional therapeutic agent, such as, for example, another antibody agent

(e.g., an antibody agent that binds to PD-1) and/or a chemotherapeutic agent (e.g., niraparib). In some embodiments, an anti-TIM-3 antibody agent is administered before, during, or after administration of an additional therapeutic agent. In some embodiments, an anti-TIM-3 antibody agent is administered before, during, or after administration of a chemotherapeutic agent (e.g., niraparib).

[00314] An anti-TIM-3 antibody agent may be administered alone or in combination with other drugs (e.g., as an adjuvant). For example, the anti-TIM-3 antibody agent can be administered in combination with other agents for the treatment or prevention of the diseases disclosed herein (e.g., cancer). In this respect, the anti-TIM-3 antibody agent can be used in combination with at least one other anticancer agent including, for example, any chemotherapeutic agent known in the art, ionization radiation, small molecule anticancer agents, cancer vaccines, biological therapies (e.g., other monoclonal antibodies, cancer-killing viruses, gene therapy, and adoptive T-cell transfer), and/or surgery.

[00315] Administration of an anti-TIM-3 antibody agent simultaneously or sequentially with an additional therapeutic agent is referred to herein as “combination therapy.” In combination therapy, an anti-TIM-3 antibody agent can be administered prior to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concurrently with, or subsequent to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of the additional therapeutic agent to a subject in need thereof. In some embodiments an anti-TIM-3 antibody agent and an additional therapeutic agent are administered 1 minute apart, 10 minutes apart, 30 minutes apart, less than 1 hour apart, 1 hour to 2 hours apart, 2 hours to 3 hours apart, 3 hours to 4 hours apart, 4 hours to 5 hours apart, 5 hours to 6 hours apart, 6 hours to 7 hours apart, 7 hours to 8 hours apart, 8 hours to 9 hours apart, 9 hours to 10 hours apart, 10 hours to 11 hours apart, 11 hours to 12 hours apart, no more than 24 hours apart, or no more than 48 hours apart.

PARP Inhibitors


[00316] In embodiments, an additional therapy is a poly (ADP-ribose) polymerase (PARP) inhibitor.

[00317] In embodiments, a PARP inhibitor inhibits PARP-1 and/or PARP-2. In some embodiments, the agent is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, or a toxin. In related embodiments, the agent is ABT-767, AZD 2461, BGB-290, BGP 15, CEP 8983, CEP 9722, DR 2313, E7016, E7449, fluzoparib (SHR 3162), IMP 4297, INO1001, JPI 289, JPI 547, monoclonal antibody B3-LysPE40 conjugate, MP 124, niraparib (ZEJULA) (MK-4827), NU 1025, NU 1064, NU 1076, NU1085, olaparib (AZD2281), ONO2231, PD 128763, R 503, R554, rucaparib (RUBRACA) (AG-014699, PF-01367338), SBP 101, SC 101914, simmiparib, talazoparib (BMN-673), veliparib (ABT-888), WW 46, 2-(4-(trifluoromethyl)phenyl)-7,8-dihydro-5H-thiopyrano[4,3-d]pyrimidin-4-ol, and salts or derivatives thereof. In some related embodiments, an agent is niraparib, olaparib, rucaparib, talazoparib, veliparib, or salts or derivatives thereof. In certain embodiments, an agent is niraparib or a salt or derivative thereof. In certain embodiments, an agent is olaparib or a salt or derivative thereof. In certain embodiments, an agent is rucaparib or a salt or derivative thereof. In certain embodiments, an agent is talazoparib or a salt or derivative thereof. In certain embodiments, an agent is veliparib or a salt or derivative thereof.

[00318] Niraparib, (3S)-3-[4-{7-(aminocarbonyl)-2H-indazol-2-yl}phenyl]piperidine, is an orally available, potent, poly (adenosine diphosphate [ADP]-ribose) polymerase (PARP)-1 and -2 inhibitor. See WO 2008/084261 (published on July 17, 2008), WO 2009/087381 (published July 16, 2009), and PCT/US17/40039 (filed June 29, 2017), the entirety of each of which is hereby incorporated by reference. Niraparib can be prepared according to Scheme 1 of WO 2008/084261.

[00319] In some embodiments, niraparib can be prepared as a pharmaceutically acceptable salt. One of skill in the art will appreciate that such salt forms can exist as solvated or hydrated polymorphic forms. In some embodiments, niraparib is prepared in the form of a hydrate.

[00320] In certain embodiments, niraparib is prepared in the form of a tosylate salt. In some embodiments, niraparib is prepared in the form of a tosylate monohydrate. The molecular structure of the tosylate monohydrate salt of niraparib is shown below:



[00321] Niraparib is a potent and selective PARP-1 and PARP-2 inhibitor with inhibitory concentration at 50% of control (IC_{50}) = 3.8 and 2.1 nM, respectively, and is at least 100-fold selective over other PARP-family members. Niraparib inhibits PARP activity, stimulated as a result of DNA damage caused by addition of hydrogen peroxide, in various cell lines with an IC_{50} and an inhibitory concentration at 90% of control (IC_{90}) of about 4 and 50 nM, respectively.

[00322] In embodiments, niraparib is administered at a dose equivalent to about 100 mg of niraparib free base (e.g., a pharmaceutically acceptable salt of niraparib such as niraparib tosylate monohydrate is administered at a dose equivalent to about 100 mg of niraparib free base). In embodiments, niraparib is administered at a dose equivalent to about 200 mg of niraparib free base (e.g., a pharmaceutically acceptable salt of niraparib such as niraparib tosylate monohydrate is administered at a dose equivalent to about 200 mg of niraparib free base). In embodiments, niraparib is administered at a dose equivalent to about 300 mg of niraparib free base (e.g., a pharmaceutically acceptable salt of niraparib such as niraparib tosylate monohydrate is administered at a dose equivalent to about 300 mg of niraparib free base).

Checkpoint Inhibitors

[00323] In embodiments, an additional therapy is an immunotherapy. In embodiments, an immunotherapy comprises administration of one or more further immune checkpoint inhibitors (e.g., administration of one, two, three, four, or more further immune checkpoint inhibitors).

[00324] Exemplary immune checkpoint targets for inhibition include: PD-1 (e.g., inhibition via anti-PD-1, anti-PD-L1, or anti-PD-L2 therapies), CTLA-4, TIM-3, TIGIT, LAGs (e.g., LAG-3), CEACAM (e.g., CEACAM-1, -3 and/or -5), VISTA, BTLA, LAIR1, CD160, 2B4, CD80,

CD86, B7-H3 (CD276), B7-H4 (VTCN1), HVEM (TNFRSF14 or CD270), KIR, A2aR, MHC class I, MHC class II, GALS, adenosine, TGFR (e.g., TGFR beta), B7-H1, B7-H4 (VTCN1), OX-40, CD137, CD40, IDO, and CSF1R. Accordingly, agents that inhibit any of these molecules can be used in combination with an anti-TIM-3 therapy described herein.

[00325] In embodiments, an immune checkpoint inhibitor is an agent that inhibits PD-1, CTLA-4, LAG-3, TIGIT, IDO, or CSF1R.

[00326] In embodiments, an immune checkpoint inhibitor is a PD-1 inhibitor. In embodiments, a PD-1 inhibitor is a PD-1 binding agent (e.g., an antibody, an antibody conjugate, or an antigen-binding fragment thereof). In embodiments, a PD-1 inhibitor is a PD-L1 or PD-L2 binding agent (e.g., an antibody, an antibody conjugate, or an antigen-binding fragment thereof). In embodiments, a PD-1 inhibitor is nivolumab, pembrolizumab, atezolizumab, durvalumab, avelumab, TSR-042, PDR-001, tislelizumab (BGB-A317), cemiplimab (REGN2810), LY-3300054, JNJ-63723283, MGA012, BI-754091, IBI-308, camrelizumab (HR-301210), BCD-100, JS-001, CX-072, BGB-A333, AMP-514 (MEDI-0680), AGEN-2034, CS1001, Sym-021, SHR-1316, PF-06801591, LZM009, KN-035, AB122, genolimzumab (CBT-501), FAZ-053, CK-301, AK 104, or GLS-010, or any of the PD-1 antibodies disclosed in WO2014/179664. In embodiments, a PD-1 inhibitor is TSR-042. In some embodiments, a PD-1-inhibitor (e.g., TSR-042) is administered according to a regimen that comprises administering an about 500 mg dose every 3 weeks for four doses followed by administering at least one about 1,000 mg dose every six weeks after the fourth dose of about 500 mg. In some embodiments, additional about 1,000 mg doses are administered every six weeks after the first about 1000 mg dose until no further clinical benefit is achieved. In some particular embodiments, a PD-1-inhibitor (e.g., TSR-042) is administered according to a dosing regimen that includes 500 mg for 4 cycles Q3W followed by 1000 mg Q6W.

[00327] In embodiments, an immune checkpoint inhibitor is a CTLA-4 inhibitor (e.g., an antibody, an antibody conjugate, or an antigen-binding fragment thereof). In embodiments, a CTLA-4 inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, or a toxin. In embodiments, a CTLA-4 inhibitor is a small molecule. In embodiments, a CTLA-4 inhibitor is a CTLA-4 binding agent. In embodiments, a CTLA-4 inhibitor is an antibody, an antibody conjugate, or an antigen-binding fragment thereof. In embodiments, a CTLA-4 inhibitor is ipilimumab (Yervoy), AGEN1884, or tremelimumab.

[00328] In embodiments, an immune checkpoint inhibitor is a LAG-3 inhibitor (e.g., an antibody, an antibody conjugate, or an antigen-binding fragment thereof). In embodiments, a LAG-3 inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, or a toxin. In embodiments, a LAG-3 inhibitor is a small molecule. In embodiments, a LAG-3 inhibitor is a LAG-3 binding agent. In embodiments, a LAG-3 inhibitor is an antibody, an antibody conjugate, or an antigen-binding fragment thereof. In embodiments, a LAG-3 inhibitor is a IMP321, BMS-986016, GSK2831781, Novartis LAG525, or a LAG-3 inhibitor described in WO 2016/126858, WO 2017/019894, or WO 2015/138920, each of which is hereby incorporated by reference in its entirety.

[00329] In embodiments, an immune checkpoint inhibitor is a TIGIT inhibitor (e.g., an antibody, an antibody conjugate, or an antigen-binding fragment thereof). In embodiments, a TIGIT inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, or a toxin. In embodiments, a TIGIT inhibitor is small molecule. In embodiments, a TIGIT inhibitor is a TIGIT binding agent. In embodiments, a TIGIT inhibitor is an antibody, an antibody conjugate, or an antigen-binding fragment thereof. In embodiments, a TIGIT inhibitor is MTIG7192A, BMS-986207, or OMP-31M32.

[00330] In embodiments, an immune checkpoint inhibitor is an IDO inhibitor. In embodiments, an IDO inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, or a toxin. In embodiments, an IDO inhibitor is small molecule. In embodiments, an IDO inhibitor is an IDO binding agent. In embodiments, an IDO inhibitor is an antibody, an antibody conjugate, or an antigen-binding fragment thereof.

[00331] In embodiments, an immune checkpoint inhibitor is a CSF1R inhibitor. In embodiments, a CSF1R inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, or a toxin. In embodiments, a CSF1R inhibitor is small molecule. In embodiments, a CSF1R inhibitor is a CSF1R binding agent. In embodiments, a CSF1R inhibitor is an antibody, an antibody conjugate, or an antigen-binding fragment thereof.

[00332] In embodiments, a method comprises administering a TIM-3 inhibitor with at least two of the immune checkpoint inhibitors. In embodiments, a method comprises administering a third checkpoint inhibitor. In embodiments, a method comprises administering a TIM-3 inhibitor with a PD-1 inhibitor, and a LAG-3 inhibitor, such that the subject receives all three. In

embodiments, a method comprises administering a TIM-3 inhibitor with a PD-1 inhibitor, a LAG-3 inhibitor, and a CTLA-4 inhibitor, such that the subject receives all four.

[00333] In embodiments, a subject has been further administered or will be administered an agent that inhibits poly (ADP-ribose) polymerase (PARP), such that the subject receives treatment with a TIM-3 inhibitor and a PARP inhibitor.

[00334] In embodiments, a subject is further administered or will be administered one or more immune checkpoint inhibitors (e.g., a PD-1 inhibitor and/or a LAG-3 inhibitor) such that the subject receives treatment with a TIM-3 inhibitor, a PARP inhibitor (e.g., niraparib), and the one or more immune checkpoint inhibitors. In embodiments, a subject is administered a TIM-3 inhibitor, a PD-1 inhibitor (e.g., TSR-042) and a PARP inhibitor (e.g., niraparib). In embodiments, a subject is administered a TIM-3 inhibitor, a PD-1 inhibitor (e.g., TSR-042), a LAG-3 inhibitor, and a PARP inhibitor (e.g., niraparib).

[00335] For female patients of childbearing potential, it is preferable that the patient have a negative serum pregnancy test within 72 hours prior to the date of administration of the first dose of an anti-TIM-3 antibody agent. It is also preferable that female patients of childbearing potential and male patients agree to use 2 adequate methods of contraception with their partner. In some embodiments, a patient agrees to use 2 methods of contraception starting with the screening visit through 150 days after the last dose of study therapy.

[00336] The present disclosure provides, in some embodiments, methods of treating cancer in a patient in need thereof, the method comprising administering a one or more compositions that deliver an anti-TIM-3 antibody agent in combination with a PD-1 inhibitor (e.g., a PD-1-binding agent such as an anti-PD-1 antibody). **Figure 1** provides an exemplary schematic of the combination of anti-TIM-3 and anti-PD-1 antibodies in order to enhance anti-tumor efficiency. In some embodiments, a patient or patient population is receiving a combination therapy that comprises administration of an anti-TIM-3 antibody agent and a PD-1-binding agent. In some embodiments, the PD-1 binding agent is nivolumab or pembrolizumab. In embodiments, a PD-1 inhibitor is a PD-1 binding agent (e.g., an antibody, an antibody conjugate, or an antigen-binding fragment thereof). In embodiments, a PD-1 binding agent is nivolumab, pembrolizumab, TSR-042, PDR-001, tislelizumab (BGB-A317), cemiplimab (REGN2810), LY-3300054, JNJ-63723283, MGA012, BI-754091, IBI-308, camrelizumab (HR-301210), BCD-100, JS-001, CX-072, AMP-514 / MEDI-0680, AGEN-2034, CS1001, TSR-042, Sym-021, PF-06801591,

LZM009, KN-035, AB122, genolimzumab (CBT-501), AK 104, or GLS-010, or derivatives thereof. In embodiments, a PD-1 inhibitor is a PD-L1 or PD-L2 binding agent (e.g., an antibody, an antibody conjugate, or an antigen-binding fragment thereof). In embodiments, a PD-1 inhibitor is a PD-L1 or PD-L2 binding agent is durvalumab, atezolizumab, avelumab, BGB-A333, SHR-1316, FAZ-053, CK-301, or, PD-L1 millamolecule, or derivatives thereof.

[00337] In some embodiments, a patient or patient population is receiving a combination therapy that comprises administration of an anti-TIM-3 antibody agent comprising an immunoglobulin heavy chain variable domain whose amino acid sequence comprises SEQ ID NO: 1 or SEQ ID NO: 7 and an immunoglobulin light chain variable domain whose amino acid sequence comprises SEQ ID NO: 2 or SEQ ID NO: 8. In some embodiments, an anti-TIM-3 antibody agent comprises an immunoglobulin heavy chain whose amino acid sequence comprises SEQ ID NO: 3 and an immunoglobulin light chain whose amino acid sequence comprises SEQ ID NO: 4.

[00338] In some embodiments, a patient or patient population is receiving a combination therapy that comprises administration of a PD-1-binding agent comprising an immunoglobulin heavy chain variable domain whose amino acid sequence comprises SEQ ID NO: 11 or SEQ ID NO: 17 and an immunoglobulin light chain variable domain whose amino acid sequence comprises SEQ ID NO: 12 or SEQ ID NO: 18. In some embodiments comprising combinations, the PD-1-binding agent comprises an immunoglobulin heavy chain whose amino acid sequence comprises SEQ ID NO: 13 and an immunoglobulin light chain whose amino acid sequence comprises SEQ ID NO: 14.

[00339] In some embodiments, an anti-TIM-3 antibody agent (e.g., an anti-TIM-3 antibody) is administered at a dose of 0.1, 1, 3 or 10 mg/kg. In some embodiments, an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of 0.1, 1, 3 or 10 mg/kg every two weeks. In some embodiments, an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of 1, 3 or 10 mg/kg every three weeks.

[00340] In some embodiments, an anti-TIM-3 antibody agent is administered according to a regimen that includes a dose of 1, 3 or 10 mg/kg every four weeks. In some embodiments, an anti-TIM-3 antibody agent at a fixed dose within a range of 200 mg to 1,500 mg. In some embodiments, an anti-TIM-3 antibody agent at a fixed dose within a range of 300 mg to 1,000 mg. In some embodiments, an anti-TIM-3 antibody agent is administered according to a regimen

that includes a fixed dose every two weeks. In some embodiments, an anti-TIM-3 antibody agent is administered according to a regimen that includes a fixed dose every three weeks. In some embodiments, an anti-TIM-3 antibody agent is administered according to a regimen that includes a fixed dose every four weeks.

[00341] In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody such as TSR-042) is administered at a dose of about 1, 3 or 10 mg/kg. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody such as TSR-042) is administered according to a regimen that includes a dose of about 1, 3 or 10 mg/kg every two weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody such as TSR-042) is administered according to a regimen that includes a dose of about 1, 3 or 10 mg/kg every three weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody) is administered according to a regimen that includes a dose of about 1, 3 or 10 mg/kg every four weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody such as TSR-042) at a dose of about 500 mg. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody such as TSR-042) is administered according to a regimen that includes a dose of about 500 mg every two weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody such as TSR-042) is administered according to a regimen that includes a dose of about 500 mg every three weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody such as TSR-042) is administered according to a regimen that includes a dose of about 500 mg every four weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody such as TSR-042) is administered according to a regimen that includes a dose of about 1000 mg every six weeks. In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody such as TSR-042) is administered according to a regimen that includes a first dose of about 500 mg every three weeks (Q3W) for the first 2-6 (e.g., the first 2, 3, 4, 5, or 6) dosage cycles and a second dose of about 1000 mg every six weeks (Q6W) until treatment is discontinued (e.g., due to disease progression, adverse effects, or as determined by a physician). In some embodiments, a PD-1-binding agent (e.g., an anti-PD-1 antibody such as TSR-042) is administered according to a regimen that includes a first dose of about 500 mg every three weeks (Q3W) for the first four dosage cycles and a second dose of about 1000 mg every six weeks (Q6W) until treatment is discontinued (e.g., due to disease progression, adverse effects, or as determined by a physician). In embodiments, a

PD-1 binding agent is an anti-PD-1 antibody. In embodiments, a PD-1 binding agent is TSR-042.

[00342] In certain methods, an anti-TIM-3 antibody agent can be administered prior to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks before), concurrently with, or subsequent to (e.g., 5 minutes, 15 minutes, 30 minutes, 45 minutes, 1 hour, 2 hours, 4 hours, 6 hours, 12 hours, 24 hours, 48 hours, 72 hours, 96 hours, 1 week, 2 weeks, 3 weeks, 4 weeks, 5 weeks, 6 weeks, 8 weeks, or 12 weeks after) the administration of a PD-1-binding agent to a subject in need thereof.

Measuring Tumor Response

[00343] In some embodiments, a clinical benefit is a complete response (“CR”), a partial response (“PR”) or a stable disease (“SD”). In some embodiments, a clinical benefit corresponds to at least SD. In some embodiments, a clinical benefit corresponds to at least a PR. In some embodiments, a clinical benefit corresponds to a CR. In some embodiments, at least 1%, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90% or 95% of patients achieve a clinical benefit. In some embodiments, at least 5% of patients achieve a clinical benefit. In some embodiments, at least 5% of patients achieve SD. In some embodiments, at least 5% of patients achieve at least a PR. In some embodiments, at least 5% of patients achieve CR. In some embodiments, at least 20% of patients achieve a clinical benefit. In some embodiments, at least 20% of patients achieve SD.

[00344] In some embodiments, the clinical benefit (e.g., SD, PR and/or CR) is determined in accordance with Response Evaluation Criteria in Solid Tumors (RECIST). In some embodiments, the clinical benefit (e.g., SD, PR and/or CR) is determined in accordance RECIST guidelines.

[00345] In some embodiments, tumor response can be measured by, for example, the RECIST v 1.1 guidelines. The guidelines are provided by E.A. Eisenhauer, *et al.*, “New response evaluation criteria in solid tumors: Revised RECIST guideline (version 1.1.),” *Eur. J. of Cancer*, 45: 228-247 (2009), which is incorporated by reference in its entirety. The guidelines require, first, estimation of the overall tumor burden at baseline, which is used as a comparator for subsequent measurements. Tumors can be measured via use of any imaging system known in

the art, for example, by a CT scan, or an X-ray. Measurable disease is defined by the presence of at least one measurable lesion. In studies where the primary endpoint is tumor progression (either time to progression or proportion with progression at a fixed date), the protocol must specify if entry is restricted to those with measurable disease or whether patients having non-measurable disease only are also eligible.

[00346] When more than one measurable lesion is present at baseline, all lesions up to a maximum of five lesions total (and a maximum of two lesions per organ) representative of all involved organs should be identified as target lesions and will be recorded and measured at baseline (this means in instances where patients have only one or two organ sites involved a maximum of two and four lesions respectively will be recorded).

[00347] Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, but in addition should be those that lend themselves to reproducible repeated measurements.

[00348] Lymph nodes merit special mention since they are normal anatomical structures which may be visible by imaging even if not involved by tumor. Pathological nodes which are defined as measurable and may be identified as target lesions must meet the criterion of a short axis of ≥ 15 mm by CT scan. Only the short axis of these nodes will contribute to the baseline sum. The short axis of the node is the diameter normally used by radiologists to judge if a node is involved by solid tumour. Nodal size is normally reported as two dimensions in the plane in which the image is obtained (for CT scan this is almost always the axial plane; for MRI the plane of acquisition may be axial, sagittal or coronal). The smaller of these measures is the short axis.

[00349] For example, an abdominal node which is reported as being 20mm· 30mm has a short axis of 20mm and qualifies as a malignant, measurable node. In this example, 20mm should be recorded as the node measurement. All other pathological nodes (those with short axis ≥ 10 mm but < 15 mm) should be considered non-target lesions. Nodes that have a short axis < 10 mm are considered non-pathological and should not be recorded or followed.

[00350] A sum of the diameters (longest for non-nodal lesions, short axis for nodal lesions) for all target lesions will be calculated and reported as the baseline sum diameters. If lymph nodes are to be included in the sum, then as noted above, only the short axis is added into the sum. The baseline sum diameters will be used as reference to further characterize any objective tumor regression in the measurable dimension of the disease.

[00351] All other lesions (or sites of disease) including pathological lymph nodes should be identified as non-target lesions and should also be recorded at baseline. Measurements are not required and these lesions should be followed as ‘present’, ‘absent’, or in rare cases ‘unequivocal progression.’ In addition, it is possible to record multiple nontarget lesions involving the same organ as a single item on the case record form (e.g., ‘multiple enlarged pelvic lymph nodes’ or ‘multiple liver metastases’).

[00352] In some embodiments, tumor response can be measured by, for example, the immune-related RECIST (irRECIST) guidelines, which include immune related Response Criteria (irRC). In irRC, measurable lesions are measured that have at least one dimension with a minimum size of 10 mm (in the longest diameter by CT or MRI scan) for nonnodal lesions and greater than or equal to 15 mm for nodal lesions, or at least 20 mm by chest X-ray.

[00353] In some embodiments, Immune Related Response Criteria include CR (complete disappearance of all lesions (measurable or not, and no new lesions)); PR (decrease in tumor burden by 50% or more relative to baseline); SD (not meeting criteria for CR or PR in the absence of PD); or PD (an increase in tumor burden of at 25% or more relative to nadir). Detailed description of irRECIST can be found at Bohnsack et al., (2014) ESMO, ABSTRACT 4958 and Nishino et al., (2013) *Clin. Cancer Res.* 19(14): 3936-43.

[00354] In some embodiments, tumor response can be assessed by either irRECIST or RECIST version 1.1. In some embodiments, tumor response can be assessed by both irRECIST and RECIST version 1.1.

Pharmacokinetics

[00355] Pharmacokinetic data can be obtained by known techniques in the art. Due to the inherent variation in pharmacokinetic and pharmacodynamic parameters of drug metabolism in human subjects, appropriate pharmacokinetic and pharmacodynamic profile components describing a particular composition can vary. Typically, pharmacokinetic and pharmacodynamic profiles are based on the determination of the mean parameters of a group of subjects. The group of subjects includes any reasonable number of subjects suitable for determining a representative mean, for example, 5 subjects, 10 subjects, 16 subjects, 20 subjects, 25 subjects, 30 subjects, 35 subjects, or more. The mean is determined by calculating the average of all subject’s measurements for each parameter measured.

[00356] In some embodiments, a patient population includes one or more subjects (“a population of subjects”) suffering from metastatic disease.

[00357] In some embodiments, a patient population includes one or more subjects that is suffering from or susceptible to cancer. In some such embodiments, the cancer is non-small cell lung cancer (NSCLC), a hepatocellular cancer, a renal cancer, a melanoma, a cervical cancer, a colorectal cancer, a squamous cell carcinoma of the anogenital region (e.g., squamous cell carcinoma of the anus, penis, cervix, vagina, or vulva), a head and neck cancer, a triple negative breast cancer, an ovarian cancer or an endometrial cancer. In embodiments, a cancer is a solid tumor (e.g., an advanced solid tumor, a metastatic solid tumor, a MSS solid tumor, a MSI-H solid tumor, or a POLE mutant solid tumor). In embodiments, a cancer is a melanoma (e.g., an advanced melanoma, a metastatic melanoma, a MSS melanoma, a MSI-H melanoma, or a POLE mutant melanoma). In embodiments, a cancer is a lung cancer such as NSCLC (e.g., advanced NSCLC, metastatic NSCLC, MSI-H NSCLC, MSS NSCLC, POLE mutant NSCLC, EGFR-mutant NSCLC, or ALK-translocated NSCLC). In embodiments, a cancer is colorectal cancer (e.g., advanced colorectal cancer, metastatic colorectal cancer, MSS colorectal cancer, MSI-H colorectal cancer, or POLE mutant colorectal cancer). In some embodiments, a patient population includes one or more subjects (e.g., comprises or consists of subjects) suffering from cancer. For example, in some embodiments, a patient population suffering from cancer may have previously been treated with a prior therapy, for example, radiation and/or chemotherapy.

[00358] In some embodiments, the pharmacokinetic parameter(s) can be any parameters suitable for describing the present composition.

[00359] In some embodiments, the pharmacokinetic parameter(s) can be any parameters suitable for describing the present composition. For example, in some embodiments, the $C_{\max}$ is about 1 $\mu\text{g/ml}$; about 5 $\mu\text{g/ml}$, about 10 $\mu\text{g/ml}$, about 15 $\mu\text{g/ml}$, about 20 $\mu\text{g/ml}$, about 25 $\mu\text{g/ml}$, about 30 $\mu\text{g/ml}$, about 35 $\mu\text{g/ml}$, about 40 $\mu\text{g/ml}$, about 45 $\mu\text{g/ml}$, about 50 $\mu\text{g/ml}$, about 55 $\mu\text{g/ml}$, about 60 $\mu\text{g/ml}$, about 65 $\mu\text{g/ml}$, about 70 $\mu\text{g/ml}$, about 75 $\mu\text{g/ml}$, about 80 $\mu\text{g/ml}$, about 85 $\mu\text{g/ml}$, about 90 $\mu\text{g/ml}$, about 95 $\mu\text{g/ml}$, about 100 $\mu\text{g/ml}$, about 150 $\mu\text{g/ml}$, about 200 $\mu\text{g/ml}$, about 250 $\mu\text{g/ml}$, about 300 $\mu\text{g/ml}$, or any other $C_{\max}$ appropriate for describing a pharmacokinetic profile of an anti-TIM-3 antibody.

[00360] In some embodiments, the $T_{\max}$ is, for example, not greater than about 0.5 hours, not greater than about 1.0 hours, not greater than about 1.5 hours, not greater than about 2.0

hours, not greater than about 2.5 hours, or not greater than about 3.0 hours, or any other $T_{\max}$ appropriate for describing a pharmacokinetic profile of an anti-TIM-3 antibody.

[00361] In general, AUC as described herein is the measure of the area under the curve that corresponds to the concentration of an analyte over a selected time period following administration of a dose of a therapeutic agent. In some embodiments, such time period begins at the dose administration (i.e., 0 hours after dose administration) and extends for about 2, about 6, about 12, about 36, about 48, about 72, about 168, about 336, about 514, about 682, or more hours after the dose administration. In some embodiments, AUC is that achieved from 0 hours to 336 hours following administration of a dose described herein.

[00362] The $AUC_{(0-336h)}$ can be, for example, about 500 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 1000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 1500 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 2000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 2500 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 3000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 3500 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 4000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 4500 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 5000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 7500 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 10,000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 15,000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 20,000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 25,000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 30,000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 35,000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 40,000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 45,000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 50,000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 65,000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 75,000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, about 90,000 $\mu\text{g}\cdot\text{hr}/\text{mL}$, or any other $AUC_{(0-336h)}$ appropriate for describing a pharmacokinetic profile of a therapeutic agent (e.g., an anti-TIM-3 antibody). In some embodiments, an anti-TIM-3 antibody is administered according to a regimen that is demonstrated to achieve an average AUC_{0-336h} of the anti-TIM-3 antibody concentration-time curve in a patient population that is within 2500 $\text{h}\cdot\mu\text{g}/\text{mL}$ to 50000 $\text{h}\cdot\mu\text{g}/\text{mL}$.

[00363] In some embodiments, the AUC from 0 hours to the end of the dosing period is determined ($AUC_{(0-T_{\text{au}})}$). In some embodiments, the dosing period is one week, two weeks, three weeks, four weeks, five weeks, six weeks, seven weeks, eight weeks, nine weeks or ten weeks. In some embodiments, the dosing period is 2 weeks. In some embodiments, the dosing period is 3 weeks.

[00364] In some embodiments, an anti-TIM-3 antibody is administered according to a regimen demonstrated to achieve a response rate in relevant patient population such that no more than 50% to 80% of patients show progressive disease after 2, 4, 6, 8, 10, 12, 14, 16, 18, or 20 weeks following initiation of treatment. In some embodiments, no more than 80% of patients show progressive disease after at least 10 weeks following initiation of treatment.

[00365] In some embodiments, an anti-TIM-3 antibody is administered according to a regimen that is sufficient to achieve an average TIM-3 receptor occupancy of at least 50% to 90% after 1, 2, 3, 4, or 5 days following a single dose of the composition. In some embodiments, administration of a composition that delivers an anti-TIM-3 antibody sufficient to achieve an average TIM-3 receptor occupancy of at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75%, at least 80%, at least 85%, or at least 90% after 3 days following a single dose of the composition.

[00366] In some embodiments, an anti-TIM-3 antibody is administered according to a regimen sufficient to achieve an average stimulation ratio of at least 1 in a functional TIM-3 receptor occupancy assay after 3 days following a single dose of the TIM-3-binding agent.

[00367] In some embodiments, an anti-TIM-3 antibody is administered according to a regimen sufficient to achieve an average TIM-3 receptor occupancy of at least 75% over a first period of time, e.g., about 14 days to about 60 days following a single dose of the anti-TIM-3 antibody. In some embodiments, an anti-TIM-3 antibody is administered according to a regimen sufficient to achieve an average TIM-3 receptor occupancy of at least 75% over the first period of time (e.g., about 15 days to about 60 days; in some embodiments about 29 days) following a single dose of the anti-TIM-3 antibody.

[00368] In some embodiments, an anti-TIM-3 antibody is administered according to a regimen sufficient to achieve an average stimulation ratio of at least 1 in a functional TIM-3 receptor occupancy assay over a first period of time, e.g., about 14 days to about 60 days following a single dose of the anti-TIM-3 antibody. In some embodiments, an anti-TIM-3 antibody is administered according to a regimen sufficient to achieve an average stimulation ratio of at least 1 in a functional TIM-3 receptor occupancy assay over the first period of time (e.g., about 15 days to about 60 days; in some embodiments about 29 days) following a single dose of the anti-TIM-3 antibody.

EXAMPLES

[00369] The following examples are provided to illustrate, but not limit the claimed invention.

Example 1. Combinatorial Blockade of TIM-3 and PD-1 in Mouse Model Systems

[00370] The effect of TIM-3 and PD-1 inhibition in a mouse T cell exhaustion assay was studied (Burkhart et al. *Int Immunol.* 1999; 11:1625-1634). In this system, in vitro stimulation of mouse CD4+ T cell receptor transgenic T cells with a super-agonist altered-peptide ligand leads to an exhausted phenotype, characterized by increased expression of PD-1 and TIM-3 (**Figure 2A**). As shown in **Figure 2B**, the combination of anti-PD-1 and anti-TIM-3 antibodies was more effective than either agent alone in enhancing the production of IFN γ in this system.

Example 2. In Vivo Efficacy Study of TSR-042 and TSR-022 Combination

[00371] In addition to evaluating the combinatorial effect of targeting PD-1 and TIM-3 in vitro, the combination was tested in an animal model of lung cancer. A humanized mouse tumor model was employed, consisting of A549 human lung cancer cells (5×10^6 cells per mouse) implanted into huNOG-EXL mice. Neonatally derived CD34+ hematopoietic stem cells were engrafted into the mouse, and animals were treated with monoclonal blocking antibodies targeting PD-1 (TSR-042) and TIM-3 (TSR-022). Here, we show that the combination of anti-PD-1 and anti-TIM-3 has a beneficial antitumor effect when compared to either agent alone (**Figure 3**).

Example 3. Dosing Regimens for an Exemplary TIM-3-binding Agent

[00372] This example describes a multicenter, open-label, first-in-human Phase 1 study evaluating a TIM-3 binding agent (an anti-TIM-3 antibody), in patients with tumors. Specifically the dosage effects in patients with advanced solid tumors treated with a particular TIM-3 binding agent. A TIM-3 binding agent (TSR-022) as described in the present study comprises a humanized monoclonal anti-TIM-3 antibody comprising a heavy chain whose amino acid sequence comprises SEQ ID NO: 3 and a light chain whose amino acid sequence comprises SEQ ID NO:4 are evaluated. This anti-TIM-3 antibody utilizes a human IGHG4*01 heavy chain

gene, and a human IGKC*01 kappa light chain gene, as scaffolds. Further, there is a single Ser to Pro point mutation in the hinge region of the IgG4 heavy chain at the canonical S228 position.

[00373] Patients were included with histologically or cytologically proven advanced (unresectable) or metastatic solid tumor and who had disease progression after treatment with available therapies that are known to confer clinical benefit or who are intolerant to other known treatment(s).

[00374] The study comprises several parts: dose escalation and cohort expansion. Part 1a of the study (dose escalation) is intended, *inter alia*, to evaluate the safety, PK, and PDy profile, tolerability and anti-cancer effect of the anti-TIM-3 antibody. A modified 3+3 design was used for dose escalation at 0.03 mg/kg, 0.1 mg/kg, 0.3 mg/kg, 1.0 mg/kg, 3 mg/kg, and 10 mg/kg and higher every 2 weeks (Q2W) with the dose of the TIM-3 antibody not to exceed 20 mg/kg. Dose escalation has continued to 1.0 mg/kg Q2W and a MTD has not yet been identified. Part 1a could also comprise the testing of a “flat” dose, or a specific fixed number of milligrams of antibody (as opposed to the weight based dosing mg/kg). Flat doses could range from 200 mg to 1500 mg of anti-TIM-3 antibody. Part 1b of the study (combination with PD1 dose escalation cohorts) is intended, *inter alia*, to evaluate the safety, PK, and PDy profile, tolerability and anti-cancer effect of the anti-TIM-3 antibody in combination with an anti-PD-1 antibody wherein the anti-TIM-3 antibody is administered as a dose escalation. Dosing will comprise anti-TIM-3 at 1.0 mg/kg, 3 mg/kg, and 10 mg/kg or higher every 2 weeks (Q2W) or every 3 weeks (Q3W) with the dose of anti-TIM-3 to not exceed the MTD defined in part 1a of the study each in combination with an anti-PD-1 (exemplary anti-PD-1 antibodies are a humanized monoclonal anti-PD-1 antibody comprising a heavy chain whose amino acid sequence comprises SEQ ID NO: 13 and a light chain whose amino acid sequence comprises SEQ ID NO:14 nivolumab, or pembrolizumab). The anti-PD-1 antibody can be dosed at an approved dose and schedule for a marketed agent and on a weight basis at a dose of 3 or 10 mg/kg Q2W or Q3W or on a flat dose basis of 500 mg antibody Q2W or Q3W by using a modified 3+3 design.

[00375] Primary endpoints include determining the safety and tolerability of TSR-022 by Common Terminology Criteria for Adverse Events (CTCAE v4) and to determine the recommended phase 2 dose (RP2D) and schedule for monotherapy and combination with an anti-PD-1 antibody. Secondary endpoints include: pharmacokinetics (PK), overall response

rate, duration of response, disease control rate, progression-free survival, overall survival, and immunogenicity. Exploratory endpoints include pharmacodynamics.

[00376] **Figure 4A** shows doses used in the dose escalation study of Part 1a for TSR-022 monotherapy. In the dose escalation study, 38 patients with advanced cancer had been enrolled as of October 27. Doses of 0.03 mg/kg, 0.1 mg/kg, 0.3 mg/kg, 1 mg/kg, 3 mg/kg, and 10 mg/kg were administered to patients.

[00377] Adult patients with advanced or metastatic solid tumors who have disease progression or treatment in tolerance after treatment with available therapies and having adequate organ function and ECOG [Eastern Cooperative Oncology Group] performance status were treated. Prior treatment with immune checkpoint inhibitors was permitted. A summary of patient demographics is shown in **Figure 5** for the 38 patients enrolled. In the Part 1a study, there were 21 male patients and 17 female patients. The mean age was 60.1 years (SD = 13.5), with a median of 61.0 years (min = 25; max = 85). The mean number of prior treatment lines was 3.2 (SD = 2.3), with a median of 2.0 (min = 1; max = 10). 10 patients had an ECOG performance status of 0, and 28 had an ECOG performance status of 1. Tumor sites included colon, skin, ovary, breast, brain, head and neck, testis, pleura, lung, rectum, thyroid, liver, or esophagus.

[00378] After patients received the first dose of TSR-022 at various levels (0.03-10 mg/kg), the TSR-022 concentration in serum was monitored for two weeks and the pharmacokinetic (PK) behaviors were characterized. **Figure 6A** shows serum concentration versus time profiles for all treatment groups in Part 1. TSR-022 demonstrated linear PK behavior for the studied doses of 0.03-10 mg/kg.

[00379] **Figure 6B** shows TIM-3 occupancy on circulating monocytes as measured by flow cytometry from whole blood samples collected from patients treated with anti-TIM-3. Receptor occupancy on peripheral monocytes was shown to correlate with TSR-022 exposure.

[00380] TIM-3 receptor occupancy (RO) by TSR-022 on circulating CD14⁺ monocytes was measured by flow cytometry. Briefly, whole blood samples from patients treated with TSR-022 were stained with anti-CD14, a non-competing anti-TIM-3 antibody (indicating total TIM-3) and a competing anti-TIM-3 antibody (indicating unbound or free TIM-3). TIM-3 occupancy by infused TSR-022 was estimated as the ratio of free TIM-3 on CD14⁺ cells to the total TIM-3 on CD14⁺ cells. A decrease in the ratio indicates an increase in TSR-022 bound TIM-3 receptor. To measure binding in the RO assay, whole blood was collected from patients at baseline (Day 1

predose), 48 hours following the first dose (Day 3) and prior to the second dose (Day 15 predose) for the Q2W schedule. Additional samples were collected on Day 22 and Day 29 following the first dose for a subset of patients who did not receive the second dose. Samples for occupancy measurement on the Q3W schedule were collected as follows: Day 1 predose, Day 5, Day 15 and Day 22. Ex vivo TSR-022 saturated healthy donor samples (“sat”) as well as controls lacking detection antibodies (“Bkgd”) were included as controls.

[00381] **Figures 7A-7C** respectively show receptor occupancy studies of the anti-TIM-3 antibody (TSR-022) administered at doses of 1 mg/kg (**Fig. 7A**), 3 mg/kg (**Fig. 7B**), and 10 mg/kg (**Fig. 7C**) every two weeks (Q2W). All samples were pre-dose, with occupancy measured after a single dose on day 1. The occupancy ratio (free TIM-3:total TIM-3) is determined at various time points (for example, timepoints can include day 1, day 3, day 15, day 22, and day 29). The receptor occupancy is maximal on day 3 across doses of TSR-022. At 3 mg/kg, maximal occupancy is maintained through day 29, and comparable results are obtained for the 10 mg/kg dose through the days of available data.

[00382] **Figure 8** provides a summary of dosages administered to patients (0.03-10 mg/kg) and the duration of treatment during the Part 1a studies.

[00383] The best response observed was also evaluated in efficacy evaluable patients (e.g., patients who received at least two doses and either had at least one post-baseline assessment or had discontinued treatment due to clinical progression prior to post-baseline tumor assessment. Stable disease (5/25 patients) and partial response (1/25 patients at 10 mg/kg) were observed as the best response in patients with rectal, thyroid, neuroendocrine, or head and neck cancer, or with soft tissue sarcoma.

[00384] **Figure 9** depicts brain scans from the patient with partial response at the 10/mg level. The patient is 42 years old with leiomyosarcoma that is metastatic to lung and kidney (kidney biopsy confirmed leiomyosarcoma: PTEN splice site, MYC amplification, ATRX mutation, CD36 mutation, RB loss, p53 loss; low mutation burden). The patient received 3 doses of TSR-022 at 10mg/kg before re-staging imaging. Additionally, the patient received gemcitabine and docetaxel for 5 months with demonstrated progression before entering the study. Pending confirmatory imaging, there appears to be a 32% tumor reduction at the 10 mg/kg dose level, and treatment is ongoing.

[00385] From the studies, TSR-022 monotherapy has been shown to be well-tolerated across multiple dose levels.

[00386] Part 2 of the study is intended, *inter alia*, are to evaluate safety and tolerability, PK, and PDy profile and anti-cancer effect of (i) the anti-TIM-3 antibody at fixed doses Q2W or Q3W or (ii) anti-TIM-3 antibody at fixed doses Q2W or Q3W in combination with anti-PD-1 antibodies at weight based or flat doses as indicated above. An anti-PD-1 antibody can be administered according to a regimen of 500 mg every three weeks (Q3W) for the initial treatment cycles (e.g., 500 mg administered Q3W for four treatment cycles) followed by administration of 1000 mg administered every six weeks (Q6W) until treatment is discontinued (e.g., due to disease progression).

[00387] Flat doses of about 100 mg to 1500 mg of the anti-TIM-3 antibody can be administered in monotherapy or in combination therapy. For example, a dose of about 100 mg, 200 mg, 300 mg, 400 mg, 500 mg, 600 mg, 700 mg, 800 mg, 900 mg, 1000 mg, 1100 mg, 1200 mg, 1300 mg, 1400 mg, or 1500 mg of the anti-TIM-3 antibody can be administered Q1W, Q2W, Q3W, Q4W, Q5W, or Q6W as monotherapy or in combination with an anti-PD-1 antibody that is administered 500 mg Q3W for four treatment cycles then as 1000 mg Q6W until treatment is discontinued (e.g., due to disease progression). In embodiments, a dose of about 100 mg, 200 mg, 300 mg, 400 mg, 500 mg, 600 mg, 700 mg, 800 mg, 900 mg, 1000 mg, 1100 mg, 1200 mg, 1300 mg, 1400 mg, or 1500 mg of the anti-TIM-3 antibody can be administered Q2W or Q3W as monotherapy or in combination with an anti-PD-1 antibody that is administered 500 mg Q3W for four treatment cycles then as 1000 mg Q6W until treatment is discontinued (e.g., due to disease progression). A 100 mg, 200 mg, 300 mg, 500 mg, 800 mg, 1000 mg, or 1200 mg dose of the anti-TIM-3 antibody can be administered as a monotherapy once each week (Q1W), once every two weeks (Q2W), once every three weeks (Q3W), once every four weeks (Q4W), once every five weeks (Q5W), or once every three weeks (Q6W). A 100 mg, 200 mg, 300 mg, 500 mg, 800 mg, 1000 mg, or 1200 mg dose of the anti-TIM-3 antibody can be administered as a monotherapy once every two weeks (Q2W) or once every three weeks (Q3W).

[00388] Receptor occupancy studies have been done based on a flat dose of 100 mg of the anti-TIM-3 antibody administered once every three weeks (Q3W). All samples were pre-dose, with occupancy measured after a single dose on day 1. As shown in **Figure 10**, target coverage

in the periphery is achieved at 100 mg for the duration of the dosing interval. Further, receptor occupancy at 100 mg Q3W is comparable to 3 mg/kg Q2W (**Figure 7B**).

[00389] Receptor occupancy studies following administration of a 300 mg flat dose of an exemplary anti-TIM-3 antibody (TSR-022) administered in combination with a 500 mg flat dose of an exemplary anti-PD-1 antibody (TSR-042) were also conducted according to methods described herein, and data are shown in **Figure 11**. A second dose of the TSR-022 was administered on day 22, with the RO sample collected prior to the second dose. Using a flat dose of 300 mg TSR-022, high RO can be maintained over the measured time period.

[00390] **Figure 12** is a composite of mean receptor occupancy data for doses of an exemplary anti-TIM-3 antibody (TSR-022) of 1 mg/kg, 3 mg/kg, 10 mg/kg and of flat doses of 100 mg, 300 mg, and 1200 mg. The figure shows that high occupancy ratios (free TIM-3:total TIM-3) can be achieved using various doses of TSR-022 as measured over a range of days.

[00391] These monotherapy or combination regimens will be tested in specific tumor types, which may include anti-PD1/L1 treated melanoma, anti-PD1/L1 treated NSCLC, colorectal cancer, ovarian cancer, renal cell carcinoma, hepatocellular carcinoma, and/or breast cancer. **Figure 4B** provides an overview of the Part 2 expansion cohort study.

EQUIVALENTS

[00392] The articles “a” and “an” as used herein in the specification and in the claims, unless clearly indicated to the contrary, should be understood to include the plural referents. Claims or descriptions that include “or” between one or more members of a group are considered satisfied if one, more than one, or all of the group members are present in, employed in, or otherwise relevant to a given product or process unless indicated to the contrary or otherwise evident from the context. The invention includes embodiments in which exactly one member of the group is present in, employed in, or otherwise relevant to a given product or process. The invention also includes embodiments in which more than one, or the entire group members are present in, employed in, or otherwise relevant to a given product or process. Furthermore, it is to be understood that the invention encompasses all variations, combinations, and permutations in which one or more limitations, elements, clauses, descriptive terms, etc., from one or more of the listed claims is introduced into another claim dependent on the same base claim (or, as relevant, any other claim) unless otherwise indicated or unless it would be evident to one of ordinary skill

in the art that a contradiction or inconsistency would arise. Where elements are presented as lists, (e.g., in Markush group or similar format) it is to be understood that each subgroup of the elements is also disclosed, and any element(s) can be removed from the group. It should be understood that, in general, where the invention, or aspects of the invention, is/are referred to as comprising particular elements, features, etc., certain embodiments of the invention or aspects of the invention consist, or consist essentially of, such elements, features, etc. For purposes of simplicity those embodiments have not in every case been specifically set forth in so many words herein. It should also be understood that any embodiment or aspect of the invention can be explicitly excluded from the claims, regardless of whether the specific exclusion is recited in the specification. The publications, websites and other reference materials referenced herein to describe the background of the invention and to provide additional detail regarding its practice are hereby incorporated by reference.

CLAIMS

What is claimed is

1. A method of treating a disorder in a subject that is responsive to T Cell Immunoglobulin and Mucin Protein 3 (TIM-3) inhibition, which method comprises administering a therapeutically effective dose of an agent that is capable of inhibiting TIM-3 signaling, wherein the therapeutically effective dose is: about 1, 3 or 10 mg/kg; a flat dose between about 100 - 1500 mg; a flat dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg; a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; a flat dose about 900 mg; a flat dose about 1000 mg; a flat dose about 1100 mg; a flat dose about 1200 mg; a flat dose about 1300 mg; a flat dose about 1400 mg; a flat dose about 1500 mg; about 1 mg/kg; about 3 mg/kg; or about 10 mg/kg.
2. A method of increasing T cell activation or T cell effector function in a subject that is responsive to T Cell Immunoglobulin and Mucin Protein 3 (TIM-3) inhibition, which method comprises administering a therapeutically effective dose of an agent that is capable of inhibiting TIM-3 signaling, wherein the therapeutically effective dose is: about 1, 3 or 10 mg/kg; a flat dose between about 100 - 1500 mg; a flat dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg; a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; a flat dose about 900 mg; a flat dose about 1000 mg; a flat dose about 1100 mg; a flat dose about 1200 mg; a flat dose about 1300 mg; a flat dose about 1400 mg; a flat dose about 1500 mg; about 1 mg/kg; about 3 mg/kg; or about 10 mg/kg.
3. A method of reducing tumors or inhibiting the growth of tumor cells in a subject that is responsive to T Cell Immunoglobulin and Mucin Protein 3 (TIM-3) inhibition, which method comprises administering a therapeutically effective dose of an agent that is capable of inhibiting TIM-3 signaling, wherein the therapeutically effective dose is: about 1, 3 or 10 mg/kg; a flat dose between about 100 - 1500 mg; a flat dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg; a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; a flat dose about 900 mg; a flat dose about 1000 mg; a flat dose about 1100 mg; a flat dose about 1200 mg; a flat dose about 1300 mg; a flat

dose about 1400 mg; a flat dose about 1500 mg; about 1 mg/kg; about 3 mg/kg; or about 10 mg/kg.

4. A method of inducing an immune response in a subject that is responsive to T Cell Immunoglobulin and Mucin Protein 3 (TIM-3) inhibition, which method comprises administering a therapeutically effective dose of an agent that is capable of inhibiting TIM-3 signaling, wherein the therapeutically effective dose is: about 1, 3 or 10 mg/kg; a flat dose between about 100 - 1500 mg; a flat dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg; a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; a flat dose about 900 mg; a flat dose about 1000 mg; a flat dose about 1100 mg; a flat dose about 1200 mg; a flat dose about 1300 mg; a flat dose about 1400 mg; a flat dose about 1500 mg; about 1 mg/kg; about 3 mg/kg; or about 10 mg/kg.

5. A method of enhancing an immune response or increasing the activity of an immune cell in a subject that is responsive to T Cell Immunoglobulin and Mucin Protein 3 (TIM-3) inhibition, which method comprises administering a therapeutically effective dose of an agent that is capable of inhibiting TIM-3 signaling, wherein the therapeutically effective dose is: about 1, 3 or 10 mg/kg; a flat dose between about 100 - 1500 mg; a flat dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg; a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; a flat dose about 900 mg; a flat dose about 1000 mg; a flat dose about 1100 mg; a flat dose about 1200 mg; a flat dose about 1300 mg; a flat dose about 1400 mg; a flat dose about 1500 mg; about 1 mg/kg; about 3 mg/kg; or about 10 mg/kg.

6. The method of claim 5, wherein the immune response is a humoral or cell mediated immune response.

7. The method of claim 6, wherein the immune response is a CD4 or CD8 T cell response.

8. The method of claim 6, wherein the immune response is a B cell response.

9. The method of any one of claims 1-8, wherein the therapeutically effective dose is about 1 mg/kg.
10. The method of any one of claims 1-8, wherein the therapeutically effective dose is about 3 mg/kg.
11. The method of any one of claims 1-8, wherein the therapeutically effective dose is about 10 mg/kg.
12. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 100 mg.
13. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 200 mg.
14. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 300 mg.
15. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 400 mg.
16. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 500 mg.
17. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 600 mg.
18. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 700 mg.

19. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 800 mg.
20. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 900 mg.
21. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 1000 mg.
22. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 1100 mg.
23. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 1200 mg.
24. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 1300 mg.
25. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 1400 mg.
26. The method of any one of claims 1-8, wherein the therapeutically effective dose is a flat dose about 1500 mg.
27. The method of any one of the preceding claims, wherein the agent is administered at the administration interval of once a week, once every 2 weeks, once every 3 weeks, once every 4 weeks, once every 5 weeks, or once every 6 weeks.
28. The method of claim 27, wherein the agent is administered at the administration interval of once every 3 weeks.

29. The method of claim 28, wherein the therapeutically effective dose is a flat dose of about 100 mg, 300 mg, 500 mg, or 900 mg.
30. The method of any one of the preceding claims, wherein the agent is administered for a period sufficient to achieve clinical benefit.
31. The method of claim 30, wherein the clinical benefit is stable disease (“SD”), a partial response (“PR”) and/or a complete response (“CR”).
32. The method of claim 31, wherein the PR or CR is determined in accordance with Response Evaluation Criteria in Solid Tumors (RECIST).
33. The method of any one of claims 30-32, wherein the agent is administered for a longer period to maintain clinical benefit.
34. The method of any one of the preceding claims, wherein the agent is administered for the period of at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 weeks, or more.
35. The method of any one of the preceding claims, wherein the disorder is a T-cell dysfunctional disorder.
36. The method of any one of the preceding claims, wherein the disorder is cancer.
37. The method of claim 36, wherein the cancer is:
- i) a cancer associated with a high tumor mutation burden (TMB);
 - ii) a cancer that is microsatellite stable (MSS),
 - iii) a cancer that is characterized by microsatellite instability,
 - iv) a cancer that has a high microsatellite instability status (MSI-H),

- v) a cancer that has a low microsatellite instability status (MSI-L),
- vi) a cancer associated with high TMB and MSI-H,
- vii) a cancer associated with high TMB and MSI-L or MSS,
- viii) a cancer that has a defective DNA mismatch repair system,
- ix) a cancer that has a defect in a DNA mismatch repair gene,
- x) a hypermutated cancer,
- xi) a cancer comprising a mutation in polymerase delta (POLD),
- xii) a cancer comprising a mutation in polymerase epsilon (POLE),
- xiii) a cancer that has homologous recombination repair deficiency/homologous repair deficiency (“HRD”);
- xiv) adenocarcinoma, endometrial cancer, breast cancer, ovarian cancer, cervical cancer, fallopian tube cancer, testicular cancer, primary peritoneal cancer, colon cancer, colorectal cancer, stomach cancer, small intestine cancer, squamous cell carcinoma of the anus, squamous cell carcinoma of the penis, squamous cell carcinoma of the cervix, squamous cell carcinoma of the vagina, squamous cell carcinoma of the vulva, soft tissue sarcoma, melanoma, renal cell carcinoma, lung cancer, non-small cell lung cancer, adenocarcinoma of the lung, squamous cell carcinoma of the lung, stomach cancer, bladder cancer, gall bladder cancer, liver cancer, thyroid cancer, laryngeal cancer, salivary gland cancer, esophageal cancer, head and neck cancer, squamous cell carcinoma of the head and neck, prostate cancer, pancreatic cancer, mesothelioma, Merkel cell carcinoma, sarcoma, glioblastoma, a hematological cancer, multiple myeloma, B-cell lymphoma, T-cell lymphoma, Hodgkin’s lymphoma/primary mediastinal B-cell lymphoma, chronic myelogenous leukemia, acute myeloid leukemia, acute lymphoblastic leukemia, non-Hodgkin’s lymphoma, neuroblastoma, a CNS tumor, diffuse intrinsic pontine glioma (DIPG), Ewing’s sarcoma, embryonal rhabdomyosarcoma, osteosarcoma, or Wilms tumor, or

xiii) a cancer of xiv), wherein the cancer is MSS or MSI-L, is characterized by microsatellite instability, is MSI-H, has high TMB, has high TMB and is MSS or MSI-L, has high TMB and is MSI-H, has a defective DNA mismatch repair system, has a defect in a DNA mismatch repair gene, is a hypermutated cancer, is an HRD cancer, comprises a mutation in polymerase delta (POLD), or comprises a mutation in polymerase epsilon (POLE).

38. The method of claim 36, wherein the cancer is a cancer that has homologous recombination repair deficiency/homologous repair deficiency (“HRD”).

39. The method of claim 35, wherein the cancer is endometrial cancer, optionally MSI-H or MSS/MSI-L endometrial cancer.

40. The method of claim 35, wherein the cancer is a MSI-H cancer comprising a mutation in POLE or POLD, optionally a MSI-H non-endometrial cancer comprising a mutation in POLE or POLD.

41. The method of claim 35, wherein the cancer is breast cancer, optionally triple negative breast cancer (TNBC).

42. The method of claim 35, wherein the cancer is ovarian cancer, optionally epithelial ovarian cancer.

43. The method of claim 35, wherein the cancer is lung cancer, optionally non-small cell lung cancer.

44. The method of claim 35, wherein the cancer is melanoma.

45. The method of claim 35, wherein the cancer is colorectal cancer.

46. The method of claim 35, wherein the cancer is squamous cell carcinoma of the anus, squamous cell carcinoma of the penis, squamous cell carcinoma of the cervix, squamous cell carcinoma of the vagina, or squamous cell carcinoma of the vulva.

47. The method of claim 35, wherein the cancer is acute myeloid leukemia.

48. The method of claim 35, wherein the cancer is acute lymphoblastic leukemia.
49. The method of claim 35, wherein the cancer is non-Hodgkin's lymphoma.
50. The method of claim 35, wherein the cancer is Hodgkin's lymphoma.
51. The method of claim 35, wherein the cancer is neuroblastoma.
52. The method of claim 35, wherein the cancer is a CNS tumor.
53. The method of claim 35, wherein the cancer is diffuse intrinsic pontine glioma (DIPG).
54. The method of claim 35, wherein the cancer is Ewing's sarcoma.
55. The method of claim 35, wherein the cancer is embryonal rhabdomyosarcoma.
56. The method of claim 35, wherein the cancer is osteosarcoma.
57. The method of claim 35, wherein the cancer is Wilms tumor.
58. The method of claim 35, wherein the cancer is soft tissue sarcoma.
59. The method of claim 48, wherein the cancer is leiomyosarcoma.
60. The method of any one of the preceding claims, wherein the subject has been further administered or will be administered an immune checkpoint inhibitor, such that the mammal receives the agent and the immune checkpoint inhibitor.
61. The method of claim 60, comprising administering one, two, or three immune checkpoint inhibitors.
62. The method of claim 60 or 61, wherein the immune checkpoint inhibitor is an agent that inhibits programmed death-1 protein (PD-1) signaling, cytotoxic T-lymphocyte-associated protein 4 (CTLA-4), lymphocyte activation gene-3 (LAG-3), T cell immunoglobulin and ITIM

domain (TIGIT), indoleamine 2,3-dioxygenase (IDO), or colony-stimulating factor 1 receptor (CSF1R).

63. The method of claim 62, wherein the immune checkpoint inhibitor is a PD-1 inhibitor.

64. The method of claim 63, wherein the PD-1 inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, a toxin, or a PD-1 binding agent.

65. The method of claim 63 or 64, wherein the PD-1 inhibitor is a PD-1-binding agent.

66. The method of claim 65, wherein the PD-1 binding agent is an antibody, an antibody conjugate, or an antigen-binding fragment thereof.

67. The method of claim 66, wherein the PD-1 inhibitor is nivolumab, pembrolizumab, PDR-001, tislelizumab (BGB-A317), cemiplimab (REGN2810), LY-3300054, JNJ-63723283, MGA012, BI-754091, IBI-308, camrelizumab (HR-301210), BCD-100, JS-001, CX-072, AMP-514 / MEDI-0680, AGEN-2034, CS1001, TSR-042, Sym-021, PF-06801591, LZM009, KN-035, AB122, genolimzumab (CBT-501), AK 104, or GLS-010, or derivatives thereof.

68. The method of claim 63 or 64, wherein the PD-1 inhibitor is a PD-L1/L2 binding agent.

69. The method of claim 68, wherein the PD-L1/L2 binding agent is an antibody, an antibody conjugate, or an antigen-binding fragment thereof.

70. The method of claim 69, wherein the PD-L1/L2 binding agent durvalumab, atezolizumab, avelumab, BGB-A333, SHR-1316, FAZ-053, CK-301, or, PD-L1 millamolecule, or derivatives thereof.

71. The method of any one of claims 63-65, wherein the PD-1 binding agent is an anti-PD-1 antibody comprising a heavy chain comprising SEQ ID NO: 13 and a light chain comprising SEQ ID NO: 14.

72. The method of any one of claims 63-71, wherein PD-1 inhibitor is administered to the subject periodically at a dose of about 500 mg or 1000 mg.

73. The method of claim 72, wherein the PD-1 inhibitor is administered to the subject periodically at a dose of about 500 mg.

74. The method of claim 72 or 73, wherein the PD-1 inhibitor is administered to the subject once every 3 weeks.

75. The method of any one of claims 71-74, wherein the PD-1 inhibitor is administered for 2, 3, 4, 5, 6, or more cycles.

76. The method of claim 75, wherein the PD-1 inhibitor is administered for 3, 4, or 5 cycles.

77. The method of claim 72, wherein the PD-1 inhibitor is administered to the subject periodically at a dose of about 1000 mg.

78. The method of claim 72 or 77, wherein the PD-1 inhibitor is administered to the subject once every 6 weeks or more.

79. The method of claim 78, wherein the PD-1 inhibitor is administered to the subject once every six weeks.

80. The method of claim 78, wherein the PD-1 inhibitor is administered at a first dose of about 500 mg once every 3 weeks for 3, 4, or 5 cycles followed by a second dose of about 1000 mg once every 6 weeks or more.

81. The method of claim 80, wherein the PD-1 inhibitor is administered at a first dose of about 500 mg once every 3 weeks for 3, 4, or 5 cycles followed by a second dose of about 1000 mg once every 6 weeks.

82. The method of claim 80, wherein the PD-1 inhibitor is administered at a first dose of about 500 mg once every 3 weeks for 3 cycles followed by a second dose of about 1000 mg once every 6 weeks or more.

83. The method of claim 80, wherein the PD-1 inhibitor is administered at a first dose of about 500 mg once every 3 weeks for 4 cycles followed by a second dose of about 1000 mg once every 6 weeks or more.

84. The method of claim 80, wherein the PD-1 inhibitor is administered at a first dose of about 500 mg once every 3 weeks for 5 cycles followed by a second dose of about 1000 mg once every 6 weeks or more.

85. The method of any one of claims 82-74, wherein the second dose of 1000 mg is administered once every 6 weeks.

86. The method of claim 60 or 61, wherein the immune checkpoint inhibitor is a CTLA-4 inhibitor.

87. The method of claim 86, wherein the CTLA-4 inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, a toxin, or a CTLA-4 binding agent.

88. The method of claim 87, wherein the CTLA-4 inhibitor is a CTLA-4 binding agent.

89. The method of claim 88, wherein the CTLA-4 binding agent is an antibody, an antibody conjugate, or an antigen-binding fragment thereof.

90. The method of claim 60 or 61, wherein the immune checkpoint inhibitor is a LAG-3 inhibitor.

91. The method of claim 90, wherein the LAG-3 inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, a toxin, or a LAG-3 binding agent.

92. The method of claim 91, wherein the LAG-3 inhibitor is a LAG-3 binding agent.
93. The method of claim 92, wherein the LAG-3 binding agent is an antibody, an antibody conjugate, or an antigen-binding fragment thereof.
94. The method of claim 60 or 61, wherein the immune checkpoint inhibitor is a TIGIT inhibitor.
95. The method of claim 94, wherein the TIGIT inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, a toxin, or a TIGIT binding agent.
96. The method of claim 95, wherein the TIGIT inhibitor is a TIGIT binding agent.
97. The method of claim 96, wherein the TIGIT binding agent is an antibody, an antibody conjugate, or an antigen-binding fragment thereof.
98. The method of claim 60 or 61, wherein the immune checkpoint inhibitor is an IDO inhibitor.
99. The method of claim 98, wherein the IDO inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, a toxin, or an IDO binding agent.
100. The method of claim 99, wherein the IDO inhibitor is a small molecule.
101. The method of claim 99, wherein the IDO inhibitor is an IDO binding agent, optionally an IDO binding agent that is an antibody, an antibody conjugate, or an antigen-binding fragment thereof.
102. The method of claim 60 or 61, wherein the immune checkpoint inhibitor is a CSF1R inhibitor.

103. The method of claim 102, wherein the CSF1R inhibitor is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, a toxin, or an IDO binding agent.

104. The method of claim 103, wherein the CSF1R inhibitor is a small molecule.

105. The method of claim 103, wherein the CSF1R inhibitor is a CSF1R binding agent, optionally a CSF1R binding agent that is an antibody, an antibody conjugate, or an antigen-binding fragment thereof.

106. The method of any one of claims 60-105, comprising administering at least two of the immune checkpoint inhibitors.

107. The method of claim 106, further comprising administering a third checkpoint inhibitor.

108. The method of claim 106 or 107, wherein the subject receives treatment with each of the agent, a PD-1 inhibitor, and a LAG-3 inhibitor, such that the subject receives all three.

109. The method of claim 108, further comprising the subject receiving treatment with a CTLA-4 inhibitor, such that the subject receives all four.

110. The method of any one of the preceding claims, wherein the subject has further been administered or will be administered an agent that inhibits poly (ADP-ribose) polymerase (PARP).

111. The method of claim 110, wherein the agent that inhibits PARP is a small molecule, a nucleic acid, a polypeptide (e.g., an antibody), a carbohydrate, a lipid, a metal, or a toxin.

112. The method of claim 110 or 111, wherein the agent that inhibits PARP is selected from the group consisting of: ABT-767, AZD 2461, BGB-290, BGP 15, CEP 8983, CEP 9722, DR 2313, E7016, E7449, fluzoparib (SHR 3162), IMP 4297, INO1001, JPI 289, JPI 547,

monoclonal antibody B3-LysPE40 conjugate, MP 124, niraparib (ZEJULA) (MK-4827), NU 1025, NU 1064, NU 1076, NU1085, olaparib (AZD2281), ONO2231, PD 128763, R 503, R554, rucaparib (RUBRACA) (AG-014699, PF-01367338), SBP 101, SC 101914, simmiparib, talazoparib (BMN-673), veliparib (ABT-888), WW 46, 2-(4-(trifluoromethyl)phenyl)-7,8-dihydro-5H-thiopyrano[4,3-d]pyrimidin-4-ol, and salts or derivatives thereof.

113. The method of any one of claims 110-112, wherein the subject receives treatment with each of the agent, a PD-1 inhibitor, and an agent that inhibits PARP, such that the subject receives all three.

114. The method of claim 113, further comprising the subject receiving treatment with a LAG-3 inhibitor, such that the mammal receives all four.

115. The method of any one of claims 60-114, wherein the therapeutically effective dose of the agent is a flat dose about 100 mg.

116. The method of any one of claims 60-114, wherein the therapeutically effective dose of the agent is a flat dose about 300 mg.

117. The method of any one of claims 60-114, wherein the therapeutically effective dose of the agent is a flat dose about 500 mg.

118. The method of any one of claims 60-114, wherein the therapeutically effective dose of the agent is a flat dose about 900 mg.

119. The method of any one of the preceding claims, wherein the subject is resistant to treatment with a PD-1 inhibitor.

120. The method of any one of the preceding claims, wherein the subject is refractory to treatment with a PD-1 inhibitor.

121. The method of any one of the preceding claims, wherein the method sensitizes the subject to treatment with a PD-1 inhibitor.
122. The method of any one of the preceding claims, wherein the subject comprises an exhausted immune cell.
123. The method of claim 122, wherein the exhausted immune cell is an exhausted T cell.
124. The method of any one of the preceding claims, wherein the subject is human.
125. The method of any one of the preceding claims, wherein the subject has previously been treated with one or more different cancer treatment modalities.
126. The method of claim 125, wherein the subject has previously been treated with one or more of surgery, radiotherapy, chemotherapy, or immunotherapy.
127. The method of claim 125 or 126, wherein the subject has previously been treated with a cytotoxic therapy.
128. The method of any one of claims 125-127, wherein the subject has previously been treated with chemotherapy.
129. The method of any one of the preceding claims, wherein the method further comprises administering another therapeutic agent or treatment.
130. The method of claim 129, wherein the method further comprises administering one or more of surgery, a radiotherapy, a chemotherapy, an immunotherapy, an anti-angiogenic agent, or an anti-inflammatory.
131. The method of claim 129, wherein the method further comprises administering chemotherapy.

132. The method of any one of the preceding claims, wherein the agent is a TIM-3 binding agent.
133. The method of claim 132, wherein the TIM-3 binding agent is an antibody, an antibody conjugate, or an antigen-binding fragment thereof.
134. The method of claim 133, wherein the TIM-3 binding agent is an antibody.
135. The method of claim 132-134, wherein the TIM-3 binding agent comprises a heavy chain comprising one or more CDR sequences having at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NOs: 21, 22, or 23.
136. The method of claim 135, wherein the TIM-3 binding agent comprises a heavy chain comprising two or three CDRs that have sequences of SEQ ID NOs: 21, 22, or 23.
137. The method of any one of claims 132-136 wherein the TIM-3 binding agent comprises a light chain comprising one or more CDR sequences having at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NOs: 24, 25, or 26.
138. The method of claim 137 wherein the TIM-3 binding agent comprises a light chain comprising two or three CDRs that have sequences of SEQ ID NOs: 24, 25, or 26.
139. The method of claim any one of claims 132-134, wherein the TIM-3 binding agent comprises a heavy chain comprising one or more CDR sequences selected from SEQ ID NOs: 21, 22, or 23; and/or a light chain variable region with one or more CDR sequences selected from SED ID NOs: 24, 25, or 26.
140. The method of claim 139, wherein the TIM-3 binding agent comprises a heavy chain comprising three CDRs that have sequences of SEQ ID NOs: 21, 22, or 23; and/or a light chain comprising three CDRs that have sequences of SED ID NOs: 24, 25, or 26.
141. The method of any one of claims 132-134, wherein the TIM-3 binding agent comprises

an immunoglobulin heavy chain variable domain comprising an amino acid sequence having at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NO:1 or SEQ ID NO:7; and/or

an immunoglobulin light chain variable domain comprising an amino acid sequence having at least about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NO:2 or SEQ ID NO:8.

142. The method of claim 141, wherein the TIM-3 binding agent comprises an immunoglobulin heavy chain variable domain comprising an amino acid sequence of SEQ ID NO:1 or SEQ ID NO:7; and an immunoglobulin light chain variable domain comprising an amino acid sequence of SEQ ID NO:2 or SEQ ID NO:8.

143. The method of any one of claims 132-134, wherein the TIM-3 binding agent comprises an immunoglobulin heavy chain polypeptide comprising an amino acid sequence having about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NO:3.

144. The method of any one of claims 132-134 and 143, wherein the TIM-3 binding agent comprises an immunoglobulin light chain polypeptide comprising an amino acid sequence having about 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, or 99% sequence identity to SEQ ID NO:4.

145. The method of any one of claims 132-134, wherein the TIM-3 binding agent comprises a heavy chain polypeptide comprising SEQ ID NO:3 and a light chain polypeptide comprising SEQ ID NO: 4.

146. The method of any one of claims 132-145, wherein the TIM-3 binding agent is administered in an amount that is about 1 mg/kg, 3 mg/kg, or 10 mg/kg; or in an amount that is about a flat dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg; a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; a flat dose about 900 mg; a flat dose about 1000 mg; a flat dose about

1100 mg; a flat dose about 1200 mg; a flat dose about 1300 mg; a flat dose about 1400 mg; or a flat dose about 1500 mg.

147. The method of claim 146, wherein the TIM-3 binding agent is administered in an amount that is a flat dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg, a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; or a flat dose about 900 mg.

148. The method of claim 146, wherein the TIM-3 binding agent is administered once every two weeks, once every three weeks, or once every four weeks.

149. The method of any one of claim 132-148, wherein the TIM-3 binding agent is administered once every three weeks.

150. The method of any one of the preceding claims, wherein the agent is administered intravenously.

151. The method of claim 150, wherein the agent is administered by intravenous infusion.

152. A method of treating cancer, the method comprising:
administering to a patient in need of treatment an anti-T Cell Immunoglobulin and Mucin Domain-3 (TIM-3) antibody at a therapeutically effective dose at an administration interval for a period sufficient to achieve clinical benefit,

wherein the antibody comprises a heavy chain comprising three CDRs that have sequences of SEQ ID NOs: 21, 22, or 23; and/or a light chain comprising three CDRs that have sequences of SED ID NOs: 24, 25, or 26.

153. A method of treating cancer, the method comprising:
administering to a patient in need of treatment an anti-T Cell Immunoglobulin and Mucin Domain-3 (TIM-3) antibody at a therapeutically effective dose at an administration interval for a period sufficient to achieve clinical benefit,

wherein the anti-TIM-3 antibody comprises an immunoglobulin heavy chain variable domain comprising SEQ ID NO: 1 or SEQ ID NO: 7 and an immunoglobulin light chain variable domain comprising SEQ ID NO: 2 or SEQ ID NO: 8.

154. A method of treating cancer, the method comprising:

administering to a patient in need of treatment an anti-T Cell Immunoglobulin and Mucin Domain-3 (TIM-3) antibody at a therapeutically effective dose at an administration interval for a period sufficient to achieve clinical benefit,

wherein the anti-TIM-3 antibody comprises a heavy chain polypeptide comprising SEQ ID NO: 3 and a light chain polypeptide comprising SEQ ID NO: 4.

155. The method of any one of claims 152-154, wherein the clinical benefit is stable disease (“SD”), a partial response (“PR”) and/or a complete response (“CR”).

156. The method of any one of claims 152-155, wherein the PR or CR is determined in accordance with Response Evaluation Criteria in Solid Tumors (RECIST).

157. The method of any one of claims 152-157, wherein the patient has a cancer associated with a POLE (DNA polymerase epsilon) or a POLD (DNA polymerase delta) mutation.

158. The method of claim 157, wherein the POLE or POLD mutation is in an exonuclease domain.

159. The method of claim 157 or 158, wherein the POLE or POLD mutation is a germline mutation.

160. The method of claim 157 or 159, wherein the POLE or POLD mutation is a sporadic mutation.

161. The method of any one of claims 157-160, wherein the method further comprises a step of first identifying the patient having the cancer with the POLE or POLD mutation.

162. The method of claim 161, wherein the POLE or POLD mutation is identified using sequencing.
163. The method of any one of claims 152-162, wherein the patient has a cancer with microsatellite instability.
164. The method of claim 163, wherein the patient has a MSI-H cancer.
165. The method of claim 164, wherein the cancer is a MSI-H cancer comprising a mutation in POLE or POLD, optionally a MSI-H non-endometrial cancer comprising a mutation in POLE or POLD.
166. The method of claim 164, wherein the patient has a MSI-L cancer.
167. The method of any one of claims 152-162, wherein the patient has a microsatellite stable (MSS) cancer.
168. The method of any one of claims 152-167, wherein the patient has a solid tumor.
169. The method of claim 168, wherein the patient has an advanced stage solid tumor.
170. The method of claim 168, wherein the patient has a metastatic solid tumor.
171. The method of any one of claims 152-170, wherein the patient has a cancer selected from: a head and neck cancer, a lung cancer, a renal cancer, a bladder cancer, a melanoma, Merkel cell carcinoma, a cervical cancer, a vaginal cancer, a vulvar cancer, a uterine cancer, an endometrial cancer, an ovarian cancer, a fallopian tube cancer, a breast cancer, a prostate cancer, a salivary gland tumor, a thymoma, an adrenocortical carcinoma, an esophageal cancer, a gastric cancer, a colorectal cancer, an appendiceal cancer, a urothelial cell carcinoma, a squamous cell carcinoma, a soft tissue sarcoma, acute myeloid leukemia (AML), acute lymphoblastic leukemia (ALL), non-Hodgkin's lymphoma, Hodgkin's lymphoma, neuroblastoma, a CNS tumor, diffuse

intrinsic pontine glioma (DIPG), Ewing's sarcoma, embryonal rhabdomyosarcoma (ERS), osteosarcoma, or Wilms tumor.

172. The method of any one of claims 152-171, wherein the cancer has homologous recombination repair deficiency/homologous repair deficiency ("HRD").

173. The method of claim 171, wherein the patient has an endometrial cancer.

174. The method of claim 173, wherein the patient has an endometrial cancer with microsatellite instability.

175. The method of claim 173, wherein the patient has a MSI-H endometrial cancer.

176. The method of claim 173, wherein the patient has MSS/MSI-L endometrial cancer.

177. The method of claim 171, wherein the patient has breast cancer.

178. The method of claim 177, wherein the patient has triple negative breast cancer (TNBC).

179. The method of claim 171, wherein the patient has ovarian cancer.

180. The method of claim 179, wherein the ovarian cancer is epithelial ovarian cancer.

181. The method of claim 171, wherein the patient has a lung cancer.

182. The method of claim 181, wherein the lung cancer is a non-small cell lung cancer (NSCLC).

183. The method of claim 171, wherein the patient has a melanoma.

184. The method of claim 171, wherein the patient has a colorectal cancer.

185. The method of claim 171, wherein the patient has a squamous cell carcinoma.
186. The method of claim 185, wherein the squamous cell carcinoma is squamous cell carcinoma of the anus, squamous cell carcinoma of the penis, squamous cell carcinoma of the cervix, squamous cell carcinoma of the vagina, or squamous cell carcinoma of the vulva.
187. The method of claim 171, wherein the patient has a soft tissue sarcoma.
188. The method of claim 187, wherein the patient has leiomyosarcoma.
189. The method of any one of claims 152-167, wherein the patient has a hematological cancer.
190. The method of claim 189, wherein the hematological cancer is DLBCL, HL, NHL, FL, AML, ALL, or MM.
191. The method of any one of claims 152-190, wherein the patient has not previously been treated with a cancer treatment modality.
192. The method of any one of claims 152-190, wherein the patient has previously been treated with one or more different cancer treatment modalities.
193. The method of claim 192, wherein the one or more different cancer treatment modalities comprise surgery, radiotherapy, chemotherapy or immunotherapy.
194. The method of any one of claims 152-193, wherein the therapeutically effective dose is about 1, 3 or 10 mg/kg.
195. The method of any one of claims 152-193, wherein the therapeutically effective dose is a flat dose between about 100 - 1500 mg.

196. The method of claim 195, wherein the therapeutically effect dose is a flat dose about 100 mg; a flat dose about 200 mg; a flat dose about 300 mg; a flat dose about 400 mg; a flat dose about 500 mg; a flat dose about 600 mg; a flat dose about 700 mg; a flat dose about 800 mg; a flat dose about 900 mg; a flat dose about 1000 mg; a flat dose about 1100 mg; a flat dose about 1200 mg; a flat dose about 1300 mg; a flat dose about 1400 mg; or a flat dose about 1500 mg.

197. The method of claim 195, wherein the therapeutically effective dose is a flat dose between about 100 – 500 mg.

198. The method of claim 195, wherein the therapeutically effective dose is a flat dose between about 1000 – 1500 mg.

199. The method of claim 195, wherein the therapeutically effective dose is a flat dose about 100 mg.

200. The method of claim 195, wherein the therapeutically effective dose is a flat dose about 200 mg.

201. The method of claim 195, wherein the therapeutically effective dose is a flat dose about 300 mg.

202. The method of claim 195, wherein the therapeutically effective dose is a flat dose about 500 mg.

203. The method of claim 195, wherein the therapeutically effective dose is a flat dose about 900 mg.

204. The method of claim 195, wherein the therapeutically effective dose is a flat dose about 1000 mg.

205. The method of claim 195, wherein the therapeutically effective dose is a flat dose about 1100 mg.

206. The method of claim 195, wherein the therapeutically effective dose is a flat dose about 1200 mg.

207. The method of claim 195, wherein the therapeutically effective dose is a flat dose about 1300 mg.

208. The method of claim 195, wherein the therapeutically effective dose is a flat dose about 1400 mg.

209. The method of claim 195, wherein the therapeutically effective dose is a flat dose about 1500 mg.

210. The method of any one of claims 152-209, wherein the anti- TIM-3 antibody is administered at the administration interval of once a week, once every 2 weeks, once every 3 weeks, once every 4 weeks, once every 5 weeks, or once every 6 weeks.

211. The method of claim 210, wherein the anti-TIM-3 antibody is administered at the administration interval of once every 2 weeks.

212. The method of claim 210, wherein the anti-TIM-3 antibody is administered at the administration interval of once every 3 weeks.

213. The method of claim 212, wherein about 100 mg of the anti- TIM-3 antibody is administered.

214. The method of claim 212, wherein about 300 mg of the anti- TIM-3 antibody is administered.

215. The method of claim 212, wherein about 500 mg of the anti- TIM-3 antibody is administered.

216. The method of claim 212, wherein about 900 mg of the anti- TIM-3 antibody is administered.

217. The method of any one of claims 152-216, wherein the anti-TIM-3 antibody is administered for the period of at least 2, 4, 6, 8, 10, 12, 14, 16, 18, or 20 weeks.
218. The method of any one of claims 152-217, wherein the anti-TIM-3 antibody is administered intravenously.
219. The method of claim 218, wherein the anti-TIM-3 antibody is administered by intravenous infusion.
220. The method of any one of claims 152-219, wherein the anti-TIM-3 antibody is administered in conjunction with an additional therapy.
221. The method of claim 220, wherein the additional therapy is surgery, radiotherapy, chemotherapy or immunotherapy.
222. The method of claim 220, wherein the additional therapy is treatment with a PARP inhibitor.
223. The method of claim 222, wherein the PARP inhibitor is niraparib, olaparib, rucaparib, talazoparib, or veliparib.
224. The method of claim 220, wherein the additional therapy comprises treatment with an anti-PD-1 antibody.
225. The method of claim 224, wherein the anti-PD-1 antibody is or comprises TSR-042, nivolumab, pembrolizumab, PD-1VR or PD-1FL.
226. The method of claim 224 or 225, wherein the anti-PD-1 antibody is or comprises TSR-042.

227. The method of any one of claims 224-226, wherein the anti-PD-1 antibody comprises a heavy chain comprising SEQ ID NO: 13 and a light chain comprising SEQ ID NO: 14.

228. The method of any one of claims 224-227, wherein the anti-PD-1 antibody is administered to the patient periodically at a dose between about 100-1000 mg.

229. The method of claim 228, wherein the anti-PD-1 antibody is administered to the patient periodically at a dose of about 500 mg.

230. The method of claim 229, wherein the anti-PD-1 antibody is administered to the patient once every 3 weeks.

231. The method of claim 229 or 230, wherein the anti-PD-1 antibody is administered for 2-6 cycles.

232. The method of claim 231, wherein the anti-PD-1 antibody is administered for 3 cycles.

233. The method of claim 231, wherein the anti-PD-1 antibody is administered for 4 cycles.

234. The method of claim 231, wherein the anti-PD-1 antibody is administered for 5 cycles.

235. The method of claim 228, wherein the anti-PD-1 antibody is administered to the patient periodically at a dose of about 1000 mg.

236. The method of claim 235, wherein the anti-PD-1 antibody is administered to the patient once every 6 weeks or more.

237. The method of claim 228, wherein the anti-PD-1 antibody is administered at a first dose of about 500 mg once every 3 weeks for 3, 4, or 5 cycles followed by a second dose of about 1000 mg once every 6 weeks or more.

238. The method of claim 237, wherein the anti-PD-1 antibody is administered at a first dose of about 500 mg once every 3 weeks for 3 cycles followed by a second dose of about 1000 mg once every 6 weeks or more.

239. The method of claim 237, wherein the anti-PD-1 antibody is administered at a first dose of about 500 mg once every 3 weeks for 4 cycles followed by a second dose of about 1000 mg once every 6 weeks or more.

240. The method of claim 237, wherein the anti-PD-1 antibody is administered at a first dose of about 500 mg once every 3 weeks for 5 cycles followed by a second dose of about 1000 mg once every 6 weeks or more.

241. The method of any one of claims 237-240, wherein the second dose of 1000 mg is administered once every 6 weeks.

242. The method of any one of claims 237-241, wherein the therapeutically effective dose of the anti-TIM-3 antibody is a flat dose about 100 mg.

243. The method of any one of claims 237-241, wherein the therapeutically effective dose of the anti-TIM-3 antibody is a flat dose about 300 mg.

244. The method of any one of claims 237-241, wherein the therapeutically effective dose of the anti-TIM-3 antibody is a flat dose about 500 mg.

245. The method of any one of claims 237-241, wherein the therapeutically effective dose of the anti-TIM-3 antibody is a flat dose about 900 mg.

246. The method of any one of claims 242-245, wherein the anti-TIM-3 antibody is administered once every three weeks.