

**NOVEL *N*-ACYL-(3-SUBSTITUTED)-(8-SUBSTITUTED)-5,6-DIHYDRO-
[1,2,4]TRIAZOLO[4,3-a]PYRAZINES AS SELECTIVE NK-3 RECEPTOR
ANTAGONISTS, PHARMACEUTICAL COMPOSITION, METHODS FOR USE
IN NK-3 RECEPTOR-MEDIATED DISORDERS**

5

FIELD OF INVENTION

The present invention relates to novel *N*-acyl-(3-substituted)-(8-substituted)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazines including their pharmaceutically acceptable solvates that are selective antagonists to neurokinin-3 receptor (NK-3) and are useful as
10 therapeutic compounds, particularly in the treatment and/or prevention of a broad array of CNS and peripheral diseases or disorders.

BACKGROUND OF INVENTION

Tachykinin receptors are the targets of a family of structurally related peptides which
15 include substance P (SP), neurokinin A (NKA) and neurokinin B (NKB), named collectively "tachykinins". Tachykinins are synthesized in the central nervous system (CNS) and peripheral tissues, where they exert a variety of biological activities. Three tachykinin receptors are known which are named neurokinin-1 (NK-1), neurokinin-2 (NK-2) and neurokinin-3 (NK-3) receptors. Tachykinin receptors belong to the
20 rhodopsin-like seven membrane G-protein coupled receptors. SP has the highest affinity and is believed to be the endogenous ligand of NK-1, NKA for NK-2 receptor and NKB for NK-3 receptor, although cross-reactivity amongst these ligands does exist. The NK-1, NK-2 and NK-3 receptors have been identified in different species. NK-1 and NK-2 receptors are expressed in a wide variety of peripheral tissues and NK-1 receptors are
25 also expressed in the CNS; whereas NK-3 receptors are primarily expressed in the CNS.

The neurokinin receptors mediate a variety of tachykinin-stimulated biological effects that include transmission of excitatory neuronal signals in the CNS and periphery (e.g.

pain), modulation of smooth muscle contractile activity, modulation of immune and inflammatory responses, induction of hypotensive effects via dilatation of the peripheral vasculature and stimulation of endocrine and exocrine gland secretions.

In the CNS, the NK-3 receptor is expressed in regions including the medial prefrontal cortex, the hippocampus, the thalamus and the amygdala. Moreover, NK-3 receptors are expressed on dopaminergic neurons. Activation of NK-3 receptors has been shown to modulate dopamine, acetylcholine and serotonin release suggesting a therapeutic utility for NK-3 receptor modulators for the treatment of a variety of disorders including psychotic disorders, anxiety, depression, schizophrenia as well as obesity, pain or inflammation (Giardina et al., *Exp. Opin. Ther. Patents*, 2000, 10(6), 939-960; *Current Opinion in Investigational Drugs*, 2001, 2(7), 950-956 and Dawson and Smith, *Current Pharmaceutical Design*, 2010, 16, 344-357).

Schizophrenia is classified into subgroups. The paranoid type is characterized by delusions and hallucinations and absence of thought disorder, disorganized behavior, and affective flattening. In the disorganized type, which is also named 'hebephrenic schizophrenia' in the International Classification of Diseases (ICD), thought disorder and flat affect are present together. In the catatonic type, prominent psychomotor disturbances are evident, and symptoms may include catatonic stupor and waxy flexibility. In the undifferentiated type, psychotic symptoms are present but the criteria for paranoid, disorganized, or catatonic types have not been met. The symptoms of schizophrenia normally manifest themselves in three broad categories, i.e. positive, negative and cognitive symptoms. Positive symptoms are those, which represent an "excess" of normal experiences, such as hallucinations and delusions. Negative symptoms are those where the patient suffers from a lack of normal experiences, such as anhedonia and lack of social interaction. The cognitive symptoms relate to cognitive impairment in schizophrenics, such as a lack of sustained attention and deficits in decision making. The current antipsychotic drugs (APDs) are fairly successful in treating the positive symptoms but fare less well for the negative and cognitive symptoms. Contrary to that, NK-3 antagonists have been shown clinically to improve on both positive and negative symptoms in schizophrenics (Meltzer et al, *Am. J.*

Psychiatry, 2004, 161, 975-984) and ameliorate cognitive behavior of schizophrenics (Curr. Opin. Invest. Drug, 2005, 6, 717-721).

In rat, morphological studies provide evidence for putative interactions between NKB neurons and the hypothalamic reproductive axis (Krajewski et al, J. Comp. Neurol., 5 2005, 489(3), 372-386). In arcuate nucleus neurons, NKB expression co-localizes with estrogen receptor α and dynorphin, implicated in progesterone feedback to Gonadotropin Releasing Hormone (GnRH) secretion (Burke et al., J. Comp. Neurol., 2006, 498(5), 712-726; Goodman et al., Endocrinology, 2004, 145(6), 2959-2967). Moreover, NK-3 receptor is highly expressed in the hypothalamic arcuate nucleus in 10 neurons which are involved in the regulation of GnRH release.

WO 00/43008 discloses a method of suppressing gonadotropin and/or androgen production with specific NK-3 receptor antagonists. More particularly, the WO 00/43008 application relates to lowering luteinizing hormone (LH) blood level by administering an NK-3 receptor antagonist. Concurrently or alternatively with 15 gonadotropin suppression, WO 00/43008 also relates to suppression of androgen production with NK-3 receptor antagonists. Recently it has been postulated that NKB acts autodynamically on kisspeptin neurons in the arcuate nucleus to synchronize and shape the pulsatile secretion of kisspeptin and drive the release of GnRH from fibers in the median eminence (Navarro et al., J. of Neuroscience, 2009, 23(38), 11859-11866). 20 All these observations suggest a therapeutic utility for NK-3 receptor modulators for sex hormone-dependent diseases.

NK-3 receptors are also found in the human myenteric and submucosal plexus of the sigmoid colon as well as in the gastric fundus (Dass et al., Gastroenterol., 2002, 122 (Suppl 1), Abstract M1033) with particular expression noted on myenteric intrinsic 25 primary afferent neurons (IPANs) (Lomax and Furness, Cell Tissue Res, 2000, 302, 59-3). Intense stimulation of IPANs changes patterns of intestinal motility and intestinal sensitivity. Electrophysiology experiments have shown that activation of the NK-3 receptor changes the voltage threshold of action potentials in IPANs and promotes the generation of long-lasting plateau potentials (Copel et al., J Physiol, 2009, 587, 1461-

1479) that may sensitize these neurons to mechanical and chemical stimuli leading to effects on gut motility and secretion. Similarly, Irritable Bowel Syndrome (IBS) is characterized by patient hypersensitivity to mechanical and chemical stimuli. Thus, NK-3 antagonists have been tested in preclinical models of IBS where they have been shown to be effective to reduce nociceptive behavior caused by colo-rectal distension (Fioramonti et al., Neurogastroenterol Motil, 2003, 15, 363-369; Shafon et al., Neurogastroenterol Motil, 2004, 16, 223-231) and, on this basis, NK-3 antagonists have been advanced into clinical development for the treatment of IBS (Houghton et al., Neurogastroenterol Motil, 2007, 19, 732-743; Dukes et al., Gastroenterol, 2007, 132, A60).

Non-peptide antagonists have been developed for each of the tachykinin receptors. Some of them have been described as dual modulators able to modulate both NK-2 and NK-3 receptors (WO 06/120478). However, known non-peptide NK-3 receptor antagonists suffer from a number of drawbacks, notably poor safety profile and limited CNS penetrability that may limit the success of these compounds in clinical development.

On this basis, new potent and selective antagonists of NK-3 receptor may be of therapeutic value for the preparation of drugs useful in the treatment and/or prevention of CNS and peripheral diseases or disorders in which NKB and the NK-3 receptors are involved.


Target potency alone, which may be demonstrated by competitive binding data, is not sufficient for drug development. Rather, efficacy *in vivo* is contingent upon achieving a relevant “free” drug concentration relative to the target potency at the physiological site of action. Drug molecules typically bind reversibly to proteins and lipids in plasma. The “free” fraction refers to the drug concentration that is unbound and therefore available to engage the biological target and elicit pharmacological activity. This free fraction is commonly determined using plasma protein binding (PPB) assays. The free drug fraction is relevant to not only achieving the desired pharmacological activity, but also potentially undesirable activities including rapid hepatic metabolism (leading to high first-pass clearance and thereby poor oral bioavailability) as well as possible off-target

activities that can lead to safety concerns (for example, inhibition of hERG ion channel activity, a widely accepted marker of cardiovascular toxicity).

The invention thus encompasses compounds of general Formula I, their pharmaceutically acceptable solvates as well as methods of use of such compounds or compositions comprising such compounds as antagonists to the NK-3 receptor. Compounds of Formula I are *N*-acyl-(3-substituted)-(8-substituted)-5,6-dihydro-[1,2,4]triazolo[4,3-*a*]pyrazines. The compounds of the invention are generally disclosed in international patent application WO2011/121137 but none is specifically exemplified therein. On another hand, unsubstituted and thus non-chiral 5,6,7,8-tetrahydro[1,2,4]triazolo[4,3-*a*]pyrazines have been disclosed in WO2010/125102 as modulators of an unrelated target, namely P2X7.

SUMMARY

In one aspect, the present disclosure provides a compound of Formula I:



wherein:

R¹ is H, F or methyl;

R^{1'} is H;

R² is H, F, Cl or methoxy;

R^{2'} is H or F;

R³ is H, F, Cl, methyl, trifluoromethyl or nitrile;

R⁴ is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl;


R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl;

X^1 is N and X^2 is S or O; or X^1 is S and X^2 is N;

$\equiv$ represents a single or a double bond depending on X^1 and X^2 ;

*--- stands for the (*R*)-enantiomer or for the racemate of compound of Formula I.

In one embodiment, the disclosure provides compounds of general Formula I:



and pharmaceutically acceptable solvates thereof, wherein:

R^1 is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;

$R^{2'}$ is H or F;

R^3 is H, F, Cl, methyl, trifluoromethyl, nitrile or R^3 is thiophen-2-yl under the condition that R^5 is not methyl;

R^4 is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

X^1 is N and X^2 is S or O; or X^1 is S and X^2 is N;

== represents a single or a double bond depending on X^1 and X^2 ;

*-- stands for the (*R*)-enantiomer or for the racemate of compound of Formula I.

In one embodiment, the present disclosure provides a pharmaceutical composition comprising at least one compound according to the disclosure or a pharmaceutically acceptable solvate thereof.

In another aspect, the present disclosure provides a pharmaceutical composition comprising a compound as disclosed above and at least one pharmaceutically acceptable carrier, diluent, excipient and/or adjuvant.

The disclosure also relates to the use of the above compounds or their pharmaceutically acceptable solvates as modulators of NK-3 receptors, preferably as antagonists of NK-3 receptors.

The disclosure also relates to the use of the above compounds or their pharmaceutically acceptable solvates as lowering agents of the circulating LH levels.

The disclosure further provides methods of treatment and/or prevention of depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic

ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis comprising the administration of a therapeutically effective amount of a compound or pharmaceutically acceptable solvate of Formula I, to a patient in need thereof. The disclosure further provides methods of treatment and/or prevention of depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis comprising the administration of a therapeutically effective amount of a compound or pharmaceutically acceptable solvate of Formula I, to a patient in need thereof. Preferably the patient is a warm-blooded animal, more preferably a human.

The disclosure further provides methods of treatment for gynecological disorders and infertility. In particular, the disclosure provides methods to lower and/or suppress the LH-surge in assisted conception comprising the administration of a therapeutically effective amount of a compound or pharmaceutically acceptable solvate of Formula I, to a patient in need thereof. Preferably the patient is a warm-blooded animal, more preferably a woman.

The disclosure further provides methods to affect androgen production to cause male castration and to inhibit the sex drive in male sexual offenders comprising the administration of a therapeutically effective amount of a compound or pharmaceutically acceptable solvate of Formula I, to a patient in need thereof. Preferably the patient is a warm-blooded animal, more preferably a man.

In another aspect, the present disclosure provides a medicament comprising a compound as disclosed above.

In another aspect, the present disclosure provides a compound as disclosed above for use in treating and/or preventing depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea,

dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian tumor or virilizing adrenal tumor), menorrhagia, or adenomyosis.

The disclosure also provides the use of a compound of Formula I or a pharmaceutically acceptable solvate thereof as a medicament. Preferably, the medicament is used for the treatment and/or prevention of depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-


producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis. Preferably, the medicament is used for the treatment and/or prevention of depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis. The medicament may also be used for the treatment of gynecologic disorders, infertility and to affect androgen production to cause male castration.

In another aspect, the present disclosure provides a compound as disclosed above for use in treating and/or preventing hot flashes.

In another aspect, the present disclosure provides a process of manufacturing a compound as disclosed above, characterized in that it comprises the following steps:

- a) reacting a compound of Formula (i)

9a



(i)


wherein:

PG represents a suitable protecting group such as for example DMB, PMB, Boc, allyl, diphenyl-phosphinamide or 2-trimethylsilylethanesulfonyl;

R^{4'} is **R⁴** as defined above or a reducible precursor of hydroxyethyl and consequently a further precursor of methoxyethyl;

*-- stands for the (*R*)-enantiomer or for the racemate;

with a compound of Formula (ii)



(ii)


wherein:

R^{5'} is **R⁵** as defined above, H or 1-((tert-butyldiphenylsilyl)oxy)ethyl;

X¹ and **X²** are as defined above;


== represents a single or a double bond depending on **X¹** and **X²**;

so as to obtain a compound of Formula (iii)




wherein **PG**, **R^{4'}**, **R^{5'}**, **X¹** and **X²** are as defined above, *--- stands for the (*R*)-enantiomer or for the racemate and == represents a single or a double bond depending on **X¹** and **X²**;

b) deprotecting compound of Formula (iii) with a suitable deprotection agent to afford compound of Formula (iv)




wherein **R^{4'}**, **R^{5'}**, **X¹** and **X²** are as defined above, *--- stands for the (*R*)-enantiomer or for the racemate and == represents a single or a double bond depending on **X¹** and **X²**;

c) when **R^{5'}** is H, introducing a trifluoromethyl or difluoromethyl group by direct C-H trifluoro- or difluoromethylation, leading to compound of Formula (v)




wherein $\mathbf{R}^{4'}$, $\mathbf{X}^1$ and $\mathbf{X}^2$ are as defined above and $\mathbf{R}^5$ is trifluoromethyl or difluoromethyl, *--- stands for the (*R*)-enantiomer or for the racemate and == represents a single or a double bond depending on $\mathbf{X}^1$ and $\mathbf{X}^2$;

d) N-acylating compound of Formula (iv) wherein $\mathbf{R}^{5'}$ is not H or compound of Formula (v), with a compound of Formula (vi)



wherein $\mathbf{R}^1$, $\mathbf{R}^{1'}$, $\mathbf{R}^2$, $\mathbf{R}^{2'}$ and $\mathbf{R}^3$ are as defined above;

leading to compound of Formula (vii)



wherein R^1 , $R^{1'}$, R^2 , $R^{2'}$, R^3 , $R^{4'}$, X^1 and X^2 are as defined above,

*--- stands for the (*R*)-enantiomer or for the racemate,

== represents a single or a double bond depending on X^1 and X^2 ; and

$R^{5''}$ is R^5 as defined above or 1-((tert-butyldiphenylsilyl)oxy)ethyl;

e) optionally further conducting one or both of the two following steps e') and e''):

e') when $R^{4'}$ is a reducible precursor of hydroxyethyl and consequently a further precursor of methoxyethyl, a step of reduction optionally followed by methyl ether formation;

e'') when $R^{5''}$ is 1-((tert-butyldiphenylsilyl)oxy)ethyl, a step of alcohol deprotection and subsequent fluorination to form 1-fluoroethyl R^5 group; or a step of alcohol deprotection, followed by an oxidation step and a subsequent fluorination step to afford 1,1-difluoroethyl R^5 group;

to afford compound of Formula I as disclosed above.

In another aspect, the present disclosure provides a compound as disclosed above for use as a lowering agent of the circulating LH levels.

In another aspect, there is provided a use of a compound as disclosed above for the manufacture of a medicament to treat and/or prevent depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen


concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian tumor or virilizing adrenal tumor), menorrhagia, or adenomyosis.

In another aspect, there is provided a use of a compound as disclosed above for the manufacture of a medicament to treat and/or prevent hot flashes.

In another aspect, there is provided a use of a compound as disclosed above for the manufacture of a medicament to lower the circulating LH levels.

DETAILED DESCRIPTION

As noted above, the invention relates to compounds of Formula I:



and pharmaceutically acceptable solvates thereof, wherein:

R¹ is H, F or methyl;

R^{1'} is H;

5 **R²** is H, F, Cl or methoxy;

R^{2'} is H or F;

R³ is H, F, Cl, methyl, trifluoromethyl, nitrile or **R³** is thiophen-2-yl under the condition that **R⁵** is not methyl;

10 **R⁴** is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

R⁵ is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably **R⁵** is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

15 **X¹** is N and **X²** is S or O; or **X¹** is S and **X²** is N;

--- represents a single or a double bond depending on **X¹** and **X²**;

*-- stands for the (*R*)-enantiomer or for the racemate of compound of Formula I.


In one specific embodiment of the invention, **R⁵** is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl. In another specific embodiment, **R⁵** is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl. In another specific embodiment, **R⁵** is 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl.

Preferred compounds of Formula I and pharmaceutically acceptable solvates thereof are those wherein:

- R^1 is H, F or methyl;
- $R^{1'}$ is H;
- 5 R^2 is H, F, Cl or methoxy;
- $R^{2'}$ is H or F;
- R^3 is H, F, Cl, methyl, trifluoromethyl, nitrile or R^3 is thiophen-2-yl under the condition that R^5 is not methyl;
- R^4 is methyl, ethyl, *n*-propyl or hydroxyethyl;
- 10 R^5 is methyl, ethyl, trifluoromethyl, difluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably R^5 is methyl, ethyl or trifluoromethyl;
- X^1 is N and X^2 is S or O, preferably X^1 is N and X^2 is S.


In an embodiment of the invention, compound of Formula I is the (*R*)-enantiomer. In
15 another embodiment, compound of Formula I is the racemate.

In one embodiment, preferred compounds of Formula I are those of Formula I':




and pharmaceutically acceptable solvates thereof, wherein R^1 , $R^{1'}$, R^2 , $R^{2'}$, R^3 ,
20 R^4 , R^5 , X^1 and X^2 are as defined in Formula I and == represents a single or a double bond depending on X^1 and X^2 .

In one embodiment, preferred compounds of Formula I are those of Formula I'':



and pharmaceutically acceptable solvates thereof, wherein **R¹**, **R^{1'}**, **R²**, **R^{2'}**, **R³**, **R⁴**, **R⁵**, **X¹** and **X²** are as defined in Formula I and == represents a single or a double bond depending on **X¹** and **X²**.

In one embodiment, preferred compounds of Formula I are those of Formula Ia:



and pharmaceutically acceptable solvates thereof, wherein:

R¹ is H, F or methyl;

R^{1'} is H;

R² is H, F, Cl or methoxy;

R^{2'} is H or F;


R³ is H, F, Cl, methyl, trifluoromethyl or nitrile;

R⁴ is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably **R⁴** is methyl, ethyl, *n*-propyl or hydroxyethyl;

R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably R^5 is methyl, ethyl, trifluoromethyl or difluoromethyl, preferably R^5 is methyl, ethyl or trifluoromethyl;

*-- stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ia.

In one embodiment, preferred compounds of Formula Ia are those of Formula Ia' and Formula Ia'':




and pharmaceutically acceptable solvates thereof, wherein $R^1, R^{1'}, R^2, R^{2'}, R^3, R^4$ and R^5 are as defined in Formula Ia.

Preferred compounds of Formula Ia' and Ia'' and pharmaceutically acceptable solvates thereof are those wherein:

- R^1 is H, F or methyl;
- $R^{1'}$ is H;
- R^2 is H, F, Cl or methoxy;
- $R^{2'}$ is H or F;
- R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;
- R^4 is methyl, ethyl, *n*-propyl or hydroxyethyl;
- R^5 is methyl, ethyl, trifluoromethyl or difluoromethyl, preferably R^5 is methyl, ethyl or trifluoromethyl.

In one embodiment, preferred compounds of Formula Ia are those of Formula Ia-1:



and pharmaceutically acceptable solvates thereof, wherein:

R³ is H, F, Cl, methyl, trifluoromethyl or nitrile, preferably **R³** is H, F or Cl;


5 **R⁴** is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably **R⁴** is methyl, ethyl, *n*-propyl or hydroxyethyl;

R⁵ is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably

10 **R⁵** is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably **R⁵** is methyl, ethyl, trifluoromethyl or difluoromethyl, preferably **R⁵** is methyl, ethyl or trifluoromethyl;

*-- stands for the (*R*)-enantiomer or for the racemate.


In one embodiment, preferred compounds of Formula Ia-1 are those of Formula Ia-1':



15

and pharmaceutically acceptable solvates thereof, wherein **R³**, **R⁴** and **R⁵** are as defined in Formula Ia-1.

In one embodiment, preferred compounds of Formula Ia are those of Formula Ia-2:



and pharmaceutically acceptable solvates thereof, wherein:

R^1 is H, F or methyl;

5 $R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;

$R^{2'}$ is H or F;


R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

10 R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably

R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably R^5 is methyl, ethyl, trifluoromethyl or difluoromethyl, preferably R^5 is methyl, ethyl or trifluoromethyl;


* -- stands for the (*R*)-enantiomer or for the racemate.

15 In one embodiment, preferred compounds of Formula Ia-2 are those of Formula Ia-2':



and pharmaceutically acceptable solvates thereof, wherein **R¹**, **R^{1'}**, **R²**, **R^{2'}**, **R³** and **R⁵** are as defined in Formula Ia-2.

In one embodiment, preferred compounds of Formula Ia are those of Formula Ia-3:



5 and pharmaceutically acceptable solvates thereof, wherein:

R¹ is H, F or methyl;

R^{1'} is H;

R² is H, F, Cl or methoxy;


R^{2'} is H or F;

10 **R³** is H, F, Cl, methyl, trifluoromethyl or nitrile;

R⁴ is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably **R⁴** is methyl, ethyl, *n*-propyl or hydroxyethyl;


*-- stands for the (*R*)-enantiomer or for the racemate.

15 In one embodiment, preferred compounds of Formula Ia-3 are those of Formula Ia-3':



and pharmaceutically acceptable solvates thereof, wherein $R^1, R^1', R^2, R^2', R^3$ and R^4 are as defined in Formula Ia-3.

In one embodiment, preferred compounds of Formula I are those of Formula Ib:




5 and pharmaceutically acceptable solvates thereof, wherein:

R^3 is F or R^3 is thiophen-2-yl under the condition that R^5 is not methyl;

R^5 is methyl, ethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably R^5 is methyl, ethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably R^5 is methyl, ethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably R^5 is methyl or ethyl;


*--stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ib.

In one embodiment, preferred compounds of Formula Ib are those of Formula Ib':




15 and pharmaceutically acceptable solvates thereof, wherein R^3 and R^5 are defined as in Formula Ib.

In one embodiment, preferred compounds of Formula Ib are those of Formula Ib'':




and pharmaceutically acceptable solvates thereof, wherein **R³** and **R⁵** are defined as in Formula Ib.

5 In one embodiment, preferred compounds of Formula Ib are those of Formula Ib-1:




and pharmaceutically acceptable solvates thereof, wherein **R⁵** is methyl, ethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably **R⁵** is methyl or ethyl.

In one embodiment, preferred compounds of Formula Ib are those of Formula Ib-2:



and pharmaceutically acceptable solvates thereof, wherein R^5 is ethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably R^5 is ethyl.

5 In one embodiment, preferred compounds of Formula I are those of Formula Ic:



and pharmaceutically acceptable solvates thereof, wherein:

R^1 is H, F or methyl;

$R^{1'}$ is H;

10 R^2 is H, F, Cl or methoxy;

$R^{2'}$ is H or F;


R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^4 is methyl, ethyl, *n*-propyl or hydroxyethyl;

R^5 is methyl, ethyl or trifluoromethyl;

15 *-- stands for the (*R*)-enantiomer or for the racemate.

In one embodiment, preferred compounds of Formula Ic are those of Formula Ic':



and pharmaceutically acceptable solvates thereof, wherein:

R¹ is H, F or methyl, preferably **R¹** is H;

5 **R^{1'}** is H;

R² is H, F, Cl or methoxy, preferably **R²** is H;

R^{2'} is H or F, preferably **R^{2'}** is H;

R³ is H, F, Cl, methyl, trifluoromethyl or nitrile, preferably **R³** is F;





R⁴ is methyl, ethyl, *n*-propyl or hydroxyethyl, preferably **R⁴** is methyl;





10 **R⁵** is methyl, ethyl or trifluoromethyl, preferably **R⁵** is methyl.





Particularly preferred compounds of Formula I of the invention are those listed in Table 1 hereafter.





TABLE 1





Cpd n°	Structure	Chemical name	MW
1		(<i>R</i>)-(3,4-dichlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	409.29





2		(R)-(3-(3-ethyl-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	372.42
3		(R)-(4-chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	374.85
4		(R)-(4-chloro-3-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	392.84
5		(R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	358.39





6		(R)-(3-chloro-4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	392.84
7		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(3,4,5-trifluorophenyl)methanone	394.37
8		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4-trifluorophenyl)methanone	394.37
9		(R)-(3,4-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	376.38




10		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4,5-tetrafluorophenyl)methanone	412.36
11		(R)-(4-fluorophenyl)(8-(2-hydroxyethyl)-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	388.42
12		(4-fluorophenyl)(8-(2-hydroxyethyl)-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	388.42
13		(R)-(3-(3-ethyl-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	356.35





14		(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	358.39
15		(R)-(3-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	358.39
16		(R)-(3-chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	374.85
17		(R)-(3,5-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	376.38





18		(R)-(2,4-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	376.38
19		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(p-tolyl)methanone	354.43
20		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(phenyl)methanone	340.4
21		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-(trifluoromethyl)phenyl)methanone	408.4





22		(R)-(8-ethyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	372.42
23		(8-ethyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	372.42
24		(R)-(4-fluorophenyl)(3-(3-methyl-1,2,4-thiadiazol-5-yl)-8-propyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	386.45
25		(R)-(4-fluoro-3-methoxyphenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	388.42


26		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(o-tolyl)methanone	354.43
27		(R)-(3-methoxyphenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	370.43
28		(R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	342.33
29		(R)-4-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6,7,8-tetrahydro-[1,2,4]triazolo[4,3-a]pyrazine-7-carbonyl)benzonitrile	365.41

30		(R)-3-(3-ethyl-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl(4-(thiophen-2-yl)phenyl)methanone	420.49
31		(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4,5-tetrafluorophenyl)methanone	412.36
32		(3,4-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	376.38

33		(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4-trifluorophenyl)methanone	394.37
34		(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(3,4,5-trifluorophenyl)methanone	394.37
35		(3-chloro-4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	392.84
36		(4-chloro-3-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	392.84

37		(4-chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	374.85
38		(3,4-dichlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	409.29
39		(3-(3-ethyl-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	372.42
40		(3-(3-ethyl-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	356.35

41		(R)-(4-fluorophenyl)(8-methyl-3-(3-(trifluoromethyl)-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	412.36
42		(R)-(3-(3-(difluoromethyl)-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	394.37
43		(R)-(3-(3-(1,1-difluoroethyl)-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	392.34
44		(R)-(4-fluorophenyl)(8-methyl-3-(3-(2,2,2-trifluoroethyl)-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone	410.33

45		((8R)-3-(3-(1-fluoroethyl)-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone	374.34
----	---	--	--------


and pharmaceutically acceptable solvates thereof.

In Table 1, the term “Cpd” means compound.

The compounds of Table 1 were named using ChemBioDraw[®] Ultra version 12.0 (PerkinElmer).

- The compounds of Formula I can be prepared by different ways with reactions known to a person skilled in the art.

The invention further relates to a process of manufacturing of compounds of Formula I:



and pharmaceutically acceptable solvates thereof, wherein:

- R^1 is H, F or methyl;
 $R^{1'}$ is H;
 R^2 is H, F, Cl or methoxy;
 $R^{2'}$ is H or F;

R^3 is H, F, Cl, methyl, trifluoromethyl, nitrile or R^3 is thiophen-2-yl under the condition that R^5 is not methyl;

R^4 is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably R^4 is methyl, ethyl, *n*-propyl or hydroxyethyl;

R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl, preferably R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl, preferably R^5 is methyl, ethyl or trifluoromethyl;


X^1 is N and X^2 is S or O; or X^1 is S and X^2 is N, preferably X^1 is N and X^2 is S or O, more preferably, X^1 is N and X^2 is S;

== represents a single or a double bond depending on X^1 and X^2 ;

*-- stands for the (*R*)-enantiomer or for the racemate of compound of Formula I;

characterized in that it comprises the following steps:

a) reacting a compound of Formula (i)



(i)

wherein:


PG represents a suitable protecting group such as for example DMB, PMB, Boc, allyl, diphenyl-phosphinamide (DPP), 2-trimethylsilylethanesulfonyl (SES), preferably **PG** is DMB;

$R^{4'}$ is R^4 as defined above or a reducible precursor of hydroxyethyl and consequently a further precursor of methoxyethyl, such as for example $-CH_2CO_2Alkyl$; where the term “reducible precursor of hydroxyethyl or consequently a further precursor of methoxyethyl” refers to any chemical group which, when reacting with reducing agents, such as for example

LiAlH₄, is reduced to hydroxyethyl and then optionally further converted to methoxyethyl;

*--- stands for the (*R*)-enantiomer or for the racemate;

with a compound of Formula (ii)



(ii)


wherein:

R^{5'} is **R⁵** as defined above, H or 1-((tert-butyldiphenylsilyl)oxy)ethyl, preferably **R^{5'}** is **R⁵** as defined above or H;

X¹ and **X²** are as defined above; and

== represents a single or a double bond depending on **X¹** and **X²**;

so as to obtain a compound of Formula (iii)




(iii)

wherein **PG**, **R^{4'}**, **R^{5'}**, **X¹** and **X²** are as defined above, *--- stands for the (*R*)-enantiomer or for the racemate and == represents a single or a double


bound depending on **X¹** and **X²**;

b) deprotecting compound of Formula (iii) with a suitable deprotection agent to afford compound of Formula (iv)



wherein $\mathbf{R}^{4'}$, $\mathbf{R}^{5'}$, $\mathbf{X}^1$ and $\mathbf{X}^2$ are as defined above, *--- stands for the (*R*)-enantiomer or for the racemate and === represents a single or a double bond depending on $\mathbf{X}^1$ and $\mathbf{X}^2$;


- 5 c) when $\mathbf{R}^{5'}$ is H, introducing a trifluoromethyl or difluoromethyl group by direct C-H trifluoro- or difluoromethylation, leading to compound of Formula (v)



wherein $\mathbf{R}^{4'}$, $\mathbf{X}^1$ and $\mathbf{X}^2$ are as defined above and $\mathbf{R}^5$ is trifluoromethyl or difluoromethyl, *--- stands for the (*R*)-enantiomer or for the racemate and === represents a single or a double bond depending on $\mathbf{X}^1$ and $\mathbf{X}^2$;

10


- d) N-acylating compound of Formula (iv) wherein $\mathbf{R}^{5'}$ is not H or compound of Formula (v), with a compound of Formula (vi)



(vi)

wherein $\mathbf{R}^1$, $\mathbf{R}^{1'}$, $\mathbf{R}^2$, $\mathbf{R}^{2'}$ and $\mathbf{R}^3$ are as defined above;

leading to compound of Formula (vii)



- 5 wherein $\mathbf{R}^1$, $\mathbf{R}^{1'}$, $\mathbf{R}^2$, $\mathbf{R}^{2'}$, $\mathbf{R}^3$, $\mathbf{R}^{4'}$, $\mathbf{X}^1$ and $\mathbf{X}^2$ are as defined above,
 *--- stands for the (*R*)-enantiomer or for the racemate,
 === represents a single or a double bond depending on $\mathbf{X}^1$ and $\mathbf{X}^2$; and
 $\mathbf{R}^{5''}$ is $\mathbf{R}^5$ as defined in Formula I or 1-((tert-butyldiphenylsilyl)oxy)ethyl;

e) optionally further conducting one or both of the two following steps:

- 10 e') when $\mathbf{R}^{4'}$ is a reducible precursor of hydroxyethyl and consequently a
 further precursor of methoxyethyl, a step of reduction optionally followed
 by methyl ether formation;
 e'') when $\mathbf{R}^{5''}$ is 1-((tert-butyldiphenylsilyl)oxy)ethyl, a step of alcohol
 deprotection and subsequent fluorination to form 1-fluoroethyl $\mathbf{R}^5$ group; or
 15 a step of alcohol deprotection, followed by an oxidation step and a
 subsequent fluorination step to afford 1,1-difluoroethyl $\mathbf{R}^5$ group;

to afford compound of Formula I.

In a preferred embodiment, the protecting group PG used in the process of the invention is DMB.

According to one embodiment, the introduction of a trifluoromethyl or difluoromethyl group at step c) may be performed by direct C-H trifluoro- or difluoromethylation as described by Ji Y. et al. in PNAS, 2011, 108(35), 14411-14415 or by Fujiwara Y. et al. in JACS, 2012, 134, 1494-1497.

According to one embodiment, the fluorination step to form 1-fluoroethyl or 1,1-difluoroethyl R^5 groups at step e”) may be performed by DAST fluorination. DAST fluorination may be performed as described in WO2004/103953, page 51.

10 Reaction schemes as described in the example section are illustrative only and should not be construed as limiting the invention in any way. According to one embodiment, compounds of Formula I can be prepared using the chiral synthesis of the invention detailed in the examples below.

The invention is further directed to the use of the compounds of the invention or
15 pharmaceutically acceptable solvates thereof as antagonists to the NK-3 receptor.

Accordingly, in a particularly preferred embodiment, the invention relates to the use of compounds of Formula I and subformulae in particular those of Table 1 above, or pharmaceutically acceptable solvates thereof, as NK-3 receptor antagonists.

Accordingly, in another aspect, the invention relates to the use of these compounds or
20 solvates thereof for the synthesis of pharmaceutical active ingredients, such as selective NK-3 receptor antagonists.

USES

The compounds of the invention are therefore useful as medicaments, in particular in
25 the prevention and/or treatment of depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion,

obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH),
5 prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS),
10 premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-
15 producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis. The compounds of the invention are therefore useful as medicaments, in particular in the prevention and/or treatment of depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion,
20 obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign
25 prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease
30 (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells

in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis.

- 5 The invention also provides for a method for delaying in patient the onset of depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia,
- 10 airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern
- 15 baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma),
- 20 other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis comprising the administration of a pharmaceutically effective amount of a compound of Formula I or pharmaceutically acceptable solvate
- 25 thereof to a patient in need thereof. The invention also provides for a method for delaying in patient the onset of depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory
- 30 bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness,

bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis comprising the administration of a pharmaceutically effective amount of a compound of Formula I or pharmaceutically acceptable solvate thereof to a patient in need thereof.

Preferably, the patient is a warm-blooded animal, more preferably a human.

The compounds of the invention are especially useful in the treatment and/or prevention of sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis. The compounds of the invention are especially useful in the treatment and/or prevention of sex hormone-dependent diseases including but not limited to

benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis.

In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of benign prostatic hyperplasia (BPH), endometriosis, uterine fibrosis, uterine fibroid tumor, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis. In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of benign prostatic hyperplasia (BPH), endometriosis, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis.

In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of endometriosis, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, polycystic ovary syndrome (PCOS) and benign prostatic hyperplasia (BPH).

- 5 In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of endometriosis.

In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of uterine fibrosis.

- 10 In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of uterine fibroid tumor.

In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of uterine leiomyoma.

In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of polycystic ovary syndrome (PCOS).

- 15 In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of benign prostatic hyperplasia (BPH).

In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of hot flashes also known as hot flushes.

- 20 In a specific embodiment, the compounds of the invention are especially useful in the treatment and/or prevention of peri-menopausal conditions (i.e. 'hot flashes'), in vitro fertilization ('IVF'), male contraceptive, female contraceptive, castration of sex offenders.

- 25 The compounds of the invention are also useful in the treatment of gynecological disorders and infertility. In particular, the invention provides methods to lower and/or suppress the LH-surge in assisted conception.

The compounds of the invention are also useful to cause male castration and to inhibit the sex drive in men. This is of particular interest in the treatment of male sexual offenders.

The invention further provides the use of a compound of Formula I or a
5 pharmaceutically acceptable solvate thereof for the manufacture of a medicament for treating and/or preventing depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory
10 bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer,
15 ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with
20 hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis in a patient. The invention further provides the use of a compound of Formula I or a
25 pharmaceutically acceptable solvate thereof for the manufacture of a medicament for treating and/or preventing depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory
30 bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness,

bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis in a patient.

Preferably, the patient is a warm-blooded animal, more preferably a human.

The invention especially provides the use of a compound of Formula I or a pharmaceutically acceptable solvate thereof for the manufacture of a medicament to treat and/or prevent sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis. The invention especially provides the use of a compound of Formula I or a pharmaceutically acceptable solvate thereof for the manufacture of a medicament to treat and/or prevent sex hormone-dependent diseases

including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian or adrenal tumor), menorrhagia and adenomyosis.

In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent endometriosis, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, polycystic ovary syndrome (PCOS) and benign prostatic hyperplasia (BPH).

In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent endometriosis.

In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent uterine fibrosis.

In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent uterine fibroid tumor.

In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent uterine leiomyoma.

In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent polycystic ovary syndrome (PCOS).

5 In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent benign prostatic hyperplasia (BPH).

In a specific embodiment, compounds of Formula I or a pharmaceutically acceptable solvate thereof may be used for the manufacture of a medicament to treat and/or prevent hot flashes also known as hot flushes.

10 The invention further provides the use of a compound of Formula I or a pharmaceutically acceptable solvate thereof for the manufacture of a medicament to lower and/or suppress the LH-surge in assisted conception in a patient. Preferably the patient is a warm-blooded animal, more preferably a woman.

15 The invention further provides the use of a compound of Formula I or a pharmaceutically acceptable solvate thereof for the manufacture of a medicament to cause male castration and to inhibit the sex drive in men. This is of particular interest in the treatment of male sexual offenders.

20 According to a further feature of the present invention there is provided a method for modulating NK-3 receptor activity, in a patient, preferably a warm blooded animal, and even more preferably a human, in need of such treatment, which comprises administering to said patient an effective amount of compound of the present invention, or a pharmaceutically acceptable solvate thereof.

25 According to one embodiment, the compounds of the invention, their pharmaceutical acceptable solvates may be administered as part of a combination therapy. Thus, are included within the scope of the present invention embodiments comprising coadministration of, and compositions and medicaments which contain, in addition to a compound of the present invention, a pharmaceutically acceptable solvate thereof as active ingredient, additional therapeutic agents and/or active ingredients. Such multiple

drug regimens, often referred to as “combination therapy”, may be used in the treatment and/or prevention of any of the diseases or conditions mediated by or associated with NK-3 receptor modulation. The use of such combinations of therapeutic agents is especially pertinent with respect to the treatment of the above-mentioned disorders
5 within a patient in need of treatment or one at risk of becoming such a patient.

In addition to the requirement of therapeutic efficacy, which may necessitate the use of active agents in addition to the NK-3 receptor modulator compounds of Formula I or pharmaceutical acceptable solvates thereof, there may be additional rationales which compel or highly recommend the use of combinations of drugs involving active
10 ingredients which represent adjunct therapy, i.e., which complement and supplement the function performed by the NK-3 receptor modulator compounds of the present invention. Suitable supplementary therapeutic agents used for the purpose of auxiliary treatment include drugs which, instead of directly treating or preventing a disease or condition mediated by or associated with NK-3 receptor modulation, treat diseases or
15 conditions which directly result from or indirectly accompany the basic or underlying NK-3 receptor modulated disease or condition.

According to a further feature of the present invention, the compound of Formula I, a pharmaceutically acceptable solvate thereof may be used in combination therapy with antipsychotic drugs (APD), to improve the efficacy and to minimize secondary effects
20 associated to APD including but not limited to Dopamine 2/3 and 5-HT₂ receptors antagonists. More particular the compound of Formula I, a pharmaceutically acceptable solvate thereof may be used as an adjunct therapy in combination with an atypical antipsychotic drug, including but not limited to risperidone, clozapine, olanzapine, where the NK-3 receptor modulator may serve a role as dose-limiting for the atypical
25 antipsychotic and therefore spare the patient from some of the side effect of those atypical antipsychotic drugs.

Thus, the methods of treatment and pharmaceutical compositions of the present invention may employ the compounds of Formula I or pharmaceutical acceptable solvates thereof in the form of monotherapy, but said methods and compositions may
30 also be used in the form of multiple therapy in which one or more compounds of

Formula I or their pharmaceutically acceptable solvates are coadministered in combination with one or more other therapeutic agents.

In the above-described embodiment combinations of the present invention, the compound of Formula I, a pharmaceutically acceptable solvate thereof and other
5 therapeutic active agents may be administered in terms of dosage forms either separately or in conjunction with each other, and in terms of their time of administration, either serially or simultaneously. Thus, the administration of one component agent may be prior to, concurrent with, or subsequent to the administration of the other component agent(s).

10 The invention also provides pharmaceutical compositions comprising a compound of Formula I or a pharmaceutically acceptable solvate thereof and at least one pharmaceutically acceptable carrier, diluent, excipient and/or adjuvant. As indicated above, the invention also covers pharmaceutical compositions which contain, in addition to a compound of the present invention, a pharmaceutically acceptable solvate
15 thereof as active ingredient, additional therapeutic agents and/or active ingredients.

Another object of this invention is a medicament comprising at least one compound of the invention, or a pharmaceutically acceptable solvate thereof, as active ingredient.

According to a further feature of the present invention there is provided the use of a compound of Formula I or a pharmaceutically acceptable solvate thereof for the
20 manufacture of a medicament for modulating NK-3 receptor activity in a patient, in need of such treatment, which comprises administering to said patient an effective amount of compound of the present invention, or a pharmaceutically acceptable solvate thereof.

Preferably, the patient is a warm-blooded animal, more preferably a human.

25 As set forth above, the compounds of the invention, their pharmaceutically acceptable solvates may be used in monotherapy or in combination therapy. Thus, according to one embodiment, the invention provides the use of a compound of the invention for the manufacture of a medicament for at least one of the purposes described above, wherein

said medicament is administered to a patient in need thereof, preferably a warm-blooded animal, and even more preferably a human, in combination with at least one additional therapeutic agent and/or active ingredient. The benefits and advantages of such a multiple drug regimen, possible administration regimens as well as suitable additional
5 therapeutic agents and/or active ingredients are those described above.

Generally, for pharmaceutical use, the compounds of the invention may be formulated as a pharmaceutical preparation comprising at least one compound of the invention and at least one pharmaceutically acceptable carrier, diluent, excipient and/or adjuvant, and optionally one or more further pharmaceutically active compounds.

10 By means of non-limiting examples, such a formulation may be in a form suitable for oral administration, for parenteral administration (such as by intravenous, intramuscular or subcutaneous injection or intravenous infusion), for topical administration (including ocular), for administration by inhalation, by a skin patch, by an implant, by a suppository, etc. Such suitable administration forms – which may be solid, semi-solid or
15 liquid, depending on the manner of administration – as well as methods and carriers, diluents and excipients for use in the preparation thereof, will be clear to the skilled person; reference is made to the latest edition of Remington's Pharmaceutical Sciences.

Some preferred, but non-limiting examples of such preparations include tablets, pills, powders, lozenges, sachets, cachets, elixirs, suspensions, emulsions, solutions, syrups,
20 aerosols, ointments, cremes, lotions, soft and hard gelatin capsules, suppositories, drops, sterile injectable solutions and sterile packaged powders (which are usually reconstituted prior to use) for administration as a bolus and/or for continuous administration, which may be formulated with carriers, excipients, and diluents that are suitable per se for such formulations, such as lactose, dextrose, sucrose, sorbitol,
25 mannitol, starches, gum acacia, calcium phosphate, alginates, tragacanth, gelatin, calcium silicate, microcrystalline cellulose, polyvinylpyrrolidone, polyethylene glycol, cellulose, (sterile) water, methylcellulose, methyl- and propylhydroxybenzoates, talc, magnesium stearate, edible oils, vegetable oils and mineral oils or suitable mixtures thereof. The formulations can optionally contain other substances that are commonly
30 used in pharmaceutical formulations, such as lubricating agents, wetting agents,

emulsifying and suspending agents, dispersing agents, desintegrants, bulking agents, fillers, preserving agents, sweetening agents, flavoring agents, flow regulators, release agents, etc.. The compositions may also be formulated so as to provide rapid, sustained or delayed release of the active compound(s) contained therein.

- 5 The pharmaceutical preparations of the invention are preferably in a unit dosage form, and may be suitably packaged, for example in a box, blister, vial, bottle, sachet, ampoule or in any other suitable single-dose or multi-dose holder or container (which may be properly labeled); optionally with one or more leaflets containing product information and/or instructions for use. Generally, such unit dosages will contain
- 10 between 0.05 and 1000 mg, and usually between 1 and 500 mg, preferably between 2 and 150 mg of at least one compound of the invention, e.g. about 2, 4, 8, 16, 32, 64 or 128 mg per unit dosage. According to another embodiment, such unit dosages will contain between 0.05 and 1000 mg, and usually between 1 and 500 mg, preferably between 2 and 400 mg, preferably between 2 and 200 mg of at least one compound of
- 15 the invention per unit dosage.

Usually, depending on the condition to be prevented or treated and the route of administration, the active compound of the invention will usually be administered between 0.001 and 10 mg per kilogram body weight, more often between 0.01 and 4 mg per kilogram body weight, preferably between 0.02 and 1.5 mg per kilogram body

20 weight, for example about 0.02, 0.04, 0.08, 0.16, 0.32, 0.64 or 1.28 mg, per kilogram body weight of the patient per day, which may be administered as a single daily dose, divided over one or more daily doses, or essentially continuously, e.g. using a drip infusion. According to another embodiment, the active compound of the invention will usually be administered between 0.001 and 10 mg per kilogram body weight, more

25 often between 0.01 and 7 mg per kilogram body weight, preferably between 0.03 and 3.5 mg per kilogram body weight of the patient per day, which may be administered as a single daily dose, divided over one or more daily doses, or essentially continuously, e.g. using a drip infusion.


According to one embodiment, the active compound of the invention will be administered as a single daily dose, divided over one, two or more daily doses, or essentially continuously, e.g. using a drip infusion.

5 DEFINITIONS

The definitions and explanations below are for the terms as used throughout the entire application, including both the specification and the claims.


When describing the compounds of the invention, the terms used are to be construed in accordance with the following definitions, unless indicated otherwise.

- 10 The term “alkyl” refers to a hydrocarbyl radical of formula C_nH_{2n+1} wherein n is a number greater than or equal to 1. Generally, alkyl groups of this invention comprise from 1 to 4 carbon atoms, preferably from 1 to 3 carbon atoms. Alkyl groups may be linear or branched. Suitable alkyl groups include but are not limited to methyl, ethyl, *n*-propyl, *i*-propyl, *n*-butyl, *i*-butyl, *s*-butyl and *t*-butyl.
- 15 The term “thiophen-2-yl” as used herein means a group of formula



wherein the arrow defines the attachment point.

The ring atoms of (3-substituted)-(8-substituted)-5,6-dihydro-[1,2,4]triazolo[4,3-*a*]pyrazines of the invention are numbered based on scheme below.



- 20 Bonds from an asymmetric carbon in compounds are generally depicted using a solid line (—), a solid wedge (▴), or a dotted wedge (▴). The use of either a


solid or dotted wedge to depict bonds from an asymmetric carbon atom is meant to indicate that only the stereoisomer shown is meant to be included.

The compounds of Formula I and subformulae thereof contain a stereogenic carbon center at position 8 and thus may exist as (*R*)- and (*S*)-enantiomers. In an embodiment of the invention, compounds of Formula I are not pure (*S*)-enantiomers relative to the C8 position.


In the compounds of the invention, a dotted wedge (⋯⋯⋯) carrying a substituent at the C8 position is used to depict the (*R*)-enantiomer, thus excluding racemic mixtures thereof.

10 In the compounds of the invention, a dotted line with a star next to the C8 position (*---) is used to depict either a dotted wedge (⋯⋯⋯) to represent the (*R*)-enantiomer or a solid line (——) to depict the racemic mixture of (*R*)- and (*S*)-enantiomer, which is called “racemate”.

For instance, (*R*)-(3,4-dichlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-*a*]pyrazin-7(8H)-yl)methanone (compound n°1) is depicted as:



The racemic mixture of this compound, (3,4-dichlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-*a*]pyrazin-7(8H)-yl)methanone
20 (compound n°38) is depicted as:



The term “solvate” is used herein to describe a compound in this invention that contains stoichiometric or sub-stoichiometric amounts of one or more pharmaceutically acceptable solvent molecule such as ethanol. The term “hydrate” refers to when the said
5 solvent is water.

All references to compounds of Formula I include references to solvates, multi-component complexes and liquid crystals thereof.

The compounds of the invention include compounds of Formula I as hereinbefore defined, including all polymorphs and crystal habits thereof, prodrugs and prodrugs
10 thereof and isotopically- labeled compounds of Formula I.

The invention also generally covers all pharmaceutically acceptable prodrugs and prodrugs of the compounds of Formula I.

The term “prodrug” as used herein means the pharmacologically acceptable derivatives of compounds of Formula I, such as for example esters, whose *in vivo* biotransformation
15 product generates the biologically active drug. Prodrugs are generally characterized by increased bio-availability and are readily metabolized into biologically active compounds *in vivo*.

The term “prodrug”, as used herein, means any compound that will be modified to form a drug species, wherein the modification may take place either inside or outside of the
20 body, and either before or after the prodrug reaches the area of the body where administration of the drug is indicated.

The term "patient" refers to a warm-blooded animal, more preferably a human, who/which is awaiting the receipt of, or is receiving medical care or is/will be the object of a medical procedure.

5 The term "human" refers to a subject of both genders and at any stage of development (i.e. neonate, infant, juvenile, adolescent, adult).

The terms "treat", "treating" and "treatment, as used herein, are meant to include alleviating, attenuating or abrogating a condition or disease and/or its attendant symptoms.

10 The terms "prevent", "preventing" and "prevention", as used herein, refer to a method of delaying or precluding the onset of a condition or disease and/or its attendant symptoms, barring a patient from acquiring a condition or disease, or reducing a patient's risk of acquiring a condition or disease.

15 The term "therapeutically effective amount" (or more simply an "effective amount") as used herein means the amount of active agent or active ingredient (e.g. NK-3 antagonist) that is sufficient to achieve the desired therapeutic or prophylactic effect in the patient to which/whom it is administered.

20 The term "administration", or a variant thereof (e.g. "administering"), means providing the active agent or active ingredient (e.g. a NK-3 antagonist), alone or as part of a pharmaceutically acceptable composition, to the patient in whom/which the condition, symptom, or disease is to be treated or prevented.

By "pharmaceutically acceptable" is meant that the ingredients of a pharmaceutical composition are compatible with each other and not deleterious to the patient thereof.

25 The term "antagonist" as used herein means a compound that competitively or non-competitively binds to a receptor at the same site as an agonist (for example, the endogenous ligand) and has reversible and competitive binding affinity to a receptor without direct modulation of receptor signaling, but that nonetheless occupies the

binding site of an agonist (for example, the endogenous ligand) to thereby block agonist-mediated receptor signaling.

The term “sex hormone-dependent disease” as used herein means a disease which is exacerbated by, or caused by, excessive, inappropriate or unregulated sex hormone production and/or an extraordinary physiological response to sex hormones. Examples of such diseases in men include but are not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, androgen dependent acne, male pattern baldness and precocious puberty in boys. Examples of such diseases in women include but are not limited to endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers (ovarian cancer, breast cancer), androgen-producing tumor (virilizing ovarian or adrenal tumor), hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), menorrhagia and adenomyosis (abnormal endometrial growth within the muscle of the uterus).

The term “Psychotic disorders” as used herein means a group of illnesses that affect the mind. These illnesses alter a patient's ability to think clearly, make good judgments, respond emotionally, communicate effectively, understand reality, and behave appropriately. When symptoms are severe, patient with psychotic disorders have difficulty staying in touch with reality and are often unable to meet the ordinary demands of daily life. Psychotic disorders include but are not limited to, schizophrenia, schizophreniform disorder, schizo-affective disorder, delusional disorder, brief psychotic disorder, shared psychotic disorder, psychotic disorder due to a general medical condition, substance-induced psychotic disorder or psychotic disorders not otherwise specified (Diagnostic and Statistical Manual of Mental Disorders, Ed. 4th, American Psychiatric Association, Washington, D.C. 1994).

The term “pharmaceutical vehicle” as used herein means a carrier or inert medium used as solvent or diluent in which the pharmaceutically active agent is formulated and/or administered. Non-limiting examples of pharmaceutical vehicles include creams, gels, lotions, solutions, and liposomes.

- 5 The present invention will be better understood with reference to the following examples. These examples are intended to be representative of specific embodiments of the invention, and are not intended as limiting the scope of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

- 10 **Figure 1** is a graph showing the plasma testosterone levels over time in intact male rats after oral administration of compound n°5 (3 mg/kg) or of a vehicle (0.5% methyl cellulose).

Figure 2 is a histogram showing the prostate weight in a rat model of Benign Prostate Hyperplasia (BPH) after oral administration of 3, 10 or 30 mg/kg of compound n°5.

- 15 **Figure 3** is a histogram showing the estradiol levels in adult, female rats tracked over the duration of consecutive estrous cycles, after oral administration of compound n°5 (10 mg/kg) or of a vehicle (0.5% methyl cellulose).

EXAMPLES

20 CHEMISTRY EXAMPLES

All reported temperatures are expressed in degrees Celsius (°C); all reactions were carried out at room temperature (RT) unless otherwise stated.

- All reactions were followed by thin layer chromatography (TLC) analysis (TLC plates, silica gel 60 F₂₅₄, Merck) was used to monitor reactions, establish silica-gel flash chromatography conditions. All other TLC developing agents/visualization techniques, experimental set-up or purification procedures that were used in this invention, when
- 25

not described in specific details, are assumed to be known to those conversant in the art and are described in such standard reference manuals as: i) Gordon, A. J.; Ford, R. A. "The Chemist's Companion – A Handbook of Practical Data, Techniques, and References", Wiley: New York, 1972; ii) Vogel's Textbook of Practical Organic
5 Chemistry, Pearson Prentice Hall: London, 1989.

HPLC-MS spectra were typically obtained on an Agilent LCMS using electrospray ionization (ESI). The Agilent instrument includes an autosampler 1100, a binary pump 1100, an ultraviolet multi-wavelength detector 1100 and a 6100 single-quad mass-spectrometer. The chromatography column used was Sunfire 3.5 μm , C18, 3.0 x 50 mm
10 in dimensions.

Eluent typically used was a mixture of solution A (0.1% TFA in H_2O) and solution B (0.1% TFA in MeCN).

Gradient was applied at a flow rate of 1.3 mL per minute as follows: gradient A (for analysis of final compounds and intermediates): held the initial conditions of 5% solution B for 0.2 min, increased linearly to 95% solution B in 6 min, held at 95%
15 during 1.75 min, returned to initial conditions in 0.25 min and maintained for 2.0 min; gradient B (for analysis of crude samples and reactions mixtures): held the initial conditions of 5% solution B for 0.2 min, increased linearly to 95% in 2.0 min, held at 95% during 1.75 min, returned to initial conditions in 0.25 min and maintained for
20 2 min.

Determination of chiral purity was made using chiral HPLC that was performed on an Agilent 1100 (binary pump and a ultraviolet multi wavelength detector) with manual or automatic (Autosampler 1100) injection capabilities. Column used is CHIRALPAK IA 5 μm , 4.6 x 250 mm 4.6 x 250 mm in isocratic mode. Choice of eluent was predicated
25 on the specifics of each separation. Further details concerning the chiral HPLC methods used are provided below.

Method A: column CHIRALPAK IA 5 μ m, 4.6 x 250 mm, eluent: EtOAc plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm; column at RT, eluent was used as sample solvent.

Method B: column CHIRALPAK IA 5 μ m, 4.6 x 250 mm, eluent: EtOAc/hexane
5 (50:50) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm; column at RT, eluent was used as sample solvent.

Method C: column CHIRALPAK IA 5 μ m 4.6 x 250mm, eluent: hexane/ethanol (80:20 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

10 Method D: column CHIRALPAK IA 5 μ m 4.6 x 250mm, eluent: hexane/ethanol (50:50 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

Method E: column CHIRALPAK ID 5 μ m 4.6 x 250mm, eluent: hexane/ethanol (80:20 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or
15 280 nm, column at RT, eluent was used as sample solvent.

Method F: column CHIRALPAK IA 5 μ m 4.6 x 250mm, eluent: DCM/ethanol (98:2 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

Method G: column CHIRALPAK IA 5 μ m 4.6 x 250mm, eluent: DCM/ethanol
20 (98:2 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

Method H: column CHIRALPAK IB 5 μ m 4.6 x 250mm, eluent: TBME plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

25 Method I: column CHIRALPAK IC 5 μ m 4.6 x 250mm, eluent: TBME/ethanol (98:2 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

Method J: column CHIRALPAK ID 5 μ m 4.6 x 250mm, eluent: EtOAc/DCM/IPAethanol (3:1:1 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

Method K: column CHIRALPAK IC 5 μ m 4.6 x 250mm, eluent: TBME/methanol (98:2 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

Method L: column CHIRALPAK IB 5 μ m 4.6 x 250mm, eluent: TBME/methanol (98:2 v/v) plus 0.1% of DEA, flow rate: 1.0 mL per minute; UV detection at 254 or 280 nm, column at RT, eluent was used as sample solvent.

- 10 Preparative HPLC purifications were typically carried out on an Agilent 1200 instrument (preparative pump 1200 and ultraviolet multi wavelength detector 1200) with manual injection. The chromatography column used was Waters Sunfire 5 μ m, C18, 19 x 100 mm, or XBridge 5 μ m, C18, 19 x 100mm depending on the type of eluent system employed, i.e. low pH or high pH conditions.
- 15 For high-pH HPLC purifications, eluent typically consisted of a mixture of solution A (0.04 M ammonium bicarbonate in H₂O plus 0.1% of conc. NH₄OH) and solution B was MeCN. The gradient was adapted depending on the impurity profile in each sample purified, thereby allowing sufficient separation between the impurities and the desired compound.
- 20 In rare cases when high-pH HPLC purification did not provide sufficient purity, low-pH HPLC was applied. For low-pH HPLC purifications, eluent typically consisted of a mixture of solution A (0.1% of TFA in H₂O) and solution B was MeCN. The gradient was adapted depending on the impurity profile in each sample purified, thereby allowing sufficient separation between the impurities and the desired compound. TFA
- 25 was removed from evaporated fractions by liquid-liquid extraction.

Chiral preparative HPLC purifications were performed on an Agilent 1200 instrument (preparative pump 1200 and ultraviolet multi wavelength detector 1200) with manual injection. The chiral columns used are CHIRALPAK IA 5 μ m, 20 x 250 mm or

CHIRALPAK IA 5 μ m, 10 x 250 mm. All chiral HPLC methods were employed in an isocratic mode. The eluent mixture was selected based on the analytical chiral HPLC experiment (see above) that provided the best chiral separation.

5 ^1H (300 MHz), ^{19}F (282 MHz) and ^{13}C NMR (75 MHz) spectra were recorded on a Bruker Avance DRX 300 instrument. Chemical shifts are expressed in parts per million, (ppm, δ units). Coupling constants are expressed in Hertz (Hz). Abbreviations for multiplicities observed in NMR spectra are as follows: s (singlet), d (doublet), t (triplet), q (quadruplet), m (multiplet), br (broad).

10 Solvents, reagents and starting materials were purchased and used as received from commercial vendors unless otherwise specified.

The following abbreviations are used:

Boc: *tert*-Butoxycarbonyl,
Cpd: Compound,
DAST: (Diethylamino)sulfur trifluoride,
15 DCM: Dichloromethane,
DEA: Diethylamine,
DMB: 2,4-Dimethoxybenzyl,
DMB-CHO: 2,4-Dimethoxybenzaldehyde,
DPP: Diphenylphosphoramidate,
20 ee: Enantiomeric excess,
eq.: Equivalent(s),
EtOAc: Ethyl acetate,
EtOH: Ethanol,
g: Gram(s),
25 h: Hour(s),
IPA: *iso*-Propylalcohol,
L: Liter(s),
MeOH: Methanol,
 μ L: Microliter(s),
30 mg: Milligram(s),


- mL: Milliliter(s),
mmol: Millimole(s),
min: Minute(s),
P: UV purity at 254 nm or 215 nm determined by HPLC-MS,
5 PMB: 4-Methoxybenzyl,
rt: Room temperature,
SES: 2-Trimethylsilylethanesulfonyl,
*t*Bu: *tert*-Butyl,
TBDPS: *tert*-Butyldiphenylsilyl,
10 TBME: *tert*-Butyl methyl ether,
TFA: Trifluoroacetic acid,
TLC: Thin layer chromatography.

The intermediates and compounds described below were named using ChemBioDraw[®] Ultra version 12.0 (PerkinElmer).

15 **I. Racemic synthesis**

I.1. General Synthetic Scheme for racemic synthesis

Compounds of the invention may be synthesized using the methodology described in Scheme 1, which represents the racemic product synthesis. The racemic products may then be subjected to chiral HPLC for chiral separation.



Scheme 1: General racemic synthetic scheme for the preparation of the compounds of the invention.

5 The general synthetic scheme comprises the following steps:

Step 1: DMB-protected_ketopiperazine **1.1** was converted to iminoether **1** by using the Meerwein reagent (Et_3OBF_4).

Step 2: Ester **2.2** was subsequently converted to acyl hydrazide **2**. Ester **2.2** may be obtained by esterification of acid **2.1**.

- 5 **Step 3:** Cyclodehydration between the acyl hydrazide **2** and the iminoether **1** furnished the protected triazolopiperazine **3.1**. Thereafter, **3.1** was subjected to acidolytic deprotection to obtain **3**. When applicable, R^5 was introduced from $\text{R}^{5'}$ affording **3'**.

- Step 4:** The thus obtained triazolopiperazine intermediate **3** (or **3'**) was acylated through reaction with the appropriate acid chloride **4.1** to obtain the racemic final target structure represented by the general Formula **4**. Optionally, R^4 may be transformed, for example by reduction of $\text{R}^{4'}$ when $\text{R}^{4'}$ contains a reducible group such as an ester group. The chiral compound **4'** was subsequently obtained by purification using preparative chiral HPLC.

1.2. Step 1: Protection and conversion to iminoether 1


- 15 **Method A: Conversion of DMB-protected ketopiperazine 1.1 to iminoether 1**

Method A is the procedure used for the synthesis of the iminoether intermediates **1** with a DMB protecting group and is detailed below:



Scheme 2: Conversion to iminoether **1**.

- 20 Method A is illustrated by the synthesis of intermediate **1a** wherein $\text{R}^{4'}$ is Me.

Synthesis of 1-(2,4-dimethoxybenzyl)-5-ethoxy-6-methyl-1,2,3,6-tetrahydropyrazine 1a


Scheme 3: Synthesis of 1-(2,4-dimethoxybenzyl)-5-ethoxy-6-methyl-1,2,3,6-tetrahydropyrazine **1a**.

- 5 Oven-dried (115°C) sodium carbonate (18.6 g, 98 mmol, 2.25 eq.) was placed in a 500 mL round-bottom flask. The round-bottom flask was backfilled with Ar and then capped with a rubber septum. A solution of 4-(2,4-dimethoxybenzyl)-3-methylpiperazin-2-one **1.1a** (20.6 g, 78 mmol, 1 eq.) in anhydrous DCM (250 mL) was added, followed by triethyloxonium tetrafluoroborate (18.6g, 98 mmol, 1.25 eq.) in one
- 10 portion. Thereafter, the reaction mixture was stirred further at RT for 1 h whereupon the reaction mixture was diluted with water (250 mL). The aqueous layer was extracted with DCM (3 x 150 mL). The organic layers were combined, dried over MgSO₄, filtered and concentrated under reduced pressure. The crude compound was then purified on silica gel (EtOAc) to afford the desired product **1a** as orange oil. Yield: 13.2 g, 58 %.
- 15 LCMS: P = 93 %, retention time = 1.8 min, (M+H+H₂O)⁺: 311; ¹H-NMR (CDCl₃): δ 7.23 (d, *J* = 8.8 Hz, 1H), 6.48 (d, *J* = 8.8 Hz, 1H), 6.44 (s, 1H), 4.02 (m, 2H), 3.92 (s, 3H), 3.91 (s, 3H), 3.86 (d, *J*_{AB} = 14.0 Hz, 1H), 3.46 (d, *J*_{AB} = 14.0 Hz, 1H), 3.44 (m, 2H), 3.10 (m, 1H), 2.79 (m, 1H), 2.32 (m, 1H), 1.35 (d, *J* = 6.8 Hz, 3H), 1.24 (t, *J* = 6.0 Hz, 3H).

20 **1.3. Step 2: Formation of acyl hydrazide 2**

Method B: Acyl hydrazide2

Method B is the procedure used for the synthesis of the acyl hydrazides **2** and is detailed below:




Scheme 4: Formation of acylhydrazide 2.

In a round-bottom flask equipped with a condenser, ester **2.2** (1 eq.) is dissolved in anhydrous EtOH and treated with hydrazine hydrate (1.2 to 20 eq., preferably 1.5 to 10 eq.) using a temperature range from RT to reflux. After allowing the reaction mixture to come to RT, the solution is concentrated under reduced pressure. Co-evaporations using a mixture of commercial DCM:MeOH (1:1) may be performed to remove residual water. The residue is then recrystallized and/or precipitated or purified on a pad of silica to afford **2**.

10 I.4. Step 3: Cyclodehydration leading to triazolopiperazine 3

Method C: Cyclodehydration and acyolysis

Method C is the procedure used for the synthesis of the triazolopiperazine **3** and is detailed below:



Scheme 5: Cyclodehydration leading to triazolopiperazine 3.

Step 1: In a round-bottom flask equipped with a condenser, imino-ether **1** (1 eq.) is dissolved in anhydrous MeOH, to which is added **2** (1 eq.) in one portion. The resulting solution is stirred at reflux overnight. Thereafter, the reaction mixture is brought to RT

and the volatiles are removed under reduced pressure. The crude compound is then purified using silica gel chromatography to afford the desired product **3.1**.

Step 2: In a round-bottom flask containing DCM is added **3.1** (1 eq.). Then, TFA (5 to 75 eq.), is added to the reaction mixture at RT. After 30 min stirring, the mixture is concentrated. Then DCM is added to the residue thus obtained, and washed with saturated NaHCO₃. The aqueous layer is extracted twice with DCM, the organic layers are washed with brine, dried over MgSO₄, filtered and concentrated under reduced pressure to obtain crude **3**. The crude **3** may be directly used in the next step without further purification.


10 In one embodiment, alternative work-up equally used involves treatment of the dried residue obtained above with 4 M HCl/dioxane (20 eq.) at RT under stirring. After 5 min, Et₂O is added to help precipitation. This precipitate is filtered off under vacuum, washed with Et₂O and dried under high vacuum to furnish **3** as hydrochloride salt.

In another embodiment, HCl could be used for **Step 2**: HCl 4M solution in 1,4-dioxane
15 (3 to 20 eq.) is added in one portion to a solution of **3.1** (1 eq.) in commercial *iso*-propanol or ethanol. The reaction mixture is stirred at 60°C. After complete conversion monitored by HPLC-MS (1 to 10 h), the reaction mixture is allowed to cool to room temperature and then further cooled to 0 °C with an ice bath. Thereupon, Et₂O is added. After 15-30 min stirring, the precipitate is filtered and dried *in vacuo* to afford **3** as
20 hydrochloride salt.

Remark: When $R^{5'} \neq R^5 = H$, introduction of groups such as trifluoro- or difluoromethyl through direct trifluoromethylation or direct difluoromethylation (Ji Y. et al., PNAS, 2011, 108(35), 14411-14415; Fujiwara Y. et al., JACS, 2012, 134, 1494-1497) may be performed.

I.5. Step 4: Acylation leading to final products**Method D: Acylation and chiral HPLC purification**

Method D is the procedure used for the synthesis of the racemic product **4** and its purification to obtain (*R*)-enantiomer **4'** compounds of general Formula I. Method D is detailed below:



Scheme 6: Acylation and chiral HPLC purification.

To a solution of crude **3** (1 eq.) in anhydrous DCM are added, at RT, **4.1** (1.17 to 1.3 eq.), followed by N-methylmorpholine (1 eq. to 3.5 eq.) dropwise over 15 sec. The reaction mixture is stirred at RT for 1 to 30 minutes and the milky suspension is poured into 1 M HCl solution or directly diluted with DCM. The aqueous phase is extracted with DCM. The organic phases are combined, optionally washed with 1 M NaOH, water, brine, dried over $MgSO_4$ and evaporated to dryness. The residue is solubilized in DCM and Et_2O and is slowly added to induce precipitation. The solid was filtered off, washed with Et_2O and dried under vacuum to afford **4**. Alternatively, the residue is preliminary purified on silica gel before precipitation or purified on silica gel only.

Substituent $R^{4'}$ may then be transformed, when applicable, into R^4 . One example of such transformation is illustrated by the synthesis of compound **12** wherein R^4 is hydroxyethyl group, obtained by reduction of $R^{4'} = -CH_2CO_2Alkyl$.


To a solution of **4** (1 eq.) in anhydrous THF is added, at $-40^\circ C$, LAH (1 eq.), The
5 reaction mixture is stirred at $-40^\circ C$ for 5 to 30 minutes and the mixture is quenched with 1 M NaOH solution. The resulting mixture is extracted with DCM twice. The organic phases are combined, dried over $MgSO_4$ and evaporated to dryness. The residue **4''** is then purified on silica gel.

Compound **4** or **4''** may be purified by chiral preparative HPLC according to the
10 abovementioned method to yield the corresponding chiral(*R*)-compound **4'**. Compounds **4**, **4''** and **4'** are compounds of Formula I of the invention.

II. Chiral synthesis

II.1. General Synthetic Scheme for chiral synthesis

Chiral compounds of the invention may be synthesized using the chiral process of the
15 invention described in Scheme 7.



Scheme 7: General synthetic scheme for the preparation of chiral compounds of the invention.

- 5 Chiral ketopiperazine **B** was protected with "PG" protecting group leading to PG-protected chiral ketopiperazine **C**. PG-protected chiral ketopiperazine **C** was converted to iminoether **D** by using the Meerwein reagent (Et_3OBF_4). Condensation reaction between the acyl hydrazide **E** and iminoether **D** was conducted under heating conditions in methanol to provide PG-protected piperazine **F** that was subsequently deprotected to
- 10 yield compound of Formula **G**.

In one embodiment, when the protecting group PG is DMB, the DMB group deprotection step (from **F** to **G**) is carried out using TFA in DCM at rt, followed by either TFA salt exchange with HCl or extraction at high pH recovering free piperazine **G**.


When applicable, R^5 was introduced from R^5 of **G**, affording **G'**.

Acylation of **G** or **G'** with the appropriate acid chloride **H** afforded the (*R*)-enantiomer of **I** typically in > 90% enantiomeric excess (chiral HPLC). When applicable, R^4 of **I** was then modified to afford R^4 , furnishing **I'**.

- 5 When applicable, $R^{5''}$ of **I** or **I'** was then modified to afford R^5 , furnishing **I''** or **I'''** respectively.

II.2. Step 1: Protection of ketopiperazine B

II.2.1. Protection of ketopiperazine B with an allyl to afford protected ketopiperazine C₁



10

Scheme 8: Allyl protection of B.


Allyl protection is illustrated by the synthesis of intermediate (*R*)-4-allyl-3-methylpiperazin-2-one (i.e. compound **C₁** wherein $R^{4'}$ is Me).

- To a solution of (*R*)-3-methylpiperazin-2-one (0.5 g, 4.38 mmol) in commercial anhydrous THF (44 mL) at rt was added K_2CO_3 (1.2 g, 8.76 mmol). 3-bromoprop-1-ene (0.41 mL, 4.82 mmol) was then added at once, and the reaction mixture was stirred under reflux for 14h.
- 15

- The reaction mixture was allowed to cool to rt, concentrated and the residue was then solubilized with water (10 mL) and DCM (10 mL). The organic layer was separated, dried over $MgSO_4$, filtered and concentrated to afford 460 mg of yellow oil. ^1H -NMR analysis shows that desired product was clearly the main product. Crude was used as-is in the following step.
- 20

LCMS: P > 90 %, retention time = 0.2 min, (M+H)⁺: 155. ¹H-NMR (CDCl₃): δ 6.2 (m, 1H), 5.8 (m, 1H), 5.3 (m, 2H), 3.4 (m, 3H), 3.3 (q, *J* = 7.2 Hz, 1H), 3.1 (m, 2H), 2.6 (m, 1H).

II.2.2. Protection of ketopiperazine B with DPP to afford protected ketopiperazine C₂




Scheme 9: DPP protection of **B**.

DPP protection is illustrated by the synthesis of intermediate (*R*)-4-(diphenylphosphoryl)-3-methylpiperazin-2-one (i.e. compound **C₂** wherein **R^{4'}** is Me).

To a solution of (*R*)-3-methylpiperazin-2-one (0.5 g, 4.38 mmol) in commercial anhydrous DCM (9 mL) under Ar atmosphere at rt was added diphenylphosphinic chloride (0.84 mL, 4.38 mmol) in one portion, followed by N-methylmorpholine (1.2 mL, 8.76 mmol) dropwise. The reaction mixture was stirred under reflux for 72h.

The reaction mixture was concentrated and the crude compound was then purified on silica gel (DCM/MeOH 99/1) to afford the desired product as colorless oil. Yield: 0.54 g, 88 %. LCMS: P = 98 %, retention time = 2.0 min, (M+H)⁺: 315; chiral HPLC retention time = 26.7 min, ee = 99.4 %; ¹H-NMR (CDCl₃): δ 7.9 (m, 4H), 7.5 (m, 6H), 6.1 (bs, 1H), 3.9 (m, 1H), 3.6 (m, 1H), 3.3 (m, 2H), 3.2 (m, 1H), 1.5 (m, 3H).

II.2.3. Protection of ketopiperazine **B** with Boc to afford protected ketopiperazine **C₃**




Scheme 10: Boc protection of **B**.

Boc protection is illustrated by the synthesis of intermediate (*R*)-tert-butyl 2-methyl-3-oxopiperazine-1-carboxylate (i.e. compound **C₃** wherein **R^{4'}** is Me).

To a solution of (*R*)-3-methylpiperazin-2-one (0.33 g, 2.87 mmol) in commercial anhydrous DCM (10 mL) at 0°C was added Boc₂O (0.77 mL, 3.30 mmol) in one portion. The reaction mixture was allowed to reach rt and stirred for 1h.

The reaction mixture was concentrated under reduced pressure and the residue was taken up in DCM (100 mL) and washed with HCl 0.5M (90 mL) and brine (120 mL), dried over MgSO₄, filtered and concentrated under reduced pressure. The crude compound was then purified on silica gel (DCM/MeOH 99/1) to afford the desired product as colorless oil. Yield: 0.45 g, 33 %. LCMS: P = 98 %, retention time = 1.9 min, (M+H)⁺: 215; ¹H-NMR (CDCl₃): δ 6.3 (bs, 1H), 4.6 (m, 1H), 4.1 (m, 1H), 3.5 (m, 1H), 3.3-3.1 (m, 2H), 1.5 (m, 3H), 1.4 (s, 9H).

II.2.4. Protection of ketopiperazine **B** with SES to afford protected ketopiperazine **C₄**



Scheme 11: SES protection of **B**.

SES protection is illustrated by the synthesis of intermediate (*R*)-3-methyl-4-((2-(trimethylsilyl)ethyl)sulfonyl)piperazin-2-one (i.e. compound **C₄** wherein **R^{4'}** is Me).


To a solution of (*R*)-3-methylpiperazin-2-one (0.25 g, 2.19 mmol) in commercial anhydrous DCM (4.5 mL) under Ar atmosphere at rt was added 2-(trimethylsilyl)ethanesulfonyl chloride (0.44 mL, 2.30 mmol) in one portion, followed by N-methylmorpholine (0.45 mL g, 4.38 mmol) dropwise. The reaction mixture was stirred at rt for 16h.

The reaction mixture was diluted with water (10 mL) and DCM (10 mL). The organic layer was separated, dried over MgSO₄, filtered and concentrated. The crude compound was then purified on silica gel (DCM/MeOH 99/1) to afford the desired product as colorless oil. Yield: 0.08 g, 13 %. LCMS: P = 95 %, retention time = 2.1 min, (M+H)⁺: 279; chiral HPLC retention time = 7.2 min, ee = 99.6 %; ¹H-NMR (CDCl₃): δ 6.1 (bs, 1H), 4.5 (m, 1H), 3.8 (m, 1H), 3.6 (m, 1H), 3.4 (m, 1H), 3.3 (m, 1H), 2.9 (m, 2H), 1.6 (m, 3H), 1.0 (m, 2H), 0.1 (s, 9H).

15 II.3. Step 2: Conversion to iminoether D

Method E: Conversion to iminoether

General Method E is the procedure used for the synthesis of intermediates **D**.



Scheme 12: Conversion to iminoether **D**.

Method E is illustrated by the synthesis of intermediate (*R*)-1-(2,4-dimethoxybenzyl)-5-ethoxy-6-methyl-1,2,3,6-tetrahydropyrazine **D₅₋₁** (i.e. compound **D** wherein **PG** is DMB and **R^{4'}** is Me). The corresponding DMB-protected ketopiperazine **C₅** is commercially available.

Oven dried (115°C) sodium carbonate (2.48 g, 23.40 mmol, 2.25 eq.) was placed in a round-bottom flask. The round-bottom flask was backfilled with Ar and then capped with a rubber septum. A solution of (*R*)-4-(2,4-dimethoxybenzyl)-3-methylpiperazin-2-one **C-1** (2.75 g, 10.40 mmol, 1 eq.) in anhydrous DCM (35 mL) was added, followed
5 by freshly prepared triethyloxonium tetrafluoroborate (2.48 g, 13.05 mmol, 1.25 eq.) in one portion. Thereafter the reaction mixture was stirred further at rt for 45 min to 1 hour, whereupon the reaction mixture was diluted with saturated aqueous NaHCO₃ (100 mL). The aqueous layer was extracted with DCM (3 x 200 mL). The organic layers were combined, dried over MgSO₄, filtered and concentrated under reduced pressure to
10 afford 3.1 g of yellow oil. The crude compound was then purified on silica gel (EtOAc/MeOH: 99/1) to afford the desired product **D-1** as a pale yellow oil. Yield: 1.44 g, 48 %. LCMS: P = 95 %, retention time = 1.8 min, (M+H₂O+H)⁺: 311; chiral HPLC retention time = 12.3 min, ee > 97 %. ¹H-NMR (CDCl₃): δ 7.23 (d, *J* = 8.8, 1H), 6.48 (d, *J* = 8.8, 1H), 6.44 (s, 1H), 4.02 (m, 2H), 3.92 (s, 6H), 3.86 (d, *J*_{AB} = 14.0, 1H),
15 3.46 (d, *J*_{AB} = 14.0, 1H), 3.44 (m, 2H), 3.10 (m, 1H), 2.79 (m, 1H), 2.32 (m, 1H), 1.35 (d, *J* = 6.8, 3H), 1.24 (t, *J* = 6.0, 3H).

The reaction mixture may alternatively be treated with brine. After stirring for about 20 min, additional water and DCM were added leading to phase separation. The organic layers were then dried over MgSO₄, filtered and concentrated under reduced pressure.
20 The crude compound was then purified on silica gel.

Method E is further illustrated by the synthesis of intermediate (*R*)-1-allyl-5-ethoxy-6-methyl-1,2,3,6-tetrahydropyrazine (i.e. compound **D₁** wherein **PG** is allyl and **R^{4'}** is Me).

To a solution of (*R*)-4-allyl-3-methylpiperazin-2-one (0.35 g, 2.27 mmol, 1 eq.) in DCM
25 (7.6 mL) at 0°C was added sodium carbonate (0.54 g, 5.11 mmol, 2.25 eq.) in one portion, followed by commercial triethyloxonium tetrafluoroborate (0.54 g, 2.84 mmol, 1.25 eq.) in one portion. Thereafter the reaction mixture was stirred further at rt for 45 min, whereupon the reaction mixture was diluted with DCM (10 mL) and brine (10 mL). The layers were separated and the aqueous layer was further extracted with

DCM (2 x 5 mL). The organic layers were combined, dried over MgSO_4 , filtered and concentrated under reduced pressure. The crude compound was then purified on silica gel (EtOAc) to afford the desired product as colorless oil. Yield: 0.19 g, 46 %. LCMS: P = 95 %, retention time = 1.5 min, $(\text{M}+\text{H})^+$: 183; $^1\text{H-NMR}$ (CDCl_3): δ 5.9 (m, 1H), 5.2 (m, 2H), 4.0 (m, 2H), 3.5 (m, 2H), 3.3 (m, 1H), 3.1-3.0 (m, 2H), 2.8 (m, 1H), 2.4 (m, 1H), 1.3 (m, 6H).

The following intermediates were also prepared from the *ad hoc* reagents:

(*R*)-(3-ethoxy-2-methyl-5,6-dihydropyrazin-1(2H)-yl)diphenylphosphine oxide in 44 % yield. LCMS: P = 98 %, retention time = 2.0 min, $(\text{M}+\text{H}_2\text{O}+\text{H})^+$: 361; chiral HPLC retention time = 4.8 min, ee = 99.4 % ; $^1\text{H-NMR}$ (CDCl_3): δ 7.9 (m, 4H), 7.5 (m, 6H), 4.0 (m, 2H), 3.7 (m, 1H), 3.6 (m, 1H), 3.5 (m, 1H), 3.1 (m, 2H), 1.4 (m, 3H), 1.2 (m, 3H).


(*R*)-tert-butyl 3-ethoxy-2-methyl-5,6-dihydropyrazine-1(2H)-carboxylate in 68% yield. LCMS: P = 98 %, retention time = 1.8 min, $(\text{M}+\text{H}_2\text{O}+\text{H})^+$: 261; $^1\text{H-NMR}$ (CDCl_3): δ 4.3 (m, 1H), 4.1 (m, 2H), 3.9 (m, 1H), 3.5 (m, 2H), 2.9 (m, 1H), 1.5 (s, 9H), 1.3 (d, J = 6.9 Hz, 3H), 1.2 (t, J = 7.0 Hz, 3H).

(*R*)-5-ethoxy-6-methyl-1-((2-(trimethylsilyl)ethyl)sulfonyl)-1,2,3,6-tetrahydropyrazine in 68 % yield. LCMS: P = 70 %, retention time = 2.0 min, $(\text{M}+\text{H}_2\text{O}+\text{H})^+$: 325; chiral HPLC retention time = 4.8 min, ee = 97.3 % ; $^1\text{H-NMR}$ (CDCl_3): δ 4.3 (m, 1H), 4.1 (m, 2H), 3.6 (m, 3H), 3.2 (m, 1H), 2.9 (m, 2H), 1.5 (m, 3H), 1.3 (m, 3H), 1.0 (m, 2H), 0.0 (s, 9H).

II.4. Step 3: Cyclodehydration leading to F

Method F: Cyclodehydration

General Method F is the general procedure used for the synthesis of chiral triazolopiperazine intermediates **F**.



Scheme 13: Formation of acylhydrazide **F**.

In a round-bottom flask equipped with a condenser, imino-ether **D** (1 eq.) was dissolved in anhydrous MeOH, to which was added **E** (1 eq.) in one portion. The resulting solution was stirred at a temperature ranging from 55°C to 70 °C for a period of time ranging from 6 hours to 8 hours. Completion of the reaction was monitored by HPLC analysis. The reaction mixture was cooled down to rt and the solvent was removed under reduced pressure. The crude compound was then purified by silica gel chromatography to afford the desired product **F**.

In an embodiment of the invention, the crude compound precipitates during cooling of the reaction mixture. In this case, the precipitate is stirred at rt in MeOH for about 5 hours before being filtered, washed with MeOH and oven dried.

Cyclodehydration is illustrated by the synthesis of intermediate (*R*)-5-(7-allyl-8-methyl-5,6,7,8-tetrahydro-[1,2,4]triazolo[4,3-*a*]pyrazin-3-yl)-3-methyl-1,2,4-thiadiazole (i.e. compound **F**₁ wherein **PG** is allyl, **R**^{4'} is Me, **X**¹ is N, **X**² is S and **R**^{5'} is methyl).

To (*R*)-1-allyl-5-ethoxy-6-methyl-1,2,3,6-tetrahydropyrazine (0.14 g, 0.77 mmol) at rt was added 3-methyl-1,2,4-thiadiazole-5-carbohydrazide (0.12 g, 0.77 mmol) at once. The mixture was diluted with commercially anhydrous MeOH (0.77 mL) to allow complete solubilization and the resulting mixture was heated to 60 °C for 16h.

The reaction mixture was then allowed to reach rt whereupon the solvent was removed under reduced pressure (1-2 mbar). The crude residue was then dissolved in DCM (10 mL), and thus-obtained organic phase washed with NaOH (1 M, 10 mL). The

organic layer was then dried over MgSO_4 , filtered and concentrated under reduced pressure (1-2 mbar) the desired product as a yellow solid. Yield: 0.09 g, 42 %. LCMS: P = 95 %, retention time = 1.6 min, $(\text{M}+\text{H})^+$: 277; chiral HPLC retention time = 21.6 min, ee = 98.9 %; $^1\text{H-NMR}$ (CDCl_3): δ 5.9 (m, 1H), 5.3 (m, 2H), 4.5 (m, 1H), 4.4 (m, 1H), 4.1 (m, 1H), 3.5 (m, 1H), 3.3 (m, 1H), 3.1 (m, 1H), 2.8 (m, 1H), 2.7 (s, 3H), 1.6 (m, 3H).

The following intermediates were also prepared from the *ad hoc* reagents:

(*R*)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)diphenylphosphine oxide in 31 % yield (reaction time: 48h and silica gel purification (EtOAc)). LCMS: P = 96 %, retention time = 2.2 min, $(\text{M}+\text{H})^+$: 437; chiral HPLC retention time = 7.5 min, ee = 98.3 %; $^1\text{H-NMR}$ (CDCl_3): δ 7.9 (m, 4H), 7.5 (m, 6H), 4.9 (m, 1H), 4.8 (dd, J = 3.1, 13.6 Hz, 1H), 4.3 (dt, J = 4.9, 12.2 Hz, 1H), 3.6 (m, 1H), 3.5 (m, 1H), 2.7 (s, 3H), 1.6 (d, J = 6.9 Hz, 3H).

(*R*)-tert-butyl 8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazine-7(8H)-carboxylate in 83% yield (reaction time: 48h). LCMS: P = 97 %, retention time = 2.3 min, $(\text{M}+\text{H})^+$: 337; chiral HPLC retention time = 19.4 min, ee = 95.1 %; $^1\text{H-NMR}$ (CDCl_3): δ 5.7 (m, 1H), 4.9 (m, 1H), 4.5 (m, 1H), 4.2 (m, 1H), 3.3 (m, 1H), 2.7 (s, 3H), 1.6 (d, J = 6.9 Hz, 3H), 1.5 (s, 9H).


(*R*)-3-methyl-5-(8-methyl-7-((2-(trimethylsilyl)ethyl)sulfonyl)-5,6,7,8-tetrahydro-[1,2,4]triazol[4,3-a]pyrazin-3-yl)-1,2,4-thiadiazole in 28 % yield (reaction time: 48h). LCMS: P = 40 %, retention time = 2.5 min, $(\text{M}+\text{H})^+$: 401; chiral HPLC retention time = 7.1 min, ee = 92.4 %; $^1\text{H-NMR}$ (CDCl_3): δ 4.9 (m, 1H), 4.3 (m, 1H), 4.1 (m, 1H), 3.6 (m, 1H), 3.0 (m, 1H), 2.7 (s, 3H), 1.6 (m, 2H), 1.4 (m, 3H), 1.0 (m, 2H), 0.0 (s, 9H).

II.5. Step 4: PG-deprotection

The methods of deprotection of above Protecting Groups (PGs) are known to those skill-in-the-art. As examples, one may refer to “Greene’s Protective Groups in Organic Synthesis”:

- Allyl: p. 806 of fourth edition;
- DPP: p. 844 of fourth edition;
- Boc: p. 725 of fourth edition;
- SES: p. 854 in fourth edition.

5 Method G: DMB deprotection – TFA/DCM




Scheme 14: DMB deprotection.

Deprotection of DMB may be performed using TFA.

When crude or precipitated **F** was used (in opposition to purified **F** on silica gel), pre-washing was performed before deprotection as follow: **F** was dissolved in DCM and optionally washed with 1M NaOH in order to remove remaining **E**. The DCM extracts were then dried over magnesium sulphate, filtered and the filter cake washed with DCM.

F was diluted with DCM and TFA (7.6 eq.) was added to the DCM solution of **F** at RT. The mixture was stirred at rt for 2h-2h30. Completion of the deprotection was monitored by HPLC. Water was added, the mixture stirred for 30 minutes and filtered. The filter cake was washed with water and DCM. The filtrate layers were separated. The pH of the aqueous layer was adjusted to 12-13 by the addition of 4M NaOH. Sodium chloride was then added and the aqueous solution was extracted with DCM. The DCM extract comprising **G** was concentrated and was used in the next step without further purification.

II.6. Optional conversion of $R^{5'}$ to R^5 in triazolopiperazine **G**

Scheme 15: conversion of $R^{5'}$ to R^5 in triazolopiperazine **G** leading to **G'**.

Substituent R^5 may then be introduced, when applicable, from $R^{5'}$ (especially when $R^{5'}$ = H). One example of such transformation is illustrated by the synthesis of intermediate **G'** wherein R^5 is trifluoromethyl.

To a solution of **G** (1 eq.) in DCM/water (3/1) are added, at rt, sodium trifluoromethanesulfinate (3 eq.) and 2-hydroperoxy-2-methylpropane (5 eq.). The reaction mixture is *not* stirred and left at rt. Monitoring conversion by HPLC-MS, extra amount of each reagent can be added if required. The resulting mixture is diluted with DCM and quenched with 4 M NaOH saturated solution. Layers were separated and aqueous layer was extracted twice with EtOAc. The organic phases are combined, dried over MgSO_4 and evaporated to dryness. The residue is purified on silica gel or used crude in next step.

A further example of such transformation is illustrated by the synthesis of intermediate **G'** wherein R^5 is difluoromethyl:

To a suspension of **G** ($R^{5'} = \text{H}$) (*R*)-5-(8-methyl-5,6,7,8-tetrahydro-[1,2,4]triazolo[4,3-*a*]pyrazin-3-yl)-1,2,4-thiadiazole (0.29 g, 1.13 mmol) and bis(difluoromethylsulfinyloxy)zinc (0.67 g, 2.26 mmol) in DCM (5 mL) and Water (2 mL), was added TFA (0.09 mL, 1.13 mmol), followed by slow addition of 2-hydroperoxy-2-methylpropane (0.77 mL, 5.64 mmol) with vigorous stirring.


When conversion was not increasing any more (HPLC-MS monitoring) bis(difluoromethylsulfinyloxy)zinc and 2-hydroperoxy-2-methylpropane were added at rt still with vigorous stirring (3 additional times (1.001 g, 3.39 mmol) and (0.773 mL, 5.64 mmol) respectively).

- 5 After 4 days in total, reaction mixture was diluted with EtOAc (50 mL) and carefully quenched with NaHCO₃ sat. solution (30 mL) and then NaHCO₃ solid until no bubbling was observed. Reaction mixture was filtered on Celite pad and phases of the filtrate were separated. Aqueous phase was filtered again on Celite pad then filtrate was extracted with EtOAc (2 x 50 mL). Organic phases were combined, dried over MgSO₄,
 10 filtered and concentrated under reduced pressure. The crude compound was then purified on silica gel (DCM/MeOH 99/1) to afford the desired product as colorless oil. Yield: 0.03 g, 10 %. LCMS: P = 97 %, retention time = 1.8 min, (M+H)⁺: 273; ¹H-NMR (CDCl₃): δ 6.8 (t, J_{H-F} = 53.5 Hz, 1H), 4.7 (m, 1H), 4.3 (m, 2H), 3.5 (m, 1H), 3.3 (m, 1H), 1.7 (d, J = 6.7 Hz, 3H); ¹⁹F-NMR (CDCl₃): δ -113.5 (dd, J = 3.2, 53.5 Hz, 2F).

15 II.7. Step 5: Acylation leading to products I

Method H: Acylation NMM/DCM

General Method H is the general procedure used for the synthesis of (*R*)-enantiomer of **Formula I** of the invention.



20

Scheme 16: Acylation.

To a solution of crude **G** or **G'** (1 eq.) in anhydrous DCM were added at 0°C **H** (1.3 eq.), followed by N-methylmorpholine (2.2 eq.) dropwise over 15 sec. The reaction

mixture was stirred at rt for 10 minutes and, the milky suspension was poured into 1 M HCl. The aqueous phase was extracted with DCM. The organic phases were combined, washed with 1 M NaOH, brine, dried over MgSO₄ and evaporated to dryness. The crude compound was purified by silica gel chromatography to afford the desired product (**R**)-

5 **I.**

Measurement of %ee confirmed that no detectable racemization occurs during the acidolytic deprotection and N-acylation steps.

Method I: Acylation – biphasic conditions

Alternatively, the reaction may be performed under biphasic conditions.

- 10 In this case, saturated sodium hydrogen carbonate solution was added to the DCM slurry of **G** or **G'** (1 eq.) at rt. **H** (1 eq.) was added and the mixture stirred for a period of time ranging from about 20 minutes to overnight at rt. Completion of the reaction was monitored by HPLC. The layers were separated and the DCM phase washed with water. The DCM extracts were dried with magnesium sulphate and filtered, washing the
- 15 filter cake with DCM. The DCM extracts were then concentrated. TBME was added and the resulting slurry stirred overnight at rt. The solid was collected by filtration, washed with TBME and pulled dry. The crude compound may be purified by silica gel chromatography or by crystallisation.

- Measurement of %ee confirmed that no detectable racemization occurs during the
- 20 acidolytic deprotection and N-acylation steps.

Substituent R^{4'} may then be transformed, when applicable, into R⁴ (see racemic synthesis).

II.8. Optional further transformation leading to products I''/I''' from I/I'

- Compound 45: From compound **I/I'** wherein **R^{5''}** = 1-((tert-butyl
- 25 butyldiphenylsilyl)oxy)ethyl, well known *tert*-butylamonium fluoride TBDPS deprotection of alkoxy was applied, followed by DAST fluorination of the latter

alcohol, leading to racemic compound 45. Both diastereomers can be separated by purification on preparative HPLC to afford 45-1 and 45-2.

Compound 43: From compound **I/I'** wherein **R^{5''}** = 1-((tert-butyldiphenylsilyl)oxy)ethyl, well known *tert*-butylammonium fluoride TBDPS deprotection of alkoxy was applied, followed by Dess-Martin oxidation, then followed by DAST fluorination of the latter ketone, leading to compound 43.

III. Chemical characterization

Compound 1: HPLC-MS: $t_R = 4.1$ min, $(M+H)^+ = 409$; Chiral HPLC (Method C): %ee = 99.0; $^1\text{H-NMR}$ (CDCl_3): δ 7.6 (m, 2H), 7.3 (m, 1H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 2.7 (s, 3H), 1.7 (d, 3H).

Compound 2: HPLC-MS: $t_R = 3.8$ min, $(M+H)^+ = 373$; Chiral HPLC (Method A): %ee = 98.0; $^1\text{H-NMR}$ (300 MHz, CDCl_3): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 3.1 (q, 2H), 1.8 (d, 3H), 1.4 (t, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -98.5.

Compound 3: HPLC-MS: $t_R = 3.8$ min, $(M+H)^+ = 375$; Chiral HPLC (Method C): %ee > 99.8; $^1\text{H-NMR}$ (CDCl_3): δ 7.5 (m, 4H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 2.7 (s, 3H), 1.7 (d, 3H).

Compound 4: HPLC-MS: $t_R = 3.9$ min, $(M+H)^+ = 393$; Chiral HPLC (Method C): %ee = 99.0; $^1\text{H-NMR}$ (CDCl_3): δ 7.6 (m, 1H), 7.3 (s, 1H), 7.2 (m, 1H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 2.7 (s, 3H), 1.7 (d, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -98.4.

Compound 5: HPLC-MS: $t_R = 3.4$ min, $(M+H)^+ = 359$; Chiral HPLC (Method C): %ee = 99.0; $^1\text{H-NMR}$ (CDCl_3): δ 7.5 (m, 2H), 7.3 (m, 2H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.5 (m, 1H), 2.7 (s, 3H), 1.7 (d, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -98.4.

Compound 6: HPLC-MS: $t_R = 3.8$ min, $(M+H)^+ = 393$; Chiral HPLC (Method C): %ee = 99.5; $^1\text{H-NMR}$ (CDCl_3): δ 7.6 (m, 1H), 7.3 (m, 2H), 5.7 (m, 1H), 4.9 (m, 1H), 4.5 (m, 1H), 4.3 (m, 1H), 3.5 (m, 1H), 2.7 (s, 3H), 1.8 (d, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -96.2.

5 Compound 7: HPLC-MS: $t_R = 3.8$ min, $(M+H)^+ = 395$; Chiral HPLC (Method C): %ee = 98.9; $^1\text{H-NMR}$ (CDCl_3): δ 7.1 (m, 2H), 5.8 (m, 1H), 5.0 (m, 1H), 4.5 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 2.7 (s, 3H), 1.8 (d, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -75.8.

Compound 8: HPLC-MS: $t_R = 3.7$ min, $(M+H)^+ = 395$; Chiral HPLC (Method C): %ee = 99.0; $^1\text{H-NMR}$ (CDCl_3): δ 7.1 (m, 2H), 6.2 (m, 1H), 5.3-5.0 (m, 2H), 4.3-3.6 (m, 2H), 2.7 (s, 3H), 1.8 (m, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -49.4, -72.0, -77.4.

10 Compound 9: HPLC-MS: $t_R = 3.6$ min, $(M+H)^+ = 377$; Chiral HPLC (Method C): %ee = 99.4; $^1\text{H-NMR}$ (CDCl_3): δ 7.3 (m, 3H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.7 (s, 3H), 1.7 (d, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -72.1, -74.4.

Compound 10: HPLC-MS: $t_R = 4.0$ min, $(M+H)^+ = 413$; Chiral HPLC (Method C): %ee = 99.0; $^1\text{H-NMR}$ (CDCl_3): δ 7.2 (m, 1H), 6.2 (m, 1H), 5.2-5.0 (m, 2H), 4.3 (m, 1H),
15 3.9-3.4 (m, 2H), 2.7 (s, 3H), 1.8 (m, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -54.2, -56.3, -67.1, -72.8.

Compound 11: HPLC-MS: $t_R = 3.2$ min, $(M+H)^+ = 389$; Chiral HPLC (Method A): %ee > 99.8; $^1\text{H-NMR}$ (CDCl_3): δ 7.5 (m, 2H), 7.2 (m, 2H), 6.1 (m, 1H), 4.9 (dd, 1H), 4.3 (m, 2H), 3.9 (m, 2H), 3.6 (m, 1H), 2.7 (s, 3H), 2.4 (m, 1H), 2.2 (m, 1H); $^{19}\text{F-NMR}$ (CDCl_3):
20 δ -97.9.

Compound 12: is racemate of compound 11.

Compound 13: HPLC-MS: $t_R = 4.1$ min, $(M+H)^+ = 357$; Chiral HPLC (Method B): %ee = 98.7; $^1\text{H-NMR}$ (CDCl_3): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.8 (m, 1H), 4.8 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.9 (q, 2H), 1.8 (d, 3H), 1.4 (t, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ
25 -98.7.

Compound 14: is racemate of compound 5.

Compound 15: HPLC-MS: $t_R = 3.4$ min, $(M+H)^+ = 359$; Chiral HPLC (Method B): %ee > 99.8; $^1\text{H-NMR}$ (CDCl_3): δ 7.5 (m, 1H), 7.2 (m, 3H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.7 (s, 3H), 1.7 (d, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -96.2.

5 Compound 16: HPLC-MS: $t_R = 3.7$ min, $(M+H)^+ = 375$; $^1\text{H-NMR}$ (CDCl_3): δ 7.5-7.3 (m, 4H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.7 (s, 3H), 1.7 (d, 3H).

Compound 17: HPLC-MS: $t_R = 3.6$ min, $(M+H)^+ = 377$; $^1\text{H-NMR}$ (CDCl_3): δ 7.3 (m, 1H), 7.0 (m, 2H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.8 (s, 3H), 1.8 (d, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -101.2.

10 Compound 18: HPLC-MS: $t_R = 3.5$ min, $(M+H)^+ = 377$; $^1\text{H-NMR}$ (CDCl_3): δ 7.5 (m, 1H), 7.0-6.9 (m, 2H), 6.2 (m, 1H), 5.2-4.9 (m, 2H), 4.3 (m, 1H), 4.0-3.7 (m, 1H), 2.7 (s, 3H), 1.7 (m, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -95.7, -102.5.

15 Compound 19: HPLC-MS: $t_R = 3.6$ min, $(M+H)^+ = 355$; $^1\text{H-NMR}$ (CDCl_3): δ 7.4 (m, 4H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.7 (s, 3H), 2.4 (s, 3H), 1.7 (d, 3H).

Compound 20: HPLC-MS: $t_R = 3.3$ min, $(M+H)^+ = 341$; Chiral HPLC (Method C): %ee = 96.8; $^1\text{H-NMR}$ (CDCl_3): δ 7.5 (m, 5H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.7 (s, 3H), 1.7 (d, 3H).

20 Compound 21: HPLC-MS: $t_R = 4.0$ min, $(M+H)^+ = 409$; $^1\text{H-NMR}$ (CDCl_3): δ 7.7 (d, 2H), 7.6 (d, 1H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (m, 1H), 2.7 (s, 3H), 1.7 (d, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -60.1.

25 Compound 22: HPLC-MS: $t_R = 3.7$ min, $(M+H)^+ = 373$; Chiral HPLC (Method B): %ee > 99.7; $^1\text{H-NMR}$ (CDCl_3): δ 7.5 (m, 2H), 7.3 (m, 2H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 2.7 (s, 3H), 2.2-2.0 (m, 2H), 1.1 (m, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -98.4.

Compound 23: is racemate of compound 22.

Compound 24: HPLC-MS: $t_R = 4.0$ min, $(M+H)^+ = 387$; Chiral HPLC (Method D): %ee = 95.5; $^1\text{H-NMR}$ (CDCl_3): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.2 (m, 1H), 3.6 (m, 1H), 2.7 (s, 3H), 2.1-2.0 (m, 2H), 1.6 (m, 2H), 1.0 (m, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -98.2.

- 5 Compound 25: HPLC-MS: $t_R = 3.5$ min, $(M+H)^+ = 389$; $^1\text{H-NMR}$ (CDCl_3): δ 7.2 (m, 2H), 7.0 (m, 2H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.9 (s, 3H), 3.5 (td, 1H), 2.7 (s, 3H), 1.8 (d, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -76.3.

- Compound 26: HPLC-MS: $t_R = 3.5$ min, $(M+H)^+ = 355$; $^1\text{H-NMR}$ (CDCl_3): δ 7.4-7.2 (m, 4H), 6.3 (m, 1H), 5.3-4.8 (m, 2H), 4.3-3.8 (m, 1H), 3.5-3.4 (m, 1H), 2.7 (2s, 3H),
10 2.3 (s, 3H). 1.7 (2s, 3H).

Compound 27: HPLC-MS: $t_R = 3.4$ min, $(M+H)^+ = 371$; $^1\text{H-NMR}$ (CDCl_3): δ 7.4 (m, 1H), 7.0 (m, 3H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.8 (s, 3H), 3.5 (td, 1H), 2.7 (s, 3H), 1.7 (d, 3H).

- Compound 28: HPLC-MS: $t_R = 3.1$ min, $(M+H)^+ = 343$; Chiral HPLC (Method C): %ee
15 = 96.1; $^1\text{H-NMR}$ (CDCl_3): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.5 (td, 1H), 2.5 (s, 3H), 1.7 (d, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -98.3.

Compound 29: HPLC-MS: $t_R = 3.2$ min, $(M+H)^+ = 366$; Chiral HPLC (Method B): %ee = 99.0; $^1\text{H-NMR}$ (CDCl_3): δ 7.8 (d, 2H), 7.6 (d, 2H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.7 (s, 3H), 1.7 (d, 3H).

- 20 Compound 30: HPLC-MS: $t_R = 4.9$ min, $(M+H)^+ = 421$; Chiral HPLC (Method A): %ee = 98.2; $^1\text{H-NMR}$ (CDCl_3): δ 7.7 (d, 2H), 7.5 (d, 2H), 7.4 (m, 2H), 7.1 (m, 1H), 5.9 (m, 1H), 4.8 (dd, 1H), 4.7 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 2.9 (q, 2H), 1.8 (d, 3H), 1.4 (t, 3H).

Compound 31: is racemate of compound 10.

- 25 Compound 32: is racemate of compound 9.

Compound 33: is racemate of compound 8.

Compound 34: is racemate of compound 7.

Compound 35: is racemate of compound 6.

Compound 36: is racemate of compound 4.

Compound 37: is racemate of compound 3.

Compound 38: is racemate of compound 1.

5 Compound 39: is racemate of compound 2.

Compound 40: is racemate of compound 13.

Compound 41: HPLC-MS: $t_R = 4.8$ min, $(M+H)^+ = 413$; Chiral HPLC (Method B): %ee = 99.7; ^1H -NMR ($CDCl_3$): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.8 (m, 1H), 4.9 (dd, 1H), 4.6 (m, 1H), 4.3 (td, 1H), 3.6 (td, 1H), 1.8 (d, 3H); ^{19}F -NMR ($CDCl_3$): δ -62.9, -98.7.

10 Compound 42: HPLC-MS: $t_R = 4.3$ min, $(M+H)^+ = 395$; Chiral HPLC (Method B): %ee = 97.4; ^1H -NMR ($CDCl_3$): δ 7.5 (m, 2H), 7.1 (m, 2H), 6.8 (t, $J_{H-F} = 53.5$ Hz, 1H), 5.8 (m, 1H), 4.9 (dd, $J = 3.1, 13.6$ Hz, 1H), 4.6 (m, 1H),), 4.3 (dt, $J = 4.6, 13.3$ Hz, 1H), 3.6 (m, 1H), 1.8 (d, $J = 6.9$ Hz, 3H); ^{19}F -NMR ($CDCl_3$): δ -105.2 (s, 1F), -113.4 (dd, $J = 9.6, 53.4$ Hz, 2F).

15 Compound 43: HPLC-MS: $t_R = 4.4$ min, $(M+H)^+ = 393$; Chiral HPLC (Method C): %ee = 96.3; ^1H -NMR ($CDCl_3$): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.9 (m, 1H), 4.8 (dd, $J = 3.3, 13.5$ Hz, 1H), 4.6 (m, 1H), 4.3 (dt, $J = 4.2, 12.7$ Hz, 1H), 3.6 (m, 1H), 2.2 (t, $J = 8.6$ Hz, 3H), 1.8 (d, $J = 6.9$ Hz, 3H); ^{19}F -NMR ($CDCl_3$): δ -88.3 (q, $J = 18.3$ Hz, 2F), -105.0 (s, 1F).

20 Compound 44: HPLC-MS: $t_R = 4.4$ min, $(M+H)^+ = 411$; Chiral HPLC (Method C): %ee = 98.6; ^1H -NMR ($CDCl_3$): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.8 (m, 1H), 4.8 (dd, $J = 3.5, 13.6$ Hz, 1H), 4.6 (m, 1H), 4.3 (dt, $J = 4.0, 12.2$ Hz, 1H), 3.7 (q, $J = 10.0$ Hz, 2H), 3.6 (m, 1H), 1.8 (d, $J = 6.9$ Hz, 3H); ^{19}F -NMR ($CDCl_3$): δ -61.1 (t, $J = 9.6$ Hz, 1F), -105.0 (s, 1F).

25 Compound 45: HPLC-MS: $t_R = 4.4$ min, $(M+H)^+ = 375$; Chiral HPLC (Method C): %ee = 98.5; ^1H -NMR ($CDCl_3$): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.9 (m, 1H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 1.9 (d, $J = 6.9$ Hz, 3H), 1.8 (m, 3H); ^{19}F -NMR ($CDCl_3$): δ -105.0 (s, 1F), -175.0 (m, 1F).

Compound 45-1: HPLC-MS: $t_R = 4.4$ min, $(M+H)^+ = 375$; Chiral HPLC (Method C): %ee = 99.2; $^1\text{H-NMR}$ (CDCl_3): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.9 (m, 1H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 1.9 (d, $J = 6.9$ Hz, 3H), 1.8 (m, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -105.0 (s, 1F), -175.0 (m, 1F).

- 5 Compound 45-2: HPLC-MS: $t_R = 4.4$ min, $(M+H)^+ = 375$; Chiral HPLC (Method C): %ee = 91.7; $^1\text{H-NMR}$ (CDCl_3): δ 7.5 (m, 2H), 7.2 (m, 2H), 5.9 (m, 1H), 5.8 (m, 1H), 4.9 (m, 1H), 4.6 (m, 1H), 4.3 (m, 1H), 3.6 (m, 1H), 1.9 (d, $J = 6.9$ Hz, 3H), 1.8 (m, 3H); $^{19}\text{F-NMR}$ (CDCl_3): δ -105.0 (s, 1F), -175.0 (m, 1F).

* * *

10 BIOLOGY EXAMPLES

Functional Assay

Aequorin assay with human NK-3 receptor

- Changes in intracellular calcium levels are a recognized indicator of G protein-coupled receptor activity. The efficacy of compounds of the invention to inhibit NKA-mediated
- 15 NK-3 receptor activation was assessed by an *in vitro* Aequorin functional assay. Chinese Hamster Ovary recombinant cells expressing the human NK-3 receptor and a construct that encodes the photoprotein apoaequorin were used for this assay. In the presence of the cofactor coelenterazine, apoaequorin emits a measurable luminescence that is proportional to the amount of intracellular (cytoplasmic) free calcium.

20 Antagonist testing

- The antagonist activity of compounds of the invention is measured following pre-incubation (3 minutes) of the compound (at various concentrations) with the cells, followed by addition of the reference agonist (NKA) at a final concentration equivalent to the EC_{80} (3 nM) and recording of emitted light (FDSS 6000 Hamamatsu) over the
- 25 subsequent 90-second period. The intensity of the emitted light is integrated using the

reader software. Compound antagonist activity is measured based on the concentration-dependent inhibition of the luminescence response to the addition of Neurokinin A.

Inhibition curves are obtained for compounds of the invention and the concentrations of compounds which inhibit 50% of reference agonist response (IC_{50}) were determined (see results in table 2below). The IC_{50} values shown in table 2 indicate that compounds of the invention are potent NK-3 antagonist compounds.

Competitive binding assays

The affinity of compounds of the invention for the human NK-3 receptor was determined by measuring the ability of compounds of the invention to competitively and reversibly displace a well-characterized NK-3 radioligand in a concentration-dependent manner.

^3H -SB222200 binding competition assay with human NK-3 receptor

The ability of compounds of the invention to inhibit the binding of the NK-3 receptor selective antagonist ^3H -SB222200 was assessed by an in vitro radioligand binding assay. Membranes were prepared from Chinese hamster ovary recombinant cells stably expressing the human NK-3 receptor. The membranes were incubated with 5nM ^3H -SB222200 (ARC) in a HEPES 25mM/ NaCl 0.1M/CaCl₂ 1mM/MgCl₂ 5mM/ BSA 0.5%/ Saponin 10μg/ml buffer at pH 7.4 and various concentrations of compounds of the invention. The amount of ^3H -SB222200 bound to the receptor was determined after filtration by the quantification of membrane associated radioactivity using the TopCount-NXT reader (Packard). Competition curves were obtained for compounds of the invention and the concentration that displaced 50% of bound radioligand (IC_{50}) were determined by linear regression analysis and then the apparent inhibition constant (K_i) values were calculated by the following equation: $K_i = IC_{50}/(1+[L]/K_d)$ where [L] is the concentration of free radioligand and K_d is its dissociation constant at the receptor, derived from saturation binding experiments (Cheng and Prusoff, 1973) (see results in table 2below).

Table 2 shows biological results obtained using the ^3H -SB222200 binding competition assay with compounds of the invention. These results indicate that compounds of the invention display potent affinity for the human NK-3 receptor.

TABLE 2

Cpd n°	Functional assay: Aequorin assay with human NK-3 receptor hNK-3 – AEQ(antagonist IC₅₀, nM)	Competitive binding assay with human NK-3 receptor hNK-3 (K_i, nM)
1	16	11
2	12	15
3	32	19
4	19	20
5	18	23
6	30	24
7	30	26
8	33	26
9	21	30
10	56	31
11	73	32
12	170	59
13	44	40
14	57	42
15	50	45
16	71	49
17	50	51
18	87	54
19	110	56
20	93	60
21	150	63
22	130	69
23	220	150
24	120	78
25	110	85

26	74	88
27	220	100
28	160	110
29	170	150
30	5	7
31	64	70
32	44	59
33	120	89
34	62	38
35	54	73
36	58	43
37	52	41
38	34	26
39	32	28
40	130	83
41	7	11
42	39	48
43	136	59
44	204	45
45	244	101
45-1	285	100
45-2	148	83

Selectivity assay

Selectivity of the compounds of the invention was determined over the other human NK receptors, namely NK-1 and NK-2 receptors.

Human NK-1

- 5 The affinity of compounds of the invention for the NK-1 receptor was evaluated in CHO recombinant cells which express the human NK-1 receptor. Membrane suspensions were prepared from these cells. The following radioligand: [³H] substance P (PerkinElmer Cat#NET111520) was used in this assay. Binding assays were

performed in a 50 mM Tris / 5 mM MnCl₂ / 150 mM NaCl / 0.1% BSA at pH 7.4. Binding assays consisted of 25 µl of membrane suspension (approximately 5 µg of protein/well in a 96 well plate), 50 µl of compound or reference ligand (Substance P) at increasing concentrations (diluted in assay buffer) and 2nM [³H] substance P. The plate
5 was incubated 60 min at 25°C in a water bath and then filtered over GF/C filters (Perkin Elmer, 6005174, presoaked in 0.5% PEI for 2h at room temperature) with a Filtration unit (Perkin Elmer). The radioactivity retained on the filters was measured by using the TopCount-NXT reader (Packard). Competition curves were obtained for compounds of the invention and the concentrations of compounds which displaced 50% of bound
10 radioligand (IC₅₀) were determined and then apparent inhibition constant K_i values were calculated by the following equation: $K_i = IC_{50}/(1+[L]/K_D)$ where [L] is the concentration of free radioligand and K_D is its dissociation constant at the receptor, derived from saturation binding experiments (Cheng and Prusoff, 1973).

Human NK-2

15 The affinity of compounds of the invention for the NK-2 receptor was evaluated in CHO recombinant cells which express the human NK-2 receptor. Membrane suspensions were prepared from these cells. The following radioligand [¹²⁵I]-Neurokinin A (PerkinElmer Cat#NEX252) was used in this assay. Binding assays were performed in a 25 mM HEPES / 1 mM CaCl₂ / 5 mM MgCl₂/ 0.5% BSA / 10µg/ml
20 saponin, at pH 7.4. Binding assays consisted of 25 µl of membrane suspension (approximately 3.75 µg of protein/well in a 96 well plate), 50 µl of compound or reference ligand (Neurokinin A) at increasing concentrations (diluted in assay buffer) and 0.1 nM [¹²⁵I]-Neurokinin A. The plate was incubated 60 min at 25°C in a water bath and then filtered over GF/C filters (Perkin Elmer, 6005174, presoaked in assay buffer
25 without saponine for 2h at room temperature) with a Filtration unit (Perkin Elmer). The radioactivity retained on the filters was measured by using the TopCount-NXT reader (Packard). Competition curves were obtained for compounds of the invention and the concentrations of compounds which displaced 50% of bound radioligand (IC₅₀) were determined and then apparent inhibition constant K_i values were calculated by the
30 following equation: $K_i = IC_{50}/(1+[L]/K_D)$ where [L] is the concentration of free

radioligand and K_D is its dissociation constant at the receptor, derived from saturation binding experiments (Cheng and Prusoff, 1973).

The compounds of the invention, which were tested in the above NK-1 and NK-2 described assays, demonstrated a low affinity at the human NK-1 and human NK-2
5 receptors: more than 200 fold shift of the K_i compared to the human NK-3 receptor (table 3). Thus, compounds according to the invention have been shown to be selective over NK-1 and NK-2 receptors.

TABLE 3

Cpd n°	hNK-3 (K_i , nM)	hNK-1 (K_i , nM)	hNK-2 (K_i , nM)
1	11	10300	7500
2	15	23800	>30000
3	19	>30000	>30000
4	20	19900	23000
5	23	>30000	>30000
6	24	22100	25000
7	26	>30000	36000
8	26	>30000	>30000
9	30	>30000	>30000
10	31	>30000	49000
11	32	22000	>30000
12	59	NA	NA
13	40	>30000	>30000
14	42	>30000	>30000
15	45	>30000	>30000
16	49	>30000	37000
17	51	>30000	>30000
18	54	>30000	>30000
19	56	>30000	>30000
20	60	>30000	>30000
21	63	>30000	>30000
22	69	>30000	>30000
23	150	NA	NA
24	78	>30000	>30000
25	85	>30000	>30000

26	88	>30000	>30000
27	100	>30000	>30000
28	110	>30000	>30000
29	150	>30000	>30000
30	7	40000	32000
31	70	>30000	>30000
32	59	>30000	>30000
33	89	>30000	>30000
34	38	>30000	>30000
35	73	>30000	30000
36	43	>30000	36000
37	41	32000	>30000
38	26	21000	28000
39	28	>30000	>30000
40	83	>30000	>30000
41	11	NA	NA
42	48	>30000	>30000
43	59	>30000	>30000
44	45	>30000	>30000
45	101	>30000	>30000
45-1	100	>30000	>30000
45-2	83	>30000	>30000

NA: not available

hERG inhibition Assay

The human ether-a-go-go related gene (hERG) encodes the inward rectifying voltage gated potassium channel in the heart (I_{Kr}) which is involved in cardiac repolarisation. I_{Kr} current inhibition has been shown to elongate the cardiac action potential, a phenomenon associated with increased risk of arrhythmia. I_{Kr} current inhibition accounts for the vast majority of known cases of drug-induced QT-prolongation. A number of drugs have been withdrawn from late stage clinical trials due to these cardiotoxic effects, therefore it is important to identify inhibitors early in drug discovery.

The hERG inhibition study aims at quantifying the *in vitro* effects of compounds of the invention on the potassium-selective IK_r current generated in normoxic conditions in stably transfected HEK 293 cells with the human ether-a-go-go-related gene (hERG).

5 Whole-cell currents (acquisition by manual patch-clamp) elicited during a voltage pulse were recorded in baseline conditions and following application of tested compounds (5 minutes of exposure). The concentrations of tested compounds (0.3 μ M; 3 μ M; 10 μ M; 30 μ M) reflect a range believed to exceed the concentrations at expected efficacy doses in preclinical models.

10 The pulses protocol applied is described as follow: the holding potential (every 3 seconds) was stepped from -80 mV to a maximum value of +40 mV, starting with -40 mV, in eight increments of +10 mV, for a period of 1 second. The membrane potential was then returned to -55 mV, after each of these incremented steps, for 1 second and finally repolarized to -80 mV for 1 second.

15 The current density recorded were normalized against the baseline conditions and corrected for solvent effect and time-dependent current run-down using experimental design in test compound free conditions.

Inhibition curves were obtained for compounds and the concentrations which decreased 50% of the current density determined in the baseline conditions (IC_{50}) were determined. All compounds for which the IC_{50} value is above 10 μ M are not considered
20 to be potent inhibitors of the hERG channel whereas compounds with IC_{50} values below 1 μ M are considered potent hERG channel inhibitors.

When tested in the hERG inhibition assay, compounds of the invention were determined to have IC_{50} values as shown in Table 4.

Determination of plasma protein binding

25 The pharmacokinetic and pharmacodynamic properties of chemicals/drugs are largely a function of the reversible binding of chemicals to plasma or serum proteins. Generally, only the unbound or "free fraction" of a drug is available for diffusion or transport

across cell membranes, and for interaction with a pharmacological/toxicological target. Consequently, the extent of the plasma protein binding (PPB) of a compound influences its action as well as its distribution and elimination.

The determination of plasma protein binding (PPB) of a compound is enabled by
5 equilibrium dialysis, an accepted and standard method for reliable estimation of the non-bound drug fraction in plasma. RED (Rapid Equilibrium Dialysis) device insert is made of two side-by-side chambers separated by an O-ring-sealed vertical cylinder of dialysis membrane (MWCO ~8,000). Plasma containing drug (at 5 μ M or blood concentrations otherwise corresponding to efficacious doses, if known) is added to one
10 chamber while buffer is added to the second. After 4 hours incubation at 37°C under shaking, an aliquot is removed from each chamber and analyzed by a LC-MS/MS procedure enables the determination of both free and bound drug.

The percentages provided in Table 4 represent for the compounds of the invention the bound drug fraction to the plasma protein. The “free fraction” may be calculated as
15 100% - % rPPB (i.e. the complementary percentage of that disclosed in Table 4, corresponding to the drug concentration that is unbound and therefore available to engage biological target and elicit pharmacological activity).

TABLE 4

Cpd n°	Exposure (%rPPB)	CardioSafety (hERG IC ₅₀ , μ M)
1	67	42
2	47	32
3	42	66
4	40	70
5	22	70
6	53	45
7	26	70
8	29	70
9	22	70
10	30	70
11	24	50

12	20	NA
13	37	70
14	21	NA
15	20	70
16	36	70
17	24	46
18	23	70
19	51	NA
20	26	50
21	38	45
22	27	70
23	34	NA
24	48	61
25	19	NA
26	19	NA
27	24	70
28	12	NA
29	10	59
30	94	32
31	31	NA
32	25	NA
33	29	NA
34	24	NA
35	52	NA
36	60	NA
37	53	NA
38	76	NA
39	43	NA
40	24	NA
41	55	NA
42	16	NA
43	47	NA
44	31	NA
45	31	NA
45-1	27	NA
45-2	33	NA

NA: not available

In vivo assay to assess compound activity in rat (oral dosing)**Castrated male rat model to assess the effect of compound of invention on circulating levels of luteinizing hormone (LH)**

The effect of compounds of the invention to inhibit luteinizing hormone (LH) secretion
5 is determined by the following biological studies.

In humans and rodents, castration is well-precedented to permit heightened, persistent GnRH signaling and consequent elevation of circulating LH. Thus, a castrated rat model is used to provide a broad index for measurement of LH inhibition as a marker of test compound inhibition of the GnRH signaling pathway.

10 Castrated adult male Sprague-Dawley (SD) rats (150-175 g,) were purchased from Janvier (St Berthevin, France). All animals were housed 2 per cage in a temperature-controlled room ($22 \pm 2^{\circ}\text{C}$) and $50 \pm 5\%$ relative humidity with a 12 hour/12 hour light/dark cycles (lights off at 6h00 pm). The animals were allowed 3 weeks of postoperative recovery prior to study. Animals were handled on a daily basis. Standard
15 diet and tap water were provided *ad libitum*. Animal cage litters were changed once a week. On the study day, animals were acclimated to the procedure room for a period of one hour prior to the initiation of the experiment.

Compounds of the invention were formulated in 0.5% methyl cellulose.

After basal sampling (T0) a single dose of compounds of the invention or vehicle was
20 administrated orally to rats. Blood samples were then collected at several time points post dosing (45, 90, 150, 300 and 420 minutes). Blood samples were obtained via tail vein bleed, drawn into EDTA-containing tubes and centrifuged immediately. Plasma samples were collected and stored in a -80°C freezer until assayed. Serum LH levels were determined using radioimmunoassay kit from RIAZEN – Rat LH, Zentech (Liège,
25 Belgium). Baseline was defined as the initial basal blood sample.

When tested in the castrated male rat model described above, compounds n° **1, 2, 4, 5, 8, 9, 11, 13, 20** and **30** of the invention significantly suppressed circulating LH levels (statistically significant, $p < 0.05$) at a dose less than or equal to 30 mg/kg).

Effect of compounds of the invention on plasma testosterone in gonad intact male rats

- 5 The study was designed to evaluate the effect of compounds of the invention on testosterone circulating levels following oral administration at 3 mg/kg on SD gonad intact male rats.

Briefly the experimental methods used for this study were as follows:

- Two groups of non-fasted rats (male, Sprague-Dawley, 200 to 225g; $n = 4$ rats /group) with jugular vein cannulation, were dosed via a single oral administration of compounds of the invention at 3 mg/kg. The control group was dosed with the vehicle. Compounds of the invention were prepared in a dose formulation of pyrogen-free water with 0.5% methylcellulose. Blood samples were collected via the catheter implanted in the jugular vein at pre-determined intervals using EDTA-3K as anti-coagulant. Samples were chilled and rapidly processed by centrifugation to obtain corresponding plasma samples. Testosterone hormone levels were determined by RIA performed on plasma samples collected for all the groups at 5 minutes before administration (basal time), and at 45, 90, 150, 300, 480 minutes and 24 hours after dosing.

- When tested in the gonad intact male rats, compound n°**5** significantly suppressed plasma testosterone level over the test period as compared to the vehicle treated group (Figure 1).

Effect of compounds of the invention on prostate weight reduction in a Benign Prostatic Hyperplasia (BPH) rat model

- Briefly, adult male rats were injected daily for four weeks with testosterone to cause an enlargement of the prostate as per methods previously described in the literature (Scolnick et al., J. Andrology, 1994, 15(4), 287-297; Rick et al., J. Urol., 2012, 187, 1498-1504; see Figure 2, Ctrl Neg vs BHP). Rats were then treated daily for three

weeks with compounds of the invention. After 21 days treatment with compounds of the invention at 3, 10 or 30 mg/kg (q.d.; PO administration), the ratio of prostate to body weight (g prostate/100g of body weight) was evaluated as an indicator of BPH. Treated groups were compared to the BPH group (Testosterone-induced BPH group followed 21 days of vehicle administration) or to the Control group (Corn oil injection for the induction phase followed by vehicle treatment rather than test compound). Comparison between groups was made by using One-Way ANOVA followed by Dunnett's test for statistical analysis.

When tested in the Benign Prostatic Hyperplasia rat model, compound n°5, demonstrated a concentration-response to reduce prostate weight to normal levels (i.e. levels in rats not exposed to exogenous testosterone; Figure 2).

Effect of compounds of the invention on Estradiol circulating level in female rats

The aim of this study was to evaluate the effect of compounds of the invention on plasma estradiol levels following oral administration at 10 mg/kg (b.i.d.) for a period of 10 days in female rats.

Briefly the experimental methods used for this study were as follows:


Two groups of adult, female rats (Sprague-Dawley, ~320g) were treated in-phase with their individual estrous cycles. Thus, treatment was started in the proestrus phase (coincident with peak estradiol levels, as shown on Day 1 in Figure 3) and rats were dosed twice daily (~9h30 and 17h30) by oral administration either with a compound of the invention at 10 mg/kg or with the vehicle for the control group. Compounds of the invention were prepared in a dose formulation of pyrogen-free water with 0.5% methylcellulose. Estradiol levels were determined for all groups by ELISA performed on plasma samples derived from blood collections taken at 30 minutes before the daily, 9h30 test article administration on all days presented in Figure 3.

In vehicle-treated, adult female rats, estradiol peaks are observed every 4-5 days consistent with the anticipated duration of the rat estrous cycle. Treatment with compound n°5, significantly decreased estradiol levels over the time-course tracked

over two consecutive estrous cycles. This finding is most apparent in the proestrus phase (i.e. for vehicle group, on Day 5 and Day 9) where estradiol levels rise coincident with ovulation.

CLAIMS

1. A compound of Formula I:



wherein:

R^1 is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;

$R^{2'}$ is H or F;

R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^4 is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl;


R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl;

X^1 is N and X^2 is S or O; or X^1 is S and X^2 is N;

== represents a single or a double bond depending on X^1 and X^2 ;


* -- stands for the (*R*)-enantiomer or for the racemate of compound of Formula I.

2. The compound according to claim 1, having Formula I' and I'':



wherein R^1 , $R^{1'}$, R^2 , $R^{2'}$, R^3 , R^4 , R^5 , X^1 and X^2 are as defined in claim 1; and
 --- represents a single or a double bond depending on X^1 and X^2 .

3. The compound according to claim 1 or claim 2, having Formula Ia:



wherein:

R^1 is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;

$R^{2'}$ is H or F;


R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^4 is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

R^5 is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl;


* --- stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ia.

4. The compound according to any one of claims 1 to 3, selected from Formulae Ia' and Ia'':



wherein $\mathbf{R}^1, \mathbf{R}^{1'}, \mathbf{R}^2, \mathbf{R}^{2'}, \mathbf{R}^3, \mathbf{R}^4$ and $\mathbf{R}^5$ are as defined in claim 3.

5. The compound according to any one of claims 1 to 4, having Formula Ia-1:



wherein:


R³ is H, F, Cl, methyl, trifluoromethyl or nitrile;

R⁴ is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

R⁵ is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

*-- stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ia-1.

6. The compound according to any one of claims 1 to 5, having Formula Ia-2:



wherein:

R¹ is H, F or methyl;

R^{1'} is H;

R² is H, F, Cl or methoxy;


R^{2'} is H or F;

R³ is H, F, Cl, methyl, trifluoromethyl or nitrile;

R⁵ is methyl, ethyl, methoxymethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

* --- stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ia-2.

7. The compound according to any one of claims 1 to 6, having Formula Ia-3:



wherein:

R¹ is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;


$R^{2'}$ is H or F;

R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^4 is methyl, ethyl, *n*-propyl, hydroxyethyl, methoxyethyl, trifluoromethyl, difluoromethyl or fluoromethyl;

*-- stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ia-3.

8. The compound according to claim 1 or claim 2, having Formula Ib:




wherein:

R^3 is F;

R^5 is methyl, ethyl, trifluoromethyl, difluoromethyl, fluoromethyl, 1-fluoroethyl, 1,1-difluoroethyl or 2,2,2-trifluoroethyl;


*-- stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ib.

9. The compound according to any one of claims 1, 2 or 8, having Formula Ib':



wherein R^3 and R^5 are defined as in claim 8.

10. The compound according to claim 1 or claim 2, having Formula Ic:



wherein:

R^1 is H, F or methyl;

$R^{1'}$ is H;

R^2 is H, F, Cl or methoxy;

$R^{2'}$ is H or F;


R^3 is H, F, Cl, methyl, trifluoromethyl or nitrile;

R^4 is methyl, ethyl, *n*-propyl or hydroxyethyl;

R^5 is methyl, ethyl or trifluoromethyl;

*-- stands for the (*R*)-enantiomer or for the racemate of compound of Formula Ic.

11. The compound according to any one of claims 1, 2 or 10, having Formula Ic':



wherein:

R¹ is H, F or methyl;

R^{1'} is H;

R² is H, F, Cl or methoxy;

R^{2'} is H or F;





R³ is H, F, Cl, methyl, trifluoromethyl or nitrile;





R⁴ is methyl, ethyl, *n*-propyl or hydroxyethyl;





R⁵ is methyl, ethyl or trifluoromethyl.





12. The compound according to any one of claims 1 to 11, selected from the group consisting of:





1		<p>(<i>R</i>)-(3,4-dichlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone</p>
---	--	---





2		(R)-(3-(3-ethyl-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone
3		(R)-(4-chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
4		(R)-(4-chloro-3-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
5		(R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone





6		(R)-(3-chloro-4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
7		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(3,4,5-trifluorophenyl)methanone
8		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4-trifluorophenyl)methanone
9		(R)-(3,4-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone



10		(<i>R</i>)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4,5-tetrafluorophenyl)methanone
11		(<i>R</i>)-(4-fluorophenyl)(8-(2-hydroxyethyl)-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
12		(4-fluorophenyl)(8-(2-hydroxyethyl)-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
13		(<i>R</i>)-(3-(3-ethyl-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone





14		(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
15		(<i>R</i>)-(3-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
16		(<i>R</i>)-(3-chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
17		(<i>R</i>)-(3,5-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone





18		(<i>R</i>)-(2,4-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
19		(<i>R</i>)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(p-tolyl)methanone
20		(<i>R</i>)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(phenyl)methanone
21		(<i>R</i>)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-(trifluoromethyl)phenyl)methanone





22		(<i>R</i>)-(8-ethyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone
23		(8-ethyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone
24		(<i>R</i>)-(4-fluorophenyl)(3-(3-methyl-1,2,4-thiadiazol-5-yl)-8-propyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
25		(<i>R</i>)-(4-fluoro-3-methoxyphenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone


26		(R)-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(o-tolyl)methanone
27		(R)-(3-methoxyphenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
28		(R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
29		(R)-4-(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6,7,8-tetrahydro-[1,2,4]triazolo[4,3-a]pyrazine-7-carbonyl)benzonitrile

31		(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4,5-tetrafluorophenyl)methanone
32		(3,4-difluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone

33		(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(2,3,4-trifluorophenyl)methanone
34		(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(3,4,5-trifluorophenyl)methanone
35		(3-chloro-4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
36		(4-chloro-3-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone

37		(4-chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
38		(3,4-dichlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
39		(3-(3-ethyl-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone
40		(3-(3-ethyl-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone

41		(R)-(4-fluorophenyl)(8-methyl-3-(3-(trifluoromethyl)-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone
42		(R)-(3-(3-(difluoromethyl)-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone
43		(R)-(3-(3-(1,1-difluoroethyl)-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone
44		(R)-(4-fluorophenyl)(8-methyl-3-(3-(2,2,2-trifluoroethyl)-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone

45		((8R)-3-(3-(1-fluoroethyl)-1,2,4-oxadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone
----	---	--

13. The compound according to any one of claims 1 to 12, selected from the group consisting of:
- (R)-(3-(3-ethyl-1,2,4-thiadiazol-5-yl)-8-methyl-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)(4-fluorophenyl)methanone;
- (R)-(4-chlorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone;
- (R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone;
- (4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone;
- (R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-oxadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone.
14. The compound according to any one of claims 1 to 13, which is (R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone.
15. A pharmaceutical composition comprising a compound according to any one of claims 1 to 14 and at least one pharmaceutically acceptable carrier, diluent, excipient and/or adjuvant.

16. Medicament comprising a compound according to any one of claims 1 to 14.
17. A compound according to any one of claims 1 to 14 for use in treating and/or preventing depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma, airway hyperresponsiveness, bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual

dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian tumor or virilizing adrenal tumor), menorrhagia, or adenomyosis.


18. The compound for use according to claim 17, wherein the compound is (R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone.
19. A compound according to any one of claims 1 to 14 for use in treating and/or preventing hot flashes.
20. The compound for use according to claim 19, wherein the compound is (R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone.
21. A compound according to any one of claims 1 to 14 for use as a lowering agent of the circulating LH levels.
22. The compound for use according to claim 21, wherein the compound is (R)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-a]pyrazin-7(8H)-yl)methanone.
23. Use of a compound according to any one of claims 1 to 14 for the manufacture of a medicament to treat and/or prevent depression, anxiety, psychosis, schizophrenia, psychotic disorders, bipolar disorders, cognitive disorders, Parkinson's disease, Alzheimer's disease, attention deficit hyperactivity disorder (ADHD), pain, convulsion, obesity, inflammatory diseases including irritable bowel syndrome (IBS) and inflammatory bowel disorders, emesis, pre-eclampsia, airway related diseases including chronic obstructive pulmonary disease, asthma,

airway hyperresponsiveness, bronchoconstriction and cough, urinary incontinence, reproduction disorders, contraception and sex hormone-dependent diseases including but not limited to benign prostatic hyperplasia (BPH), prostatic hyperplasia, metastatic prostatic carcinoma, testicular cancer, breast cancer, ovarian cancer, androgen dependent acne, male pattern baldness, endometriosis, abnormal puberty, uterine fibrosis, uterine fibroid tumor, uterine leiomyoma, hormone-dependent cancers, hyperandrogenism, hirsutism, virilization, polycystic ovary syndrome (PCOS), premenstrual dysphoric disease (PMDD), HAIR-AN syndrome (hyperandrogenism, insulin resistance and acanthosis nigricans), ovarian hyperthecosis (HAIR-AN with hyperplasia of luteinized theca cells in ovarian stroma), other manifestations of high intraovarian androgen concentrations (e.g. follicular maturation arrest, atresia, anovulation, dysmenorrhea, dysfunctional uterine bleeding, infertility), androgen-producing tumor (virilizing ovarian tumor or virilizing adrenal tumor), menorrhagia, or adenomyosis.

24. The use according to claim 23, wherein the compound is (*R*)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-*a*]pyrazin-7(8H)-yl)methanone.
25. Use of a compound according to any one of claims 1 to 14 for the manufacture of a medicament to treat and/or prevent hot flashes.
26. The use according to claim 25, wherein the compound is (*R*)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-*a*]pyrazin-7(8H)-yl)methanone.
27. Use of a compound according to any one of claims 1 to 14 for the manufacture of a medicament to lower the circulating LH levels.
28. The use according to claim 27, wherein the compound is (*R*)-(4-fluorophenyl)(8-methyl-3-(3-methyl-1,2,4-thiadiazol-5-yl)-5,6-dihydro-[1,2,4]triazolo[4,3-*a*]pyrazin-7(8H)-yl)methanone.

29. Process of manufacturing a compound according to any one of claims 1 to 14, characterized in that it comprises the following steps:

a) reacting a compound of Formula (i)



(i)


wherein:

PG represents a suitable protecting group such as for example DMB, PMB, Boc, allyl, diphenyl-phosphinamide or 2-trimethylsilylethanesulfonyl;

R^{4'} is **R⁴** as defined in claim 1 or a reducible precursor of hydroxyethyl and consequently a further precursor of methoxyethyl;

* - - stands for the (*R*)-enantiomer or for the racemate;

with a compound of Formula (ii)



(ii)


wherein:

R^{5'} is **R⁵** as defined in claim 1, H or 1-((tert-butyldiphenylsilyl)oxy)ethyl;

X¹ and **X²** are as defined in claim 1;

== represents a single or a double bond depending on **X¹** and **X²**;


so as to obtain a compound of Formula (iii)



(iii)


wherein **PG**, **R^{4'}**, **R^{5'}**, **X¹** and **X²** are as defined above, *--- stands for the (*R*)-enantiomer or for the racemate and == represents a single or a double bond depending on **X¹** and **X²**;

b) deprotecting compound of Formula (iii) with a suitable deprotection agent to afford compound of Formula (iv)



wherein $\mathbf{R}^{4'}$, $\mathbf{R}^{5'}$, $\mathbf{X}^1$ and $\mathbf{X}^2$ are as defined above, *-- stands for the (*R*)-enantiomer or for the racemate and == represents a single or a double bound depending on $\mathbf{X}^1$ and $\mathbf{X}^2$;


- 5 c) when $\mathbf{R}^{5'}$ is H, introducing a trifluoromethyl or difluoromethyl group by direct C-H trifluoro- or difluoromethylation, leading to compound of Formula (v)



wherein $\mathbf{R}^{4'}$, $\mathbf{X}^1$ and $\mathbf{X}^2$ are as defined above and $\mathbf{R}^5$ is trifluoromethyl or difluoromethyl, *-- stands for the (*R*)-enantiomer or for the racemate and == represents a single or a double bound depending on $\mathbf{X}^1$ and $\mathbf{X}^2$;

10


- d) N-acylating compound of Formula (iv) wherein $\mathbf{R}^{5'}$ is not H or compound of Formula (v), with a compound of Formula (vi)



(vi)

wherein R^1 , $R^{1'}$, R^2 , $R^{2'}$ and R^3 are as defined in claim 1;

leading to compound of Formula (vii)



(vii)

wherein R^1 , $R^{1'}$, R^2 , $R^{2'}$, R^3 , $R^{4'}$, X^1 and X^2 are as defined above,

* -- stands for the (*R*)-enantiomer or for the racemate,

== represents a single or a double bond depending on X^1 and X^2 ; and

$R^{5''}$ is R^5 as defined in claim 1 or 1-((tert-butyldiphenylsilyl)oxy)ethyl;

e) optionally further conducting one or both of the two following steps e') and e''):

e') when $R^{4'}$ is a reducible precursor of hydroxyethyl and consequently a further precursor of methoxyethyl, a step of reduction optionally followed by methyl ether formation;

e'') when $R^{5''}$ is 1-((tert-butyldiphenylsilyl)oxy)ethyl, a step of alcohol deprotection and subsequent fluorination to form 1-fluoroethyl R^5 group; or a step of alcohol deprotection, followed by an oxidation step and a subsequent fluorination step to afford 1,1-difluoroethyl R^5 group;

to afford compound of Formula I according to any one of claims 1 to 14.