

## CCR6 COMPOUNDS

5

### CROSS-REFERENCES TO RELATED APPLICATIONS

[0001] This application claims the benefit of priority to U.S. Ser. No. 61/910,838, filed December 2, 2013, the disclosure of which is incorporated herein by reference in its entirety.

10

### STATEMENT AS TO RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH AND DEVELOPMENT

[0002] NOT APPLICABLE

15

### REFERENCE TO A "SEQUENCE LISTING," A TABLE, OR A COMPUTER PROGRAM LISTING APPENDIX SUBMITTED ON A COMPACT DISK

[0003] NOT APPLICABLE

20

### BACKGROUND OF THE INVENTION

[0004] Chemokines are chemotactic cytokines that are released by a wide variety of cells to attract macrophages, T cells, eosinophils, basophils and neutrophils to sites of inflammation (reviewed in Schall, *Cytokine*, **3**:165-183 (1991), Schall, et al., *Curr Opin. Immunol.* **6**:865-873 (1994) and Murphy, *Rev. Immun.*, **12**:593-633 (1994)). In addition to stimulating chemotaxis, other changes can be selectively induced by chemokines in responsive cells, including changes in cell shape, transient rises in the concentration of intracellular free calcium ions ( $[Ca^{2+}]$ ), granule exocytosis, integrin upregulation, formation of bioactive lipids (e.g., leukotrienes) and respiratory burst, associated with leukocyte activation. Thus, the chemokines are early triggers of the inflammatory response, causing inflammatory mediator release, chemotaxis and extravasation to sites of infection or inflammation.

[0005] There are two main classes of chemokines, CXC (alpha) and CC (beta), depending on whether the first two cysteines are separated by a single amino acid (C-X-C) or are adjacent (C-C). The alpha-chemokines, such as interleukin-8 (IL-8), neutrophil-activating protein-2 (NAP-2) and melanoma growth stimulatory activity protein (MGSA) are chemotactic primarily for neutrophils, whereas beta-chemokines, such as RANTES, MIP-1a, MIP-1b, monocyte chemotactic protein-1 (MCP-1), MCP-2, MCP-3 and eotaxin are chemotactic for macrophages, T-cells, eosinophils and basophils (Deng, et al., *Nature*, **381**:661-666 (1996)). The chemokines bind specific cell-surface receptors belonging to the family of G-protein-coupled seven-transmembrane-domain proteins (reviewed in Horuk, *Trends Pharm. Sci.*, **15**:159-165 (1994)) which are termed "chemokine receptors."

[0006] On binding their cognate ligands, chemokine receptors transduce an intracellular signal through the associated trimeric G protein, resulting in a rapid increase in intracellular calcium concentration. There are at least eleven human chemokine receptors that bind or respond to beta-chemokines and at least seven human chemokine receptors that bind to the alpha chemokines. Additionally CX3CR1 (fractalkine receptor) can bind to the fractalkine chemokine, which is distinguished by a series of three amino acids between the first two cysteines. Chemokine receptors, have been implicated as being important mediators of inflammatory and immunoregulatory disorders and diseases, including asthma and allergic diseases, as well as autoimmune pathologies such as rheumatoid arthritis and atherosclerosis.

[0007] CCR6 is known to be expressed primarily in B cells, IL17 secreting T cells, regulatory T cells and dendritic cells and shows strong binding to its cognate ligand CCL20 (MIP-3 $\alpha$ ). It is expressed on approximately 30-60% of adult peripheral blood effector/memory CD4<sup>+</sup> T cells. CCR6 is involved in leukocyte homing to inflamed tissue, particularly the skin and lungs and is co-expressed on almost all T cells that have a skin homing phenotype, the CLA<sup>+</sup> T cells. Thus CCR6 may be an important player in skin pathologies in which leukocytes participate.


[0008] CCR6 expression has been linked to psoriasis in the following manner. In humans, a large majority of skin-homing CD4 T cells in the peripheral blood express CCR6 with a greater degree of CCL20-mediated chemotaxis occurring in T cells isolated from psoriatic patients (Homey, et. al., *Jl*, 2000). IL17 secreting cells are central agents in several inflammatory diseases. T cells, such as  $\gamma\delta$  T cells and TH17 T cells produce IL17 after activation. The pathogenic effects of IL17 have been associated with human diseases such as

rheumatoid arthritis (Patel DD et. al., *Ann Rheum Dis* 2013), multiple sclerosis (Zepp J, Wu L, and X Li *Trends Immunol* 2011), and psoriasis (Martin DA et. al., *J Invest Dermatol* 2012). Evidence strongly linking IL17 with psoriasis include gene wide association studies that show strong association between psoriasis and genes upstream (IL-23) or downstream (NF $\kappa$ b) of IL17 signaling pathways as well as efficacy in targeting IL17 in a clinical setting (Martin DA et. al., *J. Invest Dermat.* 2012; Papp et. al., *NEJM*, 2012; Papp et. al., *NEJM*, 2012). In addition to enhanced CCL20-mediated chemotaxis, CCR6<sup>+</sup> T cells isolated from psoriatic patients preferentially secrete IL-17A, IL22, and TNF $\alpha$  when compared to healthy controls (Kagami, et. al., *J. Invest. Dermatol.*, 2010). Lastly, ccl20 mRNA was up-regulated in lesional psoriatic skin samples (Homey, et. al., *Jl*, 2000; Dieu-Nosjean, et. al., *JEM*, 2000). In mice, CCR6 knock-out mice were protected from IL-23 driven psoriasis. Thus, a multitude of evidence in both mice and men suggest a protective role for CCR6 blockade in psoriasis and psoriasis-like models.

**[0009]** In view of the clinical importance of CCR6, the identification of compounds that modulate CCR6 function represent an attractive avenue into the development of new therapeutic agents. Such compounds and methods for their use are provided herein.

## BRIEF SUMMARY

**[0010]** Described herein are compounds having formula (I):




wherein the ring vertices a, b, c and d; Ar; A; R<sup>2</sup>, R<sup>3</sup> and the subscripts m and n have the meanings provided in the Detailed Description below. The compounds have utility in the treatment of diseases or conditions modulated at least in part by CCR6.

**[0011]** Pharmaceutical compositions of the compounds of formula (I) are also provided.

**[0012]** Further provided in the present disclosure preparative methods for the synthesis of compounds of formula (I), as well as selected intermediates useful in the preparation.

**[0012a]** In one aspect of the present disclosure, there is provided a compound having formula (I):



or a pharmaceutically acceptable salt, hydrate, solvate, or rotamer thereof,

5 wherein

A is a carboxylic acid moiety or a carboxylic acid isostere;

ring vertices a, b, c and d are independently selected from CH and  $C(R^1)$ ;

each  $R^1$  is independently selected from the group consisting of halogen, CN,  $-SF_5$ ,  $C_{1-8}$

alkyl,  $C_{3-8}$  cycloalkyl,  $C_{2-8}$  alkenyl,  $C_{2-8}$  alkynyl,  $C_{1-8}$

10 haloalkyl,  $-OR^a$ ,  $-SR^a$ ,  $-COR^a$ ,  $-NR^aR^b$ , and 5- or 6-membered heteroaryl, wherein

the alkyl portions of  $R^1$  are optionally further substituted with 1-3  $R^a$ ; and

optionally, adjacent  $R^1$  members are connected to form an additional 5- or 6-membered ring which is saturated or unsaturated having ring vertices selected

from C, O, S and N, wherein the additional 5- or 6-membered ring is optionally

15 substituted with one or two members selected from the group consisting of

halogen, hydroxyl,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy and  $C_{1-4}$  haloalkyl;

the subscript n is an integer of from 0 to 3;

each  $R^3$  is independently selected from the group consisting of halogen, CN,  $C_{1-8}$  alkyl,

20  $C_{3-8}$  cycloalkyl,  $C_{2-8}$  alkenyl,  $C_{2-8}$  alkynyl,  $C_{1-8}$  haloalkyl, aryl,  $-OR^a$ ,  $-NR^aR^b$  and  $-N(R^a)-C_1-C_4$  alkylene- $OR^b$ , and wherein the alkyl or aryl portions of  $R^2$  and

$R^3$  are optionally further substituted with 1-3  $R^a$ ;

Ar is a benzene or pyridine ring that is optionally substituted with from 1 to 5  $R^4$

substituents independently selected from the group consisting of halogen,

25 CN,  $-SF_5$ ,  $C_{1-8}$  alkyl,  $C_{3-8}$  cycloalkyl,  $C_{2-8}$  alkenyl,  $C_{2-8}$  alkynyl,  $C_{1-8}$  haloalkyl,  $C_{1-8}$  hydroxyalkyl,  $-OR^a$ ,  $-NR^aR^b$ ,

and wherein the alkyl and cycloalkyl, portions of R<sup>4</sup> are optionally further substituted with 1-3 R<sup>a</sup>;

each R<sup>a</sup> and R<sup>b</sup> is independently selected from the group consisting of hydrogen,

hydroxyl, halogen, cyano, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>3-6</sub> cycloalkyl,

amino, C<sub>1-8</sub> alkylamino, and di C<sub>1-8</sub> alkylamino, or when attached to a nitrogen

atom are optionally combined to form a 4- to 7-membered saturated ring, which is optionally substituted with oxo.

**[0012b]** In another aspect of the present disclosure, there is provided a pharmaceutical composition comprising a compound disclosed herein, or a pharmaceutically acceptable salt, hydrate, solvate, or rotamer thereof, with a pharmaceutically acceptable excipient.

**[0012c]** In a further aspect of the present disclosure, there is provided a use of a compound disclosed herein, or a pharmaceutically acceptable salt, hydrate, solvate, or rotamer thereof, in the manufacture of a medicament for treating a disease or condition modulated at least in part by CCR6, wherein said disease or condition is an inflammatory disease or condition.

**[0012d]** In another aspect of the present disclosure, there is provided a compound disclosed herein, or a pharmaceutically acceptable salt, hydrate, solvate, or rotamer thereof, for use in treating a disease or condition modulated at least in part by CCR6.

## BRIEF DESCRIPTION OF THE DRAWINGS

[0013] Figures 1A-1H provide structures and binding data for compounds prepared and evaluated by the methods described herein. Activity is shown as +,  $20000 \text{ nM} \geq \text{IC}_{50} \geq 500 \text{ nM}$ ; ++,  $500 \text{ nM} > \text{IC}_{50} \geq 100 \text{ nM}$ ; +++,  $100 \text{ nM} > \text{IC}_{50}$ .

[0014] Figures 2A-2C provide structures and migration data for compounds prepared and evaluated by the methods described herein. Activity is shown as +,  $20000 \text{ nM} \geq \text{IC}_{50} \geq 500 \text{ nM}$ ; ++,  $500 \text{ nM} > \text{IC}_{50} \geq 100 \text{ nM}$ ; +++,  $100 \text{ nM} > \text{IC}_{50}$ .

## DETAILED DESCRIPTION OF THE INVENTION

### I. ABBREVIATION AND DEFINITIONS

[0015] The term "alkyl", by itself or as part of another substituent, means, unless otherwise stated, a straight or branched chain hydrocarbon radical, having the number of carbon atoms designated (*i.e.*  $\text{C}_{1-8}$  means one to eight carbons). Examples of alkyl groups include methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, isobutyl, sec-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, and the like. The term "alkenyl" refers to an unsaturated alkyl group having one or more double bonds. Similarly, the term "alkynyl" refers to an unsaturated alkyl group having one or more triple bonds. Examples of such unsaturated alkyl groups include vinyl, 2-propenyl, crotyl, 2-isopentenyl, 2-(butadienyl), 2,4-pentadienyl, 3-(1,4-pentadienyl), ethynyl, 1- and 3-propynyl, 3-butylnyl, and the higher homologs and isomers. The term "cycloalkyl" refers to hydrocarbon rings having the indicated number of ring atoms (*e.g.*,  $\text{C}_{3-6}$ cycloalkyl) and being fully saturated or having no more than one double bond between ring vertices. "Cycloalkyl" is also meant to refer to bicyclic and polycyclic hydrocarbon rings such as, for example, bicyclo[2.2.1]heptane, bicyclo[2.2.2]octane, etc. The term "heterocycloalkane" or "heterocycloalkyl" refers to a cycloalkyl group that contain from one to five heteroatoms selected from N, O, and S, wherein the nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. The heterocycloalkane may be a monocyclic, a bicyclic or a polycyclic ring system. Non limiting examples of heterocycloalkane groups include pyrrolidine, imidazolidine, pyrazolidine, butyrolactam, valerolactam, imidazolidinone, hydantoin, dioxolane, phthalimide, piperidine, 1,4-dioxane, morpholine, thiomorpholine, thiomorpholine-S-oxide, thiomorpholine-S,S-oxide, piperazine,

pyran, pyridone, 3-pyrroline, thiopyran, pyrone, tetrahydrofuran, tetrahydrothiophene, quinuclidine, and the like. A heterocycloalkane group can be attached to the remainder of the molecule through a ring carbon or a heteroatom.

[0016] The term "alkylene" by itself or as part of another substituent means a divalent radical derived from an alkane, as exemplified by  $-\text{CH}_2\text{CH}_2\text{CH}_2\text{CH}_2-$ . Typically, an alkyl (or alkylene) group will have from 1 to 24 carbon atoms, with those groups having 10 or fewer carbon atoms being preferred in the present invention. A "lower alkyl" or "lower alkylene" is a shorter chain alkyl or alkylene group, generally having four or fewer carbon atoms. Similarly, "alkenylene" and "alkynylene" refer to the unsaturated forms of "alkylene" having double or triple bonds, respectively.

[0017] As used herein, a wavy line, " $\sim$ ", that intersects a single, double or triple bond in any chemical structure depicted herein, represent the point attachment of the single, double, or triple bond to the remainder of the molecule.

[0018] The terms "alkoxy," "alkylamino" and "alkylthio" (or thioalkoxy) are used in their conventional sense, and refer to those alkyl groups attached to the remainder of the molecule via an oxygen atom, an amino group, or a sulfur atom, respectively. Additionally, for dialkylamino groups, the alkyl portions can be the same or different and can also be combined to form a 3-7 membered ring with the nitrogen atom to which each is attached. Accordingly, a group represented as dialkylamino or  $-\text{NR}^a\text{R}^b$  is meant to include piperidinyl, pyrrolidinyl, morpholinyl, azetidiny and the like.

[0019] The term "di-( $\text{C}_{1-4}$  alkyl)amino- $\text{C}_{1-4}$  alkyl" refers to an amino group bearing two  $\text{C}_{1-4}$  alkyl groups that can be the same or different (e.g., methyl, ethyl, propyl, isopropyl, n-butyl, sec-butyl, isobutyl and tert-butyl) and which is attached to the remainder of the molecule through a  $\text{C}_{1-4}$  alkyl group (a one to four carbon alkylene linking group). Examples of di-( $\text{C}_{1-4}$  alkyl)amino- $\text{C}_{1-4}$  alkyl groups include dimethylaminomethyl, 2-(ethyl(methyl)amino)ethyl, 3-(dimethylamino)butyl, and the like.

[0020] The terms "halo" or "halogen," by themselves or as part of another substituent, mean, unless otherwise stated, a fluorine, chlorine, bromine, or iodine atom. Additionally, terms such as "haloalkyl," are meant to include monohaloalkyl and polyhaloalkyl. For example, the term " $\text{C}_{1-4}$  haloalkyl" is meant to include trifluoromethyl, 2,2,2-trifluoroethyl, 4-chlorobutyl, 3-bromopropyl, and the like.

[0021] The term "aryl" means, unless otherwise stated, a polyunsaturated, typically aromatic, hydrocarbon group which can be a single ring or multiple rings (up to three rings) which are fused together or linked covalently. The term "heteroaryl" refers to aryl groups (or rings) that contain from one to five heteroatoms selected from N, O, and S, wherein the nitrogen and sulfur atoms are optionally oxidized, and the nitrogen atom(s) are optionally quaternized. A heteroaryl group can be attached to the remainder of the molecule through a heteroatom. Non-limiting examples of aryl groups include phenyl, naphthyl and biphenyl, while non-limiting examples of heteroaryl groups include pyridyl, pyridazinyl, pyrazinyl, pyrimidinyl, triazinyl, quinoliny, quinoxaliny, quinazoliny, cinnoliny, phthalazinyl, benzotriazinyl, purinyl, benzimidazolyl, benzopyrazolyl, benzotriazolyl, benzisoxazolyl, isobenzofuryl, isoindolyl, indoliziny, benzotriazinyl, thienopyridiny, thienopyrimidinyl, pyrazolopyrimidinyl, imidazopyridines, benzothiazolyl, benzofuranyl, benzothienyl, indolyl, quinolyl, isoquinolyl, isothiazolyl, pyrazolyl, indazolyl, pteridinyl, imidazolyl, triazolyl, tetrazolyl, oxazolyl, isoxazolyl, thiadiazolyl, pyrrolyl, thiazolyl, furyl, thienyl and the like. Substituents for each of the above noted aryl and heteroaryl ring systems are selected from the group of acceptable substituents described below.

[0022] The term "arylalkyl" is meant to include those radicals in which an aryl group is attached to an alkyl group (*e.g.*, benzyl, phenethyl, and the like). Similarly, the term "heteroaryl-alkyl" is meant to include those radicals in which a heteroaryl group is attached to an alkyl group (*e.g.*, pyridylmethyl, thiazolylethyl, and the like).

[0023] As used herein, the term "heteroatom" is meant to include oxygen (O), nitrogen (N), sulfur (S) and silicon (Si).

[0024] The term "pharmaceutically acceptable salts" is meant to include salts of the active compounds which are prepared with relatively nontoxic acids or bases, depending on the particular substituents found on the compounds described herein. When compounds of the present invention contain relatively acidic functionalities, base addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired base, either neat or in a suitable inert solvent. Examples of salts derived from pharmaceutically-acceptable inorganic bases include aluminum, ammonium, calcium, copper, ferric, ferrous, lithium, magnesium, manganic, manganous, potassium, sodium, zinc and the like. Salts derived from pharmaceutically-acceptable organic bases include salts of primary, secondary and tertiary amines, including substituted amines, cyclic amines, naturally-

occurring amines and the like, such as arginine, betaine, caffeine, choline, N,N'-dibenzylethylenediamine, diethylamine, 2-diethylaminoethanol, 2-dimethylaminoethanol, ethanolamine, ethylenediamine, N-ethylmorpholine, N-ethylpiperidine, glucamine, glucosamine, histidine, hydrabamine, isopropylamine, lysine, methylglucamine, morpholine, piperazine, piperidine, polyamine resins, procaine, purines, theobromine, triethylamine, trimethylamine, tripropylamine, tromethamine and the like. When compounds of the present invention contain relatively basic functionalities, acid addition salts can be obtained by contacting the neutral form of such compounds with a sufficient amount of the desired acid, either neat or in a suitable inert solvent. Examples of pharmaceutically acceptable acid addition salts include those derived from inorganic acids like hydrochloric, hydrobromic, nitric, carbonic, monohydrogencarbonic, phosphoric, monohydrogenphosphoric, dihydrogenphosphoric, sulfuric, monohydrogensulfuric, hydriodic, or phosphorous acids and the like, as well as the salts derived from relatively nontoxic organic acids like acetic, propionic, isobutyric, malonic, benzoic, succinic, suberic, fumaric, mandelic, phthalic, benzenesulfonic, p-tolylsulfonic, citric, tartaric, methanesulfonic, and the like. Also included are salts of amino acids such as arginate and the like, and salts of organic acids like glucuronic or galactunoric acids and the like (see, for example, Berge, S.M., et al, "Pharmaceutical Salts", *Journal of Pharmaceutical Science*, **1977**, 66, 1-19). Certain specific compounds of the present invention contain both basic and acidic functionalities that allow the compounds to be converted into either base or acid addition salts.

**[0025]** The neutral forms of the compounds may be regenerated by contacting the salt with a base or acid and isolating the parent compound in the conventional manner. The parent form of the compound differs from the various salt forms in certain physical properties, such as solubility in polar solvents, but otherwise the salts are equivalent to the parent form of the compound for the purposes of the present invention.

**[0026]** In addition to salt forms, the present invention provides compounds which are in a prodrug form. Prodrugs of the compounds described herein are those compounds that readily undergo chemical changes under physiological conditions to provide the compounds of the present invention. Additionally, prodrugs can be converted to the compounds of the present invention by chemical or biochemical methods in an *ex vivo* environment. For example, prodrugs can be slowly converted to the compounds of the present invention when placed in a transdermal patch reservoir with a suitable enzyme or chemical reagent.

[0027] Certain compounds of the present invention can exist in unsolvated forms as well as solvated forms, including hydrated forms. In general, the solvated forms are equivalent to unsolvated forms and are intended to be encompassed within the scope of the present invention. Certain compounds of the present invention may exist in multiple crystalline or amorphous forms. In general, all physical forms are equivalent for the uses contemplated by the present invention and are intended to be within the scope of the present invention.

[0028] Certain compounds of the present invention possess asymmetric carbon atoms (optical centers) or double bonds; the racemates, diastereomers, geometric isomers, regioisomers and individual isomers (e.g., separate enantiomers) are all intended to be encompassed within the scope of the present invention. The compounds of the present invention may also contain unnatural proportions of atomic isotopes at one or more of the atoms that constitute such compounds. Unnatural proportions of an isotope may be defined as ranging from the amount found in nature to an amount consisting of 100% of the atom in question. For example, the compounds may incorporate radioactive isotopes, such as for example tritium ( $^3\text{H}$ ), iodine-125 ( $^{125}\text{I}$ ) or carbon-14 ( $^{14}\text{C}$ ), or non-radioactive isotopes, such as deuterium ( $^2\text{H}$ ) or carbon-13 ( $^{13}\text{C}$ ). Such isotopic variations can provide additional utilities to those described elsewhere with this application. For instance, isotopic variants of the compounds of the invention may find additional utility, including but not limited to, as diagnostic and/or imaging reagents, or as cytotoxic/radiotoxic therapeutic agents. Additionally, isotopic variants of the compounds of the invention can have altered pharmacokinetic and pharmacodynamic characteristics which can contribute to enhanced safety, tolerability or efficacy during treatment. All isotopic variations of the compounds of the present invention, whether radioactive or not, are intended to be encompassed within the scope of the present invention.

[0029] The term "and acid isosteres" means, unless otherwise stated, a group which can replace a carboxylic acid, having an acidic functionality and steric and electronic characteristics that provide a level of activity (or other compound characteristic such as solubility) similar to a carboxylic acid. Representative acid isosteres include, hydroxamic acids, sulfonic acids, sulfinic acids, sulfonamides, acyl-sulfonamides, phosphonic acids, phosphinic acids, phosphoric acids, tetrazole, and oxo-oxadiazoles.

[0030] Compounds of the invention having formula I can exist in different isomeric forms. As used herein, the terms *cis* or *trans* are used in their conventional sense in the chemical


arts, *i.e.*, referring to the position of the substituents to one another relative to a reference plane, *e.g.*, a double bond, or a ring system, such as a decalin-type ring system or a hydroquinolone ring system: in the *cis* isomer, the substituents are on the same side of the reference plane, in the *trans* isomer the substituents are on opposite sides. Additionally,  
 5 different conformers are contemplated by the present invention, as well as distinct rotamers. Conformers are conformational isomers that can differ by rotations about one or more  $\sigma$  bonds. Rotamers are conformers that differ by rotation about only a single  $\sigma$  bond.

## II. GENERAL

[0031] The present invention derives from the discovery that compounds of formula I act as potent antagonists of the CCR6 receptor. The compounds have *in vivo* anti-inflammatory activity and have superior pharmacokinetic properties. Accordingly, the compounds provided herein are useful in pharmaceutical compositions, methods for the treatment of CCR6-mediated diseases, and as controls in assays for the identification of competitive CCR6 antagonists.

## III. COMPOUNDS

[0032] In one aspect, the present invention provides for a compound of Formula I:



or a pharmaceutically acceptable salt, hydrate, solvate, N-oxide or rotamer thereof. In Formula (I), the letter  $A$  represents a carboxylic acid moiety or a carboxylic acid isostere; ring vertices  $a, b, c$  and  $d$  are independently selected from the group consisting of  $N$ ,  $CH$  and  $C(R^1)$ ; each  $R^1$  is independently selected from the group consisting of halogen,  $CN$ ,  $-SF_5$ ,  $C_{1-8}$  alkyl,  $C_{3-8}$  cycloalkyl,  $C_{2-8}$  alkenyl,  $C_{2-8}$  alkynyl,  $C_{1-8}$  haloalkyl,  $-OR^a$ ,  $-SR^a$ ,  $-COR^a$ ,  $-NR^aR^b$ ,  
 15 and 5- or 6-membered heteroaryl, wherein the alkyl portions of  $R^1$  are optionally further substituted with 1-3  $R^a$ ; and optionally, adjacent  $R^1$  members are connected to form an additional 5- or 6-membered ring which is saturated or unsaturated having ring vertices selected from  $C$ ,  $O$ ,  $S$  and  $N$ , wherein the additional 5- or 6-membered ring is optionally

substituted with one or two members selected from the group consisting of halogen, hydroxyl, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy and C<sub>1-4</sub> haloalkyl; the subscript m is an integer of from 0 to 2; the subscript n is an integer of from 0 to 3; each R<sup>2</sup> and R<sup>3</sup> is independently selected from the group consisting of halogen, CN, C<sub>1-8</sub> alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>2-8</sub> alkenyl, C<sub>2-8</sub> alkynyl, C<sub>1-8</sub> haloalkyl, aryl, -OR<sup>a</sup>, -NR<sup>a</sup>R<sup>b</sup> and -N(R<sup>a</sup>)-C<sub>1</sub>-C<sub>4</sub> alkylene-OR<sup>b</sup>, and wherein the alkyl or aryl portions of R<sup>2</sup> and R<sup>3</sup> are optionally further substituted with 1-3 R<sup>a</sup>; Ar is a 5- or 6-membered aromatic or heteroaromatic ring that is optionally substituted with from 1 to 5 R<sup>4</sup> substituents independently selected from the group consisting of halogen, CN, -SF<sub>5</sub>, C<sub>1-8</sub> alkyl, C<sub>3-8</sub> cycloalkyl, C<sub>2-8</sub> alkenyl, C<sub>2-8</sub> alkynyl, C<sub>1-8</sub> haloalkyl, C<sub>1-8</sub> hydroxyalkyl, -OR<sup>a</sup>, -NR<sup>a</sup>R<sup>b</sup>, 5- or 6-membered heteroaryl, and 3-, 4-, 5- or 6-membered heterocycloalkane wherein the heteroatoms present as ring vertices of the heteroaryl and heterocycloalkane rings are selected from N, O and S, and wherein the alkyl, cycloalkyl, aryl, heteroaryl and heterocycloalkane portions of R<sup>4</sup> are optionally further substituted with 1-3 R<sup>a</sup>; each R<sup>a</sup> and R<sup>b</sup> is independently selected from the group consisting of hydrogen, hydroxyl, halogen, cyano, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, C<sub>1-4</sub> haloalkyl, C<sub>3-6</sub> cycloalkyl, amino, C<sub>1-8</sub> alkylamino, and di C<sub>1-8</sub> alkylamino, or when attached to a nitrogen atom are optionally combined to form a 4- to 7-membered saturated ring, which is optionally substituted with oxo.

**[0033]** In some selected embodiments, the compounds of formula (I) are those wherein ring vertices a, b, c and d are each independently selected from the group consisting of CH and C(R<sup>1</sup>). In other selected embodiments, the compounds of formula (I) are those wherein ring vertex d is CH. In still other selected embodiments, the compounds of formula (I) are those wherein ring vertices c and d are each CH. In yet other selected embodiments, the compounds of formula (I) are those wherein ring vertices a, b, and c are each C(R<sup>1</sup>). In other selected embodiments, the compounds of formula (I) are those wherein ring vertices a and b are each C(R<sup>1</sup>).


**[0034]** In the compounds of formula (I), the letter A represents a carboxylic acid or a carboxylic acid isostere (or a suitable salt thereof). In certain selected embodiments of formula (I), the letter A represents a member selected from the group consisting of -CO<sub>2</sub>H, -SO<sub>3</sub>H, -PO<sub>3</sub>H<sub>2</sub> and a tetrazole. In other selected embodiments, the compounds of formula (I) are those wherein A is a carboxylic acid (-CO<sub>2</sub>H) or a tetrazole. In some embodiments A is a tetrazole. In some embodiments, A is -CO<sub>2</sub>H or a salt thereof.

[0035] Turning next to the substituents on the quinoline moiety, in certain selected embodiments m and n are both 0. In other selected embodiments, the compounds of formula (I) are those wherein m and n are independently selected from 0 and 1. For those embodiments wherein R<sup>2</sup> and R<sup>3</sup> are present, selected embodiments are those wherein each is independently selected from the group consisting of halogen, C<sub>1-8</sub> alkyl, -OR<sup>a</sup>, -NR<sup>a</sup>R<sup>b</sup> and -N(R<sup>a</sup>)-C<sub>1</sub>-C<sub>4</sub> alkylene-OR<sup>b</sup>, and wherein the alkyl portions of R<sup>2</sup> and R<sup>3</sup> are optionally further substituted with 1-3 R<sup>a</sup>.

[0036] In other selected embodiments for the compounds of formula (I), and for each of noted embodiments above, Ar is a 6-membered aromatic or heteroaromatic ring selected from the group consisting of benzene, pyridine and pyrimidine, each of which is optionally substituted with from 1-3 R<sup>4</sup> substituents. In still other selected embodiments for the compounds of formula (I), and for each of noted embodiments above, Ar is selected from the group consisting of benzene and pyridine, each of which is optionally substituted with from 1-2 R<sup>4</sup> substituents.


[0037] In one selected group of embodiments of formula (I), Ar is a benzene ring which is optionally substituted with from 1-3 R<sup>4</sup> substituents; A is -CO<sub>2</sub>H; d is CH; and each of a, b and c is independently selected from the group consisting of CH and C(R<sup>1</sup>). In this group of embodiments, further selected embodiments are those wherein m and n are each independently 0 or 1. Still further selected embodiments are those wherein m and n are each 0.

[0038] In other selected embodiments for the compounds of formula (I), and for each of noted embodiments above, the ring having a, b, c and d as ring vertices is selected from the group consisting of:












wherein the wavy line labeled y indicates the point of attachment to the NHS(O)<sub>2</sub> portion of the compound and the wavy line labeled z indicates the point of attachment to the quinoline ring.

- 5 [0039] In other selected embodiments for the compounds of formula (I), and for each of noted embodiments above, Ar is selected from the group consisting of:




wherein the wavy line labeled w indicates the point of attachment to the S(O)<sub>2</sub> moiety.


- 10 [0040] In still other selected embodiments, the compounds of formula (I) are selected from the group consisting of:



and



[0041] In still other selected embodiments, the compounds of formula (I) are selected from the group consisting of:



5 or a pharmaceutically acceptable salt, hydrate, solvate, N-oxide or rotamer thereof.

#### Preparation of Compounds

[0042] The schemes in the Examples below provide certain synthetic routes that can be followed to access certain compounds of the present invention. Other routes or modification of the routes presented below would be readily apparent to a skilled artisan and are within the scope of the present invention.

#### **IV. PHARMACEUTICAL COMPOSITIONS**

[0043] In addition the compounds provided above, the compositions for modulating CCR6, activity in humans and animals will typically contain a pharmaceutical carrier or diluent.

[0044] The term "composition" as used herein is intended to encompass a product comprising the specified ingredients in the specified amounts, as well as any product which results, directly or indirectly, from combination of the specified ingredients in the specified amounts. By "pharmaceutically acceptable" it is meant the carrier, diluent or excipient must be compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

[0045] The pharmaceutical compositions for the administration of the compounds of this invention may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy and drug delivery. All methods include the step of bringing the active ingredient into association with the carrier which constitutes one or more accessory ingredients. In general, the pharmaceutical compositions are prepared by

uniformly and intimately bringing the active ingredient into association with a liquid carrier or a finely divided solid carrier or both, and then, if necessary, shaping the product into the desired formulation. In the pharmaceutical composition the active object compound is included in an amount sufficient to produce the desired effect upon the process or condition of diseases.

[0046] The pharmaceutical compositions containing the active ingredient may be in a form suitable for oral use, for example, as tablets, troches, lozenges, aqueous or oily suspensions, dispersible powders or granules, emulsions and self emulsifications as described in U.S. Patent No. 6,451,339, hard or soft capsules, syrups, elixirs, solutions, buccal patch, oral gel, chewing gum, chewable tablets, effervescent powder and effervescent tablets. Compositions intended for oral use may be prepared according to any method known to the art for the manufacture of pharmaceutical compositions and such compositions may contain one or more agents selected from the group consisting of sweetening agents, flavoring agents, coloring agents, antioxidants and preserving agents in order to provide pharmaceutically elegant and palatable preparations. Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be for example, inert diluents, such as cellulose, silicon dioxide, aluminum oxide, calcium carbonate, sodium carbonate, glucose, mannitol, sorbitol, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; binding agents, for example PVP, cellulose, PEG, starch, gelatin or acacia, and lubricating agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated, enterically or otherwise, by known techniques to delay disintegration and absorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. They may also be coated by the techniques described in the U.S. Pat. Nos. 4,256,108; 4,166,452; and 4,265,874 to form osmotic therapeutic tablets for control release.

[0047] Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin, or olive oil. Additionally, emulsions can be prepared with a non-water miscible ingredient such as oils and stabilized with surfactants such as mono-diglycerides, PEG esters and the like.

[0048] Aqueous suspensions contain the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxy-propylmethylcellulose, sodium alginate, polyvinyl-pyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and a hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or n-propyl, p-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

[0049] Oily suspensions may be formulated by suspending the active ingredient in a vegetable oil, for example arachis oil, olive oil, sesame oil or coconut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above, and flavoring agents may be added to provide a palatable oral preparation. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

[0050] Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example sweetening, flavoring and coloring agents, may also be present.

[0051] The pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oily phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation

products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

[0052] Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents. Oral solutions can be prepared in combination with, for example, cyclodextrin, PEG and surfactants.

[0053] The pharmaceutical compositions may be in the form of a sterile injectable aqueous or oleagenous suspension. This suspension may be formulated according to the known art using those suitable dispersing or wetting agents and suspending agents which have been mentioned above. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example as a solution in 1,3-butane diol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil may be employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

[0054] The compounds of the present invention may also be administered in the form of suppositories for rectal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal temperature and will therefore melt in the rectum to release the drug. Such materials include cocoa butter and polyethylene glycols.

Additionally, the compounds can be administered via ocular delivery by means of solutions or ointments. Still further, transdermal delivery of the subject compounds can be accomplished by means of iontophoretic patches and the like. For topical use, creams, ointments, jellies, solutions or suspensions, etc., containing the compounds of the present invention are employed. As used herein, topical application is also meant to include the use of mouth washes and gargles.

[0055] The compounds of the invention may be formulated for depositing into a medical device, which may include any of variety of conventional grafts, stents, including stent grafts, catheters, balloons, baskets or other device that can be deployed or permanently implanted within a body lumen. As a particular example, it would be desirable to have devices and

methods which can deliver compounds of the invention to the region of a body which has been treated by interventional technique.

[0056] In exemplary embodiment, the inhibitory agent of this invention may be deposited within a medical device, such as a stent, and delivered to the treatment site for treatment of a portion of the body.

[0057] Stents have been used as delivery vehicles for therapeutic agents (i.e., drugs). Intravascular stents are generally permanently implanted in coronary or peripheral vessels. Stent designs include those of U.S. Pat. Nos. 4,733,655 (Palmaz), 4,800,882 (Gianturco), or 4,886,062 (Wiktor). Such designs include both metal and polymeric stents, as well as self-expanding and balloon-expandable stents. Stents may also used to deliver a drug at the site of contact with the vasculature, as disclosed in U.S. Pat. No. 5,102,417 (Palmaz) and in International Patent Application Nos. WO 91/12779 (Medtronic, Inc.) and WO 90/13332 (Cedars-Sinai Medical Center), U.S. Pat. Nos. 5,419,760 (Narciso, Jr.) and U.S. Pat. No. 5,429,634 (Narciso, Jr.), for example. Stents have also been used to deliver viruses to the wall of a lumen for gene delivery, as disclosed in U.S. patent application Ser. No. 5,833,651 (Donovan et al.).

[0058] The term “deposited” means that the inhibitory agent is coated, adsorbed, placed, or otherwise incorporated into the device by methods known in the art. For example, the inhibitory agent may be embedded and released from within (“matrix type”) or surrounded by and released through (“reservoir type”) polymer materials that coat or span the medical device. In the later example, the inhibitory agent may be entrapped within the polymer materials or coupled to the polymer materials using one or more the techniques for generating such materials known in the art. In other formulations, the inhibitory agent may be linked to the surface of the medical device without the need for a coating by means of detachable bonds and release with time, can be removed by active mechanical or chemical processes, or are in a permanently immobilized form that presents the inhibitory agent at the implantation site.

[0059] In one embodiment, the inhibitory agent may be incorporated with polymer compositions during the formation of biocompatible coatings for medical devices, such as stents. The coatings produced from these components are typically homogeneous and are useful for coating a number of devices designed for implantation.

[0060] The polymer may be either a biostable or a bioabsorbable polymer depending on the desired rate of release or the desired degree of polymer stability, but a bioabsorbable polymer is preferred for this embodiment since, unlike a biostable polymer, it will not be present long after implantation to cause any adverse, chronic local response. Bioabsorbable polymers that

5 could be used include, but are not limited to, poly(L-lactic acid), polycaprolactone, polyglycolide (PGA), poly(lactide-co-glycolide) (PLLA/PGA), poly(hydroxybutyrate), poly(hydroxybutyrate-co-valerate), polydioxanone, polyorthoester, polyanhydride, poly(glycolic acid), poly(D-lactic acid), poly(L-lactic acid), poly(D,L-lactic acid), poly(D,L-lactide) (PLA), poly(L-lactide) (PLLA), poly(glycolic acid-co-trimethylene carbonate)

10 (PGA/PTMC), polyethylene oxide (PEO), polydioxanone (PDS), polyphosphoester, polyphosphoester urethane, poly(amino acids), cyanoacrylates, poly(trimethylene carbonate), poly(iminocarbonate), copoly(ether-esters) (e.g., PEO/PLA), polyalkylene oxalates, polyphosphazenes and biomolecules such as fibrin, fibrinogen, cellulose, starch, collagen and hyaluronic acid, polyepsilon caprolactone, polyhydroxy butyric acid, polyorthoesters,

15 polyacetals, polydihydropyrans, polycyanoacrylates, cross linked or amphipathic block copolymers of hydrogels, and other suitable bioabsorbable polymers known in the art. Also, biostable polymers with a relatively low chronic tissue response such as polyurethanes, silicones, and polyesters could be used and other polymers could also be used if they can be dissolved and cured or polymerized on the medical device such as polyolefins,

20 polyisobutylene and ethylene-alphaolefin copolymers; acrylic polymers and copolymers, vinyl halide polymers and copolymers, such as polyvinyl chloride; polyvinylpyrrolidone; polyvinyl ethers, such as polyvinyl methyl ether; polyvinylidene halides, such as polyvinylidene fluoride and polyvinylidene chloride; polyacrylonitrile, polyvinyl ketones; polyvinyl aromatics, such as polystyrene, polyvinyl esters, such as polyvinyl acetate;

25 copolymers of vinyl monomers with each other and olefins, such as ethylene-methyl methacrylate copolymers, acrylonitrile-styrene copolymers, ABS resins, and ethylene-vinyl acetate copolymers; pyran copolymer; polyhydroxy-propyl-methacrylamide-phenol; polyhydroxyethyl-aspartamide-phenol; polyethyleneoxide-polylysine substituted with palmitoyl residues; polyamides, such as Nylon 66 and polycaprolactam; alkyd resins,

30 polycarbonates; polyoxymethylenes; polyimides; polyethers; epoxy resins, polyurethanes; rayon; rayon-triacetate; cellulose, cellulose acetate, cellulose butyrate; cellulose acetate butyrate; cellophane; cellulose nitrate; cellulose propionate; cellulose ethers; and carboxymethyl cellulose.

[0061] Polymers and semipermeable polymer matrices may be formed into shaped articles, such as valves, stents, tubing, prostheses and the like.

[0062] In one embodiment of the invention, the inhibitory agent of the invention is coupled to a polymer or semipermeable polymer matrix that is formed as a stent or stent-graft device.

5 [0063] Typically, polymers are applied to the surface of an implantable device by spin coating, dipping or spraying. Additional methods known in the art can also be utilized for this purpose. Methods of spraying include traditional methods as well as microdeposition techniques with an inkjet type of dispenser. Additionally, a polymer can be deposited on an implantable device using photo-patterning to place the polymer on only specific portions of  
10 the device. This coating of the device provides a uniform layer around the device which allows for improved diffusion of various analytes through the device coating.

[0064] In preferred embodiments of the invention, the inhibitory agent is formulated for release from the polymer coating into the environment in which the medical device is placed. Preferably, the inhibitory agent is released in a controlled manner over an extended time  
15 frame (e.g., months) using at least one of several well-known techniques involving polymer carriers or layers to control elution. Some of these techniques were previously described in U.S. Patent Application 20040243225A1.

[0065] Moreover, as described for example in U.S. Patent No. 6,770,729, the reagents and reaction conditions of the polymer compositions can be manipulated so that the release of the  
20 inhibitory agent from the polymer coating can be controlled. For example, the diffusion coefficient of the one or more polymer coatings can be modulated to control the release of the inhibitory agent from the polymer coating. In a variation on this theme, the diffusion coefficient of the one or more polymer coatings can be controlled to modulate the ability of an analyte that is present in the environment in which the medical device is placed (e.g. an  
25 analyte that facilitates the breakdown or hydrolysis of some portion of the polymer) to access one or more components within the polymer composition (and for example, thereby modulate the release of the inhibitory agent from the polymer coating). Yet another embodiment of the invention includes a device having a plurality of polymer coatings, each having a plurality of diffusion coefficients. In such embodiments of the invention, the release of the inhibitory  
30 agent from the polymer coating can be modulated by the plurality of polymer coatings.

[0066] In yet another embodiment of the invention, the release of the inhibitory agent from the polymer coating is controlled by modulating one or more of the properties of the polymer

composition, such as the presence of one or more endogenous or exogenous compounds, or alternatively, the pH of the polymer composition. For example, certain polymer compositions can be designed to release an inhibitory agent in response to a decrease in the pH of the polymer composition. Alternatively, certain polymer compositions can be designed to release the inhibitory agent in response to the presence of hydrogen peroxide.

## V. METHODS OF TREATING DISEASES MODULATED BY CCR6

[0067] In one aspect, the present invention provides methods of treating or preventing a CCR6-mediated condition or disease by administering to a subject having such a condition or disease, a therapeutically effective amount of any compound of the invention. Preferred compounds for use in the present methods are those compounds provided above as preferred embodiments, as well as compounds specifically exemplified in the Examples below, and provided with specific structures herein. The "subject" is defined herein to include animals such as mammals, including, but not limited to, primates (e.g., humans), cows, sheep, goats, horses, dogs, cats, rabbits, rats, mice and the like. In preferred embodiments, the subject is a human.

[0068] As used herein, the phrase "CCR6-mediated condition or disease" and related phrases and terms refer to a condition or disease characterized by inappropriate, e.g., less than or greater than normal, CCR6 functional activity. Inappropriate CCR6 functional activity might arise as the result of CCR6 expression in cells which normally do not express CCR6, increased CCR6 expression (leading to, e.g., inflammatory and immunoregulatory disorders and diseases) or decreased CCR6 expression. Inappropriate CCR6 functional activity might also arise as the result of CCL20 secretion by cells which normally do not secrete CCL20, increased CCL20 expression (leading to, e.g., inflammatory and immunoregulatory disorders and diseases) or decreased CCL20 expression. A CCR6-mediated condition or disease may be completely or partially mediated by inappropriate CCR6 functional activity. However, a CCR6-mediated condition or disease is one in which modulation of CCR6 results in some effect on the underlying condition or disease (e.g., a CCR6 antagonist results in some improvement in patient well-being in at least some patients).

[0069] The term "therapeutically effective amount" means the amount of the subject compound that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by the researcher, veterinarian, medical doctor or other clinician.

[0070] Diseases and conditions associated with inflammation, infection and cancer can be treated or prevented with the present compounds and compositions. In one group of embodiments, diseases or conditions, including chronic diseases, of humans or other species can be treated with inhibitors of CCR6 function. These diseases or conditions include: (1) allergic diseases such as systemic anaphylaxis or hypersensitivity responses, drug allergies, insect sting allergies and food allergies, (2) inflammatory bowel diseases, such as Crohn's disease, ulcerative colitis, ileitis and enteritis, (3) vaginitis, (4) psoriasis and inflammatory dermatoses such as dermatitis, eczema, atopic dermatitis, allergic contact dermatitis, urticaria and pruritus, Vitiligo (5) vasculitis, (6) spondyloarthropathies, (7) scleroderma, (8) asthma and respiratory allergic diseases such as allergic asthma, allergic rhinitis, hypersensitivity lung diseases and the like, (9) autoimmune diseases, such as arthritis (including rheumatoid and psoriatic) as well as for instance Hashimoto's thyroiditis and Grave's disease, multiple sclerosis, systemic lupus erythematosus, type I diabetes, glomerulonephritis, and the like, (10) graft rejection (including allograft rejection and graft-v-host disease), and (11) other diseases in which undesired inflammatory responses are to be inhibited, such as atherosclerosis, myositis, neurodegenerative diseases (e.g., Alzheimer's disease), encephalitis, meningitis, hepatitis, nephritis, sepsis, sarcoidosis, allergic conjunctivitis, otitis, chronic obstructive pulmonary disease, sinusitis, Behcet's syndrome and gout.

[0071] Preferably, the present methods are directed to the treatment of diseases or conditions selected from allergic diseases, psoriasis, skin conditions such as atopic dermatitis and asthma and scleroderma.

[0072] In another group of embodiments, modulation of CCR6 dependent regulatory T cell trafficking may be modulated to treat diseases or conditions including cancers, infectious diseases (viral infections, e.g., HIV infection, and bacterial infections) and immunosuppressive diseases such as organ transplant conditions and skin transplant conditions. The term "organ transplant conditions" is meant to include bone marrow transplant conditions and solid organ (e.g., kidney, liver, lung, heart, pancreas or combination thereof) transplant conditions.

[0073] Depending on the disease to be treated and the subject's condition, the compounds of the present invention may be administered by oral, parenteral (e.g., intramuscular, intraperitoneal, intravenous, ICV, intracisternal injection or infusion, subcutaneous injection, or implant), inhalation, nasal, vaginal, rectal, sublingual, or topical routes of administration

and may be formulated, alone or together, in suitable dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles appropriate for each route of administration. The present invention also contemplates administration of the compounds of the present invention in a depot formulation.

- 5 [0074] Those of skill in the art will understand that agents that modulate CCR6 activity can be combined in treatment regimens with other therapeutic agents and/or with chemotherapeutic agents or radiation. In some cases, the amount of chemotherapeutic agent or radiation is an amount which would be sub-therapeutic if provided without combination with a composition of the invention. Those of skill in the art will appreciate that
- 10 "combinations" can involve combinations in treatments (i.e., two or more drugs can be administered as a mixture, or at least concurrently or at least introduced into a subject at different times but such that both are in the bloodstream of a subject at the same time). Additionally, compositions of the current invention may be administered prior to or subsequent to a second therapeutic regimen, for instance prior to or subsequent to a dose of
- 15 chemotherapy or irradiation.

[0075] The compounds of the present invention are accordingly useful in the prevention and treatment of a wide variety of inflammatory and immunoregulatory disorders and diseases.

- [0076] In the treatment or prevention of conditions which require chemokine receptor
- 20 modulation an appropriate dosage level will generally be about 0.001 to 100 mg per kg patient body weight per day which can be administered in single or multiple doses. Preferably, the dosage level will be about 0.01 to about 25 mg/kg per day; more preferably about 0.05 to about 10 mg/kg per day. A suitable dosage level may be about 0.01 to 25 mg/kg per day, about 0.05 to 10 mg/kg per day, or about 0.1 to 5 mg/kg per day. Within this
- 25 range the dosage may be 0.005 to 0.05, 0.05 to 0.5 or 0.5 to 5.0 mg/kg per day. For oral administration, the compositions are preferably provided in the form of tablets containing 1.0 to 1000 milligrams of the active ingredient, particularly 1.0, 5.0, 10.0, 15.0, 20.0, 25.0, 50.0, 75.0, 100.0, 150.0, 200.0, 250.0, 300.0, 400.0, 500.0, 600.0, 750.0, 800.0, 900.0, and 1000.0 milligrams of the active ingredient for the symptomatic adjustment of the dosage to the
- 30 patient to be treated. The compounds may be administered on a regimen of 1 to 4 times per day, preferably once or twice per day.

[0077] It will be understood, however, that the specific dose level and frequency of dosage for any particular patient may be varied and will depend upon a variety of factors including the activity of the specific compound employed, the metabolic stability and length of action of that compound, the age, body weight, hereditary characteristics, general health, sex and diet of the subject, as well as the mode and time of administration, rate of excretion, drug combination, and the severity of the particular condition for the subject undergoing therapy.

[0078] Diseases and conditions associated with inflammation, immune disorder, infection and cancer can be treated or prevented with the present compounds, compositions, and methods.

[0079] The compounds and compositions of the present invention can be combined with other compounds and compositions having related utilities to prevent and treat the condition or disease of interest, such as inflammatory or autoimmune disorders, conditions and diseases, including inflammatory bowel disease, rheumatoid arthritis, osteoarthritis, psoriatic arthritis, polyarticular arthritis, multiple sclerosis, allergic diseases, psoriasis, atopic dermatitis and asthma, and those pathologies noted above.

[0080] For example, in the treatment or prevention of inflammation or autimmunity or for example arthritis associated bone loss, the present compounds and compositions may be used in conjunction with an anti-inflammatory or analgesic agent such as an opiate agonist, a lipoxigenase inhibitor, such as an inhibitor of 5-lipoxigenase, a cyclooxygenase inhibitor, such as a cyclooxygenase-2 inhibitor, an interleukin inhibitor, such as an interleukin-1 inhibitor, an NMDA antagonist, an inhibitor of nitric oxide or an inhibitor of the synthesis of nitric oxide, a non steroidal anti-inflammatory agent, or a cytokine-suppressing anti-inflammatory agent, for example with a compound such as acetaminophen, aspirin, codeine, fentanyl, ibuprofen, indomethacin, ketorolac, morphine, naproxen, phenacetin, piroxicam, a steroidal analgesic, sufentanyl, sunlindac, tenidap, and the like. Similarly, the instant compounds and compositions may be administered with an analgesic listed above; a potentiator such as caffeine, an H2 antagonist (*e.g.*, ranitidine), simethicone, aluminum or magnesium hydroxide; a decongestant such as phenylephrine, phenylpropanolamine, pseudoephedrine, oxymetazoline, ephinephrine, naphazoline, xylometazoline, propylhexedrine, or levo desoxy ephedrine; an antitussive such as codeine, hydrocodone, caramiphen, carbetapentane, or dextromethorphan; a diuretic; and a sedating or non sedating antihistamine.

[0081] Likewise, compounds and compositions of the present invention may be used in combination with other drugs that are used in the treatment, prevention, suppression or amelioration of the diseases or conditions for which compounds and compositions of the present invention are useful. Such other drugs may be administered, by a route and in an amount commonly used therefor, contemporaneously or sequentially with a compound or composition of the present invention. When a compound or composition of the present invention is used contemporaneously with one or more other drugs, a pharmaceutical composition containing such other drugs in addition to the compound or composition of the present invention is preferred. Accordingly, the pharmaceutical compositions of the present invention include those that also contain one or more other active ingredients or therapeutic agents, in addition to a compound or composition of the present invention. Examples of other therapeutic agents that may be combined with a compound or composition of the present invention, either administered separately or in the same pharmaceutical compositions, include, but are not limited to: (a) VLA-4 antagonists, (b) corticosteroids, such as beclomethasone, methylprednisolone, betamethasone, prednisone, prednisolone, dexamethasone, fluticasone, hydrocortisone, budesonide, triamcinolone, salmeterol, salmeterol, salbutamol, formoterol; (c) immunosuppressants such as cyclosporine (cyclosporine A, Sandimmune®, Neoral®), tacrolimus (FK-506, Prograf®), rapamycin (sirolimus, Rapamune®), Tofacitinib (Xeljanz®) and other FK-506 type immunosuppressants, and mycophenolate, *e.g.*, mycophenolate mofetil (CellCept®); (d) antihistamines (H1-histamine antagonists) such as brompheniramine, chlorpheniramine, dexchlorpheniramine, triprolidine, clemastine, diphenhydramine, diphenylpyraline, tripeleminamine, hydroxyzine, methdilazine, promethazine, trimeprazine, azatadine, cyproheptadine, antazoline, pheniramine pyrilamine, astemizole, terfenadine, loratadine, cetirizine, fexofenadine, descarboethoxyloratadine, and the like; (e) non steroidal anti asthmatics (*e.g.*, terbutaline, metaproterenol, fenoterol, isoetharine, albuterol, bitolterol and pirbuterol), theophylline, cromolyn sodium, atropine, ipratropium bromide, leukotriene antagonists (*e.g.*, zafirlukast, montelukast, pranlukast, iralukast, pobilukast and SKB-106,203), leukotriene biosynthesis inhibitors (zileuton, BAY-1005); (f) non steroidal anti-inflammatory agents (NSAIDs) such as propionic acid derivatives (*e.g.*, alminoprofen, benoxaprofen, bucloxic acid, carprofen, fenbufen, fenoprofen, fluprofen, flurbiprofen, ibuprofen, indoprofen, ketoprofen, nioprofen, naproxen, oxaprozin, piroprofen, pranoprofen, suprofen, tiaprofenic acid and tioxaprofen), acetic acid derivatives (*e.g.*, indomethacin, acetaminophen, alclofenac, clidanac, diclofenac, fenclofenac, fenclozic acid, fentiazac, furofenac,

ibufenac, isoxepac, oxpinac, sulindac, tiopinac, tolmetin, zidometacin and zomepirac),  
fenamic acid derivatives (*e.g.*, flufenamic acid, meclofenamic acid, mefenamic acid, niflumic  
acid and tolfenamic acid), biphenylcarboxylic acid derivatives (*e.g.*, diflunisal and flufenisal),  
oxicams (*e.g.*, isoxicam, piroxicam, sudoxicam and tenoxicam), salicylates (*e.g.*, acetyl  
5 salicylic acid and sulfasalazine) and the pyrazolones (*e.g.*, apazone, bezpiperylon, feprazone,  
mofebutazone, oxyphenbutazone and phenylbutazone); (g) cyclooxygenase-2 (COX-2)  
inhibitors such as celecoxib (Celebrex®) and rofecoxib (Vioxx®); (h) inhibitors of  
phosphodiesterase type IV (PDE IV); (i) gold compounds such as auranofin and  
aurothioglucose, (j) etanercept (Enbrel®), (k) antibody therapies such as orthoclone (OKT3),  
10 daclizumab (Zenapax®), basiliximab (Simulect®) and infliximab (Remicade®), adalimumab  
(Humira®), golimumab (Simponi®), rituximab (Rituxan®), tocilizumab (Actemra®),  
(l) other antagonists of the chemokine receptors, especially CCR5, CXCR2, CXCR3, CCR2,  
CCR3, CCR4, CCR7, CX<sub>3</sub>CR1 and CXCR6; (m) lubricants or emollients such as petrolatum  
and lanolin, (n) keratolytic agents (*e.g.*, tazarotene), (o) vitamin D<sub>3</sub> derivatives, *e.g.*,  
15 calcipotriene or calcipotriol (Dovonex®), (p) PUVA, (q) anthralin (Drithrocreme®),  
(r) etretinate (Tegison®) and isotretinoin and (s) multiple sclerosis therapeutic agents such as  
interferon β-1β (Betaseron®), interferon (β-1α (Avonex®), azathioprine (Imurek®,  
Imuran®), glatiramer acetate (Capoxone®), a glucocorticoid (*e.g.*, prednisolone) and  
cyclophosphamide (t) DMARDS such as methotrexate and leflunomide (u) other compounds  
20 such as 5-aminosalicylic acid and prodrugs thereof; hydroxychloroquine; D-penicillamine;  
antimetabolites such as azathioprine, 6-mercaptopurine and methotrexate; DNA synthesis  
inhibitors such as hydroxyurea and microtubule disrupters such as colchicine and proteasome  
inhibitors such as bortezomib (Velcade®). The weight ratio of the compound of the present  
invention to the second active ingredient may be varied and will depend upon the effective  
25 dose of each ingredient. Generally, an effective dose of each will be used. Thus, for  
example, when a compound of the present invention is combined with an NSAID the weight  
ratio of the compound of the present invention to the NSAID will generally range from about  
1000:1 to about 1:1000, preferably about 200:1 to about 1:200. Combinations of a compound  
of the present invention and other active ingredients will generally also be within the  
30 aforementioned range, but in each case, an effective dose of each active ingredient should be  
used.

## VI. EXAMPLES

[0082] The following examples are offered to illustrate, but not to limit the claimed invention.

[0083] Reagents and solvents used below can be obtained from commercial sources such as Aldrich Chemical Co. (Milwaukee, Wisconsin, USA). <sup>1</sup>H-NMR were recorded on a Varian Mercury 400 MHz NMR spectrometer. Significant peaks are provided relative to TMS and are tabulated in the order: multiplicity (s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet) and number of protons. Mass spectrometry results are reported as the ratio of mass over charge, followed by the relative abundance of each ion (in parenthesis). In tables, a single m/e value is reported for the M+H (or, as noted, M-H) ion containing the most common atomic isotopes. Isotope patterns correspond to the expected formula in all cases.

Electrospray ionization (ESI) mass spectrometry analysis was conducted on a Hewlett-Packard MSD electrospray mass spectrometer using the HP1100 HPLC equipped with an Agilent Zorbax SB-C18, 2.1X50 mm, 5 μ column for sample delivery. Normally the analyte was dissolved in methanol at 0.1 mg/mL and 1 microlitre was infused with the delivery solvent into the mass spectrometer, which scanned from 100 to 1500 daltons. All compounds could be analyzed in the positive ESI mode, using acetonitrile / water with 1% formic acid as the delivery solvent. The compounds provided below could also be analyzed in the negative ESI mode, using 2 mM NH<sub>4</sub>OAc in acetonitrile / water as delivery system.

[0084] The following abbreviations are used in the Examples and throughout the description of the invention:

HPLC, High Pressure Liquid Chromatography; DMF, Dimethyl formamide; TFA, Trifluoroacetic Acid; THF, Tetrahydrofuran; EtOAc, Ethyl acetate; BOC<sub>2</sub>O, di-tertbutyl dicarbonate or BOC anhydride; HPLC, High Pressure Liquid Chromatography; DIPEA, Diisopropyl ethylamine; HBTU, *O*-(benzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate; dppf, 1,1'-Bis(diphenylphosphino)ferrocene; Pd<sub>2</sub>(dba)<sub>3</sub>, Tris(dibenzylideneacetone)dipalladium(0); DIPEA, diisopropylethylamine; DMP, dimethylphthalate; Me, methyl; Et, ethyl; DCM, dichloromethane.

[0085] Compounds within the scope of this invention can be synthesized as described below, using a variety of reactions known to the skilled artisan. One skilled in the art will also recognize that alternative methods may be employed to synthesize the target compounds

of this invention, and that the approaches described within the body of this document are not exhaustive, but do provide broadly applicable and practical routes to compounds of interest.


[0086] Certain molecules claimed in this patent can exist in different enantiomeric and diastereomeric forms and all such variants of these compounds are claimed.

5 [0087] The detailed description of the experimental procedures used to synthesize key compounds in this text lead to molecules that are described by the physical data identifying them as well as by the structural depictions associated with them.

[0088] Those skilled in the art will also recognize that during standard work up procedures in organic chemistry, acids and bases are frequently used. Salts of the parent compounds are  
10 sometimes produced, if they possess the necessary intrinsic acidity or basicity, during the experimental procedures described within this patent.

### Example 1: Synthesis of Methyl 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)quinoline-8-carboxylate

15




[0089] (a) To a stirred solution of  $\text{H}_2\text{O}$  (80 mL) and  $\text{H}_2\text{SO}_4$  (120 mL) at  $0^\circ\text{C}$  was added 5-bromo-8-methylquinoline (25 g, 112.6 mmol). After obtaining a solution,  $\text{CrO}_3$  was  
20 introduced (16 g, 157.6 mmol) in portion wise while maintaining the internal temperature at  $70^\circ\text{C}$ . The reaction mixture was stirred for 1 h at  $70^\circ\text{C}$ . An additional  $\text{CrO}_3$  (16 g, 157.6 mmol) was added in portions and stirred at  $80^\circ\text{C}$  for 2.5 h. After completion of the reaction, it was cooled to  $\text{r.t.}$ , poured onto crushed ice, neutralized with aqueous ammonium hydroxide to get the solids. The solids were filtered, dried under high vacuum for 16 h to get the crude  
25 5-bromoquinoline-8-carboxylic acid (28.0 g) as a green colored solid.

[0090] (b)  $\text{K}_2\text{CO}_3$  (61.3 g, 444.0 mmol) and methyl iodide (63.1 g, 444.0 mmol) were added to a stirred suspension of 5-bromoquinoline-8-carboxylic acid (28.0 g, 111.0 mmol) in  $\text{DMF}$  (250 mL) at  $\text{r.t.}$ . The reaction mixture was heated at  $45^\circ\text{C}$  for 36 h, cooled to  $\text{r.t.}$ , filtered the solids, washed with ethyl acetate (100 mL). The filtrate was diluted with water,  
30 extracted with ethyl acetate ( $3 \times 300$  mL) and washed with water ( $3 \times 100$  mL). The ethyl

acetate layer was dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated under reduced pressure. The residue was purified by flash column chromatography on silica gel, eluting with ethyl acetate in hexanes (0-20%) to afford methyl-5-bromoquinoline-8-carboxylate as a cream color solid (20.5 g, 70 % for 2 steps).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  9.07 (dd,  $J = 4.3, 1.5$  Hz, 1 H), 8.60 (dd,  $J = 8.7, 1.6$  Hz, 1 H), 7.88 (s, 2 H), 7.58 (dd,  $J = 8.6, 4.0$  Hz, 1 H), 4.05 (s, 3 H)

[0091] (c) A solution of methyl-5-bromoquinoline-8-carboxylate (20.5 g, 77.06 mmol), 4,4,5,5-tetramethyl-2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3,2-dioxaborolane (21.4 g, 84.7 mmol), and KOAc (18.8 g, 192.6 mmol) in dry 1,4-dioxane (200 mL) was purged with nitrogen gas for 10 min. Then,  $\text{Pd}(\text{dppf})\text{Cl}_2$  (3.14 g, 7.70 mmol) was added and the resulting mixture was heated at  $95^\circ\text{C}$  for 8 h, cooled to r.t and the excess solvent was removed under vacuum. The residue was diluted with ether, filtered through a Celite plug and washed with ether (200 mL); the filtrate was concentrated under reduced pressure. The resulting residue was purified by column chromatography on silica gel, eluting with 0-50% ethyl acetate in hexanes to afford a red color thick syrup/solid which was dissolved in ether and cooled to  $-20^\circ\text{C}$  and stored for 12 h, the separated light pink color solid was filtered and dried to get pure methyl 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl) quinoline-8-carboxylate (17.5 g, 78%):  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  9.14 (dd,  $J = 8.6, 0.6$  Hz, 1 H), 9.02-9.01 (m, 1 H), 8.14 (d,  $J = 7.0$  Hz, 1 H), 7.94 (d,  $J = 7.0$  Hz, 1 H), 7.48 (dd,  $J = 8.6, 4.3$  Hz, 1 H), 4.06 (s, 3 H), 1.43 (s, 12 H).


## Example 2: Synthesis of 3,4-dichloro-2-iodoaniline



[0092] To a stirred suspension of 3,4-dichloroaniline (20 g, 123 mmol), HI (48%, 15.7 g, 123 mmol),  $\text{H}_2\text{O}_2$  (30%, 8.3 g, 246 mmol) in  $\text{H}_2\text{O}$  (62 mL) at r.t. The reaction mixture was stirred in dark at r.t for overnight. The supernatant was discarded, diluted with ethyl acetate:hexanes (1:10), quenched with saturates  $\text{NaHSO}_3$ , stirred for 1 h at ambient temperature, filtered the solids. The filtrate was washed with water, saturated aqueous  $\text{NaHCO}_3$ , dried ( $\text{Na}_2\text{SO}_4$ ), filtered and concentrated to get 1:4 regioisomeric mixture of iodo derivative (minor isomer is required). It was recrystallized from cyclohexane to afford 1:1 mixture of regioisomers enriched in the filtrate. This mixture was further purified by flash


column chromatography on silica gel, eluting with ethyl acetate:hexanes (0-2%) to get the required compound as a cream color solid (5.2 g, ~15% yield).  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.22 (d,  $J = 8.6$  Hz, 1 H), 6.60 (d,  $J = 8.6$  Hz, 1 H), 4.32 (br, 2 H).

### Example 3: 3-chloro-2-iodo-4-(trifluoromethoxy)aniline



[0093] The title compound was synthesized in a similar fashion to **Example 2** in 18% yield.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.11 (d,  $J = 8.9$  Hz, 1 H), 6.67 (d,  $J = 8.9$  Hz, 1 H), 4.30 (br, 2 H).

### Example 4: Synthesis of 2-bromo-3-fluoro-4-trifluoromethoxyaniline



[0094] (a) In a 20 mL vial was added 3-fluoro-4-trifluoromethoxyaniline (150 mg, 0.769 mmol) in acetic acid (0.5 mL) at room temperature. Bromine (118  $\mu\text{L}$ , 2.30 mmol, 3 eq.) was added and the reaction was stirred for several hours. Dichloromethane (2 mL) was added to break up the solidified reaction mixture, followed by more bromine (40  $\mu\text{L}$ , 0.781 mmol, 1 eq.) addition. After stirring for another hour, dichloromethane was removed by gently blowing a stream of nitrogen to the reaction mixture. The solid was filtered and washed thoroughly with water and then dried under vacuum to give 2, 6-dibromo-3-fluoro-4-trifluoromethoxyaniline as a white solid (172 mg, 0.489 mmol, 64% yield).


[0095] (b) In a 40 mL vial was added 2, 6-dibromo-3-fluoro-4-trifluoromethoxyaniline (172 mg, 0.489 mmol), followed by tin(II) chloride dihydrate (122 mg, 0.540 mmol, 1.1 eq.).

Acetic acid (725  $\mu\text{L}$ ) and concentrated hydrochloric acid (580  $\mu\text{L}$ ) was added to make a solution. The reaction mixture was heated at 115  $^\circ\text{C}$  for 1.5 hours. After cooling the reaction to room temperature, most of acetic acid was removed under vacuum. Water was added and the product was extracted with dichloromethane three times. The organic layer was dried over anhydrous sodium sulfate. After removing the solvent under reduced pressure, the crude

material was purified using silica gel column chromatography. The recovered 2, 6-dibromo-3-fluoro-4-trifluoromethoxyaniline was eluted using a gradient of 5 ~ 10% ethyl acetate in hexanes followed by the desired product, 2-bromo-3-fluoro-4-trifluoromethoxyaniline, which was eluted with 15% ethyl acetate in hexanes (64.8 mg, 0.236 mmol, 48%). <sup>1</sup>H NMR (400

5 MHz, CDCl<sub>3</sub>) δ 7.06 (dd, *J* = 9.0, 9.0 Hz, 1 H), 6.52 (dd, *J* = 9.0, 2.0 Hz, 1 H), 4.27 (br, 2 H).

**Example 5: Synthesis of 4-chloro-5-bromo-6-amino-2,2-difluoro-1,3-benzodioxole**



10 **[0096]** (a) To a stirred solution of 4-amino-2,2-difluoro-1,3-benzodioxole (2.5 g, 14.4 mmol) in dichloromethane at 0 °C was added N-bromosuccinimide (2.7 g, 15.2 mmol) portion wise. The solution was stirred at 0 °C for 30 min, then at room temperature for 1 h. The reaction mixture was quenched with 1M sodium thiosulfate solution and diluted with

15 deionized water, extracted with dichloromethane (2 × 50 mL). The combined organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated *in vacuo*. The crude product was purified by flash chromatography (SiO<sub>2</sub>, 0–20% ethyl acetate in hexanes) to recover starting material 1g and afford 5-bromo-4-amino-2,2-difluoro-1,3-benzodioxole as colorless oil (2.33 g, 9.25 mmol, 64%). MS: (ES) *m/z* calculated for C<sub>7</sub>H<sub>4</sub>BrF<sub>2</sub>NO [M + H]<sup>+</sup> 252.0, found 252.

20 **[0097]** (b) To a stirred solution of copper (II) chloride (2.49 g, 18.5 mmol) and *tert*-butyl nitrite (2.39 g, 23.13 mmol) in acetonitrile at 55 °C was added a solution of 5-bromo-4-amino-2,2-difluoro-1,3-benzodioxole (2.33 g, 9.25 mmol) in acetonitrile. The solution was stirred at 55 °C for 30 min, then cool down to room temperature. The reaction mixture was quenched with 5% hydrochloric acid solution and diluted with deionized water, extracted


25 with ethyl acetate (2 × 50 mL). The organic layers were dried (Na<sub>2</sub>SO<sub>4</sub>), filtered, and concentrated *in vacuo*. The crude product was purified by flash chromatography (SiO<sub>2</sub>, 0–20% ethyl acetate in hexanes) to afford 5-bromo-4-chloro-2,2-difluoro-1,3-benzodioxole as

yellowish oil (1.5 g, 5.53 mmol, 60%). MS: (ES)  $m/z$  calculated for  $C_7H_2BrClF_2O_2$   $[M + H]^+$  272.0, no MS found.

[0098] (c) The solution of 5-bromo-4-chloro-2,2-difluoro-1,3-benzodioxole (1.5 g, 5.53 mmol) in 10 mL conc. sulfuric acid was cooled to  $-10\text{ }^\circ\text{C}$ . Then a solution of 1 mL fuming nitric acid and 2 mL of conc. sulfuric acid was added drop wise. The resulting solution was stirred at  $-10\text{ }^\circ\text{C}$  for  $\sim 2$  h. Then the reaction mixture was poured into ice water and a yellow solid precipitated out. Filtered off the solid, rinsed with water and collected the solid. The crude solid was purified by flash chromatography ( $SiO_2$ , 0–5% ethyl acetate in hexanes) to afford pure 6-nitro-5-bromo-4-chloro-2,2-difluoro-1,3-benzodioxole (1.0 g, 3.16 mmol, 57%). MS: (ES)  $m/z$  calculated for  $C_7HBrClF_2NO_4$   $[M + H]^+$  316.0, found 316.


[0099] (d) To the slurry of 6-nitro-5-bromo-4-chloro-2,2-difluoro-1,3-benzodioxole (1.0 g, 3.16 mmol) in ethanol (4 mL) and water (1 mL) was added ammonium chloride (3.38 g, 63.2 mmol) and iron powder (1.06 g, 18.96 mmol). The mixture was warmed up to  $90\text{ }^\circ\text{C}$  for 1 h and then cooled down. Filtered through a Celite plug and concentrated *in vacuo*. The crude product was purified by flash chromatography ( $SiO_2$ , 0–30% ethyl acetate in hexanes) to afford 6-amino-5-bromo-4-chloro-2,2-difluoro-1,3-benzodioxole as a white solid (0.90 g, 3.16 mmol, 100%). MS: (ES)  $m/z$  calculated for  $C_7H_3BrClF_2NO_2$   $[M + H]^+$  286.0, found 286.

**Example 6: Synthesis of 2,2-difluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzo-[d][1,3]dioxol-5-amine**



[0100] The title compound was synthesized in a similar fashion to **Example 1** in 55% yield.  $^1\text{H}$  NMR (400 MHz,  $CDCl_3$ )  $\delta$  7.22 (s, 1 H), 6.31 (s, 1 H), 4.80 (br, 2 H), 1.34 (s, 12 H).


**Example 7: Synthesis of 2-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-4-(trifluoromethoxy)aniline**



[0101] The title compound was synthesized in a similar fashion to **Example 1**.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  7.43 (br, 1 H), 7.10 (dd,  $J = 4.2, 1.8$  Hz, 1 H), 6.55 (d,  $J = 7.1$  Hz, 1 H), 1.34 (s, 12 H).

5

**Example 8: Synthesis of 5-[6-[(4-*tert*-butyl-3-fluorophenyl)sulfonylamino]-2,2-difluoro-1,3-benzodioxol-5-yl]quinoline-8-carboxylic acid**



10


[0102] (a) To a solution of 2,2-difluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3-benzodioxol-5-amine (**Example 5**, 2.7 g, 9.03 mmol) in dry pyridine (12 mL) was added the 4-*tert*-butyl-3-fluorophenylsulfonyl chloride (2.82 g, 11.28 mmol) and the reaction mixture was heated at 80 °C for overnight. It was cooled to r.t. excess solvent was removed in *vacuo*. The residue was diluted with saturated aqueous ammonium chloride, extracted with DCM, purified by flash column using hexanes: ethyl acetate mixture as an eluent on silica column to get the pure compound (3.28 g, 71%)

[0103] (b) A mixture of the 4-*tert*-butyl-*N*-[2,2-difluoro-6-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)-1,3-benzodioxol-5-yl]-3-fluorobenzenesulfonamide (3.2 g, 6.22 mmol), methyl-5-bromoquinoline-8-carboxylate (**Example 1**, step-b, 1.66 g, 6.22 mmol) and 2 M aq.  $\text{K}_2\text{CO}_3$  (7.8 mL, 15.55 mmol) in 1,4-dioxane (50 mL) was purged with nitrogen for 5 minutes.  $\text{Pd}(\text{PPh}_3)_4$  (360 mg, 0.31 mmol, 5 mol%) was added and the reaction mixture was heated at 95 °C for overnight. The reaction mixture was cooled to r.t.  $\text{LiOH}\cdot\text{H}_2\text{O}$  (2.6 g, 62.2

25


mmol) was added and heated at 80 °C for 1 hour. Cooled to r.t, pH of the reaction medium was adjusted to ~6 with 2N aq.HCl, extracted with ethyl acetate, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. The residue was triturated with methanol to get pure (> 95%) title compound as an off-white solid (1.75 g, 50%). <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 8.96-8.94 (m, 1 H), 8.57 (d, *J* = 7.4 Hz, 1 H), 7.93-7.91 (m, 1 H), 7.57-7.54 (m, 2 H), 7.42 – (t, *J* = 7.8 Hz, 1 H), 7.30-7.26 (m, 1 H), 7.18-7.15 (m, 1 H), 7.05 (d, *J* = 7.5 Hz, 1 H), 6.88 (s, 1 H), 6.56 (br, 1 H), 1.43 (s, 9 H); MS: (ES) *m/z* calculated for C<sub>27</sub>H<sub>21</sub>F<sub>3</sub>N<sub>2</sub>O<sub>6</sub>S

**Example 9: Synthesis of 5-[2-[(4-*tert*-butyl-3-fluorophenyl)sulfonylamino]-5-(trifluoromethoxy)phenyl]quinoline-8-carboxylic acid**




**[0104]** The title compound was synthesized in a similar fashion to **Example 8**. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 9.02 (dd, *J* = 4.4, 1.6 Hz, 1 H), 8.60 (d, *J* = 7.5 Hz, 1 H), 8.07 (dd, *J* = 8.6, 1.7 Hz, 1 H), 7.67 (dd, *J* = 8.7, 4.4 Hz, 1 H), 7.56 (d, *J* = 8.9 Hz, 1 H), 7.47 (ddt, *J* = 9.0, 1.9, 1.1 Hz, 1 H), 7.39 – 7.24 (m, 3 H), 7.14 (dd, *J* = 8.3, 2.0 Hz, 1 H), 7.03 (dd, *J* = 12.0, 2.0 Hz, 1 H), 1.39 (s, 9 H); MS: (ES) *m/z* calculated for C<sub>27</sub>H<sub>22</sub>F<sub>4</sub>N<sub>2</sub>O<sub>5</sub>S [M+H]<sup>+</sup> 558.12 found 558.1

**Example 10: 5-[2-[(4-*tert*-butyl-3-fluorophenyl)sulfonylamino]-5-chlorophenyl]quinoline-8-carboxylic acid**



[0105] The title compound was synthesized in a similar fashion to **Example 8**. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 9.10 (dd, *J* = 4.7, 1.6 Hz, 1 H), 8.66 (d, *J* = 7.8 Hz, 1 H), 8.27 (dd, *J* = 8.6, 1.6 Hz, 1 H), 7.78 (dd, *J* = 8.6, 4.7 Hz, 1 H), 7.54 (dd, *J* = 8.6, 2.7 Hz, 1 H), 7.41–7.36 (m, 3 H), 7.16 (dd, *J* = 8.2, 1.9 Hz, 1 H), 7.06 (dd, *J* = 11.1, 1.9 Hz, 1 H), 1.40 (s, 9 H); MS: (ES) *m/z* calculated for C<sub>26</sub>H<sub>22</sub>ClFN<sub>2</sub>O<sub>4</sub>S [M+H]<sup>+</sup> 513.1 found 513.1

**Example 11: Synthesis of 5-[6-[(4-*tert*-butyl-3-fluorophenyl)sulfonylamino]-2,3-dichlorophenyl]quinoline-8-carboxylic acid**




[0106] (a) To a solution of 3,4-dichloro-2-iodoaniline (**example 2**, 2.2 g, 7.67 mmol) in dry pyridine (8 mL) was added 4-*tert*-butyl-3-fluorophenylsulfonyl chloride (2.11 g, 8.43 mmol) and the reaction mixture was heated at 80 °C for overnight. LCMS indicated presence of mono and bis-sulfonamides. 10 N aq. NaOH (3 mL) and ethanol (2 mL) were added to hydrolyze the bis-sulfonamide. The reaction mixture was then heated at 80 °C for 2 h. It was cooled to r.t, excess solvent was removed in *vacuo* to afford a dark brown solid, diluted with DCM, washed with 1N aq.HCl, and purified by flash column using hexanes: ethyl acetate mixture as an eluent on silica column to get the pure 4-*tert*-butyl-*N*-(3,4-dichloro-2-iodophenyl)-3-fluorobenzenesulfonamide as an off white solid (3.5 g, 91%)

[0107] (b) A mixture of 4-*tert*-butyl-*N*-(3,4-dichloro-2-iodophenyl)-3-fluorobenzenesulfonamide (9 g, 17.9 mmol), Methyl 5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)quinoline-8-carboxylate (7.8 g, 25.1 mmol), K<sub>3</sub>PO<sub>4</sub>·H<sub>2</sub>O (10.3 g, 44.75 mmol) and SPhos (0.73 g, 1.79 mmol) in n-BuOH:H<sub>2</sub>O (75: 25 mL) solvent system was purged with nitrogen for 15 minutes. Then Pd<sub>2</sub>(dba)<sub>3</sub> (0.65 g, 0.72 mmol) was added and heated at 85 °C for 6 h, cooled to r.t, filtered through a Celite plug, washed with ethyl acetate and the filtrate was concentrated in *vacuo*. The crude product was diluted with THF/H<sub>2</sub>O (60/10 mL) and lithium hydroxide monohydrate (7.52 g, 179 mmol) was added. The reaction mixture was then heated at 60 °C for overnight, cooled to r.t, the pH was adjusted to ~ 6 by 2N aq. HCl,

extracted with ethyl acetate (3× 500 mL), the combined ethyl acetate layer was washed with water, brine, dried (Na<sub>2</sub>SO<sub>4</sub>), filtered and concentrated. The residue was purified by flash column using ethyl acetate and hexanes as eluents to get the product which was further triturated from methanol to afford the title compound as an off-white solid (7.2 g, 73%). <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 8.98 (dd, *J* = 4.4, 1.6 Hz, 1 H), 8.69 (d, *J* = 7.5 Hz, 1 H), 7.77 – 7.69 (m, 2 H), 7.63 (d, *J* = 9.0 Hz, 1 H), 7.52 (dd, *J* = 8.6, 4.3 Hz, 1 H), 7.42 (t, *J* = 8.0 Hz, 1 H), 7.32 (dd, *J* = 4.4, 1.6 Hz, 1 H), 7.17 (dd, *J* = 11.4, 2.0 Hz, 1 H), 7.05 (d, *J* = 7.5 Hz, 1 H), 6.30 (s, 1 H), 1.48 (s, 9 H); MS: (ES) *m/z* calculated for C<sub>26</sub>H<sub>21</sub>FN<sub>2</sub>O<sub>4</sub>S [M+H]<sup>+</sup> 547.06 found 547.1


**Example 12: 5-[2-[(4-*tert*-butyl-3-fluorophenyl)sulfonylamino]-5-cyanophenyl]quinoline-8-carboxylic acid bis sodium salt**



**[0108]** The title compound was synthesized in a similar fashion to **Example 11**.


**[0109]** 0.1 N aq. NaOH (2 eq) was added to the free acid suspended in acetonitrile and water and lyophilized to get the bis sodium salt. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 9.10 (dd, *J* = 4.8, 1.6 Hz, 1 H), 8.69 (d, *J* = 7.5 Hz, 1 H), 8.19 (dd, *J* = 8.7, 1.7 Hz, 1 H), 7.89 (dd, *J* = 8.5, 2.1 Hz, 1 H), 7.81 – 7.68 (m, 3 H), 7.51 – 7.38 (m, 2 H), 7.32 (dd, *J* = 8.3, 2.0 Hz, 1 H), 7.22 (dd, *J* = 11.9, 2.0 Hz, 1 H), 1.41 (s, 9 H); MS: (ES) *m/z* calculated for C<sub>27</sub>H<sub>22</sub>FN<sub>3</sub>O<sub>4</sub>S [M+H]<sup>+</sup> 504.13 found 504

**Example 13: Synthesis of 5-[6-[(4-*tert*-butyl-3-fluorophenyl)sulfonylamino]-2-fluoro-3-(trifluoromethoxy)phenyl]quinoline-8-carboxylic acid**



[0110] The title compound was synthesized in a similar fashion to **Example 11**. <sup>1</sup>H NMR (400 MHz, CDCl<sub>3</sub>) δ 9.00 (dd, *J* = 4.3, 1.7 Hz, 1 H), 8.72 (d, *J* = 7.5 Hz, 1 H), 7.86 (dt, *J* = 8.6, 1.5 Hz, 1H), 7.64 – 7.54 (m, 2 H), 7.53 – 7.33 (m, 3 H), 7.30 – 7.18 (m, 2 H), 1.42 (s, 9 H); MS: (ES) *m/z* calculated for C<sub>27</sub>H<sub>21</sub>N<sub>2</sub>O<sub>5</sub>SF<sub>5</sub> [M - H]<sup>-</sup> 579.1, found 579.1.

**Example 14: Synthesis of 5-[6-[(4-tert-butylphenyl)sulfonylamino]-2-chloro-3-(trifluoromethoxy)phenyl]quinoline-8-carboxylic acid bis sodium salt**




10

[0111] The title compound was synthesized in a similar fashion to **Example 11**. <sup>1</sup>H NMR (400 MHz, CD<sub>3</sub>OD) δ 8.79-8.77 (m, 1 H), 7.72 (d, *J* = 7.4 Hz, 1 H), 7.57-7.54 (m, 1 H), 7.52 ((d, *J* = 9.4 Hz, 1 H), 7.38 – 7.35 (m, 2 H), 7.32 – 7.29 (m, 2 H), 7.23-7.18 (m, 2 H), 6.99 (d, *J* = 7.4 Hz, 1 H), 1.32 (s, 9 H); MS: (ES) *m/z* calculated for C<sub>27</sub>H<sub>22</sub>ClF<sub>3</sub>N<sub>2</sub>O<sub>5</sub>S [M +H]<sup>+</sup>


15 579.09, found 579.1.

**Example 15: Synthesis of 5-[6-[(4-tert-butylphenyl)sulfonylamino]-2-chloro-3-(trifluoromethoxy)phenyl]quinoline-8-carboxylic acid**




[0112] The title compound was synthesized in a similar fashion to **Example 11**.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ ) 9.06 (dd,  $J = 4.8, 1.6$  Hz, 1H), 8.61 (d,  $J = 8.0$  Hz, 1H), 7.94 (dd,  $J = 8.8, 1.6$  Hz, 1H), 7.72 – 7.60 (m, 3H), 7.42 (t,  $J = 8.1$  Hz, 1H), 7.20 (dd,  $J = 8.0, 1.6$  Hz, 1H), 7.16 – 7.06 (m, 2H), 1.42 (s, 9H); MS: (ES)  $m/z$  calculated for  $\text{C}_{27}\text{H}_{21}\text{ClF}_4\text{N}_2\text{O}_5\text{S}$   $[\text{M}+\text{H}]^+$  597.08 found 597.1

**Example 16: Synthesis of 5-[6-[(4-tert-butyl-3-fluorophenyl)sulfonylamino]-4-chloro-2,2-difluoro-1,3-benzodioxol-5-yl]quinoline-8-carboxylic acid**



[0113] The title compound was synthesized in a similar fashion to **Example 11**.  $^1\text{H}$  NMR (400 MHz,  $\text{CD}_3\text{OD}$ )  $\delta$  9.08 (dd,  $J = 4.7, 1.6$  Hz, 1 H), 8.58 (d,  $J = 7.4$  Hz, 1 H), 8.11 (dd,  $J = 8.6, 1.6$  Hz, 1 H), 7.72 (dd,  $J = 8.6, 4.7$  Hz, 1 H), 7.43–7.39 (m, 2 H), 7.18 (dd,  $J = 8.2, 2.0$  Hz, 1 H), 7.08 (dd,  $J = 10.2, 2.0$  Hz, 1 H), 1.40 (s, 9 H); MS: (ES)  $m/z$  calculated for  $\text{C}_{27}\text{H}_{20}\text{ClF}_3\text{N}_2\text{O}_6\text{S}$   $[\text{M} + \text{H}]^+$  593.07, found 593.1

**Example 17: Synthesis of 4-*tert*-butyl-*N*-[2,2-difluoro-6-[8-(1*H*-tetrazol-5-yl)-5-quinolyl]-1,3-benzodioxol-5-yl]-3-fluorobenzenesulfonamide**



5

[0114] (a) A mixture of 5-bromo-8-iodoquinoline (0.52 g, 1.55 mmol),  $\text{Zn}(\text{CN})_2$  (218 mg, 1.86 mmol), and dppf (103 mg, 0.186 mmol) in DMF (2.5 mL) was purged with  $\text{N}_2$  (gas) for 5 minutes.  $\text{Pd}_2(\text{dba})_3$  (85 mg, 0.093 mmol) was added and the resulting mixture was heated at 95 °C for 5 h. After completion of the reaction, it was cooled to r.t., diluted with water, and the aqueous solution was extracted with ethyl acetate (3 × 50 mL). The combined organic layer was washed with brine, dried over  $\text{Na}_2\text{SO}_4$  and concentrated under reduced pressure. The crude product was purified by column chromatography on silica gel, eluting with ethyl acetate-hexanes (5-25%) to get 5-bromoquinoline-8-carbonitrile as a brown solid (72 mg).

10


[0115] (b) The Suzuki reaction was carried out similar to **Example 8**. 72 mg of the starting material afforded 70 mg of the product.

15

[0116] (c) To a stirred solution of the above nitrile (35 mg, 0.064 mmol) in  $\text{H}_2\text{O}/\text{IPA}$  (5:1, 2 mL) at r.t were added  $\text{ZnBr}_2$  (43 mg, 0.324 mmol) and  $\text{NaN}_3$  (21 mg, 0.324 mmol). The reaction mixture was heated at 100 °C for 16 h. After completion of the reaction, it was cooled to r.t and neutralized with 1N aq. HCl. The aqueous solution was extracted with ethyl


acetate ( $2 \times 25$  mL) and the combined organic layers were washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated under reduced pressure. The crude product was triturated with methanol to give the title compound as a brownish solid (8 mg).  $^1\text{H}$  NMR (400 MHz,  $\text{DMSO}-d_6$ )  $\delta$  9.84 (s, 1H), 9.04 (dd,  $J = 4.2, 1.7$  Hz, 1H), 8.45 (d,  $J = 7.5$  Hz, 1H), 7.87 (dd,  $J = 8.6, 1.7$  Hz, 1H), 7.58 (dd,  $J = 8.6, 4.2$  Hz, 1H), 7.43 (d,  $J = 12.6$  Hz, 1H), 7.24 (t,  $J = 8.1$  Hz, 1H), 7.16 – 7.0 (m, 3H), 1.24 (s, 9H); MS: (ES)  $m/z$  calculated for  $\text{C}_{28}\text{H}_{22}\text{F}_3\text{N}_5\text{O}_4\text{S}$   $[\text{M}+\text{H}]^+$  582.13 found 582.1

**Example 18: Synthesis of 5-[2-[(4-*tert*-butylphenyl)sulfonylamino]-5-cyanophenyl]quinoline-8-carboxylic acid**



[0117] The title compound was synthesized in a similar fashion to **Example 11**.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ )  $\delta$  9.00 (dd,  $J = 4.3, 1.6$  Hz, 1 H), 8.68 (d,  $J = 7.4$  Hz, 1 H), 7.92 (d,  $J = 8.6$  Hz, 1 H), 7.87 (dd,  $J = 8.6, 1.2$  Hz, 1 H), 7.78 (dd,  $J = 9.0, 2.0$  Hz, 1 H), 7.62 – 7.50 (m, 5 H), 7.45 (d,  $J = 7.9$  Hz, 1 H), 7.19 (d,  $J = 7.5$  Hz, 2 H), 6.67 (s, 1 H), 1.37 (s, 9 H); MS: (ES)  $m/z$  calculated for  $\text{C}_{27}\text{H}_{23}\text{N}_3\text{O}_4\text{S}$   $[\text{M}+\text{H}]^+$  486.14 found 486.1.


**Example 19: Synthesis of 5-[6-[(4-*tert*-butylphenyl)sulfonylamino]-2-chloro-3-cyanophenyl]quinoline-8-carboxylic acid trifluoroacetic acid salt**



[0118] The title compound was synthesized in a similar fashion to **Example 11**.  $^1\text{H}$  NMR (400 MHz, DMSO- $d_6$ )  $\delta$  9.85 (bs, 1 H), 9.12-9.13 (m, 1 H), 8.48 (d,  $J = 7.4$  Hz, 1 H),  $\delta$  8.06 (bs, 1 H), 7.89-7.87 (m, 1 H), 7.70 – 7.52 (m, 7 H), 7.16 (d,  $J = 6.7$  Hz, 1 H), 1.30 (s, 9 H); MS: (ES)  $m/z$  calculated for  $\text{C}_{27}\text{H}_{22}\text{ClN}_3\text{O}_4\text{S}$   $[\text{M} + \text{H}]^+$  520.09, found 520.1

5

**Example 20: Synthesis of 5-[6-[(4-*tert*-butylphenyl)sulfonylamino]-3-cyano-2-fluorophenyl]quinoline-8-carboxylic acid hydrochloric acid salt**



10

[0119] The title compound was synthesized in a similar fashion to **Example 11**.  $^1\text{H}$  NMR (400 MHz,  $\text{CDCl}_3$ ) 9.04 (dd,  $J = 4.3, 1.6$  Hz, 1 H), 8.76 (d,  $J = 7.4$  Hz, 1 H), 7.84 – 7.75 (m, 3 H), 7.64 – 7.53 (m, 5H), 7.22 (d,  $J = 7.4$  Hz, 1 H), 6.59 (bs, 1 H) 1.37 (s, 9H); MS: (ES)  $m/z$  calculated for  $\text{C}_{27}\text{H}_{22}\text{FN}_3\text{O}_4\text{S}$   $[\text{M} + \text{H}]^+$  504.1 found 504.4

15

**Biological Example 1: Ligand binding assay**

[0120] Ligand binding assay was used to determine the ability of potential CCR6 antagonists to block the interaction between CCR6 and its ligand CCL20 (MIP3alpha). L1.2 cells stably expressing the CCR6 receptor, were centrifuged and resuspended in assay buffer (20 mM HEPES pH 7.1, 140 mM NaCl, 1 mM  $\text{CaCl}_2$ , 5 mM  $\text{MgCl}_2$ , 0.1% sodium azide and with 0.1% bovine serum albumin) to a concentration of  $5 \times 10^5$  cells/mL. Binding assays were set up as follows. First, 0.1 mL of cells ( $5 \times 10^4$  cells/well) was added to the assay plates containing the compounds, giving a final concentration of ~2-10  $\mu\text{M}$  each compound for screening (or part of a dose response for compound  $\text{IC}_{50}$  determinations). Then 0.1 mL of  $^{125}\text{I}$  labeled CCL20 (obtained from PerkinElmer; Waltham, MA) diluted in assay buffer to

a final concentration of ~50 pM, yielding ~30,000 cpm per well, was added, the plates sealed and incubated for approximately 3 hours at 25 °C on a shaker platform. Reactions were aspirated onto GF/B glass filters pre-soaked in 0.3% polyethyleneimine (PEI) solution, on a vacuum cell harvester (Packard Instruments; Meriden, CT). Scintillation fluid (50 uL; Microscint 20, Packard Instruments) was added to each well, the plates were sealed and radioactivity measured in a Top Count scintillation counter (Packard Instruments). Control wells containing either diluent only (for total counts) or 20 uM compound were used to calculate the percent of total inhibition for compound. The computer program Prism from GraphPad, Inc. (San Diego, Ca) was used to calculate IC<sub>50</sub> values. IC<sub>50</sub> values are those concentrations required to reduce the binding of labeled TARC to the receptor by 50%. Compounds in Figure 1 having an IC<sub>50</sub> value in the binding assay of less than 100 nM are labeled (+++); from 100-500 nM are labeled (++); and less than or equal to 20 μM but above 500 nM are labeled (+).

#### Biological Example 2: Migration/Chemotaxis Assay

**[0121]** A serum chemotaxis assay was used to determine the efficacy of potential receptor antagonists at blocking the migration mediated through chemokine receptors, such as CCR6. This assay was routinely performed using the ChemoTX® microchamber system with a 5-μm pore-sized polycarbonate membrane. To begin such an assay, chemokine-receptor expressing cells (such as L1.2 cells stably expressing CCR6) were collected by centrifugation at 400 x g at room temperature, then suspended at 50 million/ml in human serum. The compound being tested or an equivalent volume of its solvent (DMSO) was then added to the cell/serum mixture at a final DMSO concentration of 0.25% (v/v). Separately, recombinant human CCL20 was diluted with chemotaxis buffer (HBSS + 0.1% BSA), generally spanning a range from 0.01 nM to 500 nM, after which 29 μl of diluted chemokine was placed in the lower wells of the ChemoTX® plate. The 5-μm (pore size) polycarbonate membrane was placed onto the plate, and 20 μL of the cell/compound mixture was transferred onto each well of the membrane. The plates were incubated at 37 °C for 90 minutes, after which the polycarbonate membranes were removed and 5 μl of the DNA-intercalating agent CyQUANT (Invitrogen, Carlsbad, CA) was added to the lower wells. The amount of fluorescence, corresponding to the number of migrated cells, was measured using a Spectrafluor Plus plate reader (TECAN, San Jose, CA).

**(a) Evaluation of a compound of interest in an oxazolone induced model of DTH**

[0122] Compounds of the invention were assessed in the murine model of dermal delayed type hypersensitivity (DTH) induced by oxazolone. Briefly, 8-10 week old BALB/c mice were sensitized topically with a 1% solution of oxazolone dissolved in ethanol on their shaved abdomens on day 0. On day 6 post sensitization mice were dosed orally with either vehicle or increasing doses of compounds **1.061** and **1.033** of the invention immediately prior to and 4 hours following a topical challenge with a 0.5% solution of oxazolone in ethanol on the right ear. The following day (day 7), ear thicknesses were measured using caliper measurements. Animals treated with compound had significantly reduced ear swelling compared to vehicle treated controls indicating a compound mediated decrease in oxazolone induced dermal hypersensitivity.

**(b) Evaluation of a compound of interest in an Imiquimod induced model of psoriasis**

[0123] Multiple mouse models of psoriasis are known. For example, in the Imiquimod-induced mouse model of psoriasis, hair was removed from the back of balb/c mice three days prior to the application of 5% Imiquimod once a day. Vehicle (1% HPMC) or CCR6 compound was administered eg. orally at between 5 and 200mg/kg once/twice daily at the initiation of Imiquimod application and continued for five to ten days. Therapeutic value was assessed using the following methodologies: daily electronic caliper measurements and immunohistochemistry at the end point to assess skin thickening, flow cytometry and immunofluorescence at the end point to measure leukocyte infiltration and inflammation, and mRNA quantitation via Luminex at days two and five to assess molecular changes. Treatment of animals with compound **1.033** or **1.061** of the invention resulted in a significant improvement in disease.

**(c) Evaluation of a compound of interest in an IL-23 induced model of psoriasis.**

[0124] Another method to induce phenotypic changes associated with psoriasis involved intra-dermal injections of IL-23. Briefly, 500 ng of recombinant murine IL-23 was intra-dermally injected every other day into the right ear of C57BL6/N mice for a total of 5-6

administrations over a period of ten to twelve days. Concomitantly, PBS was intra-dermally injected every other day into the left ear of the same mice. CCR6 compound was dosed prophylactically at eg. between 5-200mg/kg once/twice daily via the oral or other route. Therapeutic dosing was performed by orally providing IL-23 challenged mice with vehicle control for four days. On the fifth day, animals were dosed with a compound of the invention at eg. 5-200 mg/kg once/twice daily via the oral or other route (at this point, two rounds of IL-23 have been intra-dermally injected into the right ear). Efficacy was assessed quantitatively by manually measuring ear thickness using a caliper daily. Treatment of animals with compound **1.033** or **1.061** of the invention resulted in a significant improvement in disease.

**(d) Evaluation of a compound of interest in a mouse model of septic shock.**

[0125] This example describes a procedure to evaluate the efficacy of CCR6 antagonists for treatment of septic shock. An animal model of endotoxic shock can be induced by injecting rodents with lipopolysaccharide (LPS). Three series of mouse groups, comprising 15 mice per group, are treated with an intra-peritoneal injection of an L.D. (lethal dose)-90 of LPS. One series of mice additionally receives phosphate buffered saline (PBS) and Tween 0.5% i.p. 30 minutes before LPS administration. A second series consists of groups of mice receiving different doses of the CCR6 antagonist given either intra-peritoneally, intra-venously, sub-cutaneously, intra-muscularly, orally, or via any other mode of administration 30 minutes before, or concurrently with, LPS administration. A third series of mice, serving as positive control, consists of groups treated with either mouse IL-10 i.p., or anti-TNF antibodies i.p., 30 minutes before LPS administration. Mice are monitored for death for 72 hours following the LPS injection.

**(e) Evaluation of a compound of interest in a rodent model of asthma/allergic lung inflammation.**

[0126] CCR6 compounds of the invention can be assessed in a murine model of allergic asthma. Asthma is induced in 8 – 10 week old BALB/c mice by sensitizing mice with OVA in Alum adjuvant on days 0 and 10. On day 20 mice are challenged with OVA in PBS intranasally to elicit airway inflammation. Groups of mice are either treated with vehicle, or increasing doses of a compound of the invention starting on day 20 and lasting until day 23. Animals are subsequently analyzed at day 23 after the intranasal OVA challenge for cellular

infiltrates in bronchoalveolar lavage (BAL). Mice treated with a compound of the invention will display significantly reduced BAL leukocyte numbers relative to vehicle treated mice.

**(f) Evaluation of a compound of interest in a mouse model of rheumatoid arthritis.**

5 [0127] This example describes a procedure to evaluate the efficacy of CCR6 antagonists for treatment of rheumatoid arthritis. An animal model of rheumatoid arthritis was induced in rodents by injecting them with type II collagen in selected adjuvants. Three series of rodent groups consisting of 15 genetically-susceptible mice or rats per group were injected sub-cutaneously or intra-dermally with type II collagen emulsified in Complete Freund's  
10 Adjuvant at days 0 and 21. One series of rodents additionally received PBS and Tween 0.5% i.p. at the initial sensitization and at different dosing schedules thereafter. A second series consists of groups of rodents received different doses of the CCR6 antagonist given either intra-peritoneally, intra-venously, sub-cutaneously, intramuscularly, orally, or via any other mode of administration at the initial sensitization, and at different dosing schedules thereafter.  
15 A third series of rodents, serving as positive control, may consist of groups treated with either mouse IL-10 i.p., or anti-TNF antibodies i.p. at the initial sensitization, and at different dosing schedules thereafter. Animals were monitored from weeks 3 till 8 for the development of swollen joints or paws, and graded on a standard disease severity scale. Disease severity was confirmed by histological analysis of joints. Animals treated with compound 1.061 of  
20 the invention displayed significantly improved scores following treatment.

**(g) Evaluation of a compound of interest in a mouse model of SLE.**

[0128] This example describes a procedure to evaluate efficacy of CCR6 antagonists for treatment of Systemic Lupus Erythematosus (SLE). Female NZB/W FI mice spontaneously develop an SLE-like pathology commencing at 6 months of age that is characterized by  
25 proteinuria, serum autoantibodies, glomerulonephritis, and eventually death. Three series of NZB/W FI mouse groups comprising 20 mice per group are tested for efficacy of CCR6 antagonist as follows: One series of mice additionally receives phosphate buffered saline (PBS) and Tween 0.5% i.p. soon after weaning, and thereafter at varying dosing schedules. A second series consists of groups of mice receiving different doses of the CCR6 antagonist  
30 given either intra-peritoneally, intra-venously, sub-cutaneously, intramuscularly, orally, or via any other mode of administration soon after weaning, and thereafter at varying dosing schedules. A third series of mice, serving as positive control, consists of groups treated with

anti-IL10 antibodies given soon after weaning, and thereafter at varying dosing schedules. Disease development is monitored in terms of eventual mortality, kidney histology, serum autoantibody levels, and proteinuria.

**(h) Evaluation of a compound of interest in a mouse model of malignancy.**

- 5 [0129] This example describes a procedure to evaluate efficacy of CCR6 antagonists for treatment of malignancy. Normal mouse strains can be transplanted with a variety of well-characterized mouse tumor lines, including a mouse thymoma EL4 which has been transfected with OVA to allow easy evaluation of tumor specific antigen responses following vaccination with OVA. Three series of mouse groups from any of these tumor models are
- 10 tested for CCR6 antagonist efficacy as follows: One series of mice additionally receives PBS and Tween 0.5% i.p. soon after tumor transplant, and thereafter at varying dosing schedules. A second series consists of groups of mice receiving different doses of the CCR6 antagonist given either intra-peritoneally, intra-venously, sub-cutaneously, intramuscularly, orally, or via any other mode of administration soon after tumor transplant, and thereafter at varying
- 15 dosing schedules. A third series of mice, serving as positive control, consists of groups treated with either anti-IL17 antibodies, anti-IFN $\gamma$  antibodies, given i.p. soon after tumor transplant, and thereafter at varying dosing schedules. Efficacy is monitored via tumor growth versus regression. In the case of the OVA-transfected EL4 thymoma model, cytolytic OVA-specific responses can be measured by stimulating draining lymph node cells with
- 20 OVA in vitro, and measuring antigen-specific cytotoxicity at 72 hours. Analysis can be performed on eg. tumor mass, IL-17 levels or infiltration of regulatory T cells and treatment with a compound of the invention would be expected to produce a beneficial effect in this model.

**(i) Evaluation of a compound of interest in a mouse carcinoma model.**

- 25 [0130] The mouse RENCA tumor model accurately mimics the progression of human adult renal cell carcinoma specifically with reference to spontaneous metastasis to lungs and serves as a model for solid tumors. Balb/c 6-8 week old female mice are inoculated with approximately 5e5 RENCA cells (mouse renal adenocarcinoma; ATCC cat# CRL-2947) under the kidney capsule and kidney tumor growth is observed over 22 days, with lung
- 30 metastasis observed as early as day 15. Animals are dosed with either vehicle or a compound of the invention eg daily subcutaneously, from the time of tumor implantation to monitor effects on primary growth, or at a later time (eg day 7) to monitor the compound effect on

metastasis. Primary tumor areas are measured twice a week using mechanical calipers. Tumor volumes are calculated by the formula  $v = \pi ab^2/6$ , where a is the longest diameter and b is the next longest diameter perpendicular to a. A reduction in tumor volume or incidence of metastasis indicates efficacy of compound in this indication.

5                    **(j) Evaluation of a compound of interest in a mouse model of IBD.**

[0131] This example describes a procedure to evaluate the efficacy of CCR6 antagonists in Inflammatory Bowel Disease (IBD). Several mouse models of IBD (including Crohn's Disease and Ulcerative Colitis) have been developed. Some of these are spontaneous models occurring in genetically engineered transgenic mice that have been depleted of certain  
 10 cytokine genes (e.g. IL-10, or IL-2). Another mouse model of IBD is obtained by transferring highly purified populations of CD4<sup>+</sup> T lymphocytes bearing a particular surface marker phenotype (namely CD45 RB hi) into SCID mice. Three series of mouse groups from anyone of these models can be used to evaluate CCR6 antagonist efficacy as follows. One group of mice additionally receives PBS and Tween 0.5% i.p. soon after weaning in the case  
 15 of the spontaneous models in transgenic mice, or at time of cell transfer into SCID mice and varying dosings thereafter for the cell transfer model. A second series consists of groups of mice receiving different doses of the CCR6 antagonist given either intraperitoneally, intravenously, sub-cutaneously, intra-muscularly, orally, or via any other mode of administration soon after weaning in the case of the spontaneous models in transgenic mice, or at time of  
 20 cell transfer into SCID mice and varying dosings thereafter for the cell transfer model. A third series of mice, serving as positive control, consists of groups treated with antibodies to either IFN $\gamma$ , or TNF, or with cytokine IL-10 soon after weaning in the case of the spontaneous models in transgenic mice, or at time of cell transfer into SCID mice and varying dosings thereafter for the cell transfer model. Mice are evaluated for 6-8 weeks for disease  
 25 development, monitored initially via weight loss and/or prolapsed rectum, and eventually by histological evaluation of the animals colon and intestinal tract.

**SEQ ID> 1 Human CCR6 mRNA transcript variant 1. ACCESSION NM\_004367**

**VERSION NM\_004367.5 GI:150417991**

30 1 agtgtatggg tgaaggaggc agcagtgtgg ccggagagga gagctgggct gggagcacag

61 gaaggteccc aggactctgt ggtcatcagt aagagagggc ccacgtgtat atgctgggtga  
 121 acagaaatgt caaccttttc aaagtctgac atttaagaga aaaaactgtg gctgttggtt  
 181 tgtggaacag acagctcctt ctttattgag tcacctctac ttctctgcta ccgctgcctg  
 241 tgagctgaag gggctgaacc atacactcct tttctacaa ccagcttgca tttttctgc  
 5 301 ccacaatgag cggggaatca atgaattca gcgatgttt cgactccagt gaagattatt  
 361 ttgtgtcagt caatacttca tattactcag ttgattctga gatgttactg tgctccttgc  
 421 aggaggtcag gcagttctcc aggctatttg taccgattgc ctactccttg atctgtgtct  
 481 ttggcctcct ggggaatatt ctggtgggtga tcaccttgc ttttataag aaggccaggt  
 541 ctatgacaga cgtctatctc ttgaacatgg ccattgcaga cactctcttt gttcttactc  
 10 601 tcccattctg ggcagtgagt catgccaccg gtgcgtgggt ttcagcaat gccacgtgca  
 661 agttgctaaa aggcactctat gccatcaact ttaactgcgg gatgctgctc ctgacttgca  
 721 ttagcatgga ccggtacatc gccattgtac aggcgactaa gtcattccgg ctccgatcca  
 781 gaacactacc gcgcagcaaa atcatctgcc ttgttgtgtg ggggctgtca gtcacatct  
 841 ccagctcaac tttgtcttc aacaaaaat acaacacca aggagcgat gtctgtgaac  
 15 901 ccaagtacca gactgtctcg gagcccatca ggtggaagct gctgatgttg gggcttgagc  
 961 tactctttgg tttcttctc cttttagtgt tcatgatatt ttgttacag ttcatgtca  
 1021 aaaccttggg gcaagctcag aattctaaaa ggcacaaagc catccgtgta atcatagctg  
 1081 tgggtcctgt gttctggct tgcagattc ctcataacat ggtcctgctt gtgacggctg  
 1141 caaatttggg taaatgaac cgatcctgcc agagcgaaaa gctaattggc tatacgaaaa  
 20 1201 ctgtcacaga agtcttggt ttctgcact gctgcctgaa ccctgtgctc tacgtttta  
 1261 ttgggcagaa gttcagaaac tactttctga agatcttgaa ggacctgtgg tgtgtgagaa  
 1321 ggaagtacaa gtctcaggc ttctcctgtg ccgggaggta ctcaaaaaac atttctcggc  
 1381 agaccagtga gaccgcagat aacgacaatg cgtcgtcctt cactatgtga tagaaagctg

1441 agtctcccta aggcatgtgt gaaacatact catagatgtt atgcaaaaaa aagtctatgg  
 1501 ccaggtatgc atggaaaatg tgggaattaa gcaaaatcaa gcaagcctct ctctgcggg  
 1561 acttaacgtg ctcatgggct gtgtgatctc ttcaggggtg ggtggtctct gataggtagc  
 1621 atttccagc actttgcaag gaatgtttg tagctctagg gtatatatcc gcctggcatt  
 5 1681 tcacaaaaca gcctttggga aatgctgaat taaagtgaat tgttgacaaa tgtaaacatt  
 1741 ttcagaaata ttcataagc gggtcacagat cacagtgtct tttggttaca gcacaaaatg  
 1801 atggcagtg tttgaaaaac taaaacagaa aaaaaaatgg aagccaacac atcactcatt  
 1861 ttaggcaaat gtttaaacat tttatctat cagaatgtt attgttgctg gttataagca  
 1921 gcaggattgg ccggctagt ttcctctca tttcccttg atacagtcaa caagcctgac  
 10 1981 cctgtaaaat ggaggtggaa agacaagctc aagtgtcac aacctggaag tgcttcggga  
 2041 agaaggggac aatggcagaa caggtgttg tgacaattgt caccaattgg ataaagcagc  
 2101 tcaggttgta gtgggccatt aggaaactgt cggtttgctt tgatttcctt gggagctgtt  
 2161 ctctgtcgtg agtgtctct gtctaaact ccattaagct gagagtgcta tgaagacagg  
 2221 atctagaata atctgtcga cagctgtgct ctgagtcct agcggagttc cagcaaaca  
 15 2281 aatggactca agagagattt gattaatgaa tcgtaatgaa gttgggggtt attgtacagt  
 2341 ttaaatgtt agatgtttt aatttttaa ataatggaa tactttttt ttttttaa  
 2401 agaaagcaac ttactgaga caatgtaga agaagtttg ttcgtttct ttaatgtgt  
 2461 tgaagagcaa tgtgtggctg aagactttg ttatgaggag ctgcagatta gctaggggac  
 2521 agctggaatt atgtggctt ctgataatta ttttaaagg gtctgaaatt tgtgatggaa  
 20 2581 tcagatttta acagctctct tcaatgacat agaaagtca tggaactcat gttttaaag  
 2641 ggctatgtaa atatatgaac attagaaaaa tagcaacttg tgttacaaaa atacaaacac  
 2701 atgttaggaa ggtactgtca tgggctaggc atggtggctc acacctgtaa tcccagcatt  
 2761 ttgggaagct aagatgggtg gatcactga ggtcaggagt ttgagaccag cctggccaac

2821 atggcgaaac ccctctctac taaaaataca aaaatttgcc aggcgtggtg gcgggtgcct  
 2881 gtaatcccag ctacttggga ggctgaggca agagaatcgc ttgaaccag gaggcagagg  
 2941 ttgcagttag ccgagatcgt gccattgcac tccagcctgg gtgacaaagc gagactccat  
 3001 ctcaaaaaaa aaaaaaaaaa aaaaggaaag aactgtcatg taaacatacc aacatgttta  
 5 3061 aacctgacaa tgggtgttatt tgaaacttta tattgttctt gtaagcttta actatatctc  
 3121 tctttaaata gcaaaataat gtcttaagat tcaaagtctg tatttttaa gcatggcttt  
 3181 ggctttgcaa aataaaaaat gtgttttgta catgaa

**SEQ ID> 2 Human CCR6 mRNA transcript variant 2. ACCESSION NM\_031409**

**10 VERSION NM\_031409.3 GI:150417990**

1 aactcacag gcctcttga aacgttccca aatcttccca gtcggcttgc agagactcct  
 61 tgctcccagg agataaccag gtaaaggagt atgaaagttt gggtaaaaac tcattgctgc  
 121 aaattgaaaa ccatgcaaag gctgtcttcc tctggggagt tcaatgcctc tctttttctt  
 181 atcactttac cattggttgg actttgattc cagggatcct acgattactc aataccctac  
 15 241 aggatataca tgggtaacca ttgcatattg ggcaaataagg cgttactttt caataggaag  
 301 tggcaatcca gaacttgctt ttgggcaatt ctagtagctc accgcttttt tcttaatgac  
 361 tgctagaagc tgcatcttat tgacagatgg tcatcacatt ggtgagctgg agtcatcaga  
 421 ttgtggggcc cggagtgagg ctgaaggagag tggatcagag cactgcctga gactcacctc  
 481 tactttcctg ctaccgctgc ctgtgagctg aaggggctga accatacact cctttttcta  
 20 541 caaccagctt gcatttttc tgcccacaat gagcggggaa tcaatgaatt tcagcgaagt  
 601 ttgcactcc agtgaagatt attttgtgtc agtcaatact tcatattact cagttgattc  
 661 tgagatgtta ctgtgtctct tgcaggaggt caggcagttc tccaggctat ttgtaccgat  
 721 tgcctactcc ttgatctgtg tctttggcct cctggggaat attctggtgg tgatcacctt

781 tgctttttat aagaaggcca ggtctatgac agacgtctat ctctgaaca tggccattgc  
 841 agacatcctc tttgttctta ctctccatt ctgggcagtg agtcatgcca ccggtgcgtg  
 901 ggttttcagc aatgccacgt gcaagttgct aaaaggcatc tatgccatca actttaactg  
 961 cgggatgctg ctctgactt gcattagcat ggaccggtac atcgccattg tacaggcgac  
 5 1021 taagtcattc cggctccgat ccagaacact accgcgagc aaaatcatct gccttgtgtg  
 1081 gtgggggctg tcagtcatca tctccagctc aactttgtc ttcaacaaa aatacaacac  
 1141 ccaaggcagc gatgtctgtg aaccaagta ccagactgtc tcggagccca tcaggtggaa  
 1201 gctgctgatg ttggggcttg agctactctt tggttcttt atcccttga tgttcattg  
 1261 atttgttac acgttcattg tcaaaacctt ggtgcaagct cagaattcta aaaggcacia  
 10 1321 agccatccgt gtaatcatag ctgtggtgct tgtgttctg gcttgcaga ttctcataa  
 1381 catggtcctg cttgtgacgg ctgcaaatg gggtaaaatg aaccgatcct gccagagcga  
 1441 aaagctaatt ggctatacga aaactgtcac agaagtctg gctttcctgc actgtgcct  
 1501 gaacctgtg ctctacgctt ttattgggca gaagtcaga aactacttc tgaagatctt  
 1561 gaaggacctg tgggtgtgga gaaggaagta caagtcctca ggcttctct gtgccgggag  
 15 1621 gtactcagaa aacatttctc ggagaccag tgagaccgca gataacgaca atgcgtcgtc  
 1681 cttcactatg tgatagaaag ctgagtctcc ctaaggcatg tgtgaaacat actcatagat  
 1741 gttatgcaaa aaaaagtcta tggccaggta tgcattgaaa atgtgggaat taagcaaat  
 1801 caagcaagcc tctctcctgc gggaactaac gtgctcatgg gctgtgtgat ctcttcaggg  
 1861 tgggggtggtc tctgataggt agcattttcc agcactttgc aaggaatgtt tttagctct  
 20 1921 agggatatata tccgcctggc atttcacaaa acagccttg ggaaatgctg aattaaagt  
 1981 aattgttgac aaatgtaaac attttcagaa atattcatga agcgggcaca gatcacagt  
 2041 tcttttggtt acagcacaaa atgatggcag tggtttgaaa aactaaaaca gaaaaaaaaa  
 2101 tggaagccaa cacatcactc attttaggca aatgtttaa cattttatc tatcagaatg

2161 ttattgttg ctggtataa gcagcaggat tggccggcta gtgttctc tcatttccct  
 2221 ttgatacagt caacaagcct gacctgtaa aatggagggtg gaaagacaag ctcaagtgtt  
 2281 cacaacctgg aagtgcctcg ggaagaaggg gacaatggca gaacagggtg tggtgacaat  
 2341 tgtcaccaat tggataaagc agctcagggt gtagtgggcc attaggaaac tgcgggttg  
 5 2401 ctttgatttc cctgggagct gttctctgtc gtgagtgtct cttgtctaaa cgtccattaa  
 2461 gctgagagtg ctatgaagac aggatctaga ataacttgc tcacagctgt gctctgagtg  
 2521 cctagcggag ttccagcaaa caaatggac tcaagagaga ttgattaat gaatcgtaat  
 2581 gaagttgggg ttattgtac agttaaaat gttagatgtt ttaatttt taaataaatg  
 2641 gaatactttt tttttttt taaagaaagc aactttactg agacaatga gaaagaagt  
 10 2701 ttgtccggt tctttaatgt ggtgaagag caatgtgtgg ctgaagactt ttgttatgag  
 2761 gagctgcaga ttgctaggg gacagctgga attatgctgg cttctgataa ttattttaa  
 2821 ggggtctgaa attgtgatg gaatcagatt ttaacagctc tctcaatga catagaaagt  
 2881 tcatggaact catgtttta aagggtatg taaatatatg aacattagaa aaatagcaac  
 2941 ttgtgttaca aaaatacaaa cacatgtag gaaggtactg tcatgggcta ggcatgggtg  
 15 3001 ctcacacctg taatcccagc attttgggaa gctaagatgg gtggatcact tgaggtcagg  
 3061 agtttgagac cagcctggcc aacatggcga aaccctctc tactaaaaat acaaaaattt  
 3121 gccaggcgtg gtggcgggtg cctgtaatcc cagctacttg ggaggctgag gcaagagaat  
 3181 cgttgaaacc caggaggcag aggttgcatg gagccgagat cgtgccattg cactccagcc  
 3241 tgggtgacaa agcgagactc catctcaaaa aaaaaaaaaa aaaaaagga aagaactgtc  
 20 3301 atgtaaacat accaacaatg ttaacctga caatggtgtt attgaaact ttattgtt  
 3361 cttgtaagct ttaactatat ctctctttaa aatgcaaaat aatgtcttaa gattcaaagt  
 3421 ctgtattttt aaagcatggc ttggctttg caaaataaaa aatgtgttt gtacatgaa

SEQ ID> 3 Human CCR6 amino acid sequence

MSGESMNFSDVFDSSDYFVSVNTSYYSVDSEMLLCSLQEVQRSLFVPIAYSLICV  
FGLLGNILVVITFAFYKKARSM TDVYLLNMAIADILFVLTLPFWAVSHATGAWVFSN  
ATCKLLKGIYAINFNCGMLLLTCISMDRYIAIVQATKSFR LRSRTLPRSKIICLVVWGL  
SVIISSTFVFNQKYNTQGSDVCEPKYQTVSEPIRWKLLMLGLELLFGFFIPLMFMIFC  
5 YTFIVKTLVQAQNSKRHKAIRVIIAVVLVFLACQIPHNMVLLVTAANLGKMNRSCQS  
EKLIGYTKTVTEVLAFLHCCLNPVLYAFIGQKFRNYFLKILKDLWCVRRKYKSSGFSC  
AGRYSENISRQTSETADNDNASSFTM


[0132] It is understood that the examples and embodiments described herein are for  
10 illustrative purposes only and that various modifications or changes in light thereof will be  
suggested to persons skilled in the art and are to be included within the spirit and purview of  
this application and scope of the appended claims.

[0133] All publications, patents, and patent applications cited herein are hereby  
incorporated by reference in their entirety for all purposes.

15

**WHAT IS CLAIMED IS:**

1. A compound having formula (I):



or a pharmaceutically acceptable salt, hydrate, solvate, or rotamer thereof,  
wherein

$A$  is a carboxylic acid moiety or a carboxylic acid isostere;

ring vertices  $a$ ,  $b$ ,  $c$  and  $d$  are independently selected from  $CH$  and  $C(R^1)$ ;

each  $R^1$  is independently selected from the group consisting of halogen,  $CN$ ,  $-SF_5$ ,  $C_{1-8}$

alkyl,  $C_{3-8}$  cycloalkyl,  $C_{2-8}$  alkenyl,  $C_{2-8}$  alkynyl,  $C_{1-8}$

haloalkyl,  $-OR^a$ ,  $-SR^a$ ,  $-COR^a$ ,  $-NR^aR^b$ , and 5- or 6-membered heteroaryl, wherein

the alkyl portions of  $R^1$  are optionally further substituted with 1-3  $R^a$ ; and

optionally, adjacent  $R^1$  members are connected to form an additional 5- or 6-

membered ring which is saturated or unsaturated having ring vertices selected

from  $C$ ,  $O$ ,  $S$  and  $N$ , wherein the additional 5- or 6-membered ring is optionally

substituted with one or two members selected from the group consisting of

halogen, hydroxyl,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy and  $C_{1-4}$  haloalkyl;

the subscript  $n$  is an integer of from 0 to 3;

each  $R^3$  is independently selected from the group consisting of halogen,  $CN$ ,  $C_{1-8}$  alkyl,

$C_{3-8}$  cycloalkyl,  $C_{2-8}$  alkenyl,  $C_{2-8}$  alkynyl,  $C_{1-8}$  haloalkyl, aryl,  $-OR^a$ ,  $-NR^aR^b$

and  $-N(R^a)-C_1-C_4$  alkylene- $OR^b$ , and wherein the alkyl or aryl portions of  $R^2$  and

$R^3$  are optionally further substituted with 1-3  $R^a$ ;

$Ar$  is a benzene or pyridine ring that is optionally substituted with from 1 to 5  $R^4$

substituents independently selected from the group consisting of halogen,

$CN$ ,  $-SF_5$ ,  $C_{1-8}$  alkyl,  $C_{3-8}$  cycloalkyl,  $C_{2-8}$  alkenyl,  $C_{2-8}$  alkynyl,  $C_{1-8}$  haloalkyl,  $C_{1-8}$

hydroxyalkyl,  $-OR^a$ ,  $-NR^aR^b$ ,

and wherein the alkyl and cycloalkyl, portions of  $R^4$  are optionally further substituted with 1-3  $R^a$ ;


each  $R^a$  and  $R^b$  is independently selected from the group consisting of hydrogen, hydroxyl, halogen, cyano,  $C_{1-4}$  alkyl,  $C_{1-4}$  alkoxy,  $C_{1-4}$  haloalkyl,  $C_{3-6}$  cycloalkyl, amino,  $C_{1-8}$  alkylamino, and di  $C_{1-8}$  alkylamino, or when attached to a nitrogen atom are optionally combined to form a 4- to 7-membered saturated ring, which is optionally substituted with oxo.

2. A compound of claim 1, wherein ring vertex d is CH.
3. A compound of claim 1, wherein ring vertices c and d are each CH.
4. A compound of claim 1, wherein ring vertices a, b, and c are each C(R<sup>1</sup>).
5. A compound of claim 1, wherein ring vertices a and b are each C(R<sup>1</sup>).
6. A compound of claim 1, wherein A is a member selected from the group consisting of -CO<sub>2</sub>H, -SO<sub>3</sub>H, -PO<sub>3</sub>H<sub>2</sub> and a tetrazole.
7. A compound of claim 1, wherein A is -CO<sub>2</sub>H,
8. A compound of claim 1, wherein n is 0 or 1.
9. A compound of claim 1, wherein n is 0.
10. A compound of claim 1, wherein Ar is a benzene ring which is optionally substituted with from 1-3 R<sup>4</sup> substituents; A is -CO<sub>2</sub>H; and d is CH.

11. A compound of claim 10, wherein the subscript n is 1.


12. A compound of claim 10, wherein the subscript n is 0.

13. A compound of claim 1, wherein the ring having a, b, c and d as ring vertices is selected from the group consisting of:




wherein the wavy line labeled y indicates the point of attachment to the NHS(O)<sub>2</sub> portion of the compound and the wavy line labeled z indicates the point of attachment to the quinoline ring.

14. A compound of claim 10, wherein Ar is selected from the group consisting of:



wherein the wavy line labeled w indicates the point of attachment to the S(O)<sub>2</sub> moiety.










15. A compound of claim 13, wherein Ar is selected from the group consisting of:




3

4 wherein the wavy line labeled w indicates the point of attachment to the S(O)<sub>2</sub> moiety.


16. A compound of claim 1, selected from the group consisting of:



and




17. A compound of claim 1, selected from the group consisting of:




or a pharmaceutically acceptable salt, hydrate, solvate, or rotamer thereof.

18. A compound of claim 1, having the structure



or a pharmaceutically acceptable salt thereof.

19. A compound of claim 1, having the structure



or a pharmaceutically acceptable salt thereof.

**20.** A pharmaceutical composition comprising a compound of any of claims **1** to **19**, or a pharmaceutically acceptable salt, hydrate, solvate, or rotamer thereof, with a pharmaceutically acceptable excipient.

**21.** Use of a compound of any of claims **1** to **19**, or a pharmaceutically acceptable salt, hydrate, solvate, or rotamer thereof, in the manufacture of a medicament for treating a disease or condition modulated at least in part by CCR6, wherein said disease or condition is an inflammatory disease or condition.

**22.** The use in accordance with claim **21**, wherein said disease or condition is selected from the group consisting of psoriasis, eczema, atopic dermatitis, allergic contact dermatitis, urticaria, puritis, allergic asthma, Crohn's disease, ulcerative colitis, ileitis, enteritis, insect sting, scleroderma, vaginitis, vasculitis, Grave's disease, Alzheimer's disease, type I diabetes, graft rejection, atherosclerosis, sepsis, sarcoidosis, chronic obstructive pulmonary disease, and sinusitis.

**23.** The use in accordance with claim **21**, wherein said disease or condition is atopic dermatitis.

**24.** The use in accordance with claim **21**, wherein said disease or condition is psoriasis.

**25.** The use in accordance with any of claims **21**, **22**, **23**, or **24**, wherein said compound is a compound of claim **13**, **15**, **16** or **17**, or a pharmaceutically acceptable salt, hydrate, solvate, or rotamer thereof.

**26.** The use in accordance with claim **25**, wherein the compound is a compound of claim **15**, or a pharmaceutically acceptable salt, hydrate, solvate, or rotamer thereof.

**27.** The use in accordance with claim **25**, wherein the compound is a compound of claim **16**, or a pharmaceutically acceptable salt, hydrate, solvate, or rotamer thereof.

**28.** A compound of any of claims **1** to **17**, or a pharmaceutically acceptable salt, hydrate, solvate, or rotamer thereof, for use in treating a disease or condition modulated at least in part by CCR6.