METHOD FOR IDENTIFYING ANTIBIOTIC TARGETS

Field of the Invention

The present invention relates to methods for identifying antibiotic targets in bacteria, to methods for identifying antibiotics and to processes for producing antibiotics and pharmaceutical compositions comprising said antibiotics.

Background to the Invention

There is an urgent need for new antibiotics to counter the emergence of new pathogens and resistance to existing antimicrobial drugs. The identification of the targets of candidate antibiotics is critical, since such information can provide access to a large number of functionally related novel drug families. For example, the discovery of the penicillin-binding proteins as targets of penicillin led to the development of a large family of antibiotics, including multiple generations of cephalosporins, penicillins and carbapenems (see Schmid (2006) Nature Biotechnology 24(4): 419-420).

Transposon directed insertion-site sequencing (TraDIS – see Langridge *et al.* (2009) Genome Research 19: 2308-2316) has recently been described and used to identify: (a) essential genes; (b) genes advantageous (but not essential) for growth; (c) genes disadvantageous for growth under particular conditions; and (d) genes involved in conferring tolerance to certain conditions ("niche-specific" essential genes). Similar techniques have been described in e.g. Gawronski *et al.* (2009) PNAS 106: 16422-16427; Goodman *et al.* (2009) Cell Host Microbe 6: 279-289; van Opijnen *et al.* (2009) Nat. Methods 6: 767-772 and Gallagher *et al.* (2011) mBio 2(1):e00315-10, and such techniques are now collectively dubbed "Tn-seq" methods.

However, an important class of antibiotic targets are gene products involved in cellular processes essential for viability in the growth conditions used. Such targets cannot be identified by Tn-seq (including TraDIS), since transposon insertions into essential genes (including those serving as antibiotic targets) are not significantly represented in the initial mutant pool. Thus, differences in transposon distribution after growth of the mutant pool with or without (or with varying amounts of) antibiotic would not arise, with the result that

Tn-seq cannot distinguish between an essential gene and an essential gene serving as an antibiotic target.

There is therefore a need for high-throughput functional screens for antibiotic targets which are capable of identifying essential genes serving as antibiotic targets.

Summary of the Invention

According to an aspect of the present invention, there is provided a method for identifying an essential gene which serves as an antibiotic target in a bacterium, the method comprising the steps of:

(a) generating a pool of mutant bacteria by transposon mutagenesis with an activating transposon (Tn_A), wherein the Tn_A comprises a promoter such that transposon insertion into bacterial DNA increases the transcription of a gene at or near the insertion site;
(b) growing bacteria from the mutant pool in the presence of different amounts of said antibiotic to produce two or more test cultures; and
(c) comparing the distribution of Tn_A insertions between test cultures to identify a putative essential gene serving as a target of said antibiotic in said bacterium.

The method may further comprise the steps of: generating an antibiotic resistant mutant of said bacterium by a method comprising the step of selecting for growth in the presence of said antibiotic to produce an antibiotic resistant mutant clone (Ab^R mutant); transforming the Ab^R mutant with: (i) one or more essential genes of said bacterium; and (ii) a transposon which insertionally inactivates bacterial DNA, to produce a pool of transposon mutants which are merodiploid for said one or more essential genes; growing bacteria from the merodiploid pool in the presence of different amounts of said antibiotic to produce two or more test cultures; and comparing the distribution of transposon insertions between test cultures to identify a putative essential gene serving as a target of said antibiotic in said bacterium.

In another aspect, there is provided a method of identifying an antibiotic comprising identifying an essential gene which serves as a target of said antibiotic according to a method of the invention.

In a further aspect, there is provided a process for producing an antibiotic comprising identifying an antibiotic by a method comprising identifying an essential gene which serves as a target of said antibiotic according to a method of the invention. Such a process may optionally further comprise the step of synthesising said antibiotic, and may optionally further comprise mixing the synthesised antibiotic with a pharmaceutically acceptable excipient to produce a pharmaceutical composition.

The use of an activating transposon ensures that transposon insertions into essential genes are represented in the initial mutant pool, since transposon insertion can now result in gene activation rather than insertional inactivation. Thus, the effect of the presence of antibiotic during subsequent culture of the mutant pool on transposon distribution can be studied (and the identity of the gene target(s) thereby determined).

Other aspects and preferred embodiments of the invention are defined and described in the other claims set out below.

Detailed Description of the Invention

All publications, patents, patent applications and other references mentioned herein are hereby incorporated by reference in their entireties for all purposes as if each individual publication, patent or patent application were specifically and individually indicated to be incorporated by reference and the content thereof recited in full.

Definitions and general preferences

Where used herein and unless specifically indicated otherwise, the following terms are intended to have the following meanings in addition to any broader (or narrower) meanings the terms might enjoy in the art:

Unless otherwise required by context, the use herein of the singular is to be read to include the plural and *vice versa*. The term "a" or "an" used in relation to an entity is to be read to refer to one or more of that entity. As such, the terms "a" (or "an"), "one or more," and "at least one" are used interchangeably herein.

As used herein, the term "comprise," or variations thereof such as "comprises" or "comprising," are to be read to indicate the inclusion of any recited integer (e.g. a feature, element, characteristic, property, method/process step or limitation) or group of integers (e.g. features, element, characteristics, properties, method/process steps or limitations) but not the exclusion of any other integer or group of integers. Thus, as used herein the term "comprising" is inclusive or open-ended and does not exclude additional, unrecited integers or method/process steps.

The term *gene* is a term describing a hereditary unit consisting of a sequence of DNA that occupies a specific location on a chromosome or plasmid and determines a particular characteristic in an organism. A gene may determine a characteristic of an organism by specifying a polypeptide chain that forms a protein or part of a protein (structural gene); or encode an RNA molecule; or regulate the operation of other genes or repress such operation; or affect phenotype by some other as yet undefined mechanism.

The terms *genomic DNA* is a term of art used herein to define chromosomal DNA as distinct from extrachromosomally-maintained plasmid DNA.

The term *genome* is a term of art used herein to define the entire genetic complement of an organism, and so includes chromosomal, plasmid, prophage and any other DNA.

The term *Gram-positive bacterium* is a term of art defining a particular class of bacteria that are grouped together on the basis of certain cell wall staining characteristics.

The term *low G+C Gram-positive bacterium* is a term of art defining a particular subclass class of evolutionarily related bacteria within the Gram-positives on the basis of the composition of the bases in the DNA. The subclass includes *Streptococcus* spp., *Staphylococcus* spp., *Listeria* spp., *Bacillus* spp., *Clostridium* spp., *Enterococcus* spp. and *Lactobacillus* spp.).

The term *high G+C Gram-positive bacterium* is a term of art defining a particular subclass class of evolutionarily related bacteria within the Gram-positives on the basis of the composition of the bases in the DNA. The subclass includes actinomycetes (actinobacteria) including *Actinomyces* spp., *Arthrobacter* spp., *Corynebacterium* spp.,

Frankia spp., *Micrococcus* spp., *Micromonospora* spp., *Mycobacterium* spp., *Nocardia* spp., *Propionibacterium* spp. and *Streptomyces* spp.

The term *Gram-negative bacterium* is a term of art defining a particular class of bacteria that are grouped together on the basis of certain cell wall staining characteristics. Examples of Gram-negative bacterial genera include *Klebsiella*, *Acinetobacter*, *Escherichia*, *Pseudomonas*, *Enterobacter* and *Neisseria*.

As used herein, the term "essential gene" is a term of art defining a particular class of genes the products of which are necessary for viability, either under all conditions or under the conditions of growth used. An important subclass of essential gene are those encoding products (e.g. proteins, peptides and regulatory polynucleotides) which contribute to metabolic processes essential for viability under important growth conditions (for example, and in the case of pathogenic bacteria, under conditions which prevail during infection or multiplication in the host).

Antibiotics and antibiotic targets

The antibiotic used to produce the test cultures of the invention is typically a novel investigational antibiotic (anti-bacterial chemotherapeutic agent), the mechanism of action (and hence biological target(s)) of which are unknown. In many applications, the antibiotic is selected from combinatorial libraries, natural product libraries, defined chemical entities, peptides, peptide mimetics and oligonucleotides.

The antibiotic target identified according to the invention is an essential gene/gene product, and may therefore be involved in one or more of the following biological processes in the bacterial host:

(a) cell division;
(b) DNA replication (including polymerization and supercoiling);
(c) transcription (including priming, elongation and termination);
(d) translation (including ribosome components, initiation, elongation and release);
(e) biosynthetic pathways (including peptidoglycan and fatty acids);
(f) plasmid addiction;
(g) cell wall assembly; and/or

(h) bacterial cell integrity.

Bacteria for use in the methods of the invention

The methods of the invention may be applied to identify an antibiotic target in any bacterium. Thus, the methods of the invention find application in the identification of antibiotic targets in: (a) Gram-positive, Gram-negative and/or Gram-variable bacteria; (b) spore-forming bacteria; (c) non-spore forming bacteria; (d) filamentous bacteria; (e) intracellular bacteria; (f) obligate aerobes; (g) obligate anaerobes; (h) facultative anaerobes; (i) microaerophilic bacteria and/or (f) opportunistic bacterial pathogens.

In certain embodiments, the methods of the invention are applied to identify an antibiotic target in bacteria of the following genera: *Acinetobacter (e.g. A. baumannii)*; *Aeromonas* (e.g. *A. hydrophila*); *Bacillus* (e.g. *B. anthracis*); *Bacteroides* (e.g. *B. fragilis*); *Bordetella* (e.g. *B. pertussis*); *Borrelia* (e.g. *B. burgdorferi*); *Brucella* (e.g. *B. abortus, B. canis, B. melitensis* and *B. suis*); *Burkholderia* (e.g. *B. cepacia* complex); *Campylobacter* (e.g. *C. jejuni*); *Chlamydia* (e.g. *C. trachomatis, C. suis* and *C. muridarum*); *Chlamydophila* (e.g. (e.g. *C. pneumoniae, C. pecorum, C. psittaci, C. abortus, C. felis and C. caviae*); *Citrobacter* (e.g. *C. freundii*); *Clostridium* (e.g. *C. botulinum, C. difficile, C. perfringens* and *C. tetani*); *Corynebacterium* (e.g. *C. diphteriae* and *C. glutamicum*); *Enterobacter* (e.g. *E. cloacae* and *E. aerogenes*); *Enterococcus* (e.g. *E. faecalis* and *E. faecium*); *Escherichia* (e.g. *E. coli*); *Flavobacterium*; *Francisella* (e.g. *F. tularensis*); *Fusobacterium* (e.g. *F. necrophorum*); *Haemophilus* (e.g. *H. somnus, H. influenzae* and *H. parainfluenzae*); *Helicobacter* (e.g. *H. pylori*); *Klebsiella* (e.g. *K. oxytoca* and *K. pneumoniae*), *Legionella* (e.g. *L. pneumophila*); Leptospira (e.g. *L. interrogans*); *Listeria* (e.g. *L. monocytogenes*); *Moraxella* (e.g. *M. catarrhalis*); *Morganella* (e.g. *M. morganii*); *Mycobacterium* (e.g. *M. leprae* and *M. tuberculosis*); *Mycoplasma* (e.g. *M. pneumoniae*); *Neisseria* (e.g. *N. gonorrhoeae* and *N. meningitidis*); *Pasteurella* (e.g. *P. multocida*); *Peptostreptococcus*; *Prevotella*; *Proteus* (e.g. *P. mirabilis* and *P. vulgaris*), *Pseudomonas* (e.g. *P. aeruginosa*); *Rickettsia* (e.g. *R. rickettsii*); *Salmonella* (e.g. serotypes . Typhi and Typhimurium); *Serratia* (e.g. *S. marcesens*); *Shigella* (e.g. *S. flexnaria, S. dysenteriae* and *S. sonnei*); *Staphylococcus* (e.g. *S. aureus, S. haemolyticus, S. intermedius, S. epidermidis* and *S. saprophyticus*); *Stenotrophomonas* (e.g. *S. maltophila*); *Streptococcus* (e.g. *S. agalactiae, S. mutans, S. pneumoniae* and *S. pyogenes*); *Treponema* (e.g. *T. pallidum*); *Vibrio* (e.g. *V. cholerae*) and *Yersinia* (e.g. *Y. pestis*).

The methods of the invention may be used to identify an antibiotic target in multi-drug resistant bacteria, including, but not limited to penicillin-resistant, methicillin-resistant, quinolone- resistant, macrolide-resistant, and/or vancomycin-resistant bacterial strains, including for example penicillin-, methicillin-, macrolide-, vancomycin-, and/or quinolone-resistant *Streptococcus pneumoniae*; penicillin-, methicillin-, macrolide-, vancomycin-, and/or quinolone-resistant *Staphylococcus aureus*; penicillin-, methicillin-, macrolide-, vancomycin-, and/or quinolone-resistant *Streptococcus pyogenes*; and penicillin-, methicillin- , macrolide-, vancomycin-, and/or quinolone-resistant enterococci.

Thus, methods of the invention may be used to identify an antibiotic target in methicillin-resistant *Staphylococcus aureus* (MRSA), for example selected from any of C-MSRA1, C-MRSA2, C-MRSA3, C-MSRA4, Belgian MRSA, Swiss MRSA and any of the EMRSA strains.

The compounds of the invention may be used to identify an antibiotic target in both high G+C Gram-positive bacteria and in low G+C Gram-positive bacteria.

The methods of the invention find particular application in the identification of an antibiotic target in a bacterium selected from *Klebsiella pneumoniae*, *Acinetobacter baumanii*, *Escherichia coli* (including ST131), *Enterococcus faecalis*, *Enterococcus faecium*, *Pseudomonas aeruginosa*, *Enterobacter cloacae*, *Enterobacter aerogenes* and *Neisseria gonorrhoeae*.

Particularly preferred are methods of identifying an antibiotic target in *Klebsiella pneumoniae*, *Acinetobacter baumanii* or *Escherichia coli*.

Mutant pools

The methods of the invention involve generating a pool of mutant bacteria by transposon mutagenesis. The size of the mutant pool affects the resolution of the method: as the pool size increases, more and more different genes with Tn_A insertions will be represented (and so effectively assayed). As the pool size decreases, the resolution of the method reduces, genes will be less effectively assayed, and more and more genes will not be assayed at all.

Ideally, the mutant pool generated in the methods of the invention is *comprehensive*, in the sense that insertions into every gene are represented. The number of Tn_A insertion mutants (i.e. the mutant pool size) required to achieve this depends on various factors, including: (a) the size of the bacterial genome; (b) the average size of the genes; and (c) any Tn_A insertion site bias.

With regard to the latter, some areas of bacterial genomes attract a low frequency of insertion (especially GC-rich regions). Thus, insertion frequencies and pool sizes large enough to ensure that insertions into insertion-refractory regions are preferred.

In general, a minimum insertion rate of one transposon per 25bp is required to achieve a comprehensive pool/library, which typically entails a minimum pool size for bacteria having a genome size of 4 to 7 Mb of 0.5×10^5 to 1×10^5, for example 5×10^5, preferably at least about 1×10^6 mutants. In many cases, 1×10^6 mutants will allow identification of ~300,000 different insertion sites and correspond to 1 transposon insertion every 13 to 23 bp (or about 40-70 different insertion sites per gene).

However, the methods of the invention do not necessarily require a comprehensive mutant pool (in the sense defined above) in order to return useful information as to the identity of antibiotic drug targets. Rather, pool sizes less than the ideal comprehensive pool may be used, provided that a reduction in resolution (and attendant failure to assay certain genes) can be tolerated. This may be the case, for example, where the method is designed to be run iteratively until the target is identified: in such embodiments the effective pool size grows with each iteration of the method.

Transposon mutagenesis

Transposons, sometimes called transposable elements, are polynucleotides capable of inserting copies of themselves into other polynucleotides. The term transposon is well known to those skilled in the art and includes classes of transposons that can be distinguished on the basis of sequence organisation, for example short inverted repeats at each end; directly repeated long terminal repeats (LTRs) at the ends; and polyA at 3'ends of RNA transcripts with 5' ends often truncated.

Transposomes are transposase-transposon complexes wherein the transposon does not encode transposase. Thus, once inserted the transposon is stable. Preferably, in order to ensure mutant pool stability, the transposon does not encode transposase and is provided in the form of a transposome (i.e. as a complex with transposase enzyme), as described below.

As used herein, the term "activating transposon" (hereinafter abbreviated "Tn_A") defines a transposon which comprises a promoter such that transposon insertion increases the transcription of a gene at or near the insertion site. Examples of such transposons are described in Troeschel *et al.* (2010) Methods Mol Biol. 668:117-39 and Kim *et al.* (2008) Curr Microbiol. 57(4): 391-394.

The activating transposon/transposome can be introduced into the bacterial genome (including chromosomal and/or plasmid DNA) by any of a wide variety of standard procedures which are well-known to those skilled in the art. For example, Tn_A transposomes can be introduced by electroporation (or any other suitable transformation method).

Preferably, the transformation method generates $1x10^3$ to $5x10^3$ transformants/ng DNA, and such transformation efficiencies are generally achievable using electroporation.

Alternatively, transposon mutagenesis using Tn_A may be performed *in vitro* and recombinant molecules transformed/transfected into bacterial cells. In such embodiments, transposomes can be prepared according to a standard protocol by mixing commercially available transposase enzyme with the transposon DNA fragment. The resulting transposomes are then mixed with plasmid DNA of the plasmid of interest to allow transposition, then the DNA introduced into a host bacterial strain using electrotransformation to generate a pool of plasmid transposon mutants.

In embodiments where mutagenesis is performed *in vitro*, it is possible to mix transposomes with genomic DNA *in vitro* and then introduce the mutagenized DNA (optionally, after fragmentation and/or circularization) into the host bacterial strain (e.g. by electroporation) whereupon endogenous recombination machinery incorporates it into the genome. Such an approach may be particularly useful in the case of bacteria which are

naturally competent (e.g. *Acinetobacter* spp.) and/or can incorporate DNA *via* homologous crossover (e.g. double crossover) recombination events.

Activating transposons for use in the methods of the invention

Any suitable activating transposon may be used in the methods of the invention. Suitable transposons include those based on Tn*3* and the Tn*3*-like (Class II) transposons including γδ (Tn*1000*), Tn*501*, Tn*2501*, Tn*21*, Tn*917* and their relatives. Also Tn*10*, Tn*5*, Tn*phoA*, Tn*903*, bacteriophage Mu and related transposable bacteriophages. A variety of suitable transposons are also available commercially, including for example the EZ-Tn5™ < R6Kγori/KAN-2> transposon.

Preferred transposons are those which carry antibiotic resistance genes (which may be useful in identifying mutants which carry a transposon) including Tn*5*, Tn*10* and Tn*phoA*. For example, Tn*10* carries a tetracycline resistance gene between its IS elements while Tn*5* carries genes encoding polypeptides conferring resistance to kanamycin, streptomycin and bleomycin. Other suitable resistance genes include those including chloramphenicol acetyltransferase (conferring resistance to chloramphenicol).

It is of course possible to generate new transposons by inserting different combinations of antibiotic resistance genes between IS elements, or by inserting combinations of antibiotic resistance genes between transposon mosaic ends (preferred), or by altering the polynucleotide sequence of the transposon, for example by making a redundant base substitution or any other type of base substitution that does not affect the transposition or the antibiotic resistance characteristics of the transposon, in the coding region of an antibiotic resistance gene or elsewhere in the transposon. Such transposons are included within the scope of the invention.


In many embodiments, a single transposon is used to generate the mutant pool. However, as explained above, the number of Tn insertion mutants (i.e. the mutant pool size) required to achieve a comprehensive pool or library depends *inter alia* on any Tn insertion site bias. Thus, in cases where the transposon insertion site bias occurs, two or more different transposons may be used in order to reduce or eliminate insertion site bias. For example, a combination of two different transposons based on Tn*5* and Tn*10* may be employed.

Promoters for use in activating transposons

The nature of the promotor present in the Tn_A is dependent on the nature of the transposon and the ultimate bacterial host. Generally, an efficient, outward-oriented promoter which drives high level transcription of DNA near or adjacent to the insertion site is chosen.

The promoter may include: (a) a Pribnow box (-10 element); (b) a –35 element and/or (c) an UP element.

For example, the *lac* promoter can be used with the EZ-Tn5™ < R6Kγori/KAN-2> transposon, and such constructs are suitable for assay of e.g. *Escherichia coli*, *Enterobacter* spp. and other members of the family *Enterobacteriaceae* such as *Klebsiella* spp. Other suitable promoters include: *rplJ* (large ribosomal subunit protein; moderate strength promoter); *tac* (artificial *lac/trp* hybrid; strong promoter) and *rrn*B (ribosomal RNA gene promoter; very strong promoter). The sequences of the latter promoters are shown below:



Determing the distribution of Tn_A insertions

The distribution of transposon insertions is preferably determined by sequencing bacterial DNA adjacent or near (5' and/or 3') the Tn_A insertion site (e.g. by sequencing DNA which comprises Tn_A -genomic DNA junctions). Typically, bacterial DNA flanking or adjacent to one or both ends of the Tn_A is sequenced.

The length of adjacent DNA sequenced need not be extensive, and is preferably relatively short (for example, less than 200 base pairs).

Various methods can be used to determine the Tn_A insertion distribution using DNA sequencing: such methods have recently been dubbed Tn-seq procedures (van Opijnen *et al.* (2009) Nat. Methods 6: 767-772). For example, Tn-seq procedures include affinity purification of amplified Tn junctions (Gawronski *et al.* (2009) PNAS 106: 16422-16427); ligation of adaptors into genome sequences distal to the end of the transposon using a specialized restriction site (Goodman *et al.* (2009) Cell Host Microbe 6: 279-289; van Opijnen *et al.* (2009) Nat. Methods 6: 767-772); selective amplification (Langridge *et al.* (2009) Genome Research 19: 2308-2316) and the generation of single-stranded DNA circles bearing Tn junctions, which serve as templates for amplification and sequencing after elimination of genomic DNA by exonuclease digestion (Gallagher *et al.* (2011) mBio 2(1):e00315-10).

Any suitable high-throughput sequencing technique can be used, and there are many commercially available sequencing platforms that are suitable for use in the methods of the invention. Sequencing-by-synthesis (SBS)-based sequencing platforms are particularly suitable for use in the methods of the invention: for example, the Illumina™ system is generates millions of relatively short sequence reads (54, 75 or 100bp) and is particularly preferred.

Other suitable techniques include methods based on reversible dye-terminators. Here, DNA molecules are first attached to primers on a slide and amplified so that local clonal colonies are formed (bridge amplification). Four types of ddNTPs are added, and non-incorporated nucleotides are washed away. Unlike pyrosequencing, the DNA can only be extended one nucleotide at a time. A camera takes images of the fluorescently labeled nucleotides then the dye along with the terminal 3' blocker is chemically removed from the DNA, allowing a next cycle.

Other systems capable of short sequence reads include SOLiD™ and Ion Torrent technologies (both sold by Applied Biosystems™). SOLiD™ technology employs sequencing by ligation. Here, a pool of all possible oligonucleotides of a fixed length are labeled according to the sequenced position. Oligonucleotides are annealed and ligated; the preferential ligation by DNA ligase for matching sequences results in a signal informative of the nucleotide at that position. Before sequencing, the DNA is amplified by emulsion PCR. The resulting bead, each containing only copies of the same DNA

molecule, are deposited on a glass slide. The result is sequences of quantities and lengths comparable to Illumina sequencing.

Ion Torrent Systems Inc. have developed a system based on using standard sequencing chemistry, but with a novel, semiconductor based detection system. This method of sequencing is based on the detection of hydrogen ions that are released during the polymerisation of DNA, as opposed to the optical methods used in other sequencing systems. A microwell containing a template DNA strand to be sequenced is flooded with a single type of nucleotide. If the introduced nucleotide is complementary to the leading template nucleotide it is incorporated into the growing complementary strand. This causes the release of a hydrogen ion that triggers a hypersensitive ion sensor, which indicates that a reaction has occurred. If homopolymer repeats are present in the template sequence multiple nucleotides will be incorporated in a single cycle. This leads to a corresponding number of released hydrogens and a proportionally higher electronic signal.

Functional assessment of putative essential genes

The putative essential gene identified by comparing the distribution of Tn_A insertions between test cultures may be further characterized by various techniques which directly or indirectly assess its function. In this way, an essential function may be definitively assigned to said putative essential gene.

Suitable techniques include bioinformatics, where the (full or partial) sequence of the putative essential gene is used to interrogate sequence databases containing information from the bacterium assayed and/or other species in order to identify genes (e.g. orthologous genes in other species) for which essential biochemical function(s) have already been assigned and/or which have been shown to be essential.

Suitable bioinformatics programs are well known to those skilled in the art and include the Basic Local Alignment Search Tool (BLAST) program (Altschul *et al.* (1990) J. Mol. Biol. 215: 403-410 and Altschul *et al.* (1997) Nucl. Acids Res. 25: 3389-3402). Suitable databases include, for example, EMBL, GENBANK, TIGR, EBI, SWISS-PROT and trEMBL.

Alternatively, or in addition, the (full or partial) sequence of the putative essential gene is used to interrogate a sequence database containing information as to the identity of essential genes which has been previously constructed using the conventional Tn-seq methods described in the prior art (e.g. as described in Gawronski *et al.* (2009) PNAS 106: 16422-16427; Goodman *et al.* (2009) Cell Host Microbe 6: 279-289; van Opijnen *et al.* (2009) Nat. Methods 6: 767-772; Langridge *et al.* (2009) Genome Research 19: 2308-2316; Gallagher *et al.* (2011) mBio 2(1):e00315-10) and/or the techniques described in WO 01/07651 (the contents of which are hereby incorporated by reference).

Alternatively, or in addition, essentiality can be imputed by eliminating the possibility that a putative essential gene acts as an antibiotic resistance gene. For example, the (full or partial) sequence of the putative essential gene is used to interrogate sequence databases containing sequence information of genes previously identified as antibiotic resistance genes using the Tn-seq methods described in e.g. Gawronski *et al.* (2009) PNAS 106: 16422-16427; Goodman *et al.* (2009) Cell Host Microbe 6: 279-289; Langridge *et al.* (2009) Genome Research 19: 2308-2316 or Gallagher *et al.* (2011) mBio 2(1):e00315-10. Antibiotic resistance genes may be identified in such methods as a class of niche-specific/conditionally essential genes.

Despite the presence of a promoter within the inserted sequence, many Tn_A insertions will disrupt gene/DNA function and allow identification of essential/important DNA regions, as in standard Tn-seq (including TraDIS). However, some transposons will be positioned appropriately with respect to specific important DNA regions, whereby transcription of those specific regions, driven by the inserted promoter, is enhanced significantly compared to endogenous transcription. By growing the mutant pool in increasing antibiotic concentrations and repeating the sequencing it is possible to observe changes in the number of reads, indicating not only which DNA region contributes to antibiotic survival, but also the relative contribution. The higher levels of specific antibiotic target transcription (driven by the transposon-inserted promoters) will favour bacterial survival in antibiotic and link insertion site to DNA region by proximity.

To identify the specific antibiotic target(s), the position of the inserted promoter can be assessed with respect to its contribution to increased transcription of relevant downstream DNA sequences. A mathematically/technically straightforward bioinformatics component of this technique permits recognition of the contribution of the inserted promoter sequence to

transcription of the putative antibiotic target gene. For example, transcription of the antibiotic target partial gene product may be enough to confer antibiotic resistance and bioinformatic analysis would allow an explanation. For example, the partial gene transcript may still encode enough information to allow translation of a truncated, but functional essential protein. Bioinformatics would allow the effects of transcriptional read through on genes downstream of the gene adjacent to the inserted transposon to be considered, where there is there no defined RNA transcription termination sequence.

For example, a transposon/promoter upstream of genes A, B and C may generate a polycistronic transcript of all three genes (A-C), upstream of B a polycistronic transcript of genes B and C and upstream of C just gene C. If the reads for the first two transposons were high and the third low in antibiotic then the antibiotic target would be gene B.

Ancillary analytic methods

The methods of the invention find particular application in cases where the antibiotic target is overexpressed to a level where it contributes and plays a significant role in the binding kinetics to antibiotic at the levels used for target deconvolution. The methods are therefore ideally suited to antibiotic targets where the antibiotic binds to and alters the function of a monomeric macromolecule. However, sensitivity may be reduced in cases where the antibiotic target is available only in a ternary complex, when additional target supplied from overexpression may not function as an effective sink and so alter the effect of the antibiotic.

Thus, in some circumstances the methods of the invention may be used in conjunction with other, complementary, techniques for identifying essential, conditionally essential, non-essential and/or essential genes serving as targets for antibiotics, as described below:

(a) Complemented sequencing

The method of the invention may optionally further comprise the steps of:

(a) generating an antibiotic resistant mutant of said bacterium by a method comprising the step of selecting for growth in the presence of said antibiotic to produce an antibiotic resistant mutant clone (Ab^R mutant);

(b) transforming the Ab^R mutant with: (i) one or more essential genes of said bacterium; and (ii) a transposon which insertionally inactivates bacterial DNA, to produce a pool of transposon mutants which are merodiploid for said one or more essential genes;

(c) growing bacteria from the merodiploid pool in the presence of different amounts of said antibiotic to produce two or more test cultures; and

(d) comparing the distribution of transposon insertions between test cultures to identify a putative essential gene serving as a target of said antibiotic in said bacterium.

The use of transposon mutant pools generated from antibiotic resistant mutants which are merodiploid for one or more essential genes ensures that transposon insertions into essential genes (including those serving as antibiotic targets) are represented in the initial mutant pool, since transposon insertions into the antibiotic target gene yield viable phenotypes under non-selective conditions (when the wild type copy of the essential gene complements the insertionally inactivated mutant copy), but not under selective conditions (when the wild type copy does not complement the insertionally inactivated mutant copy).

Thus, differences in transposon distribution after growth of the merodiploid mutant pool with or without antibiotic can be readily detected, permitting identification of essential genes which serves as antibiotic targets.

In the optional further steps set out above:

- the merodiploid pool may comprise at least 0.5×10^5 mutants, for example at least 1×10^5 mutants;
- the merodiploid pool may comprise at least 5×10^5 mutants;
- the merodiploid pool may comprise at least 1×10^6 mutants;
- the merodiploid pool may comprises 0.5×10^6 to 2×10^6 mutants;
- the merodiploid pool may comprise about 1×10^6 mutants;
- transformation with the transposon in step (b) may yield an insertion rate of at least one transposon per 50 base pairs of bacterial DNA, at least one transposon per 30 base pairs of bacterial DNA, at least one transposon per 25 base pairs of bacterial DNA, at least one transposon per 15 base pairs of bacterial DNA or at least one transposon per 10 base pairs of bacterial DNA;
- the bacterial DNA of step (b) may be chromosomal DNA, plasmid DNA or a mixture of chromosomal and plasmid DNA, and may comprise the entire bacterial genome;

- the Ab^R mutant may be generated by a method which further comprises mutagenizing said bacterium prior to selecting for growth in the presence of said antibiotic;
- the mutagenizing step may be: (a) chemical mutagenesis; and/or (b) radiation mutagenesis;
- the bacterium may be a Gram-positive bacterium;
- the bacterium may be selected from *Enterococcus faecalis*, *Enterococcus faecium* and *Neisseria gonorrhoeae;*
- the bacterium may be a Gram-negative bacterium;
- the bacterium may be selected from: *Klebsiella pneumoniae*, *Acinetobacter baumanii*, *Escherichia coli*, *E. coli* ST131 strains, *Pseudomonas aeruginosa*, *Enterobacter cloacae*, *Enterobacter aerogenes* and *Neisseria gonorrhoeae;*
- the bacteria may be grown from the merodiploid pool in step (c) by inoculating growth medium with 10^7 to 10^9, for example about 10^8, cfu from the merodiploid pool;
- in step (c) at least two test cultures may be produced, one grown in the absence of antibiotic and one grown in the presence of antibiotic (for example at a concentration of about 1 to about 4 x MIC);
- the distribution of transposon insertions between test cultures may be compared by sequencing DNA adjacent or near the transposon insertion site;
- the sequencing of DNA adjacent or near the insertion site may comprise selective amplification of transposon-bacterial DNA junctions;
- the sequencing may comprise sequencing-by-synthesis (SBS) biochemistry;
- about 25, 50, 75, 100 or greater than 100 base pairs of DNA adjacent or near the insertion site may be sequenced;
- the sequenced DNA may be 5' and/or 3' to the insertion site;
- the method may further comprise the step of assigning an essential function to said putative essential gene by sequence comparison with one or more essential gene(s) of said bacterium, for example by transposon directed insertion site sequencing;
- the method may further comprise the step of assigning an essential function to said putative essential gene by determining that it is not an antibiotic resistance gene, for example by a method comprising identifying an antibiotic resistance gene by transposon directed insertion site sequencing using a transposon which inactivates on insertion;

- the one or more essential genes of step (b) may be provided on an expression vector, on an integrative expression vector which inserts into the bacterial chromosome after transformation; on a single or low copy number expression vector which is stably maintained extrachromosomally after transformation;
- the expression vector may be a combination of plasmids containing fragments of the native chromosome of said bacterium or a bacterial artificial chromosome (BAC);
- the one or more essential genes of step (b) may be provided on linear DNA (for example, fragments of genomic DNA of said bacterium);
- the one or more essential genes of step (b) may comprise all or a defined subset of the essential genes of said bacterium;
- the one or more essential genes of step (b) may comprise at least 10, at least 20, at least 50, at least 100, at least 150, at least 200, at least 250 or at least 300 essential genes of said bacterium;
- the one or more essential genes of step (b) may be provided by a method comprising identifying one or more essential genes of said bacterium by transposon directed insertion site sequencing;
- the one or more essential gene(s) of step (b) may be selected from genes involved in: cell division; and/or DNA replication (for example, polymerization or supercoiling); and/or transcription (for example priming, elongation and/or termination); and/or translation (for example genes encoding ribosome components, genes involved in initiation, elongation and/or release); and/or biosynthetic pathways (for example peptidoglycan and/or fatty acid metabolism).
- in step (b) the Ab^R mutant may be transformed simultaneously with the one or more essential genes of said bacterium and the transposon;
- in step (b) the Ab^R mutant may be first transformed with the transposon and then with the one or more essential genes of said bacterium; and/or
- in step (b) the Ab^R mutant may be first transformed with the one or more essential genes of said bacterium and then with the transposon.

When the Ab^R mutant is transformed simultaneously with the one or more essential genes of said bacterium and the transposon, or first transformed with the one or more essential genes of said bacterium and then with the transposon, undesired transposon insertion into the introduced essential genes may occur which can reduce the efficiency of merodiploid formation and complicate the analysis of the data obtained from such libraries.

Such problems are avoided when the Ab^R mutant is first transformed with the transposon and then with the one or more essential genes of said bacterium. However, depending on the efficiency of the process used to introduce the essential gene(s), such a strategy may require a significantly larger number of transformation experiments to give a library of sufficient size. An alternative solution exploits the phenomenon of transposition immunity. Here, undesirable transposition into the introduced essential genes is eliminated (or reduced) by incorporating transposon mosaic end sequences into extrachromosomal DNA (typically, plasmid or BAC) bearing the essential genes used to create the merodiploids. Such a strategy may be used in conjunction with the transposons based on Class II (Tn*3*-like) transposons , for example, Tn3 or its relatives, as described below.

Suitable transposon systems for use in methods which exploit the phenomenon of transposition immunity (as described above) include those based on Tn*3* and its relatives. For example, the transposon delivery plasmid pAMICS2 (see Figure 2) contains the entire Tn*3*-based transposon-generating system (including genes encoding the resolvase and transposase enzymes, TnpR and TnpA) and an origin of replication (*ori*R6K) that is active only when complemented by the *pir* gene, thus preventing propagation in the recipient bacteria after transposition. The plasmid also contains the *mob*RP4 mobilisation origin to allow transfer from a suitable donor strain by conjugation, which is permissive to plasmid replication (i.e. contains the *pir* gene). Inadvertent propagation of the plasmid can also be detected by the presence of a tobramycin resistance gene (*aacA4*).

Other suitable transposons include those based on Tn*3* and the Tn*3*-like (Class II) transposons including γδ (Tn*1000*), Tn*501*, Tn*2501*, Tn*21*, Tn*917* and their relatives. Also Tn*10*, Tn*5*, Tn*phoA*, Tn*903*, bacteriophage Mu and related transposable bacteriophages. A variety of suitable transposons are also available commercially, including for example the EZ-Tn5™ < R6Kγori/KAN-2> transposon.

Preferred transposons are those which carry antibiotic resistance genes (which may be useful in identifying mutants which carry a transposon) including Tn*5*, Tn*10* and Tn*phoA*. For example, Tn*10* carries a tetracycline resistance gene between its IS elements while Tn*5* carries genes encoding polypeptides conferring resistance to kanamycin, streptomycin and bleomycin. Other suitable resistance genes include those including chloramphenicol acetyltransferase (conferring resistance to chloramphenicol).

It is of course possible to generate new transposons by inserting different combinations of antibiotic resistance genes between IS elements, or by inserting combinations of antibiotic resistance genes between transposon mosaic ends (preferred), or by altering the polynucleotide sequence of the transposon, for example by making a redundant base substitution or any other type of base substitution that does not affect the transposition or the antibiotic resistance characteristics of the transposon, in the coding region of an antibiotic resistance gene or elsewhere in the transposon. Such transposons are included within the scope of the invention.

In many embodiments, a single transposon is used to generate the mutant pool. However, as explained above, the number of Tn insertion mutants (i.e. the mutant pool size) required to achieve a comprehensive pool or library depends *inter alia* on any Tn insertion site bias. Thus, in cases where the transposon insertion site bias occurs, two or more different transposons may be used in order to reduce or eliminate insertion site bias. For example, a combination of two different transposons based on Tn*5* and Tn*10* may be employed.

Thus, the method of the invention may optionally further comprise the steps of identifying a gene (for example an essential gene) which serves as an antibiotic target in a bacterium, the method comprising the steps of:

(a) transforming bacteria with an extrachromosamal element (e.g. plasmid or BAC) comprising : (i) one or more essential genes of said bacterium; and (ii) one or more transposon repeat sequences, to produce a pool of bacteria which are merodiploid for said one or more essential genes; and

(b) transforming the merodiploids of step (a) with a transposon delivery plasmid comprising: (i) gene encoding a transposase and a resolvase; and (ii) invert repeat transposase recognition sites;

wherein the one or more transposon repeat sequences of the extrachromosomal element of step (a) confer transposon immunity against the transposon delivered by the plasmid of step (b).

In this aspect of the invention, the transposon delivery system is preferably based on Tn*3*, for example containing the Tn*3* *tnp*A and *tnp*R genes. Preferred are transposon delivery plasmids which further comprise one or more antibiotic resistance gene(s).

(b) Tn-seq

The method of the invention may optionally further comprise the steps of Tn-seq analysis as described in e.g. Gawronski *et al.* (2009) PNAS 106: 16422-16427; Goodman *et al.* (2009) Cell Host Microbe 6: 279-289; Langridge *et al.* (2009) Genome Research 19: 2308-2316 or Gallagher *et al.* (2011) mBio 2(1):e00315-10. When used in combination with Tn-seq analysis, the invention may further identify: (a) essential genes; (b) genes advantageous (but not essential) for growth; (c) genes disadvantageous for growth under particular conditions; and (d) genes involved in conferring tolerance to certain conditions ("niche-specific" essential genes), in addition to essential genes which serve as antibiotic targets.

Exemplification

The invention will now be described with reference to specific Examples. These are merely exemplary and for illustrative purposes only: they are not intended to be limiting in any way to the scope of the monopoly claimed or to the invention described. These examples constitute the best mode currently contemplated for practicing the invention.

Preparation of bacteria for electroporation

Bacteria are grown in 2 x TY broth to an OD_{600} of 0.3–0.5. Cells are then harvested and washed three times in 1/2 original culture volume 10% glycerol and resuspended in 1/1000 original culture volume 10% glycerol and stored at -80°C.

Preparation of transposomes

Transposon DNA (a derivative of EZ-Tn5™ < R6Kγori/KAN-2> possessing an internal *lac* promoter was amplified using oligonucleotides 5'-CTGTCTCTTATACACATCTCCCT and 5'-CTGTCTCTTATACACATCTCTTC with Pfu Ultra Fusion II DNA polymerase

(Stratagene). As an alternative, the internal *lac* promoter can be replaced with a *tac* promoter (as described *supra*). The resultant amplicon was then phosphorylated using T4 polynucleotide kinase (New England Biolabs). Two hundred nanograms of this DNA were then incubated with EZ-Tn5™ transposase (Epicenter Biotechnologies) at 37°C for 1 h then stored at -20°C at a DNA concentration of 20ng/µl.

Generation of mutant bacterial pools

Sixty microliters of cells (previously stored at -80°C are mixed with 0.2µl (4ng) of transposomes and 1 µl (20g) complementing plasmid comprising essential genes and electrotransformed in a 2-mm electrode gap cuvette using a Bio-Rad GenePulser II set to 2.4 kV, 25 µF, and 200 Ω. Cells are resuspended in 1 mL of SOC medium (Invitrogen) and incubated at 37°C for 2 h then spread on L-agar bacteroiological growth medium supplemented with an appropriate concentration of kanamycin. The concentration of kanamycin used is strain dependent and determined empirically

After incubation overnight at 37°C, the number of colonies on several plates is estimated by counting a proportion of them, and from this the total number of colonies on all plates is estimated conservatively. Kanamycin resistant colonies are harvested by resuspension in sterilized deionized water using a bacteriological spreader. Resuspended cells from several electroporations are then pooled to create mutant library mixtures estimated to include over 1 million mutants.

Identifying antibiotic target gene(s)

Eight cultures of 100 ml broth medium are prepared, six of which are supplemented, in duplicate, with the test antibiotic at a concentrations 0.5, 1 and 2 x MIC. Any required promoter inducer is also be added to the medium at this time to ensure active transcription directed into the chromosomal DNA from the transposon sequence.

Assuming a transposon mutant pool of 1 million mutants, 10^8 – 10^9cfu of the pool are used to inoculate each culture. Cultures are grown to stationary phase and cells harvested for genomic DNA extraction. Fresh cultures are also prepared and inoculated with 10^8 – 10^9cfu from the first cultures. These are grown to stationary phase and cells harvested for extraction of genomic DNA.

Genomic DNA is sequenced using the Illumina™ platform incorporating the TraDIS modification to obtain sequence reads initiated from the transposon insertion sites. Sequence reads are then mapped to the bacterial genome sequence and compared with the genome annotation to determine the number of sequence reads that map to each gene for the 8 cultures (6 test and 2 control). Comparison of the control data sets with each other and of test data sets with each other indicates the degree of experimental variation.

Comparison of control data with test data sets shows experimental reproducibility and indicates gene(s) targeted by the antibiotic. Illumina™ sequence reads from transposon insertion within the essential gene antibiotic target gene(s) increase in cells grown with antibiotic, where the promoter caused an increase in this specific gene transcription. Moreover, the relative read count from the target gene(s) increase with concentration of antibiotic used.

Exclusion of antibiotic resistance genes

Conventional transposon directed insertion-site sequencing (TraDIS – see Langridge *et al.* (2009) Genome Research 19: 2308-2316) can be used to identify antibiotic resistance genes which are not essential to growth under normal conditions but which confer tolerance to the antibiotic (i.e. a class of the "niche-specific" essential genes discussed in Langridge *et al.* (2009)). This permits the elimination of antibiotic resistance genes from candidate antibiotic target genes, as described below.

The MIC of the antibiotic to be tested is determined for the bacterium of interest. Four cultures of 100 ml broth medium are prepared, two of which are supplemented with the antibiotic at a concentration 0.5 to 0.75 x MIC (i.e. just below MIC). Assuming a transposon mutant pool of 1 million mutants, 10^8 – 10^9cfu of the pool are used to inoculate each culture. Cultures are grown to stationary phase and cells harvested for genomic DNA extraction. Fresh cultures are also prepared and inoculated with 10^8 – 10^9cfu from the first cultures. These are grown to stationary phase and cells harvested for extraction of genomic DNA. Genomic DNA is sequenced using the Illumina™ platform incorporating the TraDIS modification to obtain sequence reads initiated from the transposon insertion sites. Sequence reads are then mapped to the bacterial genome sequence and compared with

the genome annotation to determine the number of sequence reads that map to each gene for the 4 cultures (2 test and 2 control).

Comparison of the control data sets with each other and of test data sets with each other indicates the degree of experimental variation. Comparison of control data with test data sets shows experimental reproducibility and indicates those genes involved in resistance.

Figure 1 shows the results of a pilot study to identify genes that contribute to ciprofloxacin resistance in *Salmonella* Typhi. The graph includes data for every non-essential gene in the bacterium's genome. The transposon insertion library was grown in 4 conditions: 2 control cultures (no antibiotic) and 2 cultures with sub-optimal ciprofloxacin. Each point represents a gene and each gene is plotted 3 times (ctrl1 v ctrl 2 & CIP1 v CIP2 = black and indicates the degree of experimental variation; CIP mean v ctrl mean = grey; grey points that plot beyond the cluster of black control points represent genes for which data shows significant differences). In Figure 1, grey points below the diagonal cluster of black points are genes that contribute to resistance. The further from the black cluster the grey points are, the more significant the data. Genes that are known to contribute to ciprofloxacin resistance in *Salmonella* are found in this region of the graph, as well as genes not previously known to contribute to resistance. Grey points above the black cluster are genes that contribute to sensitivity. Again, genes known to contribute to sensitivity are found in this region of the graph, and this data identifies genes not previously known to contribute to sensitivity. Data are generally sufficiently clear so as not to require statistical analysis.

Equivalents

The foregoing description details presently preferred embodiments of the present invention. Numerous modifications and variations in practice thereof are expected to occur to those skilled in the art upon consideration of these descriptions. Those modifications and variations are intended to be encompassed within the claims appended hereto.

CLAIMS:

1. A method for identifying an essential gene which serves as an antibiotic target in a bacterium, the method comprising the steps of:

(a) generating a pool of mutant bacteria by transposon mutagenesis with an activating transposon (Tn_A), wherein the Tn_A comprises a promoter such that transposon insertion into bacterial DNA increases the transcription of a gene at or near the insertion site;

(b) growing bacteria from the mutant pool in the presence of different amounts of said antibiotic to produce two or more test cultures; and

(c) comparing the distribution of Tn_A insertions between test cultures to identify a putative essential gene serving as a target of said antibiotic in said bacterium.

2. The method of claim 1 wherein the pool of mutant bacteria comprises at least 0.5×10^5 mutants, for example at least 1×10^5 mutants.

3. The method of claim 1 wherein the pool of mutant bacteria comprises at least 5×10^5 mutants.

4. The method of claim 1 wherein the pool of mutant bacteria comprises at least 1×10^6 mutants.

5. The method of claim 1 wherein the pool of mutant bacteria comprises 0.5×10^6 to 2×10^6 mutants.

6. The method of claim 5 wherein the pool of mutant bacteria comprises about 1×10^6 mutants.

7. The method of any one of the preceding claims wherein the transposon mutagenesis step (a) yields an insertion rate of at least one transposon per 50 base pairs of bacterial DNA.

8. The method of any one of the preceding claims wherein the transposon mutagenesis step (a) yields an insertion rate of at least one transposon per 30 base pairs of bacterial DNA.

9. The method of any one of the preceding claims wherein the transposon mutagenesis step (a) yields an insertion rate of at least one transposon per 25 base pairs of bacterial DNA.

10. The method of any one of the preceding claims wherein the transposon mutagenesis step (a) yields an insertion rate of at least one transposon per 15 base pairs of bacterial DNA.

11. The method of any one of the preceding claims wherein the transposon mutagenesis step (a) yields an insertion rate of at least one transposon per 10 base pairs of bacterial DNA.

12. The method of any one of the preceding claims wherein the bacterial DNA of step (a) is chromosomal (genomic) DNA.

13. The method of any one of claims 1 to 11 wherein the bacterial DNA of step (a) is plasmid DNA or a mixture of chromosomal (genomic) and plasmid DNA.

14. The method of any one of the preceding claims wherein the transposon mutagenesis of step (a) occurs *in vivo*.

15. The method of any one of claims 1 to 13 wherein the transposon mutagenesis of step (a) occurs *in vitro*.

16. The method of any one of the preceding claims wherein the bacterium is a Gram-positive bacterium.

17. The method of claim 16 wherein the bacterium is selected from *Enterococcus faecalis*, *Enterococcus faecium* and *Neisseria gonorrhoeae*.

18. The method of any one of claims 1 to 15 wherein the bacterium is a Gram-negative bacterium.

19. The method of claim 18 wherein the bacterium is selected from: *Klebsiella pneumoniae*, *Acinetobacter baumanii*, *Escherichia coli*, *E. coli* ST131 strains, *Pseudomonas aeruginosa*, *Enterobacter cloacae*, *Enterobacter aerogenes* and *Neisseria gonorrhoeae*.

20. The method of any one of the preceding claims wherein bacteria are grown from the mutant pool in step (b) by inoculating growth medium with 10^7 to 10^9, for example about 10^8, cfu from the mutant pool.

21. The method of any one of the preceding claims wherein bacteria are grown from the mutant pool in step (b) in the presence of antibiotic at a concentration of about 0.5, about 1 and about 2 x MIC to produce at least three test cultures.

22. The method of any one of the preceding claims wherein the distribution of Tn_A insertions between test cultures is compared by sequencing DNA adjacent or near the insertion site of the Tn_A.

23. The method of claim 22 wherein the sequencing of DNA adjacent or near the insertion site of the Tn_A comprises selective amplification of transposon-bacterial DNA junctions.

24. The method of claim 22 or claim 23 wherein the sequencing comprises sequencing-by-synthesis (SBS) biochemistry.

25. The method of any one of claims 22 to 24 wherein about 25, 50, 75, 100 or greater than 100 base pairs of DNA adjacent or near the Tn_A insertion site are sequenced.

26. The method of any one of claims 22 to 25 wherein the sequenced DNA is 5' and/or 3' to the Tn_A insertion site.

27. The method of any one of the preceding claims further comprising the step of assigning an essential function to said putative essential gene by sequence comparison with one or more essential gene(s) of said bacterium.

28. The method of claim 27 wherein said essential gene(s) are identified by transposon directed insertion site sequencing using a transposon which inactivates on insertion.

29. The method of any one of the preceding claims further comprising the step of assigning an essential function to said putative essential gene by determining that it is not an antibiotic resistance gene.

30. The method of claim 29 wherein the determining step comprises identifying an antibiotic resistance gene by transposon directed insertion site sequencing using a transposon which inactivates on insertion.

31. The method of any one of the preceding claims further comprising the steps of: generating an antibiotic resistant mutant of said bacterium by a method comprising the step of selecting for growth in the presence of said antibiotic to produce an antibiotic resistant mutant clone (Ab^R mutant); transforming the Ab^R mutant with: (i) one or more essential genes of said bacterium; and (ii) a transposon which insertionally inactivates bacterial DNA, to produce a pool of transposon mutants which are merodiploid for said one or more essential genes; growing bacteria from the merodiploid pool in the presence of different amounts of said antibiotic to produce two or more test cultures; and comparing the distribution of transposon insertions between test cultures to identify a putative essential gene serving as a target of said antibiotic in said bacterium.

32. A method of identifying an antibiotic comprising identifying an essential gene which serves as a target of said antibiotic according to a method as defined in any one of the preceding claims.

33. A process for producing an antibiotic comprising the method as defined in claim 32.

34. The process of claim 33 further comprising synthesising said antibiotic.

35. The process of claim 34 further comprising mixing the synthesised antibiotic with a pharmaceutically acceptable excipient to produce a pharmaceutical composition.