

HETEROARYL ACETYLENIC SULFONAMIDE AND PHOSPHINIC ACID AMIDE HYDROXAMIC ACID TACE INHIBITORS

FIELD OF INVENTION

5

This invention relates to acetylenic aryl and heteroaryl sulfonamide and phosphinic acid amide hydroxamic acids which act as inhibitors of TNF- α converting enzyme (TACE). The compounds of the present invention are useful in disease conditions mediated by TNF- α such as rheumatoid arthritis, osteoarthritis, sepsis, AIDS, ulcerative colitis, multiple sclerosis, Crohn's disease and degenerative cartilage loss.

BACKGROUND OF THE INVENTION


15

TNF- α converting enzyme (TACE) catalyzes the formation of TNF- α from membrane bound TNF- α precursor protein. TNF- α is a pro-inflammatory cytokine that is believed to have a role in rheumatoid arthritis [Shire, M. G.; Muller, G. W. *Exp. Opin. Ther. Patents* 1998, 8(5), 531; Grossman, J. M.; Brahn, E. J. *Women's Health* 1997, 6(6), 627; Isomaki, P.; Punnonen, J. *Ann. Med.* 1997, 29, 499; Camussi, G.; Lupia, E. *Drugs*, 1998, 55(5), 613.] septic shock [Mathison, et. al. *J. Clin. Invest.* 1988, 81, 1925; Miethke, et. al. *J. Exp. Med.* 1992, 175, 91.], graft rejection [Piguet, P. F.; Grau, G. E.; et. al. *J. Exp. Med.* 1987, 166, 1280.], cachexia [Beutler, B.; Cerami, A. *Ann. Rev. Biochem.* 1988, 57, 505.], anorexia, inflammation [Ksontini, R.; MacKay, S. L. D.; Moldawer, L. L. *Arch. Surg.* 1998, 133, 558.], congestive heart failure [Packer, M. *Circulation*, 1995, 92(6), 1379; Ferrari, R.; Bachetti, T.; et. al. *Circulation*, 1995, 92(6), 1479.], post-ischaemic reperfusion injury, inflammatory disease of the central nervous system, inflammatory bowel disease, insulin resistance [Hotamisligil, G. S.; Shargill, N. S.; Spiegelman, B. M.; et. al. *Science*, 1993, 259, 87.] and HIV infection [Peterson, P. K.; Gekker, G.; et. al. *J. Clin. Invest.* 1992, 89, 574; Pallares-Trujillo, J.; Lopez-Soriano, F. J. Argiles, J. M. *Med. Res. Reviews*, 1995, 15(6), 533.], in addition to its well-documented antitumor properties [Old, L. *Science*, 1985, 230, 630.]. For example, research with anti-TNF- α antibodies and transgenic animals has demonstrated that blocking the formation of TNF- α inhibits the progression of arthritis [Rankin, E.C.; Choy, E.H.; Kassimos, D.; Kingsley, G.H.; Sopwith, A.M.; Isenberg, D.A.; Panayi, G.S. *Br. J. Rheumatol.* 1995, 34, 334; *Pharmaprojects*, 1996, Therapeutic Updates 17 (Oct.), au197-M2Z.]. This

observation has recently been extended to humans as well as described in "TNF- α in Human Diseases", *Current Pharmaceutical Design*, 1996, 2, 662.


It is expected that small molecule inhibitors of TACE would have the potential for treating a variety of disease states. Although a variety of TACE inhibitors are known, many of these molecules are peptidic and peptide-like which suffer from bioavailability and pharmacokinetic problems. In addition, many of these molecules are non-selective, being potent inhibitors of matrix metalloproteinases and, in particular, MMP-1. Inhibition of MMP-1 (collagenase 1) has been postulated to cause joint pain in clinical trials of MMP inhibitors [*Scrip*, 1998, 2349, 20]. Long acting, selective, orally bioavailable non-peptide inhibitors of TACE would thus be highly desirable for the treatment of the disease states discussed above.

Examples of sulfonamide hydroxamic acid MMP/TACE inhibitors in which a 2 carbon chain separates the hydroxamic acid and the sulfonamide nitrogen, as shown below, are disclosed in WIPO international publications WO9816503, WO9816506, WO9816514 and WO9816520 and U. S. patent 5,776,961.




U. S. patents 5,455,258, 5,506,242, 5,552,419, 5,770,624, 5,804,593 and 5,817,822 as well as European patent application EP606,046A1 and WIPO international publications WO9600214 and WO9722587 disclose non-peptide inhibitors of matrix metalloproteinases and/or TACE of which the aryl sulfonamide hydroxamic acid shown below, in which 1 carbon separates the hydroxamic acid and the sulfonamide nitrogen, is representative. Additional publications disclosing sulfonamide based MMP inhibitors which are variants of the sulfonamide-hydroxamate shown below, or the analogous sulfonamide-carboxylates, are European patent applications EP-757037-A1 and EP-757984-A1 and WIPO international publications WO9535275, WO9535276, WO9627583, WO9719068, WO9727174, WO9745402, WO9807697, and WO9831664, WO9833768, WO9839313, WO9839329, WO9842659 and WO9843963. The discovery of this type of MMP inhibitor is further detailed by MacPherson, *et. al.* in *J. Med. Chem.*, (1997), 40, 2525 and Tamura, *et. al.* in *J. Med. Chem.* (1998), 41, 640.


-3-




Publications disclosing β -sulfonamide-hydroxamate inhibitors of MMPs and/or TACE in which the carbon alpha to the hydroxamic acid has been joined in a ring to the sulfonamide nitrogen, as shown below, include U. S. patent 5,753,653, WIPO international publications WO9633172, WO9720824, WO9827069, WO9808815, WO9808822, WO9808823, WO9808825, WO9834918, WO9808827, Levin, *et. al. Bioorg. & Med. Chem. Letters* 1998, 8, 2657 and Pikul, *et. al. J. Med. Chem.* 1998, 41, 3568.




The patent applications DE19,542,189-A1, WO9718194, and EP803505 disclose additional examples of cyclic sulfonamides as MMP and/or TACE inhibitors. In this case the sulfonamide-containing ring is fused to a aromatic or heteroaromatic ring.



Analogous to the sulfonamides are the phosphinic acid amide hydroxamic acid MMP/TACE inhibitors, exemplified by the structure below, which have been disclosed in WIPO international publication WO9808853.




Sulfonamide MMP/TACE inhibitors in which a thiol is the zinc chelating group, as shown below, have been disclosed in WIPO international application 9803166.



It is an object of this invention to disclose aryl and heteroaryl sulfonamide and phosphinic acid amide hydroxamic acid MMP/TACE inhibitors in which the sulfonyl aryl group is para-substituted with a substituted butynyl moiety or a propargylic ether, amine or sulfide. These compounds provide enhanced levels of inhibition of the activity of TACE in vitro and in a cellular assay and/or selectivity over MMP-1. These compounds may therefore be used in the treatment of diseases mediated by TNF.

SUMMARY OF THE INVENTION

The TACE and MMP inhibiting ortho-sulfonamido and phosphinic acid amide aryl and heteroaryl hydroxamic acids of the present invention are represented by the formula:



where the C(=O)NHOH moiety and the -NR₅- moiety are bonded to adjacent carbons of group A; wherein

A is 5-6 membered heteroaryl having 1 to 3 heteroatoms selected from N, NR₅, S and O;

X is SO₂ or -P(O)R₁₀;

Y is aryl or 5-10 membered mono- or bi-cyclic heteroaryl having from 1 to three heteroatoms selected from N, NR₅, S and O, with the proviso that

X and Z may not be bonded to adjacent atoms of Y;

Z is O, NH, CH₂ or S;

R₅ is hydrogen or alkyl of 1-6 carbon atoms;

R_6 and R_7 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, -CN, -CCH₃;

R_8 is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, aryl, 5 to 10 membered heteroaryl having 1 to 3 heteroatoms selected from N, NR₉, S and O, or 5 to 9 membered heterocycloalkyl having 1 or 2 heteroatoms selected from N, NR₉, S and O;

R_9 is hydrogen, aryl, alkyl of 1-6 carbon atoms or cycloalkyl of 3-6 carbon atoms;

and R_{10} is alkyl of 1-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, aryl, or heteroaryl; or a pharmaceutically acceptable salt thereof.

Preferred compounds of this invention are those of structure B wherein both carbons of A adjacent the -NR₅- group have a substituent other than hydrogen.

More preferred compounds of this invention include compounds of structure B in which A is a 5-6 membered heteroaryl having 1 to 3 heteratoms selected from N, NR₉, S and O wherein:

both carbons of A adjacent the -NR₅- group have a substituent other than hydrogen;

and Y is a phenyl ring substituted at the 1- and 4-positions by X and Z, respectively.

More preferred compounds of this invention include compounds of structure B in which A is a phenyl wherein:

both carbons of A adjacent the -NR₅- group have a substituent other than hydrogen;

Y is a phenyl ring substituted at the 1- and 4-positions by X and Z, respectively;

and X is SO₂.

More preferred compounds of this invention include compounds of structure B in which A is a phenyl wherein:

-6-

both carbons of A adjacent the $-NR_5-$ group have a substituent other than hydrogen;

Y is a phenyl ring substituted at the 1- and 4-positions by X and Z, respectively;

5

X is SO_2 ;

Z is oxygen;

and R_6 and R_7 are hydrogen.

More preferred compounds of this invention include compounds of structure

10 B in which A is a phenyl wherein:

both carbons of A adjacent the $-NR_5-$ group have a substituent other than hydrogen;

Y is a phenyl ring substituted at the 1- and 4-positions by X and Z, respectively;

15


X is SO_2 ;

Z is oxygen;

R_6 and R_7 are hydrogen;


and R_8 is $-CH_2OH$ or methyl.

20 Heteroaryl, as used throughout, is a 5-10 membered mono- or bicyclic ring having from 1-3 heteroatoms selected from N, NR_9 , S and O. Heteroaryl is preferably



wherein K is NR_9 , O or S and R_9 is hydrogen, phenyl, naphthyl, alkyl of 1-6 carbon atoms, or cycloalkyl of 3-6 carbon atoms. Preferred heteroaryl rings include pyrrole, furan, thiophene, pyridine, pyrimidine, pyridazine, pyrazine, triazole, pyrazole, imidazole, isothiazole, thiazole, isoxazole, oxazole, indole, isoindole, benzofuran, benzothiophene, quinoline, isoquinoline, quinoxaline, quinazoline, benzotriazole, indazole, benzimidazole, benzothiazole, benzisoxazole, and benzoxazole.

For purposes of the definition of A, It is still more preferred that A is a heteroaryl selected from




Heteroaryl groups of the present invention may optionally be mono- or di-substituted.

Heterocycloalkyl as used herein refers to a 5 to 10 membered saturated or unsaturated mono or bi-cyclic ring having 1 or 2 heteroatoms selected from N, NR_9 , S or O. Heterocycloalkyl rings of the present invention are preferably

-8-

selected from



wherein K is NR_9 , O or S and R_9 is hydrogen, phenyl, naphthyl, alkyl of 1-6 carbon atoms, or cycloalkyl of 3-6 carbon atoms. Preferred heterocycloalkyl rings include piperidine, piperazine, morpholine, tetrahydropyran, tetrahydrofuran or pyrrolidine. Heterocycloalkyl groups of the present invention may optionally be mono- or di-substituted.

Aryl, as used herein refers phenyl or naphthyl which may, optionally be mono-, di- or tri-substituted.

Alkyl, alkenyl, alkynyl, and perfluoroalkyl include both straight chain as well as branched moieties. Alkyl, alkenyl, alkynyl, and cycloalkyl groups may be unsubstituted (carbons bonded to hydrogen, or other carbons in the chain or ring) or may be mono- or poly-substituted.

Halogen means bromine, chlorine, fluorine, and iodine.

Suitable substituents of aryl, heteroaryl, alkyl, alkenyl, alkynyl, cycloalkyl and include, but are not limited to halogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, $-\text{OR}_2$, $-\text{CN}$, $-\text{COR}_2$, perfluoroalkyl of 1-4 carbon atoms, $-\text{O}$ -perfluoroalkyl of 1-4 carbon atoms, $-\text{CONR}_2\text{R}_3$, $-\text{S}(\text{O})_n\text{R}_2$, $-\text{OPO}(\text{OR}_2)\text{OR}_3$, $-\text{PO}(\text{OR}_2)\text{R}_3$, $-\text{OC}(\text{O})\text{NR}_2\text{R}_3$, $-\text{C}(\text{O})\text{NR}_2\text{OR}_3$, $-\text{COOR}_2$,

-9-

-SO₂H, -NR₂R₃, -N[(CH₂)₂]₂NR₂, -NR₂COR₃, -NR₂COOR₃, -SO₂NR₂R₃,
-NO₂, -N(R₂)SO₂R₃, -NR₂CONR₂R₃, -NR₂C(=NR₃)NR₂R₃,
-NR₂C(=NR₃)N(SO₂R₂)R₃, NR₂C(=NR₃)N(C=OR₂)R₃, -SO₂NHCOR₄,
-CONHSO₂R₄, -tetrazol-5-yl, -SO₂NHCN, -SO₂NHCONR₂R₃, phenyl,
5 naphthyl, heteroaryl or heterocycloalkyl;

wherein -NR₂R₃ may form a pyrrolidine, piperidine, morpholine,
thiomorpholine, oxazolidine, thiazolidine, pyrazolidine, piperazine, or
azetidine ring;

R₂ and R₃ are each, independently, hydrogen, alkyl of 1-6 carbon atoms,
10 cycloalkyl of 3-6 carbon atoms, phenyl, naphthyl, heteroaryl or
heterocycloalkyl;

R₄ is alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6
carbon atoms, cycloalkyl of 3-6 carbon atoms; perfluoroalkyl of 1-4 carbon atoms,
15 phenyl, naphthyl, heteroaryl or heterocycloalkyl; and n is 0 to 2.

Suitable substituents of heterocycloalkyl groups of the present invention
include, but are not limited to alkyl of 1-6 carbon atoms, cycloalkyl of 3-6 carbon
atoms, phenyl, naphthyl, heteroaryl and heterocycloalkyl.

20

When a moiety contains more than one substituent with the same designation
each of those substituents may be the same or different.


Pharmaceutically acceptable salts can be formed from organic and inorganic
25 acids, for example, acetic, propionic, lactic, citric, tartaric, succinic, fumaric, maleic,
malonic, mandelic, malic, phthalic, hydrochloric, hydrobromic, phosphoric, nitric,
sulfuric, methanesulfonic, naphthalenesulfonic, benzenesulfonic, toluenesulfonic,
camphorsulfonic, and similarly known acceptable acids when a compound of this
invention contains a basic moiety. Salts may also be formed from organic and
30 inorganic bases, preferably alkali metal salts, for example, sodium, lithium, or
potassium, when a compound of this invention contains an acidic moiety.

The compounds of this invention may contain an asymmetric carbon atom and some of the compounds of this invention may contain one or more asymmetric centers and may thus give rise to optical isomers and diastereomers. While shown without respect to stereochemistry, the present invention includes such optical isomers and diastereomers; as well as the racemic and resolved, enantiomerically pure R and S stereoisomers; as well as other mixtures of the R and S stereoisomers and pharmaceutically acceptable salts thereof. It is recognized that one optical isomer, including diastereomer and enantiomer, or stereoisomer may have favorable properties over the other. Thus when disclosing and claiming the invention, when one racemic mixture is disclosed, it is clearly contemplated that both optical isomers, including diastereomers and enantiomers, or stereoisomers substantially free of the other are disclosed and claimed as well.

The compounds of this invention are shown to inhibit the enzymes MMP-1, MMP-9, MMP-13 and TNF- α converting enzyme (TACE) and are therefore useful in the treatment of arthritis, tumor metastasis, tissue ulceration, abnormal wound healing, periodontal disease, graft rejection, insulin resistance, bone disease and HIV infection. In particular, the compounds of the invention provide enhanced levels of inhibition of the activity of TACE in vitro and in cellular assay and/or enhanced selectivity over MMP-1 and are thus particularly useful in the treatment of diseases mediated by TNF.


This invention provides a process for preparing compounds of formula B as defined above which comprises one of the following:

- a) reacting a compound of formula V:



wherein R_5 , R_6 , R_7 , R_8 , A, X, Y and Z are as defined above and Q is COOH or a reactive derivative thereof, with hydroxylamine to give a corresponding compound of formula B; or

b) deprotecting a compound of formula VI:



(VI)

wherein R_5 , R_6 , R_7 , R_8 , A, X, Y and Z are as defined above, and R_{30} is a protecting group such as t-butyl, benzyl or trialkylsilyl, to give a compound of formula B;

c) resolving a mixture (e.g. racemate) of optically active isomers of a compound of formula B to isolate one enantiomer or diastereomer substantially free of the other enantiomer or diastereomers;

or

d) acidifying a basic compound of formula B with a pharmaceutically acceptable acid to give a pharmaceutically acceptable salt.

With regard to process a) the reaction can be carried out by processes known in the art, e.g. by reaction with a halogenating agent to form a reactive derivative (ie acid chloride) followed by reaction with the hydroxylamine.

Removal of protecting groups, as illustrated by process b) can be carried out by processes known in the art to provide the hydroxamic acid.

With regard to process c) standard separation techniques may be used to isolate particular enantiomeric or diastereomeric forms. For example a racemic mixture may be converted to a mixture of optically active diastereoisomers by reaction with a single enantiomer of a 'resolving agent' (for example by diastereomeric salt formation or formation of a covalent bond). The resulting mixture of optically active diastereoisomers may be separated by standard techniques


-12-

(e.g. crystallisation or chromatography) and individual optically active diastereoisomers then treated to remove the 'resolving agent' thereby releasing the single enantiomer of the compound of the invention. Chiral chromatography (using a chiral support, eluent or ion pairing agent) may also be used to separate enantiomeric mixtures directly.

The compounds of formula B may be isolated in the form of a salt of a pharmaceutically acceptable acid, e.g. an organic or inorganic acid by treatment with an acid such as described above.


5 The invention is further directed to a process for making compounds of structure B involving one or more reactions as follows:

1) alkylating a compound of formula I, or a salt or solvate thereof,




10

into a compound of formula II



15 2) reacting a compound of formula II above, or a salt or solvate thereof, with a chlorinating agent such as thionyl chloride, chlorosulfonic acid, oxalyl chloride, phosphorus pentachloride, or other halogenating agents such as fluorosulfonic acid or thionyl bromide to a compound of formula III:

-13-




wherein J is fluorine, bromine, chlorine.

The resultant sulfonyl chloride, fluoride or bromide, may be further converted
 5 into triazolidine, imidazolidine or benzothiazolidine derivatives, where J is 1,2,4-triazolyl,
 benzotriazolyl or imidazolyl, by reacting the compound with 1,2,4-triazole,
 imidazole or benzotriazole, respectively. R₆, R₇ and R₈ are as defined above.

The invention is still further directed to a process for making compounds of
 10 structure B involving one or more reactions as follows:

1) alkylating phenol, or a salt or solvate thereof, into a compound of formula IV:








15 2) reacting a compound of formula IV above, or a salt or solvate thereof with
 chlorosulfonic acid to prepare a compound of formula II above.

Particularly preferred intermediates are compounds of formulae II and III,
 with the proviso that R₆ is not hydrogen.

20 The invention compounds are prepared using conventional techniques known
 to those skilled in the art of organic synthesis. The starting materials used in
 preparing the compounds of the invention are known, made by known methods or are
 commercially available. The following compounds (V-IX) which may be used in
 preparing compounds of the invention are known and references are given herein
 25 below. This list is included for illustrative purposes only and is not to be construed as
 limiting in any way.

-14-

**V****VI****VII****VIII****IX**

Literature references for these materials are as follows:

Compound V:

- 5 a) Dolle, RE; Hoyer, DW; Schmidt, SJ; Ross, TM; Rinker, JM; Ator, MA
Eur. Pat. Appl. EP-628550.
- b) Wermuth, C-G; Schlewer, G; Bourguignon, J-J; Maghioros, G; Bouchet, M-
J et. al. **J. Med. Chem** (1989), 32, 528-537.
- c) Yutugi, S et. al. **Chem. Pharm. Bull.** (1971) 19, 2354-2364.
- 10 d) Dolle, RE; Hoyer, D; Rinker, JM; Ross, TM; Schmidt, SJ **Biorg. Med.**
Chem. Lett (1977) 7, 1003-1006.

Compound VI:

Camparini, A; Ponticelli, F; Tedeschi, P. **J. Chem. Soc., Perkin Trans.1** (1982),
10, 2391-4.

15 Compound VII:

Muller, C.E.; Geis, U.; Grahner, B.; Lanzner, W.; Eger, K. **J. Med. Chem.** (1996),
39, 2482.

Compound VIII:

Muller, C.E.; Geis, U.; Grahner, B.; Lanzner, W.; Eger, K. **J. Med. Chem.** (1996),
20 39, 2482.

Compound IX:

Commercially available.


Those skilled in the art will recognize that certain reactions are best carried
5 out when other potentially reactive functionality on the molecule is masked or
protected, thus avoiding undesirable side reactions and/or increasing the yield of the
reaction. To this end, those skilled in the art may use protecting groups. Examples of
these protecting group moieties may be found in T. W. Greene, P. G. M. Wuts
10 "Protective Groups in Organic Synthesis", 2nd Edition, 1991, Wiley & Sons, New
York. Reactive side chain functionalities on amino acid starting materials are
preferably protected. The need and choice of protecting groups for a particular
reaction is known to those skilled in the art and depends on the nature of the
functional group to be protected (hydroxy, amino, carboxy, etc.), the structure and
stability of the molecule of which the substituent is part and the reaction conditions.

15 When preparing or elaborating compounds of the invention containing aryl,
heteroaryl or heterocyclic rings, those skilled in the art recognize that substituents on
that ring may be prepared before, after or concomitant with construction of the ring.
For clarity, substituents on such rings have been omitted from the schemes herein
below.

20 Those skilled in the art will recognize that the nature and order of the
synthetic steps presented may be varied for the purpose of optimizing the formation
of the compounds of the invention.

The hydroxamic acid compounds of the invention, 1, are prepared according
to Scheme 1 by converting a carboxylic acid, 2, into the corresponding acid chloride
25 or anhydride, or by reacting it with a suitable peptide coupling reagent, followed by
reaction with hydroxylamine to give 1, or with a protected hydroxylamine derivative
to give 3. Compounds 3, wherein R₃₀ is a t-butyl, benzyl, trialkylsilyl or other suitable
masking group may then be deprotected by known methods to provide the
hydroxamic acid 1.


Scheme 1:



Carboxylic acids 2 may be prepared as shown in Scheme 2. Amino acid derivative 4, in which R₄₀ is hydrogen or a suitable carboxylic acid protecting group, may be sulfonylated or phosphorylated by reacting with compounds 5, in which J is a suitable leaving group including, but not limited to chlorine. The N-H compound 6 may then be alkylated with R_jJ and a base such as potassium carbonate or sodium hydride in a polar aprotic solvent such as acetone, N,N-dimethylformamide (DMF), or tetrahydrofuran (THF) to provide sulfonamide 7. Compound 7 is also available through direct reaction of 5 with an N-substituted amino acid derivative, 8. Conversion of 7 into the carboxylic acid is performed by acid, base hydrolysis, or other method consistent with the choice of protecting group R₄₀ and the presence of a carbon-carbon triple bond.

-17-

Scheme 2:




Methods of preparation of sulfonylating agents 5 are shown in Scheme 3. Thus, sulfonic acid salts 9, where ZR₅₀ is a hydroxy, thiol or substituted amino moiety may be alkylated with acetylenes 10, where J is a suitable leaving group such as halogen mesylate, tosylate, or triflate to give 11. Acetylenes 10 are commercially available or known compounds, or they may be synthesized by known methods by those skilled in the art. The sulfonic acid salts 11 may be converted into the corresponding sulfonyl chloride or other sulfonylating agent 5 by known methods, such as reaction with oxalyl chloride or other reagent compatible with substituents R₆, R₇, and R₈ and the acetylene. Alternatively, the disulfide 12 may be converted into diacetylene 13 by reaction with compounds 10, followed by reduction of the disulfide bond to provide the analogous thiols which may be converted into 5 by known methods. Alkylation of the phenol, thiophenol, aniline or protected aniline 14 with 10 to give 15, followed by reaction with chlorosulfonic acid provide sulfonic acids 16 which are readily converted into 5 with oxalyl chloride or similar reagents. Thiophenols 17 are also precursors to 5 via protection of the thiol, alkylation of ZH,

-18-

where Z is O, N or S, and deprotection of the sulfur followed by oxidation to the sulfonic acid 16.


Scheme 3:



5

The phosphorus containing analogs of 8 may be prepared using similar methodology, as shown in Scheme 4.


Scheme 4:



The acetylenic side chain may also be appended after sulfonylation or phosphorylation of the amino acid derivative, as shown in **Scheme 5**. Thus, the amino acid derivatives 4 and 8 can be sulfonylated or phosphorylated with compounds 20, where ZR_{50} is hydroxy or protected hydroxy, thiol or amine, and, if necessary, alkylated with R_7J as in **Scheme 2**, to give 21. Removal of the R_{50} masking group to give 22 and subsequent alkylation of the resulting phenol, thiol or amine with 10 provides 7. In the case where ZR_{50} is equal to OH, no deprotection step is required to give 22.

-20-


Scheme 5:



The propargylic amine analogs of 7 can be synthesized as shown in **Scheme 6** starting from the amino acid derivatives 4 and/or 8. Sulfenylation or phosphorylation with para-nitro aryl compound 23, for example 4-nitrobenzenesulfonyl chloride, followed by alkylation with R₅J (for 4) using a base such as potassium carbonate or sodium hydride in DMF provides 24. Reduction of the nitro moiety with hydrogen and palladium on carbon, tin chloride or other known method to give aniline 25 and subsequent alkylation with 10 then provides 7. Aniline 25 may be derivatized with a suitable nitrogen protecting group, such as t-butoxycarbonyl, to give 26 prior to alkylation with 10 subsequent deprotection after the alkylation step.

-21-

Scheme 6:




Acetylenic derivatives 7 are also accessible via the fluoro compounds 27, readily prepared from the amino acid derivatives 4 and/or 8 by reaction with fluoraryl

5 26, as shown in Scheme 7. Displacement of the fluorine of 27 in the presence of a base such as sodium hydride with a masked hydroxy, thiol, or amino group (HZR_{70} , where R_{70} is a suitable protecting group) in a polar aprotic solvent such as DMF, followed by deprotection gives 28, which can then be alkylated with 10 to provide 7. Conversion of 27 to 28, where Z is sulfur, might also be accomplished with Na_2S ,

10 K_2S , $NaSH$ or $KS(C=S)OEt$. The fluorine of 27 can also be displaced in a polar aprotic solvent with the propargylic derivative 29, where Z is O, S or NH, in the presence of a base such as sodium hydride, to give 7 directly.

-22-


Scheme 7:



Compound 7, wherein Z is a methylene group, is available via 30, as shown in Scheme 8. Benzylic bromination of 30 with N-bromosuccinimide in a chlorinated hydrocarbon solvent provides bromide 31. This is followed by displacement of the bromide with the appropriate propynyl cuprate to provide sulfonamide 8.

-23-

Scheme 8:




Compounds of the invention can also be prepared by modifying substituents on the acetylenic side chain at any stage after sulfonylation or phosphorylation of the starting amino acid derivatives 4 or 8. Functional groups such as halogen, hydroxy, amino, aldehyde, ester, ketone, etc. may be manipulated by standard methods to form the moieties defined by R₁-R₈ of compounds 1. It is recognized by those skilled in the art of organic synthesis that the successful use of these methods is dependent upon the compatibility of substituents on other parts of the molecule. Protecting groups and/or changes in the order of steps described herein may be required.

Some of the methods available for the derivatization of compounds of structure 32 (equivalent to compound 7 wherein R₁₂ is hydrogen) are shown in Scheme 9. Metallation of the terminal acetylene 32 followed by addition of an aldehyde or alkyl halide, sulfonate or triflate provides derivatives 33 and 34. Reaction of 32 with formaldehyde and an amine provides the Mannich addition product 35. Cyanogen bromide addition to 35 gives the propargylic bromide 36 which may be displaced with a variety of nucleophiles to give, for example, ethers, thioethers and amines 37. Palladium catalyzed coupling reactions of 32 provide the aryl or heteroaryl acetylenes 38. It is recognized by those skilled in the art of organic

synthesis that the successful use of these methods is dependent upon the compatibility of substituents on other parts of the molecule. Protecting groups and/or changes in the order of steps described herein may be required and R_{35} , R_{45} , R_{55} , R_{65} and R_{75} are alkyl, e.g. methyl.


5

Scheme 9:



Shown in Scheme 10 is the synthesis of an example of the invention wherein A is pyridyl. In this specific example, shown for illustrative purposes only, the BOC-protected amino-pyridine **39** is synthesized from 3-amino-2,6-dimethoxypyridine via reaction with BOC anhydride. The ortho-amino ester, **40**, is then constructed via metallation and subsequent carboxylation of **39**. Removal of the BOC protecting group from ester **41** provides ortho amino-ester **42**. Elaboration of **42** according to Schemes 1-9 then provides the compounds of the invention. Additional pyridyl-hydroxamates are available through the same route.

Scheme 10:



The following specific examples illustrate the preparation of representative compounds of this invention. The starting materials, intermediates, and reagents are either commercially available or can be readily prepared following standard literature procedures by one skilled in the art of organic synthesis.

Example 1

3-(4-Methoxy-benzenesulfonylamino)-thiophene-2-carboxylic acid methyl ester

To a solution of 5.00g (0.032 mol) of 3-amino-2-carbomethoxythiophene dissolved in 40 mL of chloroform was added 7.73 mL (0.032 mol) of pyridine followed by 6.57g (0.032 mol) of p-methoxybenzenesulfonyl chloride. The reaction mixture was stirred at room temperature for 5h and then washed with 3N HCl and water. The organics were then dried over Na₂SO₄, filtered and concentrated in vacuo. The resulting cream colored solid was washed with ether and dried in vacuo to

provide 6.89g (66%) of the desired sulfonamide. Electrospray Mass Spec 328.2 (M+H).

Example 2

5 **4-(4-Methoxy-benzenesulfonylamino)-thiophene-3-carboxylic acid methyl ester**

In the same manner as described in Example 1, 5.00g (0.026 mol) of 3-amino-4-carbomethoxythiophene hydrochloride provided 3.50g (41%) of the desired sulfonamide as a brown solid after trituration with ether. Electrospray Mass Spec 328.2 (M+H).

10

Example 3

5-(4-Methoxy-benzenesulfonylamino)-1-methyl-1H-pyrazole-4-carboxylic acid ethyl ester

In the same manner as described in Example 1, 2.00g (0.012 mol) of 1-methyl-2-amino-3-carboethoxy-pyrazole provided 0.923g (23%) of the desired sulfonamide as a white solid after recrystallization from ethyl acetate/Hexanes. Electrospray Mass Spec 340.2 (M+H).

15

Example 4

20 **3-(4-Methoxy-benzenesulfonylamino)-4-methyl-thiophene-2-carboxylic acid methyl ester**

In the same manner as described in Example 1, 4.14g (0.024 mol) of 3-amino-4-methyl-2-carbomethoxy thiophene provided 4.89g (47%) of the desired sulfonamide as a white solid after trituration with ether. EI Mass Spec 340.9 (M⁺).

25

Example 5

3-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-thiophene-2-carboxylic acid methyl ester

To a solution of 2.0g (6.116 mmol) of the product of Example 1 in 25 mL of DMF was added 0.257g (6.422 mmol) of 60% sodium hydride. The resulting mixture was stirred for 30 min at room temperature and then 0.76 mL (6.422 mmol) of benzyl bromide was added. This reaction mixture was stirred overnight at room temperature, poured into water and then extracted with ether. The combined organics were washed with water and brine, dried over MgSO₄, filtered and concentrated in vacuo. The residue was chromatographed on silica gel eluting with ethyl acetate/hexanes (1:3) to

30

35

provide 1.62g (65%) of the desired product as white crystals. CI Mass Spec: 418 (M+H).

Example 6

5 **4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-thiophene-3-carboxylic acid methyl ester**

In the same manner as described in Example 5, 1.50g (4.587 mmol) of the product of Example 2 provided 1.257g (66%) of the desired product as a brown oil after chromatography on silica gel eluting with ethyl acetate/hexanes (1:10). CI Mass
10 Spec: 418 (M+H).

Example 7

15 **5-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-1-methyl-1H-pyrazole-4-carboxylic acid ethyl ester**

In the same manner as described in Example 5, 0.843g (2.484 mmol) of the product of Example 3 provided 0.924g (87%) of the desired product as a white solid after trituration with ether. CI Mass Spec: 430 (M+H).

Example 8

20 **3-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-4-methyl-thiophene-2-carboxylic acid methyl ester**

In the same manner as described in Example 5, 2.00g (4.64 mmol) of the product of Example 4 provided 1.648g (68%) of the desired product as a white solid after trituration with ether. CI Mass Spec: 432 (M+H).

25

Example 9

30 **3-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-thiophene-2-carboxylic acid**

To a mixture of 1.494g (3.583 mmol) of the product of Example 5 dissolved
30 in 15 mL of methanol and 15 mL of THF was added 15 mL of 1N NaOH solution. The reaction mixture was stirred at room temperature for 36h and the organics were removed in vacuo. The resulting mixture was acidified with 10% HCl and extracted with ethyl acetate. The combined organics were washed with water and brine, dried over MgSO₄, filtered and concentrated in vacuo. The resulting residue was triturated
35 with ether and filtered to provide 1.327g (92%) of the desired carboxylic acid as a white solid. CI Mass Spec: 404 (M+H).

Example 10**4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-thiophene-3-carboxylic acid**

5 In the same manner as described in Example 9, 1.157g (2.775 mmol) of the product of Example 6 provided 0.94g (84%) of the desired carboxylic acid as a tan solid after trituration with ether. Electrospray Mass Spec: 404 (M+H).

Example 11

10 **5-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-1-methyl-1H-pyrazole-4-carboxylic acid**

To a solution of 0.799g (1.862 mmol) of the product of Example 7 in 20 mL of methanol/THF (1:1) was added 9.3 mL of 1N sodium hydroxide solution and the resulting mixture was heated to reflux for 18h. The reaction was then cooled to room
15 temperature and the organics were removed in vacuo. The resulting mixture was acidified with 10% HCl and extracted with ethyl acetate. The combined organics were washed with water and brine, dried over MgSO₄, filtered and concentrated in vacuo. The resulting residue was triturated with ether and filtered to provide 0.697g
(93%) of the desired carboxylic acid as a white solid. Electrospray Mass Spec: 402
20 (M+H).

Example 12**3-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-4-methyl-thiophene-2-carboxylic acid**

25 In the same manner as described in Example 11, 1.366g (2.622 mmol) of the product of Example 8 provided 1.16g (87%) of the desired carboxylic acid as a white solid after trituration with ether. Electrospray Mass Spec: 416 (M-H)-.

Example 13

30 **5-Bromo-4-(4-methoxy-benzenesulfonylamino)-thiophene-3-carboxylic acid methyl ester**

: To a solution of the product of Example 2 in 5.0 mL of acetic acid-chloroform (1:1) at room temperature was added 0.299g (1.682 mmol) of N-bromosuccinimide. The reaction was stirred for 18h and then diluted with ether, washed with water and
35 saturated sodium bicarbonate solution, dried over MgSO₄, filtered and concentrated in vacuo. The tan solid residue was washed with ether-hexanes (1:1) to provide

0.504g (81%) of the desired product as a tan solid. Electrospray Mass Spec: 406.1 (M+H)+

Example 14

5 **4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-5-bromo-thiophene-
3-carboxylic acid methyl ester**

In the same manner as described in Example 5, 0.424g (1.044 mmol) of the product of Example 13 gave 0.400g (77%) of the desired N-benzyl methyl ester as a white solid. Electrospray Mass Spec: 496.1 (M+H)+

10

Example 15

15 **4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-5-bromo-thiophene-
3-carboxylic acid**

In the same manner as described in Example 11, 0.356g (0.718 mmol) of the product of Example 14 gave 0.290g (84%) of the desired carboxylic acid as a white solid. Electrospray Mass Spec: 482.1 (M+H)+

Example 16

20 **4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-5-ethynyl-thiophene-
3-carboxylic acid methyl ester**

To a solution of 0.294g (0.634 mmol) of the product of Example 14 in 2.5 mL of DMF and 2.5 mL of triethylamine was added 0.448 mL (3.168 mmol) of trimethylsilylacetylene, 0.022g (0.032 mmol) of bis(triphenylphosphine)-palladium(II)-dichloride and 3mg of copper(I)iodide. The reaction mixture was then heated to 80 °C for 6h and then cooled to room temperature and diluted with ether. The organics were washed with 5% HCl solution, water and brine, dried over MgSO₄, filtered and concentrated in vacuo. The residue was dissolved in 5 mL of THF, 1 mL of 1M tetrabutylammonium fluoride-THF solution was added and the reaction was stirred at room temperature for 1h, then diluted with ether, washed with 5% HCl solution, water and brine, dried over MgSO₄, filtered and concentrated in vacuo. The residue was chromatographed on silica eluting with ethyl acetate-hexanes (1:5) to provide 0.159g (61%) of the desired product as a brown oil.

30

Electrospray Mass Spec: 442.2 (M+H)⁺

Example 17**4-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-5-ethynyl-thiophene-3-carboxylic acid**

In the same manner as described in Example 11, 0.136g (0.333 mmol) of the product of Example 16 provided 0.075g (57%) of the desired product as a tan solid after chromatography on silica eluting with ethyl acetate-hexanes (1:1). Electrospray Mass Spec: 428.1 (M+H)+

Example 18**5-Bromo-4-[(4-methoxybenzenesulfonyl)-pyridin-3-ylmethylamino]thiophene-3-carboxylic acid methyl ester**

To a solution of 4.80 g (11.82 mmol) of the product of Example 13 dissolved in 5.0 mL of DMF was added 2.04 g (12.41 mmol) of 3-picoyl chloride hydrochloride and 4.89 g (35.46 mmol) of potassium carbonate. The reaction mixture was then stirred at room temperature for 18h, diluted with water and extracted with ether. The organics were then extracted with 6N HCl solution and the aqueous acid layer was then basified with 6N NaOH solution and then extracted with ether. The resulting ether layer was dried over sodium sulfate, filtered and concentrated in vacuo to provide 4.16g (71%) of the desired product as a tan solid. Electrospray Mass Spec: 498 (M+H).

Example 19**5-Bromo-4-[(4-methoxy-benzenesulfonyl)-pyridin-3-ylmethyl-amino]-thiophene-3-carboxylic acid**

To a solution of 0.40g (0.860 mmol) of the product of Example 18 in 9.0 mL of THF-MeOH (1:1) was added 0.072g (1.72 mmol) of lithium hydroxide monohydrate. The reaction mix was heated to reflux for 18h and then concentrated in vacuo. The residue was washed with THF and filtered. The filtrate was concentrated in vacuo to provide 0.388g (100%) of the desired product as a white foam. Electrospray Mass Spec: 483 (M+H).

Example 20**tert-Butyl N-(2,6-dimethoxy-3-pyridyl)carbamate**

To a suspension of 3-amino-2,6-dimethoxypyridine (1.5 g, 7.87 mmol) was added di-tert-butyl dicarbonate (3.43 g, 15.7 mmol). The solution was heated at reflux for 36 hours, cooled to room temperature, and diluted with water. The aqueous

solution was extracted 3 times with ethyl acetate, the organic extracts were combined, washed with brine, dried over MgSO_4 , concentrated in vacuo. The residue was purified by column chromatography using hexane/ethyl acetate as eluant (gradient 100% to 4/1) to provide 2.00g (100%) of tert-butyl N-(2,6-dimethoxy-3-pyridyl)carbamate a yellow oil. Electrospray Mass Spec: 254.9 (M+H)+

Example 21

tert-Butyl N-(4-carbomethoxy-2,6-dimethoxy-3-pyridyl)carbamate

The product of Example 20 (1 g, 3.93 mmol) was dissolved in ether (35 mL) and TMEDA (1.7 mL, 1.18 mmol) and cooled to -78°C . n-Butyllithium (4.75 mL, 11.87 mmol) was added dropwise and the reaction was allowed to stir for 15 minutes at -78°C before warming to -10°C for 2.5 hours. The solution was cooled back to -78°C and methyl chloroformate (0.6 mL, 7.8 mmol) dissolved in ether (4.5 mL) was added dropwise. The reaction was held at -78°C for 10 minutes and then warmed to -10°C and allowed to stir for 1.5 hours before quenching with ammonium chloride (sat). The reaction mixture was extracted 3X with ethyl acetate. The organics were combined, washed with brine, dried over MgSO_4 , concentrated in vacuo. The residue was purified by column chromatography using hexane / ethyl acetate as eluant (gradient 9/1 to 4/1) to provide 0.423 g (34 %) of tert-butyl N-(4-carbomethoxy-2,6-dimethoxy-3-pyridyl)carbamate as a white solid. Electrospray Mass Spec: 312.8 (M+H)+

Example 22

Methyl 3-amino-2,6-dimethoxyisonicotinate

p-Toluene sulfonic acid hydrate (0.282 g, 1.48 mmol) was dissolved in toluene (11 mL) and heated to reflux overnight with azeotropic removal of water (Dean-Stark trap). The next day, the reaction was cooled to room temperature and the product of Example 21, dissolved in toluene (4 mL), was added. The reaction was heated back to reflux for 0.5 hours. The reaction was cooled to room temperature and poured into sodium bicarbonate (sat) and extracted 3 times with ether. The organics were combined, washed with brine, dried over MgSO_4 , concentrated in vacuo. The residue was purified by column chromatography using hexane / ethyl acetate as eluant (gradient 100% to 9/1) to provide 0.278 g (97%) of methyl 3-amino-2,6-dimethoxyisonicotinate as a yellow solid. Electrospray Mass Spec: 212.8 (M+H)+

Example 23**Methyl 3-(4-methoxy-benzenesulfonylamino)-2,6-dimethoxy-isonicotinate**

To a solution of the product of Example 22 (0.278 g, 1.31 mmol) in pyridine (2 mL) was added p-methoxybenzenesulfonyl chloride (0.28 g, 1.38 mmol). The reaction mixture was stirred at room temperature overnight and was then quenched with water. The mixture was extracted 3 times with ether. The organics were combined, washed with brine, dried over MgSO_4 , concentrated in vacuo to provide 0.444 g (89%) of methyl 3-(4-methoxy-benzenesulfonylamino)-2,6-dimethoxy-isonicotinate as a solid. Electrospray Mass Spec: 382.8 (M+H)+

Example 24**Methyl 3-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-2,6-dimethoxy-isonicotinate**

The product of Example 23 (0.444g, 1.16 mmol) was dissolved in DMF (4 mL) and cooled to 0 °C. Benzyl bromide (0.186 mL, 1.6 mmol) followed by sodium hydride (56 mg, 1.39 mmol, 60% dispersion in mineral oil) were added and the reaction was allowed to warm to room temperature. After 1 h, the reaction was diluted with water and extracted 4X ether. The organics were combined, washed with brine, dried over MgSO_4 , concentrated in vacuo to provide 0.545 g (100%) of pure methyl 3-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-2,6-dimethoxy-isonicotinate as an oil. Electrospray Mass Spec: 472.9 (M+H)+

Example 25**3-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-2,6-dimethoxy-isonicotinic acid**

The product of Example 24 was hydrolyzed to the corresponding carboxylic acid using the procedure of Example 19 to provide 3-[Benzyl-(4-methoxy-benzenesulfonyl)-amino]-2,6-dimethoxy-isonicotinic acid. Electrospray Mass Spec: 459.0 (M+H)+

Example 26**4-But-2-ynyloxy-benzenesulfonic acid sodium salt**

To a solution of 52.35g (0.225 mol) of 4-hydroxybenzenesulfonate sodium salt in 1L of isopropanol and 225 mL of a 1.0N solution of sodium hydroxide was added 59.96g (0.45 mol) of 1-bromo-2-butyne. The resulting mixture was heated to 70° for 15h and then the isopropanol was removed by evaporation in vacuo. The

resulting white precipitate was collected by filtration, washed with isopropanol and ether and dried in vacuo to give 56.0g (100%) of the butynyl ether as a white solid.

Example 27

5

4-But-2-ynyloxy-benzenesulfonyl chloride

To a 0° solution of 43.8 mL (0.087 mol) of 2M oxalyl chloride/dichloromethane solution in 29 mL of dichloromethane was dropwise added 6.77 mL (0.087 mol) of DMF followed by 7.24g (0.029 mol) of the product of Example 26. The reaction mixture was stirred for 10 minutes at 0° then let warm to room temperature and stirred for 2 days. The reaction was then poured into ice and extracted with 150 mL of hexanes. The organics were washed with water and brine, dried over Na₂SO₄, filtered and concentrated in vacuo to provide 6.23g (88%) of the sulfonyl chloride as a yellow solid; m.p. 63-65°C. EI Mass Spec: 243.9 (M⁺).

15

Example 28

But-2-ynyloxy-benzene

To a solution of 6.14g (0.023 mol) of triphenylphosphine dissolved in 100 mL of benzene and 40 mL of THF was added 1.75 mL (0.023 mol) of 2-butyne-1-ol. After five minutes 2.00 (0.023 mol) phenol, dissolved in 10 mL of THF, was added to the reaction followed by 3.69 mL (0.023 mol) of diethyl azodicarboxylate. The resulting reaction mixture was stirred for 18h at room temperature and then concentrated in vacuo. The residue was chromatographed on silica gel eluting with ethyl acetate/hexanes (1:10) to provide 2.18g (70%) of the butynyl ether as a clear liquid. EI Mass Spec: 146.0 MH⁺

25

Example 29

4-But-2-ynyloxy-benzenesulfonyl chloride

To a solution of 0.146g (1.0 mmol) of the product of Example 28 in 0.3 mL of dichloromethane in an acetone/ice bath under N₂ was dropwise added a solution of 0.073 mL (1.1 mmol) of chlorosulfonic acid in 0.3 mL of dichloromethane. After the addition was complete, the ice bath was removed and the reaction was stirred at room temperature for 2h. To the reaction was then dropwise added 0.113 mL (1.3 mmol)

30

-34-

of oxalyl chloride, followed by 0.015 mL DMF. The reaction was heated to reflux for 2h and then diluted with hexane and poured into ice water. The organic layer was washed with brine, dried over sodium sulfate, and concentrated in vacuo to provide 0.130mg (53%) of the desired product as a light brown solid.

5

Example 30

Methyl 3-(4-but-2-ynyloxy-benzenesulfonylamino)-2,6-dimethoxy-isonicotinate

To a solution of the product of Example 22 (0.7g, 3.3 mmol) in pyridine (6 mL) was added 4-but-2-ynyloxy-benzenesulfonyl chloride (0.8g, 3.3 mmol). The reaction mixture was stirred at room temperature overnight and was then quenched with water. The mixture was extracted 3 times with ether. The organics were combined, washed with brine, dried over MgSO₄, and concentrated in vacuo. The residue was chromatographed on silica gel eluting with ethyl acetate/hexanes (gradient 1:1 to 7:3) to provide 1.15g of the butynyloxybenzene sulfonamide as a solid. Electrospray Mass Spec: 421.1 (M+H)⁺

15

Example 31

Methyl 3-[Methyl-(4-but-2-ynyloxy-benzenesulfonyl)-amino]-2,6-dimethoxy-isonicotinate

The product of Example 30 (0.48g, 1.13 mmol) was dissolved in DMF (5 mL) and cooled to 0 °C. Iodomethane (0.1 mL, 1.58 mmol) was added, followed by sodium hydride (0.054g, 1.35 mmol, 60% dispersion in mineral oil) and the reaction was allowed to warm to room temperature. After 1 h, the reaction was diluted with water and extracted 4 times with ethyl acetate. The organics were combined, washed with brine, dried over MgSO₄, concentrated in vacuo to provide 0.23g (48%) of the N-methyl sulfonamide as a white solid. Electrospray Mass Spec: 435.2 (M+H)⁺

20
25

Example 32

3-[Methyl-(4-but-2-ynyloxy-benzenesulfonyl)-amino]-2,6-dimethoxy-isonicotinic acid

30

The product of Example 31 (0.214g, 0.49 mmol) was hydrolyzed to the corresponding carboxylic acid using the procedure of Example 19 to provide 0.198g (100%) of 3-[methyl-(4-but-2-ynyloxy-benzenesulfonyl)-amino]-2,6-dimethoxy-isonicotinic acid. Electrospray Mass Spec: 421.1 (M+H)⁺

35

Example 33**3-[Methyl-(4-but-2-ynyloxy-benzenesulfonyl)-amino]-N-hydroxy-2,6-dimethoxy-isonicotinamide**

To a solution of (0.15g, 0.35 mmol) of the product from Example 32 in 2 mL of DMF was added 0.39 mL (0.77 mmol) of a 2M solution of oxalyl chloride in dichloromethane and the resulting reaction mixture was stirred at room temperature for 2h.

In a separate flask, 0.77 mL (5.6 mmol) of triethylamine was added to a 0 °C mixture of 0.24g (3.5 mmol) of hydroxylamine hydrochloride in 4 mL of THF and 1 mL of water. After this mixture had been stirred for 15 min at 0 °C, the acid chloride solution was added to it in one portion and the resulting solution was allowed to warm to room temperature and stirred for another 3h. The dichloromethane was washed with brine, dried over MgSO₄, filtered and concentrated in vacuo. Trituration of the residue with ether provided 0.108g (75%) of the hydroxamic acid as a white powder. Electrospray Mass Spec: 436.1 (M+H)⁺

Example 34**3-(4-But-2-ynyloxy-benzenesulfonyl)-amino-2,6-dimethoxy-isonicotinic acid**

The product of Example 30 (0.400g, 0.95 mmol) was hydrolyzed to the corresponding carboxylic acid using the procedure of Example 19 to provide 0.338g (100%) of 3-(4-but-2-ynyloxy-benzenesulfonyl)-amino-2,6-dimethoxy-isonicotinic acid. Electrospray Mass Spec: 407.2 (M+H)⁺

Example 35**3-(4-But-2-ynyloxy-benzenesulfonylamino)-N-hydroxy-2,6-dimethoxy-isonicotinamide**

The product of Example 34 (86 mg, 0.21 mmol) was dissolved in DMF (2 mL). To this solution was added hydroxylamine hydrochloride (123 mg, 1.88 mmol), 1-hydroxybenzotriazole (68 mg, 0.5 mmol), triethylamine (0.3 mL, 2.1 mmol), and 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (113 mg, 0.59 mmol). The reaction mixture was stirred overnight and then filtered to remove the white precipitate. The filtrate was then diluted with dichloromethane, and washed with water, brine, dried over Na₂SO₄, filtered, concentrated in vacuo to provide an orange oil. The residue was chromatographed on silica gel eluting with ethyl acetate

to provide 32 mg (36%) of the hydroxamic acid as a white solid. Electrospray Mass Spec: 422.2 (M+H)⁺.

Pharmacology

5 The ability of the compounds of the invention, or their pharmaceutically acceptable salts, to inhibit matrix metalloproteinases or TACE and, consequently, demonstrate their effectiveness for treating diseases modulated by matrix metalloproteinases or TACE is shown by the following in vitro assays.

10 Test Procedures for Measuring MMP-1, MMP-9, and MMP-13 Inhibition

These standard pharmacological test procedures are based on the cleavage of a thiopeptide substrates such as Ac-Pro-Leu-Gly(2-mercapto-4-methyl-pentanoyl)-Leu-Gly-OEt by the matrix metalloproteinases MMP-1, MMP-13 (collagenases) or MMP-9 (gelatinase), which results in the release of a substrate product that reacts
15 colorimetrically with DTNB (5,5'-dithiobis(2-nitro-benzoic acid)). The enzyme activity is measured by the rate of the color increase. The thiopeptide substrate is made up fresh as a 20 mM stock in 100% DMSO and the DTNB is dissolved in 100% DMSO as a 100 mM stock and stored in the dark at room temperature. Both the substrate and DTNB are diluted together to 1 mM with substrate buffer (50 mM
20 HEPES pH 7.5, 5 mM CaCl₂) before use. The stock of enzyme is diluted with buffer (50 mM HEPES, pH 7.5, 5 mM CaCl₂, 0.02% Brij) to the desired final concentration. The buffer, enzyme, vehicle or inhibitor, and DTNB/substrate are added in this order to a 96 well plate (total reaction volume of 200 µl) and the increase in color is monitored spectrophotometrically for 5 minutes at 405 nm on a
25 plate reader and the increase in color over time is plotted as a linear line.

Alternatively, a fluorescent peptide substrate is used. In this test procedure, the peptide substrate contains a fluorescent group and a quenching group. Upon cleavage of the substrate by an MMP, the fluorescence that is generated is quantitated on the fluorescence plate reader. The assay is run in HCBC assay buffer (50mM
30 HEPES, pH 7.0, 5 mM Ca⁺², 0.02% Brij, 0.5% Cysteine), with human recombinant MMP-1, MMP-9, or MMP-13. The substrate is dissolved in methanol and stored frozen in 1 mM aliquots. For the assay, substrate and enzymes are diluted in HCBC

buffer to the desired concentrations. Compounds are added to the 96 well plate containing enzyme and the reaction is started by the addition of substrate. The reaction is read (excitation 340 nm, emission 444 nm) for 10 min. and the increase in fluorescence over time is plotted as a linear line.

- 5 For either the thiopeptide or fluorescent peptide test procedures, the slope of the line is calculated and represents the reaction rate. The linearity of the reaction rate is confirmed ($r^2 > 0.85$). The mean ($x \pm \text{sem}$) of the control rate is calculated and compared for statistical significance ($p < 0.05$) with drug-treated rates using Dunnett's multiple comparison test. Dose-response relationships can be generated using
- 10 multiple doses of drug and IC_{50} values with 95% CI are estimated using linear regression.

Test Procedure for Measuring TACE Inhibition

- Using 96-well black microtiter plates, each well receives a solution composed of 10 μL TACE (final concentration $1\mu\text{g/mL}$), 70 μL Tris buffer, pH 7.4 containing
- 15 10% glycerol (final concentration 10 mM), and 10 μL of test compound solution in DMSO (final concentration $1\mu\text{M}$, DMSO concentration $<1\%$) and incubated for 10 minutes at room temperature. The reaction is initiated by addition of a fluorescent peptidyl substrate (final concentration 100 μM) to each well and then shaking on a shaker for 5 sec.

- 20 The reaction is read (excitation 340 nm, emission 420 nm) for 10 min. and the increase in fluorescence over time is plotted as a linear line. The slope of the line is calculated and represents the reaction rate.

- The linearity of the reaction rate is confirmed ($r^2 > 0.85$). The mean ($x \pm \text{sem}$) of the control rate is calculated and compared for statistical significance ($p < 0.05$)
- 25 with drug-treated rates using Dunnett's multiple comparison test. Dose-response relationships can be generate using multiple doses of drug and IC_{50} values with 95% CI are estimated using linear regression.

Human Monocytic THP-1 Cell Differentiation Assay For Soluble Proteins

- 30 (THP Soluble Protein Assay)

Mitogenic stimulation of THP-1 cells cause differentiation into macrophage like cells with concomitant secretion of tumor necrosis factor ($\text{TNF-}\alpha$) and TNF receptor (TNF-R p75/80 and TNF-R p55/60) and Interleukin-8 (IL-8), among other proteins. In addition, non-stimulated THP-1 cells shed both the p75/80 and the

p55/60 receptors over time. The release of membrane bound TNF- α and possibly TNF-R p75/80 and TNF-R p55/60, but not IL-8, is mediated by an enzyme called TNF- α converting enzyme or TACE. This assay can be used to demonstrate either an inhibitory or a stimulatory compound effect on this TACE enzyme and any cytotoxic consequence of such a compound.

THP-1 cells (from ATCC) are a human monocytic cell line which were obtained from the peripheral blood of a one year old male with acute monocytic leukemia. They can be grown in culture and differentiated into macrophage like cells by stimulation with mitogens.

For the assay, THP-1 cells are seeded from an ATCC stock which was previously grown and frozen back at 5×10^6 /ml/vial. One vial is seeded into a T25-flask with 16 mls of RPMI-1640 with glutamax (Gibco) media containing 10 % fetal bovine serum, 100 units/ml penicillin, 100 μ g/ml streptomycin, and 5×10^{-5} M 2-mercapto-ethanol (THP-1 media). Each vial of cells are cultured for about two weeks prior to being used for an assay and then are used for only 4 to 6 weeks to screen compounds. Cells are subcultured on Mondays and Thursdays to a concentration of 1×10^5 /ml.

To perform an assay, the THP-1 cells are co-incubated in a 24 well plate with 50 ml/well of a 24 mg/ml stock of Lipopolysaccharide (LPS) (Calbiochem Lot# B13189) at 37°C in 5% CO₂ at a concentration of 1.091×10^6 cells/ml (1.1 ml/well) for a total of 24 hours. At the same time, 50 ml/well of drug, vehicle or THP-1 media is plated in appropriate wells to give a final volume of 1.2 ml/well. Standard and test compounds are dissolved in DMSO at a concentration of 36 mM and diluted from here to the appropriate concentrations in THP-1 media and added to the wells at the beginning of the incubation period to give final concentrations of 100 mM, 30 mM, 10 mM, 3 mM, 1 mM, 300 nM, and 100 nM. Cell exposure to DMSO was limited to 0.1 % final concentration. Positive control wells were included in the experiment which had mitogen added but no drug. Vehicle control wells were included as well, which were identical to the positive control wells, except that DMSO was added to give a final concentration of 0.083%. Negative control wells were included in the experiment which had vehicle but no mitogen or drug added to the cells. Compounds can be evaluated for their effect on basal (non-stimulated) shedding of the receptors by replacing the LPS with 50 ml/well of THP-1 media. Plates are placed into an incubator set at 5% CO₂ and at 37°C. After 4 hours of incubation, 300 ml/well of tissue culture supernatant (TCS) is removed for use in an

TNF- α ELISA. Following 24 hours of incubation, 700 μ l/well of TCS is removed and used for analysis in TNF-R p75/80, TNF-R p55/60 and IL-8 ELISAs.

In addition, at the 24 hours timepoint, and the cells for each treatment group are collected by resuspension in 500 μ l/well of THP-1 media and transferred into a FACS tube. Two ml/tube of a 0.5 mg/ml stock of propidium iodide (PI) (Boehringer Mannheim cat. # 1348639) is added. The samples are run on a Becton Dickinson FaxCaliber FLOW cytometry machine and the amount of dye taken up by each cell is measured in the high red wavelength (FL3). Only cells with compromised membranes (dead or dying) can take up PI. The percent of live cells is calculated by the number of cells not stained with PI, divided by the total number of cells in the sample. The viability values calculated for the drug treated groups were compared to the viability value calculated for the vehicle treated mitogen stimulated group ("vehicle positive control") to determine the "percent change from control". This "percent change from control" value is an indicator of drug toxicity.

The quantity of soluble TNF- α , TNF-R p75/80 and TNF-R p55/60 and IL-8 in the TCS of the THP-1 cell cultures are obtained with commercially available ELISAs from R&D Systems, by extrapolation from a standard curve generated with kit standards. The number of cells that either take up or exclude PI are measured by the FLOW cytometry machine and visualized by histograms using commercially available Cytologic software for each treatment group including all controls.

Biological variability in the magnitude of the response of THP-1 cell cultures requires that experiments be compared on the basis of percent change from "vehicle positive control" for each drug concentration. Percent change in each soluble protein evaluated from the "vehicle positive control" was calculated for each compound concentration with the following formula:

$$\% \text{ Change} = \frac{\text{pg/ml (compound)} - \text{pg/ml (veh pos control)}}{\text{pg/ml (veh pos control)} - \text{pg/ml (veh neg control)}} \times 100$$

For the soluble protein (TNF- α , p75/80, p55/60, IL-8) studies under stimulated conditions, the mean pg/ml of duplicate wells were determined and the results expressed as percent change from "vehicle positive control". For the soluble protein (p75/80 and p55/60 receptors) studies under non-stimulated conditions, the mean pg/ml of duplicate wells were determined and the results expressed as percent change from "vehicle positive control" utilizing the following formula:

-40-

$$\% \text{ Change} = \frac{\text{pg/ml (compound neg control)} - \text{pg/ml (veh neg control)}}{\text{pg/ml (veh neg control)}} \times 100$$

IC₅₀ values for each compound are calculated by non-linear regression analysis
5 using customized software utilizing the JUMP statistical package.

For the cell viability studies, the viabilities (PI exclusion) of pooled duplicate wells were determined and the results expressed as % change from "vehicle positive control". The viability values calculated for the compound treated groups were
10 compared to the viability value calculated for the "vehicle positive control" to determine "percent change from control" as below. This value "percent change from control" is an indicator of drug toxicity.


$$\% \text{ Change} = \frac{\% \text{ live cells (compound)}}{\% \text{ live cells (veh pos control)}} - 1 \times 100$$

15

References:

- Bjornberg, F., Lantz, M., Olsson, I., and Gullberg, U. Mechanisms involved in the processing of the p55 and the p75 tumor necrosis factor (TNF) receptors to soluble
20 receptor forms. *Lymphokine Cytokine Res.* 13:203-211, 1994.
- Gatanaga, T., Hwang, C., Gatanaga, M., Cappuccini, F., Yamamoto, R., and Granger, G. The regulation of TNF mRNA synthesis, membrane expression, and release by PMA- and LPS-stimulated human monocytic THP-1 cells in vitro. *Cellular Immun.* 138:1-10, 1991.
- 25 Tsuchiya, S., Yamabe, M., Yamaguchi, Y., Kobayashi, Y., Konno, T., and Tada, K. Establishment and characterization of a human acute monocytic leukemia cell line (THP-1). *Int. J. Cancer.* 26:1711-176, 1980.

Results of the above *in vitro* matrix metalloproteinase inhibition, TACE
30 inhibition, and THP standard pharmacological test procedures are given in Table 1.

Table 1

Example #	R	MMP-1 ^a	MMP-9 ^a	MMP-13 ^a	TACE ^a	THP ^b
33	Me	-10,000	607	478	11	31
35	H	-	-	-	44	-

- 5 a) IC₅₀ (nM)
 b) % Inhibition @ 3μM

Based on the standard pharmacological test procedures described above, the compounds of this invention are useful in the treatment of disorders such as arthritis, tumor metastasis, tissue ulceration, abnormal wound healing, periodontal disease, graft rejection, insulin resistance, bone disease and HIV infection.

The compounds of this invention are also useful in treating or inhibiting pathological changes mediated by matrix metalloproteinases such as atherosclerosis, atherosclerotic plaque formation, reduction of coronary thrombosis from atherosclerotic plaque rupture, restenosis, MMP-mediated osteopenias, inflammatory diseases of the central nervous system, skin aging, angiogenesis, tumor metastasis, tumor growth, osteoarthritis, rheumatoid arthritis, septic arthritis, corneal ulceration, proteinuria, aneurysmal aortic disease, degenerative cartilage loss following traumatic joint injury, demyelinating diseases of the nervous system, cirrhosis of the liver, glomerular disease of the kidney, premature rupture of fetal membranes, inflammatory bowel disease, age related macular degeneration, diabetic retinopathy, proliferative vitreoretinopathy, retinopathy of prematurity, ocular inflammation, keratoconus, Sjogren's syndrome, myopia, ocular tumors, ocular angiogenesis/neovascularization and corneal graft rejection.

Compounds of this invention may be administered neat or with a pharmaceutical carrier to a patient in need thereof. The pharmaceutical carrier may be solid or liquid.

Applicable solid carriers can include one or more substances which may also act as flavoring agents, lubricants, solubilizers, suspending agents, fillers, glidants, compression aids, binders or tablet-disintegrating agents or an encapsulating material.

In powders, the carrier is a finely divided solid which is in admixture with the finely divided active ingredient. In tablets, the active ingredient is mixed with a carrier having the necessary compression properties in suitable proportions and compacted in the shape and size desired. The powders and tablets preferably contain up to 99% of the active ingredient. Suitable solid carriers include, for example, calcium phosphate, magnesium stearate, talc, sugars, lactose, dextrin, starch, gelatin, cellulose, methyl cellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidone, low melting waxes and ion exchange resins.

Liquid carriers may be used in preparing solutions, suspensions, emulsions, syrups and elixirs. The active ingredient of this invention can be dissolved or suspended in a pharmaceutically acceptable liquid carrier such as water, an organic solvent, a mixture of both or pharmaceutically acceptable oils or fat. The liquid carrier can contain other suitable pharmaceutical additives such as solubilizers, emulsifiers, buffers, preservatives, sweeteners, flavoring agents, suspending agents, thickening agents, colors, viscosity regulators, stabilizers or osmo-regulators. Suitable examples of liquid carriers for oral and parenteral administration include water (particularly containing additives as above, e.g., cellulose derivatives, preferable sodium carboxymethyl cellulose solution), alcohols (including monohydric alcohols and polyhydric alcohols, e.g., glycols) and their derivatives, and oils (e.g., fractionated coconut oil and arachis oil). For parenteral administration the carrier can also be an oily ester such as ethyl oleate and isopropyl myristate. Sterile liquid carriers are used in sterile liquid form compositions for parenteral administration.

Liquid pharmaceutical compositions which are sterile solutions or suspensions can be utilized by, for example, intramuscular, intraperitoneal or subcutaneous injection. Sterile solutions can also be administered intravenously. Oral administration may be either liquid or solid composition form.

The compounds of this invention may be administered rectally in the form of a conventional suppository. For administration by intranasal or intrabronchial inhalation or insufflation, the compounds of this invention may be formulated into an aqueous or partially aqueous solution, which can then be utilized in the form of an aerosol. The compounds of this invention may also be administered transdermally through the use of a transdermal patch containing the active compound and a carrier that is inert to the active compound, is non-toxic to the skin, and allows delivery of the agent for systemic absorption into the blood stream via the skin. The carrier may take any number of forms such as creams and ointments, pastes, gels, and occlusive devices. The creams and ointments may be viscous liquid or semi-solid emulsions of


either the oil in water or water in oil type. Pastes comprised of absorptive powders dispersed in petroleum or hydrophilic petroleum containing the active ingredient may also be suitable. A variety of occlusive devices may be used to release the active ingredient into the blood stream such as a semipermeable membrane covering a reservoir containing the active ingredient with or without a carrier, or a matrix containing the active ingredient. Other occlusive devices are known in the literature.

The dosage to be used in the treatment of a specific patient suffering a MMP or TACE dependent condition must be subjectively determined by the attending physician. The variables involved include the severity of the dysfunction, and the size, age, and response pattern of the patient. Treatment will generally be initiated with small dosages less than the optimum dose of the compound. Thereafter the dosage is increased until the optimum effect under the circumstances is reached. Precise dosages for oral, parenteral, nasal, or intrabronchial administration will be determined by the administering physician based on experience with the individual subject treated and standard medical principles.

Preferably the pharmaceutical composition is in unit dosage form, e.g., as tablets or capsules. In such form, the composition is sub-divided in unit dose containing appropriate quantities of the active ingredient; the unit dosage form can be packaged compositions, for example packed powders, vials, ampoules, prefilled syringes or sachets containing liquids. The unit dosage form can be, for example, a capsule or tablet itself, or it can be the appropriate number of any such compositions in package form.

CLAIMS

1. Hydroxamic acids of the formula:



B

where the C(=O)NHOH moiety and the -NR₅- moiety are bonded to adjacent carbons of group A; wherein

A is 5-6 membered heteroaryl having 1 to 3 heteroatoms selected from N, NR₅, S and O;

10 X is SO₂ or -P(O)R₁₀;

Y is aryl or 5-10 membered mono- or bi-cyclic heteroaryl having from 1 to three heteroatoms selected from N, NR₅, S and O, with the proviso that X and Z may not be bonded to adjacent atoms of Y;

Z is O, NH, CH₂ or S;

15 R₅ is hydrogen or alkyl of 1-6 carbon atoms;

R₆ and R₇ are each, independently, hydrogen, alkyl of 1-6 carbon atoms, -CN, -CCH₃;

R₈ is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, aryl, 5 to 10 membered heteroaryl having 1 to 3 heteroatoms selected from N, NR₅, S and O, or 5 to 9 membered heterocycloalkyl having 1 or 2 heteroatoms selected from N, NR₅, S and O;

R₉ is hydrogen, aryl, alkyl of 1-6 carbon atoms or cycloalkyl of 3-6 carbon atoms;

25 and R₁₀ is alkyl of 1-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, aryl, or heteroaryl; or a pharmaceutically acceptable salt thereof.

This Sheet Has Been Replaced





-45-


2. A compound of structure B according to Claim 1 wherein the ring atom of A adjacent the $-NR_5-$ group is carbon and has a substituent other than hydrogen.
- 5 3. A compound according to Claim 1 or 2 wherein Y is a phenyl ring substituted at the 1- and 4-positions by X and Z, respectively.
4. A compound according to any one of Claims 1 to 3 wherein X is SO_2 .
- 10 5. A compound according to any one of Claims 1 to 4 wherein Z is oxygen.
6. A compound according to any one of Claims 1 to 5 wherein R_6 and R_7 are hydrogen.
- 15 7. A compound according to any one of Claims 1 to 6 wherein R_8 is $-CH_2OH$ or methyl.
8. A compound according to Claim 1 which is (3-[methyl-(4-but-2-ynyloxy-benzenesulfonyl-amino)]-N-hydroxy-2,6-dimethoxy-isonicotinamide.
- 20 9. A compound according to Claim 1 which is 3-(4-But-2-ynyloxy-benzenesulfonylamino)-N-hydroxy-2,6-dimethoxy-isonicotinamide.
- 25 10. A process for preparing a compound of formula I which comprises one of the following:

This Sheet Has Been Replaced



-46-


- a) reacting a compound of formula V:



(V)

- wherein R_5 , R_6 , R_7 , R_8 , A, X, Y and Z are as defined above and Q is COOH or a reactive derivative thereof, with hydroxylamine to give a corresponding compound of formula B; or

- b) deprotecting a compound of formula VI:



(VI)

10

- wherein R_5 , R_6 , R_7 , R_8 , A, X, Y and Z are as defined above, and R_{30} is a protecting group, to give a compound of formula B;

- c) resolving a mixture (e.g racemate) of optically active isomers of a compound of formula B to isolate one enantiomer or diastereomer substantially free of the other enantiomer or diastereomers;


15

or

- d) acidifying a basic compound of formula B with a pharmaceutically acceptable acid to give a pharmaceutically acceptable salt.

20

11. A compound of the formula:



II

wherein R_6 and R_7 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, -CN, -CCH₃;

This Sheet Has Been Replaced






-47-

and R_5 is alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, phenyl, naphthyl, 5 to 10 membered heteroaryl having from 1 to 3 heteroatoms selected from N, NR , O or S, or 5 to 9 membered heterocycloalkyl having 1 or 2 heteroatoms selected from N, NR , O or S.


12. A compound of the formula



wherein R_6 and R_7 are each, independently, hydrogen, alkyl of 1-6 carbon atoms, -CN, -CCH₃;

R_8 is of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, phenyl, naphthyl, 5 to 10 membered heteroaryl having from 1 to 3 heteroatoms selected from N, NR , O or S, or 5 to 9 membered heterocycloalkyl having 1 or 2 heteroatoms selected from N, NR , O or S; and
J is fluorine, bromine, chlorine, 1,2,4-triazolyl, benzotriazolyl or imidazol-yl.

13. A method of inhibiting pathological changes mediated by $TNF-\alpha$ converting enzyme (TACE) in a mammal in need thereof which comprises administering to said mammal a therapeutically effective amount of a compound having the formula



This Sheet Has Been Replaced





where the C(=O)NHOH moiety and the -NR₅- moiety are bonded to adjacent carbons of group A; wherein

A is 5-6 membered heteroaryl having 1 to 3 heteroatoms selected from N, NR₅, S and O;

5 X is SO₂ or -P(O)R₁₀;

Y is aryl or 5-10 membered mono- or bi-cyclic heteroaryl having from 1 to three heteroatoms selected from N, NR₅, S and O, with the proviso that X and Z may not be bonded to adjacent atoms of Y;

Z is O, NH, CH₂ or S;

10 R₅ is hydrogen or alkyl of 1-6 carbon atoms;

R₆ and R₇ are each, independently, hydrogen, alkyl of 1-6 carbon atoms, -CN, -CCH₃;

R₈ is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms, alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, aryl, 5 to 15 10 membered heteroaryl having 1 to 3 heteroatoms selected from N, NR₅, S and O, or 5 to 9 membered heterocycloalkyl having 1 or 2 heteroatoms selected from N, NR₅, S and O;


R₉ is hydrogen, aryl, alkyl of 1-6 carbon atoms or cycloalkyl of 3-6 carbon atoms;

20 and R₁₀ is alkyl of 1-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, aryl, or heteroaryl; or a pharmaceutically acceptable salt thereof.

14. The method according to Claim 13 wherein the condition treated is rheumatoid arthritis, graft rejection, cachexia, inflammation, fever, insulin resistance, 25 septic shock, congestive heart failure, inflammatory disease of the central nervous system, inflammatory bowel disease or HIV infection.


This Sheet Has Been Replaced





-49-

15. A pharmaceutical composition comprising a compound having the formula



B

- 5 where the C(=O)NHOH moiety and the -NR₅- moiety are bonded to adjacent carbons of group A; wherein
- A is 5-6 membered heteroaryl having 1 to 3 heteroatoms selected from N, NR₅, S and O;
- X is SO₂ or -P(O)R₁₀;
- 10 Y is aryl or 5-10 membered mono- or bi-cyclic heteroaryl having from 1 to three heteroatoms selected from N, NR₅, S and O, with the proviso that X and Z may not be bonded to adjacent atoms of Y;
- Z is O, NH, CH₂ or S;
- 15 R₅ is hydrogen or alkyl of 1-6 carbon atoms;
- R₆ and R₇ are each, independently, hydrogen, alkyl of 1-6 carbon atoms, -CN, -CCH;
- R₈ is hydrogen, alkyl of 1-6 carbon atoms, alkenyl of 2-6 carbon atoms,
- 20 alkynyl of 2-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, aryl, 5 to 10 membered heteroaryl having 1 to 3 heteroatoms selected from N, NR₅, S and O, or 5 to 9 membered heterocycloalkyl having 1 or 2 heteroatoms selected from N, NR₅, S and O;
- 25 R₉ is hydrogen, aryl, alkyl of 1-6 carbon atoms or cycloalkyl of 3-6 carbon atoms;
- and R₁₀ is alkyl of 1-6 carbon atoms, cycloalkyl of 3-6 carbon atoms, aryl, or heteroaryl; or a pharmaceutically acceptable salt thereof; and a pharmaceutically
- 30 acceptable carrier.

This Sheet Has Been Replaced

