

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
22 February 2001 (22.02.2001)

PCT

(10) International Publication Number
WO 01/12574 A1

- (51) International Patent Classification⁷: C07B 43/04, 53/00, C07C 209/62
- (21) International Application Number: PCT/GB00/02867
- (22) International Filing Date: 25 July 2000 (25.07.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
9919118.1 14 August 1999 (14.08.1999) GB
- (71) Applicant (for all designated States except US): **AVECIA LIMITED** [GB/GB]; Hexagon House, Blackley, Manchester M9 8ZS (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **MARTIN, Juliette** [GB/GB]; P.O. BOX A38, Leeds Road, Huddersfield, West Yorkshire HD2 1FF (GB). **CAMPBELL, Lynne, Alison** [GB/GB]; Earles Road, Grangemouth, Stirlingshire, FK3 8XG (GB).
- (74) Agents: **REVELL, Christopher et al.**; Intellectual Property Group, Avecia Limited, Hexagon House, P.O. Box 42, Blackley, Manchester M9 8ZS (GB).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**
— With international search report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: TRANSFER HYDROGENATION PROCESS

(57) Abstract: A catalytic transfer hydrogenation process is provided. The process can be employed to transfer hydrogenate N-substituted imines and iminium salts, which are preferably prochiral. The catalyst employed in the process is preferably a metal complex with one hydrocarbyl or cyclopentadienyl ligand and which is also coordinated to defined bidentate ligands. Preferred metals include rhodium, ruthenium and iridium. Preferred bidentate ligands are diamines and aminoalcohols, particularly those comprising chiral centres. The hydrogen donor is advantageously a mixture of triethylamine and formic acid. A process for the production of primary and secondary amines using the catalytic transfer hydrogenation of the N-substituted imines and iminium salts is also provided.

WO 01/12574 A1

TRANSFER HYDROGENATION PROCESS

This invention relates to catalytic transfer hydrogenation, particularly in the presence of a complexed transition metal, and to a process of making optically active compounds.

Eguchi et al, Tetrahedron (including Tetrahedron Reports) 1998, 54(5-6), 705-714 discusses the reduction of ketones and imines in the presence of a NADH mimic. The asymmetric reduction of certain phosphonylimines using borohydride in the presence of a cobalt catalyst is discussed in EP0901966. The asymmetric transfer hydrogenation of imines and iminium substrates in the presence of certain metal catalysts with cyclopentadienyl ligands is discussed in WO98/42463. Uematsu et al, J. Am. Chem. Soc., 1996, 118(20), 4916-7 discusses the asymmetric transfer hydrogenation of imines in the presence of certain ruthenium catalysts.

According to a first aspect of the present invention there is provided a process for the transfer hydrogenation of a substrate wherein the substrate is reacted with a hydrogen donor in the presence of a transfer hydrogenation catalyst, wherein the substrate has the general formula



(1)

wherein:

X represents NR^3 or $(\text{NR}^4\text{R}^5)^+\text{Q}^-$;

Q^- represents a monovalent anion;

R^1 and R^2 each independently represents a hydrogen atom, an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl, an optionally substituted heterocyclyl group, a substituted carbonyl functional group, a substituted thiocarbonyl functional group or substituted imino functional group, R^1 & R^2 optionally being linked in such a way as to form an optionally substituted ring;

R^3 and R^4 represents $-\text{P}(\text{O})\text{R}^6\text{R}^7$, $-\text{P}(\text{O})\text{OR}^8\text{OR}^9$, $-\text{P}(\text{O})\text{OR}^8\text{OH}$, $-\text{P}(\text{O})(\text{OH})_2$, $-\text{P}(\text{O})\text{SR}^{10}\text{SR}^{11}$, $-\text{P}(\text{O})\text{SR}^{10}\text{SH}$, $-\text{P}(\text{O})(\text{SH})_2$, $-\text{P}(\text{O})\text{NR}^{12}\text{R}^{13}\text{NR}^{14}\text{R}^{15}$, $-\text{P}(\text{O})\text{NR}^{12}\text{R}^{13}\text{NHR}^{14}$, $-\text{P}(\text{O})\text{NHR}^{12}\text{NHR}^{14}$, $-\text{P}(\text{O})\text{NR}^{12}\text{R}^{13}\text{NH}_2$, $-\text{P}(\text{O})\text{NHR}^{12}\text{NH}_2$, $-\text{P}(\text{O})(\text{NH}_2)_2$, $-\text{P}(\text{O})\text{R}^6\text{OR}^8$, $-\text{P}(\text{O})\text{R}^6\text{OH}$, $-\text{P}(\text{O})\text{R}^6\text{SR}^{10}$, $-\text{P}(\text{O})\text{R}^6\text{SH}$, $-\text{P}(\text{O})\text{R}^6\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{R}^6\text{NHR}^{12}$, $-\text{P}(\text{O})\text{R}^6\text{NH}_2$, $-\text{P}(\text{O})\text{OR}^8\text{SR}^{10}$, $-\text{P}(\text{O})\text{OR}^8\text{SH}$, $-\text{P}(\text{O})\text{OHSR}^{10}$, $-\text{P}(\text{O})\text{OHSH}$, $-\text{P}(\text{O})\text{OR}^8\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{OR}^8\text{NHR}^{12}$, $-\text{P}(\text{O})\text{OR}^8\text{NH}_2$, $-\text{P}(\text{O})\text{OHNR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{OHNHR}^{12}$, $-\text{P}(\text{O})\text{OHNH}_2$, $-\text{P}(\text{O})\text{SR}^{10}\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{SR}^{10}\text{NHR}^{12}$, $-\text{P}(\text{O})\text{SR}^{10}\text{NH}_2$, $-\text{P}(\text{O})\text{SHNR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{SHNHR}^{12}$, $-\text{P}(\text{O})\text{SHNH}_2$, $-\text{P}(\text{S})\text{R}^6\text{R}^7$, $-\text{P}(\text{S})\text{OR}^8\text{OR}^9$, $-\text{P}(\text{S})\text{OR}^8\text{OH}$, $-\text{P}(\text{S})(\text{OH})_2$, $-\text{P}(\text{S})\text{SR}^{10}\text{SR}^{11}$, $-\text{P}(\text{S})\text{SR}^{10}\text{SH}$, $-\text{P}(\text{S})(\text{SH})_2$, $-\text{P}(\text{S})\text{NR}^{12}\text{R}^{13}\text{NR}^{14}\text{R}^{15}$, $-\text{P}(\text{S})\text{NR}^{12}\text{R}^{13}\text{NHR}^{14}$, $-\text{P}(\text{S})\text{NHR}^{12}\text{NHR}^{14}$, $-\text{P}(\text{S})\text{NR}^{12}\text{R}^{13}\text{NH}_2$, $-\text{P}(\text{S})\text{NHR}^{12}\text{NH}_2$, $-\text{P}(\text{S})(\text{NH}_2)_2$, $-\text{P}(\text{S})\text{R}^6\text{OR}^8$, $-\text{P}(\text{S})\text{R}^6\text{OH}$, $-\text{P}(\text{S})\text{R}^6\text{SR}^{10}$, $-\text{P}(\text{S})\text{R}^6\text{SH}$, $-\text{P}(\text{S})\text{R}^6\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{S})\text{R}^6\text{NHR}^{12}$, $-\text{P}(\text{S})\text{R}^6\text{NH}_2$, $-\text{P}(\text{S})\text{OR}^8\text{SR}^{10}$, $-\text{P}(\text{S})\text{OHSR}^{10}$, $-\text{P}(\text{S})\text{OR}^8\text{SH}$, $-\text{P}(\text{S})\text{OHSH}$, $-\text{P}(\text{S})\text{OR}^8\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{S})\text{OR}^8\text{NHR}^{12}$, $-\text{P}(\text{S})\text{OR}^8\text{NH}_2$.

13 FEB 2004

1a


-P(S)OHNR¹²R¹³, -P(S)OHNR¹², -P(S)OHNH₂, -P(S)SR¹⁰NR¹²R¹³, -P(S)SR¹⁰NHR¹²,
 -P(S)SR¹⁰NH₂, -P(S)SHNR¹²R¹³, -P(S)SHNR¹², -P(S)SHNH₂, -PR⁶R⁷, -POR⁸OR⁹,
 -PSR¹⁰SR¹¹, -PNR¹²R¹³NR¹⁴R¹⁵, -PR⁶OR⁸, -PR⁶SR¹⁰, -PR⁶NR¹²R¹³, -POR⁸SR¹⁰,
 -POR⁸NR¹²R¹³, -PSR¹⁰NR¹²R¹³, -S(O)R¹⁶, -S(O)₂R¹⁷, -COR¹⁸, -CO₂R¹⁹, or
 SiR²⁰R²¹R²².

R⁸ represents a hydrogen atom, an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl, or an optionally substituted heterocyclyl group;

R⁶ and R⁷ independently represent an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl, an optionally substituted heterocyclyl group or -N=CR²³R²⁴ where R²³ and R²⁴ are as defined for R¹; and

R⁹ to R²² each independently represents an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, one or more of R¹ & R⁶, R² & R⁷, R⁵ & R⁷, R⁵ & R⁸, R⁵ & R¹⁰, R⁶ & R¹², R¹ & R⁹, R² & R⁹, R⁸ & R⁹, R⁸ & R¹⁰, R⁸ & R¹², R¹ & R¹⁰, R² & R¹¹, R¹⁰ & R¹¹, R¹⁰ & R¹², R¹ & R¹², R² & R¹³, R¹² & R¹³, R¹ & R¹⁴, R² & R¹⁵, R¹⁴ & R¹⁵, R¹² & R¹⁴, R¹ & R¹⁶, R¹ & R¹⁸, R¹ & R¹⁹, R¹ & R²⁰, R² & R²¹, R²⁰ & R²¹ and R²¹ & R²² optionally being linked in such a way as to form an optionally substituted ring(s),

and wherein the transfer hydrogenation catalyst has the general formula:



wherein:

R²⁷ represents a neutral optionally substituted hydrocarbyl, a neutral optionally substituted perhalogenated hydrocarbyl, or an optionally substituted cyclopentadienyl ligand;

A represents -NR²⁸-, -NR²⁹-, -NHR²⁸, -NR²⁸R²⁹ or -NR²⁹R³⁰ where R²⁸ is H, C(O)R³⁰, SO₂R³⁰, C(O)NR³⁰R³⁴, C(S)NR³⁰R³⁴, C(=NR³⁴)SR³⁵ or C(=NR³⁴)OR³⁵, R²⁹ and R³⁰ each independently represents an optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, and R³⁴ and R³⁵ are each independently hydrogen or a group as defined for R³⁰;

B represents -O-, -OH, OR³¹, -S-, -SH, SR³¹, -NR³¹-, -NR³²-, -NHR³², -NR³¹R³², -NR³¹R³³, -PR³¹- or -PR³¹R³³ where R³² is H, C(O)R³³, SO₂R³³, C(O)NR³³R³⁶, C(S)NR³³R³⁶, C(=NR³⁶)SR³⁷ or C(=NR³⁶)OR³⁷, R³¹ and R³³ each independently represents an optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, and R³⁵ and R³⁷ are each independently hydrogen or a group as defined for R³³;

E represents a linking group wherein A and B are linked through 2, 3 or 4 atoms each independently being unsubstituted or substituted;

M is a group VIII transition metal; and

13 FEB 2004

Y represents an anionic group, a basic ligand or a vacant site; provided that when Y is not a vacant site that at least one of A or B carries a hydrogen atom.

When X represents $(NR^1R^5)^+Q^-$, compounds of formula (1) are iminium salts. Iminium salts include protonated iminium salts and quaternary iminium salts. Quaternary iminium salts are represented by compounds of formula (I) in which R^5 is not hydrogen.

Anions which may be represented by Q^- include halides, optionally substituted arylsulphonates, such as optionally substituted phenyl and naphthyl sulphonates, optionally substituted alkylsulphonates including halogenated alkylsulphonates, such as C_{1-20} alkylsulphonates, optionally substituted carboxylates, such as C_{1-10} alkyl and aryl carboxylates, ions derived from the polyhalogenation of boron, phosphorous or antimony, and other common inorganic ions for example perchlorate. Particular examples of anions are bromide, chloride, iodide, hydrogen sulphate, tosylate, formate, acetate, tetrafluoroborate, hexafluorophosphate, hexafluoroantimonate, perchlorate, trifluoromethanesulphonate and trifluoroacetate. Preferred anions include bromide, chloride, iodide, formate and trifluoroacetate, and particularly preferred anions include iodide, formate and trifluoroacetate.

Hydrocarbyl groups which may be represented by one or more of R^1 , R^2 , and R^5 to R^{24} , include alkyl, alkenyl, alkynyl and aryl groups, and any combination thereof, such as aralkyl and alkaryl, for example benzyl groups.

Alkyl groups which may be represented by one or more of R^1 , R^2 , and R^5 to R^{24} include linear and branched alkyl groups comprising up to 20 carbon atoms, particularly from 1 to 7 carbon atoms and preferably from 1 to 5 carbon atoms. When the alkyl groups are branched, the groups often comprise up to 10 branched chain carbon atoms, preferably up to 4 branched chain atoms. In certain embodiments, the alkyl group may be cyclic, commonly comprising from 3 to 10 carbon atoms in the largest ring and optionally featuring one or more bridging rings. Examples of alkyl groups which may be represented by R^1 , R^2 , and R^5 to R^{24} include methyl, ethyl, propyl, 2-propyl, butyl, 2-butyl, t-butyl and cyclohexyl groups.

Alkenyl groups which may be represented by one or more of R^1 , R^2 , and R^5 to R^{24} include C_{2-20} , and preferably C_{2-6} alkenyl groups. One or more carbon - carbon double

bonds may be present. The alkenyl group may carry one or more substituents, particularly phenyl substituents. Examples of alkenyl groups include vinyl, styryl and indenyl groups. When either of R^1 or R^2 represents an alkenyl group, a carbon - carbon double bond is preferably located at the position β to the $C=X$ moiety. When either of R^1 or R^2 represents an alkenyl group, the compound of formula (1) is preferably an α,β -unsaturated iminium compound.

Alkynyl groups which may be represented by one or more of R^1 , R^2 , and R^5 to R^{24} include C_{2-20} , and preferably C_{2-10} alkynyl groups. One or more carbon - carbon triple bonds may be present. The alkynyl group may carry one or more substituents, particularly phenyl substituents. Examples of alkynyl groups include ethynyl, propyl and phenylethynyl groups. When either of R^1 or R^2 represents an alkynyl group, a carbon - carbon triple bond is preferably located at the position β to the $C=X$ moiety. When either of R^1 or R^2 represents an alkynyl group, the compound of formula (1) preferably has the triple bond in conjugation with the iminium group.

Aryl groups which may be represented by one or more of R^1 , R^2 , and R^5 to R^{24} may contain 1 ring or 2 or more fused rings which may include cycloalkyl, aryl or heterocyclic rings. Examples of aryl groups which may be represented by R^1 , R^2 , and R^5 to R^{24} include phenyl, tolyl, fluorophenyl, chlorophenyl, bromophenyl, trifluoromethylphenyl, anisyl, naphthyl and ferrocenyl groups.

Perhalogenated hydrocarbyl groups which may be represented by one or more of R^1 , R^2 , and R^5 to R^{24} independently include perhalogenated alkyl and aryl groups, and any combination thereof, such as aralkyl and alkaryl groups. Examples of perhalogenated alkyl groups which may be represented by R^1 , R^2 , and R^5 to R^{24} include $-CF_3$ and $-C_2F_5$.

Heterocyclic groups which may be represented by one or more of R^1 , R^2 , and R^5 to R^{24} independently include aromatic, saturated and partially unsaturated ring systems and may comprise 1 ring or 2 or more fused rings which may include cycloalkyl, aryl or heterocyclic rings. The heterocyclic group will contain at least one heterocyclic ring, the largest of which will commonly comprise from 3 to 7 ring atoms in which at least one atom is carbon and at least one atom is any of N, O, S or P. When either of R^1 or R^2 represents or comprises a heterocyclic group, the atom in R^1 or R^2 bonded to the $C=X$ group is preferably a carbon atom. Examples of heterocyclic groups which may be represented by R^1 , R^2 , and R^5 to R^{24} include pyridyl, pyrimidyl, pyrrolyl, thiophenyl, furanyl, indolyl, quinolyl, isoquinolyl, imidazolyl and triazolyl groups.

When any of R^1 , R^2 , and R^5 to R^{24} is a substituted hydrocarbyl or heterocyclic group, the substituent(s) should be such so as not to adversely affect the rate or stereoselectivity of the reaction. Optional substituents include halogen, cyano, nitro, hydroxy, amino, imino, thiol, acyl, hydrocarbyl, perhalogenated hydrocarbyl, heterocyclyl, hydrocarbyloxy, mono or di-hydrocarbylamino, hydrocarbylthio, esters, carboxy,

carbonates, amides, sulphonyl and sulphonamido groups wherein the hydrocarbyl groups are as defined for R^1 above. One or more substituents may be present. R^1 , R^2 , and R^5 to R^{24} may each contain one or more chiral centres.

When any of R^1 & R^2 , R^1 & R^6 , R^2 & R^7 , R^6 & R^7 , R^6 & R^8 , R^6 & R^{10} , R^6 & R^{12} , R^1 & R^8 , R^2 & R^9 , R^8 & R^9 , R^8 & R^{10} , R^8 & R^{12} , R^1 & R^{10} , R^2 & R^{11} , R^{10} & R^{11} , R^{10} & R^{12} , R^1 & R^{12} , R^2 & R^{13} , R^{12} & R^{13} , R^1 & R^{14} , R^2 & R^{15} , R^{14} & R^{15} , R^{12} & R^{14} , R^1 & R^{16} , R^1 & R^{18} , R^1 & R^{19} , R^1 & R^{20} , R^2 & R^{21} , R^{20} & R^{21} and R^{21} & R^{22} are linked in such a way that when taken together with the atoms to which they are attached that a ring is formed, it is preferred that these be 5, 6 or 7 membered rings. The rings formed in this way may additionally be fused to each other or to other ring systems. Examples of rings which may be so formed include oxazaphospholidenes, dioxaphospholans, phospholans, phosphorinans, dioxaphosphorinans and benzodioxaphospholans. The rings may be optionally substituted or may be fused to other rings.

Substituted carbonyl functional groups which may be represented by one or more of R^1 , R^2 , R^{23} and R^{24} include aldehyde, ketone, acid and ester groups, for example $-COR^{25}$, $-CO_2R^{25}$, and $-CONR^{25}R^{26}$ wherein R^{25} and R^{26} independently represent hydrogen, an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group as defined hereinbefore for R^5 .

Substituted thiocarbonyl functional groups which may be represented by one or more of R^1 , R^2 , R^{23} and R^{24} include thioaldehyde, thioketone, thioacid and thioester groups, for example $-CSR^{25}$, $-CSOR^{25}$ and $-CSNR^{25}R^{26}$ wherein R^{25} and R^{26} independently represent hydrogen, an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group as defined hereinbefore for R^5 .


Substituted imino functional groups which may be represented by one or more of R^1 , R^2 , R^{23} and R^{24} include unsubstituted imines and substituted imine and iminium groups, for example $-C(=NR^{25})R^{26}$ and $-C(=X)R^{26}$ wherein X is as hereinbefore defined and R^{25} and R^{26} independently represent hydrogen, an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group as defined hereinbefore for R^5 .

When one of R^1 or R^2 is represented by a group selected from a carbonyl functional group, a thiocarbonyl functional group and an imino functional group, it is preferred that the other of R^1 or R^2 is represented by a hydrogen atom, an optionally substituted hydrocarbyl group, a perhalogenated hydrocarbyl group or an optionally substituted heterocyclyl group.

When one of R^{23} or R^{24} is represented by a group selected from a carbonyl functional group, a thiocarbonyl functional group and an imino functional group, it is preferred that the other of R^{23} or R^{24} being represented by a hydrogen atom, an optionally

substituted hydrocarbonyl group, a perhalogenated hydrocarbonyl group or an optionally substituted heterocyclyl group.

When one of R^1 & R^6 or R^2 & R^7 is linked in such a way as to form an optionally substituted ring and the ring so formed contains more than one carbon nitrogen double bond (imino group), it is preferred that nitrogen-phosphorous heterocycle so formed has both nitrogens are attached to a common phosphorous atom. Examples of such nitrogen-phosphorous heterocycles include



In certain preferred embodiments, R^1 , R^2 , and R^5 to R^{24} are all independently C_{1-6} alkyl or are a combination of aryl, particularly phenyl, C_{1-6} alkyl and C_{6-10} aralkyl. Substituents may be present, particularly substituents para to the $C=X$ group when one or more of R^1 , R^2 , and R^5 to R^{24} is a phenyl group.

In many preferred embodiments, when R^3 or R^4 is a group selected from
 $-P(O)R^6R^7$, $-P(O)OR^8OR^9$, $-P(O)OR^8OH$, $-P(O)SR^{10}SR^{11}$, $-P(O)SR^{10}SH$,
 $-P(O)NR^{12}R^{13}NR^{14}R^{15}$, $-P(O)NR^{12}R^{13}NHR^{14}$, $-P(O)NHR^{12}NHR^{14}$, $-P(O)NR^{12}R^{13}NH_2$,
 $-P(O)NHR^{12}NH_2$, $-P(O)R^6OR^8$, $-P(O)R^6OH$, $-P(O)R^6SR^{10}$, $-P(O)R^6SH$, $-P(O)R^6NR^{12}R^{13}$,
 $-P(O)R^6NHR^{12}$, $-P(O)R^6NH_2$, $-P(O)OR^8SR^{10}$, $-P(O)OR^8SH$, $-P(O)OHSR^{10}$,
 $-P(O)OR^8NR^{12}R^{13}$, $-P(O)OR^8NHR^{12}$, $-P(O)OR^8NH_2$, $-P(O)OHNHR^{12}R^{13}$, $-P(O)OHNHR^{12}$,
 $-P(O)SR^{10}NR^{12}R^{13}$, $-P(O)SR^{10}NHR^{12}$, $-P(O)SR^{10}NH_2$, $-P(O)SHNR^{12}R^{13}$, $-P(O)SHNHR^{12}$,
 $-P(S)R^6R^7$, $-P(S)OR^8OR^9$, $-P(S)OR^8OH$, $-P(S)SR^{10}SR^{11}$, $-P(S)SR^{10}SH$, $-P(S)(SH)_2$,
 $-P(S)NR^{12}R^{13}NR^{14}R^{15}$, $-P(S)NR^{12}R^{13}NHR^{14}$, $-P(S)NHR^{12}NHR^{14}$, $-P(S)NR^{12}R^{13}NH_2$,
 $-P(S)NHR^{12}NH_2$, $-P(S)R^6OR^8$, $-P(S)R^6OH$, $-P(S)R^6SR^{10}$, $-P(S)R^6SH$, $-P(S)R^6NR^{12}R^{13}$,
 $-P(S)R^6NHR^{12}$, $-P(S)R^6NH_2$, $-P(S)OR^8SR^{10}$, $-P(S)OHSR^{10}$, $-P(S)OR^8SH$, $-P(S)OR^8NR^{12}R^{13}$,
 $-P(S)OR^8NHR^{12}$, $-P(S)OR^8NH_2$, $-P(S)OHNHR^{12}R^{13}$, $-P(S)OHNHR^{12}$, $-P(S)SR^{10}NR^{12}R^{13}$,
 $-P(S)SR^{10}NHR^{12}$, $-P(S)SR^{10}NH_2$, $-P(S)SHNR^{12}R^{13}$, $-P(S)SHNHR^{12}$, $-PR^6R^7$, $-POR^8OR^9$,
 $-PSR^{10}SR^{11}$, $-PNR^{12}R^{13}NR^{14}R^{15}$, $-PR^6OR^8$, $-PR^6SR^{10}$, $-PR^6NR^{12}R^{13}$, $-POR^8SR^{10}$,
 $-POR^8NR^{12}R^{13}$, $-PSR^{10}NR^{12}R^{13}$, or $SiR^{20}R^{21}R^{22}$, the groups R^6 to R^{15} and R^{20} to R^{22} are selected to be the same and more preferably they are selected to all be phenyl or ethyl groups. This may have the advantage that synthesis of intermediates is simplified. However, in certain embodiments it may be preferable for more than one of the groups R^6 to R^{15} or R^{20} to R^{22} present to be different, in which case it is most preferable that each R^6

to R¹⁵ or R²⁰ to R²² group present is selected to be one of methyl, ethyl, propyl, i-propyl, butyl, sec-butyl, t-butyl or phenyl.

Preferably, R³ and R⁴ are electron withdrawing groups, such as -P(O)R⁶R⁷, -P(O)OR⁸OR⁹, -P(O)OR⁸OH, -P(O)(OH)₂, -P(O)SR¹⁰SR¹¹, -P(O)SR¹⁰SH, -P(O)(SH)₂,
 5 -P(O)NR¹²R¹³NR¹⁴R¹⁵, -P(O)NR¹²R¹³NHR¹⁴, -P(O)NHR¹²NHR¹⁴, -P(O)NR¹²R¹³NH₂,
 -P(O)NHR¹²NH₂, -P(O)(NH₂)₂, -P(O)R⁶OR⁸, -P(O)R⁶OH, -P(O)R⁶SR¹⁰, -P(O)R⁶SH,
 -P(O)R⁶NR¹²R¹³, -P(O)R⁶NHR¹², -P(O)R⁶NH₂, -P(O)OR⁸SR¹⁰, -P(O)OR⁸SH, -P(O)OHSR¹⁰,
 -P(O)OHSH, -P(O)OR⁸NR¹²R¹³, -P(O)OR⁸NHR¹², -P(O)OR⁸NH₂, -P(O)OHNR¹²R¹³,
 -P(O)OHNHR¹², -P(O)OHNH₂, -P(O)SR¹⁰NR¹²R¹³, -P(O)SR¹⁰NHR¹², -P(O)SR¹⁰NH₂,
 10 -P(O)SHNR¹²R¹³, -P(O)SHNHR¹², -P(O)SHNH₂, -P(S)R⁶R⁷, -P(S)OR⁸OR⁹, -P(S)OR⁸OH, -
 P(S)(OH)₂, -P(S)SR¹⁰SR¹¹, -P(S)SR¹⁰SH, -P(S)(SH)₂, -P(S)NR¹²R¹³NR¹⁴R¹⁵,
 -P(S)NR¹²R¹³NHR¹⁴, -P(S)NHR¹²NHR¹⁴, -P(S)NR¹²R¹³NH₂, -P(S)NHR¹²NH₂, -P(S)(NH₂)₂,
 -P(S)R⁶OR⁸, -P(S)R⁶OH, -P(S)R⁶SR¹⁰, -P(S)R⁶SH, -P(S)R⁶NR¹²R¹³, -P(S)R⁶NHR¹²,
 -P(S)R⁶NH₂, -P(S)OR⁸SR¹⁰, -P(S)OHSR¹⁰, -P(S)OR⁸SH, -P(S)OHSH, -P(S)OR⁸NR¹²R¹³,
 15 -P(S)OR⁸NHR¹², -P(S)OR⁸NH₂, -P(S)OHNR¹²R¹³, -P(S)OHNHR¹², -P(S)OHNH₂,
 -P(S)SR¹⁰NR¹²R¹³, -P(S)SR¹⁰NHR¹², -P(S)SR¹⁰NH₂, -P(S)SHNR¹²R¹³, -P(S)SHNHR¹²,
 -P(S)SHNH₂, -S(O)R¹⁶, -S(O)₂R¹⁷, -COR¹⁸, and -CO₂R¹⁹. It is especially preferred that R³
 and R⁴ are groups selected from -P(O)R⁶R⁷, -P(O)OR⁸OR⁹, -P(O)R⁶OR⁸, -P(S)R⁶R⁷,
 -P(S)OR⁸OR⁹, -P(S)R⁶OR⁸, -S(O)R¹⁶, -COR¹⁸, and -CO₂R¹⁹. It is most preferred that R³
 20 and R⁴ are groups selected from -P(O)R⁶R⁷, -P(O)OR⁸OR⁹.


When either R³ or R⁴ is a group selected from -P(O)R⁶R⁷, -P(O)OR⁸OR⁹, -P(O)OR⁸OH, -P(O)SR¹⁰SR¹¹, -P(O)SR¹⁰SH, -P(O)NR¹²R¹³NR¹⁴R¹⁵, -P(O)NR¹²R¹³NHR¹⁴,
 -P(O)NHR¹²NHR¹⁴, -P(O)NR¹²R¹³NH₂, -P(O)NHR¹²NH₂, -P(O)R⁶OR⁸, -P(O)R⁶OH,
 -P(O)R⁶SR¹⁰, -P(O)R⁶SH, -P(O)R⁶NR¹²R¹³, -P(O)R⁶NHR¹², -P(O)R⁶NH₂, -P(O)OR⁸SR¹⁰,
 25 -P(O)OR⁸SH, -P(O)OHSR¹⁰, -P(O)OR⁸NR¹²R¹³, -P(O)OR⁸NHR¹², -P(O)OR⁸NH₂,
 -P(O)OHNR¹²R¹³, -P(O)OHNHR¹², -P(O)SR¹⁰NR¹²R¹³, -P(O)SR¹⁰NHR¹², -P(O)SR¹⁰NH₂,
 -P(O)SHNR¹²R¹³, -P(O)SHNHR¹², -P(S)R⁶R⁷, -P(S)OR⁸OR⁹, -P(S)OR⁸OH, -P(S)SR¹⁰SR¹¹,
 -P(S)SR¹⁰SH, -P(S)(SH)₂, -P(S)NR¹²R¹³NR¹⁴R¹⁵, -P(S)NR¹²R¹³NHR¹⁴, -P(S)NHR¹²NHR¹⁴,
 -P(S)NR¹²R¹³NH₂, -P(S)NHR¹²NH₂, -P(S)R⁶OR⁸, -P(S)R⁶OH, -P(S)R⁶SR¹⁰, -P(S)R⁶SH,
 30 -P(S)R⁶NR¹²R¹³, -P(S)R⁶NHR¹², -P(S)R⁶NH₂, -P(S)OR⁸SR¹⁰, -P(S)OHSR¹⁰, -P(S)OR⁸SH,
 -P(S)OR⁸NR¹²R¹³, -P(S)OR⁸NHR¹², -P(S)OR⁸NH₂, -P(S)OHNR¹²R¹³, -P(S)OHNHR¹²,
 -P(S)SR¹⁰NR¹²R¹³, -P(S)SR¹⁰NHR¹², -P(S)SR¹⁰NH₂, -P(S)SHNR¹²R¹³, -P(S)SHNHR¹²,
 -PR⁶R⁷, -POR⁸OR⁹, -PSR¹⁰SR¹¹, -PNR¹²R¹³NR¹⁴R¹⁵, -PR⁶OR⁸, -PR⁶SR¹⁰, -PR⁶NR¹²R¹³,
 -POR⁸SR¹⁰, -POR⁸NR¹²R¹³, -PSR¹⁰NR¹²R¹³, -S(O)R¹⁶, -S(O)₂R¹⁷, -COR¹⁸, and -CO₂R¹⁹, it is
 35 preferred that the groups represented by R⁶ to R¹⁹ are independently selected from alkyl or
 aryl groups, for example methyl, ethyl, propyl, i-propyl, butyl, sec-butyl, t-butyl and phenyl
 groups, and preferably C₁₋₆alkyl or C₆₋₁₂aryl groups.

The groups R^6 to R^{22} may contain chiral centres, or may be selected such that the atom to which they are bonded is a chiral centre.

In certain preferred embodiments, one of R^1 and R^2 is an alkyl group and the other is aryl or heterocyclyl, X is NR^3 where R^3 is either $-P(O)R^6R^7$ or $-P(O)OR^8OR^9$ wherein R^6 to R^9 is an alkyl or aryl group, more preferably a C_{1-4} alkyl group, a phenyl group or a phenyl group substituted with one or more C_{1-4} alkyl groups.

Most advantageously, the compound of formula (1) is prochiral, such that the hydrogenated product comprises a chiral atom to which R^1 , R^2 and X are each bonded. Such an asymmetric transfer hydrogenation process forms an especially preferred aspect of the present invention. Most commonly, when the compound of formula (1) is prochiral, R^1 and R^2 are different, and neither is hydrogen. Preferably, one of R^1 and R^2 is aliphatic and the other is aryl or heterocyclyl.

Examples of compounds of formula (1) include



wherein:

R^a = Ph, Naphthyl, CH_2Ph , hexyl, iPr, tBu, Et, or Me

R^b = Ph, Naphthyl, CH_2Ph , hexyl, iPr, tBu, Et, or Me

R^c = Ph, Naphthyl, CH_2Ph , hexyl, iPr, tBu, Et, or Me; and

R^d = $PO(Ph)_2$, or $PO(Et)_2$.

Hydrogen donors include hydrogen, primary and secondary alcohols, primary and secondary amines, carboxylic acids and their esters and amine salts, readily dehydrogenatable hydrocarbons, clean reducing agents, and any combination thereof.

Primary and secondary alcohols which may be employed as hydrogen donors comprise commonly from 1 to 10 carbon atoms, preferably from 2 to 7 carbon atoms, and more preferably 3 or 4 carbon atoms. Examples of primary and secondary alcohols which may be represented as hydrogen donors include methanol, ethanol, propan-1-ol, propan-2-ol, butan-1-ol, butan-2-ol, cyclopentanol, cyclohexanol, benzylalcohol, and menthol.

When the hydrogen donor is an alcohol, secondary alcohols are preferred, especially propan-2-ol and butan-2-ol.

Primary and secondary amines which may be employed as hydrogen donors comprise commonly from 1 to 20 carbon atoms, preferably from 2 to 14 carbon atoms, and more preferably 3 or 8 carbon atoms. Examples of primary and secondary amines which may be represented as hydrogen donors include ethylamine, propylamine, isopropylamine, butylamine, isobutylamine, hexylamine, diethylamine, dipropylamine, diisopropylamine, dibutylamine, diisobutylamine, dihexylamine, benzylamine, dibenzylamine and piperidine. When the hydrogen donor is an amine, primary amines are preferred, especially primary amines comprising a secondary alkyl group, particularly isopropylamine and isobutylamine.

Carboxylic acids or their esters which may be employed as hydrogen donors comprise commonly from 1 to 10 carbon atoms, preferably from 1 to 3 carbon atoms. In certain embodiments, the carboxylic acid is advantageously a beta-hydroxy-carboxylic acid. Esters may be derived from the carboxylic acid and a C₁₋₁₀ alcohol. Examples of carboxylic acids which may be employed as hydrogen donors include formic acid, lactic acid, ascorbic acid and mandelic acid. The most preferred carboxylic acid is formic acid. In certain preferred embodiments, when a carboxylic acid is employed as hydrogen donor, at least some of the carboxylic acid is preferably present as salt, preferably an amine, ammonium or metal salt. Preferably, when a metal salt is present the metal is selected from the alkali or alkaline earth metals of the periodic table, and more preferably is selected from the group I elements, such as lithium, sodium or potassium. Amines which may be used to form such salts include both aromatic and non-aromatic amines, also primary, secondary and tertiary amines and comprise typically from 1 to 20 carbon atoms. Tertiary amines, especially trialkylamines, are preferred. Examples of amines which may be used to form salts include trimethylamine, triethylamine, diisopropylethylamine and pyridine. The most preferred amine is triethylamine. When at least some of the carboxylic acid is present as an amine salt, particularly when a mixture of formic acid and triethylamine is employed, the mole ratio of acid to amine is between 1:1 and 50:1 and preferably between 1:1 and 10:1, and most preferably about 5:2. When at least some of the carboxylic acid is present as a metal salt, particularly when a mixture of formic acid and a group I metal salt is employed, the mole ratio of acid to metal ions present is between 1:1 and 50:1 and preferably between 1:1 and 10:1, and most preferably about 2:1. The ratios of acid to salts may be maintained during the course of the reaction by the addition of either component, but usually by the addition of the carboxylic acid.


Readily dehydrogenatable hydrocarbons which may be employed as hydrogen donors comprise hydrocarbons which have a propensity to aromatise or hydrocarbons which have a propensity to form highly conjugated systems. Examples of readily

dehydrogenatable hydrocarbons which may be employed by as hydrogen donors include cyclohexadiene, cyclohexene, tetralin, dihydrofuran and terpenes.

Clean reducing agents which may be represented as hydrogen donors comprise reducing agents with a high reduction potential, particularly those having a reduction potential relative to the standard hydrogen electrode of greater than about -0.1eV, often greater than about -0.5eV, and preferably greater than about -1eV. Examples of clean reducing agents which may be represented as hydrogen donors include hydrazine and hydroxylamine.


The most preferred hydrogen donors are propan-2-ol, butan-2-ol, triethylammonium formate and a mixture of triethylammonium formate and formic acid.

Transfer hydrogenation catalysts may include such catalysts such as a) the chiral Ruthenium (II) catalysts developed for ketone reduction which are disclosed in Chem. Rev., 1998, 98, 2607 see Table 2; b) the Zhang tridentate bis(oxazolinylmethyl)amine catalysts and related catalysts as disclosed in J. Am. Chem. Soc., 1998, 120, 3817, Tet. Let., 1997, 38(37), 6565 and in WO99/24410 (particularly the bis(phenyloxazolin-2-yl)amine and related catalysts discussed therein); and c) the transition metal, particularly group VIII metal, complexes with chiral ligands of formula



wherein AR is any aromatic or ring structure and R', R'' and R''' are each independently selected from aryl, alkyl, aralkyl, ring-substituted aralkyl, substituted aryl and combinations thereof as disclosed in US 5,767,276, the catalysts of a), b) and c) being incorporated herein by reference.

Preferred transfer hydrogenation catalysts for use in the process of the present invention have the general formula:



wherein:

R²⁷ represents a neutral optionally substituted hydrocarbyl, a neutral optionally substituted perhalogenated hydrocarbyl, or an optionally substituted cyclopentadienyl ligand;

A represents -NR²⁸-, -NR²⁹-, -NHR²⁸-, -NR²⁸R²⁹ or -NR²⁹R³⁰ where R²⁸ is H, C(O)R³⁰, SO₂R³⁰, C(O)NR³⁰R³⁴, C(S)NR³⁰R³⁴, C(=NR³⁴)SR³⁵ or C(=NR³⁴)OR³⁵, R²⁹ and R³⁰ each independently represents an optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, and R³⁴ and R³⁵ are each independently hydrogen or a group as defined for R³⁰;

B represents -O-, -OH, OR^{31} , -S-, -SH, SR^{31} , $-NR^{31}-$, $-NR^{32}-$, $-NHR^{32}$, $-NR^{31}R^{32}$, $-NR^{31}R^{33}$, $-PR^{31}-$ or $-PR^{31}R^{33}$ where R^{32} is H, $C(O)R^{33}$, SO_2R^{33} , $C(O)NR^{33}R^{36}$, $C(S)NR^{33}R^{36}$, $C(=NR^{36})SR^{37}$ or $C(=NR^{36})OR^{37}$, R^{31} and R^{33} each independently represents an optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, and R^{36} and R^{37} are each independently hydrogen or a group as defined for R^{33} ;

E represents a linking group;

M represents a metal capable of catalysing transfer hydrogenation; and

Y represents an anionic group, a basic ligand or a vacant site;

provided that when Y is not a vacant site that at least one of A or B carries a hydrogen atom.

The catalytic species is believed to be substantially as represented in the above formula. It may be introduced on a solid support.

Optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or optionally substituted heterocyclyl groups which may be represented by R^{29-31} or R^{33-35} are as defined hereinbefore for R^5 .

The neutral optionally substituted hydrocarbyl or perhalogenated hydrocarbyl ligand which may be represented by R^{27} includes optionally substituted aryl and alkenyl ligands.

Optionally substituted aryl ligands which may be represented by R^{27} may contain 1 ring or 2 or more fused rings which include cycloalkyl, aryl or heterocyclic rings. Preferably, the ligand comprises a 6 membered aromatic ring. The ring or rings of the aryl ligand are often substituted with hydrocarbyl groups. The substitution pattern and the number of substituents will vary and may be influenced by the number of rings present, but often from 1 to 6 hydrocarbyl substituent groups are present, preferably 2, 3 or 6 hydrocarbyl groups and more preferably 6 hydrocarbyl groups. Preferred hydrocarbyl substituents include methyl, ethyl, iso-propyl, menthyl, neomenthyl and phenyl. Particularly when the aryl ligand is a single ring, the ligand is preferably benzene or a substituted benzene. When the ligand is a perhalogenated hydrocarbyl, preferably it is a polyhalogenated benzene such as hexachlorobenzene or hexafluorobenzene. When the hydrocarbyl substituents contain enantiomeric and/or diastereomeric centres, it is preferred that the enantiomerically and/or diastereomerically purified forms of these are used. Benzene, p-cymyl, mesitylene and hexamethylbenzene are especially preferred ligands.

Optionally substituted alkenyl ligands which may be represented by R^{27} include C_{2-30} , and preferably C_{6-12} , alkenes or cycloalkenes with preferably two or more carbon-carbon double bonds, preferably only two carbon-carbon double bonds. The carbon-carbon double bonds may optionally be conjugated to other unsaturated systems which

may be present, but are preferably conjugated to each other. The alkenes or cycloalkenes may be substituted preferably with hydrocarbyl substituents. When the alkene has only one double bond, the optionally substituted alkenyl ligand may comprise two separate alkenes. Preferred hydrocarbyl substituents include methyl, ethyl, iso-propyl and phenyl. Examples of optionally substituted alkenyl ligands include cyclo-octa-1,5-diene and 2,5-norbornadiene. Cyclo-octa-1,5-diene is especially preferred.

Optionally substituted cyclopentadienyl groups which may be represented by R^{27} includes cyclopentadienyl groups capable of eta-5 bonding. The cyclopentadienyl group is often substituted with from 1 to 5 hydrocarbyl groups, preferably with 3 to 5 hydrocarbyl groups and more preferably with 5 hydrocarbyl groups. Preferred hydrocarbyl substituents include methyl, ethyl and phenyl. When the hydrocarbyl substituents contain enantiomeric and/or diastereomeric centres, it is preferred that the enantiomerically and/or diastereomerically purified forms of these are used. Examples of optionally substituted cyclopentadienyl groups include cyclopentadienyl, pentamethyl-cyclopentadienyl, pentaphenylcyclopentadienyl, tetraphenylcyclopentadienyl, ethyltetramethylpentadienyl, menthyltetraphenylcyclopentadienyl, neomenthyl-tetraphenylcyclopentadienyl, menthylcyclopentadienyl, neomenthylcyclopentadienyl, tetrahydroindenyl, menthyltetrahydroindenyl and neomenthyltetrahydroindenyl groups. Pentamethylcyclopentadienyl is especially preferred.

When either A or B is an amide group represented by $-NR^{28}$ -, $-NHR^{28}$, $NR^{28}R^{29}$, $-NR^{32}$ -, $-NHR^{32}$ or $NR^{31}R^{32}$ wherein R^{29} and R^{31} are as hereinbefore defined, and where R^{28} or R^{32} is an acyl group represented by $-C(O)R^{30}$ or $-C(O)R^{33}$, R^{30} and R^{33} independently are often linear or branched C_{1-7} alkyl, C_{1-6} -cycloalkyl or aryl, for example phenyl. Examples of acyl groups which may be represented by R^{28} or R^{33} include benzoyl, acetyl and halogenoacetyl, especially trifluoroacetyl, groups.

When either A or B is present as a sulphonamide group represented by $-NR^{28}$ -, $-NHR^{28}$, $NR^{28}R^{29}$, $-NR^{32}$ -, $-NHR^{32}$ or $NR^{31}R^{32}$ wherein R^{29} and R^{31} are as hereinbefore defined, and where R^{28} or R^{32} is a sulphonyl group represented by $-S(O)_2R^{30}$ or $-S(O)_2R^{33}$, R^{30} and R^{33} independently are often linear or branched C_{1-6} alkyl, C_{1-6} cycloalkyl or aryl, for example phenyl. Preferred sulphonyl groups include methanesulphonyl, trifluoromethanesulphonyl and especially p-toluenesulphonyl groups and naphthylsulphonyl groups.

When either of A or B is present as a group represented by $-NR^{28}$ -, $-NHR^{28}$, $NR^{28}R^{29}$, $-NR^{32}$ -, $-NHR^{32}$ or $NR^{31}R^{32}$ wherein R^{29} and R^{31} are as hereinbefore defined, and where R^{28} or R^{32} is a group represented by $C(O)NR^{30}R^{34}$, $C(S)NR^{30}R^{34}$, $C(=NR^{34})SR^{35}$, $C(=NR^{34})OR^{35}$, $C(O)NR^{33}R^{36}$, $C(S)NR^{33}R^{36}$, $C(=NR^{36})SR^{37}$ or $C(=NR^{36})OR^{37}$, R^{30} and R^{33} independently are often linear or branched C_{1-6} alkyl, such as methyl, ethyl, isopropyl, C_{1-6} cycloalkyl or aryl, for example phenyl, groups and R^{34-37} are often each independently

hydrogen or linear or branched C₁₋₈alkyl, such as methyl, ethyl, isopropyl, C₁₋₈cycloalkyl or aryl, for example phenyl, groups.

When B is present as a group represented by -OR³¹, -SR³¹, -PR³¹- or -PR³¹R³³, R³¹ and R³³ independently are often linear or branched C₁₋₈alkyl, such as methyl, ethyl, isopropyl, C₁₋₈cycloalkyl or aryl, for example phenyl.

It will be recognised that the precise nature of A and B will be determined by whether A and/or B are formally bonded to the metal or are coordinated to the metal via a lone pair of electrons.


The groups A and B are connected by a linking group E. The linking group E achieves a suitable conformation of A and B so as to allow both A and B to bond or coordinate to the metal, M. A and B are commonly linked through 2, 3 or 4 atoms. The atoms in E linking A and B may carry one or more substituents. The atoms in E, especially the atoms alpha to A or B, may be linked to A and B, in such a way as to form a heterocyclic ring, preferably a saturated ring, and particularly a 5, 6 or 7-membered ring. Such a ring may be fused to one or more other rings. Often the atoms linking A and B will be carbon atoms. Preferably, one or more of the carbon atoms linking A and B will carry substituents in addition to A or B. Substituent groups include those which may substitute R¹, as defined above. Advantageously, any such substituent groups are selected to be groups which do not coordinate with the metal, M. Preferred substituents include halogen, cyano, nitro, sulphonyl, hydrocarbyl, perhalogenated hydrocarbyl and heterocyclyl groups as defined above. Most preferred substituents are C₁₋₆ alkyl groups, and phenyl groups. Most preferably, A and B are linked by two carbon atoms, and especially an optionally substituted ethyl moiety. When A and B are linked by two carbon atoms, the two carbon atoms linking A and B may comprise part of an aromatic or aliphatic cyclic group, particularly a 5, 6 or 7-membered ring. Such a ring may be fused to one or more other such rings. Particularly preferred are embodiments in which E represents a 2 carbon atom separation and one or both of the carbon atoms carries an optionally substituted aryl group as defined above or E represents a 2 carbon atom separation which comprises a cyclopentane or cyclohexane ring, optionally fused to a phenyl ring.

E preferably comprises part of a compound having at least one stereospecific centre. Where any or all of the 2, 3 or 4 atoms linking A and B are substituted so as to define at least one stereospecific centre on one or more of these atoms, it is preferred that at least one of the stereospecific centres be located at the atom adjacent to either group A or B. When at least one such stereospecific centre is present, it is advantageously present in an enantiomerically purified state.

When B represents -O- or -OH, and the adjacent atom in E is carbon, it is preferred that B does not form part of a carboxylic group.

Compounds which may be represented by A-E-B, or from which A-E-B may be derived by deprotonation, are often aminoalcohols, including 4-aminoalkan-1-ols, 1-aminoalkan-4-ols, 3-aminoalkan-1-ols, 1-aminoalkan-3-ols, and especially 2-aminoalkan-1-ols, 1-aminoalkan-2-ols, 3-aminoalkan-2-ols and 2-aminoalkan-3-ols, and particularly 2-aminoethanols or 3-aminopropanols, or are diamines, including 1,4-diaminoalkanes, 1,3-diaminoalkanes, especially 1,2- or 2,3- diaminoalkanes and particularly ethylenediamines. Further aminoalcohols that may be represented by A-E-B are 2-aminocyclopentanols and 2-aminocyclohexanols, preferably fused to a phenyl ring. Further diamines that may be represented by A-E-B are 1,2-diaminocyclopentanes and 1,2-diaminocyclohexanes, preferably fused to a phenyl ring. The amino groups may advantageously be N-tosylated. When a diamine is represented by A-E-B, preferably at least one amino group is N-tosylated. The aminoalcohols or diamines are advantageously substituted, especially on the linking group, E, by at least one alkyl group, such as a C₁₋₄-alkyl, and particularly a methyl, group or at least one aryl group, particularly a phenyl group.

Specific examples of compounds which can be represented by A-E-B and the protonated equivalents from which they may be derived are:



Preferably, the enantiomerically and/or diastereomerically purified forms of these are used. Examples include (1S,2R)-(+)-norephedrine, (1R,2S)-(+)-cis-1-amino-2-indanol, (1S,2R)-2-amino-1,2-diphenylethanol, (1S,2R)-(-)-cis-1-amino-2-indanol, (1R,2S)-(-)-norephedrine, (S)-(+)-2-amino-1-phenylethanol, (1R,2S)-2-amino-1,2-diphenylethanol, N-tosyl-(1R,2R)-1,2-diphenylethylenediamine, N-tosyl-(1S,2S)-1,2-diphenylethylenediamine, (1R,2S)-cis-1,2-indandiamine, (1S,2R)-cis-1,2-indandiamine, (R)-(-)-2-pyrrolidinemethanol and (S)-(+)-2-pyrrolidinemethanol.

Metals which may be represented by M include metals which are capable of catalysing transfer hydrogenation. Preferred metals include transition metals, more preferably the metals in Group VIII of the Periodic Table, especially ruthenium, rhodium or


iridium. When the metal is ruthenium it is preferably present in valence state II. When the metal is rhodium or iridium it is preferably present in valence state I when R^{27} is a neutral optionally substituted hydrocarbyl or a neutral optionally substituted perhalogenated hydrocarbyl ligand, and preferably present in valence state III when R^{27} is an optionally substituted cyclopentadienyl ligand

Anionic groups which may be represented by Y include hydride, hydroxy, hydrocarbyloxy, hydrocarbylamino and halogen groups. Preferably when a halogen is represented by Y, the halogen is chloride. When a hydrocarbyloxy or hydrocarbylamino group is represented by Y, the group may be derived from the deprotonation of the hydrogen donor utilised in the reaction.

Basic ligands which may be represented by Y include water, C_{1-4} alcohols, C_{1-8} primary or secondary amines, or the hydrogen donor which is present in the reaction system. A preferred basic ligand represented by Y is water.

Most preferably, the nature of A-E-B, R^{27} and Y are chosen such that the catalyst is chiral. When such is the case, an enantiomerically and/or diastereomerically purified form is preferably employed. Such catalysts are most advantageously employed in asymmetric transfer hydrogenation processes. In many embodiments, the chirality of the catalyst is derived from the nature of A-E-B.

Examples of catalysts which may be employed in the process of the present invention include



The process is carried out preferably in the presence of a base, especially when Y is not a vacant site. The pK_a of the base is preferably at least 8.0, especially at least 10.0. Convenient bases are the hydroxides, alkoxides and carbonates of alkali metals; tertiary amines and quaternary ammonium compounds. Preferred bases are sodium 2-propoxide and triethylamine. When the hydrogen donor is not an acid, the quantity of base used can be up to 10.0, commonly up to 5.0, often up to 3.0, frequently up to 2.5 and especially in the range 1.0 to 3.5, by moles of the catalyst. When the hydrogen donor is an acid, the catalyst may be contacted with a base prior to the introduction of the hydrogen donor. In

such a case, the mole ratio of base to catalyst prior to the introduction of the hydrogen donor is often from 1:1 to 3:1, and preferably about 1:1.

Although gaseous hydrogen may be present, the process is normally operated in the absence of gaseous hydrogen since it appears to be unnecessary.

Advantageously, the process is carried out in the substantial absence of carbon dioxide.

Preferably, the process is carried out under a substantially inert atmosphere, for example nitrogen or argon.

When the product(s) from dehydrogenation of the hydrogen donor is volatile, for example boils at under 100°C, the removal of this volatile product is preferred. The removal can be accomplished by distillation preferably at less than atmospheric pressure or by use of inert gas sparging. When reduced pressure distillation is employed, the pressure is often no more than 6.66×10^4 Pa (500 mmHg), commonly no more than 2.66×10^4 Pa (200 mmHg), preferably in the range of from 6.66×10^2 to 1.33×10^4 Pa (5 to 100 mmHg), and most preferably from 1.33×10^3 to 1.06×10^4 Pa (10 to 80 mmHg). When the product(s) from dehydrogenation of the hydrogen donor is a gaseous material, for example when formic acid is present as a hydrogen donor, the removal is most preferably accomplished by the use of inert gas sparging, with for example nitrogen.

Suitably the process is carried out at temperatures in the range of from minus 78 to plus 150°C, preferably from minus 20 to plus 110°C and more preferably from minus 5 to plus 60°C. The initial concentration of the substrate, a compound of formula (1), is suitably in the range 0.05 to 1.0 and, for convenient larger scale operation, can be for example up to 6.0 more especially 0.25 to 2.0, on a molar basis. The molar ratio of the substrate to catalyst is suitably no less than 50:1 and can be up to 50000:1, preferably between 100:1 and 5000:1 and more preferably between 200:1 and 2000:1. The hydrogen donor is preferably employed in a molar excess over the substrate, especially from 5 to 50 fold or, if convenience permits, greater, for example up to 500 fold. After reaction, the mixture is worked up by standard procedures.

During the reaction a solvent may be present, preferably a polar solvent, for example methanol, ethanol or i-propanol, more preferably a polar aprotic solvent, for example acetonitrile, dimethylformamide or dichloromethane. Conveniently, the hydrogen donor may be the solvent when the hydrogen donor is a liquid at the reaction temperature, or it may be used in combination with other solvents. In certain embodiments it is preferable to use a diluent. Diluents include non-polar solvents such as toluene. Usually it is preferred to operate in substantial absence of water, but water may be present in the reaction. If the hydrogen donor or the reaction solvent is not miscible with water and the desired product is water soluble, it may be desirable to have water present as a second phase extracting the product, pushing the equilibrium and preventing loss of product optical purity as the reaction proceeds. The concentration of substrate may be chosen to optimise reaction time, yield and enantiomeric excess.

The catalytic species is believed to be substantially as represented in the above formula. It may be employed as an oligomer or metathesis product, on a solid support or may be generated in situ.

In certain embodiments it has been found that certain catalysts are preferred for the transfer hydrogenation of imines and iminium salts. Catalysts in which A-E-B is derived from N-tosyldiamines, preferably mono-N-tosyldiamines, particularly mono-N-tosylated ethylenediamines, are preferred. Especially, M is also ruthenium (II) and R^{27} is an aryl group or cyclopentadienyl group, or M is iridium (I) or rhodium (I) and R^{27} is cyclooctadiene, or M is iridium (III) or rhodium (III) and R^{27} is a cyclopentadienyl group. Further, triethylamine is preferably employed as a base, a mixture of formic acid and triethylamine in the preferred ratio of 5:2 (formic acid:triethylamine) is preferably employed as hydrogen donor. When an iminium salt is present, it is preferably a protonated imine, or is a methylated or benzylated imine with an iodide, formate or trifluoroacetate counter ion. It is believed that when Y is not a vacant site, R^{27} is a neutral ligand and when M is rhodium or iridium and is in valence state (I), A-E-B attaches to M by means of two dative bonds (the lone pairs of the heteroatoms in both A and B coordinate to M). However, when Y is not a vacant site, R^{27} is a cyclopentadienyl ligand and when M is rhodium or iridium and is in valence state (III), A-E-B attaches to M by means of one dative and one formal bond. Whereas when Y is not a vacant site, R^{27} is a neutral ligand and when M is ruthenium and is in valence state (II), A-E-B attaches to M by means of one dative and one formal bond.

The catalyst can be made by reacting a metal aryl, alkenyl or cyclopentadienyl halide complex with a compound of formula A-E-B as defined above or a protonated equivalent from which it may be derived, and, where Y represents a vacant site, reacting the product thereof with a base. The metal aryl or alkenyl halide complex preferably has the formula $[MR^{27}Z]_2$ when M is ruthenium (II) and has the formula $[MR^{27}Z]_2$ when M is iridium (I) or rhodium (I), wherein R^{27} is an aryl or alkenyl ligand as defined above, and Z represents a halide, particularly chloride. The metal cyclopentadienyl halide complex preferably has the formula $[MR^{27}Z]_2$ or $[MR^{27}Z]_4$ when M is ruthenium (II) and has the formula $[MR^{27}Z]_2$ when M is iridium (III) or rhodium (III), wherein R^{27} is an optionally substituted cyclopentadienyl ligand as defined above, and Z represents a halide, particularly chloride.

For the preparation of the catalysts according to the present invention, a solvent is preferably present. Suitable reaction temperatures are in the range 0-100, for example 20-70, °C, often giving reaction times of 0.5-24.0 h. After reaction is complete, the catalyst may if desired be isolated, but is more conveniently stored as the solution or used soon after preparation. The solution can contain the hydrogen donor and this, if a secondary alcohol, may be present in or used as the solvent for steps (a) and/or (b). The preparation

and after-handling should preferably be under an inert atmosphere, and particularly in carbon dioxide and oxygen-free conditions.

The catalyst or catalyst solution is generally treated with base either just prior to use in a transfer hydrogenation reaction, or during use. This can be accomplished by adding base to the catalyst in solution, or to the compound of formula (1) in solution, or by addition to the transfer hydrogenation reaction.

Transfer hydrogenation can be accomplished by transferring the solution of catalyst to a solution of substrate, a compound of general formula I. Alternatively a solution of substrate can be added to a solution of catalyst. Base may be pre-added to the catalyst solution and/or the substrate solution, or can be added later. The hydrogen donor if not already present in the catalyst solution may be added to the substrate solution, or may be added to the reaction mixture.

The imine and iminium salt compounds of formula (1) can generally be obtained by known literature methods. Iminium salts, for example, can be prepared by the quaternisation of imines, such as by treatment of imines with alkylating agents.

N-phosphinyl imines can be synthesised from N-hydroxy imines by treatment with a halophosphine. N-Hydroxy imines being readily available from the corresponding aldehyde or ketone by treatment with hydroxylamine. A similar approach can be used for the synthesis of certain N-sulphonylimines whereby N-hydroxy imines are treated with a halosulphoxyl compound in the presence of a base

Alternatively, N-phosphinyl imines can be synthesised from aldehydes or ketones by treatment with phosphinic amides in the presence of a condensation reagent, such as titanium tetrachloride, and a base.

Many of the N-phosphinyl, N-sulphonyl, N-sulphoxyl and N-carboxy imines can be synthesised from aldehydes or ketones by treatment with the corresponding phosphinamide, sulphonamide, sulphoxamide or carboxamides under dehydrating conditions, such azeotropic water removal, often carried out in the presence of an acid catalyst, such as p-toluene sulphonic acid or trifluoroacetic acid, or a Lewis acid catalyst, treatments with molar equivalents or excesses of drying agents, for example molecular sieves or magnesium sulphate, or treatments with combined lewis acid and dehydrating agents such as titanium tetrachloride or titanium tetraisopropoxide.

N-Carboxy and N-sulphoxyl imines may also be prepared by reaction of ketones with aza Wittig reagents.


N-Sulphoxyl imines may also be prepared by the reaction of an organometallic derivatives of imines, such as an N-lithiated imine, with chiral sulphoxides such as menthyl p-toluene sulphinate.

13 FEB 2004

N-Silylimines can be prepared from organometallic derivative of silazanes such as lithium di(trimethyl)silazamide, and ketones, or by the reaction of organometallic derivatives of imines, such as an N-lithiated imine, with a halosilane.


Furthermore, either by in situ cleavage of the R^3 or R^4 groups of the amines produced by the process of first aspect of present invention under the process conditions, or by further treatment by for example acid or base hydrolysis, a ready route to primary and secondary amines can be accessed.

According to a second aspect of the present invention there is provided a process for the production of primary or secondary amines



comprising the steps of


b) generating a substrate of general formula (1) from a carbonyl compound (2),



in which X represents NR^3 or $(\text{NR}^4\text{R}^5)^+\text{Q}^-$, and Q^- represents a monovalent anion,

d) reacting the substrate of general formula (1) with a hydrogen donor in the presence of a transfer hydrogenation catalyst, and

e) removal of R^3 or R^4 to give an amine of formula (3)



wherein:

R^1 and R^2 each independently represents a hydrogen atom, an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl, an optionally substituted heterocyclyl group, a substituted carbonyl functional group, a substituted thiocarbonyl functional group or substituted imino functional group, R^1 & R^2 optionally being linked in such a way as to form an optionally substituted ring;

R^3 and R^4 represents $-\text{P}(\text{O})\text{R}^6\text{R}^7$, $-\text{P}(\text{O})\text{OR}^8\text{OR}^9$, $-\text{P}(\text{O})\text{OR}^8\text{OH}$, $-\text{P}(\text{O})(\text{OH})_2$, $-\text{P}(\text{O})\text{SR}^{10}\text{SR}^{11}$, $-\text{P}(\text{O})\text{SR}^{10}\text{SH}$, $-\text{P}(\text{O})(\text{SH})_2$, $-\text{P}(\text{O})\text{NR}^{12}\text{R}^{13}\text{NR}^{14}\text{R}^{15}$, $-\text{P}(\text{O})\text{NR}^{12}\text{R}^{13}\text{NHR}^{14}$, $-\text{P}(\text{O})\text{NHR}^{12}\text{NHR}^{14}$, $-\text{P}(\text{O})\text{NR}^{12}\text{R}^{13}\text{NH}_2$, $-\text{P}(\text{O})\text{NHR}^{12}\text{NH}_2$, $-\text{P}(\text{O})(\text{NH}_2)_2$, $-\text{P}(\text{O})\text{R}^6\text{OR}^8$, $-\text{P}(\text{O})\text{R}^6\text{OH}$, $-\text{P}(\text{O})\text{R}^6\text{SR}^{10}$, $-\text{P}(\text{O})\text{R}^6\text{SH}$, $-\text{P}(\text{O})\text{R}^6\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{R}^6\text{NHR}^{12}$, $-\text{P}(\text{O})\text{R}^6\text{NH}_2$, $-\text{P}(\text{O})\text{OR}^8\text{SR}^{10}$, $-\text{P}(\text{O})\text{OR}^8\text{SH}$, $-\text{P}(\text{O})\text{OHSR}^{10}$, $-\text{P}(\text{O})\text{OHS}$, $-\text{P}(\text{O})\text{OR}^8\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{OR}^8\text{NHR}^{12}$, $-\text{P}(\text{O})\text{OR}^8\text{NH}_2$, $-\text{P}(\text{O})\text{OHNR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{OHNHR}^{12}$, $-\text{P}(\text{O})\text{OHNH}_2$, $-\text{P}(\text{O})\text{SR}^{10}\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{SR}^{10}\text{NHR}^{12}$, $-\text{P}(\text{O})\text{SR}^{10}\text{NH}_2$, $-\text{P}(\text{O})\text{SHNR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{SHNHR}^{12}$, $-\text{P}(\text{O})\text{SHNH}_2$, $-\text{P}(\text{S})\text{R}^6\text{R}^7$, $-\text{P}(\text{S})\text{OR}^8\text{OR}^9$, $-\text{P}(\text{S})\text{OR}^8\text{OH}$, $-\text{P}(\text{S})(\text{OH})_2$.

13 FEB 2004

18a

-P(S)SR¹⁰SR¹¹, -P(S)SR¹⁰SH, -P(S)(SH)₂, -P(S)NR¹²R¹³NR¹⁴R¹⁵, -P(S)NR¹²R¹³NHR¹⁴,
 -P(S)NHR¹²NHR¹⁴, -P(S)NR¹²R¹³NH₂, -P(S)NHR¹²NH₂, -P(S)(NH₂)₂, -P(S)R⁶OR⁹,
 -P(S)R⁶OH, -P(S)R⁶SR¹⁰, -P(S)R⁶SH, -P(S)R⁶NR¹²R¹³, -P(S)R⁶NHR¹², -P(S)R⁶NH₂,
 -P(S)OR⁹SR¹⁰, -P(S)OHSR¹⁰, -P(S)OR⁹SH, -P(S)OHSH, -P(S)OR⁹NR¹²R¹³,
 -P(S)OR⁹NHR¹², -P(S)OR⁹NH₂, -P(S)OHNR¹²R¹³, -P(S)OHNHR¹², -P(S)OHNH₂,
 -P(S)SR¹⁰NR¹²R¹³, -P(S)SR¹⁰NHR¹², -P(S)SR¹⁰NH₂, -P(S)SHNR¹²R¹³,
 -P(S)SHNHR¹², -P(S)SHNH₂, -PR⁸R⁷, -POR⁸OR⁹, -PSR¹⁰SR¹¹, -PNR¹²R¹³NR¹⁴R¹⁵,
 -PR⁸OR⁹, -PR⁸SR¹⁰, -PR⁸NR¹²R¹³, -POR⁸SR¹⁰, -POR⁸NR¹²R¹³, -PSR¹⁰NR¹²R¹³,
 -S(O)R¹⁶, -S(O)₂R¹⁷, -COR¹⁸, -CO₂R¹⁹, or SiR²⁰R²¹R²²;


R⁵ represents a hydrogen atom, an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl, or an optionally substituted heterocyclyl group;

R³⁸ represents a hydrogen atom or the group R⁵;

R⁶ and R⁷ independently represent an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl, an optionally substituted heterocyclyl group or -N=CR²³R²⁴ where R²³ and R²⁴ are as defined for R¹; and

R⁸ to R²² each independently represents an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, one or more of R¹ & R⁸, R² & R⁷, R⁵ & R⁷, R⁵ & R⁸, R⁵ & R¹⁰, R⁵ & R¹², R¹ & R⁹, R² & R⁹, R⁸ & R⁹, R⁸ & R¹⁰, R⁸ & R¹², R¹ & R¹⁰, R² & R¹¹, R¹⁰ & R¹¹, R¹⁰ & R¹², R¹ & R¹², R² & R¹³, R¹² & R¹³, R¹ & R¹⁴, R² & R¹⁵, R¹⁴ & R¹⁵, R¹² & R¹⁴, R¹ & R¹⁶, R¹ & R¹⁸, R¹ & R¹⁹, R¹ & R²⁰, R² & R²¹, R²⁰ & R²¹ and R²¹ & R²² optionally being linked in such a way as to form an optionally substituted ring(s).

and the transfer hydrogenation catalyst is of general formula:



wherein:

R²⁷ represents a neutral optionally substituted hydrocarbyl, a neutral optionally substituted perhalogenated hydrocarbyl, or an optionally substituted cyclopentadienyl ligand;

A represents -NR²⁸-, -NR²⁹-, -NHR²⁸, -NR²⁸R²⁹ or -NR²⁸R³⁰ where R²⁸ is H, C(O)R³⁰, SO₂R³⁰, C(O)NR³⁰R³⁴, C(S)NR³⁰R³⁴, C(=NR³⁴)SR³⁵ or C(=NR³⁴)OR³⁶, R²⁸ and R³⁰ each independently represents an optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, and R³⁴ and R³⁵ are each independently hydrogen or a group as defined for R³⁰;

B represents -O-, -OH, OR³¹, -S-, -SH, SR³¹, -NR³¹-, -NR³²-, -NHR³², -NR³¹R³², -NR³¹R³³, -PR³¹- or -PR³¹R³³ where R³² is H, C(O)R³³, SO₂R³³, C(O)NR³³R³⁶, C(S)NR³³R³⁶, C(=NR³⁶)SR³⁷ or C(=NR³⁶)OR³⁷, R³¹ and R³³ each independently

represents an optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, and R^{36} and R^{37} are each independently hydrogen or a group as defined for R^{33} ;

E represents a linking group wherein A and B are linked through 2, 3 or 4 atoms each independently being unsubstituted or substituted;


M is a group VIII transition metal; and

Y represents an anionic group, a basic ligand or a vacant site;

provided that when Y is not a vacant site that at least one of A or B carries a hydrogen atom.


Preferably, when a substrate of formula (1) or an amine of formula (3) comprises one or more optionally substituted ring(s), it is preferred that only one or more of R^1 & R^2 , R^6 & R^7 , R^8 & R^9 , R^{10} & R^{11} , R^{12} & R^{13} , R^{14} & R^{15} , R^{16} & R^{17} , R^{18} & R^{19} , R^{20} & R^{21} and R^{22} & R^{23} are optionally linked in such a way as to form an optionally substituted ring(s).

When the substrate of general formula (1) is an imine [i.e. $X = NR^3$], the step of generating the substrate of general formula (1) from a carbonyl compound (2),



is preferably accomplished by treating a carbonyl compound of formula (2) with a substituted amino compound R^3NH_2 .

When the substrate of general formula (1) is an iminium salt [i.e. $X = (NR^4R^5)^+Q^-$], the step of generating the substrate of general formula (1) from a carbonyl compound (2),



is preferably accomplished by treating a carbonyl compound of formula (2) with a substituted amino compound R^4NH_2 and quaternizing or protonating the resulting imine to give the iminium salt.

In certain embodiments, particularly in the generation of imines or iminium salts wherein R^3 or R^4 contain a phosphorous oxygen double bond, it is preferable in the step of generating the substrate of general formula (1) from a carbonyl compound (2) first to treat the carbonyl compound of formula (2) with hydroxylamine to generate the corresponding oxime, and then to react the oxime with an activated species containing the groups that will form R^3 or R^4 , such as halo derivative, for example $ClP(Ph)_2$.

When reacting the substrate of general formula (1) with a hydrogen donor in the presence of a transfer hydrogenation catalyst to give an amine, the step is carried out in accordance with reference to the first aspect of the present invention as described herein before.

The step of removal of the R^3 or R^4 may be carried out by any means which is generally known to cleave groups represented by R^3 or R^4 which are attached to an amine. Preferably, R^3 or R^4 are removed either by treatment with acid, treatment with base, hydrogenation or treatment with nucleophilic agents, such as fluoride, the method employed being selected according to the nature of R^3 or R^4 .

In many embodiments, R^3 or R^4 , especially when R^3 or R^4 is a group selected from $-P(O)R^6R^7$, $-P(O)OR^8OR^9$, $-P(O)OR^8OH$, $-P(O)(OH)_2$, $-P(O)SR^{10}SR^{11}$, $-P(O)SR^{10}SH$, $-P(O)(SH)_2$, $-P(O)NR^{12}R^{13}NR^{14}R^{15}$, $-P(O)NR^{12}R^{13}NHR^{14}$, $-P(O)NHR^{12}NHR^{14}$, $-P(O)NR^{12}R^{13}NH_2$, $-P(O)NHR^{12}NH_2$, $-P(O)(NH_2)_2$, $-P(O)R^6OR^8$, $-P(O)R^6OH$, $-P(O)R^6SR^{10}$, $-P(O)R^6SH$, $-P(O)R^6NR^{12}R^{13}$, $-P(O)R^6NHR^{12}$, $-P(O)R^6NH_2$, $-P(O)OR^8SR^{10}$, $-P(O)OR^8SH$, $-P(O)OHSR^{10}$, $-P(O)OHSH$, $-P(O)OR^8NR^{12}R^{13}$, $-P(O)OR^8NHR^{12}$, $-P(O)OR^8NH_2$, $-P(O)OHNHR^{12}R^{13}$, $-P(O)OHNHR^{12}$, $-P(O)OHNH_2$, $-P(O)SR^{10}NR^{12}R^{13}$, $-P(O)SR^{10}NHR^{12}$, $-P(O)SR^{10}NH_2$, $-P(O)SHNR^{12}R^{13}$, $-P(O)SHNHR^{12}$, $-P(O)SHNH_2$, $-P(S)R^6R^7$, $-P(S)OR^8OR^9$, $-P(S)OR^8OH$, $-P(S)(OH)_2$, $-P(S)SR^{10}SR^{11}$, $-P(S)SR^{10}SH$, $-P(S)(SH)_2$, $-P(S)NR^{12}R^{13}NR^{14}R^{15}$, $-P(S)NR^{12}R^{13}NHR^{14}$, $-P(S)NHR^{12}NHR^{14}$, $-P(S)NR^{12}R^{13}NH_2$, $-P(S)NHR^{12}NH_2$, $-P(S)(NH_2)_2$, $-P(S)R^6OR^8$, $-P(S)R^6OH$, $-P(S)R^6SR^{10}$, $-P(S)R^6SH$, $-P(S)R^6NR^{12}R^{13}$, $-P(S)R^6NHR^{12}$, $-P(S)R^6NH_2$, $-P(S)OR^8SR^{10}$, $-P(S)OHSR^{10}$, $-P(S)OR^8SH$, $-P(S)OHSH$, $-P(S)OR^8NR^{12}R^{13}$, $-P(S)OR^8NHR^{12}$, $-P(S)OR^8NH_2$, $-P(S)OHNHR^{12}R^{13}$, $-P(S)OHNHR^{12}$, $-P(S)OHNH_2$, $-P(S)SR^{10}NR^{12}R^{13}$, $-P(S)SR^{10}NHR^{12}$, $-P(S)SR^{10}NH_2$, $-P(S)SHNR^{12}R^{13}$, $-P(S)SHNHR^{12}$, and $-P(S)SHNH_2$, are removed by treatment with acid, such as gaseous hydrogen chloride, aqueous hydrochloric acid, hydrochloric acid/alcohol mixtures, acetic acid/formic acid/water mixture, trifluoroacetic acid, p-toluenesulphonic acid, or other mineral acids. Preferably, gaseous hydrogen chloride is bubbled through a solution of the phosphorous substituted amine, or the phosphorous substituted amine is treated with an hydrochloric acid solution to effect the cleavage of the nitrogen-phosphorous bond.

When R^3 or R^4 is a group represented by CO_2R^{19} treatments with acids or reductive methods may be used to remove the R^3 or R^4 group. When R^{19} is a benzyl group, reductive methods such as hydrogenation in the presence of palladium or charcoal and dissolving metal reductions, or treatment with strong acid, such as hydrogen bromide/acetic acid mixtures, may be employed. When R^{19} is a t-butyl group treatment with an acid such as p-toluenesulphonic or trifluoroacetic acid in an organic solvent such as alcohols, ethers or acetonitrile may be employed. When R^{19} is a methyl group, more forcing conditions may be required such as treatments with hydrazine or alkali metal hydroxides.

When R^3 or R^4 is a group represented by COR^{18} , the R^3 or R^4 group may be removed by acid or base hydrolysis.

When R^3 or R^4 is a group represented by $-S(O)_2R^{17}$, treatments with acids or reductive methods may be used to remove the R^3 or R^4 group. When R^{17} is a methyl group hydride reductions, for example using lithium aluminium hydride, or dissolving metal reductions may be employed. When R^{17} is a p-tolyl group treatment in strong acid, such as trifluoroacetic acid in methanol or 6M hydrochloric acid, may be employed. When R^{17} is a trimethylsilylethyl group treatment with an alkali metal fluoride, preferably caesium fluoride, may be employed.

When R^3 or R^4 is a group represented by $SiR^{20}R^{21}R^{22}$, treatments with acids or alkali metal fluorides may be used to remove the R^3 or R^4 group.

The invention is illustrated by the following Examples.

SYNTHESIS OF OXIME INTERMEDIATES

Oxime Intermediate 1

Synthesis of acetophenone oxime

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
1-acetophenone	1 g	170.2	5.9 mmol	1
hydroxylamine hydrochloride	0.98 g	69.5	10.0 mmol	1.7
pyridine	10 ml	79.1	-	-

In a 50 ml round-bottom flask, a mixture of acetophenone, hydroxylamine hydrochloride in pyridine was stirred, at room temperature under nitrogen blanket, for five days. The reaction mixture was then concentrated under vacuum to remove most of pyridine. The oil was then dissolved in dichloromethane. The product precipitated on standing. The product was obtained as a white solid in 65% yield.

Note : after reaction, the compound exhibit a 75/25 E/Z isomers ratio. After precipitation, one single isomer (E) was observed by 1H NMR.

Oxime Intermediate 2**Synthesis of acetophenone oxime**

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
acetophenone	1 g	120.1	8.3 mmol	1
hydroxylamine hydrochloride	0.695 g	69.5	14.1 mmol	1.7
pyridine	5 ml	79.1	-	-

In a 50 ml round-bottom flask, a mixture of acetophenone, hydroxylamine hydrochloride in pyridine was stirred, at room temperature under nitrogen blanket, for five days. The reaction mixture was then concentrated under vacuum to remove most of pyridine. The oil was then distilled at 55°C under 0.1 mbar. The product was obtained in 95% yield.

Note : the isomer E/Z ratio of the isolated compound was 87/13 as measured by ^1H NMR.

Oxime Intermediate 3**Synthesis of 2-octanone oxime**

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
2-octanone	1.2 ml	128.2	7.8 mmol	1
hydroxylamine hydrochloride	0.92 g	69.5	13.3 mmol	1.7
pyridine	10 ml	79.1	-	-

In a 50 ml round-bottom flask, a mixture of 2-octanone, hydroxylamine hydrochloride in pyridine was stirred, at room temperature under nitrogen blanket, for five days. The reaction mixture was then concentrated under vacuum to remove most of pyridine. The oil was then dissolved in dichloromethane. The product precipitated on standing. The product was obtained as a white solid in 87% yield.

SYNTHESIS OF PHOSPHINYL IMINES FROM OXIME INTERMEDIATES**Phosphinyl Imine Example 1****Synthesis of N-diphenylphosphinyl-1,1-methylnaphthyl imine**

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
acetophenone oxime	1 g	185	5.4 mmol	1
Triethylamine	0.75 ml	101.2	5.4 mmol	1
chlorodiphenylphosphine	0.97 ml	220.6	5.4 mmol	1
diethyl ether	5 ml	74.1	-	-
Dichloromethane	5 + 2 ml	84.9	-	-

13 FEB 2004

23

To a stirred solution of acetophenone oxime and triethylamine in diethyl ether/dichloromethane (1/1:10ml) was added dropwise a solution of chlorodiphenylphosphine in dichloromethane (2 ml) at -45°C . After addition, the temperature (-45°C) was maintained for 1 hour and then was allowed to slowly rise to reach room temperature without removing the ice bath (dry ice-acetone). The reaction mixture was stirred at room temperature overnight. The solution was then filtered to remove the triethylamine hydrochloride precipitate and washed with a mixture of diethyl ether/dichloromethane (1/1). The liquid solution was concentrated to give a brown viscous oil in 94% yield. Recrystallisation in toluene/diethyl ether gave the product as yellow crystals.

Phosphinyl Imine Example 2

Synthesis of N-diphenylphosphinyl-1,1-methylphenyl imine

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
acetophenone oxime	1.06 g	135	7.8 mmol	1
triethylamine	1.1 ml	101.2	7.8 mmol	1
chlorodiphenylphosphine	1.4 ml	220.6	7.8 mmol	1
diethyl ether	5 ml	74.1	-	-
dichloromethane	5 + 2 ml	84.9	-	-

To a stirred solution of acetophenone oxime and triethylamine in diethyl ether/dichloromethane (1/1:10ml) was added dropwise a solution of chlorodiphenylphosphine in dichloromethane (2 ml) at -45°C . After addition, the temperature (-45°C) was maintained for 1 hour and then was allowed to slowly rise to reach room temperature without removing the ice bath (dry ice-acetone). The reaction mixture was stirred at room temperature overnight. The solution was then filtered to remove the triethylamine hydrochloride precipitate and washed with a mixture of diethyl ether/dichloromethane (1/1). The product was obtained as a yellow solid in 80% yield after concentration, and was washed with a mixture of diethyl ether/dichloromethane (1/1).

Phosphinyl Imine Example 3

Synthesis of N-diphenylphosphinyl-1,1-methylhexyl imine

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
2-octanone oxime	0.97 g	143	6.8 mmol	1
triethylamine	0.95 ml	101.2	6.8 mmol	1
chlorodiphenylphosphine	1.2 ml	220.6	6.8 mmol	1
diethyl ether	5 ml	74.1	-	-
dichloromethane	5 + 2 ml	84.9	-	-

To a stirred solution of acetophenone oxime and triethylamine in diethyl ether/dichloromethane (1/1:10ml) was added dropwise a solution of chlorodiphenylphosphine in dichloromethane (2 ml) at -45°C . After addition, the temperature (-45°C) was maintained for 1 hour and then was allowed to slowly rise to reach room temperature without removing the ice bath (dry ice-acetone). The reaction mixture was stirred at room temperature overnight. The solution was then filtered to remove the triethylamine hydrochloride precipitate and washed with a mixture of diethyl ether/dichloromethane (1/1). The product was obtained as a colourless oil in 84% yield after concentration. Recrystallisation in toluene/diethyl ether gave the product as white crystals.

Phosphinyl Imine Example 4

Synthesis of N-diethylphosphinyl-1,1methylnaphthyl imine

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
acetophenone oxime	1 g	185	5.4 mmol	1
triethylamine	0.75 ml	101.2	5.4 mmol	1
chlorodiethylphosphine	0.66 ml	124.5	5.4 mmol	1
diethyl ether	5 ml	74.1	-	-
dichloromethane	5 + 2 ml	84.9	-	-


To a stirred solution of acetophenone oxime and triethylamine in diethyl ether/dichloromethane (1/1:10ml) was added dropwise a solution of chlorodiethylphosphine in dichloromethane (2 ml) at -45°C . After addition, the temperature (-45°C) was maintained for 1 hour and then was allowed to slowly rise to reach room temperature without removing the ice bath (dry ice-acetone). The reaction mixture was stirred at room temperature overnight. The solution was then filtered to remove the triethylamine hydrochloride precipitate and washed with a mixture of diethyl ether/dichloromethane (1/1). The product was obtained as a brown oil in 85% yield after concentration. No further purification was made.

TRANSFER HYDROGENATION OF PHOSPHINYL IMINES

Transfer Hydrogenation Of Phosphinyl Imine - Example 1

Reduction of N-diphenylphosphinyl-1,1-methylnaphthyl imine.

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
[Rh(Cp*)Cl ₂] ₂ **	1.54 mg	617.8	2.5 µmol	0.5
(R,R)-N-Tosyl-1,2-diamino-1,2-diphenylethane	1.83 mg	366	5 µmol	1
Methanol	1.6 ml	32	-	-
Et ₃ N/HCO ₂ H [2:5]	2 ml	101.2/42	24 mmol of HCO ₂ H	1200
	369 mg	369	1mmol	200



Cp* - pentamethylcyclopentadienyl

Notes: ** purchased from The Aldrich Chemical Co.

Prior to the reaction, all solvents were degassed, for example:

10ml of anhydrous methanol was added by syringe to a sealed clean dry round bottomed flask and degassed; either by reducing the pressure until the solvent began to boil and backfilling with nitrogen 3 times, or by bubbling nitrogen through the solution for at least 20 mins.

To a Schlenk flask was added the (R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane, rhodium pentamethylcyclopentadienyl dichloride dimer, N-diphenylphosphinyl-1,1-methylnaphthyl imine and methanol. The flask was stoppered with a 'Suba-seal' (RTM). Its contents were evacuated, then purged at room temperature by 3 changes of nitrogen. To the yellow/orange solution was added the formic acid/triethylamine mixture dropwise. After 30 mins, the solution had changed colour to red/brown. The mixture was stirred for 3 hours while being purged with nitrogen, the reaction was then quenched by the addition of saturated sodium carbonate solution (1 ml). The aqueous solution was then shaken with dichloromethane (5 ml), the organic phase separated and collected. The organic phase was then dried by contacting with solid anhydrous magnesium sulphate and then filtering off the solid, then the solvent was removed *in vacuo* to give product in >95% conversion and >99% ee. The sample was analysed by ¹H and ³¹P NMR.

13 FEB 2004

Transfer Hydrogenation Of Phosphinyl Imine - Example 2**Reduction of N-diphenylphosphinyl-1,1-methylnaphthyl imine with a rhodium catalyst.**

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
N-diphenylphosphinyl-1,1-methylnaphthyl imine	369 mg	369	1 mmol	50
[RhCp*Cl ₂] ₂	6.2 mg	617.8	10 μ mol	0.5
(R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane	7.3 mg	366	20 μ mol	1
acetonitrile	1.6 ml	41	-	-
Et ₃ N/HCO ₂ H [2:5]	2 ml	101.2/42	24 mmol of HCO ₂ H	1200

Cp* = pentamethylcyclopentadienyl

The (R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane and rhodium compound were weighed out into a clean dry Schlenk flask. A solution of N-diphenylphosphinyl-1,1-methylnaphthyl imine in acetonitrile was added. The flask was stoppered with a "Suba-seal" (RTM). The reaction mixture was stirred rapidly and purged at room temperature by 3 changes of nitrogen. After 10 mins, the mixture of triethylamine/formic acid was added dropwise. The orange solution became red/brown. After 3 hours, the reaction mixture was quenched with a saturated solution of Na₂CO₃ (2 ml). Extraction was then carried out by treatment with dichloromethane (5 ml). The organic layer was dried over magnesium sulphate and concentrated to give the product as a brown oil, in > 95% conversion and > 99% ee.

Note: the reaction can be carried out with or without a nitrogen purge. The ee were determined by HPLC on a chiral phase column or by ³¹P NMR with a chiral shift reagent.

Transfer Hydrogenation Of Phosphinyl Imine - Example 3**Reduction of N-diphenylphosphinyl-1,1-methylphenyl imine with a rhodium catalyst.**

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
N-diphenylphosphinyl-1,1-methylphenyl imine	319 mg	319	1 mmol	50
[RhCp*Cl ₂] ₂	6.2 mg	617.8	10 μ mol	0.5
(R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane	7.3 mg	366	20 μ mol	1
acetonitrile	1.6 ml	41	-	-
Et ₃ N/HCO ₂ H [2:5]	2 ml	101.2/42	24 mmol of HCO ₂ H	1200

Cp* = pentamethylcyclopentadienyl

The (R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane and rhodium compound were weighed out into a clean dry Schlenk flask. A solution of N-diphenylphosphinyl-1,1-methylphenyl imine in acetonitrile was added. The flask was stoppered with a "Suba-seal" (RTM). The reaction mixture was stirred rapidly and purged at room temperature by 3 changes of nitrogen. After 10 mins, the mixture of triethylamine/formic acid was added dropwise. The yellow solution became red/brown. After 3 hours, the reaction mixture was quenched with a saturated solution of sodium carbonate (2 ml). Extraction was then carried out by treatment with dichloromethane (5 ml). The organic layer was dried over magnesium sulphate and concentrated to give the product as a brown liquid, in > 95% conversion and 86% ee.

Note: The ee were determined by HPLC on a chiral phase column or by ³¹P NMR with a chiral shift reagent.

Transfer Hydrogenation Of Phosphinyl Imine - Example 4**Reduction of N-diphenylphosphinyl-1,1-methylhexyl imine with an iridium catalyst.**

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
N-diphenylphosphinyl-1,1-methylhexyl imine	327 mg	327	1 mmol	50
[IrCp*Cl ₂] ₂	7.9 mg	796.6	10 μ mol	0.5
(R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane	7.3 mg	366	20 μ mol	1
acetonitrile	1.6 ml	41	-	-
Et ₃ N/HCO ₂ H [2:5]	2 ml	101.2/42	24 mmol of HCO ₂ H	1200

Cp* = pentamethylcyclopentadienyl

1.3, FEB 2004

28

The (R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane and Iridium compound were weighed out into a clean dry Schlenk flask. A solution of N-diphenylphosphinyl-1,1-methylhexyl imine in acetonitrile was added. The flask was stoppered with a "Suba-seal" (RTM). The reaction mixture was stirred rapidly and purged at room temperature by 3 changes of nitrogen. After 10 mins, the mixture of triethylamine/formic acid was added dropwise. The yellow solution became green. After 2 hours, the reaction mixture was quenched with a saturated solution of sodium carbonate (2 ml). Extraction was then carried out by treatment with dichloromethane (5 ml). The organic layer was dried over magnesium sulphate and concentrated to give the product as a brown liquid, in > 95% conversion and 95% ee.

Note: The ee were determined by HPLC on a chiral phase column or by ^{31}P NMR with a chiral shift reagent.

Transfer Hydrogenation Of Phosphinyl Imine - Example 5

Reduction of N-diethylphosphinyl-1,1-methylnaphthyl imine with a ruthenium catalyst.

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
N-diethylphosphinyl-1,1-methylnaphthyl imine	369 mg	369	1 mmol	50
p-cymenerutheniumchloride dimer	6.1 mg	612.4	10 μmol	0.5
(R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane	7.3 mg	366	20 μmol	1
acetonitrile	1.6 ml	41	-	-
$\text{Et}_3\text{N}/\text{HCO}_2\text{H}$ [2:5]	2 ml	101.2/42	24 mmol of HCO_2H	1200

The (R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane and ruthenium compound were weighed out into a clean dry Schlenk flask. A solution of N-diethylphosphinyl-1,1-methylnaphthyl imine in acetonitrile was added. The flask was stoppered with a "Suba-seal" (RTM). The reaction mixture was stirred rapidly and purged at room temperature by 3 changes of nitrogen. After 10 mins, the mixture of triethylamine/formic acid was added dropwise. After 2 hours, the reaction mixture was quenched with a saturated solution of sodium carbonate (2 ml). Extraction was then carried out by treatment with dichloromethane (5 ml). The organic layer was dried over magnesium sulphate and concentrated to give the product as a brown oil, in > 95% conversion and 84% ee.

Note: The ee were determined by HPLC on a chiral phase column or by ^{31}P NMR with a chiral shift reagent.

13 FEB 2004

29

Transfer Hydrogenation Of Phosphinyl Imine - Example 6**Reduction of N-diphenylphosphinyl-1,1-methylnaphthyl imine with a Rhodium catalyst.**

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
N-diphenylphosphinyl-1,1-methylnaphthyl imine	369 mg	369	1 mmol	50
[RhCp*Cl ₂] ₂	6.2 mg	617.8	10 µmol	0.5
(R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane	7.3 mg	366	20 µmol	1
methanol	1.6 ml	32	-	-
Et ₃ N/HCO ₂ H [2:5]	2 ml	101.2/42	24 mmol of HCO ₂ H	1200

Cp* = pentamethylcyclopentadienyl

The (R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane and rhodium compound were weighed out into a clean dry Schlenk flask. The solution of N-diphenylphosphinyl-1,1-methylnaphthyl imine in methanol was added. The flask was stoppered with a "Suba-seal" (RTM). The reaction mixture was stirred rapidly and purged at room temperature by 3 changes of nitrogen. After 10 mins, the mixture of triethylamine/formic acid was added dropwise. The yellow/orange solution became red/brown. After 3 hours, the reaction mixture was quenched with a saturated solution of sodium carbonate (2 ml). Extraction was then carried out by treatment with dichloromethane (5 ml). The organic layer was dried over magnesium sulphate and concentrated to give the product as a brown oil, in > 95% conversion and > 99% ee.

Transfer Hydrogenation Of Phosphinyl Imine - Example 7**Reduction of N-diphenylphosphinyl-1,1-methylnaphthyl imine with a Rhodium catalyst.**

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
N-diphenylphosphinyl-1,1-methylnaphthyl imine	369 mg	369	1 mmol	50
[RhCp*Cl ₂] ₂	6.2 mg	617.8	10 µmol	0.5
(S,S)-N-tosyl-1,2-diamino-1,2-diphenylethane	7.3 mg	366	20 µmol	1
acetonitrile	1.6 ml	41	-	-
Et ₃ N/HCO ₂ H [2:5]	2 ml	101.2/42	24 mmol of HCO ₂ H	1200

Cp* = pentamethylcyclopentadienyl

The (S,S)-N-tosyl-1,2-diamino-1,2-diphenylethane and rhodium compound were weighed out into a clean dry Schlenk flask. The solution of N-diphenylphosphinyl-1,1-methylnaphthyl imine in acetonitrile was added. The flask was stoppered with a "Suba-seal" (RTM). The reaction mixture was stirred rapidly and purged at room temperature by 3 changes of nitrogen. After 10 mins, the mixture of triethylamine/formic acid was added dropwise. The orange solution became red/brown. After 3 hours, the reaction mixture was quenched with a saturated solution of sodium carbonate (2 ml). Extraction was then carried out by treatment with dichloromethane (5 ml). The organic layer was dried over magnesium sulphate and concentrated to give the product as a brown oil, in > 95% conversion and > 99% ee.

Transfer Hydrogenation Of Phosphinyl Imine - Example 8

Reduction of N-diphenylphosphinyl-1,1-methylnaphthyl imine with a Rhodium catalyst.

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
N-diphenylphosphinyl-1,1-methylnaphthyl imine	369 mg	369	1 mmol	200
[RhCp*Cl ₂] ₂	1.5 mg	617.8	2.5 µmol	0.5
(R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane	1.8 mg	366	5 µmol	1
acetonitrile	1.6 ml	41	-	-
Et ₃ N/HCO ₂ H [2:5]	2 ml	101.2/42	24 mmol of HCO ₂ H	4800

Cp* = pentamethylcyclopentadienyl

The (R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane and rhodium compound were weighed out into a clean dry Schlenk flask. The solution of N-diphenylphosphinyl-1,1-methylnaphthyl imine in acetonitrile was added. The flask was stoppered with a "Suba-seal" (RTM). The reaction mixture was stirred rapidly and purged at room temperature by 3 changes of nitrogen. After 10 mins, the mixture of Et₃N/HCO₂H was added dropwise. The orange solution became red/brown. After 3 hours, the reaction mixture was quenched with a saturated solution of sodium carbonate (2 ml). Extraction was then carried out by treatment with dichloromethane (5 ml). The organic layer was dried over magnesium sulphate and concentrated to give the product as a brown oil, in > 95% conversion and > 99% ee.

Transfer Hydrogenation Of Phosphinyl Imine - Example 9

Reduction of N-diphenylphosphinyl-1,1-methylnaphthyl imine with a Rhodium catalyst.

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
N-diphenylphosphinyl-1,1-methylnaphthyl imine	369 mg	369	1 mmol	50
[RhCp*Cl ₂] ₂	6.2 mg	617.8	10 µmol	0.5
(R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane	7.3 mg	366	20 µmol	1
acetonitrile	1.6 ml	41	-	-
HCO ₂ H	1 ml	42	24 mmol of HCO ₂ H	1200

Cp* = pentamethylcyclopentadienyl

The (R,R)-N-tosyl-1,2-diamino-1,2-diphenylethane and rhodium compound were weighed out into a clean dry Schlenk flask. The solution N-diphenylphosphinyl-1,1-methylnaphthyl imine in acetonitrile was added. The flask was stoppered with a "Suba-seal" (RTM). The reaction mixture was stirred rapidly and purged at room temperature by 3 changes of nitrogen. After 10 mins, formic acid (1 ml) was added dropwise. The orange solution became brown. After 2 hours, the reaction mixture was quenched with a saturated solution of sodium carbonate (5 ml). Extraction was then carried out by treatment with dichloromethane (5 ml). The organic layer was dried over magnesium sulphate and concentrated to give the product as a brown oil, in > 95% conversion and > 99% ee.

REMOVAL OF PHOSPHINYL GROUP FROM PHOSPHINYLAMINE**Phosphinyl group removal - Example 1****1-naphthylethylamine**

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
N-diphenylphosphinyl-1-naphthylethylamine	0.2 g	371	0.54 mmol	1
gaseous hydrogen chloride	-	36.5	-	-
ethyl alcohol	10 ml	46	-	-

Gaseous hydrogen chloride was bubbled through a stirred solution of N-diphenylphosphinyl-1-naphthylethylamine in ethyl alcohol for 2 hours at room temperature. The reaction mixture was concentrated, made basic by the addition of an aqueous solution of sodium hydroxide (2 M) and extracted with dichloromethane (3 x 10 ml).

13 FEB 2004

32

The organic layer was washed with brine (1 x 10 ml), dried over magnesium sulphate and concentrated to give the product as a yellow liquid in 80% yield.

Phosphinyl group removal – Example 2

1-phenylethylamine

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
N-diphenylphosphinyl-1-phenylethylamine	0.2 g	321	0.62 mmol	1
gaseous hydrogen chloride	-	36.5	-	-
ethyl alcohol	10 ml	46	-	-

Gaseous hydrogen chloride was bubbled through a stirred solution of N-diphenylphosphinyl-1-phenylethylamine in ethyl alcohol for 2 hours at room temperature. The reaction mixture was concentrated, made basic by the addition of an aqueous solution of sodium hydroxide (2 M) and extracted with dichloromethane (3 x 10 ml). The organic layer was washed with brine (1 x 10 ml), dried over magnesium sulphate and concentrated to give the product as a yellow liquid in 88% yield.

Phosphinyl group removal – Example 3

Synthesis of 1-methylheptylamine.

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
N-diphenylphosphinyl-1-methylheptylamine	0.2 g	329	0.61 mmol	1
gaseous hydrogen chloride	-	36.5	-	-
ethyl alcohol	10 ml	46	-	-

Gaseous hydrogen chloride was bubbled through a stirred solution of N-diphenylphosphinyl-1-methylheptylamine in ethyl alcohol, for 2 hours at room temperature. The reaction mixture was concentrated, made basic by the addition of an aqueous solution of sodium hydroxide (2 M) and extracted with dichloromethane (3 x 10 ml). The organic layer was washed with brine (1 x 10 ml), dried over magnesium sulphate and concentrated to give the product as a yellow liquid in 69% yield.

13 FEB 2004

33


Phosphinyl group removal – Example 4**1-naphthylethylamine**

Reactant	Wt/Vol	Mol.Wt	Mol	Mol ratio
N-diethylphosphinyl-1-naphthylethylamine	0.2 g	275	0.73 mmol	1
gaseous hydrogen chloride	-	36.5	-	-
ethyl alcohol	10 ml	46	-	-

Gaseous hydrogen chloride was bubbled through a stirred solution of N-diethylphosphinyl-1-naphthylethylamine in ethyl alcohol for 2 hours at room temperature. The reaction mixture was concentrated, made basic by the addition of an aqueous solution of sodium hydroxide (2 M) and extracted with dichloromethane (3 x 10 ml). The organic layer was washed with brine (1 x 10 ml), dried over magnesium sulphate and concentrated to give the product as a yellow liquid in 72% yield.

CLAIMS

1. A process for the transfer hydrogenation of a substrate wherein the substrate is reacted with a hydrogen donor in the presence of a transfer hydrogenation catalyst, wherein the substrate has the general formula



wherein:

X represents NR^3 or $(\text{NR}^4\text{R}^5)^+\text{Q}^-$;

Q^- represents a monovalent anion;

R^1 and R^2 each independently represents a hydrogen atom, an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl, an optionally substituted heterocyclyl group, a substituted carbonyl functional group, a substituted thiocarbonyl functional group or substituted imino functional group, R^1 & R^2 optionally being linked in such a way as to form an optionally substituted ring;


R^3 and R^4 represents $-\text{P}(\text{O})\text{R}^6\text{R}^7$, $-\text{P}(\text{O})\text{OR}^8\text{OR}^9$, $-\text{P}(\text{O})\text{OR}^8\text{OH}$, $-\text{P}(\text{O})(\text{OH})_2$, $-\text{P}(\text{O})\text{SR}^{10}\text{SR}^{11}$, $-\text{P}(\text{O})\text{SR}^{10}\text{SH}$, $-\text{P}(\text{O})(\text{SH})_2$, $-\text{P}(\text{O})\text{NR}^{12}\text{R}^{13}\text{NR}^{14}\text{R}^{15}$, $-\text{P}(\text{O})\text{NR}^{12}\text{R}^{13}\text{NHR}^{14}$, $-\text{P}(\text{O})\text{NHR}^{12}\text{NHR}^{14}$, $-\text{P}(\text{O})\text{NR}^{12}\text{R}^{13}\text{NH}_2$, $-\text{P}(\text{O})\text{NHR}^{12}\text{NH}_2$, $-\text{P}(\text{O})(\text{NH}_2)_2$, $-\text{P}(\text{O})\text{R}^6\text{OR}^8$, $-\text{P}(\text{O})\text{R}^6\text{OH}$, $-\text{P}(\text{O})\text{R}^6\text{SR}^{10}$, $-\text{P}(\text{O})\text{R}^6\text{SH}$, $-\text{P}(\text{O})\text{R}^6\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{R}^6\text{NHR}^{12}$, $-\text{P}(\text{O})\text{R}^6\text{NH}_2$, $-\text{P}(\text{O})\text{OR}^8\text{SR}^{10}$, $-\text{P}(\text{O})\text{OR}^8\text{SH}$, $-\text{P}(\text{O})\text{OHSR}^{10}$, $-\text{P}(\text{O})\text{OHSH}$, $-\text{P}(\text{O})\text{OR}^8\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{OR}^8\text{NHR}^{12}$, $-\text{P}(\text{O})\text{OR}^8\text{NH}_2$, $-\text{P}(\text{O})\text{OHNHR}^{12}$, $-\text{P}(\text{O})\text{OHNH}_2$, $-\text{P}(\text{O})\text{SR}^{10}\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{SR}^{10}\text{NHR}^{12}$, $-\text{P}(\text{O})\text{SR}^{10}\text{NH}_2$, $-\text{P}(\text{O})\text{SHNR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{SHNHR}^{12}$, $-\text{P}(\text{O})\text{SHNH}_2$, $-\text{P}(\text{S})\text{R}^6\text{R}^7$, $-\text{P}(\text{S})\text{OR}^8\text{OR}^9$, $-\text{P}(\text{S})\text{OR}^8\text{OH}$, $-\text{P}(\text{S})(\text{OH})_2$, $-\text{P}(\text{S})\text{SR}^{10}\text{SR}^{11}$, $-\text{P}(\text{S})\text{SR}^{10}\text{SH}$, $-\text{P}(\text{S})(\text{SH})_2$, $-\text{P}(\text{S})\text{NR}^{12}\text{R}^{13}\text{NR}^{14}\text{R}^{15}$, $-\text{P}(\text{S})\text{NR}^{12}\text{R}^{13}\text{NHR}^{14}$, $-\text{P}(\text{S})\text{NHR}^{12}\text{NHR}^{14}$, $-\text{P}(\text{S})\text{NR}^{12}\text{R}^{13}\text{NH}_2$, $-\text{P}(\text{S})\text{NHR}^{12}\text{NH}_2$, $-\text{P}(\text{S})(\text{NH}_2)_2$, $-\text{P}(\text{S})\text{R}^6\text{OR}^8$, $-\text{P}(\text{S})\text{R}^6\text{OH}$, $-\text{P}(\text{S})\text{R}^6\text{SR}^{10}$, $-\text{P}(\text{S})\text{R}^6\text{SH}$, $-\text{P}(\text{S})\text{R}^6\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{S})\text{R}^6\text{NHR}^{12}$, $-\text{P}(\text{S})\text{R}^6\text{NH}_2$, $-\text{P}(\text{S})\text{OR}^8\text{SR}^{10}$, $-\text{P}(\text{S})\text{OHSR}^{10}$, $-\text{P}(\text{S})\text{OR}^8\text{SH}$, $-\text{P}(\text{S})\text{OR}^8\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{S})\text{OR}^8\text{NHR}^{12}$, $-\text{P}(\text{S})\text{OR}^8\text{NH}_2$, $-\text{P}(\text{S})\text{OHNHR}^{12}$, $-\text{P}(\text{S})\text{OHNH}_2$, $-\text{P}(\text{S})\text{SR}^{10}\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{S})\text{SR}^{10}\text{NHR}^{12}$, $-\text{P}(\text{S})\text{SR}^{10}\text{NH}_2$, $-\text{P}(\text{S})\text{SHNR}^{12}\text{R}^{13}$, $-\text{P}(\text{S})\text{SHNHR}^{12}$, $-\text{P}(\text{S})\text{SHNH}_2$, $-\text{PR}^6\text{R}^7$, $-\text{POR}^8\text{OR}^9$, $-\text{PSR}^{10}\text{SR}^{11}$, $-\text{PNR}^{12}\text{R}^{13}\text{NR}^{14}\text{R}^{15}$, $-\text{PR}^6\text{OR}^8$, $-\text{PR}^6\text{SR}^{10}$, $-\text{PR}^6\text{NR}^{12}\text{R}^{13}$, $-\text{POR}^8\text{SR}^{10}$, $-\text{POR}^8\text{NR}^{12}\text{R}^{13}$, $-\text{PSR}^{10}\text{NR}^{12}\text{R}^{13}$, $-\text{S}(\text{O})\text{R}^{16}$, $-\text{S}(\text{O})_2\text{R}^{17}$, $-\text{COR}^{18}$, $-\text{CO}_2\text{R}^{19}$, or $\text{SiR}^{20}\text{R}^{21}\text{R}^{22}$.

R^5 represents a hydrogen atom, an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl, or an optionally substituted heterocyclyl group;

R^6 and R^7 independently represent an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl, an optionally substituted heterocyclyl group or $-\text{N}=\text{CR}^{23}\text{R}^{24}$ where R^{23} and R^{24} are as defined for R^1 ; and

R^8 to R^{22} each independently represents an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, one or more of R^1 & R^6 , R^2 & R^7 , R^3 & R^7 , R^5 & R^8 , R^5 & R^{10} , R^6 & R^{12} , R^1 & R^8 , R^2 & R^9 , R^8 & R^9 , R^9 & R^{10} , R^9 & R^{12} , R^1 & R^{10} , R^2 & R^{11} , R^{10} & R^{11} , R^{10} & R^{12} , R^1 & R^{12} , R^2 & R^{13} ,

R^{12} & R^{13} , R^1 & R^{14} , R^2 & R^{15} , R^{14} & R^{15} , R^{12} & R^{14} , R^1 & R^{16} , R^1 & R^{18} , R^1 & R^{20} , R^2 & R^{21} , R^{20} & R^{21} and R^{21} & R^{22} optionally being linked in such a way as to form an optionally substituted ring(s),
and wherein the transfer hydrogenation catalyst has the general formula:



wherein:

R^{27} represents a neutral optionally substituted hydrocarbyl, a neutral optionally substituted perhalogenated hydrocarbyl, or an optionally substituted cyclopentadienyl ligand;

A represents $-NR^{28}-$, $-NR^{29}-$, $-NHR^{28}$, $-NR^{29}R^{30}$ or $-NR^{29}R^{30}$ where R^{28} is H, $C(O)R^{30}$, SO_2R^{30} , $C(O)NR^{30}R^{34}$, $C(S)NR^{30}R^{34}$, $C(=NR^{34})SR^{35}$ or $C(=NR^{34})OR^{35}$, R^{29} and R^{30} each independently represents an optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, and R^{34} and R^{35} are each independently hydrogen or a group as defined for R^{30} ;

B represents $-O-$, $-OH$, OR^{31} , $-S-$, $-SH$, SR^{31} , $-NR^{31}-$, $-NR^{32}-$, $-NHR^{32}$, $-NR^{31}R^{32}$, $-NR^{31}R^{33}$, $-PR^{31}-$ or $-PR^{31}R^{33}$ where R^{32} is H, $C(O)R^{33}$, SO_2R^{33} , $C(O)NR^{33}R^{36}$, $C(S)NR^{33}R^{36}$, $C(=NR^{36})SR^{37}$ or $C(=NR^{36})OR^{37}$, R^{31} and R^{33} each independently represents an optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, and R^{36} and R^{37} are each independently hydrogen or a group as defined for R^{33} ;


E represents a linking group wherein A and B are linked through 2, 3 or 4 atoms each independently being unsubstituted or substituted;

M is a group VIII transition metal; and

Y represents an anionic group, a basic ligand or a vacant site;


provided that when Y is not a vacant site that at least one of A or B carries a hydrogen atom.

2. A process for the production of primary or secondary amines



comprising the steps of


- a) generating a substrate of general formula (1) from a carbonyl compound (2),



in which X represents NR^3 or $(NR^4R^5)^+Q^-$, and Q^- represents a monovalent anion,

- b) reacting the substrate of general formula (1) with a hydrogen donor in the presence of a transfer hydrogenation catalyst, and

c) removal of R^3 or R^4 to give an amine of formula (3)



wherein:

R^1 and R^2 each independently represents a hydrogen atom, an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl, an optionally substituted heterocyclyl group, a substituted carbonyl functional group, a substituted thiocarbonyl functional group or substituted imino functional group, R^1 & R^2 optionally being linked in such a way as to form an optionally substituted ring;

R^3 and R^4 represents $-P(O)R^5R^7$, $-P(O)OR^5OR^8$, $-P(O)OR^5OH$, $-P(O)(OH)_2$, $-P(O)SR^{10}SR^{11}$, $-P(O)SR^{10}SH$, $-P(O)(SH)_2$, $-P(O)NR^{12}R^{13}NR^{14}R^{15}$, $-P(O)NR^{12}R^{13}NHR^{14}$, $-P(O)NHR^{12}NHR^{14}$, $-P(O)NR^{12}R^{13}NH_2$, $-P(O)NHR^{12}NH_2$, $-P(O)(NH_2)_2$, $-P(O)R^5OR^8$, $-P(O)R^5OH$, $-P(O)R^5SR^{10}$, $-P(O)R^5SH$, $-P(O)R^5NR^{12}R^{13}$, $-P(O)R^5NHR^{12}$, $-P(O)R^5NH_2$, $-P(O)OR^5SR^{10}$, $-P(O)OR^5SH$, $-P(O)OHSR^{10}$, $-P(O)OHSH$, $-P(O)OR^5NR^{12}R^{13}$, $-P(O)OR^5NHR^{12}$, $-P(O)OR^5NH_2$, $-P(O)OHNHR^{12}R^{13}$, $-P(O)OHNHR^{12}$, $-P(O)OHNH_2$, $-P(O)SR^{10}NR^{12}R^{13}$, $-P(O)SR^{10}NHR^{12}$, $-P(O)SR^{10}NH_2$, $-P(O)SHNHR^{12}R^{13}$, $-P(O)SHNHR^{12}$, $-P(O)SHNH_2$, $-P(S)R^5R^7$, $-P(S)OR^5OR^8$, $-P(S)OR^5OH$, $-P(S)(OH)_2$, $-P(S)SR^{10}SR^{11}$, $-P(S)SR^{10}SH$, $-P(S)(SH)_2$, $-P(S)NR^{12}R^{13}NR^{14}R^{15}$, $-P(S)NR^{12}R^{13}NHR^{14}$, $-P(S)NHR^{12}NHR^{14}$, $-P(S)NR^{12}R^{13}NH_2$, $-P(S)NHR^{12}NH_2$, $-P(S)(NH_2)_2$, $-P(S)R^5OR^8$, $-P(S)R^5OH$, $-P(S)R^5SR^{10}$, $-P(S)R^5SH$, $-P(S)R^5NR^{12}R^{13}$, $-P(S)R^5NHR^{12}$, $-P(S)R^5NH_2$, $-P(S)OR^5SR^{10}$, $-P(S)OHSR^{10}$, $-P(S)OR^5SH$, $-P(S)OHSH$, $-P(S)OR^5NR^{12}R^{13}$, $-P(S)OR^5NHR^{12}$, $-P(S)OR^5NH_2$, $-P(S)OHNHR^{12}R^{13}$, $-P(S)OHNHR^{12}$, $-P(S)OHNH_2$, $-P(S)SR^{10}NR^{12}R^{13}$, $-P(S)SR^{10}NHR^{12}$, $-P(S)SR^{10}NH_2$, $-P(S)SHNHR^{12}R^{13}$, $-P(S)SHNHR^{12}$, $-P(S)SHNH_2$, $-PR^5R^7$, $-POR^5OR^8$, $-PSR^{10}SR^{11}$, $-PNR^{12}R^{13}NR^{14}R^{15}$, $-PR^5OR^8$, $-PR^5SR^{10}$, $-PR^5NR^{12}R^{13}$, $-POR^5SR^{10}$, $-POR^5NR^{12}R^{13}$, $-PSR^{10}NR^{12}R^{13}$, $-S(O)R^{16}$, $-S(O)_2R^{17}$, $-COR^{18}$, $-CO_2R^{19}$, or $SiR^{20}R^{21}R^{22}$;

R^5 represents a hydrogen atom, an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl, or an optionally substituted heterocyclyl group;


R^{3n} represents a hydrogen atom or the group R^5 ;

R^6 and R^7 independently represent an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl, an optionally substituted heterocyclyl group or $-N=CR^{23}R^{24}$ where R^{23} and R^{24} are as defined for R^1 ; and

R^8 to R^{22} each independently represents an optionally substituted hydrocarbyl, a perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, one or more of R^1 & R^6 , R^2 & R^7 , R^3 & R^7 , R^6 & R^8 , R^9 & R^{10} , R^6 & R^{12} , R^1 & R^9 , R^2 & R^9 , R^8 & R^{10} , R^9 & R^{12} , R^1 & R^{10} , R^2 & R^{11} , R^{10} & R^{11} , R^{10} & R^{12} , R^1 & R^{12} , R^2 & R^{13} , R^{12} & R^{13} , R^1 & R^{14} , R^2 & R^{15} , R^{14} & R^{15} , R^{12} & R^{14} , R^1 & R^{16} , R^1 & R^{18} , R^1 & R^{19} , R^1 & R^{20} , R^2 & R^{21} , R^{20} & R^{21} and R^{21} & R^{22} optionally being linked in such a way as to form an optionally substituted ring(s),

and the transfer hydrogenation catalyst is of general formula:

13 FEB 2004



wherein:

R^{27} represents a neutral optionally substituted hydrocarbyl, a neutral optionally substituted perhalogenated hydrocarbyl, or an optionally substituted cyclopentadienyl ligand;

A represents $-NR^{28}$, $-NR^{29}$, $-NHR^{28}$, $-NR^{28}R^{29}$ or $-NR^{29}R^{30}$ where R^{28} is H, $C(O)R^{30}$, SO_2R^{30} , $C(O)NR^{30}R^{34}$, $C(S)NR^{30}R^{34}$, $C(=NR^{34})SR^{35}$ or $C(=NR^{34})OR^{35}$, R^{29} and R^{30} each independently represents an optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, and R^{34} and R^{35} are each independently hydrogen or a group as defined for R^{30} ;

B represents $-O-$, $-OH$, OR^{31} , $-S-$, $-SH$, SR^{31} , $-NR^{31}$, $-NR^{32}$, $-NHR^{32}$, $-NR^{31}R^{32}$, $-NR^{31}R^{33}$, $-PR^{31}$ or $-PR^{31}R^{33}$ where R^{32} is H, $C(O)R^{33}$, SO_2R^{33} , $C(O)NR^{33}R^{36}$, $C(S)NR^{33}R^{36}$, $C(=NR^{36})SR^{37}$ or $C(=NR^{36})OR^{37}$, R^{31} and R^{33} each independently represents an optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, and R^{36} and R^{37} are each independently hydrogen or a group as defined for R^{33} ;

E represents a linking group wherein A and B are linked through 2, 3 or 4 atoms each independently being unsubstituted or substituted;

M is a group VIII transition metal; and

Y represents an anionic group, a basic ligand or a vacant site;

provided that when Y is not a vacant site that at least one of A or B carries a hydrogen atom.

3. A process according to Claim 1 or Claim 2 wherein M is ruthenium, rhodium or iridium.

4. A process according to any one of Claims 1 to 3 wherein R^{27} is an optionally substituted aryl; an optionally substituted alkene; or a cyclopentadienyl group substituted with between 3 and 5 substituents.

5. A process according to Claim 4 wherein R^{27} is a cyclopentadienyl group substituted with 5 substituents.

6. A process according to Claim 5 wherein R^{27} is a pentamethylcyclopentadienyl group

7. A process according to any one of Claims 1 to 6 wherein B represents $-O-$, $-OH$, OR^{31} , $-NR^{31}$, $-NR^{32}$, $-NHR^{32}$, $-NR^{31}R^{32}$, or $-NR^{31}R^{33}$ where R^{32} is H, $C(O)R^{33}$, SO_2R^{33} , $C(O)NR^{33}R^{36}$, $C(S)NR^{33}R^{36}$, $C(=NR^{36})SR^{37}$ or $C(=NR^{36})OR^{37}$, R^{31} and R^{33} each independently represents an optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or an


13 FEB 2004

optionally substituted heterocyclyl group, and R^{36} and R^{37} are each independently hydrogen or a group as defined for R^{33} .

8. A process according to Claim 7 wherein either of A or B carries an acyl or sulphonyl group.

9. A process according to Claim 8 wherein the acyl group is an acetyl group or the sulphonyl group is a toluenesulphonyl, methanesulphonyl, or trifluoromethanesulphonyl group.

10. A process according to Claim 7 in which A-E-B is one of the following:




wherein:

A represents NR^8 or NHR^8 where R^8 is hydrogen, and

B represents O or OH.

11. A process according to Claim 7 wherein A and B are linked through 2 optionally substituted carbon atoms and B represents $-NR^{31}-$, $-NR^{32}-$, $-NHR^{32}$, $-NR^{31}R^{32}$, or $-NR^{31}R^{33}$ where R^{32} is H, $C(O)R^{33}$, SO_2R^{33} , $C(O)NR^{33}R^{36}$, $C(S)NR^{33}R^{36}$, $C(=NR^{36})SR^{37}$ or $C(=NR^{36})OR^{37}$, R^{31} and R^{33} each independently represents an optionally substituted hydrocarbyl, perhalogenated hydrocarbyl or an optionally substituted heterocyclyl group, and R^{36} and R^{37} are each independently hydrogen or a group as defined for R^{33} .

12. A process according to Claim 7 in which A-E-B is one of the following:




wherein:

A represents NR^8 or NHR^8 , where R^8 is hydrogen, and

B represents NR^{12} or NHR^{12} , where R^{12} is hydrogen, tosyl or $-SO_2$ naphthyl;

or




wherein:

13 FEB 2004

A represents NR^8 or NHR^8 , where R^8 is hydrogen, and
 B represents NR^{12} or NHR^{12} , R^{12} is hydrogen or tosyl;


or



wherein:

A represents NR^8 or NHR^8 , where R^8 is hydrogen, and
 B represents NR^{12} or NHR^{12} , R^{12} is tosyl;

or



wherein:

A represents NR^8 or NHR^8 , where R^8 is hydrogen, and
 B represents NR^{12} or NHR^{12} , R^{12} is hydrogen.

13. A process according to any one of the preceding Claims wherein R^3 or R^4 is a group selected from $-\text{P}(\text{O})\text{R}^6\text{R}^7$, $-\text{P}(\text{O})\text{OR}^8\text{OR}^9$, $-\text{P}(\text{O})\text{SR}^{10}\text{SR}^{11}$, $-\text{P}(\text{O})\text{NR}^{12}\text{R}^{13}\text{NR}^{14}\text{R}^{15}$, $-\text{P}(\text{O})\text{R}^6\text{OR}^8$, $-\text{P}(\text{O})\text{R}^6\text{SR}^{10}$, $-\text{P}(\text{O})\text{R}^6\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{OR}^8\text{SR}^{10}$, $-\text{P}(\text{O})\text{OR}^8\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{O})\text{SR}^{10}\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{S})\text{R}^6\text{R}^7$, $-\text{P}(\text{S})\text{OR}^8\text{OR}^9$, $-\text{P}(\text{S})\text{SR}^{10}\text{SR}^{11}$, $-\text{P}(\text{S})\text{NR}^{12}\text{R}^{13}\text{NR}^{14}\text{R}^{15}$, $-\text{P}(\text{S})\text{R}^6\text{OR}^8$, $-\text{P}(\text{S})\text{R}^6\text{SR}^{10}$, $-\text{P}(\text{S})\text{R}^6\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{S})\text{OR}^8\text{SR}^{10}$, $-\text{P}(\text{S})\text{OR}^8\text{NR}^{12}\text{R}^{13}$, $-\text{P}(\text{S})\text{SR}^{10}\text{NR}^{12}\text{R}^{13}$, $-\text{PR}^6\text{R}^7$, $-\text{POR}^8\text{OR}^9$, $-\text{PSR}^{10}\text{SR}^{11}$, $-\text{PNR}^{12}\text{R}^{13}\text{NR}^{14}\text{R}^{15}$, $-\text{PR}^6\text{OR}^8$, $-\text{PR}^6\text{SR}^{10}$, $-\text{PR}^6\text{NR}^{12}\text{R}^{13}$, $-\text{POR}^8\text{SR}^{10}$, $-\text{POR}^8\text{NR}^{12}\text{R}^{13}$, $-\text{PSR}^{10}\text{NR}^{12}\text{R}^{13}$, $-\text{S}(\text{O})\text{R}^{16}$, $-\text{S}(\text{O})_2\text{R}^{17}$, $-\text{COR}^{18}$, $-\text{CO}_2\text{R}^{19}$, and $\text{SiR}^{20}\text{R}^{21}\text{R}^{22}$.

14. A process according to Claim 13 wherein R^3 or R^4 is $-\text{P}(\text{O})\text{R}^6\text{R}^7$ or $-\text{P}(\text{O})\text{OR}^8\text{OR}^9$.

15. A process according to any of Claims 1 to 14 wherein the groups represented by R^6 to R^{19} are independently selected from alkyl or aryl groups.

16. A process according to any of Claims 1 to 15 wherein X is NR^3 where R^3 is either $-\text{P}(\text{O})\text{R}^6\text{R}^7$ or $-\text{P}(\text{O})\text{OR}^8\text{OR}^9$ and wherein R^6 to R^8 are independently selected from C_{1-4} alkyl groups, phenyl groups or phenyl groups substituted with one or more C_{1-4} alkyl groups.

17. A process according to any one of the preceding claims wherein the compound of formula (1) is prochiral, the transfer hydrogenation catalyst is chiral, an enantiomerically and/or diastereochemically purified form of catalyst being employed, whereby the compound of formula (1) is asymmetrically hydrogenated.

18. A process according to any one of the preceding claims wherein the hydrogen donor is selected from hydrogen, primary and secondary alcohols, primary and secondary

13 FEB 2004

amines, carboxylic acids and their esters and amine salts, readily dehydrogenatable hydrocarbons, clean reducing agents, and any combination thereof.

19. A process according to Claim 18, in which the hydrogen donor is selected from propan-2-ol, butan-2-ol, triethylammonium formate and a mixture of triethylammonium formate and formic acid.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 00/02867

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07B43/04 C07B53/00 C07C209/62

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07B C07C C07F

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

BEILSTEIN Data, EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	T. EGUCHI: "Rational de novo design of NADH mimic for stereoselective reduction based on molecular orbital calculation" TETRAHEDRON, (INCL TETRAHEDRON REPORTS), vol. 54, no. 5-6, 1998, pages 705-714, XP004106597 OXFORD GB page 713, paragraph 2	1,9,11, 14
A	EP 0 901 996 A (MITSUI CHEMICALS) 17 March 1999 (1999-03-17) claims; examples	1,2
A	WO 98 42643 A (ZENECA) 1 October 1998 (1998-10-01) claims; examples 5-7	1,2
	-/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

20 September 2000

Date of mailing of the international search report

29/09/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Wright, M

INTERNATIONAL SEARCH REPORT

Inte .onal Application No

PCT/GB 00/02867

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	N. UEMATSU: "Asymmetric transfer hydrogenation of imines" JOURNAL OF THE AMERICAN CHEMICAL SOCIETY, vol. 118, no. 20, 22 May 1996 (1996-05-22), pages 4916-4917, XP002069432 DC US the whole document	1,2

INTERNATIONAL SEARCH REPORT

information on patent family members

International Application No

PCT/GB 00/02867

Patent document cited in search report		Publication date	Patent family member(s)		Publication date
EP 901996	A	17-03-1999	WO	9839276 A	11-09-1998
WO 9842643	A	01-10-1998	AU	6739198 A	20-10-1998
			CN	1257468 T	21-06-2000
			CZ	9903374 A	15-12-1999
			EP	0971865 A	19-01-2000
			NO	994677 A	24-11-1999

