




Abstract


Oximeester compounds of the formulae I, II, III and IV



(I)



(III)



R_1 is phenyl, C_1 - C_{20} alkyl or C_2 - C_{20} alkyl optionally interrupted by -O-, C_2 - C_{20} alkanoyl or benzoyl, or R_1 is C_2 - C_{12} alkoxycarbonyl or phenoxy carbonyl; R_1' is C_2 - C_{12} alkoxycarbonyl, or R_1' is phenoxy carbonyl, or R_1' is -CONR₁₀R₁₁ or CN; R_2 is C_2 - C_{12} alkanoyl, C_4 - C_6 alkenoyl, benzoyl, C_2 - C_6 alkoxycarbonyl or phenoxy carbonyl; R_3 , R_4 , R_5 , R_6 and R_7 are hydrogen, halogen, C_1 - C_{12} alkyl, cyclopentyl, cyclohexyl, phenyl, benzyl, benzoyl, C_2 - C_{12} alkanoyl, C_2 - C_{12} alkoxycarbonyl, phenoxy carbonyl or a group OR₈, SR₉, SOR₉, SO₂R₉ or NR₁₀R₁₁; R_4' , R_5' and R_6' are hydrogen, halogen, C_1 - C_{12} alkyl, cyclopentyl, cyclohexyl, phenyl, benzyl, benzoyl, C_2 - C_{12} alkanoyl, C_2 - C_{12} alkoxycarbonyl, phenoxy carbonyl, or are a group OR₈, SR₉, SOR₉, SO₂R₉, NR₁₀R₁₁; provided that at least one of R_3 , R_4 , R_5 , R_6 , R_7 , R_4' , R_5' and R_6' is OR₈, SR₉ or NR₁₀R₁₁; R_8 , R_9 , R_{10} and R_{11} are for example hydrogen, C_1 - C_{12} alkyl, phenyl; are suitable as initiators for the photopolymerization of radically polymerizable compounds.

This Sheet Has Been Replaced



9902594-2