

FLUOROPHOSPHITE CONTAINING CATALYSTS FOR HYDROFORMYLATION PROCESSES

FIELD OF THE INVENTION

[0001] This invention generally relates to fluorophosphite compounds, a catalyst solution containing the same, and a hydroformylation process using the catalyst solution.

BACKGROUND OF THE INVENTION

[0002] The hydroformylation reaction, also known as the oxo reaction, is used extensively in commercial processes for the preparation of aldehydes by the reaction of one mole of an olefin with one mole each of hydrogen and carbon monoxide. One use of the reaction is in the preparation of normal- and iso-butyraldehyde from propylene. The ratio of the amount of the normal aldehyde product to the amount of the iso aldehyde product typically is referred to as the normal-to-iso (N:I) or the normal-to-branched (N:B) ratio. In the case of propylene, the normal- and iso-butyraldehydes obtained from propylene are in turn converted into many commercially valuable chemical products such as, for example, n-butanol, 2-ethyl-hexanol, n-butyric acid, iso-butanol, neo-pentyl glycol, 2,2,4-trimethyl-1,3-pentanediol, and the mono-isobutyrate and di-isobutyrate esters of 2,2,4-trimethyl-1,3-pentanediol.

[0003] In most cases, a phosphorus ligand-containing catalyst is used for the oxo process (so called "low pressure hydroformylation process"). Phosphorus ligands not only can stabilize the metal, but can also regulate the catalyst activity and selectivity. Oxo catalyst activity often decreases as the amount of phosphorus ligands increases while the catalyst stability increases with increasing amounts of ligand. Therefore, there exists an optimum concentration of phosphorus ligands for operating an oxo reactor which is the result of a tradeoff between increased stability and reduced catalyst activity.


[0004] In reality, the gradual loss of phosphorus ligands is inevitable because of decomposition and other reasons. In some cases, such as overflow reactors, the decomposition of ligands may be worsened due to the high temperature required to separate catalysts from products. In practice, fresh phosphorus ligands have to be replenished to the reactor on a regular basis to compensate for the loss of ligands.

[0005] Thus, there is a need in the art for ligands that not only stabilize the catalyst at higher concentrations, but can also increase the metal catalyst activity at such concentrations.

SUMMARY OF THE INVENTION

[0006] A class of ligands has been discovered that can substantially increase the Rh catalyst activity by simply increasing the novel fluorophosphite ligand concentration. In some embodiments, the ligand offers one or more benefits, such as enhanced catalyst stability, increased production rate, and reduced daily operation costs due to eliminated needs for replenishing ligands.

[0007] In one aspect, the present invention provides a fluorophosphite compound having the structure of formula (I):




wherein

Ar_1 and Ar_2 are each independently selected from aryl groups containing 4 to 30 carbon atoms;

R1 to R6 are each independently selected from H and hydrocarbyl containing 1 to 40 carbon atoms; and

X is

- (i) a chemical bond directly between ring carbon atoms of each aromatic group, except when R1 to R4 are each methyl,
- (ii) a heteroatom selected from O, Si, and N, or
- (iii) a group having the formula




wherein R13 and R14 are each independently selected from hydrogen and alkyl or aryl groups having 1 to 10 carbon atoms.

[0008] In a second aspect, the present invention provides a catalyst solution comprising:

- (a) a fluorophosphite compound;
- (b) a Group VIII metal or rhenium; and
- (c) a hydroformylation solvent,

wherein the fluorophosphite compound has the structure of formula (I):




wherein

Ar₁ and Ar₂ are each independently selected from aryl groups containing 4 to 30 carbon atoms;

R1 to R6 are each independently selected from H and hydrocarbyl containing 1 to 40 carbon atoms; and

X is

- (i) a chemical bond directly between ring carbon atoms of each aromatic group;
- (ii) a heteroatom, or
- (iii) a group having the formula



wherein R13 and R14 are each independently selected from hydrogen and alkyl or aryl having 1 to 10 carbon atoms. In some embodiments, R13 and R14 are each independently selected from hydrogen and alkyl groups having one or two carbon atoms. In some embodiments, R13 and R14 are each independently selected from hydrogen and methyl groups.

[0009] In a third aspect, the present invention provides a process for preparing aldehydes, comprising contacting an olefin with hydrogen and carbon monoxide, under hydroformylation conditions, in the presence of the catalyst solution herein described.

DETAILED DESCRIPTION OF THE INVENTION


[0010] New fluorophosphite compounds have been discovered that are useful as ligands in hydroformylation reactions. The invention thus provides new fluorophosphite compounds. The invention further provides highly active catalyst solution for use in hydroformylation reactions.

[0011] The catalyst solution comprises:

- (a) a fluorophosphite compound;

- (b) a Group VIII metal or rhenium; and
- (c) a hydroformylation solvent.

[0012] The fluorophosphite compound according to the invention has the structure of formula (I):




wherein

Ar₁ and Ar₂ are each independently selected from aryl groups containing 4 to 30 carbon atoms;

R1 to R6 are each independently selected from H and hydrocarbyl containing 1 to 40 carbon atoms; and

X is

- (i) a chemical bond directly between ring carbon atoms of each aromatic group,
- (ii) a heteroatom, or
- (iii) a group having the formula






wherein R13 and R14 are each independently selected from hydrogen and alkyl or aryl having 1 to 10 carbon atoms.

[0013] In some embodiments, X is not a chemical bond when R1 to R4 are all methyl.

[0014] The heteroatom can be any atom other than carbon that can form the bonds with the two aryl rings as shown in Formula (I) without compromising the efficacy of the ligand. In some embodiments, the heteroatom is selected from sulfur, oxygen, nitrogen, and silicon, provided that the silicon and nitrogen will have additional substituents bonded thereto to complete the atoms bonding capability. Substituents which are acceptable include alkyl groups of 1 to 20 carbon atoms and aromatic groups of 6 to 20 carbon atoms. In some embodiments, the heteroatom is selected from O, Si, and N, again, with additional atoms bonded to a Si or N.

[0015] Examples of the aryl groups that Ar₁ and Ar₂ can represent include carbocyclic aryl such as phenyl, naphthyl, anthracenyl, and substituted derivatives thereof. Suitable substituents include alkyl groups of 1 to 20 carbon atoms, aromatic groups of 6 to 20 carbon atoms, alkoxy, aryloxy, halogens, carboxylic acids and derivatives thereof such as esters, amides or the carboxylate salt. Additionally suitable substituents include sulfonic acids, the salts of sulfonic acids and derivatives thereof such as esters and amides. In particular, Ar₁ and Ar₂ can each individually represent radicals having any one of the formulas (II) – (IV):






wherein R7 and R8 may represent one or more substituents independently selected from alkyl, alkoxy, halogen, cycloalkoxy, formyl, alkanoyl, cycloalkyl, aryl, aryloxy, aroyl, carboxyl, carboxylate salts, alkoxy-carbonyl, alkanoyloxy, cyano, sulfonic acid, sulfonate salts, and the like. In some embodiments the alkyl moiety of the aforesaid alkyl, alkoxy, alkanoyl, alkoxy-carbonyl and alkanoyloxy groups contains up to 8 carbon atoms. Although it is possible for m to represent 0 to 5 and for n to represent 0 to 7, in some embodiments, the value of each of m and n will not exceed 2. In some embodiments, R7 and R8 represent lower alkyl groups, i.e., straight-chain and branched-chain alkyl of up to 10 carbon atoms, and m and n each represent 0, 1, or 2. In some embodiments, R7 and R8 represent lower alkyl groups, i.e., straight-chain and branched-chain alkyl of up to 4 carbon atoms, and m and n each represent 0, 1, or 2.

[0016] The hydrocarbyl groups represented by R1 to R4 are selected from unsubstituted and substituted alkyl, cycloalkyl and aryl groups containing up to 40 carbon atoms. In some embodiments, the total carbon content of R1 to

R4 is in the range of 1 to 15 carbon atoms. Examples of the alkyl groups that R1 to R4 can represent include methyl, ethyl, butyl, pentyl, hexyl, 2-ethylhexyl, octyl, decyl, dodecyl, octadecyl, and various isomers thereof. The alkyl groups may be substituted, for example, with up to two substituents such as alkoxy, cycloalkoxy, formyl, alkanoyl, cycloalkyl, aryl, aryloxy, aroyl, carboxyl, carboxylate salts, alkoxycarbonyl, alkanoyloxy, cyano, sulfonic acid, sulfonate salts, and the like. Cyclopentyl, cyclohexyl, and cycloheptyl are examples of the cycloalkyl groups that R1 to R4 can represent. The cycloalkyl groups may be substituted with alkyl or any of the substituents described with respect to the possible substituted alkyl groups. In some embodiments, the alkyl and cycloalkyl groups that R1 to R4 can represent are alkyl of up to 8 carbon atoms, benzyl, cyclopentyl, cyclohexyl, or cycloheptyl.

[0017] Examples of the aryl groups that R1 to R4 can represent include carbocyclic aryl such as phenyl, naphthyl, anthracenyl, and substituted derivatives thereof. Examples of the carbocyclic aryl groups that R1 to R4 can represent include radicals having the formulas (V) – (VII):






wherein R9 and R10 may represent one or more substituents independently selected from alkyl, alkoxy, halogen, cycloalkoxy, formyl, alkanoyl, cycloalkyl, aryl, aryloxy, aroyl, carboxyl, carboxylate salts, alkoxy-carbonyl, alkanoyloxy, cyano, sulfonic acid, sulfonate salts, and the like. In some embodiments the alkyl moiety of the aforesaid alkyl, alkoxy, alkanoyl, alkoxy-carbonyl, and alkanoyloxy groups contains up to 8 carbon atoms. Although it is possible for p to represent 0 to 5 and for q to represent 0 to 7, in some embodiments, the value of each of p and q will not exceed 2. In some embodiments R9 and R10 represent lower alkyl groups, i.e., straight-chain and branched-chain alkyl of up to 4 carbon atoms, and m and n each represent 0, 1, or 2.

[0018] The hydrocarbonyl groups represented by R5 and R6 may be the same or different, separate or combined, and are selected from unsubstituted and substituted alkyl, cycloalkyl, and aryl groups containing a total of up to 40 carbon atoms. The total carbon content of substituents of R5 to R6 in some embodiments is in the range of 1 to 35 carbon atoms. Examples of the alkyl groups that R5 and R6 can represent include methyl, ethyl, butyl, pentyl, hexyl, 2-ethylhexyl, octyl, decyl, dodecyl, octadecyl, 1-alkylbenzyl, and various isomers thereof. The alkyl groups may be substituted, for example, with up to two substituents such as alkoxy, cycloalkoxy, formyl, alkanoyl, cycloalkyl, aryl, aryloxy, aroyl, carboxyl, carboxylate salts, alkoxy-carbonyl, alkanoyloxy, cyano, sulfonic acid, sulfonate salts, and the like. Cyclopentyl, cyclohexyl, and cycloheptyl are examples of the cycloalkyl groups that R5 and R6 individually can represent. The cycloalkyl groups may be substituted with alkyl or any of the substituents described with respect to the possible substituted alkyl groups. In some embodiments the alkyl and cycloalkyl groups that R5 and R6 represent are alkyl of up to 10 carbon atoms such as benzyl, 1-alkylbenzyl, cyclopentyl, cyclohexyl, or cycloheptyl, etc.

[0019] Examples of the aryl groups that R5 and R6 can represent include carbocyclic aryl such as phenyl, naphthyl, anthracenyl, and substituted

derivatives thereof. Examples of the carbocyclic aryl groups that R5 and R6 can represent include radicals having the formulas (VIII) – (X):




wherein R11 and R12 may represent one or more substituents independently selected from alkyl, alkoxy, halogen, cycloalkoxy, formyl, alkanoyl, cycloalkyl, aryl, aryloxy, aroyl, carboxyl, carboxylate salts, alkoxy-carbonyl, alkanoyloxy, cyano, sulfonic acid, sulfonate salts, and the like. In some embodiments, the alkyl moiety of the aforesaid alkyl, alkoxy, alkanoyl, alkoxy-carbonyl, and alkanoyloxy groups contains up to 8 carbon atoms. Although it is possible for r to represent 0 to 5 and for s to represent 0 to 7, in some embodiments, the value of each of r and s will not exceed 2. R11 and R12 represent lower alkyl groups, i.e., straight-chain and branched-chain alkyl of up to 10 carbon atoms, and r and s each represent 0, 1, or 2.


[0020] In some embodiments, R1 to R6 in combination or collectively may represent a divalent hydrocarbyl group containing up to 40 carbon atoms. In some embodiments, the divalent hydrocarbyl group contains from 12 to 36

carbon atoms. Examples of such divalent groups include alkyl of 2 to 12 carbon atoms, cyclohexyl, and aryl. Specific examples of the alkyl and cycloalkyl groups include ethyl, trimethyl, 1,3-butanediyl, 2,2-dimethyl-1,3-propanediyl, 1,1,2-triphenylethanediy, 2,2,4-trimethyl-1,3-pentanediy, 1,2-cyclohexyl, and the like.


[0021] In some embodiments, the fluorophosphite compound according to the invention has the structure of formula (XI):



Wherein R15 is selected from hydrogen and alkyl or aryl having 1 to 10 carbon atoms. In some embodiments, the fluorophosphite compound has the following structure:



In some embodiments, the fluorophosphite compound has the following structure:



[0022] The fluorophosphite compounds of this invention can be prepared by any effective method. A variety of methods for preparing fluorophosphates are reported in the literature. For example, it has been found that the fluorophosphites can be prepared by using a (benzyl)phenol starting material such as 2,2'-methylene bis(4,6-di(α,α -dimethylbenzyl)phenol) and following the procedures described in U.S. Patent No. 4,912,155; Tullock et al., *J. Org. Chem.*, 25, 2016 (1960); White et al., *J. Am. Chem. Soc.*, 92, 7125 (1970); Meyer et al., *Z. Naturforsch., Bi. Chem. Sci.*, 48, 659 (1993); or Puckette, in "Catalysis of Organic Reactions", Edited by S. R. Schmidt, CRC Press (2006), pp. 31-38.

[0023] The novel catalyst systems provided by the present invention comprise a combination of one or more transition metals selected from the Group VIII metals and rhenium and one or more of the fluorophosphite compounds described in detail hereinabove. The transition metal may be provided in the form of various metal compounds such as carboxylate salts of the transition metal. In some embodiments, the metal is rhodium.

[0024] Rhodium compounds that may be used as a source of rhodium for the active catalyst include rhodium (II) or rhodium (III) salts of carboxylic acids, examples of which include di-rhodium tetraacetate dihydrate, rhodium(II) acetate, rhodium(II) isobutyrate, rhodium(II) 2-ethylhexanoate, rhodium(II) benzoate, and rhodium(II) octanoate. Also, rhodium carbonyl species such as $\text{Rh}_4(\text{CO})_{12}$, $\text{Rh}_6(\text{CO})_{16}$, and rhodium(I) acetylacetonate dicarbonyl may be suitable rhodium feeds. Additionally, rhodium organophosphine complexes such as tris(triphenylphosphine) rhodium

carbonyl hydride may be used when the phosphine moieties of the complex fed are easily displaced by the phosphite ligands of the present invention. Other rhodium sources include rhodium salts of strong mineral acids such as chlorides, bromides, nitrates, sulfates, phosphates, and the like.

[0025] The ratio of gram moles fluorophosphite ligand to gram atoms transition metal can vary over a wide range, e.g., gram mole fluorophosphite:gram atom transition metal ratio of 1:1 to 400:1. For rhodium-containing catalyst systems, the gram mole fluorophosphite:gram atom rhodium ratio in some embodiments is in the range of 1:1 to 200:1 with ratios in the range of 1:1 to 120:1. The absolute concentration of rhodium in the reaction mixture or solution may vary from 1 mg/liter up to 5000 mg/liter or more. In some embodiments of this invention, the normal concentration of rhodium in the reaction solution is in the range of 20 to 300 mg/liter. Concentrations of rhodium lower than this range generally yield lower reaction rates with most olefin reactants and/or require reactor operating temperatures that are so high as to be detrimental to catalyst stability. Concentrations above this range involve higher rhodium costs.

[0026] No special or unusual techniques are required for preparing the catalyst systems and solutions of the present invention, although in some embodiments a catalyst of high activity is obtained if all manipulations of the rhodium and fluorophosphite ligand components are carried out under an inert atmosphere, e.g., nitrogen, argon, and the like. The desired quantities of a suitable rhodium compound and ligand are charged to the reactor in a suitable solvent. The sequence in which the various catalyst components or reactants are charged to the reactor is not critical.

[0027] The hydroformylation solvent may be selected from a wide variety of compounds, mixture of compounds, or materials that are liquid at the pressure at which the process is being operated. Such compounds and materials include various alkanes, cycloalkanes, alkenes, cycloalkenes, carbocyclic

aromatic compounds, alcohols, esters, ketones, acetals, ethers, and water. Specific examples of such solvents include alkane and cycloalkanes such as dodecane, decalin, octane, iso-octane mixtures, cyclohexane, cyclooctane, cyclododecane, methylcyclohexane; aromatic hydrocarbons such as benzene, toluene, xylene isomers, tetralin, cumene, alkyl-substituted aromatic compounds such as the isomers of diisopropylbenzene, triisopropylbenzene and tert-butylbenzene; alkenes and cycloalkenes such as 1,7-octadiene, dicyclopentadiene, 1,5-cyclooctadiene, octene-1, octene-2, 4-vinylcyclohexene, cyclohexene, 1,5,9-cyclododecatriene, 1-pentene; crude hydrocarbon mixtures such as naphtha, mineral oils and kerosene; and high-boiling esters such as 2,2,4-trimethyl-1,3-pentanediol diisobutyrate. The aldehyde product of the hydroformylation process also may be used.

[0028] In some embodiments, the solvent is the higher boiling by-products that are naturally formed during the process of the hydroformylation reaction and the subsequent steps, e.g., distillations, that may be required for aldehyde product isolation. The main criterion for the solvent is that it dissolves the catalyst and olefin substrate and does not act as a poison to the catalyst. Some examples of solvents for the production of volatile aldehydes, e.g., butyraldehydes, are those that are sufficiently high boiling to remain, for the most part, in a gas sparged reactor. Some examples of solvents and solvent combinations that are useful in the production of less volatile and non-volatile aldehyde products include 1-methyl-2-pyrrolidinone, dimethylformamide, perfluorinated solvents such as perfluoro-kerosene, sulfolane, water, and high boiling hydrocarbon liquids as well as combinations of these solvents. Non-hydroxylic compounds, in general, and hydrocarbons, in particular, may be used advantageously as the hydroformylation solvent since their use can minimize decomposition of the fluorophosphite ligands.

[0029] In another aspect, the present invention provides a process for preparing aldehydes, comprising contacting an olefin with hydrogen and

carbon monoxide, under hydroformylation conditions, in the presence of the catalyst solution herein described.

[0030] The reaction conditions for the process of the present invention can be conventional hydroformylation conditions. The process may be carried out at temperatures in the range of 20° to 200°C, in some embodiments the hydroformylation reaction temperatures are from 50° to 135°C. In some embodiments, reaction temperatures range from 75° to 125°C. Higher reactor temperatures can increase the rate of catalyst decomposition while lower reactor temperatures can result in relatively slow reaction rates. The total reaction pressure may range from ambient or atmospheric up to 70 bars absolute (1000 psig). In some embodiments, pressure ranges from 8 to 28 bars absolute (100 to 400 psig).

[0031] The hydrogen:carbon monoxide mole ratio in the reactor likewise may vary considerably ranging from 10:1 to 1:10, and the sum of the absolute partial pressures of hydrogen and carbon monoxide may range from 0.3 to 36 bars absolute. The partial pressures of the ratio of the hydrogen to carbon monoxide in the feed can be selected according to the linear:branched isomer ratio desired. Generally, the partial pressure of hydrogen and carbon monoxide in the reactor can be maintained within the range of 1.4 to 13.8 bars absolute (20 to 200 psia) for each gas. The partial pressure of carbon monoxide in the reactor can be maintained within the range of 1.4 to 13.8 bars absolute (20 to 200 psia) and can be varied independently of the hydrogen partial pressure. The molar ratio of hydrogen to carbon monoxide can be varied widely within these partial pressure ranges for the hydrogen and carbon monoxide. The ratios of hydrogen-to-carbon monoxide and the partial pressure of each in the synthesis gas (syn gas--carbon monoxide and hydrogen) can be readily changed by the addition of either hydrogen or carbon monoxide to the syn gas stream. With the fluorophosphite ligands described herein, the ratio of linear to branched products can be varied widely by changing the partial pressures of the carbon monoxide in the reactor.

[0032] Any of the known hydroformylation reactor designs or configurations such as overflow reactors and vapor take-off reactors may be used in carrying out the process provided by the present invention. Thus, a gas-sparged, vapor take-off reactor design as disclosed in the examples set forth herein may be used. In this mode of operation, the catalyst which is dissolved in a high boiling organic solvent under pressure does not leave the reaction zone while the aldehyde product is taken overhead by the unreacted gases. The overhead gases then are chilled in a vapor/liquid separator to liquefy the aldehyde product and the gases can be recycled to the reactor. The liquid product is let down to atmospheric pressure for separation and purification by conventional techniques. The process also may be practiced in a batchwise manner by contacting the olefin, hydrogen and carbon monoxide with the present catalyst in an autoclave.

[0033] A reactor design where catalyst and feedstock are pumped into a reactor and allowed to overflow with product aldehyde, i.e. liquid overflow reactor design, is also suitable. For example, high boiling aldehyde products such as nonyl aldehydes may be prepared in a continuous manner with the aldehyde product being removed from the reactor zone as a liquid in combination with the catalyst. The aldehyde product may be separated from the catalyst by conventional means such as by distillation or extraction, and the catalyst then recycled back to the reactor. Water soluble aldehyde products, such as hydroxy butyraldehyde products obtained by the hydroformylation of allyl alcohol, can be separated from the catalyst by extraction techniques. A trickle-bed reactor design is also suitable for this process. It will be apparent to those skilled in the art that other reactor schemes may be used with this invention.

[0034] The olefin used as the starting material for this invention is not particularly limiting. Specifically, the olefin can be ethylene, propylene, butene, pentene, hexene, octene, styrene, non-conjugated dienes such as

1,5-hexadiene, and blends of these olefins. Furthermore, the olefin may be substituted with functional groups so long as they do not interfere with the hydroformylation reaction. Suitable substituents on the olefin include any functional group that does not interfere with the hydroformylation reaction and includes groups such as carboxylic acids and derivatives thereof such as esters and amides, alcohols, nitriles, and ethers. Examples of substituted olefins include esters such as methyl acrylate or methyl oleate, alcohols such as allyl alcohol and 1-hydroxy-2,7-octadiene, and nitriles such as acrylonitrile.

[0035] The amount of olefin present in the reaction mixture can also vary. For example, relatively high-boiling olefins such as 1-octene may function both as the olefin reactant and the process solvent. In the hydroformylation of a gaseous olefin feedstock such as propylene, the partial pressures in the vapor space in the reactor are in the range of 0.07 to 35 bars absolute. The rate of reaction can be favored by high concentrations of olefin in the reactor. In the hydroformylation of propylene, the partial pressure of propylene in some embodiments is greater than 1.4 bars, e.g., from 1.4 to 10 bars absolute. In the case of ethylene hydroformylation, the partial pressure of ethylene in the reactor in some embodiments is greater than 0.14 bars absolute.


[0036] This invention can be further illustrated by the following examples of embodiments thereof, although it will be understood that these examples are included merely for purposes of illustration and are not intended to limit the scope of the invention. Unless otherwise indicated, all percentages are by weight. As used throughout this application, the reference to a molecule or moiety as being "substituted" means that the molecule or moiety contains one or more substituent in place of where a hydrogen atom would be on the molecule or moiety identified. Thus, a "substituted" aryl molecule having alkyl substituents would include, for example, an alkylbenzene such as toluene.

EXAMPLES

Preparation of Ligand A

[0037] Ligand A was synthesized from starting material 2,2'-methylene bis(4,6-di(α,α -dimethylbenzyl)phenol) by following the procedures described in U.S. Patent No. 4,912, 155; the entire content of which is hereby incorporated by reference. Ligand A is very soluble in common organic solvents such as toluene, acetone, cyclohexane, ethyl acetate, etc.

[0038] Spectroscopic Data of Ligand A:




[0039] Ligand A is a new compound and shows a doublet on ^{31}P NMR (in CDCl_3) centered at 121.6 ppm ($J_{\text{F-P}} = 1320$ Hz). The chemical shift of ^{19}F NMR of Ligand A shows a doublet centered at -64.9 ppm ($J_{\text{P-F}} = 1301$ Hz). The chemical shift (in ppm) of ^1H NMR of Ligand A in CDCl_3 is as follows: 1.42 (s, 6H), 1.58 (s, 6H), 1.67 (s, 12H), 3.28 (d, 1H), 3.90(dd, 1H), 6.96-7.31 (m, 24H).

Preparation of Ligand B

[0040] Ligand B was synthesized from starting material 2,4-di-(α,α -dimethylbenzyl)phenol by following the procedures described in U.S. Patent No. 4,912,155. Ligand B is slightly less soluble than Ligand A in common organic solvents.


[0041] Spectroscopic Data of Ligand B:



[0042] Ligand B is a new compound and shows a doublet on ³¹P NMR (in CDCl₃) centered at 119.89ppm ($J_{F-P} = 1265$ Hz). The chemical shift of ¹⁹F NMR of Ligand B shows a doublet centered at -57.81ppm ($J_{P-F} = 1255$ Hz). The chemical shift (in ppm) of ¹H NMR of Ligand B in CDCl₃ is as follows: 1.54 (s, 6H), 1.57 (s, 6H), 1.69 (s, 12H), 6.47 (d, 2H), 7.03 (d, 2H), 7.13-7.35 (m, 22H).


Ligand C

[0043] Ligand C is a compound taken from Puckette, in "Catalysis of Organic Reactions", Edited by S. R. Schmidt, CRC Press (2006), p. 31. The ligand described in this literature is also a fluorophosphite (Ethanox 398™, 2,2'-ethyldenebis(4,6-di-tert-butylphenyl) fluorophosphite).



Ligand D

Ligand D was synthesized from starting material 2,4-di-(α,α-dimethylbenzyl)phenol by following the procedures described in U.S. Patent No. 4,912,155.



Ligand D is a new compound and shows a doublet on ^{31}P NMR (in CDCl_3) centered at 105.9 ppm ($J_{\text{F-P}} = 1286$ Hz). The chemical shift of ^{19}F NMR of Ligand A shows a doublet centered at -61.5 ppm ($J_{\text{P-F}} = 1297$ Hz). The chemical shift (in ppm) of ^1H NMR of Ligand D in CDCl_3 is as follows: 1.39 (d, 3H), 1.50 (s, 6H), 1.62 (s, 6H), 1.70 (s, 12H), 4.31 (br s, 1H), 6.97 (s, 2H), 7.11-7.31 (m, 22H).

Hydroformylation Process Set-Up

[0044] The hydroformylation process in which propylene is allowed to react with hydrogen and carbon monoxide to produce butyraldehydes was carried out in a vapor take-off reactor made up of a vertically arranged stainless steel pipe having a 2.5 cm inside diameter and a length of 1.2 meters. The reactor was encased in an external jacket that was connected to a hot oil machine. The reactor has a filter element welded into the side down near the bottom of the reactor for the inlet of gaseous reactants. The reactor contained a thermocouple which was arranged axially with the reactor in its center for accurate measurement of the temperature of the hydroformylation reaction mixture. The bottom of the reactor has a high pressure tubing connection that was connected to a cross. One of the connections to the cross permitted the addition of non-gaseous reactants such as higher boiling alkenes or make-up solvents, another led to the high-pressure connection of a differential pressure (D/P) cell that was used to measure catalyst level in the reactor and the bottom connection was used for draining the catalyst solution at the end of the run.

[0045] In the hydroformylation of propylene in a vapor take-off mode of operation, the hydroformylation reaction mixture or solution containing the catalyst was sparged under pressure with the incoming reactants of propylene, hydrogen and carbon monoxide as well as any inert feed such as nitrogen. As butyraldehyde was formed in the catalyst solution, it and unreacted reactant gases were removed as a vapor from the top of the reactor by a side-port. The vapor removed was chilled in a high-pressure separator where the butyraldehyde product was condensed along with some of the unreacted propylene. The uncondensed gases were let down to atmospheric pressure via the pressure control valve. These gases passed through a series of dry-ice traps where any other aldehyde product was collected. The product from the high-pressure separator was combined with that of the traps, and was subsequently weighed and analyzed by standard gas/liquid phase chromatography (GC/LC) techniques for the net weight and normal/iso ratio of the butyraldehyde product.

[0046] The gaseous feeds to the reactor were fed to the reactor via twin cylinder manifolds and high-pressure regulators. The hydrogen passed through a mass flow controller and then through a commercially available "Deoxo" (registered trademark of Englehard Inc.) catalyst bed to remove any oxygen contamination. The carbon monoxide passed through an iron carbonyl removal bed (as disclosed in U.S. Patent No. 4,608,239), a similar "Deoxo" bed heated to 125°C, and then a mass flow controller. Nitrogen can be added to the feed mixture as an inert gas. Nitrogen, when added, was metered in and then mixed with the hydrogen feed prior to the hydrogen Deoxo bed. Propylene was fed to the reactor from feed tanks that were pressurized with hydrogen and was controlled using a liquid mass flow meter. All gases and propylene were passed through a preheater to ensure complete vaporization of the liquid propylene prior to entering the reactor.

Comparative Examples 1-3 (Ligand C)

[0047] Comparative Examples 1-3 were taken directly from Puckette, "Catalysis of Organic Reactions", Edited by S. R. Schmidt, CRC Press (2006), pp. 31-38, to illustrate that hydroformylation activity decreases as the molar ratio of ligand to Rh increases (the concentration of ligand increases). The ligand described in this literature is a fluorophosphite (Ethanox 398™, 2,2'-ethylidenebis(4,6-di-tert-butylphenyl) fluorophosphite) (Ligand C).

[0048] According to the literature, the reactions were conducted at 260 psig, 115°C, 1:1 H₂/CO, 54 psia C₃H₆, and 190 mL of bis-2-ethylhexylphthalate solvent (DOP) with different amounts of ligand. The data is presented in Table 1 below and catalyst activity is expressed as kilograms of butyraldehyde per gram Rh per hour.

Comparative Example 4 (Ligand B)

[0049] This example illustrates a typical hydroformylation run and the use of Ligand "B" for hydroformylation of propylene.

[0050] A catalyst solution was prepared under nitrogen using a charge of 7.5 mg of rhodium (0.075 mmol, as rhodium 2-ethylhexanoate); ligand "B" (O,O-di-2,4- α,α -dimethylbenzylphenyl) fluorophosphite), 0.80g, (1.125 mmol); 20 ml of normal butyraldehyde, and 190 ml of bis-2-ethylhexylphthalate solvent (DOP). The mixture was stirred under nitrogen until a homogeneous solution was obtained (heated, if necessary). The mixture was charged to the reactor in a manner described previously and the reactor sealed. The reactor pressure control was set at 17.9 bar (260 psig) and the external oil jacket on the reactor was heated to 85°C. Hydrogen, carbon monoxide, nitrogen, and propylene vapors were fed through the frit at the base of the reactor and the reactor was allowed to build pressure. The hydrogen and carbon monoxide (H₂/CO ratio was set to be 1:1) were fed to the reactor at a rate of 6.8 liters/min, and the nitrogen feed was set at 1.0 liter/min. The propylene was metered as a liquid and fed at a rate of 1.89 liters/min (212 grams/hour). The

temperature of the external oil was modified to maintain an internal reactor temperature of 85°C. The unit was operated for 5 hours and hourly samples taken. The hourly samples were analyzed as described above using a standard GC method. The last three samples were used to determine the N/I ratio and catalyst activity. The catalyst activity of for the last three hours averaged 13.2 kilograms butyraldehyde/gram of rhodium-hour. The product N/I ratio was 1.40.

Comparative Examples 5 and 6 (Ligand B)

[0051] Hydroformylation experiments were carried out in the same manner as Comparative Example 4, except utilizing different amounts of Ligand "B". The reaction conditions and the results of this work are presented in Table 1.

Examples 7-10 (Ligand A)

[0052] Hydroformylation experiments were carried out in the same manner as Comparative Example 4, except utilizing various amounts of Ligand "A". The reaction conditions and the results of this work are presented in Table 1.

Table 1: Effects of Ligand to Rhodium Molar Ratio on Activity and Selectivity

Example Nos.	Ligand	Temp (°C)	H ₂ /CO ratio	Ligand to Rh Molar ratio	N/I Ratio	Catalyst Activity*
C-1	C	115	1:1	14:1	1.4	15.7
C-2	C	115	1:1	30:1	3.1	6.6
C-3	C	115	1:1	50:1	3.8	3.3
C-4	B	85	1:1	15:1	1.40	13.2
C-5	B	85	1:1	30:1	1.85	7.2
C-6	B	85	1:1	60:1	2.59	5.0
7	A	85	1:1	15:1	1.61	3.0
8	A	85	1:1	30:1	1.89	3.23
9	A	85	1:1	60:1	2.50	5.49
10	A	85	1:1	90:1	2.72	6.67
11	D	95	1:1	5:1	1.41	3.72
12	D	95	1:1	15:1	1.94	9.98
13	D	95	1:1	30:1	2.72	15.46

*Activity was determined as kilograms of butyraldehydes produced per gram of rhodium per hour. All examples were done using 0.075 mmol of Rh.

[0053] Comparative Examples 1 through 3 illustrate that the hydroformylation reaction activity, under the same reaction temperature and pressure, steadily decreased from 15.7 to 3.3.

[0054] Comparative Examples 4, 5, and 6 illustrate again the typical, conventional behavior of phosphorus ligands. Under the same reaction temperature and pressure, hydroformylation activity steadily decreased from 13.2 to 5.0

[0055] Examples 7 through 10 illustrate the desirable nature of Ligand "A". The difference between Ligand "A" and "B" is that "A" has a cyclic ring structure (methylene group connecting two aromatic rings). Example 7 showed an activity of 3.0 for a 15:1 Ligand/Rh ratio. As the ratio of Ligand/Rh increased to 90:1 under the same reaction temperature and pressure, the activity increased to 6.67. The trend of changes in hydroformylation activity in Examples 7 through 10 is completely opposite to the trend showed in Comparative Examples 1 through 3 or Comparative Examples 4 through 6.


Examples 11 through 13 illustrate the desirable nature of Ligand "D". Example 11 through 13 show that as the ligand to rhodium ratio is increased from 15:1 to 30:1, the catalyst activity increases from 3.72 to 15.46 while the N/I ratio shows a corresponding increase from 1.41 to 2.72.

[0056] The invention has been described in detail with particular reference to preferred embodiments and illustrative examples thereof, but it will be understood that variations and modifications can be effected within the spirit and scope of the invention.

CLAIMS

What is claimed is:

1. A fluorophosphite compound having the structure of formula (I):




wherein

Ar₁ and Ar₂ are each independently selected from aryl groups which are substituted or unsubstituted, and which contain 4 to 30 carbon atoms;

R1 to R6 are each independently selected from H and alkyl cycloalkyl and aryl groups which are substituted or unsubstituted, and which contain 1 to 40 carbon atoms; and




X is

- (i) a chemical bond directly between ring carbon atoms of each aromatic group, except when R1 to R4 are each methyl,
- (ii) a heteroatom selected from O, Si, and N, or
- (iii) a group having the formula



wherein R13 and R14 are each independently selected from hydrogen and alkyl or aryl having 1 to 10 carbon atoms.

2. The compound according to claim 1, wherein Ar_1 and Ar_2 are each independently selected from aryl groups having the structure of formulas (II) – (IV):



wherein


R7 and R8 are independently selected from alkyl, alkoxy, halogen, cycloalkoxy, formyl, alkanoyl, cycloalkyl, aryl, aryloxy, aroyl, carboxyl, carboxylate salts, alkoxy-carbonyl, alkanoyloxy, cyano, sulfonic acid, and sulfonate salts;

m is an integer from 0 to 5; and

n is an integer from 0 to 7.


3. The compound according to claim 2, wherein R7 and R8 are independently selected from alkyl having 1 to 10 carbon atoms, and m and n are independently 0, 1, or 2.

4. The compound according to claim 1, having the structure of formula (XI):




Wherein R₁₅ is selected from hydrogen and alkyl or aryl having 1 to 10 carbon atoms.

5. The compound according to claim 1, which has the following structure:




6. The compound according to claim 1, which has the following structure:



7. A catalyst solution comprising:
- (a) a fluorophosphite compound;
 - (b) a Group VIII metal or rhenium; and
 - (c) a hydroformylation solvent,

wherein said fluorophosphite compound has the structure of formula (I):




wherein

Ar_1 and Ar_2 are each independently selected from aryl groups which are substituted or unsubstituted, and which contain 4 to 30 carbon atoms;

R_1 to R_6 are each independently selected from H and alkyl cycloalkyl and aryl groups which are substituted or unsubstituted, and which contain 1 to 40 carbon atoms; and




X is

- (i) a chemical bond directly between ring carbon atoms of each aromatic group,
- (ii) a heteroatom, or
- (iii) a group having the formula



wherein R_{13} and R_{14} are each independently selected from hydrogen and alkyl or aryl having 1 to 10 carbon atoms.

8. The catalyst solution according to claim 7, wherein Ar_1 and Ar_2 are each independently selected from aryl groups having the structure of formulas (II) – (IV):



wherein


R7 and R8 are independently selected from alkyl, alkoxy, halogen, cycloalkoxy, formyl, alkanoyl, cycloalkyl, aryl, aryloxy, aroyl, carboxyl, carboxylate salts, alkoxy-carbonyl, alkanoyloxy, cyano, sulfonic acid, and sulfonate salts;

m is an integer from 0 to 5; and

n is an integer from 0 to 7.


9. The catalyst solution according to claim 8, wherein R7 and R8 are independently selected from alkyl having 1 to 10 carbon atoms, and m and n are independently 0, 1, or 2.

10. The catalyst solution according to claim 7, wherein the fluorophosphite compound has the structure of formula (XI):




Wherein R15 is selected from hydrogen and alkyl or aryl having 1 to 10 carbon atoms.

11. The catalyst solution according to claim 7, wherein the fluorophosphite compound has the following structure:



12. The catalyst solution according to claim 7, wherein the fluorophosphite compound has the following structure:



13. The catalyst solution according to any of claims 7-12, wherein the Group VIII metal is rhodium.

14. The catalyst solution according to any of claims 7-13, which comprises from 20 to 300 mg/l of rhodium, and a ratio of gram moles of fluorophosphite to gram atom of rhodium of 1:1 to 200:1.

15. The catalyst solution according to any of claims 7-14, wherein the hydroformylation solvent is selected from alkanes, cycloalkanes, alkenes, cycloalkenes, alcohols, esters, ketones, acetals, ethers, aldehydes, water, and mixtures thereof.

16. A process for preparing aldehydes, comprising contacting an olefin with hydrogen and carbon monoxide, under hydroformylation conditions, in the presence of the catalyst solution according to any of claims 7-15.

17. The process according to claim 16, wherein the hydroformylation conditions comprise a temperature ranging from 75 to 125°C and a pressure from atmospheric to 70 bars absolute (15 to 1,000 psig).

18. The process according to claim 16 or 17, wherein the olefin is propylene, and the aldehydes comprise normal- and iso-butyraldehyde.