Polypeptide Vaccine

FIELD OF THE INVENTION

The present invention relates to the fields of cell biology, molecular biology and biochemistry. This invention also relates to the field of medicine. In particular, it relates to prevention and treatment of diseases, in particular cancer, as well as compositions for such use.

BACKGROUND TO THE INVENTION

Cancer has been the subject of intense investigation in the past two decades. However, the underlying causes responsible for cancer metastasis are poorly understood and most types of cancer preventions are still limited. Effective ways to prevent cancer metastasis are urgently needed.

Antibody based therapy has proven to be effective for cancer treatment; however, this approach has been traditionally limited to extracellular or secreted proteins expressed by cancer cells[1,2]. Thus, a number of potential cancer or tumour markers and cancer antigens have been identified in the literature and antibody therapies have been developed against some of them.

For example, the well-known cancer therapy Herceptin (Trastuzumab) is a monoclonal antibody that can kill HER2-positive cancer cells. Herceptin binds to the HER2 (human epidermal growth factor receptor 2) antigen on the cancer cell. Likewise, Bevacizumab (Avastin™) is a monoclonal antibody targeted against vascular endothelial growth factor (VEGF), one of the growth factors implicated in the formation of new blood vessels. By inhibiting angiogenesis, Bevacizumab prevents tumour cells from receiving a constant supply of blood to receive the oxygen and nutrients the tumour needs to survive.

However, the applicability of antibody therapeutics for different cancers is not universal. One of the limitations that has prevented the general use of antibody therapeutics is the large size of antibody molecules and their consequent inability to cross the plasma or cell membrane. In the absence of modification, antibodies (including monoclonal antibodies) are only generally suitable for targeting cancer antigens located at the surface or exterior of host cells [14-15]. In the examples above, HER2 receptor is located on the cell surface and is hence accessible for antibody binding by Herceptin. Likewise, VEGF is secreted into the bloodstream and is able to be bound by Bevacizumab.

Most oncogenic proteins are intracellular proteins (such as intracellular phosphatases, intracellular kinases, transcription factors, etc), and have remained under-explored by the approach of antibody therapies. The long held view that antibodies are too large to penetrate cell membrane has hampered the technology of antibody therapy used in targeting intracellular proteins.

There is therefore an urgent need for effective ways of treating and preventing cancer metastasis. Antibodies have not hitherto been used for targeting intracellular antigens or cancer markers because of the inability of the antibodies to cross the cell membrane and the consequent inaccessibility of the antigen.

Antibody-based therapies have better specificity and thus improved efficacy over standard chemotherapy regimens. Because antibodies are viewed as too large to access intracellular (inside the cell) locations, antibody therapy has traditionally targeted extracellular (outside the cell or cell surface) proteins expressed by cancer cells. However, a large pool of oncogenic proteins is found within the cell (such as intracellular phosphatases/kinases and transcription factors) and has therefore not been pursued for antibody therapies.

We previously showed three different antibodies could respectively target three intracellular proteins: PRL-3 (phosphatase of regenerating liver 3), a cancer-associated phosphatase; EGFP (enhanced green fluorescent protein), a general reporter; and mT (polyomavirus middle T), the polyomavirus middle T oncoprotein (WO2011/065923). Yet, only PRL-3 intracellular phosphatase (an enzyme) has been linked to human cancer metastasis (see Saha et al., Science 294; 1343 (2001) and Wang et al., Cancer cell 18; 52-63 (2010)), the other two intracellular proteins (EGFP and middle T) are used to elucidate the general phenomena that antibody can target intracellular proteins. However, the use of oncoproteins as cancer vaccines is controversial, and a need exists to develop improved oncoprotein cancer vaccines that are more specific, and exhibit less cross-reaction with homologous proteins to reduce side effects, whilst achieving similar, or improved, therapeutic results.

Oncogenic mutations are very common in contributing to multiple human cancers. However, these oncogenic mutations are often detected in intracellular proteins or intracellular domains of cell surface proteins. Our recent works suggest an unconventional concept that intracellular oncoproteins can be targeted by therapeutic antibodies or peptide vaccination. In this new consideration generating antibodies against those specific mutations or

vaccination using mutant peptides could specifically ablate cancer cells expressing respective mutated targets, but sparing normal tissues unharmed. However, making antibodies one by one *in vitro* specifically against countless point mutations discovered so far is very difficult and not practical. Therefore, vaccination using mutant peptides could be a choice.

SUMMARY OF THE INVENTION

In a first aspect, the present invention refers to the use of a polypeptide in the manufacture of a medicament for the treatment of cancer, wherein the treatment involves stimulating the production of anti-intracellular oncoprotein antibodies in a subject in need thereof, the polypeptide comprising a fragment of an intracellular oncoprotein having between 10 and 150 amino acids, preferably between 10 and 100 amino acids, wherein the intracellular oncoprotein is PRL3.

In a second aspect, the present invention refers to the use of a fragment of between 10 and 150 amino acids of an intracellular oncoprotein in the manufacture of a medicament for treating cancer metastasis in a patient with cancer, wherein the intracellular oncoprotein is identified to be specific to the patient's cancer, wherein the intracellular oncoprotein is PRL3.

In a third aspect, the present invention refers to the use of a fragment of an intracellular oncoprotein comprising between 10 and 150 amino acids in length in the manufacture of a medicament for treating cancer in a person, wherein the person is determined as suffering from, or likely to develop, a cancer associated with the intracellular oncoprotein before administration to that person, wherein the intracellular oncoprotein is PRL 3.

Also described herein is a pharmaceutical composition comprising an intracellular oncoprotein fragment, or a variant, homologue or derivative thereof, together with a pharmaceutically acceptable excipient, diluent or carrier.

As described herein, intracellular oncoproteins can be targeted with vaccination to prevent cancer and cancer metastasis.

As described herein, wild-type C57BL6 mice intravenously (i.v.) injected either with B16F0 or B16F10 melanoma cells, followed by i.v. injection with PRL-3 monoclonal antibody (mAb) showed that PRL-3 mAb can reduce tumors formed by B16F0 melanoma cells that express endogenous PRL-3 intracellular phosphatase,[1] but not those by B16F10 melanoma cells that do not express PRL-3.

C57BL6 mice i.v. injected with either EGFP-B 16F0 or EGFP-B16F10 cells in which both cell lines were engineered to overexpress EGFP, followed by i.v. injection with EGFP monoclonal antibody showed that regardless of PRL-3 expression status, EGFP mAb could eradicate tumors formed by both EGFP-B 16F0 and EGFP-B 16F 10 melanoma cell lines.
Further, polyomavirus middle T (Py-mT)-transgenic young females i.v. injected with mT antibody showed that the mT antibody could effectively reduce the formation of mT-expressing breast tumors.

Moreover, C57BL6 mice vaccinated with PRL-3 or EGFP antigens showed significant reduction in the formation of metastatic tumors expressing either PRL-3 or EGFP protein respectively, as compared with un-immunized mice. Furthermore, Py-mT mice vaccinated with mT antigen significantly prevented tumor formation in mammary glands. The results obtained from 194 wild type mice demonstrate that cancer metastasis caused by intracellular oncoproteins can be successfully blocked with antibody therapy and vaccination.

As shown herein, young PymT transgenic mice injected with fragments of the middle T protein exhibited reduced incidence of breast tumor than mice immunized with GST (negative control). Moreover, immunization of mice with Her2/neu fragments containing only a fragment of the intracellular domain inhibited the formation of Her2/neu expressing tumors formed by B16F0 melanoma cancer cells.

Further, MMTV-PymT mice vaccinated with Estrogen Receptor peptides or with Estrogen Receptor antibody have reduced burden of mammary tumors compared to mice immunized with GST (negative control).

We therefore provide for methods of treating cancers and preventing cancer metastasis with fragments of intracellular oncoprotein. The term “intracellular oncoprotein” as used herein may include oncoproteins that have an intracellular region. For example, membrane anchored proteins may have extracellular, transmembrane, and intracellular domains or regions.

The fragment may be a fragment of a region of the oncoprotein that comprises a mutation as compared to the wild-type, or non-oncogenic protein. The mutation may be associated with, or causative of, cancer (i.e. an oncogenic mutation).

There is provided, according to an example disclosed herein, use of a vaccine therapy comprising an intracellular oncoprotein fragment for vaccination to stimulate a host to produce its own antibodies to prevent tumor formation.

Also provided is polypeptide for use in a method of stimulating the production of anti-intracellular oncoprotein antibodies in a subject in need thereof, the polypeptide being a fragment of the intracellular oncoprotein having between 10 and 150 amino acids, preferably between 10 and 100 amino acids. A method of stimulating the production of anti-intracellular oncoprotein antibodies in a subject in need thereof, comprising administering a polypeptide which is a fragment of the intracellular oncoprotein, having between 10 and 150 amino acids is also provided.

The polypeptide may be a fragment of an oncoprotein that is intracellular, or a fragment of an intracellular region of an oncoprotein. The fragment may comprise an oncogenic mutation.

The polypetide may comprise a plurality of fragments. Thus, more than one fragment of an oncoprotein may be provided in one polypeptide. More than one polypeptide may be administered to a patient. Some or all of those polypeptides may comprise more than one oncoprotein fragment. The one or more oncoprotein fragments may be fragments of the same

oncoprotein, or may be fragments from different oncoproteins. In some cases, the polypeptide comprises more than one copy of the same fragment.

We provide, according to another example disclosed herein, a method of treatment of cancer, preferably metastatic cancer, in an individual suffering or suspected to be suffering from cancer. The method may comprise administering a therapeutically effective amount of a fragment of an intracellular oncoprotein to the individual. The intracellular oncoprotein may have been determined as being expressed in the cancer that the individual is suffering from.

The fragment may correspond to a region of the oncoprotein that comprises a mutation. That is, the fragment may comprise an oncogenic mutation. An advantage of targeting an oncogenic mutation, for example using a peptide that comprises a fragment of an oncoprotein that includes an oncogenic mutation, is that an immune response specific to a cancer may be generated, leaving normal cells unharmed. The patient may be administered a plurality of oncoprotein fragments. The plurality of oncoprotein fragments may be combined into one or more peptides that each comprises a plurality of different peptides, each optionally linked by a peptide linker. The fragments may be from the same oncoprotein, or different oncoproteins. The method may involve inducing the patient to produce anti-cancer antibodies, for example antibodies against the oncoprotein or oncogenic mutation of the oncoprotein fragment. The method may involve administering the fragment or peptide with an adjuvant.
The intracellular oncoprotein fragment may be such that it does not comprise an extracellular domain or a transmembrane domain. The intracellular oncoprotein fragment may comprise a cytoplasmic or nuclear oncoprotein fragment.

The intracellular oncoprotein fragment may comprise a portion of an Estrogen Receptor protein, for example ER (estrogen receptor), or a Hepatitis B virus protein such as HBV X-protein, PRL-3 protein, VHZ, Her2/neu or polyomavirus middle T (Py-mT) protein,, Kras, EGFR, B-raf, PI3KCA, beta-catenin, GNAS, Ret, EZH2 or GNAQ.

The cancer may be associated with or caused by over-expression of the intracellular oncoprotein of which an oncoprotein fragment is used for immunization. The cancer may be selected from the group consisting of: breast cancer, hepatocellular cancer such as hepatocellular carcinoma, ovarian cancer, liver cancer, pancreatic cancer, prostate cancer, gastric cancer, lung cancer, penis cancer, cervical cancer, brain cancer, esophageal cancer, bladder carcinoma, kidney renal cell carcinoma, ovary lymphoma and skin melanoma.

The cancer may comprise a metastatic cancer.

The method may be such that the number of metastatic tumours in a treated individual is reduced by at least 50%, at least 60%, at least 70%, at least 80% or at least 90%, compared to an untreated individual. In some cases, metastasis of the cancer is entirely prevented.

As a further example disclosed herein, there is provided an intracellular oncoprotein fragment, or a variant, homologue, or derivative thereof, for use in a method of prevention of cancer, such as metastatic cancer, the method comprising administering a prophylactically effective amount of an intracellular oncoprotein fragment, variant, homologue, or derivative thereof, to an individual suffering or suspected to be suffering from cancer.

We provide, according to another example disclosed herein, a pharmaceutical composition comprising (a) an intracellular oncoprotein fragment, or a variant, homologue orderivative thereof, together with a pharmaceutically acceptable excipient, diluent or carrier.

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of chemistry, molecular biology, microbiology, recombinant DNA and immunology, which are within the capabilities of a person of ordinary skill in the art. Such techniques are explained in the literature. See, for example, J. Sambrook, E. F. Fritsch, and T. Maniatis, 1989, Molecular Cloning: A Laboratory Manual, Second Edition, Books 1 -3, Cold Spring Harbor Laboratory Press; Ausubel, F. M. et al. (1995 and periodic supplements; Current Protocols in Molecular Biology, ch. 9, 13, and 16, John Wiley & Sons, New York, N.Y.); B. Roe, J. Crabtree, and A. Kahn, 1996, DNA Isolation and Sequencing: Essential Techniques, John Wiley & Sons; J. M. Polak and James O'D. McGee, 1990, In Situ Hybridization: Principles and Practice; Oxford University Press; M. J. Gait (Editor), 1984, Oligonucleotide Synthesis: A Practical Approach, Irl Press; D. M. J. Lilley and J. E. Dahlberg, 1992, Methods of Enzymology: DNA Structure Part A: Synthesis and Physical Analysis ofDNA Methods in Enzymology, Academic Press; Using Antibodies: A Laboratory Manual: Portable Protocol NO. I by Edward Harlow, David Lane, Ed Harlow (1999, Cold Spring Harbor Laboratory Press, ISBN 0-87969- 544-7); Antibodies: A Laboratory Manual by Ed Harlow (Editor), David Lane (Editor) (1988, Cold Spring Harbor Laboratory Press, ISBN 0-87969-314-2), 1855. Handbook of Drug Screening, edited by Ramakrishna Seethala, Prabhavathi B. Fernandes (2001 , New York, NY, Marcel Dekker, ISBN

0-8247-0562-9); and Lab Ref: A Handbook of Recipes, Reagents, and Other Reference Tools for Use at the Bench, Edited Jane Roskams and Linda Rodgers, 2002, Cold Spring Harbor Laboratory, ISBN 0-87969-630-3. Each of these general texts is herein incorporated by reference.

Everything disclosed herein includes the combination of the aspects and preferred features described except where such a combination is clearly impermissible or expressly avoided.

The section headings used herein are for organizational purposes only and are not to be construed as limiting the subject matter described.

Examples described herein will now be illustrated, by way of example, with reference to the accompanying figures. Further examples will be apparent to those skilled in the art. All documents mentioned in this text are incorporated herein by reference.

BRIEF DESCRIPTION OF THE DRAWINGS

Embodiments and experiments illustrating the principles of the invention will now be discussed with reference to the accompanying figures in which:

Figure 1 is a diagram showing that PRL-3 mAb effectively inhibits the formation of metastatic tumors formed by PRL-3 expressing cancer cells. A, Total cell lysates were prepared from B16F0 and B16F10 melanoma cancer cells and endogenous PRL-3 protein expression was determined using immunoblot. GAPDH was used as a loading control. B, C57BL6 mice were injected with lxl O^6 B16F0 cells (on day 1) followed by PRL-3 mouse antibody (mAb, clone #318) treatment with therapeutic plan. C. On day 17-20, tumors were frequently found in the lung, liver, adrenal gland, kidney, and bone in untreated mice (n=5), while the PRL-3 mAb eliminates the formation of tumors in most tissues of treated mice (n=5). D, The body weight of PRL-3 mAb ‘treated’ B 16F0 recipient mice constantly increased. Throughout the duration of the experiment. E, Parallel experiments were performed with non-PRL-3 expressing BI 6F10 melanoma cancer cell line and no therapeutic outcome was obtained with PRL-3 mAb therapy. F, The body weights of both treated and untreated B16F10 receipt mice constantly decreased over the duration of the experiment.

Figure 2 is a diagram showing that B-cells are important in mediating the antibody therapeutic event. A. Kaplan-Meier survival curves were used to compare the 'treated' and 'untreated' groups of B-cell deficient muMT mice injected with B16F0 cells. No difference in median survival rate for untreated (18.5 days) and treated (19 days) groups was evident. B. Kaplan-Meier analysis of wild-type C57BL6 mice injected with B16F0 had a statistically significant difference (.P=0.0002) in median, survival between untreated (17 days) and treated (21.5 days) groups. C. Kaplan-Meier analysis of wild-type C57BL6 mice C57BL6

mice injected with B16F10 had no statistically significant difference between untreated (16.5 days) and treated (17.5 days). D. The results between treated and untreated mice were summarized. The numbers of ineffective (or tumor-bearing) mice were placed at the numerator and the total numbers of mice were placed at the denominator. Fraction indicated at the Y-axis represents percentage of tumor- bearing mice verse total mice (n) in each group. Column represents type of mice indicated at the X-axis.

Figure 3 is a diagram showing that mT Ab effectively inhibits the formation of breast tumors expressing mT oncoprotein. A, Genotyping was used to identify transgenic MMTV-PymT females. Mice #2, #6, #7, and #10 are heterozygote transgenic females (+/-). B, #2 and #6 transgenic mice remained untreated as controls to monitor the progression of mammary gland tumor progression. #7 and #10 transgenic mice received mT antibody i.v. injection. At the end of the experiment (3-month), all mice were sacrificed. The sizes and weights of the breast tissues (tumors) were measured. The 5 pairs of mammary glands were indicated and showed as 4 groups in respective lower panel. Bar: 1 .5 cm. C, The morphology of the whole mount breast tissues was examined in breasts from untreated (a), treated (b), and wild type (c). E. Kaplan-Meier survival curves were used to compare 'treated' and 'untreated' group of MMTV-PymT heterozygous transgenic females. F. Fraction indicated at the Y-axis represents percentage of tumor-bearing mice verse total mice (n) in each group. Column represents type of mice indicated at the X-axis.

Figure 4 is a diagram showing that immunization of MMTV-PyMT mice with mT antigen could prevent the formation of breast tumors that express mT intracellular protein. A, a therapeutic plan of vaccination with mT antigen. B-D. MMTV-PymT heterozygous transgenic mice (TG) were used for the experiment. TG females (#43, #44, #45) were immunized with GST protein. By ELISA analysis, these mice had no mT antibody in their sera and showed dramatic progression of mammary gland tumors. In contrast, TG females (#37, #40, #41) were immunized with GST-mT fusion protein; these mice had high levels of mT antibodies in their sera and showed significantly repression of mammary gland tumors. The 10 mammary glands were shown as 4 groups at the bottom of each panel. Top two groups of breasts consist of the top three pairs of mammary glands, while bottom two groups represent the bottom two pairs of mammary glands. Weights of total 10 mammary glands were measured. Wild type mice (n=3) were used as controls. Bar: 1.5cm. E, Kaplan-Meier survival curves were compared between 'immunized' and 'unimmunized' MMTV-PymT mice, with median survival time (p=0.0004) clearly longer in mT- immunized group (19.5-week) versus unimmunized group (14.5-weeks). F. Summary of results between immunized and

unimmunized mice. N stands for number of mice in each group; fraction in each column represents proportion of tumor-bearing MMTV-PymT mice verse total TG females used.

Figure 5 is a diagram showing that PRL-3-, EGFP- or mT-Abs effectively inhibit the formation of PRL-3, EGFP, and mT expressing tumors.. A, Total cell lysates were prepared from EGFP-FO and EGFP-FI O cancer cells. Exogenous EGFP protein was detected in both cell lines. PRL-3 was only expressed in EGFP-FO (but low in EGFP-FIO). GAPDH was used as the loading control. B, EGFP-FO and EGFP-FI O stable populations of pooled cells were showed to be heterogeneous expressing of EGFP. C. a therapeutic plan showed C57BL6 mice which were injected with 10^6 cancer cells via tail veins (on day 1). Treated mice were i.v. injected with EGFP mAb. Untreated mice were i.v. injected with PBS. D-E, Organs were harvested, examined and imaged on 17-20 day. Tumors were found in lung, bone, adrenal, and ovary in the untreated group, but greatly reduced in treated mice. Note EGFP-Ab specifically inhibits EGFP-tumors, but not non-EGFP-tumors in lungs (panels, e). F. a. Untreated FVB/N -MMTV-PymT, b. mT mAb effectively inhibits the formation of breast tumors in transgenic MMTV-PymT mice c. non- transgenic mice show normal sizes of breast tissues as controls. G. Summary of results between untreated and treated mice with antibodies against intracellular targets. The numbers of ineffective (or tumor-bearing) mice were placed at the numerator and the total numbers of mice were placed at the denominator. 146 mice were tested by three different antibody therapies against their respective intracellular targets. N stands for number of mice in each group; fraction in each column represents proportion of mice bearing heavy tumors.

Figure 6 is a diagram showing that mice immunized with PRL-3, EGFP, or mT intracellular antigens could prevent PRL-3, EGFP and mT expressing tumors. A, a therapeutic plan of vaccination with respective antigen. B, Un-immunized, PRL-3-immunized or GFP-immunized mice were then challenged with either 1 million EGFP-FO or EGFP-F10 cancer cells via tail veins. All the organs were examined 3 weeks post-cancer cell injection and photographed with fluorescent microscopy to show morphologies of metastatic tumors. Black tumors represent ηόη-EGFP expressing melanoma cells, whereas green tumors represent those expressing EGFP. C, An overview of MMTV-PymT heterozygous transgenic females: a. Unimmunized MMTV- PymT mice b. mT-immunized mice showed reduction breast tumors. D, Summary of results between unimmunized and immunized mice with intracellular antigens. 84 mice were used in vaccination experiment. N stands for number of mice in each group; fraction in each column represents proportion of mice bearing heavy tumors.

Figure 7 is a diagram showing that generation of PRL-3 chimeric mAb specifically reacts with PRL-3 antigen. A. A schematic diagram outlining the major steps of chimeric mAb construction. B. PRL-3 chimeric mAb recognizes EGFP-PRL-3 overexpressed in DLD-I human colorectal cancer cells by indirect immunofluorescence: a, distribution of EGFP-PRL-3 in fixed DLD-I cells (green); b, PRL-3 chimeric antibody and anti-human Texas-Red to reveal the binding to PRL-3 protein; c, merged image. Bar: 20 μηι. C. Cell lysate derived from DLD-I cells overexpressing EGFP-PRL-3, and lysates derived from CHO stable cell lines overexpressing myc-PRL-3, myc-PRL-1 , and myc-PRL-2 were analyzed by western blot. PRL-3 chimeric antibody specifically recognized EGFP-PRL-3 (48 kDa) and myc-PRL-3 (20kDa), but does not react with myc-PRL-1 and myc-PRL-2.

Figure 8 is a diagram showing that PRL-3 chimeric antibody effectively inhibits the formation of metastatic tumors formed by Bl 6F0 cells that express endogenous PRL-3 A. Total cell lysates were prepared from F0 and F10 melanoma cells and analyzed by immunoblot. Abundant endogenous PRL-3 protein was detected in F0 but was almost undetectable in Fl 0 cells. B. On day 1 , nude mice (n=27) were injected with I xl O^6 F0 cells via tail vein, followed by two intravenous administrations of the PRL-3 chimeric mAb per week (day 3, 6, 9, 12, 15). C. At the end of the experiment (day 17), mice were photographed and tissues were dissected. Metastatic tumors were found in the adrenal gland, liver, bone, and abdomen in untreated mice (left), but not in treated mice (right). D. nude mice (n=22) were injected with lxl 0^6 F10 cells for the experiment. At the end of the experiment (day 17), dozens of lung metastatic tumors were found both in untreated (top panel) and treated mice (bottom panel). E. Kaplan-Meier survival curves for 'treated' and 'untreated F10 recipients were shown.

Figure 9 is a diagram showing that PRL-3 chimeric mAb inhibits the formation of metastatic tumors formed by A2780 cells and HCT-1 16 cells that express endogenous PRL-3. A. Total cell lysates were prepared from HCT 1 16-luc2, HCT-1 16, A2780, and NCI-H460 cancer cell lines. Endogenous PRL-3 protein was detected in HCT1 \ 6-luc2, HCT-1 16, and A2780 cells, but not in H460. B: On day 1 , nude mice (n=6) were injected with lxl O^6 HCT-1 \ 6-luc2 cancer cells and subsequently administered with PRL-3 chimeric mAb (n=3, treated) or PBS (n=3, untreated) on day 3, followed by two intravenous administrations of the PRL-3 chimeric mAb for 7-week. Both cancer cells and antibodies were injected via tail vein. I VIS® Imaging System was used to track and monitor tumor development in vivo at week 7. C. On day 1 , nude mice were injected with I xl O^6 cancer cells and treated as described in B. Paired experiments (untreated/treated) were terminated when mice appeared very sick. Experiment durations are indicated on the top of each panel. D. Summary of therapeutic results from

mouse PRL-3 (R3-mAb) or chimeric PRL-3 (R3- hAb) antibody therapies in nude mice injected with three human cancer cell lines. Percentages of tumor-bearing mice were averaged from each group (n=numbers of mice) and indicated at the Y- axis. 101 mice in total were used in this experiment.

Figure 10 is a diagram showing that B-cells are important in mediating therapeutic efficacy of PRL-3 chimeric antibody. A. Kaplan-Meier survival curves of 'treated' and 'untreated' HCT-1 16-injected nude and scid mice (a, b). Kaplan-Meier survival curves of 'treated' and 'untreated' Bl 6F0-injected nude and scid mice (c, d). B. Summary of results for therapeutic experiments in nude and scid mice (a-d). Percentages of tumor-bearing mice were averaged from each group (n=numbers of mice) and indicated at the Y-axis. 151 mice in total were used in this experiment.

Figure 11 is a diagram showing that A. PRL-3 is not detectable at the cell surface, a. Peak-shift was observed between A431 cells incubated with EGF-receptor antibody (black line) compared to A431 cells without incubation of EGF-receptor antibody (red line), b. Peak-shift was not observed between FO cells incubated with PRL-3 mAb (black line) and without Ab (red line). c. Peak-shift was not observed between F10 cells incubated with PRL-3 mAb (black line) and without Ab (red line). B Labeled antibodies were intravenously injected 1 hr before I VIS live imaging based working models: a. Treated mice. Early delivery of the antibody will persistently attack cancer cells to prevent them from further progression, resulting in micro-metastases in open stages, as such, fluorescent labeled PRL-3 antibody can access and bind to metastatic lung tumors in 'treated mice'. We observed strong fluorescent labeled lung in treated mice, b, Untreated mice. The uncontrolled cancer cells rapidly multiplied. They freely developed into macro-metastases and established defense territory, some kind of tumor microenvironment, which makes less accessible to antibody and immune system, as consequences, fluorescent labeled PRL- 3 antibodies were not able to label metastatic lung tumors in 'untreated mice'. We therefore are unable to observe fluorescent labeled metastatic lungs in untreated mice. In H&E stain section, black arrows showed a clear boundary ('fence') of the metastatic lung tumor from untreated B 16F0 recipients. Scale bar: 200 μιη. Red arrows indicated blood vessels.

Figure 12 is a diagram showing that PRL-3 protein is upregulated in lung cancers and AML. A. Representative IHC staining of PRL-3 expression in lung squamous cell carcinoma. B. lung adenocarcinoma. Scale bar: 100 μιη. C. The percentages of PRL-3-positive lung cancers as detected with immunohistochemistry (IHC) are summarized, grouped according

to cancer subtypes. D. AML bone marrow samples were examined by IHC, 24 out of 69 (35%) showed PRL-3 expression. Three selected images were shown.

Figure 13 is a diagram showing that NK cells in Innate Immune system are involved in the therapy. Nude mice were injected (n=2) and un-injected (n=2) with GM1 antibody 24 hrs before the experiment. On day 1 , all mice were injected with Ixl 0^6 F0 cells via tail vein, followed by two intravenous administrations of the PRL-3 chimeric mAb per week (day 3, 6, 9, 12, 15) in GM1 injected mice. On day 18, the therapeutic efficacy was examined. GMI injected nude mice showed more severe tumors (in black)-bearing burden in lung, liver, adrenal, testis, and bone than GM 1 un-injected mice, regardless PRL-3 mAb treated or untreated.

Figure 14 is a diagram showing that the outcome of the treatment is highly associated with tissue expression patterns of the targets. A, B. The protein expression patterns of PRL-3 and PRL- 2 in normal mouse tissues by western blot, GAPDH was used as a loading control. C. HCT-1 16 (PRL-2 positive cell line) recipients fail to respond to PRL-2 antibody (the antibody also cross- reacts with PRL-1) therapy, most likely due to the fact that PRL-2 is ubiquitously expressed in most of mouse tissues that we have examined.

Figure 15 shows the results of immunization of C57BL6 mice with VHZ and monitoring tumor development. 8-week old C57BL6 mice were immunized by intraperitoneal injection with a total volume of 200 ml Freund's Adjuvant: 20 mg of VHZ antigen in 100 ml saline mixed with 100 ml of complete adjuvant (Cat#77140, Pierce). The next two immunizations were injected with a total volume of 200 ml adjuvant: 20 mg of VHZ antigen in 100 ml saline mixed with 100 ml of incomplete adjuvant (Cat#77145, Pierce). The second and third injections were administrated every 2 weeks, 100-200 ml of tail bleed was collected in a heparin-coated capillary tube. Plasma was prepared from blood sample and antibody titer was measured by EL1SA. The detailed steps of ELISA were described previously. Mice with high titers of VHZ antibodies in their sera were selected and lateral tail vein injected with 1 x106 VHZ-expressing cancer cells. Unvaccinated mice serve as a negative control in this study.

Figure 16 shows sequences of fragments of Her2/neu oncoprotein and mT oncoprotein used in Examples 40 and 41. The Her2a intracellular peptide is unique for Her2, and does not cross react with EGFR (Her1).

Figure 17: Mice immunized with Her2 fragments could inhibit the formation of Her2-expressing tumors formed by B16F0 melanoma cancer cells. (A) A vaccination plan with Her2 intra-, extra-, and C-terminal fragments. (B) At week 16, unimmunized, Her2 intra–immunized, or Her2 extra–immunized mice, and Her2 C-terminal immunized mice were then challenged with 1 million B16F0 cancer cells via tail veins. All the organs were examined at ~17 days after cancer cell injection and photographed to show morphologies of metastatic tumors. Black tumors represent B16F0 melanoma cells. (C, upper) ELISA analysis was confirmed to show that Her2 fragments immunized (but not unimmunized) mice had high levels of Her2 antibodies in their sera. (C, lower). A total numbers of lung tumors from each mouse were counted and shown. N=3 in each group.

Figure 18: Mice immunized with Her2a short peptide or fragment could inhibit the formation of Her2-expressing tumors formed by B16F0 melanoma cancer cells. (A) A vaccination plan with Her2a short peptide, Her2 extra fragment. (B) At week 16, Her2 short peptide immunized, Her2 extra–immunized, and unimmunized mice, were then challenged with 1 million B16F0 cancer cells via tail veins. All the organs were examined at ~17 days after cancer cell injection and photographed to show morphologies of metastatic tumors. Black tumors represent B16F0 melanoma cells. (C, upper) ELISA analysis was confirmed to show that Her2 short peptide immunized, Her2 extra fragment immunized (but not unimmunized) mice had high levels of Her2 antibodies in their sera. (C, lower). A total numbers of lung tumors from each mouse were shown. N=3 in each group. Her2jp' is equivalent to Her2-extra.

Figure 19: MMTV-PymT TG young females vaccinated with mT peptide reduce the formation of mammary tumors. (A) A therapeutic plan of vaccination with mT peptide. (B) mT-peptide immunized mice showed marked repression of tumors compared to GST-immunized mice (unrelated immunized mice as negative control). (C, upper) the 10 mammary glands were shown as four groups at the bottom of each panel. (C, lower) An average weight of mammary glands from each mouse was shown, using WT mice as controls for normal sizes of mammary glands.

Figure 20: shows sequences for Estrogen Receptor (ER) and HBV-X protein. Estrogen Receptor peptides are used in the experiments of which the results are detailed in Figure 21. ER-red and ER-purple represent two regions (or fragments) indicated with red and purple sequences.

Figure 21: MMTV-PymT TG young females vaccinated with Estrogen Receptor peptides (ER-purple and ER-red) to reduce the burden of mammary tumors. (A) A therapeutic plan of vaccination with Estrogen Receptor peptides (ER-purple or ER-red). (B) To confirm that targeting tumors are expressing Estrogen Receptor for suitable treatment, a western blot was performed to show that Estrogen Receptor protein is expressing in tumor (T. breast and T. lung) (lane 4-5) tissues from MMTV-PymT TG adult females. (C) For a negative control, immunized mice with GST antigen, which is not expressing in the target tumors, immunized with ER-purple peptide and ER-red peptide, these two group of peptides showed marked repression of tumors compared to GST immunized control mice.

The 10 mammary glands were shown as four tissue groups at the bottom of each panel. Scale bars, 1.0 cm. (D) An average weight of mammary glands from each group of mice was shown together with wildtype (wt) mice as a control for normal breast sizes (mean ± SD).

Figure 22: The outcome of the antibody treatment is highly associated with tissue expression patterns of the targets. A. The protein expression patterns of Estrogen Receptor (ERa) in normal mouse tissues together with tumor (T. breast and T. lung) tissues were examined by western blot using ERa antibody, GAPDH was used as a loading control. B. A

therapeutic schedule of estrogen receptor 1 (ERa) antibody-treatment C. Student's t test

analysis revealed a significant difference in average breast tumor weight between untreated

and ERa-treated groups ($P = 0.0163$) (mean ± SD) $P < 0.05$ was regarded statistically

significant. D. At the end (13-week old), all mice were killed and examined. The sizes and

weights of the breast tissues (tumors) were measured. The five pairs of mammary glands were indicated and shown as four groups in respective lower panels.

Figure 23: Three possible mechanisms for antibody targeting intracellular antigens A. Antibodies may potentially enter PRL-3 expressing cells to target intracellular PRL-3 and neutralize its function. B. Some of the intracellular PRL-3 may be externalized and displayed on the surface of cancer cells by unconventional secretion. C. Proteolytic fragments of intracellular PRL-3 may be presented by MHC class I molecules to attract Cytotoxic T cells.

Figure 24: Vaccination of PRL-3 peptide (EVTYDKTPLEKDGITV**GGSG**DPHTHKTRC-KLH) could block tumors formed by PRL-3 expressing cancer cell line (B16F0) in C57BL6 mice.

Figure 25: vaccination of PRL-3 peptide could reduced tumours formed by PRL-3 expressing cells (B16F0) but not by PRL-3 non-expressing (B16F10) in C57BL6 mice.

Figure 26: Balb/c mice immunized with Ras mutated peptide (CMTEYKLVVVGA**D**GVGKSALT) could block tumours formed by CT-26 cancer cells expressing Ras (G12D) mutation.

Figure 27: A. peptides containing oncogenic mutations (mutation indicated in bold). **B.** other oncogenic mutations

Figure 28. Oncogenic mutations identified from http://share.gene.com/mutation_classification/cancer.variants.txt

DETAILED DESCRIPTION

The details of one or more embodiments of the invention are set forth in the accompanying description below including specific details of the best mode contemplated by the inventors for carrying out the invention, by way of example. It will be apparent to one skilled in the art that the present invention may be practiced without limitation to these specific details.

To explore the possibility of targeting intracellular proteins, in this study, we have chosen to focus on three representative intracellular targets for anticancer therapies in animal models.

We select a cancer associated-PRL-3 intracellular phosphatase as a target for the PRL-3 antibody therapy. PRL-3 is one of the three members (PRL-1 , -2, and -3) in the PRL (phosphatase of regenerating liver) family which was identified in 1994 and 1998[5,6]. The three PRLs form a subgroup of the protein tyrosine phosphatase (PTP) family[7].

PRL-3 was first linked to colorectal cancer metastasis in 2001. Subsequently, up- regulation of individual PRLs-PTPs was reported to be correlated with numerous types of advanced human metastatic cancers when compared with their normal counterparts[9].

The PRL phosphatases represent an intriguing group of proteins being validated as biomarkers and therapeutic targets in human cancers[10]. PRLs are intracellular C-terminally prenylated phosphatases, while mutant forms of PRLs that lack the prenylation signal are often localized in nuclei[11,12].

The localization of PRL- 1 and PRL-3 to the inner leaflet of the plasma membrane and early endosomes has been revealed by EM immunogold labeling[13]. Targeting PRLs by exogenous reagents to ablate PRLs-cancer cells requires their penetration into cells and is a challenging task.

We chose the cytosolic enhanced green fluorescent protein (EGFP), a popular reporter protein originally isolated from the jellyfish Aequorea victoria that fluoresces green when exposed to blue light. EGFP is an intracellular protein that sometimes enters the nucleus. As a neutral exogenous protein, EGFP serves as an artificial 'cancer cell specific intracellular protein' in EGFP-B 16F0 and EGFP-B16F 10 melanoma cells.

These EGFP-overexpressing melanoma cells will then be targeted by EGFP mAb. We expect that EGFP antibody should have less undesired side effect in the animal model as EGFP is not expressed in the host tissues.

We selected a well known oncoprotein, the polyomavirus middle T (mT) intracellular kinase, as the third target[14]. We used the mammary cancer model of Py-mT transgenic heterozygous females (+/-) carrying the mT oncogene under the transcriptional control of the mouse mammary tumor virus promoter/enhancer. The transgenic mice have been widely used as excellent mouse tumor models for decades in the cancer research community to assess the relative contribution of the metastatic mammary tumor phenotype[15,16].

We selected the well known oncoprotein Her2 (also known as ErbB-2). We injected C57BL6 mice with B16F0 melanoma cells as a model of cancer metastasis, and investigated whether Her2 fragments could inhibit the formation of Her2- expressing tumors.

Fragment and peptide vaccination is using shorter sequences designed and selected from whole oncoprotein sequences as antigens to stimulate host immune system to produce antibody against that specific fragment or peptide derived from the whole oncoproteins, which is expressing in the tumor cells. The 'fragment and peptide' vaccination is expected to be more specific and less cross reacting with their homology members in the same family. Compared to whole oncoprotein vaccine, fragment or peptide vaccine therefore has less side effect and is more specific. Sequences located in the whole oncoproteins, which cross react with other family members can be deleted and avoided. The resulting unique peptide is very specific to that target and does not react with its related target sequences.
To obtain a better understanding and extend our previous promising pre-clinical data[17] to future clinical applications in human, herein, we performed the entire antibody therapy study in wild type animal models to reveal a hitherto unrecognized concept that antibody therapy can be used for targeting intracellular-oncoproteins.

Oncogenic mutations are common in contributing to human cancers. Since modern technology can easily identify patients whose tumors are associated or caused by a specific oncogenic mutation, we could then design peptides corresponding to these oncogenic mutations. Vaccine with this specific mutant peptide could potentially trigger patient's immune system against the mutant proteins, the patients will produce suitable antibody "drugs" and cytotoxic T lymphocytes (CTLs) for their own anticancer therapeutics. Since vaccine could elicit long-lived immunity, when the correct antigens (peptide or gene fragment) were used to trigger patients' immune system to make their own antibody and CTLs against the correct mutated targets in ablating their tumors, they could achieve outcomes similar to antibody therapeutics.

Since dendritic cells (DCs) are antigen-presenting cells (APCs) which play a critical role in the regulation of the adaptive immune response, large numbers of DCs could be isolated and generated from CD34+ bone marrow precursors or from CD14+ monocytes *in vitro.* In vitro vaccination will educated the APCs to be able to present the peptide antigens at their cell surface. Reintroducing the educated APCs back to the same host to trigger B and T cell responses to these peptide antigens and produce antibody and CTLs to ablate cancer cells that express the specific antigens.

Alternatively, peptides synthesised to contain target antigens, optionally including oncogenic mutations, may be administered to the patient, to educate their immune system in vivo.

The Examples demonstrate that each of the above intracellular proteins can be targeted by antibodies against the cognate antigens.

If an antibody can recognize its intracellular antigen, it may be that an intracellular antigen could potentially be used for vaccination to stimulate host immune system and produce antibodies against it. This prompted us to explore vaccination with intracellular oncoproteins against tumors which express specific intracellular antigens.

Vaccination is inexpensive yet effective in stimulating host immunity. Compared with extracellular antigens, intracellular proteins used in vaccination receive much less attention because of their intracellular location. The main objective of this vaccination is to artificially activate the immune system against a particular intracellular protein to prime the antibody in some individuals against certain malignant cells expressing that intracellular protein thus preventing tumor formation in the future.

Next, we used the same three intracellular antigens (PRL-3, EGFP, and mT) to perform vaccination in C57BL6 wild type mice. We obtained interesting results, as shown in the Examples, that mice immunized with an intracellular tumor antigen are able to eradicate these cancer cells expressed that specific tumor antigen to reduce the formations of tumors. More importantly, the data suggest the possibility of vaccination to prevent disease progression for individuals who are genetically at high risk in developing certain cancers caused by known intracellular oncoproteins. Such specific active immunotherapy of cancer, if successful and effective have clear advantages over passive immunotherapy.

In this study, we used 194 mice (Figure 6) for both antibody therapies and vaccinations with intracellular proteins for anticancer. Cancer research is rapidly moving towards individualized cancer therapy, as oncologists begin to use tailoring treatment strategies to individual patients. Antibody therapy and Vaccination may fulfill the promise of personalized treatment and could be the future of individualized medicine because we all have the ability to make antibodies for our own recovery.

Currently, vaccines are mainly used to prevent bacterial and viral infections and to target a few extracellular proteins (receptors) on cancer cells. Our data indicate that strategies targeting intracellular oncoproteins hold enormous promise for cancer prevention in the future.

Oncoproteins as described herein are proteins involved in the regulation or synthesis of proteins linked to tumorigenic cell growth. Oncoproteins may be oncogenic polypeptides, involved in the transformation of normal cells into cancer cells. Oncoproteins may have higher expression in tumor cells than in normal cells. The oncoproteins are intracellular, meaning that they are located inside the cell, for example in the nucleus or cytoplasm, or attached to the intracellular surface of the cell membrane. Preferably, the oncoproteins are self-antigens, meaning that they are proteins normally found in the animal, and form part of the protein population expressed from the genome of the animal, and are not heterologous to that animal, such as viral proteins.

Alternatively, the intracellular oncoprotein may be an oncoprotein that has an intracellular region. For example it may be a membrane anchored protein that has a region which extends into the cytoplasm.

Oncoprotein can be a non-self-antigen, such as viral protein expressed by infected cells. In some cases the intracellular oncoprotein is not derived from a microorganism. For example, is not a viral oncoprotein, or is not a bacterial oncoprotein, or is not a fungal oncoprotein. In some cases the oncoprotein is not an HPV oncoprotein. Preferably, the oncoprotein is a self-antigen. One potential advantage of using intracellular self-antigens is that they may have a better chance of provoking an immune response than extracellular self-antigens because immune cells targeting extracellular self-antigens are generally eliminated during development.

PREVENTION OF CANCER BY VACCINATION WITH INTRACELLULAR ONCOPROTEINS

Previously therapeutic antibodies were thought to have access only to molecules that appear on the outer surface of cells. According to the expectation in the literature, targeting intracellular PRLs and such other intracellular oncoproteins with antibodies to ablate cancer cells and cancer metastasis has never been previously thought to be possible because of their intracellular location.

Accordingly, vaccination with intracellular oncoproteins to prevent cancer cell formation, cancer cell growth and cancer metastasis has never been previously thought to be possible. We have shown that this is not the case. Moreover, we have shown that vaccination with fragments of intracellular oncoproteins is also possible, and may have improved efficacy as compared with fragments of extracellular oncoproteins.

Vaccination with intracellular oncoproteins or fragments thereof, as disclosed herein, may result in stimulation of antibody production by the host. B cells may be stimulated to produce antibodies specific to the intracellular oncoprotein or fragment thereof. Thus, vaccination may result in stimulation of a B cell response specific to the intracellular oncoprotein or fragment thereof.

We therefore provide generally for fragments of cancer causing oncoproteins, which oncoproteins are intracellular in nature, as cancer vaccines, particularly vaccines against cancer metastasis, or to curb the spread or growth of a cancer.

We provide for use of an oncoprotein fragment, which oncoprotein is intracellular in nature, in a method of prevention of cancer including cancer metastases associated with cancer causing oncoproteins.

Vaccination may involve a single fragment from a particular oncoprotein (i.e. a homogenous population of identical peptides). Alternatively, peptides with more than one sequence may be used. For example, a mixture of peptides of sequence X and sequence Y. The peptides may be overlapping, or from different regions of an oncoprotein. In some cases the peptides do not span all, or substantially all, of the oncoprotein. That is to say that the mixture of peptides encompasses only part of the oncoprotein. In some cases, a mixture of different peptides from more than one oncoprotein may be used. Where a mixture of different peptides is used (either different peptides from the same oncoprotein, or peptides from different oncoproteins), these may be administered simultaneously or sequentially. Preferably, each of the peptides will be capable of inducing a host to produce antibodies against the intracellular oncoprotein.

According to the invention, only a single peptide may be used. Alternatively a mixture of 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more peptides may be used.

We provide for methods of preventing cancer including cancer metastases associated with cancer causing oncoproteins, which oncoproteins are intracellular in nature, by administration of a prophylactically effective amount of a fragment of the intracellular oncoprotein to a patient in need thereof.

We provide a pharmaceutical composition comprising an intracellular oncoprotein fragment, together with a pharmaceutically acceptable excipient, diluent or carrier. We provide for the

use of such a pharmaceutical composition for the prevention of cancer including cancer metastasis.

Since dendritic cells (DCs) are antigen-presenting cells (APCs) which play a critical role in the regulation of the adaptive immune response, we could isolate and generate large numbers of DCs from CD34+ bone marrow precursors or from CD14+ monocytes *in vitro*. In vitro vaccination will educate the APCs to be able to present the peptide antigens at their cell surface. The educated APCs may be reintroduced back to the same host to trigger B and T cell responses to these peptide antigens and produce antibody and CTL responses to ablate cancer cells that express the specific tumor antigens.

Alternatively, or additionally a patient's immune system may be directly educated using vaccination with an oncoprotein or fragment. Oncoproteins, antigens and/or mutations specific to a patient's cancer may be identified. This may include intracellular or extracellular oncoproteins, antigens and/or mutations. Peptides containing the relevant fragments of these oncoproteins, antigens and/or mutations may then be synthesised and used as a vaccine. Vaccination may involve the administration of the peptides in the presence or absence of one or more adjuvants. For example, Freund's complete adjuvant. Immunisation may involve administering the peptides at multiple times or locations. For example, at monthly intervals. To test the efficacy of immunisation, ELISA may be used to determine whether peptide specific antibodies have been produced by the patient.

Thus, one or more different peptides may be administered to a patient. Peptides containing fragments of particular oncoproteins, epitopes or oncogenic mutations which are known to be associated with, or a cause of, a patients cancer may be selected for administration to that peptide. Thus, the mixture of peptides chosen may be particular for (i.e. personalised to) an individual patient. A mixture of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 35, 40, 45, 50 or more peptides may be selected.

Thus, methods of treatment may involve the step of determining the presence of particular oncogenic mutations or oncoproteins in a sample obtained from a patient. The sample may be a tumor biopsy, or a blood or urine sample. Antigen specific to the tumor may be determined by any suitable method, for example Western blot, ELISA or PCT methods. Peptides may designed or chosen to correspond to oncoproteins or oncogenic mutations that are known to be present in the patient.

INTRACELLULAR ONCOPROTEINS

Intracellular oncoproteins and intracellular antigens are known in the art and may include, amongst others, any one or more of the following, or their variants, derivatives, homologues or fragments.

Suitable intracellular oncoproteins will be appreciable by the skilled person. Genes specifically up-regulated during tumor formation but poorly or not expressed in host tissues are particularly promising as tumor-specific targets. For cancers that show a genetic link, immunization of immune-competent young susceptible family members with an antigen (epitope-based peptide vaccine) that is associated with the familial cancer could prime the immune system against that oncoprotein. These endogenously stimulated antibodies could then potentially combat cancer cells expressing that particular oncoprotein. The results described herein suggest that antibody-based therapy and vaccination against cancer may be extended to a wider variety of intracellular oncoproteins as therapeutic targets. The whole class of intracellular oncoproteins previously thought to be un-targetable by therapeutic antibodies or vaccinations can now expand the scope for tailor-made cancer therapies as well as usher in a new era of cancer vaccines. We expect that one potential advantage of using intracellular self-antigens is that they may have a better chance of provoking an immune response than extracellular self-antigens because immune cells targeting extracellular self-antigens are generally eliminated during development. We found that compared to exogenously delivered antibodies, antigen-induced antibody therapy could achieve similar antitumor therapeutic efficacy. Because existing conventional clinical antibody therapy is costly, vaccination may be more useful and economical as a means of inducing high titers of antigen-induced antibodies. This concept of "cancer vaccination" is promising and challenging.

Estrogen Receptor

An oncoprotein useful in the present invention is Estrogen Receptor (ER). The oncoprotein may be human ER. It may comprise or consist of the protein sequence set out at P03372 (GI: 544257).

Estrogen Receptor and fragments thereof will be useful for treatment of breast cancer caused by, or associated with, overexpression of estrogen receptor (ER). Antibodies against ER or vaccination using ER oncoprotein or a fragment thereof could be used to prevent spreading. This is particularly useful to target ER positive breast cancer patients regardless of the expression of Her2 or other proteins.

Estrogen Receptor (ER) is a ligand-activated transcription factor composed of several domains important for hormone binding, DNA binding, and activation of transcription. Alternative splicing results in several ER mRNA transcripts, which differ primarily in their 5-prime untranslated regions. The translated receptors show less variability (see OMIM reference 133430).

Estrogen Receptor is a well known intracellular receptor (see Ross-Innes CS, Stark R, Holmes KA, et al. Cooperative interaction between retinoic acid receptor alpha and estrogen receptor in breast cancer. Genes Dev. 2010;24:171-182).

Hepatitis B Virus (HBV) Proteins

Hepatitis B proteins may be suitable for use in the invention. HBV exists as 8 genotypes.

One suitable HBV protein is the HBV X-protein. HBV X-protein is localized in the nucleus of infected cells. Most hepatocellular carcinoma (HCC) are associated with HBV infection. Accordingly, HBV proteins may be useful for treating HCC, antibody targeting viral protein to specifically destroy virally infected cells whilst leaving normal cells unharmed.

Protein sequences for HBV-X protein have been deposited at GenBank and are suitable for use in the present invention. For example, the term "HBV-X protein" may be used to refer to a protein comprising or consisting of the sequence set out at GenBank CBX46805.1 (GI: 310923520), or EMBL accession FR714506.1, or a protein encoded by a gene having a sequence as set out at Accession AB670311.1 (GI: 371919030).

PRL-3

The following text is adapted from OMIM entry 606449.

PRL-3 is also known as Protein-Tyrosine Phosphatase, Type 4A, 3; PTP4A3. The chromosomal location of PRL-3 is at gene map locus 8q24.3.

In the heart, protein kinases regulate contractility, ion transport, metabolism, and gene expression. Phosphatases, in addition to their role in dephosphorylation, are involved in cardiac hypertrophy and dysfunction.

By database searching and screening of a heart cDNA library, Matter et al. 2001 , Biochem. Biophys. Res. Commun. 283: 1061 -1068 identified a cDNA encoding PTP4A3, which they termed PRL3. The deduced PRL3 protein is 76% identical to PRLI (PTP4A1 ; 601585) and

96% identical to mouse Prl3. Northern blot analysis revealed expression of an approximately 2.3- kb PRL3 transcript predominantly in heart and skeletal muscle, with lower expression in pancreas. This expression pattern is distinct from the wider expression of PRL1 and PRL2 (PTP4A2; 601584). In situ hybridization analysis localized PRL3 expression to cardiomyocytes. Tris glycine gel analysis showed that PRL3 is expressed as a 22-kD protein. Functional and mutation analyses indicated that phosphate cleavage is dependent on cysl 04 of PRL3. Overexpression of PRL3 resulted in increased cell growth. Western blot analysis showed dephosphorylation of pl30cas (BCAR1 ; 602941) in response to angiotensin II (106150), suggesting a role for PRL3 in the modulation of intracellular calcium transients induced by angiotensin II.

To gain insights into the molecular basis for metastasis, Saha et al. 2001, Science 294: 1343-1346 compared the global gene expression profile of metastatic colorectal cancer with that of primary cancers, benign colorectal tumors, and normal colorectal epithelium. PRL3 was expressed at high levels in each of 18 cancer metastases studied but at lower levels in nonmetastatic tumors and normal colorectal epithelium. In 3 of 12 metastases examined, multiple copies of the PRL3 gene were found within a small amplicon located at chromosome 8q24.3. Saha et al. (2001) concluded that the PRL3 gene is important for colorectal cancer metastasis.

Using the Stanford G3 radiation hybrid panel and database sequence analysis, Saha et al. (2001) mapped the PRL3 gene to surrounding marker 145.20. The PRL3 gene is also tightly linked to marker SHGC-22154, which is located at 8q24.3, approximately 3 Mb from the 8q telomere.

Mouse and human PRL-3 proteins were described in detail in Li et al (2005), Clin Cancer Res; I 1 :2195-204.

PRL-3 Sequences

The methods and compositions described here make use of PRL-3 polypeptides, which are described in detail below. As used here, the term "PRL-3" is intended to refer to a sequence selected from the following.

Unigene	**Description**
AF041434.1	Homo sapiens potentially prenylated protein tyrosine phosphatase hPRL-3 mRNA, complete cds
	Homo sapiens protein tyrosine phosphatase type IVA, member 3

BT007303.1	mRNA, complete cds
AK 128380.1	Homo sapiens cDNA FLJ46523 fis, clone THYMU3034099
NM_007079.2	Homo sapiens protein tyrosine phosphatase type IVA, member 3 (PTP4A3), transcript variant 2, mRNA
AY819648.1	Homo sapiens HCV p7-transregulated protein 2 mRNA, complete cds
BC003105.1	Homo sapiens protein tyrosine phosphatase type IVA, member 3, mRNA (cDNA clone MGC: 1950 IMAGE:3357244), complete cds
NM_03261 1.1	Homo sapiens protein tyrosine phosphatase type IVA, member 3 (PTP4A3), transcript variant 1 , mRNA
AK31 1257.1	Homo sapiens cDN A, FLJ 18299
U87168.1 cds	Human protein tyrosine phosphatase homolog hPRL-R mRNA, partial
AJ276554.1	Homo sapiens mRNA for protein tyrosine phosphatase hPRL-3, short form
BC066043.1	Mus musculus protein tyrosine phosphatase 4a3, mRNA (cDNA clone MGC:90066 IMAGE:6415021), complete cds
AK190358.1	Mus musculus cDNA, clone:YI G0102103, strand:plus, reference:ENSEMBL:Mouse-Transcript-ENST:ENSMUST00000053232, based on BLAT search
CT010215.1	Mus musculus full open reading frame cDNA clone RZPDo836H0950D for gene Ptp4a3, Protein tyrosine phosphatase 4a3; complete cds, incl. stopcodon
AK 147489.1	Mus musculus adult male brain UNDEFINED CELL LINE cDNA, RIKEN full-length enriched library, clone:M5C1053F14 productprotein tyrosine phosphatase 4a3, full insert sequence
AK172192.1	Mus musculus activated spleen cDNA, RIKEN full-length enriched library, clone:F830102P03 product:protein tyrosine phosphatase 4a3, full insert sequence
AK 143702.1	Mus musculus 6 days neonate spleen cDNA, RIKEN full-length enriched library, clone:F43001 1 C20 productprotein tyrosine phosphatase 4a3, full insert sequence
AF035645.1	Mus musculus potentially prenylated protein tyrosine phosphatase mPRL-3 (Prl3) mRNA, complete cds
NM 008975.2	Mus musculus protein tyrosine phosphatase 4a3 (Ptp4a3), mRNA
AKO 14601 .1	Mus musculus 0 day neonate skin cDNA, RIKEN full-length enriched library, clone:4632430E19 productprotein tyrosine phosphatase 4a3, full insert sequence

AK004562.	Mus musculus adult male lung cDNA, RIKEN full-length enriched library, clone: 1200003F10 productprotein tyrosine phosphatase 4a3, full insert sequence
AK003954.1	Mus musculus 18-day embryo whole body cDNA, RIKEN full-length enriched library, clone: 1110029E17 productprotein tyrosine phosphatase 4a3, full insert sequence
BC027445.1	Mus musculus protein tyrosine phosphatase 4a3, mRNA (cDNA clone MGC:36146 I AGE:4482106), complete cds

A "PRL-3 polypeptide" may comprise or consist of a human PRL-3 polypeptide, such as the sequence having Unigene accession number AF041434.1.

Homologues variants and derivatives thereof of any, some or all of these polypeptides are also included. For example, PRL-3 may include Unigene Accession Number BC066043.1.

VHZ

The methods and compositions described here make use of VHZ, which is described in detail below.

VHZ is also known as DUSP23, MOSP, LDP-3, DUSP25, FLJ20442 and RP1 1 -190A12.1

As used here, the term "VHZ" may refer to a polypeptide sequence having GenBank Accession number NP_060293.2, NP_081001.1, XP_341 157.1 , XP_001 170819.1 , XP_001 170835.1 , XP_545747.2, NP_001076078.1 , NP 00101 1371.1 , NP_783859.1 , NP_001034709.1 , XP_001480730.1 , XP_001 1 17253.1 or XP 001 1 17256.1.

A "VHZ polypeptide" may comprise or consist of a human VHZ polypeptide, such as the sequence having accession number NP 060293.

With regard to nucleic acid sequences, the terms "VHZ polynucleotide", "VHZ nucleotide" and "VHZ nucleic acid" may be used interchangeably, and should be understood to specifically include both cDNA and genomic VHZ sequences. These terms are also intended to include a nucleic acid sequence capable of encoding a VHZ polypeptide and/or a fragment, derivative, homologue or variant of this.

Where reference is made to a VHZ nucleic acid, this should be taken as a reference to any member of the VHZ family of nucleic acids. Of particular interest are VHZ nucleic acids

selected from the group consisting of: NM_017823.3, NM_026725.2, XM_341 156.3, XM_001 170819.1, XM_ 170835.1 , XM_545747.2, NM_001082609.1 , NM_00101 1371.1 , N _175732.1 , NM_001039620.1 , XM_001480680.1 , XM_001 1 17253.1 or XM 001 1 17256.1.

Also included are any one or more of the nucleic acid sequences set out as "Other VHZ nucleic acid sequences" below.

For example, the VHZ nucleic acid may comprise a human VHZ sequence having GenBank Accession Number NM 017823.3.

Her2

Her2/neu (also known as ErbB-2) stands for "human epidermal growth factor receptor 2" and is a protein giving higher aggressiveness in breast cancers. It is a member of the ErbB protein family, more commonly known as the epidermal growth factor receptor family. HER2/neu has also been designated as CD340 (cluster of differentiation 340) and p185. HER2 is a cell membrane surface-bound receptor tyrosine kinase and is normally involved in the signal transduction pathways leading to cell growth and differentiation.

As described herein, HER2 may refer to a polypeptide sequence selected from GenBank accession numbers NP_004439.2, NP_001005862.1, NP_001003817.1, AAI67147.1

A "Her2 polypeptide" as referred to herein may comprise or consist of a human HER2 polypeptide sequence, such as that of accession number P04626.1

Her2 polypeptides are described in US6333169 and EP1418235.

Other oncoproteins useful in the invention include EGFR (GenBank accession numbers CAA25240 (GI:119533), ADZ75461.1 (GI326467049)), SHP1 (GenBank accession numbers NP002822.2 (GI: 18104989), NP536858.1 (GI: 18104991), NP536859.1 (GI: 18104991)), Tiam (GenBank accession numbers NP003244.2 (GI: 115583670), AAA98443.1 (GI: 897557), Q13009.2 (GI: 152031709)), Myc (GenBank accession numbers AAA59886.1 (GI: 188975), AAA59887.1 (GI: 188977), CAA25015.2 (GI: 29839758), NP002458.2 (GI: 71774083)), Ras (GenBank accession number AAA34557.1 (GI: 171374)) and Runx-1 (GenBank accession number NP001079966.1 (GI: 148232064)).

Preferred intracellular oncoprotein targets include Androgen Receptor (AAA51772; GI: 178882), Her2 (P04626.1; GI:119533, targeting the intracellular domain of the receptor), Mutant Ras (GenBank accession number AAA34557.1;GI: 171374), Jak (AAA19626.1; GI:508731), FLT3 (ITD3; CAA81393.1; GI:406323), Runx1 (AAI36381.1; GI:223459612), Tiam, Gfi (NP_005254.2 GI:71037377), Cot (MAP3K8; CAG47079.1; GI:49457560), Myc, SHP1, FGF (CAA28027.1; GI:31362).

Other preferred oncoproteins include KRas (V-Ki-ras2 Kirsten rat sarcoma viral oncogene homologue; GenBank P01116.1 GI:131875); EGFR (Epidermal growth factor receptor; P00533.2 GI:2811086); BRaf (P15056.4 GI:50403720); PI3KCA (Phosphatidylinositol 3-kinase regulatory subunit alpha; P27986.2 GI:118572681); Beta-catenin (CAA61107.1 GI:860988); GNAS (guanine nucleotide binding protein; Q5JWF2.2 GI:116248089); Ret (CAA73131.1 GI:1946207) and EZH2 (histone-lysine N-methyltransferase; enhancer of zeste homolog 2; Q15910.2 GI:3334180).

Oncogenic mutations include those listed in enclosed figures 27 and 28, and obtainable from http://share.gene.com/mutation_classification/cancer.variants.txt

Oncogenic Mutations

An oncogenic mutation is a mutation in a gene that is associated with, or causative of, cancer. The mutation may be a substitution, deletion or addition of a single amino acid residue. Alternatively, it may affect more than one amino acid residue. A number of oncogenic mutations are known in the art. Certain oncogenic mutations are listed at http://share.gene.com/mutation_classification/cancer.variants.txt. Oncogenic mutations may be identified by comparing the sequence of a protein obtained from, or associated with, a cancer, with a homologous protein obtained from an individual who is not suffering from cancer. Methods for comparing protein sequences are well known in the art.

Protein Fragments and Peptides

Protein fragments or peptides (used interchangeably herein) according to the invention are small polypeptides, peptides or peptide mimetics based on or comprising a contiguous sequence of amino acid residues from an intracellular oncogenic protein or an intracellular portion of an oncogenic protein. The protein fragments and peptides do not comprise the entire oncogenic protein. Peptides may comprise or consist of a fragment of an oncoprotein which is normally located within the cell, or a fragment of an oncoprotein corresponding to a region of that oncoprotein which is normally located within the cell.

A protein fragment or peptide according to the present invention may have a maximum length of 150, 140, 130, 120, 110 or 100 amino acids and less than the full length of the corresponding oncoprotein. More preferably the maximum peptide length is 90 amino acids, still more preferably 80 amino acids, still more preferably 70 amino acids, still more preferably 60 amino acids, still more preferably 50 amino acids, still more preferably 40 amino acids, still more preferably the maximum length is chosen from one of 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10 or 9 amino acids. For example, a peptide may have a maximum length of 150, 140, 130, 120, 110, 100, 90, 80, 70, 60, 50, 40, 30, 25, 20 or 15 amino acids.

A protein fragment or peptide according to the present invention may have a minimum length of 7 amino acids. More preferably the minimum length is chosen from one of 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, or 25 amino acids.

A peptide according to the present invention may have any length between said minimum and maximum. Thus, for example, a peptide may have a length of from 5 to 100, from 7 to 100, 7 to 80, 7 to 60, 7 to 50, 7 to 40, 8 to 30, 10 to 25, 12 to 20, 9 to 15 amino acids, 8 to 11 amino acids, 9 to 11 amino acids, 9 to 13 amino acids or 9 to 14 amino acids. In particular, the peptide may have an amino acid length chosen from one of 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 72, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100 amino acids.

In some embodiments the protein fragment may be one of between 10 and 50, between 10 and 40, between 10 and 30, between 10 and 20, between 15 and 50, between 15 and 40, between 15 and 30, between 15 and 20, between 20 and 50, between 20 and 40, between 20 and 30, between 20 and 25, between 25 and 50, between 25 and 40, between 25 and 30, between 30 and 50, between 30 and 40, or between 40 and 50 amino acids in length.

The protein fragment or peptide may have a length anywhere between the said minimum and maximum length.

The protein fragment or peptide preferably comprises at least one epitope (optionally two or more epitopes) of the oncoprotein which is recognised by the immune system of the patient, and is preferably capable of stimulating the production of anti-oncoprotein antibodies in the patient.

In some embodiments the amino acid sequence of the fragment comprises or consists of a contiguous sequence of amino acids present in the corresponding full length oncoprotein.

A protein fragment or peptide may be designed to encompass an oncogenic mutation. That is to say, that the peptide is fragment of the region of the oncoprotein that includes a mutation. The mutation may be present in the oncoprotein as compared to the wild-type or non-oncogenic form of the protein. Thus, the peptide may have a sequence specific to an oncogenic form of a protein. The peptide may have 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more amino acids on one or both sides of the mutation. The peptide may have the same number of amino acids on each side of the mutation, or may have different numbers of amino acids on each side of the mutation.

Since modern technology can easily identify patients whose tumors are associated with, or caused by, a specific oncogenic mutation, we could then design peptides corresponding to these oncogenic mutations.

In other embodiments the fragment has at least 60% amino acid sequence identity to a contiguous sequence of amino acids present in the corresponding full length protein. More preferably, the degree of sequence identity is one of 65%, 70%, 75%, 80%, 85%, 87%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% identity.

The present invention incorporates derivatives and mimetics of a contiguous sequence of amino acids from an oncoprotein that forms a protein fragment.

Peptides may be synthesised to include a single epitope or single region of the oncoprotein, and may include an oncogenic mutation. Alternatively, peptides may include a combination of more than one region of an oncoprotein or more than one epitope. For example, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more regions or epitopes of the same or different oncoproteins may be combined in the same peptide. This may involve the use of short linker peptides to connect the more than one region or epitope. The linker may consist of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or more amino acids. Peptide derivatives include variants of a given sequence of amino acids and may include naturally occurring allelic variants and synthetic variants which have substantial amino acid sequence identity to the peptide sequence as identified in the wild type full length protein allergen.

Peptide derivatives may include those peptides having at least 60% amino acid sequence identity and which are capable of stimulating generation of anti-oncoprotein antibodies.

Peptide derivatives preferably differ from the corresponding amino acid sequence in the oncoprotein by less than 5 amino acids. More preferably, the number of different amino acids is 4 amino acids or less, 3 amino acids or less, 2 amino acids or less or only 1 amino acid.

Peptide derivatives may arise through natural variations or polymorphisms which may exist between the members of a protein family from which the peptide is derived. All such derivatives are included within the scope of the invention.

Peptide derivatives may result from natural or non-natural (e.g. synthetic) interventions leading to addition, replacement, deletion or modification of the amino acid sequence.

Conservative replacements and modifications which may be found in such polymorphisms may be between amino acids within the following groups:

(i) alanine, serine, threonine;
(ii) glutamic acid and aspartic acid;
(iii) arginine and leucine;
(iv) asparagine and glutamine;
(v) isoleucine, leucine and valine;
(vi) phenylalanine, tyrosine and tryptophan;
(vii) methionine and leucine;
(viii) cysteine and valine.

Peptide derivatives may also be provided by modifying an amino acid sequence from the oncoprotein, e.g. in order to resist degradation of the peptide.

Peptide Mimetics

The designing of mimetics to a known pharmaceutically active compound is a known approach to the development of pharmaceuticals based on a "lead" compound. This might be desirable where the active compound is difficult or expensive to synthesise or where it is unsuitable for a particular method of administration, e.g. some peptides may be unsuitable active agents for oral compositions as they tend to be quickly degraded by proteases in the alimentary canal. Mimetic design, synthesis and testing is generally used to avoid randomly screening large numbers of molecules for a target property.

There are several steps commonly taken in the design of a mimetic from a compound having a given target property. Firstly, the particular parts of the compound that are critical and/or important in determining the target property are determined. In the case of a peptide, this can be done by systematically varying the amino acid residues in the peptide, e.g. by substituting each residue in turn. These parts or residues constituting the active region of the compound are known as its "pharmacophore".

Once the pharmacophore has been found, its structure is modelled according to its physical properties, e.g. stereochemistry, bonding, size and/or charge, using data from a range of sources, e.g. spectroscopic techniques, X-ray diffraction data and NMR. Computational analysis, similarity mapping (which models the charge and/or volume of a pharmacophore, rather than the bonding between atoms) and other techniques can be used in this modelling process.

In a variant of this approach, the three-dimensional structure of the ligand and its binding partner are modelled. This can be especially useful where the ligand and/or binding partner change conformation on binding, allowing the model to take account of this in the design of the mimetic.

A template molecule is then selected onto which chemical groups which mimic the pharmacophore can be grafted. The template molecule and the chemical groups grafted on to it can conveniently be selected so that the mimetic is easy to synthesise, is likely to be pharmacologically acceptable, and does not degrade *in vivo*, while retaining the biological activity of the lead compound. The mimetic or mimetics found by this approach can then be screened to see whether they have the target property, or to what extent they exhibit it. Further optimisation or modification can then be carried out to arrive at one or more final mimetics for *in vivo* or clinical testing.

With regard to the present invention, a peptide mimetic is one form of peptide derivative. A method of identifying a peptide derivative capable of stimulating an immune response may comprise the step of modifying the peptide structure to produce a peptide mimetic. This peptide mimetic may optionally be subject to testing in an antibody production assat. This process of modification of the peptide or peptide mimetic and testing may be repeated a number of times, as desired, until a peptide having the desired effect, or level of effect, on anti-oncoprotein antibody proliferation is identified.

The modification steps employed may comprise truncating the peptide or peptide mimetic length (this may involve synthesising a peptide or peptide mimetic of shorter length), substitution of one or more amino acid residues or chemical groups, and/or chemically modifying the peptide or peptide mimetic to increase stability, resistance to degradation, transport across cell membranes and/or resistance to clearance from the body.

INTRACELLULAR ONCOPROTEIN EPITOPES

The anti-intracellular oncoprotein antibodies may specifically bind to an epitope on the intracellular oncoprotein.

Methods are known in the art to determine an epitope that is bound by a particular antibody. Such epitope mapping methods are described for example in Hanson et al., (2006). Respiratory Research, 7: 126. Furthermore, a skilled person will be able to generate antibodies and screen them for particular properties.

POLYPEPTIDES

A "polypeptide" refers to any peptide or protein comprising two or more amino acids joined to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres. "Polypeptide" refers to both short chains, commonly referred to as peptides, oligopeptides or oligomers, and to longer chains, generally referred to as proteins. Polypeptides may contain amino acids other than the 20 gene-encoded amino acids.

"Polypeptides" include amino acid sequences modified either by natural processes, such as post-translational processing, or by chemical modification techniques which are well known in the art. Such modifications are well described in basic texts and in more detailed monographs, as well as in a voluminous research literature. Modifications can occur anywhere in a polypeptide, including the peptide backbone, the amino acid side-chains and the amino or carboxyl termini. It will be appreciated that the same type of modification may be present in the same or varying degrees at several sites in a given polypeptide. Also, a given polypeptide may contain many types of modifications.

Polypeptides may be branched as a result of ubiquitination, and they may be cyclic, with or without branching. Cyclic, branched and branched cyclic polypeptides may result from posttranslation natural processes or may be made by synthetic methods. Modifications include acetylation, acylation, ADP-ribosylation, amidation, covalent attachment of flavin, covalent attachment of a heme moiety, covalent attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid or lipid derivative, covalent attachment of

phosphotidylinositol, cross-inking, cyclization, disulfide bond formation, demethylation, formation of covalent cross- inks, formation of cystine, formation of pyroglutamate, formylation, gamma-carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristoylation, oxidation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and ubiquitination. See, for instance, Proteins - Structure and Molecular Properties, 2nd Ed., T. E. Creighton, W. H. Freeman and Company, New York, 1993 and Wold, F., Posttranslational Protein Modifications: Perspectives and Prospects, pgs. 1 -12 in Posttranslational Covalent Modification of Proteins, B. C. Johnson, Ed., Academic Press, New York, 1983; Seifter et al., "Analysis for protein modifications and nonprotein cofactors", Meth Enzymol (1990) 182:626- 646 and Rattan et al, "Protein Synthesis: Posttranslational Modifications and Aging", Ann NY AcadSci (1992) 663:48-62.

The term "polypeptide" includes the various synthetic peptide variations known in the art, such as a retroinverso D peptides. The peptide may be an antigenic determinant and/or a T-cell epitope. The peptide may be immunogenic in vivo. The peptide may be capable of inducing neutralising antibodies in vivo.

As applied to intracellular oncoproteins, the resultant amino acid sequence may have one or more activities, such as biological activities in common with a intracellular oncoprotein polypeptide, for example a human intracellular oncoprotein. For example, a intracellular oncoprotein homologue may have a increased expression level in cancer cells compared to normal breast cells. In particular, the term "homologue" covers identity with respect to structure and/or function providing the resultant amino acid sequence has intracellular oncoprotein activity. With respect to sequence identity (i.e. similarity), there may be at least 70%, such as at least 75%, such as at least 85%, such as at least 90% sequence identity. There may be at least 95%, such as at least 98%, sequence identity. These terms also encompass polypeptides derived from amino acids which are allelic variations of the intracellular oncoprotein nucleic acid sequence.

Where reference is made to the "activity" or "biological activity" of a polypeptide such as an intracellular oncoprotein, these terms are intended to refer to the metabolic or physiological function of the intracellular oncoprotein, including similar activities or improved activities or these activities with decreased undesirable side effects. Also included are antigenic and immunogenic activities of the intracellular oncoprotein. Examples of such activities, and methods of assaying and quantifying these activities, are known in the art, and are described in detail elsewhere in this document.

PROPHYLACTIC AND THERAPEUTIC METHODS

We disclose methods of treating an abnormal condition, such as cancer, related to excessive amounts of intracellular oncoprotein expression or activity. Methods of preventing cancer (i.e., prophylaxis) also suitably employ the same or similar approaches.

In general terms, our methods involve manipulation of cancer cells, by modulating (such as down-regulating) the expression, amount or activity of intracellular oncoprotein in the cell. The methods may involve destroying or eradicating cancer cells. The cancer cells may comprise intracellular oncoprotein expressing cancer cells. The cancer cells may be ones which over-express intracellular oncoprotein, compared to non-cancerous cells. Our methods may comprise exposing a patient to an intracellular oncoprotein fragment.

The cancer cells may be from intracellular oncoprotein positive cancer patients. Thus, our methods may comprise eradicating intracellular oncoprotein -over-expressing cancer cells from intracellular oncoprotein positive cancer patients.

A step of detecting modulated intracellular oncoprotein expression, amount or activity in a cell may be conducted before or after the manipulation step. The detection step may detect up-regulated or down-regulated intracellular oncoprotein expression, amount or activity. Any of the methods of modulating or down-regulating intracellular oncoprotein, as described in detail elsewhere in this document, or known in the art, may be used.

In particular, the method may comprise exposing the cell to an anti-intracellular oncoprotein antibody capable of specifically binding to intracellular oncoprotein. In the context of an individual suffering or suspected to be suffering from cancer, the method may comprise administering a therapeutically effective amount of intracellular oncoprotein fragment to the individual. Intracellular oncoprotein fragments and methods of administering them are described in detail elsewhere in this document.

According to our methods, a cancer cell becomes non-cancerous or the invasive or metastatic cancer cell becomes non-invasive or non-metastatic as a result of the manipulation. The cancer may in particular comprise a cancer such as an invasive or metastatic cancer selected from the group consisting of: colorectal cancer, ovarian cancer, breast cancer, liver cancer, pancreatic cancer, prostate cancer, gastric cancer, lung cancer, penis cancer, cervical cancer, brain cancer, esophageal cancer, bladder carcinoma, kidney renal cell carcinoma, ovary lymphoma and skin melanoma.

As intracellular oncoprotein is associated with aggressiveness and invasiveness of cancer, the level of intracellular oncoprotein may be detected in a cell of an individual with cancer, in a cancer or non-cancer cell, and the aggressiveness of the cancer assessed. A high level of intracellular oncoprotein amount, expression or activity compared with a normal cell indicates an aggressive or invasive cancer, and a stronger or harsher therapy may therefore be required and chosen. Similarly, a lower level may indicate a less aggressive or invasive therapy.

The approaches described here may be used for therapy of any intracellular oncoprotein related disease in general. Intracellular oncoprotein related diseases include proliferative diseases and in particular include cancer. For example, a intracellular oncoprotein related disease may include metastatic cancer, invasive cancer or aggressive cancer.

The methods of the invention are useful for treating or preventing the formation of cancer in individuals who have not yet developed a cancer, or who exhibit no symptoms or early stage symptoms of cancer. The individual may have been identified as susceptible, or likely to suffer from, a cancer through the assessment of the prevalence of that cancer in family members. For example, the presence of a cancer associated with a particular intracellular oncoprotein in one or more family members, such as siblings, maternal or paternal relatives, is indicative that the person is also likely to develop that cancer. For example, an infant or young adult whose parent(s) have developed a cancer associated with an intracellular oncoprotein may be determined to be likely to suffer from, or at risk from developing, a cancer associated with that (intracellular) oncoprotein.

The person who has been identified as likely to develop a cancer associated with an intracellular oncoprotein may be administered one or more doses of intracellular oncoprotein fragment based vaccine prior to developing the cancer, or prior to developing symptoms associated with the cancer. Alternatively or additionally, one or more of the doses may be administered after the person has developed the cancer, or has developed symptoms associated with the cancer.

The intracellular oncoprotein fragment based vaccine may be administered in a prime-boost administration regime, wherein the person is administered with a first dose of intracellular oncoprotein fragment vaccine, followed by one or more further doses of intracellular oncoprotein fragment vaccine. Suitable prime-boost administration regimes will be well known to those skilled in the art. For example, a first dose of the intracellular oncogene

fragment based vaccine may be administered at time point zero, with further doses being oncoprotein one week, one month, three months, one year, or other suitable time period later. In some cases, subsequent doses are administered one week, one month, three months, one year or other time period following administration of the previous dose.

The method may further comprise the step of analysing whether the patient produces antibodies that are specific to the intracellular oncoprotein, following administration of the intracellular oncoprotein fragment vaccine.

The methods and compositions described here suitably enable an improvement in a measurable criterion in an individual to whom the treatment is applied compared to one who has not received the treatment.

For this purpose, a number of criteria may be designated, which reflect the progress of cancer or the well-being of the patient. Useful criteria may include tumour size, tumour dimension, largest dimension of tumour, tumour number, presence of tumour markers (such as alpha-feto protein), degree or number of metastates, etc.

Thus, as an example, a treated individual may show a decrease in tumour size or number as measured by an appropriate assay or test. A treated individual may for example show a 1 %, 2%, 3%, 4%, 5%, 6%, 7%, 8%, 9%, 10%, 15%, 20%, 25%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100% or more decrease in tumour size of a particular tumour, or decrease in tumour number, or both, compared to an individual who has not been treated.

For example, a intracellular oncoprotein related disease may be defined as being "treated" if a condition associated with the disease is significantly inhibited (i.e., by 50% or more) relative to controls. The inhibition may be by at least 75% relative to controls, such as by 90%, by 95% or 100% relative to controls. The condition may comprise cell proliferation, or it may comprise cell cycle time, cell number, cell migration, cell invasiveness, tumour formation, tumour metastasis, tumour spread, etc. By the term "treatment" we mean to also include prophylaxis or alleviation of cancer.

The term proliferative disorder is used herein in a broad sense to include any disorder that requires control of the cell cycle. In particular, a proliferative disorder includes malignant and pre-neoplastic disorders. The methods and compositions described here are especially useful in relation to treatment or diagnosis of adenocarcinomas such as: small cell lung cancer, and cancer of the kidney, uterus, prostrate, bladder, ovary, colon and breast. For

example, malignancies which may be treatable include acute and chronic leukemias, lymphomas, myelomas, sarcomas such as Fibrosarcoma, myxosarcoma, liposarcoma, lymphangioendotheliosarcoma, angiosarcoma, endotheliosarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, lymphangiosarcoma, synovioma, mesothelioma, leimyosarcoma, rhabdomyosarcoma, colon carcinoma, ovarian cancer, prostate cancer, pancreatic cancer, breast cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, choriocarcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma seminoma, embryonal carcinoma, cervical cancer, testicular tumour, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, ependymoma, pinealoma, hemangioblastoma, acoustic neuoma, medulloblastoma, craniopharyngioma, oligodendroglioma, menangioma, melanoma, neutroblastoma and retinoblastoma.

The cancers treatable or preventable by the present invention are associated with expression of particular oncoproteins. Cancer cells may express, or aberrantly express (e.g. overexpress), a particular oncoprotein, or incorrectly process the oncoprotein such that it is found at higher levels in the cancer cells than in normal cells.

For example, a sample (e.g. biopsy) is obtained from a subject having a cancer, or suspected of having a cancer, or being at risk of developing a cancer. The sample is tested to determine if a selected oncoprotein, preferably an intracellular oncoprotein is aberrantly expressed, or if the patient is a carrier for a particular isoform or mutant of the oncoprotein that is correlated with a higher risk of cancer development. Accordingly, the subject can be graded for treatment based on the oncoprotein expression profile and treatment (which may be preventative) can be commenced using a protein fragment(s) or peptide(s) according to the present invention.

Over-expression of an oncoprotein comprises expression at a level that is greater than would normally be expected for a cell or tissue of a given type. As such, over-expression may be determined by comparing the level of expression of a protein between cells. For example, a comparison may be made between a cancerous cell and a non-cancerous (healthy) cell of the same type and preferably from the same tissue type (although optionally from a different subject). In another example, a comparison may be made between a cell type present in cancer tissue and a corresponding cell type present in non-cancerous tissue (optionally in the same subject).

Levels of expression may be quantitated for absolute comparison, or relative comparisons may be made.

In some embodiments over-expression of a protein may be considered to be present when the level of expression in the test sample is at least 1.1 times that in the control sample. More preferably, the level of expression may be selected from one of at least 1.2, at least 1.3, at least 1.4, at least 1.5, at least 1.6, at least 1.7, at least 1.8, at least 1.9, at least 2.0, at least 2.1, at least 2.2, at least 2.3, at least 2.4 at least 2.5, at least 2.6, at least 2.7, at least 2.8, at least 2.9, at least 3.0, at least 3.5, at least 4.0, at least 5.0, at least 6.0, at least 7.0, at least 8.0, at least 9.0, or at least 10.0 times that in the control sample.

The approach to therapy described herein involving use of intracellular oncoprotein fragments may be combined with other approaches for therapy of such disorders including expression of anti-sense constructs directed against intracellular oncoprotein polynucleotides as described here, and administering them to tumour cells, to inhibit gene function and prevent the tumour cell from growing or progressing.

Anti-sense constructs may be used to inhibit gene function to prevent growth or progression in a proliferative cell. Antisense constructs, i.e., nucleic acid, such as RNA, constructs complementary to the sense nucleic acid or mRNA, are described in detail in US 6,100,090 (Monia et a/.), and Neckers et al, 1992, Crit Rev Oncog 3(1 -2): 175-231 , the teachings of which document are specifically incorporated by reference.

In a particular example, cancer may be treated or prevented by reducing the amount, expression or activity of intracellular oncoprotein in whole or in part, for example by siRNAs capable of binding to and destroying intracellular oncoprotein mRNA.

RNA interference (RNAi) is a method of post transcriptional gene silencing (PTGS) induced by the direct introduction of double-stranded RNA (dsRNA) and has emerged as a useful tool to knock out expression of specific genes in a variety of organisms. RNAi is described by Fire et al, Nature 391 :806-81 1 (1998). Other methods of PTGS are known and include, for example, introduction of a transgene or virus. Generally, in PTGS, the transcript of the silenced gene is synthesised but does not accumulate because it is rapidly degraded. Methods for PTGS, including RNAi are described, for example, in the Ambion.com world wide web site, in the directory "/hottopics/", in the "rnai" file.

Suitable methods for RNAi in vitro are described herein. One such method involves the introduction of siRNA (small interfering RNA). Current models indicate that these 21 -23 nucleotide dsRNAs can induce PTGS. Methods for designing effective siRNAs are described, for example, in the Ambion web site described above. RNA precursors such as Short Hairpin RNAs (shRNAs) can also be encoded by all or a part of the intracellular oncoprotein nucleic acid sequence.

Alternatively, double-stranded (ds) RNA is a powerful way of interfering with gene expression in a range of organisms that has recently been shown to be successful in mammals (Wianny and Zernicka-Goetz, 2000, Nat Cell Biol 2:70-75). Double stranded RNA corresponding to the sequence of a intracellular oncoprotein polynucleotide can be introduced into or expressed in oocytes and cells of a candidate organism to interfere with intracellular oncoprotein activity.

Other methods of modulating intracellular oncoprotein gene expression are known to those skilled in the art and include dominant negative approaches. Again, these may be combined with antibody therapy using anti-intracellular oncoprotein antibodies. Thus, another approach is to use non-functional variants of intracellular oncoprotein polypeptide in this document that compete with the endogenous gene product resulting in inhibition of function.

Intracellular oncoprotein gene expression may also be modulated by as introducing peptides or small molecules which inhibit gene expression or functional activity. Such peptides or small molecules may be administered in combination with anti-intracellular oncoprotein antibodies for the treatment of cancer such as metastatic cancer.

Thus, compounds identified by assays as binding to or modulating, such as down-regulating, the amount, activity or expression of intracellular oncoprotein polypeptide may be administered to tumour or proliferative cells to prevent the function of intracellular oncoprotein polypeptide. Such a compound may be administered along with a pharmaceutically acceptable carrier in an amount effective to down-regulate expression or activity intracellular oncoprotein, or by activating or down-regulating a second signal which controls intracellular oncoprotein expression, activity or amount, and thereby alleviating the abnormal condition.

Alternatively, gene therapy may be employed to control the endogenous production of intracellular oncoprotein by the relevant cells such as cancer cells in the subject. For example, a polynucleotide encoding a intracellular oncoprotein siRNA or a portion of this

may be engineered for expression in a replication defective retroviral vector, as discussed below. The retroviral expression construct may then be isolated and introduced into a packaging cell transduced with a retroviral plasmid vector containing RNA encoding an anti-intracellular oncoprotein siRNA such that the packaging cell now produces infectious viral particles containing the sequence of interest. These producer cells may be administered to a subject for engineering cells in vivo and regulating expression of the intracellular oncoprotein polypeptide in vivo. For overview of gene therapy, see Chapter 20, Gene Therapy and other Molecular Genetic-based Therapeutic Approaches, (and references cited therein) in Human Molecular Genetics, T Strachan and A P Read, BIOS Scientific Publishers Ltd (1996).

In some embodiments, the level of intracellular oncoprotein is decreased in a cancer cell. Furthermore, in such embodiments, treatment may be targeted to, or specific to, such cancer cells. The expression of intracellular oncoprotein may be specifically decreased only in diseased cells (i.e., those cells which are cancerous), and not substantially in other non-diseased cells. In these methods, expression of intracellular oncoprotein may be not substantially reduced in other cells, i.e., cells which are not cancer cells. Thus, in such embodiments, the level of intracellular oncoprotein remains substantially the same or similar in non-cancer cells in the course of or following treatment.

PATIENTS

The subject to be treated may be any animal or human. The subject is preferably mammalian, more preferably human. The subject may be a non-human mammal, but is more preferably human. The subject may be male or female. The subject may be a patient. Therapeutic uses may be in human or animals (veterinary use).

POLYPEPTIDE SEQUENCES

It will be understood that polypeptide sequences disclosed here are not limited to the particular sequences set forth in this document, but also include homologous sequences obtained from any source, for example related cellular homologues, homologues from other species and variants or derivatives thereof, provided that they have at least one of the biological activities of an anti-intracellular oncoprotein antibody, as the case may be.

This disclosure therefore encompasses variants, homologues or derivatives of the amino acid sequences set forth in this document, as well as variants, homologues or derivatives of the amino acid sequences encoded by the nucleotide sequences disclosed here. Such sequences are generally referred to as a " intracellular oncoprotein " sequence.

BIOLOGICAL ACTIVITIES

In some embodiments, the sequences comprise at least one biological activity of an intracellular oncoprotein fragment, as the case may be.

The biological activity may comprise an immunological activity. The intracellular oncoprotein fragment may comprise an identical or similar immunological activity as compared to intracellular oncoprotein antibody or its humanised versions. By "immunological activity" we mean the capability of the anti-intracellular oncoprotein antibody, to induce a specific immune response in appropriate animals or cells on binding with a intracellular oncoprotein antigen.

The activity may include inhibition of cancer activity as for example measured by reduction of tumour size or tumour number, or inhibition of metastatic activity, such as for example measured by the assays described in the Examples. The reduction or inhibition may be conveniently assayed by causing carcinogenesis in a test animal, administering the intracellular oncoprotein fragment to the animal and determining an effect of the intracellular oncoprotein fragment as compared to a similar control animal that has not been so treated. The Examples describe such an assay in detail.

The intracellular oncoprotein fragment may have tumour inhibition or metastasis inhibition activity that is the same as, reduced from, or elevated from, the cognate fragment. For example, the intracellular oncoprotein fragment may be at least 10%, such as 20%, such as 30%, 40% 50%, 60%, 70%, 80%, 90% or more, effective compared to the cognate antibody. By this we mean that, say, if the cognate fragment is capable of reducing tumour number by 90% (see the Examples), the intracellular oncoprotein fragment may be capable of reducing tumour number by 90%, 85%, 80%, 75%, 70%, 65%, 60%, 55%, 50%, 45%, 40%, 35%, 30%, etc, as compared to an untreated animal.

Other assays that detect antibody events can also be used, instead of, or in addition to, the assays described.

HOMOLOGUES

The intracellular oncoprotein fragment polypeptides disclosed include homologous sequences obtained from any source, for example related viral/bacterial proteins, cellular homologues and synthetic peptides, as well as variants or derivatives thereof. Thus polypeptides also include those encoding homologues of intracellular oncoprotein fragment

from other species including animals such as mammals (e.g. mice, rats or rabbits), in particular humans.

In the context of the present document, a homologous sequence or homologue is taken to include an amino acid sequence which is at least 60, 70, 80 or 90% identical, such as at least 95 or 98% identical at the amino acid level over at least 30, such as 50, 70, 90 or 100 amino acids with a relevant polypeptide sequence, for example as shown in the sequence listing herein. In the context of this document, a homologous sequence is taken to include an amino acid sequence which is at least 15, 20, 25, 30, 40, 50, 60, 70, 80 or 90% identical, such as at least 95 or 98% identical at the amino acid level, such as over at least 15, 25, 35, 50 or 100, such as 200, 300, 400 or 500 amino acids with the sequence of a relevant polypeptide. Although homology can also be considered in terms of similarity (i.e. amino acid residues having similar chemical properties/functions), in the context of the present document homology may be expressed in terms of sequence identity. The sequence identity may be determined relative to the entirety of the length the relevant sequence, i.e., over the entire length or full length sequence of the relevant gene, for example.

Homology comparisons can be conducted by eye, or more usually, with the aid of readily available sequence comparison programs. These commercially available computer programs can calculate % homology between two or more sequences.

% homology may be calculated over contiguous sequences, i.e. one sequence is aligned with the other sequence and each amino acid in one sequence directly compared with the corresponding amino acid in the other sequence, one residue at a time. This is called an "ungapped" alignment. Typically, such ungapped alignments are performed only over a relatively short number of residues (for example less than 50 contiguous amino acids).

Although this is a very simple and consistent method, it fails to take into consideration that, for example, in an otherwise identical pair of sequences, one insertion or deletion will cause the following amino acid residues to be put out of alignment, thus potentially resulting in a large reduction in % homology when a global alignment is performed. Consequently, most sequence comparison methods are designed to produce optimal alignments that take into consideration possible insertions and deletions without penalising unduly the overall homology score. This is achieved by inserting "gaps" in the sequence alignment to try to maximise local homology.

However, these more complex methods assign "gap penalties" to each gap that occurs in the alignment so that, for the same number of identical amino acids, a sequence alignment with as few gaps as possible - reflecting higher relatedness between the two compared sequences - will achieve a higher score than one with many gaps. "Affine gap costs" are typically used that charge a relatively high cost for the existence of a gap and a smaller penalty for each subsequent residue in the gap. This is the most commonly used gap scoring system. High gap penalties will of course produce optimised alignments with fewer gaps. Most alignment programs allow the gap penalties to be modified.

However, the default values may be used when using such software for sequence comparisons. For example when using the GCG Wisconsin Bestfit package (see below) the default gap penalty for amino acid sequences is -12 for a gap and -4 for each extension.

Calculation of maximum % homology therefore firstly requires the production of an optimal alignment, taking into consideration gap penalties. A suitable computer program for carrying out such an alignment is the GCG Wisconsin Bestfit package (University of Wisconsin, U.S.A.; Devereux et al, 1984, Nucleic Acids Research 12:387). Examples of other software than can perform sequence comparisons include, but are not limited to, the BLAST package (see Ausubel et al, 1999 ibid - Chapter 18), FASTA (Atschul et al, 1990, J. Mol. Biol., 403-410) and the GENEWORKS suite of comparison tools. Both BLAST and FASTA are available for offline and online searching (see Ausubel et al., 1999 ibid, pages 7-58 to 7-60). The GCG Bestfit program may be used.

Although the final % homology can be measured in terms of identity, the alignment process itself is typically not based on an all-or-nothing pair comparison. Instead, a scaled similarity score matrix is generally used that assigns scores to each pairwise comparison based on chemical similarity or evolutionary distance. An example of such a matrix commonly used is the BLOSUM62 matrix - the default matrix for the BLAST suite of programs. GCG Wisconsin programs generally use either the public default values or a custom symbol comparison table if supplied (see user manual for further details). The public default values for the GCG package, or in the case of other software, the default matrix, such as BLOSUM62, may be used.

Once the software has produced an optimal alignment, it is possible to calculate % homology, such as % sequence identity. The software typically does this as part of the sequence comparison and generates a numerical result.

VARIANTS AND DERIVATIVES

The terms "variant" or "derivative" in relation to the amino acid sequences as described here includes any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) amino acids from or to the sequence. The resultant amino acid sequence may retain substantially the same activity as the unmodified sequence, such as having at least the same activity as the anti-intracellular oncoprotein antibody polypeptides shown in this document, for example in the sequence listings. Thus, the key feature of the sequences - namely ability to bind to intracellular oncoprotein polypeptides or tumour reduction activity, as described elsewhere - may be retained.

Polypeptides having the amino acid sequence shown in the Examples, or fragments or homologues thereof may be modified for use in the methods and compositions described here. Typically, modifications are made that maintain the biological activity of the sequence. Amino acid substitutions may be made, for example from 1, 2 or 3 to 10, 20 or 30 substitutions provided that the modified sequence retains the biological activity of the unmodified sequence. Amino acid substitutions may include the use of non-naturally occurring analogues, for example to increase blood plasma half-life of a therapeutically administered polypeptide.

Natural variants of intracellular oncoprotein fragments are likely to comprise conservative amino acid substitutions. Conservative substitutions may be defined, for example according to the Table below. Amino acids in the same block in the second column such as those in the same line in the third column may be substituted for each other:

ALIPHATIC	Non-polar	GAP
		ILV
	Polar-uncharged	CSTM
		NQ
	Polar-charged	DE
		KR
AROMATIC		HFWY

Intracellular oncoprotein fragments, homologues, variants and derivatives may be made by recombinant means. However, they may also be made by synthetic means using techniques well known to skilled persons such as solid phase synthesis. The proteins may also be produced as fusion proteins, for example to aid in extraction and purification. Examples of fusion protein partners include glutathione-S-transferase (GST), 6xHis, GAL4 (DNA binding and/or transcriptional activation domains) and β-galactosidase. It may also be convenient to include a proteolytic cleavage site between the fusion protein partner and the protein sequence of interest to allow removal of fusion protein sequences. The fusion protein may be such that it will not hinder the function of the protein of interest sequence. Proteins may also be obtained by purification of cell extracts from animal cells. In the present case, a fusion protein may comprise two or more fragments of one or more oncoproteins, optionally further including a linker peptide. The fragments may be from the same oncoprotein or different oncoproteins. The fusion protein may comprise two or more copies of the same fragment, or overlapping fragments of the same epitope or oncogenic mutation.

The intracellular oncoprotein polypeptides, variants, homologuesand derivatives disclosed here may be in a substantially isolated form. It will be understood that such polypeptides may be mixed with carriers or diluents which will not interfere with the intended purpose of the protein and still be regarded as substantially isolated. A intracellular oncoprotein fragment variant, homologue or derivative may also be in a substantially purified form, in which case it will generally comprise the protein in a preparation in which more than 90%, e.g. 95%, 98% or 99% of the protein in the preparation is a protein.

The anti-intracellular oncoprotein fragment polypeptides, variants, homologuesand derivatives disclosed here may be labelled with a revealing label. The revealing label may be any suitable label which allows the polypeptide , etc to be detected. Suitable labels include radioisotopes, e.g. ^{125}I, enzymes, antibodies, polynucleotides and linkers such as biotin. Labelled polypeptides may be used in diagnostic procedures such as immunoassays to determine the amount of a polypeptide in a sample. Polypeptides or labelled polypeptides may also be used in serological or cell-mediated immune assays for the detection of immune reactivity to said polypeptides in animals and humans using standard protocols.

The intracellular oncoprotein fragmentpolypeptides, variants, homologues, and derivatives disclosed here, optionally labelled, my also be fixed to a solid phase, for example the surface of an immunoassay well or dipstick. Such labelled and/or immobilised polypeptides may be packaged into kits in a suitable container along with suitable reagents, controls, instructions

and the like. Such polypeptides and kits may be used in methods of detection of antibodies to the polypeptides or their allelic or species variants by immunoassay.

Immunoassay methods are well known in the art and will generally comprise: (a) providing a polypeptide comprising an epitope bindable by an antibody against said protein; (b) incubating a biological sample with said polypeptide under conditions which allow for the formation of an antibody-antigen complex; and (c) determining whether antibody-antigen complex comprising said polypeptide is formed.

The intracellular oncoprotein fragment polypeptides, variants, homologues, and derivatives disclosed here may be used in in vitro or in vivo cell culture systems to study the role of their corresponding genes and homologues thereof in cell function, including their function in disease. For example, truncated or modified polypeptides may be introduced into a cell to disrupt the normal functions which occur in the cell. The polypeptides may be introduced into the cell by in situ expression of the polypeptide from a recombinant expression vector (see below). The expression vector optionally carries an inducible promoter to control the expression of the polypeptide.

The use of appropriate host cells, such as insect cells or mammalian cells, is expected to provide for such post-translational modifications (e.g. myristolation, glycosylation, truncation, lapidation and tyrosine, serine or threonine phosphorylation) as may be needed to confer optimal biological activity on recombinant expression products. Such cell culture systems in which the intracellular oncoprotein polypeptides, variants, homologues, and derivatives disclosed here are expressed may be used in assay systems to identify candidate substances which interfere with or enhance the functions of the polypeptides in the cell.

POLYNUCLEOTIDE SEQUENCES

The variable regions, monoclonal antibody sequences and humanised antibody sequences may comprise polynucleotides. These may comprise DNA or RNA.

They may be single-stranded or double-stranded. They may also be polynucleotides which include within them synthetic or modified nucleotides. A number of different types of modification to oligonucleotides are known in the art. These include methylphosphonate and phosphorothioate backbones, addition of acridine or poly lysine chains at the 3' and/or 5' ends of the molecule. For the purposes of the present document, it is to be understood that the polynucleotides described herein may be modified by any method available in the art.

Such modifications may be carried out in order to enhance the in vivo activity or life span of polynucleotides.

Where the polynucleotide is double-stranded, both strands of the duplex, either individually or in combination, are encompassed by the methods and compositions described here. Where the polynucleotide is single-stranded, it is to be understood that the complementary sequence of that polynucleotide is also included.

VARIANTS, DERIVATIVES AND HOMOLOGUES

The terms "variant", "homologue" or "derivative" in relation to a nucleotide sequence described in this document include any substitution of, variation of, modification of, replacement of, deletion of or addition of one (or more) nucleotides from or to the sequence. The resulting sequence may be capable of encoding a polypeptide which has intracellular oncoprotein binding activity as described elsewhere in this document.

As indicated above, with respect to sequence identity, a "homologue" has such as at least 5% identity, at least 10% identity, at least 15% identity, at least 20% identity, at least 25% identity, at least 30% identity, at least 35% identity, at least 40% identity, at least 45% identity, at least 50% identity, at least 55% identity, at least 60% identity, at least 65% identity, at least 70% identity, at least 75% identity, at least 80% identity, at least 85% identity, at least 90% identity, or at least 95% identity to a relevant sequence.

There may be at least 95% identity, such as at least 96% identity, such as at least 97% identity, such as at least 98% identity, such as at least 99% identity. Nucleotide homology comparisons may be conducted as described above. A sequence comparison program such as the GCG Wisconsin Bestfit program described above may be used for this purpose. The default scoring matrix has a match value of 10 for each identical nucleotide and -9 for each mismatch. The default gap creation penalty is -50 and the default gap extension penalty is -3 for each nucleotide.

HYBRIDISATION

We further describe nucleotide sequences that are capable of hybridising selectively to any of the sequences presented herein, such as 269, 223 and 318 variable region, antibody and humanised antibody or any variant, fragment or derivative thereof, or to the complement of any of the above. Nucleotide sequences may be at least 1 nucleotides in length, such as at least 20, 30, 40 or 50 nucleotides in length.

The term "hybridisation" as used herein shall include "the process by which a strand of nucleic acid joins with a complementary strand through base pairing" as well as the process of amplification as carried out in polymerase chain reaction technologies.

Polynucleotides capable of selectively hybridising to the nucleotide sequences presented herein, or to their complement, will be generally at least 70%, such as at least 80 or 90% and such as at least 95% or 98% homologous to the corresponding nucleotide sequences presented herein over a region of at least 20, such as at least 25 or 30, for instance at least 40, 60 or 100 or more contiguous nucleotides.

The term "selectively hybridisable" means that the polynucleotide used as a probe is used under conditions where a target polynucleotide is found to hybridize to the probe at a level significantly above background. The background hybridization may occur because of other polynucleotides present, for example, in the cDNA or genomic DNA library being screened. In this event, background implies a level of signal generated by interaction between the probe and a non- specific DNA member of the library which is less than 10 fold, such as less than 100 fold as intense as the specific interaction observed with the target DNA. The intensity of interaction may be measured, for example, by radiolabelling the probe, e.g. with ^{32}P.

Hybridisation conditions are based on the melting temperature (Tm) of the nucleic acid binding complex, as taught in Berger and Kimmel (1987, Guide to Molecular Cloning Techniques, Methods in Enzymology, Vol 152, Academic Press, San Diego CA), and confer a defined "stringency" as explained below.

Maximum stringency typically occurs at about Tm-5°C (5°C below the Tm of the probe); high stringency at about 5°C to 10°C below Tm; intermediate stringency at about 10°C to 20°C below Tm; and low stringency at about 20°C to 25°C below Tm. As will be understood by those of skill in the art, a maximum stringency hybridisation can be used to .identify or detect identical polynucleotide sequences while an intermediate (or low) stringency hybridisation can be used to identify or detect similar or related polynucleotide sequences.

We disclose nucleotide sequences that can hybridise to a nucleic acid, or a fragment, homologue, variant or derivative thereof, under stringent conditions (e.g. 65°C and O.lxSSC { lxSSC = 0.15 M NaCl, 0.015 M Na_3 Citrate pH 7.0}).

Where a polynucleotide is double-stranded, both strands of the duplex, either individually or in combination, are encompassed by the present disclosure. Where the polynucleotide is single- stranded, it is to be understood that the complementary sequence of that polynucleotide is also disclosed and encompassed.

Polynucleotides which are not 100% homologous to the sequences disclosed here but fall within the disclosure can be obtained in a number of ways. Other variants of the sequences described herein may be obtained for example by probing DNA libraries made from a range of individuals, for example individuals from different populations. In addition, other viral/bacterial, or cellular homologues particularly cellular homologues found in mammalian cells (e.g. rat, mouse, bovine and primate cells), may be obtained and such homologues and fragments thereof in general will be capable of selectively hybridising to the sequences shown in the sequence listing herein. Such sequences may be obtained by probing cDNA libraries made from or genomic DNA libraries from other animal species, and probing such libraries with probes comprising all or part of the disclosed sequences under conditions of medium to high stringency.

The polynucleotides described here may be used to produce a primer, e.g. a PCR primer, a primer for an alternative amplification reaction, a probe e.g. labelled with a revealing label by conventional means using radioactive or non-radioactive labels, or the polynucleotides may be cloned into vectors. Such primers, probes and other fragments will be at least 15, such as at least 20, for example at least 25, 30 or 40 nucleotides in length, and are also encompassed by the term polynucleotides as used herein. Fragments may be less than 500, 200, 100, 50 or 20 nucleotides in length.

Polynucleotides such as a DNA polynucleotides and probes may be produced recombinantly, synthetically, or by any means available to those of skill in the art. They may also be cloned by standard techniques.

In general, primers will be produced by synthetic means, involving a step wise manufacture of the desired nucleic acid sequence one nucleotide at a time. Techniques for accomplishing this using automated techniques are readily available in the art.

Longer polynucleotides will generally be produced using recombinant means, for example using PGR (polymerase chain reaction) cloning techniques. This will involve making a pair of primers (e.g. of about 15 to 30 nucleotides) flanking a region of the sequence which it is desired to clone, bringing the primers into contact with mRNA or cDNA obtained from an

animal or human cell, performing a polymerase chain reaction under conditions which bring about amplification of the desired region, isolating the amplified fragment (e.g. by purifying the reaction mixture on an agarose gel) and recovering the amplified DNA. The primers may be designed to contain suitable restriction enzyme recognition sites so that the amplified DNA can be cloned into a suitable cloning vector.

INTRACELLULAR ONCOPROTEIN POLYPEPTIDES AND NUCLEIC ACIDS

Intracellular oncoprotein polypeptide homologues, variants, derivatives and fragments may be defined similarly, as set out in the previous paragraphs.

Where the context permits, a reference to intracellular oncoprotein polypeptide should be taken to include reference to an intracellular oncoprotein polypeptide homologue, variant, derivative or fragment. Similarly, a reference to intracellular oncoprotein polypeptide should be taken to include reference to a intracellular oncoprotein polypeptide homologue, variant, derivative or fragment.

Similarly, where the context permits, a reference to intracellular oncoprotein nucleic acid should be taken to include reference to a intracellular oncoprotein nucleic acid homologue, variant, derivative or fragment. Similarly, a reference to intracellular oncoprotein polypeptide should be taken to include reference to a intracellular oncoprotein nucleic acid homologue, variant, derivative or fragment.

ANTI-INTRACELLULAR ONCOPROTEIN FRAGMENT PRODUCTION

The intracellular oncoprotein fragment can be produced by recombinant DN A methods or synthetic peptide chemical methods that are well known to those of ordinary skill in the art.

By way of example, the intracellular oncoprotein fragments may be synthesized by techniques well known in the art, as exemplified by "Solid Phase Peptide Synthesis: A Practical Approach" E. Atherton and R. C. Sheppard, IRL Press, Oxford England. Similarly, multiple fragments can be synthesized which are subsequently linked together to form larger fragments. These synthetic peptide fragments can also be made with amino acid substitutions at specific locations in order to test for activity in vitro and in vivo.

The intracellular oncoprotein can be synthesized in a standard microchemical facility and purity checked with HPLC and mass spectrophotometry. Methods of peptide synthesis, HPLC purification and mass spectrophotometry are commonly known to those skilled in these arts.

The intracellular oncoprotein fragments may also be expressed under in vitro and in vivo conditions in a transformed host cell into which has been incorporated the DNA sequences described here (such as variable sequences) or allelic variations thereof and which can be used in the prevention and/or treatment of cancer related diseases.

The term "vector" includes expression vectors and transformation vectors. The term "expression vector" means a construct capable of in vivo or in vitro expression. The term "transformation vector" means a construct capable of being transferred from one species to another.

Vectors which may be used for expression include recombinant viral vectors, in particular recombinant retroviral vectors (RRV) such as lentiviral vectors, adenoviral vectors including a combination of retroviral vectors.

The term 'recombinant retroviral vector" (RRV) refers to a vector with sufficient retroviral genetic information to allow packaging of an RNA genome, in the presence of packaging components, into a viral particle capable of infecting a target cell. Infection of the target cell includes reverse transcription and integration into the target cell genome. The RRV carries non-viral coding sequences which are to be delivered by the vector to the target cell. An RRV is incapable of independent replication to produce infectious retroviral particles within the final target cell. Usually the RRV lacks a functional gag pol and/or env gene and/or other genes essential for replication. Vectors which may be used include recombinant pox viral vectors such as fowl pox virus (FPV), entomopox virus, vaccinia virus such as NYVAC, canarypox virus, MVA or other non-replicating viral vector systems such as those described for example in WO9530018.

Pox viruses may be engineered for recombinant gene expression and for the use as recombinant live vaccines in a dual immunotherapeutic approach. The principal rationale for using live attenuated viruses, such as viruses, as delivery vehicles and/or vector based vaccine candidates, stems from their ability to elicit cell mediated immune responses. The viral vectors, as outlined above, are capable of being employed as delivery vehicles and as vector based vaccine candidates because of the immunogenicity of their constitutive proteins, which act as adjuvants to enhance the immune response, thus rendering a nucleotide sequence of interest (NOI) such as a nucleotide sequence encoding an intracellular oncoprotein more immunogenic.

The pox virus vaccination strategies have used recombinant techniques to introduce NOIs into the genome of the pox virus. If the NOI is integrated at a site in the viral DNA which is non- essential for the life cycle of the virus, it is possible for the newly produced recombinant pox virus to be infectious, that is to say to infect foreign cells and thus to express the integrated NOI. The recombinant pox virus prepared in this way can be used as live vaccines for the prophylaxis and/or treatment of pathologic and infectious disease and/or cancer.

Other requirements for pox viral vector delivery systems include good immunogenicity and safety. MVA is a replication-impaired vaccinia strain with a good safety record. In most cell types and normal human tissue, MVA does not replicate. Limited replication of MVA is observed in a few transformed cell types such as BHK21 cells. Carroll et al (1997 Vaccinel 5: 387-394) have shown that the recombinant MVA is equally as good as conventional recombinant vaccinia vectors at generating a protective CD8+T cell response and is an efficacious alternative to the more commonly used replication competent vaccinia virus. The vaccinia virus strains derived from MVA, or independently developed strains having the features of MVA which make MVA particularly suitable for use in a vaccine, are also suitable for use as a delivery vehicle.

The nucleotide sequence of interest, and of which expression is desired, may operably linked to a transcription unit. The term "transcription unit" as described herein are regions of nucleic acid containing coding sequences and the signals for achieving expression of those coding sequences independently of any other coding sequences. Thus, each transcription unit generally comprises at least a promoter, an optional enhancer and a polyadenylation signal. The term "promoter" is used in the normal sense of the art, e. g. an R A polymerase binding site. The promoter may contain an enhancer element. The term "enhancer" includes a DNA sequence which binds to other protein components of the transcription initiation complex and thus facilitates the initiation of transcription directed by its associated promoter. The term "cell" includes any suitable organism. The cell may comprise a mammalian cell, such as a human cell.

The term "transformed cell" means a cell having a modified genetic structure. For example, as described here, a cell has a modified genetic structure when a vector such as an expression vector has been introduced into the cell. The term "organism" includes any suitable organism. The organism may comprise a mammal such as a human.

Here the term "transgenic organism" means an organism comprising a modified genetic structure. For example, the organism may have a modified genetic structure if a vector such as an expression vector has been introduced into the organism.

INTRACELLULAR ONCOPROTEIN FRAGMENT EXPRESSION

We further describe a method comprising transforming a host cell with a or the nucleotide sequences described in this document.

We also provide a method comprising culturing a transformed host cell-which cell has been transformed with a or the such nucleotide sequences under conditions suitable for the expression of the intracellular oncoprotein fragmentencoded by said nucleotide sequences.

We further provide a method comprising culturing a transformed host cell-which cell has been transformed with a or the such nucleotide sequences under conditions suitable for the expression of the intracellular oncoprotein fragment encoded by said nucleotide sequences; and then recovering said intracellular oncoprotein fragment from the transformed host cell culture.

Thus, intracellular oncoprotein fragment encoding nucleotide sequences, fusion proteins or functional equivalents thereof, may be used to generate recombinant DNA molecules that direct the expression thereof in appropriate host cells.

By way of example, intracellular oncoprotein fragment may be produced in recombinant E. coli, yeast or mammalian expression systems, and purified with column chromatography.

The nucleotide sequences encoding the intracellular oncoprotein fragmentmay be operably linked to a promoter sequence capable of directing expression of the intracellular oncoprotein fragment encoding nucleotide sequences in a suitable host cell. When inserted into the host cell, the transformed host cell may be cultured under suitable conditions until sufficient levels of the intracellular oncoprotein fragment are achieved after which the cells may be lysed and the intracellular oncoprotein fragment is isolated.

Host cells transformed with the intracellular oncoprotein fragment encoding nucleotide sequences may be cultured under conditions suitable for the expression and recovery of the intracellular oncoprotein fragment from cell culture. The protein produced by a recombinant cell may be secreted or may be contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors

containing the intracellular oncoprotein fragment encoding nucleotide sequences can be designed with signal sequences which direct secretion of the intracellular oncoprotein fragment encoding nucleotide sequences through a particular prokaryotic or eukaryotic cell membrane. Other recombinant constructions may join the intracellular oncoprotein fragment encoding nucleotide sequence to a nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins (Kroll DJ et al(1993) DNA Cell Biol 12:441 - 5 3', see also the discussion below on vectors containing fusion proteins).

The intracellular oncoprotein fragment may also be expressed as a recombinant protein with one or more additional polypeptide domains added to facilitate protein purification. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals (Porath J (1992) Protein Expr Purif 3-26328 1), protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp, Seattle, WA). The inclusion of a cleavable linker sequence such as Factor XA or enterokinase (Invitrogen, San Diego, CA) between the purification domain and the intracellular oncoprotein fragment is useful to facilitate purification.

The nucleotide sequences described here may be engineered in order to alter a the intracellular oncoprotein fragment encoding sequences for a variety of reasons, including but not limited to alterations which modify the cloning, processing and/or expression of the gene product.

For example, mutations may be introduced using techniques which are well known in the art, e.g., site-directed mutagenesis to insert new restriction sites, to alter glycosylation patterns or to change codon preference.

In another embodiment, a or the natural, modified or recombinant intracellular oncoprotein fragment encoding nucleotide sequences may be ligated to a heterologous sequence to encode a fusion protein. By way of example, fusion proteins comprising the intracellular oncoprotein fragment or an enzymatically active fragment or derivative thereof linked to an affinity tag such as glutathione-S-transferase (GST), biotin, His6, ac-myc tag (see Emrich etal 1993 BiocemBiophys Res Commun 197(1): 21220), hemagglutinin (HA) (as described in Wilson et al (1984 Cell 37 767) or a FLAG epitope (Ford etal 1991 Protein Expr Purif Apr; 2 (2):95-107). May be produced

The fused recombinant protein may comprise an antigenic coprotein such as GST, beta-galactosidase or the lipoprotein D from Haemophilus influenzae which are relatively large co-proteins, which solubilise and facilitate production and purification thereof. Alternatively, the fused protein may comprise a carrier protein such as bovine serum albumin (BSA) or keyhole limpet haemocyanin (KLH). In certain embodiments, the marker sequence may comprise a hexa- histidine peptide, as provided in the pQE vector (Qiagen Inc) and described in Gentz et al (1989 PNAS 86: 821 -824). Such fusion proteins are readily expressable in yeast culture (as described in Mitchell et al 1993 Yeast 5:715-723) and are easily purified by affinity chromatography. A fusion protein may also be engineered to contain a cleavage site located between the nucleotide sequence encoding the intracellular oncoprotein fragment and the heterologous protein sequence, so that the intracellular oncoprotein fragment may be cleaved and purified away from the heterologous moiety. In another embodiment, an assay for the target protein may be conducted using the entire, bound fusion protein. Alternatively, the co-protein may act as an adjuvant in the sense of providing a generalised stimulation of the immune system. The co- protein may be attached to either the amino or carboxy terminus of the first protein.

Although the presence/absence of marker gene expression suggests that the nucleotide sequence for intracellular oncoprotein fragment is also present, its presence and expression should be confirmed. For example, if the intracellular oncoprotein fragment encoding nucleotide sequence is inserted within a marker gene sequence, recombinant cells containing the intracellular oncoprotein fragment coding regions may be identified by the absence of the marker gene function. Alternatively, a marker gene may be placed in tandem with an intracellular oncoprotein fragment encoding nucleotide sequence under the control of a single promoter.

Expression of the marker gene in response to induction or selection usually indicates expression of the intracellular oncoprotein fragment as well.

Additional methods to quantitate the expression of a particular molecule include radiolabeling (Melby PC etal 1993 J Immunol Methods 159:235-44) or biotinylating (Duplaa C et al 1993 Anal Biochem229-36) nucleotides, co amplification of a control nucleic acid, and standard curves onto which the experimental results are interpolated.

Quantitation of multiple samples may be speeded up by running the assay in an ELISA format where the intracellular oncoprotein fragment of interest is presented in various dilutions and a spectrophotometric or calorimetric response gives rapid quantitation.

Altered intracellular oncoprotein fragment nucleotide sequences which may be made or used include deletions, insertions or substitutions of different nucleotide residues resulting in a nucleotide sequence that encodes the same or a functionally equivalent intracellular oncoprotein fragment. By way of example, the expressed intracellular oncoprotein fragment may also have deletions, insertions or substitutions of amino acid residues which produce a silent change and result in a functionally equivalent intracellular oncoprotein fragment. Deliberate amino acid substitutions may be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity. and/or the amphipathic nature of the residues as long as the binding affinity of the anti-intracellular oncoprotein fragment is retained. For example, negatively charged amino acids include aspartic acid and glutamic acid: positively charged amino acids include lysine and arginine; and amiiio acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine, valine, glycine, alanine, asparagine, glutamine, serine, threonine, phenylalanine, and tvrosine.

Gene therapy whereby the anti-intracellular oncoprotein fragment encoding nucleotide sequences as described here is regulated in vivo may also be employed. For example, expression regulation may be accomplished by administering compounds that bind to the anti-intracellular oncoprotein antibody encoding nucleotide sequences, or control regions associated with the intracellular oncoprotein fragment encoding nucleotide sequence or its corresponding RNA transcript to modify the rate of transcription or translation.

By way of example, the intracellular oncoprotein fragment encoding nucleotide sequences described here may be under the expression control of an expression regulatory element, usually a promoter or a promoter and enhancer. The enhancer and/or promoter may be preferentially active in a hypoxic or ischaemic or low glucose environment, such that the intracellular oncoprotein fragment encoding nucleotide sequences is preferentially expressed in the particular tissues of interest, such as in the environment of a tumour cell or mass. Thus, any significant biological effect or deleterious effect of the intracellular oncoprotein fragment encoding nucleotide sequences on the individual being treated may be reduced or eliminated. The enhancer element or other elements conferring regulated expression may be present in multiple copies.

The promoter and/or enhancer may be constitutively efficient, or may be tissue or temporally restricted in their activity. Examples of suitable tissue restricted promoters/enhancers are those which are highly active in tumour cells such as a promoter/enhancer from a MUC1

gene, a CEA gene or a STV antigen gene. Examples of temporally restricted promoters/enhancers are those which are responsive to ischaemia and/or hypoxia, such as hypoxia response elements or the promoter/enhancer of agrp78 or agrp94 gene. The alpha fetoprotein (AFP) promoter is also a tumour-specific promoter. Another promoter-enhancer combination is a human cytomegalovirus (hCMV) major immediate early (MIE) promoter/enhancer combination.

The promoters may be tissue specific. That is, they may be capable of driving transcription of a intracellular oncoprotein fragment encoding nucleotide sequences in one tissue while remaining largely "silent" in other tissue types.

The term "tissue specific" means a promoter which is not restricted in activity to a single tissue type but which nevertheless shows selectivity in that they may be active in one group of tissues and less active or silent in another group. A desirable characteristic of such promoters is that they possess a relatively low activity in the absence of activated hypoxia-regulated enhancer elements, even in the target tissue. One means of achieving this is to use "silencer" elements which suppress the activity of a selected promoter in the absence of hypoxia.

The term "hypoxia" means a condition under which a particular organ or tissue receives an inadequate supply of oxygen.

The level of expression of a or the intracellular oncoprotein fragment encoding nucleotide sequences under the control of a particular promoter may be modulated by manipulating the promoter region. For example, different domains within a promoter region may possess different gene regulatory activities. The roles of these different regions are typically assessed using vector constructs having different variants of the promoter with specific regions deleted (that is, deletion analysis). This approach may be used to identify, for example, the smallest region capable of conferring tissue specificity or the smallest region conferring hypoxia sensitivity.

A number of tissue specific promoters, described above, may be used. In most instances, these promoters may be isolated as convenient restriction digestion fragments suitable for cloning in a selected vector. Alternatively, promoter fragments may be isolated using the polymerase chain reaction. Cloning of the amplified fragments may be facilitated by incorporating restriction sites at the 5' end of the primers.

COMBINATION THERAPY

The methods and compositions described here, including intracellular oncoprotein fragment vaccines, may be used in combination with other compositions and procedures for the treatment of diseases.

By way of example, the intracellular oncoprotein fragment vaccines may also be used in combination with conventional treatments of diseases such as cancer. For example, a tumor may be treated conventionally with surgery, radiation or chemotherapy combined with intracellular oncoprotein fragmentvaccine may be subsequently administered to the patient to extend the dormancy of micrometastases and to stabilize any residual primary tumor.

The intracellular oncoprotein fragment vaccine can be delivered with a therapeutically effective agent at the same moment in time and at the same site. Alternatively, the intracellular oncoprotein fragment vaccine and the therapeutically effective agent may be delivered at a different time and to a different site. The intracellular oncoprotein fragment vaccine and the therapeutically effective agent may even be delivered in the same delivery vehicle for the prevention and/or treatment of cancer.

Intracellular oncoprotein fragment vaccine may be used in combination with cytotoxic agents for the prevention and/or treatment of angiogenesis and/or cancer. Cytotoxic agents such as ricin, linked to intracellular oncoprotein fragment provides a tool for the destruction of cells that express intracellular oncoprotein or intracellular oncoprotein. These cells may be found in many locations, including but not limited to, micrometastases and primary tumours.

Intracellular oncoprotein fragment vaccine may be used in combination with a pro-drug activating enzyme in gene therapy. Instead of or as well as being selectively expressed in target tissues, the intracellular oncoprotein antibody and/or intracellular oncoprotein vaccine may be used in combination with another molecule, such as a pro-drug activation enzyme or enzymes which have no significant effect or no deleterious effect until the individual is treated with one or more pro-drugs upon which the enzyme or enzymes act. In the presence of the pro-drug activation enzyme, active treatment of an individual with the appropriate pro-drug leads to enhanced reduction in tumour growth or survival.

A pro-drug activating enzyme may be delivered to a tumour site for the treatment of a cancer. In each case, a suitable pro-drug is used in the treatment of the patient in combination with the appropriate pro-drug activating enzyme. An appropriate pro-drug is administered in conjunction with the vector. Examples of pro-drugs include: etoposide

phosphate (with alkaline phosphatase, Senter et al 1988 Proc Natl Acad Sci 85: 48424846); 5-fluorocytosine (with cytosine deaminase, Mullen et al 1994 Cancer Res 54: 1503-1506); Doxorubicin-N -p- hydroxyphenoxyacetamide (with Penicillin- V -Amidase, Kerr et al 1990 Cancer Immunol Immunother 31 : 202-206); Para-N-bis(2-chloroethyl) aminobenzoyl glutamate (with carboxypeptidase G2); Cephalosporin nitrogen mustard carbamates (with beta-lactamase); SR4233 (with P450 Reductase); Ganciclovir (with HSV thymidine kinase, Borrelli et al 1988 Proc Natl Acad Sci 85: 7572-7576); mustard pro- drugs with nitro reductase (Friedlos el al 1997 J Med Chem 40: 1270-1275) and Cyclophosphamide (with P450 Chen et a/1996 Cancer Res 56: 1331 -1340).

Examples of pro-drug activation enzymes include a thymidine phosphorylase which activates the 5-fluoro-uracil pro-drugs capcetabine and furtulon; thymidine kinase from Herpes Simplex Virus which activates ganciclovir; a cytochrome P450 which activates a pro-drug such as cyclophosphamide to a DNA damaging agent; and cytosine deaminase which activates 5- fluorocytosine. An enzyme of human origin may be used.

Other suitable molecules include those that are of therapeutic and/or diagnostic application such as, but are not limited to: sequences encoding cytokines, chemokines, hormones, antibodies, engineered immunoglobulin-like molecules, a single chain antibody, fusion proteins, enzymes, immune co-stimulatory molecules, immunomodulatory molecules, anti-sense RNA, a transdominant negative mutant of a target protein, a toxin, a conditional toxin, an antigen, a tumour suppressor protein and growth factors, membrane proteins, vasoactive proteins and peptides, anti-viral proteins and ribozymes, and derivatives thereof (such as with an associated reporter group). When included, such coding sequences may be typically operatively linked to a suitable promoter, which may be a promoter driving expression of a ribozyme(s), or a different promoter or promoters, such as in one or more specific cell types.

The molecules may be proteins which are secreted from the cell. Alternatively the molecules are not secreted and are active within the cell. In either event, the molecules may demonstrate a bystander effector or a distant bystander effect; that is the production of the expression product in one cell leading to the killing of additional, related cells, either neighbouring or distant (e.g. metastatic), which possess a common phenotype.

Suitable molecules for use in the treatment or prophylaxis of cancer include proteins (or nucleic acids encoding proteins) which: destroy the target cell (for example a ribosomal toxin), act as: tumour suppressors (such as wild-type p53); activators of anti-tumour immune mechanisms (such as cytokines, co-stimulatory molecules and immunoglobulins); inhibitors

of angiogenesis; or which provide enhanced drug sensitivity (such as pro-drug activation enzymes); indirectly stimulate destruction of target cell by natural effector cells (for example, strong antigen to stimulate the immune system or convert a precursor substance to a toxic substance which destroys the target cell (for example a prodrug activating enzyme). Encoded proteins could also destroy bystander tumour cells (for example with secreted antitumour antibody-ribosomal toxin fusion protein), indirectly stimulate destruction of bystander tumour cells (for example cytokines to stimulate the immune system or procoagulant proteins causing local vascular occlusion) or convert a precursor substance to a toxic substance which destroys bystander tumour cells (eg an enzyme which activates a prodrug to a diffusible drug).

Antisense transcripts or ribozymes which interfere with expression of cellular genes for tumour persistence (for example against aberrant myc transcripts in Burkitts lymphoma or against bcr-abl transcripts in chronic myeloid leukemia) may be delivered to enhance cancer cell killing function or metastasis preventing function of the anti-intracellular oncoprotein antibody and/or intracellular oncoprotein vaccine. The use of combinations of such molecules is also envisaged.

Examples of hypoxia regulatable therapeutic molecules can be found in PCT/GB95/00322 (WO-A-9521927).

PHARMACEUTICAL COMPOSITIONS

The intracellular oncoprotein fragment vaccines may be effective in treating cancer related diseases.

We disclose a method of preventing a cancer related disease with vaccination with an intracellular oncoprotein fragment as a vaccine.

The intracellular oncoprotein fragments for use as vaccines may be provided as isolated and substantially purified proteins and protein fragments in pharmaceutically acceptable compositions using formulation methods known to those of ordinary skill in the art.

The intracellular oncoprotein fragment vaccine may be administered in the form of a pharmaceutical composition. Such a pharmaceutical composition may include a therapeutically effective amount of anti-intracellular oncoprotein antibody and/or intracellular oncoprotein vaccine, together with a suitable excipient, diluent or carrier.

The intracellular oncoprotein fragment vaccine may in particular be introduced into the circulation of a patient, for example by being injected into a patient via, e.g., a vein.

Formulations suitable for parenteral administration include aqueous and non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation isotonic with the blood of the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example, sealed ampoules and vials, and may be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders, granules and tablets of the kind previously described.

These compositions can be administered by standard routes. These include but are not limited to: oral, rectal, ophthalmic (including intravitreal or intracameral), nasal, topical (including buccal and sublingual), intrauterine, vaginal or parenteral (including subcutaneous, intraperitoneal, intramuscular, intravenous, intradermaL intracranial, intratracheal, and epidural) transdermal, intraperitoneal, intracranial, intracerebroventricular, intracerebral, intravaginal, intrauterine, or parenteral (e.g., intravenous, intraspinal, subcutaneous or intramuscular) routes.

The intracellular oncoprotein fragment vaccine formulations may conveniently be presented in unit dosage form and may be prepared by conventional pharmaceutical techniques. Such techniques include the step of bringing into
association the active ingredient and the pharmaceutical carriers or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

In addition, the intracellular oncoprotein fragment vaccine may be incorporated into biodegradable polymers allowing for sustained release of the compound, the polymers being implanted in the vicinity of where drug delivery is desired, for example, at the site of a tumor or implanted so that the anti-intracellular oncoprotein antibody and/or intracellular oncoprotein vaccine is slowly released systemically. The biodegradable polymers and their use are described, for example, in detail in Brem et af (1. Neurosurg 1991 74:441-446). Osmotic minipumps may also be used to provide controlled delivery of high concentrations of anti-intracellular oncoprotein antibody and/or intracellular oncoprotein vaccine through

cannulae to the site of interest, such as directly into a metastatic growth or into the vascular supply to that tumor.

The intracellular oncoprotein fragment vaccine may be linked to cytotoxic agents which are infused in a manner designed to maximize delivery to the desired location. For example, ricin-linked high affinity anti-intracellular oncoprotein antibody and/or intracellular oncoprotein vaccine are delivered through a cannula into vessels supplying the target site or directly into the target. Such agents are also delivered in a controlled manner through osmotic pumps coupled to infusion cannulae.

Preferred unit dosage formulations are those containing a daily dose or unit, daily sub- dose, as herein above recited, or an appropriate fraction thereof, of the administered ingredient. It should be understood that in addition to the ingredients, particularly mentioned above, the formulations described here may include other agents conventional in the art having regard to the type of formulation in question.

The intracellular oncoprotein fragment vaccine conjugates may be administered in any suitable way, usually parenterally, for example intravenously or intraperitoneally, in standard sterile, non-pyrogenic formulations of diluents and carriers, for example isotonic saline (when administered intravenously). Once the intracellular oncoprotein fragment vaccine conjugate has bound to the target cells and been cleared from the bloodstream (if necessary), which typically takes a day or so, the pro-drug is administered, usually as a single infused dose, or the tumour is imaged. If needed, because the intracellular oncoprotein fragment vaccine conjugate may be immunogenic, cyclosporin or some other immunosuppressant can be administered to provide a longer period for treatment but usually this will not be necessary.

The dosage of the intracellular oncoprotein fragment vaccine described here will depend on the disease state or condition being treated and other clinical factors such as weight and condition of the human or animal and the route of administration of the compound.

Depending upon the half-life of the intracellular oncoprotein fragment vaccine in the particular animal or human, the intracellular oncoprotein fragment vaccine can be administered between several times per day to once a week. It is to be understood that the methods and compositions described here have application for both human and veterinary use. The methods described here contemplate single as well as multiple administrations, given either simultaneously or over an extended period of time.

The timing between administrations of the intracellular oncoprotein fragment vaccine conjugate and pro-drug may be optimised in a routine way since tumour/normal tissue ratios of conjugate (at least following intravenous delivery) are highest after about 4-6 days, whereas at this time the absolute amount of conjugate bound to the tumour, in terms of percent of injected dose per gram, is lower than at earlier times.

Therefore, the optimum interval between administration of the intracellular oncoprotein fragment vaccine conjugate and the pro-drug will be a compromise between peak tumour concentration of enzyme and the best distribution ratio between tumour and normal tissues. The dosage of intracellular oncoprotein fragment vaccine conjugate will be chosen by the physician according to the usual criteria. At least in the case of methods employing a targeted enzyme such as β- glucosidase and intravenous amygdalin as the toxic pro-drug, 1 to 50 daily doses of 0.1 to 10.0 grams per square metre of body surface area, preferably 1.0-5.0 g/m^2 are likely to be appropriate. For oral therapy, three doses per day of 0.05 to 10.0g, preferably 1.0-5.0g, for one to fifty days may be appropriate. The dosage of the intracellular oncoprotein fragment vaccine conjugate will similarly be chosen according to normal criteria, particularly with reference to the type, stage and location of the tumour and the weight of the patient. The duration of treatment will depend in part upon the rapidity and extent of any immune reaction to the intracellular oncoprotein vaccine conjugate.

The invention further provides methods of treatment in which a patient is induced to generate antibodies against an intracellular oncoprotein. Such methods involve administration of a fragment of that intracellular oncoprotein to a patient in need of such treatment. The peptide may be a fragment of an oncoprotein that comprises an oncogenic mutation. The method may comprise the step of identifying the presence of one or more oncoproteins or oncogenic mutations in a sample obtained from the patient, prior to administering the peptide comprising a fragment of the oncoprotein, or comprising the oncogenic mutation identified as present in the patient. The sample may be a tumor sample such as a biopsy, or may be another sample such as a tissue, urine, blood, mucus, or other sample. The presence of a particular oncoprotein or oncogenic mutation may be determined by western blot, or by an ELISA or PCT based method. Suitable methods are known in the art.

DISEASES

Intracellular oncoprotein fragment vaccine described here, for example in the form of pharmaceutical compositions, may be used in the prevention and/or treatment of cancer.

For the purposes of this document, the term "cancer" can comprise any one or more of the following: acute lymphocytic leukemia (ALL), acute myeloid leukemia (AML), adrenocortical cancer, anal cancer, bladder cancer, blood cancer, bone cancer, brain tumor, breast cancer, cancer of the female genital system, cancer of the male genital system, central nervous system lymphoma, cervical cancer, childhood rhabdomyosarcoma, childhood sarcoma, chronic lymphocytic leukemia (CLL), chronic myeloid leukemia (CML), colon and rectal cancer, colon cancer, endometrial cancer, endometrial sarcoma, esophageal cancer, eye cancer, gallbladder cancer, gastric cancer, gastrointestinal tract cancer, hairy cell leukemia, head and neck cancer, hepatocellular cancer, Hodgkin's disease, hypopharyngeal cancer, Kaposi's sarcoma, kidney cancer, laryngeal cancer, leukemia, leukemia, liver cancer, lung cancer, malignant fibrous histiocytoma, malignant thymoma, melanoma, mesothelioma, multiple myeloma, myeloma, nasal cavity and paranasal sinus cancer, nasopharyngeal cancer, nervous system cancer, neuroblastoma, non-Hodgkin's lymphoma, oral cavity cancer, oropharyngeal cancer, osteosarcoma, ovarian cancer, pancreatic cancer, parathyroid cancer, penile cancer, pharyngeal cancer, pituitary tumor, plasma cell neoplasm, primary CNS lymphoma, prostate cancer, rectal cancer, respiratory system, retinoblastoma, salivary gland cancer, skin cancer, small intestine cancer, soft tissue sarcoma, stomach cancer, stomach cancer, testicular cancer, thyroid cancer, urinary system cancer, uterine sarcoma, vaginal cancer, vascular system, Waldenstrom's macroglobulinemia and Wilms' tumor.

Intracellular oncoprotein vaccine described here, for example in the form of pharmaceutical compositions, can also be used in the treatment of cancer related disorders.

Such disorders include but not limited to: solid tumours; blood born tumours such as leukemias; tumor metastasis; benign tumours, for example hemangiomas, acoustic neuromas, neurofibromas, trachomas, and pyogenic granulomas; rheumatoid arthritis; psoriasis; ocular angiogenic diseases, for example, diabetic retinopathy, retinopathy of prematurity, macular degeneration, corneal graft rejection, neovascular glaucoma, retrolental fibroplasia, rubeosis; Osier- Webber Syndrome; myocardial angiogenesis; plaque neovascularization; telangiectasia; hemophiliac joints; angiofibroma; wound granulation; coronary collaterals; cerebral collateralsl arteriovenous malformations; ischemic limb angiogenesis; neovascular glaucoma; retrolental fibroplasia; diabetic neovascularisation; helicobacter related diseases, fractures, vasculogenesis, hematopoiesis, ovulation, menstruation and placentation.

EXAMPLES

Our results demonstrated that upon tumor-specific antigen challenge, host immunity could be stimulated to produce endogenous antibodies against the specific antigen leading to tumor inhibition. This concept of 'cancer vaccination' has long been a promising but challenging prospect[21]. Importantly, we found that antigen-induced-antibody therapy could achieve similar anti-tumor therapeutic efficacy compared to exogenously delivered antibodies. In addition to being more economical, we believe that the former approach may be more useful than the latter as we naturally have huge potential flexibility to generate high titers of antigen-induced anti-tumor antibodies within ourselves. Although using Oncoprotein' for vaccination does not seem a practical thought, we do believe this approach is worth for future investigation. An oncoprotein should sit properly at its native subcellular localization with its neighborhood partners for correct communication in order to preserve an oncogenic function in cancer cell signaling network. When an 'oncoprotein' was isolated from its native dynamic complexity of sub-cellular localization, it may lose its connections and unable to perform its normal biological roles in such an 'isolated' situation.

Our in vivo data reveal a hitherto unrecognized phenomenon that immune-therapies can target not only extracellular- but also intracellular-oncoproteins for anticancer activity.

Examples Section E1 (Examples 1 to 18). Proof-of-concept for targeting intracellular oncoproteins with antibody therapy or vaccination

Section E1 comprises Examples 1 to 18. Example 1 is an Introduction to Section E1 , Examples 2 to 10 are Materials and Methods for Section El , Examples 1 1 to 16 are Results for Section E1 , Example 17 is a Discussion of Section El , Example 18 is References for Section E1 .

Example 1. Introduction (Section E1)

Monoclonal antibodies (mAbs) have proven to be potent and highly specific biological agents against some of humanity's most deadly diseases[1]. To exploit the ability of antibodies to specifically bind to biological targets, significant research and development have focused on developing monoclonal antibodies against extracellular/cell surface oncogenic targets. The first wave of success from such molecular targeted cancer therapeutics was the FDA approval of the anti-vascular endothelial growth factor antibody bevacizumab (Avastin) and the anti-HER2/neu antibody trastuzumab (Herceptin) for use in treatment of colorectal and breast cancers respectively[2]. These are examples of antibodies which work conventionally against extracellular proteins. Unfortunately, the majority of cellular proteins is intracellular and has thus remained underexplored by the approach of antibody therapies[3] due to the

general view that intracellular locations are inaccessible to antibodies. However, numerous experimental findings and clinical observations since 1978 have suggested otherwise, immunologists have showed that penetration of auto-antibodies into living cells may be a common cellular phenomenon[4,5].

To prove the concept of possible immunotherapies against intracellular targets, we selected three intracellular targets for cancer treatments in this study, namely phosphatase of regenerating liver 3 (PRL-3), Enhanced Green Fluorescent Protein (EGFP), and polyomavirus middle T (mT) oncoprotein. PRL-3 is one of the three PRL-phosphatases identified between 1994 and 1998[6,7], and the PRL-1 , PRL-2, and PRL-3 form a subgroup of the protein tyrosine phosphatase (PTP) family[8]. PRLs are intracellular C-terminally prenylated phosphatases[9,10], and the localization of PRL-1 and PRL-3 to the inner leaflet of the plasma membrane and early endosomes have been revealed by EM immunogold labeling[11]. The PRL phosphatases represent an intriguing group of proteins being validated as biomarkers and therapeutic targets in human cancers. Upregulation of PRL-3 mRNA level was first found to tightly correlate with colorectal cancer metastasis in 2001 by the global gene expression profiles of metastatic colorectal cancer with that of primary cancers, benign colorectal tumors, and normal colorectal epithelium using serial analysis of gene expression (SAGE) technology[12]. PRL-3 protein levels also have been reported to be elevated in an average of 22.3 % of cancer samples (n=1008)[13]. Additionally, upregulation of individual PRLs have been reported to be correlated with numerous types of advanced human metastatic cancers when compared with their normal counterparts[14]. Thus, oncogenic PRL-3 was chosen as an ideal intracellular target for the study.

Secondly, to explore whether the antibody therapy could have a general application against other intracellular proteins, we chose a popular reporter protein, the cytosolic enhanced green fluorescent protein (EGFP) originally isolated from the jellyfish Aeguorea victoria. EGFP expresses itself as an intracellular protein with a nucleo-cytoplasmic localization pattern. Since EGFP is not expressed in the host tissues, when artificially overexpressed in cancer cells, EGFP serves as a cancer cell specific intracellular protein. This allowed us to check whether EGFP- directed antibody therapy would specifically eradicate EGFP-expressing tumors and show any non-specific undesired side effects to the host.

Finally, to test the principle of antibody therapy in another animal tumor model, we use a well known mammary cancer model of MMTV-PymT transgenic mice[15] carrying the middle T (mT) oncogene under the transcriptional control of the mouse mammary tumor virus

promoter/enhancer. Such transgenic mice have been widely used as excellent spontaneous tumor models for decades in the cancer research community to assess the relative contribution of the metastatic mammary tumor phenotype[16,17].

Vaccines are inexpensive and yet effective in treating existing cancer or prevent cancer development[18]. Such specific active immunotherapy of cancer, if successful, has clear advantages over passive immunotherapy. If an antibody could recognize its intracellular antigen we can expect that an intracellular antigen could be used to trigger natural antibody production by the host immune system to achieve a similar effect of the antibody therapy against cancer. Here, we extend our study to evaluate the reliability of vaccinations with purified PRL-3, EGFP and mT proteins for antigen-induced antibody therapies.

Example 2. Materials and Methods (Section E1): Cell lines and cell culture

Bl 6F0 (CRL-6322) and B l 6F10 (CRL-6475) murine melanoma cell lines (derived from the C57BL/6J strain) were purchased from the American Type Culture Collection (ATCC, Manassas, VA). Cells were grown in RPMI medium supplemented with 10% fetal bovine serum and antibiotics and maintained in 37°C incubator supplied with 5% $C0_2$.

Example 3. Materials and Methods (Section E1): Western blot analysis

Detailed steps were described in our previous study[22].

Example 4. Materials and Methods (Section E1): Antibodies

Mouse monoclonal antibodies (clone #318, IgGl) were generated in-house, as described previously[22]. EGFP mouse monoclonal antibody (sc-9996, IgG2a) and Polyoma virus-middle T antigen (PymT) rat monoclonal antibodies (sc-53481 , IgG2b) were from Santa Cruz biotechnology, Inc.

Example 5. Materials and Methods (Section E1): Construction of GST-PRL-3, GST-middle T plasmids, and preparation of the GST-fusion proteins

The detailed steps for generating GST-PRL-3 fusion proteins were described previously . To make GST-middle T: forward primer 5' gcggatecatggatagagttctgagcagagctgac 3' and reverse primer 5' ctgaattcctagaaatgccgggaacg 3' were used for PCR using pXJ40 vector (i.e. pXJ40-PyMT) as a template. The PCR fragments were digested with BamHI and EcoRI and ligated into respective sites of the pGEX-KG vector. The preparations of GST-fusion protein were described in detail previously[22].

Example 6. Materials and Methods (Section E1): Experimental metastasis assay[23]

All animal studies were approved by the Institutional Animal Care and Use Committee (1ACUC) and were carried out in accordance with the policies of Institute of Molecular and Cell Biology's Review Board (1RB), Singapore[18]. C57BL6 and Scid mice were from BRC (Biological Resources Centre. Agency for Science, technology and Research, Singapore), muMT mice were from The Jackson Laboratory, USA, Rag $2^{-/-}$ mice were from Taconic, USA, Rag $T^{1'}$ mice were from Taconic, USA. 8-week old mice were injected with one million cancer cells via their tail vein (day 1). The treated mice were administrated with mAbs via tail vein on day 3 and subsequent administration twice a week. 4-week old transgenic mice (PymT) were divided into two groups receiving anti-mT antibody or PBS, twice per week. Mice were sacrificed and all organs were inspected for the presence of macroscopic metastases. Metastatic tumors or organs were removed and fixed in 4% paraformaldehyde for histological analysis. Lung metastatic nodules were counted under a dissecting microscope. The numbers of mice used in antibody therapies in each experiment are summarized in Figure 5G.

Example 7. Materials and Methods (Section E1): Genotyping of MMTV-PymT mice

The MMTV-PymT mouse strain was purchased from Mouse Models of Human Cancers Consortium (MMHCC). Females fail to lactate so this strain is maintained by breeding heterozygous (Tg/+) males with FVB/N (+/+) wild type females. Tail tip DNAs were extracted with DNA digestion buffer (12 ml: 10.08 ml, 1.2 ml of I Ox GB* buffer, 0.6 ml 10 % Triton X-100, 0.12 ml of 2-mercaptoethanol). *Gitschier's Buffer (10 ml: 3.35ml of 2M Tris pH 8.8, 1.66ml of 1M $(NH_4)_2S0_4J$ .34ml of 0.5M $MgCl_2$,I .65ml MQ H_20). Forward primer P001 : 5'-cAA ATG TTG cTT GTc TGG TG-3' and reverse primer P002: 5'-GTc AGT cGA GTG cAc AGT TT-3' were used to PCR wild type 200bp fragment. Forward primer POOI : 5'-GGA AGc AAG TAc TTc AcA AGG G-3' and reverse primer P004: 5'-GGA AAG TcA cTA GGA GcA GGG-3 were used to PCR 566bp transgene fragment. The genotyping procedure has been described previously (http://inouse.ncicrf.gov/pi tocols).

Example 8. Materials and Methods (Section E1): Whole Mount Preparation and Carmine Alum Stain

All procedures below were carried out at room temperature. Abdominal mammary glands were excised during necropsy for whole mount preparations, spread on glass slides, and fixed in Carnoy's fixative (6 parts 100% ethanol, 3 parts chloroform, and 1 part glacial acetic acid) for 4 h. Subsequently, the tissue was washed in 70% ethanol for 15 min, and the ethanol was changed gradually to distilled water then was finally rinsed in distilled water for 5 min. Staining was carried out overnight in carmine alum stain. The tissue was then

dehydrated in graded alcohol solutions (70, 95, and 100%; 15 min each) and was cleared in two changes of xylene, mounted, and coverslipped using Permount. Whole mounts were observed under a Leica dissecting microscope (Leica Microsystems GmbH, Wetzlar, Germany), and digital images were recorded using a SPOT FLEX® color digital camera (Diagnostic Instruments, Inc. Sterling Heights, MI) using a SPOT software package (Version 4.5, Diagnostic Instruments, Inc. Sterling Heights, MI).

Example 9. Materials and Methods (Section E1): Immunization of C57BL6 mice and MMTV-PymT transgenic mice

Freund's Complete Adjuvant is the form that contains killed cells of Mycobacterium butyricum to enhance the immune response. The Complete Adjuvant is used in initial injections. The form that does not contain this bacterium is known as Freund's Incomplete Adjuvant that is used to boost in subsequence injections. 8-week old C57BL6 mice were immunized by intraperitoneal injection with a total volume of 200 µï Freund's Adjuvant: PRL-3 (20 µg) or EGFP (20 µg) in 100 µï saline mixed with 100 µï of complete adjuvant (Cat#77140, Pierce). 4- week old MMTV-PymT TG mice were immunized by intraperitoneal injection with a total volume of 200 µï Freund's Adjuvant: mT (10 µg) antigen in 100 µï saline mixed with 100 µï of complete adjuvant (Cat#77140, Pierce). The next two immunizations were injected with a total volume of 200 µï incomplete adjuvant (Cat#77145, Pierce). The second and third injections were done two-week intervals. Subsequently, 100-200 µï of tail bleed was collected in a heparin-coated capillary tube, plasma was prepared from the blood sample, and the antibody titer was measured by ELISA. The detailed steps of ELISA were described previously[22]. Mice with high titers of PRL-3, EGFP, or mT antibodies in their sera were selected for experiment and further analysis, The nurhbers of mice used for vacc inations are summarized in Figure 6D.

Example 10. Materials and Methods (Section E1): ELISA assay

PRL-3 and mT antigen stocks were made in carbonate-bicarbonate buffer with pH 7.4. 100 µï solution containing 50 ng of appropriate antigen was added to each 96-well plate and incubated at 4 °C overnight to coat the antigen onto the plate. The plate was then blocked with 3 % BSA in PBS containing 0.05 % Tween 20 for 1 hour at 37 °C and then washed three times with PBS. Blood serum (1.0 µï) from each mouse was diluted in 100 µï of blocking solution and added to each well, and then incubated for 2 hours at 37 °C. 100 µï of appropriate secondary antibody conjugated with HRP (Pierce, USA) diluted at 1 :5000 in PBS was added to each well and incubated at 37 °C for 1 hour. The plate was rinsed with PBS containing 0.05 % Tween-20 three times followed by 3 washes with sterile water. The substrate, 100 µï of Turbo-TMB™ (Pierce, USA)), was added to each well and incubated for

10 min at room temperature. The reaction was stopped by adding 100 µl of concentrated H2SO4. Absorbance was measured at 450 nm using a Dynatech MR7000 plate reader (Dynatech, USA).

Example 11. Materials and Methods (Section E1): Statistical analysis

Use Kaplan-Meier method to estimate cumulative survival rates for 'treated' verse 'untreated' mice. A p value of <0.05 was considered statistically significant.

Example 12. Results (Section E1): In C57BL6 wild type mice, PRL-3 mAb (IgGl) effectively retards metastatic tumors that express endogenous PRL-3 protein

We have previously demonstrated the efficacy of antibody therapy against intracellular PRL proteins in an immunocompromised nude mice metastatic tumor model[19]. To examine this therapeutic approach in a more relevant model for future clinical applications, we extended this study to examine if the antibody treatment could function in immunocompetent C57BL6 wild type mice. On day 1, mice (8-week old) were intravenously (i.v.) injected via lateral tail vein with C57BL6-derived B 16F0 (F0) melanoma cancer cells, which express endogenous PRL-3 (Figure 1 A, lane 1). On day 3, mice were divided into 'untreated' and 'treated' groups which then received two administrations of PBS or PRL-3 mAb respectively per week during the whole duration of the experiment (Figure IB). At the end of the experiment (day 17), 'untreated' mice displayed severe weight loss, multiple metastatic tumors in various organs including the lung, liver, adrenal gland, kidney and bone. In contrast, 'treated' mice appeared more active and healthier in appearance. The 'treated' mice group, but not the 'untreated' group, constantly gained body weight throughout the duration of the experiment, in line with a reduction in tumor metastases (Figure 1 C-D). To confirm the specificity of the PRL-3 mAb, an identical experiment was conducted in parallel, in which mice were intravenously injected with C57BL6-derived B16F10 (F10) cells, another melanoma cell line, which expresses very low levels of PRL-3 (Figure 1 A, lane 2). By the end of the experiment (on day 17), all F10 recipients had developed metastases in the ovary, adrenal gland, kidney, and bone, regardless of whether they had received PRL-3 mAb. Both groups of mice experienced significant loss in body weight and appeared weak/inactive (Figure I E-F), indicating that non-PRL-3 expressing F10 recipients had a poor response to PRL-3 antibody treatment. Note that F10 recipients develop more aggressive metastatic tumors in lungs (Figure IE) compared to lungs from F0 recipients (Figure 1C). These data suggest that the efficiency of the PRL-3 antibody treatment was tightly correlated with PRL- 3 expression status of the cancer cells but not with the degree of metastatic activities. The current data derived from immunocompetent wild-type C57BL6 mice thus substantiate our previous findings from inmmunocompromised nude mice[19].

Example 13. Results (Section E1): In B-cell-deficient mice, PRL-3 mAb (IgGl) fails to retard metastatic tumors that express endogenous PRL-3 protein

To investigate if lymphocytes are involved in the effect of antibody therapy observed, we employed two engineered strains of immunodeficient mice: muMT mice, which are C57BL6 mice homozygous for an inactivating mutation of the membrane exon of the mu chain gene. They cannot form a preBCR and are consequently devoid of mature B lymphocytes . We also employed Rag-2 mice that carry a germline mutation in which a large portion of the RAG-2 coding region is deleted, thus giving rise to homozygous mutants ($Rag2^{-/-}$) which are viable but fail to produce mature B or T lymphocytes[21]. Kaplan-Meier survival curves were used to compare 'treated' and 'untreated' mice among muMT-FO, C57BL6-F0, and C57BL6-F10 mice. Surprisingly, we did not observe any PRL-3 mAb therapeutic efficacy in repressing tumors in muMT-FO mice, with PRL-3 mAb therapy having no impact in extending the life-span for muMT-FO mice with a median survival of 19 days for 'treated' and 18.5 days for 'untreated' mice (Figure 2A, /? value= 0.8661). Similarly, no difference in tumor metastases was observed between 'treated' and 'untreated' groups in B- and T-cell deficient Rag2-F0 mice (data not shown). In contrast, PRL-3 mAb therapy prolonged the survival rates for C57BL6-F0 mice with a median survival of 21.5 days for 'treated' but 17 days for 'untreated' mice (Figure 2B, p= 0002). In agreement with the previous finding, PRL-3 mAb therapy did not prolong the overall survival of C57BL6-F10 mice with non-PRL-3 expressing tumors with a median survival of 17.5 days for 'treated' and 16.5 day for untreated mice (Figure 2C, >=0.535). To summarize the effectiveness of antibody therapy among the different mice backgrounds, we scored the ability for PRL-3 mAb to reduce the degree of metastatic tumor formation relative to untreated mice in each group. When the reduction of metastatic tumor formation by at least 70% (±10 %) was scored as an effective outcome, a clear beneficial effect was seen only in the C57BL6-F0 group but not in the muMT-FO (Figure 2D). Thus, PRL-3 antibody therapy likely requires mature B-cell function for anti-tumor activity.

Example 14. Results (Section E1): In C57BL6 mice, EGFP mAb (IgG2a) effectively blocks metastatic tumors that express EGFP protein

The intriguing ability of PRL-3 mAb to inhibit tumor metastasis motivated us to investigate whether the therapeutic strategy could be applied to other intracellular proteins as well. Since EGFP is a non-cancer related protein, we chose to use EGFP as an intracellular protein marker specifically expressed in cancer cells. As EGFP is not expressed in the host tissues, we predicted that EGFP antibody should have few undesired side effects in the animal model. Utilizing pooled F0 and F10 melanoma cancer cell lines heterogeneously

overexpressing exogenous EGFP protein (Figure 5A, Figure 5B), the EGFP-F0 and EGFP-F10 expressing cells were injected and treated with EGFP mAb following the same therapeutic schedule (Figure 5C). As predicted, both EGFP-FO and EGFP-F10 cells responded equally well to EGFP antibody treatment in terms of tumor regressions regardless of PRL-3 expression. Notably, discrete clusters of green fluorescent or black metastatic tumors were found in the lungs (Figure 5D-E, panels a, e). The non-EGFP expressing metastases in black thus served as internal negative controls to the green metastatic tumors in this model. Remarkably, EGFP mAb could specifically reduce numbers of EGFP-metastases but not that of non-EGFP-metastases (Figure 5D-E, panels e), indicating that only EGFP-expressing tumors respond to EGFP mAb therapy. Upon scoring the results based on metastasis tumor load reduction, we found that the efficacy of the EGFP mAb effect was tightly correlated with EGFP expression status in cancer cells (Figure 5G). Thus, antibody therapy depends on specific antibody-antigen interactions on (or within) cells for therapeutic efficacy.

Example 15. Results (Section E1): In MMTV-PymT transgenic mice, mT (rat IgG2b) mAb prevents the formation of mT-expressing mammary tumor progression

Having addressed that antibody therapy could work against both endogenous (such as PRL-3) and exogenous (such as EGFP) intracellular proteins, we asked if this approach could inhibit tumor formation in spontaneous tumor animal models. To this end, we chose popular MMTV-PymT transgenic mouse model of a spontaneous mammary tumorigenesis. MMTV-PymT transgenic mice express the polyomavirus middle T (mT) oncogenic protein, a DNA viral protein containing 421 amino acids which acts as a potent oncogene in the mammary epithelium. mT is represented linearly, tethered (21 aa) to the membrane at its carboxy-terminal end with a short KRSRHF motif likely exposed at the cell surface (too short to be an external epitope, in general). Therefore, most of this protein is facing the cytosolic compartment and has thus considered an intracellular protein[15]. Cells expressing the mT possess an enhanced metastatic potential[16]. All female carriers (genotype+/-) develop palpable mammary tumors (adenocarcinomas) at the age of 2-3 months (http://mouse.ncifcrf.gov/information/order livem ice.asp). Middle T expression is detected at high levels in male and female mammary glands, and the expression of mT oncogene is sufficient for mammary epithelial cell transformation. We used 44 heterozygous (+/-) young transgenic females-using RT-PCR to confirm their genotypes (Figure 3A) and then divided them into 'treated' (n=18) and 'untreated' (n=26) groups. To determine if mT antibody could reduce mT- expressing mammary tumor development, 'treated' mice were i.v. injected with mT antibody at 4-week of age, followed by two administrations of antibody weekly for the whole duration of the experiment for about 3-months (Figure 3B). In Figure 3C, 'untreated'

mice (#2, #6) and 'treated' mice (#7, #10) were selected as representative examples; an FVB/N non-transgenic female (#3) was used as a control for normal sizes of mammary glands (5 pairs). Histopathological examination of the mammary glands of 'untreated' mice exhibited irregular formation of side branches, enlarged terminal buds, and some large multi-iobular tumor masses (Figure 3D, a). In contrast, breast tissue from treated mice, displayed more normal growth and development compared with a normal breast from wild-type mouse (Figure 3D, b-c). Using the Kaplan-Meier method to estimate cumulative survival rates for 'treated' verse 'untreated' MMTV-PymT transgenic mice, we found that the mT Ab therapy could prolong the life-span of treated mice, leading an increased in the median survival for 'treated' MMTV-PymT mice to 19.5-week compared to 15 weeks for untreated mice (Figure 3E, /?=0.0018). In all, we found that 100% (26/26) of 'untreated' mice carried dramatic breast tumors, while only 16.6% (3/18) of mT antibody treated mice developed tumor-bearing breasts, with the rest showing significant reduction in the formation of metastatic breast tumors that express mT oncogene (Figure 3F, and Figure 5F). These results suggested that the extensive repression of spontaneous tumor formation could be achieved by treating MMTV-PymT mice with mT antibody alone.

Example 16. Results (Section E1): C57BL6 mice vaccinated with PRL-3 protein or EGFP resisted the formation of PRL-3- or EGFP-expressing tumors

Having successfully demonstrated how three distinct intracellular proteins (PRL-3, EGFP, and mT) could be targeted with their respective antibodies, we next asked if intracellular proteins could be targeted by natural antibodies generated by the immune response upon antigen challenge (vaccination). We proceeded to immunize 8-week old C57BL6 mice with three doses (20 g each) of PRL-3 antigen (n=16) or EGFP protein (n=14) at 2-week intervals (Figure 6A). At the end of the immunization, sera from 14-week old immunized C57B16 mice were tested for their antibody titers against PRL-3 antigen or EGFP protein by ELISA (data not shown). Successfully immunized mice were subsequently divided into two groups, i.v. injected with either EGFP-FO (PRL-3 positive) or EGFP-F10 (PRL-3 negative) melanoma cells. Compared to un-immunized mice (n=18) (Figure 6B, a-d), PRL-3-immunized mice displayed reduced metastatic tumors formed by EGFP-FO cancer cells in adrenal, ovary and lung (Figure 6B, e-g). This was expected as EGFP-FO melanoma cells express both EGFP and PRL-3 proteins, which should respond to both EGFP and PRL-3 antibody therapies. Remarkably, compared to metastatic lung from unimmunized-mice (Figure 6B, d), the PRL-3-immunized mice were not able to reduce the degree of aggressiveness of metastatic lung tumors formed by EGFP-F 10 melanoma cells (Figure 6B, h), which do not express PRL-3 protein (Figure 5A). Again, this suggested that antigen- induced, naturally produced PRL-3 antibodies have no impact on inhibiting non-PRL-3 expressing EGFP-F 10 tumors.

Elegantly, we showed that GFP-immunization could reduce metastatic tumors formed by both EGFP-FO and EGFPFl O cancer cells in the adrenal gland, ovary and lung (Figure 6B, i-1) regardless of PRL-3 expressing levels. Taken together, these , results suggested that the induction of anti-PRL-3 or anti-EGFP antibodies in the host immune system upon vaccination could specifically inhibit the formation of PRL-3- or EGFP-expressing tumors respectively.

Example 17. Results (Section E1): MMTV-PymT transgenic young females vaccinated with mT antigen prevented the formation of mammary gland mT tumors

To further confirm the possibility of cancer vaccination with intracellular proteins; we again employed the MMTV-PymT transgenic mammary tumor model. 38 FVB/N heterozygous females (4-week old) were divided into GST-immunized (n=3), GST-mT-immunized (n=17), and unimmunized (n=18) control groups. Mice were subjected to immunize with three doses (10 μg each) of GST or GST-mT antigen respectively, at 2-week intervals (Figure 4A). Such mice were examined by ELISA for the titers of mT antibody to confirm the existence of mT antibody in their blood circulation (see Figure 4B for representative example). Comparing to GST-immunized mice (#43, #44, #45), remarkably, mT-immunized mice (#37, #40, #41) exhibited a dramatic reduction in size and weight of mammary gland tumors (Figure 4C-D), with the average weights of breast tissues/mouse from GST-immunized, GST-mT-immunized, and wild-type control mice found to be 5.6g, 1.3g, and 0.6g, respectively (Figure 4D). The results of the vaccination experiment are summarized in Figure 6C-D. Kaplan-Meier survival curves demonstrated that the mT antigen could prolong the mean survival rates (p=0.0004) for immunized mice (19.5-week) compared to the unimmunized group (14.5-week) (Figure 4E), indicating a positive benefit of cancer vaccination in extending survival time dramatically. Collectively, the data suggest that antigen-induced host mT antibody could again effectively prevent spontaneous breast tumor formations by cancer cells expressing mT.

Example 18. Discussion (Section E1)

Despite advancements in extracellular protein targeting, intracellular protein targeting has received little attention in terms of antibody therapy. Our findings presented here document a unique proof-of-concept in which anti-cancer antibody therapy against intracellular proteins could dramatically reduce tumor progression in vivo.

Here, we demonstrated the efficacy of antibody therapy against two intracellular proteins (PRL-3 and EGFP) against murine melanoma metastases in wild type C57BL6 mice. Similar success was obtained against intracellular mT antigen in the MMTV-PymT transgenic mice

spontaneous tumor model as well. The therapeutic response observed appears to be isotype- independent, as IgG 1 (PRL-3 mAb), IgG2a (EGFP mAb), IgG2b (PymT rat mAb), and self- generated antibodies could effectively inhibit tumor progression. Proper antibody-antigen interactions seem to be the predominating factor; targeting tumors with non-related antibodies produced no beneficial response at all. Intriguingly, we also found that mature B-cells constitute an important component of the antibody therapy response. These results mirror a recent report in which B-cells were found to be essential to the anti-DR5 antibody therapy against colon adenocarcinoma[23].

Our results also demonstrated that upon tumor-specific antigen challenge, host immunity could be stimulated to produce endogenous antibodies against the specific antigen leading to tumor inhibition. This concept of 'cancer vaccination' has long been a promising but challenging prospect'[9]. Importantly, we found that compared to exogenously delivered antibodies, antigen- induced-antibody therapies can achieve similar anti-tumor therapeutic efficacy. We believe that it may be more useful and economical as we naturally have huge potential flexibility to generate high titers of antigen-induced antibodies ourselves. Although using Oncoproteins' for vaccination does not seem a practical thought, we do believe this approach is worth for future investigation. In order to preserve an oncogenic function in cancer cell signaling network, an Oncoprotein' should precisely locate at its native subcellular context coordinated with its neighborhood partners for correct communication. When an Oncoprotein' is isolated from its native dynamic complexity of sub-cellular localization, it may lose its connections and be unable to perform its normal biological roles.

The work presented here outlines a potential methodology for future clinical cancer treatments. First, primary tumors are removed and examined to identify of tumor-specific antigens. This is followed by the introduction of specific chimeric or humanized antibodies against the tumor antigen to eliminate disseminated cells, thus inhibiting micro-metastases formation[24]. This step may well prevent further spreading or relapse in cancer patients. Alternatively, for cancers that are tightly genetically predisposed, immunization of immunocompetent young susceptible family members with an antigen that is associated with the familial cancer could 'prime' the immune system against that oncoprotein. These endogenously stimulated antibodies would be long-lasting, and would neutralize cancer cells expressing that particular oncoprotein. We base this approach on our observation that a very small amount of mT antigen (10 μg) introduced into a young MMTV-PymT transgenic mice could activate host immune system and produce mT antibody to inhibit mT-expressing tumor formation, allowing asymptomatic survival for up to 4-months. Extending weeks of life-span in mice may be comparable to prolonging decades in humans.

Ultimately, the pre-clinical data presented here suggests a general application for cancer treatments by using exogenous and endogenous antibody therapy targeting both extra- and intra- cellular tumor specific-antigens. Since existing conventional clinical antibody therapy is costly, we urge researchers to look at further developing the robust antigen-induced-antibody therapy response investigated here. For greater therapeutic response, we feel that this approach could be combined together with other immune-stimulatory agents for best effect. With cancer research rapidly moving towards individualized cancer therapy, our strategy for specific targeting of intracellular (or extracellular) oncoproteins hold enormous promise for tailored cancer therapy in the future.

Example 19. References (Section E1)

1 . K. Imai and A. Takaoka, Comparing Antibody and Small-Molecule Therapies for Cancer. Nat Rev Cancer. 6(9), 714-727 (2006).

2. J. Cohen and A. Wilson, New Challenges to Medicare Beneficiary Access to mAbs. mAbs. 1(1), 56-66 (2009).

3. M. Baker, Upping the Ante on Antibodies. Nat Biotechnol. 23(9), 1065-1072 (2005).

4. A. Ruiz-Arguelles and D. Alarcon-Segovia, Penetration of Autoantibodies into Living Cells. Isr Med Assoc J. 3(2), 121 -126 (2001).

5. D. Alarcon-Segovia, A. Ruiz-Arguelles and E. Fishbein, Antibody to Nuclear Ribonucleoprotein Penetrates Live Human Mononuclear Cells through Fc Receptors. Nature. 271(5640), 67-69 (1978).

6. R. H. Diamond, D. E. Cressman, T. M. Laz, C. S. Abrams and R. Taub, Prl-1 , a Unique Nuclear Protein Tyrosine Phosphatase, Affects Cell Growth. Mol Cell Biol. 14(6), 3752-3762 (1994).

7. Q. Zeng, W. Hong and Y. H. Tan, Mouse Prl-2 and Prl-3, Two Potentially Prenylated Protein Tyrosine Phosphatases Homologous to Prl-1 . Biochem Biophys Res Comm. 244(2), 421 - 427 (1998).

8. A. Alonso, J. Sasin, N. Bottini, I. Friedberg, I. Friedberg, A. Osterman, A. Godzik, T.

Hunter, J. Dixon and T. Mustelin, Protein Tyrosine Phosphatases in the Human Genome. Cell. 117(6), 699-71 1 (2004).

9. X. Si, Q. Zeng, C. H. Ng, W. Hong and C. J. Pallen, Interaction of Farnesylated Prl-2, a Protein-Tyrosine Phosphatase, with the Beta-Subunit of Geranylgeranyltransferase II. J Bio Chem. 276(35), 32875-32882 (2001).

10. J. Wang, C. E. Kirby and R. Herbst, The Tyrosine Phosphatase Prl-1 Localizes to the Endoplasmic Reticulum and the Mitotic Spindle and Is Required for Normal Mitosis. J Biol Chem. 277(48), 46659-46668 (2002).

11. Q. Zeng, X. Si, H. Horstmann, Y. Xu, W. Hong and C. J. Pallen, Prenylation- Dependent Association of Protein-Tyrosine Phosphatases Prl-1 , -2, and -3 with the Plasma Membrane and the Early Endosome. J Biol Chem. 275(28), 21444-21452 (2000).

12. S. Saha, A. Bardelli, P. Buckhaults, V. E. Velculescu, C. Rago, B. St Croix, K. E. Romans, M. A. Choti, C. Lengauer, K. W. Kinzler and B. Vogelstein, A Phosphatase Associated with Metastasis of Colorectal Cancer. Science. 294(5545), 1343-1346 (2001).

13. H. Wang, L. A. Vardy, C P. Tan, J. M. Loo, K. Guo, J. Li, S. G. Lim, J. Zhou, W. J. Chng, S. B. Ng, H. X. Li and Q. Zeng, Pcbpl Suppresses the Translation of Metastasis- Associated Prl-3 Phosphatase. Cancer Cell. 18(1), 52-62 (2010).

14. J. A. Sager, S. Benvenuti and A. Bardelli, Prl-3: A Phosphatase for Metastasis? Cancer Biol Ther. 3(10), 952-953 (2004).

15. S. M. Dilworth, Polyoma Virus Middle T Antigen and Its Role in Identifying Cancer-Related Molecules. Nat Rev Cancer. 2(12), 951-956 (2002).

16. C. T. Guy, R. D. Cardiff and W. J. Muller, Induction of Mammary Tumors by Expression of Polyomavirus Middle T Oncogene: A Transgenic Mouse Model for Metastatic Disease. Mol Cell Biol. 12(3), 954-961 (1992).

17. Y. Husemann, J. B. Geigl, F. Schubert, P. Musiani, M. Meyer, E. Burghart, G. Forni, R. Eils, T. Fehm, G. Riethmuller and C. A. Klein, Systemic Spread Is an Early Step in Breast Cancer. Cancer Cell. 13(1), 58-68 (2008).

18. E. Gilboa, The Promise of Cancer Vaccines. Nat Rev Cancer. 4(5), 401 -41 1 (2004).

19. K. Guo, J. P. Tang, C. P. B. Tan, H. Wang and Q. Zeng, Monoclonal Antibodies Target Intracellular Prl Phosphatases to Inhibit Cancer Metastases in Mice. Cancer Biol Ther. 7(5), 750-757 (2008).

20. T. von der Weid, N. Honarvar and J. Langhorne, Gene-Targeted Mice Lacking B Cells Are Unable to Eliminate a Blood Stage Malaria Infection. J Immunol. 156(7), 2510-2516 (1996).

21. Y. Shinkai, G. Rathbun, K. P. Lam, E. M. Oltz, V. Stewart, M. Mendelsohn, J. Charron, M. Datta, F. Young and A. M. Stall, Rag-2 -Deficient Mice Lack Mature Lymphocytes Owing to Inability to Initiate V(D)J Rearrangement. Cell. 68(5), 855-867 (1992).

22. J. Li, K. Guo, V. W. C. Koh, J. P. Tang, B. Q. Gan, H. Shi, H. X. Li and Q. Zeng, Generation of Prl-3- and Prl-1 -Specific Monoclonal Antibodies as Potential Diagnostic Markers for Cancer Metastases. Clin Cancer Res. 11(6), 2195-2204 (2005).

23. Haynes NM, Hawkins ED, Li M, McLaughlin NM, Hammerling GJ, Schwendener R, Winoto A, Wensky A, Yagita H, Takeda K, Kershaw MH, Darcy PK, Smyth MJ. CD 1 1 c+ dendritic cells and B cells contribute to the tumoricidal activity of anti-DR5 antibody therapy in established tumors. J Immunol. 185(1):532-41 (2010).

24. B. Weigelt, J. L. Peterse and L. J. van 't Veer, Breast Cancer Metastasis: Markers and Models. Nat Rev Cancer. 5(8), 591 -602 (2005).

Examples Section E2 (Examples 20 to 38). Host Immunity is Crucial for the Anticancer Efficacy of a Chimeric Antibody Targeting Intracellular PRL-3 in Animal Models

Section E2 comprises Examples 20 to 38. Example 20 is an Introduction to Section E2, Examples 21 to 29 are Materials and Methods for Section E2, Examples 30 to 36 are Results for Section E2, Example 37 is a Discussion of Section E2, Example 38 is References for Section E2.

Example 20. Introduction (Section E2)

A century ago, the German chemist Paul Ehrlich proposed the concept of antibodies as "magic bullets". Indeed, monoclonal antibodies (mAbs) have proven to be nature's biological warheads against some of humanity's most deadly diseases[1,2]. In general, antibody targeted

therapy has much less toxicity than chemotherapy with small molecule inhibitors. Antibodies constitute the most rapidly growing class of human therapeutics and are ideal agents for recognizing and destroying malignant cells via the immune system. However, this therapeutic approach has been limited to surface or secreted proteins expressed by cancer cells[3,4], in part due to the assumption that antibodies are too large (150 kDa) to penetrate the cell membrane. As a consequence, a wide spectrum of intracellular oncoproteins remains unexplored in terms of an antibody therapy approach. However, the concept that intact antibodies are unable to penetrate into viable cells has been challenged by a plethora of experimental findings and clinical observations[5,6,7]. Over the past 30 years, immunologists have found that autoantibodies found in the serum of patients with different autoimmune diseases can somehow bind their respective intracellular antigens[6,7,8]. Although it is not certain that the antibodies to intracellular proteins that are found in autoimmune patients actually cause the disease, or even cell destruction, we nevertheless emphasize our main contribution of this work is that antibodies to intracellular protein can exert therapeutic effects.

PRL-1 (phosphatase of regenerating liver- 1), PRL-2, and PRL-3 represent an intriguing subgroup of the intracellular protein tyrosine phosphatases (PTP). Individual PRLs are oyerexpressed in a variety of cancer cell lines and cancer tissues when compared with their normal counterparts[9]. A recent important review well describes these PRL-PTPs in cancer progression[10]. PRLs are intracellular C-terminally prenylated phosphatases. The localization of PRL-1 and PRL-3 to the inner leaflet of the plasma membrane and early endosomes was revealed by EM immunogold labeling" [12]. In contrast, the mutant forms of PRLs that lack the prenylation signal are localized in the nuclei[13]. PRL-3 was first discovered as a metastasis-associate phosphatase linked to colorectal cancer (CRC) metastasis with the finding that it was the only gene overexpressed in 100% of liver metastasis of CRC[14]. Overexpression of PRLs has subsequently shown to have a causative role in promoting cancer metastases and they become potential unique targets for diverse cancer treatment[15]. However, as these phosphatases are intracellularly localized, the conventional approach using therapeutic antibodies would seem a daunting task. In our earlier study, we reported an unexpected observation that mouse monoclonal antibodies (mAbs) against PRL-1 and PRL-3 were able to block experimental metastasis of cancer cells overexpressing intracellular EGFP-tagged PRL-1 and PRL-3[16].

In this study, we further evaluate the reliability of such targeting strategy using a newly-generated chimeric PRL-3 antibody. Here we extend our earlier findings for potential future

clinical therapeutics against intracellular oncoproteins in five important aspects. Firstly, we generated and utilized clinically-relevant chimeric antibodies instead of mouse antibodies. Secondly, we treated mice harboring naturally-occurring human cancer cells that express endogenous PRL-3 instead of exogenous PRL-3 in Chinese hamster cells (CHO). Thirdly, we showed that depletion of nature killer (K) cells enhances tumor engraftment. Fourthly, using paired nude and scid mouse models, we discovered the crucial role of B-cells in determining the outcome of our antibody therapy. Finally, using fluorescent labeled antibodies to track antibody- tumor binding activities by IVIS live imaging system; we proposed two working models for the antibody therapy in treated and untreated mice. An evidence-based hitherto concept is proposed for a possible approach in targeting intracellular oncoproteins with antibody therapies. The results suggest that an evaluation of a wide spectrum of intracellular oncoproteins (such as phosphatases, kinases. transcription factors) as possible targets for anticancer therapy may be warranted.

Example 21. Materials and Methods (Section E2): Generation of specific PRL-3 human/mouse chimeric mAb (clone #318)

For PRL-3 chimeric mAb generation, total RNA was extracted from $6x10^6$ hybridoma cells (clone#318) [19] using the RNeasy Mini Kit (QIAGEN, cat#74104). The RNAs were then reverse-transcribed into cDNA using Superscript II RNase H (Invitrogen, Cat 18064-014). The resulting total cDNAs were used as templates to generate the 'universal variable region' using Ig- Prime Kits (Novagen, cat#69831 -3) for PCR (95°C-4^{0}C-72°C, 30 cycles). The PCR fragment was cloned into the PCRII-TOPO-Vector with a TA cloning kit (Invitrogen, part#45-0640). The PCR fragment was cut with Mfe I and Xhol, and then inserted into the respective sites of a human IgG I constant region expression vector-pCMV-human IgGI [17] to join the mouse variable region of heavy chain (clone #318) [18] with the human IgG 1 constant region. Similar PCR procedures were performed for the mouse variable region of the light chain (clone #318) with ends containing restriction sites for ApaLI and Pst 1 . The PCR fragment was cut with ApaLI and Pst I and then inserted into the respective sites of a human IgG I constant region expression vector containing the variable region of the heavy chain of clone #318. The complete construct was transiently transfected into 293T cells cultured in ultra-low IgG FBS (Gibco, 16250-078). The chimeric mAb was subsequently harvested from the culture supernatant and concentrated up to 40 times with centrifugal filter devices (Millipore, cat#UFC900596). The chimeric mAb was tested for its specificity by indirect immunofluorescence (IF) and Western blot analysis.

Example 22. Materials and Methods (Section E2): Cell lines and cell culture

HCT1 16 (CCL-247) human colorectal carcinoma cell line, H460 (NCI-H460) human non-small lung cancer cell line, A431 (CRL-1555) human epidermoid carcinoma cell line, B16F0 (CRL-6322), and B16F10 (CRL - 6475) mouse melanoma cell lines were purchased from the American Type Culture Collection (ATCC, Manassas, VA). A27.80 (Cat#931 12519) human ovarian cancer cell line was purchased from ECACC, UK. Cells were grown in appropriate media recommended by the suppliers.

Example 23. Materials and Methods (Section E2): Western blot analysis

Generation of mouse PRL-3 monoclonal antibody and Western blot procedures have been described previously[18]. GAPDH antibody was from Cell Signaling Technology (Beverly, MA).

Example 24. Materials and Methods (Section E2): Experimental Metastatic Assay in mice[20]

I $x10^6$ cancer cells were injected into the circulation of eight-week old nude mice (Jackson Labs, USA) via the tail vein on day 1 . Either chimeric PRL-3 mAb or mouse PRL-3 or PRL-1 mAbs was used to treat mice; the first antibody treatment was carried on day 3 post-cancer cell injection, followed by two administrations per week. For control untreated group, PBS was administrated via tail vein. All animal studies were approved by the Institutional Review Board of the IMCB, in strict compliance with rules and policies of the Animal Facility Center of The Agency for Science, Technology and Research (A* STAR), Singapore.

Example 25. Materials and Methods (Section E2): Depletion of NK cells

Anti-asialo GM 1 anti-serum (rabbit) was purchased from Wako Pure Chemical Industries, Ltd (Osaka 540-8605, Japan). The GM1 anti-serum (50 µl) was injected into the circulation of eight-week old nude mice (Jackson Labs, USA) via the tail vein 24 h before the experimental metastasis assay.

Example 26. Materials and Methods (Section E2): Statistical analysis

The Kaplan-Meier method was used to estimate cumulative survival rates, and differences in survival rates, p-values less than 0.01 were considered to be significant.

Example 27. Materials and Methods (Section E2): Antibody labeling and IVIS live imaging

Purified PRL-3 antibody was labeled with CF™750 Dye Antibody Labeling Kits (http://www.biotium.coin/product/product info/Newproduct/M ix-n-Stain Kits.asp). Labeled antibodies were injected via tail vein 1 hr before live imaging. Bioware Ultra Cell Line HCT

116-luc2 (www.caliperLS.com) is a luciferase expressing cell line which was stably transfected with firefly luciferase gene (luc2) under the human ubiquitin C promoter. HCT1 16-Luc2 cell line was established by using HCT1 16 human adenocarcinoma (ATCC, CCL-247™) and transducing lentivirus containing luciferase 2 gene under the control of human ubiquitin C promoter (pGL4 luc2). $1x10^6$ HCT 116-luc2 cancer cells were injected into tail veins of 8-week old nude mice (Jackson Labs, USA). Antibody was injected into treated mice via tail veins on day 3, follow by two antibody injections per week. After 7-week treatment, mice are injected by an intraperitoneal route with a luciferin solution (15 mg/ml or 30 mg/kg, in PBS, dose of 150 mg/kg) that is allowed to distribute in awaked animals for about 5-15 minutes. The mice are placed into a clear plexiglass anesthesia box (2.5-3.5 % isofluorane) that allows unimpeded visual monitoring of the animals using IVIS® Spectrum Imaging System 3D Series to track and monitor tumor development in vivo. The results between treated verse untreated mice were determined.

Example 28. Materials and Methods (Section E2): FACS analysis

The human epidermoid carcinoma cell line A431 was grown in DMEM with high glucose (4.5g/L), supplemented with 10 % FBS and 5 % antibiotic. B 16F0 and B 16F10 cells were grown in RPMI, supplemented with 10 % FBS and 5 % antibiotics. Cells ($5x10^6$) were dislodged from the dishes with non-enzymatic pre-warmed cell dissociation solution (Sigma, Cat# c-5914) and transferred to 5 ml polystyrene tubes and washed once with complete medium. The cells were then incubated with 1 µï EGFR (Genetech, USA) or 5 µï PRL-3 primary mouse antibodies[18] in 100 µï of complete medium for 1 hr at room temperature (RT). Cells were agitated every 15 mins to prevent clumping. Cells were then washed 2x with complete medium and incubated for I hr at RT with goat anti-mouse AlexaFluor 546 antibody (Invitrogen, USA), washed, and re-suspended in 1 ml complete medium prior to analysis using BD FACS Caliber. Raw data was processed using WinMDI ver2.8 software.

Example 29. Materials and Methods (Section E2): Histopathologic Analyses Using Immunohistochemistry (IHC)

Human lung tissue arrays 009-01-004; and CCOO-01 -006; CC04-01 -CC04 were purchased from Cybrdi, Inc. (Rockville, Maryland 20850 USA: http://cybrdi.com/index.php). Human AML bone marrow samples were obtained from the National University Hospital-National University of Singapore (NUH-N US) Tissue Repository with approval of the Institutional Review Board (IRB) of NUH-NUS for research uses. The use of all human tissue samples including commercial samples were approved by Institutional Review Board (IRB) of the Institute of Molecular and Cell Biology. We used Dako En Vision™ Systems K 1395 (Dako, Carpinteria, CA) to perform IHC analysis[18 19].

Example 30. Results (Section E2): Generation of PRL-3 mouse/human chimeric antibodies (clone #318)

We previously reported that PRL-3 or PRL-1 mouse mAbs could specifically target their respective intracellular PRL-3 or PRL-1 phosphatase to inhibit cancer metastases in nude mice[16]. In an attempt to translate our laboratory findings to clinical setting, we have engineered a mouse/human chimeric mAb against PRL-3 to minimize the potential antigenicity of the mouse mAb in human. Using recombinant DNA technology, we separately fused the constant domains of heavy or light chains of the human IgGI molecule[17] with the mouse variable regions of PRL-3 mAb (clone#318)[18] by transgenic fusion of the immunoglobulin genes (Figure 7A). The expression construct was transfected into Human Embryonic Kidney 293T cells to produce recombinant PRL-3 chimeric mAb that was then harvested from the culture medium and further concentrated. The antigen-binding specificity of the PRL-3 chimeric mAb was well conserved as confirmed by performing indirect immuofluorescence on DLD-1 cells that overexpress exogenous EGFP-PRL-3 (Figure 7B) and western blot analyses (Figure 7C). The PRL-3 chimeric mAb specifically recognized EGFP-PRL-3 (-48 kDa) and myc-PRL-3 (-21 kDa) (Figure 7C, lane 1 -2) but react with neither myc-PRL-1 nor myc-PRL-2 proteins (Figure 7C, lane 3-4). A 50% cell Inhibitory Cytotoxic concentration (IC50) was carried out in a mouse melanoma B 16F0 cells for the chimeric antibody and we observed no cellular toxicity in viability even at high concentrations (40 g/ml) when cells were cultured in 10% FBS medium under normal culture conditions (data not shown). Since PRL-1 , -2, and -3 are overexpressed in a broad range of human cancers[19], we anticipate that the PRL-antibodies are likely to have broad applications to block different types of PRLs-positive cancer spreading, especially in some lethal malignancies such as lung cancers and acute myeloid leukemiaJAML) that often relapse within short timeframes. Amongst lung cancers, we found PRL-3 is overexpressed in 31 % of squamous cell carcinoma and 26% of adenocarcinoma (Figure 1 A-C), two main subtypes of highly recurrent non-small cell lung carcinoma comprising 80% of human lung cancer (http://en.wikipedia.org/wiki/Lung_cancer#Non-small_cell_lung_carcinoma_.28NSCLC.29). We also found that PRL-3 is overexpressed in 35% (24 out of 69 cases) of AML (Figure 12D). A few injections of PRL-3 chimeric antibody therapies will clean up the leftover of micro-metastasis and circulating cancer cells after the surgery of PRL-3-positive cancers to prevent the recurrence.

Example 31. Results (Section E2): PRL-3 chimeric antibody effectively inhibits the metastatic tumors formed by B16F0 cancer cells that express endogenous PRL-3 but not by B16F10 cancer cells that do not express endogenous PRL-3

To find a clinically relevant animal model for PRL-3-associated cancers, we screened dozens of cancer cell lines for the expression of endogenous PRL-3 protein levels by Western blot analysis. Ideal cell line pairs for our animal models should present contrasting levels of endogenous PRL-3 and should have the ability to induce metastatic tumors in mice within short timeframes. We found two mouse melanoma cell lines B16F0 and B16F10 (F0 and F10) that could fulfill the criteria for the desired experiment. Although F10 cells are naturally more metastatic than F0 cells, we found that parental F0 cells express higher levels of endogenous PRL-3 protein than F10 cells (Figure 8A), suggesting that F10 cell metastatic activity might be no longer PRL-3 dependent. When we employed an experimental metastatic assay[20] in which cultured cancer cells were introduced into the circulation of nude mice by lateral tail vein injection, both F0 and F10 cell lines can rapidly form multiple metastatic tumors in mice within 17 days. Such aggressive in vivo metastasis models allow us to see the differences in efficacy shortly after antibody therapy between treated and untreated groups. On day 3 (the latest time we can delay in treatment) post cancer cell injections, chimeric-PRL-3 antibodies were administrated similarly via tail veins into the 'treated mice', followed by two subsequent administrations of the antibody per week (Figure 8B). At the end of the experiment, we found that the PRL-3 chimeric antibody could eradicate tumors formed by PRL-3 expressing F0 cancer cells; the metastatic tumors in multiple tissues were dramatically reduced in 'treated mice' at the end of the experiments (Figure 8C). The Kaplan-Meier survival analysis for F0 recipients will discuss later (Figure 1 OA, c). In a parallel experiment, the PRL-3 antibody had no effect in blocking metastatic tumors formed by F10 cancer cells that do not express PRL-3 protein. Hundreds of metastatic tumors were found in the lungs of both treated and untreated F10 cell recipients, and no obvious difference was seen between 'untreated mice' (Figure 8D, upper panel) and 'treated mice' (Figure 8D, lower panel). Furthermore, Kaplan-Meier survival analysis demonstrated that the PRL-3 antibody could not extend the survival time for 'treated' Fl 0 recipients (Figure 8E).

Example 32. Results (Section E2): PRL-3 chimeric antibodies effectively inhibit the formation of metastatic tumors formed by human cancer cells that express endogenous PRL-3, but not by cancer cells that do not express endogenous PRL-3

In addition to mouse F0 melanoma cells, we further found that HCT1 16-luc2, HCT-1 16 human colorectal cancer cell line, and A2780 human ovarian cancer cell line express endogenous PRL-3 protein (Figure 9A lanes: 1 -3). A2780 has also been reported as a PRL-

3 positive cell line previously [21]. As a control, we found a human non-small lung cancer cell line (NCI-H460) which does not express endogenous levels of PRL-3 (Figure 9A lane 4). Regardless PRL-3 positive or negative, the four human cancer cell lines can rapidly form metastatic tumors in nude mice within 1 -2 months respectively. HCT-1 16-luc2 is a luciferase expressing cell line which was stably transfected with firefly luciferase gene (luc2) in HCT-1 16 cells. The cell line was established by transducing lentivirus containing luciferase 2 gene under the control of human ubiquitin C promoter. This cell line can be used in vivo by Xenogen's IVIS® Spectrum Series imaging to monitor tumor formation. Using this system, we demonstrated that the PRL-3 antibody could (. inhibit metastasis of HCT1 16-luc2 cancer cells as a clear reduction of metastatic lung tumors in ' live imaging at 7 week of antibody therapy was observed (Figure 9B). Furthermore, significant differences were found between PRL-3 antibody treated and untreated mice at 2-month post- inoculation with HCT-1 16 cells (Figure 9C, a) and at 1 -month post-inoculation with A2780 cells (Figure 9C, b). The PRL-3 mAb-treated animals appeared vibrant and healthy (up to 4-month), whereas the untreated mice had all lost weight and were moribund. In paralleled, NCI-H460 (PRL-3 negative cells) recipients did not respond to PRL-3 antibody treatment (Figure 9C, c). Taken together, these results further support the conclusion that the efficiency of the PRL-3 antibody treatment is tightly correlated with PRL-3 expression status of the cancer cells. We showed that the efficiency of both mouse- and chimeric-PRL-3 antibody therapies is comparable (Figure 9D).

Example 33. Results (Section E2): NK cells are important in anticancer therapy

We then investigated if nature killer (NK) cells play a role in the PRL-3 antibody therapy since NK cells are a type of cytotoxic lymphocyte that constitute a major component of the innate immune system. To deplete the nude mice's NK cells, we pre-injected nude mice with anti-asialo GM-1 antibody[22]. Our above-mentioned procedures of the antibody therapy were performed in these GM 1 -injected nude mice. We found that the efficacy of therapies was essentially lost in GM-1 injected mice (Figure 13). Even worse, such GM 1 injected nude mice showed more severe tumors (in black)-bearing burden in lung, liver, adrenal, testis, and bone than un-injected ones, indicating that innate immune system is important in our antibody therapy. NK cells have been demonstrated to have a role in human hematopoietic stem cell graft rejection[22], removal of NK cells may result in abolishment of graft rejection of NK cell activities, leading tumor engraftment more successfully, therefore, GM1 injected 'PRL-3 -treated' mice were worse than GMI -un- injected 'PRL-3-untreated' mice in terms of tumor growth.

Example 34. Results (Section E2): B-cells may be important in mediating the therapeutic effects of PRL-3 antibodies

Thus far, we performed the PRL-3 antibody therapies in T-cell deficient nude mice. To address if B lymphocytes are critical in our antibody therapy model, we next performed antibody therapies in Severe Combined Immunodeficiency (scid) mice, which lack functional lymphocytes because they carry a deficiency that impairs rearrangement of separate gene elements of the immunoglobulin and T-cell antigen receptor genes, thus, disrupting the differentiation and maturation of both B- and T-lymphoeyte progenitor cells. They thus show a severe combined immunodeficiency affecting both B and T lymphocytes, although they have normal NK cells, macrophages, and granulocytes. For HCT-1 16-induced metastatic tumors, Kaplan-Meier analysis of survival curves of PRL-3 antibody treated versus untreated mice showed a statistically significant ($P < 0.001$) increase in life span of nude mice but not scid mice (Figure 10A, a-b). Intriguingly, we found a 45% increase in the median survival time for the treated HCT-1 16- injected nude mice (16 weeks) compared to untreated mice (1 1 weeks). In contrast, in untreated or treated groups of HCT-1 16-injected scid mice, we observed no difference in the effect of PRL- 3 antibody therapy, which showed similar median survival duration (1 1 weeks). Furthermore, we employed F0 PRL-3 positive cancer cell line to confirm this finding using similar strategies. In FO-injected nude mice (Figure 1 OA, c-d), PRL-3 antibody treatment increased median survival duration by 47% (24 days for treated, versus 17 days for untreated). Consistently, the antibody had no effect in untreated or treated FO-injected scid mice, with both groups showing similar median life-span (~17 days). The mechanism is independent of antibody species, as similar results were obtained using mouse or chimeric PRL-3 antibodies (Figure I OB, a-d). Collectively, the results from these genetic mouse models suggest that PRL-3 antibody efficacy against metastatic tumor formation is dependent on host B-cell but not T cells. We suggest that other than antibody production, B-cell may have additional function to secrete unknown factor(s) that could facilitate the actions of antibodies. Indeed, our preliminary data shows that B-cells may secrete factors that could facilitate antibody uptake by the cancer cells. We observed a substantial increase in the internalization of antibody in PRL-3 positive F0 cells when the assay was performed in presence of culture supernatant from human B cells. Parallel, PRL-3 null F10 cells were unable to harbor any antibody either in presence of the B cell conditioned media or in presence of live B cells (data not shown).

Example 35. Results (Section E2): PRL-3 antigen is insignificant at the cell surface

A possible mechanism of PRL-3 antibody action could be the binding of a cell surface antigen, triggering a B-cell dependent elimination of the PRL-3 expressing cell. However, to date, no reports describe a cell surface localization of PRL-3. To address the possibility of

PRL-3 antibody binding its antigen on the cell surface, we used a FACS assay routinely used in cell surface labeling. As a positive control, anti-EGFR antibody binding of the EGFR-overexpressing human epidermoid carcinoma A431 cell line was used for this assay. We found that incubation of A431 cells with anti-EGFR antibodies caused a distinct peak-shift in the FACS assay (Figure 1 1 A, a). However, no peak shift was observed for either F0 (PRL-3 positive) or F10 (PRL-3 negative) cells incubated with or without anti-PRL-3 antibody (Figure 1 1 A, b-c). These results imply that cell surface PRL-3 antigen, if any, was unlikely to be the major cause of antibody binding and uptake. However, since immune system constantly battles invaders, such as bacteria and viruses, and cancer cells in vivo, the cancer cells can be destructed and somehow expose their intracellular protein to immune system. If this is so, the same possibility may apply to other intracellular oncoproteins as well.

Example 36. Results (Section E2): IVIS live imaging-based working models

In our animal model systems, we found that antibody therapies were ineffective after 3- days post cancer cells injection, suggesting that the first 3-day may be sufficient for cancer cell engraftment in new tissues. Using IVIS system to track sites of metastatic tumors through labeled antibody-antigen binding, we herein proposed two working models (Figure 1 IB), a. Treated mice. 1. at early stages, cancer cells were in clusters, timely introduction of the antibodies could somehow capture antigen-specific cancer cells in the circulatory, or in the lymphatic system together with the help of lymphocytes (such as B, and NK cells) via innate immune system to destroy and remove these cancer cells. 2. Escaped cancer cells will be able to arrive at new organs for implantations; therefore, the constant delivery of the antibody to the circulation is important to unplug these incipient tumors and prevent them from further development, destructing and clearing them from the host immune system. 3. The incipient tumors in 'treated mice' have no chance to develop tumors fully and therefore were in open-stages exposed to immune system. In addition, we found higher blood vessel density in small tumors (H&E lung sections) than in large tumors. As such, fluorescent labeled antibodies could easily access via blood circulation to metastatic lung tumors in 'treated mice'. We therefore observed the 'treated' metastatic lungs with strong florescent labeled, b. Untreated mice. The above-mentioned early steps had been missing; the unchecked cancer cells were rapidly multiplied in the host. The massive cancer cells were uncontrolled implanted and had sufficient time to develop micro- to macro-metastases, building up their territory with tumor boundaries 'fences' as defense system with close-stages away from the immune system. Furthermore, we found lower blood vessel density in large areas of tumor, insufficient blood supply results in necrosis (not shown). As consequences, fluorescent labeled antibodies were less accessible via blood supply to reach metastatic lung

tumors in 'untreated mice'. We therefore found the 'untreated' metastatic lungs with weak florescent labeled.

Example 30. Discussion (Section E2)

PRL-3 is up-regulated in numerous types of human cancer. It is foreseeable that the PRL- 3 chimeric antibody will be a promising therapeutic agent in blocking diverse PRL-3-associated cancer metastases. We hope to start with our effort on a few lethal malignancies (such as lung cancer and AML) that often relapse within short timeframes. Since these cancers are aggressive, therapeutic effects are easily observed and the outcomes between treated and untreated patients can be clearly defined. Especially, leukemic cells are easily accessible and are in direct contact with antibodies in the circulating system. The terminal patients suffering from these aggressive cancers are more likely to be recruited into clinical trials. Hopefully, success in these examples may make an impact on targeting other intracellular oncoproteins with antibody therapies, which is an important area of translational research and will provide a viable alternative treatment to cancer patients.

The molecular mechanism(s) behind these unconventional findings presented here remains elusive and is actively being investigated. Currently, understanding how antibody destroys tumors in vivo is still a 'dark box' to us. Similarly, it has been more than two decades to aim at understanding the mechanism(s) of the well-know anti-HER2/neu antibody therapy, although the anti-HER2/neu antibody has been in clinical uses for years in treating breast cancers[23]. Anticipating unraveling the 'dark box' using in vitro system may be unrealistic because a major consideration of traditional in vitro cell culture system is far simplified due to the limitation of single cell type grown in an incubator, typically using medium supplemented with 10% fetal bovine serum, lack of immune system involved, which was unlikely to be able to actually mimic in vivo complexities of multiple types of cells and organs, in 100% native blood circulation and lymphatic system. Whatever mechanism is, the central part of this study is that antibodies to intracellular protein can exert therapeutic effects, and more importantly, the effect is reproducible. If we can see the same therapeutic effect in humans that could be of enormous value in cancer treatment to support for the under-recognized notion that antibody therapy can be used for targeting intracellular antigens.

Nevertheless, our hard attempts in understanding the possible events within the 'dark box' enable us to achieve several conclusions from this study:

1. The PRL-3 antibody treatment is tightly correlated with PRL-3 expression status of the cancer cells. We demonstrate that the chimeric antibody could indeed successfully

block the formation of metastatic tumors derived from several cancer cell lines (B 16F0, HCT-1 16, and A2780) that express endogenous intracellular PRL-3 phosphatase. The inhibition is specific as the antibody had no effect in blocking the formation of metastatic tumors derived from other cancer cell lines (B16F10, H460) that do not express PRL-3. Previously, we had demonstrated that mouse PRL-3 antibody had no effect in inhibiting metastatic lung tumors formed by CT26 mouse colon cells, another PRL-3 negative cancer cell line[16]. Taken together, the data support the notion that the efficiency of either mouse or chimeric PRL-3 antibody treatment is tightly correlated with PRL-3 expression status of the cancer cells. If the metastatic property of cancer cells was not due to PRL-3 overexpression (such as B16F10, H460, CT26 cells), the administration of PRL-3 mAb has no effect in blocking tumors formed by these PRL-3 negative cells. Furthermore, we previously'[6] had shown that PRL- 1 mAb specifically blocks PRL-1 (but not PRL-3) metastatic tumors; while PRL-3 mAb specifically blocks PRL-3 (but not PRL-1) metastatic tumors though PRL-1 and PRL-3 share very high homology in protein sequence. These results imply that PRL antibody therapy is highly specific to its antigen and does not involve cross-reactivity with other non-specific cell surface
proteins.

2. NK cells in 'innate immune system' are involved in the therapy. To understand if innate immune system is involved in the antibody therapy, we intravenously injected GM1 antibody via tail vein into nude mice to deplete NK cells, which are a type of cytotoxic lymphocytes that constitute a major component of the innate immune system. In the absence of NK cells, we found, anti-PRL-3 antibody did not exhibit therapeutic effects. Furthermore, tumor- engraftment was dramatically enhanced (Figure 13), indicating that NK cells normally play a critical role in graft-rejection for implantation of foreign cells. Previously, mounting evidence suggests that NK cells play an important part in the destruction of incipient tumours[24].

3. ADCC may also be involved in the therapy. The best characterized mechanism of antibody therapy is antibody-dependent cellular cytotoxicity (ADCC). In ADCC, antibodies bind to specific cell surface antigens and trigger an Fc-mediated immune response involving cytotoxic CD8 T cells, complement activation, and/or NK cell activity. Although we did not observe any peak-shift in our FACS analysis of PRL-3 cell surface antigens in both PRL-3 positive F0 cells and PRL-3 negative Fl 0 cells, we could not rule out a possibility that these cancer cells were under abnormal inflammatory pressure, which may cause the destruction of cancer cells to release and expose their intracellular proteins to be attacked by the antibody and somehow trigger specific immune response, leading lymphocyte to remove

these cancer cells. Alternatively, in vivo, cancer cells are under hypoxic stress and serum deprivation, conditions that arrest cells at G] and Go phases[25]. It is possible that these conditions may cause release of intracellular antigens for the antibody to recognize. If so, other intracellular oncoproteins may also encounter the similar situations, they therefore can be similarly targeted with antibody therapies.

4. 'Adaptive immune system' (IgM and/or B Cells) are important in the mAb therapeutic effect. We employed athymic nude mice and immunodefficient scid mice for the tumour reduction experiments and achieved the therapeutic effect only in nude mice but not in scid mice. The major differences between nude and scid mice are that nudes are T-cell deficient, but have functional IgM antibodies and B cells, while scids have no functional adaptive immune system including B-, T-cells, and IgM and other antibodies. Both nudes and scids have intact innate immune system including NK cell and complement activity, but with positive responses seen only in the nude (but not in scid) mice, indicate that mature B cells (but not T cell), and possibly IgM/serum antibodies, are important. As the antibodies were exogenously introduced into mice, the requirement of B-cells for anti- metastatic PRL-3 antibody activity is hypothesized that an alternative role for B-cell in the antibody response, possibly via secretion of unidentified factor(s) that modulates any given antibodies for the host response. Collectively, we emphasize that intricate interplay of innate and acquired immune system is crucial for the anticancer efficacy of a chimeric antibody targeting intracellular PRL-3 oncoprotein.

5. Not all intracellular oncoproteins can be targeted with antibody therapy Desirable anti-cancer therapeutic agents should specifically target cancer cells while leaving normal tissues unharmed. It should be emphasized that the PRL-3 chimeric antibody therapy has little detectable side effect in nude mice since PRL-3 expression in normal tissues is not ubiquitous. We showed that PRL-3 protein was detected in only a few organs such as the spleen, brain, and pancreas (Figure 14A). In contrast, PRL-2 is ubiquitously expressed in most of mouse tissues (Figure 14B). As expected, PRL-2 antibody therapy to PRL-2 expressing cancers was unsuccessful (Figure 14C). The PRL-2 antibody-treated mice died 1 -week earlier or showed worse outcomes than untreated mice, which might be due to PRL-2's ubiquitous expression in most of mouse tissues, implicating a possibility of PRL-2 antibody may attack normal tissues to cause this undesired side-effect. The results suggest that a good therapeutic target should be more specifically to tumor antigen without harming host normal tissues.

6. Antibody therapy against intracellular oncoprotein is clinically relevant. To generate a very aggressive cancer model for treatment, we directly injected 1 million cancer cells into the circulation of a nude mouse; the total blood volume in nude mouse is 8% of its body weight (8% of ~19g = 1 .5ml). The total blood volume in a man (50 kg) is around 4.7 liters. If 1 million cancer cells were injected into -1.5 ml blood circulation in nude mouse; which is comparable to 3133 (4.7 liter/1.5 ml) millions cancer cells in ~ 4.7 liter blood in a man. The remarkable therapeutic effect by starting antibody (intravenously injected into tail vein) treatment even 3 days post- cancer cell injection indicates the treatment is intriguingly effective. If we can achieve similar life span extension in treated patients to that of mouse system, it will be comparable in years in humans in term of the total life span and the extended life expectancy. From these preclinical studies to reality, we anticipate that a few injections of chimeric antibody is very critical to reduce recurrence rate, as the PRL-3 antibodies will clean up circulating cancer cells[26] or unplugging invisible incipient tumors after surgical removal of visible PRL-3-associated solid tumors.

At this end, the lack of any observable side effect in nude mice upon PRL-3 antibody therapy further alludes to its potential clinical benefits. Our data prompts a reevaluation of a wide spectrum of tumor-specific intracellular oncoproteins as possible targets for anti-cancer mAb therapy, thus realizing the full potential of these 'magic bullets' .

Example 38. References (Section E2)

1 . D. Hanahan and R. A. Weinberg, The Hallmarks of Cancer. Cell. 100(1), 57-70 (2000).

2. J. M. Reichert, C. J. Rosensweig, L. B. Faden and M. C. Dewitz, Monoclonal Antibody Successes in the Clinic. Nat Biotechnol. 23(9), 1073-1078 (2005).

3. K. Imai and A. Takaoka, Comparing Antibody and Small-Molecule Therapies for Cancer. Nat Rev Cancer. 6(9), 714-727 (2006).

4. M. Baker, Upping the Ante on Antibodies. Nat Biotechnol. 23(9), 1065-1072 (2005).

5. Ruiz-Arguelles and D. Alarcon-Segovia, Penetration of Autoantibodies into Living Cells. Isr Med Assoc J. 3(2), 121 -126 (2001).

6. D. AlarcoA n- Segovia, L. Llorente, A. Ruiz-Arguelles, Y. Richaud-Patin and B. Perez-Romano, Penetration of Anti-DNA Antibodies into Mononuclear Cells Causes Apoptosis. Arthritis Rheum. 38, SI 79-182 (1995).

7. D. Portales-PeArez, D. AlarcoA n-Segovia, L. Llorente, R.-A. e. A, C. Abud- Mendoza, L. Baranda, H. De-la-Fuente, T. Ternyck, R. Gonzalez-Amaro and J., Penetrating Anti- DNA Monoclonal Antibodies Induce Activation of Human Peripheral Blood Mononuclear Cells. JAutoimmun. 11(5), 563-571 (1998).

8. T. D. Golan, A. E. Gharavi and K. B. Elkon, Penetration of Autoantibodies into Living Epithelial Cells. J Invest Dermatol. 100(3), 316-322 (1993).

9. B. J. Stephens, H. Han, V. Gokhale and D. D. Von Hoff, Prl Phosphatases as Potential Molecular Targets in Cancer. Mol Cancer Ther. 4(1 1), 1653-1661 (2005).

10. D. C. Bessette, D. Qiu and C. J. Pallen, Prl Ptps: Mediators and Markers of Cancer Progression. Cancer Metastasis Rev. 27(2), 231-252 (2008).

11. J. Wang, C. E. Kirby and R. Herbst, The Tyrosine Phosphatase Prl-1 Localizes to the Endoplasmic Reticulum and the Mitotic Spindle and Is Required for Normal Mitosis. J Biol Chem. 277(48), 46659-46668 (2002).

12. Q. Zeng, X. Si, H. Horstmann, Y. Xu, W. Hong and C. J. Pallen, Prenylation-Dependent Association of Protein-Tyrosine Phosphatases Prl-1, -2, and -3 with the Plasma Membrane and the Early Endosome. J Biol Chem. 275(28), 21444-21452 (2000).

13. X. Si, Q. Zeng, C. H. Ng, W. Hong and C. J. Pallen, Interaction of Farnesylated Prl-2, a Protein-Tyrosine Phosphatase, with the Beta-Subunit of Geranylgeranyltransferase Ii. J Biol Chem. 276(35), 32875-32882 (2001).

14. S. Saha, A. Bardelli, P. Buckhaults, V. E. Velculescu, C. Rago, B. St Croix, K. E. Romans, M. A. Choti, C. Lengauer, K. W. Kinzler and B. Vogelstein, A Phosphatase Associated with Metastasis of Colorectal Cancer. Science. 294, 1343-1346 (2001).

15. Q. Zeng, J.-M. Dong, K. Guo, J. Li, H.-X. Tan, V. Koh, C. J. Pallen, E. Manser and W. Hong, Prl-3 and Prl-1 Promote Cell Migration, Invasion, and Metastasis. Cancer Res. 63(1 1), 2716-2722 (2003).

16. K. Guo, J. P. Tang, C. P. B. Tan, H. Wang and Q. Zeng, Monoclonal Antibodies Target Intracellular Prl Phosphatases to Inhibit Cancer Metastases in Mice. Cancer Biol Ther. 7(5), 750-757 (2008).

17. J. H. Brendon, C. M. Adrianus and P. C. Angeline, Passive Immunoprophylaxis and Therapy with Humanized Monoclonal Antibody Specific for Influenza a H5 Hemagglutinin in Mice. Respir Res. 7, 126-136 (2006).

18. J. Li, K. Guo, V. W. Koh, J. P. Tang, B. Q. Gan, H. Shi, H. X. Li and Q. Zeng, Generation of Prl-3- and Prl-1 -Specific Monoclonal Antibodies as Potential Diagnostic Markers for Cancer Metastases. Clin Cancer Res 11(6), 2195-2204 (2005).

19. H. Wang, L. A. Vardy, C. P. Tan, J. M. Loo, K. Guo, J. Li, S. G. Lim, J. Zhou, W. J. Chng, S. B. Ng, H. X. Li and Q. Zeng, Pcbpl Suppresses the Translation of Metastasis- Associated Prl-3 Phosphatase. Cancer cell. 18(1), 52-62 (2010).

20. B. Weigelt, L. P. Johannes and L. J. v. t. Veer, Breast Cancer Metastasis Markers and Models. Nat Rev Cancer. 5(8), 591 -602 (2005).

21. F. Polato, A. Codegoni, R. Fruscio, P. Perego, C. Mangioni, S. Saha, A. Bardelli and M. Broggini, Prl-3 Phosphatase Is Implicated in Ovarian Cancer Growth. Clin Cancer Res. 11(19 Pt 1), 6835-6839 (2005).

22. H. Yoshino, T. Ueda, M. Kawahata, K. Kobayashi, Y. Ebihara, A. Manabe, R.Tanaka, M. Ito, S. Asano, T. Nakahata and K. Tsuji, Natural Killer Cell Depletion by Anti-Asialo Gml Antiserum Treatment Enhances Human Hematopoietic Stem Cell Engraftment in Nod/Shi-Scid Mice. Bone Marrow Transplant. 26(1 1), 121 1-1216 (2000).

23. M. Kasai, T. Yoneda, S. Habu, Y. Maruyama, K. Okumura and T. Tokunaga, In Vivo Effect of Anti-Asialo Gml Antibody on Natural Killer Activity. Nature. 291(5813), 334- 335 (1981).

24. S. Park, Z. Jiang, E. D. Mortenson, L. Deng, O. Radkevich-Brown, X. Yang, H. Sattar, Y. Wang, N. K. Brown, M. Greene, Y. Liu, J. Tang, S. Wang and Y. X. Fu, The Therapeutic Effect of Anti-Her2 Neu Antibody Depends on Both Innate and Adaptive Immunity. Cancer Cell. 18(2), 160-170 (2010).

25. S. Cooper, Reappraisal of Serum Starvation, the Restriction Point, GO and Gl Phase Arrest Points. FASEB J. 17(3), 333-340 (2003).
26. Y. Hiisemann, J. B. Geigl, F. Schubert, P. Musiani, M. Meyer, E. Burghart, G. Forni, R. Eils, T. Fehm, G. Riethmuller and C. A. Klein, Systemic Spread Is an Early Step in Breast Cancer. Cancer cell. 13(1), 58-68 (2008).

Examples Section E3 (Example 39). Immunization of C57BL6 Mice with VHZ and Monitoring Tumor Development

Example 39: Section E3: Immunization of C57BL6 Mice with VHZ and Monitoring Tumor Development

8-week-old C57BL6 mice were immunized by intraperitoneal injection with a total volume of 200 ml Freund's Adjuvant: 20 mg of VHZ antigen in 100 ml saline mixed with 100 ml of complete adjuvant (Cat#77140, Pierce).

The next two immunizations were injected with a total volume of 200 ml adjuvant: 20 mg of VHZ antigen in 100 ml saline mixed with 100 ml of incomplete adjuvant (Cat#77145, Pierce).

The second and third injections were administrated every 2 weeks, 100-200 ml of tail bleed was collected in a heparin-coated capillary tube.

Plasma was prepared from blood sample and antibody titer was measured by ELISA.

The detailed steps of ELISA were described previously.

Mice with high titers of VHZ antibodies in their sera were selected and lateral tail vein injected with 1 x106 VHZ-expressing cancer cells.

The results are shown in Figure 15. Figure 15 shows that mice vaccinated with VHZ antigen can prevent VHZ-expressing tumors. Unvaccinated mice serve as a negative control in this study.

Examples Section E4 (Examples 40, 41, 42).

Example 40: Section E4: immunization with Her2/neu peptide fragments

Generally, vaccine has been associated with infectious diseases but our immune system is also able to combat with cancerous cells. As such, inducing host immunity against oncoproteins will be a paradigm shift to maximize our immune system to fight against cancer cells for the therapeutic outcome. Multiple proteins/factors can cause a normal cell to become cancerous. However, if we are able to identify at least one of them (not necessary all of them), we are able to target at that protein with its specific antibody to kill the cancerous cells expressing that specific protein. Although one may feel uncomfortable to accept an idea of using 'oncoproteins' for vaccination, it is worth noting that the purified oncoproteins used as antigens in this study are unlikely to be oncogenic on their own because they are not sitting at their native subcellular location. Therefore, they are not able to interact with their native neighbourhood partners to elicit pathway-specific responses to drive tumorigenesis. These purified forms of "oncoproteins" have lost their specific intracellular spatiotemporal localization and therefore they are only proteins. Nonetheless, a more conservative approach would be the use of specific peptide antigens (partial oncoproteins) to achieve similar therapeutic results. Epitope-based peptide (or fragment) vaccination will be more specific and less cross-reacting with homologous proteins to reduce side effects.

In this study we used 4 Her2 fragments (see Figure 16):

1. Her2 extracellular fragment (Her2 extra-)
2. Her2 intracellular fragment (Her2 intra-)
3. Her2 C-terminal fragment (Her2 C-terminal)
4. Her2a short peptide (Her2 peptide)

As shown in Figure 17, mice immunized with Her2 fragments could inhibit the formation of Her2-expressing tumors formed by B16F0 melanoma cancer cells. At week 16, unimmunized, Her2 intra–immunized, or Her2 extra–immunized mice, and Her2 C-terminal immunized mice were then challenged with 1 million B16F0 cancer cells via tail veins. All the organs were examined at ~17 days after cancer cell injection and photographed to show morphologies of metastatic tumors. As shown in Figure 17(c, upper) ELISA analysis confirmed that Her2 fragment immunized mice had high levels of Her2 antibodies in their sera. As shown in Figure 17(c, lower), the total numbers of lung tumors (black tumors in panel B represent B16F0 melanoma cells) from each mouse was counted.

As shown in Figure 18, mice immunized with Her2a short peptide could inhibit the formation of Her2-expressing tumors formed by B16F0 melanoma cancer cells. At week 16, Her2 short peptide immunized, Her2 extra–immunized (immunized with Her2-extracellular fragment), and unimmunized mice, were then challenged with 1 million B16F0 cancer cells via tail veins. All the organs were examined at ~17 days after cancer cell injection and photographed to show morphologies of metastatic tumors. ELISA analysis was confirmed to show that Her2 short peptide immunized, Her2 extra fragment immunized (but not unimmunized) mice had high levels of Her2 antibodies in their sera (see panel C, upper). A total numbers of lung tumors from each mouse were shown in panel C, lower.

Example 41: Section E4: Immunization of MMTV-PymT transgenic mice with a fragment of mT oncoprotein

To apply a general concept of using target peptide vaccination, again, we examined a second tumor model, MMTV-PymT transgenic (TG) mice spontaneous tumor model, which carry the mT intracellular DNA viral protein under the transcriptional control of the mouse mammary tumor virus promoter/enhancer as a model of oncogene-induced mammary tumorigenesis (9). All female carriers (genotype+/−) develop palpable mammary tumors (adenocarcinomas) at the age of 3 months. The mT expression is detected at high levels in mammary glands, and the expression of mT oncogene is sufficient for mammary epithelial cell transformation (10). These TG mice have been widely used as excellent spontaneous tumor models for decades by the cancer research community (10, 18).

MMTV-PymT TG young females vaccinated with mT peptide (see Figure 16) reduce the formation of mammary tumors. mT-peptide immunized mice showed marked repression of tumors compared to GST-immunized mice (unrelated immunized mice as negative control), as shown in Figure 19 a small peptide of the mT oncoprotein achieved effective prevention from breast tumors progression in the model of MMTV-PymT transgenic mice.

Example 42: Section E4: Immunization of MMTV-PymT transgenic mice with a fragment of Estrogen Receptor

We further examined the MMTV-PymT transgenic (TG) mouse spontaneous tumor model described above. Estrogen Receptor was expressed at high levels in m-T mice, and particularly in breast tissue.

MMTV-PymT TG young females vaccinated with ER-Red and ER-purple peptide (See Figure 20) which are fragments of Estrogen Receptor. Both ER-Red and ER-purple immunized mice showed marked regression of tumors compared to GST-immunized mice (unrelated immunized mice as negative control) as shown in Figure 21. A small peptide of ER achieved effective prevention from breast tumor progression in the model of MMTV-PymT transgenic mice.

MMTV-PymT mice were injected with ERa-mAb weekly for 13 weeks. Treated mice showed a statistically significant reduction in breast tumor weight compared to untreated mice (see Figure 22).

Example 43: Section E4: Immunization of mice with HBV-X protein fragment

We are examining whether Hepatitis B X protein could be used to treat liver cancer. HBV X-protein is located in the nucleus of infected cells.

Panels of liver cancer cell lines that express X-protein are examined for their ability to form tumors in vivo. The lines with the highest tumor forming capability are used.

Mice are injected with anti-X-protein antibody to investigate the ability of the anti-X-protein antibody to block tumor formation from X-protein expressing liver cancer cells. Anti-X-protein antibody is administered to the mice according to the protocols outlined above.

Mice are injected with X-protein and X-protein fragments (see Figure 20) to trigger the host immune system to produce antibodies against X-protein to prevent the formation of tumors that express X-protein. Vaccination protocols as described above are used.

The spontaneous hepatocellular carcinoma (HCC) mouse model, which mimics human liver cancer development may also be used to examine the effects of anti-X-protein antibody, X-protein and X-protein fragment vaccination on hepatocellular carcinoma. The signalling pathways involved in HCC development are discussed in Signalling pathways and treatment of hepatocellular carcinoma is discussed in Whittaker S, Marais R, Zhu AX. The role of signaling pathways in the development and treatment of hepatocellular carcinoma. Oncogene. 2010;29:4989-5005.

Example 44: Section E4: Discussion

Genes specifically up-regulated during tumor formation but poorly or not expressed in host tissues are particularly promising as tumor-specific targets. For cancers that show a genetic link, immunization of immune-competent young susceptible family members with an antigen (epitope-based peptide vaccine) that is associated with the familial cancer could prime the immune system against that oncoprotein. These endogenously stimulated antibodies could then potentially combat cancer cells expressing that particular oncoprotein.

This study suggests that antibody-based therapy and vaccination against cancer may be extended to a wider variety of intracellular oncoproteins as therapeutic targets. The whole class of intracellular oncoproteins previously thought to be un-targetable by therapeutic antibodies or vaccinations can now expand the scope for tailor-made cancer therapies as well as usher in a new era of cancer vaccines. We expect that one potential advantage of using intracellular self-antigens is that they may have a better chance of provoking an immune response than extracellular self-antigens because immune cells targeting extracellular self-antigens are generally eliminated during development. We found that compared to exogenously delivered antibodies, antigen-induced antibody therapy could achieve similar antitumor therapeutic efficacy. Because existing conventional clinical antibody therapy is costly, vaccination may be more useful and economical as a means of inducing high titers of antigen-induced antibodies. This concept of "cancer vaccination" is promising and challenging.

This study indicates that antibodies against Hepatitis B virus (HBV) proteins (such as the HBV X-protein localized in the nucleus of infected cells) (Whittaker et al; Oncogene 29;4989-5005 (2010)) could be used as therapeutic intracellular targets for hepatocellular carcinoma (HCC) as most HCC are associated with HBV infection, antibody targeting viral protein will specifically destroy virally infected cells but leave normal cell unharmed. Similarly, for breast cancer caused by overexpression of estrogen receptor (ER), antibodies against ER or vaccination using ER fragment could be used to prevent spreading. This is particular useful to target ER positive breast cancer patients regardless of the expression of Her2 or other proteins.

A potential scheme in future cancer treatment could be to remove/biopsy of primary tumor, identify at least one tumor-expressing antigen using immunohistochemistry, and administer either antibodies against that specific tumor marker or a therapeutic vaccine that stimulates antibodies against it. For patients with a strong family history of cancer, immunization of

young susceptible family members with an antigen associated with the familial cancer could also prime the immune system against tumor cells expressing that antigen. If the myriad of previously unexplored candidate target proteins are investigated, a new era of tailor-made personalized cancer therapies will soon become reality.

Although the molecular mechanism(s) behind our untraditional findings remains elusive and needs to be deciphered, a number of possible mechanisms of antibody actions can be envisaged (Figure 22).Firstly (model A), the antibody may potentially enter PRL-3 expressing cells to target intracellular PRL-3 and neutralize its function. We observed that about ~10% of PRL-3 expressing cancer cells could take up antibodies in culture and this uptake was enhanced by 6-folds upon serum starvation.(Guo et al., Cancer Therapy 2008) Serum-starvation is used to arrest cells at G1 and G0 phases.(Cooper et al., FASEB J. 2003;17:333-340). It is possible that particular stages (perhaps G1-G0) of the cell cycle can contribute to the abilities of cells to take up the antibodies. *In vivo*, cancer cells are under hypoxic stress and serum deprivation, conditions that might enhance the abilities of cancer cells to take up antibodies. It was reported that antibody can be taken into live human mononuclear cells through surface Fc receptor (Alarcon-Segovia Nature 1978: 271;67-69). We found that the phenomenon of antibody uptake was abolished if endocytosis was blocked by NH4Cl or when cells were incubated mice (unpublished data), suggesting that most likely, these antibodies bind to cell surface proteins followed by endocytosis or pinocytosis. Once in the endocytic compartments, the antibodies could be released into the cytosol.

Secondly (model B), some of the intracellular antigens may be externalized and displayed on the surface of cancer cells by unconventional secretion (24), enabling the antibodies to bind and trigger immune responses such as antibody-dependent cellular cytotoxicity (ADCC).

Thirdly (model C), proteolytic fragments of intracellular targets may be presented by major histocompatibility (MHC) class I molecules to attract Cytotoxic T cells (CTLs) to mediate cell mediated lysis of cancer cells. In addition, a small fraction of intracellular antigens is released due to necrosis or cancer cell lysis, resulting in antigen-antibody complex within the tumor and stimulating local inflammatory response to attract immune cells in targeting neighbouring viable cancer cells within tumor. The most probable mechanism could be a combination of several modes, possibly including complement-mediated events that are actually involved in achieving the final therapeutic consequence of antibodies against intracellular oncoproteins. Currently, the understanding of how antibody destroys tumors *in vivo* is still a "dark box". This is similar to anti-HER2/neu antibody therapeutic mechanism; it

has been more than two decades and we still do not fully comprehend its mechanism(s) even though the antibody has been in clinical use for years to treat breast cancers (Hüsemann et al., Cancer Cell 2008; 13:58-68).

Using *in vitro* cell culture system to unravel or reassemble the *in vivo* "dark box" may be unrealistic because the traditional *in vitro* cell culture system is over simplified. There exists the limitation of single cell type grown in an incubator which typically uses medium supplemented with 10% (hence lacking 90%) fetal bovine serum and non-involvement of immune system. The *in vitro* system is also unlikely to actually mimic *in vivo* complexities of multiple types of cells and organs, coordinating in 100% native blood circulation and lymphatic system. Whatever mechanism is used, the central finding of this study is that antibodies to intracellular protein can exert therapeutic effects. Since the therapeutic effects of antibodies to intracellular proteins are substantial and reproducible, the incomplete understanding of the underlying mechanisms should not hinder future research in this new class of therapies.

Example 45: Immunization with oncopeptides designed around oncogenic mutations

C57BL/6 mice immunized with PRL-3 synthetic peptide using the vaccination protocol shown in Figure 24B. The synthetic peptide comprised two different peptides, both fragments of the complete PRL-3 protein, and linked with a GGSG linker (indicated in bold text), and had the sequence EVTYDKTPLEKDGITV**GGSG**DPHTHKTRC-KLH. As shown in Figure 24C immunization resulted in the production of mutant peptide specific antibodies.

Following immunization, mice were challenged with B16F0 cells which express PRL-3 target protein (see Figure 24A). Immunized mice were able to block tumor formation as compared to control mice which had not been immunized, or control mice immunized with GST (see Figure 24D).

In another experiment, C57BL6 mice were challenged with B16F10 cells. Unlike B16F0 cells, B16F10 cells do not express PRL-3. Mice challenged with B16F10 cells were unable to block tumor formation, demonstrating that immunization induces a specific immune response.

Example 44: Mutation peptide vaccination against oncoproteins

Although many human tumor cell lines carry mutated oncoproteins, it is not possible to introduce these into wild type mice, since mice cannot tolerate human cancer cells, and die due to the strong immune responses that are generated.

Conversely, only limited numbers of mouse cancer cell lines carry mutated oncoproteins available for use in the mouse vaccine models. We only identified one mouse colon cancer cell line, CT26, which carries Ras (G12D) mutation. This cell line is able to induce tumors in Balb/c mice.

The immunization and challenge protocol used in this experiment is shown in Figure 26A.

Immunisation of Balb/c mice with a peptide corresponding to region of the Ras oncoprotein containing the G12D mutation (CMTEYKLVVVGA**D**GVGKSALT) was able to trigger host production of antibodies against tumors that expressed that specific point mutation in a target protein (see Figure 26B).

Immunized mice were then injected with CT26 mouse cell line in a tumorigeneisis study. Immunization was able to prevent CT-26 tumor formation in Balbc mice Figure 26C).

Antibody therapy is costly; vaccination may be more useful and economical as a means of inducing high titers of antigen-induced antibodies. This concept of "cancer vaccination" is promising and challenging.

Each of the applications and patents mentioned in this document, and each document cited or referenced in each of the above applications and patents, including during the prosecution of each of the applications and patents ("application cited documents") and any manufacturer's instructions or catalogues for any products cited or mentioned in each of the applications and patents and in any of the application cited documents, are hereby incorporated herein by reference. Furthermore, all documents cited in this text, and all documents cited or referenced in documents cited in this text, and any manufacturer's instructions or catalogues for any products cited or mentioned in this text, are hereby incorporated herein by reference.

Various modifications and variations of the described methods and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in.connection with specific preferred

embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be within the scope of the claims.

CLAIMS

1. Use of a polypeptide in the manufacture of a medicament for the treatment of cancer, wherein the treatment involves stimulating the production of anti-intracellular oncoprotein antibodies in a subject in need thereof, the polypeptide comprising a fragment of an intracellular oncoprotein having between 10 and 150 amino acids, preferably between 10 and 100 amino acids, wherein the intracellular oncoprotein is PRL3.

2. Use according to claim 1 wherein the fragment comprises an oncogenic mutation.

3. Use according to claim 1 or claim 2 wherein the polypeptide comprises a plurality of fragments.

4. Use according to claim 3 wherein the polypeptide comprises a plurality of fragments of the same oncoprotein.

5. Use according to claim 4 wherein the polypeptide comprises a plurality of fragments from two or more oncoproteins.

6. Use according to any one of the preceding claims, wherein the treatment of cancer is for reducing cancer metastasis.

7. Use to any one of the preceding claims wherein the patient has a primary cancer that expresses the intracellular oncoprotein.

8. Use of a fragment of between 10 and 150 amino acids of an intracellular oncoprotein in the manufacture of a medicament for treating cancer metastasis in a patient with cancer, wherein the intracellular oncoprotein was identified to be specific to the patient's cancer, wherein the intracellular oncoprotein is PRL3.

9. Use of a fragment of an intracellular oncoprotein comprising between 10 and 150 amino acids in length in the manufacture of a medicament for treating cancer in a person, wherein the person was determined as suffering from, or likely to develop, a cancer associated with the intracellular oncoprotein before administration of the fragment or peptide to that person, wherein the intracellular oncoprotein is PRL3.

10. The use of claim 8 or claim 9 wherein the fragment comprises an oncogenic mutation.

11. The use of any one of claims 8 to 10 wherein the patient is administered a plurality of oncoprotein fragments.

12. The use of claim 11 wherein the plurality of oncoprotein fragments are combined into one peptide, each optionally linked by a peptide linker.

13. The use of any one of claims 8 to 12 wherein the patient is induced to produce antibodies against the peptide.

14. The use of any one of claims 9 to 13 wherein the person is to be administered with the intracellular oncoprotein fragment or peptide before he/she develops the cancer.

15. The use of any one of claims 8 to 14 wherein the person is to be administered with a further dose of the intracellular oncoprotein fragment or peptide.

16. The use of any one of claims 9 to 15 wherein the person has been identified as being likely to suffer from a cancer associated with an intracellular oncoprotein by assessing the prevalence of the cancer in family members.

17. The use according to any one of the preceding claims, in which the intracellular oncoprotein comprises a cytoplasmic or nuclear oncoprotein.

18. The use according to any preceding claim, in which the cancer is associated with or caused by over-expression of the intracellular oncoprotein.

19. The use according to any preceding claim, in which the cancer is selected from the group consisting of: breast cancer, hepatocellular cancer, acute lymphocytic leukemia (ALL), acute myeloid leukemia (AML), adrenocortical cancer, anal cancer, bladder cancer, blood cancer, bone cancer, brain tumor, cancer of the female genital system, cancer of the male genital system, central nervous system lymphoma, cervical cancer, childhood rhabdomyosarcoma, childhood sarcoma, chronic lymphocytic leukemia (CLL), chronic myeloid leukemia (CML), colon and rectal cancer, colon cancer, endometrial cancer, endometrial sarcoma, esophageal cancer, eye cancer, gallbladder cancer, gastric cancer, gastrointestinal tract cancer, hairy cell leukemia, head and neck cancer, Hodgkin's disease,

hypopharyngeal cancer, Kaposi's sarcoma, kidney cancer, laryngeal cancer, leukemia, leukemia, liver cancer, lung cancer, malignant fibrous histiocytoma, malignant thymoma, melanoma, mesothelioma, multiple myeloma, myeloma, nasal cavity and paranasal sinus cancer, nasopharyngeal cancer, nervous system cancer, neuroblastoma, non-Hodgkin's lymphoma, oral cavity cancer, oropharyngeal cancer, osteosarcoma, ovarian cancer, pancreatic cancer, parathyroid cancer, penile cancer, pharyngeal cancer, pituitary tumor, plasma cell neoplasm, primary CNS lymphoma, prostate cancer, rectal cancer, respiratory system, retinoblastoma, salivary gland cancer, skin cancer, small intestine cancer, soft tissue sarcoma, stomach cancer, stomach cancer, testicular cancer, thyroid cancer, urinary system cancer, uterine sarcoma, vaginal cancer, vascular system, Waldenstrom's macroglobulinemia and Wilms' tumor.

FIGURE 1



A
B16F0
B16F10
PRL-3
GAPDH
1
2
B
Inject cells
Analysis mice
Day: 1 3 6 9 12 15 17
PRL-3 mAb

FIGURE 1 (CONTINUED)

C




FIGURE 1 (CONTINUED)

D




FIGURE 1 (CONTINUED)

E




FIGURE 1 (CONTINUED)

F




FIGURE 2





FIGURE 2 (CONTINUED)

C



D




FIGURE 3





A
1 2 3 4 5 6 7 8 9 10
Genotype
+/- mT
566bp
200bp
GAPDH
B
Select (+/-)
mice
Examed
mice
Ages: 4w 5w 6w 7w 8w 9w 11w
mT Ab
C
Transgenic MMTV-MT (+/-)
Untreated
Treated with mT Ab
WT #3 (-/-)
TG #2
TG #5
TG #7
TG #10
TG #1
TG #6
TG #7
TG #10
WT #3

FIGURE 3 (CONTINUED)









FIGURE 4

A


Select (+/-) TG
4-week old
1st immunized
2-week
2nd immunized
2-week
3rd immunized
1-week
Sera Abs titers by Elisa
Examine breasts and other organs

B




Elisa tests mT antibody titers in sera
O.D.
0.45
0.35
0.25
0.15
0.05
0
#43
#44
#45
#37
#40
#41
Immunized:
GST
GST-mT

FIGURE 4 (CONTINUED)







FIGURE 4 (CONTINUED)






FIGURE 5




A
EGFP-B16F0
EGFP-B16F10
EGFP (43 kD)
PRL-3 (20 kD)
GAPDH (36 kD)
1
2
B
EGFP-F0
EGFP-F10
a
b
C
Inject cells
Analysis mice
Day: 1 3 6 9 12 15 17
EGFP- mAb
D
EGFP-F0
Untreated
Treated (GFP mAb)
E
EGFP-F10
Untreated
Treated (GFP mAb)
Lungs
Bones
Adrenal/ Kidneys
Ovaries

FIGURE 5 (CONTINUED)

F



G





FIGURE 6

A


1stAg
2ndAg
3rdAg
Cancer cells
Analysis mice
Age: 8w
10w
12w
14w
16w
Elisa

B





Unimmunized
PRL-3-immunized
GFP-immunized
a
e
i
Kidney/Adrenal
b
f
j
Ovary
c
g
k
Lung
d
h
l
Lung
EGFP-F0 Recepients
EGFP-F10 Recepients

FIGURE 6 (CONTINUED)




D




U=Unimmunized mice
I=Immunized mice

FIGURE 7

A Outline of PRL-3 chimeric antibody construction



B



C




FIGURE 8



A
B16F0
B16F10
PRL-3
GAPDH
B
Injected cells
Analysis mice
Days: 1 3 6 9 12 15 17
PRL-3 mAb
C
B16F0 (17 days)
Untreated
Treated
Adrenal
Livers
Bones
Abdomen tumors
No tumors
D
B16F10 (17 days) Untreated
Lungs
B16F10 (17 days) Treated
Lungs

FIGURE 8 (CONTINUED)

E




FIGURE 9




A
HCT116-Luc2
HCT116
A2780
NCI-H460
PRL-3
GAPDH
1 2 3 4
B
HCT116-Luc2 (7-week)
Untreated
Treated
C
HCT116 (2-month)
A2780 (1-month)
NCI-H460 (2-month)
Untreated
Treated
a
b
c
D
% of tumor-bearing mice
100
80
60
40
20
0
n=30
n=18
n=6
n=9
n=3
n=10
n=6
n=4
HCT116
A2780
H460
untreated
m318
h318

FIGURE 10

A

a



b





FIGURE 10A (CONTINUED)






FIGURE 10B



100
80
60
40
20
0
% of tumor -bearing mice
n=30
n=18
n=6
n=7
n=5
n=5
n=21
n=18
n=12
n=10
n=9
n=7
Nude - HCT116
Scid - HCT116
Nude - B16F0
Scid - B16F0
a
b
c
d
Unreated
R3-mAb
R3-hAb

FIGURE 11A



a
A431 Cells
no Ab
EGFR Ab
Events
FL2-H
no Ab
add EGFR Ab
b
B16F0 melanoma Cells
Events
FL2-H
no Ab
add PRL-3 Ab

FIGURE 11A (CONTINUED)




FIGURE 11B






FIGURE 12

A



B



C

Summary of PRL-3 expression in human lung cancers

Subtypes of lung cancers	Squamous cell carcinoma	Adeno-carcinoma	Small cell carcinoma	Large cell carcinoma	Carcinoid	Normal lung
Total samples	74	68	12	5	4	8
PRL-3 positive samples	23	18	1	1	1	0
% of PRL-3 positive samples	31.1 %	26.4 %	8.3 %	1/5	1/4	0/8

FIGURE 12 (CONTINUED)

D PRL-3 overexpressed in 35 % of AML




FIGURE 13




FIGURE 14



A
Heart
Colon
Spleen
Adrenal
Lung
Brain
Liver
Breast
Pancreas
Kidney
Salivary
Ovary
PRL-3
GAPDH
B
Breast
Brain
Spleen
9-day embryo
Kidney
Colon
Lung
Heart
PRL-2
GAPDH
C
PRL-2 Ab
Untreated
PRL-2 Ab
PRL-3 Ab
Died 1-week earlier

FIGURE 15



Unimmunized
Immunized with VHZ
Lung
Bone
Kidney
C57BL6 WT
8-week old
1st immunized
2-week
2nd immunized
2-week
3rd immunized
1-week
Challenging with cancer cells via tail veins
ELISA test Abs
3-week
16-week old
Examine organs

FIGURE 16

Her2 Target:

1. Her2 extracellular DNA sequences and peptide sequences:

ATCACAGGTCCCCTATACATCTCAGCATGGCCGGACAGCCTGCCTGACCTCAG
CGTCTTCCAGAACCTGCAAGTAATCCGGGGACGAATTCTGCACAATGGCGCCTA
CTCGCTGACCCTGCAAGGGCTGGGCATCAGCTGGCTGGGGCTGCGCTCACTG
AGGGAACTGGGCAGTGGACTGGCCCTCATC

ITGPLYISAWPDSLPDLSVFQNLQVIRGRILHNGAYSLTLQGLGISWLGLRSLRELGS
GLALI

2. Her 2 intracellular mid region DNA and peptide

ACCCACCAGAGTGATGTGTGGAGTTATGGTGTGACTGTGTGGGAGCTGATGAC
TTTTGGGGCCAAACCTTACGATGGGATCCCAGCCCGGGAGATCCCTGACCTGC
TGGAAAAGGGGGAGCGGCTGCCCCAGCCCCCCATCTGCACCATTGATGTCTAC
ATGATCATGGTCAAATGTTGGATG

THQSDVWSYGVTVWELMTFGAKPYDGIPAREIPDLLEKGERLPQPPICTIDVYMIMV
KCWM

3. Her2 intracellular C-Terminal

GAGAACCCCGAGTACTTGACACCCCAGGGAGGAGCTGCCCCTCAGCCCCACC
CTCCTCCTGCCTTCAGCCCAGCCTTCGACAACCTCTATTACTGGGACCGGGACC
CACCAGAGCGGGGGGCTCCACCCAGCACCTTCAAAGGGACACCTACGGCAGA
GAACCCAGAGTACCTGGGTCTGGACGTGCCAGTGTAA

ENPEYLTPQGGAAPQPHPPPAFSPAFDNLYYWDRDPPERGAPPSTFKGTPTAENP
EYLGLDVPV

4. Her2a intracellular peptide: VHHRHRSSSTRSGGGDLT


mT Target:

5. MMTV-PymT mT peptide sequence:

Cys-MDRVLSRADKERLLELLKLPRQLWGD

FIGURE 17

A


C57BL6 WT
8-week old
1st immunized
2-week
2nd immunized
2-week
3rd immunized
1-week
Challenging with cancer cells via tail veins
ELISA test Abs
3-week
16-week old
Examine organs

B




Unimmunized
Her-2 intra
Her-2 extra
C-terminal

FIGURE 17 (CONTINUED)

C







FIGURE 18

A


C57BL6 WT
8-week old
1st immunized
2-week
2nd immunized
2-week
3rd immunized
1-week
Challenging with cancer cells via tail veins
ELISA test Abs
3-week
16-week old
Examine organs

B




Her2a-peptide
Her2-extra
unimmunized

FIGURE 18 (CONTINUED)

C







FIGURE 19

A


Select TG (+/-)
~5-week old
1st Ag
2-week
2nd Ag
2-week
3rd Ag
2-week
Elisa
Examine organs

B




GST-immunized
mT peptide-immunized
FVB/N Wild-Type

FIGURE 19 (CONTINUED)

c






FIGURE 20

Targeting Estrogen Receptor with antibody and vaccination in PymT-transgenic mice model

Human ESR1 full protein sequence:

MTMTLHTKASGMALLHQIQGNELEPLNRPQLKIPLERPLGEVYLDSSKPAVYNYPEGAAYEFNAAAAANAQVY GQTGLPYGPGSEAAAFGSNGLGGFPPLNSVSPSPLMLLHPPPQLSPFLQPHGQQVPYYLENEPSGYTVREAGPP AFYRPNSDNRRQGGRERLASTNDKGSMAMESAKETRYCAVCNDYASGYHYGVWSCEGCKAFFKRSIQGHNDY MCPATNQCTIDKNRRKSCQACRLRKCYEVGMMKGGIRKDRRGGRMLKHKRQRDDGEGRGEVGSAGDMRAA NLWPSPLMIKRSKKNSLALSLTADQMVSALLDAEPPILYSEYDPTRPFSEASMMGLLTNLADRELVHMINWAKRV PGFVDLTLHDQVHLLECAWLEILMIGLVWRSMEHPGKLLFAPNLLLDRNQGKCVEGMVEIFDMLLATSSRFRMM NLQGEEFVCLKSIILLNSGVYTFLSSTLKSLEEKDHIHRVLDKITDTLIHLMAKAGLTLQQQHQRLAQLLLILSHIRHMS NKGMEHLYSMKCKNVVPLYDLLLEMLDAHRLHAPTSRGGASVEETDQSHLATAGSTSSHSLQKYYITGEAEGFPAT V

ER-red area:

Peptide: CGYTVREAGPPAFYRPNSDNRRQGGRE

Fragment:NRPQLKIPLERPLGEVYLDSSKPAVYNYPEGAAYEFNAAAAANAQVYGQTGLPYGPGSEAAAFGSNGL GGFPPLNSVSPSPLMLLHPPPQLSPFLQPHGQQVPYYLENEPSGYTVREAGPPAFYRPNSDNRRQGGRERLASTN DKGSMAMESAKETR

cDNA:aacggtccgcagctcaagatccccctggagcggccccttgggcgaggtgtacctggacagcagcaagcccgccgtgtacaactaccccg agggcgccgcctacgagttcaacgccgcggccgccgccaacgcgcaggtctacggtcagaccggcctcccctacggccccgggtctggaggctgc ggcgttcggctccaacggcctgggggtttccccccactcaacagcgtgtctccgagccgctgatgctactgcacccgccgccgcagctgtcgc ctttcctgcagcccacggccagcaggtgccctactacctggagaacgagcccagcggctacacggtgcgcgaggccggccgccggcattcta caggccaaattcagataatcgacgccagggtggcagagaaagattggccagtaccaatgacaagggaagtatggctatggaatctgccaagg agactcgc

ER-purple area:

Peptide: LKHKRQRDDGEGRGEVGSAGDMRAANLC

Fragment:GGIRKDRRGGRMLKHKRQRDDGEGRGEVGSAGDMRAANLWPSPLMIKRSKKNSLALSLTADQMVS ALL cDNA:

ggtgggatacgaaaagaccgaagaggaggaggagaatgttgaaacacaagcgccagagagatgatggggagggcaggggtgaagtgggggtct gctggagacatgagagctgccaacctttggccaagcccgctcatgatcaaacgctctaagaagaacagcctggccttgtccctgacggccgacc agatggtcagtgccttg


Targeting HBV-X protein with antibody and vaccination as liver cancer model in nude mice

HBV-X protein: maarlccqld ptrdvldlrp vgaesrgrpv sgplgdlpsp saspvptidr ahlslrglpvcafssagpca lrftsarnme ttvnthmilp kvlhkrtlgl pamstidlea yfkdclfkdweelgeeirlk vfvlggcrhk lvcspapcnf ftsa

HBV-X-peptide

FKDWEELGEEIRLKVFVLGGC (conserve sequence among the genotypes).

FIGURE 21


A
elect TG (+/-)
~5-week old
1st Ag
2-week
2nd Ag
2-week
3rd Ag
2-week
Elisa
Examine organs
B
LL/2
TrampC1
TrampC2
T breast
T Lung
Estrogen receptor
GAPDH
1
2
3
4
5



C
Immunized with below antigens:
GST (as control)
ER-purple
ER-red
GST
1 2 3
wt

FIGURE 21 (CONTINUED)

D




FIGURE 22




A
Brain
Colon
Lung
Kidney
Spleen
Liver
Stomach
Thymus
Muscle
Heart
Uterus
T. Breast
T. Lung
ERa
GAPDH
B
Select mice (+/-)
Analyze mice
Ages: ~6w 7w 8w 9w 10w 11w 12w 13w
ERa-mAb

FIGURE 22 (CONTINUED)

C



D




FIGURE 23



PRL-3
Antibody
MHC class I
PRL-3 fragment
nucleus
endosome
A
B
C

FIGURE 24




A
B16F0
B16F10
PRL-3
GAPDH
1
2
B
C57BL6 WT
8-week old
1st immunized
2-week
2nd immunized
2-week
3rd immunized
1-week
Challenging with B16F0 cells
ELISA test Abs
3-week
16-week old
Examine organs

FIGURE 24 (CONTINUED)

C



D





FIGURE 25


A
B16F0
B16F10
PRL-3
GAPDH
1
2




B
C57BL6 WT
8-week old
1st immunized
2-week
2nd immunized
2-week
3rd immunized
1-week
Challenging with B16F0 or B16F10 via tail veins
ELISA test PRL-3 Abs
3-week
16-week old
Examine organs

FIGURE 25 (CONTINUED)

C



D





FIGURE 26

A


Balbc mice
8-week old
1st immunized
2-week
2nd immunized
2-week
3rd immunized
1-week
Challenging with
CT-26 cancer cells
ELISA test Abs
3-week
16-week old
Examine organs

B



ELISA
1.6
1.4
1.2
1
0.8
0.6
0.4
0.2
0
unimmunized
Ras(peptide)

FIGURE 26 (CONTINUED)

C




FIGURE 27

A

1. MTYKVVVGA**D**GVGKSATNH **(K-Ras-G12D)**
2. MTYKVVVGA**V**GVGKSATNH **(K-Ras-G12V)**
3. MTYKVVVGA**R**GVGKSATNH **(K-Ras-G12R)**
4. MTYKVVVGAG**R**VGKSATNH **(K-Ras-G13R)**
5. QHVKITDFG**R**AKLLGAEEKE **(EGFR-L858R)**
6. KIGDFGLAT**E**KSRWSGSHQ **(B-Raf-V600E)**
7. LTSTVQLI**M**QLMPFGC **(EGFR-T790M)**
8. KVKIPVAIK-----TSPKANKEI **(EGFR-E746-A750 del)**
9. KAISTRDPLS**K**ITEQEKDF **(PI3KCA-E542K)**
10. TRDPLSEIT**K**QEKDFLWSH **(PI3KCA-E545K)**
11. EYFMKQMNDA**R**HGGWTTKMD **(PI3KCA-H1047R)**
12. DSGIHSGAT**A**TAPSLSGKGN **(beta-catenin-T41A)**
13. HSGATTTAP**F**LSGKGNPEE **(beta-catenin-S45F)**
14. HSGATTTAP**P**LSGKGNPEE **(beta-catenin-S45P)**
15. SDQDLLRC**C**VLTSGIFETK **(GNAS-R201C)**
16. RSQGRIPVKW**T**AIESLFDH **(Ret-M918T)**
17. QKNEFISE**F**CGEIISQDEAD **(EZH2-Y646F)**
18. QKNEFISE**N**CGEIISQDEAD **(EZH2-Y646N)**
19. QKNEFISE**S**CGEIISQDEAD **(EZH2-Y646S)**
20. QKNEFISE**H**CGEIISQDEAD **(EZH2-Y646H)**
21. FRMVDVGG**L**RSERRKWIH **(GNA11/GNAQ-Q209L)**
22. FRMVDVGG**P**RSERRKWIH **(GNAQ-Q209P)**

FIGURE 27 (CONTINUED)

ALK (F1174L) ALK (R1275Q)	K-Ras (G12D) (LS-174T, COLO-678, GP5d, LS-513, SNU-C2B) K-Ras (G12V) (SW620, RCM-1, SK-CO-1) K-Ras (G12C) (SW1463, SW837) K-Ras (G13D) (HCT-116, HCT-15, T84) The majority of pancreatic and colon cancer
BRAF (V600E) (SK-MEL-1, 3, 5, 24 and 28, A375, A2058, HT-144, MDA-MB-435) The majority of skin cancer	EGFR (L858R) (NCI-H1975) EGFR (E746_A750del) (PC-14, NCI-H1650) EGFR (T790M) (NCI-H1975) The majority of lung cancer
PIK3CA (H1047R) (T47D, BT-20) PIK3CA (E545K) (MCF7, MDA-MB-361) PIK3CA (E542K) (CAL-51) 20% breast cancer has PI3K mutations, corresponding to basal/triple negative? Not.	HNF1A/TCF1 (W206C) HNF1A/TCF1 (W206L)
CTNNB1/beta-catenin (S45F) (LS-174T) CTNNB1/beta-catenin (S45P) (NCI-H295) CTNNB1/beta-catenin (T41A) (LXF-289, A427) 20% mutations in liver cancer	GNA11 (Q209L) GNAQ (Q209L) GNAQ (Q209P) Eye, uveal melanoma (together, GNA11 and GNAQ are mutated in over 50%)
IDH1 (R132C) (HT-1080) IDH1 (R132G) IDH1 (R132H) (mostly in glioma) IDH1 (R132L) IDH1 (R132S) Bone, Chondrosarcoma Glioma	IDH2 (R140Q) IDH2 (R172K) haematopoietic neoplasm
EZH2 (Y646F) EZH2 (Y646N) EZH2 (Y646H) EZH2 (Y646S)	PPP2R1A (R183W) PPP2R1A (R183Q) haematopoietic and lymphoid tissue
PTEN (R130G) PTEN (R130Q) PTEN (R130*)	STK11 (LKB1) mutations suggest a tumor suppressor in lung cancer
Endometrium carcinoma	Notch1 (releasing cytoplasmic domain) Lymphoid neoplasm
N-Ras (Q61R) (HT-1197, KMOE-2, SK-MEL-2) RET (M918T) Mostly in thyroid cancer	GNAS (R201C) Mostly in pituitary cancer
AKT1 (E17K)	MET (N375S)
FGFR3 (K650E) FGFR3 (K650M) FGFR3 (R248C) FGFR3 (S249C) FGFR3 (Y373C) FGFR3 (S371C) FGFR3 (G370C)	Mostly in skin cancer

FIGURE 28

ABL1	D276G
ABL1	E255K
ABL1	E255V
ABL1	E352G
ABL1	E355G
ABL1	E373G
ABL1	E459K
ABL1	F311L
ABL1	F317L
ABL1	F359A
ABL1	F359V
ABL1	F382L
ABL1	F486S
ABL1	G250E
ABL1	G321E
ABL1	H396P
ABL1	H396R
ABL1	L248V
ABL1	L387M
ABL1	M237I
ABL1	M244V
ABL1	M343T
ABL1	M351T
ABL1	Q252E
ABL1	Q252H
ABL1	Q252R
ABL1	R47G
ABL1	S417Y
ABL1	T315I
ABL1	T315N
ABL1	T389A
ABL1	V304G
ABL1	V371A
ABL1	V379I
ABL1	Y253F
ABL1	Y253H
ABL1	Y353H
ABL2	E63Q
ABL2	R483I
ALK	L560F
ALK	Y262Y
APC	S2621C
ATM	E848Q
ATM	L1794L
ATM	M1916I
ATM	P2842R
ATM	Q2441P
ATM	T2666A
ATM	T935I
BRAF	D587A
BRAF	D587E
BRAF	D594E
BRAF	D594G
BRAF	D594K
BRAF	D594V
BRAF	E586E
BRAF	E586K
BRAF	F468C
BRAF	F583F
BRAF	F595L
BRAF	F595S
BRAF	G464E
BRAF	G464R
BRAF	G464V
BRAF	G466A
BRAF	G466E
BRAF	G466R
BRAF	G466V
BRAF	G469A
BRAF	G469E
BRAF	G469R
BRAF	G469S
BRAF	G469V
BRAF	G596R
BRAF	G606E
BRAF	G615R
BRAF	H608R
BRAF	I463S
BRAF	I582M
BRAF	I592M
BRAF	I592V
BRAF	K439Q
BRAF	K439T
BRAF	K475M
BRAF	K601E
BRAF	K601N
BRAF	L584L
BRAF	L597L
BRAF	L597Q
BRAF	L597R
BRAF	L597S
BRAF	L597V
BRAF	L618S
BRAF	N581S
BRAF	Q456Q
BRAF	R444R
BRAF	R444W
BRAF	R462I
BRAF	S605F
BRAF	S605N
BRAF	S614P
BRAF	S616P
BRAF	T440P
BRAF	T599I
BRAF	V459L
BRAF	V600D
BRAF	V600E
BRAF	V600G
BRAF	V600K
BRAF	V600L
BRAF	V600M
BRAF	V600R
BRAF	W604G
CEBPA	A152A
CEBPA	H191H
CEBPA	H84L
CTNNB1	A13T
CTNNB1	A183S
CTNNB1	A183T
CTNNB1	A20A
CTNNB1	A20V
CTNNB1	A21T
CTNNB1	A39A
CTNNB1	A39G
CTNNB1	A39T
CTNNB1	A43P
CTNNB1	A43T
CTNNB1	A43V
CTNNB1	D11N
CTNNB1	D17H
CTNNB1	D32A
CTNNB1	D32G
CTNNB1	D32H
CTNNB1	D32N
CTNNB1	D32V
CTNNB1	D32Y
CTNNB1	D56N
CTNNB1	D58N
CTNNB1	D6G
CTNNB1	E53K
CTNNB1	E54K
CTNNB1	E58G
CTNNB1	E58K
CTNNB1	E65E
CTNNB1	E67K
CTNNB1	G245A
CTNNB1	G34E
CTNNB1	G34R
CTNNB1	G34V
CTNNB1	G38A
CTNNB1	G38D
CTNNB1	G48D
CTNNB1	G48V
CTNNB1	G50D
CTNNB1	H24Y
CTNNB1	H36P
CTNNB1	H36Y
CTNNB1	I199L
CTNNB1	I35I
CTNNB1	I35N
CTNNB1	I35S
CTNNB1	I35T
CTNNB1	K19D
CTNNB1	K49E
CTNNB1	K49R
CTNNB1	L31L
CTNNB1	L46L
CTNNB1	L46V
CTNNB1	M8I
CTNNB1	N287S
CTNNB1	P44L
CTNNB1	P44S
CTNNB1	P52L
CTNNB1	Q61R
CTNNB1	S23R
CTNNB1	S29A
CTNNB1	S29F
CTNNB1	S29P
CTNNB1	S33A
CTNNB1	S33C
CTNNB1	S33F
CTNNB1	S33L
CTNNB1	S33N
CTNNB1	S33P
CTNNB1	S33S
CTNNB1	S33Y
CTNNB1	S37A
CTNNB1	S37C
CTNNB1	S37F
CTNNB1	S37P
CTNNB1	S37T
CTNNB1	S37Y
CTNNB1	S46A
CTNNB1	S45C
CTNNB1	S45F
CTNNB1	S45P
CTNNB1	S45S
CTNNB1	S45T
CTNNB1	S45Y
CTNNB1	S47G
CTNNB1	S47N
CTNNB1	S47R
CTNNB1	S47T
CTNNB1	S60F
CTNNB1	T214T
CTNNB1	T40A
CTNNB1	T40I
CTNNB1	T40T
CTNNB1	T41A
CTNNB1	T41I
CTNNB1	T41N
CTNNB1	T41P
CTNNB1	T41S
CTNNB1	T41T
CTNNB1	T42A
CTNNB1	T42I
CTNNB1	T42R
CTNNB1	T76A
CTNNB1	V22A
CTNNB1	V57M
CTNNB1	W25L
CTNNB1	W66L
CTNNB1	Y30C
EGFR	A750P
EGFR	A839T
EGFR	A859I
EGFR	A864I
EGFR	D761N
EGFR	E709A
EGFR	E709G
EGFR	E709H
EGFR	E709K
EGFR	E709V
EGFR	E746K
EGFR	E746V
EGFR	E866K
EGFR	G719A
EGFR	G719C
EGFR	G719S
EGFR	G735S
EGFR	G779F
EGFR	G810S
EGFR	G873E
EGFR	H773L
EGFR	H773R
EGFR	H835L
EGFR	H850N
EGFR	I715S
EGFR	I853T
EGFR	K846R
EGFR	L688P
EGFR	L707L
EGFR	L718L
EGFR	L730F
EGFR	L792P
EGFR	L798F
EGFR	L833V
EGFR	L838V
EGFR	L858M
EGFR	L858R
EGFR	L961Q
EGFR	N826S
EGFR	P694L
EGFR	P694S
EGFR	P733L
EGFR	P753S
EGFR	P772P
EGFR	R776C
EGFR	S720F
EGFR	S752Y
EGFR	S768I
EGFR	S784F
EGFR	T725M
EGFR	T751I
EGFR	T790M
EGFR	I847I
EGFR	V742A
EGFR	V769L
EGFR	V769M
EGFR	V774M
EGFR	V815V
EGFR	V851A
EGFR	V851I
EGFR	Y727C
EGFR	Y764Y
ERBB2	E914K
ERBB2	G776S
ERBB2	L755P
ERBB2	N857S
FGFR1	P252T
FGFR1	S125L
FGFR2	D283N
FGFR2	R203C
FGFR2	R496I
FGFR2	S267P
FGFR2	W290C
FGFR3	A369A
FGFR3	A391E
FGFR3	E322K
FGFR3	F384L
FGFR3	G370C
FGFR3	G382D
FGFR3	I414I
FGFR3	K650E
FGFR3	K650M
FGFR3	K650Q
FGFR3	K650T
FGFR3	R248C
FGFR3	S249C
FGFR3	Y241C
FGFR3	Y373C
FLI3	A680V
FLI3	D835E
FLI3	D835H
FLI3	D835N
FLI3	D835V
FLI3	D835Y
FLI3	G64G
HRAS	A11A
HRAS	A18I
HRAS	A59T
HRAS	G12A
HRAS	G12C

FIGURE 28 (CONTINUED)

HRAS	G12D
HRAS	G12R
HRAS	G12S
HRAS	G12V
HRAS	G13C
HRAS	G13D
HRAS	G13G
HRAS	G13R
HRAS	G13S
HRAS	G13V
HRAS	K117E
HRAS	Q61E
HRAS	Q61H
HRAS	Q61K
HRAS	Q61L
HRAS	Q61P
HRAS	Q61Q
HRAS	Q61R
HRAS	S17G
JAK2	V617F
KIT	A829P
KIT	D52N
KIT	D572Y
KIT	D816F
KIT	D816H
KIT	D816N
KIT	D816V
KIT	D816Y
KIT	D820G
KIT	D820V
KIT	E554D
KIT	E554G
KIT	E561E
KIT	E561K
KIT	E839K
KIT	F584S
KIT	G51G
KIT	G565R
KIT	I798I
KIT	I805I
KIT	K550I
KIT	K550N
KIT	K550R
KIT	K558K
KIT	K558N
KIT	K642E
KIT	K818R
KIT	L576P
KIT	L799L
KIT	L862L
KIT	M541L
KIT	M552K
KIT	M552L
KIT	N564K
KIT	N566D
KIT	N566N
KIT	N822H
KIT	N822K
KIT	N822N
KIT	P551S
KIT	P551T
KIT	P577S
KIT	P585P
KIT	S590N
KIT	T574I
KIT	V497V
KIT	V530I
KIT	V559A
KIT	V559D
KIT	V559G
KIT	V559I
KIT	V560D
KIT	V560E
KIT	V560G
KIT	V560V
KIT	V569A
KIT	V825A
KIT	W557G
KIT	W557R
KIT	W557S
KIT	Y553N
KIT	Y553V
KIT	Y563C
KIT	Y568D
KIT	Y568Y
KRAS	A11P
KRAS	A11V
KRAS	A18D
KRAS	A18T
KRAS	A59E
KRAS	A59T
KRAS	D92Y
KRAS	E62D
KRAS	G12A
KRAS	G12C
KRAS	G12D
KRAS	G12E
KRAS	G12F
KRAS	G12G
KRAS	G12L
KRAS	G12N
KRAS	G12R
KRAS	G12S
KRAS	G12V
KRAS	G13A
KRAS	G13C
KRAS	G13D
KRAS	G13G
KRAS	G13I
KRAS	G13N
KRAS	G13R
KRAS	G13S
KRAS	G13V
KRAS	G15D
KRAS	G15S
KRAS	G60A
KRAS	H27L
KRAS	K5N
KRAS	L19F
KRAS	Q22K
KRAS	Q22Q
KRAS	Q22R
KRAS	Q61E
KRAS	Q61H
KRAS	Q61K
KRAS	Q61L
KRAS	Q61P
KRAS	Q61R
KRAS	S17G
KRAS	V8V
KRAS	V9V
MAP2K4	Q142L
MET	A1209G
MET	D1246H
MET	E168D
MET	G1137V
MET	H1112L
MET	H1112R
MET	H1112Y
MET	H1124D
MET	K1262R
MET	L1130L
MET	L1213V
MET	L229F
MET	M1149T
MET	M1268I
MET	M1268T
MET	N1118Y
MET	N375S
MET	R988C
MET	S283S
MET	T1010I
MET	T1191I
MET	V1290L
MET	Y1248C
MET	Y1248H
MET	Y1253D
MSH6	D390D
MSH6	D390N
MSH6	D575Y
MSH6	E1198A
MSH6	G141G
MSH6	P1087P
MSH6	R249R
MSH6	S315F
MSH6	V474A
MSH6	V480L
MSH6	V509A
MSH6	Y397C
NOTCH1	A1702P
NOTCH1	A2280V
NOTCH1	A2564D
NOTCH1	D1610V
NOTCH1	F1541L
NOTCH1	F1593S
NOTCH1	H2506Y
NOTCH1	I1681N
NOTCH1	L1575P
NOTCH1	L1586P
NOTCH1	L1594P
NOTCH1	L1601P
NOTCH1	L1679P
NOTCH1	P2413T
NOTCH1	R1589P
NOTCH1	R1609S
NOTCH1	R1628H
NOTCH1	T2467M
NOTCH1	V1677D
NRAS	A11T
NRAS	A18T
NRAS	A59T
NRAS	G12A
NRAS	G12C
NRAS	G12D
NRAS	G12G
NRAS	G12P
NRAS	G12R
NRAS	G12S
NRAS	G12V
NRAS	G12Y
NRAS	G13A
NRAS	G13C
NRAS	G13D
NRAS	G13G
NRAS	G13R
NRAS	G13S
NRAS	G13V
NRAS	G13Y
NRAS	Q61E
NRAS	Q61H
NRAS	Q61K
NRAS	Q61L
NRAS	Q61P
NRAS	Q61Q
NRAS	Q61R
NRAS	S65C
NRAS	Y64N
NTRK1	L213L
NTRK3	H677Y
NTRK3	R679Q
NTRK3	R721P
PDGFRA	D1071N
PDGFRA	D842I
PDGFRA	D842V
PDGFRA	D842Y
PDGFRA	D846Y
PDGFRA	F808L
PDGFRA	N870S
PDGFRA	I674I
PDGFRA	V561D
PDGFRB	I882I
PIK3CA	A1046A
PIK3CA	A1046V
PIK3CA	A1066V
PIK3CA	C378R
PIK3CA	C420R
PIK3CA	C901F
PIK3CA	D350H
PIK3CA	E1034G
PIK3CA	E110K
PIK3CA	E418K
PIK3CA	E453K
PIK3CA	E453Q
PIK3CA	E542G
PIK3CA	E542K
PIK3CA	E542V
PIK3CA	E545D
PIK3CA	E545G
PIK3CA	E545K
PIK3CA	F909L
PIK3CA	G1049R
PIK3CA	G1049S
PIK3CA	G106V
PIK3CA	G118D
PIK3CA	G122D
PIK3CA	H1047L
PIK3CA	H1047R
PIK3CA	H1047Y
PIK3CA	H1065Y
PIK3CA	H510H
PIK3CA	H701P
PIK3CA	K111N
PIK3CA	K733R
PIK3CA	M1043I
PIK3CA	M1043T
PIK3CA	M1043V
PIK3CA	N1044K
PIK3CA	N345K
PIK3CA	N345S
PIK3CA	P1011S
PIK3CA	P104R
PIK3CA	P124T
PIK3CA	P539R
PIK3CA	Q546E
PIK3CA	Q546K
PIK3CA	Q546P
PIK3CA	Q546R
PIK3CA	Q60K
PIK3CA	Q661K
PIK3CA	R108P
PIK3CA	R38C
PIK3CA	R38H
PIK3CA	R537R
PIK3CA	R88Q
PIK3CA	S1008P
PIK3CA	S405F
PIK3CA	T1025A
PIK3CA	T1025N
PIK3CA	T1025S
PIK3CA	Y1021C
PIK3CA	Y1021N
PTEN	A121E
PTEN	A121P
PTEN	A126P
PTEN	A126T
PTEN	A126V
PTEN	A137V
PTEN	A148T
PTEN	A151T
PTEN	A309A
PTEN	A34A
PTEN	A39T
PTEN	A3S
PTEN	A79T
PTEN	C105F
PTEN	C105G
PTEN	C105S
PTEN	C105Y
PTEN	C124S
PTEN	C124Y
PTEN	C136F
PTEN	C136R
PTEN	C211Y
PTEN	C304G
PTEN	C304S
PTEN	C71Y
PTEN	D107A
PTEN	D107Y
PTEN	D153D
PTEN	D153N
PTEN	D153Y
PTEN	D162H
PTEN	D19N
PTEN	D24E

FIGURE 28 (CONTINUED)

PTEN	D24G
PTEN	D24N
PTEN	D24Y
PTEN	D252D
PTEN	D252G
PTEN	D252Y
PTEN	D326G
PTEN	D331G
PTEN	D375N
PTEN	D52G
PTEN	D52N
PTEN	D68Y
PTEN	D92A
PTEN	D92G
PTEN	D92N
PTEN	D92V
PTEN	E150D
PTEN	E150G
PTEN	E150Q
PTEN	E242K
PTEN	E284K
PTEN	E288K
PTEN	E291K
PTEN	E307E
PTEN	E314K
PTEN	E73V
PTEN	E91A
PTEN	E91Q
PTEN	F154F
PTEN	F154L
PTEN	F241S
PTEN	F257L
PTEN	F341V
PTEN	F37S
PTEN	F56V
PTEN	F81C
PTEN	F90L
PTEN	F90S
PTEN	G127E
PTEN	G127V
PTEN	G129R
PTEN	G129V
PTEN	G132D
PTEN	G132S
PTEN	G132V
PTEN	G156R
PTEN	G165E
PTEN	G165R
PTEN	G20E
PTEN	G230R
PTEN	G251C
PTEN	G251V
PTEN	G36E
PTEN	G36R
PTEN	G36V
PTEN	G44D
PTEN	G44G
PTEN	H123Y
PTEN	H141Y
PTEN	H272R
PTEN	H272Y
PTEN	H397Y
PTEN	H61L
PTEN	H61P
PTEN	H61R
PTEN	H93D
PTEN	H93Q
PTEN	H93R
PTEN	H93Y
PTEN	I101I
PTEN	I101M
PTEN	I101N
PTEN	I101T
PTEN	I122S
PTEN	I135K
PTEN	I135M
PTEN	I135V
PTEN	I168F
PTEN	I253N
PTEN	I33S
PTEN	I5L
PTEN	I67K
PTEN	I67R
PTEN	K125E
PTEN	K128N
PTEN	K13E
PTEN	K13Q
PTEN	K144I
PTEN	K254E
PTEN	K342N
PTEN	K344R
PTEN	K62I
PTEN	K62R
PTEN	K66E
PTEN	K66N
PTEN	K66Q
PTEN	K6R
PTEN	K80E
PTEN	K80N
PTEN	L108R
PTEN	L112P
PTEN	L112V
PTEN	L139F
PTEN	L146V
PTEN	L152V
PTEN	L162F
PTEN	L23F
PTEN	L345Q
PTEN	L42P
PTEN	L42R
PTEN	L57S
PTEN	L57W
PTEN	L70P
PTEN	M134L
PTEN	M205I
PTEN	M35R
PTEN	M35V
PTEN	N184I
PTEN	N31N
PTEN	N323K
PTEN	N48D
PTEN	N48I
PTEN	N82T
PTEN	N94I
PTEN	N94Y
PTEN	P169H
PTEN	P204S
PTEN	P204T
PTEN	P244S
PTEN	P246L
PTEN	P339L
PTEN	P38S
PTEN	P95L
PTEN	P95S
PTEN	P96L
PTEN	P96Q
PTEN	P96S
PTEN	Q149R
PTEN	Q171E
PTEN	Q171H
PTEN	Q171P
PTEN	Q17P
PTEN	R130G
PTEN	R130L
PTEN	R130P
PTEN	R130Q
PTEN	R130R
PTEN	R142Q
PTEN	R142W
PTEN	R14G
PTEN	R159S
PTEN	R15I
PTEN	R15K
PTEN	R15S
PTEN	R173C
PTEN	R173H
PTEN	R173R
PTEN	R233R
PTEN	R234W
PTEN	R308C
PTEN	R335R
PTEN	R55G
PTEN	R84G
PTEN	S10N
PTEN	S113R
PTEN	S170G
PTEN	S170I
PTEN	S170N
PTEN	S179I
PTEN	S227F
PTEN	S338I
PTEN	S59P
PTEN	T131I
PTEN	T131N
PTEN	T160I
PTEN	T167A
PTEN	T167P
PTEN	T26P
PTEN	T348I
PTEN	T382S
PTEN	T401I
PTEN	T78A
PTEN	T78I
PTEN	V119D
PTEN	V133I
PTEN	V166L
PTEN	V216M
PTEN	V217A
PTEN	V217I
PTEN	V222A
PTEN	V255A
PTEN	V366I
PTEN	V369G
PTEN	V45V
PTEN	V53G
PTEN	V54L
PTEN	W111R
PTEN	W274G
PTEN	Y138C
PTEN	Y155C
PTEN	Y155H
PTEN	Y155N
PTEN	Y16C
PTEN	Y174N
PTEN	Y188S
PTEN	Y225C
PTEN	Y27C
PTEN	Y27N
PTEN	Y27S
PTEN	Y336F
PTEN	Y65C
PTEN	Y65D
PTEN	Y65N
PTEN	Y68H
PTEN	Y76Y
PTEN	Y88C
PTEN	Y88S
PTPN11	A16A
PTPN11	A72D
PTPN11	A72T
PTPN11	A72V
PTPN11	D61N
PTPN11	D61V
PTPN11	D61Y
PTPN11	E139D
PTPN11	E69K
PTPN11	E69V
PTPN11	E76A
PTPN11	E76G
PTPN11	E76K
PTPN11	E76Q
PTPN11	E76V
PTPN11	F71K
PTPN11	F71L
PTPN11	G503A
PTPN11	G503E
PTPN11	G503V
PTPN11	G60A
PTPN11	G60R
PTPN11	G60V
PTPN11	N58S
PTPN11	N58Y
PTPN11	P491L
PTPN11	P491S
PTPN11	Q510K
PTPN11	R138Q
PTPN11	R289G
PTPN11	S502L
PTPN11	S502P
PTPN11	T507K
PTPN11	I73I
PTPN11	W45L
PTPN11	Y63C
RB1	A14A
RB1	C706F
RB1	D718N
RB1	F839F
RB1	H673P
RB1	I1126S
RB1	I138S
RB1	K122K
RB1	L168S
RB1	L689P
RB1	M208V
RB1	M704V
RB1	N328H
RB1	Q444H
RB1	R621S
RB1	R661W
RB1	R798W
RB1	S187I
RB1	S474N
RB1	S612P
RB1	S794I
RB1	V754G
RET	A664D
RET	A876V
RET	A883F
RET	A883P
RET	A913V
RET	C609Y
RET	C618Y
RET	C630R
RET	C634A
RET	C634R
RET	C634T
RET	C634W
RET	C634Y
RET	D631G
RET	D825H
RET	E884K
RET	E901K
RET	G748C
RET	M918T
RET	P766S
RET	R348R
RET	R908K
RET	V778V
ROS1	Q865H
SMARCB1	A203T
SMARCB1	N152D
SMARCB1	P173S
SMARCB1	P48S
SMARCB1	R127G
SMARCB1	R341L
SMARCB1	R374Q
SMARCB1	R377H
SMARCB1	S284L
SMARCB1	S299S
SMARCB1	S49X
SMARCB1	T372I
SMO	A324T
SMO	A401A
SMO	A652V
SMO	A68V
SMO	L514F
SMO	P755P
SMO	R199W
SMO	R484W
SMO	R562Q
SMO	S533N
SMO	I349I
SMO	T640A
SMO	V404M

FIGURE 28 (CONTINUED)

SMO	W535L
SUFU	F197F
SUFU	P187L
TEC	R563K
IIF1	I403N
TP53	A159V
TP53	A161D
TP53	C124R
TP53	C135F
TP53	C135R
TP53	C176F
TP53	C176Y
TP53	C238Y
TP53	C242F
TP53	C242R
TP53	C242S
TP53	C242W
TP53	C275F
TP53	C275G
TP53	C275Y
TP53	D281G
TP53	D281Y
TP53	E11Q
TP53	E258K
TP53	E285K
TP53	E286K
TP53	F113C
TP53	F113S
TP53	G244A
TP53	G244C
TP53	G244D
TP53	G244S
TP53	G245C
TP53	G245S
TP53	G245V
TP53	G262V
TP53	G266E
TP53	G266V
TP53	G334V
TP53	G389W
TP53	H179Q
TP53	H179R
TP53	H193L
TP53	H193R
TP53	I162N
TP53	I195T
TP53	I232N
TP53	I251N
TP53	I251T
TP53	I254D
TP53	I255S
TP53	I255T
TP53	K132E
TP53	K132Q
TP53	K132R
TP53	K320N
TP53	L130V
TP53	L145R
TP53	L194F
TP53	L264R
TP53	L265P
TP53	M133K
TP53	N237I
TP53	M246I
TP53	N239D
TP53	N268I
TP53	P151S
TP53	P152L
TP53	P177T
TP53	P250L
TP53	P278A
TP53	P278L
TP53	P278R
TP53	P278S
TP53	P309S
TP53	Q16L
TP53	R110L
TP53	R110P
TP53	R156P
TP53	R158H
TP53	R158L
TP53	R175H
TP53	R175L
TP53	R213Q
TP53	R248L
TP53	R248Q
TP53	R248W
TP53	R249G
TP53	R249S
TP53	R267L
TP53	R267P
TP53	R267W
TP53	R273C
TP53	R273H
TP53	R273L
TP53	R280G
TP53	R280K
TP53	R280T
TP53	R282W
TP53	R283C
TP53	R283H
TP53	R283P
TP53	R337C
TP53	R337L
TP53	R379C
TP53	S241C
TP53	S241F
TP53	S241Y
TP53	V143A
TP53	V157F
TP53	V216L
TP53	V216M
TP53	V272M
TP53	V274F
TP53	Y126C
TP53	Y163C
TP53	Y220C
TP53	Y220D
TP53	Y236C
TRIM33	E811K