

DNA-PK INHIBITORS

This application claims the benefit to U.S. Provisional Application No. 61/777,816, filed on March 12, 2013, the entire content of which is incorporated herein by reference.

TECHNICAL FIELD OF THE INVENTION

[0001] The present invention relates to compounds useful as inhibitors of DNA-dependent protein kinase (DNA-PK). The invention also provides pharmaceutically acceptable compositions comprising the compounds of the invention and methods of using the compositions in the treatment of cancer,

BACKGROUND OF THE INVENTION

[0002] Ionizing radiation (IR) induces a variety of DNA damage of which double strand breaks (DSBs) are the most cytotoxic. These DSBs can lead to cell death via apoptosis and/or mitotic catastrophe if not rapidly and completely repaired. In addition to IR, certain chemotherapeutic agents including topoisomerase II inhibitors, bleomycin, and doxorubicin also cause DSBs. These DNA lesions trigger a complex set of signals through the DNA damage response network that function to repair the damaged DNA and maintain cell viability and genomic stability. In mammalian cells, the predominant repair pathway for DSBs is the Non-Homologous End Joining Pathway (NHEJ). This pathway functions regardless of the phase of the cell cycle and does not require a template to re-ligate the broken DNA ends. NHEJ requires coordination of many proteins and signaling pathways. The core NHEJ machinery consists of the Ku70/80 heterodimer and the catalytic subunit of DNA-dependent protein kinase (DNA-PKcs), which together comprise the active DNA-PK enzyme complex. DNA-PKcs is a member of the phosphatidylinositol 3-kinase-related kinase (PIKK) family of serine/threonine protein kinases that also includes ataxia telangiectasia mutated (ATM), ataxia telangiectasia and Rad3-related (ATR), mTOR, and four PI3K isoforms. However, while DNA-PKcs is in the same protein kinase family as ATM and ATR, these latter kinases function to repair DNA damage through the Homologous Recombination (HR) pathway and are restricted to the S

and G₂ phases of the cell cycle. While ATM is also recruited to sites of DSBs, ATR is recruited to sites of single stranded DNA breaks.

[0003] NHEJ is thought to proceed through three key steps: recognition of the DSBs, DNA processing to remove non-ligatable ends or other forms of damage at the termini, and finally ligation of the DNA ends. Recognition of the DSB is carried out by binding of the Ku heterodimer to the ragged DNA ends followed by recruitment of two molecules of DNA-PKcs to adjacent sides of the DSB; this serves to protect the broken termini until additional processing enzymes are recruited. Recent data supports the hypothesis that DNA-PKcs phosphorylates the processing enzyme, Artemis, as well as itself to prepare the DNA ends for additional processing. In some cases DNA polymerase may be required to synthesize new ends prior to the ligation step. The auto-phosphorylation of DNA-PKcs is believed to induce a conformational change that opens the central DNA binding cavity, releases DNA-PKcs from DNA, and facilitates the ultimate religation of the DNA ends.

[0004] It has been known for some time that DNA-PK^{-/-} mice are hypersensitive to the effects of IR and that some non-selective small molecule inhibitors of DNA-PKcs can radiosensitize a variety of tumor cell types across a broad set of genetic backgrounds. While it is expected that inhibition of DNA-PK will radiosensitize normal cells to some extent, this has been observed to a lesser degree than with tumor cells likely due to the fact that tumor cells possess higher basal levels of endogenous replication stress and DNA damage (oncogene-induced replication stress) and DNA repair mechanisms are less efficient in tumor cells. Most importantly, an improved therapeutic window with greater sparing of normal tissue will be imparted from the combination of a DNA-PK inhibitor with recent advances in precision delivery of focused IR, including image-guide RT (IGRT) and intensity-modulated RT (IMRT).


[0005] Inhibition of DNA-PK activity induces effects in both cycling and non-cycling cells. This is highly significant since the majority of cells in a solid tumor are not actively replicating at any given moment, which limits the efficacy of many agents targeting the cell cycle. Equally intriguing are recent reports that suggest a strong connection between inhibition of the NHEJ pathway and the ability to kill traditionally radioresistant cancer stem cells (CSCs). It has been shown in some tumor cells that DSBs in dormant CSCs predominantly activate DNA repair through the NHEJ pathway; it is believed that CSCs are usually in the quiescent phase of the cell cycle. This may explain why half of cancer

patients may experience local or distant tumor relapse despite treatment as current strategies are not able to effectively target CSCs. A DNA-PK inhibitor may have the ability to sensitize these potential metastatic progenitor cells to the effects of IR and select DSB-inducing chemotherapeutic agents.

[0006] Given the involvement of DNA-PK in DNA repair processes, an application of specific DNA-PK inhibitory drugs would be to act as agents that will enhance the efficacy of both cancer chemotherapy and radiotherapy. Accordingly, it would be desirable to develop compounds useful as inhibitors of DNA-PK.

SUMMARY OF THE INVENTION

[0007] It has been found that compounds of this invention, and pharmaceutically acceptable compositions thereof, are effective as inhibitors of DNA-PK. Accordingly, the invention features compounds having the general formula:



or a pharmaceutically acceptable salt thereof, where each of R¹, R², X, Ring A, Ring B and Ring C is as defined elsewhere herein.

[0008] The invention also provides pharmaceutical compositions that include a compound of formula I and a pharmaceutically acceptable carrier, adjuvant, or vehicle. These compounds and pharmaceutical compositions are useful for treating or lessening the severity of cancer.

[0009] The compounds and compositions provided by this invention are also useful for the study of DNA-PK in biological and pathological phenomena; the study of intracellular signal transduction pathways mediated by such kinases; and the comparative evaluation of new kinase inhibitors.

DETAILED DESCRIPTION OF THE INVENTION

Definitions and General Terminology

[0010] As used herein, the following definitions shall apply unless otherwise indicated. For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, and the Handbook of Chemistry and Physics, 75th Ed. 1994. Additionally, general principles of organic chemistry are described in "Organic Chemistry," Thomas Sorrell, University Science Books, Sausalito: 1999, and "March's Advanced Organic Chemistry," 5th Ed., Smith, M.B. and March, J., eds. John Wiley & Sons, New York: 2001, the entire contents of which are hereby incorporated by reference.

[0011] As described herein, compounds of the invention may optionally be substituted with one or more substituents, such as are illustrated generally above, or as exemplified by particular classes, subclasses, and species of the invention. It will be appreciated that the phrase "optionally substituted" is used interchangeably with the phrase "substituted or unsubstituted." In general, the term "substituted," whether preceded by the term "optionally" or not, refers to the replacement of one or more hydrogen radicals in a given structure with the radical of a specified substituent. Unless otherwise indicated, an optionally substituted group may have a substituent at each substitutable position of the group. When more than one position in a given structure can be substituted with more than one substituent selected from a specified group, the substituent may be either the same or different at each position.

[0012] As described herein, when the term "optionally substituted" precedes a list, said term refers to all of the subsequent substitutable groups in that list. For example, if X is halogen; optionally substituted C₁₋₃ alkyl or phenyl; X may be either optionally substituted alkyl or optionally substituted phenyl. Likewise, if the term "optionally substituted" follows a list, said term also refers to all of the substitutable groups in the prior list unless otherwise indicated. For example: if X is halogen, C₁₋₃ alkyl, or phenyl, wherein X is optionally substituted by J^X, then both C₁₋₃ alkyl and phenyl may be optionally substituted by J^X. As is apparent to one having ordinary skill in the art, groups such as H, halogen, NO₂, CN, NH₂, OH, or OCF₃ would not be included because they are not substitutable groups. As is also apparent to a skilled person, a heteroaryl or heterocyclic ring containing an NH group can be optionally substituted by replacing the

hydrogen atom with the substituent. If a substituent radical or structure is not identified or defined as “optionally substituted,” the substituent radical or structure is unsubstituted.

[0013] Combinations of substituents envisioned by this invention are preferably those that result in the formation of stable or chemically feasible compounds. The term “stable,” as used herein, refers to compounds that are not substantially altered when subjected to conditions to allow for their production, detection, and, preferably, their recovery, purification, and use for one or more of the purposes disclosed herein. In some embodiments, a stable compound or chemically feasible compound is one that is not substantially altered when kept at a temperature of 40°C or less, in the absence of moisture or other chemically reactive conditions, for at least a week.

[0014] The term “alkyl” or “alkyl group,” as used herein, means a straight-chain (i.e., unbranched) or branched, substituted or unsubstituted hydrocarbon chain that is completely saturated. Unless otherwise specified, alkyl groups contain 1-8 carbon atoms. In some embodiments, alkyl groups contain 1-6 carbon atoms, and in yet other embodiments, alkyl groups contain 1-4 carbon atoms (represented as “C₁₋₄ alkyl”). In other embodiments, alkyl groups are characterized as “C₀₋₄ alkyl” representing either a covalent bond or a C₁₋₄ alkyl chain. Examples of alkyl groups include methyl, ethyl, propyl, butyl, isopropyl, isobutyl, *sec*-butyl, and *tert*-butyl. The term “alkylene,” as used herein, represents a saturated divalent straight or branched chain hydrocarbon group and is exemplified by methylene, ethylene, isopropylene and the like. The term “alkylidene,” as used herein, represents a divalent straight chain alkyl linking group. The term “alkenyl,” as used herein, represents monovalent straight or branched chain hydrocarbon group containing one or more carbon-carbon double bonds. The term “alkynyl,” as used herein, represents a monovalent straight or branched chain hydrocarbon group containing one or more carbon-carbon triple bonds.

[0015] The term “cycloalkyl” (or “carbocycle”) refers to a monocyclic C₃-C₈ hydrocarbon or bicyclic C₈-C₁₂ hydrocarbon that is completely saturated and has a single point of attachment to the rest of the molecule, and wherein any individual ring in said bicyclic ring system has 3-7 members. Suitable cycloalkyl groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl.

[0016] The term “heterocycle,” “heterocyclyl,” “heterocycloalkyl,” or “heterocyclic” as used herein refers to a monocyclic, bicyclic, or tricyclic ring system in which at least

one ring in the system contains one or more heteroatoms, which is the same or different, and that is completely saturated or that contains one or more units of unsaturation, but which is not aromatic, and that has a single point of attachment to the rest of the molecule. In some embodiments, the “heterocycle,” “heterocyclyl,” “heterocycloalkyl,” or “heterocyclic” group has three to fourteen ring members in which one or more ring members is a heteroatom independently selected from oxygen, sulfur, nitrogen, or phosphorus, and each ring in the system contains 3 to 8 ring members.

[0017] Examples of heterocyclic rings include, but are not limited to, the following monocycles: 2-tetrahydrofuranyl, 3-tetrahydrofuranyl, 2-tetrahydrothiophenyl, 3-tetrahydrothiophenyl, 2-morpholino, 3-morpholino, 4-morpholino, 2-thiomorpholino, 3-thiomorpholino, 4-thiomorpholino, 1-pyrrolidinyl, 2-pyrrolidinyl, 3-pyrrolidinyl, 1-tetrahydropiperazinyl, 2-tetrahydropiperazinyl, 3-tetrahydropiperazinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 1-pyrazolinyl, 3-pyrazolinyl, 4-pyrazolinyl, 5-pyrazolinyl, 1-piperidinyl, 2-piperidinyl, 3-piperidinyl, 4-piperidinyl, 2-thiazolidinyl, 3-thiazolidinyl, 4-thiazolidinyl, 1-imidazolidinyl, 2-imidazolidinyl, 4-imidazolidinyl, 5-imidazolidinyl; and the following bicycles: 3-1H-benzimidazol-2-one, 3-(1-alkyl)-benzimidazol-2-one, indolinyl, tetrahydroquinolinyl, tetrahydroisoquinolinyl, benzothiolane, benzodithiane, and 1,3-dihydro-imidazol-2-one.

[0018] The term “heteroatom” means one or more of oxygen, sulfur, nitrogen, or phosphorus, including any oxidized form of nitrogen, sulfur, or phosphorus; the quaternized form of any basic nitrogen; or a substitutable nitrogen of a heterocyclic ring, for example N (as in 3,4-dihydro-2*H*-pyrrolyl), NH (as in pyrrolidinyl) or NR⁺ (as in N-substituted pyrrolidinyl).

[0019] The term “unsaturated,” as used herein, means that a moiety has one or more units of unsaturation.

[0020] The term “alkoxy,” or “thioalkyl,” as used herein, refers to an alkyl group, as previously defined, attached to the principal carbon chain through an oxygen (“alkoxy”) or sulfur (“thioalkyl”) atom.


[0021] The terms “haloalkyl,” “haloalkenyl,” and “haloalkoxy” mean alkyl, alkenyl, or alkoxy, as the case may be, substituted with one or more halogen atoms. The term “halogen” means F, Cl, Br, or I.

[0022] The term “aryl” used alone or as part of a larger moiety as in “aralkyl,” “aralkoxy,” or “aryloxyalkyl,” refers to a monocyclic, bicyclic, or tricyclic carbocyclic ring system having a total of six to fourteen ring members, wherein said ring system has a single point of attachment to the rest of the molecule, at least one ring in the system is aromatic and wherein each ring in the system contains 4 to 7 ring members. The term “aryl” may be used interchangeably with the term “aryl ring.” Examples of aryl rings include phenyl, naphthyl, and anthracene.


[0023] The term “heteroaryl,” used alone or as part of a larger moiety as in “heteroaralkyl,” or “heteroarylalkoxy,” refers to a monocyclic, bicyclic, and tricyclic ring system having a total of five to fourteen ring members, wherein said ring system has a single point of attachment to the rest of the molecule, at least one ring in the system is aromatic, at least one ring in the system contains one or more heteroatoms independently selected from nitrogen, oxygen, sulfur or phosphorus, and wherein each ring in the system contains 4 to 7 ring members. The term “heteroaryl” may be used interchangeably with the term “heteroaryl ring” or the term “heteroaromatic.”

[0024] Further examples of heteroaryl rings include the following monocycles: 2-furanyl, 3-furanyl, N-imidazolyl, 2-imidazolyl, 4-imidazolyl, 5-imidazolyl, 3-isoxazolyl, 4-isoxazolyl, 5-isoxazolyl, 2-oxazolyl, 4-oxazolyl, 5-oxazolyl, N-pyrrolyl, 2-pyrrolyl, 3-pyrrolyl, 2-pyridyl, 3-pyridyl, 4-pyridyl, 2-pyrimidinyl, 4-pyrimidinyl, 5-pyrimidinyl, pyridazinyl (e.g., 3-pyridazinyl), 2-thiazolyl, 4-thiazolyl, 5-thiazolyl, tetrazolyl (e.g., 5-tetrazolyl), triazolyl (e.g., 2-triazolyl and 5-triazolyl), 2-thienyl, 3-thienyl, pyrazolyl (e.g., 2-pyrazolyl), isothiazolyl, 1,2,3-oxadiazolyl, 1,2,5-oxadiazolyl, 1,2,4-oxadiazolyl, 1,2,3-triazolyl, 1,2,3-thiadiazolyl, 1,3,4-thiadiazolyl, 1,2,5-thiadiazolyl, pyrazinyl, 1,3,5-triazinyl, and the following bicycles: benzimidazolyl, benzofuryl, benzothiophenyl, indolyl (e.g., 2-indolyl), purinyl, quinolinyl (e.g., 2-quinolinyl, 3-quinolinyl, 4-quinolinyl), and isoquinolinyl (e.g., 1-isoquinolinyl, 3-isoquinolinyl, or 4-isoquinolinyl)..

[0025] As described herein, a bond drawn from a substituent to the center of one ring within a multiple-ring system (as shown below) represents substitution of the substituent at any substitutable position in any of the rings within the multiple ring system. For example, Structure a represents possible substitution in any of the positions shown in Structure b.




Structure a




Structure b

[0026] This also applies to multiple ring systems fused to optional ring systems (which would be represented by dotted lines). For example, in Structure c, X is an optional substituent both for ring A and ring B.



Structure c

[0027] If, however, two rings in a multiple ring system each have different substituents drawn from the center of each ring, then, unless otherwise specified, each substituent only represents substitution on the ring to which it is attached. For example, in Structure d, Y is an optional substituent for ring A only, and X is an optional substituent for ring B only.



Structure d

[0028] The term “protecting group,” as used herein, represent those groups intended to protect a functional group, such as, for example, an alcohol, amine, carboxyl, carbonyl, etc., against undesirable reactions during synthetic procedures. Commonly used protecting groups are disclosed in Greene and Wuts, *Protective Groups In Organic Synthesis*, 3rd Edition (John Wiley & Sons, New York, 1999), which is incorporated herein by reference. Examples of nitrogen protecting groups include acyl, aroyl, or carbamyl groups such as formyl, acetyl, propionyl, pivaloyl, t-butylacetyl, 2-chloroacetyl, 2-bromoacetyl, trifluoroacetyl, trichloroacetyl, phthalyl, o-nitrophenoxyacetyl, α-chlorobutyryl, benzoyl, 4-chlorobenzoyl, 4-bromobenzoyl, 4-nitrobenzoyl and chiral auxiliaries such as protected or unprotected D, L or D, L-amino acids such as alanine, leucine, phenylalanine and the like; sulfonyl groups such as benzenesulfonyl, p-toluenesulfonyl and the like; carbamate groups such as benzyloxycarbonyl, p-chlorobenzyloxycarbonyl, p-methoxybenzyloxycarbonyl, p-nitrobenzyloxycarbonyl, 2-

nitrobenzyloxycarbonyl, p-bromobenzyloxycarbonyl, 3,4-dimethoxybenzyloxycarbonyl, 3,5-dimethoxybenzyloxycarbonyl, 2,4-dimethoxybenzyloxycarbonyl, 4-methoxybenzyloxycarbonyl, 2-nitro-4,5-dimethoxybenzyloxycarbonyl, 3,4,5-trimethoxybenzyloxycarbonyl, 1-(p-biphenyl)-1-methylethoxycarbonyl, α,α -dimethyl-3,5-dimethoxybenzyloxycarbonyl, benzhydryloxycarbonyl, t-butyloxycarbonyl, diisopropylmethoxycarbonyl, isopropylloxycarbonyl, ethoxycarbonyl, methoxycarbonyl, allyloxycarbonyl, 2,2,2-trichloroethoxycarbonyl, phenoxycarbonyl, 4-nitrophenoxycarbonyl, fluorenyl-9-methoxycarbonyl, cyclopentylloxycarbonyl, adamantylloxycarbonyl, cyclohexylloxycarbonyl, phenylthiocarbonyl and the like, arylalkyl groups such as benzyl, triphenylmethyl, benzyloxymethyl and the like and silyl groups such as trimethylsilyl and the like. Preferred N-protecting groups are formyl, acetyl, benzoyl, pivaloyl, t-butylacetyl, alanyl, phenylsulfonyl, benzyl, t-butyloxycarbonyl (Boc) and benzyloxycarbonyl (Cbz). Examples of hydroxyl protecting groups include ethers, such as tetrahydropyranyl, tert butyl, benzyl, allyl, and the like; silyl ethers such as trimethyl silyl, triethyl silyl, triisopropylsilyl, tert-butyl diphenyl silyl, and the like; esters such as acetyl, trifluoroacetyl, and the like; and carbonates. Hydroxyl protecting groups also include those appropriate for the protection of phenols.


[0029] Unless otherwise depicted or stated, structures recited herein are meant to include all isomeric (e.g., enantiomeric, diastereomeric, and geometric (or conformational)) forms of the structure; for example, the R and S configurations for each asymmetric center, (Z) and (E) double bond isomers, and (Z) and (E) conformational isomers. Therefore, single stereochemical isomers as well as enantiomeric, diastereomeric, and geometric (or conformational) mixtures of the present compounds are within the scope of the invention. Compounds that have been drawn with stereochemical centers defined, usually through the use of a hatched (.....) or bolded (—) bond, are stereochemically pure, but with the absolute stereochemistry still undefined. Such compounds can have either the *R* or *S* configuration. In those cases where the absolute configuration has been determined, the chiral center(s) are labeled (*R*) or (*S*) in the drawing.

[0030] Unless otherwise stated, all tautomeric forms of the compounds of the invention are within the scope of the invention. Additionally, unless otherwise stated, structures depicted herein are also meant to include compounds that differ only in the


presence of one or more isotopically enriched atoms. For example, compounds having the present structures except for the replacement of hydrogen by deuterium or tritium, or the replacement of a carbon by a ^{13}C - or ^{14}C -enriched carbon are within the scope of this invention. Such compounds are useful, for example, as analytical tools, probes in biological assays, or as DNA-PK inhibitors with an improved therapeutic profile.

Description of Compounds of the Invention


[0031] In one aspect, the invention features compounds having the formula:



Ring A is a ring system selected from



Ring B is a ring system selected from



wherein Ring B is optionally substituted with up to 4 fluorine atoms, up to two OH or up to two C_{1-4} alkyl which is optionally substituted with up to 3 fluorine atoms, up to two OH, or up to two OC_{1-2} alkyl groups;

Ring C is a cyclohexane or a cyclobutane ring;

X is $-\text{NH}-$, $-\text{O}-$, or $-\text{OC}_{1-4}$ alkyl-;

each of R^1 and R^2 is, independently, hydrogen, $-\text{C}(\text{O})\text{NHR}^4$, $-\text{C}(\text{O})\text{OR}^4$, $-\text{NHC}(\text{O})\text{R}^4$, $-\text{NHC}(\text{O})\text{OR}^4$, $-\text{NHC}(\text{O})\text{NHR}^4$, $-\text{NHS}(\text{O})_2\text{R}^4$, $-\text{C}_{0-4}$ alkyl- NHR^4 , or $-\text{OR}^4$, wherein R^1 and R^2 cannot simultaneously be hydrogen, and wherein R^1 and R^2 and the intervening carbon atom can form a dioxane or dioxolane ring;


R^3 is hydrogen, $-C_{1-4}$ alkyl, fluoro, chloro, $-OC_{1-2}$ alkyl, $-C(O)H$, $-C(O)OH$, $-C(O)OC_{1-2}$ alkyl, $-CN$, $-C(O)NHC_{1-2}$ alkyl, or $-C(O)NH_2$, wherein each of said R^3 alkyl is optionally substituted with up to 3 fluorine atoms, up to two OH, or up to two OC_{1-2} alkyl groups;

R^4 is hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-5} cycloalkyl, phenyl, a 5-10-membered monocyclic or bicyclic heteroaryl ring selected from pyrrole, imidazole, pyrazole, triazole, thiazole, isothiazole, oxazole, pyridine, pyrimidine, pyrimidinone, pyrazine, pyridazine, or quinoline, or a 4-10-membered monocyclic or bicyclic heterocyclyl ring selected from oxetane, tetrahydrofuran, tetrahydropyran, dihydroisoxazole, pyrimidine-2,4(1H,3H)-dione, dihydrofuropyrimidine, dihydropyranopyrimidine, dihydropyrrolopyrimidine, tetrahydropteridine, or tetrahydropyridopyrimidine, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl- $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}alkyl)_2$, $C(O)NH(C_{0-4}alkyl-C_{3-5}cycloalkyl)$, CH_2OR^5 , $C_{0-4}alkyl-C(O)R^5$, $C_{0-4}alkyl-C(O)N(R^5)_2$, $C_{0-4}alkyl-C(O)OR^5$, $C_{0-4}alkyl-NHC(O)R^5$, $C_{0-4}alkyl-N(R^5)_2$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR^5 , wherein each of said optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}alkyl-C_{3-5}cycloalkyl$; and

each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, a spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.


[0032] In one embodiment, Ring C is cyclobutane.

[0033] In another aspect, the invention features compounds having the formula:




(II), wherein R^1 and R^2 are as defined for compounds of formula I.

[0034] In another aspect, the invention features compounds having the formula:




(II-A), wherein R^2 is as defined for compounds of formula I.

[0035] In another aspect, the invention features compounds having the formula:



(II-A-1), wherein R^2 is as defined for compounds of formula I.


[0036] In another aspect, the invention features compounds having the formula:



(II-B), wherein R^1 is as defined for compounds

of formula I.

[0037] In another aspect, the invention features compounds having the formula:



(II-B-1), wherein R^1 is as defined for


compounds of formula I.

[0038] In one embodiment, R^1 is $-C_{0-4}$ alkyl-NHR⁴.

[0039] In another embodiment, X is $-O-$ or $-OC_{1-4}$ alkyl-.

[0040] In one embodiment, Ring C is cyclohexane.

[0041] In another aspect, the invention features compounds having the formula:




(III), wherein R^1 is as defined for compounds of

formula I.

[0042]

[0043] In one embodiment, X is $-NH-$.


[0044] In another aspect, the invention features compounds having the formula:




(III-A), wherein R^2 is as defined for compounds of

formula I.

[0045] In another aspect, the invention features compounds having the formula:



(III-A-1) or




(III-A-2),

wherein R^2 is as defined for compounds of formula I.

[0046] In one embodiment, R^2 is $-C_{0-4}$ alkyl-NHR⁴ or $-OR^4$.

[0047] In another embodiment, R^2 is $-NHR^4$ or $-OR^4$.


[0048] In another aspect, the invention features compounds having the formula:




(III-B), wherein R^1 is as defined for compounds of

formula I.

[0049] In another aspect, the invention features compounds having the formula:



(III-B-1) or



(III-B-


2), wherein R^1 is as defined for compounds of formula I.

[0050] In one embodiment, R^1 is $-C_{0-4}$ alkyl-NHR⁴ or $-OR^4$.

[0051] In another embodiment, R^1 is $-NHR^4$ or $-OR^4$.


[0052] In another embodiment, X is $-O-$.

[0053] In another aspect, the invention features compounds having the formula:




(III-C), wherein R^2 is as defined for compounds of formula I.

[0054] In another aspect, the invention features compounds having the formula:




(III-C-1) or



(III-C-2), wherein


R^2 is as defined for compounds of formula I.

[0055] In another aspect, the invention features compounds having the formula:




(III-D), wherein R^1 is as defined for compounds of formula I.

[0056] In another aspect, the invention features compounds having the formula:



(III-D-1) or




(III-D-2), wherein

R^1 is as defined for compounds of formula I.

[0057] In one embodiment, R^1 is $-C_{0-4}$ alkyl-NHR⁴ or -OR⁴.


[0058] In another embodiment, R^1 is -NHR⁴ or -OR⁴.

[0059] In another aspect, the invention features compounds having the formula:




(III-D-3), wherein Y is -O- or -NH-.

[0060] In one embodiment of compounds having formula I, II, II-A, II-A-1, II-B, II-B-1, III, III-A, III-A-1, III-A-2, III-B, III-B-1, III-B-2, III-C, III-C-1, III-C-2, III-D, III-D-1, III-D-2, or III-D-3,



[0061] In another embodiment,



[0062] In another embodiment, R^3 is hydrogen.

[0063] In another embodiment, R^4 is hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-5} cycloalkyl, phenyl, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO₂, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-C(O)NH₂, C(O)NHC₁₋₄ alkyl, C(O)N(C_{1-4} alkyl)₂, C(O)NH(C_{0-4} alkyl- C_{3-5} cycloalkyl), CH₂OR⁵, C_{0-4} alkyl-C(O)R⁵, C_{0-4} alkyl-C(O)N(R⁵)₂, C_{0-4} alkyl-C(O)OR⁵, C_{0-4} alkyl-NHC(O)R⁵, C_{0-4} alkyl-N(R⁵)₂, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR⁵, wherein each of said optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH

groups, up to two OC₁₋₄alkyl groups, up to two SC₁₋₄alkyl groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, a spirooxetane, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

[0064] In another embodiment, R⁴ is a 5-10-membered monocyclic or bicyclic heteroaryl ring selected from pyrrole, imidazole, pyrazole, triazole, thiazole, isothiazole, oxazole, pyridine, pyrimidine, pyrimidinone, pyrazine, pyridazine, or quinoline, wherein each of said R⁴ groups is optionally substituted with up to four Br, Cl, F, or C₁₋₄alkyl, up to three CN, NO₂, C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₆cycloalkyl, C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-O-C₁₋₄ alkyl, C₀₋₄ alkyl-O-C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-C(O)NH₂, C(O)NHC₁₋₄ alkyl, C(O)N(C₁₋₄ alkyl)₂, C(O)NH(C₀₋₄ alkyl-C₃₋₅ cycloalkyl), CH₂OR⁵, C₀₋₄ alkyl-C(O)R⁵, C₀₋₄ alkyl-C(O)N(R⁵)₂, C₀₋₄ alkyl-C(O)OR⁵, C₀₋₄ alkyl-NHC(O)R⁵, C₀₋₄ alkyl-N(R⁵)₂, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR⁵, wherein each of said optional R⁴ substituents is optionally substituted with up to four fluorine atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, up to two SC₁₋₄alkyl groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and


each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, a spirooxetane, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

[0065] In yet another embodiment, R^4 is pyridine or pyrimidine, which is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl- $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}alkyl)_2$, $C(O)NH(C_{0-4}alkyl-C_{3-5}cycloalkyl)$, CH_2OR^5 , $C_{0-4}alkyl-C(O)R^5$, $C_{0-4}alkyl-C(O)N(R^5)_2$, $C_{0-4}alkyl-C(O)OR^5$, $C_{0-4}alkyl-NHC(O)R^5$, $C_{0-4}alkyl-N(R^5)_2$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR^5 , wherein each of said optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}alkyl-C_{3-5}cycloalkyl$; and each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, a spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

[0066] In another embodiment, R^4 is a 4-10-membered monocyclic or bicyclic heterocyclyl ring selected from oxetane, tetrahydrofuran, tetrahydropyran, dihydroisoxazole, pyrimidine-2,4(1H,3H)-dione, dihydrofuropyrimidine, dihydropyranopyrimidine, dihydropyrrolopyrimidine, tetrahydropteridine, or tetrahydropyridopyrimidine, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl, up to three CN, NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}alkyl-C_{3-5}cycloalkyl$, $C_{0-4}alkyl-C(O)NH_2$, $C(O)NHC_{1-4}alkyl$, $C(O)N(C_{1-4}alkyl)_2$, $C(O)NH(C_{0-4}alkyl-C_{3-5}cycloalkyl)$, CH_2OR^5 , $C_{0-4}alkyl-C(O)R^5$, $C_{0-4}alkyl-C(O)N(R^5)_2$, $C_{0-4}alkyl-C(O)OR^5$, $C_{0-4}alkyl-NHC(O)R^5$, $C_{0-4}alkyl-N(R^5)_2$, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine or piperazine, a


heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole or tetrazole, or up to two OR⁵, wherein each of said optional R⁴ substituents is optionally substituted with up to four fluorine atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, up to two SC₁₋₄alkyl groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, or pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, or tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, a spirooxetane, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

[0067] In another aspect, the invention features compounds having the formula:



R³ is hydrogen, C₁₋₄alkyl, fluoro, chloro, -OC₁₋₂alkyl, or -C(O)NH₂, -C(O)H, or -CN, wherein each of said R³ alkyl is optionally substituted with OH or up to 3 fluorine atoms;

R⁴ is



X¹ is N, CH, CF, CCl, or CC₁₋₂ alkyl optionally substituted with up to 3 fluorine atoms;

X² is N or CR^{4c}; wherein X¹ and X² cannot simultaneously be N,

each of R^{4a}, R^{4b}, and R^{4c} is, independently, hydrogen, F, Cl, Br, CN, NO₂, C₁₋₄ alkyl, C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-O-C₁₋₄ alkyl, C₀₋₄ alkyl-O-C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl,

C(O)NH₂, C(O)NHC₁₋₄ alkyl, C(O)N(C₁₋₄ alkyl)₂, C(O)NH(C₀₋₄ alkyl-C₃₋₅ cycloalkyl), a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, or piperazine, or a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, or tetrazole, or R^{4c}, R^{4a}, and the intervening atoms form a dihydrofuran, a dihydropyran, or a tetrahydropiperidine heterocyclic ring system;

wherein each of said R^{4a}, R^{4b}, or R^{4c} heterocyclic or heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C₁₋₄ alkyl, up to two OH groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and

wherein each of said R^{4a}, R^{4b}, or R^{4c} alkyl or cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.

[0068] In one embodiment, R³ is hydrogen.

[0069] In another embodiment, each of X¹ and X² is, independently, CH or N.

[0070] In another embodiment, each of R^{4a} and R^{4b} is, independently, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, or piperazine, wherein each of said R^{4a} or R^{4b} heterocyclic or heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C₁₋₄ alkyl, up to two OH groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and wherein each of said R^{4a} or R^{4b} alkyl or cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.

[0071] In another embodiment, each of R^{4a} and R^{4b} is, independently, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, or tetrazole, wherein each of said R^{4a} or R^{4b} heterocyclic or heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C₁₋₄ alkyl, up to two OH groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and wherein each of said R^{4a} or R^{4b} alkyl or cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.

[0072] In another embodiment, each of R^{4a} and R^{4b} is, independently, hydrogen, F, Cl, Br, CN, NO₂, C₁₋₄ alkyl, C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-O-C₁₋₄ alkyl, C₀₋₄ alkyl-O-C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl, C(O)NH₂, C(O)NHC₁₋₄ alkyl, C(O)N(C₁₋₄ alkyl)₂, or C(O)NH(C₀₋₄ alkyl-C₃₋₅ cycloalkyl), wherein each of said R^{4a} or R^{4b} heterocyclic or heteroaryl ring systems is

optionally substituted with up to four fluorine atoms, up to two C₁₋₄ alkyl, up to two OH groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and wherein each of said R^{4a} or R^{4b} alkyl or cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.

[0073] In another embodiment, the invention features a compound selected from the group of compounds listed in Table 1.

[0074] In another embodiment, the invention features a compound selected from the group of compounds listed in Table 2.

Compositions, Formulations, and Administration of Compounds of the Invention

[0075] In another embodiment, the invention provides a pharmaceutical composition comprising a compound of any of the formulae described herein and a pharmaceutically acceptable excipient. In a further embodiment, the invention provides a pharmaceutical composition comprising a compound of Table 1. In a further embodiment, the composition additionally comprises an additional therapeutic agent.

[0076] According to another embodiment, the invention provides a composition comprising a compound of this invention or a pharmaceutically acceptable derivative thereof and a pharmaceutically acceptable carrier, adjuvant, or vehicle. In one embodiment, the amount of compound in a composition of this invention is such that is effective to measurably inhibit a DNA-PK in a biological sample or in a patient. In another embodiment, the amount of compound in the compositions of this invention is such that is effective to measurably inhibit DNA-PK. In one embodiment, the composition of this invention is formulated for administration to a patient in need of such composition. In a further embodiment, the composition of this invention is formulated for oral administration to a patient.

[0077] The term “patient,” as used herein, means an animal, preferably a mammal, and most preferably a human.

[0078] It will also be appreciated that certain of the compounds of present invention can exist in free form for treatment, or where appropriate, as a pharmaceutically acceptable derivative thereof. According to the present invention, a pharmaceutically acceptable derivative includes, but is not limited to, pharmaceutically acceptable prodrugs, salts, esters, salts of such esters, or any other adduct or derivative which upon

administration to a patient in need is capable of providing, directly or indirectly, a compound as otherwise described herein, or a metabolite or residue thereof. As used herein, the term “inhibitory active metabolite or residue thereof” means that a metabolite or residue thereof is also an inhibitor of DNA-PK.

[0079] As used herein, the term “pharmaceutically acceptable salt” refers to those salts which are, within the scope of sound medical judgment, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response and the like.

[0080] Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge et al., describe pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences*, 66:1-19, 1977, which is incorporated herein by reference. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate, palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_{1-4} \text{ alkyl})_4$ salts. This invention also envisions the quaternization of any basic nitrogen-containing groups of the compounds disclosed herein. Water or oil-soluble or dispersable products may be obtained by such quaternization. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when

appropriate, nontoxic ammonium, quaternary ammonium, and amine cations formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, C₁₋₈ sulfonate and aryl sulfonate.

[0081] As described above, the pharmaceutically acceptable compositions of the present invention additionally comprise a pharmaceutically acceptable carrier, adjuvant, or vehicle, which, as used herein, includes any and all solvents, diluents, or other liquid vehicle, dispersion or suspension aids, surface active agents, isotonic agents, thickening or emulsifying agents, preservatives, solid binders, lubricants and the like, as suited to the particular dosage form desired. In Remington: *The Science and Practice of Pharmacy*, 21st edition, 2005, ed. D.B. Troy, Lippincott Williams & Wilkins, Philadelphia, and *Encyclopedia of Pharmaceutical Technology*, eds. J. Swarbrick and J. C. Boylan, 1988-1999, Marcel Dekker, New York, the contents of each of which is incorporated by reference herein, are disclosed various carriers used in formulating pharmaceutically acceptable compositions and known techniques for the preparation thereof. Except insofar as any conventional carrier medium is incompatible with the compounds of the invention, such as by producing any undesirable biological effect or otherwise interacting in a deleterious manner with any other component(s) of the pharmaceutically acceptable composition, its use is contemplated to be within the scope of this invention.

[0082] Some examples of materials which can serve as pharmaceutically acceptable carriers include, but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, or potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, wool fat, sugars such as lactose, glucose and sucrose; starches such as corn starch and potato starch; cellulose and its derivatives such as sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as cocoa butter and suppository waxes; oils such as peanut oil, cottonseed oil; safflower oil; sesame oil; olive oil; corn oil and soybean oil; glycols; such as propylene glycol or polyethylene glycol; esters such as ethyl oleate and ethyl laurate; agar; buffering agents such as magnesium hydroxide and aluminum

hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

[0083] The compositions of the present invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. The term "parenteral" as used herein includes subcutaneous, intravenous, intramuscular, intra-articular, intra-synovial, intrasternal, intrathecal, intraocular, intrahepatic, intralesional, epidural, intraspinal, and intracranial injection or infusion techniques. Preferably, the compositions are administered orally, intraperitoneally or intravenously. Sterile injectable forms of the compositions of this invention may be aqueous or oleaginous suspension. These suspensions may be formulated according to techniques known in the art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally acceptable diluent or solvent, for example as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium.

[0084] For this purpose, any bland fixed oil may be employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant, such as carboxymethyl cellulose or similar dispersing agents that are commonly used in the formulation of pharmaceutically acceptable dosage forms including emulsions and suspensions. Other commonly used surfactants, such as Tweens, Spans and other emulsifying agents or bioavailability enhancers which are commonly used in the manufacture of pharmaceutically acceptable solid, liquid, or other dosage forms may also be used for the purposes of formulation.

[0085] The pharmaceutically acceptable compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, aqueous suspensions or solutions. In the case of tablets for oral use, carriers commonly used include lactose and corn starch. Lubricating agents, such as magnesium stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried cornstarch. When aqueous suspensions are required for oral use, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening, flavoring or coloring agents may also be added.

[0086] Alternatively, the pharmaceutically acceptable compositions of this invention may be administered in the form of suppositories for rectal administration. These can be prepared by mixing the agent with a suitable non-irritating excipient that is solid at room temperature but liquid at rectal temperature and therefore will melt in the rectum to release the drug. Such materials include cocoa butter, beeswax and polyethylene glycols.

[0087] The pharmaceutically acceptable compositions of this invention may also be administered topically, especially when the target of treatment includes areas or organs readily accessible by topical application, including diseases of the eye, the skin, or the lower intestinal tract. Suitable topical formulations are readily prepared for each of these areas or organs.

[0088] Topical application for the lower intestinal tract can be effected in a rectal suppository formulation (see above) or in a suitable enema formulation. Topically-transdermal patches may also be used.

[0089] For topical applications, the pharmaceutically acceptable compositions may be formulated in a suitable ointment containing the active component suspended or dissolved in one or more carriers. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petrolatum, white petrolatum, propylene glycol, polyoxyethylene, polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutically acceptable compositions can be formulated in a suitable lotion or cream containing the active components suspended or dissolved in one or more pharmaceutically acceptable carriers. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol, 2-octyldodecanol, benzyl alcohol and water.

[0090] For ophthalmic use, the pharmaceutically acceptable compositions may be formulated, e.g., as micronized suspensions in isotonic, pH adjusted sterile saline or other aqueous solution, or, preferably, as solutions in isotonic, pH adjusted sterile saline or other aqueous solution, either with or without a preservative such as benzylalkonium chloride. Alternatively, for ophthalmic uses, the pharmaceutically acceptable compositions may be formulated in an ointment such as petrolatum. The pharmaceutically acceptable compositions of this invention may also be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other conventional solubilizing or dispersing agents.

[0091] Most preferably, the pharmaceutically acceptable compositions of this invention are formulated for oral administration.

[0092] Liquid dosage forms for oral administration include, but are not limited to, pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethylformamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor, and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, and mixtures thereof. Besides inert diluents, the oral compositions can also include adjuvants such as wetting agents, emulsifying and suspending agents, sweetening, flavoring, and perfuming agents.

[0093] Injectable preparations, for example, sterile injectable aqueous or oleaginous suspensions may be formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation may also be a sterile injectable solution, suspension or emulsion in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, U.S.P. and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be

employed including synthetic mono- or diglycerides. In addition, fatty acids such as oleic acid are used in the preparation of injectables.

[0094] The injectable formulations can be sterilized, for example, by filtration through a bacterial-retaining filter, or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium prior to use.

[0095] In order to prolong the effect of a compound of the present invention, it is often desirable to slow the absorption of the compound from subcutaneous or intramuscular injection. This may be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the compound then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form. Alternatively, dissolving or suspending the compound in an oil vehicle accomplishes delayed absorption of a parenterally administered compound form.

Injectable depot forms are made by forming microencapsule matrices of the compound in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of compound to polymer and the nature of the particular polymer employed, the rate of compound release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the compound in liposomes or microemulsions that are compatible with body tissues.

[0096] Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating excipients or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at ambient temperature but liquid at body temperature and therefore melt in the rectum or vaginal cavity and release the active compound.

[0097] Solid dosage forms for oral administration include capsules, tablets, pills, powders, and granules. In such solid dosage forms, the active compound is mixed with at least one inert, pharmaceutically acceptable excipient or carrier such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol, and silicic acid, b) binders such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose, and acacia, c) humectants such as glycerol, d) disintegrating agents such as agar-agar, calcium carbonate,

potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate, e) solution retarding agents such as paraffin, f) absorption accelerators such as quaternary ammonium compounds, g) wetting agents such as, for example, cetyl alcohol and glycerol monostearate, h) absorbents such as kaolin and bentonite clay, and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

[0098] Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings and other coatings well known in the pharmaceutical formulating art. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugar as well as high molecular weight polyethylene glycols and the like.

[0099] The active compounds can also be in micro-encapsulated form with one or more excipients as noted above. The solid dosage forms of tablets, dragees, capsules, pills, and granules can be prepared with coatings and shells such as enteric coatings, release controlling coatings and other coatings well known in the pharmaceutical formulating art. In such solid dosage forms the active compound may be admixed with at least one inert diluent such as sucrose, lactose or starch. Such dosage forms may also comprise, as is normal practice, additional substances other than inert diluents, e.g., tableting lubricants and other tableting aids such as magnesium stearate and microcrystalline cellulose. In the case of capsules, tablets and pills, the dosage forms may also comprise buffering agents. They may optionally contain opacifying agents and can also be of a composition that they release the active ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

[00100] Dosage forms for topical or transdermal administration of a compound of this invention include ointments, pastes, creams, lotions, gels, powders, solutions, sprays, inhalants or patches. The active component is admixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives or buffers as may be required. Ophthalmic formulation, eardrops, and eye drops are also contemplated as being within the scope of this invention. Additionally, the present invention contemplates the use of transdermal patches, which have the added advantage of providing controlled delivery of a compound to the body. Such dosage forms can be made by dissolving or dispensing the compound in the proper medium. Absorption enhancers can also be used to increase the flux of the compound across the skin. The rate can be controlled by either providing a rate controlling membrane or by dispersing the compound in a polymer matrix or gel.

[00101] The compounds of the invention are preferably formulated in dosage unit form for ease of administration and uniformity of dosage. The expression "dosage unit form" as used herein refers to a physically discrete unit of agent appropriate for the patient to be treated. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical judgment. The specific effective dose level for any particular patient or organism will depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed, and like factors well known in the medical arts.

[00102] The amount of the compounds of the present invention that may be combined with the carrier materials to produce a composition in a single dosage form will vary depending upon the host treated, the particular mode of administration. Preferably, the compositions should be formulated so that a dosage of between 0.01 - 100 mg/kg body weight/day of the inhibitor can be administered to a patient receiving these compositions.

[00103] Depending upon the particular proliferative condition or cancer to be treated, additional therapeutic agents, which are normally administered to treat or prevent that

condition, may also be present in the compositions of this invention. As used herein, additional therapeutic agents which are normally administered to treat or prevent a particular proliferative condition or cancer are known as “appropriate for the disease, or condition, being treated.” Examples of additional therapeutic agents are provided *infra*.

[00104] The amount of additional therapeutic agent present in the compositions of this invention will be no more than the amount that would normally be administered in a composition comprising that therapeutic agent as the only active agent. Preferably the amount of additional therapeutic agent in the presently disclosed compositions will range from about 50% to 100% of the amount normally present in a composition comprising that agent as the only therapeutically active agent.

Uses of the Compounds and Compositions of the Invention

[00105] In one embodiment, the invention provides a method of sensitizing a cell to a therapeutic agent or a disease state that induces a DNA lesion comprising the step of contacting the cell with one or more DNA-PK inhibitors of formulae **I**, **II**, or **III**, or subformula thereof (e.g., formulae **I-A**, **I-A-1**, **I-A-2**, **I-B**, **I-B-1**, **I-B-2**, **I-C**, **I-C-1**, **I-C-2**, **I-C-3**, **I-C-4**, **I-D**, **I-D-1**, **I-D-2**, **I-D-3**, **I-D-4**, or **I-D-5**).

[00106] The invention further provides methods of potentiating a therapeutic regimen for treatment of cancer comprising the step of administering to an individual in need thereof an effective amount of a DNA-PK inhibitor of formula **I**, **II**, or **III**, or a subformula thereof. In one aspect, the therapeutic regimen for treatment of cancer includes radiation therapy.

[00107] Compounds of the invention are useful in instances where radiation therapy is indicated to enhance the therapeutic benefit of such treatment. In addition, radiation therapy frequently is indicated as an adjuvant to surgery in the treatment of cancer. The goal of radiation therapy in the adjuvant setting is to reduce the risk of recurrence and enhance disease-free survival when the primary tumor has been controlled. Adjuvant radiation therapy is indicated in several diseases including colon, rectal, lung, gastroesophageal, and breast cancers as described below.

[00108] The invention also can be practiced by including another anti-cancer chemotherapeutic agent with a compound of the invention in a therapeutic regimen for the treatment of cancer, with or without radiation therapy. The combination of a DNA-PK

inhibitor compound of the invention with such other agents can potentiate the chemotherapeutic protocol. For example, the inhibitor compound of the invention can be administered with etoposide or bleomycin, agents known to cause DNA strand breakage.

[00109] The invention further relates to radiosensitizing tumor cells utilizing a compound of formula **I**, **II**, or **III**, or a subformula thereof. The preferred compounds are those as described for the pharmaceutical compositions of the invention. A compound that can “radiosensitize” a cell, as used herein, is defined as a molecule, preferably a low molecular weight molecule, administered to animals in therapeutically effective amount to increase the sensitivity of cells to electromagnetic radiation and/or to promote the treatment of diseases that are treatable with electromagnetic radiation (e.g., X-rays). Diseases that are treatable with electromagnetic radiation include neoplastic diseases, benign and malignant tumors, and cancerous cells.

[00110] The present invention also provides methods of treating cancer in an animal that includes administering to the animal an effective amount of a DNA-PK inhibitor such as, for example, a compound of the invention. The invention further is directed to methods of inhibiting cancer cell growth, including processes of cellular proliferation, invasiveness, and metastasis in biological systems. Methods include use of a compound of the invention as an inhibitor of cancer cell growth. Preferably, the methods are employed to inhibit or reduce cancer cell growth, invasiveness, metastasis, or tumor incidence in living animals, such as mammals. The compounds of the invention can be used, either alone or in combination with the use of IR or one or more chemotherapeutic agents, in treating cancer or inhibiting cancer cell growth. Methods of the invention also are readily adaptable for use in assay systems, e.g., assaying cancer cell growth and properties thereof, as well as identifying compounds that affect cancer cell growth.

[00111] Tumors or neoplasms include growths of tissue cells in which the multiplication of the cells is uncontrolled and progressive. Some such growths are benign, but others are termed “malignant” and can lead to death of the organism. Malignant neoplasms or “cancers” are distinguished from benign growths in that, in addition to exhibiting aggressive cellular proliferation, they can invade surrounding tissues and metastasize. Moreover, malignant neoplasms are characterized in that they show a greater loss of differentiation (greater “dedifferentiation”) and their organization relative to one another and their surrounding tissues. This property is also called “anaplasia.”

[00112] Neoplasms treatable by the present invention also include solid tumors, i.e., carcinomas and sarcomas. Carcinomas include those malignant neoplasms derived from epithelial cells which infiltrate (invade) the surrounding tissues and give rise to metastases. Adenocarcinomas are carcinomas derived from glandular tissue, or from tissues which form recognizable glandular structures. Another broad category of cancers includes sarcomas, which are tumors whose cells are embedded in a fibrillar or homogeneous substance like embryonic connective tissue. The invention also enables treatment of cancers of the myeloid or lymphoid systems, including leukemias, lymphomas, and other cancers that typically do not present as a tumor mass, but are distributed in the vascular or lymphoreticular systems.

[00113] DNA-PK activity can be associated with various forms of cancer in, for example, adult and pediatric oncology, growth of solid tumors/malignancies, myxoid and round cell carcinoma, locally advanced tumors, metastatic cancer, human soft tissue sarcomas, including Ewing's sarcoma, cancer metastases, including lymphatic metastases, squamous cell carcinoma, particularly of the head and neck, esophageal squamous cell carcinoma, oral carcinoma, blood cell malignancies, including multiple myeloma, leukemias, including acute lymphocytic leukemia, acute nonlymphocytic leukemia, chronic lymphocytic leukemia, chronic myelocytic leukemia, and hairy cell leukemia, effusion lymphomas (body cavity based lymphomas), thymic lymphoma lung cancer, including small cell lung carcinoma, cutaneous T cell lymphoma, Hodgkin's lymphoma, non-Hodgkin's lymphoma, cancer of the adrenal cortex, ACTH-producing tumors, nonsmall cell cancers, breast cancer, including small cell carcinoma and ductal carcinoma, gastrointestinal cancers, including stomach cancer, colon cancer, colorectal cancer, polyps associated with colorectal neoplasia, pancreatic cancer, liver cancer, urological cancers, including bladder cancer, including primary superficial bladder tumors, invasive transitional cell carcinoma of the bladder, and muscle-invasive bladder cancer, prostate cancer, malignancies of the female genital tract, including ovarian carcinoma, primary peritoneal epithelial neoplasms, cervical carcinoma, uterine endometrial cancers, vaginal cancer, cancer of the vulva, uterine cancer and solid tumors in the ovarian follicle, malignancies of the male genital tract, including testicular cancer and penile cancer, kidney cancer, including renal cell carcinoma, brain cancer, including intrinsic brain tumors, neuroblastoma, astrocytic brain tumors, gliomas, metastatic tumor cell invasion in

the central nervous system, bone cancers, including osteomas and osteosarcomas, skin cancers, including malignant melanoma, tumor progression of human skin keratinocytes, squamous cell cancer, thyroid cancer, retinoblastoma, neuroblastoma, peritoneal effusion, malignant pleural effusion, mesothelioma, Wilms's tumors, gall bladder cancer, trophoblastic neoplasms, hemangiopericytoma, and Kaposi's sarcoma. Methods to potentiate treatment of these and other forms of cancer are embraced by the invention.

[00114] The invention provides a method of inhibiting DNA-PK activity in a biological sample that includes contacting the biological sample with a compound or composition of the invention. The term "biological sample," as used herein, means a sample outside a living organism and includes, without limitation, cell cultures or extracts thereof; biopsied material obtained from a mammal or extracts thereof; and blood, saliva, urine, feces, semen, tears, or other body fluids or extracts thereof. Inhibition of kinase activity, particularly DNA-PK activity, in a biological sample is useful for a variety of purposes known to one of skill in the art. Examples of such purposes include, but are not limited to, biological specimen storage and biological assays. In one embodiment, the method of inhibiting DNA-PK activity in a biological sample is limited to non-therapeutic methods.

Preparation of Compounds of the Invention


[00115] As used herein, all abbreviations, symbols and conventions are consistent with those used in the contemporary scientific literature. See, e.g., Janet S. Dodd, ed., *The ACS Style Guide: A Manual for Authors and Editors*, 2nd Ed., Washington, D.C.: American Chemical Society, 1997. The following definitions describe terms and abbreviations used herein:

BPIn	pinacol boronate ester
Brine	a saturated NaCl solution in water
DCM	dichloromethane
DIAD	diisopropylazodicarboxylate
DIEA	diisopropylethylamine
DMA	dimethylacetamide
DMF	dimethylformamide
DMSO	dimethylsulfoxide
DTT	dithiothreitol

ESMS	electrospray mass spectrometry
Et ₂ O	ethyl ether
EtOAc	ethyl acetate
EtOH	ethyl alcohol
HEPES	4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid
HPLC	high performance liquid chromatography
IPA	isopropanol
LAH	lithium aluminum hydride
LC-MS	liquid chromatography-mass spectrometry
LDA	lithium diisopropylethylamide
Me	methyl
MeOH	methanol
MsCl	methanesulfonyl chloride
MTBE	methyl <i>t</i> -butyl ether
NMP	N-methylpyrrolidine
Pd ₂ (dba) ₃	tris(dibenzylideneacetone)dipalladium(0)
Pd(dppf)Cl ₂	1,1' bis(diphenylphosphino)-ferrocene dichloro-palladium
PG	protecting group
Ph	phenyl
(<i>rac</i>)-BINAP	racemic 2,2'-bis(diphenylphosphino)-1,1'-binaphthyl
RockPhos	di- <i>tert</i> -butyl(2',4',6'-triisopropyl-3,6-dimethoxy-[1,1'-biphenyl]-2-yl)phosphine
RT or rt	room temperature
SFC	supercritical fluid chromatography
SPhos	2-dicyclohexylphosphino-2',6'-dimethoxybiphenyl
TBAI	tetrabutylammonium iodide
<i>t</i> Bu	tertiary butyl
THF	tetrahydrofuran
TEA	triethylamine
TMEDA	tetramethylethylenediamine
VPhos	[3-(2-dicyclohexylphosphanylphenyl)-2,4-dimethoxy-phenyl]sulfonyloxysodium

General Synthetic Procedures

[00116] In general, the compounds of this invention may be prepared by methods described herein or by other methods known to those skilled in the art.


Example 1. General preparation of the compounds of formula **G**

Scheme 1

[00117] Compounds of formula **I**, wherein X^1 is NH (i.e., compounds of formula **I-A**), can be prepared as outlined below in Scheme 1. Accordingly, as shown in step 1-i of Scheme 1, heteroaryl compounds of formula **A** can be reacted with morpholine or a morpholine analog by heating the mixture in a polar, non-protic solvent to produce compounds of formula **B**. Utilizing a palladium-catalyzed, phosphine ligand-assisted Buchwald/Hartwig-type coupling, as shown in step 1-ii of Scheme 1, a compound of formula **B** can be reacted with aminocyclohexanes of formula **C** to produce compounds of formula **D**, wherein R^1 and R^2 are as described elsewhere herein. In one example, when monoprotected meso cyclohexane-1,4-diamines of formula **E** are prepared, removal of the protecting group forms compounds of formula **F**, as shown in step 1-iii of Scheme 1. The resulting free amine can then be reacted with various moieties having groups reactive towards amines (e.g., $R^{1a}\text{-L}$, where L is a leaving group such as chloro, bromo, iodo, toluenesulfonate, methanesulfonate, or trifluoromethanesulfonate; or where L is a reactive

carbonyl-containing moiety such as an active ester or an isocyanato group) to produce a compound of formula **G**, as shown in step 1-iv of Scheme 1.


Example 2. General preparation of the compounds of formula **M**, **N**, **R**, and **S**



Scheme 2a


[00118] Compounds of formula **I**, wherein X^1 is O can be prepared as outlined below in Schemes 2a and 2b. Accordingly, as shown in step 2-i of Scheme 2a, the hydroxyl group of heteroaryl compounds of formula **H** can be protected to produce compounds of formula **J**, which can then be reacted with morpholine or a morpholine analog by heating the mixture in a polar, non-protic solvent to produce compounds of formula **K** after removal of the protecting group, as shown in steps 2-ii and 2-iii of Scheme 2a. Subsequently, as shown in step 2-iv of Scheme 2a, a compound of formula **K** can be reacted with a compound of formula **L** under conditions sufficient to affect the S_N2 displacement of its leaving group (e.g., where L is a leaving group such as chloro, bromo, iodo, toluenesulfonate, methanesulfonate, or trifluoromethanesulfonate) to produce a compound of formula **M** or formula **M**, depending on whether R^1 or R^2 is hydrogen. In those instances when R^1 or R^2 are protected nitrogen or oxygen moieties, compounds of the invention can be produced by removal of the protecting group and subsequent synthetic manipulation of the resulting free amine/alcohol.

[00119] Alternatively, as shown in Scheme 2b, the hydroxyl group of a compound of formula **O** can be reacted with a compound of formula **L** to produce a fused bicycloheteroaryl bromide of formula **P**, which can subsequently be reacted with morpholine or a morpholine analog to produce a compound of formula **M** or formula **N**.




Scheme 2b

[00120] Alternatively, as shown in Scheme 2c, compounds of the invention in which Ring B is a dihydropyran ring can be prepared by reacting compounds of formula **Q** with dialkyl (3,6-dihydro-2*H*-pyran-4-yl)boronates to produce compounds of formula **R**. Compounds of formula **R** can then be subsequently reduced to form compounds of formula **S**.



Scheme 2c

Example 3. Preparation of ethyl (4-((7-morpholinoquinoxalin-5-yl)amino)cyclohexyl)carbamate (Compound **6**) and *N*¹-(7-morpholinoquinoxalin-5-yl)-*N*⁴-(pyrimidin-2-yl)cyclohexane-1,4-diamine (Compound **18**)



Scheme 3

[00121] As shown in step 3-i of Scheme 3, to a solution of 3-bromo-5-fluoro-benzene-1,2-diamine (compound **1001**, 1.11 g, 5.41 mmol) in methanol (11 mL) was added oxaldehyde (1.57 mL of 40 % w/v, 10.8 mmol). The reaction mixture was stirred at room temperature under nitrogen. After 2 hours a yellow solid precipitated. The reaction mixture was diluted with water (20 mL), stirred an additional 5 minutes, filtered, and the collected solid dried under high vacuum to produce 5-bromo-7-fluoroquinoxaline (compound **1002**, 868 mg, 70.6% yield): ¹H-NMR (300 MHz, DMSO-*d*₆) δ 9.06 (s, 2H), 8.36 (dd, *J* = 8.5, 2.7 Hz, 1H), 8.00 (dd, *J* = 9.2, 2.7 Hz, 1H); ESMS (*M*+*H*⁺) = 227.14.

[00122] As shown in step 3-ii of Scheme 3, to a solution of 5-bromo-7-fluoroquinoxaline (4.5 g, 19.8 mmol) in NMP (67.5 mL) was added morpholine (3.1 mL, 35.6 mmol). The reaction mixture was heated to 140°C and stirred for 15 hours. After cooling, the mixture was poured into water (200 mL), extracted with ethyl acetate (2 x

100mL), dried over magnesium sulfate, filtered, evaporated under reduced pressure, and purified by medium pressure silica gel chromatography (10 to 80% EtOAc/hexanes gradient) to provide 4-(8-bromoquinoxalin-6-yl)morpholine (compound **1003**, 3.86g, 66% yield) as a yellow solid: ^1H -NMR (400 MHz, DMSO- d_6) δ 8.82 (d, J = 1.6 Hz, 1H), 8.73 (d, J = 1.6 Hz, 1H), 8.12 (d, J = 2.5 Hz, 1H), 7.27 (d, J = 2.4 Hz, 1H), 3.87-3.69 (m, 4H), 3.44-3.34 (m, 4H); ESMS ($M+H^+$) = 227.14.

[00123] As shown in step 3-iii of Scheme 3, a mixture of 4-(8-bromoquinoxalin-6-yl)morpholine (1.57 g, 5.34 mmol), *tert*-butyl-*N*-(4-aminocyclohexyl)carbamate (1.37 g, 6.40 mmol), (*rac*)-BINAP (664 mg, 1.07 mmol), cesium carbonate (5.22 g, 16.0 mmol), and $\text{Pd}_2(\text{dba})_3$ (489 mg, 0.534 mmol) in toluene (50 mL) was heated at 100°C for 12 hours. After cooling, the mixture was diluted with ethyl acetate (150 mL) and water (25 mL), then filtered through diatomaceous earth which was subsequently washed with ethyl acetate. The combined organics were washed with brine, dried over sodium sulfate, concentrated under reduced pressure, and purified by medium pressure silica gel chromatography (0 to 60% EtOAc/hexanes gradient) to provide *tert*-butyl(-4-((7-morpholinoquinoxalin-5-yl)amino)cyclohexyl)carbamate (compound **1004**, 1.83g, 83.2% yield): ^1H -NMR (300 MHz, CDCl_3) δ 8.65 (d, J = 2.0 Hz, 1H), 8.35 (d, J = 2.0 Hz, 1H), 6.60 (d, J = 2.4 Hz, 1H), 6.34 (d, J = 2.4 Hz, 1H), 6.11 (d, J = 7.8 Hz, 1H), 4.60 (s, 1H), 3.97-3.86 (m, 4H), 3.67 (s, 2H), 3.41-3.25 (m, 4H), 1.85 (d, J = 3.0 Hz, 5H), 1.74-1.57 (m, 3H), 1.45 (s, 9H).


[00124] As shown in step 3-iv of Scheme 3, to a solution of *tert*-butyl (-4-((7-morpholinoquinoxalin-5-yl)amino)cyclohexyl)carbamate (900 mg, 2.00 mmol) in dichloromethane (16 mL) was added trifluoroacetic acid (3 mL, 38.9 mmol). The resulting black reaction mixture was stirred under an atmosphere of nitrogen at room temperature for 2 hours. Saturated aqueous sodium bicarbonate (150 mL) was added slowly until the color turned from black to orange. The mixture was extracted with dichloromethane (2 x 100 mL) and the combined organics washed with brine (50 mL), dried over sodium sulfate, and concentrated under reduced pressure to provide-*N*¹-(7-morpholinoquinoxalin-5-yl)cyclohexane-1,4-diamine, trifluoroacetate (compound **1005**): ^1H NMR (300 MHz, CDCl_3) δ 8.64 (d, J = 1.9 Hz, 1H), 8.36 (d, J = 1.9 Hz, 1H), 6.59 (d, J = 2.3 Hz, 1H), 6.34 (d, J = 2.3 Hz, 1H), 6.20 (d, J = 7.9 Hz, 1H), 3.95-3.84 (m, 4H), 3.69 (s, 1H), 3.41-3.25 (m, 4H), 2.93 (d, J = 8.9 Hz, 1H), 2.09-1.87 (m, 2H), 1.90-1.68 (m, 6H),

1.58 (dd, $J = 11.2, 8.7$ Hz, 2H); ESMS ($M+H^+$) = 328.34. This compound was used as is without further purification.

[00125] As shown in step 3-v of Scheme 3, to solution of N^1 -(7-morpholinoquinoxalin-5-yl)cyclohexane-1,4-diamine (25 mg, 0.07 mmol) and diisopropylethylamine (18.0 mg, 24.3 μ L, 0.14 mmol) in dichloromethane (750 μ L) was added ethyl chloroformate (11.4 mg, 10.0 μ L, 0.105 mmol). The reaction mixture was stirred for 12 hours, diluted with dichloromethane (10mL), washed with saturated aqueous sodium bicarbonate (5mL), dried over sodium sulfate, and concentrated under reduced pressure. The resulting residue was purified by HPLC preparative chromatography using a 10-90% acetonitrile/water (0.1% TFA) gradient as eluant to provide ethyl (4-((7-morpholinoquinoxalin-5-yl)amino)cyclohexyl)carbamate (compound **6**, 14 mg, 50% yield): $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.65 (d, $J = 2.0$ Hz, 1H), 8.36 (d, $J = 2.0$ Hz, 1H), 6.61 (d, $J = 2.4$ Hz, 1H), 6.35 (d, $J = 2.4$ Hz, 1H), 6.10 (d, $J = 7.6$ Hz, 1H), 4.72 (s, 1H), 4.12 (q, $J = 7.0$ Hz, 2H), 3.96-3.82 (m, 4H), 3.68 (s, 2H), 3.42-3.23 (m, 4H), 1.93-1.78 (m, 6H), 1.69 (dd, $J = 15.0, 6.3$ Hz, 2H), 1.25 (t, $J = 7.1$ Hz, 3H); ESMS ($M+H^+$) = 400.17.

[00126] As shown in step 3-vi of Scheme 3, A mixture of N^1 -(7-morpholinoquinoxalin-5-yl)cyclohexane-1,4-diamine (185 mg, 0.56 mmol), 2-bromopyrimidine (93 mg, 0.58 mmol), and triethylamine (143 mg, 197 μ L, 1.41 mmol) in 1-methylpyrrolidin-2-one (3 mL) was heated to 130°C and stirred for 15 hours. After cooling to room temperature, the mixture was diluted with ethyl acetate (70 mL) and methyl tert-butyl ether (20 mL), washed with water (3 x 20 mL), washed with brine (15 mL), dried over sodium sulfate, concentrated under reduced pressure, and purified by medium pressure silica gel chromatography (10 to 100% EtOAc/hexanes gradient) to provide N^1 -(7-morpholinoquinoxalin-5-yl)- N^4 -(pyrimidin-2-yl)cyclohexane-1,4-diamine (compound **18**, 102 mg, 45% yield): $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.65 (d, $J = 2.0$ Hz, 1H), 8.37 (d, $J = 2.0$ Hz, 1H), 8.27 (d, $J = 4.8$ Hz, 2H), 6.60 (d, $J = 2.4$ Hz, 1H), 6.51 (t, $J = 4.8$ Hz, 1H), 6.36 (d, $J = 2.4$ Hz, 1H), 6.15 (d, $J = 7.8$ Hz, 1H), 5.20 (d, $J = 7.7$ Hz, 1H), 4.04 (d, $J = 7.9$ Hz, 1H), 3.96-3.82 (m, 4H), 3.70 (s, 1H), 3.39-3.24 (m, 4H), 1.94 (dd, $J = 13.7, 4.4$ Hz, 6H), 1.78 (dt, $J = 28.8, 16.1$ Hz, 2H); ESMS ($M+H^+$) = 328.34.

Example 4. Preparation of 1-(4-((7-(8-oxa-3-azabicyclo[3.2.1]octan-3-yl)quinoxalin-5-yl)amino)cyclohexyl)-3-ethylurea (Compound **22**)\



Scheme 4

[00127] As shown in step 4-i of Scheme 4, to a solution of 5-bromo-7-fluoroquinoxaline (compound **1002**, 150 mg, 0.66 mmol) in NMP (2.3 mL) was added 8-oxa-3-azabicyclo[3.2.1]octane (178 mg, 1.2 mmol) at RT. The reaction mixture was sealed in a microwave vial and heated at 180°C for 20 minutes. After cooling to RT and pouring into water, the aqueous phase was extracted with EtOAc (3x). The combined extracts were dried over MgSO_4 , filtered, concentrated under reduced pressure, and purified by medium pressure silica gel chromatography (0 to 100% EtOAc/hexanes gradient) to provide 3-(8-bromoquinoxalin-6-yl)-8-oxa-3-azabicyclo[3.2.1]octane (compound **1006**, 87 mg, 41% yield) as a dark orange oil: ESMS ($\text{M}+\text{H}^+$) = 320.07.


[00128] As shown in step 4-ii of Scheme 4, a degassed solution of 3-(8-bromoquinoxalin-6-yl)-8-oxa-3-azabicyclo[3.2.1]octane (261 mg, 0.815 mmol), *tert*-butyl *N*-(4-aminocyclohexyl)carbamate (210 mg, 0.98 mmol), *rac*-BINAP (102 mg, 0.163 mmol), Cs_2CO_3 (797 mg, 2.45 mmol), and $\text{Pd}_2(\text{dba})_3$ (75 mg, 0.0815 mmol) in toluene (10.5 mL) was heated at 100°C (oil bath temp) in a sealed microwave tube for 15 hours. After cooling, the mixture was applied directly to a chromatography column and purified

by medium pressure silica gel chromatography (0 to 100% EtOAc/hexanes gradient) to afford *tert*-butyl (4-((7-(8-oxa-3-azabicyclo[3.2.1]octan-3-yl)quinoxalin-5-yl)amino)cyclohexyl)carbamate (compound **1007**, 141 mg, 36% yield) as a white solid: ¹H-NMR (400 MHz, CDCl₃) δ 8.49 (s, 1H), 8.23 (d, J = 1.5 Hz, 1H), 6.48 (s, 1H), 6.18 (d, J = 1.9 Hz, 1H), 6.06 (s, 1H), 4.52 (s, 1H), 4.47 (s, 2H), 3.60 (s, 2H), 3.45 (d, J = 11.6 Hz, 2H), 3.14-3.12 (m, 2H), 1.96-1.84 (m, 4H), 1.79 (s, 5H), 1.54 (s, 3H) and 1.38 (s, 9H) ppm; ESMS (M+H⁺) = 453.96.

[00129] As shown in step 4-iii of Scheme 4, to a solution of compound **1007** (141 mg, 0.295 mmol) in CH₂Cl₂ (2.5 mL) was added TFA (656 mg, 443 μL, 5.75 mmol) at RT. The resulting black solution was stirred for 2 hours and then the reaction was quenched by the addition of saturated NaHCO₃ until the black color gradually turned into an orange color. The reaction mixture was extracted with CH₂Cl₂ (3x) and the combined organic extracts were dried over Na₂SO₄ and evaporated to dryness to provide *N*¹-(7-(8-oxa-3-azabicyclo[3.2.1]octan-3-yl)quinoxalin-5-yl)cyclohexane-1,4-diamine, trifluoroacetate (compound **1008**): ESMS (M+H⁺) = 354.20. This material was used in subsequent reactions without any further purification.

[00130] As shown in step 4-iv of Scheme 4, to a solution of compound **1008** (45 mg, 0.071 mmol) and DIEA (36.5 mg, 49.0 μL, 0.28 mmol) in CH₂Cl₂ (1.4 mL) was added ethyl isocyanate (20 mg, 0.28 mmol) at RT. The solution was stirred at this temperature for 15 hours and then applied directly to a chromatography column and purified by medium pressure silica gel chromatograph (0 to 100% EtOAc/hexanes gradient) to afford 1-(4-((7-(8-oxa-3-azabicyclo[3.2.1]octan-3-yl)quinoxalin-5-yl)amino)cyclohexyl)-3-ethylurea (compound **22**, 8 mg, 27% yield) as a white solid: ¹H-NMR (400 MHz, CDCl₃) δ 8.54 (s, 1H), 8.22 (s, 1H), 6.43 (s, 1H), 6.19 (s, 1H), 6.02 (s, 1H), 4.47 (s, 2H), 4.38 (d, J = 5.2 Hz, 1H), 4.28 (s, 1H), 3.74 (s, 1H), 3.60 (s, 1H), 3.42 (s, 4H), 3.14-3.09 (m, 4H), 2.05-1.87 (m, 3H), 1.79 (s, 3H), 1.55 (d, J = 7.1 Hz, 2H) and 1.21-1.05 (m, 5H) ppm; ESMS (M+H⁺) = 425.35.

Example 5. Preparation of N^1 -(6-morpholinobenzo[*c*][1,2,5]oxadiazol-4-yl)- N^4 -(pyrimidin-2-yl)cyclohexane-1,4-diamine (Compound **23**)




Scheme 5

[00131] As shown in step 5-i of Scheme 5, a mixture of *tert*-butyl ((*cis*)-4-aminocyclohexyl)carbamate (compound **1009**, 490 mg, 2.3 mmol), 2-chloropyrimidine (262 mg, 2.3 mmol) and TEA (463 mg, 637 μL , 4.6 mmol) in DMF (10 mL) was subjected to microwave irradiation for 20 minutes at 150°C. The reaction mixture was diluted with EtOAc, washed with H_2O , dried over Na_2SO_4 , concentrated under reduced pressure, and purified by medium pressure silica gel chromatography (0 to 50 % EtOAc/hexanes gradient) to provide *tert*-butyl ((*cis*)-4-(pyrimidin-2-ylamino)cyclohexyl)carbamate (compound **1010**) as a white solid: ^1H -NMR (300 MHz, CDCl_3) δ 8.28 (d, $J = 4.8$ Hz, 2H), 6.53 (t, $J = 4.8$ Hz, 1H), 5.12 (s, 1H), 4.56 (s, 1H), 3.99 (dq, $J = 7.0, 3.5$ Hz, 1H), 3.65 (s, 1H), 1.83 (tq, $J = 10.2, 3.6$ Hz, 5H), 1.66 (s, 8H), 8.13–7.91 (m, 3H), 1.47 (s, 9H).

[00132] As shown in step 5-ii of Scheme 5, HCl (3 mL, 4M in THF, 12 mmol) was added to compound **1010**. The mixture was stirred for 30 min and concentrated under reduced pressure to produce (*cis*)- N^1 -(pyrimidin-2-yl)cyclohexane-1,4-diamine hydrochloride (compound **1011**). This material was used in subsequent reactions as is without further purification.

[00133] As shown in step 5-iii of Scheme 5, a mixture of 4-bromo-6-morpholinobenzo[*c*][1,2,5]oxadiazole (compound **1012**, 147 mg, 0.5 mmol), (*cis*)-*N*¹-(pyrimidin-2-yl)cyclohexane-1,4-diamine hydrochloride (120 mg, 0.6 mmol), (*rac*)-BINAP (32 mg, 0.05 mmol), Pd₂(dba)₃ (24 mg, 0.026 mmol), and cesium carbonate (506 mg, 1.55 mmol) in toluene (5 mL) was flushed with nitrogen gas and stirred overnight at 90°C under an atmosphere of nitrogen. The mixture was filtered through a layer of diatomaceous earth, concentrated under reduced pressure, and purified by medium pressure silica gel chromatography (0 to 80% EtOAc/hexanes gradient) to provide (*cis*)-*N*¹-(6-morpholinobenzo[*c*][1,2,5]oxadiazol-4-yl)-*N*⁴-(pyrimidin-2-yl)cyclohexane-1,4-diamine (compound **23**) as an orange solid: ¹H-NMR (300 MHz, CDCl₃) δ 8.20 (d, *J* = 4.9 Hz, 2H), 6.46 (t, *J* = 4.8 Hz, 1H), 6.05 (d, *J* = 1.6 Hz, 1H), 5.82 (s, 1H), 5.24 (s, 1H), 4.82 (d, *J* = 7.0 Hz, 1H), 3.98 (s, 1H), 3.85-3.72 (m, 4H), 3.60 (s, 1H), 3.23-3.06 (m, 4H), 1.95-1.62 (m, 8H).

Example 6. Preparation of 5-methoxy-N-((*cis*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)pyrimidin-2-amine (Compound **134**)



Scheme 6

[00134] As shown in step 6-i of Scheme 6, to a mixture of 5-bromo-2-fluoropyrimidine (1 g, 5.651 mmol) in iPrOH (10 mL) was added TEA (1.143 g, 1.574 mL, 11.30 mmol) and *trans*-4-aminocyclohexan-1-ol (650.8 mg, 5.651 mmol). The mixture was microwaved for 20min at 150°C, concentrated under reduced pressure, diluted with EtOAc, washed with water, and dried over Na₂SO₄. After removal of the volatiles under reduced pressure, the residue was purified by medium pressure silica gel chromatography (0-80% EtOAc/hexanes gradient) to provide (*trans*)-4-((5-bromopyrimidin-2-

yl)amino)cyclohexanol (compound **1013**, 1.2 g): $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.28 (s, 2H), 5.03 (d, $J = 8.1$ Hz, 1H), 3.91-3.49 (m, 2H), 2.31-1.90 (m, 4H), 1.56-1.19 (m, 4H)..

[00135] As shown in step 6-ii of Scheme 6, to compound **1013** (1.2 g, 4.41 mmol) in DCM (20 mL) was added TEA (1.134 g, 1.84 mL, 13.2 mmol) and MsCl (505 mg, 341 μL , 4.41 mmol). The reaction mixture was stirred for 1 hour, concentrated under reduced pressure, and purified by medium pressure silica gel chromatography (0 to 80% EtOAc/hexanes gradient) to provide *trans*-4-((5-bromopyrimidin-2-yl)amino)cyclohexyl methanesulfonate (compound **1014**): $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.29 (s, 2H), 5.03 (d, $J = 7.8$ Hz, 1H), 4.70 (tt, $J = 10.6, 3.9$ Hz, 1H), 3.80 (dt, $J = 11.2, 7.6, 3.7$ Hz, 1H), 3.04 (s, 3H), 2.30-2.12 (m, 4H), 1.93-1.69 (m, 2H), 1.51-1.33 (m, 2H).

[00136] As shown in step 6-iii of Scheme 6, to a solution of 2-amino-3-nitrophenol (5.00 g, 32.4 mmol) in dioxane (50 mL) was added bromine (6.22 g, 2.01 mL, 38.9 mmol). The mixture was stirred for 2 hours and a precipitate formed, which was collected and washed with dioxane and ether. The resulting yellow solid treated with a saturated NaHCO_3 solution, which was extracted with EtOAc (3x). The combined organics were dried over Na_2SO_4 , filtered, and concentrated under reduced pressure to yield 2-amino-5-bromo-3-nitrophenol (compound **1015**) as a brown solid. This material was carried on as is in subsequent reactions without further purification.

[00137] As shown in step 6-iv of Scheme 6, to a solution of 2-amino-5-bromo-3-nitrophenol (7.5 g, 31.8 mmol) in ethyl acetate (60 mL) was added Raney nickelTM (1.90 g, 214 μL , 32.4 mmol) and the reaction mixture was shaken for 2 hours under an atmosphere of H_2 at 30 p.s.i. After filtering and drying over Na_2SO_4 , the mixture was concentrated under reduced pressure to provide 2,3-diamino-5-bromophenol (compound **1016**), which was used as is in subsequent reactions without further purification.

[00138] As shown in step 6-v of Scheme 6, 2,3-diamino-5-bromophenol (6.0 g, 29.5 mmol) was dissolved in methanol and to this solution was added glyoxal (3.77 g, 2.98 mL, 64.9 mmol) and stirred overnight. The reaction mixture was concentrated under reduced pressure to a minimum volume and the resulting tan solid collected by filtration and dried under high vacuum to produce 7-bromoquinoxalin-5-ol (compound **1017**), which was used as is in subsequent reactions without further purification.

[00139] As shown in step 6-vi of Scheme 6, a solution of 7-bromoquinoxalin-5-ol (2.0 g, 8.89 mmol) in DCM (20 mL) was added imidazole (1.82 g, 26.7 mmol) and *tert*-


butyldimethylsilyl chloride (1.34 g, 1.65 mL, 8.89 mmol). The reaction mixture was stirred overnight at RT, concentrated under reduced pressure, and purified by medium pressure silica gel chromatography (0 to 20% EtOAc/hexanes gradient) to provide 7-bromo-5-((*tert*-butyldimethylsilyl)oxy)quinoxaline (compound **1018**) as a colorless oil: ¹H-NMR (300 MHz, CDCl₃) δ 8.69 (q, J = 1.8 Hz, 2H), 7.80 (d, J = 2.1 Hz, 1H), 7.22 (d, J = 2.1 Hz, 1H), 0.96 (s, 9H), 0.81 (s, 7H).

[00140] As shown in step 6-vii of Scheme 6, a mixture of 7-bromo-5-((*tert*-butyldimethylsilyl)oxy)quinoxaline (700 mg, 2.06 mmol), morpholine (270 mg, 270 μL, 3.09 mmol), Pd₂(dba)₃ (94.50 mg, 0.1032 mmol), (*rac*)-BINAP (129 mg, 0.206 mmol), cesium carbonate (2.02 g, 6.19 mmol) in toluene (7 mL) was flushed with nitrogen for 10 minutes. The mixture was then heated overnight at 100°C. After cooling, the reaction mixture was diluted with EtOAc, filtered through a layer of diatomaceous earth, concentrated under reduced pressure, and purified by medium pressure silica gel chromatography (0 to 30% EtOAc/hexanes gradient) to provide 7-morpholinoquinoxalin-5-ol. This compound (450 mg, 1.3 mmol) was dissolved in THF (20 mL) and tetra-*n*-butylammonium fluoride (539 mg, 2.06 mmol) was added. The reaction mixture was stirred for 0.5 hour, concentrated under reduced pressure, and purified by medium pressure silica gel chromatography (0 to 100% EtOAc/hexanes gradient) to provide 7-morpholinoquinoxalin-5-ol (compound **1019**) as a yellow solid: ¹H-NMR (300 MHz, CDCl₃) δ 8.75 (d, J = 2.0 Hz, 1H), 8.46 (d, J = 2.0 Hz, 1H), 7.70 (d, J = 41.8 Hz, 1H), 7.01 (d, J = 2.6 Hz, 1H), 6.89 (d, J = 2.5 Hz, 1H), 4.12-3.78 (m, 4H), 3.51-3.24 (m, 4H).

[00141] As shown in step 6-viii of Scheme 6, a solution of 7-morpholinoquinoxalin-5-ol (100 mg, 0.432 mmol), (*trans*)-4-((5-bromopyrimidin-2-yl)amino)cyclohexyl methanesulfonate (compound **1014**, 303 mg, 0.865 mmol), and CsCO₃ (282 mg, 0.865 mmol) in dioxane (1.0 mL) was stirred for 16 hours at 105°C. After cooling, the reaction mixture was diluted with EtOAc, filtered through diatomaceous earth, concentrated under reduced pressure, and purified by medium pressure silica gel chromatography (0 to 5% MeOH/DCM gradient) to produce 5-bromo-*N*-((*cis*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)pyrimidin-2-amine (compound **1020**, 110 mg) as a yellow foam: ¹H-NMR (400 MHz, CDCl₃) δ 8.70 (d, J = 2.0 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.29 (s, 2H), 6.98 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 5.29 (d, J = 8.3 Hz, 1H), 4.81 (s, 1H), 4.04-3.84 (m, 4H), 3.42-3.31 (m, 4H), 2.22 (s, 2H), 1.92 (d, J = 4.9 Hz, 6H).

[00142] As shown in step 6-ix of Scheme 6, to a mixture 5-bromo-*N*-((*cis*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)pyrimidin-2-amine (75 mg, 0.155 mmol), cesium carbonate (101 mg, 0.309 mmol), allylpalladium(II) chloride dimer (0.28 mg, 0.0015 mmol), RockPhos (2.17 mg, 0.0046 mmol) and MeOH (9.9 mg, 12.5 μ L, 0.31 mmol) in toluene (2 mL) was flushed with nitrogen gas and heated to 100°C for 18 hours. The reaction mixture was diluted with EtOAc, filtered through a layer of diatomaceous earth, and concentrated under reduced pressure. Purification by medium pressure silica gel chromatography (0-8% MeOH/DCM gradient) yielded 5-methoxy-*N*-((*cis*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)pyrimidin-2-amine (compound **134**, 43 mg): ¹H-NMR (300 MHz, CDCl₃) δ 8.70 (d, *J* = 1.9 Hz, 1H), 8.63 (d, *J* = 1.9 Hz, 1H), 8.07 (s, 2H), 6.96 (d, *J* = 2.5 Hz, 1H), 6.92 (d, *J* = 2.5 Hz, 1H), 5.01 (d, *J* = 8.1 Hz, 1H), 4.80 (q, *J* = 5.6, 4.2 Hz, 1H), 4.03-3.87 (m, 5H), 3.80 (s, 3H), 3.42-3.27 (m, 4H), 2.29-2.10 (m, 2H), 1.99-1.82 (m, 6H).

Example 7. Preparation of 4-(8-(((*trans*)-4-(pyrimidin-2-yloxy)cyclohexyl)oxy)quinoxalin-6-yl)morpholine (Compound **34**) and 4-(8-(((*cis*)-4-(pyrimidin-2-yloxy)cyclohexyl)oxy)-quinoxalin-6-yl)morpholine (Compound **42**)



Scheme 7

[00143] As shown in step 7-i of Scheme 7, to a solution of 1,4-dioxaspiro[4.5]decan-8-ol (compound **1021**, 1.0 g, 6.32 mmol) in DMF (10 mL) was added NaH (370 mg, 9.25 mmol). The reaction mixture was stirred for 20 minutes before the addition of 2-

chloropyrimidine (869 mg, 7.59 mmol). The mixture was stirred for 30 minute at RT and then heated to 100°C for 9 hours. After cooling, the mixture was diluted with EtOAc, washed with H₂O, dried over Na₂SO₄, concentrated under reduced pressure, and purified by medium pressure silica gel chromatography (0-40% EtOAc/hexanes) to produce 2-(1,4-dioxaspiro[4.5]decan-8-yloxy)pyrimidine (compound **1022**) as a colorless oil: ¹H NMR (300 MHz, Chloroform-d) δ 8.52 (d, J = 4.8 Hz, 2H), 6.92 (t, J = 4.8 Hz, 1H), 5.15 (ddd, J = 10.7, 6.5, 4.2 Hz, 1H), 4.05-3.87 (m, 4H), 2.14-1.85 (m, 6H), 1.79-1.65 (m, 2H); ESMS (M+H⁺) = 237.12.


[00144] As shown in step 7-ii of Scheme 7, to 2-(1,4-dioxaspiro[4.5]decan-8-yloxy)pyrimidine (620 mg, 2.624 mmol) was added HCl (4.0 mL of 6 M, 8.86 mmol) and the reaction mixture was stirred for 2 hours. The pH of the mixture was neutralized with sat. NaHCO₃(aq) and the mixture was concentrated under reduced pressure as a methanol azeotrope. To the residue was added DCM (30mL) to produce a precipitate, followed by stirring for an additional 20 minutes. The solids were filtered off and the mother liquor was concentrated under reduced pressure. The resulting residue was dissolved in methanol and sodium borohydride (151 mg, 3.99 mmol) was added as a solid. The mixture was stirred for 1 hour and the reaction quenched with HCl (6M, 0.70 mL). Stirring was continued until gas evolution ceased. The pH of the mixture was adjusted to about 8 with 1N sodium hydroxide and extracted with EtOAc (20mL). The organics were dried over sodium sulfate and concentrated under reduced pressure to produce 4-(pyrimidin-2-yloxy)cyclohexanol (compound **1023**, 248mg, 64% yield) as a mixture of (*cis*)- and (*trans*)- isomers. A 12 mg aliquot of the sample was purified via HPLC preparative reversed-phase chromatography (10-90% CH₃CN/water gradient containing 0.1% TFA) to separate the isomers: (*trans*)-4-pyrimidin-2-yloxy cyclohexanol - ¹H NMR (300 MHz, Chloroform-d) δ 8.54 (d, J = 4.8 Hz, 2H), 6.95 (t, J = 4.8 Hz, 1H), 5.05 (tt, J = 9.4, 4.0 Hz, 1H), 3.91-3.75 (m, 1H), 2.26-1.99 (m, 4H), 1.76-1.41 (m, 4H); ESMS (M+H⁺) = 195.07, (*cis*)-4-pyrimidin-2-yloxy cyclohexanol - ¹H NMR (300 MHz, Chloroform-d) δ 8.62 (d, J = 4.9 Hz, 2H), 7.04 (t, J = 4.9 Hz, 1H), 5.21 (tt, J = 5.3, 2.6 Hz, 1H), 4.56 (s, 1H), 3.85 (p, J = 5.9 Hz, 1H), 2.17-2.02 (m, 2H), 1.88-1.67 (m, 6H); ESMS (M+H⁺) = 195.07. The remaining material was used in subsequent reactions as the *cis/trans* mixture.

[00145] As shown in step 7-iii of Scheme 7, to a solution of a *cis/trans* mixture of 4-pyrimidin-2-yloxy cyclohexanol (244 mg, 1.256 mmol) and triethylamine (350 μL, 2.51

mmol) in dichloromethane (5 mL) was added methane sulfonyl chloride (145 μ L, 1.87 mmol). The reaction mixture was stirred for 2 hours, concentrated under reduced pressure, and purified by medium pressure silica gel chromatography (0- 20% EtOAc/dichloromethane gradient) to provide 4-pyrimidin-2-yloxycyclohexyl methanesulfonate (compound **1024**, 239 mg, 70% yield) as a mixture of *cis/trans* isomers: ^1H NMR (300 MHz, Chloroform-*d*) δ 8.51 (d, *J* = 4.8 Hz, 2H), 6.93 (t, *J* = 4.8 Hz, 1H), 5.13 (dq, *J* = 9.9, 3.0 Hz, 1H), 4.87 (p, *J* = 3.8 Hz, 1H), 3.04 (d, *J* = 2.4 Hz, 3H), 2.28-1.99 (m, 4H), 1.99-1.74 (m, 4H); ESMS ($\text{M}+\text{H}^+$) = 273.52.

[00146] As shown in step 7-iv of Scheme 7, a mixture of (4-pyrimidin-2-yloxycyclohexyl) methanesulfonate (105 mg, 0.386 mmol), 7-morpholinoquinoxalin-5-ol (178.3 mg, 0.7712 mmol), and Cs_2CO_3 (125.6 mg, 0.3856 mmol) in dioxane (1.5 mL) was sealed in a 5 mL microwave tube and heated to 110°C for 14 hours using an oil bath. The reaction mixture was cooled to room temperature, diluted with EtOAc, and filtered through diatomaceous earth which was subsequently washed with ethyl acetate. The filtrate was concentrated under reduced pressure and the residue purified via preparative reversed-phase HPLC (10-90% CH_3CN /water gradient containing 0.1% TFA). Fractions containing a mixture of *cis* and *trans* isomers were further purified via SFC using a chiral OJ column and eluting with 40% MeOH in CO_2 to provide 21mg of 4-(8-(((*trans*)-4-(pyrimidin-2-yloxy)cyclohexyl)oxy)quinoxalin-6-yl)morpholine (compound **34**): ^1H NMR (300 MHz, Chloroform-*d*) δ 8.69 (dd, *J* = 3.4, 1.9 Hz, 1H), 8.62 (dd, *J* = 3.6, 1.9 Hz, 1H), 8.51 (dd, *J* = 4.8, 2.2 Hz, 2H), 7.01-6.83 (m, 3H), 5.18 (tt, *J* = 7.0, 3.4 Hz, 1H), 4.79 (tt, *J* = 6.9, 3.1 Hz, 1H), 4.00-3.85 (m, 4H), 3.34 (dq, *J* = 4.8, 2.6 Hz, 4H), 2.44-2.16 (m, 4H), 1.92 (tdd, *J* = 16.4, 7.7, 2.8 Hz, 4H); ESMS ($\text{M}+\text{H}^+$) = 408.56, and 22 mg of 4-(8-(((*cis*)-4-(pyrimidin-2-yloxy)cyclohexyl)oxy)-quinoxalin-6-yl)morpholine (compound **42**): ^1H NMR (300 MHz, Chloroform-*d*) δ 8.70 (d, *J* = 1.9 Hz, 1H), 8.63 (d, *J* = 1.9 Hz, 1H), 8.52 (d, *J* = 4.8 Hz, 2H), 7.01-6.87 (m, 3H), 5.17 (ddt, *J* = 8.7, 6.7, 3.4 Hz, 1H), 4.76-4.58 (m, 1H), 4.00-3.87 (m, 4H), 3.40-3.27 (m, 4H), 2.43-2.22 (m, 4H), 2.05-1.87 (m, 2H), 1.86-1.71 (m, 2H); ESMS ($\text{M}+\text{H}^+$) = 408.56.

Example 8. *N*-[(*cis*)-4-[7-(3,6-dihydro-2*H*-pyran-4-yl)quinoxalin-5-yl]oxycyclohexyl]pyrimidin-2-amine (Compound **36**)




Scheme 8

[00147] As shown in step 8-i of Scheme 8, to a mixture of 7-bromoquinoxalin-5-ol (compound **1018**, 200 mg, 0.89 mmol) and cesium carbonate (579 mg, 1.78 mmol) in NMP (4.0 mL) was added (*trans*)-4-(pyrimidin-2-ylamino)cyclohexyl methanesulfonate (compound **1014**, 241.1 mg, 0.8887 mmol). The mixture was stirred for 18 hours at 90°C, at which time an additional 0.5 eq of compound **1014** (241 mg, 0.89 mmol) was added. After stirring at 90°C for an additional 6 hours, the reaction mixture was diluted with EtOAc, washed with H₂O, dried over Na₂SO₄, concentrated, , and purified by medium pressure silica gel chromatography (0-5% MeOH/DCM) to provide *N*-((*cis*)-4-((7-bromoquinoxalin-5-yl)oxy)cyclohexyl)pyrimidin-2-amine (compound **1025**): ¹H-NMR (300 MHz, CDCl₃) δ 9.01-8.77 (m, 2H), 8.29 (d, J = 4.8 Hz, 2H), 7.89 (d, J = 1.9 Hz, 1H), 7.25 (d, J = 2.0 Hz, 1H), 6.53 (t, J = 4.8 Hz, 1H), 5.43-5.22 (m, 1H), 4.79 (td, J = 5.2, 2.5 Hz, 1H), 4.18-3.95 (m, 1H), 3.51 (s, 1H), 2.22 (td, J = 10.2, 9.6, 5.4 Hz, 2H), 2.09-1.86 (m, 6H).

[00148] As shown in step 8-ii of Scheme 8, a mixture of *N*-((*cis*)-4-((7-bromoquinoxalin-5-yl)oxy)cyclohexyl)pyrimidin-2-amine (compound **1025**, 52 mg, 0.1299 mmol), Pd(dppf)Cl₂ (10.61 mg, 0.01299 mmol), 2-(3,6-dihydro-2H-pyran-4-yl)-4,4,5,5-tetramethyl-1,3,2-dioxaborolane (compound **1026**, 27.3 mg, 0.13 mmol), Na₂CO₃ (195 μL of 2M (aq) solution, 0.39 mmol) in DMF (1 mL) was flushed with nitrogen gas for 10 minutes. The mixture was subjected to microwave radiation for 20 min at 150 °C. After cooling, the mixture was diluted with EtOAc, washed with H₂O, dried over Na₂SO₄, concentrated under reduced pressure and purified by medium pressure silica gel chromatography (0-5% MeOH/DCM) to provide *N*-[(*cis*)-4-[7-(3,6-dihydro-2H-pyran-4-

yl]quinoxalin-5-yl]oxycyclohexyl]pyrimidin-2-amine (compound **36**) as an off-white solid: $^1\text{H-NMR}$ (300 MHz, CDCl_3) δ 8.94-8.76 (m, 2H), 8.29 (d, $J = 4.8$ Hz, 2H), 7.67 (d, $J = 1.7$ Hz, 1H), 6.53 (t, $J = 4.8$ Hz, 1H), 6.37 (tt, $J = 3.1, 1.5$ Hz, 1H), 5.30 (d, $J = 7.9$ Hz, 1H), 4.87 (dt, $J = 7.5, 3.6$ Hz, 1H), 4.43 (q, $J = 2.8$ Hz, 2H), 4.02 (t, $J = 5.5$ Hz, 3H), 2.68 (dq, $J = 6.0, 3.4, 3.0, 1.8$ Hz, 2H), 2.35-2.11 (m, 2H), 2.07-1.84 (m, 6H); ESMS ($\text{M}+\text{H}^+$) = 404.2.

Example 9. *N*-((*cis*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)pyrimidin-2-amine (Compound **28**)



Scheme 9

[00149] As shown in step 9-i of Scheme 9, 7-bromoquinoxalin-5-ol (compound **1017**, 5.4 g, 24.0 mmol), 2-((*trans*)-4-hydroxycyclohexyl)isoindoline-1,3-dione (5.607 g, 22.86 mmol), and triphenylphosphine (8.994 g, 7.945 mL, 34.29 mmol) were dissolved in anhydrous THF and the flask was cooled in an ice bath. DIAD (6.93 g, 6.64 mL, 34.3 mmol) was added dropwise and the reaction was stirred at 0°C for 5 minutes, then warmed to room temperature and stirred for 18 hours. The reaction mixture was concentrated under reduced pressure, the residue was treated with Et_2O and stirred for 0.5 hour at RT, the precipitates filtered off, the filtrate concentrated under reduced pressure, and the residue purified by medium pressure silica gel chromatography (0-50% EtOAc/hexanes

gradient) to produce 2-[(*cis*)-4-(7-bromoquinoxalin-5-yl)oxycyclohexyl]isoindoline-1,3-dione (compound **1028**, 6.2 g, 60% yield): ¹H-NMR (300 MHz, CDCl₃) δ 8.95 (d, J = 1.8 Hz, 1H), 8.86 (d, J = 1.8 Hz, 1H), 7.91 (d, J = 2.0 Hz, 1H), 7.88-7.80 (m, 2H), 7.77-7.68 (m, 2H), 7.31 (d, J = 2.0 Hz, 1H), 4.96 (t, J = 2.9 Hz, 1H), 4.29 (tt, J = 12.5, 3.8 Hz, 1H), 2.88 (qd, J = 12.9, 3.6 Hz, 2H), 2.54-2.32 (m, 2H), 1.94-1.61 (m, 4H).


[00150] As shown in step 9-ii of Scheme 9, In a round bottom flask fitted with a condenser, a mixture of 2-[4-(7-bromoquinoxalin-5-yl)oxycyclohexyl]isoindoline-1,3-dione (6.2 g, 12.34 mmol), morpholine (1.61 g, 1.62 mL, 18.5 mmol), and Cs₂CO₃ (12.06 g, 37.0 mmol) in anhydrous toluene (73 mL) was treated with rac-BINAP (768.4 mg, 1.234 mmol) and Pd₂(dba)₃ (565 mg, 0.617 mmol). The reaction mixture was heated at 110°C for 18 hours. After cooling to room temperature, the mixture was filtered through diatomaceous earth and concentrated under reduced pressure. The residue was triturated with Et₂O and the solids collected by filtration and washed with Et₂O to produce 2-[(*cis*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl]isoindoline-1,3-dione (compound **1029**, 4.2 g) as yellow solid. The filtrate was concentrated under reduced pressure and purified by medium pressure silica gel chromatography (0-100% EtOAc/hexanes gradient) to produce an additional 300 mg of compound **1029**: ¹H-NMR (300 MHz, CDCl₃) δ 8.76-8.63 (m, 2H), 7.85 (dd, J = 5.4, 3.1 Hz, 2H), 7.79-7.60 (m, 2H), 7.09 (d, J = 2.6 Hz, 1H), 6.99 (d, J = 2.5 Hz, 1H), 5.06 (t, J = 2.8 Hz, 1H), 4.27 (tt, J = 12.3, 3.8 Hz, 1H), 4.02-3.85 (m, 4H), 3.49-3.27 (m, 4H), 3.03-2.75 (m, 2H), 2.37 (d, J = 14.0 Hz, 2H), 1.83-1.56 (m, 4H).

[00151] As shown in step 9-iii of Scheme 9, to a suspension of 2-[(*cis*)-4-(7-morpholinoquinoxalin-5-yl)oxycyclohexyl]isoindoline-1,3-dione (2.3 g, 5.02 mmol) in MeOH (25 mL) was added hydrazine (321 mg, 315 μL, 10.0 mmol) and the reaction mixture stirred for 18 hours at RT, over which time the initial suspension became homogenous followed by the appearance of a precipitate. Et₂O (30 mL) was added and the reaction mixture stirred an additional 30 minutes. The precipitates were filtered off, the filtrate concentrated under reduced pressure, the residue treated with DCM (30 mL), and any remaining solids removed by filtration. The filtrate was concentrated under reduced pressure to provide (*cis*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexanamine (compound **1030**), which was used as is in subsequent reactions: ¹H-NMR (300 MHz, CDCl₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 6.95 (d, J = 2.5 Hz, 1H), 6.90

(d, $J = 2.5$ Hz, 1H), 5.00-4.67 (m, 3H), 4.03-3.81 (m, 4H), 3.49 (s, 1H), 3.43-3.25 (m, 4H), 2.88 (q, $J = 6.2$ Hz, 2H), 2.36-1.96 (m, 6H).

[00152] As shown in step 9-iv of Scheme 9, to a solution so (*cis*) 4-(7-morpholinoquinoxalin-5-yl)oxycyclohexanamine (415 mg, 1.264 mmol) and 2-methylsulfonylpyrimidine (400 mg, 2.53 mmol) was added DIEA (490 mg, 661 μ L, 3.79 mmol) and the reaction mixture was sealed in a vessel and heated to 100°C for 16 hours. After this time, the volatiles were removed under a stream of nitrogen gas and the crude residue dissolved in minimal amount of DCM. Purification by medium pressure silica gel chromatography (0-10% MeOH/DCM, 1% Et₃N] produced *N*-((*cis*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)pyrimidin-2-amine containing triethylamine hydrochloride as an impurity. Dissolved product in DCM and stirred with a silica-supported amine (Silabond amine® 40-63 μ m). The scavenger mixture was filtered, concentrated under reduced pressure, and dried under high vacuum to provide *N*-((*cis*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)pyrimidin-2-amine (Compound **28**, 435 mg): ¹H-NMR (400 MHz, CDCl₃) δ 8.68 (d, $J = 1.9$ Hz, 1H), 8.61 (d, $J = 1.9$ Hz, 1H), 8.27 (s, 1H), 8.26 (s, 1H), 6.94 (d, $J = 2.4$ Hz, 1H), 6.90 (d, $J = 2.4$ Hz, 1H), 6.50 (t, $J = 4.8$ Hz, 1H), 4.78 (s, 1H), 4.08 - 3.97 (m, 1H), 3.94 - 3.86 (m, 4H), 3.37 - 3.28 (m, 4H), 2.20 (d, $J = 9.1$ Hz, 2H), 1.95 - 1.85 (m, 6H).


Example 10. Preparation of N-[4-[7-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)quinoxalin-5-yl]oxycyclohexyl]pyrimidin-2-amine (Compound **291**)



Scheme 10a


[00153] To a mixture of 7-bromoquinoxalin-5-ol (47.53 g, 211.2 mmol), 2-(4-hydroxycyclohexyl)isoindoline-1,3-dione (52.41 g, 213.7 mmol), and PPh₃ (87.31 g, 332.9 mmol) in THF (740 mL) at 21 °C was added tert-butyl (NZ)-N-tert-butoxycarbonyliminocarbamate (DTBAD) (79.51 g, 328.0 mmol) in portions over 40 min so as to maintain the temperature below 30 °C and the resultant reaction mixture was stirred at room temperature for a further 20 h.

[00154] The reaction was evaporated in vacuo. The residual reddish-brown viscous oil was dissolved in CH₂Cl₂ and filtered through a plug of silica in a glass column using applied air pressure (plug was made with 1L of dry silica suspended in CH₂Cl₂). The plug was eluted with CH₂Cl₂, the fractions were combined and evaporated in vacuo to afford a red-brown viscous oil/foam, that was then dissolved in 700 mL of MeOH before precipitating. The mixture was stirred at room temperature for 1 h, filtered, washed with cold MeOH (500 mL) and Et₂O (100 mL), then dried in vacuo to yield a tan solid that was suspended in 300 mL MeOH and brought to reflux for 10 min. The suspension was cooled to room temperature and filtered, washed with a further MeOH and Et₂O (4:1), and dried in vacuo to provide 2-[4-(7-bromoquinoxalin-5-yl)oxycyclohexyl]isoindoline-1,3-dione (58.43 g, 126.6 mmol, 59.94%). ¹H NMR (400 MHz, CDCl₃) δ 8.96 (d, J = 1.8 Hz, 1H), 8.86 (d, J = 1.8 Hz, 1H), 7.91 (d, J = 1.9 Hz, 1H), 7.89 - 7.82 (m, 2H), 7.78 - 7.67 (m, 2H), 7.30 (d, J = 1.9 Hz, 1H), 4.95 (s, 1H), 4.29 (tt, J = 12.5, 3.7 Hz, 1H), 2.87 (qd, J = 13.1, 3.5 Hz, 2H), 2.44 (d, J = 15.2 Hz, 2H), 1.80 (t, J = 14.1 Hz, 2H), 1.67 (d, 2H). ESI-MS m/z calc. 451.05316, found 452.19 (M+1)⁺; Retention time: 0.92 minutes.



Scheme 10b


[00155] A mixture of 2-[4-(7-bromoquinoxalin-5-yl)oxycyclohexyl]isoindoline-1,3-dione (1 g, 2.211 mmol), 6-oxa-3-azabicyclo[3.1.1]heptane HCl (180 mg, 1.328 mmol), cesium carbonate (2.161 g, 6.633 mmol), Pd2(dba)3 (202.5 mg, 0.2211 mmol) and rac-BINAP (275.3 mg, 0.4422 mmol) in dioxane (5 mL) was stirred overnight at 70 °C, then heated in a microwave reactor for 15 min at 150 °C. The reaction was then diluted with methylene chloride, filtered through Celite, and concentrated. Silica gel flash column chromatography (0-5% MeOH/DCM) yielded 2-[4-[7-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)quinoxalin-5-yl]oxycyclohexyl]isoindoline-1,3-dione (750 mg, 72.1%) as a yellow solid that was carried on to the next reaction.



Scheme 10c

[00156] To a solution of 2-[4-[7-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)quinoxalin-5-yl]oxycyclohexyl]isoindoline-1,3-dione (800 mg, 1.700 mmol) in EtOH (10 mL) was added hydrazine monohydrate (85.10 mg, 83.35 μ L, 1.700 mmol) and the reaction was stirred at reflux overnight, then concentrated, diluted with DCM, and filtered. The filtrate


was concentrated, and purified on a 40 g silica gel cartridge with 0-50% (20% NH₃/MeOH) to yield 4-[7-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)quinoxalin-5-yl]oxycyclohexanamine (450 mg, 77.8%) as a yellow solid. ¹H NMR (300 MHz, Chloroform-d) δ 8.65 (d, J = 1.9 Hz, 1H), 8.53 (d, J = 1.9 Hz, 1H), 6.83 (q, J = 2.6 Hz, 2H), 4.83 (t, J = 6.0 Hz, 4H), 3.87 - 3.60 (m, 5H), 3.34 (dt, J = 8.7, 6.6 Hz, 1H), 3.01 - 2.83 (m, 1H), 2.23 (dq, J = 11.3, 5.8, 4.8 Hz, 2H), 2.07 (d, J = 8.7 Hz, 1H), 1.92 - 1.62 (m, 6H).



Scheme 10d

[00157] A mixture of 4-[7-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)quinoxalin-5-yl]oxycyclohexanamine (190 mg, 0.5581 mmol), 2-fluoropyrimidine (60 mg, 0.6118 mmol) and DIEA (200 μ L, 1.148 mmol) in 2-propanol (2 mL) was heated in a microwave reactor for 20 min at 150 °C. The reaction mixture was concentrated, and then purified from 12 g silica gel cartridge with 0-6% MeOH/DCM to yield N-[4-[7-(6-oxa-3-azabicyclo[3.1.1]heptan-3-yl)quinoxalin-5-yl]oxycyclohexyl]pyrimidin-2-amine (120.2 mg, 48.9%) as a yellow solid. Mass + 1: 419.23; Retention Time: 0.72; NMR Annotation: ¹H NMR (400 MHz, Chloroform-d) δ 8.42 (d, J = 1.9 Hz, 1H), 8.30 (d, J = 1.9 Hz, 1H), 8.04 (d, J = 4.8 Hz, 2H), 6.65 - 6.56 (m, 2H), 6.28 (t, J = 4.8 Hz, 1H), 4.99 (d, J = 8.1 Hz, 1H), 4.60 (d, J = 6.5 Hz, 3H), 3.79 (dd, J = 8.2, 4.0 Hz, 0H), 3.62 - 3.38 (m, 4H), 3.17 - 3.03 (m, 1H), 2.07 - 1.90 (m, 2H), 1.89 - 1.59 (m, 7H).


Example 11. Preparation of 6-(4-methylpiperazin-1-yl)-*N*-((1*s*,4*s*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)pyrimidin-4-amine (Compound **537**)



Scheme 11

[00158] A suspension of 4-(7-morpholinoquinoxalin-5-yl)oxycyclohexanamine (500 mg, 1.52 mmol), 4-chloro-6-(4-methylpiperazin-1-yl)pyrimidine (324 mg, 1.52 mmol), and NaOt-Bu (440 mg, 4.58 mmol) in *t*-BuOH (10.1 mL) was degassed by bubbling N₂ through the mixture for 10 min. *t*-BuXPhos Palladacycle (53 mg, 0.077 mmol) was added, and the reaction mixture was sealed hermetically and heated to 100 °C in an oil bath for 2 h. The solvent was removed *in vacuo*, and the crude residue was purified by silica gel chromatography (40 g Isco gold column, linear gradient 0% → 10% MeOH/CH₂Cl₂ [+0.1% Et₃N]) to yield 6-(4-methylpiperazin-1-yl)-*N*-((1*s*,4*s*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)pyrimidin-4-amine as a yellow solid (654.7 mg, 83.5% yield). ¹H NMR (400 MHz, Chloroform-*d*) δ 8.69 (d, *J* = 1.8 Hz, 1H), 8.61 (d, *J* = 1.8 Hz, 1H), 8.19 (s, 1H), 6.95 (d, *J* = 2.4 Hz, 1H), 6.90 (d, *J* = 2.3 Hz, 1H), 5.42 (s, 1H), 4.78 (s, 1H), 4.70 (d, *J* = 8.0 Hz, 1H), 3.97 - 3.87 (m, 4H), 3.83 (s, 1H), 3.64 - 3.50 (m, 4H), 3.40 - 3.27 (m, 4H), 2.53 - 2.42 (m, 4H), 2.34 (s, 3H), 2.19 (d, *J* = 9.2 Hz, 2H), 1.97 - 1.77 (m, 6H). ESI-MS *m/z* calc. 504.2961, found 505.44 (M+1)⁺; Retention time: 0.5 minutes.

Example 12. Preparation of 2-(4-methylpiperazin-1-yl)-*N*-((1*s*,4*s*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)pyrimidin-4-amine (Compound **535**)




Scheme 12

[00159] To a solution of 4-(7-morpholinoquinoxalin-5-yl)oxycyclohexanamine (500 mg, 1.522 mmol) and Et₃N (212 μ L, 1.522 mmol) in EtOH (7.25 mL) was added 2,4-dichloropyrimidine (216.0 mg, 1.450 mmol). The resultant reaction solution was heated to reflux for 5 h, then cooled to room temperature. The solvent was removed *in vacuo*, and the crude residue was purified by silica gel chromatography (40 g Isco gold column, linear gradient 0% $\rightarrow$ 10% MeOH/CH₂Cl₂ [+0.1% Et₃N]) to provide 2-chloro-*N*-[4-(7-morpholinoquinoxalin-5-yl)oxycyclohexyl]pyrimidin-4-amine as a yellow solid (485 mg, 75% yield). ¹H NMR (300 MHz, Chloroform-*d*) δ 8.70 (d, *J* = 2.0 Hz, 1H), 8.60 (d, *J* = 2.0 Hz, 1H), 8.00 (d, *J* = 4.8 Hz, 1H), 6.96 (d, *J* = 2.5 Hz, 1H), 6.89 (d, *J* = 2.5 Hz, 1H), 6.23 (d, *J* = 5.9 Hz, 1H), 5.18 (d, *J* = 5.7 Hz, 1H), 4.80 (s, 1H), 3.99 - 3.84 (m, 4H), 3.42 - 3.21 (m, 4H), 2.32 - 2.08 (m, 2H), 2.05 - 1.73 (m, 6H). ESI-MS *m/z* calc. 440.17276, found 441.25 (M+1)⁺; Retention time: 0.69 minutes.

[00160] A solution of 2-chloro-*N*-[4-(7-morpholinoquinoxalin-5-yl)oxycyclohexyl]pyrimidin-4-amine (236 mg, 0.5352 mmol), 1-methylpiperazine (190


μL , 1.711 mmol), and Hunig's base (315 μL , 1.808 mmol) in *i*-PrOH (3.6 mL) was heated to 170 °C in a microwave for 1 h. The solvent was removed *in vacuo*. The crude residue was purified by silica gel chromatography (40 g Isco gold column, linear gradient 0% $\rightarrow$ 10% MeOH/CH₂Cl₂ [+0.1% Et₃N]) to provide 2-(4-methylpiperazin-1-yl)-*N*-((1*s*,4*s*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)pyrimidin-4-amine as a yellow solid (219 mg, 79% yield). ¹H NMR (400 MHz, Chloroform-*d*) δ 8.69 (d, *J* = 1.9 Hz, 1H), 8.61 (d, *J* = 1.9 Hz, 1H), 7.88 (d, *J* = 5.7 Hz, 1H), 6.95 (d, *J* = 2.4 Hz, 1H), 6.89 (d, *J* = 2.4 Hz, 1H), 5.67 (d, *J* = 5.8 Hz, 1H), 4.84 - 4.63 (m, 2H), 3.99 - 3.83 (m, 5H), 3.82 - 3.73 (m, 4H), 3.37 - 3.27 (m, 4H), 2.49 - 2.40 (m, 4H), 2.33 (s, 3H), 2.23 - 2.11 (m, 2H), 1.96 - 1.82 (m, 6H). ESI-MS *m/z* calc. 504.2961, found 505.35 (M+1)⁺; Retention time: 0.51 minutes.

Example 13. Preparation of *N*-methyl-6-(((1*s*,4*s*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)amino)pyrimidine-4-carboxamide (Compound **359**)



Scheme 13a

[00161] A solution of oxalyl chloride (186.2g, 256.0 mmol) and DMF (1.7 mL, 21.96 mmol) was added to a suspension of 6-chloropyrimidine-4-carboxylic acid (9.1 g, 57.40 mmol) in dichloromethane (300 mL) dropwise via dropping funnel over 20min. Allowed to stir for 2hr and concentrated the acid chloride under reduced pressure. The acid chloride was dissolved in dichloromethane (250mL) and to it was added a solution of methylamine (30.71 mL of 40 %w/v, 395.5 mmol) in water and NaHCO₃ (188.3 mL of 1.2 M, 226.0 mmol) dropwise. The reaction mixture was allowed to stir overnight. The layers were separated and extracted the aqueous layer with dichloromethane (200mL). The combined organics were washed with brine, dried over sodium sulfate, and concentrated under reduced pressure. Chromatographed over 120g silica gel using 0-50% EtOAc/Heptane as eluent. 4.356g (44.94% yield) of 6-chloro-*N*-methylpyrimidine-4-carboxamide was obtained as a white solid. ¹H NMR (400 MHz, Chloroform-*d*) δ 8.99 (d, *J* = 1.1 Hz, 1H), 8.16 (d, *J* = 1.1 Hz, 1H), 7.90 (s, 1H), 3.06 (d, *J* = 5.2 Hz, 3H).



Scheme 13b


[00162] A solution of 4-(7-morpholinoquinoxalin-5-yl)oxycyclohexanamine (7.593 g, 23.12 mmol), 6-chloro-N-methyl-pyrimidine-4-carboxamide (4.35 g, 25.35 mmol), and t-BuXPhos palladacycle (1.519 g, 2.333 mmol) in tBuOH (100 mL) was added sodium t-butoxide (25.5 mL of 2 M, 51.00 mmol). Allowed to stir at room temperature under N₂ overnight. The reaction was diluted with dichloromethane (100mL) and filtered through Celite with dichloromethane wash. The filtrate was concentrated under reduced pressure and chromatographed over 330g silica gel using 0→14% Methanol/DCM as eluent. The resulting product was dissolve in small amount of ethanol (15mL) and the product precipitated over 1h. Filtered with cold ethanol wash to give a yellow solid. the solid was dried under vacuum over for 48h at 50°C to yield *N*-methyl-6-(((1*S*,4*S*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)amino)pyrimidine-4-carboxamide (2.58g) as a yellow solid. ¹H NMR (300 MHz, Chloroform-*d*) δ 8.70 (d, *J* = 1.9 Hz, 1H), 8.61 (d, *J* = 1.9 Hz, 1H), 8.53 - 8.44 (m, 1H), 8.00 (d, *J* = 5.6 Hz, 1H), 7.20 (d, *J* = 1.2 Hz, 1H), 6.96 (d, *J* = 2.5 Hz, 1H), 6.90 (d, *J* = 2.5 Hz, 1H), 5.50 (s, 1H), 4.83 (dt, *J* = 4.8, 2.5 Hz, 1H), 3.97 - 3.87 (m, 4H), 3.39 - 3.29 (m, 4H), 3.01 (d, *J* = 5.1 Hz, 3H), 2.31 - 2.19 (m, 2H), 2.06 - 1.70 (m, 6H). ESI-MS *m/z* calc. 463.2332, found 464.25 (*M*+1)⁺; Retention time: 0.59 minutes.

Example 14. Preparation of *N*-methyl-2-(((1*s*,4*s*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)amino)pyrimidine-4-carboxamide (Compound **350**)



Scheme 14a

[00163] A solution of 2-chloropyrimidine-4-carboxylic acid (4.425 g, 27.91 mmol) in DMF (35.40 mL) was added 1-hydroxybenzotriazole (Water (1)) (2.982 g, 19.47 mmol) and EDCI (Hydrochloric Acid (1)) (6.372 g, 33.24 mmol) at room temperature. After stirred for 10 min, to the mixture was added METHYL AMINE in tetrahydrofuran (20.93 mL of 2 M, 41.86 mmol) and stirred for 2h. The reaction was diluted with ethyl acetate, washed with water, and dried over sodium sulfate then concentrated. The resulting residue was purified from silica gel column 0-30% ethyl acetate/ hexanes to obtain 2-chloro-*N*-methylpyrimidine-4-carboxamide as a white solid. ¹H NMR (300 MHz, Chloroform-*d*) δ 8.85 (d, *J* = 4.9 Hz, 1H), 8.07 (d, *J* = 4.9 Hz, 1H), 7.82 (s, 1H), 7.27 (s, 0H), 3.06 (d, *J* = 5.2 Hz, 3H).





Scheme 14b





[00164] A solution of 4-((7-morpholinoquinoxalin-5-yl)oxycyclohexanamine (6.013 g, 18.31 mmol), 2-chloro-*N*-methyl-pyrimidine-4-carboxamide (3.41 g, 19.87 mmol) and *t*-BuXPhos palladacycle (1.220 g, 1.873 mmol) in *t*BuOH (100 mL) was added sodium *t*-butoxide (20.52 mL of 2 M, 41.04 mmol). Allowed to stir at rt under N₂ overnight. Diluted rxn with dichloromethane (100mL), filtered through celite and concentrated





filtrate under reduced pressure. The resulting residue was chromatographed over 220g silica gel using 0→14% Methanol/dichloromethane. The product was triturated with ethanol (40mL) at 70°C for 2h, filtered and dried under vacuum at 50°C overnight to obtain *N*-methyl-2-(((1*s*,4*s*)-4-((7-morpholinoquinoxalin-5-yl)oxy)cyclohexyl)amino)pyrimidine-4-carboxamide (5.03g, 62%yield) as a fine yellow powder. ¹H NMR (300 MHz, Chloroform-*d*) δ 8.69 (d, *J* = 1.9 Hz, 1H), 8.62 (d, *J* = 1.9 Hz, 1H), 8.48 (d, *J* = 4.9 Hz, 1H), 7.75 (s, 1H), 7.32 (d, *J* = 4.9 Hz, 1H), 6.96 (d, *J* = 2.5 Hz, 1H), 6.91 (d, *J* = 2.5 Hz, 1H), 5.30 (d, *J* = 7.9 Hz, 1H), 4.85 - 4.73 (m, 1H), 4.07 (dd, *J* = 8.5, 4.7 Hz, 1H), 3.99 - 3.82 (m, 4H), 3.41 - 3.20 (m, 4H), 3.01 (d, *J* = 5.1 Hz, 3H), 2.20 (q, *J* = 6.0 Hz, 2H), 2.06 - 1.82 (m, 6H). ESI-MS *m/z* calc. 463.2332, found 464.31 (M+1)⁺; Retention time: 0.71 minutes

[00165] Tables 1 and 2 provides analytical characterization data for certain compounds of formula **I** (blank cells indicate that the test was not performed).





Table 1.





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
1		370.52	(CDCl ₃) δ 8.64 (d, <i>J</i> = 2.0 Hz, 1H), 8.36 (d, <i>J</i> = 2.0 Hz, 1H), 6.61 (d, <i>J</i> = 2.4 Hz, 1H), 6.34 (d, <i>J</i> = 2.4 Hz, 1H), 5.90 (s, 1H), 5.40 (d, <i>J</i> = 7.7 Hz, 1H), 3.98 - 3.76 (m, 5H), 3.51 - 3.24 (m, 5H), 2.33 - 2.08 (m, 4H), 1.99 (s, 3H), 1.58 - 1.31 (m, 4H)
2		476.61	(CDCl ₃) δ 8.63 (d, <i>J</i> = 1.9 Hz, 1H), 8.34 (d, <i>J</i> = 1.9 Hz, 1H), 7.17 (d, <i>J</i> = 8.6 Hz, 2H), 6.90 (d, <i>J</i> = 8.5 Hz, 2H), 6.59 (d, <i>J</i> = 2.2 Hz, 1H), 6.29 (d, <i>J</i> = 2.2 Hz, 1H), 5.86 (d, <i>J</i> = 7.6 Hz, 1H), 5.29 (d, <i>J</i> = 8.0 Hz, 1H), 3.96 - 3.75 (m, 8H), 3.52 (s, 2H), 3.39 - 3.22 (m, 5H), 2.19 (d, <i>J</i> = 11.7 Hz, 2H), 2.12 - 1.89 (m, 2H), 1.43 (td, <i>J</i> = 13.0, 2.4 Hz, 2H), 1.32 - 1.15 (m, 2H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
3		428.49	(CDCl ₃) δ 8.65 (d, J = 2.0 Hz, 1H), 8.35 (d, J = 2.0 Hz, 1H), 6.60 (d, J = 2.4 Hz, 1H), 6.34 (d, J = 2.4 Hz, 1H), 6.11 (d, J = 7.8 Hz, 1H), 4.60 (s, 1H), 3.97 - 3.86 (m, 4H), 3.67 (s, 2H), 3.41 - 3.25 (m, 4H), 1.85 (d, J = 3.0 Hz, 5H), 1.74 - 1.57 (m, 3H), 1.45 (s, 9H)
4		370.46	(CDCl ₃) δ 8.64 (d, J = 1.9 Hz, 1H), 8.34 (d, J = 1.9 Hz, 1H), 6.61 (d, J = 2.2 Hz, 1H), 6.35 (d, J = 2.3 Hz, 1H), 6.12 (br s, 1H), 5.56 (d, J = 7.5 Hz, 1H), 4.11 - 3.81 (m, 5H), 3.70 (m, 1H), 3.42 - 3.24 (m, 4H), 1.99 (s, 3H), 1.86 (m, 6H), 1.75 - 1.51 (m, 2H)
5		406.12	(CDCl ₃) δ 8.65 (d, J = 2.0 Hz, 1H), 8.36 (d, J = 2.0 Hz, 1H), 6.62 (d, J = 2.4 Hz, 1H), 6.34 (d, J = 2.3 Hz, 1H), 6.09 (d, J = 7.6 Hz, 1H), 4.68 (d, J = 7.2 Hz, 1H), 3.97 - 3.82 (m, 4H), 3.76 - 3.47 (m, 2H), 3.40 - 3.23 (m, 4H), 3.01 (s, 3H), 2.04 - 1.65 (m, 8H)
6		400.17	(CDCl ₃) δ 8.65 (d, J = 2.0 Hz, 1H), 8.36 (d, J = 2.0 Hz, 1H), 6.61 (d, J = 2.4 Hz, 1H), 6.35 (d, J = 2.4 Hz, 1H), 6.10 (d, J = 7.6 Hz, 1H), 4.72 (s, 1H), 4.12 (q, J = 7.0 Hz, 2H), 3.96 - 3.82 (m, 4H), 3.68 (s, 2H), 3.42 - 3.23 (m, 4H), 1.93 - 1.78 (m, 6H), 1.69 (dd, J = 15.0, 6.3 Hz, 2H), 1.25 (t, J = 7.1 Hz, 3H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
7		442.14	(CDCl ₃) δ 8.65 (d, J = 2.0 Hz, 1H), 8.35 (d, J = 2.0 Hz, 1H), 6.61 (d, J = 2.4 Hz, 1H), 6.35 (d, J = 2.3 Hz, 1H), 6.10 (d, J = 7.7 Hz, 1H), 5.26 (s, 1H), 4.85 (d, J = 7.4 Hz, 1H), 4.03 - 3.79 (m, 8H), 3.68 (s, 2H), 3.41 - 3.24 (m, 4H), 2.16 (dd, J = 13.9, 6.0 Hz, 1H), 2.08 - 1.93 (m, 1H), 1.86 (d, J = 3.5 Hz, 6H), 1.76 - 1.54 (m, 2H)
8		416.42	(CDCl ₃) δ 8.65 (d, J = 2.0 Hz, 1H), 8.36 (d, J = 2.0 Hz, 1H), 6.61 (d, J = 2.4 Hz, 1H), 6.35 (d, J = 2.4 Hz, 1H), 6.10 (d, J = 7.6 Hz, 1H), 4.88 (d, J = 6.3 Hz, 1H), 4.21 (d, J = 4.1 Hz, 2H), 3.95 - 3.88 (m, 4H), 3.87 - 3.59 (m, 4H), 3.40 - 3.26 (m, 4H), 1.95 - 1.52 (m, 9H)
9		424.42	(CDCl ₃) δ 8.65 (d, J = 1.9 Hz, 1H), 8.37 (d, J = 2.0 Hz, 1H), 6.61 (d, J = 2.3 Hz, 1H), 6.34 (d, J = 2.3 Hz, 1H), 6.09 (d, J = 7.4 Hz, 1H), 4.81 (s, 1H), 4.66 (s, 2H), 4.01 - 3.86 (m, 4H), 3.65 (s, 2H), 3.42 - 3.24 (m, 4H), 1.87 (t, J = 2.3 Hz, 5H), 1.64 (d, J = 26.9 Hz, 6H)
10		414.44	(CDCl ₃) δ 8.65 (d, J = 1.9 Hz, 1H), 8.36 (d, J = 1.9 Hz, 1H), 6.61 (d, J = 2.3 Hz, 1H), 6.35 (d, J = 2.3 Hz, 1H), 6.10 (d, J = 7.7 Hz, 1H), 4.72 (s, 1H), 4.02 (t, J = 6.6 Hz, 2H), 3.95 - 3.82 (m, 4H), 3.68 (s, 2H), 3.43 - 3.26 (m, 4H), 1.87 (d, J = 3.7 Hz, 6H), 1.78 - 1.50 (m, 4H), 0.94 (t, J = 7.4 Hz, 3H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
11		414.44	(CDCl ₃) δ 8.65 (d, J = 2.0 Hz, 1H), 8.35 (d, J = 2.0 Hz, 1H), 6.60 (d, J = 2.4 Hz, 1H), 6.35 (d, J = 2.4 Hz, 1H), 6.10 (d, J = 7.7 Hz, 1H), 4.91 (dt, J = 12.5, 6.2 Hz, 1H), 4.69 (s, 1H), 4.01 - 3.81 (m, 4H), 3.68 (s, 2H), 3.46 - 3.24 (m, 4H), 1.93 - 1.76 (m, 6H), 1.78 - 1.56 (m, 2H), 1.25 (t, J = 9.6 Hz, 6H)
12		428.2	(CDCl ₃) δ 8.65 (d, J = 1.9 Hz, 1H), 8.35 (d, J = 1.9 Hz, 1H), 6.61 (d, J = 2.3 Hz, 1H), 6.35 (d, J = 2.3 Hz, 1H), 6.11 (d, J = 7.6 Hz, 1H), 4.74 (s, 1H), 3.99 - 3.79 (m, 6H), 3.68 (s, 2H), 3.40 - 3.24 (m, 4H), 1.87 (d, J = 3.5 Hz, 7H), 1.78 - 1.52 (m, 2H), 0.93 (d, J = 6.7 Hz, 6H)
13		418.44	(CDCl ₃) δ 8.65 (t, J = 1.6 Hz, 1H), 8.36 (t, J = 1.6 Hz, 1H), 6.61 (d, J = 2.3 Hz, 1H), 6.35 (d, J = 2.0 Hz, 1H), 6.11 (d, J = 7.6 Hz, 1H), 4.91 (d, J = 7.2 Hz, 1H), 4.80 - 4.18 (m, 4H), 4.00 - 3.84 (m, 4H), 3.81 - 3.56 (m, 2H), 3.46 - 3.21 (m, 4H), 1.87 (d, J = 3.5 Hz, 6H), 1.71 (dd, J = 16.0, 8.2 Hz, 2H)
14		410.44	(CDCl ₃) δ 8.65 (d, J = 2.0 Hz, 1H), 8.36 (d, J = 2.0 Hz, 1H), 6.61 (d, J = 2.4 Hz, 1H), 6.34 (d, J = 2.3 Hz, 1H), 6.10 (d, J = 7.1 Hz, 1H), 4.83 (s, 1H), 4.69 (s, 2H), 3.96 - 3.83 (m, 4H), 3.68 (s, 2H), 3.40 - 3.23 (m, 4H), 2.48 (t, J = 2.4 Hz, 1H), 1.87 (d, J = 4.2 Hz, 6H), 1.71 (dd, J = 15.7, 7.4 Hz, 2H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
15		478.44	(CDCl ₃) δ 8.66 (d, J = 2.0 Hz, 1H), 8.37 (d, J = 2.0 Hz, 1H), 7.10 - 6.98 (m, 2H), 6.87 (d, J = 9.0 Hz, 2H), 6.62 (d, J = 2.4 Hz, 1H), 6.36 (d, J = 2.4 Hz, 1H), 6.13 (d, J = 7.3 Hz, 1H), 5.06 (d, J = 7.9 Hz, 1H), 3.97 - 3.84 (m, 4H), 3.84 - 3.61 (m, 5H), 3.42 - 3.25 (m, 4H), 1.91 (d, J = 4.2 Hz, 6H), 1.85 - 1.68 (m, 2H)
16		453.96	(CDCl ₃) δ 8.49 (s, 1H), 8.23 (d, J = 1.5 Hz, 1H), 6.48 (s, 1H), 6.18 (d, J = 1.9 Hz, 1H), 6.06 (s, 1H), 4.52 (s, 1H), 4.47 (s, 2H), 3.60 (s, 2H), 3.45 (d, J = 11.6 Hz, 2H), 3.14 - 3.12 (m, 2H), 1.96 - 1.84 (m, 4H), 1.79 (s, 5H), 1.54 (s, 3H) and 1.38 (s, 9H) ppm
17		418.4	(CDCl ₃) δ 6.13 (d, J = 1.6 Hz, 1H), 5.89 (d, J = 1.4 Hz, 1H), 4.87 (d, J = 7.0 Hz, 1H), 4.59 (s, 1H), 3.26 (dd, J = 9.2, 4.3 Hz, 4H), 1.98 - 1.74 (m, 6H), 1.65 (dd, J = 15.9, 7.3 Hz, 3H), 1.47 (s, 9H)
18		406.48	(CDCl ₃) δ 8.65 (d, J = 2.0 Hz, 1H), 8.37 (d, J = 2.0 Hz, 1H), 8.27 (d, J = 4.8 Hz, 2H), 6.60 (d, J = 2.4 Hz, 1H), 6.51 (t, J = 4.8 Hz, 1H), 6.36 (d, J = 2.4 Hz, 1H), 6.15 (d, J = 7.8 Hz, 1H), 5.20 (d, J = 7.7 Hz, 1H), 4.04 (d, J = 7.9 Hz, 1H), 3.96 - 3.82 (m, 4H), 3.70 (s, 1H), 3.39 - 3.24 (m, 4H), 1.94 (dd, J = 13.7, 4.4 Hz, 6H), 1.78 (dt, J = 28.8, 16.1 Hz, 2H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
19		400.46	(CDCl ₃) δ 8.64 (d, J = 1.9 Hz, 1H), 8.35 (d, J = 1.9 Hz, 1H), 6.58 (t, J = 5.1 Hz, 2H), 6.34 (d, J = 2.3 Hz, 1H), 6.10 (d, J = 7.3 Hz, 1H), 4.03 (d, J = 8.3 Hz, 1H), 3.88 (t, J = 4.7 Hz, 6H), 3.68 (s, 1H), 3.42 (s, 3H), 3.37 - 3.23 (m, 4H), 1.98 - 1.78 (m, 6H), 1.69 (dd, J = 15.8, 7.5 Hz, 2H)
20		468.23	(400.0 MHz, CDCl ₃) δ 8.52 (d, J = 1.7 Hz, 1H), 8.23 (d, J = 1.6 Hz, 1H), 6.47 (d, J = 1.9 Hz, 1H), 6.19 (s, 1H), 6.03 (s, 1H), 5.19 (s, 1H), 4.76 (d, J = 7.6 Hz, 1H), 4.47 (s, 2H), 3.87 - 3.76 (m, 4H), 3.60 (s, 2H), 3.45 (d, J = 11.6 Hz, 2H), 3.13 - 3.10 (m, 2H), 2.61 (s, 1H), 2.13 - 2.04 (m, 1H), 1.95 - 1.85 (m, 5H), 1.79 (s, 5H) and 1.62 - 1.58 (m, 2H)
21		426.31	
22		425.35	(400.0 MHz, CDCl ₃) δ 8.54 (s, 1H), 8.22 (s, 1H), 6.43 (s, 1H), 6.19 (s, 1H), 6.02 (s, 1H), 4.47 (s, 2H), 4.38 (d, J = 5.2 Hz, 1H), 4.28 (s, 1H), 3.74 (s, 1H), 3.60 (s, 1H), 3.42 (s, 4H), 3.14 - 3.09 (m, 4H), 2.05 - 1.87 (m, 3H), 1.79 (s, 3H), 1.55 (d, J = 7.1 Hz, 2H) and 1.21 - 1.05 (m, 5H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
23		396.2	(CDCl ₃) δ 8.20 (d, J = 4.9 Hz, 2H), 6.46 (t, J = 4.8 Hz, 1H), 6.05 (d, J = 1.6 Hz, 1H), 5.82 (s, 1H), 5.24 (s, 1H), 4.82 (d, J = 7.0 Hz, 1H), 3.98 (s, 1H), 3.85 - 3.72 (m, 4H), 3.60 (s, 1H), 3.23 - 3.06 (m, 4H), 1.95 - 1.62 (m, 8H). [2]
24		405.59	(400 MHz, CDCl ₃) δ 8.65 (s, 1H), 8.37 (d, J = 1.8 Hz, 1H), 8.08 (d, J = 4.9 Hz, 1H), 7.41 (t, J = 7.7 Hz, 1H), 6.61 (s, 1H), 6.58 - 6.53 (m, 1H), 6.46 - 6.30 (m, 2H), 6.15 (d, J = 7.3 Hz, 1H), 4.56 (s, 1H), 3.98 - 3.78 (m, 5H), 3.76 - 3.61 (m, 1H), 3.42 - 3.24 (m, 4H), 1.97 (d, J = 29.6 Hz, 6H), 1.86 - 1.66 (m, 2H)
25		467.57	(CDCl ₃) δ 8.69 (d, J = 2.0 Hz, 1H), 8.42 (d, J = 2.0 Hz, 1H), 8.26 (s, 1H), 7.14 (d, J = 2.1 Hz, 1H), 6.64 (d, J = 2.4 Hz, 1H), 6.38 (d, J = 2.3 Hz, 1H), 6.24 (d, J = 6.8 Hz, 1H), 4.31 (q, J = 7.1 Hz, 2H), 3.97 - 3.81 (m, 5H), 3.41 - 3.24 (m, 4H), 2.22 (d, J = 7.5 Hz, 2H), 1.90 (dd, J = 22.4, 10.3 Hz, 6H), 1.35 (t, J = 7.1 Hz, 3H)
26		429.62	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 6.95 (d, J = 2.5 Hz, 1H), 6.88 (d, J = 2.5 Hz, 1H), 4.82 ? 4.60 (m, 2H), 4.00 ? 3.88 (m, 4H), 3.69 (d, J = 6.8 Hz, 1H), 3.42 ? 3.28 (m, 4H), 2.15 (p, J = 6.7, 5.6 Hz, 2H), 1.94 ? 1.75 (m, 6H), 1.46 (s, 9H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
27		406.58	(CDCl ₃) δ 8.65 (d, J = 2.0 Hz, 1H), 8.36 (d, J = 2.0 Hz, 1H), 7.98 (dd, J = 2.8, 1.5 Hz, 1H), 7.88 (d, J = 1.5 Hz, 1H), 7.79 (d, J = 2.8 Hz, 1H), 6.62 (d, J = 2.4 Hz, 1H), 6.37 (d, J = 2.4 Hz, 1H), 6.17 (s, 1H), 8.69 - 8.61 (m, 1H), 4.60 (d, J = 7.7 Hz, 1H), 4.08 - 3.83 (m, 5H), 8.71 - 8.57 (m, 1H), 3.73 (t, J = 6.9 Hz, 1H), 3.40 - 3.25 (m, 4H), 1.96 (h, J = 4.9 Hz, 6H), 1.77 (q, J = 7.4, 6.1 Hz, 2H)
28		407.3	(400 MHz, CDCl ₃) δ 8.77 - 8.59 (m, 2H), 8.29 (d, J = 4.9 Hz, 2H), 7.01 - 6.87 (m, 2H), 6.61 - 6.48 (m, 1H), 4.82 (s, 1H), 4.05 (s, 1H), 3.93 (t, J = 4.8 Hz, 4H), 3.35 (t, J = 4.8 Hz, 4H), 2.23 (d, J = 13.1 Hz, 2H), 2.05 - 1.82 (m, 6H)
29		450.61	(DMSO-d ₆) δ 12.65 (s, 1H), 8.69 (d, J = 2.0 Hz, 2H), 8.43 (d, J = 2.0 Hz, 1H), 8.00 (d, J = 7.9 Hz, 1H), 6.54 (d, J = 2.5 Hz, 1H), 6.48 (d, J = 2.3 Hz, 1H), 6.17 (d, J = 8.2 Hz, 1H), 4.06 (s, 1H), 3.77 (dd, J = 5.9, 3.8 Hz, 4H), 3.29 (s, 5H), 2.04 - 1.46 (m, 8H)
30		406.52	(CDCl ₃) δ 8.66 (d, J = 2.0 Hz, 1H), 8.58 (s, 1H), 8.37 (d, J = 2.0 Hz, 1H), 8.13 (s, 2H), 6.63 (d, J = 2.4 Hz, 1H), 6.37 (d, J = 2.4 Hz, 1H), 6.14 (d, J = 7.6 Hz, 1H), 3.96 - 3.86 (m, 4H), 3.76 (d, J = 7.7 Hz, 2H), 3.54 (d, J = 8.3 Hz, 1H), 3.39 - 3.29 (m, 4H), 2.02 - 1.86 (m, 6H), 1.76 (q, J = 8.9, 8.3 Hz, 2H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
31		422.49	(CDCl ₃) δ 12.88 - 12.45 (m, 1H), 8.66 (d, J = 2.0 Hz, 1H), 8.37 (d, J = 2.0 Hz, 1H), 7.81 (s, 1H), 6.62 (d, J = 2.3 Hz, 1H), 6.36 (d, J = 2.4 Hz, 1H), 6.12 (d, J = 7.7 Hz, 1H), 5.27 (s, 1H), 5.05 (d, J = 7.6 Hz, 1H), 4.02 - 3.83 (m, 4H), 3.63 (d, J = 47.6 Hz, 2H), 3.44 - 3.22 (m, 4H), 1.91 (q, J = 4.8, 4.3 Hz, 6H), 1.82-1.69 (m, 2H)
32		464.6	(CDCl ₃) δ 8.78 (d, J = 1.3 Hz, 1H), 8.66 (d, J = 2.0 Hz, 1H), 8.37 (d, J = 2.0 Hz, 1H), 7.91 (d, J = 1.3 Hz, 1H), 6.62 (d, J = 2.4 Hz, 1H), 6.37 (d, J = 2.4 Hz, 1H), 6.16 (d, J = 7.6 Hz, 1H), 5.13 (d, J = 7.6 Hz, 1H), 4.10 (s, 1H), 3.99 - 3.86 (m, 7H), 3.76 (s, 1H), 3.39 - 3.28 (m, 4H), 1.98 (h, J = 4.8 Hz, 6H), 1.80 (t, J = 8.7 Hz, 2H)
33		407.3	(CDCl ₃) δ 8.63 (d, J = 1.9 Hz, 1H), 7.99 (dd, J = 2.8, 1.5 Hz, 1H), 7.94 - 7.85 (m, 1H), 7.79 (d, J = 2.8 Hz, 1H), 7.14 (d, J = 1.0 Hz, 1H), 7.03 - 6.87 (m, 2H), 4.82 (d, J = 5.7 Hz, 1H), 4.70 (d, J = 8.0 Hz, 1H), 4.03 - 3.86 (m, 4H), 3.51 (s, 1H), 3.43 - 3.30 (m, 4H), 2.35 - 1.81 (m, 8H)
34		408.5	(CDCl ₃) δ 8.69 (dd, J = 3.4, 1.9 Hz, 1H), 8.62 (dd, J = 3.6, 1.9 Hz, 1H), 8.51 (dd, J = 4.8, 2.2 Hz, 2H), 7.01 - 6.83 (m, 3H), 5.18 (tt, J = 7.0, 3.4 Hz, 1H), 4.79 (tt, J = 6.9, 3.1 Hz, 1H), 4.00 - 3.85 (m, 4H), 3.34 (dq, J = 4.8, 2.6 Hz, 4H), 2.44 - 2.16 (m, 4H), 1.92 (tdd, J = 16.4, 7.7, 2.8 Hz, 4H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
35		372.23	(CDCl ₃) δ 8.68 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 6.99 - 6.85 (m, 2H), 4.70 (dq, J = 7.3, 3.5 Hz, 1H), 4.05 - 3.84 (m, 8H), 3.40 - 3.25 (m, 4H), 2.19 - 1.93 (m, 6H), 1.77 - 1.64 (m, 2H)
36		404.2	(CDCl ₃) δ 8.94 - 8.76 (m, 2H), 8.29 (d, J = 4.8 Hz, 2H), 7.67 (d, J = 1.7 Hz, 1H), 6.53 (t, J = 4.8 Hz, 1H), 6.37 (tt, J = 3.1, 1.5 Hz, 1H), 5.30 (d, J = 7.9 Hz, 1H), 4.87 (dt, J = 7.5, 3.6 Hz, 1H), 4.43 (q, J = 2.8 Hz, 2H), 4.02 (t, J = 5.5 Hz, 3H), 2.68 (dq, J = 6.0, 3.4, 3.0, 1.8 Hz, 2H), 2.35 - 2.11 (m, 2H), 2.07 - 1.84 (m, 6H)
37		425.25	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.18 (dd, J = 3.7, 0.8 Hz, 2H), 7.01 - 6.85 (m, 2H), 5.37 - 5.20 (m, 1H), 4.79 (d, J = 5.5 Hz, 1H), 4.02 - 3.85 (m, 4H), 3.43 - 3.29 (m, 4H), 2.31 - 2.15 (m, 2H), 2.02 - 1.85 (m, 6H)
38		407.25	(CDCl ₃) δ 8.61 (d, J = 2.0 Hz, 1H), 8.55 (d, J = 1.9 Hz, 1H), 8.20 (d, J = 4.8 Hz, 2H), 6.87 (d, J = 2.5 Hz, 1H), 6.81 (d, J = 2.5 Hz, 1H), 6.46 (t, J = 4.8 Hz, 1H), 4.95 (s, 1H), 4.45 (tt, J = 10.7, 3.6 Hz, 1H), 3.95 - 3.77 (m, 5H), 3.32 - 3.19 (m, 4H), 2.34 - 2.10 (m, 4H), 1.82 (dt, J = 12.9, 10.0 Hz, 2H), 1.45 - 1.20 (m, 2H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
39		441.28	(CDCl ₃) δ 8.72 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.37 (s, 1H), 6.98 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 6.36 (d, J = 1.0 Hz, 1H), 4.91 - 4.76 (m, 1H), 4.00 - 3.88 (m, 4H), 3.45 - 3.24 (m, 4H), 2.34 - 2.17 (m, 2H), 2.03 - 1.84 (m, 6H)
40		441.3	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.15 (d, J = 9.3 Hz, 1H), 7.00 - 6.87 (m, 2H), 6.64 (d, J = 9.3 Hz, 1H), 4.89 - 4.76 (m, 2H), 4.11 - 4.03 (m, 1H), 4.00 - 3.83 (m, 4H), 3.40 - 3.24 (m, 4H), 2.23 (dq, J = 12.9, 6.3, 5.6 Hz, 2H), 2.02 - 1.79 (m, 6H)
41		421.43	(CDCl ₃) δ 8.63 (d, J = 1.9 Hz, 1H), 8.55 (d, J = 1.9 Hz, 1H), 7.58 (d, J = 23.6 Hz, 2H), 6.89 (d, J = 2.4 Hz, 1H), 6.84 (d, J = 2.5 Hz, 1H), 4.73 (s, 2H), 3.93 - 3.72 (m, 5H), 3.34 - 3.18 (m, 4H), 2.29 (s, 3H), 2.15 (m, 2H), 1.84 (m, 6H)
42		408.56	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.52 (d, J = 4.8 Hz, 2H), 7.01 - 6.87 (m, 3H), 5.17 (ddt, J = 8.7, 6.7, 3.4 Hz, 1H), 4.76 - 4.58 (m, 1H), 4.00 - 3.87 (m, 4H), 3.40 - 3.27 (m, 4H), 2.43 - 2.22 (m, 4H), 2.05 - 1.87 (m, 2H), 1.86 - 1.71 (m, 2H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
43		412.48	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.36 (s, 1H), 7.12 (d, J = 3.6 Hz, 1H), 6.95 (d, J = 2.5 Hz, 1H), 6.90 (d, J = 2.5 Hz, 1H), 6.48 (d, J = 3.6 Hz, 1H), 5.18 (d, J = 8.0 Hz, 1H), 4.79 (td, J = 5.4, 2.7 Hz, 1H), 3.97 - 3.85 (m, 4H), 3.70 (q, J = 6.8 Hz, 1H), 3.39 - 3.25 (m, 4H), 2.29 - 2.12 (m, 2H), 2.07 - 1.77 (m, 6H)
44		432.6	(CDCl ₃) δ 8.71 (d, J = 2.0 Hz, 1H), 8.64 (d, J = 2.0 Hz, 1H), 8.54 (d, J = 2.9 Hz, 1H), 8.47 (d, J = 3.0 Hz, 1H), 6.99 (d, J = 2.4 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 5.81 (d, J = 8.3 Hz, 1H), 4.84 (dt, J = 5.3, 2.8 Hz, 1H), 4.13 - 4.05 (m, 1H), 4.00 - 3.84 (m, 4H), 3.43 - 3.30 (m, 4H), 2.32 - 2.17 (m, 2H), 2.02 - 1.85 (m, 6H)
45		485.26	(CDCl ₃) δ 8.70 (d, J = 2.0 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.29 (s, 2H), 6.98 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 5.29 (d, J = 8.3 Hz, 1H), 4.81 (s, 1H), 4.04 - 3.84 (m, 4H), 3.42 - 3.31 (m, 4H), 2.22 (s, 2H), 1.92 (d, J = 4.9 Hz, 6H)
46		407.57	(CDCl ₃) δ 8.63 (d, J = 2.0 Hz, 1H), 8.52 (d, J = 4.8 Hz, 2H), 8.37 (d, J = 2.0 Hz, 1H), 6.92 (t, J = 4.8 Hz, 1H), 6.62 (d, J = 2.4 Hz, 1H), 6.38 (d, J = 2.4 Hz, 1H), 6.16 (s, 1H), 5.27 (s, 1H), 4.06 - 3.78 (m, 4H), 3.64 (s, 1H), 3.48 - 3.20 (m, 4H), 2.14 (s, 2H), 2.04 - 1.80 (m, 4H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
47		372.16	(CDCl ₃) δ 8.66 (dd, J = 20.5, 1.9 Hz, 2H), 6.93 (dd, J = 17.3, 2.5 Hz, 2H), 4.87 - 4.65 (m, 1H), 4.04 - 3.83 (m, 4H), 3.72 (s, 3H), 3.46 - 3.22 (m, 4H), 2.72 - 2.40 (m, 1H), 2.35 - 1.99 (m, 4H), 1.99 - 1.51 (m, 4H)
48		358.64	(DMSO-d ₆) δ 12.13 (s, 1H), 8.72 (d, J = 1.7 Hz, 1H), 8.58 (d, J = 1.8 Hz, 1H), 7.13 (d, J = 2.1 Hz, 1H), 6.83 (d, J = 2.0 Hz, 1H), 4.94 - 4.84 (m, 1H), 3.91 - 3.68 (m, 4H), 3.51 - 3.19 (m, 4H), 2.47 - 2.33 (m, 1H), 2.04 - 1.82 (m, 4H), 1.82 - 1.60 (m, 4H)
49		425.39	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 7.99 (s, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.3 Hz, 1H), 6.20 (s, 1H), 5.19 (bs, 1H), 4.81 (bs, 1H), 3.96 - 3.84 (m, 4H), 3.40 - 3.27 (m, 4H), 2.29 - 2.14 (m, 2H), 1.99 - 1.81 (m, 6H)
50		425.39	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 8.30 (d, J = 2.1 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 5.84 (s, 1H), 5.42 (s, 1H), 4.81 (s, 1H), 3.99 - 3.82 (m, 4H), 3.39 - 3.24 (m, 4H), 2.31 - 2.19 (m, 2H), 2.08 - 1.72 (m, 8H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
51		425.33	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 8.30 (d, J = 2.1 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 5.84 (s, 1H), 5.42 (s, 1H), 4.81 (s, 1H), 3.99 - 3.82 (m, 4H), 3.39 - 3.24 (m, 4H), 2.31 - 2.19 (m, 2H), 2.08 - 1.72 (m, 8H)
52		435.19	(400 MHz, CDCl ₃) δ 8.68 (d, J = 1.4 Hz, 1H), 8.60 (d, J = 1.5 Hz, 1H), 6.93 (s, 1H), 6.90 (s, 1H), 6.28 (s, 1H), 5.06 (d, J = 7.9 Hz, 1H), 4.77 (s, 1H), 4.06 (bs, 1H), 3.97 - 3.84 (m, 4H), 3.38 - 3.25 (m, 4H), 2.27 (s, 6H), 2.18 - 2.09 (m, 2H), 1.94 - 1.83 (m, 7H)
53		433.25	(CDCl ₃) δ 8.9 Hz, 1H), 8.28 (d, J = 4.8 Hz, 2H), 6.85 (t, J = 1.9 Hz, 2H), 6.51 (t, J = 4.8 Hz, 1H), 5.29 (d, J = 6.3 Hz, 1H), 4.80 (dq, J = 5.5, 2.8 Hz, 1H), 4.57 (d, J = 3.9 Hz, 2H), 4.02 (t, J = 6.2 Hz, 1H), 3.54 - 3.43 (m, 2H), 3.19 (dd, J = 11.6, 2.6 Hz, 2H), 2.27 - 2.14 (m, 2H), 2.08 - 1.85 (m, 10H)
54		437.27	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.45 (d, J = 7.8 Hz, 2H), 7.02 - 6.81 (m, 2H), 4.78 (ddd, J = 7.3, 5.6, 3.1 Hz, 1H), 4.66 (d, J = 7.9 Hz, 1H), 4.01 - 3.78 (m, 7H), 3.41 - 3.25 (m, 4H), 2.30 - 2.08 (m, 2H), 1.94 (h, J = 8.8, 8.2 Hz, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
55		432.3	(CDCl ₃) δ 8.73 (d, J = 2.0 Hz, 1H), 8.63 (d, J = 2.0 Hz, 1H), 7.42 (d, J = 9.3 Hz, 1H), 6.96 (dd, J = 19.6, 2.5 Hz, 2H), 6.65 (d, J = 9.3 Hz, 1H), 5.40 (s, 1H), 4.86 (s, 1H), 3.94 (dd, J = 5.9, 3.8 Hz, 4H), 3.37 (dd, J = 6.0, 3.7 Hz, 4H), 2.27 (d, J = 12.7 Hz, 2H), 2.05 - 1.80 (m, 6H)
56		421.47	(CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.28 (d, J = 4.8 Hz, 2H), 6.98 - 6.86 (m, 2H), 6.53 (t, J = 4.8 Hz, 1H), 5.39 (s, 1H), 4.89 - 4.75 (m, 1H), 4.18 - 3.93 (m, 2H), 3.93 - 3.72 (m, 2H), 3.70 - 3.52 (m, 2H), 3.00 (td, J = 12.0, 3.5 Hz, 1H), 2.66 (dd, J = 12.1, 10.3 Hz, 1H), 2.21 (d, J = 8.9 Hz, 2H), 1.93 (d, J = 6.7 Hz, 6H), 1.30 (d, J = 6.2 Hz, 3H)
57		437.49	(CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.28 (d, J = 4.8 Hz, 2H), 7.02 - 6.86 (m, 2H), 6.53 (t, J = 4.8 Hz, 1H), 5.40 (s, 1H), 4.82 (d, J = 5.8 Hz, 1H), 4.20 - 3.95 (m, 2H), 3.94 - 3.53 (m, 6H), 3.03 (td, J = 12.0, 3.5 Hz, 1H), 2.86 (dd, J = 12.1, 10.4 Hz, 1H), 2.20 (d, J = 9.2 Hz, 2H), 2.11 - 1.80 (m, 7H)
58		467.16	(400 MHz, methanol-d ₄) δ 8.69 (d, J = 2.0 Hz, 1H), 8.56 (d, J = 2.0 Hz, 1H), 7.12 (d, J = 2.3 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 5.36 (s, 1H), 3.97 (s, 1H), 3.91 - 3.86 (m, 4H), 3.84 (s, 6H), 3.43 - 3.35 (m, 4H), 2.23 - 2.10 (m, 2H), 2.00 - 1.81 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
59		431.19	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.36 (d, J = 1.8 Hz, 1H), 7.54 (dd, J = 8.8, 2.2 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 6.37 (d, J = 8.7 Hz, 1H), 5.11 (s, 1H), 4.80 (s, 1H), 3.94 (s, 1H), 3.94 - 3.79 (m, 4H), 3.38 - 3.25 (m, 4H), 2.32 - 2.12 (m, 2H), 2.02 - 1.78 (m, 6H)
60		425.23	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.06 - 6.88 (m, 3H), 6.80 (dd, J = 9.4, 6.3 Hz, 1H), 4.97 - 4.71 (m, 2H), 4.14 (q, J = 7.1 Hz, 1H), 4.01 - 3.85 (m, 4H), 3.44 - 3.24 (m, 4H), 2.23 (d, J = 10.7 Hz, 2H), 1.96 (dt, J = 11.0, 7.6 Hz, 6H)
61		397.15	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.8 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 6.94 (d, J = 2.3 Hz, 1H), 6.87 (s, 1H), 5.78 - 5.64 (m, 1H), 4.73 (s, 1H), 4.01 (s, 1H), 3.97 - 3.78 (m, 4H), 3.43 - 3.18 (m, 4H), 2.26 - 2.05 (m, 2H), 1.98 - 1.73 (m, 6H), 1.36 - 1.26 (m, 1H), 1.01 - 0.92 (m, 2H), 0.78 - 0.67 (m, 2H)
62		427.23	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 6.95 (d, J = 2.3 Hz, 1H), 6.87 (d, J = 2.2 Hz, 1H), 6.79 (d, J = 8.1 Hz, 1H), 4.73 (s, 1H), 4.34 (dd, J = 8.3, 5.9 Hz, 1H), 4.04 - 3.84 (m, 7H), 3.40 - 3.28 (m, 4H), 2.35 - 2.24 (m, 1H), 2.23 - 2.11 (m, 2H), 2.09 - 1.98 (m, 1H), 1.97 - 1.76 (m, 8H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
63		463.54	(CDCl ₃) δ 8.53 (s, 1H), 8.29 (d, J = 4.8 Hz, 2H), 6.92 (d, J = 2.5 Hz, 1H), 6.85 (d, J = 2.5 Hz, 1H), 6.52 (t, J = 4.8 Hz, 1H), 5.25 (d, J = 8.3 Hz, 1H), 4.79 (s, 1H), 4.07 (d, J = 20.4 Hz, 1H), 3.98 - 3.85 (m, 4H), 3.43 - 3.21 (m, 4H), 3.05 - 2.83 (m, 2H), 2.30 - 2.14 (m, 1H), 2.03 - 1.71 (m, 7H), 1.45 (dq, J = 14.5, 7.3 Hz, 2H), 0.98 (t, J = 7.3 Hz, 3H)
64		541.26	(CDCl ₃) δ 8.55 (s, 1H), 8.29 (d, J = 1.5 Hz, 2H), 7.05 - 6.93 (m, 1H), 6.85 (d, J = 2.5 Hz, 1H), 5.49 (d, J = 8.0 Hz, 1H), 4.80 (d, J = 5.8 Hz, 1H), 4.03 - 3.81 (m, 5H), 3.45 - 3.27 (m, 4H), 2.98 (dd, J = 8.5, 7.0 Hz, 2H), 2.32 - 2.09 (m, 2H), 2.00 - 1.71 (m, 8H), 1.56 - 1.34 (m, 2H), 0.98 (td, J = 7.3, 3.3 Hz, 3H)
65		450.49	(CDCl ₃) δ 8.76 - 8.66 (m, 3H), 8.61 (d, J = 1.9 Hz, 1H), 6.99 - 6.87 (m, 2H), 5.71 (d, J = 8.1 Hz, 1H), 4.81 (s, 1H), 4.11 (s, 1H), 3.92 (t, J = 4.9 Hz, 4H), 3.34 (t, J = 4.9 Hz, 4H), 2.22 (d, J = 10.2 Hz, 2H), 2.02 - 1.85 (m, 6H)
66		495.23	(CDCl ₃) δ 8.77 (d, J = 2.3 Hz, 1H), 8.72 (d, J = 2.3 Hz, 1H), 7.12 (d, J = 2.4 Hz, 1H), 7.02 (d, J = 2.4 Hz, 1H), 4.92 (s, 1H), 4.18 - 4.08 (m, 2H), 3.93 (t, J = 4.9 Hz, 4H), 3.85 - 3.76 (m, 2H), 3.60 (q, J = 7.0 Hz, 2H), 3.47 (t, J = 4.9 Hz, 4H), 2.26 (d, J = 13.3 Hz, 2H), 2.07 (t, J = 10.5 Hz, 2H), 2.01 - 1.72 (m, 2H), 1.26 (t, J = 7.0 Hz, 4H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
67		450.3	(CDCl ₃) δ 8.71 (d, J = 2.0 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.02 (s, 1H), 6.94 (dd, J = 12.2, 2.5 Hz, 2H), 5.32 (s, 2H), 4.87 (d, J = 8.1 Hz, 1H), 4.79 (s, 1H), 4.01 - 3.86 (m, 4H), 3.36 (q, J = 5.4, 4.7 Hz, 4H), 2.83 (s, 6H), 2.19 (s, 2H), 1.92 (d, J = 4.8 Hz, 6H)
68		489.24	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.34 (s, 2H), 7.02 - 6.86 (m, 2H), 6.00 (tt, J = 3.0, 1.5 Hz, 1H), 5.41 - 5.21 (m, 1H), 4.81 (dt, J = 7.2, 3.6 Hz, 1H), 4.31 (q, J = 2.8 Hz, 2H), 4.03 (dd, J = 7.8, 4.4 Hz, 1H), 4.00 - 3.82 (m, 6H), 3.42 - 3.25 (m, 4H), 2.44 (tdd, J = 5.7, 2.9, 1.7 Hz, 2H), 2.22 (dq, J = 11.2, 6.6, 6.0 Hz, 2H), 2.02 - 1.83 (m, 6H)
69		451.21	(CDCl ₃) δ 9.05 (s, 1H), 8.88 - 8.71 (m, 3H), 8.49 (s, 1H), 7.06 (d, J = 2.3 Hz, 1H), 6.96 (d, J = 2.3 Hz, 1H), 4.85 (s, 1H), 4.17 (s, 1H), 3.93 (t, J = 4.8 Hz, 4H), 3.42 (t, J = 4.9 Hz, 4H), 2.25 - 2.10 (m, 2H), 1.95 (d, J = 11.9 Hz, 4H)
70		409.45	(methanol-d ₄) δ 8.69 (s, 1H), 8.56 (s, 1H), 7.20 (s, 1H), 7.10 (d, J = 2.2 Hz, 1H), 6.88 (d, J = 2.4 Hz, 1H), 4.93 (d, J = 14.3 Hz, 2H), 3.95 - 3.76 (m, 7H), 3.42 - 3.32 (m, 4H), 3.09 (d, J = 7.3 Hz, 1H), 2.21 - 2.09 (m, 2H), 1.85 (dd, J = 10.8, 5.6 Hz, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
71		464.4	(CDCl ₃) δ 8.79 - 8.64 (m, 3H), 8.59 (d, J = 1.9 Hz, 1H), 6.99 - 6.88 (m, 2H), 6.19 (q, J = 4.7 Hz, 1H), 5.90 (d, J = 8.2 Hz, 1H), 4.81 (dq, J = 5.3, 2.7 Hz, 1H), 4.08 (qd, J = 8.2, 6.5, 2.3 Hz, 1H), 3.97 - 3.87 (m, 4H), 3.39 - 3.29 (m, 4H), 2.93 (d, J = 4.8 Hz, 3H), 2.27 - 2.14 (m, 2H), 2.06 - 1.79 (m, 6H)
72		478.39	(CDCl ₃) δ 3.97 - 3.87 (m, 4H), 3.39 - 3.29 (m, 4H), 3.10 (s, 6H), 2.22 (dt, J = 11.3, 5.1 Hz, 2H), 1.94 (dd, J = 8.3, 3.9 Hz, 6H), 4.12 - 4.01 (m, 1H), 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.46 (s, 2H), 7.00 - 6.87 (m, 2H), 5.57 (d, J = 8.1 Hz, 1H), 4.81 (dq, J = 5.1, 2.4 Hz, 1H)
73		427.2	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.2 Hz, 1H), 8.57 (d, J = 21.7 Hz, 1H), 6.93 (d, J = 1.7 Hz, 1H), 6.86 (s, 1H), 5.75 (d, J = 7.4 Hz, 1H), 4.74 (s, 1H), 4.03 - 3.87 (m, 8H), 3.81 (dd, J = 15.2, 7.5 Hz, 1H), 3.37 - 3.25 (m, 4H), 2.96 - 2.76 (m, 1H), 2.23 - 2.07 (m, 4H), 1.84 - 1.78 (m, 6H)
74		411.25	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.86 (d, J = 2.4 Hz, 1H), 5.42 (d, J = 7.9 Hz, 1H), 4.72 (s, 1H), 3.98 (s, 1H), 3.95 - 3.86 (m, 4H), 3.40 - 3.26 (m, 4H), 3.05 - 2.87 (m, 1H), 2.32 - 2.21 (m, 2H), 2.21 - 2.07 (m, 4H), 2.03 - 1.90 (m, 1H), 1.90 - 1.77 (m, 7H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
75		562.34	(CDCl ₃) δ 8.72 - 8.51 (m, 2H), 7.01 (d, J = 2.5 Hz, 1H), 6.93 (d, J = 2.5 Hz, 1H), 4.84 (s, 1H), 4.40 (s, 2H), 3.99 (s, 1H), 3.83 (dd, J = 5.9, 3.8 Hz, 4H), 3.65 (s, 2H), 3.35 (t, J = 4.9 Hz, 4H), 2.87 (s, 2H), 2.27 - 1.91 (m, 4H), 1.90 - 1.59 (m, 2H), 1.41 (s, 9H)
76		462.23	(CDCl ₃) δ 8.70 (t, J = 2.2 Hz, 1H), 8.63 (dd, J = 2.7, 1.9 Hz, 1H), 7.97 (s, 1H), 7.00 - 6.86 (m, 2H), 5.05 (d, J = 8.2 Hz, 1H), 4.77 (s, 1H), 4.02 (s, 1H), 3.93 (t, J = 4.7 Hz, 4H), 3.86 (s, 1H), 3.35 (t, J = 4.8 Hz, 4H), 3.17 (t, J = 6.0 Hz, 1H), 2.89 (d, J = 5.5 Hz, 1H), 2.71 (t, J = 6.0 Hz, 1H), 2.21 (dd, J = 9.2, 4.9 Hz, 2H), 1.91 (d, J = 5.5 Hz, 3H), 1.75 (d, J = 5.6 Hz, 4H)
77		413.19	(400 MHz, CDCl ₃) δ 8.68 (d, J = 1.9 Hz, 1H), 8.63 - 8.54 (m, 1H), 6.93 (d, J = 2.4 Hz, 1H), 6.86 (d, J = 2.3 Hz, 1H), 5.76 - 5.55 (m, 1H), 4.93 - 4.67 (m, 4H), 3.99 (d, J = 26.0 Hz, 1H), 3.95 - 3.81 (m, 4H), 3.77 - 3.55 (m, 2H), 3.39 - 3.28 (m, 4H), 2.21 - 2.09 (m, 2H), 1.90 - 1.75 (m, 6H)
78		371.18	(400 MHz, CDCl ₃) δ 8.69 (s, 1H), 8.60 (s, 1H), 6.94 (s, 1H), 6.86 (s, 1H), 5.59 (s, 1H), 4.74 (s, 1H), 4.11 - 3.81 (m, 5H), 3.42 - 3.23 (m, 4H), 2.28 - 2.09 (m, 2H), 1.98 (s, 3H), 1.90 - 1.78 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
79		385.18	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 6.93 (d, J = 2.4 Hz, 1H), 6.86 (d, J = 2.3 Hz, 1H), 5.52 (d, J = 7.3 Hz, 1H), 4.73 (s, 1H), 3.99 (s, 1H), 3.96 - 3.87 (m, 4H), 3.38 - 3.30 (m, 4H), 2.24 - 2.08 (m, 4H), 1.88 - 1.79 (m, 6H), 1.16 (t, J = 7.6 Hz, 3H)
80		399.23	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 6.93 (d, J = 2.4 Hz, 1H), 6.86 (d, J = 2.3 Hz, 1H), 5.53 (d, J = 7.7 Hz, 1H), 4.73 (s, 1H), 4.00 (s, 1H), 3.95 - 3.84 (m, 4H), 3.39 - 3.25 (m, 4H), 2.22 - 2.06 (m, 4H), 1.88 - 1.76 (m, 6H), 1.70 - 1.61 (m, 2H), 0.94 (t, J = 7.4 Hz, 3H)
81		401.21	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.87 (d, J = 2.2 Hz, 1H), 6.65 (d, J = 8.5 Hz, 1H), 4.72 (s, 1H), 4.06 (s, 1H), 3.96 - 3.85 (m, 6H), 3.43 (s, 3H), 3.38 - 3.28 (m, 4H), 2.24 - 2.09 (m, 2H), 1.95 - 1.76 (m, 6H)
82		413.27	(400 MHz, CDCl ₃) δ 8.68 (d, J = 1.8 Hz, 1H), 8.59 (d, J = 1.8 Hz, 1H), 6.93 (d, J = 2.3 Hz, 1H), 6.86 (d, J = 2.0 Hz, 1H), 5.53 (d, J = 7.8 Hz, 1H), 4.72 (s, 1H), 4.01 (s, 1H), 3.96 - 3.81 (m, 4H), 3.37 - 3.24 (m, 4H), 2.24 - 2.06 (m, 3H), 2.01 (d, J = 7.0 Hz, 2H), 1.87 - 1.76 (m, 6H), 0.94 (d, J = 6.5 Hz, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
83		413.23	(400 MHz, CDCl ₃) δ 8.62 (d, J = 1.9 Hz, 1H), 8.55 (d, J = 1.9 Hz, 1H), 6.87 (d, J = 2.4 Hz, 1H), 6.80 (d, J = 2.3 Hz, 1H), 5.62 (d, J = 7.7 Hz, 1H), 4.64 (s, 1H), 3.99 - 3.76 (m, 5H), 3.37 - 3.14 (m, 4H), 2.21 - 1.95 (m, 2H), 1.87 - 1.63 (m, 6H), 1.13 (s, 9H)
84		431.19	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.19 (d, J = 4.8 Hz, 1H), 6.95 (d, J = 2.3 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 6.70 (dd, J = 5.1, 1.2 Hz, 1H), 6.56 (s, 1H), 4.87 (d, J = 7.6 Hz, 1H), 4.80 (s, 1H), 3.96 - 3.88 (m, 4H), 3.85 (s, 1H), 3.38 - 3.28 (m, 4H), 2.28 - 2.14 (m, 2H), 2.00 - 1.85 (m, 6H)
85		491.3	(CDCl ₃) δ 8.67 (d, J = 2.0 Hz, 1H), 8.62 (d, J = 2.0 Hz, 1H), 7.05 - 6.95 (m, 2H), 6.72 (s, 1H), 4.88 (s, 1H), 3.98 - 3.82 (m, 4H), 3.64 (t, J = 5.1 Hz, 2H), 3.41 - 3.31 (m, 4H), 3.24 (s, 3H), 3.09 (t, J = 5.2 Hz, 2H), 2.68 (s, 3H), 2.30 - 1.99 (m, 4H), 1.86 (d, J = 9.0 Hz, 4H)
86		433.2	(CDCl ₃) δ 9.68 (s, 1H), 8.76 - 8.58 (m, 2H), 7.00 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 6.41 (dd, J = 17.7, 11.1 Hz, 1H), 5.65 (d, J = 17.7 Hz, 1H), 5.33 (d, J = 11.1 Hz, 1H), 4.85 (q, J = 3.6 Hz, 1H), 4.01 (s, 1H), 3.94 - 3.70 (m, 4H), 3.47 - 3.19 (m, 4H), 2.27 - 1.94 (m, 4H), 1.93 - 1.63 (m, 4H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
87		431.2	(CDCl ₃) δ 8.63 (d, J = 1.9 Hz, 1H), 8.57 (d, J = 1.9 Hz, 1H), 8.20 (dd, J = 5.0, 1.9 Hz, 1H), 7.57 (dd, J = 7.6, 1.9 Hz, 1H), 6.89 (d, J = 2.5 Hz, 1H), 6.85 (d, J = 2.5 Hz, 1H), 6.51 (dd, J = 7.6, 4.9 Hz, 1H), 5.12 (d, J = 7.7 Hz, 1H), 4.83 - 4.71 (m, 0H), 4.24 - 4.03 (m, 1H), 3.92 - 3.77 (m, 4H), 3.35 - 3.19 (m, 4H), 2.28 - 2.10 (m, 2H), 1.88 (td, J = 8.3, 6.8, 3.9 Hz, 6H)
88		434.25	(400 MHz, CDCl ₃) δ 8.69 (s, 1H), 8.63 (s, 1H), 8.57 (s, 1H), 8.30 - 8.12 (m, 2H), 7.85 (t, J = 7.1 Hz, 1H), 7.50 - 7.37 (m, 1H), 6.94 (s, 1H), 6.89 (s, 1H), 4.89 - 4.63 (m, 1H), 4.35 - 4.13 (m, 1H), 4.01 - 3.78 (m, 4H), 3.43 - 3.19 (m, 4H), 2.37 - 2.15 (m, 2H), 2.12 - 1.82 (m, 6H)
89		434.22	(400 MHz, CDCl ₃) δ 8.99 (s, 1H), 8.73 (d, J = 3.4 Hz, 1H), 8.70 (d, J = 1.6 Hz, 1H), 8.60 (d, J = 1.7 Hz, 1H), 8.11 (d, J = 7.8 Hz, 1H), 7.40 (dd, J = 7.6, 4.8 Hz, 1H), 6.95 (d, J = 2.1 Hz, 1H), 6.90 (s, 1H), 6.30 (d, J = 7.6 Hz, 1H), 4.81 (s, 1H), 4.22 (s, 1H), 3.97 - 3.85 (m, 4H), 3.41 - 3.23 (m, 4H), 2.32 - 2.17 (m, 2H), 2.01 - 1.84 (m, 6H)
90		434.22	(400 MHz, CDCl ₃) δ 8.75 (dd, J = 4.4, 1.7 Hz, 2H), 8.70 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 7.62 (dd, J = 4.4, 1.7 Hz, 2H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.5 Hz, 1H), 6.33 (d, J = 8.2 Hz, 1H), 4.81 (s, 1H), 4.27 - 4.16 (m, 1H), 3.96 - 3.87 (m, 4H), 3.38 - 3.31 (m, 4H), 2.30 - 2.16 (m, 2H), 2.00 - 1.86 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
91		433.21	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.63 - 8.58 (m, 1H), 7.77 (dd, J = 5.2, 3.2 Hz, 2H), 7.54 - 7.40 (m, 3H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 6.23 (d, J = 8.0 Hz, 1H), 4.78 (s, 1H), 4.31 - 4.13 (m, 1H), 3.95 - 3.87 (m, 4H), 3.38 - 3.27 (m, 4H), 2.31 - 2.17 (m, 2H), 1.97 - 1.85 (m, 6H)
92		437.24	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.47 (d, J = 8.9 Hz, 1H), 7.00 (d, J = 1.0 Hz, 1H), 6.95 (d, J = 0.7 Hz, 1H), 6.94 (d, J = 2.5 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 4.81 (s, 1H), 4.07 (d, J = 12.3 Hz, 4H), 3.98 - 3.85 (m, 4H), 3.41 - 3.24 (m, 4H), 2.34 - 2.12 (m, 2H), 2.06 - 1.83 (m, 6H)
93		451.21	(methanol-d ₄) δ 8.69 (d, J = 2.2 Hz, 1H), 8.59 (d, J = 2.2 Hz, 1H), 8.53 (d, J = 4.9 Hz, 1H), 7.19 (d, J = 5.0 Hz, 1H), 7.15 (d, J = 2.3 Hz, 1H), 6.87 (d, J = 2.4 Hz, 1H), 4.95 (d, J = 7.7 Hz, 1H), 4.10 (s, 1H), 3.95 - 3.82 (m, 4H), 3.47 - 3.37 (m, 4H), 2.20 (d, J = 10.1 Hz, 2H), 2.04 - 1.81 (m, 6H)
94		437.44	(CDCl ₃) δ 8.68 (d, J = 2.0 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 8.29 (s, 2H), 6.98 - 6.87 (m, 2H), 5.36 (d, J = 8.1 Hz, 1H), 4.79 (q, J = 5.2, 4.0 Hz, 1H), 4.52 (s, 2H), 4.01 (dd, J = 8.1, 4.3 Hz, 1H), 3.95 - 3.84 (m, 4H), 3.39 - 3.28 (m, 4H), 2.25 - 2.12 (m, 2H), 1.99 - 1.82 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
95		424.21	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.7 Hz, 1H), 8.65 - 8.58 (m, 1H), 7.80 (s, 1H), 7.22 (d, J = 9.6 Hz, 1H), 7.15 (d, J = 7.6 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 4.80 (s, 1H), 4.24 - 4.09 (m, 1H), 3.98 - 3.85 (m, 4H), 3.40 - 3.29 (m, 4H), 2.29 - 2.15 (m, 2H), 2.02 - 1.86 (m, 6H)
96		435.19	(400 MHz, CDCl ₃) δ 8.89 (d, J = 4.9 Hz, 2H), 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.20 (d, J = 8.3 Hz, 1H), 7.45 (t, J = 4.9 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 4.75 (s, 1H), 4.39 - 4.21 (m, 1H), 4.00 - 3.84 (m, 4H), 3.41 - 3.21 (m, 4H), 2.29 - 2.13 (m, 2H), 2.12 - 1.87 (m, 6H)
97		435.19	(400 MHz, CDCl ₃) δ 9.25 (d, J = 1.4 Hz, 1H), 8.97 (d, J = 5.0 Hz, 1H), 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.22 - 8.04 (m, 2H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 4.81 - 4.72 (m, 1H), 4.26 - 4.15 (m, 1H), 3.96 - 3.86 (m, 4H), 3.38 - 3.29 (m, 4H), 2.28 - 2.17 (m, 2H), 2.06 - 1.86 (m, 6H)
98		435.19	(400 MHz, CDCl ₃) δ 9.33 (s, 1H), 9.12 (s, 2H), 8.70 (d, J = 1.9 Hz, 1H), 8.57 (d, J = 1.9 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 6.44 (d, J = 7.8 Hz, 1H), 4.83 (s, 1H), 4.23 (qd, J = 9.2, 4.7 Hz, 1H), 3.96 - 3.87 (m, 4H), 3.38 - 3.30 (m, 4H), 2.32 - 2.18 (m, 2H), 2.01 - 1.88 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
99		435.19	(400 MHz, CDCl ₃) δ 9.29 (dd, J = 5.0, 1.7 Hz, 1H), 8.70 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.33 (dd, J = 8.4, 1.7 Hz, 2H), 7.67 (dd, J = 8.4, 5.0 Hz, 1H), 6.95 (d, J = 2.5 Hz, 1H), 6.91 (d, J = 2.5 Hz, 1H), 4.85 (s, 1H), 4.29 - 4.13 (m, 1H), 3.98 - 3.85 (m, 4H), 3.40 - 3.26 (m, 4H), 2.32 - 2.16 (m, 2H), 2.12 - 1.98 (m, 2H), 1.99 - 1.87 (m, 4H)
100		423.13	(400 MHz, CDCl ₃) δ 11.83 (s, 1H), 8.71 (d, J = 1.9 Hz, 1H), 8.59 (d, J = 1.9 Hz, 1H), 7.55 (d, J = 2.4 Hz, 1H), 7.05 (d, J = 8.5 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.87 (dd, J = 4.7, 2.3 Hz, 1H), 6.84 (d, J = 2.4 Hz, 1H), 4.74 (s, 1H), 4.18 (s, 1H), 3.96 - 3.88 (m, 4H), 3.38 - 3.29 (m, 4H), 2.17 - 2.09 (m, 2H), 1.94 - 1.81 (m, 6H)
101		438.18	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.07 (s, 1H), 7.01 (d, J = 8.3 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.88 (d, J = 2.4 Hz, 1H), 4.76 (s, 1H), 4.23 - 4.09 (m, 1H), 3.98 - 3.84 (m, 4H), 3.40 - 3.26 (m, 4H), 2.48 (s, 3H), 2.28 - 2.12 (m, 2H), 1.99 - 1.82 (m, 6H)
102		482.36	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.59 (d, J = 2.0 Hz, 1H), 6.96 (d, J = 2.5 Hz, 1H), 6.89 (d, J = 2.5 Hz, 1H), 4.73 (d, J = 5.7 Hz, 1H), 4.18 (d, J = 7.7 Hz, 1H), 3.93 (dd, J = 5.9, 3.8 Hz, 4H), 3.69 - 3.55 (m, 1H), 3.46 (d, J = 16.5 Hz, 1H), 3.41 - 3.28 (m, 4H), 3.08 (d, J = 16.4 Hz, 1H), 2.17 (d, J = 11.6 Hz, 2H), 2.00 - 1.75 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
103		463.36	(CDCl ₃) δ 8.72 (d, J = 2.0 Hz, 1H), 8.64 (d, J = 2.0 Hz, 1H), 6.99 (d, J = 2.5 Hz, 1H), 6.93 (d, J = 2.5 Hz, 1H), 5.74 (s, 1H), 4.80 (s, 1H), 4.02 - 3.87 (m, 4H), 3.36 (t, J = 4.9 Hz, 4H), 2.19 (s, 2H), 2.08 - 1.79 (m, 6H)
104		443.38	(CDCl ₃) δ 8.73 (s, 1H), 8.68 (d, J = 2.4 Hz, 1H), 6.99 (d, J = 2.4 Hz, 1H), 6.85 (d, J = 2.4 Hz, 1H), 5.69 (d, J = 2.5 Hz, 1H), 4.76 (s, 1H), 3.84 (dd, J = 5.9, 3.9 Hz, 4H), 3.38 (dd, J = 6.0, 3.9 Hz, 4H), 2.15 (d, J = 11.1 Hz, 2H), 1.96 - 1.67 (m, 6H)
105		432.4	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.59 (d, J = 1.9 Hz, 1H), 8.30 (d, J = 1.3 Hz, 1H), 7.89 (d, J = 1.4 Hz, 1H), 6.93 (dd, J = 18.4, 2.5 Hz, 2H), 5.52 (d, J = 7.8 Hz, 1H), 4.83 (tt, J = 4.8, 2.7 Hz, 1H), 4.19 - 4.01 (m, 1H), 3.96 - 3.87 (m, 4H), 3.39 - 3.26 (m, 4H), 2.31 - 2.14 (m, 2H), 2.09 - 1.78 (m, 6H)
106		465.14	(400 MHz, CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 7.88 (dd, J = 7.4, 2.1 Hz, 1H), 7.02 (dd, J = 11.9, 8.3 Hz, 1H), 6.97 (d, J = 2.4 Hz, 1H), 6.92 (d, J = 2.4 Hz, 1H), 6.83 (dd, J = 12.4, 7.6 Hz, 1H), 4.78 (s, 1H), 4.25 (s, 1H), 4.03 - 3.85 (m, 4H), 3.44 - 3.23 (m, 4H), 2.38 (s, 3H), 2.30 - 2.13 (m, 2H), 2.06 - 1.88 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
107		485.12	(400 MHz, CDCl ₃) δ 8.73 - 8.66 (m, 1H), 8.65 - 8.59 (m, 1H), 8.05 (dd, J = 6.6, 2.8 Hz, 1H), 7.46 - 7.36 (m, 1H), 7.08 (dd, J = 11.1, 8.8 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 6.84 - 6.66 (m, 1H), 4.76 (bs, 1H), 4.22 (bs, 1H), 3.96 - 3.87 (m, 4H), 3.37 - 3.28 (m, 4H), 2.28 - 2.10 (m, 2H), 2.01 - 1.86 (m, 6H)
108		478.26	(CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 2.0 Hz, 1H), 8.49 (d, J = 1.4 Hz, 1H), 7.77 (d, J = 1.4 Hz, 1H), 6.93 (dd, J = 16.1, 2.5 Hz, 2H), 5.03 (d, J = 7.9 Hz, 1H), 4.81 (td, J = 5.3, 2.6 Hz, 1H), 4.09 - 3.98 (m, 1H), 3.98 - 3.87 (m, 4H), 3.39 - 3.26 (m, 4H), 3.19 (s, 3H), 3.12 (s, 3H), 2.22 (dt, J = 11.2, 4.9 Hz, 2H), 2.02 - 1.82 (m, 6H)
109		447.02	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.10 (d, J = 0.5 Hz, 2H), 6.97 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 5.13 (d, J = 8.3 Hz, 1H), 4.80 (s, 1H), 4.06 - 3.86 (m, 5H), 3.35 (dd, J = 5.9, 3.8 Hz, 4H), 2.27 - 2.14 (m, 2H), 1.92 (d, J = 5.1 Hz, 6H), 1.79 - 1.44 (m, 6H), 1.28 (t, J = 7.1 Hz, 1H), 1.03 - 0.83 (m, 2H), 0.68 - 0.51 (m, 2H)
110		395.19	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.64 - 8.54 (m, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.86 (d, J = 2.5 Hz, 1H), 6.06 - 5.78 (m, 1H), 4.79 - 4.68 (m, 1H), 4.11 - 3.96 (m, 1H), 3.96 - 3.86 (m, 4H), 3.40 - 3.27 (m, 4H), 2.13 (dd, J = 11.0, 5.2 Hz, 2H), 1.94 (s, 3H), 1.90 - 1.78 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
111		421.19	
112		452.18	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.66 - 8.53 (m, 3H), 7.93 (dd, J = 6.4, 5.0 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 6.78 (s, 1H), 4.78 (bs, 1H), 4.23 (bs, 1H), 3.97 - 3.85 (m, 4H), 3.38 - 3.27 (m, 4H), 2.30 - 2.15 (m, 2H), 2.02 - 1.86 (m, 6H)
113		439.24	(400 MHz, CDCl ₃) δ 8.70 (s, 1H), 8.60 (s, 1H), 6.94 (s, 1H), 6.87 (s, 1H), 5.99 (d, J = 6.8 Hz, 1H), 4.75 (bs, 1H), 4.03 (bs, 1H), 3.99 - 3.83 (m, 4H), 3.43 - 3.21 (m, 4H), 2.33 (s, 1H), 2.24 - 2.04 (m, 2H), 1.94 - 1.74 (m, 6H), 1.56 (s, 6H)
114		452.15	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.40 (dt, J = 4.3, 1.3 Hz, 1H), 8.01 (d, J = 8.5 Hz, 1H), 7.58 - 7.52 (m, 1H), 7.51 - 7.46 (m, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 4.74 (s, 1H), 4.29 - 4.17 (m, 1H), 3.97 - 3.88 (m, 4H), 3.39 - 3.30 (m, 4H), 2.25 - 2.16 (m, 2H), 2.06 - 1.88 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
115		451.16	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.8 Hz, 1H), 8.62 (d, J = 1.8 Hz, 1H), 8.09 (dd, J = 7.9, 6.1 Hz, 1H), 7.47 (d, J = 8.1 Hz, 1H), 7.13 (dd, J = 11.8, 7.9 Hz, 1H), 6.95 (d, J = 2.3 Hz, 1H), 6.90 (d, J = 2.3 Hz, 1H), 6.82 (s, 1H), 4.77 (s, 1H), 4.24 (s, 1H), 3.96 - 3.88 (m, 4H), 3.38 - 3.30 (m, 4H), 2.27 - 2.14 (m, 2H), 2.01 - 1.87 (m, 6H)
116		473.17	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.83 (d, J = 2.7 Hz, 1H), 7.19 - 6.93 (m, 4H), 6.89 (d, J = 2.4 Hz, 1H), 4.75 (s, 1H), 4.21 (s, 1H), 3.95 - 3.89 (m, 4H), 3.38 - 3.30 (m, 4H), 2.26 - 2.13 (m, 2H), 2.03 - 1.86 (m, 6H)
117		437.17	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.35 (d, J = 2.2 Hz, 1H), 6.99 (d, J = 8.2 Hz, 1H), 6.93 (d, J = 2.4 Hz, 1H), 6.88 (d, J = 2.4 Hz, 1H), 6.78 (d, J = 2.3 Hz, 1H), 4.73 (s, 1H), 4.26 - 4.14 (m, 1H), 3.96 - 3.87 (m, 7H), 3.36 - 3.29 (m, 4H), 2.23 - 2.12 (m, 2H), 2.01 - 1.82 (m, 6H)
118		437.21	(400 MHz, CDCl ₃) δ 10.00 (s, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 6.93 (s, 1H), 6.87 (d, J = 2.4 Hz, 1H), 6.81 (d, J = 2.4 Hz, 1H), 6.50 (s, 1H), 4.69 (s, 1H), 4.15 - 4.05 (m, 1H), 3.92 - 3.77 (m, 4H), 3.35 - 3.19 (m, 4H), 2.27 (d, J = 0.5 Hz, 3H), 2.18 - 2.04 (m, 2H), 1.95 - 1.74 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
119		451.25	(400 MHz, CDCl ₃) δ 8.70 (s, 1H), 8.63 (s, 1H), 6.99 - 6.50 (m, 3H), 4.76 (d, J = 20.7 Hz, 1H), 4.19 (s, 1H), 4.13 - 3.74 (m, 7H), 3.41 - 3.21 (m, 4H), 2.37 - 2.08 (m, 5H), 2.03 - 1.79 (m, 6H)
120		491.18	(400 MHz, CDCl ₃) δ 8.73 (d, J = 1.9 Hz, 1H), 8.58 (d, J = 1.9 Hz, 1H), 6.97 (d, J = 2.4 Hz, 1H), 6.93 (s, 1H), 6.88 (d, J = 2.3 Hz, 1H), 6.53 (s, 1H), 4.79 (s, 1H), 4.24 - 4.15 (m, 1H), 3.96 - 3.86 (m, 4H), 3.40 - 3.27 (m, 4H), 2.22 - 2.15 (m, 2H), 1.94 - 1.82 (m, 6H)
121		436.18	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 6.71 (d, J = 2.1 Hz, 1H), 6.54 (dd, J = 3.9, 1.7 Hz, 1H), 6.09 (dd, J = 3.9, 2.6 Hz, 1H), 5.94 (d, J = 8.1 Hz, 1H), 4.76 (s, 1H), 4.11 (s, 1H), 3.98 - 3.86 (m, 7H), 3.39 - 3.27 (m, 4H), 2.27 - 2.14 (m, 2H), 1.95 - 1.83 (m, 6H)
122		440.18	(400 MHz, CDCl ₃) δ 8.70 (s, 1H), 8.61 (s, 1H), 8.51 (s, 1H), 7.50 (s, 1H), 6.95 (s, 1H), 6.89 (s, 1H), 6.16 (d, 1H), 4.80 (s, 1H), 4.18 (s, 1H), 4.01 - 3.78 (m, 4H), 3.44 - 3.23 (m, 4H), 2.33 - 2.16 (m, 2H), 1.91 (d, J = 25.1 Hz, 6H)




Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
123		437.21	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.8 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.50 (d, J = 1.2 Hz, 1H), 7.36 (d, J = 1.0 Hz, 1H), 7.16 (d, J = 7.9 Hz, 1H), 6.94 (d, J = 2.3 Hz, 1H), 6.88 (d, J = 2.3 Hz, 1H), 4.78 (s, 1H), 4.20 - 4.09 (m, 1H), 3.95 - 3.87 (m, 4H), 3.73 (s, 3H), 3.38 - 3.28 (m, 4H), 2.25 - 2.14 (m, 2H), 2.00 - 1.82 (m, 6H)
124		473.2	(CDCl ₃) δ 8.68 (d, J = 1.9 Hz, 1H), 8.58 (d, J = 1.9 Hz, 1H), 7.85 - 7.72 (m, 2H), 6.95 (d, J = 2.5 Hz, 1H), 6.86 (d, J = 2.5 Hz, 1H), 4.89 (d, J = 7.5 Hz, 1H), 4.71 (dq, J = 5.5, 2.7 Hz, 1H), 4.00 - 3.85 (m, 7H), 3.47 - 3.27 (m, 5H), 2.19 - 2.07 (m, 2H), 1.97 - 1.65 (m, 6H)
125		473.25	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.55 (d, J = 1.9 Hz, 1H), 7.46 (d, J = 2.0 Hz, 1H), 6.97 (d, J = 2.5 Hz, 1H), 6.81 (dd, J = 22.5, 2.3 Hz, 2H), 5.65 - 5.56 (m, 1H), 4.73 (p, J = 2.5 Hz, 1H), 4.10 (s, 3H), 3.95 - 3.85 (m, 4H), 3.47 (d, J = 15.3 Hz, 1H), 3.36 - 3.27 (m, 4H), 2.23 - 2.08 (m, 2H), 1.98 - 1.62 (m, 6H)
126		473.25	(CDCl ₃) δ 8.68 (d, J = 1.9 Hz, 1H), 8.58 (d, J = 1.9 Hz, 1H), 7.41 (d, J = 2.3 Hz, 1H), 6.95 (d, J = 2.5 Hz, 1H), 6.86 (d, J = 2.5 Hz, 1H), 6.69 (d, J = 2.3 Hz, 1H), 4.95 (d, J = 7.3 Hz, 1H), 4.72 (h, J = 2.6 Hz, 1H), 4.02 - 3.82 (m, 7H), 3.52 (d, J = 9.0 Hz, 1H), 3.37 - 3.27 (m, 4H), 2.12 (dt, J = 16.1, 5.7 Hz, 2H), 1.97 - 1.65 (m, 6H)




Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
127		507.2	(CDCl ₃) δ 8.62 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 8.01 (s, 2H), 6.88 (s, 1H), 6.83 (d, J = 2.5 Hz, 1H), 4.93 (d, J = 8.1 Hz, 1H), 4.72 (d, J = 5.8 Hz, 1H), 4.53 (d, J = 6.0 Hz, 2H), 4.38 (d, J = 6.0 Hz, 2H), 3.89 - 3.75 (m, 4H), 3.35 - 3.13 (m, 4H), 2.10 (s, 3H), 1.95 - 1.72 (m, 6H), 1.36 (s, 3H)
128		477.2	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.56 (s, 2H), 7.04 - 6.87 (m, 2H), 6.19 (d, J = 6.8 Hz, 1H), 5.15 (d, J = 8.2 Hz, 1H), 5.01 (d, J = 6.8 Hz, 1H), 4.81 (s, 1H), 4.11 - 3.85 (m, 6H), 3.43 - 3.22 (m, 4H), 2.21 (d, J = 9.5 Hz, 2H), 1.94 (t, J = 3.9 Hz, 6H), 1.36 (t, J = 7.1 Hz, 3H)
129		478.93	(CDCl ₃) δ 8.60 (d, J = 1.9 Hz, 1H), 8.53 (d, J = 1.9 Hz, 1H), 7.81 (s, 2H), 6.91 - 6.75 (m, 2H), 5.08 - 4.89 (m, 2H), 4.89 - 4.77 (m, 2H), 4.67 (ddd, J = 8.2, 4.7, 1.9 Hz, 3H), 3.94 - 3.72 (m, 5H), 3.34 - 3.17 (m, 4H), 2.21 - 2.00 (m, 2H), 1.91 - 1.69 (m, 6H)
130		401.24	(400 MHz, methanol-d ₄) δ 8.69 (d, J = 2.0 Hz, 1H), 8.55 (d, J = 2.0 Hz, 1H), 7.10 (d, J = 2.3 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 4.85 - 4.82 (m, 1H), 4.07 (q, J = 6.9 Hz, 2H), 3.95 - 3.81 (m, 4H), 3.64 - 3.53 (m, 1H), 3.42 - 3.35 (m, 4H), 2.20 - 2.02 (m, 2H), 1.94 - 1.66 (m, 6H), 1.24 (t, J = 7.1 Hz, 3H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
131		429.21	(400 MHz, CDCl ₃) δ 8.73 - 8.65 (m, 1H), 8.60 (d, J = 1.9 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.87 (d, J = 2.3 Hz, 1H), 4.90 - 4.58 (m, 2H), 3.97 - 3.88 (m, 4H), 3.83 (d, J = 6.3 Hz, 2H), 3.72 (s, 1H), 3.39 - 3.27 (m, 4H), 2.20 - 2.06 (m, 2H), 1.95 - 1.76 (m, 7H), 0.93 (d, J = 6.7 Hz, 6H)
132		449.16	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.6 Hz, 1H), 8.62 (d, J = 1.7 Hz, 1H), 7.36 (t, J = 7.6 Hz, 2H), 7.19 (t, J = 7.3 Hz, 1H), 7.13 (d, J = 7.7 Hz, 2H), 6.96 (s, 1H), 6.89 (s, 1H), 5.12 (d, J = 7.2 Hz, 1H), 4.77 (s, 1H), 3.98 - 3.85 (m, 4H), 3.79 (s, 1H), 3.39 - 3.28 (m, 4H), 2.29 - 2.11 (m, 2H), 1.98 - 1.79 (m, 6H)
133		493.25	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.04 (s, 2H), 6.97 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.7 Hz, 1H), 4.79 (s, 2H), 4.07 - 3.83 (m, 6H), 3.42 - 3.25 (m, 4H), 2.29 - 2.02 (m, 2H), 1.92 (d, J = 4.6 Hz, 6H), 1.36 - 1.12 (m, 3H)
134		437.3	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.07 (s, 2H), 6.97 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.6 Hz, 1H), 4.99 (d, J = 8.0 Hz, 1H), 4.81 (d, J = 5.9 Hz, 1H), 3.93 (dd, J = 6.0, 3.7 Hz, 5H), 3.81 (s, 3H), 3.45 - 3.24 (m, 4H), 2.21 (d, J = 8.6 Hz, 2H), 2.05 - 1.78 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
135		479.2	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.19 (s, 2H), 6.96 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 5.12 (d, J = 8.1 Hz, 1H), 4.81 (d, J = 5.6 Hz, 1H), 4.09 - 3.85 (m, 5H), 3.63 - 3.41 (m, 4H), 3.43 - 3.26 (m, 4H), 2.69 (t, J = 6.6 Hz, 2H), 2.33 - 2.13 (m, 2H), 2.03 - 1.83 (m, 6H), 1.21 (t, J = 7.0 Hz, 3H)
136		494.24	(methanol-d ₄) δ 8.77 - 8.65 (m, 3H), 8.56 (d, J = 2.0 Hz, 1H), 7.11 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 4.92 (s, 1H), 4.02 (dd, J = 8.4, 4.3 Hz, 1H), 3.93 - 3.83 (m, 4H), 3.69 (t, J = 5.7 Hz, 2H), 3.47 (t, J = 5.8 Hz, 2H), 3.42 - 3.33 (m, 5H), 2.26 - 2.15 (m, 2H), 2.08 - 1.78 (m, 6H)
137		522.23	(CDCl ₃) δ 8.73 - 8.58 (m, 4H), 7.01 - 6.87 (m, 2H), 6.12 (d, J = 7.9 Hz, 1H), 5.57 (d, J = 8.1 Hz, 1H), 4.80 (s, 1H), 4.39 - 4.28 (m, 1H), 4.10 (d, J = 6.9 Hz, 1H), 3.97 - 3.87 (m, 4H), 3.54 - 3.29 (m, 9H), 2.21 (d, J = 9.9 Hz, 2H), 2.08 - 1.84 (m, 6H), 1.28 (d, J = 6.8 Hz, 3H)
138		452.92	(CDCl ₃) δ 9.04 (d, J = 3.2 Hz, 1H), 8.97 (d, J = 3.3 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.55 (d, J = 1.9 Hz, 1H), 6.90 (d, J = 2.5 Hz, 1H), 6.84 (d, J = 2.5 Hz, 1H), 6.03 (d, J = 8.2 Hz, 1H), 4.77 (td, J = 5.1, 2.6 Hz, 1H), 4.15 - 4.07 (m, 1H), 3.92 - 3.79 (m, 4H), 3.35 - 3.15 (m, 4H), 2.25 - 2.11 (m, 2H), 1.95 - 1.69 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
139		450.17	(400 MHz, CDCl ₃) δ 8.74 (d, J = 1.9 Hz, 2H), 8.70 (d, J = 1.9 Hz, 1H), 8.03 (dd, J = 8.9, 2.0 Hz, 1H), 6.97 (d, J = 2.4 Hz, 1H), 6.92 (d, J = 2.4 Hz, 1H), 6.37 (d, J = 8.9 Hz, 1H), 6.05 (s, 1H), 4.81 (s, 1H), 3.99 - 3.88 (m, 4H), 3.84 (s, 1H), 3.39 - 3.27 (m, 4H), 2.31 - 2.17 (m, 2H), 2.08 - 1.98 (m, 2H), 1.98 - 1.82 (m, 4H)
140		449.19	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.53 (d, J = 2.1 Hz, 1H), 7.88 (dd, J = 8.8, 2.4 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 6.39 (d, J = 8.7 Hz, 1H), 5.61 (s, 2H), 4.97 (d, J = 7.9 Hz, 1H), 4.80 (s, 1H), 4.02 - 3.82 (m, 5H), 3.42 - 3.26 (m, 4H), 2.27 - 2.14 (m, 2H), 2.00 - 1.81 (m, 6H)
141		492.17	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.09 (s, 2H), 7.01 - 6.86 (m, 2H), 4.88 - 4.75 (m, 1H), 4.08 - 3.80 (m, 9H), 3.43 - 3.27 (m, 4H), 3.09 - 2.96 (m, 4H), 2.31 - 2.15 (m, 2H), 2.05 - 1.81 (m, 6H)
142		421.2	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.14 (d, J = 0.8 Hz, 2H), 7.01 - 6.90 (m, 2H), 4.81 (td, J = 5.6, 2.7 Hz, 1H), 4.08 - 3.84 (m, 5H), 3.43 - 3.26 (m, 4H), 2.25 - 2.10 (m, 5H), 2.02 - 1.83 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
143		422.25	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.02 (s, 2H), 6.95 (dd, J = 11.8, 2.5 Hz, 2H), 4.81 (s, 2H), 4.08 - 3.85 (m, 5H), 3.41 - 3.30 (m, 4H), 2.20 (d, J = 10.1 Hz, 2H), 1.95 (d, J = 19.7 Hz, 6H), 1H NMR (300 MHz, Methanol-d ₄) ? 8.68 (d, J = 2.0 Hz, 1H), 8.56 (d, J = 2.1 Hz, 1H), 7.93 (s, 2H), 7.11 (d, J = 2.5 Hz, 1H), 6.88 (d, J = 2.4 Hz, 1H), 3.96 - 3.71 (m, 5H), 3.37 (dd, J = 5.8, 3.9 Hz, 4H), 2.27 - 2.04 (m, 2H), 1.98 - 1.74 (m, 6H). [2]
144		463.2	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.47 (d, J = 2.1 Hz, 1H), 7.84 (dd, J = 8.7, 2.4 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 6.37 (d, J = 8.7 Hz, 1H), 5.94 (d, J = 3.9 Hz, 1H), 4.92 (d, J = 7.8 Hz, 1H), 4.79 (s, 1H), 3.98 - 3.84 (m, 5H), 3.39 - 3.26 (m, 4H), 2.98 (d, J = 4.8 Hz, 3H), 2.27 - 2.12 (m, 2H), 1.97 - 1.83 (m, 6H)
145		477.2	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.20 (s, 1H), 7.61 (d, J = 8.2 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 6.43 (d, J = 7.9 Hz, 1H), 4.81 (s, 1H), 3.98 - 3.81 (m, 5H), 3.40 - 3.27 (m, 4H), 3.09 (s, 6H), 2.27 - 2.15 (m, 2H), 1.99 - 1.83 (m, 6H)




Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
146		464.17	(CDCl ₃) δ 8.76 (d, J = 2.0 Hz, 1H), 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.99 (dd, J = 8.8, 2.2 Hz, 1H), 6.97 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 6.38 (d, J = 8.9 Hz, 1H), 5.13 (s, 1H), 4.82 (s, 1H), 4.11 - 3.73 (m, 8H), 3.44 - 3.27 (m, 4H), 2.30 - 2.17 (m, 2H), 2.07 - 1.79 (m, 6H)
147		423.2	(400 MHz, CDCl ₃) δ 10.77 - 10.42 (m, 1H), 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.30 (d, J = 8.2 Hz, 1H), 7.15 (d, J = 5.7 Hz, 2H), 6.95 (d, J = 2.5 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 4.83 (s, 1H), 4.19 - 4.05 (m, 1H), 3.99 - 3.84 (m, 4H), 3.40 - 3.27 (m, 4H), 2.31 - 2.15 (m, 2H), 2.06 - 1.93 (m, 2H), 1.93 - 1.79 (m, 4H)
148		534.28	(DMSO-d ₆) δ 8.73 (d, J = 1.5 Hz, 3H), 8.59 (d, J = 1.9 Hz, 1H), 8.12 (d, J = 8.2 Hz, 1H), 7.84 (d, J = 7.4 Hz, 1H), 7.14 (d, J = 2.4 Hz, 1H), 6.84 (d, J = 2.3 Hz, 1H), 4.93 (s, 1H), 4.51 (ddd, J = 13.3, 8.2, 5.2 Hz, 1H), 4.01 - 3.74 (m, 6H), 3.55 (q, J = 8.0 Hz, 1H), 3.38 - 3.27 (m, 4H), 2.33 - 2.15 (m, 1H), 2.06 (d, J = 11.1 Hz, 2H), 1.96 - 1.67 (m, 8H), 1.02 (d, J = 6.3 Hz, 3H)




Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
149		520.33	(DMSO-d ₆) δ 8.76 - 8.67 (m, 3H), 8.59 (d, J = 1.9 Hz, 1H), 8.32 (d, J = 6.5 Hz, 1H), 7.84 (d, J = 7.3 Hz, 1H), 7.14 (d, J = 2.5 Hz, 1H), 6.84 (d, J = 2.3 Hz, 1H), 4.93 (s, 1H), 4.49 - 4.37 (m, 1H), 3.96 - 3.63 (m, 9H), 3.55 (dd, J = 8.9, 4.2 Hz, 1H), 3.38 - 3.27 (m, 4H), 2.23 - 2.00 (m, 4H), 1.95 - 1.71 (m, 6H)
150		520.33	(DMSO-d ₆) δ 8.76 - 8.67 (m, 3H), 8.59 (d, J = 1.9 Hz, 1H), 8.32 (d, J = 6.5 Hz, 1H), 7.84 (d, J = 7.3 Hz, 1H), 7.14 (d, J = 2.5 Hz, 1H), 6.84 (d, J = 2.3 Hz, 1H), 4.93 (s, 1H), 4.49 - 4.37 (m, 1H), 3.96 - 3.63 (m, 9H), 3.55 (dd, J = 8.9, 4.2 Hz, 1H), 3.38 - 3.27 (m, 4H), 2.23 - 2.00 (m, 4H), 1.95 - 1.71 (m, 6H)
151			(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 2.0 Hz, 1H), 7.93 (s, 2H), 7.02 - 6.90 (m, 2H), 4.82 (t, J = 6.7 Hz, 3H), 4.66 (t, J = 6.4 Hz, 2H), 4.14 (q, J = 7.1 Hz, 1H), 4.08 - 3.89 (m, 5H), 3.42 - 3.25 (m, 4H), 2.80 (s, 3H), 2.22 (s, 2H), 2.05 - 1.79 (m, 6H)
152		464.21	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 7.94 (s, 2H), 6.99 - 6.90 (m, 2H), 4.82 (s, 1H), 4.05 - 3.87 (m, 5H), 3.44 (q, J = 6.3 Hz, 1H), 3.39 - 3.26 (m, 4H), 2.23 (d, J = 12.4 Hz, 2H), 2.03 - 1.82 (m, 6H), 1.23 (d, J = 6.3 Hz, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
153		483.1	(CDCl ₃) δ 8.64 - 8.49 (m, 2H), 6.93 (d, J = 3.9 Hz, 1H), 6.83 (d, J = 2.4 Hz, 1H), 5.27 (d, J = 8.1 Hz, 1H), 4.94 (t, J = 1.9 Hz, 2H), 4.81 (t, J = 1.9 Hz, 2H), 4.72 (s, 1H), 3.96 (s, 1H), 3.89 - 3.78 (m, 4H), 3.37 - 3.25 (m, 4H), 2.11 (s, 2H), 1.82 (d, J = 5.1 Hz, 6H)
154		483.1	(CDCl ₃) δ 8.68 (dd, J = 8.7, 2.0 Hz, 1H), 8.56 (d, J = 2.0 Hz, 1H), 6.92 (d, J = 2.3 Hz, 1H), 6.84 (d, J = 2.4 Hz, 1H), 4.92 (t, J = 2.7 Hz, 2H), 4.82 (t, J = 2.7 Hz, 2H), 4.75 (s, 1H), 3.94 - 3.81 (m, 4H), 3.35 - 3.15 (m, 4H), 2.14 (d, J = 12.0 Hz, 2H), 1.82 (d, J = 17.3 Hz, 6H)
155		478.3	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 7.02 - 6.89 (m, 2H), 4.83 (s, 1H), 4.08 - 3.87 (m, 5H), 3.44 - 3.33 (m, 4H), 3.27 (d, J = 26.0 Hz, 4H), 2.23 (d, J = 9.8 Hz, 2H), 2.05 - 1.80 (m, 6H), 1.14 (s, 6H)
156		462.23	(CDCl ₃) δ 8.70 (d, J = 2.0 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.72 (s, 2H), 6.96 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 5.03 (s, 1H), 4.80 (q, J = 4.4 Hz, 1H), 4.03 - 3.89 (m, 4H), 3.83 (t, J = 7.1 Hz, 4H), 3.44 - 3.29 (m, 4H), 2.41 (dq, J = 8.6, 7.1 Hz, 2H), 2.20 (q, J = 5.9 Hz, 2H), 1.99 - 1.81 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
157		433.2	
158			(CDCl ₃) δ 8.71 (d, J = 2.0 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.41 (s, 2H), 6.97 (d, J = 2.4 Hz, 1H), 6.93 (d, J = 2.5 Hz, 1H), 5.32 (d, J = 8.1 Hz, 1H), 4.82 (s, 1H), 4.07 (s, 1H), 4.00 - 3.88 (m, 4H), 3.43 - 3.29 (m, 4H), 2.22 (s, 2H), 2.06 - 1.81 (m, 6H)
159		434.24	
160		436.2	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.05 (d, J = 2.2 Hz, 1H), 7.46 (dd, J = 8.5, 2.3 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 6.39 (d, J = 8.6 Hz, 1H), 4.77 (s, 1H), 4.53 (s, 2H), 3.95 - 3.87 (m, 5H), 3.36 - 3.32 (m, 4H), 2.25 - 2.12 (m, 2H), 1.90 (d, J = 4.4 Hz, 6H)

Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
161		451.28	(CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.26 (s, 2H), 6.95 (d, J = 2.5 Hz, 1H), 6.90 (d, J = 2.6 Hz, 1H), 5.23 (d, J = 8.1 Hz, 1H), 4.80 (d, J = 5.8 Hz, 1H), 4.26 (s, 2H), 4.03 (s, 1H), 3.97 - 3.85 (m, 4H), 3.44 - 3.21 (m, 7H), 2.32 - 2.09 (m, 2H), 2.06 - 1.70 (m, 6H)
162		492.29	(DMSO-d ₆) δ 8.81 - 8.64 (m, 3H), 8.58 (d, J = 1.9 Hz, 1H), 7.99 (d, J = 7.6 Hz, 1H), 7.76 (d, J = 7.4 Hz, 1H), 7.14 (d, J = 2.5 Hz, 1H), 6.84 (d, J = 2.3 Hz, 1H), 4.92 (s, 1H), 3.98 - 3.84 (m, 1H), 4.05 (dq, J = 13.5, 6.7 Hz, 1H), 3.79 (d, J = 9.6 Hz, 4H), 3.32 (d, J = 8.2 Hz, 4H), 2.06 (d, J = 11.8 Hz, 2H), 1.96 - 1.66 (m, 6H), 1.14 (d, J = 6.6 Hz, 6H)
163		504.26	(CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.65 - 8.53 (m, 3H), 6.93 (dd, J = 14.6, 2.5 Hz, 2H), 5.48 (d, J = 8.0 Hz, 1H), 4.81 (d, J = 6.1 Hz, 1H), 4.19 - 4.03 (m, 1H), 3.92 (dd, J = 6.0, 3.7 Hz, 4H), 3.65 - 3.54 (m, 4H), 3.34 (dd, J = 5.9, 3.7 Hz, 4H), 2.27 - 2.15 (m, 2H), 2.08 - 1.81 (m, 10H)
164		484.12	(400 MHz, CDCl ₃) δ 8.60 (d, J = 1.9 Hz, 1H), 8.52 (d, J = 1.9 Hz, 1H), 8.01 (d, J = 2.1 Hz, 1H), 7.35 (dd, J = 8.8, 2.5 Hz, 1H), 6.86 (d, J = 2.5 Hz, 1H), 6.80 (d, J = 2.5 Hz, 1H), 6.20 (d, J = 8.4 Hz, 1H), 4.68 (s, 1H), 4.47 (d, J = 8.0 Hz, 1H), 3.92 - 3.80 (m, 4H), 3.74 (s, 1H), 3.34 - 3.14 (m, 4H), 2.18 - 2.02 (m, 2H), 1.94 - 1.67 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
165		448.15	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.44 - 8.37 (m, 1H), 8.33 (d, J = 8.1 Hz, 1H), 7.58 (dd, J = 7.8, 0.9 Hz, 1H), 7.31 (dd, J = 7.7, 4.6 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 4.75 (s, 1H), 4.25 - 4.10 (m, 1H), 3.97 - 3.85 (m, 4H), 3.41 - 3.26 (m, 4H), 2.75 (s, 3H), 2.29 - 2.13 (m, 2H), 2.06 - 1.85 (m, 6H)
166		448.19	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.41 (d, J = 4.5 Hz, 1H), 8.21 (d, J = 8.0 Hz, 1H), 8.08 - 7.98 (m, 1H), 7.25 - 7.22 (m, 1H), 6.95 (d, J = 2.5 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 4.76 (s, 1H), 4.27 - 4.15 (m, 1H), 3.98 - 3.86 (m, 4H), 3.40 - 3.29 (m, 4H), 2.43 (s, 3H), 2.29 - 2.16 (m, 2H), 2.07 - 1.86 (m, 6H)
167		448.19	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.37 (dd, J = 1.4, 0.7 Hz, 1H), 8.16 (d, J = 8.4 Hz, 1H), 8.09 (d, J = 8.0 Hz, 1H), 7.66 - 7.61 (m, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 4.75 (s, 1H), 4.29 - 4.16 (m, 1H), 3.97 - 3.86 (m, 4H), 3.39 - 3.29 (m, 4H), 2.41 (s, 3H), 2.28 - 2.15 (m, 2H), 2.04 - 1.84 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
168		462.16	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.8 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.39 (d, J = 1.6 Hz, 1H), 8.17 (d, J = 8.6 Hz, 1H), 8.11 (d, J = 8.0 Hz, 1H), 7.66 (dd, J = 7.9, 2.0 Hz, 1H), 6.95 (d, J = 2.3 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 4.76 (s, 1H), 4.31 - 4.14 (m, 1H), 4.00 - 3.85 (m, 4H), 3.42 - 3.25 (m, 4H), 2.73 (q, J = 7.6 Hz, 2H), 2.30 - 2.14 (m, 2H), 1.99 (ddd, J = 34.6, 19.6, 10.3 Hz, 7H), 1.29 (t, J = 7.6 Hz, 3H)
169		448.19	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.24 (d, J = 7.8 Hz, 1H), 8.01 (d, J = 7.5 Hz, 1H), 7.72 (t, J = 7.7 Hz, 1H), 7.27 (d, J = 6.6 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 4.82 - 4.68 (m, 1H), 4.27 - 4.13 (m, 1H), 3.98 - 3.86 (m, 4H), 3.41 - 3.28 (m, 4H), 2.60 (s, 3H), 2.30 - 2.18 (m, 2H), 2.09 - 1.88 (m, 6H)
170		464.17	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.8 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.90 (d, J = 7.2 Hz, 1H), 7.83 - 7.78 (m, 1H), 7.76 - 7.69 (m, 1H), 6.96 (d, J = 2.3 Hz, 1H), 6.94 - 6.86 (m, 2H), 4.77 (s, 1H), 4.17 (s, 1H), 4.00 (s, 3H), 3.95 - 3.86 (m, 4H), 3.40 - 3.29 (m, 4H), 2.30 - 2.17 (m, 2H), 2.06 - 1.88 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
171		459.17	(400 MHz, CDCl ₃) δ 8.76 (dd, J = 4.9, 0.8 Hz, 1H), 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.48 - 8.38 (m, 1H), 8.09 (d, J = 8.0 Hz, 1H), 7.67 (dd, J = 4.9, 1.6 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 4.76 (s, 1H), 4.31 - 4.14 (m, 1H), 4.01 - 3.83 (m, 4H), 3.43 - 3.26 (m, 4H), 2.31 - 2.15 (m, 2H), 2.08 - 1.87 (m, 6H)
172		452.22	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.11 (dd, J = 7.4, 1.4 Hz, 1H), 7.96 (dd, J = 15.6, 7.7 Hz, 1H), 7.82 (d, J = 7.7 Hz, 1H), 7.09 (dd, J = 8.1, 1.8 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 4.80 (s, 1H), 4.26 - 4.09 (m, 1H), 3.99 - 3.86 (m, 4H), 3.41 - 3.25 (m, 4H), 2.31 - 2.17 (m, 2H), 2.05 - 1.86 (m, 6H)
173		452.22	
174		470.22	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.30 (d, J = 2.1 Hz, 1H), 7.80 (d, J = 7.9 Hz, 1H), 7.32 (ddd, J = 10.3, 8.1, 2.3 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 4.74 (s, 1H), 4.27 - 4.15 (m, 1H), 4.00 - 3.86 (m, 4H), 3.41 - 3.27 (m, 4H), 2.27 - 2.13 (m, 2H), 2.08 - 1.86 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
175		450.13	
176		450.2	(400 MHz, CDCl ₃) δ 12.21 (s, 1H), 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.17 (d, J = 7.4 Hz, 1H), 8.07 (dd, J = 4.2, 1.5 Hz, 1H), 7.33 (ddd, J = 10.0, 8.5, 2.9 Hz, 2H), 6.97 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.5 Hz, 1H), 4.83 - 4.74 (m, 1H), 4.24 - 4.11 (m, 1H), 3.99 - 3.86 (m, 4H), 3.41 - 3.26 (m, 4H), 2.30 - 2.18 (m, 2H), 2.08 - 1.87 (m, 6H)
177		450.17	(400 MHz, CDCl ₃) δ 11.13 (s, 1H), 8.71 (d, J = 1.9 Hz, 1H), 8.59 (d, J = 1.9 Hz, 1H), 7.64 - 7.39 (m, 2H), 7.07 (d, J = 6.8 Hz, 1H), 7.02 - 6.79 (m, 3H), 4.76 (s, 1H), 4.32 - 4.15 (m, 1H), 3.96 - 3.85 (m, 4H), 3.42 - 3.24 (m, 4H), 2.39 - 2.14 (m, 2H), 2.10 - 1.82 (m, 6H)
178		468.17	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.14 (dd, J = 7.6, 0.7 Hz, 1H), 7.91 (d, J = 8.1 Hz, 1H), 7.81 (t, J = 7.8 Hz, 1H), 7.46 (dd, J = 7.9, 0.7 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 4.80 (s, 1H), 4.24 - 4.08 (m, 1H), 3.99 - 3.83 (m, 4H), 3.41 - 3.26 (m, 4H), 2.33 - 2.18 (m, 2H), 2.11 - 1.98 (m, 2H), 1.98 - 1.87 (m, 4H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
179		464.28	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.13 (d, J = 7.7 Hz, 1H), 8.02 (d, J = 7.8 Hz, 1H), 7.86 (t, J = 7.7 Hz, 1H), 7.48 (d, J = 7.8 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 4.85 (s, 2H), 4.79 (s, 1H), 4.28 - 4.15 (m, 1H), 4.00 - 3.87 (m, 4H), 3.41 - 3.27 (m, 4H), 2.88 (s, 1H), 2.33 - 2.16 (m, 2H), 2.07 - 1.88 (m, 6H)
180		503.27	(400 MHz, CDCl ₃) δ 8.68 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.16 (d, J = 7.8 Hz, 1H), 7.56 (dd, J = 8.4, 7.3 Hz, 1H), 7.43 (d, J = 6.7 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 6.50 (d, J = 7.8 Hz, 1H), 4.74 (s, 1H), 4.21 - 4.07 (m, 1H), 3.97 - 3.86 (m, 4H), 3.54 - 3.48 (m, 4H), 3.40 - 3.29 (m, 4H), 2.26 - 2.14 (m, 2H), 2.07 - 1.88 (m, 10H)
181		501.25	(400 MHz, CDCl ₃) δ 8.91 (s, 2H), 8.71 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.30 (dd, J = 7.7, 0.7 Hz, 1H), 8.11 (t, J = 7.9 Hz, 1H), 7.74 (d, J = 8.2 Hz, 1H), 7.57 (dd, J = 8.0, 0.7 Hz, 1H), 6.97 (d, J = 2.4 Hz, 1H), 6.92 (d, J = 2.4 Hz, 1H), 4.82 (s, 1H), 4.27 - 4.11 (m, 1H), 3.99 - 3.85 (m, 4H), 3.42 - 3.27 (m, 4H), 2.34 - 2.20 (m, 2H), 2.12 - 2.00 (m, 2H), 2.00 - 1.84 (m, 4H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
182		424.21	(400 MHz, CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.15 (s, 1H), 7.43 (d, J = 8.1 Hz, 1H), 6.97 (d, J = 2.3 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 4.82 (s, 1H), 4.24 - 4.13 (m, 1H), 3.98 - 3.86 (m, 4H), 3.40 - 3.27 (m, 4H), 2.30 - 2.16 (m, 2H), 2.05 - 1.78 (m, 6H)
183		440.12	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.86 (d, J = 3.1 Hz, 1H), 7.57 (d, J = 3.1 Hz, 1H), 7.38 (d, J = 8.0 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 4.79 (s, 1H), 4.25 - 4.09 (m, 1H), 3.97 - 3.88 (m, 4H), 3.39 - 3.29 (m, 4H), 2.28 - 2.17 (m, 2H), 2.07 - 1.96 (m, 2H), 1.96 - 1.85 (m, 4H)
184		454.09	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.50 (d, J = 1.1 Hz, 1H), 7.28 (d, J = 8.3 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 4.78 (s, 1H), 4.23 - 4.07 (m, 1H), 3.99 - 3.87 (m, 4H), 3.40 - 3.27 (m, 4H), 2.53 (d, J = 1.0 Hz, 3H), 2.28 - 2.14 (m, 2H), 2.04 - 1.95 (m, 2H), 1.95 - 1.84 (m, 4H)
185		454.13	(400 MHz, CDCl ₃) δ 8.70 (t, J = 5.6 Hz, 1H), 8.61 (dd, J = 12.5, 1.9 Hz, 1H), 7.35 (d, J = 8.0 Hz, 1H), 7.11 (d, J = 0.9 Hz, 1H), 7.01 - 6.92 (m, 1H), 6.89 (d, J = 2.4 Hz, 1H), 4.85 - 4.70 (m, 1H), 4.26 - 4.11 (m, 1H), 3.90 (dd, J = 15.3, 10.4 Hz, 4H), 3.42 - 3.25 (m, 4H), 2.49 (d, J = 0.9 Hz, 3H), 2.30 - 2.16 (m, 2H), 2.12 - 1.84 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
186		437.2	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.01 (d, J = 5.8 Hz, 1H), 7.00 - 6.86 (m, 2H), 6.03 (d, J = 5.8 Hz, 1H), 4.80 (d, J = 5.8 Hz, 1H), 4.07 (s, 1H), 3.97 - 3.83 (m, 7H), 3.41 - 3.28 (m, 4H), 2.20 (dd, J = 12.3, 5.6 Hz, 2H), 1.95 (dh, J = 11.8, 5.6, 4.7 Hz, 6H)
187		492.26	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.70 (s, 1H), 6.96 (q, J = 2.6 Hz, 2H), 5.75 (d, J = 6.5 Hz, 1H), 4.82 (s, 1H), 4.65 (s, 1H), 3.93 (dd, J = 5.9, 3.8 Hz, 4H), 3.63 (s, 4H), 3.36 (dd, J = 6.1, 3.7 Hz, 4H), 2.14 (s, 2H), 1.95 (d, J = 31.2 Hz, 6H)
188		451.14	(400 MHz, CDCl ₃) δ 9.02 (d, J = 2.6 Hz, 1H), 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.17 (dd, J = 9.3, 2.6 Hz, 1H), 6.97 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 6.36 (d, J = 9.3 Hz, 1H), 5.37 (s, 1H), 4.82 (s, 1H), 4.04 (s, 1H), 3.96 - 3.87 (m, 4H), 3.40 - 3.27 (m, 4H), 2.31 - 2.18 (m, 2H), 2.03 - 1.84 (m, 6H)
189		474.147	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.6 Hz, 1H), 8.61 (d, J = 1.7 Hz, 1H), 8.32 (s, 1H), 7.56 (d, J = 6.7 Hz, 1H), 6.96 (d, J = 2.2 Hz, 1H), 6.90 (d, J = 2.3 Hz, 1H), 6.40 (d, J = 8.7 Hz, 1H), 4.96 (s, 1H), 4.80 (s, 1H), 4.01 - 3.84 (m, 5H), 3.42 - 3.24 (m, 4H), 2.21 (d, J = 8.4 Hz, 2H), 1.92 (d, J = 6.4 Hz, 6H)






Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
190			(CDCl ₃) δ 8.52 (s, 1H), 8.29 (d, J = 4.8 Hz, 2H), 6.90 (d, J = 2.5 Hz, 1H), 6.85 (d, J = 2.5 Hz, 1H), 6.52 (t, J = 4.8 Hz, 1H), 5.36 (s, 1H), 4.79 (dq, J = 5.6, 2.9 Hz, 1H), 4.04 (dp, J = 8.0, 3.8 Hz, 1H), 3.97 - 3.85 (m, 4H), 3.38 - 3.26 (m, 4H), 2.70 (s, 3H), 2.29 - 2.11 (m, 2H), 2.00 - 1.78 (m, 6H)
191		421.24	(CDCl ₃) δ 8.52 (s, 1H), 8.29 (d, J = 4.8 Hz, 2H), 6.90 (d, J = 2.5 Hz, 1H), 6.85 (d, J = 2.5 Hz, 1H), 6.53 (t, J = 4.8 Hz, 1H), 5.42 (d, J = 7.9 Hz, 1H), 4.80 (dq, J = 5.9, 2.9 Hz, 1H), 4.11 - 3.98 (m, 1H), 3.91 (dd, J = 5.9, 3.8 Hz, 4H), 3.42 - 3.21 (m, 4H), 2.71 (s, 3H), 2.31 - 2.07 (m, 2H), 2.04 - 1.77 (m, 6H)
192		463.2	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.97 (s, 1H), 6.97 (d, J = 2.5 Hz, 1H), 6.93 (d, J = 2.5 Hz, 1H), 5.36 (s, 1H), 4.80 (d, J = 6.2 Hz, 1H), 4.65 (d, J = 0.9 Hz, 2H), 4.02 (t, J = 5.8 Hz, 3H), 3.98 - 3.82 (m, 4H), 3.42 - 3.30 (m, 4H), 2.80 (t, J = 5.8 Hz, 2H), 2.29 - 2.11 (m, 2H), 2.03 - 1.77 (m, 7H)
193		524.21	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.95 (s, 2H), 7.05 - 6.88 (m, 2H), 4.92 (s, 1H), 4.79 (s, 1H), 3.93 (t, J = 4.8 Hz, 5H), 3.60 - 3.44 (m, 4H), 3.37 (d, J = 10.5 Hz, 9H), 2.20 (d, J = 6.9 Hz, 2H), 1.91 (d, J = 4.6 Hz, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
194		437.23	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.28 (d, J = 0.9 Hz, 1H), 6.97 (d, J = 2.5 Hz, 1H), 6.91 (d, J = 2.5 Hz, 1H), 5.74 - 5.58 (m, 1H), 5.16 (d, J = 8.0 Hz, 1H), 4.82 (dq, J = 5.4, 2.7 Hz, 1H), 4.06 - 3.85 (m, 7H), 3.70 (d, J = 14.0 Hz, 1H), 3.42 - 3.28 (m, 4H), 2.30 - 2.16 (m, 2H), 1.99 - 1.75 (m, 6H)
195		431.22	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 2.0 Hz, 1H), 7.43 (dd, J = 8.6, 7.2 Hz, 1H), 7.02 - 6.86 (m, 3H), 6.55 (dd, J = 8.6, 0.8 Hz, 1H), 4.80 (d, J = 6.6 Hz, 2H), 4.10 - 3.81 (m, 5H), 3.47 - 3.25 (m, 4H), 2.30 - 2.11 (m, 2H), 1.99 - 1.81 (m, 6H)
196		463.27	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.61 (d, J = 9.4 Hz, 1H), 7.41 (d, J = 7.2 Hz, 1H), 7.05 - 6.87 (m, 2H), 6.65 (s, 1H), 4.86 (s, 1H), 4.04 - 3.87 (m, 7H), 3.74 (s, 1H), 3.43 - 3.27 (m, 4H), 2.21 (d, J = 10.8 Hz, 2H), 1.98 (d, J = 23.5 Hz, 6H)
197		421.69	(400 MHz, CDCl ₃) δ 8.68 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.13 - 8.04 (m, 1H), 7.67 (d, J = 2.4 Hz, 1H), 6.99 - 6.93 (m, 2H), 6.90 (d, J = 2.4 Hz, 1H), 6.33 (d, J = 8.7 Hz, 1H), 4.77 (s, 1H), 3.99 - 3.65 (m, 7H), 3.39 - 3.27 (m, 4H), 2.25 - 2.10 (m, 2H), 1.96 - 1.82 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
198		532.11	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.6 Hz, 1H), 8.61 (d, J = 1.7 Hz, 1H), 8.57 (d, J = 2.1 Hz, 1H), 7.89 (dd, J = 8.9, 2.3 Hz, 1H), 6.95 (s, 1H), 6.90 (s, 1H), 6.62 (d, J = 9.0 Hz, 1H), 5.99 (d, J = 7.7 Hz, 1H), 4.78 (s, 1H), 4.18 (s, 1H), 4.03 - 3.81 (m, 4H), 3.76 - 3.60 (m, 4H), 3.42 - 3.25 (m, 4H), 2.62 - 2.44 (m, 4H), 2.36 (s, 3H), 2.29 - 2.14 (m, 2H), 1.99 - 1.84 (m, 6H)
199		533.01	(400 MHz, CDCl ₃) δ 8.73 - 8.65 (m, 3H), 8.60 (d, J = 1.9 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 5.93 (d, J = 8.0 Hz, 1H), 4.79 (s, 1H), 4.23 - 4.09 (m, 1H), 3.97 - 3.88 (m, 8H), 3.40 - 3.25 (m, 4H), 2.53 - 2.42 (m, 4H), 2.34 (s, 3H), 2.26 - 2.18 (m, 2H), 1.97 - 1.83 (m, 6H)
200		549.17	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.51 - 7.42 (m, 2H), 6.98 - 6.85 (m, 3H), 6.07 (d, J = 8.0 Hz, 1H), 4.77 (s, 1H), 4.25 - 4.11 (m, 1H), 3.99 - 3.85 (m, 4H), 3.40 - 3.27 (m, 4H), 3.26 - 3.14 (m, 4H), 2.67 - 2.52 (m, 4H), 2.36 (s, 3H), 2.28 - 2.13 (m, 2H), 1.98 - 1.84 (m, 6H)
201		549.1	(400 MHz, CDCl ₃) δ 9.47 (s, 1H), 8.68 (d, J = 1.7 Hz, 1H), 8.56 (d, J = 1.7 Hz, 1H), 8.21 - 8.11 (m, 1H), 6.97 - 6.85 (m, 4H), 4.82 (s, 1H), 4.23 - 4.07 (m, 1H), 3.96 - 3.87 (m, 4H), 3.38 - 3.29 (m, 4H), 3.03 (s, 4H), 2.65 (s, 4H), 2.37 - 2.25 (m, 5H), 2.02 - 1.84 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
202		464.13	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.8 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.27 (dd, J = 4.7, 1.9 Hz, 1H), 8.19 (s, 1H), 8.11 (d, J = 6.8 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.92 (d, J = 2.3 Hz, 1H), 6.51 (dd, J = 7.4, 4.9 Hz, 1H), 4.76 (s, 1H), 4.30 (s, 1H), 4.02 - 3.90 (m, 4H), 3.88 (s, 3H), 3.41 - 3.26 (m, 4H), 2.29 - 2.11 (m, 2H), 2.11 - 1.85 (m, 6H)
203		432.58	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 8.15 (s, 1H), 6.93 (dd, J = 17.9, 2.5 Hz, 2H), 6.43 (d, J = 6.1 Hz, 1H), 5.20 (s, 1H), 4.82 (s, 1H), 4.00 - 3.82 (m, 4H), 3.44 - 3.25 (m, 4H), 2.23 (d, J = 11.2 Hz, 2H), 1.91 (s, 6H)
204		421.65	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.09 (d, J = 6.0 Hz, 1H), 6.93 (dd, J = 16.1, 2.5 Hz, 2H), 6.15 (d, J = 6.0 Hz, 1H), 5.10 (d, J = 8.0 Hz, 1H), 4.81 (td, J = 5.5, 2.7 Hz, 1H), 3.97 - 3.87 (m, 4H), 3.49 (s, 1H), 3.39 - 3.27 (m, 4H), 2.49 (s, 3H), 2.27 - 2.14 (m, 2H), 2.03 - 1.80 (m, 6H)
205		436.63	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (dd, J = 5.3, 1.9 Hz, 1H), 8.19 - 8.06 (m, 1H), 6.99 - 6.84 (m, 2H), 5.34 - 5.21 (m, 1H), 4.76 (d, J = 9.7 Hz, 3H), 3.92 (t, J = 4.9 Hz, 4H), 3.81 (s, 1H), 3.33 (dd, J = 5.7, 4.1 Hz, 4H), 2.87 (d, J = 5.2 Hz, 3H), 2.19 (d, J = 8.5 Hz, 2H), 1.88 (dd, J = 13.3, 5.1 Hz, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
206		436.18	(400 MHz, CDCl ₃) δ 8.68 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.82 (d, J = 2.8 Hz, 1H), 7.09 (dd, J = 8.9, 3.0 Hz, 1H), 6.94 (d, J = 2.5 Hz, 1H), 6.90 (d, J = 2.5 Hz, 1H), 6.37 (d, J = 8.8 Hz, 1H), 4.77 (s, 1H), 4.29 (bs, 1H), 3.98 - 3.87 (m, 4H), 3.85 - 3.79 (m, 1H), 3.77 (s, 3H), 3.41 - 3.24 (m, 4H), 2.27 - 2.12 (m, 2H), 1.97 - 1.79 (m, 6H)
207			(CDCl ₃) δ 8.71 (d, J = 2.0 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.44 (s, 1H), 6.98 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 6.82 (d, J = 4.7 Hz, 1H), 5.56 (d, J = 30.8 Hz, 1H), 4.83 (d, J = 5.2 Hz, 1H), 4.04 (s, 2H), 3.97 - 3.84 (m, 4H), 3.44 - 3.29 (m, 4H), 2.23 (d, J = 8.2 Hz, 2H), 2.02 - 1.77 (m, 6H)
208		435.6	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.16 (s, 2H), 7.00 - 6.84 (m, 2H), 5.32 (s, 1H), 4.81 (s, 1H), 4.03 (s, 1H), 3.96 - 3.85 (m, 4H), 3.43 - 3.32 (m, 4H), 2.49 (q, J = 7.6 Hz, 2H), 2.21 (d, J = 8.8 Hz, 2H), 2.06 - 1.75 (m, 6H), 1.21 (t, J = 7.6 Hz, 3H)
209			(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.18 (s, 2H), 7.01 - 6.88 (m, 2H), 5.30 (d, J = 9.0 Hz, 1H), 4.81 (d, J = 5.9 Hz, 1H), 4.02 (s, 1H), 3.96 - 3.85 (m, 4H), 3.42 - 3.29 (m, 4H), 2.78 (p, J = 6.9 Hz, 1H), 2.31 - 2.14 (m, 2H), 2.01 - 1.83 (m, 6H), 1.25 (d, J = 6.9 Hz, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
210		450.17	(400 MHz, methanol-d ₄) δ 8.69 (d, J = 2.0 Hz, 1H), 8.57 (d, J = 2.0 Hz, 1H), 7.92 (d, J = 6.2 Hz, 1H), 7.17 - 7.10 (m, 2H), 7.02 (d, J = 7.1 Hz, 1H), 6.90 (d, J = 2.5 Hz, 1H), 4.93 (s, 1H), 3.93 - 3.87 (m, 4H), 3.84 - 3.79 (m, 1H), 3.44 - 3.37 (m, 4H), 2.24 - 2.15 (m, 2H), 1.97 - 1.82 (m, 6H)
211		434.19	
212		437.12	
213		434.15	
214		474.12	(400 MHz, CDCl ₃) δ 8.98 - 8.83 (m, 1H), 8.79 - 8.67 (m, 1H), 8.64 - 8.50 (m, 1H), 8.35 (d, J = 9.2 Hz, 1H), 7.04 - 6.86 (m, 2H), 6.75 (d, J = 9.0 Hz, 2H), 4.78 (s, 1H), 4.00 - 3.79 (m, 5H), 3.42 - 3.21 (m, 4H), 2.17 - 2.11 (m, 2H), 2.02 - 1.81 (m, 6H)

Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
215		451.21	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.25 (d, J = 0.9 Hz, 1H), 6.93 (dd, J = 17.1, 2.5 Hz, 2H), 5.64 (d, J = 0.9 Hz, 1H), 5.00 (d, J = 8.1 Hz, 1H), 4.80 (dq, J = 5.5, 2.7 Hz, 1H), 4.33 (q, J = 7.1 Hz, 2H), 3.97 - 3.87 (m, 4H), 3.71 (s, 1H), 3.38 - 3.29 (m, 4H), 2.27 - 2.14 (m, 2H), 2.03 - 1.80 (m, 6H), 1.37 (t, J = 7.1 Hz, 3H)
216		481.26	(CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 6.95 (d, J = 2.5 Hz, 1H), 6.90 (d, J = 2.5 Hz, 1H), 0.23 - 0.16 (m, 0H), 4.93 (s, 1H), 4.76 (s, 1H), 4.19 - 4.01 (m, 1H), 3.97 - 3.87 (m, 4H), 3.38 - 3.29 (m, 4H), 2.49 (s, 3H), 2.26 (s, 3H), 2.21 - 2.10 (m, 2H), 2.05 (d, J = 1.8 Hz, 3H), 2.00 - 1.85 (m, 6H)
217		464.17	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.15 (d, J = 5.3 Hz, 1H), 7.06 (dd, J = 5.4, 1.3 Hz, 1H), 7.00 (s, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 5.15 (s, 1H), 4.81 (d, J = 2.5 Hz, 1H), 3.95 - 3.83 (m, 8H), 3.38 - 3.30 (m, 4H), 2.27 - 2.16 (m, 2H), 1.97 - 1.85 (m, 6H)
218			(CDCl ₃) δ 8.70 (d, J = 2.0 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.17 (d, J = 0.9 Hz, 1H), 7.02 - 6.85 (m, 2H), 5.40 (s, 1H), 5.05 (td, J = 1.8, 0.8 Hz, 2H), 4.91 - 4.73 (m, 3H), 4.05 (s, 0H), 3.99 - 3.85 (m, 4H), 3.43 - 3.27 (m, 4H), 2.30 - 2.12 (m, 2H), 2.02 - 1.82 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
219		471.06	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 6.96 (d, J = 2.5 Hz, 1H), 6.89 (d, J = 2.5 Hz, 1H), 6.03 (s, 1H), 5.00 (s, 1H), 4.79 (s, 1H), 3.92 (d, J = 9.3 Hz, 7H), 3.41 - 3.25 (m, 4H), 2.28 - 2.13 (m, 2H), 1.92 (d, J = 18.6 Hz, 6H)
220		474.07	(400 MHz, CDCl ₃) δ 8.69 (s, 1H), 8.62 (s, 1H), 8.54 (s, 1H), 8.21 (d, J = 4.9 Hz, 1H), 7.22 - 7.08 (m, 2H), 6.96 (s, 1H), 6.93 (s, 1H), 4.84 (s, 1H), 3.98 - 3.86 (m, 5H), 3.38 - 3.26 (m, 4H), 2.31 - 2.19 (m, 2H), 2.03 - 1.86 (m, 6H)
221		474.12	(400 MHz, CDCl ₃) δ 8.75 (d, J = 1.9 Hz, 1H), 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.39 (s, 1H), 8.05 (dd, J = 8.8, 2.3 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 6.49 (d, J = 8.8 Hz, 1H), 5.23 (s, 1H), 4.88 - 4.71 (m, 1H), 4.05 - 3.95 (m, 1H), 3.95 - 3.85 (m, 4H), 3.39 - 3.26 (m, 4H), 2.28 - 2.16 (m, 2H), 2.00 - 1.83 (m, 6H)
222		485.14	(CDCl ₃) δ 8.63 (d, J = 1.9 Hz, 1H), 8.53 (dd, J = 1.9, 0.6 Hz, 1H), 7.71 (d, J = 9.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 6.83 (d, J = 2.5 Hz, 1H), 6.68 (d, J = 9.4 Hz, 1H), 5.33 (s, 1H), 4.76 (dq, J = 5.0, 2.5 Hz, 1H), 4.14 (s, 1H), 3.93 - 3.77 (m, 4H), 3.28 (d, J = 5.1 Hz, 7H), 2.18 (dt, J = 13.4, 4.6 Hz, 2H), 2.02 - 1.75 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
223		435.18	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 6.94 (dd, J = 15.2, 2.5 Hz, 2H), 6.07 (s, 1H), 4.82 (dt, J = 5.8, 3.0 Hz, 1H), 4.02 - 3.83 (m, 4H), 3.77 - 3.57 (m, 1H), 3.43 - 3.28 (m, 4H), 2.52 (s, 3H), 2.37 (s, 3H), 2.30 - 2.16 (m, 2H), 2.05 - 1.78 (m, 6H)
224		475.02	(CDCl ₃) δ 8.72 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.27 (s, 1H), 6.98 (d, J = 2.4 Hz, 1H), 6.92 (d, J = 2.6 Hz, 1H), 6.44 (d, J = 6.0 Hz, 1H), 4.84 (s, 1H), 3.93 (dd, J = 6.0, 3.7 Hz, 4H), 3.43 - 3.25 (m, 4H), 2.23 (s, 2H), 2.08 - 1.83 (m, 6H)
225		451.16	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.49 (d, J = 1.1 Hz, 1H), 6.93 (dd, J = 16.7, 2.5 Hz, 2H), 6.45 (d, J = 1.2 Hz, 1H), 5.01 (s, 1H), 4.81 (s, 1H), 4.39 (d, J = 0.9 Hz, 2H), 3.97 - 3.87 (m, 4H), 3.49 (s, 3H), 3.39 - 3.29 (m, 4H), 2.22 (d, J = 9.4 Hz, 2H), 1.99 - 1.87 (m, 6H)
226		463.18	(CDCl ₃) δ 8.70 (d, J = 2.0 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 6.95 (d, J = 2.5 Hz, 1H), 6.91 (d, J = 2.5 Hz, 1H), 4.75 (d, J = 5.6 Hz, 1H), 4.66 (s, 1H), 4.32 (s, 1H), 4.00 - 3.83 (m, 4H), 3.43 - 3.22 (m, 4H), 2.46 (d, J = 15.1 Hz, 5H), 2.36 (s, 3H), 2.19 (q, J = 6.3, 3.9 Hz, 2H), 1.12 (t, J = 7.6 Hz, 3H)




Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
227		453.2	(CDCl ₃) δ 8.72 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.33 (d, J = 1.8 Hz, 1H), 7.04 - 6.87 (m, 2H), 5.15 (s, 1H), 4.82 (dq, J = 5.2, 2.6 Hz, 1H), 4.24 (dt, J = 8.3, 4.7 Hz, 1H), 4.03 - 3.86 (m, 4H), 3.44 - 3.28 (m, 4H), 2.74 (qd, J = 7.6, 2.3 Hz, 2H), 2.24 (dq, J = 9.6, 4.6 Hz, 2H), 2.09 - 1.85 (m, 6H), 1.29 (t, J = 7.6 Hz, 3H)
228		462.14	(CDCl ₃) δ 8.60 (dd, J = 8.1, 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 8.00 (s, 1H), 6.92 - 6.75 (m, 2H), 5.06 (d, J = 8.0 Hz, 1H), 4.70 (s, 1H), 3.94 (s, 1H), 3.89 - 3.76 (m, 4H), 3.75 - 3.62 (m, 4H), 3.34 - 3.17 (m, 4H), 2.50 (s, 3H), 2.11 (s, 2H), 1.82 (d, J = 5.0 Hz, 5H)
229		488.57	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.44 (d, J = 2.0 Hz, 1H), 7.91 (d, J = 8.8 Hz, 1H), 6.97 (d, J = 2.4 Hz, 1H), 6.92 (d, J = 2.4 Hz, 1H), 6.60 (d, J = 8.6 Hz, 1H), 5.52 (s, 1H), 4.83 (s, 1H), 4.18 (s, 3H), 4.00 - 3.87 (m, 5H), 3.39 - 3.24 (m, 4H), 2.31 - 2.18 (m, 2H), 2.04 - 1.87 (m, 6H)
230		488.57	(400 MHz, CDCl ₃) δ 8.81 (d, J = 1.8 Hz, 1H), 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.13 (dd, J = 8.8, 2.2 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.92 (d, J = 2.4 Hz, 1H), 6.52 (d, J = 8.8 Hz, 1H), 4.82 (s, 1H), 4.36 (s, 3H), 3.96 - 3.85 (m, 5H), 3.41 - 3.28 (m, 4H), 2.28 - 2.17 (m, 2H), 2.06 - 1.83 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
231		510.2	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 2.0 Hz, 1H), 7.70 (d, J = 3.4 Hz, 1H), 6.93 (dd, J = 14.6, 2.5 Hz, 2H), 4.96 (dd, J = 17.8, 7.1 Hz, 2H), 4.76 (d, J = 6.2 Hz, 1H), 4.62 (ddt, J = 12.6, 7.1, 3.5 Hz, 1H), 4.05 - 3.82 (m, 8H), 3.75 (dd, J = 9.4, 3.2 Hz, 1H), 3.40 - 3.27 (m, 4H), 2.35 (ddt, J = 13.0, 8.2, 7.1 Hz, 1H), 2.25 - 2.08 (m, 2H), 2.00 - 1.77 (m, 6H)
232		477.18	(CDCl ₃) δ 8.61 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 7.98 (s, 2H), 6.94 - 6.75 (m, 2H), 5.01 (s, 1H), 4.71 (d, J = 5.9 Hz, 1H), 3.96 - 3.78 (m, 5H), 3.68 (d, J = 7.0 Hz, 2H), 3.37 - 3.17 (m, 4H), 2.21 - 1.96 (m, 2H), 1.92 - 1.75 (m, 6H), 1.25 - 1.06 (m, 1H), 0.66 - 0.44 (m, 2H), 0.34 - 0.18 (m, 2H)
233		451.2	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.06 (s, 2H), 7.01 - 6.86 (m, 2H), 5.00 (d, J = 8.1 Hz, 1H), 4.80 (d, J = 5.6 Hz, 1H), 4.10 - 3.82 (m, 7H), 3.42 - 3.26 (m, 5H), 2.20 (d, J = 8.2 Hz, 2H), 2.01 - 1.80 (m, 6H), 1.39 (t, J = 7.0 Hz, 3H)
234		432.17	(CDCl ₃) δ 8.72 (d, J = 1.9 Hz, 1H), 8.60 (t, J = 1.9 Hz, 2H), 6.97 (d, J = 2.5 Hz, 1H), 6.91 (d, J = 2.5 Hz, 1H), 6.70 (d, J = 1.2 Hz, 1H), 4.85 (d, J = 4.9 Hz, 1H), 3.98 - 3.84 (m, 4H), 3.42 - 3.26 (m, 4H), 2.33 - 2.17 (m, 2H), 2.01 - 1.77 (m, 5H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
235		505.04	(CDCl ₃) δ 8.68 (d, J = 2.0 Hz, 1H), 8.59 (d, J = 1.9 Hz, 1H), 8.36 (s, 2H), 6.97 - 6.86 (m, 2H), 5.53 (d, J = 8.1 Hz, 1H), 4.96 - 4.72 (m, 3H), 4.09 - 3.86 (m, 5H), 3.33 (dd, J = 5.8, 4.0 Hz, 4H), 2.28 - 2.10 (m, 2H), 1.98 - 1.78 (m, 6H)
236		505.17	(CDCl ₃) δ 8.68 (d, J = 2.0 Hz, 1H), 8.59 (d, J = 1.9 Hz, 1H), 8.36 (s, 2H), 6.97 - 6.86 (m, 2H), 5.53 (d, J = 8.1 Hz, 1H), 4.96 - 4.72 (m, 3H), 4.09 - 3.86 (m, 5H), 3.33 (dd, J = 5.8, 4.0 Hz, 4H), 2.28 - 2.10 (m, 2H), 1.98 - 1.78 (m, 6H)
237		451.16	(CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.31 (s, 2H), 6.99 - 6.87 (m, 2H), 5.34 (d, J = 8.1 Hz, 1H), 4.87 - 4.73 (m, 2H), 4.06 - 3.87 (m, 6H), 3.38 - 3.29 (m, 4H), 2.20 (q, J = 5.8 Hz, 2H), 2.04 - 1.84 (m, 6H), 1.51 (d, J = 6.5 Hz, 3H)
238		492.98	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.15 (s, 2H), 6.99 - 6.86 (m, 2H), 5.11 (td, J = 7.9, 7.4, 3.8 Hz, 1H), 4.82 - 4.55 (m, 3H), 4.10 (dd, J = 4.1, 2.7 Hz, 2H), 3.93 (dd, J = 6.0, 3.7 Hz, 5H), 3.60 - 3.43 (m, 1H), 3.36 (dd, J = 6.0, 3.7 Hz, 4H), 2.85 - 2.61 (m, 1H), 2.30 - 2.10 (m, 2H), 1.92 (s, 6H), 1.24 (q, J = 6.9 Hz, 1H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
239		507	(CDCl ₃) δ 8.61 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 7.95 (d, J = 13.9 Hz, 2H), 6.92 - 6.72 (m, 2H), 4.85 - 4.64 (m, 3H), 4.41 (t, J = 6.2 Hz, 2H), 3.83 (q, J = 6.7, 5.7 Hz, 6H), 3.33 - 3.18 (m, 5H), 2.20 - 1.97 (m, 1H), 1.82 (s, 7H)
240		493.16	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.08 (s, 2H), 7.01 - 6.87 (m, 2H), 5.11 (d, J = 8.1 Hz, 1H), 4.95 - 4.74 (m, 3H), 4.56 (t, J = 6.0 Hz, 2H), 4.17 (d, J = 6.7 Hz, 2H), 4.05 - 3.81 (m, 4H), 3.51 - 3.29 (m, 5H), 2.21 (q, J = 6.4, 5.7 Hz, 2H), 2.03 - 1.80 (m, 8H)
241		445.54	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.52 (d, J = 8.7 Hz, 1H), 6.96 (d, J = 2.5 Hz, 1H), 6.90 (d, J = 2.5 Hz, 1H), 6.23 (d, J = 8.7 Hz, 1H), 5.09 (s, 1H), 4.80 (s, 1H), 4.00 - 3.78 (m, 5H), 3.40 - 3.24 (m, 4H), 2.56 (s, 3H), 2.29 - 2.14 (m, 2H), 2.01 - 1.80 (m, 6H)
242		445.54	(400 MHz, CDCl ₃) δ 8.70 (d, J = 1.8 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.30 (d, J = 1.8 Hz, 1H), 7.37 (s, 1H), 6.96 (d, J = 2.3 Hz, 1H), 6.91 (d, J = 2.3 Hz, 1H), 4.79 (s, 1H), 4.74 (s, 1H), 4.28 (s, 1H), 4.01 - 3.83 (m, 4H), 3.40 - 3.25 (m, 4H), 2.30 - 2.17 (m, 2H), 2.11 (s, 3H), 1.99 - 1.87 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
243		465.2	(CDCl ₃) δ 8.69 (d, J = 2.0 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 5.36 (s, 1H), 4.03 (s, 1H), 1.57 (s, 6H), 2.02 - 1.80 (m, 6H), 8.42 (s, 2H), 6.99 - 6.87 (m, 2H), 4.80 (s, 1H), 3.92 (dd, J = 6.0, 3.7 Hz, 4H), 3.39 - 3.29 (m, 4H), 2.20 (d, J = 9.0 Hz, 2H)
244		475.15	(CDCl ₃) δ 8.98 - 8.86 (m, 2H), 8.66 (dd, J = 23.8, 1.9 Hz, 2H), 8.44 (s, 1H), 7.00 - 6.88 (m, 2H), 5.89 (d, J = 8.1 Hz, 1H), 4.83 (dq, J = 5.4, 2.6 Hz, 1H), 4.22 - 4.04 (m, 1H), 3.97 - 3.87 (m, 4H), 3.43 - 3.29 (m, 4H), 2.24 (td, J = 8.3, 7.4, 4.0 Hz, 2H), 2.15 - 1.81 (m, 6H)
245		467.14	(CDCl ₃) δ 8.61 (d, J = 1.9 Hz, 1H), 8.53 (d, J = 1.9 Hz, 1H), 6.84 (dd, J = 18.2, 2.5 Hz, 2H), 5.26 (s, 1H), 4.78 (d, J = 8.0 Hz, 1H), 4.68 (d, J = 5.6 Hz, 1H), 3.82 (t, J = 4.0 Hz, 10H), 3.30 - 3.15 (m, 4H), 2.10 (q, J = 6.2, 5.7 Hz, 2H), 1.94 - 1.69 (m, 6H)
246		421.23	(CDCl ₃) δ 8.61 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 8.06 (d, J = 5.0 Hz, 1H), 6.94 - 6.78 (m, 2H), 6.32 (d, J = 5.0 Hz, 1H), 5.09 (s, 1H), 4.70 (d, J = 6.0 Hz, 1H), 3.95 (s, 1H), 3.91 - 3.77 (m, 5H), 3.37 - 3.13 (m, 4H), 2.20 - 2.00 (m, 2H), 1.94 - 1.71 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
247		465.1	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.64 (t, J = 2.4 Hz, 1H), 8.05 (s, 2H), 7.04 - 6.87 (m, 2H), 5.20 (s, 1H), 4.80 (s, 1H), 4.30 (p, J = 6.1 Hz, 1H), 4.05 - 3.81 (m, 4H), 3.44 - 3.27 (m, 4H), 2.22 (t, J = 7.3 Hz, 2H), 1.93 (d, J = 4.6 Hz, 6H), 1.33 (d, J = 6.1 Hz, 6H)
248		488.48	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.66 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.05 (d, J = 8.5 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 6.51 (d, J = 8.7 Hz, 1H), 4.82 (s, 1H), 3.99 - 3.85 (m, 5H), 3.38 - 3.27 (m, 4H), 2.59 (s, 3H), 2.29 - 2.18 (m, 2H), 2.01 - 1.84 (m, 6H)
249		472.54	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.45 (d, J = 1.9 Hz, 1H), 7.86 (d, J = 7.7 Hz, 1H), 7.60 (d, J = 2.3 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.92 (d, J = 2.4 Hz, 1H), 6.50 (d, J = 2.3 Hz, 1H), 6.48 (d, J = 8.9 Hz, 1H), 4.81 (s, 1H), 3.95 - 3.85 (m, 5H), 3.39 - 3.30 (m, 4H), 2.26 - 2.17 (m, 2H), 1.99 - 1.84 (m, 6H)

Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
250		486.5	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.09 (d, J = 1.7 Hz, 1H), 7.56 - 7.45 (m, 2H), 6.96 (d, J = 2.4 Hz, 1H), 6.93 (d, J = 2.4 Hz, 1H), 6.55 (d, J = 8.5 Hz, 1H), 6.25 (d, J = 1.9 Hz, 1H), 4.84 (s, 1H), 3.96 - 3.83 (m, 8H), 3.39 - 3.30 (m, 4H), 2.28 - 2.19 (m, 2H), 2.03 - 1.83 (m, 6H)
251		491.1	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.18 (s, 2H), 7.04 - 6.87 (m, 2H), 5.21 (d, J = 8.2 Hz, 1H), 4.80 (s, 1H), 4.2-4.0 (m, 3H), 3.98 - 3.85 (m, 4H), 3.61 - 3.44 (m, 2H), 3.41 - 3.21 (m, 4H), 2.75 - 2.47 (m, 1H), 2.21 (d, J = 9.6 Hz, 2H), 1.92 (t, J = 6.4 Hz, 6H), 1.83 - 1.66 (m, 4H)
252		471.1	(CDCl ₃) δ 3.97 - 3.87 (m, 4H), 3.39 - 3.29 (m, 4H), 2.50 (s, 3H), 2.28 - 2.16 (m, 2H), 2.02 - 1.89 (m, 6H), 4.26 - 4.17 (m, 1H), 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.88 (d, J = 3.3 Hz, 1H), 6.93 (dd, J = 18.7, 2.5 Hz, 2H), 5.10 (d, J = 7.9 Hz, 1H), 4.79 (d, J = 3.7 Hz, 1H)
253		453.19	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 2.0 Hz, 1H), 8.38 (d, J = 1.1 Hz, 1H), 6.93 (dd, J = 17.3, 2.5 Hz, 2H), 6.18 (d, J = 1.2 Hz, 1H), 5.03 (s, 1H), 4.81 (d, J = 6.1 Hz, 1H), 3.97 - 3.87 (m, 4H), 3.83 (s, 1H), 3.39 - 3.29 (m, 4H), 2.51 (s, 3H), 2.21 (d, J = 8.8 Hz, 2H), 2.05 (s, 0H), 1.91 (d, J = 10.6 Hz, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
254		406.57	(400 MHz, CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.15 - 8.04 (m, 1H), 7.46 - 7.39 (m, 1H), 6.97 (d, J = 2.5 Hz, 1H), 6.93 (d, J = 2.5 Hz, 1H), 6.60 - 6.52 (m, 1H), 6.41 (d, J = 8.4 Hz, 1H), 4.80 (s, 1H), 4.68 (s, 1H), 3.98 - 3.85 (m, 5H), 3.41 - 3.30 (m, 4H), 2.27 - 2.15 (m, 2H), 1.97 - 1.86 (m, 6H)
255		450.96	(methanol-d ₄) δ 8.71 (dd, J = 7.5, 2.1 Hz, 1H), 8.57 (dd, J = 8.8, 2.1 Hz, 1H), 7.12 (d, J = 2.5 Hz, 1H), 6.88 (d, J = 2.4 Hz, 1H), 3.88 (t, J = 4.8 Hz, 4H), 3.45 - 3.35 (m, 4H), 3.19 (s, 3H), 2.25 - 2.01 (m, 2H), 2.00 - 1.80 (m, 6H)
256		420.1	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 7.80 (d, J = 1.5 Hz, 1H), 7.47 (d, J = 1.5 Hz, 1H), 7.02 - 6.86 (m, 2H), 4.80 (s, 1H), 4.27 (d, J = 8.3 Hz, 1H), 4.00 - 3.72 (m, 7H), 3.35 (dd, J = 6.0, 3.8 Hz, 4H), 2.23 (s, 2H), 2.02 - 1.80 (m, 6H)
257		437.1	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (dd, J = 1.9, 0.7 Hz, 1H), 7.99 - 7.88 (m, 1H), 7.02 - 6.87 (m, 2H), 6.34 (d, J = 8.4 Hz, 1H), 4.78 (s, 1H), 4.40 (d, J = 8.2 Hz, 1H), 4.03 - 3.75 (m, 5H), 3.35 (dd, J = 6.0, 3.7 Hz, 4H), 2.18 (s, 5H), 2.03 - 1.81 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
258		488.52	(CDCl ₃) δ 8.78 - 8.59 (m, 3H), 8.38 (s, 1H), 7.85 (s, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.3 Hz, 1H), 4.79 (s, 1H), 4.67 (d, J = 7.6 Hz, 1H), 4.32 (s, 1H), 3.99 - 3.82 (m, 4H), 3.40 - 3.27 (m, 4H), 2.33 - 2.06 (m, 5H), 2.03 - 1.87 (m, 6H)
259		488.48	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, 1H), 8.41 (s, 1H), 8.04 - 7.93 (m, 1H), 6.96 (d, J = 2.5 Hz, 1H), 6.91 (d, J = 2.1 Hz, 1H), 6.35 (d, J = 9.1 Hz, 1H), 5.07 (s, 1H), 4.82 (s, 1H), 4.02 - 3.79 (m, 5H), 3.41 - 3.27 (m, 4H), 2.83 - 2.72 (m, 3H), 2.30 - 2.17 (m, 2H), 2.02 - 1.80 (m, 6H)
260		437.24	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 6.93 (dd, J = 14.5, 2.5 Hz, 2H), 6.79 (d, J = 9.4 Hz, 1H), 6.64 (d, J = 9.4 Hz, 1H), 4.80 (d, J = 5.5 Hz, 1H), 4.27 (d, J = 7.5 Hz, 1H), 4.14 (s, 1H), 4.02 (s, 3H), 3.98 - 3.88 (m, 4H), 3.42 - 3.29 (m, 4H), 2.31 - 2.12 (m, 2H), 1.95 (ddd, J = 17.2, 9.2, 6.0 Hz, 6H)
261		477.3	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.20 (s, 2H), 7.04 - 6.84 (m, 2H), 5.16 (d, J = 8.0 Hz, 1H), 4.80 (s, 1H), 4.19 - 3.85 (m, 7H), 3.66 (dd, J = 8.5, 7.1 Hz, 1H), 3.35 (dd, J = 5.9, 3.8 Hz, 4H), 3.24 (t, J = 7.5 Hz, 1H), 2.48 - 2.11 (m, 4H), 2.05 - 1.82 (m, 6H)





Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
262		476.26	(CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.61 (d, J = 1.0 Hz, 1H), 6.98 - 6.86 (m, 2H), 4.96 (s, 1H), 4.82 - 4.71 (m, 1H), 4.63 - 4.56 (m, 1H), 4.08 - 3.87 (m, 5H), 3.38 - 3.29 (m, 4H), 2.80 (tdd, J = 6.8, 5.0, 3.1 Hz, 1H), 2.23 - 2.11 (m, 2H), 2.08 - 1.82 (m, 9H), 0.90 - 0.72 (m, 2H), 0.59 - 0.47 (m, 2H)
263		480.22	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.11 (d, J = 0.9 Hz, 1H), 6.93 (dd, J = 15.5, 2.5 Hz, 2H), 5.30 (d, J = 1.0 Hz, 1H), 5.16 (s, 1H), 4.95 (s, 1H), -0.17 - -0.23 (m, 0H), 4.80 (s, 1H), 3.92 (dd, J = 5.9, 3.7 Hz, 4H), 3.74 (s, 1H), 3.57 (dd, J = 5.6, 4.6 Hz, 2H), 3.50 - 3.29 (m, 9H), 2.20 (d, J = 9.1 Hz, 2H), 1.93 (d, J = 13.1 Hz, 6H)
264		494.1	(CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 2.0 Hz, 1H), 7.76 (d, J = 5.8 Hz, 1H), 7.02 - 6.86 (m, 2H), 5.66 (d, J = 5.9 Hz, 1H), 4.80 (d, J = 5.5 Hz, 1H), 4.63 (s, 1H), 3.97 - 3.84 (m, 5H), 1.30 - 1.20 (m, 1H), 3.64 (s, 2H), 3.38 - 3.28 (m, 4H), 1.37 (s, 6H), 5.16 - 4.86 (m, 1H), 2.24 - 2.13 (m, 2H), 1.96 - 1.82 (m, 6H)
265		421.18	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.47 (d, J = 1.1 Hz, 1H), 6.93 (dd, J = 16.7, 2.5 Hz, 2H), 6.17 (t, J = 0.9 Hz, 1H), 4.95 (s, 1H), 4.81 (td, J = 5.3, 2.5 Hz, 1H), 4.07 - 3.83 (m, 5H), 3.39 - 3.29 (m, 4H), 2.34 (s, 3H), 2.21 (dt, J = 11.1, 5.1 Hz, 2H), 2.04 - 1.76 (m, 6H)

Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
266		441.2	(CDCl ₃) δ 11.49 (s, 1H), 8.69 (d, J = 1.9 Hz, 1H), 8.59 (d, J = 1.9 Hz, 1H), 7.73 (d, J = 3.0 Hz, 1H), 7.03 - 6.89 (m, 2H), 6.56 (d, J = 7.2 Hz, 1H), 4.82 (s, 1H), 3.91 (dd, J = 6.0, 3.8 Hz, 5H), 3.39 - 3.29 (m, 4H), 2.27 - 2.14 (m, 2H), 2.07 - 1.81 (m, 6H)
267		447.11	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 8.03 (d, J = 6.1 Hz, 1H), 6.93 (dd, J = 18.6, 2.5 Hz, 2H), 6.13 (d, J = 6.1 Hz, 1H), 5.09 (s, 1H), 4.79 (s, 1H), 4.03 - 3.78 (m, 5H), 3.34 (dd, J = 6.0, 3.8 Hz, 4H), 2.20 (d, J = 8.1 Hz, 2H), 1.96 (s, 7H), 1.10 (d, J = 2.8 Hz, 2H), 1.00 (d, J = 8.0 Hz, 2H)
268		437.19	(CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 7.95 (d, J = 5.8 Hz, 1H), 6.92 (dd, J = 17.9, 2.5 Hz, 2H), 5.99 (d, J = 5.9 Hz, 1H), 5.01 (s, 1H), 4.79 (dt, J = 6.9, 3.4 Hz, 1H), 3.91 (d, J = 8.7 Hz, 8H), 3.43 - 3.25 (m, 4H), 2.32 - 2.09 (m, 2H), 2.05 - 1.74 (m, 6H)
269		468.13	(CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.70 (d, J = 6.8 Hz, 1H), 6.92 (dd, J = 14.4, 2.5 Hz, 2H), 4.86 (d, J = 8.1 Hz, 1H), 4.75 (dt, J = 8.6, 4.0 Hz, 1H), 3.97 - 3.87 (m, 5H), 3.38 - 3.28 (m, 4H), 3.14 (d, J = 2.2 Hz, 6H), 2.24 - 2.07 (m, 2H), 2.02 - 1.79 (m, 6H)

Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
270	 <chem>CC(C)(C)OC(=O)N1CCCCC1[C@H]2CCCC[C@@H]2Oc3ccc(Nc4ccnnc4)cc3</chem>		(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 2.0 Hz, 1H), 8.15 (s, 2H), 7.01 - 6.87 (m, 2H), 5.13 (d, J = 8.1 Hz, 1H), 4.80 (s, 1H), 4.24 (s, 1H), 4.01 (s, 2H), 3.93 (dd, J = 6.0, 3.7 Hz, 4H), 3.35 (dd, J = 5.9, 3.9 Hz, 4H), 2.80 (t, J = 12.7 Hz, 2H), 2.51 (t, J = 12.1 Hz, 1H), 2.21 (d, J = 8.3 Hz, 2H), 2.05 - 1.72 (m, 8H), 1.49 (s, 9H)
271	 <chem>c1ccc(Nc2ccnnc2)[C@H]3CCCC[C@@H]3Oc4ccc(Nc5ccnnc5)cc4</chem>	490.2	(CDCl ₃) δ 8.62 (d, J = 1.9 Hz, 1H), 8.55 (d, J = 1.9 Hz, 1H), 8.09 (s, 2H), 6.93 - 6.78 (m, 2H), 5.03 (d, J = 8.1 Hz, 1H), 4.72 (d, J = 5.8 Hz, 1H), 4.02 - 3.75 (m, 5H), 3.34 - 3.21 (m, 4H), 3.20 - 3.06 (m, 2H), 2.66 (td, J = 12.1, 2.5 Hz, 2H), 2.38 (ddt, J = 12.2, 7.6, 3.8 Hz, 1H), 2.12 (d, J = 8.9 Hz, 2H), 1.96 - 1.65 (m, 10H), 1.53 (qd, J = 12.3, 3.9 Hz, 2H)
272	 <chem>c1ccc(Nc2ccnnc2)[C@H]3CCCC[C@@H]3Oc4ccc(Nc5ccnnc5)cc4</chem>	406.53	(CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.04 (d, J = 2.8 Hz, 1H), 7.94 (dd, J = 4.7, 1.3 Hz, 1H), 7.11 (dd, J = 8.3, 4.7 Hz, 1H), 7.00 - 6.84 (m, 3H), 4.78 (s, 1H), 3.98 - 3.88 (m, 4H), 3.85 (d, J = 8.1 Hz, 1H), 3.38 - 3.28 (m, 4H), 2.27 - 2.15 (m, 2H), 1.97 - 1.84 (m, 6H)

Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
273		406.57	(CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.18 (d, J = 6.3 Hz, 2H), 6.96 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 6.51 - 6.42 (m, 2H), 4.79 (s, 1H), 4.47 (d, J = 7.7 Hz, 1H), 3.98 - 3.86 (m, 4H), 3.56 (s, 1H), 3.39 - 3.26 (m, 4H), 2.27 - 2.15 (m, 2H), 2.01 - 1.79 (m, 6H)
274		504.1	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.18 (s, 2H), 7.03 - 6.86 (m, 2H), 5.11 (d, J = 8.1 Hz, 1H), 4.81 (d, J = 5.9 Hz, 1H), 4.12 - 3.84 (m, 5H), 3.42 - 3.29 (m, 4H), 3.05 - 2.89 (m, 2H), 2.33 (s, 4H), 2.21 (d, J = 8.8 Hz, 2H), 2.11 - 1.61 (m, 10H)
275		519.2	(CDCl ₃) δ 8.62 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 6.93 - 6.77 (m, 2H), 5.20 (s, 1H), 4.78 - 4.60 (m, 2H), 3.92 - 3.79 (m, 5H), 3.64 (s, 0H), 3.51 (t, J = 5.1 Hz, 4H), 3.36 - 3.16 (m, 4H), 2.40 (t, J = 5.1 Hz, 4H), 2.27 (d, J = 4.8 Hz, 6H), 2.16 - 2.02 (m, 2H), 1.81 (q, J = 8.1, 5.7 Hz, 6H)
276		451.53	(CDCl ₃) δ 8.69 (d, J = 1.8 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 6.94 (d, J = 2.3 Hz, 1H), 6.90 (d, J = 2.3 Hz, 1H), 5.87 (s, 1H), 5.13 (s, 1H), 4.76 (s, 1H), 4.04 (s, 1H), 3.98 - 3.89 (m, 4H), 3.87 (s, 3H), 3.41 - 3.23 (m, 4H), 2.25 (s, 3H), 2.23 - 2.09 (m, 2H), 1.99 - 1.80 (m, 6H)

Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
277		459.08	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.17 (d, J = 0.9 Hz, 1H), 6.94 (dd, J = 15.8, 2.5 Hz, 2H), 5.31 (s, 1H), 4.83 (dp, J = 4.6, 2.5 Hz, 1H), 4.22 (qt, J = 8.5, 4.9 Hz, 1H), 4.03 - 3.87 (m, 4H), 3.39 - 3.29 (m, 4H), 2.25 (td, J = 7.9, 6.6, 3.9 Hz, 2H), 2.09 - 1.80 (m, 6H)
278		488.48	(400 MHz, CDCl ₃) δ 8.83 (d, J = 2.2 Hz, 1H), 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.02 (dd, J = 8.8, 2.3 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.5 Hz, 1H), 6.45 (d, J = 8.9 Hz, 1H), 5.12 (s, 1H), 4.81 (d, J = 2.6 Hz, 1H), 3.98 (s, 1H), 3.95 - 3.87 (m, 4H), 3.40 - 3.26 (m, 5H), 2.42 (s, 3H), 2.29 - 2.16 (m, 2H), 1.99 - 1.87 (m, 6H)
279		451.12	(CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 2.0 Hz, 1H), 7.21 - 7.11 (m, 1H), 7.00 - 6.85 (m, 2H), 5.51 (s, 1H), 4.95 - 4.70 (m, 2H), 4.53 - 4.34 (m, 1H), 4.01 - 3.74 (m, 5H), 3.45 - 3.25 (m, 4H), 2.25 - 2.09 (m, 2H), 2.03 - 1.73 (m, 6H), 1.39 - 1.25 (m, 3H)
280		423.08	(methanol-d ₄) δ 8.68 (d, J = 2.0 Hz, 1H), 8.54 (d, J = 2.0 Hz, 1H), 7.30 (d, J = 7.2 Hz, 1H), 7.10 (d, J = 2.4 Hz, 1H), 6.87 (d, J = 2.4 Hz, 1H), 5.79 (d, J = 7.2 Hz, 1H), 4.13 (qd, J = 8.5, 6.7, 2.4 Hz, 1H), 3.87 (dd, J = 5.9, 3.8 Hz, 4H), 3.42 - 3.32 (m, 5H), 2.21 - 2.07 (m, 2H), 2.01 - 1.78 (m, 6H)

Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
281		464.53	(400 MHz, CDCl ₃) δ 8.68 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.18 (d, J = 2.2 Hz, 1H), 7.58 (dd, J = 8.7, 2.5 Hz, 1H), 6.95 (d, J = 2.5 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 6.39 (d, J = 8.8 Hz, 1H), 4.78 (s, 1H), 4.69 (s, 1H), 3.96 - 3.83 (m, 5H), 3.49 (s, 1H), 3.38 - 3.28 (m, 4H), 2.26 - 2.13 (m, 2H), 1.95 - 1.84 (m, 6H), 1.56 (s, 6H)
282		421.2	(CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.93 - 7.74 (m, 2H), 7.04 - 6.82 (m, 2H), 4.80 (s, 1H), 4.50 (d, J = 8.1 Hz, 1H), 3.93 (dd, J = 6.4, 3.4 Hz, 5H), 3.41 - 3.25 (m, 4H), 2.51 - 2.33 (m, 3H), 2.23 (s, 2H), 2.04 - 1.82 (m, 6H)
283		420.2	(CDCl ₃) δ 8.62 (d, J = 1.9 Hz, 1H), 8.55 (d, J = 1.9 Hz, 1H), 7.88 (dd, J = 5.3, 0.7 Hz, 1H), 6.95 - 6.79 (m, 2H), 6.32 (ddd, J = 5.2, 1.5, 0.7 Hz, 1H), 6.12 (dt, J = 1.6, 0.8 Hz, 1H), 4.71 (d, J = 5.9 Hz, 1H), 4.40 (d, J = 8.2 Hz, 1H), 3.98 - 3.67 (m, 4H), 3.35 - 3.17 (m, 4H), 2.15 (s, 4H), 1.83 (d, J = 5.2 Hz, 5H)
284		426.1	(CDCl ₃) δ 8.62 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 6.86 (dd, J = 16.6, 2.5 Hz, 2H), 5.97 (d, J = 1.2 Hz, 1H), 5.02 (s, 1H), 4.71 (dt, J = 5.7, 3.0 Hz, 1H), 3.91 - 3.75 (m, 4H), 3.63 - 3.42 (m, 1H), 3.33 - 3.17 (m, 4H), 2.21 - 2.03 (m, 5H), 1.94 - 1.71 (m, 6H)















Cmpd. No.	Compound Structure	ESMS (M+H)	¹ H NMR (300 MHz, unless indicated otherwise) NMR peaks given as δ values
285		420.57	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.08 (d, J = 5.7 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 6.30 (s, 1H), 6.28 (dd, J = 5.7, 2.2 Hz, 1H), 4.78 (s, 1H), 4.19 (d, J = 7.7 Hz, 1H), 3.97 - 3.82 (m, 4H), 3.53 (s, 1H), 3.40 - 3.24 (m, 4H), 2.41 (s, 3H), 2.27 - 2.11 (m, 2H), 1.92 - 1.84 (m, 6H)
286		434.56	(400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 6.15 (s, 2H), 4.77 (s, 1H), 4.12 (d, J = 7.8 Hz, 1H), 3.99 - 3.81 (m, 4H), 3.53 (s, 1H), 3.40 - 3.23 (m, 4H), 2.36 (d, J = 14.5 Hz, 6H), 2.25 - 2.14 (m, 2H), 1.89 (t, J = 7.8 Hz, 6H)




Table 2.




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
287		478.3	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.33 (s, 1H), 6.94 (dd, J = 14.3, 2.5 Hz, 2H), 5.80 (d, J = 4.9 Hz, 1H), 5.42 (d, J = 8.1 Hz, 1H), 4.88 - 4.74 (m, 1H), 4.08 (s, 2H), 3.99 - 3.86 (m, 4H), 3.50 (s, 3H), 3.42 - 3.28 (m, 4H), 3.01 (dd, J = 18.4, 5.0 Hz, 3H), 2.54 (s, 3H), 2.28 - 2.08 (m, 2H), 2.03 - 1.79 (m, 6H).
288		404	¹ H NMR (400 MHz, CDCl ₃) δ 8.68 (t, J = 5.2 Hz, 1H), 8.65 (d, J = 1.7 Hz, 1H), 8.02 (d, J = 5.9 Hz, 1H), 7.26 (d, J = 1.5 Hz, 1H), 7.08 (d, J = 1.6 Hz, 1H), 7.00 (d, J = 1.6 Hz, 1H), 6.07 (d, J = 6.0 Hz, 1H), 4.91 (s, 1H), 4.75 (s, 1H), 4.54 (t, J = 9.6 Hz, 2H), 3.03 (td, J = 9.8, 1.6 Hz, 2H), 2.42 (s, 3H), 2.21 - 2.06 (m, 2H), 1.91 - 1.74 (m, 7H).
289		404	¹ H NMR (400 MHz, CDCl ₃) δ 8.79 - 8.71 (m, 2H), 8.03 (d, J = 5.8 Hz, 1H), 7.39 (s, 1H), 6.38 (s, 1H), 6.08 (d, J = 5.9 Hz, 1H), 5.06 (s, 2H), 4.83 (d, J = 29.0 Hz, 4H), 2.42 (s, 3H), 2.16 (d, J = 7.2 Hz, 2H), 1.88 (dd, J = 21.2, 8.7 Hz, 6H), 1.58 (s, 2H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
290		435	¹ H NMR (400 MHz, CDCl ₃) δ 8.62 (d, J = 1.9 Hz, 1H), 8.53 (d, J = 1.9 Hz, 1H), 8.03 (d, J = 5.9 Hz, 1H), 6.88 (td, J = 19.2, 3.6 Hz, 2H), 6.07 (d, J = 6.0 Hz, 1H), 4.87 (s, 1H), 4.74 (s, 1H), 4.06 - 3.93 (m, 1H), 3.80 - 3.69 (m, 2H), 3.58 - 3.38 (m, 2H), 2.94 (td, J = 11.9, 3.5 Hz, 1H), 2.65 - 2.52 (m, 1H), 2.42 (s, 3H), 2.13 (d, J = 10.1 Hz, 2H), 1.81 (d, J = 9.7 Hz, 6H), 1.32 - 1.08 (m, 3H).
291		419.23 [1]	¹ H NMR (400 MHz, Chloroform-d) δ 8.42 (d, J = 1.9 Hz, 1H), 8.30 (d, J = 1.9 Hz, 1H), 8.04 (d, J = 4.8 Hz, 2H), 6.65 - 6.56 (m, 2H), 6.28 (t, J = 4.8 Hz, 1H), 4.99 (d, J = 8.1 Hz, 1H), 4.60 (d, J = 6.5 Hz, 3H), 3.79 (dd, J = 8.2, 4.0 Hz, 0H), 3.62 - 3.38 (m, 4H), 3.17 - 3.03 (m, 1H), 2.07 - 1.90 (m, 2H), 1.89 - 1.59 (m, 7H).
292		413	¹ H NMR (400 MHz, CDCl ₃) δ 8.83 (dd, J = 2.3, 0.8 Hz, 1H), 8.77 - 8.68 (m, 2H), 8.54 (dd, J = 4.8, 1.6 Hz, 1H), 7.95 (d, J = 6.0 Hz, 1H), 7.86 (ddd, J = 7.9, 2.4, 1.6 Hz, 1H), 7.76 (d, J = 1.7 Hz, 1H), 7.31 (ddd, J = 7.9, 4.8, 0.8 Hz, 1H), 7.21 (d, J = 1.6 Hz, 1H), 7.11 (d, J = 0.9 Hz, 1H), 6.00 (d, J = 6.0 Hz, 1H), 4.80 (d, J = 27.2 Hz, 2H), 2.34 (s, 3H), 2.18 - 2.04 (m, 2H), 1.84 - 1.71 (m, 6H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
293		402	¹ H NMR (400 MHz, CDCl ₃) δ 11.20 (s, 1H), 8.74 (t, J = 6.1 Hz, 1H), 8.71 (d, J = 1.8 Hz, 1H), 8.03 (d, J = 5.9 Hz, 1H), 7.96 (s, 2H), 7.75 (t, J = 8.0 Hz, 1H), 7.26 (d, J = 1.5 Hz, 1H), 6.08 (d, J = 6.0 Hz, 1H), 5.09 (s, 1H), 4.82 (s, 1H), 3.77 (s, 1H), 2.43 (s, 3H), 2.24 - 2.08 (m, 2H), 1.93 - 1.76 (m, 6H).
294		410.35	¹ H NMR (300 MHz, Chloroform-d) δ 8.10 (d, J = 6.0 Hz, 1H), 7.36 - 7.28 (m, 1H), 6.65 (d, J = 2.0 Hz, 1H), 6.13 (d, J = 6.0 Hz, 1H), 4.81 (d, J = 8.0 Hz, 1H), 4.00 - 3.80 (m, 6H), 3.80 - 3.56 (m, 1H), 3.29 (dd, J = 5.8, 3.8 Hz, 4H), 3.02 - 2.84 (m, 2H), 2.48 (s, 3H), 2.36 (td, J = 11.4, 10.9, 2.5 Hz, 2H), 2.14 - 1.98 (m, 2H), 1.70 - 1.46 (m, 2H).
295		450.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.98 - 8.78 (m, 2H), 8.33 (d, J = 1.9 Hz, 1H), 7.67 (d, J = 1.8 Hz, 1H), 7.30 (d, J = 1.8 Hz, 1H), 6.49 - 6.31 (m, 1H), 5.07 (d, J = 7.6 Hz, 1H), 4.87 (d, J = 5.5 Hz, 1H), 4.43 (q, J = 2.8 Hz, 2H), 4.24 (s, 1H), 4.02 (t, J = 5.4 Hz, 2H), 2.71 (dtd, J = 13.1, 7.7, 2.7 Hz, 4H), 2.26 (dt, J = 10.4, 5.3 Hz, 2H), 1.97 (d, J = 5.3 Hz, 6H), 1.28 (t, J = 7.6 Hz, 3H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
296		550.6	¹ H NMR (300 MHz, Chloroform-d) δ 8.79 - 8.67 (m, 1H), 8.64 (d, J = 1.9 Hz, 1H), 6.93 (dd, J = 13.0, 2.6 Hz, 2H), 4.77 (s, 1H), 3.94 (q, J = 6.6, 5.0 Hz, 5H), 3.35 (t, J = 4.8 Hz, 4H), 2.16 (d, J = 33.6 Hz, 2H), 1.86 (m, J = 4.9 Hz, 6H), 1.51 (q, J = 2.0, 1.6 Hz, 9H).
297		447	¹ H NMR (400 MHz, CDCl ₃) δ 8.60 (d, J = 1.9 Hz, 1H), 8.49 (d, J = 1.9 Hz, 1H), 8.02 (d, J = 6.0 Hz, 1H), 6.78 (dd, J = 10.1, 2.5 Hz, 2H), 6.08 (d, J = 6.0 Hz, 1H), 4.96 (s, 1H), 4.74 (d, J = 2.4 Hz, 1H), 4.49 (d, J = 2.3 Hz, 2H), 3.41 (t, J = 5.4 Hz, 2H), 3.12 (dd, J = 11.6, 2.5 Hz, 2H), 2.43 (s, 3H), 2.14 (dd, J = 9.6, 4.9 Hz, 2H), 1.98 - 1.68 (m, 10H).
298		459.4	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.65 (d, J = 1.9 Hz, 1H), 7.53 (dt, J = 8.1, 1.0 Hz, 1H), 7.35 (ddd, J = 8.1, 6.8, 1.1 Hz, 1H), 7.20 (dt, J = 8.5, 0.9 Hz, 1H), 7.07 - 6.89 (m, 3H), 4.79 (td, J = 6.1, 3.1 Hz, 1H), 4.02 - 3.91 (m, 4H), 3.87 (s, 3H), 3.43 - 3.25 (m, 4H), 2.34 - 2.15 (m, 2H), 2.10 - 1.88 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
299		475.28	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.02 - 6.89 (m, 2H), 6.57 (d, J = 1.0 Hz, 1H), 4.87 - 4.76 (m, 1H), 4.76 - 4.62 (m, 1H), 4.09 - 3.88 (m, 5H), 3.45 - 3.25 (m, 4H), 2.65 (d, J = 0.8 Hz, 3H), 2.56 (s, 3H), 2.31 - 2.12 (m, 2H), 2.09 - 1.81 (m, 6H).
300		407.26	¹ H NMR (400 MHz, CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.13 (d, J = 5.9 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.85 (d, J = 2.4 Hz, 1H), 6.13 (d, J = 6.0 Hz, 1H), 5.17 (s, 1H), 4.50 (s, 1H), 4.29 (d, J = 6.7 Hz, 2H), 4.03 - 3.82 (m, 4H), 3.45 - 3.32 (m, 4H), 3.04 - 2.93 (m, 1H), 2.58 - 2.50 (m, 2H), 2.49 (s, 3H), 2.24 (ddd, J = 20.4, 10.3, 6.0 Hz, 2H).
301		407.26	¹ H NMR (400 MHz, CDCl ₃) δ 8.76 (s, 2H), 8.09 (d, J = 5.7 Hz, 1H), 6.94 (d, J = 2.3 Hz, 1H), 6.80 (d, J = 2.2 Hz, 1H), 6.73 (s, 1H), 6.19 (d, J = 5.1 Hz, 1H), 4.35 (s, 1H), 4.17 (d, J = 3.9 Hz, 2H), 4.03 - 3.77 (m, 4H), 3.45 - 3.21 (m, 4H), 2.85 (dd, J = 19.9, 9.1 Hz, 2H), 2.80 - 2.70 (m, 1H), 2.51 (s, 3H), 2.06 (dt, J = 12.7, 6.4 Hz, 2H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
302		407.21	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.13 (d, J = 5.8 Hz, 1H), 6.92 (d, J = 2.4 Hz, 1H), 6.55 (d, J = 2.4 Hz, 1H), 6.18 (d, J = 6.0 Hz, 1H), 5.07 - 4.97 (m, 1H), 4.91 (s, 1H), 3.99 - 3.82 (m, 4H), 3.50 (d, J = 5.5 Hz, 1H), 3.38 - 3.23 (m, 4H), 2.72 (dd, J = 8.1, 4.8 Hz, 1H), 2.65 (dt, J = 15.2, 7.7 Hz, 2H), 2.51 (s, 3H), 2.45 (ddd, J = 11.1, 6.8, 3.4 Hz, 2H).
303		407.26	¹ H NMR (400 MHz, CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.10 (d, J = 5.8 Hz, 1H), 6.92 (d, J = 2.4 Hz, 1H), 6.67 (d, J = 2.4 Hz, 1H), 6.15 (d, J = 6.0 Hz, 1H), 4.95 (s, 1H), 4.79 (p, J = 7.1 Hz, 1H), 3.98 - 3.83 (m, 4H), 3.45 (bt, 2H), 3.36 - 3.27 (m, 4H), 2.79 (dtd, J = 9.8, 7.2, 2.8 Hz, 2H), 2.48 (s, 3H), 2.38 (dt, J = 16.0, 8.0 Hz, 1H), 2.19 (ddd, J = 17.0, 9.7, 2.8 Hz, 2H).
304		393.21	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.27 (d, J = 4.8 Hz, 2H), 6.93 (d, J = 2.4 Hz, 1H), 6.85 (d, J = 2.4 Hz, 1H), 6.54 (t, J = 4.8 Hz, 1H), 5.36 (d, J = 7.2 Hz, 1H), 4.70 - 4.58 (m, 1H), 4.31 (d, J = 7.7 Hz, 2H), 4.01 - 3.86 (m, 4H), 3.44 - 3.30 (m, 4H), 3.04 - 2.93 (m, 1H), 2.50 (ddd, J = 13.5, 7.8, 3.1 Hz, 2H), 2.31 - 2.19 (m, 2H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
305		393.26	¹ H NMR (400 MHz, CDCl ₃) δ 8.81 (d, J = 1.9 Hz, 1H), 8.74 (d, J = 1.9 Hz, 1H), 8.29 (d, J = 4.8 Hz, 2H), 6.93 (d, J = 2.4 Hz, 1H), 6.80 (d, J = 2.4 Hz, 1H), 6.64 (d, J = 8.5 Hz, 1H), 6.50 (t, J = 4.8 Hz, 1H), 4.60 (dq, J = 15.6, 7.6 Hz, 1H), 4.19 (d, J = 4.6 Hz, 2H), 4.01 - 3.84 (m, 4H), 3.43 - 3.27 (m, 4H), 2.88 - 2.78 (m, 2H), 2.78 - 2.68 (m, 1H), 2.08 - 1.97 (m, 2H).
306		393.26	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.29 (d, J = 4.8 Hz, 2H), 6.91 (d, J = 2.4 Hz, 1H), 6.63 - 6.51 (m, 2H), 5.21 (s, 1H), 5.03 (p, J = 6.6 Hz, 1H), 4.02 - 3.79 (m, 4H), 3.63 (dd, J = 7.3, 5.9 Hz, 2H), 3.42 - 3.19 (m, 4H), 2.79 - 2.68 (m, 1H), 2.68 - 2.54 (m, 2H), 2.52 - 2.40 (m, 2H).
307		393.26	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.8 Hz, 1H), 8.61 (d, J = 1.8 Hz, 1H), 8.26 (d, J = 4.8 Hz, 2H), 6.91 (d, J = 2.0 Hz, 1H), 6.68 (d, J = 2.1 Hz, 1H), 6.53 (t, J = 4.8 Hz, 1H), 5.19 (s, 1H), 4.76 (p, J = 7.2 Hz, 1H), 3.99 - 3.79 (m, 4H), 3.56 (t, J = 6.4 Hz, 2H), 3.41 - 3.23 (m, 4H), 2.77 (dt, J = 9.7, 7.3 Hz, 2H), 2.39 (dt, J = 16.5, 8.3 Hz, 1H), 2.19 (dd, J = 19.6, 9.7 Hz, 2H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
308		478.64	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 7.79 (s, 1H), 7.23 (s, 1H), 6.95 (dd, J = 14.0, 2.5 Hz, 2H), 5.23 (d, J = 8.1 Hz, 1H), 4.80 (s, 1H), 4.09 (s, 1H), 3.94 (dd, J = 6.1, 3.6 Hz, 4H), 3.43 - 3.23 (m, 4H), 3.02 (d, J = 5.1 Hz, 3H), 2.43 (s, 3H), 2.31 - 2.12 (m, 2H), 1.95 (p, J = 10.0 Hz, 6H).
309		477.54	¹ H NMR (300 MHz, Chloroform-d) δ 8.68 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 6.85 (s, 2H), 6.30 (s, 1H), 5.06 (d, J = 16.2 Hz, 1H), 4.85 (d, J = 6.5 Hz, 3H), 4.44 - 4.30 (m, 2H), 4.14 (q, J = 7.1 Hz, 1H), 3.89 - 3.60 (m, 5H), 3.50 (s, 3H), 3.35 (q, J = 7.2 Hz, 1H), 2.49 (s, 3H), 2.24 (d, J = 8.7 Hz, 2H), 2.08 (d, J = 7.7 Hz, 2H), 1.92 (q, J = 9.6, 6.9 Hz, 6H), 1.28 (t, J = 7.1 Hz, 2H).
310		461.26	¹ H NMR (300 MHz, CDCl ₃) δ 8.72 (d, J = 1.8 Hz, 1H), 8.66 (d, J = 1.9 Hz, 1H), 7.35 (d, J = 4.3 Hz, 1H), 7.17 (d, J = 4.8 Hz, 1H), 6.97 (dd, J = 12.0, 2.2 Hz, 2H), 4.92 (s, 1H), 4.35 (s, 1H), 4.01 - 3.84 (m, 4H), 3.48 - 3.26 (m, 4H), 2.72 (s, 3H), 2.36 - 2.21 (m, 2H), 2.13 - 1.85 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
311		436.29	
312		437.33	¹ H NMR (300 MHz, CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.65 (d, J = 1.9 Hz, 1H), 7.57 (d, J = 3.2 Hz, 1H), 7.35 - 7.28 (m, 2H), 6.97 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 5.23 (s, 1H), 4.76 (s, 1H), 4.20 (s, 1H), 4.01 (s, 3H), 3.98 - 3.84 (m, 4H), 3.44 - 3.29 (m, 4H), 2.27 - 2.14 (m, 2H), 2.05 - 1.85 (m, 6H).
313		460.29	¹ H NMR (300 MHz, CDCl ₃) δ 8.68 (dd, J = 21.2, 1.9 Hz, 2H), 7.86 (s, 1H), 7.18 (d, J = 4.9 Hz, 1H), 7.00 (d, J = 4.9 Hz, 1H), 6.96 (dd, J = 12.1, 2.4 Hz, 2H), 5.18 (d, J = 7.4 Hz, 1H), 4.80 (s, 1H), 4.30 (dd, J = 11.9, 6.0 Hz, 1H), 4.05 - 3.77 (m, 4H), 3.45 - 3.14 (m, 4H), 2.75 (s, 3H), 2.37 - 2.14 (m, 2H), 2.14 - 1.87 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
314		435.32	¹ H NMR (300 MHz, CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.66 (s, 1H), 6.94 (dd, J = 14.9, 2.4 Hz, 2H), 4.84 - 4.74 (m, 1H), 4.47 (s, 1H), 4.02 - 3.87 (m, 4H), 3.87 - 3.75 (m, 1H), 3.47 - 3.21 (m, 4H), 2.38 (d, J = 4.0 Hz, 6H), 2.26 - 2.13 (m, 2H), 2.01 - 1.78 (m, 6H).
315		476.23	¹ H NMR (300 MHz, Chloroform-d) δ 8.68 (d, J = 1.9 Hz, 1H), 8.59 - 8.46 (m, 2H), 7.98 (s, 1H), 7.18 (s, 1H), 6.85 (s, 2H), 4.85 (d, J = 6.6 Hz, 2H), 3.90 - 3.62 (m, 5H), 3.35 (d, J = 8.2 Hz, 1H), 3.03 (d, J = 5.1 Hz, 3H), 2.27 (s, 2H), 2.13 - 1.80 (m, 7H), 10.58 (s, 2H).
316		411.34	¹ H NMR (300 MHz, Chloroform-d) δ 8.14 (d, J = 6.2 Hz, 1H), 6.50 (d, J = 1.7 Hz, 1H), 6.43 (d, J = 1.6 Hz, 1H), 6.21 (d, J = 6.2 Hz, 1H), 4.98 (s, 1H), 4.01 - 3.81 (m, 5H), 3.37 - 3.18 (m, 5H), 2.54 (s, 3H), 2.20 (d, J = 9.1 Hz, 3H), 2.05 - 1.71 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
317		476.55	¹ H NMR (300 MHz, Chloroform-d) δ 8.68 (d, J = 1.9 Hz, 1H), 8.62 - 8.41 (m, 2H), 7.79 (s, 1H), 7.33 (d, J = 4.9 Hz, 1H), 6.85 (q, J = 2.6 Hz, 3H), 4.85 (d, J = 6.3 Hz, 4H), 4.08 (s, 0H), 3.88 - 3.58 (m, 5H), 3.35 (q, J = 6.8 Hz, 1H), 3.03 (d, J = 5.1 Hz, 3H), 2.36 - 2.13 (m, 2H), 2.15 - 1.84 (m, 6H).
318		465.3	¹ H NMR (300 MHz, Chloroform-d) δ 8.68 (d, J = 1.9 Hz, 1H), 8.56 (d, J = 1.9 Hz, 1H), 8.33 (d, J = 1.9 Hz, 1H), 6.85 (s, 2H), 5.07 (d, J = 8.1 Hz, 1H), 4.86 (d, J = 6.4 Hz, 4H), 4.23 (s, 2H), 3.90 - 3.62 (m, 4H), 3.35 (q, J = 7.1 Hz, 1H), 2.73 (qd, J = 7.6, 2.3 Hz, 2H), 2.27 (d, J = 10.1 Hz, 3H), 2.12 - 1.82 (m, 6H), 1.28 (t, J = 7.6 Hz, 3H).
319		461.38	¹ H NMR (300 MHz, Chloroform-d) δ 8.83 (dt, J = 6.4, 1.4 Hz, 2H), 8.51 (d, J = 4.7 Hz, 1H), 7.78 (s, 1H), 7.68 (d, J = 1.7 Hz, 1H), 7.34 (dd, J = 4.9, 0.9 Hz, 1H), 6.38 (s, 1H), 5.32 (d, J = 0.9 Hz, 1H), 4.87 (s, 1H), 4.43 (q, J = 2.8 Hz, 2H), 4.17 - 3.94 (m, 3H), 3.03 (dd, J = 5.1, 1.0 Hz, 3H), 2.76 - 2.59 (m, 2H), 2.23 (d, J = 12.7 Hz, 2H), 1.98 (d, J = 7.7 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
320		451.44	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.10 (d, J = 6.0 Hz, 1H), 6.97 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.6 Hz, 1H), 6.15 (d, J = 6.0 Hz, 1H), 4.99 (s, 1H), 4.81 (s, 1H), 4.19 - 4.08 (m, 1H), 3.94 - 3.69 (m, 5H), 3.63 (t, J = 11.3 Hz, 2H), 3.04 (td, J = 11.9, 3.5 Hz, 1H), 2.92 - 2.80 (m, 1H), 2.50 (s, 3H), 2.20 (d, J = 8.9 Hz, 2H), 1.90 (d, J = 5.6 Hz, 7H).
321		461.63	¹ H NMR (300 MHz, Chloroform-d) δ 8.81 - 8.67 (m, 2H), 8.42 (s, 1H), 7.89 (s, 1H), 7.58 (d, J = 1.8 Hz, 1H), 7.09 (s, 1H), 6.28 (dq, J = 3.0, 1.6 Hz, 1H), 4.81 (s, 1H), 4.33 (q, J = 2.8 Hz, 2H), 3.92 (t, J = 5.4 Hz, 2H), 2.93 (d, J = 5.1 Hz, 3H), 2.67 - 2.48 (m, 2H), 2.18 (d, J = 11.3 Hz, 2H), 1.98 - 1.73 (m, 6H).
322		481.3	¹ H NMR (300 MHz, Chloroform-d) δ 8.62 (d, J = 1.9 Hz, 1H), 8.52 (d, J = 1.9 Hz, 1H), 6.85 (dd, J = 18.8, 2.5 Hz, 2H), 5.59 (d, J = 8.3 Hz, 1H), 5.24 (s, 1H), 4.72 (s, 1H), 3.84 (t, J = 4.9 Hz, 5H), 3.66 - 3.39 (m, 3H), 3.30 - 3.19 (m, 5H), 2.78 (t, J = 7.0 Hz, 2H), 2.11 (s, 1H), 2.00 - 1.66 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
323		435.33	
324		450.35	¹ H NMR (300 MHz, CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.79 (d, J = 5.9 Hz, 1H), 6.97 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 6.13 (dd, J = 5.9, 2.1 Hz, 1H), 5.87 (d, J = 2.0 Hz, 1H), 4.87 - 4.70 (m, 1H), 4.32 (q, J = 7.1 Hz, 2H), 4.21 (d, J = 7.8 Hz, 1H), 4.05 - 3.81 (m, 4H), 3.60 - 3.45 (m, 1H), 3.45 - 3.24 (m, 4H), 2.33 - 2.10 (m, 2H), 2.02 - 1.77 (m, 6H), 1.39 (t, J = 7.1 Hz, 3H). [2]
325		422.34	¹ H NMR (300 MHz, CDCl ₃) δ 11.03 (s, 1H), 8.71 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.09 (d, J = 7.1 Hz, 1H), 6.97 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 5.67 (dd, J = 7.2, 2.1 Hz, 1H), 5.58 (d, J = 2.1 Hz, 1H), 4.84 - 4.75 (m, 1H), 4.41 (d, J = 7.6 Hz, 1H), 4.02 - 3.87 (m, 4H), 3.54 - 3.43 (m, 1H), 3.42 - 3.27 (m, 4H), 2.28 - 2.15 (m, 2H), 2.05 - 1.78 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
326		423.4	¹ H NMR (300 MHz, Chloroform-d) δ 8.68 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.01 - 6.83 (m, 2H), 5.34 (s, 1H), 4.73 (d, J = 6.1 Hz, 1H), 3.91 (dd, J = 6.1, 3.7 Hz, 4H), 3.59 (s, 4H), 3.44 (s, 1H), 3.33 (t, J = 4.9 Hz, 4H), 2.17 (s, 5H), 1.86 (q, J = 9.1, 6.3 Hz, 6H).
327		523.35	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 2.0 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.00 (d, J = 1.3 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.5 Hz, 1H), 4.88 (d, J = 8.2 Hz, 1H), 4.77 (s, 1H), 4.63 (d, J = 9.0 Hz, 2H), 4.15 (s, 1H), 3.92 (t, J = 4.9 Hz, 4H), 3.39 - 3.26 (m, 4H), 2.91 (td, J = 8.6, 3.9 Hz, 1H), 2.72 - 2.55 (m, 2H), 2.50 - 2.10 (m, 7H), 1.91 (d, J = 5.5 Hz, 6H), 1.72 (d, J = 4.7 Hz, 1H).
328		521.69	¹ H NMR (300 MHz, Chloroform-d) δ 8.81 - 8.59 (m, 3H), 8.54 (d, J = 1.9 Hz, 1H), 6.92 - 6.73 (m, 2H), 5.50 (d, J = 8.2 Hz, 1H), 4.72 (s, 1H), 4.01 (s, 2H), 3.84 (t, J = 4.9 Hz, 5H), 3.50 (q, J = 5.3 Hz, 3H), 3.37 - 3.17 (m, 5H), 2.62 (t, J = 5.6 Hz, 2H), 2.34 (s, 7H), 2.13 (d, J = 10.7 Hz, 2H), 1.87 (d, J = 22.1 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
329		460.44	¹ H NMR (300 MHz, Chloroform-d) δ 8.62 (t, J = 1.7 Hz, 1H), 8.55 (t, J = 1.7 Hz, 1H), 8.37 (dt, J = 4.6, 1.5 Hz, 1H), 7.78 (dt, J = 7.9, 1.5 Hz, 1H), 6.93 - 6.77 (m, 3H), 4.71 (d, J = 6.1 Hz, 1H), 3.96 - 3.77 (m, 6H), 3.33 - 3.18 (m, 4H), 2.23 - 2.07 (m, 2H), 2.01 - 1.77 (m, 6H).
330		463.6	¹ H NMR (300 MHz, Chloroform-d) δ 8.62 (d, J = 1.9 Hz, 1H), 8.53 (d, J = 1.9 Hz, 1H), 8.10 (dd, J = 4.8, 1.5 Hz, 1H), 7.61 (dd, J = 8.0, 1.5 Hz, 1H), 7.12 (dd, J = 8.1, 4.8 Hz, 1H), 6.95 - 6.76 (m, 2H), 5.39 (s, 1H), 4.74 (s, 1H), 3.96 - 3.75 (m, 5H), 3.26 (t, J = 4.9 Hz, 4H), 2.16 (d, J = 12.8 Hz, 2H), 2.05 - 1.71 (m, 6H).
331		455.4	¹ H NMR (300 MHz, Chloroform-d) δ 8.62 (d, J = 1.9 Hz, 1H), 8.53 (d, J = 1.9 Hz, 1H), 7.28 (d, J = 0.7 Hz, 1H), 6.94 - 6.77 (m, 2H), 5.39 (s, 1H), 5.09 (d, J = 7.8 Hz, 1H), 4.71 (s, 1H), 3.93 - 3.77 (m, 4H), 3.58 (s, 1H), 3.26 (dd, J = 6.0, 3.7 Hz, 4H), 2.11 (s, 2H), 1.97 - 1.66 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
332		465.36	
333		435.33	
334		451.36	




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
335		421.37	
336		446.35	
337		533.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.67 - 8.58 (m, 1H), 8.58 - 8.49 (m, 1H), 8.33 (s, 2H), 6.92 - 6.78 (m, 2H), 5.41 (d, J = 8.1 Hz, 1H), 4.72 (s, 1H), 3.98 (d, J = 8.6 Hz, 1H), 3.84 (t, J = 4.8 Hz, 4H), 3.58 (s, 4H), 3.26 (t, J = 4.9 Hz, 4H), 2.36 (t, J = 4.9 Hz, 4H), 2.25 (s, 3H), 2.13 (d, J = 10.8 Hz, 2H), 1.84 (s, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
338		447.43	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.72 (d, J = 1.9 Hz, 1H), 8.59 (d, J = 1.9 Hz, 1H), 7.17 (d, J = 2.5 Hz, 1H), 6.84 (d, J = 2.4 Hz, 1H), 6.69 - 6.61 (m, 1H), 6.50 (m, 2H), 4.94 (m, 1H), 4.50 (t, J = 8.7 Hz, 2H), 4.27 (d, J = 8.2 Hz, 1H), 3.79 (m, 4H), 3.45 (m, 1H), 3.32 (m, 4H), 3.13 (t, J = 8.7 Hz, 2H), 1.98 (m, 2H), 1.74 (m, 6H).
339		449.41	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.72 (d, J = 1.9 Hz, 1H), 8.58 (d, J = 1.9 Hz, 1H), 7.14 (d, J = 2.5 Hz, 1H), 6.84 (d, J = 2.3 Hz, 1H), 6.64 (t, J = 8.0 Hz, 1H), 6.36 (d, J = 8.2 Hz, 1H), 6.22 (dd, J = 7.7, 1.0 Hz, 1H), 5.89 (s, 2H), 4.96 (m, 2H), 3.79 (m, 4H), 3.51 (m, 1H), 3.32 (m, 4H), 2.03 (m, 2H), 1.75 (m, 6H).
340		474.48	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.73 (d, J = 1.9 Hz, 1H), 8.59 (d, J = 1.9 Hz, 1H), 7.25 (m, 1H), 7.14 (d, J = 2.4 Hz, 1H), 6.89 - 6.78 (m, 3H), 5.47 (d, J = 7.7 Hz, 1H), 4.94 (m, 1H), 4.29 (s, 2H), 3.79 (m, 4H), 3.51 (m, 1H), 3.06 (s, 3H), 2.09 (m, 2H), 1.81 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
341		465.57	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.72 (d, J = 1.9 Hz, 1H), 8.58 (d, J = 1.9 Hz, 1H), 8.17 (s, 2H), 7.15 (d, J = 2.4 Hz, 1H), 6.83 (d, J = 2.3 Hz, 1H), 6.06 (d, J = 7.8 Hz, 1H), 4.92 (m, 1H), 4.77 (s, 1H), 3.79 (m, 4H), 3.50 (m, 1H), 3.30 (m, 4H), 2.02 (m, 2H), 1.80 (m, 6H), 1.42 (s, 6H).
342		364.27	¹ H NMR (400 MHz, CDCl ₃) δ 8.86 (d, J = 0.9 Hz, 1H), 8.84 (s, 1H), 8.16 (s, 1H), 7.73 - 7.63 (m, 2H), 7.17 (d, J = 7.0 Hz, 1H), 5.24 (d, J = 7.8 Hz, 1H), 5.03 (s, 2H), 4.83 (s, 2H), 4.79 (s, 1H), 4.12 - 3.99 (m, 1H), 2.26 - 2.14 (m, 2H), 2.01 - 1.84 (m, 6H).
343		523.44	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 2.0 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.00 (d, J = 1.3 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.5 Hz, 1H), 4.88 (d, J = 8.2 Hz, 1H), 4.77 (s, 1H), 4.63 (d, J = 9.0 Hz, 2H), 4.15 (s, 1H), 3.92 (t, J = 4.9 Hz, 4H), 3.39 - 3.26 (m, 4H), 2.91 (td, J = 8.6, 3.9 Hz, 1H), 2.72 - 2.55 (m, 2H), 2.50 - 2.10 (m, 7H), 1.91 (d, J = 5.5 Hz, 6H), 1.72 (d, J = 4.7 Hz, 1H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
344		441.58	
345		449.59	¹ H NMR (300 MHz, DMSO-d ₆) δ 9.18 (d, J = 1.5 Hz, 1H), 8.98 (t, 1H), 8.87 (d, J = 2.5 Hz, 1H), 8.76 - 8.69 (m, 2H), 8.57 (d, J = 1.9 Hz, 1H), 7.12 (d, J = 2.6 Hz, 1H), 6.82 (d, J = 2.3 Hz, 1H), 4.94 (m, 1H), 3.78 (m, 4H), 3.27 (m, 6H), 1.99 (m, 2H), 1.83 - 1.48 (m, 7H).
346		385.56	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.71 (d, J = 1.9 Hz, 1H), 8.59 (d, J = 1.9 Hz, 1H), 7.85 (t, 1H), 7.12 (d, J = 2.5 Hz, 1H), 6.83 (d, J = 2.4 Hz, 1H), 4.93 (m, 1H), 3.78 (m, 4H), 3.31 (m, 3H), 2.95 (t, J = 6.0 Hz, 2H), 2.03 - 1.90 (m, 2H), 1.80 (s, 3H), 1.52 (m, 7H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
347		450.63	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.72 (d, J = 1.9 Hz, 1H), 8.59 (s, 1H), 7.60 (d, J = 2.3 Hz, 1H), 7.46 (s, 1H), 7.13 (s, 1H), 6.83 (s, 1H), 6.46 (s, 1H), 5.97 (m, 1H), 4.96 (m, 1H), 3.87 - 3.70 (m, 7H), 3.30 (s, 4H), 2.95 (t, J = 5.7 Hz, 2H), 1.99 (m, 2H), 1.62 (t, J = 18.1 Hz, 7H).
348		421.6	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.72 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 8.24 (d, J = 4.7 Hz, 1H), 7.22 (t, J = 6.0 Hz, 1H), 7.11 (d, J = 2.6 Hz, 1H), 6.82 (d, J = 2.3 Hz, 1H), 6.51 (t, J = 4.7 Hz, 1H), 4.93 (m, 1H), 3.78 (m, 4H), 3.31 (m, 4H), 3.19 (t, J = 6.5 Hz, 2H), 1.97 (m, 2H), 1.55 (m, 7H).
349		464.58	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.72 (d, J = 1.9 Hz, 1H), 8.57 (d, J = 1.9 Hz, 1H), 7.59 (d, J = 5.9 Hz, 1H), 7.14 (d, J = 2.5 Hz, 1H), 6.82 (d, J = 2.3 Hz, 1H), 6.44 (d, J = 7.7 Hz, 1H), 6.22 (dd, J = 5.8, 2.1 Hz, 1H), 5.80 - 5.74 (m, 1H), 5.19 - 5.11 (m, 1H), 4.92 (m, 1H), 3.79 (m, 4H), 3.32 (m, 4H), 2.01 (m, 2H), 1.75 (m, 6H), 1.21 (d, J = 6.1 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
350		464.42	¹ H NMR (300 MHz, Chloroform-d) δ 4.79 (s, 1H), 4.24 - 3.76 (m, 5H), 3.50 - 3.20 (m, 4H), 3.02 (d, J = 5.1 Hz, 3H), 2.20 (d, J = 12.8 Hz, 2H), 2.09 - 1.75 (m, 6H), 5.29 - 5.19 (m, 1H), 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.49 (d, J = 4.8 Hz, 1H), 7.77 (s, 1H), 7.32 (d, J = 4.9 Hz, 1H), 7.02 - 6.86 (m, 2H).
351		478.38	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 2.0 Hz, 1H), 8.38 (d, J = 4.9 Hz, 1H), 6.93 (dd, J = 14.5, 2.4 Hz, 2H), 6.68 (d, J = 4.9 Hz, 1H), 5.28 (s, 1H), 4.79 (s, 1H), 4.22 - 3.86 (m, 5H), 3.34 (t, J = 4.8 Hz, 4H), 3.17 - 2.97 (m, 6H), 2.20 (s, 2H), 1.91 (d, J = 5.1 Hz, 6H).
352		418.5	¹ H NMR (300 MHz, Chloroform-d) δ 8.91 - 8.75 (m, 2H), 8.11 (d, J = 5.9 Hz, 1H), 7.68 (d, J = 1.8 Hz, 1H), 6.37 (tt, J = 3.0, 1.5 Hz, 1H), 6.16 (d, J = 6.0 Hz, 1H), 4.99 (s, 1H), 4.88 (d, J = 6.0 Hz, 1H), 4.43 (q, J = 2.8 Hz, 2H), 4.02 (t, J = 5.4 Hz, 2H), 3.85 (s, 1H), 2.77 - 2.64 (m, 2H), 2.51 (s, 3H), 2.23 (d, J = 13.2 Hz, 2H), 2.03 - 1.82 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
353		433.62	¹ H NMR (300 MHz, Chloroform-d) δ 8.67 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 8.11 (d, J = 6.0 Hz, 1H), 6.85 (s, 2H), 6.16 (d, J = 6.0 Hz, 1H), 5.00 (s, 1H), 4.85 (d, J = 6.5 Hz, 3H), 3.93 - 3.57 (m, 5H), 3.35 (q, J = 7.1 Hz, 1H), 2.62 (d, J = 8.0 Hz, 1H), 2.50 (s, 3H), 2.24 (d, J = 8.9 Hz, 2H), 2.01 - 1.81 (m, 6H).
354		407.56	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.73 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 3.1 Hz, 1H), 8.58 (d, J = 2.0 Hz, 1H), 8.47 (d, J = 6.2 Hz, 1H), 7.15 (d, J = 2.4 Hz, 1H), 6.99 (d, J = 7.8 Hz, 1H), 6.83 (d, J = 2.3 Hz, 1H), 6.65 (m, 1H), 4.94 (m, 1H), 3.79 (m, 4H), 3.55 (m, 1H), 3.35 (m, 4H), 2.03 (m, 2H), 1.78 (m, 6H).
355		421.23	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.8 Hz, 1H), 8.61 (d, J = 1.8 Hz, 1H), 8.17 (s, 1H), 6.93 (d, J = 2.2 Hz, 1H), 6.71 (d, J = 2.2 Hz, 1H), 5.36 (d, J = 7.8 Hz, 1H), 5.04 (s, 2H), 4.85 (s, 2H), 4.74 - 4.61 (m, 1H), 4.40 - 4.28 (m, 1H), 4.02 - 3.82 (m, 4H), 3.43 - 3.27 (m, 4H), 3.26 - 3.13 (m, 2H), 2.35 (dd, J = 19.4, 9.4 Hz, 2H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
356		419.19	¹ H NMR (400 MHz, CDCl ₃) δ 8.72 (s, 1H), 8.62 (d, J = 1.5 Hz, 1H), 8.50 (s, 1H), 7.74 (d, J = 1.9 Hz, 1H), 6.95 (d, J = 2.0 Hz, 1H), 6.86 (d, J = 1.9 Hz, 1H), 6.73 (d, J = 2.1 Hz, 1H), 5.44 (d, J = 7.5 Hz, 1H), 4.82 - 4.72 (m, 1H), 4.72 - 4.61 (m, 1H), 4.00 - 3.86 (m, 4H), 3.40 - 3.32 (m, 4H), 3.32 - 3.22 (m, 2H), 2.53 - 2.40 (m, 2H).
357		393.39	¹ H NMR (400 MHz, CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.12 (d, J = 5.9 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.70 (d, J = 2.4 Hz, 1H), 6.13 (d, J = 5.9 Hz, 1H), 5.06 (s, 1H), 4.72 (p, J = 6.9 Hz, 1H), 4.18 (s, 1H), 3.98 - 3.85 (m, 4H), 3.40 - 3.27 (m, 4H), 3.27 - 3.13 (m, 2H), 2.50 (s, 3H), 2.36 (ddd, J = 12.7, 10.0, 2.9 Hz, 2H).
358		463.41	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.72 (d, J = 1.9 Hz, 1H), 8.57 (d, J = 1.9 Hz, 1H), 8.22 (m, 1H), 8.00 (d, J = 2.7 Hz, 1H), 7.73 (d, J = 8.6 Hz, 1H), 7.15 (d, J = 2.3 Hz, 1H), 7.02 (dd, J = 8.7, 2.8 Hz, 1H), 6.83 (d, J = 2.4 Hz, 1H), 6.50 (d, J = 7.6 Hz, 1H), 4.93 (m, 1H), 3.79 (m, 4H), 3.52 (m, 1H), 3.34 (m, 4H), 2.76 (d, J = 4.9 Hz, 3H), 2.04 (m, 2H), 1.79 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
359		537.71	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.99 (d, J = 1.3 Hz, 1H), 6.92 (dd, J = 16.0, 2.5 Hz, 2H), 5.12 (d, J = 7.5 Hz, 1H), 4.76 (d, J = 5.4 Hz, 1H), 4.69 - 4.54 (m, 1H), 4.15 (s, 1H), 4.01 - 3.82 (m, 4H), 3.72 (ddd, J = 13.2, 7.6, 2.9 Hz, 1H), 3.44 - 3.21 (m, 5H), 3.21 - 2.99 (m, 1H), 2.46 (dq, J = 7.9, 3.9 Hz, 1H), 2.34 (s, 3H), 2.22 (t, J = 8.8 Hz, 3H), 1.86 (dd, J = 10.0, 4.8 Hz, 7H), 1.79 - 1.51 (m, 2H).
360		464.58	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.16 (d, J = 1.2 Hz, 1H), 6.93 (dd, J = 17.1, 2.5 Hz, 2H), 5.38-5.21 (m, 1H), 4.90 - 4.77 (m, 1H), 3.91 (dd, J = 6.0, 3.7 Hz, 5H), 3.44 - 3.26 (m, 4H), 3.00 (d, J = 5.1 Hz, 3H), 2.23 (d, J = 11.4 Hz, 2H), 2.04 - 1.76 (m, 6H).
361		478.57	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.38 (d, J = 4.9 Hz, 1H), 7.00 - 6.85 (m, 2H), 6.68 (d, J = 4.9 Hz, 1H), 5.29 (d, J = 7.7 Hz, 1H), 4.79 (s, 1H), 4.04 (s, 1H), 3.92 (dd, J = 6.0, 3.8 Hz, 4H), 3.41 - 3.26 (m, 4H), 3.10 (s, 3H), 3.04 (s, 3H), 2.19 (d, J = 8.7 Hz, 2H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
362		512.55	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.63 (t, J = 1.6 Hz, 1H), 8.04 (s, 1H), 6.95 (d, J = 2.3 Hz, 1H), 6.90 (d, J = 2.7 Hz, 1H), 4.88 (d, J = 8.3 Hz, 1H), 4.80 (s, 1H), 4.48 (t, J = 5.9 Hz, 2H), 4.17 (s, 1H), 4.01 - 3.81 (m, 4H), 3.34 (t, J = 4.7 Hz, 4H), 2.73 (t, J = 5.9 Hz, 2H), 2.33 (d, J = 1.2 Hz, 8H), 2.07 - 1.73 (m, 6H).
363		469.62	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 6.92 (dd, J = 21.1, 2.5 Hz, 2H), 4.74 (s, 1H), 4.35 (d, J = 7.9 Hz, 1H), 4.12 (d, J = 8.0 Hz, 1H), 4.03 - 3.67 (m, 5H), 3.63 - 3.50 (m, 1H), 3.47 - 3.20 (m, 4H), 2.77 (d, J = 11.3 Hz, 2H), 2.28 (s, 3H), 2.22 - 2.04 (m, 4H), 1.90 (dd, J = 38.6, 8.3 Hz, 8H), 1.45 (qd, J = 11.1, 3.8 Hz, 2H).
364		497.61	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.59 (d, J = 2.0 Hz, 1H), 6.92 (dd, J = 21.8, 2.5 Hz, 2H), 4.75 (s, 1H), 4.47 (dd, J = 28.7, 10.8 Hz, 2H), 4.19 (d, J = 7.7 Hz, 1H), 3.99 - 3.68 (m, 6H), 3.46 - 3.27 (m, 4H), 3.18 (t, J = 11.7 Hz, 1H), 2.85 - 2.64 (m, 1H), 2.11 (s, 5H), 1.83 (d, J = 4.9 Hz, 6H), 1.41 - 1.17 (m, 5H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
365		495.54	¹ H NMR (300 MHz, Chloroform-d) δ 8.67 (dd, J = 21.6, 1.9 Hz, 2H), 8.01 (s, 1H), 6.94 (dd, J = 16.5, 2.5 Hz, 2H), 4.79 (s, 1H), 4.67 (s, 2H), 4.18 (s, 2H), 3.93 (t, J = 4.8 Hz, 4H), 3.41 - 3.27 (m, 6H), 2.20 (d, J = 9.9 Hz, 2H), 1.93 (d, J = 5.5 Hz, 6H), 0.63 - 0.50 (m, 2H), 0.29 (t, J = 5.0 Hz, 2H).
366		475.56	
367		413.34	¹ H NMR (400 MHz, CDCl ₃) δ 8.90 (dd, J = 7.2, 1.8 Hz, 2H), 8.75 (dd, J = 4.5, 1.6 Hz, 2H), 8.10 (d, J = 6.0 Hz, 1H), 7.97 (d, J = 1.7 Hz, 1H), 7.62 (dd, J = 4.5, 1.6 Hz, 2H), 7.38 (d, J = 1.7 Hz, 1H), 6.15 (d, J = 6.0 Hz, 1H), 4.98 (s, 1H), 4.93 (s, 1H), 3.86 (s, 1H), 2.49 (s, 3H), 2.34 - 2.18 (m, 2H), 1.96 (dd, J = 20.7, 8.9 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
368		343.48	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.73 (d, J = 2.0 Hz, 1H), 8.60 (d, J = 2.0 Hz, 1H), 7.96 (s, 2H), 7.15 (d, J = 2.5 Hz, 1H), 6.85 (d, J = 2.4 Hz, 1H), 4.97 (m, 1H), 3.79 (t, J = 4.9 Hz, 4H), 3.35 (t, J = 4.9 Hz, 4H), 2.72 (t, J = 6.2 Hz, 2H), 2.65 (m, 1H), 2.01 (m, 2H), 1.77 - 1.48 (m, 6H).
369		436.5	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.71 (s, 1H), 6.94 (dd, J = 14.6, 2.5 Hz, 2H), 5.09 (d, J = 7.7 Hz, 1H), 4.80 (d, J = 5.4 Hz, 1H), 4.29 (s, 2H), 4.01 - 3.84 (m, 4H), 3.45 - 3.28 (m, 4H), 2.49 (s, 3H), 2.21 (q, J = 6.1 Hz, 2H), 1.95 (dt, J = 10.0, 4.1 Hz, 6H).
370		448.25	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (t, J = 1.9 Hz, 1H), 8.64 (dd, J = 3.9, 2.0 Hz, 1H), 8.11 (s, 1H), 7.01 - 6.90 (m, 2H), 4.75 (d, J = 20.3 Hz, 2H), 4.23 (s, 2H), 3.93 (t, J = 4.8 Hz, 4H), 3.68 (t, J = 8.7 Hz, 2H), 3.42 - 3.29 (m, 4H), 2.90 (t, J = 8.7 Hz, 2H), 2.34 - 2.12 (m, 2H), 2.11 - 1.83 (m, 6H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
371		447.5	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.79 (d, J = 5.6 Hz, 1H), 7.03 - 6.89 (m, 2H), 6.05 (d, J = 5.6 Hz, 1H), 4.77 (d, J = 5.9 Hz, 1H), 4.21 (s, 1H), 3.93 (dd, J = 6.0, 3.7 Hz, 4H), 3.68 (t, J = 8.7 Hz, 2H), 3.36 (dd, J = 5.8, 4.0 Hz, 4H), 2.85 (t, J = 8.7 Hz, 2H), 2.20 (q, J = 12.1, 9.5 Hz, 2H), 1.94 (t, J = 5.8 Hz, 6H).
372		446.44	
373		421.51	¹ H NMR (400 MHz, CDCl ₃) δ 8.71 (d, J = 1.8 Hz, 1H), 8.63 (d, J = 1.8 Hz, 1H), 7.70 (d, J = 5.0 Hz, 2H), 6.97 (d, J = 1.6 Hz, 1H), 6.92 (d, J = 2.1 Hz, 1H), 4.81 (s, 1H), 4.65 (s, 1H), 3.99 - 3.85 (m, 5H), 3.40 - 3.30 (m, 4H), 2.37 (s, 3H), 2.28 - 2.16 (m, 2H), 2.01 - 1.82 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
374		441.45	
375		420.57	¹ H NMR (300 MHz, Chloroform-d) δ 8.81 - 8.67 (m, 2H), 8.01 (d, J = 6.0 Hz, 1H), 7.46 (dd, J = 1.7, 0.7 Hz, 1H), 6.98 (d, J = 1.7 Hz, 1H), 6.06 (d, J = 6.0 Hz, 1H), 4.80 (d, J = 38.5 Hz, 2H), 4.13 - 3.97 (m, 3H), 3.51 (td, J = 11.3, 3.5 Hz, 2H), 2.99 - 2.80 (m, 1H), 2.41 (s, 3H), 2.21 - 2.03 (m, 2H), 1.94 - 1.70 (m, 7H).
376		455.33	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.07 (s, 1H), 6.93 (dd, J = 17.0, 2.5 Hz, 2H), 4.98 - 4.71 (m, 2H), 4.17 (s, 1H), 4.00 (s, 3H), 3.92 (dd, J = 5.8, 3.9 Hz, 4H), 3.42 - 3.26 (m, 4H), 2.22 (d, J = 9.5 Hz, 2H), 1.93 (dd, J = 6.1, 2.4 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
377		512.36	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.95 (d, J = 1.2 Hz, 1H), 6.92 (dd, J = 16.9, 2.5 Hz, 2H), 4.98 (s, 1H), 4.81 - 4.66 (m, 2H), 4.16 (s, 1H), 3.97 - 3.87 (m, 4H), 3.46 (d, J = 6.1 Hz, 2H), 3.39 - 3.29 (m, 4H), 2.19 (d, J = 12.5 Hz, 2H), 1.91 (d, J = 5.4 Hz, 6H), 1.26 (s, 6H), 0.94 - 0.83 (m, 1H).
378		524.38	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.99 (d, J = 1.3 Hz, 1H), 6.92 (dd, J = 17.1, 2.5 Hz, 2H), 4.77 (s, 1H), 4.66 (d, J = 8.5 Hz, 1H), 4.41 (d, J = 7.7 Hz, 1H), 4.23 - 4.10 (m, 2H), 4.05 - 3.87 (m, 6H), 3.54 (td, J = 11.7, 2.1 Hz, 2H), 3.39 - 3.29 (m, 4H), 2.19 (d, J = 11.3 Hz, 2H), 2.08 - 1.85 (m, 8H), 1.64 - 1.44 (m, 2H).
379		510.38	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.00 (d, J = 1.3 Hz, 1H), 6.95 (d, J = 2.5 Hz, 1H), 6.89 (d, J = 2.5 Hz, 1H), 4.87 - 4.56 (m, 4H), 4.16 (s, 1H), 4.04 - 3.65 (m, 8H), 3.46 - 3.26 (m, 4H), 2.43 - 2.26 (m, 1H), 2.20 (q, J = 6.6, 5.9 Hz, 2H), 1.90 (t, J = 5.5 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
380		498.36	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 2.0 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.00 - 7.91 (m, 1H), 6.92 (dd, J = 17.2, 2.5 Hz, 2H), 4.87 - 4.64 (m, 2H), 4.59 (d, J = 5.7 Hz, 1H), 4.16 (ddt, J = 9.7, 6.8, 3.7 Hz, 2H), 4.01 - 3.85 (m, 4H), 3.75 (d, J = 11.2 Hz, 1H), 3.58 (dd, J = 10.9, 7.0 Hz, 1H), 3.48 - 3.23 (m, 4H), 2.36 - 2.07 (m, 2H), 1.91 (d, J = 5.5 Hz, 6H), 1.28 (d, J = 6.8 Hz, 3H), 0.97 - 0.78 (m, 1H).
381		523.39	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.98 (d, J = 7.3 Hz, 1H), 6.92 (dd, J = 16.7, 2.5 Hz, 2H), 4.87 (d, J = 8.6 Hz, 1H), 4.77 (s, 2H), 4.61 (s, 2H), 4.15 (s, 1H), 3.91 (dd, J = 5.9, 3.8 Hz, 4H), 3.46 - 3.29 (m, 4H), 2.99 - 2.77 (m, 2H), 2.74 - 2.54 (m, 2H), 2.46 - 2.10 (m, 6H), 1.91 (d, J = 5.5 Hz, 6H).
382		498.32	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.00 (s, 1H), 6.92 (dd, J = 16.5, 2.5 Hz, 2H), 4.92 (s, 1H), 4.77 (s, 1H), 4.64 (d, J = 8.5 Hz, 1H), 4.16 (s, 1H), 4.03 - 3.82 (m, 4H), 3.65 (q, J = 5.8, 5.4 Hz, 2H), 3.56 (ddd, J = 5.6, 4.7, 1.0 Hz, 2H), 3.48 - 3.21 (m, 7H), 2.19 (d, J = 10.1 Hz, 2H), 1.91 (d, J = 5.4 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
383		476.5	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (s, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.55 (d, J = 1.9 Hz, 1H), 6.93 - 6.71 (m, 3H), 6.22 (dd, J = 3.6, 2.0 Hz, 1H), 5.01 (d, J = 8.1 Hz, 1H), 4.71 (s, 1H), 4.34 (s, 1H), 3.92 - 3.78 (m, 7H), 3.26 (dd, J = 5.9, 3.9 Hz, 4H), 2.15 (d, J = 10.8 Hz, 2H), 1.98 - 1.78 (m, 6H).
384		460.6	¹ H NMR (300 MHz, Chloroform-d) δ 8.72 (d, J = 2.0 Hz, 1H), 8.65 (d, J = 1.9 Hz, 1H), 8.39 (s, 1H), 7.01 - 6.90 (m, 3H), 6.35 (d, J = 3.5 Hz, 1H), 5.05 (t, J = 8.6 Hz, 1H), 4.83 (d, J = 5.6 Hz, 1H), 4.41 (s, 1H), 4.01 - 3.87 (m, 4H), 3.82 (d, J = 1.1 Hz, 3H), 3.44 - 3.30 (m, 4H), 2.27 (d, J = 8.9 Hz, 2H), 2.12 - 1.86 (m, 6H).
385		445.5	¹ H NMR (300 MHz, Chloroform-d) δ 8.72 (d, J = 1.9 Hz, 1H), 8.65 (d, J = 1.9 Hz, 1H), 8.03 (d, J = 5.6 Hz, 1H), 7.11 (d, J = 3.6 Hz, 1H), 7.03 - 6.91 (m, 2H), 6.41 (d, J = 3.6 Hz, 1H), 6.28 (d, J = 5.7 Hz, 1H), 4.83 (s, 1H), 4.56 (d, J = 8.0 Hz, 1H), 4.01 - 3.88 (m, 4H), 3.78 (s, 1H), 3.44 - 3.30 (m, 4H), 2.28 (d, J = 10.0 Hz, 2H), 2.11 - 1.85 (m, 6H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
386		480.41	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.49 (s, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 5.25 (s, 1H), 4.72 (s, 1H), 4.18 (s, 1H), 3.96 - 3.87 (m, 4H), 3.77 (s, 3H), 3.39 - 3.30 (m, 4H), 3.10 (s, 6H), 2.21 - 2.12 (m, 2H), 2.02 - 1.85 (m, 6H).
387		466.4	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.49 (s, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 5.25 (s, 1H), 4.72 (s, 1H), 4.18 (s, 1H), 3.96 - 3.87 (m, 4H), 3.77 (s, 3H), 3.39 - 3.30 (m, 4H), 3.10 (s, 6H), 2.21 - 2.12 (m, 2H), 2.02 - 1.85 (m, 6H).
388		480.41	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.04 (s, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 5.15 (d, J = 8.4 Hz, 1H), 4.75 (s, 1H), 4.14 (s, 1H), 3.97 - 3.83 (m, 4H), 3.59 (s, 3H), 3.38 - 3.28 (m, 4H), 3.09 (s, 6H), 2.25 - 2.13 (m, 2H), 2.00 - 1.83 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
389		504.48	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.50 (d, J = 1.1 Hz, 1H), 6.93 (dd, J = 16.5, 2.5 Hz, 2H), 6.49 (s, 1H), 4.90 (s, 1H), 4.80 (s, 1H), 4.08 - 3.87 (m, 5H), 3.44 - 3.29 (m, 6H), 2.45 (t, J = 5.3 Hz, 4H), 2.26 - 2.14 (m, 2H), 1.90 (t, J = 6.5 Hz, 6H), 1.63 (d, J = 5.7 Hz, 4H), 1.53 - 1.43 (m, 2H).
390		464.62	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.50 (d, J = 1.1 Hz, 1H), 6.93 (dd, J = 16.5, 2.5 Hz, 2H), 6.44 (d, J = 1.1 Hz, 1H), 4.93 (s, 1H), 4.80 (s, 1H), 3.97 - 3.87 (m, 4H), 3.47 - 3.29 (m, 6H), 2.31 (s, 6H), 2.21 (d, J = 9.1 Hz, 2H), 1.89 (t, J = 6.9 Hz, 6H).
391		500.63	¹ H NMR (300 MHz, Chloroform-d) δ 8.68 (d, J = 1.9 Hz, 1H), 8.58 (d, J = 1.9 Hz, 1H), 6.94 (d, J = 2.5 Hz, 1H), 6.86 (d, J = 2.5 Hz, 1H), 4.81 - 4.61 (m, 3H), 4.47 (td, J = 8.7, 5.2 Hz, 1H), 3.98 - 3.88 (m, 4H), 3.85 - 3.66 (m, 4H), 3.40 - 3.25 (m, 4H), 2.24 - 2.07 (m, 2H), 1.93 - 1.74 (m, 6H), 1.71-1.64 (m, 1H), 1.58 - 1.39 (m, 2H), 0.93 (t, J = 6.3 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
392		443.61	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.71 (d, J = 1.9 Hz, 1H), 8.58 (d, J = 1.9 Hz, 1H), 7.10 (d, J = 2.5 Hz, 1H), 6.82 (d, J = 2.6 Hz, 2H), 4.92 (m, 1H), 3.78 (m, 4H), 3.30 (m, 5H), 2.84 (t, J = 6.0 Hz, 2H), 1.96 - 1.33 (m, 18H).
393		450.58	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.73 (d, J = 1.9 Hz, 1H), 8.58 (d, J = 1.9 Hz, 1H), 8.10 (d, J = 2.7 Hz, 1H), 7.97 (d, J = 8.9 Hz, 1H), 7.38 (dd, J = 9.0, 2.7 Hz, 1H), 7.16 (d, J = 2.4 Hz, 1H), 6.84 (d, J = 2.3 Hz, 1H), 4.96 (m, 1H), 3.79 (m, 4H), 3.66 (m, 1H), 3.3 (m, 4H), 2.14 - 1.95 (m, 2H), 1.81 (m, 5H).
394		493.61	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.72 (d, J = 1.9 Hz, 1H), 8.58 (d, J = 1.9 Hz, 1H), 7.54 (d, J = 8.6 Hz, 1H), 7.14 (d, J = 2.4 Hz, 1H), 6.83 (d, J = 2.3 Hz, 1H), 6.48 (d, J = 7.7 Hz, 1H), 6.30 - 6.18 (m, 2H), 4.92 (m, 1H), 3.79 (m, 4H), 3.74 (s, 3H), 3.66 (s, 3H), 3.55 (m, 1H), 3.34 (m, 4H), 2.03 (m, 2H), 1.78 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
395		441.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.61 (d, J = 1.9 Hz, 1H), 8.53 (d, J = 2.0 Hz, 1H), 7.13 (d, J = 5.3 Hz, 1H), 6.84 (dd, J = 20.0, 2.4 Hz, 2H), 5.40 (d, J = 8.2 Hz, 1H), 4.72 (s, 1H), 4.31 (d, J = 7.0 Hz, 1H), 3.83 (dd, J = 5.8, 3.8 Hz, 4H), 3.26 (t, J = 4.9 Hz, 4H), 2.14 (d, J = 12.3 Hz, 2H), 1.95 - 1.68 (m, 6H).
396		455.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.61 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 7.72 (d, J = 2.9 Hz, 1H), 6.87 (d, J = 2.5 Hz, 1H), 6.81 (d, J = 2.5 Hz, 1H), 5.03 (d, J = 7.8 Hz, 1H), 4.71 (s, 1H), 4.16 (s, 2H), 3.83 (q, J = 3.9, 3.1 Hz, 7H), 3.32 - 3.21 (m, 4H), 2.13 (d, J = 11.2 Hz, 2H), 1.83 (t, J = 6.5 Hz, 6H).
397		422.5	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.89 (d, J = 5.8 Hz, 1H), 7.02 - 6.88 (m, 2H), 5.79 (d, J = 5.7 Hz, 1H), 4.80 (s, 1H), 4.61 (s, 2H), 4.08 - 3.88 (m, 5H), 3.35 (dd, J = 5.6, 4.1 Hz, 4H), 2.27 - 2.09 (m, 2H), 1.91 (d, J = 5.0 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
398		476.3	¹ H NMR (300 MHz, Chloroform-d) δ 8.62 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 8.38 (s, 1H), 6.94 - 6.74 (m, 2H), 4.69 (s, 1H), 4.34 (s, 1H), 4.25 (s, 1H), 3.84 (dd, J = 5.9, 3.8 Hz, 4H), 3.30 - 3.14 (m, 6H), 2.79 (d, J = 5.7 Hz, 2H), 2.71 (d, J = 5.6 Hz, 2H), 2.49 (s, 3H), 2.13 (m, 2H), 1.85 (d, J = 5.5 Hz, 6H).
399		435.35	¹ H NMR (300 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.08 (d, J = 6.3 Hz, 1H), 6.96 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.4 Hz, 1H), 6.23 (d, J = 6.3 Hz, 1H), 4.87 (s, 1H), 4.69 (s, 1H), 4.00 - 3.83 (m, 4H), 3.45 - 3.24 (m, 4H), 2.98 (s, 3H), 2.50 (s, 3H), 2.45 - 2.30 (m, 2H), 1.92 - 1.64 (m, 6H).
400		452.3	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 6.94 (dd, J = 19.8, 2.4 Hz, 2H), 5.09 (s, 1H), 4.81 (s, 1H), 4.01 - 3.88 (m, 4H), 3.42 - 3.29 (m, 4H), 2.91 (d, J = 4.8 Hz, 3H), 2.18 (m, 2H), 1.90 (s, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
401		471.32	¹ H NMR (300 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.51 (s, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 5.56 (d, J = 8.0 Hz, 1H), 4.75 (s, 1H), 4.25 (s, 1H), 3.96 - 3.89 (m, 4H), 3.88 (s, 3H), 3.47 - 3.23 (m, 4H), 2.29 - 2.06 (m, 2H), 2.06 - 1.78 (m, 6H).
402		471.36	¹ H NMR (300 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.14 (s, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.5 Hz, 1H), 5.52 (d, J = 8.6 Hz, 1H), 4.79 (s, 1H), 4.18 (s, 1H), 3.96 - 3.89 (m, 4H), 3.89 (s, 3H), 3.41 - 3.27 (m, 4H), 2.31 - 2.14 (m, 2H), 2.03 - 1.83 (m, 6H).
403		463.4	¹ H NMR (300 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.5 Hz, 1H), 4.77 (s, 1H), 4.68 (d, J = 8.4 Hz, 1H), 4.58 (t, J = 9.1 Hz, 2H), 4.25 (s, 1H), 3.98 - 3.84 (m, 4H), 3.40 - 3.27 (m, 4H), 3.19 (t, J = 9.0 Hz, 2H), 2.49 (s, 3H), 2.25 - 2.10 (m, 2H), 2.00 - 1.82 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
404		480.56	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.72 (d, J = 1.9 Hz, 1H), 8.57 (d, J = 1.9 Hz, 1H), 7.69 (s, 1H), 7.15 (d, J = 2.5 Hz, 1H), 7.01 (s, 1H), 6.84 (d, J = 2.3 Hz, 1H), 4.94 (m, 1H), 3.75 (m, 8H), 3.34 (m, 4H), 2.05 (m, 2H), 1.86 - 1.70 (m, 6H).
405		480.56	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.72 (d, J = 1.9 Hz, 1H), 8.57 (d, J = 1.9 Hz, 1H), 8.01 (m, 1H), 7.14 (s, 1H), 7.01 - 6.90 (m, 1H), 6.82 (m, 2H), 6.71 (m, 1H), 5.70 (d, J = 7.9 Hz, 1H), 4.91 (m, 1H), 3.79 (m, 4H), 3.34 (m, 5H), 2.74 (d, J = 4.6 Hz, 3H), 2.02 (m, 2H), 1.76 (m, 6H).
406		463.68	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.72 (d, J = 1.9 Hz, 1H), 8.58 (d, J = 1.9 Hz, 1H), 8.27 (d, J = 5.2 Hz, 1H), 7.46 (dd, J = 8.4, 7.1 Hz, 1H), 7.15 (d, J = 2.4 Hz, 1H), 7.10 (dd, J = 7.1, 0.9 Hz, 1H), 6.83 (d, J = 2.3 Hz, 1H), 6.75 (d, J = 7.8 Hz, 1H), 6.68 (dd, J = 8.4, 0.9 Hz, 1H), 4.91 (m, 1H), 4.14 (m, 1H), 3.80 (m, 4H), 3.35 (m, 4H), 2.82 (d, J = 4.9 Hz, 3H), 2.03 (m, 2H), 1.93 - 1.73 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
407		536.49	¹ H NMR (300 MHz, Chloroform-d) rotomers, δ 9.07 (m, 1H), 8.89 (m, 1H), 8.75 (m, 1H), 7.47 (m, 1H), 6.96 (s, 1H), 4.95 (m, 1H), 3.97 (m, 4H), 3.82 (m, 4H), 3.38 (s, 3H), 2.24 (m, 8H), 1.60 (s, 9H).
408		435.6	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.72 (d, J = 1.9 Hz, 1H), 8.57 (d, J = 1.9 Hz, 1H), 7.13 (d, J = 2.4 Hz, 1H), 6.94 (t, J = 8.0 Hz, 1H), 6.83 (d, J = 2.3 Hz, 1H), 6.26 - 6.14 (m, 2H), 6.10 - 6.03 (m, 1H), 5.57 (d, J = 7.8 Hz, 1H), 4.90 (m, 1H), 3.79 (m, 4H), 3.66 (s, 3H), 3.33 (m, 4H), 2.03 (m, 2H), 1.77 (m, 6H).
409		441.29	¹ H NMR (300 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.47 (s, 1H), 8.18 (s, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 5.47 (d, J = 7.6 Hz, 1H), 4.80 (s, 1H), 4.31 - 4.19 (m, 1H), 4.02 - 3.79 (m, 4H), 3.48 - 3.24 (m, 4H), 2.32 - 2.17 (m, 2H), 2.09 - 1.79 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
410		468.38	¹ H NMR (300 MHz, Chloroform-d) δ 8.68 (d, J = 2.0 Hz, 1H), 8.59 (d, J = 1.9 Hz, 1H), 6.94 (d, J = 2.5 Hz, 1H), 6.87 (d, J = 2.5 Hz, 1H), 4.73 (d, J = 5.5 Hz, 1H), 4.49 - 4.19 (m, 2H), 4.03 - 3.87 (m, 4H), 3.81 (q, J = 6.6 Hz, 1H), 3.44 - 3.25 (m, 4H), 2.99 (t, J = 6.3 Hz, 2H), 2.13 (q, J = 7.5, 5.0 Hz, 2H), 1.99 - 1.57 (m, 11H), 1.42 (dq, J = 10.0, 7.1, 6.7, 3.4 Hz, 1H), 1.20 (tt, J = 17.8, 10.5 Hz, 3H), 0.91 (q, J = 12.1 Hz, 2H).
411		400.37	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 6.94 (d, J = 2.5 Hz, 1H), 6.87 (d, J = 2.5 Hz, 1H), 4.70 (d, J = 2.6 Hz, 1H), 4.41 (d, J = 7.7 Hz, 1H), 3.97 - 3.81 (m, 4H), 3.41 - 3.25 (m, 4H), 2.91 (s, 6H), 2.15 (d, J = 9.2 Hz, 2H), 1.98 - 1.71 (m, 6H).
412		442.37	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 6.90 (dd, J = 21.6, 2.5 Hz, 2H), 4.72 (s, 1H), 4.48 (d, J = 7.9 Hz, 1H), 4.04 - 3.82 (m, 4H), 3.78 - 3.62 (m, 4H), 3.34 (dt, J = 5.0, 3.2 Hz, 8H), 2.15 (s, 2H), 1.84 (d, J = 4.8 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
413		428.37	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 2.0 Hz, 1H), 6.94 (d, J = 2.5 Hz, 1H), 6.87 (d, J = 2.5 Hz, 1H), 4.71 (d, J = 5.4 Hz, 1H), 4.37 (d, J = 7.9 Hz, 1H), 3.99 - 3.81 (m, 4H), 3.44 - 3.09 (m, 8H), 2.24 - 2.06 (m, 2H), 1.94 - 1.73 (m, 6H), 1.15 (t, J = 7.1 Hz, 6H).
414		455.42	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 6.94 (d, J = 2.5 Hz, 1H), 6.87 (d, J = 2.5 Hz, 1H), 4.72 (d, J = 5.6 Hz, 1H), 4.50 (d, J = 7.6 Hz, 1H), 4.02 - 3.80 (m, 4H), 3.49 - 3.18 (m, 8H), 2.41 (t, J = 5.1 Hz, 4H), 2.32 (s, 3H), 2.22 - 2.05 (m, 2H), 1.95 - 1.75 (m, 6H).
415		426.52	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 6.94 (d, J = 2.5 Hz, 1H), 6.87 (d, J = 2.5 Hz, 1H), 4.70 (s, 1H), 4.24 (d, J = 7.9 Hz, 1H), 4.02 - 3.82 (m, 5H), 3.44 - 3.23 (m, 8H), 2.61 (s, 1H), 2.26 - 2.06 (m, 2H), 1.99 - 1.73 (m, 9H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
416		440.39	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.87 (d, J = 2.5 Hz, 1H), 4.70 (d, J = 4.4 Hz, 1H), 4.48 (d, J = 7.6 Hz, 1H), 4.03 - 3.80 (m, 5H), 3.49 (d, J = 4.2 Hz, 2H), 3.45 - 3.17 (m, 8H), 2.23 - 2.04 (m, 2H), 2.02 - 1.73 (m, 6H), 1.56 - 1.49 (m, 4H).
417		475.36	¹ H NMR (300 MHz, Chloroform-d) δ 8.81 - 8.55 (m, 3H), 6.94 (dd, J = 17.2, 2.5 Hz, 2H), 6.65 (d, J = 1.1 Hz, 1H), 4.84 (td, J = 5.1, 2.4 Hz, 1H), 4.01 - 3.86 (m, 4H), 3.42 - 3.29 (m, 4H), 2.35 - 2.11 (m, 2H), 2.05 - 1.82 (m, 6H).
418		475.4	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.49 (d, J = 4.9 Hz, 1H), 7.04 - 6.89 (m, 2H), 6.81 (d, J = 4.9 Hz, 1H), 5.54 (d, J = 7.9 Hz, 1H), 4.80 (d, J = 5.2 Hz, 1H), 4.17 - 3.98 (m, 1H), 3.98 - 3.87 (m, 4H), 3.42 - 3.31 (m, 4H), 2.33 - 2.12 (m, 2H), 2.06 - 1.72 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
419		450.38	¹ H NMR (300 MHz, Chloroform-d) rotomers δ 8.82 (d, J = 2.9 Hz, 1H), 8.74 (m, 1H), 8.00 (m, 1H), 7.55 - 7.47 (m, 1H), 7.25 (m, 1H), 7.14 (m, 1H), 7.05 (m, 1H), 3.97 (m, 5H), 3.47 (m, 5H), 2.30 (s, 2H), 2.0 (m, 6H)..
420		467.48	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.73 (d, J = 1.9 Hz, 1H), 8.58 (d, J = 2.0 Hz, 1H), 7.27 - 6.96 (m, 4H), 6.83 (d, J = 2.3 Hz, 1H), 4.92 (m, 1H), 3.85 - 3.74 (m, 4H), 3.45 (m, 1H), 3.34 (m, 4H), 2.05 (m, 2H), 1.76 (m, 6H).
421		523.58	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.93 (d, J = 1.8 Hz, 1H), 8.80 (d, J = 1.8 Hz, 1H), 7.20 (s, 1H), 7.13 - 6.85 (m, 4H), 4.92 (m, 1H), 3.86 - 3.79 (m, 4H), 3.32 (m, 5H), 2.09 (m, 2H), 1.90 - 1.73 (m, 6H), 1.55 (s, 9H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
422		432.5	¹ H NMR (300 MHz, Chloroform-d) δ 8.72 - 8.52 (m, 3H), 8.38 (s, 1H), 6.87 (dd, J = 15.7, 2.5 Hz, 2H), 5.50 (d, J = 7.9 Hz, 1H), 4.79 (p, J = 3.5 Hz, 1H), 4.20 (dp, J = 14.0, 5.2, 4.7 Hz, 1H), 3.90 - 3.76 (m, 4H), 3.32 - 3.22 (m, 4H), 2.29 - 2.07 (m, 2H), 1.99 - 1.71 (m, 6H).
423		481.3	¹ H NMR (300 MHz, Chloroform-d) δ 8.74 (d, J = 1.9 Hz, 1H), 8.65 (s, 1H), 8.01 (s, 1H), 7.05 - 6.88 (m, 2H), 4.87 (s, 1H), 3.95 (d, J = 4.7 Hz, 4H), 3.77 - 3.56 (m, 1H), 3.37 (dd, J = 5.8, 3.8 Hz, 4H), 3.11 (qd, J = 7.4, 4.2 Hz, 1H), 2.28 (s, 2H), 1.98 (dd, J = 22.7, 9.4 Hz, 6H).
424		437.3	¹ H NMR (300 MHz, Chloroform-d) δ 8.62 (d, J = 1.9 Hz, 1H), 8.52 (d, J = 1.9 Hz, 1H), 8.46 - 8.35 (m, 1H), 6.84 (dd, J = 16.8, 2.5 Hz, 2H), 6.28 - 6.20 (m, 1H), 5.08 (s, 1H), 4.72 (d, J = 5.2 Hz, 1H), 4.58 - 4.43 (m, 2H), 3.93 - 3.77 (m, 4H), 3.34 - 3.19 (m, 4H), 2.14 (dd, J = 9.6, 5.4 Hz, 2H), 1.93 - 1.73 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
425		428.46	¹ H NMR (300 MHz, Chloroform-d) δ 8.68 (d, J = 2.0 Hz, 1H), 8.59 (d, J = 2.0 Hz, 1H), 6.93 (d, J = 2.5 Hz, 1H), 6.86 (d, J = 2.5 Hz, 1H), 4.72 (s, 1H), 4.20 (s, 1H), 4.11 (s, 1H), 3.98 - 3.85 (m, 4H), 3.78 (s, 1H), 3.40 - 3.24 (m, 4H), 2.14 (d, J = 9.2 Hz, 2H), 1.81 (t, J = 6.3 Hz, 6H), 1.33 (s, 9H).
426		499.35	¹ H NMR (300 MHz, Chloroform-d) δ 9.26 (m, 2H), 8.90 (m, 1H), 8.79 - 8.64 (m, 2H), 7.22 - 7.01 (m, 3H), 4.97 (m, 1H), 4.01 - 3.87 (m, 4H), 3.38 (m, 5H), 2.77 (s, 3H), 2.19 (m, 9H).
427		474.33	¹ H NMR (300 MHz, Chloroform-d) δ 8.94 - 8.63 (m, 2H), 8.07 (d, J = 2.8 Hz, 1H), 7.45 (d, J = 8.7 Hz, 1H), 7.14 - 6.87 (m, 3H), 4.84 (m, 1H), 3.99 - 3.87 (m, 4H), 3.62 - 3.48 (m, 1H), 3.35 (dt, J = 42.8, 4.8 Hz, 4H), 2.32 - 2.14 (m, 2H), 1.91 (d, J = 17.8 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
428		475.41	¹ H NMR (300 MHz, Chloroform-d) δ 8.81 - 8.64 (m, 2H), 8.20 (s, 2H), 7.08 (d, J = 2.4 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 4.90 - 4.82 (m, 1H), 4.42 (m, 1H), 3.98 - 3.88 (m, 4H), 3.57 (m, 1H), 3.45 - 3.26 (m, 4H), 2.33 - 2.18 (m, 2H), 2.04 - 1.82 (m, 6H).
429		474.37	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 2.0 Hz, 1H), 8.63 (d, J = 2.0 Hz, 1H), 8.42 (m, 1H), 8.16 - 8.08 (m, 1H), 7.20 (s, 1H), 7.04 - 6.98 (m, 1H), 6.94 (d, J = 2.4 Hz, 1H), 4.84 (m, 1H), 4.01 - 3.87 (m, 4H), 3.54 (m, 1H), 3.42 - 3.26 (m, 4H), 2.33 - 2.19 (m, 2H), 2.08 - 1.80 (m, 6H).
430		533.47	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.15 (s, 2H), 7.03 - 6.85 (m, 2H), 5.09 (d, J = 8.1 Hz, 1H), 4.78 (s, 1H), 3.92 (dd, J = 5.9, 3.7 Hz, 5H), 3.40 - 3.24 (m, 4H), 2.71 - 2.33 (m, 8H), 2.30 (s, 3H), 2.20 (s, 2H), 1.91 (d, J = 5.0 Hz, 6H), 1.26 (d, J = 3.1 Hz, 4H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
431		478.47	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.15 (s, 2H), 6.99 - 6.86 (m, 2H), 5.10 (d, J = 8.1 Hz, 1H), 4.79 (d, J = 5.9 Hz, 1H), 4.09 - 3.85 (m, 5H), 3.41 - 3.25 (m, 4H), 2.68 - 2.51 (m, 2H), 2.52 - 2.37 (m, 2H), 2.28 (s, 6H), 2.19 (q, J = 6.2, 5.8 Hz, 2H), 2.02 - 1.81 (m, 6H).
432		504.44	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.16 (s, 2H), 6.98 - 6.86 (m, 2H), 5.10 (d, J = 8.1 Hz, 1H), 4.78 (s, 1H), 4.08 - 3.83 (m, 5H), 3.39 - 3.27 (m, 4H), 2.60 (d, J = 16.9 Hz, 8H), 2.19 (d, J = 7.8 Hz, 2H), 2.01 - 1.74 (m, 10H).
433		518.72	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.15 (s, 2H), 6.99 - 6.85 (m, 2H), 5.09 (d, J = 8.1 Hz, 1H), 4.78 (s, 1H), 4.06 - 3.85 (m, 5H), 3.40 - 3.27 (m, 4H), 2.67 - 2.55 (m, 2H), 2.47 (d, J = 9.1 Hz, 6H), 2.19 (d, J = 9.6 Hz, 2H), 1.91 (d, J = 4.8 Hz, 6H), 1.62 (d, J = 5.5 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
434		520.47	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.16 (s, 2H), 6.99 - 6.84 (m, 2H), 5.11 (d, J = 8.1 Hz, 1H), 4.78 (s, 1H), 4.05 - 3.87 (m, 5H), 3.78 - 3.66 (m, 4H), 3.40 - 3.27 (m, 4H), 2.69 - 2.43 (m, 8H), 2.19 (d, J = 8.2 Hz, 2H), 1.91 (d, J = 4.8 Hz, 6H).
435		461.37	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.68 (d, J = 2.1 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 6.76 (d, J = 2.2 Hz, 1H), 5.21 (d, J = 8.4 Hz, 1H), 4.86 - 4.76 (m, 1H), 4.45 - 4.33 (m, 1H), 3.98 - 3.84 (m, 4H), 3.39 - 3.25 (m, 4H), 2.59 (s, 3H), 2.31 - 2.16 (m, 2H), 2.06 - 1.89 (m, 6H).
436		449.44	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 4.88 - 4.70 (m, 2H), 4.32 (s, 1H), 3.98 - 3.85 (m, 4H), 3.41 - 3.27 (m, 4H), 2.53 (s, 3H), 2.40 (s, 3H), 2.26 - 2.14 (m, 2H), 1.99 (s, 3H), 1.94 - 1.88 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
437		464.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.72 (d, J = 2.0 Hz, 1H), 8.65 (d, J = 1.9 Hz, 1H), 8.31 (s, 1H), 7.02 - 6.88 (m, 2H), 6.77 (d, J = 2.6 Hz, 1H), 5.42 (d, J = 8.1 Hz, 1H), 4.90 - 4.78 (m, 1H), 4.42 (d, J = 7.8 Hz, 1H), 3.99 - 3.85 (m, 4H), 3.43 - 3.27 (m, 4H), 2.37 - 2.13 (m, 2H), 2.10 - 1.88 (m, 6H).
438		437.3	¹ H NMR (300 MHz, Chloroform-d) δ 8.72 (d, J = 2.0 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.28 (s, 1H), 7.72 (s, 1H), 6.94 (dd, J = 16.4, 2.5 Hz, 2H), 5.45 (d, J = 8.2 Hz, 1H), 4.76 (d, J = 6.6 Hz, 1H), 4.26 (s, 1H), 4.01 - 3.85 (m, 7H), 3.44 - 3.28 (m, 4H), 2.30 - 2.11 (m, 2H), 2.08 - 1.82 (m, 6H).
439		435.44	¹ H NMR (400 MHz, CDCl ₃) δ 8.71 (d, J = 1.8 Hz, 1H), 8.63 (d, J = 1.8 Hz, 1H), 8.14 (d, J = 5.9 Hz, 1H), 6.98 (d, J = 2.4 Hz, 1H), 6.17 (d, J = 6.0 Hz, 1H), 5.05 (s, 1H), 4.01 - 3.70 (m, 5H), 3.42 - 3.25 (m, 4H), 2.77 (q, J = 7.6 Hz, 2H), 2.32 - 2.16 (m, 2H), 2.04 - 1.80 (m, 6H), 1.32 (t, J = 7.6 Hz, 3H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
440		436.22	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 7.81 (d, J = 6.3 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.6 Hz, 1H), 6.24 (d, J = 6.2 Hz, 1H), 5.92 (s, 1H), 4.80 (m, 1H), 3.98 (s, 3H), 3.95 - 3.87 (m, 4H), 3.54 (m, 1H), 3.38 - 3.28 (m, 4H), 2.23 (m, 2H), 2.00 - 1.81 (m, 6H).
441		434.35	¹ H NMR (300 MHz, Chloroform-d) rotomers, δ 8.80 (d, 1H), 8.73 (d, 1H), 7.73 (s, 1H), 7.16 - 6.99 (m, 3H), 4.91 (m, 1H), 3.95 (m, 4H), 3.55 (m, 1H), 3.37 (m, 4H), 2.73 (s, 3H), 2.41 (s, 3H), 2.29 (m, 2H), 2.13 (m, 6H).
442		420.4	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.89 (dd, J = 5.0, 1.4 Hz, 1H), 7.19 - 7.10 (m, 1H), 7.03 - 6.95 (m, 2H), 6.92 (d, J = 2.6 Hz, 1H), 4.80 (m, 1H), 3.96 - 3.87 (m, 4H), 3.79 (m, 1H), 3.51 (m, 1H), 3.38 - 3.29 (m, 4H), 2.51 (m, 3H), 2.25 (m, 2H), 2.00 - 1.87 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
443		420.36	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.01 (s, 1H), 7.89 (d, J = 4.8 Hz, 1H), 7.03 (d, J = 4.8 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.5 Hz, 1H), 4.78 (m, 1H), 3.98 - 3.86 (m, 4H), 3.61 (m, 2H), 3.39 - 3.28 (m, 4H), 2.30 - 2.15 (m, 5H), 1.93 (m, 6H).
444		420.28	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 8.03 (dd, J = 2.7, 0.9 Hz, 1H), 7.12 - 7.00 (m, 2H), 6.95 (d, J = 2.5 Hz, 1H), 6.90 (d, J = 2.5 Hz, 1H), 4.79 (m, 1H), 4.08 - 3.87 (m, 5H), 3.49 (m, 1H), 3.38 - 3.28 (m, 4H), 2.53 (s, 3H), 2.21 (m, 2H), 1.84 (d, J = 6.8 Hz, 6H).
445		436.43	¹ H NMR (300 MHz, Chloroform-d) δ 8.65 (m, 1H), 8.40 (m, 1H), 7.50 (dd, J = 4.5, 2.2 Hz, 1H), 7.03 (s, 1H), 6.92 (s, 1H), 6.80 (m, 2H), 4.76 (m, 1H), 4.08 (s, 3H), 3.93 (m, 5H), 3.49 (m, 1H), 3.42 - 3.30 (m, 4H), 2.20 (m, 2H), 1.92 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
446		436.17	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 7.71 (d, J = 2.4 Hz, 1H), 7.65 (d, J = 2.4 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 6.55 (t, J = 2.4 Hz, 1H), 4.79 (m, 1H), 4.13 (m, 1H), 3.92 (m, 4H), 3.85 (s, 3H), 3.48 (m, 1H), 3.38 - 3.29 (m, 4H), 2.22 (m, 2H), 1.98 - 1.88 (m, 6H).
447		436.34	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 2.0 Hz, 1H), 7.70 (s, 1H), 7.15 (m, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.92 (m, 1H), 6.64 (d, J = 8.8 Hz, 1H), 4.80 (m, 1H), 3.90 (m, 7H), 3.45 - 3.26 (m, 5H), 2.24 - 2.12 (m, 2H), 1.98 - 1.81 (m, 6H).
448		421.39	¹ H NMR (300 MHz, Chloroform-d) δ 8.78 (dd, J = 4.0, 2.0 Hz, 1H), 8.66 (d, J = 2.1 Hz, 1H), 8.55 (s, 1H), 8.06 (s, 1H), 7.02 (d, J = 2.5 Hz, 1H), 6.96 (d, J = 2.5 Hz, 1H), 4.88 (m, 1H), 3.93 (m, 4H), 3.55 (m, 1H), 3.40 (m, 4H), 3.29 (m, 1H), 2.58 (s, 3H), 2.28 (m, 2H), 2.07 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
449		435.35	¹ H NMR (300 MHz, Chloroform-d) rotomers, δ 8.85 (d, 1H), 8.67 (d, 1H), 7.92 (d, 1H), 7.07 - 6.96 (m, 2H), 4.91 (m, 1H), 4.01 - 3.88 (m, 4H), 3.53 - 3.24 (m, 5H), 2.87 (s, 3H), 2.64 (s, 3H), 2.04 - 1.76 (m, 6H).
450		421.36	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.21 (d, J = 9.2 Hz, 1H), 6.99 - 6.87 (m, 3H), 4.87 (m, 1H), 3.98 - 3.82 (m, 5H), 3.39 - 3.29 (m, 4H), 2.51 (s, 3H), 2.30 - 2.22 (m, 2H), 2.11 - 1.84 (m, 6H).
451		421.56	¹ H NMR (300 MHz, Chloroform-d) rotomers, δ 8.92 (d, J = 1.8 Hz, 1H), 8.77 (d, J = 1.8 Hz, 1H), 8.30 (m, 2H), 7.02 - 6.90 (m, 2H), 4.86 (m, 1H), 3.94 (m, 4H), 3.52 - 3.26 (m, 5H), 2.78 (s, 3H), 2.26 (d, J = 11.8 Hz, 2H), 1.90 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
452		463.35	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 5.01 (s, 2H), 4.91 (t, J = 2.5 Hz, 2H), 4.79 (s, 1H), 4.49 (s, 1H), 4.16 (s, 1H), 3.99 - 3.82 (m, 4H), 3.40 - 3.25 (m, 4H), 2.53 (s, 3H), 2.29 - 2.15 (m, 2H), 1.96 - 1.80 (m, 6H).
453		450.43	¹ H NMR (300 MHz, CDCl ₃) δ 8.69 (s, 1H), 8.62 (s, 1H), 7.89 (d, J = 5.9 Hz, 1H), 6.94 (s, 1H), 6.89 (s, 1H), 5.64 (d, J = 6.0 Hz, 1H), 4.87 - 4.66 (m, 2H), 3.98 - 3.83 (m, 4H), 3.44 - 3.24 (m, 4H), 3.13 (s, 6H), 2.26 - 2.10 (m, 2H), 2.04 - 1.75 (m, 6H).
454		436.34	¹ H NMR (300 MHz, CDCl ₃) δ 8.61 (d, J = 1.8 Hz, 1H), 8.52 (d, J = 1.9 Hz, 1H), 7.69 (s, 1H), 6.86 (d, J = 2.4 Hz, 1H), 6.80 (d, J = 2.2 Hz, 1H), 5.62 (d, J = 6.0 Hz, 1H), 4.83 (s, 1H), 4.69 (s, 1H), 3.91 - 3.74 (m, 4H), 3.36 - 3.10 (m, 4H), 2.86 (d, J = 5.0 Hz, 3H), 2.22 - 2.03 (m, 2H), 1.91 - 1.72 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
455		447.41	¹ H NMR (300 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.05 (d, J = 5.2 Hz, 1H), 6.94 (d, J = 2.5 Hz, 1H), 6.89 (d, J = 2.5 Hz, 1H), 6.39 (d, J = 5.2 Hz, 1H), 5.13 (bd, 1H), 4.76 (s, 1H), 4.10 - 3.96 (m, 1H), 3.95 - 3.83 (m, 4H), 3.40 - 3.24 (m, 4H), 2.26 - 2.09 (m, 2H), 1.97 - 1.76 (m, 7H), 1.10 - 1.00 (m, 2H), 1.00 - 0.91 (m, 2H).
456		465.29	¹ H NMR (300 MHz, CDCl ₃) δ 8.72 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.23 (d, J = 1.8 Hz, 1H), 6.97 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.4 Hz, 1H), 5.04 (d, J = 7.6 Hz, 1H), 4.81 (s, 1H), 4.21 (s, 1H), 4.02 - 3.82 (m, 4H), 3.44 - 3.26 (m, 4H), 2.33 - 2.10 (m, 3H), 2.06 - 1.82 (m, 6H), 1.19 - 1.11 (m, 2H), 1.06 - 0.97 (m, 2H).
457		437.38	¹ H NMR (300 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.21 (d, J = 5.1 Hz, 1H), 6.95 (d, J = 2.5 Hz, 1H), 6.90 (d, J = 2.5 Hz, 1H), 6.45 (d, J = 5.1 Hz, 1H), 5.33 (d, J = 7.6 Hz, 1H), 4.85 - 4.74 (m, 1H), 4.56 (s, 2H), 4.12 - 4.00 (m, 1H), 3.95 - 3.86 (m, 4H), 3.66 (s, 1H), 3.39 - 3.27 (m, 4H), 2.18 (dd, J = 14.1, 8.4 Hz, 2H), 2.01 - 1.83 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
458		489.41	¹ H NMR (300 MHz, CDCl ₃) δ 8.97 (s, 2H), 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 5.54 (d, J = 8.0 Hz, 1H), 4.81 (s, 1H), 4.38 (s, 3H), 4.16 - 4.05 (m, 1H), 4.00 - 3.81 (m, 4H), 3.42 - 3.23 (m, 4H), 2.29 - 2.15 (m, 2H), 2.05 - 1.83 (m, 6H).
459		489.41	¹ H NMR (300 MHz, CDCl ₃) δ 8.80 - 8.64 (m, 3H), 8.62 (d, J = 1.9 Hz, 1H), 6.97 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 5.69 (d, J = 7.6 Hz, 1H), 4.82 (s, 1H), 4.18 (s, 3H), 4.15 - 4.06 (m, 1H), 4.01 - 3.83 (m, 4H), 3.44 - 3.24 (m, 4H), 2.32 - 2.18 (m, 2H), 2.07 - 1.84 (m, 6H).
460		475.37	¹ H NMR (300 MHz, CDCl ₃) δ 9.02 (s, 2H), 8.84 (d, J = 2.0 Hz, 1H), 8.66 (d, J = 2.0 Hz, 1H), 7.01 (d, J = 2.3 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 5.90 (d, J = 7.1 Hz, 1H), 4.74 (s, 1H), 4.13 (s, 1H), 4.04 - 3.88 (m, 4H), 3.45 - 3.32 (m, 4H), 2.06 (d, J = 7.5 Hz, 2H), 1.96 - 1.84 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
461		475.19	¹ H NMR (300 MHz, CDCl ₃) δ 8.72 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.59 (s, 1H), 8.54 (bd, 1H), 7.37 (d, J = 4.9 Hz, 1H), 6.97 (d, J = 2.5 Hz, 1H), 6.93 (d, J = 2.1 Hz, 1H), 5.55 (bd, 1H), 4.83 (s, 1H), 4.14 (s, 1H), 4.00 - 3.89 (m, 4H), 3.42 - 3.29 (m, 4H), 2.30 - 2.18 (m, 2H), 1.96 (s, 6H).
462		473.39	¹ H NMR (300 MHz, CDCl ₃) δ 8.69 (d, J = 2.5 Hz, 3H), 8.62 (d, J = 1.9 Hz, 1H), 7.62 (d, J = 2.3 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 6.52 (d, J = 2.1 Hz, 1H), 5.37 (d, J = 7.9 Hz, 1H), 4.86 - 4.75 (m, 1H), 4.14 - 4.01 (m, 1H), 3.99 - 3.83 (m, 4H), 3.41 - 3.26 (m, 4H), 2.28 - 2.15 (m, 2H), 2.01 - 1.86 (m, 6H).
463		453.4	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 4.95 (d, J = 7.1 Hz, 1H), 4.78 (s, 1H), 4.23 (d, J = 3.8 Hz, 1H), 3.99 - 3.82 (m, 4H), 3.42 - 3.25 (m, 4H), 2.45 (d, J = 0.6 Hz, 3H), 2.30 (d, J = 2.8 Hz, 3H), 2.26 - 2.14 (m, 2H), 1.99 - 1.86 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
464		449.35	¹ H NMR (300 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.22 (s, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 4.78 (d, J = 7.8 Hz, 2H), 4.62 (t, J = 9.1 Hz, 2H), 4.23 (s, 1H), 3.97 - 3.85 (m, 4H), 3.40 - 3.30 (m, 4H), 3.25 (t, J = 9.1 Hz, 2H), 2.27 - 2.11 (m, 2H), 2.01 - 1.82 (m, 6H).
465		420.19	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.86 (d, J = 2.7 Hz, 1H), 7.78 (s, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.5 Hz, 1H), 6.78 (t, J = 2.1 Hz, 1H), 4.78 (m, 1H), 4.00 - 3.86 (m, 4H), 3.50 (s, 1H), 3.38 - 3.28 (m, 4H), 2.28 (s, 3H), 2.25 - 2.14 (m, 2H), 1.95 - 1.85 (m, 6H).
466		461.33	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (m, 1H), 8.60 (d, J = 1.9 Hz, 1H), 8.20 (m, 1H), 7.93 (s, 1H), 6.97 - 6.89 (m, 3H), 4.77 (m, 1H), 3.92 (m, 4H), 3.49 (m, 1H), 3.34 (m, 4H), 2.20 (m, 2H), 1.88 (d, J = 4.1 Hz, 6H), 1.70 (m, 4H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
467		435.35	¹ H NMR (400 MHz, CDCl ₃) δ 8.68 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.95 (s, 1H), 6.94 (d, J = 2.5 Hz, 1H), 6.90 (d, J = 2.5 Hz, 1H), 5.00 (d, J = 7.9 Hz, 1H), 4.83 - 4.74 (m, 1H), 4.07 - 3.96 (m, 1H), 3.96 - 3.87 (m, 4H), 3.38 - 3.28 (m, 4H), 2.29 (s, 3H), 2.22 - 2.12 (m, 2H), 2.07 (s, 3H), 1.93 - 1.85 (m, 6H).
468		478.21	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.91 (d, J = 2.7 Hz, 1H), 7.73 (dd, J = 8.6, 0.7 Hz, 1H), 7.11 (dd, J = 8.6, 2.8 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.5 Hz, 1H), 5.11 - 5.02 (m, 2H), 4.85 - 4.75 (m, 1H), 4.75 - 4.67 (m, 2H), 3.98 - 3.86 (m, 4H), 3.54 (s, 1H), 3.40 - 3.27 (m, 4H), 2.31 - 2.15 (m, 2H), 2.00 - 1.83 (m, 6H).
469		454.3	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.03 (s, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 4.77 (s, 1H), 4.67 (s, 1H), 4.60 (s, 1H), 4.16 (s, 1H), 4.00 - 3.82 (m, 4H), 3.41 - 3.25 (m, 4H), 3.04 (d, J = 5.0 Hz, 3H), 2.28 - 2.11 (m, 2H), 1.99 - 1.84 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
470		446.38	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.88 (d, J = 5.5 Hz, 1H), 7.61 (d, J = 2.1 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 6.84 (d, J = 5.5 Hz, 1H), 6.70 (d, J = 2.1 Hz, 1H), 4.92 (s, 1H), 4.82 - 4.74 (m, 1H), 4.44 - 4.28 (m, 1H), 3.95 - 3.86 (m, 4H), 3.39 - 3.26 (m, 4H), 2.28 - 2.17 (m, 2H), 2.06 - 1.92 (m, 6H).
471		421.35	¹ H NMR (300 MHz, Chloroform-d) δ 8.62 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 2.0 Hz, 1H), 8.42 (s, 1H), 7.91 (d, J = 13.5 Hz, 1H), 6.93 - 6.78 (m, 2H), 4.71 (s, 1H), 4.57 (d, J = 7.9 Hz, 1H), 4.22 (s, 1H), 3.91 - 3.78 (m, 4H), 3.33 - 3.19 (m, 4H), 2.16 (d, J = 6.6 Hz, 2H), 1.96 (s, 3H), 1.91 - 1.77 (m, 6H).
472		534.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.62 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 8.10 (d, J = 0.8 Hz, 1H), 6.92 - 6.77 (m, 2H), 5.36 (dd, J = 7.9, 0.9 Hz, 1H), 4.71 (q, J = 5.2, 4.2 Hz, 2H), 3.97 - 3.69 (m, 7H), 3.58 - 3.37 (m, 4H), 3.30 - 3.03 (m, 6H), 2.22 - 2.04 (m, 2H), 1.93 - 1.73 (m, 6H), 1.52 (dtd, J = 12.8, 8.9, 3.8 Hz, 2H), 1.28 - 1.16 (m, 3H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
473		462.3	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 2.0 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.16 (s, 1H), 6.92 (dd, J = 15.4, 2.5 Hz, 2H), 4.77 (d, J = 5.4 Hz, 1H), 4.25 - 4.07 (m, 2H), 3.98 - 3.86 (m, 5H), 3.51 (dd, J = 9.3, 8.2 Hz, 2H), 3.40 - 3.29 (m, 5H), 2.92 (s, 3H), 2.87 - 2.71 (m, 2H), 2.19 (d, J = 7.1 Hz, 2H), 1.91 (t, J = 4.0 Hz, 6H).
474		523.31	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.99 (d, J = 1.5 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 4.82 (d, J = 5.5 Hz, 1H), 4.77 (s, 1H), 4.15 (s, 1H), 3.96 - 3.85 (m, 4H), 3.76 - 3.65 (m, 4H), 3.38 - 3.26 (m, 4H), 2.59 - 2.45 (m, 4H), 2.34 (s, 3H), 2.27 - 2.13 (m, 2H), 1.96 - 1.85 (m, 6H).
475		449.21	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 8.52 (s, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 5.01 (d, J = 2.6 Hz, 2H), 4.98 - 4.89 (m, 2H), 4.80 (s, 1H), 4.54 (s, 1H), 4.22 (s, 1H), 3.95 - 3.86 (m, 4H), 3.39 - 3.27 (m, 4H), 2.29 - 2.16 (m, 2H), 1.98 - 1.84 (m, 6H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
476		439.4	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.03 (d, J = 1.7 Hz, 1H), 6.97 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 5.18 - 5.04 (m, 1H), 4.80 (s, 1H), 3.99 - 3.90 (m, 5H), 3.40 - 3.32 (m, 4H), 2.36 (d, J = 2.4 Hz, 3H), 2.23 - 2.15 (m, 2H), 1.95 - 1.85 (m, 6H).
477		439.36	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.26 (d, J = 1.7 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 5.07 (d, J = 7.2 Hz, 1H), 4.80 (s, 1H), 4.30 - 4.13 (m, 1H), 3.98 - 3.83 (m, 4H), 3.38 - 3.26 (m, 4H), 2.36 (d, J = 2.8 Hz, 3H), 2.29 - 2.16 (m, 2H), 1.97 - 1.85 (m, 6H).
478		424	¹ H NMR (400 MHz, CDCl ₃) δ 8.64 (t, J = 9.6 Hz, 1H), 8.53 (d, J = 1.9 Hz, 1H), 8.02 (d, J = 5.9 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 6.83 (d, J = 2.4 Hz, 1H), 6.07 (d, J = 6.0 Hz, 1H), 4.91 (s, 1H), 4.73 (s, 1H), 3.83 (dd, J = 17.8, 12.9 Hz, 5H), 3.25 (dd, J = 17.8, 13.0 Hz, 4H), 2.14 (dd, J = 8.9, 5.0 Hz, 2H), 1.91 - 1.76 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
479		449.3	¹ H NMR (300 MHz, Chloroform-d) δ 8.62 (d, J = 1.9 Hz, 1H), 8.53 (d, J = 1.9 Hz, 1H), 8.15 (s, 1H), 6.84 (dd, J = 17.3, 2.5 Hz, 2H), 4.70 (s, 1H), 4.55 (dd, J = 9.2, 8.3 Hz, 2H), 4.36 (s, 1H), 4.16 (s, 1H), 3.92 - 3.76 (m, 4H), 3.03 - 2.83 (m, 2H), 2.14 (s, 2H), 1.84 (d, J = 5.4 Hz, 6H).
480		478.26	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 2.0 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.20 (d, J = 0.9 Hz, 1H), 6.93 (dd, J = 17.6, 2.5 Hz, 2H), 5.65 (d, J = 0.9 Hz, 1H), 5.54 - 5.38 (m, 1H), 4.97 (d, J = 8.0 Hz, 1H), 4.80 (s, 1H), 4.04 - 3.85 (m, 5H), 3.81 - 3.62 (m, 3H), 3.41 - 3.24 (m, 4H), 2.62 (s, 1H), 2.31 - 2.11 (m, 2H), 1.90 (dd, J = 8.0, 3.5 Hz, 6H).
481		451.41	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.18 (s, 2H), 6.99 - 6.87 (m, 2H), 5.16 (d, J = 8.1 Hz, 1H), 4.78 (s, 1H), 4.03 - 3.87 (m, 5H), 3.80 (t, J = 6.5 Hz, 2H), 3.39 - 3.29 (m, 4H), 2.67 (t, J = 6.5 Hz, 2H), 2.26 - 2.14 (m, 2H), 1.90 (t, J = 6.5 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
482		454.35	¹ H NMR (400 MHz, CDCl ₃) δ 8.68 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.65 (d, J = 3.6 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 5.14 (s, 1H), 4.90 (s, 1H), 4.76 (s, 1H), 4.01 - 3.87 (m, 5H), 3.39 - 3.31 (m, 4H), 3.01 (d, J = 5.0 Hz, 3H), 2.23 - 2.12 (m, 2H), 1.99 - 1.81 (m, 6H).
483		464.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.81 - 1.98 (m, 6 H), 2.11 - 2.23 (m, 2 H), 2.29 (s, 6 H), 3.29 - 3.38 (m, 6 H), 3.89 - 3.94 (m, 4 H), 3.98 - 4.08 (m, 1 H), 4.75 - 4.82 (m, 1 H), 5.21 (d, J = 7.9 Hz, 1 H), 6.61 (d, J = 5.1 Hz, 1 H), 6.90 (d, J = 2.3 Hz, 1 H)
484		519.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.79 - 1.98 (m, 6 H), 2.13 - 2.26 (m, 2 H), 2.34 (s, 3 H), 2.46 - 2.78 (m, 8 H), 3.28 - 3.38 (m, 4 H), 3.60 (s, 2 H), 3.78 - 3.98 (m, 5 H), 4.76 - 4.85 (m, 1 H), 5.10 (br s, 1 H), 6.18 (d, J = 6.0 Hz, 1 H), 6.90 (d, J = 2.4 H)




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
485		441.29	¹ H NMR (400 MHz, CDCl ₃) δ 8.68 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.20 (s, 2H), 6.96 (d, J = 2.3 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 5.27 (d, J = 7.4 Hz, 1H), 4.79 (s, 1H), 4.01 - 3.84 (m, 5H), 3.40 - 3.25 (m, 4H), 2.25 - 2.12 (m, 2H), 1.97 - 1.82 (m, 6H).
486		407	¹ H NMR (400 MHz, CDCl ₃) δ 8.62 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 8.49 (s, 1H), 8.08 (d, J = 6.0 Hz, 1H), 6.89 (d, J = 2.5 Hz, 1H), 6.83 (d, J = 2.5 Hz, 1H), 6.24 (dd, J = 6.0, 1.1 Hz, 1H), 4.88 (s, 1H), 4.78 - 4.67 (m, 1H), 3.87 - 3.82 (m, 4H), 3.25 (dd, J = 13.6, 8.8 Hz, 4H), 2.15 (dd, J = 8.8, 5.1 Hz, 2H), 1.90 - 1.78 (m, 6H).
487		447.37	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.45 (s, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.87 (t, J = 9.8 Hz, 1H), 4.77 (s, 1H), 4.49 (d, J = 8.0 Hz, 1H), 4.28 (s, 1H), 3.96 - 3.86 (m, 4H), 3.40 - 3.28 (m, 4H), 2.95 - 2.85 (m, 2H), 2.73 - 2.61 (m, 2H), 2.28 - 2.17 (m, 2H), 2.17 - 2.06 (m, 2H), 1.94 - 1.89 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
488		447.37	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.48 (s, 1H), 7.73 (d, J = 2.1 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.4 Hz, 1H), 6.85 (d, J = 2.1 Hz, 1H), 5.29 (d, J = 8.3 Hz, 1H), 4.88 - 4.75 (m, 1H), 4.46 - 4.33 (m, 1H), 3.95 - 3.87 (m, 4H), 3.38 - 3.30 (m, 4H), 2.31 - 2.19 (m, 2H), 2.06 - 1.91 (m, 6H).
489		450.34	¹ H NMR (400 MHz, CDCl ₃) δ 8.68 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.85 (d, J = 6.1 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 5.81 (d, J = 6.1 Hz, 1H), 4.92 (s, 1H), 4.75 (s, 1H), 4.02 (s, 1H), 3.97 - 3.86 (m, 4H), 3.32 (dd, J = 17.6, 12.8 Hz, 4H), 3.04 (s, 6H), 2.18 (dd, J = 24.2, 17.3 Hz, 2H).
490		435.35	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.87 (d, J = 0.7 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 4.78 (s, 1H), 4.67 (s, 1H), 4.41 - 4.28 (m, 1H), 3.96 - 3.88 (m, 4H), 3.38 - 3.30 (m, 4H), 2.52 (s, 3H), 2.28 - 2.16 (m, 2H), 2.00 (s, 3H), 1.98 - 1.90 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
491		435.35	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.7 Hz, 1H), 8.61 (d, J = 1.7 Hz, 1H), 8.42 (s, 1H), 6.95 (s, 1H), 6.90 (s, 1H), 4.79 (s, 2H), 4.28 (s, 1H), 3.98 - 3.86 (m, 4H), 3.40 - 3.28 (m, 4H), 2.42 (s, 3H), 2.30 - 2.16 (m, 2H), 2.02 (s, 3H), 1.99 - 1.89 (m, 6H).
492		506.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.20 (d, J = 0.8 Hz, 1H), 6.97 (d, J = 2.4 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 5.47 (d, J = 1.0 Hz, 1H), 4.76 (d, J = 23.9 Hz, 2H), 4.12 - 3.75 (m, 7H), 3.41 - 3.28 (m, 4H), 3.20 (ddd, J = 13.2, 9.6, 3.2 Hz, 2H), 2.20 (d, J = 6.9 Hz, 2H), 1.91 (d, J = 5.2 Hz, 6H).
493		534	¹ H NMR (400 MHz, DMSO) δ 8.72 (d, J = 1.9 Hz, 1H), 8.57 (d, J = 1.9 Hz, 1H), 7.77 (s, 1H), 7.45 (s, 1H), 7.14 (d, J = 2.4 Hz, 1H), 6.83 (d, J = 2.3 Hz, 1H), 6.15 (d, J = 5.8 Hz, 1H), 4.92 (s, 1H), 4.26 (t, J = 6.1 Hz, 2H), 4.09 - 3.89 (m, 1H), 3.83 - 3.72 (m, 4H), 3.34 (d, J = 7.7 Hz, 4H), 2.57 (t, J = 6.1 Hz, 2H), 2.36 (d, J = 24.2 Hz, 4H), 2.01 (d, J = 17.5 Hz, 2H), 1.76 (d, J = 4.8 Hz, 6H), 1.52 - 1.35 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
494		439.23	¹ H NMR (300 MHz, Chloroform-d) δ 8.62 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 7.82 (d, J = 3.4 Hz, 1H), 6.88 (d, J = 2.5 Hz, 1H), 6.82 (d, J = 2.5 Hz, 1H), 4.96 (d, J = 8.0 Hz, 1H), 4.72 (d, J = 3.1 Hz, 1H), 4.18 (s, 1H), 3.90 - 3.78 (m, 4H), 3.35 - 3.19 (m, 4H), 2.41 (d, J = 0.9 Hz, 3H), 2.13 (q, J = 6.4 Hz, 2H), 1.84 (t, J = 6.3 Hz, 6H).
495		504.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.37 - 1.47 (m, 2 H), 1.51 - 1.60 (m, 4 H), 1.82 - 1.99 (m, 6 H), 2.14 - 2.24 (m, 2 H), 2.30 - 2.41 (m, 4 H), 3.29 (s, 2 H), 3.30 - 3.35 (m, 4 H), 3.89 - 3.93 (m, 4 H), 3.96 - 4.06 (m, 1 H), 4.75 - 4.81 (m, 1 H), 5.18 (d, J =
496		504.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.38 - 1.49 (m, 2 H), 1.56 - 1.64 (m, 4 H), 1.79 - 1.97 (m, 6 H), 2.15 - 2.26 (m, 2 H), 2.45 - 2.57 (m, 4 H), 3.28 - 3.38 (m, 4 H), 3.53 (s, 2 H), 3.68 - 3.99 (m, 5 H), 4.76 - 4.85 (m, 1 H), 5.11 (br s, 1 H), 6.18 (d, J = 6.0




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
497		519.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.22 (s, 2H), 7.01 - 6.85 (m, 2H), 5.21 (d, J = 8.1 Hz, 1H), 4.79 (d, J = 5.3 Hz, 1H), 4.11 - 3.87 (m, 5H), 3.35 (q, J = 3.4 Hz, 6H), 2.46 (s, 6H), 2.29 (s, 3H), 2.19 (s, 3H), 2.01 - 1.83 (m, 6H).
498		446.1	¹ H NMR (300 MHz, Chloroform-d) δ 8.72 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.23 (dd, J = 7.6, 0.8 Hz, 1H), 7.85 (d, J = 2.2 Hz, 1H), 6.97 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 6.12 (dd, J = 2.2, 0.8 Hz, 1H), 6.00 (d, J = 7.6 Hz, 1H), 4.91 (d, J = 7.9 Hz, 1H), 4.81 (s, 1H), 4.20 (s, 1H), 4.01 - 3.88 (m, 4H), 3.42 - 3.28 (m, 4H), 2.22 (s, 2H), 1.97 (dd, J = 7.8, 5.5 Hz, 6H).
499		535.24	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.19 (d, J = 0.8 Hz, 1H), 6.93 (dd, J = 16.4, 2.5 Hz, 2H), 5.42 (d, J = 1.0 Hz, 1H), 4.87 - 4.67 (m, 2H), 4.01 - 3.77 (m, 5H), 3.61 (dt, J = 27.3, 5.1 Hz, 6H), 3.41 - 3.26 (m, 4H), 2.73 (t, J = 5.3 Hz, 1H), 2.59 (ddd, J = 6.2, 5.0, 3.7 Hz, 6H), 2.19 (d, J = 6.6 Hz, 2H), 1.89 (t, J = 5.2 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
500		461.24	¹ H NMR (300 MHz, Chloroform-d) δ 8.64 (d, J = 1.9 Hz, 1H), 8.55 (d, J = 1.9 Hz, 1H), 8.21 (s, 1H), 6.88 (dd, J = 16.1, 2.5 Hz, 2H), 6.16 (d, J = 8.3 Hz, 1H), 5.98 (s, 1H), 4.80 (d, J = 2.9 Hz, 1H), 3.90 - 3.77 (m, 4H), 3.72 - 3.51 (m, 1H), 3.33 - 3.20 (m, 4H), 2.51 (s, 3H), 2.34 - 1.66 (m, 8H).
501		480.15	¹ H NMR (300 MHz, Chloroform-d) δ 8.63 (d, J = 1.9 Hz, 1H), 8.55 (d, J = 1.9 Hz, 1H), 6.95 (dd, J = 3.6, 2.3 Hz, 1H), 6.86 (dd, J = 13.7, 2.4 Hz, 2H), 6.30 (dd, J = 3.6, 2.0 Hz, 1H), 5.22 (m, 1H), 4.74 (s, 1H), 4.30 (s, 1H), 3.91 - 3.79 (m, 4H), 3.35 - 3.19 (m, 4H), 2.15 (m, 2H), 2.03 - 1.76 (m, 6H).
502		462.19	¹ H NMR (300 MHz, Chloroform-d) δ 8.63 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 2.0 Hz, 1H), 8.25 (s, 1H), 6.88 (d, J = 2.4 Hz, 1H), 6.82 (d, J = 2.5 Hz, 1H), 4.73 (s, 1H), 4.54 (s, 1H), 4.12 (s, 1H), 3.84 (dd, J = 6.0, 3.7 Hz, 4H), 3.37 - 3.19 (m, 5H), 2.17 (d, J = 10.1 Hz, 2H), 1.92 - 1.71 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
503		506.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.79 ñ 1.99 (m, 6 H), 2.15 - 2.26 (m, 2 H), 2.54 - 2.65 (m, 4 H), 3.29 - 3.37 (m, 4 H), 3.58 (s, 2 H), 3.74 - 3.83 (m, 5 H), 3.88 - 3.95 (m, 4 H), 4.75 - 4.85 (m, 1 H), 5.11 (br. s, 1 H), 6.19 (d, J = 5.9 Hz, 1 H), 6.90 (d, J
504		504.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.40 - 1.49 (m, 2 H), 1.55 - 1.65 (m, 4 H), 1.84 - 1.96 (m, 6 H), 2.12 - 2.23 (m, 2 H), 2.43 (br. s, 4 H), 3.30 - 3.40 (m, 6 H), 3.88 - 3.95 (m, 4 H), 3.97 - 4.08 (m, 1 H), 4.75 - 4.82 (m, 1 H), 5.17 (d, J = 7.9 Hz, 1 H), 6.68
505		506.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.83 - 1.96 (m, 6 H), 2.13 - 2.23 (m, 2 H), 2.53 (br. s, 4 H), 3.30 - 3.36 (m, 4 H), 3.42 (s, 2 H), 3.72 - 3.78 (m, 4 H), 3.88 - 3.95 (m, 4 H), 4.03 (br. s, 1 H), 4.75 - 4.82 (m, 1 H), 5.21 (d, J = 8.0 Hz, 1 H), 6.67 (d, J = 5)




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
506		446.3	¹ H NMR (300 MHz, DMSO-d ₆): ppm 1.71 - 1.85 (m, 6 H), 1.98 - 2.10 (m, 2 H), 3.33 - 3.38 (m, 4 H), 3.74 - 3.91 (m, 5 H), 4.87 - 4.96 (m, 1 H), 6.02 (d, J = 7.2 Hz, 1 H), 6.51 (s, 1 H), 6.83 (d, J = 1.9 Hz, 1 H), 7.14 (d, J = 1.9 Hz, 1 H), 7.94 (s, 1 H), 8.3
507		548.4	¹ H NMR (300 MHz, CDCl ₃): ppm 1.39 - 1.48 (m, 2 H), 1.54 - 1.69 (m, 4 H), 1.83 - 1.98 (m, 8 H), 2.12 - 2.24 (m, 2 H), 2.33 - 2.47 (m, 6 H), 3.31 - 3.35 (m, 4 H), 3.89 - 3.99 (m, 7 H), 4.74 - 4.81 (m, 1 H), 4.95 (d, J = 8.2 Hz, 1 H), 6.90 (d, J = 2.3 Hz, 1
508		563.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.82 - 1.98 (m, 8 H), 2.13 - 2.23 (m, 2 H), 2.29 (s, 3 H), 2.35 - 2.62 (m, 10 H), 3.29 - 3.37 (m, 4 H), 3.87 - 3.94 (m, 5 H), 3.97 (t, J = 6.3 Hz, 2 H), 4.74 - 4.81 (m, 1 H), 4.96 (d, J = 8.1 Hz, 1 H), 6.90 (d, J = 2.3 Hz, 1 H




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
509		520.3	¹ H NMR (300 MHz, CDCl ₃) ppm 1.77 - 1.96 (m, 10 H), 2.11 - 2.25 (m, 2 H), 2.57 - 2.65 (m, 4 H), 2.86 (t, J = 5.7 Hz, 2 H), 3.31 - 3.35 (m, 4 H), 3.87 - 3.98 (m, 5 H), 4.05 (t, J = 5.7 Hz, 2 H), 4.74 - 4.81 (m, 1 H), 4.98 (d, J = 8.2 Hz, 1 H), 6.90 (d, J =
510		447.16	¹ H NMR (300 MHz, Chloroform-d) δ 8.95 (d, J = 1.3 Hz, 1H), 8.72 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.14 (s, 1H), 6.95 (dd, J = 16.0, 2.5 Hz, 2H), 6.36 (d, J = 1.3 Hz, 1H), 5.11 (d, J = 7.8 Hz, 1H), 4.83 (dd, J = 5.5, 2.9 Hz, 1H), 4.01 - 3.84 (m, 4H), 3.73 - 3.55 (m, 1H), 3.40 - 3.30 (m, 4H), 2.34 - 2.21 (m, 2H), 2.08 - 1.83 (m, 6H).
511		510.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.63 (d, J = 1.9 Hz, 1H), 8.55 (d, J = 1.9 Hz, 1H), 7.93 (d, J = 1.7 Hz, 1H), 6.85 (dd, J = 16.5, 2.5 Hz, 2H), 4.74 (ddd, J = 21.7, 7.5, 3.9 Hz, 2H), 4.09 (p, J = 7.7, 7.2 Hz, 1H), 3.92 - 3.79 (m, 4H), 3.71 (dd, J = 5.7, 3.7 Hz, 4H), 3.57 (dd, J = 5.7, 3.7 Hz, 4H), 3.34 - 3.20 (m, 4H), 2.15 (td, J = 11.0, 10.0, 6.2 Hz, 2H), 1.94 - 1.79 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
512		468.27	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 7.99 (d, J = 1.8 Hz, 1H), 7.00 - 6.85 (m, 2H), 4.77 (d, J = 6.6 Hz, 2H), 4.14 (q, J = 7.2 Hz, 1H), 4.03 - 3.87 (m, 4H), 3.78 - 3.64 (m, 1H), 3.43 - 3.28 (m, 4H), 3.15 (d, J = 2.5 Hz, 6H), 2.98 - 2.84 (m, 1H), 2.20 (d, J = 11.9 Hz, 2H), 1.94 (p, J = 5.8, 5.3 Hz, 6H).
513		480.1	¹ H NMR (300 MHz, Chloroform-d) δ 10.85 (d, J = 3.0 Hz, 1H), 8.66 (d, J = 1.9 Hz, 1H), 8.40 (d, J = 2.0 Hz, 1H), 7.22 (t, J = 2.9 Hz, 1H), 6.85 (d, J = 2.4 Hz, 1H), 6.75 (d, J = 2.4 Hz, 1H), 6.36 (dd, J = 3.1, 1.9 Hz, 1H), 5.86 (d, J = 7.8 Hz, 1H), 4.59 (s, 1H), 4.25 (s, 1H), 3.82 (dd, J = 6.0, 3.7 Hz, 4H), 3.24 (dd, J = 6.0, 3.8 Hz, 4H), 1.71 (s, 11H), 1.51 (t, J = 10.1 Hz, 2H).
514		480.1	¹ H NMR (300 MHz, Chloroform-d) δ 10.96 (s, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.40 (d, J = 1.9 Hz, 1H), 6.84 (d, J = 2.4 Hz, 1H), 6.75 (d, J = 2.4 Hz, 1H), 6.34 (dd, J = 3.1, 1.9 Hz, 1H), 6.02 (d, J = 7.8 Hz, 1H), 4.59 (s, 1H), 4.23 (td, J = 8.8, 8.3, 4.2 Hz, 1H), 3.89 - 3.74 (m, 4H), 3.30 - 3.03 (m, 4H), 1.89 - 1.61 (m, 6H), 1.61 - 1.39 (m, 2H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
515		446.42	¹ H NMR (300 MHz, Chloroform-d) δ 9.53 (s, 1H), 8.71 (d, J = 2.0 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.35 (s, 1H), 7.06 (dd, J = 3.6, 2.1 Hz, 1H), 7.00 - 6.85 (m, 2H), 6.39 (dd, J = 3.6, 1.8 Hz, 1H), 5.14 (d, J = 8.3 Hz, 1H), 4.82 (s, 1H), 4.40 (d, J = 8.0 Hz, 1H), 4.01 - 3.81 (m, 4H), 3.44 - 3.25 (m, 4H), 2.26 (d, J = 9.8 Hz, 2H), 2.08 - 1.86 (m, 6H).
516		447.32	¹ H NMR (300 MHz, Chloroform-d) δ 12.29 (s, 1H), 8.71 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.43 (s, 1H), 7.95 (s, 1H), 7.00 - 6.89 (m, 2H), 6.03 (s, 1H), 4.86 (s, 1H), 4.42 (s, 1H), 3.92 (dd, J = 6.0, 3.7 Hz, 4H), 3.40 - 3.30 (m, 4H), 2.27 (d, J = 10.1 Hz, 2H), 2.07 - 1.88 (m, 6H).
517		446.33	¹ H NMR (300 MHz, Chloroform-d) δ 11.28 (s, 1H), 8.75 (d, J = 1.9 Hz, 1H), 8.53 (d, J = 2.0 Hz, 1H), 8.47 (s, 1H), 7.34 (s, 1H), 6.93 (d, J = 2.3 Hz, 1H), 6.82 (d, J = 2.5 Hz, 1H), 1.93 - 1.71 (m, 6H), 6.54 (d, J = 3.0 Hz, 1H), 5.57 (d, J = 7.9 Hz, 1H), 4.64 (s, 1H), 4.33 (s, 1H), 3.90 (t, J = 4.8 Hz, 4H), 3.31 (t, J = 4.9 Hz, 4H), 1.80-1.68 (m, 6H), 1.54 (d, J = 9.2 Hz, 2H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
518		447.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.63 (d, J = 1.9 Hz, 1H), 8.55 (d, J = 1.9 Hz, 1H), 8.32 (s, 1H), 7.42 (d, J = 2.5 Hz, 1H), 6.88 (d, J = 2.5 Hz, 1H), 6.83 (d, J = 2.5 Hz, 1H), 6.58 (d, J = 2.5 Hz, 1H), 5.07 (s, 1H), 4.74 (s, 1H), 4.26 (s, 1H), 3.84 (dd, J = 6.0, 3.7 Hz, 4H), 3.42 - 3.10 (m, 4H), 2.18 (d, J = 10.2 Hz, 2H), 2.06 - 1.70 (m, 6H).
519		543.21	¹ H NMR (300 MHz, Chloroform-d) δ 8.61 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 8.07 (s, 2H), 6.93 - 6.79 (m, 2H), 5.04 (d, J = 8.1 Hz, 1H), 4.70 (s, 1H), 4.02 - 3.76 (m, 5H), 3.34 - 3.20 (m, 4H), 3.02 - 2.87 (m, 2H), 2.66 (t, J = 6.9 Hz, 2H), 2.53 - 2.42 (m, 2H), 2.35 - 2.22 (m, 1H), 2.10 (td, J = 11.6, 3.0 Hz, 3H), 1.90 - 1.46 (m, 11H).
520		455.17	¹ H NMR (300 MHz, Chloroform-d) δ 8.61 (d, J = 2.0 Hz, 1H), 8.56 (d, J = 0.6 Hz, 2H), 8.53 (d, J = 1.9 Hz, 1H), 6.86 (d, J = 2.5 Hz, 1H), 4.67 (d, J = 6.1 Hz, 1H), 3.89 - 3.81 (m, 4H), 3.79 (d, J = 0.7 Hz, 2H), 3.31 - 3.17 (m, 4H), 2.60 (s, 1H), 2.20 - 2.03 (m, 2H), 1.79 - 1.59 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
521		465.16	¹ H NMR (300 MHz, Chloroform-d) δ 8.74 (d, J = 1.9 Hz, 1H), 8.66 (d, J = 2.0 Hz, 1H), 7.93 (s, 1H), 6.99 (d, J = 2.4 Hz, 1H), 6.93 (d, J = 2.5 Hz, 1H), 6.20 (d, J = 8.5 Hz, 1H), 4.87 (s, 1H), 4.36 (s, 1H), 3.94 (dd, J = 5.9, 3.8 Hz, 4H), 3.42 - 3.32 (m, 4H), 2.26 (s, 2H), 1.93 (s, 6H).
522		552.21	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.17 (s, 2H), 7.01 - 6.89 (m, 2H), 5.12 (d, J = 8.1 Hz, 1H), 4.80 (s, 1H), 4.09 - 3.86 (m, 5H), 3.63 (t, J = 7.1 Hz, 2H), 3.40 - 3.29 (m, 4H), 3.06 (d, J = 11.3 Hz, 2H), 2.79 (t, J = 7.1 Hz, 2H), 2.39 - 2.15 (m, 4H), 2.02 - 1.53 (m, 10H).
523		518.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.18 (s, 2H), 7.02 - 6.86 (m, 2H), 5.12 (d, J = 8.1 Hz, 1H), 4.80 (s, 1H), 4.09 - 3.87 (m, 5H), 3.42 - 3.29 (m, 4H), 3.12 (d, J = 11.2 Hz, 2H), 2.60 - 1.54 (m, 17H), 1.15 (t, J = 7.2 Hz, 3H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
524		301.24	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 6.91 (d, J = 2.5 Hz, 1H), 6.61 (d, J = 2.5 Hz, 1H), 5.19 - 5.05 (m, 1H), 3.97 - 3.85 (m, 5H), 3.37 - 3.29 (m, 4H), 2.71 (ddd, J = 16.1, 7.8, 5.0 Hz, 2H), 2.34 (ddd, J = 13.6, 7.0, 4.1 Hz, 2H).
525		379.26	¹ H NMR (400 MHz, CDCl ₃) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.31 (s, 1H), 8.30 (s, 1H), 6.93 (d, J = 2.4 Hz, 1H), 6.64 - 6.54 (m, 2H), 5.38 (d, J = 5.1 Hz, 1H), 5.17 - 5.07 (m, 1H), 4.68 - 4.56 (m, 1H), 3.95 - 3.85 (m, 4H), 3.35 - 3.27 (m, 4H), 2.94 (ddd, J = 13.4, 8.2, 5.1 Hz, 2H), 2.59 (ddd, J = 13.9, 7.0, 4.5 Hz, 2H).
526		550.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.80 - 2.07 (m, 8 H), 2.11 - 2.25 (m, 2 H), 2.39 - 2.76 (m, 6 H), 3.27 - 3.40 (m, 4 H), 3.70 - 3.84 (m, 4 H), 3.87 - 3.96 (m, 5 H), 3.99 (t, J = 6.2 Hz, 2 H), 4.73 - 4.82 (m, 1 H), 4.97 (d, J = 8.1 Hz, 1 H), 6.90 (d, J = 2.4 H




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
527		534.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.39 - 1.48 (m, 2 H), 1.56 - 1.67 (m, 4 H), 1.78 - 1.96 (m, 6 H), 2.14 - 2.25 (m, 2 H), 2.44 - 2.52 (m, 4 H), 2.72 (t, J = 6.0 Hz, 2 H), 3.31 - 3.35 (m, 4 H), 3.63 - 3.75 (m, 1 H), 3.90 - 3.93 (m, 4 H), 4.41 (t, J = 6.0 Hz, 2)
528		490.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.31 (s, 2H), 7.02 - 6.86 (m, 2H), 5.52 (d, J = 8.0 Hz, 1H), 4.81 (dt, J = 6.8, 3.4 Hz, 1H), 4.04 (td, J = 8.0, 4.0 Hz, 1H), 3.97 - 3.86 (m, 4H), 3.64 (s, 2H), 3.43 - 3.27 (m, 4H), 2.75 (t, J = 5.4 Hz, 4H), 2.32 - 2.15 (m, 2H), 2.07 (s, 2H), 1.92 (tt, J = 6.7, 3.6 Hz, 8H).
529		464.23	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.22 (s, 2H), 7.02 - 6.86 (m, 2H), 5.21 (d, J = 8.1 Hz, 1H), 4.81 (dt, J = 7.5, 3.7 Hz, 1H), 4.04 (d, J = 6.9 Hz, 1H), 3.96 - 3.82 (m, 4H), 3.41 - 3.30 (m, 4H), 3.27 (s, 2H), 2.24 (m, 8H), 2.07 - 1.77 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
530		506.25	¹ H NMR (300 MHz, Chloroform-d) δ 8.61 (d, J = 1.9 Hz, 1H), 8.53 (d, J = 1.9 Hz, 1H), 8.13 (s, 2H), 6.87 (d, J = 2.5 Hz, 1H), 6.83 (d, J = 2.5 Hz, 1H), 5.10 (d, J = 8.1 Hz, 1H), 4.71 (s, 1H), 3.94 (s, 1H), 3.88 - 3.78 (m, 4H), 3.67 - 3.56 (m, 4H), 3.32 - 3.17 (m, 6H), 2.41 - 2.26 (m, 4H), 2.12 (d, J = 8.7 Hz, 2H), 1.92 - 1.76 (m, 6H).
531		425.19	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.39 (dd, J = 2.7, 0.5 Hz, 1H), 8.04 (d, J = 3.4 Hz, 1H), 6.97 (d, J = 2.5 Hz, 1H), 6.92 (d, J = 2.5 Hz, 1H), 5.15 (d, J = 8.1 Hz, 1H), 4.83 (dq, J = 5.1, 2.6 Hz, 1H), 4.25 (dd, J = 8.1, 4.6 Hz, 1H), 3.98 - 3.88 (m, 4H), 3.39 - 3.28 (m, 4H), 2.34 - 2.17 (m, 2H), 2.03 - 1.84 (m, 6H).
532		441.29	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.13 (d, J = 4.8 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.5 Hz, 1H), 6.54 (d, J = 5.2 Hz, 1H), 5.36 (s, 1H), 4.79 (s, 1H), 4.03 (s, 1H), 3.97 - 3.84 (m, 4H), 3.41 - 3.23 (m, 4H), 2.26 - 2.12 (m, 2H), 1.98 - 1.81 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
533		441.25	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 7.99 (s, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 6.22 (d, J = 5.9 Hz, 1H), 5.22 (s, 1H), 4.81 (s, 1H), 4.03 (dt, J = 12.1, 6.2 Hz, 1H), 3.94 - 3.86 (m, 4H), 3.40 - 3.26 (m, 4H), 2.27 - 2.13 (m, 2H), 1.90 (t, J = 17.4 Hz, 6H).
534		492.39	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (s, 1H), 8.61 (s, 1H), 7.89 (d, J = 5.5 Hz, 1H), 6.95 (s, 1H), 6.89 (s, 1H), 5.71 (d, J = 5.8 Hz, 1H), 4.76 (s, 2H), 4.01 - 3.81 (m, 5H), 3.74 (s, 8H), 3.41 - 3.26 (m, 4H), 2.26 - 2.12 (m, 2H), 1.98 - 1.81 (m, 6H).
535		505.35	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.88 (d, J = 5.7 Hz, 1H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 5.67 (d, J = 5.8 Hz, 1H), 4.84 - 4.63 (m, 2H), 3.99 - 3.83 (m, 5H), 3.82 - 3.73 (m, 4H), 3.37 - 3.27 (m, 4H), 2.49 - 2.40 (m, 4H), 2.33 (s, 3H), 2.23 - 2.11 (m, 2H), 1.96 - 1.82 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
536		461.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.73 (d, J = 1.9 Hz, 1H), 8.65 (d, J = 1.9 Hz, 1H), 8.30 (s, 1H), 6.97 (dd, J = 15.8, 2.5 Hz, 2H), 6.23 (d, J = 8.2 Hz, 1H), 6.07 (s, 1H), 4.90 (dt, J = 5.1, 2.4 Hz, 1H), 3.99 - 3.88 (m, 4H), 3.72 (dq, J = 9.2, 4.7 Hz, 1H), 3.40 - 3.29 (m, 4H), 2.60 (s, 3H), 2.34 (dt, J = 14.1, 5.2 Hz, 2H), 2.23 - 1.79 (m, 6H).
537		505.35	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.7 Hz, 1H), 8.60 (d, J = 1.7 Hz, 1H), 8.18 (s, 1H), 6.95 (d, J = 2.2 Hz, 1H), 6.90 (s, 1H), 5.42 (s, 1H), 4.78 (s, 1H), 4.71 (d, J = 8.3 Hz, 1H), 3.98 - 3.87 (m, 4H), 3.83 (s, 1H), 3.65 - 3.48 (m, 4H), 3.41 - 3.24 (m, 4H), 2.54 - 2.42 (m, 4H), 2.33 (s, 3H), 2.25 - 2.11 (m, 2H), 1.97 - 1.82 (m, 6H).
538		480.19	¹ H NMR (300 MHz, Chloroform-d) δ 8.74 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.57 (d, J = 1.6 Hz, 1H), 7.38 (dd, J = 1.7, 0.8 Hz, 1H), 7.05 - 6.88 (m, 3H), 5.52 (d, J = 8.1 Hz, 1H), 4.87 (d, J = 4.0 Hz, 1H), 4.47 - 4.31 (m, 1H), 3.98 - 3.88 (m, 4H), 3.42 - 3.30 (m, 4H), 2.35 - 2.21 (m, 2H), 2.16 - 1.88 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
539		481.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.76 - 1.97 (m, 6 H), 2.15 - 2.26 (m, 2 H), 3.31 - 3.35 (m, 4 H), 3.43 (s, 3 H), 3.61 - 3.72 (m, 3 H), 3.90 - 3.93 (m, 4 H), 4.45 - 4.48 (m, 2 H), 4.76 - 4.82 (m, 1 H), 4.94 (d, J = 7.1 Hz, 1 H), 5.73 (s, 1 H), 6.89 (d, J = 2.
540		445.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.82 - 1.99 (m, 6 H), 2.13 - 2.25 (m, 2 H), 3.29 - 3.38 (m, 4 H), 3.73 (br. s, 1 H), 3.84 - 3.98 (m, 4 H), 4.45 (br. s, 1 H), 4.74 - 4.84 (m, 1 H), 6.31 (s, 1 H), 6.43 - 6.47 (m, 1 H), 6.91 (d, J = 2.3 Hz, 1 H), 6.93 - 6.99 (m
541		534.4	¹ H NMR (300 MHz, CDCl ₃): ppm 1.40 - 1.48 (m, 2 H), 1.56 - 1.67 (m, 4 H), 1.84 - 1.99 (m, 6 H), 2.11 - 2.24 (m, 2 H), 2.45 - 2.52 (m, 4 H), 2.72 (t, J = 6.2 Hz, 2 H), 3.32 - 3.35 (m, 4 H), 3.90 - 3.93 (m, 4 H), 3.96 - 4.08 (m, 1 H), 4.40 (t, J = 6.2 Hz, 2




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
542		447.25	¹ H NMR (300 MHz, Chloroform-d) δ 8.63 (d, J = 1.9 Hz, 1H), 8.56 (d, J = 1.9 Hz, 1H), 8.18 (s, 1H), 7.83 (d, J = 6.3 Hz, 1H), 6.96 - 6.80 (m, 3H), 6.23 (d, J = 8.2 Hz, 1H), 4.77 (dd, J = 5.8, 3.3 Hz, 1H), 4.23 (dq, J = 8.6, 4.3 Hz, 1H), 3.89 - 3.75 (m, 4H), 3.35 - 3.20 (m, 4H), 2.20 (d, J = 12.8 Hz, 2H), 2.10 - 1.79 (m, 6H).
543		518.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.53 (s, 1H), 8.18 (s, 2H), 6.96 - 6.80 (m, 2H), 5.14 (d, J = 8.1 Hz, 1H), 4.78 (s, 1H), 4.08 - 3.86 (m, 5H), 3.33 (dd, J = 6.0, 3.8 Hz, 4H), 3.07 - 2.94 (m, 2H), 2.71 (s, 3H), 2.35 (s, 3H), 2.26 - 1.97 (m, 4H), 1.99 - 1.57 (m, 11H).
544		518.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.61 (s, 1H), 8.18 (s, 2H), 7.02 - 6.87 (m, 2H), 5.12 (d, J = 8.2 Hz, 1H), 4.80 (s, 1H), 4.05 - 3.87 (m, 5H), 3.38 - 3.27 (m, 4H), 3.01 (d, J = 11.4 Hz, 3H), 2.74 (s, 3H), 2.36 (s, 3H), 2.20 (m, 2H), 2.06 (m, 2H), 2.00 - 1.58 (m, 11H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
545		461.24	¹ H NMR (300 MHz, Chloroform-d) δ 8.73 (d, J = 2.0 Hz, 1H), 8.64 (d, J = 2.0 Hz, 1H), 8.39 (s, 1H), 7.92 (s, 1H), 7.05 - 6.89 (m, 2H), 4.88 (s, 1H), 4.06 (s, 3H), 3.94 (dd, J = 6.0, 3.8 Hz, 4H), 3.36 (t, J = 4.9 Hz, 5H), 2.29 (s, 2H), 1.98 (d, J = 35.4 Hz, 6H).
546		461.24	¹ H NMR (300 MHz, Chloroform-d) δ 8.62 (d, J = 1.9 Hz, 1H), 8.55 (d, J = 1.9 Hz, 1H), 8.32 (s, 1H), 7.66 (s, 1H), 6.92 - 6.81 (m, 2H), 4.79 (s, 1H), 3.87 - 3.80 (m, 4H), 3.75 (s, 3H), 3.31 - 3.22 (m, 4H), 2.18 (d, J = 12.7 Hz, 2H), 2.08 - 1.75 (m, 6H).
547		505.4	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.90 (d, J = 6.0 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.92 (d, J = 2.3 Hz, 1H), 5.89 (d, J = 6.1 Hz, 1H), 4.96 (s, 1H), 4.77 (s, 1H), 4.08 - 3.98 (m, 1H), 3.96 - 3.87 (m, 4H), 3.70 - 3.53 (m, 4H), 3.43 - 3.27 (m, 4H), 2.53 - 2.42 (m, 4H), 2.35 (s, 3H), 2.25 - 2.13 (m, 2H), 2.00 - 1.85 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
548		492.39	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.8 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.93 (d, J = 5.9 Hz, 1H), 6.96 (d, J = 2.3 Hz, 1H), 6.91 (d, J = 2.3 Hz, 1H), 5.87 (d, J = 6.1 Hz, 1H), 4.97 (s, 1H), 4.77 (s, 1H), 4.02 (s, 1H), 3.97 - 3.88 (m, 4H), 3.83 - 3.70 (m, 4H), 3.63 - 3.52 (m, 4H), 3.39 - 3.27 (m, 4H), 2.25 - 2.12 (m, 2H), 2.01 - 1.85 (m, 6H).
549		504.27	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.18 (d, J = 5.1 Hz, 1H), 6.98 - 6.89 (m, 2H), 6.42 (d, J = 5.1 Hz, 1H), 5.18 (d, J = 8.0 Hz, 1H), 4.79 (dt, J = 7.9, 3.7 Hz, 1H), 4.11 - 3.87 (m, 5H), 3.42 - 3.25 (m, 4H), 3.00 (dq, J = 9.9, 3.3 Hz, 2H), 2.63 - 2.35 (m, 5H), 2.27 - 2.04 (m, 4H), 1.97 - 1.82 (m, 9H).
550		546.31	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.16 (s, 2H), 7.05 - 6.83 (m, 2H), 5.14 (d, J = 8.1 Hz, 1H), 4.79 (d, J = 5.3 Hz, 1H), 4.10 - 3.85 (m, 5H), 3.41 - 3.28 (m, 4H), 2.87 (tt, J = 12.8, 3.2 Hz, 1H), 2.20 (q, J = 6.2 Hz, 2H), 1.91 (p, J = 3.7, 2.8 Hz, 6H), 1.73 (dd, J = 13.1, 3.2 Hz, 2H), 1.29 (d, J = 24.9 Hz, 12H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
551		490.28	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.64 (d, J = 1.9 Hz, 1H), 8.20 (d, J = 5.1 Hz, 1H), 7.01 - 6.89 (m, 2H), 6.42 (d, J = 5.1 Hz, 1H), 5.15 (d, J = 7.9 Hz, 1H), 4.80 (s, 1H), 4.05 (s, 1H), 3.97 - 3.88 (m, 4H), 3.43 - 3.31 (m, 4H), 3.24 (d, J = 12.3 Hz, 2H), 2.84 - 2.69 (m, 2H), 2.59 (m, 1H), 2.20 (m, 2H), 1.92 (t, J = 8.2 Hz, 6H).
552		462.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.61 (d, J = 2.0 Hz, 1H), 8.53 (d, J = 1.9 Hz, 1H), 7.85 (s, 1H), 6.94 - 6.73 (m, 2H), 5.33 (d, J = 7.9 Hz, 1H), 4.69 (s, 1H), 4.00 - 3.78 (m, 5H), 3.38 (d, J = 1.0 Hz, 2H), 3.33 - 3.20 (m, 4H), 2.19 - 1.99 (m, 2H), 1.83 (h, J = 5.6 Hz, 6H).
553		480.19	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 6.93 (dd, J = 17.8, 2.5 Hz, 2H), 5.07 (s, 1H), 4.78 (d, J = 6.2 Hz, 1H), 4.68 (d, J = 7.9 Hz, 1H), 4.03 - 3.90 (m, 4H), 3.87 (s, 3H), 3.71 (s, 1H), 3.42 - 3.28 (m, 4H), 3.13 (s, 6H), 2.27 - 2.00 (m, 2H), 2.00 - 1.79 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
554		462.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.04 (s, 1H), 5.04 (d, J = 8.2 Hz, 1H), 4.79 (s, 1H), 4.09 - 3.90 (m, 5H), 3.85 (s, 2H), 3.42 - 3.31 (m, 4H), 3.11 (t, J = 5.8 Hz, 2H), 2.63 (t, J = 5.8 Hz, 2H), 2.19 (d, J = 8.1 Hz, 2H), 1.91 (d, J = 5.5 Hz, 6H).
555		357.15	¹ H NMR (400 MHz, CDCl ₃) δ 8.68 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.87 (d, J = 2.4 Hz, 1H), 5.47 (s, 1H), 5.30 (s, 1H), 4.50 (s, 1H), 4.00 - 3.84 (m, 4H), 3.40 - 3.27 (m, 4H), 2.45 - 2.33 (m, 2H), 2.33 - 2.18 (m, 1H), 2.15 - 2.00 (m, 2H), 1.74 (d, J = 5.4 Hz, 4H).
556		476.23	¹ H NMR (300 MHz, Chloroform-d) δ 8.61 (d, J = 1.9 Hz, 1H), 8.53 (d, J = 1.9 Hz, 1H), 7.97 (s, 1H), 6.92 - 6.77 (m, 2H), 4.95 (d, J = 8.1 Hz, 1H), 4.76 - 4.64 (m, 1H), 3.97 - 3.75 (m, 5H), 3.34 (s, 2H), 3.30 - 3.17 (m, 4H), 2.60 (dq, J = 9.7, 5.3, 4.8 Hz, 4H), 2.37 (s, 3H), 2.10 (td, J = 10.8, 10.3, 6.1 Hz, 2H), 1.90 - 1.62 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
557		590.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 2.0 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.20 (d, J = 5.1 Hz, 1H), 7.01 - 6.89 (m, 2H), 6.40 (d, J = 5.1 Hz, 1H), 5.20 (s, 1H), 4.81 (d, J = 6.1 Hz, 1H), 4.23 (s, 2H), 4.04 (s, 1H), 3.97 - 3.89 (m, 4H), 3.42 - 3.28 (m, 4H), 2.82 (t, J = 12.8 Hz, 2H), 2.67 - 2.44 (m, 1H), 2.27 - 2.11 (m, 2H), 1.98 - 1.87 (m, 8H), 1.49 (s, 9H).
558		481.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.84 - 1.99 (m, 6 H), 2.11 - 2.25 (m, 2 H), 3.32 - 3.36 (m, 4 H), 3.43 (s, 3 H), 3.70 - 3.73 (m, 2 H), 3.90 - 3.93 (m, 4 H), 3.96 - 4.07 (m, 1 H), 4.41 - 4.44 (m, 2 H), 4.72 - 4.80 (m, 1 H), 5.10 (br. s, 1 H), 6.06 (d, J = 5.6
559		494.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.84 - 1.98 (m, 6 H), 2.13 - 2.23 (m, 2 H), 2.32 (s, 6 H), 2.68 (t, J = 5.8 Hz, 2 H), 3.30 - 3.37 (m, 4 H), 3.88 - 3.95 (m, 4 H), 3.98 - 4.08 (m, 1 H), 4.37 (t, J = 5.8 Hz, 2 H), 4.73 - 4.80 (m, 1 H), 5.05 (d, J = 7.5 Hz, 1 H)




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
560		536.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.83 - 2.00 (m, 6 H), 2.10 - 2.25 (m, 2 H), 2.52 - 2.59 (m, 4 H), 2.76 (t, J = 5.8 Hz, 2 H), 3.31 - 3.38 (m, 4 H), 3.70 - 3.77 (m, 4 H), 3.89 - 3.95 (m, 4 H), 4.03 (br. s, 1 H), 4.41 (t, J = 5.8 Hz, 2 H), 4.72 - 4.80 (m, 1 H),
561		494.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.78 - 1.94 (m, 6 H), 2.14 - 2.25 (m, 2 H), 2.32 (s, 6 H), 2.67 (t, J = 5.5 Hz, 2 H), 3.28 - 3.38 (m, 4 H), 3.66 (br. s, 1 H), 3.84 - 3.98 (m, 4 H), 4.39 (t, J = 5.5 Hz, 2 H), 4.79 (br. s, 1 H), 4.90 (d, J = 7.2 Hz, 1 H), 5.73
562		549.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.81 - 1.94 (m, 6 H), 2.14 - 2.24 (m, 2 H), 2.28 (s, 3 H), 2.47 (br. s, 4 H), 2.59 (br. s, 4 H), 2.76 (t, J = 5.8 Hz, 2 H), 3.30 - 3.37 (m, 4 H), 3.69 (br. s, 1 H), 3.88 - 3.95 (m, 4 H), 4.42 (t, J = 5.8 Hz, 2 H), 4.75 - 4.83

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
563		447.37	¹ H NMR (400 MHz, CDCl ₃) δ 8.68 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.48 (d, J = 0.6 Hz, 1H), 7.75 (d, J = 2.2 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.92 - 6.88 (m, 1H), 6.67 (d, J = 1.4 Hz, 1H), 5.22 (t, J = 9.5 Hz, 1H), 4.79 (s, 1H), 4.06 (s, 1H), 4.00 - 3.84 (m, 4H), 3.40 - 3.24 (m, 4H), 2.16 (t, J = 21.2 Hz, 2H), 1.98 - 1.85 (m, 6H).
564		562.25	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.07 (s, 1H), 6.99 - 6.84 (m, 2H), 5.10 (d, J = 8.1 Hz, 1H), 4.78 (d, J = 5.7 Hz, 1H), 4.42 (s, 2H), 4.10 - 3.85 (m, 5H), 3.64 (t, J = 5.6 Hz, 2H), 3.40 - 3.29 (m, 4H), 2.65 (dt, J = 6.2, 2.8 Hz, 2H), 2.19 (q, J = 6.0 Hz, 2H), 1.98 - 1.85 (m, 6H), 1.49 (s, 9H).
565		479.1	¹ H NMR (300 MHz, Chloroform-d) δ 8.55 (d, J = 1.9 Hz, 1H), 8.47 (d, J = 1.9 Hz, 1H), 6.93 (s, 1H), 6.86 - 6.72 (m, 2H), 5.25 (s, 1H), 4.66 (s, 1H), 4.05 - 3.90 (m, 1H), 3.86 - 3.71 (m, 7H), 3.30 - 3.09 (m, 4H), 2.27 (d, J = 2.2 Hz, 3H), 2.04 (d, J = 9.2 Hz, 2H), 1.75 (d, J = 4.8 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
566		475.15	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 2.0 Hz, 2H), 8.63 (d, J = 1.9 Hz, 1H), 7.42 (s, 1H), 7.01 - 6.86 (m, 2H), 5.64 (d, J = 8.2 Hz, 1H), 4.83 (s, 1H), 3.92 (dd, J = 6.0, 3.7 Hz, 4H), 3.34 (dd, J = 5.8, 3.9 Hz, 4H), 2.11 - 1.78 (m, 6H).
567		504.1	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.74 (s, 1H), 7.00 - 6.89 (m, 2H), 4.91 (s, 1H), 4.76 (d, J = 6.1 Hz, 1H), 4.04 - 3.93 (m, 5H), 3.43 - 3.28 (m, 4H), 2.26 - 2.14 (m, 2H), 2.08 (d, J = 0.8 Hz, 3H), 2.00 - 1.92 (m, 6H), 1.69 (d, J = 21.2 Hz, 10H).
568		490.23	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 2.0 Hz, 1H), 8.22 (s, 2H), 7.04 - 6.88 (m, 2H), 5.14 (d, J = 8.1 Hz, 1H), 4.80 (d, J = 5.5 Hz, 1H), 4.09 - 3.85 (m, 5H), 3.41 - 3.31 (m, 4H), 3.29 - 3.13 (m, 1H), 3.04 - 2.92 (m, 1H), 2.81 (q, J = 8.1 Hz, 1H), 2.68 (dt, J = 9.2, 4.6 Hz, 1H), 2.52 - 2.26 (m, 5H), 2.24 - 2.13 (m, 2H), 2.02 - 1.78 (m, 7H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
569		476.06	¹ H NMR (300 MHz, Chloroform-d) δ 8.61 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 8.10 (s, 2H), 6.93 - 6.76 (m, 2H), 5.07 (d, J = 8.1 Hz, 1H), 4.68 (d, J = 15.8 Hz, 1H), 4.00 - 3.78 (m, 5H), 3.35 - 3.20 (m, 4H), 3.18 - 2.92 (m, 3H), 2.70 (dd, J = 10.7, 8.4 Hz, 1H), 2.24 - 2.02 (m, 3H), 1.94 - 1.62 (m, 7H).
570		576.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 2.2 Hz, 1H), 8.63 (d, J = 2.1 Hz, 1H), 8.19 (s, 2H), 7.03 - 6.86 (m, 2H), 4.10 - 3.86 (m, 5H), 3.46 - 3.28 (m, 4H), 3.24 - 3.13 (m, 1H), 2.35 - 2.16 (m, 1H), 2.05 - 1.84 (m, 8H), 1.81 - 1.66 (m, 1H), 1.55 - 1.41 (m, 11H).
571		476.31	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.88 (d, J = 2.4 Hz, 1H), 6.96 (d, J = 2.5 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 6.76 (s, 1H), 4.98 (s, 2H), 4.78 (s, 1H), 4.00 - 3.86 (m, 4H), 3.83 (s, 1H), 3.39 - 3.27 (m, 4H), 2.21 (d, J = 9.6 Hz, 2H), 1.89 (dd, J = 13.3, 8.6 Hz, 6H), 1.50 (s, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
572		472.2	¹ H NMR (300 MHz, CDCl ₃): ppm 1.82 - 1.99 (m, 6 H), 2.14 - 2.28 (m, 2 H), 3.32 - 3.35 (m, 4 H), 3.80 - 3.88 (m, 1 H), 3.90 - 3.93 (m, 4 H), 4.70 - 4.83 (m, 2 H), 6.37 (d, J = 8.8 Hz, 1 H), 6.38 (t, J = 73.8 Hz, 1 H), 6.90 (d, J = 2.3 Hz, 1 H), 6.95 (d, J =
573		464.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.52 (s, 6 H), 1.73 (br s, 1 H), 1.85 - 1.96 (m, 6 H), 2.11 - 2.25 (m, 2 H), 3.30 - 3.36 (m, 4 H), 3.88 - 4.00 (m, 5 H), 4.55 (d, J = 8.1 Hz, 1 H), 4.74 - 4.81 (m, 1 H), 6.52 (s, 1 H), 6.57 (dd, J = 5.4, 1.4 Hz, 1 H), 6.90 (d,
574		437.42	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.99 (s, 2H), 6.95 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.4 Hz, 1H), 4.78 (s, 1H), 4.01 - 3.84 (m, 7H), 3.41 (s, 2H), 3.37 - 3.27 (m, 4H), 2.26 - 2.14 (m, 2H), 1.93 - 1.81 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
575		407.35	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.57 (s, 1H), 8.13 (s, 2H), 6.98 (d, J = 2.4 Hz, 1H), 6.91 (d, J = 2.5 Hz, 1H), 4.86 - 4.73 (m, 1H), 3.95 - 3.88 (m, 4H), 3.56 - 3.49 (m, 1H), 3.39 - 3.31 (m, 4H), 2.30 - 2.18 (m, 2H), 1.97 - 1.84 (m, 6H).
576		450.13	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 2.0 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.13 (s, 1H), 7.01 - 6.87 (m, 2H), 4.81 (s, 1H), 4.57 (s, 2H), 4.06 (s, 1H), 3.99 - 3.87 (m, 4H), 3.35 (dd, J = 5.9, 3.8 Hz, 4H), 2.45 (s, 3H), 2.20 (d, J = 8.0 Hz, 2H), 1.93 (d, J = 5.1 Hz, 6H).
577		437.25	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 2.0 Hz, 1H), 8.02 (d, J = 1.4 Hz, 1H), 7.87 (d, J = 1.5 Hz, 1H), 6.94 (dd, J = 16.2, 2.5 Hz, 2H), 4.89 - 4.71 (m, 2H), 4.65 (s, 2H), 4.06 - 3.85 (m, 4H), 3.40 - 3.27 (m, 4H), 2.23 (dt, J = 11.5, 5.5 Hz, 2H), 2.04 - 1.80 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
578		437.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 2.0 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.01 - 6.88 (m, 2H), 6.87 - 6.66 (m, 2H), 3.94 (t, J = 4.9 Hz, 4H), 3.65 (s, 3H), 3.42 - 3.30 (m, 4H), 2.19 (s, 2H), 1.91 (d, J = 4.9 Hz, 6H).
579		505.53	¹ H NMR (400 MHz, CDCl ₃) δ 8.68 (s, 1H), 8.61 (s, 1H), 8.07 (s, 2H), 6.94 (s, 1H), 6.90 (s, 1H), 4.96 (d, J = 8.1 Hz, 1H), 4.78 (s, 1H), 4.02 - 3.81 (m, 5H), 3.47 - 3.19 (m, 5H), 3.19 - 2.83 (m, 4H), 2.76 - 2.52 (m, 3H), 2.37 (s, 3H), 2.26 - 2.11 (m, 2H), 2.03 - 1.82 (m, 6H).
580		445.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.84 - 2.17 (m, 6 H), 2.23 - 2.35 (m, 2 H), 3.31 - 3.38 (m, 4 H), 3.63 $\bar{n}$ 3.75 (m, 1 H), 3.87 - 3.96 (m, 4 H), 4.82 - 4.89 (m, 1 H), 5.88 (d, J = 7.2 Hz, 1 H), 6.11 (d, J = 7.8 Hz, 1 H), 6.44 (d, J = 2.2 Hz, 1 H), 6.90 $\bar{n}$ 6.95 (




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
581		446.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.85 - 2.18 (m, 6 H), 2.24 - 2.36 (m, 2 H), 3.31 - 3.39 (m, 4 H), 3.64 - 3.77 (m, 1 H), 3.88 - 3.97 (m, 4 H), 4.82 - 4.89 (m, 1 H), 5.91 (d, J = 7.9 Hz, 1 H), 6.12 (d, J = 7.9 Hz, 1 H), 6.93 (d, J = 2.4 Hz, 1 H), 6.98 (d, J =
582		450.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.83 - 1.96 (m, 6 H), 2.14 - 2.26 (m, 2 H), 3.30 - 3.36 (m, 4 H), 3.41 (s, 3 H), 3.87 - 3.95 (m, 5 H), 4.36 (s, 2 H), 4.59 - 4.89 (m, 2 H), 6.38 (s, 1 H), 6.48 (d, J = 5.2 Hz, 1 H), 6.90 (d, J = 2.4 Hz, 1 H), 6.95 (d, J = 2.4
583		466.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.84 - 2.00 (m, 6 H), 2.09 - 2.24 (m, 2 H), 3.30 - 3.37 (m, 4 H), 3.64 - 3.74 (m, 1 H), 3.78 (s, 3 H), 3.89 - 3.95 (m, 7 H), 4.71 - 4.79 (m, 1 H), 5.88 (d, J = 8.3 Hz, 1 H), 6.90 (d, J = 2.4 Hz, 1 H), 6.95 (d, J = 2.4 Hz, 1 H)




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
584		436.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.86 - 2.04 (m, 6 H), 2.12 - 2.26 (m, 2 H), 3.28 - 3.39 (m, 4 H), 3.88 - 3.96 (m, 4 H), 4.26 (br s, 1 H), 4.62 (s, 2 H), 4.69 - 4.78 (m, 1 H), 5.56 (br s, 1 H), 6.50 (t, J = 5.3 Hz, 1 H), 6.90 (d, J = 1.9 Hz, 1 H), 6.93 (d, J
585		536.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.81 - 1.96 (m, 6 H), 2.11 - 2.25 (m, 2 H), 2.56 (br s, 4 H), 2.76 (t, J = 5.3 Hz, 2 H), 3.30 - 3.36 (m, 4 H), 3.74 (t, J = 4.4 Hz, 4 H), 3.88 - 3.96 (m, 5 H), 4.06 (t, J = 5.3 Hz, 2 H), 4.73 - 4.81 (m, 1 H), 4.99 (d, J = 7.8
586		420.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.88 ñ 2.01 (m, 6 H), 2.10 (s, 3 H), 2.15 - 2.25 (m, 2 H), 3.30 - 3.37 (m, 4 H), 3.89 ñ 3.95 (m, 4 H), 4.25 (br s, 2 H), 4.71 - 4.78 (m, 1 H), 6.51 (dd, J = 6.8, 5.4 Hz, 1 H), 6.90 (d, J = 2.3 Hz, 1 H), 6.94 (d, J = 2.3 Hz, 1

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
587		474.2	¹ H NMR (300 MHz, CDCl ₃): ppm 1.82 - 1.98 (m, 6 H), 2.12 - 2.26 (m, 2 H), 3.25 - 3.43 (m, 4 H), 3.82 - 3.98 (m, 5 H), 4.80 (br s, 2 H), 6.52 (d, J = 8.4 Hz, 1 H), 6.87 - 6.93 (m, 2 H), 6.95 - 7.00 (m, 1 H), 7.50 (t, J = 7.9 Hz, 1 H), 8.63 (d, J = 1.2 Hz, 1 H)
588		450.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.38 (t, J = 7.0 Hz, 3 H), 1.83 - 2.00 (m, 6 H), 2.11 - 2.24 (m, 2 H), 3.30 - 3.37 (m, 4 H), 3.75 - 3.85 (m, 1 H), 3.89 - 3.94 (m, 4 H), 4.25 (q, J = 7.0 Hz, 2 H), 4.42 - 4.54 (m, 1 H), 4.72 - 4.79 (m, 1 H), 5.94 (d, J = 7.7 Hz, 1 H)
589		474.2	¹ H NMR (300 MHz, CDCl ₃): ppm 1.86 - 1.99 (m, 6 H), 2.16 - 2.27 (m, 2 H), 3.31 - 3.36 (m, 4 H), 3.88 - 3.95 (m, 5 H), 4.77 - 4.84 (m, 1 H), 4.90 (br s, 1 H), 6.55 (s, 1 H), 6.72 (d, J = 5.0 Hz, 1 H), 6.91 (d, J = 2.4 Hz, 1 H), 6.95 (d, J = 2.4 Hz, 1 H), 8.




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
590		491.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.82 - 1.98 (m, 6 H), 2.10 - 2.22 (m, 2 H), 3.31 - 3.35 (m, 4 H), 3.41 - 3.45 (m, 4 H), 3.69 - 3.82 (m, 5 H), 3.90 - 3.93 (m, 4 H), 4.43 (d, J = 7.7 Hz, 1 H), 4.70 - 4.77 (m, 1 H), 5.81 (d, J = 8.0 Hz, 1 H), 5.90 (d, J = 8.0 Hz, 1 H)
591		424.2	¹ H NMR (300 MHz, CDCl ₃): ppm 1.88 - 2.00 (m, 6 H), 2.13 - 2.28 (m, 2 H), 3.29 - 3.38 (m, 4 H), 3.88 - 3.95 (m, 4 H), 4.21 (br s, 1 H), 4.62 - 4.84 (m, 2 H), 6.45 - 6.54 (m, 1 H), 6.90 (d, J = 2.4 Hz, 1 H), 6.95 (d, J = 2.4 Hz, 1 H), 7.07 - 7.18 (m, 1 H),
592		436.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.84 - 2.03 (m, 6 H), 2.11 - 2.24 (m, 2 H), 3.30 - 3.37 (m, 4 H), 3.84 (s, 3 H), 3.89 - 3.95 (m, 4 H), 4.17 - 4.29 (m, 1 H), 4.67 - 4.76 (m, 1 H), 5.08 (d, J = 7.4 Hz, 1 H), 6.50 (dd, J = 7.3, 5.2 Hz, 1 H), 6.82 (d, J = 7.4 Hz, 1 H)

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
593		474.2	¹ H NMR (300 MHz, CDCl ₃): ppm 1.86 - 2.00 (m, 6 H), 2.13 - 2.26 (m, 2 H), 3.30 - 3.38 (m, 4 H), 3.88 - 3.97 (m, 4 H), 4.22 - 4.35 (m, 1 H), 4.73 - 4.80 (m, 1 H), 4.89 - 4.99 (m, 1 H), 6.60 (dd, J = 7.1, 5.1 Hz, 1 H), 6.91 (d, J = 2.3 Hz, 1 H), 6.96 (d, J =
594		442.2	¹ H NMR (300 MHz, CDCl ₃): ppm 1.85 - 2.00 (m, 6 H), 2.11 - 2.28 (m, 2 H), 3.29 - 3.39 (m, 4 H), 3.87 - 3.97 (m, 4 H), 4.06 - 4.18 (m, 1 H), 4.56 (d, J = 7.6 Hz, 1 H), 4.73 - 4.81 (m, 1 H), 6.90 (d, J = 2.4 Hz, 1 H), 6.95 (d, J = 2.4 Hz, 1 H), 7.05 (ddd, J =
595		434.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.88 - 2.02 (m, 6 H), 2.08 (s, 3 H), 2.15 (s, 3 H), 2.14 - 2.24 (m, 2 H), 3.31 - 3.36 (m, 4 H), 3.89 - 3.95 (m, 4 H), 4.03 - 4.16 (m, 1 H), 4.23 (br s, 1 H), 4.69 - 4.77 (m, 1 H), 6.90 (d, J = 2.4 Hz, 1 H), 6.94 (d, J = 2.4 Hz




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
596		477.68	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 2.0 Hz, 1H), 8.63 (d, J = 2.0 Hz, 1H), 6.99 (d, J = 2.4 Hz, 1H), 6.92 (t, J = 2.8 Hz, 1H), 6.08 (d, J = 1.1 Hz, 1H), 5.21 (s, 1H), 4.81 (s, 2H), 3.98 - 3.87 (m, 4H), 3.43 - 3.30 (m, 4H), 2.51 (d, J = 1.0 Hz, 3H), 2.19 (s, 2H), 2.01 - 1.77 (m, 6H).
597		449.23	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.07 - 6.87 (m, 2H), 6.02 (s, 1H), 4.83 (d, J = 5.5 Hz, 1H), 4.03 - 3.81 (m, 5H), 3.44 - 3.25 (m, 4H), 2.72 - 2.57 (m, 1H), 2.51 (s, 3H), 2.29 - 2.11 (m, 2H), 2.05 - 1.72 (m, 6H), 1.27 (t, J = 7.6 Hz, 3H).
598		450.23	¹ H NMR (300 MHz, Chloroform-d) δ 8.61 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 6.93 - 6.72 (m, 3H), 6.55 (s, 1H), 4.71 (s, 1H), 4.01 (s, 1H), 3.92 - 3.76 (m, 4H), 3.34 - 3.18 (m, 4H), 2.98 (s, 6H), 2.07 (d, J = 24.1 Hz, 2H), 1.87 (d, J = 9.2 Hz, 6H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
599		437.21	¹ H NMR (300 MHz, Chloroform-d) δ 8.62 (d, J = 1.9 Hz, 1H), 8.54 (t, J = 2.4 Hz, 1H), 6.85 (dd, J = 18.9, 2.4 Hz, 2H), 6.63 - 6.50 (m, 1H), 4.68 (s, 1H), 3.92 - 3.76 (m, 4H), 3.66 (s, 1H), 3.32 - 3.17 (m, 4H), 2.16 - 1.99 (m, 5H), 1.92 - 1.72 (m, 6H).
600		464.26	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 2.0 Hz, 1H), 8.62 (d, J = 2.0 Hz, 1H), 7.00 - 6.85 (m, 2H), 5.44 (d, J = 8.5 Hz, 1H), 4.78 (s, 1H), 3.93 (t, J = 4.8 Hz, 4H), 3.35 (t, J = 4.9 Hz, 4H), 3.06 - 2.91 (m, 1H), 2.15 (d, J = 26.6 Hz, 2H), 1.89 (s, 6H), 1.30 - 0.94 (m, 4H).
601		461.32	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.06 (d, J = 9.1 Hz, 1H), 6.98 - 6.86 (m, 2H), 6.58 (d, J = 9.1 Hz, 1H), 4.81 (dq, J = 5.5, 2.7 Hz, 1H), 4.52 (s, 1H), 4.17 (d, J = 5.0 Hz, 1H), 4.01 - 3.87 (m, 4H), 3.67 (dq, J = 8.9, 8.1 Hz, 1H), 3.43 - 3.30 (m, 4H), 2.47 - 2.29 (m, 4H), 2.22 (dt, J = 11.9, 5.5 Hz, 2H), 2.05 - 1.82 (m, 6H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
602		464.1	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.04 - 6.92 (m, 2H), 4.82 (d, J = 3.0 Hz, 1H), 3.93 (dd, J = 5.9, 3.8 Hz, 4H), 3.65 (d, J = 8.7 Hz, 1H), 3.44 - 3.29 (m, 4H), 3.08 (s, 6H), 2.39 (s, 3H), 2.21 (d, J = 14.4 Hz, 2H), 1.92 (dt, J = 17.0, 10.7 Hz, 6H).
603		450.1	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.18 (s, 1H), 7.02 - 6.86 (m, 2H), 4.81 (s, 2H), 4.00 - 3.76 (m, 5H), 3.42 - 3.26 (m, 4H), 3.07 (s, 6H), 2.20 (q, J = 6.3, 5.8 Hz, 2H), 2.01 - 1.79 (m, 6H).
604		452.1	¹ H NMR (300 MHz, Chloroform-d) δ 8.62 (d, J = 1.9 Hz, 1H), 8.53 (d, J = 1.9 Hz, 1H), 6.87 (d, J = 2.5 Hz, 1H), 6.81 (d, J = 2.6 Hz, 1H), 4.71 (s, 1H), 3.94 - 3.72 (m, 7H), 3.25 (dd, J = 5.8, 3.8 Hz, 4H), 2.29 (d, J = 10.4 Hz, 3H), 2.21 - 2.02 (m, 2H), 1.93 - 1.71 (m, 6H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
605		482.3	
606		450.17	¹ H NMR (300 MHz, Chloroform-d) δ 9.41 (s, 1H), 8.20 (dd, J = 4.8, 2.4 Hz, 2H), 6.98 - 6.77 (m, 3H), 6.44 (dt, J = 7.5, 4.8 Hz, 2H), 4.71 (s, 2H), 3.83 (dd, J = 5.7, 4.0 Hz, 6H), 3.27 (dt, J = 30.0, 4.9 Hz, 5H), 2.13 (s, 2H), 2.00 - 1.68 (m, 6H).
607		469.08	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.79 (d, J = 2.9 Hz, 1H), 6.93 (dd, J = 17.8, 2.5 Hz, 2H), 5.19 - 5.07 (m, 1H), 4.80 (dt, J = 6.7, 3.4 Hz, 1H), 4.32 (q, J = 7.1 Hz, 3H), 3.98 - 3.87 (m, 4H), 3.41 - 3.29 (m, 4H), 2.30 - 2.13 (m, 2H), 2.03 - 1.84 (m, 6H), 1.41 (t, J = 7.1 Hz, 3H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
608		468.63	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 2.0 Hz, 1H), 6.94 (dd, J = 18.9, 2.5 Hz, 2H), 5.42 (d, J = 8.4 Hz, 1H), 4.79 (s, 1H), 4.22 - 4.06 (m, 1H), 4.03 - 3.89 (m, 10H), 3.42 - 3.22 (m, 4H), 2.19 (m, 2H), 2.01 - 1.79 (m, 6H).
609		471.87	¹ H NMR (300 MHz, Chloroform-d) δ 8.72 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.03 - 6.84 (m, 2H), 5.66 (d, J = 9.1 Hz, 1H), 4.82 (s, 1H), 4.09 - 3.84 (m, 7H), 3.35 (dd, J = 5.5, 3.6 Hz, 4H), 2.20 (s, 2H), 1.91 (d, J = 6.4 Hz, 6H).
610		479.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.51 (s, 2H), 6.95 (d, J = 2.5 Hz, 1H), 6.91 (d, J = 2.6 Hz, 1H), 5.39 (d, J = 8.1 Hz, 1H), 4.88 (d, J = 1.8 Hz, 4H), 4.80 (d, J = 6.2 Hz, 1H), 4.13 - 3.98 (m, 1H), 3.96 - 3.89 (m, 4H), 3.38 - 3.30 (m, 4H), 3.06 (s, 1H), 2.28 - 2.11 (m, 2H), 1.98 - 1.84 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
611		344.19	¹ H NMR (300 MHz, Chloroform-d) δ 8.67 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 6.91 (dd, J = 16.1, 2.5 Hz, 2H), 4.81 (q, J = 3.6, 2.4 Hz, 1H), 4.03 - 3.75 (m, 4H), 3.59 (d, J = 2.9 Hz, 2H), 3.45 - 3.28 (m, 4H), 2.20 (dt, J = 9.5, 4.6 Hz, 2H), 1.74 - 1.23 (m, 7H) 1.41 (s, 1H).
612		481.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.83 - 1.96 (m, 6 H), 2.14 - 2.24 (m, 2 H), 3.31 - 3.35 (m, 4 H), 3.44 (s, 3 H), 3.69 - 3.72 (m, 2 H), 3.89 - 3.95 (m, 5 H), 4.06 - 4.09 (m, 2 H), 4.74 - 4.83 (m, 1 H), 5.04 - 5.15 (m, 1 H), 6.90 (d, J = 2.4 Hz, 1 H), 6.94 (d,
613		534.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.39 - 1.75 (m, 6 H), 1.82 - 1.95 (m, 6 H), 2.11 - 2.24 (m, 2 H), 2.37 - 2.63 (m, 4 H), 2.67 - 2.90 (m, 2 H), 3.33 (t, J = 4.7 Hz, 4 H), 3.87 - 3.97 (m, 5 H), 4.01 - 4.20 (m, 2 H), 4.74 - 4.82 (m, 1 H), 4.98 (d, J = 7.9 Hz, 1

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
614	 A quinazoline core with a morpholine group at position 8 and a cyclohexyl ether at position 6. The cyclohexane ring has a 2-fluorophenylamino group attached via its nitrogen atom.	424.2	¹ H NMR (300 MHz, CDCl ₃): ppm 1.83 - 1.95 (m, 6 H), 2.12 - 2.26 (m, 2 H), 3.30 - 3.40 (m, 4 H), 3.85 - 3.96 (m, 5 H), 4.55 - 4.68 (m, 1 H), 4.74 - 4.82 (m, 1 H), 6.10 (dd, J = 7.7, 1.8 Hz, 1 H), 6.18 (dd, J = 8.0, 1.8 Hz, 1 H), 6.90 (d, J = 1.8 Hz, 1 H), 6
615	 A quinazoline core with a morpholine group at position 8 and a cyclohexyl ether at position 6. The cyclohexane ring has a 3-(hydroxymethyl)phenylamino group attached via its nitrogen atom.	436.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.82 - 2.10 (m, 7 H), 2.14 - 2.29 (m, 2 H), 3.26 - 3.40 (m, 4 H), 3.82 - 4.05 (m, 5 H), 4.64 (s, 2 H), 4.75 - 4.84 (m, 1 H), 4.85 - 5.13 (m, 1 H), 6.46 (s, 1 H), 6.51 (d, J = 5.3 Hz, 1 H), 6.90 (d, J = 1.7 Hz, 1 H), 6.94 (d,
616	 A quinazoline core with a morpholine group at position 8 and a cyclohexyl ether at position 6. The cyclohexane ring has a 2-methoxy-5-methylphenylamino group attached via its nitrogen atom.	465.1	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 6.95 (dd, J = 16.6, 2.5 Hz, 2H), 6.30 (s, 1H), 5.02 (s, 1H), 4.82 (s, 1H), 4.38 (d, J = 0.9 Hz, 2H), 3.99 - 3.83 (m, 5H), 3.39 - 3.24 (m, 4H), 2.49 (s, 3H), 2.33 - 2.11 (m, 2H), 1.91 (q, J = 9.0, 6.9 Hz, 6H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
617		437.1	¹ H NMR (300 MHz, Chloroform-d) δ 8.63 (s, 1H), 8.20 (d, J = 4.8 Hz, 2H), 6.90 (dd, J = 16.2, 2.5 Hz, 2H), 6.44 (t, J = 4.8 Hz, 1H), 5.15 (d, J = 8.1 Hz, 1H), 4.90 (d, J = 3.9 Hz, 2H), 4.79 - 4.64 (m, 1H), 3.99 - 3.71 (m, 6H), 3.32 - 3.19 (m, 4H), 2.21 - 2.04 (m, 2H), 1.95 - 1.70 (m, 6H).
618		463.13	¹ H NMR (300 MHz, Chloroform-d) δ 8.63 (d, J = 1.9 Hz, 1H), 8.55 (d, J = 1.9 Hz, 1H), 6.86 (dd, J = 16.5, 2.5 Hz, 2H), 5.94 (s, 1H), 4.73 (s, 1H), 3.85 (dd, J = 5.9, 3.7 Hz, 4H), 3.35 - 3.20 (m, 4H), 2.67 - 2.51 (m, 2H), 2.26 (s, 3H), 2.09 (d, J = 36.8 Hz, 3H), 1.94 - 1.59 (m, 6H), 0.91 (t, J = 7.4 Hz, 3H).
619			¹ H NMR (300 MHz, Chloroform-d) δ 8.77 (s, 1H), 8.29 (d, J = 4.7 Hz, 2H), 7.02 (d, J = 2.5 Hz, 1H), 6.97 (d, J = 2.5 Hz, 1H), 6.53 (t, J = 4.8 Hz, 1H), 5.16 (dd, J = 16.9, 7.5 Hz, 2H), 4.81 (s, 1H), 3.93 (t, J = 4.8 Hz, 5H), 3.34 (t, J = 4.9 Hz, 4H), 2.27 (d, J = 21.1 Hz, 2H), 2.06 - 1.76 (m, 6H), 1.66 (d, J = 6.6 Hz, 3H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
620		435.1	¹ H NMR (300 MHz, Chloroform-d) δ 10.21 (s, 1H), 9.26 (s, 1H), 8.30 (d, J = 4.8 Hz, 2H), 7.09 - 6.88 (m, 2H), 6.54 (t, J = 4.8 Hz, 1H), 4.89 (dq, J = 4.9, 2.5 Hz, 1H), 4.04 (q, J = 6.5 Hz, 1H), 3.99 - 3.87 (m, 5H), 3.55 - 3.39 (m, 5H), 2.35 - 2.17 (m, 3H), 2.09 - 1.82 (m, 6H).
621		449.55	¹ H NMR (400 MHz, CDCl ₃) δ 8.68 (d, J = 1.7 Hz, 1H), 8.61 (d, J = 1.7 Hz, 1H), 7.95 (s, 1H), 6.94 (d, J = 2.1 Hz, 1H), 6.90 (s, 1H), 5.12 (d, J = 7.5 Hz, 1H), 4.77 (s, 1H), 4.60 (t, J = 8.5 Hz, 2H), 4.00 (s, 1H), 3.94 - 3.84 (m, 4H), 3.38 - 3.27 (m, 4H), 3.11 (t, J = 8.3 Hz, 2H), 2.25 - 2.10 (m, 2H), 1.97 - 1.82 (m, 6H).
622		424.2	¹ H NMR (300 MHz, CDCl ₃): ppm 1.81 - 1.98 (m, 6 H), 2.13 - 2.25 (m, 2 H), 3.29 - 3.38 (m, 4 H), 3.75 - 3.85 (m, 1 H), 3.88 - 3.96 (m, 4 H), 4.56 - 4.71 (m, 1 H), 4.75 - 4.83 (m, 1 H), 6.36 (dd, J = 9.1, 3.3 Hz, 1 H), 6.90 (d, J = 1.9 Hz, 1 H), 6.95 (d, J =

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
623		420.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.80 - 1.98 (m, 6 H), 2.13 - 2.23 (m, 2 H), 2.38 (s, 3 H), 3.25 - 3.37 (m, 4 H), 3.62 - 3.77 (m, 1 H), 3.88 - 3.97 (m, 4 H), 4.76 - 4.83 (m, 1 H), 6.23 (d, J = 7.9 Hz, 1 H), 6.43 (d, J = 7.3 Hz, 1 H), 6.91 (d, J = 2.5 Hz, 1 H)
624		494.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.78 - 2.01 (m, 6 H), 2.11 - 2.24 (m, 2 H), 2.33 (s, 6 H), 2.38 - 2.45 (m, 1 H), 2.69 (t, J = 5.2 Hz, 2 H), 3.26 - 3.42 (m, 4 H), 3.84 - 3.98 (m, 4 H), 4.01 (t, J = 5.2 Hz, 2 H), 4.72 - 4.83 (m, 1 H), 4.97 (d, J = 7.5 Hz, 1 H)
625		480.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.81 - 1.95 (m, 6 H), 2.11 - 2.26 (m, 2 H), 3.31 - 3.35 (m, 4 H), 3.44 (s, 3 H), 3.69 - 3.73 (m, 2 H), 3.76 - 3.84 (m, 1 H), 3.89 - 3.93 (m, 4 H), 4.04 - 4.08 (m, 2 H), 4.44 (br s, 1 H), 4.74 - 4.81 (m, 1 H), 6.37 (d, J = 8.9)




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
626		504.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.81 - 1.98 (m, 6 H), 2.12 - 2.25 (m, 2 H), 2.37 - 2.48 (m, 3 H), 2.58 - 2.77 (m, 4 H), 3.04 - 3.16 (m, 4 H), 3.28 - 3.37 (m, 4 H), 3.77 - 3.85 (m, 1 H), 3.87 - 3.97 (m, 4 H), 4.32 - 4.51 (m, 1 H), 4.73 - 4.83 (m, 1 H), 6.38 (
627		436.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.81 - 2.03 (m, 7 H), 2.12 - 2.28 (m, 2 H), 3.28 - 3.39 (m, 4 H), 3.81 (s, 3 H), 3.87 - 3.98 (m, 5 H), 4.76 - 4.84 (m, 1 H), 5.87 (d, J = 1.5 Hz, 1 H), 6.21 (dd, J = 5.9, 1.5 Hz, 1 H), 6.91 (d, J = 1.7 Hz, 1 H), 6.95 (d, J = 1
628		458.05	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 6.94 (dd, J = 14.6, 2.5 Hz, 2H), 5.57 (s, 1H), 4.86 - 4.72 (m, 1H), 4.02 - 3.86 (m, 4H), 3.50 - 3.26 (m, 5H), 2.65 (s, 3H), 2.32 - 2.06 (m, 2H), 2.06 - 1.73 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
629		518.4	¹ H NMR (300 MHz, CDCl ₃) δ 8.62 (s, 1H), 8.54 (s, 1H), 8.15 - 7.93 (m, 1H), 7.40 (d, J = 8.0 Hz, 1H), 6.88 (s, 1H), 6.84 (s, 1H), 6.34 (d, J = 8.6 Hz, 1H), 4.83 - 4.56 (m, 2H), 4.40 - 4.22 (m, 1H), 4.00 - 3.62 (m, 5H), 3.47 - 3.04 (m, 7H), 2.24 - 2.02 (m, 2H), 1.96 - 1.70 (m, 6H).
630		504.22	¹ H NMR (400 MHz, CDCl ₃) δ 8.68 (d, J = 1.8 Hz, 1H), 8.59 (d, J = 1.8 Hz, 1H), 8.06 (s, 1H), 7.55 (d, J = 8.3 Hz, 1H), 6.94 (d, J = 2.3 Hz, 1H), 6.89 (d, J = 2.2 Hz, 1H), 6.41 (d, J = 8.7 Hz, 1H), 4.88 (dd, J = 13.4, 6.7 Hz, 1H), 4.85 - 4.71 (m, 2H), 3.98 - 3.85 (m, 5H), 3.39 - 3.28 (m, 4H), 2.25 - 2.12 (m, 2H), 1.94 - 1.82 (m, 6H).
631		464.53	¹ H NMR (400 MHz, CDCl ₃) δ 8.68 (d, J = 1.7 Hz, 1H), 8.61 (d, J = 1.8 Hz, 1H), 7.01 (d, J = 8.4 Hz, 1H), 6.94 (d, J = 2.3 Hz, 1H), 6.89 (s, 1H), 5.96 (d, J = 8.4 Hz, 1H), 4.75 (s, 1H), 4.43 - 4.35 (m, 2H), 4.22 (s, 1H), 4.19 - 4.10 (m, 2H), 3.98 - 3.86 (m, 4H), 3.82 (s, 1H), 3.36 - 3.28 (m, 4H), 2.23 - 2.08 (m, 2H), 1.94 - 1.79 (m, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
632		476.44	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 7.98 (s, 1H), 7.02 - 6.85 (m, 2H), 5.04 (d, J = 8.2 Hz, 1H), 4.79 (s, 1H), 4.02 (s, 1H), 3.96 - 3.87 (m, 4H), 3.40 - 3.25 (m, 4H), 2.82 (t, J = 5.9 Hz, 2H), 2.73 (t, J = 5.8 Hz, 2H), 2.48 (s, 3H), 2.19 (d, J = 8.5 Hz, 2H), 1.91 (d, J = 5.3 Hz, 5H).
633		435.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.18 (d, J = 5.1 Hz, 1H), 7.01 - 6.88 (m, 2H), 6.42 (d, J = 5.1 Hz, 1H), 5.15 (d, J = 8.0 Hz, 1H), 4.79 (d, J = 5.7 Hz, 1H), 4.17 - 4.00 (m, 1H), 3.98 - 3.83 (m, 4H), 3.67 (td, J = 6.7, 4.0 Hz, 1H), 3.42 - 3.31 (m, 4H), 2.59 (q, J = 7.6 Hz, 2H), 2.20 (q, J = 6.1 Hz, 2H), 2.03 - 1.86 (m, 6H), 1.26 (t, J = 7.6 Hz, 3H).
634		449.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.63 (d, J = 1.9 Hz, 1H), 8.19 (d, J = 5.1 Hz, 1H), 7.01 - 6.88 (m, 2H), 6.42 (d, J = 5.2 Hz, 1H), 5.14 (d, J = 7.9 Hz, 1H), 4.80 (dt, J = 7.8, 3.8 Hz, 1H), 4.06 (d, J = 4.1 Hz, 0H), 3.99 - 3.85 (m, 4H), 3.46 - 3.24 (m, 4H), 2.78 (hept, J = 7.0 Hz, 1H), 2.34 - 2.11 (m, 2H), 2.04 - 1.81 (m, 7H), 1.25 (d, J = 7.0 Hz, 6H).




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
635		481.05	¹ H NMR (300 MHz, Chloroform-d) δ 8.63 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 6.85 (dd, J = 17.6, 2.5 Hz, 2H), 5.86 - 5.70 (m, 1H), 4.89 - 4.67 (m, 2H), 3.94 - 3.77 (m, 5H), 3.34 - 3.20 (m, 4H), 3.02 (q, J = 7.3 Hz, 2H), 2.27 - 2.04 (m, 5H), 1.93 - 1.63 (m, 6H), 1.30 (t, J = 7.3 Hz, 3H).
636		461.1	¹ H NMR (300 MHz, Chloroform-d) δ 8.63 (d, J = 1.9 Hz, 1H), 8.54 (d, J = 1.9 Hz, 1H), 6.86 (dd, J = 16.2, 2.5 Hz, 2H), 5.85 (s, 1H), 4.75 (d, J = 5.5 Hz, 1H), 3.96 - 3.76 (m, 4H), 3.34 - 3.19 (m, 4H), 2.41 (s, 3H), 2.13 (d, J = 8.5 Hz, 2H), 1.82 (q, J = 10.2, 8.8 Hz, 6H), 1.58 - 1.33 (m, 3H), 1.03 - 0.86 (m, 3H).
637		409.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.02 - 6.82 (m, 2H), 5.41 (d, J = 0.8 Hz, 1H), 4.76 (td, J = 5.9, 3.0 Hz, 1H), 4.03 - 3.84 (m, 4H), 3.49 (q, J = 5.8 Hz, 1H), 3.40 - 3.25 (m, 5H), 2.33 - 2.10 (m, 4H), 2.00 - 1.77 (m, 7H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
638		478.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.82 - 1.95 (m, 6 H), 2.14 - 2.26 (m, 2 H), 3.31 - 3.35 (m, 4 H), 3.77 - 3.85 (m, 1 H), 3.90 - 3.93 (m, 4 H), 4.71 - 4.80 (m, 3 H), 4.92 (t, J = 6.6 Hz, 2 H), 5.10 (quint., J = 5.6 Hz, 1 H), 6.39 (d, J = 8.9 Hz, 1 H), 6.90 (d,
639		535.3	¹ H NMR (300 MHz, CDCl ₃): 1.80 - 1.95 (m, 6 H), 2.10 - 2.25 (m, 2 H), 2.52 - 2.64 (m, 4 H), 2.77 (t, J = 5.6 Hz, 2 H), 3.33 (t, J = 4.7 Hz, 4 H), 3.74 (t, J = 4.7 Hz, 4 H), 3.70 - 3.85 (m, 1 H), 3.91 (t, J = 4.7 Hz, 4 H), 4.06 (t, J = 5.6 Hz, 2 H), 4.30 -
640		533.4	¹ H NMR (300 MHz, CDCl ₃): ppm 1.40 - 1.50 (m, 2 H), 1.56 - 1.70 (m, 4 H), 1.84 - 1.94 (m, 6 H), 2.11 - 2.23 (m, 2 H), 2.46 - 2.58 (m, 4 H), 2.71 - 2.80 (m, 2 H), 3.33 (t, J = 4.7 Hz, 4 H), 3.76 - 3.84 (m, 1 H), 3.91 (t, J = 4.7 Hz, 4 H), 4.07 (t, J = 5.7 H




Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
641		371.56	¹ H NMR (400 MHz, CDCl ₃) δ 8.73 - 8.65 (m, 1H), 8.60 (d, J = 1.8 Hz, 1H), 6.93 (d, J = 2.4 Hz, 1H), 6.89 (d, J = 2.3 Hz, 1H), 4.75 (s, 1H), 3.95 - 3.87 (m, 4H), 3.39 - 3.28 (m, 4H), 3.05 - 2.93 (m, 1H), 2.82 - 2.69 (m, 1H), 2.24 - 2.08 (m, 2H), 1.80 - 1.71 (m, 6H), 1.07 (d, J = 6.2 Hz, 6H).
642		424.53	¹ H NMR (400 MHz, CDCl ₃) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 7.78 (d, J = 5.8 Hz, 1H), 6.96 (d, J = 2.4 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 6.36 - 6.26 (m, 1H), 5.99 (d, J = 1.9 Hz, 1H), 4.79 (d, J = 2.4 Hz, 1H), 4.47 (d, J = 7.1 Hz, 1H), 3.99 - 3.85 (m, 4H), 3.56 - 3.49 (m, 1H), 3.39 - 3.26 (m, 4H), 2.28 - 2.16 (m, 2H), 1.98 - 1.80 (m, 6H).
643		492.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.57 - 8.45 (m, 1H), 7.81 (d, J = 8.4 Hz, 1H), 7.18 (d, J = 1.2 Hz, 1H), 6.93 (dd, J = 16.8, 2.5 Hz, 2H), 5.48 (s, 1H), 4.83 (dp, J = 4.5, 2.4 Hz, 1H), 4.23 (dp, J = 8.3, 6.6 Hz, 1H), 4.00 - 3.85 (m, 4H), 3.42 - 3.23 (m, 4H), 2.35 - 2.15 (m, 2H), 1.97 - 1.76 (m, 6H), 1.26 (d, J = 6.6 Hz, 6H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
644		465.16	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 6.93 (dd, J = 17.5, 2.5 Hz, 2H), 5.47 (s, 1H), 4.90 (s, 1H), 4.31 (q, J = 7.1 Hz, 2H), 3.92 (dd, J = 5.9, 3.7 Hz, 4H), 3.60 (s, 1H), 3.38 - 3.28 (m, 4H), 2.40 (s, 3H), 2.19 (d, J = 9.4 Hz, 2H), 1.97 - 1.80 (m, 6H), 1.36 (t, J = 7.1 Hz, 3H), 4.83 - 4.76 (m, 1H).
645		385.51	¹ H NMR (400 MHz, CDCl ₃) δ 8.68 (d, J = 1.8 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 6.94 (d, J = 2.4 Hz, 1H), 6.87 (d, J = 2.4 Hz, 1H), 4.83 (t, J = 6.9 Hz, 2H), 4.76 (s, 1H), 4.45 (t, J = 6.6 Hz, 2H), 4.14 - 4.05 (m, 1H), 3.94 - 3.88 (m, 4H), 3.35 - 3.30 (m, 4H), 2.65 (s, 1H), 2.24 - 2.15 (m, 2H), 1.75 - 1.64 (m, 6H).
646		423.16	¹ H NMR (300 MHz, Chloroform-d) δ 8.71 (d, J = 1.9 Hz, 1H), 8.57 (d, J = 1.9 Hz, 1H), 7.01 - 6.84 (m, 2H), 6.18 (d, J = 1.4 Hz, 1H), 4.80 (s, 1H), 4.43 (s, 1H), 3.93 (t, J = 4.9 Hz, 5H), 3.48 (s, 3H), 3.43 - 3.25 (m, 5H), 2.38 - 1.86 (m, 9H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
647		413.59	¹ H NMR (400 MHz, CDCl ₃) δ 8.67 (d, J = 1.9 Hz, 1H), 8.59 (d, J = 1.9 Hz, 1H), 6.94 (d, J = 2.5 Hz, 1H), 6.88 (d, J = 2.5 Hz, 1H), 4.75 (s, 1H), 4.03 - 3.94 (m, 2H), 3.94 - 3.83 (m, 4H), 3.41 (td, J = 11.7, 2.1 Hz, 2H), 3.36 - 3.25 (m, 4H), 2.89 - 2.77 (m, 2H), 2.24 - 2.13 (m, 2H), 1.85 - 1.68 (m, 8H), 1.40 (ddd, J = 16.2, 12.3, 4.4 Hz, 3H).
648		405.58	¹ H NMR (400 MHz, CDCl ₃) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.17 (dd, J = 8.5, 7.4 Hz, 2H), 6.95 (d, J = 2.5 Hz, 1H), 6.90 (d, J = 2.4 Hz, 1H), 6.69 (t, J = 7.3 Hz, 1H), 6.63 (d, J = 7.9 Hz, 2H), 4.75 (s, 1H), 3.98 - 3.71 (m, 5H), 3.52 (s, 1H), 3.40 - 3.30 (m, 4H), 2.24 - 2.11 (m, 2H), 1.94 - 1.84 (m, 6H).
649		409.17	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 7.10 (d, J = 2.3 Hz, 1H), 6.99 - 6.81 (m, 2H), 5.52 (d, J = 2.3 Hz, 1H), 4.76 (t, J = 5.9 Hz, 1H), 4.06 - 3.86 (m, 5H), 3.72 (s, 3H), 3.49 (t, J = 5.3 Hz, 1H), 3.40 - 3.18 (m, 4H), 2.19 (dq, J = 12.9, 7.1, 6.7 Hz, 2H), 1.90 (dd, J = 9.4, 5.2 Hz, 7H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
650		447.2	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.62 (d, J = 1.9 Hz, 1H), 8.47 - 8.36 (m, 1H), 6.96 (d, J = 2.5 Hz, 1H), 6.90 (d, J = 2.5 Hz, 1H), 6.18 (d, J = 1.2 Hz, 1H), 4.95 - 4.73 (m, 2H), 4.04 - 3.78 (m, 5H), 3.44 - 3.24 (m, 4H), 2.30 - 2.12 (m, 2H), 2.03 - 1.70 (m, 7H), 1.17 - 0.80 (m, 4H).
651		435.18	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.50 (d, J = 1.1 Hz, 1H), 6.93 (dd, J = 15.9, 2.5 Hz, 2H), 6.17 (d, J = 1.1 Hz, 1H), 4.98 (d, J = 8.1 Hz, 1H), 4.86 - 4.75 (m, 1H), 3.97 - 3.87 (m, 4H), 3.39 - 3.29 (m, 4H), 2.61 (q, J = 7.6 Hz, 2H), 2.21 (dt, J = 11.2, 5.0 Hz, 2H), 2.04 - 1.82 (m, 6H), 1.25 (t, J = 7.6 Hz, 3H).
652		449.23	¹ H NMR (300 MHz, Chloroform-d) δ 8.70 (d, J = 1.9 Hz, 1H), 8.61 (d, J = 1.9 Hz, 1H), 8.52 (d, J = 1.1 Hz, 1H), 6.93 (dd, J = 16.0, 2.5 Hz, 2H), 6.19 - 6.12 (m, 1H), 4.91 (d, J = 8.0 Hz, 1H), 4.81 (d, J = 5.4 Hz, 1H), 4.13 - 3.77 (m, 5H), 3.40 - 3.29 (m, 4H), 2.80 (hept, J = 6.8 Hz, 1H), 2.27 - 2.00 (m, 2H), 1.99 - 1.83 (m, 6H), 1.40 - 1.12 (m, 6H).

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
653		451.07	¹ H NMR (300 MHz, Methanol-d ₄ / CDCl ₃) δ 8.82 - 8.68 (m, 2H), 8.61 (d, J = 0.8 Hz, 1H), 7.38 (t, J = 0.8 Hz, 1H), 7.18 (s, 1H), 5.11 - 4.99 (m, 1H), 4.47 - 4.31 (m, 1H), 3.99 - 3.88 (m, 4H), 3.74 - 3.58 (m, 4H), 2.37 - 2.23 (m, 2H), 2.13 - 1.80 (m, 6H).
654		493.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.82 - 1.95 (m, 6 H), 2.11 - 2.24 (m, 2 H), 2.32 (s, 6 H), 2.68 (t, J = 5.7 Hz, 2 H), 3.29 - 3.37 (m, 4 H), 3.76 - 3.86 (m, 1 H), 3.88 - 3.95 (m, 4 H), 4.00 (t, J = 5.7 Hz, 2 H), 4.29 (d, J = 8.2 Hz, 1 H), 4.70 - 4.83 (m, 1 H)
655		548.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.81 - 1.97 (m, 6 H), 2.10 - 2.25 (m, 2 H), 2.33 (s, 3 H), 2.42 - 2.83 (m, 8 H), 2.78 (t, J = 5.6 Hz, 2 H), 3.28 - 3.39 (m, 4 H), 3.75 - 3.85 (m, 1 H), 3.86 - 3.97 (m, 4 H), 4.05 (t, J = 5.6 Hz, 2 H), 4.29 (t, J = 8.4 Hz, 1 H)

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
656		393.1	¹ H NMR (300 MHz, DMSO-d ₆) δ 8.64 (d, J = 1.9 Hz, 1H), 8.47 (d, J = 1.9 Hz, 1H), 8.25 (d, J = 4.8 Hz, 2H), 7.15 (d, J = 7.3 Hz, 1H), 6.59 - 6.47 (m, 2H), 6.31 (d, J = 2.2 Hz, 1H), 5.80 - 5.68 (m, 1H), 4.85 (s, 2H), 4.62 (d, J = 5.3 Hz, 0H), 4.24 (t, J = 7.4 Hz, 2H), 3.83 (s, 1H), 3.71 (dd, J = 8.4, 4.7 Hz, 2H), 2.14 - 1.97 (m, 2H), 1.79 (d, J = 19.4 Hz, 5H).
657		518.32	¹ H NMR (300 MHz, Chloroform-d) δ 8.59 (d, J = 1.9 Hz, 1H), 8.50 (d, J = 1.9 Hz, 1H), 8.11 (s, 2H), 6.91 - 6.70 (m, 2H), 4.66 (s, 1H), 3.92 - 3.56 (m, 7H), 3.45 (d, J = 4.3 Hz, 3H), 3.31 - 3.18 (m, 6H), 2.08 (s, 2H), 1.84 (d, J = 38.5 Hz, 8H).
658		450.3	¹ H NMR (300 MHz, CDCl ₃): ppm 1.37 (t, J = 7.0 HZ, 3 H), 1.84 - 1.94 (m, 6 H), 2.10 - 2.25 (m, 2 H), 3.29 - 3.38 (m, 4 H), 3.75 - 3.85 (m, 1 H), 3.88 - 3.95 (m, 4 H), 3.97 (q, J = 7.0 Hz, 2 H), 4.22 - 4.34 (m, 1 H), 4.73 - 4.80 (m, 1 H), 6.36 (d, J = 9.0 H

Compound No.	Compound Structure	ESMS (M+H)	¹ H NMR
659		451.34	¹ H NMR (300 MHz, Chloroform-d) δ 8.69 (d, J = 1.9 Hz, 1H), 8.60 (d, J = 1.9 Hz, 1H), 6.92 (dd, J = 19.6, 2.5 Hz, 2H), 5.25 (s, 1H), 4.76 (q, J = 8.3, 7.2 Hz, 2H), 3.97 - 3.87 (m, 4H), 3.58 - 3.28 (m, 8H), 2.40 (s, 3H), 2.24 - 2.13 (m, 2H), 2.07 - 1.78 (m, 6H).

Biological assay of compounds of the invention

DNA-PK Inhibition Assay

[00166] Compounds were screened for their ability to inhibit DNA-PK kinase using a standard radiometric assay. Briefly, in this kinase assay the transfer of the terminal ³³P-phosphate in ³³P-ATP to a peptide substrate is interrogated. The assay was carried out in 384-well plates to a final volume of 50 μL per well containing approximately 6 nM DNA-PK, 50 mM HEPES (pH 7.5), 10 mM MgCl₂, 25 mM NaCl, 0.01% BSA, 1 mM DTT, 10 μg/mL sheared double-stranded DNA (obtained from Sigma), 0.8 mg/mL DNA-PK peptide (Glu-Pro-Pro-Leu-Ser-Gln-Glu-Ala-Phe-Ala-Asp-Leu-Trp-Lys-Lys- Lys, obtained from American Peptide), and 100 μM ATP. Accordingly, compounds of the invention were dissolved in DMSO to make 10 mM initial stock solutions. Serial dilutions in DMSO were then made to obtain the final solutions for the assay. A 0.75 μL aliquot of DMSO or inhibitor in DMSO was added to each well, followed by the addition of ATP substrate solution containing ³³P-ATP (obtained from Perkin Elmer). The reaction was started by the addition of DNA-PK, peptide and ds-DNA. After 45 min, the reaction was quenched with 25 μL of 5% phosphoric acid. The reaction mixture was transferred to MultiScreen HTS 384-well PH plates (obtained from Millipore), allowed to bind for one hour, and washed three times with 1% phosphoric acid. Following the addition of 50 μL of Ultima Gold™ high efficiency scintillant (obtained from Perkin Elmer), the samples were counted in a Packard TopCount NXT Microplate Scintillation and Luminescence Counter (Packard BioScience). The K_i values were calculated using

Microsoft Excel Solver macros to fit the data to the kinetic model for competitive tight-binding inhibition.


[00167] Each of compounds **1-291, 295-331, 333-367, 369-523, 525-640, 642-644, 646,** and **648-659** has a K_i of less than 1.0 micromolar for the inhibition of DNA-PK.

Each of compounds **3, 6-14, 16-18, 23-34, 36-37, 39-41, 43-46, 49-72, 74-76, 78, 84-101, 103-123, 127-200, 202-291, 295-299, 305, 307-331, 333-341, 343, 347-366, 369-374, 376-391, 393-519, 521-523, 526-554, 556-610, 612-640, 642-644, 646, 648-655** and **657-659** has a K_i of less than 0.10 micromolar for the inhibition of DNA-PK.

[00168] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it will be readily apparent to those of ordinary skill in the art in light of the teachings of this invention that certain changes and modifications may be made thereto without departing from the spirit or scope of the appended claims.


CLAIMS

1. A compound having the formula:




or a pharmaceutically acceptable salt thereof, wherein

Ring A is



Ring B is



Ring B is optionally substituted with up to 4 fluorine atoms, up to two OH or up to two C₁₋₄alkyl which is optionally substituted with up to 3 fluorine atoms, up to two OH, or up to two OC₁₋₂alkyl groups;

Ring C is a cyclohexane or a cyclobutane ring;

X is -NH-, -O-, or -OC₁₋₄ alkyl-;


each of R¹ and R² is, independently, hydrogen, -C(O)NHR⁴, -C(O)OR⁴, -NHC(O)R⁴, -NHC(O)OR⁴, -NHC(O)NHR⁴, -NHS(O)₂R⁴, -C₀₋₄ alkyl-NHR⁴, or -OR⁴, wherein R¹ and R² cannot simultaneously be hydrogen, and wherein R¹ and R² and the intervening carbon atom can form a dioxane or dioxolane ring;

R³ is hydrogen, -C₁₋₄alkyl, fluoro, chloro, -OC₁₋₂alkyl, -C(O)H, -C(O)OH, -C(O)OC₁₋₂alkyl, -CN, -C(O)NHC₁₋₂alkyl, or -C(O)NH₂, wherein each of said R³ alkyl is optionally substituted with up to 3 fluorine atoms, up to two OH, or up to two OC₁₋₂alkyl groups;

R^4 is hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-5} cycloalkyl, phenyl, a 5-10-membered monocyclic or bicyclic heteroaryl ring selected from pyrrole, imidazole, pyrazole, triazole, thiazole, isothiazole, oxazole, pyridine, pyrimidine, pyrimidinone, pyrazine, pyridazine, and quinoline, or a 4-10-membered monocyclic or bicyclic heterocyclyl ring selected from oxetane, tetrahydrofuran, tetrahydropyran, dihydroisoxazole, pyrimidine-2,4(1H,3H)-dione, dihydrofuroypyrimidine, dihydropyranopyrimidine, dihydropyrrolopyrimidine, tetrahydropteridine, and tetrahydropyridopyrimidine, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl; up to three CN; NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl- $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}$ alkyl) $_2$, $C(O)NH(C_{0-4}$ alkyl- C_{3-5} cycloalkyl), CH_2OR^5 , C_{0-4} alkyl- $C(O)R^5$, C_{0-4} alkyl- $C(O)N(R^5)_2$, C_{0-4} alkyl- $C(O)OR^5$, C_{0-4} alkyl-NHC(O) R^5 , C_{0-4} alkyl-N(R^5) $_2$, oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine, piperazine, furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole, or tetrazole; or up to two OR^5 , wherein each of said optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl; and each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, and pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, and tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.


2. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein Ring C is cyclobutane.

3. The compound of claim 2, having the following formula:




(II), or a pharmaceutically acceptable salt thereof.

4. The compound of claim 3, having the following formula:



(II-A), or a pharmaceutically acceptable salt thereof.

5. The compound of claim 4, having the following formula:




(II-A-1), or a pharmaceutically acceptable salt

thereof.


6. The compound of claim 4 or 5, or a pharmaceutically acceptable salt thereof, wherein R² is -C₀₋₄ alkyl-NHR⁴.

7. The compound of claim 3, having the following formula:



(II-B), or a pharmaceutically acceptable salt thereof.

8. The compound of claim 7, having the following formula:




(II-B-1), or a pharmaceutically acceptable salt

thereof.


9. The compound of claim 7 or 8, or a pharmaceutically acceptable salt thereof, wherein R^1 is $-C_{0-4}$ alkyl-NHR⁴.

10. The compound of any one of claims 2-9, or a pharmaceutically acceptable salt thereof, wherein X is $-O-$ or $-OC_{1-4}$ alkyl-.

11. The compound of any one of claims 2-10, or a pharmaceutically acceptable salt thereof, wherein



12. The compound of claim 11, or a pharmaceutically acceptable salt thereof, wherein



13. The compound of any one of claims 2-12, or a pharmaceutically acceptable salt thereof, wherein R^3 is hydrogen.

14. The compound of any one of claims 2-13, or a pharmaceutically acceptable salt thereof, wherein

R^4 is hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-5} cycloalkyl, or phenyl, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl; up to three CN; NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl- $C(O)NH_2$, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4} \text{ alkyl})_2$, $C(O)NH(C_{0-4} \text{ alkyl-}C_{3-5} \text{ cycloalkyl})$, CH_2OR^5 , C_{0-4} alkyl- $C(O)R^5$, C_{0-4} alkyl- $C(O)N(R^5)_2$, C_{0-4} alkyl- $C(O)OR^5$, C_{0-4} alkyl- $NHC(O)R^5$, C_{0-4} alkyl- $N(R^5)_2$, oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine, piperazine, furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole, tetrazole, or up to two OR^5 , wherein each of said optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl; and each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, and pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, and tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

15. The compound of any one of claims 2-13, or a pharmaceutically acceptable salt thereof, wherein

R⁴ is a 5-10-membered monocyclic or bicyclic heteroaryl ring selected from pyrrole, imidazole, pyrazole, triazole, thiazole, isothiazole, oxazole, pyridine, pyrimidine, pyrimidinone, pyrazine, pyridazine, and quinoline, wherein each of said R⁴ groups is optionally substituted with up to four Br, Cl, F, or C₁₋₄alkyl; up to three CN; NO₂, C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₆cycloalkyl, C₀₋₄alkyl-C₃₋₅ cycloalkyl, C₀₋₄alkyl-O-C₁₋₄alkyl, C₀₋₄alkyl-O-C₀₋₄alkyl-C₃₋₅ cycloalkyl, C(O)OC₁₋₄alkyl, C(O)OC₀₋₄alkyl-C₃₋₅ cycloalkyl, C₀₋₄alkyl-C(O)NH₂, C(O)NHC₁₋₄alkyl, C(O)N(C₁₋₄alkyl)₂, C(O)NH(C₀₋₄alkyl-C₃₋₅ cycloalkyl), CH₂OR⁵, C₀₋₄alkyl-C(O)R⁵, C₀₋₄alkyl-C(O)N(R⁵)₂, C₀₋₄alkyl-C(O)OR⁵, C₀₋₄alkyl-NHC(O)R⁵, C₀₋₄alkyl-N(R⁵)₂, oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine, piperazine, furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole, tetrazole, or up to two OR⁵, wherein each of said optional R⁴ substituents is optionally substituted with up to four fluorine atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, up to two SC₁₋₄alkyl groups, a C(O)C₁₋₄alkyl, a C(O)OC₁₋₄alkyl, or a C(O)OC₀₋₄alkyl-C₃₋₅ cycloalkyl; and each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, and pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, and tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, spirooxetane, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.


16. The compound of any one of claims 2-13, or a pharmaceutically acceptable salt thereof, wherein

R⁴ is a 4-10-membered monocyclic or bicyclic heterocyclyl ring selected from oxetane, tetrahydrofuran, tetrahydropyran, dihydroisoxazole, pyrimidine-2,4(1H,3H)-dione, dihydrofuroypyrimidine, dihydropyranopyrimidine, dihydropyrrolopyrimidine, tetrahydropteridine, and tetrahydropyridopyrimidine, wherein each of said R⁴ groups is optionally substituted with up to four Br, Cl, F, or C₁₋₄alkyl; up to three CN; NO₂, C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₆cycloalkyl, C₀₋₄alkyl-C₃₋₅ cycloalkyl, C₀₋₄alkyl-O-C₁₋₄alkyl, C₀₋₄alkyl-O-C₀₋₄

alkyl-C₃₋₅ cycloalkyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-C(O)NH₂, C(O)NHC₁₋₄ alkyl, C(O)N(C₁₋₄ alkyl)₂, C(O)NH(C₀₋₄ alkyl-C₃₋₅ cycloalkyl), CH₂OR⁵, C₀₋₄ alkyl-C(O)R⁵, C₀₋₄ alkyl-C(O)N(R⁵)₂, C₀₋₄ alkyl-C(O)OR⁵, C₀₋₄ alkyl-NHC(O)R⁵, C₀₋₄ alkyl-N(R⁵)₂, oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine, piperazine, furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole, tetrazole, or up to two OR⁵, wherein each of said optional R⁴ substituents is optionally substituted with up to four fluorine atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, up to two SC₁₋₄alkyl groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, and pyrimidine, a 4-6-membered heterocyclcyl selected from oxetane, tetrahydrofuran, and tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, spirooxetane, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.


17. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein Ring C is cyclohexane.

18. The compound of claim 17, or a pharmaceutically acceptable salt thereof, having the following formula:




19. The compound of claim 17 or 18, or a pharmaceutically acceptable salt thereof, wherein X is -NH-.

20. The compound of any one of claims 17-19, having the following formula:




(III-A), or a pharmaceutically acceptable salt thereof.

21. The compound of claim 20, having the following formula:



(III-A-1) or



(III-A-2), or a pharmaceutically


acceptable salt thereof.

22. The compound of claim 20 or 21, or a pharmaceutically acceptable salt thereof, wherein R^2 is $-C_{0-4}$ alkyl- NHR^4 or $-OR^4$.

23. The compound of claim 22, or a pharmaceutically acceptable salt thereof, wherein R^2 is $-NHR^4$.


24. The compound of claim 22, or a pharmaceutically acceptable salt thereof, wherein R^2 is $-OR^4$.

25. The compound of claim 17-19, having the following formula:




(III-B), or a pharmaceutically acceptable salt thereof.

26. The compound of claim 25 having the following formula:



(III-B-1) or



(III-B-2), or a


pharmaceutically acceptable salt thereof.

27. The compound of claim 25 or 26, or a pharmaceutically acceptable salt thereof, wherein R^1 is $-C_{0-4}$ alkyl- NHR^4 or $-OR^4$.


28. The compound of claim 27, or a pharmaceutically acceptable salt thereof, wherein R^1 is $-NHR^4$.

29. The compound of claim 27, or a pharmaceutically acceptable salt thereof, wherein R^1 is $-OR^4$.

30. The compound of any one of claims 17-29, or a pharmaceutically acceptable salt thereof, wherein




31. The compound of claim 30, or a pharmaceutically acceptable salt thereof, wherein




32. The compound of claim 17 or 18, or a pharmaceutically acceptable salt thereof, wherein X is -O-.

33. The compound of claim 17, 18 or 32, having the following formula:




(III-C), or a pharmaceutically acceptable salt thereof.

34. The compound of claim 33, having the following formula:



(III-C-1) or



(III-C-2), or a pharmaceutically


acceptable salt thereof.

35. The compound of claim 33 or 34, or a pharmaceutically acceptable salt thereof, wherein R^2 is $-C_{0-4}$ alkyl- NHR^4 or $-OR^4$.

36. The compound of claim 35, or a pharmaceutically acceptable salt thereof, wherein R^2 is $-NHR^4$.


37. The compound of claim 35, or a pharmaceutically acceptable salt thereof, wherein R^2 is -OR⁴.

38. The compound of claim 17, 18 or 32, having the following formula:




(III-D), or a pharmaceutically acceptable salt thereof.

39. The compound according to claim 38 having the following formula:



(III-D-1) or



(III-D-2), or a pharmaceutically


acceptable salt thereof.

40. The compound of claim 38 or 39, or a pharmaceutically acceptable salt thereof, wherein R^1 is -C₀₋₄ alkyl-NHR⁴ or -OR⁴.

41. The compound of claim 40, or a pharmaceutically acceptable salt thereof, wherein R^1 is -NHR⁴.

42. The compound of claim 40, or a pharmaceutically acceptable salt thereof, wherein R^1 is -OR⁴.


43. The compound of claim 38 or 39 having the following formula:




(III-D-3), or a pharmaceutically acceptable salt thereof, wherein

Y is -O- or -NH-.

44. The compound of any one of claims 32-43, or a pharmaceutically acceptable salt thereof, wherein



45. The compound of claim 44, or a pharmaceutically acceptable salt thereof, wherein



46. The compound of any one of claims 17-45, or a pharmaceutically acceptable salt thereof, wherein R³ is hydrogen, C₁₋₄alkyl, OC₁₋₂alkyl, C(O)NH₂, or C(O)H, wherein each of said R³ alkyl is optionally substituted with OH.

47. The compound of claim 46, or a pharmaceutically acceptable salt thereof, wherein R³ is hydrogen.

48. The compound of any one of claims 17-47, or a pharmaceutically acceptable salt thereof, wherein

R⁴ is hydrogen, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₅cycloalkyl, or phenyl, wherein each of said R⁴ groups is optionally substituted with up to four Br, Cl, F, or C₁₋₄alkyl; up to three

CN; NO₂, C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₆cycloalkyl, C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-O-C₁₋₄ alkyl, C₀₋₄ alkyl-O-C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-C(O)NH₂, C(O)NHC₁₋₄ alkyl, C(O)N(C₁₋₄ alkyl)₂, C(O)NH(C₀₋₄ alkyl-C₃₋₅ cycloalkyl), CH₂OR⁵, C₀₋₄ alkyl-C(O)R⁵, C₀₋₄ alkyl-C(O)N(R⁵)₂, C₀₋₄ alkyl-C(O)OR⁵, C₀₋₄ alkyl-NHC(O)R⁵, C₀₋₄ alkyl-N(R⁵)₂, oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine, piperazine, furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole, tetrazole, or up to two OR⁵, wherein each of said optional R⁴ substituents is optionally substituted with up to four fluorine atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, up to two SC₁₋₄alkyl groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, and pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, and tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, spirooxetane, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

49. The compound of any one of claims 17-47, or a pharmaceutically acceptable salt thereof, wherein

R⁴ is a 5-10-membered monocyclic or bicyclic heteroaryl ring selected from pyrrole, imidazole, pyrazole, triazole, thiazole, isothiazole, oxazole, pyridine, pyrimidine, pyrimidinone, pyrazine, pyridazine, and quinoline, wherein each of said R⁴ groups is optionally substituted with up to four Br, Cl, F, or C₁₋₄alkyl; up to three CN; NO₂, C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₆cycloalkyl, C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-O-C₁₋₄ alkyl, C₀₋₄ alkyl-O-C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-C(O)NH₂, C(O)NHC₁₋₄ alkyl, C(O)N(C₁₋₄ alkyl)₂, C(O)NH(C₀₋₄ alkyl-C₃₋₅ cycloalkyl), CH₂OR⁵, C₀₋₄ alkyl-C(O)R⁵, C₀₋₄ alkyl-C(O)N(R⁵)₂, C₀₋₄ alkyl-C(O)OR⁵, C₀₋₄ alkyl-NHC(O)R⁵, C₀₋₄ alkyl-N(R⁵)₂, oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine, piperazine, furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole, tetrazole, or up to two OR⁵, wherein each of said optional R⁴ substituents is optionally substituted with up to four fluorine


atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, up to two SC₁₋₄alkyl groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, and pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, and tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, spirooxetane, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

50. The compound of claim 49, or a pharmaceutically acceptable salt thereof, wherein R⁴ is pyridine or pyrimidine, which is optionally substituted with up to four Br, Cl, F, or C₁₋₄alkyl; up to three CN; NO₂, C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₆cycloalkyl, C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-O-C₁₋₄ alkyl, C₀₋₄ alkyl-O-C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-C(O)NH₂, C(O)NHC₁₋₄ alkyl, C(O)N(C₁₋₄ alkyl)₂, C(O)NH(C₀₋₄ alkyl-C₃₋₅ cycloalkyl), CH₂OR⁵, C₀₋₄ alkyl-C(O)R⁵, C₀₋₄ alkyl-C(O)N(R⁵)₂, C₀₋₄ alkyl-C(O)OR⁵, C₀₋₄ alkyl-NHC(O)R⁵, C₀₋₄ alkyl-N(R⁵)₂, oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine, piperazine, furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole, tetrazole, or up to two OR⁵, wherein each of said optional R⁴ substituents is optionally substituted with up to four fluorine atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, up to two SC₁₋₄alkyl groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, and pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, and tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, spirooxetane, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

51. The compound of any one of claims 17-47, or a pharmaceutically acceptable salt thereof, wherein

R^4 is a 4-10-membered monocyclic or bicyclic heterocyclyl ring selected from oxetane, tetrahydrofuran, tetrahydropyran, dihydroisoxazole, pyrimidine-2,4(1H,3H)-dione, dihydrofuropyrimidine, dihydropyranopyrimidine, dihydropyrrolopyrimidine, tetrahydropteridine, and tetrahydropyridopyrimidine, wherein each of said R^4 groups is optionally substituted with up to four Br, Cl, F, or C_{1-4} alkyl; up to three CN; NO_2 , C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, C_{0-4} alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-O- C_{1-4} alkyl, C_{0-4} alkyl-O- C_{0-4} alkyl- C_{3-5} cycloalkyl, $C(O)OC_{1-4}$ alkyl, $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl, C_{0-4} alkyl-C(O)NH₂, $C(O)NHC_{1-4}$ alkyl, $C(O)N(C_{1-4}alkyl)_2$, $C(O)NH(C_{0-4}alkyl-C_{3-5} cycloalkyl)$, CH_2OR^5 , $C_{0-4}alkyl-C(O)R^5$, $C_{0-4}alkyl-C(O)N(R^5)_2$, $C_{0-4}alkyl-C(O)OR^5$, $C_{0-4}alkyl-NHC(O)R^5$, $C_{0-4}alkyl-N(R^5)_2$, oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, pyrrolidine, piperazine, furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, oxadiazole, tetrazole, or up to two OR^5 , wherein each of said optional R^4 substituents is optionally substituted with up to four fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, up to two SC_{1-4} alkyl groups, a $C(O)C_{1-4}$ alkyl, a $C(O)OC_{1-4}$ alkyl, or a $C(O)OC_{0-4}$ alkyl- C_{3-5} cycloalkyl; and each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, and pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, and tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, spirooxetane, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.


52. The compound of claim 32 having the formula:



(IV), or a pharmaceutically acceptable salt thereof, wherein

R³ is hydrogen, -C₁₋₄alkyl, fluoro, chloro, -OC₁₋₂alkyl, -C(O)NH₂, C(O)H, or -CN, wherein each of said R³ alkyl is optionally substituted with OH or up to 3 fluorine atoms;

R⁴ is



X¹ is N, CH, CF, CCl, or CC₁₋₂ alkyl optionally substituted with up to 3 fluorine atoms;

X² is N or CR^{4c}, wherein X¹ and X² cannot simultaneously be N;

each of R^{4a}, R^{4b}, and R^{4c} is, independently, hydrogen, F, Cl, Br, CN, NO₂, C₁₋₄ alkyl, C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-O-C₁₋₄ alkyl, C₀₋₄ alkyl-O-C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl, C(O)NH₂,

C(O)NHC₁₋₄ alkyl, C(O)N(C₁₋₄ alkyl)₂, C(O)NH(C₀₋₄ alkyl-C₃₋₅ cycloalkyl), a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, and piperazine, or a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, and tetrazole, or R^{4c}, R^{4a}, and the intervening atoms form a dihydrofuran, a dihydropyran, or a tetrahydropiperidine heterocyclic ring system;

wherein each of said R^{4a}, R^{4b}, and R^{4c} heterocyclic and heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C₁₋₄ alkyl, up to two OH groups, a

C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and


wherein each of said R^{4a}, R^{4b}, and R^{4c} alkyl and cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.

53. The compound of claim 52, or a pharmaceutically acceptable salt thereof, wherein R³ is hydrogen, -C₁₋₄alkyl, -OC₁₋₂alkyl, or -C(O)NH₂, or -C(O)H, wherein each of said R³ alkyl is optionally substituted with OH.
54. The compound of claim 52 or 53, or a pharmaceutically acceptable salt thereof, wherein R³ is hydrogen.
55. The compound of any one of claims 52-54, or a pharmaceutically acceptable salt thereof, wherein each of X¹ and X² is, independently, CH or N, wherein X¹ and X² cannot simultaneously be N.
56. The compound of any one of claims 52-55, or a pharmaceutically acceptable salt thereof, wherein each of R^{4a} and R^{4b} is, independently, a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran, tetrahydropyran, morpholine, piperidine, and piperazine,
wherein each of said R^{4a} and R^{4b} heterocyclic and heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C₁₋₄ alkyl, up to two OH groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and
wherein each of said R^{4a} and R^{4b} alkyl and cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.
57. The compound of any one of claims 52-55, or a pharmaceutically acceptable salt thereof, wherein each of R^{4a} and R^{4b} is, independently, a heteroaryl ring system selected from furan, oxazole, oxadiazole, pyrrole, pyrazole, triazole, and tetrazole,
wherein each of said R^{4a} and R^{4b} heterocyclic or heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C₁₋₄ alkyl, up to two OH groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and
wherein each of said R^{4a} and R^{4b} alkyl and cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.


58. The compound of any one of claims 52-55, or a pharmaceutically acceptable salt thereof, wherein each of R^{4a} and R^{4b} is, independently, hydrogen, F, Cl, Br, CN, NO₂, C₁₋₄ alkyl, C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₀₋₄ alkyl-O-C₁₋₄ alkyl, C₀₋₄ alkyl-O-C₀₋₄ alkyl-C₃₋₅ cycloalkyl, C₂₋₄ alkenyl, C₂₋₄ alkynyl, C(O)OC₁₋₄ alkyl, C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl, C(O)NH₂, C(O)NHC₁₋₄ alkyl, C(O)N(C₁₋₄ alkyl)₂, or C(O)NH(C₀₋₄ alkyl-C₃₋₅ cycloalkyl), wherein each of said R^{4a} and R^{4b} heterocyclic and heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C₁₋₄ alkyl, up to two OH groups, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and wherein each of said R^{4a} and R^{4b} alkyl and cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.
59. A compound selected from the list of compounds in Table 1, or a pharmaceutically acceptable salt thereof.
60. A compound selected from the list of compounds in Table 2, or a pharmaceutically acceptable salt thereof.
61. A pharmaceutical composition comprising a compound of any one of claims 1-60 or 65-85, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.
62. A compound of any one of claims 1-60 or 65-85, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition comprising said compound or a pharmaceutically acceptable salt thereof, for use in the sensitization of a cell to a therapeutic agent that induces a DNA lesion.
63. A compound of any one of claims 1-60 or 65-85, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition comprising said compound or pharmaceutically acceptable salt thereof, for use in the potentiation of a therapeutic regimen for the treatment of cancer in a patient.

64. A compound of any one of claims 1-60 or 65-85, or a pharmaceutically acceptable salt thereof, or a pharmaceutical composition comprising said compound or pharmaceutically acceptable salt thereof, in combination with one or more additional chemotherapeutic agents for use in the treatment of cancer or inhibiting cancer cell growth in a patient.


65. A compound having the formula:



Ring A is



Ring B is



wherein Ring B is optionally substituted with up to 4 fluorine atoms or up to two C_{1-4} alkyl optionally substituted with up to 3 fluorine atoms, up to two OH, or up to two OC_{1-2} alkyl groups;

X is -NH- or -O-;

each of R^1 and R^2 is, independently, hydrogen, $-C(O)NHR^4$, $-C(O)OR^4$, $-NHC(O)R^4$, $-NHC(O)OR^4$, $-NHC(O)NHR^4$, $-NHS(O)_2R^4$, $-NHR^4$, or $-OR^4$, wherein R^1 and R^2 cannot simultaneously be hydrogen, and wherein R^1 and R^2 and the intervening carbon atom can form a dioxane or dioxolane ring;


R^3 is hydrogen, $-C_{1-4}$ alkyl, fluoro, chloro, $-OC_{1-2}$ alkyl, $-C(O)OH$, $-C(O)OC_{1-2}$ alkyl, $-CN$, $-C(O)NHC_{1-2}$ alkyl, or $-C(O)NH_2$, wherein each of said R^3 alkyl is optionally substituted with up to 3 fluorine atoms, up to two OH, or up to two OC_{1-2} alkyl groups;

R^4 is hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-5} cycloalkyl, phenyl, a 5-10-membered monocyclic or bicyclic heteroaryl ring selected from pyrrole, imidazole, pyrazole, triazole,

thiazole, isothiazole, oxazole, pyridine, pyrimidine, pyrimidinone, pyrazine, pyridazine, and quinoline, or a 4-10-membered monocyclic or bicyclic heterocyclyl ring selected from oxetane, tetrahydrofuran, tetrahydropyran, dihydroisoxazole, pyrimidine-2,4(1H,3H)-dione, dihydrofuropyrimidine, dihydropyranopyrimidine, dihydropyrrolopyrimidine, tetrahydropteridine, and tetrahydropyridopyrimidine, wherein each of said R⁴ groups is optionally substituted with Br, Cl; up to three fluorine atoms, up to three C₁₋₄alkyl; C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₆cycloalkyl, an oxetane ring, a tetrahydrofuran ring, a dihydropyran ring, a tetrahydropyran ring, a pyrrolidine ring, a pyrazole ring, a triazole ring, a tetrazole ring, an oxadiazole ring, CN, CH₂OR⁵, C(O)R⁵, C(O)N(R⁵)₂, C(O)OR⁵, NO₂, NHC(O)R⁵, N(R⁵)₂; or up to two OR⁵, wherein each of said optional R⁴ substituents is optionally substituted with up to 3 fluorine atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, or up to two SC₁₋₄alkyl groups; and each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, and pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, and tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, spirooxetane, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.


66. The compound of claim 65, or a pharmaceutically acceptable salt thereof, wherein X is NH.

67. The compound of claim 66, having the following formula:




(III-A), or a pharmaceutically acceptable salt thereof.

68. The compound of claim 67, having the following formula:




(III-A-1) or



(III-A-2), or a pharmaceutically


acceptable salt thereof, wherein R^2 is NHR^4 .

69. The compound of claim 66, having the following formula:




(III-B), or a pharmaceutically acceptable salt thereof.

70. The compound of claim 69 having the following formula:



(III-B-1) or



(III-B-2), or a


pharmaceutically acceptable salt thereof, wherein R^1 is NHR^4 .

71. The compound of claim 70, or a pharmaceutically acceptable salt thereof, wherein R^3 is hydrogen, C_{1-4} alkyl, fluoro, chloro, $-OC_{1-2}$ alkyl, $-C(O)OH$, $-C(O)OC_{1-2}$ alkyl, $-CN$, $-C(O)NHC_{1-2}$ alkyl, or $-C(O)NH_2$, wherein each of said R^3 alkyl is optionally substituted with up to 3 fluorine atoms, up to two OH, or up to two OC_{1-2} alkyl groups;

R^4 is hydrogen, C_{1-4} alkyl, C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-5} cycloalkyl, phenyl, a 5-10-membered monocyclic or bicyclic heteroaryl ring selected from pyrrole, imidazole, pyrazole, triazole, thiazole, isothiazole, oxazole, pyridine, pyrimidine, pyrimidinone, pyrazine, pyridazine, and quinoline, or a 4-10-membered monocyclic or bicyclic heterocyclyl ring selected from oxetane, tetrahydrofuran, tetrahydropyran, dihydroisoxazole, pyrimidine-2,4(1H,3H)-dione, dihydrofuroypyrimidine, dihydropyranopyrimidine, dihydropyrrolopyrimidine, tetrahydropteridine, and tetrahydropyridopyrimidine, wherein each of said R^4 groups is optionally substituted with Br, Cl; up to three fluorine atoms, up to three C_{1-4} alkyl; C_{2-4} alkenyl, C_{2-4} alkynyl, C_{3-6} cycloalkyl, an oxetane ring, a tetrahydrofuran ring, a dihydropyran ring, a tetrahydropyran ring, a pyrrolidine ring, a pyrazole ring, a triazole ring, a tetrazole ring, an oxadiazole ring, CN, CH_2OR^5 , $C(O)R^5$, $C(O)N(R^5)_2$, $C(O)OR^5$, NO_2 , $NHC(O)R^5$, $N(R^5)_2$; or up to two OR^5 , wherein each of said optional R^4 substituents is optionally substituted with up to 3 fluorine atoms, up to two C_{1-4} alkyl groups, up to two OH groups, up to two OC_{1-4} alkyl groups, or up to two SC_{1-4} alkyl groups; and


each R^5 is, independently, hydrogen, C_{1-4} alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, and pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, and tetrahydropyran, and each R^5 group is optionally substituted with chloro, up to three fluorine atoms, up to two C_{1-2} alkyl, CH_2OH , CN, up to two OH, up to two OC_{1-2} alkyl, pyrrolidine, or triazole, or two R^5 groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.

72. The compound of claim 70 or 71, having the formula:




73. The compound of claim 65, or a pharmaceutically acceptable salt thereof, wherein X is O.


74. The compound of claim 73, having the following formula:




75. The compound of claim 74, having the following formula:




76. The compound of claim 74, having the following formula:




77. The compound of claim 73, having the following formula:



78. The compound of claim 77, having the following formula:




79. The compound of claim 77, having the following formula:




(III-D-3), or a pharmaceutically acceptable salt thereof, wherein

Y is -O- or -NH-.

80. The compound of claim 77 having the following formula:



(III-D-4) or



(III-D-5), or a

pharmaceutically acceptable salt thereof.

81. The compound of claim 80, or a pharmaceutically acceptable salt thereof, wherein


R³ is hydrogen, C₁₋₄alkyl, fluoro, chloro, -OC₁₋₂alkyl, -C(O)OH, -C(O)OC₁₋₂alkyl, -CN, -C(O)NHC₁₋₂alkyl, or -C(O)NH₂, wherein each of said R³ alkyl is optionally substituted with up to 3 fluorine atoms, up to two OH, or up to two OC₁₋₂alkyl groups;

R⁴ is hydrogen, C₁₋₄alkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₅cycloalkyl, phenyl, a heteroaryl ring system selected from pyrrole, imidazole, pyrazole, triazole, thiazole, isothiazole, oxazole, pyridine, pyrimidine, pyrimidinone, pyrazine, pyridazine, and quinoline, or a heterocyclic ring selected from oxetane, tetrahydrofuran, tetrahydropyran, dihydroisoxazole, pyrimidine-2,4(1H,3H)-dione, dihydrofuropyrimidine, dihydropyranopyrimidine, dihydropyrrolopyrimidine, tetrahydropteridine, and tetrahydropyridopyrimidine, wherein each of said R⁴ groups is optionally substituted with Br, Cl; up to three fluorine atoms, up to three C₁₋₄alkyl; C₂₋₄alkenyl, C₂₋₄alkynyl, C₃₋₆cycloalkyl, CN, NO₂, C(O)R⁵, C(O)N(R⁵)₂, C(O)OR⁵, NHC(O)R⁵, N(R⁵)₂; or up to two OR⁵, a heterocyclic ring system selected from oxetane, tetrahydrofuran, dihydropyran, tetrahydropyran, and pyrrolidine, a heteroaromatic

ring system selected from pyrazole, triazole, tetrazole, and oxadiazole, wherein each of said optional R⁴ substituents is optionally substituted with up to 3 fluorine atoms, up to two C₁₋₄alkyl groups, up to two OH groups, up to two OC₁₋₄alkyl groups, or up to two SC₁₋₄alkyl groups; and

each R⁵ is, independently, hydrogen, C₁₋₄alkyl, a 5-6-membered heteroaryl selected from imidazole, triazole, thiazole, pyridine, and pyrimidine, a 4-6-membered heterocyclyl selected from oxetane, tetrahydrofuran, and tetrahydropyran, and each R⁵ group is optionally substituted with chloro, up to three fluorine atoms, up to two C₁₋₂alkyl, CH₂OH, CN, up to two OH, up to two OC₁₋₂alkyl, pyrrolidine, or triazole, or two R⁵ groups together with the intervening nitrogen atom form a morpholine ring, azetidine ring, pyrrolidine ring, piperidine ring, or piperazine ring.


82. A compound having the formula:



(IV), or a pharmaceutically acceptable salt thereof, wherein

R³ is hydrogen, -C₁₋₄alkyl, fluoro, chloro, -OC₁₋₂alkyl, or -CN, wherein each of said R³ alkyl is optionally substituted with up to 3 fluorine atoms;

R⁴ is



X¹ is N, CH, CF, CCl, or CC₁₋₂alkyl optionally substituted with up to 3 fluorine atoms;


X² is N or CR^{4c};

each of R^{4a}, R^{4b}, and R^{4c} is, independently, hydrogen, F, Cl, Br, CN, NO₂, C₁₋₄alkyl, C₀₋₄alkyl-C₃₋₅cycloalkyl, C₀₋₄alkyl-O-C₁₋₄alkyl, C₀₋₄alkyl-O-C₀₋₄alkyl-C₃₋₅cycloalkyl, C₂₋₄alkenyl, C₂₋₄alkynyl, C(O)OC₁₋₄alkyl, C(O)OC₀₋₄alkyl-C₃₋₅cycloalkyl, C(O)NH₂, C(O)NHC₁₋₄alkyl, C(O)N(C₁₋₄alkyl)₂, C(O)NH(C₀₋₄alkyl-C₃₋₅cycloalkyl), a heterocyclic ring system selected from oxetane, azetidine, tetrahydrofuran, dihydropyran,


wherein each of said R^{4a}, R^{4b}, and R^{4c} heterocyclic and heteroaryl ring systems is optionally substituted with up to four fluorine atoms, up to two C₁₋₄ alkyl, a C(O)C₁₋₄ alkyl, a C(O)OC₁₋₄ alkyl, or a C(O)OC₀₋₄ alkyl-C₃₋₅ cycloalkyl; and

wherein each of said R^{4a}, R^{4b}, and R^{4c} alkyl and cycloalkyl is optionally substituted with up to 2 non-geminal OH groups or up to 3 fluorine atoms.


 is 
 or 



85. A compound, wherein the compound is



, or a pharmaceutically acceptable salt thereof.