COMPUTATIONAL METHOD FOR IDENTIFYING ADHESIN AND ADHESIN-LIKE PROTEINS OF THERAPEUTIC POTENTIAL

Field of the present Invention

A computational method for identifying adhesin and adhesin-like proteins; computer system for performing the method; and genes and proteins encoding adhesin and adhesin-like proteins

Background and object of the present Invention

The progress in genome sequencing projects has generated a large number of inferred protein sequences from different organisms. It is expected that the availability of the information on the complete set of proteins from infectious human pathogens will enable us to develop novel molecular approaches to combat them. A necessary step in the successful colonization and subsequent manifestation of disease by microbial pathogens is the ability to adhere to host cells.

Microbial pathogens encode several proteins known as adhesins that mediate their adherence to host cell surface receptors, membranes, or extracellular matrix for successful colonization. Investigations in this primary event of host-pathogen interaction over the past decades have revealed a wide array of adhesins in a variety of pathogenic microbes. Presently, substantial information on the biogenesis of adhesins and the regulation of adhesin factors is available. One of the best understood mechanisms of bacterial adherence is attachment mediated by pili or fimbriae. Several afimbrial adhesins also have been reported. In addition, limited knowledge on the target host receptors also has been gained (Finlay, B.B. and Falkow, S 1997).

New approaches to vaccine development focus on targeting adhesins to abrogate the colonization process (Wizemann, *et al* 1999). However, the specific role of particular adhesins has been difficult to elucidate. Thus, prediction of adhesins or adhesin-like proteins and their functional characterization is likely to aid not only in deciphering the molecular mechanisms of host pathogen interaction but also in developing new vaccine formulations, which can be tested in suitable experimental model systems.

One of the best understood mechanisms of bacterial adherence is attachment mediated by pili or fimbriae. For example, FimH and PapG adhesins of *Escherichia coli* (Maurer, L., Orndorff, P.(1987), Bock, K., *et al.*(1985). Other examples of pili group adhesins include type IV pili in *Pseudomonas aeruginosa*, Neisseria species, Moraxella species, Enteropathogenic *Escherichia coli* and *Vibrio cholerae* (Sperandio V *et al* (1996).

Several afimbrial adhesins are HMW proteins of *Haemophilus influenzae* (van Schilfgaarde 2000), the filamentous hemagglutinin, pertactin, of *Bordetella pertussis* (Bassinet *et al* 2000), the BabA of *H. pylori* (Yu J *et al* 2002) and the YadA adhesin of *Yersinia enterocolitica* (Neubauer *et al* 2000). The intimin receptor protein (Tir) of Enteropathogenic *E. coli* (EPEC) is another type of adhesin (Ide T *et al* 2003). Other class of adhesins includes MrkD protein of *Kleibsella pneumoniae*, Hia of *H. influenzae* (St Geme *et al* 2000), Ag I/II of *Streptococcus mutans* and SspA, SspB of *Streptococcus gordonii* (Egland *et al* 2001), FnbA, FnbB of *Staphylococcus aureus* and SfbI, protein F of *Streptococcus pyogenes* , the PsaA of *Streptococcus pneumoniae* (De *et al* 2003).

A known example of adhesins approved as vaccine is the acellular pertussis vaccine containing FHA and pertactin against *B. pertussis* the causative agent of whooping cough (Halperin, S *et al* 2003). Immunization with FimH is being evaluated for protective immunity against pathogenic *E. coli* (Langermann S *et al* 2000), in *Streptococcus pneumoniae*, PsaA is being investigated as a potential vaccine candidate against pneumococcal disease (Rapola, S *et al* 2003). Immunization results with BabA adhesin showed promise for developing a vaccine against *H. pylori* (Prinz, C *et al* 2003). A synthetic peptide sequence anti-adhesin vaccine is being evaluated for protection against *Pseudomonas aeruginosa* infections.

Screening for adhesin and adhesin like proteins by conventional experimental method is laborious, time consuming and expensive. As an alternative, homology search is used to facilitate the identification of adhesins. Although, this procedure is useful in the analysis of genome organization (Wolf *et al* 2001) and of metabolic pathways (Peregrin-Alvarez *et al* 2003, Rison *et al* 2002), it is somewhat limited in allowing functional predictions when the homologues are not functionally characterized or the sequence divergence is high. Assignment of functional roles to proteins based on this technique has been possible for only about 60% of the predicted protein sequences (Fraser *et al* 2000). Thus, we explored the possibility of developing a non-homology method based on sequence composition properties combined with the power of the Artificial Neural Networks to identify adhesins and adhesin-like proteins in species belonging to wide phylogenetic spectrum.

Twenty years ago, Nishikawa *et al* carried out some of the early attempts to classify proteins into different groups based on compositional analysis (Nishikawa *et al* 1983).

More recently, the software PropSearch was developed for analyzing protein sequences where conventional alignment tools fail to identify significantly similar sequences (Hobohm, U. and Sander, C 1995). PropSearch uses 144 compositional properties of protein sequences to detect possible structural or functional relationships between a new sequence and sequences in the database. Recently the compositional attributes of proteins have been used to develop softwares for predicting secretory proteins in bacteria and apicoplast targeted proteins in *Plasmodium falciparum* by training Artificial Neural Networks (Zuegge *et al* 2001).

Zuegge et al have used the 20 amino acid compositional properties. Their objective was to extract features of apicoplast targeted proteins in *Plasmodium falciparum*. This is distinct from our software SPAAN that focuses on adhesins and adhesin-like proteins involved in host-pathogen interaction.

Hobohm and Sander have used 144 compositional properties including isoelectric point and amino acid and dipeptide composition to generate hypotheses on putative functional role of proteins that are refractory to analysis using other sequence alignment based approaches like BLAST and FASTA. Hobohm and Sander do not specifically address the issue of adhesins and adhesin-like proteins, which is the focus of SPAAN

Nishikawa et al had originally attempted to classify proteins into various functional groups. This was a curiosity driven exercise but eventually lead to the development of a software to discriminate extra-cellular proteins from intracellular proteins. This work did not address the issue of adhesins and adhesin-like proteins, which is the focus of SPAAN.

Thus, none of the aforementioned research groups have been able to envisage the methodology of the instant application. The inventive method of this application provides novel proteins and corresponding gene sequences.

Adhesins and adhesin-like proteins mediate host-pathogen interactions. This is the first step in colonization of a host by microbial pathogens. Attempts Worldwide are focused on designing vaccine formulations comprising adhesin proteins derived from pathogens. When immunized, host will have its immune system primed against adhesins for that pathogen. When a pathogen is actually encountered, the surveillance mechanism will recognize these adhesins, bind them through antigen-antibody interactions and neutralize the pathogen through complement mediate cascade and other related clearance mechanisms. This strategy has been successfully employed in

the case of Whooping cough and is being actively pursued in the case of Pneumonia, Gastric Ulcer and Urinary tract infections.

Objects of the present Invention

The main object of the present invention is to provide a computational method for identifying adhesin and adhesin-like proteins of therapeutic potential.

Another object of invention is to provide a method for screening the proteins with unique compositional characteristics as putative adhesins in different pathogens.

Yet, another object of the invention is providing the use of gene sequences encoding the putative adhesin proteins useful as preventive therapeutics.

Summary of the present Invention

A computational method for identifying adhesin and adhesin-like proteins, said method comprising steps of computing the sequence-based attributes of protein sequences using five attribute modules of software SPAAN, (i) amino acid frequencies, (ii) multiplet frequency, (iii) dipeptide frequencies, (iv) charge composition, and (v) hydrophobic composition, training the artificial neural Network (ANN) for each of the computed five attributes, and identifying the adhesin and adhesin-like proteins having probability of being an adhesin (P_{ad}) as ≥ 0.51; a computer system for performing the method; and genes and proteins encoding adhesin and adhesin-like proteins

Detailed description of the present Invention

Accordingly, the present invention relates to a computational method for identifying adhesin and adhesin-like proteins, said method comprising steps of computing the sequence-based attributes of protein sequences using five attribute modules of software SPAAN, (i) amino acid frequencies, (ii) multiplet frequency, (iii) dipeptide frequencies, (iv) charge composition, and (v) hydrophobic composition, training the artificial neural Network (ANN) for each of the computed five attributes, and identifying the adhesin and adhesin-like proteins having probability of being an adhesin (P_{ad}) as ≥ 0.51; a computer system for performing the method; and genes and proteins encoding adhesin and adhesin-like proteins

In an embodiment of the present invention, wherein the invention relates to a computational method for identifying adhesin and adhesin-like proteins, said method comprising steps of:

a. computing the sequence-based attributes of protein sequences using five attribute modules of a neural network software, wherein the attributes

are software, (i) amino acid frequencies, (ii) multiplet frequency, (iii) dipeptide frequencies, (iv) charge composition, and (v) hydrophobic composition,

b. training the artificial neural Network (ANN) for each of the computed five attributes, and

c. identifying the adhesin and adhesin-like proteins having probability of being an adhesin (P_{ad}) as ≥ 0.51.

In another embodiment of the present invention, wherein the invention relates to a method wherein the protein sequences is obtained from pathogens, eukaryotes, and multicellular organisms.

In an embodiment of the present invention, wherein the invention relates to a method, wherein the protein sequences are obtained from the pathogens selected from a group of organisms comprising *Escherichia coli*, *Haemophilus influenzae*, *Helicobacter pylori*, *Mycoplasma pneumoniae*, *Mycobacterium tuberculosis*, *Rickettsiae prowazekii*, *Porphyromonas gingivalis*, *Shigella flexneri*, *Streptococcus mutans*, *Streptococcus pneumoniae*, *Neisseria meningitides*, *Streptococcus pyogenes*, *Treponema pallidum* and Severe Acute Respiratory Syndrome associated human coronavirus (SARS).

In yet another embodiment of the present invention, wherein the method of the invention is a non-homology method.

In still another embodiment of the present invention, wherein the invention relates to the method using 105 compositional properties of the sequences.

In still another embodiment of the present invention, wherein the invention relates to a method showing sensitivity of at least 90%.

In still another embodiment of the present invention, wherein the invention relates to the method showing specificity of 100%.

In still another embodiment of the present invention, wherein the invention relates to a method identifying adhesins from distantly related organisms.

In still another embodiment of the present invention, wherein the invention relates to the neural network has multi-layer feed forward topology, consisting of an input layer, one hidden layer, and an output layer.

In still another embodiment of the present invention, wherein the invention relates to the number of neurons in the input layer are equal to the number of input data points for each attribute.

In still another embodiment of the present invention, wherein the invention relates to the "P_{ad}" is a weighted linear sum of the probabilities from five computed attributes.

In still another embodiment of the present invention, wherein the invention relates to each trained network assigns a probability value of being an adhesin for the protein sequence.

In still another embodiment of the present invention, wherein the invention relates to a computer system for performing the method of claim 1, said system comprising a central processing unit, executing SPAAN program, giving probabilities based on different attributes using Artificial Neural Network and in built other programs of assessing attributes, all stored in a memory device accessed by CPU, a display on which the central processing unit displays the screens of the above mentioned programs in response to user inputs; and a user interface device.

In still another embodiment of the present invention, wherein the invention relates to a set of 274 annotated genes encoding adhesin and adhesin-like proteins, having SEQ ID Nos. 385 to 658.

In still another embodiment of the present invention, wherein the invention relates to a set of 105 hypothetical genes encoding adhesin and adhesin-like proteins, having SEQ ID Nos. 659 to 763.

In still another embodiment of the present invention, wherein the invention relates to a set of 279 annotated adhesin and adhesin-like proteins of SEQ ID Nos. 1 to 279.

In still another embodiment of the present invention, wherein the invention relates to a set of 105 hypothetical adhesin and adhesin-like proteins of SEQ ID Nos. 280 to 384.

One more embodiment of the present invention, wherein the invention also relates to a fully connected multilayer feed forward Artificial Neural Network based on the computational method as claimed in claim 1, comprising of an input layer, a hidden layer and an output layer which are connected in the said sequence, wherein each neuron is a binary digit number and is connected to each neuron of the subsequent layer for identifying adhesin or adhesin like proteins, wherein the program steps comprise:-

[a] feeding a protein sequence in FASTA format; [b] processing the sequence obtained in step [a] through the 5 modules named A, C, D, H and M, wherein attribute A represents an amino acid composition, attribute C represents a charge composition, attribute D represents a dipeptide composition of the 20 dipeptides [NG, RE, TN, NT, GT, TT, DE, ER, RR, RK, RI, AT, TS, IV, SG, GS, TG, GN, VI and HR], attribute H

represents a hydrophobic composition and attribute M represents amino acid frequencies in multiplets to quantify 5 types of compositional attributes of the said protein sequence to obtain numerical input vectors respectively for each of the said attributes wherein the sum of numerical input vectors is 105; [c] processing of the numerical input vectors obtained in step [b] by the input neuron layer to obtain signals, wherein the number of neurons is equal to the number of numerical input vectors for each attribute; [d] processing of signals obtained from step [c] by the hidden layer to obtain synaptic weighted signals, wherein the optimal number of neurons in the hidden layer was determined through experimentation for minimizing the error at the best epoch for each network individually; [e] delivering synaptic weighted signals obtained from step [d] to the output layer for assigning of a probability value for each protein sequence fed in step [a] as being an adhesin by each network module; [f] using the individual probabilities obtained from step [e] for computing the final probability of a protein sequence being an adhesin denoted by the P_{ad} value, which is a weighted average of the individual probabilities obtained from step [e] and the associated fraction of correlation which is a measure of the strength of the prediction.

In still another embodiment of the present invention, wherein the input neuron layer consists of a total of 105 neurons corresponding to 105 compositional properties.

In still another embodiment of the present invention, wherein the hidden layer comprises of neurons represented as 30 for amino acid frequencies, 28 for multiplet frequencies, 28 for dipeptide frequencies, 30 for charge composition and 30 for hydrophobic composition.

In still another embodiment of the present invention, wherein the output layer comprises of neurons to deliver the output values as probability value for each protein sequence.

Identification of novel adhesins and their characterization are important for studying host-pathogen interactions and testing new vaccine formulations. We have employed Artificial Neural Networks to develop an algorithm SPAAN (Software for Prediction of Adhesin and Adhesin-like proteins using Neural Networks) that can identify adhesin proteins using 105 compositional properties of a protein sequence. SPAAN could correctly predict well characterized adhesins from several bacterial species and strains. SPAAN showed 89% sensitivity and 100% specificity in a test data set that did not

contain proteins in the training set. Putative adhesins identified by the software can serve as potential preventive therapeutics.

The present invention provides a novel computational method for identifying adhesin and adhesin-like proteins of therapeutic potential. More particularly, the present invention relates to candidate genes for these adhesins. The invention further provides new leads for development of candidate genes, and their encoded proteins in their functional relevance to preventive approaches. This computational method involves calculation of several sequence attributes and their subsequent analyses lead to the identification of adhesin proteins in different pathogens. Thus, the present invention is useful for identification of the adhesin proteins in pathogenic organisms. The adhesin proteins from different genomes constitute a set of candidates for functional characterization through targeted gene disruption, microarrays and proteomics. Further, these proteins constitute a set of candidates for further testing in development of preventive therapeutics. Also, are provided the genes encoding the candidate adhesin proteins.

The present method offers novelty in the principles used and the power of Neural Networks to identify new adhesins compared to laborious and time consuming conventional methods. The present method is based on compositional properties of proteins instead of sequence alignments. Therefore this method has the ability to identify adhesin and adhesin like proteins from bacteria belonging to a wide phylogenetic spectrum. The predictions made from this method are readily verifiable through independent analysis and experimentation. The invention has the potential to accelerate the development of new preventive therapeutics, which currently requires high investment in terms of requirement of skilled labor and valuable time.

The present invention relates to a computational method for the identification of candidate adhesin proteins of therapeutic potential. The invention particularly describes a novel method to identify adhesin proteins in different genomes of pathogens. These adhesin proteins can be used for developing preventive therapeutics.

Accordingly, a computational method for identifying adhesin and adhesin-like proteins of therapeutic potential which comprises calculation of 105 compositional properties under the five sequence attributes, namely, Amino Acid frequency, Multiplet frequency, dipeptide frequency, charge composition and hydrophobic composition; and then training Artificial Neural Network (ANN, Feed Forward Error Back Propagation)

using these properties for differentiating between adhesin and non-adhesin class of proteins. This computational method involves quantifying 105 compositional attributes of query proteins and qualifying them as adhesins or non-adhesins by a P_{ad} value (Probability of being an adhesin). The present invention is useful for identification of adhesin and adhesin-like proteins in pathogenic organisms. These newly identified adhesin and adhesin-like proteins constitute a set of candidates for development of new preventive therapeutics that can be tested in suitable experimental model systems readily. In addition, the genes encoding the candidate adhesin and adhesin-like proteins are provided.

The invention provides a set of candidate adhesin and adhesin-like proteins and their coding genes for further evaluation as preventive therapeutics. The method of invention is based on the analysis of protein sequence attributes instead of sequence patterns classified to functional domains. Present method is less dependent on sequence relationships and therefore offers the potential power of identifying adhesins from distantly related organisms. The invention provides a computational method, which involves prediction of adhesin and adhesin-like proteins using Artificial Neural Networks. The proteins termed adhesin were found to be predicted with a high probability (P_{ad} 0.51) in various pathogens. Some adhesin sequences turned out to be identical or homologous to proteins that are antigenic or implicated in virulence. By this approach, proteins could be identified and short-listed for further testing in development of new vaccine formulations to eliminate diseases caused by various pathogenic organisms.

DESCRIPTION OF TABLES

Table 1: Output file format given by SPAAN.

Table 2: Organism Name, Accession number, Number of base pairs, Date of release and Total number of proteins.

Table 3. Prediction of well characterized adhesins from various bacterial pathogens using SPAAN.

Table 4. Analysis of predictions made by SPAAN on genome scans of a few selected pathogenic organisms.

Table 5: GI numbers and Gene IDs of new putative adhesins predicted by SPAAN in the genomes listed in Table 2.

Table 6: GI numbers and Gene IDs of hypothetical proteins predicted as putative adhesins by SPAAN in the genomes listed in Table 2.

Table 7: The list of 198 adhesins found in bacteria

Brief description of the accompanying drawings

Figure 1 shows the Neural Network architecture

Figure 2 shows assessment of SPAAN using defined test dataset.

Figure 3 (a) shows Histogram plots of the number of proteins in the various P_{ad} value ranges are shown. **(b)** Pairwise sequence relationships among the adhesins were determined using CLUSTAL W and plotted on X-axis. Higher scores indicate similar pairs. **(c)** plot for non-adhesins. Data are plotted in the 4 quadrant format for clear inspection.

Software program was written in C Language and operated on Red Hat Linux 8.0 operating system. The computer program accepts input protein sequences in FastA format and produces a tabulated output. The output Table contains one row for each protein listing the probability outputs of each of the five modules, a weighted average probability of these five modules (P_{ad}), and the function of the protein as described in the input sequence file. This software is called SPAAN (A Software for Prediction of Adhesins and Adhesin-like proteins using Neural Networks) and a software copyright has been filed. Although this software has multiple modules, the running of these modules have been integrated and automated. The user only needs to run one command.

AAcompo.c:

Input: File containing protein sequences in the fasta format.

Output: File containing frequencies of all 20 AAs for each protein in one row.

charge.c:

Input: File containing protein sequences in the fasta format.

Output: File containing frequency of charged amino acids (R, K, E and D) and moments (up to 18th order) of the positions of charged amino acids.

hdr.c:

Input: File containing protein sequences in the fasta format.

Output: File containing frequencies of 5 groups of amino acids formed on the basis their Hydrophobicity and moments of their positions up to 5th order.

multiplets.c:

Input: File containing protein sequences in the fasta format.

Output: File containing fractions of multiplets of each of the 20 amino acids.

querydipep.c:

Input: File.1 containing protein sequences in the fasta format.

File.2 containing list of the significant dipeptides in dipeptide analysis.

Output: File containing frequencies of the dipeptides listed in the input File.2 for each protein in the input File.1.

train.c:

Input: File containing following specifications –

1. Number of input and output parameters.
2. Number of nodes in the hidden layers.
3. Names of the training, validate and test data files.
4. Learning rate, coefficient of moment.
5. Maximum number of cycles for training.

Output: Outputs are as follows.

1. Output of the trained NN for the test data set.
2. Values of the weight connections in the trained NN.
3. Some extra information about training.

recognize.c:

Input: File containing following specifications –

1. Number of input and output parameters.
2. Number of nodes in the hidden layers.
3. Names of the query input file.
4. Name of the file containing values of the weight connections for trained NN.
5. Name of the output file.

Output: Outputs for the query entries calculated by the trained NN.

standard.c:

Input: File containing protein sequences in fasta format.

Output: File containing protein sequences in fasta format with all the new line characters removed lying within a sequence.

filter.c:

Input: File containing protein sequences in fasta format.

Output: File containing protein sequences from the input except those which are short in length (<50 AAs) and which contain any amino acid other than the 20 known amino acids.

The five attributes:

Amino Acid frequencies

Amino acid frequency f_i = (counts of ith amino acid in the sequence) / l ; i, = 1...20, l is the length of the protein.

Multiplet frequency

Multiplets are defined as homopolymeric stretches $(X)_n$ where X is any of the 20 amino acids and n is an integer > 2. After identifying all the multiplets, the frequencies of the amino acids in the multiplets were computed as

$f_i(m)$ = (counts of i^{th} amino acid occurring as multiplet) / l

Dipeptide frequencies

The frequency of a dipeptide (i, j) f_{ij} = (counts of ij^{th} dipeptide) / (total dipeptide counts); i, j ranges from 1 to 20.

It has been found that dipeptide repeats in proteins are important for functional expression of the clumping factor present on *Staphylococcus aureus* cell surface that binds to fibrinogen (Hartford *et al* 1999). Thus we included the dipeptide frequency module. The total number of dipeptides is 400. For optimal training of Neural Network, the ratio of total number of input vectors to the total number of weight connections must be around 2 to avoid over fitting (Andrea *et al*). Therefore, we identified the dipeptides whose frequencies in the adhesin data set (469 proteins, see database construction) were significantly different from that in the non-adhesin dataset (703 proteins) using *t-test*. The frequencies of top 20 dipeptides (when arranged in the descending order of the p-values of *t-test*), were fed to the Neural Network. These dipeptides were (using single letter IUPAC-IUB code) NG, RE, TN, NT, GT, TT, DE,

ER, RR, RK, RI, AT, TS, IV, SG, GS, TG, GN, VI, AND HR. With frequency inputs for 20 dipeptides and 28 neurons in the 2nd layer, the total number of weight connections is 588, and is in keeping with the criterion of avoiding over fitting.

Charge composition

The input frequency of charged amino acids (R, K, E and D considering the ionization properties of the side chains at pH 7.2) given by f_c = (counts of charged amino acids) / l

Further, information on the characteristics of the distribution of the charged amino acids in a given protein sequence was provided by computing the moments of the positions of the occurrences of the charged amino acids. Since moments characterize the patterns of distribution such as skewness and kurtosis (sharpness of the peak) we have used them to represent the distribution patterns of the charged residues in the sequence.

The general expression to compute moments of a given order; say 'i' is

M_r = r^{th} order moment of the positions of charged amino acids

$$= \sum \frac{(X_i - X_m)^r}{N}$$

Where, X_m = mean of all positions of charged amino acids

X_i = position of i^{th} charged amino acid

N = number of charged amino acids in the sequence

The moments 2^{nd} to 19^{th} order were used to train the ANN constituting a total 20 inputs in addition to frequency of charged amino acids and the length of the protein. The upper limit of 19^{th} order was set based on assessments of sensitivity and specificity on a small dataset of adhesins and non-adhesins. Moments of order greater than 19 were not useful in improvement of performance.

Hydrophobic composition

A given protein sequence was digitally transformed using the hydrophobic scores of the amino acids according to Brendel *et al.* (43). The scores for five groups of amino acids: (-8 for K, E, D, R), (-4 for S, T, N, Q), (-2 for P, H), (+1 for A, G, Y, C, W), (+2 for L, V, I, F, M).

Following inputs were given for each of the group

(a) f_i = (counts of i^{th} group) / (total counts in the protein); i ranges from 1 to 5

(b) m_{ji} = j^{th} order moment of positions of amino acids in i^{th} group; j ranges from 2 to 5.

A total of 25 inputs representing the hydrophobic composition of a protein were fed to the Neural Network. The rationale for using moments was same as described in the section on charge composition inputs.

Taken together a total of 105 compositional properties of a given protein sequence were used to predict their adhesin characteristics.

The software PropSearch uses 144 compositional properties of protein sequences to detect possible structural or functional relationships between a new sequence and sequences in the database (Hobohm and Sander 1995). The approach defines protein sequence dissimilarity (or distance) as a weighted sum of differences of compositional properties such as singlet and doublet amino acid composition, molecular weight, isoelectric point (protein property search or PropSearch). Compositional properties of proteins have also been used for predicting secretory proteins in bacteria and apicoplast targeted proteins in Plasmodium falciparum (Zuegge, et al. 2001). The properties used here are statistical methods, principal component analysis, self-organizing maps, and supervised neural networks. In SPAAN, we have used 105 compositional properties in the five modules viz. Amino Acid frequencies, Multiplet frequencies, Dipeptide frequencies, Charge composition, Hydrophobic composition. The total of 105 properties used in SPAAN are 20 for Amino acid frequencies, 20 for Multiplets frequencies, 20 for Dipeptide frequencies (Top 20 significant dipeptides are used, based on *t-test*), 20 for Charge composition (frequency of charged amino acids (R, K, E and D) and moments of 2nd to 19th order), and 25 for Hydrophobic composition (Amino acids were classified into five groups (-8 for K, E, D, R), (-4 for S, T, N, Q), (-2 for P, H), (+1 for A, G, Y, C, W), (+2 for L, V, I, F, M). A total of 25 inputs consisted of the following: Frequency of each group, Moments of positions of amino acids in each group from 2nd to 5th order.

Neural Network

A feed forward error back propagation Neural Network was used. The program is a kind gift from Charles W. Anderson, Department of Computer Science, Colorado State University, Fort Collins, CO 80523, anderson@cs.colostate.edu

Neural Network architecture

The Neural Network used here has a multi-layer feed-forward topology. It consists of an input layer, one hidden layer and an output layer. This is a 'fully-connected' Neural Network where each neuron *i* is connected to each unit *j* of the next layer (Figure 1).

The weight of each connection is denoted by w_{ij}. The state I_i of each neuron in the input layer is assigned directly from the input data, whereas the states of hidden layer neurons are computed by the sigmoid function,

$h_j = 1 / (1 + \exp -(w_{j0} + \quad w_{ij} I_i))$,

where, w_{j0} is the bias weight

The back propagation algorithm was used to minimize the differences between the computed output and the desired output. Ten thousand cycles (epochs) of iterations are performed. Subsequently, the best epoch with minimum error was identified. At this point the network produces approximate target values for a given input in the training set.

A network was trained optimally for each attribute. Thus five networks were prepared. The schematic diagram (Figure 1) shows the procedure adopted. The number of neurons in the input layer was equal to the number of input data points for each attribute (for example 20 neurons for 20 numerical input vectors of the amino acid composition attribute). The optimal number of neurons in the hidden layer was determined through experimentation for minimizing the error at the best epoch for each network individually. An upper limit for the total number of weight connections was set to half of the total number of input vectors to avoid over fitting as suggested previously (Andrea *et al*).

Computer programs to compute individual compositional attributes were written in C and executed on a PC under Red Hat Linux ver 7.3 or 8.0. The network was trained on the training set, checks error and optimizes using the validate set through back propagation. The validate set was different from the training set. Since, the number of well annotated adhesins were not many, we used the 'validate set' itself as test set for preliminary evaluation of the performance and to obtain the fraction of correlation to compute the weighted average probability (P_{ad} value) described in the next section. The training set had 367 adhesins and 580 non-adhesins. The validate set had 102 adhesins and 123 non-adhesins. The adhesins were qualified with a digit '1' and the non-adhesins were qualified with a digit '0'.

During predictions, the network is fed with new data from the sequences that were not part of training set. Each network assigns a probability value of being an adhesin to a given sequence. The final probability is computed as described in the next section.

Probability of being an adhesin, the P_{ad} value

Query proteins are processed modularly through network trained for each attribute. Thus, five probability outputs are obtained. Final prediction was computed using the following expression which is a weighted linear sum of the probabilities from five modules:

$$P_{ad} = \frac{(P_A * fc_A + P_C * fc_C + P_D * fc_D + P_H * fc_H + P_M * fc_M)}{(fc_A + fc_C + fc_D + fc_H + fc_M)}$$

P_i = Probability from i module,

fc_i = fraction of correlation of i module of the trained Neural Network,

Where i = A (Amino acid frequencies), C (Charge composition), D (Dipeptide frequencies), H (Hydrophobic composition), or M (Multiplet frequencies).

The fraction of correlation fc_i represents the fraction of total entries that were correctly predicted ($P_{i,adhesin} > 0.5$ and $P_{i,\ non\text{-}adhesin} < 0.5$) by the trained network on the test set used in preliminary evaluation (Charles Anderson).

Neural Network

A feed forward error back propagation Neural Network was used. The program was downloaded from the web site with permission from the author, Charles W. Anderson, Department of Computer Science, Colorado State University, Fort Collins, CO 80523, anderson@cs.colostate.edu

Statistical Analysis

All statistical procedures were carried out using Microsoft Excel (Microsoft Corporation Inc. USA).

Sequence analysis

Homology analysis was carried out using CLUSTAL W (Thompson *et al* 1994), BLAST (Altschul *et al* 1990), CDD (conserved domain database) search (Marchler-Bauer *et al* 2002).

The whole genome sequences of microbial pathogens present new opportunities for the development of clinical applications such as diagnostics and vaccines. The present invention provides new leads for the development of candidate genes, and their encoded proteins in their functional relevance to preventive therapeutics.

The protein sequences of both the classes, i.e. adhesin and non-adhesin, were downloaded from the existing database (National Centre for Biotechnology Information (NCBI), USA). A total of 105 compositional properties under the five sequence

attributes namely, amino acid composition, multiplet composition, dipeptide composition, charge composition and hydrophobic composition were computed by computer programs written in C language. The attributes were computed for all the proteins in both the databases. The sequence-based attributes were then used to train Artificial Neural Network for each of the protein attributes. Adhesins were qualified by the digit '1' and non-adhesins were qualified by the digit '0'. Finally each trained Artificial Neural Network was used to identify potential adhesins which can be envisaged to be useful for the development of preventive therapeutics against pathogenic infections. Accordingly, the invention provides a computational method for identifying adhesin and adhesin-like proteins of therapeutic potential, which comprises:

1. preparing two comprehensive data-sets of adhesin and non-adhesin proteins from publicly available information on protein sequences,

2. calculating computationally the sequence based attributes of the protein sequences in the publicly available protein datasets using specially developed Software for Prediction of Adhesins and Adhesin-like proteins using Neural Networks (SPAAN),

3. training the Artificial Neural Network (ANN) for the selected attributes,

4. assigning probability value suitable for an adhesin, "P_{ad}" to the query protein and identifying adhesin like property in the query proteins with the help of trained Artificial Neural Network implemented in SPAAN,

5. validating computationally the protein sequences as therapeutic potentials by comparing with the known protein sequences that are biochemically characterized in the pathogen genome.

In an embodiment of the invention the protein sequence data may be taken from an organism, specifically but not limited to organisms such as *Escherichia coli*, *Haemophilus influenzae*, *Helicobacter pylori*, *Mycoplasma pneumoniae*, *Mycobacterium tuberculosis*, *Rickettsiae prowazekii*, *Porphyromonas gingivalis*, *Shigella flexneri*, *Streptococcus mutans*, *Streptococcus pneumoniae*, *Neisseria meningitides*, *Streptococcus pyogenes*, *Treponema pallidum*, Severe Acute Respiratory Syndrome associated coronavirus.

In another embodiment to the present invention different sequence-based attributes used for identification of proteins of therapeutic potential, comprise amino acid composition, charge composition, hydrophobicity composition, multiplets frequencies, and dipeptide frequencies.

In an embodiment, the non-homologous adhesin protein sequence may be compared with that of known sequences of therapeutic applications in the selected pathogens.

In an embodiment of the invention, the sequences of adhesin or adhesin like proteins comprise sequences of sequences IDs listed in Tables 5 and 6 identified by the method of invention.

Another embodiment of the invention the computer system comprises a central processing unit, executing SPAAN program, giving probabilities based on different attributes using Artificial Neural Network and in built other programs of assessing attributes, all stored in a memory device accessed by CPU, a display on which the central processing unit displays the screens of the above mentioned programs in response to user inputs; and a user interface device.

In One embodiment of the present invention, the particulars of the organisms such as their name, strain, accession number in NCBI database and other details are given in Table 2:

The invention is further explained with the help of the following examples, which are given by illustration and should be construed to limit the scope of the present invention in any manner.

Example 1

Operating SPAAN:

The purpose of the program is to computationally calculate various sequence-based attributes of the protein sequences.

The program works as follows:

The internet downloaded FASTA format files obtained from http://www.ncbi.nlm.nih.gov were saved by the name <organism_name>.faa are converted in the standard format by C program and passed as input to another set of C programs which computes the 5 different attributes of protein sequences (a total of 105 compositional properties in all 5 modules).

The computed properties were fed as input to the 5 different Neural Networks. Each trained network assigns a probability value of being an adhesin for a query protein. The final probability (P_{ad}) was calculated as weighted average of these five individual probabilities. The weights were determined from a correlation value of correct prediction during test runs of each of the five modules.

Input/Output format:

Downloaded Files and their format:

<organism_name>.faa: file which stores the annotation and the protein sequence.

Input file Format: FASTA

">gi.vertline."<annotation>

For example,

>gi.vertline.2314605.vertline.gb.vertline.AAD08472.vertline.histidine and glutamine-rich protein

MAHHEQQQQQQANSQHHHHHHHHAHHHHYYGGEHHHHNAQQHAEQQAEQQ

AQQQQQQQAHQQQQQKAQQQNQQY

>gi.vertline.3261822.vertline.gn1.vertline.PID.vertline.e328405 PE_PGRS

MIGDGANGGPGQPGGPGGLLYGNGGHGGAGAAGQDRGAGNSAGLIGNGGAG

GAGGNGGIGGAGAPGGLGGDGGKGGFADEFTGGFAQGGRGGFGGNGNTGAS

GGMGGAGGAGGAGGAGGLLIGDGGAGGAGGIGGAGGVGGGGGAGGTGGGG

VASAFGGGNAFGGRGGDGGDGGDGGTGGAGGARGAGGAGGAGGWLSGHSG

AHGAMGSGGEGGAGGGGGARGEAGAGGGTSTGTNPGKAGAPGTQGDSGDP

GPPG

>gi.vertline.. . .

Table 1: Output file format given by SPAAN

<organism_name>.out

SN	P_A	P_C	P_D	P_H	P_M	P_{ad}-value	Protein Name
1	0.05683	0.290803	0.441338	0.50304	0.029503	0.260485	>gi.vertline.32454344.vertline.gb.vertline.AAP82966.1. vertline.orf1a polyprotein [SARS coronavirus Hong Kong ZY-2003]
2	0.639235	0.166721	0.054583	0.935385	0.453498	0.462452	>gi.vertline.32454345.vertline.gb.vertline.AAP82967.1. vertline.orf1ab polyprotein [SARS coronavirus Hong Kong ZY-2003]
3	0.651111	0.911504	0.438696	0.543944	0.924044	0.690247	>gi.vertline.32454346.vertline.gb.vertline.AAP82968.1. vertline.spike glycoprotein [SARS coronavirus Hong Kong ZY-2003]

4	0.464324	0.655003	0.179503	0.008700	0.241573	0.300970	>gi.vertline.32454347.vertline.gb.vertline.AAP82969.1. vertline.Orf3a [SARS coronavirus Hong Kong ZY-2003]

Where P_A, P_C, P_D, P_H, P_M are the outputs of the five Neural Networks.

Example 2 organisms and sequence numbers

Table 2: Organism Name, Accession number, Number of base pairs, Date of release and Total number of proteins analyzed

Organism Name	Accession Number	Number of base pairs	Date of release	Total no. of proteins
E. coli O157 H7	NC_002695	5498450	7-Mar-2001	5361
H. influenzae Rd	NC_000907	1830138	30-Sep-1996	1709
H. pylori J99	NC_000921	1643831	10-Sep-2001	1491
M. pneumoniae	NC_000912	816394	2-Apr-2001	689
M. tuberculosis H37Rv	NC_000962	4411529	7-Sep-2001	3927
R. prowazekii strain Madrid E	NC_000963	1111523	10-Sep-2001	835
P. gingivalis W83	NC_002950	2343476	9-Sep-2003	1909
S. flexneri 2a str. 2457T	NC_004741	4599354	23- Apr-2003	4072
S. mutans UA159	NC_004350	2030921	25-Oct-2002	1960
S. pneumoniae R6	NC_003098	2038615	6-Sep-2001	2043
N. meningitidis serogroup A strain Z2491	NC_003116	2184406	27-Sep-2001	2065
S. pyogenes MGAS8232	NC_003485	1895017	Jan 31, 2002	1845
T. pallidum subsp. pallidum str. Nichols	NC_000919	1138011	7-Sep-2001	1036
Severe Acute Respiratory Syndrome (SARS) associated coronavirus Frankfurt 1	AY291315	29727	11-JUN-2003	14
SARS coronavirus HSR 1	AY323977	29751	15-OCT-2003	14

SARS coronavirus ZJ01	AY297028	29715	19-MAY-2003	3
SARS coronavirus TW1	AY291451	29729	14-MAY-2003	11
SARS coronavirus CUHK-Su10	AY282752	29736	07-MAY-2003	4
SARS coronavirus Urbani	AY278741	29727	12-AUG-2003	12
SARS coronavirus	NC_004718	29751	9-Sep-2003	29
SARS coronavirus Tor2	AY274119	29751	16-MAY-2003	15
SARS coronavirus GD01	AY278489	29757	18-AUG-2003	12
SARS coronavirus CUHK-W1	AY278554	29736	31-JUL-2003	11
SARS coronavirus BJ01	AY278488	29725	01-MAY-2003	11

Example 3

The multi-layered feed forward Neural Network architecture implemented in SPAAN (figure 1). A given protein sequence in FASTA format is first processed through the 5 modules A, C, D, H, and M to quantify the five types of compositional attributes. A: Amino acid composition, C: Charge composition, D: dipeptide composition of the 20 dipeptides (NG, RE, TN, NT, GT, TT, DE, ER, RR, RK, RI, AT, TS, IV, SG, GS, TG, GN, VI, HR), H: Hydrophobic composition, M: Amino acid frequencies as Multiplets. The sequence shown is part of the FimH precursor (gi 5524634) of *E. coli*. Subsequently, these numerical data are input to the input neuron layer. The directions of arrows show data flow. The number of neurons chosen in the input layer was equal to the number of the numerical input vectors of each module. The network was optimally trained through minimization of error of detection based on validate set through back propagation. The details are described in the methods. Each network module assigns a probability value of the protein being an adhesin based on the corresponding attribute. The final probability of a protein sequence being an adhesin is the P_{ad} value a weighted average of the individual probabilities and the associated fraction of correlation which is a measure of the strength of the prediction.

Example 4

Performance of SPAAN assessed using a test set of 37 adhesins and 37 non-adhesins that were not part of the training set. Matthew's correlation coefficient (Mcc, plotted on

Y-axis) for all the proteins with P_{ad} values above a given threshold (plotted on X-axis) (figure 2). The Matthew's correlation is defined as:

$$Mcc = \frac{(TP * TN) - (FP * FN)}{\sqrt{(TN + FN)(TN + FP)(TP + FN)(TP + FP)}}$$

Where TP = True Positives, TN = True Negatives, FP = False Positives, FN = False Negatives.

Here TPs are adhesins, TNs are non-adhesins. In general, adhesins have high P_{ad} value, whereas non-adhesins have low P_{ad} value. Thus known adhesins with P_{ad} value above a given threshold are true positives whereas known non-adhesins with P_{ad} value below the given threshold are true negatives. The sensitivity, Sn is given by $\left(\frac{TP}{TP + FN}\right)$ and specificity, Sp is given by $\left(\frac{TP}{TP + FP}\right)$. False negatives are those cases, wherein a known adhesin had P_{ad} value lower than the chosen threshold. Similarly, a known non-adhesin with a P_{ad} value higher than the chosen threshold was taken as false positive. A theoretical polynomial curve of second order (dashed line) was fitted to the observed curve (smooth line) with a Karl-Pearson correlation coefficient R^2 = 0.9799. The maximum point of the theoretical curve (where first derivative vanishes and second derivative is negative) was chosen as reference (vertical dotted line) to identify the maximum Mcc = 0.94 on the observed curve (shown by arrow). The corresponding P_{ad} value threshold was 0.51. At this P_{ad} value threshold, Sn and Sp were 0.89 and 1.0 respectively. Note that the Mcc does not drop down to the x-axis because the highest P_{ad} value attained by adhesins was 0.939 in comparison to the theoretical attainable limit of 1.0.

Example 5

Assessment of SPAAN on well known adhesins from various bacterial pathogens.

Table 3. Prediction of well characterized adhesins from various bacterial pathogens using SPAAN.

Species	Disease caused	Adhesin[a]	Host ligand	P_{ad} value[b] (Range)
E. coli	Diarrhoea	PapG (27)	α-D-gal(1-4) β-D-Gal-containing receptors	0.84-0.76
		SfaS (5)	alpha-sialyl-beta-2,3-b-galactose	0.94-0.94
		FimH (63)	D-mannosides	0.96-0.23[c]

		Intimin (12)	tyrosine-phosphorylated form of host cell receptor Hp90	0.95-0.78
		PrsG (5)	Gal(alpha1-4)Gal	0.86-0.85
Nontypeable *H. influenzae*	Influenza	HMW1, HMW2	Human epithelial cells	0.97
		Hia (8)	human conjuctival cells	0.93-0.90
H. influenzae	bacterial meningitis[d]	HifE (18)	Sialylyganglioside-GM1	0.85-0.73
K. pneumoniae	Pneumonia	MrkD	type V collagen	0.82
B. pertussis	Whooping cough	FHA	Sulphated sugars on cell-surface glycoconjugates	0.85
		Pertactin	Integrins	0.43
Y. enterocolitica	Enterocolitis	YadA (5)	β_1 integrins	0.88-0.79
S. mutans	Dental Caries	SpaP (2) PAc	Salivary glycoprotein Salivary glycoprotein	0.88, 0.87 0.88
Streptococcus gordonii	Oral cavity	SspA (2)	Salivary glycoprotein	0.85, 0.84
		CshA	Fibronectin	0.78
		CshB	Fibronectin	0.63
		ScaA	Co-aggregation	0.71
		SspB (2)	Salivary glycoprotein	0.85,0.84
Streptococcus sobrinus	Tooth decay	SpaA PAg (2)	Salivary glycoprotein Salivary glycoprotein	0.89 0.89, 0.73
Streptococcus pyogenes	Scarlet Fever	Protein F	Fibronectin	0.49
Streptococcus pneumoniae	Bacterial Pneumonia	PsaA (5)	Human nasopharyngeal cells	0.82-0.78
		CbpA[e] / SpsA / PbcA/ PspC	phosphorylcholine of the teichoic acid.	0.81-0.49
Streptococcus parasanguis	Valve endocarditis	FimA	Salivary glycoprotein fibrin	0.76
Streptococcus sanguis	Tooth Decay	SsaB	Salivary glycoprotein	0.71
Enterococcus faecalis	Empyma in patients with liver disease	EfaA	Unknown	0.83
Staphylococcus aureus	Food Poisoning	FnbA FnbB (3)	Fibronectin Fibronectin	0.8 0.78, 0.77, 0.69
Helicobacter pylori	Peptic Ulcers	BabA (17)	difucosylated Lewis[b] blood group antigen	0.87-0.68

[a]: The number of sequences from different strains and homologs from related species analyzed are shown in parantheses.

[b]: Rounded off to the second decimal.

[c]: Out of 63 FimH proteins, 54 were from *E. coli*, 6 from *Shigella flexineri*, 2 from *Salmonella enterica* and 1 was from *Salmonella typhimurium*. Except 2 FimH proteins, the rest had P_{ad} 0.51. The 2 exceptions (gi numbers: 5524636, 1778448) were from *E. coli*. The gi:5524636 protein is annotated as a FimH precursor but is much shorter (129 amino acids) than other members of the family. The gi:1778448 protein is a *S. typhimurium* homolog in *E. coli*.

[d]: Other ailments include pneumonia, epiglottitis, osteomyelitis, septic arthritis and sepsis in infants and older children.

[e]: The adhesin CbpA is also known by alternative names SpsA, PbcA and PspC. A total of seven sequences were analyzed. Except 1 PspC sequence, the rest all had P_{ad} 0.51.

Example 6

Ability of SPAAN to discriminate adhesins from non-adhesins at Pad 0.51 (figure 3-a).

Example 7

The non-homology character of SPAAN assesses in both adhesins and non-adhesins (figure 3b and 3c).

Figure 3 (a – c). SPAAN is non-homology based software. A total of 130 adhesins and 130 non-adhesins were analyzed to assess whether the predictive power of SPAAN could be influenced by the sequence relationships. (a) Histogram plots of the number of proteins in the various P_{ad} value ranges are shown. Shaded bars represent adhesins whereas open bars represent non-adhesins. Note the SPAAN's ability to segregate adhesins and non-adhesins into two distinct cohesive groups. (b) Pairwise sequence relationships among the adhesins were determined using CLUSTAL W and plotted on X-axis. Higher scores indicate similar pairs. The corresponding differences in P_{ad} values in the same protein pair was plotted on the Y-axis. Each point in the diagram represents a pair. Arrow points to protein pairs of the FimH family with high ΔP_{ad} values in spite of high similarity. Since one of the FimH proteins (gi: 5524636) had very low P_{ad} value all pairs with this false negative protein show high ΔP_{ad} values. The protein (gi: 5524636) is of much shorter length compared with other members of the same family. (c) plot for non-adhesins. Data are plotted in the 4 quadrant format for clear inspection. Note that among protein pairs with CLUSTAL W score < 20 the majority (82% in adhesins and 86% in non-adhesins) have $\Delta P_{ad} < 0.2$. These data support the non-homology character of SPAAN.

Example 8

Genomescan of pathogens by SPAAN identifies well known adhesins and new adhesins and adhesin-like proteins

Table 4. Analysis of predictions made by SPAAN on genome scans of a few selected pathogenic organisms[a]

Species / Protein Class	*Escherichia coli* O157:H7	*Mycobacterium tuberculosis* H37Rv	SARS associated corona virus (11 strains)
Total number of proteins with P_{ad} 0.51	575	435	5
Known adhesins	17[b]	-	-
Putative proteins with adhesin like characteristics	92[c]	105[j]	-
Hypothetical proteins with adhesin-like characteristics	22[d]	-	-
Proteins likely to be extracytoplasmic or located at surface	190[e]	191[k]	5[m]
Phage proteins	30[f]	-	-
Others	13[g]	6[l]	-
Hypothetical proteins	157[h]	86[h]	-
Wrong predictions	54[i]	47[i]	-

[a]: SPAAN has general applicability. The three pathogens chosen here are those in which intense investigations are being conducted presently. *M. tuberculosis* is of special importance to developing countries.

[b]: Fimbrial adhesins, AidA-I, gamma intimin, curlin, translocated intimin receptor, putative adhesin and transport, Iha, prepilin peptidase dependent protein C.

[c]: These proteins have been annotated as proteins with a putative function. These sequences were analyzed using CDD (Conserved domain database, NCBI) and BLAST searches. Adhesin like domains were found in these proteins.

[d]: These proteins have been annotated as 'hypothetical'. These sequences were analyzed using CDD and BLAST searches. Adhesin like domains were found in these proteins.

[e]: These proteins are outer membrane, extracellular, transport, surface, exported, flagellar, periplasmic lipoprotein, and proteins annotated as 'hypothetical' but found to have similar functions listed here using BLAST and CDD searches.

[f]: The phage proteins were of the following functional roles – tail fiber, head decoration, DNA injection, tail, major capsid, host specificity, endolysin.

[g]: Proteins predicted by SPAAN but not readily classifiable into the classes listed here have been collectively grouped as 'Others'. However, some of these proteins are known to participate in host-pathogen interactions. The annotated functional roles are typeIII secretion, antibiotic resistance, heat shock, acid shock, structural, tellurium resistance, terminase, Hcp-like, Sec-independent translocase, uncharacterized nucleoprotein, HicB-like.

[h]: These proteins have been annotated as hypothetical. Re-analyses of these proteins using BLAST and CDD failed identify any function for these proteins.

[i]: These proteins have been annotated with functional roles that are very likely to occur within the cell. Hence these proteins may have remote possibility of functioning as adhesins or adhesin-like proteins. Therefore this set of proteins have been incorrectly predicted as adhesins or adhesin-like by SPAAN.

[j]: These proteins are PE_PGRS, PE proteins. Several reports (for example Brennan *et al.*) indicate that PE_PGRS proteins may be localized to cell surface and aid in host-pathogen interaction.

[k]: Lipoproteins (lpp, lpq, lpr), PPE, outer membrane, surface, transport, secreted, periplasmic, extracellular, ESAT-6, peptidoglycan binding, exported, mpt (with extracellular domains), and proteins annotated as 'hypothetical' but found to have similar functions listed here using BLAST and CDD searches.

[l]: These proteins were of the following functions - glutaredoxin-like thioltransferase, putative involvement in molybdate uptake, ATP synthase chain, sulphotransferases, S.erythraea rhodanese-like protein M29612|SERCYSA_5, unknown function.

[m]: These proteins were the spike glycoprotein with antigenic properties, and nsp2, nsp5, nsp6 and nsp7.

Table 5: New putative adhesins predicted by SPAAN in the genomes listed in table 2 –

(Total number = 279)

Protein GI Number	Gene ID	Protein name
Escherichia coli O157:H7		
13360742	912619	hemagglutinin/hemolysin-related protein
13362986	914770	putative ATP-binding component of a transport system
13361114	913228	putative tail fiber protein
13364757	913676	minor fimbrial subunit/D-mannose specific adhesin
13362687	915687	putative fimbrial-like protein
13360856	912599	AidA-I adhesin-like protein
13364140	915374	putative fimbrial protein
13359793	914435	putative invasin
13364768	913650	putative invasin
13364034	915471	Gamma intimin
13362703	915668	putative DNA transfer protein precursor
13364141	915376	putative fimbrial protein
13359819	914463	AidA-I adhesin-like protein
13360480	917768	putative fimbrial-like protein
13362692	915681	putative fimbrial-like protein
13362585	916824	putative ATP-binding component of a transport system
13359881	914526	putative flagellin structural protein
13361579	917311	putative type 1 fimbrial protein precursor
13360880	913991	curlin major subunit CsgA
13364036	915465	translocated intimin receptor Tir
13360740	912615	putative major pilin protein
13361582	917317	putative ATP-binding component of a transport system and adhesin protein
13364754	913683	export and assembly outer membrane protein of type 1 fimbriae
13360484	917767	homolog of Salmonella FimH protein

13364751	913688	major type 1 subunit fimbrin
13359597	913742	putative fimbrial protein
13362550	916787	putative ATP-binding component of a transport system
13359595	913739	putative fimbrial protein
13359599	913748	probable outer membrane porin protein involved in fimbrial assembly
13363900	915704	putative fimbrial protein precursor
13361575	917307	putative fimbrial-like protein
13364756	913678	fimbrial morphology
13363496	916142	truncated putative fimbrial protein
13359601	913761	putative fimbrial-like protein
13364145	915368	putative type 1 fimbrial protein
13363902	915708	putative outer membrane usher protein precursor
13361576	917309	putative outer membrane protein
13361013	913353	putative major tail subunit
13364755	913682	fimbrial morphology
13360738	912793	putative outer membrane usher protein
13363928	915608	alpha-amylase
13363495	916144	putative outer membrane protein
13362383	916617	putative type-1 fimbrial protein
13364373	914972	outer membrane vitamin B12 receptor protein BtuB
13360879	912479	minor curlin subunit precursor CsgB
13360739	912756	putative chaperone protein
13361574	917314	putative fimbrial-like protein
13361127	913212	outer membrane protease precursor
13363210	916442	putative lipoprotein
13361104	913238	major tail protein
13361709	917446	putative major tail subunit
13359725	914366	outer membrane pore protein PhoE
13360875	913765	curli production assembly/transport component CsgF
13362170	913927	putative outer membrane protein
13361473	917203	putative BigB-like protein

13364025	915286	EspF protein
13360081	916982	outer membrane receptor for ferric enterobactin (enterochelin) and colicins B and D
13362977	914779	hypothetical lipoprotein
13360351	917632	outer membrane protein X
13360696	914208	putative outer membrane precursor
13361456	917206	putative outer membrane protein
13361626	917374	putative outer host membrane protein precursor
13361698	917449	putative outer membrane protein
13362186	913421	putative outer membrane protein precursor
13362697	915676	long-chain fatty acid transport protein FadL
13360918	914188	flagellar hook protein FlgE
13360737	912506	putative outer membrane protein
13360342	917629	putative outer membrane receptor for iron transport
13363396	916248	outer membrane channel TolC
13361958	912705	putative scaffolding protein in the formation of a murein-synthesizing holoenzyme
13359921	914566	nucleoside-specific channel-forming protein TSX
13360944	913890	outer membrane receptor for ferric iron uptake
13359998	914644	putative outer membrane transport protein
13363390	916251	putative ferrichrome iron receptor precursor
13364227	915153	outer membrane phospholipase A
13361982	912846	putative outer membrane protein
13360129	917032	a minor lipoprotein
13361817	912692	putative outer membrane protein
13360233	917507	membrane spanning protein TolA
13362837	915218	putative outer membrane lipoprotein
13362328	912985	putative colanic acid biosynthesis glycosyl transferase

Haemophilus influenzae Rd

16272254	949521	prepilin peptidase-dependent protein D
16272928	950762	immunoglobin A1 protease
16272129	951072	lipoprotein

16273251	950616	hemoglobin-binding protein
30995429	950130	opacity protein
16272854	949634	protective surface antigen D15
16272283	950648	opacity associated protein
16272604	949701	hemoglobin-binding protein
Helicobacter pylori J99		
4155101	889167	putative vacuolating cytotoxin (VacA) paralog
4154798	890022	putative vacuolating cytotoxin (VacA) paralog
4155426	890036	putative vacuolating cytotoxin (VacA) paralog
4155390	890075	vacuolating cytotoxin
4155400	890058	outer membrane protein - adhesin
4155681	889718	putative Outer membrane protein
4155420	890042	Outer membrane protein/porin
4155775	889799	outer membrane protein - adhesin
4155419	890044	Outer membrane protein/porin
4154526	889066	putative Outer membrane protein
4154724	889419	putative Outer membrane protein
4155862	890404	putative Outer membrane protein
4156048	889958	putative IRON(III) DICITRATE TRANSPORT PROTEIN
4154510	889297	putative Outer membrane protein
4155432	889515	putative outer membrane protein
4155623	889671	putative Outer membrane protein
4155700	889739	putative Outer membrane function
4154740	889426	Outer membrane protein/porin
4155692	889743	putative Outer membrane protein
4155594	889648	putative outer membrane protein
4155680	889719	putative Outer membrane protein
4155217	890243	putative Outer membrane protein
4155958	889905	putative Outer membrane protein
4155201	890259	putative Outer membrane protein
4155013	889232	cag island protein
4154974	889032	putative Outer membrane protein

4155214	890244	putative Outer membrane protein
4154973	889042	Outer membrane protein
4155344	890115	putative Outer membrane protein
4155099	889160	FLAGELLIN A
4155023	888978	cag island protein
4155035	889201	cag island protein, CYTOTOXICITY ASSOCIATED IMMUNODOMINANT ANTIGEN
4155289	890164	NEURAMINYLLACTOSE-BINDING HEMAGGLUTININ PRECURSOR
Mycoplasma pneumoniae		
13507881	877207	involved in cytadherence
13507880	877268	ADP1_MYCPN adhesin P1
13508228	877211	species specific lipoprotein
13508181	877124	species specific lipoprotein
13508179	877071	Mollicute specific lipoprotein, MG307 homolog, from M. genitalium
13508178	877118	Mollicute specific lipoprotein, MG307 homolog, from M. genitalium,
13508176	876797	Mollicute specific lipoprotein, MG307 homolog, from M. genitalium
13508175	876848	Mollicute specific lipoprotein, MG307 homolog, from M. genitalium
13508106	876953	involved in cytadherence
13508350	877112	similar to phosphate binding protein Psts
Mycobacterium tuberculosis H37 Rv		
15607496	886491	PPE
15607445	886592	PPE
15610644	888270	PE_PGRS
15608588	886605	PE_PGRS
15609627	887941	PE_PGRS
15610643	888256	PE_PGRS
15607718	887725	PE_PGRS

15609054	885362	PPE
15610486	888113	PPE
15610483	888120	PPE
15610479	888033	PPE
15609771	888573	PE_PGRS
15610648	888306	PE_PGRS
15610481	888114	PE_PGRS
15608117	885264	PE_PGRS
15607973	885391	PE_PGRS
15608231	885258	PE_PGRS
15608906	885429	PE_PGRS
15608891	885544	PPE
15609990	888171	PE_PGRS
15609055	885506	PPE
15608227	887094	PE_PGRS
15610524	888151	PE_PGRS
15609490	886003	PPE
15607886	888664	PE_PGRS
15609624	887909	PE_PGRS
15607420	886621	PE_PGRS
15608897	885325	PE_PGRS(wag22)
15608590	886595	PE_PGRS
15609728	887992	PE_PGRS
15608012	885742	PE_PGRS
15608534	886745	PE_PGRS
15608940	885730	PE_PGRS
15607887	888662	PE_PGRS
15609235	888312	PE_PGRS
15610694	887822	PPE
15609533	885517	PE_PGRS
15610480		PE_PGRS

Rickettsia prowazekii strain Madrid E

15604316	883411	CELL SURFACE ANTIGEN (sca3)
15604546	883694	CELL SURFACE ANTIGEN (sca5)

Porphyromonas gingivalis W83

34541453	2551934	hemagglutinin protein HagA
34540040	2551409	lipoprotein, putative
34540364	2552375	extracellular protease, putative
34541613	2552074	hemagglutinin protein HagE
34540183	2551891	internalin-related protein

Shigella flexneri 2a str. 2457T

30065424	1080663	minor fimbrial subunit, D-mannose specific adhesin
30062726	1077662	putative adhesion and penetration protein
30063758	1078834	putative fimbrial-like protein
30065431	1080671	major type 1 subunit fimbrin (pilin)
30063366	1078379	flagellar protein FliD
30064308	1079668	outer membrane fluffing protein
30062613	1077555	flagellar hook protein FlgE
30061954	1076843	conserved hypothetical lipoprotein
30065173	1080393	putative lipase
30065425	1080664	minor fimbrial subunit, precursor polypeptide
30064485	1079637	putative fimbrial protein
30062615	1077558	flagellar basal body L-ring protein FlgH
30064307	1079452	outer membrane fluffing protein
30065601	1080859	putative glycoprotein/receptor
30062118	1077025	putative fimbrial-like protein
30064099	1079223	lipoprotein
30062616	1077559	flagellar basal body P-ring protein FlgI
30063546	1078596	putative fimbrial-like protein
30062940	1077910	putative outer membrane protein
30065426	1080665	minor fimbrial subunit, precursor polypeptide
30062779	1077721	putative outer membrane protein
30064194	1079329	putative lipoprotein
30063365	1078378	flagellin

30062298	1077222	outer membrane protein X
30064968	1080175	putative major fimbrial subunit
30061858	1076740	outer membrane pore protein E (E,Ic,NmpAB)
30062178	1080410	minor lipoprotein
30062479	1077412	putative fimbrial-like protein
30062565	1077506	minor curlin subunit precursor
30063880	1078972	putative outer membrane lipoprotein
30064531	1079686	cytoplasmic membrane protein
30065033	1080243	putative receptor protein
Streptococcus mutans UA159		
24378550	1029610	putative secreted antigen GbpB/SagA; putative peptidoglycan hydrolase
24379087	1028055	cell surface antigen SpaP
24380463	1029310	putative membrane protein
24379075	1028046	penicillin-binding protein 2b
24378955	1027967	penicillin-binding protein 1a; membrane carboxypeptidase
24379801	1028662	glucan-binding protein C, GbpC
24379528	1029536	hypothetical protein; possible cell wall protein, WapE
24379231	1028158	putative glucan-binding protein D; BglB-like protein
24380488	1029325	conserved hypothetical protein; possible transmembrane protein
24380291	1029139	putative amino acid binding protein
24379342	1028247	putative penicillin-binding protein, class C; fmt-like protein
24380047	1028904	putative ABC transporter, branched chain amino acid-binding protein
24378698	1029755	putative ABC transporter, metal binding lipoprotein; surface adhesin precursor; saliva-binding protein; lipoprotein receptor LraI (LraI family)
24378708	1029768	putative transfer protein
24379427	1028331	cell wall-associated protein precursor WapA
24379272	1028196	putative amino acid transporter, amino acid-binding protein
24379641	1028511	putative ABC transporter, amino acid binding protein

Streptococcus pneumoniae R6

15902395	934801	Choline-binding protein
15902381	934810	Choline-binding protein F
15902165	932894	Surface protein pspA precursor
15904047	934859	Choline binding protein D
15904036	933487	Choline binding protein A
15903986	933069	Choline-binding protein
15903796	933669	Autolysin (N-acetylmuramoyl-L-alanine amidase)

Neisseria meningitidis Z2491

15794121	907145	putative membrane protein
15794144	907168	putative surface fibril protein
15793284	906275	truncated pilin
15793460	906456	IgA-specific serine endopeptidase
15793282	906273	fimbrial protein precursor (pilin)
15793337	906332	adhesin
15793253	906243	putative lipoprotein
15794356	907848	putative lipoprotein
15793684	906699	putative membrane protein
15793290	906281	truncated pilin
15793283	906274	truncated pilin
15793475	906471	haemoglobin-haptoglobin-utilization protein
15793406	906401	porin, major outer membrane protein P.I
15794985	907333	adhesin MafA2
15794344	907836	putative lipoprotein
15794622	908118	hypothetical outer membrane protein
15793599	906604	pilus-associated protein
15793763	906779	putative periplasmic binding protein

Streptococcus pyogenes MGAS8232

19745214	995235	putative secreted protein
19746570	994224	putative penicillin-binding protein 1a
19745593	994771	putative 42 kDa protein
19745813	993958	putative adhesion protein

19745225	994839	putative choline binding protein
19745828	995250	streptolysin S associated protein
19746229	995021	putative minor tail protein
19746909	994105	putative laminin adhesion
19745560	995061	putative cell envelope proteinase
Treponema pallidum subsp. pallidum str. Nichols		
15639714	2611034	flagellar hook protein (flgE)
15639609	2611657	tpr protein J (tprJ)
15639111	2610909	tpr protein C (tprC)
15639125	2610968	tpr protein D (tprD)
SARS coronavirus		
31581505		spike protein S [SARS coronavirus Frankfurt 1]
32187357		spike protein S [SARS coronavirus HSR 1]
32187342		spike glycoprotein [SARS coronavirus ZJ01]
30698329		putative spike glycoprotein S [SARS coronavirus TW1]
30421454		putative spike glycoprotein [SARS coronavirus CUHK-Su10]
30027620		S protein [SARS coronavirus Urbani]
29836496	1489668	E2 glycoprotein precursor; putative spike glycoprotein [SARS coronavirus]
30795145		spike glycoprotein [SARS coronavirus Tor2]
31416295		spike glycoprotein S [SARS coronavirus GD01]
30023954		putative E2 glycoprotein precursor [SARS coronavirus CUHK-W1]
30275669		spike glycoprotein S [SARS coronavirus BJ01]
29837498		3C-like proteinase nsp5-pp1a/pp1ab (3CL-PRO) [SARS coronavirus]
29837501		putative nsp8-pp1a/pp1ab [SARS coronavirus]
29837503		putative nsp10-pp1a/pp1ab; formerly known as growth-factor-like protein [SARS coronavirus]
29837502		putative nsp9-pp1a/pp1ab [SARS coronavirus]

Table 6: Hypothetical proteins predicted as putative adhesins by SPAAN in the genomes listed in table 2 –

(Total number of proteins = 105)

Protein GI number	Gene ID
Escherichia coli O157:H7	
13363955	915578
13360000	914929
13362244	912369
13359999	914888
13361583	917316
13361172	913156
13361131	913207
13359780	914422
13360571	912499
13362197	912893
13362260	912399
13360947	913505
13361464	917196
13361635	917367
13362421	916655
13361463	917195
Haemophilus influenzae Rd	
16272115	951058
30995442	950581
Helicobacter pylori J99	
4155526	889586
4155712	889748
4155632	889684
4156035	889468
4155499	

Mycoplasma pneumoniae

13507870	877230
13508239	877245
13508109	876868
13508025	877084
13507838	876784
13507883	877183
13507871	877239
13507944	877056
13508241	876750
13507942	877055
13507840	877387
13507867	877242
13508201	877044
13507941	876985
13508114	877397

Mycobacterium tuberculosis H37Rv

15611014	886198
15610173	887320
15609513	885515
15608094	885411
15610958	886155
15607528	886436
15607678	887473
15609587	885760
15610708	887227
15609526	885246
15611033	886225
15609028	885094
15607730	887771
15609121	885813
15608255	885951

15608409 887039

15609124 885815

15607734 887797

Rickettsia prowazekii strain Madrid E

15604649 883964

15604322 883472

15604659 883996

15604417 883217

Porphyromonas gingivalis W83

34540233 2551594

Shigella flexneri 2a str. 2457T

30062687 1077638

30062956 1080449

30063681 1078754

30065435 1080675

30063891 1078983

30063211 1078195

30065233 1080463

30064387 1079531

30062638 1077590

30065236 1080466

30061839 1076721

Streptococcus mutans UA159

24378864 1029452

24380475 1029319

24380237 1029088

24379203 1028139

24380480 1029320

24379275 1029489

24379291 1028216

24379295 1028215

24379804 1028663

24379162 1029417

24378987 1029363

24379179 1028118

24379166 1028107

24378827 1029444

24380216 1029067

Streptococcus pneumoniae R6

15902140 932867

15903446 934616

15903916 934001

15903848 933609

15902832 934332

15902372 934804

15902152 932889

Neisseria meningitidis Z2491

15793668 906680

15794714 907603

Streptococcus pyogenes MGAS8232

19747011 993608

19747024 994165

19747012 994373

19746396 995057

19746651 993824

19745883 995045

19745912 994077

Treponema pallidum subsp. pallidum str. Nichols

15639844 2611061

15639720 2611059

Table 7: The list of 198 adhesins found in bacteria

PapG (E. coli)

12837502

7407201

7407207
7407205
147096
4240529
7407203
42308
7443327
78746
18265934
26111419
26250987
26109826
26249418
13506767
42301
78745
129622
147092
13506906
7407209
147080
281926
7407199
147100
78744

SfaS (E.coli)

477910
264035
42959
134449
96425

FimH (E.coli)

26251208

26111640

5524634

29422425

5524630

29422435

29422415

10946257

29422419

11120564

29422457

11120562

29422459

5524632

29422455

29422453

29422451

29422449

29422447

29422445

29422443

29422437

29422433

29422431

29422429

29422427

29422423

29422421

29422417

729494

1361011

1790775

3599571

29422441

12620398

29422439

5524628

1787779

1742472

1742463

15801636

25321294

12515169

11120566

24051859

24112911

13360484

15800801

15830279

25392018

25500156

12514120

1787173

16128908

16501811

16759519

24051219

24112354

30040724

30062478

6650093

5524636

1778448

Intimin (E.coli)

17384659

4388530

1389879

15723931

4323336

4323338

4323340

4323342

4323344

4323346

4323348

4689314

PrsG (E.coli)

42523

42529

7443328

7443329

1172645

HMW1 (Nontypeable *H. influenzae)*

282097

HMW2 (Nontypeable H. influenzae)

5929966

Hia (Nontypeable H. influenzae)

25359682

25359489

25359709

25359628

25359414

25359389

21536216

25359445

HifE (H. influenzae)

13506868
13506870
13506872
13506874
13506876
3688787
3688790
3688793
2126301
1170264
1170265
533127
535169
3025668
3025670
3025672
3025674
642038

MrkD (K. pneumoniae)

127307

FHA (B. pertussis)

17154501

Pertactin (B. pertussis)

33571840

YadA (Y. enterocolitica)

10955604
4324391
28372996
23630568
32470319

SpaP (S. mutans)

26007028

47267

PAc (S. mutans)

129552

SspA (Streptococcus gordonii)

25990270

1100971

CshA (Streptococcus gordonii)

457707

CshB (Streptococcus gordonii)

18389220

ScaA (Streptococcus gordonii)

310633

SspB (Streptococcus gordonii)

25055226

3220006

SpaA (Streptococcus sobrinus)

546643

PAg (Streptococcus sobrinus)

217036

47561

Protein F (Streptococcus pyogenes)

19224134

PsaA (Streptococcus pneumoniae)

18252614

7920456

7920458

7920460

7920462

CbpA[e] / SpsA / PbcA/ PspC

(Streptococcus pneumoniae)

14718654

2425109

2576331

2576333

3153898

9845483

19548141

FimA (Streptococcus parasanguis)

97883

SsaB (Streptococcus sanguis)

97882

EfaA (Enterococcus faecalis)

493017

FnbA (Staphylococcus aureus)

120457

FnbB (Staphylococcus aureus)

581562

21205592

13702452

BabA (Helicobacter pylori)

13309962

13309964

13309966

13309968

13309970

13309972

13309974

13309976

13309978

13309980

13309982

13309984

13309986

13309988

13309990

13309992

13309994

Advantages:

1. The method helps in discovering putative adhesins, which are of great importance in drug discoveries and preventive therapeutics.
2. The method is useful in predicting the adhesive nature of even unique proteins, because it is independent of the homology of the query proteins with other proteins.
3. This method is easy to use. For calculating the output, only the amino acid sequence is required as input. No other information is required to get the information about its adhesive nature.

REFERENCES MAY BE MADE TO

1. Andrea, T.A., Kalayeh, H.(1991) Applications of neural networks in quantitative structure-activity relationships of dihydrofolate reductase inhibitors. J. Med. Chem. 34, 2824-2836.

2. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ.(1990) Basic local alignment search tool. J Mol Biol. 215(3), 403-410.

3. Bassinet L, Gueirard P, Maitre B, Housset B, Golnon P, Guiso N.(2000) Role of adhesins and toxins in invasion of human tracheal epithelial cells by Bordetella pertussis. Infect Immun. 68(4), 1934-1941.

4. Bock, K., *et al.*(1985). Specificity of binding of a strain of uropathogenic Escherichia coli to Gal alpha 1----4Gal-containing glycosphingolipids. J. Biol. Chem. 260, 8545-8551.

5. Brendel, V., Bucher, P., Nourbakhsh, I.R., Edwin Blaisdell, B., and Karlin, S.(1992) Methods and algorithms for statistical analysis of protein sequences. Proc. Natl. Acad. Sci. USA 89, 2002-2006.

6. Brennan, M.J., Delogu, G., Chen, Y., Bardarov, S., Kriakov, J., Alavi, M., Jacobs, W.R., (2001).

7. Evidence that Mycobacterial PE_PGRS proteins are cell surface constituents that influence interactions with other cells. Infect. Immun, 69, 7326-7333.

8. De BK, Woolfitt AR, Barr JR, Daneshvar MI, Sampson JS, Ades EW, Carlone GM. (2003) Analysis of recombinant acylated pneumococcal surface adhesin A of Streptococcus pneumoniae by mass spectrometry. Arch Biochem Biophys. 15, 419(2), 147-157.

9. Egland PG, Du LD, Kolenbrander PE (2001) Identification of independent Streptococcus gordonii SspA and SspB functions in coaggregation with Actinomyces naeslundii. Infect Immun. 69(12), 7512-7516

10. Finlay, B.B. and Falkow, S.(1997) Common themes in microbial pathogenicity revisited. Microbiol. Mol. Biol. Rev. 61, 136–169,

11. Fraser, C.M., Eisen, J., Fleischmann, R.D., Ketchum, K.A., Peterson, S.(2000) Comparative genomics and understanding of microbial biology. Emerg. Infect. Dis. 6,505-6512

12. Halperin, S. A., Scheifele, D., Mills, E., Guasparini, R., Humphreys, G., Barreto, L., Smith, B.(2003) Nature, evolution, and appraisal of adverse events

and antibody response associated with the fifth consecutive dose of a five-component acellular pertussis-based combination vaccine. Vaccine 21, 2298–2306.

13. Hartford O, McDevitt D, Foster TJ.(1999) Matrix-binding proteins of Staphylococcus aureus: functional analysis of mutant and hybrid molecules. Microbiology. 145 (Pt 9), 2497-2505.

14. Hobohm, U. and Sander, C.(1995) A sequence property approach to searching protein databases. J. Mol. Biol. 251, 390-399.

15. Ide T, Michgehl S, Knappstein S, Heusipp G, Schmidt MA.(2003) Differential modulation by Ca2+ of type III secretion of diffusely adhering enteropathogenic Escherichia coli. Infect Immun. 71(4), 1725-1732.

16. Langermann S *et al.*(2000) Vaccination with FimH adhesin protects cynomolgus monkeys from colonization and infection by uropathogenic Escherichia coli. J. Infect. Dis. 181, 774-778.

17. Lowe A.M., Lambert, P.A., Smith, A.W.(1995) Cloning of an Enterococcus faecalis endocarditis antigen: homology with adhesins from some oral streptococci. Infect Immun. 63, 703-706.

18. Maurer, L., Orndorff, P.(1987). Identification and characterization of genes determining receptor binding and pilus length of Escherichia coli type 1 pili. J. Bacteriol. 169, 640-645

19. Marchler-Bauer A, Panchenko AR, Shoemaker BA, Thiessen PA, Geer LY, Bryant SH. (2002) CDD: a database of conserved domain alignments with links to domain three-dimensional structure. Nucleic Acids Res. 1, 30(1), 281-283.

20. Neubauer H, Hensel A, Aleksic S, Meyer H.(2000) Evaluation of a Yersinia adhesion gene (yadA) specific PCR for the identification of enteropathogenic Yersinia enterocolitica. Int J Food Microbiol. 15, 57(3), 225-227.

21. Nishikawa, K., Kubota, Y. and Ooi, T.(1983) Classification of proteins into groups based on amino acid composition and other characters. II. grouping into four types. J. Biochem. 94, 997-1007.

22. Peregrin-Alvarez, J.M., Tsoka, S., Ouzounis, C.A.(2003) The phylogenetic extent of metabolic enzymes and pathways. Genome Res. 13, 422-427.

23. Prinz, C., Hafsi, N. Voland, P.(2003) Helicobacter pylori virulence factors and the host immune response: implications for therapeutic vaccination. Trends in Microbiol. 11, 134-138.

24. Rapola, S., Jäntti, V., Eerola, M., Helena Mäkelä, P., Käyhty, H., Kilpi, T.(2003) Anti-PsaA and the risk of pneumococcal AOM and carriage. Vaccine 21, 3608–3613.

25. Rison. S.C., Teichmann, S.A., Thornton, J.M.(2002) Homology, pathway distance and chromosomal localization of the small molecule metabolism enzymes in Escherichia coli. J. Mol. Biol. 318, 911-932

26. Sperandio V, Bailey C, Giron JA, DiRita VJ, Silveira WD, Vettore AL, Kaper JB.(1996) Cloning and characterization of the gene encoding the OmpU outer membrane protein of Vibrio cholerae. Infect Immun. 64(12), 5406-5409.

27. St Geme JW 3rd, Cutter D.(2000) The Haemophilus influenzae Hia adhesin is an autotransporter protein that remains uncleaved at the C terminus and fully cell associated. J Bacteriol. 182(21), 6005-6013.

28. Thompson, J.D., Higgins, D.G., Gibson, T.J.(1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. Nucleic Acids Res. 22, 4673-4680

29. Van Schilfgaarde M, van Ulsen P, Eijk P, Brand M, Stam M, Kouame J, van Alphen L, Dankert J.(2000) Characterization of adherence of nontypeable Haemophilus influenzae to human epithelial cells. Infect Immun. 68(8), 4658-4665.

30. Wizemann, T.M., Adamou, J.E., Langermann, S.(1999). Adhesins as targets for vaccine development. Emerg. Infect. Dis. 5, 395-403,

31. Wolf, Y.I., Rogozin, I.B., Kondrashov, A.S., and Koonin, E.V.(2001) Genome alignment, evolution of prokaryotic genome organization and prediction of gene function using genomic context. Genome Res. 11, 356-372

32. Yu J, Leung WK, Go MY, Chan MC, To KF, Ng EK, Chan FK, Ling TK, Chung SC, Sung JJ.(2002) Relationship between Helicobacter pylori babA2 status with gastric epithelial cell turnover and premalignant gastric lesions. Gut. 51(4), 480-484.

33. Zuegge, J., Ralph, S., Schmuker, M., McFadden, G.I., Schneider, G.(2001) Deciphering apicoplast targeting signals--feature extraction from nuclear-encoded precursors of Plasmodium falciparum apicoplast proteins. Gene 280, 19-26.

5

Claims

1. A computational method for identifying adhesin and adhesin-like proteins, said method comprising steps of:
   a. computing the sequence-based attributes of protein sequences using five attribute modules of a neural network software, wherein the attributes are, (i) amino acid frequencies, (ii) multiplet frequency, (iii) dipeptide frequencies, (iv) charge composition, and (v) hydrophobic composition,
   b. training a artificial neural Network (ANN) for each of the computed five attributes, and
   c. identifying the adhesin and adhesin-like proteins having probability of being an adhesin (P_{ad}) as ≥ 0.51.
2. A method as claimed in claim 1, wherein the protein sequences are obtained from pathogens, eukaryotes, and multicellular organisms.
3. A method as claimed in claim 1, wherein the protein sequences are obtained from the pathogens selected from a group of organisms comprising *Escherichia coli*, *Haemophilus influenzae*, *Helicobacter pylori*, *Mycoplasma pneumoniae*, *Mycobacterium tuberculosis*, *Rickettsiae prowazekii*, *Porphyromonas gingivalis*, *Shigella flexneri*, *Streptococcus mutans*, *Streptococcus pneumoniae*, *Neisseria meningitides*, *Streptococcus pyogenes*, *Treponema pallidum* and Severe Acute Respiratory Syndrome associated human coronavirus (SARS).
4. A method as claimed in claim 1, wherein the method is a non-homology method.
5. A method as claimed in claim 1, wherein the method uses 105 compositional properties of the sequences.
6. A method as claimed in claim 1, wherein the method shows sensitivity of at least 90%.
7. A method as claimed in claim 1, wherein the method shows specificity of 100%.
8. A method as claimed in claim 1, wherein the method helps identifies adhesins from distantly related organisms.
9. A method as claimed in claim 1, wherein the neural network has multi-layer feed forward topology, consisting of an input layer, one hidden layer, and an output layer.

10. A method as claimed in claim 9, wherein the number of neurons in the input layer are equal to the number of input data points for each attribute.

11. A method as claimed in claim 1, wherein the "P_{ad}" is a weighted linear sum of the probabilities from five computed attributes.

12. A method as claimed in claim 1, wherein each trained network assigns a probability value of being an adhesin for the protein sequence.

13. A computer system for performing the method of claim 1, said system comprising a central processing unit, executing SPAAN program, giving probabilities based on different attributes using Artificial Neural Network and in built other programs of assessing attributes, all stored in a memory device accessed by CPU, a display on which the central processing unit displays the screens of the above mentioned programs in response to user inputs; and a user interface device.

14. A set of 274 annotated genes encoding adhesin and adhesin-like proteins, having SEQ ID Nos. 385 to 658.

15. A set of 105 hypothetical genes encoding adhesin and adhesin-like proteins, having SEQ ID Nos. 659 to 763.

16. A set of 279 annotated adhesin and adhesin-like proteins of SEQ ID Nos. 1 to 279.

17. A set of 105 hypothetical adhesin and adhesin-like proteins of SEQ ID Nos. 280 to 384.

18. A fully connected multilayer feed forward Artificial Neural Network based on the computational method as claimed in claim 1, comprising of an input layer, a hidden layer and an output layer which are connected in the said sequence, wherein each neuron is a binary digit number and is connected to each neuron of the subsequent layer for identifying adhesin or adhesin like proteins, wherein the program steps comprise:-

[a] feeding a protein sequence in FASTA format;

[b] processing the sequence obtained in step [a] through the 5 modules named A, C, D, H and M, wherein attribute A represents an amino acid composition, attribute C represents a charge composition, attribute D represents a dipeptide composition of the 20 dipeptides [NG, RE, TN, NT, GT, TT, DE, ER, RR, RK, RI, AT, TS, IV, SG, GS, TG, GN, VI

and HR], attribute H represents a hydrophobic composition and attribute M represents amino acid frequencies in multiplets to quantify 5 types of compositional attributes of the said protein sequence to obtain numerical input vectors respectively for each of the said attributes wherein the sum of numerical input vectors is 105;

[c] processing of the numerical input vectors obtained in step [b] by the input neuron layer to obtain signals, wherein the number of neurons is equal to the number of numerical input vectors for each attribute;

[d] processing of signals obtained from step [c] by the hidden layer to obtain synaptic weighted signals, wherein the optimal number of neurons in the hidden layer was determined through experimentation for minimizing the error at the best epoch for each network individually;

[e] delivering synaptic weighted signals obtained from step [d] to the output layer for assigning of a probability value for each protein sequence fed in step [a] as being an adhesin by each network module; and

[f] using the individual probabilities obtained from step [e] for computing the final probability of a protein sequence being an adhesin denoted by the P_{ad} value, which is a weighted average of the individual probabilities obtained from step [e] and the associated fraction of correlation which is a measure of the strength of the prediction.

19. A network as claimed in claim 18, wherein the input neuron layer consists of a total of 105 neurons corresponding to 105 compositional properties.

20. A network as claimed in claim 18, wherein the hidden layer comprises of neurons represented as 30 for amino acid frequencies, 28 for multiplet frequencies, 28 for dipeptide frequencies, 30 for charge composition and 30 for hydrophobic composition.

21. A network as claimed in claim 18, wherein the output layer comprises of neurons to deliver the output values as probability value for each protein sequence.