10. A method of making an oral care composition comprising a film entrained in a carrier wherein the film comprises a nonionic surfactant in an amount from 5% by weight of the film to 20% by weight of the film, the method comprising:

(a) providing the carrier;

(b) adding lamellar fragments of the film comprising the nonionic surfactant in an amount from 5% by weight of the film to 20% by weight of the film; and

(c) homogenizing the mixture.

11. The method of claim 10, wherein the nonionic surfactant is polysorbate 80.

12. The method of claim 10, wherein the film further comprises a therapeutic active.

13. The method of claim 12, wherein the therapeutic active is present in the film in an amount of from 20% by weight of the film to 65% by weight of the film.

14. The method of claim 10, wherein the film is present in an amount of from 0.1% to 5.0% by weight of the total weight of the composition.

15. Use of an oral care composition comprising a film entrained in a carrier, wherein the film comprises a nonionic surfactant in an amount from 5% by weight of the film to 20% by weight of the film, in the manufacture of a medicament for treatment of an oral condition, wherein said medicament is to be topically applied to a human or animal subject in need thereof.

16. The use of claim 15, wherein the film is to be disrupted after the topical application.

17. The use of claim 15, wherein the nonionic surfactant is polysorbate 80.

18. The use of claim 15, wherein the film further comprises a therapeutic active.

19. The use of claim 18, wherein the therapeutic active is present in the film in an amount of from 20% by weight of the film to 65% by weight of the film.

20. The use of any one of claims 15 to 19, wherein the oral condition is selected from at least one of caries, gingivitis or periodontitis.