

NOVEL HETEROCYCLIC COMPOUNDS

FIELD OF INVENTION

The present invention relates to novel heterocyclic compounds of the general formula (I) their pharmaceutically acceptable salts, pharmaceutically acceptable
5 solvates, enantiomers, diastereomers, and polymorphs. The invention also relates to processes for the preparation of the compounds of invention, pharmaceutical compositions containing the compounds and their use as selective Bruton's Tyrosine Kinase (BTK) inhibitors.

10 BACKGROUND OF THE INVENTION

Bruton's Tyrosine Kinase (BTK) is a member of the Tec family kinases with a well-characterized role in B-cell receptor (BCR)-signaling and B-cell activation (Schwartzberg, P.L., Finkelstein, L.D and Readinger, J.A., Nat. Rev. Immunol., 2005, 5, 284-295). BTK is a key signaling enzyme expressed in all hematopoietic cells
15 types, except T lymphocytes and natural killer cells. BTK is activated by the upstream Src-family kinases Blk, Lyn and Fyn and it lead to downstream activation of essential cell survival pathways such as NF- κ B and MAPK (Afar, D.E., Park, H., Howell, B. W., Rawlings, D. J., Cooper, J and Witte, O. N., Mol. Cell Biol., 1996, 16(7), 3465-3471). In turn, BTK phosphorylates and activates phospholipase-C $\otimes$
20 (PLC $\otimes$), leading to Ca²⁺ mobilization and activation of NF- κ B and MAP kinase pathways (Bajpai, U.D., Zhang, K., Teutsch, M., Sen, R and Wortis, H.H., J. Exp. Med., 2000, 191, 1735-1744).

BTK is intimately involved in multiple signal-transduction pathways regulating survival, activation, proliferation and differentiation of B-lineage lymphoid
25 cells. BTK is an upstream activator of multiple antiapoptotic signaling molecules and networks, including the signal transducer and activator of transcription 5 (STAT5) protein, phosphatidylinositol (PI) 3-kinase/AKT/mammalian target of rapamycin (mTOR) pathway, and nuclear factor kappa B (NF- κ B) (Mahajan, S., Vassilev, A.,

Sun, N., Ozer, Z., Mao, C and Uckun, F.M., J. Biol. Chem., 2001, 276, 31216-31228; Ortolano, S., Hwang, I.Y., Han, S.B and Kehrl, J.H., Eur. J. Immunol., 2006, 36, 1285-1295). This downstream signal transduction protein is a critical effector molecule that governs normal B-cell development, differentiation and functioning
5 and has also been implicated in initiation, survival and progression of mature B-cell lymphoproliferative disorders (Kuppers, R., Nat. Rev. Cancer, 2005, 5(4), 251-262).

In B-cells, BTK is important for B-cell antigen receptor-, CD40- and Toll-like receptor 4-mediated activation and proliferation. Furthermore, BTK plays a role in B-cell antigen processing and presentation (Satterthwaite, A.B., Witte, O.N., Immunol.
10 Rev., 2000, 175, 120-127). It is noteworthy that BTK is also essential in Fc γ receptor-mediated inflammatory cytokine production [tumor necrosis factor α , interleukin (IL)-1 α and IL-6] in monocytes/macrophages and therefore can contribute to immune complex-induced disease. BTK is abundantly expressed in malignant cells from patients with B-cell precursor (BCP)-acute lymphoblastic leukemia (ALL; the most
15 common form of cancer in children and adolescents), chronic lymphocytic leukemia (CLL), and non-Hodgkin's lymphoma (NHL) (Khan, W.N., Immunol. Res., 2001, 23, 147-156). Consequently, BTK has emerged as a new molecular target for treatment of B-lineage leukemia's and lymphomas.

BTK mutations in human cause an inherited disease X-linked
20 agammaglobulinemia, characterized by a lack of peripheral B-cells and low levels of serum Ig. Thus, BTK is a uniquely attractive kinase target for selective B-cell inhibition and small molecule based BTK inhibitors offers therapeutic benefit in the treatment of lymphoma and autoimmune diseases (Väliäho, J., Smith, C.I., Vihinen, M., Hum. Mutat., 2006, 27(12), 1209-1217).

25 BTK was recently identified in a siRNA screen as an essential kinase for survival in a subset of diffuse large-cell lymphomas driven by activated BCR, where an irreversible BTK inhibitor, PCI-32765 (Ibrutinib), was shown to promote apoptosis. A second study of Ibrutinib recently demonstrated in vivo clinical

responses in dogs with aggressive B-cell lymphomas (Honigberg, L. A., Smith, A.M., Sirisawad, M., Verner, E., Loury, D., Chang, B., Li, S., Pan, Z., Thamm, D. H., Miller, R.A., Buggy, J.J., Proc. Natl. Acad. Sci., U S A., 2010, 107(29), 13075-13080).

- 5 Thus BTKs are important in the regulation of many cellular processes including cell cycle regulation, proliferation, survival, apoptosis and motility and are significant components of the molecular mechanisms of diseases such as cancer, diabetes and immune inflammation (Buggy, J.J., Elias, L., Int. Rev. Immunol., 2012, 31(2), 119-132).

10 **Prior art**

- Published patent applications US2008/0076921, US2008/0108636, US2011/224235, US2012/100138/ US2012/087915, US2012/183535, US2012/129873, US2012/129821, US2012/053189, WO2008/039218, WO2008/054827, WO2008/121742, WO2010/009342, WO2011/046964, 15 WO2014/078578 and WO2011/153514, by Pharmacyclics Inc., discloses pyrazolo-pyrimidin-amine derivatives as BTK inhibitors for the treatment of autoimmune diseases or conditions, heteroimmune diseases or conditions, such as cancer, including lymphoma and inflammatory diseases or conditions. Hoffmann La Roche (WO2010/100070, US2010/222325, WO2010/122038, US2010/273768, 20 WO2012/020008, US2012/040949, WO2010/000633, US2010/004231, WO2010/006947, US2010/016301, WO2014/064131, WO2014/076104, WO2014/083026, WO2014/090715), Biogen Idec Inc (WO2011/029046, US2012/157443, WO2011/029043, US2012/157442), CGI Pharmaceuticals Inc (WO2006/099075, US2006/229337, WO2009/137596, US2011/118233, 25 WO2010/056875, US2011/301145, WO2010/068788), Avila Therapeutics Inc (US2012/0277832, WO2012/021444), BMS (WO2002/38797, US2003/040461), Boehringer (WO2014/025976), Principia (US2014/8673925), Ono Pharmaceuticals (US2013/0217880), Merck Sharp & Dohme (WO2013/010380, WO2014/093230) and Cellular Genomics Inc (WO2005/005429, US2005/101604) discloses diverse

class of heterocyclic compounds as BTK inhibitors. BASF (WO2001/019829 and WO2002/080926) disclosed certain class of heterocyclic compounds as protein kinase inhibitors. Some of the recent patent application such as WO2014/187262, WO2014/188173, WO2014/161799, CN104086551, WO2014/135473, 5 WO2014/116504, WO2014/113942, US2014/0206681, and WO2014/064131 discloses diverse class of heterocyclic compound as BTK inhibitors.

Since the ATP binding site of BTK shows a close homology to that of other Src-family kinases, such as lymphocyte-specific protein tyrosine kinase (LCK) and LYN, it is often difficult to find an ATP competitive inhibitor having sufficient 10 selectivity. A lack of selectivity of an inhibitor for BTK over these kinases could potentially have negative consequences. BTK is a member of a group of eleven tyrosine kinases (the Tec family kinases, EGFR, Jak3, ErbB2, ErbB4, and BLK) that contain a conserved cysteine residue adjacent to the ATP-binding site. This cysteine (Cys₄₈₁ in BTK) is a potential nucleophilic site which could form a covalent adduct 15 with an electrophilic inhibitor. As the inhibitory activity of such an inhibitor is dependent on the covalent interaction, this cysteine residue provides a handle for achieving the desired degree of selectivity (Singh, J., Petter, R.C., Kluge, A.F., Curr. Opin. Chem. Biol., 2010, 14, 1-6; Cohen, M.S., Zhang, C., Shokat, K.M., Taunton, J., Science, 2005, 308, 1318-1321; Leproult, E., Barluenga, S., Moras, D., Wurtz, J.M., 20 Winssinger, N., J. Med. Chem., 2011, 54, 1347-1355).

Reversible kinase inhibitors interact with the ATP-binding site. As the ATP-binding sites are highly conserved among kinases, it is difficult to develop a reversible inhibitor that selectively inhibits a desired kinase. Thus with reversible kinase inhibitors, it is difficult to achieve broad therapeutic window. Generating 25 potent, selective, oral BTK inhibitors using covalent, irreversible and electrophilic compounds is feasible approach. However, the irreversible inhibitors exhibit toxicity due to covalent bonding with off-target macromolecules. Thus to overcome undesirable off-target effects, it is essential to develop irreversible BTK inhibitors that covalently bind with BTK enzyme, without binding to off-target polypeptides.

Several BTK inhibitors are being developed as therapeutic agents for various indications. Among these, the covalent inhibitor Ibrutinib (Pharmacyclics) was developed as a selective and irreversible inhibitor of BTK, targeting the cysteine-481 residue in the active site. Ibrutinib is a potent nanomolar inhibitor of BTK and exhibited promising activity in preclinical models of BCR-driven B-lineage lymphoma and clinical testing in lymphoma patients (Pan, Z., Scheerens, H., Li, S.J., Schultz, B.E., Sprengeler, P.A., Burrill, L.C., Mendonca, R.V., Sweeney, M.D., Scott, K.C., Grothaus, P.G., Jeffery, D.A., Spoerke, J.M., Honigberg, L.A., Young, P.R., Dalrymple, S.A and Palmer, J.T., *Chem. Med. Chem.*, 2007, 2, 58-61; Honigberg, L. A., Smith, A.M., Sirisawad, M., Verner, E., Loury, D., Chang, B., Li, S., Pan, Z., Thamm, D.H., Miller, R.A and Buggy, J.J., *Proc. Natl. Acad. Sci. U S A.*, 2010, 107, 13075-13080). Likewise, dianilinopyrimidine-based irreversible BTK inhibitors with micromolar activity were developed and two lead compounds, AVL-101 and AVL-291 (Avila Therapeutics) showed promising in vitro activity against lymphoma cells (Evans, E., Ponader, S., Karp, R., et al., *Clin. Lymphoma Myeloma Leuk.*, 2011, 11 Suppl 2, S173-S174). Ibrutinib is a covalent BTK inhibitor, recently approved for the treatment of patients with various B-cell malignancies. Thus inhibition of BTK is emerging as a promising mechanism for targeting B-cell malignancies (Harrison, C., *Nat. Rev. Drug Discov.*, 2012, 11(2), 96-97).

We herein disclose novel heterocyclic compounds of general formula (I) which are selective BTK inhibitors for the prevention and treatment of disease states mediated by BTK, including cancer and inflammation. More particularly, embodiments of the present invention describe irreversible kinase inhibitors including, but not limited to, inhibition of BTK, that are useful as therapeutics in the treatment of a variety of pathological conditions including (but not limited to) cancer, lymphoma, auto-immune diseases, heteroimmune diseases, inflammatory diseases and neurodegenerative diseases or conditions.

SUMMARY OF THE INVENTION

The present invention discloses heterocyclic compounds as defined by the general formula (I) that are selective BTK inhibitors for the prevention and treatment of disease states mediated by BTK. The compounds of the present invention are useful in the treatment of human or animal body, by inhibition of BTK. The compounds of this invention are therefore suitable for the prevention and treatment of disease states mediated by BTK.

EMBODIMENT(S) OF THE INVENTION

An embodiment of the present invention provides novel heterocyclic compounds represented by the general formula (I), their tautomeric forms, their enantiomers, their diastereoisomers, their stereoisomers, their pharmaceutically acceptable salts and pharmaceutical compositions containing them or their mixtures thereof.


In a further embodiment of the present invention is provided pharmaceutical compositions containing compounds of the general formula (I), their tautomeric forms, their enantiomers, their diastereoisomers, their stereoisomers, their pharmaceutically acceptable salts, or their mixtures in combination with suitable carriers, solvents, diluents and other media normally employed in preparing such compositions.

In a still further embodiment is provided the use of heterocyclic compounds of the present invention as selective BTK inhibitors, by administering a therapeutically effective and non-toxic amount of compounds of general formula (I) or their pharmaceutically acceptable compositions to the mammals.

In a still further embodiment is provided a process for preparing the novel compounds of the present invention.

DESCRIPTION OF THE INVENTION

Accordingly, the present invention relates to the compounds of the general formula (I) represented below and their pharmaceutically acceptable salts, enantiomers and their diastereomers;




5

(I)

Wherein, V, W, X, Y & Z independently represents, 'C' or 'N'; R₁, represents groups selected from hydrogen, keto, halogen, unsubstituted or substituted groups selected from cyano, alkyl, haloalkyl, aryl, alkoxy, acyloxy, aryloxy, arylalkyl, heteroaryl, heterocyclyl, heterocycloalkyl, cycloalkyl, cycloalkylalkyl, aryloxyaryl, 10 aryloxyalkyl, aryloxyheteroaryl groups;

R₂ represent the following ring system:



Wherein R_3 at each occurrence is independently selected from hydrogen, haloalkyl, C_{1-7} alkyl, C_{2-7} alkenyl, C_{2-7} alkynyl, aryl, cycloalkyl, heterocycloalkyl, cycloalkyl(C_{1-7})alkyl, heterocycloalkyl(C_{1-7})alkyl, $C(O)NH(C_{1-7})alkyl$, $C(O)-CH=CH_2$, $C(O)-CH=CH-R_4$, $C(O)-C(CN)=CH_2$, $C(O)-C(CN)=CH-R_4$, $SO_2-NH(C_{1-7})alkyl$, $SO_2-CH=CH_2$, $SO_2-CH=CH-R_4$ groups;

R_4 is selected from $-(CH_2)_n-NR_5R_6$; wherein, $n=0-7$ and each of R_5 and R_6 are independently selected from hydrogen, haloalkyl, C_{1-7} alkyl, C_{2-7} alkenyl, C_{2-7} alkynyl, aryl, cycloalkyl, carbocycle, heterocycloalkyl, cycloalkyl(C_{1-7})alkyl, heterocycloalkyl(C_{1-7})alkyl; 'U' represent unsubstituted or substituted groups selected from alkyl, alkenyl, alkynyl, alkoxy, acyloxy, aryl, aryloxy, arylalkyl, cycloalkyl, cycloalkylalkyl, biaryl, heteroaryl, heterocycle, heterocycloalkyl, O-aryl, O-cycloalkyl, O-heteroaryl, O-heterocycle, O-heterocycloalkyl, aryloxyaryl, aryloxyalkyl, aryloxyheteroaryl, heteroaryloxyaryl, heteroaryloxyalkyl, heteroaryloxyheteroaryl, $Ph-CO-N(R_7R_8)$, $Ph-N(R_9)-CO-R_{10}$, wherein, R_7 , R_8 and R_{10} are independently selected from hydrogen, halogen, alkyl, haloalkyl, alkoxy; aryl, cycloalkyl, heteroaryl, heterocycloalkyl; further substituted with halogen, alkyl, alkoxy, haloalkoxy groups and R_9 are independently selected from hydrogen, C_{1-7} alkyl, C_{2-7} alkenyl, C_{2-7} alkynyl.

In a preferred embodiment, the groups, radicals described above may be selected from:

"Alkyl", as well as other groups having the prefix "alk", such as alkoxy and alkanoyl, means a carbon chain which may further be substituted with an oxygen atom as is well understood by a skilled artisan, which may further be either linear or branched, and combinations thereof, unless the carbon chain is defined otherwise. Examples of alkyl group include but not are limited to methyl, ethyl, propyl, isopropyl, butyl, *sec*-butyl, *tert*-butyl, pentyl, hexyl etc. Where the specified number of carbon atoms permits e.g. from C_{3-10} , the term alkyl also includes cycloalkyl groups, and combinations of linear or branched alkyl chains combined with cycloalkyl structures. When no number of carbon atoms is specified, C_{1-6} is intended.

“Alkenyl” means carbon chains which contain at least one carbon-carbon double bond, and which may be linear or branched or combinations thereof, unless the carbon chain is defined otherwise. Examples of alkenyl include but not limited to vinyl, allyl, isopropenyl, hexenyl, pentenyl, heptenyl, 1-propenyl, 2-butenyl, 2-methyl-2-butenyl etc. Where the specified number of carbon atoms permits, e. g., from C₅₋₁₀, the term alkenyl also includes cycloalkenyl groups and combinations of linear, branched and cyclic structures. When no number of carbon atoms is specified, C₍₂₋₆₎ is intended.

“Alkynyl” means carbon chains which contain at least one carbon-carbon triple bond, and which may be linear or branched or combinations thereof. Examples of alkynyl include ethynyl, propargyl, 3-methyl-1-pentynyl etc. When no number of carbon atoms is specified, C₍₂₋₆₎ is intended.

As used herein, "carbocycle" or "carbocyclic residue" is intended to mean any stable monocyclic or bicyclic or tricyclic ring, any of which may be saturated, partially unsaturated, or aromatic. Examples of such carbocycles include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, adamantyl, cyclooctyl, [3.3.0]bicyclooctane, [4.3.0]bicyclononane, [4.4.0]bicyclodecane (decalin), [2.2.2]bicyclooctane, fluorenyl, phenyl, naphthyl, indanyl, adamantyl, or tetrahydronaphthyl (tetralin). In a broader perspective, the term carbocycle is intended to include, wherever applicable, the groups representing cycloalkyl, phenyl and other saturated, partially saturated or aromatic residues;

“Cycloalkyl” is the subset of alkyl and means saturated carbocyclic ring having a specified number of carbon atoms, preferably 3-6 carbon atoms. Examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl etc. A cycloalkyl group generally is monocyclic unless otherwise stated. Cycloalkyl groups are saturated unless and otherwise stated.

The “alkoxy” refers to the straight or branched chain alkoxides of the number of carbon atoms specified.

“Aryl” means a mono- or polycyclic aromatic ring system containing carbon ring atoms. The preferred aryls are monocyclic or bicyclic 6-10 membered aromatic ring systems. Phenyl and naphthyl are preferred aryls.

The terms “Heterocycle” or “heterocyclyl” refer to saturated or unsaturated non-aromatic rings or ring systems containing at least one heteroatom selected from O, S, N further optionally including the oxidized forms of sulfur, namely SO & SO₂. Examples of heterocycles include tetrahydrofuran (THF), dihydrofuran, 1,4-dioxane, morpholine, 1,4-dithiane, piperazine, piperidine, 1,3-dioxolane, imidazoline, imidazolidine, pyrrolidine, pyrroline, tetrahydropyran, dihydropyran, oxathiolane, dithiolane, 1,3-dioxane, 1,3-dithiane, oxathiane, thiomorpholine, etc. The term “heterocycloalkyl” refers to a heterocyclic group as defined above connected to an alkyl group as defined above;

“Heteroaryl” means an aromatic or partially aromatic heterocycle that contains at least one ring heteroatom selected from O, S and N. Heteroaryls thus include heteroaryls fused to the other kinds of rings, such as aryls, cycloalkyls, and heterocycles that are not aromatic. Examples of heteroaryl groups include; pyrrolyl, isoxazolyl, isothiazolyl, pyrazolyl, pyridyl, oxazolyl, oxadiazolyl, thiadiazolyl, thiazolyl, imidazolyl, triazolyl, tetrazolyl, furyl, triazinyl, thienyl, pyrimidyl, benzisoxazolyl, benzoxazolyl, benzthiazolyl, benzothiadiazolyl, dihydrobenzofuranyl, indolyl, pyridazinyl, indazolyl, isoindolyl, dihydrobenzothienyl, indolyl, pyridazinyl, indazolyl, isoindolyl, dihydrobenzothienyl, indolizyl, cinnolyl, phthalazinyl, quinazolinyl, naphthyridinyl, carbazolyl, benzodioxolyl, quinoxalinyl, purinyl, furazanyl, isobenzylfuranyl, benzimidazolyl, benzofuranyl, benzothienyl, quinolyl, indolyl, isoquinolyl, dibenzofuranyl etc. For heterocyclyl and heteroaryl groups, rings and ring systems containing from 3-15 carbon atoms are included, forming 1-3 rings.

An “aryloxy” group used either alone or in combination with other radicals, is selected from groups containing an aryl radical, as defined above, attached directly to

an oxygen atom, more preferably groups selected from phenoxy, naphthyloxy, tetrahydronaphthyloxy, biphenyloxy, and the like;

“Cycloalkylalkyl” means an alkyl radical substituted with cycloalkyl group as defined herein. Non-limiting cycloalkylalkyl groups include cyclopropylmethyl, 5 cyclobutylmethyl, cyclopentylmethyl, cyclohexylmethyl, and the like.

An “arylalkyl” group as used herein is an aromatic substituent that is linked to an alkyl group having from one to about six carbon atoms. Non-limiting examples of arylalkyl groups include benzyl group, phenethyl and the like.

The “acyloxy” group used either alone or in combination with other radicals, 10 is selected from a suitable acyl group, directly attached to an oxygen atom; more preferably such groups are selected from acetyloxy, propionyloxy, butanoyloxy, *iso*-butanoyloxy, benzoyloxy and the like;

The term “Haloalkyl” means a alkyl structure in which at least one hydrogen is replaced with a halogen atom. In certain embodiments in which two or more 15 hydrogen atoms are replaced with halogen atoms, the halogen atoms are all the same as one another.

In certain other embodiment in which two or more hydrogen atoms are replaced with halogen atoms, the halogen atoms are not all the same as one another.

“Aryloxyalkyl” means an alkyl radical substituted with aryloxy group as 20 defined herein.

“Aryloxyaryl” means an aryl radical substituted with aryloxy group as defined herein.

“Aryloxyheteroaryl” means a heteroaryl radical substituted with aryloxy group as defined herein.

25 “Halo/ Halogen” refers to fluorine, chlorine, bromine, iodine. Chlorine and fluorine are generally preferred.

Suitable groups and substituents on the groups may be selected from those described anywhere in the specification.

The term "substituted," as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency is not exceeded, and that the substitution results in a stable compound. The term "substituted," as used herein, means that any one or more hydrogens on the designated atom is replaced with a selection from the indicated group, provided that the designated atom's normal valency is not exceeded, and that the substitution results in a stable compound.




"Pharmaceutically acceptable salts" refer to derivatives of the disclosed compounds wherein the parent compound is modified by making acid or base salts thereof. Examples of pharmaceutically acceptable salts include, but are not limited to, mineral or organic acid salts of the basic residues. Such conventional non-toxic salts include, but are not limited to, those derived from inorganic and organic acids selected from 1, 2-ethanedisulfonic, 2-acetoxybenzoic, 2-hydroxyethanesulfonic, acetic, ascorbic, benzenesulfonic, benzoic, bicarbonic, carbonic, citric, edetic, ethane disulfonic, ethane sulfonic, fumaric, glucoheptonic, gluconic, glutamic, glycolic, glycollyarsanilic, hexylresorcinic, hydrabamic, hydrobromic, hydrochloric, hydroiodide, hydroxymaleic, hydroxynaphthoic, isethionic, lactic, lactobionic, lauryl sulfonic, maleic, malic, mandelic, methanesulfonic, napsylic, nitric, oxalic, pamoic, pantothenic, phenylacetic, phosphoric, polygalacturonic, propionic, salicyclic, stearic, subacetic, succinic, sulfamic, sulfanilic, sulfuric, tannic, tartaric, and toluenesulfonic.




The term 'optional' or 'optionally' means that the subsequent described event or circumstance may or may not occur, and the description includes instances where the event or circumstance occur and instances in which it does not. For example, 'optionally substituted alkyl' means either 'alkyl' or 'substituted alkyl'. Further an optionally substituted group includes an unsubstituted group.





Unless otherwise stated in the specification, structures depicted herein are also meant to include compounds which differ only in the presence of one or more isotopically enriched atoms.




Particularly useful compounds may be selected from but not limited to the following:




Table 1:




Compd	Structures	IUPAC Names
1		1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
2		1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
3		1-(5-(5-((4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)pyridin-2-yl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)prop-2-en-1-one





4		1-(5-(2-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)prop-2-en-1-one
5		1-(5-(4-amino-3-(2-methylbenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
6		1-(5-(4-amino-3-(benzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one





7		1-(5-(4-amino-3-(2,3-dihydrobenzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
8		1-(5-(4-amino-3-(dibenzo[b,d]furan-3-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
9		N-(6-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]thiazol-2-yl)acetamide
10		1-(5-(4-amino-3-(2-methoxybenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one





11		1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-yn-1-one
12		1-(5-(4-amino-3-(3-methoxy-4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
13		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyridin-2-yl)benzamide





14		1-(5-(4-amino-3-(2-phenylbenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
15		1-(5-(4-amino-3-(benzo[d][1,3]dioxol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
16		1-(5-(4-amino-3-(4-(5-methyl-1,3,4-oxadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one





17		1-(5-(4-amino-3-(benzo[d]oxazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
18		3-(4-phenoxyphenyl)-1-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine
19		1-(5-(4-amino-3-(2-phenylbenzo[d]oxazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one





20		1-(5-(4-amino-3-(2-phenoxybenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
21		1-(5-(4-amino-3-(4-(1-methyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
22		1-(5-(4-amino-3-(benzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
23		1-(5-(4-amino-3-(2-phenylbenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one





24		(E)-1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one
25		9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-7-(4-phenoxyphenyl)-5,7-dihydro-4H-purin-8(9H)-one
26		1-(5-(4-amino-3-(4-(5-(pyridin-2-yl)-1,3,4-oxadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
27		1-(5-(4-amino-6-bromo-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one





28		(E)-2-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydrocyclopenta[c]pyrrole-2-carbonyl)-3-cyclopropylacrylonitrile
29		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-2-methoxy-N-(4-methylpyridin-2-yl)benzamide
30		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)benzamide
31		1-(5-(4-amino-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one





32		N-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)phenyl)picolinamide
33		6-amino-7-(4-phenoxyphenyl)-9-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)-7H-purin-8(9H)-one
34		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(benzo[d]thiazol-2-yl)benzamide
35		N-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)phenyl)pyrazine-2-carboxamide



36		1-(5-(4-amino-3-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
37		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyrazin-2-yl)benzamide
38		N-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)phenyl)benzamide
39		1-(5-(4-amino-3-(4-(5-(pyridin-2-yl)-1,3,4-thiadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one

40		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide
41		(Z)-methyl 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)benzimidate
42		6-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)nicotinamide
43		1-(5-(4-amino-3-(4-(pyridin-2-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one

44		(E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-en-1-yl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyridin-2-yl)benzamide
45		(E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-en-1-yl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)benzamide
46		1-(5-(4-amino-3-(4-(5-(pyrazin-2-yl)-1,3,4-thiadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
47		(E)-1-(5-(4-amino-3-(4-(pyridin-2-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one

48		(E)-1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one
49		(E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyrazin-2-yl)benzamide
50		(Z)-methyl 4-(4-amino-1-(2-((E)-4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)benzimidate
51		(E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide

52		1-(5-(4-amino-3-(4-(pyridin-3-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
53		(E)-1-(5-(4-amino-3-(4-(pyridin-3-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one
54		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-methyl-N-(4-methylpyridin-2-yl)benzamide
55		(E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-en-1-yl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-methyl-N-(4-methylpyridin-2-yl)benzamide

56		(E)-1-(5-(4-amino-3-(4-(5-(pyrazin-2-yl)-1,3,4-thiadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one
57		(E)-1-(5-(4-amino-3-(4-(5-(pyridin-2-yl)-1,3,4-oxadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one

or pharmaceutically acceptable salts of any of the compounds above.


Following is a list of abbreviations used in the description of the preparation of the compounds of the present invention:

- 5 ALL: Acute lymphoblastic leukemia
- ATP: Adenosine triphosphate
- BTK: Bruton's Tyrosine Kinase
- bs: broad singlet
- CLL: Chronic lymphocytic leukemia
- 10 CDCl₃: Deuterated chloroform
- CHCl₃: Chloroform
- d: doublet
- dd: doublet of doublet
- dt: doublet of triplet
- 15 DCM: Dichloromethane
- DMAC: N,N-(Dimethylacetamide)
- DMAP: 4-(Dimethylamino) pyridine

- DMF: *N,N*-Dimethyl formamide
DMSO: Dimethyl sulfoxide
EDTA: Ethylenediaminetetraacetic acid
EtOAc: Ethyl acetate
5 EtOH: Ethanol
HCl(g): Hydrogen chloride (gas)
K₂CO₃: Potassium carbonate
MeOH: Methanol
m: multiplet
10 mmol: millimoles
μg : microgram
MS: Mass spectrum
NHL: Non-Hodgkin's lymphoma
Na₂CO₃: Sodium carbonate
15 ng : nanogram
NIS: N-iodosuccinimide
¹H NMR: Proton nuclear magnetic resonance
Pet ether: Petroleum ether, boiling range (60-80 °C)
POCl₃ : Phosphorylchloride
20 s: singlet
t: Triplet
td: triplet of doublet
THF: Tetrahydrofuran
TLC: Thin layer chromatography
25 The novel compounds of the present invention can be prepared using the reactions and techniques described below, together with conventional techniques known to those skilled in the art of organic synthesis, or variations thereon as appreciated by those skilled in the art.

The reactions can be performed in solvents appropriate to the reagents and materials employed and suitable for the transformations being affected. Preferred methods include, but not limited to those described below, where all symbols are as defined earlier unless and otherwise defined below.

5 The compounds of the formula (I) can be prepared as described in schemes below along with suitable modifications/variations which are well within the scope of a person skilled in the art.



Scheme I:

10

Wherein 'U', R_2 and R_3 are as defined earlier. Compound of formula (I) can be prepared by variety of methods familiar to those skilled in art. Compound of formula (i) was transformed into compound (ii) by reacted with hydrazine hydrate (**Scheme-I**). Compound of formula (ii) was cyclized using formamide to afford the compound of formula (iii). Compound (iii) was reacted with N-iodosuccinimide to get compound (iv). Compound (iv) reacted with compound (v) using different base to furnish the compound of formula (vi). Compound (vi) can subjected to Suzuki type of


15

reaction, with compound (vii) using suitable catalysts, base and appropriate solvents to obtain compound of formula (viii). The deprotection of compound (viii) gives compound (ix). Compound (ix) is reacted with optionally substituted acid chlorides (x) to obtain compounds of formula (I).


5 The examples and preparations provided below further illustrate and exemplify the compounds of the present invention and methods of preparing such compounds. It is to be understood that the scope of the present invention is not limited in any way by the scope of the following examples and preparations. In the following examples molecules with a single chiral center, unless otherwise noted,
10 exist as a racemic mixture. Those molecules with two or more chiral centers, unless otherwise noted, exist as a racemic mixture of diastereomers. Single enantiomers/diastereomers may be obtained by methods known to those skilled in the art.

 The compounds of formula (I) may also be synthesized as described in
15 **Scheme II**, wherein 'U', R₂ and R₃ are as defined earlier. Compound (i) may be continuously prepared by variety of methods familiar to those skilled in art. Compound (i) was transformed into compound (ii) by reacting with dibenzyl amine using different bases. Compound of formula (ii) was reacted with different protected cycloalkyl amines (iii) using suitable bases to furnish compound (iv). Compound (iv)
20 was reduced to amine to afford the compound (v). Compound (v) was reacted with Triphosgene to get the compound (vi). Compound (vi) was deprotected to using Pd(OH)₂ to afford compound (vii). Compound (vii) was reacted with different Boronic acid to obtain compound (viii). Compound (viii) was deprotected using suitable acid to get the compound (ix). Compound (ix) was reacted with optionally substituted
25 acid chlorides using base to obtain compound of formula (I).

Scheme II



- The compounds of formula (I) may also be synthesized as described in Scheme III, wherein 'U', R_2 and R_3 are as defined earlier. Compound (i) may be
- continently prepared by variety of methods familiar to those skilled in art. Compound
- (i) was transformed into compound (ii) using Ammonia. Compound (ii) reacted with
- compound (iii) using different base to furnish the compound of formula (iv). Compound (iv) can be subjected to Suzuki type of reaction, with compound (v) using
- suitable catalysts, base and appropriate solvents to obtain compound of formula (vi).
- Compound (vi) can be halogenated to afford compound (vii). The deprotection of
- compound (vii) gives compound (viii). Compound (viii) is reacted with optionally
- substituted acid chlorides to obtain compounds of formula (I).

**Scheme III**

Compounds of the present invention can be isolated either as free amine form or as a salt corresponding to the acid used such as trifluoroacetic acid, hydrochloric acid, hydrobromic acid, oxalic acid, maleic acid, fumaric acid, succinic acid, p-toluene sulfonic acid or benzene sulfonic acid. The compounds can be purified where ever required, by recrystallization, trituration, precipitation, preparative thin layer chromatography, flash chromatography or by preparative HPLC method.

The compounds of the present invention can be used either alone or in combination with one or more therapeutic agents or pharmaceutically acceptable salts thereof. Such use will depend on the condition of the patient being treated and is well within the scope of a skilled practitioner.

The invention is further illustrated by the following non-limiting examples which describe the preferred way of carrying out the present invention. These are provided without limiting the scope of the present invention in any way.

¹H NMR spectral data given in the examples (vide infra) are recorded using a 400 MHz spectrometer (Bruker AVANCE-400) and reported in δ scale. Until and

otherwise mentioned the solvent used for NMR is CDCl_3 using TMS as the internal standard.


Example: 1

- 5 Synthesis of 1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one (compound No: 1)


Synthesis of titled compound was carried out, as described in **Scheme-IV** and step-wise procedure is described below.

10

Scheme-IV:




- 15 **Step-1:** Synthesis of tert-butyl 5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta [c]pyrrole-2(1H)-carboxylate (3)



Intermediate **1** (2.0 g, 7.66 mmol), prepared as per general process disclosed in US 2012/0088912 and triphenylphosphine (6.53 g) were mixed together, in THF
 5 (20 mL). Tert-butyl 5-hydroxyhexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate **2** (3.47 g, 15.32 mmol) was added to the mixture followed by the addition of diisopropyl diazodicarboxylate (2.26 mL, 11.49 mmol). The reaction mixture was stirred at room temperature overnight, filtered and concentrated. The residue obtained was purified by flash chromatography (CH₂Cl₂/MeOH = 98/2) to get intermediate **3**
 10 as a white solid (2.75 g, 76% Yield). ¹H NMR (DMSO-d₆, 400 MHz) δ ppm: 1.40 (s, 9H), 1.94-2.00 (m, 2H), 2.17-2.24 (m, 2H), 2.82-3.00 (m, 2H), 3.10-3.14 (m, 2H), 3.45-3.50 (m, 2H), 5.27-5.30 (m, 1H), 8.29 (s, 1H). MS (ESI-MS): m/z 471.10 (M+H)⁺.


15 **Step-2:** Synthesis of tert-butyl 5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate (**5**)



To a stirred solution Intermediate **3** (2.7 g, 5.74 mmol), dissolved in dry DMF (27 mL), PdCl₂(PPh₃)₂ (0.4 g, 0.57 mmol), 4-phenoxyphenylboronic acid **4** (1.84 g,

8.61 mmol) and KHCO_3 (3.44 g, 34.46 mmol) was added. The reaction mixture was heated at 90°C for 2 hrs, under N_2 atmosphere. Mixture was cooled to room temperature, diluted with water (50 mL) and extracted with EtOAc (3 x 50mL). The combined organic layer was washed with water (2 x 25 mL) and brine solution (25 mL), dried over Na_2SO_4 , and concentrated to dryness. The residue obtained was purified by column chromatography (using 0-5% methanol in DCM as a mobile phase) to obtain Intermediate 5 as an off white solid (2.2 g, 74% Yield). ^1H NMR (DMSO- d_6 , 400 MHz) δ ppm: 1.48 (s, 9H), 1.98-2.04 (m, 2H), 2.27-2.34 (m, 2H), 2.89 (s, 2 H), 3.13-3.17 (m, 2 H), 3.47 (q, 2H, $J=8.0$ Hz), 5.36 (q, 1H, $J=8.0$ Hz), 7.10-7.14 (m, 4H), 7.144-7.20 (m, 1H), 7.40-7.43 (m, 2H), 7.65-7.68 (m, 2H), 8.23 (s, 1 H), ESI-MS (ESI-MS): m/z 535.23 ($\text{M}+\text{Na}$) $^+$.


Step-3: Synthesis of 1-(octahydrocyclopenta[c]pyrrol-5-yl)-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine (6)



15

To a solution of Intermediate 5 (2.1 g, 4.09 mmol) in CH_2Cl_2 (40 mL) was added TFA (1.25 mL, 16.37 mmol). After stirring 2 hrs at room temperature, the solvent was removed and the residues were dissolved in a mixture of ethyl acetate (50 mL) and dilute aq. K_2CO_3 . The organic layer was separated, dried over MgSO_4 , filtered and concentrated to provide Intermediate 6 as a white solid (1.2 g, 71% Yield). ^1H NMR (400 MHz) δ ppm: 1.92-1.96 (m, 2H), 2.31-2.39 (m, 2H), 2.74-2.78 (m, 2H), 2.89-2.30 (m, 2H), 3.12-3.20 (m, 2H), 5.43-5.37 (m, 1H), 7.11-7.20 (m, 5H), 7.41-7.45 (m, 2H), 7.64-7.66 (m, 2H), 8.24 (s, 1H); MS (ESI-MS): m/z 413.20 ($\text{M}+\text{H}$) $^+$.

Step-4: Synthesis of 1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one, compound No: 1



5

To a solution of Intermediate 6 (1.1 g, 2.66 mmol), dissolved in CH₂Cl₂ (30 mL), tri-ethyl amine (1.11 mL, 8.00 mmol) was added followed by addition of acryl chloride (0.2 mL, 2.53 mmol). The reaction was stopped after 2 hrs. The reaction mixture was washed with water and then with brine. The organic layer was separated, dried over MgSO₄, filtered and concentrated. Residue obtained was purified by flash chromatography (using CH₂Cl₂/MeOH=25/1, as a mobile phase) to get compound 1 as a white solid (0.75 g, 60% Yield). ¹H NMR: (CDCl₃, 400 MHz): δ 8.36 (s, 1H), 7.66-7.62 (m, 2H), 7.37-7.41 (m, 2H), 7.13-7.20 (s, 3H), 7.07-7.09 (m, 2H), 6.36-6.50 (m, 2H), 5.68-5.71 (m, 1H), 5.53-5.59 (m, 3H), 3.82-3.87 (m, 2H), 3.45-3.53 (m, 2H), 3.10-3.21 (m, 2H), 2.50-2.58 (m, 2H), 2.11-2.17 (m, 2H); ESI-MS: (+ve mode) 467.20 (M+H)⁺ (100 %); UPLC: 98.09 %.

15


Example: 2

Synthesis of 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyridin-2-yl)benzamide (compound No: 13)


20

Synthesis of titled compound was carried out, as described in **Scheme-V** and step-wise procedure is described below.

Scheme-V:




Step-1: Synthesis of tert-butyl 5-(4-amino-3-iodo-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta [c]pyrrole-2(1H)-carboxylate (**3**)



Intermediate **1** (0.22 g, 0.851 mmol) and triphenylphosphine (0.71 g) were mixed together in THF (10 mL). Tert-butyl 5-hydroxyhexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate **2** (0.38 g, 1.7 mmol) was added to the reaction mixture followed by the addition of diisopropyl diazodicarboxylate (0.24 mL, 1.22 mmol).

The reaction mixture was stirred at room temperature overnight, filtered and concentrated. The residue obtained was purified by flash chromatography ($\text{CH}_2\text{Cl}_2/\text{MeOH} = 98/2$) to get intermediate **3** as a white solid (0.3 g, 76% Yield). ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ ppm: 1.40 (s, 9H), 1.94-2.00 (m, 2H), 2.17-2.24 (m, 2H), 2.82-3.00 (m, 2H), 3.10-3.14 (m, 2H), 3.45-3.50 (m, 2H), 5.27-5.30 (m, 1H), 8.29 (s, 1H). MS (ESI-MS): m/z 471.10 ($\text{M}+\text{H}$) $^+$.

Step-2: Synthesis of tert-butyl 5-(4-amino-3-(4-(pyridin-2-ylcarbamoyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrole-2(1H)-carboxylate (**5**)




10

To a stirred solution of intermediate **3** (0.3 g, 0.638 mmol), dissolved in dry DMF (3 mL) were added $\text{PdCl}_2(\text{PPh}_3)_2$ (0.089 g, 0.127 mmol), (4-(pyridin-2-ylcarbamoyl)phenyl)boronic acid **4** (0.31 g, 0.95 mmol) and KHCO_3 (0.340 g, 3.56 mmol). The reaction mixture was heated at 90 °C for 2hrs, under N_2 atmosphere. Mixture was cooled to room temperature, diluted with water (50 mL) and extracted with EtOAc (3 x 50mL). The combined organic layer was washed with water (2 x 25 mL) and brine solution (25 mL), dried over Na_2SO_4 and concentrated to dryness. The residue obtained was purified by column chromatography (silica gel, 0-5% methanol in DCM) to obtain intermediate **5** as an off white solid (0.25 g, 72.56 % Yield). ^1H NMR ($\text{DMSO}-d_6$, 400 MHz) δ ppm: 1.59 (s, 9H), 2.11-2.17 (m, 2H), 2.49-2.57 (m, 2H), 3.07-3.09 (m, 2 H), 3.28 (bs, 2H), 3.64 (bs, 2H), 5.55 (q, 1H, $J=8.0$ Hz), 7.11 (q,

20


1H, J =8.0 Hz), 7.78-7.81 (m, 1H), 7.82 (m, 2H), 8.10 (d, 2H, J=8.0Hz), 8.35 (m, 1H), 8.41-8.43 (m, 2H), 8.63 (s, 1H). ESI-MS (ESI-MS): m/z 541.41 (M+H)⁺.

Step-3: Synthesis of 4-(4-amino-1-(octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyridin-2-yl)benzamide (6)



To a solution of Intermediate 5 (0.25 g, 0.462 mmol) in CH₂Cl₂ (10 mL), TFA (1.0 mL, 15.87 mmol) was added and the reaction mixture was stirred for 2 hrs at room temperature. The solvent was removed and the residue obtained was dissolved in a mixture of ethyl acetate (50 mL) and dilute aq. K₂CO₃. The organic layer was dried over MgSO₄, filtered and concentrated to get Intermediate 6 as a white solid (0.13 g, 63.85% Yield). ¹H NMR (DMSO-d₆, 400 MHz) δ ppm: 1.15-1.23 (m, 2H), 2.32-2.37 (m, 2H), 2.54-2.58 (m, 2H), 2.93-2.97 (m, 2H), 3.24-3.29 (m, 2H), 5.33-5.37 (m, 1H), 7.16-7.19 (m, 1H), 7.77 (q, 2H, J=12.0Hz), 7.84-7.88 (m, 1H), 8.18-8.20 (m, 2H), 8.22-8.24 (m, 1H), 8.25-8.30 (m, 1H), 8.40-8.41 (m, 1H), 10.83 (s, 1H); MS (ESI-MS): m/z 441.15 (M+H)⁺.


Step-4: Synthesis of 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyridin-2-yl)benzamide; **compound No: 13**



To a solution of Intermediate **6** (0.13 g, 0.295 mmol), dissolved in CH₂Cl₂ (30 mL) and tri-ethyl amine (0.090 g, 0.886 mmol), acryl chloride (0.026 g, 0.295 mmol) was added and the reaction mixture was stirred for 2 hrs. The reaction mixture was washed with water and brine solution. The organic layer was dried over MgSO₄, filtered, concentrated and residue obtained was purified by flash chromatography, using CH₂Cl₂/MeOH (25/1) to get compound **13** as a white solid (0.03 g, 20.58 % Yield). ¹H NMR: (CDCl₃, 400 MHz): δ 8.72 (s, 1H), 8.43 (d, 1H, *J* = 6.4 Hz), 8.39 (s, 1H), 8.35-8.34 (m, 1H), 8.13 (d, 2H, *J* = 8.4 Hz), 7.88 (d, 2H, *J* = 8.4 Hz), 7.83-7.79 (m, 1H), 7.14-7.11 (m, 1H), 6.49 (dd, 1H, *J*₁ = 10.0 Hz, *J*₂ = 16.8 Hz), 6.42 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 16.8 Hz), 5.72 (dd, 1H, *J*₁ = 2.8 Hz, *J*₂ = 10.0 Hz), 5.61-5.55 (m, 3H), 3.89-3.84 (m, 2H), 3.57-3.47 (m, 2H), 3.24-3.21 (m, 1H), 3.15-3.12 (m, 1H), 2.60-2.52 (m, 2H), 2.21-2.14 (m, 1H); ESI-MS: (+ve mode) 495.4 (M+H)⁺ (100 %); HPLC: 99.09.

Example: 3


Compound 2: (1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



¹H NMR: (CDCl₃, 400 MHz): δ 9.11 (s, 1H), 8.41 (s, 1H), 8.34-8.30 (m, 2H), 7.87 (dd, 1H, $J_1 = 1.6$ Hz, $J_2 = 8.4$ Hz), 6.49 (dd, 1H, $J_1 = 9.6$ Hz, $J_2 = 16.8$ Hz), 6.42 (dd, 1H, $J_1 = 2.8$ Hz, $J_2 = 16.8$ Hz), 5.72 (dd, 1H, $J_1 = 2.4$ Hz, $J_2 = 9.6$ Hz), 5.64-5.57 (m, 1H), 5.50 (bs, 2H), 3.89-3.84 (m, 2H), 3.57-3.47 (m, 2H), 3.25-3.20 (m, 1H), 3.17-3.11 (m, 1H), 2.62-2.54 (m, 2H), 2.22-2.13 (m, 2H) ; **ESI-MS:** (+ve mode) 431.9 (M+H)⁺ (100 %); **HPLC:** 96.04 %.

Example:4


- 10 **Compound 3:** 1-(5-(5-((4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)pyridin-2-yl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)prop-2-en-1-one



¹H NMR: (DMSO-d₆, 400 MHz): δ 8.27 (s, 1H), 8.11-8.09 (m, 1H), 7.63-7.61 (m, 2H), 7.44-7.40 (m, 1H), 7.48-7.44 (m, 2H), 7.19-7.17 (m, 1H), 7.13-7.09 (m, 5H), 6.56-6.49 (m, 1H), 6.44-6.41 (m, 1H), 6.11-6.06 (m, 1H), 5.64-5.61 (m, 1H), 5.39 (s, 2H), 4.41-4.39 (m, 2H), 3.81-3.80 (m, 1H), 3.64-3.57 (m, 2H), 3.46-3.45 (m, 2H), 3.19-3.16 (m, 2H); **ESI-MS:** (+ve mode) 559.35 (M+H)⁺ (100 %); **HPLC:** 95.82 %.

Example:5


Compound 4: 1-(5-(2-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)prop-2-en-1-one



- 5 ¹H NMR: (CDCl₃, 400 MHz): δ 8.37 (s, 1H), 7.65 (dd, 2H, *J*₁ = 2.0 Hz, *J*₂ = 6.4 Hz), 7.40 (t, 2H, *J* = 4.4 Hz), 7.18-7.13 (m, 3H), 7.09 (d, 2H, *J* = 7.6 Hz), 6.37-6.27 (m, 2H), 5.61 (dd, 1H, *J*₁ = 3.6 Hz, *J*₂ = 9.2 Hz), 5.41 (bs, 2H), 3.79-3.68 (m, 2H), 3.35 (dd, 1H, *J*₁ = 4.8 Hz, *J*₂ = 12.8 Hz), 3.27 (dd, 1H, *J*₁ = 4.8 Hz, *J*₂ = 10.4 Hz), 3.08-3.05 (m, 1H), 2.96 (t, 1H, *J* = 6.0 Hz), 2.89-2.86 (m, 1H), 2.77-2.75 (m, 1H), 2.70-2.57 (m, 4H); **ESI-MS:** (+ve mode) 496.15 (M+H)⁺ (100 %); **HPLC:** 96.62 %.
- 10

Example:6

Compound 5: 1-(5-(4-amino-3-(2-methylbenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one




15

¹H NMR: (CDCl₃, 400 MHz): δ 8.40 (s, 1H), 8.19 (d, 1H, *J* = 2.0 Hz), 8.12 (d, 1H, *J* = 8.4 Hz), 7.77 (dd, 1H, *J*₁ = 2.0 Hz, *J*₂ = 8.4 Hz), 6.49 (dd, 1H, *J*₁ = 10.0 Hz, *J*₂ = 16.8 Hz), 6.42 (dd, 1H, *J*₁ = 2.8 Hz, *J*₂ = 16.8 Hz), 5.72 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 9.6 Hz), 5.61-5.57 (m, 1H), 5.31 (bs, 2H), 3.89-3.83 (m, 2H), 3.56-3.47 (m, 2H), 3.24-3.22 (m, 1H), 3.14-3.12 (m, 1H), 2.91 (s, 3H), 2.59-2.55 (m, 2H), 2.19-2.14 (m, 2H); **ESI-MS:** (+ve mode) 446.0 (M+H)⁺ (100 %); **HPLC:** 95.09 %.

Example:7


Compound 6: 1-(5-(4-amino-3-(benzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



¹H NMR: (DMSO, 400 MHz): δ 8.24 (s, 1H), 8.16 (dd, 2H, *J*₁ = 6.0 Hz, *J*₂ = 4.4 Hz), 7.85 (d, 1H, *J* = 5.6 Hz), 7.65 (dd, 1H, *J*₁ = 8.4 Hz, *J*₂ = 1.6 Hz), 7.56 (d, 1H, *J* = 5.2 Hz), 7.62 (dd, 1H, *J*₁ = 10.4 Hz, *J*₂ = 16.8 Hz), 6.14 (dd, 1H, *J*₁ = 16.8 Hz, *J*₂ = 2.4 Hz), 5.67 (dd, 1H, *J*₁ = 10.0 Hz, *J*₂ = 2.4 Hz), 5.45-5.41 (m, 1H), 3.81-3.76 (m, 1H), 3.66-3.60 (m, 1H), 3.54-3.50 (m, 1H), 3.42-3.35 (m, 1H), 3.00-3.08 (m, 1H), 2.30-2.98 (m, 1H), 2.38-2.31 (m, 2H), 2.09-2.04 (m, 2H); **ESI-MS:** (+ve mode) 431.0 (M+H)⁺ (100 %), 453.2 (M+ Na)⁺ (25%); **UPLC:** 98.53 %.

Example:8


Compound 7: 1-(5-(4-amino-3-(2,3-dihydrobenzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



¹H NMR (DMSO-d₆, 400 MHz) δ ppm: 8.22 (s, 1H), 7.55-7.51 (m, 2H), 7.42-7.38 (m, 2H), 6.65-6.58 (m, 1H), 6.16-6.11 (dd, 1H, $J_1 = 2.8\text{ Hz}$, $J_2 = 17.2\text{ Hz}$), 5.68-5.66 (m, 1H), 5.64-5.39 (m, 1H), 3.42-3.40 (m, 1H), 3.39-3.37 (m, 1H), 3.35-3.35 (m, 1H), 3.32-3.30 (m, 3H), 3.10-2.83 (m, 2H), 2.82-2.80 (m, 2H), 2.33-2.29 (m, 3H), 2.04-2.03 (m, 2H); (ESI-MS): (+ve mode) 433.05 (M+H)⁺.(100%), UPLC: 95.80%.

Example:9


Compound 8: 1-(5-(4-amino-3-(dibenzo[b,d]furan-3-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



¹H NMR: (DMSO, 400 MHz): δ 8.30-8.28 (m, 1H), 8.26 (s, 1H), 8.22-8.20 (m, 1H), 7.92 (d, 1H, $J = 0.8\text{ Hz}$), 7.75-7.69 (m, 2H), 7.58-7.54 (m, 1H), 7.46-5.43 (m, 1H), 6.66-6.59 (m, 1H), 6.14 (dd, 1H, $J_1 = 2.4\text{ Hz}$, $J_2 = 16.8\text{ Hz}$), 5.67 (dd, 1H, $J_1 = 2.4\text{ Hz}$, $J_2 = 10.4\text{ Hz}$), 5.46-5.43 (m, 1H), 3.82-3.77 (m, 1H), 3.65-3.61 (m, 1H), 3.55-3.51 (m, 1H), 3.39-3.35 (m, 1H), 3.17-2.92 (m, 2H), 2.41-2.33 (m, 2H), 2.11-2.05 (m, 2H); ESI-MS: (+ve mode) 465.1 (M+H)⁺ (100 %), 487.3 (M+ Na)⁺ (10%); UPLC: 95.50.

Example:10

Compound 9: N-(6-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo [3,4-d]pyrimidin-3-yl)benzo[d]thiazol-2-yl)acetamide




5

¹H NMR: (DMSO , 400 MHz): δ 12.41 (s, 1H), 8.24 (s, 2H), 8.87-8.85 (m, 1H), 7.69-7.67 (m, 1H), 6.65-6.58 (m, 1H), 6.14 (dd, 1H, $J_1 = 2.4$ Hz, $J_2 = 14.4$ Hz), 5.66 (dd, 1H, $J_1 = 1.2$ Hz, $J_2 = 10.4$ Hz), 5.44-5.41 (m, 1H), 3.81-3.38 (m, 3H), 3.40-3.33 (m, 1H), 3.11-2.99 (m, 2H), 2.50-2.37 (m, 2H), 2.20 (s, 3H), 2.12-1.90 (m, 2H); **ESI-**
 10 **MS:** (+ve mode) 489.3 (M+H)⁺ (100 %), 511.0 (M+ Na)⁺ (10%); **UPLC:** 95.29 %.

Example:11

Compound 10: 1-(5-(4-amino-3-(2-methoxybenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d] pyrimidin -1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one




15

¹H NMR (DMSO-d₆, 400 MHz) δ ppm: 8.23 (s, 1H), 8.17 (s, 1H), 7.79 (m, 1H, *J* = 8.4 Hz), 7.67-7.65 (m, 1H), 6.64-6.58 (m, 1H), 6.30-6.11 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 16.8 Hz), 5.70-5.65 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 10.4 Hz), 5.45-5.42 (m, 1H), 4.19 (s, 3H), 3.80-3.76 (m, 1H), 3.62-3.53 (m, 1H), 3.40-3.38 (m, 1H), 3.10-2.83 (m, 1H),
 5 2.82-2.80 (m, 2H), 2.36-2.32 (m, 2H), 2.06-2.05 (m, 2H); (ESI-MS): (+ve mode) 462.05 (M+H)⁺.(100%), UPLC: 95.22%, Ret. time = 3.09 min.

Example:12

Compound 11: 1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-yn-1-one
 10




¹H NMR: (DMSO-d₆, 400 MHz): δ 8.23 (s, 1H), 7.67-7.64 (m, 2H), 7.44-7.40 (m, 2H), 7.19-7.10 (m, 5H), 5.4 (s, 1H), 4.21-4.18 (m, 1H), 3.83-3.74 (m, 1H), 3.65-3.61 (m, 2H), 3.05-3.03 (m, 2H), 2.34-2.31 (m, 2H), 2.05-2.03 (m, 2H); **ESI-MS:** (+ve
 15 mode) 465.50 (M+H)⁺ (100 %); HPLC: 99.12 %.

Example:13

Compound 12: 1-(5-(4-amino-3-(3-methoxy-4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one


20



¹H NMR: (CDCl₃, 400 MHz): δ 8.38 (s, 1H), 7.37-7.33 (m, 3H), 7.22 (dd, 1H, *J*₁ = 8.0 Hz, *J*₂ = 2.0 Hz), 7.14-7.08 (m, 2H), 7.03 (d, 2H, *J* = 8.0 Hz), 6.51-6.37 (m, 2H), 5.70 (dd, 1H, *J*₁ = 10.0 Hz, *J*₂ = 2.4 Hz), 5.65 (bs, 2H), 5.60-5.53 (m, 1H), 3.95 (s, 3H), 3.89-3.84 (m, 2H), 3.55-3.51 (m, 2H), 3.24-3.21 (m, 1H) 3.15-3.11 (m, 1H), 2.63-2.54 (m, 2H), 2.21-2.12 (m, 2H); **ESI-MS:** (+ve mode) 497.1 (M+H)⁺ (100 %), 519.25 (M+ Na)⁺ (50%); **UPLC:** 95.90 %.

Example: 14

10 **Compound 14:** 1-(5-(4-amino-3-(2-phenylbenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one




¹H NMR: (CDCl₃, 400 MHz): δ 8.41 (s, 1H), 8.27 (d, 1H, *J* = 1.6 Hz), 8.23 (d, 1H, *J* = 8.0 Hz), 8.15-8.13 (m, 2H), 7.82 (dd, 1H, *J*₁ = 8.0 Hz, *J*₂ = 2.0 Hz), 7.56-7.53 (m, 3H), 6.52-6.38 (m, 2H), 5.70 (dd, 1H, *J*₁ = 10.0 Hz, *J*₂ = 2.0 Hz), 5.63-5.59 (m, 1H), 5.49 (bs, 2H), 3.90-3.84 (m, 2H), 3.57-3.48 (m, 2H), 3.25-3.22 (m, 1H), 3.17-3.14

(m, 1H), 2.63-2.55 (m, 2H), 2.22-2.13 (m, 2H); **ESI-MS**: (+ve mode) 507.6 (M+H)⁺ (100 %), 530.1 (M+ Na)⁺ (30%); **UPLC**: 97.51 %.

Example:15


- 5 **Compound 15**: 1-(5-(4-amino-3-(benzo[d][1,3]dioxol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



- ¹H NMR: (DMSO-d₆, 400 MHz): δ 8.21 (s, 1H), 7.15-7.13 (m, 1H), 7.12-7.10 (m, 1H), 7.09-7.07 (m, 1H), 6.65-6.58 (m, 1H), 6.16-6.12 (m, 1H), 6.11 (s, 2H), 5.68-5.64 (m, 1H), 5.42-5.35 (m, 1H), 3.83-3.81 (m, 1H), 3.80-3.75 (m, 1H), 3.65-3.60 (m, 1H), 3.50-3.49 (m, 1H), 3.08-3.06 (m, 1H), 2.99-2.96 (m, 1H), 2.36-2.82 (m, 2H), 2.07-2.04 (m, 2H); **ESI-MS**: (+ve mode) 419.58 (M+H)⁺ (100 %); **HPLC**: 96.33 %.

Example: 16


- 15 **Compound 16**: 1-(5-(4-amino-3-(4-(5-methyl-1,3,4-oxadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



¹H NMR: (DMSO-d₆, 400 MHz): δ 8.25 (s, 1H), 8.13-8.11 (d, 2H, *J* = 8.0 Hz), 7.89-7.87 (d, 2H, *J* = 8.0 Hz), 6.65-6.58 (m, 1H), 6.16-6.11 (dd, 1H, *J*₁ = 2.8 Hz, *J*₂ = 16.8 Hz), 5.68-6.65 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 20 Hz), 5.46-5.41 (m, 1H), 3.78-3.76 (m, 1H), 3.64-3.61 (m, 1H), 3.54-3.50 (m, 1H), 3.39-3.34 (m, 1H), 3.23-3.08 (m, 1H), 3.07-3.00 (m, 1H), 2.61 (s, 3H), 2.38-2.32 (m, 2H), 2.07-2.05 (m, 2H); **ESI-MS:** (+ve mode) 457.10 (M+H)⁺ (100 %); **UPLC:** 95.87 %.

Example:17


Compound 17: 1-(5-(4-amino-3-(benzo[d]oxazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



¹H NMR: (CDCl₃, 400 MHz): δ 8.40 (s, 1H), 8.22 (s, 1H), 7.99-7.97 (m, 2H), 7.74-7.72 (m, 1H), 6.47-6.42 (m, 1H), 5.73-5.70 (m, 1H), 5.78-5.60 (m, 2H), 3.86-3.84 (m, 2H), 3.57-3.55 (m, 2H), 3.22-3.19 (m, 2H), 2.56-2.54 (m, 2H), 2.18-2.16 (m, 2H); **ESI-MS:** (+ve mode) 416.78 (M+H)⁺ (100 %); **HPLC:** 96.12 %.

Example:18


Compound 18: 3-(4-phenoxyphenyl)-1-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine



¹H NMR: (DMSO, 400 MHz): δ 8.24 (s, 1H), 7.64 (dd, 2H, $J_1 = 6.8$ Hz, $J_2 = 2.0$ Hz), 7.44-7.40 (m, 2H), 7.19-7.10 (m, 5H), 6.94-6.87 (m, 1H), 6.21 (d, 1H, $J = 10.0$ Hz),
 5 6.15 (d, 1H, $J = 16.8$ Hz), 5.41-6.53 (m, 1H), 3.29-3.24 (m, 2H), 3.04-3.01 (m, 4H), 2.34-2.32 (m, 2H), 2.10-1.90 (m, 2H); **ESI-MS**: (+ve mode) 503.15 (M+H)⁺ (100 %); **UPLC**: 95.16 %.

Example:19


10 **Compound 19**: 1-(5-(4-amino-3-(2-phenylbenzo[d]oxazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



¹H NMR: (CDCl₃, 400 MHz): δ 8.41 (s, 1H), 8.32-8.30 (m, 2H), 7.96-7.93 (m, 2H), 7.73-7.70 (m, 1H), 7.59-7.57 (m, 3H), 6.52-6.43 (m, 1H), 5.72-5.69 (m, 1H), 5.62-
 15 5.59 (m, 1H), 5.50-5.49 (m, 1H), 3.90-3.84 (m, 2H), 3.58-3.48 (m, 2H), 3.23-3.19 (m, 2H), 2.60-2.58 (m, 2H), 2.20-2.17 (m, 2H); **ESI-MS**: (+ve mode) 492.35 (M+H)⁺ (100 %); **HPLC**: 95.63 %.

Example: 20


Compound 20 : 1-(5-(4-amino-3-(2-phenoxybenzo[d]thiazol-6-yl)-1H-pyrazolo [3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



- 5 ¹H NMR (DMSO-d₆, 400 MHz) δ ppm: 8.22 (s, 1H), 7.82 (d, 1H, *J* = 8.4 Hz), 7.70-7.68 (m, 1H), 7.56-7.53 (m, 2H), 7.49-7.47 (m, 2H), 7.42-7.40 (m, 1H), 6.65-6.61 (m, 1H), 6.16 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 14.4 Hz), 5.68 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 10.4 Hz), 5.64-5.40 (m, 1H), 3.58-3.50 (m, 1H), 3.38-3.36 (m, 1H), 3.35-3.33 (m, 1H), 3.25-2.84 (m, 2H), 2.82-2.80 (m, 2H), 2.36-2.32 (m, 2H), 2.08-2.00 (m, 2H); (ESI-MS):
- 10 (+ve mode) 524.15 (M+H)⁺. (100%), UPLC: 95.74%.

Example: 21


Compound 21: 1-(5-(4-amino-3-(4-(1-methyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazolo [3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



¹H NMR (DMSO-d₆, 400 MHz) δ ppm: 8.38 (s, 1H), 7.86-7.83 (m, 1H), 7.71-7.70 (m, 3H), 7.69-7.63 (m, 2H), 6.51-6.37 (m, 2H), 5.72-5.69 (m, 1H), 5.59-5.44 (m, 2H), 3.99 (s, 1H), 3.50-3.46 (m, 2H), 3.23-3.14 (m, 2H), 2.57-2.55 (m, 2H), 2.18-2.14 (m, 2H), 1.68-1.59 (m, 2H); (ESI-MS): (+ve mode) 455.10 (M+H)⁺ (100%), HPLC: 95.98%.

Example:22

Compound 22: 1-(5-(4-amino-3-(benzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



10


¹H NMR: (DMSO-d₆, 400 MHz): δ 8.84 (s, 1H), 8.24 (s, 1H), 8.01 (s, 1H), 7.93-7.91 (d, 1H, *J* = 8.0 Hz), 7.73-7.71 (d, 1H, *J* = 8.0 Hz), 6.65-6.58 (m, 1H), 6.16-6.11 (dd, 1H, *J*₁ = 2.8 Hz, *J*₂ = 16.8 Hz), 5.68-5.65 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 11.2 Hz), 5.44-5.41 (m, 1H), 3.78-3.66 (m, 2H), 3.63-3.60 (m, 2H), 3.53-3.50 (m, 1H), 3.40-3.38 (m, 1H), 3.15-2.85 (m, 2H), 2.07-2.05 (m, 2H); ESI-MS: (+ve mode) 416.10 (M+H)⁺ (100 %); UPLC: 95.64 %.

15

Example: 23

Compound 23: (5-(4-amino-3-(2-phenylbenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one


20



¹H NMR: (DMSO-*d*₆, 400 MHz): δ 8.24-8.27 (m, 3H), 8.02-8.01 (d, 1H, *J* = 4.0 Hz), 7.95-7.93 (d, 1H, *J* = 8.0 Hz), 7.73-7.70 (m, 1H), 7.67-7.63 (m, 3H), 6.66-6.59 (m, 1H), 6.16-6.12 (m, 1H), 5.68-5.65 (m, 1H), 5.45-5.42 (m, 1H), 3.81-3.66 (m, 1H),
 5 3.64-3.61 (m, 1H), 3.55-3.50 (m, 1H), 3.39-3.35 (m, 1H), 3.10-3.00 (m, 2H), 2.44-2.31 (m, 2H), 2.09-2.04 (m, 2H); **ESI-MS:** (+ve mode) 492.05 (M+H)⁺ (100 %); **UPLC:** 97.40 %.

Example:24

10 **Compound 24:** (E)-1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one




¹H NMR: (D₂O, 400 MHz): δ 8.38 (s, 1H), 7.65 (d, 2H, *J* = 6.8 Hz), 7.62-7.47 (m, 2H), 7.46-7.45 (m, 1H), 7.29-7.16 (m, 2H), 6.81-6.69 (m, 2H), 5.56-5.52 (m, 1H),
 15 3.99-3.91 (m, 3H), 3.80 (dd, 1H, *J*₁ = 8.4 Hz, *J*₂ = 13.2 Hz), 3.63 (dd, 1H, *J*₁ = 4.4 Hz, *J*₂ = 11.2 Hz), 3.50 (dd, 1H, *J*₁ = 4.8 Hz, *J*₂ = 13.2 Hz), 3.20-3.11 (m, 2H), 2.93

(s, 6H), 2.47 -2.41 (m, 2H), 2.23-2.16 (m, 2H). ; ESI-MS: (+ve mode) 525.7 (M+H)⁺ (100 %); HPLC: 97.25 %.

Example:25


- 5 **Compound 25:** 9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-7-(4-phenoxyphenyl)-5,7-dihydro-4H-purin-8(9H)-one



- ¹H NMR (DMSO-d₆, 400 MHz) δ ppm: 8.12 (s, 1H), 7.45-7.41 (m, 4H), 7.20-7.17 (m, 1H), 7.15-7.11 (m, 1H), 6.63-6.56 (m, 1H), 6.15 (dd, 1H, *J*₁ = 4.0 Hz, *J*₂ = 16.0 Hz), 5.74-5.72 (m, 2H), 5.97-5.64 (m, 1H), 5.01-4.93 (m, 1H), 3.62-3.46 (m, 3H), 3.40-3.35 (m, 2H), 3.20-2.90 (m, 3H), 1.90-1.97 (m, 2H); (ESI-MS): (+ve mode) 483.10 (M+H)⁺.(100%); HPLC: 98.02%.

Example:26


- 15 **Compound 26:** 1-(5-(4-amino-3-(4-(5-(pyridin-2-yl)-1,3,4-oxadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



¹H NMR: (DMSO-d₆, 400 MHz): δ 8.84-8.83 (d, 1H, *J* = 4.0 Hz), 8.31-8.26 (m, 4H), 8.12-8.08 (m, 1H), 7.95-7.93 (d, 2H, *J* = 8.0 Hz), 7.69-7.66 (m, 1H), 6.65-6.59 (m, 1H), 6.16-6.12 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 16.8 Hz), 5.65-5.68 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 10.4 Hz), 5.47-5.44 (m, 1H), 3.81-3.77 (m, 1H), 3.65-3.61 (m, 1H), 3.55-3.50 (m, 1H), 3.39-3.33 (m, 1H), 3.12-2.90 (m, 2H), 2.42-2.31 (m, 2H), 2.09-2.07 (m, 2H); **ESI-MS:** (+ve mode) 520.20 (M+H)⁺ (85 %); **UPLC:** 95.96 %.

Example:27


Compound 27: 1-(5-(4-amino-6-bromo-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



¹H NMR (DMSO-d₆, 400 MHz) δ ppm: 8.13 (s, 1H), 7.46-7.41 (m, 4H), 7.21-7.19 (m, 1H), 7.15-7.13 (m, 2H), 7.11-7.09 (m, 2H), 6.65-6.54 (m, 1H), 6.16 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 16.8 Hz), 5.68 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 10.4 Hz), 5.35-5.33 (m, 1H), 3.52-3.50 (m, 1H), 3.38-3.34 (m, 1H), 3.33-3.31 (m, 1H), 3.12-2.83 (m, 1H), 2.81-2.80 (m, 1H), 2.67-2.65 (m, 2H), 2.37-2.35 (m, 1H), 2.33-2.00 (m, 2H); **(ESI-MS):** (+ve mode) 546.15 (M+H)⁺ (100%); **UPLC:** 95.60%.

Example:28


Compound 28: 2-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydrocyclopenta[c]pyrrole-2-carbonyl)-3-cyclopropylacrylonitrile



¹H NMR: (CDCl₃-d₁, 400 MHz): δ 8.38 (s, 1H), 7.67-7.65 (m, 2H), 7.42-7.32 (m, 2H), 7.19-7.15 (m, 3H), 7.11-7.09 (m, 2H), 6.87-6.84 (d, 1H, *J* = 11.6 Hz), 5.59-5.53 (m, 1H), 5.41 (s, 2H), 4.05-3.88 (m, 2H), 3.68-3.54 (m, 2H), 3.23-3.12 (m, 2H), 2.62-2.52 (m, 2H), 2.17-2.08 (m, 2H), 1.44-1.26 (m, 2H), 0.98-0.93 (m, 2H), 0.89-0.87 (m, 1H); **ESI-MS:** (+ve mode) 532.25 (M+H)⁺ (100 %); **UPLC:** 95.05 %.

Example:29


Compound 29: 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-2-methoxy-N-(4-methylpyridin-2-yl)benzamide



¹H NMR: (DMSO, 400 MHz): δ 10.42 (s, 1H), 8.26 (s, 1H), 8.23-8.21 (m, 1H), 8.16-8.10 (m, 1H), 8.07 (d, 1H, *J* = 8.0 Hz), 7.45-7.42 (m, 2H), 7.03-7.02 (m, 1H), 6.66-6.59 (m, 1H), 6.15 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 16.4 Hz), 5.67 (dd, 1H, *J*₁ = 2.8 Hz, *J*₂ = 10.4 Hz), 5.46-5.42 (m, 1H), 4.08 (s, 3H), 3.85-3.75 (m, 1H), 3.70-3.57 (m, 1H), 3.56-3.45 (m, 2H), 3.15-2.90 (m, 2H), 2.45-2.38 (m, 5H), 2.18-2.06 (m, 2H); **ESI-MS:** (+ve mode) 539.2 (M+H)⁺ (100 %); **UPLC:** 96.93 %.

Example:30

Compound 30: 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)benzamide




5

¹H NMR: (CDCl₃, 400 MHz): δ 8.59 (bs, 1H), 8.39 (s, 1H), 8.26 (s, 2H), 8.19 (d, 1H, *J* = 5.2 Hz), 8.11 (d, 2H, *J* = 8.0 Hz), 7.87 (d, 2H, *J* = 8.0 Hz), 6.95 (d, 1H, *J* = 5.2 Hz), 6.48 (dd, 1H, *J*₁ = 10.0 Hz, *J*₂ = 16.8 Hz), 6.40 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 16.8 Hz), 5.70 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 12.4 Hz), 5.60-5.356 (m, 1H), 5.44 (bs, 2H), 3.88-3.83 (m, 2H), 3.56-3.46 (m, 2H), 3.23-3.21 (m, 1H), 3.14-3.12 (m, 1H), 2.59-2.54 (m, 2H), 2.43 (s, 3H), 2.20-2.09 (m, 2H); **ESI-MS:** (+ve mode) 509.1 (M+H)⁺ (100 %); **HPLC:** 96.67 %.

10

Example:31


Compound 31: 1-(5-(4-amino-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



¹H NMR (DMSO-d₆, 400 MHz) δ ppm: 8.16 (s, 1H), 7.45-7.41 (m, 3H), 7.15-7.13 (m, 2H), 6.70-6.66 (m, 5H), 6.86-6.59 (m, 1H), 6.16 (dd, 1H, $J_1 = 2.8$ Hz, $J_2 = 10.4$ Hz), 5.68 (dd, 1H, $J_1 = 2.4$ Hz, $J_2 = 10.4$ Hz), 5.30-5.24 (m, 1H), 3.80-3.78 (m, 1H), 3.66-3.51 (m, 2H), 3.51-3.35 (m, 1H), 3.10-2.95 (m, 1H), 2.37-2.34 (m, 1H), 2.32-2.25 (m, 3H), 2.07-2.05 (m, 2H); (ESI-MS): (+ve mode) 466.05 (M+H)⁺ (100%); UPLC: 97.65%.

Example:32


Compound 32: N-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)phenyl)picolinamide



¹H NMR: (DMSO-d₆, 400 MHz): δ 10.82 (s, 1H), 8.77-8.76 (m, 1H), 8.23 (s, 1H), 8.18-8.12 (m, 1H), 8.11-8.09 (m, 3H), 7.71-7.65 (m, 3H), 6.65-6.59 (m, 1H), 6.16-6.11 (m, 1H), 5.68-5.65 (m, 1H), 5.43-5.41 (m, 1H), 3.82-3.79 (m, 1H), 3.76-3.73 (m, 1H), 3.50-3.48 (m, 1H), 3.37-3.35 (m, 1H), 3.23-3.19 (m, 2H), 2.35-2.32 (m, 2H), 2.08-2.04 (m, 2H); ESI-MS: (+ve mode) 495.15 (M+H)⁺ (100 %); HPLC: 98.31 %.

Example:33


Compound 33: 6-amino-7-(4-phenoxyphenyl)-9-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)-7H-purin-8(9H)-one



¹H NMR: (DMSO, 400 MHz): δ 8.12 (s, 1H), 7.45-7.40 (m, 4H), 7.21-7.19 (m, 1H), 7.17-7.13 (m, 4H), 6.94-6.87 (m, 1H), 6.21-6.12 (m, 2H), 5.71 (m, 1H), 4.97 (m, 1H), 3.34-3.33 (m, 1H), 3.00-2.96 (m, 4H), 2.61-2.59 (m, 2H), 1.86-1.81 (m, 2H);
 5 ESI-MS: (+ve mode) 519.15 (M+H)⁺ (100%); 541.35 (M+Na)⁺ (10%);
 UPLC: 95.21 %.

Example:34


Compound 34: 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(benzo[d]thiazol-2-yl)benzamide
 10



¹H NMR: (DMSO-d₆, 400 MHz): δ 13.0 (s, 1H), 8.32-8.30 (m, 2H), 8.26 (s, 1H), 8.19-8.17 (m, 1H), 7.86-7.84 (m, 2H), 7.82-7.80 (m, 1H), 7.49-7.47 (m, 1H), 7.36-7.34 (m, 1H), 6.66-6.59 (m, 1H), 6.17-6.12 (m, 1H), 5.68-5.65 (m, 1H), 5.49-5.47 (m, 1H), 3.83-3.81 (m, 1H), 3.76-3.73 (m, 1H), 3.09-3.06 (m, 1H), 2.45-2.44 (m, 2H), 2.37-2.35 (m, 2H); ESI-MS: (+ve mode) 551.78 (M+H)⁺ (100 %); HPLC: 97.74 %.
 15

Example:35


Compound 35: N-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)phenyl)pyrazine-2-carboxamide



¹H NMR: (DMSO-d₆, 400 MHz): δ 10.92 (s, 1H), 9.32 (s, 1H), 8.95 (s, 1H), 8.84-8.83 (m, 1H), 8.23 (s, 1H), 8.11-8.09 (d, 2H, *J* = 8.0 Hz), 7.69-7.67 (d, 2H, *J* = 8.0 Hz), 6.65-6.59 (m, 1H), 6.16-6.11 (dd, 1H, *J*₁ = 2.8 Hz, *J*₂ = 16.8 Hz), 5.68-5.43 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 10.4 Hz), 5.43-5.40 (m, 1H), 3.81-3.76 (m, 1H), 3.66-3.61 (m, 1H), 3.54-3.50 (m, 1H), 3.38-3.34 (m, 1H), 3.10-3.08 (m, 1H), 3.00-2.98 (m, 1H), 2.37-2.32 (m, 2H), 2.08-2.03 (m, 2H); **ESI-MS:** (+ve mode) 496.15 (M+H)⁺ (100 %);
 10 **UPLC:** 95.55 %.

Example:36

Compound 36: 1-(5-(4-amino-3-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one




15

¹H NMR: (DMSO-d₆, 400 MHz): δ 8.19 (s, 1H), 7.21-7.19 (m, 1H), 7.17-7.14 (m, 1H), 7.07-7.05 (m, 1H), 6.59-6.55 (m, 1H), 6.15-6.12 (m, 1H), 5.67-5.64 (m, 1H), 5.40-5.32 (m, 1H), 3.81-3.79 (m, 1H), 3.78-3.75 (m, 1H), 3.59-3.57 (m, 1H), 3.51-

3.48 (m, 1H), 3.03-3.00 (m, 1H), 2.97-2.93 (m, 1H), 2.36-2.82 (m, 2H), 2.02-2.00 (m, 2H); **ESI-MS**: (+ve mode) 455.78 (M+H)⁺ (100 %); **HPLC**: 96.22 %.

Example:37


- 5 **Compound 37**: 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyrazin-2-yl)benzamide



¹H NMR: (DMSO-d₆, 400 MHz): δ 11.22 (s, 1H), 9.45 (s, 1H), 8.50-8.49 (m, 1H), 8.44-8.43 (m, 1H), 8.23-8.22 (m, 2H), 7.83-7.81 (m, 2H), 6.66-6.59 (m, 1H), 6.16-6.12 (m, 1H), 5.68-5.65 (m, 1H), 5.47-5.44 (m, 1H), 3.81-3.80 (m, 1H), 3.79-3.76 (m, 1H), 3.54-3.53 (m, 1H), 3.39-3.38 (m, 2H), 3.08-3.01 (m, 2H), 2.39-2.31 (m, 2H), 2.09-2.06 (m, 2H); **ESI-MS**: (+ve mode) 496.25 (M+H)⁺ (100 %); **HPLC**: 96.38 %.

Example:38


- 15 **Compound 38**: N-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)phenyl)benzamide



¹H NMR: (DMSO-*d*₆, 400 MHz): δ 10.44 (s, 1H), 8.24 (s, 1H), 7.98 (d, 4H, *J* = 8.0 Hz), 7.65 (d, 2H, *J* = 8.4 Hz), 7.60-7.53 (m, 3H), 6.62 (dd, 1H, *J*₁ = 16.8 Hz, *J*₂ = 10.2 Hz), 6.14 (dd, 1H, *J*₁ = 16.8 Hz, *J*₂ = 2.4 Hz), 5.67 (dd, 1H, *J*₁ = 10.2 Hz, *J*₂ = 2.4 Hz), 5.42-5.38 (m, 1H), 3.78-3.75 (m, 1H), 3.66-3.60 (m, 1H), 3.55-3.50 (m, 1H),
 5 3.37-3.33 (m, 1H), 3.10-3.06 (m, 1H), 3.01-2.98 (m, 1H), 2.37-2.32 (m, 2H), 2.08-2.05 (m, 2H); **ESI-MS:** (+ve mode) 494.1 (M+H)⁺ (100 %); **UPLC:** 96.83 %.

Example:39


Compound 39: 1-(5-(4-amino-3-(4-(5-(pyridin-2-yl)-1,3,4-thiadiazol-2-yl)phenyl)-1H-pyrazolo [3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



¹H NMR: (DMSO-*d*₆, 400 MHz): δ 8.74-8.76 (m, 1H), 8.33-8.35 (m, 1H), 8.22-8.26 (m, 3H), 8.08-8.15 (m, 1H), 7.87-7.89 (m, 2H), 7.62-7.64 (m, 1H), 6.59-6.66 (m, 1H),
 15 6.12-6.17 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 16.8 Hz), 5.65-5.68 (m, 1H), 5.41-5.48 (m, 1H), 3.75-3.90 (m, 1H), 3.58-3.68 (m, 1H), 3.55-3.58 (m, 1H), 3.35-3.37 (m, 1H), 2.90-3.10 (m, 2H), 2.35-2.37 (m, 2H), 2.07-2.08 (m, 2H); **ESI-MS:** (+ve mode) 536.05 (M+H)⁺ (100 %); **UPLC:** 97.81 %.

20 Example: 40


Compound 40: 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide



¹H NMR: (CDCl₃, 400 MHz): δ 9.02 (bs, 1H), 8.73 (s, 1H), 8.51 (d, 1H, *J* = 5.2 Hz), 8.39 (s, 1H), 8.14 (d, 2H, *J* = 8.4 Hz), 7.89 (d, 2H, *J* = 8.4 Hz), 7.33 (dd, 1H, *J*₁ = 0.8 Hz, *J*₂ = 5.2 Hz), 6.48 (dd, 1H, *J*₁ = 10.0 Hz, *J*₂ = 16.8 Hz), 6.40 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 16.8 Hz), 5.69 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 9.6 Hz), 5.60-5.56 (m, 3H), 3.87-3.83 (m, 2H), 3.56-3.47 (m, 2H), 3.25-3.20 (m, 1H), 3.18-3.09 (m, 1H), 2.57-2.53 (m, 2H), 2.18-2.15 (m, 2H); ESI-MS: (+ve mode) 563.3 (M+H)⁺ (100 %),; HPLC: 99.55 %.

Example:41

10 **Compound 41:** (1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one




¹H NMR: (CDCl₃, 400 MHz): δ 8.47 (d, 2H, *J* = 8.4 Hz), 8.32 (s, 2H), 7.75 (d, 2H, *J* = 8.4 Hz), 7.53 (d, 1H, *J* = 6.4 Hz), 6.52-6.45 (m, 2H), 6.42 (dd, 1H, *J*₁ = 2.4 Hz, *J*₂ = 16.8 Hz), 5.72 (dd, 1H, *J*₁ = 2.8 Hz, *J*₂ = 10.0 Hz), 5.59-5.55 (m, 1H), 3.91 (s, 3H), 3.87-3.83 (m, 2H), 3.56-3.47 (m, 2H), 3.26-3.21 (m, 1H), 3.16-3.11 (m, 1H), 2.58-

2.53 (m, 2H), 2.36 (s, 3H), 2.19-2.15 (m, 2H) ; **ESI-MS**: (+ve mode) 523.2 (M+H)⁺ (100 %),; **HPLC**: 98.58 %.

Example:42


- 5 **Compound 42**: 6-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)nicotinamide



- 10 **¹H NMR**: (DMSO-d₆, 400 MHz): δ 10.77 (s, 1H), 8.26-8.25 (m, 2H), 8.19-8.17 (m, 2H), 8.08 (m, 1H), 7.80-7.78 (m, 2H), 7.03-7.02 (m, 1H), 6.66-6.59 (m, 1H), 6.16-6.12 (m, 1H), 5.68-5.65 (m, 1H), 5.48-5.43 (m, 1H), 3.81-3.79 (m, 1H), 3.77-3.74 (m, 1H), 3.61-3.58 (m, 1H), 3.22-3.18 (m, 2H), 3.13-3.07 (m, 2H), 2.35 (s, 3H), 2.08-2.06 (m, 2H); **ESI-MS**: (+ve mode) 509.35 (M)⁺ (100 %); **HPLC**: 97.99 %.

Example:43


- 15 **Compound 43**: 1-(5-(4-amino-3-(4-(pyridin-2-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one



¹H NMR: (DMSO-d₆, 400 MHz): δ 8.24 (s, 1H), 8.20-8.18 (dd, 1H, $J_1 = 8.0$ Hz, $J_2 = 8.0$ Hz), 7.90-7.86 (m, 1H), 7.71-7.69 (d, 2H, $J = 8.0$ Hz), 7.29-7.27 (d, 2H, $J = 16$ Hz), 7.18-7.15 (m, 1H), 7.11-7.09 (d, 1H, $J = 8.0$ Hz), 6.65-6.58 (m, 1H), 6.16-6.11 (dd, 1H, $J_1 = 4.0$ Hz, $J_2 = 8.0$ Hz), 5.68-5.65 (dd, 1H, $J_1 = 2.4$ Hz, $J_2 = 10$ Hz), 5.44-5.40 (m, 1H), 3.80-3.76 (m, 1H), 3.65-3.50 (m, 3H), 3.08-3.07 (m, 1H), 3.00-2.97 (m, 1H), 2.38-2.32 (m, 2H), 2.08-2.03 (m, 2H); **ESI-MS:** (+ve mode) 468.00 (M+H)⁺ (100 %); **UPLC:** 95.99 %.

Example:44


10 **Compound 44:** (E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl) octahydrocyclopenta [c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyridin-2-yl) benzamide



¹H NMR (DMSO-d₆, 400 MHz) δ ppm: 10.89 (s, 1H), 9.83-9.81 (m, 1H), 8.35-8.29 (m, 1H), 8.28-8.23 (m, 1H), 8.22-8.20 (m, 3H), 7.89-7.81 (m, 1H), 7.80-7.78 (m, 1H), 6.72-6.70 (m, 1H), 6.66-6.64 (m, 1H), 5.53-5.50 (m, 1H), 3.91-3.89 (m, 2H), 2.79 (d, 6H, $J = 4.4$ Hz); **(ESI-MS):** (+ve mode) 552.40 (M+H)⁺ (100%); **UPLC:** 98.02%.

Example:45


20 **Compound 45:** (E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl) octahydrocyclopenta [c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl) benzamide



¹H NMR: (D₂O, 400 MHz): δ 8.47 (s, 1H), 8.34 (d, 1H, *J* = 6.4 Hz), 8.23 (d, 2H, *J* = 8.4 Hz), 7.96 (d, 2H, *J* = 8.4 Hz), 7.59-7.56 (m, 2H), 6.83-6.70 (m, 2H), 5.64-5.51 (m, 1H), 4.01-3.95 (m, 3H), 3.83 (dd, 1H, *J*₁ = 8.4 Hz, *J*₂ = 13.2 Hz), 3.67 (dd, 1H, *J*₁ = 4.8 Hz, *J*₂ = 11.2 Hz), 3.55 (dd, 1H, *J*₁ = 4.8 Hz, *J*₂ = 13.2 Hz), 3.27-3.23 (m, 1H), 3.21-3.18 (m, 1H), 2.91 (s, 6H), 2.65 (s, 3H), 2.54-2.47 (m, 2H), 2.30-2.24 (m, 2H); **ESI-MS:** (+ve mode) 566.3 (M+H)⁺ (100 %); **HPLC:** 96.24 %.

Example:46

10 **Compound 46:** 1-(5-(4-amino-3-(4-(5-(pyrazin-2-yl)-1,3,4-thiadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one




¹H NMR: (DMSO-d₆, 400 MHz): δ 9.53 (s, 1H), 8.87-8.85 (m, 2H), 8.27-8.25 (m, 3H), 7.90-7.88 (d, 2H, *J* = 8.0 Hz), 6.66-6.59 (m, 1H), 6.17-6.12 (m, 1H), 5.68-5.65 (m, 1H), 5.47-5.42 (m, 1H), 3.80-3.78 (m, 1H), 3.56-3.52 (m, 1H), 3.37-3.33 (m, 1H),

3.10-2.90 (m, 3H), 2.39-2.32 (m, 2H), 2.10-2.07 (m, 2H); **ESI-MS**: (+ve mode) 537.20 (M+H)⁺ (100 %); **HPLC**: 97.71 %.

Example:47

- 5 **Compound 47**: (E)-1-(5-(4-amino-3-(4-(pyridin-2-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one




10 **¹H NMR**: (DMSO-d₆, 400 MHz): δ 8.24 (s, 1H), 8.21-8.19 (m, 1H), 7.90-7.86 (m, 1H), 7.71-7.69 (m, 2H), 7.30-7.28 (m, 2H), 7.19-7.15 (m, 1H), 7.12-7.10 (d, 1H, *J* = 8.0 Hz), 6.66-6.60 (m, 1H), 6.42-6.38 (d, 1H, *J* = 16 Hz), 5.44-5.41 (m, 1H), 3.77-3.74 (m, 1H), 3.62-3.59 (m, 1H), 3.52-3.48 (m, 1H), 3.37-3.36 (m, 1H), 3.09-2.99 (m, 4H), 2.36-2.31 (m, 2H), 2.15 (s, 6H), 2.08-2.04 (m, 2H); **ESI-MS**: (+ve mode) 525.45 (M+H)⁺ (100 %); **HPLC**: 96.91 %.

15

Example:48


Compound 48: (E)-1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one



¹H NMR: (CD₃OD, 400 MHz): δ 8.43 (s, 1H), 8.09 (d, 2H, *J* = 6.4 Hz), 7.45-7.41 (m, 2H), 7.22-7.16 (m, 3H), 7.13-7.11 (m, 2H), 6.87 (d, 1H, *J* = 15.2 Hz), 6.77 (dd, 1H, *J*₁ = 6.8 Hz, *J*₂ = 14.0 Hz), 5.69-5.64 (m, 1H), 3.98 (d, 2H, *J* = 6.8 Hz), 3.97-3.92 (m, 1H), 3.83-3.78 (m, 1H), 3.67-3.64 (m, 4H), 3.54 (dd, 1H, *J*₁ = 4.8 Hz, *J*₂ = 13.2 Hz), 3.32-3.28 (m, 1H), 3.21-3.17 (m, 1H), 2.93 (s, 6H), 2.55-2.50 (m, 2H), 2.24-2.19 (m, 2H); **ESI-MS**: (+ve mode) 524.3 (M+H)⁺ (100 %),; **HPLC**: 97.39 %.

Example:49

10 **Compound 49**: (E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyrazin-2-yl)benzamide




¹H NMR: (DMSO-d₆, 400 MHz): δ 11.25 (s, 1H), 9.45 (s, 1H), 8.51-8.50 (m, 1H), 8.44 (m, 1H), 8.26 (s, 1H), 8.23-8.21 (m, 2H), 7.83-7.81 (m, 2H), 6.64-6.61 (m, 1H), 6.43-6.39 (m, 1H), 5.48-5.45 (m, 1H), 3.81-3.78 (m, 1H), 3.76-3.72 (m, 1H), 3.68-3.62 (m, 1H), 3.20-3.18 (m, 1H), 3.04-3.02 (m, 3H), 2.99-2.97 (m, 1H), 2.37-2.15 (m,

2H), 2.15 (m, 6H), 2.08 (m, 2H); **ESI-MS**: (+ve mode) 553.45 (M+H)⁺ (100.%); **HPLC**: 95.44 %.

Example:50


- 5 **Compound 50**: (Z)-methyl 4-(4-amino-1-(2-((E)-4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl) benzimidate



- ¹H NMR**: (CD₃OD, 400 MHz): δ 8.73 (d, 1H, *J* = 6.4 Hz), 8.47 (s, 1H), 8.31 (d, 2H, *J* = 8.4 Hz), 8.13 (s, 1H), 7.98 (d, 2H, *J* = 8.4 Hz), 7.77-7.76 (m, 1H), 6.87 (d, 1H, *J* = 15.2 Hz), 6.78 (dd, 1H, *J*₁ = 6.8 Hz, *J*₂ = 13.6), 5.72-5.68 (m, 1H), 4.31 (s, 3H), 3.98 (d, 2H, *J* = 7.2 Hz), 3.96-3.93 (m, 1H), 3.81 (dd, 1H, *J*₁ = 8.0 Hz, *J*₂ = 12.8 Hz), 3.68 (dd, 1H, *J*₁ = 4.8 Hz, *J*₂ = 11.2 Hz), 3.53 (dd, 1H, *J*₁ = 4.8 Hz, *J*₂ = 13.2 Hz), 3.31-3.26 (m, 2H), 2.70 (s, 3H), 2.57-2.52 (m, 2H), 2.26-2.24 (m, 2H); **ESI-MS**: (+ve mode) 580.5 (M+H)⁺ (100 %);, **HPLC**: 96.62 %.

Example:51


- Compound 51**: (E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta [c] pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-(trifluoromethyl)pyridine-2-yl)benzamide



¹H NMR: (CD₃OD, 400 MHz): δ 8.68 (d, 1H, *J* = 5.6 Hz), 8.55 (s, 1H), 8.48 (s, 1H), 8.30 (d, 2H, *J* = 8.4 Hz), 7.94 (d, 2H, *J* = 8.4 Hz), 7.65 (dd, 1H, *J*₁ = 1.2 Hz, *J*₂ = 5.6 Hz), 6.9 (d, 1H, *J* = 15.2 Hz), 6.80-6.75 (m, 1H), 5.69-5.66 (m, 1H), 4.09 (d, 2H, *J* = 7.2 Hz), 4.00-3.94 (m, 1H), 3.81 (dd, 1H, *J*₁ = 8.0 Hz, *J*₂ = 12.8 Hz), 3.69 (dd, 1H, *J*₁ = 4.4 Hz, *J*₂ = 11.2 Hz), 3.53 (dd, 1H, *J*₁ = 4.4 Hz, *J*₂ = 13.2 Hz), 3.32-3.27 (m, 2H), 2.93 (s, 6H), 2.57-2.52 (m, 2H), 2.28-2.23 (m, 2H); **ESI-MS**: (+ve mode) 620.4 (M+H)⁺ (100 %); **HPLC**: 97.87 %.

10 Example:52

Compound 52: 1-(5-(4-amino-3-(4-(pyridin-3-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one




¹H NMR: (CDCl₃-d₁, 400 MHz): δ 8.50-8.46 (m, 2H), 8.35 (s, 1H), 7.70-7.68 (d, 2H, *J* = 8.0 Hz), 7.43-7.734 (m, 2H), 7.21-7.19 (d, 2H, *J* = 8.0 Hz), 6.47-6.39 (m, 2H), 5.73-5.70 (m, 1H), 5.59-5.55 (m, 1H), 3.88-3.83 (m, 2H), 3.61-3.47 (m, 4H), 3.23-3.13 (m, 2H), 2.57-2.52 (m, 2H), 2.18-2.16 (m, 2H); **ESI-MS**: (+ve mode) 468.15 (M+H)⁺ (100 %); **HPLC**: 95.64 %.

Example:53

Compound 53: (E)-1-(5-(4-amino-3-(4-(pyridin-3-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-


5 1-one



¹H NMR: (DMSO-d₆, 400 MHz): δ 8.47-8.48 (m, 1H), 8.41-8.39 (m, 1H), 8.23 (s, 1H), 7.70-7.67 (m, 2H), 7.58-7.55 (m, 1H), 7.48-7.45 (m, 1H), 7.23-7.21 (m, 2H), 6.64-6.59 (m, 1H), 6.42-6.38 (m, 1H), 5.43-5.40 (m, 1H), 3.77-3.74 (m, 1H), 3.64-3.61 (m, 1H), 3.52-3.49 (m, 1H), 3.09-2.96 (m, 5H), 2.35-2.30 (m, 2H), 2.15 (s, 6H), 2.06-2.04 (m, 2H); **ESI-MS:** (+ve mode) 525.45 (M+H)⁺ (100 %); **HPLC:** 95.44 %.

Example:54


Compound 54: 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-methyl-N-(4-methylpyridin-2-yl)benzamide



¹H NMR: (DMSO-d₆, 400 MHz): δ 8.23 (s, 1H), 8.21-8.20 (m, 1H), 7.58 (d, 1H, *J* = 8.4 Hz), 7.12-7.10 (m, 1H), 7.04-7.02 (m, 1H), 6.65-6.58 (m, 1H), 6.16-6.11 (m, 1H), 5.40-5.38 (m, 1H), 3.77-3.75 (m, 1H), 3.62-3.59 (m, 1H), 3.53-3.49 (m, 1H), 3.43 (s, 3H), 3.11-3.08 (m, 1H), 2.98-2.96 (m, 1H), 2.34-2.29 (m, 3H), 2.21 (s, 3H), 2.06-2.03 (m, 2H); ESI-MS: (+ve mode) 523.35 (M+H)⁺ (100 %); HPLC: 98.29 %.

Example:55


Compound 55: (E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl) octahydrocyclopenta [c] pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-methyl-N-(4-methylpyridin-2-yl)benzamide



¹H NMR: (DMSO-d₆, 400 MHz): δ 8.22 (s, 1H), 8.20 (d, 1H, *J* = 8.4 Hz), 7.57-7.55 (m, 2H), 7.43-7.41 (m, 2H), 7.11-7.09 (m, 1H), 7.03-7.02 (m, 1H), 6.63-6.59 (m, 1H), 6.41 (m, 1H), 5.39 (m, 1H), 3.79-3.72 (m, 1H), 3.59-3.57 (m, 2H), 3.43-3.42 (m, 2H), 3.08 (s, 3H), 3.05-3.03 (m, 2H), 2.32-2.30 (m, 2H), 2.16-2.09 (m, 3H), 2.09-2.06 (m, 3H), 2.04-2.02 (m, 6H); ESI-MS: (+ve mode) 580.55 (M+H)⁺ (100 %); HPLC: 96.27 %.

Example:56


Compound 56: (E)-1-(5-(4-amino-3-(4-(5-(pyrazin-2-yl)-1,3,4-thiadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one



¹H NMR: (DMSO-d₆, 400 MHz): δ 9.54 (s, 1H), 8.88-8.86 (m, 2H), 8.27-8.25 (m, 3H), 7.90-7.88 (d, 2H, *J* = 8.0 Hz), 6.66-6.59 (m, 1H), 6.43-6.39 (d, 2H, *J* = 16 Hz), 5.47-5.43 (m, 1H), 3.80-3.75 (m, 1H), 3.65-3.60 (m, 1H), 3.53-3.51 (m, 1H), 3.83-3.33 (m, 1H), 3.09-3.00 (m, 4H), 2.40-2.32 (m, 2H), 2.15 (s, 6H), 2.09-2.07 (m, 2H); **ESI-MS:** (+ve mode) 594.40 (M+H)⁺ (100 %); **HPLC:** 97.57 %.

Example:57

Compound 57: (E)-1-(5-(4-amino-3-(4-(5-(pyridin-2-yl)-1,3,4-oxadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one










¹H NMR: (DMSO-d₆, 400 MHz): δ 8.85-8.84 (m, 1H), 8.32-8.27 (m, 4H), 8.13-8.09 (m, 1H), 7.95-7.93 (m, 2H), 7.69-7.67 (m, 1H), 6.50-6.61 (m, 1H), 6.43-6.39 (m, 1H), 5.47-5.44 (m, 1H), 3.77-3.75 (m, 1H), 3.62-3.60 (m, 1H), 3.52-3.50 (m, 1H), 3.17-3.03 (m, 5H), 2.37-2.32 (m, 2H), 2.15 (s, 6H), 2.12-2.08 (m, 2H); **ESI-MS:** (+ve mode) 577.55 (M+H)⁺ (100 %); **HPLC:** 99.24 %.






Using the above procedures, following compounds (**Table-2**) can be prepared, using different boronic acids and finally reacting with optionally substituted acid chlorides.






5






Table-2:






Compd	Structures	IUPAC Names
58		1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)prop-2-en-1-one
59		2-acryloyl-5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3-dihydrocyclopenta[c]pyrrole-4,6(1H,5H)-dione

60		2-acryloyl-5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrole-1,3(2H,4H)-dione
61		1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[b]pyrrol-1(2H)-yl)prop-2-en-1-one
62		1-(4-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,5,6-tetrahydrocyclopenta[b]pyrrol-1(4H)-yl)prop-2-en-1-one
63		1-(6-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-cyclopenta[c]pyridin-2(3H)-yl)prop-2-en-1-one
64		1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-4,5,6,7-tetrahydro-1H-isoindol-2(3H)-yl)prop-2-en-1-one






65		1-(6-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3-azabicyclo[3.2.0]heptan-3-yl)prop-2-en-1-one
66		1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-6-azabicyclo[3.2.0]heptan-6-yl)prop-2-en-1-one
67		N-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydropentalen-2-yl)acrylamide
68		1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one
69		1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-isoindol-2(3H)-yl)prop-2-en-1-one







70		1-(4-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
71		1-(4-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[b]pyrrol-1(2H)-yl)prop-2-en-1-one
72		1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,5,6-tetrahydrocyclopenta[b]pyrrol-1(4H)-yl)prop-2-en-1-one
73		1-(4-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)prop-2-en-1-one
74		1-(6-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one






75		1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,4,5,6,7-hexahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one
76		1-(7-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,4,5,6,7-hexahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one
77		1-(6-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one
78		1-(7-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one
79		1-(7-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,4,5,6,7-hexahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one






80		1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)prop-2-en-1-one
81		2-acryloyl-5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3-dihydrocyclopenta[c]pyrrole-4,6(1H,5H)-dione
82		2-acryloyl-5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrole-1,3(2H,4H)-dione
83		1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[b]pyrrol-1(2H)-yl)prop-2-en-1-one
84		1-(4-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,5,6-tetrahydrocyclopenta[b]pyrrol-1(4H)-yl)prop-2-en-1-one






85		1-(6-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-cyclopenta[c]pyridin-2(3H)-yl)prop-2-en-1-one
86		1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-4,5,6,7-tetrahydro-1H-isindol-2(3H)-yl)prop-2-en-1-one
87		1-(6-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3-azabicyclo[3.2.0]heptan-3-yl)prop-2-en-1-one
88		1-(3-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-6-azabicyclo[3.2.0]heptan-6-yl)prop-2-en-1-one
89		N-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydropentalen-2-yl)acrylamide

90		1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one
91		1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-isoindol-2(3H)-yl)prop-2-en-1-one
92		1-(4-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
93		1-(4-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[b]pyrrol-1(2H)-yl)prop-2-en-1-one
94		1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,5,6-tetrahydrocyclopenta[b]pyrrol-1(4H)-yl)prop-2-en-1-one






95		1-(4-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)prop-2-en-1-one
96		1-(6-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one
97		1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,4,5,6,7-hexahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one
98		1-(7-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,4,5,6,7-hexahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one
99		1-(6-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one
100		1-(7-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one





101		1-(7-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,4,5,6,7-hexahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one
102		1-(5-(4-amino-3-(benzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)prop-2-en-1-one
103		1-(6-(4-amino-3-(benzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-cyclopenta[c]pyridin-2(3H)-yl)prop-2-en-1-one
104		1-(6-(4-amino-3-(benzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3-azabicyclo[3.2.0]heptan-3-yl)prop-2-en-1-one
105		N-(5-(4-amino-3-(benzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydropentalen-2-yl)acrylamide





106		1-(5-(4-amino-3-(benzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one
107		1-(5-(4-amino-3-(2,3-dihydrobenzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)prop-2-en-1-one
108		1-(6-(4-amino-3-(2,3-dihydrobenzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-cyclopenta[c]pyridin-2(3H)-yl)prop-2-en-1-one
109		1-(6-(4-amino-3-(2,3-dihydrobenzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3-azabicyclo[3.2.0]heptan-3-yl)prop-2-en-1-one
110		N-(5-(4-amino-3-(2,3-dihydrobenzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydropentalen-2-yl)acrylamide





111		1-(5-(4-amino-3-(2,3-dihydrobenzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one
112		N-(6-(1-(2-acryloyl-1,2,3,4,5,6-hexahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]thiazol-2-yl)acetamide
113		N-(6-(1-(2-acryloyloctahydro-1H-cyclopenta[c]pyridin-6-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]thiazol-2-yl)acetamide
114		N-(5-(3-(2-acetamidobenzo[d]thiazol-6-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydropentalen-2-yl)acrylamide
115		(E)-1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)-4-(dimethylamino)but-2-en-1-one





Blank page at the time of filing





116		(E)-1-(6-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-cyclopenta[c]pyridin-2(3H)-yl)-4-(dimethylamino)but-2-en-1-one
117		(E)-1-(6-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3-azabicyclo[3.2.0]heptan-3-yl)-4-(dimethylamino)but-2-en-1-one
118		(E)-1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)-4-(dimethylamino)but-2-en-1-one
119		(E)-1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one
120		(E)-1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)-4-(dimethylamino)but-2-en-1-one





121		(E)-1-(6-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-cyclopenta[c]pyridin-2(3H)-yl)-4-(dimethylamino)but-2-en-1-one
122		(E)-1-(6-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3-azabicyclo[3.2.0]heptan-3-yl)-4-(dimethylamino)but-2-en-1-one
123		(E)-1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)-4-(dimethylamino)but-2-en-1-one
124		1-(5-(4-amino-3-(4-benzoylphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one





125		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-phenylbenzamide
126		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-methyl-N-phenylbenzamide
127		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-2-methyl-N-phenylbenzamide
128		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-2-methoxy-N-phenylbenzamide





129		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N,2-dimethyl-N-phenylbenzamide
130		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-2-methoxy-N-methyl-N-phenylbenzamide
131		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-2-methyl-N-(pyridin-2-yl)benzamide
132		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-2-methoxy-N-(pyridin-2-yl)benzamide





133		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-2-methyl-N-(4-methylpyridin-2-yl)benzamide
134		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-2-methyl-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide
135		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-2-methoxy-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide
136		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-6-chloro-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide

137		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-6-fluoro-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide
138		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-6-bromo-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide
139		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-6-(difluoromethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide
140		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-6-(trifluoromethyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide





141		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-6-methyl-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide
142		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide
143		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-methyl-N-phenylbenzamide
144		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-(pyridin-2-yl)benzamide





145		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-methyl-N-(pyridin-2-yl)benzamide
146		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-(4-methylpyridin-2-yl)benzamide
147		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-methyl-N-(4-methylpyridin-2-yl)benzamide
148		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-methyl-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide





149		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide
150		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methyl-N-phenylbenzamide
151		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methoxy-N-phenylbenzamide
152		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N,2-dimethyl-N-phenylbenzamide





153		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methoxy-N-methyl-N-phenylbenzamide
154		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methyl-N-(pyridin-2-yl)benzamide
155		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methoxy-N-(pyridin-2-yl)benzamide
156		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N,N-dimethyl-N-(pyridin-2-yl)benzamide




157		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methoxy-N-methyl-N-(pyridin-2-yl)benzamide
158		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methyl-N-(4-methylpyridin-2-yl)benzamide
159		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methoxy-N-(4-methylpyridin-2-yl)benzamide
160		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N,2-dimethyl-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide





161		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methoxy-N-methyl-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide
162		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methyl-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide
163		4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methoxy-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide
164		4-(3-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-8-aminoimidazo[1,5-a]pyrazin-1-yl)-N-phenylbenzamide




165		4-(3-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-8-aminoimidazo[1,5-a]pyrazin-1-yl)-N-methyl-N-phenylbenzamide
166		4-(3-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-8-aminoimidazo[1,5-a]pyrazin-1-yl)-N-(pyridin-2-yl)benzamide
167		4-(3-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-8-aminoimidazo[1,5-a]pyrazin-1-yl)-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide
168		4-(3-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-8-aminoimidazo[1,5-a]pyrazin-1-yl)-2-methoxy-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide




169		4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-phenylbenzamide
170		4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-methyl-N-phenylbenzamide
171		4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-(pyridin-2-yl)benzamide
172		4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-(4-methylpyridin-2-yl)benzamide



173		4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide
174		4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-2-methoxy-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide
175		4-(1-(2-acryloyl-1,2,3,4,5,6-hexahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-phenylbenzamide
176		4-(1-(1-acryloyloctahydrocyclopenta[b]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-phenylbenzamide




177		4-(4-amino-1-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-phenylbenzamide
178		(E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-phenylbenzamide
179		(E)-4-(4-amino-1-(2-(4-(cyclopropyl(methyl)amino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-phenylbenzamide




180		(Z)-4-(4-amino-1-(2-(2-cyano-3-cyclopropylacryloyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-phenylbenzamide
181		4-(7-(2-acryloyl-1,2,3,4,5,6-hexahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide
182		4-(7-(1-acryloyloctahydrocyclopenta[b]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide
183		4-(4-amino-7-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide




184		(E)-4-(4-amino-7-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide
185		(E)-4-(4-amino-7-(2-(4-(cyclopropyl(methyl)amino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide
186		(Z)-4-(4-amino-7-(2-(2-cyano-3-cyclopropylacryloyl)octahydrocyclopenta[c]pyrrol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide





187		4-(3-(2-acryloyl-1,2,3,4,5,6-hexahydrocyclopenta[c]pyrrol-5-yl)-8-aminoimidazo[1,5-a]pyrazin-1-yl)-N-phenylbenzamide
188		4-(3-(1-acryloyloctahydrocyclopenta[b]pyrrol-5-yl)-8-aminoimidazo[1,5-a]pyrazin-1-yl)-N-phenylbenzamide
189		4-(8-amino-3-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)imidazo[1,5-a]pyrazin-1-yl)-N-phenylbenzamide





190		(E)-4-(8-amino-3-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)imidazo[1,5-a]pyrazin-1-yl)-N-phenylbenzamide
191		(E)-4-(8-amino-3-(2-(4-(cyclopropyl(methyl)amino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)imidazo[1,5-a]pyrazin-1-yl)-N-phenylbenzamide






192		(Z)-4-(8-amino-3-(2-(2-cyano-3-cyclopropylacryloyl)octahydrocyclopenta[c]pyrrol-5-yl)imidazo[1,5-a]pyrazin-1-yl)-N-phenylbenzamide
193		4-(9-(2-acryloyl-1,2,3,4,5,6-hexahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-phenylbenzamide
194		4-(9-(1-acryloyloctahydrocyclopenta[b]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-phenylbenzamide





195		4-(6-amino-8-oxo-9-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)-8,9-dihydro-7H-purin-7-yl)-N-phenylbenzamide
196		(E)-4-(6-amino-9-(2-(4-(dimethylamino)but-2-en-1-yl)octahydrocyclopenta[c]pyrrol-5-yl)-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-phenylbenzamide
197		(E)-4-(6-amino-9-(2-(4-(cyclopropyl(methyl)amino)but-2-en-1-yl)octahydrocyclopenta[c]pyrrol-5-yl)-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-phenylbenzamide





198		(Z)-4-(6-amino-9-(2-(2-cyano-3-cyclopropylacryloyl)octahydrocyclopenta[c]pyrrol-5-yl)-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-phenylbenzamide
199		1-(5-(4-amino-3-(4-(5-phenyl-1,3,4-oxadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
200		1-(5-(4-amino-6-chloro-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one






201		1-(5-(4-amino-6-(difluoromethyl)-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
202		1-(5-(8-amino-1-(4-phenoxyphenyl)imidazo[1,5-a]pyrazin-3-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
203		1-(4-phenoxyphenyl)-3-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)imidazo[1,5-a]pyrazin-8-amine
204		7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidine-6-carboxylic acid

205		4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(5-chloropyridin-2-yl)benzamide
206		1-(5-(4-amino-3-(4-(morpholinomethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
207		1-(5-(4-amino-3-(4-(piperidin-1-ylmethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
208		1-(5-(4-amino-3-(4-((3,4-dihydroisoquinolin-2(1H)-yl)methyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one

209		1-(5-(4-amino-3-(4-((tetrahydrofuran-2-yl)oxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
210		1-(5-(4-amino-3-(4-(1-morpholinoethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
211		1-(5-(4-amino-3-(4-(pyrrolidin-1-ylmethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
212		1-(5-(4-amino-3-(4-((dimethylamino)methyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
213		N-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzyl)acetamide

214		1-(5-(4-amino-3-(4-(2-morpholinoethoxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
215		2-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzyl)isoindoline-1,3-dione
216		1-(5-(4-amino-3-(4-((3,4-dihydroisoquinolin-2(1H)-yl)methyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
217		1-(5-(4-amino-3-(4-(azepan-1-ylmethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one

218		1-(5-(4-amino-3-(4-((hexahydro-1H-isoindol-2(3H)-yl)methyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
219		1-(5-(4-amino-3-(4-((thiomorpholinomethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
220		1-(5-(4-amino-3-(4-((1,1-dioxidobenzo[d]isothiazol-2(3H)-yl)methyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one
221		2-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzyl)isoindolin-1-one

222		9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-7-(benzo[d]thiazol-6-yl)-7H-purin-8(9H)-one
223		4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-methyl-N-(4-methylpyridin-2-yl)benzamide
224		9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-7-(4-((methyl(4-methylpyridin-2-yl)amino)methyl)phenyl)-7H-purin-8(9H)-one
225		4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-(pyridin-2-yl)benzamide
226		9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-7-(4-benzoylphenyl)-7H-purin-8(9H)-one

Testing of Compounds of the invention

Biological studies:

Invitro BTK inhibitory activity assay:

- 5 In vitro BTK inhibitory activity of test compounds were screened using BTK kinase assay on ADP Glo platform (Li, H., Totoritis, R.D., Lor, L.A., Schwartz, B., Caprioli, P., Jurewicz, A.J and Zhang, G., Assay Drug Dev. Technol., 2009, 7(6), 598-605). Briefly, fixed amount of recombinant purified human BTK (3 ng/reaction from SignalChem, USA) were incubated with increasing concentration of test
- 10 compounds, in 1X kinase reaction buffer (40 mM Tris-Cl, pH7.5, 20 mM MgCl₂, 2 mM MnCl₂, 0.1mg/ml BSA and 50 μ M DTT). Enzymatic reaction was initiated by adding a substrate cocktail containing 50 μ M of ATP (final concentration) and 5 μ g of polyGln4Tyr1 (Signal Chem) in total 25 μ l of reaction, in round bottom white 96 well plate. The reaction mixture was incubated at room temperature for 2 hr. After 2 hr of
- 15 incubation, 10 μ l of the reaction mix was mixed with 10 μ l of ADP Glo reagent, in another round bottom white 96 well plate and incubated at room temperature for 40 min. This was followed by addition of kinase detection reagent (20 μ l per reaction) and incubation at room temperature for 30 min. Finally, plate was read for luminescence at an integration time of 500 millisecond per well. Data were plotted
- 20 taking Enzyme with no inhibitor set as the 100% kinase activity and for dose response curve, % Kinase activity was plotted against conc on Log scale and IC₅₀ was determined by non linear curve fitting method using GraphPad Prism software 6. The invitro BTK inhibitory activity (IC₅₀) for representative compounds are listed in Table 3.

Table 3: Invitro (IC₅₀) BTK inhibitory & CYP inhibition data of representative compounds

compd No.	In-vitro BTK inhibitory activity: IC ₅₀ (nM)	CYP inhibition @ 10 μ M concn					
		1A2	2C8	2C9	2C19	2D6	3A4
1	1.3	A	A	A	A	A	A

2	1.1	A	A	A	A	A	A
3	8	A	A	B	A	A	A
4	2	A	A	A	A	A	A
5	8	A	A	A	A	A	B
6	1.7	A	A	A	A	A	A
7	6	A	A	B	A	A	B
8	4	A	A	A	B	A	A
9	28	A	B	B	A	A	B
10	25	A	B	A	A	B	B
11	13	A	A	A	A	A	A
12	37	A	B	B	A	A	B
13	1.8	A	A	A	B	A	A
14	1.4	A	A	A	A	A	A
15	18	A	A	A	A	A	B
16	1.7	A	A	A	A	A	A
17	5	A	A	A	A	A	B
18	218	A	B	B	A	A	B
19	4	A	A	A	A	A	A
20	223	A	B	B	A	A	B
21	50	B	A	A	A	A	A
22	40	A	A	A	A	B	A
23	48	A	B	B	A	A	B
24	2.5	A	A	A	A	A	A
25	5	A	A	A	A	A	A
26	1.1	A	A	A	A	A	A
27	1.6	A	A	A	A	A	A
28	16	A	A	B	A	A	B

29	1.2	A	A	A	A	A	A
30	6	A	A	A	A	A	A
31	0.6	A	A	A	A	A	A
32	4	A	A	A	A	A	B
33	1.5	A	A	A	A	A	A
34	359	B	A	A	B	A	B
35	1.2	A	A	A	A	A	A
36	1.1	A	A	A	A	A	A
37	1	A	A	A	A	A	A
38	17	A	A	A	A	B	A
39	1.4	A	A	A	A	A	A
40	6	B	A	B	A	A	A
41	3.5	A	A	A	A	A	B
42	0.1	A	A	A	A	A	A
43	2	A	A	A	A	A	A
44	1.4	A	A	A	A	A	A
45	10	B	A	A	A	A	A
46	1	A	A	A	A	A	A
47	3	A	A	A	A	A	A
48	50	B	A	A	A	A	B
49	2.5	A	A	A	A	A	A
50	19	A	A	A	A	B	B
51	100	A	B	B	A	A	B
52	162	B	A	A	B	A	B
53	10	A	A	A	A	A	B
54	0.1	A	A	A	A	A	A
55	80	A	B	B	B	A	B

56	50	B	B	B	A	A	B
57	100	A	B	B	A	A	B
A: <50% CYP inhibition @ 10 μ M concentrations; B: >50% CYP inhibition @ 10 μ M concentrations							

CYP inhibition studies:

CYP inhibition studies were performed with test compounds, at two concentrations (2 μ M and 10 μ M), using human liver microsomes (Yao, M., Zhu, M., Sinz, M.W., Zhang, H., Humphreys, W. F., Rodrigues, A. D and Dai, R., Journal of Pharmaceutical and Biomedical Analysis, 2007, 44, 211-223; Walsky, R. L and Obach, R.S., Drug Metab. Dispos., 2004, 32, 647-660). Human liver microsomes were mixed with 100 mM phosphate buffer (pH 7.4) and probe substrate and warmed to 37° in microcentrifuge tubes. Aliquots of this mixture (499 μ L) were transferred to each pre-labeled microcentrifuge tubes, followed by addition of the 1 μ L of inhibitors (test compound / CYP-specific positive control inhibitor) or control solvent (DMSO). Aliquots of this mixture (90 μ L) were transferred to each pre-labeled microcentrifuge tubes in duplicate. Final solvent concentrations were 0.2% (v/v) or less. Incubations were commenced with the addition of 10 μ L NADPH stock (assay concentration, 1 mM) to a final incubation volume of 100 μ L and incubated in shaking water bath (at 37°C and 100 rpm), for the period defined in Tables 1. Incubations were terminated by addition of 400 μ L of termination solvent (CH₃CN) containing internal standard. The terminated samples were vortex-mixed, centrifuged at 10000 rpm for 5 min and supernatant transferred into HPLC vials for LC-MS/MS analysis to monitor metabolites produced by marker CYP reactions. CYP inhibitory activity (% inhibition) of test compounds is listed in **Table 3**.

In vivo efficacy studies:

Demonstration of in vivo efficacy of test compounds in rats mice, oral routes of administration.

Animals

All the animal experiments were carried out in female rats and mice, bred in-house. Animals were housed in groups of 6 animals per cage, for a week, in order to habituate them to vivarium conditions (25 ± 4 °C, 60-65 % relative humidity, 12: 12 h light: dark cycle, with lights on at 7.30 am). All the animal experiments were carried out according to the internationally valid guidelines following approval by the 'Zydus Research Center animal ethical committee'.

SCW Arthritis

Female Sprague dawley (SD) rats were primed with an intra-articular injection of 20 μ l of peptidoglycan polysaccharide (PGPS), at 0.5 mg/ml of rhamnose in the right ankle. At 2 weeks the paw swelling were measured using a plethysmometer and rats assigned to groups based on initial paw swelling. On day 14 after model initiation, rats were dosed orally (po) with the test compounds. Following the dose administration, 1 h later, the rats received a booster dose of 0.5 ml of PGPS (0.5 mg/ml of rhamnose) via i.v. injection using their tail vein. Compounds were dosed for the following two more days and their paw volumes were measured for 3 more days. The efficacy of the compound was determined as percentage inhibition of paw swelling verses the control (untreated) group. Representative data of some of the test compounds are listed in **Table-4**.

20

Diffused large B cell lymphoma Xenografts using TMD-8 cell line

Female SCID mice were inoculated sc with 10×10^6 TMD-8 cells in 0.1 mL of PBS to the right flank. Animals were observed twice weekly for occurrence of tumor. Once the tumors became palpable (around 100 mm³) around 14 days after injection, treatment was initiated via oral route. Tumor volume was determined every alternate day using digital calipers and the tumor volume was calculated using the formula: $[\text{length}/2] \times [\text{width}^2]$. Body weights of the animals were also recorded 3 times a week as a measure of treatment related side effect. Treatment was continued

25

for two more weeks and inhibition of tumor volume compared to vehicle control was considered as efficacy endpoint. Representative data of some of the test compounds are listed in **Table-4**.

5 Protocol for Collagen Induced Arthritis (CIA) Study in Mice

CIA is a frequently used animal model of human RA (Courtenay, J.S., Dallman, M.J., Dayan, A.D., Martin, A. and Mosedale, B., Nature, 1980, 283, 666-668; Bevaart, L., Vervoordeldonk, M.J., Tak, P.P., Methods Mol. Biol., 2010, 602, 181-192). Following 7 days acclimation, mice were randomly assigned to groups according to body weight. Mice were immunized subcutaneously in the tail using bovine type II collagen mix in complete Freund's adjuvant (CFA). Twenty-one days after the first immunization, mice were given booster dose of collagen in incomplete Freund's adjuvant (IFA). Mice were monitored every other day after the booster dose for the development of arthritis. Mice were recruited for the study once clinical signs were visible. Eight animals were assigned each of three groups [vehicle, positive control and test compounds] and treatment was continued for four weeks and percentage inhibition in clinical score is recorded as per graded score. Body weights of the animals were also recorded 3 times a week as a measure of treatment related side effect, paw thickness measured twice a week and blood serum are collected at termination for cytokines profile. Representative data of some of the test compounds are listed in **Table-4**.

Table-4: Invivo efficacy data of representative compounds

Compounds (Dose:10 mpk, po)	% inhibition of paw inflammation, Day-16, in SCW Arthritis Model	% Inhibition in tumor volume, Day- 16, in TMD-8 Xenografts Model	% inhibition of clinical score, Day- 28, in CIA Arthritis Model
25	69.6±12.7	50.2±10.1	90.1±6.3

32	75.3±6.2	56.1±8.2	100±3.9
42	51.3±9.7	62.2±8.8	100±4.2
46	81.8±11.2	54.3±9.1	88±7.1
54	89.2±10.3	92.1±13.1	98±4.5
Data represent Mean ± SD, n=8 animals in each group			

The novel compounds of the present invention can be formulated into suitable pharmaceutically acceptable compositions by combining with suitable excipients by techniques and processes and concentrations as are well known.

- 5 The compounds of formula (I) or pharmaceutical compositions containing them are useful as a medicament for the inhibition of BTK activity and suitable for humans and other warm blooded animals, and may be administered either by oral, topical or parenteral administration.

Thus, a pharmaceutical composition comprising the compounds of the present invention may comprise a suitable binder, suitable bulking agent &/or diluent and any
10 other suitable agents as may be necessary. Optionally, the pharmaceutical composition may be suitably coated with suitable coating agents.

The compounds of the present invention (I) are BTK inhibitors and are useful in the treatment of disease states mediated by BTK enzyme, preferably cancer,
15 arthritis and related disorders.

In one of the embodiments the present invention of formula (I) in combination with one or more suitable pharmaceutically active agents selected from following therapeutic agents in any combination. Immunosuppressants (e.g., Methotrexate, mercaptopurine, cyclophosphamide), glucocorticoids, non-steroidal anti-
20 inflammatory drugs, Cox-2 specific inhibitors, TNF- α binding proteins (eg., Infliximab, etanercept), interferon- γ , interferon- α , interleukin-2, antihistamines, beta-agonist, anticholinergics, anti-cancer agents or their suitable pharmaceutically acceptable salts. Further examples of anticancer agents for use in combination with


BTK inhibitors include chemotherapy or a targeted therapy, alkylating agents, platinum compounds, DNA altering agents, Topoisomerase inhibitors, microtubule modifiers, antimetabolites, anticancer antibiotics, hormones, Aromatase inhibitors, antibodies, cytokines, vaccines, drug conjugates, inhibitors of mitogen-activated protein kinase signaling (ex: BAY 43-9006), Syk inhibitors, mTOR inhibitors, antibodies (Rituxan), other anticancer agents that can be employed in combination include, Vinblastin, Bleomycin, Cisplatin, Acivicin, Azacitidine, Decitabine, Doxorubicin, Enloplatin, Flurouracil, Methotrexate, Vinblastin, Vincristine and BCR/ABL antagonist

10 The quantity of active component, that is, the compounds of Formula (I) according to this invention, in the pharmaceutical composition and unit dosage form thereof may be varied or adjusted widely depending upon the particular application method, the potency of the particular compound and the desired concentration. Generally, the quantity of active component will range between 0.5% to 90% by weight of the composition.

15 While the present invention has been described in terms of its specific embodiments, certain modifications and equivalents will be apparent to those skilled in the art and are intended to be included within the scope of the present invention.

We claim:

1. Compound having the structure of general formula (I)




(I)

5 and their pharmaceutically acceptable salts, enantiomers and their diastereomers, wherein, V, W, X, Y & Z independently selected from, 'C' or 'N';

R₁ is selected from hydrogen, keto, halogen, unsubstituted or substituted groups selected from cyano, alkyl, haloalkyl, aryl, alkoxy, acyloxy, aryloxy, 10 arylalkyl, heteroaryl, heterocyclyl, heterocycloalkyl, cycloalkyl, cycloalkylalkyl, aryloxyaryl, aryloxyalkyl, aryloxyheteroaryl, groups;

R₂ is selected from the following ring system:




R_3 at each occurrence is independently selected from hydrogen, haloalkyl, C_{1-7} alkyl, C_{2-7} alkenyl, C_{2-7} alkynyl, aryl, cycloalkyl, heterocycloalkyl, cycloalkyl(C_{1-7})alkyl, heterocycloalkyl(C_{1-7})alkyl, $C(O)NH(C_{1-7})$ alkyl, $C(O)-CH=CH_2$, $C(O)-CH=CH-R_4$, $C(O)-C(CN)=CH_2$, $C(O)-C(CN)=CH-R_4$, $SO_2-NH(C_{1-7})$ alkyl, $SO_2-CH=CH_2$, $SO_2-CH=CH-R_4$ groups;

R_4 at each occurrence is selected from $-(CH_2)_n-NR_5R_6$; wherein, $n=0-7$ and each of R_5 and R_6 are independently selected from hydrogen, haloalkyl, C_{1-7} alkyl, C_{2-7} alkenyl, C_{2-7} alkynyl, aryl, cycloalkyl, carbocycle, heterocycloalkyl, cycloalkyl(C_{1-7})alkyl, heterocycloalkyl(C_{1-7})alkyl;

'U' is selected from unsubstituted or substituted groups selected from alkyl, alkenyl, alkynyl, alkoxy, acyloxy, aryl, aryloxy, arylalkyl, cycloalkyl, cycloalkylalkyl, biaryl, heteroaryl, heterocycle, heterocycloalkyl, O-aryl, O-cycloalkyl, O-heteroaryl, O-heterocycle, O-heterocycloalkyl, aryloxyaryl, aryloxyalkyl, aryloxyheteroaryl, heteroaryloxyaryl, heteroaryloxyalkyl, heteroaryloxyheteroaryl, $Ph-CO-N(R_7R_8)$, $Ph-N(R_9)-CO-R_{10}$, wherein, R_7 , R_8

and R_{10} are independently selected from hydrogen, halogen, alkyl, haloalkyl, alkoxy; aryl, cycloalkyl, heteroaryl, heterocycloalkyl; further substituted with halogen, alkyl, alkoxy, haloalkoxy groups and R_9 are independently selected from hydrogen, C_{1-7} alkyl, C_{2-7} alkenyl, C_{2-7} alkynyl.

- 5 2. The compound as claimed in claim 1, wherein R_1 is selected from hydrogen, keto, halogen, unsubstituted or substituted groups selected from cyano, alkyl, haloalkyl groups.
3. The compound as claimed in claim 1, wherein R_2 is selected from the following ring system.



10

4. The compound as claimed in claim 1, wherein R_3 is selected from $C(O)-CH=CH_2$, $C(O)-CH=CH-R_4$, $SO_2-NH(C_{1-7})alkyl$, $SO_2-CH=CH_2$, $SO_2-CH=CH-R_4$ groups, wherein R_4 is as defined earlier.
5. The compound as claimed in claim 1, wherein U is selected from unsubstituted or substituted groups selected from aryloxyaryl, aryloxyalkyl, aryloxyheteroaryl, heteroaryloxyaryl, heteroaryloxyalkyl, heteroaryloxyheteroaryl, $Ph-CO-N(R_7R_8)$, $Ph-N(R_9)-CO-R_{10}$, wherein R_7 to R_{10} are as defined earlier.
6. A compound as claimed in claim 1 selected from the group comprising of:

15

- 1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 5 1-(5-(5-((4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)methyl)pyridin-2-yl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(2-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 10 1-(5-(4-amino-3-(2-methylbenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(benzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(2,3-dihydrobenzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-
- 15 d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(dibenzo[b,d]furan-3-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- N-(6-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]thiazol-2-yl)acetamide;
- 20 1-(5-(4-amino-3-(2-methoxybenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-yn-1-one;
- 1-(5-(4-amino-3-(3-methoxy-4-phenoxyphenyl)-1H-pyrazolo[3,4-
- 25 d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyridin-2-yl)benzamide;
- 1-(5-(4-amino-3-(2-phenylbenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;

- 1-(5-(4-amino-3-(benzo[d][1,3]dioxol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(4-(5-methyl-1,3,4-oxadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 5 1-(5-(4-amino-3-(benzo[d]oxazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 3-(4-phenoxyphenyl)-1-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine;
- 1-(5-(4-amino-3-(2-phenylbenzo[d]oxazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 10 1-(5-(4-amino-3-(2-phenoxybenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(4-(1-methyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 15 1-(5-(4-amino-3-(benzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(2-phenylbenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- (E)-1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one;
- 20 one;
- 9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-7-(4-phenoxyphenyl)-5,7-dihydro-4H-purin-8(9H)-one;
- 1-(5-(4-amino-3-(4-(5-(pyridin-2-yl)-1,3,4-oxadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 25 2-en-1-one;
- 1-(5-(4-amino-6-bromo-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;

- (E)-2-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydrocyclopenta[c]pyrrole-2-carbonyl)-3-cyclopropylacrylonitrile;
 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-2-methoxy-N-(4-methylpyridin-2-yl)benzamide;
 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)benzamide;
 1-(5-(4-amino-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
 N-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)phenyl)picolinamide;
 6-amino-7-(4-phenoxyphenyl)-9-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)-7H-purin-8(9H)-one;
 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(benzo[d]thiazol-2-yl)benzamide;
 N-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)phenyl)pyrazine-2-carboxamide;
 1-(5-(4-amino-3-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyrazin-2-yl)benzamide;
 N-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)phenyl)benzamide;
 1-(5-(4-amino-3-(4-(5-(pyridin-2-yl)-1,3,4-thiadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide;

- Z)-methyl 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)benzimidate;
6-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)nicotinamide;
- 5 1-(5-(4-amino-3-(4-(pyridin-2-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
(E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyridin-2-yl)benzamide;
- 10 (E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)benzamide;
1-(5-(4-amino-3-(4-(5-(pyrazin-2-yl)-1,3,4-thiadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-
- 15 2-en-1-one;
(E)-1-(5-(4-amino-3-(4-(pyridin-2-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one;
(E)-1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-
- 20 one;
(E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyrazin-2-yl)benzamide;
- 25 (Z)-methyl 4-(4-amino-1-(2-((E)-4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)benzimidate;

(E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta
[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-(trifluoromethyl)
pyridin-2-yl)benzamide;

5 1-(5-(4-amino-3-(4-(pyridin-3-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-
yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;

(E)-1-(5-(4-amino-3-(4-(pyridin-3-yloxy)phenyl)-1H-pyrazolo[3,4-d]
pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)
but-2-en-1-one;

10 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo
[3,4-d]pyrimidin-3-yl)-N-methyl-N-(4-methylpyridin-2-yl)benzamide;

(E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta
[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-methyl-N-(4-methyl
pyridin-2-yl)benzamide;

15 (E)-1-(5-(4-amino-3-(4-(5-(pyrazin-2-yl)-1,3,4-thiadiazol-2-yl)phenyl)-1H-
pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-
(dimethylamino)but-2-en-1-one;

(E)-1-(5-(4-amino-3-(4-(5-(pyridin-2-yl)-1,3,4-oxadiazol-2-yl)phenyl)-1H-
pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-
(dimethylamino)but-2-en-1-one

20

7. A compound as claimed in claim 1 selected from the group comprising of:

1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-
yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;

25 1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-
yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;

1-(5-(5-((4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-
yl)methyl)pyridin-2-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-
one;

- 1-(5-(2-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)ethyl)hexahydropyrrolo[3,4-c]pyrrol-2(1H)-yl)prop-2-en-1-one;
1-(5-(4-amino-3-(2-methylbenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
5 1-(5-(4-amino-3-(benzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
1-(5-(4-amino-3-(2,3-dihydrobenzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
1-(5-(4-amino-3-(dibenzo[b,d]furan-3-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
10 N-(6-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]thiazol-2-yl)acetamide;
1-(5-(4-amino-3-(2-methoxybenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
15 1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
1-(5-(4-amino-3-(3-methoxy-4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyridin-2-yl)benzamide;
20 1-(5-(4-amino-3-(2-phenylbenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
1-(5-(4-amino-3-(benzo[d][1,3]dioxol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
25 1-(5-(4-amino-3-(4-(5-methyl-1,3,4-oxadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
1-(5-(4-amino-3-(benzo[d]oxazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;

- 3-(4-phenoxyphenyl)-1-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-4-amine;
- 1-(5-(4-amino-3-(2-phenylbenzo[d]oxazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 5 1-(5-(4-amino-3-(2-phenoxybenzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(4-(1-methyl-1H-pyrazol-4-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(benzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 10 1-(5-(4-amino-3-(2-phenylbenzo[d]oxazol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- (E)-1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one;
- 15 9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-7-(4-phenoxyphenyl)-5,7-dihydro-4H-purin-8(9H)-one;
- 1-(5-(4-amino-3-(4-(5-(pyridin-2-yl)-1,3,4-oxadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 20 1-(5-(4-amino-6-bromo-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- (E)-2-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydrocyclopenta[c]pyrrole-2-carbonyl)-3-cyclopropylacrylonitrile;
- 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-2-methoxy-N-(4-methylpyridin-2-yl)benzamide;
- 25 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)benzamide;
- 1-(5-(4-amino-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;

- N-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)phenyl)picolinamide;
 6-amino-7-(4-phenoxyphenyl)-9-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)-7H-purin-8(9H)-one;
- 5 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(benzo[d]thiazol-2-yl)benzamide;
 N-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)phenyl)pyrazine-2-carboxamide;
 1-(5-(4-amino-3-(2,2-difluorobenzo[d][1,3]dioxol-5-yl)-1H-pyrazolo[3,4-
- 10 d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyrazin-2-yl)benzamide;
 N-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)phenyl)benzamide;
- 15 1-(5-(4-amino-3-(4-(5-(pyridin-2-yl)-1,3,4-thiadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-(trifluoromethyl)pyridin-2-
- 20 yl)benzamide;
 Z)-methyl 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)benzimidate;
 6-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)nicotinamide;
- 25 1-(5-(4-amino-3-(4-(pyridin-2-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
 (E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyridin-2-yl)benzamide;

- (E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)benzamide;
- 5 1-(5-(4-amino-3-(4-(5-(pyrazin-2-yl)-1,3,4-thiadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- (E)-1-(5-(4-amino-3-(4-(pyridin-2-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one;
- 10 (E)-1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one;
- (E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(pyrazin-2-yl)benzamide;
- 15 (Z)-methyl 4-(4-amino-1-(2-((E)-4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-methylpyridin-2-yl)benzimidate;
- (E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(4-(trifluoromethyl)pyridine-2-yl)benzamide;
- 20 1-(5-(4-amino-3-(4-(pyridin-3-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- (E)-1-(5-(4-amino-3-(4-(pyridin-3-yloxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one;
- 25 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-methyl-N-(4-methylpyridin-2-yl)benzamide;

- (E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-methyl-N-(4-methylpyridin-2-yl)benzamide;
- (E)-1-(5-(4-amino-3-(4-(5-(pyrazin-2-yl)-1,3,4-thiadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one;
- (E)-1-(5-(4-amino-3-(4-(5-(pyridin-2-yl)-1,3,4-oxadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one;
- 1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)prop-2-en-1-one;
- 2-acryloyl-5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3-dihydrocyclopenta[c]pyrrole-4,6(1H,5H)-dione;
- 2-acryloyl-5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrole-1,3(2H,4H)-dione;
- 1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[b]pyrrol-1(2H)-yl)prop-2-en-1-one;
- 1-(4-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,5,6-tetrahydrocyclopenta[b]pyrrol-1(4H)-yl)prop-2-en-1-one;
- 1-(6-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-cyclopenta[c]pyridin-2(3H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-4,5,6,7-tetrahydro-1H-isoindol-2(3H)-yl)prop-2-en-1-one;
- 1-(6-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3-azabicyclo[3.2.0]heptan-3-yl)prop-2-en-1-one;
- 1-(3-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-6-azabicyclo[3.2.0]heptan-6-yl)prop-2-en-1-one;
- N-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydropentalen-2-yl)acrylamide;

- 1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one;
 1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-isoindol-2(3H)-yl)prop-2-en-1-one;
 5 1-(4-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
 1-(4-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[b]pyrrol-1(2H)-yl)prop-2-en-1-one;
 1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,5,6-tetrahydrocyclopenta[b]pyrrol-1(4H)-yl)prop-2-en-1-one;
 10 1-(4-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)prop-2-en-1-one;
 1-(6-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one;
 15 1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,4,5,6,7-hexahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one;
 1-(7-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,4,5,6,7-hexahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one;
 1-(6-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one;
 20 1-(7-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one;
 1-(7-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,4,5,6,7-hexahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one;
 25 1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)prop-2-en-1-one;
 2-acryloyl-5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3-dihydrocyclopenta[c]pyrrole-4,6(1H,5H)-dione;

- 2-acryloyl-5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrole-1,3(2H,4H)-dione;
 1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[b]pyrrol-1(2H)-yl)prop-2-en-1-one;
 5 1-(4-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,5,6-tetrahydrocyclopenta[b]pyrrol-1(4H)-yl)prop-2-en-1-one;
 1-(6-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-cyclopenta[c]pyridin-2(3H)-yl)prop-2-en-1-one;
 10 1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-4,5,6,7-tetrahydro-1H-isoindol-2(3H)-yl)prop-2-en-1-one;
 1-(6-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3-azabicyclo[3.2.0]heptan-3-yl)prop-2-en-1-one;
 1-(3-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-6-azabicyclo[3.2.0]heptan-6-yl)prop-2-en-1-one;
 15 N-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydropentalen-2-yl)acrylamide;
 1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one;
 1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-isoindol-2(3H)-yl)prop-2-en-1-one;
 20 1-(4-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
 1-(4-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[b]pyrrol-1(2H)-yl)prop-2-en-1-one;
 25 1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,5,6-tetrahydrocyclopenta[b]pyrrol-1(4H)-yl)prop-2-en-1-one;
 1-(4-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)prop-2-en-1-one;

- 1-(6-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,4,5,6,7-hexahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one;
- 5 1-(7-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,4,5,6,7-hexahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one;
- 1-(6-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one;
- 1-(7-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one;
- 10 1-(7-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-2,3,4,5,6,7-hexahydro-1H-cyclopenta[b]pyridin-1-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(benzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)prop-2-en-1-one;
- 15 1-(6-(4-amino-3-(benzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-cyclopenta[c]pyridin-2(3H)-yl)prop-2-en-1-one;
- 1-(6-(4-amino-3-(benzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3-azabicyclo[3.2.0]heptan-3-yl)prop-2-en-1-one;
- N-(5-(4-amino-3-(benzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydropentalen-2-yl)acrylamide;
- 20 1-(5-(4-amino-3-(benzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(2,3-dihydrobenzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)prop-2-en-
- 25 1-one;
- 1-(6-(4-amino-3-(2,3-dihydrobenzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-cyclopenta[c]pyridin-2(3H)-yl)prop-2-en-1-one

- 1-(6-(4-amino-3-(2,3-dihydrobenzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3-azabicyclo[3.2.0]heptan-3-yl)prop-2-en-1-one;
N-(5-(4-amino-3-(2,3-dihydrobenzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydropentalen-2-yl)acrylamide;
- 5 1-(5-(4-amino-3-(2,3-dihydrobenzo[b]thiophen-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)prop-2-en-1-one;
N-(6-(1-(2-acryloyl-1,2,3,4,5,6-hexahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]thiazol-2-yl)acetamide;
- 10 N-(6-(1-(2-acryloyloctahydro-1H-cyclopenta[c]pyridin-6-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzo[d]thiazol-2-yl)acetamide;
N-(5-(3-(2-acetamidobenzo[d]thiazol-6-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-1-yl)octahydropentalen-2-yl)acrylamide;
- (E)-1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)-4-(dimethylamino)but-2-en-1-one;
- 15 (E)-1-(6-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-cyclopenta[c]pyridin-2(3H)-yl)-4-(dimethylamino)but-2-en-1-one;
- (E)-1-(6-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3-azabicyclo[3.2.0]heptan-3-yl)-4-(dimethylamino)but-2-en-1-one;
- 20 (E)-1-(5-(4-amino-3-(4-phenoxyphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)-4-(dimethylamino)but-2-en-1-one;
- (E)-1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)-4-(dimethylamino)but-2-en-1-one;
- 25 one;

- (E)-1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-5,6-dihydrocyclopenta[c]pyrrol-2(1H,3H,4H)-yl)-4-(dimethylamino)but-2-en-1-one;
- 5 (E)-1-(6-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydro-1H-cyclopenta[c]pyridin-2(3H)-yl)-4-(dimethylamino)but-2-en-1-one;
- (E)-1-(6-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3-azabicyclo[3.2.0]heptan-3-yl)-4-(dimethylamino)but-2-en-1-one;
- 10 (E)-1-(5-(4-amino-3-(benzo[d]thiazol-6-yl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)-3,4,6,7-tetrahydro-1H-cyclopenta[c]pyridin-2(5H)-yl)-4-(dimethylamino)but-2-en-1-one;
- 1-(5-(4-amino-3-(4-benzoylphenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo
- 15 [3,4-d]pyrimidin-3-yl)-N-phenylbenzamide;
- 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo [3,4-d]pyrimidin-3-yl)-N-methyl-N-phenylbenzamide;
- 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo [3,4-d]pyrimidin-3-yl)-2-methyl-N-phenylbenzamide
- 20 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo [3,4-d]pyrimidin-3-yl)-2-methoxy-N-phenylbenzamide;
- 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo [3,4-d]pyrimidin-3-yl)-N,2-dimethyl-N-phenylbenzamide;
- 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo
- 25 [3,4-d]pyrimidin-3-yl)-2-methoxy-N-methyl-N-phenylbenzamide;
- 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo [3,4-d]pyrimidin-3-yl)-2-methyl-N-(pyridin-2-yl)benzamide;
- 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo [3,4-d]pyrimidin-3-yl)-2-methoxy-N-(pyridin-2-yl)benzamide;

- 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo
[3,4-d]pyrimidin-3-yl)-2-methyl-N-(4-methylpyridin-2-yl)benzamide;
4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3
,4-d]pyrimidin-3-yl)-2-methyl-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide;
5 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4
-d]pyrimidin-3-yl)-2-methoxy-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide;
4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-6-chloro-7H-
pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide;
4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-6-fluoro-7H-
10 pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide;
4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-6-bromo-7H-
pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide;
4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-6-(difluoro
methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide;
15 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-6-(trifluoro
methyl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide;
4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-6-methyl-7H-
pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide;
4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-
20 d]pyrimidin-5-yl)-N-phenylbenzamide;
4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-
d]pyrimidin-5-yl)-N-methyl-N-phenylbenzamide;
4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-
d]pyrimidin-5-yl)-N-(pyridin-2-yl)benzamide;
25 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-
d]pyrimidin-5-yl)-N-methyl-N-(pyridin-2-yl)benzamide;
4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-
d]pyrimidin-5-yl)-N-(4-methylpyridin-2-yl)benzamide;

- 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-methyl-N-(4-methylpyridin-2-yl)benzamide;
- 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-methyl-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide;
- 5 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide;
- 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methyl-N-phenylbenzamide;
- 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methoxy-N-phenylbenzamide;
- 10 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N,2-dimethyl-N-phenylbenzamide;
- 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methoxy-N-methyl-N-phenylbenzamide;
- 15 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methyl-N-(pyridin-2-yl)benzamide;
- 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methoxy-N-(pyridin-2-yl)benzamide;
- 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N,2-dimethyl-N-(pyridin-2-yl)benzamide;
- 20 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methoxy-N-methyl-N-(pyridin-2-yl)benzamide;
- 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methyl-N-(4-methylpyridin-2-yl)benzamide;
- 25 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methoxy-N-(4-methylpyridin-2-yl)benzamide;
- 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N,2-dimethyl-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide;

- 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methoxy-N-methyl-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide;
- 5 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methyl-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide;
- 4-(7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-2-methoxy-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide;
- 4-(3-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-8-aminoimidazo[1,5-a]pyrazin-1-yl)-N-phenylbenzamide;
- 10 4-(3-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-8-aminoimidazo[1,5-a]pyrazin-1-yl)-N-methyl-N-phenylbenzamide;
- 4-(3-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-8-aminoimidazo[1,5-a]pyrazin-1-yl)-N-(pyridin-2-yl)benzamide;
- 15 4-(3-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-8-aminoimidazo[1,5-a]pyrazin-1-yl)-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide;
- 4-(3-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-8-aminoimidazo[1,5-a]pyrazin-1-yl)-2-methoxy-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide;
- 4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-phenylbenzamide;
- 20 4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-methyl-N-phenylbenzamide;
- 4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-(pyridin-2-yl)benzamide;
- 25 4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-(4-methylpyridin-2-yl)benzamide;
- 4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide;

- 4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-2-methoxy-N-(4-(trifluoromethyl)pyridin-2-yl)benzamide;
- 5 4-(1-(2-acryloyl-1,2,3,4,5,6-hexahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-phenylbenzamide;
- 4-(1-(1-acryloyloctahydrocyclopenta[b]pyrrol-5-yl)-4-amino-1H-pyrazolo [3,4-d]pyrimidin-3-yl)-N-phenylbenzamide;
- 4-(4-amino-1-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-phenylbenzamide;
- 10 (E)-4-(4-amino-1-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-phenylbenzamide;
- (E)-4-(4-amino-1-(2-(4-(cyclopropyl(methyl)amino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-phenylbenzamide;
- 15 (Z)-4-(4-amino-1-(2-(2-cyano-3-cyclopropylacryloyl)octahydrocyclopenta[c]pyrrol-5-yl)-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-phenylbenzamide;
- 4-(7-(2-acryloyl-1,2,3,4,5,6-hexahydrocyclopenta[c]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide;
- 4-(7-(1-acryloyloctahydrocyclopenta[b]pyrrol-5-yl)-4-amino-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide;
- 20 4-(4-amino-7-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide;
- (E)-4-(4-amino-7-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide;
- 25 (E)-4-(4-amino-7-(2-(4-(cyclopropyl(methyl)amino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide;

- (Z)-4-(4-amino-7-(2-(2-cyano-3-cyclopropylacryloyl)octahydrocyclopenta[c]pyrrol-5-yl)-7H-pyrrolo[2,3-d]pyrimidin-5-yl)-N-phenylbenzamide;
 4-(3-(2-acryloyl-1,2,3,4,5,6-hexahydrocyclopenta[c]pyrrol-5-yl)-8-aminoimidazo[1,5-a]pyrazin-1-yl)-N-phenylbenzamide;
 5 4-(3-(1-acryloyloctahydrocyclopenta[b]pyrrol-5-yl)-8-aminoimidazo[1,5-a]pyrazin-1-yl)-N-phenylbenzamide;
 4-(8-amino-3-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)imidazo[1,5-a]pyrazin-1-yl)-N-phenylbenzamide;
 (E)-4-(8-amino-3-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)imidazo[1,5-a]pyrazin-1-yl)-N-phenylbenzamide;
 10 (E)-4-(8-amino-3-(2-(4-(cyclopropyl(methyl)amino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)imidazo[1,5-a]pyrazin-1-yl)-N-phenylbenzamide;
 (Z)-4-(8-amino-3-(2-(2-cyano-3-cyclopropylacryloyl)octahydrocyclopenta[c]pyrrol-5-yl)imidazo[1,5-a]pyrazin-1-yl)-N-phenylbenzamide;
 15 4-(9-(2-acryloyl-1,2,3,4,5,6-hexahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-phenylbenzamide;
 4-(9-(1-acryloyloctahydrocyclopenta[b]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-phenylbenzamide;
 4-(6-amino-8-oxo-9-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)-8,9-dihydro-7H-purin-7-yl)-N-phenylbenzamide;
 20 (E)-4-(6-amino-9-(2-(4-(dimethylamino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-phenylbenzamide;
 (E)-4-(6-amino-9-(2-(4-(cyclopropyl(methyl)amino)but-2-enoyl)octahydrocyclopenta[c]pyrrol-5-yl)-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-phenylbenzamide;
 25 benzamide;
 (Z)-4-(6-amino-9-(2-(2-cyano-3-cyclopropylacryloyl)octahydrocyclopenta[c]pyrrol-5-yl)-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-phenylbenzamide;
 1-(5-(4-amino-3-(4-(5-phenyl-1,3,4-oxadiazol-2-yl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;

- 1-(5-(4-amino-6-chloro-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-6-(difluoromethyl)-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidin-7-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 5 1-(5-(8-amino-1-(4-phenoxyphenyl)imidazo[1,5-a]pyrazin-3-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(4-phenoxyphenyl)-3-(2-(vinylsulfonyl)octahydrocyclopenta[c]pyrrol-5-yl)imidazo[1,5-a]pyrazin-8-amine;
- 7-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-5-(4-phenoxyphenyl)-7H-pyrrolo[2,3-d]pyrimidine-6-carboxylic acid;
- 10 4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)-N-(5-chloropyridin-2-yl)benzamide;
- 1-(5-(4-amino-3-(4-(morpholinomethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 15 1-(5-(4-amino-3-(4-(piperidin-1-ylmethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(4-((3,4-dihydroisoquinolin-2(1H)-yl)methyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 20 1-(5-(4-amino-3-(4-((tetrahydrofuran-2-yl)oxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(4-(1-morpholinoethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(4-(pyrrolidin-1-ylmethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 25 1-(5-(4-amino-3-(4-((dimethylamino)methyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- N-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzyl)acetamide;

- 1-(5-(4-amino-3-(4-(2-morpholinoethoxy)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
2-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzyl)isoindoline-1,3-dione;
- 5 1-(5-(4-amino-3-(4-((3,4-dihydroisoquinolin-2(1H)-yl)methyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(4-(azepan-1-yl)methyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 10 1-(5-(4-amino-3-(4-((hexahydro-1H-isoindol-2(3H)-yl)methyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 1-(5-(4-amino-3-(4-(thiomorpholinomethyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 15 1-(5-(4-amino-3-(4-((1,1-dioxidobenzo[d]isothiazol-2(3H)-yl)methyl)phenyl)-1H-pyrazolo[3,4-d]pyrimidin-1-yl)hexahydrocyclopenta[c]pyrrol-2(1H)-yl)prop-2-en-1-one;
- 2-(4-(1-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-4-amino-1H-pyrazolo[3,4-d]pyrimidin-3-yl)benzyl)isoindolin-1-one;
- 20 9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-7-(benzo[d]thiazol-6-yl)-7H-purin-8(9H)-one;
- 4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-methyl-N-(4-methylpyridin-2-yl)benzamide;
- 9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-7-(4-((methyl(4-methylpyridin-2-yl)amino)methyl)phenyl)-7H-purin-8(9H)-one;
- 25 4-(9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-8-oxo-8,9-dihydro-7H-purin-7-yl)-N-(pyridin-2-yl)benzamide;
- 9-(2-acryloyloctahydrocyclopenta[c]pyrrol-5-yl)-6-amino-7-(4-benzoylphenyl)-7H-purin-8(9H)-one

- 5 8. A pharmaceutical composition comprising a therapeutically effective amount of a compound of Formula (I) as claimed in any of the preceding claims and optionally one or more pharmaceutically acceptable carriers, diluents or excipients.
- 10 9. The pharmaceutical composition which is useful for treating a disease, disorder or condition, which would benefit from inhibition of tyrosine kinase (s), such as BTK activity to provide clinical benefit in diseases such as autoimmune disorders, an inflammatory disease, or an allergy disease selected from rheumatoid arthritis or lupus erythromatosis, asthma, allergic rhinitis, B-cell proliferative disorder such as B-cell lymphoma, multiple sclerosis, mantle cell lymphoma, chronic lymphocytic lymphoma / leukemia, diffuse large B-cell lymphoma, Follicular lymphoma or chronic lymphocytic leukemia, B-cell
15 prolymphocytic leukemia.
- 20 10. A method of treating a disease, disorder or condition, which would benefit from inhibition of tyrosine kinase (s), such as BTK activity comprising administering to a patient in need thereof an effective amount of a compound of Formula (I) according to any of the preceding claims or its suitable pharmaceutical composition.
- 25 11. A medicine for the treatment of a disease, disorder or condition, which would benefit from inhibition of tyrosine kinase (s), such as BTK activity, which comprises administering a therapeutically effective amount of compound of Formula (I) or its pharmaceutical composition as defined in any of the preceding claims to a patient or subject in need thereof.

12. A pharmaceutical composition comprising the compound of the present invention in combination with one or more suitable pharmaceutically active agents selected from following therapeutic agents in any combination. Immunosuppressants (eg., Methotrexate, mercaptopurine, cyclophosphamide),
5 glucocorticoids, non-steroidal anti-inflammatory drugs, Cox-2 specific inhibitors, TNF- α binding proteins (eg., Infliximab, etanercept), interferon- γ , interferon- α , interleukin-2, antihistamines, beta-agonist, anticholinergics, anticancer agents or their suitable pharmaceutically acceptable salts. Further examples of anticancer agents for use in combination with BTK inhibitors
10 include chemotherapy or a targeted therapy, alkylating agents, platinum compounds, DNA altering agents, Topoisomerase inhibitors, microtubule modifiers, antimetabolites, anticancer antibiotics, hormones, Aromatase inhibitors, antibodies, cytokines, vaccines, drug conjugates, inhibitors of mitogen-activated protein kinase signaling (ex: BAY 43-9006), Syk
15 inhibitors, mTOR inhibitors, antibodies (Rituxan), other anticancer agents that can be employed in combination include, Vinblastin, Bleomycin, Cisplatin, Acivicin, Azacitidine, Decitabine, Doxorubicin, Etoposide, Flurouracil, Methotrexate, Vinblastin, Vincristine and BCR/ABL antagonist. Use of a compound of Formula (I) or its pharmaceutical composition according to any
20 of the preceding claims for the manufacture of a medicament for treating a disease, disorder or condition, which would benefit from inhibition of tyrosine kinase (s), such as BTK activity.
13. Use of the compound of formula (I) and a suitable pharmaceutically
25 acceptable agent selected from anti-cancer and anti-inflammatory agents or their pharmaceutically acceptable salts for the treatment of autoimmune and allergic disorders, such as rheumatoid arthritis, systemic lupus erythematosus, scleroderma, asthma, allergic rhinitis, allergic eczema, B-cell lymphoma,

5

multiple sclerosis, juvenile rheumatoid arthritis, juvenile idiopathic arthritis, inflammatory bowel disease, graft versus host disease, psoriatic arthritis, ankylosing spondylitis and uveritis, B cell lymphoma, multiple sclerosis, diffuse large B-cell lymphoma, follicular lymphoma or chronic lymphocytic leukemia and B-cell prolymphocytic leukemia.