

T 1632 FF

IED

HERBICIDAL 2,6-SUBSTITUTED PYRIDINES

The present invention is concerned with certain 2,6-substituted pyridines, their preparation and use as herbicides.

Pyridines and pyridine derivatives have many uses, for example, as pharmaceuticals, pesticides (herbicides, acaricides, anthelmintics, bird repellants), feed supplements, solvents, reagents, intermediates, and chemicals for the polymer and textile industry. Various 2,6-diaryloxy- or -diarylmethoxy-pyridine derivatives have been investigated for such uses, the compounds often having additional substitution of the central pyridine ring.

10 EP-A2-180188 is concerned with 2-hydroxypyridine derivatives, for example 2,6-dihydroxypyridines, as potentiators for increasing the anticancer activity of 5-fluorouracil and related compounds. The preparation and debenzoylation of 2,6-dibenzyloxy-3-chloro-pyridine is disclosed. In J. Biol. Chem., 72, Vol. 247(23), pages

15 7628 to 7633, (which is concerned with the synthesis of 2,3,6-trihydroxypyridine and accumulation and partial characteristics of the product of 2,6-dihydroxypyridine oxidation) the Grignard reaction of 3-bromo-2,6-dibenzyloxypyridine with isopropyl bromide is disclosed. There is no disclosure of

20 biological activity for 2,6-dibenzyloxy-3-chloropyridine nor its bromo analogue.

2,6-Di(3-methylphenoxy)-pyridine is disclosed in French Patent Specification No. 1527714 which is concerned with polyethers containing a 2,6-linked pyridine group. US-A-4830846 is concerned

25 with a separation process for anhydrous HCl and HBr by thermal cleavage utilising a hindered pyridine, one example of which being 2,6-diphenoxypyridine. Again no biological activity is disclosed for these aryloxy pyridines.

The dibenzyloxy analogue of 2,6-diphenoxypyridine in addition

30 to being mentioned in JP-A-63250224, EP-A-180188 and J. Med. Chem,


10(2), pages 320 to 325, all medical research publications, was prepared as an example of the herbicidal compounds proposed in US-A-3 535 328 and the divisional published as US-A-3 687 959.

Both texts are predominantly directed to the herbicidal, fungicidal, and, for the compounds claimed in US-A-3 535 328, nematocidal and insecticidal activity, of amido or aminothioethoxy derivatives. No herbicidal activity data is given for the disclosed 2,6-dibenzyloxypyridine.

Herbicidal evaluation of 2,6-dibenzyloxypyridine and its analogue 2,6-diphenoxypyridine now undertaken has revealed that the compounds have herbicidal activity primarily against representative broadleaf plant species in post-emergence foliar spray application but little or no post-emergence action against representative grass-type plants and a complete absence of pre-emergence herbicidal activity.

Surprisingly, it has now been found that certain substituted forms of 2,6-diphenoxypyridine and 2,6-dibenzyloxypyridine, and also the related assymetric 2-phenoxy-6-dibenzyloxy-pyridines, have a significantly different spectrum of activity and not only possess high activity against broadleaf plants in foliar spray applications but also have significant action against grass-type plants, particularly against the important weed, barnyard grass; a number of examples additionally being effective against both plant types pre-emergence.

Accordingly, the present invention provides a herbicidal composition which comprises a carrier and, as an active ingredient, a compound of the general formula



in which

35 each of n and m independently is 0 or 1;

each of Ar^1 and Ar^2 independently represents an aryl group, at least one of Ar^1 and Ar^2 being substituted by one or more of the same or different substituents selected from halogen atoms, alkyl groups, alkoxy groups, haloalkyl groups and haloalkoxy groups;

5 and

R^1 represents a hydrogen atom, or an alkyl, alkylthio or haloalkyl group having from 1 to 4 carbon atoms, and R^2 represents a hydrogen or a halogen atom provided that at least one of R^1 and R^2 represents a hydrogen atom.

10 An alkyl group, unless otherwise specified may be a straight chain or branched chain group suitably containing up to 12 carbon atoms, preferably from 1 to 4 carbon atoms. An alkyl portion of a haloalkyl, alkylthio, alkoxy or haloalkoxy group suitably has from 1 to 4 carbon atoms, preferably 1 or 2 carbon atoms.

15 Halogen is fluorine, chlorine, bromine or iodine. Haloalkyl and haloalkoxy are preferably mono-, di- or tri-fluoro-alkyl and -alkoxy, especially trifluoromethyl and trifluoromethoxy.

An aryl group is preferably a phenyl group.

20 The $\text{Ar}^1-(\text{CH}_2)_n-\text{O}-$ and $\text{Ar}^2-(\text{CH}_2)_m-\text{O}-$ portions of compounds of formula I may be identical or different; thus the compounds may be symmetrical or assymetrical diaryloxypyridines or diarylmethoxypyridines, or may be aryloxyarylmethoxypyridines.

Preferably one of n and m is 1 and the other is 0 or 1, but most preferably each of n and m is 1.

25 At least one of the groups Ar^1 and Ar^2 is substituted.

Suitable substituents include methoxy and trifluoromethoxy groups and, preferably, methyl groups, fluorine atoms, chlorine atoms, and trifluoromethyl groups. From 1 to 5 substituents may be present; preferably 1 or 2 substituents.

30 R^1 is preferably a hydrogen atom or a methyl, methylthio or trifluoromethyl group. Especially preferred is hydrogen or methyl.

R^2 is suitably a hydrogen or chlorine atom, especially a hydrogen atom.

35 Preferred compounds fall in the following categories, with reference to the symbols used in general formula I:

- (i) n is 0; m is 0; R^1 and R^2 each represents a hydrogen atom; Ar^1 represents a trifluoromethylphenyl group, especially a 3-trifluoromethylphenyl group; and Ar^2 represents an unsubstituted or chloro- or trifluoromethyl-substituted phenyl group;
- (ii) n is 0; m is 1; R^1 and R^2 each represents a hydrogen atom; Ar^1 represents a substituted phenyl group, especially a 3-trifluoromethylphenyl group; and Ar^2 represents an unsubstituted phenyl group or a fluoro-substituted group;
- (iii) n is 1; m is 1; R^1 and R^2 each represents a hydrogen atom; and each of Ar^1 and Ar^2 represents a fluoro- or methyl- substituted phenyl group, especially ortho and/or para substituted, e.g. a 2- or 4- fluorophenyl group or a 2-methyl group.

Most of the compounds of general formula I are novel and accordingly the present invention provides a compound of general formula I in which n, m, Ar^1 , Ar^2 , R^1 and R^2 have the meanings given above with the proviso that

- a) when n and m are both 0, then Ar^1 and Ar^2 do not each represent a 3-methylphenyl group; and
- b) when n and m are both 1, then R^2 does not represent a chlorine or bromine atom.


The compounds of formula I may be prepared by appropriate adaptation of conventional methods of obtaining substituted pyridine compounds, the basic technique being, for example, reaction of a metal salt of the appropriate alcohol with an appropriate 2,6-dihalopyridine, in a solvent and suitably at elevated temperature, ideally at reflux. For symmetrical pyridine compounds preparation, the reaction can be carried out in one step by using a molar ratio of salt to pyridine of at least 2:1. For assymetrical compounds of formula I, separate introduction of the two substituents is required in a two-stage process.

The present invention therefore provides a process for the preparation of a compound of formula I, which comprises

- a) to obtain a compound of formula I in which Ar^1 and Ar^2 are the same, reacting a metal salt of a compound of general formula



in which Ar^1 and n are as defined in claim 1, with a 2,6-halopyridine of general formula



(III)

in which Hal^1 and Hal^2 each, independently, represents a halogen atom, and R^1 and R^2 are as defined in claim 1, in a molar ratio of at least 2:1

or

- b) to obtain a compound of formula I in which Ar^1 and Ar^2 are different, reacting a compound of formula II with a 2,6-dihalopyridine of formula III in a molar ratio of 1:1 and then reacting the resulting product with a metal salt of a compound of general formula IV



in which Ar^2 and n are as defined in claim 1, in a molar ratio of at least 1:1

each reaction stage being carried out in the presence of an organic solvent.

The metal salt of the alcohol of formula II is conveniently an alkali metal salt, for example a sodium or potassium salt, and is conveniently generated by reaction of the alcohol with a suitable metal base, for example a metal carbonate or hydride. Preferably the metal salt is a sodium salt generated using sodium hydride.

The organic solvent utilised in the reaction would be selected depending on the nature of the reactants involved. Generally any polar organic solvent is suitable, for example dimethylformamide and tetrahydrofuran.

5 Hal¹ and Hal² conveniently are both the same, each representing a chlorine or bromine atom.

The prepared compounds of formula I may, if desired, be isolated and purified using conventional techniques.

10 Compounds of general formula II, III and IV are generally known and/or are easily preparable by standard techniques.

The present invention also provides the use of a compound of formula I as a herbicide. Further, in accordance with the invention there is provided a method of combating undesired plant growth at a locus by treating the locus with a composition
15 according to the invention or a compound of formula I. As the most useful action is by foliar spray application, the locus is most suitably the plants in a crop area, typical crops being cereals. However, application may also be to the soil for those compounds having pre-emergence herbicidal action. The dosage of active
20 ingredient used may, for example, be in the range of from 0.01 to 10 kg/ha, preferably 0.1 to 1.0 kg/ha.

Whilst all compounds of the present invention have useful activity against broadleaf and narrowleaf plants, exceptionally good activity is shown by compounds in which at least one of n and
25 m is 1, particularly in the phenoxy/benzyloxy pyridines and, most preferably, in the dibenzyloxypyridines of the present invention.

The present invention also extends to a method of making a herbicidal composition of the invention which comprises bringing a compound of formula I into association with at least one carrier.

30 Preferably there are at least two carriers in a composition of the present invention, at least one of which is a surface-active agent.

A carrier in a composition according to the invention is any material with which the active ingredient is formulated to
35 facilitate application to the locus to be treated, which may be, as

appropriate, a plant, seed or soil, or to facilitate storage, transport or handling. A carrier may be a solid or a liquid, including a material which is normally gaseous but which has been compressed to form a liquid, and any of the carriers normally used
5 in formulating herbicidal compositions may be used. Preferably compositions according to the invention contain 0.5 to 95% by weight of active ingredient.

Suitable solid carriers include natural and synthetic clays and silicates, for example natural silicas such as diatomaceous
10 earths; magnesium silicates, for example talcs; magnesium aluminium silicates, for example attapulgites and vermiculites; aluminium silicates, for example kaolinites, montmorillonites and micas; calcium carbonate; calcium sulphate; ammonium sulphate; synthetic hydrated silicon oxides and synthetic calcium or aluminium
15 silicates; elements, for example carbon and sulphur; natural and synthetic resins, for example coumarone resins, polyvinyl chloride, and styrene polymers and copolymers; solid polychlorophenols; bitumen; waxes; and solid fertilisers, for example superphosphates.

Suitable liquid carriers include water; alcohols, for example
20 isopropanol and glycols; ketones, for example acetone, methyl ethyl ketone, methyl isobutyl ketone and cyclohexanone; ethers; aromatic or araliphatic hydrocarbons, for example benzene, toluene and xylene; petroleum fractions, for example kerosine and light mineral oils; chlorinated hydrocarbons, for example carbon tetrachloride,
25 perchloroethylene and trichloroethane. Mixtures of different liquids are often suitable.

Agricultural compositions are often formulated and transported in a concentrated form which is subsequently diluted by the user before application. The presence of small amounts of a carrier
30 which is a surface-active agent facilitates this process of dilution. Thus preferably at least one carrier in a composition according to the invention is a surface-active agent. For example the composition may contain at least two carriers, at least one of which is a surface-active agent.

A surface-active agent may be an emulsifying agent, a dispersing agent or a wetting agent; it may be nonionic or ionic. Examples of suitable surface-active agents include the sodium or calcium salts of polyacrylic acids and lignin sulphonic acids; the
5 condensation products of fatty acids or aliphatic amines or amides containing at least 12 carbon atoms in the molecule with ethylene oxide and/or propylene oxide; fatty acid esters of glycerol, sorbitol, sucrose or pentaerythritol; condensates of these with ethylene oxide and/or propylene oxide; condensation products of
10 fatty alcohol or alkyl phenols, for example p-octylphenol or p-octylcresol, with ethylene oxide and/or propylene oxide; sulphates or sulphonates of these condensation products; alkali or alkaline earth metal salts, preferably sodium salts, of sulphuric or sulphonic acid esters containing at least 10 carbon atoms in the
15 molecule, for example sodium lauryl sulphate, sodium secondary alkyl sulphates, sodium salts of sulphonated castor oil, and sodium alkylaryl sulphonates such as dodecylbenzene sulphonate; and polymers of ethylene oxide and copolymers of ethylene oxide and propylene oxide.

20 The herbicidal composition of the invention may also contain other active ingredients, for example compounds possessing insecticidal or fungicidal properties, or other herbicides.

The following Examples illustrate the invention. The structures of the compounds prepared in the following Examples were
25 additionally confirmed by mass spectrometry and NMR.

Preparation of 2,6-diaryloxy pyridines

Example 1

30 Preparation of 2,6-di-[3-trifluoromethoxyphenoxy]pyridine

3-Trifluoromethoxyphenol (19g; 0.106 mol) was added to oil-free sodium hydride (2.7g) in dry dimethylformamide (120 ml). 2,6-Dichloropyridine (8g; 0.054 mol) was then added and the reaction mixture refluxed for 4 hours. The symmetrical phenoxy
35 pyridine 2,6-di-[3-trifluoromethoxyphenoxy]-pyridine was isolated,

following conventional work up, chromatography and distillation procedures, as a colourless oil (17g; yield 73%).

Boiling point: 145°C at 3 mm Hg.

5. Example 2

Preparation of 2,6-di-(3-trifluoromethylphenoxy)-4-methylthio-pyridine

To a solution of 2,6-dichloro-4-nitropyridine (3.8g; 0.02 mol) in dimethylformamide (50 ml) cooled with an ice-bath was sodium
10 thiomethoxide (1.4g; 0.02 mol) slowly added. After stirring for 3 hours, the reaction mixture was quenched with water (300 ml) and extracted 3 times each with a hexane/ethyl acetate mixture (100 ml; 1/1). The combined extracts were dried with anhydrous magnesium sulphate and the solvent was removed in vacuo. Purification by
15 silica gel flash column chromatography afforded 2,6-dichloro-4-methylthiopyridine (2.9g; 74%) as a colourless oil.

A mixture of 2,6-dichloro-4-methylthiopyridine (0.97 g; 0.005 mol), 3-hydroxybenzotrifluoride (1.3 ml; 10.4 mmol) and sodium hydride (0.2 g) was reacted and worked up according to the
20 procedure of Example 1 above to give 2,6-di-(3-trifluoromethylphenoxy)-4-methylthiopyridine (1.9 g; 85%)

Example 3

Preparation of 2,6-di-(3-trifluoromethylphenoxy)-4-methylpyridine

25 To a solution of 2-methyl-2-chloromethyloxirane (50 g; 0.47 mol) in concentrated hydrochloric acid (23 ml) at ice-bath temperature was added a solution of sodium cyanide (27.4 g; 0.56 mol) in hydrochloric acid (23 ml). After stirring for 10 hours at that temperature, the reaction mixture was warmed to 40°C
30 and a solution of potassium cyanide (33.8 g; 0.52 mol) in water (50 ml) was added. The reaction mixture was warmed to 50°C and stirred for 4 hours. After cooling, the mixture was neutralised and extracted 3 times each with ethyl acetate (150 ml). The combined extracts were dried with anhydrous magnesium sulphate.
35 Removal of the solvent in vacuo afforded 1,3-dicyano-2-methyl-2-

hydroxypropane (56.4 g, 96%). This compound was added carefully to a 33% solution of hydrogen bromide in glacial acetic acid at ice-bath temperature. The reaction mixture was stirred for 3 days at ambient temperature. The solvent was removed in vacuo and the residual oil brought to pH 12 with a 10 molar aqueous solution of sodium hydroxide. The alkaline solution was extracted 3 times each with ethyl acetate (100 ml). The extracts were combined and the solvent removed in vacuo to afford 6-amino-2-bromo-4-methylpyridine (56 g; 66%) as colourless crystals. Melting point 99°C.

A solution of 6-amino-2-bromo-4-methylpyridine (45.2 g, 0.24 mol) in water (100 ml) and concentrated sulphuric acid (43 g) at 0°C was stirred and a solution of sodium nitrite (13.2 g; 0.19 mol) in water (20 ml) was added. After 2 hours, the reaction mixture was warmed to 60°C and stirred for 1 hour. After cooling, the mixture was extracted with dichloromethane (200 ml). The solvent was removed in vacuo to afford 2-bromo-4-methyl-6-hydroxypyridine (20.2 g; 56%) as colourless crystals. Melting point 152°C. This material was mixed with tribromomethane (100 ml) and phosphoryl bromide (24 g) and heated to reflux for 3 hours. After cooling, the reaction mixture was quenched carefully with an aqueous 50% solution of sodium hydroxide and the mixture extracted twice each with dichloromethane (100 ml). The solvent was removed in vacuo and the crude product purified by flash silica gel column chromatography using hexane/ethyl acetate (7/3). 2,6-dibromo-4-methylpyridine (6.9 g; 25%) was obtained. Melting point 77°C. This compound (1.3 g; 5.2 mmol) was then converted using 3-hydroxybenzotrifluoride (1.4 ml; 10.7 mmol) and sodium hydride according to the procedure of Example 1 above to give 2,6-di-(3-trifluoromethylphenoxy)-4-methylpyridine as an oil (1.8g; 84%).

Example 4

Preparation of 2-(3-trifluoromethylphenoxy)-6-(4-fluorophenoxy)-pyridine

a) Preparation of 2-(3-trifluoromethylphenoxy)-6-chloropyridine

5 3-Trifluoromethylphenol (65g; 0.40 mol) was added dropwise and with stirring to a suspension of oil-free sodium hydride (10g) in dry dimethylformamide (250 ml) maintained under a nitrogen atmosphere. After 30 minutes
10 2,6-dichloropyridine (59.5g; 0.40 mol) was added and the reaction mixture refluxed with stirring for 1½ hours. Dimethylformamide was removed in vacuo, and solvent extraction carried out using a chloroform/water mixture (11; 50/50). Following purification and then chromatography, a colourless oil (88g; 80%) was obtained from the organic extraction layer,
15 and identified as 2-(3-trifluoromethylphenoxy)-6-chloropyridine.

Boiling point: 120°C at 2mm Hg.

b) Preparation of 2-(3-trifluoromethylphenoxy)-6-(4-fluorophenoxy)pyridine

20 The procedure of a) above was followed using 4-fluorophenol (8g; 0.07 mol) in dry dimethylformamide (20ml), sodium hydride (2g) also in dry dimethylformamide (60 ml), 2-(3-trifluoromethylphenoxy)-6-chloropyridine from a) above (19.5g; 0.07 mol) with a 2 hour reflux. The assymetrical
25 phenoxypyridine 2-(3-trifluoromethylphenoxy)-6-(4-fluorophenoxy)pyridine (18.9g; 76%) was derived as a colourless oil from the organic layer of chloroform/water (500ml; 50/50) following solvent extraction.

Boiling point: 145°C at 1mm Hg

30 Preparation of aryloxy-arylmethoxy-pyridines

Example 5

Preparation of 2-(3-trifluoromethylphenoxy)-6-benzyloxypyridine

35

a) Preparation of 2-benzyloxy-6-chloropyridine

Following the procedure of Example 4a) above, benzyl alcohol (5g; 46 mmols) reacted with sodium hydride (1.19g), was refluxed for 1 hour with 2,6-dichloropyridine (6.9g; 46mmols) using sodium hydride and dry dimethylformamide as solvent. 2-Benzyloxy-6-chloropyridine (6.1g; 60%) was obtained as a colourless oil following solvent extraction with chloroform/water (500ml; 50/50), purification and chromatographic separation of the product from the organic layer, identified by elemental analysis and mass spectrum.

b) Preparation of 2-(3-trifluoromethylphenoxy)-6-benzyloxy-pyridine

The procedure of Example 4a) was also used to introduce the 3-trifluoromethyl phenoxy group by reaction of 2-benzyloxy-6-chloropyridine (5g; 23mmols) from a) above with 3-trifluoromethylphenol (3.3g; 0.018 mols) using dry sodium hydride (0.55g) and dry dimethylformamide (200ml) as solvent, with a 1 hour reflux stage. 2-(3-Trifluoromethylphenoxy)-6-benzyloxy-pyridine (4.9g; 51%) was obtained as a yellow oil following solvent extraction (chloroform/water - 500ml; 50/50), and chromatography plus distillation of the organic layer.

Boiling point: 145°C at 2mm Hg.

25 Preparation of 2,6-diarylmethoxypyridines

Example B

Preparation of 2,6-dibenzyloxypyridine


Benzyl alcohol (5.1g; 0.05 mols) was reacted with 2,6-dichloropyridine (3.5g; 0.024 mols) utilising sodium hydride (1.2g) and, as solvent, dimethylformamide by the procedure of Example 1 above. Usual work up, chromatography and recrystallisation gave 2,6-dibenzyloxypyridine as a colourless solid (5.5g; 81%).

35 Melting point: 74°C

Examples Nos 6 to 25

Compounds according to the invention were prepared following the relevant preparation technique of the above Examples 1 to 5 and B and using appropriate starting materials.

5 Details of Examples 6 to 25 are given in Table 1 below, identification being by reference to the substituents shown in the following formula:



15 In Table 1 'bp' indicates boiling point and 'mp' indicates melting point. Elemental analysis data is given in Table 1a following.

Table 1

20

Ex. No.	R	n	R ¹	R ²	m	R ³	Identification Data	
25	6	3-CF ₃	0	H	H	0	3-CF ₃	bp 150°C/1mmHg
	7	3-OCF ₃	0	H	H	0	H	bp 150°C/1mmHg
	8	3-CF ₃	0	H	H	0	2-F	bp 165°C/2mmHg
	9	3-CF ₃	0	H	H	0	3-F	bp 180°C/2mmHg
	10	3-CF ₃	0	H	H	0	2,4-F ₂	bp 150°C/1mmHg
	11	3-CF ₃	0	H	H	0	3-CH ₃	bp 160°C/2mmHg
30	12	3-CF ₃	0	H	H	0	3-Cl	bp 160°C/2mmHg
	13	3-CF ₃	0	H	H	0	4-Cl	bp 150°C/3mmHg
	14	3-CF ₃	0	H	Cl	0	3-CF ₃	bp 175°C/2mmHg
	15	3-CF ₃	0	H	H	0	3-OCH ₃	-
	16	3-CF ₃	0	H	H	0	4-CH ₃	-
35								

Table 1 (continued)

Ex. No.	R	n	R ¹	R ²	m	R ³	Identification Data
17	3-CF ₃	0	CF ₃	H	0	3-CF ₃	bp 160°C/2mmHg
18	3-CF ₃	0	H	H	1	3-F	bp 135°C/3mmHg
19	2-CH ₃	1	H	H	1	2-CH ₃	bp 165°C/3mmHg
20	3-Cl	1	H	H	1	3-Cl	bp 180°C/2mmHg
21	4-Cl	1	H	H	1	4-Cl	mp 51-52°C
22	2-F	1	H	H	1	2-F	bp 170°C/3mmHg
23	3-F	1	H	H	1	3-F	bp 180°C/2mmHg
24	4-F	1	H	H	1	4-F	mp 82-84°C
25	3-CF ₃	1	H	H	1	3-CF ₃	bp 155°C/2mmHg

Table 1a

Example No.	Elemental Analysis (%)					
	Calculated			Found		
	C	H	N	C	H	N
1	52.9	2.5	3.3	53.6	3.0	3.6
2	53.9	2.9	3.1	53.7	2.7	3.1
3	58.1	3.2	3.4	57.9	3.1	3.4
4	61.9	3.2	4.0	62.1	3.2	4.4
5	66.1	4.1	4.1	66.3	5.0	4.5
6	57.1	2.8	3.5	56.6	3.1	3.8
7	65.3	3.6	4.2	65.7	3.1	3.5
9	61.9	3.2	4.0	62.3	3.7	3.9
10	58.9	2.7	3.8	58.6	2.9	4.3
11	66.1	4.1	4.3	66.3	4.3	4.2
12	59.1	3.0	3.8	60.1	3.0	3.5
13	59.1	3.0	3.8	59.3	3.5	4.1
14	52.6	2.3	3.2	54.6	3.1	4.0

Table 1a (continued)

Example	Elemental Analysis (%)						
No.	Calculated			Found			
	C	H	N	C	H	N	
5							
	15	63.2	3.9	3.9	63.0	3.8	4.0
	16	66.1	4.1	4.1	65.9	4.0	3.8
	17	51.4	2.1	3.0	51.2	2.1	3.2
10	18	61.4	3.3	4.2	61.0	3.4	4.4
	19	79.0	6.6	4.4	78.7	6.3	4.3
	21	63.3	4.2	3.9	63.7	4.3	3.8
	22	69.7	4.6	4.3	69.5	3.9	3.9
	23	69.7	4.5	4.3	70.9	5.0	3.5
15	24	69.7	4.6	4.3	69.6	4.6	5.0
	25	59.0	3.5	3.3	60.2	3.7	3.3

Example 26

20 Herbicidal Activity

To evaluate their herbicidal activity, compounds were tested using, as a representative range of monocotyledon plants: maize, Zea mays (Mz); rice, Oryza sativa (R); barnyard grass, Echinochloa crus-galli (BG); and oat, Avena sativa (O). As a representative range of dicotyledon plants, the following were also used: linseed, Linum usitatissimum (L); mustard, Sinapsis alba (M); sugar beet, Beta vulgaris (SB) and soya bean, Glycine max (S).

The tests fall into two categories, pre-emergence and post-emergence. The pre-emergence tests involved spraying a liquid formulation of the compound onto the soil in which the seeds of the plant species mentioned above had recently been sown. The post-emergence tests involved two types of test, viz., soil drench and foliar spray tests. In the soil drench tests the soil in which the seedling plants of the above species were growing was drenched with a liquid formulation containing a test compound, and in the

foliar spray tests the seedling plants were sprayed with such a formulation.

The soil used in the tests was a prepared horticultural loam.

The formulations used in the tests were prepared from
5 solutions of the test compounds in acetone containing 0.4% by weight of an alkylphenol/ethylene oxide condensate available under the trade mark TRITON X-155. These acetone solutions were diluted with water and the resulting formulations applied at dosage levels corresponding to 5 kg or 1 kg of active material per hectare in a
10 volume equivalent to 600 litres per hectare in the soil spray and foliar spray test, and at a dosage of level equivalent to 10 kilograms of active material per hectare in a volume equivalent to approximately 3,000 litres per hectare in the soil drench tests.

In the pre-emergence tests untreated sown soil and in the
15 post-emergence tests untreated soil bearing seedling plants were used as controls.

The herbicidal effects of the test compounds were assessed visually twelve days after spraying the foliage and the soil, and thirteen days after drenching the soil, and were recorded on a 0-9
20 scale. A rating 0 indicates growth as untreated control, a rating 9 indicates death. An increase of 1 unit on the linear scale approximates to a 10% increase in the level of effect.

The results of the tests are set out in Table 2 below. In the Table, a blank space indicates a rating 0; an asterisk
25 indicates that no result was obtained.

Compound B is the compound 2,6-dibenzoyloxy pyridine, melting point 74°C, disclosed as the compound of Example No. 81 of Table I(b) of US-A-3 535 328 and its divisional US-A-3 687 959. Compound A is the corresponding diphenoxy compound, 2,6-diphenoxy pyridine,
30 boiling point 170°C at 5 mmHg. The herbicidal activity of these compounds was assessed by the procedure given above, and the results included in Table 2: the data for compound A precedes that for the diphenoxy compounds 1 to 16 of the present invention and the data for compound B precedes that for the dibenzoyloxy compounds
35 19 to 25 of the present invention.

Table 2

Example No.	Soil drench 10 kg/ha								Dosage kg/ha	Foliar spray								Pre-emergence							
	Mz	R	BG	O	L	M	SB	S		Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S
A									5	3	2	2	3	5	9	6	2								
									1	1		1	1	3	7	2	1								
1			1	2	1	3	2		5	4	1	5	5	4	8	9	6	1		3			5	1	
									1	3		1	2	2	8	8	3			1			2		
2			5	2		6			5	6		8	6	6	8	9	7			8	4	1	8	7	
									1	4		8	5	4	8	8	6			7			7	6	
4			3	1	1				5	5	2	7	5	4	9	8	6	3	1	6	3	2	5	3	
									1	3		3	2	3	8	8	3	1		2	1		3		
5	6	4	7	5	3	7	7	3	5	5	5	8	6	7	9	9	7	5	1	8	5	3	8	6	
									1	2	1	4	3	5	9	8	4	1		6	1	1	5	3	
6	2	1	4	3	2	4	3	1	5	5	3	6	5	5	9	9	6	4	3	7	5	4	8	7	1
									1	2	1	4	4	4	9	9	4	2	1	5	1		6	4	

Table 2 (continued)

Example No.	Soil drench 10 kg/ha								Dosage kg/ha	Foliar spray								Pre-emergence							
	Mz	R	BG	O	L	M	SB	S		Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S
7	2	1	2	1					5	6	2	6	4	5	8	8	5	1		6	4	2	4	2	
									1	4		2	2	4	8	8	4		3			2			
8									5	6	4	8	6	7	9	9	7		4	3		7	3	2	
									1	3		3	3	6	9	8	6				2				
9									5	5	4	8	8	8	9	9	8								
									1	3	2	6	4	7	9	9	7								
10		1	1	1					5	6	2	7	4	5	9	9	6	1		4	1		6	2	
									1	4		2	2	2	9	8	5	1		1		2			
11									5	5	2	7	5	7	9	9	5		1	4	2		4	3	
									1	3		4	3	6	9	9	4				4	1			
12									5	7	5	8	7	8	9	9	7	2	2	7	6	2	8	7	3
									1	5	2	6	5	7	9	9	7	1		5	4		6	3	1

Table 2 (continued)

Example No.	Soil drench 10 kg/ha								Dosage kg/ha	Foliar spray								Pre-emergence							
	Mz	R	BG	O	L	M	SB	S		Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S
13				3	3	4	3	1	5	5	4	6	6	7	9	9	6		2	6	4	2	7	6	
									1	3	3	3	4	6	9	8	5						6		
14									5	6	2	5	5	4	9	9	6	1		5			4		
									1	3		1	2	2	8	7	4						2		
15									5	5		7	4	5	9	9	7	1		5			1		
									1	2		5	3	4	8	9	6								
16									5	5	2	7	3	6	9	9	6								
									1	3		2		2	6	3	5								
17									5	5		6	3	6	7	6	6			5			1		
									1					2	1		2			2					
18	2	2	5	5	3	3	1		5	6	5	8	7	8	9	9	7						*	*	
									1	4	3	6	6	6	9	8	6						4	2	

Table 2 (continued)

Example No.	Soil drench 10 kg/ha								Dosage kg/ha	Foliar spray								Pre-emergence							
	Mz	R	BG	O	L	M	SB	S		Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S
B						5			5	3		4	2	4	9	8	4								
									1	1		2		3	9	7	4								
19									5	4	5	8	5	7	9	9	5	2		8	4		7	4	
									1	2	1	6	3	6	8	9	3			2			3		
20									5	6		6	4	6	9	9	5								
									1	3		2	2	3	8	7	3								
21									5	5		7	4	6	8	9	6						2		
									1	3		5	2	5	8	7	5								
22	4	2	5	5					5	4	6	8	6	7	9	9	6	3	2	8	6	3	8	8	
									1	2	2	7	5	6	9	8	5			6	3		7	5	
23									5	7	2	8	4	6	9	9	7			6			1		
									1	4		6	1	4	9	8	5								

Table 2 (continued)


Example No.	Soil drench 10 kg/ha								Dosage kg/ha	Foliar spray								Pre-emergence							
	Mz	R	BG	O	L	M	SB	S		Mz	R	BG	O	L	M	SB	S	Mz	R	BG	O	L	M	SB	S
24					1	1	3		5	4	2	9	4	6	9	9	5			3			1	2	
									1	2		8	2	6	9	8	5								
25									5	5		5	4	5	9	7	6								
									1	2		2	1	3	7	5	4								



T 1632 FF

CLAIMS

1. A herbicidal composition which comprises a carrier and, as active ingredient, a compound of the general formula



in which

- each of n and m independently is 0 or 1;
- each of Ar^1 and Ar^2 independently represents an aryl group, at least one of Ar^1 and Ar^2 being substituted by one or more of the same or different substituents selected from halogen atoms, alkyl groups, alkoxy groups, haloalkyl groups and haloalkoxy groups;
- and
- R^1 represents a hydrogen atom, or an alkyl, alkylthio or haloalkyl group having from 1 to 4 carbon atoms, and R^2 represents a hydrogen or a halogen atom provided that at least one of R^1 and R^2 represents a hydrogen atom.
2. A composition as claimed in claim 1, in which, in the compound of formula I, at least one of n and m is 1.
3. A composition as claimed in claim 2, in which, in a compound of formula I, each of n and m is 1.
4. A composition as claimed in any one of claims 1 to 3, in which, in the compound of formula I, the substituents of Ar^1 and/or Ar^2 are independently selected from methyl groups, methoxy groups, fluorine atoms, chlorine atoms, trifluoromethyl and trifluoromethoxy groups.
5. A composition as claimed in any one of claims 1 to 4, in which R^1 represents a hydrogen atom and R^2 represents a hydrogen or



chlorine atom, or R^1 represents a trifluoromethyl, methylthio or methyl group and R^2 represents a hydrogen atom.

6. A composition as claimed in any one of claims 1 to 5, which comprises at least two carriers, at least one of which is a surface-active agent.

7. A compound of the general formula I as defined in any one of the claims 1 to 5, with the proviso that


- a) when n and m are both 0, then Ar^1 and Ar^2 do not each represent a 3-methylphenyl group; and
- b) when n and m are both 1, then, R^2 does not represent a chlorine or bromine atom.

8. A process for the preparation of a compound as claimed in claim 7, which comprises

- a) to obtain a compound of formula I in which Ar^1 and Ar^2 are the same, reacting a metal salt of a compound of general formula




in which Ar^1 and n are as defined in claim 1 with a 2,6-halopyridine of general formula




in which Hal^1 and Hal^2 each, independently, represent a halogen atom, and R^1 and R^2 are as defined in claim 1, in a molar ratio of at least 2:1

or

- b) to obtain a compound of formula I in which Ar^1 and Ar^2 are different, reacting a compound of formula II with a 2,6-dihalopyridine of formula III in a molar ratio of 1:1



and then reacting the resulting product with a metal salt
of a compound of general formula IV



5

in which Ar^2 and n are as defined in claim 1, in a molar
ratio of at least 1:1

each reaction stage being carried out in the presence of an organic
solvent.

10

9. A method of combating undesired plant growth at a locus which
comprises treating the locus with a composition as claimed in any
one of claims 1 to 6, or a compound of formula I as defined in
claim 1.

15

10. The use of a composition as claimed in any one of claims 1 to
6, or a compound of formula I as defined in claim 1, as a
herbicide.