

NOVEL COMPOUNDS AS CANNABINOID
RECEPTOR LIGANDS AND USES THEREOF

This application claims priority to the provisional application Serial No.
5 60/809,712 filed on May 31, 2006.

Technical Field

The present invention relates to compounds that are cannabinoid receptor
ligands, compositions comprising such compounds, and methods of treating
conditions and disorders using such compounds and compositions.
10

Background

(-)- Δ^9 -Tetrahydrocannabinol (Δ^9 -THC), the major psychoactive constituent of
marijuana, exerts a broad range of therapeutic effects through its interactions with two
cannabinoid (CB) receptor subtypes, CB₁ and CB₂. CB₁ receptors are highly
15 expressed in the central nervous system and to a lesser degree in the periphery in a
variety of tissues of the cardiovascular and gastrointestinal systems. By contrast, CB₂
receptors are most abundantly expressed in multiple lymphoid organs and cells of the
immune system, including spleen, thymus, tonsils, bone marrow, pancreas and mast
cells.

20 The psychotropic side effects caused by Δ^9 -THC and other nonselective CB
agonists are mediated by CB₁ receptors. These CB₁ receptor-mediated effects, such
as euphoria, sedation, hypothermia, catalepsy, and anxiety, have limited the
development and clinical utility of nonselective CB agonists. Recent studies have
demonstrated that CB₂ modulators are analgesic in preclinical models of nociceptive
25 and neuropathic pain without causing the adverse side effects associated with CB₁
receptor activation. Therefore, compounds that selectively target CB₂ receptors are an
attractive approach for the development of novel analgesics.

Pain is the most common symptom of disease and the most frequent complaint
with which patients present to physicians. Pain is commonly segmented by duration
30 (acute vs. chronic), intensity (mild, moderate, and severe), and type (nociceptive vs.
neuropathic).

Nociceptive pain is the most well known type of pain, and is caused by tissue

injury detected by nociceptors at the site of injury. After the injury, the site becomes a source of ongoing pain and tenderness. This pain and tenderness are considered "acute" nociceptive pain. This pain and tenderness gradually diminish as healing progresses and disappear when healing is complete. Examples of acute nociceptive pain include surgical procedures (post-op pain) and bone fractures. Even though there may be no permanent nerve damage, "chronic" nociceptive pain results from some conditions when pain extends beyond six months. Examples of chronic nociceptive pain include osteoarthritis, rheumatoid arthritis, and musculoskeletal conditions (e.g., back pain), cancer pain, etc.

Neuropathic pain is defined as "pain initiated or caused by a primary lesion or dysfunction in the nervous system" by the International Association for the Study of Pain. Neuropathic pain is not associated with nociceptive stimulation, although the passage of nerve impulses that is ultimately perceived as pain by the brain is the same in both nociceptive and neuropathic pain. The term neuropathic pain encompasses a wide range of pain syndromes of diverse etiologies. The three most commonly diagnosed pain types of neuropathic nature are diabetic neuropathy, cancer neuropathy, and HIV pain. In addition, neuropathic pain is diagnosed in patients with a wide range of other disorders, including trigeminal neuralgia, post-herpetic neuralgia, traumatic neuralgia, phantom limb, as well as a number of other disorders of ill-defined or unknown origin.


Managing the spectrum of pain etiologies remains a major public health problem and both patients and clinicians are seeking improved strategies to effectively manage pain. No currently available therapies or drugs effectively treat all types of nociceptive and neuropathic pain states. The compounds of the present invention are novel CB₂ receptor modulators that have utility in treating pain, including nociceptive and neuropathic pain.

The location of CB₂ receptors on the surface of immune cells suggests a role for these receptors in immunomodulation and inflammation. Recent studies have demonstrated that CB₂ receptor ligands have immunomodulatory and anti-inflammatory properties. Therefore, compounds that interact with CB₂ receptors offer a unique pharmacotherapy for the treatment of immune and inflammatory disorders.

Summary

One aspect of the invention is directed towards compounds of formula (I), or

pharmaceutical salts, prodrugs, salts of prodrugs, or combinations thereof,



(I),

R₁ is alkoxyalkyl, alkoxyalkoxyalkyl, hydroxyalkyl, A, or A-alkylene-;

5 R₂ is hydrogen, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkyl, alkyl-S(O)₂-, aryl, arylalkyl, arylalkenyl, azidoalkyl, cyano, cycloalkyl, halo, haloalkyl, heteroaryl, heterocycle, -(CR₂₁R₂₂)_m-OH, R_aR_bN-, R_aR_bN-alkyl-, R_cR_dNC(O)-, or R₈-R₇-;

10 R₃ is hydrogen, alkoxy, alkoxyalkyl, alkyl, alkylcarbonyl, alkyl-S(O)₂-, aryl, arylalkyl, arylalkenyl, cyano, cycloalkyl, halo, haloalkyl, heteroaryl, heterocycle, -(CR₃₁R₃₂)_m-OH, R_aR_bN-, R_aR_bN-alkyl-, or R₈-R₇-; or

R₂ and R₃, together with the carbon atoms to which they are attached, form a 4-, 5-, 6-, or 7-membered monocyclic ring, optionally fused to a benzo or a monocyclic heteroaryl, said monocyclic ring contains zero, one, or two additional
15 double bonds, zero or one oxygen atom, and zero or one nitrogen atom as ring atoms; two non-adjacent atoms of said monocyclic ring can be optionally linked by an alkenylene bridge of 2, 3, or 4 carbon atoms, or optionally linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms, said monocyclic ring is independently unsubstituted or substituted with 1, 2, 3, 4, or 5 substituents independently selected
20 from the group consisting of oxo, alkyl, halo, -OH, -O(alkyl), and haloalkyl; two substituents on the same carbon atom of said monocyclic ring, together with the carbon atom to which they are attached, optionally form a 3-, 4-, 5-, or 6-membered monocyclic cycloalkyl ring, wherein the monocyclic cycloalkyl ring is optionally substituted with 1, 2, 3, 4, 5, or 6 substituents independently selected from the group
25 consisting of alkyl and haloalkyl;

with the proviso that when R₂ and R₃ are other than forming a ring with the carbon atoms to which they are attached, and R₁ is A or A-alkylene-, then R₄ is alkyl wherein the alkyl is optionally substituted with one substituent selected from the group consisting of alkoxy, alkoxycarbonyl, carboxy, halo, -OH, and R_eR_fN-, alkynyl,
30 cycloalkyl, cycloalkenyl, aryl, cycloalkylalkyl, cycloalkenylalkyl, arylalkyl, or R₁₀-L₂-R₉- wherein R₉ is aryl, cycloalkyl, or cycloalkenyl;

R₄ is alkyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, heterocycle, cycloalkylalkyl, cycloalkenylalkyl, arylalkyl, heteroarylalkyl, heterocyclealkyl, or R₁₀-L₂-R₉-; wherein the alkyl group is optionally substituted with one substituent selected from the group consisting of alkoxy, alkoxycarbonyl, carboxy, halo, -OH, and R_eR_fN-;

R₇, R₈, R₉, and R₁₀ are each independently aryl, cycloalkyl, cycloalkenyl, heteroaryl, or heterocycle;

R_a and R_b, at each occurrence, are each independently hydrogen, alkoxycarbonyl, alkyl, alkylcarbonyl, alkyl-S(O)₂-, or arylalkyl;

R_c and R_d, are each independently hydrogen or alkyl;

R_e and R_f, are each independently hydrogen, alkyl, or alkylcarbonyl;

A is a 4-, 5-, 6-, 7-, 8-, or 9-membered monocyclic heterocycle containing zero or one double bond, and one or two oxygen, and zero or one nitrogen as ring atoms, optionally fused with a monocyclic ring selected from benzo, cycloalkyl, cycloalkenyl, heterocycle and heteroaryl; wherein two non-adjacent atoms of each A can be optionally linked by an alkenylene bridge of 2, 3, or 4 carbon atoms, or optionally linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms; each A is independently unsubstituted or substituted with 1, 2, 3, 4, 5, or 6 substituents independently selected from the group consisting of oxo, alkyl, halo, -OH, -O(alkyl), and haloalkyl;

L₁ is a single bond or -NR_g-;


L₂ is a single bond, alkylene, or -O-;

R_g is hydrogen or alkyl;

R₂₁, R₂₂, R₃₁, and R₃₂, at each occurrence, are each independently hydrogen, alkyl, haloalkyl, or halo; and

m, at each occurrence, is independently 1, 2, 3, or 4.

Another aspect of the invention relates to compounds of formula (II), or pharmaceutical salts, prodrugs, salts of prodrugs, or combinations thereof,



(II)

wherein

R_{1a} is alkyl, haloalkyl, or cycloalkylalkyl;

R_{2a} is hydrogen or alkyl;

R_x represents optional substituent of phenyl, selected from the group consisting of alkyl, alkoxy, cyano, halo, haloalkoxy, hydroxyl, and haloalkyl; and

5 n is 1, 2, 3, 4, or 5.

Also comprised in the present application are pharmaceutical compositions comprising a therapeutically effective amount of a compound of the invention or a pharmaceutically acceptable salt thereof, in combination with a pharmaceutically acceptable carrier.

10 The present application also relates to a method of treating pain, nociceptive pain, and neuropathic pain in a mammal in need of such treatment comprising administering to the mammal a therapeutically effective amount of a compound of the invention or a pharmaceutically acceptable salt thereof.

Included in the present application is a method of treating a disorder selected
15 from the group consisting of inflammatory disorders, immune disorders, neurological disorders, cancers of the immune system, respiratory disorders, and cardiovascular disorders in a mammal in need of such treatment comprising administering to the mammal a therapeutically effective amount of a compound of the invention or a pharmaceutically acceptable salt thereof.

20 Included in the present application is a method of providing neuroprotection in a mammal in need comprising administering to the mammal a therapeutically effective amount of a compound of the invention or a pharmaceutically acceptable salt thereof.

25 Definition of Terms

For a variable that occurs more than one time in any substituent or in the compound of the invention or any other formulae herein, its definition on each occurrence is independent of its definition at every other occurrence. Combinations of substituents are permissible only if such combinations result in stable compounds.
30 Stable compounds are compounds which can be isolated from a reaction mixture.

All patents, patent applications, and literature references cited in the specification are herein incorporated by reference in their entirety. In the case of inconsistencies, the present disclosure, including definitions, will prevail.

As used in the specification and the appended claims, unless specified to the

contrary, the following terms have the meaning indicated:

The term "alkenyl" as used herein, means a straight or branched chain hydrocarbon containing from 2 to 10 carbons and containing at least one carbon-carbon double bond formed by the removal of two hydrogens. Representative
5 examples of alkenyl include, but are not limited to, ethenyl, 2-propenyl, 2-methyl-2-propenyl, 3-butenyl, 4-pentenyl, 5-hexenyl, 2-heptenyl, 2-methyl-1-heptenyl, and 3-decenyl.

The term "alkenylene" denotes a divalent group derived from a straight or branched chain hydrocarbon of 2, 3, or 4 carbon atoms and contains at least one
10 carbon-carbon double. Representative examples of alkylene include, but are not limited to, $-\text{CH}=\text{CH}-$ and $-\text{CH}_2\text{CH}=\text{CH}-$.

The term "alkoxy" as used herein, means an alkyl group, as defined herein, appended to the parent molecular moiety through an oxygen atom. Representative examples of alkoxy include, but are not limited to, methoxy, ethoxy, propoxy, 2-
15 propoxy, butoxy, tert-butoxy, pentyloxy, and hexyloxy.

The term "alkoxyalkoxy" as used herein, means an alkoxy group, as defined herein, appended to the parent molecular moiety through another alkoxy group, as defined herein. Representative examples of alkoxyalkoxy include, but are not limited to, tert-butoxymethoxy, 2-ethoxyethoxy, 2-methoxyethoxy, and methoxymethoxy.

20 The term "alkoxyalkoxyalkyl" as used herein, means an alkoxyalkoxy group, as defined herein, appended to the parent molecular moiety through an alkylene group, as defined herein. Representative examples of alkoxyalkoxyalkyl include, but are not limited to, tert-butoxymethoxymethyl, ethoxymethoxymethyl, (2-methoxyethoxy)methyl, and 2-(2-methoxyethoxy)ethyl.

25 The term "alkoxyalkyl" as used herein, means an alkoxy group, as defined herein, appended to the parent molecular moiety through an alkylene group, as defined herein. Representative examples of alkoxyalkyl include, but are not limited to, tert-butoxymethyl, 2-ethoxyethyl, 2-methoxyethyl, methoxymethyl, 2-methoxy-2-methylpropyl, and 3-methoxypropyl.

30 The term "alkoxycarbonyl" as used herein, means an alkoxy group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of alkoxycarbonyl include, but are not limited to, methoxycarbonyl, ethoxycarbonyl, and tert-butoxycarbonyl.

The term "alkoxycarbonylalkyl" as used herein, means an alkoxycarbonyl

group, as defined herein, appended to the parent molecular moiety through an alkylene group, as defined herein. Representative examples of alkoxycarbonylalkyl include, but are not limited to, 3-methoxycarbonylpropyl, 4-ethoxycarbonylbutyl, and 2-tert-butoxycarbonylethyl.

- 5 The term "alkyl" as used herein, means a straight or branched chain hydrocarbon containing from 1 to 10 carbon atoms. Representative examples of alkyl include, but are not limited to, methyl, ethyl, n-propyl, iso-propyl, n-butyl, sec-butyl, iso-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, n-hexyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 2,2-
- 10 dimethylpropyl, 1-methylpropyl, 1-ethylpropyl, 1,2,2-trimethylpropyl, 3-methylhexyl, 2,2-dimethylpentyl, 2,3-dimethylpentyl, n-heptyl, n-octyl, n-nonyl, and n-decyl.

- The term "alkylcarbonyl" as used herein, means an alkyl group, as defined herein, appended to the parent molecular moiety through a carbonyl group, as defined herein. Representative examples of alkylcarbonyl include, but are not limited to,
- 15 acetyl, 1-oxopropyl, 2,2-dimethyl-1-oxopropyl, 1-oxobutyl, and 1-oxopentyl.

- The term "alkylcarbonylalkyl" as used herein, means an alkylcarbonyl group, as defined herein, appended to the parent molecular moiety through an alkylene group, as defined herein. Representative examples of alkylcarbonylalkyl include, but are not limited to, 2-oxopropyl, 3,3-dimethyl-2-oxopropyl, 3-oxobutyl, and 3-
- 20 oxopentyl.

- The term "alkylcarbonyloxy" as used herein, means an alkylcarbonyl group, as defined herein, appended to the parent molecular moiety through an oxygen atom. Representative examples of alkylcarbonyloxy include, but are not limited to, acetyloxy, ethylcarbonyloxy, and tert-butylcarbonyloxy.

- 25 The term "alkylene" means a divalent group derived from a straight or branched chain hydrocarbon of from 1 to 10 carbon atoms. Representative examples of alkylene include, but are not limited to, -CH₂-, -CH(CH₃)-, -CH(C₂H₅)-, -CH(CH(CH₃)(C₂H₅))-, -C(H)(CH₃)CH₂CH₂-, -C(CH₃)₂-, -CH₂CH₂-, -CH₂CH₂CH₂-, -CH₂CH₂CH₂CH₂-, and -CH₂CH(CH₃)CH₂-.

- 30 The term "alkylsulfinyl" as used herein, means an alkyl group, as defined herein, appended to the parent molecular moiety through a sulfinyl group, as defined herein. Representative examples of alkylsulfinyl include, but are not limited to, methylsulfinyl and ethylsulfinyl.

 The term "alkylsulfinylalkyl" as used herein, means an alkylsulfinyl group, as

defined herein, appended to the parent molecular moiety through an alkylene group, as defined herein. Representative examples of alkylsulfinylalkyl include, but are not limited to, methylsulfinylmethyl and ethylsulfinylmethyl.

The term "alkynyl" as used herein, means a straight or branched chain
5 hydrocarbon group containing from 2 to 10 carbon atoms and containing at least one carbon-carbon triple bond. Representative examples of alkynyl include, but are not limited, to acetylenyl, 1-propynyl, 2-propynyl, 1-propyl-pent-3-ynyl, 3-butynyl, 2-pentynyl, and 1-butynyl.

The term "aryl," as used herein, means phenyl, a bicyclic aryl or a tricyclic
10 aryl. The bicyclic aryl is naphthyl, or a phenyl fused to a cycloalkyl, or a phenyl fused to a cycloalkenyl. Representative examples of the bicyclic aryl include, but are not limited to, dihydroindanyl, indenyl, naphthyl, dihydronaphthalenyl, and tetrahydronaphthalenyl. The tricyclic aryl is exemplified by a bicyclic aryl fused to a monocyclic cycloalkyl, or a bicyclic aryl fused to a monocyclic cycloalkenyl, or a
15 bicyclic aryl fused to a phenyl. Representative examples of tricyclic aryl ring include, but are not limited to, anthracene, phenanthrene, dihydroanthracenyl, fluorenyl, 1,2-dihydroacenaphthylenyl, and tetrahydrophenanthrenyl. The phenyl, bicyclic and tricyclic aryls are attached to the parent molecular moiety through any carbon atom contained within the phenyl, bicyclic and tricyclic aryls respectively.

20 The term "arylalkyl" as used herein, means an aryl group, as defined herein, appended to the parent molecular moiety through an alkylene group, as defined herein. Representative examples of arylalkyl include, but are not limited to, benzyl, 2-phenylethyl, 1-methyl-3-phenylpropyl, 2-methyl-1-phenylbutyl, 1-phenylpropyl, 3-phenylpropyl, and 2-naphth-2-ylethyl.

25 The term "azido" as used herein, means a $-N_3$ group.

The term "azidoalkyl" as used herein, means an azido group, as defined herein, appended to the parent molecular moiety through an alkylene group, as defined herein.

The term "carbonyl" as used herein, means a $-C(O)-$ group.

30 The term "carboxy" as used herein, means a $-CO_2H$ group.

The term "carboxyalkyl" as used herein, means a carboxy group, as defined herein, appended to the parent molecular moiety through an alkylene group, as defined herein. Representative examples of carboxyalkyl include, but are not limited to, carboxymethyl, 2-carboxyethyl, and 3-carboxypropyl.

The term "cyano" as used herein, means a -CN group.

The term "cyanoalkyl" as used herein, means a cyano group, as defined herein, appended to the parent molecular moiety through an alkylene group, as defined herein. Representative examples of cyanoalkyl include, but are not limited to, cyanomethyl, 2-cyanoethyl, and 3-cyanopropyl.

The term "cycloalkenyl" as used herein, means a monocyclic or bicyclic ring system containing zero heteroatoms in the ring. The monocyclic cycloalkenyl has three-, four-, five-, six-, seven- or eight carbon atoms and zero heteroatoms. The three or four-membered ring systems have one double bond, the five- or six-membered ring systems have one or two double bonds, and the seven- or eight-membered ring systems have one, two or three double bonds. Representative examples of monocyclic ring systems include, but are not limited to, 2-cyclohexen-1-yl, 3-cyclohexen-1-yl, 2,4-cyclohexadien-1-yl and 3-cyclopenten-1-yl. Bicyclic ring systems are exemplified by a monocyclic cycloalkenyl ring fused to a monocyclic cycloalkyl ring, or a monocyclic cycloalkenyl ring fused to a monocyclic cycloalkenyl ring. Representative examples of bicyclic ring systems include, but are not limited to 3a, 4, 5, 6, 7, 7a-hexahydro-1H-indenyl, 4,5,6,7-tetrahydro-3aH-indene, and octahydronaphthalenyl. The monocyclic or the bicyclic cycloalkenyl ring can be appended to the parent molecular moiety through any substitutable carbon atom within the monocyclic or the bicyclic cycloalkenyl.

The term "cycloalkenylalkyl" as used herein, means a cycloalkenyl group as defined herein, appended to the parent molecular moiety through an alkylene group, as defined herein.

The term "cycloalkyl" as used herein, means a monocyclic, or a bicyclic ring system, or a spirocyclic cycloalkyl. The monocyclic cycloalkyl is a carbocyclic ring system containing 3, 4, 5, 6, 7, or 8 carbon atoms, zero heteroatoms and zero double bonds. Examples of monocyclic ring systems include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, and cyclooctyl. Bicyclic ring systems are exemplified by a monocyclic cycloalkyl ring fused to a monocyclic cycloalkyl ring. Representative examples of bicyclic ring systems include, but are not limited to, bicyclo[4.1.0]heptane, bicyclo[6.1.0]nonane, octahydroindene, and decahydronaphthalene. Spirocyclic cycloalkyl is exemplified by a monocyclic cycloalkyl ring wherein two of the substituents on the same carbon atom of the ring, form a 4-, 5-, or 6-membered monocyclic cycloalkyl. An example of a spirocyclic

cycloalkyl is spiro[2.5]octane. The monocyclic, bicyclic and spirocyclic cycloalkyl groups of the present invention can be appended to the parent molecular moiety through any substitutable carbon atom of the groups.

The term "cycloalkylalkyl" as used herein, means a cycloalkyl group, as defined herein, appended to the parent molecular moiety through an alkylene group, as defined herein. Representative examples of cycloalkylalkyl include, but are not limited to, cyclopentylmethyl, cyclohexylmethyl, cyclopropylmethyl, and 1-cyclopropylethyl.

The term "formyl" as used herein, means a -C(O)H group.

The term "formylalkyl" as used herein, means a formyl group, as defined herein, appended to the parent molecular moiety through an alkylene group, as defined herein. Representative examples of formylalkyl include, but are not limited to, formylmethyl and 2-formylethyl.

The term "halo" or "halogen" as used herein, means -Cl, -Br, -I or -F.

The term "haloalkoxy" as used herein, means an alkoxy group, as defined herein, in which one, two, three, four, five or six hydrogen atoms are replaced by halogen. Representative examples of haloalkoxy include, but are not limited to, trifluoromethoxy, difluoromethoxy, 2,2,2-trifluoroethoxy, and 2,2-difluoroethoxy.

The term "haloalkyl" as used herein, means an alkyl group, as defined herein, in which one, two, three, four, five, six, or seven hydrogen atoms are replaced by halogen. Representative examples of haloalkyl include, but are not limited to, chloromethyl, 2-fluoroethyl, 2,2-difluoroethyl, trifluoromethyl, 2,2,2-trifluoroethyl, difluoromethyl, pentafluoroethyl, 2-chloro-3-fluoropentyl, and 2-iodoethyl.

The term "heteroaryl," as used herein, means a monocyclic heteroaryl or a bicyclic heteroaryl. The monocyclic heteroaryl is a 5 or 6 membered ring containing at least one heteroatom independently selected from the group consisting of O, N, and S. The 5-membered ring contains two double bonds and one, two, three, or four heteroatoms. The 6 membered ring contains three double bonds and one, two, three or four heteroatoms. Representative examples of monocyclic heteroaryl include, but are not limited to, furanyl, imidazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, oxazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrazolyl, pyrrolyl, tetrazolyl, thiadiazolyl, thiazolyl, thienyl, triazolyl, and triazinyl. The bicyclic heteroaryl is exemplified by a monocyclic heteroaryl fused to phenyl, or a monocyclic heteroaryl fused to a monocyclic cycloalkyl, or a monocyclic heteroaryl fused to a monocyclic

cycloalkenyl, or a monocyclic heteroaryl fused to a monocyclic heteroaryl, or a monocyclic heteroaryl fused to a monocyclic heterocycle. Representative examples of bicyclic heteroaryl include, but are not limited to, benzofuranyl, benzoxadiazolyl, 1,3-benzothiazolyl, benzimidazolyl, benzodioxolyl, benzothienyl, chromenyl, 5 cinnolinyl, furopyridine, indolyl, indazolyl, isoindolyl, isoquinolinyl, naphthyridinyl, oxazolopyridine, quinolinyl, thienopyridine and thienopyridinyl. The monocyclic and the bicyclic heteroaryl groups are connected to the parent molecular moiety through any substitutable carbon atom or any substitutable nitrogen atom contained within the groups.

10 The term "heteroarylalkyl" as used herein, means a heteroaryl group as defined herein, appended to the parent molecular moiety through an alkylene group, as defined herein. An example of heteroarylalkyl is 3-thienylpropyl.

The term "heterocycle" or "heterocyclic" as used herein, refers to a monocyclic, bicyclic, tricyclic, or a spirocyclic ring system which contains at least 15 one heteroatom. The monocyclic heterocycle is a 3, 4, 5, 6, 7, or 8- membered ring containing at least one heteroatom independently selected from the group consisting of O, N, and S. The 3 or 4 membered ring contains 1 heteroatom selected from the group consisting of O, N and S, and optionally one double bond. The 5-membered ring contains zero or one double bond, and one, two or three heteroatoms in the ring 20 selected from the group consisting of O, N and S. The 6, 7, or 8-membered ring contains zero, one, or two double bonds, and one, two, or three heteroatoms in the ring selected from the group consisting of O, N and S. Representative examples of monocyclic heterocycle include, but are not limited to, azetidiny, azepanyl, aziridinyl, diazepanyl, 1,3-dioxanyl, 1,4-dioxanyl, 1,3-dioxolanyl, 4,5- 25 dihydroisoxazol-5-yl, 3,4-dihydropyran-6-yl, 1,3-dithiolanyl, 1,3-dithianyl, imidazoliny, imidazolidiny, isothiazoliny, isothiazolidiny, isoxazoliny, isoxazolidiny, morpholiny, oxadiazoliny, oxadiazolidiny, oxazoliny, oxazolidiny, oxetanyl, piperaziny, piperidiny, pyranyl, pyrazoliny, pyrazolidiny, pyrroliny, pyrrolidinyl, tetrahydrofuranyl, tetrahydropyrany, tetrahydrothienyl, thiadiazoliny, 30 thiadiazolidiny, thiazoliny, thiazolidiny, thiomorpholiny, 1,1-dioxidothiomorpholiny (thiomorpholine sulfone), thiopyrany, and trithianyl. The bicyclic heterocycle of the present invention is exemplified by a monocyclic heterocycle fused to a phenyl group, or a monocyclic heterocycle fused to a monocyclic cycloalkylgroup, or a monocyclic heterocycle fused to a monocyclic

cycloalkenyl group, or a monocyclic heterocycle fused to a monocyclic heterocycle group. Representative examples of bicyclic heterocycle include, but are not limited to, 1,3-benzodioxol-4-yl, 1,3-benzodithiolyl, 2,3-dihydro-1,4-benzodioxinyl, 2,3-dihydro-1-benzofuranyl, 2,3-dihydro-1-benzothienyl, 2,3-dihydro-1H-indolyl, and
5 1,2,3,4-tetrahydroquinolyl. Spirocyclic heterocycle means a 4-, 5-, 6-, 7-, or 8-membered monocyclic heterocycle ring wherein two of the substituents on the same carbon atom form a 4-, 5-, or 6-membered monocyclic cycloalkyl, wherein the cycloalkyl is optionally substituted with 1, 2, 3, 4, or 5 alkyl groups. One example of a spiroheterocycle is 5-oxaspiro[3,4]octane. The tricyclic heterocycle is a bicyclic
10 heterocycle fused to a phenyl, or a bicyclic heterocycle fused to a monocyclic cycloalkyl, or a bicyclic heterocycle fused to a monocyclic cycloalkenyl, or a bicyclic heterocycle fused to a monocyclic heterocycle. Representative examples of tricyclic heterocycle include, but are not limited to, 2,3,4,4a,9,9a-hexahydro-1H-carbazolyl, 5a,6,7,8,9,9a-hexahydrodibenzo[b,d]furanyl, and 5a,6,7,8,9,9a-
15 hexahydrodibenzo[b,d]thienyl. The monocyclic, bicyclic, tricyclic, and spirocyclic heterocycle groups are connected to the parent molecular moiety through any substitutable carbon atom or any substitutable nitrogen atom contained within the group.

The term "heterocyclealkyl" as used herein, means a heterocycle group as
20 defined herein, appended to the parent molecular moiety through an alkylene group, as defined herein.

The term "hydroxy" as used herein, means an -OH group.

The term "hydroxyalkyl" as used herein, means at least one hydroxy group, as defined herein, is appended to the parent molecular moiety through an alkylene group,
25 as defined herein. Representative examples of hydroxyalkyl include, but are not limited to, hydroxymethyl, 2-hydroxyethyl, 3-hydroxypropyl, 3-hydroxy-3-methylbutyl, 2,3-dihydroxypentyl, and 2-ethyl-4-hydroxyheptyl.

The aryl, cycloalkyl, cycloalkenyl, heterocycle, or heteroaryl moieties of this invention, as a substituent, or as part of a substituent, is each independently
30 unsubstituted or substituted with 1, 2, 3, 4, 5, or 6 substituents as described herein below, unless otherwise noted. The optional substituents are selected from the group consisting of alkenyl, alkoxy, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkoxysulfonyl, alkyl, alkylcarbonyl, alkylcarbonylalkyl, alkylcarbonyloxy, alkylsulfinyl, alkylsulfinylalkyl, alkyl-S(O)₂-,

alkyl-S(O)₂-alkyl-, alkyl-S-, alkyl-S-alkyl-, alkynyl, carboxy, carboxyalkyl, cyano, cyanoalkyl, formyl, formylalkyl, halogen, haloalkyl, haloalkoxy, hydroxy, hydroxyalkyl, oxo, -SH, N(O)₂, NZ₁Z₂-alkylene-O-, -NZ₁Z₂, and (NZ₃Z₄)carbonyl. The exceptions are the cycloalkenyl, the aryl, the heteroaryl, and the heterocycle

5 moieties as represented by R₄ and R₉ wherein the optional substituents are selected from the group consisting of alkenyl, alkoxy, alkoxyalkoxy, alkoxyalkoxyalkyl, alkoxyalkyl, alkyl, alkylcarbonyl, alkylcarbonylalkyl, alkylcarbonyloxy, alkyl-S-, alkyl-S-alkyl-, alkynyl, cyano, cyanoalkyl, formyl, formylalkyl, halogen, haloalkyl, haloalkoxy, hydroxy, hydroxyalkyl, oxo, -SH, N(O)₂, NZ₁Z₂-alkylene-O-, and

10 -NZ₁Z₂.

The term "hydroxy-protecting group" or "O-protecting group" means a substituent which protects hydroxy groups against undesirable reactions during synthetic procedures. Examples of hydroxy-protecting groups include, but are not limited to, substituted methyl ethers, for example, methoxymethyl, benzyloxymethyl,

15 2-methoxyethoxymethyl, 2-(trimethylsilyl)-ethoxymethyl, benzyl, and triphenylmethyl; tetrahydropyranyl ethers; substituted ethyl ethers, for example, 2,2,2-trichloroethyl and t-butyl; silyl ethers, for example, trimethylsilyl, t-butyldimethylsilyl and t-butyldiphenylsilyl; cyclic acetals and ketals, for example, methylene acetal, acetone and benzylidene acetal; cyclic ortho esters, for example,

20 methoxymethylene; cyclic carbonates; and cyclic boronates. Commonly used hydroxy-protecting groups are disclosed in T.W. Greene and P.G.M. Wuts, Protective Groups in Organic Synthesis, 3rd edition, John Wiley & Sons, New York (1999).

The term "nitrogen protecting group" as used herein, means those groups intended to protect an amino group against undesirable reactions during synthetic

25 procedures. Preferred nitrogen protecting groups are acetyl, benzoyl, benzyl, benzyloxycarbonyl (Cbz), formyl, phenylsulfonyl, tert-butoxycarbonyl (Boc), tert-butylacetyl, trifluoroacetyl, and triphenylmethyl (trityl).

The term "NZ₁Z₂" as used herein, means two groups, Z₁ and Z₂, which are appended to the parent molecular moiety through a nitrogen atom. Z₁ and Z₂ are each

30 independently hydrogen, alkyl, alkoxyalkyl, alkylcarbonyl, haloalkyl, or formyl. In certain instances, Z₁ and Z₂ taken together with the nitrogen atom to which they are attached form a heterocyclic ring, optionally substituted with 1, 2, 3, or 4 substituents selected from alkyl, hydroxy and haloalkyl. Representative examples of NZ₁Z₂ include, but are not limited to, amino, methylamino, acetylamino, acetylmethylamino,

phenylamino, benzylamino, azetidiny, pyrrolidinyl and piperidinyl.

The term "NZ₃Z₄" as used herein, means two groups, Z₃ and Z₄, which are appended to the parent molecular moiety through a nitrogen atom. Z₃ and Z₄ are each independently hydrogen, alkyl, haloalkyl, phenyl or benzyl wherein the phenyl moiety is optionally substituted with 1, 2, 3, or 4 substituents selected from alkyl, hydroxy and haloalkyl. Representative examples of NZ₃Z₄ include, but are not limited to, amino, methylamino, phenylamino and benzylamino.

The term "oxo" as used herein, means a =O moiety.

The term "sulfinyl" as used herein, means a -S(O)- group.

The term "tautomer" as used herein means a proton shift from one atom of a compound to another atom of the same compound wherein two or more structurally distinct compounds are in equilibrium with each other.

The term "pharmaceutically acceptable prodrug" or "prodrug," as used herein, represents those prodrugs of the compounds of the present invention which are, within the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response, and the like, commensurate with a reasonable benefit/risk ratio, and effective for their intended use. Prodrugs of the present invention may be rapidly transformed in vivo to compounds of formula (I), for example, by hydrolysis in blood.


Detailed Description

Compounds of the invention have the formula (I) as described above.

Particular values of variable groups in compounds of formula (I) are as follows. Such values may be used where appropriate with any of the other values, definitions, claims or embodiments defined hereinbefore or hereinafter.

In compounds of formula (I), R₁ is alkoxyalkyl, alkoxyalkoxyalkyl, hydroxyalkyl, A, or A-alkylene- wherein A is as disclosed in the Summary. Embodiments of the present invention include compounds wherein R₁ is A or A-alkylene- wherein A is as disclosed in the Summary. Some examples of A are those that are represented by formula (i), (ii), (iii), (iv), (v) and (vi), wherein each ring is independently unsubstituted or substituted as described in the Summary. Certain examples of the optional substituents of A include, but are not limited to, alkyl such as C₁₋₆ alkyl, and oxo. The alkylene moiety of A-alkylene-, for example, is C₁₋₆ alkylene. Further example of the alkylene moiety of A-alkylene- is C₁₋₃ alkylene.

Yet further example is C₁-C₂ alkylene.




Other examples of compounds of formula (I) include those wherein R₁ is alkoxyalkyl, alkoxyalkoxyalkyl, or hydroxyalkyl

5 R₂ is hydrogen, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkyl, alkyl-S(O)₂-, aryl, arylalkyl, arylalkenyl, azidoalkyl, cyano, cycloalkyl, halo, haloalkyl, heteroaryl, heterocycle, -(CR₂₁R₂₂)_m-OH, R_aR_bN-, R_aR_bN-alkyl-, R_cR_dNC(O)-, or R₈-R₇-; wherein R₂₁, R₂₂, m, R_a, R_b, R_c, R_d, R₇, and R₈ are as described in the Summary, and the optional substituents of aryl, cycloalkyl, heteroaryl
 10 and heterocycle moieties are as disclosed in the Definition of Terms. Certain examples of compounds of formula (I) include those wherein R₂ is hydrogen, alkoxycarbonyl, alkyl, aryl (for example, optionally substituted phenyl), halo, haloalkyl, or -(CR₂₁R₂₂)_m-OH; wherein R₂₁, R₂₂, and m are as described in the Summary, and the optional substituents of the aryl moiety are as disclosed in the
 15 Definition of Terms, for example, the optional substituents of the aryl moiety are selected from the group consisting of alkyl and halo. Included, but are not limited to, are compounds in which R₂₁ and R₂₂ are hydrogen and m is 1. Embodiments of the present invention include compounds in which R₂ is hydrogen or alkyl.

R₃ is hydrogen, alkoxy, alkoxyalkyl, alkyl, alkylcarbonyl, alkyl-S(O)₂-, aryl,
 20 arylalkyl, arylalkenyl, cyano, cycloalkyl, halo, haloalkyl, heteroaryl, heterocycle, -(CR₃₁R₃₂)_m-OH, R_aR_bN-, R_aR_bN-alkyl-, or R₈-R₇-; wherein R₃₁, R₃₂, m, R_a, and R_b, are as described in the Summary, and the optional substituents of aryl, cycloalkyl, heteroaryl and heterocycle moieties are as disclosed in the Definition of Terms. Examples of compounds of formula (I) include, but are not limited to, those wherein
 25 R₃ is hydrogen, alkyl (for example, methyl, tert-butyl), alkylcarbonyl, aryl (for example, phenyl), cycloalkyl (for example, cyclopropyl), halo, haloalkyl, heterocycle, or -(CR₃₁R₃₂)_m-OH, wherein R₃₁, R₃₂, and m are as disclosed in the Summary. The optional substituents of aryl, cycloalkyl, and heterocycle moieties are as disclosed in the Definition of Terms, for example, the optional substituents are selected from the
 30 group consisting of alkyl and halo. Non limiting examples of R₃₁ and R₃₂ (R₃₁ and R₃₂ can be the same or different) are alkyl (for example, methyl) or haloalkyl (for


example, 2-iodoethyl or trifluoromethyl). m , for example, is 1. Embodiments of the present invention include compounds in which R_3 is alkyl (for example, methyl or tert-butyl) or $-(CR_{31}R_{32})_m-OH$. Other examples include those wherein R_3 is $-(CR_{31}R_{32})_m-OH$, wherein m is 1, and R_{31} and R_{32} are alkyl (such as, but not limited to, methyl) or haloalkyl (such as, but not limited to, trifluoromethyl).

In another embodiment, R_2 and R_3 , together with the carbon atoms to which they are attached, form a 4-, 5-, 6-, or 7-membered monocyclic ring as described in the Summary. Embodiments of the present invention include compounds of formula (I) wherein R_2 and R_3 , together with the carbon atoms to which they are attached, form a monocyclic ring as described in the Summary, containing zero heteroatoms in the ring. Formulae (vii), (viii), (ix), (x), (xi), (xii), (xiii), and (xiv) represent some of these rings that can be formed by R_2 , R_3 , together with the carbon atoms to which they are attached.



Yet other compounds of the present invention include those wherein R_2 and R_3 , together with the carbon atoms to which they are attached, form a monocyclic ring as described in the Summary, containing zero or one additional double bond, zero oxygen atom and zero nitrogen atom as ring atoms; and two non-adjacent atoms of said monocyclic ring are linked by an alkenylene bridge of 2, 3, or 4 carbon atoms, or linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms. Examples include, but are not limited to, (xii), (xiii) and (xiv).

Yet other compounds of the present invention include those wherein R_2 and R_3 , together with the carbon atoms to which they are attached, form a monocyclic ring as described in the Summary, containing zero or one additional double bond, and one oxygen atom and zero or one nitrogen atom as ring atoms. Examples of such monocyclic ring include, but are not limited to, formula (xv) – (xxix)



Each monocyclic ring formed by R_2 , R_3 , and the carbon atoms to which they are attached is independently unsubstituted or substituted as described in the Summary, for example, these rings can be independently unsubstituted or substituted with 1, 2, 3, 4, 5 or 6 substituents independently selected from alkyl such as C_{1-6} alkyl, hydroxy, and oxo. Such rings are optionally fused with benzo or a monocyclic heteroaryl (for example, 1,3,4-oxadiazole, pyrrole, furan, and the like).

R_4 is alkyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, heterocycle, cycloalkylalkyl, cycloalkenylalkyl, arylalkyl, heteroarylalkyl, heterocyclealkyl, or R_{10} - L_2 - R_9 -; wherein the alkyl group is optionally substituted with one substituent selected from the group consisting of alkoxy, alkoxy carbonyl, carboxy, halo, -OH, and $R_e R_f N$ -. In one embodiment, R_4 is optionally substituted aryl. In another embodiment, R_4 is optionally substituted phenyl. Examples of the optional substituents of R_4 include, but are not limited to, alkyl, alkoxy, cyano, halo, haloalkoxy, hydroxy, and haloalkyl. Other examples of compounds of formula (I) are those wherein R_4 is heteroaryl or heterocycle, each of which is optionally substituted as described in the Definition of Terms. Yet other examples are those wherein R_4 is alkyl optionally substituted with $R_e R_f N$ - wherein R_e and R_f are as disclosed in the Summary. Further examples are those wherein R_4 is optionally substituted cycloalkyl wherein the optional substituents are as discussed in Definition of Terms.

L_1 is a single bond or $-NR_g-$ wherein R_g is hydrogen or alkyl. Certain compounds of the present invention include those wherein L_1 is a single bond. Yet others are those wherein L_1 is $-NR_g-$ wherein R_g is hydrogen or alkyl. Other examples include those wherein L_1 is $-NH-$.

It is appreciated that the present invention contemplates compounds of formula (I) with combinations of the above embodiments, including particular, more particular and preferred embodiments.

Accordingly, one aspect of the invention relates to a group of compounds of

formula (I), or pharmaceutically acceptable salts thereof, wherein R_1 is A or A-alkenyl-, R_2 is hydrogen, alkoxycarbonyl, alkyl, aryl, halo, haloalkyl, or $-(CR_{21}R_{22})_m-OH$; R_3 is hydrogen, alkyl, aryl, cycloalkyl, halo, haloalkyl, heterocycle, or $-(CR_{31}R_{32})_m-OH$; R_4 is alkyl wherein the alkyl is optionally substituted with one

5 substituent selected from the group consisting of alkoxy, alkoxycarbonyl, carboxy, halo, -OH, and R_eR_fN- , alkynyl, cycloalkyl, cycloalkenyl, aryl, cycloalkylalkyl, cycloalkenylalkyl, arylalkyl, or $R_{10}-L_2-R_9-$, wherein R_9 is aryl, cycloalkyl, or cycloalkenyl; and A, L_1 , R_{21} , R_{22} , R_{31} , R_{32} , m, R_e , R_f , L_2 , and R_{10} are as disclosed in the Summary, and the optional substituents of the aryl, cycloalkyl, and cycloalkenyl

10 moieties are as disclosed in the Definition of Terms. A, for example, is formula (i), (ii), (iii), (iv), (v), or (vi), wherein each ring is independently unsubstituted or substituted as described in the Summary. R_{21} and R_{22} , for example, are hydrogen. m, for example, is 1. R_{31} and R_{32} are, for example, independently alkyl (such as methyl) or haloalkyl (such as 2-iodoethyl or trifluoromethyl). In one embodiment, R_3 is alkyl

15 (for example, methyl or tert-butyl) or $-(CR_{31}R_{32})_m-OH$, and R_2 is hydrogen or alkyl wherein R_{31} , R_{32} and m are as described in the Summary. Examples include those wherein R_2 is hydrogen or alkyl (for example, methyl), and R_3 is $-(CR_{31}R_{32})_m-OH$ wherein m is 1, and R_{31} and R_{32} are alkyl (such as, but not limited to, methyl) or haloalkyl (such as, but not limited to, trifluoromethyl). Other examples include those

20 wherein R_2 is hydrogen or alkyl (for example, methyl), and R_3 is alkyl (for example, tert-butyl). Yet other examples include those wherein R_2 is hydrogen or alkyl (for example, methyl), and R_3 is $-(CR_{31}R_{32})_m-OH$, wherein R_{31} and R_{32} are alkyl (for example, methyl), and m is 1. R_4 , for example, is aryl, cycloalkyl, arylalkyl, cycloalkylalkyl, or $R_{10}-L_2-R_9-$, wherein R_9 is aryl or cycloalkyl. In other

25 embodiments, R_4 is aryl or $R_{10}-L_2-R_9-$, wherein R_9 is aryl. The optional substituents of aryl and cycloalkyl moieties are as disclosed in the Definition of Terms. An example of the aryl moiety is phenyl. Other examples of the aryl moiety include, but are not limited to, naphthyl and 1,2-dihydroacenaphthylenyl. The optional substituents of aryl and cycloalkyl moieties are as disclosed in the Definition of

30 Terms. Examples of the optional substituents of aryl and cycloalkyl moieties include, but are not limited to, alkyl, alkoxy, cyano, halo, haloalkoxy, hydroxy, and haloalkyl.

Another aspect of the invention provides a group of compounds of formula (I), or pharmaceutically acceptable salts thereof, wherein wherein R_1 is A or A-alkenyl-, R_2 and R_3 , together with the carbon atoms to which they are attached form a

monocyclic ring, and A, R₄, L₁, and said monocyclic ring are as described in the Summary. Some examples of A are as described herein above. Certain examples of the monocyclic ring formed by R₂, R₃, and the carbon atoms to which they are attached are represented by formulae (vii), (viii), and (x)-(xxix), each of which is optionally substituted with substituents as described in the Summary, and each of which is optionally fused with benzo or a monocyclic heteroaryl (for example, oxadiazole, furan or pyrrole). One example of such fused ring is represented by formula (ix). Examples of the optional substituents on the rings formed by R₂, R₃, and the carbon atoms to which they are attached include, but are not limited to, alkyl such as C₁₋₆ alkyl, hydroxy, and oxo.

Yet another aspect of the invention relates to a group of compounds of formula (I) or pharmaceutically acceptable salts thereof, wherein R₁ is alkoxyalkyl, alkoxyalkoxyalkyl, or hydroxyalkyl, R₂ and R₃, together with the carbon atoms to which they are attached, form a 4-, 5-, 6-, or 7-membered monocyclic ring, optionally fused to a benzo or a monocyclic heteroaryl, said monocyclic ring contains zero or one additional double bond, zero oxygen atom and zero nitrogen atom as ring atoms; two non-adjacent atoms of said monocyclic ring are linked by an alkenylene bridge of 2, 3, or 4 carbon atoms, or linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms, said monocyclic ring is independently unsubstituted or substituted with 1, 2, 3, 4, or 5 substituents independently selected from the group consisting of oxo, alkyl, halo, -OH, -O(alkyl), and haloalkyl; two substituents on the same carbon atom of said monocyclic ring, together with the carbon atom to which they are attached, optionally form a 3-, 4-, 5-, or 6-membered monocyclic cycloalkyl ring, wherein the monocyclic cycloalkyl ring is optionally substituted with 1, 2, 3, 4, 5, or 6 substituents independently selected from the group consisting of alkyl and haloalkyl; and R₄, and L₁, are as described in the Summary. Some examples of the ring formed by R₂, R₃, and the carbon atoms to which they are attached are represented by formulae (xii), (xiii) and (xiv), each of which is optionally substituted as described in the Summary. Examples of the optional substituents include, but are not limited to, alkyl such as C₁₋₆ alkyl, hydroxy, and oxo.

A further aspect of the invention provides a group of compounds of formula (I) or pharmaceutically acceptable salts thereof, wherein R₁ is alkoxyalkyl, alkoxyalkoxyalkyl, or hydroxyalkyl, R₂ and R₃, together with the carbon atoms to which they are attached, form a 4-, 5-, 6-, or 7-membered monocyclic ring as

described in the Summary, containing zero or one additional double bond, and one oxygen atom, and zero or one nitrogen atom as ring atoms; and R_4 , and L_1 , are as described in the Summary. Some examples of the monocyclic ring formed by R_2 , R_3 , and the carbon atoms to which they are attached are represented by formula (xv)-

5 (xxix), each of which is optionally substituted as described in the Summary.

Examples of the optional substituents include, but are not limited to, alkyl such as C_{1-6} alkyl, hydroxy, and oxo.

For the above three groups of compounds of formula (I) described, R_4 is alkyl, alkynyl, or alkyl substituted with NR_eR_f , wherein R_e and R_f are as described in the
10 Summary.

For the above three groups of compounds of formula (I) described, other examples of R_4 are aryl, arylalkyl, cycloalkyl, cycloalkylalkyl, or R_{10} - L_2 - R_9 - wherein R_9 is aryl or cycloalkyl, L_2 and R_{10} are as disclosed in the Summary, and the optional substituents of the aryl moiety is as disclosed in the Definition of Terms.

15 Embodiments of the present invention include compounds of formula (I) wherein R_4 is aryl or R_{10} - L_2 - R_9 -, wherein R_9 is aryl wherein the aryl moiety is optionally substituted as disclosed in the Definition of Terms. Preferably, R_4 is optionally substituted aryl. An example of the aryl moiety is phenyl. Other examples of the aryl moiety include naphthyl and 1,2-dihydroacenaphthylene. Examples of the optional
20 substituents include, but are not limited to, alkyl, alkoxy, cyano, halo, haloalkoxy, hydroxy, and haloalkyl.

For the above three groups of compounds of formula (I) described, other examples of R_4 are heteroaryl, heterocycle, heteroarylalkyl, heterocyclealkyl, or R_{10} - L_2 - R_9 - wherein R_9 is heteroaryl or heterocycle, L_2 and R_{10} are as disclosed in the
25 Summary, and the optional substituents of the heteroaryl and heterocycle moieties are as disclosed in the Definition of Terms. Examples of the heteroaryl moiety include pyridinyl, thienyl, benzofuranyl, and quinolinyl. Examples of the heterocycle moiety include 3,4-dihydropyran-6-yl, tetrahydrofuranyl, tetrahydropyranyl, 1,3-benzodioxol-4-yl, and 2,3-dihydro-1-benzofuranyl. Examples of the optional
30 substituents include, but are not limited to, alkyl, alkoxy, oxo, cyano, halo, haloalkoxy, hydroxy, and haloalkyl.


For all the foregoing embodiments described, examples of a subgroup include those wherein L_1 is a single bond. Yet other examples of a subgroup include those wherein L_1 is $-NR_g$ - wherein R_g is hydrogen or alkyl. Yet other examples of a

subgroup include those wherein L_1 is $-NH-$.

A further aspect of the invention relates to compounds of formula (I) or pharmaceutically acceptable salts thereof, wherein L_1 is a single bond; R_1 is alkoxyalkyl, alkoxyalkoxyalkyl, or hydroxyalkyl, R_2 is hydrogen or alkyl, R_3 is

5 $-(CR_{31}R_{32})_m-OH$; R_4 is phenyl; R_{31} and R_{32} are methyl; and m is 1.

Yet a further aspect of the invention provides compounds of formula (II) or pharmaceutically acceptable salts thereof,



(II)

10 wherein

R_{1a} is alkyl, haloalkyl, or cycloalkylalkyl;

R_{2a} is hydrogen or alkyl;

R_x represents optional substituent of phenyl, selected from the group consisting of alkyl, alkoxy, cyano, halo, haloalkoxy, hydroxyl, and haloalkyl; and

15 n is 1, 2, 3, 4, or 5.

In compounds of formula (II), R_{2a} is hydrogen or alkyl such as C_{1-6} alkyl. In one embodiment, R_{2a} is hydrogen. In another embodiment, R_{2a} is C_{1-6} alkyl such as, but not limited to, methyl. R_{1a} is cycloalkylalkyl wherein the cycloalkyl moiety is cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl, each of which is independently
20 unsubstituted or substituted as described in the Definition of Terms. One example of the cycloalkyl moiety is cyclobutyl. Examples of the optional substituents of the cycloalkyl moiety include, but are not limited to, alkyl, haloalkyl, hydroxy, oxo, alkoxy, and haloalkoxy. In other embodiments, R_{1a} is alkyl (such as, but not limited to, butyl). In yet another embodiment, R_{1a} is haloalkyl.

25 Compounds of the present invention may exist as stereoisomers wherein, asymmetric or chiral centers are present. These stereoisomers are "R" or "S" depending on the configuration of substituents around the chiral carbon atom. The terms "R" and "S" used herein are configurations as defined in IUPAC 1974 Recommendations for Section E, Fundamental Stereochemistry, Pure Appl. Chem.,
30 1976, 45: 13-30. The present invention contemplates various stereoisomers and mixtures of various ratio thereof and are included within the scope of this invention.

Individual stereoisomers of compounds of the present invention may be prepared synthetically from commercially available starting materials which contain asymmetric or chiral centers or by preparation of racemic mixtures followed by resolution well-known to those of ordinary skill in the art. These methods of resolution are exemplified by (1) attachment of a mixture of enantiomers to a chiral auxiliary, separation of the resulting mixture of diastereomers by recrystallization or chromatography and liberation of the optically pure product from the auxiliary or (2) direct separation of the mixture of optical enantiomers on chiral chromatographic columns.

Geometric isomers can exist in the present compounds. The invention contemplates the various geometric isomers and mixtures thereof resulting from the disposition of substituents around a carbon-carbon double bond, a carbon-nitrogen double bond, a cycloalkyl group, or a heterocycle group. Substituents around a carbon-carbon double bond or a carbon-nitrogen bond are designated as being of Z or E configuration and substituents around a cycloalkyl or heterocycle are designated as being of cis or trans configuration.

Within the present invention it is to be understood that compounds disclosed hererin may exhibit the phenomenon of tautomerism

Thus, the formulae drawings within this specification can represent only one of the possible tautomeric or stereoisomeric forms. It is to be understood that the invention encompasses any tautomeric or stereoisomeric form, and mixtures thereof, and is not to be limited merely to any one tautomeric or stereoisomeric form utilized within the naming of the compounds or formulae drawings.

Abbreviations


Abbreviations which have been used in the descriptions of the Schemes and the Examples that follow are: DME for dimethoxyethane, DMF for N,N-dimethylformamide; EtOAc for ethyl acetate, EtOH for ethanol, HATU for O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate, Et₂O for diethyl ether, Et₃N for triethylamine, HPLC for high pressure liquid chromatography, MeOH for methanol, min for minute or minutes DMSO for dimethylsulfoxide; TFA for trifluoroacetic acid, THF for tetrahydrofuran; Ts for p-CH₃PhS(O)₂O-; and Tf or triflate for CF₃S(O)₂O-.

Methods for Preparing Compounds

The compounds and processes of the present invention will be better understood in connection with the following synthetic Schemes and Examples which illustrate a means by which the compounds of the present invention can be prepared.

5

Scheme 1




As shown in Scheme 1, compounds of formula (1) containing an amine group when treated with compounds of formula (2), wherein X is chloro or -OH under coupling conditions known to one skilled in the art, will provide compounds of formula (3). Typical conditions for the reaction of compounds of formula (2) wherein X is chloro and compounds of formula (1) include but are not limited to stirring an equimolar mixture of the compounds in solvents such as chloroform, dichloromethane or THF in the presence of a base such as but not limited to diisopropylethylamine at 0-30 °C for 8-24 hours. Acid coupling conditions of compounds of formula (2), wherein X is -OH and compounds of formula (1), include stirring an equimolar mixture of the compounds with a coupling reagent such as but not limited to bis(2-oxo-3-oxazolidinyl)phosphinic chloride (BOPCl), 1,3-dicyclohexylcarbodiimide (DCC), polymer supported 1,3-dicyclohexylcarbodiimide (PS-DCC), O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU), O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU) along with a coupling auxiliary such as but not limited to 1-hydroxy-7-azabenzotriazole (HOAT) or 1-hydroxybenzotriazole hydrate (HOBt) in the presence or absence of a base such as but not limited to N-methyl morpholine, diisopropylethylamine in solvents such as but not limited to THF, N,N-dimethylacetamide, N,N-dimethylformamide, pyridine and chloroform. Typical reactions can be carried out between 0-65 °C or may be carried out in a microwave reactor to facilitate the coupling.

20


25

Scheme 2




As shown in Scheme 2, compounds of formula (3) may be converted into compounds of formula (4) which are representative compounds of the present invention. Typical conditions include, but are not limited to, the treatment of compounds of formula (3) with sodium hydride in DMF at 0 °C, followed by the addition of reagents such as R₁-Y, wherein R₁ is as defined in formula (I) and Y is chloro, bromo, iodo, mesyl or triflate. Alternatively, other bases such as potassium hydroxide or potassium tert-butoxide in a mixture of THF and DMF, followed by treatment with R₁-Y will also provide compounds of formula (4).

Scheme 3



Alternatively, compounds of formula (4) may also be prepared according to the methods outlined in Scheme 3. Compounds of formula (1) when treated with sodium hydride in DMF at 0 °C, followed by the addition of reagents such as R₁-Y, wherein R₁ is as defined in formula (I) and Y is chloro, bromo, iodo, tosyl, mesyl or triflate will provide compounds of formula (6). Alternatively, compounds of formula (1) may be heated neat or in the presence of a minimal amount of solvent to facilitate mixing with compounds of formula R₁-Y to obtain compounds of formula (6). Compounds of formula (6) may be isolated as a salt or a free base. The treatment of compounds of formula (6) with compounds of formula (2), wherein X is chloro or -OH, under coupling conditions as outlined in Scheme 1 will generate compounds of formula (4), which are representative of compounds of the present invention.

Scheme 4




As outlined in Scheme 4, compounds of formula (7), wherein R_{100} is halide or triflate and which can be prepared according to the methods outlined in Schemes 1-3, when treated with a boronic acid of formula $R_3B(OH)_2$, wherein R_3 is aryl, arylalkenyl, cycloalkyl, heterocycle or heteroaryl, a palladium catalyst such as

5 dichlorobis(triphenyl)phosphine)palladium (II) and sodium carbonate in a mixture of solvents which include but are not limited to various mixtures of DME, ethanol and water under heated conditions will provide compounds of formula (4) which contain alkenyl, aryl, arylalkenyl, cycloalkyl, heterocycle or heteroaryl substituents in the R_3 position.

10


Scheme 5



15

As outlined in Scheme 5, compounds of formula (9) wherein R_{100} is halide or triflate and which can be prepared according to the methods outlined in Schemes 1-3, when treated with a boronic acid of formula $R_2B(OH)_2$, wherein R_2 is aryl, arylalkenyl, cycloalkyl, heterocycle or heteroaryl and a palladium catalyst according to the methods outlined in Scheme 4 will provide compounds of formula (4) which contain the aryl, arylalkenyl, cycloalkyl, heterocycle or heteroaryl in the R_2 position.

Scheme 6




20

Scheme 6 describes another alternative method of preparation of compounds of formula (4). Compounds of formula (11) when treated with oxalyl chloride in dichloromethane containing a catalytic amount of DMF will provide the acid chloride of formula (12). The acid chloride of formula (12) when treated with potassium thiocyanate in acetone will provide compounds of formula (13). Compounds of


formula (13) when treated with an amine of formula R_1-NH_2 in solvents such as but not limited to THF will provide compounds of formula (14). Compounds of formula (14) when treated with substituted alpha-bromo-ketones of formula (15) in ethanol or mixtures of ethanol and toluene under heated conditions will provide compounds of formula (4).

Scheme 7




Compounds of formula (I) containing a L_1 group that is $-NH-$, may be prepared as outlined in Scheme 7. Compounds of formula (16) when treated with an amine of formula (17), wherein R_4 is defined in formula (I), will provide compounds of formula (18). Compounds of formula (18) when treated with compounds of formula (6) will provide compounds of formula (19) which are representative of compounds of formula (I).

Scheme 8




Alternatively, compounds of formula (6) when treated with an isocyanate of formula (20) wherein R_4 is defined in formula (I), will provide compounds of formula (19).

Scheme 9




Similarly, compounds of formula (6) when treated with carbonyl diimidazole, followed by treatment with methyl iodide, will provide the imidazolide compounds of formula (21). Compounds of formula (21) when treated with an amine of formula (22), wherein R_4 and R_g are defined in formula (I), will provide compounds of formula (19A) which are representative of compounds of formula (I).

Scheme 10



As shown in Scheme 10, compounds of formula (23) wherein R_{31} is defined herein, when treated at low temperatures with an organolithium reagent such as but not limited to $R_{32}Li$ or a Grignard reagent such as but not limited to $R_{32}MgBr$ can be converted to compounds of formula (24). The reaction is typically conducted in a solvent such as but not limited to diethyl ether.

Scheme 11



As shown in Scheme 11, compounds of formula (25) wherein R_3 is as defined in formula (I), when treated with pyrrolidine and *p*-toluenesulfonic acid monohydrate in a solvent such as but not limited to cyclohexane at reflux followed by treatment with sulfur and cyanamide in a solvent such as methanol, will provide compounds of formula (1) wherein R_2 is hydrogen.

It will be appreciated that the synthetic schemes and specific examples as illustrated in the Examples section are illustrative and are not to be read as limiting

the scope of the invention as it is defined in the appended claims. All alternatives, modifications, and equivalents of the synthetic methods and specific examples are included within the scope of the claims.

Optimum reaction conditions and reaction times for each individual step may vary depending on the particular reactants employed and substituents present in the reactants used. Unless otherwise specified, solvents, temperatures and other reaction conditions may be readily selected by one of ordinary skill in the art. Specific procedures are provided in the Examples section. Reactions may be worked up in the conventional manner, e.g. by eliminating the solvent from the residue and further purified according to methodologies generally known in the art such as, but not limited to, crystallization, distillation, extraction, trituration and chromatography. Unless otherwise described, the starting materials and reagents are either commercially available or may be prepared by one skilled in the art from commercially available materials using methods described in the chemical literature.

Routine experimentations, including appropriate manipulation of the reaction conditions, reagents and sequence of the synthetic route, protection of any chemical functionality that may not be compatible with the reaction conditions, and deprotection at a suitable point in the reaction sequence of the method are included in the scope of the invention. Suitable protecting groups and the methods for protecting and deprotecting different substituents using such suitable protecting groups are well known to those skilled in the art; examples of which may be found in T. Greene and P. Wuts, *Protecting Groups in Chemical Synthesis* (3rd ed.), John Wiley & Sons, NY (1999), which is incorporated herein by reference in its entirety. Synthesis of the compounds of the invention may be accomplished by methods analogous to those described in the synthetic schemes described hereinabove and in specific examples.

Starting materials, if not commercially available, may be prepared by procedures selected from standard organic chemical techniques, techniques that are analogous to the synthesis of known, structurally similar compounds, or techniques that are analogous to the above described schemes or the procedures described in the synthetic examples section.

When an optically active form of a compound of the invention is required, it may be obtained by carrying out one of the procedures described herein using an optically active starting material (prepared, for example, by asymmetric induction of a suitable reaction step), or by resolution of a mixture of the stereoisomers of the

compound or intermediates using a standard procedure (such as chromatographic separation, recrystallization or enzymatic resolution).

Similarly, when a pure geometric isomer of a compound of the invention is required, it may be obtained by carrying out one of the above procedures using a pure
5 geometric isomer as a starting material, or by resolution of a mixture of the geometric isomers of the compound or intermediates using a standard procedure such as chromatographic separation.

Examples

10 The compounds and processes of the present invention will be better understood by reference to the following Examples, which are intended as an illustration of and not a limitation upon the scope of the invention.

Example 1

15 N-[(2Z)-3-(2-methoxyethyl)-1,3-thiazol-2(3H)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide

Example 1A

2,2,3,3-tetramethylcyclopropanecarbonyl chloride.

20 To a solution of 2,2,3,3-tetramethylcyclopropanecarboxylic acid (0.50 g, 3.5 mmol) in 18 mL of methylene chloride at 0 °C was added oxalyl chloride (0.61 mL, 7.0 mmol) and a catalytic amount of dimethylformamide (2 drops). The solution was stirred at ambient temperature for 1 hour, and then concentrated under reduced pressure to provide 0.56 g of the title compound.

25

Example 1B

2,2,3,3-tetramethyl-N-1,3-thiazol-2-ylcyclopropanecarboxamide

To a solution of 2-aminothiazole (0.39 g, 3.9 mmol) in 10 mL of methylene chloride at 0 °C was added a solution of the product from Example 1A in 8 mL of
30 chloroform, followed by triethylamine (1.0 mL, 7.7 mmol). The mixture was stirred for 7 hours at 35 °C, cooled to ambient temperature and diluted with water. The phases were separated and the aqueous phase was extracted with methylene chloride. The combined organic extracts were washed twice with water and then brine, dried over magnesium sulfate, filtered, and concentrated under reduced pressure.

Purification by column chromatography (SiO₂, 30-60% ethyl acetate/hexanes gradient) afforded 0.11 g (14%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.18 (s, 1 H), 1.25 (s, 6 H), 1.35 (s, 6 H), 6.92 (d, J = 3.4 Hz, 1 H), 7.39 (d, J = 3.4 Hz, 1 H), 10.7 (s, 1 H); MS (DCI/NH₃) m/z 225 (M+H)⁺. Anal. Calculated for C₁₁H₁₆N₂OS: C, 58.90; H, 7.19; N, 12.49. Found: C, 59.03; H, 7.34; N, 12.34.

Example 1C

N-[(2Z)-3-(2-methoxyethyl)-1,3-thiazol-2(3H)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide

To a solution of Example 1B (0.16 g, 0.71 mmol) in 3.5 mL of 4:1 tetrahydrofuran:dimethylformamide at 0 °C was added potassium hydroxide (90 mg, 1.7 mmol). After stirring for 1 hour at room temperature, 2-bromoethyl methyl ether (73 μL, 7.1 mmol) was added and the solution was heated to 65 °C for 14 hours. The solution was allowed to cool to ambient temperature and then diluted with ethyl acetate and washed twice with water and then brine. The organic extract was dried over magnesium sulfate, filtered, and concentrated under reduced pressure. Purification by column chromatography (SiO₂, 30-50% ethyl acetate/hexanes gradient) afforded 22 mg (11%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ 1.25 (s, 12H), 1.75 (s, 1H), 3.32 (s, 3H), 3.74 (t, J = 5.6 Hz, 2H), 4.44 (t, J = 5.4 Hz, 2H), 6.95 (d, J = 3.4 Hz, 1H), 7.48 (t, J = 3.7 Hz, 1H); MS (DCI/NH₃) m/z 283 (M+H)⁺. Anal. Calculated for C₁₄H₂₂N₂O₂S: C, 59.54; H, 7.85; N, 9.92. Found: C, 59.76; H, 7.97; N, 9.91.

Example 2

5-chloro-2-methoxy-N-[(2Z)-3-(2-methoxyethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

Example 2A

3-(2-methoxyethyl)-1,3-thiazol-2(3H)-imine hydrobromide

A mixture of 2-aminothiazole (15 g, 0.15 mol) and 2-bromoethyl methyl ether (17 mL, 0.18 mol) were heated at 85 °C for 16 hours. After cooling to ambient temperature the resulting solid was triturated twice with isopropyl alcohol to afford 26 g (72%) of the title compound. ¹H NMR (DMSO-*d*₆, 300 MHz) δ ppm 3.27 (s, 3 H),

3.63 (t, $J = 5.1$ Hz, 2 H), 4.23 (t, $J = 4.9$ Hz, 2 H), 7.02 (d, $J = 4.7$ Hz, 1 H), 7.38 (d, $J = 4.4$ Hz, 1 H), 9.52 (s, 1H); MS (DCI/NH₃) m/z 159 (M+H)⁺.

Example 2B

5 5-Chloro-2-methoxy-*N*-[(2*Z*)-3-(2-methoxyethyl)-1,3-thiazol-2(3*H*)-ylidene]benzamide

To a solution of the product from Example 2A (0.77 g, 3.2 mmol) and 5-chloro-2-methoxybenzoic acid (0.50 g, 2.7 mmol) in 14 mL of THF at 0 °C was added *O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate (HATU, 1.24 g, 3.2 mmol) and *N,N*-diisopropylethylamine (1.1 mL, 6.2 mmol). The mixture was heated to 65 °C for 2.5 hours, cooled to ambient temperature and then diluted with ethyl acetate. The mixture was washed twice with water, then saturated aqueous sodium bicarbonate, and brine. The organic extract was dried over magnesium sulfate, and concentrated under reduced pressure. Purification by column chromatography (SiO₂, 20-35% ethyl acetate/hexanes gradient) afforded 0.38 g (43%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 3.35 (s, 3 H), 3.72-3.81 (m, 2 H), 3.91 (s, 3 H), 4.41-4.48 (m, 2 H), 6.65 (d, $J=4.7$ Hz, 1 H), 6.92 (d, $J=8.8$ Hz, 1 H), 7.16 (d, $J=4.7$ Hz, 1 H), 7.34 (dd, $J=8.8, 3.1$ Hz, 1 H), 7.99 (d, $J=2.7$ Hz, 1 H); MS (DCI/NH₃) m/z 349 (M+H)⁺. Anal. Calculated for C₁₈H₂₄N₂O₃S: C, 62.04; H, 6.94; N, 8.04. Found: C, 62.24; H, 7.08; N, 8.04.

Example 3

N-[(2*Z*)-3-(2-methoxyethyl)-1,3-thiazol-2(3*H*)-ylidene]cycloheptanecarboxamide
Cycloheptanecarboxylic acid (29 mg, 0.20 mmol), 3 equiv of polymer bound dicyclohexylcarbodiimide (PS-DCC), 1-hydroxybenzotriazole hydrate (HOBT, 22 mg, 0.16 mmol), *N,N*-diisopropylethylamine (62 mg, 0.50 mmol), and the product of Example 2A (39 mg, 0.16 mmol) were combined in dimethylacetamide (DMA, 2.8 mL) and heated in a microwave to 100 °C for 420 seconds. The mixture was filtered through Si-Carbonate (6 mL-1g cartridge from Silicycle Chemical Division) and then concentrated to dryness. The residue was dissolved in 1:1 DMSO/MeOH and purified by reverse phase HPLC to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 1.39 - 1.72 (m, 11 H) 1.83 - 1.95 (m, 2 H) 2.46 - 2.52 (m, 1 H) 3.23 - 3.26 (m, 3 H) 3.67 (t, 2 H) 4.29 (t, 2 H) 6.76 - 6.97 (d, 1 H) 7.30 - 7.43 (d, 1 H); MS (ESI) m/z

283 (M+H)⁺

Example 4

N-[(2Z)-3-(3-methoxypropyl)-1,3-thiazol-2(3H)-ylidene]-2,2,3,3-
tetramethylcyclopropanecarboxamide

Example 4A

3-(3-methoxypropyl)-1,3-thiazol-2(3H)-imine hydrobromide

A mixture of 2-aminothiazole (1.0 g, 10 mmol) and 1-bromo-3-methoxypropane (1.8 g, 12 mmol) were heated at 85 °C for 16 hours. The solid was cooled to ambient temperature, triturated with ethanol, and then collected by filtration to provide 1.2 g (48%) of the title compound. MS (DCI/NH₃) m/z 173 (M+H)⁺.

Example 4B

N-[(2Z)-3-(3-methoxypropyl)-1,3-thiazol-2(3H)-ylidene]-2,2,3,3-
tetramethylcyclopropanecarboxamide

The product of Example 4A (0.60 g, 2.4 mmol) and 2,2,3,3-tetramethylcyclopropanecarboxylic acid (0.34 g, 2.4 mmol) were processed using the method described in Example 2B. Purification by column chromatography (SiO₂, 20-40% ethyl acetate/hexanes gradient) afforded 0.33 g (47%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.21 (s, 6 H), 1.34 (s, 6 H), 1.56 (d, J=5.4 Hz, 1 H), 2.00-2.13 (m, 2 H), 3.31-3.39 (m, 5 H), 4.23 (t, J=6.8 Hz, 2 H), 6.50 (d, J=4.7 Hz, 1 H), 6.88 (d, J=4.7 Hz, 1 H); MS (DCI/NH₃) m/z 297 (M+H)⁺. Anal. Calculated for C₁₅H₂₄N₂O₂S: C, 60.78; H, 8.27; N 9.45. Found: C, 60.78; H, 8.27; N, 9.34.

Example 5

N-[(2Z)-3-(2-methoxyethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2,2,3,3-
tetramethylcyclopropanecarboxamide

Example 5A

3-(2-methoxyethyl)-4-methyl-1,3-thiazol-2(3H)-imine hydrobromide

A mixture of 4-methylthiazol-2-ylamine (0.75 g, 6.5 mmol) and 2-bromoethyl methyl ether (730 μL, 7.8 mmol) was heated at 85 °C for 15 hours. The mixture was cooled to ambient temperature and the resulting solid was triturated with isopropanol.

Recrystallization from hot ethanol afforded 0.56 g (34%) of the title compound. ¹H NMR (DMSO-*d*₆, 300 MHz) δ ppm 2.25 (d, *J*=1.4 Hz, 3 H) 3.25 (s, 3 H) 3.57 (t, *J*=5.1 Hz, 2 H) 4.15 (t, *J*=5.1 Hz, 2 H) 6.68 (d, *J*=1.4 Hz, 1 H) 9.40 (s, 1 H); MS (DCI/NH₃) *m/z* 173 (M+H)⁺.

5

Example 5B

N-[(2Z)-3-(2-methoxyethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide

The product of Example 5A (0.30 g, 1.2 mmol) and 2,2,3,3-tetramethylcyclopropanecarboxylic acid (0.19 g, 1.3 mmol) were processed using the method described in Example 2B. Purification by column chromatography (SiO₂, 20-40% ethyl acetate/hexanes gradient) afforded 0.14 g (41%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.21 (s, 6 H), 1.34 (s, 6 H), 1.59 (s, 1 H), 2.30 (s, 3 H), 3.30 (s, 3 H), 3.70 (t, *J*=5.09 Hz, 2 H), 4.25 (t, *J*=5.26 Hz, 2 H), 6.09 (s, 1 H); MS (DCI/NH₃) *m/z* 297 (M+H)⁺. Anal. Calculated for C₁₅H₂₄N₂O₂S: C, 60.78; H, 8.16; N, 9.45. Found: C, 60.79; H, 7.82; N, 9.36.

15

Example 6

ethyl ((2Z)-3-(2-methoxyethyl)-2-[(2,2,3,3-tetramethylcyclopropyl)carbonyl]imino)-2,3-dihydro-1,3-thiazol-4-yl)acetate

20

Example 6A

ethyl [2-imino-3-(2-methoxyethyl)-2,3-dihydro-1,3-thiazol-4-yl]acetate hydrobromide

A mixture of (2-aminothiazol-4-yl)acetic acid ethyl ester (18.6 g, 100 mmol) and 2-bromoethyl methyl ether (15.3 g, 110 mmol) were processed using the method described in Example 2A to afford 14.1 g (83%) of the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.14 - 1.28 (m, 3 H) 3.24 (s, 3 H) 3.54 (t, *J*=5 Hz, 2 H) 3.91 (s, 2 H) 4.04 - 4.25 (m, 4 H) 6.92 (s, 1 H) 9.50 (s, 1 H); MS (DCI/NH₃) *m/z* 231(M+H)⁺.

30

Example 6B

ethyl ((2Z)-3-(2-methoxyethyl)-2-[(2,2,3,3-tetramethylcyclopropyl)carbonyl]imino)-2,3-dihydro-1,3-thiazol-4-yl)acetate

The product of Example 6A (2.3 g, 10 mmol) and 2,2,3,3-

tetramethylcyclopropane carboxylic acid (1.6 g, 11 mmol) were processed as described using the method described in Example 2B to afford 2.1 g (54%) of the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.06 - 1.33 (m, 15 H) 1.48 (s, 1 H) 3.22 (s, 3 H) 3.59 (t, *J*=5 Hz, 2 H) 3.91 (s, 2 H) 4.12 (t, *J*=7 Hz, 2 H) 4.14 - 4.24 (m, 2 H) 6.69 (s, 1 H); MS (DCI/NH₃) *m/z* 369 (M+H)⁺. Anal. Calculated for C₁₈H₂₈N₂O₄S: C, 68.67; H, 7.66; N, 7.62. Found: C, 68.67; H, 7.66; N, 7.60.

Example 7

ethyl (2Z)-3-(2-methoxyethyl)-2-[(2,2,3,3-tetramethylcyclopropyl)carbonyl]imino)-
2,3-dihydro-1,3-thiazole-4-carboxylate

Example 7A

ethyl 2-imino-3-(2-methoxyethyl)-2,3-dihydro-1,3-thiazole-4-carboxylate
hydrobromide

A mixture of 2-aminothiazole-4-carboxylic acid ethyl ester (17.2 g, 100 mmol) and 2-bromoethyl methyl ether (15.3 g, 110 mmol) were processed using the method described in Example 2A to afford 17.1 g (83%) of the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.30 (t, *J*=7 Hz, 3 H) 3.22 (s, 3 H) 3.60 (t, *J*=5 Hz, 2 H) 4.32 (t, *J*=7 Hz, 2 H) 4.35 - 4.61 (m, 2 H) 7.84 (s, 1 H) 9.76 (s, 1 H); MS (DCI/NH₃) *m/z* 231 (M+H)⁺.

Example 7B

ethyl (2Z)-3-(2-methoxyethyl)-2-[(2,2,3,3-tetramethylcyclopropyl)carbonyl]imino)-
2,3-dihydro-1,3-thiazole-4-carboxylate

The product of Example 7A (2.3 g, 10 mmol) and 2,2,3,3-tetramethylcyclopropane carboxylic acid (1.6 g, 11 mmol) were processed using the method described in Example 2B to afford 1.9 g (53 %) of the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.11 - 1.36 (m, 12 H) 1.53 (s, 1 H) 3.21 (s, 3 H) 3.31 (s, 3 H) 3.53 - 3.62 (m, 2 H) 4.30 (q, *J*=7 Hz, 2 H) 4.62 - 4.75 (m, 2 H) 7.77 (s, 1 H); MS (DCI/NH₃) *m/z* 355 (M+H); Anal. Calculated for C₁₇H₂₆N₂O₄S: C, 57.61; H, 7.39; N, 7.86. Found: C, 57.86; H, 7.67; N, 7.85.

Example 8

N-[(2Z)-4-(hydroxymethyl)-3-(2-methoxyethyl)-1,3-thiazol-2(3H)-ylidene]-2,2,3,3-

tetramethylcyclopropanecarboxamide

To a solution of the product of Example 7B (0.355 g, 1.00 mmol) in 100 mL of THF at 0 °C was added lithium borohydride (10 mL of a 2.0 M solution in THF) and the resulting solution was allowed to warm to ambient temperature and stirred overnight. The mixture was quenched with water and then diluted with saturated aqueous Na₂CO₃ and extracted twice with ethyl acetate. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered, and concentrated. Purification by column chromatography (SiO₂, 0-50% ethyl acetate/hexanes gradient) afforded 0.278 g (89%) of the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.20 (d, 12 H) 1.47 (s, 1 H) 3.24 (s, 3 H) 3.64 (t, *J*=6 Hz, 2 H) 4.30 (t, *J*=6 Hz, 2 H) 4.50 (d, *J*=5 Hz, 2 H) 5.75 (s, 1 H) 6.68 (s, 1 H); MS (DCI/NH₃) *m/z* 313(M+H)⁺; Anal. Calculated for C₁₅H₂₄N₂O₃S•0.2H₂O: C, 57.01; H, 7.78; N, 8.86. Found: C, 56.90; H, 7.61; N, 8.86.

Example 92-ethoxy-N-[(2*Z*)-3-(2-methoxyethyl)-4-(trifluoromethyl)-1,3-thiazol-2(3*H*)-ylidene]benzamideExample 9A2-ethoxy-N-(4-trifluoromethyl-thiazol-2-yl)-benzamide

A solution of 2-ethoxybenzoic acid (0.75 g, 4.5 mmol) in 23 mL of methylene chloride at 0 °C was treated with oxalyl chloride (0.44 mL, 4.9 mmol) followed by 2 drops of dimethylformamide. The solution was stirred at ambient temperature for 1 hour and then concentrated under reduced pressure to provide 0.83 g of 2-ethoxybenzoyl chloride. To a solution of 4-trifluoromethylthiazol-2-ylamine (0.50 g, 3.0 mmol) in 10 mL THF at 0 °C was added a solution of the freshly prepared acid chloride in 5 mL of THF and 2 mL of methylene chloride, followed by triethylamine (1.0 mL, 6.6 mmol). The reaction mixture was warmed to 65 °C and stirred 8 hours. The mixture was diluted with ethyl acetate and washed twice with water, then brine. The organic extract was dried over magnesium sulfate, filtered, and concentrated under reduced pressure. Purification by column chromatography (SiO₂, 30-50% ethyl acetate/hexanes gradient) afforded 0.47 g (50%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.66 (t, *J*=6.95 Hz, 3 H), 4.38 (q, *J*=6.89 Hz, 2 H), 7.03 -

7.10 (m, 2 H), 7.17 (d, $J=7.80$ Hz, 1 H), 7.42 (s, 1 H), 8.29 (dd, $J=7.97$, 1.86 Hz, 1 H).
MS (DCI/NH₃) m/z 317 (M+H)⁺.

Example 9B

2-ethoxy-N-[(2Z)-3-(2-methoxyethyl)-4-(trifluoromethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 9A (0.47 g, 1.5 mmol) and 1-bromo-2-methoxyethane (0.16 ml, 1.6 mmol) were processed using the method described in Example 1B. Purification by column chromatography (SiO₂, 30-40% ethyl acetate/hexanes gradient) afforded 0.06 g (11%) of the title compound. ¹H NMR (DMSO-*d*₆, 300 MHz) δ ppm 1.24 (t, $J=6.95$ Hz, 3 H), 3.02 (s, 3 H), 3.36 - 3.63 (m, 2 H), 3.86 - 4.02 (m, 1 H), 4.13 (q, $J=7.12$ Hz, 2 H), 4.21 - 4.33 (m, 1 H), 7.08 (t, $J=7.46$ Hz, 1 H), 7.17 (d, $J=8.14$ Hz, 1 H), 7.40 (dd, $J=7.46$, 1.70 Hz, 1 H), 7.46 - 7.56 (m, 1 H), 8.09 (s, 1 H); MS (DCI/NH₃) m/z 375 (M+H)⁺. Anal. Calculated for C₁₆H₁₇F₃N₂O₃S•0.2 H₂O: C, 50.84; H, 4.64; N, 7.41. Found: C, 50.62; H, 4.35; N, 7.61.

Example 10

N-[(2Z)-3-(2-methoxyethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide

Example 10A

3-(2-Methoxyethyl)-5-methyl-3H-thiazol-2-ylideneamine hydrobromide

A mixture of 5-methyl-thiazol-2-ylamine (1.0 g, 8.8 mmol) and 2-bromoethyl methyl ether (1.0 mL, 11 mmol) were heated at 85 °C for 16 hours. The mixture was cooled to ambient temperature, triturated with ethanol and the solid was collected by filtration to afford 0.90 g (40%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 2.25 (d, $J=1.4$ Hz, 3 H), 3.36 (s, 3 H), 3.72-3.81 (m, 2 H), 4.36-4.43 (m, 2 H), 6.61 (d, $J=1.7$ Hz, 1 H), 9.54 (s, 1 H); MS (DCI/NH₃) m/z 173 (M+H)⁺.

Example 10B

N-[(2Z)-3-(2-methoxyethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide

The product of Example 10A (0.40 g, 1.6 mmol) and 2,2,3,3-tetramethylcyclopropanecarboxylic acid (0.25 g, 1.8 mmol) were processed using the method described in Example 2B. Purification by column chromatography (SiO₂, 20-40% ethyl acetate/hexanes gradient) afforded 0.30 g (63%) of the title compound. ¹H

NMR (CDCl₃, 300 MHz) δ ppm 1.21 (s, 6 H), 1.33 (s, 6 H), 1.52 (s, 1 H), 2.22 (s, 3 H), 3.35 (s, 3 H), 3.68 (t, $J=5.09$ Hz, 2 H), 4.24 (t, $J=4.92$ Hz, 2 H), 6.67 (s, 1 H); MS (DCI/NH₃) m/z 297 (M+H)⁺. Anal. Calculated for C₁₅H₂₄N₂O₂S: C, 60.78; H, 8.16; N, 9.45. Found: C, 60.69; H, 8.31; N, 9.19.

5

Example 11

2-ethoxy-N-[(2Z)-3-(2-methoxyethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]benzamide

To the product of Example 10A (0.55 g, 2.2 mmol) and 2-ethoxybenzoyl chloride (0.33 g, 1.8 mmol) in 10 mL of THF at 0 °C was added triethylamine (0.55 mL, 4.0 mmol). The solution was stirred at 65 °C for 4 hours then allowed to cool to ambient temperature and diluted with ethyl acetate. The solution was washed twice with water and then brine. The combined aqueous washings were extracted with ethyl acetate. The combined organic extracts were dried over magnesium sulfate, filtered, and concentrated under reduced pressure. Purification by column chromatography (SiO₂, 20-30% ethyl acetate/hexanes gradient) afforded 0.28 g (42%) of the title compound. ¹H NMR (DMSO-*d*₆, 300 MHz) δ ppm 1.31 (t, $J=7.0$ Hz, 3 H), 2.26 (d, $J=1.4$ Hz, 3 H), 3.25 (s, 3 H), 3.69 (t, $J=5.3$ Hz, 2 H), 4.05 (q, $J=7.1$ Hz, 2 H), 4.30 (t, $J=5.3$ Hz, 2 H), 6.95 (t, $J=7.5$ Hz, 1 H), 7.04 (d, $J=7.8$ Hz, 1 H), 7.20 (d, $J=1.4$ Hz, 1 H), 7.32-7.41 (m, 1 H), 7.68 (dd, $J=7.6, 1.9$ Hz, 1 H); MS (DCI/NH₃) m/z 321 (M+H)⁺. Anal. Calculated for C₁₆H₂₀N₂O₃S•0.2 H₂O: C, 59.31; H, 6.35; N, 8.65. Found: C, 59.18; H, 6.02; N, 8.29.

20

Example 12

3-fluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-

25

methylbenzamide

Example 12A

3-(2-Methoxyethyl)-4,5-dimethyl-3H-thiazol-2-ylideneamine hydrobromide

A mixture of 4,5-dimethylthiazol-2-ylamine (9.0 g, 70 mmol) and 2-bromoethyl methyl ether (7.9 mL, 84 mmol) were heated at 85 °C for 12 hours. The mixture was cooled to ambient temperature and then triturated with isopropanol. The solid was collected by filtration and dried under vacuum to provide 10 g (56%) of the title compound. ¹H NMR (DMSO-*d*₆, 300 MHz) δ ppm 2.17 (s, 3 H), 2.19 (s, 3 H),

30

3.25 (s, 3 H) 3.56 (t, $J=5.1$ Hz, 2 H) 4.16 (t, $J=5.1$ Hz, 2 H) 9.41 (s, 1 H); MS (DCI/ NH_3) m/z 129 ($\text{M}+\text{H}$)⁺.

Example 12B

3-fluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-methylbenzamide

5 The product of Example 12A (39 mg, 0.15 mmol) and 3-fluoro-2-methylbenzoic acid (31 mg, 0.22 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, $\text{DMSO}-d_6$) δ ppm 2.21 - 2.24 (m, 3 H), 2.26 - 2.28 (m, 3 H), 2.44 - 2.47 (m, 3 H), 3.24 (s, 3 H), 3.66 - 10 3.71 (m, 2 H), 4.35 (t, 2 H), 7.21 - 7.31 (m, 2 H), 7.76 (d, 1 H); MS (ESI) m/z 324 ($\text{M}+\text{H}$)⁺.

Example 13

5-fluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-methylbenzamide

Example 13A

5-Fluoro-2-methyl-benzoyl chloride

A solution of 5-fluoro-2-methylbenzoic acid (380 mg, 2.47 mmol) in thionyl chloride (5 mL) was heated to reflux for 3 hours. The solution was cooled to ambient temperature and the volatile components were removed under reduced pressure. The residue was dissolved in fresh toluene (10 mL) and concentrated under reduced pressure twice and then placed under high vacuum to afford the title compound (420 mg).

Example 13B

5-fluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-methylbenzamide

To a suspension of the product of Example 12A (549 mg, 2.05 mmol) and triethylamine (0.859 mL, 6.16 mmol) in THF (6 mL) were added a solution of the product from Example 13A in THF (2 mL). The mixture was heated at reflux for 14 hours, then cooled to ambient temperature and diluted with water and CH_2Cl_2 . The phases were separated and the organic extract was washed with water and brine, dried (Na_2SO_4) and concentrated. The residue was dissolved in warm EtOAc (10 mL) and

allowed to stand at room temperature for 14 hours. The crystals were isolated by filtration (EtOAc wash) to afford the title compound (450 mg, 68%). MS (ESI) m/z 324 (M+H)⁺.

5 Example 14

3-methoxy-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-4-methylbenzamide

The product of Example 12A (39 mg, 0.15 mmol) and 3-methoxy-4-methylbenzoic acid (37 mg, 0.22 mmol) were processed using the method described
10 in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.21 (d, 6 H), 2.26 (s, 3 H), 3.26 (s, 3 H), 3.74 (t, 2 H), 3.85 (s, 3 H), 4.40 (t, 2 H), 7.22 (d, 1 H), 7.68 - 7.72 (m, 2 H); MS (ESI) m/z 335 (M+H)⁺.

Example 15

15 2-ethoxy-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 12A (0.35 g, 1.3 mmol) and 2-ethoxybenzoic acid (0.43 g, 2.6 mmol) were processed using the methods described in Example 13. Purification by column chromatography (SiO₂, 30-50% ethyl acetate/hexanes
20 gradient) afforded 0.078 g (18%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.46 (t, *J*=7.0 Hz, 3 H), 2.18-2.31 (m, 6 H), 3.30 (s, 3 H), 3.78 (t, *J*=5.3 Hz, 2 H), 4.17 (d, *J*=7.1 Hz, 2 H), 4.37 (s, 2 H), 6.89-7.04 (m, 2 H), 7.36 (t, *J*=7.6 Hz, 1 H), 7.96 (dd, *J*=7.8, 1.7 Hz, 1 H); MS (DCI/NH₃) m/z 335 (M+H)⁺. Anal. Calculated for C₁₇H₂₂N₂O₃S·0.1 H₂O: C, 60.73; H, 6.65; N, 8.33. Found: C, 60.37; H, 6.42; N, 8.31.

25

Example 16

N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-1-naphthamide

The product of Example 12A (39 mg, 0.15 mmol) and 1-naphthoic acid (39 mg, 0.22 mmol) were processed using the method described in Example 3 to afford
30 the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.25 (s, 3 H), 2.28 (s, 3 H), 3.26 (s, 3 H), 3.72 - 3.77 (m, 2 H), 4.37 - 4.43 (m, 2 H), 7.52 - 7.60 (m, 3 H), 7.95 - 7.99 (m, 1 H), 8.02 - 8.06 (m, 1 H), 8.28 - 8.31 (m, 1 H), 9.03 - 9.07 (m, 1 H); MS (ESI) m/z 341 (M+H)⁺.

Example 17

N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-naphthamide

The product of Example 12A (39 mg, 0.15 mmol) and 2-naphthoic acid (39 mg, 0.22 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.24 (s, 3 H) 2.29 (s, 3 H) 3.28 (s, 3 H) 3.80 (t, 2 H) 4.49 (t, 2 H) 7.55 - 7.62 (m, 2 H) 7.95 - 7.99 (m, 2 H) 8.08 (d, 1 H) 8.26 - 8.29 (m, 1 H) 8.76 (s, 1 H); MS (ESI) *m/z* 341 (M+H)⁺.

10

Example 18

5-chloro-2-methoxy-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 12A (39 mg, 0.15 mmol) and 5-chloro-2-methoxybenzoic acid (41 mg, 0.22 mmol) were processed using the methods described in Example 13 to afford the title compound. MS (ESI) *m/z* 355 (M+H)⁺.

15

Example 19

1-hydroxy-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-naphthamide

The product of Example 12A (39 mg, 0.15 mmol) and 1-hydroxy-2-naphthoic acid (41 mg, 0.22 mmol) were processed using the methods described in Example 13 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.27 (s, 3 H), 2.31 (s, 3 H), 3.28 (s, 3 H), 3.80 (t, 2 H), 4.44 (t, 2 H), 7.36 (d, 1 H), 7.53 (t, 1 H), 7.61 (t, 1 H), 7.86 (d, 1 H), 8.06 (d, 1 H), 8.28 (d, 1 H), 14.38 (s, 1 H); MS (ESI) *m/z* 357 (M+H)⁺.

20

25

Example 20

4-fluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-1-naphthamide

The product of Example 12A (39 mg, 0.15 mmol) and 4-fluoro-1-naphthoic acid (42 mg, 0.22 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.25 (s, 3 H) 2.28 (s, 3 H) 3.26 (s, 3 H) 3.74 (t, 2 H) 4.41 (t, 2 H) 7.38 - 7.44 (m, 1 H) 7.65 - 7.72 (m, 2

30

H) 8.12 (d, 1 H) 8.37 - 8.41 (m, 1 H) 9.23 (d, 1 H) MS (ESI) m/z 359 (M+H)⁺.

Example 21

2-methoxy-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-4-
5 (methylthio)benzamide

The product of Example 12A (39 mg, 0.15 mmol) and 2-methoxy-4-methylsulfanylbzoic acid (44 mg, 0.22 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.20 (s, 3 H) 2.23 (s, 3 H) 2.53 (s, 3 H) 3.24 (s, 3 H) 3.68 (t, 2 H) 3.81 (s, 3 H)
10 4.29 (t, 2 H) 6.83 - 6.87 (m, 1 H) 6.87 - 6.90 (m, 1 H) 7.75 (d, 1 H) MS (ESI) m/z 367 (M+H)⁺.

Example 22

2-chloro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-5-
15 (methylthio)benzamide

The product of Example 12A (39 mg, 0.15 mmol) and 2-chloro-5-methylsulfanylbzoic acid (44 mg, 0.22 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.23 (s, 3 H) 2.27 (s, 3 H) 3.24 (s, 3 H) 3.29 (s, 3 H) 3.69 (t, 2 H) 4.33 (t, 2 H)
20 7.30 - 7.33 (m, 1 H) 7.39 - 7.42 (m, 1 H) 7.67 (d, 1 H) MS (ESI) m/z 371 (M+H)⁺.

Example 23

N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-4-
(trifluoromethyl)nicotinamide

25 The product of Example 12A (0.30 g, 1.1 mmol) and 4-trifluoromethylnicotinic acid (0.43 g, 2.2 mmol) were processed as in the methods of Example 13. Purification by column chromatography (SiO₂, 0-20% methanol/methylene chloride gradient) afforded 0.23 g (28%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 2.26 (s, 3 H), 2.29 (s, 3 H), 3.30 (s, 3 H), 3.72 (t, *J*=5.1 Hz, 2 H), 4.33 (t, *J*=5.1 Hz, 2 H), 7.59 (d, *J*=5.1 Hz, 1 H), 8.79 (d, *J*=5.1 Hz, 1 H), 9.23 (s, 1 H); MS (DCI/NH₃) m/z 360 (M+H)⁺. Anal. Calculated for C₁₅H₁₆F₃N₃O₂S: C, 50.13; H, 4.49; N, 11.69. Found: C, 50.12; H, 4.33; N, 11.75.

30

Example 242-hydroxy-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 12A (1.7 g, 9.4 mmol) and 2-hydroxybenzoic acid (1.6 g, 11 mmol) were processed using the method described in Example 2B. Purification by column chromatography (SiO₂, 20-40% ethyl acetate/hexanes gradient) afforded 0.91 g (32%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 2.26 (d, *J*=1.0 Hz, 3 H), 2.29 (d, *J*=0.7 Hz, 3 H), 3.31 (s, 3 H), 3.78-3.86 (m, 2 H), 4.34 (t, *J*=5.1 Hz, 2 H), 6.89 (dt, *J*=7.9, 7.0, 1.0 Hz, 1 H), 6.95 (dd, *J*=8.1, 1.0 Hz, 1 H), 7.37 (dt, *J*=7.7, 1.9 Hz, 1 H), 8.15 (dd, *J*=8.0, 1.9 Hz, 1 H); MS (DCI/NH₃) *m/z* 307 (M+H)⁺. Anal. Calculated for C₁₅H₁₈N₂O₃S: C, 58.80; H, 5.92; N, 9.14. Found: C, 58.60; H, 5.86; N, 9.01.

Example 252-(2-methoxyethoxy)-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]benzamideExample 25AMethyl-2-(2-Methoxy-ethoxy)-benzoate

To a solution of triphenylphosphine (0.36 g, 1.40 mmol) in 10 mL of THF at 0 °C was added diisopropyl azodicarboxylate (275 μ L, 1.40 mmol). The mixture was stirred for 0.5 hours and then methyl-2-hydroxybenzoate (400 mg, 1.3 mmol) and 2-methoxyethanol (110 μ L, 1.40 mmol) were added. The mixture was allowed to warm to ambient temperature and stirred for 16 hours. The mixture was concentrated under reduced pressure and the residue was used without purification. MS (DCI/NH₃) *m/z* 211 (M+H)⁺.

Example 25B2-(2-Methoxy-ethoxy)-benzoic acid

A mixture of the product of Example 25A (0.27 g, 1.3 mmol) in 40% aqueous potassium hydroxide was stirred for 6 hours. The mixture was then diluted with water, made slightly acidic by the addition of 2 N aqueous HCl, and then extracted three times with ethyl acetate. The combined organic extracts were dried over

magnesium sulfate, filtered, and concentrated under reduced pressure to afford 0.25 g of the title compound

Example 25C

5 2-(2-Methoxy-ethoxy)-N-[3-(2-methoxyethyl)-4,5-dimethyl-3H-thiazol-2-ylidene]-benzamide

The product of Example 25B (0.25 g, 1.3 mmol) and the product of Example 12A (0.28 g, 1.5 mmol) were processed using the methods described in Example 13. Purification by column chromatography (SiO₂, 30-60% ethyl acetate/hexanes
10 gradient) afforded 35 mg (7%) of the title compound. MS (DCI/NH₃) m/z 365 (M+H)⁺. Anal. Calculated for C₁₈H₂₄N₂O₄S 0.5 H₂O: C, 57.89; H, 6.75; N, 7.50. Found: C, 57.77; H, 6.59; N, 7.44.

Example 26

15 5-chloro-2-ethoxy-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 12A (97 mg, 0.52 mmol) and 5-chloro-2-ethoxybenzoic acid (95 mg, 0.47 mmol) were processed using the methods described in Example 13 to afford the title compound. ¹H NMR (CDCl₃, 400 MHz) δ ppm 1.45
20 (t, J=6.9 Hz, 3 H), 2.24 (s, 3 H), 2.28 (s, 3 H), 3.31 (s, 3 H), 3.78 (t, J=4.8 Hz, 2 H), 4.15 (q, J=7.1 Hz, 2 H), 4.45 (s, 2 H), 6.90 (d, J=8.6 Hz, 1 H), 7.30 (dd, J=8.9, 2.8 Hz, 1 H), 7.89 (d, J=2.8 Hz, 1 H); MS (DCI/NH₃) m/z 369 (M+H)⁺. Anal. Calculated for C₁₇H₂₁ClN₂O₃S: C, 55.35; H, 5.74; N, 7.59. Found: C, 55.13; H, 5.59; N, 7.54.

Example 27

25 2-ethoxy-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]nicotinamide

The product of Example 12A (0.40 g, 2.2 mmol) and 2-ethoxynicotinic acid (0.40 g, 2.4 mmol) were processed using the method described in Example 2B.
30 Purification by column chromatography (SiO₂, 0-30% methanol/methylene chloride gradient) afforded 0.34 g (45%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.45 (t, J=7.1 Hz, 3 H), 2.23 (s, 3 H), 2.26 (s, 3 H), 3.31 (s, 3 H), 3.78 (t, J=5.3 Hz, 2 H), 4.37 (d, J=4.7 Hz, 2 H), 4.52 (q, J=7.0 Hz, 2 H), 6.91 (dd, J=7.5, 4.7 Hz, 1

H), 8.21 (dd, $J=4.7, 2.0$ Hz, 1 H), 8.32 (dd, $J=7.5, 2.0$ Hz, 1 H); MS (DCI/ NH_3) m/z 336 ($\text{M}+\text{H}$)⁺. Anal. Calculated for $\text{C}_{16}\text{H}_{21}\text{N}_3\text{O}_3\text{S}\cdot 0.2\text{H}_2\text{O}$: C, 56.68; H, 6.36; N, 12.39. Found: C, 56.65; H, 6.32; N, 12.38.

5

Example 28

2-chloro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]nicotinamide

A mixture of 2-chloronicotinic acid (0.42 g, 2.7 mmol) and 1,1'-carbonyldiimidazole (0.43 g, 2.7 mmol) in 3 mL of ethyl acetate was stirred at ambient temperature for 4 hours. The mixture was treated with water (3 mL) and the product of Example 12A (0.45 g, 2.4 mmol) and then heated at 65 °C for 13 hours. The mixture was cooled to ambient temperature, diluted with ethyl acetate and the layers separated. The organic phase was washed with twice with water and then brine, dried over magnesium sulfate, filtered and concentrated under reduced pressure. The residue was recrystallized from methylene chloride and afforded 0.14 g (18%) of the title compound. ¹H NMR (CDCl_3 , 300 MHz) δ ppm 2.26 (s, 3 H), 2.28 (s, 3 H), 3.30 (s, 3 H), 3.76 (t, $J=5.1$ Hz, 2 H), 4.36 (t, $J=5.1$ Hz, 2 H), 7.28-7.32 (m, 1 H), 8.28 (dd, $J=7.5, 2.0$ Hz, 1 H), 8.42 (dd, $J=4.7, 2.0$ Hz, 1 H); MS (DCI/ NH_3) m/z 326 ($\text{M}+\text{H}$)⁺. Anal. Calculated for $\text{C}_{14}\text{H}_{16}\text{ClN}_3\text{O}_2\text{S}$: C, 51.61; H, 4.95; N, 12.90. Found: C, 51.57; H, 4.76; N, 12.74.

20

Example 29

N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-(trifluoromethoxy)benzamide

A mixture of 2-trifluoromethoxybenzoyl chloride (0.59 g, 2.6 mmol) and the product of Example 12A were processed using the method described in Example 11 to afford the title compound. ¹H NMR (CDCl_3 , 300 MHz) δ ppm 2.24 (s, 3 H), 2.28 (s, 3 H), 3.29 (s, 3 H), 3.76 (t, $J=4.9$ Hz, 2 H), 4.45 (t, $J=4.7$ Hz, 2 H), 7.27-7.39 (m, 2 H), 7.46 (td, $J=7.7, 1.9$ Hz, 1 H), 8.06 (dd, $J=7.6, 1.9$ Hz, 1 H); Anal. Calculated for $\text{C}_{16}\text{H}_{17}\text{F}_3\text{N}_2\text{O}_3\text{S}$: C, 51.33; H, 4.58; N, 7.48. Found: C, 51.29; H, 4.40; N, 7.37.

30

Example 30

5-bromo-2-ethoxy-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-

ylidene]benzamide

The product of Example 12A (0.20 g, 1.1 mmol) and 5-bromo-2-ethoxybenzoyl chloride (0.28 g, 1.1 mmol) were processed using the method described in Example 11. Purification by column chromatography (SiO₂, 30-60% ethyl acetate/hexanes gradient) afforded 149 mg (38%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.46 (t, *J*=7.0 Hz, 3 H), 2.25 (s, 3 H), 2.29 (s, 3 H), 3.31 (s, 3 H), 3.79 (t, *J*=4.6 Hz, 2 H), 4.09-4.23 (m, 2 H), 4.44-4.61 (m, 2 H), 6.85 (d, *J*=8.8 Hz, 1 H), 7.42-7.48 (m, 1 H), 8.02 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 415 (M+H)⁺. Anal. Calculated for C₁₇H₂₁BrN₂O₃S: C, 49.40; H, 5.12; N, 6.78. Found: C, 49.68; H, 5.03; N, 6.71.

Example 31

N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-(trifluoromethyl)benzamide

The product of Example 12A (0.50 g, 2.7 mmol) and 2-trifluoromethylbenzoyl chloride (0.62 g, 3.0 mmol) were processed using the method described in Example 11. Purification by column chromatography (SiO₂, 20-40% ethyl acetate/hexanes gradient) afforded 0.43 g (44%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 2.26 (s, 3 H), 2.28 (s, 3 H), 3.29 (s, 3 H), 3.72 (t, *J*=4.7 Hz, 2 H), 4.41 (t, *J*=4.6 Hz, 2 H), 7.45-7.62 (m, 2 H), 7.71 (d, *J*=7.1 Hz, 1 H), 7.86 (d, *J*=7.1 Hz, 1 H); MS (DCI/NH₃) *m/z* 359 (M+H)⁺. Anal. Calculated for C₁₆H₁₇F₃N₂O₂S: C, 53.62; H, 4.78; N, 7.82. Found: C, 53.58; H, 4.51; N, 7.70.

Example 32

2-iodo-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 12A (0.25 g, 1.3 mmol) and 2-iodobenzoyl chloride (0.37 g, 1.4 mmol) were processed using the method described in Example 11. Purification by preparative HPLC on a Waters Symmetry C8 column (40 mm X 100 mm, 7 μm particle size) using a gradient of 20% to 95% acetonitrile:0.1% aqueous TFA over 12 minutes (15 minutes run time) at a flow rate of 70 mL/min afforded 0.12 g (23%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 2.30 (s, 3 H), 2.31 (s, 3 H), 3.29 (s, 3 H), 3.72 (t, *J*=4.7 Hz, 2 H), 4.41 (t, *J*=4.6 Hz, 2 H), 7.45-7.62 (m, 2

H), 7.71 (d, $J=7.1$ Hz, 1 H), 7.86 (d, $J=7.1$ Hz, 1 H); MS (DCI/ NH_3) m/z 417 ($\text{M}+\text{H}$)⁺.

Example 33

5 2-fluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-5-
 (trifluoromethyl)benzamide

The product of Example 12A (0.25 g, 1.3 mmol) and 2-fluoro-5-trifluoromethylbenzoyl chloride (0.32 g, 1.4 mmol) were processed using the method described in Example 11. Purification by preparative HPLC on a Waters Symmetry
10 C8 column (40 mm X 100 mm, 7 μm particle size) using a gradient of 20% to 95% acetonitrile:0.1% aqueous TFA over 12 minutes (15 minutes run time) at a flow rate of 70 mL/min afforded 70 mg (14%) of the title compound. ¹H NMR (CDCl_3 , 300 MHz) δ ppm 2.29 (s, 3 H), 2.31 (s, 3 H), 3.33 (s, 3 H), 3.73-3.89 (m, 2 H), 4.44-4.57 (m, 2 H), 7.18-7.24 (m, 1 H), 7.68 (d, $J=9.2$ Hz, 1 H), 8.38 (d, $J=6.8$ Hz, 1 H); MS
15 (DCI/ NH_3) m/z 377 ($\text{M}+\text{H}$)⁺. Anal. Calculated for $\text{C}_{16}\text{H}_{16}\text{F}_4\text{N}_2\text{O}_2\text{S}\cdot 0.1\text{H}_2\text{O}$: C, 51.06; H, 4.28; N, 7.44. Found: C, 50.54; H, 4.05; N, 7.27.

Example 34

20 2-bromo-5-methoxy-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-
 ylidene]benzamide

The product of Example 12A (0.20 g, 1.1 mmol) and 2-bromo-5-methoxybenzoic acid (0.25 g, 1.1 mmol) were processed using the methods described in Example 13. Purification by preparative HPLC on a Waters Symmetry C8 column (40 mm X 100 mm, 7 μm particle size) using a gradient of 20% to 95%
25 acetonitrile:0.1% aqueous TFA over 12 minutes (15 minutes run time) at a flow rate of 70 mL/min afforded 0.13 g (29%) of the title compound. ¹H NMR (CDCl_3 , 300 MHz) δ ppm 2.27 (s, 3 H), 2.29 (s, 3 H), 3.30 (s, 3 H), 3.80 (t, $J=4.7$ Hz, 2 H), 3.83 (s, 3 H), 4.50-4.59 (m, 2 H), 6.82 (dd, $J=8.8, 3.1$ Hz, 1 H), 7.44 (d, $J=3.1$ Hz, 1 H), 7.50 (d, $J=8.8$ Hz, 1 H); MS (DCI/ NH_3) m/z 401 ($\text{M}+\text{H}$)⁺. Anal. Calculated for
30 $\text{C}_{16}\text{H}_{19}\text{BrN}_2\text{O}_3\text{S}$: C, 48.13; H, 4.80; N, 7.02. Found: C, 47.88; H, 4.55; N, 6.89.

Example 35

5-fluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-

(trifluoromethyl)benzamide

The product of Example 12A (0.20 g, 1.1 mmol) and 5-fluoro-2-trifluoromethylbenzoyl chloride (0.18 mL, 1.2 mmol) were processed using the method described in Example 11. Purification by preparative HPLC on a Waters Symmetry C8 column (40 mm X 100 mm, 7 μ m particle size) using a gradient of 20% to 95% acetonitrile:0.1% aqueous TFA over 12 minutes (15 minutes run time) at a flow rate of 70 mL/min afforded 43 mg (11%) of the title compound. ^1H NMR (CDCl_3 , 300 MHz) δ ppm 2.27 (s, 3 H), 2.29 (s, 3 H), 3.30 (s, 3 H), 3.65-3.76 (m, 2 H), 4.37-4.48 (m, 2 H), 7.13-7.20 (m, 1 H), 7.56 (d, $J=7.5$ Hz, 1 H), 7.71 (dd, $J=9.2$, 5.4 Hz, 1 H); MS (DCI/ NH_3) m/z 377 ($\text{M}+\text{H}$) $^+$. Anal. Calculated for $\text{C}_{16}\text{H}_{16}\text{F}_4\text{N}_2\text{O}_2\text{S}\cdot 0.3\text{H}_2\text{O}$: C, 50.34; H, 4.38; N, 7.34. Found: C, 49.95; H, 4.02; N, 7.09.

Example 36

15 N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2,5-bis(trifluoromethyl)benzamide

The product of Example 12A (0.20 g, 1.1 mmol) and 2,5-bis-trifluoromethylbenzoyl chloride (0.33 g, 1.2 mmol) were processed using the method described in Example 11. Purification by preparative HPLC on a Waters Symmetry C8 column (40 mm X 100 mm, 7 μ m particle size) using a gradient of 20% to 95% acetonitrile:0.1% aqueous TFA over 12 minutes (15 minutes run time) at a flow rate of 70 mL/min afforded 0.14 g (31%) of the title compound. ^1H NMR (CDCl_3 , 300 MHz) δ ppm 2.27 (s, 3 H), 2.29 (s, 3 H), 3.30 (s, 3 H) 3.71 (t, $J=4.9$ Hz, 2 H) 4.36 (t, $J=5.1$ Hz, 2 H) 7.72-7.77 (m, 1 H) 7.82-7.87 (m, 1 H) 8.15 (s, 1 H); MS (DCI/ NH_3) m/z 427 ($\text{M}+\text{H}$) $^+$. Anal. Calculated for $\text{C}_{17}\text{H}_{16}\text{F}_6\text{N}_2\text{O}_2\text{S}$: C, 47.89; H, 3.78; N, 6.57. Found: C, 47.49; H, 3.42; N, 6.38.

Example 37

30 2-fluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-6-(trifluoromethyl)benzamide

The product of Example 12A (0.20 g, 1.1 mmol) and 2-fluoro-6-trifluoromethylbenzoyl chloride (0.17 mL, 1.2 mmol) were processed using the method described in Example 11. Purification by preparative HPLC on a Waters

Symmetry C8 column (40 mm X 100 mm, 7 μ m particle size) using a gradient of 20% to 95% acetonitrile:0.1% aqueous TFA over 12 minutes (15 minutes run time) at a flow rate of 70 mL/min afforded 0.13 g (32%) of the title compound. ^1H NMR (CDCl_3 , 300 MHz) δ ppm 2.27 (s, 3 H), 2.28 (s, 3 H), 3.27 (s, 3 H), 3.69 (t, $J=4.9$ Hz, 2 H), 4.37 (t, $J=4.6$ Hz, 2 H), 7.28-7.33 (m, 1 H), 7.38-7.50 (m, 2 H); MS (DCI/ NH_3) m/z 377 ($\text{M}+\text{H}$) $^+$. Anal. Calculated for $\text{C}_{16}\text{H}_{16}\text{F}_4\text{N}_2\text{O}_2\text{S}$: C, 51.06; H, 4.28; N, 7.44. Found: C, 50.98; H, 4.07; N, 7.36.

Example 38

10 2-chloro-6-fluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 12A (0.20 g, 1.1 mmol) and 2-chloro-6-fluorobenzoyl chloride (0.23 g, 1.2 mmol) were processed using the method described in Example 11. Purification by preparative HPLC on a Waters Symmetry C8 column (40 mm X 100 mm, 7 μ m particle size) using a gradient of 20% to 95% acetonitrile:0.1% aqueous TFA over 12 minutes (15 minutes run time) at a flow rate of 70 mL/min afforded 66 mg (18%) of the title compound. ^1H NMR (CDCl_3 , 300 MHz) δ ppm 2.26 (s, 3 H), 2.27 (s, 3 H), 3.28 (s, 3 H), 3.73 (t, $J=4.9$ Hz, 2 H), 4.35 (t, $J=4.7$ Hz, 2 H), 6.97-7.05 (m, 1 H), 7.19-7.24 (m, 2 H); MS (DCI/ NH_3) m/z 343 ($\text{M}+\text{H}$) $^+$. Anal. Calculated for $\text{C}_{15}\text{H}_{16}\text{ClFN}_2\text{O}_2\text{S}\cdot 0.2\text{C}_2\text{HF}_3\text{O}_2$: C, 50.59; H, 4.47; N, 7.66. Found: C, 50.70; H, 4.34; N, 7.55.

Example 39

25 3-fluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-(trifluoromethyl)benzamide

The product of Example 12A (0.335 g, 1.9 mmol) and 3-fluoro-2-trifluoromethylbenzoyl chloride (0.47 g, 2.1 mmol) were processed using the method described in Example 11. Purification by preparative HPLC on a Waters Symmetry C8 column (40 mm X 100 mm, 7 μ m particle size) using a gradient of 20% to 95% acetonitrile:0.1% aqueous TFA over 12 minutes (15 minutes run time) at a flow rate of 70 mL/min afforded 0.14 g (20%) of the title compound. ^1H NMR (CDCl_3 , 300 MHz) δ ppm 2.25 (d, $J=0.7$ Hz, 3 H), 2.27 (s, 3 H), 3.28 (s, 3 H), 3.68 (t, $J=5.1$ Hz, 2 H), 4.31 (t, $J=5.1$ Hz, 2 H), 7.18 (dd, $J=11.0, 8.3$ Hz, 1 H), 7.38-7.42 (m, 1 H), 7.52

(td, $J=8.0, 5.1$ Hz, 1 H); MS (DCI/ NH_3) m/z 377 ($\text{M}+\text{H}$)⁺. Anal. Calculated for $\text{C}_{16}\text{H}_{16}\text{F}_4\text{N}_2\text{O}_2\text{S}$: C, 51.06; H, 4.28; N, 7.44. Found: C, 51.15; H, 3.96; N, 7.38

Example 40

5 2-chloro-5-fluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 12A (0.20 g, 1.1 mmol) and 2-chloro-5-fluorobenzoyl chloride (0.23 g, 1.2 mmol) were processed using the method described in Example 11. Purification by preparative HPLC on a Waters Symmetry C8 column (40 mm X 100 mm, 7 μm particle size) using a gradient of 20% to 95% acetonitrile:0.1% aqueous TFA over 12 minutes (15 minutes run time) at a flow rate of 70 mL/min afforded 17 mg (4%) of the title compound. ¹H NMR ($\text{DMSO}-d_6$, 500 MHz) δ ppm 2.23 (s, 3 H), 2.26 (s, 3 H), 3.22 (s, 3 H), 3.68 (t, $J=5.3$ Hz, 2 H), 4.33 (t, $J=5.3$ Hz, 2 H), 7.30 (td, $J=8.4, 3.1$ Hz, 1 H), 7.52 (dd, $J=8.9, 5.2$ Hz, 1 H), 7.64 (dd, $J=9.1, 3.3$ Hz, 1 H); MS (DCI/ NH_3) m/z 343 ($\text{M}+\text{H}$)⁺. Anal. Calculated for $\text{C}_{15}\text{H}_{16}\text{ClFN}_2\text{O}_2\text{S} \cdot 0.1\text{C}_2\text{HF}_3\text{O}_2$: C, 51.54; H, 4.58; N, 7.91. Found: C, 51.68; H, 4.35; N, 7.95.

Example 41

20 N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide

The product of Example 12A (1.5 g, 8.0 mmol) and 2,2,3,3-tetramethylcyclopropanecarboxylic acid (0.77 g, 5.4 mmol) were processed using the methods described in Example 13. Recrystallization from ethyl acetate afforded 0.99 g (60%) of the title compound. ¹H NMR ($\text{DMSO}-d_6$, 300 MHz) δ ppm 1.16 (s, 6 H), 1.23 (s, 6 H), 1.44 (s, 1 H), 2.13 (s, 3 H), 2.19 (s, 3 H), 3.24 (s, 3 H), 3.61 (t, $J=5.4$ Hz, 2 H), 4.21 (t, $J=5.4$ Hz, 2 H); MS (DCI/ NH_3) m/z 311 ($\text{M}+\text{H}$)⁺. Anal. Calculated for $\text{C}_{16}\text{H}_{26}\text{N}_2\text{O}_2\text{S}$: C, 61.92; H, 8.44; N, 9.02. Found: C, 61.89; H, 8.38; N, 8.81.

30 Example 42

N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-methylpentanamide

The product of Example 12A (39 mg, 0.15 mmol) and 2-methylvaleric acid

(26 mg, 0.22 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (DMSO-*d*₆, 300 MHz) δ ppm 0.85 (t, 3 H) 1.09 (d, 3 H) 1.20 - 1.28 (m, 2 H) 1.32 - 1.40 (m, 1 H) 1.59 - 1.67 (m, 1 H) 2.19 (s, 3 H) 2.22 (s, 3 H) 2.52 - 2.57 (m, 1 H) 3.24 (s, 3 H) 3.63 (t, 2 H) 4.26 - 4.33 (m, 2 H) MS (ESI) m/z 285 (M+H)⁺.

Example 43

N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2,2-dimethylbutanamide

10 The product of Example 12A (39 mg, 0.15 mmol) and 2,2-dimethylbutyric acid (26 mg, 0.22 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (DMSO-*d*₆, 300 MHz) δ ppm 0.72 (t, 3 H) 1.11 (s, 6 H) 1.53 - 1.59 (m, 2 H) 2.15 (s, 3 H) 2.19 (s, 3 H) 3.24 (s, 3 H) 3.63 (t, 2 H) 4.22 (t, 2 H) MS (ESI) m/z 285 (M+H)⁺.

Example 44

2-ethyl-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]butanamide

20 The product of Example 12A (39 mg, 0.15 mmol) and 2-ethylbutyric acid (26 mg, 0.22 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (DMSO-*d*₆, 300 MHz) δ ppm 0.80 (t, 6 H) 1.48 (s, 2 H) 1.56 - 1.65 (m, 2 H) 2.19 (s, 3 H) 2.23 (s, 3 H) 2.27 - 2.33 (m, 1 H) 3.23 (s, 3 H) 3.63 (t, 2 H) 4.24 - 4.33 (m, 2 H) MS (ESI) m/z 285 (M+H)⁺.

Example 45

N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]cyclohexanecarboxamide

25 The product of Example 12A (39 mg, 0.15 mmol) and cyclohexanecarboxylic acid (28 mg, 0.22 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (DMSO-*d*₆, 300 MHz) δ ppm 1.23 - 1.32 (m, 2 H) 1.34 - 1.43 (m, 2 H) 1.58 - 1.65 (m, 1 H) 1.67 - 1.74 (m, 2 H) 1.83 - 1.89 (m, 2 H) 2.19 (s, 3 H) 2.23 (s, 3 H) 2.35 - 2.42 (m, 1 H) 3.24 (s, 3 H) 3.64 (t, 2 H) 4.31 (t, 3 H) MS (ESI) m/z 297 (M+H)⁺.

Example 46N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-1-methylcyclohexanecarboxamide

5 The product of Example 12A (0.30 g, 1.1 mmol) and 1-methylcyclohexanecarboxylic acid (0.32 g, 2.2 mmol) were processed using the methods described in Example 13. Purification by column chromatography (SiO₂, 30-50% ethyl acetate/hexanes gradient) afforded 80 mg (23%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.16 (s, 3 H), 1.19-1.58 (m, 10 H), 2.18 (s, 3 H), 2.20 (s, 3 H), 3.30 (s, 3 H), 3.69 (t, *J*=5.3 Hz, 2 H), 4.19-4.31 (m, 2 H); MS (DCI/NH₃) *m/z* 311 (M+H)⁺. Anal. Calculated for C₁₆H₂₆N₂O₂S: C, 61.90; H, 8.44; N, 9.02. Found: C, 61.86; H, 8.80; N, 9.02.

Example 47

15 cis-N-[(2Z)-3-(2-Methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-methylcyclohexanecarboxamide

 The product of Example 12A (0.30 g, 1.1 mmol) and (cis)-2-methylcyclohexanecarboxylic acid (0.32 g, 2.2 mmol) were processed using the methods described in Example 13. Purification by column chromatography (SiO₂, 30-50% ethyl acetate/hexanes gradient) afforded 0.24 g (68%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 0.85 (d, *J*=7.1 Hz, 3 H), 1.36-1.84 (m, 8 H), 2.17 (s, 3 H), 2.20 (s, 3 H), 2.31-2.42 (m, 1 H), 2.53-2.65 (m, 1 H), 3.29 (s, 3 H), 3.69 (t, *J*=4.2 Hz, 2 H), 4.17-4.29 (m, 2 H); MS (DCI/NH₃) *m/z* 311 (M+H)⁺. Anal. Calculated for C₁₆H₂₆N₂O₂S: C, 61.90; H, 8.44; N, 9.02. Found: C, 62.15; H, 8.70; N, 8.73.

Example 48N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-4-methylcyclohexanecarboxamide

30 The product of Example 12A (39 mg, 0.15 mmol) and 4-methylcyclohexanecarboxylic acid (31 mg, 0.22 mmol) were processed using the methods described in Example 13 to afford the title compound. ¹H NMR (DMSO-*d*₆, 300 MHz) δ ppm 0.84 – 0.89 (m, 3 H) 1.15 – 1.23 (m, 2 H) 1.47 – 1.56 (m, 4 H) 1.95

– 2.03 (m, 2 H) 2.19 (s, 3 H) 2.23 (s, 3 H) 3.24 (s, 3 H) 3.60 – 3.67 (m, 2 H) 4.26 – 4.36 (m, 4 H) MS (ESI) m/z 311 (M+H)⁺.

Example 49

5 N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]cycloheptanecarboxamide

The product of Example 12A (39 mg, 0.15 mmol) and cycloheptanecarboxylic acid (31 mg, 0.22 mmol) were processed using the methods described in Example 13 to afford the title compound. ¹H NMR (DMSO-*d*₆, 300 MHz) δ ppm 1.45 – 1.52 (m, 4
10 H) 1.54 – 1.56 (m, 2 H) 1.60 – 1.70 (m, 4 H) 1.85 – 1.92 (m, 2 H) 2.19 (s, 3 H) 2.23 (s, 3 H) 2.55 – 2.61 (m, 1 H) 3.24 (s, 3 H) 3.61 – 3.66 (m, 2 H) 4.27 – 4.34 (m, 2 H) MS (ESI) m/z 311 (M+H)⁺.

Example 50

15 (1S)-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]spiro[2.5]octane-1-carboxamide

The product of Example 12A and (1S)-spiro[2.5]octane-1-carboxylic acid (Bennani, Y.L., et al. US 20042043961) were processed using the methods described
20 in Example 13 to provide the title compound. MS (DCI/NH₃) m/z 323 (M+H)⁺.

Example 51

(2R)-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-propylhex-4-ynamide

25 The product of Example 12A (0.30 g, 1.1 mmol) and (2R)-propyl-hex-4-ynoic acid (0.35 g, 2.2 mmol) were processed using the methods described in Example 13. Purification by column chromatography (SiO₂, 20-30% ethyl acetate/hexanes gradient) afforded 0.30 g (82%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 0.90 (t, *J*=7.5 Hz, 3 H), 1.27-1.39 (m, 2 H), 1.62-1.72 (m, 2 H), 1.75 (t, *J*=2.4
30 Hz, 3 H), 2.19 (s, 3 H), 2.22 (s, 3 H), 2.31-2.74 (m, 3 H), 3.29 (s, 3 H), 3.65-3.75 (m, 2 H), 4.16-4.33 (m, 2 H); MS (DCI/NH₃) m/z 323 (M+H)⁺. Anal. Calculated for C₁₇H₂₆N₂O₂S: C, 63.32; H, 8.13; N, 8.69. Found: C, 63.12; H, 8.35; N, 8.51.

Example 52(1S,3R,5S)-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-3,5-dimethylcyclohexanecarboxamide

The product of Example 12A (0.30 g, 1.1 mmol) and (1S,3R,5S)-
 5 dimethylcyclohexanecarboxylic acid (0.35 g, 2.2 mmol) were processed using the
 methods described in Example 13. Purification by column chromatography (SiO₂,
 20-40% ethyl acetate/hexanes gradient) afforded 0.12 g (33%) of the title compound.
¹H NMR (CDCl₃, 300 MHz) δ ppm 0.91 (s, 3 H), 0.92-0.94 (m, 3 H), 1.01-1.13 (m, 2
 H), 1.61-1.69 (m, 3 H), 1.90-2.00 (m, 3 H), 2.19 (s, 3 H), 2.21-2.25 (m, 3 H), 2.26-
 10 2.30 (m, 1 H), 3.30 (s, 3 H), 3.71 (t, *J*=5.3 Hz, 2 H), 4.19-4.44 (m, 2 H); MS
 (DCI/NH₃) *m/z* 325 (M+H)⁺. Anal. Calculated for C₁₇H₂₈N₂O₂S: C, 62.93; H, 8.70;
 N, 8.63. Found: C, 63.29; H, 8.91; N, 8.71.

Example 53

15 (9R,1R,8S)-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-
ylidene]bicyclo[6.1.0]nonane-9-carboxamide

(endo)-Bicyclo[6.1.0]nonane-9-carboxylic acid (0.38 g, 2.2 mmol, Bennani,
 Y.L., et al., US2004077617) and the product of Example 12A (0.30 g, 1.1 mmol)
 were processed using the methods described in Example 13. Purification by column
 20 chromatography (SiO₂, 20-35% ethyl acetate/hexanes gradient) afforded 0.27 g (72%)
 of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.03-1.20 (m, 1 H), 1.29-
 1.51 (m, 6 H), 1.52-1.77 (m, 6 H), 2.07 (dd, *J*=14.1, 2.9 Hz, 2 H), 2.17 (s, 3 H), 2.20
 (s, 3 H), 3.31 (s, 3 H), 3.70 (t, *J*=4.9 Hz, 2 H), 4.20-4.30 (m, 2 H); MS (DCI/NH₃) *m/z*
 337 (M+H)⁺. Anal. Calculated for C₁₈H₂₈N₂O₂S: C, 64.25; H, 8.39; N, 8.32. Found:
 25 C, 64.06; H, 8.54; N, 8.22.

Example 54

(9S,1R,8S)-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-
ylidene]bicyclo[6.1.0]nonane-9-carboxamide

30 (exo)-Bicyclo[6.1.0]nonane-9-carboxylic acid (0.38 g, 2.2 mmol, Bennani,
 Y.L., et al., US2004077617) and the product of Example 12A (0.30 g, 1.1 mmol)
 were processed using the methods described in Example 13. Purification by column
 chromatography (SiO₂, 20-30% ethyl acetate/hexanes gradient) afforded 70 mg (19%)

of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.17-1.29 (m, 2 H), 1.34-1.49 (m, 6 H), 1.57-1.76 (m, 6 H), 1.95-2.04 (m, 1 H), 2.15 (s, 3 H), 2.20 (s, 3 H), 3.30 (s, 3 H), 3.70 (t, *J*=5.3 Hz, 2 H), 4.24 (t, *J*=5.3 Hz, 2 H); MS (DCI/NH₃) *m/z* 337 (M+H)⁺. Anal. Calculated for C₁₈H₂₈N₂O₂S: C, 64.25; H, 8.39; N, 8.32. Found: C, 64.33; H, 8.52; N, 8.23.

Example 55

(1R,6R,7R)-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-1-methylbicyclo[4.1.0]heptane-7-carboxamide

The product of Example 12A (0.30 g, 1.1 mmol) and 1-methylbicyclo[4.1.0]heptane-7-carboxylic acid (0.35 g, 2.2 mmol, Bennani, Y L., et al., US2004077617) were processed using the methods described in Example 13. Purification by column chromatography (SiO₂, 20-35% ethyl acetate/hexanes gradient) afforded 40 mg (11%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.15-1.42 (m, 7 H), 1.58-1.72 (m, 3 H), 1.77 (d, *J*=5.4 Hz, 1 H), 1.84-2.04 (m, 2 H), 2.15 (s, 3 H), 2.19 (s, 3 H), 3.28-3.33 (m, 3 H), 3.69 (t, *J*=5.3 Hz, 2 H), 4.12-4.39 (m, 2 H); MS (DCI/NH₃) *m/z* 323 (M+H)⁺. Anal. Calculated for C₁₇H₂₆N₂O₂S: C, 63.32; H, 8.13; N, 8.69. Found: C, 63.35; H, 8.3; N, 8.56.

Example 57

2,2,3,3-Tetramethylcyclopropanecarboxylic acid [4,5-dimethyl-3-(2-phenoxy-ethyl)-3H-thiazol-2-ylidene]-amide

Example 57A

4,5-Dimethyl-3-(2-phenoxy-ethyl)-3H-thiazol-2-ylideneamine hydrobromide

A mixture of 4,5-dimethylthiazol-2-ylamine (1.0 g, 7.8 mmol) and (2-bromoethoxy)benzene (1.9 g, 9.4 mmol) were heated neat to 85 °C for 19 hours. The mixture was cooled to ambient temperature and the residue was crystallized from isopropanol. The solid was collected by filtration and dried under vacuum to afford 1.3 g (50%) of the title compound. MS (DCI/NH₃) *m/z* 249 (M+H)⁺.

Example 57B

2,2,3,3-Tetramethylcyclopropanecarboxylic acid [4,5-dimethyl-3-(2-phenoxy-ethyl)-

3H-thiazol-2-ylidene]-amide

The product of Example 57A (0.40 g, 1.2 mmol) and 2,2,3,3-tetramethylcyclopropanecarboxylic acid (0.19 g, 1.3 mmol) were processed using the methods described in Example 13. Purification by column chromatography (SiO₂, 20-40% ethyl acetate/hexanes gradient) afforded 0.14 g (34%) of the title compound. ¹H NMR (CDCl₃, 500 MHz) δ ppm 1.22 (s, 6 H), 1.32-1.35 (m, 6 H), 1.56 (s, 1 H), 2.15 (s, 3 H), 2.27 (s, 3 H), 4.32 (t, *J*=5.5 Hz, 2 H), 4.44 (t, *J*=5.3 Hz, 2 H), 6.90 (d, *J*=8.1 Hz, 2 H), 6.95 (t, *J*=7.3 Hz, 1 H), 7.25-7.29 (m, 2 H); MS (DCI/NH₃) *m/z* 373 (M+H)⁺. Anal. Calculated for C₂₁H₂₈N₂O₂S: C, 67.71; H, 7.58; N, 7.52. Found: C, 67.31; H, 7.70; N, 7.30.

Example 58

N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2,2-dimethyltetrahydro-2H-pyran-4-carboxamide

A mixture of the product of Example 12A (150 mg, 0.56 mmol), 2,2-dimethyltetrahydro-pyran-4-carboxylic acid (127 mg, 0.56 mmol), N-(3-dimethylaminopropyl)-N-ethylcarbodiimide hydrochloride (133 mg, 0.70 mmol), 1-hydroxybenzotriazole (94.5 mg, 0.70 mmol) and triethylamine (312 μL, 2.24 mmol) in 5 mL of THF were stirred overnight at room temperature. The reaction mixture was diluted with ethyl acetate and washed with 1 M aqueous NaHCO₃ and brine. The organic extract was dried (Na₂SO₄), filtered and concentrated. Purification of the residue by chromatography afforded the title compound: ¹H NMR (400 MHz, DMSO-*d*₆) δ 1.13 (s, 3 H) 1.16 (s, 3 H) 1.33 - 1.53 (m, 2 H) 1.67 - 1.77 (m, 2 H) 2.16 (s, 3 H) 2.20 (s, 3 H) 2.59 - 2.72 (m, 1 H) 3.24 (s, 3 H) 3.53 - 3.61 (m, 2 H) 3.63 (t, *J*=5.1 Hz, 2 H) 4.24 (t, *J*=5.4 Hz, 2 H); MS (ESI+) *m/z* 327 (M+H)⁺.

Example 59

2,2,3,3-tetrafluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-1-methylcyclobutanecarboxamide

The product of 12A (0.30 g, 1.6 mmol) and 2,2,3,3-tetrafluoro-1-methylcyclobutanecarboxylic acid (0.37 g, 1.8 mmol) were processed using the methods described in Example 13. Purification by column chromatography (SiO₂, 30-45% ethyl acetate/hexanes gradient) afforded 0.15 g (27%) of the title compound. ¹H

NMR (CDCl₃, 300 MHz) δ ppm 1.59 (s, 3 H), 2.23 (s, 3 H), 2.25 (s, 3 H), 2.29-2.43 (m, 1 H), 3.29 (s, 3 H), 3.31-3.46 (m, 1 H), 3.70 (t, $J=5.1$ Hz, 2 H), 4.26-4.48 (m, 2 H); MS (DCI/NH₃) m/z 355 (M+H)⁺. Anal. Calculated for C₁₄H₁₈F₄N₂O₂S: C, 47.45; H, 5.12; N, 7.91. Found: C, 47.41; H, 5.04; N, 7.81.

5

Example 60

1-hydroxy-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]cyclohexanecarboxamide

Commercially available 1-hydroxy-cyclohexanecarboxylic acid and the product of Example 12A were processed using the method described in Example 58 to afford the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ ppm 1.09 - 1.32 (m, 1 H), 1.36 - 1.68 (m, 7 H), 1.72 - 1.88 (m, 2 H), 2.18 (s, 3 H), 2.22 (s, 3 H), 3.23 (s, 3 H), 3.64 (t, $J=5.3$ Hz, 2 H), 4.29 (t, $J=5.4$ Hz, 2 H), 4.34 (s, 1 H); MS (ESI⁺) m/z 335 (M+Na)⁺; Anal. Calculated for C₁₅H₂₄N₂O₃S: C, 57.66; H, 7.74; N, 8.97. Found: C, 57.76; H, 7.80; N, 8.88.

15

Example 61

1-([(2Z)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]amino)carbonyl)cyclohexyl propionate

Propionyloxy-cyclohexanecarboxylic acid (Hartmann, Willy et al., *Synthesis* (1989), 4, 272-4) and the product from Example 12A were processed using the method described in Example 58 to afford the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ ppm 1.03 (t, $J=7.5$ Hz, 3 H), 1.15 - 1.34 (m, 1 H), 1.36 - 1.66 (m, 5 H), 1.75 (td, $J=13.1, 4.1$ Hz, 2 H), 2.01 - 2.13 (m, 2 H), 2.16 (s, 3 H), 2.20 (s, 3 H), 2.32 (q, $J=7.5$ Hz, 2 H), 3.22 (s, 3 H), 3.58 (t, $J=5.4$ Hz, 2 H), 4.19 (t, $J=5.4$ Hz, 2 H); MS (ESI⁺) m/z 369 (M+H)⁺; Anal. Calculated for C₁₈H₂₈N₂O₄S: C, 58.67; H, 7.66; N, 7.60. Found: C, 58.46; H, 7.64; N, 7.75.

25

Example 62

N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide

30

Example 62A

2,2,3,3-Tetramethylcyclopropanecarboxylic acid benzothiazol-2-ylamide

A mixture of 2,2,3,3-tetramethylcyclopropanecarboxylic acid (0.50 g, 3.5 mmol) and benzothiazol-2-ylamine (0.58 g, 3.9 mmol) were processed as in Example 9A.

Purification by column chromatography (SiO₂, 20-40% ethyl acetate/hexanes gradient) afforded 0.26 g (27%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.22 - 1.27 (m, 6 H), 1.36 (s, 6 H), 1.67 (s, 1 H), 2.07 - 2.18 (m, 2 H), 3.34 (s, 3 H), 3.41 (t, *J*=5.8 Hz, 2 H), 4.39 - 4.53 (m, 2 H), 7.20 - 7.26 (m, 1 H), 7.37 - 7.42 (m, 2 H), 7.60 (d, *J*=7.5 Hz, 1 H); MS (DCI/NH₃) *m/z* 275 (M+H)⁺.

Example 62B

2,2,3,3-Tetramethylcyclopropanecarboxylic acid [3-(2-methoxyethyl)-3H-

benzothiazol-2-ylidene]-amide

The product of Example 62A (0.12 g, 0.43 mmol), 2-bromoethyl methyl ether (0.44 mL, 4.7 mmol) and potassium hydroxide (56 mg, 1.0 mmol) were processed using the method described in Example 1B. Purification by column chromatography (SiO₂, 20-50% ethyl acetate/hexanes gradient) afforded 12 mg (8%) of the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.25 (s, 6 H), 1.36 (s, 6 H), 1.66 (s, 1 H), 3.34 (s, 3 H), 3.79 (t, *J*=5.6 Hz, 2 H), 4.53 (t, *J*=5.6 Hz, 2 H), 7.21-7.25 (m, 1 H), 7.37-7.42 (m, 2 H), 7.58 (d, *J*=7.8 Hz, 1 H); MS (DCI/NH₃) *m/z* 371 (M+H)⁺. Anal. Calculated for C₁₈H₂₄N₂O₂S: C, 65.03; H, 7.28; N, 8.43. Found: C, 64.94; H, 7.10; N, 8.40.

Example 63

N-[(2Z)-3-(3-methoxypropyl)-1,3-benzothiazol-2(3H)-ylidene]-2,2,3,3-

tetramethylcyclopropanecarboxamide

Example 63A

3-(3-Methoxy-propyl)-3H-benzothiazol-2-ylideneamine hydrobromide

Benzothiazol-2-ylamine (1.0 g, 6.6 mmol) and 1-bromo-3-methoxy-propane (1.2 g, 7.9 mmol) were processed using the method described in Example 12A.

Recrystallization from ethyl acetate provided 1.7 g (89%) of the title compound. ¹H NMR (DMSO-d₆, 300 MHz) δ ppm 1.90-2.02 (m, 2 H), 3.18 (s, 3 H), 3.39 (t, *J*=5.9 Hz, 2 H), 4.31 (t, *J*=7.1 Hz, 2 H), 7.37-7.48 (m, 1 H), 7.53-7.69 (m, 2 H), 8.00 (dd, *J*=8.0, 0.8 Hz, 1 H), 10.08 (s, 1 H); MS (DCI/NH₃) *m/z* 233 (M+H)⁺.

Example 63B

2,2,3,3-Tetramethylcyclopropanecarboxylic acid [3-(3-methoxy-propyl)-3H-

benzothiazol-2-ylidene)-amide

The product of Example 63A (0.40 g, 1.3 mmol) and 2,2,3,3-Tetramethylcyclopropanecarboxylic acid (0.19 g, 1.3 mmol) were processed using the methods described in Example 13. Purification by column chromatography (SiO₂, 20-30% ethyl acetate/hexanes gradient) afforded 0.32 g (70%) of the title compound.

- 5 ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.26 (s, 6 H), 1.36 (s, 6 H), 1.64 (br s, 1 H), 1.71 (s, 1 H), 3.87 (s, 3 H), 7.27-7.32 (m, 2 H), 7.40-7.47 (m, 1 H), 7.62 (d, *J*=7.5 Hz, 1 H); MS (DCI/NH₃) *m/z* 289 (M+H)⁺. Anal. Calculated for C₁₆H₂₀N₂O₂S: C, 65.86; H, 7.56; N, 8.08. Found: C, 65.54; H, 7.65; N, 7.81.

10 Example 64

N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-3-phenylpropanamide

Example 64A

3-(2-Methoxyethyl)-3H-benzothiazol-2-ylideneamine hydrobromide

- 15 Benzothiazol-2-ylamine (10.0 g, 66.6 mmol) and 2-bromoethyl methyl ether (9.39 mL, 99.9 mmol) were combined and heated at 85 °C for 6 hours. The dark solid was triturated with EtOH then filtered and dried under vacuum to afford the title compound (15.8 g, 82%). ¹H NMR (DMSO-*d*₆, 300 MHz) δ ppm 3.23 (s, 3 H), 3.69 (t, *J*=5.1 Hz, 2 H), 4.51 (t, *J*=5.1 Hz, 2 H), 7.42 (dt, *J*=1.0, 8.0 Hz, 1 H), 7.56 (m, 1 H), 7.72 (d, *J*=8.0 Hz, 1 H), 8.00 (dd, *J*=1.1, 8.0 Hz, 1H), 10.16 (br s, 2 H); MS (DCI/NH₃) *m/z* 209 (M+H)⁺.
- 20

Example 64B

N-[3-(2-Methoxyethyl)-3H-benzothiazol-2-ylidene]-3-phenylpropionamide

- 25 The product of Example 64A (39 mg, 0.14 mmol) and hydrocinnamic acid (26 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.77 - 2.85 (m, 2 H) 2.92 - 3.05 (m, 2 H) 3.19 - 3.25 (m, 3 H) 3.76 (t, 2 H) 4.52 (t, 2 H) 7.08 - 7.19 (m, 1 H) 7.22 - 7.37 (m, 5 H) 7.41 - 7.53 (m, 1 H) 7.59 - 7.74 (m, 1 H) 7.75 - 8.03 (m, 1 H); MS(ESI) *m/z* 341(M+H)⁺.
- 30

Example 65

(2S)-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-2-

phenylbutanamide

The product of Example 64A (39 mg, 0.14 mmol) and (S)-(+)-2-phenylbutyric acid (26 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.89 (t, 3 H) 1.69 - 1.89 (m, 1 H) 2.04 - 2.23 (m, 1 H) 3.11 - 3.20 (m, 3 H) 3.57 - 3.76 (m, 3 H) 4.54 (t, 2 H) 7.17 - 7.23 (m, 1 H) 7.27 - 7.41 (m, 5 H) 7.43 - 7.53 (m, 1 H) 7.61 - 7.71 (m, 1 H) 7.74 - 7.88 (m, 1 H); MS(ESI) *m/z* 355 (M+H)⁺.

Example 6610 N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-4-thien-2-ylbutanamide

The product of Example 64A (39 mg, 0.14 mmol) and 4-(2-thienyl)butyric acid (29 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 1.86 - 2.08 (m, 2 H) 2.55 (t, 2 H) 2.87 (t, 2 H) 3.17 - 3.24 (m, 3 H) 3.72 (t, 2 H) 4.54 (t, 2 H) 6.83 - 6.89 (m, 1 H) 6.90 - 7.01 (m, 1 H) 7.22 - 7.37 (m, 2 H) 7.42 - 7.55 (m, 1 H) 7.64 - 7.72 (m, 1 H) 7.75 - 7.88 (m, 1 H); MS(ESI) *m/z* 361 (M+H)⁺.

Example 67N²-acetyl-N¹-[3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-L-leucinamide

20 The product of Example 64A (39 mg, 0.14 mmol) and *N*-acetyl-L-leucine (29 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.78 - 1.07 (m, 6 H) 1.43 - 1.57 (m, 1 H) 1.57 - 1.75 (m, 2 H) 1.81 - 1.92 (m, 3 H) 3.22 - 3.26 (m, 3 H) 3.72 (t, 2 H) 4.44 - 4.55 (m, 1 H) 4.55 - 4.67 (m, 2 H) 7.24 - 7.40 (m, 1 H) 7.43 - 7.54 (m, 1 H) 7.62 - 7.73 (m, 1 H) 7.79 - 7.92 (m, 1 H) 7.95 - 8.07 (m, 1 H); MS(ESI) *m/z* 364 (M+H)⁺.

Example 6830 3-(2-chlorophenyl)-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]propanamide

The product of Example 64A (39 mg, 0.14 mmol) and 3-(2-chlorophenyl)propionic acid (31 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆)

δ ppm 2.82 (t, 2 H) 3.08 (t, 2 H) 3.18 - 3.25 (m, 3 H) 3.79 (t, 2 H) 4.54 (t, 2 H) 7.17 - 7.29 (m, 2 H) 7.30 - 7.44 (m, 3 H) 7.45 - 7.54 (m, 1 H) 7.59 - 7.75 (m, 1 H) 7.76 - 7.93 (m, 1 H).

5

Example 69N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-3-methyl-2-phenylpentanamide

The product of Example 64A (39 mg, 0.14 mmol) and 3-methyl-2-phenylvaleric acid (33 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.59 - 0.68 (m, 2 H) 0.72 - 0.82 (m, 2 H) 0.84 - 0.98 (m, 3 H) 1.08 - 1.27 (m, 1 H) 2.21 - 2.39 (m, 1 H) 3.11 - 3.24 (m, 3 H) 3.38 - 3.50 (m, 1 H) 3.65 - 3.82 (m, 2 H) 4.57 (t, 2 H) 7.16 - 7.23 (m, 1 H) 7.25 - 7.36 (m, 3 H) 7.37 - 7.54 (m, 3 H) 7.60 - 7.73 (m, 1 H) 7.75 - 7.88 (m, 1 H); MS(ESI) *m/z* 383(M+H)⁺.

15

Example 704-ethyl-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]benzamide

The product of Example 64A (39 mg, 0.14 mmol) and 4-ethylbenzoic acid (26 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 1.22 (t, 3 H) 2.66 - 2.72 (m, 2 H) 3.25 - 3.26 (m, 3 H) 3.85 (t, 2 H) 4.75 (t, 2 H) 7.31 - 7.42 (m, 3 H) 7.47 - 7.60 (m, 1 H) 7.68 - 7.79 (m, 1 H) 7.85 - 7.94 (m, 1 H) 8.12 - 8.23 (m, 2 H); MS (ESI) *m/z* 341 (M+H)⁺.

25

Example 713-fluoro-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-2-methylbenzamide

The product of Example 64A (39 mg, 0.14 mmol) and 3-fluoro-2-methylbenzoic acid (26 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.51 - 2.53 (m, 3 H) 3.22 - 3.25 (m, 3 H) 3.80 (t, 2 H) 4.69 (t, 2 H) 7.26 - 7.43 (m, 3 H) 7.50 - 7.61 (m, 1 H) 7.70 - 7.79 (m, 1 H) 7.86 - 7.99 (m, 2 H); MS (ESI) *m/z* 345 (M+H)⁺.

Example 725-fluoro-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-2-methylbenzamide

5 The product of Example 64A (39 mg, 0.14 mmol) and 5-fluoro-2-methylbenzoic acid (26 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.51 - 2.53 (m, 3 H) 3.22 - 3.25 (m, 3 H) 3.80 (t, 2 H) 4.69 (t, 2 H) 7.26 - 7.43 (m, 3 H) 7.50 - 7.61 (m, 1 H) 7.70 - 7.79 (m, 1 H) 7.86 - 7.99 (m, 2 H); MS (ESI) m/z 345 (M+H)⁺.

Example 733-fluoro-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-4-methylbenzamide

15 The product of Example 64A (39 mg, 0.14 mmol) and 5-fluoro-4-methylbenzoic acid (26 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.31 - 2.34 (m, 3 H) 3.23 - 3.26 (m, 3 H) 3.83 (t, 2 H) 4.77 (t, 2 H) 7.33 - 7.40 (m, 1 H) 7.41 - 7.49 (m, 1 H) 7.49 - 7.59 (m, 1 H) 7.69 - 7.78 (m, 1 H) 7.86 - 7.95 (m, 2 H) 7.97 - 8.02 (m, 1 H); MS (ESI) m/z 345 (M+H)⁺.

Example 742,3-difluoro-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]benzamide

25 The product of Example 64A (39 mg, 0.14 mmol) and 2,3-difluorobenzoic acid (27 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 3.20 - 3.26 (m, 3 H) 3.82 (t, 2 H) 4.72 (t, 2 H) 7.28 - 7.36 (m, 1 H) 7.37 - 7.43 (m, 1 H) 7.51 - 7.59 (m, 1 H) 7.59 - 7.67 (m, 1 H) 7.73 - 7.81 (m, 1 H) 7.90 - 8.00 (m, 2 H); MS (ESI) m/z 349 (M+H)⁺.

Example 752,5-difluoro-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]benzamide

The product of Example 64A (39 mg, 0.14 mmol) and 2,5-difluorobenzoic

acid (27 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 3.23 (s, 3 H) 3.82 (t, 2 H) 4.72 (t, 2 H) 7.38 (s, 2 H) 7.46 (s, 1 H) 7.54 (s, 1 H) 7.77 (s, 1 H) 7.89 (s, 1 H) 7.92 - 7.98 (m, 1 H); MS (ESI) *m/z* 349 (M+H)⁺.

5

Example 76

2-acetyl-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]benzamide

The product of Example 64A (39 mg, 0.14 mmol) and 2-acetylbenzoic acid (28 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.44 - 2.48 (m, 3 H) 3.21 - 3.25 (m, 3 H) 3.77 (t, 2 H) 4.67 (t, 2 H) 7.30 - 7.44 (m, 2 H) 7.51 - 7.66 (m, 3 H) 7.70 - 7.82 (m, 1 H) 7.89 - 8.00 (m, 1 H) 8.11 - 8.25 (m, 1 H); MS (ESI) *m/z* 355 (M+H)⁺.

15

Example 77

3-methoxy-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-4-methylbenzamide

The product of Example 64A (39 mg, 0.14 mmol) and 3-methoxy-4-methylbenzoic acid (28 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.21 - 2.26 (m, 3 H) 3.25 - 3.28 (m, 3 H) 3.85 (t, 2 H) 3.88 - 3.90 (m, 3 H) 4.76 (t, 2 H) 7.26 - 7.31 (m, 1 H) 7.32 - 7.39 (m, 1 H) 7.47 - 7.57 (m, 1 H) 7.71 - 7.78 (m, 2 H) 7.79 - 7.84 (m, 1 H) 7.88 - 7.93 (m, 1 H); MS (ESI) *m/z* 357 (M+H)⁺.

25

Example 78

2-ethoxy-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]benzamide

The product of Example 64A (39 mg, 0.14 mmol) and 2-ethoxybenzoic acid (28 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.21 - 2.26 (m, 3 H) 3.25 - 3.28 (m, 3 H) 3.85 (t, 2 H) 3.88 - 3.90 (m, 3 H) 4.76 (t, 2 H) 7.26 - 7.31 (m, 1 H) 7.32 - 7.39 (m, 1 H) 7.47 - 7.57 (m, 1 H) 7.71 - 7.78 (m, 2 H) 7.79 - 7.84 (m, 1 H) 7.88 - 7.93 (m, 1 H); MS (ESI) *m/z* 357 (M+H)⁺.

Example 79N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-4-(methylthio)benzamide

5 The product of Example 64A (39 mg, 0.14 mmol) and 4-methylsulfanylbobenzoic acid (29 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.53 - 2.58 (m, 3 H) 3.22 - 3.27 (m, 3 H) 3.84 (t, 2 H) 4.73 (t, 2 H) 7.31 - 7.44 (m, 3 H) 7.47 - 7.61 (m, 1 H) 7.67 - 7.79 (m, 1 H) 7.82 - 8.02 (m, 1 H) 8.08 - 8.22 (m, 2 H); MS (ESI) m/z 359 (M+H)⁺.

10

Example 80N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-1-naphthamide

15 The product of Example 64A (39 mg, 0.14 mmol) and 1-naphthoic acid (29 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 3.23 - 3.27 (m, 3 H) 3.90 (t, 2 H) 4.71 (t, 2 H) 7.32 - 7.42 (m, 1 H) 7.45 - 7.68 (m, 4 H) 7.71 - 7.79 (m, 1 H) 7.83 - 8.22 (m, 3 H) 8.40 - 8.56 (m, 1 H) 9.05 - 9.17 (m, 1 H); MS (ESI) m/z 363 (M+H)⁺.

20

Example 81N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-2-naphthamide

25 The product of Example 64A (39 mg, 0.14 mmol) and 2-naphthoic acid (29 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 3.28 - 3.29 (m, 3 H) 3.91 (t, 2 H) 4.78 (t, 2 H) 7.25 - 7.44 (m, 1 H) 7.50 - 7.58 (m, 1 H) 7.58 - 7.67 (m, 2 H) 7.71 - 7.82 (m, 1 H) 7.88 - 7.97 (m, 1 H) 7.99 - 8.07 (m, 2 H) 8.08 - 8.21 (m, 1 H) 8.26 - 8.40 (m, 1 H) 8.72 - 8.93 (m, 1 H); MS (ESI) m/z 363 (M+H)⁺.

30

Example 825-chloro-2-hydroxy-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]benzamide

The product of Example 64A (39 mg, 0.14 mmol) and 5-chloro-2-

hydroxybenzoic acid (29 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 3.22 - 3.25 (m, 3 H) 3.85 (t, 2 H) 4.73 (t, 2 H) 6.94 - 7.08 (m, 2 H) 7.36 - 7.53 (m, 2 H) 7.57 - 7.67 (m, 1 H) 7.79 - 7.92 (m, 1 H) 7.96 - 8.09 (m, 2 H); MS (ESI) *m/z* 363 (M+H)⁺.

Example 83

5-chloro-2-methoxy-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]benzamide

10 The product of Example 64A (39 mg, 0.14 mmol) and 5-chloro-2-methoxybenzoic acid (32 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 3.22 - 3.26 (m, 3 H) 3.80 (t, 2 H) 3.83 - 3.88 (m, 3 H) 4.62 (t, 2 H) 7.13 - 7.23 (m, 1 H) 7.32 - 7.43 (m, 1 H) 7.46 - 7.59 (m, 2 H) 7.69 - 7.77 (m, 1 H) 7.79 - 7.88 (m, 1 H) 7.88 - 7.98 (m, 1 H); MS (ESI) *m/z* 377 (M+H)⁺.

Example 84

1-hydroxy-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-2-naphthamide

20 The product of Example 64A (39 mg, 0.14 mmol) and 1-hydroxy-2-naphthoic acid (32 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 3.25 - 3.28 (m, 3 H) 3.89 (t, 2 H) 4.75 (t, 2 H) 7.35 - 7.47 (m, 2 H) 7.53 - 7.63 (m, 2 H) 7.63 - 7.70 (m, 1 H) 7.79 - 7.93 (m, 2 H) 7.96 - 8.06 (m, 1 H) 8.04 - 8.19 (m, 1 H) 8.23 - 8.40 (m, 1 H); MS (ESI) *m/z* 379 (M+H)⁺.

Example 85

4-fluoro-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-1-naphthamide

30 The product of Example 64A (39 mg, 0.14 mmol) and 4-fluoro-1-naphthoic acid (32 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 3.22 - 3.26 (m, 3 H) 3.83 (t, 2 H) 4.67 (t, 2 H) 7.35 - 7.59 (m, 3 H) 7.65 - 7.82 (m, 3 H) 7.83 - 8.03 (m,

Example 892-benzyl-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]benzamide

The product of Example 64A (39 mg, 0.14 mmol) and α -phenyl-*o*-toluic acid (36 mg, 0.17 mmol) were processed using the method described in Example 3 to

5 afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 3.18 - 3.20 (m, 3 H) 3.72 (t, 2 H) 4.48 - 4.52 (m, 2 H) 4.63 (t, 2 H) 7.07 - 7.30 (m, 6 H) 7.30 - 7.41 (m, 2 H) 7.40 - 7.47 (m, 1 H) 7.49 - 7.57 (m, 1 H) 7.67 - 7.79 (m, 1 H) 7.85 - 7.96 (m, 1 H) 8.02 - 8.12 (m, 1 H); MS (ESI) *m/z* 403 (M+H)⁺.

Example 902-chloro-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-5-(trifluoromethyl)benzamide

The product of Example 64A (39 mg, 0.14 mmol) and 2-chloro-5-trifluoromethylbenzoic acid (38 mg, 0.17 mmol) were processed using the method

15 described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 3.20 - 3.25 (s, 3 H) 3.80 (t, 2 H) 4.70 (t, 2 H) 7.35 - 7.45 (m, 1 H) 7.50 - 7.65 (m, 1 H) 7.76 - 7.83 (m, 2 H) 7.84 - 7.91 (m, 1 H) 7.94 - 8.03 (m, 1 H) 8.22 - 8.32 (m, 1 H); MS (ESI) *m/z* 415 (M+H)⁺.

Example 91N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-2-(2-phenylethyl)benzamide

The product of Example 64A (39 mg, 0.14 mmol) and 2-phenethylbenzoic acid (38 mg, 0.17 mmol) were processed using the method described in Example 3 to

25 afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.78 - 2.95 (m, 2 H) 3.16 - 3.22 (m, 3 H) 3.32 - 3.37 (m, 2 H) 3.76 (t, 2 H) 4.65 (t, 2 H) 6.99 - 7.46 (m, 9 H) 7.49 - 7.57 (m, 1 H) 7.67 - 7.80 (m, 1 H) 7.82 - 7.97 (m, 1 H) 7.97 - 8.15 (m, 1 H); MS (ESI) *m/z* 417 (M+H)⁺.

Example 922-bromo-5-methoxy-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]benzamide

The product of Example 64A (39 mg, 0.14 mmol) and 2-bromo-5-

methoxybenzoic acid (39 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 3.20 - 3.26 (m, 3 H) 3.72 - 3.87 (m, 5 H) 4.67 (t, 2 H) 6.91 - 7.08 (m, 1 H) 7.30 - 7.43 (m, 1 H) 7.45 - 7.51 (m, 1 H) 7.51 - 7.65 (m, 2 H) 7.70 - 7.83 (m, 1 H) 7.86 - 7.98 (m, 1 H); MS (ESI) *m/z* 423 (M+H)⁺.

Example 93

2-iodo-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]benzamide

The product of Example 64A (39 mg, 0.14 mmol) and 2-iodobenzoic acid (42 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 3.20 - 3.24 (m, 3 H) 3.81 (t, 2 H) 4.69 (t, 2 H) 7.15 - 7.25 (m, 1 H) 7.34 - 7.41 (m, 1 H) 7.47 - 7.62 (m, 2 H) 7.70 - 7.81 (m, 1 H) 7.87 - 7.97 (m, 2 H) 7.97 - 8.03 (m, 1 H); MS (ESI) *m/z* 439 (M+H)⁺.

Example 94

3-iodo-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]benzamide

The product of Example 64A (39 mg, 0.14 mmol) and 3-iodobenzoic acid (42 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 3.26 - 3.27 (m, 3 H) 3.79 (t, 2 H) 4.76 (t, 2 H) 7.26 - 7.43 (m, 2 H) 7.48 - 7.61 (m, 1 H) 7.69 - 7.81 (m, 1 H) 7.89 - 8.00 (m, 2 H) 8.18 - 8.31 (m, 1 H) 8.49 - 8.59 (m, 1 H); MS (ESI) *m/z* 439 (M+H)⁺.

Example 95

4-iodo-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]benzamide

The product of Example 64A (39 mg, 0.14 mmol) and 4-iodobenzoic acid (42 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 3.23 - 3.25 (m, 3 H) 3.80 (t, 2 H) 4.75 (t, 2 H) 7.29 - 7.45 (m, 1 H) 7.47 - 7.62 (m, 1 H) 7.70 - 7.80 (m, 1 H) 7.85 - 7.96 (m, 3 H) 7.98 - 8.05 (m, 2 H); MS (ESI) *m/z* 439 (M+H)⁺.

Example 96

N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-3-methylbutanamide

The product of Example 64A (39 mg, 0.14 mmol) and isovaleric acid (17 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.93 (d, 6 H) 2.10 - 2.23 (m, 1 H) 2.37 (d, 2 H) 3.21 - 3.25 (m, 3 H) 3.74 (t, 2 H) 4.57 (t, 2 H) 7.31 (m, 1 H) 7.48 (m, 1 H) 7.66 (m, 1 H) 7.83 (m, 1 H); MS (ESI) *m/z* 393 (M+H)⁺.

Example 97N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-2-methylpentanamide

The product of Example 64A (39 mg, 0.14 mmol) and 2-methylvaleric acid (20 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.88 (t, 3 H) 1.14 (d, 3 H) 1.21 - 1.34 (m, 2 H) 1.35 - 1.47 (m, 1 H) 1.63 - 1.76 (m, 1 H) 2.53 - 2.62 (m, 1 H) 3.22 - 3.26 (m, 3 H) 3.75 (t, 2 H) 4.58 (t, 2 H) 7.31 (m, 1 H) 7.49 (m, 1 H) 7.66 (m, 1 H) 7.83 (m, 1 H); MS (ESI) *m/z* 307 (M+H)⁺.

Example 98N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-3-methylpentanamide

The product of Example 64A (39 mg, 0.14 mmol) and 3-methylvaleric acid (20 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.82 - 0.96 (m, 6 H) 1.14 - 1.27 (m, 1 H) 1.31 - 1.44 (m, 1 H) 1.87 - 2.05 (m, 1 H) 2.25 - 2.34 (m, 1 H) 2.43 - 2.49 (m, 1 H) 3.21 - 3.26 (m, 3 H) 3.74 (t, 2 H) 4.56 (t, 2 H) 7.31 (m, 1 H) 7.48 (m, 1 H) 7.66 (m, 1 H) 7.84 (m, 1 H); MS (ESI) *m/z* 307 (M+H)⁺.

Example 99N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-4-methylpentanamide

The product of Example 64A (39 mg, 0.14 mmol) and 4-methylvaleric acid (20 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.90 (d, 6 H) 1.49 - 1.61 (m, 3 H) 2.45 - 2.50 (m, 2 H) 3.21 - 3.26 (m, 3 H) 3.74 (t, 2 H) 4.58 (t, 2 H) 7.32 (m, 1 H) 7.48 (m, 1 H) 7.66 (m, 1 H) 7.84 (m, 1 H); MS (ESI) *m/z* 307 (M+H)⁺.

Example 100N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-2,2-dimethylbutanamide

The product of Example 64A (39 mg, 0.14 mmol) and 2,2-dimethylbutyric acid (20 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.76 (t, 3 H) 1.11 - 1.23 (m, 6 H) 1.62 (t, 2 H) 3.18 - 3.27 (m, 3 H) 3.75 (t, 2 H) 4.58 (t, 2 H) 7.26 (m, 1 H) 7.46 (m, 1 H) 7.66 (m, 1 H) 7.84 (m, 1 H); MS (ESI) *m/z* 307 (M+H)⁺.

10

Example 101N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-3,3-dimethylbutanamide

The product of Example 64A (0.64 g, 2.2 mmol) and 3,3-dimethyl-butyric acid (0.26 mL, 2.0 mmol) were processed using the method described in Example 3. Purification by column chromatography (SiO₂, 20-30% ethyl acetate/hexanes gradient) afforded 0.44 g (71%) of the title compound. ¹H NMR (DMSO-*d*₆, 300 MHz) δ ppm 1.03 (s, 9 H), 2.36-2.43 (s, 2 H), 3.22 (s, 3 H), 3.73 (t, *J*=5.4 Hz, 2 H), 4.56 (t, *J*=5.4 Hz, 2 H), 7.32 (m, 1 H), 7.49 (td, *J*=7.8, 1.0 Hz, 1 H), 7.67 (d, *J*=8.5 Hz, 1 H), 7.84 (dd, *J*=8.0, 1.2 Hz, 1 H); MS (DCI/NH₃) *m/z* 331 (M+H)⁺. Anal. Calculated for C₁₆H₂₂N₂O₂S: C, 62.71; H, 7.24; N, 9.14. Found: C, 62.79; H, 7.41; N, 9.06.

20

Example 1022-ethyl-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]butanamide

The product of Example 64A (39 mg, 0.14 mmol) and 2-ethylbutyric acid (20 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.84 (t, 6 H) 1.47 - 1.59 (m, 2 H) 1.60 - 1.72 (m, 2 H) 2.27 - 2.39 (m, 1 H) 3.21 - 3.26 (m, 3 H) 3.75 (t, 2 H) 4.58 (t, 2 H) 7.32 (m, 1 H) 7.48 (m, 1 H) 7.65 (m, 1 H) 7.82 (m, 1 H); MS (ESI) *m/z* 307 (M+H)⁺.

30

Example 103N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-

ylidene]cyclopentanecarboxamide

The product of Example 64A (39 mg, 0.14 mmol) and cyclopentanecarboxylic acid (19 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 1.50 - 1.62 (m, 2 H) 1.62 - 1.72 (m, 2 H) 1.76 - 1.95 (m, 4 H) 2.85 - 2.98 (m, 1 H) 3.22 - 3.26 (m, 3 H) 3.74 (t, 2 H) 4.56 (m, 2 H) 7.31 (m, 1 H) 7.48 (m, 1 H) 7.66 (m, 1 H) 7.83 (d, 1 H); MS (ESI) m/z 305 (M+H)⁺.

Example 10410 2-cyclopentyl-N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]acetamide

The product of Example 64A (39 mg, 0.14 mmol) and cyclopentylacetic acid (22 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 1.09 - 1.25 (m, 2 H) 1.44 - 1.56 (m, 2 H) 1.55 - 1.65 (m, 2 H) 1.70 - 1.84 (m, 2 H) 2.20 - 2.38 (m, 1 H) 2.46 - 2.49 (m, 1 H) 3.21 - 3.24 (m, 3 H) 3.24 - 3.29 (m, 1 H) 3.74 (t, 2 H) 4.56 (t, 2 H) 7.32 (m, 1 H) 7.49 (m, 1 H) 7.65 (m, 1 H) 7.83 (m, 1 H); MS (ESI) m/z 319 (M+H)⁺.

Example 10520 N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]cyclohexanecarboxamide

The product of Example 64A (39 mg, 0.14 mmol) and cyclohexanecarboxylic acid (22 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 1.13 - 1.36 (m, 3 H) 1.37 - 1.52 (m, 2 H) 1.58 - 1.67 (m, 1 H) 1.66 - 1.80 (m, 2 H) 1.84 - 2.00 (m, 2 H) 2.32 - 2.46 (m, 1 H) 3.22 - 3.25 (m, 3 H) 3.75 (t, 2 H) 4.56 (t, 2 H) 7.30 (m, 1 H) 7.48 (m, 1 H) 7.65 (m, 1 H) 7.83 (m, 1 H); MS (ESI) m/z 319 (M+H)⁺.

Example 10630 N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-1-methylcyclohexanecarboxamide

The product of Example 64A (39 mg, 0.14 mmol) and 1-methylcyclohexanecarboxylic acid (24 mg, 0.17 mmol) were processed using the method described in

Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 1.10 - 1.16 (m, 3 H) 1.21 - 1.39 (m, 5 H) 1.40 - 1.48 (m, 1 H) 1.48 - 1.58 (m, 2 H) 2.06 - 2.21 (m, 2 H) 3.21 - 3.25 (m, 3 H) 3.76 (t, 2 H) 4.58 (t, 2 H) 7.30 (m, 1 H) 7.47 (m, 1 H) 7.65 (m, 1 H) 7.82 (m, 1 H); MS (ESI) *m/z* 333 (M+H)⁺.

5

Example 107

cis-N-[(2*Z*)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3*H*)-ylidene]-2-methylcyclohexanecarboxamide

The product of Example 64A (39 mg, 0.14 mmol) and (*cis*)-2-methylcyclohexanecarboxylic acid (24 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.79 (d, 3 H) 1.14 - 1.87 (m, 8 H) 2.24 - 2.39 (m, 1 H) 2.56 - 2.65 (m, 1 H) 3.19 - 3.23 (m, 3 H) 3.72 (t, 2 H) 4.54 (t, 2 H) 7.22 - 7.37 (m, 1 H) 7.42 - 7.50 (m, 1 H) 7.58 - 7.67 (m, 1 H) 7.75 - 7.88 (m, 1 H); MS (ESI) *m/z* 333 (M+H)⁺.

15

Example 108

N-[(2*Z*)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3*H*)-ylidene]-4-methylcyclohexanecarboxamide

The product of Example 64A (39 mg, 0.14 mmol) and 4-methylcyclohexanecarboxylic acid (24 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.78 - 0.91 (m, 3 H) 0.94 - 1.26 (m, 2 H) 1.38 - 1.63 (m, 4 H) 1.69 - 2.01 (m, 1 H) 2.04 - 2.40 (m, 2 H) 2.53 - 2.65 (m, 1 H) 3.19 - 3.26 (m, 3 H) 3.76 (t, 2 H) 4.56 (t, 2 H) 7.23 - 7.37 (m, 1 H) 7.40 - 7.56 (m, 1 H) 7.59 - 7.70 (m, 1 H) 7.76 - 7.86 (m, 1 H); MS (ESI) *m/z* 333 (M+H)⁺.

25

Example 109

2-cyclohexyl-N-[(2*Z*)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3*H*)-ylidene]acetamide

The product of Example 64A (39 mg, 0.14 mmol) and cyclohexylacetic acid (24 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.86 - 1.05 (m, 2 H) 1.06 - 1.35 (m, 3 H) 1.54 - 1.75 (m, 5 H) 1.77 - 1.92 (m, 1 H) 2.36 (d, 2 H) 3.19 - 3.25 (m, 3 H) 3.75 (t, 2 H) 4.54 (t, 2 H) 7.25 - 7.36 (m, 1 H) 7.43 - 7.56 (m, 1 H) 7.60

30

- 7.69 (m, 1 H) 7.77 - 7.88 (m, 1 H); MS (ESI) m/z 333 (M+H)⁺.

Example 110

N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]cycloheptanecarboxamide

5

The product of Example 64A (39 mg, 0.14 mmol) and cycloheptylacetic acid (24 mg, 0.17 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 1.42 - 1.62 (m, 6 H) 1.63 - 1.82 (m, 4 H) 1.85 - 2.05 (m, 2 H) 2.54 - 2.72 (m, 1 H) 3.18 - 3.26 (m, 3 H) 3.72 (t, 2 H) 4.55 (t, 2 H) 7.23 - 7.40 (m, 1 H) 7.42 - 7.55 (m, 1 H) 7.60 - 7.72 (m, 1 H) 7.75 - 7.88 (m, 1 H); MS (ESI) m/z 333 (M+H)⁺.

10

Example 111

N-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-5-methylthiophene-2-carboxamide

15

The product of Example 64A (39 mg, 0.14 mmol) and 5-methylthiophene-2-carboxylic acid (28 mg, 0.20 mmol) were processed using the method described in Example 3 to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 2.51 (s, 3 H) 3.26 (s, 3 H) 3.82 (t, 2 H) 4.66 (t, 2 H) 6.91 - 6.93 (m, 1 H) 7.35 (t, 1 H) 7.51 (t, 1 H) 7.68 - 7.72 (m, 2 H) 7.89 (d, 1 H); MS (ESI) m/z 332 (M+H)⁺.

20

Example 112

N-[(2E)-6-fluoro-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide

25

Example 112A

N-(6-fluoro-1,3-benzothiazol-2-yl)-2,2,3,3-tetramethylcyclopropanecarboxamide

30

A mixture of 6-fluoro-benzothiazol-2-ylamine (1 equiv), N-(3-dimethylaminopropyl)-N-ethylcarbodiimide hydrochloride (1 equiv), 1-hydroxybenzotriazole, triethylamine (1.1 equiv), and 2,2,3,3-tetramethylcyclopropanecarboxylic acid (0.8 equiv) in 3:1 THF/Et₃N (1 M) and were stirred overnight at room temperature. The mixture was diluted with EtOAc, washed with 1 M aqueous NaHCO₃, dried (Na₂SO₄), filtered, and concentrated. Purified by

silica gel chromatography afforded the title compound. MS (ESI⁺) m/z 293 (M+H)⁺.

Example 112B

2,2,3,3-Tetramethylcyclopropanecarboxylic acid [6-fluoro-3-(2-methoxyethyl)-3H-benzothiazol-2-ylidene]-amide

To a solution of the product of Example 112A (1 equiv) in 1:1 DMF/THF (0.1 M) was added sodium hydride (60% dispersion in mineral oil, 1.2 equiv) and 2-bromoethyl methyl ether (1.2 equiv). The mixture was stirred at 65 °C overnight then cooled to ambient temperature and diluted with EtOAc. The mixture was washed with 1 M saturated aqueous NaHCO₃, dried (Na₂SO₄), filtered and concentrated. Purified by silica gel chromatography afforded the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ ppm 1.20 (s, 6 H), 1.27 (s, 6 H), 1.60 (s, 1 H), 3.23 (s, 3 H), 3.72 (t, J=5.4 Hz, 2 H), 4.54 (t, J=5.4 Hz, 2 H), 7.33 (td, J=9.0, 2.7 Hz, 1 H), 7.65 (dd, J=8.8, 4.4 Hz, 1 H), 7.75 (dd, J=8.1, 2.7 Hz, 1 H); MS (ESI⁺) m/z 351 (M+H)⁺.

Example 113

2-cyclopentyl-N-[(2Z)-6-fluoro-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]acetamide

Example 113A

6-Fluoro-3-(2-methoxyethyl)-3H-benzothiazol-2-ylideneamine hydrobromide

Commercially available 6-fluoro-benzothiazol-2-ylamine and 2-bromoethyl methyl ether were processed as described for example 12A to afford the title compound. MS (ESI⁺) m/z 227 (M+H)⁺.

Example 113B

2-cyclopentyl-N-[(2Z)-6-fluoro-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]acetamide

The product from Example 113A and cyclopentylacetyl chloride were processed as described for example 11 to afford the title compound. ¹H NMR (400 MHz, DMSO-d₆) δ ppm 4.54 (d, J=5.76 Hz, 2 H) 7.65 (t, J=8.14 Hz, 1 H) 7.75 (dd, J=8.81, 2.71 Hz, 1 H) 7.83 (dd, J=7.97, 1.53 Hz, 2 H) 7.97 (dd, J=8.14, 1.36 Hz, 1 H) 8.41 (d, J=3.05 Hz, 1 H); MS (ESI⁺) m/z 337 (M+H)⁺; Anal. Calculated for C₁₇H₂₁FN₂O₂S: C, 60.69; H, 6.29; N, 8.33. Found: C, 60.67; H, 6.41; N, 8.25.

Example 114

N-[(2Z)-6-fluoro-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-2-tetrahydro-
2H-pyran-4-ylacetamide

5

Example 114A

(Tetrahydro-pyran-4-yl)-acetyl chloride

Commercially available (tetrahydro-pyran-4-yl)-acetic acid and oxalyl chloride were processed as described for example 9A to afford the title compound.

10 MS (DCI/NH₃) m/z 159 (M+H)⁺.

Example 114B

N-[6-Fluoro-3-(2-methoxyethyl)-3H-benzothiazol-2-ylidene]-2-(tetrahydro-pyran-4-
yl)-acetamide

15 The product from Example 113A and the product from Example 114A were processed as described for Example 11 to afford the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ ppm 1.13 - 1.35 (m, 2 H), 1.52 - 1.66 (m, 2 H), 2.05 (d, 1 H), 2.43 (d, J=7.1 Hz, 2 H), 3.22 (s, 3 H), 3.72 (t, J=5.3 Hz, 2 H), 3.78 - 3.88 (m, 4 H), 4.56 (t, J=5.3 Hz, 2 H), 7.36 (td, J=9.0, 2.7 Hz, 1 H), 7.70 (dd, J=9.0, 4.2 Hz, 1 H), 7.81 (dd, J=8.5, 2.7 Hz, 1 H); MS (ESI⁺) m/z 353 (M+H)⁺; Anal. Calculated for C₁₇H₂₁FN₂O₃S: C, 57.94; H, 6.01; N, 7.95. Found: C, 58.03; H, 5.97; N, 7.87.

20

Example 115

5-fluoro-N-[(2Z)-6-fluoro-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-2-
methoxybenzamide

25

The product from Example 113A and 5-fluoro-2-methoxy-benzoic acid were processed using the methods described in Example 13 to afford the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ ppm 3.23 (s, 3 H), 3.72 - 3.87 (m, 5 H), 4.65 (t, J=5.4 Hz, 2 H), 7.15 (dd, J=9.3, 4.2 Hz, 1 H), 7.37 (dd, 2 H), 7.64 (dd, J=9.2, 3.4 Hz, 1 H), 7.77 (dd, J=9.2, 4.4 Hz, 1 H), 7.88 (dd, J=8.1, 2.7 Hz, 1 H); MS (ESI⁺) m/z 379 (M+H)⁺; Anal. Calculated for C₁₈H₁₆F₂N₂O₃S: C, 57.13; H, 4.26; N, 7.40. Found: C, 57.05; H, 4.08; N, 7.35.

30

Example 1165-chloro-N-[(2Z)-6-fluoro-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]-2-methoxybenzamide

The product from Example 113A and 5-chloro-2-methoxybenzoic acid were
 5 processed using the methods described in Example 13 to afford the title compound.
¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 3.24 (s, 3 H), 3.79 (t, *J*=5.3 Hz, 2 H), 3.83 (s,
 3 H), 4.65 (t, *J*=5.3 Hz, 2 H), 7.17 (d, *J*=9.2 Hz, 1 H), 7.41 (td, *J*=9.1, 2.5 Hz, 1 H),
 7.52 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.78 (dd, *J*=9.0, 4.2 Hz, 1 H), 7.83 (d, *J*=2.7 Hz, 1 H),
 7.89 (dd, *J*=8.5, 2.7 Hz, 1 H); MS (ESI⁺) *m/z* 395 (M+H)⁺; Anal. Calculated for
 10 C₁₈H₁₆ClFN₂O₃S: C, 54.75; H, 4.08; N, 7.09. Found: C, 54.29; H, 3.94; N, 6.99.

Example 117N-[(2Z)-3-(2-methoxyethyl)-4-methyl-5-morpholin-4-yl-1,3-thiazol-2(3H)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide

15

Example 117A2,2,3,3-Tetramethylcyclopropanecarboxylic acid (4-methyl-5-morpholin-4-yl-thiazol-2-yl)-amide

20 A mixture of 4-methyl-5-morpholin-4-yl-thiazol-2-ylamine (prepared using
 the method described in Christopher et al., *Bioorganic and Medicinal Chemistry
 Letters* 2004, 14(22), 5521-5525), N-(3-dimethylaminopropyl)-N-ethylcarbodiimide
 hydrochloride, 1-hydroxybenzotriazole, triethylamine, and 2,2,3,3-
 tetramethylcyclopropanecarboxylic acid were processed using the method described
 25 in Example 58 to afford the title compound. MS (ESI⁺) *m/z* 324 (M+H)⁺.

Example 117B2,2,3,3-Tetramethylcyclopropanecarboxylic acid [3-(2-methoxyethyl)-4-methyl-5-morpholin-4-yl-3H-thiazol-2-ylidene]-amide

30 The product from Example 117A was processed using the method described in
 Example 112B to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm
 1.16 (s, 6 H), 1.23 (s, 6 H), 1.44 (s, 1 H), 2.22 (s, 3 H), 2.70 - 2.79 (m, 4 H), 3.24 (s, 3
 H), 3.62 (t, *J*=5.3 Hz, 2 H), 3.66 - 3.72 (m, 4 H), 4.21 (t, *J*=5.4 Hz, 2 H); MS (ESI⁺)

m/z 382 (M+H)⁺.

Example 118

N-[(2Z)-5-chloro-3-(2-methoxyethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2,2,3,3-
tetramethylcyclopropanecarboxamide

Example 118A

2,2,3,3-Tetramethylcyclopropanecarboxylic acid (5-chloro-4-methyl-thiazol-2-yl)-
amide

10 A mixture of 2-amino-5-chloro-2-methyl-thiazole (Matsuo, Masaaki; Ogino, Takashi; Igari, Norihiro; Seno, Hachiro; Shimomura, Kyoichi., EP 412404) (150 mg, 0.81 mmol), 2,2,3,3-tetramethylcyclopropanecarbonyl chloride (143 mg, 0.89 mmol), 4-dimethylaminopyridine (50.0 mg, 0.41 mmol) and triethylamine (226 μ L, 1.62 mmol) in 15 mL of THF heated at reflux for 48 hours. The mixture was cooled to
 15 ambient temperature, diluted with EtOAc and washed with brine. The layers were separated and the aqueous phase was extracted with EtOAc (2X). The combined organic extracts were dried (Na₂SO₄), filtered, and concentrated. Purification by silica gel chromatography afforded the title compound: MS (LC/MS) m/z 273 (M+H)⁺.

20

Example 118B

2,2,3,3-Tetramethylcyclopropanecarboxylic acid (5-chloro-3- (2-methoxy-ethy)- 4-
methyl-3H-thiazol-2-ylidene]-amide

A mixture of the product of Example 118A (150 mg, 0.55 mmol), NaH (60%
 25 dispersion in mineral oil, 290 mg, 0.71 mmol) and 2-bromoethyl methyl ether (57 μ L, 0.61 mmol) in 20 mL of 2:1 THF/DMF was processed according to the method described in Example 112B to provide the title compound: ¹H NMR (300 MHz, DMSO-*d*₆) δ 1.18 (s, 6 H) 1.23 (s, 6 H) 1.50 (s, 1 H) 2.29 (s, 3 H) 3.25 (s, 3 H) 3.63 (t, *J*=5.3 Hz, 2 H) 4.27 (t, *J*=5.3 Hz, 2 H); MS (DCI/NH₃) m/z 331 (M+H)⁺.

30

Example 119

N-[(2Z)-3-(2-methoxyethyl)-5-methyl-4-phenyl-1,3-thiazol-2(3H)-ylidene]-2,2,3,3-
tetramethylcyclopropanecarboxamide

Example 119A3-(2-methoxyethyl)-5-methyl-4-phenyl-3H-thiazol-2-ylideneamine hydrobromide

A mixture of 5-methyl-4-phenyl-thiazol-2-ylamine (300 mg, 1.58 mmol) and
5 2-bromoethyl methyl ether (300 μ L, 3.20 mmol) was processed using the method
described in Example 12A to provide the title compound.

Example 119B

10 2,2,3,3-Tetramethylcyclopropanecarboxylic acid [3-(2-methoxy-ethyl)-5-methyl-4-phenyl-3H-thiazol-2-ylidene]-amide

A mixture of the product of Example 119A (290 mg, 1.16 mmol), 2,2,3,3-tetramethylcyclopropanecarboxylic acid (182 mg, 1.28 mmol), HATU (661 mg, 1.74 mmol) and triethylamine (0.97 mL, 6.96 mmol) in 15 mL of DMF was processed according to the method of Example 2B to provide the title compound: ^1H NMR (400
15 MHz, DMSO- d_6) δ 1.17 (s, 6 H) 1.26 (s, 6 H) 1.49 (s, 1 H) 2.01 (s, 3 H) 3.02 (s, 3 H) 3.43 (t, $J=6.0$ Hz, 2 H) 4.06 (t, $J=5.8$ Hz, 2 H) 7.39 - 7.43 (m, 2 H) 7.50 - 7.57 (m, 3 H); MS (DCI/ NH_3) m/z 373 ($\text{M}+\text{H}$) $^+$.

Example 120

20 N-[(2Z)-4-(4-chlorophenyl)-3-(2-methoxyethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide

Example 120A

25 4-(4-chlorophenyl)-3-(2-methoxyethyl)-5-methyl-3H-thiazol-2-ylideneamine hydrobromide

A mixture of 4-(4-chlorophenyl)-5-methyl-thiazol-2-ylamine (420 mg, 1.87 mmol) and 2-bromoethyl methyl ether (600 μ L, 6.40 mmol) was processed according to the method of Example 12A to provide the title compound: MS (DCI/ NH_3) m/z 283
30 ($\text{M}+\text{H}$) $^+$.

Example 120B2,2,3,3-Tetramethylcyclopropanecarboxylic acid [4-(4-chlorophenyl)-3-(2-methoxy-

ethy)-5-methyl-3H-thiazol-2-ylidene]-amide

A mixture of the product of Example 120A (156 mg, 0.55 mmol), 2,2,3,3-tetramethylcyclopropanecarboxylic acid (94 mg, 0.66 mmol), HATU (479 mg, 0.83 mmol) and triethylamine (0.46 mL, 3.30 mmol) in 10 mL of DMF was processed according to the method of Example 2B to provide the title compound: ¹H NMR (500 MHz, DMSO-*d*₆) δ 1.17 (s, 6 H) 1.25 (s, 6 H) 1.49 (s, 1 H) 2.00 (s, 3 H) 3.04 (s, 3 H) 3.44 (t, *J*=5.8 Hz, 2 H) 4.04 (t, *J*=5.8 Hz, 2 H) 7.45 (d, *J*=8.2 Hz, 2 H) 7.60 (d, *J*=8.5 Hz, 2 H); MS (DCI/NH₃) *m/z* 407 (M+H)⁺.

10

Example 121N-[(2Z)-3-(2-methoxyethyl)-4,5,6,7-tetrahydro-1,3-benzothiazol-2(3H)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamideExample 121A

15

3-(2-methoxy-ethy)-4,5,6,7-tetrahydro-3H-benzothiazol-2-ylideneamine hydrobromide

20

A mixture of 4,5,6,7-tetrahydro-benzothiazol-2-ylamine (300 mg, 1.94 mmol) and 2-bromoethyl methyl ether (600 μL, 6.40 mmol) was processed according to the method of Example 12A to provide the title compound as crude product: MS (LC/MS) *m/z* 213 (M+H)⁺.

Example 121B2,2,3,3-Tetramethylcyclopropanecarboxylic acid [3-(2-methoxy-ethy)-4,5,6,7-tetrahydro-3H-benzothiazol-2-ylidene]-amide

25

A mixture of the product of Example 121A (156 mg, 0.55 mmol), 2,2,3,3-tetramethylcyclopropanecarboxylic acid (94 mg, 0.66 mmol), HATU (479 mg, 0.83 mmol) and triethylamine (0.46 mL, 3.30 mmol) in 10 mL of DMF was processed according to the method of Example 2B to provide the title compound: ¹H NMR (500 MHz, DMSO-*d*₆) δ 1.16 (s, 6 H) 1.23 (s, 6 H) 1.45 (s, 1 H) 1.69 - 1.82 (m, 4 H) 2.43 - 2.48 (m, 2 H) 2.54 - 2.59 (m, 2 H) 3.24 (s, 3 H) 3.60 (t, *J*=5.3 Hz, 2 H) 4.16 (t, *J*=5.3 Hz, 2 H); MS (DCI/NH₃) *m/z* 337 (M+H)⁺.

30

Example 122

N-[(2Z)-3-(2-methoxyethyl)-3,4,5,6-tetrahydro-2H-cyclopenta[d][1,3]thiazol-2-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide

Example 122A

5 2,2,3,3-Tetramethylcyclopropanecarboxylic acid (5,6-dihydro-4H-cyclopentathiazol-2-yl)-amide

A mixture of 5,6-dihydro-4H-cyclopentathiazole-2-ylamine (177 mg, 1.26 mmol), 2,2,3,3-tetramethylcyclopropanecarboxylic acid (244 mg, 1.52 mmol), 4-dimethylaminopyridine (50.0 mg, 0.41 mmol) and triethylamine (351 μ L, 2.52 mmol)
 10 in 20 mL of THF was processed according to the method of Example 118A to provide the title compound: MS (DCI/NH₃) m/z 265 (M+H)⁺.

Example 122B

15 2,2,3,3-Tetramethylcyclopropanecarboxylic acid [3-(2-methoxy-ethyl)- (3,4,5,6-tetrahydro-cyclopentathiazol-2-ylidene)-amide

A mixture of the product of Example 122A (254 mg, 0.95 mmol), NaH (60% dispersion in mineral oil, 50.0 mg, 1.22 mmol) and 2-bromoethyl methyl ether (100 μ L, 1.07 mmol) in 30 mL of THF/DMF (2/1) was processed according to the method of Example 112B to provide the title compound: ¹H NMR (500 MHz, DMSO-*d*₆)
 20 δ 1.16 (s, 6 H) 1.23 (s, 6 H) 1.45 (s, 1 H) 2.31 - 2.38 (m, 2 H) 2.72 (t, *J*=7.0 Hz, 2 H) 2.78 (t, *J*=7.2 Hz, 2 H) 3.24 (s, 3 H) 3.61 (t, *J*=5.3 Hz, 2 H) 4.15 (t, *J*=5.2 Hz, 2 H); MS (DCI/NH₃) m/z 323 (M+H)⁺.

Example 123

25 N-[(7Z)-8-(2-methoxyethyl)-5,8-dihydro[1,3]thiazolo[4,5-e][2,1,3]benzoxadiazol-7(4H)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide

Example 123A

4,5-dihydro[1,3]thiazolo[4,5-e][2,1,3]benzoxadiazol-7-amine hydrobromide
 30 To a solution of commercially available 5-bromo-6,7-dihydro-5H-benzo[1,2,5]oxadiazol-4-one (1.1 g, 5.1 mmol) in absolute ethanol (60 mL) was added thiourea. The reaction mixture was stirred at 60 °C for overnight and then concentrated. The residue was triturated in hexanes to afford 1.3 g (90%) of the title

compound. MS (ESI⁺) m/z 195 (M+H)⁺.

Example 123B

N-4,5-dihydro[1,3]thiazolo[4,5-*e*][2,1,3]benzoxadiazol-7-yl-2,2,3,3-

5 tetramethylcyclopropanecarboxamide

Example 123A and 2,2,3,3-tetramethylcyclopropanecarbonyl chloride were processed as described for example 118A to afford the title compound. MS (ESI⁺) m/z 319 (M+H)⁺.

Example 123C

N-[(7Z)-8-(2-methoxyethyl)-5,8-dihydro[1,3]thiazolo[4,5-*e*][2,1,3]benzoxadiazol-
10 7(4*H*)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide

The product of Example 123B (1 equiv), potassium tert-butoxide (1.1 equiv) and 2-bromoethyl methyl ether (1 equiv) were combined in DMF (0.1 M) and heated
15 in a SmithSynthesizerTM microwave at 250 °C for 15 minutes. The mixture was diluted with EtOAc and washed with 1 M aqueous NaHCO₃. The phases were separated and the aqueous phase was extracted with EtOAc (3X). The combined organic extracts were dried over Na₂SO₄, filtered, and concentrated. Purification by silica gel chromatography afforded the title compound and the product of Example
20 124. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.20 (s, 6 H), 1.26 (s, 6 H), 1.57 (s, 1 H), 3.10 (t, *J*=7.5 Hz, 2 H), 3.23 (s, 3 H), 3.26 (t, *J*=7.4 Hz, 2 H), 3.71 (t, *J*=5.8 Hz, 2 H), 4.67 (t, *J*=5.8 Hz, 2 H); MS (ESI⁺) m/z 377 (M+H)⁺.

Example 124

25 N-[(7Z)-8-(2-methoxyethyl)[1,3]thiazolo[4,5-*e*][2,1,3]benzoxadiazol-7(8*H*)-ylidene]-
2,2,3,3-tetramethylcyclopropanecarboxamide

The title compound was obtained as byproduct during the synthesis of example 123C. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 1.23 (s, 6 H), 1.29 (s, 6 H), 1.67 (s, 1 H), 3.24 (s, 3 H), 3.85 (t, *J*=5.7 Hz, 2 H), 4.96 (t, *J*=5.5 Hz, 2 H), 7.94 (d, *J*=9.2 Hz, 1 H), 8.06 (d, *J*=9.5 Hz, 1 H); MS (ESI⁺) m/z 374 (M+H)⁺.
30

Example 125

2-Ethoxy-N-[(2Z)-3-(2-methoxyethyl)-4,6-dihydrofuro[3,4-d]thiazol-2(3H)-ylidene]-benzamide

Example 125A

5 4-Bromo-dihydrofuran-3-one

The title compound was prepared according to the procedure using the method described in Baker, Tracy J, Wiemer, David F, *J. Org. Chem.*, 1998, 63(8), 2613-2618 and was used immediately using the method described in Example 125B.

10 Example 125B

3a-Ethoxy-3a,4,6,6a-tetrahydrofuro[3,4-d]thiazol-2-ylamine

The product of Example 125A and thiourea were processed using the method described in Example 123A to afford the title compound. MS (ESI⁺) m/z 189 (M+H)⁺.

15 Example 125C

3a-Ethoxy-3-(2-methoxy-ethyl)-tetrahydro-furo[3,4-d]thiazol-2-ylideneamine

The product from Example 125B and 1-bromo-2-methoxyethane were processed using the method described in Example 12A to afford the title compound. MS (ESI⁺) m/z 247 (M+H)⁺.

20

Example 125D

2-Ethoxy-N-[3a-ethoxy-3-(2-methoxy-ethyl)-tetrahydro-furo[3,4-d]thiazol-2-ylidene]-benzamide

25 The product from Example 125B and 2-ethoxybenzoyl chloride were processed as described for example 118A to afford the title compound. MS (ESI⁺) m/z 395 (M+H)⁺.

Example 125E

30 2-Ethoxy-N-[3-(2-methoxyethyl)-4,6-dihydro-3H-furo[3,4-d]thiazol-2-ylidene]-benzamide

To a solution of the product from Example 125D (15 mg, 0.04 mmol) in toluene (10 mL) was added *p*-toluenesulfonic acid monohydrate (2 mg). The mixture was refluxed for 3 hours and then cooled to room temperature, diluted with ethyl acetate, washed with 1M NaHCO₃, dried (Na₂SO₄), filtered and concentrated.

Purification by preparative HPLC on a waters Symmetry C8 column (40 mm X 100 mm, 7 μ m particle size) using a gradient of 10 % to 100 % acetonitrile: ammonium acetate (10 mM) over 15 min at a flow rate of 70 mL/min afforded the title compound. ^1H NMR (300 MHz, DMSO- d_6) δ ppm 1.33 (t, $J=7.0$ Hz, 3 H), 3.24 (s, 3 H), 3.65 (t, $J=4.9$ Hz, 2 H), 4.07 (q, $J=6.8$ Hz, 2 H), 4.25 (t, $J=5.1$ Hz, 2 H), 4.96 (s, 4 H), 6.92 - 7.01 (m, 1 H), 7.07 (d, $J=8.1$ Hz, 1 H), 7.36 - 7.44 (m, 1 H), 7.74 (dd, $J=7.5, 1.7$ Hz, 1 H); MS (ESI $^+$) m/z 349 (M+H) $^+$.

Example 126

10 3-chloro-2-fluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-6-(trifluoromethyl)benzamide

Example 126A

3-(2-Methoxy-ethyl)-4,5-dimethyl-3H-thiazol-2-ylideneamine

15 The product of Example 12A was purified via flash column chromatography (SiO $_2$, 9:1:0.1 CH $_2$ Cl $_2$: CH $_3$ OH : NH $_4$ OH) to provide the title compound. ^1H NMR (300 MHz, CDCl $_3$) δ ppm 2.18 (s, 6 H) 3.32 (s, 3 H) 3.78 (t, $J=5.10$ Hz, 2 H) 4.39 (t, $J=4.70$ Hz, 2 H) 9.45 (s, 2 H); MS (DCI/NH $_3$) m/z 187 (M+H) $^+$.

Example 126B

20 3-Chloro-2-fluoro-N-[3-(2-methoxy-ethyl)-4,5-dimethyl-3H-thiazol-2-ylidene]-6-trifluoromethyl-benzamide

To a suspension of the product of Example 126A (0.20 g, 1.1 mmol) in 35 mL THF was added Et $_3$ N (0.37 mL, 2.7 mmol). This mixture was cooled to 0 $^\circ$ C and 3-chloro-2-fluoro-6-trifluoromethyl benzoyl chloride (Alfa Aesar, 0.35 g, 1.3 mmol) in 25 5 mL THF was added dropwise via syringe. The mixture was allowed to stir at ambient temperature for 1 hour, then was warmed to reflux and was allowed to stir for 8 hours. The mixture was then cooled to ambient temperature and filtered. The filtrate was concentrated under reduced pressure and purified via flash column chromatography (SiO $_2$, 7:3 hexanes:EtOAc) to provide the title compound (0.20 g, 30 0.50 mmol, 46% yield). ^1H NMR (300 MHz, CDCl $_3$) δ ppm 2.27 (s, 3 H), 2.28 (s, 3 H), 3.27 (s, 3 H), 3.68 (t, $J=4.7$ Hz, 2 H), 4.28-4.37 (m, 2 H), 7.37-7.43 (m, 1 H), 7.44-7.52 (m, 1 H); MS (DCI/NH $_3$) m/z 411 (M+H) $^+$; Anal. calculated for C $_{16}$ H $_{15}$ ClF $_4$ N $_2$ O $_2$ S: C, 46.78; H, 3.68; N, 6.82. Found: C, 46.83; H, 3.30; N, 6.65.

Example 1275-chloro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-(trifluoromethyl)benzamide

- 5 To the product of Example 126A (0.20 g, 1.1 mmol) in 35 mL THF at 0 °C was added Et₃N (0.37 mL, 2.7 mmol) followed by 5-chloro-2-trifluoromethyl-benzoyl chloride (Matrix, 0.26 g, 1.3 mmol) in 5 mL THF dropwise via syringe. This mixture was stirred at ambient temperature for 1 hour then was warmed to reflux and allowed to stir for 8 hours. The mixture was then cooled to ambient temperature and filtered.
- 10 The filtrate was concentrated under reduced pressure and the residue was purified via flash column chromatography (SiO₂, 4:1 hexanes:EtOAc) to provide the title compound (0.23 g, 0.57 mmol, 53% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 2.26 (s, 3 H), 2.28 (s, 3 H), 3.30 (s, 3 H), 3.71 (t, *J*=4.9 Hz, 2 H), 4.36 (t, *J*=4.9 Hz, 2 H), 7.45 (ddd, *J*=8.5, 2.0, 0.7 Hz, 1 H), 7.64 (d, *J*=8.5 Hz, 1 H), 7.84 (d, *J*=2.0 Hz, 1 H);
- 15 MS (DCI/NH₃) *m/z* 393 (M+H)⁺; Anal. calculated for C₁₆H₁₆ClF₃N₂O₂S: C, 48.92; H, 4.11; N, 7.13. Found: C, 48.66; H, 3.81; N, 7.01.

Example 1282,3-dichloro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]benzamide

- 20 To the product of Example 126A (0.20 g, 1.1 mmol) in 35 mL THF was added Et₃N (0.37 mL, 2.7 mmol). This mixture was cooled to 0 °C and 2,3-dichlorobenzoyl chloride (Lancaster, 0.27 g, 1.3 mmol) in 5 mL THF was added dropwise via syringe. This mixture stirred at ambient temperature for 1 hour then was warmed to reflux and
- 25 was allowed to stir for 3 hours. The mixture was cooled to ambient temperature and was quenched with 5 mL saturated, aqueous NH₄Cl and diluted with 10 mL EtOAc. The layers were separated and the aqueous layer was extracted 2 X 5 mL EtOAc and 2 X 5 mL CH₂Cl₂. The combined organics were washed 1 X 5 mL saturated, aqueous NaCl then were dried over anhydrous Na₂SO₄, filtered, concentrated under reduced
- 30 pressure and purified via flash column chromatography (SiO₂, 10% CH₃OH:EtOAc). The material was still impure so it was purified again via flash column chromatography (SiO₂, 1:1 hexanes:EtOAc) to provide the title compound (0.105 g, 0.29 mmol, 27% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 2.26 (s, 3 H), 2.28 (s, 3

H), 3.30 (s, 3 H), 3.76 (t, $J = 4.6$ Hz, 2 H), 4.40 (t, $J = 4.2$ Hz, 2 H), 7.23 (t, $J = 7.7$ Hz, 1 H), 7.49 (dd, $J = 8.0, 1.5$ Hz, 1 H), 7.72 (dd, $J = 7.6, 1.5$ Hz, 1 H); MS (DCI/ NH_3) m/z 359 ($M+H$)⁺; Anal. calculated for $\text{C}_{15}\text{H}_{16}\text{Cl}_2\text{N}_2\text{O}_2\text{S}$: C, 50.15; H, 4.49; N, 7.80. Found: C, 50.17; H, 4.26; N, 7.69.

5

Example 129

N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2,2-dimethyl-2,3-dihydro-1-benzofuran-7-carboxamide

To a solution of the product of Example 126A (0.20 g, 1.1 mmol) and Et_3N (0.45 mL, 3.2 mmol) in 30 mL THF was added 2,2-dimethyl-2,3-dihydro-1-benzofuran-7-carbonyl chloride (Acros, 0.34 g, 1.6 mmol). This mixture was warmed to reflux and allowed to stir for 2 hours. The mixture was then cooled to ambient temperature and concentrated under reduced pressure. The residue was diluted with 10 mL EtOAc and washed with 5 mL saturated, aqueous NH_4Cl . The layers were separated and the aqueous layer was extracted 2 x 5 mL EtOAc. The combined organics were dried over anhydrous Na_2SO_4 , filtered, concentrated under reduced pressure and purified via flash column chromatography (SiO_2 , 4:1 hexanes:EtOAc) to provide the title compound (0.21 mmol, 0.57 mmol, 53% yield). ^1H NMR (300 MHz, CDCl_3) δ ppm 1.55 (s, 6 H), 2.22 (s, 3 H), 2.27 (s, 3 H), 3.02 (s, 2 H), 3.31 (s, 3 H), 3.77-3.89 (m, 2 H), 4.33-4.50 (m, 2 H), 6.85 (t, $J = 7.5$ Hz, 1 H), 7.22 (d, $J = 7.5$ Hz, 1 H), 8.00 (d, $J = 8.1$ Hz, 1 H); MS (DCI/ NH_3) m/z 361 ($M+H$)⁺; Anal. calculated for $\text{C}_{19}\text{H}_{24}\text{N}_2\text{O}_3\text{S}$: C, 63.31; H, 6.71; N, 7.77. Found: C, 63.19; H, 6.50; N, 7.66.

20

Example 130

2,2-difluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-1,3-benzodioxole-4-carboxamide

To the product of Example 126A (0.20 g, 1.1 mmol) in 30 mL THF was added Et_3N (0.37 mL, 2.7 mmol) followed by 2,2-difluoro-1,3-benzodioxole-4-carbonyl chloride (Lancaster, 0.29 g, 1.3 mmol). This mixture was stirred at ambient temperature for 17 hours then was warmed to reflux and allowed to stir for an additional 4 hours. The mixture was then cooled to ambient temperature and additional 2,2-difluoro-1,3-benzodioxole-4-carbonyl chloride (73 mg, 0.33 mmol) and Et_3N (0.37 mL, 2.7 mmol) were added. This mixture was warmed to reflux at which

30

temperature it stirred for 2 hours. The mixture was then cooled to ambient temperature, diluted with 10 mL EtOAc and washed with 5 mL saturated, aqueous NH_4Cl . The layers were separated and the aqueous layer was extracted 2 x 5 mL EtOAc. The combined organics were dried over anhydrous Na_2SO_4 , filtered, concentrated under reduced pressure and purified via flash column chromatography (50% hexanes:EtOAc) to afford the title compound (0.22 g, 0.59 mmol, 55% yield). ^1H NMR (300 MHz, CDCl_3) δ ppm 2.26 (s, 3 H), 2.28 (s, 3 H), 3.32 (s, 3 H), 3.83 (t, $J=5.1$ Hz, 2 H), 4.44 (t, $J=5.1$ Hz, 2 H), 7.12 (d, $J=4.4$ Hz, 1 H), 7.14 (s, 1 H), 7.89 (dd, $J=6.6, 2.9$ Hz, 1 H); MS (DCI/ NH_3) m/z 371 ($\text{M}+\text{H}$) $^+$; Anal. calculated for $\text{C}_{16}\text{H}_{16}\text{F}_2\text{N}_2\text{O}_4\text{S}$: C, 51.89; H, 4.35; N, 7.56. Found: C, 52.27; H, 4.24; N, 7.53.

Example 131

5-bromo-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2,3-dihydro-1-benzofuran-7-carboxamide

The product of Example 126A (0.20 g, 1.1 mmol), Et_3N (0.45 mL, 3.2 mmol) and 5-bromo-2,3-dihydrobenzo[b]-furan-7-carbonyl chloride (Maybridge, 0.42 g, 1.6 mmol) in 35 mL THF were processed as in Example 126B. The resulting crude material was purified via flash column chromatography (SiO_2 , 7:3 hexanes:EtOAc) to afford the title compound (0.16 g, 0.39 mmol, 36% yield). ^1H NMR (300 MHz, CDCl_3) δ ppm 2.24 (s, 3 H), 2.28 (s, 3 H), 3.23 (t, $J=8.5$ Hz, 2 H), 3.33 (s, 3 H), 3.78 - 3.87 (m, 2 H), 4.45 - 4.56 (m, 2 H), 4.70 - 4.81 (m, 2 H), 7.35 - 7.39 (m, 1 H), 8.09 (d, $J=2.0$ Hz, 1 H); MS (DCI/ NH_3) m/z 411, 413 ($\text{M}+\text{H}$) $^+$; Anal. calculated for $\text{C}_{17}\text{H}_{19}\text{BrN}_2\text{O}_3\text{S}\cdot 0.3\text{H}_2\text{O}$: C, 49.00; H, 4.74; N, 6.72. Found: C, 48.91; H, 4.36; N, 6.57.

Example 132

2-bromo-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]benzamide

To a solution of the product of Example 126A (0.20 g, 1.1 mmol) and Et_3N (0.3 mL, 2.2 mmol) in 25 mL THF was added 2-bromobenzoyl chloride (Aldrich, 0.18 mL, 1.4 mmol). This mixture stirred at ambient temperature for 20 hours then was concentrated under reduced pressure and the residue was diluted with 10 mL EtOAc and washed with 5 mL NH_4Cl . The layers were separated and the aqueous

layer was extracted (2 X 5 mL EtOAc). The combined organic layers were dried over anhydrous Na₂SO₄, filtered, concentrated under reduced pressure and purified via recrystallization with 50% hexanes/EtOAc to afford 0.18 g of the title compound (0.49 mmol, 46% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 2.26 (s, 3 H), 2.28 (s, 3 H), 3.30 (s, 3 H), 3.78 (t, *J*=5.1 Hz, 2 H), 4.42 (t, *J*=5.1 Hz, 2 H), 7.22 (dt, *J*=7.8, 1.7 Hz, 1 H), 7.34 (dt, *J*=7.5, 1.4 Hz, 1 H), 7.63 (dd, *J*=8.0, 1.2 Hz, 1 H), 7.90 (dd, *J*=7.6, 1.9 Hz, 1 H); MS (DCI/NH₃) *m/z* 369, 371 (M+H)⁺; Anal. calculated for C₁₅H₁₇BrN₂O₂S: C, 48.79; H, 4.64; N, 7.59. Found: C, 48.84; H, 4.49; N, 7.40.

10

Example 1332,6-dichloro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]benzamide

15

The product of Example 126A (0.20 g, 1.1 mmol), Et₃N (0.3 mL, 2.2 mmol) and 2,6-dichlorobenzoyl chloride (Aldrich, 0.2 mL, 1.4 mmol) in 25 mL THF were processed as in Example 132 to afford the title compound (0.12 g, 0.32 mmol, 30% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 2.28 (s, 6 H), 3.27 (s, 3 H), 3.74 (t, *J*=4.6 Hz, 2 H), 4.34-4.44 (m, 2 H), 7.19 (dd, *J*=8.8, 6.8 Hz, 1 H), 7.28-7.34 (m, 2 H); MS (DCI/NH₃) *m/z* 359 (M+H)⁺.

20

Example 134N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]quinoline-4-carboxamideExample 134AQuinoline-4-carbonyl chloride

25

A solution of 4-quinolinecarboxylic acid (Aldrich, 0.25 g, 1.4 mmol) in 5 mL of thionyl chloride was warmed to reflux and allowed to stir for 1 hour. The mixture was then cooled to ambient temperature and concentrated under reduced pressure. This material was dissolved in 10 mL toluene and concentrated under reduced pressure (3X) to afford the title compound.

30

Example 134BN-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]quinoline-4-carboxamide

To a suspension of the product of Example 126A (0.20 g, 1.1 mmol) in 25 mL

THF was added Et₃N (0.3 mL, 2.2 mmol) followed by the freshly prepared quinoline-4-carbonyl chloride. This mixture was warmed to reflux and allowed to stir for 18 hours. The material was then concentrated under reduced pressure and 10 mL EtOAc, 5 mL H₂O and 5 mL NH₄OH were added. The layers were separated and the aqueous layer was extracted 2 X 5 mL EtOAc. The combined organic layers were dried over anhydrous Na₂SO₄, filtered, concentrated under reduced pressure and purified via flash column chromatography (SiO₂, 5% CH₃OH in EtOAc) to afford the title compound (0.12 g, 0.35 mmol, 33% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 2.32 (d, *J*=0.7 Hz, 3 H), 2.35 (d, *J*=0.7 Hz, 3 H), 3.31 (s, 3 H), 3.78 (t, *J*=5.3 Hz, 2 H), 4.48 (t, *J*=5.1 Hz, 2 H), 7.65 (ddd, *J*=8.5, 7.0, 1.2 Hz, 1 H), 7.80 (ddd, *J*=8.4, 6.9, 1.4 Hz, 1 H), 8.00 (d, *J*=4.4 Hz, 1 H), 8.06-8.11 (m, 1 H), 8.84 (ddd, *J*=8.5, 1.4, 0.7 Hz, 1 H), 8.93 (d, *J*=4.4 Hz, 1 H); MS (DCI/NH₃) *m/z* 342 (M+H)⁺; Anal. calculated for C₁₈H₁₉N₃O₂S: C, 63.32; H, 5.61; N, 12.31. Found: C, 63.23; H, 5.46; N, 12.10.

15 Example 135

N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]quinoline-5-carboxamide

The quinoline-5-carboxylic acid (Lancaster, 0.25 g, 1.4 mmol) was converted to quinoline-5-carbonyl chloride using 5 mL SOCl₂ using the method described in Example 134A. The product of Example 126A (0.20 g, 1.1 mmol), Et₃N (0.3 mL, 2.2 mmol) and quinoline-5-carbonyl chloride in 30 mL THF were processed as in Example 134B to afford the title compound (0.18 g, 0.53 mmol, 49% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 2.30 (d, *J*=0.7 Hz, 3 H), 2.34 (s, 3 H), 3.31 (s, 3 H), 3.80 (t, *J*=5.1 Hz, 2 H), 4.49 (t, *J*=5.3 Hz, 2 H), 7.60 (dd, *J*=8.6, 4.2 Hz, 1 H), 7.83 (dd, *J*=8.5, 7.1 Hz, 1 H), 8.15 (dt, *J*=8.5, 1.0 Hz, 1 H), 8.40 (dd, *J*=7.3, 1.2 Hz, 1 H), 8.87 (dd, *J*=4.1, 1.7 Hz, 1 H), 9.55 (ddd, *J*=8.8, 1.7, 0.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 342 (M+H)⁺; Anal. calculated for C₁₈H₁₉N₃O₂S: C, 63.32; H, 5.61; N, 12.31. Found: C, 63.44; H, 5.05; N, 12.10.

30 Example 136

N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]isoquinoline-5-carboxamide

The isoquinoline-5-carboxylic acid (Lancaster, 0.25 g, 1.4 mmol) was

converted to the corresponding acid chloride with 5 mL of thionyl chloride using the method described in Example 134A. The product of Example 126A (0.20 g, 1.1 mmol), Et₃N (0.3 mL, 2.2 mmol) and the acid chloride in 30 mL THF were processed as in Example 134B to afford the title compound (98 mg, 0.29 mmol, 27% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 2.31 (s, 3 H), 2.35 (s, 3 H), 3.31 (s, 3 H), 3.81 (t, *J*=5.3 Hz, 2 H), 4.50 (t, *J*=5.3 Hz, 2 H), 7.76 (dd, *J*=7.5, 7.5 Hz, 1 H), 8.23 (d, *J*=8.1 Hz, 1 H), 8.48 (d, *J*=6.4 Hz, 1 H), 8.61 (dd, *J*=7.1, 1.4 Hz, 1 H), 9.03 (d, *J*=6.1 Hz, 1 H), 9.27 (d, *J*=1.0 Hz, 1 H); MS (DCI/NH₃) *m/z* 342 (M+H)⁺; Anal. calculated for C₁₈H₁₉N₃O₂S: C, 63.32; H, 5.61; N, 12.31. Found: C, 62.97; H, 5.54; N, 12.07.

10

Example 137

N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2,3-dihydro-1-benzofuran-7-carboxamide

The 2,3-dihydrobenzofuran-7-carboxylic acid (TCI-US, 0.25 g, 1.4 mmol) was converted to 2,3-dihydrobenzofuran-7-carbonyl chloride with 5 mL of thionyl chloride using the method described in Example 134A. The product of Example 126A (0.20 g, 1.1 mmol), Et₃N (0.3 mL, 2.2 mmol) and the freshly prepared 2,3-dihydrobenzofuran-7-carbonyl chloride in 30 mL THF were processed as in Example 134B. Purification of the crude material via recrystallization with 50% hexanes:EtOAc gave the title compound (0.12 g, 0.36 mmol, 34% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 2.25 (s, 3 H), 2.30 (s, 3 H), 3.22 (t, *J*=8.8 Hz, 2 H), 3.31 (s, 3 H), 3.81 (t, *J*=5.3 Hz, 2 H), 4.44 (t, *J*=5.3 Hz, 2 H), 4.64 (t, *J*=8.8 Hz, 2 H), 6.86 (t, *J*=7.5 Hz, 1 H), 7.32 (dd, *J*=7.3, 1.2 Hz, 1 H), 7.93 (dd, *J*=8.0, 0.8 Hz, 1 H); MS (DCI/NH₃) *m/z* 333 (M+H)⁺; Anal. calculated for C₁₇H₂₀N₂O₃S•0.1H₂O: C, 61.09; H, 6.09; N, 8.38. Found: C, 60.99; H, 5.91; N, 8.25.

25

Example 138

2-chloro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]quinoline-4-carboxamide

The 2-chloroquinoline-4-carboxylic acid (TCI-JP, 0.29 g, 1.4 mmol) was converted to 2-chloroquinoline-4-carbonyl chloride with 5 mL of thionyl chloride using the method described in Example 134A. The product of Example 126A (0.20 g, 1.1 mmol), Et₃N (0.3 mL, 2.2 mmol) and 2-chloroquinoline-4-carbonyl chloride in 30

30

mL THF were processed as in Example 134B to afford the title compound (0.14 g, 0.37 mmol, 35% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 2.33 (s, 3 H), 2.35 (s, 3 H), 3.31 (s, 3 H), 3.78 (t, *J*=5.3 Hz, 2 H), 4.49 (t, *J*=5.3 Hz, 2 H), 7.66 (ddd, *J*=8.5, 7.1, 1.4 Hz, 1 H), 7.81 (ddd, *J*=8.5, 7.1, 1.4 Hz, 1 H), 7.94 (s, 1 H), 7.99 (dt, *J*=8.5, 0.7 Hz, 1 H), 8.83 (dd, *J*=8.6, 1.2 Hz, 1 H); MS (DCI/NH₃) *m/z* 376 (M+H)⁺; Anal. calculated for C₁₈H₁₈ClN₃O₂S: C, 57.52; H, 4.83; N, 11.18. Found: C, 57.44; H, 4.59; N, 10.97.

Example 139

10 N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-1,2-dihydroacenaphthylene-5-carboxamide

The acenaphthene-5-carboxylic acid (Aldrich, 0.29 g, 1.4 mmol) in 5 mL of thionyl chloride was converted to acenaphthene-5-carbonyl chloride using the method described in Example 134A. The product of Example 126A (0.20 g, 1.1 mmol), Et₃N
15 (0.3 mL, 2.2 mmol) and acenaphthene-5-carbonyl chloride in 30 mL THF were processed as in Example 134B to afford the title compound (87 mg, 0.24 mmol, 22% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 2.28 (d, *J*=0.7 Hz, 3 H), 2.33 (d, *J*=0.7 Hz, 3 H), 3.33 (s, 3 H), 3.42 (s, 4 H), 3.84 (t, *J*=5.3 Hz, 2 H), 4.51 (t, *J*=5.3 Hz, 2 H), 7.30-7.36 (m, 2 H), 7.52 (dd, *J*=8.6, 7.0 Hz, 1 H), 8.43 (d, *J*=7.5 Hz, 1 H), 8.81 (d,
20 *J*=8.8 Hz, 1 H); MS (DCI/NH₃) *m/z* 367 (M+H)⁺; Anal. calculated for C₂₁H₂₂N₂O₂S: C, 68.82; H, 6.05; N, 7.64. Found: C, 68.63; H, 5.72; N, 7.40.

Example 140

25 2,3-dichloro-N-[(2Z)-3-(2-methoxyethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]benzamide

Example 140 A

3-(2-Methoxy-ethyl)-5-methyl-3H-thiazol-2-ylideneamine
The product of Example 10A (17.6 g, 70 mmol) was treated with ~50 mL 20%
30 aqueous K₂CO₃ then the mixture was extracted with EtOAc (3 X 25 mL). The combined organics were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford the title compound (5.9 g, 34 mmol, 49% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 2.03 (d, *J*=1.70 Hz, 3 H) 3.36 (s, 3 H) 3.62 (t, *J*=5.10

Hz, 2 H) 3.83 (t, $J=4.80$ Hz, 2 H) 6.15 - 6.21 (m, 1 H); MS (DCI/ NH_3) m/z 173 ($M+H$)⁺.

Example 140 B

2,3-dichloro-N-[(2Z)-3-(2-methoxyethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]benzamide

To the product of Example 140A (0.20 g, 1.2 mmol) in 15 mL THF was added Et_3N (0.48 mL, 3.5 mmol) followed by a solution of 2,3-dichlorobenzoyl chloride (Lancaster, 0.31 g, 1.5 mmol) in 5 mL THF. This mixture stirred at ambient temperature for 1 hour then was concentrated under reduced pressure, quenched with 5 mL saturated, aqueous NH_4Cl and extracted with EtOAc (3 X 5 mL). The combined organic extracts were dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. The resulting solids were recrystallized from 50% hexanes/EtOAc to afford the title compound (0.27 g, 0.78 mmol, 68% yield). ^1H NMR (300 MHz, CDCl_3) δ ppm 2.32 (d, $J=1.4$ Hz, 3 H), 3.34 (s, 3 H), 3.71 (t, $J=5.1$ Hz, 2 H), 4.34 (t, $J=4.8$ Hz, 2 H), 6.82-6.86 (m, 1 H), 7.22 (t, $J=7.8$ Hz, 1 H), 7.49 (dd, $J=8.1, 1.7$ Hz, 1 H), 7.73 (dd, $J=7.5, 1.7$ Hz, 1 H); MS (DCI/ NH_3) m/z 345 ($M+H$)⁺; Anal. calculated for $\text{C}_{14}\text{H}_{14}\text{Cl}_2\text{N}_2\text{O}_2\text{S}$: C, 48.70; H, 4.09; N, 8.11. Found: C, 48.39; H, 3.70; N, 7.94.

Example 141

5-chloro-N-[(2Z)-3-(2-methoxyethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]-2-(trifluoromethyl)benzamide

The product of Example 140A (0.20 g, 1.2 mmol), Et_3N (0.48 mL, 3.5 mmol) and 5-chloro-2-(trifluoromethyl)benzoyl chloride (Matrix, 0.31 g, 1.5 mmol) in 20 mL THF were processed using the method described in Example 140B to afford the title compound (0.16 g, 0.42 mmol, 36% yield). ^1H NMR (300 MHz, CDCl_3) δ ppm 2.33 (d, $J=0.7$ Hz, 3 H), 3.34 (s, 3 H), 3.69 (t, $J=5.1$ Hz, 2 H), 4.34 (t, $J=4.6$ Hz, 2 H), 6.85 (s, 1 H), 7.42-7.48 (m, 1 H), 7.64 (d, $J=8.5$ Hz, 1 H), 7.85 (d, $J=2.0$ Hz, 1 H); MS (DCI/ NH_3) m/z 379 ($M+H$)⁺; Anal. calculated for $\text{C}_{15}\text{H}_{14}\text{ClF}_3\text{N}_2\text{O}_2\text{S}$: C, 47.56; H, 3.73; N, 7.40. Found: C, 47.31; H, 3.30; N, 7.33.

Example 142

2-chloro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-

ylidene]benzamide

The product of Example 126A (0.20 g, 1.1 mmol), Et₃N (0.45 mL, 3.2 mmol) and 2-chlorobenzoyl chloride (Aldrich, 0.26 g, 1.4 mmol) in 15 mL THF were processed as in Example 132 to afford the title compound (0.13 g, 0.40 mmol, 37% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 2.28 (s, 3 H), 2.32 (s, 3 H), 3.31 (s, 3 H), 3.76 (t, *J*=5.3 Hz, 2 H), 4.42 (t, *J*=5.1 Hz, 2 H), 7.30-7.46 (m, 3 H), 7.77-7.83 (m, 1 H); MS (DCI/NH₃) *m/z* 325 (M+H)⁺; Anal. calculated for C₁₅H₁₇ClN₂O₂S: C, 55.46; H, 5.28; N, 8.62. Found: C, 55.59; H, 4.81; N, 8.47.

Example 143N-[(2Z)-5-tert-butyl-3-(2-methoxyethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-ethoxybenzamideExample 143A5-tert-Butyl-3-(2-methoxyethyl)-4-methyl-1,3-thiazol-2(3H)-ylideneamine

A mixture of 5-tert-butyl-4-methylthiazole-2-ylamine (1.5 g, 8.8 mmol) and 2-bromoethyl methyl ether (0.91 mL, 9.7 mmol) was warmed to 85 °C and allowed to stir for 24 hours. The crude material was dissolved in ~5 mL of a 1:1 mixture of CH₂Cl₂ and CH₃OH and a small amount of silica gel was added. This mixture was concentrated to dryness and the residue was purified via flash column chromatography (SiO₂, 9:1:0.1 CH₂Cl₂:CH₃OH:NH₄OH) to afford the title compound (1.0 g, 4.4 mmol, 50% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 1.41 (s, 9 H) 2.38 (s, 3 H) 3.35 (s, 3 H) 3.66 (t, *J*=4.70 Hz, 2 H) 4.16 (t, *J*=4.70 Hz, 2 H); MS (DCI/NH₃) *m/z* 229 (M+H)⁺.

Example 143BN-[(2Z)-5-tert-butyl-3-(2-methoxyethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-ethoxybenzamide

A mixture of the product of Example 143A (0.15 g, 0.66 mmol), Et₃N (0.28 mL, 2.0 mmol) and 2-ethoxybenzoyl chloride (Aldrich, 0.16 g, 0.86 mmol) in 15 mL THF were processed using the method described in Example 129 to afford the title compound (0.12 g, 0.33 mmol, 50% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 1.41 (t, *J*=7.1 Hz, 3 H), 1.45 (s, 9 H), 2.47 (s, 3 H), 3.31 (s, 3 H), 3.77 (t, *J*=5.3 Hz, 2 H), 4.11 (q, *J*=6.8 Hz, 2 H), 4.43 (t, *J*=5.3 Hz, 2 H), 6.95 (dt, *J*=7.5, 1.0 Hz, 1 H), 7.04 (d,

$J=8.1$ Hz, 1 H), 7.37 (ddd, $J=8.7, 6.9, 1.7$ Hz, 1 H), 7.78 (dd, $J=7.6, 1.9$ Hz, 1 H); MS (DCI/NH₃) m/z 377 (M+H)⁺; Anal. calculated for C₂₀H₂₈N₂O₃S: C, 63.80; H, 7.50; N, 7.44. Found: C, 64.19; H, 7.44; N, 7.19.

5

Example 144N-[(2Z)-5-tert-butyl-3-(2-methoxyethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2,3-dichlorobenzamide

A mixture of the product of Example 143A (0.15 g, 0.66 mmol), Et₃N (0.28 mL, 2.0 mmol) and 2,3-dichlorobenzoyl chloride (Lancaster, 0.18 g, 0.86 mmol) in 15 mL THF were processed using the method described in Example 129 to afford the title compound (0.11 g, 0.27 mmol, 42% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 1.43 (s, 9 H), 2.43 (s, 3 H), 3.30 (s, 3 H), 3.74 (t, $J=5.3$ Hz, 2 H), 4.36 (t, $J=5.3$ Hz, 2 H), 7.22 (dd, $J=8.0, 8.0$ Hz, 1 H), 7.48 (dd, $J=8.1, 1.7$ Hz, 1 H), 7.72 (dd, $J=7.6, 1.5$ Hz, 1 H); MS (DCI/NH₃) m/z 401 (M+H)⁺; Anal. calculated for C₁₈H₂₂Cl₂N₂O₂S: C, 53.87; H, 5.53; N, 6.98. Found: C, 53.86; H, 5.37; N, 6.76.

15

Example 145N-[(2Z)-5-tert-butyl-3-(2-methoxyethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-(trifluoromethyl)benzamide

A mixture of the product of Example 143A (0.15 g, 0.66 mmol), Et₃N (0.28 mL, 2.0 mmol) and 5-chloro-2-trifluoromethylbenzoyl chloride (Matrix, 0.17 g, 0.86 mmol) in 15 mL THF were processed using the method described in Example 129 to afford the title compound (0.16 g, 0.37 mmol, 56% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 1.46 (s, 9 H), 2.49 (s, 3 H), 3.31 (s, 3 H), 3.70 (t, $J=5.3$ Hz, 2 H), 4.41 (t, $J=5.3$ Hz, 2 H), 7.56-7.62 (m, 1 H), 7.71-7.76 (m, 2 H); MS (DCI/NH₃) m/z 435 (M+H)⁺; Anal. calculated for C₁₉H₂₂ClF₃N₂O₂S: C, 52.47; H, 5.10; N, 6.44. Found: C, 52.52; H, 4.94; N, 6.05.

25

Example 146N-[(2Z)-3-(2-methoxyethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]-1-naphthamide

The product of Example 140A (0.20 g, 1.2 mmol), Et₃N (0.48 mL, 3.5 mmol) and 1-naphthoyl chloride (Aldrich, 0.22 g, 1.5 mmol) in 15 mL THF were processed using the method described in Example 140B to afford the title compound (0.23 g,

30

0.69 mmol, 60% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 2.36 (d, *J*=1.0 Hz, 3 H), 3.35 (s, 3 H), 3.78 (t, *J*=5.4 Hz, 2 H), 4.45 (t, *J*=5.1 Hz, 2 H), 7.10-7.15 (m, 1 H), 7.46-7.58 (m, 3 H), 7.88-7.94 (m, 1 H), 7.97 (d, *J*=8.1 Hz, 1 H), 8.17 (dd, *J*=7.3, 1.2 Hz, 1 H), 8.83-8.90 (m, 1 H); MS (DCI/NH₃) *m/z* 327 (M+H)⁺; Anal. calculated for C₁₈H₁₈N₂O₂S: C, 66.23; H, 5.56; N, 8.58. Found: C, 66.10; H, 5.64; N, 8.51.

Example 147

N-[(2Z)-5-tert-butyl-3-(2-methoxyethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

The 5-chloro-2-methoxybenzoic acid (Aldrich, 0.21 g, 1.1 mmol) was converted to 5-chloro-2-methoxybenzoyl chloride with 5 mL of thionyl chloride using the method described in Example 134A. A mixture of the product of Example 143A (0.20 g, 0.88 mmol), Et₃N (0.48 mL, 3.4 mmol) and 5-chloro-2-methoxybenzoyl chloride in 15 mL THF were processed using the method described in Example 129 to afford the title compound (0.13 g, 0.32 mmol, 37% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 1.41 (s, 9 H), 2.42 (s, 3 H), 3.32 (s, 3 H), 3.77 (t, *J*=5.3 Hz, 2 H), 3.90 (s, 3 H), 4.36 (t, *J*=5.1 Hz, 2 H), 6.90 (d, *J*=8.8 Hz, 1 H), 7.31 (dd, *J*=8.8, 3.1 Hz, 1 H), 7.94 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 397 (M+H)⁺; Anal. calculated for C₁₉H₂₅ClN₂O₃S: C, 57.49; H, 6.35; N, 7.06. Found: C, 57.51; H, 6.30; N, 6.85.

Example 148

N-[(2Z)-3-(2-methoxyethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]-2-methylbenzamide

The product of Example 140A (0.20 g, 1.2 mmol), Et₃N (0.48 mL, 3.5 mmol) and *o*-toloyl chloride (Aldrich, 0.23 g, 1.2 mmol) in 15 mL THF were processed using the method described in Example 140B to afford the title compound (0.26 g, 0.90 mmol, 78% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 2.30 (s, 3 H), 2.70 (s, 3 H), 3.35 (s, 3 H), 3.74 (t, *J*=5.1 Hz, 2 H), 4.36 (t, *J*=4.8 Hz, 2 H), 6.77-6.81 (m, 1 H), 7.23 (t, *J*=7.3 Hz, 2 H), 7.28-7.35 (m, 1 H), 8.08-8.13 (m, 1 H); MS (DCI/NH₃) *m/z* 291 (M+H)⁺; Anal. calculated for C₁₅H₁₈N₂O₂S: C, 62.04; H, 6.25; N, 9.65. Found: C, 62.40; H, 6.11; N, 9.70.

Example 149

2,3-dichloro-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-

ylidene]benzamideExample 149A5-Methyl-3-(tetrahydro-furan-2-ylmethyl)-3H-thiazol-2-ylideneamine

- 5 A mixture of 2-amino-5-methylthiazole (1.0 g, 8.7 mmol) and 2-(bromomethyl)tetrahydrofuran (Maybridge, 1.1 mL, 10 mmol) was warmed to 85 °C and allowed to stir for 24 hours. The mixture was then cooled to ambient temperature and purified via column chromatography (SiO₂, 9:1:0.1 CH₂Cl₂:CH₃OH:NH₄OH) to afford the title compound (1.5 g, 7.5 mmol, 86% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 1.55 - 1.70 (m, 1 H) 1.89 - 2.01 (m, 2 H) 2.07 - 2.20 (m, 1 H) 2.28 (d, J=1.36 Hz, 3 H) 3.72 - 3.82 (m, 1 H) 3.86 - 4.00 (m, 2 H) 4.08 - 4.24 (m, 2 H) 6.98 - 7.04 (m, 1 H); MS (DCI/NH₃) m/z 199 (M+H)⁺.
- 10

Example 149B

2,3-dichloro-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

15

- To a solution of the product of Example 149A (0.17 g, 0.86 mmol) in 10 mL THF and 1 mL DMF was added Et₃N (0.36 mL, 2.6 mmol) followed by 2,3-dichlorobenzoyl chloride (Lancaster, 0.27 g, 1.3 mmol). This mixture was warmed to 50 °C and allowed to stir for 2 hours. The mixture was cooled to ambient temperature, diluted with 10 mL EtOAc, and quenched with 10 mL NH₄Cl. The layers were separated and the aqueous layer was extracted 2 X 5 mL EtOAc. The combined organics were dried over anhydrous Na₂SO₄, filtered, concentrated under reduced pressure and purified via column chromatography (SiO₂, 20% hexanes/EtOAc) to afford the title compound (0.24 g, 0.64 mmol, 75% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 1.58-1.68 (m, 1 H), 1.78-1.94 (m, 2 H), 2.00-2.13 (m, 1 H), 2.32 (s, 3 H), 3.72-3.91 (m, 2 H), 4.06-4.15 (m, 1 H), 4.24 (ddd, J=14.0, 7.0, 2.7 Hz, 1 H), 4.47 (dd, J=13.6, 2.7 Hz, 1 H), 6.91-6.95 (m, 1 H), 7.22 (t, J=8.0 Hz, 1 H), 7.49 (dd, J=8.1, 1.7 Hz, 1 H), 7.72 (dd, J=7.5, 1.7 Hz, 1 H); MS (DCI/NH₃) m/z 371 (M+H)⁺; Anal. calculated for C₁₆H₁₆Cl₂N₂O₂S: C, 51.76; H, 4.34; N, 7.55.
- 20
- 25
- 30 Found: C, 51.66; H, 4.17; N, 7.46.

Example 150N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-

methylbenzamide

The product of Example 126A (0.20 g, 1.1 mmol), Et₃N (0.45 mL, 3.2 mmol) and o-toloyl chloride (Aldrich, 0.22 g, 1.4 mmol) in 15 mL THF were processed using the method described in Example 129 to afford the title compound (0.15 g, 0.49 mmol, 46% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 2.23 (d, J=0.7 Hz, 3 H), 2.26 (s, 3 H), 2.70 (s, 3 H), 3.31 (s, 3 H), 3.77 (t, J=5.3 Hz, 2 H), 4.35 (t, J=5.3 Hz, 2 H), 7.23 (t, J=7.5 Hz, 2 H), 7.27-7.35 (m, 1 H), 8.08-8.14 (m, 1 H); MS (DCI/NH₃) m/z 305 (M+H)⁺; Anal. calculated for C₁₆H₂₀N₂O₂S: C, 63.13; H, 6.62; N, 9.20. Found: C, 63.43; H, 6.53; N, 9.14.

Example 151N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 126A (0.20 g, 1.1 mmol), Et₃N (0.45 mL, 3.2 mmol) and benzoyl chloride (Aldrich, 0.16 mL, 1.4 mmol) in 15 mL THF were processed using the method described in Example 129 to afford the title compound (0.11 g, 0.38 mmol, 35% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 2.24 (s, 3 H), 2.27 (d, J=0.7 Hz, 3 H), 3.32 (s, 3 H), 3.82 (t, J=5.3 Hz, 2 H), 4.41 (t, J=5.3 Hz, 2 H), 7.38-7.51 (m, 3 H), 8.27-8.34 (m, 2 H); MS (DCI/NH₃) m/z 291 (M+H)⁺; Anal. calculated for C₁₅H₁₈N₂O₂S: C, 62.04; H, 6.25; N, 9.65. Found: C, 62.02; H, 6.05; N, 9.56.

Example 1522-chloro-4-fluoro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 126A (0.20 g, 1.1 mmol), Et₃N (0.37 mL, 2.7 mmol) and 2-chloro-4-fluorobenzene-1-carbonyl chloride (Acros, 0.25 g, 1.3 mmol) in 15 mL THF were processed using the method described in Example 132 to afford the title compound (0.19 g, 0.56 mmol, 52% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 2.24 (s, 3 H), 2.27 (s, 3 H), 3.30 (s, 3 H), 3.76 (t, J=5.1 Hz, 2 H), 4.35 (t, J=5.1 Hz, 2 H), 7.00 (ddd, J=8.5, 7.8, 2.4 Hz, 1 H), 7.15 (dd, J=8.8, 2.4 Hz, 1 H), 8.04 (dd, J=8.8, 6.4 Hz, 1 H); MS (DCI/NH₃) m/z 343 (M+H)⁺; Anal. calculated for C₁₅H₁₆ClFN₂O₂S: C, 52.55; H, 4.70; N, 8.17. Found: C, 52.60; H, 4.38; N, 8.06.

Example 153

2-chloro-4-fluoro-N-[(2Z)-3-(2-methoxyethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 140A (0.17 g, 1.0 mmol), Et₃N (0.35 mL, 2.5 mmol) and 2-chloro-4-fluorobenzene-1-carbonyl chloride (Acros, 0.23 g, 1.2 mmol) in 15 mL THF were processed using the method described in Example 140B to afford the title compound (0.15 g, 0.46 mmol, 46% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 2.32 (d, J=1.4 Hz, 3 H), 3.35 (s, 3 H), 3.73 (t, J=5.4 Hz, 2 H), 4.36 (t, J=4.7 Hz, 2 H), 6.80-6.84 (m, 1 H), 7.00 (ddd, J=8.6, 7.8, 2.5 Hz, 1 H), 7.16 (dd, J=8.6, 2.5 Hz, 1 H), 8.04 (dd, J=8.8, 6.4 Hz, 1 H); MS (DCI/NH₃) m/z 329 (M+H)⁺; Anal. calculated for C₁₄H₁₄ClFN₂O₂S: C, 51.15; H, 4.29; N, 8.52. Found: C, 51.11; H, 3.90; N, 8.43.

Example 154

2,5-dichloro-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]benzamide

The 2,5-dichlorobenzoic acid (Aldrich, 0.28 g, 1.4 mmol) was converted to 2,5-dichlorobenzoyl chloride with 5 mL of thionyl chloride using the method described in Example 134A. The product of Example 126A (0.20 g, 1.1 mmol), Et₃N (0.45 mL, 3.2 mmol) and 2,5-dichlorobenzoyl chloride in 15 mL THF were processed using the method described in Example 132 to afford the title compound (0.10 g, 0.28 mmol, 26% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 2.25 (d, J=1.0 Hz, 3 H), 2.28 (d, J=1.0 Hz, 3 H), 3.31 (s, 3 H), 3.76 (t, J=5.3 Hz, 2 H), 4.36 (t, J=5.3 Hz, 2 H), 7.28 (d, J=2.4 Hz, 1 H), 7.32-7.36 (m, 1 H), 7.94 (d, J=2.4 Hz, 1 H); MS (DCI/NH₃) m/z 359 (M+H)⁺; Anal. calculated for C₁₅H₁₆Cl₂N₂O₂S: C, 50.15; H, 4.49; N, 7.80. Found: C, 50.22; H, 4.15; N, 7.63.

Example 155

2,5-dichloro-N-[(2Z)-3-(2-methoxyethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]benzamide

The 2,5-dichlorobenzoic acid (Aldrich, 0.28 g, 1.4 mmol) was converted to 2,5-dichlorobenzoyl chloride with 5 mL of thionyl chloride using the method described in Example 134A. The product of Example 140A (0.20 g, 1.1 mmol), Et₃N (0.45 mL, 3.2 mmol) and 2,5-dichlorobenzoyl chloride in 15 mL THF were processed using the method described in Example 129 to afford the title compound (0.24 g, 0.70

mmol, 65% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 2.32 (d, *J*=1.4 Hz, 3 H), 3.35 (s, 3 H), 3.73 (t, *J*=5.1 Hz, 2 H), 4.37 (t, *J*=4.7 Hz, 2 H), 6.83-6.86 (m, 1 H), 7.28 (dd, *J*=8.5, 2.4 Hz, 1 H), 7.35 (d, *J*=8.5 Hz, 1 H), 7.94 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 345 (M+H)⁺; Anal. calculated for C₁₄H₁₄Cl₂N₂O₂S: C, 48.70; H, 4.09; N, 8.11.

5 Found: C, 48.60; H, 3.78; N, 8.02.

Example 156

5-chloro-2-methoxy-N-[(2Z)-3-(2-methoxyethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]benzamide

10 The 5-chloro-2-methoxybenzoic acid (Aldrich, 0.28 g, 1.5 mmol) was converted to 5-chloro-2-methoxybenzoyl chloride with 5 mL of thionyl chloride using the method described in Example 134A. The product of Example 140A (0.19 g, 1.1 mmol), Et₃N (0.45 mL, 3.2 mmol) and 5-chloro-2-methoxybenzoyl chloride in 15 mL THF were processed using the method described in Example 129 to afford the title
15 compound (0.25 g, 0.72 mmol, 67% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 2.30 (d, *J*=1.0 Hz, 3 H), 3.36 (s, 3 H), 3.74 (t, *J*=5.1 Hz, 2 H), 3.90 (s, 3 H), 4.36 (t, *J*=4.7 Hz, 2 H), 6.82 (s, 1 H), 6.90 (d, *J*=8.8 Hz, 1 H), 7.33 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.96 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 341 (M+H)⁺; Anal. calculated for C₁₅H₁₇ClN₂O₃S: C, 52.86; H, 5.03; N, 8.22. Found: C, 52.84; H, 4.72; N, 8.13.

20

Example 157

2,3-dichloro-N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

25

Example 157A

5-Methyl-3-(tetrahydro-pyran-2-ylmethyl)-3H-thiazol-2-ylideneamine

A mixture of 2-amino-5-methylthiazole (1.2 g, 10.5 mmol) and 2-(bromomethyl)tetrahydro-2H-pyran (Aldrich, 1.5 mL, 11.6 mmol) was warmed to 85 °C and allowed to stir for 18 hours. The mixture was cooled to ambient temperature
30 and purified by flash column chromatography (SiO₂, first 10% CH₃OH:EtOAc then 9:1:0.1 CH₂Cl₂:CH₃OH:NH₄OH) to afford the title compound (1.1 g, 5.2 mmol, 49% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 1.12 - 1.32 (m, 1 H) 1.44 - 1.63 (m, 2 H) 1.79 - 1.95 (m, 2 H) 2.06 (d, *J*=12.89 Hz, 1 H) 2.25 (d, *J*=1.36 Hz, 3 H) 3.33 - 3.45

(m, 1 H) 3.65 - 3.78 (m, 1 H) 3.86 - 4.01 (m, 2 H) 4.44 (dd, $J=14.92$, 2.03 Hz, 1 H)
6.56 - 6.65 (m, 1 H) 9.48 (s, 1 H); MS (DCI/NH₃) m/z 213 (M+H)⁺.

Example 157B

2,3-dichloro-N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol- 2(3H)-ylidene]benzamide

The product of Example 157A (0.15 g, 0.71 mmol), Et₃N (0.30 mL, 2.1 mmol) and 2,3-dichlorobenzoyl chloride (Lancaster, 0.19 g, 0.92 mmol) in 15 mL THF were processed as in Example 129 to afford the title compound (0.14 g, 0.36 mmol, 51% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 1.16-1.33 (m, 1H), 1.46-1.56 (m, 3 H), 1.64-1.73 (m, 1 H), 1.80-1.92 (m, 1 H), 2.32 (d, $J=1.4$ Hz, 3 H), 3.30-3.43 (m, 1 H), 3.69 (qt, $J=11.5, 8.8, 2.0$ Hz, 1 H), 3.91-4.01 (m, 2 H), 4.41 (dd, $J=13.9, 2.7$ Hz, 1 H), 6.86 (d, $J=1.4$ Hz, 1 H), 7.23 (t, $J=7.8$ Hz, 1 H), 7.49 (dd, $J=8.0, 1.5$ Hz, 1 H), 7.74 (dd, $J=7.6, 1.5$ Hz, 1 H); MS (DCI/NH₃) m/z 385 (M+H)⁺; Anal. calculated for C₁₇H₁₈Cl₂N₂O₂S: C, 52.99; H, 4.71; N, 7.27. Found: C, 53.15; H, 4.72; N, 7.14.

Example 158

2-ethoxy-N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)- ylidene]benzamide

The product of Example 157A (0.15 g, 0.71 mmol), Et₃N (0.30 mL, 2.1 mmol) and 2-ethoxybenzoyl chloride (Aldrich, 0.18 g, 0.92 mmol) in 15 mL THF were processed as in Example 129 to afford the title compound (0.14 g, 0.39 mmol, 55% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 1.20-1.32 (m, 1 H), 1.46 (t, $J=7.0$ Hz, 3 H), 1.50-1.54 (m, 3 H), 1.63-1.79 (m, 1 H), 1.80-1.90 (m, 1 H), 2.29 (s, 3 H), 3.32-3.43 (m, 1 H), 3.64-3.76 (m, 1 H), 3.92-4.00 (m, 2 H), 4.16 (q, $J=6.8$ Hz, 2 H), 4.37-4.47 (m, 1 H), 6.76-6.84 (m, 1 H), 6.93-7.00 (m, 2 H), 7.35 (t, $J=8.1$ Hz, 1 H), 7.96 (dd, $J=8.0, 1.9$ Hz, 1 H); MS (DCI/NH₃) m/z 361 (M+H)⁺; Anal. calculated for C₁₉H₂₄N₂O₃S: C, 63.61; H, 6.71; N, 7.77. Found: C, 63.56; H, 6.73; N, 7.26.

Example 159

5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol- 2(3H)-ylidene]benzamide

The 5-chloro-2-methoxybenzoic acid (Aldrich, 0.66 g, 3.5 mmol) was converted to 5-chloro-2-methoxybenzoyl chloride with 5 mL of thionyl chloride using

the method described in Example 134A. The product of Example 149A (0.35 g, 0.18 mmol), Et₃N (0.74 mL, 5.3 mmol) and 5-chloro-2-methoxybenzoyl chloride in 20 mL THF were processed using the method described in Example 149B to afford the title compound (0.25 g, 0.68 mmol, 20% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 1.67 - 1.78 (m, 1 H), 1.84 - 1.96 (m, 2 H), 2.00 - 2.13 (m, 1 H), 2.34 (d, *J*=1.4 Hz, 3 H), 3.71 - 3.81 (m, 1 H), 3.85 (s, 3 H), 3.86 - 3.93 (m, 1 H), 4.20 - 4.42 (m, 3 H), 7.07 (d, *J*=9.2 Hz, 1 H), 7.12 - 7.16 (m, 1 H), 7.39 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.81 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 367 (M+H)⁺; Anal. calculated for C₁₇H₁₉ClN₂O₃S•0.2H₂O : C, 55.12; H, 5.28; N, 7.56. Found: C, 54.90; H, 4.95; N, 7.55.

10

Example 160

5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

The 5-chloro-2-methoxybenzoic acid (Aldrich, 0.17 g, 0.92 mmol) was converted to 5-chloro-2-methoxybenzoyl chloride with 5 mL of thionyl chloride using the method described in Example 134A. A mixture of the product of Example 157A (0.15 g, 0.71 mmol), Et₃N (0.30 mL, 2.1 mmol) and 5-chloro-2-methoxybenzoyl chloride in 15 mL THF were processed using the method described in Example 129 to afford the title compound (0.11 g, 0.29 mmol, 41% yield). ¹H NMR (300 MHz, CDCl₃) δ ppm 1.20-1.36 (m, 1 H), 1.48-1.61 (m, 3 H), 1.66-1.76 (m, 1 H), 1.83-1.92 (m, 1 H), 2.30 (d, *J*=1.0 Hz, 3 H), 3.33-3.44 (m, 1 H), 3.67-3.77 (m, 1 H), 3.90 (s, 3 H), 3.93-4.05 (m, 2 H), 4.40 (dd, *J*=13.9, 2.4 Hz, 1 H), 6.83 (s, 1 H), 6.90 (d, *J*=8.8 Hz, 1 H), 7.32 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.97 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 381 (M+H)⁺; Anal. calculated for C₁₈H₂₁ClN₂O₃S•0.15H₂O: C, 56.36; H, 5.60; N, 7.30. Found: C, 56.70; H, 5.41; N, 6.91.

25

Example 161

2-ethoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 149A (0.20 g, 1.0 mmol), Et₃N (0.42 mL, 3.0 mmol) and the 2-ethoxybenzoyl chloride (Aldrich, 0.23 mL, 1.5 mmol) in 15 mL THF were processed using the method described in Example 149B to afford the title compound (0.18 g, 0.52 mmol, 52% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 1.40 (t, *J*=7.0

30

Hz, 3 H), 1.64-1.75 (m, 1 H), 1.83-1.94 (m, 2 H), 2.00-2.12 (m, 1 H), 2.33 (d, $J=1.4$ Hz, 3 H), 3.70-3.80 (m, 1 H), 3.83-3.93 (m, 1 H), 4.11 (q, $J=7.1$ Hz, 2 H), 4.18-4.27 (m, 1 H), 4.28-4.42 (m, 2 H), 6.96 (dt, $J=7.5, 1.0$ Hz, 1 H), 7.04 (d, $J=8.1$ Hz, 1 H), 7.09-7.13 (m, 1 H), 7.38 (ddd, $J=8.4, 7.4, 1.9$ Hz, 1 H), 7.77 (dd, $J=7.6, 1.9$ Hz, 1 H);

- 5 MS (DCI/ NH_3) m/z 347 ($\text{M}+\text{H}$)⁺; Anal. calculated for $\text{C}_{18}\text{H}_{22}\text{N}_2\text{O}_3\text{S}\cdot 0.1\text{H}_2\text{O}$: C, 62.40; H, 6.40; N, 8.09. Found: C, 63.49; H, 5.90; N, 7.84.

Example 162

10 5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

Example 162A

Toluene-4-sulfonic acid tetrahydro-furan-3-ylmethyl ester

To a solution of tetrahydro-3-furanmethanol (Aldrich, 1.0 mL, 10.4 mmol) in
 15 5 mL CH_2Cl_2 and 5 mL pyridine was added para-toluenesulfonyl chloride (3.0 g, 15.6 mmol) portion-wise over 15 minutes. This mixture stirred at ambient temperature for 3 hours then 5 mL H_2O was added. The layers were separated and the aqueous layer was extracted 2 X 5 mL CH_2Cl_2 . The combined organics were dried over Na_2SO_4 , filtered, concentrated under reduced pressure and dried under vacuum (~1 mm Hg) to
 20 afford the title compound (2.62 g, 10.2 mmol, 98% yield). ^1H NMR (300 MHz, CDCl_3) δ ppm 1.49 - 1.63 (m, 1 H) 1.94 - 2.08 (m, 1 H) 2.46 (s, 3 H) 2.52 - 2.68 (m, 1 H) 3.49 (dd, $J=9.16, 5.09$ Hz, 1 H) 3.64 - 3.84 (m, 3 H) 3.88 - 4.03 (m, 2 H) 7.36 (d, $J=8.14$ Hz, 2 H) 7.76 - 7.82 (m, 2 H); MS (DCI/ NH_3) m/z 257 ($\text{M}+\text{H}$)⁺.

25

Example 162B

5-Methyl-3-(tetrahydro-furan-3-ylmethyl)-3H-thiazol-2-ylideneamine

A mixture of the product of Example 162A (1.62 g, 6.3 mmol), 2-amino-5-methylthiazole (0.72 g, 6.3 mmol) and LiBr (55 mg, 0.63 mmol) in 2 mL DMF was warmed to 85 °C and allowed to stir for 16 hours. The mixture was then allowed to
 30 cool to ambient temperature, diluted with 10 mL CH_2Cl_2 and washed with 1 X 5 mL 10% aqueous Na_2CO_3 solution. The layers were separated and the aqueous layer was extracted 2 X 5 mL CH_2Cl_2 . The combined organics were dried over anhydrous Na_2SO_4 , filtered, concentrated under reduced pressure and purified by column

chromatography (SiO₂, 9:1:0.1 CH₂Cl₂:CH₃OH:NH₄OH) to afford the title compound (0.31 g, 1.6 mmol, 25% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 1.61 - 1.74 (m, 1 H) 1.96 - 2.04 (m, 1 H) 2.05 (d, *J*=1.36 Hz, 3 H) 2.69 - 2.84 (m, 1 H) 3.53 (dd, *J*=8.82, 5.76 Hz, 1 H) 3.63 (dd, *J*=7.63, 2.20 Hz, 2 H) 3.69 - 3.81 (m, 2 H) 3.89 (ddd, *J*=8.31, 5.42 Hz, 1 H) 6.36 - 6.42 (m, 1 H); MS (DCI/NH₃) *m/z* 199 (M+H)⁺.

Example 162C

5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-3-yl)methyl]-1,3-thiazol-2(3H)-ylidene]benzamide

The 5-chloro-2-methoxybenzoic acid (Aldrich, 0.40 g, 2.1 mmol) was converted to 5-chloro-2-methoxybenzoyl chloride with 5 mL of thionyl chloride using the method described in Example 134A. The product of Example 162B (0.21 g, 1.1 mmol), Et₃N (0.44 mL, 3.2 mmol) and 5-chloro-2-methoxybenzoyl chloride in 15 mL THF were processed using the method described in Example 149B to afford the title compound (0.28 g, 0.76 mmol, 72% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 1.69 - 1.83 (m, 1 H), 1.96 - 2.09 (m, 1 H), 2.34 (d, *J*=1.4 Hz, 3 H), 2.84 - 3.00 (m, 1 H), 3.64 (dd, *J*=8.8, 5.1 Hz, 1 H), 3.72 - 3.81 (m, 2 H), 3.85 (s, 3 H), 3.89 - 3.99 (m, 1 H), 4.17 - 4.33 (m, 2 H), 7.06 (d, *J*=8.8 Hz, 1 H), 7.14 - 7.17 (m, 1 H), 7.39 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.84 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 367 (M+H)⁺; Anal. calculated for C₁₇H₁₉ClN₂O₃S: C, 55.66; H, 5.22; N, 7.64. Found: C, 55.77; H, 4.85; N, 7.26.

Example 163

2-ethoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-3-yl)methyl]-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 162B (0.17 g, 0.86 mmol), Et₃N (0.36 mL, 2.6 mmol) and the 2-ethoxybenzoyl chloride (Aldrich, 0.15 mL, 0.94 mmol) in 10 mL THF were processed using the method described in Example 149B to afford the title compound (0.26 g, 0.74 mmol, 86% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 1.39 (t, *J*=7.0 Hz, 3 H), 1.68-1.82 (m, 1 H), 1.92-2.05 (m, 1 H), 2.34 (d, *J*=1.4 Hz, 3 H), 2.85-3.02 (m, 1 H), 3.63 (dd, *J*=8.8, 5.4 Hz, 1 H), 3.70-3.80 (m, 2 H), 3.88-3.97 (m, 1 H), 4.12 (q, *J*=6.9 Hz, 2 H), 4.17-4.33 (m, 2 H), 6.96 (dt, *J*=7.5, 1.0 Hz, 1 H), 7.05 (d, *J*=7.8 Hz, 1 H), 7.11-7.16 (m, 1 H), 7.38 (ddd, *J*=8.6, 7.0, 1.7 Hz, 1 H), 7.79 (dd, *J*=7.8, 1.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 347 (M+H)⁺; Anal. calculated for C₁₈H₂₂N₂O₃S: C,

62.40; H, 6.40; N, 8.09. Found: C, 62.43; H, 6.29; N, 7.96.

Example 164

5 2-ethoxy-N-[(2Z)-3-[2-(2-methoxyethoxy)ethyl]-5-methyl-1,3-thiazol-2(3H)-ylidene]benzamide

Example 164A

3-[2-(2-Methoxyethoxy)ethyl]-5-methyl-1,3-thiazol-2(3H)-ylideneamine

A mixture of 2-amino-5-methylthiazole (1.5 g, 13.0 mmol) and 1-bromo-2-(2-methoxyethoxy)ethane (Aldrich, 2.0 mL, 14.5 mmol) was processed as in Example 10 157A to afford the title compound (2.2 g, 10.9 mmol, 78% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 2.29 (d, J=1.36 Hz, 3 H) 3.34 (s, 3 H) 3.49 - 3.54 (m, 2 H) 3.60 - 3.66 (m, 2 H) 3.80 (t, J=5.10 Hz, 2 H) 4.13 (t, J=4.80 Hz, 2 H) 6.99 - 7.04 (m, 1 H); MS (DCI/NH₃) m/z 217 (M+H)⁺.

15 Example 164B

2-ethoxy-N-[(2Z)-3-[2-(2-methoxyethoxy)ethyl]-5-methyl-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 164A (0.22 g, 1.0 mmol), Et₃N (0.23 mL, 2.0 mmol) and 2-ethoxybenzoyl chloride (Aldrich, 0.25 g, 1.3 mmol) in 15 mL THF were 20 processed as in Example 129 to afford the title compound (0.15 g, 0.40 mmol, 40% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 1.40 (t, J=7.0 Hz, 3 H), 2.33 (d, J=1.4 Hz, 3 H), 3.31 (s, 3 H), 3.46-3.51 (m, 2 H), 3.58-3.63 (m, 2 H), 3.85 (t, J=5.3 Hz, 2 H), 4.11 (q, J=6.8 Hz, 2 H), 4.40 (t, J=5.3 Hz, 2 H), 6.96 (dt, J=7.5, 1.0 Hz, 1 H), 7.04 (d, J=8.5 Hz, 1 H), 7.11-7.15 (m, 1 H), 7.38 (ddd, J=8.4, 7.4, 1.9 Hz, 1 H), 7.79 (dd, 25 J=7.6, 1.9 Hz, 1 H); MS (DCI/NH₃) m/z 365 (M+H)⁺; Anal. calculated for C₁₈H₂₄N₂O₄S: C, 59.88; H, 6.86; N, 7.51. Found: C, 60.05; H, 6.81; N, 7.60.

Example 166

30 2,3-dichloro-N-[(2Z)-3-[2-(2-methoxyethoxy)ethyl]-5-methyl-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 164A (0.22 g, 1.0 mmol), Et₃N (0.23 mL, 2.0 mmol) and 2,3-dichlorobenzoyl chloride (Lancaster, 0.27 g, 1.3 mmol) in 15 mL THF were processed as in Example 129 to afford the title compound (95 mg, 0.24 mmol, 24%

yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 2.36 (d, *J*=1.4 Hz, 3 H), 3.31 (s, 3 H), 3.46-3.50 (m, 2 H), 3.57-3.62 (m, 2 H), 3.83 (t, *J*=5.4 Hz, 2 H), 4.40 (t, *J*=5.1 Hz, 2 H), 7.18-7.21 (m, 1 H), 7.33 (t, *J*=7.8 Hz, 1 H), 7.58 (dd, *J*=8.0, 1.5 Hz, 1 H), 7.64 (dd, *J*=7.6, 1.5 Hz, 1 H); MS (DCI/NH₃) *m/z* 389 (M+H)⁺; Anal. calculated for C₁₆H₁₈Cl₂N₂O₃S: C, 49.36; H, 4.66; N, 7.20. Found: C, 48.98; H, 4.60; N, 6.99

Example 167

5-chloro-2-methoxy-N-[(2Z)-3-[2-(2-methoxyethoxy)ethyl]-5-methyl-1,3-thiazol-2(3H)-ylidene]benzamide

The 5-chloro-2-methoxybenzoic acid (Aldrich, 0.24 g, 1.3 mmol) was converted to the 5-chloro-2-methoxybenzoyl chloride with 5 mL of thionyl chloride using the method described in Example 134A. The product of Example 164A (0.22 g, 1.0 mmol), Et₃N (0.42 mL, 3.0 mmol) and 5-chloro-2-methoxybenzoyl chloride in 15 mL THF were processed using the method described in Example 129 to afford the title compound (0.21 g, 0.55 mmol, 55% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 2.34 (d, *J*=1.4 Hz, 3 H), 3.31 (s, 3 H), 3.47-3.52 (m, 2 H), 3.59-3.64 (m, 2 H), 3.86 (s, 3 H), 3.88 (t, *J*=5.4 Hz, 2 H), 4.41 (t, *J*=4.8 Hz, 2 H), 7.07 (d, *J*=8.8 Hz, 1 H), 7.14-7.17 (m, 1 H), 7.39 (dd, *J*=9.0, 2.9 Hz, 1 H), 7.83 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 385 (M+H)⁺; Anal. calculated for C₁₇H₂₁ClN₂O₄S: C, 53.05; H, 5.50; N, 7.28. Found: C, 52.93; H, 5.61; N, 7.26.

Example 168

2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

The product of Example 149A (0.15 g, 0.76 mmol), Et₃N (0.32 mL, 2.3 mmol) and the *o*-anisoyl chloride (Aldrich, 0.15 mL, 1.1 mmol) in 10 mL THF were processed using the method described in Example 149B to afford the title compound (88 mg, 0.26 mmol, 35% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 1.66-1.76 (m, 1 H), 1.84-1.95 (m, 2 H), 2.02-2.12 (m, 1 H), 2.33 (d, *J*=1.4 Hz, 3 H), 3.71-3.79 (m, 1 H), 3.85 (s, 3 H), 3.84-3.92 (m, 1 H), 4.22-4.28 (m, 1 H), 4.29-4.41 (m, 2 H), 6.98 (dt, *J*=7.6, 1.0 Hz, 1 H), 7.07 (dd, *J*=8.5, 1.0 Hz, 1 H), 7.10-7.12 (m, 1 H), 7.42 (ddd, *J*=9.2, 7.5, 2.0 Hz, 1 H), 7.85 (dd, *J*=7.8, 1.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 333 (M+H)⁺; Anal. calculated for C₁₇H₂₀N₂O₃S: C, 61.42; H, 6.06; N, 8.43. Found: C,

61.35; H, 6.10; N, 8.28.

Example 169

1-(1,1-Dimethylpropyl)-3-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]urea hydrochloride

Example 169A

1-(1,1-Dimethylpropyl)-3-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]urea

To a solution of 1,1-dimethylpropylamine (174mg, 2.0 mmole) in 19 mL of THF and 1 mL of *N,N*-diisopropylethyl amine was added 4-nitrophenyl chloroformate (403 mg, 2.0 mmole). The solution was irradiated in a sealed tube placed in a single node microwave at 70 °C for 300 sec (maximum power 300W) with stirring. The resulting solution was cooled to room temperature and 3-(2-methoxyethyl)-4,5-dimethyl-3H-thiazol-2-ylideneamine hydrobromide (587 mg, 2.2 mmole) from Example 12A was added. The sealed tube was irradiated at 120 °C for 1800 sec with stirring. The mixture was cooled and the volatile components were removed under reduced pressure. The residue was partitioned between water and ethyl acetate. The phases were separated and organic extract was dried over anhydrous Na₂SO₄, filtered, and concentrated. Purification by column chromatography (SiO₂, 0-70% ethyl acetate/hexanes gradient) afforded the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.76 (t, *J*=7 Hz, 3 H) 1.20 (s, 6 H) 1.66 (d, *J*=7 Hz, 2 H) 2.06 (s, 3 H) 2.11 (s, 3 H) 3.30 (s, 3 H) 3.55 (t, *J*=5 Hz, 2 H) 4.06 (t, *J*=5 Hz, 2 H) 6.13 (s, 1 H), MS (DCI/NH₃) *m/z* 300 (M+H)⁺.

Example 169B

1-(1,1-Dimethylpropyl)-3-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]urea hydrochloride

To a solution of the product from Example 169A in MeOH was added a solution of HCl in Et₂O. The title compound was isolated by filtration. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.84 (t, *J*=7 Hz, 3 H) 1.26 (s, 6 H) 1.65 (q, *J*=7 Hz, 2 H) 2.24 (s, 6 H) 3.24 (s, 3 H) 3.64 (t, *J*=5 Hz, 2 H) 4.45 (s, 2 H), MS (DCI/NH₃) *m/z* 300 (M+H)⁺. Anal. Calculated for C₁₄H₂₅ClN₃O₂S: C, 50.06 H, 7.80; N, 12.51. Found: C, 50.11; H, 7.87; N, 12.35.

Example 1701-(1,1-Dimethyl-propyl)-3-[(2Z)-3-(2-methoxyethyl)-1,3-benzothiazol-2(3H)-ylidene]urea

5 To a solution of 1,1-dimethyl-propylamine (0.60 mL, 5.2 mmol) and triethylamine (0.40 mL, 2.9 mmol) in 12 mL of a 1:1 mixture of THF:DMF at 0 °C was added *p*-nitrophenylchloroformate (0.58 g, 2.9 mmol). After 30 minutes, the product of Example 64A (0.75 g, 2.6 mmol) and another aliquot of triethylamine (0.40 mL, 2.9 mmol) were added and the solution stirred at ambient temperature for 9

10 hours. The mixture was diluted with ethyl acetate then washed twice with water and brine. The organic extract was dried over magnesium sulfate, filtered, and concentrated under reduced pressure. Purification by column chromatography (SiO₂, 20-40% ethyl acetate/hexanes gradient) afforded 0.06 g (8%) of the title compound.

¹H NMR (CDCl₃, 300 MHz) δ ppm 0.91 (t, *J*=7.46 Hz, 3 H), 1.39 (s, 6 H), 1.78 (q, *J*=7.46 Hz, 2 H), 3.28 (s, 3 H), 3.86 (t, *J*=4.92 Hz, 2 H), 4.82 (t, *J*=4.58 Hz, 2 H), 7.46

15 (t, *J*=7.63 Hz, 1 H), 7.58 (t, *J*=7.63 Hz, 1 H), 7.64 - 7.70 (m, 1 H), 7.75 (d, *J*=7.80 Hz, 1 H), 9.06 (s, 1H). MS (DCI/NH₃) *m/z* 322 (M+H)⁺.

Example 171

20 1-[(2Z)-3-(2-Methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-3-(3,3,5,5-tetramethylcyclohexyl)urea

The product of Example 12A and 3,3,5,5-tetramethylcyclohexylamine were processed using the method described in Example 169A to afford the title compound.

¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.90 (s, 6 H) 1.02 (s, 6 H) 1.17 - 1.27 (m, 1 H)

25 1.61 (d, *J*=12 Hz, 1 H) 2.18 - 2.27 (m, 6 H) 3.24 (s, 3 H) 3.62 (t, *J*=5 Hz, 2 H) 3.76 - 3.87 (m, *J*=5 Hz, 2 H) 3.98 (s, 2 H) 4.40 (s, 2 H), MS (DCI/NH₃) *m/z* 366 (M+H)⁺.

Anal. Calculated for C₁₉H₁₄ClN₃O₂S•0.7CH₄O: C, 55.49 H, 8.7 N, 9.85. Found C, 55.81 H, 8.37 N, 9.52.

30

Example 172

1-[(2Z)-3-(2-Methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-3-(1-methyl-3-phenylpropyl)urea

The product of Example 12A and 1-methyl-3-phenylpropylamine were processed using the method described in Example 169A to afford the title compound.

¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.13 (d, *J*=6 Hz, 3 H) 1.60 - 1.82 (m, 2 H) 2.18 (s, 3 H) 2.20 (s, 3 H) 2.55 - 2.68 (m, 2 H) 3.25 (s, 3 H) 3.53 - 3.80 (m, 3 H) 4.31 (s, 2 H) 7.08 - 7.35 (m, 5 H), MS (DCI/NH₃) *m/z* 362 (M+H)⁺.

Example 173

ethyl *N*-({[(2*Z*)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3*H*)-ylidene]amino}carbonyl)-L-leucinate

The product of Example 12A and (2*S*)-ethyl-2-amino-4-methylpentanoate were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.87 (dd, *J*=9, 6 Hz, 6 H) 1.37 - 1.50 (m, 1 H) 1.51 - 1.65 (m, 2 H) 1.84 (s, 3 H) 2.17 (s, 3 H) 2.21 (s, 3 H) 3.22 - 3.23 (m, 3 H) 3.62 (t, *J*=5 Hz, 2 H) 4.18 - 4.33 (m, 2 H) 4.37 - 4.48 (m, 1 H) 7.88 (d, *J*=9 Hz, 1 H), MS (DCI/NH₃) *m/z* 342 (M+H)⁺. Anal. Calculated for C₁₆H₂₇N₃O₃S: C, 56.26 H, 7.97 N, 12.31. Found C, 56.20 H, 8.04 N, 12.31.

Example 174

1-(1,1-Dimethylpropyl)-3-[(2*Z*)-5-methyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3*H*)-ylidene]urea

The product of Example 157A and 2,2-dimethylpropylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.76 (t, 3 H) 1.12 - 1.17 (m, 1 H) 1.20 (s, 6 H) 1.38 - 1.49 (m, *J*=3 Hz, 3 H) 1.50 - 1.59 (m, 1 H) 1.66 (q, *J*=7 Hz, 2 H) 1.75 - 1.82 (m, 1 H) 2.12 (s, 3 H) 3.22 - 3.30 (m, 1 H) 3.54 - 3.65 (m, 1 H) 3.80 - 3.93 (m, 3 H) 6.19 (s, 1 H) 6.79 (s, 1 H), MS (DCI/NH₃) *m/z* 326 (M+H)⁺. Anal. Calculated for C₁₆H₂₇N₃O₂S: C, 59.04 H, 8.36 N, 12.91. Found C, 59.06 H, 8.36 N, 12.91.

Example 175

1-(1,2-Dimethyl-propyl)-3-[(2*Z*)-5-methyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3*H*)-ylidene]urea hydrochloride

The product of Example 157A and 1,2-dimethylpropylamine were processed according to the method described in Example 169A to afford the title compound. ¹H

NMR (300 MHz, DMSO-*d*₆) δ ppm 0.81 (dd, *J*=7, 3 Hz, 1 H) 0.84 - 0.96 (m, 6 H) 1.06 (d, *J*=7 Hz, 3 H) 1.20 (t, 1 H) 1.34 - 1.54 (m, 3 H) 1.60 - 1.76 (m, 1 H) 1.69 - 1.95 (m, 2 H) 2.28 (s, 3 H) 3.22 - 3.37 (m, 1 H) 3.84 (d, *J*=11 Hz, 2 H) 4.10 - 4.30 (m, 1 H) 4.31 - 4.47 (m, 1 H) 7.23 - 7.41 (m, 1 H) 7.83 - 8.06 (m, 1 H),), MS (DCI/NH₃)
 5 m/z 326 (M+H)⁺. Anal. Calculated for C₁₆H₂₈ClN₃O₂S: C, 53.10 H, 7.80 N, 11.16. Found C, 52.73 H, 7.96 N, 10.82.

Example 176

10 1-Cyclohexyl-3-[(2Z)-4,5-dimethyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea

Example 176A

4,5-Dimethyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylideneamine hydrobromide
 15 A mixture of 2-amino-4,5-dimethylthiazole and 2-(bromomethyl)tetrahydro-2H-pyran were processed using the method described in Example 12A to afford the title compound ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.13 - 1.31 (m, 1 H) 1.36 - 1.52 (m, 3 H) 1.64 - 1.85 (m, 2 H) 2.18 (d, *J*=4 Hz, 6 H) 3.19 - 3.33 (m, 1 H) 3.49 - 3.63 (m, 1 H) 3.77 - 3.89 (m, 1 H) 3.94 - 4.02 (m, 2 H) 9.34 (s, 2 H),), MS
 20 (DCI/NH₃) m/z 227 (M+H)⁺.

Example 176B

1-Cyclohexyl-3-[(2Z)-4,5-dimethyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea
 The product of Example 176A and cyclohexylamine were processed using the
 25 method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.05 - 1.29 (m, 6 H) 1.37 - 1.49 (m, 3 H) 1.51 - 1.63 (m, 2 H) 1.76 (m, 5 H) 2.06 (s, 3 H) 2.11 (s, 3 H) 3.19 - 3.30 (m, 1 H) 3.56 - 3.68 (m, 1 H) 3.70 - 3.88 (m, 3 H) 3.99 (dd, *J*=14, 3 Hz, 1 H) 6.59 (d, *J*=8 Hz, 1 H),), MS (DCI/NH₃) m/z 352 (M+H)⁺. Anal. Calculated for C₁₈H₂₉N₃O₂S•0.1H₂O: C, 61.19 H, 8.33 N, 11.89. Found C, 61.03 H, 8.45 N, 11.69.
 30

Example 177

1-(4-Methylcyclohexyl)-3-[(2Z)-5-methyl-3-(tetrahydropyran-2-ylmethyl)-1,3-

thiazol-2(3H)-ylidene]urea

The product of Example 157A and 4-methylcyclohexylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.88 (dd, *J*=12, 7 Hz, 3 H) 0.96 (d, *J*=3 Hz, 1 H) 1.09 - 1.31 (m, 3 H) 1.31 - 1.70 (m, 9 H) 1.69 - 1.85 (m, 2 H) 2.12 (s, 3 H) 3.21 - 3.29 (m, 1 H) 3.55 - 3.64 (m, *J*=3 Hz, 2 H) 3.79 - 3.95 (m, 3 H) 6.62 (t, *J*=8 Hz, 1 H) 6.77 - 6.81 (m, *J*=1 Hz, 1 H), MS (DCI/NH₃) *m/z* 352 (M+H)⁺.

Example 178

10 1-(1,1-Dimethylpropyl)-3-[(2Z)-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea

Example 178A3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylideneamine hydrobromide

15 A mixture of 2-aminothiazole and 2-(bromomethyl)tetrahydro-2H-pyran were processed using the method described in Example 12A to afford the title compound. MS (DCI/NH₃) *m/z* 199 (M+H)⁺

Example 178B

20 1-(1,1-Dimethylpropyl)-3-[(2Z)-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea

The product of Example 178A and 1,1-dimethylpropylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.77 (t, *J*=7 Hz, 3 H) 1.14 - 1.19 (m, 1 H) 1.21 (s, 6 H) 1.39 - 1.49 (m, 3 H) 1.50 - 1.59 (m, 1 H) 1.61 - 1.72 (m, 2 H) 1.79 (s, 1 H) 3.19 - 3.28 (m, 1 H) 3.56 - 3.68 (m, 1 H) 3.78 - 3.89 (m, 1 H) 3.92 - 4.02 (m, 2 H) 6.23 (s, 1 H) 6.55 (d, *J*=5 Hz, 1 H) 7.08 (d, *J*=5 Hz, 1 H), MS (DCI/NH₃) *m/z* 312 (M+H)⁺. Anal. Calculated for C₁₅H₂₅N₃O₂S: C, 57.85 H, 8.09 N, 13.49 Found C, 58.01 H, 8.23 N, 13.30.

30

Example 179N-[(2Z)-4,5-dimethyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-N-[(1S)-1,2,2-trimethylpropyl]urea

The product of Example 176A and (1S)-1,2,2-trimethylpropylamine were

processed using the method described in Example 169A to afford the title compound.

¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.84 (s, 9 H) 0.98 (d, *J*=7 Hz, 3 H) 1.12 - 1.28 (m, 1 H) 1.40 - 1.49 (m, 3 H) 1.47 - 1.64 (m, *J*=13 Hz, 1 H) 1.73 - 1.84 (m, 1 H) 2.12 (s, 3 H) 3.23 - 3.30 (m, 1 H) 3.50 - 3.68 (m, 2 H) 3.80 - 4.03 (m, 3 H) 6.46 (dd, *J*=9, 3 Hz, 1 H) 6.80 (dd, *J*=5, 2 Hz, 1 H), MS (DCI/NH₃) *m/z* 354 (M+H)⁺. Anal. Calculated for C₁₈H₃₁N₃O₂S, 61.15 H, 8.84 N, 11.69. Found C, 60.80 H, 8.88 N, 11.69.

Example 180

10 1-(2,2-Dimethylpropyl)-3-[(2*Z*)-4,5-dimethyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3*H*)-ylidene]urea

The product of Example 176A and 2,2-dimethylpropylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.84 (s, 9 H) 1.16 - 1.28 (m, 1 H) 1.45 (s, 3 H) 1.56 - 1.68 (m, 1 H) 1.76 - 1.87 (m, 1 H) 2.06 (s, 3 H) 2.11 (s, 3 H) 2.77 - 2.87 (m, 1 H) 2.90 - 3.02 (m, 1 H) 3.15 - 3.28 (m, 1 H) 3.60 - 3.70 (m, 1 H) 3.72 - 3.86 (m, 2 H) 3.94 - 4.11 (m, 1 H) 6.58 - 6.78 (m, 1 H), MS (DCI/NH₃) *m/z* 339 (M+H)⁺. Anal. Calculated for C₁₇H₂₉N₃O₂S: C, 60.14 H, 8.61 N, 12.38. Found C, 60.22 H, 8.71 N, 12.35.

20

Example 181

N-[(2*Z*)-3-(tetrahydro-2*H*-pyran-2-ylmethyl)-1,3-thiazol-2(3*H*)-ylidene]-*N'*-[(1*S*)-1,2,2-trimethylpropyl]urea

The product of Example 178A and (1*S*)-1,2,2-trimethylpropylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.84 (s, 9 H) 0.99 (d, *J*=7 Hz, 3 H) 1.14 - 1.26 (m, *J*=11 Hz, 1 H) 1.36 - 1.49 (m, 3 H) 1.52 - 1.59 (m, 1 H) 1.74 - 1.82 (m, 1 H) 3.20 - 3.28 (m, 1 H) 3.52 - 3.70 (m, 2 H) 3.78 - 4.08 (m, 3 H) 6.46 - 6.60 (m, 2 H) 7.10 (t, *J*=5 Hz, 1 H), MS (DCI/NH₃) *m/z* 326 (M+H)⁺. Anal. Calculated for C₁₆H₂₇N₃O₂S: C, 59.04 H, 8.36 N, 12.91. Found C, 59.08 H, 8.28 N, 12.80.

30

Example 182

N-[(2*Z*)-5-methyl-3-(tetrahydro-2*H*-pyran-2-ylmethyl)-1,3-thiazol-2(3*H*)-ylidene]-*N'*-

[(1S)-1,2,2-trimethylpropyl]urea

The product of Example 157A and (1S)-1,2,2-trimethylpropylamine were processed using the method described in Example 169A to afford the title compound.

¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.84 (s, 9 H) 0.98 (d, *J*=7 Hz, 3 H) 1.11 - 1.30 (m, 1 H) 1.45 (s, 3 H) 1.51 (d, 2 H) 2.12 (s, 4 H) 3.21 - 3.30 (m, 1 H) 3.48 - 3.70 (m, 2 H) 3.78 - 4.05 (m, 3 H) 6.46 (dd, *J*=9, 3 Hz, 1 H) 6.80 (dd, *J*=5, 2 Hz, 1 H), MS (DCI/NH₃) *m/z* 339 (M+H)⁺. Anal. Calculated for C₁₇H₂₉N₃O₂S C, 60.14 H, 8.61 N, 12.38. Found C, 60.10 H, 8.81 N, 12.02.

Example 1831-(4-Methylcyclohexyl)-3-[(2Z)-4,5-dimethyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea

The product of Example 176A and 4-methylcyclohexylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.82 - 1.02 (m, 4 H) 1.15 - 1.29 (m, 3 H) 1.38 - 1.51 (m, 4 H) 1.52 - 1.70 (m, 4 H) 1.78 (s, 3 H) 2.06 (s, 3 H) 2.11 (s, 3 H) 3.19 - 3.28 (m, 1 H) 3.61 (s, 1 H) 3.69 - 3.86 (m, 3 H) 3.98 (dd, *J*=14, 3 Hz, 1 H) 6.57 (d, *J*=8 Hz, 1 H), MS (DCI/NH₃) *m/z* 366 (M+H)⁺. Anal. Calculated for C₁₈H₂₃N₃O₂S: C, 62.45 H, 8.55 N, 11.50. Found C, 62.77 H, 8.86 N, 11.50.

Example 1841-(2,2-Dimethylpropyl)-3-[(2Z)-5-methyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea

The product of Example 157A and 2,2-dimethylpropylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.81 (s, 9 H) 1.09 - 1.26 (m, 1 H) 1.34 - 1.59 (m, 4 H) 1.69 - 1.87 (m, 1 H) 2.12 (d, *J*=1 Hz, 3 H) 2.80 - 2.98 (m, 2 H) 3.08 - 3.27 (m, 1 H) 3.52 - 3.67 (m, 1 H) 3.80 - 4.06 (m, 3 H) 6.68 - 6.76 (m, 1 H) 6.78 - 6.90 (m, 1 H), MS (DCI/NH₃) *m/z* 326 (M+H)⁺. Anal. Calculated for C₁₆H₂₇N₃O₂S: C, 59.04 H, 8.36 N, 12.91. Found C, 58.89 H, 8.56 N, 12.71.

Example 1851-tert-Butyl-3-[(2Z)-4,5-dimethyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-

ylidene]urea

The product of Example 176A and *tert*-butylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.13 - 1.22 (m, 1 H) 1.27 (s, 9 H) 1.44 (d, *J*=4 Hz, 3 H) 1.60 (d, *J*=12 Hz, 1 H) 1.77 (s, 1 H) 2.06 (s, 3 H) 2.11 (s, 3 H) 3.16 - 3.28 (m, 1 H) 3.55 - 3.68 (m, 1 H) 3.68 - 3.87 (m, 2 H) 4.00 (dd, *J*=14, 3 Hz, 1 H) 6.90 (d, *J*=9 Hz, 1 H), MS (DCI/NH₃) *m/z* 326 (M+H)⁺. Anal. Calculated for C₁₆H₂₇N₃O₂S•0.3H₂O: C, 58.02 H, 8.41 N, 12.70. Found C, 58.44 H, 8.12 N, 12.41.

Example 186

1-(1,1-Dimethylpropyl)-3-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea

Example 186A

4,5-Dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylideneamine hydrobromide

A mixture of 2-amino-4,5-dimethylthiazole and 2-(bromomethyl)tetrahydrofuran were processed using the method described in Example 12A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 1.43 - 1.64 (m, 1 H) 1.73 - 2.00 (m, 3 H) 2.17 (s, 3 H) 2.19 (s, 3 H) 3.53 - 3.68 (m, 1 H) 3.71 - 3.85 (m, 1 H) 3.91 - 4.17 (m, 3 H) 9.34 (s, 1 H), MS (DCI/NH₃) *m/z* 212 (M+H)⁺.

Example 186B

1-(1,1-Dimethylpropyl)-3-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea

The product of Example 186A and 1,1-dimethylpropylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.84 (t, *J*=7 Hz, 3 H) 1.22 - 1.31 (m, 6 H) 1.53 - 1.72 (m, 3 H) 1.75 - 1.86 (m, 1 H) 1.91 - 2.05 (m, 1 H) 2.08 - 2.20 (m, 1 H) 2.24 (s, 6 H) 3.55 - 3.71 (m, 1 H) 3.76 - 3.88 (m, 1 H) 4.09 - 4.20 (m, 1 H) 4.30 (s, 1 H) 4.40 - 4.54 (m, 1 H) 6.94 (d, *J*=9 Hz, 1 H), MS (DCI/NH₃) *m/z* 326 (M+H)⁺.

Example 187

1-(2,2-Dimethylpropyl)-3-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea

The product of Example 186A and 2,2-dimethylpropylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.86 (s, 9 H) 1.54 - 1.65 (m, 1 H) 1.77 - 1.97 (m, 3 H) 2.07 (s, 3 H) 2.10 - 2.14 (m, 3 H) 2.78 - 2.97 (m, 2 H) 3.54 - 3.65 (m, 1 H) 3.71 - 3.85 (m, 2 H) 3.98 - 4.11 (m, 1 H) 4.14 - 4.31 (m, 1 H) 6.66 (t, *J*=7 Hz, 1 H), MS (DCI/NH₃) *m/z* 326 (M+H)⁺. Anal. Calculated for C₁₆H₂₇N₃O₂S: C, 59.04 H, 8.36 N, 12.91. Found C, 58.91 H, 8.64 N, 12.77.

10

Example 188

1-[(2Z)-4,5-Dimethyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-3-(3,3,5,5-tetramethylcyclohexyl)urea

The product of Example 176A and 3,3,5,5-tetramethylcyclohexylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.88 (s, 6 H) 0.91 - 1.01 (m, 2 H) 1.01 - 1.05 (m, 6 H) 1.20 (d, *J*=14 Hz, 2 H) 1.37 - 1.62 (m, 7 H) 1.77 (d, *J*=4 Hz, 1 H) 2.04 - 2.09 (m, 3 H) 2.09 - 2.15 (m, 3 H) 3.09 - 3.28 (m, 1 H) 3.55 - 3.87 (m, 4 H) 3.98 (dd, *J*=14, 3 Hz, 1 H) 6.56 (d, *J*=8 Hz, 1 H), MS (DCI/NH₃) *m/z* 408 (M+H)⁺. Anal. Calculated for C₂₂H₃₇N₃O₂S•0.4H₂O: C, 63.7 H, 89.18N, 10.13. Found C, 63.49 H, 8.93 N, 10.12.

20

Example 189

N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-*N'*-[(1*S*)-1,2,2-trimethylpropyl]urea

25

The product of Example 186A and (1*S*)-1,2,2-trimethylpropylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.83 (s, 9 H) 0.98 (dd, *J*=7, 2 Hz, 3 H) 1.53 - 1.64 (m, 1 H) 1.76 - 1.99 (m, 3 H) 2.06 (s, 3 H) 2.11 (s, 3 H) 3.53 - 3.65 (m, 2 H) 3.73 - 3.84 (m, 2 H) 4.05 - 4.14 (m, 1 H) 4.17 - 4.22 (m, 1 H) 6.34 - 6.42 (m, 1 H), MS (DCI/NH₃) *m/z* 339 (M+H)⁺. Anal. Calculated for C₁₇H₂₉N₃O₂S: C, 60.14 H, 8.61 N, 12.38. Found C, 60.06 H, 8.95 N, 12.29.

30

Example 190*N*-[(2*Z*)-4,5-dimethyl-3-(tetrahydro-2*H*-pyran-2-ylmethyl)-1,3-thiazol-2(3*H*)-ylidene]-*N'*-[(1*R*)-1,2,2-trimethylpropyl]urea

The product of Example 176A and (1*R*)-1,2,2-trimethylpropylamine were
 5 processed using the method described in Example 169A to afford the title compound.
¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.84 (s, 9 H) 0.99 (dd, *J*=7, 5 Hz, 3 H) 1.13 -
 1.30 (m, 1 H) 1.38 - 1.50 (m, 3 H) 1.54 - 1.68 (m, *J*=11 Hz, 1 H) 1.74 - 1.85 (m, 1 H)
 2.06 (s, 3 H) 2.11 (s, 3 H) 3.19 - 3.27 (m, 1 H) 3.53 - 3.66 (m, 1 H) 3.71 - 3.88 (m, 3
 H) 3.95 - 4.12 (m, 1 H) 6.37 (d, *J*=9 Hz, 1 H), MS (DCI/NH₃) *m/z* 354 (M+H)⁺.
 10 Anal. Calculated for C₁₈H₃₁N₃O₂S: C, 61.15 H, 9.04 N, 11.89. Found C, 61.36 H,
 9.08 N, 11.80.

Example 191

15 1-*tert*-Butyl-3-[(2*Z*)-5-methyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3*H*)-ylidene]urea

The product of Example 157A and *tert*-butyl amine were processed using the
 method described in Example 169A to afford the title compound. ¹H NMR (300
 MHz, DMSO-*d*₆) δ ppm 1.11 - 1.22 (m, 1 H) 1.27 (s, 9 H) 1.42 - 1.47 (m, *J*=1 Hz, 3
 H) 1.47 - 1.60 (m, 1 H) 1.78 (d, *J*=5 Hz, 1 H) 2.12 (s, 3 H) 3.17 - 3.26 (m, 1 H) 3.55 -
 20 3.66 (m, *J*=6 Hz, 1 H) 3.80 - 3.95 (m, 3 H) 6.34 (s, 1 H) 6.79 (s, 1 H), MS (DCI/NH₃)
m/z 326 (M+H)⁺. Anal. Calculated for C₁₆H₂₇N₃O₂S•0.3H₂O: C, 58.08 H, 8.41 N,
 12.70. Found C, 58.44 H, 8.12 N, 12.41.

Example 192

25 1-(2,3-Dichlorophenyl)-3-[(2*Z*)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3*H*)-ylidene]urea

The product of Example 186A and 2,3-dichlorophenylamine were processed
 using the method described in Example 169A to afford the title compound. ¹H NMR
 (300 MHz, DMSO-*d*₆) δ ppm 1.50 - 1.67 (m, 1 H) 1.75 - 2.06 (m, 3 H) 2.15 (s, 3 H)
 30 2.18 (s, 3 H) 3.57 - 3.69 (m, 1 H) 3.72 - 3.83 (m, 1 H) 3.83 - 3.96 (m, 1 H) 4.10 - 4.30
 (m, 2 H) 7.30 (s, 1 H) 7.32 (d, *J*=1 Hz, 1 H) 7.97 - 8.03 (m, 1 H) 8.22 (s, 1 H), MS
 (DCI/NH₃) *m/z* 400, 401 (M+H)⁺. Anal. Calculated for C₁₃H₂₃Cl₂N₃O₂S•0.3MeOH:
 C, 54.15 H, 8.23 N, 14.20. Found C, 54.47 H, 7.91N, 13.99.

Example 193

1-(2,2-Dimethylpropyl)-3-[(2Z)-3-(2-methoxyethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]urea

5 The product of Example 5A and 2,2-dimethylpropylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.77 (t, *J*=7 Hz, 3 H) 1.21 (s, 6 H) 1.66 (q, *J*=7 Hz, 2 H) 2.21 (s, 3 H) 3.23 (s, 3 H) 3.57 (t, *J*=5 Hz, 2 H) 4.07 (t, *J*=5 Hz, 2 H) 6.21 (s, 1 H) 8.10 (d, *J*=9 Hz, 1 H), MS (DCI/NH₃) *m/z* 286 (M+H)⁺. Anal. Calculated for C₁₃H₂₃N₃O₂S•0.1MeOH: C, 54.70 H, 8.15 N, 14.26. Found C, 54.47 H, 7.91 N, 13.99.

Example 194

N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-N'-[(1R)-1,2,2-trimethylpropyl]urea

15 The product of Example 186A and (1R)-1,2,2-trimethylpropylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.84 (s, 9 H) 0.98 (dd, *J*=7, 2 Hz, 3 H) 1.49 - 1.66 (m, 1 H) 1.74 - 1.99 (m, 3 H) 2.06 (s, 3 H) 2.11 (s, 3 H) 3.52 - 3.67 (m, 2 H) 3.72 - 3.88 (m, 2 H) 4.10 (dt, *J*=14, 4 Hz, 1 H) 4.16 - 4.27 (m, 1 H) 6.37 (dd, *J*=10, 4 Hz, 1 H), *m/z* 339 (M+H)⁺. Anal. Calculated for C₁₇H₂₉N₃O₂S: C, 60.14 H, 8.61 N, 12.38. Found C, 60.18 H, 8.88 N, 12.33.

Example 195

25 N-[(1S)-1,2-dimethylpropyl]-N'-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea

30 The product of Example 186A and (1S)-1,2,2-trimethylpropylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.83 (s, 9 H) 0.98 (dd, *J*=7, 2 Hz, 3 H) 1.52 - 1.66 (m, 1 H) 1.75 - 1.97 (m, 3 H) 2.06 (s, 3 H) 2.11 (s, 3 H) 3.51 - 3.66 (m, 2 H) 3.72 - 3.86 (m, 2 H) 4.10 (dt, *J*=14, 4 Hz, 1 H) 4.19 (s, 1 H) 6.37 (dd, *J*=10, 4 Hz, 1 H), *m/z* 339 (M+H)⁺. Anal. Calculated for C₁₇H₂₉N₃O₂S: C, 60.14 H, 8.61 N, 12.38. Found C, 60.06 H, 8.95 N, 12.29.

Example 196

1-(1-Cyclopropylethyl)-3-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea

5 The product of Example 12A and 1-cyclopropylethylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (300 MHz, DMSO-*d*₆) δ ppm 0.05 - 0.15 (m, 1 H) 0.18 - 0.55 (m, 3 H) 0.78 - 0.92 (m, 1 H) 1.11 (dd, *J*=7, 2 Hz, 3 H) 1.51 - 1.64 (m, 1 H) 1.75 - 1.99 (m, 3 H) 2.07 (s, 3 H) 2.12 (s, 3 H) 3.06 - 3.22 (m, 1 H) 3.61 (dd, 1 H) 3.73 - 3.85 (m, 2 H) 4.02 - 4.14 (m, 1
10 H) 4.15 - 4.25 (m, 1 H) 6.57 - 6.70 (m, 1 H); *m/z* 324 (M+H)⁺. Anal. Calculated for C₁₆H₂₅N₃O₂S: C, 59.41 H, 7.79 N, 12.99. Found C, 59.13 H, 7.78 N, 12.88.

Example 197

15 1-[(2Z)-5-(2,4-Difluorophenyl)-3-(2-methoxyethyl)-1,3-thiazol-2(3H)-ylidene]-3-(1,1-dimethylpropyl)urea

Example 197A

N-[5-chloro-3-(2-methoxyethyl)-1,3-thiazol-2(3H)-ylidene]acetamide

20 A flask was charged with 2-acetamido-5-chlorothiazole (Lancaster, 19.3 g, 110 mmol) in 200 mL of 2:1 THF/DMF. To the solution was added sodium hydride (60% dispersion in mineral oil, 5.44 g, 142 mmol). The mixture was stirred at room temperature for 15 min and then 2-bromoethyl methyl ether (18.3 g, 131 mmol) was added. The reaction mixture was warmed to 85 °C and stirred overnight. After cooling to room temperature, the mixture was diluted with ethyl acetate and washed
25 with water. The organic extract was dried (MgSO₄), filtered, and concentrated. The residue was purified by flash chromatography on SiO₂ using a gradient of 0% to 100 % ethyl acetate: hexane to provide 10.3 g (42%) of the title compound as the more polar regioisomer: ¹H NMR (300 MHz, CDCl₃) δ 2.28 (s, 3 H) 3.35 (s, 3 H) 3.65 - 3.71 (m, 2 H) 4.28 - 4.36 (m, 2 H) 7.00 (s, 1 H); MS (ESI⁺) *m/z* 235 (M+H)⁺.

30

Example 197B

N-[5-(2,4-difluorophenyl)-3-(2-methoxyethyl)-1,3-thiazol-2(3H)-ylidene]acetamide

A flask was charged with the product from Example 197A (10.2 g, 42.6

mmol), 2,6-difluorophenylboronic acid (8.08 g, 51.1 mmol), Na₂CO₃ (64.0 mL of a 2 M aqueous solution, 128 mmol) and PdCl₂(PPh₃)₂ (1.5 g, 2.13 mmol) in 100 mL of DME/H₂O/ethanol (7:3:2). The mixture was warmed to 85 °C and stirred overnight. After cooling to room temperature, the mixture was diluted with ethyl acetate and washed with water. The organic extract was dried (MgSO₄), filtered and concentrated. The residue was purified by flash chromatography on SiO₂ using a gradient of 0% to 100 % ethyl acetate: hexane to provide 11.5 g (86 %) of the title compound: ¹H NMR (300 MHz, CDCl₃) δ 2.17 (s, 3 H) 3.27 (s, 3 H) 3.71 (t, J=5.3 Hz, 2 H) 4.37 (t, J=5.4 Hz, 2 H) 7.17 - 7.24 (m, 1 H) 7.38 - 7.48 (m, 1 H) 7.64 - 7.74 (m, 1 H) 7.88 (s, 1 H); MS (ES I⁺) m/z 313 (M+H)⁺.

Example 197C

N-5-(2,4-difluoro-phenyl)-3-(2-methoxyethyl)-1,3-thiazol-2(3H)-ylideneamine

To a solution of the product from Example 197B (11.5 g, 36.8 mmol) in 100 mL of THF was added 25 mL of 5 N aqueous HCl. The mixture was warmed to 40 °C and stirred overnight. After cooling to room temperature, the solvent was removed under reduced pressure and the residue diluted with ethyl acetate. The mixture was neutralized to pH 7 with saturated aqueous NaHCO₃ and then washed with water. The organic extract was dried (MgSO₄), filtered and concentrated. The residue was purified by flash chromatography on SiO₂ using a gradient of 0% to 100 % ethyl acetate: hexane to provide 8.5 g (85 %) of the title compound: ¹H NMR (300 MHz, DMSO-*d*₆) δ 3.27 (s, 3 H) 3.57 (t, J=5.3 Hz, 2 H) 3.86 (t, J=5.4 Hz, 2 H) 7.06 - 7.14 (m, Hz, 1 H) 7.25 (s, 1 H) 7.29 (dd, J=9.2, 2.7 Hz, 2 H) 7.34 (dd, J=5.9, 3.2 Hz, 1 H) 7.94 (s, 1 H); MS (ES I⁺) m/z 271 (M+H)⁺.

Example 197D

1-[(2Z)-5-(2,4-Difluorophenyl)-3-(2-methoxyethyl)-1,3-thiazol-2-ylidene]-3-(1,1-dimethylpropyl)urea

A mixture of the product of Example 197C and 1,1-dimethylpropylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 0.78 (t, J=7.06 Hz, 3 H) 1.24 (s, 6 H) 1.68 (q, J=7.67 Hz, 2 H) 3.27 (s, 3 H) 3.66 (t, J=5.52 Hz, 2 H) 4.20 (t, J=5.22 Hz, 2 H) 6.49 (m, 1 H) 7.16 (td, J=7.98, 1.84 Hz, 1 H) 7.38 (m, 1 H) 7.58 (m, 1 H) 7.61 (s, 1 H); MS (DCI/NH₃) m/z 384 (M+H)⁺.

Example 1981-[(2Z)-5-(2,4-Difluorophenyl)-3-(2-methoxyethyl)-1,3-thiazol-2-ylidene]-3-(1-methylpropyl)urea

5 A mixture of the product of Example 197C and *sec*-butylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 0.84 (t, *J*=7.32 Hz, 3 H) 1.05 (d, *J*=6.41 Hz, 3 H) 1.42 (m, 2 H) 3.27 (s, 3 H) 3.59 (m, 1 H) 3.66 (t, *J*=5.49 Hz, 2 H) 4.21 (t, *J*=5.80 Hz, 2 H) 6.93 (d, *J*=8.54 Hz, 1 H) 7.18 (td, *J*=8.24, 2.44 Hz, 1 H) 7.39 (m, 1 H) 7.57
10 (td, *J*=8.85, 6.41 Hz, 1 H) 7.62 (brs, 1 H); MS (DCI/NH₃) *m/z* 370 (M+H)⁺.

Example 1991-Cyclopentyl-3-[(2Z)-5-(2,4-difluorophenyl)-3-(2-methoxyethyl)-1,3-thiazol-2(3H)-ylidene]urea

15 A mixture of the product of Example 197C and cyclopentylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (500 MHz, DMSO-*d*₆) δ ppm 1.47 (m, 4 H) 1.65 (m, 2 H) 1.80 (m, 2 H) 3.27 (s, 3 H) 3.66 (t, *J*=5.49 Hz, 2 H) 3.95 (m, 1 H) 4.20 (t, *J*=5.19 Hz, 2 H) 7.10 (d, *J*=7.63 Hz, 1 H) 7.18 (td, *J*=8.24, 2.44 Hz, 1 H) 7.39 (m, 1 H) 7.57 (m, 1 H) 7.63 (s, 1
20 H); MS (DCI/NH₃) *m/z* 382 (M+H)⁺.

Example 2001-[(2Z)-5-(2,4-Difluorophenyl)-3-(2-methoxyethyl)-1,3-thiazol-2(3H)-ylidene]-3-(4-methylcyclohexyl)urea

25 A mixture of the product of Example 197C and 4-methylcyclohexylamine were processed using the method described in Example 169A to afford the title compound. ¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 0.90 (m, 3 H) 0.98 (m, 2 H) 1.49 (m, 8 H) 3.27 (s, 3 H) 3.66 (m, 2 H) 4.21 (m, 2 H) 6.93 (t, *J*=8.90 Hz, 1 H) 7.17 (td, *J*=7.06, 2.45 Hz, 1 H) 7.38 (m, 1 H) 7.56 (m, 1 H) 7.62 (m, 1 H); MS (DCI/NH₃) *m/z*
30 410 (M+H)⁺.

Example 202N-[(2Z)-5-*tert*-butyl-3-(2-methoxyethyl)-1,3-thiazol-2(3H)-ylidene]-2-ethoxybenzamide

Example 202A5-tert-butylthiazol-2-amine

To a flask equipped with a Dean-Stark trap was added 3,3-dimethylbutanal
5 (Aldrich, 5.0 g, 50 mmol), pyrrolidine (Aldrich, 4.4 mL, 52 mmol) and p-
toluenesulfonic acid monohydrate (10 mg) in cyclohexane (70 mL). The mixture was
heated to reflux for 3 hours, the water was removed and the organic phase was
concentrated under reduced pressure. The residue was dissolved in methanol (20 mL)
and cooled to 0 °C. Sulfur (Aldrich, 1.6 g, 50 mmol) and a solution of cyanamide
10 (Aldrich, 2.1 g, 50 mmol) in methanol (5 mL) were added. The reaction mixture was
allowed to warm to ambient temperature, stirred for 12 hours, and was concentrated
under reduced pressure. The residue was purified by column chromatography (SiO₂,
2% methanol in CH₂Cl₂) to afford the title compound. MS (ESI⁺) m/z 157 (M+H)⁺.

Example 202B5-tert-butyl-3-(2-methoxyethyl)thiazol-2(3H)-imine hydrobromide

A mixture of Example 202A and commercially available 2-bromoethyl methyl
ether (Aldrich) were processed according to the method described in Example 12A to
afford the title compound. MS (ESI⁺) m/z 215 (M+H)⁺.

Example 202C

20 N-[(2Z)-5-tert-butyl-3-(2-methoxyethyl)-1,3-thiazol-2(3H)-ylidene]-2-
ethoxybenzamide

Commercially available 2-ethoxybenzoic acid (Aldrich) and Example 202B
were processed using the method described in Example 58 to afford the title
compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.28 - 1.36 (m, 12 H),
25 3.26 (s, 3 H), 3.71 (t, J=5.4 Hz, 2 H), 4.06 (q, J=6.9 Hz, 2 H), 4.31 (t, J=5.4 Hz, 2 H),
6.95 (td, J=7.4, 0.8 Hz, 1 H), 7.05 (dd, J=8.5, 0.7 Hz, 1 H), 7.21 (s, 1 H), 7.32 - 7.42
(m, 1 H), 7.67 (dd, J=7.6, 1.9 Hz, 1 H); MS (ESI⁺) m/z 363 (M+H)⁺.

Example 203

30 2-ethoxy-N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-4-ylmethyl)-1,3-thiazol-2(3H)-
ylidene]benzamide

Example 203A(tetrahydro-2H-pyran-4-yl)methyl 4-methylbenzenesulfonate

To a solution of tetrahydro-2H-pyran-4-ylmethanol (Combi-Blocks, 2.0 g,

17.2 mmol) in 10 mL of CH_2Cl_2 and 10 mL of pyridine was added *p*-toluenesulfonyl chloride (3.5 g, 18.1 mmol) in portions over 15 minutes. The mixture stirred at ambient temperature for 16 hours and was quenched with 10 mL of saturated, aqueous NaHCO_3 . The layers were separated and the aqueous layer was
5 extracted with three 10 mL portions of CH_2Cl_2 . The combined organic extracts were dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure to afford the title compound. ^1H NMR (300 MHz, dimethylsulfoxide- d_6) δ ppm 1.05 - 1.25 (m, 2 H), 1.40 - 1.53 (m, 2 H), 1.73 - 1.94 (m, 1 H), 2.43 (s, 3 H), 3.14 - 3.28 (m, 2 H), 3.71 - 3.84 (m, 2 H), 3.88 (d, $J=6.4$ Hz, 2 H), 7.48 (d, $J=8.5$ Hz, 2 H), 7.79 (d,
10 $J=8.5$ Hz, 2 H); MS (DCI/ NH_3) m/z 288 ($\text{M}+\text{NH}_4$) $^+$.

Example 203B

5-methyl-3-((tetrahydro-2H-pyran-4-yl)methyl)thiazol-2(3H)-imine

A mixture of Example 203A (1.9 g, 7.0 mmol), 2-amino-5-methylthiazole (0.80 g, 7.0 mmol) and tetrabutylammonium iodide (1.3 g, 3.5 mmol) in 3 mL of *N,N*-
15 dimethylformamide was warmed to 85 °C and was allowed to stir for 24 hours. The mixture was diluted with 10 mL of CH_2Cl_2 , washed with 10% aqueous NaHCO_3 , dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. Purification via column chromatography (SiO_2 , 10% methanol in ethyl acetate then 9:1:0.1 CH_2Cl_2 : methanol : NH_4OH) afforded the title compound. MS (DCI/ NH_3)
20 m/z 213 ($\text{M}+\text{H}$) $^+$.

Example 203C

2-ethoxy-N-[(2Z)-5-methyl-3-((tetrahydro-2H-pyran-4-yl)methyl)-1,3-thiazol-2(3H)-ylidene]benzamide

To a solution of Example 203B (0.11 g, 0.52 mmol) in 10 mL of tetrahydrofuran and 1 mL of *N,N*- dimethylformamide at ambient temperature was
25 added triethylamine (0.22 mL, 1.6 mmol) followed by 2-ethoxybenzoyl chloride (0.11 g, 0.57 mmol). This mixture was warmed to 50 °C, stirred for 3 hours, was quenched with 10 mL of saturated aqueous NH_4Cl , and diluted with 10 mL of CH_2Cl_2 . The layers were separated and the aqueous layer was extracted with three 5 mL portions of
30 CH_2Cl_2 . The combined organic extracts were dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. Purification via flash column chromatography (SiO_2 , 50% hexanes in ethyl acetate) afforded the title compound. ^1H NMR (300 MHz, CDCl_3) δ ppm 1.46 (t, $J=7.0$ Hz, 3 H), 1.52 - 1.61 (m, 1 H), 1.57

(s, 3 H), 2.14 - 2.26 (m, 1 H), 2.29 (d, $J=1.4$ Hz, 3 H), 3.36 (dt, $J=11.7, 2.4$ Hz, 2 H), 3.98 (ddd, $J=11.4, 4.1, 1.5$ Hz, 2 H), 4.06 (d, $J=7.5$ Hz, 2 H), 4.17 (q, $J=6.8$ Hz, 2 H), 6.59 - 6.62 (m, 1 H), 6.93 - 7.01 (m, 2 H), 7.36 (ddd, $J=8.3, 7.5, 1.9$ Hz, 1 H), 7.97 (dd, $J=8.0, 1.9$ Hz, 1 H); MS (DCI/ NH_3) m/z 361 ($\text{M}+\text{H}$)⁺. Anal. Calculated for $\text{C}_{19}\text{H}_{24}\text{N}_2\text{O}_3\text{S}$: C, 63.31; H, 6.71; N, 7.77. Found: C, 63.27; H, 6.57; N, 7.48.

Example 204

2,4-dimethoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

Example 204A

10 5-methyl-3-((tetrahydrofuran-2-yl)methyl)thiazol-2(3H)-imine

A mixture of 2-amino-5-methylthiazole (1 g, 8.7 mmol) and 2-(bromomethyl)tetrahydrofuran (1.1 mL, 10 mmol) was warmed to 85 °C and was allowed to stir for 24 hours. The mixture was cooled to ambient temperature and purified via flash column chromatography (SiO_2 , 10% methanol in ethyl acetate then
15 9:1:0.1 CH_2Cl_2 : methanol : NH_4OH) to afford the title compound. MS (DCI/ NH_3) m/z 199 ($\text{M}+\text{H}$)⁺.

Example 204B

2,4-dimethoxybenzoyl chloride

A solution of 2,4-dimethoxybenzoic acid (0.25 g, 1.4 mmol) in 5 mL of SOCl_2
20 was warmed to reflux and was allowed to stir for 2 hours. The mixture was cooled to ambient temperature and concentrated under reduced pressure. The crude material was diluted with 5 mL of toluene and concentrated under reduced pressure. This dilution with toluene and concentration was repeated two additional times to give the crude title compound which was used without additional purification or
25 characterization.

Example 204C

2,4-dimethoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

To a solution of Example 204A (0.18 g, 0.91 mmol) in 10 mL of
30 tetrahydrofuran at ambient temperature was added triethylamine (0.38 mL, 2.7 mmol) followed by Example 204B (1.4 mmol) in 3 mL of tetrahydrofuran via cannula. This mixture was warmed to 50 °C, stirred for 3 hours, then quenched with 10 mL of saturated aqueous NH_4Cl , and diluted with 10 mL of CH_2Cl_2 . The layers were separated and the aqueous phase was extracted three 5 mL portions of CH_2Cl_2 . The

combined organic extracts were dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. Purification via column chromatography (SiO_2 , 50% hexanes in ethyl acetate) afforded the title compound. ^1H NMR (300 MHz, CD_3OD) δ ppm 1.65 - 1.79 (m, 1 H), 1.84 - 1.96 (m, 2 H), 1.99 - 2.15 (m, 1 H), 2.31 (d, $J=1.4$ Hz, 3 H), 3.71 - 3.81 (m, 1 H), 3.84 - 3.93 (m, 1 H), 3.85 (s, 3 H), 3.86 (s, 3 H), 4.19 - 4.42 (m, 3 H), 6.51 - 6.62 (m, 2 H), 7.05 - 7.10 (m, 1 H), 8.03 (d, $J=8.8$ Hz, 1 H); MS (DCI/ NH_3) m/z 363 ($\text{M}+\text{H}$) $^+$. Anal. Calculated for $\text{C}_{18}\text{H}_{22}\text{N}_2\text{O}_4\text{S}$: C, 59.65; H, 6.12; N, 7.73. Found: C, 59.47; H, 6.01; N, 7.62.

Example 205

10 5-chloro-2-methoxy-N-[(2Z)-4-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

Example 205A

4-methyl-3-((tetrahydro-2H-pyran-2-yl)methyl)thiazol-2(3H)-imine

A mixture of 2-amino-4-methylthiazole (1.0 g, 8.8 mmol) and 2-(bromomethyl)tetrahydropyran (1.1 mL, 8.8 mmol) was warmed to 85 °C and was allowed to stir for 24 hours. The mixture was cooled to ambient temperature and the crude material was purified via column chromatography (SiO_2 , 10% methanol in ethyl acetate then 9:1:0.1 CH_2Cl_2 : methanol : NH_4OH) to afford the title compound. MS (DCI/ NH_3) m/z 213 ($\text{M}+\text{H}$) $^+$.

Example 205B

5-chloro-2-methoxybenzoyl chloride

A solution of 2-methoxy-5-chlorobenzoic acid (0.37 g, 2.0 mmol) in 10 mL of SOCl_2 was warmed to reflux and was allowed to stir for 2 hours. The mixture was cooled to ambient temperature and concentrated under reduced pressure. The crude material was diluted with 5 mL of toluene and concentrated under reduced pressure. This dilution with toluene and concentration was repeated two additional times to afford the title compound. ^1H NMR (300 MHz, dimethylsulfoxide- d_6) δ ppm 3.82 (s, 3 H), 7.16 (d, $J=8.8$ Hz, 1 H), 7.49 - 7.59 (m, 1 H), 7.61 (d, $J=2.7$ Hz, 1 H).

Example 205C

30 5-chloro-2-methoxy-N-[(2Z)-4-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

To a solution of Example 205A (0.21 g, 1.0 mmol) in 10 mL of tetrahydrofuran at ambient temperature was added triethylamine (0.41 mL, 3.0 mmol)

followed by Example 205B (2.0 mmol) in 5 mL of tetrahydrofuran via cannula. This mixture was warmed to 50 °C, stirred for 2 hours, then quenched with 10 mL of saturated aqueous NH₄Cl, and diluted with 10 mL of CH₂Cl₂. The layers were separated and the aqueous phase was extracted with three 5 mL portions of CH₂Cl₂.

- 5 The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 50% hexanes in ethyl acetate) afforded the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.31 - 1.48 (m, 1 H), 1.47 - 1.67 (m, 3 H), 1.70 - 1.82 (m, 1 H), 1.84 - 1.98 (m, 1 H), 2.40 (d, *J*=1.0 Hz, 3 H), 3.26 - 3.40 (m, 1 H), 3.82 - 3.94 (m, 2 H), 3.87
10 (s, 3 H), 4.00 - 4.11 (m, 1 H), 4.42 (dd, *J*=13.9, 2.7 Hz, 1 H), 6.55 (d, *J*=1.0 Hz, 1 H), 7.07 (d, *J*=8.8 Hz, 1 H), 7.40 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.87 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 381 (M+H)⁺. Anal. Calculated for C₁₈H₂₁ClN₂O₃S: C, 56.76; H, 5.56; N, 7.35. Found: C, 56.58; H, 5.43; N, 7.19.

Example 206

- 15 5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-4-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

- To Example 203B (0.11 g, 0.52 mmol) and triethylamine (0.22 mL, 1.6 mmol) in 10 mL of tetrahydrofuran and 1 mL of N,N-dimethylformamide was added Example 205B (0.68 mmol) in 2 mL of tetrahydrofuran. This mixture was warmed to
20 50 °C, stirred for 2 hours, then quenched with 10 mL of saturated aqueous NH₄Cl, and diluted with 10 mL of CH₂Cl₂. The layers were separated and the aqueous phase was extracted with three 5 mL portions of CH₂Cl₂. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via flash column chromatography (SiO₂, 50% hexanes in ethyl acetate)
25 resulted in the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.34 - 1.60 (m, 4 H), 2.18 - 2.32 (m, 1 H), 2.34 (d, *J*=1.4 Hz, 3 H), 3.38 (dt, *J*=11.6, 2.5 Hz, 2 H), 3.86 (s, 3 H), 3.94 (ddd, *J*=11.6, 4.2, 1.9 Hz, 2 H), 4.14 (d, *J*=7.5 Hz, 2 H), 7.07 (d, *J*=8.8 Hz, 1 H), 7.12 (q, *J*=1.4 Hz, 1 H), 7.40 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.83 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 381 (M+H)⁺. Anal. Calculated for C₁₈H₂₁ClN₂O₃S: C, 56.76;
30 H, 5.56; N, 7.35. Found: C, 56.48; H, 5.46; N, 7.23.

Example 207

5-chloro-2-methoxy-N-[(2Z)-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

Example 207A3-((tetrahydro-2H-pyran-2-yl)methyl)thiazol-2(3H)-imine

A mixture of 2-aminothiazole (1.0 g, 10 mmol) and 2-(bromomethyl)tetrahydro-2H-pyran (1.3 mL, 10 mmol) was warmed to 85 °C and was
5 allowed to stir for 24 hours. The mixture was cooled to ambient temperature and the crude material was purified via column chromatography (SiO₂, 10% methanol in ethyl acetate then 9:1:0.1 CH₂Cl₂ : methanol : NH₄OH) to afford the title compound. MS (DCI/NH₃) m/z 199 (M+H)⁺.

Example 207B

10 5-chloro-2-methoxy-N-[(2Z)-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

To a solution of Example 207A (0.19 g, 0.96 mmol) in 10 mL of tetrahydrofuran at ambient temperature was added triethylamine (0.40 mL, 2.9 mmol) followed by Example 205B (2.0 mmol) in 5 mL of tetrahydrofuran via cannula. This
15 mixture was warmed to 50 °C, stirred for 2 hours, then quenched with 10 mL of saturated aqueous NH₄Cl, and diluted with 10 mL of CH₂Cl₂. The layers were separated and the aqueous phase was extracted with three 5 mL portions of CH₂Cl₂. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂,
20 50% hexanes in ethyl acetate) afforded the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.20 - 1.41 (m, 1 H), 1.46 - 1.63 (m, 3 H), 1.67 - 1.77 (m, 1 H), 1.82 - 1.96 (m, 1 H), 3.35 - 3.45 (m, 1 H), 3.75 - 3.85 (m, 1 H), 3.86 (s, 3 H), 3.89 - 4.00 (m, 1 H), 4.19 - 4.27 (m, 1 H), 4.38 - 4.46 (m, 1 H), 6.91 (d, J=4.7 Hz, 1 H), 7.08 (d, J=8.8 Hz, 1 H), 7.38 (d, J=4.7 Hz, 1 H), 7.41 (dd, J=8.8, 3.1 Hz, 1 H), 7.85 (d, J=3.1
25 Hz, 1 H); MS (DCI/NH₃) m/z 367 (M+H)⁺. Anal. Calculated for C₁₇H₁₉ClN₂O₃S: C, 55.66; H, 5.22; N, 7.64. Found: C, 55.72; H, 5.08; N, 7.55.

Example 208

5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

30

Example 208A(R)-(tetrahydrofuran-2-yl)methyl 4-methylbenzenesulfonate

To a solution of (R)-tetrahydrofurfuryl alcohol (Lancaster, 1.0 g, 9.8 mmol) in 5 mL of CH₂Cl₂ and 5 mL of pyridine was added *p*-toluenesulfonyl chloride (2.8 g,

14.7 mmol) in portions over 15 minutes. The mixture was stirred at ambient temperature for 3 hours and was quenched with 10 mL of saturated, aqueous NaHCO₃. The layers were separated and the aqueous layer was extracted with three 5 mL portions of CH₂Cl₂. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford the title compound. MS (DCI/NH₃) m/z 257 (M+H)⁺, 274 (M+NH₄)⁺.

Example 208B

(R)-5-methyl-3-((tetrahydrofuran-2-yl)methyl)thiazol-2(3H)-imine

A mixture of Example 208A (1.5 g, 5.9 mmol), 2-amino-5-methylthiazole (0.68 g, 5.9 mmol) and tetrabutylammonium iodide (1.1 g, 3.0 mmol) in 3 mL of N,N-dimethylformamide was warmed to 85 °C and was allowed to stir for 48 hours. The mixture was diluted with 10 mL of CH₂Cl₂ and the solution was quenched with 10 mL of saturated, aqueous NaHCO₃. The layers were separated and the aqueous layer was extracted twice with 10 mL portions of CH₂Cl₂. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 10% methanol in ethyl acetate then 9:1:0.1 CH₂Cl₂ : methanol : NH₄OH) afforded the title compound. MS (DCI/NH₃) m/z 199 (M+H)⁺.

Example 208C

5-chloro-2-methoxybenzoyl chloride

A solution of 2-methoxy-5-chlorobenzoic acid (0.22 g, 1.2 mmol) in 10 mL of SOCl₂ was warmed to reflux and was allowed to stir for 2 hours. The mixture was cooled to ambient temperature and concentrated under reduced pressure. The crude material was diluted with 5 mL of toluene and concentrated under reduced pressure. This dilution with toluene and concentration was repeated two additional times to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 3.82 (s, 3 H), 7.16 (d, J=8.8 Hz, 1 H), 7.49 - 7.59 (m, 1 H), 7.61 (d, J=2.7 Hz, 1 H).

Example 208D

5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

To a solution of Example 208B (0.23 g, 1.2 mmol) in 10 mL of tetrahydrofuran at ambient temperature was added triethylamine (0.49 mL, 3.5 mmol) followed by Example 208C (1.2 mmol) in 5 mL of tetrahydrofuran via cannula. This

mixture was warmed to 50 °C and was allowed to stir for 3 hours and was quenched with 10 mL of NH₄Cl and diluted with 10 mL of CH₂Cl₂. The layers were separated and the aqueous layer was extracted with three 5 mL portions of CH₂Cl₂. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via flash column chromatography (SiO₂, 1 : 1 : 1 hexanes : ethyl acetate : CH₂Cl₂) afforded the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.64 - 1.79 (m, 1 H), 1.84 - 1.96 (m, 2 H), 2.00 - 2.14 (m, 1 H), 2.34 (d, *J*=1.4 Hz, 3 H), 3.69 - 3.81 (m, 1 H), 3.84 - 3.93 (m, 1 H), 3.85 (s, 3 H), 4.20 - 4.43 (m, 3 H), 7.07 (d, *J*=8.8 Hz, 1 H), 7.14 (q, *J*=1.1 Hz, 1 H), 7.39 (dd, *J*=9.0, 2.9 Hz, 1 H), 7.81 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 367 (M+H)⁺. Anal. Calculated for C₁₇H₁₉ClN₂O₃S: C, 55.66; H, 5.22; N, 7.64. Found: C, 55.42; H, 5.08; N, 7.58.

Example 209

5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-[(2S)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

Example 209A

(S)-(tetrahydrofuran-2-yl)methyl 4-methylbenzenesulfonate

To a solution of (S)-tetrahydrofurfuryl alcohol (Codexis, 1.6 g, 15.2 mmol) in 5 mL of CH₂Cl₂ and 5 mL of pyridine was added *p*-toluenesulfonyl chloride (4.3 g, 22.8 mmol) in portions over 15 minutes. The mixture stirred at ambient temperature for 3 hours and was quenched with 10 mL of saturated, aqueous NaHCO₃. The layers were separated and the aqueous layer was extracted with three 5 mL portions of CH₂Cl₂. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford the title compound. MS (DCI/NH₃) *m/z* 257 (M+H)⁺, 274 (M+NH₄)⁺.

Example 209B

(S)-5-methyl-3-((tetrahydrofuran-2-yl)methyl)thiazol-2(3H)-imine

A mixture of Example 209A (1.6 g, 6.1 mmol), 2-amino-5-methylthiazole (0.7 g, 6.1 mmol) and tetrabutylammonium iodide (2.3 g, 6.1 mmol) in 5 mL of N,N-dimethylformamide was warmed to 85 °C and was allowed to stir for 18 hours. The mixture was diluted with 10 mL of CH₂Cl₂, washed with 10% aqueous NaHCO₃, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 10% methanol in ethyl acetate then

9:1:0.1 CH₂Cl₂ : methanol : NH₄OH) afforded the title compound. MS (DCI/NH₃) m/z 199 (M+H)⁺.

Example 209C

5 5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-[(2S)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

Example 209B (0.32 g, 1.6 mmol), triethylamine (0.67 mL, 4.8 mmol) and Example 205B (1.9 mmol) in 20 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.64 - 1.79 (m, 1 H), 1.84 - 1.96 (m, 2 H), 2.00 - 2.13 (m, 1 H), 2.34 (d, J=1.4 Hz, 3 H), 3.71 - 3.81 (m, 1 H), 3.84 - 3.93 (m, 1 H), 3.85 (s, 3 H), 4.20 - 4.43 (m, 3 H), 7.07 (d, J=8.8 Hz, 1 H), 7.14 (q, J=1.4 Hz, 1 H), 7.39 (dd, J=9.0, 2.9 Hz, 1 H), 7.81 (d, J=2.7 Hz, 1 H); MS (DCI/NH₃) m/z 367 (M+H)⁺. Anal. Calculated for C₁₇H₁₉ClN₂O₃S: C, 55.66; H, 5.22; N, 7.64. Found: C, 55.48; H, 4.96; N, 7.52.

Example 210

15 2,2,3,3-tetrafluoro-1-methyl-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]cyclobutanecarboxamide

Example 204A (0.20 g, 1.0 mmol), triethylamine (0.42 mL, 3.0 mmol) and 2,2,3,3-tetrafluoro-1-(methyl)cyclobutanecarbonyl chloride (ABCR, 0.27 g, 1.3 mmol) in 15 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.53 (s, 3 H), 1.62 - 1.73 (m, 1 H), 1.81 - 1.93 (m, 2 H), 1.95 - 2.10 (m, 1 H), 2.27 - 2.44 (m, 1 H), 2.32 (d, J=1.4 Hz, 3 H), 3.33 - 3.43 (m, 1 H), 3.69 - 3.79 (m, 1 H), 3.80 - 3.90 (m, 1 H), 4.20 - 4.36 (m, 3 H), 7.11 (dd, J=1.4, 0.7 Hz, 1 H); MS (DCI/NH₃) m/z 367 (M+H)⁺. Anal. Calculated for C₁₅H₁₈F₄N₂O₂S: C, 49.17; H, 4.95; N, 7.65. Found: C, 49.27; H, 4.88; N, 7.58.

25 Example 211

5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(oxetan-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

Example 211A

oxetan-2-ylmethyl 4-methylbenzenesulfonate

30 To a solution of 2-hydroxymethyloxetane (TCI-US, 2.0 g, 23 mmol) in 10 mL of CH₂Cl₂ and 10 mL of pyridine was added *p*-toluenesulfonyl chloride (6.5 g, 34 mmol) in portions over 15 minutes. The mixture was stirred at ambient temperature for 3 hours and was quenched with 10 mL of saturated, aqueous NaHCO₃. The layers

were separated and the aqueous layer was extracted with three 5 mL portions of CH₂Cl₂. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 70% hexanes in ethyl acetate) afforded the title compound. MS (DCI/NH₃)
 5 m/z 243 (M+H)⁺, 260 (M+NH₄)⁺.

Example 211B

5-methyl-3-(oxetan-2-ylmethyl)thiazol-2(3H)-imine

A mixture of Example 211A (1.1 g, 4.6 mmol), 2-amino-5-methylthiazole (0.53 g, 4.6 mmol) and tetrabutylammonium iodide (0.85 g, 2.3 mmol) in 5 mL of N,N-dimethylformamide was warmed to 85 °C and was allowed to stir for 18 hours.
 10 The mixture was diluted with 10 mL of CH₂Cl₂, washed with 10% aqueous NaHCO₃, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 10% methanol in ethyl acetate then 9:1:0.1 CH₂Cl₂ : methanol : NH₄OH) afforded the title compound. MS (DCI/NH₃)
 15 m/z 185 (M+H)⁺.

Example 211C

5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(oxetan-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

Example 211B (0.26 g, 1.4 mmol), triethylamine (0.59 mL, 4.2 mmol) and
 20 Example 205B (1.7 mmol) in 15 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 2.35 (d, J=1.4 Hz, 3 H), 2.43 - 2.59 (m, 1 H), 2.70 - 2.85 (m, 1 H), 3.85 (s, 3 H), 4.37 - 4.51 (m, 2 H), 4.57 - 4.71 (m, 2 H), 5.15 - 5.25 (m, 1 H), 7.06 (d, J=9.2 Hz, 1 H), 7.18 (q, J=1.1 Hz, 1 H), 7.39 (dd, J=8.8, 2.7 Hz, 1 H), 7.79 (d, J=2.7 Hz, 1 H); MS (DCI/NH₃)
 25 m/z 353 (M+H)⁺. Anal. Calculated for C₁₆H₁₇ClN₂O₃S: C, 54.46; H, 4.86; N, 7.94. Found: C, 54.41; H, 4.88; N, 7.80.

Example 212

5-chloro-N-[(2Z)-3-(1,3-dioxolan-2-ylmethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

30 Example 212A

3-((1,3-dioxolan-2-yl)methyl)-5-methylthiazol-2(3H)-imine

A mixture of 2-amino-5-methylthiazole (1 g, 8.7 mmol) and 2-bromomethyl-1,3-dioxolane (0.98 mL, 9.6 mmol) was warmed to 85 °C and was allowed to stir for

18 hours. The mixture was cooled to ambient temperature and purified via column chromatography (SiO₂, 10% methanol in ethyl acetate then 9:1:0.1 CH₂Cl₂ : methanol : NH₄OH) to afford the title compound. MS (DCI/NH₃) m/z 201 (M+H)⁺.

Example 212B

5 5-chloro-N-[(2Z)-3-(1,3-dioxolan-2-ylmethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 212A (0.25 g, 1.3 mmol), triethylamine (0.52 mL, 3.8 mmol) and Example 205B (1.5 mmol) in 15 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 2.33 (d, J=1.4 Hz, 3 H), 3.86 (s, 3 H), 3.87 - 3.99 (m, 4 H), 4.41 (d, J=4.1 Hz, 2 H), 5.27 (t, J=4.1 Hz, 1 H), 7.07 (d, J=8.8 Hz, 1 H), 7.12 (q, J=1.4 Hz, 1 H), 7.40 (dd, J=8.8, 2.7 Hz, 1 H), 7.87 (d, J=2.7 Hz, 1 H); MS (DCI/NH₃) m/z 369 (M+H)⁺. Anal. Calculated for C₁₆H₁₇ClN₂O₄S: C, 52.10; H, 4.65; N, 7.60. Found: C, 52.15; H, 4.42; N, 7.44.

Example 213

15 5-chloro-N-[(2Z)-3-[2-(1,3-dioxolan-2-yl)ethyl]-5-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 213A

3-(2-(1,3-dioxolan-2-yl)ethyl)-5-methylthiazol-2(3H)-imine

A mixture of 2-amino-5-methylthiazole (1.0 g, 8.7 mmol) and 2-(2-bromoethyl)-1,3-dioxolane (1.1 mL, 8.7 mmol) was warmed to 85 °C and was allowed to stir for 18 hours. The mixture was cooled to ambient temperature and purified via column chromatography (SiO₂, 10% methanol in ethyl acetate then 9:1:0.1 CH₂Cl₂ : methanol : NH₄OH) to afford the title compound. MS (DCI/NH₃) m/z 215 (M+H)⁺.

25 Example 213B

5-chloro-N-[(2Z)-3-[2-(1,3-dioxolan-2-yl)ethyl]-5-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 213A (0.25 g, 1.2 mmol), triethylamine (0.49 mL, 3.5 mmol) and Example 205B (1.3 mmol) in 10 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 2.18 - 2.25 (m, 2 H), 2.33 (d, J=1.0 Hz, 3 H), 3.82 - 3.87 (m, 2 H), 3.86 (s, 3 H), 3.93 - 4.01 (m, 2 H), 4.36 (dd, J=7.1 Hz, 2 H), 4.93 (t, J=4.4 Hz, 1 H), 7.07 (d, J=8.8 Hz, 1 H), 7.10 (q, J=1.4 Hz, 1 H), 7.40 (dd, J=8.8, 2.7 Hz, 1 H), 7.96 (d, J=3.1 Hz, 1 H); MS (DCI/NH₃)

m/z 383 (M+H)⁺. Anal. Calculated for C₁₇H₁₉ClN₂O₄S: C, 53.33; H, 5.00; N, 7.32. Found: C, 53.02; H, 4.52; N, 7.22.

Example 214

N-[(2Z)-3-(1,3-dioxolan-2-ylmethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]-2-ethoxybenzamide

Example 212A (0.20 g, 1.0 mmol), triethylamine (0.42 mL, 3.0 mmol) and 2-ethoxybenzoyl chloride (0.17 g, 1.1 mmol) in 10 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.41 (t, J=7.0 Hz, 3 H), 2.33 (d, J=1.4 Hz, 3 H), 3.82 - 4.01 (m, 4 H), 4.12 (q, J=6.9 Hz, 2 H), 4.40 (d, J=4.4 Hz, 2 H), 5.27 (t, J=4.2 Hz, 1 H), 6.96 (dt, J=7.5, 0.8 Hz, 1 H), 7.05 (d, J=8.5 Hz, 1 H), 7.09 (q, J=1.4 Hz, 1 H), 7.39 (ddd, J=8.7, 6.9, 1.7 Hz, 1 H), 7.83 (dd, J=7.6, 1.9 Hz, 1 H); MS (DCI/NH₃) m/z 349 (M+H)⁺. Anal. Calculated for C₁₇H₂₀N₂O₄S: C, 58.60; H, 5.79; N, 8.04. Found: C, 58.22; H, 5.32; N, 7.93.

Example 215

5-bromo-2-ethoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

Example 215A

5-bromo-2-ethoxybenzoic acid

To a solution of 2-ethoxybenzoic acid (3.3 g, 20.0 mmol) in 75 mL of acetonitrile at 0 °C was added N-bromosuccinimide (3.7 g, 21 mmol) in 15 mL of acetonitrile. The reaction mixture was warmed to ambient temperature and the mixture was allowed to stir for 48 hours. The mixture was quenched with 20 mL of H₂O and the layers were separated. The aqueous layer was extracted with three 15 mL portions of CH₂Cl₂ and the combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 10% hexanes in ethyl acetate) afforded the title compound. MS (DCI/NH₃) m/z 262, 264 (M+NH₄)⁺.

Example 215B

5-bromo-2-ethoxybenzoyl chloride

A solution of Example 215A (0.21 g, 0.86 mmol) in 5 mL of SOCl₂ was warmed to reflux and was allowed to stir for 2 hours. The mixture was cooled to ambient temperature and concentrated under reduced pressure. The crude material

was diluted with 5 mL of toluene and concentrated under reduced pressure. This dilution with toluene and concentration was repeated two additional times to give the crude title compound which was used without additional purification or characterization.

5

Example 215C

5-bromo-2-ethoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

Example 208B (0.17 g, 0.86 mmol), triethylamine (0.36 mL, 2.6 mmol) and Example 215B (0.86 mmol) in 10 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.39 (t, *J*=7.0 Hz, 3 H), 1.63 - 1.78 (m, 1 H), 1.84 - 1.96 (m, 2 H), 2.00 - 2.15 (m, 1 H), 2.34 (d, *J*=1.4 Hz, 3 H), 3.69 - 3.81 (m, 1 H), 3.84 - 3.95 (m, 1 H), 4.10 (q, *J*=7.0 Hz, 2 H), 4.19 - 4.43 (m, 3 H), 6.99 (d, *J*=8.8 Hz, 1 H), 7.14 (q, *J*=1.2 Hz, 1 H), 7.49 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.87 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 425, 427 (M+H)⁺. Anal. Calculated for C₁₈H₂₁BrN₂O₃S: C, 50.83; H, 4.98; N, 6.59. Found: C, 50.89; H, 4.87; N, 6.51.

15

Example 216

5-chloro-2-ethoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

20

Example 216A

5-chloro-2-ethoxybenzoic acid

To a solution of 2-ethoxybenzoic acid (4.4 g, 26.6 mmol) in 80 mL of acetonitrile at 0 °C was added *N*-chlorosuccinimide (3.7 g, 28 mmol) in 20 mL of acetonitrile dropwise over 30 minutes. The reaction mixture was warmed to ambient temperature and the mixture was allowed to stir for 70 hours. The mixture was quenched with 20 mL of H₂O and the layers were separated. The aqueous layer was extracted with three 15 mL portions of CH₂Cl₂ and the combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via recrystallization with ether and hexanes afforded the title compound. MS (DCI/NH₃) *m/z* 201 (M+H)⁺, 218 (M+NH₄)⁺.

30

Example 216B

5-chloro-2-ethoxybenzoyl chloride

A solution of Example 216A (0.25 g, 1.0 mmol) in 5 mL of SOCl₂ was

warmed to reflux and was allowed to stir for 2 hours. The mixture was cooled to ambient temperature and concentrated under reduced pressure. The crude material was diluted with 5 mL of toluene and concentrated under reduced pressure. This dilution with toluene and concentration was repeated two additional times to give the crude title compound which was used without additional purification or characterization.

Example 216C

5-chloro-2-ethoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

Example 208B (0.20 g, 1.0 mmol), triethylamine (0.42 mL, 3.0 mmol) and Example 216B (1.0 mmol) in 10 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.39 (t, J=7.0 Hz, 3 H), 1.62 - 1.77 (m, 1 H), 1.83 - 1.97 (m, 2 H), 1.99 - 2.14 (m, 1 H), 2.34 (d, J=1.4 Hz, 3 H), 3.70 - 3.80 (m, 1 H), 3.84 - 3.93 (m, 1 H), 4.10 (q, J=6.9 Hz, 2 H), 4.20 - 4.44 (m, 3 H), 7.04 (d, J=8.8 Hz, 1 H), 7.14 (q, J=1.4 Hz, 1 H), 7.35 (dd, J=8.8, 2.7 Hz, 1 H), 7.73 (d, J=2.7 Hz, 1 H); MS (DCI/NH₃) m/z 381 (M+H)⁺. Anal. Calculated for C₁₈H₂₁ClN₂O₃S: C, 56.76; H, 5.56; N, 7.35. Found: C, 56.36; H, 5.28; N, 7.25.

Example 217

4-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

Example 217A

4-chloro-2-methoxybenzoyl chloride

A solution of the 4-chloro-2-methoxybenzoic acid (0.24 g, 1.3 mmol) in 7 mL of SOCl₂ was warmed to reflux and was allowed to stir for 2 hours. The mixture was cooled to ambient temperature and concentrated under reduced pressure. The crude material was diluted with 5 mL of toluene and concentrated under reduced pressure. This dilution with toluene and concentration was repeated two additional times to give the crude title compound which was used without additional purification or characterization.

Example 217B

4-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

Example 204A (0.20 g, 1.0 mmol), triethylamine (0.42 mL, 3.0 mmol) and Example 217A (1.3 mmol) in 15 mL of tetrahydrofuran and 1 mL of N,N-dimethylformamide were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CDCl₃) δ ppm 1.61 - 1.74 (m, 1 H), 1.77 - 1.98 (m, 2 H), 1.99 - 2.14 (m, 1 H), 2.30 (d, *J*=1.4 Hz, 3 H), 3.72 - 3.82 (m, 1 H), 3.82 - 3.90 (m, 1 H), 3.91 (s, 3 H), 4.11 - 4.20 (m, 1 H), 4.27 (ddd, *J*=13.7, 6.8, 2.9 Hz, 1 H), 4.41 - 4.51 (m, 1 H), 6.87 - 6.91 (m, 1 H), 6.93 - 6.99 (m, 2 H), 7.97 (d, *J*=8.8 Hz, 1 H); MS (DCI/NH₃) *m/z* 367 (M+H)⁺. Anal. Calculated for C₁₇H₁₉ClN₂O₃S: C, 55.66; H, 5.22; N, 7.64. Found: C, 55.40; H, 5.31; N, 7.48.

10

Example 218

5-chloro-2-methoxy-N-[(2Z)-1-(2-methoxyethyl)-1,4,6,7-tetrahydro-2H-pyrano[4,3-d][1,3]thiazol-2-ylidene]benzamide

Example 218A

6,7-dihydro-4H-pyrano[4,3-*d*][1,3]thiazol-2-amine

15

To a solution of tetrahydro-4H-pyran-4-one (Aldrich) (7.22 g, 72.11 mmol) in cyclohexane (70 mL) were added pyrrolidine (6.26 mL, 7.57 mmol) and *p*-toluenesulfonic acid monohydrate (13.72 mg, 0.07 mmol). The reaction mixture was refluxed for 3 hours with a Dean-Stark trap, cooled and concentrated. The residue was dissolved in methanol (20 mL) and then sulfur (2.31 g, 72.11 mmol) was added followed by a solution of cyanamide (3.03 g, 72.11 mmol) in methanol (5 mL) at 0 °C. The reaction mixture was stirred at room temperature overnight, filtered, concentrated and purified by column chromatography using an Analogix® Intelliflash280TM (SiO₂, 0-5% methanol in dichloromethane) to afford the title compound. MS (ESI⁺) *m/z* 157 (M+H)⁺.

25

Example 218B

1-(2-methoxyethyl)-6,7-dihydro-1H-pyrano[4,3-*d*]thiazol-2(4H)-imine hydrobromide

A mixture of product of Example 218A (1.0 g, 6.4 mmol) and 2-bromoethyl methyl ether (3.0 mL, 32.0 mmol) was processed according to the method of Example 2A to afford the title compound: MS (LC/MS) *m/z* 213 (M+H)⁺.

30

Example 218C

5-chloro-2-methoxy-N-[(2Z)-1-(2-methoxyethyl)-1,4,6,7-tetrahydro-2H-pyrano[4,3-*d*][1,3]thiazol-2-ylidene]benzamide

To a solution of Example 218B (150.0 mg, 0.51 mmol) in tetrahydrofuran (10

mL) was added *N*-(3-dimethylaminopropyl)-*N*-ethylcarbodiimide hydrochloride (97.0 mg, 0.51 mmol), 1-hydroxybenzotriazole (69.0 mg, 0.51 mmol), triethylamine (178.0 μ L, 1.28 mmol), and 5-chloro-2-methoxybenzoic acid (Aldrich) (95.0 mg, 0.51 mmol). The mixture was stirred overnight at 80 °C, and then diluted with ethyl acetate, washed with 1 M aqueous NaHCO₃, dried (Na₂SO₄), filtered and concentrated. Purification by preparative HPLC on a Waters Symmetry C8 column (40 mm X 100 mm, 7 μ m particle size) using a gradient of 10 % to 100 % acetonitrile: ammonium acetate (10 mM) over 15 minutes at a flow rate of 70 mL/minutes afforded the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-*d*₆) δ 2.75 (t, *J*=5.4 Hz, 2 H), 3.25 (s, 3 H), 3.69 (t, *J*=5.3 Hz, 2 H), 3.80 (s, 3 H), 3.95 (t, *J*=5.4 Hz, 1 H), 4.27 (t, *J*=5.3 Hz, 2 H), 4.58 (s, 2 H), 7.12 (d, *J*=8.8 Hz, 1 H), 7.46 (dd, *J*=9.0, 2.9 Hz, 1 H), 7.69 (d, *J*=2.7 Hz, 1 H); MS (ESI⁺) *m/z* 383 (M+H)⁺. Anal. Calculated for C₁₇H₁₉ClN₂O₄S: C, 53.33; H, 5.00; N, 7.32. Found: C, 53.21; H, 4.80; N, 7.27.

Example 219

15 5-bromo-2-methoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

Example 219A

5-bromo-2-methoxybenzoic acid

To a solution of 2-methoxybenzoic acid (6 g, 39.4 mmol) in 80 mL of acetonitrile was added *N*-bromosuccinimide (7.4 g, 41.4 mmol) in 20 mL of acetonitrile. The reaction mixture was warmed to ambient temperature and the mixture was allowed to stir for 16 hours. Additional *N*-bromosuccinimide (14.8 g, 82.8 mmol) was added and the reaction mixture stirred for an additional 48 hours. The mixture was quenched with 25 mL of H₂O and the layers were separated. The aqueous layer was extracted with three 15 mL of portions of CH₂Cl₂ and the combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 50% hexanes in ethyl acetate) afforded the title compound. MS (DCI/NH₃) *m/z* 248, 250 (M+NH₄)⁺.

30

Example 219B

5-bromo-2-methoxybenzoyl chloride

A solution of Example 219A (0.28 g, 1.4 mmol) in 5 mL of SOCl₂ was warmed to reflux and was allowed to stir for 2 hours. The mixture was cooled to

ambient temperature and concentrated under reduced pressure. The crude material was diluted with 5 mL of toluene and concentrated under reduced pressure. This dilution with toluene and concentration was repeated two additional times to give the crude title compound which was used without additional purification or
5 characterization.

Example 219C

5-bromo-2-methoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

Example 208B (0.25 g, 1.3 mmol), triethylamine (0.53 mL, 3.8 mmol) and
10 Example 219B (1.4 mmol) in 15 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.65 - 1.80 (m, 1 H), 1.85 - 1.96 (m, 2 H), 2.01 - 2.14 (m, 1 H), 2.34 (d, J=1.4 Hz, 3 H), 3.71 - 3.80 (m, 1 H), 3.85 (s, 3 H), 3.85 - 3.93 (m, 1 H), 4.21 - 4.41 (m, 3 H), 7.02 (d, J=9.2 Hz, 1 H), 7.14 (q, J=1.1 Hz, 1 H), 7.53 (dd, J=8.8, 2.7 Hz, 1 H), 7.95 (d, J=2.4 Hz, 1
15 H); MS (DCI/NH₃) m/z 411, 413 (M+H)⁺. Anal. Calculated for C₁₇H₁₉BrN₂O₃S: C, 49.64; H, 4.66; N, 6.81. Found: C, 49.48; H, 4.53; N, 6.72.

Example 220

5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(2-tetrahydro-2H-pyran-4-ylethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

Example 220A

2-(tetrahydro-2H-pyran-4-yl)ethyl 4-methylbenzenesulfonate

To a solution of 2-(tetrahydropyran-4-yl)-ethanol (1.5 g, 11.5 mmol) in 10 mL of CH₂Cl₂ and 7 mL of pyridine was added *p*-toluenesulfonyl chloride (2.4 g, 12.7 mmol) portion wise over 15 minutes. The mixture stirred at ambient temperature for
25 3 hours and was quenched with 10 mL of saturated, aqueous NaHCO₃. The layers were separated and the aqueous layer was extracted with three 5 mL of portions of CH₂Cl₂. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 70% hexanes in ethyl acetate) afforded the title compound. MS (DCI/NH₃)
30 m/z 302 (M+NH₄)⁺.

Example 220B

5-methyl-3-(2-(tetrahydro-2H-pyran-4-yl)ethyl)thiazol-2(3H)-imine

A mixture of Example 220A (1.9 g, 6.7 mmol), 2-amino-5-methylthiazole

(0.77 g, 6.7 mmol) and tetrabutylammonium iodide (1.1 g, 3.3 mmol) in 2 mL of N,N-dimethylformamide was warmed to 85 °C and was allowed to stir for 18 hours. The mixture was diluted with 10 mL of CH₂Cl₂, washed with 10 mL of 10% aqueous NaHCO₃, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 10% methanol in ethyl acetate then 9:1:0.1 CH₂Cl₂ : methanol : NH₄OH) afforded the title compound. MS (DCI/NH₃) m/z 227 (M+H)⁺.

Example 220C

5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(2-tetrahydro-2H-pyran-4-ylethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

Example 220B (0.20 g, 0.9 mmol), triethylamine (0.37 mL, 0.26 mmol) and Example 205B (0.9 mmol) in 10 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.26 - 1.44 (m, 2 H), 1.47 - 1.65 (m, 1 H), 1.71 - 1.89 (m, 4 H), 2.34 (d, J=1.4 Hz, 3 H), 3.32 - 3.41 (m, 2 H), 3.86 (s, 3 H), 3.86 - 3.94 (m, 2 H), 4.26 - 4.35 (m, 2 H), 7.07 (d, J=8.8 Hz, 1 H), 7.14 (q, J=1.4 Hz, 1 H), 7.40 (dd, J=9.0, 2.9 Hz, 1 H), 7.89 (d, J=3.1 Hz, 1 H); MS (DCI/NH₃) m/z 395 (M+H)⁺. Anal. Calculated for C₁₉H₂₃ClN₂O₃S: C, 57.79; H, 5.87; N, 7.09. Found: C, 57.54; H, 5.67; N, 7.07.

Example 221

5-chloro-N-[(2Z)-5-ethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 221A

5-ethyl-3-((tetrahydrofuran-2-yl)methyl)thiazol-2(3H)-imine

A mixture of 5-ethylthiazol-2-amine and 2-(bromomethyl)tetrahydrofuran were processed using the method described in Example 2A to afford the title compound. MS (ESI) m/z 213 (M+H)⁺.

Example 221B

5-chloro-N-[(2Z)-5-ethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 221A and 5-chloro-2-methoxybenzoic acid were processed using the method described in Example 2B to afford the title compound. ¹H NMR (500 MHz, dimethylsulfoxide-*d*₆) δ ppm 1.21 (t, J=7.63 Hz, 3 H) 1.59 - 1.68 (m, 1 H) 1.77 - 1.85 (m, 2 H) 1.89 - 1.97 (m, 1 H) 2.67 (dd, J=15.56, 7.63 Hz, 2 H) 3.65 (dd, J=14.95, 7.02

Hz, 1 H) 3.75 - 3.82 (m, 1 H) 3.78 (s, 3 H) 4.13 - 4.24 (m, 2 H) 4.24 - 4.30 (m, 1 H) 7.10 (d, $J=8.85$ Hz, 1 H) 7.26 (t, $J=1.22$ Hz, 1 H) 7.44 (dd, $J=8.85, 2.75$ Hz, 1 H) 7.64 (d, $J=2.75$ Hz, 1 H); MS (ESI) m/z 381(M+H)⁺.

Example 222

5 5-chloro-2-methoxy-N-[(2Z)-5-propyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

Example 222A

5-propyl-3-((tetrahydrofuran-2-yl)methyl)thiazol-2(3H)-imine

10 A mixture of 5-propylthiazol-2-amine and 2-(bromomethyl)tetrahydrofuran were processed using the method described in Example 2A to afford the title compound. MS (ESI) m/z 227 (M+H)⁺.

Example 222B

5-chloro-2-methoxy-N-[(2Z)-5-propyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

15 Example 222A and 5-chloro-2-methoxybenzoic acid were processed using the method described in Example 2B to afford the title compound. ¹H NMR (500 MHz, dimethylsulfoxide-*d*₆) δ ppm 0.93 (t, $J=7.32$ Hz, 3 H) 1.56 - 1.67 (m, 3 H) 1.77 - 1.85 (m, 2 H) 1.93 (dt, $J=19.22, 7.02$ Hz, 1 H) 2.62 (t, $J=7.02$ Hz, 2 H) 3.65 (dd, $J=14.95, 6.71$ Hz, 1 H) 3.74 - 3.80 (m, 1 H) 3.77 - 3.79 (m, 3 H) 4.15 - 4.24 (m, 2 H) 4.24 - 4.30 (m, 1 H) 7.10 (d, $J=8.85$ Hz, 1 H) 7.25 - 7.28 (m, 1 H) 7.44 (dd, $J=8.85, 2.75$ Hz, 1 H) 7.64 (d, $J=2.75$ Hz, 1 H); MS (ESI) m/z 395(M+H)⁺.

Example 223

5-chloro-N-[(2Z)-5-chloro-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

25 Example 223A

5-chloro-N-(5-chlorothiazol-2-yl)-2-methoxybenzamide

30 A mixture of 5-chlorothiazol-2-amine hydrochloride (513 mg, 3 mmol), 5-chloro-2-methoxybenzoic acid (670 mg, 3.6mmol), 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (1.15 g, 6 mmol), 1-hydroxybenzotriazole hydrate (810 mg, 6 mmol) and 4-(dimethylamino)pyridine (73 mg, 0.6 mmol) in pyridine was stirred at room temperature for 2 hours. The volatiles were removed under vacuum, and the resulting mixture was diluted with water and extracted with ethyl acetate. The organic extract was dried (Na₂SO₄), filtered and

concentrated. The residue was washed with a small amount of ethyl acetate, and filtered to afford the title compound. MS (ESI) m/z 303 ($M+H$)⁺.

Example 223B

5-chloro-N-[(2Z)-5-chloro-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

5

Example 223A (250 mg, 0.83 mmol) in tetrahydrofuran/N,N-dimethylformamide (1:2) (9 mL) was treated with NaH (60%) (40 mg, 1.0 mmol) for 10 minutes then 2-(bromomethyl)tetrahydrofuran (164 mg, 1.0 mmol) was added. The mixture was heated at 150 °C for 2 hours. After cooling to ambient temperature, the mixture was diluted with water and extracted with ethyl acetate. The organic extract was dried (Na₂SO₄), filtered and concentrated. Purification by reverse phase HPLC afforded the title compound. ¹H NMR (400 MHz, dimethylsulfoxide-*d*₆) δ ppm 1.60 - 1.70 (m, 1 H) 1.79 - 1.89 (m, 2 H) 1.92 - 2.01 (m, 1 H) 3.66 (dd, *J*=15.04, 7.06 Hz, 1 H) 3.76 - 3.83 (m, 1 H) 3.79 - 3.81 (m, 3 H) 4.15 - 4.23 (m, 1 H) 4.24 - 4.33 (m, 2 H) 7.14 (d, *J*=8.90 Hz, 1 H) 7.49 (dd, *J*=8.90, 2.76 Hz, 1 H) 7.74 (d, *J*=3.07 Hz, 1 H) 7.77 (s, 1 H) MS (ESI) m/z 387 ($M+H$)⁺. 81887-154-2

10

15

Example 224

4,5-dichloro-2-methoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

20

Example 224A

4,5-dichloro-2-methoxybenzoic acid

25

To a solution of 4-chloro-2-methoxybenzoic acid (5 g, 26.8 mmol) in 200 mL of acetonitrile was added *N*-chlorosuccinimide (17.9 g, 134 mmol). The mixture was allowed to stir for 72 hours at ambient temperature and was quenched with 50 mL of H₂O. The layers were separated and the aqueous layer was extracted with three 25 mL portions of CH₂Cl₂. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 25% hexanes in ethyl acetate) afforded the title compound. MS (DCI/NH₃) m/z 238 ($M+NH_4$)⁺.

30

Example 224B

4,5-dichloro-2-methoxybenzoyl chloride

A solution of Example 224A (0.18 g, 0.81 mmol) in 5 mL of SOCl₂ was warmed to reflux and was allowed to stir for 2 hours. The mixture was cooled to

ambient temperature and concentrated under reduced pressure. The crude material was diluted with 5 mL of toluene and concentrated under reduced pressure. This dilution with toluene and concentration was repeated two additional times to afford the title compound which was used without additional purification or characterization.

5

Example 224C

4,5-dichloro-2-methoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

Example 208B (0.16 g, 0.81 mmol), triethylamine (0.34 mL, 2.4 mmol) and Example 224B (0.81 mmol) in 10 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.64 - 1.79 (m, 1 H), 1.83 - 1.96 (m, 2 H), 2.03 - 2.15 (m, 1 H), 2.34 (d, *J*=1.4 Hz, 3 H), 3.70 - 3.80 (m, 1 H), 3.83 - 3.93 (m, 1 H), 3.87 (s, 3 H), 4.20 - 4.44 (m, 3 H), 7.15 (q, *J*=1.1 Hz, 1 H), 7.26 (s, 1 H), 7.99 (s, 1 H); MS (DCI/NH₃) *m/z* 401 (M+H)⁺. Anal. Calculated for C₁₇H₁₈Cl₂N₂O₃S: C, 50.88; H, 4.52; N, 6.98. Found: C, 50.63; H, 4.41; N, 6.83.

15

Example 225

5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

Example 225A

20 (tetrahydro-2H-pyran-3-yl)methyl 4-methylbenzenesulfonate

To a solution of (tetrahydropyran-3-yl)-methanol (Matrix, 1.67 g, 14.4 mmol) in 15 mL of CH₂Cl₂ and 15 mL of pyridine was added *p*-toluenesulfonyl chloride (2.9 g, 15.1 mmol) in portions over 10 minutes. The mixture stirred at ambient temperature for 18 hours and was quenched with 10 mL of saturated, aqueous NaHCO₃. The layers were separated and the aqueous phase was extracted three 5 mL of portions of CH₂Cl₂. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 70% hexanes in ethyl acetate) afforded the title compound. MS (DCI/NH₃) *m/z* 288 (M+NH₄)⁺.

25

Example 225B

5-methyl-3-((tetrahydro-2H-pyran-3-yl)methyl)thiazol-2(3H)-imine

A mixture of Example 225A (1.0 g, 3.7 mmol), 2-amino-5-methylthiazole (0.42 g, 3.7 mmol) and tetrabutylammonium iodide (0.68 g, 1.85 mmol) in 1 mL of

N,N- dimethylformamide was warmed to 85 °C and was allowed to stir for 16 hours. The mixture was diluted with 10 mL of CH₂Cl₂, washed with 10 mL of 10% aqueous NaHCO₃, dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 10% methanol in ethyl acetate then 9:1:0.1 CH₂Cl₂ : methanol : NH₄OH) afforded the title compound. MS (DCI/NH₃) m/z 213 (M+H)⁺.

Example 225C

5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide

Example 225B (0.19 g, 0.89 mmol), triethylamine (0.29 mL, 2.1 mmol) and Example 205B (0.63 mmol) in 10 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.33 - 1.49 (m, 1 H), 1.50 - 1.67 (m, 1 H), 1.69 - 1.86 (m, 2 H), 2.20 - 2.32 (m, 1 H), 2.34 (d, J=1.4 Hz, 3 H), 3.32 - 3.38 (m, 1 H), 3.50 (ddd, J=11.5, 9.5, 3.1 Hz, 1 H), 3.72 - 3.82 (m, 2 H), 3.86 (s, 3 H), 4.17 (d, J=7.5 Hz, 2 H), 7.07 (d, J=8.8 Hz, 1 H), 7.11 (q, J=1.4 Hz, 1 H), 7.40 (dd, J=8.8, 2.7 Hz, 1 H), 7.85 (d, J=2.7 Hz, 1 H); MS (DCI/NH₃) m/z 381 (M+H)⁺. Anal. Calculated for C₁₈H₂₁ClN₂O₃S: C, 56.76; H, 5.56; N, 7.35. Found: C, 56.84; H, 5.32; N, 7.29.

Example 226

2-chloro-N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]nicotinamide

A mixture of Example 186A (0.15 g), 2-chloronicotinic acid (99 mg), 1-hydroxybenzotriazole hydrate (80 mg), 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (120 mg) and triethylamine (0.15 mL) in N,N-dimethylformamide was stirred overnight at room temperature, poured into water and extracted with ether (3x). The combined organic extracts were dried (Na₂SO₄), filtered and the solvent evaporated. The crude material was purified by gradient flash chromatography over silica gel eluting with ethyl acetate:hexane (1:4 to 1:1) to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.65 (m, 1H), 1.85 (m, 2H), 2.0 (m, 1H), 2.25 (s, 3H), 2.28 (s, 3H), 3.62 (dd, 1H), 3.77 (dd, 1H), 4.08 (dd, 1H), 4.28 (m, 1H), 4.38 (dd, 1H), 7.50 (dd, 1H), 8.24 (dd, 1H), 8.45 (dd, 1H); MS (ESI+) m/z 352 (M+H)⁺.

Example 227

5-chloro-N-[(2Z)-4,5-dimethyl-3-[(3-methyl-4,5-dihydroisoxazol-5-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 227A

3-allyl-4,5-dimethylthiazol-2(3H)-imine hydrobromide

- 5 A mixture of 4,5-dimethyl-1,3-thiazol-2-amine (1 g) and allylbromide (0.95 g) in toluene (5 mL) was heated to 85 °C for 12 hours, cooled, diluted with ether, filtered and the solvent was evaporated to afford crude product that was taken on to the next step without further characterization.

Example 227B

- 10 (Z)-N-(3-allyl-4,5-dimethylthiazol-2(3H)-ylidene)-5-chloro-2-methoxybenzamide

- A mixture of Example 227A (1.3 g), Example 205B (1.36 g) and triethylamine (1.0 g) in tetrahydrofuran (40 mL) was heated to 60 °C for 4 hours, cooled and solvent was evaporated. The crude material was triturated with ether, filtered and solvent evaporated. The crude was flash chromatographed over silica gel gradient eluting
15 with ethyl acetate:hexane (2:3 to 3:2) to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 2.22 (s, 3H), 2.23 (s, 3H), 3.78 (s, 3H), 4.85 (m, 2H), 4.96, (dq, J = 17.3, 1.3 Hz, 1H), 5.19 (dq, J = 10.5, 1.3 Hz, 1H), 5.92-6.05 (m, 1H), 7.10 (d, 1H), 7.44 (dd, 1H), 7.66 (d, 1H); MS (ESI+) m/z 337 (M+H)⁺.

Example 227C

- 20 5-chloro-N-[(2Z)-4,5-dimethyl-3-[(3-methyl-4,5-dihydroisoxazol-5-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

- To a solution of acetaldoxime (56 mg, 1.48 mmol) in CHCl₃ (10 mL) under N₂ was added N-chlorosuccinimide (200 mg) and pyridine (10 μL). After 4.5 hours at room temperature, Example 227B (100 mg) was added, followed by triethylamine
25 (0.15 g) and the reaction continued to stir at room temperature for 21 hours. The reaction mixture was washed with water and partitioned. The aqueous layer was extracted again with CH₂Cl₂ and the combined organic extracts were dried (MgSO₄), filtered and solvent evaporated. The crude was purified by flash chromatography over silica gel eluting with ethyl acetate:hexane (1:1) to afford the title compound. ¹H
30 NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.93 (s, 3H), 2.23 (s, 6H), 2.90 (dd, 1H), 3.14 (dd, 1H), 3.79 (s, 3H), 4.11 (dd, 1H), 4.27, (dd, 1H), 4.98 (m, 1H), 7.11 (d, 1H), 7.44 (dd, 1H), 7.67 (d, 1H); MS (ESI+) m/z 394 (M+H)⁺.

Example 228

N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-4-(trifluoromethyl)nicotinamide

Example 186A (0.15 g) and 4-(trifluoromethyl)nicotinic acid (0.12 g) were processed according to the method of Example 226. The crude was purified by reverse phase preparative HPLC on a Waters Symmetry C8 column (25 mm x 100 mm, 7 μ m particle size) using a gradient of 10% to 100% acetonitrile:0.1% aqueous trifluoroacetic acid over 8 minutes (10 minutes run time) at a flow rate of 40 mL/minutes) to afford the title compound. ^1H NMR (300 MHz, dimethylsulfoxide- d_6) δ ppm 1.63 (m, 1H), 1.85 (m, 2H), 1.96 (m, 1H), 2.25 (s, 3H), 2.29 (s, 3H), 3.62 (dd, 1H), 3.77 (dd, 1H), 4.07 (dd, 1H), 4.25 (m, 1H), 4.38 (dd, 1H), 7.80 (d, 1H), 8.89 (d, 1H), 9.12 (s, 1H); MS (ESI+) m/z 386 (M+H) $^+$

Example 229

N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-ethoxynicotinamide

Example 186A (0.15 g) and 2-ethoxynicotinic acid (0.1 g) were processed according to the method of Example 226. The crude was purified by reverse phase preparative HPLC on a Waters Symmetry C8 column (25 mm x 100 mm, 7 μ m particle size) using a gradient of 10% to 100% acetonitrile:0.1% aqueous trifluoroacetic acid over 8 minutes (10 minutes run time) at a flow rate of 40 mL/minutes) to afford the title compound. ^1H NMR (300 MHz, dimethylsulfoxide- d_6) δ ppm 1.32 (t, 3H), 1.66 (m, 1H), 1.85 (m, 2H), 1.97 (m, 1H), 2.22 (s, 3H), 2.26 (s, 3H), 3.62 (dd, 1H), 3.78 (dd, 1H), 4.06 (dd, 1H), 4.3-4.42 (m, 3H), 7.02 (dd, 1H), 8.13 (dd, 1H), 8.22 (dd, 1H); MS (ESI+) m/z 362 (M+H) $^+$

Example 230

N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2,3,6-trifluoroisonicotinamide

Example 186A (0.15 g) and 2,3,6-trifluoroisonicotinic acid (0.11 g) were processed according to the method of Example 226. The crude was purified by reverse phase preparative HPLC on a Waters Symmetry C8 column (25 mm x 100 mm, 7 μ m particle size) using a gradient of 10% to 100% acetonitrile:0.1% aqueous trifluoroacetic acid over 8 minutes (10 minutes run time) at a flow rate of 40 mL/minutes) to afford the title compound. ^1H NMR (300 MHz, dimethylsulfoxide- d_6) δ ppm 1.67 (m, 1H), 1.86 (m, 2H), 2.00 (m, 1H), 2.27 (s, 3H), 2.30 (s, 3H), 3.62

(dd, 1H), 3.78 (dd, 1H), 4.15 (dd, 1H), 4.28 (m, 1H), 4.43 (dd, 1H), 7.57 (t, 1H); MS (ESI+) m/z 372 (M+H)⁺

Example 231

6-chloro-N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-4-(trifluoromethyl)nicotinamide

5

Example 186A (0.15 g) and 6-chloro-4-(trifluoromethyl)nicotinic acid (0.17 g) were processed according to the method of Example 226. The crude was purified by reverse phase preparative HPLC on a Waters Symmetry C8 column (25 mm x 100 mm, 7 µm particle size) using a gradient of 10% to 100% acetonitrile:0.1% aqueous trifluoroacetic acid over 8 minutes (10 minutes run time) at a flow rate of 40 mL/minutes) to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.62 (m, 1H), 1.85 (m, 2H), 1.96 (m, 1H), 2.25 (s, 3H), 2.29 (s, 3H), 3.62 (dd, 1H), 3.77 (dd, 1H), 4.08 (dd, 1H), 4.24 (m, 1H), 4.38 (dd, 1H), 7.99 (s, 1H), 9.00 (s, 1H); MS (ESI+) m/z 420 (M+H)⁺

15

Example 232

5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-[(3-methyl-4,5-dihydroisoxazol-5-yl)methyl]-1,3-thiazol-2(3H)-ylidene]benzamide

Example 232A

3-allyl-5-methylthiazol-2(3H)-imine hydrobromide

20

A mixture of 2-amino-5-methylthiazole (2.5 g) and allylbromide (3.31 g) was processed according to the method of Example 227A to afford the title compound that was taken directly to the next step. MS (ESI+) m/z 155 (M+H)⁺.

Example 232B

(Z)-N-(3-allyl-5-methylthiazol-2(3H)-ylidene)-5-chloro-2-methoxybenzamide

25

A mixture of Example 232A (0.5 g) and 5-chloro-2-methoxybenzoylchloride (0.57 g) were processed according to the method of Example 227B to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 2.28 (d, J = 1.4 Hz, 3H), 3.78 (s, 3H), 4.77 (d, 2H), 5.13 (dq, J = 16.9, 1.3 Hz, 1H), 5.24 (dq, J = 10.5, 1.4 Hz, 2H), 5.94-6.07 (m, 1H), 7.11 (d, 1H), 7.23 (q, J = 1.4 Hz, 1H), 7.43 (dd, 1H), 7.68 (d, 1H); MS (ESI+) m/z 323 (M+H)⁺

30

Example 232C

5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-[(3-methyl-4,5-dihydroisoxazol-5-yl)methyl]-1,3-thiazol-2(3H)-ylidene]benzamide

Example 232B was processed according to the method of Example 227C to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.90 (s, 3H), 2.28 (s, 3H), 2.87 (dd, 1H), 3.11 (dd, 1H), 4.23 (d, 2H), 4.94 (m, 1H), 7.11 (d, 1H), 7.25 (s, 1H), 7.45 (dd, 1H), 7.67 (d, 1H); MS (ESI+) m/z 380 (M+H)⁺

5

Example 233

N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-(ethylamino)benzamide

Example 186A (0.15 g) and 2-ethylaminobenzoic acid (Pellon *Syn. Lett.* **2005**, 10, 1606) (0.1 g) were processed according to the method of Example 226. The crude
10 was purified by flash chromatography over silica gel gradient eluting with ethyl acetate:hexane (1:9 to 1:3) to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.25 (t, 3H), 1.70 (m, 1H), 1.87 (m, 2H), 2.01 (m, 1H), 2.20 (s, 3H), 2.26 (s, 3H), 3.20 (m, 2H), 3.62 (dd, 1H), 3.79 (dd, 1H), 4.12 (dd, 1H), 4.28 (m, 1H), 4.38 (dd, 1H), 6.55 (t, 1H), 6.67 (d, 1H), 7.27 (t, 1H), 8.22 (dd, 1H),
15 8.52 (t, 1H); MS (ESI+) m/z 360 (M+H)⁺.

Example 234

N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-iodo-2-(methylamino)benzamide

Example 186A (0.15 g) and 5-iodo-2-methylaminobenzoic acid (0.17 g) were
20 processed according to the method of Example 226. The crude was purified by flash chromatography over silica gel gradient eluting with ethyl acetate:hexane (1:19 to 1:10) to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.7-1.82 (m, 1H), 1.84-1.95 (m, 2H), 1.95-2.08 (m, 1H), 2.22-2.29 (s, 3H), 2.27 (s, 3H), 2.84 (d, 3H), 3.65 (dd, 1H), 3.82 (dd, 1H), 4.12 (dd, 1H), 4.28 (m, 1H), 4.36 (dd, 1H),
25 6.52 (d, 1H), 7.53 (dd, 1H), 8.46 (d, 1H), 8.52 (q, 1H); MS (ESI+) m/z 472 (M+H)⁺.

Example 235

5-bromo-N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

To a mixture of 3,3-dimethylbutyraldehyde (Aldrich) (5 mL, 39.8 mmol) and
30 Example 278A (641.0 mg, 4.0 mmol) was added a mixture of dimethylsulfoxide (560 μL, 8 mmol) and 12 N aqueous HCl (667 μL, 8 mmol). The reaction mixture was heated at 40 °C overnight. The mixture was concentrated and the residue was dried under vacuum for 2 hours. The residue (252 mg, 0.9 mmol) was dissolved in

tetrahydrofuran (10 mL). To this solution was added 5-bromo-2-methoxy-benzoic acid (209.0 mg, 0.9 mmol), N-(3-dimethylaminopropyl)-N-ethylcarbodiimide hydrochloride (73 mg, 0.9 mmol), 1-hydroxybenzotriazole (122.0 mg, 0.9 mmol) and triethylamine (315.0 μ L, 2.3 mmol). The mixture was stirred overnight at 80 °C, and cooled to room temperature. The mixture was diluted with ethyl acetate, washed with 1 M aqueous NaHCO₃, dried (Na₂SO₄), filtered and concentrated. The residue was purified by flash chromatography using an Analogix® Intelliflash280TM (SiO₂, 0-75 % ethyl acetate in hexanes) to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-*d*₆) δ 1.32 (s, 9 H), 1.58 - 1.71 (m, 1 H), 1.75 - 1.86 (m, 2 H), 1.87 - 2.00 (m, 1 H), 3.64 (dd, *J*=15, 6.8 Hz, 1 H), 3.78 (s, 3 H), 3.79 - 3.83 (m, 1 H), 4.19 (d, *J*=5.8 Hz, 2 H), 4.23 - 4.35 (m, 1 H), 7.05 (d, *J*=8.8 Hz, 1 H), 7.25 (s, 1 H), 7.56 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.76 (d, *J*=2.7 Hz, 1 H); MS (ESI⁺) *m/z* 453 (M)⁺, 455 (M+2)⁺.

Example 236

5-chloro-2-(cyclopropyloxy)-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

Example 236A

methyl 5-chloro-2-(2-chloroethoxy)benzoate

A mixture of methyl-5-chlorosalicylate (19.5 g, 105 mmol), 2-chloroethyl *p*-toluenesulfonate (19.3 mL, 107 mmol) and K₂CO₃ (28.9 g, 210 mmol) in 105 mL of N,N-dimethylformamide was warmed to 50 °C and allowed to stir for 18 hours. The mixture was cooled to ambient temperature, diluted with 25 mL of ethyl acetate and 25 mL of H₂O. The layers were separated and the organic layer was dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 75% hexanes in ethyl acetate) afforded the title compound. MS (DCI/NH₃) *m/z* 249 (M+H)⁺, 266 (M+NH₄)⁺.

Example 236B

5-chloro-2-(vinylxy)benzoic acid

To Example 236A (15 g, 60 mmol) in 100 mL of tetrahydrofuran at 0 °C was added potassium *t*-butoxide (8.9 g, 75.6 mmol) portion wise with the internal reaction temperature being maintained below 5 °C. After the addition was complete, the mixture was allowed to warm to ambient temperature and was allowed to stir for 18 hours. The mixture was diluted with 25 mL of H₂O and 25 mL of ethyl acetate and

the layers were separated. The aqueous layer was acidified with 1 N aqueous HCl to pH 7 and was extracted with three 15 mL portions of ethyl acetate. These organic extracts (excluding the original organic layer before acidification) were combined and dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford the title compound. MS (DCI/NH₃) m/z 216 (M+NH₄)⁺.

Example 236C

methyl 5-chloro-2-(vinylloxy)benzoate

To Example 236B (5.1 g, 26 mmol) in 30 mL of N,N- dimethylformamide was added K₂CO₃ (10.7 g, 78 mmol) followed by CH₃I (1.8 mL, 29 mmol). The mixture stirred at ambient temperature for 3 hours and was diluted with 20 mL of H₂O and 20 mL of ether. The layers were separated and the aqueous layer was extracted twice with 10 mL of ether. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 75% hexanes in ethyl acetate) afforded the title compound. MS (DCI/NH₃) m/z 213 (M+H)⁺, 230 (M+NH₄)⁺.

Example 236D

methyl 5-chloro-2-cyclopropoxybenzoate

To a solution of Example 236C (1.29 g, 6.1 mmol) in 15 mL of dichloroethane at -5 °C was added chloro-iodomethane (1.4 mL, 19.4 mmol). A solution of diethylzinc (1M solution in hexanes, 9.7 mL, 9.7 mmol) was added dropwise over 1 hour using a syringe pump. After the addition was complete, the mixture was allowed to warm to ambient temperature and was stirred for 45 minutes. The mixture was cooled to 0 °C and quenched with 5 mL of saturated, aqueous NH₄Cl and 1 mL of concentrated NH₄OH. This mixture was diluted with 10 mL of ethyl acetate and the layers were separated. The aqueous layer was extracted twice with 10 mL of ethyl acetate and the combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 75% hexanes in ethyl acetate) afforded the title compound. MS (DCI/NH₃) m/z 227 (M+H)⁺, 244 (M+NH₄)⁺.

Example 236E

5-chloro-2-cyclopropoxybenzoic acid

To a solution of Example 236D (1.24 g, 5.5 mmol) in 10 mL of ethanol at ambient temperature was added 5 mL of 40% aqueous KOH. The mixture was stirred at ambient temperature for 2 hours and was partially concentrated to remove the

ethanol. The aqueous residue was extracted with three 10 mL of portions of CH₂Cl₂. The aqueous layer was acidified with 10 % aqueous HCl to ~pH 1 and then extracted with three 10 mL of portions of CH₂Cl₂. The combined organic extracts (from both extractions) were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure to afford the title compound. MS (DCI/NH₃) m/z 213 (M+H)⁺, 230 (M+NH₄)⁺.

Example 236F

5-chloro-2-(cyclopropyloxy)-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

10 A mixture of Example 236E (0.30 g, 1.4 mmol) and 1,1'-carbonyldiimidazole (0.27 g, 1.7 mmol) in 10 mL of ethyl acetate was stirred at ambient temperature for 4 hours. Example 208B (0.28 g, 1.4 mmol) in 2 mL of ethyl acetate and 2 mL of tetrahydrofuran was added and the mixture was warmed to reflux for 16 hours. The mixture was cooled to ambient temperature and was quenched with 10 mL of H₂O and 5 mL 5% aqueous HCl and was diluted with 10 mL of ethyl acetate. The layers were separated and the aqueous layer was extracted twice with 5 mL of ethyl acetate. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 40% hexanes in ethyl acetate) afforded the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 0.76 - 0.82 (m, 2 H), 1.63 - 1.77 (m, 1 H), 1.84 - 1.96 (m, 2 H), 2.01 - 2.14 (m, 1 H), 2.33 (d, J=1.4 Hz, 3 H), 3.69 - 3.93 (m, 4 H), 4.18 - 4.43 (m, 4 H), 7.13 (q, J=1.4 Hz, 1 H), 7.38 - 7.41 (m, 2 H), 7.74 (dd, J=2.0, 1.0 Hz, 1 H); MS (DCI/NH₃) m/z 393 (M+H)⁺. Anal. Calculated for C₁₉H₂₁ClN₂O₃S: C, 58.08; H, 5.39; N, 7.13. Found: C, 57.77; H, 5.45; N, 7.09.

25

Example 237

5-chloro-N-[(2Z)-3-(1,4-dioxan-2-ylmethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 237A

3-((1,4-dioxan-2-yl)methyl)-5-methylthiazol-2(3H)-imine

30 A mixture of 2-amino-5-methylthiazole (0.77 g, 6.7 mmol) and 2-iodomethyl-1,4-dioxane (Synchem-OHG, 1.5 g, 6.7 mmol) was warmed to 85 °C and was allowed to stir for 18 hours. The mixture was cooled to ambient temperature and the crude material was purified via column chromatography (SiO₂, 10% methanol in ethyl

acetate then 9:1:0.1 CH₂Cl₂ : methanol : NH₄OH) to afford the title compound. MS (DCI/NH₃) m/z 215 (M+H)⁺.

Example 237B

5-chloro-N-[(2Z)-3-(1,4-dioxan-2-ylmethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

5

Example 237A (0.20 g, 0.93 mmol), triethylamine (0.39 mL, 2.8 mmol) and Example 205B (0.93 mmol) in 10 mL of tetrahydrofuran were processed as described in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 2.33 (d, J=1.4 Hz, 3 H), 3.34 (dd, J=11.5, 9.5 Hz, 1 H), 3.50 - 3.71 (m, 3 H), 3.82 (dt, J=10.5, 3.1 Hz, 2 H), 3.86 (s, 3 H), 3.99 - 4.09 (m, 1 H), 4.19 - 4.36 (m, 2 H), 7.08 (d, J=8.8 Hz, 1 H), 7.09 (q, J=1.0 Hz, 1 H), 7.40 (dd, J=9.0, 2.9 Hz, 1 H), 7.81 (d, J=2.7 Hz, 1 H); MS (DCI/NH₃) m/z 383 (M+H)⁺. Anal. Calculated for C₁₇H₁₉ClN₂O₄S: C, 53.33; H, 5.00; N, 7.32. Found: C, 53.51; H, 4.93; N, 7.29.

10

Example 238

N-[(2Z)-5-acetyl-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

15

Example 238A

N-(5-acetyl-4-methylthiazol-2-yl)-5-chloro-2-methoxybenzamide

To a solution of 5-acetyl-2-amino-4-methylthiazole (5.0 g, 32 mmol) in 50 mL of tetrahydrofuran was added triethylamine (13.4 mL, 96 mmol) followed by Example 205B (32 mmol) in 10 mL of tetrahydrofuran via cannula. The mixture was warmed to 50 °C and was allowed to stir for 18 hours. The mixture was cooled to ambient temperature quenched with 15 mL of NH₄Cl and diluted with 15 mL of ethyl acetate. The layers were separated and the aqueous layer was extracted with three 10 mL of portions of ethyl acetate. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was washed with ethyl acetate and the remaining solids were pure title compound. MS (DCI/NH₃) m/z 325 (M+H)⁺.

20

25

Example 238B

N-[(2Z)-5-acetyl-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

30

A mixture of Example 238A (1.1 g, 3.2 mmol), Example 208A (1.0 g, 3.9 mmol), tetrabutylammonium iodide (0.36 g, 0.98 mmol) and potassium t-butoxide

(0.58 g, 4.9 mmol) in 12 mL of N,N-dimethylformamide was warmed to 65 °C and was allowed to stir for 16 hours. The mixture was cooled to ambient temperature quenched with 10 mL of NH₄Cl and diluted with 10 mL of ethyl acetate. The layers were separated and the aqueous layer was extracted twice with 10 mL of ethyl acetate.

- 5 The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 30% hexanes in ethyl acetate) afforded the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.73 - 1.87 (m, 1 H), 1.89 - 2.05 (m, 2 H), 2.09 - 2.22 (m, 1 H), 2.52 (s, 3 H), 2.79 (s, 3 H), 3.69 - 3.78 (m, 1 H), 3.85 - 3.94 (m, 1 H), 3.88 (s, 3 H), 4.21 - 4.31 (m, 1 H), 4.38 - 4.49 (m, 1 H), 4.59 (dd, *J*=13.9, 3.1 Hz, 1 H), 7.10 (d, *J*=9.2 Hz, 1 H), 7.44 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.94 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 409 (M+H)⁺. Anal. Calculated for C₁₉H₂₁ClN₂O₄S·0.12H₂O: C, 55.52; H, 5.21; N, 6.81. Found: C, 55.83; H, 5.21; N, 6.41.

Example 239

- 15 5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

- To Example 238B (0.11 g, 0.27 mmol) in 5 mL of tetrahydrofuran at -78 °C was added a solution of methyl lithium (1.6 M in ether, 0.50 mL, 0.81 mmol) dropwise over 5 minutes. The mixture stirred at -78 °C for 1 hour and was slowly warmed to ambient temperature and was allowed to stir for 18 hours. The mixture was quenched with 5 mL of NH₄Cl and diluted with 5 mL of ethyl acetate. The layers were separated and the aqueous layer was extracted twice with 5 mL of ethyl acetate. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 20% hexanes in ethyl acetate) afforded the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.26 (none, 1 H), 1.61 (s, 6 H), 1.73 - 1.85 (m, 1 H), 1.88 - 2.01 (m, 2 H), 2.05 - 2.19 (m, 1 H), 2.52 (s, 3 H), 3.69 - 3.78 (m, 1 H), 3.85 (s, 3 H), 3.87 - 3.93 (m, 1 H), 4.02 - 4.19 (m, 2 H), 4.36 - 4.45 (m, 1 H), 4.50 (dd, *J*=13.9, 3.1 Hz, 1 H), 7.06 (d, *J*=8.8 Hz, 1 H), 7.39 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.83 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 425 (M+H)⁺. Anal. Calculated for C₂₀H₂₅ClN₂O₄S·0.1H₂O: C, 56.29; H, 5.95; N, 6.56. Found: C, 55.95; H, 5.87; N, 6.47.

Example 240

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-

5-chloro-2-methoxybenzamideExample 240A5-tert-butylthiazol-2-amine

To a solution of 3,3-dimethylbutyraldehyde (10 g, 99.8 mmol) in 200 mL of
5 cyclohexane was added pyrrolidine (8.7 mL, 0.11 mol) followed by *p*-toluenesulfonic
acid monohydrate (0.95 g, 5.0 mmol). This reaction flask was equipped with a Dean-
Stark trap and the mixture was warmed to reflux and was allowed to stir for 3 hours.
The mixture was cooled to ambient temperature, filtered and concentrated under
reduced pressure. The residue was dissolved in 75 mL of CH₃OH, sulfur was added
10 (3.2 g, 99.8 mmol), and the mixture was cooled to 0 °C. Cyanamide (4.2 g, 99.8
mmol) was added portion wise over 10 minutes and the mixture was allowed to warm
to ambient temperature and stir for 18 hours. The reaction mixture was concentrated
under reduced pressure and purified by column chromatography (SiO₂, ethyl acetate
then 10% methanol in ethyl acetate) to afford the title compound. MS (DCI/NH₃) *m/z*
15 157 (M+H)⁺.

Example 240B(R)-(tetrahydrofuran-2-yl)methyl 4-methylbenzenesulfonate

To a solution of (R)-tetrahydrofurfuryl alcohol (Lancaster, 1.0 g, 9.8 mmol) in
5 mL of CH₂Cl₂ and 5 mL of pyridine was added *p*-toluenesulfonyl chloride (2.8 g,
20 14.7 mmol) in portions over 15 minutes. The mixture was stirred at ambient
temperature for 3 hours and was quenched with 5 mL of saturated, aqueous NaHCO₃.
The layers were separated and the aqueous layer was extracted with three 5 mL of
portions of CH₂Cl₂. The combined organic extracts were dried over anhydrous
Na₂SO₄, filtered and concentrated under reduced pressure to afford the title
25 compound. MS (DCI/NH₃) *m/z* 257 (M+H)⁺, 274 (M+NH₄)⁺.

Example 240C(R)-5-tert-butyl-3-((tetrahydrofuran-2-yl)methyl)thiazol-2(3H)-imine 4-
methylbenzenesulfonate

A mixture of Example 240A (9.8 g, 62.7 mmol), Example 240B (23.5 g, 91.7
30 mmol) and tetrabutylammonium iodide (11.6 g, 31.4 mmol) in 35 mL of *N,N*-
dimethylformamide was warmed to 85 °C and was allowed to stir for 72 hours. The
mixture was diluted with 50 mL of CH₂Cl₂ and the layers were separated. The
organic layer was washed with 15 mL of saturated, aqueous NaHCO₃ and the
combined aqueous layers were extracted with three 10 mL of portions of CH₂Cl₂.

Example 241A3-(((1,3-dioxolan-2-yl)methyl)-5-tert-butylthiazol-2(3H)-imine

A mixture of Example 240A (0.17 g, 1.1 mmol) and 2-bromomethyl-1,3-dioxolane (0.21 g, 1.2 mmol) was warmed to 85 °C and was allowed to stir for 18 hours. The mixture was cooled to ambient temperature and the crude material was purified via flash column chromatography (SiO₂, 10% methanol in ethyl acetate then 9:1:0.1 CH₂Cl₂ : methanol : NH₄OH) to afford the title compound. MS (DCI/NH₃) m/z 243 (M+H)⁺.

Example 241B

10 N-[(2Z)-5-tert-butyl-3-((1,3-dioxolan-2-yl)methyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 241A (64 mg, 0.26 mmol), triethylamine (0.11 mL, 0.79 mmol) and Example 205B (0.26 mmol) in 2 mL of tetrahydrofuran and 0.5 mL of N,N-dimethylformamide were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.38 (s, 9 H), 3.86 (s, 3 H), 3.88 - 3.98 (m, 4 H), 4.42 (d, J=4.1 Hz, 2 H), 5.29 (dd, J=4.1 Hz, 1 H), 7.07 (d, J=8.8 Hz, 1 H), 7.12 (s, 1 H), 7.40 (dd, J=9.0, 2.9 Hz, 1 H), 7.87 (d, J=2.7 Hz, 1 H); MS (DCI/NH₃) m/z 411 (M+H)⁺. Anal. Calculated for C₁₉H₂₃ClN₂O₄S: C, 55.54; H, 5.64; N, 6.82. Found: C, 55.43; H, 5.60; N, 6.62.

Example 242

20 5-chloro-N-[(2Z)-5-chloro-3-((2R)-tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 242A5-chloro-N-(5-chlorothiazol-2-yl)-2-methoxybenzamide

25 A mixture of 2-amino-5-chlorothiazole hydrochloric acid (1.0 g, 5.9 mmol), 5-chloro-2-methoxybenzoic acid (1.3 g, 7.0 mmol), 1-ethyl-(3-dimethylaminopropyl)carbodiimide hydrochloride (Chem-Impex International, 2.2 g, 12 mmol), 1-hydroxybenzotriazole (0.95 g, 7.0 mmol) and 4-dimethylaminopyridine (0.14 g, 1.2 mmol) in 6 mL of pyridine was allowed to stir at ambient temperature for 30 72 hours. The reaction mixture was concentrated under reduced pressure and 10 mL of H₂O was added. The resulting solids were isolated by filtration, washed with 5 mL of H₂O and twice with 5 mL of ethyl acetate, and dried to afford the title compound. MS (DCI/NH₃) m/z 303 (M+H)⁺.

Example 242B5-chloro-N-[(2Z)-5-chloro-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

To a slurry of sodium hydride (40 mg of a 60% dispersion, 1.0 mmol) in 4 mL of N,N- dimethylformamide at 0 °C was added Example 242A (0.20 g, 0.66 mmol). This mixture was allowed to warm to ambient temperature and stirred for 1 hour. The mixture was cooled to 0 °C and Example 208A (0.19 g, 0.73 mmol) was added. The mixture was warmed to 80 °C and allowed to stir for 24 hours and cooled to ambient temperature. The reaction mixture was quenched with ice and 5 mL of saturated, aqueous NH₄Cl and diluted with 5 mL of ethyl acetate. The layers were separated and the aqueous layer was extracted with three 5 mL of portions of ethyl acetate. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via flash column chromatography (SiO₂, 50% hexanes : ethyl acetate) afforded the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.65 - 1.80 (m, 1 H), 1.85 - 1.97 (m, 2 H), 2.02 - 2.18 (m, 1 H), 3.72 - 3.82 (m, 1 H), 3.84 - 3.94 (m, 1 H), 3.87 (s, 3 H), 4.21 - 4.38 (m, 2 H), 4.40 - 4.48 (m, 1 H), 7.09 (d, *J*=8.8 Hz, 1 H), 7.43 (dd, *J*=9.0, 2.9 Hz, 1 H), 7.50 (s, 1 H), 7.90 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 387 (M+H)⁺. Anal. Calculated for C₁₆H₁₆Cl₂N₂O₃S: C, 49.62; H, 4.16; N, 7.23. Found: C, 50.49; H, 4.03; N, 6.70.

Example 2435-chloro-N-[(2Z)-5-chloro-3-(tetrahydro-2H-pyran-4-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 242A (0.20 g, 0.66 mmol), Example 203A (0.20, 0.73 mmol) and NaH (40 mg, 1 mmol) in 4 mL of tetrahydrofuran and 1 mL of N,N- dimethylformamide were processed as in Example 242B to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.36 - 1.52 (m, 2 H), 1.53 - 1.61 (m, 2 H), 2.19 - 2.38 (m, 1 H), 3.40 (dt, *J*=11.7, 2.4 Hz, 2 H), 3.88 (s, 3 H), 3.92 - 4.00 (m, 2 H), 4.19 (d, *J*=7.1 Hz, 2 H), 7.10 (d, *J*=8.8 Hz, 1 H), 7.44 (dd, *J*=9.0, 2.9 Hz, 1 H), 7.55 (s, 1 H), 7.91 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 401 (M+H)⁺.

Example 244N-[(2Z)-5-tert-butyl-3-[(3-methyl-4,5-dihydroisoxazol-5-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamideExample 244A

N-(5-tert-butylthiazol-2-yl)-5-chloro-2-methoxybenzamide

To a solution of Example 202A (0.94 g, 6.0 mmol) in tetrahydrofuran (40 mL) was added Example 205B (1.23 g, 6.0 mmol), triethylamine (2.4 mL, 18 mmol), and 4-dimethylaminopyridine (7.5 mg, 0.06 mmol). The reaction mixture was stirred at 60 °C for 14 hours and then cooled to ambient temperature, diluted with saturated aqueous NaHCO₃ (20 mL) and extracted with ethyl acetate (3 x 30 mL). The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography using an Analogix® Intelliflash280™ (SiO₂, 0-100 % ethyl acetate in hexanes) to afford the title compound. MS (ESI⁺) m/z 325 (M+H)⁺

Example 244B

N-[(2Z)-3-allyl-5-tert-butyl-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

To a solution of Example 244A (410 mg, 1.3 mmol) in 5 mL of tetrahydrofuran:N,N-dimethylformamide (4/1) at 0 °C was added potassium *tert*-butoxide (230 mg, 1.9 mmol). The reaction mixture was stirred for 1 hour then allyl bromide (0.16 mL, 1.9 mmol) was added. The mixture was warmed to 65 °C and stirred overnight. The mixture was cooled to ambient temperature, concentrated, diluted with CH₂Cl₂ and washed with water and brine. The organic layer was dried over magnesium sulfate, filtered, and concentrated under reduced pressure. Purification by chromatography (20-50% ethyl acetate/hexane gradient) afforded the title compound. ¹H NMR (CDCl₃, 300 MHz) δ ppm 1.34 (s, 9 H), 3.90 (s, 3 H), 4.78 - 4.85 (m, 3 H), 5.22 - 5.36 (m, 2 H), 5.90 - 6.09 (m, *J*=17.0, 10.17 Hz, 1 H), 6.62 (s, 1 H), 6.90 (d, *J*=9.2 Hz, 1 H), 7.32 (dd, *J*=8.8, 3.1 Hz, 1 H), 8.00 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) m/z 365 (M+H)⁺.

Example 244C

N-[(2Z)-5-tert-butyl-3-[(3-methyl-4,5-dihydroisoxazol-5-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 244B was processed according to the method of Example 227C to afford the title compound. MS (DCI/NH₃) m/z 422 (M+H)⁺.

Example 245

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-2,2-dimethyl-4-oxo-3,4-dihydro-2H-pyran-6-carboxamide

To a solution of the product from Example 240C (300 mg, 0.94 mmol) and 2,2-dimethyl-4-oxo-3,4-dihydro-2H-pyran-6-carboxylic acid (180 mg, 1.0 mmol) in 5 mL of N,N-dimethylformamide were added 1-hydroxybenzotriazole hydrate (190 mg, 1.4 mmol), triethylamine (0.30 mL, 2.1 mmol), and 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (260 mg, 1.4 mmol). The mixture was warmed to 65 °C and stirred overnight. The mixture was cooled to ambient temperature, diluted with CH₂Cl₂ and washed with water and brine. The organic extract was dried (MgSO₄), filtered, and concentrated under reduced pressure. Purification by column chromatography (SiO₂, 20-40% ethyl acetate/hexanes gradient) afforded the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.34 (s, 9H), 1.57 (s, 6H), 1.72 - 1.84, (m, 1H), 1.83 - 1.94 (m, 1H), 2.00 - 2.12 (m, 2H), 2.55 (s, 2H), 3.73 - 3.88 (m, 2H), 4.10 - 4.27 (m, 2H), 4.39 (dd, *J*=13.7, 2.8 Hz, 1H), 6.48 (s, 1H), 6.91 (s, 1H); MS (DCI/NH₃) *m/z* 393 (M+H)⁺.

Example 246

N-[(2Z)-5-tert-butyl-3-(tetrahydro-2H-pyran-4-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

To a solution of Example 244A (1.0 g, 3.1 mmol) in 4:1 N,N-dimethylformamide/tetrahydrofuran (20 mL) were added potassium tert-butoxide (Aldrich, 0.42 g, 3.7 mmol) and 4-(iodomethyl)tetrahydro-2H-pyran (Maybridge, 0.97 g, 4.3 mmol). The reaction mixture was stirred at 80 °C for 16 hours, cooled to room temperature, quenched with saturated aqueous NaHCO₃ (20 mL) and extracted with ethyl acetate (3 x 20 mL). The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography using an Analogix® Intelliflash280TM (SiO₂, 0-100 % ethyl acetate in hexanes) to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.21 - 1.51 (m, 4 H), 1.32 (s, 9 H), 2.06 - 2.35 (m, 1 H), 3.20 - 3.30 (m, 2 H), 3.79 (s, 3 H), 3.80 - 3.91 (m, *J*=9.3, 2.2, 2.0 Hz, 2 H), 4.06 (d, *J*=7.1 Hz, 2 H), 7.11 (d, *J*=8.8 Hz, 1 H), 7.30 (s, 1 H), 7.45 (dd, *J*=8.8, 3.1 Hz, 1 H), 7.64 (d, *J*=2.7 Hz, 1 H); MS (ESI⁺) *m/z* 423 (M+H)⁺; Anal. Calculated for C₂₁H₂₇ClN₂O₃S: C, 59.63; H, 6.43; N, 6.62. Found: C, 59.66; H, 6.36; N, 6.56.

Example 247

5-chloro-2-methoxy-N-[(2Z)-3-(tetrahydro-2H-pyran-4-ylmethyl)-4,5,6,7-tetrahydro-

1,3-benzothiazol-2(3H)-ylidene]benzamideExample 247A5-chloro-2-methoxy-N-(4,5,6,7-tetrahydrobenzo[d]thiazol-2-yl)benzamide

Commercially available 4,5,6,7-tetrahydrobenzo[d]thiazol-2-amine (Aldrich) and 5-chloro-2-methoxybenzoic acid (Aldrich) were processed using the method described in Example 58 to afford the title compound. MS (ESI⁺) m/z 323 (M+H)⁺

Example 247B5-chloro-2-methoxy-N-[(2Z)-3-(tetrahydro-2H-pyran-4-ylmethyl)-4,5,6,7-tetrahydro-1,3-benzothiazol-2(3H)-ylidene]benzamide

Example 203A and Example 247A were processed using the method described in Example 246 to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.25 - 1.55 (m, 4 H), 1.69 - 1.94 (m, 4 H), 2.07 - 2.30 (m, 1 H), 2.52 - 2.59 (m, 2 H), 2.58 - 2.66 (m, 2 H), 3.18 - 3.30 (m, 2 H), 3.80 (s, 3 H), 3.81 - 3.89 (m, 2 H), 4.03 (d, J=7.1 Hz, 2 H), 7.11 (d, J=8.8 Hz, 1 H), 7.45 (dd, J=8.8, 2.7 Hz, 1 H), 7.69 (d, J=3.1 Hz, 1 H); MS (ESI⁺) m/z 421 (M+H)⁺.

Example 2485-chloro-2-methoxy-N-[(2Z)-3-(2-methoxyethyl)-4,4,6,6-tetramethyl-4,6-dihydrofuro[3,4-d][1,3]thiazol-2(3H)-ylidene]benzamideExample 248A4-bromo-2,2,5,5-tetramethyldihydrofuran-3(2H)-one

To a solution of commercially available 2,2,5,5-tetramethyldihydrofuran-3(2H)-one (Aldrich, 10.0 g, 0.07 mol) in CH₂Cl₂ (100 mL) was added bromine (Aldrich, 3.6 mL, 0.07 mol,) dropwise at room temperature. The reaction mixture was stirred for 2 hours over which time the reaction mixture became clear. Then, the reaction mixture was cooled to 0 °C, quenched with NaHCO₃ powder in small portions, filtered and concentrated under reduced pressure to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.26 (s, 3 H), 1.27 (s, 3 H), 1.30 (s, 3 H), 1.39 (s, 3 H), 5.22 (s, 1 H).

Example 248B4,4,6,6-tetramethyl-4,6-dihydrofuro[3,4-d]thiazol-2-amine

To a solution of Example 248A (10.0 g, 0.045 mol) in ethanol (100 mL) were added thiourea (3.8 g, 0.05 mol) and triethylamine (6.3 mL, 0.045 mol). The reaction mixture was heated at reflux overnight, then cooled, and concentrated under reduced

pressure. The residue was diluted with water (25 mL) and extracted with ethyl acetate (3x50 mL). The combined organic extracts were dried (Na₂SO₄), filtered and concentrated. The residue was purified by column chromatography (SiO₂, 0-5 % methanol in CH₂Cl₂) to afford the title compound. MS (ESI⁺) m/z 199 (M+H)⁺.

5 Example 248C

5-chloro-2-methoxy-N-(4,4,6,6-tetramethyl-4,6-dihydrofuro[3,4-d]thiazol-2-yl)benzamide

Example 248B and Example 205B were processed using the method described in Example 244A to afford the title compound. MS (ESI⁺) m/z 367 (M+H)⁺.

10 Example 248D

5-chloro-2-methoxy-N-[(2Z)-3-(2-methoxyethyl)-4,4,6,6-tetramethyl-4,6-dihydrofuro[3,4-d][1,3]thiazol-2(3H)-ylidene]benzamide

Example 248C and commercially available 2-bromoethyl methyl ether (Aldrich) were processed using the method described in Example 246 to afford the
15 title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.48 (s, 6 H), 1.55 (s, 6 H), 3.25 (s, 3 H), 3.71 - 3.90 (m, 2 H), 3.81 (s, 3 H), 4.20 (t, J=5.6 Hz, 2 H), 7.13 (d, J=9.2 Hz, 1 H), 7.48 (dd, J=9.0, 2.9 Hz, 1 H), 7.68 (d, J=2.7 Hz, 1 H); MS (ESI⁺) m/z 425 (M+H)⁺

Example 249

20 5-chloro-2-methoxy-N-[(2Z)-3-(2-methoxyethyl)-6,6-dimethyl-4-oxo-4,6-dihydrofuro[3,4-d][1,3]thiazol-2(3H)-ylidene]benzamide

Example 249A

tert-butyl 6,6-dimethyl-4-oxo-4,6-dihydrofuro[3,4-d]thiazol-2-ylcarbamate

To a solution of diisopropylamine (23.5 mL, 165 mmol) in tetrahydrofuran
25 (200 mL) was added butyllithium (103 mL, 1.6 M in hexanes, 165 mmol) dropwise at -78 °C. The solution was stirred at -78 °C for 30 minutes then transferred via cannula into a solution of methyl 2-(tert-butoxycarbonylamino)thiazole-4-carboxylate (Combi-Blocks, 14.2 g, 55 mmol) in tetrahydrofuran (300 mL) at -78 °C. After stirring at -78 °C for 30 minutes, dry acetone (Acros, 16.2 mL, 220 mmol) was added
30 dropwise and the reaction mixture was allowed to warm to room temperature and stirred overnight. The reaction mixture was quenched with saturated aqueous NH₄Cl solution (200 mL) and the aqueous layer was extracted with ethyl acetate (4 x 200 mL). The combined organic extracts were dried (Na₂SO₄), filtered and concentrated.

The residue was purified by column chromatography (SiO₂, 0-100 % ethyl acetate in hexanes) to afford the title compound. MS (ESI⁺) m/z 285 (M+H)⁺

Example 249B

2-amino-6,6-dimethylfuro[3,4-d]thiazol-4(6H)-one

5 To a solution of Example 249A (7.4 g, 26.0 mmol) in CH₂Cl₂ (20 mL) was added trifluoroacetic acid (20.0 mL, 260 mmol) slowly at 0 °C. The reaction mixture was stirred at room temperature for 3 hours, and then concentrated under vacuum. The residue was diluted with ethyl acetate (100 mL) and neutralized with saturated aqueous NaHCO₃ solution. The layers were separated and the aqueous phase was
10 extracted with ethyl acetate (5 x 100 mL). The combined organic extracts were dried (Na₂SO₄), filtered and concentrated to afford the title compound. MS (ESI⁺) m/z 185 (M+H)⁺

Example 249C

2-imino-3-(2-methoxyethyl)-6,6-dimethyl-2,3-dihydrofuro[3,4-d]thiazol-4(6H)-one

15 hydrobromide

A mixture of Example 249B and commercially available 2-bromoethyl methyl ether (Aldrich) was processed at 120 °C using the method described in Example 12A to afford the title compound. MS (ESI⁺) m/z 243 (M+H)⁺

Example 249D

20 5-chloro-2-methoxy-N-[(2Z)-3-(2-methoxyethyl)-6,6-dimethyl-4-oxo-4,6-dihydrofuro[3,4-d][1,3]thiazol-2(3H)-ylidene]benzamide

Example 249C and Example 205B were processed using the method described in Example 244A to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.71 (s, 6 H), 3.26 (s, 3 H), 3.78 (t, J=5.6 Hz, 2 H), 3.82
25 (s, 3 H), 4.46 (t, J=5.6 Hz, 2 H), 7.17 (d, J=8.8 Hz, 1 H), 7.53 (dd, J=9.0, 2.9 Hz, 1 H), 7.77 (d, J=2.7 Hz, 1 H); MS (ESI⁺) m/z 411 (M+H)⁺; Anal. Calculated for C₁₈H₁₉ClN₂O₅S: C, 52.62; H, 4.66; N, 6.82. Found: C, 52.72; H, 4.49; N, 6.90.

Example 250

30 N-[(2Z)-5-acetyl-4-methyl-3-(oxetan-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 238A and Example 211A were processed using the method described in Example 246 to afford the title compound. ¹H NMR (300 MHz, CDCl₃) δ ppm 2.49
(s, 3 H), 2.54 - 2.67 (m, 1 H), 2.74 - 2.82 (m, 1 H), 2.84 (s, 3 H), 3.93 (s, 3 H), 4.45 -

4.74 (m, 4 H), 5.30 (d, 1 H), 6.93 (d, $J=9.2$ Hz, 1 H), 7.38 (dd, $J=8.8, 2.7$ Hz, 1 H), 7.97 (d, $J=2.7$ Hz, 1 H); MS (ESI⁺) m/z 493 (M+H)⁺; Anal. Calculated for C₁₈H₁₉ClN₂O₄S: C, 54.75; H, 4.85; N, 7.09. Found: C, 54.68; H, 4.70; N, 7.07.

Example 251

5 5-chloro-N-[(2Z)-4,4-dimethyl-1-(oxetan-2-ylmethyl)-1,4,6,7-tetrahydro-2H-pyrano[4,3-d][1,3]thiazol-2-ylidene]-2-methoxybenzamide

Example 251A

ethyl 2-(2-(tert-butoxycarbonylamino)thiazol-4-yl)acetate

The title compound was obtained from commercially available ethyl 2-(2-aminothiazol-4-yl)acetate (Aldrich) as per the procedure described in JP 06345736. MS (ESI⁺) m/z 287 (M+H)⁺.

Example 251B

tert-butyl 4-(2-hydroxyethyl)thiazol-2-ylcarbamate

To a cooled solution of Example 251A in tetrahydrofuran (100 mL) was added lithium borohydride (Aldrich, 100 mL, 2 M solution in tetrahydrofuran) at 0 °C. The reaction mixture was heated at reflux overnight, then cooled to 0 °C, quenched with water and extracted with ethyl acetate (3x100 mL). The combined organic extracts were dried (Na₂SO₄), filtered and concentrated. The residue was purified by column chromatography (SiO₂, 0-5 % methanol in CH₂Cl₂) to afford the title compound. MS (ESI⁺) m/z 245 (M+H)⁺.

Example 251C

tert-butyl 4-(2-(tetrahydro-2H-pyran-2-yloxy)ethyl)thiazol-2-ylcarbamate

To a solution of Example 251B (6.3 g, 27.4 mmol) in CH₂Cl₂ (100 mL) was added commercially available 3,4-dihydro-2H-pyran (Aldrich, 21 g, 250 mmol) and pyridinium-*p*-toluenesulfonic acid (Aldrich, 3.5 g, 14.0 mmol). The reaction mixture was stirred overnight at room temperature and was diluted with CH₂Cl₂, washed with water, dried (Na₂SO₄), filtered and concentrated. The residue was purified by column chromatography (SiO₂, 0-100 % ethyl acetate in hexanes) to afford the title compound. MS (ESI⁺) m/z 329 (M+H)⁺.

30 Example 251D

tert-butyl 5-(2-hydroxypropan-2-yl)-4-(2-(tetrahydro-2H-pyran-2-yloxy)ethyl)thiazol-2-ylcarbamate

Example 251C, diisopropylamine, butyllithium, and dry acetone (Acros) were processed as described for Example 249A to obtain the title compound. MS (ESI⁺)

m/z 387 (M+H)⁺.

Example 251E

4,4-dimethyl-6,7-dihydro-4H-pyrano[4,3-d]thiazol-2-amine

To a solution of Example 251D (4.6 g, 11 mmol) in tetrahydrofuran was added
 5 concentrated HCl (6.9 mL). The reaction mixture was heated at reflux for overnight
 and then cooled. The mixture was basified with 5N NaOH (17 mL) and extracted with
 ethyl acetate (3 x 50 mL). The combined organic extracts were dried (Na₂SO₄),
 filtered and concentrated. The residue was purified by column chromatography (SiO₂,
 0-10% methanol in CH₂Cl₂) to afford the title compound. MS (ESI⁺) m/z 185
 10 (M+H)⁺.

Example 251F

5-chloro-N-(4,4-dimethyl-6,7-dihydro-4H-pyrano[4,3-d]thiazol-2-yl)-2-methoxybenzamide

Example 251E and Example 205B were processed using the method described
 15 in Example 244A to afford the title compound. MS (ESI⁺) m/z 353 (M+H)⁺

Example 251G

5-chloro-N-[(2Z)-4,4-dimethyl-1-(oxetan-2-ylmethyl)-1,4,6,7-tetrahydro-2H-pyrano[4,3-d][1,3]thiazol-2-ylidene]-2-methoxybenzamide

Example 251F and Example 211A were processed using the method described
 20 in Example 246 to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.45 (s, 3 H), 1.46 (s, 3 H), 2.67 - 2.75 (m, 2 H), 2.74 - 2.83 (m, 2 H), 3.79 (s, 3 H), 3.90 - 4.01 (m, 2 H), 4.32 - 4.44 (m, 2 H), 4.45 - 4.58 (m, 2 H), 5.06 (d, 1 H), 7.11 (d, J=8.8 Hz, 1 H), 7.45 (dd, J=8.8, 2.7 Hz, 1 H), 7.62 (d, J=2.7 Hz, 1 H); MS (ESI⁺) m/z 423 (M+H)⁺; Anal. Calculated for C₂₀H₂₃ClN₂O₄S·0.2C₄H₈O₂: C, 56.71; H, 5.63; N, 6.36. Found: C, 56.33; H, 5.39; N, 6.41.

Example 252

5-chloro-N-[(2Z)-4,4-dimethyl-1-[(2R)-tetrahydrofuran-2-ylmethyl]-1,4,6,7-tetrahydro-2H-pyrano[4,3-d][1,3]thiazol-2-ylidene]-2-methoxybenzamide

Example 251F and Example 208A were processed using the method described
 30 in Example 246 to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.45 (s, 6 H), 1.61 - 2.04 (m, 4 H), 2.71 (t, J=5.8 Hz, 2 H), 3.56 - 3.70 (m, 1 H), 3.73 - 3.84 (m, 1 H), 3.79 (s, 3 H), 3.96 (t, J=5.3 Hz, 2 H), 4.00 - 4.05 (m, 1 H), 4.20 - 4.38 (m, 2 H), 7.11 (d, J=8.8 Hz, 1 H), 7.46 (dd, J=9.0, 2.9 Hz, 1 H), 7.66 (d,

$J=3.1$ Hz, 1 H); MS (ESI⁺) m/z 437 (M+H)⁺; Anal. Calculated for C₂₁H₂₅ClN₂O₄S: C, 57.72; H, 5.77; N, 6.41. Found: C, 57.58; H, 5.86; N, 6.33.

Example 253

N-[(2Z)-5-acetyl-4-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Commercially available 2-(bromomethyl)tetrahydro-2H-pyran (Aldrich) and Example 238A were processed using the method described in Example 246 to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.28 - 1.38 (m, 1 H), 1.41 - 1.52 (m, 3 H), 1.62 - 1.71 (m, 1 H), 1.78 - 1.88 (m, 1 H), 2.50 (s, 3 H), 2.61 - 2.84 (m, 3 H), 3.64 - 3.93 (m, 3 H), 3.83 (s, 3 H), 4.14 (dd, $J=14.1$, 8.6 Hz, 1 H), 4.37 (dd, $J=14.2$, 3.1 Hz, 1 H), 7.16 (d, $J=8.8$ Hz, 1 H), 7.51 (dd, $J=9.0$, 2.9 Hz, 1 H), 7.79 (d, $J=2.7$ Hz, 1 H); MS (ESI⁺) m/z 423 (M+H)⁺; Anal. Calculated for C₂₀H₂₃ClN₂O₄S: C, 56.80; H, 5.48; N, 6.62. Found: C, 56.53; H, 5.27; N, 6.55.

Example 254

N-[(2Z)-5-acetyl-4-methyl-3-(tetrahydro-2H-pyran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 225A and Example 238A were processed using the method described in Example 246 to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.47 (s, 6 H) 1.52 (s, 6 H) 3.72 (s, 3 H) 5.35 (s, 2 H) 7.07 (d, 3 H) 7.32 (d, 1 H) 7.37 (d, $J=2.71$ Hz, 1 H) 7.40 - 7.50 (m, $J=8.81$ Hz, 1 H); MS (ESI⁺) m/z 423 (M+H)⁺; Anal. Calculated for C₂₀H₂₃ClN₂O₄S·0.2H₂O: C, 56.32; H, 5.53; N, 6.57. Found: C, 56.19; H, 5.50; N, 6.62.

Example 255

5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-4-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

A solution of methyllithium (Aldrich, 1.6 M in diethyl ether, 0.41 mL, 0.66 mmol) was added slowly a solution of Example 253 (0.14 g, 0.33 mmol) in tetrahydrofuran (3 mL) at -78 °C. The reaction mixture was stirred at -78 °C for 30 minutes and was allowed to reach room temperature. The reaction mixture was quenched with water (6 mL) and extracted with ethyl acetate (2x10 mL). The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography using an Analogix® Intelliflash280TM (SiO₂, 0-5 % methanol in

CH₂Cl₂) to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.21 - 1.36 (m, 1 H), 1.41 - 1.48 (m, 3 H), 1.49 (s, 3 H), 1.50 (s, 3 H), 1.57 - 1.67 (m, 1 H), 1.74 - 1.85 (m, 1 H), 2.42 (s, 3 H), 3.68 - 3.89 (m, 3 H), 3.79 (s, 3 H), 3.92 - 4.09 (m, 1 H), 4.17 - 4.38 (m, 1 H), 5.60 (s, 1 H), 7.10 (d, *J*=8.8 Hz, 1 H), 7.44 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.68 (d, *J*=2.7 Hz, 1 H); MS (ESI⁺) *m/z* 439 (M+H)⁺; Anal. Calculated for C₂₁H₂₇ClN₂O₄S: C, 57.46; H, 6.20; N, 6.38. Found: C, 57.44; H, 5.88; N, 6.06.

Example 256

5-chloro-2-methoxy-N-[(2Z)-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-5-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

Example 256A

2-(2-amino-4-methylthiazol-5-yl)-1,1,1,3,3,3-hexafluoropropan-2-ol

The title compound was prepared from commercially available of 4-methylthiazol-2-amine (Aldrich) and hexafluoroacetone trihydrate (Aldrich) using the procedure described in European Journal of Organic Chemistry, (21), 4286-4291; 2003. MS (ESI⁺) *m/z* 281 (M+H)⁺.

Example 256B

5-chloro-N-(5-(1,1,1,3,3,3-hexafluoro-2-hydroxypropan-2-yl)-4-methylthiazol-2-yl)-2-methoxybenzamide

Commercially available 5-chloro-2-methoxybenzoic acid (Aldrich) and Example 256A were processed using the method described in Example 58 to afford the title compound. MS (ESI⁺) *m/z* 449 (M+H)⁺.

Example 256C

5-chloro-2-methoxy-N-[(2Z)-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-5-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

Example 256B and Example 208A were processed using the method described in Example 246 to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.62 - 1.76 (m, 1 H), 1.78 - 1.93 (m, 2 H), 1.93 - 2.09 (m, 1 H), 2.57 (s, 3 H), 3.57 - 3.70 (m, 1 H), 3.74 - 3.88 (m, 1 H), 3.80 (s, 3 H), 4.12 - 4.26 (m, 1 H), 4.39 (d, 2 H), 7.14 (d, *J*=8.8 Hz, 1 H), 7.49 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.74 (d, *J*=2.7 Hz, 1 H), 9.34 (s, 1 H); MS (ESI⁺) *m/z* 533 (M+H)⁺.

Example 257

5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-4-methyl-3-(tetrahydro-2H-pyran-3-

ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 254 and commercially available methyllithium (Aldrich, 1.6 M in diethyl ether) were processed using the method described in Example 255 to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.32 - 1.48 (m, 2 H), 1.51 (s, 6 H), 1.57 - 1.80 (m, 2 H), 2.02 - 2.23 (m, 1 H), 2.41 (s, 3 H), 3.19 - 3.28 (m, 1 H), 3.35 - 3.43 (m, 1 H), 3.55 - 3.73 (m, 2 H), 3.79 (s, 3 H), 3.95 - 4.29 (m, 2 H), 5.60 (s, 1 H), 7.10 (d, *J*=8.8 Hz, 1 H), 7.44 (dd, *J*=8.8, 3.1 Hz, 1 H), 7.67 (d, *J*=2.7 Hz, 1 H); MS (ESI⁺) *m/z* 439 (M+H)⁺; Anal. Calculated for C₂₁H₂₇ClN₂O₄S: C, 57.46; H, 6.20; N, 6.38. Found: C, 57.14; H, 6.23; N, 6.53.

Example 258

5-chloro-N-[(2Z)-3-{[(4R)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}-5-(1-hydroxy-1-methylethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 258A

5-chloro-N-(5-(2-hydroxypropan-2-yl)-4-methylthiazol-2-yl)-2-methoxybenzamide

Example 238A and methyllithium (Aldrich, 1.6M in diethyl ether) were processed using the method described in Example 255 to afford the title compound. MS (ESI⁺) *m/z* 341 (M+H)⁺.

Example 258B

5-chloro-N-[(2Z)-3-{[(4R)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}-5-(1-hydroxy-1-methylethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 258A and commercially available (S)-(2,2-dimethyl-1,3-dioxolan-4-yl)methyl 4-methylbenzenesulfonate (Aldrich) were processed using the method described in Example 246 to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.25 (s, 3 H), 1.31 (s, 3 H), 1.50 (s, 6 H), 2.45 (s, 3 H), 3.78 (s, 3 H), 3.84 (dd, *J*=8.8, 6.1 Hz, 1 H), 4.04 (dd, *J*=8.5, 6.4 Hz, 1 H), 4.18 - 4.29 (m, 1 H), 4.32 - 4.42 (m, 1 H), 4.46 - 4.58 (m, 1 H), 5.63 (s, 1 H), 7.10 (d, *J*=8.8 Hz, 1 H), 7.43 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.64 (d, *J*=2.7 Hz, 1 H); MS (ESI⁺) *m/z* 455 (M+H)⁺; Anal. Calculated for C₂₁H₂₇ClN₂O₅S: C, 55.44; H, 5.98; N, 6.16. Found: C, 55.34; H, 5.79; N, 6.21.

Example 259

5-chloro-N-[(2Z)-6,6-dimethyl-4-oxo-3-[(2R)-tetrahydrofuran-2-ylmethyl]-4,6-dihydrofuro[3,4-d][1,3]thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 259A

5-chloro-N-(6,6-dimethyl-4-oxo-4,6-dihydrofuro[3,4-d]thiazol-2-yl)-2-methoxybenzamide

Example 249B, triethylamine, 4-dimethylaminopyridine, and Example 205B were processed as described for Example 244A to obtain the title compound. MS

5 (ESI⁺) m/z 353 (M+H)⁺.

Example 259B

5-chloro-N-[(2Z)-6,6-dimethyl-4-oxo-3-[(2R)-tetrahydrofuran-2-ylmethyl]-4,6-dihydrofuro[3,4-d][1,3]thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 259A and Example 208A were processed using the method described
10 in Example 246 to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.71 (s, 6 H), 1.73 - 2.05 (m, 4 H), 3.60 - 3.69 (m, 1 H), 3.72 - 3.81 (m, 1 H), 3.82 (s, 3 H), 4.18 - 4.35 (m, 2 H), 4.35 - 4.47 (m, 1 H), 7.17 (d, J=8.8 Hz, 1 H), 7.53 (dd, J=8.8, 2.7 Hz, 1 H), 7.76 (d, J=2.7 Hz, 1 H); MS (ESI⁺) m/z 437 (M+H)⁺; Anal. Calculated for C₂₀H₂₁ClN₂O₅S·0.4H₂O: C, 54.09; H, 4.95; N, 6.31. Found: C,
15 53.81; H, 4.55; N, 5.99.

Example 260

5-chloro-N-[(2Z)-3-([(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl)]-5-(1-hydroxy-1-methylethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 258A and commercially available (R)-(2,2-dimethyl-1,3-dioxolan-4-yl)methyl 4-methylbenzenesulfonate (Aldrich) were processed using the method
20 described in Example 246 to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.25 (s, 3 H), 1.31 (s, 3 H), 1.50 (s, 6 H), 2.45 (s, 3 H), 3.78 (s, 3 H), 3.84 (dd, J=8.5, 6.1 Hz, 1 H), 4.04 (dd, J=8.8, 6.4 Hz, 1 H), 4.14 - 4.30 (m, 1 H), 4.32 - 4.43 (m, 1 H), 4.43 - 4.63 (m, 1 H), 5.63 (s, 1 H), 7.10 (d, J=9.2 Hz, 1
25 H), 7.43 (dd, J=9.0, 2.9 Hz, 1 H), 7.64 (d, J=2.7 Hz, 1 H); MS (ESI⁺) m/z 455 (M+H)⁺; Anal. Calculated for C₂₁H₂₇ClN₂O₅S: C, 55.44; H, 5.98; N, 6.16. Found: C, 55.73; H, 6.07; N, 6.07.

Example 261

N-[(2Z)-5-acetyl-3-(1,4-dioxan-2-ylmethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

30

Example 238A and commercially available 2-(iodomethyl)-1,4-dioxane (Synchem) were processed using the method described in Example 246 to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 2.50 (s, 3 H), 2.72

(s, 3 H), 3.33 - 3.42 (m, 1 H), 3.45 - 3.56 (m, 2 H), 3.59 - 3.77 (m, 2 H), 3.78 - 3.87 (m, 1 H), 3.83 (s, 3 H), 4.00 - 4.11 (m, 1 H), 4.12 - 4.25 (m, 1 H), 4.31 - 4.43 (m, 1 H), 7.16 (d, $J=8.8$ Hz, 1 H), 7.52 (dd, $J=8.8, 3.1$ Hz, 1 H), 7.79 (d, $J=2.7$ Hz, 1 H); MS (ESI⁺) m/z 425 (M+H)⁺; Anal. Calculated for C₁₉H₂₁ClN₂O₅S: C, 53.71; H, 4.98; N, 6.59. Found: C, 53.32; H, 4.73; N, 6.59.

Example 262

5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-4-methyl-3-(oxetan-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 258A and Example 211A were processed using the method described in Example 246 to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.50 (s, 6 H), 2.47 (s, 3 H), 2.59 - 2.84 (m, 1 H), 3.78 (s, 3 H), 4.25 - 4.64 (m, 5 H), 5.00 - 5.19 (m, 1 H), 5.63 (s, 1 H), 7.09 (d, $J=9.2$ Hz, 1 H), 7.43 (dd, $J=8.8, 3.1$ Hz, 1 H), 7.60 (d, $J=2.7$ Hz, 1 H); MS (ESI⁺) m/z 411 (M+H)⁺; Anal. Calculated for C₁₉H₂₃ClN₂O₄S: C, 55.54; H, 5.64; N, 6.82. Found: C, 55.41; H, 5.51; N, 6.78.

Example 263

5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-4-methyl-3-(tetrahydrofuran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 258A and Example 162A were processed using the method described in Example 246 to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.51 (s, 6 H), 1.65 - 1.82 (m, 1 H), 1.84 - 2.03 (m, 1 H), 2.43 (s, 3 H), 2.68 - 2.89 (m, 1 H), 3.48 - 3.71 (m, 3 H), 3.78 (s, 3 H), 3.80 - 3.93 (m, 1 H), 4.08 - 4.34 (m, 2 H), 5.62 (s, 1 H), 7.09 (d, $J=8.8$ Hz, 1 H), 7.43 (dd, $J=8.8, 3.1$ Hz, 1 H), 7.65 (d, $J=2.7$ Hz, 1 H); MS (ESI⁺) m/z 425 (M+H)⁺; Anal. Calculated for C₂₀H₂₅ClN₂O₄S: C, 56.53; H, 5.93; N, 6.59. Found: C, 56.35; H, 5.55; N, 6.56.

Example 264

5-chloro-N-[(2Z)-3-(1,4-dioxan-2-ylmethyl)-5-(1-hydroxy-1-methylethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 258A and commercially available 2-(iodomethyl)-1,4-dioxane (Synchem) were processed using the method described in Example 246 to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 0.93 (t, $J=7.46$ Hz, 3 H) 1.28 - 1.38 (m, 2 H) 1.45 (s, 6 H) 1.63 - 1.78 (m, 2 H) 2.69 (t, $J=5.42$ Hz, 2 H) 3.79 (s, 3 H) 3.97 (t, $J=5.42$ Hz, 2 H) 4.07 - 4.17 (m, 2 H) 7.11 (d, $J=8.81$ Hz, 1 H) 7.46 (dd, $J=8.81, 2.71$ Hz, 1 H) 7.69 (d, $J=2.71$ Hz, 1 H); LCMS (ESI⁺) m/z 441

(M+H)⁺; Anal. Calculated for C₂₀H₂₅ClN₂O₅S: C, 54.48; H, 5.71; N, 6.35. Found: C, 54.54; H, 5.38; N, 6.43.

Example 265

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-

5 3-methoxy-2-naphthamide

Commercially available 3-methoxy-2-naphthoic acid (Aldrich) and Example 240C were processed using the method described in Example 240E to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.32 (s, 9 H), 1.58 - 1.74 (m, 1 H), 1.75 - 1.88 (m, 2 H), 1.88 - 1.97 (m, 1 H), 3.60 - 3.71 (m, 1 H), 3.74 - 3.85 (m, 1 H), 3.88 (s, 3 H), 4.15 - 4.26 (m, 2 H), 4.27 - 4.41 (m, 1 H), 7.24 (s, 1 H), 7.31 - 7.43 (m, 2 H), 7.45 - 7.55 (m, 1 H), 7.86 (dd, J=13.6, 8.5 Hz, 2 H), 8.12 (s, 1 H); MS (ESI⁺) m/z 425 (M+H)⁺; Anal. Calculated for C₂₀H₂₃ClN₂O₄S·0.2C₄H₈O₂·0.2H₂O: C, 66.82; H, 6.78; N, 6.28. Found: C, 66.70; H, 6.65; N, 6.33.

15 Example 266

N-[(2Z)-5-tert-butyl-3-[(3-methyloxetan-3-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

To a solution of Example 244A (0.75g, 2.31 mmol) in N,N-dimethylformamide/tetrahydrofuran (1:4, 20 mL) were added potassium tert-butoxide (0.39 g, 3.46 mmol), tetrabutylammonium iodide (0.09 mg, 0.23 mmol) and commercially available 3-(chloromethyl)-3-methyloxetane (TCI, 0.28 g, 2.31 mmol). The reaction mixture was stirred at 80 °C for 16 hours, cooled, diluted with ethyl acetate (20 mL) and quenched with saturated aqueous NaHCO₃ (20 mL). The aqueous layer was extracted with ethyl acetate (2 x 20 mL). The combined organic layers were washed with water (1 x 25 mL). The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. The residue was purified by column chromatography using an Analogix® Intelliflash280TM (SiO₂, 0-100 % ethyl acetate in hexanes) to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.24 (s, 3 H), 1.33 (s, 9 H), 3.77 (s, 3 H), 4.19 (d, J=6.1 Hz, 2 H), 4.38 (s, 2 H), 4.69 (d, J=6.1 Hz, 2 H), 7.10 (d, J=9.2 Hz, 1 H), 7.29 (s, 1 H), 7.44 (dd, J=9.0, 2.9 Hz, 1 H), 7.58 (d, J=2.7 Hz, 1 H); MS (ESI⁺) m/z 409 (M+H)⁺; Anal. Calculated for C₂₀H₂₅ClN₂O₃S: C, 58.74; H, 6.16; N, 6.85. Found: C, 58.92; H, 6.04; N, 6.84.

Example 267

5-chloro-2-methoxy-N-[(2Z)-3-(tetrahydrofuran-2-ylmethyl)-3,4,5,6-tetrahydro-2H-cyclopenta[d][1,3]thiazol-2-ylidene]benzamide

Example 267A5 5,6-dihydro-4H-cyclopenta[d]thiazol-2-amine

A mixture of 2-chlorocyclopentanone (5.0 g, 40 mmol) and thiourea (3.0 g, 40 mmol) was heated at 70 °C for 3 hours. After cooling, the solid was triturated with ethanol and collected by filtration to afford the title compound. MS (ESI) m/z 141 (M+H)⁺.

10 Example 267B

Example 267A and 5-chloro-2-methoxybenzoic acid were processed using the method described in Example 223A to afford the title compound. MS (ESI) m/z 309 (M+H)⁺.

Example 267C15 5-chloro-2-methoxy-N-[(2Z)-3-(tetrahydrofuran-2-ylmethyl)-3,4,5,6-tetrahydro-2H-cyclopenta[d][1,3]thiazol-2-ylidene]benzamide

Example 267B (150mg, 0.49 mmol) in tetrahydrofuran/N,N-dimethylformamide (2:1)(10 mL) was treated with NaH (60%) (24 mg, 0.58 mmol). After 10 minutes, 2-(bromomethyl)tetrahydrofuran (96 mg, 0.58 mmol) was added and the mixture was heated at 95 °C for 12 hours. After cooling to ambient temperature, the mixture was diluted with water, and extracted with ethyl acetate. The organic extract was dried (Na₂SO₄), filtered and concentrated. Purification by chromatography afforded the title compound. ¹H NMR (400 MHz, dimethylsulfoxide-*d*₆) δ ppm 1.62 - 1.73 (m, 1 H) 1.79 - 1.88 (m, 2 H) 1.91 - 2.02 (m, 1 H) 2.35 - 2.44 (m, 2 H) 2.77 - 2.91 (m, 4 H) 3.64 (dd, *J*=14.7, 7.1 Hz, 1 H) 3.74 - 3.81 (m, 1 H) 3.79 (s, 3 H) 4.00 (dd, *J*=13.5, 8.0 Hz, 1 H) 4.23 (dd, *J*=13.5, 3.7 Hz, 1 H) 4.26 - 4.33 (m, 1 H) 7.11 (d, *J*=8.9 Hz, 1 H) 7.44 (dd, *J*=8.9, 2.8 Hz, 1 H) 7.68 (d, *J*=2.8 Hz, 1 H); MS (ESI) m/z 393 (M+H)⁺.

Example 26830 5-chloro-2-methoxy-N-[(2Z)-3-(tetrahydro-2H-pyran-4-ylmethyl)-3,4,5,6-tetrahydro-2H-cyclopenta[d][1,3]thiazol-2-ylidene]benzamide

Example 267B and 4-(bromomethyl)tetrahydro-2H-pyran were processed using the method described in Example 267C to afford the title compound. ¹H NMR

(500 MHz, dimethylsulfoxide- d_6) δ ppm 1.33 (ddd, $J=24.7, 11.90, 4.3$ Hz, 2 H) 1.40 - 1.54 (m, 2 H) 1.83 - 1.86 (m, 1 H) 2.15 - 2.26 (m, 1 H) 2.36 - 2.45 (m, 2 H) 2.82 (dt, $J=20.8, 6.7$ Hz, 4 H) 3.26 (td, $J=11.9, 1.8$ Hz, 1 H) 3.80 (s, 3 H) 3.84 (dd, $J=11.6, 2.4$ Hz, 2 H) 3.99 (d, $J=7.3$ Hz, 2 H) 7.11 (d, $J=8.9$ Hz, 1 H) 7.45 (dd, $J=8.9, 2.8$ Hz, 1 H) 7.70 (d, $J=2.8$ Hz, 1 H); MS (ESI) m/z 407 (M+H)⁺.

Example 269

N-[(2Z)-4,5-dimethyl-3-(tetrahydro-2H-pyran-4-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2,2-dimethyltetrahydro-2H-pyran-4-carboxamide

Example 269A

10 A mixture of 4,5-dimethylthiazol-2-amine (Aldrich) and 2,2-dimethyltetrahydro-2H-pyran-4-carboxylic acid (Chembridge Building Block Library) were processed using the method described in Example 223A to afford the title compound. MS (ESI) m/z 269 (M+H)⁺.

Example 269B

15 N-[(2Z)-4,5-dimethyl-3-(tetrahydro-2H-pyran-4-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2,2-dimethyltetrahydro-2H-pyran-4-carboxamide

Example 269A and 4-(bromomethyl)tetrahydro-2H-pyran were processed using the method described in Example 267C to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide d_6) δ ppm 1.11 - 1.14 (m, 3 H) 1.15 - 1.18 (m, 3 H) 1.30 - 1.43 (m, 4 H) 1.43 - 1.56 (m, 2 H) 1.68 - 1.80 (m, 2 H) 2.14 - 2.18 (m, 3 H) 2.19 - 2.23 (m, 3 H) 2.55 - 2.71 (m, 1 H) 3.16 - 3.26 (m, 2 H) 3.35 - 3.41 (m, 1 H) 3.60 (dd, $J=11.9, 2.4$ Hz, 1 H) 3.63 - 3.68 (m, 1 H) 3.79 - 3.83 (m, 1 H) 3.83 - 3.87 (m, 1 H) 4.01 - 4.03 (m, 1 H) 4.03 - 4.06 (m, 1 H); MS (ESI) m/z 376 (M+H)⁺.

Example 270

25 N-[(2Z)-5-tert-butyl-3-(oxetan-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 244A and the product from Example 211A were processed using the method described in Example 244B to afford the title compound. ¹H NMR (400 MHz, CDCl₃) δ ppm 1.35 (s, 9 H) 2.39 - 2.56 (m, 1 H) 2.68 - 2.83 (m, 1 H) 3.86 - 3.90 (s, 3 H) 4.34 - 4.38 (m, 1 H) 4.37 - 4.43 (m, 1 H) 4.55 (dd, $J=14.1, 5.8$ Hz, 1 H) 4.62 - 4.68 (m, 1 H) 5.15 - 5.22 (m, 1 H) 6.89 (d, $J=8.9$ Hz, 1 H) 6.93 - 6.94 (m, 1 H) 7.31 (dd, $J=8.9, 2.8$ Hz, 1 H) 7.89 (d, $J=2.8$ Hz, 1 H); MS (ESI) m/z 395 (M+H)⁺.

Example 271

N-[(2Z)-5-tert-butyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-
5-chloro-2-methoxybenzamide

Example 244A and 2-(bromomethyl)tetrahydro-2H-pyran were processed using the method described in Example 246 to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.20 - 1.34 (m, 2 H) 1.34 (s, 9 H) 1.51 - 1.55 (m, 2 H) 1.71 (dt, *J*=12.8, 1.8 Hz, 1 H) 1.85 - 1.92 (m, 1 H) 3.39 (td, *J*=14.3, 11.3, 3.4 Hz, 1 H) 3.70 - 3.77 (m, 1 H) 3.90 (s, 3 H) 3.94 - 4.00 (m, 2 H) 4.40 (dd, *J*=14.0, 3.0 Hz, 1 H) 6.78 (s, 1 H) 6.90 (d, *J*=8.9 Hz, 1 H) 7.32 (dd, *J*=8.9, 2.8 Hz, 1 H) 7.95 (d, *J*=2.8 Hz, 1 H); MS (ESI) *m/z* 423 (M+H)⁺.

10

Example 272

N-[(2Z)-5-tert-butyl-3-(1,4-dioxan-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-
2-methoxybenzamide

Example 244A and 2-(iodomethyl)-1,4-dioxane were processed using the method described in Example 246 to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.34 (s, 9 H) 3.32 (dd, *J*=11.60, 10.1 Hz, 1 H) 3.52 - 3.62 (m, 1 H) 3.68 - 3.76 (m, 2 H) 3.80 (dd, *J*=11.0, 2.8 Hz, 1 H) 3.85 - 3.91 (m, 1 H) 3.90 (s, 3 H) 4.00 - 4.06 (m, 1 H) 4.05 (dd, *J*=19.5, 6.71 Hz, 1 H) 4.33 (dd, *J*=10.7, 3.1 Hz, 1 H) 6.74 (s, 1 H) 6.90 (d, *J*=8.9 Hz, 1 H) 7.33 (dd, *J*=8.9, 2.8 Hz, 1 H) 7.92 (d, *J*=2.8 Hz, 1 H); MS (ESI) *m/z* 425 (M+H)⁺.

20

Example 273

N-[(2Z)-5-tert-butyl-3-{{(4S)-2,2-dimethyl-1,3-dioxolan-4-yl}methyl}-1,3-thiazol-
2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 244A and (*R*)-(2,2-dimethyl-1,3-dioxolan-4-yl)methyl 4-methylbenzenesulfonate were processed using the method described in Example 246 to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.34 (s, 9 H) 1.36 (s, 3 H) 1.37 (s, 3 H) 3.76 (dd, *J*=8.9, 6.7 Hz, 1 H) 3.90 (s, 3 H) 4.11 (dd, *J*=8.9, 6.4 Hz, 1 H) 4.31 - 4.42 (m, 2 H) 4.47 - 4.53 (m, 1 H) 6.80 (s, 1 H) 6.90 (d, *J*=8.9 Hz, 1 H) 7.32 (dd, *J*=8.9, 2.75 Hz, 1 H) 7.90 (d, *J*=3.1 Hz, 1 H); MS (ESI) *m/z* 439 (M+H)⁺.

25

Example 274

N-[(2Z)-5-tert-butyl-3-{{(4R)-2,2-dimethyl-1,3-dioxolan-4-yl}methyl}-1,3-thiazol-
2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 244A and (*S*)-(2,2-dimethyl-1,3-dioxolan-4-yl)methyl 4-methylbenzenesulfonate were processed using the method described in Example 246

30

to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.35 (s, 9 H) 1.36 (s, 3 H) 1.37 (s, 3 H) 3.76 (dd, *J*=8.9, 6.71 Hz, 1 H) 3.90 (s, 3 H) 4.11 (dd, *J*=8.9, 6.4 Hz, 1 H) 4.33 - 4.43 (m, 2 H) 4.48 - 4.54 (m, 1 H) 6.81 (s, 1 H) 6.90 (d, *J*=8.9 Hz, 1 H) 7.33 (dd, *J*=8.9, 2.8 Hz, 1 H) 7.89 (d, *J*=2.8 Hz, 1 H); MS (ESI) *m/z* 439 (M+H)⁺.

5

Example 275

N-[(2Z)-5-tert-butyl-3-(tetrahydrofuran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 244A and the product from Example 162A were processed using the method described in Example 246 to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.35 (s, 9 H) 1.71 - 1.79 (m, 1 H) 1.98 - 2.10 (m, 1 H) 2.90 - 3.02 (m, 1 H) 3.64 (dd, *J*=9.15, 5.19 Hz, 1 H) 3.76 - 3.83 (m, 2 H) 3.90 (s, 3 H) 3.95 - 4.01 (m, 1 H) 4.08 (dd, *J*=13.43, 7.93 Hz, 1 H) 4.24 (dd, *J*=13.12, 7.32 Hz, 1 H) 6.63 (s, 1 H) 6.91 (d, *J*=8.85 Hz, 1 H) 7.33 (dd, *J*=8.85, 3.05 Hz, 1 H) 7.97 (d, *J*=3.05 Hz, 1 H); MS (ESI) *m/z* 409 (M+H)⁺.

15

Example 276

N-[(2Z)-5-tert-butyl-3-(tetrahydro-2H-pyran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 244A and the product from Example 225A were processed using the method described in Example 246 to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.34 (s, 9 H) 1.39 - 1.48 (m, 1 H) 1.56 - 1.67 (m, 1 H) 1.69 - 1.77 (m, 1 H) 1.78 - 1.89 (m, 1 H) 2.23 - 2.34 (m, 1 H) 3.34 (dd, *J*=11.0, 8.2 Hz, 1 H) 3.53 (ddd, *J*=9.2, 3.1 Hz, 1 H) 3.74 - 3.85 (m, 2 H) 3.90 (s, 3 H) 4.04 - 4.16 (m, 2 H) 6.60 (s, 1 H) 6.90 (d, *J*=8.9 Hz, 1 H) 7.33 (dd, *J*=8.9, 2.75 Hz, 1 H) 7.98 (d, *J*=2.8 Hz, 1 H); MS (ESI) *m/z* 423 (M+H)⁺.

25

Example 277

N-[(2Z)-5-tert-butyl-3-[(2S)-5-oxotetrahydrofuran-2-yl]methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 277A

(S)-(5-oxotetrahydrofuran-2-yl)methyl 4-methylbenzenesulfonate

30

A mixture of (S)-5-(hydroxymethyl)tetrahydrofuran-2(3H)-one, para-toluenesulfonyl chloride and pyridine were processed using the method described in Example 162A to afford the title compound. MS (ESI) *m/z* 288 (M+18)⁺.

Example 277B

N-[(2Z)-5-tert-butyl-3-[(2S)-5-oxotetrahydrofuran-2-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 244A and the product from Example 277A were processed using the method described in Example 246 to afford the title compound. ¹H NMR (400 MHz, CDCl₃) δ ppm 1.34 (s, 9 H) 2.07 - 2.20 (m, 1 H) 2.34 - 2.48 (m, 2 H) 2.49 - 2.61 (m, 1 H) 3.90 (s, 3 H) 4.39 (dd, J=14.4, 6.1 Hz, 1 H) 4.55 (dd, J=14.4, 3.1 Hz, 1 H) 4.92 - 5.01 (m, 1 H) 6.75 (s, 1 H) 6.92 (d, J=8.9 Hz, 1 H) 7.35 (dd, J=8.9, 2.8 Hz, 1 H) 7.92 (d, J=2.8 Hz, 1 H); MS (ESI) m/z 423 (M+H)⁺.

Example 278

N-[(2Z)-5-acetyl-4-methyl-3-[(2R)-tetrahydrofuran-2-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-1-benzofuran-5-carboxamide

Example 278A

(R)-1-((tetrahydrofuran-2-yl)methyl)thiourea

To a 0 °C solution of (R)-(tetrahydrofuran-2-yl)methanamine (5.0 g, 49.5 mmol) and triethylamine (690 uL, 4.95 mmol) in tetrahydrofuran (100 mL) was added carbon disulfide (5.65 g, 74.3 mmol). Stirring was continued for 0.5 hour followed by the dropwise addition of 30% hydrogen peroxide (5.6 g, 49.5 mmol) so that the temperature was maintained below 40 °C. The reaction mixture was poured into water, and extracted with ethyl acetate. The organic extract was dried (Na₂SO₄), filtered and concentrated to afford an oil. The residue was dissolved in tetrahydrofuran, and treated with 7 N ammonia in methanol (14.3 mL, 100 mmol). The precipitate was collected by filtration and washed with water to afford the title compound.

Example 278B

(R)-1-(2-imino-4-methyl-3-((tetrahydrofuran-2-yl)methyl)-2,3-dihydrothiazol-5-yl)ethanone

To a solution of pentane-2,4-dione (451 mg, 4.5 mmol) and the product from Example 278A (786 mg, 4.5 mmol) in tetrahydrofuran (5 mL) was added a mixture of dimethylsulfoxide (0.64 mL, 9.0 mmol) and concentrated HCl (0.75 mL, 9.0 mmol). The reaction mixture was heated at 40 °C for 12 hours. After cooling to ambient temperature, the mixture was diluted with ethyl acetate and washed with saturated aqueous NaHCO₃. The organic extract was dried (Na₂SO₄), filtered and concentrated to afford the title compound. MS (ESI) m/z 241 (M+H)⁺.

Example 278C

N-[(2Z)-5-acetyl-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-1-benzofuran-5-carboxamide

Example 298B and benzofuran-5-carboxylic acid were processed using the method described in Example 223A to afford the title compound. ¹H NMR (400 MHz, CDCl₃) δ ppm 1.73 - 1.84 (m, 1 H) 1.91 - 2.03 (m, 2 H) 2.16 - 2.26 (m, 1 H) 2.50 (s, 3 H) 2.81 (s, 3 H) 3.75 (dd, J=14.12, 6.44 Hz, 1 H) 3.90 (dd, J=14.1, 7.4 Hz, 1 H) 4.16 (dd, J=13.8, 8.0 Hz, 1 H) 4.41 - 4.51 (m, 1 H) 4.71 (dd, J=13.8, 3.4 Hz, 1 H) 6.88 (s, 1 H) 7.55 (d, J=8.6 Hz, 1 H) 7.68 (t, J=2.2 Hz, 1 H) 8.30 (dt, J=8.6, 1.5 Hz, 1 H) 8.57 - 8.59 (m, 1 H); MS (ESI) m/z 385 (M+H)⁺.

Example 279

N-[(2Z)-5-(1-hydroxy-1-methylethyl)-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-1-benzofuran-5-carboxamide

Example 278C (300 mg, 0.78 mmol) in tetrahydrofuran (20 mL) was treated with a solution of methyllithium in diethyl ether (1.56 mL, 1.56 mmol) at -78 °C. The reaction mixture was stirred at -78 °C for 4 hours and quenched with saturated aqueous NH₄Cl, warmed to ambient temperature and extracted with ethyl acetate. The organic extract was dried (Na₂SO₄), filtered and concentrated. Purification by reverse phase HPLC afforded the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.65 (s, 3 H) 1.67 (s, 3 H) 1.74 - 1.84 (m, 1 H) 1.89 - 2.00 (m, 2 H) 1.99 - 2.02 (m, 1 H) 2.12 - 2.22 (m, 1 H) 2.55 (s, 3 H) 3.76 (dd, J=14.3, 7.6 Hz, 1 H) 3.90 (dd, J=15.0, 6.7 Hz, 1 H) 4.06 (dd, J=13.7, 7.3 Hz, 1 H) 4.40 - 4.49 (m, 1 H) 4.64 (dd, J=14.0, 3.7 Hz, 1 H) 6.85 (dd, J=2.1, 0.9 Hz, 1 H) 7.52 (d, J=8.5 Hz, 1 H) 7.66 (d, J=2.1 Hz, 1 H) 8.30 (dd, J=8.5, 1.5 Hz, 1 H) 8.57 (d, J=1.5 Hz, 1 H); MS (ESI) m/z 401 (M+H)⁺.

Example 280

N-[(2Z)-5-acetyl-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-(2,2,2-trifluoroethoxy)benzamide

Example 280A

N-[(2Z)-5-acetyl-4-methyl-3-(((2R)-tetrahydrofuran-2-yl)methyl)thiazol-2(3H)-ylidene)-5-chloro-2-fluorobenzamide

Example 278B and 5-chloro-2-fluorobenzoic acid were processed using the method described in Example 223A to afford the title compound. MS (ESI) m/z 397 (M+H)⁺.

Example 280B

N-[(2Z)-5-acetyl-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-(2,2,2-trifluoroethoxy)benzamide

To the solution of the product from Example 280A (360 mg, 0.91 mmol) in tetrahydrofuran (4 mL) was added 2,2,2-trifluoroethanol (227 mg, 2.27 mmol) and a 1.0 M solution of potassium tert-butoxide in tetrahydrofuran (2.27 mL, 2.27 mmol). The reaction mixture was stirred at room temperature for 24 hours. The mixture was diluted with water, and extracted with ethyl acetate. The organic extract was dried (Na₂SO₄), filtered and concentrated. Purification by reverse phase HPLC afforded the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.64 - 1.75 (m, 1 H) 1.90 - 2.00 (m, 2 H) 2.10 - 2.20 (m, 1 H) 2.51 (s, 3 H) 2.80 (s, 3 H) 3.73 (dd, J=15.9, 7.3 Hz, 1 H) 3.88 (dd, J=15.3, 6.7 Hz, 1 H) 4.07 (dd, J=13.7, 8.2 Hz, 1 H) 4.33 - 4.41 (m, 1 H) 4.47 (dd, J=16.8, 8.2 Hz, 2 H) 4.62 (dd, J=13.7, 2.8 Hz, 1 H) 7.03 (d, J=8.5 Hz, 1 H) 7.39 (dd, J=8.5, 2.8 Hz, 1 H) 8.02 (d, J=2.8 Hz, 1 H); MS (ESI) m/z 477 (M+H)⁺.

Example 281

N-[(2Z)-5-tert-butyl-3-[(5-methyltetrahydrofuran-2-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 281Ahex-5-en-2-ol

A solution of hex-5-en-2-one (10 g, 102 mmol) in ether (60 mL) was treated with lithium aluminum hydride (4.0 g, 110 mmol) and the mixture was stirred at room temperature for 1 hour. The reaction mixture was quenched with saturated aqueous NH₄Cl, and extracted with ether. The organic extract was dried (Na₂SO₄), filtered and concentrated to afford the title compound.

Example 281B2-(bromomethyl)-5-methyltetrahydrofuran

A solution of the product from Example 281A (6.9 g, 69 mmol) in CH₂Cl₂ (100 mL) was treated with N-bromosuccinimide (14.7, 83 mmol). The reaction mixture was stirred at room temperature for 48 hours, poured into water, and extracted with CH₂Cl₂. The organic extract was dried (Na₂SO₄), filtered and concentrated to afford the title compound. MS (ESI) m/z 179 (M+H)⁺.

Example 281C

N-[(2Z)-5-tert-butyl-3-[(5-methyltetrahydrofuran-2-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

ylidene]-5-chloro-2-methoxybenzamide

Example 244A and the product from Example 281B were processed using the method described in Example 246 to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.22 (d, *J*=6.1 Hz, 3 H) 1.33 - 1.37 (m, 9 H) 1.46 - 1.54 (m, 2 H) 1.67 - 1.76 (m, 1 H) 1.91 - 2.00 (m, 1 H) 3.90 (s, 3 H) 4.01 - 4.08 (m, 1 H) 4.29 - 4.34 (m, 2 H) 4.39 - 4.46 (m, 1 H) 6.89 - 6.90 (m, 1 H) 6.90 - 6.91 (m, 1 H) 7.32 (dd, *J*=8.5, 3.1 Hz, 1 H) 7.94 (d, *J*=2.8 Hz, 1 H); MS (ESI) *m/z* 423 (M+H)⁺.

Example 282N-[(2Z)-5-tert-butyl-3-[(5,5-dimethyltetrahydrofuran-2-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamideExample 282A2-methylhex-5-en-2-ol

A 0 °C solution of hex-5-en-2-one (10 g, 102 mmol) in ether (200 mL) was treated dropwise with a 3.0 M solution of methylmagnesium iodide in ether (102 mL, 306 mmol) over 20 minutes. The reaction mixture was gradually warmed to room temperature, and stirred for 1 hour, quenched with water, and filtered through Celite (ether wash). The filtrate was concentrated and the resulting residue was distilled (27-30 °C at 5 mm Hg) to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.23 (s, 6 H) 1.54 - 1.62 (m, 2 H) 2.11 - 2.19 (m, 2 H) 4.96 (dq, *J*=11.29, 1.83, 1.22 Hz, 1 H) 5.05 (dq, *J*=17.39, 1.53 Hz, 1 H) 5.80 - 5.91 (m, 1 H).

Example 282B5-(bromomethyl)-2,2-dimethyltetrahydrofuran

The product from 282A was processed using the method described in Example 281B to afford the title compound. ¹H NMR (400 MHz, CDCl₃) δ ppm 1.23 (s, 3 H) 1.29 (s, 3 H) 1.73 - 1.82 (m, 2 H) 1.81 - 1.91 (m, 1 H) 2.10 - 2.22 (m, 1 H) 3.32 (dd, *J*=10.13, 7.06 Hz, 1 H) 3.43 (dd, *J*=9.82, 4.60 Hz, 1 H) 4.16 - 4.27 (m, 1 H).

Example 282CN-[(2Z)-5-tert-butyl-3-[(5,5-dimethyltetrahydrofuran-2-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 244A and the product from Example 282B were processed using the method described in Example 246 to afford the title compound. ¹H NMR (400 MHz, dimethylsulfoxide-*d*₆) δ ppm 1.13 (s, 3 H) 1.17 (s, 3 H) 1.31 (s, 9 H) 1.54 - 1.62 (m, 1 H) 1.64 - 1.72 (m, 1 H) 1.72 - 1.80 (m, 1 H) 1.93 - 2.04 (m, 1 H) 3.78 (s, 3 H) 4.12

(dd, $J=15.34, 6.75$ Hz, 1 H) 4.29 - 4.39 (m, 2 H) 7.10 (d, $J=8.59$ Hz, 1 H) 7.20 (s, 1 H) 7.44 (dd, $J=8.59, 2.76$ Hz, 1 H) 7.64 (d, $J=2.76$ Hz, 1 H); MS (ESI) m/z 437 (M+H)⁺.

Example 283

5 N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-(2-methoxyethoxy)benzamide

Example 283A

Example 240C and 5-chloro-2-fluorobenzoic acid were processed using the method described in Example 223A to afford the title compound. MS (ESI) m/z 397 (M+H)⁺.

Example 283B

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-(2-methoxyethoxy)benzamide

Example 283A and 2-methoxyethanol were processed using the method described in Example 280B to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.36 (s, 9 H) 1.62 - 1.71 (m, 1 H) 1.74 - 1.83 (m, 1 H) 1.84 - 1.93 (m, 1 H) 2.02 - 2.11 (m, 1 H) 3.42 (s, 3 H) 3.77 (m, 3 H) 3.84 (dd, $J=15.0, 6.71$ Hz, 1 H) 4.21 (t, $J=5.2$ Hz, 2 H) 4.22 - 4.31 (m, 2 H) 4.43 (dd, $J=10.4$ Hz, 1 H) 6.90 (s, 1 H) 6.97 (d, $J=8.9$ Hz, 1 H) 7.30 (dd, $J=8.5, 2.8$ Hz, 1 H) 7.89 (d, $J=2.8$ Hz, 1 H) MS (ESI) m/z 477 (M+H)⁺.

Example 284

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-6-chloroquinoline-8-carboxamide

Example 240C and 6-chloroquinoline-8-carboxylic acid were processed using the method described in Example 223A to afford the title compound. ¹H NMR (400 MHz, CDCl₃) δ ppm 1.38 (s, 9 H) 1.63 - 1.74 (m, 1 H) 1.74 - 1.83 (m, 1 H) 1.82 - 1.93 (m, 1 H) 1.98 - 2.08 (m, 1 H) 3.72 - 3.88 (m, 2 H) 4.18 - 4.30 (m, 2 H) 4.40 (dd, $J=13.2, 2.5$ Hz, 1 H) 6.90 (s, 1 H) 7.40 (dd, $J=8.3, 4.0$ Hz, 1 H) 7.82 (d, $J=2.5$ Hz, 1 H) 7.99 (d, $J=2.2$ Hz, 1 H) 8.06 (dd, $J=8.3, 1.84$ Hz, 1 H) 9.02 (dd, $J=4.3, 1.84$ Hz, 1 H); MS (ESI) m/z 430 (M+H)⁺.

Example 285

5-chloro-2-methoxy-N-[(2Z)-5-(1-methylcyclopropyl)-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

Example 285A2-(2-aminothiazol-5-yl)propan-2-ol

To a -78 °C solution of thiazol-2-amine (7.0 g, 69.9 mmol) in tetrahydrofuran (200 mL) was added a 10.0 M solution of butyllithium in hexane (14 mL, 140 mmol).
5 The mixture was stirred at -78 °C for 1 hour and chlorotrimethylsilane (15.2 g, 140 mmol) was added dropwise. The mixture was allowed to warm up to -40 °C, cooled to -78 °C and a 10.0 M solution of butyllithium in hexane (7.0, 70 mmol) was added. After 10 minutes, propan-2-one (8.12 g, 140 mmol) was added and the mixture was stirred for 12 hours. The reaction mixture was quenched with saturated aqueous
10 NH₄Cl and extracted with ether. The organic extract was dried (Na₂SO₄), filtered and concentrated. The residue was purified by column chromatography using an Analogix® Intelliflash280™ (SiO₂, 0-100 % ethyl acetate in hexanes) to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.61 (s, 6 H) 5.04 - 5.19 (brs, 2 H) 6.89 (s, 1 H); MS (ESI) m/z 159 (M+H)⁺.

15

Example 285B(R)-5-(prop-1-en-2-yl)-3-((tetrahydrofuran-2-yl)methyl)thiazol-2(3H)-imine

Example 285A and the product from Example 208A were processed using the method described in Example 240C to afford the title compound. MS (ESI) m/z 225 (M+H)⁺.

20

Example 285C(R,Z)-5-chloro-2-methoxy-N-(5-(prop-1-en-2-yl)-3-((tetrahydrofuran-2-yl)methyl)thiazol-2(3H)-ylidene)benzamide

Example 285B and the product from Example 205B were processed using the method described in Example 223A to afford the title compound. MS (ESI) m/z 393 (M+H)⁺.

25

Example 285D5-chloro-2-methoxy-N-[(2Z)-5-(1-methylcyclopropyl)-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide

A 20 mL vial was charged with 1 mL of CH₂Cl₂ and 1,2-dimethoxyethane
30 (110 mg, 1.22 mmol). The solution was cooled to -10 °C and diethylzinc (151 mg, 1.22 mmol) was added. To this mixture was added dropwise diiodomethane (654 mg, 2.44 mmol). After the addition was complete, the resulting clear solution was stirred for 10 minutes at -10 °C. A solution of Example 285C (80mg, 0.204 mmol) was

added. The mixture was allowed to warm to room temperature, stirred for 16 hours, then concentrated. Purification by reverse phase preparative HPLC on a Waters Symmetry C8 column (25 mm x 100 mm, 7 μ m particle size) using a gradient of 10% to 100% acetonitrile:0.1% aqueous trifluoroacetic acid over 8 minutes (10 minutes run
 5 time) at a flow rate of 40 mL/minutes) afforded the title compound for Example 285D and the title compound for Example 286. Characterization for Example 285D: ^1H NMR (400 MHz, CDCl_3) δ ppm 0.97 - 1.02 (m, 2 H) 1.02 - 1.08 (m, 2 H) 1.50 (s, 3 H) 1.62 - 1.72 (m, 1 H) 1.82 - 1.91 (m, 1 H) 1.93 - 2.04 (m, 1 H) 2.21 - 2.31 (m, 1 H) 3.72 - 3.86 (m, 2 H) 4.04 (s, 3 H) 4.29 - 4.39 (m, 1 H) 4.48 - 4.61 (m, 1 H) 4.80 - 4.93
 10 (m, 1 H) 7.02 (d, $J=9.2$ Hz, 1 H) 7.54 (dd, $J=8.9, 2.8$ Hz, 1 H) 7.53 - 7.55 (m, 1 H) 8.04 (d, $J=2.8$ Hz, 1 H); MS (ESI) m/z 407 ($\text{M}+\text{H}$) $^+$.

Example 286

5-chloro-N-[(2Z)-5-(1-hydroxy-3-iodo-1-methylpropyl)-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

15 See Example 285D for experimental details: ^1H NMR (500 MHz, CDCl_3) δ ppm 1.61 (d, $J=1.53$ Hz, 3 H) 1.63 - 1.72 (m, 1 H) 1.79 - 1.96 (m, 2 H) 2.03 - 2.14 (m, 2 H) 2.40 - 2.47 (m, 2 H) 3.02 - 3.11 (m, 1 H) 3.19 (dd, $J=17.4, 9.2$ Hz, 1 H) 3.78 (dd, $J=15.0, 7.0$ Hz, 1 H) 3.83 - 3.89 (m, 1 H) 3.91 (s, 3 H) 4.15 - 4.22 (m, 1 H) 4.25 - 4.31 (m, 1 H) 6.91 (d, $J=8.9$ Hz, 1 H) 7.11 (d, $J=5.5$ Hz, 1 H) 7.34 (dd, $J=8.5, 3.1$ Hz, 1 H)
 20 7.96 (d, $J=2.8$ Hz, 1 H); MS (ESI) m/z 550 ($\text{M}+\text{H}$) $^+$.

Example 287

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-[(1-methylcyclopropyl)methoxy]benzamide

The product from 283A and (1-methylcyclopropyl)methanol were processed
 25 using the method described in Example 280B to afford the title compound. ^1H NMR (500 MHz, CDCl_3) δ ppm 0.36 (t, $J=4.8$ Hz, 2 H) 0.55 (t, $J=4.6$ Hz, 2 H) 1.22 (s, 3 H) 1.36 (s, 9 H) 1.61 - 1.70 (m, 1 H) 1.74 - 1.83 (m, 1 H) 1.83 - 1.92 (m, 1 H) 2.01 - 2.10 (m, 1 H) 3.77 (dd, $J=14.3, 7.6$ Hz, 1 H) 3.81 (s, 2 H) 3.84 (dd, $J=15.9, 8.2$ Hz, 1 H) 4.21 (dd, $J=13.4, 6.41$ Hz, 1 H) 4.23 - 4.29 (m, 1 H) 4.42 (dd, $J=13.4, 2.4$ Hz, 1 H)
 30 6.86 (d, $J=9.5$ Hz, 1 H) 6.87 - 6.88 (m, 1 H) 7.26 (dd, $J=7.9, 3.7$ Hz, 1 H) 7.81 (d, $J=2.8$ Hz, 1 H); MS (ESI) m/z 463 ($\text{M}+\text{H}$) $^+$.

Example 288

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-

5-chloro-2-isopropoxybenzamide

Example 283A and propan-2-ol were processed using the method described in Example 280B to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.35 (s, 9 H) 1.35 (d, *J*=5.8 Hz, 6 H) 1.63 - 1.71 (m, 1 H) 1.75 - 1.82 (m, 1 H) 1.83 - 1.92 (m, 1 H) 2.01 - 2.10 (m, 1 H) 3.77 (dd, *J*=14.65, 7.32 Hz, 1 H) 3.85 (dd, *J*=14.7, 6.7 Hz, 1 H) 4.20 (dd, *J*=13.7, 6.4 Hz, 1 H) 4.24 - 4.30 (m, 1 H) 4.42 (dd, *J*=13.4, 2.8 Hz, 1 H) 4.49 - 4.57 (m, 1 H) 6.86 (s, 1 H) 6.89 (d, *J*=8.9 Hz, 1 H) 7.26 (dd, *J*=8.9, 2.8 Hz, 1 H) 7.84 (d, *J*=2.8 Hz, 1 H); MS (ESI) *m/z* 437 (M+H)⁺.

Example 289

10 N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-ethoxybenzamide

Example 283A and ethanol were processed using the method described in Example 280B to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.35 (s, 9 H) 1.44 (t, *J*=7.02 Hz, 3 H) 1.62 - 1.73 (m, 1 H) 1.75 - 1.83 (m, 1 H) 1.84 - 1.92 (m, 1 H) 2.01 - 2.11 (m, 1 H) 3.77 (dd, *J*=13.7, 7.6 Hz, 1 H) 3.84 (dd, *J*=14.7, 6.7 Hz, 1 H) 4.13 (dd, *J*=14.0, 7.2 Hz, 2 H) 4.22 (dd, *J*=13.4, 6.0 Hz, 1 H) 4.25 - 4.31 (m, 1 H) 4.41 (dd, *J*=13.4, 2.8 Hz, 1 H) 6.86 (s, 1 H) 6.89 (d, *J*=8.9 Hz, 1 H) 7.29 (dd, *J*=8.9, 2.8 Hz, 1 H) 7.91 (d, *J*=2.8 Hz, 1 H); MS (ESI) *m/z* 423 (M+H)⁺.

Example 290

20 N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-(tetrahydrofuran-3-yloxy)benzamide

Example 283A and tetrahydrofuran-3-ol were processed using the method described in Example 280B to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm ¹H NMR (500 MHz, CDCl₃) δ ppm 1.36 (s, 9 H) 1.61 - 1.70 (m, 1 H) 1.76 - 1.83 (m, 1 H) 1.84 - 1.93 (m, 1 H) 2.03 - 2.10 (m, 1 H) 2.11 - 2.25 (m, 2 H) 3.77 (dd, *J*=14.0, 7.3 Hz, 1 H) 3.82 - 3.91 (m, 2 H) 3.98 (dd, *J*=15.3, 8.5 Hz, 1 H) 4.02 (d, *J*=3.7 Hz, 2 H) 4.20 (ddd, *J*=6.4, 1.5 Hz, 1 H) 4.23 - 4.30 (m, 1 H) 4.41 (dd, *J*=13.7, 3.1 Hz, 1 H) 4.88 - 5.00 (m, 1 H) 6.83 (dd, *J*=8.9, 0.6 Hz, 1 H) 6.88 (d, *J*=0.9 Hz, 1 H) 7.28 (dd, *J*=8.9, 2.75 Hz, 1 H) 7.87 (t, *J*=3.1 Hz, 1 H); MS (ESI) *m/z* 423 (M+H)⁺.

Example 291

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-[(2-methoxyethyl)(methyl)amino]benzamide

A mixture of the product from Example 283A (120 mg, 0.3 mmol), 2-methoxy-N-methylethanamine (54 mg, 0.6 mmol) and triethylamine (127 μ L, 0.91 mmol) in tetrahydrofuran (1 mL) was heated at 120 °C in a microwave (CEM) for 2 hours. The reaction mixture was diluted with water and extracted with ethyl acetate. The organic extract was dried (Na_2SO_4), filtered, and concentrated. Purification by reverse phase preparative HPLC on a Waters Symmetry C8 column (25 mm x 100 mm, 7 μ m particle size) using a gradient of 10% to 100% acetonitrile:0.1% aqueous trifluoroacetic acid over 8 minutes (10 minutes run time) at a flow rate of 40 mL/minutes) to afford the title compound.

¹H NMR (500 MHz, CDCl_3) δ ppm 1.36 (s, 9 H) 1.62 - 1.71 (m, 1 H) 1.78 - 1.86 (m, 1 H) 1.86 - 1.93 (m, 1 H) 2.01 - 2.11 (m, 1 H) 2.89 (s, 3 H) 3.26 (s, 3 H) 3.29 (t, $J=6.1$ Hz, 2 H) 3.54 (t, $J=6.4$ Hz, 2 H) 3.78 (dd, $J=14.7, 7.3$ Hz, 1 H) 3.85 (dd, $J=15.0, 8.2$ Hz, 1 H) 4.18 (dd, $J=13.7, 6.4$ Hz, 1 H) 4.23 - 4.30 (m, 1 H) 4.40 (dd, $J=13.7, 3.1$ Hz, 1 H) 6.85 (s, 1 H) 6.92 (d, $J=8.9$ Hz, 1 H) 7.20 (dd, $J=8.9, 2.8$ Hz, 1 H) 7.67 (d, $J=2.4$ Hz, 1 H); MS (ESI) m/z 466 ($\text{M}+\text{H}$)⁺.

Example 292

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-(difluoromethoxy)benzamide

Example 240C and 5-chloro-2-(difluoromethoxy)benzoic acid were processed using the method described in Example 223A to afford the title compound.

¹H NMR (500 MHz, CDCl_3) δ ppm 1.36 (s, 9 H) 1.60 - 1.69 (m, 1 H) 1.76 - 1.84 (m, 1 H) 1.85 - 1.93 (m, 1 H) 2.02 - 2.12 (m, 1 H) 3.78 (dd, $J=13.7, 7.3$ Hz, 1 H) 3.85 (dd, $J=15.0, 6.7$ Hz, 1 H) 4.20 (dd, $J=13.4, 6.4$ Hz, 1 H) 4.24 - 4.30 (m, 1 H) 4.44 (dd, $J=13.7, 2.8$ Hz, 1 H) 6.73 (t, $J=6.3$ Hz, 1 H) 6.91 (s, 1 H) 7.17 (d, $J=8.5$ Hz, 1 H) 7.38 (dd, $J=8.5, 2.8$ Hz, 1 H) 8.05 (d, $J=2.8$ Hz, 1 H); MS (ESI) m/z 445 ($\text{M}+\text{H}$)⁺.

Example 293

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-(trifluoromethoxy)benzamide

Example 240C and 5-chloro-2-(trifluoromethoxy)benzoic acid were processed using the method described in Example 223A to afford the title compound.

¹H NMR (500 MHz, CDCl_3) δ ppm 1.36 (s, 9 H) 1.58 - 1.67 (m, 1 H) 1.75 - 1.83 (m, 1 H) 1.84 - 1.93 (m, 1 H) 2.01 - 2.11 (m, 1 H) 3.77 (dd, $J=13.7, 6.4$ Hz, 1 H) 3.84 (dd, $J=15.0, 6.7$ Hz, 1 H) 4.18 (dd, $J=13.7, 6.7$ Hz, 1 H) 4.22 - 4.29 (m, 1 H) 4.46 (dd, $J=13.7, 2.8$

Hz, 1 H) 6.91 (s, 1 H) 7.23 (dd, $J=8.5$, 1.2 Hz, 1 H) 7.40 (dd, $J=8.9$, 2.8 Hz, 1 H) 8.06 (d, $J=2.8$ Hz, 1 H); MS (ESI) m/z 463 (M+H)⁺.

Example 294

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-

5 5-chloro-2-(2,2,2-trifluoroethoxy)benzamide

The product from 283A and 2,2,2-trifluoroethanol were processed using the method described in Example 280B to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.37 (s, 9 H) 1.60 - 1.69 (m, 1 H) 1.76 - 1.84 (m, 1 H) 1.85 - 1.93 (m, 1 H) 2.02 - 2.11 (m, 1 H) 3.78 (dd, $J=13.7$, 7.3 Hz, 1 H) 3.85 (dd, $J=15.0$, 7.0 Hz, 1 H) 4.21 (dd, $J=13.4$, 6.4 Hz, 1 H) 4.24 - 4.29 (m, 1 H) 4.43 (dd, $J=11.0$, 2.8 Hz, 1 H) 4.47 (dd, $J=17.1$, 8.5 Hz, 2 H) 6.91 (s, 1 H) 7.01 (d, $J=8.9$ Hz, 1 H) 7.33 (dd, $J=8.5$, 2.8 Hz, 1 H) 7.99 (d, $J=2.8$ Hz, 1 H); MS (ESI) m/z 477 (M+H)⁺.

Example 295

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-

15 5-chloro-2-[3-(dimethylamino)propoxy]benzamide

The product from 283A and 3-(dimethylamino)propan-1-ol were processed using the method described in Example 280B to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.38 (s, 9 H) 1.62 - 1.71 (m, 1 H) 1.84 - 1.91 (m, 1 H) 1.91 - 1.98 (m, 1 H) 2.09 - 2.19 (m, 1 H) 2.29 - 2.36 (m, 2 H) 3.04 - 3.09 (m, $J=4.9$ Hz, 2 H) 3.81 (dd, $J=13.7$, 6.4 Hz, 1 H) 3.87 (dd, $J=15.0$, 7.0 Hz, 1 H) 4.25 - 4.34 (m, 4 H) 4.51 (d, $J=12.8$ Hz, 1 H) 6.93 (d, $J=9.2$ Hz, 1 H) 7.00 - 7.02 (m, 1 H) 7.44 (dd, $J=8.9$, 2.9 Hz, 1 H) 8.20 (d, $J=2.8$ Hz, 1 H) 10.35 - 10.51 (m, 1 H); MS (ESI) m/z 480 (M+H)⁺.

Example 296

25 5,6-dichloro-N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]quinoline-8-carboxamide

A mixture of Example 186A (145 mg), 5,6-dichloroquinoline-8-carboxylic acid (99.65 mg) (Bailey J. Heterocycl. Chem. 1974, 11, 229), 1-hydroxybenzotriazole hydrate (55 mg), 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (79 mg) and triethylamine (0.14 mL) in tetrahydrofuran (3 mL) was heated at 70 °C on a shaker overnight, cooled, quenched in saturated NaHCO₃ and extracted with ethyl acetate (2x). The organic extracts were dried (Na₂SO₄), filtered and the solvent was evaporated. The crude material was purified by reverse phase preparative HPLC

on a Waters Symmetry C8 column (25 mm x 100 mm, 7 μ m particle size) using a gradient of 10% to 100% acetonitrile:10 mM ammonium acetate over 8 minutes (10 minutes run time) at a flow rate of 40 mL/minutes) to afford the title compound. ^1H NMR (300 MHz, CDCl_3) δ ppm 1.62-1.75 (m, 1H), 1.85-1.95 (m, 2H), 2.0-2.1 (m, 1H), 2.26 (s, 3H), 2.30 (s, 3H), 3.70-3.77 (m, 1H), 3.83-3.90 (m, 1H), 3.97 (dd, 1H), 4.35-4.42 (m, 1H), 4.55 (dd, 1H), 7.53 (dd, 1H), 8.60 (dd, 1H), 9.07 (m, 1H); MS (DCI/ NH_3) m/z 436, 438 ($\text{M}+\text{H}$) $^+$.

Example 297

6-chloro-N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]quinoline-8-carboxamide

A mixture of Example 186A (106 mg), 6-chloro-quinoline-8-carboxylic acid (73 mg) (Weyer et al, *Arzneim. Forsch* 1974, 24, 269), 1-hydroxybenzotriazole hydrate (47 mg), 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (69 mg), and triethylamine (0.15 mL) in tetrahydrofuran (2 mL) was heated to 70 $^\circ\text{C}$ on a shaker overnight, cooled, poured into saturated NaHCO_3 and extracted with ethyl acetate (2x). The organic extracts were dried (Na_2SO_4), filtered and the solvent was evaporated. The crude material was dissolved in warm methanol and allowed to cool overnight. The solid precipitate was discarded, the filtrate was concentrated to dryness, triturated with cold methanol, and collected to afford the title compound. ^1H NMR (300 MHz, dimethylsulfoxide- d_6) δ ppm 1.56-1.64 (m, 1H), 1.70-1.95 (m, 3H), 2.26 (s, 3H), 2.27 (s, 3H), 3.57-3.65 (m, 1H), 3.73-3.80 (m, 1H), 3.97-4.05 (m, 1H), 4.24-4.35 (m, 2H), 7.60 (dd, 1H), 7.78 (d, 1H), 8.15 (d, 1H), 8.38 (dd, 1H), 8.91 (dd, 1H); MS (DCI/ NH_3) m/z 402 ($\text{M}+\text{H}$) $^+$.

Example 298

6-chloro-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]quinoline-8-carboxamide

A mixture of Example 149A (142 mg), 6-chloro-quinoline-8-carboxylic acid (105 mg) (Weyer et al, *Arzneim. Forsch* 1974, 24, 269), 1-hydroxybenzotriazole hydrate (67 mg), 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (94 mg), and triethylamine (0.15 mL) in tetrahydrofuran (3 mL) was heated to 70 $^\circ\text{C}$ for 24 hours, cooled, poured into saturated NaHCO_3 and extracted with ethyl acetate (2x). The organic extracts were dried (Na_2SO_4), filtered and the solvent was evaporated. The crude material was purified by reverse phase preparative HPLC on a

Waters Symmetry C8 column (25 mm x 100 mm, 7 μ m particle size) using a gradient of 10% to 100% acetonitrile: 10 mM ammonium acetate over 8 minutes (10 minutes run time) at a flow rate of 40 mL/minutes) to afford the title compound. ^1H NMR (300 MHz, CDCl_3) δ ppm 1.65-1.72 (m, 1H), 1.80-1.95 (m, 2H), 2.0-2.1 (m, 1H), 2.34 (d, $J = 1.4$ Hz, 3H), 3.74-3.90 (m, 2H), 4.14 (dd, 1H), 4.2-4.3 (m, 1H), 4.47 (dd, 1H), 6.95 (q, $J = 1.4$ Hz, 1H), 7.42 (dd, 1H), 7.84 (d, 1H), 8.03 (d, 1H), 8.08 (dd, 1H), 9.04 (dd, 1H); MS (DCI/ NH_3) m/z 388 ($\text{M}+\text{H}$) $^+$.

Example 299

5,6-dichloro-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]quinoline-8-carboxamide

A mixture of Example 149A (117 mg), 5,6-dichloroquinoline-8-carboxylic acid (71 mg), 1-hydroxybenzotriazole hydrate (51 mg), 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (79 mg) and triethylamine (0.15 mL) in tetrahydrofuran was processed and purified according to the method of Example 298 to afford the title compound. ^1H NMR (300 MHz, CDCl_3) δ ppm 1.6-1.71 (m, 1H), 1.78-1.95 (m, 2H), 2.0-2.1 (m, 1H), 2.34 (d, $J = 1.4$ Hz, 3H), 3.72-3.90 (m, 2H), 4.12 (dd, 1H), 4.2-4.3 (m, 1H), 4.46 (dd, 1H), 6.94 (q, $J = 1.4$ Hz, 1H), 7.53 (dd, 1H), 8.60 (dd, 1H), 9.06 (dd, 1H); MS (DCI/ NH_3) m/z 422 ($\text{M}+\text{H}$) $^+$.

Example 300

3-methoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-naphthamide

A mixture of Example 149A (113 mg), 3-methoxy-2-naphthoic acid (Aldrich) (69 mg), 1-hydroxybenzotriazole hydrate (50 mg), 1-ethyl-3-[3-(dimethylamino)propyl]-carbodiimide hydrochloride (79 mg) and triethylamine (0.15 mL) in tetrahydrofuran (2 mL) was shaken over the weekend at room temperature, poured into saturated NaHCO_3 and extracted with ethyl acetate (2x). The organic extracts were dried (Na_2SO_4), filtered and the solvent was evaporated. The crude material was purified by flash chromatography over silica gel eluting with ethyl acetate:hexane (7:3) to afford the title compound. ^1H NMR (300 MHz, CDCl_3) δ ppm 1.64-1.78 (m, 1H), 1.80-1.95 (m, 2H), 2.02-2.15 (m, 1H), 2.32 (d, $J = 1.4$ Hz, 3H), 3.74-3.92 (m, 2H), 4.01 (s, 3H), 4.2 (dd, 1H), 4.27-4.35 (m, 1H), 4.51 (d, 1H), 6.91 (br s, 1H), 7.20 (s, 1H), 7.33 (m, 1H), 7.46 (m, 1H), 7.73 (d, 1H), 7.81 (d, 1H), 8.42 (s, 1H); MS (DCI/ NH_3) m/z 383 ($\text{M}+\text{H}$) $^+$.

Example 301

N-[(2Z)-5-tert-butyl-3-[(cis)-(3-methoxycyclobutyl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 301A5 cis-3-benzyloxymethylcyclobutanol methyl ether

To a solution of cis-3-benzyloxymethylcyclobutanol (Albany Molecular Research Institute, 1.0 g, 5.2 mmol) in 10 mL of tetrahydrofuran at 0 °C was added NaH (0.62 g, 15.6 mmol). The mixture stirred for 15 minutes and iodomethane (0.49 mL, 7.8 mmol) was added and the mixture was allowed to warm to ambient
10 temperature and stirred for 16 hours. Some starting material remained by TLC so additional NaH (0.21, 5.2 mmol) and iodomethane (0.32 mL, 5.2 mmol) were added and the mixture stirred for an additional 2 hours. The mixture was quenched with 10 mL of NH₄Cl and diluted with 10 mL of ethyl acetate. The layers were separated and the aqueous layer was extracted twice with 5 mL portions of ethyl acetate. The
15 combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 75% hexanes in ethyl acetate) afforded the title compound. MS (DCI/NH₃) m/z 207 (M+H)⁺.

Example 301B20 (cis-3-methoxycyclobutyl)methanol

A solution of Example 301A (1.05 g, 5.2 mmol) in 10 mL of ethanol was degassed and the flask was filled with N₂. This was repeated two additional times. Pd/C (0.1 g, 10 wt%) was added and the mixture was degassed again and flushed with N₂. This was repeated two additional times and the flask was put under 1 atm. of H₂
25 and the mixture was allowed to stir at ambient temperature for 72 hours. The mixture was degassed and the flask was filled with N₂. The reaction mixture was filtered, concentrated under reduced pressure and purified by column chromatography (SiO₂, 25% hexanes in ethyl acetate) to afford the title compound. MS (DCI/NH₃) m/z 134 (M+NH₄)⁺.

30 Example 301C(cis-3-methoxycyclobutyl)methyl 4-methylbenzenesulfonate

Example 301B (0.49 g, 4.2 mmol) and *p*-toluenesulfonyl chloride (0.80 g, 4.2 mmol) in 5 mL of CH₂Cl₂ and 5 mL of pyridine were processed as in Example 203A to afford the title compound. MS (DCI/NH₃) m/z 288 (M+NH₄)⁺.

Example 301D5-tert-butyl-3-((cis-3-methoxycyclobutyl)methyl)thiazol-2(3H)-imine

Example 240A (0.25 g, 1.6 mmol), Example 301C (0.44 g, 1.6 mmol) and tetrabutylammonium iodide (0.30 g, 0.81 mmol) in 0.5 mL of N,N-dimethylformamide were processed as in Example 240C to afford the title compound. MS (DCI/NH₃) m/z 266 (M+H)⁺.

Example 301EN-[(2Z)-5-tert-butyl-3-[(cis)-(3-methoxycyclobutyl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 301D (0.19 g, 0.75 mmol), triethylamine (0.31 mL, 2.2 mmol) and Example 205B (0.75 mmol) in 10 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound (0.105 g, 0.25 mmol, 33% yield). ¹H NMR (300 MHz, CD₃OD) δ ppm 1.38 (s, 9 H), 1.71 - 1.84 (m, 2 H), 2.31 - 2.47 (m, 3 H), 3.21 (s, 3 H), 3.71 - 3.83 (m, 1 H), 3.86 (s, 3 H), 4.29 (d, J=6.4 Hz, 2 H), 7.07 (d, J=8.8 Hz, 1 H), 7.14 (s, 1 H), 7.40 (dd, J=8.8, 3.1 Hz, 1 H), 7.83 (d, J=2.7 Hz, 1 H); MS (DCI/NH₃) m/z 423 (M+H)⁺. Anal. Calculated for C₂₁H₂₇ClN₂O₃S: C, 59.63; H, 6.43; N, 6.62. Found: C, 59.66; H, 6.28; N, 6.44.

Example 302N-[(2Z)-5-tert-butyl-3-[(cis)-(3-hydroxycyclobutyl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamideExample 302A(cis-3-(benzyloxymethyl)cyclobutoxy)(tert-butyl)dimethylsilane

To a solution of cis-3-benzyloxymethylcyclobutanol (Albany Molecular Research Institute, 1.0 g, 5.2 mmol) in 50 mL of CH₂Cl₂ was added imidazole (2.7 g, 39 mmol) followed by *tert*-butyldimethylsilyl chloride (3.9 g, 26 mmol). This mixture stirred at ambient temperature for 2 hours and was quenched with 10 mL of saturated aqueous NH₄Cl. The layers were separated and the aqueous layer was extracted with three 5 mL of portions of CH₂Cl₂. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 75% hexanes in ethyl acetate) afforded the title compound. MS (DCI/NH₃) m/z 307 (M+H)⁺.

Example 302B(cis-3-(tert-butyldimethylsilyloxy)cyclobutyl)methanol

A solution of Example 302A (3.7 g, 12 mmol) in 20 mL of ethanol was degassed and the flask was flushed with N₂. This was repeated two additional times. Pd/C (0.37 g, 10 wt%) was added and the mixture was degassed again and flushed with N₂. This was repeated two additional times then the flask was put under 1 atm. of H₂ and the reaction mixture was allowed to stir at ambient temperature for 48 hours. The mixture was degassed and the flask was filled with N₂ then the reaction mixture was filtered, concentrated under reduced pressure and purified by column chromatography (SiO₂, 50% hexanes in ethyl acetate) to afford the title compound. MS (DCI/NH₃) m/z 217 (M+NH₄)⁺.

10

Example 302C

(cis-3-(tert-butyldimethylsilyloxy)cyclobutyl)methyl 4-methylbenzenesulfonate

Example 302B (1.2 g, 5.5 mmol) and *p*-toluenesulfonyl chloride (1.1 g, 5.5 mmol) in 7 mL of CH₂Cl₂ and 7 mL of pyridine were processed as in Example 203A to afford the title compound. MS (DCI/NH₃) m/z 371 (M+H)⁺, 388 (M+NH₄)⁺.

15

Example 302D

5-tert-butyl-3-((cis-3-(tert-butyldimethylsilyloxy)cyclobutyl)methyl)thiazol-2(3H)-imine

20

Example 240A (0.72 g, 4.6 mmol), Example 302C (1.7 g, 4.6 mmol) and tetrabutylammonium iodide (0.85 g, 2.3 mmol) in 1.5 mL of N,N-dimethylformamide were processed as in Example 240C to afford the title compound. MS (DCI/NH₃) m/z 355 (M+H)⁺.

Example 302E

(Z)-N-(5-tert-butyl-3-((cis-3-(tert-butyldimethylsilyloxy)cyclobutyl)methyl)thiazol-2(3H)-ylidene)-5-chloro-2-methoxybenzamide

25

Example 302D (0.57 g, 1.6 mmol), triethylamine (0.67 mL, 4.8 mmol) and Example 205B (1.6 mmol) in 10 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound. MS (DCI/NH₃) m/z 523 (M+H)⁺.

Example 302F

N-[(2Z)-5-tert-butyl-3-[(cis)-(3-hydroxycyclobutyl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

30

To Example 302E (0.78 g, 1.5 mmol) in 10 mL of tetrahydrofuran at ambient temperature was added tetrabutylammonium fluoride (1 M in tetrahydrofuran, 1.8 mL, 1.8 mmol) dropwise via syringe pump over 30 minutes. The reaction mixture was stirred at ambient temperature for 2 hours and was concentrated under reduced

pressure and purified by column chromatography (SiO₂, 50% hexanes in ethyl acetate then 100% ethyl acetate) to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.38 (s, 9 H), 1.70 - 1.82 (m, 2 H), 2.25 - 2.42 (m, 3 H), 3.86 (s, 3 H), 4.00 - 4.10 (m, 1 H), 4.28 (d, *J*=6.4 Hz, 2 H), 7.07 (d, *J*=9.2 Hz, 1 H), 7.12 (s, 1 H), 7.39 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.80 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 409 (M+H)⁺.

Example 303

N-[(2Z)-5-tert-butyl-3-(((cis)-3-hydroxy-3-methylcyclobutyl)methyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 303A

10 (Z)-N-(5-tert-butyl-3-((3-oxocyclobutyl)methyl)thiazol-2(3H)-ylidene)-5-chloro-2-methoxybenzamide

To Example 302F (0.57 g, 1.4 mmol) in 15 mL of CH₂Cl₂ was added 4-methylmorpholine *N*-oxide (0.82 g, 7.0 mmol) followed by approximately 0.5 g powdered 4 Å molecular sieves. This mixture was stirred at ambient temperature for 15 minutes and was cooled to 0 °C and tetrapropylammonium perruthenate (49 mg, 0.14 mmol) was added in portions over 5 minutes. The mixture was stirred at 0 °C for 5 minutes and was allowed to warm to ambient temperature and stirred for an additional 90 minutes. The mixture was filtered through Celite, concentrated under reduced pressure and purified by column chromatography (SiO₂, 20% hexanes in ethyl acetate) to afford the title compound. MS (DCI/NH₃) *m/z* 407 (M+H)⁺.

Example 303B

N-[(2Z)-5-tert-butyl-3-(((cis)-3-hydroxy-3-methylcyclobutyl)methyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

25 To Example 302A (0.21 g, 0.52 mmol) in 10 mL of tetrahydrofuran at -78 °C was added a 1.6 M solution of methyllithium in diethyl ether (1.0 mL, 1.6 mmol) dropwise over 5 minutes. The mixture was stirred at -78 °C for 2 hours and was slowly warmed to ambient temperature and was allowed to stir for 18 hours. The mixture was quenched with 5 mL of saturated aqueous NH₄Cl and diluted with 5 mL of ethyl acetate. The layers were separated and the aqueous layer was extracted twice 30 with 5 mL of ethyl acetate. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 30% hexanes in ethyl acetate) afforded the title compound.

¹H NMR (300 MHz, CD₃OD) δ ppm 1.32 (s, 3 H), 1.38 (s, 9 H), 1.90 - 2.00 (m, 2 H), 2.06 - 2.16 (m, 2 H), 2.35 - 2.52 (m, 1 H), 3.86 (s, 3 H), 4.30 (d, *J*=7.1 Hz, 2 H), 7.07 (d, *J*=8.8 Hz, 1 H), 7.12 (s, 1 H), 7.39 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.80 (d, *J*=3.1 Hz, 1 H); MS (DCI/NH₃) *m/z* 423 (M+H)⁺. Anal. Calculated for C₂₁H₂₇ClN₂O₃S·0.1H₂O:

5 C, 59.38; H, 6.45; N, 6.59. Found: C, 59.17; H, 6.62; N, 6.28.

Example 304

5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-3-[(*cis*)-3-methoxycyclobutyl)methyl]-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

10

Example 304A

(Z)-N-(5-acetyl-3-[(*cis*)-3-methoxycyclobutyl)methyl]-4-methylthiazol-2(3H)-ylidene)-5-chloro-2-methoxybenzamide

Example 238A (0.57 g, 1.8 mmol), Example 301C (0.48 g, 1.8 mmol) and potassium *tert*-butoxide (0.42 g, 3.5 mmol) were processed as described in the procedure for Example 238B to afford the title compound. MS (DCI/NH₃) *m/z* 423 (M+H)⁺.

15

Example 304B

5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-3-[(*cis*)-3-methoxycyclobutyl)methyl]-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-

20

methoxybenzamide

Example 304A (20 mg, 0.047 mmol) and methyllithium (1.6 M in ether, 88 μL, 0.14 mmol) in 1 mL of tetrahydrofuran were processed as described in Example 239 to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.61 (s, 6 H), 1.80 - 1.87 (m, 2 H), 2.32 - 2.43 (m, 2 H), 2.49 (s, 3 H), 3.20 - 3.24 (m, 1 H), 3.20 (s, 3 H), 3.34 - 3.41 (m, 1 H), 3.68 - 3.78 (m, 1 H), 3.86 (s, 3 H), 4.38 (d, *J*=6.1 Hz, 2 H), 7.06 (d, *J*=8.8 Hz, 1 H), 7.39 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.86 (d, *J*=2.7 Hz, 1 H) MS (DCI/NH₃) *m/z* 439 (M+H)⁺.

25

Example 305

N-[(2Z)-5-*tert*-butyl-3-[2-(2-methoxyethoxy)ethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

30

Example 305A

5-*tert*-butyl-3-(2-(2-methoxyethoxy)ethyl)thiazol-2(3H)-imine hydrobromide

A mixture of Example 240A (0.20 g, 1.3 mmol) and 1-bromo-2-(2-

methoxyethoxy)ethane (0.27 g, 1.4 mmol) was warmed to 85 °C and was allowed to stir for 24 hours. The mixture was cooled to ambient temperature and the crude solids were triturated with ethanol and ether to afford the title compound. MS (DCI/NH₃) m/z 259 (M+H)⁺.

5

Example 305B

N-[(2Z)-5-tert-butyl-3-[2-(2-methoxyethoxy)ethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 305A (0.3 g, 0.88 mmol), triethylamine (0.49 mL, 3.5 mmol) and Example 205B (0.88 mmol) in 10 mL of tetrahydrofuran and 1.5 mL of N,N-dimethylformamide were processed as described in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.38 (s, 9 H), 3.31 (s, 3 H), 3.47 - 3.52 (m, 2 H), 3.59 - 3.65 (m, 2 H), 3.87 (dd, J=5.4 Hz, 2 H), 3.86 (s, 3 H), 4.42 (dd, J=5.1 Hz, 2 H), 7.07 (d, J=8.8 Hz, 1 H), 7.18 (s, 1 H), 7.39 (dd, J=8.8, 2.7 Hz, 1 H), 7.82 (d, J=2.7 Hz, 1 H); MS (DCI/NH₃) m/z 427 (M+H)⁺. Anal. Calculated for C₂₀H₂₇ClN₂O₄S: C, 56.26; H, 6.37; N, 6.56. Found: C, 56.06; H, 5.50; N, 6.43.

15

Example 306

N-[(2Z)-5-tert-butyl-3-(3-methoxypropyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 306A

20

5-tert-butyl-3-(3-methoxypropyl)thiazol-2(3H)-imine

A mixture of Example 240A (0.20 g, 1.3 mmol) and 1-bromo-3-methoxypropane (0.22 g, 1.4 mmol) was warmed to 85 °C and was allowed to stir for 24 hours. The mixture was cooled to ambient temperature, concentrated under reduced pressure and purified via column chromatography (SiO₂, 10% methanol in ethyl acetate then 9:1:0.1 CH₂Cl₂ : methanol : NH₄OH) to afford the title compound. MS (DCI/NH₃) m/z 229 (M+H)⁺.

25

Example 306B

N-[(2Z)-5-tert-butyl-3-(3-methoxypropyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

30

Example 306A (0.25 g, 0.81 mmol), triethylamine (0.34 mL, 2.4 mmol) and Example 205B (0.81 mmol) in 10 mL of tetrahydrofuran and 1 mL of N,N-dimethylformamide were processed as in Example 208D to afford the title compound.

¹H NMR (300 MHz, CD₃OD) δ ppm 1.38 (s, 9 H), 2.05 - 2.17 (m, 2 H), 3.32 (s, 3 H), 3.41 (t, *J*=5.9 Hz, 2 H), 3.86 (s, 3 H), 4.33 (t, *J*=7.0 Hz, 2 H), 7.07 (d, *J*=8.8 Hz, 1 H), 7.11 (s, 1 H), 7.40 (dd, *J*=9.0, 2.9 Hz, 1 H), 7.89 (d, *J*=2.7 Hz, 1 H); MS (DCI/NH₃) *m/z* 397 (M+H)⁺. Anal. Calculated for C₁₉H₂₅ClN₂O₃S: C, 57.49; H, 6.35; N, 7.06.

5 Found: C, 57.18; H, 6.21; N, 6.94.

Example 307

N-[(2Z)-5-tert-butyl-3-(2-ethoxyethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 307A

10 5-tert-butyl-3-(2-ethoxyethyl)thiazol-2(3H)-imine

A mixture of Example 240A (0.17 g, 1.1 mmol) and 2-(bromoethyl)ether (0.20 g, 1.2 mmol) was warmed to 85 °C and was allowed to stir for 24 hours. The mixture was cooled to ambient temperature, concentrated and purified via column chromatography (SiO₂, 10% methanol in ethyl acetate then 9:1:0.1 CH₂Cl₂ : methanol : NH₄OH) to
15 afford the title compound. MS (DCI/NH₃) *m/z* 229 (M+H)⁺.

Example 307B

N-[(2Z)-5-tert-butyl-3-(2-ethoxyethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

20 Example 307A (0.24 g, 1.05 mmol), triethylamine (0.44 mL, 3.2 mmol) and Example 205B (1.05 mmol) in 15 mL of tetrahydrofuran were processed as in Example 208D to afford the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.14 (t, *J*=7.0 Hz, 3 H), 1.38 (s, 9 H), 3.52 (q, *J*=6.9 Hz, 2 H), 3.81 (t, *J*=5.3 Hz, 2 H), 3.86 (s, 3 H), 4.41 (t, *J*=5.3 Hz, 2 H), 7.07 (d, *J*=8.8 Hz, 1 H), 7.14 (s, 1 H), 7.39 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.83 (d, *J*=3.1 Hz, 1 H); MS (DCI/NH₃) *m/z* 397 (M+H)⁺. Anal.
25 Calculated for C₁₉H₂₅ClN₂O₃S: C, 57.49; H, 6.35; N, 7.06. Found: C, 57.34; H, 6.04; N, 6.94.

Example 308

N-[(2Z)-5-tert-butyl-3-(3-hydroxy-3-methylbutyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

30

Example 308A

3-hydroxy-3-methylbutyl 4-methylbenzenesulfonate

To a solution of 3-methylbutane-1,3-diol (2.13 mL, 20 mmol) in pyridine (20

mL) at 0 °C was added para-toluenesulfonyl chloride (3.8 g, 20 mmol) in pyridine (10 mL) drop-wise over 15 minutes. This mixture stirred at ambient temperature for 3 hours and 35 mL H₂O was added and the mixture was extracted with ethyl acetate (2 x 50 mL). The combined organics were washed with H₂O (2 x 50 mL), dried over
5 MgSO₄, filtered, concentrated under reduced pressure and dried under vacuum (~1 mm Hg) to afford the title compound. MS (ESI) m/z 276 (M+18)⁺.

Example 308B

N-[(2Z)-5-tert-butyl-3-(3-hydroxy-3-methylbutyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

10 A solution of Example 244A (75 mg, 0.23 mmol) in N,N-dimethylformamide (2 mL) was treated with NaH 60% dispersion in oil (9.5 mg, 0.23 mmol) followed by Example 308A (60 mg, 0.23 mmol). The reaction mixture was stirred at room temperature for 18 hours, poured into water and extracted with ethyl acetate. The organic layer was washed with H₂O (2 x 50 mL), dried over MgSO₄, filtered,
15 concentrated under reduced pressure and dried under vacuum (~1 mm Hg) to afford the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-*d*₆) δ ppm 1.17 (s, 6 H), 1.32 (s, 9 H), 1.76 - 1.91 (m, 2 H), 3.79 (s, 3 H), 4.14 - 4.31 (m, 2 H), 4.43 (s, 1 H), 7.10 (d, *J*=8.8 Hz, 1 H), 7.31 (s, 1 H), 7.44 (dd, *J*=9.0, 2.9 Hz, 1 H), 7.76 (d, *J*=3.1 Hz, 1 H); MS (ESI) m/z 411 (M+H)⁺. Anal. Calculated for C₂₀H₂₇ClN₂O₃S, 58.45
20 H, 6.62 N, 6.82. Found C, 58.30 H, 6.51 N, 6.71.

Example 309

N-[(2Z)-5-tert-butyl-3-(3-hydroxy-3-methylbutyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 309A

25 (Z)-N-(5-tert-butyl-4-methylthiazol-2(3H)-ylidene)-5-chloro-2-methoxybenzamide
A mixture of 5-tert-butyl-4-methylthiazol-2(3H)-imine (250 mg, 1.5 mmol) in tetrahydrofuran (10 mL) was treated with triethylamine (0.25 mL, 1.8 mmol) and 5-chloro-2-methoxybenzoylchloride (307 mg, 1.5 mmol). The reaction mixture was stirred at 60 °C for 18 hours then concentrated under reduced pressure. The residue
30 was diluted with ethyl acetate and H₂O. The organic extract was dried over MgSO₄, filtered, concentrated under reduced pressure and dried under vacuum to afford the title compound (490 mg, 96% yield). MS (ESI) m/z 339 (M+H)⁺.

Example 309B

N-[(2Z)-5-tert-butyl-3-(3-hydroxy-3-methylbutyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 308A and Example 309A were processed using the method described in Example 308B to afford the title compound. ¹H NMR (300 MHz,

5 dimethylsulfoxide-*d*₆) δ ppm 1.19 (s, 6 H), 1.38 (s, 9 H), 1.65 - 1.81 (m, 2 H), 2.41 (s, 3 H), 3.78 (s, 3 H), 4.17 - 4.33 (m, 2 H), 4.49 (s, 1 H), 7.09 (d, *J*=8.8 Hz, 1 H), 7.43 (dd, *J*=8.8, 3.1 Hz, 1 H), 7.78 (d, *J*=2.7 Hz, 1 H). MS (ESI) *m/z* 425 (M+H)⁺. Anal. Calculated for C₂₁H₂₉ClN₂O₃S C, 59.35 H, 6.88 N, 6.55. Found C, 58.83 H, 7.13 N, 6.41.

10 Example 310

N-[(2Z)-5-tert-butyl-3-(2-methoxyethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Commercially available 2-bromoethyl methyl ether (Aldrich) and Example 244A were processed using the method described in Example 246 to afford the title
15 compound. ¹H NMR (300 MHz, dimethylsulfoxide-*d*₆) δ ppm 1.35 (s, 9 H), 3.36 (s, 3 H), 3.68 - 3.84 (m, 2 H), 3.90 (s, 3 H), 4.36 (t, *J*=5.1 Hz, 2 H), 6.77 (s, 1 H), 6.90 (d, *J*=8.8 Hz, 1 H), 7.32 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.95 (d, *J*=2.7 Hz, 1 H); MS (ESI⁺) *m/z* 383 (M+H)⁺; Anal. Calculated for C₁₈H₂₃ClN₂O₃S: C, 56.46; H, 6.05; N, 7.32. Found: C, 56.69; H, 6.02; N, 7.42.

20 Example 311

5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-3-(2-methoxyethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 311A

25 (Z)-N-(5-acetyl-3-(2-methoxyethyl)-4-methylthiazol-2(3H)-ylidene)-5-chloro-2-methoxybenzamide

Commercially available 2-bromoethyl methyl ether (Aldrich) and Example 238A were processed using the method described in Example 246 to afford the title compound. MS (ESI⁺) *m/z* 383 (M+H)⁺.

Example 311B

30 5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-3-(2-methoxyethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

Example 311A and commercially available methyllithium (Aldrich, 1.6 M in diethyl ether) were processed using the method described in Example 255 to afford

the title compound. ¹H NMR (300 MHz, dimethylsulfoxide-d₆) δ ppm 1.50 (s, 6 H), 2.42 (s, 3 H), 3.25 (s, 3 H), 3.68 (t, *J*=5.4 Hz, 2 H), 3.78 (s, 3 H), 4.30 (t, *J*=5.4 Hz, 2 H), 5.62 (s, 1 H), 7.09 (d, *J*=9.2 Hz, 1 H), 7.43 (dd, 1 H), 7.63 (d, *J*=2.7 Hz, 1 H); MS (ESI⁺) *m/z* 399 (M+H)⁺; Anal. Calculated for C₁₈H₂₃ClN₂O₄S: C, 54.20; H, 5.81; N, 7.02. Found: C, 54.30; H, 5.68; N, 6.91.

Example 312

N-[(2Z)-5-tert-butyl-3-(2-methoxy-2-methylpropyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 244A and 1-bromo-2-methoxy-2-methylpropane were processed using the method described in Example 246 to afford the title compound. ¹H NMR (500 MHz, CDCl₃) δ ppm 1.23 (s, 6 H) 1.36 (s, 9 H) 3.22 (s, 3 H) 3.89 (s, 3 H) 4.32 (s, 2 H) 6.89 (s, 1 H) 6.91 (s, 1 H) 7.33 (dd, *J*=8.9, 2.8 Hz, 1 H) 7.92 (d, *J*=2.8 Hz, 1 H); MS (ESI) *m/z* 411 (M+H)⁺.

Example 313

N-[(2Z)-3-butyl-5-(1-hydroxy-1-methylethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide

Example 313A

(Z)-N-(5-acetyl-3-butyl-4-methylthiazol-2(3H)-ylidene)-5-chloro-2-methoxybenzamide

A mixture of Example 238A (0.40 g, 1.2 mmol), 1-bromobutane (0.16 mL, 1.5 mmol), and potassium t-butoxide (0.22 g, 1.9 mmol) in 5 mL N,N-dimethylformamide was warmed to 65 °C and stirred for 20 hours. The mixture was cooled to ambient temperature quenched with 5 mL of saturated aqueous NH₄Cl and diluted with 5 mL of ethyl acetate. The layers were separated and the aqueous phase was extracted twice with 5 mL of ethyl acetate. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 40% hexanes in ethyl acetate) afforded the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.04 (t, *J*=7.3 Hz, 3 H), 1.42 - 1.55 (m, 2 H), 1.77 - 1.90 (m, 2 H), 2.51 (s, 3 H), 2.76 (s, 3 H), 3.88 (s, 3 H), 4.32 - 4.41 (m, 2 H), 7.10 (d, *J*=8.8 Hz, 1 H), 7.45 (dd, *J*=8.8, 2.7 Hz, 1 H), 7.99 (d, *J*=3.1 Hz, 1 H); MS (DCI/NH₃) *m/z* 381 (M+H)⁺. Anal. Calculated for C₁₈H₂₁ClN₂O₃S: C, 56.76; H, 5.56; N, 7.35. Found: C, 56.68; H, 5.49; N, 7.26.

Example 313B

N-[(2*Z*)-3-butyl-5-(1-hydroxy-1-methylethyl)-4-methyl-1,3-thiazol-2(3*H*)-ylidene]-5-chloro-2-methoxybenzamide

To the product of Example 313A (90 mg, 0.24 mmol) in 5 mL of THF at -78 °C was added a solution of methyllithium (1.6 M in diethyl ether, 0.44 mL, 0.71 mmol) dropwise over 5 minutes. The mixture was stirred at -78 °C for 4 hours and was slowly warmed to ambient temperature and allowed to stir for 12 hours. The mixture was quenched with 3 mL of saturated aqueous NH₄Cl and diluted with 5 mL of ethyl acetate. The layers were separated and the aqueous phase was extracted twice with 5 mL ethyl acetate. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 20% hexanes in ethyl acetate) afforded the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.02 (t, *J*=7.3 Hz, 3 H), 1.39 - 1.54 (m, 2 H), 1.61 (s, 6 H), 1.71 - 1.84 (m, 2 H), 2.51 (s, 3 H), 3.85 (s, 3 H), 4.23 - 4.31 (m, 2 H), 7.06 (d, *J*=8.8 Hz, 1 H), 7.39 (dd, *J*=9.0, 2.9 Hz, 1 H), 7.87 (d, *J*=3.1 Hz, 1 H); MS (DCI/NH₃) *m/z* 397 (M+H)⁺. Anal. Calculated for C₁₉H₂₅ClN₂O₃S: C, 57.49; H, 6.35; N, 7.06. Found: C, 57.36; H, 6.33; N, 6.85.

Example 314

(5-chloro-*N*-[(2*Z*)-3-(cyclobutylmethyl)-5-(1-hydroxy-1-methylethyl)-4-methyl-1,3-thiazol-2(3*H*)-ylidene]-2-methoxybenzamide

Example 314A

(*Z*)-*N*-(5-acetyl-3-(cyclobutylmethyl)-4-methylthiazol-2(3*H*)-ylidene)-5-chloro-2-methoxybenzamide

A mixture of the product of Example 238A (0.75 g, 2.3 mmol), (bromomethyl)cyclobutane (0.31 mL, 2.8 mmol), and potassium *t*-butoxide (0.41 g, 3.5 mmol) in 7 mL *N,N*-dimethylformamide was warmed to 65 °C and stirred for 16 hours. The mixture was cooled to ambient temperature, quenched with 5 mL of saturated aqueous NH₄Cl and diluted with 10 mL of ethyl acetate. The layers were separated and the aqueous phase was extracted twice with 7 mL of ethyl acetate. The combined organic extracts were dried over anhydrous Na₂SO₄, filtered and concentrated under reduced pressure. Purification via column chromatography (SiO₂, 40% hexanes in ethyl acetate) afforded the title compound. ¹H NMR (300 MHz, CD₃OD) δ ppm 1.88 - 2.15 (m, 6 H), 2.51 (s, 3 H), 2.75 (s, 3 H), 2.83 - 2.96 (m, 1 H), 3.89 (s, 3 H), 4.46 (d, *J*=7.5 Hz, 2 H), 7.11 (d, *J*=9.2 Hz, 1 H), 7.45 (dd, *J*=8.8, 2.7

Hz, 1 H), 7.99 (d, $J=3.1$ Hz, 1 H); MS (DCI/ NH_3) m/z 393 ($\text{M}+\text{H}$)⁺. Anal. Calculated for $\text{C}_{19}\text{H}_{21}\text{ClN}_2\text{O}_3\text{S}$: C, 58.08; H, 5.39; N, 7.13. Found: C, 58.06; H, 5.20; N, 7.06.

Example 314B

(5-chloro-N-[(2Z)-3-(cyclobutylmethyl)-5-(1-hydroxy-1-methylethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide

To the product of Example 314A (0.13 g, 0.33 mmol) in 5 mL of tetrahydrofuran at -78°C was added a solution of methyllithium (1.6 M in diethyl ether, 0.62 mL, 0.99 mmol) dropwise over 5 minutes. The mixture was stirred at -78°C for 1 hour then slowly warmed to ambient temperature and allowed to stir for 16 hours. The mixture was quenched with 5 mL of saturated aqueous NH_4Cl and diluted with 5 mL of ethyl acetate. The layers were separated and the aqueous phase was extracted twice with 5 mL ethyl acetate. The combined organic extracts were dried over anhydrous Na_2SO_4 , filtered and concentrated under reduced pressure. Purification via column chromatography (SiO_2 , 30% hexanes in ethyl acetate) afforded the title compound. ^1H NMR (300 MHz, CD_3OD) δ ppm 1.61 (s, 6 H), 1.86 - 2.13 (m, 6 H), 2.49 (s, 3 H), 2.79 - 2.93 (m, 1 H), 3.86 (s, 3 H), 4.37 (d, $J=7.1$ Hz, 2 H), 7.06 (d, $J=8.8$ Hz, 1 H), 7.39 (dd, $J=8.8, 2.7$ Hz, 1 H), 7.87 (d, $J=2.7$ Hz, 1 H); MS (DCI/ NH_3) m/z 409 ($\text{M}+\text{H}$)⁺. Anal. Calculated for $\text{C}_{20}\text{H}_{25}\text{ClN}_2\text{O}_3\text{S}$: C, 58.74; H, 6.16; N, 6.85. Found: C, 58.70; H, 6.12; N, 6.74.

In Vitro Methods

The CB_1 and CB_2 radioligand binding assays described herein are utilized to ascertain the selectivity of compounds of the present application for binding to CB_2 relative to CB_1 receptors.

Human CB_2 Radioligand Binding Assays:

HEK293 cells stably expressing human CB_2 receptors were grown until a confluent monolayer was formed. Briefly, the cells were harvested and homogenized in TE buffer (50 mM Tris-HCl, 1 mM MgCl_2 , and 1 mM EDTA) using a polytron for 2 X 10 second bursts in the presence of protease inhibitors, followed by centrifugation at 45,000Xg for 20 minutes. The final membrane pellet was re-homogenized in storage buffer (50 mM Tris-HCl, 1 mM MgCl_2 , and 1 mM EDTA and 10% sucrose) and frozen at -78°C until used. Saturation binding reactions were initiated by the addition of membrane preparation (protein concentration of 5 μg /well for human

CB₂) into wells of a deep well plate containing [³H]CP-55,940 (120 Ci/mmol, a nonselective CB agonist commercially available from Tocris) in assay buffer (50 mM Tris, 2.5 mM EDTA, 5 mM MgCl₂, and 0.5 mg/mL fatty acid free BSA, pH 7.4). After 90 min incubation at 30 °C, binding reaction was terminated by the addition of 300 µl/well of cold assay buffer followed by rapid vacuum filtration through a UniFilter-96 GF/C filter plates (pre-soaked in 1 mg/mL BSA for 2 hours). The bound activity was counted in a TopCount using Microscint-20. Saturation experiments were conducted with twelve concentrations of [³H]CP-55,940 ranging from 0.01 to 8 nM. Competition experiments were conducted with 0.5 nM [³H]CP-55,940 and five concentrations (1 nM to 10 µM) of displacing ligands. The addition of 10 µM unlabeled CP-55,940 (Tocris, Ellisville, MO) was used to assess nonspecific binding.

The majority of the compounds of the present application bound to CB₂ receptors with K_i of less than about 1,000 nM, preferably less than 400 nM, more preferably less than 200 nM, and most preferably lower than 100 nM.

15 Human CB₁ Radioligand Binding Assay:

HEK293 human CB₁ membranes were purchased from Perkin Elmer. Binding was initiated by the addition of membranes (8-12 µg per well) into wells (Scienceware 96-well DeepWell plate, VWR, West Chester, PA) containing [³H]CP-55,940 (120 Ci/mmol, Perkin Elmer, Boston, MA) and a sufficient volume of assay buffer (50 mM Tris, 2.5 mM EDTA, 5 mM MgCl₂, and 0.5 mg/mL fatty acid free BSA, pH 7.4) to bring the total volume to 250 µL. After incubation (30 °C for 90 minutes), binding was terminated by the addition of 300 µL per well of cold assay buffer and rapid vacuum filtration (FilterMate Cell Harvester, Perkin Elmer, Boston, MA) through a UniFilter-96 GF/C filter plate (Perkin Elmer, Boston, MA) (pre-soaked in 0.3% PEI at least 3 hours), followed by five washes with cold assay buffer. The bound activity was counted in the TopCount using Microscint-20 (both from Perkin Elmer, Boston, MA). Competition experiments were conducted with 1 nM [³H]CP-55,940 and five concentrations (1 nM to 10 µM) of displacing ligands. The addition of 10 µM unlabeled CP-55,940 (Tocris, Ellisville, MO) was used to assess nonspecific binding. The majority of the compounds of the present application tested for CB₁ binding, bound to CB₁ receptors with K_i 10x – 1000x higher than the K_i for CB₂. These results show that the compounds of the present application preferably bind to CB₂ receptors, therefore are selective ligands for the CB₂ receptor.

In Vivo Methods:Animals

Adult male Sprague-Dawley rats (250-300 g body weight, Charles River Laboratories, Portage, MI) were used. Animal handling and experimental protocols were approved by the Institutional Animal Care and Use Committee (IACUC) at Abbott Laboratories. For all surgical procedures, animals were maintained under halothane anesthesia (4% to induce, 2% to maintain), and the incision sites were sterilized using a 10% povidone-iodine solution prior to and after surgeries.

10 Incision Model of Postoperative Pain

A skin incision model of postoperative pain was produced using the procedures previously described (Brennan et al., 1996, Pain, 64, 493). All rats were anesthetized with isoflurane delivered via a nose cone. Right hind paw incision was performed following sterilization procedures. The plantar aspect of the left hind paw was placed through a hole in a sterile plastic drape. A 1-cm longitudinal incision was made through the skin and fascia of the plantar aspect of the hind paw, starting 0.5 cm from the proximal edge of the heel and extending towards the toes, the plantar muscle was elevated and incised longitudinally leaving the muscle origin and insertion points intact. The skin was then closed with two mattress sutures (5-0 nylon). After surgery, animals were then allowed to recover for 2 hours, at which time tactile allodynia was assessed as described below. To evaluate the anti-nociceptive effects, animals were i.p. administered vehicle or test compound 90 minutes following skin incision and tactile allodynia was assessed 30 minutes after compound administration.

Tactile allodynia was measured using calibrated von Frey filaments (Stoelting, Wood Dale, IL) as previously described (Chaplan, S.R., F.W. Bach, J.W. Pogrel, J.M. Chung and T.L. Yaksh, 1994, Quantitative assessment of tactile allodynia in the rat paw, J. Neurosci. Methods, 53, 55). Rats were placed into inverted individual plastic cage (20 x 12.5 x 20 cm) on top of a suspended wire mesh grid, and acclimated to the test chambers for 20 minutes. The von Frey filaments were applied perpendicularly from underneath the cage through openings in the wire mesh floor directly to an area within 1-3 mm (immediately adjacent) of the incision, and then held in this position for approximately 8 seconds with enough force to cause a slight bend in the filament. Positive responses included an abrupt withdrawal of the hind paw from the stimulus, or flinching behavior immediately following removal of the stimulus. A 50%

withdrawal threshold was determined using an up-down procedure (Dixon, W.J., 1980, Efficient analysis of experimental observations, Ann. Rev. Pharmacol. Toxicol. 20, 441).

Representative compounds of the present application showed a statistically significant change in paw withdrawal latency versus a saline vehicle at less than about 300 micromoles/kg in the incision model of postoperative pain. In a more preferred embodiment, compounds of the present application showed efficacy at less than about 50 micromoles/kg in the incision model of postoperative pain.

Complete Freund's Adjuvant (CFA) Model of Inflammatory Pain

Chronic inflammatory thermal hyperalgesia was induced by injection of 150 μ l of a 50% solution of CFA in phosphate buffered saline (PBS) into the plantar surface of the right hind paw in rats; control animals received only PBS treatment. Thermal hyperalgesia was assessed 48 hours post CFA injection. Thermal hyperalgesia was determined using a commercially available thermal paw stimulator (University Anesthesiology Research and Development Group (UARDG), University of California, San Diego, CA) described by Hargreaves et al. (Hargreaves, et. al., 1988, Pain 32, 77). Rats were placed into individual plastic cubicles mounted on a glass surface maintained at 30 °C, and allowed a 20 min habituation period. A thermal stimulus, in the form of radiant heat emitted from a focused projection bulb, was then applied to the plantar surface of each hind paw. The stimulus current was maintained at 4.50 ± 0.05 amp, and the maximum time of exposure was set at 20.48 sec to limit possible tissue damage. The elapsed time until a brisk withdrawal of the hind paw from the thermal stimulus was recorded automatically using photodiode motion sensors. The right and left hind paw of each rat was tested in three sequential trials at approximately 5-minute intervals. Paw withdrawal latency (PWL) was calculated as the mean of the two shortest latencies.

Representative compounds of the present invention showed a statistically significant change in paw withdrawal latency versus a saline vehicle at less than about 300 micromoles/kg in the Complete Freund's Adjuvant (CFA) model of inflammatory pain. In a more preferred embodiment, compounds of the present invention showed efficacy at less than about 50 micromoles/kg in the Complete Freund's Adjuvant (CFA) model of inflammatory pain.

Spinal Nerve Ligation Model of Neuropathic Pain

A model of spinal nerve ligation-induced (SNL model) neuropathic pain was produced using the procedure originally described by Kim and Chung (Kim, S.H. and J.M. Chung, 1992, An experimental model for peripheral neuropathy produced by segmental spinal nerve ligation in the rat, *Pain* 50, 355). The left L5 and L6 spinal nerves of the rat were isolated adjacent to the vertebral column and tightly ligated with a 5-0 silk suture distal to the DRG, and care was taken to avoid injury of the L4 spinal nerve. Sham rats underwent the same procedure, but without nerve ligation. All animals were allowed to recover for at least one week and not more than three weeks prior to assessment of tactile allodynia.

Tactile allodynia was measured using calibrated von Frey filaments (Stoelting, Wood Dale, IL) as previously described (Chaplan, S.R., F.W. Bach, J.W. Pogrel, J.M. Chung and T.L. Yaksh, 1994, Quantitative assessment of tactile allodynia in the rat paw, *J. Neurosci. Methods* 53, 55). Rats were placed into inverted individual plastic containers (20 x 12.5 x 20 cm) on top of a suspended wire mesh grid, and acclimated to the test chambers for 20 minutes. The von Frey filaments were presented perpendicularly to the plantar surface of the selected hind paw, and then held in this position for approximately 8 sec with enough force to cause a slight bend in the filament. Positive responses included an abrupt withdrawal of the hind paw from the stimulus, or flinching behavior immediately following removal of the stimulus. A 50% withdrawal threshold was determined using an up-down procedure (Dixon, W.J., 1980, Efficient analysis of experimental observations, *Ann. Rev. Pharmacol. Toxicol.* 20, 441). Only rats with a baseline threshold score of less than 4.25 g were used in this study, and animals demonstrating motor deficit were excluded. Tactile allodynia thresholds were also assessed in several control groups, including naive, sham-operated, and saline infused animals as well as in the contralateral paws of nerve-injured rats.

Representative compounds of the present invention showed efficacy at less than about 300 micromoles/kg in the spinal nerve ligation model of neuropathic pain. In a more preferred embodiment, compounds of the present invention showed efficacy at less than about 100 micromoles/kg in the spinal nerve ligation model of neuropathic pain.

The data contained herein demonstrates that compounds of the present invention bind to the CB₂ receptor. Certain compounds of the present invention were shown to have an analgesic effect in two types of animal pain models relating to

neuropathic and nociceptive pain.

In addition to the data contained herein, several lines of evidence support the assertion that CB₂ receptors play a role in analgesia. For example, Zimmer et al. have reported that the nonselective cannabinoid agonist Δ^9 -THC retains some analgesic efficacy in CB₁ receptor knockout mice (Zimmer, A., et al., Proc. Nat. Acad. Sci., 1999, 96, 5780-5785). HU-308 is one of the first highly selective CB₂ agonists identified that elicits an antinociceptive response in the rat formalin model of persistent pain (Hanus, L., et al., Proc. Nat. Acad. Sci., 1999, 96, 14228-14233). The CB₂-selective cannabinoid ligand AM-1241 exhibits robust analgesic efficacy in animal models of acute thermal pain (Malan, T. P., et al., Pain, 2001, 93, 239-245; Ibrahim, M. M., et al., Proc. Nat. Acad. Sci., 2005, 102(8), 3093-3098), persistent pain (Hohmann, A. G., et al., J. Pharmacol. Exp. Ther., 2004, 308, 446-453), inflammatory pain (Nackley, A. G., et al., Neuroscience, 2003, 119, 747-757; Quartilho, A. et al., Anesthesiology, 2003, 99, 955-60), and neuropathic pain (Ibrahim, M. M., et al., Proc. Nat. Acad. Sci., 2003, 100, 10529-10533). The CB₂-selective partial agonist GW405833, also known as L768242, is efficacious in rodent models of neuropathic, incisional, and both chronic and acute inflammatory pain (Valenzano, K. J., et al., Neuropharmacology, 2005, 48, 658-672 and Clayton, N., et al., Pain, 2002, 96, 253-260). The analgesic effects induced by these CB₂-selective ligands are blocked by CB₂ and not by CB₁ receptor antagonists. Furthermore, at fully efficacious doses, AM-1241 and GW405833 are devoid of typical CB₁ receptor-mediated CNS side effects, providing evidence that modulation of CB₂ receptors can produce broad-spectrum pain relief with reduced side-effect liability.

The potential exists for CB₂ modulators to have opioid sparing effects. A synergy between the analgesic effects of morphine and the nonselective CB agonist Δ^9 -THC has been documented (Cichewicz, D. L., Life Sci. 2004, 74, 1317-1324). Therefore, CB₂ ligands have additive or synergistic analgesic effects when used in combination with lower doses of morphine or other opioids, providing a strategy for reducing adverse opioid events, such as tolerance, constipation, and respiratory depression, without sacrificing analgesic efficacy.

CB₂ receptors are present in tissues and cell types associated with immune functions and CB₂ receptor mRNA is expressed by human B cells, natural killer cells, monocytes, neutrophils, and T cells (Galiegue et al., Eur. J. Biochem., 1995, 232, 54-

61). Studies with CB₂ knockout mice have suggested a role for CB₂ receptors in modulating the immune system (Buckley, N. E., et al., *Eur. J. Pharmacol.* 2000, 396, 141-149). Although immune cell development and differentiation are similar in knockout and wild type animals, the immunosuppressive effects of Δ^9 -THC are absent
5 in the CB₂ receptor knockout mice, providing evidence for the involvement of CB₂ receptors in immunomodulation. As such, selective CB₂ modulators are useful for the treatment of autoimmune diseases including but not limited to multiple sclerosis, rheumatoid arthritis, systemic lupus, myasthenia gravis, type I diabetes, irritable bowel syndrome, psoriasis, psoriatic arthritis, and hepatitis; and immune related
10 disorders including but not limited to tissue rejection in organ transplants, gluten-sensitive enteropathy (Celiac disease), asthma, chronic obstructive pulmonary disease, emphysema, bronchitis, acute respiratory distress syndrome, allergies, allergic rhinitis, dermatitis, and Sjogren's syndrome.

Microglial cells are considered to be the immune cells of the central nervous
15 system (CNS) where they regulate the initiation and progression of immune responses. They are quiescent and resting having a ramified morphology as long as the CNS is healthy. Microglia express a variety of receptors enabling them to survey the CNS and respond to pathological events. Insult or injury to the CNS leads to microglial cell activation, which is characterized by various morphological changes
20 allowing response to the lesion. Ramifications are retracted and microglia are transformed into amoeboid-like cells with phagocytic function. They can proliferate, rapidly migrate to the site of injury, and produce and release cytokines, chemokines and complement components (Watkins L. R., et al., *Trends in Neuroscience*, 2001, 24(8), 450; Kreutzberg, G. W., *Trends Neurosci.*, 1996, 19, 312-318). CB₂ receptor
25 expression on microglia is dependent upon inflammatory state with higher levels of CB₂ found in primed, proliferating, and migrating microglia relative to resting or fully activated microglial (Carlisle, S. J., et al. *Int. Immunopharmacol.*, 2002, 2, 69). Neuroinflammation induces many changes in microglia cell morphology and there is an upregulation of CB₂ receptors and other components of the endocannabinoid
30 system. It is conceivable that CB₂ receptors may be more susceptible to pharmacological effects during neuroinflammation (Walter, L., Stella, N., *Br. J. Pharmacol.* 2004, 141, 775-785). Neuroinflammation occurs in several neurodegenerative diseases, and induction of microglial CB₂ receptors has been

observed (Carrier, E. J., et al., *Current Drug Targets – CNS & Neurological Disorders*, 2005, 4, 657-665). Thus, CB₂ ligands may be clinically useful for the treatment of neuroinflammation.

CB₂ receptor expression has been detected in perivascular microglial cells
5 within normal, healthy human cerebellum (Nunez, E., et al., *Synapse*, 2004, 58, 208-213). Perivascular cells are immunoregulatory cells located adjacent to CNS blood vessels and, along with parenchymal microglia and astrocytes, they play a pivotal role in maintaining CNS homeostasis and blood-brain barrier functionality (Williams, K., et al., *Glia*, 2001, 36, 156-164). CB₂ receptor expression has also been detected on
10 cerebromicrovascular endothelial cells, which represent a main component of the blood-brain barrier (Golech, S. A., et al., *Mol. Brain Res.*, 2004, 132, 87-92). A recent report demonstrated that CB₂ receptor expression is up-regulated in the brains of macaques with simian immunodeficiency virus-induced encephalitis (Benito, C., et al., *J. Neurosci.* 2005, 25(10), 2530-2536). Thus, compounds that affect CB₂
15 signaling may protect the blood-brain barrier and be clinically useful in the treatment of neuroinflammation and a variety of neuroinflammatory disorders including retroviral encephalitis, which occurs with human immunodeficiency virus (HIV) infection in the CNS.

Multiple sclerosis is common immune-mediated disease of the CNS in which
20 the ability of neurons to conduct impulses becomes impaired through demyelination and axonal damage. The demyelination occurs as a consequence of chronic inflammation and ultimately leads to a broad range of clinical symptoms that fluctuate unpredictably and generally worsen with age. These include painful muscle spasms, tremor, ataxia, motor weakness, sphincter dysfunction, and difficulty speaking
25 (Pertwee, R. G., *Pharmacol. Ther.* 2002, 95, 165-174). The CB₂ receptor is up-regulated on activated microglial cells during experimental autoimmune encephalomyelitis (EAE) (Maresz, K., et al., *J. Neurochem.* 2005, 95, 437-445). CB₂ receptor activation prevents the recruitment of inflammatory cells such as leukocytes into the CNS (Ni, X., et al., *Multiple Sclerosis*, 2004, 10, 158-164) and plays a
30 protective role in experimental, progressive demyelination (Arevalo-Martin, A.; et al., *J. Neurosci.*, 2003, 23(7), 2511-2516), which are critical features in the development of multiple sclerosis. Thus, CB₂ receptor modulators provide a unique treatment for demyelinating pathologies.

Alzheimer's disease is a chronic neurodegenerative disorder accounting for

the most common form of elderly dementia. Recent studies have revealed that CB₂ receptor expression is upregulated in neuritic plaque-associated microglia from brains of Alzheimer's disease patients (Benito, C., et al., J. Neurosci., 2003, 23(35), 11136-11141). *In vitro*, treatment with the CB₂ agonist JWH-133 abrogated β -amyloid-induced microglial activation and neurotoxicity, effects that can be blocked by the CB₂ antagonist SR144528 (Ramirez, B. G., et al., J. Neurosci. 2005, 25(8), 1904-1913). CB₂ modulators possess both anti-inflammatory and neuroprotective actions and thus have clinical utility in treating neuroinflammation and in providing neuroprotection associated with the development of Alzheimer's disease.

Increased levels of epithelial CB₂ receptor expression are observed in human inflammatory bowel disease tissue (Wright, K., et al., Gastroenterology, 2005, 129, 437-453). Activation of CB₂ receptors re-established normal gastrointestinal transit after endotoxic inflammation was induced in rats (Mathison, R., et al., Br. J. Pharmacol. 2004, 142, 1247-1254). CB₂ receptor activation in a human colonic epithelial cell line inhibited TNF- α -induced interleukin-8 (IL-8) release (Ihenetu, K. et al., Eur. J. Pharmacol. 2003, 458, 207-215). Chemokines released from the epithelium, such as the neutrophil chemoattractant IL-8, are upregulated in inflammatory bowel disease (Warhurst, A. C., et al., Gut, 1998, 42, 208-213). Thus, administration of CB₂ receptor modulators represents a novel approach for the treatment of inflammation and disorders of the gastrointestinal tract including but not limited to inflammatory bowel disease, irritable bowel syndrome, secretory diarrhea, ulcerative colitis, Crohn's disease and gastroesophageal reflux disease (GERD).

Hepatic fibrosis occurs as a response to chronic liver injury and ultimately leads to cirrhosis, which is a major worldwide health issue due to the severe accompanying complications of portal hypertension, liver failure, and hepatocellular carcinoma (Lotersztajn, S., et al., Annu. Rev. Pharmacol. Toxicol., 2005, 45, 605-628). Although CB₂ receptors were not detectable in normal human liver, CB₂ receptors were expressed liver biopsy specimens from patients with cirrhosis. Activation of CB₂ receptors in cultured hepatic myofibroblasts produced potent antifibrogenic effects (Julien, B., et al., Gastroenterology, 2005, 128, 742-755). In addition, CB₂ knockout mice developed enhanced liver fibrosis after chronic administration of carbon tetrachloride relative to wild-type mice. Administration of CB₂ receptor modulators represents a unique approach for the treatment of liver

fibrosis.

CB₂ receptors are involved in the neuroprotective and anti-inflammatory mechanisms induced by the interleukin-1 receptor antagonist (IL-1ra) (Molina-Holgado, F., et al., *J. Neurosci.*, 2003, 23(16), 6470-6474). IL-1ra is an important anti-inflammatory cytokine that protects against ischemic, excitotoxic, and traumatic brain insults. CB₂ receptors play a role in mediating these neuroprotective effects indicating that CB₂ ligands are useful in the treatment of traumatic brain injury, stroke, and in mitigating brain damage.

Cough is a dominant and persistent symptom of many inflammatory lung diseases, including asthma, chronic obstructive pulmonary disease, viral infections, and pulmonary fibrosis (Patel, H. J., et al., *Brit. J. Pharmacol.*, 2003, 140, 261-268). Recent studies have provided evidence for the existence of neuronal CB₂ receptors in the airways, and have demonstrated a role for CB₂ receptor activation in cough suppression (Patel, H. J., et al., *Brit. J. Pharmacol.*, 2003, 140, 261-268 and Yoshihara, S., et al., *Am. J. Respir. Crit. Care Med.*, 2004, 170, 941-946). Both exogenous and endogenous cannabinoid ligands inhibit the activation of C-fibers via CB₂ receptors and reduce neurogenic inflammatory reactions in airway tissues (Yoshihara, S., et al., *J. Pharmacol. Sci.* 2005, 98(1), 77-82; Yoshihara, S., et al., *Allergy and Immunology*, 2005, 138, 80-87). Thus, CB₂-selective modulators have utility as antitussive agents for the treatment pulmonary inflammation, chronic cough, and a variety of airway inflammatory diseases including but not limited to asthma, chronic obstructive pulmonary disease, and pulmonary fibrosis.

Osteoporosis is a disease characterized by reduced bone mass, which leads to deterioration of bone microstructure and increased susceptibility to fracture. Age is associated with bone loss and it is estimated that 50% of all Caucasian women will have osteoporosis by the age of 80 (Ralston, S. H., *Curr. Opin. Pharmacol.*, 2003, 3, 286-290). There is a substantial genetic contribution to bone mass density and the CB₂ receptor gene is associated with human osteoporosis (Karsak, M., et al., *Human Molecular Genetics*, 2005, 14(22), 3389-3396). Osteoclasts and osteoblasts are largely responsible for maintaining bone structure and function through a process called remodeling, which involves resorption and synthesis of bone (Boyle, W. J., et al., *Nature*, 2003, 423, 337-342). CB₂ receptor expression has been detected on osteoclasts and osteoblastic precursor cells, and administration of a CB₂ agonist in mice caused a dose-dependent increase in bone formation (Grotenhermen, F. and

Muller-Vahl, K., Expert Opin. Pharmacother., 2003, 4(12), 2367-2371). Cannabinoid inverse agonists, including the CB₂-selective inverse agonist SR144528, have been shown to inhibit osteoclast activity and reverse ovariectomy-induced bone loss in mice, which is a model for post-menopausal osteoporosis (Ralston, S. H., et al.,
5 Nature Medicine, 2005, 11, 774-779). Thus, CB₂ modulators are useful for the treatment and prevention of osteoporosis, osteoarthritis, and bone disorders.

Artherosclerosis is a chronic inflammatory disease and is a leading cause of heart disease and stroke. CB₂ receptors have been detected in both human and mouse atherosclerotic plaques. Administration of low doses of THC in apolipoprotein E
10 knockout mice slowed the progression of atherosclerotic lesions, and these effects were inhibited by the CB₂-selective antagonist SR144528 (Steffens, S., et al., Nature, 2005, 434, 782-786). Thus, compounds with activity at the CB₂ receptor are clinically useful for the treatment of atherosclerosis.

CB₂ receptors are expressed on malignant cells of the immune system and
15 targeting CB₂ receptors to induce apoptosis may constitute a novel approach to treating malignancies of the immune system. Selective CB₂ agonists induce regression of malignant gliomas (Sanchez, C., et al., Cancer Res., 2001, 61, 5784-5789), skin carcinomas (Casanova, M. L., et al., J. Clin. Invest., 2003, 111, 43-50), and lymphomas (McKallip, R. J., et al., Blood, 2002, 15(2), 637-634). Thus, CB₂
20 modulators have utility as anticancer agents against tumors of immune origin.

Activation of CB₂ receptors has been demonstrated to protect the heart against the deleterious effects of ischemia and reperfusion (Lepicier, P., et al., Brit. J. Pharm. 2003, 139, 805-815; Bouchard, J.-F., et al., Life Sci. 2003, 72, 1859-1870; Filippo, C. D., et al., J. Leukoc. Biol. 2004, 75, 453-459). Thus, CB₂ modulators have utility for
25 the treatment or prophylaxis of cardiovascular disease and the development of myocardial infarction.

The present invention also provides pharmaceutical compositions that comprise compounds of the present invention. The pharmaceutical compositions comprise compounds of the present invention formulated together with one or more
30 non-toxic pharmaceutically acceptable carriers.

The pharmaceutical compositions of this invention can be administered to humans and other mammals orally, rectally, parenterally, intracisternally, intravaginally, topically (as by powders, ointments or drops), buccally or as an oral or nasal spray. The term "parenterally," as used herein, refers to modes of

administration that include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intraarticular injection and infusion.

The term "pharmaceutically acceptable carrier," as used herein, means a non-toxic, inert solid, semi-solid or liquid filler, diluent, encapsulating material or formulation auxiliary of any type. Some examples of materials which can serve as pharmaceutically acceptable carriers are sugars such as, but not limited to, lactose, glucose and sucrose; starches such as, but not limited to, corn starch and potato starch; cellulose and its derivatives such as, but not limited to, sodium carboxymethyl cellulose, ethyl cellulose and cellulose acetate; powdered tragacanth; malt; gelatin; talc; excipients such as, but not limited to, cocoa butter and suppository waxes; oils such as, but not limited to, peanut oil, cottonseed oil, safflower oil, sesame oil, olive oil, corn oil and soybean oil; glycols; such as propylene glycol; esters such as, but not limited to, ethyl oleate and ethyl laurate; agar; buffering agents such as, but not limited to, magnesium hydroxide and aluminum hydroxide; alginic acid; pyrogen-free water; isotonic saline; Ringer's solution; ethyl alcohol, and phosphate buffer solutions, as well as other non-toxic compatible lubricants such as, but not limited to, sodium lauryl sulfate and magnesium stearate, as well as coloring agents, releasing agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants can also be present in the composition, according to the judgment of the formulator.

Pharmaceutical compositions of this invention for parenteral injection comprise pharmaceutically acceptable sterile aqueous or nonaqueous solutions, dispersions, suspensions or emulsions as well as sterile powders for reconstitution into sterile injectable solutions or dispersions just prior to use. Examples of suitable aqueous and nonaqueous carriers, diluents, solvents or vehicles include water, ethanol, polyols (such as glycerol, propylene glycol, polyethylene glycol and the like), vegetable oils (such as olive oil), injectable organic esters (such as ethyl oleate) and suitable mixtures thereof. Proper fluidity can be maintained, for example, by the use of coating materials such as lecithin, by the maintenance of the required particle size in the case of dispersions and by the use of surfactants.

These compositions may also contain adjuvants such as preservatives, wetting agents, emulsifying agents and dispersing agents. Prevention of the action of microorganisms can be ensured by the inclusion of various antibacterial and antifungal agents, for example, paraben, chlorobutanol, phenol sorbic acid and the

like. It may also be desirable to include isotonic agents such as sugars, sodium chloride and the like. Prolonged absorption of the injectable pharmaceutical form can be brought about by the inclusion of agents, which delay absorption such as aluminum monostearate and gelatin.

- 5 In some cases, in order to prolong the effect of the drug, it is desirable to slow the absorption of the drug from subcutaneous or intramuscular injection. This can be accomplished by the use of a liquid suspension of crystalline or amorphous material with poor water solubility. The rate of absorption of the drug then depends upon its rate of dissolution that, in turn, may depend upon crystal size and crystalline form.
- 10 Alternatively, delayed absorption of a parenterally administered drug form is accomplished by dissolving or suspending the drug in an oil vehicle.

 Injectable depot forms are made by forming microencapsule matrices of the drug in biodegradable polymers such as polylactide-polyglycolide. Depending upon the ratio of drug to polymer and the nature of the particular polymer employed, the

15 rate of drug release can be controlled. Examples of other biodegradable polymers include poly(orthoesters) and poly(anhydrides). Depot injectable formulations are also prepared by entrapping the drug in liposomes or microemulsions, which are compatible with body tissues.

 The injectable formulations can be sterilized, for example, by filtration

20 through a bacterial-retaining filter or by incorporating sterilizing agents in the form of sterile solid compositions which can be dissolved or dispersed in sterile water or other sterile injectable medium just prior to use.

 Solid dosage forms for oral administration include capsules, tablets, pills, powders and granules. In such solid dosage forms, the active compound may be

25 mixed with at least one inert, pharmaceutically acceptable carrier or excipient, such as sodium citrate or dicalcium phosphate and/or a) fillers or extenders such as starches, lactose, sucrose, glucose, mannitol and silicic acid; b) binders such as carboxymethylcellulose, alginates, gelatin, polyvinylpyrrolidone, sucrose and acacia; c) humectants such as glycerol; d) disintegrating agents such as agar-agar, calcium

30 carbonate, potato or tapioca starch, alginic acid, certain silicates and sodium carbonate; e) solution retarding agents such as paraffin; f) absorption accelerators such as quaternary ammonium compounds; g) wetting agents such as cetyl alcohol and glycerol monostearate; h) absorbents such as kaolin and bentonite clay and i) lubricants such as talc, calcium stearate, magnesium stearate, solid polyethylene

glycols, sodium lauryl sulfate and mixtures thereof. In the case of capsules, tablets and pills, the dosage form may also comprise buffering agents.

Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such carriers as lactose or milk sugar as well as
5 high molecular weight polyethylene glycols and the like.

The solid dosage forms of tablets, dragees, capsules, pills and granules can be prepared with coatings and shells such as enteric coatings and other coatings well-known in the pharmaceutical formulating art. They may optionally contain opacifying agents and may also be of a composition such that they release the active
10 ingredient(s) only, or preferentially, in a certain part of the intestinal tract, optionally, in a delayed manner. Examples of embedding compositions that can be used include polymeric substances and waxes.

The active compounds can also be in micro-encapsulated form, if appropriate, with one or more of the above-mentioned carriers.

Liquid dosage forms for oral administration include pharmaceutically
15 acceptable emulsions, solutions, suspensions, syrups and elixirs. In addition to the active compounds, the liquid dosage forms may contain inert diluents commonly used in the art such as, for example, water or other solvents, solubilizing agents and emulsifiers such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate,
20 benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, dimethyl formamide, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofurfuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan and mixtures thereof.

Besides inert diluents, the oral compositions may also include adjuvants such
25 as wetting agents, emulsifying and suspending agents, sweetening, flavoring and perfuming agents.

Suspensions, in addition to the active compounds, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-
30 agar, tragacanth and mixtures thereof.

Compositions for rectal or vaginal administration are preferably suppositories which can be prepared by mixing the compounds of this invention with suitable non-irritating carriers or carriers such as cocoa butter, polyethylene glycol or a suppository wax which are solid at room temperature but liquid at body temperature and therefore

melt in the rectum or vaginal cavity and release the active compound.

Compounds of the present invention can also be administered in the form of liposomes. As is known in the art, liposomes are generally derived from phospholipids or other lipid substances. Liposomes are formed by mono- or multi-
5 lamellar hydrated liquid crystals that are dispersed in an aqueous medium. Any non-toxic, physiologically acceptable and metabolizable lipid capable of forming liposomes can be used. The present compositions in liposome form can contain, in addition to a compound of the present invention, stabilizers, preservatives, excipients and the like. The preferred lipids are natural and synthetic phospholipids and
10 phosphatidyl cholines (lecithins) used separately or together.

Methods to form liposomes are known in the art. See, for example, Prescott, Ed., *Methods in Cell Biology*, Volume XIV, Academic Press, New York, N.Y. (1976), p. 33 et seq.

Dosage forms for topical administration of a compound of this invention
15 include powders, sprays, ointments and inhalants. The active compound may be mixed under sterile conditions with a pharmaceutically acceptable carrier and any needed preservatives, buffers or propellants, which may be required. Ophthalmic formulations, eye ointments, powders and solutions are also contemplated as being within the scope of this invention.

20 Actual dosage levels of active ingredients in the pharmaceutical compositions of this invention can be varied so as to obtain an amount of the active compound(s) that is effective to achieve the desired therapeutic response for a particular patient, compositions and mode of administration. The selected dosage level will depend upon the activity of the particular compound, the route of administration, the severity
25 of the condition being treated and the condition and prior medical history of the patient being treated.

When used in the above or other treatments, a therapeutically effective amount of one of the compounds of the present invention can be employed in pure form or, where such forms exist, in pharmaceutically acceptable salt, ester or prodrug form.
30 The phrase "therapeutically effective amount" of the compound of the invention means a sufficient amount of the compound to treat disorders, at a reasonable benefit/risk ratio applicable to any medical treatment. It will be understood, however, that the total daily usage of the compounds and compositions of the present invention will be decided by the attending physician within the scope of sound medical

judgement. The specific therapeutically effective dose level for any particular patient will depend upon a variety of factors including the disorder being treated and the severity of the disorder; activity of the specific compound employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and rate of excretion of the specific compound employed; the duration of the treatment; drugs used in combination or coincidental with the specific compound employed; and like factors well known in the medical arts.

The term "pharmaceutically acceptable salt," as used herein, means acid addition salts or basic addition salts. The salts can be prepared in situ during the final isolation and purification of the compounds of the present application or separately by reacting the free base of the compounds of the present invention with an inorganic or organic acid. Representative acid addition salts include, but are not limited to, acetate, adipate, alginate, citrate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, camphorate, camphorsulfonate, digluconate, glycerophosphate, hemisulfate, heptanoate, hexanoate, fumarate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate (isethionate), lactate, malate, maleate, fumarate, methanesulfonate, nicotinate, 2-naphthalenesulfonate, oxalate, pamoate, pectinate, persulfate, 3-phenylpropionate, picrate, pivalate, propionate, succinate, sulfate, (L) tartrate, (D) tartrate, (DL) tartrate, thiocyanate, phosphate, glutamate, bicarbonate, p-toluenesulfonate, and undecanoate.

Basic addition salts can be prepared during the final isolation and purification of the present compounds by reaction of a carboxyl group with a suitable base such as the hydroxide, carbonate, or bicarbonate of a metal cation such as lithium, sodium, potassium, calcium, magnesium, or aluminum, or an organic primary, secondary, or tertiary amine. Quaternary amine salts derived from methylamine, dimethylamine, trimethylamine, triethylamine, diethylamine, ethylamine, tributylamine, pyridine, N,N-dimethylaniline, N-methylpiperidine, N-methylmorpholine, dicyclohexylamine, procaine, dibenzylamine, N,N-dibenzylphenethylamine, 1-phenamine, and N,N'-dibenzylethylenediamine, ethylenediamine, ethanolamine, diethanolamine, piperidine, piperazine, and the like, are contemplated as being within the scope of the present invention.

The term "pharmaceutically acceptable prodrug" or "prodrug," as used herein, represents those prodrugs of the compounds of the present invention which are, within

the scope of sound medical judgement, suitable for use in contact with the tissues of humans and lower animals without undue toxicity, irritation, allergic response, and the like. Prodrugs of the present invention may be rapidly transformed in vivo to compounds of the invention, for example, by hydrolysis in blood.


5 The present invention contemplates compounds of the invention formed by synthetic means or formed by in vivo biotransformation.

 The compounds of the invention can exist in unsolvated as well as solvated forms, including hydrated forms, such as hemi-hydrates. In general, the solvated forms, with pharmaceutically acceptable solvents such as water and ethanol among
10 others, are equivalent to the unsolvated forms for the purposes of the invention.

 The total daily dose of the compounds of this invention administered to a human or lower animal may range from about 0.003 to about 30 mg/kg/day. For purposes of oral administration, more preferable doses can be in the range of from about 0.01 to about 10 mg/kg/day. If desired, the effective daily dose can be divided
15 into multiple doses for purposes of administration; consequently, single dose compositions may contain such amounts or submultiples thereof to make up the daily dose.

What is claimed is:

1. A compound according to formula (I),



(I),

or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein

R₁ is alkoxyalkyl, alkoxyalkoxyalkyl, hydroxyalkyl, A, or A-alkylene-;

10 R₂ is hydrogen, alkoxy, alkoxyalkyl, alkoxycarbonyl, alkoxycarbonylalkyl, alkyl, alkyl-S(O)₂-, aryl, arylalkyl, arylalkenyl, azidoalkyl, cyano, cycloalkyl, halo, haloalkyl, heteroaryl, heterocycle, -(CR₂₁R₂₂)_m-OH, R_aR_bN-, R_aR_bN-alkyl-, R_cR_dNC(O)-, or R₈-R₇-;

15 R₃ is hydrogen, alkoxy, alkoxyalkyl, alkyl, alkylcarbonyl, alkyl-S(O)₂-, aryl, arylalkyl, arylalkenyl, cyano, cycloalkyl, halo, haloalkyl, heteroaryl, heterocycle, -(CR₃₁R₃₂)_m-OH, R_aR_bN-, R_aR_bN-alkyl-, or R₈-R₇-; or

R₂ and R₃, together with the carbon atoms to which they are attached, form a 4-, 5-, 6-, or 7-membered monocyclic ring, optionally fused to a benzo or a monocyclic heteroaryl, said monocyclic ring contains zero, one, or two additional double bonds, zero or one oxygen atom, and zero or one nitrogen atom as ring atoms; two non-adjacent atoms of said monocyclic ring can be optionally linked by an alkenylene bridge of 2, 3, or 4 carbon atoms, or optionally linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms, said monocyclic ring is independently unsubstituted or substituted with 1, 2, 3, 4, or 5 substituents independently selected from the group consisting of oxo, alkyl, halo, -OH, -O(alkyl), and haloalkyl; two substituents on the same carbon atom of said monocyclic ring, together with the carbon atom to which they are attached, optionally form a 3-, 4-, 5-, or 6-membered monocyclic cycloalkyl ring, wherein the monocyclic cycloalkyl ring is optionally substituted with 1, 2, 3, 4, 5, or 6 substituents independently selected from the group consisting of alkyl and haloalkyl;

20

25

30

with the proviso that when R₂ and R₃ are other than forming a ring with the

carbon atoms to which they are attached, and R_1 is A or A-alkylene-, then R_4 is alkyl wherein the alkyl is optionally substituted with one substituent selected from the group consisting of alkoxy, alkoxycarbonyl, carboxy, halo, -OH, and R_eR_fN -, alkynyl, cycloalkyl, cycloalkenyl, aryl, cycloalkylalkyl, cycloalkenylalkyl, arylalkyl, or

5 $R_{10}-L_2-R_9$ - wherein R_9 is aryl, cycloalkyl, or cycloalkenyl;

R_4 is alkyl, alkynyl, cycloalkyl, cycloalkenyl, aryl, heteroaryl, heterocycle, cycloalkylalkyl, cycloalkenylalkyl, arylalkyl, heteroarylalkyl, heterocyclealkyl, or $R_{10}-L_2-R_9$ -; wherein the alkyl group is optionally substituted with one substituent selected from the group consisting of alkoxy, alkoxycarbonyl, carboxy, halo, -OH,

10 and R_eR_fN -;

R_7 , R_8 , R_9 , and R_{10} are each independently aryl, cycloalkyl, cycloalkenyl, heteroaryl, or heterocycle;

R_a and R_b , at each occurrence, are each independently hydrogen, alkoxycarbonyl, alkyl, alkylcarbonyl, alkyl-S(O)₂-, or arylalkyl;

15 R_c and R_d , are each independently hydrogen or alkyl;

R_e and R_f , are each independently hydrogen, alkyl, or alkylcarbonyl;

A is a 4-, 5-, 6-, 7-, 8-, or 9-membered monocyclic heterocycle containing zero or one double bond, and one or two oxygen, and zero or one nitrogen as ring atoms, optionally fused with a monocyclic ring selected from benzo, cycloalkyl, cycloalkenyl, heterocycle and heteroaryl; wherein two non-adjacent atoms of each A can be optionally linked by an alkenylene bridge of 2, 3, or 4 carbon atoms, or optionally linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms; each A is independently unsubstituted or substituted with 1, 2, 3, 4, 5, or 6 substituents independently selected from the group consisting of oxo, alkyl, halo, -OH, -O(alkyl),

25 and haloalkyl;

L_1 is a single bond or -NR_g-;

L_2 is a single bond, alkylene, or -O-;


R_g is hydrogen or alkyl;

R_{21} , R_{22} , R_{31} , and R_{32} , at each occurrence, are each independently hydrogen, alkyl, haloalkyl, or halo; and

30 m , at each occurrence, is independently 1, 2, 3, or 4.

2. The compound of claim 1, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein R_1 is A or A-alkylene-.


3. The compounds of claim 2, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein A is (i), (ii), (iii), (iv), (v), or (vi)



5

4. The compound of claim 1, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein R_2 and R_3 , together with the carbon atoms to which they are attached, form a 4-, 5-, 6-, or 7-membered monocyclic ring, optionally fused to a benzo or a monocyclic heteroaryl, said monocyclic ring contains zero or one additional double bond, and zero oxygen atom and zero nitrogen atom as ring atoms; two non-adjacent atoms of said monocyclic ring are linked by an alkenylene bridge of 2, 3, or 4 carbon atoms, or linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms, said monocyclic ring is independently unsubstituted or substituted with 1, 2, 3, 4, or 5 substituents independently selected from the group consisting of oxo, alkyl, halo, -OH, -O(alkyl), and haloalkyl; two substituents on the same carbon atom of said monocyclic ring, together with the carbon atom to which they are attached, optionally form a 3-, 4-, 5-, or 6-membered monocyclic cycloalkyl ring, wherein the monocyclic cycloalkyl ring is optionally substituted with 1, 2, 3, 4, 5, or 6 substituents independently selected from the group consisting of alkyl and haloalkyl.

5. The compound of claim 4, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein R_2 and R_3 , together with the carbon atoms to which they are attached, form a ring as represented by (ix), (xii), (xiii), or (xiv)





6. The compound of claim 1, or a pharmaceutically acceptable salt, prodrug, salt

of a prodrug, or a combination thereof, wherein R_2 and R_3 , together with the carbon atoms to which they are attached, form a 4-, 5-, 6-, or 7-membered monocyclic ring, optionally fused to a benzo or a monocyclic heteroaryl, said monocyclic ring contains zero or one additional double bond, and one oxygen atom and zero or one nitrogen atom as ring atoms; two non-adjacent atoms of said monocyclic ring can be optionally linked by an alkenylene bridge of 2, 3, or 4 carbon atoms, or optionally linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms, said monocyclic ring is independently unsubstituted or substituted with 1, 2, 3, 4, or 5 substituents independently selected from the group consisting of oxo, alkyl, halo, -OH, -O(alkyl), and haloalkyl; two substituents on the same carbon atom of said monocyclic ring, together with the carbon atom to which they are attached, optionally form a 3-, 4-, 5-, or 6-membered monocyclic cycloalkyl ring, wherein the monocyclic cycloalkyl ring is optionally substituted with 1, 2, 3, 4, 5, or 6 substituents independently selected from the group consisting of alkyl and haloalkyl.

15

7. The compound of claim 6, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein the monocyclic ring is formula (xv), (xvi), (xvii), (xviii), (xix), (xx), (xxi), (xxii), (xxiii), (xxiv), (xxv), (xxvi), (xxvii), (xxviii), or (xxix)



20

8. The compound of claim 1, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein

R_1 is A or A-alkenyl-,

25 R_2 is hydrogen, alkoxycarbonyl, alkyl, aryl, halo, haloalkyl, or $-(CR_{21}R_{22})_m-OH$;

R_3 is hydrogen, alkyl, aryl, cycloalkyl, halo, haloalkyl, heterocycle, or

-(CR₃₁R₃₂)_m-OH; and

R₄ is alkyl wherein the alkyl is optionally substituted with one substituent selected from the group consisting of alkoxy, alkoxycarbonyl, carboxy, halo, -OH, and R_cR_iN-, alkynyl, cycloalkyl, cycloalkenyl, aryl, cycloalkylalkyl, cycloalkenylalkyl, arylalkyl, or R₁₀-L₂-R₉, wherein R₉ is aryl, cycloalkyl, or cycloalkenyl.

9. The compound of claim 8, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein R₄ is aryl or R₁₀-L₂-R₉ wherein R₉ is aryl.

10. The compound of claim 8, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein

R₂ is hydrogen or alkyl; and
R₃ is alkyl or -(CR₃₁R₃₂)_m-OH.

11. The compound of claim 10, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein

R₄ is phenyl, optionally substituted with alkyl, alkoxy, cyano, halo, haloalkoxy, hydroxy, and haloalkyl;
R₃₁ and R₃₂ are each independently alkyl or haloalkyl; and
m is 1.

12. The compound of claim 1, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein

R₁ is A or A-alkenyl; and
R₂ and R₃, together with the carbon atoms to which they are attached form a 4-, 5-, 6-, or 7-membered monocyclic ring, optionally fused to a benzo or a monocyclic heteroaryl, said monocyclic ring contains zero, one, or two additional double bonds, zero or one oxygen atom, and zero or one nitrogen atom as ring atoms; two non-adjacent atoms of said monocyclic ring can be optionally linked by an alkenylene bridge of 2, 3, or 4 carbon atoms, or optionally linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms, said monocyclic ring is independently unsubstituted or substituted with 1, 2, 3, 4, or 5 substituents independently selected

from the group consisting of oxo, alkyl, halo, -OH, -O(alkyl), and haloalkyl; two substituents on the same carbon atom of said monocyclic ring, together with the carbon atom to which they are attached, optionally form a 3-, 4-, 5-, or 6-membered monocyclic cycloalkyl ring, wherein the monocyclic cycloalkyl ring is optionally substituted with 1, 2, 3, 4, 5, or 6 substituents independently selected from the group consisting of alkyl and haloalkyl.

13. The compound of claim 12, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein R_4 is aryl or R_{10} - L_2 - R_9 -, wherein R_9 is aryl.

14. The compound of claim 1, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein

R_1 is alkoxyalkyl, alkoxyalkoxyalkyl, or hydroxyalkyl, and
15 R_2 and R_3 , together with the carbon atoms to which they are attached, form a 4-, 5-, 6-, or 7-membered monocyclic ring, optionally fused to a benzo or a monocyclic heteroaryl, said monocyclic ring contains zero or one additional double bond, zero oxygen atom and zero nitrogen atom as ring atoms; two non-adjacent atoms of said monocyclic ring are linked by an alkenylene bridge of 2, 3, or 4 carbon
20 atoms, or linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms, said monocyclic ring is independently unsubstituted or substituted with 1, 2, 3, 4, or 5 substituents independently selected from the group consisting of oxo, alkyl, halo, -OH, -O(alkyl), and haloalkyl; two substituents on the same carbon atom of said monocyclic ring, together with the carbon atom to which they are attached, optionally form a 3-, 4-, 5-,
25 or 6-membered monocyclic cycloalkyl ring, wherein the monocyclic cycloalkyl ring is optionally substituted with 1, 2, 3, 4, 5, or 6 substituents independently selected from the group consisting of alkyl and haloalkyl.

15. The compound of claim 1, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein

R_1 is alkoxyalkyl, alkoxyalkoxyalkyl, or hydroxyalkyl, and
 R_2 and R_3 , together with the carbon atoms to which they are attached, form a 4-, 5-, 6-, or 7-membered monocyclic ring, optionally fused to a benzo or a monocyclic heteroaryl, said monocyclic ring contains zero or one additional double

bonds, one oxygen atom, and zero or one nitrogen atom as ring atoms; two non-adjacent atoms of said monocyclic ring can be optionally linked by an alkenylene bridge of 2, 3, or 4 carbon atoms, or optionally linked by an alkylene bridge of 1, 2, 3, or 4 carbon atoms, said monocyclic ring is independently unsubstituted or substituted with 1, 2, 3, 4, or 5 substituents independently selected from the group consisting of oxo, alkyl, halo, -OH, -O(alkyl), and haloalkyl; two substituents on the same carbon atom of said monocyclic ring, together with the carbon atom to which they are attached, optionally form a 3-, 4-, 5-, or 6-membered monocyclic cycloalkyl ring, wherein the monocyclic cycloalkyl ring is optionally substituted with 1, 2, 3, 4, 5, or 6 substituents independently selected from the group consisting of alkyl and haloalkyl.

16. The compound of claim 1, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein

L_1 is a single bond;
 R_1 is alkoxyalkyl, alkoxyalkoxyalkyl, or hydroxyalkyl,
 R_2 is hydrogen or alkyl,
 R_3 is $-(CR_{31}R_{32})_m-OH$;
 R_4 is phenyl;
 R_{31} and R_{32} are methyl; and
 m is 1.

17. The compound of claim 16 is 5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-3-(2-methoxyethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide, or a pharmaceutically acceptable salt thereof.

18. The compound of claim 1 selected from the group consisting of
 (2R)-N-[(2Z)-3-(2-methoxyethyl)-4,5-dimethyl-1,3-thiazol-2(3H)-ylidene]-2-propylhex-4-ynamide;
 N-[(7Z)-8-(2-methoxyethyl)-5,8-dihydro[1,3]thiazolo[4,5-e][2,1,3]benzoxadiazol-7(4H)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide;
 N-[(7Z)-8-(2-methoxyethyl)[1,3]thiazolo[4,5-e][2,1,3]benzoxadiazol-7(8H)-ylidene]-2,2,3,3-tetramethylcyclopropanecarboxamide;
 2-Ethoxy-N-[(2Z)-3-(2-methoxyethyl)-4,6-dihydrofuro[3,4-d]thiazol-2(3H)-ylidene]-benzamide;

- 2,3-dichloro-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 2,3-dichloro-N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 5 2-ethoxy-N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 10 2-ethoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 15 2-ethoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 2-ethoxy-N-[(2Z)-3-[2-(2-methoxyethoxy)ethyl]-5-methyl-1,3-thiazol-2(3H)-ylidene]benzamide;
- 2,3-dichloro-N-[(2Z)-3-[2-(2-methoxyethoxy)ethyl]-5-methyl-1,3-thiazol-2(3H)-ylidene]benzamide;
- 20 2(3H)-ylidene]benzamide;
- 5-chloro-2-methoxy-N-[(2Z)-3-[2-(2-methoxyethoxy)ethyl]-5-methyl-1,3-thiazol-2(3H)-ylidene]benzamide;
- 2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 25 1-(1,1-dimethylpropyl)-3-[(2Z)-5-methyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea;
- 1-(1,2-dimethyl-propyl)-3-[(2Z)-5-methyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea hydrochloride;
- 1-cyclohexyl-3-[(2Z)-4,5-dimethyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea;
- 30 1-(4-methylcyclohexyl)-3-[(2Z)-5-methyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea;
- 1-(1,1-dimethylpropyl)-3-[(2Z)-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea;

- N-[(2Z)-4,5-dimethyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-N'-[(1S)-1,2,2-trimethylpropyl]urea;
- 1-(2,2-dimethylpropyl)-3-[(2Z)-4,5-dimethyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea;
- 5 N-[(2Z)-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-N'-[(1S)-1,2,2-trimethylpropyl]urea;
- N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-N'-[(1S)-1,2,2-trimethylpropyl]urea;
- 1-(4-methylcyclohexyl)-3-[(2Z)-4,5-dimethyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea;
- 10 1-(2,2-dimethylpropyl)-3-[(2Z)-5-methyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea;
- 1-tert-butyl-3-[(2Z)-4,5-dimethyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea;
- 15 1-(1,1-dimethylpropyl)-3-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea;
- 1-(2,2-dimethylpropyl)-3-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea;
- 1-[(2Z)-4,5-dimethyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-3-(3,3,5,5-tetramethylcyclohexyl)urea;
- 20 N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-N'-[(1S)-1,2,2-trimethylpropyl]urea;
- N-[(2Z)-4,5-dimethyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-N'-[(1R)-1,2,2-trimethylpropyl]urea;
- 25 1-tert-butyl-3-[(2Z)-5-methyl-3-(tetrahydropyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea;
- 1-(2,3-dichlorophenyl)-3-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea;
- N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-N'-[(1R)-1,2,2-trimethylpropyl]urea;
- 30 N-[(1S)-1,2-dimethylpropyl]-N'-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea;
- 1-(1-cyclopropylethyl)-3-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]urea;

- 2-ethoxy-N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-4-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 2,4-dimethoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 5 5-chloro-2-methoxy-N-[(2Z)-4-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-4-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 10 5-chloro-2-methoxy-N-[(2Z)-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide;
- 5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-[(2S)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide;
- 15 2,2,3,3-tetrafluoro-1-methyl-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]cyclobutanecarboxamide;
- 5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(oxetan-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 5-chloro-N-[(2Z)-3-(1,3-dioxolan-2-ylmethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- 20 5-chloro-N-[(2Z)-3-[2-(1,3-dioxolan-2-yl)ethyl]-5-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- N-[(2Z)-3-(1,3-dioxolan-2-ylmethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]-2-ethoxybenzamide;
- 25 5-bromo-2-ethoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide;
- 5-chloro-2-ethoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide;
- 4-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 30 5-chloro-2-methoxy-N-[(2Z)-1-(2-methoxyethyl)-1,4,6,7-tetrahydro-2H-pyrano[4,3-d][1,3]thiazol-2-ylidene]benzamide;
- 5-bromo-2-methoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide;

- 5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(2-tetrahydro-2H-pyran-4-ylethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 5-chloro-N-[(2Z)-5-ethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- 5 5-chloro-2-methoxy-N-[(2Z)-5-propyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 5-chloro-N-[(2Z)-5-chloro-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- 4,5-dichloro-2-methoxy-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide;
- 10 ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide;
- 5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-(tetrahydro-2H-pyran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]benzamide;
- 2-chloro-N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]nicotinamide;
- 15 5-chloro-N-[(2Z)-4,5-dimethyl-3-[(3-methyl-4,5-dihydroisoxazol-5-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-4-(trifluoromethyl)nicotinamide;
- N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-ethoxynicotinamide;
- 20 ylidene]-2,3,6-trifluoroisonicotinamide;
- N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2,3,6-trifluoroisonicotinamide;
- 6-chloro-N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-4-(trifluoromethyl)nicotinamide;
- 25 5-chloro-2-methoxy-N-[(2Z)-5-methyl-3-[(3-methyl-4,5-dihydroisoxazol-5-yl)methyl]-1,3-thiazol-2(3H)-ylidene]benzamide;
- N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-(ethylamino)benzamide;
- N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-iodo-2-(methylamino)benzamide;
- 30 ylidene]-5-iodo-2-(methylamino)benzamide;
- 5-bromo-N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- 5-chloro-2-(cyclopropyloxy)-N-[(2Z)-5-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide;

- 5-chloro-N-[(2Z)-3-(1,4-dioxan-2-ylmethyl)-5-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- N-[(2Z)-5-acetyl-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;
- 5 5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;
- N-[(2Z)-5-tert-butyl-3-(1,3-dioxolan-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-10 5-chloro-2-methoxybenzamide;
- 5-chloro-N-[(2Z)-5-chloro-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- 5-chloro-N-[(2Z)-5-chloro-3-(tetrahydro-2H-pyran-4-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- 15 N-[(2Z)-5-tert-butyl-3-[(3-methyl-4,5-dihydroisoxazol-5-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;
- N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-2,2-dimethyl-4-oxo-3,4-dihydro-2H-pyran-6-carboxamide;
- N-[(2Z)-5-tert-butyl-3-(tetrahydro-2H-pyran-4-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;
- 20 5-chloro-2-methoxy-N-[(2Z)-3-(tetrahydro-2H-pyran-4-ylmethyl)-4,5,6,7-tetrahydro-1,3-benzothiazol-2(3H)-ylidene]benzamide;
- 5-chloro-2-methoxy-N-[(2Z)-3-(2-methoxyethyl)-4,4,6,6-tetramethyl-4,6-dihydrofuro[3,4-d][1,3]thiazol-2(3H)-ylidene]benzamide;
- 25 5-chloro-2-methoxy-N-[(2Z)-3-(2-methoxyethyl)-6,6-dimethyl-4-oxo-4,6-dihydrofuro[3,4-d][1,3]thiazol-2(3H)-ylidene]benzamide;
- N-[(2Z)-5-acetyl-4-methyl-3-(oxetan-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;
- 5-chloro-N-[(2Z)-4,4-dimethyl-1-(oxetan-2-ylmethyl)-1,4,6,7-tetrahydro-2H-pyrano[4,3-d][1,3]thiazol-2-ylidene]-2-methoxybenzamide;
- 30 5-chloro-N-[(2Z)-4,4-dimethyl-1-[(2R)-tetrahydrofuran-2-ylmethyl]-1,4,6,7-tetrahydro-2H-pyrano[4,3-d][1,3]thiazol-2-ylidene]-2-methoxybenzamide;
- N-[(2Z)-5-acetyl-4-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;

- N-[(2Z)-5-acetyl-4-methyl-3-(tetrahydro-2H-pyran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;
- 5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-4-methyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- 5-chloro-2-methoxy-N-[(2Z)-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-5-[2,2,2-trifluoro-1-hydroxy-1-(trifluoromethyl)ethyl]-1,3-thiazol-2(3H)-ylidene]benzamide;
- 5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-4-methyl-3-(tetrahydro-2H-pyran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- 5-chloro-N-[(2Z)-3-{[(4R)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}-5-(1-hydroxy-1-methylethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- 5-chloro-N-[(2Z)-6,6-dimethyl-4-oxo-3-[(2R)-tetrahydrofuran-2-ylmethyl]-4,6-dihydrofuro[3,4-d][1,3]thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- 5-chloro-N-[(2Z)-3-{[(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl}-5-(1-hydroxy-1-methylethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- N-[(2Z)-5-acetyl-3-(1,4-dioxan-2-ylmethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;
- 5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-4-methyl-3-(oxetan-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- 5-chloro-N-[(2Z)-5-(1-hydroxy-1-methylethyl)-4-methyl-3-(tetrahydrofuran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- 5-chloro-N-[(2Z)-3-(1,4-dioxan-2-ylmethyl)-5-(1-hydroxy-1-methylethyl)-4-methyl-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;
- N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-3-methoxy-2-naphthamide;
- N-[(2Z)-5-tert-butyl-3-[(3-methyloxetan-3-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;
- 5-chloro-2-methoxy-N-[(2Z)-3-(tetrahydrofuran-2-ylmethyl)-3,4,5,6-tetrahydro-2H-cyclopenta[d][1,3]thiazol-2-ylidene]benzamide;
- 5-chloro-2-methoxy-N-[(2Z)-3-(tetrahydro-2H-pyran-4-ylmethyl)-3,4,5,6-tetrahydro-2H-cyclopenta[d][1,3]thiazol-2-ylidene]benzamide;
- N-[(2Z)-4,5-dimethyl-3-(tetrahydro-2H-pyran-4-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2,2-dimethyltetrahydro-2H-pyran-4-carboxamide;
- N-[(2Z)-5-tert-butyl-3-(oxetan-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-

chloro-2-methoxybenzamide;

N-[(2Z)-5-tert-butyl-3-(tetrahydro-2H-pyran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;

N-[(2Z)-5-tert-butyl-3-(1,4-dioxan-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;

N-[(2Z)-5-tert-butyl-3-([(4S)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;

N-[(2Z)-5-tert-butyl-3-([(4R)-2,2-dimethyl-1,3-dioxolan-4-yl]methyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;

N-[(2Z)-5-tert-butyl-3-(tetrahydrofuran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;

N-[(2Z)-5-tert-butyl-3-(tetrahydro-2H-pyran-3-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;

N-[(2Z)-5-tert-butyl-3-([(2S)-5-oxotetrahydrofuran-2-yl]methyl)-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;

N-[(2Z)-5-acetyl-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-1-benzofuran-5-carboxamide;

N-[(2Z)-5-(1-hydroxy-1-methylethyl)-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-1-benzofuran-5-carboxamide;

N-[(2Z)-5-acetyl-4-methyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-(2,2,2-trifluoroethoxy)benzamide;

N-[(2Z)-5-tert-butyl-3-[(5-methyltetrahydrofuran-2-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;

N-[(2Z)-5-tert-butyl-3-[(5,5-dimethyltetrahydrofuran-2-yl)methyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-(2-methoxyethoxy)benzamide;

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-6-chloroquinoline-8-carboxamide;


5-chloro-2-methoxy-N-[(2Z)-5-(1-methylcyclopropyl)-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]benzamide;

5-chloro-N-[(2Z)-5-(1-hydroxy-3-iodo-1-methylpropyl)-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-2-methoxybenzamide;

N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-

- ylidene]-5-chloro-2-[(1-methylcyclopropyl)methoxy]benzamide;
 N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-isopropoxybenzamide;
 N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-ethoxybenzamide;
 N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-(tetrahydrofuran-3-yloxy)benzamide;
 N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-[(2-methoxyethyl)(methyl)amino]benzamide;
 N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-(difluoromethoxy)benzamide;
 N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-(trifluoromethoxy)benzamide;
 N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-(2,2,2-trifluoroethoxy)benzamide;
 N-[(2Z)-5-tert-butyl-3-[(2R)-tetrahydrofuran-2-ylmethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-[3-(dimethylamino)propoxy]benzamide;
 5,6-dichloro-N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]quinoline-8-carboxamide;
 6-chloro-N-[(2Z)-4,5-dimethyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]quinoline-8-carboxamide;
 6-chloro-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]quinoline-8-carboxamide;
 5,6-dichloro-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]quinoline-8-carboxamide;
 3-methoxy-N-[(2Z)-5-methyl-3-(tetrahydrofuran-2-ylmethyl)-1,3-thiazol-2(3H)-ylidene]-2-naphthamide; and
 N-[(2Z)-5-tert-butyl-3-[2-(2-methoxyethoxy)ethyl]-1,3-thiazol-2(3H)-ylidene]-5-chloro-2-methoxybenzamide;
 or a pharmaceutical acceptable salt thereof.

19. A compound of formula (II), or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof



(II)

wherein

R_{1a} is alkyl, haloalkyl, or cycloalkylalkyl;

5 R_{2a} is hydrogen or alkyl;

R_x is an optional substituent of phenyl, selected from the group consisting of alkyl, alkoxy, cyano, halo, haloalkoxy, hydroxyl, and haloalkyl; and n is 1, 2, 3, 4, or 5.

10 20. The compound of claim 19, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein R_{1a} is cyclobutylmethyl.

21. The compound of claim 19, or a pharmaceutically acceptable salt, prodrug, salt of a prodrug, or a combination thereof, wherein R_{1a} is alkyl.

15

22. The compound of claim 19 selected from the group consisting of 5-chloro-*N*-[(2*Z*)-5-(1-hydroxy-1-methylethyl)-3-(((*cis*)-3-methoxycyclobutyl)methyl)-4-methyl-1,3-thiazol-2(3*H*)-ylidene]-2-methoxybenzamide,

20 *N*-[(2*Z*)-3-butyl-5-(1-hydroxy-1-methylethyl)-4-methyl-1,3-thiazol-2(3*H*)-ylidene]-5-chloro-2-methoxybenzamide; or (5-chloro-*N*-[(2*Z*)-3-(cyclobutylmethyl)-5-(1-hydroxy-1-methylethyl)-4-methyl-1,3-thiazol-2(3*H*)-ylidene]-2-methoxybenzamide; or a pharmaceutically acceptable salt thereof.

25

23. A pharmaceutical composition comprising therapeutically effective amount of a compound of claim 1 having formula (I) or a pharmaceutically acceptable salt thereof, in combination with a pharmaceutically acceptable carrier.

30 24. A method of treating neuropathic pain, nociceptive pain, and inflammatory pain in a mammal in need of such treatment comprising administering to the mammal

a therapeutically effective amount of a compound of claim 1 having formula (I) or a pharmaceutically acceptable salt thereof.

25. A method of treating a disorder selected from the group consisting of
5 inflammatory disorders, immune disorders, neurological disorders, cancers of the immune system, respiratory disorders, and cardiovascular disorders in a mammal in need of such treatment, comprising administering to the mammal a therapeutically effective amount of a compound of claim 1 having formula (I), or a pharmaceutically acceptable salt thereof.

10

26. A method of providing neuroprotection in a mammal in need of such treatment comprising administering to the mammal a therapeutically effective amount of a compound of claim 1 having formula (I) or a pharmaceutically acceptable salt thereof.

15 27. A pharmaceutical composition comprising a therapeutically effective amount of a compound of claim 19 having formula (II) or a pharmaceutically acceptable salt thereof, in combination with a pharmaceutically acceptable carrier.

28. A method of treating neuropathic pain, nociceptive pain, and inflammatory
20 pain in a mammal in need of such treatment comprising administering to the mammal a therapeutically effective amount of a compound of claim 19 having formula (II) or a pharmaceutically acceptable salt thereof.

29. A method of treating a disorder selected from the group consisting of
25 inflammatory disorders, immune disorders, neurological disorders, cancers of the immune system, respiratory disorders, and cardiovascular disorders in a mammal in need of such treatment, comprising administering to the mammal a therapeutically effective amount of a compound of claim 19 having formula (II), or a pharmaceutically acceptable salt thereof.

30

30. A method of providing neuroprotection in a mammal in need of such treatment comprising administering to the mammal a therapeutically effective amount of a compound of claim 19 having formula (II) or a pharmaceutically acceptable salt thereof.