

QUINOLINE DERIVATIVES AS ANTIBACTERIAL AGENTS

The present invention relates to the use of quinoline derivatives for the manufacture of a medicament for the treatment of a bacterial infection.

5

Resistance to first-line antibiotic agents is an emerging problem. Some important examples include penicillin-resistant *Streptococcus pneumoniae*, vancomycin-resistant enterococci, methicillin-resistant *Staphylococcus aureus*, multi-resistant salmonellae.

10

The consequences of resistance to antibiotic agents are severe. Infections caused by resistant microbes fail to respond to treatment, resulting in prolonged illness and greater risk of death. Treatment failures also lead to longer periods of infectivity, which increase the numbers of infected people moving in the community and thus exposing the general population to the risk of contracting a resistant strain infection.

15

Hospitals are a critical component of the antimicrobial resistance problem worldwide. The combination of highly susceptible patients, intensive and prolonged antimicrobial use, and cross-infection has resulted in infections with highly resistant bacterial pathogens.

20

Self-medication with antimicrobials is another major factor contributing to resistance. Self-medicated antimicrobials may be unnecessary, are often inadequately dosed, or may not contain adequate amounts of active drug.

25

Patient compliance with recommended treatment is another major problem. Patients forget to take medication, interrupt their treatment when they begin to feel better, or may be unable to afford a full course, thereby creating an ideal environment for microbes to adapt rather than be killed.

30

Because of the emerging resistance to multiple antibiotics, physicians are confronted with infections for which there is no effective therapy. The morbidity, mortality, and financial costs of such infections impose an increasing burden for health care systems worldwide.

35


Therefore, there is a high need for new compounds to treat bacterial infections, especially for the treatment of infections caused by resistant strains.

WO 2004/011436 discloses substituted quinoline derivatives having activity against Mycobacteria, in particular against *Mycobacterium tuberculosis*. One particular compound of these substituted quinoline derivatives is described in Science (2005), 307, 223-227.

- 5 It has now been found that quinoline derivatives described in WO 2004/011436 also show activity against other bacteria than Mycobacteria.

Therefore, the present invention relates to the use of a compound for the manufacture of a medicament for the treatment of a bacterial infection, said compound being a

- 10 compound of formula



a pharmaceutically acceptable acid or base addition salt thereof, a stereochemically isomeric form thereof or a *N*-oxide form thereof, wherein

R¹ is hydrogen, halo, polyhaloC₁₋₆alkyl, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, Ar or Het;

- 15 p is an integer equal to 1 or 2;

R² is C₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyloxy or C₁₋₆alkylthio;

R³ is C₁₋₆alkyl, Ar, Het or Het¹;

R⁴ and R⁵ each independently are hydrogen, C₁₋₆alkyl or benzyl; or

R⁴ and R⁵ together and including the N to which they are attached may form a radical

- 20 selected from the group of pyrrolidinyl, 2-pyrrolinyl, 3-pyrrolinyl, pyrrolyl, imidazolidinyl, pyrazolidinyl, 2-imidazolinyl, 2-pyrazolinyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with
- 25 C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkylthioC₁₋₆alkyl or pyrimidinyl;

R⁶ is hydrogen, halo, polyhaloC₁₋₆alkyl, C₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio; or

two vicinal R⁶ radicals may be taken together to form a bivalent radical of formula -CH=CH-CH=CH- ;

5 r is an integer equal to 1 or 2;

R⁷ is hydrogen, C₁₋₆alkyl, Ar, Het or Het¹;

Ar is a homocycle selected from the group of phenyl, naphthyl, acenaphthyl, 1,2-dihydro-acenaphthyl, tetrahydronaphthyl, each homocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of hydroxy, halo, cyano, nitro, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, polyhaloC₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl, carboxyl, C₁₋₆alkyloxycarbonyl, aminocarbonyl, morpholinyl and mono- or di(C₁₋₆alkyl)aminocarbonyl;

15 Het is a monocyclic heterocycle selected from the group of *N*-phenoxypiperidinyl, piperidinyl, pyrrolyl, pyrazolyl, imidazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl; each monocyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, hydroxy, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl or Ar-C(=O)-;

20 Het¹ is a bicyclic heterocycle selected from the group of quinolinyl, quinoxalinyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl, benzothienyl, 2,3-dihydrobenzo[1,4]dioxinyl or benzo[1,3]dioxolyl ; each bicyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, hydroxy, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl or Ar-C(=O)-;

provided that the bacterial infection is other than a Mycobacterial infection.

30

The present invention also relates to a method of treating a bacterial infection in a mammal, in particular a warm-blooded mammal, more in particular a human, comprising administering an effective amount of a compound of the invention to the mammal.

35

As used hereinbefore or hereinafter C₁₋₆alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 6 carbon

atoms such as methyl, ethyl, propyl, 1-methylethyl, butyl, pentyl, hexyl, 2-methylbutyl and the like.

As used hereinbefore or hereinafter, the term (=O) forms a carbonyl moiety when
5 attached to a carbon atom.

The term halo is generic to fluoro, chloro, bromo and iodo. As used in the foregoing or hereinafter, polyhaloC₁₋₆alkyl as a group or part of a group is defined as mono- or polyhalosubstituted C₁₋₆alkyl, for example, methyl with one or more fluoro atoms, for
10 example, difluoromethyl or trifluoromethyl, 1,1-difluoro-ethyl and the like. In case more than one halogen atoms are attached to an alkyl group within the definition of polyhaloC₁₋₆alkyl, they may be the same or different.

In the definition of Het or Het¹, or when R⁴ and R⁵ are taken together, it is meant to
15 include all the possible isomeric forms of the heterocycles, for instance, pyrrolyl comprises 1*H*-pyrrolyl and 2*H*-pyrrolyl.

The Ar, Het or Het¹ listed in the definitions of the substituents of the compounds of formula (I) (see for instance R³) as mentioned hereinbefore or hereinafter may be
20 attached to the remainder of the molecule of formula (I) through any ring carbon or heteroatom as appropriate, if not otherwise specified. Thus, for example, when Het is imidazolyl, it may be 1-imidazolyl, 2-imidazolyl, 4-imidazolyl and the like.

Lines drawn from substituents into ring systems indicate that the bond may be attached
25 to any of the suitable ring atoms.

When two vicinal R⁶ radicals are taken together to form a bivalent radical of formula -CH=CH-CH=CH-, this means that the two vicinal R⁶ radicals form together with the phenyl ring to which they are attached a naphthyl.
30

For therapeutic use, salts of the compounds of formula (I) are those wherein the counterion is pharmaceutically acceptable. However, salts of acids and bases which are non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound. All salts, whether
35 pharmaceutically acceptable or not, are included within the ambit of the present invention.

The pharmaceutically acceptable addition salts as mentioned hereinabove or hereinafter are meant to comprise the therapeutically active non-toxic acid addition salt forms which the compounds of formula (I) are able to form. The latter can conveniently be obtained by treating the base form with such appropriate acids as inorganic acids, for example, hydrohalic acids, e.g. hydrochloric, hydrobromic and the like; sulfuric acid; 5 nitric acid; phosphoric acid and the like; or organic acids, for example, acetic, propanoic, hydroxyacetic, 2-hydroxypropanoic, 2-oxopropanoic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, 2-hydroxy-1,2,3-propanetricarboxylic, methanesulfonic, ethanesulfonic, benzenesulfonic, 4-methylbenzenesulfonic, 10 cyclohexanesulfamic, 2-hydroxybenzoic, 4-amino-2-hydroxybenzoic and the like acids. Conversely the salt form can be converted by treatment with alkali into the free base form.

The compounds of formula (I) containing acidic protons may be converted into their 15 therapeutically active non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. primary, secondary and tertiary aliphatic and aromatic amines such as methylamine, 20 ethylamine, propylamine, isopropylamine, the four butylamine isomers, dimethylamine, diethylamine, diethanolamine, dipropylamine, diisopropylamine, di-*n*-butylamine, pyrrolidine, piperidine, morpholine, trimethylamine, triethylamine, tripropylamine, quinuclidine, pyridine, quinoline and isoquinoline, the benzathine, *N*-methyl-D-glucamine, 2-amino-2-(hydroxymethyl)-1,3-propanediol, hydrabamine 25 salts, and salts with amino acids such as, for example, arginine, lysine and the like. Conversely the salt form can be converted by treatment with acid into the free acid form.

The term addition salt also comprises the hydrates and solvent addition forms which the compounds of formula (I) are able to form. Examples of such forms are e.g. hydrates, 30 alcoholates and the like.

The *N*-oxide forms of the present compounds are meant to comprise the compounds of formula (I) wherein one or several tertiary nitrogen atoms are oxidized to the so-called *N*-oxide. 35

The compounds of formula (I) may be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide

form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate
5 organic peroxides may comprise peroxy acids such as, for example, benzenecarboperoxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chlorobenzenecarboperoxoic acid, peroalkanoic acids, e.g. peroacetic acid, alkylhydroperoxides, e.g. *t*.butyl hydro-peroxide. Suitable solvents are, for example, water, lower alcohols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated
10 hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

It will be appreciated that some of the compounds of formula (I) and their *N*-oxides or addition salts may contain one or more centres of chirality and exist as stereochemically isomeric forms.

15 The term "stereochemically isomeric forms" as used hereinbefore or hereinafter defines all the possible stereoisomeric forms which the compounds of formula (I), and their *N*-oxides, addition salts or physiologically functional derivatives may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the
20 mixture of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure.


In particular, stereogenic centers may have the *R*- or *S*-configuration; substituents on bivalent cyclic (partially) saturated radicals may have either the *cis*- or *trans*-configuration. Compounds encompassing double bonds can have an *E* (*entgegen*) or *Z* (*zusammen*) -stereochemistry at said double bond.
25

Stereochemically isomeric forms of the compounds of formula (I) are obviously intended to be embraced within the scope of this invention.

Following CAS-nomenclature conventions, when two stereogenic centers of known
30 absolute configuration are present in a molecule, an *R* or *S* descriptor is assigned (based on Cahn-Ingold-Prelog sequence rule) to the lowest-numbered chiral center, the reference center. The configuration of the second stereogenic center is indicated using relative descriptors [*R**,*R**] or [*R**,*S**], where *R** is always specified as the reference center and [*R**,*R**] indicates centers with the same chirality and [*R**,*S**] indicates
35 centers of unlike chirality. For example, if the lowest-numbered chiral center in the molecule has an *S* configuration and the second center is *R*, the stereo descriptor would be specified as *S*-[*R**,*S**]. If "α" and "β" are used : the position of the highest priority

substituent on the asymmetric carbon atom in the ring system having the lowest ring number, is arbitrarily always in the "α" position of the mean plane determined by the ring system. The position of the highest priority substituent on the other asymmetric carbon atom in the ring system relative to the position of the highest priority substituent on the reference atom is denominated "α", if it is on the same side of the mean plane determined by the ring system, or "β", if it is on the other side of the mean plane determined by the ring system.

In the formula below, the two asymmetric carbon atoms on the lateral chain of the present compounds are indicated by *.



The stereodescriptors cis, trans, E, Z, R, S are well known to a person skilled in the art. Reference is also made to J.Am.Chem.Soc., 1982, 104, 5521-5523, which is incorporated herein by reference.


When a specific stereoisomeric form is indicated, this means that said form is substantially free, *i.e.* associated with less than 50 %, preferably less than 20 %, more preferably less than 10 %, even more preferably less than 5 %, further preferably less than 2 % and most preferably less than 1 % of the other isomer(s). Thus, when a compound of formula (I) is for instance specified as (α S, β R), this means that the compound is substantially free of the (α R, β S) isomer.

The compounds of formula (I) may be synthesized in the form of racemic mixtures of enantiomers which can be separated from one another following art-known resolution procedures. The racemic compounds of formula (I) may be converted into the corresponding diastereomeric salt forms by reaction with a suitable chiral acid. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali. An alternative manner of separating the enantiomeric forms of the compounds of formula (I) involves liquid chromatography using a chiral stationary phase. Said pure

stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

The invention also comprises derivative compounds (usually called "pro-drugs") of the pharmacologically-active compounds according to the invention, which are degraded *in vivo* to yield the compounds according to the invention. Pro-drugs are usually (but not always) of lower potency at the target receptor than the compounds to which they are degraded. Pro-drugs are particularly useful when the desired compound has chemical or physical properties that make its administration difficult or inefficient. For example, the desired compound may be only poorly soluble, it may be poorly transported across the mucosal epithelium, or it may have an undesirably short plasma half-life. Further discussion on pro-drugs may be found in Stella, V. J. *et al.*, "Prodrugs", *Drug Delivery Systems*, 1985, pp. 112-176, and *Drugs*, 1985, **29**, pp. 455-473.

Pro-drug forms of the pharmacologically-active compounds according to the invention will generally be compounds according to either formula (I), the pharmaceutically acceptable acid or base addition salts thereof, the stereochemically isomeric forms thereof, the tautomeric forms thereof and the *N*-oxide forms thereof, having an acid group which is esterified or amidated. Included in such esterified acid groups are groups of the formula $-\text{COOR}^x$, where R^x is a C_{1-6} alkyl, phenyl, benzyl or one of the following groups :




Amidated groups include groups of the formula $-\text{CONR}^y\text{R}^z$, wherein R^y is H, C_{1-6} alkyl, phenyl or benzyl and R^z is $-\text{OH}$, H, C_{1-6} alkyl, phenyl or benzyl.

Compounds according to the invention having an amino group may be derivatised with a ketone or an aldehyde such as formaldehyde to form a Mannich base. This base will hydrolyze with first order kinetics in aqueous solution.

Whenever used herein, the term "compounds of formula (I)" is meant to also include their *N*-oxide forms, their addition salts or their stereochemically isomeric forms. Of special interest are those compounds of formula (I) which are stereochemically pure.

- 5 A first interesting embodiment of the present invention relates to the use of a compound for the manufacture of a medicament for the treatment of a bacterial infection, said compound being a compound of formula



a pharmaceutically acceptable acid or base addition salt thereof, a stereochemically isomeric form thereof or a *N*-oxide form thereof, wherein

- 10 R^1 is hydrogen, halo, polyhaloC₁₋₆alkyl, C₁₋₆alkyl, Ar or Het;
 p is an integer equal to 1 or 2;
 R^2 is C₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyloxy or C₁₋₆alkylthio;
 R^3 is Ar, Het or Het¹;
 R^4 and R^5 each independently are hydrogen, C₁₋₆alkyl or benzyl; or
 15 R^4 and R^5 together and including the N to which they are attached may form a radical selected from the group of pyrrolidinyl, 2-pyrrolinyl, 3-pyrrolinyl, pyrrolyl, imidazolidinyl, pyrazolidinyl, 2-imidazolinyl, 2-pyrazolinyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and
 20 thiomorpholinyl, each of said rings may optionally be substituted with C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkylthioC₁₋₆alkyl or pyrimidinyl;
 R^6 is hydrogen, halo, polyhaloC₁₋₆alkyl, C₁₋₆alkyl, C₁₋₆alkyloxy,
 25 C₁₋₆alkylthio; or
 two vicinal R^6 radicals may be taken together to form a bivalent radical of formula -CH=CH-CH=CH- ;
 r is an integer equal to 1 or 2;

R⁷ is hydrogen, C₁₋₆alkyl, Ar, Het or Het¹;


Ar is a homocycle selected from the group of phenyl, naphthyl, acenaphthyl, tetrahydronaphthyl, each homocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of hydroxy, halo, cyano, nitro, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, polyhaloC₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl, carboxyl, C₁₋₆alkyloxycarbonyl, aminocarbonyl, morpholinyl and mono- or di(C₁₋₆alkyl)aminocarbonyl;

Het is a monocyclic heterocycle selected from the group of *N*-phenoxypiperidiny, piperidiny, pyrrolyl, pyrazolyl, imidazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl; each monocyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, hydroxy, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl or Ar-C(=O)-;

Het¹ is a bicyclic heterocycle selected from the group of quinolinyl, quinoxalinyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl, benzothienyl, 2,3-dihydrobenzo[1,4]dioxinyl or benzo[1,3]dioxolyl ; each bicyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, hydroxy, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl or Ar-C(=O)-;


provided that the bacterial infection is other than a Mycobacterial infection.

A second interesting embodiment of the present invention or any subgroup thereof as mentioned hereinbefore as interesting embodiment, relates to a compound of formula (I-a)



a pharmaceutically acceptable acid or base addition salt thereof, a stereochemically isomeric form thereof or a *N*-oxide form thereof.

A third interesting embodiment of the present invention or any subgroup thereof as mentioned hereinbefore as interesting embodiment, relates to a compound of formula (I-b)



- 5 a pharmaceutically acceptable acid or base addition salt thereof, a stereochemically isomeric form thereof or a *N*-oxide form thereof.

- A fourth interesting embodiment relates to the compounds of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R¹ represents hydrogen, halo, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, Ar or Het; preferably hydrogen, halo, C₁₋₆alkyl, Ar or Het; more preferably hydrogen, halo, optionally substituted phenyl or Het; even more preferably hydrogen, halo, optionally substituted phenyl, optionally substituted furanyl, or pyridinyl; still further preferred hydrogen, halo or optionally substituted phenyl; most preferred halo, such as for example bromo or chloro, in particular bromo.

- A fifth interesting embodiment relates to the compounds of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R¹ is halo, polyhaloC₁₋₆alkyl, Ar or Het; preferably halo, Ar or Het; more preferably halo, optionally substituted phenyl, or Het; even more preferably halo, phenyl, optionally substituted furanyl, or pyridinyl; most preferred halo or phenyl.

- A sixth interesting embodiment relates to the compounds of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R¹ is hydrogen, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, hydroxyC₁₋₆alkyl, Ar or Het, wherein Ar is preferably optionally substituted phenyl, even more preferably Ar is phenyl, and wherein Het is preferably optionally substituted furanyl or pyridinyl.

- A seventh interesting embodiment relates to the compounds of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R² is

C₁₋₆alkyloxy or C₁₋₆alkylthio, in particular methoxy or methylthio; preferably C₁₋₆alkyloxy; more preferably methoxy.

5 An eight interesting embodiment relates to the compounds of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R³ is C₁₋₆alkyl, Ar or Het or wherein R³ is Ar or Het or wherein R³ is Ar or Het¹; preferably R³ is Ar; more preferably optionally substituted phenyl or optionally substituted naphthyl; even more preferably phenyl optionally substituted with halo or C₁₋₆alkyloxy, or naphthyl optionally substituted with halo or C₁₋₆alkyloxy; most preferably phenyl
10 optionally substituted with 1 or 2 halo, in particular fluoro, or naphthyl, in particular 1-naphthyl or 2-naphthyl.

A ninth interesting embodiment relates to the compounds of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R⁴ and
15 R⁵ each independently are hydrogen or C₁₋₆alkyl; or R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from the group of pyrrolidinyl, 2-pyrrolinyl, 3-pyrrolinyl, pyrrolyl, imidazolidinyl, pyrazolidinyl, 2-imidazolinyl, 2-pyrazolinyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and
20 thiomorpholinyl, each of said rings may optionally be substituted with C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkylthioC₁₋₆alkyl or pyrimidinyl; more preferably R⁴ is C₁₋₆alkyl and R⁵ is hydrogen or C₁₋₆alkyl; or R⁴ and R⁵ together and including the N to which they are attached may form a radical selected
25 from the group of pyrrolidinyl, 2-pyrrolinyl, 3-pyrrolinyl, pyrrolyl, imidazolidinyl, pyrazolidinyl, 2-imidazolinyl, 2-pyrazolinyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, amino,
30 mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkylthioC₁₋₆alkyl or pyrimidinyl; even more preferably R⁴ is C₁₋₆alkyl and R⁵ is hydrogen or C₁₋₆alkyl; or R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from the group of pyrrolyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl,
35 triazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl,

C₁₋₆alkylthioC₁₋₆alkyl or pyrimidinyl; most preferably R⁴ is C₁₋₆alkyl, in particular methyl or ethyl, more in particular methyl, and R⁵ is hydrogen or C₁₋₆alkyl, in particular methyl or ethyl, more in particular methyl.

- 5 A tenth interesting embodiment relates to the compounds of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R⁴ and R⁵ each independently are hydrogen, C₁₋₆alkyl or benzyl; preferably R⁴ and R⁵ each independently are hydrogen or C₁₋₆alkyl; more preferably R⁴ is C₁₋₆alkyl and R⁵ is hydrogen or C₁₋₆alkyl; most preferred R⁴ and R⁵ are C₁₋₆alkyl, in particular methyl or ethyl, more in particular methyl.
- 10

- An eleventh interesting embodiment are the compounds of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R⁴ and R⁵ together and including the N to which they are attached may form a radical selected
- 15 from the group of pyrrolidinyl, 2-pyrrolinyl, 3-pyrrolinyl, pyrrolyl, imidazolidinyl, pyrazolidinyl, 2-imidazolinyl, 2-pyrazolinyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, amino,
- 20 mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkylthio C₁₋₆alkyl or pyrimidinyl; preferably R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from the group of pyrrolyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and thiomorpholinyl, each of said rings may
- 25 optionally be substituted with C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkylthioC₁₋₆alkyl or pyrimidinyl; more preferably R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from the group of pyrrolyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, piperazinyl,
- 30 morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkylthioC₁₋₆alkyl or pyrimidinyl; even more preferably R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from the group
- 35 of imidazolyl, triazolyl, piperidinyl, piperazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, amino, mono- or di(C₁₋₆alkyl)amino,

- C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkylthioC₁₋₆alkyl or pyrimidinyl; or R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from the group of pyrrolidinyl, pyrazinyl, imidazolyl, triazolyl, piperidinyl, piperazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with
- 5 C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkylthioC₁₋₆alkyl or pyrimidinyl; most preferred R⁴ and R⁵ together and including the N to which they are attached form imidazolyl.
- 10 Preferably, the substituents on the ring system when R⁴ and R⁵ are taken together, are selected from amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy, C₁₋₆alkylthio and pyrimidinyl; more preferably the substituents are selected from C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, C₁₋₆alkyloxy, C₁₋₆alkylthio and pyrimidinyl; even more preferably the substituents are selected from
- 15 C₁₋₆alkyl or pyrimidinyl.

- A twelfth interesting embodiment relates to the compounds of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R⁶ is hydrogen, halo, polyhaloC₁₋₆alkyl or C₁₋₆alkyl; or two vicinal R⁶ radicals may be taken
- 20 together to form a bivalent radical of formula -CH=CH-CH=CH-; or R⁶ is hydrogen, halo, C₁₋₆alkyloxy or C₁₋₆alkyl; or two vicinal R⁶ radicals may be taken together to form a bivalent radical of formula -CH=CH-CH=CH-; preferably R⁶ is hydrogen, halo, polyhaloC₁₋₆alkyl or C₁₋₆alkyl; more preferably R⁶ is hydrogen, halo or C₁₋₆alkyl; even more preferably R⁶ is hydrogen or halo; most preferred R⁶ is hydrogen.

25

A thirteenth interesting embodiment relates to the compounds of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein R⁷ is hydrogen.

- 30 A fourteenth interesting embodiment relates to the compounds of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein r is an integer equal to 1.

- A fifteenth interesting embodiment relates to the compounds of formula (I) or any
- 35 subgroup thereof as mentioned hereinbefore as interesting embodiment wherein p is an integer equal to 1.

A sixteenth interesting embodiment relates to the compounds of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment provided that when one R^1 is C_{1-6} alkyl then p is an integer equal to 2 and the other R^1 substituent is selected from halo, polyhalo C_{1-6} alkyl, C_{1-6} alkyl, Ar or Het.

5

A seventeenth interesting embodiment are the compounds of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein the compound is other than (α S, β R)-6-bromo- α -[2-(dimethylamino)ethyl]-2-methoxy- α -1-naphthalenyl- β -phenyl-3-quinolineethanol, a pharmaceutically acceptable acid or base addition salt thereof or a *N*-oxide form thereof.

10

An eighteenth interesting embodiment is the use of a compound of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment, for the manufacture of a medicament for the treatment of an infection with a gram-positive and/or a gram-negative bacterium.

15

A nineteenth interesting embodiment is the use of a compound of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment for the manufacture of a medicament for the treatment of an infection with a gram-positive bacterium.

20

A twentieth interesting embodiment is the use of the compounds of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment for the manufacture of a medicament for the treatment of an infection with a gram-negative bacterium.


25

A twenty first interesting embodiment is the use of a compound of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment for the manufacture of a medicament for the treatment of a bacterial infection wherein the compound of formula (I) has a $IC_{90} < 15 \mu\text{l/ml}$ against at least one bacterium, in particular a gram-positive bacterium; preferably a $IC_{90} < 10 \mu\text{l/ml}$; more preferably a $IC_{90} < 5 \mu\text{l/ml}$; the IC_{90} value being determined as described hereinafter.

30

A twenty second interesting embodiment relates to the use of a compound for the manufacture of a medicament for the treatment of a bacterial infection, said compound being a compound of formula

35



a pharmaceutically acceptable acid or base addition salt thereof, a stereochemically isomeric form thereof or a *N*-oxide form thereof, wherein

- R^1 is hydrogen, halo, polyhaloC₁₋₆alkyl, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, Ar or Het;
- 5 p is an integer equal to 1 or 2;
- R^2 is C₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyloxy or C₁₋₆alkylthio;
- R^3 is C₁₋₆alkyl, Ar, Het or Het¹;
- R^4 and R^5 each independently are hydrogen, C₁₋₆alkyl or benzyl; or
- R^4 and R^5 together and including the N to which they are attached may form a radical
- 10 selected from the group of pyrrolidinyl, 2-pyrrolinyl, 3-pyrrolinyl, pyrrolyl, imidazolidinyl, pyrazolidinyl, 2-imidazolinyl, 2-pyrazolinyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with
- 15 C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkylthioC₁₋₆alkyl or pyrimidinyl;
- R^6 is hydrogen, halo, polyhaloC₁₋₆alkyl, C₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio; or
- 20 two vicinal R^6 radicals may be taken together to form a bivalent radical of formula -CH=CH-CH=CH- ;
- r is an integer equal to 1 or 2;
- R^7 is hydrogen, C₁₋₆alkyl, Ar, Het or Het¹;
- Ar is a homocycle selected from the group of phenyl, naphthyl, acenaphthyl, 1,2-dihydro-acenaphthyl, tetrahydronaphthyl, each homocycle may optionally be
- 25 substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of hydroxy, halo, cyano, nitro, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, hydroxyC₁₋₆alkyl,

C₁₋₆alkyloxy, polyhaloC₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl, carboxyl, C₁₋₆alkyloxycarbonyl, aminocarbonyl, morpholinyl and mono- or di(C₁₋₆alkyl)aminocarbonyl;

5 Het is a monocyclic heterocycle selected from the group of *N*-phenoxypiperidinyl, piperidinyl, pyrrolyl, pyrazolyl, imidazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl; each monocyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, hydroxy, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, 10 C₁₋₆alkyloxyC₁₋₆alkyl or Ar-C(=O)-;

Het¹ is a bicyclic heterocycle selected from the group of quinolinyl, quinoxalinyl, indolyl, benzimidazolyl, benzoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl, benzothienyl, 2,3-dihydrobenzo[1,4]dioxinyl or benzo[1,3]dioxolyl ; each bicyclic heterocycle may optionally be substituted 15 with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, hydroxy, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl or Ar-C(=O)-;

provided that when R³ is Ar, Het or Het¹; then R¹ is hydroxyC₁₋₆alkyl or C₁₋₆alkyloxy; and

20 provided that the bacterial infection is other than a Mycobacterial infection.

Preferably, the compounds of formula (I') are compounds wherein one or more, preferably all, of the following definitions apply :

R¹ is halo, in particular bromo; hydroxyC₁₋₆alkyl, in particular hydroxymethyl; or 25 C₁₋₆alkyloxy, in particular methoxy;
R² is C₁₋₆alkyloxy, in particular methoxy;
R³ is C₁₋₆alkyl, phenyl or naphthyl;
R⁴ and R⁵ are C₁₋₆alkyl;
R⁶ is hydrogen;
30 R⁷ is hydrogen;
p is 1;
r is 1.

35 Interesting compounds of formula (I') are compounds K, L, M, N, P and Q.

Also interesting compounds of the present invention are those compounds of formula (I) wherein one or more, preferably all, of the following definitions apply :

- a) R¹ is hydrogen; C₁₋₆alkyl; halo, in particular bromo or chloro; hydroxyC₁₋₆alkyl; C₁₋₆alkyloxy; optionally substituted phenyl, in particular phenyl; or Het, in particular furanyl optionally substituted with hydroxyC₁₋₆alkyl, or pyridyl;
- b) R² is C₁₋₆alkyloxy, in particular methoxy or ethoxy; C₁₋₆alkylthio, in particular methylthio; or C₁₋₆alkyloxyC₁₋₆alkyloxy;
- 5 c) R³ is phenyl optionally substituted with 1 or 2 halo, in particular fluoro or chloro; naphthyl optionally substituted with 1 or 2 halo or C₁₋₆alkyloxy; thienyl; piperidinyl substituted with Ar-C(=O); 2,3-dihydrobenzo[1,4]dioxinyl; benzo[1,3]dioxolyl; or acenaphthyl; furanyl; or C₁₋₆alkyl;
- 10 d) R⁴ and R⁵ are each independently hydrogen; C₁₋₆alkyl; benzyl; or R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from imidazolyl; pyrazinyl substituted with C₁₋₆alkyl; piperazinyl substituted with C₁₋₆alkyl; piperazinyl substituted with pyrimidinyl; piperidinyl; piperidinyl substituted with dimethylamino; thiomorpholinyl; morpholinyl; pyrrolidinyl; or triazolyl;
- 15 e) R⁶ is hydrogen; halo, in particular chloro, fluoro or bromo; C₁₋₆alkyl; C₁₋₆alkyloxy or two vicinal R⁶ radicals may be taken together to form a bivalent radical of formula -CH=CH-CH=CH-;
- f) R⁷ is hydrogen.
- 20 Preferred compounds of the present invention are compound 50, 206, 31, 26, 27, 32, 33, 109, 39, 44, 41, compound A, E and F, a pharmaceutically acceptable acid or base addition salt thereof or a *N*-oxide form thereof.

Another group of preferred compounds are the compounds defined in claim 30, i.e.

25 compounds 36, 46, 206, 31, 26, 33, 13, 39, 44, Compound A, E, F, R and S, a pharmaceutically acceptable acid or base addition salt thereof or a *N*-oxide form thereof.

Yet other preferred compounds are compound R and compound S, a pharmaceutically

30 acceptable acid or base addition salt thereof or a *N*-oxide form thereof.

The present invention also relates to a compound as claimed in claim 35, i.e.

compounds A to I, K to Z, A' to R', a pharmaceutically acceptable acid or base addition salt thereof or a *N*-oxide form thereof, in particular to compounds K to Z, and

35 A' to R', a pharmaceutically acceptable acid or base addition salt thereof or a *N*-oxide form thereof.

The present invention also relates to a compound of formula (I) or any subgroup thereof as mentioned hereinbefore as interesting embodiment wherein Ar is 1,2-dihydro-acenaphthyl.

- 5 The compounds of formula (I) can be prepared according to the methods described in WO 2004/011436, which is incorporated herein by reference.


In general, the compounds according to the invention can be prepared by a succession of steps, each of which is known to the skilled person.

10

In particular, the compounds according to formula (I) can be prepared by reacting an intermediate compound of formula (II) with an intermediate compound of formula (III) according to the following reaction scheme (1) :

15

Scheme 1




using BuLi in a mixture of diisopropyl amine and tetrahydrofuran, and wherein all variables are defined as in formula (I). Stirring may enhance the rate of the reaction. The reaction may conveniently be carried out at a temperature ranging between -20 and -70 °C.

20

The starting materials and the intermediate compounds of formula (II) and (III) are compounds that are either commercially available or may be prepared according to conventional reaction procedures generally known in the art. For example, intermediate compounds of formula (II-a) may be prepared according to the following reaction scheme (2):

25

Scheme 2




- wherein all variables are defined as in formula (I). Reaction scheme (2) comprises step
- 5 (a) in which an appropriately substituted aniline is reacted with an appropriate acylchloride such as 3-phenylpropionyl chloride, 3-fluorobenzenepropanoyl chloride or *p*-chlorobenzenepropanoyl chloride, in the presence of a suitable base, such as triethylamine and a suitable reaction-inert solvent, such as methylene chloride or ethylene dichloride. The reaction may conveniently be carried out at a temperature
- 10 ranging between room temperature and reflux temperature. In a next step (b) the adduct obtained in step (a) is reacted with phosphoryl chloride (POCl_3) in the presence of *N,N*-dimethylformamide (Vilsmeier-Haack formylation followed by cyclization). The reaction may conveniently be carried out at a temperature ranging between room temperature and reflux temperature. In a next step (c) a specific R^2 -group, wherein R^2
- 15 is for example an C_{1-6} alkyloxy or C_{1-6} alkylthio radical is introduced by reacting the intermediate compound obtained in step (b) with a compound H-X-C_{1-6} alkyl wherein X is S or O.

- It is evident that in the foregoing and in the following reactions, the reaction products
- 20 may be isolated from the reaction medium and, if necessary, further purified according to methodologies generally known in the art, such as extraction, crystallization and chromatography. It is further evident that reaction products that exist in more than one enantiomeric form, may be isolated from their mixture by known techniques, in particular preparative chromatography, such as preparative HPLC. Typically,
- 25 compounds of formula (I) may be separated into their isomeric forms.

The intermediate compounds of formula (III) are compounds that are either commercially available or may be prepared according to conventional reaction procedures generally known in the art. For example, intermediate compounds of formula (III-a) in which R^3 is Ar substituted with s substituents R^{10} , wherein each R^{10} is independently selected from the group of hydroxy, halo, cyano, nitro, amino, mono- or di(C_{1-6} alkyl)amino, C_{1-6} alkyl, polyhalo C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkyloxy, polyhalo C_{1-6} alkyloxy, C_{1-6} alkyloxy C_{1-6} alkyl, carboxyl, C_{1-6} alkyloxycarbonyl, aminocarbonyl, morpholinyl and mono- or di(C_{1-6} alkyl)aminocarbonyl and s is an integer equal to zero, 1, 2 or 3, may be prepared according to the following reaction scheme (3):

Scheme 3



Reaction scheme (3) comprises step (a) in which an appropriately substituted Ar, in particular an appropriately substituted phenyl, is reacted by Friedel-Craft reaction with an appropriate acylchloride such as 3-chloropropionyl chloride, in the presence of a suitable Lewis acid, such as for example $AlCl_3$, $FeCl_3$, $SnCl_4$, $TiCl_4$ or $ZnCl_2$ and a suitable reaction-inert solvent, such as methylene chloride or ethylene dichloride. The reaction may conveniently be carried out at a temperature ranging between room temperature and reflux temperature. In a next step (b) an amino group ($-NR^4R^5$) is introduced by reacting the intermediate compound obtained in step (a) with HNR^4R^5 .

In general, bacterial pathogens may be classified as either gram-positive or gram-negative pathogens. Antibiotic compounds with activity against both gram-positive and gram-negative pathogens are generally regarded as having a broad spectrum of activity. The compounds of the present invention are regarded as active against gram-positive and/or gram-negative bacterial pathogens, in particular against gram-positive bacterial pathogens. In particular, the present compounds are active against at least one gram-positive bacterium, preferably against several gram-positive bacteria, more preferably against one or more gram-positive bacteria and/or one or more gram-negative bacteria.

The present compounds have bactericidal or bacteriostatic activity.

Examples of gram-positive and gram-negative aerobic and anaerobic bacteria, include Staphylococci, for example *S. aureus*; Enterococci, for example *E. faecalis*; Streptococci, for example *S. pneumoniae*, *S. mutans*, *S. pyogenes*; Bacilli, for example
5 *Bacillus subtilis*; Listeria, for example *Listeria monocytogenes*; Haemophilus, for example *H. influenza*; Moraxella, for example *M. catarrhalis*; Pseudomonas, for example *Pseudomonas aeruginosa*; and Escherichia, for example *E. coli*.

Gram-positive pathogens, for example Staphylococci, Enterococci and Streptococci are particularly important because of the development of resistant strains which are both
10 difficult to treat and difficult to eradicate from for example a hospital environment once established. Examples of such strains are methicillin resistant *Staphylococcus aureus* (MRSA), methicillin resistant coagulase negative staphylococci (MRCNS), penicillin resistant *Streptococcus pneumoniae* and multiple resistant *Enterococcus faecium*.

15 The compounds of the present invention also show activity against resistant bacterial strains.

The compounds of the present invention are especially active against *Staphylococcus aureus*, including resistant *Staphylococcus aureus* such as for example methicillin
20 resistant *Staphylococcus aureus* (MRSA), and *Streptococcus pneumoniae*.

In particular, the compounds of the present invention are active on those bacteria of which the viability depends on proper functioning of F1F0 ATP synthase. Without being bound to any theory, it is taught that the activity of the present compounds lies in
25 inhibition of the F1F0 ATP synthase, in particular the inhibition of the F0 complex of the F1F0 ATP synthase, more in particular the inhibition of subunit c of the F0 complex of the F1F0 ATP synthase, leading to killing of the bacteria by depletion of the cellular ATP levels of the bacteria.

30 Whenever used hereinbefore or hereinafter, that the compounds can treat a bacterial infection it is meant that the compounds can treat an infection with one or more bacterial strains.

Whenever used hereinbefore or hereinafter, that the bacterial infection is other than a Mycobacterial infection it is meant that the bacterial infection is other than an infection
35 with one or more Mycobacterial strains.

The exact dosage and frequency of administration of the present compounds depends on the particular compound of formula (I) used, the particular condition being treated, the severity of the condition being treated, the age, weight, gender, diet, time of administration and general physical condition of the particular patient, the mode of administration as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that the effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds of the instant invention.

10

The compound of the present invention may be administered in a pharmaceutically acceptable form optionally in a pharmaceutically acceptable carrier. The compounds and compositions comprising the compounds can be administered by routes such as topically, locally or systemically. Systemic application includes any method of introducing the compound into the tissues of the body, e.g., intrathecal, epidural, intramuscular, transdermal, intravenous, intraperitoneal, subcutaneous, sublingual, rectal, and oral administration. The specific dosage of antibacterial to be administered, as well as the duration of treatment, may be adjusted as needed.

20

Bacterial infections which may be treated by the present compounds include, for example, central nervous system infections, external ear infections, infections of the middle ear, such as acute otitis media, infections of the cranial sinuses, eye infections, infections of the oral cavity, such as infections of the teeth, gums and mucosa, upper respiratory tract infections, lower respiratory tract infections, genitourinary infections, gastrointestinal infections, gynecological infections, septicemia, bone and joint infections, skin and skin structure infections, bacterial endocarditis, burns, antibacterial prophylaxis of surgery, and antibacterial prophylaxis in immunosuppressed patients, such as patients receiving cancer chemotherapy, or organ transplant patients.

25

30

Given the fact that the compounds of formula (I) are active against gram-positive and/or gram-negative bacteria, the present compounds may be combined with other antibacterial agents in order to effectively combat bacterial infections.

35

Therefore, the present invention also relates to a combination of (a) a compound of formula (I), and (b) one or more other antibacterial agents provided that the one or more other antibacterial agents are other than antimycobacterial agents.

The present invention also relates to a combination of (a) a compound of formula (I), and (b) one or more other antibacterial agents provided that the one or more other antibacterial agents are other than antimycobacterial agents, for use as a medicine.

- 5 A pharmaceutical composition comprising a pharmaceutically acceptable carrier and, as active ingredient, a therapeutically effective amount of (a) a compound of formula (I), and (b) one or more other antibacterial agents provided that the one or more other antibacterial agents are other than antimycobacterial agents, is also comprised by the present invention.

10

The present invention also relates to the use of a combination or pharmaceutical composition as defined above for the treatment of a bacterial infection, in particular a bacterial infection other than a Mycobacterial infection.

- 15 The present pharmaceutical composition may have various pharmaceutical forms for administration purposes. As appropriate compositions there may be cited all compositions usually employed for systemically administering drugs. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compounds, optionally in addition salt form, as the active ingredient is combined in
20 intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, in particular, for administration orally or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media
25 may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs, emulsions and solutions; or solid carriers such as starches, sugars, kaolin, diluents, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules and tablets. Because of their ease in administration, tablets and capsules represent the most
30 advantageous oral unit dosage forms in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and
35 glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also

included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations.

Depending on the mode of administration, the pharmaceutical composition will
5 preferably comprise from 0.05 to 99 % by weight, more preferably from 0.1 to 70 % by weight of the active ingredients, and, from 1 to 99.95 % by weight, more preferably from 30 to 99.9 weight % of a pharmaceutically acceptable carrier, all percentages being based on the total composition.

10 The weight to weight ratio's of the compound of formula (I) and (b) the other antibacterial agent(s) when given as a combination may be determined by the person skilled in the art. Said ratio and the exact dosage and frequency of administration depends on the particular compound of formula (I) and the other antibacterial agent(s) used, the particular condition being treated, the severity of the condition being treated,
15 the age, weight, gender, diet, time of administration and general physical condition of the particular patient, the mode of administration as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that the effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician
20 prescribing the compounds of the instant invention.

The compounds of formula (I) and the one or more other antibacterial agents may be combined in a single preparation or they may be formulated in separate preparations so that they can be administered simultaneously, separately or sequentially. Thus, the
25 present invention also relates to a product or a kit containing (a) a compound of formula (I), and (b) one or more other antibacterial agents provided that the one or more other antibacterial agents are other than antimycobacterial agents, as a combined preparation for simultaneous, separate or sequential use in the treatment of a bacterial infection.

30 The pharmaceutical composition may additionally contain various other ingredients known in the art, for example, a lubricant, stabilising agent, buffering agent, emulsifying agent, viscosity-regulating agent, surfactant, preservative, flavouring or colorant.

35 It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage.

Unit dosage form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, suppositories, injectable solutions or suspensions and the like, and segregated multiples thereof. The daily dosage of the compound according to the invention will, of course, vary with the compound employed, the mode of administration, the treatment desired and the bacterial disease indicated.

The other antibacterial agents which may be combined with the compounds of formula (I) are antibacterial agents known in the art. The other antibacterial agents comprise antibiotics of the β -lactam group such as natural penicillins, semisynthetic penicillins, natural cephalosporins, semisynthetic cephalosporins, cephamycins, 1-oxacephems, clavulanic acids, penems, carbapenems, nocardicins, monobactams; tetracyclines, anhydrotetracyclines, anthracyclines; aminoglycosides; nucleosides such as *N*-nucleosides, *C*-nucleosides, carbocyclic nucleosides, blasticidin S; macrolides such as 12-membered ring macrolides, 14-membered ring macrolides, 16-membered ring macrolides; ansamycins; peptides such as bleomycins, gramicidins, polymyxins, bacitracins, large ring peptide antibiotics containing lactone linkages, actinomycins, amphomycin, capreomycin, distamycin, enduracidins, mikamycin, neocarzinostatin, stendomycin, viomycin, virginiamycin; cycloheximide; cycloserine; variotin; sarkomycin A; novobiocin; griseofulvin; chloramphenicol; mitomycins; fumagillin; monensins; pyrrolnitrin; fosfomicin; fusidic acid; D-(*p*-hydroxyphenyl)glycine; D-phenylglycine; enediynes.

Specific antibiotics which may be combined with the present compounds of formula (I) are for example benzylpenicillin (potassium, procaine, benzathine), phenoxymethylpenicillin (potassium), phenethicillin potassium, propicillin, carbenicillin (disodium, phenyl sodium, indanyl sodium), sulbenicillin, ticarcillin disodium, methicillin sodium, oxacillin sodium, cloxacillin sodium, dicloxacillin, flucloxacillin, ampicillin, mezlocillin, piperacillin sodium, amoxicillin, ciclacillin, hecicillin, sulbactam sodium, talampicillin hydrochloride, bacampicillin hydrochloride, pivmecillinam, cephalixin, cefaclor, cephaloglycin, cefadroxil, cephadrine, cefroxadine, cephapirin sodium, cephalothin sodium, cephacetrile sodium, cefsulodin sodium, cephaloridine, cefatrizine, cefoperazone sodium, cefamandole, vefotiam hydrochloride, cefazolin sodium, ceftizoxime sodium, cefotaxime sodium, cefmenoxime hydrochloride, cefuroxime, ceftriaxone sodium, ceftazidime, cefoxitin,

5 cefmetazole, cefotetan, latamoxef, clavulanic acid, imipenem, aztreonam, tetracycline, chlortetracycline hydrochloride, demethylchlortetracycline, oxytetracycline, methacycline, doxycycline, rolitetracycline, minocycline, daunorubicin hydrochloride, doxorubicin, aclarubicin, kanamycin sulfate, bekanamycin, tobramycin, gentamycin
10 sulfate, dibekacin, amikacin, micronomicin, ribostamycin, neomycin sulfate, paromomycin sulfate, streptomycin sulfate, dihydrostreptomycin, destomycin A, hygromycin B, apramycin, sisomicin, netilmicin sulfate, spectinomycin hydrochloride, astromicin sulfate, validamycin, kasugamycin, polyoxin, blasticidin S, erythromycin, erythromycin estolate, oleandomycin phosphate, tracetyloleandomycin, kitasamycin,
15 josamycin, spiramycin, tylosin, ivermectin, midecamycin, bleomycin sulfate, peplomycin sulfate, gramicidin S, polymyxin B, bacitracin, colistin sulfate, colistinmethanesulfonate sodium, enramycin, mikamycin, virginiamycin, capreomycin sulfate, viomycin, enviomycin, vancomycin, actinomycin D, neocarzinostatin, bestatin, pepstatin, monensin, lasalocid, salinomycin, amphotericin B, nystatin, natamycin,
20 trichomycin, mithramycin, lincomycin, clindamycin, clindamycin palmitate hydrochloride, flavophospholipol, cycloserine, pecilocin, griseofulvin, chloramphenicol, chloramphenicol palmitate, mitomycin C, pyrrolnitrin, fosfomycin, fusidic acid, bicozamycin, tiamulin, siccanin.

20 EXPERIMENTAL PART


Of some compounds the absolute stereochemical configuration of the stereogenic carbon atom(s) therein was not experimentally determined. In those cases the stereochemically isomeric form which was first isolated is designated as "A" and the second as "B", without further reference to the actual stereochemical configuration.
25 However, said "A" and "B" isomeric forms can be unambiguously characterized by a person skilled in the art, using art-known methods such as, for example, X-ray diffraction.
In case "A" and "B" are stereoisomeric mixtures, they can be further separated whereby the respective first fractions isolated are designated "A1" respectively "B1"
30 and the second as "A2" respectively "B2", without further reference to the actual stereochemical configuration. However, said "A1, A2" and "B1, B2" isomeric forms can be unambiguously characterized by a person skilled in the art, using art-known methods such as, for example, X-ray diffraction.
35 Whenever hereinabove or hereinafter, a compound is designated as "A" or "B" this means that the compound is a mixture of two enantiomers. Whenever hereinabove or

hereinafter, a compound is designated as "A1", "A2", "B1" or "B2" this means that the compound is an enantiomer.

Present compounds which are also described in WO2004/011436, bear the same
5 compound number as in WO2004/011436. The Ex. Nr. in the below Tables refer to the Example numbers of WO 2004/011436 indicating according to which procedure the compounds were prepared.

Additional compounds are indicated by way of letters.

10 Compound A, B and C were prepared according to the procedures described in WO 2004/011436. The scheme below indicates the synthesis pathway of compound A, B and C. The Example numbers A8, A9, B12 and B13 correspond to the procedures of WO 2004/011436.



15 Present intermediate 1 was prepared in the same way as intermediate 12 of WO 2004/011436, i.e. according to Example A8 of WO 2004/011436, but starting from 1,3-difluorobenzene. Yield : 57 % of present intermediate 1.

Present intermediate 2 was prepared in the same way as intermediate 14 of WO
20 2004/011436, i.e. according to Example A9 of WO 2004/011436, but starting from present intermediate 1 and reacting it with N-ethylbenzylamine. Yield : 88% of present intermediate 2.

Compound C was prepared as follows :

25 present intermediate 2 was reacted with intermediate compound 3 of WO 2004/011436 in the same way as described in Example B12 of WO 2004/011436. The residue (5.4g)

obtained according to the procedure of B12, which is a mixture of diastereoisomers, was purified by column chromatography over silica gel (eluent: CH₂Cl₂/Cyclohexane: 60/40). Two fractions were collected and the solvent was evaporated. The second fraction was crystallized from diisopropylether yielding 0.83g of compound C (diastereoisomer B) (yield:13 %).


Compounds A and B were prepared as follows :

The residue obtained in the synthesis of compound C was reacted with 1-chloroethyl chloroformate in the same way as described in Example B13 of WO 2004/011436.

The residue (1.7g) obtained according to the procedure of Example B13, which is mixture of diastereoisomers, was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH; 98/2/0.1). Two fractions were collected and the solvent was evaporated. The fractions were crystallized separately from diisopropylether yielding 0.31g of compound B (diastereoisomer A) (yield:27 %) and 0.52g of compound A (diastereoisomer B) (yield:45 %).

Compound D was prepared according to the procedures described in WO 2004/011436.

The scheme below indicates the synthesis pathway of compound D. The Example numbers A9, B12 and B13 correspond to the procedures of WO 2004/011436.



Present intermediate 3 was prepared in the same way as intermediate 14 of WO 2004/011436, i.e. according to Example A9 of WO 2004/011436, but starting from 3-chloropropiophenone. Yield : 98 % of present intermediate 3.

Compound J was prepared as follows :

present intermediate 3 was reacted with intermediate compound 3 of WO 2004/011436 in the same way as described in Example B12 of WO 2004/011436. The obtained residue (4.9g) was purified by column chromatography over silica gel (eluent :
5 CH_2Cl_2). One fraction was collected and the solvent was evaporated. Yield : 1.43g of compound J, which is a mixture of diastereoisomers.

Compound D was prepared as follows :

compound J was reacted with 1-chloroethyl formate in the same way as described in
10 Example B13 of WO 2004/011436. The residue (1.2g) obtained according to the procedure of Example B13 was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}/\text{NH}_4\text{OH}$; 95/5/0.5). Two fractions were collected and the solvent was evaporated. The second fraction was crystallized from diisopropylether yielding 0.08g of compound D (diastereoisomer B) (yield:10 %).

15

Compounds E and F were prepared by separating compound 4 (diastereoisomer B) of WO 2004/011436 into its enantiomers by the following procedure :

Final compound 4 of WO 2004/011436 (2.5g) was separated into its enantiomers by column chromatography (eluent: hexane/ethanol 99.95/0.05; column: CHIRACEL
20 OD). Two pure fractions were collected and their solvents were evaporated. Yield: 0.5g of compound E (enantiomer B1) (mp. 180°C) and 0.12g of compound F (enantiomer B2) (mp. 175°C).

Compound G was prepared as follows :


25 A mixture of compound 115 of WO 2004/011436 (prepared according to B15 of WO 2004/011436) (0.00028 mol), pyridine-3-boronic acid 1,3-propanediol cyclic ester (0.00055 mol), $\text{Pd}[\text{P}(\text{Ph}_3)]_4$ (0.00003 mol) and Na_2CO_3 2M (0.0011 mol) in dimethoxyethane (4ml) was stirred at 90°C for 1.5 hours, then poured out into H_2O and extracted with CH_2Cl_2 . The organic layer was separated, dried (MgSO_4), filtered and
30 the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: $\text{CH}_2\text{Cl}_2/\text{CH}_3\text{OH}$ 95/5; $5\mu\text{m}$). The pure fractions were collected and the solvent was evaporated. Yield: 0.09g of compound G (60 %) (mp. 201°C).

Compound H was prepared as follows :

35 A mixture of compound 15 of WO 2004/011436 (prepared according to B7 of WO 2004/011436) (0.0009 mol), 2-furanboronic acid (0.0018 mol), $\text{Pd}[\text{P}(\text{Ph}_3)]_4$ (0.00009 mol) and Na_2CO_3 2M (0.0036 mol) in dimethoxyethane (10ml) was stirred at 90°C for

6 hours, then poured out into H₂O and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered and the solvent was evaporated. The residue (0.57g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 99/1/0.1; 10μm). The pure fractions were collected and the solvent was evaporated, yielding 0.23g of residue. This fraction was crystallized from diisopropylether/acetonitrile. The precipitate was filtered off and dried. Yield : 0.071g of compound H (15 %) (mp. 215°C).

Compound I was prepared according to the procedures described in WO 2004/011436. The scheme below indicates the synthesis pathway of compound I. The Example numbers A10 and B1 correspond to the procedures of WO 2004/011436.




A mixture of commercially available 5-acetyl-1,2-dihydroacenaphthylene (0.0407 mol) and dimethylamine hydrochloride (0.0858 mol) in paraformaldehyde (37 % in water, 4ml), HCl/iPrOH (1ml) and ethanol (100ml) was stirred and refluxed for 48 hours. The solvent was evaporated. The residue was taken up in H₂O/HCl 3N/CH₂Cl₂. The aqueous layer was basified and extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered, and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂/CH₃OH/NH₄OH 95/5/0.2; 15-40μm). The pure fractions were collected and the solvent was evaporated. Yield: 4.9g of intermediate 4 (48 %).

n-BuLi 1.6M (0.0102 mol) was added dropwise at -20°C to a solution of diisopropylamine (0.0091 mol) in tetrahydrofuran (15ml) under N₂ flow. The mixture was stirred at -20°C for 20 minutes, then cooled to -70°C. A solution of intermediate compound 3 of WO 2004/011436 (0.0091 mol) in tetrahydrofuran (10ml) was added dropwise. The mixture was stirred at -70°C for 2 hours. A solution of present intermediate 4 (0.01 mol) in tetrahydrofuran (20ml) was added dropwise. The mixture was stirred at -70°C for 3 hours, poured out into ice water and extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered, and the solvent was evaporated. The residue (6.5g) was purified by column chromatography over silica gel

(eluent: CH₂Cl₂/CH₃OH/NH₄OH 99/1/0.2; 15-40μm). Two fractions were collected and the solvent was evaporated. Yield: 0.96g F1, 0.72g F2. F1 was crystallized from diethyl ether. The precipitate was filtered off and dried. Yield: 0.87g of compound I (17 %).

5 Compounds Q and P were prepared according to the synthesis pathway shown below.



Compound Q was synthesized as follows:

n-BuLi (1.6M in hexane, 14.1ml, 0.0225mol) was added dropwise at -70°C under nitrogen flow to a solution of compound 14 of WO 2004/011436 (prepared according to B7 of WO 2004/011436) (5.0g, 9.0mmol) in tetrahydrofuran (50ml). The mixture was stirred for 90 minutes at -70°C and then *N,N*-dimethylformamide (5.5ml, 0.072mol) was added. The resulting mixture was stirred for 2 hours at -70°C. Then water was added. The mixture was extracted with EtOAc. The organic layer was washed with water and then brine, dried over MgSO₄, filtered and evaporated till dryness. The residue was crystallized from diisopropylether and CH₃OH. Yield: 1.7g of intermediate 5 (A) (38%). Intermediate 6 was prepared from compound 15 according to the same procedure as described for intermediate 5.

Sodium borohydride (7.6mg, 0.2mmol) was added at 0°C to a solution of present intermediate 5 (0.1g, 0.2mmol) in CH₃OH (2ml) and tetrahydrofuran (2ml). The mixture was stirred for 2 hours at 0°C, hydrolyzed with water and extracted with


EtOAc. The organic layer was washed with water then brine, was dried over MgSO_4 , filtered and evaporated till dryness. The residue (0.09g) was crystallized from CH_3OH . Yield: compound Q (A) (15%; mp: 187 °C).

Compound P (B) (196 °C) was synthesized from present intermediate 6 according to the same procedure as described for compound Q.

Compound R was prepared from compound 56 as follows :

Compound 56 (0.0016 mol) was purified by column chromatography over Chiralpak AD (eluent : ethanol 100%). Two fractions were collected and the solvent was evaporated. The second fraction was compound R. Yield : 0.397g.

Compound S was prepared as follows :



A mixture of intermediate compound 13 of WO2004/011436 (0.016 mol), N-benzylmethyl amine (0.016 mol) and K_2CO_3 (0.0176 mol) in acetonitrile (50 ml) was stirred at 80 °C for 4 hours, then brought to room temperature, poured out into water and extracted with methylene chloride. The organic layer was washed with saturated NaCl, dried (MgSO_4), filtered and the solvent was evaporated. Yield : 4.85 g of present intermediate 7.

nBuLi 1.6M in hexane (0.0159 mol) was added dropwise at -20°C under N_2 flow to a solution of N-(1-methylethyl)-2-propanamine (0.0159 mol) in tetrahydrofuran (30ml) . The mixture was stirred at -20°C for 20 minutes, then cooled to -70°C. A solution of intermediate compound 3 of WO2004/011436 (0.0132 mol) in tetrahydrofuran (45ml) was added. The mixture was stirred for 1 hour and 30 minutes. A solution of present intermediate 7 (0.0159 mol) in tetrahydrofuran (45ml) was added. The mixture was stirred at -70°C for 3 hours, then poured out on ice at -30°C and extracted with EtOAc.

The organic layer was washed with saturated NaCl, dried (MgSO₄), filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH₂Cl₂ 100 then cyclohexane/EtOAc 90/10; 15-40μm). Two fractions were collected and the solvent was evaporated. The second fraction yielded 3.3 g of compound S'.

1-chloroethyl chloroformate (0.0048mol) was added at room temperature to a solution of compound S' (0.0048mol) in dichloromethane (34.5 ml). The mixture was stirred at 80°C for 1 hour, then brought to room temperature and evaporated till dryness.


Methanol (34.5ml) was added. The mixture was stirred at 80°C for 30 minutes, then

brought to room temperature and evaporated till dryness. The residue (2.9g) was purified by column chromatography over silica gel (eluent:CH₂Cl₂/CH₃OH/NH₄OH : 96/4/0.1; 15-40μm). Four fractions were collected and the solvent was evaporated. Yield : 0.11g F1; 0.13g F2; 0.12g F3 and 0.17g F4. F1, F2 and F3 were purified by column chromatography over kromasil (eluent: CH₂Cl₂/CH₃OH/NH₄OH : 98/2/0.1; 10μm). Two fractions were collected and the solvent was evaporated. Yield : 0.113g F1 and 0.13g F2. Both fractions were crystallized from diisopropyl ether/diethyl ether. The precipitate was filtered off and dried. Yield : 0.05g of F1 (the A diastereoisomer) and 0.12 g of compound S (B) (198°C).




Compounds K to O, R to Z, A' to R' were prepared according to the procedures described in WO 2004/011436. In the below Tables, the Ex.nr. given for said compounds refer to the corresponding Ex.nr. of WO 2004/011436 indicating according to which procedure the compounds were prepared.







Tables 1 to 5 list compounds of formula (I) according to the present invention.

Table 1 :




Comp. nr.	Ex. nr.	R ¹	R ²	R ³	R ⁶	Stereochemistry and melting points
41	B1	H	OCH ₃	phenyl	H	(B); 160°C
206	B7	Br	OCH ₃	1-naphthyl	F	(B); 213°C

Comp. nr.	Ex. nr.	R ¹	R ²	R ³	R ⁶	Stereochemistry and melting points
31	B1	Cl	OCH ₃	phenyl	H	(B); 181°C
26	B1	phenyl	OCH ₃	phenyl	H	(A); 174°C
27	B1	phenyl	OCH ₃	phenyl	H	(B); 192°C
32	B1	Br	SCH ₃	phenyl	H	(A); 208°C
33	B1	Br	SCH ₃	phenyl	H	(B); 196°C
109	B9	Br	OCH ₃	3-fluorophenyl	H	(A2); 156°C
39	B9	Br	OCH ₃	3-fluorophenyl	H	(B); 207°C
36	B1	Br	OCH ₃	phenyl	Cl	(A); 197°C
14	B7	Br	OCH ₃	1-naphthyl	H	(A); 210°C
15	B7	Br	OCH ₃	1-naphthyl	H	(B); 244°C
46	B7	Br	OCH ₃	2-naphthyl	H	(B); 162°C
4	B1	Br	OCH ₃	phenyl	H	(B); 190°C
174	B9		OCH ₃	3-fluorophenyl	H	(A); 159°C
24	B1	Br	OCH ₃	3-thienyl	H	(A); 162°C
5	B2	Br	OCH ₂ CH ₃	phenyl	H	(A); 162°C
38	B9	Br	OCH ₃	3-fluorophenyl	H	(A); 198°C
69	B1	Br	OCH ₃	2-fluorophenyl	H	(A); oil
70	B1	Br	OCH ₃	2-fluorophenyl	H	(B); oil
16	B1	Br	OCH ₃	4-chlorophenyl	H	(A); 200°C
17	B1	Br	OCH ₃	4-chlorophenyl	H	(B); 190°C
67	B8	Br	OCH ₃	2,5-difluorophenyl	H	(A); 60°C
116	B15	Br	OCH ₃	3,5-difluorophenyl	H	(B); 224°C
159	B8	Br	OCH ₃	2,5-difluorophenyl	H	(A1); 208°C
160	B8	Br	OCH ₃	2,5-difluorophenyl	H	(A2); 167°C
113	B14	Br	OCH ₃	2,3-difluorophenyl	H	(A); 128°C
196	B15	Br	OCH ₃	3,4-difluorophenyl	H	(B); 184°C
76	B1	Br	OCH ₃		H	(B); 130°C
84	B1	phenyl	OCH ₃	1-naphthyl	H	(A); 248°C
45	B7	Br	OCH ₃	2-naphthyl	H	(A); 262°C
77	B1	Br	OCH ₃		H	(A); 202°C

Comp. nr.	Ex. nr.	R ¹	R ²	R ³	R ⁶	Stereochemistry and melting points
78	B1	Br	OCH ₃		H	(B); 202°C
165	B9	Br		3-fluorophenyl	H	(B); oil
72	B1	Br	OCH ₃	1-naphthyl	CH ₃	(B); 178°C
212	B7		OCH ₃	1-naphthyl	H	(B); 220°C
163	B7	Br	OCH ₃	6-methoxy-2-naphthyl	H	(B); 206°C
73	B1	Br	OCH ₃	1-naphthyl	Cl	(B); 174°C
201	B1	Br	OCH ₃		H	(A); 214°C
30	B1	Cl	OCH ₃	phenyl	H	(A); 170°C
21	B1	Br	OCH ₃	2-thienyl	H	(B); 176°C
25	B1	Br	OCH ₃	3-thienyl	H	(B); 160°C
20	B1	Br	OCH ₃	2-thienyl	H	(A); 96°C
6	B2	Br	OCH ₂ CH ₃	phenyl	H	(B); 74°C
108	B9	Br	OCH ₃	3-fluorophenyl	H	(A1); 160°C
37	B1	Br	OCH ₃	phenyl	Cl	(B); 221°C
13	B7	Br	OCH ₃	1-naphthyl	H	(A2); (1S,2R)120°C
130	B7	Br	OCH ₃	5-bromo-2-naphthyl	H	(B); 220°C
G		3-pyridyl	OCH ₃	3,5-difluorophenyl	H	(A); 201°C
H		2-furanyl	OCH ₃	1-naphthyl	H	(B); 215°C
I		Br	OCH ₃		H	(A) ; >260°C
F	B1	Br	OCH ₃	phenyl	H	(B2); 175°C
E	B1	Br	OCH ₃	phenyl	H	(B1); 180°C
K	B1	Br	OCH ₃		H	(A); 116°C
L	B1	Br	OCH ₃		H	(B); 143°C
M	B1	Br	OCH ₃		H	(A)
N	B1	Br	OCH ₃		H	(A); 76°C

Comp. nr.	Ex. nr.	R ¹	R ²	R ³	R ⁶	Stereochemistry and melting points
O	B1	Br	OCH ₃	phenyl	F	(B); 162-163°C
P		CH ₂ OH	OCH ₃	1-naphthyl	H	(B); 196°C
Q		CH ₂ OH	OCH ₃	1-naphthyl	H	(A); 187°C


Table 2:



5

Comp. nr.	Ex. nr.	R ¹	R ³	R ⁴	R ⁵	R ^{6a}	R ^{6b}	Phys. data (salt/melting points) and stereochemistry
44	B4	Br	phenyl	H	H	H	H	(A); 190°C
19	B1	Br	phenyl	CH ₂ CH ₃	CH ₂ CH ₃	H	H	.ethanedioate (2:3), (B); 150°C
18	B1	Br	phenyl	CH ₂ CH ₃	CH ₂ CH ₃	H	H	.ethanedioate (2:3); (A); 230°C
R	B7	Br	1-naphthyl	CH ₃	H	H	H	(A2)
S	B13	Br	2-naphthyl	CH ₃	H	H	H	(B); 198°C
T	B13	Br	2-naphthyl	CH ₃	H	H	H	(A); 178°C
U	B1	Br	phenyl	CH ₂ CH ₃	CH ₃	H	H	(A)
V	B13	Br	phenyl	CH ₃	H	H	H	(B); 197°C
W	B13	CH ₃	phenyl	CH ₃	H	H	H	(B)
X	B1	OCH ₃	phenyl	CH ₂ CH ₃	CH ₂ CH ₃	H	H	(A); 140°C
Y	B1	Br	phenyl	CH ₂ CH ₃	CH ₃	Cl	H	(A)
Z	B13	H		CH ₃	H	H	H	(B)
A'	B1	Br	phenyl	CH ₂ CH ₃	CH ₂ CH ₃	H	Cl	(B); 180°C
B'	B1	Br	phenyl		CH ₃	H	Cl	(A); 204°C
C'	B1	Br	phenyl		CH ₃	H	Cl	(B); 183°C

Table 3



Comp nr.	Ex. nr.	R ¹	R ²	R ³	L	Phys.data (NMR/melting points) and stereochemistry
50	B1	Br	OCH ₃	phenyl	1-imidazolyl	(B); 230°C
137	B7	Br	OCH ₃	2-naphthyl	4-methylpyrazinyl	(B); 232°C
154	B7	Br	OCH ₃	6-bromo-2-naphthyl	4-methylpiperazinyl	(B); 254°C
49	B1	Br	OCH ₃	phenyl	1-imidazolyl	(A); 216°C
136	B7	Br	OCH ₃	2-naphthyl	4-methylpyrazinyl	(A); 188°C
48	B1	Br	OCH ₃	phenyl	1-piperidinyl	(B); 210°C
55	B1	Br	OCH ₃	phenyl	thiomorpholinyl	(A); oil
171	B7	Br	OCH ₃	2-naphthyl	1-pyrrolidinyl	(B); 218°C
129	B1	Br	OCH ₃	2-naphthyl	1-piperidinyl	(B); 212°C
149	B7	Br	OCH ₃	1-naphthyl	4-methylpyrazinyl	(B); 232°C
166	B7	Br	OCH ₃	6-bromo-2-naphthyl	1-piperidinyl	(B); 218°C
151	B7	Br	OCH ₃	3-bromo-1-naphthyl	4-methylpiperazinyl	(A); 178°C
53	B1	Br	OCH ₃	phenyl	1-(1,2,4-triazolyl)	(A); 180°C
47	B1	Br	OCH ₃	phenyl	1-piperidinyl	(A); 190°C
51	B1	Br	OCH ₃	phenyl	1-(4-methyl)piperazinyl	(A); 150°C
170	B7	Br	OCH ₃	2-naphthyl	1-pyrrolidinyl	(A); 238°C
128	B1	Br	OCH ₃	2-naphthyl	1-piperidinyl	(A); 254°C
145	B7	Br	OCH ₃	1-naphthyl	1-piperidinyl	(A); 212°C
139	B7	Br	OCH ₃	2-naphthyl	morpholinyl	(A); 258°C
140	B7	Br	OCH ₃	2-naphthyl	morpholinyl	(B); 214°C
155	B7	Br	OCH ₃	6-bromo-2-naphthyl	1-piperidinyl	(A); 224°C
A	B13	Br	OCH ₃	2,4-difluorophenyl	NH(CH ₂ CH ₃)	(B); 171°C
B	B13	Br	OCH ₃	2,4-difluorophenyl	NH(CH ₂ CH ₃)	(A)
C	B12	Br	OCH ₃	2,4-difluorophenyl	N(CH ₂ CH ₃)(CH ₂ phenyl)	(B)






Comp. nr.	Ex. nr.	R ¹	R ²	R ³	L	Phys.data (NMR/melting points) and stereochemistry
D	B13	Br	OCH ₃	phenyl	NH(CH ₃)	(B); 197°C
209	B7	Br	OCH ₃	2-naphthyl		(B); 198°C
58	B6	Br	OCH ₃	phenyl		(A); 208°C
11	B6	Br	OCH ₃	phenyl		(B); 208°C
D'	B1	Br	OCH ₃	phenyl		(A)
E'	B1	H	SCH ₃	phenyl	1-imidazolyl	(B1)
F'	B1	H	SCH ₃	phenyl	1-imidazolyl	(B2)


Table 4:



Comp. nr.	Ex. nr.	R ^{1a}	R ^{1b}	R ³	R ⁶	stereochemistry and melting points
215	B9	Br	CH ₃	3-fluorophenyl	H	(A); 197°C
226	B7	Br	CH ₃	1-naphthyl	H	(B); 208°C
229	B7	CH ₃	CH ₃	1-naphthyl	H	(A); 238°C
227	B15	Br	CH ₃	3,5-difluorophenyl	H	(A); 195°C

Comp nr.	Ex. nr.	R ^{1a}	R ^{1b}	R ³	R ⁶	stereochemistry and melting points
223	B7	Br	H	1-naphthyl		(B); 205°C

Table 5



Comp nr.	Ex. nr.	R ^{6a}	R ^{6b}	R ^{6c}	stereochemistry and melting points
G'	B1	H	F	OCH ₃	(A); 154-155°C
H'	B1	H	Cl	CH ₃	(A); 120-122°C
I'	B1	H	Cl	Cl	(B); 169-170°C
J'	B1	OCH ₃	OCH ₃	H	(A); 106-108°C
K'	B1	Cl	H	Cl	(A); 124-125°C
L'	B1	H	F	OCH ₃	(B); 191-192°C
M'	B1	H	CH ₃	H	(B); 146-147°C
N'	B1	H	F	H	(B); 159-160°C
O'	B1	H	F	H	(A); 147-148°C
P'	B1	H	Cl	H	(A); 102-104°C
Q'	B1	H	Cl	H	(B); 137-139°C
R'	B1	H	CH ₃	H	(A); 150-151°C

Analytical methods

The mass of some compounds was recorded with LCMS (liquid chromatography mass spectrometry). The methods used are described below. The data are gathered in Table 6 below.

General procedure 1:

The HPLC gradient was supplied by an Alliance HT 2795 (Waters) system consisting of a quaternary pump with degasser, an autosampler, and DAD detector. Flow from the column was split to the MS detector. MS detectors were configured with an electrospray ionization source. The capillary needle voltage was 3 kV and the source temperature was maintained at 100 °C. Nitrogen was used as the nebulizer gas. Data acquisition was performed with a Waters-Micromass MassLynx-Openlynx data system.

General procedure 2:

The HPLC gradient was supplied by an Alliance HT 2790 (Waters) system consisting of a quaternary pump with degasser, an autosampler, a column oven (set at 40 °C) and DAD detector. Flow from the column was split to the MS detector. MS detectors were configured with an electrospray ionization source. Mass spectra were acquired by scanning from 100 to 1000 in 1 second using a dwell time of 0.1 second. The capillary needle voltage was 3 kV and the source temperature was maintained at 140 °C. Nitrogen was used as the nebulizer gas. Data acquisition was performed with a Waters-Micromass MassLynx-Openlynx data system.

LCMS-method 1

In addition to general procedure 1: LCMS analysis was carried out (electrospray ionization in positive mode, scanning mode from 100 to 900 amu) on a Kromasil C18 column (Interchim, Montluçon, FR; 5 µm, 4.6 x 150 mm) with a flow rate of 1 ml/minute. Two mobile phases (mobile phase A: 30 % 6.5mM ammonium acetate + 40 % acetonitrile + 30 % formic acid (2ml/l); mobile phase B: 100 % acetonitrile) were employed to run a gradient condition from 100 % A for 1 minute to 100 % B in 4 minutes, 100 % B for 5 minutes to 100 % A in 3 minutes, and reequilibrate with 100 % A for 2 minutes.

LCMS-method 2

In addition to general procedure 1: Reversed phase HPLC was carried out on a Sunfire C18 column (3.5 μ m, 4.6 x 100 mm) with a flow rate of 0.8 ml/min. Three mobile phases (mobile phase A: 100 % 7 mM ammonium acetate ; mobile phase B: 100 % acetonitrile; mobile phase C: 0.2 % formic acid + 99.8 % ultra-pure water) were employed to run a gradient condition from 25 % A , 50% B and 25% C (hold for 1 minute) to 100 % B in 4 minutes, 100% B for 4 minutes and reequilibrate with initial conditions for 2 minutes. An injection volume of 10 μ l was used. Cone voltage was 20 V for positive and negative ionization mode. Mass spectra were acquired by scanning from 100 to 1000 in 0.4 seconds using an interscan delay of 0.3 seconds.

LCMS-method 3

In addition to general procedure 2: Reversed phase HPLC was carried out on an Xterra MS C18 column (3.5 mm, 4.6 x 100 mm) with a flow rate of 1.6 ml/min. Two mobile phases (mobile phase A: 70 % methanol + 30 % H₂O; mobile phase B: 0.1 % formic acid in H₂O/methanol 95/5) were employed to run a gradient condition from 100 % B to 5 % B + 95 % A in 12 minutes. An injection volume of 10 μ l was used. Cone voltage was 10 V for positive ionization mode and 20 V for negative ionization mode.

Table 6 : LCMS parent peak

Compound No	LCMS method	LCMS parent peak (MH ⁺)	R _t (min)
A	1	541	5.06
B	1	541	5.76
C	1	631	11.17
M	1	471	5.6
R	1	541	5.87
U	2	521	2.87
W	1	427	5.06
Y	2	555	3.98
Z	1	403	4.17
E'	3	234	8.93
F'	3	234	8.94

Pharmacological examples

Preparation of bacterial suspensions for susceptibility testing:

The bacteria used in this study were grown overnight in flasks containing 100 ml Mueller-Hinton Broth (Becton Dickinson - cat. no. 275730) in sterile de-ionized water, with shaking, at 37 °C. Stocks (0.5 ml/tube) were stored at -70 °C until use. Bacteria titrations were performed in microtiter plates to detect the TCID₅₀, in which the TCID₅₀ represents the dilution that gives rise to bacterial growth in 50 % of inoculated cultures.

In general, an inoculum level of approximately 100 TCID₅₀ was used for susceptibility testing.

Anti bacterial Susceptibility testing: IC₉₀ determination

Microtitre plate assay

Flat-bottom, sterile 96-well plastic microtiter plates were filled with 180 µl of sterile deionized water, supplemented with 0.25 % BSA. Subsequently, stock solutions (7.8 x final test concentration) of compounds were added in 45 µl volumes in column 2. Serial five-fold dilutions (45 µl in 180 µl) were made directly in the microtiter plates from column 2 to reach column 11. Untreated control samples with (column 1) and without (column 12) inoculum were included in each microtiter plate. Depending on the bacteria type, approximately 10 to 60 CFU per well of bacteria inoculum (100 TCID₅₀), in a volume of 100 µl in 2.8x Mueller-Hinton broth medium, was added to the rows A to H, except column 12. The same volume of broth medium without inoculum was added to column 12 in row A to H. The cultures were incubated at 37°C for 24 hours under a normal atmosphere (incubator with open air valve and continuous ventilation). At the end of incubation, one day after inoculation, the bacterial growth was quantitated fluorometrically. Therefore resazurin (0.6 mg/ml) was added in a volume of 20 µl to all wells 3 hours after inoculation, and the plates were re-incubated overnight. A change in colour from blue to pink indicated the growth of bacteria. The fluorescence was read in a computer-controlled fluorometer (Cytofluor Biosearch) at an excitation wavelength of 530 nm and an emission wavelength of 590 nm. The % growth inhibition achieved by the compounds was calculated according to standard methods. The IC₉₀ (expressed in µg/ml) was defined as the 90 % inhibitory concentration for bacterial growth. The results are shown in Table 6.

Agar dilution method.

MIC₉₉ values (the minimal concentration for obtaining 99 % inhibition of bacterial growth) can be determined by performing the standard Agar dilution method according to NCCLS standards* wherein the media used includes Mueller-Hinton agar.

- 5 * Clinical laboratory standard institute. 2005. Methods for dilution Antimicrobial susceptibility tests for bacteria that grows Aerobically: approved standard -sixth edition

Time kill assays

- Bactericidal or bacteriostatic activity of the compounds may be determined in a time
10 kill assay using the broth microdilution method *. In a time kill assay on
Staphylococcus aureus and methicillin resistant *S. aureus* (MRSA), the starting
inoculum of *S. aureus* and MRSA is 10⁶ CFU / ml in Muller Hinton broth. The
antibacterial compounds are used at the concentration of 0.1 to 10 times the MIC (i.e.
IC₉₀ as determined in microtitre plate assay). Wells receiving no antibacterial agent
15 constitute the culture growth control. The plates containing the microorganism and the
test compounds are incubated at 37 °C. After 0, 4, 24, and 48 hrs of incubation samples
are removed for determination of viable counts by serial dilution (10⁻¹ to 10⁻⁶) in sterile
PBS and plating (200 µl) on Mueller Hinton agar. The plates are incubated at 37 °C for
24 hrs and the number of colonies are determined. Killing curves can be constructed by
20 plotting the log₁₀CFU per ml versus time. A bactericidal effect is commonly defined as
3-log₁₀ decrease in number of CFU per ml as compared to untreated inoculum. The
potential carryover effect of the drugs is removed by serial dilutions and counting the
colonies at highest dilution used for plating.

- 25 * Zurenko,G.E. *et al.* In vitro activities of U-100592 and U-100766, novel oxazolidinone antibacterial
agents. *Antimicrob. Agents Chemother.* **40**, 839-845 (1996).

Determination of cellular ATP levels

- In order to analyse the change in the total cellular ATP concentration (using ATP
bioluminescence Kit, Roche), assays are carried out by growing a culture of *S. aureus*
30 (ATCC29213) stock in 100 ml Mueller Hinton flasks and incubate in a shaker-
incubator for 24 hrs at 37 °C (300 rpm). Measure OD₄₀₅ nm and calculate the CFU/ml.
Dilute the cultures to 1 x 10⁶ CFU/ml (final concentration for ATP measurement: 1 x
10⁵ CFU/100 µl per well) and add test compound at 0.1 to 10 times the MIC (i.e. IC₉₀
as determined in microtitre plate assay). Incubate these tubes for 0, 30 and 60 minutes
35 at 300 rpm and 37°C. Use 0.6 ml bacterial suspension from the snap-cap tubes and add
to a new 2 ml eppendorf tubes. Add 0.6 ml cell lysis reagent (Roche kit), vortex at
max speed and incubate for 5 minutes at room temperature. Cool on ice. Let the

luminometer warm up to 30°C (Luminoskan Ascent Labsystems with injector). Fill one column (= 6 wells) with 100 µl of the same sample. Add 100 µl Luciferase reagent to each well by using the injector system. Measure the luminescence for 1 sec.

5 Table 7 : IC₉₀ values (µg/ml) determined according to the Microtitre plate assay.

Comp. No.	IC90 (µg/ml)												
	BSU 43639	ECO 25922	EFA 14506	EFA 29212	LMO 49594	PAE 27853	SMU 33402	SPN 6305	SPY 8668	STA 43300	STA 25923	STA 29213	STA RMETH
A							10.8	1.1	10.8		10.8	12.1	
50	10.5			10.5	23.6		21.0		26.5			26.5	10.5
41					17.0								
F							22.6	25.3	22.6				
206								12.8					
31	10.3			4.6	4.1	10.3	10.3	5.2	1.8	11.6			
44	7.6										7.6		
26	10.0			4.5	4.5	2.0	11.3	5.6	2.5	11.3	8.9		
27	10.0				8.9			12.6					
E	12.7			11.3	7.1	10.1	10.1	5.7	5.1	12.7			
32		11.7		4.6	4.6	10.4	4.6	13.1	4.6	13.1	9.3		13.1
33	10.4		11.7	4.6	2.3	11.7	10.4	5.9	2.6	5.9	13.1		
109				11.7	9.3	11.7		13.1	13.1	13.1	10.4		
39				10.4	4.7	10.4	11.7	5.9	1.9	13.1	10.4		
137						21.2		23.8			8.4		21.2
36	24.1			4.8	5.4	4.8	3.8	2.7	0.4	4.8			
14	11.1		9.9	5.0	2.5	5.0	12.4	6.2	2.8	5.6	9.9		14.0
15	24.8		24.8	5.0	2.5	12.4	12.4	6.2	2.2	5.0			
46				5.0	2.8	5.0	5.6	2.8	0.6	5.6			
154	10.9			27.4	27.4				27.4		13.8		24.5
4								12.7					
174					10.2								
175					10.2			12.8					
24								10.2					
5							10.4						
38					10.4								
69					10.4	10.4		13.1					

Comp. No.	IC90 (µg/ml)												
	BSU	ECO	EFA	EFA	LMO	PAE	SMU	SPN	SPY	STA	STA	STA	STA
	43639	25922	14506	29212	49594	27853	33402	6305	8668	43300	25923	29213	RMETH
70					10.4			13.1					
49					10.5		10.5	13.3	10.5		5.9		11.8
136	10.6			10.6	10.6								
19								13.4					
215					10.7								
G	13.6			10.8	10.8								
16						13.6							
17					10.8			0.5					
67					10.8			13.6					
116					10.8								
110					10.8			13.6					
111					10.8								
113					10.8			13.6					
196							10.8						
B											8.6		
H					13.6			13.6					
48					10.9	10.9	10.9	13.7	13.7				
76					11.0	11.0							
84					11.0		11.0	7.0	11.0				
45	11.1		12.4										
77						11.2		14.2					
78							11.2	11.2	11.2				
55								0.6					
165							14.3	14.3	11.3				
72							9.0	14.3	11.4				
226								14.3					
212	11.4												
171								5.18	11.6				
163								14.7					
73							11.8		14.8				
129							11.9	15.0	11.9				
149					12.2								
201	12.3			12.3	12.3	12.3							


IC90 (µg/ml)													
Comp.	BSU	ECO	EFA	EFA	LMO	PAE	SMU	SPN	SPY	STA	STA	STA	STA
No.	43639	25922	14506	29212	49594	27853	33402	6305	8668	43300	25923	29213	RMETH
C								15.9					
18				13.3	13.3			0.7					
166								16.9					
151			15.4										15.4
30				10.3	10.3	9.2	10.3	5.2	10.3				
D								4.9					
229							22.5	5.0					
21				11.5	4.6		11.5	12.8		12.8			
25				11.5	11.5		11.5	12.8	4.1	12.8			
20								2.6					
6				11.6	11.6	11.6	4.6	13.0	4.6				
108								13.1					
53								5.9					
37								12.1					
47								6.9					
227								5.0					
13	11.1		11.1	8.8	11.1	2.2	3.5	2.8	8.8	11.1	11.1		
51	28.1			25.0	25.0			5.6					
I								11.6					
170								26.0					
128					9.4								
145			23.7					9.4					
139							18.9	26.7					
140							9.5						
223							5.4	13.6	5.4				
130							10.1						
155			26.9										
209								15.1					
58							5.9		8.3				
11					11.7		10.4		11.7				
K								1.77				7.92	
L								8.88					
M								7.47					

Comp. No.	IC90 (µg/ml)												
	BSU	ECO	EFA	EFA	LMO	PAE	SMU	SPN	SPY	STA	STA	STA	STA
	43639	25922	14506	29212	49594	27853	33402	6305	8668	43300	25923	29213	RMETH
N								7.69				7.69	
O								9.31					
P								8.03					
Q								10.11				8.03	
R								0.43				1.71	
S								0.68				1.71	
T								1.92				8.58	
U								2.07					
V								4.91					
W								6.76				16.98	
X								8.62					
Y								8.78					
Z								9.01					
A'								10.1					
B'								10.1					
C'								10.1					
D'								2.34				9.33	
E'								7.38				7.38	
F'								8.28				16.52	
G'								1.75				15.6	
H'								1.75					
I'								1.82					
J'								2.01				8.96	
K'								4.07					
L'								4.4				19.64	
M'								9.24					
N'								2.08					
O'								2.08				26.23	
P'								1.92					
Q'								1.92					
R'								2.07					

- BSU 43639 means *Bacillus subtilis* (ATCC43639); ECO 25922 means *Escherichia coli* (ATCC25922); EFA 14506 means *Enterococcus faecalis* (ATCC14506); EFA 29212 means *Enterococcus faecalis* (ATCC29212); LMO 49594 means *Listeria*
- 5 *monocytogenes* (ATCC49594); PAE 27853 means *Pseudomonas aeruginosa* (ATCC27853); SMU 33402 means *Streptococcus mutans* (ATCC33402); SPN 6305 means *Streptococcus pneumoniae* (ATCC6305); SPY 8668 means *Streptococcus pyogenes* (ATCC8668); STA 43300 means *Staphylococcus aureus* (ATCC43300); STA 25923 means *Staphylococcus aureus* (ATCC25923); STA 29213 means
- 10 *Staphylococcus aureus* (ATCC29213); STA RMETH means methicilline resistant *Staphylococcus aureus* (MRSA) (a clinical isolate from the University of Antwerp). ATCC means American type tissue culture.

Claims

1. Use of a compound for the manufacture of a medicament for the treatment of a bacterial infection caused by *Staphylococci*, *Enterococci* or *Streptococci*, said compound being a compound of formula



a pharmaceutically acceptable acid or base addition salt thereof, a stereochemically isomeric form thereof or a N-oxide form thereof, wherein

R¹ is hydrogen, halo, polyhaloC₁₋₆alkyl, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, Ar or Het;

p is an Integer equal to 1 or 2;

R² is C₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyloxy or C₁₋₆alkylthio;

R³ is C₁₋₆alkyl, Ar, Het or Het¹;

R⁴ and R⁵ each independently are hydrogen, C₁₋₆alkyl or benzyl; or

R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from the group of pyrrolidinyl, 2-pyrrolinyl, 3-pyrrolinyl, pyrrolyl, imidazolidinyl, pyrazolidinyl, 2-imidazolinyl, 2-pyrazolinyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkylthioC₁₋₆alkyl or pyrimidinyl;

R⁶ is hydrogen, halo, polyhaloC₁₋₆alkyl, C₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio; or

two vicinal R⁶ radicals may be taken together to form a bivalent radical of formula -CH=CH-CH=CH- ;

r is an integer equal to 1 or 2;

R⁷ is hydrogen, C₁₋₆alkyl, Ar, Het or Het¹;

Ar is a homocycle selected from the group of phenyl, naphthyl, acenaphthyl, 1,2-dihydro-acenaphthyl, tetrahydronaphthyl, each homocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of hydroxy, halo, cyano, nitro, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, polyhaloC₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl, carboxyl, C₁₋₆alkyloxycarbonyl, aminocarbonyl, morpholinyl and mono- or di(C₁₋₆alkyl)aminocarbonyl;

Het is a monocyclic heterocycle selected from the group of N-phenoxypiperidinyl, piperidinyl, pyrrolyl, pyrazolyl, imidazolyl, furanyl, thienyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyridinyl, pyrimidinyl, pyrazinyl and pyridazinyl; each monocyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, hydroxy, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl or Ar-C(=O)-;


Het¹ is a bicyclic heterocycle selected from the group of quinolinyl, quinoxalinyl, indolyl, benzimidazolyl, ben-

zoxazolyl, benzisoxazolyl, benzothiazolyl, benzisothiazolyl, benzofuranyl, benzothienyl, 2,3-dihydrobenzo[1,4]dioxinyl or benzo[1,3]dioxolyl; each bicyclic heterocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of halo, hydroxy, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl or Ar-C(=O)-;

and


provided that the compound is other than (α S, β R)-6-bromo- α -[2-(dimethylamino) ethyl]-2-methoxy- α -1-naphthalenyl- β -phenyl-3-quinolineethanol, a pharmaceutically acceptable acid or base addition salt thereof or a N-oxide form thereof.

2. Use according to claim 1 wherein R¹ is hydrogen, halo, polyhaloC₁₋₆alkyl, C₁₋₆alkyl, Ar or Het; R³ is Ar, Het or Het¹, and Ar is a homocycle selected from the group of phenyl, naphthyl, acenaphthyl, tetrahydronaphthyl, each homocycle may optionally be substituted with 1, 2 or 3 substituents, each substituent independently selected from the group of hydroxy, halo, cyano, nitro, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkyl, polyhaloC₁₋₆alkyl, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, polyhaloC₁₋₆alkyloxy, C₁₋₆alkyloxyC₁₋₆alkyl, carboxyl, C₁₋₆alkyloxycarbonyl, aminocarbonyl, morpholinyl and mono- or di(C₁₋₆alkyl)aminocarbonyl.
3. Use according to claim 1 or 2 wherein the compound of formula (I) is a compound having the following formula



a pharmaceutically acceptable acid or base addition salt thereof, a stereochemically isomeric form thereof or a N-oxide form thereof.

4. Use according to any one of the preceding claims wherein the compound of formula (I) or (I-a) is a compound having the following formula




a pharmaceutically acceptable acid or base addition salt thereof, a stereochemically isomeric form thereof or a N-oxide form thereof.

5. Use according to any one of the preceding claims wherein R¹ is hydrogen; C₁₋₆alkyl; halo; hydroxyC₁₋₆alkyl; C₁₋₆alkyloxy; optionally substituted phenyl; or Het.
6. Use according to claim 5 wherein R¹ is hydrogen, halo, optionally substituted phenyl, or Het.
7. Use according to claim 6 wherein R¹ is hydrogen, halo or optionally substituted phenyl.

8. Use according to claim 7 wherein R¹ is halo.
9. Use according to any one of the preceding claims wherein R² is C₁₋₆alkyloxy or C₁₋₆alkylthio.
10. Use according to any one of the preceding claims wherein R³ is C₁₋₆alkyl, Ar or Het.
11. Use according to claim 10 wherein R³ is Ar or Het.
12. Use according to any one of claims 1 to 9 wherein R³ is Ar or Het¹.
13. Use according to claim 10, 11 or 12 wherein R³ is Ar.
14. Use according to claim 13 wherein R³ is optionally substituted phenyl or optionally substituted naphthyl.
15. Use according to any one of the preceding claims wherein R⁴ and R⁵ each independently are hydrogen, C₁₋₆alkyl or benzyl.
16. Use according to claim 15 wherein R⁴ and R⁵ each independently are hydrogen or C₁₋₆alkyl.
17. Use according to any one of claims 1 to 14 wherein R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from the group of pyrrolidinyl, 2-pyrrolinyl, 3-pyrrolinyl, pyrrolyl, imidazolidinyl, pyrazolidinyl, 2-imidazolyl, 2-pyrazolyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkylthioC₁₋₆alkyl or pyrimidinyl.
18. Use according to claim 17 wherein R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from the group of pyrrolidinyl, pyrrolyl, imidazolyl, pyrazolyl, triazolyl, piperidinyl, pyridinyl, piperazinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, morpholinyl and thiomorpholinyl, each of said rings may optionally be substituted with C₁₋₆alkyl, halo, polyhaloC₁₋₆alkyl, hydroxy, hydroxyC₁₋₆alkyl, C₁₋₆alkyloxy, amino, mono- or di(C₁₋₆alkyl)amino, C₁₋₆alkylthio, C₁₋₆alkyloxyC₁₋₆alkyl, C₁₋₆alkylthioC₁₋₆alkyl or pyrimidinyl.
19. Use according to any one of the preceding claims wherein R⁶ is hydrogen, halo, polyhaloC₁₋₆alkyl, or C₁₋₆alkyl; or two vicinal R⁶ radicals may be taken together to form a bivalent radical of formula -CH=CH-CH=CH-.
20. Use according to any one of claims 1 to 18 wherein R⁶ is hydrogen, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, or two vicinal R⁶ radicals may be taken together to form a bivalent radical of formula -CH=CH-CH=CH-.
21. Use according to claim 19 or 20 wherein R⁶ is hydrogen or halo.
22. Use according to any one of the preceding claims wherein R⁷ is hydrogen.
23. Use according to any one of the preceding claims wherein r is an integer equal to 1.
24. Use according to any one of claims 1, 2, 5 to 23 wherein p is an integer equal to 1.
25. Use according to claim 3 provided that when one R¹ is C₁₋₆alkyl then the other R¹ substituent is selected from halo, polyhaloC₁₋₆alkyl, C₁₋₆alkyl, Ar or Het.
26. Use according to any one of the preceding claims wherein the bacterial infection is an infection with a gram-positive bacterium.
27. Use according to claim 1 wherein R¹ is hydrogen; C₁₋₆alkyl; halo; hydroxyC₁₋₆alkyl; C₁₋₆alkyloxy; phenyl; furanyl optionally substituted with hydroxyC₁₋₆alkyl; or pyridyl; R² is C₁₋₆alkyloxy; C₁₋₆alkylthio; or C₁₋₆alkyloxyC₁₋₆alkyloxy; R³ is phenyl optionally substituted with 1 or 2 halo; naphthyl optionally substituted with 1 or 2 halo or C₁₋₆alkyloxy; thienyl; piperidinyl substituted with Ar-C(=O); 2,3-dihydrobenzo[1,4]dioxinyl; benzo[1,3]dioxolyl; acenaphthyl; furanyl; or C₁₋₆alkyl; R⁴ and R⁵ are each independently hydrogen; C₁₋₆alkyl; benzyl; or R⁴ and R⁵ together and including the N to which they are attached may form a radical selected from imidazolyl; pyrazinyl substituted with C₁₋₆alkyl;

piperazinyl substituted with C₁₋₆alkyl; piperazinyl substituted with pyrimidinyl; piperidinyl; piperidinyl substituted with dimethylamino; thiomorpholinyl; morpholinyl; pyrrolidinyl; or triazolyl; R⁶ is hydrogen; halo, C₁₋₆alkyl, C₁₋₆alkyloxy or two vicinal R⁶ radicals may be taken together to form a bivalent radical of formula -CH=CH-CH=CH-; R⁷ is hydrogen

28. Use according to claim 1 provided that when R³ is Ar, Het or Het¹; then R¹ is hydroxyC₁₋₆alkyl or C₁₋₆alkyloxy.
29. Use according to claim 28 wherein R¹ is halo, hydroxyC₁₋₆alkyl, or C₁₋₆alkyloxy, R² is C₁₋₆alkyloxy; R³ is C₁₋₆alkyl, phenyl or naphthyl; R⁴ and R⁵ are C₁₋₆alkyl; R⁶ is hydrogen, R⁷ is hydrogen, p is 1; r is 1
30. Use according to claim 1 wherein the compound of formula (I) is selected from the following compounds



R ¹	R ²	R ³	R ⁴	R ⁵	R ⁶	Stereochemistry
Cl	OCH ₃	phenyl	CH ₃	CH ₃	H	(B)
phenyl	OCH ₃	phenyl	CH ₃	CH ₃	H	(A)
Br	SCH ₃	phenyl	CH ₃	CH ₃	H	(B)
Br	OCH ₃	3-fluorophenyl	CH ₃	CH ₃	H	(B)
Br	OCH ₃	phenyl	CH ₃	CH ₃	Cl	(A)
Br	OCH ₃	2-naphthyl	CH ₃	CH ₃	H	(B)
Br	OCH ₃	1-naphthyl	CH ₃	CH ₃	F	(B)
Br	OCH ₃	phenyl	H	H	H	(A)
Br	OCH ₃	2,4-difluorophenyl	H	CH ₂ CH ₃	H	(B)
Br	OCH ₃	1-naphthyl	CH ₃	CH ₃	H	(A2) ; (1S,2R)
Br	OCH ₃	phenyl	CH ₃	CH ₃	H	(B1) ; (1S,2S)
Br	OCH ₃	phenyl	CH ₃	CH ₃	H	(B2) ; (1R,2R)
Br	OCH ₃	1-naphthyl	CH ₃	H	H	(A2)
Br	OCH ₃	2-naphthyl	CH ₃	H	H	(B)

a pharmaceutically acceptable acid or base addition salt thereof or a N-oxide form thereof, and wherein:

(A) or (B) means that the compound is a mixture of two enantiomers and (A) is isolated first and (B) second, wherein the isolation is by chromatography over silica gel using eluent CH₂Cl₂/CH₃OH/NH₄OH, CH₂Cl₂/CH₃OH or CH₂Cl₂/iPrOH/NH₄OH;


(A1), (A2), (B1) and (B2) means that the compound is an enantiomer;



(A1) and (A2) respectively represent the first and second fractions isolated from (A); and

(B1) and (B2) respectively represent the first and second fractions isolated from (B),


and wherein the isolation of the enantiomers is by chiral column (chiracel OD, chiracel AD) chromatography using eluent hexane/ethanol, heptane/iPrOH or hexane/2-propanol.

31. A combination of (a) a compound of formula (I) as defined in any one of claims 1 to 30, and (b) one or more other antibacterial agents provided that the one or more other antibacterial agents are other than antimycobacterial agents, wherein the combination is for use in the treatment of a bacterial infection caused by *Staphylococci*, *Enterococci* or *Streptococci*.
32. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and, as active ingredient, a therapeutically effective amount of (a) a compound of formula (I) as defined in any one of claims 1 to 30, and (b) one or more other antibacterial agents provided that the one or more other antibacterial agents are other than antimycobacterial agents, wherein the composition is for use in the treatment of a bacterial infection caused by *Staphylococci*, *Enterococci* or *Streptococci*.
33. A compound as defined in any one of claims 1 to 30, for use in the treatment of a bacterial infection caused by *Staphylococci*, *Enterococci* or *Streptococci*.
34. A product containing (a) a compound of formula (I) as defined in any one of claims 1 to 30, and (b) one or more other antibacterial agents provided that the one or more other antibacterial agents are other than antimycobacterial agents, as a combined preparation for simultaneous, separate or sequential use in the treatment of a bacterial infection caused by *Staphylococci*, *Enterococci* or *Streptococci*.
35. A compound of formula (I) wherein the compound is selected from the following compounds



R ¹	R ²	R ³	R ⁴	R ⁵	R ^{6a}	R ^{6b}	R ^{6c}	Stereo-chemistry
3-pyridyl	OCH ₃	3,5-difluorophenyl	CH ₃	CH ₃	H	H	H	(A)
2-furanyl	OCH ₃	1-naphthyl	CH ₃	CH ₃	H	H	H	(B)
Br	OCH ₃		CH ₃	CH ₃	H	H	H	(A)
Br	OCH ₃		CH ₃	CH ₃	H	H	H	(B)
Br	OCH ₃		CH ₃	CH ₃	H	H	H	(A)
Br	OCH ₃		CH ₃	CH ₃	H	H	H	(A)
Br	OCH ₃	phenyl	CH ₃	CH ₃	H	H	H	(B)

(continued)

R ¹	R ²	R ³	R ⁴	R ⁵	R ^{6a}	R ^{6b}	R ^{6c}	Stereo-chemistry
-CH ₂ OH	OCH ₃	1-naphthyl	CH ₃	CH ₃	H	H	H	(B)
-CH ₂ OH	OCH ₃	1-naphthyl	CH ₃	CH ₃	H	H	H	(A)
Br	OCH ₃	2-naphthyl	CH ₃	H	H	H	H	(B)
Br	OCH ₃	2-naphthyl	CH ₃	H	H	H	H	(A)
Br	OCH ₃	phenyl	CH ₂ CH ₃	CH ₃	H	H	H	(A)
OCH ₃	OCH ₃	phenyl	CH ₂ CH ₃	CH ₂ CH ₃	H	H	H	(A)
Br	OCH ₃	phenyl	CH ₂ CH ₃	CH ₃	H	Cl	H	(A)
Br	OCH ₃	phenyl	CH ₂ CH ₃	CH ₂ CH ₃	H	H	Cl	(B)
Br	OCH ₃	phenyl		CH ₃	H	H	Cl	(A)
Br	OCH ₃	phenyl		CH ₃	H	H	Cl	(B)
Br	OCH ₃	phenyl	CH ₃	CH ₃	H	F	OCH ₃	(A)
Br	OCH ₃	phenyl	CH ₃	CH ₃	H	Cl	CH ₃	(A)
Br	OCH ₃	phenyl	CH ₃	CH ₃	H	Cl	Cl	(B)
Br	OCH ₃	phenyl	CH ₃	CH ₃	OCH ₃	OCH ₃	H	(A)
Br	OCH ₃	phenyl	CH ₃	CH ₃	Cl	H	Cl	(A)
Br	OCH ₃	phenyl	CH ₃	CH ₃	H	F	OCH ₃	(B)
Br	OCH ₂	phenyl	CH ₃	CH ₃	H	CH ₃	H	(B)
Br	OCH ₃	phenyl	CH ₃	CH ₃	H	F	H	(B)
Br	OCH ₃	phenyl	CH ₃	CH ₃	H	F	H	(A)
Br	OCH ₃	phenyl	CH ₃	CH ₃	H	Cl	H	(A)
Br	OCH ₃	phenyl	CH ₃	CH ₃	H	Cl	H	(B)
Br	OCH ₃	phenyl	CH ₃	CH ₃	H	CH ₃	H	(A)
Br	OCH ₃	phenyl	$NR^4R^5 =$ 		H	H	H	(A)
H	SCH ₃	phenyl	NR ⁴ R ⁵ = 1-imidazolyl		H	H	H	(B1)
H	SCH ₃	phenyl	NR ⁴ R ⁵ = 1-imidazolyl		H	H	H	(B2)

a pharmaceutically acceptable acid or base addition salt thereof or a *N*-oxide form thereof, and wherein:

(A) or (B) means that the compound is a mixture of two enantiomers and (A) is isolated first and (B) second, wherein the isolation is by chromatography over silica gel using eluent CH₂Cl₂/CH₃OH/NH₄OH, CH₂Cl₂/CH₃OH or CH₂Cl₂/*i*PrOH/NH₄OH;

(A1), (A2), (B1) and (B2) means that the compound is an enantiomer,

(A1) and (A2) respectively represent the first and second fractions isolated from (A), and

(B1) and (B2) respectively represent the first and second fractions isolated from (B),

and wherein the isolation of the enantiomers is by chiral column (chiracel OD, chiracel AD) chromatography using eluent hexane/ethanol, heptane/iPrOH or hexane/2-propanol

36. Use as claimed in any of claims 1 to 30, combination as claimed in claim 31, composition as claimed in claim 32, compound as claimed in claim 33 or product as claimed in claim 34, wherein the bacterial infection is an infection with methicillin resistant *Staphylococcus aureus* (MRSA), methicillin resistant coagulase negative staphylococci (MRCNS), penicillin resistant *Streptococcus pneumoniae* or multiple resistant *Enterococcus faecium*.
37. Use, combination, composition, compound or product as claimed in claim 36, wherein the bacterial infection is an infection with *Staphylococcus aureus* or *Streptococcus pneumoniae*.
38. Use, combination, composition, compound or product as claimed in claim 37, wherein the bacterial infection is an infection with *Staphylococcus aureus* (MRSA).