What is claimed is:

1. The use of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol in the manufacture of a medicament for the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification.

2. The use of Claim 1, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is in the form of an oral dosage form.

3. The use of Claims 1 or 2, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is in the form of a tablet.

4. The use of any one of Claims 1 to 3, wherein the use is in combination with another therapeutic agent.

5. The use of Claim 4, wherein the therapeutic agent is selected from trastuzumab, capecitabine, bevacizumab, paclitaxel and docetaxel.

6. The use of any one of Claims 1 to 5, wherein the use is after previous treatment for brain cancer.

7. The use of Claim 6, wherein the previous treatment for brain cancer is selected from surgery, radiation therapy and chemotherapy or mixtures thereof.

8. The use of Claims 6, wherein the previous treatment for brain cancer is selected from surgery, conventional external radiation therapy, three-dimensional conformal radiation therapy, intensity modulated radiation therapy, stereotactic radiosurgery, fractionated stereotactic radiation therapy, proton radiation therapy, internal or implant radiation therapy, temozolomide, bevacizumab, carmustine, lomustine, procarbazine, vincristine, tumor treating fields therapy, everolimus, procarbazine, lomustine, cisplatin, carboplatin and methotrexate or mixtures thereof.

9. The use of any one of Claims 1 to 5, wherein the use is after previous treatment for breast cancer.

10. The use of Claim 9, wherein the previous treatment for breast cancer is selected from surgery, sentinel lymph node biopsy followed by surgery, radiation therapy, chemotherapy, hormone therapy and targeted therapy.

11. The use of Claim 9, wherein the previous treatment for breast cancer is selected from lumpectomy, partial mastectomy, segmental mastectomy, total mastectomy, modified radical mastectomy, external radiation, internal radiation, methotrexate, paclitaxel albumin-stabilized nanoparticle formulation, ado-trastuzumab emtansine, doxorubicin, fluorouracil, everolimus, anastrozole, exemestane, capecitabine, cyclophosphamide, docetaxel, epirubicin, exemestane, toremifene, fulvestrant, letrozole, gemcitabine, trastuzumab, ixabepilone, lapatinib, tamoxifen, pertuzumab and toremifene, or mixtures thereof.

12. The use of any one of Claims 1 to 11, wherein *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is used.

13. The use of Claim 12, wherein about 550 to about 650 mg twice per day of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is used.

14. The use of any one of Claims 1 to 11, wherein (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is used.

15. The use of Claim 14, wherein about 25 to about 1800 mg per day of (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is used.

## CLAIMS

1. The use of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol in the manufacture of a medicament for the treatment of local or metastatic brain cancer that is caused by ErbB2 over-expression or amplification, wherein the use is after previous treatment for brain cancer.

2. The use of Claim 1, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is in the form of an oral dosage form.

3. The use of Claims 1 or 2, wherein the *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine or (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is in the form of a tablet.

4. The use of any one of Claims 1 to 3, wherein the use is in combination with another therapeutic agent.

5. The use of Claim 4, wherein the therapeutic agent is selected from trastuzumab, capecitabine, bevacizumab, paclitaxel and docetaxel.

6. The use of Claim 1, wherein the previous treatment for brain cancer is selected from surgery, radiation therapy and chemotherapy or mixtures thereof.

7. The use of Claim 1, wherein the previous treatment for brain cancer is selected from surgery, conventional external radiation therapy, three-dimensional conformal radiation therapy, intensity modulated radiation therapy, stereotactic radiosurgery, fractionated stereotactic radiation therapy, proton radiation therapy, internal or implant radiation therapy, temozolomide, bevacizumab, carmustine, lomustine, procarbazine, vincristine, tumor treating fields therapy, everolimus, procarbazine, lomustine, cisplatin, carboplatin and methotrexate or mixtures thereof.

8. The use of any one of Claims 1 to 5, wherein the use is after previous treatment for breast cancer.

9. The use of Claim 8, wherein the previous treatment for breast cancer is selected from surgery, sentinel lymph node biopsy followed by surgery, radiation therapy, chemotherapy, hormone therapy and targeted therapy.

10. The use of Claim 8, wherein the previous treatment for breast cancer is selected from lumpectomy, partial mastectomy, segmental mastectomy, total mastectomy, modified radical mastectomy, external radiation, internal radiation, methotrexate, paclitaxel albumin-stabilized nanoparticle formulation, ado-trastuzumab emtansine, doxorubicin, fluorouracil, everolimus, anastrozole, exemestane, capecitabine, cyclophosphamide, docetaxel, epirubicin, exemestane, toremifene, fulvestrant, letrozole, gemcitabine, trastuzumab, ixabepilone, lapatinib, tamoxifen, pertuzumab and toremifene, or mixtures thereof.

11. The use of any one of Claims 1 to 10, wherein *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is used.

12. The use of Claim 11, wherein about 550 to about 650 mg twice per day of *N*4-(4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)-*N*6-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)quinazoline-4,6-diamine is used.

13. The use of any one of Claims 1 to 10, wherein (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is used.

14. The use of Claim 13, wherein about 25 to about 1800 mg per day of (2-((4-((4-([1,2,4]triazolo[1,5-*a*]pyridin-7-yloxy)-3-methylphenyl)amino)quinazolin-6-yl)amino)-4-methyl-4,5-dihydrooxazol-4-yl)methanol is used.