PCT WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 7 : A61M 15/00 — A2

(11) International Publication Number: WO 00/21594

(43) International Publication Date: 20 April 2000 (20.04.00)

(21) International Application Number: PCT/US99/23698

(22) International Filing Date: 7 October 1999 (07.10.99)

(30) Priority Data:
60/103,702 9 October 1998 (09.10.98) US

(71) Applicant: INHALE THERAPEUTIC SYSTEMS, INC. [US/US]; 150 Industrial Road, San Carlos, CA 94070 (US).


(72) Inventors: CLARK, Andrew; 436 Valdez Avenue, Half Moon Bay, CA 94019 (US). SCHULER, Carlos; 10344 Denison Avenue, Cupertino, CA 95014 (US). PABOOJIAN, Steve; 2133 Avy Avenue, Menlo Park, CA 94025 (US).

(74) Agents: CAGAN, Felissa, H. et al.; Inhale Therapeutic Systems, Inc., 150 Industrial Road, San Carlos, CA 94070 (US).

(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published
Without international search report and to be republished upon receipt of that report.

(54) Title: FLOW RESISTANCE MODULATED AEROSOLIZED ACTIVE AGENT DELIVERY



(57) Abstract

The present invention is directed to methods and devices for delivering an active agent formulation to the lung of a human patient. The active agent formulation may be in dry powder form, it may be nebulized, or it may be in admixture with a propellant. The active agent formulation is delivered to a patient at a low inspiratory flow rate for an initial period of time to increase bioavailability of the active agent.