
SUMMARY ~~OF THE INVENTION~~

[0006] The present ~~invention~~ disclosure provides crystalline forms of salts of Compound 1, N-(4-fluorophenyl)-N-(4-((7-methoxy-6-(methylcarbamoyl)quinolin-4-yl)oxy)phenyl)cyclopropane-1,1-dicarboxamide, which has the structure:




Compound 1

[0007] Compound 1 is disclosed in WO 2019/148044, the contents of which is incorporated herein by reference in its entirety.

[0008] Specific crystalline salt forms of an active pharmaceutical ingredient (API), such as Compound 1, can have several advantages over other polymorphic or amorphous forms, such as increased stability during storage or processing, more favorable solubility, and increased bioavailability. Reported herein are several stable crystalline salt forms of Compound 1.

[0008a] In one aspect of the present disclosure, there is provided a crystalline salt form of Compound 1



Compound 1

selected from the group consisting of Compound 1 Besylate Form A, Compound 1 Besylate Form B, Compound 1 Benzoate Form A, Compound 1 Camsylate Form A, Compound 1 Camsylate Form B, Compound 1 Citrate Form A, Compound 1 Citrate Form B, Compound 1 Esylate Form A, Compound 1 Esylate Form B, Compound 1 HBr Form A, Compound 1 HBr Form B, Compound 1 Lactate Form A, Compound 1 Malate Form A, Compound 1 Maleate Form A, Compound 1 Maleate Form B, Compound 1 Mesylate Form A, Compound 1 Mesylate Form B, Compound 1 Mesylate Form C, Compound 1 Oxalate Form A, Compound 1 Oxalate

Form B, Compound 1 Oxalate Form C, Compound 1 Propionate Form A, Compound 1 Succinate Form A, Compound 1 Sulfate Form A, Compound 1 Sulfate Form B, Compound 1 Sulfate Form C, Compound 1 Tartrate Form A, Compound 1 Hemitartrate Form B, Compound 1 Tartrate Form C, Compound 1 Tosylate Form A, and Compound 1 Tosylate Form B.


[0008b] In another aspect of the present disclosure, there is provided a pharmaceutical composition comprising a crystalline salt form disclosed herein and a pharmaceutically acceptable excipient.

[0008c] In a further aspect of the present disclosure, there is provided a crystalline salt form disclosed herein, or a pharmaceutical composition disclosed herein, for use in therapy.

[0008d] In another aspect of the present disclosure, there is provided a use of a crystalline salt form disclosed herein, or a pharmaceutical composition disclosed herein, in the manufacture of a medicament for treating a disease, disorder, or syndrome mediated at least in part by modulating *in vivo* activity of a protein kinase.

[0008e] In a further aspect of the present disclosure, there is provided a use of a crystalline salt form disclosed herein, in the manufacture of a medicament for inhibiting a protein kinase.

[0009] In one ~~aspect~~example, the ~~invention~~ present disclosure includes a crystalline salt form of Compound 1



,

Compound 1

wherein the crystalline salt form is selected from Compound 1 Besylate Form A, Compound 1 Besylate Form B, Compound 1 Benzoate Form A, Compound 1 Camsylate Form A, Compound 1 Camsylate Form B, Compound 1 Citrate Form A, Compound 1 Citrate Form B, Compound 1 Esylate Form A, Compound 1 Esylate Form B, Compound 1 HBr Form A, Compound 1 HBr Form B, Compound 1 Lactate Form A, Compound 1 Malate Form A, Compound 1 Maleate Form A, Compound 1 Maleate Form B, Compound 1 Mesylate Form A, Compound 1 Mesylate Form B, Compound 1 Mesylate Form C, Compound 1 Oxalate Form A, Compound 1 Oxalate Form B, Compound 1 Oxalate Form C, Compound 1 Propionate Form A, Compound 1 Succinate Form A, Compound 1 Sulfate Form A, Compound 1 Sulfate Form B, Compound 1 Sulfate Form C, Compound 1 Tartrate Form A,

Compound 1 Hemitartrate Form B, Compound 1 Tartrate Form C, Compound 1 Tosylate Form A, and Compound 1 Tosylate Form B.

[0010] In one example~~aspect~~, the present disclosure ~~invention~~ includes a pharmaceutical composition comprising a crystalline salt form described herein and a pharmaceutically acceptable excipient.

[0011] In another example~~aspect~~, the present disclosure ~~invention~~ includes a method of treating a disease, disorder, or syndrome mediated at least in part by modulating *in vivo* activity of a protein kinase, comprising administering to a subject in need thereof a crystalline salt form or pharmaceutical composition described herein.

[0012] In one example~~embodiment of this aspect~~, the disease, disorder, or syndrome mediated at least in part by modulating *in vivo* activity of a protein kinase is cancer.

[0013] In another example~~aspect~~, the present disclosure ~~invention~~ includes a method for inhibiting a protein kinase, the method comprising contacting the protein kinase with a crystalline salt form or pharmaceutical composition described herein.

[0014] In one example~~embodiment of this aspect~~, the protein kinase is Axl, Mer, c-Met, KDR, or a combination thereof.

[0015] A further example~~aspect~~ provides processes for making the solid forms described herein.

BRIEF DESCRIPTION OF THE FIGURES

[0016] **FIG. 1** is an XRPD pattern of Compound 1 Besylate Form A.

[0017] **FIG. 2** is a DSC thermogram of Compound 1 Besylate Form A.

[0018] **FIG. 3** is a TGA thermogram of Compound 1 Besylate Form A.

[0019] **FIG. 4** is an XRPD pattern of Compound 1 Besylate Form B (mixture with Compound 1 Besylate Form A).

[0020] **FIG. 5** is an XRPD pattern of Compound 1 Benzoate Form A.

[0021] **FIG. 6** is an XRPD pattern of Compound 1 Camsylate Form A.

[0022] **FIG. 7** is a DSC thermogram of Compound 1 Camsylate Form A.

[0023] **FIG. 8** is a TGA thermogram of Compound 1 Camsylate Form A.

[0024] **FIG. 9** is an XRPD pattern of Compound 1 Camsylate Form B.

[0025] **FIG. 10** is an XRPD pattern of Compound 1 Citrate Form A (mixture with Compound 1 Form A).

[0026] **FIG. 11** is an XRPD pattern of Compound 1 Citrate Form B.


[0027] **FIG. 12** is a DSC thermogram of Compound 1 Citrate Form B.

[0028] **FIG. 13** is a TGA thermogram of Compound 1 Citrate Form B.

[0029] **FIG. 14** is a DVS isotherm plot of Compound 1 Citrate Form B.

WHAT IS CLAIMED IS:

1. A crystalline salt form of Compound 1



,

Compound 1

selected from the group consisting of Compound 1 Besylate Form A, Compound 1 Besylate Form B, Compound 1 Benzoate Form A, Compound 1 Camsylate Form A, Compound 1 Camsylate Form B, Compound 1 Citrate Form A, Compound 1 Citrate Form B, Compound 1 Esylate Form A, Compound 1 Esylate Form B, Compound 1 HBr Form A, Compound 1 HBr Form B, Compound 1 Lactate Form A, Compound 1 Malate Form A, Compound 1 Maleate Form A, Compound 1 Maleate Form B, Compound 1 Mesylate Form A, Compound 1 Mesylate Form B, Compound 1 Mesylate Form C, Compound 1 Oxalate Form A, Compound 1 Oxalate Form B, Compound 1 Oxalate Form C, Compound 1 Propionate Form A, Compound 1 Succinate Form A, Compound 1 Sulfate Form A, Compound 1 Sulfate Form B, Compound 1 Sulfate Form C, Compound 1 Tartrate Form A, Compound 1 Hemitartrate Form B, Compound 1 Tartrate Form C, Compound 1 Tosylate Form A, and Compound 1 Tosylate Form B.

~~2. The crystalline salt form of claim 1, characterized as Compound 1 Besylate Form A.~~

~~3.~~2. The crystalline salt form of ~~any one of~~ claim~~s~~ 1 ~~or 2~~, wherein:

~~the~~ Compound 1 Besylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 3.78, 7.56, 10.08, 10.42, 11.70, 11.78, 12.00, 12.71, 13.20, 13.92, 14.06, 14.44, 14.93, 15.18, 15.96, 16.21, 16.40, 16.73, 16.86, 17.38, 18.32, 19.07, 20.07, 20.26, 20.51, 20.94, 21.45, 21.60, 22.30, 22.90, 23.05, 23.28, 23.71, 24.28, 24.97, 25.30, 25.69, 26.33, 26.71, 27.29, 27.65, 28.07, 29.22, and 29.50;

Compound 1 Besylate Form B is characterized by an XRPD pattern substantially identical to Figure 4;

Compound 1 Benzoate Form A is characterized by an XRPD pattern substantially identical to Figure 5;

Compound 1 Camsylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.55, 7.90, 8.25, 8.38, 9.55, 9.87, 10.26, 11.12, 11.34, 12.50, 12.94, 13.40, 13.65, 14.79, 15.83, 16.16, 16.41, 16.54, 16.82, 17.10, 17.69, 18.02, 18.22, 18.47, 18.91, 19.10, 19.47, 20.04, 20.87, 21.05, 22.16, 22.36, 22.93, 23.71, 24.55, 25.33, 26.02, 26.40, 27.09, 27.33, 27.97, 28.32, 28.68, and 28.98;

Compound 1 Camsylate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.25, 8.32, 8.51, 8.76, 9.05, 9.58, 9.88, 10.00, 10.52, 10.80, 11.06, 11.96, 12.24, 12.58, 13.98, 15.62, 15.81, 16.15, 16.70, 17.12, 17.42, 17.71, 18.41, 19.37, 19.55, 20.63, 21.13, 21.56, 21.89, 22.50, 23.04, 24.43, 26.08, 27.07, 27.97, 28.22, and 28.73;

Compound 1 Citrate Form A is characterized by an XRPD pattern substantially identical to Figure 10;

Compound 1 Citrate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.58, 5.95, 7.61, 8.88, 9.19, 9.94, 11.93, 13.26, 13.79, 14.29, 14.68, 15.26, 15.95, 16.64, 17.11, 17.87, 18.21, 18.43, 18.90, 19.36, 19.67, 20.14, 20.57, 21.47, 22.30, 22.64, 23.06, 23.80, 24.25, 24.55, 24.81, 25.10, 25.45, 25.54, 25.85, 26.35, 26.51, 26.83, 27.16, 27.57, 27.85, 28.08, and 28.47;

Compound 1 Esylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 3.97, 7.95, 9.89, 10.62, 10.92, 11.03, 11.34, 11.77, 12.03, 12.53, 12.71, 13.27, 14.00, 14.70, 15.08, 15.94, 16.17, 16.92, 17.01, 17.19, 17.35, 17.61, 18.15, 18.76, 19.03, 19.66, 19.89, 20.26, 20.65, 20.78, 21.44, 22.08, 23.14, 23.27, 23.99, 24.32, 25.43, 25.66, 25.85, 26.52, 26.87, 27.61, 27.96, 28.18, 28.40, and 28.91;

Compound 1 Esylate Form B is characterized by an XRPD pattern substantially identical to Figure 18;

Compound 1 HBr Form A is characterized by an XRPD pattern substantially identical to Figure 19;

Compound 1 HBr Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.26, 8.29, 9.75, 9.91, 10.56, 10.72, 12.25, 12.50, 12.75, 13.31, 13.54, 14.11, 14.72, 14.89, 15.87, 16.29, 16.50, 17.36, 17.78, 18.09, 18.44, 18.75, 19.12, 19.24, 19.56, 19.92, 20.40, 20.55, 20.69, 21.18, 22.09, 22.64, 23.24, 24.56, 25.80, 26.52, 27.23, 27.43, 28.13, 28.46, 28.76, 29.08, 29.66, and 30.07;

Compound 1 Lactate Form A is characterized by an XRPD pattern substantially identical to Figure 21;

Compound 1 Malate Form A is characterized by an XRPD pattern substantially identical to Figure 22;

Compound 1 Maleate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 7.58, 8.34, 10.92, 12.38, 12.87, 14.17, 15.13, 15.78, 16.43, 16.91, 17.51, 18.09, 18.71, 20.30, 21.60, 21.99, 22.30, 22.49, 23.32, 23.56, 23.92, 24.79, 26.04, 26.41, 26.92, 27.62, 28.18, 28.54, and 29.01;

Compound 1 Maleate Form B is characterized by an XRPD pattern substantially identical to Figure 28.

Compound 1 Mesylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.06, 6.65, 7.66, 8.35, 10.16, 10.72, 11.98, 12.62, 13.32, 14.41, 14.79, 15.42, 15.72, 15.89, 16.23, 16.77, 17.20, 17.59, 18.08, 18.40, 18.81, 19.33, 19.53, 19.71, 20.14, 20.38, 20.73, 21.05, 21.56, 21.83, 22.16, 23.14, 24.06, 24.54, 24.77, 25.75, 26.82, 27.71, 27.94, 28.71, 29.27, and 29.85;

Compound 1 Mesylate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.14, 6.62, 7.72, 7.89, 10.33, 10.73, 12.74, 13.31, 13.88, 14.23, 14.61, 15.21, 15.52, 15.87, 16.17, 17.18, 17.41, 17.88, 18.15, 18.46, 18.73, 19.13, 19.54, 20.15, 20.53, 20.75, 20.98, 21.27, 21.60, 22.03, 22.28, 23.07, 23.32, 24.16, 24.42, 24.87, 25.13, 25.64, 26.24, 26.95, 27.17, 27.49, 27.77, 28.42, and 29.54;

Compound 1 Mesylate Form C is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 7.55, 9.53, 10.25, 11.04, 11.49, 11.65, 12.24, 12.51, 13.88, 14.44, 15.36, 15.66, 16.29, 16.58, 17.16, 17.54, 19.04, 19.20, 19.79, 20.54, 21.12, 21.24, 21.62, 22.22, 23.15, 23.57, 23.99, 24.61, 24.77, 25.22, 25.61, 25.99, 26.60, 26.89, 27.29, 27.91, 29.03, and 29.29;

Compound 1 Oxalate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.90, 5.25, 6.20, 8.05, 9.87, 10.54, 12.29, 14.79, 15.35, 15.84, 16.49, 16.89, 17.55, 18.27, 18.67, 19.12, 19.40, 19.81, 20.26, 20.50, 21.06, 21.74, 22.17, 22.93, 23.26, 23.50, 23.81, 24.66, 25.06, 25.95, 26.90, 27.82, and 28.24;

Compound 1 Oxalate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.20, 7.59, 8.02, 8.42, 10.53, 12.36, 12.66, 13.09, 13.35, 13.88, 14.15, 14.46, 15.06, 15.23, 17.42, 17.75, 18.28, 19.07, 19.52, 20.33,

21.18, 21.43, 21.74, 22.08, 22.59, 23.53, 24.21, 24.52, 24.71, 25.24, 25.85, 26.24, 26.55, 26.90, 27.43, 28.56, 29.30, and 29.51;

Compound 1 Oxalate Form C is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.13, 7.70, 10.29, 10.59, 13.09, 13.70, 14.13, 15.33, 15.87, 16.57, 17.19, 17.50, 17.83, 18.15, 18.61, 19.24, 19.42, 20.11, 20.34, 20.64, 21.07, 21.47, 21.59, 21.91, 22.56, 22.92, 23.37, 23.87, 24.35, 25.35, 25.91, 26.47, 26.68, 27.74, 28.47, and 29.14;

Compound 1 Propionate Form A is characterized by an XRPD pattern substantially identical to Figure 38;

Compound 1 Succinate Form A is characterized by an XRPD pattern substantially identical to Figure 39;

Compound 1 Sulfate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.74, 5.85, 9.47, 10.04, 11.74, 11.95, 13.77, 14.99, 15.42, 15.71, 16.48, 17.36, 17.66, 17.92, 18.59, 18.95, 19.36, 19.74, 20.22, 20.77, 21.38, 21.86, 22.74, 23.30, 23.70, 24.33, 25.07, 25.32, 25.67, 26.23, and 27.00;

Compound 1 Sulfate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.27, 8.57, 9.32, 10.95, 11.29, 12.07, 12.87, 13.25, 13.82, 14.13, 14.67, 16.94, 17.21, 17.64, 19.99, 20.26, 21.03, 21.44, 22.00, 22.71, 23.85, 25.20, and 26.02;

Compound 1 Sulfate Form C is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 3.19, 5.35, 9.04, 9.62, 10.73, 11.31, 11.60, 12.27, 12.67, 13.76, 14.31, 14.65, 15.21, 15.63, 16.11, 16.33, 16.63, 17.11, 18.32, 18.75, 18.99, 19.36, 19.47, 20.16, 20.55, 21.01, 21.52, 21.62, 21.89, 22.19, 22.53, 23.19, 23.79, 24.23, 24.81, 25.07, 25.41, 26.06, 26.73, 27.58, 28.16, 28.63, 28.90, 29.22, 29.47, and 29.73;

Compound 1 Tartrate Form A is characterized by an XRPD pattern substantially identical to Figure 43;

Compound 1 Hemitartrate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.17, 5.66, 7.22, 9.27, 10.10, 10.40, 11.37, 11.55, 12.77, 13.94, 14.51, 15.60, 16.06, 16.48, 17.09, 17.69, 17.90, 18.27, 18.94, 19.15, 19.59, 19.81, 20.06, 20.32, 21.23, 21.83, 22.34, 22.84, 23.53, 24.26, 24.70, 25.26, 25.80, 26.78, 27.29, 27.90, 29.04, and 29.45;

Compound 1 Tartrate Form C is characterized by an XRPD pattern substantially identical to Figure 69;

Compound 1 Tosylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.81, 7.05, 8.52, 9.74, 10.40, 10.99, 11.26, 11.66, 12.58, 12.81, 14.28, 14.62, 15.30, 15.80, 16.03, 16.34, 17.12, 17.55, 18.03, 18.94, 19.26, 19.67, 20.31, 20.65, 21.02, 21.43, 22.11, 22.44, 22.66, 22.94, 23.07, 23.54, 23.79, 24.55, 24.88, 25.42, 25.83, 26.10, 26.53, 27.70, 28.11, 28.62, 29.18, 29.40, 29.66, and 29.83; and

Compound 1 Tosylate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.61, 8.69, 9.25, 10.68, 12.44, 12.69, 13.76, 14.42, 14.81, 15.47, 16.08, 16.57, 17.45, 17.76, 18.44, 18.83, 19.01, 19.35, 20.42, 20.84, 21.15, 21.69, 22.09, 22.86, 23.12, 24.08, 24.53, 24.86, 25.22, 25.91, 26.33, 26.82, 27.63, 28.61, 29.11, and 29.77.

~~4.~~3. The crystalline salt form of ~~any one of~~ claim~~s~~ 1 or 2 ~~3~~, wherein:

~~the~~ Compound 1 Besylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 3.78, 7.56, 10.42, 20.51, 20.94, 21.60, and 25.30;

Compound 1 Camsylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.55, 9.87, 10.26, 16.16, 16.82, 20.87, 22.36, and 23.71;

Compound 1 Camsylate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.25, 10.52, 16.15, 21.13, 21.56, and 21.89;

Compound 1 Citrate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.58, 5.95, 8.88, 11.93, 13.79, 14.29, 16.64, 18.90, 21.47, 23.80, 24.25, and 25.85;

Compound 1 Esylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 3.97, 7.95, 9.89, 10.62, 13.27, 15.08, 16.17, 16.92, 17.01, 20.65, 20.78, 21.44, 22.08, 23.99, 25.43, and 26.52;

Compound 1 HBr Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.26, 9.91, 10.72, 20.55, 20.69, 21.18, 22.09, 23.24, 24.56, and 25.80;

Compound 1 Maleate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 10.92, 15.13, 15.78, 16.43, 18.09, 18.71, 20.30, 21.99, 22.30, 22.49, 23.32, 23.56, 24.79, 26.41, 26.92, and 29.01;

Compound 1 Mesylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.06, 7.66, 14.79, 15.42, 16.23, 17.20, 17.59, 18.40, 19.33, 19.53, 20.38, 20.73, 21.83, 22.16, 23.14, 24.06, and 24.77;

Compound 1 Mesylate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 14.61, 15.21, 16.17, 17.18, 17.41, 18.46, 19.13, 19.54, 20.15, 20.53, 20.75, 20.98, 21.27, 21.60, 22.03, 23.32, 24.16, 24.42, 24.87, and 25.13;

Compound 1 Mesylate Form C is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 7.55, 9.53, 11.49, 12.51, 14.44, 19.04, 20.54, 21.12, 21.24, and 23.15;

Compound 1 Oxalate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.90, 5.25, 9.87, 10.54, 12.29, 15.35, 18.67, 19.40, 19.81, 20.26, 22.93, and 24.66;

Compound 1 Oxalate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.20, 7.59, 10.53, 15.06, 15.23, 19.52, 21.18, 21.43, 22.59, 24.21, and 25.85;

Compound 1 Oxalate Form C is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 7.70, 10.29, 10.59, 13.09, 13.70, 15.33, 17.83, 18.15, 20.11, 21.07, 21.91, and 22.56;

Compound 1 Sulfate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.74, 5.85, 9.47, 10.04, 11.74, 11.95, 16.48, 20.22, and 22.74;

Compound 1 Sulfate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.27, 8.57, 9.32, 16.94, 19.99, 21.03, 21.44, 22.00, and 26.02;

Compound 1 Sulfate Form C is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 3.19, 5.35, 9.62, 12.27, 16.33, 17.11, 18.75, 18.99, 19.36, 19.47, 20.55, 21.01, 21.52, 21.62, 21.89, 22.19, 23.79, 25.07, and 26.06;

Compound 1 Hemitartrate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.17, 5.66, 10.10, 10.40, 11.37, 11.55, 14.51, 17.09, 21.23, 21.83, 23.53, 24.26, 25.26, and 26.78;

Compound 1 Tosylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 8.52, 9.74, 14.28, 16.03, 17.55, 18.94, 19.26, 20.31, 21.02, 21.43, 22.11, 22.44, 22.66, 23.07, 24.55, and 27.70; and

Compound 1 Tosylate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.61, 9.25, 12.44, 15.47, 17.76, 19.01, 19.35, 25.91, 26.33, and 27.63.

~~5.~~4. The crystalline salt form of any one of claims 1-~~4~~3, wherein the Compound 1 Besylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 3.78, 7.56, 10.42, 20.51, 20.94, 21.60, and 25.30;

Compound 1 Camsylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.55, 9.87, 10.26, 16.16, 16.82, 20.87, 22.36, and 23.71;

Compound 1 Camsylate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.25, 10.52, 16.15, 21.13, 21.56, and 21.89;

Compound 1 Citrate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.58, 5.95, 8.88, 11.93, 13.79, 14.29, 16.64, 18.90, 21.47, 23.80, 24.25, and 25.85;

Compound 1 Esylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 3.97, 7.95, 9.89, 10.62, 13.27, 15.08, 16.17, 16.92, 17.01, 20.65, 20.78, 21.44, 22.08, 23.99, 25.43, and 26.52;

Compound 1 HBr Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.26, 9.91, 10.72, 20.55, 20.69, 21.18, 22.09, 23.24, 24.56, and 25.80;

Compound 1 Maleate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 10.92, 15.13, 15.78, 16.43, 18.09, 18.71, 20.30, 21.99, 22.30, 22.49, 23.32, 23.56, 24.79, 26.41, 26.92, and 29.01;

Compound 1 Mesylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.06, 7.66, 14.79, 15.42, 16.23, 17.20, 17.59, 18.40, 19.33, 19.53, 20.38, 20.73, 21.83, 22.16, 23.14, 24.06, and 24.77;

Compound 1 Mesylate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 14.61, 15.21, 16.17, 17.18, 17.41, 18.46, 19.13, 19.54, 20.15, 20.53, 20.75, 20.98, 21.27, 21.60, 22.03, 23.32, 24.16, 24.42, 24.87, and 25.13;

Compound 1 Mesylate Form C is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 7.55, 9.53, 11.49, 12.51, 14.44, 19.04, 20.54, 21.12, 21.24, and 23.15;

Compound 1 Oxalate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.90, 5.25, 9.87, 10.54, 12.29, 15.35, 18.67, 19.40, 19.81, 20.26, 22.93, and 24.66;

Compound 1 Oxalate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.20, 7.59, 10.53, 15.06, 15.23, 19.52, 21.18, 21.43, 22.59, 24.21, and 25.85;

Compound 1 Oxalate Form C is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 7.70, 10.29, 10.59, 13.09, 13.70, 15.33, 17.83, 18.15, 20.11, 21.07, 21.91, and 22.56;

Compound 1 Sulfate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.74, 5.85, 9.47, 10.04, 11.74, 11.95, 16.48, 20.22, and 22.74;

Compound 1 Sulfate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.27, 8.57, 9.32, 16.94, 19.99, 21.03, 21.44, 22.00, and 26.02;

Compound 1 Sulfate Form C is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 3.19, 5.35, 9.62, 12.27, 16.33, 17.11, 18.75, 18.99, 19.36, 19.47, 20.55, 21.01, 21.52, 21.62, 21.89, 22.19, 23.79, 25.07, and 26.06;

Compound 1 Hemitartrate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.17, 5.66, 10.10, 10.40, 11.37, 11.55, 14.51, 17.09, 21.23, 21.83, 23.53, 24.26, 25.26, and 26.78;

Compound 1 Tosylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 8.52, 9.74, 14.28, 16.03, 17.55, 18.94, 19.26, 20.31, 21.02, 21.43, 22.11, 22.44, 22.66, 23.07, 24.55, and 27.70; and

Compound 1 Tosylate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.61, 9.25, 12.44, 15.47, 17.76, 19.01, 19.35, 25.91, 26.33, and 27.63.

~~6.~~5. The crystalline salt form of any one of claims 1-~~5~~4, wherein the Compound 1 Besylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 3.78, 7.56, 10.08, 10.42, 11.70, 11.78, 12.00, 12.71, 13.20, 13.92, 14.06, 14.44, 14.93, 15.18, 15.96, 16.21, 16.40, 16.73, 16.86, 17.38, 18.32, 19.07, 20.07, 20.26, 20.51, 20.94, 21.45, 21.60, 22.30, 22.90, 23.05, 23.28, 23.71, 24.28, 24.97, 25.30, 25.69, 26.33, 26.71, 27.29, 27.65, 28.07, 29.22, and 29.50;

Compound 1 Camsylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.55, 7.90, 8.25, 8.38, 9.55, 9.87, 10.26, 11.12, 11.34, 12.50, 12.94, 13.40, 13.65, 14.79, 15.83, 16.16, 16.41, 16.54, 16.82, 17.10, 17.69, 18.02, 18.22, 18.47, 18.91, 19.10, 19.47, 20.04, 20.87, 21.05, 22.16, 22.36, 22.93, 23.71, 24.55, 25.33, 26.02, 26.40, 27.09, 27.33, 27.97, 28.32, 28.68, and 28.98;

Compound 1 Camsylate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.25, 8.32, 8.51, 8.76, 9.05, 9.58, 9.88, 10.00, 10.52, 10.80, 11.06, 11.96, 12.24, 12.58, 13.98, 15.62, 15.81, 16.15, 16.70, 17.12, 17.42, 17.71, 18.41, 19.37, 19.55, 20.63, 21.13, 21.56, 21.89, 22.50, 23.04, 24.43, 26.08, 27.07, 27.97, 28.22, and 28.73;

Compound 1 Citrate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.58, 5.95, 7.61, 8.88, 9.19, 9.94, 11.93, 13.26, 13.79, 14.29, 14.68, 15.26, 15.95, 16.64, 17.11, 17.87, 18.21, 18.43, 18.90, 19.36, 19.67, 20.14, 20.57, 21.47, 22.30, 22.64, 23.06, 23.80, 24.25, 24.55, 24.81, 25.10, 25.45, 25.54, 25.85, 26.35, 26.51, 26.83, 27.16, 27.57, 27.85, 28.08, and 28.47;

Compound 1 Esylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 3.97, 7.95, 9.89, 10.62, 10.92, 11.03, 11.34, 11.77, 12.03, 12.53, 12.71, 13.27, 14.00, 14.70, 15.08, 15.94, 16.17, 16.92, 17.01, 17.19, 17.35, 17.61, 18.15, 18.76, 19.03, 19.66, 19.89, 20.26, 20.65, 20.78, 21.44, 22.08, 23.14, 23.27, 23.99, 24.32, 25.43, 25.66, 25.85, 26.52, 26.87, 27.61, 27.96, 28.18, 28.40, and 28.91;

Compound 1 HBr Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.26, 8.29, 9.75, 9.91, 10.56, 10.72, 12.25, 12.50, 12.75, 13.31, 13.54, 14.11, 14.72, 14.89, 15.87, 16.29, 16.50, 17.36, 17.78, 18.09, 18.44, 18.75, 19.12, 19.24, 19.56, 19.92, 20.40, 20.55, 20.69, 21.18, 22.09, 22.64, 23.24, 24.56, 25.80, 26.52, 27.23, 27.43, 28.13, 28.46, 28.76, 29.08, 29.66, and 30.07;

Compound 1 Maleate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 7.58, 8.34, 10.92, 12.38, 12.87, 14.17, 15.13, 15.78, 16.43, 16.91, 17.51, 18.09, 18.71, 20.30, 21.60, 21.99, 22.30, 22.49, 23.32, 23.56, 23.92, 24.79, 26.04, 26.41, 26.92, 27.62, 28.18, 28.54, and 29.01;

Compound 1 Mesylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.06, 6.65, 7.66, 8.35, 10.16, 10.72, 11.98, 12.62, 13.32, 14.41, 14.79, 15.42, 15.72, 15.89, 16.23, 16.77, 17.20, 17.59, 18.08, 18.40, 18.81, 19.33, 19.53, 19.71, 20.14, 20.38, 20.73, 21.05, 21.56, 21.83, 22.16, 23.14, 24.06, 24.54, 24.77, 25.75, 26.82, 27.71, 27.94, 28.71, 29.27, and 29.85;

Compound 1 Mesylate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.14, 6.62, 7.72, 7.89, 10.33, 10.73, 12.74, 13.31, 13.88, 14.23, 14.61, 15.21, 15.52, 15.87, 16.17, 17.18, 17.41, 17.88, 18.15, 18.46, 18.73, 19.13, 19.54, 20.15, 20.53, 20.75, 20.98, 21.27, 21.60, 22.03, 22.28, 23.07, 23.32, 24.16, 24.42, 24.87, 25.13, 25.64, 26.24, 26.95, 27.17, 27.49, 27.77, 28.42, and 29.54;

Compound 1 Mesylate Form C is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 7.55, 9.53, 10.25, 11.04, 11.49, 11.65, 12.24, 12.51, 13.88, 14.44, 15.36, 15.66, 16.29, 16.58, 17.16, 17.54, 19.04, 19.20, 19.79, 20.54, 21.12, 21.24, 21.62, 22.22, 23.15, 23.57, 23.99, 24.61, 24.77, 25.22, 25.61, 25.99, 26.60, 26.89, 27.29, 27.91, 29.03, and 29.29;

Compound 1 Oxalate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.90, 5.25, 6.20, 8.05, 9.87, 10.54, 12.29, 14.79, 15.35, 15.84, 16.49, 16.89, 17.55, 18.27, 18.67, 19.12, 19.40, 19.81, 20.26, 20.50, 21.06, 21.74, 22.17, 22.93, 23.26, 23.50, 23.81, 24.66, 25.06, 25.95, 26.90, 27.82, and 28.24;

Compound 1 Oxalate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.20, 7.59, 8.02, 8.42, 10.53, 12.36, 12.66, 13.09, 13.35, 13.88, 14.15, 14.46, 15.06, 15.23, 17.42, 17.75, 18.28, 19.07, 19.52, 20.33, 21.18, 21.43, 21.74, 22.08, 22.59, 23.53, 24.21, 24.52, 24.71, 25.24, 25.85, 26.24, 26.55, 26.90, 27.43, 28.56, 29.30, and 29.51;

Compound 1 Oxalate Form C is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.13, 7.70, 10.29, 10.59, 13.09, 13.70, 14.13, 15.33, 15.87, 16.57, 17.19, 17.50, 17.83, 18.15, 18.61, 19.24, 19.42, 20.11, 20.34, 20.64, 21.07, 21.47, 21.59, 21.91, 22.56, 22.92, 23.37, 23.87, 24.35, 25.35, 25.91, 26.47, 26.68, 27.74, 28.47, and 29.14;

Compound 1 Sulfate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.74, 5.85, 9.47, 10.04, 11.74, 11.95, 13.77, 14.99, 15.42, 15.71, 16.48, 17.36, 17.66, 17.92, 18.59, 18.95, 19.36, 19.74, 20.22, 20.77, 21.38, 21.86, 22.74, 23.30, 23.70, 24.33, 25.07, 25.32, 25.67, 26.23, and 27.00;

Compound 1 Sulfate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.27, 8.57, 9.32, 10.95, 11.29, 12.07, 12.87, 13.25, 13.82, 14.13, 14.67, 16.94, 17.21, 17.64, 19.99, 20.26, 21.03, 21.44, 22.00, 22.71, 23.85, 25.20, and 26.02;

Compound 1 Sulfate Form C is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 3.19, 5.35, 9.04, 9.62, 10.73, 11.31, 11.60, 12.27, 12.67, 13.76, 14.31, 14.65, 15.21, 15.63, 16.11, 16.33, 16.63, 17.11, 18.32, 18.75, 18.99, 19.36, 19.47, 20.16, 20.55, 21.01, 21.52, 21.62, 21.89, 22.19, 22.53, 23.19, 23.79, 24.23, 24.81, 25.07, 25.41, 26.06, 26.73, 27.58, 28.16, 28.63, 28.90, 29.22, 29.47, and 29.73;

Compound 1 Hemitartrate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.17, 5.66, 7.22, 9.27, 10.10, 10.40, 11.37, 11.55, 12.77, 13.94, 14.51, 15.60, 16.06, 16.48, 17.09, 17.69, 17.90, 18.27, 18.94, 19.15, 19.59, 19.81, 20.06, 20.32, 21.23, 21.83, 22.34, 22.84, 23.53, 24.26, 24.70, 25.26, 25.80, 26.78, 27.29, 27.90, 29.04, and 29.45;

Compound 1 Tosylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.81, 7.05, 8.52, 9.74, 10.40, 10.99, 11.26, 11.66, 12.58, 12.81, 14.28, 14.62, 15.30, 15.80, 16.03, 16.34, 17.12, 17.55, 18.03, 18.94, 19.26, 19.67, 20.31, 20.65, 21.02, 21.43, 22.11, 22.44, 22.66, 22.94, 23.07, 23.54, 23.79, 24.55, 24.88, 25.42, 25.83, 26.10, 26.53, 27.70, 28.11, 28.62, 29.18, 29.40, 29.66, and 29.83; and

Compound 1 Tosylate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.61, 8.69, 9.25, 10.68, 12.44, 12.69, 13.76, 14.42, 14.81, 15.47, 16.08, 16.57, 17.45, 17.76, 18.44, 18.83, 19.01, 19.35, 20.42, 20.84, 21.15, 21.69, 22.09, 22.86, 23.12, 24.08, 24.53, 24.86, 25.22, 25.91, 26.33, 26.82, 27.63, 28.61, 29.11, and 29.77.

~~7.~~6. The crystalline salt form of any one of claims 1-~~6~~5, wherein the Compound 1 Besylate Form A is characterized by a first endotherm at a temperature of about 144 ºC and a second endotherm at a temperature of about 170 ºC in a DSC thermogram;

Compound 1 Camsylate Form A is characterized by a first endotherm at a temperature of about 110 ºC and a second endotherm at a temperature of about 174 ºC in a DSC thermogram;

Compound 1 Citrate Form B is characterized by an endotherm with an onset temperature at about 175 ºC in a DSC thermogram;

Compound 1 Esylate Form A is characterized by a first endotherm at a temperature of about 69 ºC, a second endotherm at about 156 ºC, and a third endotherm at a temperature of about 190 ºC, in a DSC thermogram;

Compound 1 Maleate Form A is characterized by an endotherm at a temperature of about 117 ºC in a DSC thermogram;

Compound 1 Mesylate Form C is characterized by a first endotherm with a peak at a temperature of about 69 °C, wherein the first endotherm has a lower limit of about 36 and an upper limit of a temperature of about 96 °C, and a second endotherm with a peak at a temperature of about 208 °C, wherein the second endotherm has a lower limit of a temperature of about 169 and an upper limit of a temperature of about 219 °C;

Compound 1 Hemitartrate Form B is characterized by a first endotherm at a temperature of about 111 ºC, and a second endotherm at a temperature of about 148 ºC in a DSC thermogram;

Compound 1 Tosylate Form A is characterized by an endotherm with an onset temperature at about 214 ºC in a DSC thermogram; and

Compound 1 Tosylate Form B is characterized by a sharp endotherm with an onset temperature at ~183 °C in a DSC thermogram.

~~8.~~7. The crystalline salt form of any one of claims 1-~~7~~6, wherein the Compound 1 Besylate Form A is characterized by a weight loss of about 7.9% over a temperature range of 39-177 ºC in a TGA thermogram;

Compound 1 Camsylate Form A is characterized by a weight loss of about 5.9% over a temperature range of 38-170 ºC in a TGA thermogram;

Compound 1 Citrate Form B is characterized by a weight loss beginning at a temperature of about 159 ºC and ending at about 220 ºC in a TGA thermogram;

Compound 1 Esylate Form A is characterized by a weight loss of about 8.5 wt% over a temperature range of 38-188 ºC in a TGA thermogram;

Compound 1 Maleate Form A is characterized by a weight loss of about 3.1 wt% in the temperature range of 42-146 ºC in a TGA thermogram;

Compound 1 Mesylate Form C is characterized by a weight loss of from 0.5 to 3.0 wt% in the temperature range of 36-130 ºC in a TGA thermogram;

Compound 1 Hemitartrate Form B is characterized by a weight loss of about 7.4 wt% in the temperature range of 39-157 ºC in a TGA thermogram;

Compound 1 Tosylate Form A is characterized by a weight loss of about 0.9 wt% in the temperature range of 72-231 ºC in a TGA thermogram; and

Compound 1 Tosylate Form B is characterized by a weight loss of about 0.7 wt% in the temperature range of 39-214 ºC in a TGA thermogram.

~~9. The crystalline salt form of claim 1, characterized as Compound 1 Besylate Form B.~~

~~10. The crystalline salt form of claim 9, wherein the Compound 1 Besylate Form B is characterized by an XRPD pattern substantially identical to Figure 4.~~

~~11. The crystalline salt form of claim 1, characterized as Compound 1 Benzoate Form A.~~

~~12. The crystalline salt form of claim 11, wherein the Compound 1 Benzoate Form A is characterized by an XRPD pattern substantially identical to Figure 5.~~

~~13. The crystalline salt form of claim 1, characterized as Compound 1 Camsylate Form A.~~

~~14. The crystalline salt form of claim 13, wherein the Compound 1 Camsylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.55, 7.90, 8.25, 8.38, 9.55, 9.87, 10.26, 11.12, 11.34, 12.50, 12.94, 13.40, 13.65, 14.79, 15.83, 16.16, 16.41, 16.54, 16.82, 17.10, 17.69, 18.02, 18.22, 18.47, 18.91, 19.10, 19.47, 20.04, 20.87, 21.05, 22.16, 22.36, 22.93, 23.71, 24.55, 25.33, 26.02, 26.40, 27.09, 27.33, 27.97, 28.32, 28.68, and 28.98.~~

~~15. The crystalline salt form of any one of claims 13-14, wherein the Compound 1 Camsylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.55, 9.87, 10.26, 16.16, 16.82, 20.87, 22.36, and 23.71.~~

~~16. The crystalline salt form of any one of claims 13-15, wherein the Compound 1 Camsylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.55, 9.87, 10.26, 16.16, 16.82, 20.87, 22.36, and 23.71.~~

~~17. The crystalline salt form of any one of claims 13-16, wherein the Compound 1 Camsylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.55, 7.90, 8.25, 8.38, 9.55, 9.87, 10.26, 11.12, 11.34, 12.50, 12.94, 13.40, 13.65, 14.79, 15.83, 16.16, 16.41, 16.54, 16.82, 17.10, 17.69, 18.02, 18.22, 18.47, 18.91, 19.10, 19.47, 20.04, 20.87, 21.05, 22.16, 22.36, 22.93, 23.71, 24.55, 25.33, 26.02, 26.40, 27.09, 27.33, 27.97, 28.32, 28.68, and 28.98.~~

18. The crystalline salt form of any one of claims 13-17, wherein the Compound 1 Camsylate Form A is characterized by a first endotherm at a temperature of about 110 ºC and a second endotherm at a temperature of about 174 ºC in a DSC thermogram.

19. The crystalline salt form of any one of claims 13-18, wherein the Compound 1 Camsylate Form A is characterized by a weight loss of about 5.9% over a temperature range of 38-170 ºC in a TGA thermogram.

20. The crystalline salt form of claim 1, characterized as Compound 1 Camsylate Form B.

21. The crystalline salt form of claim 20, wherein the Compound 1 Camsylate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.25, 8.32, 8.51, 8.76, 9.05, 9.58, 9.88, 10.00, 10.52, 10.80, 11.06, 11.96, 12.24, 12.58, 13.98, 15.62, 15.81, 16.15, 16.70, 17.12, 17.42, 17.71, 18.41, 19.37, 19.55, 20.63, 21.13, 21.56, 21.89, 22.50, 23.04, 24.43, 26.08, 27.07, 27.97, 28.22, and 28.73.

22. The crystalline salt form of any one of claims 20-21, wherein the Compound 1 Camsylate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.25, 10.52, 16.15, 21.13, 21.56, and 21.89.

23. The crystalline salt form of any one of claims 20-22, wherein the Compound 1 Camsylate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.25, 10.52, 16.15, 21.13, 21.56, and 21.89.

24. The crystalline salt form of any one of claims 20-23, wherein the Compound 1 Camsylate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.25, 8.32, 8.51, 8.76, 9.05, 9.58, 9.88, 10.00, 10.52, 10.80, 11.06, 11.96, 12.24, 12.58, 13.98, 15.62, 15.81, 16.15, 16.70, 17.12, 17.42, 17.71, 18.41, 19.37, 19.55, 20.63, 21.13, 21.56, 21.89, 22.50, 23.04, 24.43, 26.08, 27.07, 27.97, 28.22, and 28.73.

25. The crystalline salt form of claim 1, characterized as Compound 1 Citrate Form A.

26. The crystalline salt form of claim 25, wherein the Compound 1 Citrate Form A is characterized by an XRPD pattern substantially identical to Figure 10.

27. The crystalline salt form of claim 1, characterized as Compound 1 Citrate Form B.

28. The crystalline salt form of claim 27, wherein the Compound 1 Citrate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.58, 5.95, 7.61, 8.88, 9.19, 9.94, 11.93, 13.26, 13.79, 14.29, 14.68, 15.26, 15.95, 16.64, 17.11, 17.87, 18.21, 18.43, 18.90, 19.36, 19.67, 20.14, 20.57, 21.47, 22.30,

~~22.64, 23.06, 23.80, 24.25, 24.55, 24.81, 25.10, 25.45, 25.54, 25.85, 26.35, 26.51, 26.83, 27.16, 27.57, 27.85, 28.08, and 28.47.~~

~~29. The crystalline salt form of any one of claims 27-28, wherein the Compound 1 Citrate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.58, 5.95, 8.88, 11.93, 13.79, 14.29, 16.64, 18.90, 21.47, 23.80, 24.25, and 25.85.~~

~~30. The crystalline salt form of any one of claims 27-29, wherein the Compound 1 Citrate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.58, 5.95, 8.88, 11.93, 13.79, 14.29, 16.64, 18.90, 21.47, 23.80, 24.25, and 25.85.~~

~~31. The crystalline salt form of any one of claims 27-30, wherein the Compound 1 Citrate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.58, 5.95, 7.61, 8.88, 9.19, 9.94, 11.93, 13.26, 13.79, 14.29, 14.68, 15.26, 15.95, 16.64, 17.11, 17.87, 18.21, 18.43, 18.90, 19.36, 19.67, 20.14, 20.57, 21.47, 22.30, 22.64, 23.06, 23.80, 24.25, 24.55, 24.81, 25.10, 25.45, 25.54, 25.85, 26.35, 26.51, 26.83, 27.16, 27.57, 27.85, 28.08, and 28.47.~~

~~32. The crystalline salt form of any one of claims 27-31, wherein the Compound 1 Citrate Form B is characterized by an endotherm with an onset temperature at about 175 ºC in a DSC thermogram.~~

~~33. The crystalline salt form of any one of claims 27-32, wherein the Compound 1 Citrate Form B is characterized by a weight loss beginning at a temperature of about 159 ºC and ending at about 220 ºC in a TGA thermogram.~~

~~34.~~8. The crystalline salt form of any one of claims ~~27~~1-~~33~~7, wherein the Compound 1 Citrate Form B is characterized by about a 4.1 wt% increase, as measured by DVS, when taken from an environment of 5% relative humidity to an environment of 95% relative humidity;

Compound 1 Maleate Form A is characterized by about a 2.5 wt% increase, as measured by DVS, when taken from an environment of 5% relative humidity to an environment of 95% relative humidity;

Compound 1 Mesylate Form C is characterized by about a 5.1 wt% increase, as measured by DVS, when taken from an environment of 5% relative humidity to an environment of 95% relative humidity;

Compound 1 Tosylate Form A is characterized by about a 0.9 wt% increase, as measured by DVS, when taken from an environment of 5% relative humidity to an environment of 95% relative humidity; and

Compound 1 Tosylate Form B is characterized by about a 0.9 wt% increase, as measured by DVS, when taken from an environment of 5% relative humidity to an environment of 95% relative humidity.

~~35. The crystalline salt form of claim 1, characterized as Compound 1 Esylate Form A.~~

~~36. The crystalline salt form of claim 35, wherein the Compound 1 Esylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 3.97, 7.95, 9.89, 10.62, 10.92, 11.03, 11.34, 11.77, 12.03, 12.53, 12.71, 13.27, 14.00, 14.70, 15.08, 15.94, 16.17, 16.92, 17.01, 17.19, 17.35, 17.61, 18.15, 18.76, 19.03, 19.66, 19.89, 20.26, 20.65, 20.78, 21.44, 22.08, 23.14, 23.27, 23.99, 24.32, 25.43, 25.66, 25.85, 26.52, 26.87, 27.61, 27.96, 28.18, 28.40, and 28.91.~~

~~37. The crystalline salt form of any one of claims 35-36, wherein the Compound 1 Esylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 3.97, 7.95, 9.89, 10.62, 13.27, 15.08, 16.17, 16.92, 17.01, 20.65, 20.78, 21.44, 22.08, 23.99, 25.43, and 26.52.~~

~~38. The crystalline salt form of any one of claims 35-37, wherein the Compound 1 Esylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 3.97, 7.95, 9.89, 10.62, 13.27, 15.08, 16.17, 16.92, 17.01, 20.65, 20.78, 21.44, 22.08, 23.99, 25.43, and 26.52.~~

~~39. The crystalline salt form of any one of claims 35-38, wherein the Compound 1 Esylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 3.97, 7.95, 9.89, 10.62, 10.92, 11.03, 11.34, 11.77, 12.03, 12.53, 12.71, 13.27, 14.00, 14.70, 15.08, 15.94, 16.17, 16.92, 17.01, 17.19, 17.35, 17.61, 18.15, 18.76, 19.03, 19.66, 19.89, 20.26, 20.65, 20.78, 21.44, 22.08, 23.14, 23.27, 23.99, 24.32, 25.43, 25.66, 25.85, 26.52, 26.87, 27.61, 27.96, 28.18, 28.40, and 28.91.~~

~~40. The crystalline salt form of any one of claims 35-39, wherein the Compound 1 Esylate Form A is characterized by a first endotherm at a temperature of about 69 ºC, a second endotherm at about 156 ºC, and a third endotherm at a temperature of about 190 ºC, in a DSC thermogram.~~

41. The crystalline salt form of any one of claims 35-40, wherein the Compound 1 Esylate Form A is characterized by a weight loss of about 8.5 wt% over a temperature range of 38-188 ^{o}C in a TGA thermogram.

42. The crystalline salt form of claim 1, characterized as Compound 1 Esylate Form B.

43. The crystalline salt form of claim 42, wherein the Compound 1 Esylate Form B is characterized by an XRPD pattern substantially identical to Figure 18.

44. The crystalline salt form of claim 1, characterized as Compound 1 HBr Form A.

45. The crystalline salt form of claim 44, wherein the Compound 1 HBr Form A is characterized by an XRPD pattern substantially identical to Figure 19.

46. The crystalline salt form of claim 1, characterized as Compound 1 HBr Form B.

47. The crystalline salt form of claim 46, wherein the Compound 1 HBr Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.26, 8.29, 9.75, 9.91, 10.56, 10.72, 12.25, 12.50, 12.75, 13.31, 13.54, 14.11, 14.72, 14.89, 15.87, 16.29, 16.50, 17.36, 17.78, 18.09, 18.44, 18.75, 19.12, 19.24, 19.56, 19.92, 20.40, 20.55, 20.69, 21.18, 22.09, 22.64, 23.24, 24.56, 25.80, 26.52, 27.23, 27.43, 28.13, 28.46, 28.76, 29.08, 29.66, and 30.07.

48. The crystalline salt form of any one of claims 46-47, wherein the Compound 1 HBr Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.26, 9.91, 10.72, 20.55, 20.69, 21.18, 22.09, 23.24, 24.56, and 25.80.

49. The crystalline salt form of any one of claims 46-48, wherein the Compound 1 HBr Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.26, 9.91, 10.72, 20.55, 20.69, 21.18, 22.09, 23.24, 24.56, and 25.80.

50. The crystalline salt form of any one of claims 46-49, wherein the Compound 1 HBr Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.26, 8.29, 9.75, 9.91, 10.56, 10.72, 12.25, 12.50, 12.75, 13.31, 13.54, 14.11, 14.72, 14.89, 15.87, 16.29, 16.50, 17.36, 17.78, 18.09, 18.44, 18.75, 19.12, 19.24, 19.56, 19.92, 20.40, 20.55, 20.69, 21.18, 22.09, 22.64, 23.24, 24.56, 25.80, 26.52, 27.23, 27.43, 28.13, 28.46, 28.76, 29.08, 29.66, and 30.07.

51. The crystalline salt form of claim 1, characterized as Compound 1 Lactate Form A.

52. The crystalline salt form of claim 51, wherein the Compound 1 Lactate Form A is characterized by an XRPD pattern substantially identical to Figure 21.

~~53. The crystalline salt form of claim 1, characterized as Compound 1 Malate Form A.~~

~~54. The crystalline salt form of claim 53, wherein the Compound 1 Malate Form A is characterized by an XRPD pattern substantially identical to Figure 22.~~

~~55. The crystalline salt form of claim 1, characterized as Compound 1 Maleate Form A.~~

~~56. The crystalline salt form of claim 55, wherein the Compound 1 Maleate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 7.58, 8.34, 10.92, 12.38, 12.87, 14.17, 15.13, 15.78, 16.43, 16.91, 17.51, 18.09, 18.71, 20.30, 21.60, 21.99, 22.30, 22.49, 23.32, 23.56, 23.92, 24.79, 26.04, 26.41, 26.92, 27.62, 28.18, 28.54, and 29.01.~~

~~57. The crystalline salt form of any one of claims 55-56, wherein the Compound 1 Maleate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 10.92, 15.13, 15.78, 16.43, 18.09, 18.71, 20.30, 21.99, 22.30, 22.49, 23.32, 23.56, 24.79, 26.41, 26.92, and 29.01.~~

~~58. The crystalline salt form of any one of claims 55-57, wherein the Compound 1 Maleate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 10.92, 15.13, 15.78, 16.43, 18.09, 18.71, 20.30, 21.99, 22.30, 22.49, 23.32, 23.56, 24.79, 26.41, 26.92, and 29.01.~~

~~59. The crystalline salt form of any one of claims 55-58, wherein the Compound 1 Maleate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 7.58, 8.34, 10.92, 12.38, 12.87, 14.17, 15.13, 15.78, 16.43, 16.91, 17.51, 18.09, 18.71, 20.30, 21.60, 21.99, 22.30, 22.49, 23.32, 23.56, 23.92, 24.79, 26.04, 26.41, 26.92, 27.62, 28.18, 28.54, and 29.01.~~

~~60. The crystalline salt form of any one of claims 55-59, wherein the Compound 1 Maleate Form A is characterized by an endotherm at a temperature of about 117 ºC in a DSC thermogram.~~

~~61. The crystalline salt form of any one of claims 55-60, wherein the Compound 1 Maleate Form A is characterized by a weight loss of about 3.1 wt% in the temperature range of 42-146 ºC in a TGA thermogram.~~

~~62. The crystalline salt form of any one of claims 55-61, wherein the Compound 1 Maleate Form A is characterized by about a 2.5 wt% increase, as measured by DVS, when taken from an environment of 5% relative humidity to an environment of 95% relative humidity.~~

~~63. The crystalline salt form of claim 1, characterized as Compound 1 Maleate Form B.~~

64. The crystalline salt form of claim 63, wherein the Compound 1 Maleate Form B is characterized by an XRPD pattern substantially identical to Figure 28.

65. The crystalline salt form of claim 1, characterized as Compound 1 Mesylate Form A.

66. The crystalline salt form of claim 65, wherein the Compound 1 Mesylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.06, 6.65, 7.66, 8.35, 10.16, 10.72, 11.98, 12.62, 13.32, 14.41, 14.79, 15.42, 15.72, 15.89, 16.23, 16.77, 17.20, 17.59, 18.08, 18.40, 18.81, 19.33, 19.53, 19.71, 20.14, 20.38, 20.73, 21.05, 21.56, 21.83, 22.16, 23.14, 24.06, 24.54, 24.77, 25.75, 26.82, 27.71, 27.94, 28.71, 29.27, and 29.85.

67. The crystalline salt form of any one of claims 65-66, wherein the Compound 1 Mesylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.06, 7.66, 14.79, 15.42, 16.23, 17.20, 17.59, 18.40, 19.33, 19.53, 20.38, 20.73, 21.83, 22.16, 23.14, 24.06, and 24.77.

68. The crystalline salt form of any one of claims 65-67, wherein the Compound 1 Mesylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.06, 7.66, 14.79, 15.42, 16.23, 17.20, 17.59, 18.40, 19.33, 19.53, 20.38, 20.73, 21.83, 22.16, 23.14, 24.06, and 24.77.

69. The crystalline salt form of any one of claims 65-68, wherein the Compound 1 Mesylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.06, 6.65, 7.66, 8.35, 10.16, 10.72, 11.98, 12.62, 13.32, 14.41, 14.79, 15.42, 15.72, 15.89, 16.23, 16.77, 17.20, 17.59, 18.08, 18.40, 18.81, 19.33, 19.53, 19.71, 20.14, 20.38, 20.73, 21.05, 21.56, 21.83, 22.16, 23.14, 24.06, 24.54, 24.77, 25.75, 26.82, 27.71, 27.94, 28.71, 29.27, and 29.85.

70. The crystalline salt form of claim 1, characterized as Compound 1 Mesylate Form B.

71. The crystalline salt form of claim 70, wherein the Compound 1 Mesylate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.14, 6.62, 7.72, 7.89, 10.33, 10.73, 12.74, 13.31, 13.88, 14.23, 14.61, 15.21, 15.52, 15.87, 16.17, 17.18, 17.41, 17.88, 18.15, 18.46, 18.73, 19.13, 19.54, 20.15, 20.53, 20.75, 20.98, 21.27, 21.60, 22.03, 22.28, 23.07, 23.32, 24.16, 24.42, 24.87, 25.13, 25.64, 26.24, 26.95, 27.17, 27.49, 27.77, 28.42, and 29.54.

72. The crystalline salt form of any one of claims 70-71, wherein the Compound 1 Mesylate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein

~~the one or more peaks is selected from 14.61, 15.21, 16.17, 17.18, 17.41, 18.46, 19.13, 19.54, 20.15, 20.53, 20.75, 20.98, 21.27, 21.60, 22.03, 23.32, 24.16, 24.42, 24.87, and 25.13.~~

~~73. The crystalline salt form of any one of claims 70-72, wherein the Compound 1 Mesylate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 14.61, 15.21, 16.17, 17.18, 17.41, 18.46, 19.13, 19.54, 20.15, 20.53, 20.75, 20.98, 21.27, 21.60, 22.03, 23.32, 24.16, 24.42, 24.87, and 25.13.~~

~~74. The crystalline salt form of any one of claims 70-73, wherein the Compound 1 Mesylate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.14, 6.62, 7.72, 7.89, 10.33, 10.73, 12.74, 13.31, 13.88, 14.23, 14.61, 15.21, 15.52, 15.87, 16.17, 17.18, 17.41, 17.88, 18.15, 18.46, 18.73, 19.13, 19.54, 20.15, 20.53, 20.75, 20.98, 21.27, 21.60, 22.03, 22.28, 23.07, 23.32, 24.16, 24.42, 24.87, 25.13, 25.64, 26.24, 26.95, 27.17, 27.49, 27.77, 28.42, and 29.54.~~

~~75. The crystalline salt form of claim 1, characterized as Compound 1 Mesylate Form C.~~

~~76. The crystalline salt form of claim 75, wherein the Compound 1 Mesylate Form C is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 7.55, 9.53, 10.25, 11.04, 11.49, 11.65, 12.24, 12.51, 13.88, 14.44, 15.36, 15.66, 16.29, 16.58, 17.16, 17.54, 19.04, 19.20, 19.79, 20.54, 21.12, 21.24, 21.62, 22.22, 23.15, 23.57, 23.99, 24.61, 24.77, 25.22, 25.61, 25.99, 26.60, 26.89, 27.29, 27.91, 29.03, and 29.29.~~

~~77. The crystalline salt form of any one of claims 75-76, wherein the Compound 1 Mesylate Form C is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 7.55, 9.53, 11.49, 12.51, 14.44, 19.04, 20.54, 21.12, 21.24, and 23.15.~~

~~78. The crystalline salt form of any one of claims 75-77, wherein the Compound 1 Mesylate Form C is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 7.55, 9.53, 11.49, 12.51, 14.44, 19.04, 20.54, 21.12, 21.24, and 23.15.~~

~~79. The crystalline salt form of any one of claims 75-78, wherein the Compound 1 Mesylate Form C is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 7.55, 9.53, 10.25, 11.04, 11.49, 11.65, 12.24, 12.51, 13.88, 14.44, 15.36, 15.66, 16.29, 16.58, 17.16, 17.54, 19.04, 19.20, 19.79, 20.54, 21.12, 21.24, 21.62, 22.22, 23.15, 23.57, 23.99, 24.61, 24.77, 25.22, 25.61, 25.99, 26.60, 26.89, 27.29, 27.91, 29.03, and 29.29.~~

80. The crystalline salt form of any one of claims 75-79, wherein the Compound 1 Mesylate Form C is characterized by a first endotherm with a peak at a temperature of about 69 °C, wherein the first endotherm has a lower limit of about 36 and an upper limit of a temperature of about 96 °C, and a second endotherm with a peak at a temperature of about 208 °C, wherein the second endotherm has a lower limit of a temperature of about 169 and an upper limit of a temperature of about 219 °C.

81. The crystalline salt form of any one of claims 75-80, wherein the Compound 1 Mesylate Form C is characterized by a weight loss of from 0.5 to 3.0 wt% in the temperature range of 36-130 ºC in a TGA thermogram.

82. The crystalline salt form of any one of claims 75-81, wherein the Compound 1 Mesylate Form C is characterized by about a 5.1 wt% increase, as measured by DVS, when taken from an environment of 5% relative humidity to an environment of 95% relative humidity.

83. The crystalline salt form of claim 1, characterized as Compound 1 Oxalate Form A.

84. The crystalline salt form of claim 83, wherein the Compound 1 Oxalate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.90, 5.25, 6.20, 8.05, 9.87, 10.54, 12.29, 14.79, 15.35, 15.84, 16.49, 16.89, 17.55, 18.27, 18.67, 19.12, 19.40, 19.81, 20.26, 20.50, 21.06, 21.74, 22.17, 22.93, 23.26, 23.50, 23.81, 24.66, 25.06, 25.95, 26.90, 27.82, and 28.24.

85. The crystalline salt form of any one of claims 83-84, wherein the Compound 1 Oxalate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.90, 5.25, 9.87, 10.54, 12.29, 15.35, 18.67, 19.40, 19.81, 20.26, 22.93, and 24.66.

86. The crystalline salt form of any one of claims 83-85, wherein the Compound 1 Oxalate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.90, 5.25, 9.87, 10.54, 12.29, 15.35, 18.67, 19.40, 19.81, 20.26, 22.93, and 24.66.

87. The crystalline salt form of any one of claims 83-86, wherein the Compound 1 Oxalate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.90, 5.25, 6.20, 8.05, 9.87, 10.54, 12.29, 14.79, 15.35, 15.84, 16.49, 16.89, 17.55, 18.27, 18.67, 19.12, 19.40, 19.81, 20.26, 20.50, 21.06, 21.74, 22.17, 22.93, 23.26, 23.50, 23.81, 24.66, 25.06, 25.95, 26.90, 27.82, and 28.24.

88. The crystalline salt form of claim 1, characterized as Compound 1 Oxalate Form B.

89. The crystalline salt form of claim 88, wherein the Compound 1 Oxalate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.20, 7.59, 8.02, 8.42, 10.53, 12.36, 12.66, 13.09, 13.35, 13.88, 14.15, 14.46, 15.06, 15.23, 17.42, 17.75, 18.28, 19.07, 19.52, 20.33, 21.18, 21.43, 21.74, 22.08, 22.59, 23.53, 24.21, 24.52, 24.71, 25.24, 25.85, 26.24, 26.55, 26.90, 27.43, 28.56, 29.30, and 29.51.

90. The crystalline salt form of any one of claims 88-89, wherein the Compound 1 Oxalate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.20, 7.59, 10.53, 15.06, 15.23, 19.52, 21.18, 21.43, 22.59, 24.21, and 25.85.

91. The crystalline salt form of any one of claims 88-90, wherein the Compound 1 Oxalate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.20, 7.59, 10.53, 15.06, 15.23, 19.52, 21.18, 21.43, 22.59, 24.21, and 25.85.

92. The crystalline salt form of any one of claims 88-91, wherein the Compound 1 Oxalate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.20, 7.59, 8.02, 8.42, 10.53, 12.36, 12.66, 13.09, 13.35, 13.88, 14.15, 14.46, 15.06, 15.23, 17.42, 17.75, 18.28, 19.07, 19.52, 20.33, 21.18, 21.43, 21.74, 22.08, 22.59, 23.53, 24.21, 24.52, 24.71, 25.24, 25.85, 26.24, 26.55, 26.90, 27.43, 28.56, 29.30, and 29.51.

93. The crystalline salt form of claim 1, characterized as Compound 1 Oxalate Form C.

94. The crystalline salt form of claim 93, wherein the Compound 1 Oxalate Form C is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.13, 7.70, 10.29, 10.59, 13.09, 13.70, 14.13, 15.33, 15.87, 16.57, 17.19, 17.50, 17.83, 18.15, 18.61, 19.24, 19.42, 20.11, 20.34, 20.64, 21.07, 21.47, 21.59, 21.91, 22.56, 22.92, 23.37, 23.87, 24.35, 25.35, 25.91, 26.47, 26.68, 27.74, 28.47, and 29.14.

95. The crystalline salt form of any one of claims 93-94, wherein the Compound 1 Oxalate Form C is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 7.70, 10.29, 10.59, 13.09, 13.70, 15.33, 17.83, 18.15, 20.11, 21.07, 21.91, and 22.56.

96. The crystalline salt form of any one of claims 93-95, wherein the Compound 1 Oxalate Form C is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 7.70, 10.29, 10.59, 13.09, 13.70, 15.33, 17.83, 18.15, 20.11, 21.07, 21.91, and 22.56.

~~97. The crystalline salt form of any one of claims 93-96, wherein the Compound 1 Oxalate Form C is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.13, 7.70, 10.29, 10.59, 13.09, 13.70, 14.13, 15.33, 15.87, 16.57, 17.19, 17.50, 17.83, 18.15, 18.61, 19.24, 19.42, 20.11, 20.34, 20.64, 21.07, 21.47, 21.59, 21.91, 22.56, 22.92, 23.37, 23.87, 24.35, 25.35, 25.91, 26.47, 26.68, 27.74, 28.47, and 29.14.~~

~~98. The crystalline salt form of claim 1, characterized as Compound 1 Propionate Form A.~~

~~99. The crystalline salt form of claim 98, wherein the Compound 1 Propionate Form A is characterized by an XRPD pattern substantially identical to Figure 38.~~

~~100. The crystalline salt form of claim 1, characterized as Compound 1 Succinate Form A.~~

~~101. The crystalline salt form of claim 100, wherein the Compound 1 Succinate Form A is characterized by an XRPD pattern substantially identical to Figure 39.~~

~~102. The crystalline salt form of claim 1, characterized as Compound 1 Sulfate Form A.~~

~~103. The crystalline salt form of claim 102, wherein the Compound 1 Sulfate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.74, 5.85, 9.47, 10.04, 11.74, 11.95, 13.77, 14.99, 15.42, 15.71, 16.48, 17.36, 17.66, 17.92, 18.59, 18.95, 19.36, 19.74, 20.22, 20.77, 21.38, 21.86, 22.74, 23.30, 23.70, 24.33, 25.07, 25.32, 25.67, 26.23, and 27.00.~~

~~104. The crystalline salt form of any one of claims 102-103, wherein the Compound 1 Sulfate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.74, 5.85, 9.47, 10.04, 11.74, 11.95, 16.48, 20.22, and 22.74.~~

~~105. The crystalline salt form of any one of claims 102-104, wherein the Compound 1 Sulfate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.74, 5.85, 9.47, 10.04, 11.74, 11.95, 16.48, 20.22, and 22.74.~~

~~106. The crystalline salt form of any one of claims 102-105, wherein the Compound 1 Sulfate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.74, 5.85, 9.47, 10.04, 11.74, 11.95, 13.77, 14.99, 15.42, 15.71, 16.48, 17.36, 17.66, 17.92, 18.59, 18.95, 19.36, 19.74, 20.22, 20.77, 21.38, 21.86, 22.74, 23.30, 23.70, 24.33, 25.07, 25.32, 25.67, 26.23, and 27.00.~~

~~107. The crystalline salt form of claim 1, characterized as Compound 1 Sulfate Form B.~~

~~108. The crystalline salt form of claim 107, wherein the Compound 1 Sulfate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or~~

more peaks is selected from 4.27, 8.57, 9.32, 10.95, 11.29, 12.07, 12.87, 13.25, 13.82, 14.13, 14.67, 16.94, 17.21, 17.64, 19.99, 20.26, 21.03, 21.44, 22.00, 22.71, 23.85, 25.20, and 26.02.

109. The crystalline salt form of any one of claims 107-108, wherein the Compound 1 Sulfate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.27, 8.57, 9.32, 16.94, 19.99, 21.03, 21.44, 22.00, and 26.02.

110. The crystalline salt form of any one of claims 107-109, wherein the Compound 1 Sulfate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.27, 8.57, 9.32, 16.94, 19.99, 21.03, 21.44, 22.00, and 26.02.

111. The crystalline salt form of any one of claims 107-110, wherein the Compound 1 Sulfate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.27, 8.57, 9.32, 10.95, 11.29, 12.07, 12.87, 13.25, 13.82, 14.13, 14.67, 16.94, 17.21, 17.64, 19.99, 20.26, 21.03, 21.44, 22.00, 22.71, 23.85, 25.20, and 26.02.

112. The crystalline salt form of claim 1, characterized as Compound 1 Sulfate Form C.

113. The crystalline salt form of claim 112, wherein the Compound 1 Sulfate Form C is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 3.19, 5.35, 9.04, 9.62, 10.73, 11.31, 11.60, 12.27, 12.67, 13.76, 14.31, 14.65, 15.21, 15.63, 16.11, 16.33, 16.63, 17.11, 18.32, 18.75, 18.99, 19.36, 19.47, 20.16, 20.55, 21.01, 21.52, 21.62, 21.89, 22.19, 22.53, 23.19, 23.79, 24.23, 24.81, 25.07, 25.41, 26.06, 26.73, 27.58, 28.16, 28.63, 28.90, 29.22, 29.47, and 29.73.

114. The crystalline salt form of any one of claims 112-113, wherein the Compound 1 Sulfate Form C is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 3.19, 5.35, 9.62, 12.27, 16.33, 17.11, 18.75, 18.99, 19.36, 19.47, 20.55, 21.01, 21.52, 21.62, 21.89, 22.19, 23.79, 25.07, and 26.06.

115. The crystalline salt form of any one of claims 112-114, wherein the Compound 1 Sulfate Form C is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 3.19, 5.35, 9.62, 12.27, 16.33, 17.11, 18.75, 18.99, 19.36, 19.47, 20.55, 21.01, 21.52, 21.62, 21.89, 22.19, 23.79, 25.07, and 26.06.

116. The crystalline salt form of any one of claims 112-115, wherein the Compound 1 Sulfate Form C is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 3.19, 5.35, 9.04, 9.62, 10.73, 11.31, 11.60, 12.27, 12.67, 13.76, 14.31, 14.65, 15.21, 15.63, 16.11, 16.33, 16.63, 17.11, 18.32, 18.75, 18.99, 19.36, 19.47, 20.16, 20.55, 21.01,

21.52, 21.62, 21.89, 22.19, 22.53, 23.19, 23.79, 24.23, 24.81, 25.07, 25.41, 26.06, 26.73, 27.58, 28.16, 28.63, 28.90, 29.22, 29.47, and 29.73.

117. The crystalline salt form of claim 1, characterized as Compound 1 Tartrate Form A.

118. The crystalline salt form of claim 117, wherein the Compound 1 Tartrate Form A is characterized by an XRPD pattern substantially identical to Figure 43.

119. The crystalline salt form of claim 1, characterized as Compound 1 Hemitartrate Form B.

120. The crystalline salt form of claim 119, wherein the Compound 1 Hemitartrate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.17, 5.66, 7.22, 9.27, 10.10, 10.40, 11.37, 11.55, 12.77, 13.94, 14.51, 15.60, 16.06, 16.48, 17.09, 17.69, 17.90, 18.27, 18.94, 19.15, 19.59, 19.81, 20.06, 20.32, 21.23, 21.83, 22.34, 22.84, 23.53, 24.26, 24.70, 25.26, 25.80, 26.78, 27.29, 27.90, 29.04, and 29.45.

121. The crystalline salt form of any one of claims 119-120, wherein the Compound 1 Hemitartrate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.17, 5.66, 10.10, 10.40, 11.37, 11.55, 14.51, 17.09, 21.23, 21.83, 23.53, 24.26, 25.26, and 26.78.

122. The crystalline salt form of any one of claims 119-121, wherein the Compound 1 Hemitartrate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.17, 5.66, 10.10, 10.40, 11.37, 11.55, 14.51, 17.09, 21.23, 21.83, 23.53, 24.26, 25.26, and 26.78.

123. The crystalline salt form of any one of claims 119-122, wherein the Compound 1 Hemitartrate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.17, 5.66, 7.22, 9.27, 10.10, 10.40, 11.37, 11.55, 12.77, 13.94, 14.51, 15.60, 16.06, 16.48, 17.09, 17.69, 17.90, 18.27, 18.94, 19.15, 19.59, 19.81, 20.06, 20.32, 21.23, 21.83, 22.34, 22.84, 23.53, 24.26, 24.70, 25.26, 25.80, 26.78, 27.29, 27.90, 29.04, and 29.45.

124. The crystalline salt form of any one of claims 119-123, wherein the Compound 1 Hemitartrate Form B is characterized by a first endotherm at a temperature of about 111 ºC, and a second endotherm at a temperature of about 148 ºC in a DSC thermogram.

125. The crystalline salt form of any one of claims 119-124, wherein the Compound 1 Hemitartrate Form B is characterized by a weight loss of about 7.4 wt% in the temperature range of 39-157 ºC in a TGA thermogram.

126. The crystalline salt form of claim 1, characterized as Compound 1 Tartrate Form C.

127. The crystalline salt form of claim 126, wherein the Compound 1 Tartrate Form C is characterized by an XRPD pattern substantially identical to Figure 69.

128. The crystalline salt form of claim 1, characterized as Compound 1 Tosylate Form A.

129. The crystalline salt form of claim 128, wherein the Compound 1 Tosylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 5.81, 7.05, 8.52, 9.74, 10.40, 10.99, 11.26, 11.66, 12.58, 12.81, 14.28, 14.62, 15.30, 15.80, 16.03, 16.34, 17.12, 17.55, 18.03, 18.94, 19.26, 19.67, 20.31, 20.65, 21.02, 21.43, 22.11, 22.44, 22.66, 22.94, 23.07, 23.54, 23.79, 24.55, 24.88, 25.42, 25.83, 26.10, 26.53, 27.70, 28.11, 28.62, 29.18, 29.40, 29.66, and 29.83.

130. The crystalline salt form of any one of claims 128-129, wherein the Compound 1 Tosylate Form A is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 8.52, 9.74, 14.28, 16.03, 17.55, 18.94, 19.26, 20.31, 21.02, 21.43, 22.11, 22.44, 22.66, 23.07, 24.55, and 27.70.

131. The crystalline salt form of any one of claims 128-130, wherein the Compound 1 Tosylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 8.52, 9.74, 14.28, 16.03, 17.55, 18.94, 19.26, 20.31, 21.02, 21.43, 22.11, 22.44, 22.66, 23.07, 24.55, and 27.70.

132. The crystalline salt form of any one of claims 128-131, wherein the Compound 1 Tosylate Form A is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 5.81, 7.05, 8.52, 9.74, 10.40, 10.99, 11.26, 11.66, 12.58, 12.81, 14.28, 14.62, 15.30, 15.80, 16.03, 16.34, 17.12, 17.55, 18.03, 18.94, 19.26, 19.67, 20.31, 20.65, 21.02, 21.43, 22.11, 22.44, 22.66, 22.94, 23.07, 23.54, 23.79, 24.55, 24.88, 25.42, 25.83, 26.10, 26.53, 27.70, 28.11, 28.62, 29.18, 29.40, 29.66, and 29.83.

133. The crystalline salt form of any one of claims 128-132, wherein the Compound 1 Tosylate Form A is characterized by an endotherm with an onset temperature at about 214 ºC in a DSC thermogram.

134. The crystalline salt form of any one of claims 128-133, wherein the Compound 1 Tosylate Form A is characterized by a weight loss of about 0.9 wt% in the temperature range of 72-231 ºC in a TGA thermogram.

135. The crystalline salt form of any one of claims 128-134, wherein the Compound 1 Tosylate Form A is characterized by about a 0.9 wt% increase, as measured by DVS, when taken from an environment of 5% relative humidity to an environment of 95% relative humidity.

~~136. The crystalline salt form of claim 1, characterized as Compound 1 Tosylate Form B.~~

~~137. The crystalline salt form of claim 136, wherein the Compound 1 Tosylate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.61, 8.69, 9.25, 10.68, 12.44, 12.69, 13.76, 14.42, 14.81, 15.47, 16.08, 16.57, 17.45, 17.76, 18.44, 18.83, 19.01, 19.35, 20.42, 20.84, 21.15, 21.69, 22.09, 22.86, 23.12, 24.08, 24.53, 24.86, 25.22, 25.91, 26.33, 26.82, 27.63, 28.61, 29.11, and 29.77.~~

~~138. The crystalline salt form of any one of claims 136-137, wherein the Compound 1 Tosylate Form B is characterized by one or more peaks in an XRPD pattern on a 2 Theta scale, wherein the one or more peaks is selected from 4.61, 9.25, 12.44, 15.47, 17.76, 19.01, 19.35, 25.91, 26.33, and 27.63.~~

~~139. The crystalline salt form of any one of claims 136-138, wherein the Compound 1 Tosylate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.61, 9.25, 12.44, 15.47, 17.76, 19.01, 19.35, 25.91, 26.33, and 27.63.~~

~~140. The crystalline salt form of any one of claims 136-139, wherein the Compound 1 Tosylate Form B is characterized by all of the following peaks in an XRPD pattern on a 2 Theta scale, selected from 4.61, 8.69, 9.25, 10.68, 12.44, 12.69, 13.76, 14.42, 14.81, 15.47, 16.08, 16.57, 17.45, 17.76, 18.44, 18.83, 19.01, 19.35, 20.42, 20.84, 21.15, 21.69, 22.09, 22.86, 23.12, 24.08, 24.53, 24.86, 25.22, 25.91, 26.33, 26.82, 27.63, 28.61, 29.11, and 29.77.~~

~~141. The crystalline salt form of any one of claims 136-140, wherein the Compound 1 Tosylate Form B is characterized by a sharp endotherm with an onset temperature at ~183 °C in a DSC thermogram.~~

~~142. The crystalline salt form of any one of claims 136-141, wherein the Compound 1 Tosylate Form B is characterized by a weight loss of about 0.7 wt% in the temperature range of 39-214 ºC in a TGA thermogram.~~

~~143. The crystalline salt form of any one of claims 136-142, wherein the Compound 1 Tosylate Form B is characterized by about a 0.9 wt% increase, as measured by DVS, when taken from an environment of 5% relative humidity to an environment of 95% relative humidity.~~

~~144.~~9. A pharmaceutical composition comprising a crystalline salt form of any one of claims 1-~~143~~ 8 and a pharmaceutically acceptable excipient.

10. A crystalline salt form of any one of claims 1-8, or a pharmaceutical composition of claim 9, for use in therapy.

~~145.~~11. Use of a crystalline salt form of any one of claims 1-8, or a pharmaceutical composition of claim 9, in the manufacture of a medicament for ~~A method of~~ treating a disease, disorder, or syndrome mediated at least in part by modulating *in vivo* activity of a protein kinase~~, comprising administering to a subject in need thereof a crystalline salt form of any one of claims 1-143, or a pharmaceutical composition of claim 144~~.

~~146.~~12. The ~~method~~ use of claim ~~145~~10, wherein the disease, disorder, or syndrome mediated at least in part by modulating *in vivo* activity of a protein kinase is cancer.

~~147.~~13. Use of a crystalline salt form of any one of claims 1-8, in the manufacture of a medicament ~~A method~~ for inhibiting a protein kinase~~, the method comprising contacting the protein kinase with a crystalline salt form of any one of claims 1-143~~.

~~148.~~14. The ~~method~~ use of any of claims ~~145~~10-~~147~~12, wherein the protein kinase is Axl, Mer, c-Met, KDR, or a combination thereof.