

DESCRIPTION

ALIPHATIC ACID AMIDE DERIVATIVE

5 TECHNICAL FIELD

[0001]

The present invention relates to an aliphatic acid amide derivative having antagonist activity for the serotonin 5-HT_{2A} receptor and the serotonin 5-HT₇ receptor, or a pharmaceutically acceptable salt thereof, and a medicament for treating a
10 neuropsychiatric disease comprising the same as an active ingredient.

BACKGROUND ART

15 [0002]

The serotonin (5-hydroxytryptamine; hereinafter, also referred to as "5-HT") is known as one of main neurotransmitters in central nervous system, and it is also known that serotonin is involved in various brain functions
20 such as emotional reaction and cognitive function.

[0003]

The 5-HT_{2A} receptor which is one of 5-HT receptor subtypes is a Gq/11 protein-coupled receptor, and is highly expressed in, for example, cerebral cortex, hippocampus, and raphe
25 nucleus. Drugs having antagonist activity for the 5-HT_{2A} receptor include antidepressant drugs, mianserin and mirtazapine. Atypical antipsychotic drugs which have

antagonist activity for the 5-HT_{2A} receptor are used as a medicament for treating diseases such as schizophrenia, bipolar disorder, major depression, and autistic spectrum disorder (Non-Patent Literature 1, Non-Patent Literature 2).

5 [0004]

The 5-HT₇ receptor is a Gs protein-coupled receptor, and is extensively expressed in, for example, hypothalamus, thalamus, hippocampus, and raphe nucleus (Non-patent Literature 9).

Drugs having antagonist activity for the 5-HT₇ receptor include
10 Lurasidone which is used as a medicament for treating schizophrenia and a bipolar disorder and Vortioxetine which is used for treating major depression. There is, however, no commercial drug having selective antagonist activity for the 5-HT₇ receptor. Lurasidone is used for treating schizophrenia and
15 a bipolar disorder, and Vortioxetine is used for treating major depression (Non-patent Literature 3, Non-patent Literature 4, and Non-patent Literature 5). It is also known that in some animal models, antagonizing the 5-HT₇ receptor leads to antidepressant and anxiolytic effect, and improvement effect in
20 cognition function (Non-patent Literature 6, Non-patent Literature 7). It is also known that 5-HT₇ receptor deficient mice exhibit antidepressant effect (Non-patent Literature 8).

[0005]

As described above, it is shown that antagonists for the
25 5-HT_{2A} receptor and the 5-HT₇ receptor are separately useful for many neuropsychiatric diseases, but no drugs having antagonist activity for the 5-HT_{2A} receptor and the 5-HT₇ receptor in a

selective and potent manner have been reported.

[0006]

CITATION LIST

NON PATENT LITERATURE

5 [Non Patent Literature 1] P. Seeman, Can. J. Psychiatry. 4: 27-38, 2002

[Non Patent Literature 2] C. J. Schmidt, Life Science. 56(25): 2209-2222, 1995

10 [Non Patent Literature 3] L. Citrome, J. Clinical Practice. 65(2): 189-210, 2011

[Non Patent Literature 4] YS. Woo, Neuropsychiatric Disease and Treatment. 9: 1521-1529, 2013

[Non Patent Literature 5] C. Sanchez, Pharmacology & Therapeutics. 145: 43-57, 2015

15 [Non Patent Literature 6] Bonaventure P, J Pharmacol Exp Ther. 321: 690-8, 2007

[Non Patent Literature 7] Horisawa T, Behavioural Brain Research. 220: 83-90, 2011

20 [Non Patent Literature 8] M. Guscott, Neuropharmacology. 48: 492-502, 2005

[Non Patent Literature 9] T W. Lovenberg, Neuron. 11: 449-458, 1993

SUMMARY OF INVENTION

25 PROBLEMS TO BE SOLVED BY THE INVENTION

[0007]

An object of the present invention is the provision of

novel compounds which have antagonist activity for both of the serotonin 5-HT_{2A} receptor and the 5-HT₇ receptor, and are useful as a medicament for treating neuropsychiatric diseases.

5 MEANS OF SOLVING THE PROBLEMS


[0008]

The present inventors have extensively studied to accomplish the above object, and then found that a compound of Formula (1) as below, or a pharmaceutically acceptable salt thereof (hereinafter, also referred to as "the present compound") has antagonist activity for both of the serotonin 5-HT_{2A} receptor and the 5-HT₇ receptor. Based upon these new findings, the present invention has been achieved.

15 [0009]

The present invention is illustrated as follows.

[Item 1] A compound of Formula (1):



20 or a pharmaceutically acceptable salt thereof,
wherein

Z is a nitrogen atom or -CR^A-;

Y is carbonyl or sulfonyl;

m is 1, 2, 3, or 4;

n is 0, 1, 2, or 3;

provided that when Y is sulfonyl, n is not 0;

R^A is hydrogen, hydroxy, C₁₋₆ alkyl, or C₁₋₆ alkoxy;

5 R^{1a}, R^{1b}, R^{1c}, and R^{1d} are each independently hydrogen, hydroxy, halogen, or C₁₋₆ alkyl optionally-substituted with the same or different 1 to 3 halogen;

R^{2a}, R^{2b}, R^{2c}, and R^{2d} are each independently, and each of R^{2cs} or each of R^{2ds} is independently when R^{2c} or R^{2d} exists plurally,
10 hydrogen, halogen, or C₁₋₆ alkyl (wherein the alkyl may be optionally substituted with the same or different 1 to 5 substituents selected from the group consisting of halogen, hydroxy, C₃₋₈ cycloalkyl, and C₁₋₆ alkoxy), provided that when R^{2a} and R^{2b}, or R^{2c} and R^{2d} which attach to the same carbon atom are
15 each independently the said C₁₋₆ alkyl, they may be combined together with the carbon atom to which they attach to form a 3- to 6-membered saturated carbocycle;


R³ is C₁₋₆ alkyl, wherein the alkyl may be optionally substituted with the same or different 1 to 5 substituents
20 selected from the group consisting of halogen, hydroxy, cyano, C₃₋₈ cycloalkyl, and C₁₋₆ alkoxy;

R^{4a}, R^{4b}, R^{4c}, and R^{4d} are each independently, and each of R^{4cs} or each of R^{4ds} is independently when R^{4c} or R^{4d} exists plurally,
hydrogen, halogen, C₁₋₆ alkyl (wherein the alkyl may be optionally
25 substituted with the same or different 1 to 5 substituents selected from the group consisting of halogen, hydroxy, cyano, C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkyl ester, and amino optionally-

substituted with the same or different 1 or 2 C₁₋₆ alkyl (wherein the alkyl may be optionally substituted with oxo)), C₂₋₆ alkynyl, or amino (wherein the amino may be optionally substituted with the same or different 1 or 2 C₁₋₆ alkyl); provided that when any
 5 two of R^{4a}, R^{4b}, R^{4c}, and R^{4d} are each independently the said C₁₋₆ alkyl, they may be combined together with the carbon atom or the carbon atoms to which they attach to form a 3- to 6-membered saturated carbocycle or a 4- to 6-membered saturated heterocycle wherein when the said C₁₋₆ alkyl groups have a substituent, the
 10 substituent may be included as a ring member of the carbocycle or heterocycle;

provided that when Y is sulfonyl and n is 1, any one of R^{4c} or R^{4d} is hydrogen; and when Y is sulfonyl and n is 2 or 3, any one of R^{4c} or R^{4d} which binds to the carbon atom adjacent to Y is
 15 hydrogen; and

Ring Q is a group of the following Formula (2a), (2b), (2c), (2d), (2e), or (2f):



wherein

R^{5a} , R^{5b} , R^{5c} , and R^{5d} are each independently hydrogen, halogen, cyano, C_{1-6} alkyl, C_{1-6} alkoxy (wherein the alkyl and the alkoxy may be each independently optionally substituted with the same or different 1 to 3 halogen), or amino optionally-substituted with the same or different 1 or 2 C_{1-6} alkyl; and

R^6 is hydrogen, C_{1-6} alkyl, or C_{3-8} cycloalkyl, wherein the alkyl and the cycloalkyl may be each independently optionally substituted with the same or different 1 to 3 halogen.

10

[0010]

[Item 2] The compound of Item 1, or a pharmaceutically acceptable salt thereof, wherein R^{1a} , R^{1b} , R^{1c} , and R^{1d} are hydrogen.

15 [0011]

[Item 3] The compound of Item 1 or 2, or a pharmaceutically acceptable salt thereof, wherein R^{2a} , R^{2b} , R^{2c} , and R^{2d} are hydrogen.

[0012]

20 [Item 4] The compound of any one of Items 1 to 3, or a pharmaceutically acceptable salt thereof, wherein m is 1.

[0013]

[Item 5] The compound of any one of Items 1 to 4, or a pharmaceutically acceptable salt thereof, wherein R^3 is C_{1-6} alkyl.

25

[0014]

[Item 6] The compound of any one of Items 1 to 5, or a pharmaceutically acceptable salt thereof, wherein Z is $-CR^A-$.

[0015]

5 [Item 7] The compound of any one of Items 1 to 6, or a pharmaceutically acceptable salt thereof, wherein R^A is hydrogen.

[0016]

10 [Item 8] The compound of any one of Items 1 to 7, or a pharmaceutically acceptable salt thereof, wherein R^{5a} , R^{5b} , R^{5c} , and R^{5d} are each independently hydrogen, halogen, or C_{1-6} alkyl.

[0017]

15 [Item 9] The compound of any one of Items 1 to 8, or a pharmaceutically acceptable salt thereof, wherein R^{5a} , R^{5c} , and R^{5d} are hydrogen.

[0018]

20 [Item 10] The compound of any one of Items 1 to 9, or a pharmaceutically acceptable salt thereof, wherein R^{5b} is hydrogen, halogen, or C_{1-6} alkyl.

[0019]

25 [Item 11] The compound of any one of Items 1 to 10, or a pharmaceutically acceptable salt thereof, wherein R^6 is hydrogen.

[0020]

[Item 12] The compound of any one of Items 1 to 11, or a pharmaceutically acceptable salt thereof, wherein R^{4a} , R^{4b} , R^{4c} , and R^{4d} are each independently, and each of R^{4c} s or each of R^{4d} s is independently when R^{4c} or R^{4d} exists plurally, hydrogen, halogen, or C_{1-6} alkyl, wherein the alkyl may be optionally substituted with the same or different 1 to 5 substituents selected from the group consisting of halogen, hydroxy, and C_{1-6} alkoxy.

[0021]

[Item 13] The compound of any one of Items 1 to 12, or a pharmaceutically acceptable salt thereof, wherein n is 0 or 1.

[0022]


[Item 14] The compound of any one of Items 1 to 13, or a pharmaceutically acceptable salt thereof, wherein Y is carbonyl.

[0023]

[Item 15] The compound of any one of Items 1 to 13, or a pharmaceutically acceptable salt thereof, wherein Y is sulfonyl.

[0024]

[Item 16] The compound of Item 1, or a pharmaceutically acceptable salt thereof, wherein the compound is selected from the group consisting of:



[0025]

[Item 17] A drug comprising a compound of any one of Items 1 to 16, or a pharmaceutically acceptable salt thereof, as an active ingredient.

[0026]

[Item 18] A medicament for treating a mental disease or central nervous system disease, comprising a compound of any one of Items 1 to 16, or a pharmaceutically acceptable salt thereof, as an active ingredient.

[0027]

[Item 19] The medicament of Item 18, wherein the mental disease

or central nervous system disease is schizophrenia, a bipolar disorder, a sleep disorder, an autism spectrum disorder, major depression, treatment-resistant depression, or a psychopathic symptom or dementia associated with Alzheimer's disease or
5 Parkinson's disease.

[0028]

[Item 20] A method for treating a mental disease or central nervous system disease, comprising administering a
10 therapeutically effective amount of a compound of any one of Items 1 to 16, or a pharmaceutically acceptable salt thereof, to a patient in need thereof.

[0029]

15 [Item 21] Use of a compound of any one of Items 1 to 16, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating a mental disease or central nervous system disease.

20 [0030]

[Item 22] A compound of any one of Items 1 to 16, or a pharmaceutically acceptable salt thereof, for use in the treatment of a mental disease or central nervous system disease.

25 [0031]

[Item 23] A medicament for treating a mental disease or central nervous system disease, comprising a compound of any one of Items

1 to 16, or a pharmaceutically acceptable salt thereof, in combination with at least one drug selected from the group consisting of a drug for treating a developmental disorder such as an autism spectrum disorder and an attention deficit hyperactivity disorder, an antipsychotic drug and a schizophrenic drug, a drug for treating a bipolar disorder, an antidepressant drug, an antianxiety drug, a drug for treating an obsessive-compulsive disorder, a drug for treating a stress disorder such as post-traumatic stress disorder, a drug for treating a mood disorder, a drug for treating an eating disorder, a drug for treating a sleep disorder such as insomnia, narcolepsy, a sleep apnea syndrome, and a circadian rhythm disorder, a drug for treating a sexual dysfunction, a drug for treating drug dependence, a drug for treating dementia such as Alzheimer's disease, a drug for treating a behavioral and psychological symptom associated with dementia, a cerebral metabolism and circulation improving drug, a drug for treating a movement disorder such as Parkinson's disease, an analgesic drug, an antiepileptic drug, an anticonvulsant, a migraine drug, an anesthetic, and a central stimulant.

[0032]

[Item 24] A medicament for treatment of a mental disease or central nervous system disease, comprising a compound of any one of Items 1 to 16, or a pharmaceutically acceptable salt thereof, as an active ingredient, wherein the medicament is used in the treatment concomitantly with at least one drug selected from the

group consisting of a drug for treating a developmental disorder such as an autism spectrum disorder and an attention deficit hyperactivity disorder, an antipsychotic drug and a schizophrenic drug, a drug for treating a bipolar disorder, an antidepressant drug, an antianxiety drug, a drug for treating an obsessive-compulsive disorder, a drug for treating a stress disorder such as a post-traumatic stress disorder, a drug for treating a mood disorder, a drug for treating an eating disorder, a drug for treating a sleep disorder such as insomnia, narcolepsy, a sleep apnea syndrome, and a circadian rhythm disorder, a drug for treating a sexual dysfunction, a drug for treating drug dependence, a drug for treating dementia such as Alzheimer's disease, a drug for treating a behavioral and psychological symptom associated with dementia, a cerebral metabolism and circulation improving drug, a drug for treating a movement disorder such as Parkinson's disease, an analgesic drug, an antiepileptic drug, an anticonvulsant, a migraine drug, an anesthetic, and a central stimulant.

[0033]

[Item 25] A crystal of a compound of any one of Examples 17, 77, 78, 79, and 80, characterized by powder XRD patterns comprising powder XRD peaks of 4 or more (preferably 10 or more) $2\theta \pm 0.2$ values selected from those measured in each of the Examples.

EFFECT OF THE INVENTION

[0034]

The present compound has antagonist activity for the 5-HT_{2A} receptor and the 5-HT₇ receptor. The present compound is useful as a medicament for treating a neuropsychiatric disease and a central nervous system disease.

5

BRIEF DESCRIPTION OF DRAWINGS

[0035]

Fig. 1 shows results of fear conditioning test using the compound of Example 17 (Test 6).

10

Fig. 2 shows results of measurement of the amount of released glutamic acid in rat brain using the compound of Example 17 (Test 7).

Fig. 3 shows a chart of powder XRD measurement of the compound of Example 78.

15

DESCRIPTION OF EMBODIMENTS

[0036]

Hereinafter, the present invention is described in detail. In the description, the number of carbon atoms in a definition of "substituent" may be indicated as, for example, "C₁₋₆". Specifically, the term "C₁₋₆ alkyl" means an alkyl group having 1 to 6 carbon atoms.

20

[0037]

The "halogen" used herein includes, for example, fluorine atom, chlorine atom, bromine atom, and iodine atom. Preferably, it includes fluorine atom and chlorine atom.

25

[0038]

The "C₁₋₆ alkyl" used herein means a straight- or branched-chain saturated hydrocarbon group having 1 to 6 carbon atoms. Preferably, the "C₁₋₆ alkyl" includes "C₁₋₄ alkyl", and more preferably, it includes "C₁₋₃ alkyl". The "C₁₋₃ alkyl" includes, 5 for example, methyl, ethyl, propyl, and 1-methylethyl. The "C₁₋₄ alkyl" includes, for example, butyl, 1,1-dimethylethyl, 1-methylpropyl, and 2-methylpropyl, besides the above-listed examples of the "C₁₋₃ alkyl". The "C₁₋₆ alkyl" includes, for example, pentyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1- 10 methylbutyl, 2-methylbutyl, 4-methylpentyl, 3-methylpentyl, 2-methylpentyl, 1-methylpentyl, and hexyl, besides the above-listed examples of the "C₁₋₄ alkyl".

The phrase "alkyl is substituted with oxo" used herein means that a keto group (C=O) is formed with any one of carbon atoms 15 of which the alkyl consists.

[0039]

The "C₃₋₈ cycloalkyl" used herein means a cyclic saturated hydrocarbon group having 3 to 8 carbon atoms, and it includes those which have a partially-unsaturated bond or a bridged 20 structure. The "C₃₋₈ cycloalkyl" preferably includes "C₃₋₆ cycloalkyl". The "C₃₋₆ cycloalkyl" includes, for example, cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl. The "C₃₋₈ cycloalkyl" includes, for example, cycloheptyl and cyclooctyl, besides the above-listed examples of the "C₃₋₆ cycloalkyl".

25 [0040]

The "C₁₋₆ alkoxy" is used interchangeably with "C₁₋₆ alkyloxy" herein, and the "C₁₋₆ alkyl" part is the same as the above "C₁₋₆

alkyl". The "C₁₋₆ alkoxy" preferably includes "C₁₋₄ alkoxy", and more preferably "C₁₋₃ alkoxy". The "C₁₋₃ alkoxy" includes, for example, methoxy, ethoxy, propoxy, and 1-methylethoxy. The "C₁₋₄ alkoxy" includes, for example, butoxy, 1,1-dimethylethoxy, 1-methylpropoxy, and 2-methylpropoxy, besides the above-listed examples of the "C₁₋₃ alkoxy". The "C₁₋₆ alkoxy" includes, for example, pentyloxy, 1,1-dimethylpropoxy, 1,2-dimethylpropoxy, 1-methylbutoxy, 2-methylbutoxy, 4-methylpentyloxy, 3-methylpentyloxy, 2-methylpentyloxy, 1-methylpentyloxy, and hexyloxy, besides the above-listed examples of the "C₁₋₄ alkoxy".

[0041]

The "C₁₋₆ alkyl ester" used herein means an ester (-COOR') of which R' part is the above "C₁₋₆ alkyl". The "C₁₋₆ alkyl ester" preferably includes "C₁₋₄ alkyl ester", and more preferably "C₁₋₃ alkyl ester". The "C₁₋₃ alkyl ester" includes, for example, methyl ester, ethyl ester, propyl ester, and 1-methylethyl ester. The "C₁₋₄ alkyl ester" includes, for example, butyl ester, 1,1-dimethylethyl ester, 1-methylpropyl ester, and 2-methylpropyl ester, besides the above-listed examples of the "C₁₋₃ alkyl ester".

The "C₁₋₆ alkyl ester" includes, for example, pentyl ester, 1,1-dimethylpropyl ester, 1,2-dimethylpropyl ester, 1-methylbutyl ester, 2-methylbutyl ester, 4-methylpentyl ester, 3-methylpentyl ester, 2-methylpentyl ester, 1-methylpentyl ester, and hexyl ester, besides the above-listed examples of the "C₁₋₄ alkyl ester".

[0042]

The "C₂₋₆ alkynyl" used herein means a straight- or branched-chain unsaturated hydrocarbon group having a triple bond and 2

to 6 carbon atoms. The "C₂₋₆ alkynyl" preferably includes "C₂₋₄ alkynyl", and more preferably "C₂₋₃ alkynyl". The "C₂₋₃ alkynyl" includes, for example, ethynyl and propynyl. The "C₂₋₄ alkynyl" includes, for example, butynyl, besides the above-listed examples
5 of the "C₂₋₃ alkynyl". The "C₂₋₆ alkynyl" includes, for example, pentynyl and hexynyl, besides the above-listed examples of the "C₂₋₄ alkynyl".

[0043]

The "3- to 6-membered saturated carbocycle" used herein
10 means a cyclic saturated hydrocarbon group having 3 to 6 carbon atoms that includes those which have a partially-unsaturated bond or a bridged structure. The "3- to 6-membered saturated carbocycle" preferably includes "5- or 6-membered monocyclic saturated carbocycle". The "5- or 6-membered monocyclic
15 saturated carbocycle" includes, for example, cyclopentane and cyclohexane. The "3- to 6-membered saturated carbocycle" includes, for example, cyclopropane and cyclobutane, besides the above-listed examples of the "5- or 6-membered monocyclic saturated carbocycle".

20 [0044]

The "4- to 6-membered saturated heterocycle" group means a saturated ring composed of 4 to 6 atoms comprising 1 or 2 atoms independently selected from the group consisting of nitrogen atom, oxygen atom, and sulfur atom, besides carbon atoms. The "4- to
25 6-membered saturated heterocycle" group includes those which have a partially-unsaturated bond or a bridged structure. The "4- to 6-membered saturated heterocycle" group preferably includes "5-

or 6-membered monocyclic saturated heterocycle" group. The "5-
 or 6-membered monocyclic saturated heterocycle" group includes,
 for example, tetrahydrofuryl, pyrrolidinyl, imidazolidinyl,
 piperidinyl, morpholinyl, thiomorpholinyl, dioxothiomorpholinyl,
 5 hexamethyleneiminyl, oxazolidinyl, thiazolidinyl,
 oxoimidazolidinyl, dioxoimidazolidinyl, oxooxazolidinyl,
 dioxooxazolidinyl, dioxothiazolidinyl, tetrahydrofuranyl,
 tetrahydropyranyl. The "4- to 6-membered saturated heterocycle"
 group includes, for example, oxetanyl and azetidiny, besides the
 10 above-listed examples of the "5- or 6-membered monocyclic
 saturated heterocycle" group.

[0045]

In a compound of Formula (1), preferred examples of Z, Y, m,
 n, R^A, R^{1a}, R^{1b}, R^{1c}, R^{1d}, R^{2a}, R^{2b}, R^{2c}, R^{2d}, R³, R^{4a}, R^{4b}, R^{4c}, R^{4d}, Q,
 15 R^{5a}, R^{5b}, R^{5c}, R^{5d}, and R⁶ are shown below, but the scope of the
 present invention is not limited to the scope of the following
 compounds.

[0046]

One embodiment of Z includes -CR^A-. Another embodiment of
 20 Z includes a nitrogen atom.

[0047]

One embodiment of Y includes carbonyl. Another embodiment
 of Y includes sulfonyl.

[0048]

25 A preferred embodiment of m includes 1 and 2. A more
 preferred embodiment of m includes 1.

[0049]

A preferred embodiment of n includes 0, 1, and 2. A more preferred embodiment of n includes 0 and 1. An especially preferred embodiment of n includes 1.

[0050]

5 A preferred embodiment of R^A includes hydrogen, hydroxy, and C_{1-6} alkyl. A more preferred embodiment includes hydrogen and C_{1-3} alkyl. A further preferred embodiment includes hydrogen, ethyl, and methyl. The most preferred embodiment includes hydrogen.

[0051]

10 Preferred embodiments of R^{1a} , R^{1b} , R^{1c} , and R^{1d} include, each independently, hydrogen, hydroxy, halogen, and C_{1-6} alkyl. More preferred embodiments include hydrogen, halogen, and C_{1-3} alkyl. Further preferred embodiments include hydrogen, fluorine atom, chlorine atom, ethyl, and methyl. Especially preferred
15 embodiments include hydrogen, fluorine atom, and methyl. The most preferred embodiments include hydrogen.

[0052]

Preferred embodiments of R^{2a} , R^{2b} , R^{2c} , and R^{2d} include, each independently, hydrogen, halogen, and C_{1-6} alkyl. More preferred
20 embodiments include hydrogen, halogen, and C_{1-3} alkyl. Further preferred embodiments include hydrogen, fluorine atom, chlorine atom, ethyl, and methyl. Especially preferred embodiments include hydrogen, fluorine atom, and methyl. The most preferred embodiments include hydrogen.

25 [0053]

A preferred embodiment of R^3 includes C_{1-6} alkyl. More preferred embodiment includes C_{1-3} alkyl. A further preferred

embodiment includes ethyl and methyl. An especially preferred embodiment includes ethyl.

[0054]

Preferred embodiments of R^{4a} , R^{4b} , R^{4c} , and R^{4d} include, each
5 independently, hydrogen, halogen, C_{1-6} alkyl, C_{2-6} alkynyl, and
amino optionally-substituted with the same or different 1 or 2
 C_{1-6} alkyl. More preferred embodiments include hydrogen, halogen,
 C_{1-6} alkyl, and amino optionally-substituted with the same or
different 1 or 2 C_{1-6} alkyl. Further preferred embodiments include
10 hydrogen, halogen, and C_{1-6} alkyl. Especially preferred
embodiments include hydrogen and C_{1-6} alkyl. Especially preferred
embodiments include hydrogen, methyl, and methoxymethyl. The
most preferred embodiments include hydrogen.

[0055]

15 A preferred embodiment of Q includes (2a), (2c), (2d), (2e),
and (2f). A more preferred embodiment includes (2a), (2d), and
(2e). An especially preferred embodiment includes (2a).

[0056]

Preferred embodiments of R^{5a} , R^{5b} , R^{5c} , and R^{5d} include, each
20 independently, hydrogen, halogen, cyano, C_{1-6} alkyl, C_{1-6} alkoxy,
and amino optionally-substituted with the same or different 1 or
2 C_{1-6} alkyl. More preferred embodiments include hydrogen, halogen,
cyano, C_{1-6} alkyl, and C_{1-6} alkoxy. Further preferred embodiments
include hydrogen, fluorine atom, and chlorine atom. Especially
25 preferred embodiments include hydrogen and fluorine atom.

[0057]

A preferred embodiment of R^6 includes hydrogen, C_{1-6} alkyl,

and C₃₋₈ cycloalkyl. A more preferred embodiment includes hydrogen and C₁₋₆ alkyl. A further preferred embodiment includes hydrogen and C₁₋₃ alkyl. An especially preferred embodiment includes hydrogen.

5 [0058]

One embodiment of a compound of Formula (1) includes the following embodiment (A):

(A) A compound, or a pharmaceutically acceptable salt thereof, wherein

10 Z is a nitrogen atom or -CR^A-;

Y is carbonyl or sulfonyl;

m is 1 or 2;

n is 0, 1, 2, or 3;

provided that when Y is sulfonyl, n is not 0;

15 R^A is hydrogen or C₁₋₆ alkyl;

R^{1a}, R^{1b}, R^{1c}, and R^{1d} are each independently hydrogen or C₁₋₆ alkyl;

R^{2a}, R^{2b}, R^{2c}, and R^{2d} are each independently, and each of R^{2c}s or each of R^{2d}s is independently when R^{2c} or R^{2d} exists plurally, hydrogen or C₁₋₆ alkyl;


20 R³ is C₁₋₆ alkyl;

R^{4a}, R^{4b}, R^{4c}, and R^{4d} are each independently, and each of R^{4c}s or each of R^{4d}s is independently when R^{4c} or R^{4d} exists plurally, hydrogen, halogen, C₁₋₆ alkyl (wherein the alkyl may be optionally substituted with the same or different 1 to 5 substituents selected from the group consisting of halogen, hydroxy, C₁₋₆ alkoxy, C₁₋₆ alkyl ester, and amino optionally-substituted with the same

or different 1 or 2 C₁₋₆ alkyl (wherein the alkyl may be optionally substituted with oxo)), C₂₋₆ alkynyl, or amino (wherein the amino may be optionally substituted with the same or different 1 or 2 C₁₋₆ alkyl); provided that when any two of R^{4a}, R^{4b}, R^{4c}, and R^{4d} are each independently the above C₁₋₆ alkyl, they may be combined together with the carbon atom or the carbon atoms to which they attach to form a 3- to 6-membered saturated carbocycle or a 4- to 6-membered saturated heterocycle wherein when the C₁₋₆ alkyl groups have a substituent, the substituent may be included as a ring member of the carbocycle or heterocycle;

provided that when Y is sulfonyl and n is 1, either R^{4c} or R^{4d} is hydrogen; and when Y is sulfonyl and n is 2 or 3, either R^{4c} or R^{4d} that binds to the carbon atom adjacent to Y is hydrogen; and

Ring Q is a group of the following Formula (2a), (2b), (2c), (2d), (2e), or (2f):



wherein

R^{5a}, R^{5b}, R^{5c}, and R^{5d} are each independently hydrogen,

halogen, or C₁₋₆ alkyl; and

R⁶ is hydrogen or C₁₋₆ alkyl.

[0059]

Another embodiment of a compound of Formula (1) includes the
5 following embodiment (B):

(B) A compound, or a pharmaceutically acceptable salt thereof,
wherein

Z is a nitrogen atom or -CR^A-;

Y is carbonyl;

10 m is 1 or 2;

n is 0, 1, 2, or 3;

R^A is hydrogen or C₁₋₆ alkyl;

R^{1a}, R^{1b}, R^{1c}, and R^{1d} are each independently hydrogen or C₁₋₆
alkyl;


15 R^{2a}, R^{2b}, R^{2c}, and R^{2d} are each independently, and each of R^{2c}s
or each of R^{2d}s is independently when R^{2c} or R^{2d} exists plurally,
hydrogen or C₁₋₆ alkyl;

R³ is C₁₋₆ alkyl;

R^{4a}, R^{4b}, R^{4c}, and R^{4d} are each independently, and each of R^{4c}s
20 or each of R^{4d}s is independently when R^{4c} or R^{4d} exists plurally,
hydrogen, halogen, C₁₋₆ alkyl (wherein the alkyl may be optionally
substituted with the same or different 1 to 5 substituents
selected from the group consisting of halogen, hydroxy, C₁₋₆ alkoxy,
C₁₋₆ alkyl ester, and amino optionally-substituted with the same
25 or different 1 or 2 C₁₋₆ alkyl (wherein the alkyl may be optionally
substituted with oxo)), C₂₋₆ alkynyl, or amino (wherein the amino
may be optionally substituted with the same or different 1 or 2

C₁₋₆ alkyl); provided that when any two of R^{4a}, R^{4b}, R^{4c}, and R^{4d} are each independently the above C₁₋₆ alkyl, they may be combined together with the carbon atom or the carbon atoms to which they attach to form a 3- to 6-membered saturated carbocycle or a 4-
 5 to 6-membered saturated heterocycle wherein when the C₁₋₆ alkyl groups have a substituent, the substituent may be included as a ring member of the carbocycle or heterocycle;

Ring Q is a group of the following Formula (2a), (2b), (2c), (2d), (2e), or (2f):



wherein

R^{5a}, R^{5b}, R^{5c}, and R^{5d} are each independently hydrogen, halogen, or C₁₋₆ alkyl; and

R⁶ is hydrogen or C₁₋₆ alkyl.

[0060]

Another embodiment of a compound of Formula (1) includes the following embodiment (C):

(C) A compound, or a pharmaceutically acceptable salt thereof, wherein

Z is a nitrogen atom or $-\text{CR}^{\text{A}}-$;

Y is sulfonyl;

m is 1 or 2;

n is 1, 2, or 3;

5 R^{A} is hydrogen or C_{1-6} alkyl;

$\text{R}^{1\text{a}}$, $\text{R}^{1\text{b}}$, $\text{R}^{1\text{c}}$, and $\text{R}^{1\text{d}}$ are each independently hydrogen or C_{1-6} alkyl;


$\text{R}^{2\text{a}}$, $\text{R}^{2\text{b}}$, $\text{R}^{2\text{c}}$, and $\text{R}^{2\text{d}}$ are each independently, and each of $\text{R}^{2\text{cs}}$ or each of $\text{R}^{2\text{ds}}$ is independently when $\text{R}^{2\text{c}}$ or $\text{R}^{2\text{d}}$ exists plurally,
10 hydrogen or C_{1-6} alkyl;

R^3 is C_{1-6} alkyl;

$\text{R}^{4\text{a}}$, $\text{R}^{4\text{b}}$, $\text{R}^{4\text{c}}$, and $\text{R}^{4\text{d}}$ are each independently, and each of $\text{R}^{4\text{cs}}$ or each of $\text{R}^{4\text{ds}}$ is independently when $\text{R}^{4\text{c}}$ or $\text{R}^{4\text{d}}$ exists plurally, hydrogen, halogen, C_{1-6} alkyl (wherein the alkyl may be optionally
15 substituted with the same or different 1 to 5 substituents selected from the group consisting of halogen, hydroxy, C_{1-6} alkoxy, C_{1-6} alkyl ester, and amino optionally-substituted with the same or different 1 or 2 C_{1-6} alkyl (wherein the alkyl may be optionally substituted with oxo)), C_{2-6} alkynyl, or amino (wherein the amino
20 may be optionally substituted with the same or different 1 or 2 C_{1-6} alkyl); provided that when any two of $\text{R}^{4\text{a}}$, $\text{R}^{4\text{b}}$, $\text{R}^{4\text{c}}$, and $\text{R}^{4\text{d}}$ are each independently the above C_{1-6} alkyl, they may be combined together with the carbon atom or the carbon atoms to which they
25 attach to form a 3- to 6-membered saturated carbocycle or a 4- to 6-membered saturated heterocycle wherein when the C_{1-6} alkyl groups have a substituent, the substituent may be included as a ring member of the carbocycle or heterocycle;

provided that when n is 1, either R^{4c} or R^{4d} is hydrogen; and when n is 2 or 3, either R^{4c} or R^{4d} that binds to the carbon atom adjacent to Y is hydrogen; and

Ring Q is a group of the following Formula (2a), (2b), (2c),
 5 (2d), (2e), or (2f):



wherein

R^{5a} , R^{5b} , R^{5c} , and R^{5d} are each independently hydrogen, halogen, or C_{1-6} alkyl; and

10 R^6 is hydrogen or C_{1-6} alkyl.

[0061]

Another embodiment of a compound of Formula (1) includes the following embodiment (D):

(D) A compound, or a pharmaceutically acceptable salt thereof,

15 wherein

Z is $-CR^A-$;

Y is carbonyl;

m is 1;

n is 0 or 1;

R^A is hydrogen;

R^{1a}, R^{1b}, R^{1c}, and R^{1d} are hydrogen;

R^{2a}, R^{2b}, R^{2c}, and R^{2d} are hydrogen;

R³ is C₁₋₆ alkyl;

5 R^{4a}, R^{4b}, R^{4c}, and R^{4d} are each independently, and each of R^{4c}s
or each of R^{4d}s is independently when R^{4c} or R^{4d} exists plurally,
hydrogen, halogen, C₁₋₆ alkyl (wherein the alkyl may be optionally
substituted with the same or different 1 to 5 substituents
selected from the group consisting of halogen, hydroxy, C₁₋₆ alkoxy,
10 C₁₋₆ alkyl ester, and amino optionally-substituted with the same
or different 1 or 2 C₁₋₆ alkyl (wherein the alkyl may be optionally
substituted with oxo)), C₂₋₆ alkynyl, or amino (wherein the amino
may be optionally substituted with the same or different 1 or 2
C₁₋₆ alkyl); provided that when any two of R^{4a}, R^{4b}, R^{4c}, and R^{4d}
15 are each independently the above C₁₋₆ alkyl, they may be combined
together with the carbon atom or the carbon atoms to which they
attach to form a 3- to 6-membered saturated carbocycle or a 4-
to 6-membered saturated heterocycle wherein when the C₁₋₆ alkyl
groups have a substituent, the substituent may be included as a
20 ring member of the carbocycle or heterocycle; and

Ring Q is a group of formula (2a).

[0062]

Another embodiment of a compound of Formula (1) includes the
following embodiment (E):

25 (E) A compound, or a pharmaceutically acceptable salt thereof,
wherein

Z is -CR^A-;

Y is carbonyl;

m is 1;

n is 0 or 1;

R^A is hydrogen;

5 R^{1a}, R^{1b}, R^{1c}, and R^{1d} are each independently hydrogen;

R^{2a}, R^{2b}, R^{2c}, and R^{2d} are each independently hydrogen;

R³ is C₁₋₆ alkyl;

R^{4a}, R^{4b}, R^{4c}, and R^{4d} are each independently, and each of R^{4c}s or each of R^{4d}s is independently when R^{4c} or R^{4d} exists plurally,
 10 hydrogen, halogen, C₁₋₆ alkyl (wherein the alkyl may be optionally substituted with the same or different 1 to 5 substituents selected from the group consisting of halogen, hydroxy, C₁₋₆ alkoxy, C₁₋₆ alkyl ester, and amino optionally-substituted with the same or different 1 or 2 C₁₋₆ alkyl (wherein the alkyl may be optionally
 15 substituted with oxo)), C₂₋₆ alkynyl, or amino (wherein the amino may be optionally substituted with the same or different 1 or 2 C₁₋₆ alkyl); provided that when any two of R^{4a}, R^{4b}, R^{4c}, and R^{4d} are each independently the above C₁₋₆ alkyl, they may be combined together with the carbon atom or the carbon atoms to which they
 20 attach to form a 3- to 6-membered saturated carbocycle or a 4- to 6-membered saturated heterocycle wherein when the C₁₋₆ alkyl groups have a substituent, the substituent may be included as a ring member of the carbocycle or heterocycle; and

Ring Q is a group of formula (2d).

25 [0063]

Another embodiment of a compound of Formula (1) includes the following embodiment (F):

(F) A compound, or a pharmaceutically acceptable salt thereof, wherein

Z is $-CR^A-$;

Y is carbonyl;

5 m is 1;

n is 0 or 1;

R^A is hydrogen;

R^{1a} , R^{1b} , R^{1c} , and R^{1d} are hydrogen;

R^{2a} , R^{2b} , R^{2c} , and R^{2d} are hydrogen;

10 R^3 is C_{1-6} alkyl;

R^{4a} , R^{4b} , R^{4c} , and R^{4d} are each independently, and each of R^{4c} s or each of R^{4d} s is independently when R^{4c} or R^{4d} exists plurally, hydrogen, halogen, C_{1-6} alkyl (wherein the alkyl may be optionally substituted with the same or different 1 to 5 substituents
15 selected from the group consisting of halogen, hydroxy, C_{1-6} alkoxy, C_{1-6} alkyl ester, and amino optionally-substituted with the same or different 1 or 2 C_{1-6} alkyl (wherein the alkyl may be optionally substituted with oxo)), C_{2-6} alkynyl, or amino (wherein the amino may be optionally substituted with the same or different 1 or 2
20 C_{1-6} alkyl); provided that when any two of R^{4a} , R^{4b} , R^{4c} , and R^{4d} are each independently the above C_{1-6} alkyl, they may be combined together with the carbon atom or the carbon atoms to which they attach to form a 3- to 6-membered saturated carbocycle or a 4- to 6-membered saturated heterocycle wherein when the C_{1-6} alkyl
25 groups have a substituent, the substituent may be included as a ring member of the carbocycle or heterocycle; and

Ring Q is a group of formula (2e).

[0064]

Another embodiment of a compound of Formula (1) includes the following embodiment (G):

(G) A compound, or a pharmaceutically acceptable salt thereof,
5 wherein

Z is $-CR^A-$;

Y is sulfonyl;

m is 1;

n is 1;

10 R^A is hydrogen;

R^{1a} , R^{1b} , R^{1c} , and R^{1d} are hydrogen;

R^{2a} , R^{2b} , R^{2c} , and R^{2d} are hydrogen;

R^3 is C_{1-6} alkyl;

R^{4a} , R^{4b} , R^{4c} , and R^{4d} are each independently, and each of R^{4c} s
15 or each of R^{4d} s is independently when R^{4c} or R^{4d} exists plurally,
hydrogen, halogen, C_{1-6} alkyl (wherein the alkyl may be optionally
substituted with the same or different 1 to 5 substituents
selected from the group consisting of halogen, hydroxy, C_{1-6} alkoxy,
 C_{1-6} alkyl ester, and amino optionally-substituted with the same
20 or different 1 or 2 C_{1-6} alkyl (wherein the alkyl may be optionally
substituted with oxo)), C_{2-6} alkynyl, or amino (wherein the amino
may be optionally substituted with the same or different 1 or 2
 C_{1-6} alkyl); provided that when any two of R^{4a} , R^{4b} , R^{4c} , and R^{4d}
are each independently the above C_{1-6} alkyl, they may be combined
25 together with the carbon atom or the carbon atoms to which they
attach to form a 3- to 6-membered saturated carbocycle or a 4-
to 6-membered saturated heterocycle wherein when the C_{1-6} alkyl

groups have a substituent, the substituent may be included as a ring member of the carbocycle or heterocycle;

provided that either R^{4c} or R^{4d} is hydrogen; and

Ring Q is a group of formula (2a).

5 [0065]

Another embodiment of a compound of Formula (1) includes the following embodiment (H):

(H) A compound, or a pharmaceutically acceptable salt thereof, wherein

10 Z is $-CR^A-$;

Y is sulfonyl;

m is 1;

n is 1;

R^A is hydrogen;

15 R^{1a} , R^{1b} , R^{1c} , and R^{1d} are hydrogen;

R^{2a} , R^{2b} , R^{2c} , and R^{2d} are hydrogen;

R^3 is C_{1-6} alkyl;

R^{4a} , R^{4b} , R^{4c} , and R^{4d} are each independently, and each of R^{4cs} or each of R^{4ds} is independently when R^{4c} or R^{4d} exists plurally,
20 hydrogen, halogen, C_{1-6} alkyl (wherein the alkyl may be optionally substituted with the same or different 1 to 5 substituents selected from the group consisting of halogen, hydroxy, C_{1-6} alkoxy, C_{1-6} alkyl ester, amino optionally-substituted with the same or different 1 or 2 C_{1-6} alkyl (wherein the alkyl may be optionally
25 substituted with oxo)), C_{2-6} alkynyl, or amino (wherein the amino may be optionally substituted with the same or different 1 or 2 C_{1-6} alkyl); provided that when any two of R^{4a} , R^{4b} , R^{4c} , and R^{4d}

are each independently the above C₁₋₆ alkyl, they may be combined together with the carbon atom or the carbon atoms to which they attach to form a 3- to 6-membered saturated carbocycle or a 4- to 6-membered saturated heterocycle wherein when the C₁₋₆ alkyl groups have a substituent, the substituent may be included as a ring member of the carbocycle or heterocycle;

provided that either R^{4c} or R^{4d} is hydrogen; and

Ring Q is a group of formula (2e).

[0066]

A compound of Formula (1) may have at least one asymmetric carbon atom. The present compound, therefore, includes racemates of a compound of Formula (1), as well as optically-active isomers of a compound of Formula (1). When a compound of Formula (1) has two or more asymmetric carbon atoms, it may have stereoisomerism. The present compound, therefore, includes stereoisomers of a compound of Formula (1) and mixtures thereof.

[0067]

Further, deuterated compounds in which any one, two, or more ¹H in a compound of Formula (1) are replaced with ²H(D) are included in a compound of Formula (1).

[0068]

A compound of Formula (1) or a pharmaceutically acceptable salt thereof may exist in the form of a hydrate and/or a solvate, and the hydrate and solvate such as an ethanol solvate are included in the present compound. Further, the present compound also includes those in crystalline forms of its all embodiments.

[0069]

When a compound of Formula (1) has an acidic group, a pharmaceutically acceptable salt thereof includes, for example, an alkali metal salt such as a sodium salt and a potassium salt; an alkaline earth metal salt such as a calcium salt and a magnesium salt; an inorganic metal salt such as a zinc salt; and an organic basic salt such as triethylamine, triethanolamine, trihydroxymethylaminomethane, and an amino acid;.

[0070]

When a compound of Formula (1) has a basic group, a pharmaceutically acceptable salt thereof includes, for example, an inorganic acid salt such as hydrochloride, hydrobromide, sulfate, phosphate, and nitrate; an organic acid salt such as acetate, propionate, succinate, lactate, malate, tartrate, citrate, maleate, fumarate, methanesulfonate, p-toluenesulfonate, benzenesulfonate, ascorbate, and orotate.

[0071]

Hereinafter, the processes to prepare the present compound are explained along with examples, but the present invention shall not be limited thereto.

[0072]

Preparation Process

The present compound can be prepared by means of any of the preparation processes mentioned below and those combined with known processes.

Each compound appearing in the following schemes may also be in its salt form, and such salt may include, for example, a corresponding salt exemplified as a salt of a compound of Formula

(1). The reactions mentioned below are just examples, and thus the present compound may be optionally prepared by other means based on the knowledge of a person skilled in organic synthesis.

[0073]

5 If there is a functional group that needs to be protected in preparation processes mentioned below, the functional group may be protected as appropriate and then deprotected after completing a reaction or reaction sequences to obtain a desired compound, even though the use of any protecting group is not
10 specifically indicated.

[0074]

 The protecting group used herein includes, for example, general protecting groups described in *T. W. Greene and P. G. M. Wuts, "Protective Groups in Organic Synthesis", 3rd Ed., John*
15 *Wiley and Sons, inc., New York (1999)*. In more detail, a protecting group for an amino group includes, for example, benzyloxycarbonyl, tert-butoxycarbonyl, acetyl, and benzyl. A protecting group for a hydroxy group includes, for example, trialkylsilyl, acetyl, and benzyl.


20 [0075]

 The protection and deprotection can be carried out by conventional means in organic synthetic chemistry (for example, means described in *T. W. Greene and P. G. M. Wuts, "Protective Groups in Organic Synthesis", 3rd Ed., John Wiley and Sons, inc.,*
25 *New York (1999)*), or similar means to them.

[0076]

Preparation Process 1

The compound of Formula (1a), among compounds of Formula (1), can be prepared, for example, by the following process:



wherein Z , m , n , R^{1a} , R^{1b} , R^{1c} , R^{1d} , R^{2a} , R^{2b} , R^{2c} , R^{2d} , R^3 , R^{4a} , R^{4b} ,
5 R^{4c} , R^{4d} , and Ring Q are as defined in the above Item [1].

[0077]

Compound (1a) is prepared by reacting Compound (4) and carboxylic acid of Formula (5) in a suitable inert solvent in the presence of a suitable condensing agent. The reaction may be carried out in the presence of a suitable base. The reaction temperature generally ranges from about -20°C to the boiling point of a solvent used herein. The reaction time depends on reaction conditions such as the reaction temperature, a condensing agent used herein, starting materials, and a solvent used herein, and generally ranges from 10 minutes to 48 hours.

[0078]

Examples of condensing agents used herein include dicyclohexylcarbodiimide (DCC), diisopropylcarbodiimide (DIPC), 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide (WSC),
20 benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (BOP), diphenylphosphonyl amide (DPPA), N,N-carbonyldiimidazole (CDI), benzotriazol-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU), and O-(7-

azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium
hexafluorophosphate (HATU).

As necessary, additives may be added, such as N-
hydroxysuccinimide (HOSu), 1-hydroxybenzotriazole (HOBt), 3-
5 hydroxy-4-oxo-3,4-dihydro-1,2,3-benzotriazine (HOOBt).

[0079]

Examples of bases used herein include organic bases such as
triethylamine, diisopropylethylamine, and pyridine; inorganic
bases such as potassium carbonate, sodium carbonate, cesium
10 carbonate, potassium bicarbonate, sodium bicarbonate, potassium
dihydrogenphosphate, dipotassium hydrogenphosphate, potassium
phosphate, sodium dihydrogenphosphate, disodium
hydrogenphosphate, sodium phosphate, potassium hydroxide, sodium
hydroxide, and sodium hydride; and metal alkoxides such as sodium
15 methoxide and potassium tert-butoxide.

[0080]


Examples of inert solvents used herein include halogenated
hydrocarbons such as chloroform and dichloromethane; aromatic
hydrocarbons such as benzene and toluene; ether solvents such as
20 diethyl ether, tetrahydrofuran (THF), and 1,4-dioxane; aprotic
polar solvents such as acetonitrile, acetone, methylethyl ketone,
dimethylformamide, N-methyl-2-pyrrolidinone, and
dimethylsulfoxide; basic solvents such as pyridine; and mixture
solvents thereof.

25 [0081]

Preparation Process 2

Among compounds of Formula (4), the compound of Formula (4a)

can be prepared, for example, by the following process.



In the scheme, Z, R^{1a}, R^{1b}, R^{1c}, R^{1d}, R^{2a}, R^{2b}, R^{2c}, R^{2d}, R³, and Ring Q are as defined in the above Item [1]; and l is 0, 1, 2, or 3.

5 [0082]

Compound (8) can be prepared by reacting Compound (6) and an aldehyde of Formula (7) under reductive amination with a suitable reducing agent in a suitable inert solvent. The reaction may be carried out in the presence of a suitable base or acid, as necessary. The reaction temperature generally ranges from about -20°C to the boiling point of a solvent used herein. The reaction time depends on reaction conditions such as the reaction temperature, a reducing agent used herein, starting materials, and a solvent used herein, and generally ranges from 10 minutes to 48 hours.

[0083]

Examples of reducing agents used herein include complex hydride compounds such as sodium triacetoxyborohydride, lithium aluminum hydride, sodium borohydride, and sodium cyanoborohydride; and borane complexes such as borane-dimethyl sulfide complex and borane-tetrahydrofuran complex.

[0084]

Examples of bases used herein include organic bases such as triethylamine, diisopropylethylamine, and pyridine; inorganic

bases such as potassium carbonate, sodium carbonate, cesium carbonate, potassium bicarbonate, sodium bicarbonate, potassium dihydrogenphosphate, dipotassium hydrogenphosphate, potassium phosphate, sodium dihydrogenphosphate, disodium hydrogenphosphate, sodium phosphate, potassium hydroxide, sodium hydroxide, and sodium hydride; and metal alkoxides such as sodium methoxide and potassium tert-butoxide.

[0085]

Examples of acids used herein include organic acids such as acetic acid, trifluoroacetic acid, and methanesulfonic acid; and inorganic acids such as hydrochloric acid and sulfuric acid.

[0086]

Examples of inert solvents used herein include water; acetonitrile; halogenated hydrocarbons such as chloroform and dichloromethane; aromatic hydrocarbons such as benzene and toluene; ether solvents such as 1,2-dimethoxyethane, tetrahydrofuran, and 1,4-dioxane; alcohol solvents such as methanol, ethanol, and 2-propanol; aprotic polar solvents such as dimethylformamide and N-methyl-2-pyrrolidinone; and mixture solvents thereof.

[0087]

Compound (4a) can be prepared by treating Compound (8) with a suitable acid in a suitable inert solvent. The reaction temperature generally ranges from -20°C to the boiling point of a solvent used herein. The reaction time depends on reaction conditions such as the reaction temperature, an acid used herein, starting materials, and a solvent used herein, and generally

ranges from 10 minutes to 48 hours.

[0088]

Examples of acids used herein include organic acids such as trifluoroacetic acid; and inorganic acids such as hydrochloric
5 acid and sulfuric acid.


[0089]

Examples of inert solvents used herein include halogenated hydrocarbons such as chloroform and dichloromethane; aromatic hydrocarbons such as benzene and toluene; ether solvents such as
10 diethyl ether, tetrahydrofuran, 1,4-dioxane, and 1,2-dimethoxyethane; lower alcohols such as methanol, ethanol, and 2-propanol; and aprotic polar solvents such as acetonitrile, dimethylformamide, N-methyl-2-pyrrolidinone, and dimethylsulfoxide; and mixture solvents thereof.

15 [0090]

Preparation Process 3

The compound of Formula (8) can be prepared, for example, by the following process.



In the scheme, Z, m, R^{1a}, R^{1b}, R^{1c}, R^{1d}, R^{2a}, R^{2b}, R^{2c}, R^{2d}, R³, and Ring Q are as defined in the above Item [1]; LG is a leaving group such as iodine, bromine, chlorine, substituted sulfonyl (e.g., methanesulfonyl and p-toluenesulfonyl).

[0091]

Compound (8) can be prepared by reacting Compound (6) with the alkylating agent of Formula (9) in a suitable inert solvent. The reaction may be carried out in the presence of a suitable base, as necessary, and further in the presence of a suitable phase transfer catalyst. The reaction temperature generally ranges from about -20°C to the boiling point of a solvent used herein. The reaction time depends on reaction conditions such as the reaction temperature, a base used herein, starting materials, and a solvent used herein, and generally ranges from 10 minutes to 48 hours.

[0092]

Examples of bases used herein include organic bases such as triethylamine, diisopropylethylamine, and pyridine; inorganic bases such as potassium carbonate, sodium carbonate, cesium carbonate, potassium bicarbonate, sodium bicarbonate, potassium dihydrogenphosphate, dipotassium hydrogenphosphate, potassium phosphate, sodium dihydrogenphosphate, disodium hydrogenphosphate, sodium phosphate, potassium hydroxide, sodium hydroxide, and sodium hydride; and metal alkoxides such as sodium methoxide and potassium tert-butoxide.

[0093]

Examples of phase transfer catalysts used herein include tetrabutylammonium hydrogen sulfate.

[0094]

Examples of inert solvents used herein include halogenated hydrocarbons such as chloroform and dichloromethane; aromatic hydrocarbons such as benzene and toluene; ether solvents such as diethyl ether, tetrahydrofuran (THF), and 1,4-dioxane; lower alcohols such as methanol, ethanol, and 2-propanol; aprotic polar solvents such as acetonitrile, acetone, methylethyl ketone, dimethylformamide, N-methyl-2-pyrrolidinone, and dimethylsulfoxide; and mixture solvent thereof.

[0095]

Compound (11) can be prepared by reacting Compound (6) with the alkylating agent of Formula (10) in a suitable inert solvent. The reaction may be carried out in the presence of a suitable base as necessary, and further in the presence of a suitable phase transfer catalyst. The reaction temperature generally

ranges from about -20°C to the boiling point of a solvent used herein. The reaction time depends on reaction conditions such as the reaction temperature, a base used herein, starting materials, and a solvent used herein, and generally ranges from 10 minutes to 48 hours.

[0096]

Examples of bases used herein include organic bases such as triethylamine, diisopropylethylamine, and pyridine; inorganic bases such as potassium carbonate, sodium carbonate, cesium carbonate, potassium bicarbonate, sodium bicarbonate, potassium dihydrogenphosphate, dipotassium hydrogenphosphate, potassium phosphate, sodium dihydrogenphosphate, disodium hydrogenphosphate, sodium phosphate, potassium hydroxide, sodium hydroxide, and sodium hydride; and metal alkoxides such as sodium methoxide and potassium tert-butoxide.

[0097]

Examples of phase transfer catalysts used herein include tetrabutylammonium hydrogen sulfate.

[0098]

Examples of inert solvents used herein include halogenated hydrocarbons such as chloroform and dichloromethane; aromatic hydrocarbons such as benzene and toluene; ether solvents such as diethyl ether, tetrahydrofuran (THF), and 1,4-dioxane; lower alcohols such as methanol, ethanol, and 2-propanol; aprotic polar solvents such as acetonitrile, acetone, methylethyl ketone, dimethylformamide, N-methyl-2-pyrrolidinone, and dimethylsulfoxide; and mixture solvents thereof.

[0099]

Compound (8) can be prepared by reacting Compound (11) with the alkylating agent of Formula (12) in a suitable inert solvent. The reaction may be carried out in the presence of a suitable
5 base as necessary, and further in the presence of a suitable phase transfer catalyst. The reaction temperature generally ranges from about -20°C to the boiling point of a solvent used herein. The reaction time depends on reaction conditions such as the reaction temperature, a base used herein, starting
10 materials, and a solvent used herein, and generally ranges from 10 minutes to 48 hours.

[0100]

Examples of bases used herein include organic bases such as triethylamine, diisopropylethylamine, and pyridine; inorganic
15 bases such as potassium carbonate, sodium carbonate, cesium carbonate, potassium bicarbonate, sodium bicarbonate, potassium dihydrogenphosphate, dipotassium hydrogenphosphate, potassium phosphate, sodium dihydrogenphosphate, disodium hydrogenphosphate, sodium phosphate, potassium hydroxide, sodium
20 hydroxide, and sodium hydride; and metal alkoxides such as sodium methoxide and potassium tert-butoxide.

[0101]

Examples of phase transfer catalysts used herein include tetrabutylammonium hydrogen sulfate.

25 [0102]


Examples of inert solvents used herein include halogenated hydrocarbons such as chloroform and dichloromethane; aromatic

hydrocarbons such as benzene and toluene; ether solvents such as diethyl ether, tetrahydrofuran (THF), and 1,4-dioxane; lower alcohols such as methanol, ethanol, and 2-propanol; aprotic polar solvents such as acetonitrile, acetone, methylethyl ketone, dimethylformamide, N-methyl-2-pyrrolidinone, and dimethylsulfoxide; and mixture solvents thereof.

[0103]

Preparation Process 4

Among compounds of Formula (1), the compound of Formula (1f) can be prepared, for example, by the following process.



In the scheme, Z, m, R^{1a}, R^{1b}, R^{1c}, R^{1d}, R^{2a}, R^{2b}, R^{2c}, R^{2d}, R³, R^{4a}, R^{4b}, R^{4c}, and Ring Q are as defined in the above Item [1].

[0104]

Compound (14) can be prepared by reacting Compound (4) and sulfonyl chloride of Formula (13) in the presence of a suitable inert solvent. The reaction may be carried out in the presence of a suitable base. The reaction temperature generally ranges

from about -20°C to the boiling point of a solvent used herein. The reaction time depends on reaction conditions such as the reaction temperature, starting materials used herein, and a solvent, and generally ranges from 10 minutes to 48 hours.

5 [0105]

Examples of bases used herein include organic bases such as triethylamine, diisopropylethylamine, and pyridine; inorganic bases such as potassium carbonate, sodium carbonate, cesium carbonate, potassium bicarbonate, sodium bicarbonate, potassium
10 dihydrogenphosphate, dipotassium hydrogenphosphate, potassium phosphate, sodium dihydrogenphosphate, disodium hydrogenphosphate, sodium phosphate, potassium hydroxide, sodium hydroxide, and sodium hydride; and metal alkoxides such as sodium methoxide and potassium tert-butoxide.

15 [0106]

Examples of inert solvents used herein include halogenated hydrocarbons such as chloroform and dichloromethane; aromatic hydrocarbons such as benzene and toluene; ether solvents such as diethyl ether, tetrahydrofuran (THF), and 1,4-dioxane; aprotic
20 polar solvents such as acetonitrile, acetone, methylethyl ketone, dimethylformamide, N-methyl-2-pyrrolidinone, and dimethylsulfoxide; basic solvents such as pyridine; and mixture solvents thereof.

[0107]

25 Compound (1f) can be prepared by reacting Compound (14) with a hydroxide such as tetrabutylammonium hydroxide and sodium hydroxide in a suitable inert solvent. The reaction temperature

generally ranges from about -20°C to the boiling point of a solvent used herein. The reaction time depends on reaction conditions such as the reaction temperature, starting materials used herein, and a solvent used herein, and generally ranges from


[0108]

Examples of inert solvents used herein include aromatic hydrocarbons such as benzene and toluene; ether solvents such as diethyl ether, tetrahydrofuran (THF), and 1,4-dioxane; aprotic polar solvents such as acetonitrile, acetone, methylethyl ketone, dimethylformamide, N-methyl-2-pyrrolidinone, and dimethylsulfoxide; and mixture solvents thereof.

[0109]

Preparation Process 5

Among compounds of Formula (6), the compound of Formula (6a) can be prepared, for example, by the following process.



In the scheme, R^{5a} , R^{5b} , R^{5c} , and R^{5d} are as defined in the above Item [1]; and A is iodine, bromine, or chlorine.

[0110]

Compound (15) can be prepared by reacting 1-(tert-butoxycarbonyl)piperidine-4-carboxylic acid with N,O-dimethylhydroxylamine or its hydrochloride in the presence of a suitable condensing agent in a suitable inert solvent. The reaction may be carried out in the presence of a suitable base. The reaction temperature generally ranges from about -20°C to the boiling point of a solvent used herein. The reaction time depends on reaction conditions such as the reaction temperature, a condensing agent used herein, starting materials, and a solvent used herein, and generally ranges from 10 minutes to 48 hours.

[0111]

Compound (15) can also be prepared by reacting N,O-dimethylhydroxylamine or its salt with an acid halide or acid anhydride which is derived from 1-(tert-butoxycarbonyl)piperidine-4-carboxylic acid, in the presence of a suitable base, in a suitable inert solvent. The reaction temperature generally ranges from about -20°C to the boiling point of a solvent used herein. The reaction time depends on reaction conditions such as the reaction temperature, a condensing agent used herein, starting materials, and a solvent used herein, and generally ranges from 10 minutes to 48 hours.

[0112]

Examples of condensing agents used herein include dicyclohexylcarbodiimide (DCC), diisopropylcarbodiimide (DIPC), 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide (WSC), benzotriazol-1-yl-tris(dimethylamino)phosphonium hexafluorophosphate (BOP), diphenylphosphonyl amide (DPPA), N,N-

carbonyldiimidazole (CDI), and benzotriazol-1-yl-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU). The reaction may be carried out with addition of an additive such as N-hydroxysuccinimide (HOSu), 1-hydroxybenzotriazole (HOBt), and 3-hydroxy-4-oxo-3,4-dihydro-1,2,3-benzotriazine (HOOBt), as necessary.

[0113]

Examples of bases used herein include organic bases such as triethylamine, diisopropylethylamine, and pyridine; inorganic bases such as potassium carbonate, sodium carbonate, cesium carbonate, potassium bicarbonate, sodium bicarbonate, potassium dihydrogenphosphate, dipotassium hydrogenphosphate, potassium phosphate, sodium dihydrogenphosphate, disodium hydrogenphosphate, sodium phosphate, potassium hydroxide, sodium hydroxide, and sodium hydride; and metal alkoxides such as sodium methoxide and potassium tert-butoxide.

[0114]

Examples of inert solvents used herein include halogenated hydrocarbons such as chloroform and dichloromethane; aromatic hydrocarbons such as benzene and toluene; ether solvents such as diethyl ether, tetrahydrofuran (THF), and 1,4-dioxane; aprotic polar solvents such as acetonitrile, acetone, methylethyl ketone, dimethylformamide, N-methyl-2-pyrrolidinone, and dimethylsulfoxide; basic solvents such as pyridine; and mixture solvent thereof.

[0115]

Compound (17) can be prepared by reacting Compound (15) with

a lithiated compound which is produced by treating Compound (16) with an organolithium such as n-butyllithium in a suitable inert solvent. The reaction temperature generally ranges from about -78°C to the boiling point of a solvent used herein. The reaction time depends on reaction conditions such as the reaction temperature, a reagent used herein, starting materials, and a solvent used herein, and generally ranges from 10 minutes to 48 hours.

[0116]

Examples of inert solvents used herein include aromatic hydrocarbons such as benzene and toluene; ether solvents such as diethyl ether, tetrahydrofuran (THF), and 1,4-dioxane; and mixture solvents thereof.

[0117]

Compound (18) can be prepared by reacting Compound (17) with hydroxylamine or a salt thereof in the presence of a suitable base as necessary. The reaction temperature generally ranges from about -20°C to the boiling point of a solvent used herein. The reaction time depends on the reaction conditions such as the reaction temperature, a base used herein, starting materials, and a solvent used herein, and generally ranges from 10 minutes to 48 hours.

[0118]

Examples of bases used herein include organic bases such as triethylamine, diisopropylethylamine, and pyridine; inorganic bases such as potassium carbonate, sodium carbonate, cesium carbonate, potassium bicarbonate, sodium bicarbonate, potassium

dihydrogenphosphate, dipotassium hydrogenphosphate, potassium phosphate, sodium dihydrogenphosphate, disodium hydrogenphosphate, sodium phosphate, potassium hydroxide, sodium hydroxide, and sodium hydride; metal alkoxides such as sodium methoxide and potassium tert-butoxide; and sodium acetate.

[0119]

Examples of solvents used herein include aromatic hydrocarbons such as benzene and toluene; ether solvents such as diethyl ether, tetrahydrofuran (THF), and 1,4-dioxane; lower alcohols such as methanol, ethanol, and 2-propanol; aprotic polar solvents such as dimethylformamide and N-methyl-2-pyrrolidinon; water; and mixture solvents thereof.

[0120]

Compound (19) can be prepared by treating Compound (18) with a suitable base in a suitable inert solvent. The reaction temperature generally ranges from about -20°C to the boiling point of a solvent used herein. The reaction time depends on reaction conditions such as the reaction temperature, a base used herein, starting materials, and a solvent used herein, and generally ranges from 10 minutes to 48 hours.

[0121]

Examples of bases used herein include organic bases such as triethylamine, diisopropylethylamine, and pyridine; inorganic bases such as potassium carbonate, sodium carbonate, cesium carbonate, potassium bicarbonate, sodium bicarbonate, potassium dihydrogenphosphate, dipotassium hydrogenphosphate, potassium phosphate, sodium dihydrogenphosphate, disodium

hydrogenphosphate, sodium phosphate, potassium hydroxide, sodium hydroxide, and sodium hydride; and metal alkoxides such as sodium methoxide and potassium tert-butoxide.

[0122]

5 Examples of inert solvents used herein include halogenated hydrocarbons such as chloroform and dichloromethane; aromatic hydrocarbons such as benzene and toluene; ether solvents such as diethyl ether, tetrahydrofuran (THF), and 1,4-dioxane; lower alcohols such as methanol, ethanol, and 2-propanol; aprotic polar
10 solvents such as acetonitrile, acetone, methylethyl ketone, dimethylformamide, N-methyl-2-pyrrolidinone, and dimethylsulfoxide; and mixture solvents thereof.

[0123]

15 Compound (6a) can be prepared by processing Compound (19) with a suitable acid in a suitable inert solvent. The reaction temperature generally ranges from -20°C to the boiling point of a solvent used herein. The reaction time depends on reaction conditions such as the reaction temperature, an acid used herein, starting materials, and a solvent used herein, and generally
20 ranges from 10 minutes to 48 hours.

[0124]

25 Examples of inert solvents used herein include halogenated hydrocarbons such as chloroform and dichloromethane; aromatic hydrocarbons such as benzene and toluene; ether solvents such as diethyl ether, tetrahydrofuran, 1,4-dioxane, and 1,2-dimethoxyethane; lower alcohols such as methanol, ethanol, and 2-propanol; aprotic polar solvents such as acetonitrile,

dimethylformamide, N-methyl-2-pyrrolidinone, and dimethylsulfoxide; and mixture solvents thereof.

[0125]

Examples of acids used herein include inorganic acids such as hydrochloric acid and sulfuric acid; and organic acids such as trifluoroacetic acid.

[0126]

The present compound which has a desired functional group at a desired position can be prepared by suitably combining the above preparation processes. The isolation and purification of each intermediate or product in the above preparation processes can be carried out by conventional methods in organic synthesis, for example, by suitably combining filtration, extraction, washing, drying, concentration, crystallization, and various chromatography. Some intermediates may be used in the next step without purification.

[0127]

Some starting compounds or intermediates in the above preparation processes can exist in a salt form such as hydrochloride depending on reaction conditions, but can be used as it is or in a free form thereof. When a starting compound or an intermediate is obtained in a salt form and it needs to be used or obtained as a free form thereof, it can be transformed to a free form thereof by dissolving or suspending it in a suitable solvent and neutralizing the resulted solution or suspension with a base such as aqueous sodium bicarbonate.

[0128]

Some compounds of Formula (1) or pharmaceutically acceptable salts thereof can exist as isomers such as tautomer (for example, keto-enol form), regioisomer, geometrical isomer, and optical isomer. The present invention encompasses every possible isomer including the above, and mixtures thereof which have various mixture proportions.

[0129]

Optical isomers can be resolved by a known isolation method such as a method with an optically-active column and fractional crystallization at a suitable step in the above-mentioned preparation processes. In addition, an optically-active starting material can also be used for starting materials.

[0130]

In order to obtain a compound of Formula (1) as a salt thereof, when the product is a salt of a compound of Formula (1), the product should be directly purified; or when the product is in a free form of a compound of Formula (1), the product should be dissolved or suspended in an appropriate solvent and then an acid or a base should be added thereto to form a salt thereof.

[0131]

The present compound has antagonist activity for the 5-HT_{2A} receptor and antagonist activity for the 5-HT₇ receptor, and has different mechanism from other existing medicaments for treating mental diseases. The present compound can provide a new option in medication for various mental diseases. Specifically, the present compound is beneficial for the treatment of mental diseases. The present compound is also beneficial for the

treatment of central nervous system diseases.

[0132]

The mental diseases or central nervous system diseases which are expected to be treated effectively include, for example, F00-
 5 F09: organic, including symptomatic, mental disorders, F10-F19: mental and behavioural disorders due to use of psychoactive substance, F20-F29: schizophrenia, schizotypal disorders, and delusional disorders, F30-F39: mood [affective] disorders, F40-F48: neurotic disorders, stress-related disorders, and somatoform
 10 disorders, F51: nonorganic sleep disorders, F52: sexual dysfunction, not caused by organic disorder or disease, F84: pervasive developmental disorders, F90-F98: behavioural and emotional disorders with onset usually in childhood and adolescence, G20-G26: extrapyramidal and movement disorders, G30-
 15 G32: other degenerative diseases of nervous system, and G47: sleep disorders in International Classification of Diseases, 10th edition (ICD-10).

F00-F09: Organic, including symptomatic, mental disorders include, for example, Alzheimer's disease, vascular dementia,
 20 dementia with Lewy bodies, dementia in Parkinson's disease, mental disorders due to other diseases such as brain damage, and other mental disorders due to brain dysfunction and to physical disease.

F10-F19: Mental and behavioural disorders due to use of
 25 psychoactive substance include delirium tremens, psychotic disorder, and amnestic syndrome, due to use of various substance.

F20-F29: Schizophrenia, schizotypal disorders, and

delusional disorders include paranoid schizophrenia, simple schizophrenia, and delusional disorders.

F30-F39: Mood [affective] disorders include manic episode, bipolar affective disorder, and depressive episode.

5 F40-F48: Neurotic disorders, stress-related disorders, and somatoform disorders include phobic anxiety disorders, obsessive-compulsive disorder, and somatoform disorders.

F51: Nonorganic sleep disorders include nonorganic insomnia, sleepwalking, and nightmares.

10 F52: Sexual dysfunction, not caused by organic disorder or disease, includes lack or loss of sexual desire and unspecified sexual dysfunction.

F84: Pervasive developmental disorders include, for example, autism and overactive disorder associated with mental retardation
15 and stereotyped movements.

F90-F98: Hyperkinetic disorders and behavioural and emotional disorders with onset usually in childhood and adolescence include hyperkinetic disorders, conduct disorders, and mixed disorders of conduct and emotions.

20 G20-G26: Extrapyrarnidal and movement disorders include Parkinson's disease and secondary Parkinsonism.

G30-G32: Other degenerative diseases of nervous system include Alzheimer's disease, frontotemporal dementia, frontotemporal lobar degeneration, dementia with Lewy bodies, and
25 senile degeneration of brain.

G47: Sleep disorders include disorders of initiating and maintaining sleep [insomnias], disorders of the sleep-wake

schedule, narcolepsy, and cataplexy.

The present compound is useful for treatment or prevention of relapse of various symptoms associated with these diseases such as psychopathic symptoms, sleep disorders, depressive
5 symptoms, anxiety symptoms, and cognitive dysfunction.

[0133]

Serotonin (5-hydroxytryptamine: 5-HT), one of main neurotransmitters in central nervous system, is known to be involved in various brain functions such as emotional reaction
10 and cognitive function. The 5-HT_{2A} receptor, one of 5-HT receptor subtypes, is highly expressed in, for example, cerebral cortex, hippocampus, and raphe nucleus. The 5-HT_{2A} which is expressed in prefrontal cortex is also known to positively regulate a dopamine neural pathway in the ventral tegmental area (Non-Patent
15 Literature 2). That is, inhibiting the 5-HT_{2A} receptor in prefrontal cortex is considered to lead to an inhibitory effect to psychotic symptoms.

[0134]

The 5-HT₇ receptor is extensively expressed in, for example,
20 hypothalamus, thalamus, hippocampus, and raphe nucleus, and involves in a regulation of circadian rhythm in mammals (Non-Patent Literature 9). It is known that defect in circadian rhythm relates to various CNS disorders, especially for depression, seasonal affective disorder, a sleep disorder, shift work
25 syndrome, and jet lag. Drugs having antagonist activity for the 5-HT₇ receptor include Lurasidone which is used as a medicament for treating schizophrenia and a bipolar disorder and

Vortioxetine which is used for treating major depression. There is, however, no drug having selective antagonist activity for the 5-HT₇ receptor.

[0135]

5 It is also known that in some animal models, antagonizing the 5-HT₇ receptor leads to antidepressant and anxiolytic effect, and improvement effect in cognition function (Non-patent Literature 6, Non-patent Literature 7).

[0136]

10 From the above pharmacological knowledge, it is expected that inhibiting the 5-HT_{2A} receptor, along with inhibiting the 5-HT₇ receptor would be useful in various neuropsychiatric diseases such as depression, a sleep disorder, and a psychotic symptom. There is no report for a drug which have a selective and potent
15 antagonist activity on both of the 5-HT_{2A} receptor and the 5-HT₇ receptor.

[0137]

 The present compound has potent binding affinity to the 5-HT_{2A} receptor and the 5-HT₇ receptor (Test 1), and it shows an
20 antagonist activity on both of the 5-HT_{2A} receptor and the 5-HT₇ receptor. In a preferred embodiment of the present invention, the present compound can exert a pharmacological effect based on antagonist ability for the 5-HT_{2A} receptor and the 5-HT₇ receptor at the blood concentration which do not cause side effect such
25 as extrapyramidal symptom and hyperprolactinemia due to D₂ antagonist activity because a binding affinity of the present compound to the 5-HT_{2A} receptor and the 5-HT₇ receptor is 100

times higher more more than that to the D₂ receptor.

[0138]

In another preferred embodiment of the present invention, the present compound is expected to have a small effect for cardiovascular system because there is a big difference between the inhibitory concentration of hERG channel which is an express indicator of arrhythmia in long QT, and the express concentration of the expected pharmacological effect (Test 5).

[0139]

The disappearance half-life (" $T_{1/2}$ ") of a drug is a factor for determining the frequency of administration to retain its effect. It is thought that plural administrations of a drug per day having short $T_{1/2}$ can cause forgetting to take a medication or unfinishing taking a medication, and can lead to hindering a suitable medication. Furthermore, if the frequency of administration increases, it is concerned that the incidence rate of side effects can increase or the tolerability can decrease in association with high-dose administration. From the viewpoint mentioned above, if a drug having long $T_{1/2}$ is found out, the drug is expected to be a long-acting drug with little concern mentioned above, which can bring in liability relief of medicated patients.

In a preferred embodiment of the present compound, the estimated human disappearance half-life (" $T_{1/2}$ ") of the present compound is 8 hours or more (Test 4). Thus, it is expected that the drug efficacy can be retained for a long period in human body, the medication adherence of medicated patients can be improved, and a high tolerability can be exhibited at the administration.

[0140]

The present compound can be orally or parenterally administered. In the case of oral administration, the compound can be administered in commonly-used dosage form. In the case
5 of parenteral administration, the compound can be administered, for example, in a topical administration form, an injection form, a transdermal form, and a nasal form. The oral form or the rectal administration form include, for example, capsule, tablet, pill, powder, cachet, suppository, and liquid. The injection form
10 includes, for example, aseptic solution and suspension. The topical administration form includes, for example, cream, ointment, lotion, and transdermal formulation (e.g., normal patch and matrix).

[0141]

15 The above-mentioned dosage forms are prepared with a pharmaceutically acceptable excipient and an additive in a conventional manner. The pharmaceutically acceptable excipient and additive include carrier, binder, flavor, buffer, thickener, colorant, stabilizing agent, emulsifier, dispersant, suspending
20 agent, and preservative.

The pharmaceutically acceptable carrier includes, for example, magnesium carbonate, magnesium stearate, talc, sugar, lactose, pectin, dextrin, starch, gelatin, tragacanth, methylcellulose, sodium carboxymethylcellulose, low-melting-
25 point wax, and cocoa butter. The capsule form can be prepared by filling a capsule with the present compound and a pharmaceutically acceptable carrier. The present compound can

be put into a capsule with or without a pharmaceutically acceptable excipient. The cachet can also be prepared in a similar manner.

[0142]

5 The injectable liquid form includes solution, suspension, and emulsion, including, for example, water solution and water-propylene glycol solution. The liquid form may comprise water, and also it may be prepared in a solution of polyethylene glycol or/and propylene glycol. The liquid form suitable for oral
10 administration may be prepared by adding the present compound to water and also adding colorant, flavor, stabilizing agent, sweetener, solubilizer, or thickener thereto, as appropriate. Alternatively, the liquid form suitable for oral administration may be prepared by adding the present compound and a dispersant
15 to water and rendering the liquid sticky. The thickener used herein includes, for example, pharmaceutically acceptable natural or synthetic gum, resin, methylcellulose, sodium carboxymethylcellulose, and a known suspending agent.

[0143]

20 The dose of each compound can depend on patient's disease, age, body weight, gender, symptom, and the administration route. In general, the present compound is administered to an adult (body weight: 50 kg) by 0.1 to 1000 mg/day, preferably 1 to 300 mg/day, once a day or in 2 to 3 divided doses per day. Or, it
25 may be administered once in a few days to a few weeks.

[0144]

In order to enhance the effect and/or reduce side effect

thereof, the present compound may be used in combination with another drug. For example, the present compound may be used in combination with an antianxiety drug such as a selective serotonin reuptake inhibitor. Hereinafter, a drug with which the present
5 compound may be used in combination is abbreviated as a "concomitant drug".

[0145]

Examples of the concomitant drug used herein include a drug for treating a developmental disorder such as an autism spectrum
10 disorder and an attention deficit hyperactivity disorder, an antipsychotic drug and a schizophrenic drug, a drug for treating a bipolar disorder, an antidepressant drug, an antianxiety drug, a drug for treating an obsessive-compulsive disorder, a drug for treating a stress disorder such as a post-traumatic stress
15 disorder, a drug for treating a mood disorder, a drug for treating an eating disorder, a drug for treating a sleep disorder such as insomnia, narcolepsy, a sleep apnea syndrome, and a circadian rhythm disorder, a drug for treating a sexual dysfunction, a drug for treating drug dependence, a drug for treating dementia such
20 as Alzheimer's disease, a drug for treating a behavioral and psychological symptom associated with dementia, a cerebral metabolism and circulation improving drug, a drug for treating a movement disorder such as Parkinson's disease, an analgesic drug, an antiepileptic drug, an anticonvulsant, a migraine drug, an
25 anesthetic, and a central stimulant.

[0146]

The administration interval between the present compound and

its concomitant drug is not limited; i.e., the concomitant drug may be administered to a subject patient at the same time as the present compound or at a suitable interval. Or, the present compound and its concomitant drug can be formulated into a combination drug comprising these. The dose of the concomitant drug can be suitably determined based on the clinically-used dose thereof. The combination ratio of the present compound and its concomitant drug can be suitably determined based on its subject patient to be administered, administration route, disease, pathology, and combinations thereof. For example, when the subject patient is a human being, the concomitant drug may be used in 0.01 to 100 parts by weight per one part by weight of the present compound. For the purpose of reducing side effects, a concomitant drug such as an antiemetic drug, a sleep-inducing drug, and an anticonvulsant may be used in combination.

EXAMPLES

[0147]

The present invention is explained in more detail in the following by referring to Reference examples, Examples, and Tests; however, the scope of the present invention is not limited thereto. Compound names used in Reference examples and Examples are not always based on IUPAC nomenclature system. Abbreviations may be used for brevity, and these abbreviations have same meanings as those described above.

Compounds were identified by proton nuclear magnetic resonance absorption spectroscopy (^1H -NMR) or LC-MS. Amino

chromatography in Reference examples and Examples were conducted with an amino column from Yamazen Corporation. LC-MS was carried out with various conditions shown in the following table. Retention time (R.T.) denotes the time when a peak of a mass spectrum appears in LC-MS measurement.

Analytical apparatus	Shimadzu LCMS-2020
Column	Phenomenex Kinetex 1.7 μ m C18 (50 mm $\times$ 2.10 mm)
Eluent	A: MeOH, B: 0.05% TFA/H ₂ O
Gradient condition	0.0 min; A/B = 30:70 0.0 to 1.90 min; A/B = 99:1 1.91 to 3.00 min; A/B = 30:70
Flow rate	0.5 mL/min
Wavelength (UV)	220 nm
Column temperature	40°C

[0148]

Powder x-ray diffraction (powder XRD) measurement was carried out with various conditions shown in the following table.

Analytical apparatus	Empyrian (Spectris Co., Ltd.)
X ray	CuK α /45 kV/40 mA
Divergence slit	1/4°
Solar slit	0.04 rad
Anti-scattering slit	5.5 mm
Step size	0.013°
Scan range	4 to 40° (2 θ)
Cumulative time	100 second/step
Measurement temperature	23°C (296 K)

[0149]

Differential scanning calorimetry (DSC) measurement was carried out with various conditions shown in the following table.

Analytical apparatus	DSC2500 or DSCQ1000 (TA Instruments Inc.)
Measurement temperature range	10 to 250°C
Heating rate	10°C/minutes
Container	TzeroPan or Aluminium hermetic pan (Pinhole)
Atmosphere gas flow rate	Dry nitrogen: about 50 mL/minutes

[0150]

5 The following abbreviations may be used in the specification.

In NMR data of Reference examples and Examples, the following abbreviation are used.

Me: Methyl

DMF: N,N-Dimethylformamide

THF: Tetrahydrofuran

tert-: Tertiary

CDCl₃: Deuterated chloroform

DMSO-d₆: Deuterated dimethylsulfoxide


5 [0151]

Proton nuclear magnetic resonance spectra were measured with FT-NMR spectrometer (300 MHz or 400 MHz, JEOL). The chemical shifts were shown in δ value (ppm). The signs used in NMR denote the following meanings; s is singlet, d is doublet, dd is double
 10 doublet, dt is double triplet, t is triplet, q is quartet, m is multiplet, br is broad, brs is broad singlet, and J is the coupling constant..

[0152]

Example 1

15 (2R)-N-Ethyl-N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}-2-hydroxypropanamide



To a solution of the compound of Reference example 1 (20.0 mg) in N,N-dimethylformamide (0.5 mL) were added D-lactic acid
 20 (6.18 mg), triethylamine (20.8 mg), and 2-(1H-benzo[d][1,2,3]triazol-1-yl)-1,1,3,3-tetramethylisouronium hexafluorophosphate (33.8 mg). The mixture was stirred at room temperature for 16 hours, and water (4.0 mL) was added thereto. The mixture was extracted with chloroform (10 mL x 2), dried with
 25 anhydrous sodium sulfate, filtered, and concentrated. The


concentrated residue was purified by high performance liquid chromatography (column; SEMIPREPARATIVE C-18 column, separation condition; acetonitrile/trifluoroacetic acid: water/trifluoroacetic acid), and desalted with MP-Carbonate resin
 5 to obtain the titled compound (9.73 mg).

LC - MS: R.T. = 1.433 min ObsMS = 364 [M+1]

[0153]

Example 2

N-Ethyl-N-{2-[4-(5-fluoro-1H-indazol-1-yl)piperidin-1-yl]ethyl}-3-hydroxypropanamide
 10




To a suspension of the compound of Reference example 2 (50.0 mg) in dichloromethane (1.0 mL) were added triethylamine (0.077 mL), 30 % aqueous 3-hydroxypropionic acid (0.046 mL), and 2-(1H-benzo[d][1,2,3]triazol-1-yl)-1,1,3,3-tetramethylisouronium
 15 hexafluorophosphate (67.9 mg). The mixture was stirred at room temperature for 2 hours, and purified by silica gel column chromatography (chloroform/methanol) to obtain the titled compound (43.2 mg).

¹H-NMR (400 MHz, CDCl₃) δ: 7.89-7.86 (1H, m), 7.37-7.30 (1H, m), 7.29-7.24 (1H, m), 7.11-7.04 (1H, m), 4.44-4.27 (1H, m), 3.87-3.77 (2H, m), 3.72-3.57 (1H, m), 3.51-3.42 (1H, m), 3.42-3.27 (3H, m), 3.17-2.96 (2H, m), 2.59-2.47 (4H, m), 2.36-2.17 (4H, m), 2.05-1.89 (2H, m), 1.18-1.05 (3H, m).

[0154]

Example 3

(2R)-N-Ethyl-N-{2-[4-(5-fluoro-1H-indazol-1-yl)piperidin-1-yl]ethyl}-2-hydroxypropanamide














To a suspension of the compound of Reference example 2 (50.0 mg) in dichloromethane (1.0 mL) were added triethylamine (0.077 mL), D-lactic acid (14.9 mg), and 2-(1H-benzo[d][1,2,3]triazol-1-yl)-1,1,3,3-tetramethylisouronium hexafluorophosphate (67.9 mg). The mixture was stirred at room temperature for 4 hours, and purified by silica gel column chromatography (chloroform/methanol) to obtain the titled compound (13.9 mg).
¹H-NMR (400 MHz, CDCl₃) δ: 7.93 (1H, s), 7.42-7.28 (2H, m), 7.18-7.08 (1H, m), 4.50-4.20 (2H, m), 3.80-3.62 (1H, m), 3.59-2.88 (4H, m), 2.68-2.53 (1H, m), 2.44-2.20 (3H, m), 2.11-1.94 (1H, m), 1.67-1.44 (5H, m), 1.37-1.10 (6H, m).


[0155]

Examples 4 to 15

According to the method of Example 3, the compounds of Examples 4 to 15 were prepared from the corresponding Reference example compounds.

Example	Chemical structure	Instrumental analysis data
4		LC-MS: R.T. = 1.417 min ObsMS = 364 [M+1]
5		LC-MS: R.T. = 1.517 min ObsMS = 436 [M+1]
6		LC-MS: R.T. = 1.500 min ObsMS = 378 [M+1]
7		LC-MS: R.T. = 1.517 min ObsMS = 392 [M+1]
8		LC-MS: R.T. = 1.567 min ObsMS = 392 [M+1]
9		LC-MS: R.T. = 1.258 min ObsMS = 418 [M+1]
10		LC-MS: R.T. = 1.467 min ObsMS = 378 [M+1]


11		LC-MS: R.T. = 1.483 min ObsMS = 420 [M+1]
12		LC-MS: R.T. = 1.150 min ObsMS = 350 [M+1]
13		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.58-7.53 (1H, m), 7.16-7.11 (1H, m), 4.48-4.32 (1H, m), 3.75-3.59 (1H, m), 3.56-2.80 (5H, m), 2.65-2.43 (1H, m), 2.31 (3H, s), 2.29-1.85 (5H, m), 1.57-1.44 (4H, m), 1.32-1.27 (3H, m), 1.22-1.06 (3H, m).
14		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.52-7.40 (1H, m), 7.37 (1H, d, $J = 8.5$ Hz), 7.27 (1H, d, $J = 8.5$ Hz), 4.47-4.26 (1H, m), 3.75-3.57 (1H, m), 3.56-2.88 (6H, m), 2.62-2.45 (2H, m), 2.40 (3H, s), 2.34-1.93 (5H, m), 1.57-1.40 (2H, m), 1.33-1.25 (3H, m), 1.21-1.07 (3H, m).

15		¹ H-NMR (400 MHz, CDCl ₃) δ: 7.71-7.60 (1H, m), 7.57-7.54 (1H, m), 7.29-7.25 (1H, m), 4.58-4.28 (1H, m), 3.79-3.62 (1H, m), 3.58-2.87 (6H, m), 2.72-2.49 (2H, m), 2.40-1.93 (5H, m), 1.61-1.43 (2H, m), 1.37-1.29 (3H, m), 1.26-1.08 (3H, m).
----	---	--

[0156]

Example 16

(2S)-N-Ethyl-N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}-2-hydroxypropanamide



5


To a suspension of the compound of Reference example 1 (400 mg) in tetrahydrofuran (1.1 mL) were added triethylamine (0.763 mL), L-lactic acid (0.098 mL), and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (0.501 mg). The mixture was stirred at room temperature for 4 hours. To the mixture were added methanol (6.0 mL), 2 mol/L aqueous sodium hydroxide (3.0 mL), and water (1.0 mL). The mixture was stirred at room temperature for 1 hour. Then, water (30 mL) was added thereto, and the mixture was extracted with chloroform (30 mL x 2), dried over anhydrous sodium sulfate, filtered, and concentrated. The concentrated residue was purified by silica gel column chromatography (chloroform/methanol) to obtain the titled compound (279 mg).

^1H -NMR (400 MHz, CDCl_3) δ : 7.72-7.56 (1H, m), 7.17 (1H, dd, J = 8.5, 1.8 Hz), 6.99 (1H, td, J = 8.8, 1.8 Hz), 4.48-4.32 (1H, m), 3.75-3.59 (1H, m), 3.52-2.86 (7H, m), 2.58-2.44 (2H, m), 2.32-2.11 (2H, m), 2.09-1.87 (4H, m), 1.32-1.25 (3H, m), 1.17 (2H, t, J = 7.0 Hz), 1.09 (1H, t, J = 7.0 Hz).

[0157]

Example 17

N-Ethyl-N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}-3-hydroxypropanamide



To a suspension of the compound of Reference example 1 (3.97 g) in acetonitrile (50 mL) were added triethylamine (6.08 mL), 3-hydroxypropionic acid (4.91 g), and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (6.22 g). The mixture was stirring at room temperature for 2 hours. To the mixture were added methanol (50 mL) and cesium carbonate (14.2 g). The mixture was stirred at 70°C for 2 hours, and then filtered and concentrated. To the residue was added chloroform (50 mL), and insolubles were filtered out. The eluent was concentrated. The concentrated residue was purified by amino silica gel chromatography (hexane/ethyl acetate and chloroform/methanol), and further purified by silica gel column chromatography (chloroform/methanol) to obtain the titled compound (2.97 g).

^1H -NMR (400 MHz, CD_3OD) δ : 7.93-7.84 (1H, m), 7.42-7.33 (1H, m),

7.20-7.12 (1H, m), 3.88-3.79 (2H, m), 3.62-3.37 (4H, m), 3.23-3.04 (4H, m), 2.69-2.54 (4H, m), 2.42-2.26 (2H, m), 2.15-1.95 (4H, m), 1.29-1.07 (3H, m).

Powder XRD D(°, 2θ±0.2)

5 8.32, 9.72, 13.30, 13.62, 13.83, 14.24, 16.77, 17.81, 19.89, 19.95, 21.58, 22.02, 24.03, 26.77, 26.84 (Among them, the characteristic 10 peaks were 8.32, 9.72, 13.83, 14.24, 16.77, 19.89, 19.95, 21.58, 22.02, 24.03, and the more characteristic 4 peaks were 8.32, 9.72, 13.83, 16.77.)

10 DSC


Enthalpy (normalized): 71.71 J/g

Onset x: 67.55°C

[0158]

Example 18

15 N-Ethyl-N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}-2-hydroxy-3-methoxypropanamide



To a suspension of the compound of Reference 1 (70.0 mg) in tetrahydrofuran (1.9 mL) were added triethylamine (0.134 mL), 2-hydroxy-3-methoxypropionic acid (27.7 mg), and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (88.0 mg). The reaction mixture was stirred at room temperature for 3 hours and concentrated. The residue was purified by silica gel column chromatography


(chloroform/methanol) to obtain the titled compound (51.7 mg).

$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.78-7.62 (1H, m), 7.24-7.19 (1H, m), 7.03 (1H, td, $J = 8.9, 2.2$ Hz), 4.58-4.47 (1H, m), 3.82-2.89 (13H, m), 2.64-2.48 (2H, m), 2.39-1.96 (6H, m), 1.26-1.08 (3H, m).

5 [0159]

Example 19

N-Ethyl-N-{2-[4-(6-fluoro-1H-indazol-3-yl)piperidin-1-yl]ethyl}-2-hydroxypropanamide



10 To a solution of the compound of Reference example 6 (400 mg) in N,N-dimethylformamide (4.0 mL) were added diisopropylethylamine 0.577 mL), DL-lactic acid (0.082 mL), and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (628 mg). The mixture was stirred at room
15 temperature for 1 hour. To the reaction mixture was added 2 mol/L aqueous sodium hydroxide (3.0 mL), and the mixture was stirred at room temperature for 1 hour. Then, water (10 mL) was added to the reaction mixture. The mixture was extracted with chloroform (5.0 mL x 3), dried over anhydrous sodium sulfate,
20 filtered, and concentrated. The concentrated residue was purified by silica gel column chromatography (chloroform/methanol) to obtain the titled compound (217 mg).


$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 10.01-9.79 (1H, m), 7.74-7.63 (1H, m), 7.06 (1H, dd, $J = 9.1, 1.8$ Hz), 6.88 (1H, t, $J = 7.9$ Hz), 4.52-

4.37 (1H, m), 3.88-3.74 (2H, m), 3.74-3.59 (2H, m), 3.58-3.14 (3H, m), 3.11-2.93 (3H, m), 2.65-2.49 (2H, m), 2.36-1.95 (5H, m), 1.38-1.30 (3H, m), 1.26-1.10 (3H, m).

[0160]

5 Example 20

(2S)-N-Ethyl-N-{2-[4-(6-fluoro-1H-indol-3-yl)piperidin-1-yl]ethyl}-2-hydroxypropanamide



To a suspension of the compound of Reference example 7 (30.0
10 mg) in tetrahydrofuran (0.83 mL) were added triethylamine (0.0575 mL), L-lactic acid (8.95 mg), and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (37.8 mg). The mixture was stirred at room temperature for 2 hours. To the reaction mixture were added methanol (0.8 mL) and 2 mol/L aqueous
15 sodium hydroxide (0.4 mL). The mixture was stirred at room temperature for 30 minutes. The reaction mixture was concentrated, and chloroform (10 mL) was added thereto. The mixture was dried over anhydrous sodium sulfate, filtered, and concentrated. The concentrated residue was purified by silica gel column
20 chromatography (chloroform/methanol) to obtain the titled compound (24.3 mg).


¹H-NMR (400 MHz, CDCl₃) δ: 8.01 (1H, s), 7.45 (1H, dd, J = 8.5, 5.5 Hz), 6.97 (1H, dd, J = 9.8, 2.4 Hz), 6.87 (1H, s), 6.82-6.77 (1H, m), 4.47-4.32 (1H, m), 3.77-2.87 (6H, m), 2.79-2.68 (1H, m),

2.60-2.44 (2H, m), 2.30-2.13 (2H, m), 2.06-1.67 (4H, m), 1.33-1.25 (3H, m), 1.22-1.06 (3H, m).

[0161]

Example 21

5 (2S)-N-Ethyl-N-{2-[4-(5-fluoro-1H-indazol-1-yl)piperidin-1-yl]ethyl}-2-hydroxypropanamide




To a suspension of the compound of Reference example 2 (400 mg) in tetrahydrofuran (4.0 mL) were added triethylamine (0.768 mL), L-lactic acid (161 mg), and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (786 mg). The mixture was stirred at room temperature for 3 hours. To the reaction mixture was added 15 % aqueous sodium hydroxide (4.0 mL), and the mixture was stirred at room temperature for 1 hour. Then, concentrated hydrochloric acid was added thereto until pH of the reaction mixture reached 7, and the mixture was concentrated. The concentrated residue was purified by silica gel column chromatography (chloroform/methanol) to obtain the titled compound (269 mg).

20 ¹H-NMR (400 MHz, CDCl₃) δ: 7.88 (1H, s), 7.38-7.24 (2H, m), 7.12-7.04 (1H, m), 4.48-4.29 (2H, m), 3.77-3.56 (2H, m), 3.50-2.93 (5H, m), 2.72-2.48 (2H, m), 2.38-2.18 (3H, m), 2.12-1.88 (2H, m), 1.32-1.25 (3H, m), 1.23-1.06 (3H, m).

[0162]

Example 22

N-Ethyl-2-hydroxy-N-{2-[4-(1H-indol-3-yl)piperidin-1-yl]ethyl}propanamide




5 To a suspension of the compound of Reference example 8 (300 mg) in tetrahydrofuran (8.7 mL) were added triethylamine (0.606 mL), DL-lactic acid (94.0 mg), and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (398 mg). The mixture was stirred at room temperature for 3 hours, and methanol
10 (6.0 mL) and 2 mol/L aqueous sodium hydroxide (3.0 mL) were added thereto. The mixture was stirred at room temperature for 1 hour. Then, the reaction mixture was concentrated, and chloroform (30 mL) was added to the residue. The mixture was dried over anhydrous sodium sulfate, filtered, and concentrated. The
15 concentrated residue was purified by silica gel column chromatography (chloroform/methanol) to obtain the titled compound (258 mg).

¹H-NMR (400 MHz, CDCl₃) δ: 7.93-7.85 (1H, m), 7.57 (1H, d, J = 7.9 Hz), 7.29 (1H, d, J = 8.5 Hz), 7.12 (1H, t, J = 7.3 Hz), 7.03
20 (1H, t, J = 7.0 Hz), 6.91 (1H, s), 4.47-4.33 (1H, m), 3.80-3.08 (4H, m), 3.05-2.87 (2H, m), 2.83-2.73 (1H, m), 2.59-2.43 (2H, m), 2.30-2.13 (2H, m), 2.06-1.93 (2H, m), 1.83-1.66 (2H, m), 1.29 (3H, t, J = 6.4 Hz), 1.21-1.06 (3H, m).

[0163]

Example 23

(2S)-N-Ethyl-2-hydroxy-N-{2-[4-(1H-indol-3-yl)piperidin-1-yl]ethyl}propanamide



5 To a suspension of the compound of Reference example 8 (300 mg) in tetrahydrofuran (8.7 mL) were added triethylamine (0.606 mL), L-lactic acid (94.0 mg), and O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (398 mg). The mixture was stirred at room temperature for 3 hours, and methanol
10 (6.0 mL) and 2 mol/L aqueous sodium hydroxide (3.0 mL) were added thereto. The mixture was stirred at room temperature for 1 hour. Then, the reaction mixture was concentrated, and chloroform (30 mL) was added to the residue. The mixture was dried over anhydrous sodium sulfate, filtered, and concentrated. The
15 concentrated residue was purified by silica gel column chromatography (chloroform/methanol) to obtain the titled compound (258 mg).







¹H-NMR (400 MHz, CDCl₃) δ: 8.06-7.92 (1H, m), 7.61 (1H, d, J = 7.9 Hz), 7.35 (1H, d, J = 7.9 Hz), 7.17 (1H, t, J = 7.3 Hz), 7.08
20 (1H, t, J = 7.6 Hz), 6.99-6.94 (1H, m), 4.51-4.36 (1H, m), 3.82-1.74 (16H, m), 1.34 (3H, t, J = 6.1 Hz), 1.26-1.11 (3H, m).








[0164]






Examples 24 to 50






According to the method of Example 23, the compounds of





Examples 24 to 50 were prepared from the corresponding Reference example compounds.

Example	Chemical structure	Instrumental analysis data
24		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.74 (1H, dd, $J = 8.8, 5.2$ Hz), 7.21 (1H, dd, $J = 8.5, 1.8$ Hz), 7.05-7.00 (1H, m), 3.83-3.74 (2H, m), 3.45-3.31 (2H, m), 3.19-3.01 (3H, m), 2.68-2.59 (2H, m), 2.51-2.37 (2H, m), 2.25-2.09 (2H, m), 2.05-1.95 (2H, m), 1.19-1.06 (6H, m), 0.96-0.90 (2H, m).
25		LC-MS : R.T. = 1.142 min ObsMS = 346 [M+1]
26		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.65 (1H, d, $J = 7.9$ Hz), 7.49-7.42 (2H, m), 7.24-7.19 (1H, m), 4.43 (1H, td, $J = 13.3, 6.5$ Hz), 3.75-3.19 (8H, m), 2.78-2.57 (5H, m), 1.36-1.30 (3H, m), 1.25-1.11 (3H, m).
27		LC-MS : R.T. = 1.167 min ObsMS = 365 [M+1]
28		LC-MS : R.T. = 1.450 min ObsMS = 374 [M+1]
29		LC-MS : R.T. = 1.258 min ObsMS = 378 [M+1]

30		LC-MS : R.T. = 1.467 min ObsMS = 390 [M+1]
31		LC-MS : R.T. = 1.450 min ObsMS = 378 [M+1]
32		LC-MS : R.T. = 1.314 min ObsMS = 375 [M+1]
33		LC-MS : R.T. = 1.400 min ObsMS = 418 [M+1]
34		LC-MS : R.T. = 1.333 min ObsMS = 418 [M+1]
35		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.70-7.63 (1H, m), 7.23-7.19 (1H, m), 7.07-7.00 (1H, m), 4.06-3.82 (3H, m), 3.42-3.30 (2H, m), 3.15-2.99 (3H, m), 2.62-2.48 (4H, m), 2.30-2.21 (2H, m), 2.09-2.01 (5H, m), 1.24-1.12 (6H, m).
36		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.69-7.63 (1H, m), 7.24-7.20 (1H, m), 7.07-7.00 (1H, m), 4.03-3.92 (2H, m), 3.68-3.57 (2H, m), 3.55-3.22 (3H, m), 3.12-3.00 (3H, m), 2.66-2.55 (2H, m), 2.31-2.20 (2H, m), 2.10-1.98 (4H, m), 1.26-1.15 (3H, m).

37		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.78-7.70 (1H, m), 7.23-7.19 (1H, m), 7.07-7.00 (1H, m), 4.11-4.00 (1H, m), 3.76-3.51 (4H, m), 3.37-2.92 (6H, m), 2.67-1.93 (8H, m), 1.24-1.07 (6H, m).
38		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.72-7.63 (1H, m), 7.23-7.20 (1H, m), 7.07-7.00 (1H, m), 4.58-4.36 (1H, m), 3.81-2.94 (7H, m), 2.43-2.33 (2H, m), 2.20-1.98 (6H, m), 1.84-1.63 (3H, m), 1.35-1.30 (3H, m), 1.24-1.10 (3H, m).
39		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.70-7.56 (1H, m), 7.19-7.14 (1H, m), 7.03-6.95 (1H, m), 4.49-4.33 (1H, m), 3.75-3.43 (2H, m), 3.31-2.81 (6H, m), 2.59-2.44 (2H, m), 2.32-1.88 (6H, m), 1.64-1.47 (2H, m), 1.33-1.22 (3H, m), 0.90-0.80 (3H, m).
40		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.75-7.62 (1H, m), 7.23-7.20 (1H, m), 7.07-7.00 (1H, m), 3.75-2.92 (8H, m), 2.62-2.51 (2H, m), 2.37-1.94 (6H, m), 1.37-1.27 (3H, m), 1.27-1.11 (3H, m).
41		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.74-7.63 (1H, m), 7.23-7.18 (1H, m), 7.08-6.99 (1H, m), 4.53-4.38 (1H, m), 3.97-3.82 (2H, m), 3.54-3.21 (2H, m), 3.19-2.95 (3H, m), 2.63-2.49 (2H, m), 2.34-2.18 (2H, m), 2.15-1.97 (4H, m), 1.38-1.28 (3H, m), 1.28-1.11 (6H, m).


42		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.72-7.63 (1H, m), 7.23-7.18 (1H, m), 7.07-7.00 (1H, m), 4.52-4.39 (1H, m), 3.97-3.84 (1H, m), 3.53-3.21 (2H, m), 3.16-2.96 (3H, m), 2.61-2.50 (2H, m), 2.34-2.20 (2H, m), 2.12-1.96 (4H, m), 1.78-1.62 (1H, m), 1.36-1.28 (3H, m), 1.28-1.17 (6H, m).
43		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.68-7.61 (1H, m), 7.23-7.19 (1H, m), 7.07-6.99 (1H, m), 3.73-3.58 (1H, m), 3.49-3.31 (4H, m), 3.11-2.94 (3H, m), 2.57-2.49 (2H, m), 2.44-2.16 (3H, m), 2.11-1.97 (6H, m), 1.84-1.52 (6H, m), 1.35-1.05 (4H, m).
44		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.69-7.61 (1H, m), 7.23-7.18 (1H, m), 7.07-7.00 (1H, m), 4.05-3.97 (1H, m), 3.50-3.31 (4H, m), 3.13-2.97 (3H, m), 2.59-2.42 (3H, m), 2.31-2.17 (2H, m), 2.10-1.81 (7H, m), 1.71-1.42 (6H, m), 1.23-1.05 (3H, m).
45		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.52-7.46 (1H, m), 7.44-7.31 (1H, m), 7.30-7.25 (1H, m), 4.57-4.37 (1H, m), 3.80-2.91 (8H, m), 2.74-2.49 (2H, m), 2.39-1.98 (6H, m), 1.37-1.29 (3H, m), 1.26-1.10 (3H, m).
46		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.77-7.61 (1H, m), 7.24-7.20 (1H, m), 7.09-7.01 (1H, m), 4.24-4.13 (1H, m), 3.72-1.94 (20H, m), 1.75-1.51 (1H, m), 1.20-1.07 (3H, m).

47		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 8.05-7.94 (1H, m), 7.61 (1H, d, $J = 7.9$ Hz), 7.35 (1H, d, $J = 7.9$ Hz), 7.17 (1H, t, $J = 7.0$ Hz), 7.09 (1H, t, $J = 7.3$ Hz), 6.97 (1H, dd, $J = 6.1, 2.4$ Hz), 3.92-3.56 (4H, m), 3.46-1.43 (15H, m), 1.24-1.11 (3H, m).
48		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 8.12-7.97 (1H, m), 7.50 (1H, dd, $J = 8.5, 5.5$ Hz), 7.02 (1H, dd, $J = 9.5, 2.1$ Hz), 6.96-6.91 (1H, m), 6.85 (1H, td, $J = 9.2, 2.0$ Hz), 3.95-3.52 (4H, m), 3.50-1.45 (16H, m), 1.22-1.10 (3H, m).
49		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 8.00 (1H, br s), 7.61 (1H, d, $J = 7.9$ Hz), 7.35 (1H, d, $J = 7.9$ Hz), 7.17 (1H, t, $J = 7.6$ Hz), 7.08 (1H, t, $J = 7.3$ Hz), 6.99-6.94 (1H, m), 4.74-4.34 (2H, m), 3.87-1.71 (15H, m), 1.34 (3H, t, $J = 6.4$ Hz), 1.25-1.11 (3H, m).
50		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.29-7.22 (2H, m), 6.94 (1H, td, $J = 9.2, 2.4$ Hz), 6.47 (1H, d, $J = 3.7$ Hz), 4.53-4.12 (2H, m), 3.80-3.64 (2H, m), 3.56-2.88 (5H, m), 2.69-2.54 (2H, m), 2.42-2.25 (2H, m), 2.15-1.98 (4H, m), 1.38-1.33 (3H, m), 1.29-1.13 (3H, m).

[0165]

Example 51

N-Ethyl-N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}-2-hydroxyethane-1-sulfonamide



To a solution of the compound of Reference example 23 (636 mg) in tetrahydrofuran (20 mL) was added 10 % aqueous tetrabutylammonium hydroxide (17.3 mL). The mixture was stirred at 60°C for 2 hours, and concentrated. The residue was purified by silica gel column chromatography (chloroform/methanol), and further purified by amino silica gel column chromatography (chloroform/methanol) to obtain the titled compound (330 mg).





¹H-NMR (400 MHz, CDCl₃) δ: 7.79 (1H, dd, J = 8.9, 5.2 Hz), 7.22 (1H, dd, J = 8.5, 2.4 Hz), 7.04 (1H, td, J = 8.9, 2.2 Hz), 5.94 (1H, br s), 4.03-3.95 (2H, m), 3.60-3.53 (2H, m), 3.33-3.24 (4H, m), 3.21-3.09 (3H, m), 2.59-2.52 (2H, m), 2.31-2.13 (4H, m), 2.11-2.00 (2H, m), 1.25 (3H, t, J = 7.0 Hz).






[0166]




Examples 52 to 64

According to the method of Example 51, the compounds of Examples 52 to 64 were prepared from the corresponding Reference example compounds.

Example	Chemical structure	Instrumental analysis data
52		¹ H-NMR (400 MHz, CDCl ₃) δ: 7.90-7.81 (1H, m), 7.17 (1H, dd, J = 8.2, 2.1 Hz), 7.07 (1H, td, J = 9.0, 2.2 Hz), 3.65-2.21 (17H, m), 1.67-1.56 (1H, m), 1.43-1.31 (1H, m), 1.26 (3H, d, J = 6.7 Hz), 1.09 (3H, t, J = 7.0 Hz).

53		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.69–7.60 (1H, m), 7.19–7.15 (1H, m), 7.04–6.96 (1H, m), 4.03–3.95 (2H, m), 3.32–3.22 (4H, m), 3.16–3.09 (2H, m), 3.09–2.96 (3H, m), 2.92–2.81 (1H, m), 2.47–2.36 (2H, m), 2.27–1.93 (6H, m), 1.87–1.74 (2H, m), 1.17 (3H, t, $J = 7.3$ Hz).
54		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 7.91–7.88 (1H, m), 7.61–7.58 (2H, m), 7.40–7.32 (1H, m), 3.91 (1H, dd, $J = 11.6, 4.9$ Hz), 3.68–3.62 (1H, m), 3.39–3.08 (9H, m), 2.52–2.45 (2H, m), 2.31–2.22 (2H, m), 2.16–2.03 (4H, m), 1.92–1.82 (2H, m), 1.35 (3H, d, $J = 6.7$ Hz), 1.24 (3H, t, $J = 7.3$ Hz).
55		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 7.89 (1H, dd, $J = 8.9, 5.2$ Hz), 7.25 (1H, dd, $J = 8.9, 2.1$ Hz), 7.08 (1H, td, $J = 9.2, 2.4$ Hz), 3.91 (1H, dd, $J = 11.0, 4.9$ Hz), 3.67–3.61 (1H, m), 3.58–3.51 (4H, m), 3.40–3.23 (6H, m), 2.72–2.63 (4H, m), 2.52–2.44 (2H, m), 1.92–1.80 (2H, m), 1.35 (3H, d, $J = 6.7$ Hz), 1.27–1.19 (3H, m).
56		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 7.91 (1H, dd, $J = 8.9, 5.2$ Hz), 7.38 (1H, dd, $J = 8.9, 2.1$ Hz), 7.17 (1H, td, $J = 9.0, 2.0$ Hz), 3.91 (1H, dd, $J = 11.6, 4.9$ Hz), 3.64 (1H, dd, $J = 11.6, 6.7$ Hz), 3.39–3.05 (8H, m), 2.51–2.44 (2H, m), 2.30–2.21 (2H, m), 2.15–1.99 (5H, m), 1.91–1.81 (2H, m), 1.35 (3H, d, $J = 6.7$ Hz), 1.23 (3H, t, $J = 7.0$ Hz).


57		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.96 (1H, s), 7.59 (1H, d, $J = 7.9$ Hz), 7.30 (1H, d, $J = 7.9$ Hz), 7.15-7.09 (1H, m), 7.07-7.01 (1H, m), 6.96-6.91 (1H, m), 3.96-3.87 (2H, m), 3.72-3.51 (3H, m), 3.35-3.12 (6H, m), 2.93-1.85 (9H, m), 1.25-1.13 (3H, m).
58		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.65 (1H, d, $J = 7.3$ Hz), 7.18 (1H, d, $J = 9.1$ Hz), 5.98 (1H, br s), 4.04-3.96 (2H, m), 3.60-3.54 (2H, m), 3.34-3.26 (4H, m), 3.21-3.06 (3H, m), 2.61-2.51 (2H, m), 2.36-2.32 (3H, m), 2.30-2.14 (4H, m), 2.10-1.98 (2H, m), 1.25 (3H, t, $J = 7.0$ Hz).
59		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.61 (1H, s), 7.42 (1H, d, $J = 8.5$ Hz), 7.32 (1H, dd, $J = 8.5, 1.8$ Hz), 6.13-5.86 (1H, m), 4.04-3.98 (2H, m), 3.61-3.55 (2H, m), 3.37-3.24 (4H, m), 3.23-3.09 (3H, m), 2.62-2.53 (2H, m), 2.45 (3H, s), 2.37-2.17 (4H, m), 2.13-1.99 (2H, m), 1.25 (3H, t, $J = 7.0$ Hz).
60		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.72 (1H, d, $J = 8.5$ Hz), 7.53-7.49 (1H, m), 7.24-7.20 (1H, m), 5.80 (1H, br s), 4.00-3.90 (2H, m), 3.57-3.48 (2H, m), 3.31-3.18 (4H, m), 3.18-3.05 (3H, m), 2.63-2.44 (2H, m), 2.36-2.08 (4H, m), 2.08-1.96 (2H, m), 1.20 (3H, t, $J = 7.0$ Hz).
61		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.78 (1H, d, $J = 7.9$ Hz), 7.52-7.44 (2H, m), 7.26-7.20 (1H, m), 5.88 (1H, br s), 3.98-3.92 (2H, m), 3.58-3.49 (2H, m), 3.31-

		3.19 (4H, m), 3.19-3.05 (3H, m), 2.57-2.49 (2H, m), 2.33-2.12 (4H, m), 2.10-1.98 (2H, m), 1.20 (3H, t, J = 7.0 Hz).
62		¹ H-NMR (400 MHz, CDCl ₃) δ: 8.06 (1H, s), 7.53 (1H, dd, J = 8.9, 5.2 Hz), 7.03 (1H, dd, J = 9.5, 2.1 Hz), 6.97-6.92 (1H, m), 6.89-6.81 (1H, m), 4.00-3.92 (2H, m), 3.65-3.55 (2H, m), 3.34-3.15 (6H, m), 2.91-1.58 (10H, m), 1.29-1.20 (3H, m).
63		¹ H-NMR (400 MHz, CDCl ₃) δ: 7.92 (1H, s), 7.43 (1H, dd, J = 9.2, 3.7 Hz), 7.33 (1H, dd, J = 8.5, 2.4 Hz), 7.13 (1H, td, J = 9.2, 2.4 Hz), 4.58-4.38 (1H, m), 4.08-4.00 (2H, m), 3.64-3.50 (2H, m), 3.40-3.13 (6H, m), 2.76-2.53 (2H, m), 2.51-2.22 (4H, m), 2.18-1.98 (1H, m), 1.79-1.47 (2H, m), 1.25 (3H, t, J = 7.3 Hz).
64		¹ H-NMR (400 MHz, CDCl ₃) δ: 7.92 (1H, s), 7.47-7.39 (1H, m), 7.33 (1H, dd, J = 8.5, 2.4 Hz), 7.17-7.10 (1H, m), 4.55-4.36 (1H, m), 3.98-3.68 (3H, m), 3.52-2.98 (5H, m), 2.79-2.14 (4H, m), 2.14-1.96 (1H, m), 1.69-1.46 (5H, m), 1.34 (3H, d, J = 7.3 Hz), 1.25 (3H, t, J = 7.0 Hz).

[0167]

Example 65

3-Amino-N-ethyl-N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}-2-hydroxypropanamide



To a solution of 3-((tert-butoxycarbonyl)amino)-2-hydroxypropionic acid (40.6 mg) in N,N-dimethylformamide (2.0 mL) was added O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (75.0 mg). The mixture was stirred at room temperature for 1 hour, and triethylamine (0.053 mL) and the compound of Reference example 1 (60.0 mg) were added thereto. The mixture was stirred at room temperature for 3 hours, and water (10 mL) was added thereto. The mixture was extracted with chloroform/methanol (10 mL x 6), dried over anhydrous sodium sulfate, filtered, and concentrated. The concentrated residue was purified by silica gel column chromatography (chloroform/methanol). Then, to the resulted compound (50 mg) was added 4 mol/L hydrochloric acid-ethyl acetate (2.0 mL). The mixture was stirred at room temperature for 1 hour, and concentrated. The concentrated residue was purified by amino silica gel column chromatography (chloroform/methanol) to obtain the titled compound (37.0 mg).






¹H-NMR (400 MHz, CDCl₃) δ: 7.77-7.58 (1H, m), 7.20 (1H, d, J = 7.9 Hz), 7.08-6.91 (1H, m), 4.76-4.24 (1H, m), 3.86-2.69 (9H, m), 2.69-2.41 (4H, m), 2.41-1.74 (7H, m), 1.35-0.89 (3H, m).

[0168]

Examples 66 to 70

According to the method of Example 65, the compounds of


Examples 66 to 70 were prepared by the corresponding Reference example compounds.

Example	Chemical structure	Instrumental analysis data
66		LC-MS: R.T. = 1.167 min ObsMS = 405 [M+1]
67		LC-MS: R.T. = 1.200 min ObsMS = 419 [M+1]
68		LC-MS: R.T. = 1.200 min ObsMS = 405 [M+1]
69		LC-MS: R.T. = 1.300 min ObsMS = 419 [M+1]
70		LC-MS: R.T. = 1.100 min ObsMS = 405 [M+1]

[0169]

Example 71

- 5 3-Acetamide-N-ethyl-N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}-2-hydroxypropanamide






To a solution of the compound of Example 65 (20.0 mg) in tetrahydrofuran (2.0 mL) was added anhydrous acetic acid (0.00598 mL). The mixture was stirred at room temperature for 1 hour. Then, 2 mol/L aqueous sodium hydroxide (0.1 mL) was added thereto, and the mixture was stirred at room temperature for 10 minutes. The reaction mixture was concentrated, and purified by amino silica gel column chromatography (chloroform/methanol) to obtain the titled compound (10.1 mg).

¹H-NMR (400 MHz, CDCl₃) δ: 7.77-7.55 (1H, m), 7.19-7.14 (1H, m), 7.05-6.97 (1H, m), 6.19-6.01 (1H, m), 4.49-4.34 (1H, m), 4.03-3.47 (4H, m), 3.47-2.84 (6H, m), 2.72-2.48 (2H, m), 2.43-2.19 (2H, m), 2.16-1.97 (4H, m), 1.96-1.90 (3H, m), 1.24-1.06 (3H, m).
[0170]

Examples 72 to 74


According to the method of Example 71, the compounds of Examples 72 to 74 were prepared from the corresponding Reference example compounds.

Example	Chemical structure	Instrumental analysis data
72		LC-MS: R.T. = 1.258 min ObsMS = 461 [M+1]
73		LC-MS: R.T. = 1.350 min ObsMS = 447 [M+1]
74		LC-MS: R.T. = 1.270 min ObsMS = 461 [M+1]

[0171]

Example 75

N-Ethyl-N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}-4-hydroxy-1-methylpyrrolidine-3-carboxamide



5

To a solution of the compound of Example 66 (10.0 mg) in methanol (0.5 mL) were 37 % aqueous formaldehyde (0.002 mL), sodium cyanoborohydride (1.09 mg), and acetic acid (0.003 mL). The mixture was stirred at room temperature for 16 hours. Then, water (10 mL) was added thereto, and the mixture was extracted with chloroform (10 mL x 2) and concentrated. The concentrated residue was purified by high performance liquid chromatography


10

(column; semipreparative C-18 column, separation condition; acetonitrile/trifluoroacetic acid: water/trifluoroacetic acid), and desalted with MP-Carbonate resin to obtain the titled compound (6.88 mg).

5 LC-MS: R.T. = 1.083 min ObsMS = 419 [M+1]
[0172]

Example 76

N-Ethyl-N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}-4-hydroxy-1-methylpiperidine-3-carboxamide



10

According to a similar method to Example 75, the titled compound was prepared from the compound of Example 67.

LC-MS: R.T. = 1.092 min ObsMS = 433 [M+1]
[0173]

15 Example 77

N-Ethyl-N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}-3-hydroxypropanamide orotate

To a solution of the compound of Example 17 (30 mg) in methanol (0.4 mL) were added orotic acid (14.4 mg). The mixture
20 was stirred at 60°C for 1 hour, and then ethyl acetate (2.0 mL) was added thereto. The mixture was stirred at room temperature for 1 hour, and the precipitated solid was collected by filtration to obtain the titled compound (27.3 mg).

¹H-NMR (400 MHz, DMSO-D₆) δ: 11.02 (1H, s), 9.94 (1H, brs), 8.03

(1H, dd, J = 8.8, 5.2 Hz), 7.74-7.68 (1H, m), 7.35-7.27 (1H, m),
 5.79 (1H, d, J = 1.8 Hz), 4.53 (1H, brs), 3.65 (2H, t, J = 6.7
 Hz), 3.62-3.49 (3H, m), 3.48-3.19 (6H, m), 3.18-2.62 (4H, m),
 2.28-2.13 (2H, m), 2.13-1.91 (2H, m), 1.12 (2H, t, J = 7.0 Hz),
 1.02 (1H, t, J = 7.0 Hz).

Powder XRD ($^{\circ}$, $2\theta \pm 0.2$)

4.87, 7.24, 15.27, 15.75, 16.02, 16.39, 16.73, 17.62, 19.51,
 20.04, 21.24, 22.05, 22.34, 24.23, 24.80 (Among them, the
 characteristic 10 peaks were 4.87, 7.24, 15.27, 15.75, 16.02,
 17.62, 19.51, 20.04, 21.24, 22.05, and the more characteristic 4
 peaks were 4.87, 19.51, 20.04, 21.24.)

DSC

Enthalpy (normalized): 62.81 J/g

Onset x: 157.09 $^{\circ}$ C

[0174]

Example 78

N-Ethyl-N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-
 yl]ethyl}-3-hydroxypropanamide hydrobromide in Form A

To a solution of the compound of Example 17 (6.03 g) in
 acetonitrile (77 mL) were added dropwise 10 to 20 % HBr-EtOH
 (14.1 g). The mixture was stirred at room temperature for 1 hour,
 and precipitation of a solid was confirmed. The mixture was
 stirred at 80 $^{\circ}$ C. After confirming that the precipitated solid
 was completely dissolved, the mixture was stirred for 1 hour.
 Then, the mixture was gradually cooled, and stirred at room
 temperature for 12 hours. The precipitated solid was collected
 by filtration, washed with acetonitrile (10 mL), and dried to

obtain a white solid (7.0 g). The resulted white solid (7.0 g) was mixed with tetrahydrofuran (15 mL), and the mixture was stirred at 80°C for 3.5 hours. Then, the mixture was gradually cooled, and stirred at room temperature for 12 hours. The solid
5 was collected by filtration, washed with tetrahydrofuran (10 mL), and dried to obtain the titled compound (6.20 g).

¹H-NMR (400 MHz, DMSO-D₆) δ: 9.48 (0.2H, brs), 9.19 (0.8H, brs), 8.11-7.99 (1H, m), 7.78-7.70 (1H, m), 7.40-7.30 (1H, m), 4.53 (1H, brs), 3.78-3.56 (6H, m), 3.55-3.09 (8H, m), 2.40-2.18 (3H, m),
10 m), 2.15-1.97 (2H, m), 1.19-0.97 (3H, m).

Powder XRD(°, 2θ±0.2)

4.48, 8.98, 12.40, 13.63, 15.23, 16.35, 17.07, 17.88, 18.06, 21.19, 23.13, 23.83, 24.48, 25.22, 25.98 (Among them, the characteristic 10 peaks were 4.48, 12.40, 15.23, 17.07, 17.88,
15 21.19, 23.13, 24.48, 25.22, 25.98, and the more characteristic 4 peaks were 12.40, 15.23, 17.07, 21.19.)

DSC

Enthalpy (normalized): 82.83 J/g

Onset x: 163.64°C

20 [0175]

Example 79

N-Ethyl-N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}-3-hydroxypropanamide hydrobromide in Form B

To a solution of the compound of Example 17 (100 mg) in
25 acetonitrile (2.0 mL) was added dropwise 10-20 % HBr-EtOH (0.2 mL). The mixture was stirred at room temperature for 1 hour, and the precipitated solid was collected by filtration to obtain the

titled compound (67.0 mg).

Powder XRD($^{\circ}$, $2\theta \pm 0.2$)

7.25, 7.65, 9.90, 11.17, 12.48, 15.34, 15.71, 22.36, 23.36, 23.74, 24.62, 25.05, 25.18, 25.26, 29.36 (Among them, the characteristic
5 10 peaks were 7.25, 7.65, 9.90, 15.34, 15.71, 23.36, 24.62, 25.05, 25.18, 25.26, and the more characteristic 4 peaks were 7.25, 7.65, 9.90, 15.34.)

DSC

Enthalpy (normalized): 86.51 J/g

10 Onset x: 164.04 $^{\circ}$ C

[0176]

Example 80

N-Ethyl-N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}-3-hydroxypropanamide tosylate

15 To a solution of the compound of Example 17 (66.7 g) in acetone (200 mL) was added a solution of p-toluenesulfonic acid monohydrate (41.9 g) in acetone (150 mL). The mixture was stirred at room temperature for 2 hours. The precipitated solid was collected by filtration, washed with acetone (60 mL), and dried
20 to obtain a white solid (78.8 g). The resulted white solid (78.8 g) and a mixture of acetone (608 mL) and water (33.4 mL) were stirred at 70 $^{\circ}$ C. After confirming that the white solid was completely dissolved, the mixture was cooled gradually, and stirred at room temperature for 12 hours. After stirring at 0 $^{\circ}$ C
25 for 1 hour, the resulted solid was collected by filtration, washed with a mixture solvent of 5 % water-acetone (70 mL) which was cooled to 0 $^{\circ}$ C, and dried to obtain the titled compound (69.4 g).

¹H-NMR (400 MHz, DMSO-D₆) δ: 9.39-9.00 (1H, m), 8.08-7.99 (1H, m),
 7.77-7.71 (1H, m), 7.46 (2H, d, J = 7.9 Hz), 7.39-7.30 (1H, m),
 7.10 (2H, d, J = 7.9 Hz), 4.53 (1H, brs), 3.76-3.56 (6H, m),
 3.54-3.11 (8H, m), 2.38-2.17 (6H, m), 2.12-1.96 (2H, m), 1.16-
 5 0.98 (3H, m).

Powder XRD(°, 2θ±0.2)

6.87, 7.03, 9.64, 13.74, 15.19, 15.37, 15.68, 16.22, 19.82, 21.60,
 21.67, 22.10, 23.15, 23.88, 27.67 (Among them, the characteristic
 10 peaks were 7.03, 9.64, 15.19, 15.37, 15.68, 16.22, 21.60,
 10 21.67, 23.15, 23.88, and the more characteristic 4 peaks were
 7.03, 9.64, 15.19, 15.68.)

DSC


Enthalpy (normalized): 99.47 J/g

Onset x: 162.26°C

15 [0177]

Reference example 1

N-Ethyl-2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethane-1-amine dihydrochloride



20 To a solution of 6-fluoro-3-(4-piperidinyl)-1,2-benzisoxazole (5.78 g) in chloroform (131 mL) were added tert-butyl ethyl(2-oxoethyl)carbamate (4.91 g) and sodium triacetoxyborohydride (11.1 g). The mixture was stirred at room temperature for 5 hours. Saturated aqueous sodium bicarbonate


(100 mL) were added thereto, and the mixture was extracted with chloroform (100 mL x 2), dried over anhydrous sodium sulfate, filtered, and concentrated. The residue was purified by silica gel column chromatography (chloroform/methanol). To a solution of the resulted compound in chloroform (131 mL) was added 4 mol/L hydrochloric acid-ethyl acetate (60 mL). The mixture was stirred at room temperature for 2 hours. The precipitated solid was collected by filtration, washed with ethyl acetate (20 mL x 2), and dried to obtain the titled compound (6.24 g).







¹H-NMR (400 MHz, DMSO-D₆) δ: 9.31 (2H, brs), 8.18 (1H, dd, J = 8.5, 5.5 Hz), 7.74 (1H, dd, J = 8.9, 2.1 Hz), 7.35 (1H, ddd, J = 9.0, 9.0, 2.2 Hz), 3.81-3.72 (2H, m), 3.55-3.46 (5H, m), 3.28-3.15 (2H, m), 3.08-2.97 (2H, m), 2.43-2.23 (4H, m), 1.25 (3H, t, J = 7.3 Hz).






[0178]

Reference examples 2 to 13

According to a similar method of Reference example 1, the compounds of Reference examples 2 to 13 were prepared from the corresponding Reference example compounds or reagents.

Reference example	Chemical structure	Instrumental analysis data
2		¹ H-NMR (400 MHz, CD ₃ OD) δ: 8.07-7.98 (1H, m), 7.76-7.60 (1H, m), 7.48-7.39 (1H, m), 7.25 (1H, ddd, J = 9.1, 9.1, 1.8 Hz), 5.06-4.94 (1H, m), 3.94-3.84 (2H, m), 3.64-3.55 (4H, m), 3.49-3.37 (2H, m), 3.19 (2H, q, J = 7.3 Hz), 2.77-2.64 (2H, m), 2.36-2.25 (2H, m), 1.39 (3H, t, J = 7.3 Hz).


3		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.55-7.47 (1H, m), 7.24 (1H, s), 7.18 (1H, d, J = 9.2 Hz), 3.47-2.94 (8H, m), 2.66-2.45 (2H, m), 2.34 (3H, d, J = 1.8 Hz), 2.30-2.16 (1H, m), 2.15-1.94 (4H, m), 1.10 (3H, t, J = 7.0 Hz).
4		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 7.81 (1H, s), 7.53-7.43 (2H, m), 3.92-3.83 (2H, m), 3.72-3.42 (5H, m), 3.40-3.32 (2H, m), 3.20 (2H, q, J = 7.1 Hz), 2.56-2.36 (7H, m), 1.39 (3H, t, J = 7.3 Hz).
5		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.00 (1H, d, J = 8.5 Hz), 7.74 (1H, s), 7.41 (1H, dd, J = 8.5, 1.8 Hz), 3.92-3.81 (2H, m), 3.71-3.44 (5H, m), 3.40-3.32 (2H, m), 3.20 (2H, q, J = 6.9 Hz), 2.64-2.36 (4H, m), 1.39 (3H, t, J = 7.3 Hz).
6		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.06 (1H, dd, J = 8.8, 5.2 Hz), 7.26 (1H, d, J = 9.1 Hz), 7.06 (1H, dd, J = 9.1, 9.1 Hz), 3.92-3.80 (2H, m), 3.67-3.50 (5H, m), 3.42-3.33 (2H, m), 3.20 (2H, q, J = 7.1 Hz), 2.57-2.43 (2H, m), 2.41-2.27 (2H, m), 1.40 (3H, t, J = 7.3 Hz).
7		$^1\text{H-NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ : 10.98 (1H, s), 10.72 (1H, s), 9.20 (2H, s), 7.67 (1H, dd, J = 8.5, 5.5 Hz), 7.15-7.09 (2H, m), 6.88-6.80 (1H, m), 3.74-3.65 (2H, m), 3.49-3.39 (4H, m), 3.21-3.09 (2H, m), 3.08-2.93 (3H, m), 2.18-2.07 (4H, m), 1.23 (3H, t, J = 7.0 Hz).
8		$^1\text{H-NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ : 10.89 (1H, s), 10.70 (1H, s), 9.20 (2H, s), 7.67 (1H, d, J = 7.9 Hz), 7.35 (1H, d, J = 8.5 Hz), 7.12 (1H, d, J = 2.4 Hz), 7.09-7.04 (1H, m), 6.99-6.94 (1H, m), 3.75-3.65 (2H, m), 3.52-3.40 (4H, m), 3.22-3.11 (2H, m), 3.07-2.94 (3H, m), 2.22-2.07 (4H, m), 1.23 (3H, t, J = 7.3 Hz).

9		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 8.03-7.92 (1H, m), 7.67-7.61 (2H, m), 7.43-7.37 (1H, m), 3.92-3.73 (2H, m), 3.64-3.42 (5H, m), 3.38-3.31 (2H, m), 3.19 (2H, q, $J = 7.3$ Hz), 2.55-2.33 (4H, m), 1.38 (3H, t, $J = 7.3$ Hz).
10		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 7.90 (1H, d, $J = 7.9$ Hz), 7.63-7.58 (1H, m), 7.53 (1H, d, $J = 8.5$ Hz), 7.34 (1H, dd, $J = 7.6, 7.6$ Hz), 4.16-3.72 (4H, m), 3.69-3.40 (8H, m), 3.19 (2H, q, $J = 7.1$ Hz), 1.38 (3H, t, $J = 7.3$ Hz).
11		$^1\text{H-NMR}$ (400 MHz, CD_3OD) δ : 7.93 (1H, dd, $J = 9.2, 4.9$ Hz), 7.33 (1H, dd, $J = 8.9, 2.1$ Hz), 7.14 (1H, ddd, $J = 9.0, 9.0, 2.0$ Hz), 4.33-3.71 (4H, m), 3.70-3.41 (8H, m), 3.19 (2H, q, $J = 7.3$ Hz), 1.38 (3H, t, $J = 7.3$ Hz).
12		$^1\text{H-NMR}$ (400 MHz, $\text{DMSO}-d_6$) δ : 11.01 (1H, s), 9.20 (2H, s), 8.04-7.98 (1H, m), 7.81 (1H, dd, $J = 9.1, 4.3$ Hz), 7.57 (1H, ddd, $J = 9.1, 9.1, 2.4$ Hz), 3.82-3.70 (2H, m), 3.55-3.40 (4H, m), 3.26-3.12 (3H, m), 3.09-2.96 (2H, m), 2.42-2.22 (4H, m), 1.23 (3H, t, $J = 7.3$ Hz).
13		LC-MS: R.T. = 0.443 min ObsMS = 290 [M+1]

[0179]

Reference example 14

6-Fluoro-5-methyl-3-(piperidin-4-yl)-1,2-benzisoxazole
monohydrochloride



a) Preparation of tert-butyl 4-[methoxy(methyl)carbamoyl]piperidine-1-carboxylate (Compound IN-1-1)

The mixture of 1-(tert-butoxycarbonyl)piperidine-4-carboxylic acid (5.00 g), N,O-dimethylhydroxylamine hydrochloride (3.19 g), N1-((ethylimino)methylene)-N3,N3-dimethylpropane-1,3-diamine hydrochloride (5.02 g), triethylamine (4.41 g), and N,N-dimethylformamide (100 mL) was stirred at room temperature for 1.5 hours. Then, saturated aqueous ammonium chloride was added thereto, and the mixture was extracted with ethyl acetate. The combined organic layer was washed twice with saturated aqueous ammonium chloride, and with saturated aqueous sodium bicarbonate and brine, and dried over anhydrous sodium sulfate. The mixture was filtered and concentrated to obtain the titled compound (4.52 g).

¹H-NMR (400 MHz, CDCl₃) δ: 4.03-4.24 (2H, m), 3.72 (3H, s), 3.19 (3H, s), 2.70-2.86 (3H, m), 1.63-1.76 (4H, m), 1.46 (9H, s).

[0180]

b) Preparation of tert-butyl 4-(2,4-difluoro-5-methylbenzoyl)piperidine-1-carboxylate (Compound IN-1-2)

To a solution of 1-bromo-2,4-difluoro-5-methylbenzene (2.28 g) in tetrahydrofuran (36 mL) was added dropwise 1.63 mol/L n-butyllithium/hexane (7.43 mL) over 3 minutes at -78°C. After stirring at -78°C for 1 hour, Compound IN-1-1 (1.50 g) was added thereto, and the mixture was stirred for 2.5 hours at -78°C. Then, saturated aqueous ammonium chloride was added thereto, and the mixture was extracted with ethyl acetate, dried over anhydrous sodium sulfate, filtered, and concentrated. The concentrated residue was purified by silica gel column chromatography (hexane/ethyl acetate) to obtain the titled compound (2.01 g).

¹H-NMR (400 MHz, CDCl₃) δ: 7.17-7.25 (1H, m), 6.80-6.87 (1H, m), 1.83-1.92 (2H, m), 1.59-1.69 (2H, m), 1.45 (9H, s).

[0181]

c) Preparation of tert-butyl 4-(6-fluoro-5-methyl-1,2-benzoxazol-3-yl)piperidine-1-carboxylate (Compound IN-1-3)

A mixture of Compound IN-1-2 (731 mg), hydroxylamine hydrochloride (599 mg), and sodium acetate (707 mg) in ethanol (10 mL) was stirred at 60°C for 4 hours. Then, water was added thereto, and the mixture was extracted with ethyl acetate, washed with brine, dried over anhydrous sodium sulfate, filtered, and concentrated. The concentrated residue was purified by silica gel column chromatography (hexane/ethyl acetate). The resulted

product (335 mg) was mixed with cesium carbonate (615 mg) and acetonitrile (9.0 mL), and the mixture was stirred in a sealed tube at 130°C for 3.5 hours. Then, the reaction mixture was filtered, concentrated, and purified by silica gel column chromatography (hexane/ethyl acetate) to obtain the titled compound (90.8 mg).

¹H-NMR (400 MHz, CDCl₃) δ: 7.47 (1H, d, J = 7.1 Hz), 7.21 (1H, d, J = 9.0 Hz), 4.11-4.36 (2H, m), 3.16-3.26 (1H, m), 2.89-3.03 (2H, m), 2.38 (3H, d, J = 1.7 Hz), 2.01-2.10 (2H, m), 1.87-1.99 (2H, m).

[0182]

d) Preparation of 6-fluoro-5-methyl-3-(piperidin-4-yl)-1,2-benzisoxazole monohydrochloride (Reference example 14)


To a solution of Compound IN-1-3 (131 mg) in dichloromethane (1.0 mL) was added 4 mol/L hydrochloric acid/ethyl acetate (1.0 mL), and the mixture was stirred at room temperature for 1.5 hours. Then, the reaction mixture was concentrated to obtain the titled compound (114 mg).

¹H-NMR (400 MHz, CDCl₃) δ: 7.30 (1H, d, J = 7.3 Hz), 7.11 (1H, s), 7.06 (1H, d, J = 9.1 Hz), 4.12-4.04 (2H, m), 3.09-3.01 (1H, m), 2.84-2.75 (2H, m), 2.22 (3H, d, J = 1.8 Hz), 1.93-1.86 (2H, m), 1.84-1.71 (2H, m).

[0183]

Reference example 15

5-Methyl-3-(piperidin-4-yl)-1,2-benzisoxazole monohydrochloride




According to a similar method to Reference example 14, the titled compound was prepared from 1-fluoro-2-iodo-4-methylbenzene. ¹H-NMR (400 MHz, CDCl₃) δ: 7.48-7.43 (2H, m), 7.38-7.34 (1H, m), 4.29-4.20 (2H, m), 3.28-3.18 (1H, m), 3.02-2.92 (2H, m), 2.48 (3H, s), 2.11-2.03 (2H, m), 2.03-1.90 (2H, m).

[0184]

Reference example 16

N-(Cyclopropylmethyl)-2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethane-1-amine




To a solution of 2-(4-(6-fluorobenzo[d]isoxazol-3-yl)piperidin-1-yl)ethane-1-amine (50.0 mg) in chloroform (2.0 mL) was added cyclopropanecarbaldehyde (16.0 mg), and the mixture was stirred at room temperature for 30 minutes. Then, sodium triacetoxyborohydride (60.4 mg) was added thereto, and the mixture was stirred at room temperature for 1 hour. Saturated aqueous sodium bicarbonate (30 mL) was added thereto. The mixture was extracted with chloroform (30 mL x 2), dried over anhydrous sodium sulfate, filtered, and concentrated to obtain the titled compound (26.0 mg).

LC-MS: R.T. = 1.250 min ObsMS = 318 [M+1]

[0185]

Reference example 17

N-{2-[4-(6-Fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}propane-2-amine




According to a similar method to Reference example 16, the titled compound was prepared from acetone.

LC-MS: R.T. = 1.176 min ObsMS = 306 [M+1]

[0186]

Reference example 18

((2-[4-(6-Fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl)amino)acetonitrile



To a suspension of 55 % sodium hydride (7.92 mg) in N,N-dimethylformamide (2.0 mL) was added the compound of Reference example 20 (60.0 mg), and the mixture was stirred at room temperature for 30 minutes. 2-Iodoacetonitrile (41.3 mg) was added thereto. The reaction mixture was stirred at room temperature for 1 hour, and water (30 mL) was added thereto. The mixture was extracted with chloroform (30 mL x 2), dried over


anhydrous sodium sulfate, filtered, and concentrated. To the concentrated residue was added 4 mol/L hydrochloric acid/ethyl acetate (3.0 mL), the mixture was stirred at room temperature for 1 hour, and concentrated. The concentrated residue was purified
 5 by amino silica gel column chromatography (hexane/ethyl acetate) to obtain the titled compound (25.0 mg).

LC-MS: R.T. = 0.530 min ObsMS = 303 [M+1]

[0187]

Reference example 19

10 N-Ethyl-3-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]
 propane-1-amine dihydrochloride




According to a similar method to Reference example 18, the titled compound was prepared from the compound of Reference
 15 example 21.

[0188]

Reference example 20

Tert-butyl {2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}carbamate



To a solution (60 mL) of 6-fluoro-3-(piperidin-4-yl)benzo[d]isoxazole (2.89 g) in chloroform was added tert-butyl (2-oxoethyl)carbamate (2.30 g), and the mixture was stirred at


room temperature for 30 minutes. Sodium triacetoxyborohydride (3.34 g) was added thereto, and the mixture was stirred at room temperature for 3 hours. Saturated aqueous sodium bicarbonate (100 mL) was added thereto, and the mixture was extracted with
 5 chloroform (100 mL x 2), dried over anhydrous sodium sulfate, filtered, and concentrated. The concentrated residue was purified by silica gel column chromatography (chloroform/methanol) to obtain the titled compound (3.98 g).

¹H-NMR (400 MHz, CDCl₃) δ: 7.67 (1H, dd, J = 8.8, 5.2 Hz), 7.24-
 10 7.20 (1H, m), 7.04 (1H, ddd, J = 8.8, 8.8, 2.0 Hz), 5.02 (1H, s), 3.29-3.20 (2H, m), 3.10-2.98 (3H, m), 2.53-2.46 (2H, m), 2.22-2.13 (2H, m), 2.09-2.00 (4H, m), 1.45 (9H, s).

[0189]

Reference example 21

15 Tert-butyl {3-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]propyl}carbamate




To a solution of 6-fluoro-3-(piperidin-4-yl)benzo[d]isoxazole (2.00 g) in acetonitrile (50 mL) were added
 20 water (12.5 mL), tert-butyl(3-bromopropyl)carbamate (2.59 g), and potassium carbonate (3.76 g). The mixture was stirred at 60°C for 3.5 hours, and water (200 mL) was added thereto. The mixture was extracted with ethyl acetate (200 mL x 2), dried over anhydrous sodium sulfate, filtered, and concentrated. The

concentrated residue was purified by silica gel column chromatography (chloroform/methanol) to obtain the titled compound (3.46 g).

¹H-NMR (400 MHz, CDCl₃) δ: 7.67 (1H, dd, J = 8.8, 5.2 Hz), 7.22 (1H, dd, J = 8.5, 2.4 Hz), 7.02 (1H, ddd, J = 8.8, 8.8, 2.2 Hz), 5.50 (1H, s), 3.23-3.17 (2H, m), 3.12-3.00 (3H, m), 2.46 (2H, t, J = 6.7 Hz), 2.16-1.99 (6H, m), 1.72-1.64 (2H, m), 1.43 (9H, s).
[0190]

Reference example 22

10 N-{2-[4-(6-Fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}propane-1-amine



a) Preparation of 3-[1-(2-chloroethyl)piperidin-4-yl]-6-fluoro-1,2-benzisoxazole (Compound IN-2-1)

15 To a solution (25 mL) of 6-fluoro-3-(piperidin-4-yl)benzo[d]isoxazole (3.00 g) in tetrahydrofuran were added water (6.3 mL), potassium hydroxide (1.68 g), and 1-bromo-2-chloroethane (5.65 mL), and the mixture was stirred at room temperature for 24 hours. Water (15 mL) was added thereto. The mixture was extracted with chloroform (10 mL x 3), dried over anhydrous sodium sulfate, filtered, and concentrated. The concentrated residue was purified by silica gel column

20

chromatography (ethyl acetate/methanol) to obtain the titled compound (1.94 g).

¹H-NMR (400 MHz, CDCl₃) δ: 7.70 (1H, dd, J = 8.5, 5.3 Hz), 7.25 (1H, dd, J = 8.9, 2.5 Hz), 7.06 (1H, ddd, J = 8.8, 8.8, 2.1 Hz),
5 3.64 (2H, t, J = 6.9 Hz), 3.01-3.15 (3H, m), 2.81 (2H, t, J = 6.9 Hz), 2.24-2.39 (2H, m), 2.02-2.18 (4H, m).

[0191]

b) Preparation of N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}propane-1-amine (Reference example 22)


10 To a solution of Compound IN-2-1 (80.0 mg) in acetonitrile (1.4 mL) were added n-propylamine (0.0349 mL), potassium carbonate (117 mg), potassium iodide (9.39 mg), and the mixture was stirred at 90°C for 4 hours. The reaction mixture was filtered and concentrated. The concentrated residue was purified
15 by amino silica gel column chromatography (chloroform/methanol) to obtain the titled compound (50.9 mg).

¹H-NMR (400 MHz, CDCl₃) δ: 7.66 (1H, dd, J = 8.8, 5.2 Hz), 7.22 (1H, dd, J = 8.5, 2.4 Hz), 7.03 (1H, ddd, J = 8.8, 8.8, 2.2 Hz),
3.10-2.99 (3H, m), 2.72 (2H, t, J = 6.1 Hz), 2.58 (2H, t, J = 7.0
20 Hz), 2.53 (2H, t, J = 6.4 Hz), 2.21-1.99 (6H, m), 1.56-1.46 (2H, m), 0.91 (3H, t, J = 7.3 Hz).

[0192]

Reference example 23

N-Ethyl-N-{2-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperidin-1-yl]ethyl}ethenesulfonamide
25



To a solution of the compound of Reference example 1 (1.00 g) in dichloromethane (40 mL) were added triethylamine (2.39 mL) and 2-chloroethanesulfonyl chloride (0.433 mL), and the mixture was stirred at room temperature for 4 hours. The reaction mixture was concentrated, and the concentrated residue was purified by silica gel column chromatography (chloroform/methanol) to obtain the titled compound (0.670 g).





$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.65 (1H, dd, $J = 8.5, 4.9$ Hz), 7.22 (1H, dd, $J = 8.5, 2.4$ Hz), 7.04 (1H, ddd, $J = 8.9, 8.9, 2.2$ Hz), 6.54 (1H, dd, $J = 16.5, 9.8$ Hz), 6.20 (1H, d, $J = 16.5$ Hz), 5.89 (1H, d, $J = 9.8$ Hz), 3.31 (2H, t, $J = 7.0$ Hz), 3.25 (2H, q, $J = 7.1$ Hz), 3.11-3.00 (3H, m), 2.61 (2H, t, $J = 6.7$ Hz), 2.32-2.17 (2H, m), 2.10-1.97 (4H, m), 1.20 (3H, t, $J = 7.3$ Hz).





[0193]




Reference examples 24 to 36


According to the method of Reference example 23, the compounds of Reference examples 24 to 36 were prepared from the corresponding Reference example compounds.

Reference example	Chemical structure	Instrumental analysis data
24		LC-MS : R.T. = 1.566 min ObsMS = 396 [M+1]

25		¹ H-NMR (400 MHz, CDCl ₃) δ: 7.67 (1H, dd, J = 8.5, 5.5 Hz), 7.22 (1H, dd, J = 8.5, 1.8 Hz), 7.04 (1H, ddd, J = 8.9, 8.9, 2.0 Hz), 6.42 (1H, dd, J = 16.5, 9.8 Hz), 6.19 (1H, d, J = 16.5 Hz), 5.89 (1H, d, J = 9.8 Hz), 3.28-3.14 (4H, m), 3.12-2.96 (3H, m), 2.50-2.38 (2H, m), 2.24-1.99 (6H, m), 1.88-1.78 (2H, m), 1.19 (3H, t, J = 7.3 Hz).
26		¹ H-NMR (400 MHz, CDCl ₃) δ: 7.73 (1H, d, J = 7.9 Hz), 7.57-7.47 (2H, m), 7.30-7.24 (1H, m), 5.95 (1H, s), 5.57-5.54 (1H, m), 3.32-3.22 (4H, m), 3.17-2.99 (3H, m), 2.51-2.36 (2H, m), 2.28-2.05 (6H, m), 2.02 (3H, s), 1.90-1.77 (2H, m), 1.19 (3H, t, J = 7.0 Hz).
27		¹ H-NMR (400 MHz, CDCl ₃) δ: 7.60 (1H, dd, J = 8.8, 5.2 Hz), 7.11 (1H, dd, J = 8.5, 2.4 Hz), 6.96 (1H, ddd, J = 8.8, 8.8, 2.0 Hz), 5.96-5.93 (1H, m), 5.57-5.54 (1H, m), 3.59-3.48 (4H, m), 3.31-3.23 (4H, m), 2.68-2.57 (4H, m), 2.50-2.39 (2H, m), 2.02 (3H, s), 1.89-1.77 (2H, m), 1.19 (3H, t, J = 7.3 Hz).
28		¹ H-NMR (400 MHz, CDCl ₃) δ: 7.62 (1H, dd, J = 8.5, 4.9 Hz), 7.17 (1H, dd, J = 8.2, 1.5 Hz), 6.99 (1H, ddd, J = 8.8, 8.8, 2.0 Hz), 5.92-5.88 (1H, m), 5.53-5.48 (1H, m), 3.27-3.16 (4H, m), 3.09-2.93 (3H, m), 2.43-2.32 (2H, m), 2.16-1.92 (9H, m), 1.86-1.69 (2H, m), 1.14 (3H, t, J = 7.3 Hz).

29		¹ H-NMR (400 MHz, CDCl ₃) δ: 7.99 (1H, s), 7.61 (1H, d, J = 7.9 Hz), 7.35 (1H, d, J = 8.5 Hz), 7.17 (1H, ddd, J = 7.6, 7.6, 1.2 Hz), 7.12-7.06 (1H, m), 6.97 (1H, d, J = 2.4 Hz), 6.54 (1H, dd, J = 16.5, 9.8 Hz), 6.21 (1H, d, J = 17.1 Hz), 5.89 (1H, d, J = 10.4 Hz), 3.46-3.34 (2H, m), 3.26 (2H, q, J = 7.1 Hz), 3.20-3.05 (2H, m), 2.93-2.81 (1H, m), 2.76-2.61 (2H, m), 2.43-2.22 (2H, m), 2.14-2.02 (2H, m), 1.99-1.79 (2H, m), 1.22 (3H, t, J = 7.0 Hz).
30		¹ H-NMR (400 MHz, CDCl ₃) δ: 7.57-7.46 (1H, m), 7.19 (1H, d, J = 9.1 Hz), 6.52 (1H, dd, J = 15.2, 9.1 Hz), 6.21 (1H, d, J = 16.5 Hz), 5.91 (1H, d, J = 8.5 Hz), 3.47-3.00 (8H, m), 2.79-2.60 (1H, m), 2.42-1.94 (9H, m), 1.22 (3H, t, J = 7.0 Hz).
31		¹ H-NMR (400 MHz, CDCl ₃) δ: 7.46 (1H, s), 7.42 (1H, d, J = 8.5 Hz), 7.34-7.29 (1H, m), 6.55 (1H, dd, J = 16.8, 10.1 Hz), 6.20 (1H, d, J = 16.5 Hz), 5.89 (1H, d, J = 9.8 Hz), 3.37-3.29 (2H, m), 3.25 (2H, q, J = 7.1 Hz), 3.12-3.00 (3H, m), 2.68-2.57 (2H, m), 2.44 (3H, s), 2.34-2.19 (2H, m), 2.14-2.01 (4H, m), 1.21 (3H, t, J = 7.3 Hz).
32		¹ H-NMR (400 MHz, CDCl ₃) δ: 7.62 (1H, d, J = 8.5 Hz), 7.56 (1H, d, J = 1.8 Hz), 7.26 (1H, dd, J = 8.2, 1.5 Hz), 6.53 (1H, dd, J = 16.8, 10.1 Hz), 6.20 (1H, d, J = 17.1 Hz), 5.89 (1H, d, J = 9.8 Hz), 3.31 (2H, t, J = 7.0 Hz), 3.25 (2H, q, J = 7.1 Hz), 3.12-2.98


		(3H, m), 2.66-2.56 (2H, m), 2.33-2.18 (2H, m), 2.12-1.98 (4H, m), 1.20 (3H, t, $J = 7.0$ Hz).
33		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.73-7.69 (1H, m), 7.57-7.49 (2H, m), 7.30-7.25 (1H, m), 6.55 (1H, dd, $J = 16.5, 9.8$ Hz), 6.20 (1H, d, $J = 16.5$ Hz), 5.89 (1H, d, $J = 9.8$ Hz), 3.32 (2H, t, $J = 7.0$ Hz), 3.25 (2H, q, $J = 7.1$ Hz), 3.16-3.00 (3H, m), 2.61 (2H, t, $J = 6.7$ Hz), 2.34-2.19 (2H, m), 2.14-2.01 (4H, m), 1.21 (3H, t, $J = 7.0$ Hz).
34		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.99 (1H, s), 7.50 (1H, dd, $J = 8.8, 5.2$ Hz), 7.02 (1H, dd, $J = 9.8, 1.8$ Hz), 6.93 (1H, d, $J = 1.8$ Hz), 6.85 (1H, ddd, $J = 9.1, 9.1, 2.4$ Hz), 6.53 (1H, dd, $J = 16.8, 10.1$ Hz), 6.20 (1H, d, $J = 16.5$ Hz), 5.90 (1H, d, $J = 10.4$ Hz), 3.45-3.32 (2H, m), 3.26 (2H, q, $J = 7.1$ Hz), 3.21-3.00 (2H, m), 2.91-2.55 (3H, m), 2.39-1.99 (4H, m), 1.98-1.75 (2H, m), 1.22 (3H, t, $J = 7.0$ Hz).
35		$^1\text{H-NMR}$ (400 MHz, CDCl_3) δ : 7.93 (1H, s), 7.37 (1H, dd, $J = 9.2, 4.3$ Hz), 7.32 (1H, dd, $J = 8.5, 2.4$ Hz), 7.12 (1H, ddd, $J = 8.9, 8.9, 2.4$ Hz), 6.54 (1H, dd, $J = 16.8, 10.1$ Hz), 6.20 (1H, d, $J = 16.5$ Hz), 5.90 (1H, d, $J = 10.4$ Hz), 4.43-4.31 (1H, m), 3.36-3.22 (4H, m), 3.17-3.04 (2H, m), 2.67-2.57 (2H, m), 2.38-2.22 (4H, m), 2.08-1.95 (2H, m), 1.21 (3H, t, $J = 7.0$ Hz).

36		¹ H-NMR (400 MHz, CDCl ₃) δ: 7.93 (1H, s), 7.40-7.35 (1H, m), 7.34-7.29 (1H, m), 7.16-7.09 (1H, m), 5.98 (1H, br s), 5.56 (1H, br s), 4.44-4.29 (1H, m), 3.42-3.26 (4H, m), 3.17-2.99 (2H, m), 2.72-2.55 (2H, m), 2.38-2.18 (4H, m), 2.08-1.93 (5H, m), 1.22 (3H, t, J = 7.3 Hz).
----	---	--

[0194]

Reference example 37

3-[4-(1,2-Benzisoxazol-3-yl)piperidin-1-yl]-N-ethylpropane-1-amine dihydrochloride




According to a similar method of Reference examples 18 and 21, the titled compound was prepared from 3-(piperidin-4-yl)benzo[d]isoxazole.

[0195]

10 Reference example 38

N-Ethyl-3-[4-(6-fluoro-1,2-benzisoxazol-3-yl)piperazin-1-yl]propane-1-amine dihydrochloride



15 According to a similar method to Reference examples 18 and 21, the titled compound was prepared from 6-fluoro-3-(piperazin-1-yl)benzo[d]isoxazole monohydrochloride.

[0196]

Test 1: Evaluation of binding activity for human 5-HT_{2A} receptor,

human 5-HT₇ receptor, and human D₂ receptor

Binding affinity of the present compound for human 5-HT_{2A} receptor, human 5-HT₇ receptor, and human D₂ receptor was measured by the following procedures.

5 CHO cell membrane fraction in which human 5-HT_{2A} receptor, human 5-HT₇ receptor, or human D₂ receptor was expressed was purchased from PerkinElmer, Inc. In a test for evaluating binding affinity, a test compound dissolved in dimethylsulfoxide (DMSO) and each receptor membrane sample diluted in buffer were mixed
10 with [3H]Ketanserin, [3H]SB-269970, or [3H]Spiperone (all purchased from PerkinElmer, Inc.) for 5-HT_{2A} receptor, 5-HT₇ receptor, or D₂ receptor, respectively. Each mixture was incubated at room temperature for 60 minutes. Non specific binding to receptors was obtained from a competitive binding test
15 in the presence of 10 µmol/L 8-OH-DPAT, 10 µmol/L Mianserin, or 10 µmol/L Spiperone, respectively. Radioactivity caused by binding to receptors was measured with a liquid scintillation counter (PerkinElmer, Inc.), and 50% inhibition concentration was calculated. Ki value was evaluated from dissociation constant
20 which was calculated from saturated binding tests, and a substrate concentration, and it was used as an index for binding affinity. The results are shown in the following table.

Example	5-HT _{2A} Ki (nmol/L)	5-HT ₇ Ki (nmol/L)	D ₂ Ki (nmol/L)	D ₂ Ki/ 5-HT _{2A} Ki	D ₂ Ki/ 5-HT ₇ Ki
1	2.0	2.5	226	113	89
2	2.1	9.3	>1000	>483	>108
3	4.7	5.2	>1000	>212	>192
4	0.4	0.7	267	738	390
5	0.6	2.4	116	191	48
6	0.8	5.7	278	364	49
7	0.2	1.0	79	519	80
8	1.4	6.2	294	216	47
9	0.4	1.6	<100	<231	<64
10	1.7	2.7	<100	<59	<38
11	1.5	5.2	<100	<68	<19
12	1.7	6.9	711	425	103
13	4.6	20.3	>1000	>217	>49
14	5.5	14.6	>1000	>183	>68
15	4.8	26.4	>1000	>208	>38
16	2.5	5.9	389	157	66
17	0.8	4.9	155	187	31
18	2.5	3.7	198	79	53
19	1.3	2.3	105	81	46
20	0.5	1.5	46	96	31
21	3.4	3.4	>1000	>296	>295
22	2.1	5.8	528	252	91
23	1.8	7.9	>1000	>552	>127
24	1.1	1.1	<100	<95	<91
25	6.4	7.7	1389	217	182
26	58.0	22.1	>10000	>172	>453
27	10.3	11.1	2164	210	195
28	2.2	10.0	<100	<45	<10
29	1.1	2.6	146	135	55

30	0.8	6.6	153	184	23
31	1.0	1.2	89	91	77
32	97.0	109.7	>10000	>103	>91
33	0.3	1.6	55	200	34
34	0.6	4.6	247	386	53
35	1.5	3.8	<100	<67	<27
36	0.9	1.0	80	93	84

37	0.9	1.1	<100	<110	<94
38	8.3	9.6	352	42	36
39	0.8	0.7	<100	<126	<137
40	1.1	1.2	214	202	185
41	4.6	1.8	<100	<22	<57
42	2.1	2.5	74	35	30
43	1.1	4.3	<100	<87	<23
44	0.5	3.2	<100	<194	<31
45	15.0	22.6	>300	>20	>13
46	0.9	4.2	75	80	18
47	2.8	8.9	457	166	51
48	0.6	2.7	80	134	30
49	9.8	22.6	>1000	>102	>44
50	4.0	13.5	>1000	>252	>74
51	2.7	1.9	309	113	164
52	298.2	332.6	>10000	>34	>30
53	1.1	1.4	<100	<92	<71
54	2.0	2.6	116	58	45
55	1.1	0.1	78	69	890
56	1.2	0.4	<100	<86	<254
57	1.6	8.7	210	128	24
58	53.2	19.4	>1000	>19	>52
59	44.7	13.6	>1000	>22	>74
60	19.9	24.3	>1000	>50	>41
61	10.9	12.7	514	47	41
62	0.5	2.8	104	192	37
63	3.1	6.9	532	172	77
64	1.5	2.4	348	228	147
65	4.0	7.6	195	48	26
66	0.8	5.4	212	273	40
67	6.1	0.9	<100	<16	<110

68	9.9	48.5	910	92	19
69	0.6	1.5	<100	<165	<65
70	2.0	3.7	52	26	14
71	2.2	6.2	223	99	36
72	1.6	10.6	<100	<61	<9
73	5.7	14.4	205	36	14
74	2.4	6.8	138	57	20
75	1.0	3.6	124	127	34
76	7.2	14.4	193	27	13

[0197]

Test 2: Evaluation of antagonist activity for human 5-HT_{2A} receptor and human 5-HT₇ receptor

Aequorin, Gα 16 protein, and each receptor were transiently expressed in CHO-K1 cells (Chinese hamster ovary). The cells were cultivated in a CO₂ incubator at 37°C overnight, seeded into a 384-well plate, and stood at room temperature for 2 hours. Each compound dissolved in DMSO was added thereto, and changes in luminescence were measured by FDSS/μCELL drug discovery screening support system (Hamamatsu Photonics K.K.). As for antagonist activity, inhibitory activity of each compound was calculated by setting the luminescence level of wells to which 10 μmol/L endogenous ligand was added as 100%. The results are shown in the following table.

Example	5-HT _{2A} antagonist activity	5-HT ₇ antagonist activity
	IC ₅₀ (nmol/L)	IC ₅₀ (nmol/L)
1	8	37
2	12	94
3	62	95
4	44	31
5	9	66
6	48	66
7	8	11
8	26	8
9	15	8
10	40	9
11	9	67
12	78	82
13	76	81
14	79	82
15	60	86
16	59	90
17	49	66
18	58	72
19	76	61
20	19	48
21	73	68
22	738	40
23	277	68
24	7	41
25	90	26
26	843	71
27	64	73
28	95	80
29	39	99

30	45	39
31	55	64
32	890	73
33	8	59
34	6	78
35	33	81
36	13	82

37	21	66
38	756	59
39	8	9
40	9	2
41	7	7
42	8	9
43	9	82
44	10	77
45	640	49
46	9	83
47	94	43
48	10	18
49	633	16
50	49	81
51	57	62
52	946	177
53	72	44
54	31	1
55	7	3
56	7	6
57	56	100
58	3643	103
59	5888	62
60	819	72
61	94	15
62	53	28
63	65	46
64	44	101
65	103	81
66	37	70
67	94	72

68	949	91
69	745	85
70	67	9
71	25	78
72	42	8
73	68	81
74	68	77
75	306	80
76	551	65

[0198]

Test 3-1: Metabolic stability test in human hepatic microsome

The metabolic stability of the present compound in human hepatic microsome was evaluated as mentioned below. Human hepatic microsome was obtained from Xenontech. Human hepatic microsome, NADPH, and each test compound were mixed in 25 mmol/L phosphate buffer solution (pH 7.4) to reach the following concentrations as shown below, and the mixture was incubated at 37°C for 30 minutes.

- Human hepatic microsome: 0.1 mg/mL
- NADPH: 3.2 mmol/L
- Test compound: 0.1 µmol/L

The residual ratio of test compound in each sample after 30 minutes was measured by LC-MS, and the metabolic stability in human hepatic microsome was calculated from the following formula.

Metabolic stability in human hepatic microsome (mL/min/mg protein) = $-\text{LN}(\text{residual ratio})/30/0.1$

The results are shown in the following table.

Example	Metabolic stability in human hepatic microsome (mL/min/mg protein)		Example	Metabolic stability in human hepatic microsome (mL/min/mg protein)
1	<0.05		40	<0.05
2	<0.05		41	<0.05
3	<0.05		42	<0.05
4	<0.05		43	<0.05
5	0.186		44	<0.05
6	0.053		45	<0.05
7	0.08		46	<0.05
8	0.108		47	<0.05
9	<0.05		48	<0.05
10	<0.05		49	<0.05
11	<0.05		50	<0.05
12	0.089		51	<0.05
13	0.054		53	<0.05
16	<0.05		54	<0.05
17	<0.05		55	0.059
18	<0.05		56	<0.05
19	<0.05		57	0.06
20	<0.05		58	<0.05
21	<0.05		59	0.079
22	<0.05		61	<0.05
23	<0.05		62	<0.05
24	<0.05		63	<0.05
25	<0.05		64	0.097
26	<0.05		65	<0.05
27	<0.05		66	<0.05
29	<0.05		67	<0.05
30	<0.05		68	<0.05

31	<0.05		69	<0.05
33	0.268		70	<0.05
34	<0.05		71	<0.05
35	<0.05		72	0.057
36	<0.05		74	<0.05
37	<0.05		75	<0.05
38	<0.05		76	<0.05
39	<0.05			

[0199]

Test 3-2: Metabolic stability test in human hepatic microsome

For more accurate evaluation of the metabolic stability in human hepatic microsome, the metabolic stability of the present compound in human hepatic microsome was evaluated at an appropriate concentration of human hepatic microsome as mentioned below. Human hepatic microsome was obtained from Xenontech. Human hepatic microsome, NADPH, and each test compound were mixed in 25 mmol/L phosphate buffer solution (pH 7.4) to reach the following concentrations as shown below, and the mixture was incubated at 37°C for 60 minutes.

- Human hepatic microsome: 0.5 or 1.0 mg/mL
- NADPH: 3.2 mmol/L
- Test compound: 0.1 µmol/L

The residual ratio of a test compound in each sample after 30 or 60 minutes was measured by LC-MS, and the metabolic stability in human hepatic microsome was calculated from the following formula.

Metabolic stability in human hepatic microsome (mL/min/mg protein) = $-\text{LN} (\text{residual ratio}) / \text{reaction time} / \text{concentration of}$

human hepatic microsome

The results are shown in the following table.

Example	Metabolic stability in human hepatic microsome (mL/min/mg protein)
16	0.018
17	0.0025
51	0.016

[0200]

Test 4: Predictive test of human half life

5 The disappearance half life of the present compound in human was predicted as mentioned below.

 The present compound was intravenously administered to a rat as an aqueous solution of 0.01 mol/L hydrochloric acid. Blood was collected on 5 minutes, 15 minutes, 30 minutes, 1 hour, 2
10 hours, 4 hours, 6 hours, and 24 hours after the administration. Plasma was obtained from the collected blood, the drug concentration in plasma was measured by LC-MS, and the distribution volume of rat was calculated from the transition of the concentration.

15 The unbound fraction rate of the present compound in human or rat serum was measured by equilibrium dialysis method.

 The half-life in human was calculated according to the following formula using the results of the distribution volume of rat, the unbound fraction rate in human or rat serum, and the
20 metabolic stability in human hepatic microsome obtained in Test 3-2.

- Distribution volume of human = Distribution volume of rat x

Unbound fraction rate in human serum / Unbound fraction rate in rat serum

- Human hepatic clearance = (Hepatic blood flow of human x Unbound fraction rate in human serum x 56.7 x Metabolic stability in human hepatic microsome) / (Hepatic blood flow of human + Unbound fraction rate in human serum x 56.7 x Metabolic stability in human hepatic microsome)

- Half-life in human = 0.693 x Distribution volume of human / Human hepatic clearance

10 The results are shown in the following table.

Example	Predicted half-life (h)
16	3
17	27
51	6

[0201]

Test 5: Evaluation of inhibitory activity for hERG channel

Inhibitory activity of the present compound for hERG channel was measured by whole cell patch clamp method with an auto patch clamp system using CHO cells in which hERG channel involved in human rapidly activating delayed rectifier potassium current (I_{Kr}) was forcibly expressed.

(Preparation of cell suspension)

hERG-CHO cells purchased from ChanTest Cop. were cultured in a CO₂ incubator at 37°C, and dissociated from a flask with trypsin shortly before the measurement of hERG current to prepare a cell suspension.

(Preparation of solution)

Extracellular and intracellular fluids used in the measurement were prepared as follows.

Extracellular fluid: 2 mmol/L CaCl_2 , 1 mmol/L MgCl_2 , 10 mmol/L HEPES, 4 mmol/L KCl, 145 mmol/L NaCl, 10 mmol/L Glucose

5 Intracellular fluid: 10 mmol/L HEPES, 10 mmol/L EGTA, 20 mmol/L KCl, 130 mmol/L KF

Test compound solution: A test compound was dissolved in DMSO to reach the concentration of 2 mmol/L or 20 mmol/L in order to prepare a test compound solution. The test compound solution was
10 further diluted 200 times with the extracellular fluid, and serial-diluted with the extracellular fluid to prepare a test compound solution in each concentration which is necessary to calculate IC_{50} value of hERG inhibition.

(Measurement of current value and data analysis)

15 The cell suspension, the extracellular fluid, the intracellular fluid, and the measurement plate were set in an auto patch clamp system, and hERG current was measured by whole cell patch clamp method. A voltage-protocol was as follows: the holding potential was adjusted to -80 mV, the depolarizing pulse
20 was provided at -50 mV to +20 mV for 5 seconds, the repolarizing pulse was provided at -50 mV for 5 seconds, then the potential was returned to the holding potential. Each pulse interval was 15 seconds. The data analysis was carried out with Qube Assay Software (Sophion Sophion). The test was carried out by applying
25 incrementally 4 concentrations of each test compound, and the average of the maximum peak tail currents which were obtained from the last 3 stimulations in each concentration was determined

as the evaluated data. Based on the current inhibition rate for a pre-applied current at each concentration of each test compound, IC_{50} value was calculated by Hill equation with the software.

The results are shown in the following table.

Example	hERG inhibitory IC ₅₀ (μ mol/L)	hERG inhibitory IC ₅₀ (nmol/L) / 5-HT _{2A} Ki (nmol/L)	hERG inhibitory IC ₅₀ (nmol/L) / 5-HT ₇ Ki (nmol/L)
1	4.8	2401	1896
2	7.4	3575	796
3	>10	>2122	>1922
4	7.1	19617	10367
5	1.1	1810	455
6	1.5	1958	265
7	3.5	22975	3526
8	4.2	3088	677
9	5.4	12480	3453
10	3.6	2135	1351
11	10.5	7172	2034
12	2.2	1313	319
13	2.9	629	143
14	9.6	1754	656
16	4.8	1939	814
17	2.8	3381	567
18	4.1	1637	1097
19	>10	>7705	>4405
20	4.2	8787	2849
21	15.3	4529	4520
22	>10	>4475	>1726
23	>10	>5520	>1266
24	5.5	5203	4988
25	>10	>1563	>1307
26	>10	>172	>453
27	>10	>968	>899
29	0.9	830	341
30	1.2	1441	182

31	3.1	3169	2694
33	2.2	8013	1374
34	7.3	11440	1578
35	4.3	2888	1143
36	1.3	1504	1361
37	>10	>10968	>9375
38	>10	>1200	>1036
39	4.7	5934	6462
40	9.9	9377	8589

41	6.7	1461	3821
42	6.4	3022	2583
43	7.0	6091	1613
44	6.4	12428	1977
45	>10	>665	>442
46	5.7	6053	1373
47	5.4	1954	607
48	2.9	4870	1090
49	>10	>1017	>442
50	6.2	1563	458
51	6.9	2536	3665
53	7.2	6616	5125
54	7.4	3685	2855
55	7.4	6575	84634
56	2.1	1805	5341
57	6.2	3785	717
58	>10	>188	>516
59	>10	>224	>738
61	>10	>916	>790
62	5.4	9970	1944
63	14.5	4677	2101
64	>10	>6551	>4214
65	>10	>2477	>1314
66	>10	>12871	>1865
67	>10	>1632	>11030
69	>10	>16548	>6545
70	>10	>4913	>2739
71	>10	>4452	>1624
72	9.9	6029	933
74	6.7	2795	991
75	3.8	3881	1051

76	>10	>1394	>697
----	-----	-------	------

[0202]

Test 6: Fear conditioning Test

Male SD rats were used. For preparation of an administration solution, a test compound was dissolved in 0.01 mol/L hydrochloric acid, and the aqueous solution was used.

Fear conditioning test was conducted as mentioned below using Image J FZ2 for Contextual and cued fear conditioning test from O'HARA & CO., LTD.

An animal was placed into a chamber quipped in the Image J FZ2 for Contextual and cued fear conditioning test, and electric stimulations were applied 7 times at 0.5 mA for 10 seconds. Then, a rat was gently removed from a chamber, and on the next day, the administration solution (a solvent or test compound solution) was subcutaneously administered, the animal was placed in a chamber on 30 minutes later. The percentage of freezing reaction time from when an animal was placed in a chamber to 300 seconds later was used as a test result.

Analysis of the test results were carried out as follows.

Parametric Dunnett's multiple comparison was conducted on both of the test compound administration group and the solvent administration group (significance level: 5% on both sides). When the test compound administration group showed a significant suppression of the freezing reaction time, compared with the solvent administration group, it was regarded as showing an anxiolytic effect.

The results of this test using the compound of Example 17

are shown in Figure 1.

[0203]

Test 7: Measurement of the amount of released glutamic acid in rat brain

5 A cannula was indwelled in a prefrontal cortex position in a male Wistar rat, and the rat was subjected to the test after a recovery period of at least 1 week.

 For calibration of a biosensor, 5 mM solution of glutamic acid and 100 mM solution of ascorbic acid were used. Released
10 glutamic acid was measured with the biosensor which had calibrated. The measurement was started 12 hours or more after the biosensor was inserted into a guide cannula. After administration of a test compound, the measurement was performed for 2 hours or more.

 The average amount of released glutamic acid on 30 minutes
15 before administration was used as a standard, and the area under the curve was calculated on 2 hours after administration. The data of the compound administration group was compared with that of the solvent administration group.

 The average values of each group were compared by parametric
20 Dunnett's multiple comparison test, and when the test compound administration group showed a significant high value compared with the solvent administration group, it was regarded as showing a promoting effect of releasing glutamic acid (significance level: 5% on both sides).

25 The results of this test using the compound of Example 17 are shown in Figure 2.

[0204]

Test 8: Evaluation of binding activity for side-effect-related receptor

Binding affinity of the present compound for side-effect-related receptor, such as adrenergic $\alpha 1A$ receptor, can be measured
5 by the following method.

Evaluation test for binding is carried out as follows with the CHO cell membrane fraction in which human target receptor is expressed. A test compound dissolved in dimethylsulfoxide (DMSO), each receptor membrane sample diluted with buffer, and [3H]-
10 labelled ligand which has strong binding affinity to each target receptor are mixed. Each mixture is incubated at room temperature, added quickly on a glassfiber filter plate (Multiscreen FB, Millipore, Inc.), and vacuum-filtered. Radioactivity remaining on the filter is measured with a liquid scintillation counter
15 (PerkinElmer, Inc.). Binding inhibition rate is calculated from the following formula. A control compound which has strong binding affinity to a target receptor is used to calculate the non-specific binding amount to the receptor membrane sample, instead of a test compound.

20 [0205]

Binding inhibition rate to target receptor (%) = $100 - 100 \times \frac{\{(\text{Binding amount of } [^3\text{H}]\text{-labelled ligand in the presence of test compound}) - (\text{Binding amount of } [^3\text{H}]\text{-labelled ligand in the presence of } 10 \mu\text{mol/L control compound})\}}{\{(\text{Binding amount of } [^3\text{H}]\text{-labelled ligand in the absence of test compound}) - (\text{Binding amount of } [^3\text{H}]\text{-labelled ligand in the presence of } 10 \mu\text{mol/L control compound})\}}$
25

[0206]

Test 9: Evaluation of intracerebral transferability (test for intracerebral transferability in rats)

In this test, intracerebral transferability of the present compound was evaluated by the following method. The present compounds were subcutaneously administered as a solution in saline, or orally administered as a suspension in methyl cellulose to a 7-week-old SD or WKY rat. Plasma and brain were collected on 0.5 hour, 1 hour, or 2 hours after the administration in order to measure the drug concentrations in plasma and brain by LC-MS.

Binding rates of the present compound to plasma and brain protein were measured by equilibrium dialysis method.

K_{p,uu,brain} (unbound drug concentration ratio between brain/plasma) can be calculated by applying the compound concentrations in plasma and brain and the binding rates to plasma and brain protein obtained from the above test into the following formula.

$$K_{p,uu,brain} = \frac{(\text{Compound concentration in brain} \times (100 - \text{Binding rate to brain protein} (\%)) / 100)}{(\text{Compound concentration in plasma} \times (100 - \text{Binding rate to plasma protein} (\%)) / 100)}$$

[0207]

The results of Test 9 are shown in the following table.

Example	K _{p,uu,brain}	Example	K _{p,uu,brain}
2	0.35	21	0.54
16	1.32	23	1.47
17	0.58	41	0.63
18	0.56	51	1.26
19	0.33	63	0.35

[0208]

Test 10: Evaluation of hepatotoxic risk (Dansyl glutathione
(dGSH) trapping assay)

The present compound was metabolized in hepatic microsome,
5 and from the resulted metabolite, reactive metabolite which
reacts with dansyl glutathione (dGSH) was detected and quantified
as the following method. Measurement was carried out with a
screening robot (Tecan) for metabolic reaction, and with a
fluorescence detection UPLC system (Waters) for metabolite-dGSH
10 binding concentration.

[0209]

(Preparation of solution)

The present compound was dissolved in DMSO to prepare 10
mmol/L test compound solution. 7.6 mL of potassium phosphate
15 buffer (500 mmol/L, pH 7.4), 1.9 mL of human hepatic microsome
(Xenotech, 20 mg protein/mL), and 1.27 mL of pure water were
mixed to prepare a microsome solution. To 3.78 mL of the
microsome solution was added 0.67 mL of pure water to prepare a
microsome (dGSH(-)) solution. To 6.48 mL of the microsome
20 solution was added 1.14 mL of the dGSH solution (20 mmol/L) to
prepare a microsome (dGSH(+)) solution. 80.9 mg of NADPH was
dissolved in 30 mL of pure water to prepare a cofactor solution.
33 mg of tris(2-carboxyethyl)phosphine (TECP) was dissolved in
115 mL of methanol to prepare a reaction stop solution.

25 [0210]

(Reaction)

12 μ L of the test compound solution was mixed with 388 μ L

of pure water, and the mixture was dispensed in 50 µL each into 6 wells of a 96-well plate. The 6 wells were divided into 3 groups of 2 wells, and each was named as "reaction group", "unreacted group", and "dGSH-free group". To the "reaction group" and "unreacted group" was added the microsome (dGSH(+)) solution, and to the "dGSH-free group" was added the microsome (dGSH(-)) solution in 50 µL each. To the "reaction group" and "dGSH-free group" was added to the cofactor solution, and to the "unreacted group" was added pure water in 50 µL each. After incubated at 37°C for 60 minutes, the reaction stop solution was added in 450 µL each to stop the reaction. To the "reaction group" and "dGSH-free group" was added pure water, and to the "unreacted group" was added the cofactor solution in 50 µL each. The plate was cooled at -20°C for 1 hour, and the solutions were centrifuged (4000 rpm, 10 minutes). Supernatants were collected into another plate and subjected to analysis.

[0211]

(Analysis)

Metabolite-dGSH binding concentration was measured by the following conditions, using a fluorescence detection UPLC system. Column: Waters ACQUITY UPLC BEHC18 1.7 µm 2.1 x 10 mm
Eluent: A, 0.2% aqueous formic acid; B, 0.2% formic acid/acetonitrile
Gradient: B, 20% (0 min) -> 70% (9.33 min) -> 90% (10.63 min) -> 20% (11 min) -> 20% (14 min)

Fluorescence intensity was corrected with the composition of organic solvent at the time of elution because fluorescence

intensity changes depending on the composition of organic solvent.

[0212]

The results of Test 10 are shown in the following table.

Example	Metabolite-dGSH binding concentration (μM)		Example	Metabolite-dGSH binding concentration (μM)
1	N.D.		39	N.D.
2	0.127		40	0.125
3	N.D.		41	N.D.
4	0.165		42	N.D.
5	N.D.		43	0.177
6	N.D.		44	0.096
7	N.D.		45	N.D.
8	0.223		46	0.156
9	N.D.		47	N.D.
10	N.D.		48	N.D.
11	N.D.		49	0.45
12	N.D.		50	N.D.
13	0.214		51	N.D.
14	0.089		53	0.063
16	N.D.		54	N.D.
17	N.D.		55	0.057
18	N.D.		56	0.514
19	0.175		57	N.D.
20	1.661		58	N.D.
21	0.133		59	N.D.
22	0.509		61	N.D.
23	0.173		62	0.857
24	7.645		63	0.268
25	N.D.		64	0.432
26	N.D.		65	0.112
27	N.D.		66	0.222
29	0.091		67	0.113
30	0.101		69	N.D.
31	N.D.		70	0.257

33	N.D.		71	0.132
34	0.268		72	0.147
35	0.153		74	N.D.
36	0.161		75	N.D.
37	0.108		76	N.D.
38	N.D.			
(N.D. means below detection limit.)				


INDUSTRIAL APPLICABILITY

[0213]

The present compound shows antagonist activity for the serotonin 5-HT_{2A} receptor and the serotonin 5-HT₇ receptor, and therefore, the present compound is useful as a medicament for treating neuropsychiatric diseases.

CLAIMS

1. A compound of Formula (1):



5 or a pharmaceutically acceptable salt thereof,
wherein

Z is a nitrogen atom or $-CR^A-$;

Y is carbonyl or sulfonyl;

m is 1, 2, 3, or 4;

10 n is 0, 1, 2, or 3;

provided that when Y is sulfonyl, n is not 0;

R^A is hydrogen, hydroxy, C_{1-6} alkyl, or C_{1-6} alkoxy;

15 R^{1a} , R^{1b} , R^{1c} , and R^{1d} are each independently hydrogen, hydroxy, halogen, or C_{1-6} alkyl optionally-substituted with the same or different 1 to 3 halogen;

20 R^{2a} , R^{2b} , R^{2c} , and R^{2d} are each independently, and each of R^{2c} s or each of R^{2d} s is independently when R^{2c} or R^{2d} exists plurally, hydrogen, halogen, or C_{1-6} alkyl (wherein the alkyl may be optionally substituted with the same or different 1 to 5 substituents selected from the group consisting of halogen, hydroxy, C_{3-8} cycloalkyl, and C_{1-6} alkoxy), provided that when R^{2a} and R^{2b} , or R^{2c} and R^{2d} which

attach to the same carbon atom are each independently the said C₁₋₆ alkyl, they may be combined together with the carbon atom to which they attach to form a 3- to 6-membered saturated carbocycle;

5 R³ is C₁₋₆ alkyl, wherein the alkyl may be optionally substituted with the same or different 1 to 5 substituents selected from the group consisting of halogen, hydroxy, cyano, C₃₋₈ cycloalkyl, and C₁₋₆ alkoxy;


10 R^{4a}, R^{4b}, R^{4c}, and R^{4d} are each independently, and each of R^{4c}s or each of R^{4d}s is independently when R^{4c} or R^{4d} exists plurally, hydrogen, halogen, C₁₋₆ alkyl (wherein the alkyl may be optionally substituted with the same or different 1 to 5 substituents selected from the group consisting of halogen, hydroxy, cyano, C₃₋₈ cycloalkyl, C₁₋₆ alkoxy, C₁₋₆ alkyl ester, and amino optionally-substituted with the same or different 1 or 2 C₁₋₆ alkyl (wherein the alkyl may be optionally substituted with oxo)), C₂₋₆ alkynyl, or amino (wherein the amino may be optionally substituted with the same or different 1 or 2 C₁₋₆ alkyl); provided that

20 when any two of R^{4a}, R^{4b}, R^{4c}, and R^{4d} are each independently the said C₁₋₆ alkyl, they may be combined together with the carbon atom or the carbon atoms to which they attach to form a 3- to 6-membered saturated carbocycle or a 4- to 6-membered saturated heterocycle wherein when the said C₁₋₆

25 alkyl groups have a substituent, the substituent may be included as a ring member of the carbocycle or heterocycle;

provided that when Y is sulfonyl and n is 1, any one of R^{4c} or R^{4d} is hydrogen; and when Y is sulfonyl and n is 2 or 3, any one of R^{4c} or R^{4d} which binds to the carbon atom adjacent to Y is hydrogen; and

5 Ring Q is a group of the following Formula (2a), (2b), (2c), (2d), (2e), or (2f):



wherein

10 R^{5a}, R^{5b}, R^{5c}, and R^{5d} are each independently hydrogen, halogen, cyano, C₁₋₆ alkyl, C₁₋₆ alkoxy (wherein the alkyl and the alkoxy may be optionally each independently substituted with the same or different 1 to 3 halogen), or amino optionally-substituted with the same or
 15 different 1 or 2 C₁₋₆ alkyl; and

R⁶ is hydrogen, C₁₋₆ alkyl, or C₃₋₈ cycloalkyl, wherein the alkyl and the cycloalkyl may be optionally each

independently substituted with the same or different 1 to 3 halogen.

2. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein R^{1a} , R^{1b} , R^{1c} , and R^{1d} are hydrogen.

3. The compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein R^{2a} , R^{2b} , R^{2c} , and R^{2d} are hydrogen.

4. The compound of any one of claims 1 to 3, or a pharmaceutically acceptable salt thereof, wherein m is 1.

5. The compound of any one of claims 1 to 4, or a pharmaceutically acceptable salt thereof, wherein R^3 is C_{1-6} alkyl.

6. The compound of any one of claims 1 to 5, or a pharmaceutically acceptable salt thereof, wherein Z is $-CR^A-$.

7. The compound of any one of claims 1 to 6, or a pharmaceutically acceptable salt thereof, wherein R^A is hydrogen.

8. The compound of any one of claims 1 to 7, or a pharmaceutically acceptable salt thereof, wherein R^{5a} , R^{5b} , R^{5c} , and R^{5d} are each independently hydrogen, halogen, or C_{1-6} alkyl.

5

9. The compound of any one of claims 1 to 8, or a pharmaceutically acceptable salt thereof, wherein R^{5a} , R^{5c} , and R^{5d} are hydrogen.

10 10. The compound of any one of claims 1 to 9, or a pharmaceutically acceptable salt thereof, wherein R^{5b} is hydrogen, halogen, or C_{1-6} alkyl.

11 11. The compound of any one of claims 1 to 10, or a
15 pharmaceutically acceptable salt thereof, wherein R^6 is hydrogen.


12. The compound of any one of claims 1 to 11, or a pharmaceutically acceptable salt thereof, wherein R^{4a} , R^{4b} ,
20 R^{4c} , and R^{4d} are each independently, and each of R^{4c} s or each of R^{4d} s is independently when R^{4c} or R^{4d} exists plurally, hydrogen, halogen, or C_{1-6} alkyl, wherein the alkyl may be optionally substituted with the same or different 1 to 5 substituents selected from the group consisting of halogen,
25 hydroxy, and C_{1-6} alkoxy.

13. The compound of any one of claims 1 to 12, or a pharmaceutically acceptable salt thereof, wherein n is 0 or 1.

5 14. The compound of any one of claims 1 to 13, or a pharmaceutically acceptable salt thereof, wherein Y is carbonyl.

15 15. The compound of any one of claims 1 to 13, or a pharmaceutically acceptable salt thereof, wherein Y is sulfonyl.

16. The compound of claim 1, or a pharmaceutically acceptable salt thereof, wherein the compound is selected
15 from the group consisting of:



17. A drug comprising a compound of any one of claims 1 to 16, or a pharmaceutically acceptable salt thereof, as an active ingredient.

18. A medicament for treating a mental disease or central nervous system disease, comprising a compound of any one of claims 1 to 16, or a pharmaceutically acceptable salt thereof, as an active ingredient.

19. The medicament of claim 18, wherein the mental disease or central nervous system disease is organic, including symptomatic, mental disorders.

20. The medicament of claim 19, wherein the organic,
including symptomatic, mental disorders is Alzheimer's
disease, vascular dementia, dementia with Lewy bodies,
5 dementia in Parkinson's disease, mental disorders due to
other diseases such as brain damage, or other mental
disorders due to brain dysfunction and to physical disease.

21. The medicament of claim 18, wherein the mental disease
10 or central nervous system disease is schizophrenia, a
bipolar disorder, a sleep disorder, an autism spectrum
disorder, major depression, treatment-resistant depression,
or a psychopathic symptom or dementia associated with
Alzheimer's disease or Parkinson's disease.

15

22. Use of a compound of any one of claims 1 to 16, or a
pharmaceutically acceptable salt thereof, in the
manufacture of a medicament for treating a mental disease
or central nervous system disease.

20

23. A compound of any one of claims 1 to 16, or a
pharmaceutically acceptable salt thereof, for use in the
treatment of a mental disease or central nervous system
disease.

25

24. A medicament for treating a mental disease or central nervous system disease, comprising a compound of any one of claims 1 to 16, or a pharmaceutically acceptable salt thereof, in combination with at least one drug selected from the group consisting of a drug for treating a developmental disorder such as an autism spectrum disorder and an attention deficit hyperactivity disorder, an antipsychotic drug and a schizophrenic drug, a drug for treating a bipolar disorder, an antidepressant drug, an antianxiety drug, a drug for treating an obsessive-compulsive disorder, a drug for treating a stress disorder such as post-traumatic stress disorder, a drug for treating a mood disorder, a drug for treating an eating disorder, a drug for treating a sleep disorder such as insomnia, narcolepsy, a sleep apnea syndrome, and a circadian rhythm disorder, a drug for treating a sexual dysfunction, a drug for treating drug dependence, a drug for treating dementia such as Alzheimer's disease, a drug for treating a behavioral and psychological symptom associated with dementia, a cerebral metabolism and circulation improving drug, a drug for treating a movement disorder such as Parkinson's disease, an analgesic drug, an antiepileptic drug, an anticonvulsant, a migraine drug, an anesthetic, and a central stimulant.

25. A medicament for treatment of a mental disease or central nervous system disease, comprising a compound of any one of claims 1 to 16, or a pharmaceutically acceptable salt thereof, as an active ingredient, wherein the medicament is used in the treatment concomitantly with at least one drug selected from the group consisting of a drug for treating a developmental disorder such as an autism spectrum disorder and an attention deficit hyperactivity disorder, an antipsychotic drug and a schizophrenic drug, a drug for treating a bipolar disorder, an antidepressant drug, an antianxiety drug, a drug for treating an obsessive-compulsive disorder, a drug for treating a stress disorder such as a post-traumatic stress disorder, a drug for treating a mood disorder, a drug for treating an eating disorder, a drug for treating a sleep disorder such as insomnia, narcolepsy, a sleep apnea syndrome, and a circadian rhythm disorder, a drug for treating a sexual dysfunction, a drug for treating drug dependence, a drug for treating dementia such as Alzheimer's disease, a drug for treating a behavioral and psychological symptom associated with dementia, a cerebral metabolism and circulation improving drug, a drug for treating a movement disorder such as Parkinson's disease, an analgesic drug, an antiepileptic drug, an anticonvulsant, a migraine drug, an anesthetic, and a central stimulant.