METHODS OF DETERMINING PATIENT RESPONSE BY MEASUREMENT OF HER-3

Cross-Reference To Related Applications

This application claims the benefit of and priority under 35 USC §119(e) to U.S. Provisional Application No. 61/176,630, filed May 8, 2009, U.S. Provisional Application No. 61/187,962, filed June 17, 2009, and U.S. Provisional Application 61/145,029, filed January 15, 2009, which are incorporated by reference in their entireties.

Background of the Invention

A biomarker generally has a characteristic that can be objectively measured and evaluated as an indicator of normal biological processes, pathogenic processes or pharmacological responses to a therapeutic intervention. *See* Atkinson *et al.*, 2001, *Clin. Pharmacol. Ther.* 69:89-95. Biomarkers vary widely in nature, ease of measurement and correlation with physiological states of interest. *See, e.g.*, Frank *et al.*, 2003, *Nature Reviews Drug Discovery* 2:566-580. It is widely believed that the development of new validated biomarkers will lead both to significant reductions in healthcare and drug development costs and to significant improvements in treatment for a wide variety of diseases and conditions. Thus, a great deal of effort has been directed to using new technologies to find new classes of biomarkers. *See, e.g.*, Petricoin *et al.*, 2002, *Nature Reviews Drug Discovery*, 1:683-695; and Sidransky, 2002, *Nature Reviews Cancer* 2:210-219; Ludwig and Weinstein, 2005, *Nature Reviews Cancer* 5:845-856; Lee *et al.*, 2007, *Adv. Cancer. Res.*, 96:269-298; Dhani and Siu, 2008, *Cancer Metastasis Rev.* 27:339-349; Carden *et al.*, 2009, *Clin. Pharmacol. Ther.* 85:131- 133.

The interactions of cell surface membrane components play crucial roles in transmitting extracellular signals to a cell in normal physiology and in disease conditions. In particular, many types of cell surface receptors undergo dimerization, oligomerization or clustering in connection with the transduction of an extracellular event or signal into a cellular response, such as, *e.g.*, proliferation, increased or decreased gene expression or the like. *See, e.g.*, George *et al.*, 2002, *Nature Reviews Drug Discovery* 1:808-820; Mellado *et al*, 2001, *Ann. Rev. Immunol.* 19:397-421; Schlessinger, 2000, *Cell* 103:211-225; and Yarden, 2001, *Eur. J. Cancer* 37:S3-S8. The role of such events in diseases, such as cancer, has been the object of intense research and has led to the development of several new drugs and drug candidates. *See, e.g.*, Herbst and Shin, 2002, *Cancer* 94:1593-1611; Yarden and Sliwkowski, 2001, *Nature Reviews Molecular Cell Biology* 2:127-137; McCormick, 1999,

Trends in Cell Biology 9:53-56 (1999); and Blume-Jensen and Hunter, 2001, *Nature* 411:355-365.

Expression levels of individual cell surface receptors, such as Her-2 in breast cancer, have been used as biomarkers, especially to determine patient prognosis or whether a patient will or will not respond to certain treatments. In addition, oncogenic tyrosine kinases such as members of the epidermal growth factor receptor family have provided targets for drug development. However, the tyrosine kinase inhibitors targeted to EGFR and Her-2 have shown less clinical efficacy than anticipated from promising preclinical studies, which has led to interest in other EGFR-family members, such as Her-3, in part for prognostic value as biomarkers and in part because of interactions with other family members, leading to potential new drug targets. *See* Menendez and Lupu, 2007, *Breast Cancer Research* 9:111; Lee-Hoeflich *et al.,* 2008, *Cancer Res* 68:5878-5887; Fuchs *et al.*, 2006, *Anticancer Res.* 26:4397-4402; Sergina *et al.*, 2007, *Nature* 445:437-441; and Tovey *et al.,* 2006, *J. Pathol.* 210:358-362.

Her-3 is sometimes over-expressed in breast cancer, colorectal cancer, ovarian cancer, bladder cancer, prostate cancer, non-small cell lung cancer, melanoma, pharyngeal cancer, pancreatic cancer, esophageal cancer, glioma, biliary tract carcinoma, cholangiocarcinoma, gastric cancer, endometrial cancer, gall bladder cancer, squamous cell carcinoma or basal cell carcinoma. Conventional immunohistochemical (IHC) or fluorescence *in situ* hybridization (FISH) analyses have been used to detect Her-3 over-expression. Unfortunately, IHC and FISH have certain limitations as diagnostic tools in that they are not necessarily accurate and also prone to different interpretations by different laboratory personnel. There are currently no methods for accurately assessing the level of Her-3. The advent of a quantitative method for measuring Her-3 would facilitate the ability to accurately determine a cancer patient's prognosis and/or whether a patient is likely to respond to a certain treatment. See Mosesson *et al.*, 2004, *Semin. Cancer. Biol.* 14:262-270.

Summary of the Invention

In a first aspect, the invention is drawn to a method of measuring and/or quantifying the presence and/or amount of Her-3 and/or Her-3 in a complex in a sample, the method comprising providing a sample and determining the presence and/or quantity of Her-3 and/or Her-3 in a complex in the sample. In certain embodiments, the amount of Her-3 is above a first threshold, such that the sample is stratified as having a "high" amount of Her-3 (e.g., either total Her-3 and/or Her-3 homodimers and/or Her-3 heterodimers). In

some embodiments the first threshold value for Her-3 is a total Her-3 (H3T) of ≥ 0.158 and a low Her-3 value is below this threshold. Or, other ranges may be used depending upon the patient cohort and/or the significant event being monitored. Thus, each of the threshold values and/or threshold ranges described herein may vary by about 0.5 log units or less on a log scale and/or 25% or less on a linear scale (i.e., be $\leq 25\%$ larger and/or $\leq 25\%$ smaller than the specific ranges disclosed herein), or by about 20% or less, or by about 15% or less, or by about 10% or less, or by about 5% or less.

In a preferred embodiment, the sample is a biological sample. In a preferred embodiment, the sample is a tissue sample. In a preferred embodiment, the sample is a fresh tissue sample, a fixed sample, a frozen sample or a lysate. In a preferred embodiment, the sample is a tumor sample. In a preferred embodiment, the sample is a frozen tumor tissue sample. In a preferred embodiment, the sample comprises a tumor lysate from a fresh or frozen tumor sample. In a preferred embodiment, the sample is an FFPE or solubilized FFPE sample. In a preferred embodiment, the sample comprises a breast cancer sample. In a certain embodiments, the breast cancer is early stage (i.e., adjuvant) breast cancer or metastatic breast cancer.

In certain embodiments of each of the methods and/or aspects of the invention as disclosed herein, the method comprises detection of other biomarkers in the sample. For example, other biomarkers such as Her-2 and/or p95 may be measured. Or, other biomarkers such as can be at least one of FOXM1, PRAME, Bc12, STK15, CEGP1, Ki-67, GSTM1, CA9, PR, BBC3, NME1, SURV, GATA3, TFRC, YB-l, DPYD, GSTM3, RPS6KB1, Src, Chkl, ID1, EstR1, p27, CCNB1, XIAP, Chk2, CDC25B, IGF1R, AK055699, P13KC2A, TGFB3, BAGI1, CYP3A4, EpCAM, VEGFC, pS2, hENT1, WISP1, HNF3A, NFKBp65, BRCA2, EGFR, TK1, VDR, Contig51037, pENT1, EPHX1, IF1A, CDH1, HIFlα, IGFBP3; CTSB, Her3 or DIABLO. In certain embodiments, the other biomarker can be VEGF, CD31, KDR, p95, or Her-2.

In certain embodiments of each of the methods and/or aspects of the invention as disclosed herein, the level of Her-2 expression in the breast cancer is high. In certain embodiments, high Her-2 expression is a log10H2T $\geq$ about 1.14 - 1.25. In certain embodiments, the high Her-2 expression comprises expression that is very high and/or moderately high. In certain embodiments, the very high Her-2 expression is a log10H2T $\geq$ about 1.84 - 2.21. In certain embodiments of each of the methods disclosed herein, the moderately high expression is between 1.14 – 1.25 and 1.84-2.21 (i.e., $\geq$1.14 – 1.25 and $\leq$ 1.84 – 2.21). Or, other ranges may be used depending upon the patient cohort and/or

the significant event being monitored. Thus, each of the threshold values and/or threshold ranges described herein may vary by about 0.5 log units or less on a log scale and/or 25% or less on a linear scale (i.e., be $\leq$ 25% larger and/or $\leq$ 25% smaller than the specific ranges disclosed herein), or by about 20% or less, or by about 15% or less, or by about 10% or less, or by about 5% or less.

Also, in certain embodiments of each of the methods and/or aspects of the invention as disclosed herein, the level of p95 may be evaluated as either high or low. In some embodiments the first threshold value for p95 is a total p95 value of $\geq$ 90 (on a linear scale) and a low p95 value is below this threshold. Or, other ranges may be used depending upon the patient cohort and/or the significant event being monitored. Thus, each of the threshold values and/or threshold ranges described herein may vary by about 0.5 log units or less on a log scale and/or 25% or less on a linear scale (i.e., be $\leq$ 25% larger and/or $\leq$ 25% smaller than the specific ranges disclosed herein), or by about 20% or less, or by about 15% or less, or by about 10% or less, or by about 5% or less.

In certain embodiments, if the level of Her-3 is high, the patient is less likely or unlikely to respond to the targeted therapy. In certain embodiments, if the level of Her-3 is low, the patient is more likely to respond to the targeted therapy. In certain embodiments, the therapy is a Her-acting agent. In certain embodiments, the therapy is at least one of a Her-2 acting agent or a Her-3-targeted agent.

Thus, in certain embodiments of each of the methods and aspects of the invention as disclosed herein, the method comprises measuring in a biological sample from the subject's cancer an amount of Her-2 and/or Her-2 homodimers, wherein if the amount of Her-2 and/or Her-2 homodimers is moderately high and Her-3 expression is low, then the patient is likely to respond to the Her-2 acting agent and/or the patient has a long time course. In certain embodiments, the method comprises measuring in a biological sample from the subject's cancer an amount of Her-2 and/or Her-2 homodimers, wherein if the amount of Her-2 and/or Her-2 homodimers is moderately high and Her-3 expression is high, then the patient is unlikely to respond to the Her-2 acting agent and/or the patient has a short time course.

Additionally and/or alternatively, in certain embodiments of each of the methods and aspects of the invention as disclosed herein, the method comprises measuring in a biological sample from the subject's cancer an amount p95, wherein if the amount of p95 and Her-3 expression is low, then the patient is likely to respond to the therapeutically acting agent and/or the patient has a long time course. In an embodiment, the patient also has a high (or moderately high) level of Her-2. In certain embodiments, the method comprises measuring

in a biological sample from the subject's cancer an amount of Her-2 and/or Her-2 homodimers, wherein if the amount of Her-2 and/or Her-2 homodimers is high and/or moderately high and Her-3 expression and/or p95 expression is high, then the patient is unlikely to respond to the Her-2 acting agent and/or the patient has a short time course.

In a preferred embodiment, the sample is a blood, plasma or lymph sample. In a preferred embodiment, the blood or plasma sample contains circulating tumor cells. In a preferred embodiment, the sample comprises cell lines. In a preferred embodiment, the measurement may be quantitative across a wide dynamic range.

In a preferred embodiment, the method comprises mixing the sample with a binding compound and determining the presence and/or quantity of the binding compound bound to Her- 3 and/or Her-3 in a complex. In a preferred embodiment, the binding compound binds specifically to Her-3. In a preferred embodiment, the binding compound comprises an antibody. In a preferred embodiment, the antibody is raised against one of the peptides having SEQ ID NOs:1-8, as set forth in Example 2 and shown in Figure 2A. In a preferred embodiment, the antibody is a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:13, 14 and 15, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:16, 17 and 18, respectively; and/or a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:19, 20 and 21, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:22, 23 and 24, respectively. In a preferred embodiment, the antibody is the antibody with the amino acid sequence having SEQ ID NOs:9 and 11 as set forth in Table 1 (see Detailed Description) for the light and heavy chains, respectively, and/or SEQ ID NOs:10 and 12 as set forth in Table 1 (see Detailed Description) for the light and heavy chains, respectively. In a preferred embodiment, the method comprises mixing (i) a sample that may contain Her-3 and/or Her-3 in a complex; (ii) a proximity probe that is capable of binding Her-3, the proximity probe having an effective proximity and (iii) at least one binding compound, the at least one binding compound being capable of binding Her-3 and having one or more signaling molecules attached, wherein binding of the proximity probe and the binding compound within the effective proximity produces a signal from the molecular tags that correlates with the presence and/or quantity of Her-3 and/or Her-3 in a complex. In a preferred embodiment, the proximity probe and/or binding compound is capable of binding specifically to Her-3 or the

at least one other analyte. In a preferred embodiment, the proximity probe and/or binding compound further comprises an antibody and each antibody can bind to a specific epitope on Her-3. In a preferred embodiment, the antibody is raised against one of the peptides having SEQ ID NOs:1-8, as set forth in Example 2 and shown in Figure 2A. In a preferred embodiment, the antibody is a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:13, 14 and 15, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:16, 17 and 18, respectively; and/or a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:19, 20 and 21, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:22, 23 and 24, respectively. In a preferred embodiment, the antibody is the antibody with the amino acid sequence having SEQ ID NOs:9 and 11 as set forth in Table 1 (see Detailed Description) for the light and heavy chains, respectively, and/or SEQ ID NOs:10 and 12 as set forth in Table 1 (see Detailed Description) for the light and heavy chains, respectively. In a preferred embodiment, the sample is a biological sample. In a preferred embodiment, the sample is a tissue sample. In a preferred embodiment, the sample is a fixed sample, a frozen sample or a lysate. In a preferred embodiment, the sample is a tumor sample. In a preferred embodiment, the sample is a frozen tumor tissue sample. In a preferred embodiment, the sample comprises a tumor lysate. In a preferred embodiment, the sample comprises a breast cancer sample. In a preferred embodiment, the sample is an FFPE sample or solubilized FFPE sample. In a preferred embodiment, the sample is a blood, plasma or lymph sample. In a preferred embodiment, the blood or plasma sample contains circulating tumor cells. In a preferred embodiment, the sample comprises cell lines. In a preferred embodiment, the measurement may be quantitative across a wide dynamic range. In a preferred embodiment, the wide dynamic range is approximately 2 logs. In a more preferred embodiment, the wide dynamic range is about 1-1.5 logs in breast cancer samples. In a preferred embodiment, the method provides a quantitative continuum of Her-3 expression. In a preferred embodiment, the measurement or quantity is sensitive to at least about 1000 receptors per cell to about 200,000 receptors per cell as determined by accuracy studies utilizing well-characterized cell line models and cross-validating technologies such as ELISA and flow cytometry. In a preferred embodiment, the measurement or quantity is sensitive to at least about 5000 receptors per cell to about 200,000 receptors per cell. In a preferred

embodiment, the measurement or quantity is sensitive to at least about 10,000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement or quantity is sensitive to at least about 25,000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement is specific as determined using isotype control antibodies and comparison with conventional IHC methods.

In a further preferred embodiment, the proximity probe comprises an antibody and a first nucleic acid and the binding compound comprises an antibody and a second nucleic acid, wherein the first and the second nucleic acids are complementary to each other and able to hybridize to determine the effective proximity and produce the signal, directly or indirectly, through hybridization. In a preferred embodiment, the proximity probe and/or binding compound is capable of binding specifically to Her-3. In a preferred embodiment, the binding compound and/or the proximity probe further comprises an antibody and each antibody binds to a different epitope on Her-3. In a preferred embodiment, the antibody is raised against one of the peptides having SEQ ID NOs: 1-8, as set forth in Example 2 and shown in Figure 2A. In a preferred embodiment, the antibody is a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:13, 14 and 15, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:16, 17 and 18, respectively; and/or a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:19, 20 and 21, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:22, 23 and 24, respectively. In a preferred embodiment, the antibody is the antibody with the amino acid sequence having SEQ ID NOs:9 and 11 as set forth in Table 1 (see Detailed Description) for the light and heavy chains, respectively, and/or SEQ ID NOs:10 and 12 as set forth in Table 1 (see Detailed Description) for the light and heavy chains, respectively. In a preferred embodiment, the sample is a biological sample. In a preferred embodiment, the sample is a tissue sample. In a preferred embodiment, the sample is a fixed sample, a frozen sample or a lysate. In a preferred embodiment, the sample is a tumor sample. In a preferred embodiment, the sample is a frozen tumor tissue sample. In a preferred embodiment, the sample comprises a tumor lysate. In a preferred embodiment, the sample comprises a breast cancer sample. In a preferred embodiment, the sample is an FFPE sample or solubilized FFPE sample. In a preferred embodiment, the sample is a blood, plasma or lymph sample. In a preferred

embodiment, the blood or plasma sample contains circulating tumor cells. In a preferred embodiment, the sample comprises cell lines. In a preferred embodiment, the measurement may be quantitative across a wide dynamic range. In a preferred embodiment, the wide dynamic range is about 2 logs. In a more preferred embodiment, the wide dynamic range is about 1-1.5 logs in breast cancer samples. In a preferred embodiment, the method provides a quantitative continuum of Her-3 expression. In a preferred embodiment, the measurement or quantity is sensitive to at least about 1000 receptors per cell to about 200,000 receptors per cell as determined by accuracy studies utilizing well-characterized cell line models and cross-validating technologies such as ELISA and flow cytometry. In a preferred embodiment, the measurement or quantity is sensitive to at least about 5000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement or quantity is sensitive to at least about 10,000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement or quantity is sensitive to at least about 25,000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement is specific as determined by using isotype control antibodies and comparison with conventional IHC methods.

In a preferred embodiment, the proximity probe comprises a cleaving probe that has a cleavage-inducing moiety and the at least one binding compound has one or more molecular tags attached to the binding compound by a cleavable linkage, wherein the cleavable linkage may be cleaved within the effective proximity, producing a signal that correlates with the presence and/or quantity of Her-3. In a preferred embodiment, the cleaving probe and/or binding compound is capable of binding specifically to Her-3. In a preferred embodiment, the binding compound and/or the proximity probe further comprises an antibody and each antibody binds to a different epitope on Her-3. In a preferred embodiment, the antibody is raised against one of the peptides having SEQ ID NOs: 1-8, as set forth in Example 2 and shown in Figure 2A. In a preferred embodiment, the antibody is a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:13, 14 and 15, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:16, 17 and 18, respectively; and/or a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:19, 20 and 21, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:22, 23 and 24, respectively. In a preferred embodiment, the antibody is the antibody with the

amino acid sequence having SEQ ID NOs:9 and 11 as set forth in Table 1 (see Detailed Description) for the light and heavy chains, respectively, and/or SEQ ID NOs:10 and 12 as set forth in Table 1 (see Detailed Description) for the light and heavy chains, respectively. In a preferred embodiment, the sample is a biological sample. In a preferred embodiment, the sample is a tissue sample. In a preferred embodiment, the sample is a fixed sample, a frozen sample or a lysate. In a preferred embodiment, the sample is a tumor sample. In a preferred embodiment, the sample is a frozen tumor tissue sample. In a preferred embodiment, the sample comprises a tumor lysate. In a preferred embodiment, the sample comprises a breast cancer sample. In a preferred embodiment, the sample is an FFPE sample or solubilized FFPE sample. In a preferred embodiment, the sample is a blood, plasma or lymph sample. In a preferred embodiment, the blood or plasma sample contains circulating tumor cells. In a preferred embodiment, the sample comprises cell lines. In a preferred embodiment, the measurement may be quantitative across a wide dynamic range. In a preferred embodiment, the wide dynamic range is about 2 logs. In a more preferred embodiment, the wide dynamic range is about 1-1.5 logs in breast cancer samples. In a preferred embodiment, the method provides a quantitative continuum of Her-3 expression. In a preferred embodiment, the measurement or quantity is sensitive to at least about 1000 receptors per cell to about 200,000 receptors per cell as determined by accuracy studies utilizing well-characterized cell line models and cross-validating technologies such as ELISA and flow cytometry. In a preferred embodiment, the measurement or quantity is sensitive to at least about 5000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement or quantity is sensitive to at least about 10,000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement or quantity is sensitive to at least about 25,000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement is specific as determined by using isotype control antibodies and comparison with conventional IHC methods.

In a second aspect, the invention is drawn to a method for determining whether a subject with a cancer is likely to respond to treatment with a targeted therapy, for predicting a time course of disease and/or for predicting the probability of a significant event in the time course of the subject's cancer, comprising measuring in a biological sample from the subject's cancer an amount of Her-3, wherein the method is dependent on the level of Her-3. In certain embodiments, if the level of Her-3 is high, the patient is less likely or unlikely to respond to the targeted therapy. In certain embodiments, if the level of Her-3

is low, the patient is more likely to respond to the targeted therapy. In certain embodiments, as as described in more detail herein, the therapy is a Her acting agent. In further embodiments, the therapy is at least one of a Her-2 acting agent or a Her-3-targeted agent. In certain embodiments, the amount of Her-3 is above a first threshold, such that the sample is stratified as having a "high" amount of Her-3 (e.g., either total Her-3 and/or Her-3 homodimers and/or Her-3 heterodimers). In some embodiments the first threshold value for her-3 is a total Her-3 (H3T) of ≥ 0.158 and a low Her-3 value is below this threshold. Or, other ranges may be used depending upon the patient cohort and/or the significant event being monitored. Thus, each of the threshold values and/or threshold ranges described herein may vary by about 0.5 log units or less on a log scale and/or 25% or less on a linear scale (i.e., be $\leq 25\%$ larger and/or $\leq 25\%$ smaller than the specific ranges disclosed herein), or by about 20% or less, or by about 15% or less, or by about 10% or less, or by about 5% or less.

In a preferred embodiment, the subject's cancer is breast cancer, colorectal cancer, ovarian cancer, bladder cancer, prostate cancer, non-small cell lung cancer, melanoma, pharyngeal cancer, pancreatic cancer, esophageal cancer, glioma, biliary tract carcinoma, cholangiocarcinoma, gastric cancer, endometrial cancer, gall bladder cancer, squamous cell carcinoma or basal cell carcinoma. In a preferred embodiment, the subject's cancer is breast cancer, melanoma, synovial carcinoma, colorectal cancer or ovarian cancer. In a preferred embodiment, the subject's cancer is a Her-2 positive breast cancer. In a certain embodiments, the breast cancer is early stage (i.e., adjuvant) breast cancer or metastatic breast cancer.

As noted above, in certain embodiments, the method comprises detection of other biomarkers in the sample. For example, other biomarkers such as Her-2 and/or p95 may be measured. Or, other biomarkers such as can be at least one of FOXM1, PRAME, Bc12, STK15, CEGP1, Ki-67, GSTM1, CA9, PR, BBC3, NME1, SURV, GATA3, TFRC, YB-1, DPYD, GSTM3, RPS6KB1, Src, Chkl, ID1, EstR1, p27, CCNB1, XIAP, Chk2, CDC25B, IGF1R, AK055699, P13KC2A, TGFB3, BAGI1, CYP3A4, EpCAM, VEGFC, pS2, hENT1, WISP1, HNF3A, NFKBp65, BRCA2, EGFR, TK1, VDR, Contig51037, pENT1, EPHX1, IF1A, CDH1, HIFlα, IGFBP3; CTSB, Her3 or DIABLO. In certain embodiments, the other biomarker can be VEGF, CD31, KDR, p95, or Her-2.

In certain embodiments, the level of Her-2 expression in the breast cancer is high. In certain embodiments, high Her-2 expression is a log10H2T $\geq$ about 1.14-1.25. In certain embodiments, the high Her-2 expression comprises expression that is very high

and/or moderately high. In certain embodiments, the very high Her-2 expression is a log10H2T ≥ about 1.84 -2.21. In certain embodiments of each of the methods disclosed herein, the moderately high expression is between 1.14 – 1.25 and 1.84-2.21 (i.e., ≥1.14 – 1.25 and ≤ 1.84 – 2.21). Or, other ranges may be used depending upon the patient cohort and/or the significant event being monitored. Thus, each of the threshold values and/or threshold ranges described herein may vary by about 0.5 log units or less on a log scale and/or 25% or less on a linear scale (i.e., be ≤ 25% larger and/or ≤ 25% smaller than the specific ranges disclosed herein), or by about 20% or less, or by about 15% or less, or by about 10% or less, or by about 5% or less.

Also, in certain embodiments, the level of p95 may be evaluated as either high or low. In some embodiments the first threshold value for p95 is a total p95 value of > or ≥ 90 (on a linear scale) and a low p95 value is below this threshold. Or, other ranges may be used depending upon the patient cohort and/or the significant event being monitored. Thus, each of the threshold values and/or threshold ranges described herein may vary by about 25% or less (i.e., be ≤ 25% larger and/or ≤ 25% smaller than the specific ranges disclosed herein), or by about 20% or less, or by about 15% or less, or by about 10% or less, or by about 5% or less.

In certain embodiments, if the level of Her-3 is high, the patient is less likely or unlikely to respond to the targeted therapy. In certain embodiments, if the level of Her-3 is low, the patient is more likely to respond to the targeted therapy. In certain embodiments, the therapy is a Her-acting agent. In certain embodiments, the therapy is at least one of a Her-2 acting agent or a Her-3-targeted agent.

Thus, in certain embodiments, the method comprises measuring in a biological sample from the subject's cancer an amount of Her-2 and/or Her-2 homodimers, wherein if the amount of Her-2 and/or Her-2 homodimers is moderately high and Her-3 expression is low, then the patient is likely to respond to the Her-2 acting agent and/or the patient has a long time course. In certain embodiments, the method comprises measuring in a biological sample from the subject's cancer an amount of Her-2 and/or Her-2 homodimers, wherein if the amount of Her-2 and/or Her-2 homodimers is moderately high and Her-3 expression is high, then the patient is unlikely to respond to the Her-2 acting agent and/or the patient has a short time course.

Additionally and/or alternatively, in certain embodiments, the method comprises measuring in a biological sample from the subject's cancer an amount p95, wherein if the amount of p95 and Her-3 expression is low, then the patient is likely to respond to the

therapeutically acting agent and/or the patient has a long time course. In an embodiment, the patient also has a high (or moderately high) level of Her-2. In certain embodiments, the method comprises measuring in a biological sample from the subject's cancer an amount of Her-2 and/or Her-2 homodimers, wherein if the amount of Her-2 and/or Her-2 homodimers is high and/or moderately high and Her-3 expression and/or p95 expression is high, then the patient is unlikely to respond to the Her-2 acting agent and/or the patient has a short time course.

In a preferred embodiment, the targeted therapy is at least one Her family-targeted agent. In a preferred embodiment, the Her family-targeted agent is a multi- or single-targeted agent. In a preferred embodiment, the multi-targeted agent is a dual kinase inhibitor or a bispecific antibody. In a preferred embodiment, the Her family-targeted agent is trastuzumab, lapatinib or pertuzumab. In a preferred embodiment, the at least one Her family-targeted agent is at least two agents, wherein the at least two agents are one or more Her-2-targeted monoclonal antibodies and/or EGFR-targeted monoclonal antibodies and/or an EGFR and Her-2 dual kinase inhibitor. In a preferred embodiment, the monoclonal antibody is trastuzumab. In a preferred embodiment the EGFR-targeted monoclonal antibody is cetuximab or panitumumab. In a preferred embodiment, the dual kinase inhibitor is lapatinib, erlotinib or gefitinib. In a preferred embodiment, the targeted therapy is a Her-3 or Her-3 signaling pathway acting agent. In a preferred embodiment, the Her-3 or Her-3 signaling pathway targeted agent is a Her-3 monoclonal antibody, a Her-3 dimerization inhibitor, a Her-3 phosphorylation inhibitor and/or an inhibitor of a Her-3 signaling pathway member selected from the group consisting of PI3K, Akt, mTOR and ERK1/2. In a preferred embodiment, likeliness to respond, likeliness to have a long time course and/or likeliness to have a significant event is measured as an overall survival rate, as time to progression, as disease-free survival, as progression-free survival, and/or as objective tumor response using the RECIST criteria.

In a preferred embodiment, whether the cancer is Her-2 positive is determined by IHC, FISH, CISH, quantitative mRNA, a hybridization array, or VERATAG®. In a preferred embodiment, determining the level of Her-3 is performed using IHC, FISH, CISH, quantitative mRNA, hybridization array, or VERATAG®.

In a preferred embodiment, the method further comprises determining that a subject is afflicted with a Her-2 positive cancer that is unlikely to respond to treatment according to a method of the invention, then advising a medical professional of the treatment option of administering to the subject an effective amount of a different therapeutic agent.

In a third aspect, the invention is drawn to a purified antibody that binds to Her-3. In a preferred embodiment, the antibody is a polyclonal antibody or a monoclonal antibody. In a preferred embodiment, the antibody is a monoclonal antibody. In a preferred embodiment, the antibody is raised against one of the peptides having SEQ ID NOs:1-8, as set forth in Example 2 and shown in Figure 2A. In a preferred embodiment, the antibody is a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:13, 14 and 15, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:16, 17 and 18, respectively; and/or a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs: 19, 20 and 21, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:22, 23 and 24, respectively. In a preferred embodiment, the antibody is the antibody with the amino acid sequence having SEQ ID NOs:9 and 11 as set forth in Table 1 (see Detailed Description) for the light and heavy chains, respectively, and/or SEQ ID NOs:10 and 12 as set forth in Table 1 (see Detailed Description) for the light and heavy chains, respectively.

Brief Description of the Drawings

Figure 1 shows the levels of Her-3 expression as determined by a Her-3 ELISA kit (R&D Systems, Inc.) in several stably-transfected clones of HEK 293 (human embryonic kidney cells) transfected with an HER3 expression vector. The construction of the expression vector is described in Example 1. One clone, 293H3-Clone 1, expressed high levels of HER3 and was selected as a control for use in the optimized HER3 VERATAG® assay.

Figure 2A lists peptide sequences used to immunize mice to raise Her-3-specific antibodies, wherein SEQ ID NO: 1 is LGSALSLPVLNRPRGTGQSLLSP; SEQ ID NO: 2 is SAYHSQRHSLLTPVTPLSP; SEQ ID NO: 3 is VGSDLSASLGSTQSCPLHPVPI; SEQ ID NO; 4 is CQGPGHQAPHVHYARLKTLRS; SEQ ID NO; 5 is LEEVELEPELDLDLDLEAE; SEQ ID NO; 6 is CFDNPDYWHSRLFPKANA; SEQ ID NO: 7 is CPDYWHSRLFPKANAQRT; and SEQ ID NO; 8 is CFPKANAQRT. The peptide sequences represent different epitopes from the C-terminal region of Her-3 (the length of each peptide and the position relative to the N-terminus of the protein are shown). Antibodies raised against each peptide are listed in the fourth column. Each antibody has been confirmed to test positive in ELISA, IHC, and VERATAG® assays. Figure 2B shows

the results of IHC studies in which two cell lines were screened with B9A11, an Her-3-specific antibody raised against one of the Her-3 peptides, an Her-2-specific antibody Ab8 (HerceptTest™), and a control antibody ITC-IgG2a. One cell line, SKOV3 (upper panels), is known to express high levels of HER2 and low levels of HER3. The other cell line, 293H3-clone1 (lower panels), is the stablytransfected cell line described in Example 1 and shown in Figure 1 to over-express HER3 but which expresses low levels of HER2. The IHC results show a strong signal of B9A11 with the 293H3-clone 1 cells but not the SKOV3 cells, as expected.

Figure 3 shows HER3 levels as determined by VERATAG® in FFPE blocks from 4 different cell lines, cross-validated with data from three other assays (IHC, ELISA, and flow cytometry). Cell lines expressing varying levels of HER3 were chosen for these studies: 293H3-Clone 1, MDA-MB-453, MDAMB-468 and SKOV3 (the latter 3 from ATCC). The cell lines were chosen to represent HER3 receptor levels that spanned greater than 2 logs. FFPE blocks were prepared as described in Example 3 for testing with the HER3 VERATAG® assay and IHC. A portion of the cells from the same lot were tested for HER3 receptor number using flow cytometry. Further, a whole cell lysate was prepared from the same lot of cells for quantifying HER3 with an ELISA kit (Human ErbB3-DuoSet ELISA: R&D Systems, Inc.). The data shows a wide dynamic range for the VERATAG® assay, with results consistent with all three other methodologies.

Figure 4 shows examples of patient tumor samples in which a Board-certified pathologist has circled the tumor area.

Figure 5 shows the equipment used and the work flow of the HER3 VERATAG® assay. FFPE samples are first deparaffinized and rehydrated using a series of solvents (top panel 1). Antigen retrieval is accomplished using 1x DAKO (Lab Vision) in a pressure cooker (top panel 2). The samples are then rinsed with water and a hydrophobic pen is used to draw a circle around the sample, retaining reagents on the slide. The samples are then blocked and treated with a mixture of VERATAG®-conjugated Ab-6 (Lab Vision) and biotin-conjugated B9A11 (top panel 3). After incubation and washing, streptavidin-conjugated methylene blue reagent is added, incubated and washed, and then illumination buffer containing fluorescein and two capillary electrophoresis internal markers (MF and ML, marker first and last, respectively) is added. The bound VERATAG® is released using an LED array, which photoactivates cleavage of the VERATAG® (top panel 4). VERATAG® intermediates are reduced to a quantifiable form

using sodium borohydride and the VERATAG® reporters are separated and detected using capillary electrophoresis (ABI3130 CE instrument, top panel 5).

Figure 6 shows the results from an experiment designed to identify the optimal antibody concentration for maximizing the dynamic range of the VERATAG® assay. Cell lines spanning the entire dynamic range of the assay were chosen: 293H3-Clone 1, MDA-MB-453, MDA-MB-468, MDA-MB-231 and SKOV3. The concentration of the antibodies B9A11-biotin and Ab-6 Pro-11 were varied (column 1 of the table) as follows: 1 mg/mL B9A11-biotin and 1 mg/mL Ab-6 Pro-11, 1 mg/mL B9A11-biotin and 2 mg/mL Ab-6 Pro-11 and 2 mg/mL B9A11- biotin and 1 mg/mL Ab-6 Pro-11, in rows 4, 5 and 6, respectively. The results for each cell line are shown in the bar graph. Expected fold changes for pair-wise comparisons were based on HER3 flow cytometry and ELISA results from the same cell line FFPE block preparation. An optimal concentration of 2 mg/mL B9A1 1-biotin and 1 mg/mL Ab-6 Pro-11 was chosen (circled) for best performance based on the accurate detection of HER3 as compared with the expected fold changes shown in row 2 of the table. The dynamic range shown here is approximately 2 logs.

Figure 7 shows the accuracy of the HER3 VERATAG® assay using three successful replicates from four well-characterized cell lines (293H3-Clone 1, MDA-MB-453, MDA-MB-468 and SKOV3). The VERATAG® measurements were compared with in-house generated flow cytometry and ELISA data. 100% of the results matched the in-house data from flow cytometry and ELISA in that 293H3-clone 1> MDA-MB-453>MDA-MB-468>SKOV3. No overlap was observed between signal levels for any of the four cell line samples.

Figure 8 demonstrates the sensitivity of the HER3 VERATAG® assay. One batch containing 8 replicates of the low HER3 expression control cell line, MDA-MB-468, was compared with 8 replicates of the low/negative HER3 expression control cell line, SKOV3, to determine sensitivity. All of the pairwise comparisons (64/64) between MDA-MB-468 and SKOV3 resulted in MDA-MB-468 having higher levels of HER3 than SKOV3.

Figure 9 shows the inter-assay reproducibility of the HER3 VERATAG® assay. Eight separate HER3 total VERATAG® assays were performed on the four well-characterized cell lines, 293H3-Clone 1, MDA-MB-453, MDA-MB-468 and SKOV3, using different CE illuminators, several operators and on different days over a 4 week period. Following batch normalization procedures, the data was compared across the 8 batches to ascertain reproducibility. The coefficient of variability across the dynamic range was between 8 and

15%. Values are represented as the Log_{10} normRPA, which is the log of the normalized relative peak area/tumor area and then batch normalized using expected values.

Figure 10 shows the precision of the HER3 VERATAG® assay. The intra-assay reproducibility of the HER3 VERATAG® assay was demonstrated by comparing the performance of 15 replicates of each of the 3 control cell lines, 293H3-Clone 1, MDA-MB-453 and MDA-MB468. Pairwise comparisons were made of the 15 replicates in each batch to determine precision. 95% of the 293H3-Clone 1 data was within 1.2-fold and 95% of the MDA-MB-468 data is within 1.37-fold. The VERATAG® data (shown in normalized RPA) for the 15 replicates of each cell line is shown by the 15 bars on the right of each panel. The control data for the 3 cell lines expressing moderate to high levels of HER3 and a low/negative HER3 expressing cell line, SKOV3, are shown on the left of each panel.

Figure 11 shows the linearity of the HER3 VERATAG® assay using different sample sizes. Samples of diminishing size (1x, 1/2x, 1/4x, 1/16x) from each of the 3 well-characterized cell lines, 293H3-Clone 1, MDA-MB-453 and MDA-MB-468, were tested in the VERATAG® assay and the data was compared in a pairwise manner to assess the linearity of the assay. The MDA-MB-453 cell line shows linearity down to approximately $1/16^{th}$ of the original sample size; MDA-MB-468 shows linearity down to approximately $1/2^{th}$ of the original sample size.

Figure 12 shows the specificity of the HER3 VERATAG® assay as determined by isotype controls. Isotype control antibodies were tested in the VERATAG® assay format to ascertain the non-specific background of the assay. For the HER3 Ab-6-Pro11 antibody, the isotype control was IgG1-Pro11. For the HER3 B9A11-biotin antibody, the isotype control was IgG1-biotin. Signal derived using these isotype controls is not antigen-specific and therefore represents non-specific background. In the each panel, the VERATAG® results are shown for the normal assay format using the HER3 Ab6- Pro11 and B9A11-biotin antibodies in the bars labeled "control." In the left panel, the VERATAG® data is shown for the HER3 Ab6-Pro1 1 and IgG1-biotin antibodies in the bars labeled "IgG-bio." In the right panel, the VERATAG® data is shown for the HER3 B9A11-biotin and IgG1-Pro11 antibodies in the bars labeled "IgG-Pro11." Each antibody pairing was tested on an array of FFPE samples including standard cell line controls (293H3-Clone 1, MDA-MB-453, MDA-MB468, SKOV3 and T47D) as well as several tumor samples (41776B1, 32712A2, 30345C2, 106305A2, and 106341A2). Units are normalized RPA*IB/TA.

Figure 13 shows VERATAG® data on tumor samples from the International Serum Her-2/neu Study Group trial. This cohort of patients (n=105) was prospectively observed

during trastuzumab treatment between 1999 and 2006. All patients were determined to be Her-2 positive by either IHC or FISH and had not been exposed to trastuzumab prior to the study. The samples were evaluated for HER3 levels using the HER3 VERATAG® assay in eight separate batches. The results are shown in the 8 panels in this figure. Each panel also includes results for 5 control cell lines: 293H3-Clone 1, MDA-MB-453, MDA-MB-468, SKOV3 and T47D (shown on the left of each panel). Results are shown in $\log_{10}$ normalized RPA units.

Figure 14 shows the data from positional scanning analyses used to determine the optimal cut-off for trastuzumab (i.e., herceptin) response. In the left panel, patients above the statistically significant cut-off (see arrow) had an unfavorable time to progression (TTP) compared to patients below the cut-off (hazard ratio = 2.3; p = 0.0004). In the right panel, a significant cutoff could not be determined, but using the cut-off found for TTP, a trend for worse overall survival (OS) in patients above the cut-off (indicated by the arrow) was observed (hazard ratio =1.7; p = 0.059).

Figure 15 shows Kaplan-Meier plots for a cohort of 82 trastuzumab-treated patients stratified first by total HER2 levels (H2T) and then further stratified by total HER3 levels (H3T). In the upper left panel, a Kaplan-Meier plot shows the percent of patients with progression-free survival (months) for two groups of patients subdivided (based on a previously reported cut-off) into HER2-normal and HER-2 over-expressing groups. The HER2 over-expressing group was then further subdivided, using the cut-off shown in Figure 14, into two subgroups based on the level of HER3. The Kaplan-Meier plot of these two subgroups is shown in the upper right panel. The lower panel shows three sets of results from the upper panels: the normal HER2 group (log(H2T)<1.14), the HER2-high, HER3-low group (log(H2T)>1.14, H3T<0.158) and the HER2-high, HER3-high group (log(H2T)>1.14, H3T>0.158). Univariate Cox proportional hazards analyses examining the HER3-over-expressing subgroup identified H3T (high vs low) as the most significant predictor of time to progression (TTP; HR = 2.98, p = 0.0004).

Figure 16 shows the results of HER3 VERATAG® assays in synovial carcinoma, colon cancer and ovarian cancer. The top panel shows the results for a panel of synovial carcinoma samples. The 4 bars on the left are control samples (293H3-Clone 13, MDA-MB-453, MDAMB-468 and SKOV3, from left to right, respectively). The middle panel shows the results for a panel of colon cancer samples. The same control cell lines are used (shown in the left 4 bars). The lower panel shows results comparing HER2 expression to HER3 expression in a panel of ovarian tumor samples. Results are shown in normalized

RPA units. The dynamic range of these tumor samples ranges from 0.5-1.5, depending on the cancer.

Figure 17 shows that with the High HER2 (H2T) group (log10H2T >1.25, or >13.8 on the linear scale), the ability to subgroup patients into different groups based on High or low p95 (p95 >90 or <=90) and high or low HER3 (H3T) allows further stratification of clinical outcomes as measured by median TTP. Univariate KM analysis with the p95 and H3T subgroups combined, gives the results in the KM plots in this Figure. These data suggest that HER2-positive breast cancer patients as assessed by HERmark/VERATAG® (i.e., high H2T, ie log10H2T>1.25, or >13.8 on the linear scale) can be classified into at least 4 sub-groups with different outcomes following trastuzumab treatment. In the KM, the group with low H3T (in generation 1 of the HER3 assay, H3T< 0.158) and low p95 (p95<90) has a median TTP of 15.0 months, compared with 9.3 months for the group with low H3T (H3T<0.158) and high p95 (p95>90); compared with 6.4 months for the high H3T group (H3T>0.158 in generation 1 of the HER3 assay) and low p95 group, and compared with 3.2 months for the high H3T (H3T>0.158) and high p95 (p95>90). The trend for the differences among the groups is significant (p<0.0001).

Figure 18 shows Kaplan-Meier (KM) analyses comparing the percent progression free on the y axis over time on the x axis (time to progression, TTP) of various subgroups from the Lipton cohort, as defined by the combined VERATAG® measurements of HER2 total (H2T high or low), p95HER2 (p95 high or low), and HER3 Total (H3T) high (H3T>0.158, in Generation 1 of the H3T assay) or H3T low (H3T<=0.158 in Generation 1 of the assay). Cut-offs were identified by lowest p-value in a positional scanning analysis. H2T high = (log10H2T > 1.25 or on a linear scale, >13.8). Low H2T = log10H2T <= 1.25 or on a linear scale, <=13.8. p95 low = p95<=90 and p95 high = p95>90 (on a linear scale), and H3T hi >0.158 on a linear scale, and H3Tlo <=0.158 on a linear scale). KM analyses demonstrated that patients who were FISH positive, H2T high, p95 lo (low) and H3T lo (low) had a median TTP of 14.7 months, compared with the 4 other groups that did not fare as well. Three groups with nearly superimposable lines (i.e., FISHnegative/H2Tlo group – median TTP = 4.5, FISH positive/H2Tlo(low) group-median TTP = 3.7, and FISHpositive/H2Thi (high)/p95hi (high)/H3T hi (high)- median TTP = 3.2) all had shorter median TTP than the group FISH positive/H2Thigh/p95lo/H3Tlo-median TTP = 15). The group defined as FISHpositigve/H2Thi(high)/and p95orH3Thigh had a median TTP = 9.3, which was in between the group with the best median TTP (FISH positive/H2Thigh/p95lo/H3Tlo) and the 3 groups in red/blue and black. Thus, HERmark

assay identified multiple subgroups of HER2 positive patients with varying clinical outcomes as measured by TTP on trastuzumab-based therapy. Neither the magnitude of HER2 over-expression nor the outcome for these subgroups was predictable by FISH/CEP17 copy number. HER2 FISH positive MBC patients with high p95 and/or high H3T may represent subsets of patients with *de novo* resistance to trastuzumab. While the applicants do not wish to be confined to any particular mechanistic theory, possible mechanisms that may account for the poor response to trastuzumab observed in these subgroups may include insufficient trastuzumab and/or lack of trastuzumab binding target (i.e., p95) and increased signaling via formation of heterodimers that are not completely suppressed by trastuzumab.

Detailed Description of the Invention

As used herein, the terms "embodiment" and "aspect" are used interchangeably.

"Antibody" means an immunoglobulin that binds to, and is thereby defined as complementary with, a particular spatial and polar organization of another molecule. The antibody can be monoclonal, polyclonal or recombinant and can be prepared by techniques that are well known in the art such as immunization of a host and collection of sera (polyclonal) or by preparing continuous hybrid cell lines and collecting the secreted protein (monoclonal) or by cloning and expressing nucleotide sequences or mutagenized versions thereof coding at least for the amino acid sequences required for binding. Antibodies may include a complete immunoglobulin or fragment thereof, which immunoglobulins include the various classes and isotypes, such as IgA, IgD, IgE, IgG1, IgG2a, IgG2b and IgG3, IgM, etc. Fragments thereof may include Fab, Fv and F(ab')2, Fab' and the like. Antibodies may also be single-chain antibodies, chimeric antibodies, humanized antibodies or any other antibody derivative known to one of skill in the art that retains binding activity that is specific for a particular binding site. In addition, aggregates, polymers and conjugates of immunoglobulins or their fragments can be used where appropriate so long as binding affinity for a particular binding site is maintained. Guidance in the production and selection of antibodies and antibody derivatives for use in immunoassays, including such assays employing releasable molecular tags (as described below) can be found in readily available texts and manuals, *e.g.*, Harlow and Lane, 1988, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, New York; Howard and Bethell, 2001, *Basic Methods in Antibody Production and Characterization*, CRC Press; Wild, ed., 1994, *The Immunoassay Handbook*, Stockton Press, New York.

"Binding compound" shall refer to a molecule capable of binding to another molecule of interest. A binding compound may be an antibody, a peptide, a peptide or non-peptide ligand for a cell surface receptor, a protein, an oligonucleotide, an oligonucleotide analog, such as a peptide nucleic acid, a lectin or any other molecular entity that is capable of specifically binding to a target molecule or complex. In one embodiment, the target molecule is a protein or protein complex. In another embodiment, a binding compound further comprises a proximity probe. In one embodiment, a binding compound comprises one or more molecular tags attached to a binding moiety. In another embodiment, a second binding compound may be bound to the binding compound and measured or quantified as a correlative for the presence of the binding compound, which is bound to the target protein. As another specific example, either the first or second binding compound may generate an effector molecule that acts in conjunction with a proximity probe with an effective proximity, producing a signal that correlates with the presence of the target protein. Further, in another embodiment, binding compounds may have molecular tags that interact with one another within an effective proximity to form a complex that generates a signal or can be detected and measured in a manner that correlates with the presence of the target protein. More specifically, the target protein or complex may be Her-3 or Her-3 in a complex.

"Binding moiety" means any molecule to which molecular tags can be directly or indirectly attached that is capable of binding to an analyte. Binding moieties include, but are not limited to, antibodies, peptides, proteins, nucleic acids and organic molecules having a molecular weight of up to about 1000 daltons and containing atoms selected from the group consisting of hydrogen, carbon, oxygen, nitrogen, sulfur and phosphorus. Preferably, binding moieties are antibodies.

"Cell lines" refers to cells that have been separated from their original tissue, clonally multiplied and/or maintained in culture. As specific examples, cell lines may be derived from each type of cancer and multiple different cell lines may be derived from samples of the same type of cancer. Examples of different types of cell lines include, but are not limited to, breast cancer cell lines, such as MCF-7, MDA-MB-453, MDA-MB-468, or T-47D or cell lines derived from other tissues, such as SKOV3 or HEK293.

"Chemotherapeutic agent" means a chemical substance that is used to treat a condition, particularly cancer.

A "cleavable linkage," as used herein, refers to a chemical linking group that may be cleaved to release a detectable molecular tag connected to a binding moiety with the cleavable linkage.

A "cleavage-inducing moiety," or "cleaving agent," as used herein, is a group that produces an active species that is capable of cleaving a cleavable linkage. Preferably, the active species is a chemical species that exhibits short-lived activity so that its cleavage-inducing effects are only in the proximity of the site of its generation.

A "cleaving probe," as used herein, refers to a reagent that comprises a cleavage-inducing moiety, as defined herein, and a binding compound such as an antibody, a peptide, a peptide or non-peptide ligand for a cell surface receptor, a protein, such as streptavidin, a small molecule, such as biotin, an oligonucleotide, an oligonucleotide analog, such as a peptide nucleic acid, a lectin or any other molecular entity that is capable of binding to a target protein or molecule or stable molecular complex.

"Dual kinase inhibitor" refers to molecules that inhibit more than one kinase, for example but not limited to, an inhibitor of both EGFR and Her-2 kinase activity, such as lapatinib.

"Effective proximity," as used herein, describes the distance between two binding compounds that is sufficient to generate a detectable signal, indicating the presence of the target molecule. For example, a proximity probe and a binding compound that are bound on Her-3 (or with another analyte of interest) within an effective proximity will generate a detectable signal, indicating and/or quantifying the presence of Her-3 and/or a Her-3 complex. Preferably, the effective proximity range for many detection systems is less than 200 nM, more preferably, less than 50 nM.

"EGFR", "ErbB1", "erbB-1", "HER1", "her-1", "Her-1"and refers to the epidermal growth factor receptor and allelic variants thereof, as described, for example, by Ono and Kuwano (*see* Ono and Kuwano (2006) *Cuin. Cancer Res.* 12:7242-7251) and Genbank accession number P00533. Unless indicated otherwise, the terms "EGFR", "ErbB1", "erbB-1", "HER1", "her-1 ", "Her-1" when used herein refer to the human protein.

"Epitope" refers to a site on the surface of a molecule, usually a protein, to which an antibody molecule or other binding compound binds. Generally, a protein has several or many different epitopes, also called antigenic determinants, and reacts with antibodies of different specificities.

"FFPE" shall refer to a group of cells or quantity of tissue that are fixed, particularly conventional formalin-fixed paraffin-embedded samples. Such samples are typically, for

example, without limitation, used in an assay for receptor complexes in the form of thin sections, *e.g.* 3-10 μm thick, of fixed tissue mounted on a microscope slide or equivalent surface. Such samples also typically undergo a conventional re-hydration procedure, and optionally, an antigen retrieval procedure as a part of, or preliminary to, assay measurements.

As used herein, "greater than or equal to" (i.e., ≥ or > =) can in certain alternative embodiments mean "greater than" (>). Also, as used herein, "less than or equal to" (i.e., ≤ or < =) can in certain alternative embodiments mean "less than" (<).

"Her-2", "ErbB2", "c-Erb-B2", "HER2", "Her2" and "*neu*" are used interchangeably herein and refer to native Her-2, and allelic variants thereof, as described, for example, in Semba *et al.*, 1985, *P.N.A.S. USA* 82:6497-650 and Yamamoto *et al.*, 1986, *Nature* 319:230-234 and Genebank accession number X03363. Unless indicated otherwise, the terms "Her-2", "ErbB2", "c-Erb-B2", "HER2" and "Her2" when used herein refer to the human protein. The gene encoding Her2 is referred to herein as "erbB2."

"Her-2-acting agent," as used herein, refers to a compound that can alter a biological activity of Her-2 or a Her-2 expressing cell or a Her-2 positive cancer cell. Such biological activities include, but are not limited to, dimerization, autophosphorylation, phosphorylation of another receptor, signal transduction and the like. Biological activities can include, without limitation, cell survival and cell proliferation and inhibition of such activities by a Her-2 acting agent could be direct or indirect cell killing (eg, ADCC), disruption of protein complexes or complex formation, modulation of protein trafficking or enzyme inhibition. Biological activities can also include patient response as set forth in this application. Exemplary Her-2-acting agents include, but are not limited to, the large molecules 4D5, pertuzumab, and trastuzumab and small molecules such as AEE-788 and lapatinib.A Her-2 complex is used to describe complexes of proteins, such as heterodimers, in which Her-2 is a component. A Her-2 complex may include a Her-2 homodimer, or a heterodimer that includes Her-2 (e.g., a Her-2/Her-3 heterodimer)

"Her-3", "ErbB3", "c-erb-B3", "erbB-3", "HER3" and "Her3" are used interchangeably herein and refer to native Her-3, and allelic variants thereof, as described, for example, in Kraus MH, *et al.* (1989) *Proc Natl Acad Sci U S A* 86:9193-9197 and Plowman GD, *et al.* (1990) *Proc Natl Acad Sci U S A.* 87:4905-4909 and Genbank accession number P21860. Unless indicated otherwise, the terms "Her-3", "ErbB3", "c-erb-B3", "erbB-3", "HER3" and "Her3" when used herein refer to the human protein. The gene encoding Her3 is referred to herein as "erbB3."

"Her-3 complex" is used to describe complexes of proteins, such as heterodimers, in which Her-3 is a component. Examples of heterodimers including Her-3 include but are not limited to Her-1/Her-3 and Her-2/Her-3.

"Her-3-targeted agent" or "Her-3 signaling pathway targeted agent" refers to therapeutics that alter the biological activity of Her-3 or members of the Her family signaling pathway. Such biological activities include, but are not limited to, dimerization, autophosphorylation, phosphorylation of another receptor, signal transduction and the like. Biological activities can include, without limitation, cell survival and cell proliferation and inhibition of such activities by a Her-3 or Her-3 signaling pathway member acting agent could be direct or indirect cell killing (eg, ADCC), disruption of protein complexes or complex formation, modulation of protein trafficking or enzyme inhibition. Biological activities can also include patient response as set forth in this application. Exemplary Her-3 or Her-3 signaling pathway member acting agents might include, but are not limited to, large molecules (such as antibodies) or small molecules (such as small molecule kinase inhibitors) targeted to Her-3, PI3K, Akt, mTOR ,ERK1/2, or PYK2.

"High" refers to a measure that is greater than normal, greater than a standard such as a predetermined measure or a subgroup measure or that is relatively greater than another subgroup measure. For example, high Her-3 refers to a measure of Her-3 that is greater than a normal Her-3 measure. A normal Her-3 measure may be determined according to any method available to one skilled in the art. High Her-3 may also refer to a measure that is equal to or greater than a predetermined measure, such as a predetermined cutoff. High Her-3 may also refer to a measure of Her-3 wherein a high Her-3 subgroup has relatively greater levels of Her-3 than another subgroup. For example, without limitation, according to the present specification, two distinct patient subgroups can be created by dividing samples around a mathematically determined point, such as, without limitation, a median, thus creating a subgroup whose measure is high (ie, higher than the median) and another subgroup whose measure is low. Her-3 can be measured by any method known to one skilled in the art such as, for example, without limitation, using VERATAG® or using any standard immunohistochemical (IHC) method . In some cases, a "high" expression level may comprise a range of expression that is very high and a range of expression that is "moderately high" where moderately high is a level of expression that is greater than normal, but less than "very high". Example ranges for high (including very high and moderately high) Her-2 expression and/or high Her-3 and/or high p95 expression are provided herein.

"IHC, FISH and CISH" are methods (immunohistochemistry, fluorescence *in situ* hybridization, and chromogenic *in situ* hybridization, respectively) used to detect the presence of molecular entities in cells or tissues. For example, membrane receptors such as Her-3 and/or other members of the EGFR family of receptors can be detected using these methods.

"Isotype control antibodies" refers to antibodies that have the same underlying immunoglobulin structure as a specific antibody used as a binding compound but that do not have specificity for the targeted epitope. The use of isotype control antibodies allows one to observe any binding that is due to non-specific binding.

"Likely to" (and "unlikely to"), as used herein, refers to an increased (or decreased) probability that an item, object, thing or person will occur. Thus, in one example, a subject that is likely to respond to treatment with trastuzumab has an increased probability of responding to treatment with trastuzumab relative to a reference subject or group of subjects.

"Long," as used herein, refers to a time measure that is greater than normal, greater than a standard such as a predetermined measure or a subgroup measure that is relatively longer than another subgroup measure. For example, with respect to a patient's longevity, a long time progression refers to time progression that is longer than a normal time progression. Whether a time progression is long or not may be determined according to any method available to one skilled in the art. In one embodiment, "long" refers to a time that is greater than the median time course required for a significant event to occur in a disease.

"Low" is a term that refers to a measure that is less than normal, less than a standard such as a predetermined measure or a subgroup measure that is relatively less than another subgroup measure. For example, low Her-3 means a measure of Her-3 that is less than a normal Her-3 measure in a particular set of samples of patients. A normal Her-3 measure may be determined according to any method available to one skilled in the art. Low Her-3 may also mean a measure that is less than a predetermined measure, such as a predetermined cutoff. Low Her-3 may also mean a measure wherein a low Her-3 subgroup is relatively lower than another subgroup. For example, without limitation, according to the present specification, two distinct patient subgroups can be created by dividing samples around a mathematically determined point, such as, without limitation, a median, thus creating a group whose measure is low (i.e., less than the median) with respect to another group whose measure is high (i.e., greater than the median). Her-3 can be measured by any method known to one skilled in the art such as, for example, without limitation, using the

VERATAG® method or using any standard immunohistochemical (IHC) method. Example ranges for low values of Her-3, Her-2, and p95 expression are provided herein.

"Lysate" refers to the solution produced when the cell membranes of cells are disrupted, whether by physical or chemical methods. For example, "tumor lysates" typically contain representative components of the cells comprising the tumor, including but not limited to, protein markers, enzymes, nucleic acids and complexes of proteins and other molecules that can subsequently be measured in various assays.

A "molecular tag," as used herein, refers to a molecule that can be measured directly or indirectly, can be distinguished from other molecules based on one or more physical, chemical or optical differences among the molecules being separated, including but not limited to, electrophoretic mobility, molecular weight, shape, solubility, pKa, hydrophobicity, charge, charge/mass ratio, polarity or the like. In one embodiment, molecular tags in a plurality or set differ in electrophoretic mobility and optical detection characteristics and can be separated by electrophoresis. In another embodiment, molecular tags in a plurality or set may differ in molecular weight, shape, solubility, pKa, hydrophobicity, charge, polarity and can be separated by normal phase or reverse phase HPLC, ion exchange HPLC, capillary electrochromatography, mass spectroscopy, gas phase chromatography or a like technique.

Measurement of molecular tags may also involve using secondary molecular interactions, with or without further modification, to detect, enhance or amplify a measurable signal that acts as a correlative for the presence and/or quantity of an analyte, such as Her-3 or a Her-3 in a complex. In one embodiment, a set of two or more molecular tags may interact within an effective proximity to produce a measurable signal. As molecular tags, a measurable signal may be generated, for example, by detection of two complementary nucleic acid sequences that will hybridize when the complementary sequences are within an effective proximity. Other examples that either generate a measurable signal or that can be measured using detection methods know in the art include, but are not limited to, FRET, BRET, BiFC, LCI and QPCR.

"Overall survival" or "OS" refers to a time as measured from the start of treatment to death or censor. Censoring may come from a study end or change in treatment. Overall survival can refer to a probability as, for example, a probability when represented in a Kaplan-Meier plot of being alive at a particular time, that time being the time between the start of the treatment to death or censor.

"Pre-determined cutoff" as used herein, refers to the value of a predetermined measure on subjects exhibiting certain attributes that allow the best discrimination between two or more categories of an attribute. For example, a pre-determined cutoff that allows one to discriminate between two categories such as high Her-3 expression and low Her-3 expression for determining overall survival may be used. Pre-determined cutoffs may be used to separate the subjects with values lower than or higher than the pre-determined cutoff to optimize the prediction model.

A "proximity probe," as used herein, refers to a reagent that comprises a moiety capable of acting within effective proximity to a molecular tag on a binding compound to generate a detectable signal and an antibody, a peptide, a peptide or non-peptide ligand for a cell surface receptor, a protein, such as streptavidin, a small molecule, such as biotin, an oligonucleotide, an oligonucleotide analog, such as a peptide nucleic acid, a lectin or any other molecular entity that is capable of specifically binding to a target protein or molecule or stable complex. For example, a proximity probe comprised of a Her-3-targeted antibody with a molecular tag may be capable of binding to Her-3 within an effective proximity to one or more Her-3 binding compounds, or a binding compound of another protein of interest, that has one or more molecular tags attached. In one embodiment, a proximity probe comprises a binding molecule and a first nucleic acid and a binding molecule comprises an antibody and a second nucleic acid, wherein the first and second nucleic acids are complementary to each other and each is a predetermined length so that when the nucleic acids are within an effective proximity of one another, they hybridize. Hybridization may be measured by any method known to one skilled in the art. For example, fluorophores may be attached to the nucleic acids as indicators of hybridization. In a preferred embodiment, hybridization is measured with a nucleic acid amplification method such as, for example, without limitation, the rolling circle amplification method (*see*, for example, Lizardi *et al,.*, (1998) *Nat Genet.* 19: 225-232).

"RECIST" shall mean "Response Evaluation Criteria in Solid Tumours" and is a set of published rules that define when cancer patients improve ("respond"), stay the same ("stable") or worsen ("progression") during treatments. Response as defined by RECIST criteria have been published, for example, at *Journal of the National Cancer Institute*, Vol. 92, No. 3, February 2, 2000 and RECIST criteria may include other similar published definitions and rule sets.

"Respond" to treatment, and other forms of this verb, as used herein, refer to the reaction of a subject to treatment with an agent. As an example, a subject responds to

treatment if growth of a tumor in the subject is retarded about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more. In another example, a subject responds to treatment if a tumor in the subject shrinks by about 5%, 10%, 20%, 30%, 40%, 50% or more as determined by any appropriate measure, *e.g.*, by mass or volume. In another example, a subject responds to treatment with a Her2-acting agent if the subject experiences a life expectancy extended by about 5%, 10%, 20%, 30%, 40%, 50% or more beyond the life expectancy predicted if no treatment is administered. In another example, a subject responds to treatment with an agent if the subject has an overall survival or increased time to progression. Several methods may be used to determine if a patient responds to a treatment including the RECIST criteria, as set forth herein.

"Sample" or "tissue sample" or "patient sample" or "patient cell or tissue sample" or "specimen" each refer to a collection of similar cells obtained from a tissue of a subject or patient. The source of the tissue sample may be solid tissue as from a fresh tissue, frozen and/or preserved organ or tissue or biopsy or aspirate; blood or any blood constituents, bodily fluids such as cerebral spinal fluid, amniotic fluid, peritoneal fluid or interstitial fluid or cells from any time in gestation or development of the subject. The tissue sample may contain compounds that are not naturally intermixed with the tissue in nature such as preservatives, anticoagulants, buffers, fixatives, nutrients, antibiotics or the like. Cells may be fixed in a conventional manner, such as in an FFPE manner.

"Short," as used herein, refers to a time measure that is shorter than normal, shorter than a standard such as a predetermined measure or a subgroup measure that is relatively shorter than another subgroup measure. For example, with respect to a patient's longevity, a short time progression refers to time progression that is shorter than a normal time progression or shorter than predicted. Whether a time progression is short or not may be determined according to any method available to one skilled in the art. In one embodiment, "short" refers to a time that is less than the median time course required for a significant event to occur in a disease.

"Signaling pathway", as used herein, refers to a process in which the binding of extracellular signaling molecules to cell-surface receptors trigger events inside the cell and/or a process in which intracellular signaling cascades can be triggered through intracellular interactions. For example, receptor tyrosine kinases are transmembrane proteins that propagate growth factor signals from the cell surface to intracellular processes that control critical functions such as growth, differentiation, angiogenesis and inhibition of apoptosis. In cancer, these signaling pathways are often exploited to facilitate tumor growth and

metastasis. One such family of receptor tyrosine kinases is the epidermal growth factor receptor (EGFR) family. EGFR family members, EGFR, HER2, HER3 and HER4, are over-expressed in a wide variety of tumor types.

"Significant event," as used herein, shall refer to an event in a patient's disease that is important as determined by one skilled in the art. Examples of significant events include, for example, without limitation, primary diagnosis, death, recurrence, the determination that a patient's disease is metastatic, relapse of a patient's disease or the progression of a patient's disease from any one of the above noted stages to another. A significant event may be any important event used to assess OS, TTP and/or using the RECIST or other response criteria, as determined by one skilled in the art.

As used herein, the terms "subject" and "patient" are used interchangeably. As used herein, the terms "subject" and "subjects" refer to an animal, preferably a mammal including a non-primate (*e.g.*, a cow, pig, horse, donkey, goat, camel, cat, dog, guinea pig, rat, mouse or sheep) and a primate (*e.g.*, a monkey, such as a cynomolgus monkey, gorilla, chimpanzee or a human).

"Targeted therapy" refers to therapeutic treatment that attempts to identify and treat specific cells involved in disease without harming or altering normal cells. Targeted therapeutics may be comprised of, but not limited to, small molecules, such as lapatinib and iressa/gleevec, monoclonal antibodies, such as trastuzumab or nucleic acids, such as siRNAs used to block expression of gene products involved in disease processes. Targeted therapies are useful in the treatment of many disease processes, such as cancer.

As used herein, "time course" shall refer to the amount of time between an initial event and a subsequent event. For example, with respect to a patient's cancer, time course may relate to a patient's disease and may be measured by gauging significant events in the course of the disease, wherein the first event may be diagnosis and the subsequent event may be metastasis, for example.

"Time to progression" or "TTP" refers to a time as measured from the start of the treatment to progression or a cancer or censor. Censoring may come from a study end or from a change in treatment. Time to progression can also be represented as a probability as, for example, in a Kaplan-Meier plot where time to progression may represent the probability of being progression free over a particular time, that time being the time between the start of the treatment to progression or censor.

"Treatment," and other forms of this word refer to the administration of an agent to impede a disease, such as the growth of a cancer, to cause a cancer to shrink by weight or

volume, to extend the expected survival time of the subject and/or time to progression of the tumor or the like. Treatment may also refer to any course which one skilled, for example, a treating physician, deems expedient.

The term "VERATAG®" refers to single and multiplexed and multi-label assays, materials, methods and techniques for performing and utilizing such assays, including but not limited to reagents, analytical procedures and software related to those assays. The terms VERATAG®, vTag and ETAG® shall be used interchangeably.

In a first aspect, the invention is drawn to a method of measuring and/or quantifying the presence and/or amount of Her-3 and/or Her-3 in a complex in a sample, the method comprising providing a sample and determining the presence and/or quantity of Her-3 and/or Her-3 in a complex in the sample. In a preferred embodiment, the sample is a biological sample. In a preferred embodiment, the sample is a tissue sample. In a preferred embodiment, the sample is a fresh tissue sample, a fixed sample, a frozen sample or a lysate. In a preferred embodiment, the sample is a tumor sample. In a preferred embodiment, the sample is a frozen tumor tissue sample. In a preferred embodiment, the sample comprises a tumor lysate from a fresh or frozen tumor sample. In a preferred embodiment, the sample is an FFPE sample or solubilized FFPE sample. In a preferred embodiment, the sample comprises a breast cancer sample. In a certain embodiments, the breast cancer is early stage (i.e., adjuvant) breast cancer or metastatic breast cancer. In certain embodiments, the level of Her-2 expression in the breast cancer is high. In certain embodiments, high Her-2 expression is a log10H2T $\geq$ about 1.14-1.25. In certain embodiments, the high Her-2 expression comprises expression that is very high and/or moderately high. In certain embodiments, the very high Her-2 expression is a log10H2T $\geq$ about 1.84 -2.21. Or, other ranges may be used depending upon the patient cohort. In certain embodiments, if the level of Her-3 is high, the patient is less likely or unlikely to respond to the targeted therapy. In certain embodiments, if the level of Her-3 is low, the patient is more likely to respond to the targeted therapy. In certain embodiments, as as described in more detail herein, the therapy is a Her acting agent. In further embodiments, the therapy is at least one of a Her-2 acting agent or a Her-3-targeted agent.

In a preferred embodiment, the sample is a blood, plasma or lymph sample. In a preferred embodiment, the blood or plasma sample contains circulating tumor cells. In a preferred embodiment, the sample comprises cell lines. In a preferred embodiment, the measurement may be quantitative across a wide dynamic range.

In a preferred embodiment, the method comprises mixing the sample with a binding compound and determining the presence and/or quantity of the binding compound bound to Her- 3 and/or Her-3 in a complex. In a preferred embodiment, the binding compound binds specifically to Her-3. In a preferred embodiment, the binding compound comprises an antibody. In a preferred embodiment, the antibody is raised against one of the peptides having SEQ ID NOs:1-8, as set forth in Example 2 and shown in Figure 2A. In a preferred embodiment, the antibody is a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:13, 14 and 15, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:16, 17 and 18, respectively; and/or a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:19, 20 and 21, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:22, 23 and 24, respectively, as shown in Table 1B. In a preferred embodiment, the antibody is the antibody with the amino acid sequence having SEQ ID NOs:9 and 11 as set forth in Table 1A for the light and heavy chains, respectively, and/or SEQ ID NOs:10 and 12 as set forth in Table 1B for the light and heavy chains, respectively.

TABLE 1A

Her3 light chain sequences:

>Her3.B9A11.H1 _LC (clone 6-6)

SEQ ID NO:9
MDSQAQVLILLLLWVSGTCGDIVMSQSPSSLAVSAGEKVTLSCKSSQSLLNSRTRKNYLAWYQQKPGQS
PKLLIYWASTRESGVPD RFTGSGSGTDFTLTVSSVQAEDLAVYYCKQSYNLWTFGGGTKLEIK

>Her3.F9B10.3_LC (clone 7-3)

SEQ ID NO:10
MRCLAEFLGLLVLWIPGAIGDIVMTQGAPSVPVTPGESVSISCRSSKSLLQNNGNTYLYWFLQRPGQSP
QLLIYRMSNLASGVPDRFSGSGSGTAFTLRISRVEAEDVGVYYCMQHLEYPLTFGAGTKLGLK

Her3 heavy chain sequences:

>Her3.B9A11.H1_HC (clone 1-14)

SEQ ID NO:11
MECNWILPFILSVTSGVYSEVQLQQPGTVLARPGASVRMSCKASGYTFTSYWMHWVKQRPGQGLEWIGA
IYPGNSDTRDNQKFKGKAELTAVTSASTAYMELSSLTNEDSAVYYCTSYYFDGAGYFDFWGQGTTLTVS
S

>Her3.F9B10.3_HC (clone 2-1)

SEQ ID NO:12
MEWSWVFLFLLSVIASVQSQVQLQQSGAEVVRPGASVTLSCKASAYTFTDYELHWMRQTPVHGLEWI
GASDPETGGSAYNQKFKGKAILTADKSSSTAFMELRSLTSEDSAVYFCTRRIFYFGSRGDFFDYWGQGTS
LTVSS

TABLE 1B

Complementarity-Determining Regions (CDRs)				
	Her3 . B9A11 . H1_LC	Her3.B9A11.H1_HC	Her3.F9B10.3_LC	Her3.F9B10.3_HC
CDR1	KSSQSLLNSRTRKNYLA SEQ ID NO:13	SYWMH SEQ ID NO: 16	RSSKSLLQNNGNTYLY SEQ ID NO: 19	DYELH SEQ ID NO: 22
CDR2	WASTRES SEQ ID NO: 14	AIYPGNSDTRDNQKFKG SEQ ID NO: 17	RMSNLAS SEQ ID NO: 20	ASDPETGGSAYNQKFKG SEQ ID NO: 23
CDR3	KQSYNLWT SEQ ID NO: 15	YYFDGAGYFDF SEQ ID NO: 18	MQHLEYPLT SEQ ID NO: 21	RIFYFGSRGDFFDY SEQ ID NO: 24

TABLE 1A and 1B. The amino acid sequences of the light and heavy chains of two antibodies that bind Her3, Her3.B9A11.H1 and Her3.F9B10.3, are shown in Table 1A. The clonal isolates from which the sequences were derived are shown and the complementarity-determining regions (CDRs) are underlined. The light and heavy chains are denoted by "_LC" or "_HC", respectively. Table 1B shows the three CDRs for the B9A11.H1 light and heavy chains and the F9B10.3 light and heavy chains, respectively.

In a preferred embodiment, the method comprises mixing (i) a sample that may contain Her-3 and/or Her-3 in a complex; (ii) a proximity probe that is capable of binding Her-3 and/or at least one other analyte, the proximity probe having an effective proximity and (iii) at least one binding compound, the at least one binding compound being capable of binding Her-3 and having one or more signaling molecules attached, wherein binding of the proximity probe and the binding compound within the effective proximity produces a signal from the molecular tags that correlates with the presence and/or quantity of Her-3 and/or Her-3 in a complex. In a preferred embodiment, the proximity probe and/or binding compound is capable of binding specifically to Her-3. In a preferred embodiment, the proximity probe and/or binding compound further comprises an antibody and each antibody may bind to a specific epitope on Her-3. In a preferred embodiment, the antibody is raised against one of the peptides having SEQ ID NOs: 1- 8, as set forth in Example 2 and shown in Figure 2A. In a preferred embodiment, the antibody is a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:13, 14 and 15, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ

ID NOs:16, 17 and 18, respectively; and/or a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:19, 20 and 21, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:22, 23 and 24, respectively (Table 1B). In a preferred embodiment, the antibody is the antibody with the amino acid sequence having SEQ ID NOs:9 and 11 as set forth in Table 1A for the light and heavy chains, respectively, and/or SEQ ID NOs:10 and 12 as set forth in Table 1A for the light and heavy chains, respectively. In a preferred embodiment, the sample is a biological sample. In a preferred embodiment, the sample is a tissue sample. In a preferred embodiment, the sample is a fixed sample, a frozen sample or a lysate. In a preferred embodiment, the sample is a tumor sample. In a preferred embodiment, the sample is a frozen tumor tissue sample. In a preferred embodiment, the sample comprises a tumor lysate. In a preferred embodiment, the sample comprises a breast cancer sample as discussed herein. For example, in a certain embodiments, the breast cancer is early stage (i.e., adjuvant) breast cancer or metastatic breast cancer. In certain embodiments, the level of Her-2 expression in the breast cancer is high. In certain embodiments, high Her-2 expression is a log10H2T $\geq$ about 1.14-1.25. In certain embodiments, the high Her-2 expression comprises expression that is very high and/or moderately high. In certain embodiments, the very high Her-2 expression is a log10H2T $\geq$ about 1.84 -2.21. Or, other ranges may be used depending upon the patient cohort.

In a preferred embodiment, the sample is an FFPE sample or solubilized FFPE sample. In a preferred embodiment, the sample is a blood, plasma or lymph sample. In a preferred embodiment, the blood or plasma sample contains circulating tumor cells. In a preferred embodiment, the sample contains exosomes and/or other vesicles. In a preferred embodiment, the sample comprises cell lines. In a preferred embodiment, the measurement may be quantitative across a wide dynamic range. In a preferred embodiment, the wide dynamic range is approximately 2 logs. In a more preferred embodiment, the wide dynamic range is about 1-1.5 logs in breast cancer samples. In a preferred embodiment, the method provides a quantitative continuum of Her-3 expression. In a preferred embodiment, the measurement or quantity is sensitive to at least about 1000 receptors per cell to about 200,000 receptors per cell as determined by accuracy studies utilizing well-characterized cell line models and cross-validating technologies such as ELISA and flow cytometry. In a preferred embodiment, the measurement or quantity is sensitive to at least about 5000 receptors per cell to about 200,000 receptors per cell. In a

preferred embodiment, the measurement or quantity is sensitive to at least about 10,000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement or quantity is sensitive to at least about 25,000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement is specific as determined using isotype control antibodies and comparison with conventional IHC methods.

In a preferred embodiment, determining the presence and/or quantity of binding compound bound to Her-3 further comprises providing a second binding compound, the second binding compound being able to specifically bind the binding compound bound to Her-3 and determining the presence and/or quantity of the second binding compound as correlative of the presence and/or quantity of the binding compound bound to Her-3. In a preferred embodiment, the second binding compound is an antibody.

The use of a second binding compound that is capable of specifically binding the first binding compound and has one or more molecular tags may have practical advantages. For example, multiple Her-3-specific first binding compounds may be tested using a single second binding compound to which is attached one or more molecular tags, abrogating the need for attaching molecular tags to each of the multiple Her-3-specific first binding compounds. In a preferred embodiment, the first binding compound is a mouse antibody and the second binding compound is an anti-mouse antibody raised in a non-mouse species (*e.g.*, goat anti-mouse antibodies) to which cleavable molecular tags have been attached.

Second binding compounds are typically labeled with probes useful for detection. Detection systems commonly used for detecting second binding compounds include but are not limited to cleavable molecular tags, as described herein; radiolabels (*i.e.*, radioisotopes such as I-125); enzymes that convert a chemical into a measurable colorimetric, fluorescent or electrochemical signal (*e.g.*, peroxidases) and fluorescent proteins (*e.g.*, green fluorescent protein and its many derivatives).

The antibody can be monoclonal, polyclonal or recombinant and can be prepared by techniques that are well known in the art. Antibodies may include a complete immunoglobulin or fragment thereof, which immunoglobulins include the various classes and isotypes, such as IgA, IgD, IgE, IgG1, IgG2a, IgG2b and IgG3, IgM, etc. Fragments thereof may include Fab, Fv and F(ab')2, Fab' and the like. Antibodies may also be single-chain antibodies, chimeric antibodies, humanized antibodies or any other antibody derivative known to one of skill in the art that retains binding activity that is specific for a particular binding site. In addition, aggregates, polymers and conjugates of

immunoglobulins or their fragments can be used where appropriate so long as binding affinity is maintained.

To facilitate the development of methods to measure Her-3 in biological samples, Her-3-specific monoclonal antibodies were created. Mice were immunized against peptides from Her-3 (shown in Figure 2A) and standard methods as set forth further herein and as known to one skilled in the art were used to create hybridomas. Many methods are known for the creation and production of monoclonal antibodies, for example, the hybridoma method as first described by Koehler et al. (1975) *Nature* 256:495-497 or other methods described in the literature (see Goding, JW (1980) *J. Immunol. Methods* 34:285-308; Harlow E and Lane D (1988) in *Antibodies: A Laboratory Manual*, Chapter 6; Kennett RH et al.(1980) *Monoclonal Antibodies*, Plenum Press; Zola H (1987) *Monoclonal Antibodies: A Manual of Techniques*, CRC Press).

In one embodiment, the method of creating hybridomas begins with immunizing a host animal, such as a mouse, to elicit the production of lymphocytes that produce antibodies targeted to the peptide or protein(s) of interest. Lymphocytes may also be immunized *in vitro*. The antigen used may be a peptide, a protein or a cell displaying the antigen on the cell surface. Lymphocytes are collected then fused by chemical (*e.g.,* with PEG) or electrical (*e.g.,* by electrofusion) methods with myeloma cells to form hybridoma cells, typically under conditions that prevent the growth and/or survival of the parent myeloma cells. Fused cells are allowed to grow because they contain enzymes that facilitate survival in the culture medium. In a preferred embodiment, the culture medium contains hypoxanthine, aminopterin and thymidine (HAT medium), which prevents the growth of cells lacking hypoxanthine quinine phosphoribosyl transferase (HPRT). The HPRT is supplied to the fused cell by the lymphocyte partner, allowing survival of the hybridoma but preventing survival of the parent myeloma cells, which lack HPRT.

Culture media in which hybridomas are grown (*i.e.,* conditioned media) are typically assayed for the production of monoclonal antibodies directed against the antigen using a variety of techniques (see Voller, *et al.* (1978) *J. Clin. Pathol.* 31:507-520), including but not limited to, immunoprecipitation or an *in vitro* binding assay such as enzyme-linked immunosorbant assay (ELISA; *see* Engvall E (1977) in *Biomedical Applications of Immobilized Enzymes and Proteins*, edited by TMS Chang, 2:87-96, Plenum Press), radioimmunoassay (RIA; *see* Sonksen PH (1974) *Brit. Med. Bull.* 30:1-103), Western blots or flow cytometry. Conditioned media from the hybridomas were profiled in a series of assays including ELISA (Figure 1), Western blot (Figure 2) and flow cytometry (Figure 3).

In preferred embodiments, studies using both native and permeabilized and fixed cells are performed to identify antibodies that may perform well in applications that use fixed cells or tissues, such as immunohistochemistry (IHC). Clones of interest may be subcloned by limiting dilution or single cell flow cytometry.

As will be known to those skilled in the art, monoclonal antibodies secreted by hybridoma clones (or subclones) can be purified using conventional purification procedures such as, but not limited to, dialysis, affinity chromatography, gel electrophoresis or protein A-sepharose (or protein L-agarose) chromatography.

One antibody generated, B9A11 (i.e., SEQ ID NOs: 9 and 11), which was raised against the peptide CFDNPDYWHSRLFPKANA (SEQ ID NO: 6) from Her-3, was chosen for use in the Her3- VERATAG® assay experiments described herein. The hybridoma cell lines that produce antibodies B9A11 and F9B10 were deposited with the American Type Culture Collection, 10801 University Boulevard, Manassas, Va. 20110, on January 12, 2010, under the terms of the Budapest Treaty and were accorded the ATCC accession numbers PTA-10574 and PTA-10575.

Many methods and reagents are commonly used to prepare biological samples for analysis. Several methods are outlined or referenced herein and many others are known to those skilled in the art. Samples containing Her-3 suitable for use as biomarkers may come from a wide variety of sources, including cell cultures, animal or plant tissues, patient biopsies, blood or the like. Preferably, samples are human patient samples. Samples are prepared for assays of the invention using conventional techniques, which may depend on the source from which a sample is taken. For biopsies and medical specimens, guidance is provided in the following references: Bancroft JD & Stevens A, eds. 1977, *Theory and Practice of Histological Techniques*, Churchill Livingstone, Edinburgh,; Pearse, 1980, *Histochemistry. Theory and applied. 4th ed.*, Churchill Livingstone, Edinburgh.

Examples of patient tissue samples that may be used include, but are not limited to tissues of breast, prostate, ovary, colon, lung, endometrium, stomach, salivary gland or pancreas. The tissue sample can be obtained by a variety of procedures including surgical excision, aspiration or biopsy. The tissue may be fresh or frozen. In one embodiment, the biological sample may be cells cultured *in vitro* and collected by centrifugation as a cell pellet. In one embodiment, the samples may be patient blood samples or specific blood cell types or subsets of blood cell types (*e.g.*, buffy coats). In one embodiment, the biological sample may be exosomes or samples containing exosomes. Exosomes are small (30-200 nm) vesicles that can be secreted by most cell types, including tumor cells (*see* Mignot *et al* (2006) *J. Cell. Mol. Med.* 10:376- 3 88), *in vivo* and *in vitro*. Tumor-derived exosomes are

thought to play a role in the ability of tumors to evade the immune system and have potential for both diagnostic and therapeutic applications (*see* Taylor and Black (1985) *J. Natl. Cancer Inst.* 74:859-867) and are therefore biological samples of interest.

In a preferred embodiment, the sample is a tumor sample. Examples of types of tumor samples include cancers such as, without limitation, carcinomas, sarcomas, myelomas, leukemias, lymphomas and mixed type cancers. In one embodiment, the cancer is a bone cancer, for example, Ewing's sarcoma, osteosarcoma and rhabdomyosarcoma and other soft-tissue sarcomas. In another embodiment, the cancer is a brain tumor, for example, oligodendroglioma, ependymoma, menengioma, lymphoma, schwannoma or medulloblastoma. In another embodiment, the cancer is breast cancer. In another embodiment, the cancer is an endocrine system cancer, for example, adrenal, pancreatic, parathyroid, pituitary and thyroid cancers. In another embodiment, the cancer is a gastrointestinal cancer, for example, anal, colorectal, esophageal, gall bladder, gastric, liver and small intestine cancers. In another embodiment, the cancer is a gynecological cancer, for example, cervical, endometrial, uterine, fallopian tube, gestational trophoblastic disease, choriocarcinoma, ovarian, vaginal or vulvar cancer. In another embodiment, the cancer is a head and neck cancer, for example, laryngeal, oropharyngeal, parathryroid or thyroid cancer. In another embodiment, the cancer is melanoma, squamous cell carcinoma or basal cell carcinoma. In another embodiment, the cancer is a leukemic cancer, for example, acute lymphocytic leukemia, acute myelogenous leukemia, chronic lymphocytic leukemia, chronic myelogenous leukemia, hairy cell leukemia or a myeloproliferative disorder. In another embodiment, the cancer is a lung cancer, for example, a mesothelioma or non-small cell lung cancer. In another embodiment, the cancer is a lymphoma, such as cutaneous T cell lymphoma, Hodgkin's disease or non-Hodgkin's disease. In another embodiment, the cancer is metastatic cancer. In another embodiment, the cancer is a myeloma, for example, a multiple myeloma. In another embodiment, the cancer is penile cancer. In another embodiment, the cancer is prostate cancer. In another embodiment, the cancer is testicular cancer. In another embodiment, the cancer is thyroid cancer, for example, papillary, follicular, medullary or anaplastic or undifferentiated thyroid carcinoma. In another embodiment, the cancer is a urinary tract cancer, for example, bladder, kidney or urethral cancer.

Methods for preparing cells cultured *in vitro* as fresh, frozen or fixed samples are known to those with skill in the art and exemplary methods are described herein. In one embodiment, assays of the invention are carried out on tissue samples that have been fixed and

embedded in paraffin and a step of deparaffination may be carried out. A tissue sample may be fixed (*i.e.*, preserved) by conventional methodology. *See, e.g.*, Lee G. Luna, HT (ASCP) Ed., 1960, *Manual of Histological Staining Method of the Armed Forces Institute of Pathology* 3rd edition, The Blakston Division McGraw-Hill Book Company, New York; Ulreka V. Mikel, Ed., 1994, *The Armed Forces Institute of Pathology Advanced Laboratory Methods in Histology and Pathology*, Armed Forces Institute of Pathology, American Registry of Pathology, Washington, D.C. One of skill in the art will appreciate that the choice of a fixative is determined by the purpose for which the tissue is to be histologically stained or otherwise analyzed. One of skill in the art will also appreciate that the length of fixation depends upon the size of the tissue sample and the fixative used.

Generally, a tissue sample is first fixed and is then dehydrated through an ascending series of alcohols, infiltrated and embedded with paraffin or other sectioning media so that the tissue sample may be sectioned. Alternatively, one may section the tissue and fix the sections obtained. By way of example, the tissue sample may be embedded and processed in paraffin by conventional methodology according to conventional techniques or as described herein. Once the tissue sample is embedded, the sample may be sectioned by a microtome according to conventional techniques. Sections may have a thickness in a range from about three microns to about twelve microns, and preferably, a thickness from about 5 microns to about 10 microns. In one embodiment, a section may have an area from about 10 mm^2 to about 1 cm^2. Once cut, the sections may be attached to slides by several standard methods. Examples of slide adhesives include, but are not limited to, silane, gelatin and poly-L-lysine. Paraffin-embedded sections may be attached to positively charged slides and/or slides coated with poly-L-lysine.

If paraffin has been used as the embedding material, the tissue sections are generally deparaffinized and rehydrated prior to detection of biomarkers. Tissue sections may be deparaffinized by several conventional standard methodologies. For example, xylenes and a gradually descending series of alcohols may be used according to conventional techniques described by the references provided herein. Alternatively, commercially available deparaffinizing non-organic agents such as Hemo-De® (CMS, Houston, Tex.) may be used.

Cell lysates of mammalian tissue culture cells or fresh or frozen tissues may be prepared by conventional cell lysis techniques (*e.g.*, 0.14 M NaCl, 1.5 mM $MgCl_2$, 10 mM Tris-Cl (pH 8.6), 0.5% Nonidet P-40, and protease and/or phosphatase inhibitors as

required). For fresh mammalian tissues, sample preparation may also include a tissue disaggregation step, such as crushing, mincing, grinding or sonication.

Stable cell lines expressing varying levels of Her-3 were generated. Cell lines stably expressing varying levels of a protein of interest are useful in validating new assays, such as the Her-3 VERATAG® assay, with respect to many parameters such as optimal antibody concentrations, accuracy, sensitivity, reproducibility, precision, linearity, specificity and dynamic range. HEK 293 cells were used to create the stable Her-3-expressing cell lines. HEK 293 cells are a specific cell line originally derived from human embryonic kidney cells transformed with adenovirus DNA (*see* Graham et al. (1977) *J. Gen. Virol.* **36**: 59–74). HEK 293 cells are easy to grow in culture, transfect readily and have been widely used in cell biology research as well as in protein production for the biotechnology industry for many years. The generation of the Her-3- expressing cell lines is described in Example 1 and the results of ELISA assays to determine the level of Her-3 in each of these cell lines is shown in Figure 1.

In a further preferred embodiment, the proximity probe comprises an antibody and a first nucleic acid and the binding compound comprises an antibody and a second nucleic acid, wherein the first and the second nucleic acids are complementary to each other and able to hybridize to determine the effective proximity and produce the signal, directly or indirectly, through hybridization. In a preferred embodiment, the proximity probe and/or binding compound is capable of binding specifically to Her-3. In a preferred embodiment, the binding compound and/or the proximity probe further comprises an antibody and each antibody binds to a different epitope on Her-3. In a preferred embodiment, the antibody is raised against one of the peptides having SEQ ID NOs: 1-8, as set forth in Example 2 and shown in Figure 2A. In a preferred embodiment, the antibody is a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:13, 14 and 15, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:16, 17 and 18, respectively; and/or a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:19, 20 and 21, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:22, 23 and 24, respectively (Table 1B). In a preferred embodiment, the antibody is the antibody with the amino acid sequence having SEQ ID NOs:9 and 11 as set forth in Table 1A for the light and heavy chains, respectively, and/or SEQ ID NOs:10 and 12 as set forth in Table 1A for the light

and heavy chains, respectively. In a preferred embodiment, the sample is a biological sample. In a preferred embodiment, the sample is a tissue sample. In a preferred embodiment, the sample is a fixed sample, a frozen sample or a lysate. In a preferred embodiment, the sample is a tumor sample. In a preferred embodiment, the sample is a frozen tumor tissue sample. In a preferred embodiment, the sample comprises a tumor lysate. In a preferred embodiment, the sample comprises a breast cancer sample as described herein. In a preferred embodiment, the sample is an FFPE sample or solubilized FFPE sample. In a preferred embodiment, the sample is a blood, plasma or lymph sample. In a preferred embodiment, the blood or plasma sample contains circulating tumor cells. In a preferred embodiment, the sample contains exosomes and/or other vesicles. In a preferred embodiment, the sample comprises cell lines. In a preferred embodiment, the measurement may be quantitative across a wide dynamic range. In a preferred embodiment, the wide dynamic range is about 2 logs. In a more preferred embodiment, the wide dynamic range is about 1-1.5 logs in breast cancer samples. In a preferred embodiment, the method provides a quantitative continuum of Her-3 expression. In a preferred embodiment, the measurement or quantity is sensitive to at least about 1000 receptors per cell to about 200,000 receptors per cell as determined by accuracy studies utilizing well-characterized cell line models and cross-validating technologies such as ELISA and flow cytometry. In a preferred embodiment, the measurement or quantity is sensitive to at least about 5000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement or quantity is sensitive to at least about 10,000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement or quantity is sensitive to at least about 25,000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement is specific as determined using isotype control antibodies and comparison with conventional IHC methods. Examples of proximity probes and binding compounds, as set forth herein, can be found, for example, in United States patent applications 7,306,904; 7,320,860 and 7,351,528, each of which is incorporated by reference herein, including any drawings.

Proximity assays are increasingly useful for the understanding of the biological role of molecular complexes, as well as in the study of biomarkers. For example, binding compounds that specifically bind Her-3 or Her-3 in a complex can be coupled with many different detection systems to measure the presence and/or quantity of Her-3 or Her-3 in a complex. Any method known to one of skill in the art to be useful for determining an amount of Her-3 or Her-3 in a complex can be used in accordance with the present invention.

Such methods include but are not limited to Foerster resonance energy transfer (FRET), bioluminescence resonance energy transfer (BRET), biomolecular fluoresence complementation, proximity ligation assay (PLA), scintillation proximity assasy (SPA) and rolling circle amplification (RCA) or any other method for detecting nucleic acid duplexes formed by the proximity of a binding probe and a proximity probe with complementary strands of nucleic acids.

In conducting the methods of the invention, a combination of the assay components is made, including the sample being tested, the binding compounds and optionally the proximity probe. Generally, assay components may be combined in any order. In certain applications, however, the order of addition may be relevant. For example, one may wish to monitor competitive binding, such as in a quantitative assay. Or one may wish to monitor the stability of an assembled complex. In such applications, reactions may be assembled in stages.

The amounts of each reagent can generally be determined empirically. The amount of sample used in an assay will be determined by the predicted number of target complexes present and the means of separation and detection used to monitor the signal of the assay. In general, the amounts of the binding compounds and the proximity probe can be provided in molar excess relative to the expected amount of the target molecules in the sample, generally at a molar excess of at least about 1.5, more desirably about 10-fold excess, or more. In specific applications, the concentration used may be higher or lower, depending on the affinity of the binding compound or proximity probe and the expected number of target molecules present on a single cell.

The assay mixture can be combined and incubated under conditions that provide for binding of the probes to the cell surface molecules, usually in an aqueous medium, generally at a physiological pH (comparable to the pH at which the cells are cultured), maintained by a buffer at a concentration in the range of about 10 to 200 mM. Conventional buffers may be used, as well as other conventional additives as necessary, such as salts, growth medium, stabilizers, etc. Physiological and constant temperatures are normally employed. Incubation temperatures normally range from about 4° to 70°C, usually from about 15° to 45°C, more usually about 25° to 37°C.

In a preferred embodiment, the proximity probe comprises a cleaving probe that has a cleavage-inducing moiety and the at least one binding compound has one or more molecular tags attached to the binding compound by a cleavable linkage, wherein the cleavable linkage may be cleaved within the effective proximity, producing a signal that

correlates with the presence and/or quantity of Her-3. In a preferred embodiment, the cleaving probe and/or binding compound is capable of binding specifically to Her-3. In a preferred embodiment, the binding compound and/or the proximity probe further comprises an antibody and each antibody binds to a different epitope on Her-3. In a preferred embodiment, the antibody is raised against one of the peptides having SEQ ID NOs: 1-8, as set forth in Example 2 and shown in Figure 2A. In a preferred embodiment, the antibody is a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:13, 14 and 15, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:16, 17 and 18, respectively; and/or a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:19, 20 and 21, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:22, 23 and 24, respectively. In a preferred embodiment, the antibody is the antibody with the amino acid sequence having SEQ ID NOs:9 and 11 as set forth in Table 1A for the light and heavy chains, respectively, and/or SEQ ID NOs:10 and 12 as set forth in Table 1A for the light and heavy chains, respectively. In a preferred embodiment, the sample is a biological sample. In a preferred embodiment, the sample is a tissue sample. In a preferred embodiment, the sample is a fixed sample, a frozen sample or a lysate. In a preferred embodiment, the sample is a tumor sample. In a preferred embodiment, the sample is a frozen tumor tissue sample. In a preferred embodiment, the sample comprises a tumor lysate. In a preferred embodiment, the sample comprises a breast cancer sample as described herein. In a preferred embodiment, the sample is an FFPE sample or solubilized FFPE sample. In a preferred embodiment, the sample is a blood, plasma or lymph sample. In a preferred embodiment, the blood or plasma sample contains circulating tumor cells. In a preferred embodiment, the sample comprises cell lines. In a preferred embodiment, the measurement may be quantitative across a wide dynamic range. In a preferred embodiment, the wide dynamic range is about 2 logs. In a more preferred embodiment, the wide dynamic range is about 1-1.5 logs in breast cancer samples. In a preferred embodiment, the method provides a quantitative continuum of Her-3 expression. In a preferred embodiment, the measurement or quantity is sensitive to at least about 1000 receptors per cell to about 200,000 receptors per cell as determined by accuracy studies utilizing well-characterized cell line models and cross- validating technologies such as ELISA and flow cytometry. In a preferred embodiment, the measurement or quantity is sensitive to at least about 5000 receptors per cell

to about 200,000 receptors per cell. In a preferred embodiment, the measurement or quantity is sensitive to at least about 10,000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement or quantity is sensitive to at least about 25,000 receptors per cell to about 200,000 receptors per cell. In a preferred embodiment, the measurement is specific as determined using isotype control antibodies and comparison with conventional IHC methods.

A two antibody proximity assay was optimized. Ab-6 (LabVision), a Her-3-specific monoclonal antibody with epitope specificity to the cytoplasmic terminus of Her-3, was conjugated to the VERATAG® reporter (Pro 11) for use as the proximity probe (Ab-6-Pro11). The proprietary monoclonal antibody, B9A11, was conjugated to biotin for use as the cleaving probe (B9A11-biotin) when complexed with streptavidin-methylene blue ("molecular scissors"). The assay methods are described in Example 5. Several cell lines that expressed varying levels of Her-3 ranging from very high levels (in one of the stably transfected HEK 293 cell lines, called 293H3-Clone 1) to moderate to low levels (MDA-MB-468 and MDA-MB -453, respectively) to low to no detectable Her-3 (in SKOV3 cells) were used in the optimization studies. The optimization process included determining the optimal antibody concentration for maximizing dynamic range (see Example 7 and Figure 6), determining the accuracy of the assay (see Example 8 and Figure 7), testing the sensitivity, reproducibility, and precision of the assay (see Example 9/Figure 8, Example 10/Figure 9 and Example 11/Figure 10, respectively), the linearity of the assay in different sample sizes (see Example 12 and Figure 11) and the specificity of the assay by testing for non-specific binding using isotype controls (see Example 13 and Figure 12).

Isotype controls are typically performed to eliminate the possibility that the binding results are due to the particular isotype of the antibody rather than the individual antibody. Additionally, one skilled in the art will appreciate that any signal "noise" seen in the isotype controls can be subtracted from the total signal, potentially yielding a more refined result. When an isotype control experiment was performed, the contribution of non-specific binding was observed to be very low (see Figure 12).

Many advantages are provided by measuring Her-3 or the Her-3 in a complex using releasable molecular tags, including separation of released molecular tags from an assay mixture providing greatly reduced background and a significant gain in sensitivity and separation and detection providing a convenient multiplexing capability so that multiple receptor complex components may be readily measured simultaneously in the same assay. Assays employing such tags can have a variety of forms and are disclosed in the following references: U.S.

Patent Numbers 7,105,308; 6,627,400; 7,402,397; 7,402,398 and 7,402,399, as well as International Patent Publication No. WO 2004/011900, each of which is incorporated herein by reference in its entirety. A wide variety of separation techniques may be employed that can distinguish molecules based on one or more physical, chemical or optical differences among molecules being separated including electrophoretic mobility, molecular weight, shape, solubility, pKa, hydrophobicity, charge, charge/mass ratio or polarity. In one embodiment, molecular tags in a plurality or set differ in electrophoretic mobility and optical detection characteristics and are separated by electrophoresis. In another embodiment, molecular tags in a plurality or set may differ in molecular weight, shape, solubility, pKa, hydrophobicity, charge, polarity and are separated by normal phase or reverse phase HPLC, ion exchange HPLC, capillary electrochromatography, mass spectroscopy or gas phase chromatography.

Sets of molecular tags are provided that can be separated into distinct bands or peaks by a separation technique after they are released from binding compounds. Identification and quantification of such peaks provides a measure or profile of the presence and/or amounts of Her-3. Molecular tags within a set may be chemically diverse; however, for convenience, sets of molecular tags are usually chemically related. For example, they may all be peptides or they may consist of different combinations of the same basic building blocks or monomers or they may be synthesized using the same basic scaffold with different substituent groups for imparting different separation characteristics. The number of molecular tags in a plurality may vary depending on several factors including the mode of separation employed, the labels used on the molecular tags for detection, the sensitivity of the binding moieties and the efficiency with which the cleavable linkages are cleaved.

Measurements made directly on tissue samples may be normalized by including measurements on cellular or tissue targets that are representative of the total cell number in the sample and/or the numbers of particular subtypes of cells in the sample (*see*, for example, United States Patent Application Publication No. US 2009/0191559 which is incorporated by reference herein in its entirety, including any drawings). The additional measurement may be preferred, or even necessary, because of the cellular and tissue heterogeneity in patient samples, particularly tumor samples, which may comprise substantial fractions of normal cells.

In one embodiment, a binding compound can be represented by the following formula:

$$B\text{-}(L\text{-}E)_k$$

wherein B is binding moiety; L is a cleavable linkage and E is a molecular tag. In homogeneous assays, cleavable linkage, L, may be an oxidation-labile linkage, and more preferably, it is a linkage that may be cleaved by singlet oxygen. The moiety "-(L-E)$_k$" indicates that a single binding compound may have multiple molecular tags attached via cleavable linkages. In one aspect, k is an integer greater than or equal to one, but in other embodiments, k may be greater than several hundred, *e.g.* 100 to 500 or k is greater than several hundred to as many as several thousand, *e.g.* 500 to 5000. Usually each of the plurality of different types of binding compounds has a different molecular tag, E. Cleavable linkages, *e.g.* oxidation-labile linkages, and molecular tags, E, are attached to B by way of conventional chemistries.

Preferably, B is an antibody that specifically binds to a target, such as Her-3. Antibodies specific for Her-3 epitopes are provided in the examples set forth herein. Antibody compositions may be readily formed from a wide variety of commercially available antibodies, either monoclonal or polyclonal or by methods disclosed herein.

Cleavable linkage, L, can be virtually any chemical linking group that may be cleaved under conditions that do not degrade the structure or affect detection characteristics of the released molecular tag, E. Whenever a cleaving probe is used in a homogeneous assay format, cleavable linkage, L, is cleaved by a cleavage agent generated by the cleaving probe that acts over a short distance so that only cleavable linkages within an effective proximity of the proximity probe are cleaved. Typically, such an agent must be activated by making a physical or chemical change to the reaction mixture so that the agent produces a short lived active species that diffuses to a cleavable linkage to affect cleavage.

In a non-homogeneous format, because specifically-bound binding compounds are separated from unbound binding compounds, a wider selection of cleavable linkages and cleavage agents are available for use. Cleavable linkages may not only include linkages that are labile to reaction with a locally acting reactive species, such as hydrogen peroxide, singlet oxygen or the like, but also linkages that are labile to agents that operate throughout a reaction mixture, such as base-labile linkages, photocleavable linkages, linkages cleavable by reduction, linkages cleaved by oxidation, acid-labile linkages and peptide linkages cleavable by specific proteases. References describing many such linkages include Greene and Wuts, 1991, *Protective Groups in Organic Synthesis, Second Edition*, John Wiley & Sons, New York; Hermanson,1996, *Bioconjugate Techniques*, Academic Press, New York; and U.S. Patent No. 5,565,324, each of which is incorporated by reference herein.

Molecular tag, E, in the present invention may comprise an electrophoric tag as described in the following references when separation of pluralities of molecular tags are carried out by gas chromatography or mass spectrometry: *See, e.g.*, Zhang *et al.*, 2002, *Bioconjugate Chem.* 13:1002-1012; Giese, 1983, *Anal. Chem.* 2:165-168; and U.S. Patent Nos. 4,650,750; 5,360,819; 5,516,931; and 5,602,273, each of which is hereby incorporated by reference in its entirety.

Molecular tag, E, is preferably a water-soluble organic compound that is stable with respect to the active species, especially singlet oxygen, and that includes a detection or reporter group. Otherwise, E may vary widely in size and structure. In one embodiment, E has a molecular weight in the range of from about 50 to about 2500 daltons, more preferably, from about 50 to about 1500 daltons. E may comprise a detection group for generating an electrochemical, fluorescent or chromogenic signal. In embodiments employing detection by mass, E may not have a separate moiety for detection purposes. Preferably, the detection group generates a fluorescent signal.

Molecular tags within a plurality are selected so that each has a unique separation characteristic and/or a unique optical property with respect to the other members of the same plurality. In one embodiment, the chromatographic or electrophoretic separation characteristic is retention time under a set of standard separation conditions conventional in the art, *e.g.*, voltage, column pressure, column type, mobile phase or electrophoretic separation medium. In another embodiment, the optical property is a fluorescence property, such as emission spectrum, fluorescence lifetime or fluorescence intensity at a given wavelength or band of wavelengths. Preferably, the fluorescence property is fluorescence intensity. One or two or more of the molecular tags of a plurality may have identical migration or retention times, but they will have unique fluorescent properties, *e.g.* spectrally resolvable emission spectra, so that all the members of the plurality are distinguishable by the combination of molecular separation and fluorescence measurement.

Preferably, released molecular tags are detected by electrophoretic separation and the fluorescence of a detection group. In such embodiments, molecular tags having substantially identical fluorescence properties have different electrophoretic mobilities so that distinct peaks in an electropherogram are formed under separation conditions. Preferably, pluralities of molecular tags of the invention are separated by a conventional capillary electrophoresis apparatus, either in the presence or absence of a conventional sieving matrix. During or after electrophoretic separation, the molecular tags are detected or identified by recording fluorescence signals and migration times (or migration distances) of the

separated compounds or by constructing a chart of relative fluorescent and order of migration of the molecular tags (*e.g.*, as an electropherogram). Preferably, the presence, absence and/or amounts of molecular tags are measured by using one or more standards.

A cleavage-inducing moiety, or cleaving agent, is a group that produces an active species that is capable of cleaving a cleavable linkage, preferably by oxidation. Preferably, the active species is a chemical species that exhibits short-lived activity so that its cleavage-inducing effects are only in the proximity of the site of its generation. Either the active species is inherently short lived, so that it will not create significant background beyond the proximity of its creation, or a scavenger is employed that efficiently scavenges the active species, so that it is not available to react with cleavable linkages beyond a short distance from the site of its generation. Illustrative active species include singlet oxygen, hydrogen peroxide, NADH and hydroxyl radicals, phenoxy radical, superoxide and the like. Illustrative quenchers for active species that cause oxidation include polyenes, carotenoids, vitamin E, vitamin C, amino acid-pyrrole N-conjugates of tyrosine, histidine and glutathione. *See, e.g.* Beutner *et al.*, 2000, *Meth. Enzymol.* 319:226-241.

One consideration in designing assays employing a cleavage-inducing moiety and a cleavable linkage is that they not be so far removed from one another when bound to a receptor complex that the active species generated by the cleavage-inducing moiety cannot efficiently cleave the cleavable linkage. In one embodiment, cleavable linkages preferably are within about 1000 nm and preferably within about 20-200 nm of a bound cleavage-inducing moiety. More preferably, for photosensitizer cleavage-inducing moieties generating singlet oxygen, cleavable linkages are within about 20-100 nm of a photosensitizer in a receptor complex. One of ordinary skill in the art will recognize that the effective proximity of a particular sensitizer may depend on the details of a particular assay design and may be determined or modified by routine experimentation.

A sensitizer is a compound that can be induced to generate a reactive intermediate, or species, usually singlet oxygen. Preferably, a sensitizer used in accordance with the invention is a photosensitizer. Other sensitizers included within the scope of the invention are compounds that on excitation by heat, light, ionizing radiation or chemical activation will release a molecule of singlet oxygen. The best known members of this class of compounds include the endoperoxides such as 1,4-biscarboxyethyl-1,4-naphthalene endoperoxide, 9,10- diphenylanthracene-9, 1 0-endoperoxide and 5,6,11,12-tetraphenyl naphthalene 5,12- endoperoxide. Heating or direct absorption of light by these compounds releases singlet oxygen. Further sensitizers are disclosed by Di Mascio *et al.*, 1994, *FEBS*

Lett. 355:287 and Kanofsky, 1983, *J.Biol. Chem.* 258:5991-5993; Pierlot *et al.*, 2000, *Meth. Enzymol.* 319:3-20.

Photosensitizers may be attached directly or indirectly, via covalent or non-covalent linkages, to the antibodies. Guidance for constructing such compositions are available in the literature, *e.g.* in the fields of photodynamic therapy, immunodiagnostics and the like. Exemplary guidance may be found in Ullman *et al.*, 1994, *Proc. Natl. Acad. Sci. USA* 91, 5426- 5430; Strong *et al.*, 1994, *Ann. New York Acad. Sci.* 745: 297-320; Yarmush *et al.*, 1993, *Crit. Rev. Therapeutic Drug Carrier Syst.* 10: 197-252; and U.S. Patent Nos. 5,709,994, 5,340,716, 6,251,581, and 5,516,636.

A large variety of light sources are available to photo-activate photosensitizers to generate singlet oxygen. Both polychromatic and monochromatic sources may be used as long as the source is sufficiently intense to produce enough singlet oxygen in a practical time duration. The length of the irradiation depends on the nature of the photosensitizer, the nature of the cleavable linkage, the power of the source of irradiation and its distance from the sample. In general, the period for irradiation may be less than about a second to as long as about 3 hours and is usually in the range of 15 minutes to 2 hours. Exemplary light sources include lasers such as, *e.g.*, helium-neon lasers, argon lasers, YAG lasers, He/Cd lasers and ruby lasers; photodiodes; mercury, sodium and xenon vapor lamps and incandescent lamps such as, *e.g.*, tungsten and tungsten/halogen and flashlamps. An exemplary photoactivation device suitable for use in the methods of the invention is disclosed International Patent Publication No. WO 03/051669, which is incorporated by reference herein, including any drawings. In such embodiments, the photoactivation device is an array of light emitting diodes (LEDs) mounted in housing that permits the simultaneous illumination of all the wells in a 96-well plate.

Examples of photosensitizers that may be utilized in the present invention are those that have the above properties and those disclosed by U.S. Patent Nos. 5,536,834, 5,763,602, 5,565,552, 5,709,994, 5,340,716, 5,516,636, 6,251,581 and 6,001,673; published European Patent Application No. 0484027; Martin *et al.*, 1990, *Methods Enzymol.* 186:635-645 and Yarmush *et al.*, 1993, *Crit. Rev. Therapeutic Drug Carrier Syst.* 10:197-252, all of which are incorporated by reference herein, including any drawings. As with sensitizers, in certain embodiments, a photosensitizer may be associated with a solid phase support by being covalently or non-covalently attached to the surface of the support or incorporated into the body of the support. In general, the photosensitizer is associated with the support in an amount necessary to achieve the necessary amount of singlet oxygen.

Generally, the amount of photosensitizer is determined empirically according to routine methods.

Following cleavage, the sample can then be analyzed to determine the identity of molecular tags that have been released. Where an assay employing a plurality of binding compounds is employed, separation of the molecular tags will generally precede their detection. The methods for both separation and detection are determined in the process of designing the molecular tags for the assay. A preferred mode of separation employs electrophoresis, in which the various tags are separated based on known differences in their electrophoretic mobilities.

In a second aspect, the invention is drawn to a method for determining whether a subject with a cancer is likely to respond to treatment with a targeted therapy, for predicting a time course of disease and/or for predicting the probability of a significant event in the time course of the subject's cancer, comprising measuring in a biological sample from the subject's cancer an amount of Her-3, wherein the method is dependent on the level of Her-3.

In certain embodiments, if the level of Her-3 is high, the patient is less likely or unlikely to respond to the targeted therapy. In certain embodiments, if the level of Her-3 is low, the patient is more likely to respond to the targeted therapy. In certain embodiments, as as described in more detail herein, the therapy is a Her acting agent. In further embodiments, the therapy is at least one of a Her-2 acting agent or a Her-3-targeted agent.

In a certain embodiments, the breast cancer is early stage (i.e., adjuvant) breast cancer or metastatic breast cancer. In certain embodiments, the level of Her-2 expression in the breast cancer is high. In certain embodiments, high Her-2 expression is a log10H2T $\geq$ about 1.14-1.25. In certain embodiments, the high Her-2 expression comprises expression that is very high and/or moderately high. In certain embodiments, the very high Her-2 expression is a log10H2T $\geq$ about 1.84 -2.21. Or, other ranges may be used depending upon the patient cohort.

In a preferred embodiment, a time course is measured by determining the time between significant events in the course of a patient's disease, wherein the measurement is predictive of whether a patient has a long time course. In a preferred embodiment, the significant event is the progression from primary diagnosis to death. In a preferred embodiment, the significant event is the progression from primary diagnosis to metastatic disease. In a preferred embodiment, the significant event is the progression

from primary diagnosis to relapse. In a preferred embodiment, the significant event is the progression from surgery to death. In a preferred embodiment, the significant event is the progression from surgery to relapse. In a preferred embodiment, the significant event is from surgery to metastases. In a preferred embodiment, the significant event is the progression from metastatic disease to death. In a preferred embodiment, the significant event is the progression from metastatic disease to relapse. In a preferred embodiment, the significant event is the progression from relapse to death. In a preferred embodiment, the time course is measured with respect to overall survival rate, time to progression and/or using the RECIST or other response criteria.

The Her-3 VERATAG® assay was used to examine the Her-3 levels in a cohort of patients from the International Serum Her-3/neu Study Group trial. These patients (n=105) were selected primarily by IHC for Her-2 positivity performed at a central location by a single pathologist and all received trastuzumab. Only patients with Her-2 over-expressing tumors (>10% of tumor cells IHC 3+ as determined by HercepTest) and/or ErbB2-amplified (with positive FISH testing mandatory on all IHC 2+ cases) metastatic breast cancer were included in this study (see Example 14 for additional details). The Her3 VERATAG® assay was performed on these patient samples and out of 105 samples on which the assay was performed, 85 had measurable Her-3 levels above the limit of detection (8 had no detectable tumor, 1 sample had a fluorescein failure, and 8 samples were below the limit of detection and considered low/negative in the assay). The results are shown in Figure 13.

Her-3 levels and in particular Her-3 levels in Her-2 positive tumors have been implicated in having prognostic value with respect to the time course of disease progression and overall survival as well as response to therapy and particularly with respect to escape from EGFR-family-targeted therapeutics. See Sergina *et al.* (2007) *Nature* 445:437-441, Osipo *et al.* (2007) *Int J Oncol.* 30:509-520, De Alava *et al.* (2007) *Clinical Oncol.* 25:2656-2663, Ma and Bose (2008) *E-Updates* in HER1 and HER2 Targeting in Breast Cancer, Volume 2, Tovey *et al.* (2006) *J. Pathol.* 210:358-362, Menendez and Lupu (2007) *Breast Cancer Res.* 9:111, Fuchs *et al.* (2006) *Anticancer Res.* 26:4397-4402, Lee-Hoeflich *et al.* (2008) *Cancer Res.* 68:5878-5886.

Her-3 expression has been examined in many cancers, including tumors of patients treated with therapeutics targeted to EGFR family members (e.g., trastuzumab, pertuzumab, lapitinib, cetuximab, gefitinib and erlotinib) as well as chemotherapeutics. In comparing Her- family member levels with clinical outcome, data suggest specifically that Her-3 expression and/or the relative amounts of Her-2 and Her-3 may be of use as a prognostic and

predictive diagnostic biomarker in ovarian cancer (*see* Amier *et al.* (2008) *J Clin. Oncol.* 26:abstract 5552, Amier *et al.* (2008) Meeting: *2008 Molecular Markers*, abstract 25 and Xu *et al.* (1999) *Clin. Cancer Res.* 5:3652-3660) and non-small-cell lung cancer (Cappuzzo *et al.* (2005) *Brit. J. Cancer* 93:1334-1340). In a preferred embodiment, the method further comprises determining whether the level of Her-3 is high or low by dividing a sample group of Her-2-positive patients into at least two subgroups comprising one subgroup with a high amount of Her-3 and at least one other subgroup with a low amount of Her-3, wherein if the Her-3 is low, then the patient is likely to respond to Her-2-targeted therapy, the time course of the disease is likely to be long and the patient is not likely to have a significant event. In a preferred embodiment, the subject's cancer is breast cancer, colorectal cancer, ovarian cancer, bladder cancer, prostate cancer, non-small cell lung cancer, melanoma, pharyngeal cancer, pancreatic cancer, esophageal cancer, glioma, bile duct carcinoma, biliary tract carcinoma, cholangiocarcinoma, gastric cancer, endometrial cancer, gall bladder cancer, squamous cell carcinoma or basal cell carcinoma. In a preferred embodiment, the subject's cancer is breast cancer, melanoma, colorectal cancer or ovarian cancer. In a preferred embodiment, the subject's cancer is a Her-2 positive breast cancer. In a preferred embodiment, the breast cancer is early stage (i.e., adjuvant) breast cancer or metastatic breast cancer.

In a preferred embodiment, the targeted therapy is at least one Her family-targeted agent. In a preferred embodiment, the Her family-targeted agent is a multi- or single-targeted agent. In a preferred embodiment, the multi-targeted agent is a dual kinase inhibitor or a bispecific antibody. In a preferred embodiment, the Her family targeted agent is trastuzumab, lapatinib or pertuzumab. In a preferred embodiment, the at least one Her family-targeted agent is at least two agents, wherein the at least two agents are one or more Her-2-targeted monoclonal antibodies and/or EGFR-targeted monoclonal antibodies and/or an EGFR and Her-2 dual kinase inhibitor. In a preferred embodiment, the monoclonal antibody is trastuzumab. In a preferred embodiment the EGFR-targeted monoclonal antibody is cetuximab or panitumumab. In a preferred embodiment, the EGFR-targeted monoclonal antibody is zalutumumab, nimotuzumab, and matuzumab. In a preferred embodiment, the dual kinase inhibitor is lapatinib, erlotinib or gefitinib. In a preferred embodiment, the targeted therapy is a Her-3 or Her-3 signaling pathway acting agent. In a preferred embodiment, the Her-3 or Her-3 signaling pathway targeted agent is a Her-3 monoclonal antibody, a Her-3 dimerization inhibitor, a Her-3 phosphorylation inhibitor and/or an inhibitor of a Her-3 signaling pathway member selected from the group consisting of PI3K, Akt,

mTOR, ERK1/2, or PYK2. In a preferred embodiment, likeliness to respond, likeliness to have a long time course and/or likeliness to have a significant event is measured as an overall survival rate, as time to progression, as disease-free survival, as progression-free survival, and/or as objective tumor response using the RECIST criteria. Signal transduction refers to any process by which cells convert one kind of signal into another. Typically, this involves some type of signal on the cell surface (for example, the binding of a ligand to a cell surface receptor), followed by a cascade of biochemical reactions inside the cell, which are carried out by enzymes, resulting in a signal transduction pathway or signaling pathway, effecting a multitude of cellular functions. In the case of the EGFR family of tyrosine kinase receptors, each of the four receptors in the family has an extracellular domain, comprising both a dimerization domain and a ligand-binding domain, as well as a trans-membrane domain and an intracellular domain with tyrosine kinase activity (*see* Burgess *et al.* (2003) *Mol Cell.*12:541- 542). In Her-3, the kinase domain is not functional but through dimerization with other family members, Her-3 can exert significant signaling pathway effects. Evidence suggests that cooperation of multiple ErbB receptors and ligands is required for initiating cell transformation. When activated, this family of receptors sustains a complex network of signaling pathways. All EGFR family members have been found to be expressed and/or altered in a variety of cancers and may play a significant role in tumor development, including proliferation, apoptosis, and metastasis (*see* Burgess (2008) *Growth Factors* 26:263-274 and Normanno *et al.* (2006) *Gene* 366:2-16). Intense interest in targeting the EGFR family members (*see* Bianco *et al.* (2007) *Int. J. Biochem. Cell. Biol.* 39:1416-1431), particularly EGFR and Her-2, has resulted in several approved targeted therapeutics. Based on promising preclinical data in both *in vitro* and *in vivo* test models, the results in clinical trials have been somewhat disappointing, resulting in increased interest in other family members such as Her-3, as well as downstream signaling pathway members.

Her-3 signaling has been linked to cancer and, in particular, Her-2/Her-3 dimer formation may be crucial for increased aggression in tumors that over-express Her-2, leading to interest in targeted therapeutics that inhibit dimer formation as well as downstream pathways activated by Her-3. Her-3 is particularly adept at signaling because it has 6 binding sites for phosphoinositide 3'-kinases (PI3K) which in turn, activate protein kinase B (also called AKT). The "PI3K/AKT" signaling pathway has been shown to be required for an extremely diverse array of cellular activities - most notably cellular proliferation and survival – fueling the interest in targeting both Her-3 as well as downstream signaling pathway

members to create novel targeted therapeutics or to potentiate the therapeutic value of current EGFR-family targeted therapeutics. (For review, *see* Stern (2008) *J Mammary Gland Biol Neoplasia* 13:215-223, Sithanandam and Anderson (2008) *Cancer Gene Ther.* 15:413-448 and Arkin and Moasser (2008) *Curr. Opin. Investig. Drugs* 9:1264-1276).

In a preferred embodiment, whether the cancer is Her-2 positive is determined by IHC, FISH, CISH, quantitative mRNA, hybridization array or VERATAG®. In a preferred embodiment, determining the level of Her-3 is performed using IHC, FISH, CISH, quantitative mRNA, hybridization array or VERATAG®. In a preferred embodiment, the method further comprises determining whether an amount of Her-3 protein is low by comparing the amount of Her-3 in the subject's cancer to a pre-determined cutoff. In a preferred embodiment, the method further comprises determining the level of Her-3 by dividing a sample group of Her-2-positive patients into at least two subgroups comprising one subgroup with a high amount of Her-3 and at least one other subgroup with a low amount of Her-3, wherein if the Her-3 is high, then the patient is unlikely to respond to Her-2-targeted therapy, the time course of the disease is likely to be short and/or the patient is likely to have a significant event.

In a preferred embodiment, the method further comprises determining that a subject is afflicted with a Her-2 positive cancer that is unlikely to respond to treatment according to a method of the invention, then advising a medical professional of the treatment option of administering to the subject an effective amount of a different therapeutic agent.

In a third aspect, the invention is drawn to a purified antibody that binds to Her-3. In a preferred embodiment, the antibody is a polyclonal antibody or a monoclonal antibody. In a preferred embodiment, the antibody is a monoclonal antibody. In a preferred embodiment, the antibody is raised against one of the peptides having SEQ ID NOs:1-8, as set forth in Example 2 and shown in Figure 2A. In a preferred embodiment, the antibody is a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:13, 14 and 15, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:16, 17 and 18, respectively; and/or a monoclonal antibody comprising (a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs: 19, 20 and 21, respectively, and (b) a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:22, 23 and 24, respectively (Table 1B). In a preferred embodiment, the antibody is the antibody with the amino acid sequence having SEQ ID NOs:9 and 11 as set

forth in Table 1A for the light and heavy chains, respectively, and/or SEQ ID NOs:10 and 12 as set forth in Table 1A for the light and heavy chains, respectively.

In a preferred embodiment, the invention is drawn to the DNA encoding the antibody. The DNA encoding the monoclonal antibodies is isolated and sequenced using techniques commonly known to those skilled in the art of cloning. Once isolated, the DNA can be ligated into expression vectors and transfected into appropriate host cells to obtain recombinant antibodies from cultured cells (*see* Plueckthun (1992) *Immunological Rev.* 130: 151-188).

Those with skill in the art will appreciate that the amino acid sequence of an antibody can be modified and that modifications may be desirable to enhance the properties of the antibody for therapeutic, analytical or diagnostic use. Further it will be appreciated that one or more amino acids in these antibodies may be changed by insertion, deletion or substitution without appreciably diminishing the binding characteristics of the antibody. Exemplary amino acid changes would be substitutions using amino acids with similar molecular characteristics (*i.e.*, conservative substitutions, *e.g.*, changing amino acids from within the following subgroups of aromatic amino acids, acidic amino acids, basic amino acids or amino acids with amides or sulphurs). Other non-conservative substitutions or insertions may be made without appreciably altering molecular integrity or binding characteristics. Further, some amino acid changes or collection of amino acid changes will enhance properties of the antibody, including but not limited to, better binding affinity, greater stability, (*e.g.*, resistance to proteases) selectivity and/or ease of production. Methods for changing amino acid sequences and/or selecting for molecules with better properties are known to those with skill in the art. Preferably, in intact antibodies, the degree of sequence identity after modification is at least 50% and more preferably, at least 75% and most preferably at least 90-95%. Each of these antibodies is intended to be within the scope of the contemplated invention.

In a preferred embodiment, antibodies targeted to Her-3 may be used to develop additional Her-3-targeted molecules. Modifications of the antibodies described herein may be desirable to improve qualities including, but not limited to, increasing effector function, decreasing immunogenicity, increasing stability, improving pharmacologic properties such as serum half-life and aiding in ease and yield of production. Each of these targeted molecules is intended to be within the scope of the contemplated invention.

In a preferred embodiment, humanized antibodies comprising the antigen binding regions of the antibodies described herein (see Table 1A) in a human framework may be used

for therapeutic applications. Several methods for humanizing antibodies have been reported (*see* Jones *et al.* (1986) *Nature* 321:522-525, Riechmann *et al.* (1988) *Nature* 332:323-327 and Verhoeyen *et al.* (1988) *Science* 239:1534-1536). Typically, the non-human sequences of the variable domain are screened computationally against the entire repertoire of human light and heavy chain variable domain sequences to find the human variable framework sequences closest to the rodent sequences (*see* Sims *et al.*(1993) *J. Immunol.* 151:2296-2308, Chothia *et al.* (1987) *J. Mol. Biol.* 186:901-917). Alternatively, consensus frameworks can be used (*see* Carter *et al.* (1992) *Proc. Natl. Acad. Sci. USA* 89:4285-4289 and Presta *et al.* (1993) *J. Immunol.* 151:2623- 2632). In a preferred embodiment, computer-aided design is used to select sequences that confer stability and retain or improve binding characteristics. Each of these is intended to be within the scope of the contemplated invention.

In another embodiment, the antibody CDRs may be used to create targeted binding molecules that bind the same epitope in Her-3 but are contained within a framework that is not a native antibody. For example, one skilled in the art would appreciate that methods are available for creating binding molecules in which the framework may be a portion of an antibody, for example, an scFv or F(ab')$_2$ (*see* WO 93/16185 and Carter *et al.*(1992) *Bio/Technology* 10:163-167, respectively), each of which is incorporated by reference herein. One skilled in the art may also appreciate that a completely unrelated protein (such as a bacterial beta-lactamase) can properly display the binding domain(s) to form a binding compound. In this sense, related antibodies, as defined herein, are intended to be within the scope of the invention.

The antibody may act therapeutically through binding alone or through other properties (e.g., enzymatic activity or toxic warheads). In one embodiment , the targeted protein may be modified to exert a therapeutic effect or a greater therapeutic effect via antigen- dependent cell-mediated cytotoxicity (ADCC) or complement-dependent cytotoxicity (CDC). In another embodiment, toxins may be conjugated to the antibody or targeted protein. Exemplary small molecule toxins include but are not limited to maytansine, calicheamicin and CC-1065 (*see*, e.g., Carter and Senter (2008) *Cancer J.* 14:154-169). Additionally, radiolabels can be linked to antibodies to create targeted therapeutics. Biologic toxins may also be linked to targeted proteins and include, but not be limited to, diphtheria toxin, *Pseudomonas* exotoxin, abrin and ricin (*see* Kreitman (2006) *AAPS J.* 18:E532-551)

In a further embodiment, the targeted antibodies (or fragments thereof) may be fused to enzymes for use in antibody-directed enzyme prodrug therapy (ADEPT; *see* Bagashawe (1987) *Br. J. Cancer* 58:700-703 and Senter *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:4842- 4846). In another embodiment, the antibodies or targeted proteins may be fused to molecules such as polyethylene glycol, that enhance pharmacologic properties, such as serum half-life (*see* Harris and Chess (2003) *Nat. Rev. Drug Discov.* 2:214-221).

Examples

The present invention may be better understood by reference to the following non-limiting examples.

Example 1: Generation of stable cell lines expressing varying levels of HER3 protein

A commercially available cDNA (Origene Technologies, Inc.) for full-length HER3 (NM_001982.2) was digested with the restriction enzymes Not I and Xba I and the resulting fragment was subcloned into pcDNA.3 1 Zeocin selectable expression vector. The resulting plasmid was transformed into bacteria and screened to verify the correct insert. Positive clones were sequence verified, expanded in bacteria and the plasmid purified using the Qiagen Maxi-prep kit. Human embryonic kidney cells (HEK-293) were purchased from the American Type Culture Collection and maintained in DMEM supplemented with 10% FBS, 1x penicillin- streptomycin (100X is 10,000 U/ml penicillin-G and 10,000 ug/ml streptomycin), and Glutamax (GIBCO) at 37C in 5% CO_2. The day prior to transfection, the cells were split to approximately 25-30% confluence and incubated overnight in media without pen-strep. The cells were then transfected with Fugene HD (Roche) according to the manufacturer's instructions. The next day the media was replaced with fresh complete media and the cells were incubated for 48 hours prior to the addition of 400mg/mL Zeocin (Invitrogen) in complete media. The concentration of Zeocin was determined by performing a killing curve using varying concentrations of Zeocin on wild-type 293 cells that do not contain the transfected plasmid. Approximately 16 Zeocin-resistant clones were isolated using cloning rings and frozen in liquid nitrogen. A subset of the original clones could be expanded successfully and tested using an HER3-specific ELISA kit (R & D Systems, Inc), according to the manufacturers instructions and verified as over-expressing HER3 (Figure 1). One clone with high expression of HER3 as demonstrated by ELISA (293-H3 clone 1) was selected as a control for use in the optimized assay.

Example 2: Generation and Screening of antibodies against HER3

A HER3 -specific monoclonal antibody (Ab-6) with epitope specificity to the cytoplasmic terminus of HER3 was purchased from LabVision. The VERATAG®

reporter (Pro11) and streptavidin-conjugated methylene blue ("molecular scissors") were synthesized and purified according to protocol described previously (see, for example, above and United States Patent 7,105,308, which is incorporated by reference herein, including any drawings). Antibody-VERATAG® and antibody-biotin conjugates, i.e., Ab6-Pro11 and B9A11-biotin, were made using sulfo-NHS-LC-LC-biotin (Pierce) as linker according to manufacturer's protocol and conjugation products purified by HPLC (Agilent). A series of proprietary antibodies were generated as follows. Mice were immunized with fixed 293 clone 13 cells or a series of peptides representing different epitopes contained within the c-terminal region of the HER3 protein (Figure 2A). During the immunization period, each mouse received several immunizations over a 4 week period. Hybridomas were produced and clones isolated using limiting dilution. Conditioned media from individual clones were profiled by a series of assays, including CellSpot™ or ELISA screening, followed by scale-up and then further screening by immunohistochemistry (IHC) (Figure 2B), and then finally by VERATAG® Technology using either a chemical release method (Methylene Blue) (Figure 2C) or the dual-antibody light release method (Figure 2D). Several antibodies that performed well using these methods and are described in Figure 2A. One in particular, B9A11, was the most robust and gave the best dynamic range and sensitivity and was carried forward to develop the final format of the assay as described below in Example 5.

Example 3: Generation of blocks and FFPE sections from a panel of cell lines expressing varying levels of HER3 by FACS and ELISA.

Three cell lines with varying expression of HER3 protein, MDA-MB-453, MDA-MB-468 and SKOV-3, were purchased from American Type Cell Culture Collection. MDA-MB-453 and MDA-MB-468 cell lines were maintained at 37°C and 5% CO_2 in Dulbecco's modified Eagle medium (DMEM), 10% FBS, 1x penicillin-streptomycin and 1x Glutamax. SKOV3 cells were were maintained at 37°C and 5% CO_2 in McCoy's 5a Media supplemented with 10% FBS, 1x penicillin-streptomycin and 1x Glutamax (GIBCO). 293-H3 clone 1 cells were maintained at 37°C and 5% CO_2 in DMEM supplemented with 10% FBS, 1x penicillin-streptomycin and 1x Glutamax. Cells were grown to near confluence on at least ten 500mm culture plates for each cell line. After removal of medium, the cells were washed once with cold 1x PBS and 15 mL of 1x Pen-fix (Thermo Scientific) was added to each plate. Cells were scraped and the cell solution fixed overnight (>16 hrs) at 4°C. Following the overnight fixation the cells were centrifuged at 3200xg for 15 min. The cell pellet was transferred to a rubber O- ring, wrapped with filter paper and placed in a processing cassette. An automatic Tissue-Tek

processor was used for processing. Briefly, the cell pellet was exposed to increasing concentrations of alcohol, Clear-rite (xylene substitute) and paraffin. After processing, the pellet was embedded in a block using a paraffin embedding station. All solvents used for cell pellet processing were obtained from Richard-Allen Scientific. The proportion of the same lot of cells was tested for HER3 receptor number using flow cytometry by the following method. A whole cell lysate was prepared from the preparation of cells for quantifying levels of HER3 receptor by using a commercially available ELISA kit and following manufacturer's recommendations (Human ErbB3-DuoSet ELISA; R & D systems). Sections of 7 um in thickness were sliced with a microtome (LEICA) and placed on positively charged glass slides (VWR). Slides were air-dried for 30 min and then baked in a heated oven set at 70°C for 1.5 hr. All sample slides were stored at 4°C for future assays. Results of the ELISA, flow cytometry, IHC and VERATAG® comparison are shown in Figure 3.

Example 4: Generation of sections from commercially available breast cancer FFPE sections.

FFPE breast cancer blocks were purchased from Asterand. Sections of 5 um in thickness were sliced with a microtome (LEICA) and placed on positively charged glass slides (VWR). Sections were air-dried for 30 min and then baked in an oven at 70°C for 1.5 hr. All sample slides were stored at 4°C for future assays. Previously sectioned breast tumors from clinical material were also used for these studies. Examples of H & E stained tumors on glass slides are shown in Figure 4.

Example 5: Her-3 VERATAG® Assay in Formalin Fixed, Paraffin Embedded (FFPE) Cell Lines and Breast Tissue

FFPE samples were deparaffinized/rehydrated using a series of solvents. Briefly, slides were sequentially soaked in xylene (2×, 5 min), 100% ethanol (2×, 5 min), 70% ethanol (2×, 5 min) and deionized water (2×, 5 min). Heat-induced epitope retrieval of the rehydrated samples was performed in a slide holder containing 250 mL of 1× DAKO (pH 9.0) (Lab Vision) using a pressure cooker (Biocare). After being cooled for 30 min at room temperature, the slides were rinsed once with deionized water. A hydrophobic circle was drawn on slide using a hydrophobic pen (Zymed) to retain reagents on slides. The samples were then blocked for 1hr with blocking buffer that contains 1% mouse serum, 1.5% BSA and a cocktail of protease and phosphatase inhibitors (Roche) in 1×PBS. After removal of the blocking buffer with aspiration, a mixture of VERATAG®-conjugated (Ab-6: LabVision, 1ug/mL) and biotin-conjugated (B9A1 1; Monogram proprietary, 2ug/mL) antibodies prepared

in blocking buffer was added and binding reactions were incubated overnight in a humidified chamber at 4°C with shaking. The antibody mix was aspirated and samples were washed with wash buffer containing 0.25% TritonX-100 in 1×PBS and streptavidin-conjugated methylene blue at concentration of 2.5ug/mL in 1×PBS was added. After 1 hr incubation at room temperature, the excess streptavidin-methylene blue reagent was aspirated and the samples were washed in wash buffer once followed by 3 changes of deionized water. Illumination buffer containing 3 pM fluorescein and two CE internal markers (MF and ML) in 0.01×PBS was added on sample sections. The bound VERATAG® was released at ~4°C by photo-activated cleavage using an in-house LED array illuminator equipped with an electronic chiller block (Torrey Pine Scientific). After illumination, VERATAG® intermediates are reduced to a quantifiable form by the addition of sodium borohydride. The CE sample containing the released VERATAG® reporters was collected from above the tissue section on the slides and the released VERATAG® reporters in the CE samples were separated and detected on ABI3130 CE instrument (22-cm capillary array; Applied Biosystems) under CE injection condition of 6kV and 50 sec at 30°C. The general workflow of the H3T assay in the clinical lab is illustrated in Figure 5.

Example 6: CE Peak analysis, tumor area normalization and batch normalization

The identification and quantification of VERATAG® was carried out using VERATAG® Informer software (*see,* for example, United States publication number 2007-0203408-A1). To analyze the VERATAG® signals in a raw CE electropherogram, two CE internal markers, MF (first marker) and ML (last marker), were used to identify the VERATAG® peaks according to their electrophoretic mobility or migration time, t, relative to the two markers, *i.e.*, [t(VERATAG®)-t(MF)]/[t(ML)-t(MF)] . The identified VERATAG® peaks were then quantified by peak area calculation for each VERATAG®. To correct for variability in VERATAG® recovery from the tissue section, and the run variability in CE injection efficiency and/or detection sensitivity across capillary array, fluorescein (3 pM) was included in the illumination buffer, and co-electrophoresed as an internal reference control in each sample run. The area of each VERATAG® peak is then reported as RFU or RPA by area normalization of the VERATAG® peak (VERATAG® peak area) to the internal fluorescein peak (fluorescein peak area/3 pM. This is quantified as RPA*IB vol/TA for variable tumor samples (=Relative peak area multiplied by the illumination buffer volume (IB) loaded onto sample section; divided by the tumor area in mm^2 (RPA*IB vol/TA = pmole/L*L/mm^2 = pmole/mm^2). Specifically, the CE fluorescence signal intensity of a VERATAG® reporter, or

the peak area ($PA_{VeraTag®}$), is given in relative fluorescent units (signal height) integrated across time (RFU-S/S). The relative peak area ($RPA_{VeraTag®}$ is measured by normalizing the VERATAG® peak area ($PA_{VeraTag®}$) with respect to the internal fluorescein standard of known concentration (PA_F), and is therefore proportional to the initial concentration of the analyte being measured. As the VERATAG®assay signal is a quantitative readout that scales with tumor content, accurate comparison of VERATAG® assay signals across clinical samples requires adjustment for differences in this parameter. Therefore, tumor content is measured on the sample VERATAG® assay signal data collection, and the tumor content is used to normalize the VERATAG® assay result. It should be noted that, if the tumor sample is either very small or very large the reaction volumes (i.e., volume of antibody, streptavidin-conjugated methylene blue and illumination buffer reagents) are adjusted, and this adjustment is reflected in the tumor content normalization. After adjusting VERATAG® peak areas for migration time, fluorescein, illumination buffer and tumor area, controls and samples are normalized by multiplying their adjusted peak area with the respective calculated Batch Normalization Factor (BNF). Each adjusted RPA value is multiplied by the respective BNF to obtain a normalized RPA value. Because this normalized RPA value is unitless by definition, its value is referred to in VERATAG® units. All reportable (not failed, and not saturated) normalized values for a given sample are averaged to determine a final value for that sample.

Various quality control checks have been developed to ensure data is of highest quality.

1. Fluorescein out of range. A sample trace with fluorescein out of range is failed. The fluorescein range is calculated by using the median of the adjusted fluorescein value +/- XX%. The XX% value, typically 20-40%, is adjustable and is associated with individual templates. Sample peaks are failed if the absolute height of the peak is >7000 units, referred to as a saturated peak.
2. RPA must be greater than or equal to 0.03.
3. If an undiluted converted peak sample is failed due to RPA <0.03, then all of the corresponding converted peak diluted samples are also failed.
4. At least 2 converted peak controls must have called values for converted peak batch normalization to proceed.
5. A sample or control with poor trace quality is failed.
6. If illumination buffer-only controls have contamination, samples may be failed if trace quality may be affected, e.g., if the contaminating peak overlaps with

the released peak

7. A batch with poor batch normalization is failed.

8. A batch with an abnormally high batch normalization factor is failed.

For each sample, following the VERATAG® assay, an H&E (Hematoxylin and Eosin) staining and evaluation is performed to assure presence of tumor cells and to enable an estimation of tumor area. These slides are deparaffinized, hydrated, stained, and then dehydrated and mounted using standard procedures before microscopic examination.

Example 7: Optimization of antibody concentration to increase dynamic range

Optimal concentrations of the Ab-6 and the B9A1 1 antibody were determined by varying the final concentrations in the VERATAG® HER3 total assay (as described above) on a cell line panel spanning the entire dynamic range of the assay (293H3-clone 1, MDA-MB-453, MDA-MB-468, and SKOV3). These results were then compared this with expected fold changes based on HER3 flow cytometry and ELISA results from the same cell line FFPE block preparation. An optimal concentration of 2mg/mL B9A11-biotin and 1mg/mL of Ab-6 Pro-11 was selected for performance based on the accurate detection of HER3 as compared with the expected fold changes as compared to ELISA and flow cytometry in this same set of cell lines as shown in Figure 6 where a 2:1 ratio of B9A11 to Ab-6 is circled.

Example 8: Accuracy of HER3 total VERATAG® assay

One batch of the HER3 total VERATAG® assay was performed using three successful replicates from four well-characterized cell lines (293H3-clone 1, MDA-MB-453, MDA-MB468, MDA-MB-231 and SKOV3) and then compared for accuracy of VERATAG® measurement with in-house generated flow cytometry and ELISA data. 100% of the results matched the in- house data from flow cytometry and ELISA in that 293H3-clone 1> MDA-MB-453>MDA-MB468>SKOV3. No overlap was observed between signal levels for any of the four cell line samples, *i.e.*, each cell line separated completely. Internal datasets on HER3 Total levels were generated by both ELISA and flow cytometry. Results for the four accuracy cell lines are presented in Figure 7. Results from HER3 flow cytometry were very similar to results from HER3 ELISA in that the same rank order preservation was demonstrated by using these cross- validating technologies.

Example 9: Sensitivity of HER3 total VERATAG® assay

The sensitivity of the HER3 total VERATAG® assay was determined by comparing one batch containing 8 replicates of the low HER3 expression control cell line, MDA-MB-468 with 8 replicates of the low/negative HER3 expression control cell line, SKOV3. The values

for each replicate of MDA-MB-468 were compared with SKOV3 replicates by pairwise comparisons to determine sensitivity. 100% (64/64) of the pairwise comparisons between MDA-MB-468 and SKOV3 resulted in MDA-MB-468>SKOV3 (Figure 8).

Example 10: Reproducibility of HER3 total VERATAG® assay

Inter-assay reproducibility was determined by performing 8 separate batches of the HER3 total VERATAG® assay as described above on 4 well characterized cell lines (293H3-clone 1, MDA-MB-453, MDA-MB-468 and SKOV3) using different instrumentation (CE, illuminators), several operators, and performing the assay over a 4 week period. Following batch normalization procedures, the data was compared across the 8 batches to determine reproducibility. The reproducibility across the dynamic range was between 8-15% (see Figure 9).

Example 11: Precision of HER3 total VERATAG® assay

Intra-assay reproducility was determined by performing 1 batch/cell line of the HER3 total VERATAG® assay and comparing the performance of 15 replicates of each of the 3 control cell lines (293H3-clone 1, MDA-MB-453, MDA-MB-468). Pairwise comparisons were made of the 15 replicates in each batch to determine precision of the assay. 95% of the 293H3-clone 1 data was within 1.2 fold, 100% of the MDA-MB-453 data was within 1.23-fold, and 95% of the MDA-MB-468 data is within 1.37-fold. Results are shown in Figure 10.

Example 12: Linearity of the HER3 total VERATAG® Assay.

The linearity of the HER3 VERATAG® result was determined by taking well-characterized cell line controls 293H3-clone 1, MDA-MB-453, MDA-MB-468) and performing successive "cut-down" experiments to create FFPE sections with the following dimensions 1, ½, ¼, 1/16. These "cut-down" sections were then run in the H3T VERATAG® assay and the final section area normalized data was compared in a pairwise manner to understand the linearity of the assay with respect to section size (Figure 11). MDA-MB-453 is linear down to approximately 1/16 of the original section size, while MDA-MB-468 is linear down to approximately should be 1/2 of the original section size. Results are shown in Figure 11.

Example 13: Specificity of the HER3 total VERATAG® Assay

Patient-derived tumor samples and cell line controls were run using the VERATAG® HER3 Total Assay and using isotype control antibodies. For the HER3 Ab-6-Pro11, the matched isotype control was IgG1-Pro11. For the HER3 B9A11-biotin, the matched isotype control was also IgG1-biotin. Signal from these reactions is not antigen-

specific, and would represent non-specific background. Samples were run in each of the following three formats and run within the same batch side by side:

Format 1: HER3 Ab6-Pro11/B9A11 -biotin (normal format)

Format 2: HER2 Ab6-Pro11/IgG1-biotin

Format 3: IgG1-Pro11/B9A11-biotin

Sample results from each IgG1 format (Format 2 and Format 3) were compared to the negative control, SKOV3, present in each batch (comparison parameter A). Samples results were also compared to the respective actual HER3 Total signals (Format 1). Results are shown in Figure 12.

Example 14: Measurement of clinical breast tumors and dynamic range

The study population comprised patients (n=105) that were prospectively observed during trastuzumab-based therapy at a single institution between 1999 and 2006 (the International Serum Her-2/neu Study Group trial). Only outpatients (ECOG PS 0–2, age >18 years, estimated life expectancy >12 weeks) with HER-2/neu-overexpressing (>10% of tumor cells IHC 3+ as determined by the HercepTest; DAKO Diagnostics, Austria) and/or ERBB2- amplified (with positive FISH testing mandatory in all IHC 2+ cases) MBC were included. In addition, patients were required to be trastuzumab-naïve and have bi-dimensionally measurable disease progressing within 4 weeks before initiation of treatment (excluding previously irradiated lesions). Samples from this study group trial were run in the H3T VERATAG® assay in eight separate batches. For each batch, CE peak analyses were performed, as well as tumor area analyses and the batch was normalized accordingly. Out of the 105 patient samples run in the assay, 85 samples had measurable H3T levels above the limit of detection, 8 patient samples had no detectable tumor, 1 sample demonstrated a fluorescein failure excluding the data, and finally there were 8 patient samples that were below the limit of detection of the assay and considered low/negative in H3T expression. The results are shown in figure 13.

Example 15: Determination of optimal cutoff for trastuzumab response

Using positional scanning analysis an optimal cut-off was determined whereby patients above statistically significant cut-off had an unfavorable time to progression (TTP) compared to patients that were below this cut-off (Figure 14). The cut-off was ~ the median of the results of the population tested (0.158, HR=2.3, p=0.0004. TTP was defined as the time from the initiation of trastuzumab-containing treatment to progression (SWOG) or censor, and OS was defined as the time from initiation of trastuzumab-containing treatment to death or censor. When looking at overall survival in this population of patients no

significant cut-off could be determined, however there was a trend in OS using the 0.158 cut-off (HR=1 .7; p=0.059).

Example 16: Kaplan Meier (KM) analysis for TTP

A previously reported H2T cutoff (logH2T $\geq$ 1.14) was used to sub-divide the patients into HER2- normal (N=26, median TTP = 4.1 mos) and HER2-overexpressing (N=55, median TTP = 11.1 mos, HR=0.43, p=0.0002) groups. In the HER2-overexpressing group, levels of H3T expression above an optimal cutoff (H3T > 0.158; Figure 14), as defined by a positional scanning predicted shorter median time to progression (N=25, median TTP = 6.1 mos) compared with H3T expression below the cutoff (N=30, median TTP=13.1 mos, HR=2.7, p=.0002). Univariate Cox proportional hazards analyses examining the HER2-overexpressing sub-group identified H3T (above vs below a particular cutoff) as the most significant predictor of TTP (HR=2.98, p=0.0004). These results are shown in Figure 15.

Example 17: HER3 total expression by VERATAG® in other malignancies

H3T VERATAG® assay was performed on a number of different malignancies other than breast cancer, including colon, ovarian, synovial tumors (Figure 16). Similar dynamic ranges were observed for all of these cancers with the following rank order of range: Ovarian>Synovial Carcinoma>/=Colon. The dynamic range in these tumors ranges from 0.5-1.5 logs depending on the cancer.

Example 18: Measurement of HER3 in conjunction with P95 and correlation with trastuzumab response

A previously reported P95 cut-off (US Patent Application 12/629,037) was used to further stratify the HER-2 over-expressing patients described above after initial subdivision based on their level of HER3 to produce 4 patient subgroups as shown in Figure 17. Patients with a low level of P95 and HER3, defined as below an optimal cut-off predicted by positional scanning, had a longer median time to progression (TTP=15.0 mos) than any of the other subgroups (logrank test for trend p<0.0001). Patients with a high level of P95 and HER3 had the shortest median time-to-progression (TTP=3.2 mos). The results are shown in Figure 17

Example 19: Measurement of HER2, HER3 and P95 and correlation with trastuzumab response

The patient population in Example 17 was subdivided by their level of HER2, P95 and HER3 as shown in Figure 18. Patients who are HER2-high, P95-low, and HER3-low, as defined by a positional scanning methodology, had a longer time to progression than patients who have a high level of HER2 and a high level of HER3 or P95 or both with the latter

showing a similar time to progression as patients who have a low HER2 value as determined by the H2T VERATAG® assay. Within the subgroup with normal HER2 expression levels (H2Tlo) FISH-positive and FISH-negative groups experienced a similar time to progress.

All publications and other materials described herein are used to illuminate the invention or provide additional details respecting the practice and are incorporated by reference in their entirety.

Claims

1. A monoclonal antibody comprising:

(a) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:13, 14 and 15, respectively, and a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:16, 17 and 18; and/or

(b) a light chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs: 19, 20 and 21, respectively, and a heavy chain variable region comprising CDR1, CDR2 and CDR3 having sequences as set forth in SEQ ID NOs:22, 23 and 24, respectively; and/or

(c) an amino acid sequence comprising SEQ ID NOs: 9 and 11 for the light and heavy chains, respectively; and/or

(d) an amino acid sequence comprising SEQ ID NOs:10 and 12 for the light and heavy chains, respectively.

2. A method of measuring and/or quantifying the presence and/or amount of Her-3 or Her-3 in a complex in a sample from a subject, the method comprising mixing the sample with an antibody of claim 1 and determining the presence and/or quantity of the antibody bound to Her-3 and/or Her-3 in a complex.

3. A method for determining whether a subject with a cancer is likely to respond to treatment with a Her family-targeted agent, for predicting a time course of disease and/or for predicting the probability of a significant event in the time course of the subject's cancer, comprising the method according to claim 2.

4. The method of claim 3, wherein if the level of Her-3 is high, the patient is less likely to respond to the Her family-targeted agent; and if the level of Her-3 is low, the patient is more likely to respond to the Her family-targeted agent.

5. The method of any of claims 2 to 4, wherein the method comprises measuring and/or quantifying the presence and/or amount of at least one of total Her-3, Her-3 homodimers, or Her-3 heterodimers.

6. The method of any one of claims 2 to 5, further comprising measuring the amount p95 in the sample.

7. The method of claim 6, wherein if the amount of p95 and Her-3 expression is low, then the patient is likely to respond to the Her-family targeted agent and/or the patient has a long time course.

8. The method of any one of claims 2 to 7, wherein the sample comprises a tumor sample.

9. The method of any one of any one of claims 2 to 8, wherein the sample comprises a tumor sample; and/or the sample comprises a breast cancer sample, a colorectal cancer sample, an ovarian cancer sample, a non-small cell lung cancer sample, or a gastric cancer sample.

10. The method of any one of claims 2 to 9, wherein the sample comprises a Her-2 positive sample.

11. The method of any one of claims 2 to 10, wherein the sample comprises breast cancer.

12. The method of any one of claims 2 to 11, wherein the sample comprises a Her-2 positive breast cancer sample.

13. The method of any of one of claims 2 to 12, wherein the sample comprises an early stage breast cancer sample or a metastatic breast cancer sample

14. The method of any one of claims 2 to 13, wherein the sample is an FFPE sample or solubilized FFPE sample.

15. The method of any one of claims 2 to 14, wherein the method comprises:

(a) mixing (i) the sample from the subject; (ii) a proximity probe that is capable of binding Her-3, the proximity probe having an effective proximity and (iii) at least one binding compound, the at least one binding compound being capable of

binding Her-3 and having one or more signaling molecules attached, wherein binding of the proximity probe and the binding compound within the effective proximity produces a signal from the molecular tags that correlates with the presence and/or quantity of Her-3 and/or Her-3 in a complex;

(b) detecting the signal from the molecular tags to determine the presence and/or quantity of Her-3 and/or Her-3 in a complex;,

wherein the proximity probe and/or the binding compound comprises the antibody of claim 1.

16. The method of claim 15, wherein the proximity probe comprises a cleaving probe that has a cleavage-inducing moiety and the at least one binding compound has one or more molecular tags attached to the binding compound by a cleavable linkage, wherein the cleavable linkage may be cleaved within the effective proximity, producing a signal that correlates with the presence and/or quantity of Her-3.

17. The method of any one of claims 2 to 14, wherein the method comprises the steps of:

(a) contacting the sample with a tagged Her-3 binding composition that specifically binds to Her-3 and has one or more molecular tags attached thereto via a cleavable linkage;

(b) contacting the sample with a cleaving agent;

(c) cleaving the cleavable linkage of the tagged Her-3 binding composition, thereby releasing the one or more molecular tag; and

(d) quantitating the released molecular tags to determine the presence and/or amount of Her-3 in the sample,

wherein the a tagged Her-3 binding composition comprises the Her-3 antibody of claim 1.

18. The method of any one of claims 3 to 17, wherein the Her family-targeted agent comprises a multi- or single-targeted agent.

19. The method of claim 18, wherein the Her family-targeted agent comprises a dual kinase inhibitor or a bispecific antibody.

20. The method of any one of claims 3 to 19, wherein the Her family-targeted agent comprises trastuzumab, lapatinib, pertuzumab, cetuximab, panitumumab, erlotinib, or gefitinib.

21. The method of claim 20, wherein the Her family-targeted agent comprises trastuzumab.

22. The method of any of claims 3 to 21, wherein likeliness to respond, likeliness to have a long time course and/or likeliness to have a significant event is measured as at least one of an overall survival rate, as time to progression, as disease-free survival, as progression-free survival, time to distant reoccurrence, hazard ratio and/or as objective tumor response or clinical benefit using the RECIST criteria.

23. An antibody according claim 1 for use in any one of the methods of claims 2 to 22.