



166166



G00002

TRANSGENIC ANIMALS FOR
PRODUCING SPECIFIC ISOTYPES OF
HUMAN ANTIBODIES VIA NON-COGNATE SWITCH REGIONS

BACKGROUND OF THE INVENTION

5 A quarter century after the discovery of
monoclonal antibodies (mAbs) [G. Kohler and C.
Milstein, Nature 256:495-497 (1975)], their therapeutic
utility is finally being realized. Monoclonal
antibodies have now been approved as therapies in
10 transplantation, cancer, infectious disease,
cardiovascular disease and inflammation. Many
monoclonal antibodies are in late stage clinical trials
to treat a broad range of disease indications. As a
result, mAbs represent one of the largest classes of
15 drugs currently in development.

 The utility of mAbs stems from their specific
recognition of a complex target followed by high
affinity binding to that target. Because different C_H
isotypes have different effector functions, it is
20 desirable to tailor the mAb isotype to the desired
effector function. For, example, a mAb bearing a
constant region with effector functions, e.g., human
IgG₁, can be used to direct complement dependent
cytotoxicity or antibody-dependent cytotoxicity to a

- 2 -

target cell. Alternatively, a mAb with a constant region essentially lacking effector function, e.g., human IgG₂ or IgG₄, can be used to block signal transduction, either by binding to and neutralizing a
5 ligand, or by blocking a receptor binding site.

Many therapeutic applications for monoclonal antibodies require repeated administrations, especially for chronic diseases such as autoimmunity or cancer. Because mice are convenient for immunization and
10 recognize most human antigens as foreign, mAbs against human targets with therapeutic potential have typically been of murine origin. However, murine mAbs have inherent disadvantages as human therapeutics. They require more frequent dosing to maintain a therapeutic
15 level of mAb because of a shorter circulating half-life in humans than human antibodies. More critically, repeated administration of murine immunoglobulin creates the likelihood that the human immune system will recognize the mouse protein as foreign, generating
20 a human anti-mouse antibody (HAMA) response. At best, a HAMA response will result in a rapid clearance of the murine antibody upon repeated administration, rendering the therapeutic useless. More likely is that a HAMA response can cause a severe allergic reaction. This
25 possibility of reduced efficacy and safety has lead to the development of a number of technologies for reducing the immunogenicity of murine mAbs.

In order to reduce the immunogenicity of antibodies generated in mice, various attempts have
30 been made to replace murine protein sequences with human protein sequences in a process now known as humanization. The first humanization attempts utilized molecular biology techniques to construct recombinant

- 3 -

antibodies. For example, the complementarity determining regions (CDR) from a mouse antibody specific for a hapten were grafted onto a human antibody framework, effecting a CDR replacement. The new antibody retained the binding specificity conveyed by the CDR sequences. [See P.T. Jones et al. *Nature* 321: 522-525 (1986)]. The next level of humanization involved combining an entire mouse VH region (HuVnp) with a human constant region such as $\gamma 1$. [S.L. Morrison et al., *Proc. Natl. Acad. Sci.*, 81, pp. 6851-6855 (1984)]. Such chimeric antibodies, which still contain greater than 30% xenogeneic sequences, are sometimes only marginally less immunogenic than totally xenogeneic antibodies. [M. Bruggemann et al., *J. Exp. Med.*, 170, pp. 2153-2157 (1989)].

Subsequently, attempts were carried out to introduce human immunoglobulin genes into the mouse, thus creating transgenic mice capable of responding to antigens with antibodies having human sequences. [See Bruggemann et al. *Proc. Nat'l. Acad. Sci. USA* 86:6709-6713 (1989)]. These attempts were thought to be limited by the amount of DNA which could be stably maintained by available cloning vehicles. As a result, many investigators concentrated on producing mini-loci containing limited numbers of V region genes and having altered spatial distances between genes as compared to the natural or germline configuration. [See United States Patent 5,569,825 to Lonberg et al., (1996)]. These studies indicated that producing human sequence antibodies in mice is possible, but serious obstacles remained regarding obtaining sufficient diversity of binding specificities and effector functions (isotypes)

- 4 -

from these transgenic animals to meet the growing demand for antibody therapeutics.

In order to provide additional diversity, work has been conducted to add large germline fragments of the human Ig locus into transgenic mammals. For example, a majority of the human V, D, and J region genes arranged with the same spacing found in the unrearranged germline of the human genome and the human C_μ and C_δ constant regions was introduced into mice using yeast artificial chromosome (YAC) cloning vectors. [See PCT patent application WO 94/02602 to Kucherlapati et al.]. A 22 kb DNA fragment comprising sequences encoding a human gamma-2 constant region and the upstream sequences required for class-switch recombination was latter appended to the foregoing transgene. In addition, a portion of a human kappa locus comprising V_k, J_k and C_k region genes, also arranged with substantially the same spacing found in the unrearranged germline of the human genome, was introduced into mice using YACS. Gene targeting was used to inactivate the murine IgH & kappa light chain immunoglobulin gene loci and such knockout strains were bred with the above transgenic strains to generate a line of mice having the human V, D, J, C_μ, C_δ and Cγ2 constant regions as well as the human V_k, J_k and C_k region genes all on an inactivated murine immunoglobulin background. [See PCT patent application WO 94/02602 to Kucherlapati et al.; see also Mendez et al., *Nature Genetics* 15:146-156 (1997)].

Yeast artificial chromosomes as cloning vectors in combination with gene targeting of endogenous loci and breeding of transgenic strains

- 5 -

provided one solution to the problem of antibody diversity. Several advantages were obtained by this approach. One advantage was that YACs can be used to transfer hundreds of kilobases of DNA into a host cell.

5 Therefore, use of YAC cloning vehicles allows inclusion of substantial portions of the entire human Ig Heavy and light chain regions into a transgenic animal thus approaching the level of potential diversity available in the human. Another advantage of this approach is

10 that the large number of V genes has been shown to restore full B cell development in mice deficient in murine immunoglobulin production. This ensures that these reconstituted mice are provided with the requisite cells for mounting a robust human antibody

15 response to any given immunogen. [See PCT patent application WO 94/02602 to Kucherlapati et al.; L.Green and A. Jakobovits, *J. Exp. Med.* 188:483-495 (1998)]. A further advantage is that sequences can be deleted or inserted onto the YAC by utilizing high frequency

20 homologous recombination in yeast. This provides for facile engineering of the YAC transgenes.

As mentioned above, there are several strategies that exist for the generation of mammals that produce human antibodies. In particular, there is

25 the "minilocus" approach that is typified by work of GenPharm International, Inc. and the Medical Research Council, YAC introduction of large and substantially germline fragments of the Ig loci that is typified by work of Abgenix, Inc. (formerly Cell Genesys), and

30 introduction of entire or substantially entire loci through the use microcell fusion as typified by work of Kirin Beer Kabushiki Kaisha.

- 6 -

In the minilocus approach, an exogenous Ig locus is mimicked through the inclusion of pieces (individual genes) from the Ig locus. Thus, one or more V_H genes, one or more D_H genes, one or more J_H genes, a mu
5 constant region, and a second constant region (preferably a gamma constant region) are formed into a construct for insertion into an animal. This approach is described or related to work in U.S. Patent No. 5,545,807 to Surani et al. and U.S. Patent Nos.
10 5,545,806, 5,625,825, 5,625,126, 5,633,425, 5,661,016, 5,770,429, 5,789,650, and 5,814,318 each to Lonberg and Kay, U.S. Patent No. 5,591,669 to Krimpenfort and Berns, U.S. Patent Nos. 5,612,205, 5,721,367, 5,789,215 to Berns et al., and U.S. Patent No. 5,643,763 to Choi
15 and Dunn, and GenPharm International U.S. Patent Application Serial Nos. 07/574,748, filed August 29, 1990, 07/575,962, filed August 31, 1990, 07/810,279, filed December 17, 1991, 07/853,408, filed March 18, 1992, 07/904,068, filed June 23, 1992, 07/990,860,
20 filed December 16, 1992, 08/053,131, filed April 26, 1993, 08/096,762, filed July 22, 1993, 08/155,301, filed November 18, 1993, 08/161,739, filed December 3, 1993, 08/165,699, filed December 10, 1993, 08/209,741, filed March 9, 1994, the disclosures of which are
25 hereby incorporated by reference. See also European Patent No. 0 546 073 B1, International Patent Application Nos. WO 92/03918, WO 92/22645, WO 92/22647, WO 92/22670, WO 93/12227, WO 94/00569, WO 94/25585, WO 96/14436, WO 97/13852, and WO 98/24884, the disclosures
30 of which are hereby incorporated by reference in their entirety. See further Taylor et al. "A transgenic mouse that expresses a diversity of human sequence

- 7 -

heavy and light chain immunoglobulins." *Nucleic Acids Research* 20:6287-6295 (1992), Chen et al.

"Immunoglobulin gene rearrangement in B-cell deficient mice generated by targeted deletion of the J_H locus"

5 *International Immunology* 5:647-656 (1993), Tuailon et al. "Analysis of direct and inverted DJ_H rearrangements in a human Ig heavy chain transgenic minilocus" *J.*

Immunol. 154:6453-6465 (1995), Choi et al. "Transgenic mice containing a human heavy chain immunoglobulin gene
10 fragment cloned in a yeast artificial chromosome"

Nature Genetics 4:117-123 (1993), Lonberg et al.

"Antigen-specific human antibodies from mice comprising four distinct genetic modifications." *Nature*

368:856-859 (1994), Taylor et al. "Human immunoglobulin
15 transgenes undergo rearrangement, somatic mutation and class switching in mice that lack endogenous IgM."

International Immunology 6:579-591 (1994), Tuailon et al. "Analysis of direct and inverted DJ_H rearrangements in a human Ig heavy chain transgenic minilocus" *J.*

20 *Immunol.* 154:6453-6465 (1995), and Fishwild et al.

"High-avidity human IgG monoclonal antibodies from a novel strain of minilocus transgenic mice." *Nature Biotech.* 14:845-851 (1996), the disclosures of which are hereby incorporated by reference in their entirety.

25 In connection with YAC introduction, Green et al. *Nature Genetics* 7:13-21 (1994) describes the generation of YACs containing 245 kb and 190 kb-sized germline configuration fragments of the human heavy chain locus and kappa light chain locus, respectively,
30 which contained core variable and constant region sequences. *Id.* The work of Green et al. was recently extended to the introduction of greater than

- 8 -

approximately 80% of the human antibody repertoire through introduction of megabase sized, germline configuration YAC fragments of the human heavy chain loci and kappa light chain loci, respectively, to

5 produce XenoMouse™ mice. See Mendez et al. *Nature Genetics* 15:146-156 (1997), Green and Jakobovits *J. Exp. Med.* 188:483-495 (1998), and U.S. Patent Application Serial No. 08/759,620, filed December 3, 1996, the disclosures of which are hereby incorporated

10 by reference. Such approach is further discussed and delineated in U.S. Patent Application Serial Nos. 07/466,008, filed January 12, 1990, 07/610,515, filed November 8, 1990, 07/919,297, filed July 24, 1992, 07/922,649, filed July 30, 1992, filed 08/031,801,

15 filed March 15, 1993, 08/112,848, filed August 27, 1993, 08/234,145, filed April 28, 1994, 08/376,279, filed January 20, 1995, 08/430, 938, April 27, 1995, 08/464,584, filed June 5, 1995, 08/464,582, filed June 5, 1995, 08/463,191, filed June 5, 1995, 08/462,837,

20 filed June 5, 1995, 08/486,853, filed June 5, 1995, 08/486,857, filed June 5, 1995, 08/486,859, filed June 5, 1995, 08/462,513, filed June 5, 1995, 08/724,752, filed October 2, 1996, and 08/759,620, filed December 3, 1996. See also Mendez et al. *Nature Genetics*

25 15:146-156 (1997) and Green and Jakobovits *J. Exp. Med.* 188:483-495 (1998). See also European Patent No., EP 0 463 151 B1, grant published June 12, 1996, International Patent Application No., WO 94/02602, published February 3, 1994, International Patent

30 Application No., WO 96/34096, published October 31, 1996, and WO 98/24893, published June 11, 1998. The disclosures of each of the above-cited patents,

- 9 -

applications, and references are hereby incorporated by reference in their entirety.

In connection with the microcell fusion approach, portions or whole human chromosomes can be introduced into mice as described in European Patent Application No. EP 0 843 961 A1, the disclosure of which is hereby incorporated by reference. It will be understood that mice generated using this approach and containing the human Ig heavy chain locus will generally possess more than one, and potentially all, of the human constant region genes. Such mice will produce, therefore, antibodies that bind to particular antigens having a number of different constant regions. Thus, there is no way to preselect the desired constant region for particular effector function.

Also, the transchromosomes are mitotically and meiotically unstable. As a result, either the human IgH, the human IgK or both transchromosomes are lost with a frequency approaching 80%. This results in aberrantly high recovery of mouse Ig λ mAbs and hybridoma instability.

Technology exists for in vitro isotype switching of antibodies. Antibodies produced from transgenic mice that produce only IgG1 isotypes, from transgenic mice that produce multiple IgG isotypes, or from phage display technologies may have the desired antigen-specificity and affinity, but not have the desired effector function. In this instance, the variable region of the heavy chain, at the least, and most likely, the entire light chain of the antibody must be cloned.

Methods for cloning include recovery of genomic DNA from a library, recovery of cDNA from a

- 10 -

library, recovery of genomic DNA using specific oligonucleotide primers, and PCR using specific oligonucleotide primers and cDNA as template (RT-PCR). Each method, especially PCR-based methods, require that
5 the clone be sequenced to verify faithful reproduction of the antibody coding sequences. Then the variable region of the heavy chain must be operably linked via DNA ligation to the desired constant region gene. Then, the engineered VH-CH gene must be operably linked
10 to expression controlling regions such as a promoter-enhancer and a polyadenylation site. Such an expression construct might also be needed for the Ig light chain of the antibody.

The expression construct(s) must be stably
15 transfected into a suitable host cell for transcription and translation to produce a secreted form of the engineered mAb. Typically, at the least, extensive screening must be performed to find a clone of the cell line that expresses sufficient levels of mAb for
20 further experiments and subsequent manufacturing. More likely, methodologies such as DNA amplification must be employed to raised the copy number of the antibodies expression constructs and consequently, the expression level of the mAb.

25 Finally, the re-engineered mAb must be re-tested to confirm that it has retained the desired qualities and has the desire function, including specificity, affinity, and presence or absence of effector function. Other technologies for isotype
30 switching exist, but all such programs to re-engineer the mAb isotype require experimentation and expertise in molecular biology and tissue culture, and are labor intensive, slow, expensive, and covered by issued and

- 11 -

pending intellectual property, requiring additional
licensing fees, if even available for licensing. Thus,
re-engineering of mAb from one isotype to another
requires expertise, extra monetary expenditure and
5 slows down the development of the monoclonal antibody
for pre-clinical and clinical trials.

Having a technology that would produce the
mAb with the desired Cy isotype a priori would obviate
the need for antibody re-engineering. By having three
10 different XenoMouse strains, one each capable of making
only Cy2, Cy4 or Cy1, a transgenic mouse can be pulled
off the shelf, and then can be immunized to produce
mAbs with the desired affinity, antigen-specificity and
the desired isotype and with the desired effector
15 function a priori. This increases the efficiency and
user-friendliness for development of monoclonal
antibody based therapeutics. No expertise in molecular
biology or antibody engineering is required. The
antigen-specific mAb can be taken directly into pre-
20 clinical studies without the extra expenditure of money
and time, resulting in a decrease in the development
cost and an acceleration of the timeline for
development of the therapeutic mAb.

The present invention is directed to solving
25 the problem of obtaining a pre-selected human antibody
isotype from a transgenic mouse, in addition to the
desired specificity, which is compatible with the
therapeutic goals for which the antibody will be used.

SUMMARY OF THE INVENTION

30 The present invention solves the problems
referred to above by providing, in one aspect of the
invention, transgenic non-human animals capable of

- 12 -

producing high affinity, fully human antibodies of a desired isotype in response to immunization with any virtually any desired antigen. The aforementioned transgenic non-human animals have in their somatic and
5 germline cells an unrearranged human immunoglobulin heavy chain transgene that encodes, on rearrangement, a fully human immunoglobulin heavy chain of the desired isotype.

The human immunoglobulin heavy chain transgene in the
10 foregoing animals comprises a human constant region gene segment comprising exons encoding the desired heavy chain isotype, operably linked to switch segments from a constant region of a different heavy chain isotype, i.e., a non-cognate switch region.

15 The foregoing transgenic non-human animal also has in its somatic and germ cells a human immunoglobulin light chain transgene. In a preferred embodiment, the endogenous immunoglobulin heavy and light chain loci of the transgenic non-human animal are
20 inactivated so that the animal is incapable of producing endogenous heavy or light chains. In a particularly preferred embodiment, the non-human transgenic animal is a mouse.

In another aspect, the invention provides an
25 unrearranged human immunoglobulin heavy chain transgene that encodes, on rearrangement, for a human heavy chain of a desired isotype. The transgenes of the invention comprise a DNA sequence identical to the DNA sequence of human chromosome 14 starting at least from the first
30 D segment gene of the human immunoglobulin heavy chain locus, continuing through the J segment genes and the constant region genes through C_μ of that locus. In the transgenes of the invention, the aforementioned DNA

- 13 -

fragment is operably linked to and is capable of isotype switching to an additional constant region segment. Said additional constant region segment comprises a switch region and human constant region coding segment, wherein the constant region coding segment is operably linked to a switch region that it is not normally associated with, i.e., a non-cognate switch region. In transgenes of the invention, the foregoing DNA fragment and constant region segment is operably linked to at least one human V segment gene. In one embodiment of the invention, the transgene is a yeast artificial chromosome (YAC).

In the transgenes of the invention, the non-cognate switch region may be a switch region from a different species than the constant region coding segment. In one embodiment, the non-cognate switch region is a mouse switch region operably linked to a human constant region coding segment encoding a human gamma, alpha or epsilon constant region. In a preferred embodiment, the switch region is a mouse gamma-1 switch region. In more preferred embodiments, the switch region is a mouse gamma-1 switch region and the human constant region coding segment encodes a gamma-1 or a gamma-4 constant region. In a particularly preferred embodiment, the transgene is the yH2Bm yeast artificial chromosome (YAC) or the yH2Cm YAC.

In another embodiment, both the non-cognate switch region and the constant region coding segment are human sequences, the non-cognate switch region being from a human constant region of a different isotype than the constant region coding segment. In a preferred embodiment, the switch region is a human

- 14 -

gamma-2 switch region and the constant region coding segment is an isotype other than gamma-2. In a more preferred embodiment, a transgene of the invention comprises a human gamma-2 switch region and a human
5 gamma-1 or human gamma-4 constant region coding segment. In particularly preferred embodiments, the transgene is the yHG1 YAC or the yHG4 YAC.

In still another embodiment, a transgene of the invention comprises a human non-cognate switch
10 region and a human constant region coding segment, wherein the switch region and the membrane exons of the constant region coding segment are from the same human constant region isotype and the secreted constant region exons are from a different isotype. The
15 transgenes of the invention also may comprise a human non-cognate switch region and a human constant region coding segment wherein the switch region is from one isotype, the secreted constant region exons are from a second isotype and the membrane constant region exons
20 are from yet a third isotype.

In a preferred embodiment, the switch region and membrane exons are from a human gamma-2 constant region. In particularly preferred embodiments, the switch region and membrane exons are from a human
25 gamma-2 constant region and the secreted constant region exons are from a human gamma-1 or a human gamma-4 constant region. In preferred embodiments, the transgene is the yHG1/2 YAC or the yHG4/2 YAC.

In another embodiment, any of the foregoing
30 transgenes of the invention comprise a plurality of different human VH genes. In a preferred embodiment, the transgene comprises at least 50% of the human germline VH genes. In another embodiment, the

- 15 -

transgene comprises at least 40 different human VH genes. Preferably, the transgene comprises at least 66 different human VH genes. Most preferably, the transgene comprises the entire human VH region of a human heavy chain locus. In another embodiment, the transgene comprises a sufficient number of different human VH genes so that the transgene is capable of encoding at least 1×10^5 different functional human immunoglobulin heavy chain sequence combinations, without taking into account junctional diversity or somatic mutation events. In still another embodiment, the number of human VH genes in the transgene is sufficient to produce at least 50% of the B-cell population of a wild-type mouse in a transgenic mouse containing the transgene.

A transgene of the invention further comprises a murine 3' enhancer, positioned 3' of the constant region gene containing the non-cognate switch region. In one embodiment the murine 3' enhancer is an approximately 0.9 kb core region of the native enhancer. In an alternative embodiment, the 3' enhancer is an approximately 4 kb region of the murine enhancer that includes the core region. In still another embodiment, the transgene includes the mouse major enhancer locus.

In another aspect, the invention provides methods for producing the transgenic non-human animals of the invention. According to the methods, an unrearranged human immunoglobulin heavy chain transgene is introduced into the germline of a non-human animal to produce a transgenic non-human animal having the transgene in its somatic and germ cells. Breeding of the human heavy chain transgenic animals with

- 16 -

transgenic non-human animals containing a human immunoglobulin light chain transgene produces transgenic non-human animals containing a human heavy chain transgene of the invention and a human light chain transgene. Either of the aforementioned transgenic non-human animals can be bred with animals having inactivated heavy and/or light chain loci to produce a transgenic non-human animal that produces a fully human antibody and is incapable of producing an endogenous antibody.

In one embodiment, a transgene of the invention is introduced into an embryonic stem (ES) cell which is then inserted into a blastocyst. The blastocyst with the ES cell containing the transgene of the invention is then surgically inserted into the uterus of the non-human animal to produce a chimeric non-human animal. The chimeric animal is bred to obtain germline transmission of the transgene of the invention to produce a transgenic, non-human animal having somatic and germ cells containing the transgene of the invention. Accordingly, a further aspects of the invention are an ES cell comprising a transgene of the invention and non-human animals having the transgene in some or all of its cells.

In still another aspect, the invention provides a method for producing high affinity, fully human antibodies of a desired isotype that are specific for an antigen of interest in a transgenic non-human animal of the invention. According to the method, a transgenic non-human animal of the invention is contacted with an antigen of interest under conditions that induce the production of an antibody by the B-cells of the animal. High affinity, fully human,

- 17 -

antigen-specific antibodies of the desired isotype can be collected from the blood stream of the transgenic non-human animal.

Alternatively, according to the methods of the invention, the antibody producing B-cells can be harvested from the animal and immortalized by any means known in the art, for the continuous production of antibodies. In one embodiment, the B-cells are fused with a mouse myeloma cell-line to produce antibody-secreting hybridomas. Such hybridomas can be screened to select those secreting high affinity, fully human, antigen-specific antibodies.

In a further aspect, the invention provides hybridomas derived from antibody producing B-cells harvested from a transgenic animal of the invention.

The antibodies of this invention may also be by the expression of B-cells expressing a desired antibody, by cloned human immunoglobulin genes, by phage-display, or by any other method known in the art.

20

- 18 -

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a schematic depiction of the yH1C and yH2Bm (or yH2Cm) yeast artificial chromosomes (YACs).

Figure 2 is a schematic depiction of the yH1C and yHG1
5 yeast artificial chromosomes (YACs).

Figure 3 is a schematic depiction of the yH1C and yHG1/2 yeast artificial chromosomes (YACs).

Figure 4 is a schematic depiction of the yH1C and yHG4/2 yeast artificial chromosomes (YACs).

10 Figure 5 is a schematic diagram of the targeting vectors (TV1 and TV4) for retrofitting yH1C to yHG1 and yHG4 YACs.

Figure 6 illustrates the construction of the targeting vectors for retrofitting yH1C to yHG1/2 and yHG4/2
15 YACs.

Figure 7 is a schematic diagram of the targeting vectors (TV G1/2 and TV G4/2) for retrofitting yH1C to yHG1/2 and yHG4/2 YACs.

Figure 8 shows Southern blot analyses of ES clones
20 fused with yH3B YAC (Clone Z 70.17.1)

- 19 -

DETAILED DESCRIPTION OF THE INVENTION

This invention relates to novel transgenes for the production of human immunoglobulin heavy chains of a desired isotype and to embryonic stem (ES) cells and transgenic non-human animals comprising the transgenes. This invention also relates to methods for producing such transgenic non-human animals and for producing fully human antibodies of a desired isotype in response to an antigen of interest in a transgenic animal of the invention.

The transgenes and transgenic non-human animals of the invention are useful in the production of fully human antibodies of various isotypes or classes. For therapeutic uses of such antibodies, the different effector functions of the individual antibody isotypes permits the use of a particular isotype to achieve a desired therapeutic effect. It is desirable, thus, to produce strains of transgenic non-human animals that produce antibodies of a single isotype following immunization with an antigen of interest.

In order that the invention herein described may be more fully understood, the following detailed description is set forth. In the description, the following terms are employed:

Gene regions - the DNA involved in producing or selecting a particular polypeptide chain; including promoters, enhancers, any switch regions preceding a constant gene as well as upstream and downstream preceding and following coding regions, and intervening sequences such as introns between coding segments or exons.

Gene segments - the coding segments in a multi-exon gene such as an immunoglobulin heavy chain

- 20 -

constant region. For example, the gene for the secreted form of the human immunoglobulin heavy chain gamma constant region contains 4 gene segments: CH1, H, CH2, and CH3.

5 Germline configuration - the arrangement and spacing of immunoglobulin gene segments before any somatic gene rearrangement has occurred.

Klenow Fragment - A large fragment of the enzyme Polymerase I, usually from *E. coli*. This
10 fragment does not contain any 5' to 3' exonuclease activity and only has polymerase activity. It can be used for end-filling of DNA molecules to create blunt ends.

Library - A mixture of cloned DNA fragments
15 usually propagated on DNA-based vectors, e.g., plasmids in bacteria, lambda bacteriophage in *E. coli*, P1 bacteriophage in *E. coli*, bacterial artificial chromosomes in *E. coli*, yeast artificial chromosomes in *Saccharomyces cerevisiae*, mammalian artificial
20 chromosomes in cultured cells, mammalian chromosome fragments in somatic cell hybrids.

Linker - Synthetic DNA fragments that are designed to contain restriction sites and other properties which can be added to larger DNA molecules,
25 e.g., to facilitate cloning and/or build back portions of DNA fragments encoding desired polypeptides.

Screening the Library - The process of searching for a specific sequence of cloned DNA in a library.

30 Sterile transcripts - Transcripts produced from the Ig loci thought not to be translated into required somatic gene segment rearrangement or class

- 21 -

switch recombination. In a B cell or a pre-B cell producing IgM there may be, for example, germline mRNA transcripts corresponding to the CH genes which potentially indicate which isotype the cell will switch to and produce.

Vector - A DNA molecule used to transport a foreign DNA into a host and being replicated in that host, to transform that host. available vectors include but are not limited to viruses (prokaryotes and eukaryotes), bacterial plasmids or artificial chromosomes.

Yeast artificial chromosomes (YACS) - cloning vehicles constructed from elements of yeast chromosomes which allow the vector to be replicated and maintained in yeast cells in vivo. Yeast elements include a centromere, an autonomous replication sequence, a pair of telomeres, yeast selectable markers, and usually a bacterial origin of replication and selectable marker for replication and selection of the YAC vector arms in bacteria. DNA inserts of up to at least 2000 kb can be cloned and maintained using YACs.

XENOMOUSE DEVELOPMENT

XenoMouse is a mouse which has inactivated mouse IgH and Igk loci and is transgenic for functional megabase-sized human IgH and Igk transgenes. The generation and characterization of XenoMouse has been described [See Mendez et al., *Genomics* 26:294-307 (1995); Mendez et al., *Nature Genetics*, 15, pp. 146-156 (1997); Green et al., *Nature Genetics* 7:13-21 (1994); International Patent application WO 94/02602, by Kucherlapati et al., published on February 3, 1994].

- 22 -

More particularly, there have been deletions of key elements of the mouse IgH and Igk loci by homologous recombination in mouse embryonic stem cells, followed by germline transmission of the mutations and

5 subsequent breeding to produce mice which are homozygous for both inactivated loci (DI mice). Such mice are incapable of making mouse IgH and Igk chains and display an arrest in B cell development in the bone marrow at the proB/preB-I stage. [Green et al, *Nature*

10 *Genetics*, 7, pp. 13-21 (1994); Green and Jakobovits, *J. Exp. Med.*, 188:483 (1998)]. The human IgH and Igk loci, cloned on yeast artificial chromosomes, were introduced into ES cells via yeast spheroplast-ES cell fusion [Jakobovits et al., *Nature* 362:255-258 (1993)].

15 After germline transmission and subsequent breeding onto the DI background, the human IgH and Igk YAC transgenes, yH1C and yK2, were able to functionally substitute for their murine counterparts and support B cell development. In addition, these mice produced

20 fully human IgMk and IgG2k antibodies, and ultimately, hybridomas secreting antigen-specific, high affinity fully human IgG2k monoclonal antibodies with therapeutic potential were generated.

THE yH1C TRANSGENE

25 The human IgH transgene, yH1C, is composed of 66 VH, all the D elements, all J elements, C μ and C δ , all regulatory elements, all in germline configuration. By using homologous recombination in yeast, the 3' end of yH1C was appended with a 22 kb fragment containing

30 the human γ 2 gene, including its switch regulatory elements, and a 4 kb fragment containing the mouse 3'

- 23 -

enhancer element [See Mendez et al., *Nature Genetics* 15:146-156 (1997) the disclosure of which is hereby incorporated by reference]. The left YAC arm carries expression cassettes for the yeast selectable marker ADE2, the mammalian selectable marker and the right YAC arm carries expression cassettes for the yeast selectable marker LYS2 and a mammalian selectable marker Neo, encoding resistance to the drug, G418. The latter is non-functional in ES cells as its promoter, MMT (mouse metallothionine), is probably non-functional in ES cells. In other cell types, the MMT promoter drives transcription at only very low levels, if at all, under normal physiological conditions and requires heavy metals, e.g., Cd, for higher level transcription. Indeed, ES cells transfected with this construct never became resistant to even low levels of G418.

B-CELL DEVELOPMENT

B cell development initiates in the bone marrow with a deletional recombination between a D and J gene. Subsequently, a V gene recombines with the DJ to make a VDJ, which is transcribed, producing a spliced VDJC μ transcript. If the transcript is in-frame, then a μ chain is synthesized upon translation. Similarly, and generally after V_HDJ_H recombination and successful pairing of the μ chain with surrogate light chain, the Ig light chain loci rearrange their V and J gene segments. Successful B cell development in the bone marrow results in B cells expressing IgM κ or IgM λ on the cell surface. In the mouse, 95% of the B cells express IgM κ ; in the human, approximately 60% of the B cells express IgM κ .

- 24 -

These IgM producing B cells form the primary immune repertoire and perform immune surveillance for recognition of foreign antigens. In the mouse or in humans, these IgM producing B cells can subsequently
5 undergo isotype class-switching from IgM to the IgG or IgA, or IgE isotypes. The frequency of class switching increases during an immune response. Mice and humans each have genes for four different isotypes of IgG. They are IgG1, IgG2a, IgG2b, and IgG3 in the mouse, and
10 IgG1, IgG2, IgG3, IgG4 in the human. Humans have two IgA isotypes, IgA1 and IgA2, and one IgE isotype. In a mouse, there is, on average, 6500, 4200 and 1200 µg/ml of IgG1, IgG2a, and IgG2B respectively, and 260 µg/ml IgA. In the human, of the total IgG, about 70% is
15 IgG1, 18% is IgG2, 8% is IgG3 and 3% is IgG4. In the total IgA in humans, about 80% is IgA1 and 20% is IgA2.

EFFECTOR FUNCTIONS OF ANTIBODIES

Different isotypes have different effector functions. Such differences in function are reflected
20 in distinct 3-dimensional structures for the various immunoglobulin isotypes [P.M. Alzari et al., Annual Rev. Immunol. 6:555-580 (1988)]. For example, the human IgG1 and IgG3 isotypes are involved in complement mediated-lysis or antibody-dependent cellular
25 cytotoxicity (ADCC) and the IgG2 and IgG4 have little or no known effector functions. [Snapper and F.D. Finkelman, Fundamental Immunology 3d Ed., pp. 837-863]. Since different effector functions are associated with different IgG isotypes, it is therefore desirable to be
30 able to select the isotype and the binding specificity of the mAb to produce optimal therapeutic benefit. For

- 25 -

example, if a mAb is desired to neutralize a cytokine response or block the activity of a receptor, then a mAb lacking effector functions, such as an IgG2 or an IgG4 might be desired. On the other hand, if the

5 killing of a cell via binding of a mAb to an antigen on the cell surface is desired, then a mAb such as an IgG1, with its specific effector functions, either ADCC or CML, is desired. Thus, a transgenic mouse engineered for the generation of fully human monoclonal

10 antibodies would be desirable to control the isotype of the resulting monoclonal antibodies. In this case, one could select a particular antibody isotype by immunizing a particular transgenic mouse strain which produces only the desired human antibody isotype. Such

15 mice would ensure that any resulting antigen-specific IgG mAbs would possess the desired effector functions. This would preclude subsequent re-engineering of the antibody gene to change the constant region including the isolation (cloning) of the variable region and the

20 ligation of said VH region to the desired CH gene.

In one embodiment of the present invention, the sole C γ gene on the yH1C human IgH YAC, C γ 2, is replaced by another CH gene. For example, instead of the 22 kb fragment carrying the complete human C γ 2 gene

25 other inserts carrying human CH genes could be cloned into the targeting vector of Mendez et al. [See Mendez et al., *Nature Genetics* 15:146-156 (1997)]. The human C γ 1-4 genes have been sequenced and can be isolated from bacteriophage lambda libraries of human genomic

30 DNA and subsequently recovered on EcoRI fragments of about 20-25 kb [See J.W. Ellison et al., *Nucleic Acids Res.*, 13:4071-4079 (1982); J. Ellison et al., *Proc.*

- 26 -

Natl. Acad. Sci. USA, 79:1984-1985 (1982); S. Huck et al., *Nucleic Acids Res.*, 14:1779-1789 (1986); J. Ellison et al., *DNA*, 1:11-18 (1981) the disclosures of which are hereby incorporated by reference].

- 5 Similarly, the sequences for mouse Cy1, Cy2a, Cy2b, and Cy3 are all known [See H. Hayashida et al., *EMBO Journal*, 3:2047-2053 (1984) the disclosure of which is hereby incorporated by reference].

CLASS SWITCHING

- 10 Class switch recombination (CSR) from IgM to IgG, IgA or IgE is mediated through a deletional recombination event occurring between tandem directly repetitive switch regions present 5' of all IgH constant region genes except C δ . Switch regions are
15 known to be composed of the I promoter, the I exon and a set of direct repeats flanked by inverted repeat sequences. Enhancers and cytokine response sequences are known to lie in the region near the I promoter. At least one transcriptional enhancer, located immediately
20 3' of the downstream inverted repeat, in the mouse Cy1 gene has been hypothesized [J.P. Manis et al., *J. Exp. Med.* 188:1421-1431 (1998)]. Also required is iEm, an enhancer located between JH and Cm. Transcription initiates at the I promoter and proceeds through the I
25 exon to the end of the C gene. This transcript is processed to yield a non-coding sterile transcript with the I exon spliced to the CH exons. Transcription through the switch region is required for class switch recombination. The human and mouse S μ and S γ regions
30 have been sequenced, the sequences of which are publicly available from the Genbank database.

- 27 -

In the mouse, different combinations of lymphokines and activators have profoundly different effects on class switching from IgM to individual CH genes. For example, *in vitro* the combination of LPS and interleukin-4 induces class switching to IgG1 and IgE and suppresses switching to IgG2b and IgG3. Other lymphokines affecting CSR include but are not limited to, IL-5, TGF- β , interferon- γ . These lymphokines are secreted *in vivo* by helper cells such as the antigen presenting T- and follicular dendritic cells in the germinal centers of secondary lymphoid tissues. These lymphokines modulate transcription of their responding CH genes prior to CSR, probably through activation of the corresponding I promoter. For example, the IL-4 response element in the mouse C γ 1 I promoter has been mapped. [Rothman et al., *Int. Immunol* 2, pp. 621-627 (1990).] The lymphokine responsiveness of the human switch regions is not yet as well-characterized as that of the mouse. However, the different human switch (S) regions may also have different responses to different lymphokines and activators. This may in part be the source of the different levels of the IgG subclasses in human serum.

NON-COGNATE SWITCHING

In view of the real and possible differential responsiveness of mouse and human S regions, respectively, to lymphokines and other activators, it is desirable to have heterologous switch regions controlling CSR in human antibody producing transgenic mice. For example, Ig γ 1 is the most abundant class of IgG in the mouse. It is known also that CSR can occur

- 28 -

from human $S\mu$ regions to mouse $S\gamma 1$. [Taylor et al.,
Int. Immunol., 6, pp. 579-591 (1994).] Using standard
tools of molecular biology and the well-characterized
and cloned mouse $S\gamma 1$ sequence [Mowatt and Dunnick, *J.*
5 *Immuno.*, 136, pp. 2674-2683 (1986), Genbank accession #
M12389], it is possible to engineer a DNA vector having
the mouse $S\gamma 1$ functionally linked to a human CH coding
sequence, e.g., human $C\gamma 1$. Included downstream of the
human CH coding sequences would be a sequence
10 encompassing the mouse 3' enhancer. The m3'E sequence
could be a 4 kb *Xba*I fragment or a 900 bp *Stu* I
fragment, both of which encompass the core DNase I
hypersensitive sites, HS1,2. [Dariavach et al., *Eur. J.*
Immunol. 21, pp. 1499-1504 (1991); Petterson et al.,
15 *Immunobiol.*, 198, pp. 236-248 (1997)]. By having 5' and
3' flanking homology to $\gamma H1C$ and an appropriate
selectable marker, such a vector can be recombined *in*
vivo in yeast to replace the human $C\gamma 2$ gene on $\gamma H1C$. A
YAC engineered in this way would retain intact all of
20 the VH , D_H , J_H , C_μ , and C_δ of $\gamma H1C$, but would have a
chimeric CH gene: the mouse $S\gamma 1$ elements would control
switching from human IgM to the downstream human CH
coding sequences.

In another embodiment, the human $C\gamma 2$ coding
25 sequences, including all of the exons for the secreted
and membrane-bound forms of the C_H gene are replaced by
another human C_H gene. In this way, the human $S\gamma 2$
sequences control CSR from C_μ to the downstream C_H gene.
It is known that the hSg2 sequences are stable in $\gamma H1C$
30 while other human S sequences, some of which have
longer tandem arrays of S repeats may be less stable.
It is also known that CSR in transgenic mice with the

- 29 -

human $\text{Cy}2$ gene is efficient and generates high serum levels of human IgG2 and results in efficient production of fully human IgG2 mAbs. Thus, it may be preferable to retain the human $\text{Sy}2$ with their favorable stability and *in vivo* response to antigen challenge while engineering CSR to occur to another isotype, e.g., either $\text{Cy}1$ or $\text{Cy}4$. To accomplish this, a vector with the following elements would be constructed: 5' homology located between human $\text{Sy}2$ and the human $\text{Cy}2$ coding exon 1, a human CH gene other than $\text{Cy}2$, the mouse 3' enhancer, a yeast selectable marker, and 3' targeting homology in the YAC arm for example. Such a vector would be introduced into yeasts carrying yH1C and targeted recombinants would be selected and screened. It should be understood that in these examples many variations can be created by one skilled in the art and that these examples are not meant to indicate that these are the only means to achieve the end of a transgenic mouse having CSR driven by heterologous S regions.

THE ROLE OF ENHANCERS

In addition to S regions, other *cis* regulatory elements are known to be or may be required for CSR. The requirement for iEm has been mentioned. Also, an enhancer required for expression of normal levels of IgG has been hypothesized to be between the 3' inverted repeat of mouse $\text{Sy}1$ and the $\text{C}_{\text{H}}1$ exon. This enhancer could be conserved in other C_{H} genes in the mouse and humans and this interval should be retained in any vector designed for CSR via heterologous switch sequences. [Elenich et al., *J. Immunol.* 157, pp. 176-

- 30 -

182 (1996); Cunningham et al., *Int. Immunol.*, 10, pp. 1027-1037 (1998)]. Also important is a cluster of enhancers 3' of the C α gene in mouse and humans. In the mouse the 40 kb region downstream of C α contains
5 four enhancer elements, hallmarks of which are Dnase I hypersensitive sites (HS). These enhancers are in 5' to 3' order: HS3a, 4 kb downstream of C α ; HS1,2 (known in the literature and in this application as m3'E), 15 kb 3' of C α ; HS3b 25 kb 3' of C α ; and HS4, ca; 30 kb 3'
10 of C α ; HS1,2, HS3a, and HS3b enhance expression in activated B cells and plasma cells. HS4 is active over the course of B cell development, but is apparently dispensable as the yH1C YAC lacks HS4 and yet supports efficient B cell development in mice. Together, these
15 elements can act synergistically to enhance transcription and are hypothesized to form a locus control region (LCR) for the IgH locus in mouse and humans. It has been hypothesized that there is some redundancy of function of the individual HS units. The
20 unimpaired activity of these elements may be required for CSR although HS1,2 and HS3a are separately dispensable CSR [See J.P. Manis et al., *J. Exp. Med.* 188:1421-1431 (1998)].

HS1,2 (3'E) was the first discovered enhancer
25 of this set. The HS1,2 sites and sequences homologous to consensus binding domains for transcription factors such as AP-1 can be isolated on a 900 bp Stu-I fragment. [Dariavich et al., *Eur. J. Immunol.*, 21, pp. 1499-1504 (1991); Genbank accession #X62778]. The 3'E
30 in the mouse is oriented opposite to the 3'E of the rat, suggesting that like other enhancers, its function is orientation independent. However, the 3'E has been

- 31 -

shown to have position dependent activity and enhance transcription more effectively when positioned further from the promoter. *Gene*, 136, pp. 349-353 (1993). It is known that a 4 kb XbaI fragment encompassing the 900 bp StuI fragment with HS1,2, positioned 3' of the human Cg2 gene, can support CSR and high level expression of IgG2 in transgenic mice [See Mendez et al., *Nature Genetics* 15:146-156 (1997)].

The insertion of a strong promoter (PGK) into the mouse IgH 3' LCR can abrogate class switching to some IgG isotypes (IgG2a, IgG3, IgG2b) and lower expression of others (IgG1, IgA). Curiously, the promoter and its expressed gene also come under the control of the LCR: the PGK-driven expression construct is down-regulated and can be up-regulated in activated B cells [J.P. Manis et al., *J. Exp. Med.* 188:1421-1431 (1998)]. Thus, in YAC transgenes carrying a β -gal expression construct driven by the strong constitutive promoter, PGK, with construct 3' and adjacent to the mouse 3'E core construct (900 bp StuI), it may be advantageous to screen for YACs that have integrated into the ES cell genome with concomitant loss of the PGK- β -gal construct. This can be accomplished by PCR using primers for β -gal, or by Southern blots probed with the β -gal gene.

IMMUNOGLOBULIN MEMBRANE EXONS

Two forms of each IgH isotype and class, secreted (s) and membrane (m), can be made by a B cell.

Ig(s) and Ig(m) are synthesized through alternative splicing of IgH transcripts. Two membrane exons lie 2 kb downstream of the CH3 exon of each human IgG gene.

- 32 -

Encoded by the membrane exons are a hydrophobic transmembrane sequence and a short approximately 3 amino acid cytoplasmic tail. Alternative splicing from CH3 to the first membrane exon results in membrane bound IgG. The membrane bound Ig interacts with other proteins in the B cell, e.g., Ig α , Ig β and CD45, among others, to form a complex called the B cell receptor (BCR) that is capable of signal transduction. Binding of antigen by the V region of the IgG displayed in the extracellular environment, e.g., soluble or on antigen presenting cells, can lead to signal transduction. This signal transduction by the BCR leads to activation of the B cells and ultimately efficient affinity maturation and germinal center formation in the secondary immune response.

Additionally, binding of antigen by the Ig of the BCR may lead to internalization, processing of and presentation of antigen fragments by MHC molecules for presentation to helper cells. Clearly, efficient assembly of a functional BCR is required for an efficient primary and secondary immune response.

The human IgG1 membrane exons may not complex well with the other components of the BCR, resulting in a chimeric BCR that may not signal as efficiently as that of the mouse. G. Pluschke et al., *J. Immunolog. Methods*, 215, pp. 27-37 (1998). A human IgG1 construct with all of the human exons encoding the secreted and membrane forms of IgG1 was inserted into the mouse IgG2a locus such that all of the mouse Cy2a exons were replaced and CSR to the human coding exons was under the control of the mouse Sy2a region. Chimeric human IgG1 (mouse VDJ-human IgG1) was expressed at levels

- 33 -

100x less than mouse IgG2a and antigen specific mAbs were not recovered. Thus, although class switching driven by the mouse Sy2a did occur, the normal immune response was compromised. Alternatively, exons coding for secreted human IgG1 have been used to replace only the exons encoding the secreted form of mouse IgG1. This construct produced a chimeric IgG1 heavy chain gene that contained all of the human exons for secreted IgG1 but with the downstream mouse membrane exons intact. Class switching would have been driven by the mouse Sy1 regions. Membrane bound Ig would be mouse V-human $\gamma 1$ CH1-CH3-mouse C $\gamma 1$ (mem). In this transgenic mouse, the serum levels of human IgG1 were equivalent to mouse IgG1 in normal naïve mice. Thus, the mouse Sy1 can drive efficient class switching and the mouse IgG1 membrane exons can function with at least the human $\gamma 1$ CH1-CH3 exons. The authors did not test the mice for production of antigen-specific mAbs. As in the previous construct, the resulting IgG1 mAbs would have been chimeric: mouse VDJ functionally linked to the secreted form of human C $\gamma 1$.

Given these results, an intact set of human C $\gamma 1$ exons, coding both the secreted and membrane forms of C $\gamma 1$, and functionally linked to a human IgH locus (V_H, D_H, J_H, C μ , C δ and S γ regions) may function sub-optimally because of inefficient assembly of the membrane-bound human IgG1 may not yield a fully functional BCR. Thus, it may be preferable to replace the human C $\gamma 1$ membrane exons with those from another isotype known to assemble efficiently into a functional BCR. Such exons may include the mouse C $\gamma 1$ exons or other murine C membrane exons. Alternatively, the human C $\gamma 2$ membrane exons would be expected to function

- 34 -

well in the BCR of the mouse because the XenoMouse G2 has high levels of secreted IgG2 and produces high affinity antigen-specific mAbs efficiently. Thus, the human Cy2 membrane exons could be functionally linked
5 to the human Cy1 CH1-CH3 exons. The sequence for the membrane exons is known (X52847 for hyl; AB006775 for hy2).

VECTOR CONSTRUCTION

In one embodiment, a targeting vector is
10 generated to introduce only the CH1-CH3 exons into the yH1C YAC. The sequence of all of the human CH1-CH3 as well as introns and flanking DNA is available [See J.W. Ellison et al., *Nucleic Acids Res.*, 13:4071-4079 (1982); J. Ellison et al., *Proc. Natl. Acad. Sci. USA*,
15 79:1984-1985 (1982); S. Huck et al., *Nucleic Acids Res.*, 14:1779-1789 (1986); J. Ellison et al., *DNA*, 1:11-18 (1981) the disclosures of which are hereby incorporated by reference], allowing all restriction sites to be mapped electronically and a targeting
20 vector to be constructed. One such vector would contain 5' homology upstream of human Cy2 CH1 exon, an expression construct for a positive/negative selectable marker in yeast (URA3), a direct repeat of the 5' targeting homology, sequence containing the human Cy1
25 exons CH1-CH3, and 3' targeting homology. This vector would be transfected into yeasts carrying yH1C and homologous recombinants positively selected on plate lacking uracil and then screened by Southern blot hybridization or PCR to test for the loss of human Cy2
30 CH1-CH3 exons and the concomitant gain of human Cy1 CH1-CH3. Once identified, the deletion of the URA3

- 35 -

gene can be selected with 5'-florouracil. Such loss would be expected to occur at high frequency (10^{-4} - 10^{-5}) because of efficient intra-chromosomal recombination between direct repeat sequences in yeast. Deletion of the URA3 gene restores a fully human IgH in a configuration functional for class switch recombination and the expression of antibodies. It is obvious that there are other strategies for accomplishing such engineering. Also, there may be motivation to engineer other human Cy genes, e.g., human Cy4, into the human Cy2 locus.

CRE-LOX MEDIATED CLASS SWITCHING

The CRE-lox system allows the targeted insertion of DNA into pre-defined sites. Derived from P1 bacteriophage, the CRE recombinase drives intra-DNA or inter-DNA recombination between loxP sites [B. Sauer et al., *New Biologist* 2:441-449 (1990); S. Fukushige et al., *Proc. Natl. Acad. Sci. USA*, 89:7905-7909 (1992); Y.-R.Zou et al., *Current Biol.*, 4:1099-1103 (1994)]. A lox P site (sequence:TA ACT TCG TAT AGC ATA CAT TAT ACG AAG TTA TA (SEQ ID NO: 1)) is introduced into the DNA of a yH YAC. The sequence is positioned 3' of the 3' inverted repeat of the downstream S region, e.g., Sg2, and 5' of the splice acceptor sequence of the CH1 exon of the downstream Cg gene, e.g., Cg1. The lox P site can be inserted directly into the YAC via homologous recombination in yeast, or it can be incorporated into a larger targeting vector, such as the ones described earlier in this description. When incorporating the site into such YAC targeting vectors, the loxP site can be introduced on a PCR primer used for amplifying the

- 36 -

targeting homology, e.g., 5' targeting homology, or can be inserted as an oligonucleotide ligated *in vitro*. As homologous recombination between 2 loxP sites is orientation dependent, it is important to note the orientation in which the first loxP site is inserted into the YAC.

In the second phase, a plasmid vector for insertion of the alternative C γ gene is generated. At the core of this vector is a cassette carrying the C γ gene to be introduced and a loxP site to enable the introduction: this cassette starts with a lox P site in the same 5'-3' orientation as in the YAC, followed by the DNA upstream of CH1, corresponding to the site of lox P insertion upstream of the C γ on the YAC, and continuing in germline configuration through CH1, with the CH1 exon splice acceptor intact, through downstream of the polyadenylation site 3' of the second membrane exon. For example, an approximately 7 kb Hind III fragment will capture all of the required DNA for all human C γ genes. Alternatively, only the CH1-CH3 exons including appropriate 3'signals for transcription and translation (untranslated region, polyadenylation site) could be used to generate only the secreted form of the mAb. To abrogate possible read through transcription, a eucaryotic transcriptional terminator sequence can be appended downstream of the CH gene on the vector. To facilitate selection of transformants, an expression cassette for a selectable marker such as puromycin or hygromycin may be appended downstream of the CH gene.

Once hybridomas are generated from the transgenic mouse carrying the yH transgene engineered with the loxP site, CRE-lox mediate class switching can

- 37 -

be induced by co-transfecting, e.g., by electroporation or lipofection, the circularized insertion vector, and either purified CRE recombinase or a CRE expression vector. In co-transfected cells, CRE will mediate
5 insertion of the novel CH gene into the locus, where it would be transcribed and spliced *in cis* to the upstream VHDJH encoding the desired mAb specificity. The transcriptional terminator would preclude run on transcription into the downstream CH gene. If the
10 vector has a selectable marker, then transfected hybridomas can be selected with the appropriate drug, and then pools or individual clones screened by ELISA for mAbs of the desired novel isotype. If the vector lacks a selectable marker, then pools of transfected
15 hybridomas can be screened by ELISA and hybridomas producing the desired isotype can be subcloned from the pool. If the replacement CH gene encodes membrane bound IgH also, then the hybridomas can be screened and sorted by flow cytometry.

20 In some instances, it may be preferable to possess two different isotypes of a single antigen-specific mAb, with one isotype having one activity, such as ADCC or CML, and the other isotype lacking effector function, but with identical antigen-binding
25 characteristics such as epitope specificity and affinity. This goal could be achieved by molecularly cloning the variable regions of the heavy chain and light chain and then functionally linking them to the appropriate constant regions, followed by transfection
30 into cells for production of the mAb. However, this process can be labor and time intensive. Alternatively, using the CRE-lox process described

- 38 -

above, the monoclonal antibody can be efficiently class-switched *in vivo* in the hybridoma.

Mouse Strains

The following mouse strains are described
5 and/or utilized herein:

Double Inactivated (DI) Strain

The DI strain of mice are mice that do not produce functional endogenous mouse Ig. In preferred embodiments, the DI mice possess an inactivated mouse J_H
10 region and an inactivated mouse C_k region. The construction of this strain is discussed extensively elsewhere. For example, the techniques utilized for generation of the DI strains are described in detail in U.S. Patent Application Serial Nos. 07/466,008, filed
15 January 12, 1990, 07/610,515, filed November 8, 1990, 07/919,297, filed July 24, 1992, 08/031,801, filed March 15, 1993, 08/112,848, filed August 27, 1993, 08/234,145, filed April 28, 1994, 08/724,752, filed October 2, 1996. See also European Patent No., EP 0
20 463 151 B1, grant published June 12, 1996, International Patent Application No., WO 94/02602, published February 3, 1994, International Patent Application No., WO 96/34096, published October 31, 1996, and PCT Application No. PCT/US96/05928, filed
25 April 29, 1996. The disclosures of each of the above-cited patent and patent applications are hereby incorporated by reference in their entirety. DI mice possess a very immature B-cell development. The mice do not produce mature B-cells, only pro-B-cells. [Green
30 and Jakobovits, *J. Exp. Med.*, 188, pp. 483-495 (1998)].

- 39 -

XenoMouse I Strain

The design, construction, and analysis of the XenoMouse I strain was discussed in detail in Green et al., *Nature Genetics*, 7:13-21 (1994). Such mice
5 produced human IgM κ antibodies against a DI background. The mice showed improved B-cell function when compared to the DI strain of mice which have little to no B-cell development. While XenoMouse I strains of mice mount a sizeable immune response to antigenic challenge, their
10 production of B-cells was only 20-25% of wild-type mice and they possessed a limited response to antigens. Both characteristics appear to be related to their limited V-gene repertoire.

L6 Strain

15 The L6 strain is a mouse producing human IgM κ antibodies against a DI background of endogenous mouse Ig. L6 mice contain an inserted human heavy chain and an inserted human kappa light chain. The L6 strain is generated through breeding of a mouse containing a
20 heavy chain insert against a double inactivated background (L6H) and a mouse having a kappa light chain insert against a double inactivated background (L6L). The heavy chain insert comprises an intact approximately 970 kb human DNA insert from a YAC
25 containing approximately 66 V $_H$ segments, starting at V $_H$ 6-1 and ending at V $_H$ 3-65, and including the major D gene clusters (approximately 32), J $_H$ genes (6), the intronic enhancer (Em), C μ , and through about 25 kb past C δ , in germline configuration. The light chain
30 insert, yK2, comprises an intact approximately 800 kb human DNA insert from a YAC which contains

- 40 -

approximately 32 V_k genes starting at V_{k-B3} and ending at V_{k-Op11} . The 800 kb insert contains a deletion of approximately 100 kb starting at $V_{k-Lp-13}$ and ending at V_{k-Lp-5} . However, the DNA is in germline configuration
5 from $V_{k-Lp-13}$ to 100 kb past V_{k-Op-1} , and also contains the J_k genes, the intronic and 3' enhancers, the constant C_k gene, and Kde . [Mendez et al., *Nature Genetics*, 15, pp. 146-156 (1997)]. Furthermore, L6 mice exhibit predominant expression of human kappa light chain, a
10 large population of mature B-cells, and normal levels of IgM_k human antibodies. [Green and Jakobovits, *J. Exp. Med.*, 188, pp. 483-495 (1998)].

XenoMouse IIa Strain:

The XenoMouse IIa mice represent second
15 generation XenoMouse™ strains equipped with germline configuration megabase-sized human Ig loci, against a DI background, such that the mice do not produce functional endogenous Ig. Essentially, the mice are equivalent in construction to the L6 strain, but
20 additionally include the human $C\gamma 2$ gene with its entire switch and regulatory sequences and the mouse 3' enhancer *in cis*. The mice contain an approximately 1020 kb heavy and an approximately 800 kb kappa light chain loci, which include the majority of the human
25 variable region genes, including heavy chain genes (approximately 66 V_H) and kappa light chain genes (approximately 32 V_k), human heavy constant region genes (μ , δ , and γ) and kappa constant region genes (C_k), and all of the major identified regulatory elements. These
30 mice have been shown to access the full spectrum of the variable genes incorporated into their genome.

- 41 -

Furthermore, they exhibit efficient class switching and somatic hypermutation, predominant expression of human kappa light chain, a large population of mature B-cells, and normal levels of IgM_k and IgG_k human antibodies. Such mice mount a vigorous human antibody response to multiple immunogens, including human IL-8, human EGF receptor (EGFR), and human tumor necrosis factor- α (TNF- α), ultimately yielding antigen-specific fully human mAbs with sub-nanomolar affinities. This last result conclusively demonstrates XenoMouse™ as an excellent source for rapid isolation of high affinity, fully human therapeutic mAbs against a broad spectrum of antigens with any desired specificity.

As will be appreciated from the above introduction, the XenoMouse II strain appears to undergo mature B-cell development and mount powerful adult-human-like immune responses to antigenic challenge. The L6 strain also appear to undergo mature B-cell development. When XenoMouse II strains, a markedly different B-cell development profile is observed. Owing to this difference, it appears that the quantity and complexity of variable region sequences introduced into the animals are essential to the induction of B-cell maturation and development and the generation of an adult-human-like immune response. Thus, in addition to the strains' utility to generate human antibodies, the strains provide a valuable tool for studying the production and function of human antibodies in the normal immune response, as well as the abnormal response characteristic of autoimmune disease and other disorders.

- 42 -

Variable Region - Quantitative Diversity

It is predicted that the specificity of antibodies (i.e., the ability to generate antibodies to a wide spectrum of antigens and indeed to a wide spectrum of independent epitopes thereon) is dependent upon the variable region genes on the heavy chain (V_H) and kappa light chain (V_K) genome. The human heavy chain genome includes approximately 95 V_H genes of which 41 are functional genes which encode variable regions of the human heavy chain of immunoglobulin molecules. In addition, the human light chain genome includes approximately 40 V_K genes on its proximal end of which 25 are functional which encode variable regions of the human kappa light chain of immunoglobulin molecules. We have demonstrated that the specificity of antibodies can be enhanced through the inclusion of a plurality of genes encoding variable light and heavy chains.

Provided in accordance with the present invention are transgenic mice having a substantial portion of the human Ig locus, preferably including both a human heavy chain locus and a human kappa light chain locus. In preferred embodiments, therefore, greater than 10% of the human V_H and V_K genes are utilized. More preferably, greater than about 20%, 30%, 40%, 50%, 60%, or even 70% or greater of V_H and V_K genes are utilized. In a preferred embodiment, heavy and light chain constructs that include 32 genes from the proximal region of the human V_K light chain genome and/or 66 genes from the V_H portion of the human IgH locus, respectively, are utilized. As will be appreciated, genes may be included either sequentially, i.e., in the order found in the human genome, or out of

- 43 -

sequence, i.e., in an order other than that found in the human genome, or a combination thereof. Thus, by way of example, an entirely sequential portion of either the human V_H or human V_K region of the locus can
5 be utilized, or various V genes in either the V_H or V_K genome can be skipped while maintaining an overall sequential arrangement, or V genes within either the V_H or V_K genome can be reordered, and the like. In a preferred embodiment, the entire human loci are
10 inserted in the mouse genome in substantially germline configuration as found in humans. In any case, it is expected and the results described herein demonstrate that the inclusion of a diverse array of genes from the V_H and V_K genome leads to enhanced antibody specificity
15 and ultimately to enhanced antibody affinities.

Such mice preferably further include the entire human D_H region, the entire human J_H region and the human mu constant region, and can additionally be equipped with other human constant regions for the
20 coding and generation of additional isotypes of antibodies. Such isotypes can include genes encoding γ_1 , γ_2 , γ_3 , γ_4 , α , ϵ , and δ and other constant region encoding genes with appropriate switch and regulatory sequences. As will be appreciated, and as discussed in
25 more detail below, a variety of switch and regulatory sequences can be utilized in connection with any particular constant region selection.

The following Table indicates the diversity of antibody combinations that are possible in humans,
30 based strictly on random V-D-J joining and combination with kappa light chains, without consideration of N-addition, deletions or somatic mutation events. Based on these considerations, there are greater than 7

- 44 -

$\times 10^5$ possible antibody combinations in humans, of any particular isotype.

TABLE 1

	Region	Heavy Chain	Kappa Light Chain
5	Functional Variable "V"	~41	25
	Functional Diversity "D"	≥ 23	--
	Joining "J"	6	5
10	Combinations (VxDxJ)	5,658	125
15	Total Combinations (HC Combinations x LC Combinations)	7.1×10^5	

In connection with a preferred embodiment of the invention, through the inclusion of about 34 functional V_H genes and 18 functional V_K genes in a mouse with a full complement of D_H , J_H , and J_K genes, the possible diversity of antibody production is on the order of 4.2×10^5 different antibodies. As before, such calculation does not take into account N-addition or somatic mutation events. Therefore, it will be appreciated that mice in accordance with the invention, such as the L6 and the XenoMouse II strains, offer substantial antibody diversity. In preferred embodiments, mice are designed to have the capability of producing greater than 2×10^5 different heavy chain V-D-J combinations

- 45 -

and kappa light chain V-J combinations, without accounting for N-additions or somatic mutation events.

Variable Region - Qualitative Diversity

In addition to quantitative diversity,
5 quantitative selection of V-genes (i.e., large numbers of diverse V-genes) and/or qualitative selection of V-genes (i.e., selection of particular V-genes) appears to play a role in what we refer to herein as "qualitative diversity." Qualitative diversity, as used
10 herein, refers to diversity in V-D-J rearrangements wherein junctional diversity and/or somatic mutation events are introduced. During heavy chain rearrangement, certain enzymes (RAG-1, RAG-2, and possibly others) are responsible for the cutting of the
15 DNA representing the coding regions of the antibody genes. Terminal deoxynucleotidyl transferase (Tdt) activity, which is responsible for N-terminal additions of nucleotides between the V-D and D-J gene exons is up-regulated. Similar enzymes and others (SCID and
20 other DNA repair enzymes) are responsible for the deletion(s) that occurs at the junctions of these coding segments. Junctional diversity refers to both N-addition events and formation of the complementarity determining region 3 (CDR3) . As will be appreciated,
25 CDR3 is located across the D region and includes the V-D and D-J junctional events. Thus, N-additions and deletions during both D-J rearrangement and V-D rearrangement are responsible for CDR3 diversity.

The junctional diversity created by
30 N-additions and CDR3 additions play a clear role developing antibody specificity.

- 46 -

In accordance with the invention, rearranged V-D-J gene sequences show N-addition lengths that are comparable to expected adult-human N-addition lengths. Further, amino acid sequences across the open reading
5 frame (ORF) corresponding to CDR3 sequences show CDR3 lengths that are comparable to expected adult-human CDR3 lengths. Such data is indicative that quantitative variable region diversity and/or qualitative variable region diversity results in
10 human-like junctional diversity. Such junctional diversity is expected to lead to a more human-like antibody specificity.

Variable Region - Affinities

While we have not conclusively demonstrated a
15 direct causal connection between the increased variable region inclusion and antibody specificity, it appears, and it is expected that through providing such diversity, the ability of the mouse to mount an immune response to a wide array of antigens is possible and
20 enhanced. Additionally, such mice appear more equipped to mount immune responses to a wide array of epitopes upon individual antigens or immunogens. From our data it also appears that antibodies produced in accordance with the present invention possess enhanced affinities.
25 Such data includes comparisons between mice in accordance with the invention and the XenoMouse I strains, as well as consideration of the published results of GenPharm International and the MRC. In connection with the XenoMouse I strains, as mentioned
30 above, such mice possessed sub-normal B-cell production and a only limited response to antigens. Such result appeared related in part to the limited V-gene

- 47 -

repertoire. Similarly, results reported by GenPharm International and the MRC indicate a limited response to diverse antigens.

Without wishing to be bound to any particular theory or mode of operation of the invention, it would appear that enhanced affinities appear to result from the provision of the large number and complexity of V regions. From our data, the provision of greater numbers and/or selection of qualities of V-gene sequences, enhances junctional diversity (N-additions and formation of complementarity determining region 3 ("CDR3") diversity), which is typical of an adult-human-like immune response, and which play a substantial role in affinity maturation of antibodies. It may also be that such antibodies are more effective and efficient in somatic mutation events that lead to enhanced affinities. Each of junctional diversity and somatic mutation events are discussed in additional detail below.

With respect to affinities, antibody affinity rates and constants derived through utilization of plural V_H and V_K genes (i.e., the use of 32 genes on the proximal region of the V_K light chain genome and 66 genes on the V_H portion of the genome) results in association rates (k_a in $M^{-1} S^{-1}$) of greater than about 0.50×10^{-6} , preferably greater than 2.00×10^{-6} , and more preferably greater than about 4.00×10^{-6} ; dissociation rates (k_d in S^{-1}) of greater than about 1.00×10^{-4} , preferably greater than about 2.00×10^{-4} , and more preferably greater than about 4.00×10^{-4} ; and dissociation constant (in M) of greater than about 1.00×10^{-10} , preferably greater than about 2.00×10^{-10} , and more preferably greater than about 4.00×10^{-10} .

- 48 -

Preferably, such mice additionally do not produce functional endogenous immunoglobulins. This is accomplished in a preferred embodiment through the inactivation (or knocking out) of endogenous heavy and light chain loci. For example, in a preferred embodiment, the mouse heavy chain J-region and mouse kappa light chain J-region and C_κ- region are inactivated through utilization of homologous recombination vectors that replace or delete the region.

VARIABLE REGION - B-CELL DEVELOPMENT

B-cell development is reviewed in Klaus B *Lymphocytes* (IRL Press (1990)) and Chapters 1-3 of T. Honjo *et al.*, *Immunoglobulin Genes* (Academic Press Ltd. San Diego, CA (1989)). Generally, in mammals, blood cell development, including B- and T-cell lymphocytes, originate from a common pluripotent stem cell. The lymphocytes, then, evolve from a common lymphoid progenitor cell. Following an early gestational period, B-cell initiation shifts from the liver to the bone marrow where it remains throughout the life of the mammal.

In the life cycle of a B-cell, the first generally recognizable cell is a pro-pre-B-cell which is found in the bone marrow. Such a cell has begun heavy chain V-D-J rearrangement, but does not yet make protein. The cell then evolves into a large, rapidly dividing, pre-B-cell I which is a cytoplasmically μ^- cell. This pre-B-cell I then stops dividing, shrinks, and undergoes light chain V-J rearrangement becoming a pre-B-cell II which expresses surface IgM, which leave

- 49 -

the marrow as immature B-cells. Most of the emerging immature B-cells continue to develop and to produce surface IgD, indicative of their completion of differentiation and development as fully mature immunocompetent peripheral B-cells, which reside primarily in the spleen. [Hardy and Rolink, *Ann. NY Acad. Sci.*, 764, pp. 19-24 (1995); Rolink and Melchers, *Immunol. Lett.*, 54, pp. 157-161 (1996)]. However, it is possible to eliminate the delta constant region and still obtain immunocompetent cells.

B-cell differentiation and development can be monitored and/or tracked through the use of surface markers. For example, the B220 antigen is expressed in relative abundance on mature B-cells in comparison to pre-B-cells I or II. Thus, cells that are B220⁺ and surface IgM⁺ (μ^+) can be utilized to determine the presence of mature B-cells. Additionally, cells can be screened for surface IgD expression (δ^+). Another antigen, heat stable antigen, is expressed by pre-B-cells I and later developmental stages.

TABLE 2

Marker	Bone Marrow			Spleen	
	pro-pre-B-cell	pre-B-cell I	pre-B-cell II emerging B-cell	immature B-cell	mature B-cell
B220	+	+	+	+	++
HSA	-	+	+	hi	lo
μ	-	-	+	+	+
δ^*	-	-	-	-	+

* Assuming the presence of a functional copy of the C δ gene on the transgene.

- 50 -

Through use of B-cell markers, such as those mentioned above, development and differentiation of B-cells can be monitored and assessed.

We have previously demonstrated that DI mice
5 (mice that do not undergo heavy chain V-D-J rearrangement or light chain V-J rearrangement) do not produce mature B-cells. In fact, such mice arrest at the production of pro-pre-B-cells and B-cells never move from the bone marrow to peripheral tissues,
10 including the spleen. Thus, both B-cell development and antibody production are completely arrested. The same result is seen in mice that are only heavy chain inactivated -- B-cell development and differentiation arrests in the bone marrow.

15 Our XenoMouse I strain produced functional, somewhat mature B-cells. However, the numbers of B-cells, in both the bone marrow and peripheral tissues, were significantly reduced relative to wild type mice.

20 In contrast, our XenoMouse II strains and L6 strains, unexpectedly possess almost complete B-cell reconstitution. Therefore, in accordance with the invention, we have demonstrated that through the quantitative inclusion or qualitative inclusion of
25 variable region genes B-cell differentiation and development can be greatly reconstituted. Reconstitution of B-cell differentiation and development is indicative of immune system reconstitution. In general, B-cell reconstitution is
30 compared to wild type controls. Thus, in preferred embodiments of the invention, populations of mice having inserted human variable regions possess greater

- 51 -

than about 50% B-cell reconstitution when compared to populations of wild type mice.

ISOTYPE SWITCHING BY XENOMOUSE

As is discussed in detail herein, as
5 expected, XenoMouse II mice undergo efficient and effective isotype switching from the human transgene encoded mu isotype to the transgene encoded gamma-2 isotype. As mentioned above, mice in accordance with the invention can additionally be equipped with other
10 human constant regions for the generation of additional isotypes. Such isotypes can include genes encoding γ_1 , γ_2 , γ_3 , γ_4 , α , ϵ , δ , and other constant region encoding genes. Alternative constant regions can be included on the same transgene, i.e., downstream from the human mu
15 constant region, or, alternatively, such other constant regions can be included on another chromosome. It will be appreciated that where such other constant regions are included on the same chromosome as the chromosome including the human mu constant region encoding
20 transgene, cis-switching to the other isotype or isotypes can be accomplished. On the other hand, where such other constant region is included on a different chromosome from the chromosome containing the mu constant region encoding transgene, trans-switching to
25 the other isotype or isotypes can be accomplished. Such arrangement allows tremendous flexibility in the design and construction of mice for the generation of antibodies to a wide array of antigens.

It will be appreciated that constant regions
30 have known switch and regulatory sequences that they are associated with. All of the murine and human constant region genes had been sequenced and published

- 52 -

by 1989. See Honjo et al. "Constant Region Genes of the Immunoglobulin Heavy Chain and the Molecular Mechanism of Class Switching" in *Immunoglobulin Genes* (Honjo et al. eds., Academic Press (1989)), the disclosure of which is hereby incorporated by reference. For example, in U.S. Patent Application Serial No. 07/574,748, the disclosure of which is hereby incorporated by reference, the cloning of the human gamma-1 constant region was predicted based on known sequence information from the prior art. It was set forth that in the unrearranged, unswitched gene, the entire switch region was included in a sequence beginning less than 5 kb from the 5' end of the first γ -1 constant exon. Therefore the switch region was also included in the 5' 5.3 kb HindIII fragment that was disclosed in Ellison et al. *Nucleic Acids Res.* 10:4071-4079 (1982). Similarly, Takahashi et al. *Cell* 29:671-679 (1982) also reported that the fragment disclosed in Ellison contained the switch sequence, and this fragment together with the 7.7 kb HindIII to BamHI fragment must include all of the sequences necessary for the heavy chain isotype switching transgene construction.

Thus, it will be appreciated that any human constant region of choice can be readily incorporated into mice in accordance with the invention without undue experimentation. Such constant regions can be associated with their native switch sequences (i.e., a human γ_1 , γ_2 , γ_3 , or γ_4 constant region with a human γ_1 , γ_2 , γ_3 , or γ_4 switch, respectively) or can be associated with other switch sequences (i.e., a human γ_4 constant region with a human γ_2 switch). Various 3' enhancer sequences can

- 53 -

also be utilized, such as mouse, human, or rat, to name a few. Similarly other regulatory sequences can also be included.

As an alternative to, and/or in addition to,
5 isotype switching *in vivo*, B-cells can be screened for secretion of "chimeric" antibodies. For example, the L6 mice, in addition to producing fully human IgM antibodies, produce antibodies having fully human heavy chain V, D, J regions coupled to mouse constant
10 regions, such as a variety of gammas (i.e., mouse IgG1, 2, 3, 4) and the like. Such antibodies are highly useful in their own right. For example, human constant regions can be included on the antibodies through *in vitro* isotype switching techniques well known in the
15 art. Alternatively, and/or in addition, fragments (i.e., F(ab) and F(ab')₂ fragments) of such antibodies can be prepared which contain little or no mouse constant regions.

As discussed above, the most critical factor
20 to antibody production is specificity to a desired antigen or epitope on an antigen. Class of the antibody, thereafter, becomes important according to the therapeutic need. In other words, will the therapeutic index of an antibody be enhanced by
25 providing a particular isotype or class? Consideration of that question raises issues of complement fixation and the like, which then drives the selection of the particular class or isotype of antibody. Gamma constant regions assist in affinity maturation of
30 antibodies. However, the inclusion of a human gamma constant region on a transgene is not required to achieve such maturation. Rather, the process appears

- 54 -

to proceed as well in connection with mouse gamma constant regions which are trans-switched onto the mu encoded transgene.

EXAMPLE 1

5 YAC VECTORS FOR MURINE γ 1-HUMAN γ 4
 OR MURINE γ 1-HUMAN γ 1

Replacement vectors for targeting the parent YAC yH1C to replace the human γ 2 switch element and human CH γ 2 exons with the murine γ 1 switch element and
10 either the human C γ 1 exons or the human C γ 4 exons were prepared (Figures 1). The vectors were designated as pMuShu1 and pMuShu4 (Figure 5). The vectors were constructed using a low copy number cloning vector known as pACYC177. This vector pACYC177 is available
15 from New England Biolabs, Inc., Beverly, MA and the sequence can be found in Genbank sequence database under the sequence accession number Genebank #X 06402. A low copy number origin of replication is useful to prevent unwanted rearrangements or deletions of the
20 plasmid DNA when propagated in *E coli*.

The first step was to introduce a linker into pACYC177 in order to accommodate the elements needed for the targeting vector. The linker contained the following restriction sites: NheI-SalI-SmaI-
25 NotI-EcoRI-XbaI-SacI-BamHI.

The nucleotide sequence of the linker (SEQ ID NO: 2) is shown below:

5' - cta gtc gac aaa tat tcc ccg ggc ggc cgc tta
cgt atg aat tca gcg cgc ttc tag aac tcg agt gag
30 ctc

- 55 -

The nucleotide sequence of the complimentary strand of the linker (SEQ ID NO: 3) is shown below:

5' - gat cga gct cac tcg agt tct aga agc gcg ctg
aat tca tac gta agc ggc cgc ccg ggg aat att tgt
5 cga

Restriction Enzymes

Unless otherwise stated all restriction enzymes were purchased from New England Biolabs Inc. (Beverly, MA). Furthermore, all restriction digestion
10 conditions were standardized according to the following conditions: 1 microgram of DNA was digested in 20 µl of the appropriate restriction buffer and using 5 units of restriction enzyme for 1 hour. Restriction buffers are specified by the manufacturer for particular enzymes
15 and the compositions are provided in the product catalog from New England Biolabs Inc. (Beverly, MA).

To introduce the linker, pACYC177 was digested with restriction enzymes NheI/BamHI according to the manufacturer's instructions. The linker as
20 shown above was ligated with a 2208 bp fragment of pACYC177 isolated on an agarose gel and purified using GeneClean kit (Bio 101) (Vista, CA). This process removed only non-essential regions of the vector including NheI and BamHI restriction sites.

25 The next step was to introduce the yeast URA3 gene, with its promoter and coding sequences as a marker for selection of yeast cells containing the YAC. A DNA 1971 bp fragment of the URA3 gene containing the promoter and coding sequence was obtained from pYAC4,
30 which is available from the American Type Culture Collection (ATCC) catalog no. 67379 (Manassas, VA) and the sequence can be obtained from Genbank using

- 56 -

accession no. #U01086. The URA3 fragment also provides sufficient 3' homology for targeting. The plasmid pYAC4 was digested with the restriction enzymes, SalI and MscI, according to the manufacturers instructions. Likewise, the vector-linker combination, pACYC177/linker, was digested with SalI and SmaI according to the manufacturers instructions. Subsequently, the two restriction nucleotide digested DNAs pACYC177 and URA3 were ligated together to produce Int 2.

The next step was to introduce the beta-galactosidase gene (beta Gal) by first digesting Int 2 with XbaI and SacI according to the manufacturers instructions. The beta Gal gene was cloned from the vector pGK beta Gal which can be obtained from Cell Genesys, Inc. (Foster City, CA). The DNA pGK beta Gal was digested with the restriction enzymes, XbaI and SacI. Linearized Int 2 and the 2553 kb fragment from pGK beta Gal were ligated together to produce the next intermediate called Int 3.

The above beta Gal expression construct is incomplete. The missing portion of beta Gal is obtained by digesting the pGK beta Gal plasmid with restriction enzyme SacI and NcoI. The 1165 kb fragment from the pGK beta gal digestion was isolated by agarose gel eletrophoresis, wherein the fragment was excised from the ethidium bromide stained gel and purified like described above using a Geneclean kit (Bio 101) (Vista, CA). Similarly, Int 3 was digested with the restriction enzyme SacI, according to the manufacturers instructions. The linearized Int 3 was isolated by agarose gel eletrophoresis. The 1165 bp fragment from pGK beta gal and the linearized Int 3 were ligated

- 57 -

together using the enzyme T4 DNA ligase, purchased from New England Biolabs, Inc. The DNA fragment was reisolated and then treated with Klenow fragment to blunt the ends. The linearized DNA was circularised
5 using the enzyme NcoI which blunt ligates to SacI blunt ends to produce Int 4.

INTRODUCTION OF 5' HOMOLOGY

The region of 5' homology for targeted recombination was isolated from the sequence of A-287-C10 YAC by rescue of the 3' end and had been previously
10 cloned into plasmid ppKMlc [See Mendez et al., *Nature Genetics* 15:146-156 (1997)]. The A287-C10 YAC was isolated by screening DNA pools from the Washington University human YAC library (Washington University,
15 St. Louis, MO) using PCR primers for the human V_H6 gene. Isolation and characterization of the A287-C10 YAC was described in detail in International Patent application WO 94/02602, by Kucherlapati et al., published on February 3, 1994, and that disclosure is hereby
20 incorporated by reference.

Int 4 was digested with the restriction enzymes NotI and SnaBI and then treated with Calf Intestine Phosphatase as follows: 1 microgram of DNA in 20 microliters of restriction digest reaction, 5 units
25 of calf intestine phosphatase (New England Biolabs., Beverly, MA). The enzyme and DNA were incubated for 30 minutes at 30 °C, then heated to 65 °C for another 30 minutes to denature the phosphatase. The vector ppKMlc was digested with the restriction enzyme, EcoRI, and
30 then treated with Klenow Fragment to create a blunt end to remove the EcoRI site. The linearized ppKMlc was

- 58 -

isolated and digested with the restriction enzyme NotI, according to the manufacturers instructions. A fragment of approximately 1 kb 5' homology was isolated. Next, the 1 kb fragment was ligated together
5 with the NotI and SnaBI digested Int 4. This DNA preparation was named Int 5.

The γ 1 and γ 4 CH1, Hinge, CH2, and CH3 coding exons, transmembrane exons and approximately 3 kb of downstream sequence (~7kb each) were introduced into
10 the replacement vectors through two intermediate DNAs derived from pBR322.

First, pBR322 was digested with Hind III, treated with Calf Intestine Phosphatase (CIP) and ligated with the approximately 7kb Hind III fragment
15 containing the γ 1 sequences from P1 clone #1737 (G1). P1 phage clone was purchased from Genome Systems, Inc. (St Louis, MO). This resulted in intermediate plasmid pCG12.

The second intermediate was constructed by
20 digesting pBR322 with the restriction enzymes, HindIII and BamHI. The 3986 kb fragment was treated with calf intestine phosphatase and ligated with approximately 7kb HindIII/BamHI fragment containing the human γ 4 sequences from BAC clone #176E10, purchased from Genome
25 Systems, Inc. (St. Louis, MO). This intermediate plasmid was called pCG43.

In order to complete construction of targeting vectors, a new linker was cloned into the XbaI restriction site of Int 5. The linker had the
30 following restriction sites: XbaI•kill -MfeI-SspI-HindIII-SnaBI-BclI-XhoI-MluI-Xba•kill. Int 5 with the linker cloned into it was called Int6. The linker sequence ID NO: 4) is shown below:

- 59 -

5' cta ggc aat tga taa tat taa gct tta cgt atc tga
tca tcc tcg aga cgc gtg

Complementary strand sequence (SEQ ID NO: 5):

5' cgt taa cta tta taa ttc gaa atg cat aga cta gta
5 gga gct ctg cgc acg atc

The Linker was oriented in Int 6 as such:

A287 - SnaBI(ex)-EcoRI-BssHII-XbaI(ex)-MfeI-
HingIII-BclI-XhoI-MluI-pGK-beta Gal

The restriction site XbaI•kill indicates that
10 the particular XbaI site will be eliminated upon
ligation into the larger DNA. The linker is
conveniently designed so that it can ligate into an
XbaI site but the site does not survive the ligation.
The particular XbaI site which contained the linker was
15 determined by first cloning the linker and then
digesting the DNA with the following pairs of
restriction enzymes separately: NotI and HindIII; XbaI
and SphI; and MluI and SphI. Introduction of the
linker eliminates one XbaI site. The position of the
20 linker in Int 6 was determined by the distance between
the newly introduced Hind III site and the NotI site
which was present in Int 5.

CLONING OF MOUSE γ 1 SWITCH REGION:

The plasmid EH10 was obtained from the
25 University of Michigan and is a pBR322 based plasmid
containing a murine γ 1 switch region on a HindIII/EcoRI
fragment [M.R. Mowatt et al., *J. Immunol.* 136:2647-2683
(1983)]. The plasmid was digested with the restriction
enzymes EcoRI and HindIII and the 10kb fragment

- 60 -

containing the mouse $\gamma 1$ switch was isolated and purified as above.

CONSTRUCTION OF pMSL4

Construction of pMSL1 involved a three way
5 ligation. The first element was the 10kb fragment
containing the mouse $\gamma 1$ switch was isolated from EcoRI
and HindIII digested EH10 as described above. The
second element was pBR322 digested with BamHI and
EcoRI. The final element was the pBR322 based plasmid,
10 pCG43, containing human $\gamma 4$ on an approximately 7kb
HindIII and BamHI fragment. All three were ligated
together to create pMSL4.

CONSTRUCTION OF pMSL1

Construction of pMSL1 also involved a three
15 way ligation. The first element was the 10kb fragment
containing the mouse $\gamma 1$ switch was isolated from EcoRI
and HindIII digested EH10 as described above. The
second element was pBR322 digested with EcoRI and
BamHI. The final element was pBR322 based plasmid
20 pCG12 containing approximately 7 kb fragment of human
 $\gamma 1$ which was modified by introduction of Hind III kill-
HamHI linker into HindIII site on 3' end of the 7 kb
fragment. Thusly modified plasmid after double digest
with BamHI and HindIII releases 7 kb HindIII/BamHI
25 fragment which is subsequently used in three piece
ligation.

MOUSE 3' ENHANCER

We isolated the 0.9 kb core part of the
enhancer by StuI restriction digestion from the 4kb

- 61 -

MluI fragment of the pIBgamma2 targeting vector containing the murine 3' enhancer (HSIg2) cloned into pIB [M.J. Mendez et al., *Nature Genetics*, 15:146-156 (1997)].

- 5 We digested Int 6 with the restriction enzyme XhoI, followed by treatment with the Klenow fragment to create blunt ends. We ligated the resulting linearized Int 6 with the 0.9 kb StuI fragment of murine 3' enhancer to create Int 7. We verified the cloning
- 10 reaction by performing restriction digestion on sample clones with using EcoRI. In addition, we confirmed the desired orientation of the fragment by digests with NcoI; NcoI and HindIII; and HindIII and PvuII and the known restriction map of the 0.9 bp Stu I fragment.
- 15 We introduced another linker into plasmid Int 7. We digested Int 7 with MfeI and SnaBI (double digest), followed by treatment with calf intestine phosphatase. Next, we introduced the following linker by performing a ligation reaction and creating
- 20 intermediate Int 8. The restriction sites inserted by the linker into Int 7 are as follows: MfeI•kill-HindIII-SnaBI-BclII-BglIII-Bam HI-BglIII-NheI•kill. Again, MfeI•kill indicates that the MfeI site was eliminated upon ligation into the larger DNA.

- 25 The nucleotide sequence of the linker (SEQ ID NO: 6):
- 5' aat taa gct tgt acg tac tga tca aga tct gga tcc
aga tct

The nucleotide sequence of the complementary strand
(SEQ ID NO: 7):

- 62 -

5' aga tct gga tcc aga tct tga tca gta cgt aca agt
t

TARGETING VECTORS

The complete targeting vectors were
5 constructed by digesting Int8 with the restriction
enzymes SceI and HindIII, followed by treatment with
calf intestine phosphatase. The plasmids pMSL1 and
pMSL4 were partially digested with the restriction
enzymes SceI and HindIII. A 17 kb fragment was
10 isolated by polyacrylamide gel electrophoresis. The
purified 17 kb fragment was ligated with Int 8 to
create the final targeting vectors as shown in Figure
5.

EXAMPLE 2

15 TARGETING OF γ 1 OR γ 4 CONSTRUCTS ON γ H1C YAC

The TV1 or TV4 vectors (5 ug DNA) were
linearized by digestion with the restriction enzyme
NotI (Figure 5). The DNA was purified by phenol
extraction followed by phenol/chloroform extraction.
20 Next, the DNA was precipitated with ethanol and then
used to transform a yeast clone containing the γ H1C
YAC using a LiAc transformation protocol. [See
Schiestl, R.H. et al., Curr. Genet. 16,339-346 (1989)].
Transformants were plated onto SC-URA agar media plates
25 and incubated at 22 °C until colonies appeared or
approximately 5-6 days. SC-URA plates contain a media
for growth of yeast which lacks uracil and therefore
selects for yeast colonies that can produce their own
uracil. Similarly, SC-LYS plates contain a media for
30 growth of yeast which lacks lysine and selects for

- 63 -

yeast colonies that can produce their own lysine. The resulting colonies were repicked onto SC-URA plates and on SC-LYS plates — for genetic testing — to look for the loss of the LYS marker. Only clones which grew on SC-URA and did not grow on SC-LYS were grown in YPDA media for 48 hours at 22 °C. YAC DNA was isolated and analyzed by polymerase chain reaction (PCR) to evaluate whether the desired isotype replacements occurred as expected. In this case, human $\gamma 1$ or $\gamma 4$ CH exons should have replaced human $\gamma 2$ CH coding exons (Figure 1). The yeast media used here was prepared from supplements obtained from BIO 101 (Vista, CA).

PCR Primers used for this assay are as follows:

HG1: 5' cac acc gcg gtc aca tgg c (SEQ ID NO: 8)
HG3: 5' cta ctc tag ggc acc tgt cc (SEQ ID NO: 9)

The PCR reaction consisted of 35 cycles of the following: 94°C for 15 seconds followed by 60°C for 45 seconds and then 72°C for 90 seconds per cycle. HG1 primer was positioned at nucleotide 181 on consensus human $\text{Cy}1$, $\text{Cy}2$, $\text{Cy}4$ alignment and primer HG3 was positioned at nucleotide 994 of this alignment. These primers will amplify DNA from $\text{Cy}1$, $\text{Cy}2$, and $\text{Cy}4$ isotypes.

Due to restriction site polymorphism in the human Cy genes, the particular isotype of the template DNA could be determined by restriction digestion of the PCR products to yield unique sets of DNA fragments. For example, the restriction enzyme PvuII restricts the PCR product into two fragments of 621 bp and 196 bp when $\text{Cy}2$ DNA is the template for the PCR products, but does not cut the product if $\text{Cy}1$ or $\text{Cy}4$ is the template.

- 64 -

Similarly, the restriction enzyme Eco47III restricts the PCR product into two fragments of 438 bp and 379 bp when Cy1 is obtained. Finally, the restriction enzyme BglIII restricts the PCR product into two fragments of 686 bp and 125 bp when Cy4 is obtained. In this way all three isotypes of IgG could be distinguished.

In the next level of characterization, all yeast clones which exhibited correct genetics as well as the desired IgG isotype were further screened by Southern blot assay [J. Sambrook et al. Molecular Cloning: A Laboratory Manual, Chapter 9, pages 31-45, Cold Spring Harbor Laboratory Press (1989)]. A 5 microgram sample of DNA for each clone was digested overnight with the restriction enzymes Hind III, EcoRI and Bam HI. yHIC YAC DNA, which served as the original target on the replacement vectors was used as control. The digested DNAs were separated on 0.8% agarose gel, stained with ethidium bromide and photographed and then transferred onto nylon membrane (Gene Screen Hybridization Membrane, NEN Life Sciences). Next, the YAC candidates were checked on Southern blots using hybridization probes from the following Ig genes: D, mu, J, delta, murine 3' enhancer, Cg1,4,V1-6. [See M.J. Mendez et al., Genomics 26, 294-307 (1995); M.J Mendez et al., Nature genetics vol 15, 146-156 (1997) (V3 probe)]

The following probes were used for Southern blotting:

HG1 : CAC ACC GCG GTC ACA TGG C (SEQ ID NO: 8)

HG3: CTA CTC TAG GGC ACC TGT CC (SEQ ID NO: 9)

These primers will amplify ~820 bp fragment for gamma 1,2 and 4 . Either one can be used as a probe as they are highly homologous.

- 65 -

To amplify VH5 following primers were used:

VH5A: 5' GTC GAC GGG CTC GGG GCT GGT TTC TCT (SEQ ID NO: 10)

5 VH5B 5' GGG CCC TGA TTC AAA TTT TGT GTC TCC (SEQ ID NO: 11)

For HPRT following primers were used :

REP3: 5' CTG GAG TCC TAT TGA CAT CGC C (SEQ ID NO: 12)

REP4: 5' GGT TCT TTC CGC CTC AGA AGG (SEQ ID NO: 13)

10 And, finally, to amplify Cmu following primers were used ;

Jm1: 5' GCT GAC ACG TGT CCT CAC TGC (SEQ ID NO: 14)

Jm4: 5' CCC CAG TTG CCC AGA CAA CGG (SEQ ID NO: 15)

15 Finally, we confirmed the general structural integrity of the YACs using CHEF gel pulse-field gel electrophoresis (CHEF DR-II, Bio Rad Life Sciences, Hercules, CA).

We designated the YAC encoding Cy1 as yH2Bm and the YAC encoding human Cy4 as yH2Cm.

EXAMPLE 3

20 CONSTRUCTION OF VECTORS FOR RETROFITTING yH1C YAC TO y1 (TV G1) AND y4 (TV G4)

25 Vector construction for preparing the targeting vectors to retrofit the yH1C YAC to yHG1 and yHG4 is schematically shown in Figure 6. The targeting vectors were built on a backbone of pACYC177 (Genbank #X06402) available from New England Biolabs Inc. (Beverly, MA). We introduced a linker into pACYC177 to facilitate cloning of a murine 3' enhancer.

- 66 -

We called the pACYC177 vector containing the linker Int 9. The arrangement of restriction enzyme cloning sites in the linker was as follows: HindIII-SalI-MluI-PacI-FseI-HindIII. The linker nucleotide sequence (SEQ ID NO: 16) is shown below:

5' agc ttg tcg aca cgc gtt taa tta agg ccg gcc a

The nucleotide sequence of the complementary strand (SEQ ID NO: 17):

5' agc ttg gcc ggc ctt aat taa acg cgt gtc gac a

10 We cloned the murine 3' enhancer from the yH1C targeting vector as an approximately 4kb MluI fragment. The 4 kb enhancer fragment was cloned into the MluI site of pACYC177 modified with the linker shown above and the DNA is called Int 10. Proper
15 orientation of murine 3' enhancer was determined by digesting prospective clones with the restriction enzymes NgoMI and EagI.

Amplification of a 5' homology region

20 A region of 5' homology was obtained by PCR amplification of the relevant portion of pIBgamma2 targeting vector [M.R. Mowatt, et al., J. Immunol. 136:2647-2683 (1983)]. The nucleotide sequence of the primers used for amplifying the 5' homology region
25 were[See also Genbank Accession no. M12389];

Primer 1 :5' tgg tgg ccg aga agg cag gcc a (SEQ ID NO: 18)

- 67 -

Primer 2 :5' ccg cgg gca tgc aac ttc gta taa tgt atg
cta tac gaa gtt att gtg gga cag agc tgg gcc
cag g (SEQ ID NO: 19)

Primer 2 contains SacII and SphI sites as well as a lox
5 p element. The 5' homology region was PCR amplified
using the following PCR conditions: 20 cycles of 94°C
for 3 seconds, followed by 55°C for 30 seconds, and then
72°C for 60 seconds. The region was then sequenced
after cloning it into TA-TOPO vector. TA-TOPO is
10 available from Invitrogen, Inc. (Carlsbad, CA).

The primers for 5' homology sequencing are shown below:
seq1: gtc tgg ccc ctc tgc tgc (SEQ ID NO: 20)
seq2: cac cca taa aag gct gga (SEQ ID NO: 21)
rev. seq1: acg gct cat gcc cat tgg (SEQ ID NO: 22)
15 rev. seq2: tag tga gtg ggc ctg act (SEQ ID NO: 23)

We compared the resulting sequence to the
human switch γ 2 sequence (Genebank #U39934) and
determined that it is identical.

Cloning of γ 4 coding and 3' homology regions

20 We obtained human γ 4 coding exons and a
region of 3' homology by performing partial enzymatic
digestion of the plasmid pCG43 using the restriction
enzyme SacII. We then digested plasmid pGS43 with the
restriction enzyme BamHI and we cloned a purified
25 approximately 7 kb fragment into the TA vector
containing 5' homology. We called this intermediate
recombinant DNA molecule IntI G4.

IntI G4 was digested with BamHI, treated with
calf intestine phosphatase and then the vector were

- 68 -

isolated using agarose gel electrophoresis. The isolated vector was ligated as described before with 3.4 kb BamHI fragment from pIB gamma2. Orientation of the insert was determined by double digest NotI/Hind

5 III. The HindIII site was determined to be at the 5' end of 3.4 kb BamHI fragment as determined by the sizes of fragments after digests.

Next we needed to determine orientation of the linker in pACYC177/enhancer plasmid. This was done

10 by preparing a panel of double digestions of pACYC177/linker with SmaI and with one of each of the following second enzymes, SalI, MluI, PacI and FseI. Linker orientation was determined by sizes of resulting fragments.

15 The positions of restriction sites in pACYC177/enhancer plasmid is as follows: ClaI/SmaI (pACYC177)-HindIII-FseI-PacI-MluI- ((enhancer:PstI-PvuII-EcoRI-NcoI-NheI-ApaI)) -MluI-SalI)- PfmII (pACYC177).

20 The next step was to clone URA 3 gene into pACYC177/enhancer plasmid. The purpose of this is to retrofit the targeting vector with a selectable yeast marker, as well as 3' homology to drive homologous recombination. In order to clone URA3 gene, Int 2

25 (constructed for original TV1 and 4, described in Example 1) was digested using SacII/SalI. Similarly, the pACYC177/enhancer plasmid was digested with SacII/SalI. 3.8 kb fragment from Int 2 digestion and a 5kb fragment from pACYC177/enhancer were isolated on

30 agarose gel electrophoresis and ligated together. The resulting plasmid contains enhancer and URA3 gene in pACYC177 backbone. The next step was to introduce two more linkers into Int1 G4 with cloned 3.4 kb Bam HI

- 69 -

fragment. The resulting intermediate was called Int 2 G4.

The Linkers were as follows: NotI-kill-FseI-NotI-kill and Linker sequences:

- 5 GGCCATGGCCGGCCAT (SEQ ID NO: 24)
TACCGGCCGGTACCGG (SEQ ID NO: 25)

The second linker had the following restriction sites:
BamHI-KpnI-EcoRV-MfeI-FseI-SfiI-BamHI-kill:
GATCCGGTACCGATATCCAATTGGGCCGGCCGGCCATATAGGCCT (SEQ ID
10 NO: 26)
GCCATGGCTATAGGTAAACCCGGCCGGCCGGTATATCCGGACTAG (SEQ
ID NO: 27)

The purpose of introducing linker 1 was to provide an FseI site for the final cloning step, as well as eliminating one of NotI sites. In addition, this leaves the final targeting vector with a unique NotI site, which was used to linearize the targeting vector before transformation. The second linker was used to clone the last fragment needed to restore the complete downstream region of yH1C YAC, a 1.5 kb BamHI/EcoRI fragment.

Linker 2 was introduced via partial digest with Bam HI. IntI G4 (with cloned 3.4 kb BamHI fragment) was partially digested with BamHI and the partial digest was isolated on agarose gel as a 13kb fragment. The 13 kb fragment was treated with calf intestine phosphatase and ligated with linker. The position of the linker and its orientation (whether it went into correct BamHI site at 3' end of 3.4 BamHI fragment) was determined by digesting clones with MfeI and NotI (double digest). The MfeI site is introduced with a

- 70 -

linker and the NotI site is present in the vector. Relative fragment sizes permit allowed identification of the position and orientation of the linker. The plasmids retrofitted with linker are now Int 3 G1 and
5 Int 3 G4.

The next step was to clone into Int 3 G4 plasmid, a 1.5 kb fragment obtained by BamHI/EcoRI double digest of pIBgamma2 plasmid. The 1.5 kb fragment was cloned into BamHI partial digest/MfeI
10 digest of Int 3 G4. Since it was directional cloning, no orientation determination was needed in this step. The resulting plasmid was called Int 4 G4.

The next step was to introduce the linker with the restriction sites NotI-kill-FseI-NotI-kill. Int
15 4 G4 was digested with NotI, treated with calf intestine phosphatase, isolated on agarose gel and ligated with the linker. The resulting plasmid was called Int 5 G4. Again, there was no need to determine linker orientation in this step. As a result, a unique
20 Not I site was eliminated and one FseI site was added. The purpose of introducing the FseI site was to allow cloning of a fragment spanning from 5' homology region to the 1.5 kb fragment to pACYC177/enhancer/URA3 plasmid.

25 The final cloning step was a partial digest of Int 5 G4 with FseI, followed by isolation of a 13 kb fragment on agarose gel and ligation into the unique FseI site of pACYC177/enhancer/URA3. The orientation of this insert was determined by a double restriction
30 digest with NotI/FseI. The final targeting vectors were called TV G4.

To construct the TVG1 targeting vector to retrofit the yH1C YAC with human Cyl coding exons and a

- 71 -

a region of 3' homology, we use the procedure described above for the construction of TV G4. We obtain the Cyl coding exons and 3' homology region from plasmid pCG12.

5 **EXAMPLE 4** CONSTRUCTION OF TARGETING VECTORS TV
 G1/2 AND TV G4/2 FOR TARGETING OF yH1C
 YAC FOR NON-COGNATE SWITCHING

 Next, we constructed a vector, TV G1/2, which has a chimeric construct of human γ 1 CH coding exons attached downstream of 5 kb of human switch γ 2 region
10 DNA (Figure 3). In addition, this vector contains human γ 2 transmembrane exons located 3' of the γ 1 CH coding exons (Figures 3). The vector is based on pACYC177 vector, which is available from New England Biolabs (Beverly, MA). We constructed the vector,
15 which we call TV G1/2, using the following procedure:

1. First, 5 ug of pIBgamma2 was digested with the restriction enzymes, HindIII and BamHI, and a 6.5 kb fragment was isolated on an agarose gel. The vector, pIBgamma2, contains 22 kb of human genomic
20 DNA with γ 2 flanked by two EcoRI restriction enzyme sites and was previously used to generate yH1C. The 6.5 kb fragment was then ligated into the pCR™2.1 vector from Invitrogen, Inc. (Carlsbad, CA). pCR2.1 was prepared by digesting
25 1 μ g of plasmid with BamHI/HindIII and treating with calf intestine phosphatase and then isolated on an agarose gel.
2. The resulting plasmid (6.5 kb fragment + pCR™2.1 vector) was subjected to partial digestion with

- 72 -

- restriction enzyme XmnI followed by digestion with the restriction enzyme HindIII. The digestion with XmnI occurs 75 bp upstream of the $\gamma 2$ stop codon. Therefore, the 4th exon of the $\gamma 1/\gamma 2$ chimeric gene initially contains 75 bp of $\gamma 2$ in addition to the two 3' $\gamma 2$ membrane exons. The coding region for $\gamma 1$ and $\gamma 2$ are identical throughout this 75 bp region and there is no effect.
- 5
- 10 3. Next, 5 μ g of pCG12 was digested with the restriction enzymes HindIII and XmnI and a 1.7 kb fragment was isolated by agarose gel electrophoresis. The vector pCG12, which is described in Example 1, contains approximately 7kb
- 15 of human $\gamma 1$.
4. The 1.7 kb HindIII/XmnI fragment comprising most of the coding sequences of $\gamma 1$ obtained from pCG12 was ligated into the HindIII/XmnI partially digested vector (6.5 kb fragment + pCRTM2.1 vector)
- 20 described in step 2.
5. The resulting plasmid contains chimeric sequences of coding regions of $\gamma 1$ attached to a downstream region of $\gamma 2$ that contains transmembrane exons. We verified the composition of the plasmid by
- 25 restriction digestion with Eco47III.
6. The pCRTM2.1 vector having cloned 5' homology, described in Example 3, was digested with SacII and BamHI. Likewise, we digested the plasmid described in step 5 with SacII and Bam HI (double
- 30 digest) and cloned an approximately 5kb fragment

- 73 -

from the plasmid into the pCR2.1 vector with the 5' homology. Then we digested the resulting vector with SacII and SacII and cloned into this site fragments obtained from the vector described in step 5). We determined the Orientation of the SacII insert by SphI digests. The resulting plasmid contained chimeric $\gamma 1/\gamma 2$ C_H exons downstream from a region of 5' homology in a pCR2.1 vector. We called this vector TA G1/2, indicating that it is derived from pCR2.1, known as TA cloning vector, and contain chimeric $\gamma 1/\gamma 2$ C_H exons.

7. We retrofitted the pCR[™]2.1 vector with cloned 5' homology described in step 6 with the yeast selectable marker, URA3, gene as follows:
 - a. We obtained the URA3 gene by digesting Int 2 (described in Example 1) with SalI. The products of the SalI digestion are subjected to an additional reaction with Klenow fragment to create blunt ends. These products are consecutively digested with SacI.
 - b. The pCR[™]2.1 vector produced in step 6 was digested with BamHI, then blunted with Klenow fragment and further digested with SacI.
 - c. The URA3 gene obtained in 7(a) was ligated into the pCR[™]2.1 vector prepared as in 7(b). The resulting vector contains a 5' homology region with a downstream URA3 gene, flanked

- 74 -

by NotI sites (one from the linker in Int2 and another originally present in pCR2.1).

8. We excised the NotI fragment from the plasmid produced in step 7 and cloned it into the NotI site of Int 1 (also used for TV1 and 4 cloning). We determined the correct orientation of the NotI fragment by using an EcoRI digest. The resulting plasmid, which we designated Int 3, has a 5' homology region with a URA3 gene downstream in the low copy origin background (pACYC177).
9. To finish construction of the final targeting vector for the G1/2 chimeric YAC, we partially digested the vector having the cloned 5' homology and G1/2 chimeric IgG constant region in the pCR2.1 background (produced in step 6) with EcoRI restriction enzyme and subsequently recut with SacI restriction enzyme. We isolated an 8Kb fragment containing the 5' homology and the chimeric G1/2 IgG constant region and cloned it into EcoRI and SacI sites of Int 3, which resulted in the TV G1/2 targeting vector. There was no need to determine the insert orientation, since we introduced the fragment by directional cloning.
- Alternatively, after step 7 we remove the 5' homology and URA3 gene from the plasmid described in step 7 by digesting with XbaI. Then we clone the 5' homology and URA3 gene into the XbaI site of the plasmid described in step 6 to produce the TV G1/2 vector shown in Figure 7.

- 75 -

We construct the TV G4/2 targeting vector, which has a chimeric construct with human γ 4 coding exons attached downstream of 5 kb of human switch γ 2 region DNA and human γ 2 CH coding exons (Figure 4) utilizing the above-described procedures with some modifications.

The plasmid produced in step 1 is subjected to partial digestion with restriction enzyme XmnI followed by digestion with the restriction enzyme HindIII. The digestion with XmnI occurs 75 bp upstream of the γ 2 stop codon. Therefore, the 4th exon of the γ 4/ γ 2 chimeric gene initially contains 75 bp of γ 2 in addition to the two 3' γ 2 membrane exons. There is a single base pair difference between γ 4 and γ 2 in this region, which results in a single amino acid change. To correct this, we perform site directed mutagenesis to a C to T at the human gamma 2 gene using a Directed Mutagenesis Kit from Clontech Laboratories Inc. (Palo Alto, CA).

The we perform the repair using two synthetic oligonucleotides: one to replace the nucleotide necessary to switch sequence from γ 2 to γ 4 and an auxiliary oligonucleotide to eliminate a NotI site in the pCR2.1 vector with the cloned chimeric G4/2 7kb fragment.

For replacing C with T, we use the following oligonucleotide:

CCTCTCCCTGTCTCTGGGTAAATGAGTGCC

The T residue in **bold** is replacing C in original plasmid.

The second oligonucleotide (to eliminate NotI site

- 76 -

) is:

TATCCATCACACTGGCGACCGCTCGAGCAT

This oligo contains a Not I site (shown in bold) in
5 which a G has been replaced with an A to disrupt the
site. We sequence part of the plasmid to confirm that
the correct nucleotide was replaced.

EXAMPLE 5TARGETING STRATEGY

10 We linearized the above described vectors,
TV1, TV4, TV G1, TV G4, TVG 1/2 and TV G4/2 and used
them to transform yeast cell cultures with yH1C YAC by
lithium acetate transformation. We subsequently used
the linearized vectors to replace the targeted genes on
15 yH1C as described below, to produce the new YACs yH2Bm,
yH2Cm, yHG1, yHG4, yHG1/2 and yHG4/2 respectively
(Figures 1-4). We plated yeast cells on SC-URA media
after transformation to select for the integration of
the URA3 marker. We checked any resulting clones for
20 YAC integrity using pulse field gel electrophoresis.
In addition, we analyzed clones by Southern blot to
validate the structure and identity.

In the case of yHG1/2, the URA3 gene in
resulting YACs was flanked by 5' homology sequences
25 which we removed as follows.

We plated yeast cultures containing yHG1/2
YAC on agar plates with 5 FOA (negative selection for
URA). We checked the resulting 5 FOA-resistant clones
for integrity by pulse field gel and Southern blots as
30 outlined above.

- 77 -

We use the TVG1 vector and the TV G4/2 vector to produce the yHG1 and yHG4/2 YACs, respectively, following the above-described strategies.

EXAMPLE 65 INTRODUCTION OF THE yH2BM YAC INTO ES CELLS

We introduced the YAC, yH2BM, into mouse embryonic stem (ES) cells through yeast spheroplast fusion as described in detail below. [See B. Birren *et al.*, *Genome Analysis: A Laboratory Manual*, Volume 3, 10 Cloning Systems, "Chapter 5: Introduction of YACs into mammalian cells by spheroplast fusion", pages 548-550, Cold Spring Harbor Laboratory Press, Plainview, NY]. yH2BM containing yeast cells were spheroplasted using zymolase 20T at 0.15 mg/ml. The yH2BM spheroplasts 15 were fused with HPRT-deficient E 14.TG3B1 mouse ES cells which had been cultured as described below [See Tsuda *et al.*, *Genomics* 42:413-421 (1997)]. HAT selection was initiated 48 hours after fusion. HPRT-positive ES cell clones were selected at a 20 frequency of 1 clone/15-20x 10⁶ fused cells. Twenty-one HAT resistant colonies were expanded for genome analysis and were analyzed for YAC integrity by Southern and CHEF blot analyses. In control experiments fusing ES cells alone and yeast 25 spheroplasts alone, no colonies were detected.

The detailed procedure is as follows:

PRODUCING YEAST SPHEROPLASTS

Excess yeast cells were prepared because up to 50% will be lost during the spheroplasting 30 procedure. For fusing 5 x 10⁷ ES cells, approximately 5 x 10⁹ yeast cells are needed.

- 78 -

We inoculated selective medium (SC-) with freezer stock to give a starting inoculum of approximately 5×10^6 cells/ml. We determined the cell density in the culture by use of a hemocytometer. We
5 grew the cells at 23°C with shaking at 250 rpm overnight. Incubation can also be at 14°C or 18°C to increase YAC stability. Culturing at 30°C may result in deletion of some Ig gene segments in the YAC. In the morning, the culture density was be 2×10^7 cells/ml.
10 We stepped down to 1×10^7 cells/ml with YPDA (rich) medium, and incubated for 2-3 hours. Culture density at this step should not exceed 2×10^7 cells/ml because exponentially growing yeasts are needed for efficient and complete spheroplasting.
15 We poured the desired amount of culture (to provide 5×10^9 cells) into sterile 50ml tubes, centrifuged at 1000-1200g (2300-2500 rpm in a Jouan GR4-22 centrifuge) at room temperature for 5-10 minutes, and discarded the supernatant. Alternatively,
20 large volumes of cells can be pelleted in large conical centrifuge tubes.

We added 20 ml of sterile H₂O to each tube of yeast cells, resuspended the cells by vortexing (or with a pipette), centrifuged as above, and discarded
25 the supernatant. Next, we added 20 ml of 1 M sorbitol to each tube of yeast cells, resuspended the cells by vortexing (or with a pipette), centrifuged as above, and discarded the supernatant. We resuspended the cells in SPE buffer (1 M sorbitol, 10 mM sodium
30 phosphate, 10 mM EDTA) containing a 1:500 dilution of freshly added 2-mercaptoethanol to a final cell concentration of 5×10^8 cells/ml.

- 79 -

We combined 10 μ l of the previous cell suspension with 90 μ l of 5% (w/v) SDS and another 10 μ l of the cell suspension with 90 μ l of 1 M sorbitol. We determined the cell concentration of each mixture with a hemocytometer. Warm the yeast cell suspension from the previous paragraph to 30°C. For each 1 ml of yeast cell suspension, add 1.5 μ l of a 100 mg/ml stock of Zymolyase-20T (ICN). We incubated stationary at 30°C. At 5 minute intervals, combine 10 μ l of the cell suspension with 90 μ l of 5% (w/v) SDS and determine the cell concentration with a hemocytometer. We monitored the decrease in the number of cells that remain in the presence of SDS treatment (compared with the initial cell concentration in the presence of sorbitol).

When 95% of the cells become lysed in SDS, immediately centrifuged the sample at 200-300g (1000-1200 rpm in a Jouan GR4-22 centrifuge) at room temperature for 5 minutes and poured off the supernatant carefully (the pellet should be very loose and some loss of cells will occur). The total time for spheroplasting (steps after zymolase is added) is typically 5-20 minutes.

We gently resuspended the spheroplasts in 20 ml of STC buffer (0.98 M sorbitol, 10mM Tris, 10mM CaCl₂) by inversion or careful pipetting, centrifuged the sample at 200-300g (1000-1200 rpm in a Jouan GR4-22 centrifuge) at room temperature for 5 minutes, and carefully poured off the supernatant. We repeated this procedure one time. We resuspended the spheroplasts at 2.5×10^8 cells/ml in STC and kept them at room temperature (or on ice) until used in step 14.

- 80 -

FUSION WITH ES CELLS

We transferred 1 ml containing 2.5×10^8 spheroplasts of the spheroplast suspension to a 15-ml tube and centrifuged 1 ml at 200-300g (1000-1200 rpm in a Jouan GR4-22 centrifuge) at room temperature for 5 minutes. We removed all of the supernatant by slow aspiration with a drawn-out glass pipette. With the tube in a semihorizontal position, we gently added 1 ml of ES cells (at 5×10^6 cells/ml) without disturbing the spheroplast pellet. The spheroplast:ES cell ratio can vary from 25:1 to 50:1.

We prepared the ES cells in advance as follows: we started ES cultures in plates coated with mouse primary feeders, with a starting density of 6×10^6 ES cells per 100-mm plate and standard ES medium (DMEM high glucose, 100 units/ml of penicillin, 100 µg/ml of streptomycin, 2 mM L-glutamine, 100 µM 2-mercaptoethanol, 1000 units/ml of murine Leukemia inhibitory Factor [ESGRO[™]], and 15% heat-inactivated fetal calf serum). Following 48 hours of standard growth conditions, we trypsinized the cultures and used the resulting cells to start cultures on gelatin-coated plates at 10^7 ES cells per 100-mm plate. We allowed growth to continue for 16-24 hours. Four hours before fusion, we replaced the medium on the ES plates with fresh medium. Immediately before fusion, we trypsinized the cells, washed three times with serum-free ES medium at room temperature, and resuspended in serum-free ES medium at 5×10^6 cells/ml.

We centrifuged the combined spheroplast/ES cell sample at 300g (1200 rpm in a Jouan GR4-22 centrifuge) at room temperature for 3 minutes and

- 81 -

carefully aspirated off all medium with a drawn-out glass pipette.

We gently tapped the tube to loosen the cell pellet and used a P1000 tip to slowly add 0.5 ml of 50% (w/v) PEG 1500 (pH 8.0; e.g., Boeringer Mannheim 783641) containing 10 mM CaCl_2 (prewarmed to 37°C). While this solution was being added, we gently mixed the cells with the pipette tip. Once all of the solution was added, we slowly pipetted the cell suspension up and down one time. We incubated the cell suspension at room temperature for 90 seconds. We slowly added 5 ml of serum-free ES medium by pipetting it from the bottom of the tube. We incubated cells at room temperature for 30 minutes and, after 30 minutes, centrifuged the cell suspension at 300g (1200 rpm in a Jouan GR4-22 centrifuge) at room temperature for 3 minutes and carefully aspirated off all medium with a drawn-out glass pipette.

We resuspended the cells in 10 ml of standard ES medium and plated the entire sample ($\sim 5 \times 10^6$ ES cells) on a 100-mm mouse primary-feeder-coated plate. If initial attempts result in the generation of too many colonies, the amount of the sample plated on each 100-mm plate may need to be adjusted downward.

We incubated the plates under standard ES cell growth conditions overnight and then replaced the medium with fresh ES medium.

Following 48 hours of culturing (after the spheroplast fusion), we began growth under the appropriate selective conditions (i.e., dictated by the specific mammalian selectable marker present on the YAC). We replaced the medium every 2 days. We picked

- 82 -

and plate ES colonies on mouse primary-feeder-coated plates for expansion per standard procedures. We typically observed ES colonies 10-15 days following spheroplast fusion.

5 Here, seven ES cell clones (referred to as 1 through 7 in Table 3) derived from ES cell fusion with yH2Bm-containing yeast were found to contain all expected EcoRI and BamHI yH2 fragments detected by probes spanning the entire insert. As shown in Table
10 3, the following human genes were detected in the ES cell genome as part of characterization of the ES cell DNA prior to transgenic mouse generation: all the different V_H families could be detected V_{H1} , V_{H2} , V_{H3} , V_{H4} , V_{H5} , and V_{H6} ; human D_H , and J_H ; human C_μ and C_δ
15 constant regions; mouse switch $\gamma 1$ (mS $\gamma 1$) and human C $\gamma 1$ C_H exons.

- 83 -

TABLE 3

h or ¹ m genes	ES Cell Clone						
	1	2	3	4	5	6	7
5	V _H 1	+	+	+	+	+	+
	V _H 2	+	+	+	+	+	+
	V _H 3	+	+	+	+	+	+
	V _H 4	+	+	+	+	+	+
	V _H 5	+	+	+	+	+	+
	V _H 6	+	+	+	+	+	+
10	D _H	+	+	+	+	+	+
	J _H	+	+	+	+	+	+
	C _u	+	+	+	+	+	+
	C _δ	+	+	+	+	+	+
15	mSy1	+	+	+	+	+	+
	Cy1	+	+	+	+	+	+

¹m - denotes mouse genes (mSy1)

EXAMPLE 7

INTRODUCTION OF ES CELLS CONTAINING
THE yH2BM YAC INTO MICE

20 In order to generate chimeric mice from ES cells containing the YAC yH2BM DNA, we microinjected of blastocysts, followed by breeding. We isolated ES cells containing the YAC yH2BM DNA as described in Example 5, and expanded for the generation of chimeric

25 mice. Next, we microinjected yH2BM-bearing ES cell clones into mouse C57Bl/6 blastocysts [See B. Hogan *et al.*, "Manipulating the Mouse Embryo: A Laboratory Manual", Section D, Introduction of New Genetic Information, "Injection of Cells into the Blastocyst"

- 84 -

pages 188-196, (1986) (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY)]. Chimeric offspring were identified by coat color.

Table 4

Clone	# embryos Injected	# live born pups	# of chimeras	# chimera breedings	# germline
2BM-1	162	16	15	8	4
2BM-2	144	5	5	3	2
2BM-3	335	20	15	6	0
2BM-4	261	6	4	3	0
2BM-5	344	15	15	7	1
2BM-6	382	27	27	8	6
2BM-7	201	29	27	7	5

Table 4 summarizes the data for generating transgenic mice using seven different yH2BM containing ES cell lines. Five out of seven clones were transmitted through the mouse germline.

EXAMPLE 8**BREEDING MICE CONTAINING yH2BM
YAC DNA WITH yK2:DI MICE**

In order to generate mice that produce human antibodies in the absence of endogenous antibodies, we bred yK2-transgenic mice with double-inactivated (DI) mouse strains. The DI mouse strains are homozygous for gene targeted-inactivated mouse heavy and kappa light chain loci and, thus, lack in antibody production [see Jakobovits *et al.*, *Nature* 362:255-258 (1993); Green *et al.*, *Nature Genetics* 7:13-21 (1994)]. We bred one of the yK2-transgenic mouse strains, J23.1, with DI mice to generate mice hemizygous or homozygous for yK2 YACs on a homozygous inactivated mouse heavy and kappa chain

- 85 -

background (yK2;DI). The breeding scheme for generating a new XenoMouse, which is hemizygous for the yH2BM YAC is shown below. Subsequent breeding of XenoMouse males to XenoMouse females yields XenoMouse progeny who are either hemizygous or homozygous for yH2BM and/or yK2. From these progeny, breeding of males and females, both of which are homozygous for both yH2BM and yK2, will yield a true breeding line of XenoMouse yH2BM.

10 XenoMouse H2BM Breeding scheme

Generation 1 : (Chimera or Transgenic bred to yK2:DI)

yH2BM⁺;yK2⁻;mJ_H^{+/+};mCk^{+/+} **X** yH2BM⁻;yK2⁺; mJ_H^{-/-};mCk^{-/-}

Generation 2 : (Xenohet x yK2:DI)

yH2BM⁺;yK2⁺;mJ_H^{+/-};mCk^{+/-} **X** yH2BM⁻;yK2⁺;mJ_H^{-/-};mCk^{-/-}

15 Generation 3 (Almost Xenomouse x yK2:DI)
or Xenomouse x yK2:DI)

yH2BM⁺;yK2⁻;mJ_H^{+/-};mCk^{-/-} **X** yH2BM⁻;yK2⁺;mJ_H^{-/-};mCk^{-/-}

yH2BM⁺;yK2⁺;mJ_H^{-/-};mCk^{-/-} **X** yH2BM⁻;yK2⁺;mJ_H^{-/-};mCk^{-/-}

XenoMouse: yH2BM⁺; yK2⁺; mJ_H^{-/-}; mCk^{-/-}

20 We confirmed the integrity of the human heavy and kappa chain YACs in XenoMouse H2BM strains by Southern blot analysis. In all XenoMouse H2BM strains analyzed, yH2BM was transmitted unaltered through multiple generations with no apparent deletions or
25 rearrangements.

- 86 -

EXAMPLE 9

FLOW CYTOMETRY ANALYSIS

To further characterize Xenomouse H2BM transgenic mice, peripheral blood and spleen lymphocytes were isolated from 8-10 week old mice and controls. The cells were purified on Lympholyte M (Accurate) (San Diego, CA) and treated with purified anti-mouse CD32/CD16 Fc receptor (Pharmingen, 01241D) (San Diego, CA) to block non-specific binding to Fc receptors. Next, the cells were stained with various antibodies and analyzed on a FACStar^{PLUS} (Becton Dickinson, CELLQuest software). The panel of antibodies used to stain XenoMouse H2BM cells included: Cychrome (Cyc) anti-B220 (Pharmingen, 01128A); fluorescein isothiocyanate (FITC) anti-human IgM (Pharmingen, 34154X); FITC anti-mouse IgM (Pharmingen, 02204D).

Lymphocytes from four animals from three different XenoMouse H2BM strains were evaluated and compared to wild type B6/129 mice using flow cytometry, as shown in Table 5 below.

Strains XM-2BM-1, XM-2BM-2 and XM-2BM-6 showed about a 60-80% reconstitution in the B-cell compartment compared to wild-type mice (Table 5). Transgenic mice having the yH2BM YAC DNA show significant human antibody and immune system development.

- 87 -

Table 5

	strain	% of B220 ⁺ IgM ⁺	AVG. % B CELLS
5	XM-2BM-1 #1	34	24
	XM-2BM-1 #2	24	
	XM-2BM-1 #3	19	
	XM-2BM-1 #4	17	
10	XM-2BM-2 #1	28	28
	XM-2BM-2 #2	27	
	XM-2BM-2 #3	29	
	XM-2BM-2 #4	27	
15	XM-2BM-6 #1	27	25
	XM-2BM-6 #2	28	
	XM-2BM-6 #3	22	
	XM-2BM-6 #4	23	
	WT B6X129 #1	40	39
	WT B6X129 #2	38	
	WT B6X129 #3	35	
	WT B6X129 #4	43	

EXAMPLE 10

20 SERUM LEVELS OF HUMAN ANTIBODIES
 IN UNIMMUNIZED MICE

An ELISA for determination of human antibodies in unimmunized mouse serum was carried out. For more detailed information and procedures on immunoassays see E. Harlow et al., *Antibodies: A Laboratory Manual*, Chapter 14, "Immunoassay", pages 553-614, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). The concentration of human immunoglobulins were determined using the following capture antibodies: mouse anti-human IgM (CGI/ATCC, HB-57) (Manassas, VA). The detection antibodies used in ELISA experiments were mouse anti-human IgG1-HRP (Southern Biotechnology, 9050-05) (Birmingham, AL), mouse anti-human IGM-HRP (Southern

- 88 -

Biotechnology, 9020-05) (Birmingham, AL). Standards used for quantitation of human Ig were: human IgM κ (Cappel, 13000) (Costa Mesa, CA) and human IgG1 (Calbiochem 400126) (San Diego, CA).

- 5 As shown in Table 6, XenoMouse H2BM mice produced significant baseline levels of both human IgM and human IgG in the absence of immunization.

Table 6

10

Quantitation of hIgY1 and hIgM in XenoMouse H2BM			
		IgY1	IgM
		$\mu\text{g/ml}$	$\mu\text{g/ml}$
XM-2BM-7	H-781-2	298	140
	H-850-1	172	101
	H-850-2	250	110
XM-2BM-1	H-908-1	1.3	70
	H-908-5	0.35	51
	H-953-8	3.7	81
XM-2BM-2	H-873-2	0.8	38
	H-873-3	1.5	52
	H-873-4	1.7	90
XM-2BM-6	H-910-4	1	68
	H-911-3	0.8	47
	H-912-4	0.3	44

15 EXAMPLE 11

IMMUNIZATION AND HYBRIDOMA GENERATION

- Groups of six 8 to 10 weeks old XenoMice H2BM were immunized subcutaneously at the base of the tail (or other route of administration (IP, footpad, etc.) with 10 μg of either recombinant human IL-8, 5 μg TNF- α or CEM cells (for CD147). The antigen is emulsified in complete Freund's adjuvant for the primary immunization and in incomplete Freund's adjuvant for the additional

- 89 -

immunization. For more detailed information and procedures on animal immunizations see E. Harlow et al., *Antibodies: A Laboratory Manual*, Chapter 5, "Immunizations" pages 53-138, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). Immunizations are carried out at 3-4 week intervals for at least 3 booster immunizations (boosts).

When making monoclonal antibodies, the mice receive a final injection of antigen or cells in PBS four days before the fusion. For more detailed information and procedures on making monoclonal antibodies see E. Harlow et al., *Antibodies: A Laboratory Manual*, Chapter 6, "Monoclonal Antibodies", pages 139-244, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). Lymph node lymphocytes from immunized mice are fused with the non-secretory myeloma NSO cell line [S. Ray, et al., *Proc. Natl. Acad. Sci. USA*, 91:5548-5551 (1994)] or the P3-X63-Ag8.653 myeloma and are subjected to HAT selection as previously described [G. Galfre, et al., *Methods Enzymol.* 73:3-46 (1981)].

Table 7 shows that transgenic mice produced according to Examples 6-8 above, and immunized with recombinant human IL-8, 5 μ g TNF- α or CEM cells (for CD147) yielded human IgG1 monoclonal antibodies.

EXAMPLE 12

EVALUATION OF ANTIBODY SPECIFICITY AND ISOTYPE

We performed an ELISA for the determination of whether transgenic mice were producing antigen-specific antibodies (Table 7). We determined the antigen specificity and isotype of antibodies

- 90 -

isolated from mouse serum and from hybridoma supernatants as described [Coligan et al., Unit 2.1, "Enzyme-linked immunosorbent assays," in *Current Protocols in Immunology* (1994).], using recombinant human IL-8, CD147 and TNF- α to capture the antigen-specific antibodies. We determined the concentration of human immunoglobulins using the following capture antibodies: rabbit anti-human IgG (Southern Biotechnology, 6145-01). The detection antibodies used in ELISA experiments were mouse anti-human IgG1-HRP (Caltag, MH1015) (Burlingame, CA), mouse anti-human IGM-HRP (Southern Biotechnology, 9020-05), and goat anti-human kappa-biotin (Vector, BA-3060). Standards used for quantitation of human and mouse Ig are: human IgG₁ (Calbiochem, 400122), human IgM κ (Cappel, 13000), human IgG₂ κ (Calbiochem, 400122), mouse IgG κ (Cappel 55939), mouse IgM κ (Sigma, M-3795), and mouse IgG₄ λ (Sigma, M-9019).

Table 7 further shows that transgenic mice produced according to Examples 6-8 above, and immunized with recombinant human IL-8, 5 μ g TNF- α or CEM cells (for CD147) yielded human IgG1 monoclonal antibodies that were antigen specific and of the predicted isotype.

- 91 -

Table 7

Fusion	# animals	XM mice yH2BM mice =G1	# cells x 10 ⁶	# Hybridomas			
				IgG2	IgG4	IgG1	IgM
	IL-8						
(CEM21)CD147	10	G1-1	325			1	1
(CEM22)CD147	14	G1-1	198				15
IL8-16	10	G1-5	243			20	5
IL8-17	12	G1-6	268			31	5
IL8-18	10	G1-1	213			10	
IL8-19	10	G1-2	136			4	
	TNF- α						
TNF-38	10	G1-1/2	179 ing.			1	3
			116 popl.			2	6
TNF-39	10	G1-6	180			4	6

EXAMPLE 13

INTRODUCTION OF THE yH2CM YAC INTO ES CELLS

We introduced the YAC, yH2CM, into mouse embryonic stem (ES) cells by yeast spheroplast fusion as described in detail in Example 6. [See B. Birren *et al.*, *Genome Analysis: A Laboratory Manual*, Volume 3, Cloning Systems, Chapter 5: "Introduction of YACs into mammalian cells by spheroplast fusion", pages 548-550, Cold Spring Harbor Laboratory Press, Plainview, NY].

Generally, yH2CM containing yeast cells were spheroplasted using zymolase 20T at 1.5 mg/ml. The yH2CM spheroplasts were fused with HPRT-deficient E 14.TG3B1 mouse ES cells which had been cultured as described [see Jakobovits *et al.*, *Nature* 362:255-258 (1993); Green *et al.*, *Nature Genetics* 7:13-21 (1994); E. Robertson in *Teratocarcinomas and Embryonic Stem Cells*, pages 71-112, IRL, Oxford (1987)] HAT selection

- 92 -

was initiated 48 hours after fusion. HPRT-positive ES cell clones were selected at a frequency of 1 clone/15-20x 10⁶ fused cells. HAT resistant colonies were expanded for genome analysis and were analyzed for YAC integrity by Southern and CHEF blot analyses. In control experiments with mock fusions of ES cells and yeast spheroplasts, no colonies were detected.

Ten ES cell clones (referred to as Clones 1-10 in Table 8) derived from ES cell fusion with yH2CM-containing yeast contained all expected EcoRI and BamHI yH2 fragments detected by probes spanning the entire insert. As shown in Table 8, we detected the following human genes in the ES cell genome as part of characterization of the ES cell DNA prior to transgenic mouse generation: all the different V_H families could be detected V_H1, V_H2, V_H3, V_H4, V_H5, and V_H6; human D_H, and J_H; human C_μ and C_δ constant regions; mouse switch γ1 (mSγ1) and human Cγ4 C_H exons.

- 93 -

Table 8

h or 'm genes	ES Cell Clone									
	1	2	3	4	5	6	7	8	9	10
5	V _H 1	+	+	+	+	+	+	+	+	+
	V _H 2	+	+	+	+	+	+	+	+	+
	V _H 3	+	+	+	+	+	+	+	+	+
	V _H 4	+	+	+	+	+	+	+	+	+
	V _H 5	+	+	+	+	+	+	+	+	+
10	V _H 6	+	+	+	+	+	+	+	+	+
	D _H	+	+	+	+	+	+	+	+	+
	J _H	+	+	+	+	+	+	+	+	+
	C _μ	+	+	+	+	+	+	+	+	+
	C _δ	+	+	+	+	+	+	+	+	+
15	mSy1	+	+	+	+	+	+	+	+	+
	Cy4	+	+	+	+	+	+	+	+	+

'm - denotes mouse genes (mSy1)

EXAMPLE 14INTRODUCTION OF ES CELLS CONTAINING
THE yH2CM YAC INTO MICE

20 In order to generate chimeric mice from the
YAC yH2CM DNA containing ES cells, microinjection of
blastocysts was conducted, followed by breeding. ES
cells containing the YAC yH2CM DNA were isolated as
described in Example 6, and expanded for the generation
25 of chimeric mice. Next, yH2CM-bearing ES cell clones
were microinjected into mouse C57Bl/6 blastocysts [See
B. Hogan et al., "Manipulating the Mouse Embryo: A
Laboratory Manual", Section D, Introduction of New
Genetic Information, "Injection of Cells into the
30 Blastocyst" pages 188-196, (1986) (Cold Spring Harbor

- 94 -

Laboratory Press, Cold Spring Harbor, NY)]. Chimeric offspring were identified by coat color.

Table 9

Clone	# embryos Injected	# live born pups	# of chimeras	# chimera breedings	# germline
2CM-1	381	26	24	8	4
2CM-2	399	26	14	10	5
2CM-3	224	21	10	5	0
2CM-4	217	21	14	4	0
2CM-5	276	19	15	9	0
2CM-6	296	18	12	2	0
2CM-7	269	22	16	6	0
2CM-8	133	12	12	9	0
2CM-9	177	5	3	1	0

Table 9 summarized the data for generating transgenic mice using nine different yH2CM containing ES cell lines. Two out of nine clones were transmitted through the mouse germline.

EXAMPLE 15

**BREEDING MICE CONTAINING yH2CM YAC
DNA WITH yK2:DI MICE**

In order to generate mice that produced human antibodies in the absence of endogenous antibodies, yK2-transgenic mice were previously bred with double-inactivated (DI) mouse strains. The DI mouse strains were homozygous for gene targeted-inactivated mouse heavy and kappa chain loci and thus were deficient in antibody production [see Jakobovits *et al.*, *Nature* 362:255-258 (1993); Green *et al.*, *Nature Genetics* 7:13-21 (1994)]. One of the yK2-transgenic mouse strains, J23.1, was bred with DI mice to generate mice hemizygous or homozygous for yK2 YACs on a

- 95 -

homozygous inactivated mouse heavy and kappa chain background (yK2;DI). The breeding scheme for generating a new Xenomouse, which is hemizygous for the yH2CM YAC is shown below. Subsequent breeding of

5 XenoMouse males to XenoMouse females yields XenoMouse progeny who are either hemizygous or homozygous for yH2CM and/or yK2. From these progeny, breeding of males and females, both of which are homozygous for both yH2CM and yK2, will yield a true breeding line of

10 XenoMouse H2CM.

XenoMouse H2CM Breeding scheme

Generation 1 : (Chimera or Transgenic bred to YK2;DI)

yH2CM⁺;yK2⁻;mJ_H^{+/+};mCκ^{+/+} X yH2CM⁻;yK2⁺; mJ_H^{-/-};mCκ^{-/-}

Generation 2 : (Xenohet x YK2;DI)

15 yH2CM⁺;yK2⁺;mJ_H^{+/-};mCκ^{+/-} X yH2CM⁻;yK2⁺;mJ_H^{-/-};mCκ^{-/-}

Generation 3 (Almost Xenomouse x yK2;DI)
or Xenomouse x yK2;DI)

yH2CM⁺;yK2⁺;mJ_H^{+/-};mCκ^{-/-} X yH2CM⁻;yK2⁺;mJ_H^{-/-};mCκ^{-/-}

yH2CM⁺;yK2⁺;mJ_H^{-/-};mCκ^{-/-} X yH2CM⁻;yK2⁺;mJ_H^{-/-};mCκ^{-/-}

20 XenoMouse: yH2CM⁺; yK2⁺; mJ_H^{-/-}; mCκ^{-/-}

The integrity of the human heavy and kappa chain YACs in XenoMouse H2CM strains was confirmed by Southern blot analysis. In all XenoMouse H2CM strains analyzed, yH2CM was transmitted unaltered through

25 multiple generations with no apparent deletions or rearrangements.

- 96 -

EXAMPLE 16

FLOW CYTOMETRY ANALYSIS

To further characterize Xenomouse H2CM transgenic mice, peripheral blood and spleen lymphocytes were isolated from 8-10 week old mice and controls. The cells were purified on Lympholyte M (Accurate) (San Diego, CA) and treated with purified anti-mouse CD32/CD16 Fc receptor (Pharmingen, 01241D) (San Diego, CA) to block non-specific binding to Fc receptors. Next, the cells were stained with various antibodies and analyzed on a FACStar^{PLUS} (Becton Dickinson, CELLQuest software). The panel of antibodies used to stain XenoMouse H2CM cells included: Cychrome (Cyc) anti-B220 (Pharmingen, 01128A); fluorescein isothiocyanate (FITC) anti-human IgM (Pharmingen, 34154X); FITC anti-mouse IgM (Pharmingen, 02204D).

Lymphocytes from two animals from two different XenoMouse H2CM strains were evaluated and compared to wild type B6/129 mice using flow cytometry as shown in Table 10 below.

Strain XM2Cm-2 homo showed about a 80-100% reconstitution in the B-cell compartment (Table 10). Transgenic mice having the yH2CM YAC DNA show significant human antibody and immune system development. Control 129xB6, DI, Xenomouse 2a heterozygous and homozygous were compared to mice heterozygous and homozygous for the yH2CM YAC.

- 97 -

Table 10

	ID	%B220 ⁺ _μ ⁺
	129xB6	22.2
	129xB6	24.8
5	129xB6	24.5
	DI	0.6
	XM2A-5 het	29.2
	XM2A-5 het	23.7
	XM2A-5 homo	23.4
10	XM2A-5 homo	25.5
	XM2Cm-2 het	19.3
	XM2Cm-2 het	19.2
	XM2Cm-2 homo	29.8
	XM2Cm-2 homo	23.6
15	Avg. 129xB6	23.8 ± 1.4
	DI	0.6
	Avg. XM2A-5 het	26.5 ± 3.9
	Avg. XM2A-5 homo	24.5 ± 1.5
	Avg. XM2Cm-2 het	19.3 ± 0.1
20	Avg. XM2Cm-2 homo	26.7 ± 4.4

EXAMPLE 17**SERUM LEVELS OF HUMAN ANTIBODIES
IN UNIMMUNIZED MICE**

An ELISA for determination of human
 25 antibodies in unimmunized mouse serum was carried out.
 For more detailed information and procedures on
 immunoassays see E. Harlow et al., *Antibodies: A
 Laboratory Manual*, Chapter 14, "Immunoassay", pages

- 98 -

553-614, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). The concentration of human immunoglobulins were determined using the following capture antibodies: mouse anti-human IgM
5 (CGI/ATCC, HB-57) (Manassas, VA). The detection antibodies used in ELISA experiments were mouse anti-human IgG₄-HRP (Southern Biotechnology, 9050-05) (Birmingham, AL), mouse anti-human IGM-HRP (Southern Biotechnology, 9020-05) (Birmingham, AL). Standards
10 used for quantitation of human Ig were: human IgMk (Cappel, 13000) (Costa Mesa, CA) and human IgG1 (Calbiochem 400126) (San Diego, CA).

As shown in Table 11, rows 15-30, Xenomouse H2CM produced significant baseline levels of both human
15 IgM and IgG4 in the absence of immunization.

- 99 -

Table 11

	Mouse ID	hlgM ($\mu\text{g/ml}$)	hlgG2 ($\mu\text{g/ml}$)	hlgG4 ($\mu\text{g/ml}$)
5	1 129 x B6			
	2 129 x B6			
	3 129 x B6			
	4 DI			
	5 DI			
	6 DI			
10	7 XM2A-5	89.5	37.6	
	8 XM2A-5	97.1	37.6	
	9 XM2A-5	98.0	409.7	
	10 XM2A-5	85.1	18.2	
	11 XM2A-5	72.0	423.1	
	12 XM2A-5	74.3	273.3	
15	13 XM2A-5	98.6	16.8	
	14 XM2A-5	126.8	28.8	
	Xenomouse H2CM			
	15 XM2Cm-1	109.4		33.2
	16 XM2Cm-1	83.6		187.1
	17 XM2Cm-1	84.9		665.3
20	18 XM2Cm-1	88.7		61.3
	19 XM2Cm-1	93.1		177.2
	20 XM2Cm-1	79.4		36.9
	21 XM2Cm-1	80.4		91.2
	22 XM2Cm-1	76.9		238.6
	23 XM2Cm-2	35.2		20.9
25	24 XM2Cm-2	35.4		88.8
	25 XM2Cm-2	28.0		42.5
	26 XM2Cm-2	25.0		20.6
	27 XM2Cm-2	66.8		23.0
	28 XM2Cm-2	28.1		14.8
	29 XM2Cm-2	27.3		30.1
30	30 XM2Cm-2	32.6		69.3
	129xB6	N.D.	N.D.	N.D.
	DI	N.D.	N.D.	N.D.
	XM2A-5	92.7 $\pm$ 17.2	155.6 $\pm$ 182.1	N.D.
	XM2Cm-1	87.1 $\pm$ 10.4	N.D.	186.4 $\pm$ 208.0
	XM2Cm-2	34.8 $\pm$ 13.5	N.D.	38.8 $\pm$ 26.7

- 100 -

EXAMPLE 18

IMMUNIZATION AND HYBRIDOMA GENERATION

Groups of six 8 to 10 weeks old XenoMice H2CM were immunized subcutaneously at the base of the tail with 10 µg of either recombinant human IL-6 or IL-8. The antigen is emulsified in complete Freund's adjuvant for the primary immunization and in incomplete Freund's adjuvant for the additional immunizations. For more detailed information and procedures on animal immunizations see E. Harlow et al., *Antibodies: A Laboratory Manual*, Chapter 5, "Immunizations" pages 53-138, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). Immunizations are carried out at 3-4 week intervals for at least 3 booster immunizations (boosts).

When making monoclonal antibodies, the mice receive a final injection of antigen or cells in PBS four days before the fusion. For more detailed information and procedures on making monoclonal antibodies see E. Harlow et al., *Antibodies: A Laboratory Manual*, Chapter 6, "Monoclonal Antibodies", pages 139-244, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). Lymph node lymphocytes from immunized mice fused with the non-secretory myeloma NSO cell line [S. Ray, et al., *Proc. Natl. Acad. Sci. USA*, 91:5548-5551 (1994)] or P3-X63-Ag8.653 myeloma cells and subjected to HAT selection as previously described [G. Galfre, et al., *Methods Enzymol.* 73:3-46 (1981)].

Table 12 shows that transgenic mice produced according to Examples 14-16 above, and immunized with

- 101 -

recombinant human IL-6 or IL-8, yielded human IgG4 monoclonal antibodies.

EXAMPLE 19EVALUATION OF ANTIBODY SPECIFICITY AND ISOTYPE

5 We performed an ELISA to determine whether transgenic mice were producing antigen-specific antibodies (Table 12). We also determined the human antibody isotype produced (Table 12). Antigen specificity and isotype determinations were performed
10 on antibodies isolated from mouse serum and from hybridoma supernatants as described [Coligan et al., Unit 2.1, "Enzyme-linked immunosorbent assays," in *Current protocols in immunology* (1994).] using recombinant human IL-6 or IL-8, to capture the antigen-
15 specific antibodies. The concentration of human and mouse immunoglobulins were determined using the following capture antibodies: rabbit anti-human IgG (Southern Biotechnology, 6145-01). The detection antibodies used in ELISA experiments was mouse
20 anti-human IgG1-HRP (Caltag, MH1015) (Burlingame, CA), mouse anti-human IGM-HRP (Southern Biotechnology, 9020-05), and goat anti-human kappa-biotin (Vector, BA-3060). Standards used for quantitation of human and mouse Ig were: human IgG₁ (Calbiochem, 400122), human
25 IgM_k (Cappel, 13000), human IgG₂k (Calbiochem, 400122), mouse IgGk (Cappel 55939), mouse IgMk (Sigma, M-3795), and mouse IgG₄λ (Sigma, M-9019).

Table 12 further shows that transgenic mice produced according to Examples 14-16 above, and
30 immunized with recombinant human IL-6 or IL-8 yielded

- 102 -

human IgG4 monoclonal antibodies that were antigen specific and of the predicted isotype.

Table 12

Fusion	# animals	XM mice yH2GM mice =G4	# cells x 10 ⁶	# Hybridomas			
				IgG2	IgG4	IgG1	IgM
IL6-1	12	G4-2	132		13		
IL8-13	10	G4 1-2	66		27		
IL8-14	10	G4 1-2	41		7		
IL8-15		G4 1-2	41		14		

EXAMPLE 20

10 INTRODUCTION OF THE yHG1/2 YAC INTO ES CELLS

We introduced the YAC, yHG1/2, into mouse embryonic stem (ES) cells by yeast spheroplast fusion as described in Example 6. [See B. Birren et al., *Genome Analysis: A Laboratory Manual*, Volume 3, Cloning Systems, Chapter 5: "Introduction of YACs into mammalian cells by spheroplast fusion", pages 548-550, Cold Spring Harbor Laboratory Press, Plainview, NY]. Generally, yHG1/2 containing yeast cells were spheroplasted using zymolase 20T at 0.15 mg/ml. The yHG1/2 spheroplasts were fused with HPRT-deficient E 14.TG3B1 mouse ES cells which were cultured as described [see Jakobovits et al., *Nature* 362:255-258 (1993); Green et al., *Nature Genetics* 7:13-21 (1994); E. Robertson in *Teratocarcinomas and Embryonic Stem Cells*, pages 71-112, IRL, Oxford (1987)] HAT selection

- 103 -

was initiated 48 hours after fusion. HPRT-positive ES cell clones were selected. HAT resistant colonies are expanded for genome analysis and were analyzed for YAC integrity by Southern and CHEF blot analyses. Control experiments were "mock" fusions of ES cells alone and yeast spheroplasts alone.

We examined four ES cell clones derived from ES cell fusion with yHG1/2-containing yeast using Southern blot with probes spanning the entire insert to determine whether the clones contain all expected EcoRI, HindIII and BamHI yH2 fragments. We found that all four ES cell clones contained intact YACs. We detected the following human genes in the ES cell genome as part of characterization of the ES cell DNA prior to transgenic mouse generation: all the different V_H families (V_{H1} , V_{H2} , V_{H3} , V_{H4} , V_{H5} , and V_{H6}); human D_H , and J_H ; human C_μ and C_δ constant regions; human switch $\gamma 2$ (hSy2) and human $C\gamma 1 C_H$ exons (see Figure 8 and Table 13).

TABLE 13

ES CLONE	D	μ	C γ	Cd	Sg	3'e	J	V 2	V 3	V 4	V 5	V 6
#1 Z 72.12.1	+	+	+	+	+	+	+	+	+	+	+	+
#2 Z 72.7.1	+	+	+	+	+	+	+	+	+	+	+	+
#3 Z 72.8.1	+	+	+	+	+	+	+	+	+	+	+	+
#4 Z 70.17.1	+	+	+	+	+	+	+	+	+	+	+	+

YAC Southern blot analysis of ES clones after fusion with yH3B

- 105 -

EXAMPLE 21**INTRODUCTION OF ES CELLS CONTAINING
THE yHG1/2 YAC INTO MICE**

5 To generate chimeric mice from ES cells
containing YAC yHG1/2 DNA we used blastocyst
microinjection followed by breeding. We isolated ES
cells containing the YAC yHG1/2 DNA as described in
Example 6, and expanded for the generation of chimeric
10 mice. Next, we microinjected yHG1/2-bearing ES cell
clones into mouse C57Bl/6 blastocysts [See B. Hogan et
al., *Manipulating the Mouse Embryo: A Laboratory*
Manual, Section D, Introduction of New Genetic
Information, "Injection of Cells into the Blastocyst"
15 pages 188-196, (1986) (Cold Spring Harbor Laboratory
Press, Cold Spring Harbor, NY)].

EXAMPLE 22**BREEDING MICE CONTAINING yHG1/2 YAC
DNA WITH yK2:DI MICE**

20 In order to generate mice that produce human
antibodies in the absence of endogenous antibodies,
yK2-transgenic mice are previously bred with
double-inactivated (DI) mouse strains. The DI mouse
strains are homozygous for gene targeted-inactivated
25 mouse heavy and kappa chain loci and thus are deficient
in antibody production [see Jakobovits et al., *Nature*
362:255-258 (1993); Green et al., *Nature Genetics* 7:13-
21 (1994)]. One of the yK2-transgenic mouse strains,
J23.1, is bred with DI mice to generate mice hemizygous
30 or homozygous for yK2 YACs on a homozygous inactivated
mouse heavy and kappa chain background (yK2;DI). The
breeding scheme for generating a new Xenomouse, which
is hemizygous for the yHG1/2 YAC is shown below.

- 106 -

Subsequent breeding of XenoMouse males to XenoMouse females yields XenoMouse progeny who are either hemizygous or homozygous for yHG1/2 and/or yK2. From these progeny, breeding of males and females, both of which are homozygous for both yHG1/2 and yK2, will yield a true breeding line of XenoMouse HG1/2.

XenoMouse yHG1/2 Breeding scheme

Generation 1 : (Chimera or Transgenic bred to YK2:DI)

yHG1/2⁺;yK2⁻;mJ_H^{+/+};mCk^{+/+} **X** yHG1/2⁻;yK2⁺; mJ_H^{-/-};mCk^{-/-}

10 Generation 2 : (Xenohet x YK2:DI)

yHG1/2⁺;yK2⁺;mJ_H^{+/+};mCk^{+/+} **X** yHG1/2⁻;yK2⁺;mJ_H^{-/-};mCk^{-/-}

Generation 3 (Almost Xenomouse x yK2:DI)
or Xenomouse x yK2:DI)

yHG1/2⁺;yK2⁺;mJ_H^{+/+};mCk^{-/-} **X** yHG1/2⁻;yK2⁺;mJ_H^{-/-};mCk^{-/-}

15 yHG1/2⁺;yK2⁺;mJ_H^{-/-};mCk^{-/-} **X** yHG1/2⁻;yK2⁺;mJ_H^{-/-};mCk^{-/-}

XenoMouse: yHG1/2⁺; yK2⁺; mJ_H^{-/-}; mCk^{-/-}

The integrity of the human heavy and kappa chain YACs in XenoMouse H2CM strains is confirmed by Southern blot analysis. In all XenoMouse HG1/2M strains analyzed, yHG1/2 is transmitted unaltered through multiple generations with no apparent deletions or rearrangements.

EXAMPLE 23

FLOW CYTOMETRY ANALYSIS

25 To further characterize Xenomouse HG1/2 transgenic mice, peripheral blood and spleen lymphocytes are isolated from 8-10 week old mice and

- 107 -

controls. The cells are purified on Lympholyte M (Accurate) (San Diego, CA) and treated with purified anti-mouse CD32/CD16 Fc receptor (Pharmingen, 01241D) (San Diego, CA) to block non-specific binding to Fc
5 receptors. Next, the cells are stained with various antibodies and analyzed on a FACStar^{PLUS} (Becton Dickinson, CELLQuest software). The panel of antibodies used to stain XenoMouse HG1/2M cells include: Cychrome (Cyc) anti-B220 (Pharmingen, 01128A);
10 fluorescein isothiocyanate (FITC) anti-human IgM (Pharmingen, 34154X); FITC anti-mouse IgM (Pharmingen, 02204D).

Lymphocytes from four animals from different XenoMouse H2G1/2M strains are evaluated and compared to
15 wild type B6/129 mice using flow cytometry.

Transgenic mice having the yHG1/2 YAC DNA will show significant human antibody and immune system development. Control 129xB6, DI, Xenomouse 2a heterozygous and homozygous are compared to mice
20 heterozygous and homozygous for the yHG1/2 YAC.

EXAMPLE 24

SERUM LEVELS OF HUMAN ANTIBODIES IN UNIMMUNIZED MICE

An ELISA for determination of human
25 antibodies in unimmunized mouse serum is carried out. For more detailed information and procedures on immunoassays see E. Harlow et al., *Antibodies: A Laboratory Manual*, Chapter 14, "Immunoassay", pages 553-614, Cold Spring Harbor Laboratory Press, Cold
30 Spring Harbor, New York (1988). The concentration of human immunoglobulins are determined using the

- 108 -

following capture antibodies: mouse anti-human IgM (CGI/ATCC, HB-57) (Manassas, VA). The detection antibodies used in ELISA experiments are mouse anti-human IgG1-HRP (Southern Biotechnology, 9050-05) 5 (Birmingham, AL), mouse anti-human IGM-HRP (Southern Biotechnology, 9020-05) (Birmingham, AL). Standards used for quantitation of human Ig are: human IgMx (Cappel, 13000) (Costa Mesa, CA) and human IgG1 (Calbiochem 400126) (San Diego, CA).

10 **EXAMPLE 25**IMMUNIZATION AND HYBRIDOMA GENERATION

Groups of six 8 to 10 weeks old XenoMice yHG1/2 are immunized subcutaneously at the base of the tail (or other route of administration (IP, footpad, 15 etc.) with 10 µg of antigen of choice. The antigen is emulsified in complete Freund's adjuvant for the primary immunization and in incomplete Freund's adjuvant for the additional immunizations. For more detailed information and procedures on animal 20 immunizations see E. Harlow et al., *Antibodies: A Laboratory Manual*, Chapter 5, "Immunizations" pages 53-138, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). Immunizations are carried out at 3-4 week intervals for at least 3 booster 25 immunizations (boosts).

When making monoclonal antibodies, the mice receive a final injection of antigen or cells in PBS four days before the fusion. For more detailed information and procedures on making monoclonal 30 antibodies see E. Harlow et al., *Antibodies: A*

- 109 -

Laboratory Manual, Chapter 6, "Monoclonal Antibodies", pages 139-244, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). Lymph node lymphocytes from immunized mice are fused with the
5 non-secretory myeloma NSO cell line [S. Ray, et al., *Proc. Natl. Acad. Sci. USA*, 91:5548-5551 (1994)] or the P3-X63-Ag8.653 myeloma cells and are subjected to HAT selection as previously described [G. Galfre, et al., *Methods Enzymol.* 73:3-46 (1981)].

10 **EXAMPLE 26**EVALUATION OF ANTIBODY SPECIFICITY AND ISOTYPE

To determine of whether transgenic mice are producing antigen-specific antibodies, we performed an ELISA. We also determine the human antibody isotype
15 produced. We perform antigen specificity and isotype determinations on antibodies isolated from mouse serum and from hybridoma supernatants as described [Coligan et al., Unit 2.1, "Enzyme-linked immunosorbent assays," in *Current Protocols in Immunology* (1994).], using
20 antigen to capture the antigen-specific antibodies. The concentration of human and mouse immunoglobulins are determined using the following capture antibodies: rabbit anti-human IgG (Southern Biotechnology, 6145-01). The detection antibodies used in ELISA
25 experiments are: mouse anti-human IgG1-HRP (Caltag, MH1015) (Burlingame, CA), mouse anti-human IGM-HRP (Southern Biotechnology, 9020-05), and goat anti-human kappa-biotin (Vector, BA-3060). Standards used for quantitation of human and mouse Ig are: human IgG₁
30 (Calbiochem, 400122), human IgMk (Cappel, 13000), human

- 110 -

IgG₂K (Calbiochem, 400122), mouse IgG₁K (Cappel 55939), mouse IgM₁K (Sigma, M-3795), and mouse IgG₄λ (Sigma, M-9019).

Transgenic mice produced according to
5 Examples 20-22 above, and immunized with an antigen yield human IgG₁ monoclonal antibodies that are antigen specific and of the predicted isotype.

EXAMPLE 27

INTRODUCTION OF THE yHG4 YAC INTO ES CELLS

10 We introduced the YAC, yHG4, into mouse embryonic stem (ES) cells by yeast spheroplast fusion as described in Example 6. [See B. Birren et al., *Genome Analysis: A Laboratory Manual*, Volume 3, Cloning Systems, Chapter 5: "Introduction of YACs into
15 mammalian cells by spheroplast fusion", pages 548-550, Cold Spring Harbor Laboratory Press, Plainview, NY]. Generally, yHG4 containing yeast cells were spheroplasted using zymolase 20T at 1.5 mg/ml. The yHG4 spheroplasts were fused with HPRT-deficient E
20 14.TG3B1 mouse ES cells which were cultured as described [see Jakobovits et al., *Nature* 362:255-258 (1993); Green et al., *Nature Genetics* 7:13-21 (1994); E. Robertson in *Teratocarcinomas and Embryonic Stem Cells*, pages 71-112, IRL, Oxford (1987)] HAT selection
25 was initiated 48 hours after fusion. HPRT-positive ES cell clones were selected. HAT resistant colonies were expanded for genome analysis and were analyzed for YAC integrity by Southern and CHEF blot analyses. Control experiments included "mock" fusions of ES cells alone
30 and yeast spheroplasts alone.

- 111 -

We examined eight ES cell clones derived from ES cell fusion with yHG4-containing yeast using Southern blot with probes spanning the entire insert to determine whether the clones contain all expected EcoRI and BamHI yH2 fragments. We found that the yHG4 YAC was intact in all eight clones. We detected the following human genes in the ES cell genome as part of characterization of the ES cell DNA prior to transgenic mouse generation: all the different V_H families (V_H1, V_H2, V_H3, V_H4, V_H5, and V_H6); human D_H, and J_H; human C_μ and C_δ constant regions; human switch γ2 (hSγ2) and human Cγ4 C_H exons.

EXAMPLE 28

15 INTRODUCTION OF ES CELLS CONTAINING
 THE yHG4 YAC INTO MICE

To generate chimeric mice from the YAC yHG4 DNA containing ES cells, we used microinjection of blastocysts followed by breeding. We isolated ES cells containing the YAC yHG4 DNA as described in Example 6, and expanded for the generation of chimeric mice. Next, we microinjected yHG4-bearing ES cell clones into mouse C57Bl/6 blastocysts [See B. Hogan et al., *Manipulating the Mouse Embryo: A Laboratory Manual*, Section D, Introduction of New Genetic Information, "Injection of Cells into the Blastocyst" pages 188-196, (1986) (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY)]. We identified chimeric offspring by coat color.

Germline transmission was obtained. We identified yHG4 transgenic mice by PCR using primers specific for human V6.

To generate mice that produce human antibodies in the absence of endogenous antibodies, yK2-transgenic mice were previously bred with double-inactivated (DI) mouse strains, which are homozygous for gene targeted-inactivated mouse heavy and kappa chain loci and, thus, are deficient in antibody production [see Jakobovits et al., *Nature* 362:255-258 (1993); Green et al., *Nature Genetics* 7:13-21 (1994)]. One of the yK2-transgenic mouse strains, J23.1, was bred with DI mice to generate mice hemizygous or homozygous for yK2 YACs on a homozygous inactivated mouse heavy and kappa chain background (yK2;DI). The breeding scheme used to generate a new Xenomouse, which is hemizygous for the yHG4 YAC is shown below. Subsequent breeding of hemizygous XenoMouse males to hemizygous XenoMouse females yields XenoMouse progeny that are homozygous for yHG4 and/or yK2. From these progeny, breeding of males and females, both of which are homozygous for both yHG4 and yK2, will yield a true breeding line of XenoMouseG4.

25 **Generation 1 :** (Chimera or Transgenic bred to YK2:DI)
yHG4⁺; yK2⁻; mJ_H^{+/+}; mCk^{+/-} **X** yHG4⁻; yK2⁻; mJ_H^{-/-}; mCk^{-/-}
Generation 2 : (Xenohet x YK2:DI)
yHG4⁺; yK2⁺; mJ_H^{+/-}; mCk^{+/-} **X** yHG4⁻; yK2⁺; mJ_H^{-/-}; mCk^{-/-}
Generation 3 (Almost Xenomouse x yK2:DI)
30 or Xenomouse x yK2:DI)

- 113 -

yHG4⁺; yK2⁺; mJ_H^{+/-}; mCκ^{-/-} **X** yHG4⁻; yK2⁺; mJ_H^{-/-}; mCκ^{-/-}
yHG4⁺; yK2⁺; mJ_H^{-/-}; mCκ^{-/-} **X** yHG4⁻; yK2⁺; mJ_H^{-/-}; mCκ^{-/-}

XenoMouse: yHG4⁺; yK2⁺; mJ_H^{-/-}; mCκ^{-/-}

The integrity of the human heavy and kappa
5 chain YACs in XenoMouse G4 strains was confirmed by
Southern blot analysis. In all XenoMouse G4 strains
analyzed, yHG4 was transmitted unaltered through
multiple generations with no apparent deletions or
rearrangements.

10 EXAMPLE 30

FLOW CYTOMETRY ANALYSIS

To further characterize Xenomouse G4
transgenic mice, peripheral blood and spleen
lymphocytes are isolated from 8-10 week old mice and
15 controls. The cells are purified on Lympholyte M
(Accurate) (San Diego, CA) and treated with purified
anti-mouse CD32/CD16 Fc receptor (Pharmingen, 01241D).
(San Diego, CA) to block non-specific binding to Fc
receptors. Next, the cells are stained with various
20 antibodies and analyzed on a FACStar^{PLUS} (Becton
Dickinson, CELLQuest software). The panel of
antibodies used to stain XenoMouse G4 cells include:
Cychrome (Cyc) anti-B220 (Pharmingen, 01128A);
fluorescein isothiocyanate (FITC) anti-human IgM
25 (Pharmingen, 34154X); FITC anti-mouse IgM (Pharmingen,
02204D).

Lymphocytes from four animals from three
different XenoMouse G4 strains are evaluated and
compared to wild type B6/129 mice using flow cytometry.

- 114 -

Trangenic mice having the G4 YAC DNA will show significant human antibody and immune system development. Control 129xB6, DI, Xenomouse 2a heterozygous and homozygous are compared to mice
5 heterozygous and homozygous for the G4 YAC.

EXAMPLE 31**SERUM LEVELS OF HUMAN ANTIBODIES
IN UNIMMUNIZED MICE**

An ELISA for determination of human
10 antibodies in unimmunized mouse serum is carried out. For more detailed information and procedures on immunoassays see E. Harlow et al., *Antibodies: A Laboratory Manual*, Chapter 14, "Immunoassay", pages 553-614, Cold Spring Harbor Laboratory Press, Cold
15 Spring Harbor, New York (1988). The concentration of human immunoglobulins are determined using the following capture antibodies: mouse anti-human IgM (CGI/ATCC, HB-57) (Manassas, VA). The detection
antibodies used in ELISA experiments are mouse
20 anti-human IgG1-HRP (Southern Biotechnology, 9050-05) (Birmingham, AL), mouse anti-human IGM-HRP (Southern Biotechnology, 9020-05) (Birmingham, AL). Standards used for quantitation of human Ig are: human IgMk (Cappel, 13000) (Costa Mesa, CA) and human IgG1
25 (Calbiochem 400126) (San Diego, CA).

- 115 -

EXAMPLE 32

IMMUNIZATION AND HYBRIDOMA GENERATION

Groups of six 8 to 10 weeks old XenoMice yHG4 are immunized subcutaneously at the base of the tail or other route of administration (IP, footpad, etc.) with 10 µg of antigen. The antigen is emulsified in complete Freund's adjuvant for the primary immunization and in incomplete Freund's adjuvant for the additional immunizations. For more detailed information and procedures on animal immunizations see E. Harlow et al., *Antibodies: A Laboratory Manual*, Chapter 5, "Immunizations" pages 53-138, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). Immunizations are carried out at 3-4 week intervals for at least 3 booster immunizations (boosts).

When making monoclonal antibodies, the mice receive a final injection of antigen or cells in PBS four days before the fusion. For more detailed information and procedures on making monoclonal antibodies see E. Harlow et al., *Antibodies: A Laboratory Manual*, Chapter 6, "Monoclonal Antibodies", pages 139-244, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York (1988). Lymph node lymphocytes from immunized mice are fused with the non-secretory myeloma NSO line [S. Ray, et al., *Proc. Natl. Acad. Sci. USA*, 91:5548-5551 (1994)] or the P3-X63-Ag8.653 myeloma and are subjected to HAT selection as previously described [G. Galfre, et al., *Methods Enzymol.* 73:3-46 (1981)].

- 116 -

EXAMPLE 33EVALUATION OF ANTIBODY SPECIFICITY AND ISOTYPE

An ELISA for the determination of whether transgenic mice are producing antigen-specific antibodies is performed. It is further desired to confirm the human antibody isotype produced. Antigen specificity and isotype determination are performed on antibodies isolated from mouse serum and from hybridoma supernatants as described [Coligan et al., Unit 2.1, "Enzyme-linked immunosorbent assays," in *Current protocols in immunology* (1994).] using recombinant antigen to capture the antigen-specific antibodies. The concentration of human and mouse immunoglobulins are determined using the following capture antibodies:

15 rabbit anti-human IgG (Southern Biotechnology, 6145-01). The detection antibodies used in ELISA experiments is mouse anti-human IgG1-HRP (Caltag, MH1015) (Burlingame, CA), mouse anti-human IGM-HRP (Southern Biotechnology, 9020-05), and goat anti-human kappa-biotin (Vector, BA-3060). Standards used for

20 quantitation of human and mouse Ig are: human IgG₁ (Calbiochem, 400122), human IgMκ (Cappel, 13000), human IgG₂κ (Calbiochem, 400122), mouse IgGκ (Cappel 55939), mouse IgMκ (Sigma, M-3795), and mouse IgG₄λ (Sigma,

25 M-9019).

Transgenic mice produced according to Examples 27-29 above, and immunized with antigen yield human IgG₄ monoclonal antibodies that are antigen specific and of the predicted isotype.

- 117 -

Biological Materials:

The following biological materials are disclosed and discussed in connection with the above Examples and are exemplary of materials that can be utilized and prepared in accordance with the present invention:

- ppKM1C (yH1C targeting vector) (
- p1B (targeting vector)
- TV1 (mSg1-hCg1 plasmid DNA vector for targeting yH1C to make yH2Bm)
- 10 • TV4 (mSg1-hCg4 plasmid DNA vector for targeting yH1C to make yH2Cm)
- TV G1 (hCg1 plasmid DNA vector for targeting yH1C to make yHG1)
- TV G4 (hCg1 plasmid DNA vector for targeting yH1C to make yHG4)
- 15 • yH2Cm (mSg1-hCg4 YAC) (deposited with the ATCC on _____ and having accession number _____)
- yH2Bm (mSg1-hCg1 YAC) (deposited with the ATCC on _____ and having accession number _____)
- 20 • yHG1 (hSg2-hCg1 YAC)
- yHG4 (also referred to as yH3C) (hSg2-hCg4 YAC) (deposited with the ATCC on _____ and having accession number _____)
- yH3B (also referred to as yHG1/2) (hSg2-hCg1-hCg2(TM) YAC) (deposited with the ATCC on _____ and having accession number _____)
- 25 • ES-yH2Cm clone 1
- ES-yH2Cm clone 2
- ES-yH2Bm clone 1
- 30 • ES-yH2Bm clone 2
- ES-yH2Bm clone 3
- ES-yH2Bm clone 4

- 118 -

- ES-yH2Bm clone 5
- ES-yH2Bm clone 6
- ES-yH2Bm clone 7
- ES-yH2Bm clone 8
- 5 • ES-yH2Bm clone 9

- 119 -

INCORPORATION BY REFERENCE

All references cited herein, including patents, patent applications, papers, text books and the like, and the references cited therein, to the extent that they are not already, are hereby incorporated by reference in their entirety. In addition, the following references are also incorporated by reference herein in their entirety, including references cited in such references.

- | | |
|----|--|
| 10 | M.J. Mendez <u>et al.</u> , "Functional Transplant of Megabase Human Immunoglobulin Loci Recapitulates Human Antibody Response in Mice," <i>Nature Genetics</i> , 15:146-156 (1997) |
| 15 | G.T. Williams <u>et al.</u> , "Membrane Immunoglobulin Without Sheath or Anchor," <i>Molecular Immunology</i> , 30:1427-1432, (1993) |
| 20 | G.T. Williams <u>et al.</u> , "The α/β Sheath and Its Cytoplasmic Tyrosine Are Required For Signaling By The B-cell Antigen Receptor But Not for Capping or For Serine/Threonine-Kinase Recruitment," <i>Immunology</i> , 91:474-478 (1994) |
| 25 | A.R. Venkitaraman <u>et al.</u> , "The B-cell Antigen Receptor of the Five Immunoglobulin Classes," <i>Nature</i> , 352:777-781 (1991) |
| 30 | G.T. Williams <u>et al.</u> , "The Sequence of The μ Transmembrane Segment Determines the Tissue Specificity of the Transport of Immunoglobulin M to The Cell Surface," <i>J. Exp. Med.</i> 171:947-952 (1990) |
| | S. Pettersson <u>et al.</u> , "A Second B cell-specific Enhancer 3' of The Immunoglobulin Heavy-chain Locus," <i>Nature</i> , 344:165-168 (1990) |

- 120 -

- P. Dariavach et al., "The Mouse IgH 3' -Enhancer," *Eur. J. Immunol.*, 21:1499-1504 (1991)
- 5 G.P. Cook et al., "Regulated Activity of the IgH Intron Enhancer (Ep) in the T Lymphocyte Lineage," *International Immunology*, 7:89-95 (1995)
- K.B. Meyer et al., "The Igk 3' -Enhancer Triggers Gene Expression in Early B Lymphocytes but Its Activity is Enhanced on B cell Activation," *International Immunology*, 8:1561-1568 (1996)
- 10 M.S. Neuberger et al., "Recombinant Antibodies Possessing Novel Effector Functions," *Nature*, 312:604-608 (1984)
- C.I. Bindon et al., "Human Monoclonal IgG Isotypes Differ in Complement Activating Function at the Level of C4 As Well AS Clq," *J. Exp. Med.*, 168:127-142 (1988)
- 15 S. Huck et al., "Sequence of a Human Immunoglobulin Gamma 3 Heavy Chain Constant Region Gene: Comparison With the Other Human Cy Genes," *Nucleic Acids Research*, 14: 1779-1789 (1986)
- 20 J. Ellison et al., "Nucleotide Sequence of a Human Immunoglobulin Cy₄ Gene," *DNA*, 1: 11-18 (1981)
- J.W. Ellison et al., "The Nucleotide Sequence of a Human Immunoglobulin Cy₁ Gene," *Nucleic Acids Research*, 10:4071-4079 (1982)
- 25 J. Ellison et al., "Linkage and Sequence Homology of Two Human Immunoglobulin γ Heavy Chain Constant Region Genes," *Immunology*, 79:1984-1988 (1982)
- H. Hayashida et al., "Concerted Evolution of the Mouse Immunoglobulin Gamma Chain Genes," *The EMBO Journal*, 3:2047-2053 (1984)
- 30

- 121 -

- A. Jakobovits et al., "Analysis of Homozygous Mutant Chimeric Mice: Deletion of the Immunoglobulin Heavy-chain Joining Region Blocks B-cell Development and Antibody Production," *Genetics*, 90:2551-2555 (1993)
- 5 A. Jakobovits, "The Long-Awaited Magic Bullets: Therapeutic Human Monoclonal Antibodies From Transgenic Mice," *Exp. Opin. Invest. Drugs*, 7:607-614 (1998)
- 10 P.T. Jones et al., "Replacing The Complementarity-determining Regions in a Human Antibody with Those From a Mouse," *Nature*, 321:522-525 (1986)
- M.S. Neuberger et al., "Isotype Exclusion and Transgene Down-regulation in Immunoglobulin- λ Transgenic Mice," *Nature*, 338:350-352 (1989)
- 15 M. Brüggenmann et al., "(A Repertoire of Monoclonal Antibodies with Human Heavy Chains from Transgenic Mice," *Immunology*, 86:6709-6713 (1989)
- 20 C.J. Jolly et al., "Rapid Methods for the Analysis of Immunoglobulin Gene Hypermutation: Application to Transgenic and Gene Targeted Mice," *Nucleic Acids Research*, 25:1913-1919 (1997)
- C. J. Jolly et al., "The Targeting of Somatic Hypermutation," *Immunology*, 8:159-168 (1996)
- 25 S. Pettersson et al., "Cellular Selection Leads to Age-Dependent and Reversible Down-regulation of Transgenic Immunoglobulin Light Chain Genes," *International Immunology*, 1:509-516 (1989)
- 30 K.B. Meyer et al., "The Igk 3' -Enhancer Triggers Gene Expression in Early B Lymphocytes but its Activity is enhanced on B Cell Activation," *International Immunology*, 8:1561-1568 (1996)

- 122 -

- G.P. Cook et al., "Regulated Acticity of the IgH Intron Enhancer (Ep) in the T Lymphocyte Lineage," *International Immunology*, 7:89-95 (1995)
- 5 C.J. Jolly et al., "The Targeting of Somatic Hypermutation," *Immunology*, 8:159-168 (1996)
- S. Pettersson et al., "Cellular Selection Leads to Age-dependent and Reversible Down-regulation of Transgenic Immunoglobulin Light Chain Genes," *International Immunology*, 1:509-516 (1989)
- 10 L.E. Reid et al., "A Single DNA response element can confer inducibility by both α - and γ -interferon," *Biochemistry*, 86:840-844 (1989)
- J. Stavnezer et al., "Immunoglobulin Heavy-chain Switching May be Directed by Prior Induction of Transcripts from Constant-region genes," *Immunology*, 85:7704-7708 (1988)
- 15 F.C. Mills et al., "Sequences of Human Immunoglobulin Switch Regions: Implications for Recombination and Transcription," *Nucleic Acids Research*, 18:7305-7316 (1990)
- 20 P. Rothman et al., (Structure and Expression of Germline Immunoglobulin $\gamma 3$ Heavy Chain Gene Transcripts: Implications for Mitogen and Lymphokine Directed class-switching," *International Immunology*, 2:621-627 (1990)
- 25 P. Sideras et al., "Production of Sterile Transcripts of Cy Genes in an IgM-producing Human Neoplastic B Cell Line that Switches to IgG-producing Cells," *International Immunology*, 1:632-642 (1989)
- 30 J.P. Manis et al., "Class Switching in B Cells Lacking 3' Immunoglobulin Heavy Chain Enhancers," *J. Exp. Med.*, 188: 1421-1431 (1998)

- 123 -

- J. Durdik et al., "Isotype Switching by a Microinjected μ Immunoglobulin Heavy Chain Gene in Transgenic Mice," *Immunology*, 86:2346-2350 (1989)
- 5 N. Takahashi et al., "Structure of Human Immunoglobulin Gamma Genes: Implications for Evolution of a Gene Family," *Cell*, 29:671-679 (1982)
- 10 J. Zhang et al., "A Selective Defect in IgG2b Switching as a Result of Targeted Mutation of the I γ 2b Promoter and Exon," *The EMBO Journal*, 12:3529-3537: (1993)
- S. Jung et al., "Shutdown of Class Switch Recombination by Deletion of a Switch Region Control Element," *Science*, 259: 984-987 (1993)
- 15 L. Xu et al., "Replacement of Germ-line ϵ promoter by Gene Targeting Alters Control of Immunoglobulin Heavy Chain Class Switching," *Immunology*, 90:3705-3709 (1993)
- 20 A. Bottaro et al., "S Region Transcription *per se* Promotes Basal IgE Class Switch Recombination But Additional Factors Regulate the Efficiency of the Process," *The EMBO Journal*, 13:665-674 (1994)
- F.C. Mills et al., "Human IgS γ Regions and Their Participation in Sequential Switching to IgE," *The Journal of Immunology*, 155:3021-3036 (1995)
- 25 S.C. Li et al., "Expression of I μ -C γ Hybrid Germline Transcripts Subsequent to Immunoglobulin Heavy Chain Class Switching," *International Immunology*, 6:491-497 (1994)
- 30 Q. Pan et al., "Characterization of Human γ 4 Switch Region Polymorphisms Suggest a Meiotic Recombinational Hot Spot Within the Ig Locus: Influence of S Region Length on IgG4 Production," *The Journal of Immunology*, 161:3520-3526 (1998)

- 124 -

- T. Honjo et al., "Constant-Region Genes of the Immunoglobulin Heavy Chain and the Molecular Mechanism of Class Switching," *Immunoglobulin Genes*, (1989)
- 5 N. Lonberg et al., "Human Antibodies from Transgenic Mice," *Intern. Rev. Immunol.*, 13:65-93 (1995)
- G. Pluschke et al., "Generation of Chimeric Monoclonal Antibodies from Mice that Carry Human Immunoglobulin C γ 1 Heavy or C κ Light Chain Gene Segments," *Journal of Immunological Methods*, 215:27-37 (1998)
- 10 M. Brüggemann et al., "The Immunogenicity of Chimeric Antibodies," *J. Exp. Med.*, 170:2153-2157 (1989)
- F.A. Harding et al., "Class Switching in Human Immunoglobulin Transgenic Mice," *GenPharm International*,
- 15 N. Lonberg et al., "Antigen-specific Human Antibodies from Mice Comprising Four Distinct Genetic Modifications," *Nature*, 368:856-859 (1994)
- A. Cattaneo et al., "Polymeric Immunoglobulin M is Secreted by Transfectants of Non-lymphoid Cells in the Absence of Immunoglobulin J Chain
- 20 A.L. Defranco, "The Complexity of Signaling Pathways Activated by the BCR," *Current Opinion in Immunology*, 9:296-308 (1997)
- V. Arulampalam et al., "The Enhancer Shift: A Model to Explain the Developmental Control of IgH Gene Expression in B-lineage Cells," *Immunology Today*, 18:549-554 (1997)
- 25 S. Pettersson et al., "Temporal Control of IgH Gene Expression in Developing B Cells by the 3' Locus Control Region," *Immunobiol.*, 198:236-248 (1997)
- 30

- 125 -

- K. Kuze et al., "Characterization of the Enhancer Region for Germline Transcription of the Gamma 3 Constant Region Gene of Human Immunoglobulin," *International Immunology*, 3:647-655 (1991)
- 5 R. Mocikat et al., "The effect of the Rat Immunoglobulin Heavy-chain 3' Enhancer is position Dependent," *Gene*, 136:349-353 (1993)
- P. Dariavach et al., "The Mouse IgH 3' -Enhancer," *Eur. J. Immunol.*, 21:1499-1504 (1991)
- 10 J.S. Michaelson et al., "Identification of 3' α -hs4, a Novel Ig Heavy Chain Enhancer Element Regulated at Multiple Stages of B Cell Differentiation," *Nucleic Acids Research*, 23:975-981 (1995)
- 15 S. Pettersson et al., "A Second B Cell-specific Enhancer 3' of the Immunoglobulin Heavy-chain Locus," *Nature*, 344:165-168 (1990)
- V. Arulampalam et al., "Elevated Expression Levels of an Ig Transgene in Mice Links the IgH 3' Enhancer to the Regulation of IgH Expression," *International Immunology*, 8:1149-1157 (1996)
- 20 S. Delphin et al., "Characterization of an Interleukin 4 (IL-4) Responsive Region in the Immunoglobulin Heavy Chain Germline ϵ Promoter: Regulation by NF-IL-4, a C/EBP Family Member and Nf- κ B/p50," *J. Exp. Med.*, 181:181-192 (1995)
- 25 P. Matthias et al., "The Immunoglobulin Heavy Chain Locus Contains Another B-Cell-Specific 3' Enhancer Close to the α Constant Region," *Molecular and Cellular Biology*, 13:1547-1553 (1993)
- 30 R. Lieberman et al., "An Enhancer at the 3' End of the Mouse Immunoglobulin Heavy Chain Locus," *Nucleic Acids Research*, 19:933-937 (1991)

- 126 -

- P. Dariavach et al., "The Mouse IgH 3' -Enhancer," *Eur. J. Immunol.*, 21:1499-1504 (1991)
- 5 J.S. Michaelson et al., "Regulation of 3' IgH Enhancers by a Common Set of Factor, Including κ B-Binding Proteins," *The Journal of Immunology*, 156:2828-2839 (1996)
- 10 J. Chen et al., "Mutations of the Intronic IgH Enhancer and its Flanking Sequences Differentially Affect Accessibility of the J_H Locus," *The EMBC Journal*, 12:4635-4645 (1993)
- M. Cogné et al., "A Class Switch Control Region at the 3' End of the Immunoglobulin Heavy Chain Locus," *Cell*, 77: 737-747 (1994)
- 15 S. Huck et al., "Sequence of a Human Immunoglobulin Gamma 3 Heavy Chain Constant Region Gene: Comparison with the Other Human C γ Genes," *Nucleic Acids Research*, 13:1779-1789 (1986)
- 20 J.W. Ellison et al., "The Nucleotide Sequence of a Human Immunoglobulin C γ_1 Gene," *Nucleic Acids Research*, 10:4071-4079 (1982)
- J. Ellison et al., "Linkage and Sequence Homology of Two Human Immunoglobulin γ Heavy Chain Constant Region Genes," *Immunology*, 79:1984-1988 (1982)
- 25 J.B. Bolen, "Protein Tyrosine Kinases in the Initiation of Antigen Receptor Signaling," *Current Opinion in Immunology*, 7:306-311 (1995)
- L. O'Rourke et al., "Co-receptors of B Lymphocytes," *Current Opinion in Immunology*, 9:324-329 (1994)
- 30 T. Kurosaki, "Molecular Mechanisms in B Cell Antigen Receptor Signaling," *Current Opinion in Immunology*, 9:309-318 (1997)

- 127 -

- B.E. Pearson et al., "Expression of the Human β -amyloid Precursor Protein Gene from a Yeast Artificial Chromosome in Transgenic Mice," *Genetics*, 90:10578-10582 (1993)
- 5 J.F. Loring et al., "Rational Design of an Animal Model for Alzheimer's Disease: Introduction of Multiple Human Genomic Transgenes to Reproduce AD Pathology in a Rodent," *Neurobiology of Aging*, 17:173-182 (1996)
- 10 J.J. MacQuitty, "The Real Implications of Dolly," *Nature Biotechnology*, 15:294 (1997)
- M.T.F. Huang, "Gene Targeting Technology for Creating Transgenic Models of Lymphopoiesis," *Laboratory Animal Science*, 43:156-159 (1993)
- 15 J.J. MacQuitty, "GenPharm's Knockout Mice," *Science*, 257:1188 (1992)
- M.T.F. Huang, "T Cell Development in CD3- ζ Mutant Mice," *Intern. Rev. Immunol.*, 13:29-41 (1995)
- 20 L.D. Taylor et al., "A Transgenic Mouse That Expresses a Diversity of Human Sequence Heavy and Light Chain Immunoglobulins," *Nucleic Acids Research*, 20:6287-6295 (1992)
- D.M. Fishwild et al., "High-avidity Human IgGk Monoclonal Antibodies from a Novel Strain of Minilocus Transgenic Mice," *Nature Biotechnology*, 14:845-851 (1996)
- 25 L.D. Taylor et al., "Human Immunoglobulin Transgenes Undergo Rearrangement, Somatic Mutation and Class Switching in Mice that Lack Endogenous IgM,"
- 30 *International Immunology*, 6:579-591 (1994)

- 128 -

- N. Tuailon et al., "Human Immunoglobulin Heavy-chain Minilocus Recombination in Transgenic Mice: Gene-segment Use in μ and γ Transcripts," *Immunology*, 90:3720-3724 (1993)
- 5 N. Lonberg et al., "Antigen-specific Human Antibodies from Mice Comprising Four Distinct Genetic Modifications," *Nature*, 368:856-859 (1994)
- 10 M.J. Shlomchic et al., "The Role of B Cells in *lpr/lpr*-induced Autoimmunity," *J. Exp. Med.*, 180:1295-1306 (1994)
- L. Pricop et al., "Antibody Response Elicited by T-dependent and T-independent Antigens in Gene Targeted κ -deficient Mice," *International Immunology*, 6:1839-1847 (1994)
- 15 Y. Liu et al., "Gene-targeted B-deficient Mice Reveal a Critical Role for B Cells in the CD4 T Cell Response," *International Immunology*, 7:1353-1362 (1995)
- 20 T.K. Choi et al., "Transgenic Mice Containing a Human Heavy Chain Immunoglobulin Gene Fragment Cloned in a Yeast Artificial Chromosome," *Nature Genetics*, 4:117-123 (1993)
- 25 S.D. Wager et al., "Antibody Expression from the Core Region of the Human IgH Locus Reconstructed in Transgenic Mice Using Bacteriophage P1 Clones," *Genomics*, 35:405-414 (1996)
- 30 S.D. Wagner et al., "The Diversity of Antigen-specific Monoclonal Antibodies from Transgenic Mice Bearing Human Immunoglobulin Gene Miniloci," *Eur. J. Immunol.*, 24:2672-2681 (1994)
- S.D. Wagner et al., "Antibodies Generated from human Immunoglobulin Miniloci in Transgenic Mice," *Nucleic Acids Research*, 22:1389-1393 (1994)

- 129 -

- 5 N.P. Davies et al., "Creation of Mice Expressing Human Antibody Light Chains by Introduction of a Yeast Artificial Chromosome Containing the Core Region of the Human Immunoglobulin κ Locus," *Bio/Technology*, 11:91-914 (1993)
- F.D. Batista et al., "Affinity Dependence of the B Cell Response to Antigen: A Threshold, a Ceiling, and the Importance of Off-Rate," *Immunity*, 8:751-759 (1998)
- 10 M. R. Ehrenstein et al., "Targeted Gene Disruption Reveals a Role for Natural Secretory IgM in the Maturation of the Primary Immune Response," *Immunology*, 95:10089-10093 (1998)
- 15 C. Milstein et al., "Both DNA Strands of Antibody Genes are Hypermutation Targets," *Immunology*, 95:8791-8794 (1998)
- J.E. Sale et al., "TdT-Accessible Breaks Are Scattered over the Immunoglobulin V Domain in a Constitutively Hypermutating B Cell Line," *Immunity*, 9:859-869 (1998)
- 20 M.S. Neuberger, "Antigen Receptor Signaling Gives Lymphocytes a Long Life," *Cell*, 90:971:973 (1997)
- B. Goyenechea et al., "Cells Strongly Expressing Ig κ Transgenes Show Clonal Recruitment of Hypermutation: A Role for Both MAR and the Enhancers," *The EMBO Journal*, 16:3987-3994 (1997)
- 25 Y.M. The et al., "The Immunoglobulin (Ig) α and Ig β Cytoplasmic Domains Are Independently Sufficient to Signal B Cell Maturation and Activation in Transgenic Mice," *J. Exp. Med.*, 185:1753-1758 (1997)
- 30 J. Yélamos et al., "Targeting of Non-Ig Sequences in Place of the V Segment by Somatic Hypermutation," *Nature*, 376:225-229 (1995)

- 130 -

- S.D. Wagner et al., "Codon bias Targets Mutation," *Nature*, 376:732 (1995)
- 5 N. Klix et al., "Multiple Sequences from Downstream of the J α Cluster Can Combine to Recruit Somatic Hypermutation to a Heterologous, Upstream Mutation Domain," *Eur. J. Immunol.*, 28:317-326 (1996)
- M. Neuberger et al., "Mice Perform a Human Repertoire," *Nature*, 386:25-26 (1997)
- 10 N.P. Davies et al., "Targeted Alterations in Yeast Artificial Chromosomes for Inter-species Gene Transfer," *Nucleic Acids Research*, 20:2693-2698 (1992)
- M. Brüggemann et al., "Strategies for Expressing Human Antibody Repertoires in Transgenic Mice," *Immunology Today*, 17:391-397 (1996)
- 15 X. Zou et al., "Dominant Expression of a 1.3 Mb Human Igk Locus Replacing Mouse Light Chain Production," *The FASEB Journal*, 10:1227-1232 (1996)
- I.K. Jarmer et al., "Chimaeric Monoclonal Antibodies Encoded by the Human V H 26 Gene From Naïve Transgenic Mice Display a Wide Range of Antigen-binding Specificities," *Immunology*, 88:174-182 (1996)
- 20 X. Zou et al., "Subtle Differences in Antibody Responses and Hypermutation of λ Light Chains in Mice with a Disrupted α Constant Region," *Eur. J. Immunol.*, 25:2154-2162 (1995)
- 25 A.V. Popov et al., "Yeast Colony Size Reflects YAC Copy Number," *Nucleic Acids Research*, 25:2039-2040 (1997)
- 30 N.P. Davies et al., "Extension of Yeast Artificial Chromosomes by Cosmid Multimers," *Nucleic Acids Research*, 21:767-768 (1993)

- 131 -

- C. Bützler et al., "Rapid Induction of B-cell Lymphomas in Mice Carrying a Human IgH/c-MYC^{YAC}," *Oncogene*, 14:1383-1388 (1997)
- 5 A.V. Popov et al., "Assembly and Extension of Yeast Artificial Chromosomes to Build Up a Large Locus," *Gene*, 177:195-201 (1996)
- H. Waldmann et al., "Monoclonal Antibodies for Immunosuppression," *Monoclonal Antibody Therapy Prog Allergy*, 45:16-30 (1988)
- 10 M. Brüggemann et al., "Designer Mice: The Production of Human Antibody Repertoires in Transgenic Animals," *generation of Antibodies by Cell and Gene Immobilization*, 7:33-40 (1993)
- 15 A.G. Betz et al., "Discriminating Intrinsic and Antigen-selected Mutational hotspots in Immunoglobulin V Genes," *Immunology Today*, 14:405-409
- M. Brüggemann et al., "Construction, Function and Immunogenicity of Recombinant Monoclonal Antibodies," *Behring, Inst. Mitt.*, 87:21-14 (1990)
- 20 M. Brüggemann et al., "Production of Human Antibody Repertoires in Transgenic Mice," *Current Opinion in Biotechnology*, 8:455-458 (1997)
- K.B. Meyer et al., "The Immunoglobulin κ Locus Contains a Second, Stronger B-cell-specific Enhancer Which is Located Downstream of the Constant Region," *The EMBO Journal*, 8:1959-1964 (1989)
- 25 M. Brüggemann et al., "Sequence of a Rat Immunoglobulin γ_{2c} Heavy Chain Constant Region cDNA: Extensive Homology to Mouse γ_3 ," *Eur. J. Immunol.*, 18:317-319 (1988)
- 30 M. Brüggemann et al., "Human Antibody Production in Transgenic Mice: Expression from 100 kb of the Human IgH Locus," *Eur. J. Immunol.*, 21:1323-1326 (1991)

- 132 -

- M. Brüggemann, "Evolution of the Rate Immunoglobulin Gamma Heavy-chain Gene Family," *Gene*, 74:473-482 (1988)
- 5 R. Sitia et al., "Regulation of Membrane IgM Expression in Secretory B Cells: Translational and Post-transnational Events," *The EMBO Journal*, 6:3969-3977 (1987)
- 10 M.J. Sharpe et al., "Somatic Hypermutation of Immunoglobulin κ may depend on Sequences 3' of C κ and Occurs on Passenger Transgenes," *The EMBO Journal*, 10:2139-2145 (1991)
- S. Biocca et al., "Expression and Targeting of Intracellular Antibodies in Mammalian Cells," *The EMBO Journal*, 9:101-108 (1990)
- 15 K.J. Patel et al., "Antigen Presentation by the B Cell Antigen Receptor Is Driven by the α/β Sheath and Occurs Independently of Its Cytoplasmic Tyrosines," *Cell*, 74:939-946 (1993)
- 20 K.B. Meyer et al., "The Importance of the 3' -Enhancer Region in Immunoglobulin κ Gene Expression," *Nucleic Acids Research*, 18:5609-56115 (1990)
- A.G. Betz et al., "Elements Regulating Somatic Hypermutation of an Immunoglobulin κ Gene: Critical Role for the Intron Enhancer/Matrix Attachment Region," *Cell*, 77:239-248 (1994)
- 25 J.O. Mason et al., "Transcription Cell Type Specificity Is Conferred by an Immunoglobulin V_H Gene Promoter That Includes a Functional Consensus Sequence," *Cell*, 41:479-487 (1985)
- 30 M.S. Neuberger et al., "Activation of Mouse Complement by Monoclonal Mouse Antibodies," *Eur. J. Immunol.*, 11:1012-1016 (1981)

- 133 -

- 5 M.R. Walker et al., "Interaction of Human IgG Chimeric Antibodies with the Human FcRI and FcRII Receptors: Requirements for Antibody-Mediated Host Cell-Target Cell Interaction," *Molecular Immunology*, 26:403-411 (1989)
- M. Brüggemann et al., "Comparison of the Effector Functions of Human Immunoglobulins Using a Matched Set of Chimeric Antibodies," *J.Exp. Med.* 166:1351-1361 (1987)
- 10 R. Sherman-Gold, "Monoclonal Antibodies: The Evolution from '80s Magic Bullets to Mature, Mainstream Applications as Clinical Therapeutics," *Genetic Engineering News*, 17 (1997)

We claim:

1. A transgene comprising a DNA fragment comprising a DNA sequence of a contiguous region of the immunoglobulin heavy chain locus on human chromosome 14 from the D segment genes, continuing through the J segment genes and the constant region gene C μ of that locus, wherein said DNA fragment is operably linked to at least one human V segment gene, and wherein said DNA fragment further is operably linked to an additional constant region gene, said additional constant region gene comprising a mouse gamma-1 switch region and human constant region coding exons, wherein said transgene is a yeast artificial chromosome (YAC).

2. The transgene according to claim 1, wherein said human constant region coding exons encode a human constant region selected from the group consisting of a human gamma constant region, a human alpha constant region and a human epsilon constant region.

3. The transgene according to claim 2, wherein said human constant region coding exons encode a human gamma constant region.

4. The transgene according to claim 3, wherein said human gamma constant region is a gamma-1 constant region.

5. The transgene according to claim 4 which is the yH2Bm yeast artificial chromosome (YAC) of Figure 1.

6. The transgene according to claim 3, wherein said human gamma constant region is a gamma-2 constant region.

7. The transgene according to claim 3, wherein said human gamma constant region is a gamma-3 constant region.

8. The transgene according to claim 3, wherein said human
5 gamma constant region is a gamma-4 constant region.

9. The transgene according to claim 8 which is the yH2Cm yeast artificial chromosome (YAC) of Figure 1.

10. The transgene according to claim 2, wherein said human
10 constant region coding exons encode a human alpha constant region.

11. The transgene according to claim 10, wherein said
15 human alpha constant region is an alpha-1 constant region.

12. The transgene according to claim 10, wherein said human alpha constant region is an alpha-2 constant region.

13. The transgene according to claim 2, wherein said human
20 constant region coding exons encode a human epsilon constant region.

14. The transgene according to claim 1, wherein said DNA
25 fragment is operably linked to a plurality of human VH genes.

15. The transgene according to claim 14, wherein said DNA fragment is operably linked to at least 50% of the human germline VH genes.

30

16. The transgene according to claim 14, wherein said DNA fragment is operably linked to at least 40 different human VH genes.

17. The transgene according to claim 14, wherein said DNA
fragment is operably linked to a sufficient number of different
5 human VH genes so that the transgene is capable of encoding at
least 1×10^5 different functional human immunoglobulin heavy
chain sequences through V(D)J recombination, without taking into
account junctional diversity or somatic mutation events.

10 18. The transgene according to claim 14, wherein the
number of human VH genes is sufficient to produce at least 50%
of the B-cell population of a wild-type mouse in a transgenic
mouse containing the transgene.

15 19. A transgene which is the
yHG4 artificial chromosome (YAC) of Figure 2.

20 20. A transgene comprising a DNA fragment comprising a DNA
sequence of a contiguous region of the immunoglobulin heavy
chain locus on human chromosome 14, from the D segment
genes, continuing through the J segment genes and the
25 constant region gene C μ of that locus, wherein said DNA
fragment is operably linked to at least one human V segment
gene, and wherein said DNA fragment further is operably
linked to an additional constant region gene, said
additional constant region gene comprising a human switch
30 region, human CH1, C hinge , CH2 and CH3 exons and human
membrane exons, wherein said human switch region and said
human membrane exons are from the same isotype and the
human CH1, C hinge , CH2 and CH3 exons are from a different

isotype than said human switch region and said human membrane exons.

21. The transgene according to claim 20, wherein said
5 human switch region and said human membrane exons are human gamma-2 sequences.

22. The transgene according to claim 21, wherein said
10 human CH1, C_{hinge}, CH2 and CH3 exons encode a human constant region selected from the group consisting of: a human gamma-1 constant region, a human gamma-3 constant region, a human gamma-4 constant region, a human alpha-1 constant region, a human alpha-2 constant region and a human epsilon constant region.

15 23. The transgene according to claim 22, wherein the human CH1, C_{hinge}, CH2 and CH3 exons encode a human gamma-1 constant region.

24. The transgene according to claim 23 which is the
20 yHG1/2 yeast artificial chromosome (YAC) of Figure 3.

25 25. The transgene according to claim 20, wherein the human CH1, C_{hinge}, CH2 and CH3 exons encode a human gamma-4 constant region.

26. The transgene according to claim 20, wherein said DNA
fragment is operably linked to a plurality of human VH genes.

27. The transgene according to claim 20, wherein said DNA
30 fragment is operably linked to at least 50% of the human germline VH genes.

28. The transgene according to claim 20, wherein said DNA fragment is operably linked to at least 40 different human VH genes.

5 29. The transgene according to claim 20, wherein said DNA fragment is operably linked to a sufficient number of different human VH genes so that the transgene is capable of encoding at least 1×10^5 different functional human immunoglobulin heavy chain sequences through V(D)J recombination, without taking into
10 account junctional diversity or somatic mutation events.

30. The transgene according to claim 20, wherein the number of human VH genes is sufficient to produce at least 50% of the B-cell population of a wild-type mouse in a transgenic
15 mouse containing the transgene.

31. The transgene according to claim 1 or claim 20, further comprising a mouse 3' enhancer.

20 32. The transgene according to claim 31, wherein said mouse 3' enhancer is an approximately 0.9 kb core fragment of a mouse germline 3' enhancer.

33. The transgene according to claim 31, wherein said
25 mouse 3' enhancer is an approximately 4 kb fragment of a mouse germline 3' enhancer.

34. The transgene according to claim 31, wherein said
mouse 3' enhancer is a locus control region.

30

35. A non-human embryonic stem (ES) cell comprising a transgene according to any one of claims 1-34.

36. The embryonic stem (ES) cell according to claim 35, which is a mouse ES cell.

37. A transgenic non-human animal and its progeny,
5 wherein the somatic and germ cells comprise the transgene according to any one of claims 1-34.

38. The transgenic non-human animal and progeny according to claim 37, further comprising a human immunoglobulin light
10 chain transgene.

39. The transgenic non-human animal and progeny according to claim 37 or claim 38, wherein the animal is a mouse.

15 40. The transgenic non-human animal and progeny according to claim 37 or claim 38, further comprising inactivated endogenous immunoglobulin heavy and light chain loci.

41. The transgenic non-human animal according to claim 40, wherein the animal is a mouse.

20 42. A method for producing a transgenic non-human animal and its progeny whose somatic and germ cells comprise a transgene according to any one of claims 1-34 and that produces high affinity, fully human antibodies of a desired isotype
25 specific for an antigen of interest following immunization with the antigen of interest, comprising:

(a) introducing said transgene into an embryonic stem cell;
(b) producing from the embryonic stem cell a transgenic non-human animal comprising a germ cell that contains said
30 transgene; and

(c) breeding said transgenic non-human animal as needed to produce a transgenic non-human animal and its progeny that produces high affinity, fully human antibodies of a desired

isotype specific for an antigen of interest following immunization with the antigen of interest.

43. The method according to claim 42, wherein the
5 transgenic non-human animal is a mouse.

44. A method for producing a high-affinity, fully human antibody of a desired isotype, wherein said antibody is specific for an antigen of interest, comprising the steps of contacting a
10 transgenic, non-human animal according to any one of claims 37-41 with the antigen of interest to induce antibody production in a B-cell of said animal, and collecting said antibody.

45. An antibody producing B-cell harvested from a
15 transgenic non-human animal according to any one of claims 37-41 immunized with an antigen of interest.

46. The B-cell according to claim 45 which has been immortalized.
20

47. The method according to claim 44, wherein the antibody is collected from the blood stream of said transgenic non-human animal.

25 48. The method according to claim 44, wherein the antibody is collected from an immortalized B-cell according to claim 46.

49. The method according to claim 44, wherein said antibody is collected from a host cell transfected with DNA
30 isolated from a B-cell according to claim 45.

50. A transgene comprising a DNA fragment comprising a DNA sequence identical to the DNA sequence of human chromosome 14 from the D segment genes of the human immunoglobulin heavy chain

locus, continuing through the J segment genes and the constant region genes through C_μ of that locus, wherein said DNA fragment is operably linked to at least one human V segment gene, and wherein said DNA fragment further is operably linked to an additional constant region, said additional constant region comprising: (1) a human switch region; (2) a region comprising human CH1, C_{hinge}, CH2 and CH3 exons; and (3) human membrane exons, wherein said human switch region, said region comprising human CH1, C_{hinge}, CH2 and CH3 exons, and said human membrane exons are from different isotypes.

51. A transgene comprising a DNA fragment comprising a DNA sequence identical to the DNA sequence of human chromosome 14 from the D segment genes of the human immunoglobulin heavy chain locus, continuing through the J segment genes and the constant region genes through C_μ of that locus, wherein said DNA fragment is operably linked to at least one human V segment gene, and wherein said DNA fragment further is operably linked to an additional constant region, said additional constant region comprising a human switch region, human CH1, C_{hinge}, CH2 and CH3 exons, and non-human membrane exons, wherein said human switch region, said region comprising human CH1, C_{hinge}, CH2 and CH3 exons, and said non-human membrane exons are from the same or from different isotypes.

52. A transgene comprising a DNA fragment comprising a DNA sequence identical to the DNA sequence of human chromosome 14 from the D segment genes of the human immunoglobulin heavy chain locus, continuing through the J segment genes and the constant region genes through C_μ of that locus, wherein said DNA fragment is operably linked to at least one human V segment gene, and wherein said DNA fragment further is operably linked to an additional constant region, said additional constant region comprising a non-human switch region, human CH1, C_{hinge}, CH2 and

CH3 exons, and non-human membrane exons, wherein said non-human switch region and said non-human membrane exons are from the same species.

- 5 53. The transgene according to any one of claims 1-34 or 50-52, wherein a loxP site is inserted 3' of the switch region and 5' of the CH1 exon.