

IP 67


1

DESCRIPTION

NOVEL PEPTIDE DERIVATIVES

5 Technical Field

This invention relates to a novel peptide derivative having an antitumor activity, and, more detailedly, relates to a peptide derivative represented by the following formula or a salt thereof



wherein A and B each represent either of the following (a) and (b),

(a) A represents a hydrogen atom, and B represents a phenyl group substituted with a halogen atom, hydroxyl group, lower alkyl group or lower alkoxy group, or a heteroaryl group,

(b) A represents $-\text{CONH}-\text{R}^1$, $-\text{CSNH}-\text{R}^1$, a hydroxymethyl group, a lower alkoxycarbonyl group or a carboxyl group, wherein, R^1 represents a lower alkyl group or a heteroaryl group, and B represents a phenyl group optionally substituted with a halogen atom, hydroxyl group, lower alkyl group or lower alkoxy group.

Background Art

Peptides having a cytostatic activity and/or an antineoplasm activity have been isolated from marine molluscs, sea hare Dolabella auricularia and these peptides are called dolastatins 1 to 15. Among them,


2-21-5030

dolastatin 10 is a pentapeptide extracted from Dolabella auricularia from the Indian Ocean in 1987 by G. R.

Pettit, et al. and having the following structural formula, and is said to be the strongest cytostatic

5 substance presently known (see, G. R. Pettit, et al., J. Am. Chem. Soc., 109, 6883 (1987) and U.S. Patent No. 4,816,444).

10



[Dolastatin 10]

15

Further, recently, publication was made on the total synthesis of dolastatin 10 itself (see, U.S. Patent No. 4,978,744).

In this connection, the present inventors
20 previously disclosed certain dolastatin 10 derivatives (see, WO93/03054 Pamphlet).

The present inventors found that certain dolastatin 10 derivatives wherein the dolaphenine (which means an α -(thiazolyl)phenethylamino group) at the
25 C-terminus of dolastatin 10 is substituted with another substituent have a much stronger antitumor activity than that of dolastatin 10.

Disclosure of Invention

30 In the present description, the term "lower" means that the number of the carbon atoms of a group or compound to which this term is attached is 6 or less, preferably 4 or less.

In the above formula (I), as the "lower alkyl group", there can, for example, be mentioned methyl,
35 ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-

butyl, tert-butyl, n-hexyl groups, etc. and as the "lower alkoxy group", there can, for example, be mentioned methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy groups, etc. Further, the "halogen atom" includes
5 fluorine, chlorine, bromine and iodine atoms.

The "heteroaryl group" means an aromatic heterocyclic group containing hetero atom(s) selected from O, S and N, preferably, 5 or 6-membered heterocyclic group containing 1 to 4 hetero atoms, such as
10 thienyl, furyl, pyrrolyl, imidazolyl, thiazolyl, oxazolyl, thiadiazolyl, tetrazolyl, pyridyl, pyrimidinyl, triazinyl groups, etc.

The "phenyl group substituted with a halogen atom, hydroxyl group, lower alkyl group or lower alkoxy group" represented by the symbol B includes a phenyl
15 group substituted with one halogen atom, hydroxyl group, lower alkyl group or lower alkoxy group, and there can, for example, be mentioned 2-fluorophenyl, 2-chlorophenyl, 2-bromophenyl, 3-fluorophenyl, 3-iodophenyl,
20 4-chlorophenyl, 4-bromophenyl, 2-hydroxyphenyl, 2-methylphenyl, 4-ethylphenyl, 2-methoxyphenyl, 4-ethoxyphenyl groups, etc. Further, the "phenyl group optionally substituted with a halogen atom, hydroxyl group, lower alkyl group or lower alkoxy group" includes
25 an unsubstituted phenyl group besides the above substituted phenyl groups.

A group of preferred compounds in the invention are compounds of the above formula (I) wherein A represents a hydrogen atom and B represents a phenyl
30 group substituted with a halogen atom, hydroxyl group, lower alkyl group or lower alkoxy group, or a heteroaryl group, particularly compounds of the above formula (I) wherein B represents a phenyl group substituted with a halogen atom, hydroxyl group, lower alkyl group or lower
35 alkoxy group; a thienyl group; or a pyridyl group.

Another group of preferred compounds are

compounds of the above formula (I) wherein A represents
-CONH-R¹, -CSNH-R¹, a hydroxymethyl group, a lower
alkoxycarbonyl group or a carboxyl group, wherein R¹
represents a lower alkyl group or a heteroaryl group,
5 and B represents a phenyl group optionally substituted
with a halogen atom, hydroxyl group, lower alkyl group
or lower alkoxy group, particularly compounds of the
above formula (I) wherein A represents -CONH-R¹,
-CSNH-R¹, a hydroxymethyl group, a lower alkoxycarbonyl
10 group or a carboxyl group, wherein R¹ represents a lower
alkyl group, a thiazolyl group or thiadiazolyl group,
and B represents an unsubstituted phenyl group.


In the compounds of the above formula (I) of
the invention, the carbon atoms to which an isopropyl
15 group, a sec-butyl group, a methoxy group and a methyl
group bind respectively are asymmetric carbon atoms, and
therefore, they can arbitrary have an R- or S-configu-
ration. All those compounds are included in the scope
of the invention, but in view of pharmacological activi-
20 ty, compounds having the same configuration as
dolastatin 10 are preferred.

The peptide compounds of the above formula (I)
can, further, exist as salts, and as examples of such
salts, there can be mentioned hydrochlorides, hydro-
25 bromides, trifluoroacetates, p-toluenesulfonates, ace-
tates, etc.


According to the invention, a peptide compound
of the above formula (I) can be prepared by condensing
the respective amino acids or peptide fragments, for
30 example, according to a liquid phase synthesis method
(see, E. Schröder and K. Lübke, "The Peptides", volume
1, pages 76-136, 1965, published by Academic Press)
known in the field of peptide chemistry.

For example, for avoiding racemization at the
35 condensation reaction, it is preferred to conduct syn-
thesis by condensing a tripeptide fragment of the fol-

following formula (II)




with a fragment of the following formula (III)




wherein A and B are as defined above.

15 Further, for synthesizing many compounds of the invention efficiently, it is preferred to conduct the synthesis by condensing a tetrapeptide fragment of the following formula (IV)



25 with a fragment of the following formula (V)



wherein A and B are as defined above.

The condensation reaction can be conducted, generally, by treating the fragments with a condensing agent, e.g. dicyclohexylcarbodiimide (DCC), diphenyl phosphoryl azide (DPPA) or diethyl phosphorocyanidate (DEPC), a so-called BOP reagent, or the like in an inert

solvent such as, for example, chloroform, ethyl acetate, tetrahydrofuran (THF), dimethylformamide (DMF) or acetonitrile, if necessary in the presence of an organic base such as, for example, triethylamine, N-methylmorpholine
5 or diisopropylethylamine (DIEA).

The reaction temperature is usually -10°C to room temperature, preferably around 0°C . The ratios of the compound of the formula (III), the organic base and the condensing agent to the compound of the formula (II)
10 are not strictly limited, but, usually, it is advantageous to use the compound of the formula (III) of at least one mole, preferably of the order of 1.0 to 1.1 moles, the organic base of the order of 2 moles, and the condensing agent of the equimolar order, respectively
15 per mole of the compound of the formula (II).

A compound of the formula (I) wherein A represents a carboxyl group can also be prepared by alkali hydrolysis of the compound of the formula (I) wherein A represents a lower alkoxycarbonyl group.

20 The isolation and purification of thus obtained peptide compound of the formula (I) from the reaction mixture can be conducted by methods known per se, for example by recrystallization, ion exchange chromatography, gel filtration, high performance liquid
25 chromatography, etc.

The compounds of the above formula (III) and (IV) used as starting materials in the above reaction are novel compounds not disclosed in prior literatures, and can easily be prepared by condensing amino acids,
30 which are constituents thereof, according to a liquid phase synthesis method.

The peptide compounds of the formula (I) of the invention have a higher antitumor activity than dolastatin 10, and have a large therapeutic index, and
35 are useful for treatment of acute myelocytic leukemia, acute lymphocytic leukemia, chronic melanoma, pulmonary

adenocarcinoma, neuroblastoma, pulmonary small cell carcinoma, breast cancer, colon cancer, ovary cancer, bladder cancer, etc.

The antitumor activity of the compounds can be
5 assayed as follows.

(1) Assay of antitumor activity

0.1 ml (10^6 cells/mouse) portions of mouse leukemia P388 cells were implanted intraperitoneally into 7-week-old CDF1 mice. A compound was intraperito-
10 neally administered thereinto on the first day (the day after implantation) and the fifth day after implanta-
tion, and the life or death of the mice was observed for 60 days. From the results were calculated increases in life span (ILS, %) according to the following equation.
15 In the following equation, T means median survival days of the chemical administration group, and C means median survival days of the control group.

$$\text{ILS} = \frac{T - C}{C} \times 100$$

20

The results are shown in the following Table. Antitumor activity is shown as the relative ratio in the case where the ILS of dolastatin 10 is supposed to be 100.

25

Example No. of compound	Antitumor activity
8	190
12	190
14	190
dolastatin 10	100

30

The compound of the invention, when used as a drug, can be used by formulating them into any dosage form of solid forms (e.g., tablets, hard capsules, soft
35 capsules, granules, powders, fine granules, pills, troches, etc.), semi-solid forms (e.g., suppositories,

ointments, etc.) and liquid forms (e.g., injections, emulsions, suspensions, lotions, sprays, etc.). As nontoxic additives usable in the above formulations, there can, for example, be mentioned starches, gelatin, 5 glucose, lactose, fructose, maltose, magnesium carbonate, talc, magnesium stearate, methylcellulose, carboxymethylcellulose or salts thereof, gum arabic, polyethylene glycol, p-hydroxybenzoic acid alkyl esters, syrups, ethanol, propylene glycol, vaseline, carbowax, 10 glycerol, sodium chloride, sodium sulfite, sodium phosphate, citric acid, etc. The drug can also contain another therapeutically effective drug.

The content of the compound of the invention in the drug varies depending on the dosage form, but it 15 is generally preferred that the drug contains the compound at a concentration of 0.1 to 50 wt % in the case of solid and semi-solid forms, and at a concentration of 0.05 to 10 wt % in the case of liquid form.


The dose of the compound of the invention can 20 widely be varied depending on the kind of warm-blooded animals including human beings as a subject, administration routes, the seriousness of symptoms, the diagnoses of doctors, etc., but can generally be on the order of 0.01 to 50 mg/kg per day. However, it is of course 25 possible to administer the compound in an amount smaller than the lower limit of the above range or in an amount larger than the upper limit thereof in accordance with the seriousness of symptom of the patient and the diagnosis of the doctor as mentioned above. The above dose 30 can be administered once a day or in divided several portions per day.


Examples

The invention is further described below 35 according to Referential Examples and Examples.

As for the structure of compounds correspond-

ing to compound numbers used in Referential Examples and Examples, please refer to the following Flow Sheets 1 and 2. Therein, Bu^t represents a tert-butyl group, Boc a tert-butoxycarbonyl group, Bzl a benzyl group, Me a methyl group, B a phenyl group substituted with a halogen atom, hydroxyl group, lower alkyl group or lower alkoxy group, or a heteroaryl group, and A $-\text{CONH}-\text{R}^1$, $-\text{CSNH}-\text{R}^1$; a hydroxymethyl group, a lower alkoxycarbonyl group or a carboxyl group, wherein R^1 represents a lower alkyl group or a heteroaryl group.

Flow Sheet 1

Flow Sheet 2

Referential Example 1

Preparation of compound 2

30 ml of a tetrahydrofuran - n-hexane (1 : 1) solution of 23.8 % lithium diisopropylamide (LDA, 66.4 mmoles) is gradually poured into 40 ml of anhydrous tetrahydrofuran under stirring at -20° in an atmosphere of nitrogen, the mixture is cooled to -78°, and 9.84 g (60 mmoles) of benzyl propionate is added dropwise over a period of 30 minutes. 5 minutes later, a solution of 7.96 g (40 mmoles) of Boc-prolinal in 40 ml of tetrahydrofuran is added dropwise at the same temperature over a period of 1 hour. The mixture is stirred at the same temperature for 15 minutes, 150 ml of ice-cooled 1N-hydrochloric acid is added, and the mixture is warmed to room temperature. The mixture is extracted with ethyl acetate, the ethyl acetate layer is washed with water and dried, the solvent is distilled off under reduced pressure, and the remaining oily matter is purified by silica gel flash chromatography using ethyl acetate - n-hexane (1 : 5) as an eluent to obtain the desired compound 2 as colorless oily matter. 3.86 g (26.6 %).

$$[\alpha]_D^{27} -28.4^\circ \text{ (c = 0.82, MeOH)}$$

¹H-NMR (CDCl₃, δ) 1.30 (3H, d, J=7.0Hz), 1.45 (9H, s), 1.6-2.1 (m), 2.61 (1H, quintet, J=7.0Hz), 3.0-3.6 (m), 3.7-4.1 (m), 5.13 (2H, s), 7.34 (5H, s)

Referential Example 2

Preparation of compound 3

730 mg (2.01 mmoles) of compound 2 obtained in Referential Example 1 is dissolved in 10 ml of dimethylformamide, 0.7 ml (11.22 mmoles) of methyl iodide is added therein under stirring, and 0.16 g (4.00 mmoles) of sodium hydride (60 % in mineral oil) is added therein. Stirring is continued at 0° for 1 hour, ice water is added, and the mixture is extracted with ethyl acetate - benzene (4 : 1). The organic layer is washed

with 5 % potassium hydrogen sulfate, saturated aqueous sodium bicarbonate, 5 % sodium thiosulfate and saturated saline in this order, and dried. The resultant crude product is purified by silica gel flash chromatography using ethyl acetate - n-hexane (1 : 10) as an eluent to obtain the desired compound 3 as colorless oily matter. 530 mg (72.5 %).

$[\alpha]_D^{27} -25.7^\circ$ (c = 0.389, MeOH)

$^1\text{H-NMR}$ (CDCl_3 , δ) 1.26 (3H, d, $J=6.8\text{Hz}$), 1.45 (9H, s), 1.65-2.1 (m), 2.56 (1H, quintet, $J=7.0\text{Hz}$), 3.0-4.0 (m), 3.38 (3H, s), 5.14 (2H, s), 7.34 (5H, s)

Referential Example 3

Preparation of compound 4

(a) 1 ml of concentrated hydrochloric acid is added to 97.1 mg (0.2 mmole) of compound 1 (known compound) under ice cooling, and the mixture is stirred at 0° for 1 hour and evaporated to dryness under reduced pressure. The residue is dissolved in 2 ml of dimethylformamide, 0.15 ml of triethylamine is added dropwise at 0° , and the mixture is again evaporated to dryness under reduced pressure.

(b) On the other hand, 76 mg (0.2 mmole) of compound 3 obtained in Referential Example 2 is dissolved in 0.5 ml of ethyl acetate, 2.0 ml of 2N-hydrogen chloride/ethyl acetate is added under ice cooling, and the mixture is brought to room temperature, stirred for 1.5 hours, evaporated to dryness under reduced pressure and then dried.

The products obtained in (a) and (b) are combined and dissolved in 0.8 ml of dimethylformamide, 34.3 mg (1.1 equivalents) of DEPC is added, the mixture is ice-cooled, 56 μl (2 equivalents) of triethylamine is added, and stirring is continued under ice cooling for 1 hour and then at room temperature overnight. The solvent is evaporated under reduced pressure, the residue is dissolved in dichloromethane, and the solution is

washed with saturated aqueous sodium bicarbonate and saturated saline and dried. The resultant crude product is purified by silica gel flash chromatography using dichloromethane - methanol (20 : 1) as an eluent, and then by Sephadex LH-20 chromatography using n-hexane - dichloromethane - methanol (2 : 7.5 : 2.5) as an eluent to obtain the desired compound 4 as an amorphous solid. 117 mg (85.0 %).

$[\alpha]_D^{26} -44.0^\circ$ (c = 0.80, MeOH)

$^1\text{H-NMR}$ (CDCl_3 , δ) 0.7-1.5 (m), 1.27 (3H, d, J=7.0Hz), 1.5-2.25 (m), 2.25-2.9 (m), 3.01 (3H, s), 3.29 (3H, s), 3.35 (3H, s), 3.8-4.3 (m), 4.5-5.0 (m), 5.13 (2H, s), 7.34 (5H, s)

Referential Example 4-A

Preparation of compound 6-A

(in compound 6, A = CONH-Et, * = S)

1.33 g (5 mmoles) of Boc-phenylalanine is dissolved in 20 ml of tetrahydrofuran, and while the solution is stirred at -15° , 0.56 ml (5 mmoles) of N-methylmorpholine and then 0.67 ml (5 mmoles) of isobutyl chloroformate are added. After stirring the mixture at -15° for 5 minutes, 0.64 g (2 equivalents) of aqueous 70 % ethylamine solution is added, and stirring is continued at -15° for 15 minutes and then at room temperature for 1.5 hours. The reaction solution is concentrated under reduced pressure, the residue is dissolved in ethyl acetate, the solution is washed with ice-cooled 2N-hydrochloric acid and saturated aqueous sodium bicarbonate and dried, the solvent is distilled off, and the residue is crystallized from ethyl acetate-ether-n-hexane to obtain the desired compound 6-A as needle crystals. 1.12 g (76.7 %).

Melting point $123-4^\circ$.

$^1\text{H-NMR}$ (CDCl_3 , δ) 0.99 (3H, t, J=7.3Hz), 1.41 (9H, s), 2.9-3.2 (m), 3.22 (2H, q, J=7.3Hz), 4.25 (1H, dd, J=14.3Hz, J=7.5Hz), 5.04 (1H, br. d) 5.61 (1H, br.

s), 7.25 (5H, s)

Referential Example 4-B

Preparation of compound 6-B

(in compound 6, A = CONH-Et, * = R)

5 The desired compound 6-B is obtained from BOC-D-phenylalanine in all the same manner as in Referential Example 4-A.

Referential Example 4-C

Preparation of compound 6-C

10

(in compound 6, A = , * = S)

133 mg (0.5 mmole) of Boc-phenylalanine and 50 mg (0.5 mmole) of 2-aminothiazole are dissolved in 1 ml of dimethylformamide, and while the solution is stirred at 0°, 86 mg (1 equivalent) of DEPC and 70 µl (1 equivalent) of triethylamine are added. After stirring is continued at 0° for 3 hours and then at room temperature overnight, the mixture is evaporated to dryness under reduced pressure, the residue is dissolved in dichloromethane, and the solution is washed with 10 % citric acid and saturated aqueous sodium bicarbonate and dried. The crude product is purified by preparative TLC using ethyl acetate - n-hexane (3 : 4) as a developing solvent to obtain the desired compound 6-C as sand-like crystals. 128 mg (73.6 %). Melting point 158-160°.

[α]_D²⁵ -11.0° (c = 0.2, CHCl₃)
¹H-NMR (CDCl₃, δ) 1.41 (9H, s), 3.0-3.3 (2H, m), 4.5-4.8 (1H, m), 5.0-5.2 (1H, br. d), 7.23 (5H, m), 7.26 (2H, dd, J=41.3Hz, J=3.7Hz)

30

Referential Example 4-D

Preparation of compound 6-D

(in compound 6, A = , * = S)

35

The desired compound 6-D is obtained from BOC-phenylalanine and 2-amino-1,3,4-thiadiazole in all

the same manner as in Referential Example 4-C.

$$[\alpha]_D^{27} +34.1^\circ \text{ (c = 0.960, MeOH)}$$

¹H-NMR (CDCl₃, δ) 1.28 (9H, s), 3.0-3.3 (2H, m), 4.6-4.9 (1H, m), 6.27 (1H, d, J=7.3Hz), 7.26 (5H, s), 8.84 (1H, s), 13.5 (1H, br. s)

Referential Example 4-E

Preparation of compound 6-E


(in compound 6, A = CSNH-Et, * = S)

0.217 g (0.745 mmole) of compound 6-A obtained in Referential Example 4-A and 151 mg (0.5 equivalent) of Lawesson reagent are dissolved in 5 ml of benzene, and the solution is refluxed with heating for 45 minutes. The reaction solution is evaporated to dryness under reduced pressure, and the residue is purified by preparative TLC using dichloromethane - methanol (40 : 1) as a developing solvent to obtain the desired thioamide (compound 6-E) as a yellow waxy solid. 0.230 g (quantitative).

¹H-NMR (CDCl₃, δ) 1.01 (3H, t, J=7.3Hz), 1.41 (9H, s), 3.0-3.2 (2H, m), 3.3-3.7 (2H, m), 4.48 (1H, dd, J=14.5Hz, J=7.9Hz), 5.25-5.55 (1H, br. d), 7.24 (5H, s)

Example 1

Preparation of compound 5-A

(in compound 5, B = )

(a) 400 mg (0.58 mmole) of compound 4 obtained in Referential Example 3 is dissolved in 6 ml of t-butanol - water (9 : 1), 80 mg of 5 % palladium-carbon is added, and the solution is stirred under a stream of hydrogen for 5 hours. The catalyst is filtered and washed, and the filtrate and the washings are evaporated to dryness under reduced pressure and dried to obtain compound 4', a carboxylic acid, as a colorless glassy solid. 337 mg (quantitative).

(b) 35 mg (60 μmoles) of the carboxylic acid obtained in (a) and 14 mg (1.5 equivalents) of





p-chlorophenethylamine are dissolved in 0.5 ml of dimethylformamide, and under ice cooling and stirring, 12.4 mg (1.2 equivalents) of DEPC and 16 μ l (1.88 equivalents) of triethylamine are added, and stirring is continued at 0° for at least 3 hours and then overnight allowing the ice to melt. The reaction solution is concentrated under reduced pressure, the residue is dissolved in dichloromethane, and the solution is washed with saturated aqueous sodium bicarbonate and saturated saline and dried. The resultant crude product is purified by preparative TLC using dichloromethane - methanol (10 : 1) as a developing solvent and then Sephadex LH-20 chromatography using n-hexane - dichloromethane - methanol (2 : 7.5 : 2.5) as an eluent to obtain the desired compound 5-A as amorphous powder. 35.2 mg (79.6 %).





$$[\alpha]_D^{28} -32.9^\circ \text{ (c = 0.292, MeOH)}$$





¹H-NMR (CDCl₃, δ) 0.7-1.1 (m), 1.22 (3H, d, J=7.0Hz), 2.26 (6H, s), 3.03 (3H, s), 3.31 (3H, s), 3.36 (3H, s), 3.7-4.2 (m), 4.79 (1H, dd, J=9.2Hz, 6.6Hz), 6.86 (1H, br. d), 7.1-7.3 (4H, m)



Examples 2 to 15

The following compounds are obtained by reacting compound 4' with the corresponding phenethylamine derivative in the same manner as in Example 1.

Example	Compound	B	$[\alpha]_D$ (MeOH)	$^1\text{H-NMR}$ (CDCl ₃ , δ)
2	5 - B		-39.3° (C=0.293) (28°)	0.7~1.1 (m)、1.21 (3H, d, J=7.0Hz)、 1.5~2.2 (m)、2.2~2.65 (m)、 3.02 (3H, s)、3.31 (3H, s)、 3.36 (3H, s)、3.7~4.25 (m)、 4.76 (1H, dd, J=8.6Hz, J=6.7Hz)、 6.9~7.3 (4H, m)
3	5 - C		-31.3° (C=0.364) (27°)	0.65~1.3 (m)、1.21 (3H, d, J=7.3Hz)、 1.5~2.3 (m)、2.52 (6H, s)、 3.05 (3H, s)、3.30 (3H, s)、 3.34 (3H, s)、3.7~4.2 (m)、 4.5~4.8 (m)、 6.89 (4H, dd, J=24.4Hz, J=8.6Hz)
4	5 - D		-23.7° (C=0.351) (27°)	0.7~1.15 (m)、1.21 (3H, d, J=7.0Hz)、 2.90 (6H, s)、3.01 (3H, s)、 3.31 (3H, s)、3.36 (3H, s)、 3.78 (3H, s)、4.5~4.85 (m)、 6.97 (4H, dd, J=27.9Hz, J=8.8Hz)
5	5 - E		-38.0° (C=0.277) (28°)	0.65~1.2 (m)、1.22 (3H, d, J=7.3Hz)、 1.5~2.3 (m)、2.55 (6H, br. s)、 3.02 (3H, s)、3.31 (3H, s)、 3.36 (3H, s)、3.7~4.25 (m)、 4.75 (1H, dd, J=8.6Hz, J=6.8Hz) 6.7~7.3 (4H, m)

Example	Compound	B	$[\alpha]_D$ (MeOH)	$^1\text{H-NMR}$ (CDCl ₃ , δ)
6	5 - F		-31.2° ($C=0.284$) (28°)	0.7~1.2 (m)、1.22 (3H, d, $J=7.1\text{Hz}$)、 1.5~2.3 (m)、2.5~2.9 (m)、 3.02 (3H, s)、3.31 (3H, s)、 3.36 (3H, s)、3.7~4.25 (m)、 4.5~4.9 (m)、7.18 (4H, s)
7	5 - G		-30.3° ($C=0.307$) (25°)	0.7~1.1 (m)、1.19 (3H, d, $J=7.3\text{Hz}$)、 1.5~2.2 (m)、2.39 (6H, s)、 2.65~2.9 (m)、3.08 (3H, s)、 3.33 (3H, s)、3.37 (3H, s)、 3.6~4.25 (m)、4.6~5.05 (m)、 5.9~6.2 (m)、6.55~7.2 (5H, m)
8	5 - H		-44.6° ($C=0.435$) (23°)	0.65~1.2 (m)、1.21 (3H, d, $J=7.0\text{Hz}$)、 1.5~2.3 (m)、2.5 (6H, br. s)、 3.01 (3H, s)、3.31 (3H, s)、 3.36 (3H, s)、4.0~4.3 (m)、 4.76 (1H, dd, $J=8.6\text{Hz}$, $J=6.4\text{Hz}$)、 6.9~7.3 (4H, m)
9	5 - I		-44.5° ($C=0.339$) (22°)	0.7~1.2 (m)、1.22 (3H, d, $J=6.8\text{Hz}$)、 1.5~2.2 (m)、2.3~2.7 (m)、 3.01 (3H, s)、3.31 (3H, s)、 3.37 (3H, s)、3.75~3.95 (m)、 4.0~4.25 (m)、 4.77 (1H, dd, $J=8.8\text{Hz}$, $J=6.6\text{Hz}$)、 7.0~7.4 (4H, m)

Example	Compound	B	$[\alpha]_D$ (MeOH)	$^1\text{H-NMR}$ (CDCl ₃ , δ)
10	5 - J		-35.2° ($C=0.328$) (28°)	0.7~1.2 (m)、1.23 (3H, d, $J=7.3\text{Hz}$)、 1.4~2.25 (m)、2.63 (6H, br. s)、 3.01 (3H, s)、3.31 (3H, s)、 3.37 (3H, s)、3.75~3.95 (m)、 3.95~4.3 (m)、4.5~4.9 (m)、 7.0~7.4 (4H, m)
11	5 - K		-33.7° ($C=0.348$) (26°)	0.65~1.1 (m)、1.23 (3H, d, $J=6.8\text{Hz}$)、 1.5~2.2 (m)、2.34 (3H, s)、 2.92 (6H, s)、3.00 (3H, s)、 3.31 (3H, s)、3.37 (3H, s)、 3.95~4.3 (m)、4.5~4.85 (m)、 7.13 (4H, s)
12	5 - L		-35.0° ($C=0.345$) (25°)	0.7~1.2 (m)、1.24 (3H, d, $J=7.0\text{Hz}$)、 1.5~2.3 (m)、2.47 (6H, br. s)、 3.06 (3H, s)、3.33 (3H, s)、 3.34 (3H, s)、3.9~4.3 (m)、 4.77 (1H, dd, $J=8.8\text{Hz}$, $J=6.7\text{Hz}$)、 6.65~7.2 (4H, m)
13	5 - M		-37.5° ($C=0.311$) (28°)	0.65~1.1 (m)、1.20 (3H, d, $J=7.0\text{Hz}$)、 1.5~2.25 (m)、2.48 (6H, br. s)、 3.01 (3H, s)、3.31 (3H, s)、 3.35 (3H, s)、3.83 (3H, s)、 3.95~4.25 (m)、 4.76 (1H, dd, $J=8.4\text{Hz}$, $J=6.6\text{Hz}$)、 6.7~7.3 (4H, m)

Example	Compound	B	$[\alpha]_D$ (MeOH)	$^1\text{H-NMR}$ (CDCl ₃ , δ)
14	5 - N		n. d.	0.6 ~ 1.15 (m)、1.24 (3H, d, $J=6.8\text{Hz}$)、 1.5 ~ 2.2 (m)、2.30 (6H, s)、 3.02 (3H, s)、3.32 (3H, s)、 3.37 (3H, s)、4.0 ~ 4.3 (m)、 4.79 (1H, dd, $J=9.0\text{Hz}$, $J=6.7\text{Hz}$)、 6.7 ~ 7.2 (3H, m)
15	5 - O		-41.2° ($C=1.68$) (25°)	0.7 ~ 1.1 (m)、1.21 (3H, d, $J=7.3\text{Hz}$)、 1.5 ~ 2.3 (m)、2.54 (6H, br. s)、 3.01 (3H, s)、3.31 (3H, s)、 3.36 (3H, s)、3.5 ~ 3.95 (m)、 4.0 ~ 4.3 (m)、 4.75 (1H, dd, $J=9.2\text{Hz}$, $J=6.4\text{Hz}$)、 7.0 ~ 7.75 (4H, m)

Example 16

Preparation of compound 7-A

(in compound 7, A = , * = S)

5 70 mg (0.2 mmole) of compound 6-C obtained in Referential Example 4-C is dissolved in 1 ml of 50 % trifluoroacetic acid-dichloromethane at 0°, and the solution is brought to room temperature, stirred for 3
10 hours and evaporated under reduced pressure. The residue is washed thoroughly with ether and dried under reduced pressure.


The above compound and 88 mg (0.15 mmole) of compound 4' obtained in (a) of Example 1 are dissolved
15 in 1.5 ml of dimethylformamide, and under ice cooling and stirring, 30 mg (0.184 mmole) of DEPC and 41 mg (0.406 mmole) of triethylamine are added. Stirring is continued at 0° for 3 hours and then at room temperature overnight, the reaction solution is evaporated under re-
20 duced pressure, the residue is dissolved in dichloromethane, and the solution is washed with saturated aqueous sodium bicarbonate and saturated saline and dried. The resultant product is purified by preparative TLC using dichloromethane - methanol (10 : 1) as a
25 developing solvent to obtain the desired compound 7-A as white powder. 86.5 mg (69.6 %).

$[\alpha]_D^{28} -26.1^\circ$ (c = 0.3185, MeOH)
 $^1\text{H-NMR}$ (CDCl_3 , δ) 0.7-1.3 (m), 1.11 (3H, d, J=6.6Hz), 2.98 (3H, s), 2.99 (3H, s), 3.34 (6H, s),
 30 3.5-4.2 (m), 4.5-5.1 (m), 7.23 (5H, s), 7.41 (2H, dd, J=56.5Hz, J=4.0Hz)

Examples 17 to 24

Compounds 6-A, 6-B, 6-D and 6-E obtained in Referential Examples 4-A, 4-B, 4-D and 4-E are
 35 deprotected respectively according to Example 16, and then reacted with compound 4' to obtain the desired

compounds 7-B, 7-C, 7-D and 7-E. Likewise, by reaction of L- or D-phenylalanine methyl ester hydrochloride or phenylalanine ethyl ester or phenylalaninol with compound 4', the corresponding compounds 7-F, 7-G, 7-H and 5 7-I are obtained. The results are shown in the following Table.

Example	Compound	A	*	$[\alpha]_D$ (MeOH)	$^1\text{H-NMR}$ (CDCl ₃ , δ)
17	7 - B	-CONH-Et	S	- 51.2° (C=0.3475) (27°)	0.6 ~ 1.2 (m), 1.5 ~ 2.0 (m), 2.42 (6H, s), 3.02 (3H, s), 3.31 (3H, s), 3.35 (3H, s), 3.6 ~ 4.2 (m), 4.5 ~ 4.9 (m), 7.25 (5H, s)
18	7 - C	-CONH-Et	R	- 50.8° (C=0.323) (27°)	0.6 ~ 1.1 (m), 1.1 ~ 1.4 (m), 1.5 ~ 2.5 (m), 2.96 (3H, s), 3.24 (3H, s), 3.33 (3H, s), 3.6 ~ 4.2 (m), 4.5 ~ 4.9 (m), 7.23 (5H, s)
19	7 - D	-CONH- 	S	- 30.6° (C=0.310) (23°)	0.7 ~ 1.2 (m), 1.5 ~ 2.2 (m), 2.36 (6H, br. s), 3.10 (3H, s), 3.34 (3H, s), 3.36 (3H, s), 4.5 ~ 5.1 (m), 5.90 (1H, br. d), 7.25 (5H, s), 8.82 (1H, s)
20	7 - E	-CSNH-Et	S	- 47.8° (C=0.347) (22°)	0.6 ~ 1.2 (m), 1.5 ~ 2.2 (m), 2.34 (6H, s), 3.08 (3H, s), 3.31 (3H, s), 3.37 (3H, s), 3.8 ~ 4.2 (m), 4.5 ~ 5.1 (m), 7.25 (5H, s)
21	7 - F	-COOMe	S	n. d.	n. d.

Example	Compound	A	*	$[\alpha]_D$ (MeOH)	$^1\text{H-NMR}$ (CDCl ₃ , δ)
22	7 - G	-COOMe	R	- 48.7° (C=0.3285) (28°)	0.7~1.2 (m)、1.21 (3H, d, J=7.3Hz)、 1.5~2.3 (m)、2.50 (6H, br. s)、 3.02 (3H, s)、3.32 (3H, s)、 3.34 (3H, s)、3.70 (3H, s)、 3.8~4.3 (m)、4.5~5.0 (m)、 7.0~7.4 (5H, m)、
23	7 - H	-COOEt	S	- 56.4° (C=0.3265) (27°)	0.7~1.1 (m)、1.21 (3H, t, J=7.3Hz) 2.70 (6H, br. s)、3.02 (3H, s)、 3.32 (3H, s)、3.34 (3H, s)、 4.15 (2H, q, J=7.3Hz)、 4.5~4.9 (m)、7.24 (5H, s)、
24	7 - I	-CH ₂ OH	S	- 51.6° (C=0.2735) (27°)	0.6~1.2 (m)、1.13 (3H, d, J=7.0Hz)、 1.5~2.3 (m)、2.44 (6H, br. s)、 3.02 (3H, s)、3.32 (3H, s)、 3.40 (3H, s)、3.9~4.4 (m)、 4.76 (1H, dd, J=8.8Hz, J=6.8Hz)、 7.25 (5H, s)

Example 25

Preparation of compound 7-J

(in compound 7, A = COOH, * = S)

38.1 mg (50 μ moles) of compound 7-F obtained
 5 in Example 21 is dissolved in 0.5 ml of methanol, 55 μ l
 (55 μ moles) of 1N-sodium hydroxide is added, and the
 mixture is stirred at room temperature for 3 hours. The
 mixture is ice-cooled, 55 μ l of 1N-hydrochloric acid is
 added, the mixture is evaporated to dryness under re-
 10 duced pressure, and the tetrahydrofuran-soluble part of
 the residue is purified by Sephadex LH-20 chromatography
 using n-hexane - dichloromethane - methanol (2 : 7.5 :
 2.5) as an eluent to obtain powder of the desired com-
 pound 7-J. 37.4 mg (100 %).

15 $[\alpha]_D^{30} -33.4^\circ$ (c = 0.3265, MeOH)
 $^1\text{H-NMR}$ (CDCl_3 , δ) 0.6-1.3 (m), 1.5-2.2 (m),
 2.6-2.8 (6H, m), 3.07 (3H, s), 3.33 (3H, s), 3.38
 (3H, s), 3.6-4.2 (m), 4.5-4.9 (m), 7.21 (5H, s)

Example 26

20 Preparation of compound 7-K

(in compound 7, A = COOH, * = R)


Compound 7-K is obtained from compound 7-G in
 all the same manner as in Example 25.

25 $[\alpha]_D^{29} -63.4^\circ$ (c = 0.330, MeOH)
 $^1\text{H-NMR}$ (CDCl_3 , δ) 0.7-1.4 (m), 1.5-2.3 (m),
 2.71 (6H, br. s), 3.04 (3H, s), 3.32 (3H, s), 3.37
 (3H, s), 4.6-5.0 (m), 7.22 (5H, s)



CLAIMS

1. A peptide derivative represented by the following formula or a salt thereof



wherein A and B each represent either of the following (a) and (b),

(a) A represents a hydrogen atom, and B represents a phenyl group substituted with a halogen atom, hydroxyl group, lower alkyl group or lower alkoxy group, or a heteroaryl group

(b) A represents -CONH-R¹, -CSNH-R¹, a hydroxymethyl group, a lower alkoxycarbonyl group or a carboxyl group, wherein, R¹ represents a lower alkyl group or a heteroaryl group, and B represents a phenyl group optionally substituted with a halogen atom, hydroxyl group, lower alkyl group or lower alkoxy group.

2. The peptide derivative or salt thereof according to claim 1 wherein A represents a hydrogen atom, and B represents a phenyl group substituted with a halogen atom, hydroxyl group, lower alkyl group or lower alkoxy group, or a heteroaryl group.
3. The peptide derivative or salt thereof according to claim 1 wherein A represents a hydrogen atom, and B represents a phenyl group substituted with a halogen atom, hydroxyl group, lower alkyl group or lower alkoxy group; a thienyl group; or a pyridyl group.
4. The peptide derivative or salt thereof accord-



ing to claim 1 wherein A represents $-\text{CONH}-\text{R}^1$, $-\text{CSNH}-\text{R}^1$, a hydroxymethyl group, a lower alkoxy carbonyl group or a carboxyl group, wherein R^1 represents a lower alkyl group or a heteroaryl group, and B represents a phenyl group optionally substituted with a halogen atom, hydroxyl group, lower alkyl group or lower alkoxy group.

5. The peptide derivative or salt thereof according to claim 1 wherein A represents $-\text{CONH}-\text{R}^1$, $-\text{CSNH}-\text{R}^1$, a hydroxymethyl group, a lower alkoxy carbonyl group or a carboxyl group, wherein R^1 represents a lower alkyl group, a thiazolyl group or a thiadiazolyl group, and B represents an unsubstituted phenyl group.