
ABSTRACT



NEW IMIDAZOPYRIDINE COMPOUNDS, A PROCESS FOR THEIR PREPARATION AND PHARMACEUTICAL COMPOSITIONS CONTAINING THEM

Compounds of formula (I):



(I)

wherein:

- R^1 represents a hydrogen atom, a halogen atom or an alkyl, polyhaloalkyl, cyano, nitro, hydroxycarbonyl, alkoxycarbonyl, aminocarbonyl, alkylaminocarbonyl or dialkylaminocarbonyl group,
- R^2 represents a hydrogen atom, an alkyl group, an optionally substituted aryl group, an optionally substituted heteroaryl group, or a group R^{20}–C(X)– wherein:
  - R^{20} represents an alkyl group, an alkoxy group, an amino group, an alkylamino group, a dialkylamino group, an optionally substituted aryl group or an optionally substituted heteroaryl group,
  - X represents an oxygen atom, a sulphur atom, or a group NR^{21} wherein R^{21} represents a hydrogen atom or an alkyl group,
- R^3 represents a hydrogen atom or an alkyl group,
- n represents an integer from 1 to 6 inclusive,
- the representation  means: 

their enantiomers, diastereoisomers and also addition salts thereof with a pharmaceutically acceptable acid or base.

Medicaments.