

DESCRIPTION

METHOD FOR PRODUCING *cis*-5-HYDROXY-2-PIPERIDINECARBOXYLIC
ACID DERIVATIVE, AND METHOD FOR PURIFYING *cis*-5-HYDROXY-2-
5 PIPERIDINECARBOXYLIC ACID

TECHNICAL FIELD

[0001]

The present invention relates to an industrial method for producing a
10 derivative of *cis*-5-hydroxy-2-piperidinecarboxylic acid (which is also called *cis*-5-hydroxypipicolic acid). The present invention also relates to a method for purifying *cis*-5-hydroxy-2-piperidinecarboxylic acid.

BACKGROUND ART

15 [0002]

cis-5-Hydroxy-2-piperidinecarboxylic acid is a useful intermediate for
pharmaceuticals. However, since this compound has 2 asymmetric carbons, 4 kinds
of isomers are present, and selective synthesis of a single kind of isomer, or single
kind of diastereomer, is very difficult. Thus, improvement of the purity by
20 separation of unnecessary isomers, or purification, by chemical conversion is
necessary.

[0003]

For example, methods for obtaining *cis*-5-hydroxy-2-piperidinecarboxylic
acid by introduction of a hydroxyl group to 2-piperidinecarboxylic acid by
25 hydroxylation reaction using a microorganism or enzyme have been reported.
However, it has been reported that a compound having a substituted hydroxyl group
at another position, such as a 3-position-hydroxylated compound, may be produced

as a by-product in addition to the compound of interest, *cis*-5-hydroxy-2-piperidinecarboxylic acid (Non-patent Document 1). Separation of the by-product is not described in the document. A method in which 5-hydroxy-2-piperidinecarboxylic acid is synthesized from 5-hydroxylysine by enzymatic reaction has also been reported (Patent Document 1), but the document does not describe separation of the generated *cis/trans* isomers. In a report describing synthesis of 5-hydroxy-2-piperidinecarboxylic acid from 5-hydroxylysine by a method similar to the method of Patent Document 1 (Patent Document 2), isomers are separated using an ion-exchange column. Since this method requires use of excessive amounts of a filler and eluent with respect to the substrate, the method is not realistic from the viewpoint of industrial production.

Thus, there is no known method by which a necessary stereoisomer can be selectively obtained with high purity from *cis*-5-hydroxy-2-piperidinecarboxylic acid synthesized using a microorganism or enzyme.

[0004]

There are known chemical synthesis methods using, as a material, L-amino acid, in which the stereochemistry of one of the two asymmetric carbons, the 2-position carbon, can be fixed. For example, a method using L-pyroglutamic acid as a material (Patent Document 3) has been reported. This method requires use of an iridium catalyst, which is expensive, for formation of a piperidine ring, which is problematic. Methods using L-glutamic acid as a material (Non-patent Documents 2 and 3) and a method using a proline derivative as a material (Non-patent Document 4) have also been reported, but all of these methods require use of a diazo compound, which is highly risky, and also require a multistep complex process. Moreover, for separation of (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid, the reaction is carried out via 5-oxo-2-piperidinecarboxylic acid to preferentially obtain the (2*S*,5*S*) compound by its reduction (Non-patent Document 4). This requires separation

using a silica gel column for removal of impurities such as isomers. Because of the load of this process, this method is not industrially satisfactory.

[0005]

It is also known that (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid and its esters are lactonized to give benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate (Non-patent Documents 4 and 5).

These documents on lactonization do not mention about impurities such as isomers, and the effects of purification by these methods have been unclear. Non-patent

Document 4 describes that benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-

carboxylate was obtained as an oily substance after purification with a silica gel

column. Non-patent Document 5 describes that crystallization of benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate occurs, but, since the

lactonization is carried out using *cis*-5-hydroxy-2-piperidinecarboxylic acid as a material, behavior of impurities such as isomers and the effect of purification are not

clear. There is a report describing separation of a *cis*/*trans* mixture of methyl *N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylate by acid treatment (Non-patent Document 3), but, since the compound was a carboxylic acid ester, it was impossible to separate lactone obtained from the *cis* isomer from the ester of the *trans* isomer by a simple method such as solvent extraction. Moreover, Non-patent Document 3

does not describe the yield in this process. When the present inventors studied this method, the yield of the lactone was low because of occurrence of a remarkable side reaction between lactone obtained from the *cis* isomer (benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate)) and the residual *trans* isomer ((2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid ester).

As described above, there are problems to be solved for industrial production of highly pure *cis*-5-hydroxy-2-piperidinecarboxylic acid by an inexpensive method.

PRIOR ART DOCUMENTS

[Patent Documents]

[0006]

Patent Document 1: JP 4590981 B

5 Patent Document 2: JP 2010-88395 A

Patent Document 3: WO 2010/126820

[Non-patent Documents]

[0007]

Non-patent Document 1: Adv. Synth. Catal., 2011, 353, 1375.

10 Non-patent Document 2: Chem. Commun., 1996, 349.

Non-patent Document 3: Tetrahedron Lett., 1988, 29, 2231.

Non-patent Document 4: Tetrahedron: Asym., 2006, 17, 2479.

Non-patent Document 5: Rec. Trav. Chim. Pays-Bas, 1959, 78, 648.

15 SUMMARY OF THE INVENTION

PROBLEMS TO BE SOLVED BY THE INVENTION

[0008]

The present invention aims to provide a method for purifying *cis*-5-hydroxy-2-piperidinecarboxylic acid, which is a useful intermediate for pharmaceuticals, and a
20 method for producing its derivative.

MEANS FOR SOLVING THE PROBLEMS

[0009]

The present inventors intensively studied to solve the problems described
25 above. As a result, the present inventors solved the problems by a process of reacting low-purity *cis*-5-hydroxy-2-piperidinecarboxylic acid containing impurities with an acid halide and/or acid anhydride to induce a *cis*-5-hydroxy-2-

piperidinecarboxylic acid derivative, thereby solving the problems.

In a more specific mode, the present inventors discovered that impurities such as isomers can be separated by reacting *cis*-5-hydroxy-2-piperidinecarboxylic acid with an acid halide and/or acid anhydride to convert the *cis*-5-hydroxy-2-
5 piperidinecarboxylic acid into a lactone whose amino group is protected, and then performing crystallization and/or solvent extraction. That is, studies by the present inventors revealed that the *cis*-5-hydroxy-2-piperidinecarboxylic acid derivative in the present invention can be selectively converted to lactone by reaction with an acid halide and/or acid anhydride, and that this lactone can be crystallized under specific
10 conditions. The present inventors then developed a method in which the lactone is crystallized to achieve removal of impurities such as isomers having low crystallizability.

In another specific mode, the present inventors discovered that impurities such as isomers having a carboxyl group can be removed by solvent extraction after
15 reacting *cis*-5-hydroxy-2-piperidinecarboxylic acid with an acid halide and/or acid anhydride, and then with an alcohol in the presence of an acid catalyst to convert the *cis*-5-hydroxy-2-piperidinecarboxylic acid into an ester whose amino group is protected. That is, the present inventors studied to develop a method in which, in the presence of an acid catalyst, a *cis*-5-hydroxy-2-piperidinecarboxylic acid
20 derivative is selectively converted into lactone, and the lactone is reacted with an alcohol present in the reaction system to efficiently and selectively convert the *cis* isomer into an ester while suppressing a side reaction, followed by removing residual impurities such as isomers having a carboxyl group by solvent extraction.

The present inventors then discovered that highly pure *cis*-5-hydroxy-2-
25 piperidinecarboxylic acid can be obtained by hydrolyzing the resulting lactone and/or ester to give *cis*-*N*-protected-5-hydroxy-2-piperidinecarboxylic acid, and then removing the protective group for the amino group, thereby completing the present


invention.

[0010]

That is, the present invention can be summarized as follows.

[1] A method for producing a *cis*-5-hydroxy-2-piperidinecarboxylic acid
 5 derivative, the method comprising a step of converting *cis*-5-hydroxy-2-
 piperidinecarboxylic acid into a compound(s) of Formula (1) and/or Formula (2)
 (wherein R¹ represents a protective group for an amino group, and R² represents a
 C₁-C₆ alkyl group).

[0011]



[0012]

[2] The method for producing a *cis*-5-hydroxy-2-piperidinecarboxylic acid
 derivative according to [1], comprising a step of reacting *cis*-5-hydroxy-2-
 piperidinecarboxylic acid with an acid halide and/or acid anhydride to convert the
 15 *cis*-5-hydroxy-2-piperidinecarboxylic acid into a compound of Formula (1).

[3] The method for producing a *cis*-5-hydroxy-2-piperidinecarboxylic acid
 derivative according to [1], comprising a step of reacting *cis*-5-hydroxy-2-
 piperidinecarboxylic acid with an acid halide and/or acid anhydride, and then with an
 alcohol in the presence of an acid catalyst, to convert the *cis*-5-hydroxy-2-
 20 piperidinecarboxylic acid into a compound of Formula (2).


[4] A method for regenerating *cis*-5-hydroxy-2-piperidinecarboxylic acid,
 the method comprising the steps of:

converting *cis*-5-hydroxy-2-piperidinecarboxylic acid into a compound(s) of
 Formula (1) and/or Formula (2) (wherein R¹ represents a protective group for an


amino group, and R^2 represents a C_1 - C_6 alkyl group); and

converting the compound(s) of Formula (1) and/or Formula (2) into *cis*-5-hydroxy-2-piperidinecarboxylic acid.

[0013]



(1)



(2)


[0014]

[5] The method for producing a *cis*-5-hydroxy-2-piperidinecarboxylic acid derivative according to any one of [1] to [3], wherein the *cis*-5-hydroxy-2-piperidinecarboxylic acid is *cis*-5-hydroxy-2-piperidinecarboxylic acid synthesized by bacterial reaction and/or enzymatic reaction.


[6] The method for regenerating *cis*-5-hydroxy-2-piperidinecarboxylic acid according to [4], wherein the *cis*-5-hydroxy-2-piperidinecarboxylic acid is *cis*-5-hydroxy-2-piperidinecarboxylic acid synthesized by bacterial reaction and/or enzymatic reaction.

[7] A method for purifying *cis*-5-hydroxy-2-piperidinecarboxylic acid, the method comprising a step of reacting a mixture containing *cis*-5-hydroxy-2-piperidinecarboxylic acid and an impurity with an acid halide and/or acid anhydride, or with an acid halide and/or acid anhydride and an alcohol, to convert the *cis*-5-hydroxy-2-piperidinecarboxylic acid into a compound(s) of Formula (1) and/or Formula (2), separating the compound(s) and then converting the separated compound(s) into *cis*-5-hydroxy-2-piperidinecarboxylic acid.

[0015]



(1)



(2)

[0016]

[8] The method for purifying *cis*-5-hydroxy-2-piperidinecarboxylic acid according to [7], wherein the impurity is 2-piperidinecarboxylic acid or an analogue thereof.

[9] The method for purifying *cis*-5-hydroxy-2-piperidinecarboxylic acid according to [8], wherein the 2-piperidinecarboxylic acid or an analogue thereof is *trans*-5-hydroxy-2-piperidinecarboxylic acid.

[10] The method for purifying *cis*-5-hydroxy-2-piperidinecarboxylic acid according to any one of [7] to [9], wherein the step of separating the compound(s) of Formula (1) and/or Formula (2) is carried out by crystallization or solvent extraction.

[11] The method for purifying *cis*-5-hydroxy-2-piperidinecarboxylic acid according to any one of [7] to [10], wherein the mixture containing *cis*-5-hydroxy-2-piperidinecarboxylic acid and an impurity is a mixture synthesized by bacterial reaction and/or enzymatic reaction.

EFFECT OF THE INVENTION

[0017]

By the method of the present invention, highly pure *cis*-5-hydroxy-2-piperidinecarboxylic acid and its derivatives can be provided.

BRIEF DESCRIPTION OF THE DRAWINGS

[0018]

Fig. 1 is a schematic diagram illustrating routes for purifying *cis*-5-hydroxy-2-

piperidinecarboxylic acid from *cis*-5-hydroxy-2-piperidinecarboxylic acid that may contain an impurity(s). The route 1 is a route in which *cis*-5-hydroxy-2-piperidinecarboxylic acid is reacted with an acid halide and/or acid anhydride to give a compound of Formula (1) directly or via *cis*-*N*-protected-5-hydroxy-2-piperidinecarboxylic acid, and the compound is then converted into *cis*-5-hydroxy-2-piperidinecarboxylic acid. The route 2 is a route in which *cis*-*N*-protected-5-hydroxy-2-piperidinecarboxylic acid is reacted with an acid catalyst to give a compound of Formula (1), and the compound is then converted into *cis*-5-hydroxy-2-piperidinecarboxylic acid by hydrolysis. The route 3 is a route in which *cis*-*N*-protected-5-hydroxy-2-piperidinecarboxylic acid is reacted with an alcohol in the presence of an acid catalyst to give a compound of Formula (2), and the compound is then converted to *cis*-5-hydroxy-2-piperidinecarboxylic acid by hydrolysis.

Fig. 2. is a diagram showing the result of HPLC analysis of hydroxylysines obtained with recombinant lysine hydroxylases.

Fig. 3 is a diagram showing the result of HPLC analysis of hydroxylysines obtained with recombinant lysine hydroxylases.

DESCRIPTION OF THE EMBODIMENTS

[0019]


The present invention is described below in detail.

In the present invention, examples of the *cis*-5-hydroxy-2-piperidinecarboxylic acid include (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid, (2*R*,5*R*)-5-hydroxy-2-piperidinecarboxylic acid, and mixtures thereof. The *cis*-5-hydroxy-2-piperidinecarboxylic acid may be a racemic compound. The *cis*-5-hydroxy-2-piperidinecarboxylic acid may be forming a salt with an acid or base.


[0020]

The method for producing a *cis*-5-hydroxy-2-piperidinecarboxylic acid

derivative of the present invention comprises a step of converting *cis*-5-hydroxy-2-piperidinecarboxylic acid into a compound(s) of Formula (1) and/or Formula (2).



(1)



(2)

[0021]

5 In Formula (1) and Formula (2), R¹ represents a protective group for an amino group, and specific examples of the protective group include the following groups. However, the protective group is not limited to these examples.

Examples of the protective group for the amino group include acyl groups such as formyl, acetyl, chloroacetyl, dichloroacetyl, trichloroacetyl, trifluoroacetyl, propionyl, benzoyl, and 4-chlorobenzoyl; alkoxycarbonyl groups such as methoxycarbonyl, ethoxycarbonyl, *tert*-butoxycarbonyl, benzyloxycarbonyl, and allyloxycarbonyl; arylalkyl groups such as benzyl, 4-methoxybenzyl, 4-bromobenzyl, and 1-phenethyl; and sulfonyl groups such as methanesulfonyl, *p*-toluenesulfonyl, and 2-nitrobenzenesulfonyl.

15 [0022]

Among these, acyl groups and alkoxycarbonyl groups are preferred since these groups can be easily removed; acetyl, chloroacetyl, trifluoroacetyl, benzoyl, *tert*-butoxycarbonyl, and benzyloxycarbonyl are more preferred; and acetyl, *tert*-butoxycarbonyl, and benzyloxycarbonyl are still more preferred since these are industrially inexpensive. The protective group is especially preferably benzyloxycarbonyl since benzyloxycarbonyl can be removed by hydrogenation without leaving a non-volatile component, and the load of purification of *cis*-5-hydroxy-2-piperidinecarboxylic acid can therefore be reduced.

[0023]

In Formula (2), R^2 represents a C_1 - C_6 alkyl, and specific examples of R^2 include primary alkyl groups such as methyl, ethyl, propyl, butyl, pentyl, and hexyl; secondary alkyl groups such as isopropyl, isobutyl, isopentyl, cyclopentyl, and cyclohexyl; and tertiary alkyl groups such as *tert*-butyl. Among these, methyl, ethyl, isopropyl, butyl, and *tert*-butyl are preferred, and ethyl and isopropyl are more preferred.

[0024]

In the present invention, the *cis*-5-hydroxy-2-piperidinecarboxylic acid derivative may be the compound(s) of Formula (1) and/or Formula (2) itself/themselves.

Specific examples of the compound of Formula (1) include benzyl 5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate, *tert*-butyl 5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate, and 5-acetyl-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane.

Specific examples of the compound of Formula (2) include methyl *cis-N*-(benzyloxycarbonyl)-5-hydroxy-2-piperidinecarboxylate, ethyl *cis-N*-(benzyloxycarbonyl)-5-hydroxy-2-piperidinecarboxylate, isopropyl *cis-N*-(benzyloxycarbonyl)-5-hydroxy-2-piperidinecarboxylate, methyl *cis-N*-(*tert*-butyloxycarbonyl)-5-hydroxy-2-piperidinecarboxylate, ethyl *cis-N*-(*tert*-butyloxycarbonyl)-5-hydroxy-2-piperidinecarboxylate, and isopropyl *cis-N*-(*tert*-butyloxycarbonyl)-5-hydroxy-2-piperidinecarboxylate.

[0025]

The *cis*-5-hydroxy-2-piperidinecarboxylic acid derivative may be a compound obtained by chemical conversion of a compound(s) of Formula (1) and/or Formula (2).

Specific examples of such a *cis*-5-hydroxy-2-piperidinecarboxylic acid derivative include carboxylic acids such as *cis-N*-(benzyloxycarbonyl)-5-hydroxy-2-

piperidinecarboxylic acid and *cis-N*-(*tert*-butyloxycarbonyl)-5-hydroxy-2-piperidinecarboxylic acid; esters such as benzyl *cis-N*-(benzyloxycarbonyl)-5-hydroxy-2-piperidinecarboxylate and benzyl *cis-N*-(*tert*-butyloxycarbonyl)-5-hydroxy-2-piperidinecarboxylate; carboxylic acid amides such as *cis-N*-(benzyloxycarbonyl)-5-hydroxy-2-piperidinecarboxylic acid amide, *cis-N*-(*tert*-butyloxycarbonyl)-5-hydroxy-2-piperidinecarboxylic acid amide, benzyl *cis*-5-hydroxy-2-(methylcarbamoyl)piperidine-1-carboxylate, *tert*-butyl *cis*-5-hydroxy-2-(methylcarbamoyl)piperidine-1-carboxylate, benzyl *cis*-2-(1-*tert*-butyloxycarbamoylpyrrolidine-3-ylcarbamoyl)-5-hydroxypiperidine-1-carboxylate, *tert*-butyl *cis*-2-(1-benzyloxycarbamoylpyrrolidine-3-ylcarbamoyl)-5-hydroxypiperidine-1-carboxylate, benzyl *cis*-2-(1-*tert*-butyloxycarbamoylpiperidine-4-ylcarbamoyl)-5-hydroxypiperidine-1-carboxylate, and *tert*-butyl *cis*-2-(1-benzyloxycarbamoylpiperidine-4-ylcarbamoyl)-5-hydroxypiperidine-1-carboxylate.

[0026]

In one mode of the method of the present invention, *cis*-5-hydroxy-2-piperidinecarboxylic acid is reacted with an acid halide and/or acid anhydride to protect the amino group of the *cis*-5-hydroxy-2-piperidinecarboxylic acid, and subsequent lactonization of the resultant causes its conversion into a compound of Formula (1).

[0027]

The acid halide and/or acid anhydride is/are not limited as long as the acid halide and/or acid anhydride can protect the amino group and allow(s) the lactonization. Examples of the acid halide include acid chlorides and acid bromides. In view of ease of handling, acid chlorides are preferred. Examples of the acid anhydride include carboxylic acid anhydrides, and sulfonic acid anhydrides. In view of ease of handling and the cost, carboxylic acid anhydrides are preferred.

Specific examples of the acid halide and/or acid anhydride include acylating

agents such as formic acid-acetic anhydride, acetic anhydride, acetyl chloride, chloroacetyl chloride, dichloroacetyl chloride, trichloroacetyl chloride, trifluoroacetic anhydride, propionyl chloride, benzoyl chloride, 4-chlorobenzoyl chloride, acetyl bromide, propionyl bromide, and benzoyl bromide; alkoxycarbonylating agents such as di-*tert*-butyldicarbonate, benzyloxycarbonyl chloride, allyloxycarbonyl chloride, benzyloxycarbonyl bromide, and allyloxycarbonyl bromide; and sulfonylating agents such as methanesulfonyl chloride, *p*-toluenesulfonyl chloride, 2-nitrobenzenesulfonyl chloride, methanesulfonyl bromide, *p*-toluenesulfonyl bromide, and 2-nitrobenzenesulfonyl bromide.

10 Among these, acylating agents and alkoxycarbonylating agents are preferred. Acetic anhydride, chloroacetyl chloride, trichloroacetyl chloride, trifluoroacetic anhydride, di-*tert*-butyldicarbonate, benzyloxycarbonyl chloride, and allyloxycarbonyl chloride are more preferred since these can be easily removed after the protection of the amino group. Acetic anhydride, di-*tert*-butyldicarbonate, and benzyloxycarbonyl chloride are still more preferred since these are industrially inexpensive. Benzyloxycarbonyl chloride is especially preferred since it can be removed by hydrogenation without leaving a non-volatile component, and the load of purification of *cis*-5-hydroxy-2-piperidinecarboxylic acid can therefore be reduced. [0028]

20 In the present invention, two or more kinds of acid halide and/or acid anhydride may be used. In cases where two or more kinds of acid halide and/or acid anhydride are used, these may be added to the reaction system at once, but it is preferred to use different kinds of acid halide and/or acid anhydride for in each the process of amino group protection and the process of lactonization. For example, it is preferred to carry out a method in which benzyloxycarbonyl chloride is used for the protection of the amino group, and acetic anhydride is used for the lactonization. In cases where the protection of the amino group and the lactonization are carried out

in a single reactor, a single kind of acid halide and/or acid anhydride is preferably used for suppression of by-products having different protective groups. In cases where the protection of the amino group and the lactonization are carried out in separate reactors, two or more kinds of acid halide and/or acid anhydride are preferably used in view of optimization of the reactions and cost reduction. In particular, the protection of the amino group and the lactonization are preferably carried out in a single reactor since the number of reactors required for the production can be decreased, and the cost can therefore be reduced.

[0029]

The amount of the acid halide and/or acid anhydride to be used is usually 1 to 10 molar equivalents, preferably 1.2 to 5 molar equivalents, more preferably 1.5 to 3 molar equivalents relative to the total amount of the *cis*-5-hydroxy-2-piperidinecarboxylic acid and the amine compound to be protected. The acid halide and/or acid anhydride may be added dividedly in a plurality of times.

[0030]

In the reaction with the acid halide and/or acid anhydride, a base may be used, if necessary. Specific examples of the base include, but are not limited to, tertiary amines such as triethylamine, diisopropylethylamine, *N*-methylmorpholine, quinuclidine, and 1,4-diazabicyclo[2.2.2]octane; pyridines such as pyridine, 4-dimethylaminopyridine, and 2,6-lutidine; organic strong bases such as 1,8-diazabicyclo[5.4.0]undeca-7-ene and tetramethylguanidine; metal amides such as lithium diisopropylamide and sodium hexamethyldisilazide; alkyl metals such as *n*-butyllithium, *sec*-butyllithium, *tert*-butyllithium, and isopropylmagnesium bromide; metal hydrides such as sodium hydride and calcium hydride; metal alkoxides such as sodium methoxide, sodium ethoxide, sodium *tert*-butoxide, and potassium *tert*-butoxide; carbonates such as sodium hydrogen carbonate and potassium carbonate; phosphates such as potassium phosphate and sodium hydrogen phosphate;

hydroxides such as sodium hydroxide, potassium hydroxide, lithium hydroxide, and calcium hydroxide; and cyanides such as sodium cyanide and potassium cyanide.

Preferred bases vary depending on the protective reagent to be used. In cases where a preferred protective reagent, that is, acetic anhydride, di-*tert*-butyldicarbonate, or benzyloxycarbonyl chloride, is used, the base is preferably tertiary amines, pyridines, carbonates, or hydroxides. In cases where the especially preferred protective reagent, that is, benzyloxycarbonyl chloride, is used, the base is preferably a hydroxide, which is inexpensive.

In cases where a base is used, the amount of base is preferably added such that the pH of the reaction mixture becomes 7 to 12. In cases where water is used as the solvent, the pH of the reaction mixture herein means the pH of the layer containing water. In cases where a solvent other than water is used, the pH of the reaction mixture herein means the pH of the layer which contains water when an equal volume of water is added to the reaction mixture. In cases where water is used as the solvent, hydrolysis of the compound of Formula (1) may occur when the basicity of the reaction mixture is too strong. The pH is therefore more preferably 7 to 12, especially preferably 7 to 11.

[0031]

Examples of the reaction solvent include water; esters such as ethyl acetate and butyl acetate; alcohols such as methanol, ethanol, 1-propanol, 2-propanol, 1-butanol, 2-ethyl-1-hexanol, and 2-butanol; ethers such as diethyl ether, di-*n*-butyl ether, diisopropyl ether, di-*n*-butyl ether, *tert*-butyl methyl ether, tetrahydrofuran, and dioxane; ketones such as acetone, methyl ethyl ketone, and methyl isobutyl ketone; nitriles such as acetonitrile; amides such as dimethylformamide, dimethylacetamide, and *N*-methylpyrrolidinone; sulfones such as dimethylsulfoxide and sulfolane; hydrocarbons such as hexane, heptane, and toluene; and mixed solvents of two or more of these. Among these, water, and mixed solvents containing water and one

or more of the above solvents, are preferred. In particular, in cases where the *cis*-5-hydroxy-2-piperidinecarboxylic acid is synthesized by bacterial reaction and/or enzymatic reaction, the *cis*-5-hydroxy-2-piperidinecarboxylic acid is obtained as a solution whose solvent is mainly water. Therefore, in view of simplicity of the production process, it is preferred to use the solution as it is, or to concentrate part of the solution, for carrying out the reaction using water as a main solvent.

[0032]

The reaction temperature is usually -20°C to 100°C, preferably -10°C to 50°C. Since reaction at high temperature may cause decomposition of reagents and products, the reaction temperature is more preferably 0°C to 30°C.

[0033]

The protection of the amino group and the lactonization are preferably carried out by adding an acid halide and/or acid anhydride (the route 1 in Fig. 1). It is especially preferred to carry out the protection of the amino group and the lactonization using a single acid halide and/or acid anhydride in a single reactor in view of simplicity of the operation (the upper formula in the route 1 in Fig. 1).

[0034]

The protection of the amino group and the lactonization may also be carried out under different conditions (the route 2 in Fig. 1). That is, the amino group may be protected by addition of the acid halide and/or acid anhydride, and then an acid catalyst may be used to perform the lactonization. In cases where the protection of the amino group and the lactonization are carried out under different reaction conditions, it is preferred to suppress production of the compound of Formula (1) at the stage of protection of the amino group. The protection of the amino group can be selectively and effectively carried out by using the acid halide and/or acid anhydride in the same molar amount as that of the amino group, or by adding the acid halide and/or acid anhydride dividedly such that excessive presence of the acid halide

and/or acid anhydride relative to the amino group is avoided, and then stopping the reaction when the material *cis*-5-hydroxy-2-piperidinecarboxylic acid has almost disappeared. In cases where the protection of the amino group is carried out in a solvent containing water, the compound of Formula (1) generated by lactonization, if any, can be hydrolyzed by a method such as making the pH of the reaction mixture strongly basic, increasing the temperature of the reaction mixture to a temperature of not less than room temperature, performing the reaction for long time, or a combination of one or more of these methods, and, as a result, the amino-protected compound can be obtained. By subsequently reacting the amino-protected compound with an acid catalyst, the compound of Formula (1) can be obtained.

[0035]

The compound of Formula (1) produced can be isolated by a method such as the crystallization described below.

[0036]

In another mode of the method of the present invention, *cis*-5-hydroxy-2-piperidinecarboxylic acid is reacted with an acid halide and/or acid anhydride, and then with an alcohol in the presence of an acid catalyst, to convert the *cis*-5-hydroxy-2-piperidinecarboxylic acid into a compound of Formula (2) (the route 3 in Fig. 1). By the reaction with an alcohol in the presence of an acid catalyst, the compound of Formula (1) produced by the reaction with an acid halide and/or acid anhydride is converted to the compound of Formula (2). In this process, in cases where compounds having a carboxyl group other than the compound of Formula (1), especially the 2-piperidinecarboxylic acid or an analogue thereof described later, are contained as impurities, these compounds are not converted to lactone like the compound of Formula (1). Therefore, the compound derived from *cis*-5-hydroxy-2-piperidinecarboxylic acid can be selectively converted to the compound of Formula (2), which is an ester, while the other compounds are left in the reaction system as

carboxylic acids. Thereafter, by extraction with a basic aqueous solution, the compound of Formula (2) can be separated into the organic layer, while the impurities other than the compound of Formula (2) can be separated into the aqueous layer.

5 [0037]

The alcohol used herein is not limited as long as it is a C₁-C₆ alcohol. Examples of the alcohol include primary alcohols such as methanol, ethanol, 1-propanol, 1-butanol, 1-pentanol, and 1-hexanol; secondary alcohols such as 2-propanol, 2-butanol, cyclopentanol, and cyclohexanol; and tertiary alcohols such as
10 *tert*-butanol. Among these, secondary alcohols are preferred, and 2-propanol is more preferred since it is inexpensive, has a low boiling point, and can be easily removed. In cases where a primary alcohol is used, a side reaction may occur to cause direct conversion of a carboxylic acid into its ester in the presence of an acid catalyst, resulting in low effect of removal of impurities. In cases where a tertiary
15 alcohol is used, reactivity with the compound of Formula (1) is low. Therefore, the compound of Formula (2), which is a secondary alcohol, may react with the compound of Formula (1) to partially form a dimer, resulting in low yield of the compound of Formula (2). Thus, in the present invention, a secondary alcohol is preferably used.

20 An ester of the above-described alcohol may be added in the presence of an acid catalyst and water so that hydrolysis reaction of the ester allows the alcohol to be present in the reaction system, and the alcohol may be reacted with the *cis*-*N*-protected-5-hydroxy-2-piperidinecarboxylic acid in the presence of an acid catalyst.

The alcohol, or the ester which generates the alcohol, may be used in an
25 excess amount, and the molar amount of the alcohol or ester to be used is usually 1 to 500 times the molar amount of the *cis*-5-hydroxy-2-piperidinecarboxylic acid. In cases where the amount of the alcohol or ester is too small, a side reaction in which

the compound of Formula (2) reacts with the compound of Formula (1) may occur to form a dimer, while in cases where the amount of the alcohol or ester is too large, the effect of the acid catalyst decreases. Thus, the molar amount of the alcohol or ester is preferably 2 to 100 times, more preferably 3 to 50 times, especially preferably 5 to 20 times the molar amount of the *cis*-5-hydroxy-2-piperidinecarboxylic acid

[0038]

Examples of the acid catalyst include inorganic acids such as hydrochloric acid, sulfuric acid, nitric acid, phosphoric acid, and polyphosphoric acid; sulfonic acids such as *p*-toluenesulfonic acid, methanesulfonic acid, and trifluoromethanesulfonic acid; and carboxylic acids such as acetic acid, formic acid, trifluoroacetic acid, trichloroacetic acid, benzoic acid, and oxalic acid. Among these, sulfonic acids are preferred since they have high solubility in organic solvents and sufficiently strong acidity for allowing the reaction to proceed. *p*-Toluenesulfonic acid and methanesulfonic acid are more preferred since they are industrially inexpensive.

The molar amount of the acid catalyst to be added is usually 0.001 to 10 times the molar amount of the *cis*-5-hydroxy-2-piperidinecarboxylic acid. In cases where the molar amount of the acid catalyst is too small, the reaction proceeds slowly, while the molar amount of the acid catalyst is too large, the load of the post treatment is large. The molar amount of the acid catalyst is therefore preferably 0.01 to 5 times, more preferably 0.02 to 1 times, especially preferably 0.05 to 0.5 times the molar amount of the *cis*-5-hydroxy-2-piperidinecarboxylic acid.

[0039]

Examples of the reaction solvent include organic acids such as formic acid and acetic acid; esters such as ethyl acetate and butyl acetate; alcohols such as methanol, ethanol, 1-propanol, 2-propanol, 1-butanol, 2-ethyl-1-hexanol, and 2-butanol; ethers such as diethyl ether, di-*n*-butyl ether, diisopropyl ether, di-*n*-butyl

ether, *tert*-butyl methyl ether, tetrahydrofuran, and dioxane; ketones such as acetone, methyl ethyl ketone, and methyl isobutyl ketone; nitriles such as acetonitrile; amides such as dimethylformamide, dimethylacetamide, and *N*-methylpyrrolidinone; sulfones such as dimethylsulfoxide and sulfolane; hydrocarbons such as hexane, heptane, and toluene; and mixed solvents of two or more of these. Among these, solvents which do not show coordination of a proton are preferred since they do not weaken the activity of the acid catalyst. More specifically, hydrocarbons are preferred. In addition, organic acids, which themselves are acids, and esters, which produce acids by hydrolysis, are preferred. Toluene is more preferred because of its low cost, high reaction rate, and high solubility of acid catalysts therein.

[0040]

The reaction temperature is usually 0°C to 150°C, preferably 20°C to 120°C. Reaction at higher temperature may cause a side reaction, but reaction at lower temperature is slow and takes a long time. The reaction temperature is therefore more preferably 40°C to 80°C.

[0041]

The compound of Formula (2) generated can be isolated by a method such as the extraction method using an organic solvent described later.

[0042]

The compounds of Formulae (1) and (2) can be converted to *cis-N*-protected-5-hydroxy-2-piperidinecarboxylic acid by treatment with a base.

[0043]

Examples of the base to be used herein include hydroxides such as sodium hydroxide, potassium hydroxide, lithium hydroxide, and calcium hydroxide; phosphates such as sodium phosphate, potassium phosphate, and calcium phosphate; carbonates such as lithium carbonate, sodium carbonate, potassium carbonate, magnesium carbonate, and calcium carbonate; and hydrogen carbonates such as

sodium hydrogen carbonate and potassium hydrogen carbonate. The base is preferably a hydroxide.

The molar amount of the base to be used is usually 0.1 to 10 times, preferably 0.5 to 5 times, more preferably 0.9 to 3 times the total molar amount of the compounds of Formulae (1) and (2).

[0044]

By carrying out deprotection of the amino group of the *cis*-*N*-protected-5-hydroxy-2-piperidinecarboxylic acid, *cis*-5-hydroxy-2-piperidinecarboxylic acid (*cis*-5-hydroxypipericolic acid) can be produced. In particular, in cases where R¹ in the Formulae (1) and (2) is a benzyloxycarbonyl group, deprotection of the amino group can be carried out by hydrogenation reaction. The hydrogenation reaction can be carried out by, for example, using a palladium carbon catalyst. By such a reaction, the *cis*-5-hydroxy-2-piperidinecarboxylic acid can be regenerated.

[0045]

The method for purifying *cis*-5-hydroxy-2-piperidinecarboxylic acid of the present invention is a method in which highly pure *cis*-5-hydroxy-2-piperidinecarboxylic acid is obtained by purification from *cis*-5-hydroxy-2-piperidinecarboxylic acid containing an impurity(s).

[0046]

Examples of the impurity include 2-piperidinecarboxylic acid or an analogue thereof, amino acids, peptides, sugars, and fatty acids. Since, among these, 2-piperidinecarboxylic acid or an analogue thereof, amino acids, and peptides have an amino group(s) and carboxyl group(s), these can be hardly separated from the desired *cis*-5-hydroxy-2-piperidinecarboxylic acid of interest by an ordinary purification method such as ion-exchange resin purification. In particular, since 2-piperidinecarboxylic acid or an analogue thereof has the same skeleton as *cis*-5-hydroxy-2-piperidinecarboxylic acid, its separation is very difficult, and

establishment of removal process is very important.

Examples of the 2-piperidinecarboxylic acid or an analogue thereof include isomers of *cis*-5-hydroxy-2-piperidinecarboxylic acid; and 2-piperidinecarboxylic acid and analogues thereof. Examples of the isomers of *cis*-5-hydroxy-2-
 5 piperidinecarboxylic acid include stereoisomers such as *trans* isomers, and structural isomers having hydroxyl groups at different positions. It should be noted that optical isomers of *cis*-5-hydroxy-2-piperidinecarboxylic acid are not included in the impurity of the present invention.

Examples of the *trans* isomers include *trans*-5-hydroxy-2-
 10 piperidinecarboxylic acid, that is, (2*S*,5*R*)-5-hydroxy-2-piperidinecarboxylic acid, (2*R*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid, and mixtures of these. Examples of the structural isomers include *cis*-2-hydroxy-2-piperidinecarboxylic acid, *trans*-2-hydroxy-2-piperidinecarboxylic acid, *cis*-3-hydroxy-2-piperidinecarboxylic acid, *trans*-3-hydroxy-2-piperidinecarboxylic acid, and *trans*-4-hydroxy-2-
 15 piperidinecarboxylic acid. It should be noted that *cis*-4-hydroxy-2-piperidinecarboxylic acid is not included in the impurity of the present invention.

Examples of the 2-piperidinecarboxylic acid and analogues thereof include 2-piperidinecarboxylic acid and 1,2,3,4-tetrahydro-2-pyridinecarboxylic acid.

Examples of the amino acids include essential amino acids such as proline,
 20 lysine, and isoleucine, and unnatural amino acids such as 3-hydroxyproline, 4-hydroxyproline, and 5-hydroxylysine.

Examples of the peptides include dipeptides, oligopeptides, and proteins.

Examples of the sugars include glucose, gluconic acid, sodium gluconate, potassium gluconate, and calcium gluconate.

25 Examples of the fatty acids include C₂-C₄ short-chain fatty acids, C₅-C₁₂ medium-chain fatty acids, long-chain fatty acids having not less than 13 carbon atoms, and glycerol esters.

[0047]

A mixture containing *cis*-5-hydroxy-2-piperidinecarboxylic acid and an impurity(s) is reacted with an acid halide and/or acid anhydride, or with an acid halide and/or acid anhydride and an alcohol, to convert the *cis*-5-hydroxy-2-piperidinecarboxylic acid into a compound(s) of Formula (1) and/or Formula (2).
5 Subsequently, the compound(s) is/are separated by crystallization, solvent extraction, and/or the like and then treated with a base to convert the compound(s) into *cis*-*N*-protected-5-hydroxy-2-piperidinecarboxylic acid. By subsequently carrying out deprotection of the amino group, *cis*-5-hydroxy-2-piperidinecarboxylic acid which
10 the impurity has been removed can be obtained.

That is, *cis*-5-hydroxy-2-piperidinecarboxylic acid can be separated from the impurity utilizing specific lactonization of the *cis*-5-hydroxy-2-piperidinecarboxylic acid. Even in cases where the impurity other than the above-described impurities is contained, the *cis*-5-hydroxy-2-piperidinecarboxylic acid can be separated also from
15 the impurity.

In the present description, the crystallization includes not only ordinary crystallization by adding a poor solvent, acid, base, or the like to a solution, or azeotropically removing a good solvent such as water, to decrease solubility of the desired product and recover the product as crystals; but also recrystallization in
20 which crude crystals that have once been obtained are dissolved in an appropriate solvent and then recrystallized. The crystallization may be promoted by addition of seed crystals to the solution.

In the crystallization of the compound of Formula (1), crystallization is preferably carried out for a crude product of the compound of Formula (1) obtained
25 by the reaction with the acid halide and/or acid anhydride. This is because a small amount of acidic component(s) generated from the acid halide and/or acid anhydride, and/or other components derived from reagents, may remain to induce crystallization.

[0048]

Examples of the crystallization solvent for the crystallization treatment include water; organic acids such as acetic acid and propionic acid; esters such as ethyl acetate, isopropyl acetate, and butyl acetate; alcohols such as methanol, ethanol, 5 1-propanol, 2-propanol, 1-butanol, 2-ethyl-1-hexanol, and 2-butanol; ethers such as diethyl ether, di-*n*-butyl ether, diisopropyl ether, methyl-*tert*-butyl ether, di-*n*-butyl ether, tetrahydrofuran, and dioxane; ketones such as acetone, methyl ethyl ketone, and methyl isobutyl ketone; nitriles such as acetonitrile; aliphatic hydrocarbons such as hexane and heptane; aromatic hydrocarbons such as toluene and xylene; and 10 mixed solvents of two or more of these. Among these, alcohols, aliphatic hydrocarbons, aromatic hydrocarbons, and mixed solvents of two or more of these are preferred since solubility of the compound(s) of Formula (1) and/or Formula (2) in these solvents is sufficiently low. Aliphatic hydrocarbons, aromatic hydrocarbons, and mixed solvents of two or more of these are more preferred, and 15 hexane, heptane, toluene, and mixed solvents of two or more of these are especially preferred.

[0049]

Examples of the organic solvent to be used for the extraction include water-insoluble solvents, for example, esters such as ethyl acetate, isopropyl acetate, and 20 butyl acetate; ethers such as diethyl ether, methyl-*tert*-butyl ether, di-*n*-butyl ether, diisopropyl ether, di-*n*-butyl ether, *tert*-butyl methyl ether, and tetrahydrofuran; aliphatic hydrocarbons such as hexane and heptane; and aromatic hydrocarbons such as toluene and xylene.

[0050]

25 The above-described method is especially useful in cases where *cis*-5-hydroxy-2-piperidinecarboxylic acid is to be purified from a mixture of *cis*-5-hydroxy-2-piperidinecarboxylic acid and *trans*-5-hydroxy-2-piperidinecarboxylic


acid.

[0051]

That is, in a mode of the purification method of the present invention (the route 1 or route 2 in Fig. 1), a mixture of *cis*-5-hydroxy-2-piperidinecarboxylic acid and *trans*-5-hydroxy-2-piperidinecarboxylic acid is reacted with an acid halide and/or
5 acid anhydride to convert the *cis*-5-hydroxy-2-piperidinecarboxylic acid into the compound of Formula (1). The resulting compound is then separated by crystallization and/or extraction with an organic solvent, and then treated with a base to convert the compound into *cis*-*N*-protected-5-hydroxy-2-piperidinecarboxylic acid.
10 By subsequently carrying out deprotection of the amino group, *cis*-5-hydroxy-2-piperidinecarboxylic acid separated from the *trans* isomer can be obtained.

In such cases, the acid halide and/or acid anhydride is/are preferably used in an excessive amount over 1 molar equivalent relative to the total amount of the *cis*-5-hydroxy-2-piperidinecarboxylic acid and the *trans*-5-hydroxy-2-piperidinecarboxylic
15 acid. The amount of the acid halide and/or acid anhydride is usually 1 to 10 molar equivalents, preferably 1.2 to 5 molar equivalents, more preferably 1.5 to 3 molar equivalents. In cases where both an acid halide and an acid anhydride are used, these are added such that their total amount exceeds 1 molar equivalent relative to the total amount of the *cis*-5-hydroxy-2-piperidinecarboxylic acid and the *trans*-5-
20 hydroxy-2-piperidinecarboxylic acid.

By the reaction with the acid halide and/or acid anhydride, the *cis*-5-hydroxy-2-piperidinecarboxylic acid is converted to the compound of Formula (1). On the other hand, the *trans*-5-hydroxy-2-piperidinecarboxylic acid remains in the reaction system without being converted, or is converted to the mixed acid anhydride of *trans*-
25 5-hydroxy-2-piperidinecarboxylic acid described below. This mixed acid anhydride can be converted into the *trans*-5-hydroxy-2-piperidinecarboxylic acid again by post treatment. Thus, its separation by crystallization reaction is possible.




[0052]

In another mode of the purification method of the present invention (the route 3 in Fig. 1), a mixture of *cis*-5-hydroxy-2-piperidinecarboxylic acid and *trans*-5-hydroxy-2-piperidinecarboxylic acid is reacted with an acid halide and/or acid anhydride to protect the amino group. Subsequently, the *cis*-*N*-protected-5-hydroxy-2-piperidinecarboxylic acid generated is reacted with an alcohol (or an ester that generates an alcohol) in the presence of an acid catalyst to allow conversion into the compound of Formula (2) via the compound of Formula (1). The resulting compound is separated by extraction with an organic solvent, and then treated with a base to allow conversion into *cis*-*N*-protected-5-hydroxy-2-piperidinecarboxylic acid. By subsequently carrying out deprotection of the amino group, *cis*-5-hydroxy-2-piperidinecarboxylic acid separated from the *trans* isomer can be obtained.

In such cases, the acid halide and/or acid anhydride is/are preferably used in an amount of about 1 molar equivalent relative to the total amount of the *cis*-5-hydroxy-2-piperidinecarboxylic acid and the *trans*-5-hydroxy-2-piperidinecarboxylic acid.

By performing the reaction with the acid halide and/or acid anhydride to protect the amino group, and then with the alcohol (or ester that generates an alcohol) in the presence of the acid catalyst, the *cis*-5-hydroxy-2-piperidinecarboxylic acid is converted into the compound of Formula (2), while the *trans*-5-hydroxy-2-piperidinecarboxylic acid is converted to the *trans*-*N*-protected-5-hydroxy-2-piperidinecarboxylic acid described below. Thus, their separation is possible by extraction with an organic solvent.



[0053]

The “*cis*-5-hydroxy-2-piperidinecarboxylic acid containing an impurity(s)” and “mixture of *cis*-5-hydroxy-2-piperidinecarboxylic acid and *trans*-5-hydroxy-2-piperidinecarboxylic acid” may be those synthesized by bacterial reaction and/or enzymatic reaction. For example, by the method described in Example 29 of JP 4590981 B, a “mixture of *cis*-5-hydroxy-2-piperidinecarboxylic acid and *trans*-5-hydroxy-2-piperidinecarboxylic acid” can be obtained. The “*cis*-5-hydroxy-2-piperidinecarboxylic acid containing an impurity(s)” and “mixture of *cis*-5-hydroxy-2-piperidinecarboxylic acid and *trans*-5-hydroxy-2-piperidinecarboxylic acid” may also be products recovered after separation of a *cis*-5-hydroxy-2-piperidinecarboxylic acid derivative by the method of the present invention.

[0054]

In cases where *cis*-5-hydroxy-2-piperidinecarboxylic acid is purified by the method of the present invention from the “*cis*-5-hydroxy-2-piperidinecarboxylic acid containing an impurity(s)” or “mixture of *cis*-5-hydroxy-2-piperidinecarboxylic acid and *trans*-5-hydroxy-2-piperidinecarboxylic acid” synthesized by bacterial reaction and/or enzymatic reaction, the obtained *cis*-5-hydroxy-2-piperidinecarboxylic acid is preferably further subjected to adsorption purification using activated carbon and/or crystallization using a solvent containing water to remove impurities such as a colored substance, to further increase purity of the *cis*-5-hydroxy-2-piperidinecarboxylic acid.

[0055]

As the activated carbon, an arbitrary known activated carbon may be used, and examples of the known activated carbon include coal-based, wood-based,

coconut shell-based, and resin-based activated carbons. The activated carbon may also be prepared by activating a material activated carbon such as a coal-based, wood-based, coconut shell-based, or resin-based material activated carbon by a method such as gas activation, steam activation, or chemical activation using zinc chloride, phosphoric acid, or the like.

Specific examples of the activated carbon include Calgon CPG, Calgon CAL, Calgon SGL, Diasorb W, Diahope MS10, Diahope M010, Diahope MS16, Diahope 6MD, Diahope 6MW, Diahope 8ED, Diahope ZGN4, and CENTUR, manufactured by Calgon Mitsubishi Chemical Corporation; GAC, GAC PLUS, GCN PLUS, C GRAN, RO, ROX, DARCO, CN, SX, SX PLUS, SA, SX, PK, and W, manufactured by Norit Japan Co., Ltd.; GW, GWH, GLC, 4GC, KW, PW, and PK, manufactured by Kuraray Chemical Co., Ltd.; HC-30S, GL-30S, 4G-3S, PS, and PC, manufactured by Tsurumicoal Co., Ltd.; P, W, CW, SG, SGP, S, GB, CA, and K, manufactured by Futamura Chemical Co., Ltd.; Shirasagi KL, Shirasagi W2C, Shirasagi WH2C, Shirasagi W5C, Shirasagi WH5C, Shirasagi WH5X, Shirasagi XS7100H-3, Carboraffin, Shirasagi A, Shirasagi C, and Shirasagi M, manufactured by Japan EnviroChemicals Ltd.; and Hokuetsu CL-K, Hokuetsu HS, and Hokuetsu KS, manufactured by Ajinomoto Fine-Techno Co., Inc.

By this operation, impurities showing absorption within the wavelength range of 400 nm to 800 nm can be removed.

[0056]

As the solvent containing water to be used for the crystallization, a water-soluble organic solvent can be used. Specific examples of the solvent include organic acids such as acetic acid and propionic acid; esters such as ethyl acetate; alcohols such as methanol, ethanol, 1-propanol, 2-propanol, and 1-butanol; ethers such as tetrahydrofuran and dioxane; ketones such as acetone, methyl ethyl ketone, and diethyl ketone; nitriles such as acetonitrile; amides such as dimethylformamide,

dimethylacetamide, and *N*-methylpyrrolidinone; sulfones such as dimethylsulfoxide and sulfolane; and mixed solvents of two or more of these. It is also preferred to further increase purity of the *cis*-5-hydroxy-2-piperidinecarboxylic acid by carrying out crystallization of the *cis*-5-hydroxy-2-piperidinecarboxylic acid using the above-described solvent. Among these, organic acids, alcohols, and ketones, whose solubility in water is higher, are preferred. Alcohols and ketones are more preferred, and ethanol and acetone are especially preferred.

EXAMPLES

10 [0057]

The present invention is described below in more detail by way of Examples. However, the present invention is not limited by these Examples.

[0058]

The quantitative analyses in the Examples were carried out by HPLC (High Performans Liquid Chromatography) under the following conditions.

<HPLC-1>

Column: Astec CLC-D (4.6 mm × 150 mm, 5 μm), manufactured by
SUPELCO

Mobile phase: 2 mmol/L aqueous copper sulfate solution

20 Flow rate: 1.0 mL/minute

Column temperature: 45°C

Detection wavelength: UV 254 nm

<HPLC-2>

Column: L-column (4.6 mm × 250 mm, 5 μm), manufactured by Chemicals
25 Evaluation and Research Institute, Japan

Mobile phase:

A: 0.1 wt% aqueous trifluoroacetic acid solution

B: methanol

gradient (concentration of B): 0 minute, 20% → 2 minute, 20% → 10 minute, 80% → 20 minute, 80%

Flow rate: 1.0 mL/minute

5 Column temperature: 40°C

Detection wavelength: UV 200 nm

<HPLC-3>

Column: ZORBAX Eclipse Plus C18 (4.6 mm × 150 mm, 1.8 μm),
manufactured by Agilent Technologies

10 Mobile phase:

A: 0.1 wt% aqueous phosphoric acid solution

B: acetonitrile

gradient (concentration of B): 0 minute, 55% → 6 minute, 55% → 9 minute, 80% → 12 minute, 80%

15 Flow rate: 1.0 mL/minute

Column temperature: 40°C

Detection wavelength: UV 210 nm

<HPLC-4>

Column: COSMOSIL 5C₁₈-AR-II (4.6 mm × 150 mm), manufactured by
20 Nacalai Tesque, Inc.

Mobile phase: 50 mmol/L phosphate buffer (pH 2.7)

Flow rate: 1.0 mL/minute

Column temperature: 40°C

Detection wavelength: UV 340 nm

25 <HPLC-5>

Column: SUMICHIRAL OA-6100 (4.6 mm × 250 mm), manufactured by
Sumika Chemical Analysis Service, Ltd.

Mobile phase: 1 mmol/L copper sulfate

Flow rate: 1.0 mL/minute

Column temperature: 30°C

Detection wavelength: UV 254 nm

5 <HPLC-6>

Column: CLC-D (4.6 mm × 150 mm), manufactured by SUPELCO

Mobile phase: 2 mmol/L copper sulfate

Flow rate: 1.0 mL/minute

Column temperature: 30°C

10 Detection wavelength: UV 254 nm

[0059]

[Reference Example 1]

[Preparation Example of Recombinant *Escherichia coli* (*E. coli*) JM109/pKW32

15 (dpkA, aip, gdh, kr), in Which *N*-Methyl-L-amino Acid Dehydrogenase (Hereinafter Referred to as DpkA), L-Amino Acid Oxidase (Hereinafter Referred to as AIP), Glucose-1-dehydrogenase (Hereinafter Referred to as GDH), and Amino Acid Racemase (hereinafter referred to as KR) Are Co-expressed]

(1) Cloning of Genes

20 Based on a gene sequence of dpkA (SEQ ID NO:1) encoding DpkA (GenBank Accession No. BAD89743, SEQ ID NO:2) derived from *Pseudomonas putida* (*P. putida*), primers for amplifying the full-length sequence of the dpkA gene, dpkA_F (SEQ ID NO:9) and dpkA_R (SEQ ID NO:10), were designed and synthesized. PCR was carried out using chromosomal DNA of *P. putida* as a
25 template according to a conventional method, to obtain a DNA fragment of about 1.0 kbp.

A gene sequence of aip (SEQ ID NO:3) encoding a protein AIP (SEQ ID

NO:4), which has the amino acid sequence of L-amino acid oxidase derived from *Scomber japonicus* (GenBank Accession No.CAC00499) with the exception that the signal peptide is removed and methionine is added, was designed and artificially synthesized. Primers for amplifying the full-length sequence of the aip gene, aip_F (SEQ ID NO:11) and aip_R (SEQ ID NO:12), were designed and synthesized. PCR was carried out according to a conventional method, to obtain a DNA fragment of about 1.5 kbp.

Based on a gene sequence of gdh (SEQ ID NO:5) encoding a protein (SEQ ID NO:6) having the amino acid sequence of GDH derived from *Bacillus subtilis* (GenBank Accession No.NP_388275) with the exception that the 96th amino acid residue, glutamic acid, is substituted with alanine, primers for amplifying the full-length sequence of the gdh gene, gdh_F (SEQ ID NO:13) and gdh_R(SEQ ID NO:14), were designed and synthesized. PCR was carried out according to a conventional method, to obtain a DNA fragment of about 0.8 kbp.

Based on a gene sequence of kr (SEQ ID NO:7) encoding KR derived from *P. putida* (GenBank Accession No. NP_745855, SEQ ID NO:8), primers for amplifying the full-length sequence of the rk gene, kr_F (SEQ ID NO:15) and kr_R (SEQ ID NO:16), were designed and synthesized. PCR was carried out using chromosomal DNA of *P. putida* as a template according to a conventional method, to obtain a DNA fragment of about 1.2 kbp.

[0060]

(2) Preparation of Expression Plasmid

Each of the DNA fragments obtained in (1) was digested with restriction enzymes *Eco*RI and *Xba*I, and introduced downstream of the trc promoter in a *Mun*I/*Xba*I digest of a plasmid pKW32, which is described in WO 2012/029819, using a Ligation-Convenience Kit (manufactured by Nippon Gene Co., Ltd.), to obtain pKW32dpkA, pKW32aip, pKW32gdh, and pKW32kr, respectively.

Subsequently, pKW32aip was digested with *SpeI* and *NdeI* to obtain a DNA fragment of about 2.4 kbp containing *aip*, and the resulting DNA fragment was introduced downstream of *dpkA* in the ring-opened (linear) plasmid of about 4.2 kbp obtained by digesting pKW32dpkA with *XbaI* and *NdeI*, to obtain pKW32 (*dpkA*, *aip*).

pKW32gdh was digested with *SpeI* and *NdeI* to obtain a DNA fragment of about 1.7 kbp containing *gdh*, and the resulting DNA fragment was introduced downstream of *aip* in the ring-opened (linear) plasmid of about 5.7 kbp obtained by digesting pKW32 (*dpkA*, *aip*) with *XbaI* and *NdeI*, to obtain pKW32 (*dpkA*, *aip*, *gdh*).

Finally, pKW32kr was digested with *SpeI* and *NdeI* to obtain a DNA fragment of about 2.1 kbp containing *kr*, and the resulting DNA fragment was introduced downstream of *gdh* in the ring-opened (linear) plasmid of about 6.5 kbp obtained by digesting pKW32 (*dpkA*, *aip*, *gdh*) with *XbaI* and *NdeI*, to obtain pKW32 (*dpkA*, *aip*, *gdh*, *kr*).

[0061]

(3) Preparation of Expressing Strain

Using the plasmid pKW32 (*dpkA*, *aip*, *gdh*, *kr*) obtained in (2), *E. coli* JM109 (manufactured by Takara Bio Inc.) was transformed according to a conventional method, to obtain recombinant *E. coli* JM109/pKW32 (*dpkA*, *aip*, *gdh*, *kr*).

[0062]

[Reference Example 2]

[Preparation Example of 5-Hydroxy-2-piperidinecarboxylic Acid Reaction Mixture]

To a 1-L jar fermenter (manufactured by ABLE Corporation, type BMJ), 45 g of 5-hydroxylysine hydrochloride (230 mmol, prepared according to a method described in Bull. Chem. Soc. Jpn., 1962, 35, 2006), 2.27 mL of Adekanol LG-109,

and 67.34 g (374 mmol) of glucose were added, and dissolved in water. Thereafter, 20 wt% aqueous sodium hydroxide solution was added dropwise thereto until the pH became 8. To the resulting liquid, bovine-liver-derived catalase (manufactured by Wako Pure Chemical Industries, Ltd.) was added at 2000 U/L; NADP⁺ (manufactured by Oriental Yeast Co., Ltd.) was added at 0.2 mmol/L; and wet cells of the recombinant *E. coli* JM109/pKW32 (dpkA, aip, gdh, kr) prepared in Reference Example 1 were added at 25 g/L; followed by adding water to the resulting mixture to a liquid volume of 566 mL. The reaction was allowed to proceed at 30°C, a stirring rate of 500 rpm, and an aeration rate of about 1 L/min for 43 hours. During the reaction, the pH was kept at 8 by adding 20 wt% aqueous sodium hydroxide solution dropwise. Exactly the same operations were carried out twice thereafter, and the reaction mixtures obtained by the 3 rounds of operations (using 135 g of 5-hydroxylysine hydrochloride, 690 mmol) were combined. The pH of the reaction mixture was adjusted to 2.5 using 6 mol/L sulfuric acid. Thereafter, 20 wt% aqueous sodium hydroxide solution was added dropwise to adjust the pH to 4, and the resulting liquid was centrifuged at 10,000 rpm for 20 minutes to remove insoluble impurities. The obtained supernatant was passed through a microfiltration membrane (manufactured by Asahi Kasei Corporation), and then through an ultrafiltration membrane (manufactured by Asahi Kasei Corporation), to remove impurities. Analysis of 5-hydroxy-2-piperidinecarboxylic acid contained in the resulting solution was carried out by HPLC under the conditions of <HPLC-1>. As a result, the amount of (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid produced was 43.9g (302 mmol; yield, 44%), and the amount of (2*S*,5*R*)-5-hydroxy-2-piperidinecarboxylic acid produced was 29.7g (205 mmol; yield, 30%).

Example 1

[0063]

<1-1> Production of Benzyl (1*S*,4*S*)-5-Aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate

To 1153 g of a reaction mixture containing 20.7 g (143 mmol) of (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid and 18.0 g (124 mmol) of (2*S*,5*R*)-5-hydroxy-2-piperidinecarboxylic acid [(2*S*,5*S*):(2*S*,5*R*) = 53.6:46.4 (molar ratio)] obtained according to the method of Reference Example 2, 10 mol/L aqueous sodium hydroxide solution (60 mL) was added to adjust the pH from 3.65 to 10.81. The resulting liquid was concentrated under reduced pressure, to obtain 422.4 g of a slurry solution. In each of 2 flasks, 211.2 g of the obtained slurry was placed, and the internal temperature was decreased to 5°C to 7°C, followed by carrying out the following operations for each flask.

[The following is description of the operations for one of the flasks. Each flask contained 10.4 g (72 mmol) of (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid and 9.0 g (62 mmol) of (2*S*,5*R*)-5-hydroxy-2-piperidinecarboxylic acid.]

[0064]

To the flask, 18.8 mL (133 mmol) of benzyloxycarbonyl chloride was added dropwise, and 13.3 mL (133 mmol) of 10 mol/L aqueous sodium hydroxide solution was then added to adjust the pH from about 9 to 10, followed by allowing the reaction to proceed at a temperature within the range of 10°C to 20°C for 10 minutes. To the resulting liquid, 18.8 mL (133 mmol) of benzyloxycarbonyl chloride was added dropwise, and 13.5 mL (135 mmol) of 10 mol/L aqueous sodium hydroxide solution was then added to adjust the pH from 9 to 11, followed by allowing the reaction to proceed at a temperature within the range of 10°C to 20°C for 3 hours. To the resulting liquid, 19 mL of water was added, and 6.2 mL (44 mmol) of benzyloxycarbonyl chloride was then added dropwise, followed by allowing the liquid to gradually warm to room temperature. To the resulting liquid, 200 mL of toluene was added, and 4 mL of 10 mol/L aqueous sodium hydroxide solution (40

mmol) was then added to adjust the pH to 11, followed by separating the organic layer. To the aqueous layer, 100 mL of toluene was added to perform re-extraction, and the aqueous layer was separated to obtain the organic layer. This organic layer was combined with the previously obtained organic layer. The resulting mixture
 5 was washed with 10 mL of water, followed by separating the resulting organic layer. From the organic layer obtained from each of the two flasks, the solvent was removed by distillation under reduced pressure at 35°C, to obtain 64.9 g of a yellow oily substance.

As a result of ¹H-NMR analysis, this oily substance was found to be a mixture
 10 containing 45 wt% benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate (111 mmol; yield, 78% with respect to the (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid), 15 wt% (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid=benzyloxyformic acid=anhydride (23 mmol; yield, 19% with respect to the (2*S*,5*R*)-5-hydroxy-2-piperidinecarboxylic acid), 36 wt% benzyl
 15 alcohol, and 5 wt% toluene. Thus, the conversion of the material 5-hydroxy-2-piperidinecarboxylic acid, which contained (2*S*,5*S*) and (2*S*:5*R*) at a ratio of (2*S*,5*S*): (2*S*:5*R*) = 53.6:46.4 (molar ratio), to benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate allowed effective removal of compounds having the stereochemistry of (2*S*,5*R*), and, by this, the purity of compounds having
 20 the stereochemistry of (2*S*,5*S*) could be increased to (2*S*,5*S*): (2*S*:5*R*) = 82.8:17.2 (molar ratio).

[0065]

The resulting crude product was cooled to 10°C to 15°C, and 36 mL of toluene and 36 mL of hexane were added thereto. As a result, precipitation of white
 25 solids occurred. The precipitated white solids were collected by filtration, and washed by sprinkling 100 mL of hexane thereon. The resulting white solids were dried under reduced pressure at room temperature, to obtain 20.7 g of white solids.

As a result of ^1H -NMR analysis, the white solids were found to be a mixture containing 92.5 wt% benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate (yield, 51% with respect to the (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid) and 7.5 wt% (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid=benzyloxyformic acid=anhydride. By the crystallization, the purity of compounds having the stereochemistry of (2*S*,5*S*) could be increased to (2*S*,5*S*):(2*S*,5*R*) = 92.5:7.5 (molar ratio).

^1H -NMR (400 MHz, CDCl_3)

δ 1.77-1.85 (1H, m), 2.02-2.25 (3H, m), 3.52 (1H, d, $J=11.9$ Hz), 3.70 (1H, dt, $J=12.2, 3.3$ Hz), 4.70-4.86 (2H, m), 5.12-5.21 (2H, m), 7.32-7.40 (5H, m).

[0066]

<1-1'> Production of Benzyl (1*S*,4*S*)-5-Aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate

To 80 g of a reaction mixture containing 1.74 g (12.0 mmol) of (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid and 1.01 g (7.0 mmol) of (2*S*,5*R*)-5-hydroxy-2-piperidinecarboxylic acid [(2*S*,5*S*):(2*S*,5*R*)=63.2:36.8 (molar ratio)] obtained according to the method of Reference Example 2, 10 mol/L aqueous sodium hydroxide solution was added to adjust the pH from 3.65 to 9.8. The resulting liquid was concentrated under reduced pressure, and 38 g of a solution was obtained.

To the solution, 3.1 mL of 2-propanol was added. The internal temperature of the obtained solution was set to about 25°C. To the solution, 3.2 g (18.8 mmol) of benzyloxycarbonyl chloride was added dropwise, and 2.3 g (22.6 mmol) of 40 wt% aqueous sodium hydroxide solution was further added, followed by allowing the reaction to proceed at a temperature within the range of 20°C to 25°C for 30 minutes.

To the resulting liquid, 3.2 g (18.8 mmol) of benzyloxycarbonyl chloride was added dropwise, and 2.3 g (22.6 mmol) of 40 wt% aqueous sodium hydroxide solution was further added, followed by allowing the reaction to proceed at a temperature within

the range of 20°C to 25°C for 30 minutes. Thereafter, the operation of adding 1.6 g (9.4 mmol) of benzyloxycarbonyl chloride dropwise to the resulting liquid, adding 1.1 g (11.3 mmol) of 40 wt% aqueous sodium hydroxide solution thereto, and then allowing the reaction to proceed at a temperature within the range of 20°C to 25°C for 30 minutes was repeated 3 times. Subsequently, the reaction mixture was cooled to 0°C to 5°C, and crystals precipitated in the reaction system were collected by filtration. The obtained crystals were dried under reduced pressure at 40°C to 45°C to obtain 2 g of white crystals.

As a result of ¹H-NMR analysis, the obtained crystals were found to be benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate (80 wt%; yield, 56% with respect to the (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid).

[0067]

<1-1''> Production of Benzyl (1*S*,4*S*)-5-Aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate

To 160 g of a reaction mixture containing 3.26 g (22.5 mmol) of (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid and 2.24 g (15.4 mmol) of (2*S*,5*R*)-5-hydroxy-2-piperidinecarboxylic acid [(2*S*,5*S*): (2*S*,5*R*)=59.4:40.6 (molar ratio)] obtained according to the method of Reference Example 2, 10 mol/L aqueous sodium hydroxide solution was added to adjust the pH from 3.65 to 9.5. The resulting liquid was concentrated under reduced pressure, and 74 g of a solution was obtained. To the solution, 6.0 mL of 2-propanol was added. The internal temperature of the obtained solution was set to about 25°C. To the solution, 6.4 g (37.6 mmol) of benzyloxycarbonyl chloride was added dropwise, and 4.5 g (45.2 mmol) of 40 wt% aqueous sodium hydroxide solution was further added, followed by allowing the reaction to proceed at a temperature within the range of 20°C to 25°C for 30 minutes. To the resulting liquid, 6.4 g (37.6 mmol) of benzyloxycarbonyl chloride was added dropwise, and 4.5 g (45.2 mmol) of 40 wt% aqueous sodium hydroxide solution was

further added, followed by allowing the reaction to proceed at a temperature within the range of 20°C to 25°C for additional 30 minutes. Thereafter, 3.2 g (18.8 mmol) of benzyloxycarbonyl chloride was added dropwise to the reaction mixture, and 2.3 g (22.6 mmol) of 40 wt% aqueous sodium hydroxide solution was further added dropwise thereto, followed by allowing the reaction to proceed at a temperature within the range of 20°C to 25°C for 30 minutes. Thereafter, 1.9 g (11.1 mmol) of benzyloxycarbonyl chloride was added dropwise to the reaction mixture, and 1.4 g (14.0 mmol) of 40 wt% aqueous sodium hydroxide solution was added dropwise thereto, followed by allowing the reaction to proceed at a temperature within the range of 20°C to 25°C for 30 minutes. To the resulting reaction mixture, 71 mL of toluene was added, and liquid separation was carried out. To the extracted organic layer, 13 mL of water and 6.2 g of 40 wt% aqueous sodium hydroxide solution were added, and liquid separation was carried out again. The obtained aqueous layer, in an amount of 32 g, was divided into two aliquots, and subjected to the purification methods A and B.

[0068]

Purification Method A: Addition of Acetic Anhydride

To 16 g of the aqueous layer obtained in <1-1'>, 9.1 g (89 mmol) of acetic anhydride was added. The obtained crystals were collected by filtration, and dried under reduced pressure at 40°C to 45°C. The dried crystals appeared white, and their weight was 1.5 g.

The obtained crystals were analyzed by HPLC under the conditions of <HPLC-3>. As a result, the obtained crystals were found to be benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate (100 wt%; yield, 50% with respect to the (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid).

[0069]

Purification Method B: Extraction under Acidic Conditions

To 16 g of the aqueous layer obtained in <1-1'>, 30 mL of water was added, and 5 mol/L hydrochloric acid was added thereto to adjust the pH in the system to 1.07. Thereafter, 55 mL of ethyl acetate was added to the resulting mixture, and liquid separation was carried out. Subsequently, 55 mL of ethyl acetate was added to the obtained aqueous layer, and liquid separation was carried out again. The obtained organic layers were combined, and concentrated under reduced pressure until the weight decreased to half of the original weight. To the obtained organic layer, 2.9 g (28.1 mmol) of triethylamine and 2.9 g (28.1 mmol) of acetic anhydride were added at 20°C to 25°C. After stirring the resulting mixture for 30 minutes, 0.57 g (5.6 mmol) of triethylamine and 0.57 g (5.6 mmol) of acetic anhydride were added thereto at 20°C to 25°C. After stirring the resulting reaction mixture for 30 minutes, 20 g of saturated aqueous sodium hydrogen carbonate solution was added thereto. After liquid separation, 71 mL of toluene was added to the obtained aqueous layer, and the operation of extraction was carried out again.

The obtained organic layers were combined, and concentrated under reduced pressure. Cooling of the obtained concentrate to 0°C to 5°C caused precipitation of crystals. The obtained crystals were collected by filtration, and then dried under reduced pressure at 40°C to 45°C. The dried crystals were white, and their weight was 2.4 g. The obtained crystals were analyzed by HPLC under the conditions of <HPLC-3>. As a result, the obtained crystals were found to be benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate (98 wt%; yield, 70% with respect to the (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid).

[0070]

<1-2> Production of (2*S*,5*S*)-*N*-Benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic Acid

To a flask, 20.0 g (92.5 wt%, 71 mmol) of the benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate obtained in <1-1> and 60 mL of ethanol were

added, and 76.5 mL of 1 mol/L aqueous sodium hydroxide solution was added thereto at room temperature, followed by stirring the mixture for 1 hour. The solvent was removed under reduced pressure at 35°C to 40°C, to obtain 61.2 g of a yellow oily substance. To the resulting substance, 100 mL of ethyl acetate was added, and the organic layer was separated. To the obtained aqueous layer, 14 mL of 5 mol/L hydrochloric acid was added to adjust the pH to 1.2. The aqueous layer was subjected to extraction using 200 mL of ethyl acetate twice, and the obtained organic layer was washed with 2 mL of water. After separation of the organic layer, the solvent was removed under reduced pressure at 35°C to 40°C, to obtain 24.0 g of a pale yellow oily substance.

As a result of ¹H-NMR analysis, this pale yellow oily substance was found to be (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid (purity, 76 wt%; 65 mmol; yield, 92%). The peak for (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid was not found at all by ¹H-NMR.

¹H-NMR (400 MHz, CDCl₃, rotational isomer mixture)
 δ 1.27-1.39 (1H, m), 1.70-1.85 (1H, m), 1.98-2.06 (1H, m), 2.27-2.39 (1H, m), 2.80 (0.4H, t, *J*=11.8 Hz), 2.87 (0.6H, t, *J*=11.6 Hz), 3.61-3.72 (1H, m), 4.18-4.33 (1H, m), 4.81-4.88 (0.4H, m), 4.93-4.98 (0.6H, m), 5.12-5.21 (2H, m), 7.28-7.40 (5H, m).

[0071]

<1-3> Production of (2*S*,5*S*)-5-Hydroxy-2-piperidinecarboxylic Acid

To a flask, 18.0 g (64.5 mmol) of the (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid obtained in <1-2>, 90 mL of ethanol, and 2.5 g of 10% palladium carbon (manufactured by N. E. Chemcat Corporation, PE-type, 55.3%, containing water) were added, and hydrogenation was carried out under normal temperature and pressure. Three hours later, 22 mL of water was added thereto, and hydrogenation was carried out under normal temperature and pressure. Disappearance of the starting material could be confirmed 8.5 hours later. The

palladium carbon was removed by filtration, and 100 mL of water was sprinkled thereon for washing. From the obtained filtrate, the solvent was removed under reduced pressure at 45°C to 55°C, to obtain 11.6 g of a white slurry.

The obtained slurry was analyzed by HPLC under the conditions of <HPLC-1>. As a result, the slurry was found to contain 5-hydroxy-2-piperidinecarboxylic acid with a purity of 65 wt% in an amount of 64.5 mmol and yield of 81%. The isomer ratio was as follows: (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid:(2*S*,5*R*)-5-hydroxy-2-piperidinecarboxylic acid=98.8:1.2. The material used in <1-1> had a purity of as low as (2*S*,5*S*):(2*S*,5*R*)=53.7:46.3 (molar ratio), and the purity was increased to (2*S*,5*S*):(2*S*,5*R*)=98.8:1.2 (molar ratio) by the method of the present invention.

[0072]

<1-4> Activated Carbon Treatment, and Crystallization

To a flask, 0.38 g of the slurry obtained in <1-3> (purity, 65 wt%; 1.7 mmol), 2.5 mL of water, 0.025 g of an activated carbon manufactured by Calgon Mitsubishi Chemical Corporation, CAL, and 0.025 g of an activated carbon manufactured by Calgon Mitsubishi Chemical Corporation, 6ED, were added. The resulting mixture was stirred at room temperature for 1 hour, and the active carbons were then removed by filtration, followed by concentrating the aqueous solution to 0.5 mL. While the concentrate was stirred, 5.0 mL of ethanol was added dropwise thereto, and the resulting mixture was cooled to 5°C. As a result, crystals gradually precipitated. While the temperature was kept at 5°C, the stirring was continued for 1 hour, and 2.0 mL of acetone was added dropwise thereto at the constant temperature of 5°C, followed by stirring the resulting mixture for additional 0.5 hour. Thereafter, the precipitated crystals were collected by filtration, and 1.0 mL of an ice-cooled ethanol/acetone solution (=5/2 volume ratio) was sprinkled thereon for washing. The crystals were then dried under reduced pressure at 60°C, to obtain 0.25 g (purity,

98 wt%; yield, 99.2%) of crystals.

The obtained crystals were analyzed by HPLC under the conditions of <HPLC-1>. Based on the result, the isomer ratio of 5-hydroxy-2-piperidinecarboxylic acid contained in the obtained crystals was as follows: (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid:(2*S*,5*R*)-5-hydroxy-2-piperidinecarboxylic acid=99.5:0.5.

Example 2

[0073]

<2-1> Production of (2*S*,5*S*)-*N*-Benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic Acid and (2*S*,5*R*)-*N*-Benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic Acid

To 124 mL of a reaction mixture containing 0.723 g (4.98 mmol) of (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid and 0.830 g (5.72 mmol) of (2*S*,5*R*)-5-hydroxy-2-piperidinecarboxylic acid obtained according to the method of Reference Example 2 [(2*S*,5*S*): (2*S*,5*R*)=46.5:53.5 (molar ratio)], 2.88 mL (36 mmol) of 50 wt% aqueous sodium hydroxide solution was added, and the resulting mixture was concentrated at a bath temperature of 40°C under a reduced pressure of 50 hPa, to obtain 132 g of a solution. In a water bath, 3.1 mL (22 mmol) of benzyloxycarbonyl chloride and 0.44 mL (5.5 mmol) of 50 wt% aqueous sodium hydroxide solution were dividedly added to the obtained solution while the pH was adjusted to about 9. After leaving the resulting solution to stand overnight, 60 mL of ethyl acetate and 0.88 mL (11 mmol) of 50 wt% aqueous sodium hydroxide solution were added thereto, followed by carrying out filtration through Celite. The organic layer was separated, and the aqueous layer was washed with ethyl acetate. To the aqueous layer, 2.9 mL of concentrated hydrochloric acid (33 mmol) was added to adjust the pH to 3. Thereafter, extraction with ethyl acetate was carried out 3 times. The solvent was removed from the resulting organic layer under reduced pressure at 35°C,

to obtain 2.44 g of a yellow oily substance.

As a result of $^1\text{H-NMR}$ analysis, this oily substance was found to be a mixture containing 80 wt% (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid and (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid (6.91 mmol; yield, 65%; (2*S*,5*S*): (2*S*,5*R*)=4:6 (molar ratio)), 12 wt% ethyl acetate, and 8 wt% acetic acid.

[0074]

<2-2> Production of Benzyl (2*S*,5*S*)-3-Oxo-2-oxa-5-azabicyclo[2.2.2]octane-5-carboxylate

To a flask, 2.35 g of the (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid and (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid obtained in <2-1> [6.72 mmol, (2*S*,5*S*): (2*S*,5*R*)=4:6 (molar ratio)], 10 mL of toluene, and 116 mg (0.67 mmol) of *p*-toluenesulfonic acid monohydrate were added, and the reaction was allowed to proceed at 60°C for 2 hours. After cooling the resulting reaction product to room temperature, 30 mL of ethyl acetate, 5 mL of water, and 7 mL of 1 mol/L sodium hydroxide solution were added thereto. As a result, the pH of the aqueous layer became 9. After separating the aqueous layer, re-extraction was carried out with ethyl acetate. The obtained organic layer was washed with saturated aqueous sodium bicarbonate solution and saturated brine. The resulting organic layer was concentrated to obtain 0.44 g of a brown oily substance.

As a result of $^1\text{H-NMR}$ analysis, this oily substance was found to be a mixture containing 75 wt% benzyl (2*S*,5*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate (1.27 mmol; yield, 19%), 17 wt% ethyl acetate, and 9 wt% toluene.

In the material *N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid, (2*S*,5*S*) and (2*S*,5*R*) were contained at (2*S*,5*S*): (2*S*,5*R*)=4:6 (molar ratio). By its conversion to benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate,

compounds having the stereochemistry of (2*S*,5*R*) could be effectively removed.

However, since the yield of benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate was 19%, the (2*S*,5*S*) isomer was obtained in a moderate yield of about 50%. It has been observed that a side reaction occurred between benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate generated by the reaction and (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid left unreacted in the system, to cause esterification. This is assumed to be the cause of the moderate yield.

[0075]

10 <2-3> Production of (2*S*,5*S*)-*N*-Benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic Acid

To a flask, 0.44 g (purity, 75 wt%; 1.27 mmol) of the benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate obtained in <2-2>, 2.6 mL of methanol, and 2.6 mL of 1 mol/L aqueous sodium hydroxide solution were added, and left to stand overnight. To the resulting reaction mixture, 1.3 mL of 2 mol/L hydrochloric acid was added to adjust the pH to 4, and extraction with ethyl acetate was then carried out. The resulting organic layer was concentrated to obtain 0.33 g of a brown oily substance. The substance was then purified by silica gel chromatography to obtain 0.25 g of (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid (0.88 mmol; yield, 69%).

Example 3

[0076]

25 <3-1> Production of (2*S*,5*S*)-*N*-Benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic Acid and (2*S*,5*R*)-*N*-Benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic Acid

To a flask, 0.76 g of (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid and (2*S*,5*R*)-5-hydroxy-2-piperidinecarboxylic acid [5.24 mmol, (2*S*,5*S*): (2*S*,5*R*)=7:3

(molar ratio)], and 6.3 mL of 1 mol/L aqueous sodium hydroxide solution were added, and 0.93 mL (5.3 mmol) of benzyloxycarbonyl chloride was added to the resulting mixture under ice-cooling. The temperature of the mixture was then slowly increased to room temperature, and 2 mL of tetrahydrofuran and 5.2 mL of 1 mol/L aqueous sodium hydroxide solution were dividedly added thereto while the pH was adjusted to about 9. After leaving the resulting mixture to stand overnight, the organic layer was separated, and the aqueous layer was washed with ethyl acetate. To the aqueous layer, 3.2 mL of 2 mol/L hydrochloric acid was added to adjust the pH to 3, and the resulting mixture was extracted 3 times with ethyl acetate. The resulting organic layer was concentrated to obtain 0.79 g of a yellow oily substance.

As a result of ^1H -NMR analysis, this oily substance was found to be a mixture containing 76 wt% (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid and (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid (2.15 mmol; yield, 41%; (2*S*,5*S*): (2*S*,5*R*)=7:3 (molar ratio)), and 24 wt% ethyl acetate.

[0077]

<3-2> Production of Benzyl (2*S*,5*S*)-5-Aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate

To a flask, 0.79 g of the (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid and (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid obtained in <3-1> (purity, 76 wt%; 2.15 mmol; (2*S*,5*S*): (2*S*,5*R*)=7:3 (molar ratio)), 5 mL of toluene, and 49 mg (0.28 mmol) of *p*-toluenesulfonic acid monohydrate were added, and the reaction was allowed to proceed at 60°C for 2 hours. After cooling the resulting reaction product to room temperature, 10 mL of ethyl acetate, 3 mL of water, and 1.2 mL of 1 mol/L sodium hydroxide solution were added thereto. As a result, the pH of the aqueous layer became 7. After separating the aqueous layer, re-extraction was carried out with ethyl acetate. The obtained organic layer was washed with saturated aqueous

sodium bicarbonate solution and saturated brine. The resulting organic layer was concentrated to obtain 0.34 g of a brown oily substance.

As a result of ^1H -NMR analysis, this oily substance was found to be a mixture containing 62 wt% benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate (0.80 mmol; yield, 37%), 21 wt% (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid (0.25 mmol; yield, 12%), and 17 wt% toluene.

In the material *N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid, (2*S*,5*S*) and (2*S*,5*R*) were contained at (2*S*,5*S*):(2*S*,5*R*)=7:3 (molar ratio). By its conversion to benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate, compounds having the stereochemistry of (2*S*,5*R*) could be effectively removed. The total yield of benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate and (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid was 49%, and the (2*S*,5*S*) isomer was obtained in a moderate yield of about 70%.

Similarly to <2-1>, a side reaction occurred between benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate generated by the reaction and (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid left unreacted in the system, to cause esterification. However, since the amount of (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid in the starting material was small, the (2*S*,5*S*) isomer could be selectively, and relatively efficiently recovered.

[0078]

<3-3> Production of (2*S*,5*S*)-*N*-Benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic Acid

To a flask, 0.34 g of the oily substance obtained in <3-2> [containing 0.80 mmol benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate and 0.25 mmol (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid], 2 mL of methanol, and 1.9 mL of 1 mol/L aqueous sodium hydroxide solution were added,

and left to stand overnight. To the resulting reaction mixture, 2 mL of 2 mol/L hydrochloric acid was added to adjust the pH to 4, and extraction with ethyl acetate was then carried out. The resulting organic layer was concentrated to obtain 0.39 g of a brown oily substance.

5 As a result of ^1H -NMR analysis, this oily substance was found to be a mixture containing 76 wt% (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid (1.05 mmol; yield, 100%; (2*S*,5*S*):(2*S*,5*R*)=87:13 (molar ratio)), 23 wt% ethyl acetate, and 1 wt% acetic acid.

[0079]

10 <3-4> Production of (2*S*,5*S*)-5-Hydroxy-2-piperidinecarboxylic acid

To a flask, 0.39g of the (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid obtained in <3-3> (purity, 77 wt%; 1.06 mmol), 2 mL of methanol, and 61 mg of 10% palladium carbon (manufactured by N. E. Chemcat Corporation, PE-type, 55.3%, containing water) were added, and hydrogenation was

15 carried out under normal temperature and pressure for 3 hours. The palladium carbon was removed by filtration through Celite, and washing with methanol-water was carried. The obtained filtrate was concentrated to obtain 0.21 g of (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid as a pale yellow oily substance.

Example 4

20 [0080]

<4-1> Production of Benzyl (1*S*,4*S*)-5-Aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate and Ethyl (2*S*,5*S*)-*N*-Benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylate

To a flask, 0.26 g of (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid and (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid obtained according to the method of <2-1> (purity, 82 wt%; 0.76 mmol; (2*S*,5*S*):(2*S*,5*R*)=5:5 (molar ratio)), 1 mL of ethyl acetate, and 5 μL

25

(0.08 mmol) of methanesulfonic acid were added, and the reaction was allowed to proceed at 60°C for 3 hours. After cooling the resulting reaction product to room temperature, 10 mL of ethyl acetate and 1 mL of 1 mol/L sodium hydroxide solution were added thereto. As a result, the pH of the aqueous layer became about 10.

5 After separating the aqueous layer, re-extraction was carried out with ethyl acetate. The obtained organic layer was dried over magnesium sulfate. The resulting organic layer was concentrated to obtain 0.12 g of a pale brown oily substance.

As a result of ¹H-NMR analysis, this oily substance was found to be a mixture containing 26 wt% benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-
10 carboxylate (0.11 mmol; yield, 15%), 52 wt% ethyl (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid (0.20 mmol; yield, 26%), 13 wt% ethyl acetate, and 8 wt% toluene.

It is thought that, in the present Example, a small amount of ethanol generated from ethyl acetate reacted with benzyl (1*S*,4*S*)-5-aza-2-oxa-3-
15 oxobicyclo[2.2.2]octane-5-carboxylate, to generate ethyl (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylate. In the material *N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid, (2*S*,5*S*) and (2*S*,5*R*) were contained at (2*S*,5*S*):(2*S*,5*R*)=5:5 (molar ratio). As a result of ¹H-NMR analysis and HPLC analysis (under the conditions of <HPLC-2>), it could be confirmed that the
20 isomer ratio of the ethyl *N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylate was as follows: (2*S*,5*S*):(2*S*,5*R*)>10:1 (molar ratio). Thus, compounds having the stereochemistry of (2*S*,5*R*) could be effectively removed. Since the total yield of benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate and ethyl (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylate was 41%, the yield
25 of the (2*S*,5*S*) isomer was about 80%. Thus, the (2*S*,5*S*) isomer could be selectively and efficiently recovered.

[0081]

<4-2> Production of (2*S*,5*S*)-*N*-Benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic Acid

To a flask, 0.12 g of the oily substance obtained in <4-1> (containing 0.11 mmol of benzyl (1*S*,4*S*)-5-aza-2-oxa-3-oxobicyclo[2.2.2]octane-5-carboxylate and 5 0.20 mmol of ethyl (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylate), 1 mL of ethanol, and 0.66 mL of 1 mol/L aqueous sodium hydroxide solution were added, and left to stand overnight. The reaction mixture was then concentrated, and 0.7 mL of 1 mol/L hydrochloric acid was added thereto to adjust the pH to 3, followed by carrying out extraction with ethyl acetate. The 10 resulting organic layer was concentrated under reduced pressure to obtain 0.11 g of a brown oily substance.

As a result of ¹H-NMR analysis, this oily substance was found to be a mixture containing 84 wt% (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid (0.33 mmol, quantitative, (2*S*,5*S*):(2*S*,5*R*)=10:1 (molar ratio)), 11 wt% ethyl 15 acetate, and 5 wt% toluene.

[0082]

<4-3> Production of (2*S*,5*S*)-5-Hydroxy-2-piperidinecarboxylic Acid

To a flask, 0.11 g of the (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid obtained in <4-2> (purity, 82 wt%, 0.33 mmol), 1 mL of 20 methanol, and 18 mg of 10% palladium carbon (manufactured by N. E. Chemcat Corporation, PE-type, 55.3%, containing water) were added, and hydrogenation was carried out under normal temperature and pressure for 9 hours. The palladium carbon was removed by filtration through Celite, and washing with methanol-water was carried. The obtained filtrate was concentrated to obtain 58 mg of (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid as a pale yellow oily substance 25 ((2*S*,5*S*):(2*S*,5*R*)=10:1 (molar ratio), ¹H-NMR analysis).

Example 5

[0083]

<5-1> Production of Ethyl (2*S*,5*S*)-*N*-Benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylate

5 To a flask, 0.55 g of (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid and (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid obtained according to the method of <2-1> (purity, 78 wt%; 1.52 mmol; (2*S*,5*S*): (2*S*,5*R*)=3:7 (molar ratio)), 2 mL of toluene, 0.5 mL of ethanol, and 10 μ L (0.15 mmol) of methanesulfonic acid were added, and the
10 reaction was allowed to proceed at 40°C for 5 hours.

The reaction mixture was analyzed by HPLC under the conditions of <HPLC-2>, and the following result was obtained: ethyl (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylate:ethyl (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylate:(2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-
15 piperidinecarboxylic acid:(2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid=24:8:7:57.

Under the assumption that the material and products have the same absorption coefficient, the degree of conversion of the (2*S*,5*S*) isomers was about 77%, and the degree of conversion of the (2*S*,5*R*) isomers was about 12%. Thus, the (2*S*,5*S*)
20 isomers showed a higher degree of esterification.

[0084]

<5-1'> Production of Isopropyl (2*S*,5*S*)-*N*-Benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylate

To a flask, 0.62 g of (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-
25 piperidinecarboxylic acid and (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid obtained according to the method of <2-1> (purity, 78 wt%; 1.72 mmol; (2*S*,5*S*): (2*S*,5*R*)=3:7 (molar ratio)), 2 mL of toluene, 0.5 mL of 2-

propanol, and 11 μL (0.17 mmol) of methanesulfonic acid were added, and the reaction was allowed to proceed at 50°C for 5 hours. After cooling the resulting reaction product to room temperature, 3 mL of toluene and 1.2 mL of 1 mol/L sodium hydroxide solution were added thereto. As a result, the pH of the aqueous layer became about 9. After separating the aqueous layer, re-extraction was carried out with toluene. The resulting organic layer was concentrated to obtain 0.17 g of a pale brown oily substance.

As a result of ^1H -NMR analysis, this oily substance was found to be a mixture containing 91 wt% isopropyl *N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylate (0.47 mmol; yield, 27%; (2*S*,5*S*):(2*S*,5*R*)=10:1 (molar ratio)), and 9 wt% toluene.

In the present Example, the material *N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid contained (2*S*,5*S*) and (2*S*,5*R*) at (2*S*,5*S*):(2*S*,5*R*)=3:7 (molar ratio). Based on the result of ^1H -NMR analysis, the isomer ratio of the isopropyl *N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylate was follows: (2*S*,5*S*):(2*S*,5*R*)=10:1 (molar ratio). Thus, compounds having the stereochemistry of (2*S*,5*R*) could be effectively removed. Since the yield of the isopropyl (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylate was 27%, the yield of the (2*S*,5*S*) isomer was about 80%. Thus, the (2*S*,5*S*) isomer could be selectively and efficiently recovered.

[0085]

<5-2> Production of (2*S*,5*S*)-*N*-Benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic Acid

To a flask, 0.17 g of the isopropyl (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylate obtained in <5-1'> (purity, 91 wt%; 0.47 mmol), 0.7 mL methanol, and 0.71 mL of 1 mol/L aqueous sodium hydroxide solution were added, and the reaction was allowed to proceed at 60°C for 4 hours. After washing the

reaction mixture with toluene, 0.45 mL of 2 mol/L hydrochloric acid was added thereto to adjust the pH to 3, and extraction with ethyl acetate was carried out. The resulting organic layer was concentrated to obtain 0.17 g of a brown oily substance. As a result of ¹H-NMR analysis, this oily substance was found to be a mixture
 5 containing 70 wt% *N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid (0.41 mmol; yield, 88%; (2*S*,5*S*):(2*S*,5*R*)=7:1 (molar ratio)), 26 wt% ethyl acetate, and 4 wt% toluene.

[0086]

<5-3> Production of (2*S*,5*S*)-5-Hydroxy-2-piperidinecarboxylic Acid

10 To a flask, 0.17 g of the (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid obtained in <5-2> (purity, 70 wt%; 0.41 mmol; (2*S*,5*S*):(2*S*,5*R*)=7:1 (molar ratio)), 1 mL of methanol, and 22 mg of 10% palladium carbon (manufactured by N. E. Chemcat Corporation, PE-type, 55.3%, containing water) were added, and hydrogenation was carried out under normal temperature and
 15 pressure for 3 hours. The palladium carbon was removed by filtration through Celite, and washing with methanol-water was carried. The obtained filtrate was concentrated to obtain 89 mg of (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid as a pale yellow oily substance.

20 [0087]

[Reference Example 3]

<3-1> Cloning of Lysine Hydroxylase Gene

Based on a gene sequence (hyl-1, SEQ ID NO:17) encoding an L-arginine-β hydroxylase VioC homologue Hyl-1 derived from the *Flavobacterium johnsoniae*
 25 NBRC14942 strain (GenBank Accession No. ABQ06186, SEQ ID NO:18), primers for amplifying the full-length sequence of the hyl-1 gene, hyl1_F (SEQ ID NO:29) and hyl1_R (SEQ ID NO:30), were designed and synthesized. Using chromosomal

DNA of *Flavobacterium johnsoniae* as a template, PCR was carried out according to a conventional method, to obtain a DNA fragment of about 1.0 kbp.

In addition, VioC homologues derived from the *Kineococcus radiotolerans* NBRC101839 strain, *Chitinophaga pinensis* NBRC15968 strain, *Chryseobacterium*
 5 *gleum* NBRC15054 strain, and *Niastella koreensis* NBRC106392 strain were designated Hyl-2 (GenBank Accession No.ABS05421, SEQ ID NO:20), Hyl-3 (GenBank Accession No.ACU60313, SEQ ID NO:22), Hyl-4 (GenBank Accession No.EFK34737, SEQ ID NO:24), and Hyl-5 (GenBank Accession No.AEV99100, SEQ ID NO:26), respectively. Based on gene sequences encoding the enzymes,
 10 (hyl-2 (SEQ ID NO:19), hyl-3 (SEQ ID NO:21), hyl-4 (SEQ ID NO:23), and hyl-5 (SEQ ID NO:25)), primers for amplifying the full-length sequence of each gene were designed and synthesized. Primers hyl2_f (SEQ ID NO:31) and hyl2_r (SEQ ID NO:32) for hyl-2, primers hyl3_f (SEQ ID NO:33) and hyl3_r (SEQ ID NO:34) for hyl-3, primers hyl4_f (SEQ ID NO:35) and hyl4_r (SEQ ID NO:36) for hyl-4, and
 15 primers hyl5_f (SEQ ID NO:37) and hyl5_r (SEQ ID NO:38) for hyl-5 were synthesized, and PCR was carried out using chromosomal DNA of each strain according to a conventional method. Each reaction produced a DNA fragment of about 1.0 kbp.

Each of the 5 kinds of DNA fragments obtained was digested with restriction
 20 enzymes *Nde*I and *Xho*I, and ligated into *Nde*I/*Xho*I-digested pET21a (Novagen) according to a conventional method, to obtain pEHYL1, pEHYL2, pEHYL3, pEHYL4, and pEHYL5, respectively.

A gene sequence (hyl-6, SEQ ID NO:27) encoding Hyl-6 (GenBank Accession No. ABS05421, SEQ ID NO:28), which was derived from a marine
 25 actinobacterium PHSC20C1, was artificially synthesized, and inserted into pJExpress401 (DNA2.0) to prepare a plasmid pJHYL6.

Subsequently, *E. coli* (*Escherichia coli*) BL21(DE3) (manufactured by

Invitrogen) was transformed with each of the resulting plasmids according to a conventional method, to obtain recombinant *E. coli* BL21(DE3)/pEHYL1, BL21(DE3)/pEHYL2, BL21(DE3)/pEHYL3, BL21(DE3)/pEHYL4, BL21(DE3)/pEHYL5, and BL21(DE3)/pJHYL6. In order to obtain bacterial cells expressing the genes, each type of recombinant *E. coli* was cultured at 30°C using liquid LB medium supplemented with ampicillin and a lac promoter inducer, and collected after 20 hours of the culture.

[0088]

<3-2> Confirmation of Lysine Hydroxylase Activity by Resting-cell Reaction

In a plastic tube, 5 mmol/L L-lysine, 10 mmol/L 2-oxoglutaric acid, 1 mmol/L L-ascorbic acid, 0.1 mmol/L iron sulfate, and recombinant *E. coli* obtained by the method according to Reference Example <3-1> were mixed to provide a reaction mixture such that the turbidity OD₆₀₀ was 10. The reaction was allowed to proceed in 0.5 mL of the prepared mixture at 30°C at pH 7.0 for 3 hours. The reaction product was derivatized with 1-fluoro-2,4-dinitrophenyl-5-L-alaninamide (FDAA), and hydroxylysine was analyzed by HPLC under the conditions of <HPLC-4>. As a result, as shown in Fig. 2 and Fig. 3, it could be confirmed that BL21(DE3)/pEHYL2 and BL21(DE3)/pJHYL6 produced a compound corresponding to the retention time of a standard product of 3-hydroxylysine, 8.04 minutes. It could also be confirmed that BL21(DE3)/pEHYL1, BL21(DE3)/pEHYL3, BL21(DE3)/pEHYL4, and BL21(DE3)/pEHYL5 produced a compound corresponding to the retention time of a standard product of 4-hydroxylysine, 8.16 minutes.

[0089]

<3-3> Synthesis of (2*S*,3*S*)-3-Hydroxylysine

To a 1-L jar fermenter, 35 mL of 1 mol/L potassium phosphate buffer (pH 7.0), 304 mL of desalted water, 1.28 g of L-lysine hydrochloride, 2.05 g of 2-oxoglutaric acid, 0.14 g of sodium L-ascorbate, 0.02 g of iron sulfate, 0.35 g of

Adekanol LG109, and 8 g of wet cells of recombinant *E. coli* BL21(DE3)/pEHL2 obtained by the method according to Reference Example <3-1> were mixed together, and the reaction was allowed to proceed at 30°C, pH 7.0, a stirring rate of 500 rpm, and an aeration rate of 2.0 vvm for 17 hours. Completion of the reaction was
 5 judged by confirming disappearance of the peak for L-lysine by carrying out HPLC analysis under the conditions of <HPLC-5>. From the liquid after the completion of the reaction, bacterial cells and bacterial debris were removed by centrifugation and microfiltration, to obtain 390 g of a filtrate.

After passing 390 g of the filtrate through an ion-exchange resin column
 10 (DIAION (registered trademark) SK-1B (Type H), 60.0 g), washing was carried out with water, followed by elution with an aqueous solution containing 150 mmol of ammonia. The ammonia eluate was concentrated to obtain 1.0 g of (2*S*,3*S*)-3-hydroxylysine (6.17 mmol; yield, 88%).

¹H-NMR (400MHz, D₂O) δ, 1.45-1.58 (2H, m), 1.63-1.73 (1H, m), 1.74-1.88 (1H,
 15 m), 2.93-3.04 (2H, m), 3.47 (1H, d, *J*=4.3 Hz), 3.89 (1H, dt, *J*=8.4, 4.5 Hz)
 [0090]


<3-4> Determination of Stereochemistry of (2*S*,3*S*)-3-Hydroxylysine

To a flask, 8.3 mg (0.051 mmol) of (2*S*,3*S*)-3-hydroxylysine obtained by the method according to Reference Example <3-3>, 0.26 ml of 1 mol/L aqueous sodium
 20 hydroxide solution, and 18 μL (0.13 mmol) of benzyloxycarbonyl chloride were added, and the resulting mixture was stirred at room temperature for 1 hour. To the mixture, 0.26 mL of 1 mol/L aqueous sodium hydroxide solution and 18 μL (0.13 mmol) of benzyloxycarbonyl chloride were further added, and the reaction was allowed to proceed overnight at room temperature. Subsequently, 0.5 mL of
 25 tetrahydrofuran was added to the reaction mixture, and the reaction was allowed to proceed at 60°C for additional 2 hours. After cooling the reaction mixture to room temperature, 95 mg of sodium hydroxide was added thereto, and the reaction was

allowed to proceed overnight at room temperature. The reaction mixture was washed twice with toluene-tetrahydrofuran (1:1), and 250 μ L of concentrated hydrochloric acid was added to the reaction mixture to make the reaction mixture strongly acidic. After 3 times of washing with ethyl acetate, the aqueous layer was extracted 4 times with 1-butanol. The 1-butanol layer was dried over anhydrous magnesium sulfate, and then concentrated, to obtain 14.6 mg of (4*S*,5*S*)-5-(3-benzyloxycarbonylamino-
 5 propyl)-2-oxo-4-oxazolidinecarboxylic acid (0.045 mmol; yield, 89%).

The stereochemistry of the obtained (4*S*,5*S*)-5-(3-benzyloxycarbonylamino-
 10 propyl)-2-oxo-4-oxazolidinecarboxylic acid was determined by NOESY measurement. The NOESY measurement was carried out using an AVANCE DRX500 spectrometer manufactured by Bruker (equipped with a CryoProbe) at 25°C with a mixing time of 0.8 msec. As a chemical shift reference, 3.31 ppm for methanol was used. The result of NOESY measurement was as
 15 follows.

[0091]



[0092]

Since a cross peak was found between the 3-position hydrogen atom (H^4) and
 20 4-position hydrogen atom (H^5), but was not found between the 4-position and the 1'-position, the substituents at the 4-position and the 5-position could be confirmed to have the *cis* configuration. Since the absolute configuration of the lysine used in the enzymatic reaction was *S*, it could be confirmed that the 5-(3-benzyloxycarbonylamino-
 20 propyl)-2-oxo-4-oxazolidinecarboxylic acid obtained in the

present Reference Example has the stereochemistry of (4*S*,5*S*), and that 3-hydroxylysine as its material has the stereochemistry of (2*S*,3*S*).

¹H-NMR (400 MHz, MeOH-d₄) δ, 1.39-1.53 (3H, m, H^{1'}, H^{2'} x2), 1.59-1.68 (1H, m, H^{1'}), 3.02-3.08 (2H, m, H^{3'}), 3.60-3.64 (1H, m, H⁴), 3.93-4.00 (1H, m, H⁵), 4.90-5.02 (2H, m, Bn), 7.18-7.28 (5H, m, Bn).

[0093]

[Reference Example 4]

Production of (2*S*,3*S*)-3-Hydroxypipicollic Acid [(2*S*,3*S*)-3-Hydroxy-2-piperidinecarboxylic Acid]

In a plastic tube, 0.75 mL of 1 mol/L tris hydroxymethyl aminomethane buffer (pH 8.0), 9.21 mL of desalted water, 86 mg of the (2*S*,3*S*)-3-hydroxylysine obtained in Reference Example <3-3>, 0.083 mL of 50 mmol/L NADPH, 0.7 mL of 1.0 mol/L glucose, and 1.25 mL of a 100-g/L suspension of the recombinant *E. coli* JM109/pKW32 (dpkA, aip, gdh, kr) obtained in Reference Example 1 were mixed together, and the reaction was allowed to proceed at 30°C at pH 8.0 at a stirring rate of 1000 rpm for 20 hours. Completion of the reaction was judged by confirming disappearance of the peak for (2*S*,3*S*)-3-hydroxylysine by HPLC analysis under the conditions of <HPLC-6>. From the liquid after the completion of the reaction, bacterial cells and bacterial debris were removed by centrifugation, to obtain 10.5 g of a supernatant.

After passing 10.5 g of the supernatant through an ion-exchange resin column (DIAION (registered trademark) SK-1B (Type H), 4.0g), washing was carried out with water, followed by elution with an aqueous solution containing 16.4 mmol of ammonia. The ammonia eluate was concentrated to obtain 255 mg of a solid brown substance. As a result of NMR analysis, this solid substance was found to be a mixture containing 20 wt% (2*S*,3*R*)-3-hydroxypipicollic acid (0.35 mmol; yield,

66.3%) and 80 wt% tris hydroxymethyl aminomethane.

¹H-NMR (400 MHz, D₂O) δ , 1.38-1.56 (2H, m), 1.73-1.85 (2H, m), 2.71-2.79 (1H, m), 3.04-3.11 (1H, m), 3.23 (1H, d, $J=7.6$ Hz), 3.79-3.86 (1H, m).

5 Example 6

[0094]

Production of (2*S*,5*S*)-*N*-Benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic Acid

To a flask, 78.7 mg (0.542 mmol) of (2*S*,5*S*)-5-hydroxy-2-piperidinecarboxylic acid and 2.8 mg (0.019 mmol) of (2*S*,5*R*)-5-hydroxy-2-piperidinecarboxylic acid obtained according to the method of Example 1, and 98.5 mg of (2*S*,3*S*)-3-hydroxy-2-piperidinecarboxylic acid obtained according to the method of Reference Example 4 (purity, 20 wt%, 0.136 mmol; containing 78.8 mg of tris hydroxymethyl aminomethane) were added, and 0.200 mL of water and 0.674 mL of 2 mol/L aqueous sodium hydroxide solution were added thereto at room temperature. After adding 0.190 mL (1.35 mmol) of benzyloxycarbonyl chloride dropwise to the resulting mixture, 0.500 mL of 2 mol/L aqueous sodium hydroxide solution was added thereto to adjust the pH from 9 to 10. The reaction was allowed to proceed at room temperature for 25 minutes, and 0.098 mL (0.697 mmol) of benzyloxycarbonyl chloride was then added dropwise to the resulting reaction mixture, followed by adding 0.400 mL of 2 mol/L aqueous sodium hydroxide solution thereto to adjust the pH from 10 to 11. The reaction was further allowed to proceed at room temperature for 40 minutes, and 0.098 mL (0.697 mmol) of benzyloxycarbonyl chloride was added dropwise thereto, followed by allowing the reaction to proceed at room temperature for 2 hours. To the resulting reaction mixture, 3 mL of toluene was added, and liquid separation was then carried out. To the extracted organic layer, 0.500 mL of water was added, and liquid separation was carried out again. The resulting organic layer was concentrated to obtain 330 mg of

a pale yellow oily substance. To this oily substance, 2 mL of ethanol, 1.1 mL of 1 mol/L aqueous sodium hydroxide solution, and 0.3 mL of water were added, and the reaction was allowed to proceed at room temperature for about 1 hour. The resulting reaction mixture was concentrated, and ethanol was evaporated.

5 Thereafter, 1 mL of ethanol, 0.3 mL of water, 0.1 mL of 1 mol/L aqueous sodium hydroxide solution, and 0.1 mL of 2 mol/L aqueous sodium hydroxide solution were added thereto, and the reaction was allowed to proceed at room temperature for 2 hours. The reaction mixture was concentrated, and ethanol was evaporated, followed by carrying out liquid separation using 3 mL of toluene. To the obtained
10 aqueous layer, 1 mL of water and 0.75 mL of 1 mol/L hydrochloric acid were added to adjust the pH of the aqueous solution to 1. The aqueous layer was subjected twice to extraction with 4 mL of ethyl acetate, and the resulting organic layer was concentrated to obtain 92.0 mg of a pale yellow oily substance.

As a result of ^1H -NMR analysis, this pale yellow oily substance was found to
15 contain 80.5 wt% (2*S*,5*S*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid (0.265 mmol; yield, 49%) and 19.5 wt% ethyl acetate. No peak was found at all for either (2*S*,5*R*)-*N*-benzyloxycarbonyl-5-hydroxy-2-piperidinecarboxylic acid or (2*S*,3*S*)-*N*-benzyloxycarbonyl-3-hydroxy-2-piperidinecarboxylic acid by ^1H -NMR.

20 INDUSTRIAL APPLICABILITY


[0095]

The present invention can be used as a method for purifying *cis*-5-hydroxy-2-piperidinecarboxylic acid, which is useful as an intermediate for pharmaceuticals, and can also be used as a method for producing its derivatives.


25

CLAIMS

1. A method for producing a *cis*-5-hydroxy-2-piperidinecarboxylic acid derivative, said method comprising a step of converting *cis*-5-hydroxy-2-piperidinecarboxylic acid into a compound(s) of Formula (1) and/or Formula (2) (wherein R¹ represents a protective group for an amino group, and R² represents a C₁-C₆ alkyl group).




(1)




(2)

2. The method for producing a *cis*-5-hydroxy-2-piperidinecarboxylic acid derivative according to claim 1, comprising a step of reacting *cis*-5-hydroxy-2-piperidinecarboxylic acid with an acid halide and/or acid anhydride to convert said *cis*-5-hydroxy-2-piperidinecarboxylic acid into a compound of Formula (1).
3. The method for producing a *cis*-5-hydroxy-2-piperidinecarboxylic acid derivative according to claim 1, comprising a step of reacting *cis*-5-hydroxy-2-piperidinecarboxylic acid with an acid halide and/or acid anhydride, and then with an alcohol in the presence of an acid catalyst, to convert said *cis*-5-hydroxy-2-piperidinecarboxylic acid into a compound of Formula (2).
4. A method for regenerating *cis*-5-hydroxy-2-piperidinecarboxylic acid, said method comprising the steps of:
- converting *cis*-5-hydroxy-2-piperidinecarboxylic acid into a compound(s) of Formula (1) and/or Formula (2) (wherein R¹ represents a protective group for an amino group, and R² represents a C₁-C₆ alkyl group); and
- converting the compound(s) of Formula (1) and/or Formula (2) into *cis*-5-hydroxy-2-piperidinecarboxylic acid.



(1)




(2)


5. The method for producing a *cis*-5-hydroxy-2-piperidinecarboxylic acid derivative according to any one of claims 1 to 3, wherein said *cis*-5-hydroxy-2-piperidinecarboxylic acid is *cis*-5-hydroxy-2-piperidinecarboxylic acid synthesized by bacterial reaction and/or enzymatic reaction.

6. The method for regenerating *cis*-5-hydroxy-2-piperidinecarboxylic acid according to claim 4, wherein said *cis*-5-hydroxy-2-piperidinecarboxylic acid is *cis*-5-hydroxy-2-piperidinecarboxylic acid synthesized by bacterial reaction and/or enzymatic reaction.

7. A method for purifying *cis*-5-hydroxy-2-piperidinecarboxylic acid, said method comprising a step of reacting a mixture containing *cis*-5-hydroxy-2-piperidinecarboxylic acid and an impurity with an acid halide and/or acid anhydride, or with an acid halide and/or acid anhydride and an alcohol, to convert the *cis*-5-hydroxy-2-piperidinecarboxylic acid into a compound(s) of Formula (1) and/or Formula (2), separating said compound(s) and then converting the separated compound(s) into *cis*-5-hydroxy-2-piperidinecarboxylic acid.



(1)



(2)

8. The method for purifying *cis*-5-hydroxy-2-piperidinecarboxylic acid according to claim 7, wherein said impurity is 2-piperidinecarboxylic acid or an analogue thereof.

9. The method for purifying *cis*-5-hydroxy-2-piperidinecarboxylic acid

according to claim 8, wherein said 2-piperidinecarboxylic acid or an analogue thereof is *trans*-5-hydroxy-2-piperidinecarboxylic acid.

10. The method for purifying *cis*-5-hydroxy-2-piperidinecarboxylic acid according to any one of claims 7 to 9, wherein said step of separating the
5 compound(s) of Formula (1) and/or Formula (2) is carried out by crystallization or solvent extraction.

11. The method for purifying *cis*-5-hydroxy-2-piperidinecarboxylic acid according to any one of claims 7 to 10, wherein said mixture containing *cis*-5-hydroxy-2-piperidinecarboxylic acid and an impurity is a mixture synthesized by
10 bacterial reaction and/or enzymatic reaction.

JPOXMLDOC01-seql.app
SEQUENCE LISTING

<110> API Corporation

<120> Method of producing cis-5-hydroxy-2-piperidine carboxylate derivatives and method of purifying cis-5-hydroxy-2-piperidine carboxylate derivatives

<130> AP340-13366

<150> JP2012-278314

<151> 2012-12-20

<160> 38

<170> PatentIn version 3.5

<210> 1

<211> 1026

<212> DNA

<213> Pseudomonas putida

<220>

<221> CDS

<222> (1)..(1023)

<400> 1

atg tcc gca cct tcc acc agc acc gtt gtg cgc gtg cct ttt acc gag	48
Met Ser Ala Pro Ser Thr Ser Thr Val Val Arg Val Pro Phe Thr Glu	
1 5 10 15	
ctg caa agc ctg ttg cag gcc att ttc cag cgc cat ggg tgc agc gag	96
Leu Gln Ser Leu Leu Gln Ala Ile Phe Gln Arg His Gly Cys Ser Glu	
20 25 30	
gcc gtg gcc cgg gtg ctg gcc cac aac tgc gcc agc gcc cag cgt gat	144
Ala Val Ala Arg Val Leu Ala His Asn Cys Ala Ser Ala Gln Arg Asp	
35 40 45	
ggc gcc cat agc cat ggg gtg ttc cgc atg ccc ggt tat gtc tcg acc	192
Gly Ala His Ser His Gly Val Phe Arg Met Pro Gly Tyr Val Ser Thr	
50 55 60	
ttg gcc agc ggc tgg gtc gat ggc cag gcc acg cca cag gtc agc gac	240
Leu Ala Ser Gly Trp Val Asp Gly Gln Ala Thr Pro Gln Val Ser Asp	
65 70 75 80	
gtg gcc gcc ggc tat gtg cgt gtc gat gct gcg ggc ggt ttt gcc cag	288
Val Ala Ala Gly Tyr Val Arg Val Asp Ala Ala Gly Gly Phe Ala Gln	
85 90 95	
cca gca ctg gcg gcg gcc cgt gag ctg ttg gtg gcg aag gcg cgc agc	336
Pro Ala Leu Ala Ala Arg Glu Leu Val Ala Lys Ala Arg Ser	
100 105 110	
gca ggc att gcc gtg ctg gcg atc cac aac tcg cac cac ttc gcc gcg	384
Ala Gly Ile Ala Val Leu Ala Ile His Asn Ser His His Phe Ala Ala	
115 120 125	
cta tgg ccg gat gtc gag ccg ttc gcc gaa gag ggc ctg gta gcc ctc	432
Leu Trp Pro Asp Val Glu Pro Phe Ala Glu Glu Gly Leu Val Ala Leu	
130 135 140	
agc gtg gtc aac agc atg acc tgc gtg gtg ccg cat ggt gca cgc aag	480
Ser Val Val Asn Ser Met Thr Cys Val Val Pro His Gly Ala Arg Lys	
145 150 155 160	
ccg ctg ttc ggt acc aac ccc atc gct ttt gct gcg cct tgc gcc gag	528
Pro Leu Phe Gly Thr Asn Pro Ile Ala Phe Ala Ala Pro Cys Ala Glu	
165 170 175	
cat gac ccg atc gtt ttc gac atg gcc acc agt gcc atg gcc cat ggc	576
His Asp Pro Ile Val Phe Asp Met Ala Thr Ser Ala Met Ala His Gly	
180 185 190	

JPOXMLDOC01-seql.app

gat	gtg	cag	att	gcc	gcg	cgc	gcc	ggc	cag	caa	ttg	ccg	gag	ggc	atg	624
Asp	Val	Gln	Ile	Ala	Ala	Arg	Ala	Gly	Gln	Gln	Leu	Pro	Glu	Gly	Met	
	195						200				205					
ggg	gtg	gat	gcc	gat	ggc	cag	ccg	acc	acc	gac	ccg	aag	gcg	atc	ctg	672
Gly	Val	Asp	Ala	Asp	Gly	Gln	Pro	Thr	Thr	Asp	Pro	Lys	Ala	Ile	Leu	
	210					215					220					
gaa	ggc	ggc	gcc	ttg	ctg	cca	ttt	ggc	ggg	cac	aag	ggc	tcg	gcg	ttg	720
Glu	Gly	Gly	Ala	Leu	Leu	Pro	Phe	Gly	Gly	His	Lys	Gly	Ser	Ala	Leu	
	225				230					235					240	
tcg	atg	atg	gtc	gag	ctg	ctg	gcg	gcg	gcg	ctg	acc	ggc	ggt	cat	ttc	768
Ser	Met	Met	Val	Glu	Leu	Leu	Ala	Ala	Ala	Leu	Thr	Gly	Gly	His	Phe	
			245						250					255		
tcc	tgg	gag	ttc	gat	tgg	tcc	ggg	cat	ccg	ggg	gcg	aaa	acg	cca	tgg	816
Ser	Trp	Glu	Phe	Asp	Trp	Ser	Gly	His	Pro	Gly	Ala	Lys	Thr	Pro	Trp	
			260					265					270			
acc	ggg	cag	ttg	atc	atc	gtc	atc	aac	cca	ggc	aag	gcc	gag	ggc	gag	864
Thr	Gly	Gln	Leu	Ile	Ile	Val	Ile	Asn	Pro	Gly	Lys	Ala	Glu	Gly	Glu	
		275					280					285				
cgc	ttt	gcc	cag	cgc	agc	cgc	gag	ctg	gtg	gag	cac	atg	cag	gcg	gtg	912
Arg	Phe	Ala	Gln	Arg	Ser	Arg	Glu	Leu	Val	Glu	His	Met	Gln	Ala	Val	
	290					295					300					
ggg	ctg	acg	cgc	atg	ccg	ggc	gag	cgg	cgc	tac	cgt	gag	cgc	gag	gtg	960
Gly	Leu	Thr	Arg	Met	Pro	Gly	Glu	Arg	Arg	Tyr	Arg	Glu	Arg	Glu	Val	
	305				310					315					320	
gcc	gag	gag	gag	ggg	gtg	gcg	gtg	acc	gag	cag	gag	ttg	caa	ggc	ctg	1008
Ala	Glu	Glu	Glu	Gly	Val	Ala	Val	Thr	Glu	Gln	Glu	Leu	Gln	Gly	Leu	
			325						330					335		
aaa	gag	ctg	ctt	ggc	tga											1026
Lys	Glu	Leu	Leu	Gly												
			340													

<210> 2
 <211> 341
 <212> PRT
 <213> Pseudomonas putida

<400> 2

Met	Ser	Ala	Pro	Ser	Thr	Ser	Thr	Val	Val	Arg	Val	Pro	Phe	Thr	Glu	
1				5					10					15		
Leu	Gln	Ser	Leu	Leu	Gln	Ala	Ile	Phe	Gln	Arg	His	Gly	Cys	Ser	Glu	
			20					25					30			
Ala	Val	Ala	Arg	Val	Leu	Ala	His	Asn	Cys	Ala	Ser	Ala	Gln	Arg	Asp	
		35					40					45				
Gly	Ala	His	Ser	His	Gly	Val	Phe	Arg	Met	Pro	Gly	Tyr	Val	Ser	Thr	
	50					55				60						
Leu	Ala	Ser	Gly	Trp	Val	Asp	Gly	Gln	Ala	Thr	Pro	Gln	Val	Ser	Asp	
	65				70					75				80		
Val	Ala	Ala	Gly	Tyr	Val	Arg	Val	Asp	Ala	Ala	Gly	Gly	Phe	Ala	Gln	
			85					90					95			
Pro	Ala	Leu	Ala	Ala	Ala	Arg	Glu	Leu	Leu	Val	Ala	Lys	Ala	Arg	Ser	
			100					105					110			
Ala	Gly	Ile	Ala	Val	Leu	Ala	Ile	His	Asn	Ser	His	His	Phe	Ala	Ala	
		115					120					125				
Leu	Trp	Pro	Asp	Val	Glu	Pro	Phe	Ala	Glu	Glu	Gly	Leu	Val	Ala	Leu	
	130					135					140					
Ser	Val	Val	Asn	Ser	Met	Thr	Cys	Val	Val	Pro	His	Gly	Ala	Arg	Lys	
	145				150					155				160		
Pro	Leu	Phe	Gly	Thr	Asn	Pro	Ile	Ala	Phe	Ala	Ala	Pro	Cys	Ala	Glu	
				165					170					175		
His	Asp	Pro	Ile	Val	Phe	Asp	Met	Ala	Thr	Ser	Ala	Met	Ala	His	Gly	
			180					185					190			
Asp	Val	Gln	Ile	Ala	Ala	Arg	Ala	Gly	Gln	Gln	Leu	Pro	Glu	Gly	Met	
		195					200					205				

JPOXMLDOC01-seql.app

Gly	Val	Asp	Ala	Asp	Gly	Gln	Pro	Thr	Thr	Asp	Pro	Lys	Ala	Ile	Leu
210						215					220				
Glu	Gly	Gly	Ala	Leu	Leu	Pro	Phe	Gly	Gly	His	Lys	Gly	Ser	Ala	Leu
225					230					235					240
Ser	Met	Met	Val	Glu	Leu	Leu	Ala	Ala	Ala	Leu	Thr	Gly	Gly	His	Phe
				245						250				255	
Ser	Trp	Glu	Phe	Asp	Trp	Ser	Gly	His	Pro	Gly	Ala	Lys	Thr	Pro	Trp
			260					265					270		
Thr	Gly	Gln	Leu	Ile	Ile	Val	Ile	Asn	Pro	Gly	Lys	Ala	Glu	Gly	Glu
		275					280					285			
Arg	Phe	Ala	Gln	Arg	Ser	Arg	Glu	Leu	Val	Glu	His	Met	Gln	Ala	Val
	290					295					300				
Gly	Leu	Thr	Arg	Met	Pro	Gly	Glu	Arg	Arg	Tyr	Arg	Glu	Arg	Glu	Val
305					310					315					320
Ala	Glu	Glu	Glu	Gly	Val	Ala	Val	Thr	Glu	Gln	Glu	Leu	Gln	Gly	Leu
				325					330					335	
Lys	Glu	Leu	Leu	Gly											
			340												

<210> 3
 <211> 1488
 <212> DNA
 <213> Scomber japonicus

<220>
 <221> CDS
 <222> (1)..(1485)

<400>	3																		
atg	gag	cac	ttg	gct	gat	tgc	ttg	gaa	gat	aag	gat	tac	gat	aca	ctt				48
Met	Glu	His	Leu	Ala	Asp	Cys	Leu	Glu	Asp	Lys	Asp	Tyr	Asp	Thr	Leu				
1				5					10					15					
ttg	cag	aca	ttg	gac	aac	ggg	ttg	cct	cac	att	aac	act	tct	cat	cac				96
Leu	Gln	Thr	Leu	Asp	Asn	Gly	Leu	Pro	His	Ile	Asn	Thr	Ser	His	His				
			20					25					30						
gtt	gtc	att	gtt	ggg	gcc	gga	atg	gca	ggg	ttg	acc	gct	gcc	aaa	ttg				144
Val	Val	Ile	Val	Gly	Ala	Gly	Met	Ala	Gly	Leu	Thr	Ala	Ala	Lys	Leu				
			35				40					45							
ctt	caa	gat	gca	gga	cac	act	gtt	aca	atc	ttg	gag	gct	aac	gac	aga				192
Leu	Gln	Asp	Ala	Gly	His	Thr	Val	Thr	Ile	Leu	Glu	Ala	Asn	Asp	Arg				
			50			55				60									
gtt	ggg	gga	aga	gtc	gaa	act	tac	aga	aat	gaa	aag	gag	ggg	tgg	tat				240
Val	Gly	Gly	Arg	Val	Glu	Thr	Tyr	Arg	Asn	Glu	Lys	Glu	Gly	Trp	Tyr				
65				70					75					80					
gct	gaa	atg	gga	gcc	atg	aga	atc	cca	tct	tct	cat	aga	atc	gtt	cag				288
Ala	Glu	Met	Gly	Ala	Met	Arg	Ile	Pro	Ser	His	Arg	Ile	Val	Gln					
				85				90					95						
tgg	ttt	gtc	aag	aaa	ttg	ggg	gtt	gaa	atg	aac	gag	ttt	gtc	atg	aca				336
Trp	Phe	Val	Lys	Lys	Leu	Gly	Val	Glu	Met	Asn	Glu	Phe	Val	Met	Thr				
			100					105					110						
gat	gac	aac	acc	ttc	tac	ttg	gtt	aat	gga	gtc	aga	gaa	aga	act	tat				384
Asp	Asp	Asn	Thr	Phe	Tyr	Leu	Val	Asn	Gly	Val	Arg	Glu	Arg	Thr	Tyr				
			115				120					125							
gtt	gtc	caa	gag	aac	cct	gat	gtt	ttg	aag	tac	aat	gtc	tct	gaa	tcc				432
Val	Val	Gln	Glu	Asn	Pro	Asp	Val	Leu	Lys	Tyr	Asn	Val	Ser	Glu	Ser				
			130			135					140								
gag	aaa	ggg	att	tct	gct	gat	gac	ttg	ctt	gac	aga	gcc	ttg	cag	aag				480
Glu	Lys	Gly	Ile	Ser	Ala	Asp	Asp	Leu	Leu	Asp	Arg	Ala	Leu	Gln	Lys				
145					150					155				160					
gtt	aaa	gaa	gag	gtc	gaa	gct	aat	ggg	tgt	aag	gca	gct	ttg	gag	aag				528
Val	Lys	Glu	Glu	Val	Glu	Ala	Asn	Gly	Cys	Lys	Ala	Ala	Leu	Glu	Lys				
				165					170					175					

JPOXMLDOC01-seql.app

tac gat aga tac tct gtt aag gaa tat ttg aaa gaa gag ggt gga ctt	576
Tyr Asp Arg Tyr Ser Val Lys Glu Tyr Leu Lys Glu Glu Gly Gly Leu	
180 185 190	
tcc cca ggt gct gtt aga atg att gga gat ttg ctt aac gag caa tct	624
Ser Pro Gly Ala Val Arg Met Ile Gly Asp Leu Leu Asn Glu Gln Ser	
195 200 205	
ttg atg tac act gcc ctt tcc gaa atg atc tat gat cag gca gac gtt	672
Leu Met Tyr Thr Ala Leu Ser Glu Met Ile Tyr Asp Gln Ala Asp Val	
210 215 220	
aat gat tca gtc agt tac cac gag gtt aca ggt gga tcc gat ttg ctt	720
Asn Asp Ser Val Ser Tyr His Glu Val Thr Gly Gly Ser Asp Leu Leu	
225 230 235 240	
cca gaa gct ttc ttg tca gtt ctt gac gtc cct atc ttg ctt aac tcc	768
Pro Glu Ala Phe Leu Ser Val Leu Asp Val Pro Ile Leu Leu Asn Ser	
245 250 255	
aag gtt aag cat atc aga caa tca gat aag ggt gtt atc gtc agt tat	816
Lys Val Lys His Ile Arg Gln Ser Asp Lys Gly Val Ile Val Ser Tyr	
260 265 270	
cag act gga aat gaa tca agt ttg atg gac ctt tct gct gat att gtt	864
Gln Thr Gly Asn Glu Ser Ser Leu Met Asp Leu Ser Ala Asp Ile Val	
275 280 285	
ttg gtc act aca acc gcc aaa gcc gca ttg ttt att gat ttc gac cca	912
Leu Val Thr Thr Thr Ala Lys Ala Ala Leu Phe Ile Asp Phe Asp Pro	
290 295 300	
cct ttg tct atc tcc aag atg gag gct ttg aga tct gtt cac tac gat	960
Pro Leu Ser Ile Ser Lys Met Glu Ala Leu Arg Ser Val His Tyr Asp	
305 310 315 320	
tct tcc act aag atc ttg ctt act ttt aga gac aag ttc tgg gaa gat	1008
Ser Ser Thr Lys Ile Leu Leu Thr Phe Arg Asp Lys Phe Trp Glu Asp	
325 330 335	
gac ggt att aga ggt gga aag tca atc aca gat gga cca agt aga tac	1056
Asp Gly Ile Arg Gly Gly Lys Ser Ile Thr Asp Gly Pro Ser Arg Tyr	
340 345 350	
atc tac tat cct tca cat agt ttt cac acc aac gag act atc ggt gtt	1104
Ile Tyr Tyr Pro Ser His Ser Phe His Thr Asn Glu Thr Ile Gly Val	
355 360 365	
ttg ctt gcc tca tat act tgg tct gac gaa tcc ttg ctt ttc ttg gga	1152
Leu Leu Ala Ser Tyr Thr Trp Ser Asp Glu Ser Leu Leu Phe Leu Gly	
370 375 380	
gct tct gat gaa gag ttg aag gag ttg gcc ctt aga gac ttg gca aaa	1200
Ala Ser Asp Glu Glu Leu Lys Glu Leu Ala Leu Arg Asp Leu Ala Lys	
385 390 395 400	
att cac ggt gaa caa gtt tgg gat aag tgc aca gga gtt atc gtc aag	1248
Ile His Gly Glu Gln Val Trp Asp Lys Cys Thr Gly Val Ile Val Lys	
405 410 415	
aaa tgg tct gct gac cca tac tcc ttg ggt gca ttt gct ctt ttc acc	1296
Lys Trp Ser Ala Asp Pro Tyr Ser Leu Gly Ala Phe Ala Leu Phe Thr	
420 425 430	
cct tac caa cat ttg gaa tat gca cag gag ttg ttt tct tct gaa ggt	1344
Pro Tyr Gln His Leu Glu Tyr Ala Gln Glu Leu Phe Ser Ser Glu Gly	
435 440 445	
aga gtt cat ttt gct gga gag cac acc gca ttc cct cat gct tgg att	1392
Arg Val His Phe Ala Gly Glu His Thr Ala Phe Pro His Ala Trp Ile	
450 455 460	
gaa act tca atg aaa agt gct atc aga gct gcc aca aac att aac aaa	1440
Glu Thr Ser Met Lys Ser Ala Ile Arg Ala Ala Thr Asn Ile Asn Lys	
465 470 475 480	
gtc gct aac gaa gaa tcc acc att gaa cac aca aaa gat gag ttg taa	1488
Val Ala Asn Glu Glu Ser Thr Ile Glu His Thr Lys Asp Glu Leu	
485 490 495	

<210> 4
<211> 495

<212> PRT

<213> Scomber japonicus

<400> 4

```

Met Glu His Leu Ala Asp Cys Leu Glu Asp Lys Asp Tyr Asp Thr Leu
1      5      10      15
Leu Gln Thr Leu Asp Asn Gly Leu Pro His Ile Asn Thr Ser His His
20      25      30
Val Val Ile Val Gly Ala Gly Met Ala Gly Leu Thr Ala Ala Lys Leu
35      40      45
Leu Gln Asp Ala Gly His Thr Val Thr Ile Leu Glu Ala Asn Asp Arg
50      55      60
Val Gly Gly Arg Val Glu Thr Tyr Arg Asn Glu Lys Glu Gly Trp Tyr
65      70      75      80
Ala Glu Met Gly Ala Met Arg Ile Pro Ser Ser His Arg Ile Val Gln
85      90      95
Trp Phe Val Lys Lys Leu Gly Val Glu Met Asn Glu Phe Val Met Thr
100     105     110
Asp Asp Asn Thr Phe Tyr Leu Val Asn Gly Val Arg Glu Arg Thr Tyr
115     120     125
Val Val Gln Glu Asn Pro Asp Val Leu Lys Tyr Asn Val Ser Glu Ser
130     135     140
Glu Lys Gly Ile Ser Ala Asp Asp Leu Leu Asp Arg Ala Leu Gln Lys
145     150     155     160
Val Lys Glu Glu Val Glu Ala Asn Gly Cys Lys Ala Ala Leu Glu Lys
165     170     175
Tyr Asp Arg Tyr Ser Val Lys Glu Tyr Leu Lys Glu Glu Gly Gly Leu
180     185     190
Ser Pro Gly Ala Val Arg Met Ile Gly Asp Leu Leu Asn Glu Gln Ser
195     200     205
Leu Met Tyr Thr Ala Leu Ser Glu Met Ile Tyr Asp Gln Ala Asp Val
210     215     220
Asn Asp Ser Val Ser Tyr His Glu Val Thr Gly Gly Ser Asp Leu Leu
225     230     235     240
Pro Glu Ala Phe Leu Ser Val Leu Asp Val Pro Ile Leu Leu Asn Ser
245     250     255
Lys Val Lys His Ile Arg Gln Ser Asp Lys Gly Val Ile Val Ser Tyr
260     265     270
Gln Thr Gly Asn Glu Ser Ser Leu Met Asp Leu Ser Ala Asp Ile Val
275     280     285
Leu Val Thr Thr Thr Ala Lys Ala Ala Leu Phe Ile Asp Phe Asp Pro
290     295     300
Pro Leu Ser Ile Ser Lys Met Glu Ala Leu Arg Ser Val His Tyr Asp
305     310     315     320
Ser Ser Thr Lys Ile Leu Leu Thr Phe Arg Asp Lys Phe Trp Glu Asp
325     330     335
Asp Gly Ile Arg Gly Gly Lys Ser Ile Thr Asp Gly Pro Ser Arg Tyr
340     345     350
Ile Tyr Tyr Pro Ser His Ser Phe His Thr Asn Glu Thr Ile Gly Val
355     360     365
Leu Leu Ala Ser Tyr Thr Trp Ser Asp Glu Ser Leu Leu Phe Leu Gly
370     375     380
Ala Ser Asp Glu Glu Leu Lys Glu Leu Ala Leu Arg Asp Leu Ala Lys
385     390     395     400
Ile His Gly Glu Gln Val Trp Asp Lys Cys Thr Gly Val Ile Val Lys
405     410     415
Lys Trp Ser Ala Asp Pro Tyr Ser Leu Gly Ala Phe Ala Leu Phe Thr
420     425     430
Pro Tyr Gln His Leu Glu Tyr Ala Gln Glu Leu Phe Ser Ser Glu Gly
435     440     445
Arg Val His Phe Ala Gly Glu His Thr Ala Phe Pro His Ala Trp Ile
450     455     460
Glu Thr Ser Met Lys Ser Ala Ile Arg Ala Ala Thr Asn Ile Asn Lys
465     470     475     480

```

Val Ala Asn Glu Glu Ser Thr Ile Glu His Thr Lys Asp Glu Leu
 485 490 495

<210> 5
 <211> 786
 <212> DNA
 <213> Bacillus subtilis

<220>
 <221> CDS
 <222> (1)..(783)

<400> 5
 atg tat ccg gat tta aaa gga aaa gtc gtc gct att aca gga gct gct 48
 Met Tyr Pro Asp Leu Lys Gly Lys Val Val Ala Ile Thr Gly Ala Ala
 1 5 10 15
 tca ggg ctc gga aag gcg atg gcc att cgc ttc ggc aag gag cag gca 96
 Ser Gly Leu Gly Lys Ala Met Ala Ile Arg Phe Gly Lys Glu Gln Ala
 20 25 30
 aaa gtg gtt atc aac tat tat agt aat aaa caa gat ccg aac gag gta 144
 Lys Val Val Ile Asn Tyr Tyr Ser Asn Lys Gln Asp Pro Asn Glu Val
 35 40 45
 aaa gaa gag gtc atc aag gcg ggc ggt gaa gct gtt gtc gtc caa gga 192
 Lys Glu Glu Val Ile Lys Ala Gly Gly Glu Ala Val Val Val Gln Gly
 50 55 60
 gat gtc acg aaa gag gaa gat gta aaa aat atc gtg caa acg gca att 240
 Asp Val Thr Lys Glu Glu Asp Val Lys Asn Ile Val Gln Thr Ala Ile
 65 70 75 80
 aag gag ttc ggc aca ctc gat att atg att aat aat gcc ggt ctt gca 288
 Lys Glu Phe Gly Thr Leu Asp Ile Met Ile Asn Asn Ala Gly Leu Ala
 85 90 95
 aat cct gtg cca tct cac gaa atg ccg ctc aag gat tgg gat aaa gtc 336
 Asn Pro Val Pro Ser His Glu Met Pro Leu Lys Asp Trp Asp Lys Val
 100 105 110
 atc ggc acg aac tta acg ggt gcc ttt tta gga agc cgt gaa gcg att 384
 Ile Gly Thr Asn Leu Thr Gly Ala Phe Leu Gly Ser Arg Glu Ala Ile
 115 120 125
 aaa tat ttc gta gaa aac gat atc aag gga aat gtc att aac atg tcc 432
 Lys Tyr Phe Val Glu Asn Asp Ile Lys Gly Asn Val Ile Asn Met Ser
 130 135 140
 agt gtg cac gaa gtg att cct tgg ccg tta ttt gtc cac tat gcg gca 480
 Ser Val His Glu Val Ile Pro Trp Pro Leu Phe Val His Tyr Ala Ala
 145 150 155 160
 agt aaa ggc ggg ata aag ctg atg aca gaa aca tta gcg ttg gaa tac 528
 Ser Lys Gly Gly Ile Lys Leu Met Thr Glu Thr Leu Ala Leu Glu Tyr
 165 170 175
 gcg ccg aag ggc att cgc gtc aat aat att ggg cca ggt gcg atc aac 576
 Ala Pro Lys Gly Ile Arg Val Asn Asn Ile Gly Pro Gly Ala Ile Asn
 180 185 190
 acg cca atc aat gct gaa aaa ttc gct gac cct aaa cag aaa gct gat 624
 Thr Pro Ile Asn Ala Glu Lys Phe Ala Asp Pro Lys Gln Lys Ala Asp
 195 200 205
 gta gaa agc atg att cca atg gga tat atc ggc gaa ccg gag gag atc 672
 Val Glu Ser Met Ile Pro Met Gly Tyr Ile Gly Glu Pro Glu Glu Ile
 210 215 220
 gcc gca gta gca gcc tgg ctt gct tcg aag gaa gcc agc tac gtc aca 720
 Ala Ala Val Ala Ala Trp Leu Ala Ser Lys Glu Ala Ser Tyr Val Thr
 225 230 235 240
 ggc atc acg tta ttc gcg gac ggc ggt atg aca caa tat cct tca ttc 768
 Gly Ile Thr Leu Phe Ala Asp Gly Gly Met Thr Gln Tyr Pro Ser Phe
 245 250 255
 cag gca ggc cgc ggt taa 786
 Gln Ala Gly Arg Gly

260

<210> 6
 <211> 261
 <212> PRT
 <213> Bacillus subtilis

<400> 6
 Met Tyr Pro Asp Leu Lys Gly Lys Val Val Ala Ile Thr Gly Ala Ala
 1 5 10 15
 Ser Gly Leu Gly Lys Ala Met Ala Ile Arg Phe Gly Lys Glu Gln Ala
 20 25 30
 Lys Val Val Ile Asn Tyr Tyr Ser Asn Lys Gln Asp Pro Asn Glu Val
 35 40 45
 Lys Glu Glu Val Ile Lys Ala Gly Gly Glu Ala Val Val Val Gln Gly
 50 55 60
 Asp Val Thr Lys Glu Glu Asp Val Lys Asn Ile Val Gln Thr Ala Ile
 65 70 75 80
 Lys Glu Phe Gly Thr Leu Asp Ile Met Ile Asn Asn Ala Gly Leu Ala
 85 90 95
 Asn Pro Val Pro Ser His Glu Met Pro Leu Lys Asp Trp Asp Lys Val
 100 105 110
 Ile Gly Thr Asn Leu Thr Gly Ala Phe Leu Gly Ser Arg Glu Ala Ile
 115 120 125
 Lys Tyr Phe Val Glu Asn Asp Ile Lys Gly Asn Val Ile Asn Met Ser
 130 135 140
 Ser Val His Glu Val Ile Pro Trp Pro Leu Phe Val His Tyr Ala Ala
 145 150 155 160
 Ser Lys Gly Gly Ile Lys Leu Met Thr Glu Thr Leu Ala Leu Glu Tyr
 165 170 175
 Ala Pro Lys Gly Ile Arg Val Asn Asn Ile Gly Pro Gly Ala Ile Asn
 180 185 190
 Thr Pro Ile Asn Ala Glu Lys Phe Ala Asp Pro Lys Gln Lys Ala Asp
 195 200 205
 Val Glu Ser Met Ile Pro Met Gly Tyr Ile Gly Glu Pro Glu Glu Ile
 210 215 220
 Ala Ala Val Ala Ala Trp Leu Ala Ser Lys Glu Ala Ser Tyr Val Thr
 225 230 235 240
 Gly Ile Thr Leu Phe Ala Asp Gly Gly Met Thr Gln Tyr Pro Ser Phe
 245 250 255
 Gln Ala Gly Arg Gly
 260

<210> 7
 <211> 1230
 <212> DNA
 <213> Pseudomonas putida

<220>
 <221> CDS
 <222> (1)..(1227)

<400> 7
 atg ccc ttt cgc cgt acc ctt ctg gct gca tcc ctg gca ctt ctg atc 48
 Met Pro Phe Arg Arg Thr Leu Leu Ala Ala Ser Leu Ala Leu Leu Ile
 1 5 10 15
 acc gga cag gcc ccc ctg tat gcg gca cca ccg ttg tcg atg gac aac 96
 Thr Gly Gln Ala Pro Leu Tyr Ala Ala Pro Pro Leu Ser Met Asp Asn
 20 25 30
 ggc acc aac acc ctg acc gtg caa aac agc aat gcc tgg gtc gaa gtc 144
 Gly Thr Asn Thr Leu Thr Val Gln Asn Ser Asn Ala Trp Val Glu Val

JPOXMLDOC01-seq1.app

	35		40		45												
agc	gcc	agc	gcc	ctg	cag	cac	aac	atc	cgc	acg	ctg	cag	gcc	gag	ctg		192
Ser	Ala	Ser	Ala	Leu	Gln	His	Asn	Ile	Arg	Thr	Leu	Gln	Ala	Glu	Leu		
	50						55				60						
gcc	ggc	aag	tcc	aag	ctg	tgc	gcc	gtg	ctc	aag	gcc	gat	gcc	tat	ggc		240
Ala	Gly	Lys	Ser	Lys	Leu	Cys	Ala	Val	Leu	Lys	Ala	Asp	Ala	Tyr	Gly		
	65				70					75					80		
cac	ggt	atc	ggc	ctg	gta	atg	cca	tcg	atc	atc	gcc	caa	ggc	gtg	ccc		288
His	Gly	Ile	Gly	Leu	Val	Met	Pro	Ser	Ile	Ile	Ala	Gln	Gly	Val	Pro		
				85					90					95			
tgc	gtg	gcg	gtg	gcc	agc	aac	gag	gag	gcc	cgc	gtg	gtc	cgc	gcc	agt		336
Cys	Val	Ala	Val	Ala	Ser	Asn	Glu	Glu	Ala	Arg	Val	Val	Arg	Ala	Ser		
				100				105				110					
ggc	ttc	acc	ggg	caa	ctg	gtg	cgg	gta	cgc	ctg	gcc	agc	ctc	agc	gag		384
Gly	Phe	Thr	Gly	Gln	Leu	Val	Arg	Val	Arg	Leu	Ala	Ser	Leu	Ser	Glu		
		115					120					125					
ctg	gaa	gat	ggc	ttg	cag	tac	gac	atg	gaa	gag	ctg	gtg	ggc	agc	gcg		432
Leu	Glu	Asp	Gly	Leu	Gln	Tyr	Asp	Met	Glu	Glu	Leu	Val	Gly	Ser	Ala		
	130					135					140						
gaa	ttt	gcc	cgc	cag	gcc	gat	gcc	atc	gcc	gcg	cgc	cat	ggc	aag	acc		480
Glu	Phe	Ala	Arg	Gln	Ala	Asp	Ala	Ile	Ala	Ala	Arg	His	Gly	Lys	Thr		
	145				150				155						160		
ttg	cgc	att	cac	atg	gcg	ctc	aac	tcc	agc	ggc	atg	agc	cgc	aac	ggg		528
Leu	Arg	Ile	His	Met	Ala	Leu	Asn	Ser	Ser	Gly	Met	Ser	Arg	Asn	Gly		
			165					170					175				
gtg	gag	atg	gcc	acc	tgg	tcc	ggc	cgt	ggc	gaa	gcg	ctg	cag	atc	acc		576
Val	Glu	Met	Ala	Thr	Trp	Ser	Gly	Arg	Gly	Glu	Ala	Leu	Gln	Ile	Thr		
			180				185					190					
gac	cag	aag	cac	ctc	aag	ctg	gtc	gcg	ctg	atg	acc	cac	ttc	gcc	gtg		624
Asp	Gln	Lys	His	Leu	Lys	Leu	Val	Ala	Leu	Met	Thr	His	Phe	Ala	Val		
	195						200				205						
gaa	gac	aag	gac	gat	gta	cgc	aag	ggc	ctg	gcg	gca	ttc	aac	gag	cag		672
Glu	Asp	Lys	Asp	Asp	Val	Arg	Lys	Gly	Leu	Ala	Ala	Phe	Asn	Glu	Gln		
	210				215				220								
acc	gac	tgg	ttg	atc	aag	cac	gcc	agg	ctg	gac	cgc	agc	aag	ctc	acc		720
Thr	Asp	Trp	Leu	Ile	Lys	His	Ala	Arg	Leu	Asp	Arg	Ser	Lys	Leu	Thr		
	225				230				235						240		
ctg	cac	gcc	gcc	aac	tcg	ttc	gct	acg	ctg	gaa	gtg	ccg	gaa	gcg	cgc		768
Leu	His	Ala	Ala	Asn	Ser	Phe	Ala	Thr	Leu	Glu	Val	Pro	Glu	Ala	Arg		
				245					250				255				
ctg	gac	atg	gta	cga	acg	ggt	ggc	gcg	ctg	ttc	ggc	gac	acc	gtg	ccg		816
Leu	Asp	Met	Val	Arg	Thr	Gly	Gly	Ala	Leu	Phe	Gly	Asp	Thr	Val	Pro		
	260						265					270					
gcg	cgc	acc	gag	tac	aaa	cgt	gcg	atg	cag	ttc	aaa	tcg	cac	gtg	gcg		864
Ala	Arg	Thr	Glu	Tyr	Lys	Arg	Ala	Met	Gln	Phe	Lys	Ser	His	Val	Ala		
	275					280					285						
gcg	gtg	cac	agc	tat	ccg	gcc	ggc	aac	acc	gtg	ggc	tat	gac	cgc	acc		912
Ala	Val	His	Ser	Tyr	Pro	Ala	Gly	Asn	Thr	Val	Gly	Tyr	Asp	Arg	Thr		
	290				295				300								
ttc	acc	ctg	gcc	cgt	gat	tcg	cgg	ctg	gcc	aac	att	acg	gtc	ggg	tac		960
Phe	Thr	Leu	Ala	Arg	Asp	Ser	Arg	Leu	Ala	Asn	Ile	Thr	Val	Gly	Tyr		
	305				310				315						320		
tcc	gat	ggc	tac	cgc	cgg	gta	ttc	acc	aac	aag	ggc	cat	gtg	ctg	atc		1008
Ser	Asp	Gly	Tyr	Arg	Arg	Val	Phe	Thr	Asn	Lys	Gly	His	Val	Leu	Ile		
				325					330				335				
aac	ggc	cac	cgt	gtg	ccg	gtc	gtg	ggc	aag	gtg	tcg	atg	aac	acg	ctg		1056
Asn	Gly	His	Arg	Val	Pro	Val	Val	Gly	Lys	Val	Ser	Met	Asn	Thr	Leu		
			340				345					350					
atg	gtc	gat	gtc	acc	gac	ttc	cct	gat	gtg	aag	ggg	ggt	aac	gaa	gtg		1104
Met	Val	Asp	Val	Thr	Asp	Phe	Pro	Asp	Val	Lys	Gly	Gly	Asn	Glu	Val		
	355					360						365					
gtg	ctg	ttc	ggc	aag	cag	gcc	ggg	ggc	gaa	atc	acc	cag	gcc	gag	atg		1152
Val	Leu	Phe	Gly	Lys	Gln	Ala	Gly	Gly	Glu	Ile	Thr	Gln	Ala	Glu	Met		
	370					375					380						

```

gaa gaä atc aac ggc gcg ttg ctc gcc gat ttg tac acc gta tgg ggc      1200
Glu Glu Ile Asn Gly Ala Leu Leu Ala Asp Leu Tyr Thr Val Trp Gly
385                               390                               395           400
aat tcc aac ccg aag ata ctc gtc gac tga                                1230
Asn Ser Asn Pro Lys Ile Leu Val Asp
          405

```

```
<210> 8
<211> 409
<212> PRT
<213> Pseudomonas putida
```

<400>	8														
Met 1	Pro	Phe	Arg	Arg 5	Thr	Leu	Leu	Ala	Ala 10	Ser	Leu	Ala	Leu 15	Leu	Ile
Thr	Gly	Gln	Ala 20	Pro	Leu	Tyr	Ala	Ala 25	Pro	Pro	Leu	Ser	Met 30	Asp	Asn
Gly	Thr	Asn 35	Thr	Leu	Thr	Val	Gln 40	Asn	Ser	Asn	Ala	Trp 45	Val	Glu	Val
Ser	Ala	Ser	Ala	Leu	Gln	His 55	Asn	Ile	Arg	Thr	Leu	Gln	Ala	Glu	Leu
Ala 65	Gly	Lys	Ser	Lys	Leu	Cys 70	Ala	Val	Leu	Lys 75	Ala	Asp	Ala	Tyr	Gly 80
His	Gly	Ile	Gly	Leu 85	Val	Met	Pro	Ser	Ile 90	Ile	Ala	Gln	Gly	Val 95	Pro
Cys	Val	Ala	Val	Ala 100	Ser	Asn	Glu	Glu 105	Ala	Arg	Val	Val	Arg	Ala	Ser
Gly	Phe	Thr 115	Gly	Gln	Leu	Val	Arg 120	Val	Arg	Leu	Ala	Ser	Leu 125	Ser	Glu
Leu	Glu 130	Asp	Gly	Leu	Gln	Tyr 135	Asp	Met	Glu	Glu	Leu	Val	Gly	Ser	Ala
Glu 145	Phe	Ala	Arg	Gln	Ala 150	Asp	Ala	Ile	Ala	Ala 155	Arg	His	Gly	Lys	Thr 160
Leu	Arg	Ile	His	Met 165	Ala	Leu	Asn	Ser	Ser 170	Gly	Met	Ser	Arg	Asn	Gly 175
Val	Glu	Met	Ala 180	Thr	Trp	Ser	Gly	Arg 185	Gly	Glu	Ala	Leu	Gln	Ile	Thr
Asp	Gln	Lys 195	His	Leu	Lys	Leu	Val 200	Ala	Leu	Met	Thr	His	Phe	Ala	Val
Glu	Asp 210	Lys	Asp	Asp	Val	Arg	Lys 215	Gly	Leu	Ala	Ala	Phe	Asn	Glu	Gln
Thr 225	Asp	Trp	Leu	Ile	Lys	His	Ala 230	Arg	Leu	Asp	Arg	Ser	Lys	Leu	Thr 240
Leu	His	Ala	Ala	Asn 245	Ser	Phe	Ala	Thr	Leu	Glu	Val	Pro	Glu	Ala	Arg
Leu	Asp	Met	Val 260	Arg	Thr	Gly	Gly 265	Ala	Leu	Phe	Gly	Asp	Thr	Val	Pro
Ala	Arg	Thr	Glu	Tyr	Lys	Arg	Ala 280	Met	Gln	Phe	Lys	Ser	His	Val	Ala
Ala	Val	His	Ser	Tyr	Pro	Ala	Gly 295	Asn	Thr	Val	Gly	Tyr	Asp	Arg	Thr
Phe 305	Thr	Leu	Ala	Arg	Asp	Ser	Arg	Leu	Ala	Asn 315	Ile	Thr	Val	Gly	Tyr 320
Ser	Asp	Gly	Tyr	Arg 325	Arg	Val	Phe	Thr	Asn	Lys 330	Gly	His	Val	Leu	Ile
Asn	Gly	His	Arg	Val 340	Pro	Val	Val	Gly	Lys	Val	Ser	Met	Asn	Thr	Leu
Met	Val	Asp	Val	Thr	Asp	Phe	Pro 360	Asp	Val	Lys	Gly	Gly	Asn	Glu	Val
Val	Leu	Phe	Gly	Lys	Gln	Ala	Gly 375	Gly	Glu	Ile	Thr	Gln	Ala	Glu	Met
Glu 385	Glu	Ile	Asn	Gly	Ala	Leu	Leu	Ala	Asp	Leu	Tyr	Thr	Val	Trp	Gly 400

Asn Ser Asn Pro Lys Ile Leu Val Asp
405

```

<210> 9
<211> 36
<212> DNA
<213> Artificial sequence

<220>
<223> PCR primer

<400> 9
gcgaattcat gtccgcacct tccaccagca ccgttg 36

<210> 10
<211> 36
<212> DNA
<213> Artificial sequence

<220>
<223> PCR primer

<400> 10
gctctagatc agccaagcag ctctttcagg ccttgc 36

<210> 11
<211> 41
<212> DNA
<213> Artificial sequence

<220>
<223> PCR primer

<400> 11
gcgaattcat ggagcacttg gctgattgct tggaagataa g 41

<210> 12
<211> 39
<212> DNA
<213> Artificial sequence

<220>
<223> PCR primer

<400> 12
gctctagatt acaactcatc ttttgtgtgt tcaatgggtg 39

<210> 13
<211> 37
<212> DNA
<213> Artificial sequence

<220>
<223> PCR primer

<400> 13
ggggaattca tgtatccgga tttaaaagga aaagtcg 37

```

```

<210> 14
<211> 28
<212> DNA
<213> Artificial sequence

<220>
<223> PCR primer

<400> 14
ggggtctaga ttaaccgcgg cctgcctg 28

<210> 15
<211> 36
<212> DNA
<213> Artificial sequence

<220>
<223> PCR primer

<400> 15
gcgaattcat gccctttcgc cgtacccttc tggctg 36

<210> 16
<211> 39
<212> DNA
<213> Artificial sequence

<220>
<223> PCR primer

<400> 16
gctctagatc agtcgacgag tatcttcggg ttggaattg 39

<210> 17
<211> 1113
<212> DNA
<213> Flavobacterium johnsoniae NBRC14942

<220>
<221> CDS
<222> (1)..(1110)

<400> 17
atg aaa tca caa tca tta att gaa gat gag ata cca gta aaa gaa aac 48
Met Lys Ser Gln Ser Leu Ile Glu Asp Glu Ile Pro Val Lys Glu Asn
1 5 10 15
tat gct tat caa att cct aca agc ccg ctg ata gtg gag gtt acg cct 96
Tyr Ala Tyr Gln Ile Pro Thr Ser Pro Leu Ile Val Glu Val Thr Pro
20 25 30
cag gaa aga aac att ttg tct aat gtg ggc gct ctg ctg gaa aag gca 144
Gln Glu Arg Asn Ile Leu Ser Asn Val Gly Ala Leu Leu Glu Lys Ala
35 40 45
ttt aag agc tat gaa aac cca gat tat ata gaa gcg ctt cat ctg tat 192
Phe Lys Ser Tyr Glu Asn Pro Asp Tyr Ile Glu Ala Leu His Leu Tyr
50 55 60
tct ttt cag ctt ctt cca gaa aga ata gcc aga att tta agc cgt ttt 240
Ser Phe Gln Leu Leu Pro Glu Arg Ile Ala Arg Ile Leu Ser Arg Phe
65 70 75 80
gga aca gat ttc tca gct gat cag tat ggc gct att att ttt aga ggt 288
Gly Thr Asp Phe Ser Ala Asp Gln Tyr Gly Ala Ile Ile Phe Arg Gly
85 90 95
ctt ctt gaa gtt gat cag gat cat ctg gga cca act cct gcg aat tgg 336

```

JPOXMLDOC01-seql.app

```

Leu Leu Glu Val Asp Gln Asp His Leu Gly Pro Thr Pro Ala Asn Trp
100 105 110
cag agc gct gat tac tca aaa ctc aat aaa tac ggc ttt att tgt tcc 384
Gln Ser Ala Asp Tyr Ser Lys Leu Asn Lys Tyr Gly Phe Ile Cys Ser
115 120 125
ttg ctg cat ggt gca gtt cct tca aaa cca gta caa tat tat gcg cag 432
Leu Leu His Gly Ala Val Pro Ser Lys Pro Val Gln Tyr Tyr Ala Gln
130 135 140
aga aag ggc ggg gga att ctt cat gct gtt att cca gat gag aaa atg 480
Arg Lys Gly Gly Gly Ile Leu His Ala Val Ile Pro Asp Glu Lys Met
145 150 155 160
gca gct acg caa aca ggt tcg gga tca aaa aca aat ttg tat gtt cat 528
Ala Ala Thr Gln Thr Gly Ser Gly Ser Lys Thr Asn Leu Tyr Val His
165 170 175
aca gaa gat gct ttt ctt tta cat cag gct gat ttt tta agt ttt cta 576
Thr Glu Asp Ala Phe Leu Leu His Gln Ala Asp Phe Leu Ser Phe Leu
180 185 190
tat ctg cga aat gaa gaa aga gtt cct tct aca ctt tac tca gta agg 624
Tyr Leu Arg Asn Glu Glu Arg Val Pro Ser Thr Leu Tyr Ser Val Arg
195 200 205
tcg cat ggt aag gtg aat aag ata atg gaa aag ctt ttt gat cca att 672
Ser His Gly Lys Val Asn Lys Ile Met Glu Lys Leu Phe Asp Pro Ile
210 215 220
tat caa tgt cct aaa gat gct aat tat cag gaa gaa att aat gat ggt 720
Tyr Gln Cys Pro Lys Asp Ala Asn Tyr Gln Glu Glu Ile Asn Asp Gly
225 230 235 240
ccg ctg gct tct gtt tta tat gga aat aaa aag ctg cct ttt att aga 768
Pro Leu Ala Ser Val Leu Tyr Gly Asn Lys Lys Leu Pro Phe Ile Arg
245 250 255
ttt gat gca gca gag cag ata ttt aat gaa aac gcc gga cag act ccc 816
Phe Asp Ala Ala Glu Gln Ile Phe Asn Glu Asn Ala Gly Gln Thr Pro
260 265 270
gaa gct ctt tac aat tta act gaa ttt tgg aat gaa gct aaa gag ttg 864
Glu Ala Leu Tyr Asn Leu Thr Glu Phe Trp Asn Glu Ala Lys Glu Leu
275 280 285
att aat agt gat tat atc cca gat tct ggt gat gtt ata ttt gta aat 912
Ile Asn Ser Asp Tyr Ile Pro Asp Ser Gly Asp Val Ile Phe Val Asn
290 295 300
aat cat ttg tgt gct cac gga aga agt gct ttt aca gca ggg cag aaa 960
Asn His Leu Cys Ala His Gly Arg Ser Ala Phe Thr Ala Gly Gln Lys
305 310 315 320
gag gag aat ggt aag ctt gtg cca tgt gag aga cga caa atg tta aga 1008
Glu Glu Asn Gly Lys Leu Val Pro Cys Glu Arg Arg Gln Met Leu Arg
325 330 335
atg atg agc aaa acc agt cta att cat ata aga tca atg aca cat acc 1056
Met Met Ser Lys Thr Ser Leu Ile His Ile Arg Ser Met Thr His Thr
340 345 350
gat gat ccg tat ttt gtt atg gaa gaa cat tta gga aaa gtt ttt gat 1104
Asp Asp Pro Tyr Phe Val Met Glu Glu His Leu Gly Lys Val Phe Asp
355 360 365

cag gct taa 1113
Gln Ala
370

```

<210> 18

<211> 370

<212> PRT

<213> Flavobacterium johnsoniae NBRC14942

<400> 18

Met Lys Ser Gln Ser Leu Ile Glu Asp Glu Ile Pro Val Lys Glu Asn

1

5

10

15

Tyr Ala Tyr Gln Ile Pro Thr Ser Pro Leu Ile Val Glu Val Thr Pro

JPOXMLDOC01-seq1.app

			20					25					30				
Gln	Glu	Arg	Asn	Ile	Leu	Ser	Asn	Val	Gly	Ala	Leu	Leu	Glu	Lys	Ala		
		35					40					45					
Phe	Lys	Ser	Tyr	Glu	Asn	Pro	Asp	Tyr	Ile	Glu	Ala	Leu	His	Leu	Tyr		
	50					55					60						
Ser	Phe	Gln	Leu	Leu	Pro	Glu	Arg	Ile	Ala	Arg	Ile	Leu	Ser	Arg	Phe		
65					70					75					80		
Gly	Thr	Asp	Phe	Ser	Ala	Asp	Gln	Tyr	Gly	Ala	Ile	Ile	Phe	Arg	Gly		
			85						90					95			
Leu	Leu	Glu	Val	Asp	Gln	Asp	His	Leu	Gly	Pro	Thr	Pro	Ala	Asn	Trp		
		100						105					110				
Gln	Ser	Ala	Asp	Tyr	Ser	Lys	Leu	Asn	Lys	Tyr	Gly	Phe	Ile	Cys	Ser		
	115					120						125					
Leu	Leu	His	Gly	Ala	Val	Pro	Ser	Lys	Pro	Val	Gln	Tyr	Tyr	Ala	Gln		
	130					135					140						
Arg	Lys	Gly	Gly	Gly	Ile	Leu	His	Ala	Val	Ile	Pro	Asp	Glu	Lys	Met		
145					150					155					160		
Ala	Ala	Thr	Gln	Thr	Gly	Ser	Gly	Ser	Lys	Thr	Asn	Leu	Tyr	Val	His		
			165					170						175			
Thr	Glu	Asp	Ala	Phe	Leu	Leu	His	Gln	Ala	Asp	Phe	Leu	Ser	Phe	Leu		
	180							185					190				
Tyr	Leu	Arg	Asn	Glu	Glu	Arg	Val	Pro	Ser	Thr	Leu	Tyr	Ser	Val	Arg		
	195					200						205					
Ser	His	Gly	Lys	Val	Asn	Lys	Ile	Met	Glu	Lys	Leu	Phe	Asp	Pro	Ile		
	210				215						220						
Tyr	Gln	Cys	Pro	Lys	Asp	Ala	Asn	Tyr	Gln	Glu	Glu	Ile	Asn	Asp	Gly		
225					230					235					240		
Pro	Leu	Ala	Ser	Val	Leu	Tyr	Gly	Asn	Lys	Lys	Leu	Pro	Phe	Ile	Arg		
			245					250						255			
Phe	Asp	Ala	Ala	Glu	Gln	Ile	Phe	Asn	Glu	Asn	Ala	Gly	Gln	Thr	Pro		
	260							265					270				
Glu	Ala	Leu	Tyr	Asn	Leu	Thr	Glu	Phe	Trp	Asn	Glu	Ala	Lys	Glu	Leu		
	275					280						285					
Ile	Asn	Ser	Asp	Tyr	Ile	Pro	Asp	Ser	Gly	Asp	Val	Ile	Phe	Val	Asn		
	290					295					300						
Asn	His	Leu	Cys	Ala	His	Gly	Arg	Ser	Ala	Phe	Thr	Ala	Gly	Gln	Lys		
305					310					315					320		
Glu	Glu	Asn	Gly	Lys	Leu	Val	Pro	Cys	Glu	Arg	Arg	Gln	Met	Leu	Arg		
			325						330					335			
Met	Met	Ser	Lys	Thr	Ser	Leu	Ile	His	Ile	Arg	Ser	Met	Thr	His	Thr		
		340						345					350				
Asp	Asp	Pro	Tyr	Phe	Val	Met	Glu	Glu	His	Leu	Gly	Lys	Val	Phe	Asp		
	355						360					365					
Gln	Ala																
	370																

<210> 19
 <211> 996
 <212> DNA
 <213> Kineococcus radiotolerans NBRC101839

<220>
 <221> CDS
 <222> (1)..(993)

<400>	19																
atg	tcc	tcg	ctg	ttc	ctc	gac	tcc	tcc	gcc	cac	gtg	ccg	acc	ctg	ttc		48
Met	Ser	Ser	Leu	Phe	Leu	Asp	Ser	Ser	Ala	His	Val	Pro	Thr	Leu	Phe		
1			5					10				15					
gag	ctg	ccc	gcg	ccc	cag	cgg	gcc	gcg	ctg	gcc	gcg	ctg	ggc	gcg	cgc		96
Glu	Leu	Pro	Ala	Pro	Gln	Arg	Ala	Ala	Leu	Ala	Ala	Leu	Gly	Ala	Arg		
		20					25					30					
ctg	acc	gcg	gac	ccg	gtg	acc	gag	ccc	gac	gcc	ttc	ggc	cgc	cag	gcg		144

JPOXMLDOC01-seq1.app

Leu	Thr	Ala	Asp	Pro	Val	Thr	Glu	Pro	Asp	Ala	Phe	Gly	Arg	Gln	Ala		
		35					40					45					
cgc	ctg	ctg	gcc	cgc	gaa	ctg	tcc	gtc	gag	gtc	acc	gag	gcc	ctg	tgg	192	
Arg	Leu	Leu	Ala	Arg	Glu	Leu	Ser	Val	Glu	Val	Thr	Glu	Ala	Leu	Trp		
		50					55					60					
gcg	ttc	gag	gaa	cgg	gga	tcg	gac	tcc	ggg	gtc	ctc	gtc	ctg	cgc	ggc	240	
Ala	Phe	Glu	Glu	Arg	Gly	Ser	Asp	Ser	Gly	Val	Leu	Val	Leu	Arg	Gly		
		65					70					75			80		
ctg	gag	gtc	ggt	gag	ctg	ccg	ccc	acc	ccg	gcc	gac	aac	acc	ggc	ggg	288	
Leu	Glu	Val	Gly	Glu	Leu	Pro	Pro	Thr	Pro	Ala	Asp	Asn	Thr	Gly	Gly		
				85					90					95			
atc	ggc	ggg	cgc	acc	ctg	ctc	gcc	cgc	cag	cag	gcg	atc	gtc	agc	cac	336	
Ile	Gly	Gly	Arg	Thr	Leu	Leu	Ala	Arg	Gln	Gln	Ala	Ile	Val	Ser	His		
			100					105					110				
gcg	ctg	ggg	cac	atg	gtc	ggc	tac	gcc	gcc	gag	ggc	cac	ggg	cac	ctc	384	
Ala	Leu	Gly	His	Met	Val	Gly	Tyr	Ala	Ala	Glu	Gly	His	Gly	His	Leu		
		115					120					125					
ctg	cag	gac	atg	gtc	ccc	aac	gcc	agg	ctc	gcc	gcg	acc	cag	cag	tcg	432	
Leu	Gln	Asp	Met	Val	Pro	Asn	Ala	Arg	Leu	Ala	Ala	Thr	Gln	Gln	Ser		
		130					135					140					
cag	ggc	tcc	cgg	gtg	gag	ctg	gag	gcg	cac	acc	gag	cag	tgc	ttc	tcc	480	
Gln	Gly	Ser	Arg	Val	Glu	Leu	Glu	Ala	His	Thr	Glu	Gln	Cys	Phe	Ser		
		145				150			155					160			
gac	ctg	cgc	ccc	gac	tac	gtc	gtc	ctg	ggc	tgc	ctg	cgc	ggg	gac	gcc	528	
Asp	Leu	Arg	Pro	Asp	Tyr	Val	Val	Leu	Gly	Cys	Leu	Arg	Gly	Asp	Ala		
			165					170					175				
gac	gcc	gcc	acc	tac	gcg	ttc	cgc	gcc	ctg	gac	ctg	ctg	gcc	cac	gtg	576	
Asp	Ala	Ala	Thr	Tyr	Ala	Phe	Arg	Ala	Leu	Asp	Leu	Leu	Ala	His	Val		
			180				185					190					
gac	ccc	acc	gac	gtc	atg	gag	ctg	ttc	cgg	ccg	ctg	tgg	acg	acg	ctg	624	
Asp	Pro	Thr	Asp	Val	Met	Glu	Leu	Phe	Arg	Pro	Leu	Trp	Thr	Thr	Leu		
		195					200					205					
gtc	gac	gag	tcc	ttc	gcc	gac	ttc	ctc	gac	acc	cgc	gag	gtg	cgc	ggg	672	
Val	Asp	Glu	Ser	Phe	Ala	Asp	Phe	Leu	Asp	Thr	Arg	Glu	Val	Arg	Gly		
		210				215					220						
ccg	ttc	tcc	atc	ctc	tcc	ggc	gac	gtc	gac	gac	ccg	acg	atg	ctc	gtc	720	
Pro	Phe	Ser	Ile	Leu	Ser	Gly	Asp	Val	Asp	Asp	Pro	Thr	Met	Leu	Val		
		225			230				235					240			
gac	cag	gac	ctc	atg	cac	ggc	atc	acc	aag	cac	gcc	cag	gcc	ctg	ctg	768	
Asp	Gln	Asp	Leu	Met	His	Gly	Ile	Thr	Lys	His	Ala	Gln	Ala	Leu	Leu		
			245					250				255					
gag	cgc	gtg	ctg	gag	atc	tac	gtc	gcc	cac	cgc	cac	gcc	gtc	gtc	ctc	816	
Glu	Arg	Val	Leu	Glu	Ile	Tyr	Val	Ala	His	Arg	His	Ala	Val	Val	Leu		
		260					265					270					
cag	ccc	ggg	gac	gtg	ctg	ctg	ctg	gac	aac	ctg	cgc	gcc	atg	cac	ggc	864	
Gln	Pro	Gly	Asp	Val	Leu	Leu	Leu	Asp	Asn	Leu	Arg	Ala	Met	His	Gly		
		275				280					285						
cgc	tcg	ccg	ttc	gcc	ccg	cgc	ttc	gac	ggc	acc	gac	ccg	ttc	atc	tcc	912	
Arg	Ser	Pro	Phe	Ala	Pro	Arg	Phe	Asp	Gly	Thr	Asp	Arg	Phe	Ile	Ser		
		290				295					300						
cgg	ggt	ttc	gtc	gtc	cgc	gac	ctg	cgc	cgc	tcc	cgc	ttc	gcc	cgc	ccc	960	
Arg	Gly	Phe	Val	Val	Arg	Asp	Leu	Arg	Arg	Ser	Arg	Phe	Ala	Arg	Pro		
		305			310				315						320		
ggc	ggg	aac	cgc	gtc	gtg	cag	gcc	agc	ttc	agc	tga					996	
Gly	Gly	Asn	Arg	Val	Val	Gln	Ala	Ser	Phe	Ser							
			325					330									

<210> 20
 <211> 331
 <212> PRT
 <213> Kineococcus radiotolerans NBRC101839

<400> 20

JPOXMLDOC01-seq1.app

```

Met Ser Ser Leu Phe Leu Asp Ser Ser Ala His Val Pro Thr Leu Phe
1      5      10      15
Glu Leu Pro Ala Pro Gln Arg Ala Ala Leu Ala Ala Leu Gly Ala Arg
20      25      30
Leu Thr Ala Asp Pro Val Thr Glu Pro Asp Ala Phe Gly Arg Gln Ala
35      40      45
Arg Leu Leu Ala Arg Glu Leu Ser Val Glu Val Thr Glu Ala Leu Trp
50      55      60
Ala Phe Glu Glu Arg Gly Ser Asp Ser Gly Val Leu Val Leu Arg Gly
65      70      75      80
Leu Glu Val Gly Glu Leu Pro Pro Thr Pro Ala Asp Asn Thr Gly Gly
85      90      95
Ile Gly Gly Arg Thr Leu Leu Ala Arg Gln Gln Ala Ile Val Ser His
100     105     110
Ala Leu Gly His Met Val Gly Tyr Ala Ala Glu Gly His Gly His Leu
115     120     125
Leu Gln Asp Met Val Pro Asn Ala Arg Leu Ala Ala Thr Gln Gln Ser
130     135     140
Gln Gly Ser Arg Val Glu Leu Glu Ala His Thr Glu Gln Cys Phe Ser
145     150     155     160
Asp Leu Arg Pro Asp Tyr Val Val Leu Gly Cys Leu Arg Gly Asp Ala
165     170     175
Asp Ala Ala Thr Tyr Ala Phe Arg Ala Leu Asp Leu Leu Ala His Val
180     185     190
Asp Pro Thr Asp Val Met Glu Leu Phe Arg Pro Leu Trp Thr Thr Leu
195     200     205
Val Asp Glu Ser Phe Ala Asp Phe Leu Asp Thr Arg Glu Val Arg Gly
210     215     220
Pro Phe Ser Ile Leu Ser Gly Asp Val Asp Asp Pro Thr Met Leu Val
225     230     235     240
Asp Gln Asp Leu Met His Gly Ile Thr Lys His Ala Gln Ala Leu Leu
245     250     255
Glu Arg Val Leu Glu Ile Tyr Val Ala His Arg His Ala Val Val Leu
260     265     270
Gln Pro Gly Asp Val Leu Leu Leu Asp Asn Leu Arg Ala Met His Gly
275     280     285
Arg Ser Pro Phe Ala Pro Arg Phe Asp Gly Thr Asp Arg Phe Ile Ser
290     295     300
Arg Gly Phe Val Val Arg Asp Leu Arg Arg Ser Arg Phe Ala Arg Pro
305     310     315     320
Gly Gly Asn Arg Val Val Gln Ala Ser Phe Ser
325     330

```

```

<210> 21
<211> 1149
<212> DNA
<213> Chitinophaga pinensis NBRC15968

```

```

<220>
<221> CDS
<222> (1)..(1146)

```

```

<400> 21
atg aga ccc tta gac gtg aca ccc aca att agc cca gga gcc cag gac      48
Met Arg Pro Leu Asp Val Thr Pro Thr Ile Ser Pro Gly Ala Gln Asp
1      5      10      15
ctt ccg cgc act atg cat ttt gct gct gaa cct cct tta cag cct ttg      96
Leu Pro Arg Thr Met His Phe Ala Ala Glu Pro Pro Leu Gln Pro Leu
20      25      30
ata ata gat att act gaa gaa gaa aaa ctg gaa att acc tat atc ggg      144
Ile Ile Asp Ile Thr Glu Glu Glu Lys Leu Glu Ile Thr Tyr Ile Gly
35      40      45
aaa aag cta aaa aga aag tat aaa agc tat gat gat ccc ggt ttt att      192

```

JPOXMLDOC01-seql.app

Lys	Lys	Leu	Lys	Arg	Lys	Tyr	Lys	Ser	Tyr	Asp	Asp	Pro	Gly	Phe	Ile	
50						55				60						
tca	atg	ctg	cac	tta	aat	gcc	tat	acg	ctg	cta	ccg	gag	cgt	ata	gca	240
Ser	Met	Leu	His	Leu	Asn	Ala	Tyr	Thr	Leu	Leu	Pro	Glu	Arg	Ile	Ala	
65					70					75					80	
aag	gtg	ctg	agt	aat	ttc	ggg	aca	gac	ttt	tcc	gac	cag	caa	tac	gga	288
Lys	Val	Leu	Ser	Asn	Phe	Gly	Thr	Asp	Phe	Ser	Asp	Gln	Gln	Tyr	Gly	
				85					90					95		
gct	gtc	gta	ttg	cgt	gga	ctg	ata	gaa	ata	ggg	cag	gat	gaa	tta	ggc	336
Ala	Val	Val	Leu	Arg	Gly	Leu	Ile	Glu	Ile	Gly	Gln	Asp	Glu	Leu	Gly	
			100					105					110			
cca	acc	cca	cgt	tcc	tgg	cag	gaa	acc	gac	cat	gaa	aag	att	atg	gaa	384
Pro	Thr	Pro	Arg	Ser	Trp	Gln	Glu	Thr	Asp	His	Glu	Lys	Ile	Met	Glu	
			115					120				125				
tat	ggc	ttc	att	tcc	tcc	tta	tta	cat	ggc	gct	gta	cca	tcc	aaa	ccc	432
Tyr	Gly	Phe	Ile	Ser	Ser	Leu	Leu	His	Gly	Ala	Val	Pro	Ser	Lys	Pro	
			130			135					140					
gtc	gag	tat	ttc	gcg	cag	cga	aaa	ggg	ggg	ggc	tta	atg	cac	gcg	att	480
Val	Glu	Tyr	Phe	Ala	Gln	Arg	Lys	Gly	Gly	Gly	Leu	Met	His	Ala	Ile	
			145		150				155						160	
att	cct	gat	gag	aat	atg	agc	ttt	aca	caa	aca	ggc	tca	ggg	tcc	cgt	528
Ile	Pro	Asp	Glu	Asn	Met	Ser	Phe	Thr	Gln	Thr	Gly	Ser	Gly	Ser	Arg	
			165					170						175		
aca	gat	ctt	ttt	gta	cat	aca	gaa	gat	gct	ttc	ctg	cat	aat	gcg	gct	576
Thr	Asp	Leu	Phe	Val	His	Thr	Glu	Asp	Ala	Phe	Leu	His	Asn	Ala	Ala	
			180					185					190			
gat	ttt	ctg	agt	ttt	ctt	ttc	ctg	cgg	aat	gaa	gaa	cgt	gtg	cct	tcc	624
Asp	Phe	Leu	Ser	Phe	Leu	Phe	Leu	Arg	Asn	Glu	Glu	Arg	Val	Pro	Ser	
			195					200					205			
acc	tta	tat	tct	atc	cgc	tct	cat	ggc	aga	ccg	gat	gcg	ata	tta	cag	672
Thr	Leu	Tyr	Ser	Ile	Arg	Ser	His	Gly	Arg	Pro	Asp	Ala	Ile	Leu	Gln	
			210			215					220					
gag	ctt	ttc	aag	cct	atc	tat	aag	tgt	ccg	aag	gat	gcg	aac	tat	gct	720
Glu	Leu	Phe	Lys	Pro	Ile	Tyr	Lys	Cys	Pro	Lys	Asp	Ala	Asn	Tyr	Ala	
			225		230					235					240	
tcc	gaa	gaa	gcc	ctg	gga	gat	gac	atc	cgt	act	tct	gtt	tta	tat	ggg	768
Ser	Glu	Glu	Ala	Leu	Gly	Asp	Asp	Ile	Arg	Thr	Ser	Val	Leu	Tyr	Gly	
			245					250						255		
agc	aga	tcc	gct	ccc	ttc	atg	cgc	ttt	gat	gct	gcg	gaa	cag	att	tat	816
Ser	Arg	Ser	Ala	Pro	Phe	Met	Arg	Phe	Asp	Ala	Ala	Glu	Gln	Ile	Tyr	
			260					265					270			
aat	gaa	gac	gcc	aat	cag	gat	cct	gaa	gct	tta	cat	aat	ctg	aaa	aga	864
Asn	Glu	Asp	Ala	Asn	Gln	Asp	Pro	Glu	Ala	Leu	His	Asn	Leu	Lys	Arg	
			275					280					285			
ttc	tgg	gaa	gag	gcg	cgc	aaa	ctg	ata	tat	aat	gac	ttc	gtt	cct	gag	912
Phe	Trp	Glu	Glu	Ala	Arg	Lys	Leu	Ile	Tyr	Asn	Asp	Phe	Val	Pro	Glu	
			290			295					300					
tca	ggg	gac	ctg	atc	ttt	gtg	aat	aat	cat	ctt	tgt	gcc	cat	ggc	cgg	960
Ser	Gly	Asp	Leu	Ile	Phe	Val	Asn	Asn	His	Leu	Cys	Ala	His	Gly	Arg	
			305		310					315					320	
aat	gct	ttc	ctg	gca	ggc	ttc	aga	gag	gaa	aat	ggg	cag	ctg	gta	aaa	1008
Asn	Ala	Phe	Leu	Ala	Gly	Phe	Arg	Glu	Glu	Asn	Gly	Gln	Leu	Val	Lys	
			325					330						335		
tgc	gaa	cgc	cgt	ctt	atg	tta	cgt	atg	atg	agc	aaa	acc	agc	ctg	att	1056
Cys	Glu	Arg	Arg	Leu	Met	Leu	Arg	Met	Met	Ser	Lys	Thr	Ser	Leu	Ile	
			340					345					350			
aac	atc	cgt	gaa	gta	acc	cac	ccc	gaa	aac	cct	tat	ctc	atc	atg	gaa	1104
Asn	Ile	Arg	Glu	Val	Thr	His	Pro	Glu	Asn	Pro	Tyr	Leu	Ile	Met	Glu	
			355				360						365			
gag	cac	tac	gga	aaa	gta	tat	agc	gct	cac	ctg	gca	aac	ctt	taa		1149
Glu	His	Tyr	Gly	Lys	Val	Tyr	Ser	Ala	His	Leu	Ala	Asn	Leu			
			370			375					380					

<210> 22
 <211> 382
 <212> PRT
 <213> Chitinophaga pinensis NBRC15968

<400> 22
 Met Arg Pro Leu Asp Val Thr Pro Thr Ile Ser Pro Gly Ala Gln Asp
 1 5 10 15
 Leu Pro Arg Thr Met His Phe Ala Ala Glu Pro Pro Leu Gln Pro Leu
 20 25 30
 Ile Ile Asp Ile Thr Glu Glu Glu Lys Leu Glu Ile Thr Tyr Ile Gly
 35 40 45
 Lys Lys Leu Lys Arg Lys Tyr Lys Ser Tyr Asp Asp Pro Gly Phe Ile
 50 55 60
 Ser Met Leu His Leu Asn Ala Tyr Thr Leu Leu Pro Glu Arg Ile Ala
 65 70 75 80
 Lys Val Leu Ser Asn Phe Gly Thr Asp Phe Ser Asp Gln Gln Tyr Gly
 85 90 95
 Ala Val Val Leu Arg Gly Leu Ile Glu Ile Gly Gln Asp Glu Leu Gly
 100 105 110
 Pro Thr Pro Arg Ser Trp Gln Glu Thr Asp His Glu Lys Ile Met Glu
 115 120 125
 Tyr Gly Phe Ile Ser Ser Leu Leu His Gly Ala Val Pro Ser Lys Pro
 130 135 140
 Val Glu Tyr Phe Ala Gln Arg Lys Gly Gly Gly Leu Met His Ala Ile
 145 150 155 160
 Ile Pro Asp Glu Asn Met Ser Phe Thr Gln Thr Gly Ser Gly Ser Arg
 165 170 175
 Thr Asp Leu Phe Val His Thr Glu Asp Ala Phe Leu His Asn Ala Ala
 180 185 190
 Asp Phe Leu Ser Phe Leu Phe Leu Arg Asn Glu Glu Arg Val Pro Ser
 195 200 205
 Thr Leu Tyr Ser Ile Arg Ser His Gly Arg Pro Asp Ala Ile Leu Gln
 210 215 220
 Glu Leu Phe Lys Pro Ile Tyr Lys Cys Pro Lys Asp Ala Asn Tyr Ala
 225 230 235 240
 Ser Glu Glu Ala Leu Gly Asp Asp Ile Arg Thr Ser Val Leu Tyr Gly
 245 250 255
 Ser Arg Ser Ala Pro Phe Met Arg Phe Asp Ala Ala Glu Gln Ile Tyr
 260 265 270
 Asn Glu Asp Ala Asn Gln Asp Pro Glu Ala Leu His Asn Leu Lys Arg
 275 280 285
 Phe Trp Glu Glu Ala Arg Lys Leu Ile Tyr Asn Asp Phe Val Pro Glu
 290 295 300
 Ser Gly Asp Leu Ile Phe Val Asn Asn His Leu Cys Ala His Gly Arg
 305 310 315 320
 Asn Ala Phe Leu Ala Gly Phe Arg Glu Glu Asn Gly Gln Leu Val Lys
 325 330 335
 Cys Glu Arg Arg Leu Met Leu Arg Met Met Ser Lys Thr Ser Leu Ile
 340 345 350
 Asn Ile Arg Glu Val Thr His Pro Glu Asn Pro Tyr Leu Ile Met Glu
 355 360 365
 Glu His Tyr Gly Lys Val Tyr Ser Ala His Leu Ala Asn Leu
 370 375 380

<210> 23
 <211> 1098
 <212> DNA
 <213> Chryseobacterium gleum NBRC15054

<220>
 <221> CDS
 <222> (1)..(1095)

```

<400> 23
atg aat tct aca caa att tta gat aaa gac tgt tta aca gca gta aag      48
Met Asn Ser Thr Gln Ile Leu Asp Lys Asp Cys Leu Thr Ala Val Lys
1      5      10      15
ctg ctt cct acg gtc ata gaa gtc act tct cag gag aga aga atg ata      96
Leu Leu Pro Thr Val Ile Glu Val Thr Ser Gln Glu Arg Arg Met Ile
20      25      30
aaa gat gca gcg ctg cat ctt cag aaa aaa tat ggc act tat gaa aac      144
Lys Asp Ala Ala Leu His Leu Gln Lys Lys Tyr Gly Thr Tyr Glu Asn
35      40      45
cgg gat ttt ata aaa cat gtt cat caa ctg gcc tct tat ttt ctt ccg      192
Arg Asp Phe Ile Lys His Val His Gln Leu Ala Ser Tyr Phe Leu Pro
50      55      60
gaa agg att cta aat ata gca gct gat ttt gca agt gac ttt tct gaa      240
Glu Arg Ile Leu Asn Ile Ala Ala Asp Phe Ala Ser Asp Phe Ser Glu
65      70      75      80
aat cag tat gga gcg ctg gtt ttt aca gga ttg atg gag ata gac cag      288
Asn Gln Tyr Gly Ala Leu Val Phe Thr Gly Leu Met Glu Ile Asp Gln
85      90      95
gaa gaa ata ggt tct act cca ccc aac tgg caa tcg gca gat tat tca      336
Glu Glu Ile Gly Ser Thr Pro Pro Asn Trp Gln Ser Ala Asp Tyr Ser
100      105      110
aag ttt aat tta tat ggt ttt gcg tgt gcg ctt att cat ggg gca ctt      384
Lys Phe Asn Leu Tyr Gly Phe Ala Cys Ala Leu Ile His Gly Ala Leu
115      120      125
ccc tca aag cct gta caa tat tat tca cag cgt aaa ggc ggt gga ttg      432
Pro Ser Lys Pro Val Gln Tyr Tyr Ser Gln Arg Lys Gly Gly Gly Leu
130      135      140
atc cac gct att att cct gat gaa aaa atg aaa gaa aca cag aca gga      480
Ile His Ala Ile Ile Pro Asp Glu Lys Met Lys Glu Thr Gln Thr Gly
145      150      155      160
tca gga tcc tca acg gat ctg tat gta cat aca gaa gat gct ttt ctg      528
Ser Gly Ser Ser Thr Asp Leu Tyr Val His Thr Glu Asp Ala Phe Leu
165      170      175
aaa cat cag gct gac ttt tta agc ttt atg tat gtc cga aat gaa gag      576
Lys His Gln Ala Asp Phe Leu Ser Phe Met Tyr Val Arg Asn Glu Glu
180      185      190
cag gta cct tca act ctt tat tct atc cgt tct cat gag tct att ggg      624
Gln Val Pro Ser Thr Leu Tyr Ser Ile Arg Ser His Glu Ser Ile Gly
195      200      205
gaa aag tac agg aca ctt ttt gag cct att tat aaa atc cct aaa gat      672
Glu Lys Tyr Arg Thr Leu Phe Glu Pro Ile Tyr Lys Ile Pro Lys Asp
210      215      220
gcc aat ctg gaa acg gga gaa aat gaa gaa gaa act ctg gat tct gta      720
Ala Asn Leu Glu Thr Gly Glu Asn Glu Glu Glu Thr Leu Asp Ser Val
225      230      235      240
ttg tat gga aat acc aac ctt cct ttt atg cga ttt gat gcg gcg gaa      768
Leu Tyr Gly Asn Thr Asn Leu Pro Phe Met Arg Phe Asp Ala Ala Glu
245      250      255
cag ctt ttc aat tcc agt atc aga cag tca gaa gaa gcg cag cat aca      816
Gln Leu Phe Asn Ser Ser Ile Arg Gln Ser Glu Glu Ala Gln His Thr
260      265      270
ctg cat gag ttc tgg gaa gaa gcc aga cat ttg att tat tca gga ttt      864
Leu His Glu Phe Trp Glu Glu Ala Arg His Leu Ile Tyr Ser Gly Phe
275      280      285
acg cct cag gcc gga gat gtt att ctg gtt aat aat cat tta tgt gct      912
Thr Pro Gln Ala Gly Asp Val Ile Leu Val Asn Asn His Leu Cys Ala
290      295      300
cac gga aga tct gct ttc cgt gcg gga gta aga aat att gac ggt ata      960
His Gly Arg Ser Ala Phe Arg Ala Gly Val Arg Asn Ile Asp Gly Ile
305      310      315      320
gaa cag ccg tgc gaa cga aga att atg ctt cgg atg atg agt aaa gtg      1008
Glu Gln Pro Cys Glu Arg Arg Ile Met Leu Arg Met Met Ser Lys Val

```

JPOXMLDOC01-seql.app

	325		330		335	
agc ctt att gat atg aga gca cat acc ctt aca gaa gat cct ttt ttt						1056
Ser Leu Ile Asp Met Arg Ala His Thr Leu Thr Glu Asp Pro Phe Phe						
	340		345		350	
gtc ata gaa gaa cat ctg ggt aaa aac ttt caa cat ttt taa						1098
Val Ile Glu Glu His Leu Gly Lys Asn Phe Gln His Phe						
	355		360		365	

<210> 24
 <211> 365
 <212> PRT
 <213> Chryseobacterium gleum NBRC 15054

<400> 24

Met Asn Ser Thr Gln Ile Leu Asp Lys Asp Cys Leu Thr Ala Val Lys																
1	5						10						15			
Leu Leu Pro Thr Val Ile Glu Val Thr Ser Gln Glu Arg Arg Met Ile							25					30				
	20															
Lys Asp Ala Ala Leu His Leu Gln Lys Lys Tyr Gly Thr Tyr Glu Asn							40					45				
	35															
Arg Asp Phe Ile Lys His Val His Gln Leu Ala Ser Tyr Phe Leu Pro							55				60					
	50															
Glu Arg Ile Leu Asn Ile Ala Ala Asp Phe Ala Ser Asp Phe Ser Glu											75				80	
	65															
Asn Gln Tyr Gly Ala Leu Val Phe Thr Gly Leu Met Glu Ile Asp Gln																
	85						90								95	
Glu Glu Ile Gly Ser Thr Pro Pro Asn Trp Gln Ser Ala Asp Tyr Ser																
	100						105								110	
Lys Phe Asn Leu Tyr Gly Phe Ala Cys Ala Leu Ile His Gly Ala Leu																
	115						120								125	
Pro Ser Lys Pro Val Gln Tyr Tyr Ser Gln Arg Lys Gly Gly Gly Leu																
	130						135								140	
Ile His Ala Ile Ile Pro Asp Glu Lys Met Lys Glu Thr Gln Thr Gly																
	145															160
Ser Gly Ser Ser Thr Asp Leu Tyr Val His Thr Glu Asp Ala Phe Leu																
	165															175
Lys His Gln Ala Asp Phe Leu Ser Phe Met Tyr Val Arg Asn Glu Glu																
	180															190
Gln Val Pro Ser Thr Leu Tyr Ser Ile Arg Ser His Glu Ser Ile Gly																
	195						200									205
Glu Lys Tyr Arg Thr Leu Phe Glu Pro Ile Tyr Lys Ile Pro Lys Asp																
	210						215									220
Ala Asn Leu Glu Thr Gly Glu Asn Glu Glu Glu Thr Leu Asp Ser Val																
	225						230									240
Leu Tyr Gly Asn Thr Asn Leu Pro Phe Met Arg Phe Asp Ala Ala Glu																
	245															255
Gln Leu Phe Asn Ser Ser Ile Arg Gln Ser Glu Glu Ala Gln His Thr																
	260															270
Leu His Glu Phe Trp Glu Glu Ala Arg His Leu Ile Tyr Ser Gly Phe																
	275						280									285
Thr Pro Gln Ala Gly Asp Val Ile Leu Val Asn Asn His Leu Cys Ala																
	290						295									300
His Gly Arg Ser Ala Phe Arg Ala Gly Val Arg Asn Ile Asp Gly Ile																
	305						310									320
Glu Gln Pro Cys Glu Arg Arg Ile Met Leu Arg Met Met Ser Lys Val																
	325															335
Ser Leu Ile Asp Met Arg Ala His Thr Leu Thr Glu Asp Pro Phe Phe																
	340															350
Val Ile Glu Glu His Leu Gly Lys Asn Phe Gln His Phe																
	355						360									365

<210> 25
 <211> 1116

<212> DNA

<213> Niastella koreensis NBRC106392

<220>

<221> CDS

<222> (1)..(1113)

<400> 25

atg gaa act atc att gaa tcc aga caa cgc att aac agt ccc ggc gta	48
Met Glu Thr Ile Ile Glu Ser Arg Gln Arg Ile Asn Ser Pro Gly Val	
1 5 10 15	
tta ccg cca cct ttg agt cca ttg atc gtt gac gtt aca ccg aag gag	96
Leu Pro Pro Pro Leu Ser Pro Leu Ile Val Asp Val Thr Pro Lys Glu	
20 25 30	
cgt gcc tcc att tca aac gtg gcc aat atc tta tta aaa gcc ttt ggc	144
Arg Ala Ser Ile Ser Asn Val Ala Asn Ile Leu Leu Lys Ala Phe Gly	
35 40 45	
cat tat gaa cat cct gat ttc atc tcc gct ttg cac ctg aat gct ttt	192
His Tyr Glu His Pro Asp Phe Ile Ser Ala Leu His Leu Asn Ala Phe	
50 55 60	
cag tta tta ccg gaa cgt att gcg ggg ata ctg agc cgt ttt ggt acc	240
Gln Leu Leu Pro Glu Arg Ile Ala Gly Ile Leu Ser Arg Phe Gly Thr	
65 70 75 80	
gac ttc tcg cgc cac caa tac ggc gcg ttg gtg ttc aga ggc ctt aca	288
Asp Phe Ser Arg His Gln Tyr Gly Ala Leu Val Phe Arg Gly Leu Thr	
85 90 95	
gaa gta gat cag gag gcg ctt ggc cct acc ccg ccc tcg tgg aaa gaa	336
Glu Val Asp Gln Glu Ala Leu Gly Pro Thr Pro Pro Ser Trp Lys Glu	
100 105 110	
acc gat tac agc aag ctt gtt aaa tat gga ttt att tgc tcg ctg ctg	384
Thr Asp Tyr Ser Lys Leu Val Lys Tyr Gly Phe Ile Cys Ser Leu Leu	
115 120 125	
cat ggc gcc att cca tca aaa cca gta caa tat tat gcg cag cga aaa	432
His Gly Ala Ile Pro Ser Lys Pro Val Gln Tyr Tyr Ala Gln Arg Lys	
130 135 140	
ggc ggt ggt tta ctg cat gcc gtt att ccc gat gaa aaa atg agt cat	480
Gly Gly Gly Leu Leu His Ala Val Ile Pro Asp Glu Lys Met Ser His	
145 150 155 160	
acg caa acc ggc tcc ggc tcg cgc acc gat ctt ttt gtg cat acc gaa	528
Thr Gln Thr Gly Ser Gly Ser Arg Thr Asp Leu Phe Val His Thr Glu	
165 170 175	
gat gcg ttc tta ttt aac cag gcc gat ttt ctc agc ttc ctg ttc ctg	576
Asp Ala Phe Leu Phe Asn Gln Ala Asp Phe Leu Ser Phe Leu Phe Leu	
180 185 190	
cgg aat gaa gaa cag gtg cca tct acg tta tat tcg atc cgg tcg cat	624
Arg Asn Glu Glu Gln Val Pro Ser Thr Leu Tyr Ser Ile Arg Ser His	
195 200 205	
ggc gat acc aac gcc atc atg gcg gag ctg ttc aaa ccc att tat aag	672
Gly Asp Thr Asn Ala Ile Met Ala Glu Leu Phe Lys Pro Ile Tyr Lys	
210 215 220	
tgt ccg aag gat gcg aat tat gcc gac gat gaa aat gcc ggc gag gaa	720
Cys Pro Lys Asp Ala Asn Tyr Ala Asp Asp Glu Asn Ala Gly Glu Glu	
225 230 235 240	
gtg acc act tct atc tta tac ggt aac cgc gaa cgg ccc ttt atc cgc	768
Val Thr Thr Ser Ile Leu Tyr Gly Asn Arg Glu Arg Pro Phe Ile Arg	
245 250 255	
ttc gat gcc gcg gaa cag atc tac aac gaa aag gcc gga caa acg ccg	816
Phe Asp Ala Ala Glu Gln Ile Tyr Asn Glu Lys Ala Gly Gln Thr Pro	
260 265 270	
gaa gcc atg cac aac ctg gtg cgt ttt tgg gac gaa gcc aaa caa ctt	864
Glu Ala Met His Asn Leu Val Arg Phe Trp Asp Glu Ala Lys Gln Leu	
275 280 285	
atc tac aat gat ttc gtg ccc gat tcg ggc gat ctc att ttt gta aac	912
Ile Tyr Asn Asp Phe Val Pro Asp Ser Gly Asp Leu Ile Phe Val Asn	

JPOXMLDOC01-seql.app

290	295	300	
aac cat ttg tgc gcg cat ggc cgg aat tca ttt gtg gcc ggt tat cgt			960
Asn His Leu Cys Ala His Gly Arg Asn Ser Phe Val Ala Gly Tyr Arg			
305	310	315	320
aat gaa aac ggt cag ctg gta aaa tgt gaa cgc cgg ttg atg tta cgc			1008
Asn Glu Asn Gly Gln Leu Val Lys Cys Glu Arg Arg Leu Met Leu Arg			
	325	330	335
atg atg agc aag acc agc ctc atc aat att cag tcg gtg acc cag tta			1056
Met Met Ser Lys Thr Ser Leu Ile Asn Ile Gln Ser Val Thr Gln Leu			
	340	345	350
aac gac ccg tat ttc att atg gaa gaa cac tac ggc aaa ttg ttt cat			1104
Asn Asp Pro Tyr Phe Ile Met Glu Glu His Tyr Gly Lys Leu Phe His			
	355	360	365
tca caa caa taa			1116
Ser Gln Gln			
370			

<210> 26

<211> 371

<212> PRT

<213> Niastella koreensis NBRC106392

<400> 26

Met Glu Thr Ile Ile Glu Ser Arg Gln Arg Ile Asn Ser Pro Gly Val	
1	5 10 15
Leu Pro Pro Pro Leu Ser Pro Leu Ile Val Asp Val Thr Pro Lys Glu	
	20 25 30
Arg Ala Ser Ile Ser Asn Val Ala Asn Ile Leu Leu Lys Ala Phe Gly	
	35 40 45
His Tyr Glu His Pro Asp Phe Ile Ser Ala Leu His Leu Asn Ala Phe	
	50 55 60
Gln Leu Leu Pro Glu Arg Ile Ala Gly Ile Leu Ser Arg Phe Gly Thr	
65	70 75 80
Asp Phe Ser Arg His Gln Tyr Gly Ala Leu Val Phe Arg Gly Leu Thr	
	85 90 95
Glu Val Asp Gln Glu Ala Leu Gly Pro Thr Pro Pro Ser Trp Lys Glu	
	100 105 110
Thr Asp Tyr Ser Lys Leu Val Lys Tyr Gly Phe Ile Cys Ser Leu Leu	
	115 120 125
His Gly Ala Ile Pro Ser Lys Pro Val Gln Tyr Tyr Ala Gln Arg Lys	
	130 135 140
Gly Gly Gly Leu Leu His Ala Val Ile Pro Asp Glu Lys Met Ser His	
145	150 155 160
Thr Gln Thr Gly Ser Gly Ser Arg Thr Asp Leu Phe Val His Thr Glu	
	165 170 175
Asp Ala Phe Leu Phe Asn Gln Ala Asp Phe Leu Ser Phe Leu Phe Leu	
	180 185 190
Arg Asn Glu Glu Gln Val Pro Ser Thr Leu Tyr Ser Ile Arg Ser His	
	195 200 205
Gly Asp Thr Asn Ala Ile Met Ala Glu Leu Phe Lys Pro Ile Tyr Lys	
	210 215 220
Cys Pro Lys Asp Ala Asn Tyr Ala Asp Asp Glu Asn Ala Gly Glu Glu	
225	230 235 240
Val Thr Thr Ser Ile Leu Tyr Gly Asn Arg Glu Arg Pro Phe Ile Arg	
	245 250 255
Phe Asp Ala Ala Glu Gln Ile Tyr Asn Glu Lys Ala Gly Gln Thr Pro	
	260 265 270
Glu Ala Met His Asn Leu Val Arg Phe Trp Asp Glu Ala Lys Gln Leu	
	275 280 285
Ile Tyr Asn Asp Phe Val Pro Asp Ser Gly Asp Leu Ile Phe Val Asn	
	290 295 300
Asn His Leu Cys Ala His Gly Arg Asn Ser Phe Val Ala Gly Tyr Arg	
305	310 315 320
Asn Glu Asn Gly Gln Leu Val Lys Cys Glu Arg Arg Leu Met Leu Arg	

				325					330					335			
Met	Met	Ser	Lys	Thr	Ser	Leu	Ile	Asn	Ile	Gln	Ser	Val	Thr	Gln	Leu		
			340					345					350				
Asn	Asp	Pro	Tyr	Phe	Ile	Met	Glu	Glu	His	Tyr	Gly	Lys	Leu	Phe	His		
		355					360					365					
Ser	Gln	Gln															
	370																

<210> 27
 <211> 1068
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> artificial synthesis

<220>
 <221> CDS
 <222> (1)..(1065)

<400> 27

atg gaa aca atg tca gca atc gcc cca tct aaa tca cac tta tcg aat	48
Met Glu Thr Met Ser Ala Ile Ala Pro Ser Lys Ser His Leu Ser Asn	
1 5 10 15	
agc tta cgt gtc gca cgc agc aaa gag agc gat att acg gtc cac gaa	96
Ser Leu Arg Val Ala Arg Ser Lys Glu Ser Asp Ile Thr Val His Glu	
20 25 30	
ctg cag agc agc ctg ttt acg ctg gat tct gct agc gcg gaa gcg atc	144
Leu Gln Ser Ser Leu Phe Thr Leu Asp Ser Ala Ser Ala Glu Ala Ile	
35 40 45	
cat acc gcg gct gag cgc att acc gcc cac ccg aac gaa aac ccg gac	192
His Thr Ala Ala Glu Arg Ile Thr Ala His Pro Asn Glu Asn Pro Asp	
50 55 60	
gat ttc ggc cgt cag gcg ctg gca gcg gcg ttt agc ttg ccg gaa gag	240
Asp Phe Gly Arg Gln Ala Leu Ala Ala Ala Phe Ser Leu Pro Glu Glu	
65 70 75 80	
gtg cgt gcg gcg gtc ttg aat ttt gcc gag gtg ggt agc gag agc ggc	288
Val Arg Ala Ala Val Leu Asn Phe Ala Glu Val Gly Ser Glu Ser Gly	
85 90 95	
atc atg gtt gtt cgt ggt ctg tac gtg gat gag gac ctg gcc gac acc	336
Ile Met Val Val Arg Gly Leu Tyr Val Asp Glu Asp Leu Ala Asp Thr	
100 105 110	
ccg ctg gat aac aag agc ggc ctg ggt gcg cgt acc gtt ttt gcg aaa	384
Pro Leu Asp Asn Lys Ser Gly Leu Gly Ala Arg Thr Val Phe Ala Lys	
115 120 125	
gag atg gcc atg ctg gcg cat ctg ctg ggc agc atg gtg gcg tac gag	432
Glu Met Ala Met Leu Ala His Leu Leu Gly Ser Met Val Ala Tyr Glu	
130 135 140	
gcg gaa ggc aac ggt cat ctg att caa gac atg gtg ccg aat ccg aag	480
Ala Glu Gly Asn Gly His Leu Ile Gln Asp Met Val Pro Asn Pro Lys	
145 150 155 160	
ctg gcg gtc acg caa caa agc cag ggt agc aag gtt gag ctg gaa gca	528
Leu Ala Val Thr Gln Gln Ser Gln Gly Ser Lys Val Glu Leu Glu Ala	
165 170 175	
cat acc gag cag tgc ttc agc gac ttc aaa ccg gat tat gtt att ctg	576
His Thr Glu Gln Cys Phe Ser Asp Phe Lys Pro Asp Tyr Val Ile Leu	
180 185 190	
ggt gct ctg cgt ggc gac gaa aac gcc aac acc tat gca ttc tcc ggt	624
Gly Ala Leu Arg Gly Asp Glu Asn Ala Asn Thr Tyr Ala Phe Ser Gly	
195 200 205	
cgc aaa ctg gtt cag cac atg tcc gcc gaa gaa gtg gcg aaa ctg cgc	672
Arg Lys Leu Val Gln His Met Ser Ala Glu Glu Val Ala Lys Leu Arg	
210 215 220	

JPOXMLDOC01-seql.app

```

caa cct ctg tgg gca act acc atc gat gag agc ttt caa ccg tac att      720
Gln Pro Leu Trp Ala Thr Thr Ile Asp Glu Ser Phe Gln Pro Tyr Ile
225                230                235                240
ccg aat ccg gac gac gtt cgc ggt ccg ttc ccg att ctg acg ggc cca      768
Pro Asn Pro Asp Asp Val Arg Gly Pro Phe Pro Ile Leu Thr Gly Pro
                245                250                255
gag gac gat ccg tac atc cgt gta gac cag gag ctg atg cac ggt atc      816
Glu Asp Asp Pro Tyr Ile Arg Val Asp Gln Glu Leu Met His Gly Ile
                260                265                270
acc gcg gac gct caa cgc ctg ctg cgc aag gtt gtg gat acg tat atc      864
Thr Ala Asp Ala Gln Arg Leu Leu Arg Lys Val Val Asp Thr Tyr Ile
                275                280                285
gag cac cgc gac gcg cat gtc ttg cag cct ggt gat ctg ctg atg ctg      912
Glu His Arg Asp Ala His Val Leu Gln Pro Gly Asp Leu Leu Met Leu
                290                295                300
gac aat ctg cgt gca atg cac ggt cgt agc atg ttc gct ccg cgt ttt      960
Asp Asn Leu Arg Ala Met His Gly Arg Ser Met Phe Ala Pro Arg Phe
305                310                315                320
gac ggc aag gat cgt ttc att gca cgt ggt ttc gtc gtg cgt gat cgt      1008
Asp Gly Lys Asp Arg Phe Ile Ala Arg Gly Phe Val Val Arg Asp Arg
                325                330                335
cgt aag ttg tgg ccg cag ctg ttg gaa gat cgt cgc acc ctg ggt gca      1056
Arg Lys Leu Trp Pro Gln Leu Leu Glu Asp Arg Arg Thr Leu Gly Ala
                340                345                350
gtc cac tcc taa
Val His Ser
                355

```

<210> 28

<211> 355

<212> PRT

<213> marine actinobacterium PHSC20C1

<400> 28

```

Met Glu Thr Met Ser Ala Ile Ala Pro Ser Lys Ser His Leu Ser Asn
1          5          10          15
Ser Leu Arg Val Ala Arg Ser Lys Glu Ser Asp Ile Thr Val His Glu
20        25        30
Leu Gln Ser Ser Leu Phe Thr Leu Asp Ser Ala Ser Ala Glu Ala Ile
35        40        45
His Thr Ala Ala Glu Arg Ile Thr Ala His Pro Asn Glu Asn Pro Asp
50        55        60
Asp Phe Gly Arg Gln Ala Leu Ala Ala Ala Phe Ser Leu Pro Glu Glu
65        70        75        80
Val Arg Ala Ala Val Leu Asn Phe Ala Glu Val Gly Ser Glu Ser Gly
85        90        95
Ile Met Val Val Arg Gly Leu Tyr Val Asp Glu Asp Leu Ala Asp Thr
100       105       110
Pro Leu Asp Asn Lys Ser Gly Leu Gly Ala Arg Thr Val Phe Ala Lys
115       120       125
Glu Met Ala Met Leu Ala His Leu Leu Gly Ser Met Val Ala Tyr Glu
130       135       140
Ala Glu Gly Asn Gly His Leu Ile Gln Asp Met Val Pro Asn Pro Lys
145       150       155       160
Leu Ala Val Thr Gln Gln Ser Gln Gly Ser Lys Val Glu Leu Glu Ala
165       170       175
His Thr Glu Gln Cys Phe Ser Asp Phe Lys Pro Asp Tyr Val Ile Leu
180       185       190
Gly Ala Leu Arg Gly Asp Glu Asn Ala Asn Thr Tyr Ala Phe Ser Gly
195       200       205
Arg Lys Leu Val Gln His Met Ser Ala Glu Glu Val Ala Lys Leu Arg
210       215       220
Gln Pro Leu Trp Ala Thr Thr Ile Asp Glu Ser Phe Gln Pro Tyr Ile

```

JPOXMLDOC01-seql.app

225		230		235		240									
Pro	Asn	Pro	Asp	Asp	Val	Arg	Gly	Pro	Phe	Pro	Ile	Leu	Thr	Gly	Pro
			245				250							255	
Glu	Asp	Asp	Pro	Tyr	Ile	Arg	Val	Asp	Gln	Glu	Leu	Met	His	Gly	Ile
		260					265						270		
Thr	Ala	Asp	Ala	Gln	Arg	Leu	Leu	Arg	Lys	Val	Val	Asp	Thr	Tyr	Ile
		275					280					285			
Glu	His	Arg	Asp	Ala	His	Val	Leu	Gln	Pro	Gly	Asp	Leu	Leu	Met	Leu
	290					295					300				
Asp	Asn	Leu	Arg	Ala	Met	His	Gly	Arg	Ser	Met	Phe	Ala	Pro	Arg	Phe
305				310					315					320	
Asp	Gly	Lys	Asp	Arg	Phe	Ile	Ala	Arg	Gly	Phe	Val	Val	Arg	Asp	Arg
			325				330						335		
Arg	Lys	Leu	Trp	Pro	Gln	Leu	Leu	Glu	Asp	Arg	Arg	Thr	Leu	Gly	Ala
		340					345						350		
Val	His	Ser													
	355														

<210> 29
 <211> 37
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer

<400> 29
 ttatcatatg aaatcacaat cattaattga agatgag 37

<210> 30
 <211> 41
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer

<400> 30
 tgtaatatgct cgagagcctg atcaaaaact tttcctaaat g 41

<210> 31
 <211> 31
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer

<400> 31
 attcacatat gtcctcgctg ttcctcgact c 31

<210> 32
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer

<400> 32

agcttctcga ggctgaagct ggcctgcacg 30

<210> 33
 <211> 32
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer

<400> 33
 ataatcatat gagaccctta gacgtgacac cc 32

<210> 34
 <211> 38
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer

<400> 34
 aatagctcga gaagggttgc caggtgagcg ctatatac 38

<210> 35
 <211> 30
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer

<400> 35
 ataatcatat gaattctaca caaatttttag 30

<210> 36
 <211> 32
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer

<400> 36
 aatagctcga gaaaatgttg aaagttttta cc 32

<210> 37
 <211> 29
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> PCR primer

<400> 37
 ataatcatat ggaaactatc attgaatcc 29

<210> 38

<211> 33

<212> DNA

<213> Artificial Sequence

<220>

<223> PCR primer

<400> 38

aatagctcga gttggttgtga atgaaacaat ttg

33