

COMPOUNDS AND METHODS FOR ^{18}F LABELED AGENTS

Field of invention:

5 This invention relates to novel compounds suitable for radiolabeling or already being radiolabeled with halogen, more specifically with ^{18}F , a method of making such compounds, a composition comprising such compounds, their use for diagnostic imaging, a kit comprising a sealed vial containing a predetermined quantity of such novel compounds, such compounds for use as medicament, as diagnostic imaging agent and
10 most specifically as imaging agent for Positron Emission Tomography.

Background art:

Over the last few years, *in-vivo* scanning using Positron Emission Tomography (PET) has
15 increased. PET is both a medical and research tool. It is used heavily in clinical oncology for medical imaging of tumors and the search for metastasis, and for clinical diagnosis of certain diffuse brain diseases such as those causing various types of dementias. Radiotracers consisting of a radionuclide stably bound to a biomolecule are used for *in vivo* imaging of disorders.

20

In designing an effective radiopharmaceutical tracer for use as a diagnostic agent, it is imperative that the drug has appropriate *in vivo* targeting and pharmacokinetic properties. Fritzberg *et al.* (*J. Nucl. Med.*, 1992, 33:394) state further that radionuclide chemistry and associated linkages underscore the need to optimize the attachment and labelling of
25 chemical modifications of the biomolecule carrier, diluent, excipient or adjuvant. Hence the type of radionuclide, the type of biomolecule and the method used for linking them to one another may have a crucial effect onto the radiotracer properties.

Peptides are biomolecules that play a crucial role in many physiological processes
30 including actions as neurotransmitters, hormones, and antibiotics. Research has shown their importance in such fields as neuroscience, immunology, pharmacology and cell biology. Some peptides can act as chemical messenger. They bind to receptor on the target cell surface and the biological effect of the ligand is transmitted to the target tissue. Hence the specific receptor binding property of the ligand can be exploited by labelling the
35 ligand with a radionuclide. Theoretically, the high affinity of the ligand for the receptor facilitates retention of the radio labeled ligand in receptor expressing tissues. However, it

is still under investigation which peptides can efficiently be labeled and under which conditions the labelling shall occur. It is well known that receptor specificity of ligand peptide may be altered during chemical reaction. Therefore an optimal peptidic construct has to be determined.

5

Tumors overexpress various receptor types to which peptide bound specifically. Boerman *et al.* (*Seminar in Nuclear Medicine*, 30(3) July, 2000; pp195-208) provide a non exhaustive list of peptides binding to receptor involved in tumor, *i.e.*, somatostatin, vasoactive intestinal peptide (VIP), bombesin binding to gastrin-releasing peptide (GRP) receptor, gastrin, cholecystokinin (CCK) and calcitonin.

10

The radionuclides used in PET scanning are typically isotopes with short half lives such as ^{11}C (~20 min), ^{13}N (~10 min), ^{15}O (~2 min), ^{68}Ga (~68 min) or ^{18}F (~110 min). Due to their short half lives, the radionuclides must be produced in a cyclotron which is not too far away in delivery-time from the PET scanner. These radionuclides are incorporated into biologically active compounds or biomolecules that have the function to vehicle the radionuclide into the body though the targeted site, for example a tumor.

15

The linkage of the radionuclide to the biomolecule is done by various methods resulting in the presence or not of a linker between the radionuclide and the biomolecule. Hence, various linkers are known. C.J.Smith *et al.* ("*Radiochemical investigations of ^{177}Lu -DOTA-8-Aoc-BBN[7-14] NH_2 : an in vitro / in vivo assessment of the targeting ability of this new radiopharmaceutical for PC-3 human prostate cancer cells.*" *Nucl. Med. Bio.*, 30(2):101-9; 2003) disclose radiolabeled bombesin wherein the linker is DOTA-X where X is a carbon tether. However, the radiolabel ^{177}Lu (half life 6,5 days) does not match the biological half-life of the native bombesin what makes the ^{177}Lu -DOTA-X-bombesin a non-appropriate radiotracer for imaging tumor.

20

25

E.Garcia Garayoa *et al.* ("*Chemical and biological characterization of new $\text{Re}(\text{CO})_3[^{99\text{m}}\text{Tc}](\text{CO})_3$ bombesin Analogues.*" *Nucl. Med. Biol.*, 17-28; 2007) disclose a spacer between the radionuclide [$^{99\text{m}}\text{Tc}$] and the bombesin wherein the spacer is - β -Ala- β -Ala- and 3,6-dioxa-8-aminooctanoic acid. E.Garcia Garayoa *et al.* conclude that the different spacer does not have a significant effect on stability or on receptor affinity.

30

Listed above linkers have been specifically designed for a specific type of radionuclide and determine the type and chemical conditions of the radiobinding method.

35

More recently, peptides have been conjugated to a macrocyclic chelator for labelling with ^{64}Cu , ^{86}Y , and ^{68}Ga for PET application. However, such radionuclides interact with the *in vivo* catabolism resulting in unwanted physiologic effects and chelate attachment.

5

The nucleophilic aromatic ^{18}F -fluorination reaction is of great importance for ^{18}F -labeled radiopharmaceuticals which are used as *in vivo* imaging agents for targeting and visualizing diseases, e.g., solid tumors or diseases of brain. A very important technical goal in using ^{18}F -labeled radiopharmaceuticals is the quick preparation and administration of the radioactive compound due to the fact that the ^{18}F isotopes have a short half-life of about only 111 minutes.

^{18}F -labeled compounds are gaining importance due to the availability thereof as well as due to the development of methods for labeling biomolecules. It has been shown that some compounds labeled with ^{18}F produce images of high quality. Additionally, the longer lifetime of ^{18}F would permit longer imaging times and allow preparation of radiotracer batches for multiple patients and delivery of the tracer to other facilities, making the technique more widely available to clinical investigators. Additionally, it has been observed that the development of PET cameras and availability of the instrumentation in many PET centers is increasing. Hence, it is increasingly important to develop new tracers labeled with ^{18}F .

The nucleophilic aromatic ^{18}F -fluorination reaction is of great importance for ^{18}F -labeled radiopharmaceuticals which are used as *in vivo* imaging agents targeting and visualizing diseases, e.g., solid tumors.

Various methods of radiofluorination have been published using different precursors or starting material for obtaining ^{18}F -labeled peptides. Due to the smaller size of peptides, both higher target-to-background ratios and rapid blood clearance can often be achieved with radiolabeled peptides. Hence, short-lived positron emission tomography (PET) isotopes are potential candidates for labelling peptides. Among a number of positron-emitting nuclides, fluorine-18 appears to be the best candidate for labelling bioactive peptides by virtue of its favourable physical and nuclear characteristics. The major disadvantage of labelling peptides with ^{18}F is the laborious and time-consuming preparation of the ^{18}F labelling agents. Due to the complex nature of peptides and several functional groups associated with the primary structure, ^{18}F -labeled peptides are not

prepared by direct fluorination. Hence, difficulties associated with the preparation of ^{18}F -labeled peptide were alleviated with the employment of prosthetic groups as shown below. Several such prosthetic groups have been proposed in the literature, including N-succinimidyl-4- ^{18}F fluorobenzoate, m-maleimido-N-(p- ^{18}F fluorobenzyl)-benzamide, N-(p- ^{18}F fluorophenyl) maleimide, and 4- ^{18}F fluorophenacylbromide. Almost all of the methodologies currently used today for the labeling of peptides and proteins with ^{18}F utilize active esters of the fluorine labeled synthon.



$\bigcirc$ = aliphatic, aromatic or hetero-aromatic, alicyclic

${}^{18}\text{F}-\bigcirc-\text{RM}$ = PROSTHETIC GROUP

RM = reactive moiety

LG = Leaving group that can be replaced by ^{18}F

X = functional group for reaction with RM

10

Okarvi *et al.* ("Recent progress in fluorine-18 labeled peptide radiopharmaceuticals." *Eur. J. Nucl. Med.*, 2001 Jul; 28(7):929-38)) present a review of the recent developments in ^{18}F -labeled biologically active peptides used in PET.

15 Xianzhong Zhang *et al.* (" ^{18}F -labeled bombesin analogs for targeting GRP receptor-expressing prostate cancer." *J. Nucl. Med.*, 47(3):492-501 (2006)) relate to the 2-step method detailed above. [Lys3]Bombesin ([Lys3]BBN) and aminocaproic acid-bombesin(7-14) (Aca-BBN(7-14)) were labeled with ^{18}F by coupling the Lys3 amino group and Aca amino group, respectively, with N-succinimidyl-4- ^{18}F -fluorobenzoate (^{18}F -SFB) under slightly basic condition (pH 8.5). Unfortunately, the obtained ^{18}F -FB-[Lys3]BBN is metabolically relatively unstable having for result to reduce the extent of use of the ^{18}F -FB-[Lys3]BBN for reliable imaging of tumor.

25

Thorsten Poethko *et al.* ("Two-step methodology for high-yield routine radiohalogenation of peptides: ^{18}F -labeled RGD and octreotide analogs." *J. Nucl. Med.*, 2004 May; 45(5):892-902) relate to a 2-step method for labelling RGD and octreotide analogs. The method discloses the steps of radiosynthesis of the ^{18}F -labeled aldehyde or ketone and


the chemoselective ligation of the ^{18}F -labeled aldehyde or ketone to the aminooxy functionalized peptide.

- 5 Thorsten Poethko *et al.* ("First ^{18}F -labeled tracer suitable for routine clinical imaging of somatostatin receptor-expressing tumors using positron emission tomography." *Clin. Cancer Res.*, 2004 Jun 1;10(11):3593-606) apply the 2-step method for the synthesis of ^{18}F -labeled carbohydrate Tyr(3)-octreotate (TOCA) analogs with optimized pharmacokinetics suitable for clinical routine somatostatin-receptor (sst) imaging.
- 10 WO 2003/080544 A1 and WO 2004/080492 A1 relate to radiofluorination methods of bioactive peptides for diagnostics imaging using the 2-step method shown above.

The most crucial aspect in the successful treatment of any cancer is early detection. Likewise, it is crucial to properly diagnose the tumor and metastasis.

- 15 Routine application of ^{18}F -labeled peptides for quantitative *in vivo* receptor imaging of receptor-expressing tissues and quantification of receptor status using PET is limited by the lack of appropriate radiofluorination methods for routine large-scale synthesis of ^{18}F -labeled peptides. There is a clear need for radiofluorination method that can be conducted
- 20 rapidly without loss of receptor affinity by the peptide and leading to a positive imaging (with reduced background), wherein the radiotracer is stable and shows enhanced clearance properties

- The conversions of *mono*- (mainly *para*-) substituted trimethylammonium benzene derivatives (1) to substituted [^{18}F]-fluoro-benzene derivatives (2) which may serve as radiopharmaceutical itself or as prosthetic group for the F-18 labeling of small and large
- 25 molecules have been reported in the literature (Irie *et al.* 1982, *Fluorine Chem.*, 27, (1985), 117-191; Haka *et al.* 1989) (see Scheme 1).



Scheme 1

There are only a few publications about nucleophilic aromatic ^{18}F -fluorination reactions of trimethyl ammonium substituted aromatic derivatives which contain two or more substituents beside the trimethylammonium moiety:

- 5 Oya *et al.* treated [2-Chloro-5-(2-dimethylcarbamoyl-phenylsulfanyl)-4-nitro-phenyl]-trimethyl-ammonium triflate with [^{18}F] potassium fluoride and obtained the desired ^{18}F -labeled compound (*J. Med. Chem.*, 2002, 45(21):4716-4723).

- 10 Li *et al.* report on the ^{18}F -fluorination reaction of 4-(N,N,N-trimethylammonium)-3-cyano-3'-iodobenzophenone triflate (*Bioconjugate Chemistry*, 2003, 14(2):287-294).

Enas *et al.* convert (2,2-Dimethyl-1,3-dioxo-indan-5-yl)-trimethyl-ammonium triflate into the desired ^{18}F -labeled compound (*J. Fluorine Chem.*, 1993, 63(3):233-41).

- 15 Seimbille *et al.* and other groups have labeled (2-Chloro-4-nitro-phenyl)-trimethyl-ammonium triflate successfully with ^{18}F (*J. Labeled Compd. Radiopharm.*, 2005, 48, 11:829-843).

- 20 (2-Benzoyloxy-4-formyl-phenyl)-trimethyl-ammonium triflate has successfully been labeled with ^{18}F at high temperature (130°C) by Langer *et al.* (*Bioorg. Med. Chem.*, EN; 9; 3; 2001:677-694).

- 25 Lang *et al.* have radiolabeled trimethyl-(2-methyl-4-pentamethylphenyl methoxycarbonyl-phenyl)-ammonium triflate by using [^{18}F] potassium fluoride (*J. Med. Chem.*, 42, 9, 1999:1576-1586).

Trimethyl-(4-nitro-naphthalen-1-yl)-ammonium triflate has been labeled with ^{18}F by Amokhtari *et al.* (*J. Labeled Compd. Radiopharm.*; S42, 1 (1999):S622-S623).

- 30 Lemaire *et al.* have converted (2-formyl-5-methoxy-phenyl)-trimethyl-ammonium triflate into the desired ^{18}F -labeled product (*J. Labeled Compd. Radiopharm.*, 44, 2001:S857-S859).

- 35 VanBrocklin *et al.* describe the ^{18}F labeling of (2-bromo-4-nitro-phenyl)-trimethyl-ammonium triflate (*J. Labeled Compd. Radiopharm.*, 44; 2001:S880-S882) and

Cetir Centre Medic report on the successful ^{18}F -labeling of (5-Chloro-8-hydroxy-quinolin-7-yl)-trimethyl-ammonium triflate (EP 1 563 852 A1).

5 D.A.Sutton *et al.* ("*Evaluation of 1-fluoro-2-nitro-4-trimethylammoniumbenzene iodide, a protein-solubilizing agent*", *Biochem. J.*, 1972, 130:589-595) disclose model derivatives which consist of a benzene substituted with trimethylammonium, an electron withdrawing nitro group and a glycine, phenylalanine or acetyltyrosine.

10 C.Lemaire *et al.* ("*Highly enantioselective synthesis of no-carrier-added 6-[^{18}F]fluoro-L-dopa by chiral phase-transfer alkylation*", *Eur. J. Org. Chem.*, 2004:2899-2904) disclose 2-[^{18}F]fluoro-4,5-dimethoxybenzaldehyde to be used to prepare 6-[^{18}F]fluoro-L-dopa.

15 L.Lang *et al.* ("*Development of fluorine-18-labeled 5-HT_{1A} antagonists*", *J. Med. Chem.*, 1999, 42(9):1576-1586) disclose conversion of pentamethyl 4-(trimethylammonium trifluoromethanesulfonate)benzoate and pentamethyl 3-methyl-4-(trimethylammonium trifluoromethanesulfonate)benzoate to the respective ^{18}F substituted benzoyl chloride which is then coupled with WAY 100635 (N-{2-[4-(2-methoxyphenyl)-piperazino]ethyl}-N-(2-pyridyl)cyclohexanecarboxamide).

20 S.Oya *et al.* ("*New PET imaging agent for the serotonin transporter: [^{18}F]ACF (2-[(*l*-amino-4-chloro-5-fluorophenyl)thio]-N,N-dimethyl-benzenmethanamine)*", *J. Med. Chem.*, 2002, 45:4716-4723) disclose conversion of [2-chloro-5-(2-dimethylaminocarbonyl-phenylthio)-4-nitro-phenyl]trimethylammonium trifluoromethanesulfonate to the respective ^{18}F substituted compound.

25 M.J.Al-Darwich *et al.* ("*Enantioselective synthesis of non-carrier-added (n.c.a.) (S)-4-chloro-2-[^{18}F]fluorophenylalanine and (S)-(α -methyl)-4-chloro-2-[^{18}F]fluorophenylalanine*", *J. Fluorine Chem.*, 1996, 80:117-124) disclose 4-chloro-2-trimethylammoniumbenzaldehyde triflate to be reacted to 4-chloro-2-[^{18}F]fluorobenzaldehyde which are then
30 further reacted to yield the title compounds.

Y.Seimbille *et al.* ("*Fluorine-18 labeling of 6,7-disubstituted anilinoquinazoline derivatives for positron emission tomography (PET) imaging of tyrosine kinase receptors: synthesis of ^{18}F -Iressa and related molecular probes*", *J. Labeled Compd. Radiopharm.*, 2005, 48:829-
35 843, *i.a.*, report on the reaction of 3-chloro-4-trimethylammonium nitrobenzene trifluoro-

methanesulfonate to give 3-chloro-4-[^{18}F]fluoroaniline via 3-chloro-4-[^{18}F]fluoro-nitrobenzene.


WO 2002/44144 A1 relates to nucleophilic reaction for preparing radiolabeled imaging agents using [^{18}F]fluoride to react with trimethylammoniumbenzene compounds.

WO 2006/083424 A2 relates to [^{18}F]-radiolabeled compounds and the manufacture thereof.

Most of these mentioned ^{18}F -labeled aromatic derivatives which contain two or more additional substituents cannot be coupled to chemical functionalities like amines, thiols, carboxylic acids, phenols or other chemicals groups of complex molecules like peptides without further transformations.

^{18}F labeling of more complex radiopharmaceuticals like peptides takes place in all known publications in a two- or multi-step strategy (see Scheme 2, review article: *Eur. J. Nucl. Med.*, 2001, 28:929-938).

For these kinds of ^{18}F -labeling also mono-substituted trimethylammonium benzene derivatives are used and react in a first step with [^{18}F] potassium fluoride to obtain substituted [^{18}F]-fluoro-benzene derivatives. These compounds are then coupled in a second step to larger and more complex molecules like peptides, small molecules or nucleotides (see scheme 2).



Scheme 2

Especially 4-[^{18}F]fluorobenzaldehyde has been used in many examples for F-18 labelling of complex molecules (e.g., *J. Nucl. Med.*, 2004, 45(5):892-902). But also N-succinimidyl-8-[4'-[^{18}F]fluorobenzylamino]suberate (*Bioconjugate Chem.*, 1991, 2:44-49), 4-[^{18}F]fluorophenacyl bromide and 3-[^{18}F]fluoro-5-nitrobenzimidate (*J. Nucl. Med.*, 1987,

28:462-470), m-maleimido-N-(p-[^{18}F]fluorobenzyl)-benzamide (*J. Labeled Compd. Radiopharm.*, 1989, 26:287-289), N-{4-[4-[^{18}F]fluorobenzylidene(aminooxy)-butyl]-maleimide (*Bioconjugate Chem.*, 2003, 14:1253-1259), [^{18}F]N-(4-fluorobenzyl)-2-bromoacetamide (*Bioconjugate Chem.*, 2000, 11:627-636) and [^{18}F]-3,5-difluorophenyl azide (and 5 derivatives) (*J. Org. Chem.*, 1995, 60:6680-6681) are known examples. F-18 labeling of peptides via para-[^{18}F]-fluorobenzoates is also a very common method either by coupling of the corresponding acid with additional activating agents (such as 1,3-dicyclohexylcarbodiimide/1-hydroxy-7-azabenzotriazole (DCC/HOAt) or N-[(dimethyl-amino)-1H-1,2,3-triazolyl[4,5]pyridine-1-yl-methylene]-N-methyl-methan-aminium hexafluorophosphate N-oxide (HATU/DIPEA, *Eur. J. Nucl. Med. Mol. Imaging.*, 2002, 29:754-759) or by isolated N-succinimidyl 4-[^{18}F]fluorobenzoate (*Nucl. Med. Biol.*, 1996, 23:365).

None of these compounds and none of other published compounds allow a direct (one-step) labeling of peptides with ^{18}F -fluoride.

Therefore is an object of the present invention the development of a practical and mild technique for ^{18}F labeling of molecules like, e.g., peptides, oligonucleotides or small molecule targeting agents and to provide radiofluorination methods for obtaining radiotracer based on receptor specific peptides for the detection of tumors.

Summary of the Invention:

The object of the present invention is solved as detailed herein below.

25

A first aspect of the present invention refers to novel compounds having general chemical Formula A (general chemical Formula I), wherein $K = \text{N}^+(\text{R}^1)(\text{R}^2)(\text{R}^3) \text{X}^-$ and to pharmaceutically acceptable salts of inorganic or organic acids thereof, hydrates, complexes, esters, amides, solvates and prodrugs thereof. These compounds are precursors for a single step labeling more preferably radiolabeling to yield the novel compounds according to the second aspect of the present invention.

30

A second aspect of the present invention refers to novel compounds (radiopharmaceuticals, labeled ^{18}F), having general chemical Formula A, wherein $K = \text{W}$ (general chemical Formula II) and to pharmaceutically acceptable salts of inorganic or

35

organic acids thereof, hydrates, complexes, esters, amides, solvates and prodrugs thereof.

5 Compounds having general chemical Formula A, wherein $K = N^+(R^1)(R^2)(R^3) X^-$ (general chemical Formula I), can be converted into compounds having general chemical Formula A, wherein $K = W$ (general chemical Formula II), by means of a one-step labeling more preferably radiolabeling reaction with a fluoine isotope, more specifically with ^{18}F .

10 A third aspect of the present invention refers to a one-step method of labeling more preferably radiolabeling compounds having general chemical Formula A, wherein $K = N^+(R^1)(R^2)(R^3) X^-$, in order to arrive at compounds having general chemical Formula A, wherein $K = W$.

15 A fourth aspect of the present invention refers to compositions, more preferably to diagnostic compositions, comprising a compound having general chemical Formula A, wherein $K = N^+(R^1)(R^2)(R^3) X^-$, or a pharmaceutically acceptable salts of inorganic or organic acids thereof, hydrates, complexes, esters, amides, solvates and prodrugs thereof and a pharmaceutically acceptable carrier, diluent, excipient or adjuvant. According to this fourth aspect the present invention further refers to compositions, more preferably
20 diagnostic compositions, comprising a radiolabelled compound having general chemical Formula A, wherein $K = W$, or a pharmaceutically acceptable salts of inorganic or organic acids thereof, hydrates, complexes, esters, amides, solvates and prodrugs thereof and a pharmaceutically acceptable carrier, diluent, excipient or adjuvant.

25 A fifth aspect, the present invention refers to a method of imaging diseases, the method comprising introducing into a patient a detectable quantity of a labeled compound having general chemical Formula A, wherein $K = W$, or a pharmaceutically acceptable salts of inorganic or organic acids thereof, hydrates, complexes, esters, amides, solvates and prodrugs thereof.

30 A sixth aspect of the present invention refers to a kit for preparing a radiopharmaceutical preparation, said kit comprising a sealed vial containing a predetermined quantity of the compound of Formula A, wherein $K = N^+(R^1)(R^2)(R^3) X^-$, or a pharmaceutically acceptable salts of inorganic or organic acids thereof, hydrates, complexes, esters, amides, solvates
35 and prodrugs thereof and further optionally an acceptable carrier, diluent, excipient or adjuvant supplied as a mixture with the compound having general chemical Formula I or

independently for the manufacture of a compound having general chemical Formula II. More preferably, the present invention relates to a kit comprising a compound or composition, as defined herein above, in powder form, and a container containing an appropriate solvent for preparing a solution of the compound or composition for administration to an animal, including a human.

A seventh aspect of the present invention refers to a compound having general chemical Formula A, wherein $K = N^+(R^1)(R^2)(R^3) X^-$ or W, or of a pharmaceutically acceptable salts of inorganic or organic acids thereof, hydrates, complexes, esters, amides, solvates and prodrugs thereof for use as medicament and, if $K = W$, for use as diagnostic imaging agent and more specifically for use as imaging agent for PET.

An eighth aspect of the present invention refers to a use of a compound having general chemical Formula A, wherein $K = N^+(R^1)(R^2)(R^3) X^-$ or W, or of a pharmaceutically acceptable salts of inorganic or organic acids thereof, hydrates, complexes, esters, amides, solvates and prodrugs thereof for the manufacture of a medicament, more specifically for the manufacture of a diagnostic imaging agent and most specifically for the manufacture of a diagnostic imaging agent for imaging tissue at a target site using the imaging agent.

A ninth aspect of the present invention refers to bombesin analogs that bind specifically to human GRP receptors present in prostate tumor, breast tumor and metastasis. In a preferred embodiment, the bombesin analog is a peptide having sequence from Seq ID 1 to Seq ID 102 or those sequences disclosed below.

Further aspects of the present invention are directed to methods and intermediates useful for synthesizing the tumor imaging compounds having general chemical Formulae I and II as described herein below.

Detailed Description of the Invention:

As used hereinafter in the description of the invention and in the claims, the term "alkyl", by itself or as part of another group, refers to a straight chain or branched chain alkyl group with 1 to 20 carbon atoms such as, for example methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, isopentyl, neopentyl, heptyl, hexyl, decyl. Alkyl groups can also be substituted, such as by halogen atoms, hydroxyl groups, C_1 - C_4 alkoxy groups

or C₆-C₁₂ aryl groups (which, intern, can also be substituted, such as by 1 to 3 halogen atoms). More preferably alkyl is C₁-C₁₀ alkyl, C₁-C₆ alkyl or C₁-C₄ alkyl.

As used hereinafter in the description of the invention and in the claims, the term

5 "cycloalkyl" by itself or as part of another group, refers to mono- or bicyclic chain of alkyl group with 3 to 20 carbon atoms such as, for example cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl. More preferably cycloalkyl is C₃-C₁₀ cycloalkyl or C₅-C₈ cycloalkyl, most preferably C₆ cycloalkyl.

10 As used hereinafter in the description of the invention and in the claims, the term "heterocycloalkyl", by itself or as part of another group, refers to groups having 3 to 20 mono- or bi-ring atoms of a cycloalkyl; and containing carbon atoms and 1, 2, 3 or 4 oxygen, nitrogen or sulfur heteroatoms. More preferably heterocycloalkyl is C₃-C₁₀ heterocycloalkyl, C₅-C₈ heterocycloalkyl or C₅-C₁₄ heterocycloalkyl, most preferably C₆ heterocycloalkyl.

20 As used hereinafter in the description of the invention and in the claims, the term "aralkyl" refers to aryl- substituted alkyl radicals such as benzyl, diphenylmethyl, triphenylmethyl, phenylethyl, phenylbutyl and diphenylethyl.

As used hereinafter in the description of the invention and in the claims, the terms "aryloxy" refers to aryl groups having an oxygen through which the radical is attached to a nucleus, examples of which are phenoxy.

25

As used hereinafter in the description of the invention and in the claims, the terms "alkenyl" and "alkynyl" are similarly defined as for alkyl, but contain at least one carbon-carbon double or triple bond, respectively. More preferably C₂-C₆ alkenyl and C₂-C₆ alkynyl.

30

As used hereinafter in the description of the invention and in the claims, the term "lower unbranched or branched alkyl" shall have the following meaning: a substituted or unsubstituted, straight or branched chain monovalent or divalent radical consisting substantially of carbon and hydrogen, containing no unsaturation and having from one to eight carbon atoms, e.g., but not limited to methyl, ethyl, n-propyl, n-pentyl, 1,1-dimethylethyl (t-butyl), n-heptyl and the like.

35

As used hereinafter in the description of the invention and in the claims, the terms "aralkenyl" refers to aromatic structure (aryl) coupled to alkenyl as defined above.

5 As used hereinafter in the description of the invention and in the claims, the terms "alkoxy (or alkyloxy), aryloxy, and aralkenyloxy" refer to alkyl, aryl, and aralkenyl groups respectively linked by an oxygen atom, with the alkyl, aryl, and aralkenyl portion being as defined above.

10 As used hereinafter in the description of the invention and in the claims, the terms "salts of inorganic or organic acids", "inorganic acid" and "organic acid" refer to mineral acids, including, but not being limited to: acids such as carbonic, nitric, phosphoric, hydrochloric, perchloric or sulphuric acid or the acidic salts thereof such as potassium hydrogen sulphate, or to appropriate organic acids which include, but are not limited to: acids such
15 as aliphatic, cycloaliphatic, aromatic, araliphatic, heterocyclic, carboxylic and sulphonic acids, examples of which are formic, acetic, trifluoroacetic, propionic, succinic, glycolic, gluconic, lactic, malic, fumaric, pyruvic, benzoic, anthranilic, mesylic, fumaric, salicylic, phenylacetic, mandelic, embonic, methanesulfonic, ethanesulfonic, benzenesulfonic, phanthothenic, toluenesulfonic, trifluormethanesulfonic and sulfanilic acid, respectively.

20

As used hereinafter in the description of the invention and in the claims, the term "aryl" by itself or as part of another group refers to monocyclic or bicyclic aromatic groups containing from 6 to 12 carbon atoms in the ring portion, preferably 6-10 carbons in the ring portion, such as phenyl, naphthyl or tetrahydronaphthyl.

25

As used hereinafter in the description of the invention and in the claims, the term "heteroaryl" by itself or as part of another group, refers to groups having 5 to 14 ring atoms; 6, 10 or 14 π (pi) electrons shared in a cyclic array; and containing carbon atoms and 1, 2, 3 or 4 oxygen, nitrogen or sulfur heteroatoms (where examples of heteroaryl
30 groups are: thienyl, benzo[b]thienyl, naphtho[2,3-b]thienyl, thianthrenyl, furyl, pyranly, isobenzofuranyl, benzoxazolyl, chromenyl, xanthenyl, phenoxathiinyl, 2*H*-pyrrolyl, pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyrazinyl, pyrimidinyl, pyridazinyl, indoliziny, isoindolyl, 3*H*-indolyl, indolyl, indazolyl, purinyl, 4*H*-quinoliziny, isoquinolyl, quinolyl, phthalazinyl, naphthyridinyl, quinazolinyl, cinnolinyl, pteridinyl, 4*aH*-carbazolyl, carbazolyl, carbolinyl,
35 phenanthridinyl, acridinyl, perimidinyl, phenanthrolinyl, phenazinyl, isothiazolyl, phenothiazinyl, isoxazolyl, furazanyl and phenoxazinyl groups).

Whenever the term substituted is used, it is meant to indicate that one or more hydrogens on the atom indicated in the expression using "substituted" is replaced with a selection from the indicated group, provided that the indicated atom's normal valency is not exceeded, and that the substitution results in a chemically stable compound, i. e. a compound that is sufficiently robust to survive isolation to a useful degree of purity from a reaction mixture, and formulation into a pharmaceutical composition. The substituent groups may be selected from halogen atoms, hydroxyl groups, C₁-C₄ alkoxy groups or C₆-C₁₂ aryl groups (which, intern, can also be substituted, such as by 1 to 3 halogen atoms).

As used hereinafter in the description of the invention and in the claims, the term "fluorine isotope" (F) refers to all isotopes of the fluorine atomic element. Fluorine isotope (F) is selected from radioactive or non-radioactive isotope. The radioactive fluorine isotope is selected from ¹⁸F. The non-radioactive "cold" fluorine isotope is selected from ¹⁹F.

As used hereinafter in the description of the invention and in the claims, the term "prodrug" means any covalently bonded compound, which releases the active parent pharmaceutical according to formula II.

The term "prodrug" as used throughout this text means the pharmacologically acceptable derivatives such as esters, amides and phosphates, such that the resulting in vivo biotransformation product of the derivative is the active drug as defined in the compounds of formula (I). The reference by Goodman and Gilman (The Pharmacological Basis of Therapeutics, 8 ed, McGraw-Hill, Int. Ed. 1992, "Biotransformation of Drugs", p 13-15) describing prodrugs generally is hereby incorporated. Prodrugs of a compound of the present invention are prepared by modifying functional groups present in the compound in such a way that the modifications are cleaved, either in routine manipulation or in vivo, to the parent compound. Prodrugs of the compounds of the present invention include those compounds wherein for instance a hydroxy group, such as the hydroxy group on the asymmetric carbon atom, or an amino group is bonded to any group that, when the prodrug is administered to a patient, cleaves to form a free hydroxyl or free amino, respectively.

Typical examples of prodrugs are described for instance in WO 99/33795, WO 99/33815, WO 99/33793 and WO 99/33792 all incorporated herein by reference.

Prodrugs are characterized by excellent aqueous solubility, increased bioavailability and are readily metabolized into the active inhibitors in vivo.


As used hereinafter in the description of the invention and in the claims, the terms "amino acid sequence" and "peptide" are defined herein as a polyamide obtainable by (poly)condensation of at least two amino acids.

5

As used hereinafter in the description of the invention and in the claims, the term "amino acid" means any molecule comprising at least one amino group and at least one carboxyl group, but which has no peptide bond within the molecule. In other words, an amino acid is a molecule that has a carboxylic acid functionality and an amine nitrogen having at least one free hydrogen, preferably in alpha position thereto, but no amide bond in the molecule structure. Thus, a dipeptide having a free amino group at the N-terminus and a free carboxyl group at the C-terminus is not to be considered as a single "amino acid" in the above definition. The amide bond between two adjacent amino acid residues which is obtained from such a condensation is defined as "peptide bond". Optionally, the nitrogen atoms of the polyamide backbone (indicated as NH above) may be independently alkylated, e.g., with C₁-C₆-alkyl, preferably CH₃.

An amide bond as used herein means any covalent bond having the structure

20



wherein the carbonyl group is provided by one molecule and the NH-group is provided by the other molecule to be joined. The amide bonds between two adjacent amino acid residues which are obtained from such a polycondensation are defined as "peptide bonds". Optionally, the nitrogen atoms of the polyamide backbone (indicated as NH above) may be independently alkylated, e.g., with -C₁-C₆-alkyl, preferably -CH₃.

As used hereinafter in the description of the invention and in the claims, an amino acid residue is derived from the corresponding amino acid by forming a peptide bond with another amino acid.

As used hereinafter in the description of the invention and in the claims, an amino acid sequence may comprise naturally occurring and/or synthetic / artificial amino acid residues, proteinogenic and/or non-proteinogenic amino acid residues. The non-

35

proteinogenic amino acid residues may be further classified as (a) homo analogues of proteinogenic amino acids, (b) β -homo analogues of proteinogenic amino acid residues and (c) further non-proteinogenic amino acid residues.

5 Accordingly, the amino acid residues may be derived from the corresponding amino acids, *e.g.*, from

- proteinogenic amino acids, namely Ala, Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr and Val; or

10 • non-proteinogenic amino acids, such as

- homo analogues of proteinogenic amino acids wherein the sidechain has been extended by a methylene group, *e.g.*, homoalanine (Hal), homoarginine (Har), homocysteine (Hcy), homoglutamine (Hgl), homohistidine (Hhi), homoisoleucine (Hil), homoleucine (Hle), homolysine (Hly), homomethionine (Hme),
- 15 homophenylalanine (Hph), homoproline (Hpr), homoserine (Hse), homothreonine (Hth), homotryptophane (Htr), homotyrosine (Hty) and homovaline (Hva);
- β -homo analogues of proteinogenic amino acids wherein a methylene group has been inserted between the α -carbon and the carboxyl group yielding β -amino acids, *e.g.*, β -homoalanine (β Hal), β -homoarginine (β Har), β -homoasparagine (β Has), β -
- 20 homocysteine (β Hcy), β -homoglutamine (β Hgl), β -homohistidine (β Hhi), β -homoisoleucine (β Hil), β -homoleucine (β Hle), β -homolysine (β Hly), β -homomethionine (β Hme), β -homophenylalanine (β Hph), β -homoproline (β Hpr), β -homoserine (β Hse), β -homothreonine (β Hth), β -homotryptophane (β Htr), β -homotyrosine (β Hty) and β -homovaline (β Hva);

- 25 ○ further non-proteinogenic amino acids, *e.g.*, α -aminoadipic acid (Aad), β -aminoadipic acid (β Aad), α -aminobutyric acid (Abu), α -aminoisobutyric acid (Aib), β -alanine (β Ala), 4-aminobutyric acid (4-Abu), 5-aminovaleric acid (5-Ava), 6-aminohexanoic acid (6-Ahx), 8-aminooctanoic acid (8-Aoc), 9-aminononanoic acid (9-Anc), 10-aminodecanoic acid (10-Adc), 12-aminododecanoic acid (12-Ado), α -
- 30 aminosuberic acid (Asu), azetidine-2-carboxylic acid (Aze), β -cyclohexylalanine (Cha), aitrulline (Cit), dehydroalanine (Dha), γ -carboxyglutamic acid (Gla), α -cyclohexylglycine (Chg), propargylglycine (Pra), pyroglutamic acid (Glp), α -tert-butylglycine (Tle), 4-benzoylphenylalanine (Bpa), δ -hydroxylysine (Hyl), 4-hydroxyproline (Hyp), allo-isoleucine (alle), lanthionine (Lan), (1-naphthyl)alanine (1-Nal),
- 35 (2-naphthyl)alanine (2-Nal), norleucine (Nle), norvaline (Nva), ornithine (Orn), phenylglycin (Phg), pipercolic acid (Pip), sarcosine (Sar), selenocysteine (Sec),

- statine (Sta), β -thienylalanine (Thi), 1,2,3,4-tetrahydroisochinoline-3-carboxylic acid (Tic), allo-threonine (aThr), thiazolidine-4-carboxylic acid (Thz), γ -aminobutyric acid (GABA), iso-cysteine (iso-Cys), diaminopropionic acid (Dpr), 2,4-diaminobutyric acid (Dab), 3,4-diaminobutyric acid ($\gamma\beta$ Dab), biphenylalanine (Bip), phenylalanine substituted in para-position with $-C_1-C_6$ alkyl, -halide, $-NH_2$, $-CO_2H$ or Phe(4-R) (wherein R = $-C_1-C_6$ alkyl, -halide, $-NH_2$, or $-CO_2H$); peptide nucleic acids (PNA, *cf.*, P.E. Nielsen, *Acc. Chem. Res.*, 32, 624-30);
- or their N-alkylated analogues, such as their N-methylated analogues.

- Cyclic amino acids may be proteinogenic or non-proteinogenic, such as Pro, Aze, Glp, Hyp, Pip, Tic and Thz.

For further examples and details reference can be made to, *e.g.*, J.H. Jones, *J. Peptide Sci.*, 2003, 9, 1-8 which is herein incorporated by reference.

- As used hereinafter in the description of the invention and in the claims, the terms "non-proteinogenic amino acid" and "non-proteinogenic amino acid residue" also encompass derivatives of proteinogenic amino acids. For example, the side chain of a proteinogenic amino acid residue may be derivatized thereby rendering the proteinogenic amino acid residue "non-proteinogenic". The same applies to derivatives of the C-terminus and/or the N-terminus of a proteinogenic amino acid residue terminating the amino acid sequence.

- As used hereinafter in the description of the invention and in the claims, a proteinogenic amino acid residue is derived from a proteinogenic amino acid selected from the group consisting of Ala, Arg, Asn, Asp, Cys, Gln, Glu, Gly, His, Ile, Leu, Lys, Met, Phe, Pro, Ser, Thr, Trp, Tyr and Val either in L- or D-configuration; the second chiral center in Thr and Ile may have either R- or S-configuration. Therefore, for example, any posttranslational modification of an amino acid sequence, such as N-alkylation, which might naturally occur renders the corresponding modified amino acid residue "non-proteinogenic", although in nature said amino acid residue is incorporated in a protein. Preferably modified amino acids are selected from N-alkylated amino acids, β -amino acids, γ -amino acids, lanthionines, dehydro amino acids, and amino acids with alkylated guanidine moieties.

- As used hereinafter in the description of the invention and in the claims, the term "peptidomimetic" relates to molecules which are related to peptides, but with different properties. A peptidomimetic is a small protein-like chain designed to mimic a peptide.

They typically arise from modification of an existing peptide in order to alter the molecule's properties. For example, they may arise from modifications to change the molecule's stability or biological activity. This can have a role in the development of drug-like compounds from existing peptides. These modifications involve changes to the peptide
5 that will not occur naturally.

As used hereinafter in the description of the invention and in the claims, the term "peptide analogs", by itself refers to synthetic or natural compounds which resemble naturally occurring peptides in structure and/or function.

10

As used hereinafter in the description of the invention and in the claims, the term "pharmaceutically acceptable salt" relates to salts of inorganic and organic acids, such as mineral acids, including, but not limited to, acids such as carbonic, nitric or sulfuric acid, or organic acids, including, but not limited to acids such as aliphatic, cycloaliphatic, aromatic,
15 araliphatic, heterocyclic, carboxylic and sulphonic acids, examples of which are formic, acetic, trifluoroacetic, propionic, succinic, glycolic, gluconic, lactic, malic, fumaric, pyruvic, benzoic, anthranilic, mesylic, salicylic, phenylacetic, mandelic, embonic, methansulfonic, ethanesulfonic, benzenesulfonic, phantothenic, toluenesulfonic and sulfanilic acid.

20 If a chiral center or another form of an isomeric center is present in a compound having general chemical Formulae A, I, II, III or IV of the present invention, as given hereinafter, all forms of such isomers, including enantiomers and diastereoisomers, are intended to be covered herein. Compounds containing a chiral center may be used as a racemic mixture or as an enantiomerically enriched mixture, or the racemic mixture may be separated
25 using well-known techniques and an individual enantiomer maybe used alone. In cases in which compounds have unsaturated carbon-carbon double bonds, both the cis-isomer and trans-isomers are within the scope of this invention. In cases in which compounds may exist in tautomeric forms, such as keto-enol tautomers, each tautomeric form is contemplated as being included within the scope of the present invention whether existing
30 in equilibrium or predominantly in one form.

As used hereinafter in the description of the invention and in the claims, the term "oligonucleotide" shall have the following meaning: short sequences of nucleotides, typically with twenty or fewer bases. Examples are, but are not limited to, molecules
35 named and cited in the book: *"The aptamers handbook. Functional oligonuclides and their*

application" by Svenn Klussmann, Wiley-VCH, 2006. An example for such an oligonucleotide is TTA1 (*J. Nucl. Med.*, 2006, April, 47(4):668-78).

As used hereinafter in the description of the invention and in the claims, the term

5 "aptamer" refers to an oligonucleotide, comprising from 4 to 100 nucleotides, wherein at least two single nucleotides are connected to each other via a phosphodiester linkage. Said aptamers have the ability to bind specifically to a target molecule (see, e.g., M Famulok, G Mayer, "Aptamers as Tools in Molecular Biology and Immunology", in: "Combinatorial Chemistry in Biology, Current Topics in Microbiology and Immunology" (M

10 Famulok, CH Wong, EL Winnacker, Eds.), Springer Verlag Heidelberg, 1999, Vol. 243, 123-136). There are many ways known to the skilled person of how to generate such aptamers that have specificity for a certain target molecule. An example is given in WO 01/09390 A, the disclosure of which is hereby incorporated by reference. Said aptamers may comprise substituted or non-substituted natural and non-natural nucleotides.

15 Aptamers can be synthesized *in vitro* using, e.g., an automated synthesizer. Aptamers according to the present invention can be stabilized against nuclease degradation, e.g., by the substitution of the 2'-OH group versus a 2'-fluoro substituent of the ribose backbone of pyrimidine and versus 2'-O-methyl substituents in the purine nucleic acids. In addition, the 3' end of an aptamer can be protected against exonuclease degradation by inverting the

20 3' nucleotide to form a new 5'-OH group, with a 3' to 3' linkage to a penultimate base.

For the purpose of this invention, the term "nucleotide" refers to molecules comprising a nitrogen-containing base, a 5-carbon sugar, and one or more phosphate groups. Examples of said base comprise, but are not limited to, adenine, guanine, cytosine, uracil,


25 and thymine. Also non-natural, substituted or non-substituted bases are included. Examples of 5-carbon sugar comprise, but are not limited to, D-ribose, and D-2-desoxyribose. Also other natural and non-natural, substituted or non-substituted 5-carbon sugars are included. Nucleotides as used in this invention may comprise from one to three phosphates.

30

As used hereinafter in the description of the invention and in the claims, the term "halogen" refers to F, Cl, Br and I.

In a **first aspect** the present invention refers to compounds having general chemical

35 Formula A (general chemical Formula I), wherein $K = N^+(R^1)(R^2)(R^3) X^-$:



wherein:

5

-G is selected from the group comprising -F, -Cl, -Br, -I, -NO, -NO₂, -NR⁴COCF₃,
 -NR⁴SO₂CF₃, -N(R⁴)SO₂R⁵, -N(CF₃)₂, -NHCSNHR⁴, -N(SO₂R⁵)₂, -N(O)=NCONH₂,
 -NR⁴CN, -NHCSR⁵, -N≡C, -N=C(CF₃)₂, -N=NCF₃, -N=NCN, -NR⁴COR⁴,
 -NR⁴COOR⁵, -OSO₂CF₃, -OSO₂C₆H₅, -OCOR⁵, -ONO₂, -OSO₂R⁵, -O-C=CH₂,
 10 -OCF₂CF₃, -OCOCF₃, -OCN, -OCF₃, -C≡N, -C(NO₂)₃, -COOR⁴, -CONR⁴R⁵,
 -C(S)NH₂, -CH=NOR⁴, -CH₂SO₂R⁴, -COCF₃, -CF₃, -CF₂Cl-CBr₃, -CCIF₂, -CCl₃,
 -CF₂CF₃, -C≡CR⁴, -CH=NSO₂CF₃, -CH₂CF₃, -COR⁵, -CH=NOR⁵, -CH₂CONH₂,
 -CSNHR⁵, -CH=NNHCSNH₂, -CH=NNHCONHNH₂, -C≡C-CF₃, -CF=CFCF₃,
 -CF₂-CF₂-CF₃, -CR⁴(CN)₂, -COCF₂CF₂CF₃, -C(CF₃)₃, -C(CN)₃, -CR⁴=C(CN)₂,
 15 -1-pyrryl, -C(CN)=C(CN)₂, -C-pyridyl, -COC₆H₅, -COOC₆H₅, -SOCF₃, -SO₂CF₃,
 -SCF₃, -SO₂CN, -SCOCF₃, -SOR⁵, -S(OR⁵), -SC≡CR⁴, -SO₂R⁵, -SSO₂R⁵, -SR⁵,
 -SSR⁴, -SO₂CF₂CF₃, -SCF₂CF₃, -S(CF₃)=NSO₂CF₃, -SO₂C₆H₅, -SO₂N(R⁵)₂,
 -SO₂C(CF₃)₃, -SC(CF₃)₃, -SO(CF₃)=NSO₂CF₃, -S(O)(=NH)CF₃, -S(O)(=NH)R⁵,
 -S-C=CH₂, -SCOR⁵, -SOC₆H₅, -P(O)C₃F₇, -PO(OR⁵)₂, -PO(N(R⁵)₂)₂, -P(N(R⁵)₂)₂,
 20 -P(O)R⁵, and -PO(OR⁵)₂ or another electron-drawing group wherein the respective
 substituent can be in ortho, para or meta position in respect of the K group,

For the purposes of the present invention, the term “electron-drawing group” or
 “electron withdrawing group” refers to a chemical moiety (substituent) which is
 25 attached to the benzene ring, which is able to decrease the electron density of the
 benzene ring and which is listed in Chem. Rev. (1991), 91, 165-195, Table1 (and
 references therein) with values of σ_m or $\sigma_p > 0$;

-Q is hydrogen, lower unbranched or branched alkyl, aryl, heteroaryl,
 30 -O-(C₁-C₄ alkyl), -CN, -halogen, -SO₂-R⁴, -NO₂ or a condensed aryl or condensed

heteroaryl wherein the respective substituent can be in ortho, para or meta position in respect of the K group,

wherein R^4 is hydrogen or lower unbranched or branched alkyl and
 R^5 is lower unbranched or branched alkyl,

-L- is a bond, -CO-, -SO₂-, -(CH₂)_d-CO-, -SO-, -C≡C-CO-, -[CH₂]_m-E-[CH₂]_n-CO-,
 -[CH₂]_m-E-[CH₂]_n-SO₂-, -C(=O)-O-, -NR¹⁰-, -O-, -(S)_p-, -C(=O)NR¹²-, -C(=S)NR¹²-,
 -C(=S)O-, C₁-C₆ cycloalkyl, alkenyl, heterocycloalkyl, unsubstituted or substituted
 aryl or unsubstituted or substituted heteroaryl, aralkyl, heteroaralkyl, alkyleneoxy,
 arylenoxy, aralkyloxy, -SO₂NR¹³-, -NR¹³SO₂-, -NR¹³C(=O)O-, -NR¹³C(=O)NR¹²-, -
 NH-NH- and -NH-O-,

wherein

d is an integer of from 1 to 6,,

m and n, independently, are any integer of from 0 to 5,

-E- is a bond, -S-, -O- or -NR⁹-,

wherein R⁹ is H, C₁-C₁₀ alkyl, aryl, heteroaryl or aralkyl,

p is any integer of from 1 to 3 ,

R¹⁰, R¹¹ and R¹², independently, are selected from the group comprising H, C₁-
 C₁₀ alkyl, aryl, heteroaryl or aralkyl and

R¹³ is H, substituted or nonsubstituted, linear or branched C₁-C₆ alkyl, aryl,
 cycloalkyl, heterocycloalkyl, heteroaryl, aralkyl or heteroaralkyl,

-Y- is a bond or a spacer,

U is a targeting agent, preferably selected from the group comprising peptides,
 peptidomimetics, small molecules and oligonucleotides.

K is N⁺(R¹)(R²)(R³) X⁻,

X⁻ is CF₃S(O)₂O⁻, C₄F₉S(O)₂O⁻, iodide anion, bromide anion, chloride anion,
 perchlorate anion (ClO₄⁻), phosphate anion, trifluoroacetate anion (CF₃-
 C(O)O⁻), or the anion of another salt of an inorganic or organic acid,

wherein

R¹, R² and R³ are independently from each other selected from the group comprising substituted or unsubstituted alkyl and aralkyl.

The invention further refers to pharmaceutically acceptable salts of inorganic or organic acids, hydrates, complexes, esters, amides, solvates and prodrugs of the compounds having general chemical Formula I.

In a preferred embodiment of the present invention, in the compounds having general chemical Formula I, **-G** is selected from the group comprising -F, -Cl, -Br, -NO₂,
 10 -NR⁴SO₂R⁵, -NHCSNHR⁴, -NR⁴CN, -NR⁴SO₂CF₃, -N≡C, -NR⁴COR⁴, -NR⁴COOR⁵,
 -OSO₂R⁵, -OCF₃, -C≡N, -COOR⁴, -CONR⁴R⁵, -COCF₃, -CF₂CF₃, -C≡CR⁴, -COR⁵,
 -CH₂CONH₂, -CF₃, -C≡C-CF₃, -CF₂-CF₂-CF₃, -C(CN)=C(CN)₂, -COC₆H₅, -SO₂CF₃, -
 SCOCF₃, -SO₂R⁵, -SO₂CF₂CF₃, -SO₂C₆H₅, -SO₂N(R⁵)₂ and -PO(OR⁵)₂ wherein the
 respective substituent can be in ortho, para or meta position in respect of the K group.

15

In a further preferred embodiment of the present invention, in the compounds having general chemical Formula I, **-G** is selected from the group comprising -F, -Cl, -Br, -NO₂,
 -NR⁴SO₂R⁵, -NR⁴COR⁴, -NR⁴COOR⁵, -C≡N, -CONR⁴R⁵, -C≡CR⁴, -COR⁵, -CF₃, -COC₆H₅,
 -SO₂CF₃, -SO₂R⁵, -SO₂C₆H₅ and -SO₂N(R⁵)₂ wherein the respective substituent can be in
 20 ortho, para or meta position in respect of the K group.

In a further alternative embodiment of the present invention, in the compounds having general chemical Formula I, **-G** is selected from the group comprising -F, -Cl, -Br, -NO₂,
 -NR⁴SO₂R⁵, -NR⁴COR⁴, -NR⁴COOR⁵, -C≡N, -CONR⁴R⁵, -C≡CR⁴, -COR⁵, -CF₃ and
 25 -SO₂R⁵ wherein the respective substituent can be in ortho, para or meta position in respect of the K group.

In a further more preferred alternative embodiment of the present invention, in the compounds having general chemical Formula I, **-G** is selected from a group comprising
 30 those members which have a value of the Hammet constant $\sigma \geq 0.35$ (compare *Chem. Rev.*, 1991, 91:165, Table 1) and which contain a fluoro or a nitrogen atom:

-F, -NO, -NO₂, -NR⁴SO₂CF₃, -N(CF₃)₂, -N(SO₂R⁵)₂, -N(O)=NCONH₂, -N≡C, -N=NCF₃,
 -N=NCN, -NR⁴COR⁴, -OSO₂CF₃, -OCOR⁵, -ONO₂, -OCF₂CF₃, -OCOCF₃, -OCN, -OCF₃,
 35 -C≡N, -C(NO₂)₃, -CONR⁴R⁵, -CH=NOR⁴, -COCF₃, -CF₃, -CF₂Cl-CBr₃, -CClF₂, -CF₂CF₃,

- CH=NSO₂CF₃, -CH=NNHCSNH₂, -CF=CFCF₃, -CF₂-CF₂-CF₃, -CR⁴(CN)₂,
 -COCF₂CF₂CF₃, -C(CF₃)₃, -C(CN)₃, -CR⁴=C(CN)₂, -C(CN)=C(CN)₂, -SOCF₃, -SO₂CF₃,
 -SCF₃, -SO₂CN, -SCOCF₃, -SO₂CF₂CF₃, -SCF₂CF₃, -S(CF₃)=NSO₂CF₃, -SO₂N(R⁵)₂,
 -SO₂C(CF₃)₃, -SC(CF₃)₃, -SO(CF₃)=NSO₂CF₃, -S(O)(=NH)CF₃, -S(O)(=NH)R⁵ and
 5 -P(O)C₃F₇ wherein the respective substituent can be in ortho, para or meta position in respect of the K group.

In a further even more preferred alternative embodiment of the present invention, in the compounds having general chemical Formula I, -G is selected from a group comprising
 10 those members of the embodiment herein above which have a value of the Hammett constant $\sigma \geq 0.50$ (compare *Chem. Rev.*, 1991, 91:165, Table 1) or which contain a fluoro atom:

- F, -NO, -NO₂, -NR⁴SO₂CF₃, -N(CF₃)₂, -N(O)=NCONH₂, -N=NCF₃, -N=NCN, -OSO₂CF₃,
 15 -ONO₂, -OCF₂CF₃, -OCOCF₃, -OCN, -OCF₃, -C $\equiv$ N, -C(NO₂)₃, -COCF₃, -CF₃, -CF₂Cl-CBr₃,
 -CClF₂, -CF₂CF₃, -CH=NSO₂CF₃, -CF=CFCF₃, -CF₂-CF₂-CF₃, -CR⁴(CN)₂, -COCF₂CF₂CF₃,
 -C(CF₃)₃, -C(CN)₃, -CR⁴=C(CN)₂, -C(CN)=C(CN)₂, -SOCF₃, -SO₂CF₃, -SCF₃, -SO₂CN,
 -SCOCF₃, -SO₂CF₂CF₃, -SCF₂CF₃, -S(CF₃)=NSO₂CF₃, -SO₂N(R⁵)₂, -SO₂C(CF₃)₃,
 -SC(CF₃)₃, -SO(CF₃)=NSO₂CF₃, -S(O)(=NH)CF₃ and -P(O)C₃F₇ wherein the respective
 20 substituent can be in ortho, para or meta position in respect of the K group.

In a further even more preferred alternative embodiment of the present invention, in the compounds having general chemical Formula I, -G is selected from a group comprising -
 F, -NO₂, -OCF₂CF₃-OCF₃, -C $\equiv$ N, -COCF₃, -CF₃, -CF₂CF₃, -CF₂-CF₂-CF₃, -COCF₂CF₂CF₃,
 25 -SO₂CF₃, -SO₂CN, -SO₂CF₂CF₃, -SO₂N(R⁵)₂ and SC(CF₃)₃ wherein the respective substituent can be in ortho, para or meta position in respect of the K group.

In a further even more preferred alternative embodiment of the present invention, in the compounds having general chemical Formula I, -G is selected from the group comprising
 30 those members with a value of the Hammett constant $\sigma \geq 0.50$ (compare *Chem. Rev.*, 1991, 91:165, Table 1) or which contain a sulfur or a fluoro atom:

- F, -NR⁴SO₂CF₃, -N(CF₃)₂, -N=NCF₃, -OSO₂CF₃, -OCF₂CF₃, -OCOCF₃, -OCF₃, -COCF₃,
 -CF₃, -CF₂Cl-CBr₃, -CClF₂, -CF₂CF₃, -CH=NSO₂CF₃, -CF=CFCF₃, -CF₂-CF₂-CF₃,
 35 -COCF₂CF₂CF₃, -C(CF₃)₃, -SOCF₃, -SO₂CF₃, -SCF₃, -SO₂CN, -SO₂R⁵, -SCOCF₃,
 -SO₂CF₂CF₃, -SCF₂CF₃, -S(CF₃)=NSO₂CF₃, -SO₂N(R⁵)₂, -SO₂C(CF₃)₃, -SC(CF₃)₃,

-SO(CF₃)=NSO₂CF₃, -S(O)(=NH)CF₃ and -P(O)C₃F₇ wherein the respective substituent can be in ortho, para or meta position in respect of the K group.

In a further even more preferred alternative embodiment of the present invention, in the compounds having general chemical Formula I, -G is selected from the group comprising -F, -NR⁴SO₂CF₃, -OSO₂CF₃-OCF₂CF₃, -OCF₃, -COCF₃, -CF₃, -SO₂CF₃, SO₂R⁵ and -SO₂N(R⁵)₂ wherein the respective substituent can be in ortho, para or meta position in respect of the K group.

A positive value of a Hammett constant is a measure of electron deficiency. It seems that certain combinations of substituents with particular atoms (nitrogen, sulfur and/or fluoro) are favourable over others. For example nitrogen or fluoro substituents combined with positive Hammett constants allow a ¹⁸F radiolabeling with relative high radiochemical yields whereas sulfur or fluoro atoms seem to guarantee radiolabeling reactions with only minor side reactions. It is for example known from literature that the choice of substituent can influence the ratio of ring fluorination versus methyl fluoride formation at trimethylammonium benzene derivatives with two substituents in total (review Coenen, "Fluorine-18 Labeling Methods: Features and Possibilities of Basic Reactions", 2006, in: P.A.Schubiger, M.Friebe, L.Lehmann, (eds), *PET-Chemistry - The Driving Force in Molecular Imaging*. Springer, Berlin Heidelberg, p.15-50, in particular p.23-26).

In a further alternative embodiment of the present invention -G is selected from the group comprising -F, -Cl, -Br, -NO₂, -OSO₂R⁴, -OCF₃, -C≡N, -COOR⁴, -CONR⁴R⁵, -COCF₃, -CF₂CF₃, -COR⁵, -CF₃, -C≡C-CF₃, -CF₂-CF₂-CF₃, -COC₆H₅, -SO₂CF₃, -SCOCF₃, -SO₂R⁵, -SO₂CF₂CF₃, -SO₂C₆H₅, -SO₂N(R⁵)₂, and -PO(OR⁵)₂ and more preferably from -F, -Cl, -Br, -NO₂, -C≡N, -CF₃, -SO₂CF₃, -SO₂R⁵, -SO₂C₆H₅, or -SO₂N(R⁵)₂ wherein the respective substituent can be in ortho, para or meta position in respect of the K group.

In a further preferred embodiment of the present invention, -Q is -H, lower unbranched or branched alkyl, aryl, heteroaryl, -O-(C₁-C₄alkyl), -CN, -halogen, -SO₂-R⁴-, -NO₂ or a condensed aryl or heteroaryl wherein the respective substituent can be in ortho, para or meta position in respect of the K group.

In a more preferred embodiment of the present invention, -Q is -H, -C₁-C₄ alkyl, -O-(C₁-C₄ alkyl), -CN, -F, -Cl, -Br or -NO₂ wherein the respective substituent can be in ortho, para or meta position in respect of the K group.

In an even more preferred embodiment of the present invention, -Q is -H, -CH₃, -O-CH₃, -CN, -F, -Cl or -NO₂ wherein the respective substituent can be in ortho, para or meta position in respect of the K group.

5

In an alternative embodiment -Q is -H, -CN, -halogen, -SO₂-R⁴ or -NO₂ wherein the respective substituent can be in ortho, para or meta position in respect of the K group.

10 In a further alternative embodiment of the present invention -Q is selected from -H, -CN, -F, -Cl, -Br or -NO₂, more preferably from -H, -CN, -F or -NO₂ wherein the respective substituent can be in ortho, para or meta position in respect of the K group.

15 In a further alternative embodiment of the present invention, in the compounds having general chemical Formula I, -Q is -H, -CN or -F wherein the respective substituent can be in ortho, para or meta position in respect of the K group.

In a very preferred embodiment of the present invention, any of -G and -Q, more preferred -G, is in *ortho*, or less preferred, in *para* position relative to $K = N^+(R^1)(R^2)(R^3) X^-$.

20 In a preferred embodiment of compounds of Formula I, -G and -Q are independently from each other selected from -H, -CN, CF₃, and -Cl.

In a more preferred embodiment -G and -Q are independently from each other H, -CF₃, or CN.

25

In a even more preferred embodiment in a more preferred embodiment -G and -Q are independently from each other H, -CF₃, or -CN, whereas at least -G or -Q is -CF₃ or -CN.

30 In a further preferred embodiment of the present invention, in the compounds having general chemical Formula I, R⁴ is hydrogen or unbranched or branched C₁-C₆ alkyl, more preferably C₁-C₄ alkyl, most preferably hydrogen or methyl.

In a further preferred embodiment of the present invention, in the compounds having general chemical Formula I, R⁵ is unbranched or branched C₁-C₆ alkyl, more preferably C₁-C₄ alkyl, most preferably methyl.

35

In a preferred embodiment of the present invention, in compounds having general chemical Formula I, R^1 , R^2 and R^3 are independently from each other selected from the group comprising lower unbranched (linear) or branched alkyl or aralkyl.

- 5 In another preferred embodiment of the present invention, in the compounds having general chemical Formula I, R^1 , R^2 and R^3 are independently from each other selected from the group comprising aralkyl or lower alkyl, whereas one of the three moieties (R^1 , R^2 , R^3) can be resin-bound.
- 10 In another preferred embodiment of the present invention, in the compounds having general chemical Formula I, R^1 , R^2 and R^3 are independently from each other selected from the group comprising aralkyl or lower alkyl, preferably C_1 - C_6 alkyl, whereas at least two moieties of the three moieties (R^1 , R^2 , R^3) are alkyl.
- 15 In another preferred embodiment of the present invention, in the compounds having general chemical Formula I, R^1 , R^2 and R^3 are independently from each other selected from the group comprising C_1 - C_6 alkyl moieties.

In another preferred embodiment of the present invention, in the compounds having
20 general chemical Formula I, R^1 is aralkyl and R^2 and R^3 are each methyl.

In another preferred embodiment of the present invention, in the compounds having general chemical Formula I, R^1 , R^2 and R^3 are each methyl.

- 25 In a further preferred embodiment of the present invention, in the compound having general chemical Formula I, X^- is $CF_3-C(O)O^-$, $CF_3S(O)_2O^-$, $C_4F_9S(O)_2O^-$.

In an even more preferred embodiment of the present invention, in the compound having general chemical Formula I, X^- is $CF_3-C(O)O^-$ or $CF_3S(O)_2O^-$.

30

In a further preferred embodiment of the present invention, $-L-$ is a bond, $-CO-$, $-SO_2-$, $-(CH_2)_d-CO-$, $-SO-$, or $-C\equiv C-CO-$, wherein d is an integer of from 1 to 6.


- In a more preferred embodiment of the present invention, in the compounds having
35 general chemical Formula I, $-L-$ is $-CO-$, $-SO_2-$ or $-C\equiv C-CO-$.

In an even more preferred embodiment of the present invention, in the compounds having general chemical Formula I, -L- is -CO- or -SO₂-.

5 In a further preferred embodiment of the present invention, in the compound having general chemical Formula I, the spacer -Y- is a natural or unnatural amino acid sequence or mixture thereof or a non-amino acid group.

10 In a more preferred embodiment of the present invention, in the compounds having general chemical Formula I, the spacer -Y- is an amino acid sequence with two (2) to twenty (20) amino acid residues.

15 In an even more preferred embodiment of the present invention, in the compounds having general chemical Formula I, the spacer -Y- is Arg-Ser, Arg-Ava, Lys(Me)₂-β-ala, Lys(Me)₂-ser, Arg-β-ala, Ser-Ser, Ser-Thr, Arg-Thr, S-alkylcysteine, cysteic acid, thioalkylcysteine (S-S-Alkyl) or



wherein k and l are independently selected in the range of from 0 to 4.

20 In an even more preferred embodiment of the present invention, in the compounds having general chemical Formula I, the spacer -Y- is a non-amino acid moiety selected from the group comprising

-NH-(CH₂)_p-CO-, wherein p is an integer of from 2 to 10,

-NH-(CH₂-CH₂-O)_q-CH₂-CH₂-CO-, wherein q is an integer of from 0 to 5.

25 -NH-cycloalkyl-CO- wherein cycloalkyl is selected from C₅-C₈ cycloalkyl, more preferably C₆ atom cycloalkyl, and

-NH-heterocycloalkyl-(CH₂)_v-CO- wherein heterocycloalkyl is selected from C₅-C₈ heterocycloalkyl containing carbon atoms and 1, 2, 3 or 4 oxygen, nitrogen or sulfur heteroatoms more preferably 1 to 2 heteroatom even more preferably 1 heteroatom and v

30 is an integer of from 1 to 4, more preferably v is an integer of from 1 to 2

U is a targeting agent.

For the purposes of the present invention, the term "targeting agent" shall have the following meaning: The targeting agent is a compound or moiety that targets or directs the radionuclide attached to it to a specific site in a biological system. A targeting agent can be any compound or chemical entity that binds to or accumulates at a target site in a mammalian body, *i.e.*, the compound localizes to a greater extent at the target site than to surrounding tissue.

The compounds of this invention are useful for the imaging of a variety of cancers including but not limited to: carcinoma such as bladder, breast, colon, kidney, liver, lung, including small cell lung cancer, esophagus, gall-bladder, ovary, pancreas, stomach, cervix, thyroid, prostate and skin, hematopoietic tumors of lymphoid and myeloid lineage, tumors of mesenchymal origin, tumors of central peripheral nervous systems, other tumors, including melanoma, seminoma, teratocarcinoma, osteosarcoma, xeroderma pigmentosum, keratoxanthoma, thyroid follicular cancer and Kaposi's sarcoma.

Most preferably, the use is not only for imaging of tumors, but also for imaging of inflammatory and/or neurodegenerative diseases, such as multiple sclerosis or Alzheimer's disease, or imaging of angiogenesis-associated diseases, such as growth of solid tumors, and rheumatoid arthritis.

Preferably the targeting agent **U** is a peptide, a peptidomimetic, or an oligonucleotide, particularly one which has specificity to target the complex to a specific site in a biological system. Small molecules effective for targeting certain sites in a biological system can also be used as the targeting agent.

Small molecules effective for targeting certain sites in a biological system can be used as the targeting agent **U**. Smaller organic molecules may be "small chemical entities". As used in this application, the term "small chemical entity" shall have the following meaning: A small chemical entity is a compound that has a molecular mass of from 200 to 800 or of from 150 to 700, more preferably from 200 to 700, more preferably from 250 to 700, even more preferably from 300 to 700, even more preferably from 350 to 700 and most preferably from 400 to 700. A small chemical entity as used herein may further contain at least one aromatic or heteroaromatic ring and may also have a primary or secondary amine, a thiol or hydroxyl group coupled via which the benzene ring structure in the compounds of general chemical Formulae I and II is coupled via -L-Y-. Such targeting moieties are known in the art, so are methods for preparing them.

- The small molecule targeting agents may preferably be selected from those described in the following references: P.L.Jager, M.A.Korte, M.N.Lub-de Hooge, A. van Waarde, K.P.Koopmans, P.J.Perik and E.G.E. de Vries, *Cancer Imaging*, (2005) 5, 27-32; W.D.Heiss and K.Herholz, *J. Nucl. Med.*, (2006) 47(2), 302-312; and T.Higuchi and M.Schwaiger, *Curr. Cardiol. Rep.*, (2006) 8(2), 131-138. More specifically examples of
- 5 small molecule targeting agents are listed hereinafter:

Name	Abbr.	target
18F-2b-Carbomethoxy-3b-(4-fluorophenyl)tropane	CFT	DAT (dopamine transporter)
18F-Fluoroethylspiperone	FESP	D2 (dopamine 2 receptor), 5-HT ₂ (5-hydroxytryptamine receptor)
18F-Fallypride		D2 (dopamine 2 receptor)
18F-Altanserlin		5-HT _{2A} receptor
18F-Cyclofoxy		Opioid receptors
18F-CPFPX		Adenosine A1 receptor
Batimastat		MMP
Fatty acids and analogues		
Choline analogues (metabolism)		
Flumazenil		Benzodiazepine receptors
Raclopride		D2 receptors
Dihydrotestosterone and analogues		AR
Tamoxifen and analogues		
Deoxyglucose		
Thymidine		Proliferation marker- thymidine kinase
DOPA		
Benzazepines		D ₁ antagonists
N-methyl spiperone and derivatives thereof		dopamine receptors
Benzamide raclopride; benzamide derivatives, e.g., fallopride, iodo benzamide; clozapine, quetiapine		D ₂ receptors
Nomifensine, substituted analogs of cocaine, e.g., tropane type derivatives of cocaine, methyl phenidate		DAT

2 β -Carboxymethoxy-3 β -(4-iodophenyl)tropane	CIT	DAT
	CIT-FE, CIT-FM	DAT
Altanserin, setoperon, ketanserin		5-HT _{2A}
	McN5652, 403U76 derivative ADAM, DASP, MADAM	5-HTT
Acetylcholine analogues	MP3A, MP4A, PMP; QNB, TKB, NMPB,	acetylcholine receptors
Scopolamine, benztropine		acetylcholine receptors
Flumazenil		GABA receptor
	RO-15-4513, FDG	GABA receptor
	PK-11195	benzodiazepine receptor
Xanthine analogues	CPFPX, MPDX	adenosine receptor
Carfentanyl, diprenorphine		opoid receptor

Further various small molecule targeting agents and the targets thereof are given in Table 1 in W.D.Heiss and K.Herholz, *ibid.* and in Figure 1 in T.Higuchi, M.Schwaiger, *ibid.*

- 5 Further preferred biomolecules are sugars, oligosaccharides, polysaccharides, aminoacids, nucleic acids, nucleotides, nucleosides, oligonucleotides, proteins, peptides, peptidomimetics, antibodies, aptamers, lipids, hormones (steroid and nonsteroid), neurotransmitters, drugs (synthetic or natural), receptor agonists and antagonists, dendrimers, fullerenes, virus particles and other targeting molecules / biomolecules (*e.g.*,
10 cancer targeting molecules).

Preferably the targeting agent **U** is a peptide.

- 15 The targeting agent **U** may be a peptide comprising from 4 to 100 amino acids wherein the amino acids may be selected from natural and non-natural amino acids and also may comprise modified natural and non-natural amino acids.

- 20 Examples for peptides as targeting agent targeting agent **U** are, but are not limited to, somatostatin and derivatives thereof and related peptides, somatostatin receptor specific peptides, neuropeptide Y and derivatives thereof and related peptides, neuropeptide Y₁ and the analogs thereof, bombesin and derivatives thereof and related peptides, gastrin, gastrin releasing peptide and the derivatives thereof and related peptides, epidermal

growth factor (EGF of various origin), insulin growth factor (IGF) and IGF-1, integrins ($\alpha_3\beta_1$, $\alpha_v\beta_3$, $\alpha_v\beta_5$, $\alpha_{IIb}\beta_3$), LHRH agonists and antagonists, transforming growth factors, particularly TGF- α ; angiotensin; cholecystokinin receptor peptides, cholecystokinin (CCK) and the analogs thereof; neurotensin and the analogs thereof, thyrotropin releasing hormone, pituitary adenylate cyclase activating peptide (PACAP) and the related peptides thereof, chemokines, substrates and inhibitors for cell surface matrix metalloproteinase, prolactin and the analogs thereof, tumor necrosis factor, interleukins (IL-1, IL-2, IL-4 or IL-6), interferons, vasoactive intestinal peptide (VIP) and the related peptides thereof. Such peptides comprise from 4 to 100 amino acids, wherein the amino acids are selected from natural and non-natural amino acids and also comprise modified natural and non-natural amino acids. Preferably targeting agent **U** is not insulin.

More preferably targeting agent **U** may be selected from the group comprising bombesin and bombesin analogs, preferably those having the sequences listed herein below, somatostatin and somatostatin analogs, preferably those having the sequences listed herein below, neuropeptide Y_1 and the analogs thereof, preferably those having the sequences listed herein below, vasoactive intestinal peptide (VIP) and the analogs thereof.

Even more preferably targeting agent **U** may be selected from the group comprising bombesin, somatostatin, neuropeptide Y_1 and the analogs thereof.

Even more preferably targeting agent **U** may be bombesin and the analogs thereof.

Bombesin is a fourteen amino acid peptide that is an analog of human Gastrin releasing peptide (GRP) that binds with high specificity to human GRP receptors present in prostate tumor, breast tumor and metastasis. Accordingly, as to a ninth aspect of the present invention, bombesin analogs are provided.

In a more preferred embodiment, bombesin analogs have the following sequence having Formula III:

$AA_1-AA_2-AA_3-AA_4-AA_5-AA_6-AA_7-AA_8-NT_1T_2$ (type A) Formula III, with:

$T_1 = T_2 = H$ or $T_1 = H, T_2 = OH$ or $T_1 = CH_3, T_2 = OH$

$AA_1 = \text{Gln, Asn, Phe(4-CO-NH}_2\text{)}$

$AA_2 = \text{Trp, D-Trp}$

$AA_3 = \text{Ala, Ser, Val}$

AA₄ = Val, Ser, Thr

AA₅ = Gly, (N-Me)Gly

AA₆ = His, His(3-Me), (N-Me)His, (N-Me)His(3-Me)

5 AA₇ = Sta, Statine analogs and isomers, 4-Am,5-MeHpA, 4-Am,5-MeHxA, γ-substituted aminoacids

AA₈ = Leu, Cpa, Cba, CpnA, Cha, t-buGly, tBuAla, Met, Nle, iso-Bu-Gly

In a more preferred embodiment, bombesin analogs have the following sequence of formula IV:

10 AA₁-AA₂-AA₃-AA₄-AA₅-AA₆-AA₇-AA₈-NT₁T₂ (type B) Formula IV, with:

T₁ = T₂ = H or T₁ = H, T₂ = OH or T₁ = CH₃, T₂ = OH


AA₁ = Gln, Asn or Phe(4-CO-NH₂)

AA₂ = Trp, D-Trp

AA₃ = Ala, Ser, Val

15 AA₄ = Val, Ser, Thr

AA₅ = βAla, β²- and β³-amino acids as shown herein after



20 wherein SC represents a side chain found in proteinogenic amino acids and homologs of proteinogenic amino acids,

AA₆ = His, His(3-Me), (N-Me)His, (N-Me)His(3-Me)

AA₇ = Phe, Tha, Nal,

AA₈ = Leu, Cpa, Cba, CpnA, Cha, t-buGly, tBuAla, Met, Nle, iso-Bu-Gly.

25 Therefore, in an even more preferred embodiment of the present invention targeting agent U may be selected from the group comprising bombesin analogs having sequence III or IV.

In a more preferred embodiment, bombesin analogs have the following sequences:

30

• Seq ID

P

• Seq ID 1 Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂

• Seq ID 2 Gln-Trp-Ala-Val-Gly-His(Me)-Sta-Leu-NH₂

- Seq ID 3 Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂
- Seq ID 4 Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂
- Seq ID 7 Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa-NH₂
- Seq ID 8 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂
- 5 • Seq ID 12 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂
- Seq ID 17 Gln-Trp-Ala-Val-Gly-His-4-Am,5-MeHpA-Leu-NH₂
- Seq ID 23 Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Cpa-NH₂
- Seq ID 27 Gln-Trp-Ala-Val-NMeGly-His-FA02010-Cpa-NH₂
- Seq ID 28 Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-tbuGly-NH₂
- 10 • Seq ID 30 Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-tBuGly-NH₂
- Seq ID 32 Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂
- Seq ID 33 Gln-DTrp-Ala-Val-Gly-His-4-Am,5-MeHpA-tbuGly-NH₂
- Seq ID 34 Gln-DTrp-Ala-Val-Gly-His-4-Am-5-MeHxA-Cpa-NH₂
- Seq ID 35 Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa-NH₂
- 15 • Seq ID 36 Gln-DTrp-Ala-Val-Gly-His-Sta-tbuAla-NH₂
- Seq ID 42 Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Cpa-NH₂
- Seq ID 43 Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-tBuGly-NH₂
- Seq ID 46 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂
- Seq ID 48 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂
- 20 • Seq ID 49 Gln-Trp-Ala-Val-Gly-NMeHis-4-Am,5-MeHpA-Cpa-NH₂
- Seq ID 49 Gln-Trp-Ala-Val-Gly-NMeHis(3Me)-4-Am,5-MeHpA-Leu-NH₂
- Seq ID 50 Gln-Trp-Ala-Val-Gly-NMeHis-4-Am,5-MeHpA-Leu-NH₂
- Seq ID 51 Gln-Trp-Ala-Val-NMeGly-His-AHMHxA -Leu-NH₂
- Seq ID 52 Gln-Trp-Ala-Val-βAla-NMeHis-Tha-Cpa-NH₂
- 25 • Seq ID 53 Gln-Trp-Ala-Val-βAla-NMeHis-Phe-Cpa-NH₂
- Seq ID 54 Gln-Trp-Ala-Val-βAla-NMeHis-Phe-Leu-NH₂
- Seq ID 55 Gln-Trp-Ala-Val-βAla-DHis-Phe-Leu-NH₂
- Seq ID 56 Gln-Trp-Ala-Val-βAla-His-βhLeu-Leu-NH₂
- Seq ID 57 Gln-Trp-Ala-Val-βAla-His-βhIle-Leu-NH₂
- 30 • Seq ID 58 Gln-Trp-Ala-Val-βAla-His-βhLeu-tbuGly-NH₂
- Seq ID 59 Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Tha-NH₂
- Seq ID 60 Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Nle-NH₂
- Seq ID 61 Gln-Trp-Ala-Val-βAla-NMeHis-Phe-tbuGly-NH₂
- Seq ID 62 Gln-Trp-Ala-Val-βAla-NMeHis-Tha-tbuGly-NH₂
- 35 • Seq ID 63 Gln-Trp-Ala-Val-βAla-His(3Me)-Tha-tbuGly-NH₂

- Seq ID 64 Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Cpa-NH₂
- Seq ID 65 Gln-Trp-Ala-NMeVal-βAla-His-Phe-Leu-NH₂
- Seq ID 66 Gln-Trp-Ala-Val-βAla-His-NMePhe-Leu-NH₂
- Seq ID 67 Gln-DTrp-Ala-Val-βAla-His-Phe-Leu-NH₂
- 5 • Seq ID 68 Gln-Trp-DAla-Val-βAla-His-Phe-Leu-NH₂
- Seq ID 69 Gln-Trp-Ala-DVal-βAla-His-Phe-Leu-NH₂
- Seq ID 70 Gln-Trp-Ala-Val-βAla-His-DPhe-Leu-NH₂
- Seq ID 71 Gln-Trp-Ala-Val-βAla-His-βHle-tbuGly-NH₂
- Seq ID 72 Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-Cpa-NH₂
- 10 • Seq ID 73 Gln-Trp-Ala-Val-NMeGly-His-Sta-Cpa-NH₂
- Seq ID 74 Gln-Trp-Ala-Val-NMeGly-His-Sta-tbuAla-NH₂
- Seq ID 75 Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-tbuAla-NH₂
- Seq ID 77 Gln-Trp-Ala-Val-His(Me)-Sta-Leu-NH₂
- Seq ID 82 Gln-Trp-Ala-Val-Gly-His(3Me)-FA4-Am,5-MeHpA-Leu-NH₂
- 15 • Seq ID 90 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂
- Seq ID 91 Gln-Trp-Ala-Val-Gly-His-4-Am,5-MeHpA-Leu-NH₂
- Seq ID 101 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am-5-MeHpA – 4-amino-5-methylheptanoic acid -Leu-NH₂
- Seq ID 102 Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am-5-MeHpA – 4-amino-5-methylheptanoic acid -Cpa-NH₂
- 20

Thus, the invention also refers to bombesin analogs that bind specifically to human GRP receptors present in prostate tumor, breast tumor and metastasis. In a preferred embodiment, the bombesin analogs are peptides having sequences from Seq ID 1 to Seq ID 102 and preferably have one of them. More preferably a bombesin analog is additionally labeled with a fluorine isotope (F) wherein fluorine isotope (F) is selected from ¹⁸F or ¹⁹F. More preferably the bombesin analog is radiolabeled with ¹⁸F. The bombesin analog is preferably radiolabeled using the radiofluorination method of the present invention.

30

In a more preferred embodiment, somatostatin analogs have the following sequences:

- Seq ID 104 ----c[Lys-(NMe)Phe-1Nal-D-Trp-Lys-Thr]
- Seq ID 105 ----c[Dpr-Met-(NMe)Phe-Tyr-D-Trp-Lys]

35

In a more preferred embodiment, neuropeptide Y₁ analogs have the following sequences:

- Seq ID 106 -DCys-Leu-Ile-Thr-Arg-Cys-Arg-Tyr-NH₂
- Seq ID 107 -DCys-Leu-Ile-Val-Arg-Cys-Arg-Tyr-NH₂

5 (_ indicates disulfide bridge)

In a more preferred embodiment, peptide is tetrapeptide of the following sequences:

valyl-β-alanyl-phenylalanyl-glycine amide

valyl-β-alanyl-histidyl(π-Me)-glycine amide

10

In a further preferred embodiment targeting agent (P) may be selected from the group comprising oligonucleotides comprising from 4 to 100 nucleotides.

In other preferred embodiments the targeting agent U is selected to be an oligonucleotide.

15 In a further preferred embodiment the targeting agent U may be selected from the group comprising oligonucleotides comprising from 4 to 100 nucleotides.

Preferred as targeting agents U are peptides comprising from 4 to 100 amino acids or oligonucleotides comprising from 4 to 100 nucleotides.

20 Preferred oligonucleotide is TTA1 (see experimental part).

In a further preferred embodiment of the present invention, the targeting agent U may comprise a combination of any of the aforementioned bioactive molecules suitable to bind to a target site together with a reacting moiety which serves the linking between the

25 bioactive molecule and the rest of the compound of the invention (Formulae I, II, III), wherein reacting moiety is selected from -NR⁷, -NR⁷-(CH₂)_n-, -O-(CH₂)_n- or -S-(CH₂)_n-, wherein R⁷ is hydrogen or unbranched or branched alkyl and n is an integer from 1 to 6 and wherein the suitable bioactive molecule is selected from peptide, peptidomimetic, oligonucleotide, or small molecule.

30

In a preferred embodiment of R⁷ the alkyl is unbranched or branched C₁-C₆ alkyl, more preferably a methyl.

35 In a preferred embodiment U is NR⁷-peptide, or -(CH₂)_n-peptide, -O-(CH₂)_n- peptide or -S-(CH₂)_n- peptide, NR⁷- small-molecule, or -(CH₂)_n- small-molecule, -O-(CH₂)_n- small-molecule or -S-(CH₂)_n- small-molecule, NR⁷- oligonucleotide, or -(CH₂)_n- oligonucleotide, -

O-(CH₂)_n- oligonucleotide or -S-(CH₂)_n- oligonucleotide - wherein n is an integer of from 1 to 6.

In a more preferred embodiment **U** is -NR⁷-peptide, -(CH₂)_n-peptide, wherein n is an integer of from 1 to 6.

In another more preferred embodiment **U** is -NR⁷-oligonucleotide or -(CH₂)_n-oligonucleotide, wherein n is an integer of from 1 to 6.

In another more preferred embodiment **U** is -NR⁷-small-molecule or -(CH₂)_n-small molecule, wherein n is an integer of from 1 to 6.

In a preferred embodiment, the precursor for a single step radiolabeling is selected from the following list wherein **U** is a bombesin analog:

- Ia-1 4-(Trimethylammonium)-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂,
- Ia-2 4-(Trimethylammonium)-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(Me)-Sta-Leu-NH₂,
- Ia-3 4-(Trimethylammonium)-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- Ia-4 4-(Trimethylammonium)-3-cyano-benzoyl-1,4-cis-Achc-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-5 4-(Trimethylammonium)-3-cyano-benzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-6 4-(Trimethylammonium)-3-cyano-benzoyl-AOC-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-7 4-(Trimethylammonium)-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa-NH₂,
- Ia-8 4-(Trimethylammonium)-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- Ia-9 4-(Trimethylammonium)-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-10 4-(Trimethylammonium)-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,

- Ia-11 4-(Trimethylammonium)-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-12 4-(Trimethylammonium)-3-cyano-benzoyl-Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- 5 • Ia-13 4-(Trimethylammonium)-3-cyano-benzoyl-Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- Ia-14 4-(Trimethylammonium)-3-cyano-benzoyl-Lys(Me)₂-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-AM-5-MeHpA-Leu-NH₂,
- Ia-15 4-(Trimethylammonium)-3-cyano-benzoyl-Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 10 • Ia-16 4-(Trimethylammonium)-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- Ia-17 4-(Trimethylammonium)-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His-4-Am,5-MeHpA--Leu-NH₂,
- 15 • Ia-18 4-(Trimethylammonium)-3-trifluoromethyl-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-19 4-(Trimethylammonium)-3-trifluoromethyl-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- Ia-20 4-(Trimethylammonium)-3-trifluoromethyl-benzoyl-1,4-cis-Achc-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 20 • Ia-21 4-(Trimethylammonium)-3-trifluoromethyl-benzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-22 4-(Trimethylammonium)-3-trifluoromethyl-benzoyl-Arg-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- 25 • Ia-23 4-(Trimethylammonium)-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Cpa-NH₂,
- Ia-24 4-(Trimethylammonium)-3-cyano-benzoyl -Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-25 4-(Trimethylammonium)-3-cyano-benzoyl-DOA-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 30

In a preferred embodiment, the precursor for a single step radiolabeling is selected from the following list wherein U is somatostatin analogs:

- 1a-66: 4-(Trimethylammonium)-3-cyano-benzoyl-Ava-ε-c[Lys-(NMe)Phe-1Nal-D-Trp-Lys-Thr]
- 1a-67: 4-(Trimethylammonium)-3-cyano-benzoyl-Ava-β-c[Dpr-Met-(NMe)Phe-Tyr-D-Trp-Lys]

5

In a preferred embodiment, the precursor for a single step radiolabeling is selected from the following list wherein U is neuropeptide Y₁ analogs:


- 1a-68: 4-(Trimethylammonium)-3-cyano-benzoyl-Ava-DCys-Leu-Ile-Thr-Arg-Cys-Arg-Tyr-NH₂]
- 1a-69: 4-(Trimethylammonium)-3-cyano-benzoyl-Ava-DCys-Leu-Ile-Val-Arg-Cys-Arg-Tyr-NH₂]

10


In a preferred embodiment, the precursor for a single step radiolabeling is selected from the following list wherein U is a small molecule:

15


20




25



30





35



The $N^+(R^1)(R^2)(R^3)X^-$ group attached to the benzene ring can be displaced with fluorine isotope, to provide a chemically and biologically stable bond,

In a **second aspect** the present invention refers to compounds having general chemical Formula A (general chemical Formula II) wherein $K = W$:

It has been found that compounds according to Formula I can be ^{18}F -labeled surprisingly in a one-step radiofluorination reaction in order to arrive at compounds according to Formula A (general chemical Formula II), wherein $K = W$:



A

II

wherein the residues and substituents K , $N^+(R^1)(R^2)(R^3)$, X^- , $-G$, $-Q$, $-L-$, $-Y-$ and $-U$ have the same meaning as depicted above for compounds having general chemical Formula I. This includes in particular all preferred embodiments mentioned above with regard to the residues and substituents K , R^1 , R^2 , R^3 , W , X^- , $-G$, $-Q$, $-L-$, $-Y-$ and $-U$ and all residues used to define these residues and substituents, such as R^4 , R^5 and the like.

W is a fluorine isotope, preferably radioactive or non-radioactive ("cold") fluorine isotope, more preferably W is ^{18}F or ^{19}F , even more preferably W is ^{18}F .

Surprisingly, It has been found that when Q or G are selected from the group of substituents listed in Chem Rev. (1999), Vol. 42, No. 9, 165-195, Table1 and have a σ_m and σ_p value < 0 for compound of formula I then one-step labelling is less or not suited for good or high radio-chemical yields.


The invention further refers to pharmaceutically acceptable salts of inorganic or organic acids, hydrates, complexes, esters, amides, solvates and prodrugs of the compounds having general chemical Formula II.

Unless otherwise specified, when referring to the compounds having general chemical Formula I per se as well as to any pharmaceutical composition thereof the present invention includes all of the salts of inorganic or organic acids thereof, hydrates, complexes, esters, amides, solvates and prodrugs of the compounds of the invention.

W is a fluorine isotope, preferably radioactive or non-radioactive ("cold") fluorine isotope, more preferably W is ^{18}F or ^{19}F , even more preferably W is ^{18}F .


If W is ^{18}F , the compound of the invention having general chemical Formula II being radiopharmaceutically labelled with ^{18}F has the following general chemical Formula IIA:

41



IIA

More preferably, when $W = {}^{19}\text{F}$ then the compound having general chemical Formula II
 5 being radiopharmaceutically labelled with ${}^{19}\text{F}$ has general chemical Formula IIB:



IIB

10 In a preferred embodiment of compounds of Formula II, -G and -Q are independently from each other selected from -H, -CN, CF_3 , and -Cl.

In a more preferred embodiment -G and -Q are independently from each other H, - CF_3 , or CN.

15

In a even more preferred embodiment In a more preferred embodiment -G and -Q are independently from each other H, - CF_3 , or -CN, whereas at least -G or -Q is - CF_3 or -CN.

20 In a preferred embodiment, the radiopharmaceutical compound labelled with ${}^{18}\text{F}$ or ${}^{19}\text{F}$ is selected from the following list, wherein U is a bombesin analog:

- IIA-a-1 4-[${}^{18}\text{F}$]Fluoro-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂,
- 25 • IIA-a-2 4-[${}^{18}\text{F}$]Fluoro-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(Me)-Sta-Leu-NH₂,
- IIA-a-3 4-[${}^{18}\text{F}$]Fluoro-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,

- IIA-a-4 4-[18]Fluoro-3-cyano-benzoyl-1,4-cis-Achc-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIA-a-5 4-[18]Fluoro-3-cyano-benzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIA-a-6 4-[18]Fluoro-3-cyano-benzoyl-AOC-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 5 • IIA-a-7 4-[18]Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa-NH₂,
- IIA-a-8 4-[18]Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-FA4-Am,5-MeHpA-Leu-NH₂,
- 10 • IIA-a-9 4-[18]Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIA-a-10 4-[18]Fluoro-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIA-a-11 4-[18]Fluoro-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 15 • IIA-a-12 4-[18]Fluoro-3-cyano-benzoyl-Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIA-a-13 4-[18]Fluoro-3-cyano-benzoyl-Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- 20 • IIA-a-14 4-[18]Fluoro-3-cyano-benzoyl-Lys(Me)₂-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIA-a-15 4-[18]Fluoro-3-cyano-benzoyl-Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIA-a-16 4-[18]Fluoro-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- 25 • IIA-a-17 4-[18]Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His-4-Am,5-MeHpA-Leu-NH₂,
- IIA-a-18 4-[18]Fluoro-3-trifluoromethyl-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-LeuNH₂,
- 30 • IIA-a-19 4-[18]Fluoro-3-trifluoromethyl-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- IIA-a-20 4-[18]Fluoro-3-trifluoromethyl-benzoyl-1,4-cis-Achc-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIA-a-21 4-[18]Fluoro-3-trifluoromethyl-benzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂ ,
- 35

- IIA-a-22 4-[18]Fluoro-3-trifluoromethyl-benzoyl-Arg-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-23 4-[18]-Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Cpa-NH₂,
- 5 • IIB-a-24 4-[18]-Fluoro-3-cyano-benzoyl -Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-25 4-[18]-Fluoro-3-cyano-benzoyl-DOA-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-26 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂,
- 10 • IIB-a-27 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His-FA02010-Cpa-NH₂,
- IIB-a-28 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-tbuGly-NH₂,
- IIB-a-29 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-30 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-tBuGly-NH₂,
- 15 • IIB-a-31 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-32 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-33 3,4-[18]-Difluorobenzoyl-Ava-Gln-DTrp-Ala-Val-Gly-His-4-Am,5-MeHpA-tbuGly-NH₂,
- 20 • IIB-a-34 3,4-[18]-Difluorobenzoyl-Ava-Gln-DTrp-Ala-Val-Gly-His-4-Am-5-MeHxA-Cpa-NH₂,
- IIB-a-35 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa-NH₂,
- IIB-a-36 3,4-[18]-Difluorobenzoyl-Ava-Gln-DTrp-Ala-Val-Gly-His-Sta-tbuAla-NH₂,
- 25 • IIB-a-37 3,4-[18]-Difluorobenzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂,
- IIB-a-38 3,4-[18]-Difluorobenzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-39 3,4-[18]-Difluorobenzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-40 3,4-[18]-Difluorobenzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 30 • IIB-a-41 3,4-[18]-Difluorobenzoyl-Arg-βAla-Arg-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-42 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Cpa-NH₂,
- IIB-a-43 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-tBuGly-NH₂,
- IIB-a-44 3,4-[18]-Difluorobenzoyl-Arg-Arg-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- 35

- IIB-a-45 3,4-[18]-Difluorobenzoyl-Arg-βAla-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-46 3,4-[18]-Difluorobenzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- 5 • IIB-a-47 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Cpa-NH₂,
- IIB-a-48 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-49 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-NMeHis-4-Am,5-MeHpA-Cpa-NH₂,
- 10 • IIB-a-49 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-NMeHis(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-50 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-NMeHis-4-Am,5-MeHpA-Leu-NH₂,
- 15 • IIB-a-51 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His-AHMHxA -Leu-NH₂,
- IIB-a-52 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Tha-Cpa-NH₂,
- IIB-a-53 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Phe-Cpa-NH₂,
- IIB-a-54 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Phe-Leu-NH₂,
- 20 • IIB-a-55 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-DHis-Phe-Leu-NH₂,
- IIB-a-56 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-βhLeu-Leu-NH₂,
- IIB-a-57 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-βhIle-Leu-NH₂,
- IIB-a-58 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-βhLeu-tbuGly-NH₂,
- IIB-a-59 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Tha-NH₂,
- 25 • IIB-a-60 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Nle-NH₂,
- IIB-a-61 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Phe-tbuGly-NH₂,
- IIB-a-62 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Tha-tbuGly-NH₂,
- IIB-a-63 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His(3Me)-Tha-tbuGly-NH₂,
- IIB-a-64 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Cpa-NH₂,
- 30 • IIB-a-65 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-NMeVal-βAla-His-Phe-Leu-NH₂,
- IIB-a-66 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-NMePhe-Leu-NH₂,
- IIB-a-67 3,4-[18]-Difluorobenzoyl-Ava-Gln-DTrp-Ala-Val-βAla-His-Phe-Leu-NH₂,
- IIB-a-68 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-DAla-Val-βAla-His-Phe-Leu-NH₂,
- IIB-a-69 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-DVal-βAla-His-Phe-Leu-NH₂,
- 35 • IIB-a-70 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-DPhe-Leu-NH₂,

- IIB-a-71 3,4-[18]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-βhIle-tbuGly-NH₂,
- IIB-a-72 4-[18]-Fluoro-3-cyano-phenylsulfonyl -Ava-Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-Cpa-NH₂,
- IIB-a-73 4-[18]-Fluoro-3-cyano-phenylsulfonyl -Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-
5 Cpa-NH₂,
- IIB-a-74 4-[18]-Fluoro-3-cyano-phenylsulfonyl -Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-tbuAla-NH₂,
- IIB-a-75 4-[18]-Fluoro-3-cyano-phenylsulfonyl -Ava-Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-tbuAla-NH₂.
- 10 • 4-[18]Fluoro-3-cyano-benzoyl-(piperidyl-4-carbonyl)-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 4-[18]Fluoro-3-cyano-benzoyl-(piperazin-1-yl-acetyl)-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 4-[18]Fluoro-3-cyano-benzoyl-1,4-trans-Achc-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-
15 NH₂,
- IIB-a-1 4-[19]-Fluoro-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-
20 NH₂,
- IIB-a-2 4-[19]-Fluoro-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-His(Me)-Sta-Leu-NH₂,
- IIB-a-3 4-[19]-Fluoro-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-4 4-[19]-Fluoro-3-cyano-benzoyl-1,4-cis-Achc-Gln-Trp-Ala-Val-Gly-His(3Me)-
25 Sta-Leu-NH₂,
- IIB-a-5 4-[19]-Fluoro-3-cyano-benzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-6 4-[19]-Fluoro-3-cyano-benzoyl-AOC-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-7 4-[19]-Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-
30 Cpa-NH₂,
- IIB-a-8 4-[19]-Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-9 4-[19]-Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,

- IIB-a-10 4-[19]-Fluoro-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-11 4-[19]-Fluoro-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 5 • IIB-a-12 4-[19]-Fluoro-3-cyano-benzoyl-Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-13 4-[19]-Fluoro-3-cyano-benzoyl-Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-14 4-[19]-Fluoro-3-cyano-benzoyl-Lys(Me)₂-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-10 Am,5-MeHpA-Leu-NH₂,
- IIB-a-15 4-[19]-Fluoro-3-cyano-benzoyl-Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-16 4-[19]-Fluoro-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- 15 • IIB-a-17 4-[19]-Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-18 4-[19]-Fluoro-3-trifluoromethyl-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-19 4-[19]-Fluoro-3-trifluoromethyl-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,20
- IIB-a-20 4-[19]-Fluoro-3-trifluoromethyl-benzoyl-1,4-cis-Achc-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-21 4-[19]-Fluoro-3-trifluoromethyl-benzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 25 • IIB-a-22 4-[19]-Fluoro-3-trifluoromethyl-benzoyl-Arg-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-23 4-[19]-Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Cpa-NH₂,
- IIB-a-24 4-[19]-Fluoro-3-cyano-benzoyl -Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,30
- IIB-a-25 4-[19]-Fluoro-3-cyano-benzoyl-DOA-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-26 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂,
- IIB-a-27 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His-FA02010-Cpa-NH₂,

- IIB-a-28 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-tbuGly-NH₂,
- IIB-a-29 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-30 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-tBuGly-NH₂,
- IIB-a-31 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-32 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-33 3,4-[19]-Difluorobenzoyl-Ava-Gln-DTrp-Ala-Val-Gly-His-4-Am,5-MeHpA-tbuGly-NH₂,
- IIB-a-34 3,4-[19]-Difluorobenzoyl-Ava-Gln-DTrp-Ala-Val-Gly-His-4-Am-5-MeHxA-Cpa-NH₂,
- IIB-a-35 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa-NH₂,
- IIB-a-36 3,4-[19]-Difluorobenzoyl-Ava-Gln-DTrp-Ala-Val-Gly-His-Sta-tbuAla-NH₂,
- IIB-a-37 3,4-[19]-Difluorobenzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂,
- IIB-a-38 3,4-[19]-Difluorobenzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-39 3,4-[19]-Difluorobenzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-40 3,4-[19]-Difluorobenzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-41 3,4-[19]-Difluorobenzoyl-Arg-βAla-Arg-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-42 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Cpa-NH₂,
- IIB-a-43 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-tBuGly-NH₂,
- IIB-a-44 3,4-[19]-Difluorobenzoyl-Arg-Arg-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-45 3,4-[19]-Difluorobenzoyl-Arg-βAla-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-46 3,4-[19]-Difluorobenzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-47 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Cpa-NH₂,
- IIB-a-48 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-49 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-NMeHis-4-Am,5-MeHpA-Cpa-NH₂,

- IIB-a-49 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-NMeHis(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-50 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-NMeHis-4-Am,5-MeHpA-Leu-NH₂,
- 5 • IIB-a-51 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His-AHMHxA -Leu-NH₂,
- IIB-a-52 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Tha-Cpa-NH₂,
- IIB-a-53 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Phe-Cpa-NH₂,
- IIB-a-54 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Phe-Leu-NH₂,
- 10 • IIB-a-55 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-DHis-Phe-Leu-NH₂,
- IIB-a-56 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-βhLeu-Leu-NH₂,
- IIB-a-57 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-βhIle-Leu-NH₂,
- IIB-a-58 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-βhLeu-tbuGly-NH₂,
- IIB-a-59 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Tha-NH₂,
- 15 • IIB-a-60 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Nle-NH₂,
- IIB-a-61 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Phe-tbuGly-NH₂,
- IIB-a-62 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Tha-tbuGly-NH₂,
- IIB-a-63 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His(3Me)-Tha-tbuGly-NH₂,
- IIB-a-64 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Cpa-NH₂,
- 20 • IIB-a-65 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-NMeVal-βAla-His-Phe-Leu-NH₂,
- IIB-a-66 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-NMePhe-Leu-NH₂,
- IIB-a-67 3,4-[19]-Difluorobenzoyl-Ava-Gln-DTrp-Ala-Val-βAla-His-Phe-Leu-NH₂,
- IIB-a-68 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-DAla-Val-βAla-His-Phe-Leu-NH₂,
- IIB-a-69 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-DVal-βAla-His-Phe-Leu-NH₂,
- 25 • IIB-a-70 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-DPhe-Leu-NH₂,
- IIB-a-71 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-βhIle-tbuGly-NH₂,
- IIB-a-72 4-[19]-Fluoro-3-cyano-phenylsulfonyl -Ava-Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-Cpa-NH₂,
- IIB-a-73 4-[19]-Fluoro-3-cyano-phenylsulfonyl -Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Cpa-NH₂,
- 30 • IIB-a-74 4-[19]-Fluoro-3-cyano-phenylsulfonyl -Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-tbuAla-NH₂,
- IIB-a-75 4-[19]-Fluoro-3-cyano-phenylsulfonyl -Ava-Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-tbuAla-NH₂.

- 4-[19]Fluoro-3-cyano-benzoyl-(piperidyl-4-carbonyl)-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
 - 4-[19]Fluoro-3-cyano-benzoyl-(piperazin-1-yl-acetyl)-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 5 • 4-[19]Fluoro-3-cyano-benzoyl-1,4-trans-Achc-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂,


10 In a preferred embodiment, the radiopharmaceutical labelled with ¹⁸F or ¹⁹F is selected from the following list, wherein **U** is a somatostatin analog:

- IIA-a-76: 4-[18]Fluoro-3-cyano-benzoyl- Ava-ε-c[Lys-(NMe)Phe-1Nal-D-Trp-Lys-Thr]
 - IIA-a-77: 4-[18]Fluoro-3-cyano-benzoyl- Ava-β-c[Dpr-Met-(NMe)Phe-Tyr-D-Trp-Lys]
 - IIB-a-76: 4-[19]Fluoro-3-cyano-benzoyl- Ava-ε-c[Lys-(NMe)Phe-1Nal-D-Trp-Lys-Thr]
- 15 • IIB-a-77: 4-[19]Fluoro-3-cyano-benzoyl- Ava-β-c[Dpr-Met-(NMe)Phe-Tyr-D-Trp-Lys]


In a preferred embodiment, the radiopharmaceutical labelled with ¹⁸F or ¹⁹F is selected from the following list, wherein **U** is a neuropeptide Y₁ analog:

- 20 • IIA-a-78: 4-[18]Fluoro-3-cyano-benzoyl- Ava-DCys-Leu-Ile-Thr-Arg-Cys-Arg-Tyr-NH₂
- IIA-a-79: 4-[18]Fluoro-3-cyano-benzoyl- Ava- DCys-Leu-Ile-Val-Arg-Cys-Arg-Tyr-NH₂
- IIA-a-78: 4-[19]Fluoro-3-cyano-benzoyl- Ava-DCys-Leu-Ile-Thr-Arg-Cys-Arg-Tyr-NH₂
- IIA-a-79: 4-[19]Fluoro-3-cyano-benzoyl- Ava- DCys-Leu-Ile-Val-Arg-Cys-Arg-Tyr-NH₂


25 In a preferred embodiment, the radiopharmaceutical labelled with ¹⁸F or ¹⁹F is selected from the following list, wherein **U** is a small molecule:



3-Cyano-4-fluoro-N-{6-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-thiophen-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-hexyl}-benzamide




3-Cyano-4-fluoro-N-{4-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-ylcarbonyl]-butyl}-benzamide




3-Cyano-4-fluoro-N-{2-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-ylcarbonyl]-ethyl}-benzamide

5



10

3-Cyano-4-fluoro-N-{2-[2-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-ethanesulfonyl]-ethyl}-benzamide




5

3-Cyano-4-fluoro-N-(2-{2-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-ethanesulfonyl}-ethyl)-benzamide

10

15




20

3-Cyano-4-fluoro-N-(2-{2-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-ethylsulfanyl}-ethyl)-benzamide

25


30



35

3-Cyano-N-((2R,3R)-2,3-dihydroxy-4-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-butyl)-4-fluoro-benzamide


5



10

3-Cyano-4-[F]fluoro-N-(thymidiny-propyl)-benzamide


15



20


2-Chloro-4-fluoro-N-{3-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-propyl}-benzamide

25



30


3-Cyano-4-[F]fluoro-N-(2-[2-thymidiny-ethoxy]-ethyl)-benzamide





10 3-Cyano-4-fluoro-N-{6-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2,6-dioxo-5-trifluoromethyl-3,6-dihydro-2H-pyrimidin-1-yl]-hexyl}-benzamide

15

20



30 3-Cyano-4-fluoro-N-(2-{2-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-ethylamino}-ethyl)-benzamide



5

In a **fourth aspect**, the present invention also provides compositions comprising a compound having general chemical Formulae I or II or the pharmaceutically acceptable salts of inorganic or organic acids thereof, hydrates, complexes, esters, amides, solvates and prodrugs thereof and a pharmaceutically acceptable carrier, diluent, excipient or adjuvant. Pharmaceutically acceptable carriers, diluents, excipients or adjuvants may include any and all solvents, dispersion media, antibacterial and antifungal agents, isotonic agents, enzyme inhibitors, transfer ligands such as glucoheptonate, tartrate, citrate, or mannitol, and the like. Such compositions may be formulated as sterile, pyrogen-free, parenterally acceptable aqueous solution which may optionally be supplied in lyophilized form. The compositions of the invention may be provided as components of kits which may include buffers, additional vials, instructions for use, and the like.

In a **fifth aspect**, the present invention refers to a method of imaging diseases, the method comprising introducing into a patient a detectable quantity of a labeled compound having general chemical Formula A, wherein $K = W$, or a pharmaceutically acceptable salts of inorganic or organic acids thereof, hydrates, complexes, esters, amides, solvates and prodrugs thereof.

In a **sixth aspect**, the present invention provides a kit comprising a sealed vial containing a predetermined quantity of a compound having general chemical Formula I or the pharmaceutically acceptable salts of inorganic or organic acids thereof, hydrates,

complexes, esters, amides, solvates and prodrugs thereof and optionally a pharmaceutically acceptable carrier, diluent, excipient or adjuvant.

In a **seventh aspect** of the invention, compounds according to Formula II, including
5 compounds having general chemical Formulae IIA and IIB, are provided for use as medicament. Accordingly, the invention relates to compounds having general chemical Formula IIA for use as a positron emitting tomography (PET) diagnostic agent, wherein the fluorine isotope is ^{18}F and to compounds having general chemical Formula IIB for use as a precursor to prepare a positron emitting tomography (PET) diagnostic agent, wherein
10 the fluorine isotope is ^{19}F . More preferably, the invention relates to the use of compound having general chemical Formula I for the manufacture of compounds having general chemical Formula IIA as a diagnostic agent. Most preferably, the use is for imaging of tumors, imaging of inflammatory and/or neurodegenerative diseases, such as multiple sclerosis or Alzheimer's disease, or imaging of angiogenesis-associated diseases, such
15 as growth of solid tumors, and rheumatoid arthritis.

In another aspect, the present invention relates to compound having general chemical Formula II for use in biological assays and chromatographic identification. More preferably, the invention relates to compounds having general chemical Formula IIB for
20 use in biological assays and chromatographic identification, wherein the fluorine isotope is ^{19}F . More preferably, the invention relates to the use of compounds having general chemical Formula I for the manufacture of compounds having general chemical Formula IIB as a measurement agent.

25 In another aspect of the invention compounds according to Formula II are provided for use as diagnostic imaging agent, preferably as imaging agent for PET applications.

In an **eighth aspect** of the invention, compounds having general chemical Formulae I or II of pharmaceutically acceptable salts of inorganic or organic acids thereof, hydrates,
30 complexes, esters, amides, solvates and prodrugs thereof are provided for the use in manufacturing of a medicament, more specifically for the manufacture of a diagnostic imaging agent and most specifically for the manufacture of a diagnostic imaging agent for imaging tissue at a target site using the imaging agent.

The compounds of this invention are useful for the imaging of a variety of cancers including but not limited to: carcinoma such as bladder, breast, colon, kidney, liver, lung, including small cell lung cancer, esophagus, gall-bladder, ovary, pancreas, stomach, cervix, thyroid, prostate and skin, hematopoietic tumors of lymphoid and myeloid lineage, tumors of mesenchymal origin, tumors of central peripheral nervous systems, other tumors, including melanoma, seminoma, teratocarcinoma, osteosarcoma, xeroderma pigmentosum, keratoxanthoma, thyroid follicular cancer and Kaposi's sarcoma.

Most preferably, the use is for only for imaging of tumors, but also for imaging of inflammatory and/or neurodegenerative diseases, such as multiple sclerosis or Alzheimer's disease, or imaging of angiogenesis-associated diseases, such as growth of solid tumors, and rheumatoid arthritis.

In a preferred embodiment, the bombesin analog is a peptide of sequence from Seq ID 1 to Seq ID 102 and preferably one of them, as far as the compound having general chemical Formula A comprises bombesin or bombesin analogs, this compound binds specifically to human GRP receptors present in prostate tumor, breast tumor and metastasis. More preferably the bombesin analog is additionally radiolabeled with a fluorine isotope (F) wherein F is ^{18}F or ^{19}F . More preferably the bombesin analog is radiolabeled using the radiofluorination method of the present invention.

Therefore, according to a ninth aspect, the present invention refers to bombesin analogs that bind specifically to human GRP receptors present in prostate tumor, breast tumor and metastasis.

Further, the compounds having general chemical Formula II, in which W is ^{19}F or other non-radioactive ("cold") halogen elements may be used in biological assays and chromatographic identification. More preferably, the invention relates to the use of a compound having general chemical Formula I for the manufacture of a compound having general chemical Formula IIB as a measurement agent.

The radioactively labeled compounds according to Formula II provided by the invention may be administered intravenously in any pharmaceutically acceptable carrier, *e.g.*, conventional medium such as an aqueous saline medium, or in blood plasma medium, as a pharmaceutical composition for intravenous injection. Such medium may also contain conventional pharmaceutical materials such as, for example, pharmaceutically acceptable

salts to adjust the osmotic pressure, buffers, preservatives and the like. Among the preferred media are normal saline and plasma. Suitable pharmaceutical acceptable carriers are known to the person skilled in the art. In this regard reference can be made to e.g., Remington's Practice of Pharmacy, 11th ed. and in J. of. Pharmaceutical Science & Technology, Vol. 52, No. 5, Sept-Oct., p. 238-311 see table page 240 to 311, both publication include herein by reference.

The concentration of the compound having general chemical Formula II and the pharmaceutically acceptable carrier, for example, in an aqueous medium, varies with the particular field of use. A sufficient amount is present in the pharmaceutically acceptable carrier when satisfactory visualization of the imaging target (e.g., a tumor) is achievable.

In accordance with the invention, the radiolabeled compounds having general chemical Formula II either as a neutral composition or as a salt with a pharmaceutically acceptable counter-ion are administered in a single unit injectable dose. Any of the common carriers known to those with skill in the art, such as sterile saline solution or plasma, can be utilized after radiolabelling for preparing the injectable solution to diagnostically image various organs, tumors and the like in accordance with the invention. Generally, the unit dose to be administered for a diagnostic agent has a radioactivity of about 0.1 mCi to about 100 mCi, preferably 1 mCi to 20 mCi. For a radiotherapeutic agent, the radioactivity of the therapeutic unit dose is about 10 mCi to 700 mCi, preferably 50 mCi to 400 mCi. The solution to be injected at unit dosage is from about 0.01 ml to about 30 ml. For diagnostic purposes after intravenous administration, imaging of the organ or tumor *in vivo* can take place in a matter of a few minutes. However, imaging takes place, if desired, in hours or even longer, after injecting into patients. In most instances, a sufficient amount of the administered dose will accumulate in the area to be imaged within about 0.1 of an hour to permit the taking of scintigraphic images. Any conventional method of scintigraphic imaging for diagnostic purposes can be utilized in accordance with this invention.

30

In a **third aspect**, the present invention refers to a method of preparing a compound having general chemical Formula II (method for fluorination and preferably radiofluorination) using an appropriate fluorination agent. The method comprises the (single) step of coupling a compound having general chemical Formula I with a fluorination agent, more preferably the fluorination agent is a radioactive or non-radioactive ("cold") fluorine isotope derivate. In the latter case the reagent to convert the compound having general chemical Formula I to the

35

compound having general chemical Formula II is a fluorination agent. More preferably the compound having general chemical Formula II may thereafter be converted into a pharmaceutically acceptable salts of inorganic or organic acids thereof, hydrates, complexes, esters, amides, solvates and prodrugs thereof if desired. The reagents, solvents and
5 conditions which can be used for this fluorination are common and well-known to the skilled person in the field. See, *e.g.*, *J. Fluorine Chem.*, 27 (1985):117-191.

In a preferred embodiment of the method of preparing a compound having general chemical Formula A, wherein K = W comprises reacting a compound of general chemical
10 Formula A, wherein K = $-N^+(R^1)(R^2)(R^3) X^-$ with a fluorine isotope.

In a preferred embodiment of the method, the compound having general chemical Formula I and its pharmaceutically acceptable salts of inorganic or organic acids thereof, hydrates, complexes, esters, amides, solvates and prodrugs thereof is any preferred
15 compound described above for obtaining any preferred compound having general chemical Formula II, more specifically any preferred compound having general chemical Formulae IIA and IIB, or pharmaceutically acceptable salt, hydrate, ester, amide, solvate or prodrug thereof as described above.

20 In a preferred method of preparing a compound having general chemical Formula II, the step of fluorination more preferably radiofluorination of a compound having general chemical Formula I is carried out at a temperature at or below 90°C.

In a preferred method of preparing a compound of Formula II, the step of fluorination
25 more preferably radiofluorination of a compound of Formula I is carried out at a temperature selected from a range from 10°C to 90°C.

In a preferred embodiment, the method of fluorination more preferably radiofluorination occurs at a reaction temperature of from room temperature to 80°C.

30

In a preferred method of preparing a compound of Formula II, the step of fluorination more preferably radiofluorination of a compound of Formula I is carried out at a temperature selected from a range from 10°C to 70°C.

In a preferred method of preparing a compound of Formula II, the step of fluorination more preferably radiofluorination of a compound of Formula I is carried out at a temperature selected from a range from 30°C to 60°C.

- 5 In a preferred method of preparing a compound of Formula II, the step of fluorination more preferably radiofluorination of a compound of Formula I is carried out at a temperature selected from a range from 45 to 55°C.

- 10 In a preferred method of preparing a compound of Formula II, the step of fluorination more preferably radiofluorination of a compound of Formula I is carried out at a temperature at 50°C.

- More preferably, the radioactive fluorine isotope derivate is 4,7,13,16,21,24-Hexaoxa-1,10-diazabicyclo[8.8.8]-hexacosane K18F (crownether salt Kryptofix K18F), $K^{18}F$, $H^{18}F$,
15 $KH^{18}F_2$, $Cs^{18}F$, $Na^{18}F$ or tetraalkylammonium salt of ^{18}F (e.g. [F-18] tetrabutylammonium fluoride). Most preferably, the a radioactive fluorine isotope derivate is $K^{18}F$, $H^{18}F$, or $KH^{18}F_2$.


- In a preferred embodiment, the fluorination agent is a non-radioactive fluorine isotope.
20 More preferably, the non-radioactive fluorine isotope is ^{19}F derivative, most preferably ^{19}F .

In a preferred embodiment the solvents used in the present method may be DMF, DMSO, MeCN, DMA, DMAA, or mixture thereof, preferably the solvent is DMSO.

- 25 A new method is warranted in which the final product is prepared in a single step from the precursor. Only one purification step is necessary thereby the preparation can be accomplished in a short time (considering the half-life of ^{18}F). In a typical prosthetic group preparation, very often temperatures of 100°C and above are employed. The invention provides methods to accomplish the preparation at temperatures (80°C or below) that
30 preserve the biological properties of the final product. Additionally, single purification step is optionally carried out, thereby the preparation can be accomplished in a short time (considering the half-life of ^{18}F).

- In a tenth aspect the present invention refers to compounds having the general chemical
35 Formula V:

60



wherein $N^+(R^1)(R^2)(R^3)$, X^- , $-G$, and $-Q$, have the same meaning as depicted above for compounds having general chemical Formula I. This includes in particular all preferred embodiments mentioned above with regard to the residues and substituents R^1 , R^2 , R^3 , X^- , $-G$, and $-Q$, and all residues used to define these residues and substituents, such as R^4 , R^5 and the like;

R^6 is selected from the group comprising $-S(O)_2-N(H)-CH_2-C(O)OH$, $-S(O)_2-N(Me)-CH_2-C(O)OH$ and $C(O)OH$.

In a preferred embodiment R^6 is selected from the group comprising $-S(O)_2-N(Me)-CH_2-C(O)OH$ and $C(O)OH$.


In a more preferred embodiment R^6 is $C(O)OH$.

In a preferred embodiment of compounds of Formula V, $-G$ and $-Q$ are independently from each other selected from $-H$, $-CN$, CF_3 , and $-Cl$.


In a more preferred embodiment of compounds of Formula V, $-G$ and $-Q$ are independently from each other H , $-CF_3$, or CN .

In a even more preferred embodiment of compounds of Formula V, $-G$ and $-Q$ are independently from each other H , $-CF_3$, or $-CN$, whereas at least one member of the group comprising $-G$ or $-Q$ is $-CF_3$ or $-CN$.


Preferred compounds of Formula V are selected from the group comprising



Trifluoro-methanesulfonate(4-carboxy-2-cyano-phenyl)-trimethyl-ammonium;




Trifluoro-methanesulfonate(4-carboxy-2-trifluoromethyl-phenyl)-trimethyl-ammonium;



Trifluoro-methanesulfonate[4-(carboxymethyl-methyl-sulfamoyl)-2-cyano-phenyl]-trimethyl-ammonium;

Compounds of Formula V are suited to be coupled to targeting agents towards compounds of Formula I which are starting materials for the radio labeling reaction

5 towards compounds of Formula I or Formula A.



10 In a eleventh aspect the present invention refers to a method to synthesize compounds of Formula I (Formula A) from compounds of Formula V.

Compounds of Formula V can be condensed to targeting agents equipped with or without a spacer to obtain compounds of Formula I as defined above (Formula A) by using typical condensing agents which are known to persons skilled in the art. Suited condensing agents are for example DCC, DIC and 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-

15

methylpiperidinium tetrafluoroborate (J. Am. Chem. Soc. 2005, 127, 48, 16912-16920). Examples for such a reaction are depicted in scheme 3 and 4.

Example for Labeling:

5 First example:

¹⁸F-fluoride (up to 40 GBq) was azeotropically dried in the presence of Kryptofix 222 (5 mg in 1.5 ml MeCN) and cesium carbonate (2,3 mg in 0.5 ml water) by heating under a stream of nitrogen at 110-120°C for 20-30 minutes. During this time 3 × 1 ml MeCN were added and evaporated. After drying, a solution of the precursor (2 mg) in 150 µl DMSO
10 was added. The reaction vessel was sealed and heated at 50-70°C for 5-15 mins to effect labeling. The reaction was cooled to room temperature and dilute with water (2.7 ml). The crude reaction mixture was analyzed using an analytical HPLC. The product was obtained by preparative radio HPLC to give to desired ¹⁸F labeled peptide.

15 Second example:

Model compounds and peptides containing unnatural Histidine analogues

¹⁸F-fluoride (up to 100GBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and potassium carbonate (1mg in 0.5ml water) or cesium carbonate (2.3mg in 0.5ml water) by heating under nitrogen at 100-120°C for 20-30 minutes. During
20 this time 2-3 × 1ml MeCN were added and evaporated under vacuum or with a stream of nitrogen. After drying, a solution of the precursor (100-300µl of 0.0025 – 0.08M mg in DMSO) was added. The reaction vessel was sealed and heated at 50-90°C for 5-15mins to effect labeling. The crude reaction mixture was analyzed by HPLC. The product peak was confirmed by co-injection of the reaction mixture with the F19 cold standard. The
25 reaction mixture was diluted with water (4ml) and transferred to a semi-prep HPLC for purification. The product peak was collected diluted with water (10-20ml) and immobilized on a C18 separation cartridge. The pure F18-labeled product was eluted from the cartridge with EtOH (1-2ml).

30 Peptides containing natural Histidine

¹⁸F-fluoride (up to 100GBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and cesium carbonate (2.3mg in 0.5ml water) by heating under nitrogen at 100-120°C for 20-30 minutes. During this time 2-3 × 1ml MeCN were added and evaporated under vacuum or with a stream of nitrogen. After drying, a solution of the
35 precursor (100-300µl of 0.0025 – 0.08M mg in DMSO) was added. The reaction vessel was sealed and heated at 50-90°C for 5-15mins to effect labeling. The crude reaction

mixture was analyzed by HPLC. The product peak was confirmed by co-injection of the reaction mixture with the F19 cold standard. The reaction mixture was diluted with water (4ml) and transferred to a semi-prep HPLC for purification. The product peak was collected diluted with water (10-20ml) and immobilized on a C18 separation cartridge. The
5 pure F18-labeled product was eluted from the cartridge with EtOH (1-2ml).

Without further elaboration, it is believed that one skilled in the art can, using the preceding description, utilize the present invention to its fullest extent. The following
10 preferred specific embodiments are, therefore, to be construed as merely illustrative, and not limitative of the remainder of the disclosure in any way whatsoever.

The entire disclosure[s] of all applications, patents and publications, cited herein are incorporated by reference herein.
15

The following examples can be repeated with similar success by substituting the generically or specifically described reactants and/or operating conditions of this invention for those used in the preceding examples.

20 From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention and, without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions.

25 **General Method for the Preparation of Compounds**


The peptide portion of the molecule -L-Y-U can be conveniently prepared according generally established techniques known in the art of peptide synthesis, such as solid-phase peptide synthesis. They are amenable Fmoc-solid phase peptide synthesis,
30 employing alternate protection and deprotection. These methods are well documented in peptide literature. (Reference: "*Fmoc Solid Phase Peptide Synthesis*" A practical approach", Edited by W.C.Chan and P.D.White, Oxford University Press 2000) (For Abbreviations see Descriptions).

35 In the following two schematic examples are given of how to prepare a compound having general chemical Formula II using a compound having general chemical Formula I. The

methods presented as schemes below are in principle suitable to generate compounds over the whole breadth of Formula II using compounds over the whole breadth of Formula I. The examples presented below are given merely to illustrate a way of labeling a compound having general chemical Formula I to arrive at a compound having general chemical Formula II and is not to be understood as to limit the invention to the methods exemplified herein.

Scheme 3 depicts an example of a synthetic route for forming a trimethylammonium substituted aromatic moiety containing peptide having general chemical Formula I and subsequent direct radiolabeling towards the corresponding ^{18}F -labeled compound having general chemical formula II.

The synthesis starts with commercially available 2-Fluoro-5-formyl-benzonitrile (**3**) which is converted to the corresponding acid (**4**) by known oxidation methods, e.g., use of chromium reagents, manganese reagents or other typical reagents which are well-known to experts and which can be taken from, but are not limited to, the methods described and cited in the book: "*Modern oxidation methods*" by Jan-Erling Bäckvall, Wiley-VCH, 2004. A useful method is, e.g., the oxidation with sodium chlorite in phosphate-buffered tert-butanol solution.




Scheme 3

Acid (4) can be converted to the corresponding methyl ester with methanol and acetyl chloride (*Helv. Chim. Acta*, 2005, 88, 7:1630-1657). But the preparation of other alkyl esters (e.g., ethyl ester) and other esterification methods including variants under basic

conditions are also possible and useful and are well known to experts. The displacement of fluoride (5) with dimethylamine or dimethylamine hydrochloride by nucleophilic aromatic substitution reaction can be carried out, e.g., in a suspension of DMSO and potassium carbonate (e.g., *Bioorg. Med. Chem. Lett.*, 10, 23, 2000:2603-2606). Useful other solvents can be selected from, but are not limited to, acetonitrile and DMF. The quarternisation of aniline (6) with methyl iodide, methyl triflate or other alkylating agents is carried out in boiling dichloromethane with normal or increased pressure (e.g., 1-10 bar). Other useful solvents for this reaction can be selected from, but are not limited to, acetone and dichloroethane. The crude product can be purified by reverse-phase column chromatography. The methyl ester (7) can be cleaved in boiling trifluoro acetic acid and water (*Bioorg. Med. Chem.*, 2003, 11:4189-4206) or under other acidic conditions. The acid (8) is coupled by solid phase synthesis (as shown in scheme 3) to solid-phase bound peptide to obtain amide (9) (or esters) by methods which are well known to experts. Typical condensating agents for those kinds of couplings are diisopropylcarbodiimid or dicyclohexylcarbodiimid, but also other condensating agents (see, e.g., Chan and White (*Fmoc Solid Phase Peptide Synthesis - A Practical Approach*)), are possible. The resin-bound peptide is then cleaved from the resin by acidification to obtain liberated peptide (10). The cleavage is also possible by other appropriate methods which are very much dependent on the kind of linker. The methods of peptide cleavage from resin are very well known to experts and described in literature (e.g., Chan and White - *Fmoc Solid Phase Peptide Synthesis - A Practical Approach*). The purified peptide (10) is converted at 70°C $\pm$ 45°C with [¹⁸F] potassium fluoride, potassium carbonate and Kryptofix (4,7,13,16,21,24-Hexaoxa-1,10-diaza-bicyclo[8.8.8]hexacosane) in dimethylsulfoxide to obtain the desired ¹⁸F-labeled peptide (11). The reagents and solvents which are used for this radiofluorination are common, well-known to experts since many years and described in many publications (e.g., *J. Fluorine Chem.*, 27 (1985):117-191). It was surprisingly found that the radiofluorination does take place and that the temperature can be decreased to <90°C so that the peptide is not harmed or decomposed. The preferred temperature range for conducting the radioflurination of compounds according to Formula I is from 10°C to 90°C. More preferred is the temperature range of 10°C to 70°C. Even more preferred is the range of temperature from 35°C to 65°C. Further preferred is a range of temperature from 45°C to 55°C. Most preferred is a temperature of 50°C for carrying out the radiofluorination of compounds having general chemical Formula I to arrive at compounds having general chemical Formula II.

Another example of a trimethyl ammonium derivative was prepared as shown in Scheme 4: Commercially available sulfonylchloride **12** was coupled with the secondary amine sarcosin methyl ester to the sulfonamide **13** in dichlormethane and diisopropyl ethyl amine to scavenge hydrogen chloride. Other useful solvents for this reaction can be, but are not limited to, DMF, THF, dioxane, dichloroethane, diethylether and tert butyl ether. Other useful bases can be, but are not limited to, trimethylamine, N-methyl morpholine, NMP and sodium hydrogen carbonate.




Scheme 4

- 5 Sulfonamide **13** was then treated in an aromatic nucleophilic substitution reaction with dimethylammonium hydrochloride and potassium carbonate in dimethylsulfoxide to obtain dimethylaniline **14**. The aniline derivative **14** was quaternized with methyltriflate or methyl iodide in dichloromethane or dichloroethane towards ammonium salt **15**. The resulting ester **15** was cleaved under acidic conditions, e.g., in boiling trifluoro acetic acid and water

(*Bioorg. Med. Chem.*, 2003, 11:4189-4206). The acid **16** was condensed to the resin-bound peptide by methods which are well known to experts. Typical condensating agents for those kinds of couplings are diisopropylcarbodiimid or dicyclohexylcarbodiimid, but also other condensating agents are possible and described. Examples are given in, but are not limited to, the methods described and cited in the book: Chan and White - "*Fmoc Solid Phase Peptide Synthesis - A Practical Approach*". The resin-bound trimethylammonium peptide **17** is then cleaved from the resin by acidification to obtain liberated peptide (**18**). The cleavage is also possible by other appropriate methods which are very much dependent on the kind of linker. The methods of peptide cleavage from resin are very well known to experts. Examples are given in, but are not limited to, the methods described and cited in the book: Chan and White - "*Fmoc Solid Phase Peptide Synthesis - A Practical Approach*". The purified peptide (**18**) is converted at 70°C ± 45°C with [¹⁸F] potassium fluoride, potassium carbonate and Kryptofix (4,7,13,16,21,24-Hexaoxa-1,10-diaza-bicyclo[8.8.8]hexacosane) in dimethylsulfoxide to obtain the desired ¹⁸F-labeled peptide (**19**). The reagents and solvents which are used for this radiofluorination are common, well-known to experts and are described in many publications (e.g., *J. Fluorine Chem.*, 27 (1985):117-191). Beside potassium carbonate as base also tetraalkyl ammonium carbonate is possible. It was surprisingly found that the radiofluorination does take place at the given mild temperatures so that the peptide is not harmed or decomposed. The preferred temperature range for conducting the radiofluorination of compounds having general chemical Formula I is from 10°C to 90°C. More preferred is the temperature range of 10°C to 70°C. Even more preferred is the range of temperature from 35°C to 65°C. Further preferred is a range of temperature from 45°C to 55°C. Most preferred is a temperature of 50°C for carrying out the radiofluorination of compounds according to Formula I to arrive at compounds according to Formula II.


More complex peptides can also be labeled directly in a one-step F-18 labeling procedure: The following fluorination-reaction (20 → 21; Scheme 5) is carried out in DMSO at 70°C at 15 min reaction time.



- 5 It is surprising that molecules with such a complex pattern of unprotected functionalities can be F-18 labeled easily under these mild conditions with high incorporation of F-18 fluoride. The chromatogram of Fig. 6 shows the desired product as peak in the radioactive version, whereas the non-radioactive standard co-elutes in the UV-chromatogram (HPLC chromatogram of reaction mixture with co-injection of the cold standard).


10

Another example for the F-18 labelling of complex peptides is shown in the following Scheme 6:




Again, the following HPLC-chromatogram represents the radioactive and the UV-version of the corresponding F-18 and F-19 fluoro-peptides (23 and 22), respectively. The Fluorination-reaction is also carried out in DMSO at 70°C at 15 min reaction time. The respective chromatogram is shown in Fig. 7 (HPLC chromatogram of reaction mixture with co-injection of the cold standard).

The F-18 labeling of small molecules is also possible. For example compound 24 (WO 2007/16538 A2) can be condensed with trimethylammonium benzoic acid derivative 25 to obtain the precursor 26 for F-18 labeling (Scheme 7). The incorporation of F-18 fluoride is achieved to obtain F-18 labeled compound 27 (the corresponding non-radioactive reference standard is published (WO 2007/16538 A2)).



Scheme 7

An example of the F-18 labeling of an oligonucleotide is shown in scheme 8. TTA1 (Nucleic Acids Research, 2004, Vol. 32, No. 19, 5757–5765) is equipped with a trimethylammonium derivative of Formula III by use of a triazine condensating agent (J. Am. Chem. Soc. 2005, 127, 48, 16912-16920). The subsequent F-18 radiolabeling is obtained in reasonable yield, although the specific activity was relatively low due to the fact that the purification of the F-18 labeled compound is achieved under non-optimal circumstances..


**Examples:****5 Example A:**

Synthesis of $X^- N^+(R^1)(R^2)(R^3)-(C_6H_4(-G;-Q))--L--Y--U$ (Ia)


Synthesis of H-Y--U: Solid-phase peptide synthesis (SPPS) involves the stepwise addition of amino acid residues to a growing peptide chain that is linked to an insoluble support or matrix, such as polystyrene. The C-terminal residue of the peptide is first anchored to a commercially available support (e.g., Rink amide resin) with its amino group protected with an N-protecting agent, fluorenylmethoxycarbonyl (Fmoc) group. The amino protecting group is removed with suitable deprotecting agent such as piperidine for Fmoc and the next amino acid residue (in N-protected form) is added with a coupling agent such as dicyclohexylcarbodiimide (DCC), di-isopropyl-cyclohexylcarbodiimide (DCCI), hydroxybenzotriazole (HOBt). Upon formation of a peptide bond, the reagents are washed from the support. After addition of the final residue of (Y), the peptide is attached to the solid support is ready for the coupling of $X^- N^+(R^1)(R^2)(R^3)-(C_6H_4(-G))-L-OH$.

Synthesis of $X^- N^+(R^1)(R^2)(R^3)-(C_6H_4(-G))-L-Y-U$ (Ia)


Ia: L = CO, $X^- = CF_3SO_3^-$



-G = -CN, -CF₃, -F and others specified



5 (G = CN), **8** (G = CF₃), **11** (G = F)



To a suspension of the resin -Y--U--RESIN in DMF (0.1 to 0.25 mmol), $X^- N^+(R^1)(R^2)(R^3)-(C_6H_4(-G))-L-OH$ (**5**, **8** or **11**, 2 – 4 equivalents) was added along with 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium tetrafluoroborate.

Example B describes the preparation of **5**, **8** and **11**.


After 3-8 hours, the resin was washed with DMF and dichloromethane. The peptide Ia) was isolated from the resin using TFA:diisopropylsilane:phenol:water cocktail with concomitant removal of the protecting groups of amino acids. The product was purified by HPLC using appropriate TFA:H₂O: 0.1 TFA gradient using C₁₈-reverse phase column. The products were identified by mass spectra.

Ia-1: 4-(Trimethylammonium)-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂; Molecular wt: Calculated: 1422.73, Found: 711.86 ((M⁺+1)/2)

Table 1 lists all the trimethylammonium compounds precursors Ia-1 – Ia-25 used for F-18 labeling

Example B

Preparation of X⁻ N⁺(R¹)(R²)(R³)-(C₆H₄(-G))--L-OH 5 (-G = -CN)



a) Synthesis of 3-Cyano-4-fluoro-benzoic acid:

To a stirred solution of 15.0 g (97,6 mmol) 2-fluoro-5-formyl-benzonitrile (Aldrich), 150 ml dest. water and 630 ml t-butanol were added 40.8 g (361 mmol) sodium chlorite and 35.9 g (230 mmol) sodium hydrogen phosphate dihydrate. The reaction mixture was stirred over night and poured into a diluted aqueous hydrogen chloride solution (pH=3.5). The pH value was readjusted to pH=3.5 by aqueous hydrogen chloride. The aqueous solution was extracted trice with dichloromethane/isopropanol (10:1). The combined organic phases were dried (sodium sulfate) and concentrated. The residue was purified by extraction with sodium hydrogen carbonate solution and dichloromethane, acidification with aqueous

solution and subsequent filtering. The solid crude product 1 was obtained in 90% yield (14.5 g, 87.8 mmol) and was used for the next step without purification.

MS-ESI: 166 ($M^+ + 1$, 77),

Elementary analysis:

5 Calculated: C 58.19% H 2.44% F 11.51% N 8.48%

Found: C 58.81% H 2.42% F 11.41% N 8.47%

b) Synthesis of 3-Cyano-4-fluoro-benzoic acid methyl ester 2:

10 To a stirred suspension of 16.0 g (96.9 mmol) 1 and 161 ml methanol were added 30.4 g (387.6 mmol) acetyl chloride drop wisely at 0°C. The reaction mixture was stirred over night, filtered and concentrated. The residue was diluted with dichloromethane, washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The residue was purified by column chromatography (hexane :

15 ethylacetate). The desired product 2 was obtained in 78.1% yield (13.5 g; 75.7 mmol)

MS-ESI: 180 ($M^+ + 1$, 57),

Elementary analysis:

Calculated: C 60.34% H 3.38% F 10.60% N 7.82%

Determined: C 60.51% H 3.39% F 10.57% N 7.80%

20

c) Synthesis of 3-Cyano-4-dimethylamino-benzoic acid methyl ester 3:

To a stirred solution of 24.0 g (134 mmol) 2 and 240 ml dimethylsulphoxid were added 13.2 g (161 mmol) dimethylamine hydrochloride and 38.9 g (281 mmol) potassium carbonate. The reaction mixture was stirred over night and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude product 3 was

25

30 obtained in 94% yield (25.7 g, 126 mmol) and was used for the next step without purification.

MS-ESI: 205 ($M^+ + 1$, 59),

Elementary analysis:

Calculated: C 64.69% H 5.92% N 13.72%

35 Found: C 64.79% H 5.95% N 13.69%

d) Synthesis of (2-Cyano-4-methoxycarbonyl-phenyl)-trimethyl-ammonium trifluoro-methanesulfonate 4:

To a stirred solution of 6.16 g (30.2 mmol) 3 and 110 ml dichloromethane were added
5 50.0 g (302 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was stirred
over night and diethylether was added. After evaporation of one third of the solvent
volume the desired compound precipitates and the rest of the solvent were decanted. The
solid was washed extensively (ten times) with large amounts of diethylether. The solid
was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography
10 (acetonitril/water - gradient 1:99 to 80:20). The desired compound 4 was obtained in 69 %
yield (20.8 mmol, 7.68 g).

MS-ESI: 219 (M^+ , 100),

Elementary analysis:

Calculated: C 42.39% H 4.10% F 15.47% N 7.61%

15 Found: C 42.42% H 4.12% F 15.41% N 7.59%

e) Synthesis of Trifluoro-methanesulfonate(4-carboxy-2-cyano-phenyl)-trimethyl-ammonium 5:

20 A solution of 4.01 g (10.9 mmol) 4, 95 ml dest. water and 95 ml trifluoroacetic acid was
refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump
vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed
extensively with diethyl ether and dried by oil pump vacuum. The solid crude product 5
was obtained in 93% yield (3.59 g, 10.1 mmol) and crude compound 5 was used for the
25 next step without purification.

MS-ESI: 205 (M^+ , 100),


Elementary analysis:

Calculated: C 40.68% H 3.70% F 16.09% N 7.91%

Found: C 40.72% H 3.71% F 16.06% N 7.91%

30

Preparation of RG-L₁-B₁-OH 8 (G = CF₃)



a) 4-Dimethylamino-2-trifluoromethyl-benzoic acid methyl ester 6:

- 5 To a stirred solution of 4,48 g (22.5 mmol) 4-Fluoro-2-trifluoromethyl-benzoic acid methyl ester (Rarechem) and 60.0 ml dimethylsulfoxide were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 8 h at 60°C in an autoclave and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice
- 10 with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product 6 was obtained in 72 % yield (4,00 g, 16,2 mmol).
- 15 MS-ESI: 248 ($M^+ + 1$, 78).
 Elementary analysis: C 53.44% H 4.89% F 23.05% N 5.67%
 Found: C 53.46% H 4.91% F 23,04% N 5.64%

b) Trifluoro-methanesulfonate(4-methoxycarbonyl-3-trifluoromethyl-phenyl)-trimethyl-ammonium 7:

- 20 To a stirred solution of 3,09 g (12,5 mmol) 6 and 50 ml dichloromethane were added 20,5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2 days then cooled to room temperature. Diethylether was added. The desired compound
- 25 precipitated and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and

purified by (C-18) RP-column chromatography (acetonitrile/water - gradient 1:99 to 80:20).

The desired compound 7 was obtained in 69% yield (3,55 g, 8,63 mmol).

MS-ESI: 262 (M^+ , 87),

Elementary analysis:

5 Calculated: C 37.96% H 3.68% F 27.71% N 3.41%

Determined: C 38.01% H 3.63% F 27.69% N 3.41%

c) Trifluoro-methanesulfonate(4-carboxy-3-trifluoromethyl-phenyl)-trimethylammonium 8:

- 10 A solution of 2,84 g (6,92 mmol) 7, 60 ml dist. water and 60 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude was obtained in 89% yield (2,45 g; 6.16 mmol) and crude compound 8 was used for the next step
- 15 without purification.

MS-ESI: 248 (M^+ , 59),


Elementary analysis:

Calculated: C 36.28% H 3.30% F 28.69% N 3.53%

Determined: C 36.30% H 3,32% F 28.67% N 3,52%

20

Preparation of $X^- N^+(R^1)(R^2)(R^3)--(C_6H_4(-G))--L-OH$ 10 (-G = -F)



- 25 a) 3-Fluoro-4-dimethylamino-benzoic acid methyl ester 9: To a stirred solution of 38,7g (225 mmol) 3,4-difluoro-benzoic acid methyl ester (Apollo) and 600 ml dimethylsulphoxide

were added 22.3 g (270 mmol) dimethylamine hydrochloride and 65.4 g (473 mmol) potassium carbonate. The reaction mixture was stirred for 5 h at 55°C in an autoclave and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude product 9 was obtained in 71% yield (31.5 g, 160.0 mmol) and was used for the next step without purification.

MS-ESI: 198 ($M^+ + 1$, 72).

10 Elementary analysis:

Calculated: C 60.91% H 6.13% F 9.63% N 7.10%

Found: C 60.99% H 6.15% F 9.60% N 7.07%

15 b) Synthesis of (2-fluoro-4-methoxycarbonyl-phenyl)-trimethyl-ammonium trifluoro-methanesulfonate 10:

To a stirred solution of 3,90 g (19,8 mmol) 9 and 70 ml dichloromethane were added 32.5 g (198 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was stirred for 2.5 days at room temperature and diethylether was added. The desired compound precipitates and the solvent were decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound 10 was obtained in 80% yield (5,72 g, 15.84 mmol).

MS-ESI: 212 (M^+ , 76),

25 Elementary analysis:

Calculated: C 39.89% H 4.18% F 21.03% N 3.88%

Found: C 39.93% H 4.20% F 21.01% N 3.84%

30 c) Synthesis of (4-carboxy-2-fluoro-phenyl)-trimethyl-ammonium trifluoro-methanesulfonate 11:

A solution of 4,00 g (11.1 mmol) 10, 96 ml dest. water and 96 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude product 11

was obtained in 92% yield (3.54 g, 10.2 mmol) and crude compound 11 was used for the next step without purification.

MS-ESI: 198 (M^+ , 76),

Elementary analysis:

5 Calculated: C 38.04% H 3.77% F 21.88% N 4.03%




Found: C 38.10% H 3.79% F 21.81% N 4.00%

Example C:


10 Synthesis of $F-(C_6H_4(-G))--L--Y--U$ (IIB-a)

For the identification $^{18}F-(C_6H_4(-G))--L--Y--U$, reference standards

$^{19}F-(C_6H_4(-G))--L--Y--U$ were prepared according to the scheme shown below.



1 ($G = CN$), **12** ($G = CF_3$), **13** ($G = F$)



15

Compounds 1, 12 and 13 were purchased commercially.

20 Synthesis of $H-Y--U--RESIN$ and coupling of $^{19}F-(C_6H_4(-G))--L-OH$ were accomplished according to the methods described in Example A.

IIB-a-1: 4-[19]-Fluoro-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂, Molecular Weight, Calculated: 1382.62, Found: 691.9 ($(M^++1)/2$)

Table 2 lists all the F-19 compounds (IIB-a-1 to IIB-a-25) prepared for the chromatographic identification for products resulting from Ia-1 – Ia-25 used for F-18 labeling as well as binding affinities. The Table lists all other F-19 compounds used in the SA analysis for the selection of high affinity analogs. Measurement of binding constants is described in Example D.

Example D:

In vitro binding affinity and specificity of Bombesin analogs for the human bombesin 2 receptor (GRPR) were assessed via a competitive receptor-binding assay using ^{125}I -[Tyr⁴]-Bombesin (Perkin Elmer; specific activity 81.4 TBq/mmol) as GRPR-specific radioligand. The assay was performed based on the scintillation proximity assay (SPA) technology (J.W.Carpenter *et al.*, *Meth. Mol. Biol.*, 2002; 190:31-49) using GRPR-containing cell membranes (Perkin Elmer) and wheat germ agglutinin (WGA)-coated PVT beads (Amersham Bioscience).

Briefly, GRPR-containing membranes and WGA-PVT beads were mixed in assay buffer (50 mM Tris/HCl pH 7.2, 5 mM MgCl₂, 1 mM EGTA, Complete protease inhibitor (Roche Diagnostics GmbH) and 0.3% PEI) to give final concentrations of approximately 100 µg/ml protein and 40 mg/ml PVT-SPA beads. The ligand ^{125}I -[Tyr⁴]-Bombesin was diluted to 0.5 nM in assay buffer. The test compounds were dissolved in DMSO to give 1 mM stock solutions later on, they were diluted in assay buffer to 8 pM – 1.5 µM.

The assay was then performed as follows: First, 10 µl of compound solution to be tested for binding were placed in white 384 well plates (Optiplate-384, Perkin-Elmer). At next, 20 µl GRPR/WGA-PVT bead mixture and 20 µl of the ligand solution were added. After 90 minutes incubation at room temperature, another 50 µl of assay buffer were added, the plate sealed and centrifuged for 10 min at 520xg at room temperature. Signals were measured in a TopCount (Perkin Elmer) for 1 min integration time per well. The IC₅₀ was calculated by nonlinear regression using the GraFit data analysis software (Erithacus Software Ltd.). Furthermore, the K_i was calculated based on the IC₅₀ for test compound as well as the K_D and the concentration of the ligand ^{125}I -[Tyr⁴]-Bombesin. Experiments were done with quadruple samples.


The Binding affinities measured for all the cold F-19 compounds are listed Table 2.

Example E:General Radiolabeling Method

- 5 In a 5 mL Wheaton vial, ^{18}F -fluoride (up to 40GBq) was azeotropically dried in the presence of Kryptofix 222 (5 mg in 1.5 ml CH_3CN) and potassium carbonate (1 mg in 0.5 ml water) or cesium carbonate (2,3 mg in 0.5 ml water) by heating under a stream of nitrogen at 110-120°C for 20-30 minutes. During this time 3×1 ml CH_3CN were added and evaporated. After drying, a solution of Ia-1 (2 mg) in 150 μl DMSO was added. The
- 10 reaction vessel was sealed and heated at 50-70°C for 5-15 mins to effect labeling. The reaction was cooled to room temperature and diluted with water (2.7 ml). The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6 mm, 1.8 μ , 2 ml/min, solvent A: H_2O , solvent B: MeCN, gradient: 5%-95% B in 7 min or Column Econosphere C18, 53 $\times$ 7 mm, 3 μ , 3 ml/min (Alltech), solvent A: H_2O + 0.1 %
- 15 TFA, solvent B: MeCN/Water 9/1 + 0.1 % TFA, gradient: 5-95 % B in 7 min. The F-18 labeled product IIA-a-1 was confirmed by co-injection with the cold F-19 fluoro standard (IIB-a-1) on the Econsphere analytical HPLC.

The product IIA-a-1 was obtained by preparative radio HPLC to give to desired F-18 labeled peptide.

20



HPLC of Ia-1, IIA-a-1 and IIB-a-1 are shown in Figs. 1 and 2.

Conditions for HPLC according to Fig. 1:

5

Column: C-18, Gradient: A: 10 % CH₃CN: 90 % H₂O, B: 90 % CH₃CN: 10 % H₂O containing 0.1 % TFA (UPLC); Flow rate: 2 mL, 1 % A to 99 % B in 2 min.

Conditions for HPLC according to Fig. 2:

10

Column: C-18, Gradient: A: 10mM K₂HPO₄ pH: 9,04 ; B: 10mM K₂HPO₄ pH: 9,04/ACN 3/ 7; Flow rate: 2 mL, 5 % A to 95 % B in 7 min. Retention time IIA-a-1 – 5.01 min. IIB-a-1 – 4.89 min.

15 In a similar manner, Compounds shown in Table 1 (Ia-2 to Ia-22) were labeled with F-18 to yield F-18 labeled peptides, IIA-a-2 to IIA-a-22 (Table 3) respectively. The chromatographic behavior of IIA-a-2 to IIA-a-22 were compared with IIB-a-2 to IIB-a-22 respectively for complete characterization in rodent studies and imaging.

20 Human serum Stability of IIA-a-1:

To qualify for pre-clinical and clinical use, it is necessary to establish the stability of the compounds in human serum. The compound a 70 µL of human serum containing the F-18 labeled peptide (5.89 MBq/mL) was incubated at 37°C for 90 min. An aliquot was
25 withdrawn at various intervals and the purity assessed by HPLC. The purity was evaluated in ... (?) under two different conditions. Potassium hydrogen phosphate buffer system was used as a mobile phase to measure the stability of attached F-18 label.

HPLC chromatograms are shown in Fig. 3. Conditions for HPLC: Column: C-18, Gradient:
30 A: 10mM K₂HPO₄ pH: 9,04 ; B: 10mM K₂HPO₄ pH: 9,04/ACN 3/ 7; Flow rate: 2 mL, 5% A to 95 % B in 7 min.

Trifluoroacetic acid system was used as a mobile phase to measure the stability of the entire molecule.

35

Column: C-18, Gradient: A: 10% CH₃CN: 90 % H₂O, B: 90 % CH₃CN: 10 % H₂O containing 0.1 % TFA (chromatograms in Fig. 4).

The above experiments clearly show that in the compounds of the present invention, the F-18 isotope and the $[^{18}\text{F}]\text{-C}_6\text{H}_4(-\text{G},-\text{Q})\text{--L--Y--(IIA)}$, targeting molecule of the invention are stable in human serum.

- 5 In a similar manner, the compounds shown in Table 1 (Ia-1 to Ia-22) were labelled with F-18 to yield F-18 labelled peptides, IIA-a-1 to IIA-a-22 (Table 3) respectively. The chromatographic behaviour of IIA-a-1 to IIA-a-22 were compared with IIB-a-1 to IIB-a-22 respectively for complete characterization in rodent studies and imaging.

10 Example 1

a) 2-Chloro-4-dimethylamino-benzoic acid methyl ester (1a)

- To a stirred solution of 4,00g (20.6 mmol) 2-chloro-4-fluoro-benzoic acid methyl ester
15 (Apollo) and 60 ml dimethylsulphoxid were added 2.03 g (24,7 mmol) dimethylamine hydrochloride and 5,97 g (43.2 mmol) potassium carbonate. The reaction mixture was stirred over night and is reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases
20 were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude product 1a was obtained in 99 % yield (4,36 g, 20.4 mmol) and was used for the next step without purification.

MS-ESI: 213 / 215 ($\text{M}^+ + 1$, 64 / 48).

25 Elementary analysis:

Calculated: C 56.21% H 5.66% N 6.56%

Determined: C 56.39% H 5,67% N 6.54%

30 b) Synthesis of (3-chloro-4-methoxycarbonyl-phenyl)-trimethyl-ammonium trifluoromethanesulfonate (1b)

- To a stirred solution of 4.49 g (21.0 mmol) 1a and 75 ml dichloromethane were added
34,5 g (210 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was stirred for 2 days at room temperature. 17 g (10 mmol) methyltriflate (Aldrich) were added and the
35 reaction mixture was stirred at 40°C for 20 h. The reaction mixture was cooled to 20°C and diethylether was added. The desired compound precipitates and the solvent was

decanted and the solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **1b** was obtained in 86% yield (6,86 g, 18,1 mmol).

5

MS-ESI: 228/230 (M^+ , 81),

Elementary analysis:

Calculated: C 38.15% H 4.00% F 15.09% N 3.71%

Determined: C 38.18% H 4.02% F 15.04% N 3.70%

10

c) Synthesis of (4-carboxy-3-chloro-phenyl)-trimethyl-ammonium trifluoromethanesulfonate (**1c**).

15 A solution of 0.5 g (1.32 mmol) **1b**, 12 ml dest. water and 12 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude **1c** was obtained in 98% yield (471 mg, 1.3 mmol) and crude compound **1c** was used for the next
20 step without purification.

MS-ESI: 214 / 216 (M^+ , 89 / 56),

Elementary analysis:

Calculated: C 36.32% H 3.60% F 15.67% N 3.85%

25 Determined: C 36.37% H 3.63% F 15.61% N 3,83%

d) Synthesis of (4-Trimethylammonium-2-chloro-benzoyl)-Trp-Ala-Val-Leu-NH₂- triflate salt (**1d**).

30 To a stirred solution of 72.8 mg (0.2 mmol) **1c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Trp(Boc)-Ala-Val-Leu-NH₂ (loading 0.68 mmol/g) which was prepared by standard
35 protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 69,4 mg (0.2 mmol) **1c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was

added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophilized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**1d**) in 30.0 % yield - 16.2 mg (0.0195 mmol). MS-ESI: 683/685 (M^+ , 39 / 26)

e) Synthesis of [^{18}F]- (4-Fluoro-2-chloro-benzoyl)-Trp-Ala-Val-Leu-NH₂ (**1e**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (316MBq, 33 μ l) were added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **1d** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 80°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **1e** was confirmed by co-injection with the cold F-19 fluoro standard on the Econsphere analytical HPLC.

Example 2

a) Synthesis of 3-Cyano-4-fluoro-benzoic acid (**2a**).

To a stirred solution of 15.0 g (97,6 mmol) 2-fluoro-5-formyl-benzonitrile (Aldrich), 150 ml dest. water and 630 ml t-butanol were added 40.8 g (361 mmol) sodium chlorite and 35.9 g (230 mmol) sodium hydrogen phosphate dihydrate. The reaction mixture was stirred over night and poured into a diluted aqueous hydrogen chloride solution (pH=3.5). The pH value was readjusted to pH=3.5 by aqueous hydrogen chloride. The aqueous solution was extracted trice with dichloromethane/isopropanol (10:1). The combined organic phases

were dried (sodium sulphate) and concentrated. The residue was purified by extraction with sodium hydrogen carbonate solution and dichloromethane, acidification with aqueous solution and subsequent filtering. The solid crude product **2a** was obtained in 90% yield (14.5 g, 87.8 mmol) and was used for the next step without purification.

5

MS-ESI: 166 ($M^+ + 1$, 77),

Elementary analysis:

Calculated: C 58.19% H 2.44% F 11.51% N 8.48%

Determined: C 58.81% H 2.42% F 11.41% N 8.47%

10

b) Synthesis of 3-Cyano-4-fluoro-benzoic acid methyl ester (**2b**).

15

To a stirred suspension of 16.0 g (96.9 mmol) **2a** and 161 ml methanol were added 30.4 g (387.6 mmol) acetyl chloride drop wisely at 0°C. The reaction mixture was stirred over night, filtered and concentrated. The residue was diluted with dichloromethane, washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The residue was purified by column chromatography (hexane : ethylacetate). The desired product **2b** was obtained in 78,1% yield (13.5 g; 75.7 mmol)

20

MS-ESI: 180 ($M^+ + 1$, 57),

Elementary analysis:

Calculated: C 60.34% H 3.38% F 10.60% N 7.82%

Determined: C 60.51% H 3.39% F 10.57% N 7.80%

25

c) Synthesis of 3-Cyano-4-dimethylamino-benzoic acid methyl ester (**2c**).

30

To a stirred solution of 24.0 g (134 mmol) **2b** and 240 ml dimethylsulphoxid were added 13.2 g (161 mmol) dimethylamine hydrochloride and 38,9 g (281 mmol) potassium carbonate. The reaction mixture was stirred over night and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude product **2c** was obtained in 94% yield (25,7 g, 126 mmol) and was used for the next step without purification.

35

MS-ESI: 205 ($M^+ + 1$, 59),

Elementary analysis:

5 Calculated: C 64.69% H 5.92% N 13.72%

 Determined: C 64.79% H 5.95% N 13.69%

10 d) Synthesis of (2-Cyano-4-methoxycarbonyl-phenyl)-trimethyl-ammonium trifluoro-methanesulfonate (2d).

To a stirred solution of 6.16 g (30.2 mmol) **2c** and 110 ml dichloromethane were added 50.0 g (302 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was stirred over night and diethylether was added. After evaporation of one third of the solvent
15 volume the desired compound precipitates and the rest of the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **2d** was obtained in 69 % yield (20.8 mmol, 7.68 g).

20

MS-ESI: 219 (M^+ , 100),

Elementary analysis:

Calculated: C 42.39% H 4.10% F 15.47% N 7.61%

Determined: C 42.42% H 4.12% F 15.41% N 7.59%

25

e) Synthesis of Trifluoro-methanesulfonate(4-carboxy-2-cyano-phenyl)-trimethyl-ammonium; (**2e**).

A solution of 4.01 g (10.9 mmol) **2d**, 95 ml dest. water and 95 ml trifluoroacetic acid was
30 refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude product **2e** was obtained in 93% yield (3.59 g, 10.1 mmol) and crude compound **2e** was used for the next step without purification.

35

MS-ESI: 205 (M^+ , 100),

Elementary analysis:

Calculated: C 40.68% H 3.70% F 16.09% N 7.91%

Determined: C 40.72% H 3.71% F 16.06% N 7.91%

5 f) Synthesis of peptide

(4-Trimethylammonium-3-cyano-benzoyl)-Arg-Ala-His(π -Me)-Leu-NH₂ – triflate salt, (2f).

To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Arg(Pbf)-Ala-His(π -Me)-Leu-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was lyophilized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**2f**) in 27,0 % yield - 14,9 mg (0.0176 mmol). MS-ESI: 698 (M⁺, 100).

g) Synthesis of

[¹⁸F]-(4-Fluoro-3-cyano-benzoyl)- Arg-Ala-His(π -Me)-Leu-NH₂ (2g).

30

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (356 MBq, 38 μ l) were added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **2f** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 50°C for 10 min. The crude reaction mixture was analyzed using an analytical

35

HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8μ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3μ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product (**2g**) was confirmed by co-injection with the cold F-19 fluoro standard on the Econsphere analytical HPLC.

Example 3

a) 3-Fluoro-4-dimethylamino-benzoic acid methyl ester (**3a**)

To a stirred solution of 38,7g (225 mmol) 3,4-difluoro-benzoic acid methyl ester (Apollo) and 600 ml dimethylsulphoxid were added 22.3 g (270 mmol) dimethylamine hydrochloride and 65.4 g (473 mmol) potassium carbonate. The reaction mixture was stirred for 5h at 55°C in an autoclave and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude product **3a** was obtained in 71% yield (31.5 g, 160.0 mmol) and was used for the next step without purification.

MS-ESI: 198 (M⁺ +1, 72).

Elementary analysis:

Calculated: C 60.91% H 6.13% F 9.63% N 7.10%

Determined: C 60.99% H 6.15% F 9.60% N 7.07%

b) Synthesis of (2-fluoro-4-methoxycarbonyl-phenyl)-trimethyl-ammonium trifluoromethanesulfonate (**3b**)

To a stirred solution of 3,90 g (19,8 mmol) **3a** and 70 ml dichloromethane were added 32.5 g (198 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was stirred for 2.5 days at roomtemperature and diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **3b** was obtained in 80% yield (5,72 g, 15.84 mmol).

MS-ESI: 212 (M⁺, 76),

Elementary analysis:

Calculated: C 39.89% H 4.18% F 21.03% N 3.88%

Determined: C 39.93% H 4.20% F 21.01% N 3.84%

- 5 c) Synthesis of (4-carboxy-2-fluoro-phenyl)-trimethyl-ammonium trifluoro-methanesulfonate (3c).

A solution of 4,00 g (11.1 mmol) **3b**, 96 ml dest. water and 96 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude product **3c** was obtained in 92% yield (3.54 g, 10.2 mmol) and crude compound **3c** was used for the next step without purification.

- 15 MS-ESI: 198 (M^+ , 76),

Elementary analysis:

Calculated: C 38.04% H 3.77% F 21.88% N 4.03%

Determined: C 38.10% H 3.79% F 21.81% N 4,00%

- 20 d) Synthesis of (4-Trimethylammonium-3-fluoro-benzoyl)-Gly-Thr-Tyr-Ala-NH₂— triflate salt (3d).

To a stirred solution of 69,4 mg (0.2 mmol) **3c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Gly-Thr(OtBu)-Tyr(O-tBu)-Ala-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 69,4 mg (0.2 mmol) **3c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the

supernatant was lyophilized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**3d**) in 41.6 % yield - 20.0 mg (0.027 mmol).

MS-ESI: 590 (M^+ , 100)

5

e) Synthesis of

(3-Fluoro-4- ^{18}F -fluoro-benzoyl)-Gly-Thr-Tyr-Ala-NH₂ (**3e**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 10 500µl water) and MeCN (1.5ml) the fluorine containing water (405 MBq, 35µl) were added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **3d** (2mg) in anhydrous DMSO (300µl) was added. After heating at 80°C for 15 min. The crude reaction mixture was analyzed using an analytical 15 HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3µ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **3e** was confirmed by co-injection with the cold F-19 fluoro standard on the Econsphere analytical HPLC.

20

Example 4

a) 2-Fluoro-4-dimethylamino-benzoic acid methyl ester (**4a**)

To a stirred solution of 3.87 g (22.5 mmol) 2,4-difluoro-benzoic acid methyl ester (Apollo) 25 and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 5h at 55°C in an autoclave and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined 30 dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **4a** was obtained in 55% yield (2.44 g, 12.4 mmol).

35 MS-ESI: 198 ($M^+ + 1$, 86).

Elementary analysis:

Calculated: C 60.91% H 6.13% F 9.63% N 7.10%

Determined: C 60.95% H 6.14% F 9.59% N 7.08%

5 b) Synthesis of (3-fluoro-4-methoxycarbonyl-phenyl)-trimethyl-ammonium trifluoro-methanesulfonate (4b)

To a stirred solution of 2.46 g (12.5 mmol) **4a** and 50 ml dichloromethane were added 20.5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2 days. The solvent was carefully substituted by dichloroethan. The reaction mixture
10 was refluxed for 2 days and then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **4b** was obtained in 80% yield (3.61 g,
15 10.0 mmol).

MS-ESI: 212 (M^+ , 77),

Elementary analysis:

Calculated: C 39.89% H 4.18% F 21.03% N 3.88%

20 Determined: C 39.94% H 4.21% F 21.00% N 3.85%

c) Synthesis of (4-carboxy-3-fluoro-phenyl)-trimethyl-ammonium trifluoro-methanesulfonate (4c).

25 A solution of 2.50 g (6,92 mmol) **4b**, 60 ml dest. water and 60 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude product **4c** was obtained in 100% yield (2.40 g; 6,92 mmol) and crude compound **4c** was used for the
30 next step without purification.

MS-ESI: 198 (M^+ , 76),

Elementary analysis:

Calculated: C 38.04% H 3.77% F 21.88% N 4.03%

35 Determined: C 38.09% H 3,80% F 21.82% N 4,01%

d) Synthesis of (4-Trimethylammonium-2-fluoro-benzoyl)-Lys(N-dimethyl)-Ala-Gly-Leu-NH₂- triflate salt (4d).

To a stirred solution of 69,4 mg (0.2 mmol) **4c** in 1.5 ml dichloromethane and 0.25 ml
5 dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Lys(N-dimethyl)-Ala-Gly-Leu-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of
10 69,4 mg (0.2 mmol) **4c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a
15 mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the
20 desired product (**4d**) in 45,0 % yield - 24,7 mg (0.0293 mmol).

MS-ESI: 596 (M⁺, 100)

e) Synthesis of (2-Fluoro-4-[¹⁸F]-fluoro-benzoyl)-Lys(N-dimethyl)-Ala-Gly-Leu-NH₂ (**4e**).

25

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500µl water) and MeCN (1.5ml) the fluorine containing water (385 MBq, 39µl) were added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was
30 repeated again. A solution of **4d** (2mg) in anhydrous DMSO (300µl) was added. After heating at 90°C for 10 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18 , 53×7mm, 3µ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA,
35 gradient: 5-95% B in 7 mins. The F-18 labeled product **4e** was confirmed by co-injection with the cold F-19 fluoro standard on the Econsphere analytical HPLC.

Example 5

a) 4-Dimethylamino-2,3-difluoro-benzoic acid methyl ester (5a)

5

To a stirred solution of 4,28 g (22.5 mmol) 2,3,4-Trifluoro-benzoic acid methyl ester (Apollo) and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 5h at 55°C in an autoclave and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **5a** was obtained in 69 % yield (3.34 g, 15,5 mmol).

15

MS-ESI: 216 ($M^+ + 1$, 81).

Elementary analysis: C 55.81% H 5.15% F 17.66% N 6.51%

Determined: C 55.90% H 5.19% F 17.63% N 6,48%

20

b) Trifluoro-methanesulfonate(2,3-difluoro-4-methoxycarbonyl-phenyl)-trimethyl-ammonium (5b)

To a stirred solution of 2.69 g (12.5 mmol) **5a** and 50 ml dichloromethane were added 20.5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2 days then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **5b** was obtained in 82% yield (3,88 g, 10.3 mmol).

30

MS-ESI: 230 (M^+ , 34),

Elementary analysis:

Calculated: C 38.00% H 3.72% F 25.04% N 3.69%

35 Determined: C 38.04% H 3.74% F 25.00% N 3.67%

c) Trifluoro-methanesulfonate(4-carboxy-2,3-difluoro-phenyl)-trimethyl-ammonium; (**5c**).

A solution of 2.63 g (6,92 mmol) **5b**, 60 ml dest. water and 60 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude product **5c** was obtained in 89% yield (2.24 g; 6.16 mmol) and crude compound **5c** was used for the next step without purification.

10 MS-ESI: 216 (M^+ , 77),

Elementary analysis:

Calculated: C 36.17% H 3.31% F 26.01% N 3.83%

Determined: C 36.21% H 3.32% F 26.00% N 3,81%

15 d) Synthesis of (4-Trimethylammonium-2,3-difluoro-benzoyl)-Val-Arg-Ser-Gly-NH₂— triflate salt (**5d**).

To a stirred solution of 73 mg (0.2 mmol) **5c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val-Arg(Pbf)-Ser(OtBu)-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 73 mg (0.2 mmol) **5c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was lyophilized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**5d**) in 39,5 % yield - 19,6 mg (0.0256 mmol).

35 MS-ESI: 616 (M^+ , 100)

e) Synthesis of

(2,3-Difluoro-4-[¹⁸F]-fluoro-benzoyl)-Val-Arg-Ser-Gly-NH₂ (5e).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500µl water) and MeCN (1.5ml) the fluorine containing water (319 MBq, 35µl) were added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **5d** (2mg) in anhydrous DMSO (300µl) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3µ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **5e** was confirmed by co-injection with the cold F-19 fluoro standard on the Econsphere analytical HPLC.

15 Example 6

a) 4-Dimethylamino-2,6-difluoro-benzoic acid methyl ester (6a)

To a stirred solution of 4.28 g (22.5 mmol) 2,4,6-Trifluoro-benzoic acid methyl ester (Apollo) and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 5h at 55°C in an autoclave and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **6a** was obtained in 74 % yield (3.59 g, 16.7 mmol).

30 MS-ESI: 216 (M⁺ +1, 69).

Elementary analysis: C 55.81% H 5.15% F 17.66% N 6.51%

Determined: C 55.89% H 5.18% F 17.64% N 6.49%

b) Trifluoro-methanesulfonate(2,6-difluoro-4-methoxycarbonyl-phenyl)-trimethyl-ammonium (6b)

To a stirred solution of 2.69 g (12.5 mmol) **6a** and 50 ml dichloromethane were added 20.5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2 days then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **6b** was obtained in 78% yield (3.70 g, 9,75 mmol).

MS-ESI: 230 (M^+ , 55),

Elementary analysis:

Calculated: C 38.00% H 3.72% F 25.04% N 3.69%

Determined: C 38.05% H 3.73% F 25.01% N 3.68%

c) Trifluoro-methanesulfonate(4-carboxy-2,6-difluoro-phenyl)-trimethyl-ammonium; (**6c**).

A solution of 2.63 g (6,92 mmol) **6b**, 60 ml dest. water and 60 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude product **6c** was obtained in 92% yield (2.38 g; 6.37 mmol) and crude compound **6c** was used for the next step without purification.

MS-ESI: 216 (M^+ , 70),

Elementary analysis:

Calculated: C 36.17% H 3.31% F 26.01% N 3.83%

Determined: C 36.20% H 3.33% F 25.99% N 3,82%

d) Synthesis of (4-Trimethylammonium-2,6-difluoro-benzoyl)-Gly-Pro-Phe-Val-NH₂— triflate salt (**6d**).

To a stirred solution of 73 mg (0.2 mmol) **6c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Gly-Pro-Phe-Val-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol.

The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 73 mg

(0.2 mmol) **6c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with

5 dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml dwastilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water :
10 acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**6d**) in 19,6 % yield - 9,75 mg (0.0127 mmol).

MS-ESI: 616 (M^+ , 100)

e) Synthesis of (2,6-Difluoro-4-[^{18}F]-fluoro-benzoyl)-Gly-Pro-Phe-Val-NH₂ (**6e**).

15 To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500µl water) and MeCN (1.5ml) the fluorine containing water (320 MBq, 37µl) were added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was
20 repeated again. A solution of **6d** (2mg) in anhydrous DMSO (300µl) was added. After heating at 90°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18 , 53×7mm, 3µ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA,
25 gradient: 5-95% B in 7 mins. The F-18 labeled product **6e** was confirmed by co-injection with the cold F-19 fluoro standard on the Econsphere analytical HPLC.

Example 7

30 a) 2-Bromo-4-dimethylamino-benzoic acid methyl ester (**7a**)

To a stirred solution of 5,24 g (22.5 mmol) 2-Bromo-4-fluoro-benzoic acid methyl ester (Rarechemicals) and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction
35 mixture was stirred for 11h at 70°C in an autoclave and was concentrated with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane,

washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **7a** was
5 obtained in 70 % yield (4,08 g, 15.8 mmol).

MS-ESI: 258/560 ($M^+ + 1$, 88/83).

Elementary analysis: C 46.53% H 4.69% N 5.43%

Determined: C 46,60% H 4.71% N 5.42%

10

b) Trifluoro-methanesulfonate(3-bromo-4-methoxycarbonyl-phenyl)-trimethyl-ammonium (7b)

To a stirred solution of 2,69 g (12,5 mmol) **7a** and 50 ml dichloromethane were added
15 20,5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2 days then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient
20 1:99 to 80:20). The desired compound **7b** was obtained in 69% yield (3,66 g, 8,63 mmol).

MS-ESI: 273/275 ($M^+ + 1$, 78/72),

Elementary analysis:

Calculated: C 34.14% H 3.58% F 13.50% N 3.32%

25 Determined: C 34.17% H 3.59% F 13.47% N 3.31%

c) Trifluoro-methanesulfonate(3-bromo-4-carboxy-phenyl)-trimethyl-ammonium; (7c).

A solution of 2,92 g (6,92 mmol) **7b**, 60 ml dest. water and 60 ml trifluoroacetic acid was
30 refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude product **7c** was obtained in 87% yield (2,46 g; 6.02 mmol) and crude compound **7c** was used for the next step without purification.

35

MS-ESI: 258/260 (M^+ , 64/59),

Elementary analysis:

Calculated: C 32.37% H 3.21% F 13.96% N 3.43%

Determined: C 32.41% H 3,22% F 13.94% N 3,42%

- 5 d) Synthesis of (4-Trimethylammonium-2-bromo-benzoyl)-Gly-Phe-Ile-Gly-NH₂-triflate salt (7d).

To a stirred solution of 81.6 mg (0.2 mmol) **7c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Gly-Phe-Ile-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 81.6 mg (0.2 mmol) **7c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**7d**) in 24,6 % yield - 12,5 mg (0,016 mmol).

25

MS-ESI: 633/635 (M⁺, 100/88)

- e) Synthesis of [¹⁸F]-(2-Bromo-4-fluoro-benzoyl)-Gly-Phe-Ile-Gly-NH₂ (**7e**).

30 To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500µl water) and MeCN (1.5ml) the fluorine containing water (336 MBq, 35µl) were added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **7d** (2mg) in anhydrous DMSO (300µl) was added. After heating at 90°C for 12 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B:

35

MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18 , 53×7mm, 3μ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **7e** was confirmed by co-injection with the cold F-19 fluoro standard on the Econosphere analytical HPLC.

5

Example 8

a) 4-Dimethylamino-2-nitro-benzoic acid methyl ester (**8a**)

- 10 To a stirred solution of 4,48 g (22.5 mmol) 4-fluoro-2-nitro-benzoic acid methyl ester (J. Fluorine Chem.; 63; 1-2; (1993); 25-30) and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 7h at 60°C in an autoclave and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane,
- 15 washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **8a** was obtained in 61 % yield (3,08 g, 13,7 mmol).

20

MS-ESI: 225 (M⁺ +1, 71).

Elementary analysis: C 53.57% H 5.39% N 12.49%

Determined: C 53.60% H 5.40% N 12,47%

- 25 b) Trifluoro-methanesulfonate(4-methoxycarbonyl-3-nitro-phenyl)-trimethylammonium (**8b**)

- To a stirred solution of 2,80 g (12,5 mmol) **8a** and 50 ml dichloromethane were added 20,5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2 days then cooled to room temperature. Diethylether was added. The desired
- 30 compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **8b** was obtained in 45% yield (2,18 g, 5,63 mmol).

- 35 MS-ESI: 239 (M⁺ , 89),

Elementary analysis:

Calculated: C 37.12% H 3.89% F 14.68% N 7.21%

Determined: C 37.15% H 3.90% F 14.67% N 7.19%

c) Trifluoro-methanesulfonate(4-carboxy-3-nitro-phenyl)-trimethyl-ammonium (8c).

5

A solution of 2.68 g (6,92 mmol) **8b**, 60 ml dest. water and 60 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude was obtained in 96% yield (2,48 g; 6.64 mmol) and crude compound **8c** was used for the next step without purification.

MS-ESI: 225 (M^+ , 66),

Elementary analysis:

15 Calculated: C 35.30% H 3.50% F 15.23% N 7.48%

Determined: C 35.31% H 3,50% F 15.23% N 7,47%

d) Synthesis of (4-Trimethylammonium-2-nitro-benzoyl)-Ser-Thr-Val-Gly-NH₂-triflate salt (8d).

20

To a stirred solution of 75,0 mg (0.2 mmol) **8c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Ser(Ot-Bu)-Thr(OtBu)-Val-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 75,0 mg (0.2 mmol) **8c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS

with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**8d**) in 35% yield (16,3 mg, 0,0228 mmol).

5 MS-ESI: 569 (M^+ , 100)

e) Synthesis of [^{18}F]- (2-Nitro-4-fluoro-benzoyl)-Ser-Thr-Val-Gly- NH_2 (**8e**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in
10 500 μl water) and MeCN (1.5ml) the fluorine containing water (376 MBq, 35 μl) was added.
The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen.
Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated
again. A solution of **8d** (2mg) in anhydrous DMSO (300 μl) was added. After heating at
80°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC
15 (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN,
gradient: 5%-95% B in 7 mins or Column Econosphere C18 , 53 $\times$ 7mm, 3 μ , 3ml/min
(Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient:
5-95% B in 7 mins. The F-18 labeled product **8e** was confirmed by co-injection with the
cold F-19 fluoro standard on the Econsphere analytical HPLC.

20

Example 9

a) 2-Cyano-4-dimethylamino-benzoic acid methyl ester (**9a**)

25 To a stirred solution of 4,03 g (22.5 mmol) 2-Cyano-4-fluoro-benzoic acid methyl ester (J.
Med. Chem., 35; 24; (1992); 4613-4627) and 60.0 ml dimethylsulphoxid were added 2.23
g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate.
The reaction mixture was stirred for 9h at 60°C in an autoclave and was concentrated with
high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane,
30 washed twice with water. The combined water phases were extracted with
dichloromethane. The combined dichloromethane phases were washed with diluted
sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The
oily crude was purified by column chromatography and the desired product **9a** was
obtained in 85% yield (3,90 g, 19.1 mmol).

35

MS-ESI: 205 ($M^+ + 1$, 81).

Elementary analysis: C 64.69% H 5.92% N 13.72%

Determined: C 64.72% H 5.95% N 13.70%

- 5 b) Trifluoro-methanesulfonate(3-cyano-4-methoxycarbonyl-phenyl)-trimethylammonium; (9b)

To a stirred solution of 2,55 g (12,5 mmol) **9a** and 50 ml dichloromethane were added 20,5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed
10 for 2 days then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **9b** was obtained in 78% yield (3,59 g, 9,75 mmol).

15

MS-ESI: 219 ($M^+ + 1$, 79),

Elementary analysis:

Calculated: C 42.39% H 4.10% F 15.47% N 7.61%

Determined: C 42.41% H 4.11% F 15.42% N 7.60%

20

- c) Trifluoro-methanesulfonate(4-carboxy-3-cyano-phenyl)-trimethyl-ammonium; (9c).

A solution of 2,55 g (6,92 mmol) **9b**, 60 ml dest. water and 60 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump
25 vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude was obtained in 86% yield (2,10 g; 5.95 mmol) and crude compound **9c** was used for the next step without purification.

- 30 MS-ESI: 205 (M^+ , 76),

Elementary analysis:

Calculated: C 40.68% H 3.70% F 16.09% N 7.91%

Determined: C 40.69% H 3,71% F 16.07% N 7,90%

- 35 d) Synthesis of (4-Trimethylammonium-2-cyano-benzoyl)-Arg-Val-Gly-Phe-NH₂- triflate salt (9d).

To a stirred solution of 71.0 mg (0.2 mmol) **9c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Arg(Pbf)-Val-Gly-Phe-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 71.0 mg (0.2 mmol) **9c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**9d**) obtained in 38% yield (20.0 mg, 0.0247 mmol).

MS-ESI: 665 (M⁺, 100)

e) Synthesis of [¹⁸F]-(2-Cyano-4-fluoro-benzoyl)-Arg-Val-Gly-Phe-NH₂ (**9e**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500µl water) and MeCN (1.5ml) the fluorine containing water (309 MBq, 35µl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **9d** (2mg) in anhydrous DMSO (300µl) was added. After heating at 80°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3µ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **9e** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

35

Example 10

a) 4-Dimethylamino-3-methanesulfonyl-benzoic acid methyl ester (10a)

To a stirred solution of 5,23 g (22.5 mmol) 4-Fluoro-3-methanesulfonyl-benzoic acid methyl ester (J. Med. Chem.; 40; 13; 1997; 2017-2034) and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 7h at 60°C in an autoclave and was concentrated with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **10a** was obtained in 56 % yield (3,24 g, 12,6 mmol).

MS-ESI: 258 ($M^+ + 1$, 81).

Elementary analysis: C 51.35% H 5.88% N 5.44%

Determined: C 51.37% H 5.90% N 5.42%

b) Trifluoro-methanesulfonate(2-methanesulfonyl-4-methoxycarbonyl-phenyl)-trimethyl-ammonium, (10b)

To a stirred solution of 3,22 g (12,5 mmol) **10a** and 50 ml dichloromethane were added 20,5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2 days then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **10b** was obtained in 58% yield (3,05 g, 7,25 mmol).

30

MS-ESI: 272 (M^+ , 88),

Elementary analysis:

Calculated: C 37.05% H 4.31% F 13.52% N 3.32%

Determined: C 37.09% H 4.33% F 13.50% N 3.31%

35

c) Trifluoro-methanesulfonate(4-carboxy-2-methanesulfonyl-phenyl)-trimethylammonium; (10c).

A solution of 2,91 g (6,92 mmol) **10b**, 60 ml dest. water and 60 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude was obtained in 96% yield (2,70 g; 6.64 mmol) and crude compound **10c** was used for the next step without purification.

MS-ESI: 258 (M^+ , 93),

Elementary analysis:

Calculated: C 35.38% H 3.96% F 13.99% N 3.44%

Determined: C 35.39% H 3,96% F 13.97% N 3,44%

d) Synthesis of (4-Trimethylammonium-3-methanesulfonyl-benzoyl)-Gly-Phe-Val-Leu-NH₂- triflate salt (10d).

To a stirred solution of 81.0 mg (0.2 mmol) **10c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Gly-Phe-Val-Leu-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 81.0 mg (0.2 mmol) **10c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml dwastilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**10d**) in 33 % yield - 17,7 mg (0.021 mmol).

MS-ESI: 675 (M^+ , 100)

e) Synthesis of [^{18}F]- (3-Methanesulfonyl-4-fluoro-benzoyl)-Gly-Phe-Val-Leu-NH₂ (10e).

- 5 To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μl water) and MeCN (1.5ml) the fluorine containing water (399 MBq, 35 μl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **10d** (2mg) in anhydrous DMSO (300 μl) was added. After heating at
- 10 80°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **10e** was confirmed by co-injection with the
- 15 non-radioactive F-19 fluoro standard on the Econosphere analytical HPLC.

Example 11

a) 4-Dimethylamino-3-nitro-benzoic acid methyl ester (11a)

20

- To a stirred solution of 4.48 g (22.5 mmol) 4-Fluoro-3-nitro-benzoic acid methyl ester (Bioorg. Med. Chem.; 6; 8; 1998; 1185 - 1208) and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 8h at 60°C in an autoclave and was
- 25 reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **11a** was
- 30 obtained in 69 % yield (3.48 g, 15.5 mmol).

MS-ESI: 225 ($M^+ + 1$, 74).

Elementary analysis: C 53.57% H 5.39% N 12.49%

Determined: C 53.62% H 5.42% N 12.46%

35

b) Trifluoro-methanesulfonate(4-methoxycarbonyl-2-nitro-phenyl)-trimethylammonium (11b)

To a stirred solution of 2,80 g (12,5 mmol) **11a** and 50 ml dichloromethane were added
5 20,5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed
for 2 days then cooled to room temperature. Diethylether was added. The desired
compound precipitates and the solvent was decanted. The solid was washed extensively
(ten times) with large amounts of diethylether. The solid was dried by use of oil pump
vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient
10 1:99 to 80:20). The desired compound **11b** was obtained in 71% yield (3,44 g, 8,88
mmol).

MS-ESI: 239 (M^+ , 82),

Elementary analysis:

15 Calculated: C 37.12% H 3.89% F 14.68% N 7.21%
Determined: C 37.14% H 3.91% F 14.67% N 7.20%

c) Trifluoro-methanesulfonate(4-carboxy-2-nitro-phenyl)-trimethyl-ammonium (11c).

20 A solution of 2.68 g (6,92 mmol) **11b**, 60 ml dest. water and 60 ml trifluoroacetic acid was
refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump
vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed
extensively with diethyl ether and dried by oil pump vacuum. The solid crude was obtained
in 93% yield (2,41 g; 6.44 mmol) and crude compound **11c** was used for the next step
25 without purification.

MS-ESI: 225 (M^+ , 66),

Elementary analysis:

Calculated: C 35.30% H 3.50% F 15.23% N 7.48%
30 Determined: C 35.32% H 3,51% F 15.21% N 7,46%

d) Synthesis of (4-Trimethylammonium-3-nitro-benzoyl)-Thr-Val-Phe-Leu-NH₂- triflate salt (11d).

35 To a stirred solution of 75,0 mg (0.2 mmol) **11c** in 1.5 ml dichloromethane and 0.25 ml
dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2

mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Thr(OtBu)-Val-Phe-Leu-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of
5 75,0 mg (0.2 mmol) **11c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a
10 mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the
15 desired product (**11d**) in 56% yield (30.1mg, 0.0364 mmol).

MS-ESI: 686 (M⁺, 100)

e) Synthesis of [¹⁸F]-(3-Nitro-4-fluoro-benzoyl)- Thr-Val-Phe-Leu-NH₂ (**11e**).

20

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500μl water) and MeCN (1.5ml) the fluorine containing water (344 MBq, 35μl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated
25 again. A solution of **11d** (2mg) in anhydrous DMSO (300μl) was added. After heating at 65°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8μ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3μ, 3ml/min (Alltech), solvent A: H₂O + 0.1% TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient:
30 5-95% B in 7 mins. The F-18 labeled product **11e** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

Example 12

35 a) 4-Dimethylamino-3-trifluoromethyl-benzoic acid methyl ester (**12a**)

To a stirred solution of 4,48 g (22.5 mmol) 4-Fluoro-3-trifluoromethyl-benzoic acid methyl ester (Rarechem) and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 8h at 60°C in an autoclave and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **12a** was obtained in 72 % yield (4,00 g, 16,2 mmol).

MS-ESI: 248 (M^+ +1, 100).

Elementary analysis: C 53.44% H 4.89% F 23.05% N 5.67%

Determined: C 53.48% H 4.90% F 23.03% N 5.65%

b) Trifluoro-methanesulfonate(4-methoxycarbonyl-2-trifluoromethyl-phenyl)-trimethyl-ammonium (**12b**)

To a stirred solution of 3.09 g (12.5 mmol) **12a** and 50 ml dichloromethane were added 20.5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2 days then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **12b** was obtained in 69% yield (3.55 g, 8,63 mmol).

MS-ESI: 262 (M^+ , 67),

Elementary analysis:

Calculated: C 37.96% H 3.68% F 27.71% N 3.41%

Determined: C 38.00% H 3.62% F 27.68% N 3.40%

c) Trifluoro-methanesulfonate(4-carboxy-2-trifluoromethyl-phenyl)-trimethyl-ammonium (**12c**).

A solution of 2.84 g (6,92 mmol) **12b**, 60 ml dest. water and 60 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude was obtained
5 in 89% yield (2.45 g; 6.16 mmol) and crude compound **12c** was used for the next step without purification.

MS-ESI: 248 (M^+ , 100),

Elementary analysis:

10 Calculated: C 36.28% H 3.30% F 28.69% N 3.53%
Determined: C 36.29% H 3.31% F 28.67% N 3.51%

d) Synthesis of (4-Trimethylammonium-3-trifluormethyl-benzoyl)-Val- β Ala-Phe-Gly-NH₂-triflate salt (**12d**).

15

To a stirred solution of 79,4 mg (0.2 mmol) **12c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-Phe-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol.

20 The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 79,4 mg (0.2 mmol) **12c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was
25 again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was
30 liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**12d**) in 21.5 % yield - 10.9 mg (0.014 mmol).

MS-ESI: 636 (M^+ , 100)

35 e) Synthesis of [¹⁸F]-(3-Trifluormethyl-4-fluoro-benzoyl)-Val- β Ala-Phe-Gly-NH₂ (**12e**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500µl water) and MeCN (1.5ml) the fluorine containing water (356 MBq, 35µl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. The step was repeated
5 again. A solution of **12d** (2mg) in anhydrous DMSO (300µl) was added. After heating at 50°C for 18 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3µ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-
10 95% B in 7 mins. The F-18 labeled product **12e** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econosphere analytical HPLC.

Example 13

15 a) 4-Dimethylamino-2-trifluoromethyl-benzoic acid methyl ester (13a)

To a stirred solution of 4,48 g (22.5 mmol) 4-Fluoro-2-trifluoromethyl-benzoic acid methyl ester (Rarechem) and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction
20 mixture was stirred for 8h at 60°C in an autoclave and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by
25 column chromatography and the desired product **13a** was obtained in 72 % yield (4,00 g, 16,2 mmol).

MS-ESI: 248 (M⁺ +1, 78).

Elementary analysis:	C	53.44%	H	4.89%	F	23.05%	N	5.67%
30 Determined:	C	53.46%	H	4.91%	F	23,04%	N	5.64%

b) Trifluoro-methanesulfonate(4-methoxycarbonyl-3-trifluoromethyl-phenyl)-trimethyl-ammonium (13b)

35 To a stirred solution of 3,09 g (12,5 mmol) **13a** and 50 ml dichloromethane were added 20,5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed

for 2 days then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **13b** was obtained in 69% yield (3,55 g, 8,63 mmol).

MS-ESI: 262 (M^+ , 87),

Elementary analysis:

Calculated: C 37.96% H 3.68% F 27.71% N 3.41%

Determined: C 38.01% H 3.63% F 27.69% N 3.41%

c) Trifluoro-methanesulfonate(4-carboxy-3-trifluoromethyl-phenyl)-trimethylammonium (**13c**).

15

A solution of 2,84 g (6,92 mmol) **13b**, 60 ml dest. water and 60 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude was obtained in 89% yield (2,45 g; 6.16 mmol) and crude compound **13c** was used for the next step without purification.

MS-ESI: 248 (M^+ , 59),

Elementary analysis:

Calculated: C 36.28% H 3.30% F 28.69% N 3.53%

Determined: C 36.30% H 3,32% F 28.67% N 3,52%

d) Synthesis of (4-Trimethylammonium-2-trifluormethyl-benzoyl)-Val- β Ala-His(π -Me)-Gly-NH₂- triflate salt (**13d**)

30

To a stirred solution of 79,4 mg (0.2 mmol) **13c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-His(π -Me)-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of

35

79,4 mg (0.2 mmol) **13c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed
5 extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS
10 with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**13d**).
MS-ESI: 640 (M^+ , 100)

e) Synthesis of ($[^{18}F]$ -4-Fluoro-2-trifluormethyl-benzoyl)-Val- β Ala-His(π -Me)-Gly-NH₂ (**13e**)
15
To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (321 MBq, 35 μ l) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated
20 again. A solution of **13d** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 75°C for 20 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18 , 53 $\times$ 7mm, 3 μ , 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient:
25 5-95% B in 7 mins. The F-18 labeled product **13e** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econosphere analytical HPLC.

Example 14

30 a) 4-Fluoro-3-trifluoromethoxy-benzoic acid methyl ester (**14a**).

To a stirred suspension of 21,2g (96,9 mmol) 4-Fluoro-3-trifluoromethoxy-benzoic acid (JRD-Fluoro) and 161 ml methanol were added 30.4 g (387,6 mmol) acetyl chloride drop
wisely at 0°C. The reaction mixture was stirred over night, filtered and concentrated. The
35 residue was diluted with dichloromethane, washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The residue was

purified by column chromatography (hexane : ethylacetate). The desired product **14a** was obtained in 75% yield (17.3 g; 72.7 mmol)

MS-ESI: 239 ($M^+ + 1$, 66),

5 Elementary analysis:

Calculated: C 45.39% H 2.54% F 31.91%

Determined: C 45.41% H 2,52% F 31.89%

b) 4-Dimethylamino-3-trifluoromethoxy-benzoic acid methyl ester (**14b**)

10

To a stirred solution of 5.36 g (22.5 mmol) **14a** and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 8h at 60°C in an autoclave and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with
15 dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **14b** was obtained in 69 % yield (4,09 g, 15.5 mmol).

20

MS-ESI: 264 ($M^+ + 1$, 100).

Elementary analysis: C 50.20% H 4.60% F 21.65% N 5.32%

Determined: C 50.22% H 4.61% F 21,64% N 5.31%

25 c) Trifluoro-methanesulfonate(4-methoxycarbonyl-2-trifluoromethoxy-phenyl)-trimethyl-ammonium (**14c**)

To a stirred solution of 3,29 g (12,5 mmol) **14b** and 50 ml dichloroethane were added 20,5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2
30 days then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **14c** was obtained in 57% yield (3,06 g, 7,13 mmol).

35

MS-ESI: 278 (M^+ , 82),

Elementary analysis:

Calculated: C 36.54% H 3.54% F 26.68% N 3.28%

Determined: C 36.56% H 3.56% F 27.67% N 3.26%

- 5 d) Trifluoro-methanesulfonate(4-carboxy-2-trifluoromethoxy-phenyl)-trimethylammonium (14d).

A solution of 2,95g (6,92 mmol) **14c**, 60 ml dest. water and 60 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump
10 vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude was obtained in 94% yield (2,68 g; 6.50 mmol) and crude compound **14d** was used for the next step without purification.

- 15 MS-ESI: 264 (M^+ , 100),

Elementary analysis:

Calculated: C 34.87% H 3.17% F 27.58% N 3.39%

Determined: C 34.89% H 3,19% F 27.56% N 3,38%

- 20 e) Synthesis of (4-Trimethylammonium-3-trifluormethoxy-benzoyl)-Val- β Ala-Phe-Gly-NH₂-triflate salt (14e)

To a stirred solution of 82.6 mg (0.2 mmol) **14d** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2
25 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-Phe-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 82.6 mg (0.2 mmol) **14d** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added
30 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for
35 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was

liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**14e**) in 28 % yield - 14,6 mg (0.0182 mmol).

MS-ESI: 653 (M^+ , 100)

5

f) Synthesis of

($[^{18}F]$ -4-Fluoro-3-trifluormethoxy-benzoyl)-Val- β Ala-Phe-Gly-NH₂ (**14f**)

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (321 MBq, 35 μ l) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **14e** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 65°C for 10 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **14f** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

20

Example 15

a) 5-Dimethylamino-2-trifluoromethyl-benzoic acid methyl ester (**15a**)

To a stirred solution of 4,48 g (22.5 mmol) 5-Fluoro-2-trifluoromethyl-benzoic acid methyl ester (Rarechem) and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 8h at 60°C in an autoclave and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **15a** was obtained in 72 % yield (4,00 g, 16,2 mmol).

35

MS-ESI: 248 ($M^+ + 1$, 100).

Elementary analysis: C 53.44% H 4.89% F 23.05% N 5.67%

Determined: C 53.45% H 4.90% F 23.05% N 5.65%

5 b) Trifluoro-methanesulfonate(3-methoxycarbonyl-4-trifluoromethyl-phenyl)-trimethyl-ammonium (15b)

To a stirred solution of 3,09 g (12,5 mmol) **15a** and 50 ml dichloromethane were added 20,5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2 days then cooled to room temperature. Diethylether was added. The desired
10 compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **15b** was obtained in 69% yield (3,55 g, 8,63 mmol).

15

MS-ESI: 262 (M^+ , 100),

Elementary analysis:

Calculated: C 37.96% H 3.68% F 27.71% N 3.41%

Determined: C 38.00% H 3.69% F 27.68% N 3.40%

20

c) Trifluoro-methanesulfonate(3-carboxy-4-trifluoromethyl-phenyl)-trimethyl-ammonium (15c).

A solution of 2,84 g (6,92 mmol) **15b**, 60 ml dest. water and 60 ml trifluoroacetic acid was
25 refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude was obtained in 89% yield (2,45 g; 6.16 mmol) and crude compound **15c** was used for the next step without purification.

30

MS-ESI: 248 (M^+ , 45),

Elementary analysis:

Calculated: C 36.28% H 3.30% F 28.69% N 3.53%

Determined: C 36.31% H 3,31% F 28.68% N 3,51%

35

d) Synthesis of (5-Trimethylammonium-2-trifluormethyl-benzoyl)-Val-βAla-Trp-Gly-NH₂-triflate salt (**15d**)

To a stirred solution of 79,4 mg (0.2 mmol) **15c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val-βAla-Trp(N-Boc)-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 79,4 mg (0.2 mmol) **15c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**15d**) in 26 % yield - 13,9 mg (0.017 mmol).
MS-ESI: 675 (M⁺, 100)

e) Synthesis of ([¹⁸F]-5-Fluoro-2-trifluormethyl-benzoyl)-Val-βAla-Trp-Gly-NH₂ (**15e**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500μl water) and MeCN (1.5ml) the fluorine containing water (316MBq, 33μl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **15d** (2mg) in anhydrous DMSO (300μl) was added. After heating at 75°C for 18 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8μ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3μ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **15e** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econosphere analytical HPLC.

Example 16

a) 2-Bromo-5-dimethylamino-benzoic acid methyl ester (16a)

5 To a stirred solution of 5,24 g (22.5 mmol) 2-Bromo-5-fluoro-benzoic acid methyl ester (Rarechemicals) and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 11h at 70°C in an autoclave and was concentrated with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with
10 dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **16a** was obtained in 70 % yield (4,08 g, 15.8 mmol).

15

MS-ESI: 258/560 ($M^+ + 1$, 90/81).

Elementary analysis: C 46.53% H 4.69% N 5.43%

Determined: C 46,59% H 4.72% N 5.41%

20 b) Trifluoro-methanesulfonate(4-bromo-3-methoxycarbonyl-phenyl)-trimethylammonium (16b)

To a stirred solution of 2,69 g (12,5 mmol) **16a** and 50 ml dichloromethane were added 20,5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed
25 for 2 days then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **16b** was obtained in 79% yield (4,17 g, 9,88
30 mmol).

MS-ESI: 272/274 (M^+ , 89/80),

Elementary analysis:

Calculated: C 34.14% H 3.58% F 13.50% N 3.32%

35 Determined: C 34.16% H 3.60% F 13.48% N 3.30%

c) Trifluoro-methanesulfonate(4-bromo-3-carboxy-phenyl)-trimethyl-ammonium (16c).

A solution of 2,92 g (6,92 mmol) **16b**, 60 ml dest. water and 60 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude product **16c** was obtained in 87% yield (2,46 g; 6.02 mmol) and crude compound **16c** was used for the next step without purification.

MS-ESI: 258/260 (M^+ , 78/69),

Elementary analysis:

Calculated: C 32.37% H 3.21% F 13.96% N 3.43%

Determined: C 32.40% H 3,22% F 13.95% N 3,41%

d) Synthesis of (5-Trimethylammonium-2-bromo-benzoyl)-Val- β Ala-Arg-Gly-NH₂- triflate salt (16d).

To a stirred solution of 81.4 mg (0.2 mmol) **16c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-Arg(Pbf)-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 81.4 mg (0.2 mmol) **16c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**16d**) in 52% yield (27,2 mg, 0,0338 mmol).

MS-ESI: 655/657 (M^+ , 100/82)

e) Synthesis of [^{18}F]- (2-Bromo-5-fluoro-benzoyl)-Val- β Ala-Arg-Gly-NH₂ (**16e**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (334 MBq, 35 μ l) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **16d** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 90°C for 20 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **16e** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econosphere analytical HPLC.

15

Example 17

a) 5-Dimethylamino-2-methanesulfonyl-benzoic acid methyl ester (**17a**)

To a stirred solution of 5.23 g (22.5 mmol) 5-Fluoro-2-methanesulfonyl-benzoic acid methyl ester (J. Med. Chem.; 40; 13; 1997; 2017-2034) and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 7h at 60°C in an autoclave and was concentrated with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **17a** was obtained in 56 % yield (3.24 g, 12.6 mmol).

30

MS-ESI: 257 ($\text{M}^+ + 1$, 75).

Elementary analysis: C 51.35% H 5.88% N 5.44%

Determined: C 51.37% H 5.88% N 5.42%

35 b) Trifluoro-methanesulfonate(4-methanesulfonyl-3-methoxycarbonyl-phenyl)-trimethyl-ammonium (**17b**).

To a stirred solution of 3,22 g (12,5 mmol) **17a** and 50 ml dichloromethane were added 20,5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2 days then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **17b** was obtained in 58% yield (3,05 g, 7,25 mmol).

MS-ESI: 272 (M^+ , 69),

Elementary analysis:

Calculated: C 37.05% H 4.31% F 13.52% N 3.32%

Determined: C 37.07% H 4.33% F 13.48% N 3.31%

c) Trifluoro-methanesulfonate(3-carboxy-4-methanesulfonyl-phenyl)-trimethyl-ammonium; (**17c**).

A solution of 2,91 g (6,92 mmol) **17b**, 60 ml dest. water and 60 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude was obtained in 96% yield (2,70 g; 6.64 mmol) and crude compound **17c** was used for the next step without purification.

MS-ESI: 258 ($M^+ + 1$, 69),

Elementary analysis:

Calculated: C 35.38% H 3.96% F 13.99% N 3.44%

Determined: C 35.40% H 3,97% F 13.96% N 3,43%

d) Synthesis of (5-Trimethylammonium-2-methanesulfonyl-benzoyl)-Val- β Ala-Arg-Gly-NH₂- triflate salt (**17d**).

To a stirred solution of 81.0 mg (0.2 mmol) **17c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-

Val-βAla-Arg(Pbf)-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 81.0 mg (0.2 mmol) **17c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**17d**) in 33% yield (17,2 g, 0.0214 mmol).

MS-ESI: 656 (M⁺ , 100)

e) Synthesis of [¹⁸F]-(2-Methanesulfonyl-5-fluoro-benzoyl)-Val-βAla-Arg-Gly-NH₂ (**17e**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500μl water) and MeCN (1.5ml) the fluorine containing water (366 MBq, 35μl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **17d** (2mg) in anhydrous DMSO (300μl) was added. After heating at 70°C for 18 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8μ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18 , 53×7mm, 3μ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **17e** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

Example 18

a) 2-Chloro-5-dimethylamino-benzoic acid methyl ester (**18a**)

To a stirred solution of 4,00g (20,6 mmol) 2-Chloro-5-fluoro-benzoic acid methyl ester (Rarechem) and 60 ml dimethylsulphoxid were added 2,03 g (24,7 mmol) dimethylamine hydrochloride and 5,97 g (43,2 mmol) potassium carbonate. The reaction mixture was stirred over night and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude product **18a** was obtained in 99% yield (4,46 g, 20,9 mmol) and was used for the next step without purification.

MS-ESI: 213 / 215 ($M^+ + 1$, 78 / 53).

Elementary analysis:

Calculated: C 56.21% H 5.66% N 6.56%

Determined: C 56.29% H 5,68% N 6.55%

b) Synthesis of (4-chloro-3-methoxycarbonyl-phenyl)-trimethyl-ammonium trifluoromethanesulfonate (**18b**)

To a stirred solution of 4.49 g (21.0 mmol) **18a** and 75 ml dichloromethane were added 34,5 g (21.0 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was stirred for 2 days at room temperature. 17 g (10 mmol) methyltriflate (Aldrich) were added and the reaction mixture was stirred at 40°C for 20 h. The reaction mixture was cooled to 20 °C and diethylether was added. The desired compound precipitates and the solvent was decanted and the solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **18b** was obtained in 86% yield (6,78 g, 18,1 mmol).

MS-ESI: 227/229 (M^+ , 78/21),

Elementary analysis:

Calculated: C 38.15% H 4.00% F 15.09% N 3.71%

Determined: C 38.17% H 4.03% F 15.05% N 3.70%

c) Synthesis of (3-carboxy-4-chloro-phenyl)-trimethyl-ammonium trifluoromethanesulfonate (**18c**).

A solution of 0.5 g (1,32 mmol) **18b**, 12 ml dest. water and 12 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude product **18c** was obtained in 98% yield (471 mg, 1,3 mmol) and crude compound **18c** was used for the next step without purification.

MS-ESI: 214 / 216 (M^+ , 89 / 51),

Elementary analysis:

Calculated:	C	36.32%	H	3.60%	F	15.67%	N	3.85%
Determined:	C	36.37%	H	3,63%	F	15.61%	N	3,83%

d) Synthesis of (5-Trimethylammonium-2-chloro-benzoyl)-Val- β Ala-Arg-Gly-NH₂- triflate salt (**18d**).

15

To a stirred solution of 73.0 mg (0.2 mmol) **18c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-Arg(Pbf)-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 73.0 mg (0.2 mmol) **18c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**18d**) in 31 % yield - 15,3 mg (0.020 mmol).

MS-ESI: 611/613 ($M^+ + 1$, 100/41).

35

e) Synthesis of [¹⁸F]-(5-Fluoro-2-chloro-benzoyl)-Val- β Ala-Arg-Gly-NH₂ (**18e**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500µl water) and MeCN (1.5ml) the fluorine containing water (384 MBq, 35µl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen.

- 5 Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **18d** (2mg) in anhydrous DMSO (300µl) was added. After heating at 80°C for 20 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3µ, 3ml/min
- 10 (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **18e** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

Example19

15

a) 5-Dimethylamino-2-nitro-benzoic acid methyl ester (19a)

- To a stirred solution of 4.48 g (22.5 mmol) 5-Fluoro-5-nitro-benzoic acid methyl ester (Rarechem) and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine
- 20 hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 8h at 60°C in an autoclave and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution,
- 25 dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **19a** was obtained in 69 % yield (3.49 g, 15.5 mmol).

MS-ESI: 225 (M⁺ +1, 52).

- 30 Elementary analysis: C 53.57% H 5.39% N 12.49%
Determined: C 53.61% H 5.40% N 12.47%

b) Trifluoro-methanesulfonate(3-methoxycarbonyl-4-nitro-phenyl)-trimethylammonium (19b)

35

To a stirred solution of 2,80 g (12,5 mmol) **19a** and 50 ml dichloromethane were added 20,5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2 days then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **19b** was obtained in 71% yield (3,44 g, 8,88 mmol).

MS-ESI: 239 (M^+ , 69),

Elementary analysis:

Calculated: C 37.12% H 3.89% F 14.68% N 7.21%

Determined: C 37.14% H 3.91% F 14.66% N 7.20%

c) Trifluoro-methanesulfonate(3-carboxy-4-nitro-phenyl)-trimethyl-ammonium (**19c**).

A solution of 2,46 g (6,92 mmol) **19b**, 60 ml dest. water and 60 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude was obtained in 93% yield (2,41 g; 6.44 mmol) and crude compound **19c** was used for the next step without purification.

MS-ESI: 225 (M^+ , 96),

Elementary analysis:

Calculated: C 35.30% H 3.50% F 15.23% N 7.48%

Determined: C 35.34% H 3,52% F 15.23% N 7,47%

d) Synthesis of (2-Nitro-5-trimethylammonium-benzoyl)-Val- β Ala-Phe-Gly-NH₂ – triflate salt (**19d**).

To a stirred solution of 75,0 mg (0.2 mmol) **19c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-Phe-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with

dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 75,0 mg (0.2 mmol) **19c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken
5 intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was
10 liophilized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent **19d** was obtained in 42% yield (20,8 mg, 0.0273 mmol).

MS-ESI: 614 (M^+ , 100).

15

e) Synthesis of

[^{18}F]- (5-Fluoro-2-nitro-benzoyl)-Val- β Ala-Phe-Gly-NH₂ (**19e**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in
20 500 μ l water) and MeCN (1.5ml) the fluorine containing water (311 MBq, 35 μ l) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **19d** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 70°C for 12 min. The crude reaction mixture was analyzed using an analytical HPLC
25 (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **19e** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

30

Example 20

a) 2-Chloro-4-dimethylamino-5-methanesulfonyl-benzoic acid methyl ester (**20a**)

35 To a stirred solution of 6,00 g (22.5 mmol) 2-Chloro-4-fluoro-5-methanesulfonyl-benzoic acid methyl ester (J. Med. Chem.; 40; 13; 1997; 2017-2034) and 60.0 ml

dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 15h at 65°C in an autoclave and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **20a** was obtained in 59 % yield (3,87 g, 13,3 mmol).

10 MS-ESI: 292/294 ($M^+ + 1$, 69/23).

Elementary analysis: C 45.29% H 4.84% N 4.80%

Determined: C 45.31% H 4.86% N 4,78%

15 b) Trifluoro-methanesulfonate(5-chloro-2-methanesulfonyl-4-methoxycarbonyl-phenyl)-trimethyl-ammonium (**20b**)

To a stirred solution of 3,65 g (12,5 mmol) **20a** and 50 ml dichloromethane were added 20,5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2 days then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **20b** was obtained in 58% yield (3,31 g, 7,25 mmol).

25

MS-ESI: 307 (M^+ , 100),

Elementary analysis:

Calculated: C 34.25% H 3.76% F 12.50% N 3.07%

Determined: C 34.24% H 3.80% F 12.47% N 3.06%

30

c) Trifluoro-methanesulfonate(4-carboxy-5-chloro-2-methanesulfonyl-phenyl)-trimethylammonium (**20c**).

35 A solution of 3,16 g (6,92 mmol) **20b**, 60 ml dest. water and 60 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed

extensively with diethyl ether and dried by oil pump vacuum. The solid crude was obtained in 83% yield (2,53 g; 5.74 mmol) and crude compound **20c** was used for the next step without purification.

5 MS-ESI: 293 (M^+ , 48),

Elementary analysis:

Calculated: C 32.62% H 3.42% F 12.90% N 3.17%

Determined: C 32.64% H 3.44% F 12.89% N 3.16%

10 d) Synthesis of (2-Chloro-5-methanesulfonyl-4-trimethylammonium-benzoyl)-Val- β Ala-Phe-Gly-NH₂- triflate salt (**20d**).

To a stirred solution of 88,4 mg (0.2 mmol) **20c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-Phe-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 88,4 mg (0.2 mmol) **20c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound H-Val- β Ala-Phe-Gly-NH₂ and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophilized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent. The desired compound **20d** was obtained in 43% yield (23,2 mg, 280 mmol).

30 MS-ESI: 681/683 (M^+ , 100).

e) Synthesis of [¹⁸F]-(2-Chloro-4-fluoro-5-methanesulfonyl-benzoyl)-Val- β Ala-Phe-Gly-NH₂ (**20e**).

35

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500µl water) and MeCN (1.5ml) the fluorine containing water (322 MBq, 35µl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated
5 again. A solution of **20d** (2mg) in anhydrous DMSO (300µl) was added. After heating at 90°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3µ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient:
10 5-95% B in 7 mins. The F-18 labeled product **20e** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econosphere analytical HPLC.

Example 21

15 a) Synthesis of 2-Cyano-5-fluoro-benzoic acid methyl ester (**21a**).

To a stirred suspension of 16,0 g (96,9 mmol) 2-Cyano-5-fluoro-benzoic acid (Apollo) and 161 ml methanol were added 30.4 g (387,6 mmol) acetyl chloride drop wisely at 0°C. The reaction mixture was stirred over night, filtered and concentrated. The residue was diluted
20 with dichloromethane, washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The residue was purified by column chromatography (hexane : ethylacetate). The desired product **21a** was obtained in 86.0 % yield (14,9 g; 83.3 mmol)

25 MS-ESI: 180 (M⁺ +1,100),

Elementary analysis:

Calculated: C 60.34% H 3.38% F 10.60% N 7.82%

Determined: C 60.41% H 3,39% F 10.58% N 7,79%

30 b) Synthesis of 2-Cyano-5-dimethylamino-benzoic acid methyl ester (**21b**).

To a stirred solution of 4.03 g (22,5 mmol) **21a** and 60.0 ml dimethylsulphoxid were added 2.23 g (27.0 mmol) dimethylamine hydrochloride and 6.54 g (47.3 mmol) potassium carbonate. The reaction mixture was stirred for 15h at 65°C in an autoclave and was
35 reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted

with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **21b** was obtained in 89.0 % yield (4.09 g, 20.0 mmol).

5

MS-ESI: 205 ($M^+ + 1$, 100),

Elementary analysis:

Calculated: C 64.69% H 5.92% N 13.72%

10 Determined: C 64.75% H 5.94% N 13.68%

c) Trifluoro-methanesulfonate(4-cyano-3-methoxycarbonyl-phenyl)-trimethylammonium (21c).

15

To a stirred solution of 2.55 g (12.5 mmol) **21b** and 50 ml dichloromethane were added 20.5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2 days then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **21c** was obtained in 88% yield (4.05 g, 11.0 mmol).

20

25 MS-ESI: 219 (M^+ , 71),

Elementary analysis:

Calculated: C 42.39% H 4.10% F 15.47% N 7.61%

Determined: C 42.41% H 4.13% F 15.45% N 7.60%

30 d) Synthesis of Trifluoro-methanesulfonate(3-carboxy-4-cyano-phenyl)-trimethylammonium (21d).

A solution of 4.01 g (10.9 mmol) **21c**, 95 ml dest. water and 95 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude product **21d**

35

was obtained in 96% yield (3,70 g, 10,5 mmol) and crude compound **21d** was used for the next step without purification.

MS-ESI: 205 (M^+ , 76),

5 Elementary analysis:

Calculated: C 40.68% H 3.70% F 16.09% N 7.91%

Determined: C 40.70% H 3,72% F 16.07% N 7,90%

10 e) Synthesis of peptide(5-Trimethylammonium-2-cyano-benzoyl)-Val- β Ala-Arg-Gly-NH₂ – triflate salt (**21e**).

To a stirred solution of 70.8 mg (0.2 mmol) **21d** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-Arg(Pbf)-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard
15 protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) **21d** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol)
20 diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl
25 ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water: acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent. The desired compound **21e** was obtained in 38% yield (18,6 mg, 0.0247 mmol).

30

MS-ESI: 603 (M^+ , 100),

f) Synthesis of [¹⁸F]-(5-Fluoro-2-cyano-benzoyl)-Val- β Ala-Arg-Gly-NH₂ (**21f**).

35 To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (345 MBq, 35 μ l) was added.

The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **21e** (2mg) in anhydrous DMSO (300µl) was added. After heating at 90°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3µ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **21f** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econosphere analytical HPLC.

10

Example 22

a) Synthesis of 2-Chloro-4,5-difluoro-benzoic acid methyl ester (**22a**).

- 15 To a stirred suspension of 5,0 g (26 mmol) 2-Chloro-4,5-difluoro-benzoic acid (Apollo) and 50 ml methanol were added 7,41 ml (104 mmol) acetyl chloride drop wisely at 0°C. The reaction mixture was stirred over night, filtered and concentrated. The residue was diluted with dichloromethane, washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The residue was purified by column
- 20 chromatography (hexane : ethylacetate). The desired product **22a** was obtained in 84% yield (4,51 g, 21.8 mmol).

MS-ESI: 207 / 209 (M⁺ +1, 64 / 22).

Elementary analysis:

- 25 Calculated: C 46.51% H 2.44% F 18.39%
Determined: C 46.59% H 2.46% F 18.35%

b) 2-Chloro-4-dimethylamino-5-fluoro-benzoic acid methyl ester (**22b**).

- 30 To a stirred solution of 23.1 g (112 mmol) **22a** and 231 ml dimethylsulphoxid were added 10.0 g (123 mmol) dimethylamine hydrochloride and 32.4 g (234 mmol) potassium carbonate. The reaction mixture was stirred for 24h at 60°C in an autoclave and was reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted
- 35 with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The

oily crude was purified by column chromatography and the desired product **22b** was obtained in 89,5% yield (23.1 g 100 mmol).

MS-ESI: 232 / 234 (M^+ +1, 55 / 18).

5 Elementary analysis:

Calculated: C 51.85% H 4.79% F 8.20% N 6.05%

Determined: C 51.89% H 4,81% F 8,18% N 6,03%

10 c) Trifluoro-methanesulfonate(5-chloro-2-fluoro-4-methoxycarbonyl-phenyl)-trimethyl-ammonium (**22c**).

To a stirred solution of 7.06 g (30.5 mmol) **22b** and 100 ml dichloroethane were added 50 g (305 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was stirred for 24 hours at 90°C and then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **22c** was obtained in 91.1% yield (11.0 g; 27.8 mmol).

MS-ESI: 246 / 248 (M^+ , 100 / 32).

Calculated: C 36.42% H 3.57% F 19.20% N 3.54%

Determined: C 36,46% H 3.58% F 19.18% N 3.51%

25 d) Synthesis of Trifluoro-methanesulfonate(4-carboxy-5-chloro-2-fluoro-phenyl)-trimethyl-ammonium (**22d**).

A solution of 2.0 g (5.05 mmol) **22c**, 45 ml dest. water and 45 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude product **22d** was obtained and crude compound **22d** was used for the next step without purification. The desired crude compound **22d** was obtained in 76% yield (1,46 g, 3,84 mmol).

MS-ESI: 232 / 234 (M^+ , 68 / 21).

Calculated: C 34.61% H 3.17% F 19.91% N 3.67%

Determined: C 34,66% H 3.19% F 19.94% N 3.66%

e) Synthesis of peptide(4-Trimethylammonium-2-chloro-5-fluoro-benzoyl)-Val-βAla-Phe-Gly-NH₂- triflate salt (22e).

5

To a stirred solution of 76 mg (0.2 mmol) **22d** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid were added 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val-βAla-Phe-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 10 4h. The mixture was filtered and washed with dimethylformamide The coupling step was repeated. Thus, to a stirred solution of 76 mg (0.2 mmol) **22d** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound H-Val-βAla-Phe-Gly-NH₂ and the mixture was again shaken intensively for 15 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was 20 purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain **22e** in 37% yield (18.6 mg, 0,0241 mmol).

MS-ESI: 620/622 (M⁺, 100/34).

25 f) Synthesis of [¹⁸F]-(4-Trimethylammonium-2-chloro-5-fluoro-benzoyl)-Val-βAla-Phe-Gly-NH₂ (22f).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500μl water) and MeCN (1.5ml) the fluorine containing water (319 MBq, 33μl) was added.

30 The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **22e** (2mg) in anhydrous DMSO (300μl) was added. After heating at 90°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8μ, 2ml/min, solvent A: H₂O, solvent B: MeCN, 35 gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3μ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient:

5-95% B in 7 mins. The F-18 labeled product **22f** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

Example 23

5

Synthesis of (4-Trimethylammonium-3-cyano-benzoyl)-Gly-Tyr-βAla-Val-NH₂, (**23a**).

To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Gly-Tyr(OtBu)-βAla-Val-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was lyophilized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product **23a** in 40% yield (19,8 mg (0.0176 mmol)).

MS-ESI: 609 (M⁺, 67).

Synthesis of [¹⁸F]-4-Fluoro-3-cyano-benzoyl)-Gly-Tyr-βAla-Val-NH₂, (**23b**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500μl water) and MeCN (1.5ml) the fluorine containing water (356 MBq, 35μl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **23a** (2mg) in anhydrous DMSO (300μl) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8μ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3μ, 3ml/min

(Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **23b** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

5 Example 24

Synthesis of (4-Trimethylammonium-3-cyano-benzoyl)-Ava-His(π -Me)-Sta-Leu-NH₂-triflate salt, (**24a**).

- 10 To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Ava-His(π -Me)-Sta-Leu-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed
- 15 with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed
- 20 extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS
- 25 with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product **24a** in 38% yield (21.6 mg (0.023 mmol)).

MS-ESI: 727 (M⁺, 77).

- 30 Synthesis of [¹⁸F]-4-Fluoro-3-cyano-benzoyl)-Ava-His(π -Me)-Sta-Leu-NH₂ , (**24b**).

- To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (377 MBq, 35 μ l) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen.
- 35 Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **24a** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at

50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8μ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3μ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **24b** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econosphere analytical HPLC.

Example 25

10 Synthesis of (4-Trimethylammonium-3-cyano-benzoyl)-N-MeGly-His(π -Me)-Sta-Leu-NH₂-triflate salt, (25a).

To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-N-MeGly-His(Tr)-Sta-Leu-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophilized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product **25a** in 29% yield (16,0 mg (0.0189 mmol)).

30 MS-ESI: 698 (M⁺, 75).

Synthesis of [¹⁸F]-4-Fluoro-3-cyano-benzoyl)-N-MeGly-His(π -Me)-Sta-Leu-NH₂, (25b).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500μl water) and MeCN (1.5ml) the fluorine containing water (382 MBq, 35μl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen.

Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **25a** (2mg) in anhydrous DMSO (300µl) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3µ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **25b** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

10 Example 26

Synthesis of (4-Trimethylammonium-3-cyano-benzoyl)-Val-βAla-Arg-Gly-NH₂-triflate salt, (**26a**).

- 15 To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val-βAla-Arg(Pbf)-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed
- 20 with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed
- 25 extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS
- 30 with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product **26a** in 47% yield (20.3 mg, 0.0305 mmol).
(19,5 mg (0.026 mmol)).
MS-ESI: 603 (M⁺, 100).

- 35 Synthesis of [¹⁸F]-4-Fluoro-3-cyano-benzoyl)-Val-βAla-Arg-Gly-NH₂, (**26b**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500µl water) and MeCN (1.5ml) the fluorine containing water (344 MBq, 35µl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated
5 again. A solution of **26a** (2mg) in anhydrous DMSO (300µl) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3µ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient:
10 5-95% B in 7 mins. The F-18 labeled product **26b** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econosphere analytical HPLC.

Example 27

15 Synthesis of (4-Trimethylammonium-3-cyano-benzoyl)-Val-βAla-His(II-Me)-Gly-NH₂ triflate salt, (**27a**).

To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2
20 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val-βAla-His(II-Me)-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was
25 added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml
30 triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product **27a** in 33% yield (16,0 mg (0.0215 mmol)).
35 MS-ESI: 597 (M⁺, 100).

Synthesis of [^{18}F]-4-Fluoro-3-cyano-benzoyl)-Val- β Ala-His(π -Me)-Gly-NH₂, (27b).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (367 MBq, 35 μ l) was added.

- 5 The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of 27a (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 50°C for 15 min the crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min
10 (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product 27b was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC (Fig. 5; HPLC chromatogram of reaction mixture with co-injection of the cold standard).

15

Example 28

Synthesis of (4-Trimethylammonium-3-cyano-benzoyl)-Val- β Ala-His(π -Me)-Leu-NH₂ triflate salt, (28a).

20

- To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-His(π -Me)-Leu-NH₂ (loading 0.68 mmol/g) which was prepared by standard
25 protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the
30 mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the
35 supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS

with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product **28a** in 57% yield (29,8 mg (0.037 mmol)).

MS-ESI: 654 (M^+ , 100).

5 Synthesis of [^{18}F]-4-Fluoro-3-cyano-benzoyl)-Val- β Ala-His(π -Me)-Leu-NH₂, (**28b**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (356 MBq, 35 μ l) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen.

10 Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **28a** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min
15 (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **28b** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

Example 29

20

Synthesis of (4-Trimethylammonium-3-cyano-benzoyl)-Val- β Ala-Phe-Gly-NH₂ triflate salt, (**29a**).

To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml
25 dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-Phe-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8
30 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml
35 trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was

separated by centrifugation. Water was added to the solid and the supernatant was lyophilized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product **29a** in 46% yield (22,2 mg (0.0299 mmol)).

5 MS-ESI: 593 (M^+ , 100).

Synthesis of [^{18}F]- (4-Fluoro-3-cyano-benzoyl)-Val- β Ala-Phe-Gly-NH₂, (**29b**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 10 500 μ l water) and MeCN (1.5ml) the fluorine containing water (333 MBq, 35 μ l) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **29a** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC 15 (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **29b** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econosphere analytical HPLC.

20

Example 30

Synthesis of (4-Trimethylammonium-3-cyano-benzoyl)-Val- β Ala-Trp-Gly-NH₂-triflate salt, (30a).

25

To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-Trp(Boc)-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard 30 protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the 35 mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a

mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was lyophilized. The residue was purified by preparative RP-18 HPLC-MS
5 with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product **30a** in 41% yield (20,8 mg (0.021 mmol)).
MS-ESI: 632 (M^+ , 100).

Synthesis of [^{18}F]-4-Fluoro-3-cyano-benzoyl)-Val- β Ala-Trp-Gly-NH₂, (**30b**).

10 To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (368 MBq, 35 μ l) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated
15 again. A solution of **30a** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient:
20 5-95% B in 7 mins. The F-18 labeled product **30b** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

Example 31

25 Synthesis of (4-Trimethylammonium-3-cyano-benzoyl)-Val- β Ala-Tyr-Gly-NH₂ triflate salt, (**31a**).

To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2
30 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-Tyr(OtBu)-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was
35 added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the

mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was lyophilized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product **31a** in 36% yield (17,7 mg (0.0234 mmol)).

MS-ESI: 609 (M^+ , 100).

Synthesis of [^{18}F]-4-Fluoro-3-cyano-benzoyl)-Val- β Ala-Tyr-Gly-NH₂, (**31b**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (339 MBq, 35 μ l) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **31a** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **31b** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econosphere analytical HPLC.

Example 32

a) Synthesis of (4-Trimethylammonium-3-trifluormethyl-benzoyl)-Val- β Ala-His(π -Me)-Gly-NH₂- triflate salt (**32a**).

To a stirred solution of 79,4 mg (0.2 mmol) **12c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-His(π -Me)-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 79.4 mg (0.2 mmol) **12c** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was

added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was lyophilized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**32a**) in 59% yield (24,2 mg (0.0384 mmol)). MS-ESI: 598 (M^+ , 88)

b) Synthesis of [^{18}F]-4-Fluoro-3-cyano-benzoyl)-Val- β Ala-His(π -Me)-Gly-NH₂, (**32b**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (318 MBq, 35 μ l) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **32a** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **32b** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econosphere analytical HPLC.

Example 33

a) Synthesis of (4-Trimethylammonium-3-cyano-benzoyl)-Val- β Ala-Lys-Gly-NH₂– triflate salt (**33a**).

To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-Lys(Boc)-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed

with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product **33a** in 52% yield (24,5 mg (0.0338 mmol)). MS-ESI: 575 (M^+ , 100).

b) Synthesis of [^{18}F]-4-Fluoro-3-cyano-benzoyl)-Val- β Ala-Lys-Gly-NH₂, (**33b**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (301 MBq, 35 μ l) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **33a** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **33b** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

Example 34

30

a) Synthesis of (4-Trimethylammonium-3-cyano-benzoyl)-Val- β Ala-Met-Gly-NH₂— triflate salt (**34a**).

To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-

Val-βAla-Met-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed peptide resin and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product **34a** in 37% yield (17,4 mg (0.024 mmol)).

MS-ESI: 577 (M⁺, 100).

b) Synthesis of [¹⁸F]-4-Fluoro-3-cyano-benzoyl)-Val-βAla-Met-Gly-NH₂, (**34b**).

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500μl water) and MeCN (1.5ml) the fluorine containing water (383 MBq, 35μl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **34a** (2mg) in anhydrous DMSO (300μl) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8μ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18 , 53×7mm, 3μ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **34b** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

Example 35

a) Synthesis of (3-Cyano-4-fluoro-benzenesulfonyl)-methyl-amino]-acetic acid methyl ester (**35a**)

To a suspension of 6.90 g (0.05 mol) sarcosine hydrochloride and 19.5 g (0.15 mol) diisopropylethyl amine in 70.0 ml dichloromethane were added 11.5 g (0.055 mol) 3-cyano-4-fluoro-benzenesulfonyl chloride (Aldrich) in 50.0 ml dichloromethane dropwisely at 0°C. The suspension was stirred for 4h. The suspension was poured on 150 ml stirred
5 ice/water mixture. The water phase was separated and extracted twice with dichloromethane. The combined dichloromethane phases were washed twice with diluted hydrogen chloride solution and subsequently with sodium hydrogen carbonate solution, dried with sodium sulphate and concentrated. The oily crude product **35a** was purified by column chromatography (hexane : ethylacetate). The desired product **35a** was obtained in
10 58,2% yield (8.32 g ; 29.1 mmol)

MS-ESI: 287 ($M^+ + 1$, 100).

Elementary analysis: C 46.15% H 3.87% F 6.64% N 9.79%

Determined: C 46.16% H 3.88% F 6.65% N 9.80%

15

b) Synthesis of (3-Cyano-4-dimethylamino-benzenesulfonyl)-methyl-amino]-acetic acid methyl ester (**35b**)

To a stirred solution of 5.72 g (20.0 mmol) **35a** and 60 ml dimethylsulphoxid were added
20 2.03 g (24,7 mmol) dimethylamine hydrochloride and 5,97 g (43.2 mmol) potassium carbonate. The reaction mixture was stirred over night and was reduced with high vacuum rotation evaporator at 60°C. The residue was diluted with dichloromethane, washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with diluted sodium hydrogen carbonate
25 solution, dried with sodium sulphate and concentrated. The oily crude product **35b** was obtained in 95% yield (5.92 g, 19.9 mmol) and was used for the next step without purification.

MS-ESI: 312 ($M^+ + 1$, 100).

30 Elementary analysis: C 50.15% H 5.50% N 13.50%

Determined: C 50.18% H 5.52% N 13.48%

c) Synthesis of Trifluoro-methanesulfonate[2-cyano-4-(methoxycarbonylmethyl-methyl-sulfamoyl)-phenyl]-trimethyl-ammonium (**35c**)

35

To a stirred solution of 3.89 g (12.5 mmol) **35b** and 50 ml dichloromethane were added 20.5 g (125 mmol) methyltriflate (Aldrich) drop wisely. The reaction mixture was refluxed for 2 days then cooled to room temperature. Diethylether was added. The desired compound precipitates and the solvent was decanted. The solid was washed extensively (ten times) with large amounts of diethylether. The solid was dried by use of oil pump vacuum and purified by (C-18) RP-column chromatography (acetonitril/water - gradient 1:99 to 80:20). The desired compound **35c** was obtained in 43 % yield (2,55 g, 5,375 mmol).

MS-ESI: 326 (M^+ , 100).

Elementary analysis: C 37.89% H 4.24% F 11.99% N 8.84%

Determined: C 37.92% H 4.26% F 11.96% N 8.86%

d) Trifluoro-methanesulfonate[4-(carboxymethyl-methyl-sulfamoyl)-2-cyano-phenyl]-trimethyl-ammonium (35d)

A solution of 2,38g (5,0 mmol) **35c**, 50 ml dest. water and 50 ml trifluoroacetic acid was refluxed for 2 days. The reaction mixture was evaporated, dried by use of oil pump vacuum over night and treated with diethyl ether. The resulting solid was filtered, washed extensively with diethyl ether and dried by oil pump vacuum. The solid crude was obtained in 79% yield (1.82 g; 3,95 mmol) and crude compound **35d** was used for the next step without purification.

MS-ESI: 312 (M^+ , 100).

Elementary analysis: C 36.44% H 3.93% F 12.35% N 9.11%

Determined: C 36.47% H 3.95% F 12.33% N 9.10%

e) Synthesis of (4-Trimethylammonium-3-cyano-benzenesulfonyl)-Gly-Val- β Ala-His(π -Me)-Gly-NH₂- triflate salt (35e)

30

To a stirred solution of 82,4 mg (0.2 mmol) **35d** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.065 mmol Rink-resin-bound H-Val- β Ala-His(π -Me)-Gly-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 82

35

mg (0.2 mmol) **35d** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water : acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent to obtain the desired product (**35e**) in 55% yield – 30.5 mg (0.036 mmol).

MS-ESI: 705 (M^+ , 100).

f) Synthesis of [^{18}F]- (4-Fluoro-3-cyano-benzenesulfonyl)-Val- β Ala-His(π -Me)-Gly-NH₂-triflate salt (**35f**)

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (344 MBq, 33 μ l) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **35e** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **35f** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econosphere analytical HPLC.

Example 36

a) Synthesis of N-(4-Trimethylammonium-3-cyano-benzoyl)-6-fluoro-dopamin - triflate salt (**36a**)

To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2

mmol) diisopropylcarbodiimid. The solution was stirred for 20 min and 34 mg (0.2 mmol) symphatomimetic 6-fluorodopamin (J. Fluorine Chem.; 74; 1; 1995; 113-122, CAS Nr. 71144-39-3) was added. The reaction mixture was stirred intensively for 8 h. The reaction mixture was evaporated in vacuum, diluted with dichloromethane : iso-propanol mixture (10:1) and washed twice with water. The combined water phases were extracted with dichloromethane. The combined organic phases were washed with brine, dried with sodium sulphate and concentrated. The oily crude was purified by RP column chromatography (water : acetonitril gradient) and the desired product **36a** was obtained in 44 % yield (45 mg, 0.088 mmol).

MS-ESI: 358 (M^+ , 100).

Elementary analysis: C 47.34% H 4.17% F 14.98% N 8.28%

Determined: C 47.36% H 4.19% F 14.97% N 8.27%

b) Synthesis of [^{18}F]-N-(4-Fluoro-3-cyano-benzoyl)-6-fluoro-dopamin - triflate salt (**36b**)

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μl water) and MeCN (1.5ml) the fluorine containing water (356 MBq, 35 μl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen.

Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **36a** (2mg) in anhydrous DMSO (300 μl) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min

(Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **36b** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

Example 37

a) Synthesis of N-(4-Trimethylammonium-3-cyano-benzoyl)-didemethyltamoxifen- triflate salt (**37a**)

To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was stirred for 20 min and 0.2 mmol estrogen

antagonist didemethyltamoxifen (J. Pharm. Sci.; 82; 9; (1993); 927-933, CAS Nr. 80234-20-4) was added. The reaction mixture was stirred intensively for 8 h. The reaction mixture was evaporated in vacuum, diluted with dichloromethane and washed twice with water. The combined water phases were extracted with dichloromethane. The combined
5 dichloromethane phases were washed with brine, dried with sodium sulphate and concentrated. The oily crude was purified by column chromatography and the desired product **37b** was obtained in 56 % yield (76 mg, 0.112 mmol)

MS-ESI: 531 (M^+ , 100).

10 Elementary analysis: C 63.61% H 5.34% F 8.38% N 6.18%
Determined: C 63.64% H 5.35% F 8.37% N 6.17%

b) Synthesis of [^{18}F]-N-(4-Fluoro-3-cyano-benzoyl) didemethyltamoxifen (**37b**)

15 To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (337 MBq, 35 μ l) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **37a** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at
20 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **37b** was confirmed by co-injection with the
25 non-radioactive F-19 fluoro standard on the Econosphere analytical HPLC.

Example 38

a) Synthesis of N-(4-Trimethylammonium-3-cyano-benzoyl)-alaphen - triflate salt (**38a**)

30

To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was stirred for 20 min and 49 mg (0.2 mmol) alaphen (Pharm.Chem.J.(Engl.Transl.); 9; 3; (1975); p. 158; CAS Nr. 15269-42-8) was
35 added. The reaction mixture was stirred intensively for 8 h. The reaction mixture was evaporated in vacuum, diluted with dichloromethane:isopropanol mixture (10:1) and

washed twice with water. The combined water phases were extracted with dichloromethane. The combined dichloromethane phases were washed with brine, dried with sodium sulphate and concentrated. The oily crude was purified by RP column chromatography and the desired product **38a** was obtained in 64 % yield (77 mg, 0.13 mmol).

MS-ESI: 394 (M^+ , 100)

Elementary analysis: C 53.38% H 5.84% F 11.01% N 8.12%

Determined: C 53.41% H 5.85% F 11.00% N 8.11%

b) Synthesis of [^{18}F]-N-(4-Fluoro-3-cyano-benzoyl) alaphen (**38b**)

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500 μ l water) and MeCN (1.5ml) the fluorine containing water (364 MBq, 35 μ l) was added.

The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen.

Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of **38a** (2mg) in anhydrous DMSO (300 μ l) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC

(Column Zorbax SB C18, 50 $\times$ 4.6mm, 1.8 μ , 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53 $\times$ 7mm, 3 μ , 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product **38b** was confirmed by co-injection with the non-radioactive F-19 fluoro standard on the Econsphere analytical HPLC.

Example 39


a) Synthesis of 3-Cyano-4-(trimethylammonium)-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His-(4-amino-5-methyl-heptanoic acid)-Leu-NH₂ trifluoroacetic acid salt (**39a**)

To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.025 mmol Rink-resin-bound H-Ava-Gln-Trp-Ala-Val-Gly-His-(4-amino-5-methyl-heptanoic acid)-Leu-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) **2e** in 1.5 ml dichloromethane

and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water: acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent. The desired compound 39 was obtained in 35% yield (12,2 mg, 0.012 mmol).

MS-ESI: 1237 (M^+ , 100),

b) "Synthesis of 3-cyano-4-[18F]fluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His-(4-amino-5-methyl-heptanoic acid)-Leu-NH₂ (39b)



18F-fluoride (2475MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and cesium carbonate (2.3mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. A solution of 3-cyano-4-trimethylammoniumbenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His-(4-amino-5-methyl-heptanoic acid)-Leu-NH₂ trifluoroacetate salt (2mg) (which was synthesized by standard solid phase Fmoc-peptide methods described and cited, *e.g.*, in the book: Chan and White - "Fmoc Solid Phase Peptide Synthesis - A Practical Approach") in anhydrous DMSO (150μl) was added. After heating at 70°C for 15 min. The reaction mixture was transferred to a vial containing water (4ml). The reaction vial washed with 150 μl DMSO and this was also transferred to the vial containing water. This solution was transferred to a semi-prep HPLC (column: ACE 5μ C18, 250x10 mm, solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 30%B for 5 min at 2 ml/min, then 30-70% B in 10 mins at 3 ml/min) and the desired F18 product peak was collected (253MBq).

20.4% d.c.). The F-18 labeled product was confirmed by co-injection with the F-19 cold standard on an analytical HPLC (Column ACE 3 μ C18 50 $\times$ 4.6mm, 1ml/min (Agilent), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 0% for 2min, then 0%B to 95%B in 7 mins).

5

Example 40

a) Synthesis of 3-Cyano-4-(trimethylammonium)-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His($\square$ Me)-Sta-Leu-NH₂ trifluoroacetic acid salt (40a)

10

To a stirred solution of 70.8 mg (0.2 mmol) 2e in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.025 mmol Rink-resin-bound H-Arg-Ava-Gln-Trp-Ala-Val-Gly-His($\square$ Me)-Sta-Leu-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) 2e in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water: acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent. The desired compound 40a was obtained in 28% yield (10,8 mg, 0.007 mmol).

15


20

25

MS-ESI: 1423 (M⁺, 100),

30

b) Synthesis of 3-Cyano-4-[¹⁸F]fluorobenzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His($\square$ Me)-Sta-Leu-NH₂ (40b)



¹⁸F-fluoride (2475MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and cesium carbonate (2.3mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. A solution of 3-cyano-4-trimethylammoniumbenzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂ trifluoroacetate salt (2mg) (which was synthesized by standard solid phase Fmoc-peptide methods described and cited, *e.g.*, in the book: Chan and White - "Fmoc Solid Phase Peptide Synthesis - A Practical Approach") in anhydrous DMSO (150μl) was added. After heating at 70°C for 15 min. The reaction mixture was transferred to a vial containing water (4ml). The reaction vial washed with 150μl DMSO and this was also transferred to the vial containing water. This solution was transferred to a semi-prep HPLC (column: ACE 5μ C18, 250x10 mm, solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 30%B for 5min at 2ml/min, then 30-70% B in 10 mins at 3ml/min) and the desired F18 product peak was collected (144MBq, 11.5% d.c.). The F-18 labeled product was confirmed by co-injection with the F-19 cold standard on an analytical HPLC (Column ACE 3μ C18 50x4.6mm, 1ml/min (Agilent), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 0% for 2min, then 0%B to 95%B in 7 mins).

Example 41


a) Synthesis of 3-Cyano-4-(trimethylammonium)-benzoyl-Arg-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂ trifluoroacetic acid salt (41a)

To a stirred solution of 70.8 mg (0.2 mmol) 2e in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.025 mmol Rink-resin-bound H-Arg-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus,

to a stirred solution of 70.8 mg (0.2 mmol) 2e in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS with a water: acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent. The desired compound 41a was obtained in 24% yield (9,0 mg, 0.006 mmol).

MS-ESI: 1394 (M^+ , 100)

b) Synthesis of 3-cyano-4-[^{18}F]fluorobenzoyl-Arg- β Ala-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂ (41b)



^{18}F -fluoride (1419MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and potassium carbonate (1mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. A solution of 3-cyano-4-trimethylammoniumbenzoyl-Arg- β Ala-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂ trifluoroacetate salt (2mg) (which was synthesized by standard solid phase Fmoc-peptide methods described and cited, e.g., in the book: Chan and White - "Fmoc Solid Phase Peptide Synthesis - A Practical Approach") in anhydrous DMSO (150 μ l) was added. After heating at 50°C for 15 min. The reaction mixture was diluted with water (4ml) and transferred to a semi-prep HPLC (column: Zorbax Bonus 5 μ C18, 250x9.2 mm, solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 29-34% B in 20 mins at 3ml/min) and the desired F18 product peak was collected (150MBq, 21.1% d.c.). The F-18 labeled product was confirmed by co-injection with the F-19 cold standard

on an analytical HPLC (Column Zorbax Bonus 5 μ C18 250 $\times$ 4.6mm, 1ml/min (Agilent), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 29%B to 34%B in 20 mins).


5 Example 42

a) Synthesis of 3-Cyano-4-(trimethylammonium)-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂ trifluoroacetic acid salt (42a)

- 10 To a stirred solution of 70.8 mg (0.2 mmol) 2e in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.025 mmol Rink-resin-bound H-Arg- β Ala-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture
- 15 was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 70.8 mg (0.2 mmol) 2e in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and
- 20 washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was liophylized. The residue was purified by preparative RP-18 HPLC-MS
- 25 with a water: acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent. The desired compound 42a was obtained in 32% yield (11,1 mg, 0.008 mmol).

MS-ESI: 1267 (M⁺, 100)

- 30 b) Synthesis of 3-cyano-4-[18F]fluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂ (42b)



18F-fluoride (869MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and potassium carbonate (1mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. A solution of 3-cyano-4-trimethylammoniumbenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu -NH₂ trifluoroacetate salt (2mg) (which was synthesized by standard solid phase Fmoc-peptide methods described and cited, e.g., in the book: Chan and White - "Fmoc Solid Phase Peptide Synthesis - A Practical Approach") in anhydrous DMSO (150μl) was added. After heating at 50°C for 15 min. The reaction mixture was diluted with water (4ml) and transferred to a semi-prep HPLC (column: Zorbax Bonus 5μ C18, 250x9.2 mm, solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 34-38% B in 20 mins at 3ml/min) and the desired F18 product peak was collected (184MBq, 37.8% d.c.). The F-18 labeled product was confirmed by co-injection with the F-19 cold standard on an analytical HPLC (Column Zorbax Bonus 5μ C18 250x4.6mm, 1ml/min (Agilent), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 34%B to 38%B in 20 mins).

Example 43


a) Synthesis of 3-Trifluormethyl-4-(trimethylammonium)-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂ (43a)

To a stirred solution of 79,4 mg (0.2 mmol) 12c in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2 mmol) diisopropylcarbodiimid. The solution was added to 0.025 mmol Rink-resin-bound H-Arg-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂ (loading 0.68 mmol/g) which was prepared by standard protocol. The mixture was shaken intensively for 4h. The mixture was filtered and washed with dimethylformamide. The coupling step was repeated. Thus, to a stirred solution of 79,4 mg (0.2 mmol) 12c in 1.5 ml dichloromethane and 0.25 ml dimethylformamid was added 65 mg (0.5 mmol) diisopropylethylamin and 0.031 ml (0.2

mmol) diisopropylcarbodiimid. The solution was added to the washed Rink-resin-bound peptide and the mixture was again shaken intensively for 4h. The mixture was filtered and washed extensively with dimethylformamide and dichloromethane. The resin was treated with a mixture of 0.85 ml trifluoroacetic acid, 0.05 ml distilled water, 0.05 ml phenol, 0.05 ml triisopropylsilane for 3h. The mixture was added in ca. 9 ml ice cold methyl tert-butyl ether. The solid was separated by centrifugation. Water was added to the solid and the supernatant was lyophilized. The residue was purified by preparative RP-18 HPLC-MS with a water: acetonitril gradient and 0.1% trifluoro acetic acid as co-solvent. The desired compound 43a was obtained in 29% yield (11,5 mg, 0.0072 mmol).

MS-ESI: 1480 (M^+ , 100)

b) Synthesis of 3-trifluoromethyl-4-[^{18}F]fluorobenzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂ (43b)



^{18}F -fluoride (835MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and cesium carbonate (2.3mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. A solution of 3-trifluoromethyl-4-trimethylammoniumbenzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂ trifluoroacetate salt (2mg) (which was synthesized by standard solid phase Fmoc-peptide methods described and cited, e.g., in the book: Chan and White - "Fmoc Solid Phase Peptide Synthesis - A Practical Approach") in anhydrous DMSO (150μl) was added. After heating at 70°C for 15 min the reaction mixture was diluted with water (4ml) and transferred to a semi-prep HPLC (column: ACE 5μ C18 250x10 mm, solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 20% for 2min, then 20-60% B in 20 mins at 3ml/min) and the desired F18 product peak was collected (78MBq, 29.0% d.c.). The F-18 labeled product was confirmed by co-injection with the F-19 cold standard on an analytical HPLC (Column ACE C18,

50×4.6mm, 3μ, 2ml/min, solvent A: 10mM K₂HPO₄ in H₂O, solvent B: 10mM K₂HPO₄ in MeCN/H₂O (7:3), gradient: 5%-95% B in 7 mins).

Example 44

5

a) Synthesis of Trifluoro-methanesulfonate[4-(1-benzyl-2,3-dihydro-1H-pyrrolo[2,3-b]quinolin-4-ylcarbamoyl)-2-trifluoromethyl-phenyl]-trimethyl-ammonium (44a)

To a suspension 30 mg (1.3 mmol) sodium hydride in 2 ml dry THF 275 mg (1 mmol) 1-Benzyl-2,3-dihydro-1H-pyrrolo[2,3-b]quinolin-4-ylamine (J. Med. Chem. 2004, 47, 1413) in 1 ml dry THF were added drop wisely. The solution was stirred for 30 min at room temperature. A solution of 794 mg (2 mmol) 12c, 650 mg (5 mmol) diisopropylethylamin and 0.31 ml (2.0 mmol) diisopropylcarbodiimid which was stirred in 8 ml DMF was added. The reaction mixture was stirred for 4 hours at room temperature and reduced with high vacuum rotation evaporator at 65°C. The residue was diluted with diethyl ether and filtered. The filter cake solid was purified by RP column chromatography (MeCN:water) to obtain 44a in 25% yield (163 mg, 0.25 mmol) after lyophilisation.

MS-ESI: 505 (M⁺, 100)

20 Elementary analysis:

Calculated: C 55.04% H 4.31% N 8.56%

Determined: C 55.02% H 4.32% N 8.55%

b) Synthesis of [F-19]-N-(1-Benzyl-2,3-dihydro-1H-pyrrolo[2,3-b]quinolin-4-yl)-4-fluoro-3-trifluoromethyl-benzamide (44b)

25

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500μl water) and MeCN (1.5ml) the fluorine containing water (364 MBq, 35μl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of 44a (2mg) in anhydrous DMSO (300μl) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8μ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3μ, 3ml/min (Alltech), solvent A: H₂O + 0.1% TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product 44b was confirmed by co-injection with the

non-radioactive F-19 fluoro standard (J. Med. Chem. 2004, 47, 1413-1422) on the Econsphere analytical HPLC.

Example 45

5

a) Synthesis of trimethyl-(2-nitro-4-{2-[(2-pyridin-2-yl-quinoline-4-carbonyl)-amino]-ethylcarbamoyl}-phenyl) trifluoro-methanesulfonate ammonium salt (45a)

10 To a solution of 328 mg (1 mmol) 2-[(2-Pyridin-2-yl-quinoline-4-carbonyl)-amino]-ethyl-ammonium chloride (Tetrahedron, (2004), 8729-8738) and 325 mg (2.5 mmol) diisopropylethylamine was added a solution of 748 mg (2 mmol) 12c, 650 mg (5 mmol) diisopropylethylamin and 0.31 ml (2.0 mmol) diisopropylcarbodiimid which was stirred in 8 ml DMF 20 min before. The reaction mixture was stirred for 20 hours and was concentrated in vacuum. The residue was treated with diethyl ether – the supernatant was
15 decanted and the solid was solved in water-acetonitrile. The product was purified by RP-column chromatography. The desired product 45a was obtained in 31% yield (201 mg, 0.31 mmol).

MS-ESI: 500 (M^+ , 100)

20 Elementary analysis:

Calculated: C 51.85% H 4.20% N 12.96%

Determined: C 51.86% H 4.19% N 12.95%

25 b) Synthesis of [F-18]-2-Pyridin-2-yl-quinoline-4-carboxylic acid [2-(4-fluoro-3-nitro-benzoylamino)-ethyl]-amide (45b)

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500µl water) and MeCN (1.5ml) the fluorine containing water (387 MBq, 35µl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen.
30 Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of 45a (2mg) in anhydrous DMSO (300µl) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econsphere C18, 53×7mm, 3µ, 3ml/min
35 (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product 45b was confirmed by co-injection with the

non-radioactive F-19 fluoro standard (Tetrahedron 60 (2004) 8729–8738) on the Econsphere analytical HPLC.

Example 46

5

a) Synthesis of Trifluoro-methanesulfonate{2-fluoro-4-[4'-((S)-1-methoxycarbonyl-2-methyl-propylsulfamoyl)-biphenyl-4-ylcarbamoyl]-phenyl}-trimethyl-ammonium (46a)

To a solution of 362 mg (1 mmol) (S)-2-(4'-Amino-biphenyl-4-sulfonylamino)-3-methyl-
10 butyric acid methyl ester (WO2007/16538A2) and 325 mg (2.5 mmol) diisopropylethylamine was added a solution of 694 mg (2 mmol) 3c, 650 mg (5 mmol) diisopropylethylamin and 0.31 ml (2.0 mmol) diisopropylcarbodiimid which was stirred for 20 min in 8 ml DMF before. The reaction mixture was stirred for 20 hours and was concentrated in vacuum. The residue was treated with diethyl ether; the supernatant was
15 decanted and the solid crude product was solved in water-acetonitrile. The product was purified by RP-column chromatography. The desired product 46a was obtained in 38% yield (263 mg, 0.38 mmol).

MS-ESI: 543 (M⁺, 100)

20 Elementary analysis:

Calculated: C 50.36% H 4.81% N 6.07%

Determined: C 50.38% H 4.80% N 6.07%

b) Synthesis of [F-18]-(S)-2-[4'-(3,4-Difluoro-benzoylamino)-biphenyl-4-sulfonylamino]-3-
25 methyl-butyric acid methyl ester (46b)

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500µl water) and MeCN (1.5ml) the fluorine containing water (321 MBq, 35µl) was added. The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen.
30 Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of 46a (2mg) in anhydrous DMSO (300µl) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H2O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53×7mm, 3µ, 3ml/min
35 (Alltech), solvent A: H2O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins. The F-18 labeled product 46b was confirmed by co-injection with the

non-radioactive F-19 fluoro standard (Tetrahedron 60 (2004) 8729–8738) on the Econsphere analytical HPLC.

Example 47

5

Synthesis of [F-18](4-fluoro-3-cyano-benzoyl)-TTA1 (47)


To a stirred solution of 70.8 mg (0.2 mmol) **2e** in 0.25 ml acetonitril was added 33 mg (0.25 mmol) diisopropylethylamin and 66 mg (0.2 mmol) 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylpiperidinium tetrafluoroborate (J. Am. Chem. Soc. 2005, 127, 48, 16912-16920). The reaction was stirred for 40 min. Five microliters of the reaction solution were added without further purification to 1.2 mg (100 nmol) TTA1 (Nucleic Acids Research, 2004, Vol. 32, No. 19, 5757–5765) dissolved in 20 µl buffer (pH ~7). A solution of citric acid was added (pH~6). After incubation for 1 hr at 37°C, the product was purified by spin-
10 filtration using a spin filter with a 10 kDa cut-off membrane (Microcon® MY-10, Amicon bioseparations). The residue on the filter was washed three times with acidified water (pH ~ 6 - citric acid). The purity was determined by HPLC analyses.

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500µl water) and MeCN (1.5ml) the fluorine containing water (316MBq, 33µl) were added.
20 The solvent was removed by heating at 120°C for 10 mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again.

The above mentioned TTA1-trimethylammonium-solution was added. After heating at 80°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC
25 (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 0.1%-25% B in 7 mins or Column Econosphere C18, 53×7mm, 3µ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 1/1 + 0.1% TFA, gradient: 0.1-25% B in 7 mins. The F-18 labeled product was approached by co-injection with the cold F-19 fluoro standard on the Econsphere analytical HPLC.


30

Example A1. Radiosynthesis of Methyl 3-Cyano-4-[¹⁸F]fluorobenzoate




¹⁸F-fluoride (63MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and potassium carbonate (1mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. After drying, a 0.04M solution of methyl 3-cyano-4-trimethylammoniumbenzoate triflate salt in 300µl DMSO was added. The reaction vessel was sealed and heated at 50°C for 15mins to effect labeling. The crude reaction mixture was analyzed using an analytical HPLC (Column ACE C18, 50×4.6mm, 3µ, 2ml/min, solvent A: 10mM K₂HPO₄ in H₂O, solvent B: 10mM K₂HPO₄ in MeCN/H₂O (7:3), gradient: 5%-95% B in 7 mins), the incorporation yield was 93.5%. The F-18 labeled product was confirmed by co-injection with the F-19 cold standard on an analytical HPLC (Column Altech Econsphere C18 RP, 53×7mm, 3µ, 3ml/min (Agilent), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5%B to 95%B in 7 mins).

Example B1. Radiosynthesis of Methyl 3-Trifluoromethyl-4-[¹⁸F]fluorobenzoate




¹⁸F-fluoride (73MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and potassium carbonate (1mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. After drying, a 0.04M solution of methyl 3-trifluoromethyl-4-trimethylammoniumbenzoate triflate salt in 300µl DMSO was added. The reaction vessel was sealed and heated at 50°C for 15mins to effect labeling. The crude reaction mixture was analyzed using an analytical HPLC (Column ACE C18, 50×4.6mm, 3µ, 2ml/min, solvent A: 10mM K₂HPO₄ in H₂O, solvent B: 10mM K₂HPO₄ in MeCN/H₂O (7:3), gradient: 5%-95% B in 7 mins), the incorporation yield was 86.6%. The F-18 labeled product was confirmed by co-injection with the F-19 cold standard on an analytical HPLC (Column Zorbax SB, 50×4.6mm, 1.8µ,

3ml/min (Agilent), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5%B to 95%B in 7 mins).


Example C1. Radiosynthesis of Methyl 2-Chloro-4-[¹⁸F]fluorobenzoate

- 5 ¹⁸F-fluoride (307MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and potassium carbonate (1mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. After drying, a solution of 1mg methyl 2-chloro-4-trimethylammoniumbenzoate triflate salt in 100μl DMSO was added. The reaction vessel was sealed and heated at 50°C for 15mins
- 10 to effect labeling. The crude reaction mixture was analyzed using an analytical HPLC (Column ACE C18, 50×4.6mm, 3μ, 2ml/min, solvent A: 10mM K₂HPO₄ in H₂O, solvent B: 10mM K₂HPO₄ in MeCN/H₂O (7:3), gradient: 5%-95% B in 7 mins), the incorporation yield was 70.0%. The F-18 labeled product was confirmed by co-injection with the F-19 cold standard on an analytical HPLC (Column Zorbax SB, 250×4.6mm, 5μ, 1ml/min (Agilent),
- 15 solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5%B to 95%B in 7 mins).

Figure 9: HPLC chromatogram of reaction mixture with co-injection of the cold standard

Example D1. Radiosynthesis of Methyl 2-Fluoro-4-[¹⁸F]fluorobenzoate


- 5 ¹⁸F-fluoride (839MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and cesium carbonate (2.3mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. After drying, a solution of 1mg methyl 2-fluoro-4-trimethylammoniumbenzoate triflate salt in 100μl DMSO was added. The reaction vessel was sealed and heated at 70°C for 15mins
- 10 to effect labeling. The crude reaction mixture was analyzed using an analytical HPLC (Column ACE C18, 50×4.6mm, 3μ, 2ml/min, solvent A: 10mM K₂HPO₄ in H₂O, solvent B: 10mM K₂HPO₄ in MeCN/H₂O (7:3), gradient: 5%-95% B in 7 mins), the incorporation yield was 86.1%. The F-18 labeled product was confirmed by co-injection with the F-19 cold standard on an analytical HPLC (Column Zorbax SB, 250×4.6mm, 5μ, 1ml/min (Agilent),
- 15 solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5%B to 95%B in 7 mins).

Example E. Radiosynthesis of Methyl 3-Fluoro-4-[¹⁸F]fluorobenzoate

20

- ¹⁸F-fluoride (751MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and potassium carbonate (1mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. After
- 25 drying, a solution of 1mg methyl 3-fluoro-4-trimethylammoniumbenzoate triflate salt in 100μl DMSO was added. The reaction vessel was sealed and heated at 50°C for 15mins to effect labeling. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8μ, 2ml/min, solvent A: H₂O, solvent B: water, gradient: 5%-95% B in 7 mins), the incorporation yield was 85.4%.

Example F1. Radiosynthesis of Methyl (3-Cyano-4-[¹⁸F]fluoro-benzenesulfonylamino)-acetate




5

¹⁸F-fluoride (123MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and cesium carbonate (2.3mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. After drying, a solution of 1mg methyl (3-cyano-4-trimethylammoniumbenzenesulfonyl-amino)-acetate triflate salt in 300μl DMSO was added. The reaction vessel was sealed and heated at 70°C for 15mins to effect labeling. The crude reaction mixture was analyzed using an analytical HPLC (Column ACE C18, 50×4.6mm, 3μ, 2ml/min, solvent A: 10mM K₂HPO₄ in H₂O, solvent B: 10mM K₂HPO₄ in MeCN/H₂O (7:3), gradient: 5%-95% B in 7 mins), the incorporation yield was 77.6%.

15

Example G1. Radiosynthesis of 3-cyano-4-[¹⁸F]fluorobenzoyl-D-Ala-D-Phe-NH₂




20

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500μl water) and MeCN (1.5ml) the fluorine containing water (1160MBq, 80μl) were added. The solvent was removed by heating at 120°C for 10mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of 3-cyano-4-trimethylammoniumbenzoyl-D-Ala- D-Phe-NH₂ trifluoroacetate salt (2mg) in anhydrous DMSO (300μl) was added. After heating at 50°C for 15 min. The reaction was cooled to room temperature and dilute with water (2.7ml). The crude reaction mixture was analyzed using an analytical HPLC (Column ACE C18, 50×4.6mm, 3μ, 2ml/min, solvent A: 10mM K₂HPO₄ in H₂O, solvent B: 10mM K₂HPO₄ in

25

MeCN/H₂O (7:3), gradient: 5%-95% B in 7 mins). The product was obtained by preparative radio HPLC to give 301MBq (51% d.c.) [column: Phenomenex Luna C18, 250x10mm, 5μ, solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5%B for 5min at 2ml/min, 5%B for 1min at 3ml/min, then 5-60% B in 19 mins at 3ml/min].

Example H1. Radiosynthesis of 3-cyano-4-[¹⁸F]fluorobenzoyl-Val-βAla-Phe-Gly-NH₂




10

To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500μl water) and MeCN (1.5ml) the fluorine containing water (454MBq, 50μl) were added. The solvent was removed by heating at 120°C for 10mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of 3-cyano-4-trimethylammoniumbenzoyl-Val-βAla-Phe-Gly-NH₂ trifluoroacetate salt (2mg) in anhydrous DMSO (300μl) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical HPLC (Column Zorbax SB C18, 50x4.6mm, 1.8μ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18, 53x7mm, 3μ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins). The F-18 labeled product was confirmed by co-injection with the F-19 cold standard on the Econsphere analytical HPLC.

20


Example I1. Radiosynthesis of 3-cyano-4-[¹⁸F]fluorobenzoyl-Val-βAla-Arg-Gly-NH₂

25




To a Wheaton vial (5ml) charged with Kryptofix 222 (5mg), potassium carbonate (1mg in 500µl water) and MeCN (1.5ml) the fluorine containing water (316MBq, 33µl) were added.

- 5 The solvent was removed by heating at 120°C for 10mins under a stream of nitrogen. Anhydrous MeCN (1ml) was added and evaporated as before. This step was repeated again. A solution of 3-cyano-4-trimethylammoniumbenzoyl-Val-□Ala-Arg-Gly-NH₂ trifluoroacetate salt (ZK6005341, 2mg) in anhydrous DMSO (300µl) was added. After heating at 50°C for 15 min. The crude reaction mixture was analyzed using an analytical
- 10 HPLC (Column Zorbax SB C18, 50×4.6mm, 1.8µ, 2ml/min, solvent A: H₂O, solvent B: MeCN, gradient: 5%-95% B in 7 mins or Column Econosphere C18 , 53×7mm, 3µ, 3ml/min (Alltech), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 5-95% B in 7 mins.

Example J1. Radiosynthesis of 3-cyano-4-[¹⁸F]fluorobenzoyl-Val-βAla-His(Me)-Gly-NH₂

- 5 ¹⁸F-fluoride (123MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and cesium carbonate (2.3mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. After drying, a solution of 3-cyano-4-trimethylammoniumbenzoyl-Val-□Ala-Arg-Gly-NH₂ trifluoroacetate salt (ZK6012623, 2mg) in anhydrous DMSO (300μl) was added. After
- 10 heating at 70°C for 5 min. The crude reaction mixture was analyzed using an analytical HPLC (Column ACE C18, 50×4.6mm, 3μ, 2ml/min, solvent A: 10mM K₂HPO₄ in H₂O, solvent B: 10mM K₂HPO₄ in MeCN/H₂O (7:3), gradient: 5%-95% B in 7 mins), the incorporation yield was 77.0%. The F-18 labeled product was confirmed by co-injection with the F-19 cold standard on the same analytical column for analyzing the reaction
- 15 mixture.

Figure 10: HPLC chromatogram of reaction mixture with co-injection of the cold standard


Example K1. Radiosynthesis of 3-cyano-4-[¹⁸F]fluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His-FA01010-Leu-NH₂

- 25 ¹⁸F-fluoride (2475MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and cesium carbonate (2.3mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. A solution of 3-cyano-4-trimethylammoniumbenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His-FA01010-

Leu-NH₂ trifluoroacetate salt (2mg) in anhydrous DMSO (150μl) was added. After heating at 70°C for 15 min. The reaction mixture was transferred to a vial containing water (4ml). The reaction vial washed with 150μl DMSO and this was also transferred to the vial containing water. This solution was transferred to a semi-prep HPLC (column: ACE 5μ C18, 250x10 mm, solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 30%B for 5min at 2ml/min, then 30-70% B in 10 mins at 3ml/min) and the desired F18 product peak was collected (253MBq, 20.4% d.c.). The F-18 labeled product was confirmed by co-injection with the F-19 cold standard on an analytical HPLC (Column ACE 3μ C18 50x4.6mm, 1ml/min (Agilent), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 0% for 2min, then 0%B to 95%B in 7 mins).

Figure 11: HPLC chromatogram of reaction mixture with co-injection of the cold standard

Example L1. Radiosynthesis of 3-cyano-4-[¹⁸F]fluorobenzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂




20

¹⁸F-fluoride (2475MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and cesium carbonate (2.3mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. A solution of 3-cyano-4-trimethylammoniumbenzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂ trifluoroacetate salt (2mg) in anhydrous DMSO (150μl) was added. After heating at 70°C for 15 min. The reaction mixture was transferred to a vial containing water (4ml). The reaction vial washed with 150μl DMSO and this was also transferred to the vial containing water. This solution was transferred to a semi-prep HPLC (column: ACE 5μ C18, 250x10 mm, solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 30%B for 5min at 2ml/min, then 30-70% B in 10 mins at 3ml/min) and the desired F18 product peak was collected (144MBq, 11.5% d.c.). The F-18 labeled product

was confirmed by co-injection with the F-19 cold standard on an analytical HPLC (Column ACE 3 μ C18 50 $\times$ 4.6mm, 1ml/min (Agilent), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 0% for 2min, then 0%B to 95%B in 7 mins).


5 Figure 12:HPLC chromatogram of reaction mixture with co-injection of the cold standard

10 Example M1. Radiosynthesis of 3-cyano-4-[¹⁸F]fluorobenzoyl-Arg- β Ala-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂




15 ¹⁸F-fluoride (1419MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and potassium carbonate (1mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. A solution of 3-cyano-4-trimethylammoniumbenzoyl-Arg- β Ala-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂ trifluoroacetate salt (2mg) in anhydrous DMSO (150 μ l) was added. After
 20 heating at 50°C for 15 min. The reaction mixture was diluted with water (4ml) and transferred to a semi-prep HPLC (column: Zorbax Bonus 5 μ C18, 250 $\times$ 9.2 mm, solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 29-34% B in 20 mins at 3ml/min) and the desired F18 product peak was collected (150MBq, 21.1% d.c.). The F-18 labeled product was confirmed by co-injection with the F-19 cold standard on an
 25 analytical HPLC (Column Zorbax Bonus 5 μ C18 250 $\times$ 4.6mm, 1ml/min (Agilent), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 29%B to 34%B in 20 mins).

30 Example N1. Radiosynthesis of 3-cyano-4-[¹⁸F]fluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂



¹⁸F-fluoride (869MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and potassium carbonate (1mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. A solution of 3-cyano-4-trimethylammoniumbenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu -NH₂ trifluoroacetate salt (2mg) in anhydrous DMSO (150μl) was added. After heating at 50°C for 15 min. The reaction mixture was diluted with water (4ml) and transferred to a semi-prep HPLC (column: Zorbax Bonus 5μ C18, 250x9.2 mm, solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 34-38% B in 20 mins at 3ml/min) and the desired F18 product peak was collected (184MBq, 37.8% d.c.). The F-18 labeled product was confirmed by co-injection with the F-19 cold standard on an analytical HPLC (Column Zorbax Bonus 5μ C18 250x4.6mm, 1ml/min (Agilent), solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 34%B to 38%B in 20 mins).

Example O1. Radiosynthesis of 3-trifluoromethyl-4-[¹⁸F]fluorobenzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂




20

¹⁸F-fluoride (835MBq) was azeotropically dried in the presence of Kryptofix 222 (5mg in 1.5ml MeCN) and cesium carbonate (2.3mg in 0.5ml water) by heating under nitrogen at 120°C for 30 minutes. During this time 2 x 1ml MeCN were added and evaporated. A solution of 3-trifluoromethyl-4-trimethylammoniumbenzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu NH₂ trifluoroacetate salt (2mg) in anhydrous DMSO (150μl) was added. After heating at 70°C for 15 min. The reaction mixture was diluted with water

25

(4ml) and transferred to a semi-prep HPLC (column: ACE 5 μ C18 250x10 mm, solvent A: H₂O + 0.1%TFA, solvent B: MeCN/Water 9/1 + 0.1% TFA, gradient: 20% for 2min, then 20-60% B in 20 mins at 3ml/min) and the desired F18 product peak was collected (78MBq, 29.0% d.c.). The F-18 labeled product was confirmed by co-injection with the F-19 cold standard on an analytical HPLC (Column ACE C18, 50x4.6mm, 3 μ , 2ml/min, solvent A: 10mM K₂HPO₄ in H₂O, solvent B: 10mM K₂HPO₄ in MeCN/H₂O (7:3), gradient: 5%-95% B in 7 mins).


Generic Structure of 3-Cyano activated Bombesin peptides precursors



Precursor	Sequence
	Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa-NH2
	Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-FA1010-Leu-NH2
	Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH2
	Arg-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH2
	Lys(Me)2-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH2
	Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-FA01010-Leu-NH2
	Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-FA01010-Leu-NH2
	Lys(Me)2-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-FA01010-Leu-NH2
	Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH2
	Lys(Me)2-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-FA01010-Leu-NH2
	Ava-Gln-Trp-Ala-Val-Gly-His-FA01010-Leu-NH2
	Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH2
	Arg-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH2
	Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH2
	1,4-cis-Achc-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH2
	Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Cpa-NH2
	Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Fa01010-Cpa-NH2
	Dioxa-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH2
	Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH2

5

Generic Structure of 3-Cyano activated Bombesin peptides labeled with F-18




5

10

15

F18 labeled	Sequence
	Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa-NH ₂
	Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-FA01010-Leu-NH ₂
	Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH ₂
	Arg-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH ₂
	Lys(Me) ₂ -βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH ₂
	Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-FA01010-Leu-NH ₂
	Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-FA01010-Leu-NH ₂
	Lys(Me) ₂ -Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-FA01010-Leu-NH ₂
	Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH ₂
	Lys(Me) ₂ -βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-FA01010-Leu-NH ₂
	Ava-Gln-Trp-Ala-Val-Gly-His-FA01010-Leu-NH ₂
	Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH ₂
	Arg-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH ₂
	Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH ₂
	1,4-cis-Achc-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH ₂
	Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Cpa-NH ₂
	Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Fa01010-Cpa-NH ₂
	Dioxa-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH ₂
	Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH ₂

Generic Structure of 3-Trifluoromethyl activated Bombesin peptides precursors



20


Precursor	Sequence
	Arg-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-FA01010-Leu-NH ₂
	Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH ₂
	Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH ₂
	Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH ₂

25

30

35

40

Generic Structure of 3-Trifluoromethyl activated Bombesin peptides labeled with F-18

5

F18 labeled	Sequence
	Arg- β Ala-Gln-Trp-Ala-Val-Gly-His(3Me)-FA01010-Leu-NH ₂
	Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH ₂
	Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH ₂
	Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH ₂


Aminoacid abbreviations

All natural amino acids were represented by 3-letter codes. Unless otherwise stated all the aminoacids have L-configurations.

5

Sta – Statine

His(3Me) – 3 – methylhistidine



Ava – 5-aminovaleric acid

AOC – 8-aminooctanoic acid

15

tBuGly – t-butylglycine

tBuAla – t-butylalanine

20 β hLeu – β -homoleucine β hIle – β -homoisoleucineLys(Me)₂ – N,N-dimethyllysine

25

DOA – 3,6-dioxo-8-aminooctanoic acid

4-Am-5-MeHpA – 4-amino-5-methylheptanoic acid

30 4-Am-5-MeHxA – 4-amino-5-methylhexanoic acid

1,4-cis-ACHC – 1,4-cis-aminocyclohexanecarboxylic acid

AHMHxA – (3R,4S)-4-amino-3-hydroxy-5-methylhexanoic acid

Biodistribution of F-18-Bombesin analog

see figure 13

5 wherein Bombesin analogue is Gln-Trp-Ala-Val-Gly-His-FA01010-Leu-NH₂

Radiolabeling of this bombesin analogue with F-18 was carried out via the method. The radiochemical yield was approx. 27% (decay corrected) giving 76 MBq in 50 µl ethanol
 10 with a radiochemical purity of >99% by HPLC and a specific activity of ~480 GBq/ mmol.

Nude mice bearing human prostate cancer PC-3 were injected with 100 µl radioactive peptide dissolved in PBS containing 135 kBq per animal. For blocking 100 µg unlabeled gastrin-releasing peptide was co-injected. One hour post injection the animals were
 15 sacrificed and organs dissected for counting in a gamma-counter. Values are expressed as percent of the injected dose per gram organ weight.

	1h %ID/g	1h Blocking %ID/g
Tumor (% ID/g)	1,00 ±0,01	0,18 ±0,03
Blood (% ID/g)	0,05 ±0,01	0,12 ±0,00
Muscle (% ID/g)	0,02 ±0,00	0,03 ±0,02
Pancreas (% ID/g)	0,34 ±0,03	0,10 ±0,02
Liver (% ID/g)	0,35 ±0,13	0,39 ±0,05
Kidneys (% ID/g)	0,24 ±0,02	0,71 ±0,12
Tumor/ Tissue-Ratios		
T/ Blood	21,03 ±11,92	1,57 ±0,22
T/ Muscle	59,99 ±29,53	6,31 ±3,27

20 It can be seen that ¹⁸F-labelled bombesin analog accumulates in tumor and the targeting agent ¹⁸F- labelled bombesin is specific since the blocking values are low in case of tumor and unchanged for the other tissue.

Comparison of ^{18}F -labelled bombesin analogs

Protocol as above

Table 1

5

Table 1 shows biodistribution in Nude mice bearing human prostate cancer PC-3 were injected with 100 μl radioactive peptide dissolved in PBS containing 135 kBq per animal.

10

Bombesin Analogs for PET: Comparison with ^{18}F -choline (FCH) and ^{18}F -FB-Lys-BN

5

Figure 8 shows that tumor – tissue ratio of Bombesin analog Gln-Trp-Ala-Val-Gly-His-FA01010-Leu-NH₂ is 2,5 time higher than the tumor – tissue ratio of ^{18}F -choline (FCH) and ^{18}F -FB-Lys-BN.

10


Synthesis of H-Y-E: Solid-phase peptide synthesis (SPPS) involves the stepwise addition of amino acid residues to a growing peptide chain that is linked to an insoluble support or matrix, such as polystyrene. The C-terminal residue of the peptide is first anchored to a commercially available support (e.g., Rink amide resin) with its amino group protected with an N-protecting agent, fluorenylmethoxycarbonyl (Fmoc) group. The amino protecting group is removed with suitable deprotecting agent such as piperidine for Fmoc and the next amino acid residue (in N-protected form) is added with a coupling agents such as dicyclohexylcarbodiimide (DCC), di-isopropyl-cyclohexylcarbodiimide (DCCI), hydroxybenzotriazole (HOBt). Upon formation of a peptide bond, the reagents are washed from the support. After addition of the final residue of (Y), the peptide is attached to the solid support is ready for the coupling of RG--L₁--B₁-OH.

It is understood that the examples and embodiments described herein are for illustrative purpose only and that various modifications and changes in light thereof as well as combinations of features described in this application will be suggested to persons skilled in the art and are to be included within the spirit and purview of the described invention and within the scope of the appended claims. From the foregoing description, one skilled in the art can easily ascertain the essential characteristics of this invention and, without departing from the spirit and scope thereof, can make various changes and modifications of the invention to adapt it to various usages and conditions. The entire disclosure[s] of all applications, patents and publications, cited herein are incorporated by reference herein.

Claims:

1. A compound having general chemical Formula I

5



A

wherein

10

-G is selected from the group comprising -F, -Cl, -Br, -I, -NO, -NO₂, -NR⁴COCF₃, -NR⁴SO₂CF₃, -N(R⁴)SO₂R⁵, -N(CF₃)₂, -NHCSR⁵, -N(SO₂R⁵)₂, -N(O)=NCONH₂, -NR⁴CN, -NHCSR⁵, -N≡C, -N=C(CF₃)₂, -N=NCF₃, -N=NCN, -NR⁴COR⁴, -NR⁴COOR⁵, -OSO₂CF₃, -OSO₂C₆H₅, -OCOR⁵, -ONO₂, -OSO₂R⁵, -O-C=CH₂, -OCF₂CF₃, -OCOCF₃, -OCN, -OCF₃, -C≡N, -C(NO₂)₃, -COOR⁴, -CONR⁴R⁵, -C(S)NH₂, -CH=NOR⁴, -CH₂SO₂R⁴, -COCF₃, -CF₃, -CF₂Cl-CBr₃, -CCIF₂, -CCl₃, -CF₂CF₃, -C≡CR⁴, -CH=NSO₂CF₃, -CH₂CF₃, -COR⁵, -CH=NOR⁵, -CH₂CONH₂, -CSNHR⁵, -CH=NNHCSR⁵, -CH=NNHCONHNH₂, -C≡C-CF₃, -CF=CFCF₃, -CF₂-CF₂-CF₃, -CR⁴(CN)₂, -COCF₂CF₂CF₃, -C(CF₃)₃, -C(CN)₃, -CR⁴=C(CN)₂, -1-pyrryl, -C(CN)=C(CN)₂, -C-pyridyl, -COC₆H₅, -COOC₆H₅, -SOCF₃, -SO₂CF₃, -SCF₃, -SO₂CN, -SCOCF₃, -SOR⁵, -S(OR⁵), -SC≡CR⁴, -SO₂R⁵, -SSO₂R⁵, -SR⁵, -SSR⁴, -SO₂CF₂CF₃, -SCF₂CF₃, -S(CF₃)=NSO₂CF₃, -SO₂C₆H₅, -SO₂N(R⁵)₂, -SO₂C(CF₃)₃, -SC(CF₃)₃, -SO(CF₃)=NSO₂CF₃, -S(O)(=NH)CF₃, -S(O)(=NH)R⁵, -S-C=CH₂, -SCOR⁵, -SOC₆H₅, -P(O)C₃F₇, -PO(OR⁵)₂, -PO(N(R⁵)₂)₂, -P(N(R⁵)₂)₂, -P(O)R⁵₂, and -PO(OR⁵)₂ or another electron-drawing group wherein the respective substituent can be in ortho, para or meta position in respect of the K group,

25

-Q is hydrogen, unbranched or branched lower alkyl, aryl, heteroaryl, -O-(C₁-C₄ alkyl), -CN, -halogen, -SO₂-R⁴, -NO₂ or a condensed aryl or condensed heteroaryl wherein the respective substituent can be in ortho, para or meta position in respect of the K group,

30

wherein R^4 is hydrogen or unbranched or branched lower alkyl and
 R^5 is unbranched or branched lower alkyl,

5 $-L-$ is a bond, $-CO-$, $-SO_2-$, $-(CH_2)_d-CO-$, $-SO-$, $-C\equiv C-CO-$, $-[CH_2]_m-E-[CH_2]_n-CO-$,
 $-[CH_2]_m-E-[CH_2]_n-SO_2-$, $-C(=O)-O-$, $-NR^{10}-$, $-O-$, $-(S)_p-$, $-C(=O)NR^{12}-$, $-C(=S)NR^{12}-$,
 $-C(=S)O-$, C_1-C_6 cycloalkyl, alkenyl, heterocycloalkyl, unsubstituted or substituted
aryl or unsubstituted or substituted heteroaryl, aralkyl, heteroaralkyl, alkyloxy,
aryloxy, aralkyloxy, $-SO_2NR^{13}-$, $-NR^{13}SO_2-$, $-NR^{13}C(=O)O-$, $-NR^{13}C(=O)NR^{12}-$, -
NH-NH- and -NH-O-,

10 wherein

d is an integer of from 1 to 6,

m and n , independently, are any integer of from 0 to 5,

$-E-$ is a bond, $-S-$, $-O-$ or $-NR^9-$,

wherein R^9 is H, C_1-C_{10} alkyl, aryl, heteroaryl or aralkyl,

15 p is any integer of from 1 to 3 ,

R^{10} and R^{12} , independently, are selected from the group comprising H, C_1-
 C_{10} alkyl, aryl, heteroaryl or aralkyl and

R^{13} is H, substituted or non substituted, linear or branched C_1-C_6 alkyl, aryl,
cycloalkyl, heterocycloalkyl, heteroaryl, aralkyl or heteroaralkyl,

20

Y is a bond or a spacer,

U is a targeting agent,

25

X^- is $CF_3S(O)_2O^-$, $C_4F_9S(O)_2O^-$, iodide anion, bromide anion, chloride anion,
perchlorate anion (ClO_4^-), phosphate anion, trifluoroacetate anion (CF_3-
 $C(O)O^-$), or the anion of another salt of an inorganic or organic acid,

K is $N^+(R^1)(R^2)(R^3) X^-$ or W ,

30

wherein

R^1 , R^2 and R^3 are independently from each other selected from the group
comprising substituted or unsubstituted alkyl and aralkyl,


35

W is a fluorine isotope,

and a pharmaceutically acceptable salt of an inorganic or organic acid thereof, a hydrate, complex, ester, amide, solvate and prodrug thereof.

2. The compound according to claim 1, wherein **W** = ^{18}F or ^{19}F , more preferably ^{18}F .
- 5
3. The compound according to any one of the preceding claims, wherein **-L-** is a bond, $-\text{CO}-$, $-\text{SO}_2-$, $-(\text{CH}_2)_d-\text{CO}-$, $-\text{SO}-$ or $-\text{C}\equiv\text{C}-\text{CO}-$ and wherein *d* is an integer of from 1 to 6.
- 10
4. The compound according to any one of the preceding claims, wherein **-L-** is $-\text{CO}-$, $-\text{SO}_2-$, or $-\text{C}\equiv\text{C}-\text{CO}-$.
5. The compound according to any one of the preceding claims, wherein **-L-** is $-\text{CO}-$, or $-\text{SO}_2-$.
- 15
6. The compound according to any one of the preceding claims, wherein **-G** is selected from the group comprising $-\text{F}$, $-\text{Cl}$, $-\text{Br}$, $-\text{NO}_2$, $-\text{NR}^4\text{SO}_2\text{R}^5$, $-\text{NR}^4\text{COR}^4$, $-\text{NR}^4\text{COOR}^5$, $-\text{C}\equiv\text{N}$, $-\text{CONR}^4\text{R}^5$, $-\text{C}\equiv\text{CR}^4$, $-\text{COR}^5$, $-\text{CF}_3$ and $-\text{SO}_2\text{R}^5$ wherein the respective substituent can be in ortho, para or meta position in respect of the K group.
- 20
7. The compound according to any one of the preceding claims, wherein **-Q** is $-\text{H}$, $-\text{C}_1-\text{C}_4$ alkyl, $-\text{O}-(\text{C}_1-\text{C}_4\text{alkyl})$, $-\text{CN}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$ or $-\text{NO}_2$ wherein the respective substituent can be in ortho, para or meta position in respect of the K group.
- 25
8. The compound according to any one of the preceding claims, wherein **-Q** is $-\text{H}$, methyl, $-\text{O}-\text{methyl}$, $-\text{CN}$, $-\text{F}$, $-\text{Cl}$, $-\text{Br}$ or $-\text{NO}_2$ wherein the respective substituent can be in ortho, para or meta position in respect of the K group.
- 30
9. The compound according to any one of the preceding claims, wherein **-Q** is $-\text{H}$, $-\text{CN}$ or $-\text{F}$ wherein the respective substituent can be in ortho, para or meta position in respect of the K group.
- 35
10. The compound according to any one of the preceding claims, wherein **-G** and **-Q** are independently from each other H , $-\text{CF}_3$, or $-\text{CN}$, whereas at least $-\text{G}$ or $-\text{Q}$ is $-\text{CF}_3$ or $-\text{CN}$.

11. The compound according to any one of the preceding claims, wherein R^4 is hydrogen or unbranched or branched C_1 - C_6 alkyl.
- 5 12. The compound according to any one of the preceding claims, wherein R^5 is unbranched or branched C_1 - C_6 alkyl.
- 10 13. The compound according to any one of the preceding claims, wherein R^1 , R^2 and R^3 are independently from each other selected from the group comprising aralkyl or lower alkyl, whereas at least two moieties of R^1 , R^2 , R^3 are alkyl.
14. The compound according to any one of the preceding claims, wherein R^1 is aralkyl and R^2 and R^3 are each methyl.
- 15 15. The compound according to any one of the preceding claims, wherein R^1 , R^2 and R^3 are each methyl.
16. The compound according to any one of the preceding claims, wherein X^- is $CF_3-C(O)O^-$, $CF_3S(O)_2O^-$ or $C_4F_9S(O)_2O^-$.
- 20 17. The compound according to any one of the preceding claims, wherein X^- is $CF_3-C(O)O^-$ or $CF_3S(O)_2O^-$.
- 25 18. The compound according to any one of the preceding claims, wherein the spacer Y is a natural or unnatural amino acid sequence or mixture thereof or a non-amino acid group.
19. The compound according to any one of the preceding claims, wherein the spacer Y is an amino acid sequence with two (2) to twenty (20) amino acid residues.
- 30 20. The compound according to any one of the preceding claims, wherein Y is Arg-Ser, Arg-Ava, Lys(Me) $_2$ - β -ala, Lys(Me) $_2$ -ser, Arg- β -ala, Ser-Ser, Ser-Thr, Arg-Thr, S-alkylcysteine, Cysteic acid, thioalkylcysteine (S-S-Alkyl) or



wherein k and l are independently selected in the range of from 0 to 4.

21. The compound according to any of the preceding claims, wherein -Y- is a non-amino acid moiety selected from the group comprising

-NH-(CH₂)_p-CO-, wherein p is an integer of from 2 to 10,

-NH-(CH₂-CH₂-O)_q-CH₂-CH₂-CO-, wherein q is an integer of from 0 to 5,

-NH-cycloalkyl-CO- wherein cycloalkyl is selected from C₅-C₈ cycloalkyl, more preferably C₆ atom cycloalkyl, and

-NH-heterocycloalkyl-(CH₂)_v-CO- wherein heterocycloalkyl is selected from C₅-C₈ heterocycloalkyl containing carbon atoms and 1, 2, 3 or 4 oxygen, nitrogen or sulfur heteroatoms and v is an integer of from 1 to 4, more preferably v is an integer of from 1 to 2.

22. The compound according to any one of the preceding claims, wherein U is a targeting agent selected from the group comprising peptides, peptidomimetics, small molecules and oligonucleotides.

23. The compound according to any one of the preceding claims, wherein U is a peptide comprising from 4 to 100 amino acids.

24. The compound according to any one of the preceding claims, wherein U is selected from the group comprising somatostatin and derivatives thereof and related peptides, somatostatin receptor specific peptides, neuropeptide Y and derivatives thereof and related peptides, neuropeptide Y₁ and the analogs thereof, bombesin and derivatives thereof and related peptides, gastrin, gastrin releasing peptide and the derivatives thereof and related peptides, epidermal growth factor (EGF of various origin), insulin growth factor (IGF) and IGF-1, integrins (α₃β₁, α_vβ₃, α_vβ₅, αIIbβ₃), LHRH agonists and antagonists, transforming growth factors, particularly TGF-α; angiotensin; cholecystokinin receptor peptides, cholecystokinin (CCK) and the analogs thereof; neurotensin and the analogs thereof, thyrotropin releasing hormone, pituitary adenylate cyclase activating peptide (PACAP) and the related

peptides thereof, chemokines, substrates and inhibitors for cell surface matrix metalloproteinase, prolactin and the analogs thereof, tumor necrosis factor, interleukins (IL-1, IL-2, IL-4 or IL-6), interferons, vasoactive intestinal peptide (VIP) and the related peptides thereof

5

25. The compound according to any one of the preceding claims, wherein **U** is selected from the group comprising bombesin, somatostatin, neuropeptide Y₁ and analogs thereof.

10 26. The compound according to any one of the preceding claims, wherein **U** is selected from the group comprising bombesin analogs having sequence III or IV:

AA₁-AA₂-AA₃-AA₄-AA₅-AA₆-AA₇-AA₈-NT₁T₂ (type A) III, with:

T₁ = T₂ = H or T₁ = H, T₂ = OH or T₁ = CH₃, T₂ = OH

AA₁ = Gln, Asn, Phe(4-CO-NH₂)

15

AA₂ = Trp, D-Trp

AA₃ = Ala, Ser, Val

AA₄ = Val, Ser, Thr

AA₅ = Gly, (N-Me)Gly

AA₆ = His, His(3-Me), (N-Me)His, (N-Me)His(3-Me)

20

AA₇ = Sta, Statine analogs and isomers, 4-Am,5-MeHpA, 4-Am,5-MeHxA, γ-substituted aminoacids

AA₈ = Leu, Cpa, Cba, CpnA, Cha, t-buGly, tBuAla, Met, Nle, iso-Bu-Gly

AA₁-AA₂-AA₃-AA₄-AA₅-AA₆-AA₇-AA₈-NT₁T₂ (type B) IV, with:

25

T₁ = T₂ = H or T₁ = H, T₂ = OH or T₁ = CH₃, T₂ = OH

AA₁ = Gln, Asn or Phe(4-CO-NH₂)


AA₂ = Trp, D-Trp

AA₃ = Ala, Ser, Val

AA₄ = Val, Ser, Thr

30

AA₅ = βAla, β²- and β³-amino acids as shown herein after



wherein SC represents a side chain found in proteinogenic amino acids and homologs of proteinogenic amino acids,

35

AA₆ = His, His(3-Me), (N-Me)His, (N-Me)His(3-Me)

AA₇ = Phe, Tha, Nal,

AA₈ = Leu, Cpa, Cba, CpnA, Cha, t-buGly, tBuAla, Met, Nle, iso-Bu-Gly.

27. The compound according to any one of the preceding claims, wherein **U** is
 5 NR⁷-peptide, or -(CH₂)_n-peptide, -O-(CH₂)_n- peptide or -S-(CH₂)_n- peptide, NR⁷-
 small-molecule, or -(CH₂)_n- small-molecule, -O-(CH₂)_n- small-molecule or -S-(CH₂)_n-
 small-molecule, NR⁷- oligonucleotide, or -(CH₂)_n- oligonucleotide, -O-(CH₂)_n-
 oligonucleotide or -S-(CH₂)_n- oligonucleotide, wherein n is an integer of from 1 to 6.

10 28. The compound according to any one of the preceding claims, wherein **R**⁷ is
 hydrogen or unbranched or branched C₁-C₆ alkyl.

29. The compound according to any one of the preceding claims, wherein **R**⁷ is
 hydrogen or methyl.

15 30. The compound according to any one of the preceding claims, wherein **U** is a small
 molecule having a molecular mass of from 200 to 800.

20 31. The compound according to any one of the preceding claims, wherein **U** is a
 oligonucleotide.

32. The compound according to any one of the preceding claims, comprising


- Ia-1 4-(Trimethylammonium)-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂,
- 25 • Ia-2 4-(Trimethylammonium)-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(Me)-Sta-Leu- NH₂,
- Ia-3 4-(Trimethylammonium)-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu- NH₂,
- Ia-4 4-(Trimethylammonium)-3-cyano-benzoyl-1,4-cis-Achc-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 30 • Ia-5 4-(Trimethylammonium)-3-cyano-benzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-6 4-(Trimethylammonium)-3-cyano-benzoyl-AOC-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 35 • Ia-7 4-(Trimethylammonium)-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa-NH₂,

- Ia-8 4-(Trimethylammonium)-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- Ia-9 4-(Trimethylammonium)-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 5 • Ia-10 4-(Trimethylammonium)-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-11 4-(Trimethylammonium)-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-12 4-(Trimethylammonium)-3-cyano-benzoyl-Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- 10 • Ia-13 4-(Trimethylammonium)-3-cyano-benzoyl-Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- Ia-14 4-(Trimethylammonium)-3-cyano-benzoyl-Lys(Me)₂-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-AM-5-MeHpA-Leu-NH₂,
- 15 • Ia-15 4-(Trimethylammonium)-3-cyano-benzoyl-Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-16 4-(Trimethylammonium)-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- Ia-17 4-(Trimethylammonium)-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His-4-Am,5-MeHpA- -Leu-NH₂,
- 20 • Ia-18 4-(Trimethylammonium)-3-trifluoromethyl-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-19 4-(Trimethylammonium)-3-trifluoromethyl-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- 25 • Ia-20 4-(Trimethylammonium)-3-trifluoromethyl-benzoyl-1,4-cis-Achc-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-21 4-(Trimethylammonium)-3-trifluoromethyl-benzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- Ia-22 4-(Trimethylammonium)-3-trifluoromethyl-benzoyl-Arg-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- 30 • Ia-23 4-(Trimethylammonium)-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Cpa-NH₂,
- Ia-24 4-(Trimethylammonium)-3-cyano-benzoyl -Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 35 • Ia-25 4-(Trimethylammonium)-3-cyano-benzoyl-DOA-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,


- 1a-66: 4-(Trimethylammonium)-3-cyano-benzoyl- Ava- ϵ -c[Lys-(NMe)Phe-1Nal-D-Trp-Lys-Thr]
- 1a-67: 4-(Trimethylammonium)-3-cyano-benzoyl- Ava- β -c[Dpr-Met-(NMe)Phe-Tyr-D-Trp-Lys]
- 5 • 1a-68: 4-(Trimethylammonium)-3-cyano-benzoyl- Ava-DCys-Leu-Ile-Thr-Arg-Cys-Arg-Tyr-NH₂]
- 1a-69: 4-(Trimethylammonium)-3-cyano-benzoyl- Ava-DCys-Leu-Ile-Val-Arg-Cys-Arg-Tyr-NH₂]

10


15



20





25




30

35





and



33. The compound according to any one of claims 1 – 32, selected from

- IIA-a-1 4-[18]Fluoro-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂,
- IIA-a-2 4-[18]Fluoro-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(Me)-Sta-Leu-NH₂,
- IIA-a-3 4-[18]Fluoro-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- IIA-a-4 4-[18]Fluoro-3-cyano-benzoyl-1,4-cis-Achc-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIA-a-5 4-[18]Fluoro-3-cyano-benzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIA-a-6 4-[18]Fluoro-3-cyano-benzoyl-AOC-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIA-a-7 4-[18]Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa-NH₂,
- IIA-a-8 4-[18]Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-FA4-Am,5-MeHpA-Leu-NH₂,
- IIA-a-9 4-[18]Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,

- IIA-a-10 4-[18]Fluoro-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIA-a-11 4-[18]Fluoro-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 5 • IIA-a-12 4-[18]Fluoro-3-cyano-benzoyl-Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIA-a-13 4-[18]Fluoro-3-cyano-benzoyl-Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIA-a-14 4-[18]Fluoro-3-cyano-benzoyl-Lys(Me)₂-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- 10 • IIA-a--15 4-[18]Fluoro-3-cyano-benzoyl-Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIA-a-16 4-[18]Fluoro-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- 15 • IIA-a-17 4-[18]Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His-4-Am,5-MeHpA-Leu-NH₂,
- IIA-a-18 4-[18]Fluoro-3-trifluoromethyl-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-LeuNH₂,
- IIA-a-19 4-[18]Fluoro-3-trifluoromethyl-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- 20 • IIA-a-20 4-[18]Fluoro-3-trifluoromethyl-benzoyl-1,4-cis-Achc-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIA-a-21 4-[18]Fluoro-3-trifluoromethyl-benzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 25 • IIA-a-22 4-[18]Fluoro-3-trifluoromethyl-benzoyl-Arg-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- 4-[18]Fluoro-3-cyano-benzoyl-(piperidyl-4-carbonyl)-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 4-[18]Fluoro-3-cyano-benzoyl-(piperazin-1-yl-acetyl)-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 30 • 4-[18]Fluoro-3-cyano-benzoyl-1,4-trans-Achc-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂,
- IIB-a-1 4-[19]-Fluoro-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂,
- 35


- IIB-a-2 4-[19]-Fluoro-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-His(Me)-Sta-Leu-NH₂,
- IIB-a-3 4-[19]-Fluoro-3-cyano-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- 5 • IIB-a-4 4-[19]-Fluoro-3-cyano-benzoyl-1,4-cis-Achc-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-5 4-[19]-Fluoro-3-cyano-benzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-6 4-[19]-Fluoro-3-cyano-benzoyl-AOC-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 10 • IIB-a-7 4-[19]-Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa-NH₂,
- IIB-a-8 4-[19]-Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-9 4-[19]-Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 15 • IIB-a-10 4-[19]-Fluoro-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-11 4-[19]-Fluoro-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 20 • IIB-a-12 4-[19]-Fluoro-3-cyano-benzoyl-Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-13 4-[19]-Fluoro-3-cyano-benzoyl-Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-14 4-[19]-Fluoro-3-cyano-benzoyl-Lys(Me)₂-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- 25 • IIB-a-15 4-[19]-Fluoro-3-cyano-benzoyl-Arg-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-16 4-[19]-Fluoro-3-cyano-benzoyl-Lys(Me)₂-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- 30 • IIB-a-17 4-[19]-Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-Gly-His-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-18 4-[19]-Fluoro-3-trifluoromethyl-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-19 4-[19]-Fluoro-3-trifluoromethyl-benzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- 35

- IIB-a-20 4-[19]-Fluoro-3-trifluoromethyl-benzoyl-1,4-cis-Achc-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-21 4-[19]-Fluoro-3-trifluoromethyl-benzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 5 • IIB-a-22 4-[19]-Fluoro-3-trifluoromethyl-benzoyl-Arg-βAla-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-23 4-[19]-Fluoro-3-cyano-benzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Cpa-NH₂,
- IIB-a-24 4-[19]-Fluoro-3-cyano-benzoyl -Ser-Ser-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-
10 Leu-NH₂,
- IIB-a-25 4-[19]-Fluoro-3-cyano-benzoyl-DOA-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-26 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂,
- IIB-a-27 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His-FA02010-Cpa-
15 NH₂,
- IIB-a-28 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-tbuGly-NH₂,
- IIB-a-29 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- 20 • IIB-a-30 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-tBuGly-NH₂,
- IIB-a-31 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-32 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- 25 • IIB-a-33 3,4-[19]-Difluorobenzoyl-Ava-Gln-DTrp-Ala-Val-Gly-His-4-Am,5-MeHpA-tbuGly-NH₂,
- IIB-a-34 3,4-[19]-Difluorobenzoyl-Ava-Gln-DTrp-Ala-Val-Gly-His-4-Am-5-MeHxA-Cpa-NH₂,
- IIB-a-35 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa-
30 NH₂,
- IIB-a-36 3,4-[19]-Difluorobenzoyl-Ava-Gln-DTrp-Ala-Val-Gly-His-Sta-tbuAla-NH₂,
- IIB-a-37 3,4-[19]-Difluorobenzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu-NH₂,
- IIB-a-38 3,4-[19]-Difluorobenzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 35 • IIB-a-39 3,4-[19]-Difluorobenzoyl-Arg-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,


- IIB-a-40 3,4-[19]-Difluorobenzoyl-Arg-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-41 3,4-[19]-Difluorobenzoyl-Arg-βAla-Arg-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu-NH₂,
- 5 • IIB-a-42 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Cpa-NH₂,
- IIB-a-43 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-tBuGly-NH₂,
- IIB-a-44 3,4-[19]-Difluorobenzoyl-Arg-Arg-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- 10 • IIB-a-45 3,4-[19]-Difluorobenzoyl-Arg-βAla-Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu-NH₂,
- IIB-a-46 3,4-[19]-Difluorobenzoyl-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-47 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Cpa-NH₂,
- 15 • IIB-a-48 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-49 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-NMeHis-4-Am,5-MeHpA-Cpa-NH₂,
- 20 • IIB-a-49 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-NMeHis(3Me)-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-50 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-Gly-NMeHis-4-Am,5-MeHpA-Leu-NH₂,
- IIB-a-51 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-NMeGly-His-AHMHxA -Leu-NH₂,
- 25 • IIB-a-52 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Tha-Cpa-NH₂,
- IIB-a-53 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Phe-Cpa-NH₂,
- IIB-a-54 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Phe-Leu-NH₂,
- IIB-a-55 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-DHis-Phe-Leu-NH₂,
- 30 • IIB-a-56 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-βhLeu-Leu-NH₂,
- IIB-a-57 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-βhIle-Leu-NH₂,
- IIB-a-58 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-βhLeu-tbuGly-NH₂,
- IIB-a-59 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Tha-NH₂,
- IIB-a-60 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Nle-NH₂,
- 35 • IIB-a-61 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Phe-tbuGly-NH₂,

- IIB-a-62 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-NMeHis-Tha-tbuGly-NH₂,
- IIB-a-63 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His(3Me)-Tha-tbuGly-NH₂,
- 5 • IIB-a-64 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Cpa-NH₂,
- IIB-a-65 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-NMeVal-βAla-His-Phe-Leu-NH₂,
- IIB-a-66 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-NMePhe-Leu-NH₂,
- IIB-a-67 3,4-[19]-Difluorobenzoyl-Ava-Gln-DTrp-Ala-Val-βAla-His-Phe-Leu-NH₂,
- IIB-a-68 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-DAla-Val-βAla-His-Phe-Leu-NH₂,
- 10 • IIB-a-69 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-DVal-βAla-His-Phe-Leu-NH₂,
- IIB-a-70 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-DPhe-Leu-NH₂,
- IIB-a-71 3,4-[19]-Difluorobenzoyl-Ava-Gln-Trp-Ala-Val-βAla-His-βhIle-tbuGly-NH₂,
- IIB-a-72 4-[19]-Fluoro-3-cyano-phenylsulfonyl -Ava-Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-Cpa-NH₂,
- 15 • IIB-a-73 4-[19]-Fluoro-3-cyano-phenylsulfonyl -Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-Cpa-NH₂,
- IIB-a-74 4-[19]-Fluoro-3-cyano-phenylsulfonyl -Ava-Gln-Trp-Ala-Val-NMeGly-His-Sta-tbuAla-NH₂,
- IIB-a-75 4-[19]-Fluoro-3-cyano-phenylsulfonyl -Ava-Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-tbuAla-NH₂.
- 20 • IIA-a-76:4-[18]Fluoro-3-cyano-benzoyl- Ava-ε-c[Lys-(NMe)Phe-1Nal-D-Trp-Lys-Thr]
- IIA-a-77:4-[18]Fluoro-3-cyano-benzoyl- Ava-β-c[Dpr-Met-(NMe)Phe-Tyr-D-Trp-Lys]
- IIB-a-76:4-[19]Fluoro-3-cyano-benzoyl- Ava-ε-c[Lys-(NMe)Phe-1Nal-D-Trp-Lys-Thr]
- IIB-a-77:4-[19]Fluoro-3-cyano-benzoyl- Ava-β-c[Dpr-Met-(NMe)Phe-Tyr-D-Trp-Lys]
- 25 • IIA-a-78:4-[18]Fluoro-3-cyano-benzoyl- Ava-DCys-Leu-Ile-Thr-Arg-Cys-Arg-Tyr-NH₂
- IIA-a-79:4-[18]Fluoro-3-cyano-benzoyl- Ava- DCys-Leu-Ile-Val-Arg-Cys-Arg-Tyr-NH₂
- IIA-a-78:4-[19]Fluoro-3-cyano-benzoyl- Ava-DCys-Leu-Ile-Thr-Arg-Cys-Arg-Tyr-NH₂
- IIA-a-79:4-[19]Fluoro-3-cyano-benzoyl- Ava- DCys-Leu-Ile-Val-Arg-Cys-Arg-Tyr-NH₂


204



3-Cyano-4-fluoro-N-{6-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-thiophen-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-hexyl}-benzamide




3-Cyano-4-fluoro-N-{4-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-butyl}-benzamide




3-Cyano-4-fluoro-N-{2-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-ethyl}-benzamide

5




10


3-Cyano-4-fluoro-N-(2-(2-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-ethanesulfinyl)-ethyl)-benzamide




10 3-Cyano-4-fluoro-N-(2-{2-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-ethanesulfonyl}-ethyl)-benzamide



20 3-Cyano-4-fluoro-N-(2-{2-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-ethylsulfanyl}-ethyl)-benzamide




30 3-Cyano-N-((2R,3R)-2,3-dihydroxy-4-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-butyl)-4-fluoro-benzamide




3-Cyano-4-[F]fluoro-N-(thymidiny-propyl)-benzamide

5




15

2-Chloro-4-fluoro-N-{3-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-5-methyl-2,6-dioxo-3,6-dihydro-2H-pyrimidin-1-yl]-propyl}-benzamide



3-Cyano-4-[F]fluoro-N-(2-[2-thymidinyl-ethoxy]-ethyl)-benzamide;

20




25


3-Cyano-4-fluoro-N-{6-[3-((2R,4S,5R)-4-hydroxy-5-hydroxymethyl-tetrahydro-furan-2-yl)-2,6-dioxo-5-trifluoromethyl-3,6-dihydro-2H-pyrimidin-1-yl]-hexyl}-benzamide

30

5



or



wherein fluoro (F) is ^{18}F and ^{19}F .

10

34. The compound according to claims 1 to 32, wherein U is selected from the group comprising

15

- Seq ID 1 Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu- NH_2
- Seq ID 2 Gln-Trp-Ala-Val-Gly-His(Me)-Sta-Leu- NH_2
- Seq ID 3 Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu- NH_2
- Seq ID 4 Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu- NH_2
- Seq ID 7 Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa- NH_2
- Seq ID 8 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu- NH_2
- Seq ID 12 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu- NH_2
- Seq ID 17 Gln-Trp-Ala-Val-Gly-His-4-Am,5-MeHpA- -Leu- NH_2
- Seq ID 23 Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Cpa- NH_2
- Seq ID 27 Gln-Trp-Ala-Val-NMeGly-His-FA02010-Cpa- NH_2
- Seq ID 28 Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-tbuGly- NH_2
- Seq ID 30 Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-tBuGly- NH_2
- Seq ID 32 Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Leu- NH_2
- Seq ID 33 Gln-DTrp-Ala-Val-Gly-His-4-Am,5-MeHpA-tbuGly- NH_2
- Seq ID 34 Gln-DTrp-Ala-Val-Gly-His-4-Am-5-MeHxA-Cpa- NH_2

25

- Seq ID 35 Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa- NH₂
- Seq ID 36 Gln-DTrp-Ala-Val-Gly-His-Sta-tbuAla- NH₂
- Seq ID 42 Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Cpa- NH₂
- Seq ID 43 Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-tBuGly- NH₂
- 5 • Seq ID 46 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu- NH₂
- Seq ID 48 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu- NH₂
- Seq ID 49 Gln-Trp-Ala-Val-Gly-NMeHis-4-Am,5-MeHpA-Cpa- NH₂
- Seq ID 49 Gln-Trp-Ala-Val-Gly-NMeHis(3Me)-4-Am,5-MeHpA-Leu- NH₂
- Seq ID 50 Gln-Trp-Ala-Val-Gly-NMeHis-4-Am,5-MeHpA-Leu- NH₂
- 10 • Seq ID 51 Gln-Trp-Ala-Val-NMeGly-His-AHMHxA -Leu- NH₂
- Seq ID 52 Gln-Trp-Ala-Val-βAla-NMeHis-Tha-Cpa- NH₂
- Seq ID 53 Gln-Trp-Ala-Val-βAla-NMeHis-Phe-Cpa- NH₂
- Seq ID 54 Gln-Trp-Ala-Val-βAla-NMeHis-Phe-Leu- NH₂
- Seq ID 55 Gln-Trp-Ala-Val-βAla-DHis-Phe-Leu- NH₂
- 15 • Seq ID 56 Gln-Trp-Ala-Val-βAla-His-βhLeu-Leu- NH₂
- Seq ID 57 Gln-Trp-Ala-Val-βAla-His-βhIle-Leu- NH₂
- Seq ID 58 Gln-Trp-Ala-Val-βAla-His-βhLeu-tbuGly- NH₂
- Seq ID 59 Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Tha- NH₂
- Seq ID 60 Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Nle- NH₂
- 20 • Seq ID 61 Gln-Trp-Ala-Val-βAla-NMeHis-Phe-tbuGly- NH₂
- Seq ID 62 Gln-Trp-Ala-Val-βAla-NMeHis-Tha-tbuGly- NH₂
- Seq ID 63 Gln-Trp-Ala-Val-βAla-His(3Me)-Tha-tbuGly- NH₂
- Seq ID 64 Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Cpa- NH₂
- Seq ID 65 Gln-Trp-Ala-NMeVal-βAla-His-Phe-Leu- NH₂
- 25 • Seq ID 66 Gln-Trp-Ala-Val-βAla-His-NMePhe-Leu- NH₂
- Seq ID 67 Gln-DTrp-Ala-Val-βAla-His-Phe-Leu- NH₂
- Seq ID 68 Gln-Trp-DAla-Val-βAla-His-Phe-Leu- NH₂
- Seq ID 69 Gln-Trp-Ala-DVal-βAla-His-Phe-Leu- NH₂
- Seq ID 70 Gln-Trp-Ala-Val-βAla-His-DPhe-Leu- NH₂
- 30 • Seq ID 71 Gln-Trp-Ala-Val-βAla-His-βhIle-tbuGly- NH₂
- Seq ID 72 Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-Cpa- NH₂
- Seq ID 73 Gln-Trp-Ala-Val-NMeGly-His-Sta-Cpa- NH₂
- Seq ID 74 Gln-Trp-Ala-Val-NMeGly-His-Sta-tbuAla- NH₂
- Seq ID 75 Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-tbuAla- NH₂
- 35 • Seq ID 77 Gln-Trp-Ala-Val-His(Me)-Sta-Leu- NH₂

- Seq ID 82 Gln-Trp-Ala-Val-Gly-His(3Me)-FA4-Am,5-MeHpA-Leu- NH₂
- Seq ID 90 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu- NH₂
- Seq ID 91 Gln-Trp-Ala-Val-Gly-His-4-Am,5-MeHpA-Leu- NH₂
- Seq ID 101 Gln-Trp-Ala-Val-Gly-His(3Me)- 4-Am-5-MeHpA – 4-amino-5-methylheptanoic acid -Leu- NH₂
- Seq ID 102 Gln-Trp-Ala-Val-NMeGly-His(3Me)- 4-Am-5-MeHpA – 4-amino-5-methylheptanoic acid -Cpa- NH₂

35. Method of preparing a compound having general chemical Formula A, wherein K = W, according to any one of claims 1 – 34, in which method a compound having general chemical Formula A, wherein K = -N⁺(R¹)(R²)(R³) X⁻, according to any one of claims 1 – 32 is labelled with a fluorine isotope.

36. Method according to claim 35, comprising the step of coupling a compound having general chemical Formula A, wherein K = -N⁺(R¹)(R²)(R³) X⁻, according to any one of claims 1 – 32, with a fluorine isotope to form a compound having general chemical Formula A, wherein K = W according to anyone of claims 1 -35, Wherein K, R¹, R², R³, X⁻ and W are defined as in the claims above.

37. A composition comprising a compound having general chemical Formula A, wherein K = -N⁺(R¹)(R²)(R³) X⁻ or W, according to any one of claims 1 – 35, and a pharmaceutically acceptable carrier, diluent, adjuvant or excipient.

38. A method of imaging diseases, the method comprising introducing into a patient a detectable quantity of a labelled compound having general chemical Formula A, wherein K = W, according to any one of claims 1 – 35, or of a pharmaceutically acceptable salt of an inorganic or organic acid thereof, a hydrate, complex, ester, amide, solvate and prodrug thereof.


39. A kit comprising a sealed vial containing a predetermined quantity of a compound having general chemical Formula A, wherein K = -N⁺(R¹)(R²)(R³) X⁻, according to any one of claims 1 – 35, or a pharmaceutically acceptable salt of an inorganic or organic acid thereof, a hydrate, complex, ester, amide, solvate and prodrug thereof.

40. A compound having general chemical Formula A, wherein K = -N⁺(R¹)(R²)(R³) X⁻ or W, according to any one of claims 1 – 35, or a pharmaceutically acceptable salt of

an inorganic or organic acid thereof, a hydrate, complex, ester, amide, solvate and prodrug thereof for use as medicament.

41. A compound having general chemical Formula A, wherein $K = W$, according to any one of claims 1 – 35, or a pharmaceutically acceptable salt of an inorganic or organic acid thereof, a hydrate, complex, ester, amide, solvate and prodrug thereof for use as diagnostic imaging agent.
42. A compound having general chemical Formula A, wherein $K = W$, according to any one of claims 1 – 35, or a pharmaceutically acceptable salt of an inorganic or organic acid thereof, a hydrate, complex, ester, amide, solvate and prodrug thereof for use as imaging agent for positron emission tomography (PET).
43. Use of a compound having general chemical Formula A, wherein $K = -N^+(R^1)(R^2)(R^3) X^-$ or W , according to any one of claims 1 – 35, or of a pharmaceutically acceptable salt of an inorganic or organic acid thereof, a hydrate, complex, ester, amide, solvate and prodrug thereof for the manufacture of a medicament.
44. Use of a compound having general chemical Formula A, wherein $K = -N^+(R^1)(R^2)(R^3) X^-$ or W , according to any one of claims 1 – 35, or of a pharmaceutically acceptable salt of an inorganic or organic acid thereof, a hydrate, complex, ester, amide, solvate and prodrug thereof for the manufacture of a diagnostic imaging agent.
45. The use according to claim 44 for the manufacture of a diagnostic imaging agent for PET imaging.

46. A compound having general chemical Formula V:



wherein $N^+(R^1)(R^2)(R^3)$, X^- , $-G$, and $-Q$, have the same meaning as depicted above for compounds having general chemical Formula A and

R^6 is selected from the group comprising $-S(O)_2-N(H)-CH_2-C(O)OH$, $-S(O)_2-N(Me)-CH_2-C(O)OH$ and $C(O)OH$.

5

47. A method of preparing compound of Formula A wherein $K = -N^+(R^1)(R^2)(R^3) X^-$ by reacting a compound of Formula V with a targeting agent.

48. The method according to claim 47 wherein the compound of Formula A wherein $K = -N^+(R^1)(R^2)(R^3) X^-$ and the targeting agent are reacted optionally with a condensing agent.

10

49. Peptide sequence selected from

- Seq ID 1 Gln-Trp-Ala-Val-NMeGly-His-Sta-Leu- NH_2
- 15 • Seq ID 2 Gln-Trp-Ala-Val-Gly-His(Me)-Sta-Leu- NH_2
- Seq ID 3 Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Leu- NH_2
- Seq ID 4 Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Leu- NH_2
- Seq ID 7 Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa- NH_2
- Seq ID 8 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu- NH_2
- 20 • Seq ID 12 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu- NH_2
- Seq ID 17 Gln-Trp-Ala-Val-Gly-His-4-Am,5-MeHpA-Leu- NH_2
- Seq ID 23 Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Cpa- NH_2
- Seq ID 27 Gln-Trp-Ala-Val-NMeGly-His-FA02010-Cpa- NH_2
- Seq ID 28 Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-tbuGly- NH_2
- 25 • Seq ID 30 Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-tBuGly- NH_2
- Seq ID 32 Gln-Trp-Ala-Val-NMeGly-His(3Me)-4-Am,5-MeHpA-Leu- NH_2
- Seq ID 33 Gln-DTrp-Ala-Val-Gly-His-4-Am,5-MeHpA-tbuGly- NH_2
- Seq ID 34 Gln-DTrp-Ala-Val-Gly-His-4-Am-5-MeHxA-Cpa- NH_2
- Seq ID 35 Gln-Trp-Ala-Val-NMeGly-His(3Me)-Sta-Cpa- NH_2
- 30 • Seq ID 36 Gln-DTrp-Ala-Val-Gly-His-Sta-tbuAla- NH_2
- Seq ID 42 Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-Cpa- NH_2
- Seq ID 43 Gln-Trp-Ala-Val-Gly-His(3Me)-Sta-tBuGly- NH_2
- Seq ID 46 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu- NH_2
- Seq ID 48 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu- NH_2
- 35 • Seq ID 49 Gln-Trp-Ala-Val-Gly-NMeHis-4-Am,5-MeHpA-Cpa- NH_2
- Seq ID 49 Gln-Trp-Ala-Val-Gly-NMeHis(3Me)-4-Am,5-MeHpA-Leu- NH_2

- Seq ID 50 Gln-Trp-Ala-Val-Gly-NMeHis-4-Am,5-MeHpA-Leu- NH₂
- Seq ID 51 Gln-Trp-Ala-Val-NMeGly-His-AHMHxA -Leu- NH₂
- Seq ID 52 Gln-Trp-Ala-Val-βAla-NMeHis-Tha-Cpa- NH₂
- Seq ID 53 Gln-Trp-Ala-Val-βAla-NMeHis-Phe-Cpa- NH₂
- 5 • Seq ID 54 Gln-Trp-Ala-Val-βAla-NMeHis-Phe-Leu- NH₂
- Seq ID 55 Gln-Trp-Ala-Val-βAla-DHis-Phe-Leu- NH₂
- Seq ID 56 Gln-Trp-Ala-Val-βAla-His-βhLeu-Leu- NH₂
- Seq ID 57 Gln-Trp-Ala-Val-βAla-His-βhIle-Leu- NH₂
- Seq ID 58 Gln-Trp-Ala-Val-βAla-His-βhLeu-tbuGly- NH₂
- 10 • Seq ID 59 Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Tha- NH₂
- Seq ID 60 Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Nle- NH₂
- Seq ID 61 Gln-Trp-Ala-Val-βAla-NMeHis-Phe-tbuGly- NH₂
- Seq ID 62 Gln-Trp-Ala-Val-βAla-NMeHis-Tha-tbuGly- NH₂
- Seq ID 63 Gln-Trp-Ala-Val-βAla-His(3Me)-Tha-tbuGly- NH₂
- 15 • Seq ID 64 Gln-Trp-Ala-Val-βAla-His(3Me)-Phe-Cpa- NH₂
- Seq ID 65 Gln-Trp-Ala-NMeVal-βAla-His-Phe-Leu- NH₂
- Seq ID 66 Gln-Trp-Ala-Val-βAla-His-NMePhe-Leu- NH₂
- Seq ID 67 Gln-DTrp-Ala-Val-βAla-His-Phe-Leu- NH₂
- Seq ID 68 Gln-Trp-DAla-Val-βAla-His-Phe-Leu- NH₂
- 20 • Seq ID 69 Gln-Trp-Ala-DVal-βAla-His-Phe-Leu- NH₂
- Seq ID 70 Gln-Trp-Ala-Val-βAla-His-DPhe-Leu- NH₂
- Seq ID 71 Gln-Trp-Ala-Val-βAla-His-βhIle-tbuGly- NH₂
- Seq ID 72 Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-Cpa- NH₂
- Seq ID 73 Gln-Trp-Ala-Val-NMeGly-His-Sta-Cpa- NH₂
- 25 • Seq ID 74 Gln-Trp-Ala-Val-NMeGly-His-Sta-tbuAla- NH₂
- Seq ID 75 Gln-Trp-Ala-Val-NMeGly-His-4-Am,5-MeHpA-tbuAla- NH₂
- Seq ID 77 Gln-Trp-Ala-Val-His(Me)-Sta-Leu- NH₂
- Seq ID 82 Gln-Trp-Ala-Val-Gly-His(3Me)-FA4-Am,5-MeHpA-Leu- NH₂
- Seq ID 90 Gln-Trp-Ala-Val-Gly-His(3Me)-4-Am,5-MeHpA-Leu- NH₂
- 30 • Seq ID 91 Gln-Trp-Ala-Val-Gly-His-4-Am,5-MeHpA-Leu- NH₂
- Seq ID 101 Gln-Trp-Ala-Val-Gly-His(3Me)- 4-Am-5-MeHpA – 4-amino-5-methylheptanoic acid -Leu- NH₂
- Seq ID 102 Gln-Trp-Ala-Val-NMeGly-His(3Me)- 4-Am-5-MeHpA – 4-amino-5-methylheptanoic acid -Cpa- NH₂