


Description


The present invention relates to novel benzopyrans having pharmacological activity, to processes and intermediates for use in their preparation, to pharmaceutical compositions and their preparation, and to their use in the treatment of mammals.

US-A-4110347 describes compounds having blood pressure lowering activity which are of formula (A'):




(A')

and acid addition salts thereof wherein R_a is a hydrogen atom or a C_{1-6} hydrocarbon group optionally substituted by a hydroxyl or C_{1-6} alkoxy group; R_b is a hydrogen atom or C_{1-6} alkyl group, or NR_aR_b is a 3-8 membered heterocyclic group optionally substituted by one or two methyl groups; R_c is a hydrogen or halogen atom or a C_{1-6} alkyl, C_{2-6} alkenyl, C_{1-6} alkoxy, C_{2-6} alkenoxy, C_{1-6} alkylthio, hydroxyl, amino, C_{1-6} alkylamino, C_{1-6} dialkylamino, nitro, trifluoromethyl, C_{2-7} acylamino, C_{1-6} alkoxy sulphonylamino, carboxyl, nitrile or AOR_g , ASR_g , ASO_2R_g , $ANHR_g$, ANR_gCOR_h , $ANR_gSO_2R_h$ or $ANR_gCO_2R_h$ group wherein A is an alkylene group of 1-4 carbon atoms, R_g is an alkyl group of 1-4 carbon atoms and, R_h is an alkyl group of 1 to 4 carbon atoms; and R_d is a hydrogen or halogen atom or methyl or methoxy, or R_e together with R_d forms a



system; R_e is a hydrogen atom or a C_{1-6} alkyl or phenyl group; and R_f is a hydrogen atom or a C_{1-6} alkyl or phenyl group.


US-A-4251532 describes compounds having useful anti-hypertensive activity, which are of formula (B'):



(B')

wherein R_i is a hydrogen atom or an alkyl group of up to 4 carbon atoms optionally substituted by a chlorine or bromine atom or by a hydroxyl group or by an alkoxy group of up to 4 carbon atoms or by an acyloxy group of up to 4 carbon atoms and R_j is a hydrogen atom or an alkyl group of up to 4 carbon atoms or R_i is joined to R_j so that together with the nitrogen atom to which they are attached they form a 5-, 6- or 7-membered heterocyclic ring which is optionally substituted by methyl; Y is a group COR_k , CO_2R_k , SOR_k , SO_2R_k , $SOOR_k$, SP_2OR_k , $CH(OH)R_k$, $C(R_k)=NOH$, $C(R_k)=NNH_2$, $CONH_2$, $CONR_1R_m$, $SONR_1R_m$ or $SO_2NR_1R_m$ where R_k and R_l are each independently a hydrocarbon group of up to 8 carbon atoms or such a group inertly substituted by a chlorine or bromine atom or by a hydroxyl group or by an alkoxy group of 1-4 carbon atoms or by an acyloxy group of up to 4 carbon atoms or by 3 fluorine atoms attached to the same carbon atom and R_m is a hydrogen atom or an alkyl group of up to 4 carbon atoms; and salts thereof and O-acyl derivatives thereof wherein the O-acyl moiety contains up to 18 carbon atoms.


EP—A—9912 describes anti-hypertensive compounds of formula (C'):



(C')


wherein the pyrrolidino and OR_n is a hydrogen atom, an alkyl group of 1 to 3 carbon atoms or an acyl group of 1 to 8 carbon atoms; or a pharmaceutically acceptable acid addition salt thereof.

EP—A—28449 describes compounds having blood pressure lowering activity, with low levels of unwanted cardiac effects, which compounds are of formula (D'):



(D')

R_o is a hydrogen atom or a lower alkyl group;
 R_p is a hydrogen atom or a lower alkyl group;
 R_q is a hydrogen atom or a lower alkyl group;
 R_r is a hydrogen atom or a lower alkyl group;
 R_s is a lower alkyl or a substituted alkyl group; or
 R_r and R_s are joined so that together with the nitrogen atom to which they are attached they form a 5-
 6- or 7-membered ring optionally containing an oxygen or sulphur atom;
 R_t is an electron withdrawing group;
 R_u is an electron donating group; and the NR_s, R_s and OR_q moieties are trans.
 EP—A—28064 describes compounds having blood pressure lowering activity, with low levels of
 unwanted cardiac effects, which compounds are of formula (E'):



(E')


and salts and pro-drugs thereof, wherein:

R_o, R_p, R_q, R_r and R_s are as defined for formula (D'), and R_v is an electron donating group and R_w is an
 electron withdrawing group; and the NR_s, R_s and OR_q moieties are trans.

British Patent No.'s 1511187, 1548221 and 1548222 also describe classes of benzopyrans having blood pressure lowering activity.

It has now been found that a further class of benzopyrans have blood pressure lowering activity. Such compounds are characterised by the presence of an oxo group in the nitrogen-containing ring which
 substitutes the benzopyran in the 4-position.

Accordingly, the present invention provides a compound of formula (I):



(I)

wherein:

One of R_1 and R_2 is hydrogen and the other is selected from the class of alkylcarbonyl, alkoxycarbonyl, alkylcarbonyloxy, alkylhydroxymethyl, nitro, cyano, chloro, trifluoromethyl, alkylsulphinyl, alkylsulphonyl, alkoxysulphinyl, alkoxysulphonyl, alkoxycarbonylamino or aminosulphinyl, aminosulphonyl or aminocarbonyl, the amino moiety being optionally substituted by one or two alkyl groups, or alkylsulphinylamino, alkylsulphonylamino, alkoxysulphinylamino or alkoxysulphonylamino or ethylenyl terminally substituted by alkylcarbonyl, nitro or cyano, or $-C(alkyl)NOH$ or $-C(alkyl)NNH_2$, the alkyl groups or alkyl moieties of alkyl-containing groups having from 1 to 6 carbon atoms;

one of R_3 and R_4 is hydrogen or alkyl having from 1 to 4 carbon atoms and the other is alkyl having from 1 to 4 carbon atoms, or R_3 and R_4 together with the carbon atom to which they are attached are spiroalkyl having from 3 to 6 atoms;

R_5 is hydrogen, alkyl having from 1 to 3 carbon atoms or acyl having from 1 to 8 carbon atoms; and n is 1 or 2; the lactam group being trans to the OR_5 group.


The other of R_1 and R_2 , when one of them is hydrogen, is preferably selected from the class of alkylcarbonyl, alkoxycarbonyl, alkylcarbonyloxy, alkylhydroxymethyl, nitro or cyano.

The alkyl groups or alkyl moieties of alkyl-containing groups, in respect of the other of R_1 and R_2 , are preferably methyl or ethyl.

R_3 and R_4 are preferably both alkyl having from 1 to 4 carbon atoms. In particular they are both methyl or ethyl, preferably both methyl.

When R_5 is alkyl, preferred examples thereof include methyl, ethyl and n-propyl, of which methyl is most preferred. When R_5 is acyl, a preferred class is unsubstituted carboxylic acyl, such as aliphatic acyl or benzoyl. R_5 however is preferably hydrogen.

Within formula (I) is a sub-group of preferred compounds of formula (II):



(II)

wherein one of R_1 and R_2 is hydrogen and the other is selected from the class of alkylcarbonyl, alkoxycarbonyl, alkylcarbonyloxy, alkylhydroxymethyl, nitro or cyano, the alkyl groups or alkyl moieties of alkyl-containing group being methyl or ethyl, and n is 1 or 2.

Compounds of formula (II), wherein one of R_1 and R_2 is hydrogen and the other is nitro or cyano, are greatly preferred. Additionally, compounds of formula (II), wherein R_2 is hydrogen and R_1 is one of the substituents as defined hereinbefore, are preferred. Consequently the most preferred compounds are those wherein R_1 is nitro or cyano and R_2 is hydrogen.


The compounds of formula (I) and (II) cover both a piperidone substituent (when $n = 2$) and a pyrrolidone substituent (when $n = 1$).

It is preferred that the compounds of formula (I) and (II) are in substantially pure form.

The present invention extends to the compounds of formulae (I) and (II) whenever prepared synthetically.

The compounds of formulae (I) and (II) have asymmetric centres and therefore exist in optically active forms. The present invention extends to all such forms individually and to mixtures of them.

The present invention also provides a process for the preparation of a compound of formula (I), which comprises cyclising a compound of formula (III), or when R_5 is hydrogen a metal salt thereof:



wherein one of R_1' and R_2' is hydrogen or a group or atom convertible into hydrogen and the other is one of the class of substituents as defined hereinbefore for the other of R_1 and R_2 or a group or atom convertible thereto, R_3 to R_5 and n are as defined hereinbefore, L_1 is a leaving group; and wherein the substituted amino group is trans to the OR_5 group; in the case when one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group or atom convertible into one of the class of substituents as defined hereinbefore for the other of R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_5 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.

The leaving group L_1 is a group that is displaceable by a secondary amino nucleophile. Preferred examples of such groups include hydroxy and, in particular, C_{1-4} alkoxy, such as ethoxy.

The cyclisation is normally carried out by heating the compound of formula (III) under reflux in an inert solvent, such as xylene or toluene.

When a metal salt of formula (III) is used, the sodium salt is preferred. However, it is even more preferred not to use a metal salt at all, especially as any elimination side reactions are thereby avoided.

Conversions of an aromatic group or atom into one of the class of substituents as defined hereinbefore for the other of R_1 and R_2 , when one of them is hydrogen, are generally known. For example, it is preferred, when carrying out the reaction, to protect any unsubstituted terminal amino moieties, such as when the other of R_1 and R_2 is aminosulphinyl, aminosulphonyl or aminocarbonyl, with a protecting agent. Examples of protecting agents include acyl groups, such as acetyl. Protection of the unsubstituted terminal amino moiety is carried out by reacting a compound of formula (III), wherein one of R_1' and R_2' is hydrogen and the other is aminosulphinyl, aminosulphonyl or aminocarbonyl, with, for example, an acyl chloride. Removal of the acyl protecting agent is carried out by base hydrolysis after the cyclising reaction.


Conversions of an aromatic group or atom into one of the class of substituents as defined hereinbefore for the other R_1 and R_2 , when one of them is a group or atom convertible into hydrogen, are generally known as well. For example, a hydrogen atom may be replaced by a nitro group by nitrating in a known manner a compound of formula (III); wherein one of R_1' and R_2' is hydrogen and the other is acetamido, followed by hydrolysing the compound, converting the resulting amine into a diazonium salt, and finally decomposing it, leaving a compound of formula (I), wherein one of R_1 and R_2 is hydrogen and the other is nitro.

It is however preferred that any conversions are carried out at an earlier stage as mentioned hereinafter.

Preferably, the alkylating agent is an alkyl iodide, the reaction being carried out in an inert solvent, such as toluene, in the presence of a base, such as potassium *t*-butoxide.


Preferably, the acylating agent is a carboxylic acid or a derivative thereof, such as an anhydride, the reaction being carried out in a non-hydroxylic solvent in the presence of a condensation promoting agent, such as dicyclohexyl carbodiimide.

Compounds of formula (III) can be prepared by reacting a compound of formula (IV):



(IV)

wherein R_1' and R_2' and R_3 and R_4 are as defined hereinbefore, with a compound of formula (V):



(V)


wherein n and L_1 are as defined hereinbefore.

The reaction is normally carried out in a solvent at low, medium or high temperature. The solvent may be an alcohol, such as methanol or ethanol.

When L_1 is hydroxy the reaction proceeds well if carried out in refluxing ethanol in the presence of aqueous sodium carbonate. When L_1 is C_{1-4} alkoxy, the reaction is preferably carried out in the presence of sodium hydroxide in ethanol.

Under some conditions, the compound of formula (III) spontaneously cyclises to form a compound of formula (I).


Compounds of formula (IV) can be prepared, preferably *in situ*, by reacting a compound of formula (VI):



(VI)


wherein R_1' and R_2' and R_3 and R_4 are as defined hereinbefore and the hydroxy group is trans to the bromo atom, with a base, such as potassium hydroxide, in ether or aqueous dioxan.

Alternatively, compounds of formula (III) can be prepared by reacting a compound of formula (VII):



(VII)

wherein R_1' , R_2' , R_3 and R_4 are as defined hereinbefore, and the amino group is trans to the hydroxy group, with a compound of formula (VIII):




(VIII)

wherein n and L_1 are as defined hereinbefore and L_2 is a leaving group.

The leaving group, L_2 , is a group that is displaceable by a primary amino nucleophile. Preferred examples of such groups include halo, such as chloro and bromo.

Compounds of formula (VII) can be prepared by a reaction of a compound of formula (IV) with ethanolic ammonium hydroxide solution. Alternatively, they can be prepared by reduction with zinc and hydrochloric acid of a compound of formula (IX):




(IX)

wherein R_1' , R_2' , R_3 and R_4 are as defined hereinbefore and wherein the azide group is trans to the hydroxy group.

Compounds of formulas (IX) can in turn be prepared from a compound of formula (IV) by reaction with sodium azide in the presence of boric acid in for example dimethylformamide.

The present invention provides another process for the preparation of a compound of formula (I), which comprises oxidising a compound of formula (X), or when R_5 is hydrogen a metal salt thereof:




(X)

wherein R_1' , R_2' , R_3 to R_5 and n are as defined hereinbefore, and wherein the nitrogen containing ring is trans to the OR_5 group; in the case when one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group or atom convertible into one of the class of substituents as defined hereinbefore for the other or R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_5 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.

The oxidation is preferably carried out in a solvent such as aqueous methanol with a metal periodate such as potassium periodate.

Compounds of formula (X) can be prepared by cyclising in the presence of an acid a compound of formula (XI):



(XI)

wherein R is methoxy or ethoxy and R_1' , R_2' , R_3 to R_5 and n are as defined hereinbefore and wherein the substituted amino group is trans to the OR_5 group.


Formulae X and XI have been disclosed in EP—A—93534 which was published after the priority date of the present application but which claim a priority with respect to these formulae which is earlier than that of the application. However, this arises only in respect of the Designated Contracting States CH, DE, FR, GB, IT and NL.

- 5 Compounds of the formula (XI) can in turn be prepared by reacting a compound of formula (IV) with a compound of formula (XII):



- 10 wherein n and R_6 are as defined hereinbefore.

The present invention provides a further process for the preparation of a compound of formula (I), which comprises reacting a compound of formula (IV) with an anion of formula (XIII):




wherein n is as defined hereinbefore; in the case when one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group or atom convertible into one of the class of substituents as defined hereinbefore for the other of R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_6 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.

The reaction is preferably carried out in a solvent such as dimethyl sulphoxide in the presence of a base, such as sodium hydride.

- 30 The compound of formula (IV) can be prepared *in situ* from the corresponding compound of formula (VI). In such circumstances, it is advantageous not to add the lactam of formula (XIII) until sufficient time has elapsed for the epoxide of formula (IV) to be produced.

The present invention provides yet another process for the preparation of a compound of formula (I), which comprises cyclising a compound of formula (XIV):



- wherein R_1' , R_2' , R_3 to R_5 and n are as defined hereinbefore and L_3 is a leaving group, and wherein the substituted amino group is trans to the OR_5 group; in the case when one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group or atom convertible into one of the class of substituents as defined hereinbefore for the other of R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_6 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.

- 50 The leaving group L_3 is a group that is displaceable by a secondary amino nucleophile adjacent to a carbonyl function. A preferred example is chloro.

The cyclisation reaction is preferably carried out in a solvent such as dimethylformamide in the presence of a base, such as sodium hydride.

Compounds of formula (XIV) can be prepared by reacting a compound of formula (VII) with a compound of formula (XV):




- wherein L_3 and n are as defined hereinbefore and L_4 is a leaving group; and optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.
- 65

The leaving group, L_4 , is a group that, when adjacent a carbonyl function, is displaceable by a primary amino nucleophile.

The reaction is preferably carried out in a solvent, such as chloroform or methylene chloride, in the presence of aqueous base, such as aqueous sodium hydroxide.

In the reactions with the epoxide of formula (IV), the trans isomer is specifically formed.

Compounds of formula (VI) can be prepared in accordance with known processes, for example the process described in the aforementioned U.S. patents and European patent publications. Schematically, such process can be depicted thus:



(a) Room temperature; NaOH/40% benzyltrimethylammonium hydroxide in methanol;

(b) Heat in o-dichlorobenzene;

(c) N-bromosuccinimide/dimethylsulphoxide/water;

(d) Bromine in carbon tetrachloride; and

(e) Acetone/water.

The above process can produce mixtures of compounds during reaction (b) owing to the two sites available for ring formation. It is therefore advisable to remove any of the undesired compound by, for example, chromatography, before reaction (c) or (d).

Instead of carrying out the conversion of a group or atom into hydrogen or group or atom into one of the class of substituents defined for the other of R_1 and R_2 after cyclising a compound of formula (III) or (XIV), or after oxidising a compound of formula (X), or after reacting a compound of formula (IV) with an anion of formula (XIII), it is greatly preferred that any such conversions are carried out at an earlier stage, preferably on the chromene produced after reaction (b) above. In other words, it is preferred that, for the processes of the invention R_1' and R_2' are R_1 and R_2 respectively.

As mentioned previously, the compounds of formula (I) exist in optically active forms, and the processes of the present invention produce mixtures of such forms. The individual isomers may be separated one from another by chromatography using a chiral phase. Alternatively, an asymmetric synthesis would offer a route to the individual form.

It is preferred that the compounds of formula (I) are isolated in substantially pure form.

The intermediates of formula (III), (IX), (X), (XI) and (XIV) are novel and constitute part of the invention as set forth in the different sets of claims.

As mentioned previously, the compounds of formula (I), and especially those of formula (II), have been found to have blood-pressure lowering activity. They are therefore potentially useful as a pharmaceutical in the treatment of hypertension.

The present invention accordingly provides a pharmaceutical composition which comprises a compound of this invention and a pharmaceutically acceptable carrier. In particular, the present invention provides an antihypertensive pharmaceutical composition which comprises an anti-hypertensive effective amount of a compound of this invention and a pharmaceutically acceptable carrier.

The composition are preferably adapted for oral administration. However, they may be adapted for other modes of administration, for example parenteral administration for patients suffering from heart failure.

In order to obtain consistency of administration it is preferred that a composition of the invention is in the form of a unit-dose. Suitable unit dose forms include tablets, capsules and powders in sachets or vials. Such unit dose forms may contain from 1 to 100 mg of a compound of the invention and more usually from 2 to 50 mg, for example 5 to 25 mg such as 6, 10, 15 or 20 mg. Such compositions may be administered from 1 to 6 times a day, more usually from 2 to 4 times a day, in a manner such that the daily dose is from 5 to 200 mg for a 70 kg human adult and more particularly from 10 to 100 mg.

The compositions of the invention may be formulated with conventional excipients, such as a filler, a disintegrating agent, a binder, a lubricant, a flavouring agent and the like. They are formulated in conventional manner, for example in a manner similar to that used for known anti-hypertensive agents, diuretics and B-blocking agents.

The present invention further provides a compound of formula (I) and especially of formula (II) for use in the treatment of hypertension.

The present invention yet further provides a method of treating hypertension in mammals including man, which comprises administering to the suffering mammal an anti-hypertensive effective amount of a compound of formula (I) or a pharmaceutical composition of the invention.

The following descriptions relate to the preparation of intermediates and the following examples relate to the preparation of a compound of formula (I).

Description 1

6-Cyano-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo[b]pyran

4-Cyanophenol (19.60 g), sodium hydroxide (9.90 g), 40% benzyltrimethylammonium hydroxide in methanol (34.50 g) and 3-methyl-3-chlorobutylene (25.50 g) were stirred in water (150 ml) and dichloromethane (150 ml) for 5.5 days at room temperature. After separation of the layers, the aqueous layer was extracted twice with chloroform, and the combined organic phase evaporated leaving a gum which was taken up in ether and washed three times with 10% sodium hydroxide solution and with water before drying over magnesium sulphate. Removal of drying agent and solvent gave a viscous liquid having absorptions in the IR (film) at 2100, 2220, 3290 cm^{-1} . This liquid (20.91 g) was heated in o-dichlorobenzene (40 ml) at reflux temperature for 1.5 hours under nitrogen. After distillation of the solvent the fraction boiling at 110–114°/0.02 mmHg (16.57 g) was collected, which on standing formed a low melting solid, having an IR absorption at 2230 cm^{-1} . (See M. Harfenist and E. Thom, *J. Org. Chem.*, 841 (1972) who quote m.p. 36–37°).

Addition to this 6-cyanochromene (16.50 g) dissolved in dimethyl sulphoxide (150 ml) containing water (3.24 ml) of N-bromosuccinimide (31.90 g) with vigorous stirring and colling, followed by dilution with water and extraction *via* ethyl acetate gave a mixture which was boiled in acetone (300 ml) and water (100 ml) for 5 hours to hydrolyse the small amount of 3,4-dibromide present. Evaporation of solvents gave 6-cyano-trans-3-bromo-3,4-dihydro-2,2-dimethyl-2H-benzo[b]pyran-4-ol as white crystals (24.37 g). A small sample had m.p. 128–128.5° from 60–80° petroleum ether, nmr (CDCl_3) δ 1.43 (3H), 1.62 (3H), 7.48 (1H, exchangeable), 4.07 (1H, d, J = 9), 4.87 (1H, d, J = 9), 6.80 (1H, d, J = 8), 7.43 (1H, q, J = 8, 2), 7.78 (1H, d, J = 2). Analysis calculated for $\text{C}_{12}\text{H}_{12}\text{NO}_2\text{Br}$: C, 51.07; H, 4.26; N, 4.96; Br, 28.37. Found: C, 50.95; H, 4.38; N, 5.03; Br, 28.39%.

The bromohydrin (24.30 g) was stirred with sodium hydroxide pellets (5.00 g) in water (250 ml) and dioxan (200 ml) for 3 hours at room temperature. The solvents were removed by distillation under high vacuum and the residue taken up in ether and washed with water and brine before drying over magnesium sulphate. Removal of drying agent and solvent and gave crude 6-cyano-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo[b]pyran (16.02 g) as a gum, having an absorption at 2230 cm^{-1} in the IR and Nmr (CCl_4) δ 1.26 (3H), 1.54 (3H), 3.40 and 3.80 (each 1H, d, J = 4), 6.77 (1H, d, J = 8), 7.43 (1H, q, J = 8, 2), 7.58 (1H, d, J = 2).

Description 2

6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-amino-2H-benzo[b]pyran-3-ol

The title compound was prepared by stirring 6-cyano-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo[b]pyran in ethanolic ammonium hydroxide solution at room temperature until thin layer chromatography showed consumption of the starting epoxide.

Description 3

6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(1-keto-4-chlorobutyrylamino)-2H-benzo[b]pyran-3-ol

The amino chromanol (1.40 g), as obtained in Description 2, was stirred in chloroform (20 ml) and water (10 ml) containing sodium hydroxide pellets (0.26 g) at room temperature. 4-Chlorobutyryl chloride

(0.72 ml) was added and the reaction stirred vigorously for 0.5 hours. Separation of the layers and washing the organic layer with water, then brine, drying over magnesium sulphate, filtration and evaporation gave the title compound as a pale yellow solid.

5

Description 4

6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(4,4-diethoxybutylamino)-2H-benzo[b]pyran-3-ol

6-Cyano-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo[b]pyran (200 mg) and 4-aminobutyraldehyde diethylacetal (200 mg) were heated to 100°C for 1.5 hours, a clear yellow solution forming during this time. After cooling, dilution with ether, and washing successively with water and brine, drying over sodium sulphate and evaporation, the aminoacetal was obtained as a pale yellow oil (291 mg).

10

Description 5

6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(2-hydroxy-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol

The oily acetal, as obtained in Description 4, was dissolved in dioxan (2 ml) and treated with 2.5M HCl (1 ml). After 30 minutes the reaction was diluted with ether and neutralised with sodium carbonate solution.

15

The two phases were separated, the aqueous layer further extracted with ether and the combined extracts washed with water and brine and dried over sodium sulphate. The organic phase was filtered and applied to Kieselgel 60 (10 g) and diluted with ethyl acetate-heptane-triethylamine (10:20:2). Three fractions were obtained (total 128 mg) containing the title compound. TLC (silica gel; ethyl acetate-heptane-triethylamine (10:20:2) showed the presence of varying amounts of the two positional isomers in each fraction.

20

IR (KBr disc) 3450, 2230 cm^{-1} for all three fractions.

Mass spectrum (isobutane and ammonium C.I.) showed m/z 271 ($\text{MH}^+ - \text{H}_2\text{O}$) for all three fractions.

25

Description 6

6-Nitro-3,4-dihydro-2-methyl-3,4-epoxy-2H-benzo[b]pyran

p-Nitrophenol (49.3 g), 3-bromobut-1-yne (39.0 g), potassium carbonate (66 g) and potassium iodide (3.1 g) were heated and stirred under nitrogen for 20 hours. The mixture was cooled, filtered, and evaporated and the residue taken up in ether and washed with sodium hydroxide solution (10%) before drying over magnesium sulphate. Filtration and evaporation gave the phenoxybutyne as a yellow oil (39.03 g). This crude phenoxybutyne (30 g) was heated in o-dichlorobenzene (1 litre) for 24 hours under nitrogen. Removal of solvent, and recrystallisation from 60–80° petroleum ether gave 2-methyl-6-nitrochromene as an oily solid. The nitrochromene (6.10 g) dissolved in dimethyl sulphoxide (50 ml) containing water (1.12 ml) was treated with N-bromosuccinimide (11.40 g) in one portion with vigorous stirring. After 0.5 hours the reaction mixture was poured into water (500 ml) and extracted with ethyl acetate to yield the bromohydrin (7.4 g) as a sticky solid. Recrystallisation from ethyl acetate-petroleum ether gave a sample of m.p. 159°. The bromohydrin (2.27 g) was stirred with potassium hydroxide pellets (2.2 g) in ether (500 ml) for 48 hours. Filtration gave the 3,4-epoxy-2-methylchroman as a yellow crystalline solid (1.05 g).

30

35

40

Description 7

6-Nitro-3,4-dihydro-2-methyl-trans-4-(ethoxycarbonylpropylamino)-2H-benzo[b]pyran-3-ol

The epoxide of Description 6 (1.03 g) was boiled with ethyl 4-aminobutyrate hydrochloride (0.84 g) and sodium hydroxide pellets (0.20 g) in ethanol (50 ml) for 10 hours. Filtration and evaporation and chromatographic purification gave a gum (1.10 g) which was dissolved in 2N hydrochloric acid and extracted three times with ethyl acetate. Basification of the aqueous phase and extraction with ethyl acetate gave a crude ester which was used as such in Example 9.

45

Description 8

7-Nitro-3,4-dihydro-2,2-dimethyl-trans-4-(3-carethoxypropylamino)-2H-benzo[b]pyran-3-ol

7-Nitro-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo[b]pyran (0.48 g, the preparation of which is described in Example 3 of U.K. Patent No. 1,548,221), ethyl 4-aminobutyrate hydrochloride (0.34 g) and sodium hydroxide pellets (0.08 g) were refluxed in ethanol (50 ml) for 12 hours. Filtration and evaporation and chromatography on a chromatotron (2 mm silica gel HF₂₅₄, gradient elution with pentane-ethyl acetate) gave recovered epoxide (0.20 g) and trans-4-(3-carbethoxypropylamino)-7-nitro-2,2-dimethyl-2H-benzo[b]pyran-3-ol (0.21 g) as a gum, which was used in Example 10.

50

55

Description 9

6-Chloro-3,4-dihydro-2,2-dimethyl-trans-3-bromo-4-hydroxy-2H-benzo[b]pyran

The title compound was prepared analogously to the preparation of the 3-bromo-4-hydroxy compound of Description 1 giving a crude crystalline solid.

60

NMR (CDCl_3): δ 1.35 (3H, s); 1.53 (3H, s); 3.22 (1H, m); 4.00 (1H, d, $J = 9\text{Hz}$); 4.77 (1H, d, $J = 9\text{Hz}$); 6.51 (1H, d, $J = 8\text{Hz}$); 7.03 (1H, q, $J = 8.2\text{Hz}$); 7.30 (1H, narrow m).


65

Description 10

6-Chloro-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo[b]pyran

The crude crystalline solid (10.27 g) of Description 9 was dissolved in dimethyl sulphoxide (50 ml) and treated with sodium hydride (1.06 g, 80% dispersion on oil) over a period of an hour. The resulting material was used as such immediately in Example 12.

Example 1

6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidiny)-2H-benzo-[b]-pyran-3-ol

(E1)

6-Cyano-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo-[b]-pyran (0.50 g), as obtained in Description 1, 4-aminobutyric acid (1.25 g) and sodium bicarbonate (1.00 g) were refluxed in ethanol (15 cc) and distilled water (2.5 cc) for 10 hours. The reaction was filtered and evaporated and the residue chromatographed on 25 g Kieselgel 60. Elution with MeOH-chloroform (1:3) gave 132 mg of the most polar product. This was refluxed in toluene (10 cc) for 2 hours, cooled and the solvent evaporated. The residue was chromatographed on 5 g Kieselgel 60 and eluted with MeOH-chloroform (1:3) to give the title compound as a white solid (90 mg), m.p. 230—231°.

IR (KBr disc): 3260, 2220, 1651 cm^{-1} ;

NMR (CDCl_3): δ 1.28 (3H); 1.55 (3H); 2.11 (2H, m); 2.57 (2H, m); 3.22 (3H, 1 exchangeable H, broad m); 3.64 (1H, d, $J = 10$); 5.26 (1H, d, $J = 10$); 6.87 (1H, d, $J = 9$); 7.24 (1H, narrow m); 7.45 (1H, 1, $J = 9, 2$).

Analysis calculated for $\text{C}_{18}\text{H}_{18}\text{N}_2\text{O}_3$: C, 67.12; H, 6.34; N, 9.78%

Found: C, 66.83; H, 6.17; N, 9.50%

Example 2

6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidiny)-2H-benzo-[b]-pyran-3-ol of formula (E1)

6-Cyano-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo-[b]-pyran (1.00 g), as obtained in Description 1, ethyl 4-aminobutyrate hydrochloride (0.84 g), ethanol (50 ml) and sodium hydroxide pellets (0.20 g) were stirred at room temperature for 8 days, then at 40° for 3 hours. After cooling and evaporation the residue was taken up in ethyl acetate and filtered. Evaporation of the filtrate gave a gum (1.46 g) which was chromatographed using a chromatotron (2 mm silica gel HF₂₅₄ plate; 2 runs; solvent flow rate 6 ml/min.). Elution with 2% methanol-chloroform mixture gave starting epoxide (0.23 g) followed by a more polar ester fraction (0.64 g), and a mixture (0.15 g) which on further chromatography under identical conditions gave 6-cyano-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidiny)-2H-benzo-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidiny)-2H-benzo[b]pyran-3-ol having an identical NMR spectrum to that obtained in Example 1.

A portion of the ester fraction (150 mg) as dissolved in ether containing a little ethanol and treated with anhydrous ethanolic HCl. The precipitate was collected and triturated with ether to give trans-4-(3-carbethoxy-propylamino)-6-cyano-2,2-dimethyl-2H-benzo[b]pyran-3-ol hydrochloride (138 mg) of m.p. 198—200°.

NMR (CD_3OD): 1.23 (s, 3H) overlapped with 1.26 (t, $J = 8, 8$ 3H); 1.58 (s, 3H); 2.19 (m, 2H); 2.53 (m, 2H); 2.85—3.45 (irreg. m, 2H); 4.02 (d, $J = 10$, 1H) overlapped with 4.16 (q, $J = 8, 8, 8$ 2H) and 3.75—4.65 (m, 3H, exchangeable); 4.53 (3, $J = 10$, 1H); 7.00 (d, $J = 9$, 1H); 7.60 (q, $J = 9, 2$, 1H); 8.15 (d, $\alpha = 2$, 1H);

Analysis calculated for $\text{C}_{18}\text{H}_{23}\text{N}_2\text{O}_4$: C, 58.61; H, 6.83; N, 7.59%

Found: C, 58.55; H, 6.80; N, 7.29%

The remainder of the ester fraction was heated under reflux in xylene (50 ml) for 7.25 hours. The solution was cooled and filtration gave 6-cyano-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidiny)-2H-benzo[b]pyran-3-ol (425 mg) as crystals of m.p. 226° having an identical NMR spectrum and t.l.c. characteristics as the compound of Example 1.

Example 3

6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidiny)-2H-benzo(b)pyran-3-ol of formula (E1)

6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(1-keto-4-chlorobutylamino)-2H-benzo(b)pyran-3-ol (0.76 g), as obtained in Description 3, in dry tetrahydrofuran (10 ml) was added to a suspension of sodium hydride

(0.15 g) in tetrahydrofuran (20 ml) and the reaction stirred under nitrogen for 3 hours. Addition of water and extraction via ethyl acetate gave 540 mg of the title compound having an identical NMR spectrum and tlc characteristics as the compound of Example 1.


Example 4

6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol of formula (E1)

A solution of 6-Cyano-3,4-dihydro-2,2-dimethyl-trans-3-bromo-4-hydroxy-2H-benzo[b]pyran (4g, 14.2 mM) in dimethylsulphoxide (20 ml) was stirred and sodium hydride (60% dispersion in oil, 0.6 g, 15 mM) added. The suspension was stirred for 1 hour when a solution of 6-cyano-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo[b]pyran resulted. 2-Pyrrolidone (1.8 g, 21 mM) and further sodium hydride (0.8 g, 21 mM) were introduced and the mixture stirred at room temperature for an additional 16 hours. Water (40 ml) was slowly added to the mixture to induce crystallisation of the product after which it was cooled in ice and filtered under suction. Crystallisation from ethanol (20 ml) gave the title compound as a cream coloured solid in 60% yield. Recrystallisation from ethyl acetate afforded the pure product as needles, m.p. 226.5—227.5° having an nmr spectrum and t.l.c. characteristics identical to those of the compound of Example 1.

Example 5

6-Carbomethoxy-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol (E5)




(E5)

6-Carbomethoxy-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo[b]pyran (2.42 g, the preparation of which is described in Example 4 of U.K. Patent No. 1,511,187) and 2-pyrrolidone (0.88 g), were stirred in dimethyl sulphoxide (40 ml) under nitrogen room temperature. Sodium hydride (0.31 g, 81% dispersion in mineral oil) was added during 5 mins. and the reaction stirred for a further 6 hours. Addition of water, extraction with ethyl acetate, drying of the organic phase with magnesium sulphate, filtration, evaporation and recrystallisation from ethyl acetate-pentane gave 6-Carbomethoxy-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol of m.p. 190—192°.

NMR (CDCl₃ soln + 1 drop D₂O): δ 1.30 (3H, s); 1.55 (3H, s); 1.75—2.30 (2H, m); 2.60 (2H, irregular t, J = 8Hz); 2.80—3.40 (2H, m); 3.74 (1H, d, J = 10 Hz); 3.87 (3H, s); 5.33 (1H, d, J = 10Hz); 6.86 (1H, d, J = 8Hz); 7.67 (1H, narrow m); 7.88 (1H, q, J = 8, 2Hz).

Example 6

6-Carbomethoxy-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-piperidinyl)-2H-benzo[b]pyran-3-ol (E6)




(E6)

In a similar manner to that described in Example 5 employing 2-piperidone in place of 2-pyrrolidone, 6-carbomethoxy-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-piperidinyl)-2H-benzo[b]pyran-3-ol was prepared as crystals of m.p. 249—250° from ethyl acetate.

NMR (CDCl₃ soln + 1 drop D₂O): δ 1.27 (3H, s); 1.52 (3H, s); 1.65—2.05 (4H, m); 2.60 (2H, irregular t, J = 7Hz); 2.85—3.15 (2H, m); 3.77 (1H, d, J = 10Hz); 3.88 (3H, s); 5.94 (1H, d, J = 10Hz); 6.86 (1H, d, J = 8Hz); 7.71 (1H, narrow m); 7.87 (1H, q, J = 8, 2Hz).

Example 7

6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-piperidiny)-2H-benzo[b]pyran-3-ol (E7)



(E7)

In a similar manner to that described in Example 5 employing 2-piperidone in place of 2-pyrrolidone and 6-cyano-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo[b]pyran in place of the corresponding 6-carbomethoxy compound, 6-cyano-3,4-dihydro-2,2-dimethyl *trans*-4-(2-oxo-1-piperidiny)-2H-benzo[b]pyran-3-ol was prepared, as crystals of m.p. 155° from ethyl acetate.

IR (KBr disc) 3480, 2212, 1612 cm^{-1} ;

NMR (CDCl_3 soln): δ 1.25 (3H, s); 1.50 (3H, s); 1.63—2.10 (4H, m); 2.36—2.76 (2H, m); 3.72 (1H, d, $J = 10\text{Hz}$); 3.90—4.20 (1H, exchangeable, m); 5.72 (1H, d, $J = 10\text{Hz}$); 6.76 (1H, d, $J = 8\text{Hz}$); 7.17 (1H, m, narrow); 7.42 (1H, q, $J = 8, 2\text{Hz}$).


Example 8

6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidiny)-2H-benzo[b]pyran-3-ol of formula (E1)

6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(2-hydroxy-1-pyrrolidiny)-2H-benzo[b]pyran-3-ol (5 mg), as obtained in Description 5, dissolved in methanol-water (1 ml) was treated with an excess of sodium periodate with stirring during 15 hours at room temperature. Evaporation of solvents and extraction by ethyl acetate gave material having identical thin layer characteristics when applied to silica gel plates developed in either chloroform-methanol (15:1) or heptane-ethyl acetate-triethylamine and infra red spectrum to the compound of Example 1.

Example 9

6-Nitro-3,4-dihydro-2-methyl-trans-4-(2-oxo-1-pyrrolidiny)-2H-benzo[b]pyran-3-ol (E9)



(E9)


The crude ester (0.27 g) of Description 7 was heated under reflux in an atmosphere of nitrogen for 72 hours. Evaporation of solvent and trituration with ethanol and recrystallisation from ethyl acetate gave 6-nitro-3,4-dihydro-2-methyl-trans-4-(2-oxo-1-pyrrolidiny)-2H-benzo[b]pyran-3-ol as a pale yellow solid (104 mg) of m.p. 238—242°.

IR (KBr disc) 1650 cm^{-1} ;

NMR ($\text{DMSO}-d_6$): δ 1.47 (3H, d, $J = 7\text{Hz}$); 1.80—2.23 (2H, m); 2.23—4.00 (5H, series of m); 4.25 (1H, q, $J = 10, 7\text{Hz}$); 5.14 (1H, d, $J = 10\text{Hz}$); 7.02 (1H, d, $J = 9\text{Hz}$); 7.68 (1H, narrow m); 8.06 (1H, q, $J = 3\text{Hz}$).

Example 10

7-Nitro-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol (E10)




(E10)

The gum of Description 8 was heated under reflux in xylene (30 ml) under nitrogen for 3 days. Evaporation of solvent gave a complex mixture which was purified on a chromatotron (conditions as in Description 8) to give 7-nitro-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol (0.04 g) after recrystallisation from ethyl acetate m.p. 211—211.5°C.

NMR (CDCl₃): δ 1.30 (3H, s); 1.55 (3H, s); 1.80—3.60 (6H, series of m); 3.80 (1H, d, J = 10Hz); 5.30 (1H, d, J = 10Hz); 7.00—7.85 (3H, m).

Example 11

6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-yl acetate (E11)



(E11)


6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol (0.5 g), acetic anhydride (10 ml) and pyridine (0.2 ml) were refluxed for 24 hours. After cooling the reaction mixture was poured onto ice and extracted with ethyl acetate. The organic layer was washed several times with water, then with NaHCO₃ solution and brine before drying over magnesium sulphate. Filtration and evaporation and chromatography of the resulting gum using a chromatotron (2 mm silica gel HF₂₅₄; elution with ethyl acetate), gave a solid (0.35 g) which was recrystallised from ethyl acetate-pentane m.p. 152-153°, as the title compound.

IR (KBr disc) 2225, 1745, 1695 cm⁻¹;

NMR (CDCl₃): δ 1.35 (3H, s); 1.42 (3H, s); 2.10 (3H, s) overlapped by 1.80—3.40 (6H, m); 5.08 (1H, d, J = 10Hz); 5.45 (1H, d, J = 10Hz); 6.85 (1H, d, J = 8Hz); 7.22 (1H, d, J = 2Hz); 7.40 (1H, q, J = 8, 2Hz).

Example 12

6-Chloro-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol (E12)




(E12)

2-Pyrrolidone (4.5 g) and sodium hydride (1.59 g) were added to the material of Description 10, and the

mixture stirred for 20 hours. Cautious addition of water and filtration of the resulting solid, followed by two crystallisations from ethyl acetate gave the title compound, m.p. 202—203°.

Example 13

6-Acetyl-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol (E13)




(E13)

6-Acetyl-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo[b]pyran (0.33 g, prepared as described in Example 1 of U.K. Patent No. 1,511,187), and 2-pyrrolidone (0.15 g), were stirred in dimethylsulphoxide (25 ml) under nitrogen at room temperature. Sodium hydride (0.05 g, 80%) was added during 2 mins and the reaction stirred for a further 22 hours. Addition of water, extraction with ethyl acetate, drying of the organic layer with magnesium sulphate, filtration, evaporation and recrystallisation from ethyl acetate gave the title compound (0.04 g) of m.p. 218—219°.

NMR (CDCl₃): δ (3H, s); 1.55 (3H, s); 1.85—2.25 (2H, m); 2.55 (3H, s) overlapped by 2.45—2.75 (2H, m); 2.80—3.45 (3H, m); 3.75 (1H, d, J = 10Hz); 5.36 (1H, d, J = 10Hz); 6.96 (1H, d, J = 8Hz); 7.63 (1H, narrow m); 7.83 (1H, q, J = 8, 2Hz).

Example 14

6-Acetyl-3,4-dihydro-2,2-dimethyl-trans-4-(2-oxo-1-piperidiny)-2H-benzo[b]pyran-3-ol (E14)



(E14)

In a similar manner to that described in Example 13 employing 2-piperidone in place of 2-pyrrolidone, the title compound was prepared.

NMR (CDCl₃): δ (3H, s); 1.53 (3H, s); 1.65—2.20 (4H, m); 2.53 (3H, s) overlapped by 2.40—2.75 (2H, m); 2.85—3.20 (2H, m); 3.78 (1H, d, J = 10Hz) overlapped by 3.50—4.00 (1H, m); 5.94 (1H, d, J = 10Hz); 6.88 (1H, d, J = 8Hz); 7.66 (1H, narrow m); 7.80 (1H, q, J = 8, 2Hz).

Pharmacological Data

Systolic blood pressures were recorded by a modification of the tail cuff method described by I. M. Claxton, M. G. Palfreyman, R. H. Poyser, R. L. Whiting, European Journal of Pharmacology, 37, 179 (1976). W+W BP recorder, model 8005 was used to display pulses. Prior to all measurements rats were placed in a heated environment (33.5±0.5°C) before transfer to a restraining cage. Each determination of blood pressure was the mean of at least 6 readings. Spontaneously hypertensive rats (ages 12—18 weeks) with systolic blood pressure 170 mmHg were considered hypertensive.

	Compound of Example 1	Time Post Dose Hours	% Change in Systolic Blood Pressure	% Change in Heart Rate
5	6 rats	1	-47±1	-2±2
	Dose 1 mg/kg po	2	-35±4	-5±3
10	Initial Blood Pressure	4*	-36±4	-5±2
	211±5 mmHg	6*	-34±3	-8±2
15	Initial Hearts Rate	24	+3±2	-9±2
	497±3 beats/min			

* At 4 and 6 hours one rat had no measurable pulse.

Compounds of the other examples were also tested and were found also to be active.

20

Toxicity

No toxic effects were observed in the above test.

Claims for the Contracting States: BE LU SE


25

1. A compound of formula (I):-

30

35

40



(I)

45

wherein:

50

one of R_1 and R_2 is hydrogen and the other is selected from the class of alkylcarbonyl, alkoxy carbonyl, alkylcarbonyloxy, alkylhydroxymethyl, nitro, cyano, chloro, trifluoromethyl, alkylsulphanyl, alkylsulphonyl, alkoxysulphanyl, alkoxysulphonyl, alkylcarbonylamino, alkoxy carbonylamino, or aminosulphanyl, aminosulphonyl or aminocarbonyl, the amino moiety being optionally substituted by one or two alkyl groups, or alkylsulphanylamino, alkylsulphonylamino, alkoxysulphanylamino or alkoxysulphonylamino, or ethylenyl terminally substituted by alkylcarbonyl, nitro or cyano or $-C(alkyl)NOH$ or $-C(alkyl)NNH_2$, the alkyl groups or alkyl moieties of alkyl-containing groups having from 1 to 6 carbon atoms;

55

one of R_3 and R_4 is hydrogen or alkyl having from 1 to 4 carbon atoms and the other is alkyl having from 1 to 4 carbon atoms, or R_3 and R_4 together with the carbon atom to which they are attached are spiroalkyl having from 3 to 6 carbon atoms;

R_5 is hydrogen, alkyl having from 1 to 3 carbon atoms or acyl having from 1 to 8 carbon atoms; and n is 1 or 2; the lactam group being trans to the OR_5 group.

60

2. A compound according to claim 1, characterised in that one of R_1 and R_2 is hydrogen and the other is selected from the class of alkylcarbonyl, alkoxy carbonyl, alkylcarbonyloxy, alkylhydroxymethyl, nitro, chloro or cyano.

3. A compound according to claim 1 or 2, characterised in that the alkyl groups or alkyl moieties of alkyl-containing groups, in respect of the other of R_1 and R_2 , are methyl or ethyl.

65

4. A compound according to any of the preceding claims, characterised in that R_3 and R_4 are both alkyl groups having from 1 to 4 carbon atoms.

5. A compound according to any one of the preceding claims, characterised in that R_5 is hydrogen.

	Compound of Example 1	Time Post Dose Hours	% Change in Systolic Blood Pressure	% Change in Heart Rate
5	6 rats	1	-47±1	-2±2
	Dose 1 mg/kg po	2	-35±4	-5±3
10	Initial Blood Pressure	4*	-36±4	-5±2
	211±5 mmHg	6*	-34±3	-8±2
15	Initial Hearts Rate	24	+3±2	-9±2
	497±3 beats/min			

* At 4 and 6 hours one rat had no measurable pulse.


Compounds of the other examples were also tested and were found also to be active.

Toxicity

No toxic effects were observed in the above test.

Claims for the Contracting States: BE LU SE

1. A compound of formula (I):-



(I)

wherein:

one of R_1 and R_2 is hydrogen and the other is selected from the class of alkylcarbonyl, alkoxycarbonyl, alkylcarbonyloxy, alkylhydroxymethyl, nitro, cyano, chloro, trifluoromethyl, alkylsulphanyl, alkylsulphonyl, alkoxysulphanyl, alkoxysulphonyl, alkylcarbonylamino, alkoxycarbonylamino, or aminosulphanyl, aminosulphonyl or aminocarbonyl, the amino moiety being optionally substituted by one or two alkyl groups, or alkylsulphanylamino, alkylsulphonylamino, alkoxysulphanylamino or alkoxysulphonylamino, or ethylenyl terminally substituted by alkylcarbonyl, nitro or cyano or $-C(\text{alkyl})\text{NOH}$ or $-C(\text{alkyl})\text{NNH}_2$, the alkyl groups or alkyl moieties of alkyl-containing groups having from 1 to 6 carbon atoms;

one of R_3 and R_4 is hydrogen or alkyl having from 1 to 4 carbon atoms and the other is alkyl having from 1 to 4 carbon atoms, or R_3 and R_4 together with the carbon atom to which they are attached are spiroalkyl having from 3 to 6 carbon atoms;

R_5 is hydrogen, alkyl having from 1 to 3 carbon atoms or acyl having from 1 to 8 carbon atoms; and n is 1 or 2; the lactam group being trans to the OR_5 group.

2. A compound according to claim 1, characterised in that one of R_1 and R_2 is hydrogen and the other is selected from the class of alkylcarbonyl, alkoxycarbonyl, alkylcarbonyloxy, alkylhydroxymethyl, nitro, chloro or cyano.


3. A compound according to claim 1 or 2, characterised in that the alkyl groups or alkyl moieties of alkyl-containing groups, in respect of the other of R_1 and R_2 , are methyl or ethyl.

4. A compound according to any of the preceding claims, characterised in that R_3 and R_4 are both alkyl groups having from 1 to 4 carbon atoms.

5. A compound according to any one of the preceding claims, characterised in that R_5 is hydrogen.



6. A compound of formula (II):



(II)

characterised in that one of R_1 and R_2 is hydrogen and the other is selected from the class of alkylcarbonyl, alkoxy, alkoxycarbonyl, alkylcarbonyloxy, alkylhydroxymethyl, nitro or cyano, the alkyl groups or alkyl moieties of alkyl containing groups being methyl or ethyl, and n is 1 or 2; the lactam group being trans to the OH group.

7. A compound according to any one of the preceding claims, characterised in that one of R_1 and R_2 is hydrogen and the other is nitro or cyano.

8. A compound according to any one of the preceding claims characterised in that R_2 is hydrogen.

9. A compound according to claim 8 wherein R_1 is cyano and R_3 and R_4 are both methyl.

10. 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran - 3 - ol; 7 - nitro - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - cyano - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b] - pyran - 3 - yl acetate; 6 - nitro - 3,4 - dihydro - 2 - methyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - carbomethoxy - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - carbomethoxy - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - chloro - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b] - pyran - 3 - ol; 6 - acetyl - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; or 6 - acetyl - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran - 3 - ol.

11. 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b] - pyran - 3 - ol.

12. The compound of claim 11 in the form of a solid.

13. The compound of claim 12 in the form of a crystalline solid.


14. The compound of claim 12 or claim 13 having a melting point range between 226 and 231°C inclusive.

15. The compound of claim 14 having a melting point from 230 to 231°C.

16. A compound according to any one of the preceding claims characterised in that it is in a pharmaceutically acceptable form.

17. A compound according to any one of the preceding claims characterised in that it is prepared synthetically.

18. A process for the preparation of a compound of formula (I), as defined in any one of claims 1 to 11, characterised by cyclising a compound of formula (III), or, when R_5 is hydrogen, a metal salt thereof:




(III)

wherein one of R_1 and R_2 is hydrogen or a group or atom convertible into hydrogen and the other is one of the class of substituents as defined in claim 1 for the other of R_1 and R_2 , or a group or atom convertible thereto, R_3 to R_5 and n are as defined in claim 1, L is a leaving group, and wherein the substituted amino



group is trans to the OR_5 group; in the case where one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group or atom convertible into one of the class of substituents as defined in claim 1 for the other of R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_5 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.


19. A process for the preparation of a compound of formula (I), as defined in any one of claims 1 to 11, characterised by oxidising a compound of formula (X), or when R_5 is hydrogen, a metal salt thereof:



(X)


wherein R_1' and R_2' are as defined in claim 18 and R_3 to R_5 and n are as defined in claim 1, and wherein the nitrogen containing ring is trans to the OR_5 group; in the case when one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group or atom convertible into one of the class of substituents as defined in claim 1 for the other of R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_5 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.

20. A process for the preparation of a compound of formula (I), as defined in any one of claims 1 to 11, characterised by reacting a compound of formula (IV):



(IV)

wherein R_1' and R_2' are as defined in claim 18 and R_3 and R_4 are as defined in claim 1, with an anion of formula (XIII):




(XIII)

wherein n is as defined in claim 1; in the case when one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group or atom convertible into one of the class of substituents as defined in claim 1 for the other of R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_5 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.



21. A process for the preparation of a compound of formula (I), as defined in any one of claims 1 to 11, characterised by cyclising a compound of formula (XIV):



wherein R_1' and R_2' are as defined in claim 18 and R_3 to R_5 and n as defined in claim 1, and L_3 is a leaving group, and wherein the substituted amino group is trans to the OR_5 group; in the case when one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group or atom convertible into one of the class of substituents as defined in claim 1 for the other of R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_5 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.

22. A process according to any one of claims 18 to 21, characterised in that R_1' and R_2' are R_1 and R_2 respectively.

23. A process according to any of claims 18 to 21, characterised in that the compound of formula (I) is isolated in pharmaceutically acceptable form.

24. A pharmaceutical composition characterised by a compound of formula (I), as defined in any one of claims 1 to 17, and a pharmaceutically acceptable carrier.


25. A compound of formula (I), as defined in any one of claims 1 to 17 for use as an active therapeutic substance.

26. A compound of formula (I), as defined in any one of claims 1 to 17 or a composition as defined in claim 24, for use in the treatment of hypertension.

27. A compound of formula (XIV), as defined in claim 21.


28. A compound of formula (III), as defined in claim 18.

29. A compound of formula (IX):




wherein R_1' , R_2' , R_3 and R_4 are as defined in claims 1 and 18, the azide group being trans to the hydroxy group.

30. A compound of formula (XI):



wherein R_5 is methoxy or ethoxy, R_1' , R_2' , R_3 to R_5 and n are as defined in claims 1 to 18, the substituted amino group being trans to the OR_5 group.

31. 6-Cyano-3,4-dihydro-2,2-dimethyl-*trans*-4-amino-2H-benzo[b]pyran-3-ol.
32. 6-Cyano-3,4-dihydro-2,2-dimethyl-*trans*-4-(1-keto-4-chlorobutylamino)-2H-benzo[b]pyran-3-ol.
33. 6-Cyano-3,4-dihydro-2,2-dimethyl-*trans*-4-(4,4-diethoxybutylamino)-2H-benzo[b]pyran-3-ol.
34. 6-Nitro-3,4-dihydro-2-methyl-3,4-epoxy-2H-benzo[b]pyran.
35. 6-Nitro-3,4-dihydro-2-methyl-*trans*-4-(ethoxycarbonylpropylamino)-2H-benzo[b]pyran-3-ol.
36. 7-Nitro-3,4-dihydro-2,2-dimethyl-*trans*-4-(3-carbethoxypropylamino)-2H-benzo[b]pyran-3-ol.
37. 6-Chloro-3,4-dihydro-2,2-dimethyl-*trans*-3-bromo-4-hydroxy-2H-benzo[b]pyran.
38. 6-Chloro-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo[b]pyran.
39. A compound of formula (X):




(X)

wherein R_1' , R_2' , R_3 to R_5 and n are as defined in claims 1 and 18, the nitrogen-containing ring being trans to the OR_5 group.

40. 6-Cyano-3,4-dihydro-2,2-dimethyl-*trans*-4-(2-hydroxy-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol.

Claims for the Contracting States: CH DE FR GB IT NL LI

1. A compound of formula (I):-



(I)

wherein:

- one of R_1 and R_2 is hydrogen and the other is selected from the class of alkylcarbonyl, alkoxycarbonyl, alkylcarbonyloxy, alkylhydroxymethyl, nitro, cyano, chloro, trifluoromethyl, alkylsulphinyl, alkylsulphonyl, alkoxysulphinyl, alkoxysulphonyl, alkylcarbonylamino, alkoxy carbonylamino, or aminosulphinyl, aminosulphonyl or aminocarbonyl, the amino moiety being optionally substituted by one or two alkyl groups, or alkylsulphinylamino, alkylsulphonylamino, alkoxysulphinylamino or alkoxysulphonylamino, or ethylenyl terminally substituted by alkylcarbonyl, nitro or cyano or $-C(alkyl)NOH$ or $-C(alkyl)NNH_2$, the alkyl groups or alkyl moieties of alkyl-containing groups having from 1 to 6 carbon atoms;
- one of R_3 and R_4 is hydrogen or alkyl having from 1 to 4 carbon atoms and the other is alkyl having from 1 to 4 carbon atoms, or R_3 and R_4 together with the carbon atom to which they are attached are spiroalkyl having from 3 to 6 carbon atoms;
- R_5 is hydrogen, alkyl having from 1 to 3 carbon atoms or acyl having from 1 to 8 carbon atoms; and
- n is 1 or 2; the lactam group being trans to the OR_5 group.




2. A compound according to claim 1, characterised in that one of R_1 and R_2 is hydrogen and the other is selected from the class of alkylcarbonyl, alkoxy carbonyl, alkylcarbonyloxy, alkylhydroxymethyl, nitro, chloro or cyano.

3. A compound according to claim 1 or 2, characterised in that the alkyl groups or alkyl moieties of alkyl-containing groups, in respect of the other of R_1 and R_2 , are methyl or ethyl.

4. A compound according to any one of the preceding claims, characterised in that R_3 and R_4 are both alkyl groups having from 1 to 4 carbon atoms.

5. A compound according to any one of the preceding claims, characterised in that R_5 is hydrogen.

6. A compound of formula (II):



(II)

characterised in that one of R_1 and R_2 is hydrogen and the other is selected from the class of alkylcarbonyl, alkoxy carbonyl, alkylcarbonyloxy, alkylhydroxymethyl, nitro or cyano, the alkyl groups or alkyl moieties of alkyl containing groups being methyl or ethyl, and n 1 is or 2; the lactam group being trans to the OH group.

7. A compound according to any one of the preceding claims, characterised in that one of R_1 and R_2 is hydrogen and the other is nitro or cyano.

8. A compound according to any one of the preceding claims characterised in that R_2 is hydrogen.

9. A compound according to claim 8 wherein R_1 is cyano and R_3 and R_4 are both methyl.

10. 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran - 3 - ol; 7 - nitro - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 -

cyano - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - acetate; 6 - nitro - 3,4 - dihydro - 2 - methyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - yl

3 - ol; 6 - carbomethoxy - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 3 - ol; 6 - carbomethoxy - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran -

benzo[b]pyran - 3 - ol; 6 - chloro - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - acetyl - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H -

benzo[b]pyran - 3 - ol; or 6 - acetyl - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran - 3 - ol.

11. 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol.

12. The compound of claim 11 in the form of a solid.

13. The compound of claim 12 in the form of a crystalline solid.


14. The compound of claim 12 or claim 13 having a melting point range between 226 and 231°C inclusive.

15. The compound of claim 14 having a melting point from 230 to 231°C.

16. A compound according to any one of the preceding claims characterised in that it is in a pharmaceutically acceptable form.

17. A compound according to any one of the preceding claims characterised in that it is prepared synthetically.

18. A process for the preparation of a compound of formula (I), as defined in any one of claims 1 to 11, characterised by cyclising a compound of formula (III), or, when R_5 is hydrogen, a metal salt thereof:




(III)



wherein one of R_1' and R_2' is hydrogen or a group or atom convertible into hydrogen and the other is one of the class of substituents as defined in claim 1 for the other of R_1 and R_2 , or a group or atom convertible thereto; R_3 to R_5 and n are as defined in claim 1, L_1 is a leaving group, and wherein the substituted amino group is trans to the OR_5 group; in the case where one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group of atom convertible into one of the class of substituents as defined in claim 1 for the other of R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_5 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.


19. A process for the preparation of a compound of formula (I), as defined in any one of claims 1 to 11, characterised by oxidising a compound of formula (X), or when R_5 is hydrogen, a metal salt thereof:



(X)


wherein R_1' and R_2' are as defined in claim 18 and R_3 to R_5 and n are as defined in claim 1, and wherein the nitrogen containing ring is trans to the OR_5 group; in the case when one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group or atom convertible into one of the class of substituents as defined in claim 1 for the other of R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_5 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.

20. A process for the preparation of a compound of formula (I), as defined in any one of claims 1 to 11, characterised by reacting a compound of formula (IV):



(IV)

wherein R_1' and R_2' are as defined in claim 18 and R_3 and R_4 are as defined in claim 1, with an anion of formula (XIII):




(XIII)

wherein n is as defined in claim 1; in the case when one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group or atom convertible into one of the class of substituents as defined in claim 1 for the other of R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_5 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.



21. A process for the preparation of a compound of formula (I), as defined in any one of claims 1 to 11, characterised by cyclising a compound of formula (XIV):



wherein R_1' and R_2' are as defined in claim 18 and R_3 to R_5 and n is defined in claim 1, and L_3 is a leaving group, and wherein the substituted amino group is trans to the OR_5 group; in the case when one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group or atom convertible into one of the class of substituents as defined in claim 1 for the other of R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_5 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.

22. A process according to any one of claims 18 to 21, characterised in that R_1' and R_2' are R_1 and R_2 respectively.

23. A process according to any of claims 18 to 21, characterised in that the compound of formula (I) is isolated in pharmaceutically acceptable form.

24. A pharmaceutical composition characterised by a compound of formula (I), as defined in any one of claims 1 to 17, and a pharmaceutically acceptable carrier.


25. A compound of formula (I), as defined in any one of claims 1 to 17 for use as an active therapeutic substance.

26. A compound of formula (I), as defined in any one of claims 1 to 17 or a composition as defined in claim 24, for use in the treatment of hypertension.

27. A compound of formula (XIV), as defined in claim 21.

28. A compound of formula (III), as defined in claim 18.

29. A compound of formula (IX):



wherein R_1' , R_2' , R_3 and R_4 are as defined in claims 1 and 18, the azide group being trans to the hydroxy group.

30. 6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-amino-2H-benzo[b]pyran-3-ol.

31. 6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(1-keto-4-chlorobutylamino)-2H-benzo[b]pyran-3-ol.

32. 6-Nitro-3,4-dihydro-2-methyl-3,4-epoxy-2H-benzo[b]pyran.

33. 6-Nitro-3,4-dihydro-2-methyl-trans-4-(ethoxycarbonylpropylamino)-2H-benzo[b]pyran-3-ol.

34. 7-Nitro-3,4-dihydro-2,2-dimethyl-trans-4-(3-carbethoxypropylamino)-2H-benzo[b]pyran-3-ol.


35. 6-Chloro-3,4-dihydro-2,2-dimethyl-trans-3-bromo-4-hydroxy-2H-benzo[b]pyran.

36. 6-Chloro-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo[b]pyran.



Claims for the Contracting State: AT

1. A process for the preparation of a compound of formula (I):-




(I)

wherein:

one of R_1 and R_2 is hydrogen and the other is selected from the class of alkylcarbonyl, alkoxycarbonyl, alkylcarbonyloxy, alkylhydroxymethyl, nitro, cyano, chloro, trifluoromethyl, alkylsulphinyl, alkylsulphonyl, alkoxysulphinyl, alkoxysulphonyl, alkylcarbonylamino, alkoxycarbonylamino, or aminosulphinyl, aminosulphonyl or aminocarbonyl, the amino moiety being optionally substituted by one or two alkyl groups, or alkylsulphinylamino, alkylsulphonylamino, alkoxysulphinylamino or alkoxysulphonylamino, or ethylenyl terminally substituted by alkylcarbonyl, nitro or cyano or $-C(alkyl)NOH$ or $-C(alkyl)NNH_2$, the alkyl groups or alkyl moieties of alkyl-containing groups having from 1 to 6 carbon atoms;

one of R_3 and R_4 is hydrogen or alkyl having from 1 to 4 carbon atoms and the other is alkyl having from 1 to 4 carbon atoms, or R_3 and R_4 together with the carbon atom to which they are attached are spiroalkyl having from 3 to 6 carbon atoms;

R_5 is hydrogen, alkyl having from 1 to 3 carbon atoms or acyl having from 1 to 8 carbon atoms; and n is 1 or 2; the lactam group being trans to the OR_5 group characterised by cyclising a compound of formula (III), or, when R_5 is hydrogen, a metal salt thereof:




(III)

wherein one of R_1' and R_2' is hydrogen or a group or atom convertible into hydrogen and the other is one of the class of substituents as defined in for the other of R_1 and R_2 , or a group or atom convertible thereto, R_3 to R_5 and n are as defined, L is a leaving group, and wherein the substituted amino group is trans to the OR_5 group; in the case where one of R_1' and R_2' is a group or atom convertible into hydrogen; in the case when the other of R_1' and R_2' is a group of atom convertible into one of the class of substituents as defined in claim 1 for the other of R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_5 is hydrogen in the compound of formula (I); optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.




2. A process for the preparation of a compound of formula (I), as defined in claim 1 characterised by oxidising a compound of formula (X), or when R_5 is hydrogen, a metal salt thereof:



20 wherein R_1' and R_3 to R_5 and n are as defined in claim 1, and wherein the nitrogen containing ring is trans to the OR_5 group; in the case when one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group or atom convertible into one of the class of substituents as defined in claim 1 for the other R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_5 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.

3. A process for the preparation of a compound of formula (I), as defined in claim 1, characterised by reacting a compound of formula (IV):




wherein R_1' and R_2' and R_3 and R_4 are as defined in claim 1, with an anion of formula (XIII):



50 wherein n is as defined in claim 1; in the case when one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group or atom convertible into one of the class of substituents as defined in claim 1 for the other of R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_5 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.

55 4. A process for the preparation of a compound of formula (I), as defined in claim 1 characterised by cyclising a compound of formula (XIV):





wherein R_1' and R_2' and R_3 to R_5 and n as defined in claim 1, and L_3 is a leaving group, and wherein the substituted amino group is trans to the OR_5 group; in the case when one of R_1' and R_2' is a group or atom convertible into hydrogen, converting the group or atom into hydrogen; in the case when the other of R_1' and R_2' is a group or atom convertible into one of the class of substituents as defined in claim 1 for the other of R_1 and R_2 , converting the group or atom into one of the class of substituents as defined; and, in the case when R_5 is hydrogen in the compound of formula (I), optionally alkylating or acylating with an alkylating agent having from 1 to 3 carbon atoms or an acylating agent having from 1 to 8 carbon atoms.


5. A process according to any one of claims 1 to 4, characterised in that one of R_1 and R_2 is hydrogen and the other is selected from the class of alkylcarbonyl, alkoxy-carbonyl, alkylcarbonyloxy, alkylhydroxymethyl, nitro, chloro or cyano.

6. A process according to any one of claims 1 to 5, characterised in that the alkyl groups or alkyl moieties of alkyl-containing groups, in respect of the other of R_1 and R_2 , are methyl or ethyl.

7. A process according to any one of the preceding claims, characterised in that R_3 and R_4 are both alkyl groups having from 1 to 4 carbon atoms.

8. A process according to any one of the preceding claims, characterised in that R_5 is hydrogen.

9. A process according to any one of claims 1 to 4, for the preparation of a compound of formula (II):



(II)

characterised in that one of R_1 and R_2 is hydrogen and the other is selected from the class of alkylcarbonyl, alkoxycarbonyl, alkylcarbonyloxy, alkylhydroxymethyl, nitro or cyano, the alkyl groups or alkyl moieties of alkyl containing groups being methyl or ethyl, and n is 1 or 2; the lactam group being trans to the OH group.

10. A process according to any one of the preceding claims, characterised in that one of R_1 and R_2 is hydrogen and the other is nitro or cyano.

11. A process according to any one of the preceding claims, characterised in that R_2 is hydrogen.

12. A process according to claim 11 wherein R_1 is cyano and R_3 and R_4 are both methyl.

13. A process according to any one of claims 1 to 4 for the preparation of 6-Cyano-3,4-dihydro-2,2-dimethyl-*trans*-4-(2-oxo-1-piperidinyl)-2H-benzo[b]pyran-3-ol; 7-nitro-3,4-dihydro-2,2-dimethyl-*trans*-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol; 6-cyano-3,4-dihydro-2,2-dimethyl-*trans*-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-yl acetate; 6-nitro-3,4-dihydro-2-methyl-*trans*-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol; 6-carbomethoxy-3,4-dihydro-2,2-dimethyl-*trans*-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol; 6-carbomethoxy-3,4-dihydro-2,2-dimethyl-*trans*-4-(2-oxo-1-piperidinyl)-2H-benzo[b]pyran-3-ol; 6-chloro-3,4-dihydro-2,2-dimethyl-*trans*-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol; 6-acetyl-3,4-dihydro-2,2-dimethyl-*trans*-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol; or 6-acetyl-3,4-dihydro-2,2-dimethyl-*trans*-4-(2-oxo-1-piperidinyl)-2H-benzo[b]pyran-3-ol.

14. A process according to any of claims 1 to 4 for the preparation of 6-Cyano-3,4-dihydro-2,2-dimethyl-*trans*-4-(2-oxo-1-pyrrolidinyl)-2H-benzo[b]pyran-3-ol.

15. A process according to claim 14 wherein the resulting compound is in the form of a solid.

16. A process according to claim 15 wherein the resulting compound is in the form of a crystalline solid.

17. A process according to claim 15 or 16 wherein the resulting compound has a melting point range between 226 and 231°C inclusive.


18. A process according to claim 17 wherein the resulting compound has a melting point from 230 to 231°C.

19. A process according to any one of claims 1 to 18, characterised in that R_1' and R_2' are R_1 and R_2 respectively.

20. A process according to any of claims 1 to 19, characterised in that the compound of formula (I) is isolated in pharmaceutically acceptable form.



1. Eine Verbindung der Formel (I):



(I)

in der

einer von R_1 und R_2 Wasserstoff ist und der andere aus der Klasse Alkylcarbonyl, Alkoxy carbonyl, Alkylcarbonyloxy, Alkylhydroxymethyl, Nitro, Cyano, Chlor, Trifluormethyl, Alkylsulfinyl, Alkylsulfonyl, Alkoxy sulfinyl, Alkoxy sulfonyl, Alkylcarbonylamino, Alkoxy carbonylamino oder Aminosulfinyl, Aminosulfonyl oder Aminocarbonyl, wobei der Amino teil gegebenenfalls durch eine oder zwei Alkylgruppen substituiert ist, oder Alkylsulfinylamino, Alkylsulfonylamino, Alkoxy sulfinylamino oder Alkoxy sulfonylamino, oder Ethylenyl, das endständig durch Alkylcarbonyl, Nitro oder Cyano oder $-C(Alkyl)NOH$ oder $-C(Alkyl)NNH_2$ substituiert ist, ausgewählt ist, wobei die Alkylgruppen oder Alkylteile von Alkyl enthaltenden Gruppen 1 bis 6 Kohlenstoffatome haben;

einer von R_3 und R_4 Wasserstoff oder Alkyl mit 1 bis 4 Kohlenstoffatomen und der andere Alkyl mit 1 bis 4 Kohlenstoffatomen ist oder R_3 und R_4 zusammen mit dem Kohlenstoffatom, an das sie gebunden sind, Spiroalkyl mit 3 bis 6 Kohlenstoffatomen sind;

R_5 Wasserstoff, Alkyl mit 1 bis 3 Kohlenstoffatomen oder Acyl mit 1 bis 8 Kohlenstoffatomen ist; und n 1 oder 2 ist;

wobei die Lactamgruppe in trans-Stellung zur OR_5 -Gruppe steht.


2. Eine Verbindung nach Anspruch 1, dadurch gekennzeichnet, daß einer von R_1 und R_2 Wasserstoff ist und der andere aus der Klasse Alkylcarbonyl, Alkoxy carbonyl, Alkylcarbonyloxy, Alkylhydroxymethyl, Nitro, Chlor oder Cyano ausgewählt ist.

3. Eine Verbindung nach Anspruch 1 oder 2, dadurch gekennzeichnet, daß die Alkylgruppen oder Alkylteile von Alkyl enthaltenden Gruppen hinsichtlich des anderen von R_1 und R_2 Methyl oder Ethyl sind.

4. Eine Verbindung nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß R_3 und R_4 Alkylgruppen mit 1 bis 4 Kohlenstoffatomen sind.

5. Eine Verbindung nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß R_5 Wasserstoff ist.

6. Eine Verbindung der Formel (II):



(II)



dadurch gekennzeichnet, daß einer von R_1 und R_2 Wasserstoff ist und der andere aus der Klasse Alkylcarbonyl, Alkoxycarbonyl, Alkylcarbonyloxy, Alkylhydroxymethyl, Nitro oder Cyano ausgewählt ist, wobei die Alkylgruppen oder Alkylteile von Alkyl enthaltenden Gruppen Methyl oder Ethyl sind, und n 1 oder 2 ist; wobei die Lactamgruppe in trans-Stellung zur OH-Gruppe steht.

7. Eine Verbindung nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß einer von R_1 und R_2 Wasserstoff ist und der andere Nitro oder Cyano ist.

8. Eine Verbindung nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß R_2 Wasserstoff ist.

9. Eine Verbindung nach Anspruch 8, wobei R_1 Cyano ist und R_3 und R_4 beide Methyl sind.

10. 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran - 3 - ol; 7 - Nitro - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b] - pyran - 3 - yl - acetat; 6 - Nitro - 3,4 - dihydro - 2 - methyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - Carbomethoxy - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - Carbomethoxy - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - Chlor - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b] - pyran - 3 - ol; 6 - Acetyl - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; oder 6 - Acetyl - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran - 3 - ol.

11. 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b] - pyran - 3 - ol.

12. Die Verbindung von Anspruch 11 in der Form eines Feststoffes.

13. Die Verbindung von Anspruch 12 in der Form eines kristallinen Feststoffes.


14. Die Verbindung nach Anspruch 12 oder Anspruch 13 mit einem Schmelzbereich zwischen 226 und 231°C einschließlich.

15. Die Verbindung von Anspruch 14 mit einem Schmelzpunkt von 230 bis 231°C.

16. Eine Verbindung nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß sie in einer pharmazeutisch annehmbaren Form vorliegt.

17. Eine Verbindung nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß sie synthetisch hergestellt ist.

18. Ein Verfahren zur Herstellung einer Verbindung der Formel (II), wie sie in irgendeinem der Ansprüche 1 bis 11 definiert ist, gekennzeichnet durch Cyclisieren einer Verbindung der Formel (III) oder, wenn R_5 Wasserstoff ist, deren Metallsalz




(III)


in der einer von R_1' und R_2' Wasserstoff oder eine Gruppe oder ein Atom ist, die in Wasserstoff unwandelbar sind, und der andere einer von der Klasse von Substituenten, wie in Anspruch 1 für den anderen von R_1 und R_2 definiert ist, oder eine Gruppe oder ein Atom, die dazu umwandelbar sind, ist R_3 bis R_5 und n wie in Anspruch 1 definiert sind, L_1 eine nucleofuge Abgangsgruppe ist und wobei die substituierte Aminogruppe in trans-Stellung zur OH-Gruppe steht; in dem Fall, wo einer von R_1' und R_2' eine Gruppe oder ein Atom ist, die in Wasserstoff umwandelbar sind, Umwandeln der Gruppe oder des Atoms in Wasserstoff; in dem Fall, wenn der andere von R_1' und R_2' eine Gruppe oder ein Atom ist, die in eine der Klasse von Substituenten wie in Anspruch 1 für den anderen von R_1 und R_2 definiert umwandelbar sind, Umwandeln der Gruppe oder des Atoms in einer der Klasse von Substituenten wie definiert; und in dem Fall, wenn R_5 Wasserstoff bei der Verbindung der Formel (I) ist, gegebenenfalls Alkylieren oder Acylieren mit einem Alkylierungsmittel mit 1 bis 3 Kohlenstoffatomen oder mit einem Acylierungsmittel mit 1 bis 8 Kohlenstoffatomen.



19. Ein Verfahren zur Herstellung einer Verbindung der Formel (I), wie sie in irgendeinem der Ansprüche 1 bis 11 definiert ist, gekennzeichnet durch Oxidieren einer Verbindung der Formel (X) oder, wenn R_5 Wasserstoff ist, deren Metallsalz



- 20 in der R_1' und R_2' wie in Anspruch 18 definiert sind und R_3 bis R_5 und n wie in Anspruch 1 definiert sind, und wobei der Stickstoff enthaltende Ring in trans-Stellung zur OR_5 -Gruppe steht;
 in dem Fall, wenn einer von R_1' und R_2' eine Gruppe oder ein Atom ist, die in Wasserstoff umwandelbar sind, Umwandeln der Gruppe oder des Atoms in Wasserstoff;
 in dem Fall, wenn der andere von R_1' und R_2' eine Gruppe oder ein Atom ist, die in eine der Klasse von Substituenten wie in Anspruch 1 für den anderen von R_1 und R_2 definiert umwandelbar sind, Umwandeln der Gruppe oder des Atoms in eine der Klasse von Substituenten wie definiert; und
 in dem Fall, wenn R_5 Wasserstoff bei der Verbindung der Formel (I) ist, gegebenenfalls Alkylieren oder Acylieren mit einem Alkylierungsmittel mit 1 bis 3 Kohlenstoffatomen oder mit einem Acylierungsmittel mit 1 bis 8 Kohlenstoffatomen.
- 30 20. Ein Verfahren zur Herstellung einer Verbindung der Formel (I), wie sie in irgendeinem der Ansprüche 1 bis 11 definiert ist, gekennzeichnet durch Umsetzen einer Verbindung der Formel (IV)




- 45 in der R_1' und R_2' wie in Anspruch 18 definiert sind und R_3 und R_4 wie in Anspruch 1 definiert sind, mit einem Anion der Formel (XIII)



- in der n wie in Anspruch 1 definiert ist, in dem Fall, wenn einer von R_1' und R_2' eine Gruppe oder ein Atom ist, die in Wasserstoff umwandelbar sind, Umwandeln der Gruppe oder des Atoms in Wasserstoff;
 in dem Fall, wenn der andere von R_1' und R_2' eine Gruppe oder ein Atom ist, die in eine der Klasse von Substituenten wie in Anspruch 1 für den anderen von R_1 und R_2 definiert umwandelbar sind, Umwandeln der Gruppe oder des Atoms in eine der Klasse von Substituenten wie definiert; und
 in dem Fall, wenn R_5 Wasserstoff bei der Verbindung der Formel (I) ist, gegebenenfalls Alkylieren oder Acylieren mit einem Alkylierungsmittel mit 1 bis 3 Kohlenstoffatomen oder mit einem Acylierungsmittel mit 1 bis 8 Kohlenstoffatomen.



21. Ein Verfahren zur Herstellung einer Verbindung der Formel (I), wie sie in irgendeinem der Ansprüche 1 bis 11 definiert ist, gekennzeichnet durch Cyclisieren einer Verbindung der Formel (XIV)



(XIV)

in der R_1' und R_2' wie in Anspruch 18 definiert sind und R_3 bis R_5 und n wie in Anspruch 1 definiert sind und L_3 eine nucleofuge Abgangsgruppe ist und wobei die substituierte Aminogruppe in trans-Stellung zur OR_5 -Gruppe steht,

in dem Fall, wenn einer von R_1' und R_2' eine Gruppe oder ein Atom ist, die in Wasserstoff umwandelbar sind, Umwandeln der Gruppe oder des Atoms in Wasserstoff;

in dem Fall, wenn der andere von R_1' und R_2' eine Gruppe oder ein Atom ist, die in eine der Klasse von Substituenten wie in Anspruch 1 für den anderen von R_1 und R_2 definiert umwandelbar sind, Umwandeln der Gruppe oder des Atoms in eine der Klasse von Substituenten wie definiert; und

in dem Fall, wenn R_5 Wasserstoff bei der Verbindung der Formel (I) ist, gegebenenfalls Alkylieren oder Acylieren mit einem Alkylierungsmittel mit 1 bis 3 Kohlenstoffatomen oder mit einem Acylierungsmittel mit 1 bis 8 Kohlenstoffatomen.

22. Ein Verfahren nach irgendeinem der Ansprüche 18 bis 21, dadurch gekennzeichnet, daß R_1' und R_2' , R_1 und R_2 sind.

23. Ein Verfahren nach irgendeinem der Ansprüche 18 bis 21, dadurch gekennzeichnet, daß die Verbindung der Formel (I) in pharmazeutisch annehmbarer Form isoliert wird.

24. Eine pharmazeutische Zusammensetzung, gekennzeichnet durch eine Verbindung der Formel (I), wie sie in irgendeinem der Ansprüche 1 bis 17 definiert ist, und einem pharmazeutisch annehmbaren Träger.


25. Eine Verbindung der Formel (I), wie sie in irgendeinem der Ansprüche 1 bis 17 definiert ist, zur Verwendung als eine aktive therapeutische Substanz.

26. Eine Verbindung der Formel (I), wie sie in irgendeinem der Ansprüche 1 bis 17 definiert ist, oder eine Zusammensetzung, wie sie in Anspruch 24 definiert ist, zur Verwendung bei der Behandlung von hohem Blutdruck.

27. Eine Verbindung der Formel (XIV), wie sie in Anspruch 21 definiert ist.

28. Eine Verbindung der Formel (III), wie sie in Anspruch 18 definiert ist.


29. Eine Verbindung der Formel (IX)



(IX)

in der R_1' , R_2' , R_3 und R_4 wie in den Ansprüchen 1 und 18 definiert sind, wobei die Azidogruppe in trans-Stellung zur Hydroxygruppe steht.

30. Eine Verbindung der Formel (XI)



(XI)



in der R_6 Methoxy oder Ethoxy ist, R_1' , R_2' , R_3 bis R_5 und n wie in den Ansprüchen 1 bis 18 definiert sind, wobei die substituierte Aminogruppe in *trans*-Stellung zur OR_5 -Gruppe steht.

31. 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - amino - 2H - benzo[b]pyran - 3 - ol.

32. 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (1 - keto - 4 chlorbutylamino) - 2H - benzo[b]pyran - 3 - ol.

33. 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (4,4 - diethoxybutylamino) - 2H - benzo[b]pyran - 3 - ol.

34. 6 - Nitro - 3,4 - dihydro - 2 - methyl - 3,4 - epoxy - 2H - benzo[b]pyran.


35. 6 - Nitro - 3,4 - dihydro - 2 - methyl - *trans* - 4 - (ethoxycarbonylpropylamino) - 2H - benzo[b]pyran - 3 - ol.

36. 7 - Nitro - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (3 - carbethoxypropylamino) - 2H - benzo[b]pyran - 3 - ol.

37. 6 - Chlor - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 3 - brom - 4 - hydroxy - 2H - benzo[b]pyran.

38. 6 - Chlor - 3,4 - dihydro - 2,2 - dimethyl - 3,4 - epoxy - 2H - benzo[b]pyran.

39. Eine Verbindung der Formel (X)




(X)

in der R_1' , R_2' , R_3 bis R_5 und n wie in den Ansprüchen 1 und 18 definiert sind, wobei der Stickstoff enthaltende Ring in *trans*-Stellung zur OR_5 -Gruppe steht.

40. 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - hydroxy - 1 - pyrrolidiny) - 2H - benzo[b]pyran - 3 ol.

Patentansprüche für die Vertragsstaaten: CH DE FR GB IT LI NL

1. Eine Verbindung der Formel (I):



(I)

in der

einer von R_1 und R_2 Wasserstoff ist und der andere aus der Klasse Alkylcarbonyl, Alkoxy carbonyl, Alkylcarbonyloxy, Alkylhydroxymethyl, Nitro, Cyano, Chlor, Trifluormethyl, Alkylsulfinyl, Alkylsulfonyl, Alkoxy sulfinyl, Alkoxy sulfonyl, Alkylcarbonylamino, Alkoxy carbonylamino oder Aminosulfinyl, Aminosulfonyl oder Aminocarbonyl, wobei der Aminoteil gegebenenfalls durch eine oder zwei Alkylgruppen substituiert ist; oder Alkylsulfinylamino, Alkylsulfonylamino, Alkoxy sulfinylamino oder Alkoxy sulfonylamino, oder Ethylenyl, das endständig durch Alkylcarbonyl, Nitro oder Cyano oder $-C(Alkyl)NOH$ oder $-C(Alkyl)NNH_2$ substituiert ist, ausgewählt ist, wobei die Alkylgruppen oder Alkylteile von Alkyl enthaltenden Gruppen 1 bis 6 Kohlenstoffatome haben;



einer von R_3 und R_4 Wasserstoff oder Alkyl mit 1 bis 4 Kohlenstoffatomen und der andere Alkyl mit 1 bis 4 Kohlenstoffatomen ist oder R_3 und R_4 zusammen mit dem Kohlenstoffatom, an das sie gebunden sind, Spiroalkyl mit 3 bis 6 Kohlenstoffatomen sind;

R_5 Wasserstoff, Alkyl mit 1 bis 3 Kohlenstoffatomen oder Acyl mit 1 bis 8 Kohlenstoffatomen ist; und n 1 oder 2 ist;

wobei die Lactamgruppe in trans-Stellung zur OR_5 -Gruppe steht.


2. Eine Verbindung nach Anspruch 1, dadurch gekennzeichnet, daß einer von R_1 und R_2 Wasserstoff ist und der andere aus der Klasse Alkylcarbonyl, Alkoxy carbonyl, Alkylcarbonyloxy, Alkylhydroxymethyl, Nitro, Chlor oder Cyano ausgewählt ist.

3. Eine Verbindung nach Anspruch 1 oder 2, dadurch gekennzeichnet, daß die Alkylgruppen oder Alkylteile von Alkyl enthaltenden Gruppen hinsichtlich des anderen von R_1 und R_2 Methyl oder Ethyl sind.

4. Eine Verbindung nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß R_3 und R_4 Alkylgruppen mit 1 bis 4 Kohlenstoffatomen sind.

5. Eine Verbindung nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß R_5 Wasserstoff ist.

6. Eine Verbindung der Formel (II):



(II)

dadurch gekennzeichnet, daß einer von R_1 und R_2 Wasserstoff ist und der andere aus der Klasse Alkylcarbonyl, Alkoxy carbonyl, Alkylcarbonyloxy, Alkylhydroxymethyl, Nitro oder Cyano ausgewählt ist, wobei die Alkylgruppen oder Alkylteile von Alkyl enthaltenden Gruppen Methyl oder Ethyl sind, und n 1 oder 2 ist; wobei die Lactamgruppe in trans-Stellung zur OH-Gruppe steht.

7. Eine Verbindung nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß einer von R_1 und R_2 Wasserstoff ist und der andere Nitro oder Cyano ist.

8. Eine Verbindung nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß R_2 Wasserstoff ist.

9. Eine Verbindung nach Anspruch 8, wobei R_1 Cyano ist und R_3 und R_4 beide Methyl sind.

10. 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran - 3 - ol; 7 - Nitro - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - yl - acetate; 6 - Nitro - 3,4 - dihydro - 2 - methyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - Carbomethoxy - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - Carbomethoxy - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - Chlor - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - Acetyl - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; oder 6 - Acetyl - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran - 3 - ol.

11. 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol.

12. Die Verbindung von Anspruch 11 in der Form eines Feststoffes.

13. Die Verbindung von Anspruch 12 in der Form eines kristallinen Feststoffes.

14. Die Verbindung von Anspruch 12 oder Anspruch 13 mit einem Schmelzbereich zwischen 226 und 231°C einschließend.


15. Die Verbindung von Anspruch 14 mit einem Schmelzpunkt von 230 bis 231°C.

16. Eine Verbindung nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß sie in einer pharmazeutisch annehmbaren Form vorliegt.

17. Eine Verbindung nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß sie synthetisch hergestellt ist.




18. Ein Verfahren zur Herstellung einer Verbindung der Formel (I), wie sie in irgendeinem der Ansprüche 1 bis 11 definiert ist, gekennzeichnet durch Cyclisieren einer Verbindung der Formel (III) oder, wenn R_5 Wasserstoff ist, deren Metallsalz



(III)

- 15 in der einer von R_1' und R_2' Wasserstoff oder eine Gruppe oder ein Atom ist, die in Wasserstoff unverwandelbar sind, und der andere einer von der Klasse von Substituenten, wie in Anspruch 1 für den anderen von R_1 und R_2 definiert ist, oder eine Gruppe oder ein Atom, die dazu umwandelbar sind, ist R_3 bis R_5 und n wie in Anspruch 1 definiert sind, L_1 eine nucleofuge Abgangsgruppe ist und wobei die substituierte Aminogruppe in trans-Stellung zur OH-Gruppe steht; in dem Fall, wo einer von R_1' und R_2' eine Gruppe oder ein Atom ist, die in Wasserstoff umwandelbar sind, Umwandeln der Gruppe oder des Atoms in Wasserstoff; in dem Fall, wenn der andere von R_1' und R_2' eine Gruppe oder ein Atom ist, die in eine der Klasse von Substituenten wie in Anspruch 1 für den anderen von R_1 und R_2 definiert umwandelbar sind, Umwandeln der Gruppe oder des Atoms in eine der Klasse von Substituenten wie definiert; und in dem Fall, wenn R_5 Wasserstoff bei der Verbindung der Formel (I) ist, gegebenenfalls Alkylieren oder
- 20 Acylieren mit einem Alkylierungsmittel mit 1 bis 3 Kohlenstoffatomen oder mit einem Acylierungsmittel mit 1 bis 8 Kohlenstoffatomen.


19. Ein Verfahren zur Herstellung einer Verbindung der Formel (I), wie sie in irgendeinem der Ansprüche 1 bis 11 definiert ist, gekennzeichnet durch Oxidieren einer Verbindung der Formel (X) oder, wenn R_5 Wasserstoff ist, deren Metallsalz



(X)

- 45 in der R_1' und R_2' wie in Anspruch 18 definiert sind und R_3 bis R_5 und n wie in Anspruch 1 definiert sind, und wobei der Stickstoff enthaltende Ring in trans-Stellung zur OR_5 -Gruppe steht; in dem Fall, wenn einer von R_1' und R_2' eine Gruppe oder ein Atom ist, die in Wasserstoff umwandelbar sind, Umwandeln der Gruppe oder des Atoms in Wasserstoff; in dem Fall, wenn der andere von R_1' und R_2' eine Gruppe oder ein Atom ist, die in eine der Klasse von Substituenten wie in Anspruch 1 für den anderen von R_1 und R_2 definiert umwandelbar sind, Umwandeln der Gruppe oder des Atoms in eine der Klasse von Substituenten wie definiert; und
- 50 in dem Fall, wenn R_5 Wasserstoff bei der Verbindung der Formel (I) ist, gegebenenfalls Alkylieren oder Acylieren mit einem Alkylierungsmittel mit 1 bis 3 Kohlenstoffatomen oder mit einem Acylierungsmittel mit 1 bis 8 Kohlenstoffatomen.


20. Ein Verfahren zur Herstellung einer Verbindung der Formel (I), wie sie in irgendeinem der Ansprüche 1 bis 11 definiert ist, gekennzeichnet durch Umsetzen einer Verbindung der Formel (IV)



(IV)




in der R_1' und R_2' wie in Anspruch 18 definiert sind und R_3 und R_4 wie in Anspruch 1 definiert sind, mit einem Anion der Formel (XIII)



(XIII)

- in der n wie in Anspruch 1 definiert ist, in dem Fall, wenn einer von R_1' und R_2' eine Gruppe oder ein Atom ist, die in Wasserstoff umwandelbar sind, Umwandeln der Gruppe oder des Atoms in Wasserstoff; in dem Fall, wenn der andere von R_1' und R_2' eine Gruppe oder ein Atom ist, die in eine der Klasse von Substituenten wie in Anspruch 1 für den anderen von R_1 und R_2 definiert umwandelbar sind, Umwandeln der Gruppe oder des Atoms in eine der Klasse von Substituenten wie definiert; und in dem Fall, wenn R_5 Wasserstoff bei der Verbindung der Formel (I) ist, gegebenenfalls Alkylieren oder Acylieren mit einem Alkylierungsmittel mit 1 bis 3 Kohlenstoffatomen oder mit einem Acylierungsmittel mit 1 bis 8 Kohlenstoffatomen.

21. Ein Verfahren zur Herstellung einer Verbindung der Formel (I), wie sie in irgendeinem der Ansprüche 1 bis 11 definiert ist, gekennzeichnet durch Cyclisieren einer Verbindung der Formel (XIV)



(XIV)

- in der R_1' und R_2' wie in Anspruch 18 definiert sind und R_3 bis R_5 und n wie in Anspruch 1 definiert sind und L_3 eine nucleofuge Abgangsgruppe ist und wobei die substituierte Aminogruppe in trans-Stellung zur OR_5 -Gruppe steht, in dem Fall, wenn einer von R_1' und R_2' eine Gruppe oder ein Atom ist, die in Wasserstoff umwandelbar sind, Umwandeln der Gruppe oder des Atoms in Wasserstoff; in dem Fall, wenn der andere von R_1' und R_2' eine Gruppe oder ein Atom ist, die in eine der Klasse von Substituenten wie in Anspruch 1 für den anderen von R_1 und R_2 definiert umwandelbar sind, Umwandeln der Gruppe oder des Atoms in eine der Klasse von Substituenten wie definiert; und in dem Fall, wenn R_5 Wasserstoff bei der Verbindung der Formel (I) ist, gegebenenfalls Alkylieren oder Acylieren mit einem Alkylierungsmittel mit 1 bis 3 Kohlenstoffatomen oder mit einem Acylierungsmittel mit 1 bis 8 Kohlenstoffatomen.

22. Ein Verfahren nach irgendeinem der Ansprüche 18 bis 21, dadurch gekennzeichnet, daß R_1' und R_2' , R_1 und R_2 sind.

23. Ein Verfahren nach irgendeinem der Ansprüche 18 bis 21, dadurch gekennzeichnet, daß die Verbindung der Formel (I) in pharmazeutisch annehmbarer Form isoliert wird.

24. Eine pharmazeutische Zusammensetzung, gekennzeichnet durch eine Verbindung der Formel (I), wie sie in irgendeinem der Ansprüche 1 bis 17 definiert ist, und einem pharmazeutisch annehmbaren Träger.


25. Eine Verbindung der Formel (I), wie sie in irgendeinem der Ansprüche 1 bis 17 definiert ist, zur Verwendung als eine aktive therapeutische Substanz.

26. Eine Verbindung der Formel (I), wie sie in irgendeinem der Ansprüche 1 bis 17 definiert ist; oder eine Zusammensetzung, wie sie in Anspruch 24 definiert ist, zur Verwendung bei der Behandlung von hohem Blutdruck.

27. Eine Verbindung der Formel (XIV), wie sie in Anspruch 21 definiert ist.

28. Eine Verbindung der Formel (III), wie sie in Anspruch 18 definiert ist.

29. Eine Verbindung der Formel (IX)



(IX)



in der R_1 , R_2 , R_3 und R_4 wie in den Ansprüchen 1 und 18 definiert sind, wobei die Azidogruppe in trans-Stellung zur Hydroxygruppe steht.

30. 6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-amino-2H-benzo[b]pyran-3-ol.

31. 6-Cyano-3,4-dihydro-2,2-dimethyl-trans-4-(1-keto-4-chlorbutylamino)-2H-

benzo[b]pyran-3-ol.

32. 6-Nitro-3,4-dihydro-2-methyl-3,4-epoxy-2H-benzo[b]pyran.

33. 6-Nitro-3,4-dihydro-2-methyl-trans-4-(ethoxycarbonylpropylamino)-2H-benzo[b]pyran-3-ol.


34. 7-Nitro-3,4-dihydro-2,2-dimethyl-trans-4-(3-carbethoxypropylamino)-2H-benzo[b]pyran-3-ol.

35. 6-Chlor-3,4-dihydro-2,2-dimethyl-trans-3-brom-4-hydroxy-2H-benzo[b]pyran.

36. 6-Chlor-3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzo[b]pyran.

Patentansprüche für den Vertragsstaat: AT

1. Eine Verfahren zur Herstellung einer Verbindung der Formel (I):



(I)


in der

einer von R_1 und R_2 Wasserstoff ist und der andere aus der Klasse Alkylcarbonyl, Alkoxy-carbonyl, Alkylcarbonyloxy, Alkylhydroxymethyl, Nitro, Cyano, Chlor, Trifluormethyl, Alkylsulfinyl, Alkylsulfonyl, Alkoxy-sulfinyl, Alkoxy-sulfonyl, Alkylcarbonylamino, Alkoxy-carbonylamino oder Aminosulfinyl, Aminosulfonyl oder Aminocarbonyl, wobei der Aminoteil gegebenenfalls durch eine oder zwei Alkylgruppen substituiert ist, oder Alkylsulfinylamino, Alkylsulfonylamino, Alkoxy-sulfinylamino oder Alkoxy-sulfonylamino, oder Ethylenyl, das endständig durch Alkylcarbonyl, Nitro oder Cyano oder $-C(Alkyl)NOH$ oder $-C(Alkyl)NNH_2$ substituiert ist, ausgewählt ist, wobei die Alkylgruppen oder Alkylteile von Alkyl enthaltenden Gruppen 1 bis 6 Kohlenstoffatome haben;

einer von R_3 und R_4 Wasserstoff oder Alkyl mit 1 bis 4 Kohlenstoffatomen und der andere Alkyl mit 1 bis 4 Kohlenstoffatomen ist oder R_3 und R_4 zusammen mit dem Kohlenstoffatom, an das sie gebunden sind, Spiroalkyl mit 3 bis 6 Kohlenstoffatomen sind;

R_5 Wasserstoff, Alkyl mit 1 bis 3 Kohlenstoffatomen oder Acyl mit 1 bis 8 Kohlenstoffatomen ist; und $n=1$ oder 2 ist;

wobei die Lactamgruppe in trans-Stellung zur OR_5 -Gruppe steht, gekennzeichnet durch Cyclisieren einer Verbindung der Formel (III) oder, wenn R_5 Wasserstoff ist, deren Metallsalz



(III)

in der einer von R_1' und R_2' Wasserstoff oder eine Gruppe oder ein Atom ist, die in Wasserstoff umwandelbar sind, und der andere einer von der Klasse von Substituenten, wie für den anderen von R_1 und R_2 definiert ist, oder eine Gruppe oder ein Atom, die dazu umwandelbar sind, ist, R_3 bis R_5 und n wie definiert sind, L_1 eine nucleofuge Abgangsgruppe ist und wobei die substituierte Aminogruppe in trans-Stellung zur OH-Gruppe steht;






in dem Fall, wo einer von R_1' und R_2' eine Gruppe oder ein Atom ist, die in Wasserstoff umwandelbar sind, Umwandeln der Gruppe oder des Atoms in Wasserstoff; in dem Fall, wenn der andere von R_1' und R_2' eine Gruppe oder ein Atom ist, die in eine der Klasse von Substituenten wie in Anspruch 1 für den anderen von R_1 und R_2 definiert umwandelbar sind, Umwandeln der Gruppe oder des Atoms in eine der Klasse von Substituenten wie definiert; und

in dem Fall, wenn R_5 Wasserstoff bei der Verbindung der Formel (I) ist, gegebenenfalls Alkylieren oder Acylieren mit einem Alkylierungsmittel mit 1 bis 3 Kohlenstoffatomen oder mit einem Acylierungsmittel mit 1 bis 8 Kohlenstoffatomen.

2. Ein Verfahren zur Herstellung einer Verbindung der Formel (I), wie sie in Anspruch 1 definiert ist, gekennzeichnet durch Oxidieren einer Verbindung der Formel (X) oder, wenn R_5 Wasserstoff ist, deren Metallsalz



(X)


in der R_1' und R_2' und R_3 bis R_5 und n wie in Anspruch 1 definiert sind, und wobei der Stickstoff enthaltende Ring in trans-Stellung zur OR_5 -Gruppe steht;

in dem Fall, wenn einer von R_1' und R_2' eine Gruppe oder ein Atom ist, die in Wasserstoff umwandelbar sind, Umwandeln der Gruppe oder des Atoms in Wasserstoff;

in dem Fall, wenn der andere von R_1' und R_2' eine Gruppe oder ein Atom ist, die in eine der Klasse von Substituenten wie in Anspruch 1 für den anderen von R_1 und R_2 definiert umwandelbar sind, Umwandeln der Gruppe oder des Atoms in eine der Klasse von Substituenten wie definiert; und


in dem Fall, wenn R_5 Wasserstoff bei der Verbindung der Formel (I) ist, gegebenenfalls Alkylieren oder Acylieren mit einem Alkylierungsmittel mit 1 bis 3 Kohlenstoffatom oder mit einem Acylierungsmittel mit 1 bis 8 Kohlenstoffatomen.

3. Ein Verfahren zur Herstellung einer Verbindung der Formel (II), wie sie in Anspruch 1 definiert ist, gekennzeichnet durch Umsetzen einer Verbindung der Formel (IV)



(IV)

in der R_1' und R_2' und R_3 und R_4 wie in Anspruch 1 definiert sind, mit einem Anion der Formel (XIII)




(XIII)

in der n wie in Anspruch 1 definiert ist, in dem Fall, wenn einer von R_1' und R_2' eine Gruppe oder ein Atom ist, die in Wasserstoff umwandelbar sind, Umwandeln der Gruppe oder des Atoms in Wasserstoff;



in dem Fall, wenn der andere von R_1' und R_2' eine Gruppe oder ein Atom ist, die in eine der Klasse von Substituenten wie in Anspruch 1 für den anderen von R_1 und R_2 definiert umwandelbar sind, Umwandeln der Gruppe oder des Atoms in eine der Klasse von Substituenten wie definiert; und
 5 in dem Fall, wenn R_5 Wasserstoff bei der Verbindung der Formel (I) ist, gegebenenfalls Alkylieren oder Acylieren mit einem Alkylierungsmittel mit 1 bis 3 Kohlenstoffatomen oder mit einem Acylierungsmittel mit 1 bis 8 Kohlenstoffatomen.

4. Ein Verfahren zur Herstellung einer Verbindung der Formel (II), wie sie in Anspruch 1 definiert ist, gekennzeichnet durch Cyclisieren einer Verbindung der Formel (XIV)



in der R_1' und R_2' und R_3 bis R_5 und n wie in Anspruch 1 definiert sind und L_3 eine nucleofuge Abgangsgruppe ist und wobei die substituierte Aminogruppe in trans-Stellung zur OR_5 -Gruppe steht, in dem Fall, wenn einer von R_1' und R_2' eine Gruppe oder ein Atom ist, die in Wasserstoff umwandelbar sind, Umwandeln der Gruppe oder des Atoms in Wasserstoff;

in dem Fall, wenn der andere von R_1' und R_2' eine Gruppe oder ein Atom ist, die in eine der Klasse von Substituenten wie in Anspruch 1 für den anderen von R_1 und R_2 definiert umwandelbar sind, Umwandeln der Gruppe oder des Atoms in eine der Klasse von Substituenten wie definiert; und

in dem Fall, wenn R_5 Wasserstoff bei der Verbindung der Formel (I) ist, gegebenenfalls Alkylieren oder Acylieren mit einem Alkylierungsmittel mit 1 bis 3 Kohlenstoffatomen oder mit einem Acylierungsmittel mit 1 bis 8 Kohlenstoffatomen.


5. Ein Verfahren nach irgendeinem der Ansprüche 1 bis 4, dadurch gekennzeichnet, daß einer von R_1 und R_2 Wasserstoff ist und der andere aus der Klasse Alkylcarbonyl, Alkoxy carbonyl, Alkylcarbonyloxy, Alkylhydroxymethyl, Nitro, Chlor oder Cyano ausgewählt ist.

6. Ein Verfahren nach irgendeinem der Ansprüche 1 bis 5, dadurch gekennzeichnet, daß die Alkylgruppen oder Alkylteile von Alkyl enthaltenden Gruppen hinsichtlich des anderen von R_1 und R_2 Methyl oder Ethyl sind.

7. Ein Verfahren nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß R_3 und R_4 Alkylgruppen mit 1 bis 4 Kohlenstoffatomen sind.

8. Ein Verfahren nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß R_5 Wasserstoff ist.

9. Ein Verfahren nach irgendeinem der Ansprüche 1 bis 4 zur Herstellung einer Verbindung der Formel (II):



dadurch gekennzeichnet, daß
 einer von R_1 und R_2 Wasserstoff ist und der andere aus der Klasse Alkylcarbonyl, Alkoxy carbonyl, Alkylcarbonyloxy, Alkylhydroxymethyl, Nitro oder Cyano ausgewählt ist, wobei die Alkylgruppen oder Alkylteile von Alkyl enthaltenden Gruppen Methyl oder Ethyl sind, und
 n 1 oder 2 ist;

wobei die Lactamgruppe in trans-Stellung zur OH-Gruppe steht.



10. Ein Verfahren nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß einer von R_1 und R_2 Wasserstoff ist und der andere Nitro oder Cyano ist.

11. Ein Verfahren nach irgendeinem der vorstehenden Ansprüche, dadurch gekennzeichnet, daß R_2 Wasserstoff ist.

12. Ein Verfahren nach Anspruch 11, wobei R_1 Cyano ist und R_3 und R_4 beide Methyl sind.

13. Ein Verfahren nach irgendeinem der Ansprüche 1 bis 4 zur Herstellung von 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran - 3 - ol; 7 - Nitro - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b] - pyran - 3 - yl - acetat; 6 - Nitro - 3,4 - dihydro - 2 - methyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - Carbomethoxy - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - Carbomethoxy - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - Chlor - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b] - pyran - 3 - ol; 6 - Acetyl - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 ol; oder 6 - Acetyl - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - piperidinyl) - 2H - benzo[b]pyran - 3 - ol.

14. Ein Verfahren nach irgendeinem der Ansprüche 1 bis 4 zur Herstellung von 6 - Cyano - 3,4 - dihydro - 2,2 - dimethyl - *trans* - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b] - pyran - 3 - ol.

15. Ein Verfahren nach Anspruch 14, wobei die erhaltene Verbindung in der Form eines Feststoffes vorliegt.

16. Ein Verfahren nach Anspruch 15, wobei die erhaltene Verbindung in der Form eines kristallinen Feststoffes vorliegt.

17. Ein Verfahren nach Anspruch 15 oder 16, wobei die erhaltene Verbindung einen Schmelzbereich zwischen 226 und 231°C, einschließlich, hat.


18. Ein Verfahren nach Anspruch 17, wobei die erhaltene Verbindung einen Schmelzpunkt von 230 bis 231°C hat.

19. Ein Verfahren nach irgendeinem der Ansprüche 1 bis 18, dadurch gekennzeichnet, daß R_1' und R_2' R_1 und R_2 sind.

20. Ein Verfahren nach irgendeinem der Ansprüche 1 bis 19, dadurch gekennzeichnet, daß die Verbindung der Formel (I) in einer pharmazeutisch annehmbaren Form isoliert wird.

Revendications pour les Etats contractants: BE LU SE

1. Composé de formule (I):



(I)

dans laquelle:

l'un des R_1 et R_2 est un atome d'hydrogène et l'autre est choisi dans la classe des groupes alcoylcarbonyle, alcoylcarbonyle, alcoylcarbonyle, alcoylhydroxyméthyle, nitro, cyano, chloro, trifluorométhyle, alcoylsulfinyle, alcoylsulfonyle, alcoylsulfonyle, alcoylsulfonylamine, alcoylcarbonylamine, alcoylcarbonylamine, ou aminosulfinyle, aminosulfonyle ou aminocarbonyle, la partie amino étant éventuellement substituée par un ou deux groupes alcoyles, ou alcoylsulfinylamine, alcoylsulfonylamine, alcoylsulfonylamine ou alcoylsulfonylamine, éthylényle substitué en position terminale par un alcoylcarbonyle, nitro ou cyano, ou $-C(\text{alcoyl})NOH$ ou $-C(\text{alcoyl})NNH_2$, les groupes alcoyles ou les parties alcoyles des radicaux contenant le groupe alcoyle ayant 1 à 6 atomes de carbone;

l'une des R_3 et R_4 est un atome d'hydrogène ou un groupe alcoyle ayant 1 à 4 atomes de carbone et l'autre est le groupe alcoyle ayant 1 à 4 atomes de carbone, ou R_3 et R_4 forment, avec l'atome de carbone auquel ils sont attachés, un groupe spiroalcoyle de 3 à 6 atomes de carbone;

R_5 est un atome d'hydrogène, un groupe alcoyle ayant 1 à 3 atomes de carbone ou acyle ayant 1 à 8 atomes de carbone; et

n est 1 ou 2; le groupe lactame étant en position *trans* par rapport au groupe OR_5 .




2. Composé suivant la revendication 1, caractérisé en ce que l'un des R_1 et R_2 est un atome d'hydrogène et l'autre est choisi dans la classe des groupes alcoylcarbonyle, alcoxycarbonyle, alcoylcarbonyloxy, alcoylhydroxyméthyle, nitro, chloro ou cyano.

3. Composé suivant la revendication 1 ou 2, caractérisé en ce que les groupes alcoyles ou les parties alcoyles des radicaux contenant un groupe alcoyle concernant l'autre des R_1 et R_2 , sont des groupes méthyle ou éthyle.

4. Composé suivant l'une quelconque des revendications précédentes caractérisé en ce que R_3 et R_4 sont tous deux des groupes alcoyles ayant 1 à 4 atomes de carbone.

5. Composé suivant l'une quelconque des revendications précédentes caractérisé en ce que R_5 est un atome d'hydrogène.

6. Composé de formule (II):



(II)

caractérisé en ce que l'une des R_1 et R_2 est un atome d'hydrogène et l'autre est choisi dans la classe des groupes alcoylcarbonyle, alcoxycarbonyle, alcoylcarbonyloxy, alcoylhydroxyméthyle, nitro ou cyano, les groupes alcoyles ou les parties alcoyles des radicaux contenant un groupe alcoyle étant un groupe méthyle ou éthyle, et n est 1 ou 2; le groupe lactame étant trans par rapport au groupe OH.

7. Composé suivant l'une quelconque des revendications précédentes, caractérisé en ce que l'un des R_1 et R_2 est un atome d'hydrogène et l'autre est un groupe nitro ou cyano.

8. Composé suivant l'une quelconque des revendications précédentes, caractérisé en ce que R_2 est un atome d'hydrogène.

9. Composé suivant la revendication 8, caractérisé en ce que R_1 est un groupe cyano et R_3 et R_4 sont tous deux un groupe méthyle.

10. 6 - Cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pipéridinyl) - 2H - benzo[b]pyran - 3 - ol; 7 - nitro - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; acétate de 6 - cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - yle; 6 - nitro - 3,4 - dihydro - 2 - méthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - carbométhoxy - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - carbométhoxy - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pipéridinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - chloro - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - acétyl - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; ou 6 - acétyl - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pipéridinyl) - 2H - benzo[b]pyran - 3 - ol.

11. 6 - Cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol.

12. Composé suivant la revendication 11, sous la forme d'un solide.

13. Composé suivant la revendication 12, sous la forme d'un solide cristallin.

14. Composé suivant la revendication 12 ou la revendication 13, ayant une gamme de points de fusion entre 226° et 231°C inclus.


15. Composé suivant la revendication 14 ayant un point de fusion de 230 à 231°C.

16. Composé suivant l'une quelconque des revendications précédentes, caractérisé en ce qu'il est sous une forme acceptable du point de vue pharmaceutique.

17. Composé suivant l'une quelconque des revendications précédentes, caractérisé en ce qu'il est préparé par synthèse.




18. Procédé pour la préparation d'un composé de formule (II), tel que défini dans l'une quelconque des revendications 1 à 11, caractérisé en ce qu'on cyclise un composé de formule (III), ou lorsque R_5 est un atome d'hydrogène, un sel métallique de celui-ci:



(III)

où l'un des R_1' et R_2' est un atome d'hydrogène ou un groupe ou un atome pouvant être converti en un atome d'hydrogène et l'autre est l'un des substituants de la classe définie dans la revendication 1 pour l'autre des R_1 et R_2 , ou un groupe ou un atome pouvant être converti en ceux-ci, R_3 à R_5 et n sont tels que définis dans la revendication 1, L_1 est un groupe mobile, et où le groupe amino substitué est trans par rapport au groupe OR_5 ; dans le cas où l'un des R_1' et R_2' est un groupe ou un atome pouvant être converti en un atome d'hydrogène, on convertit le groupe ou l'atome en un atome d'hydrogène; dans le cas où l'autre des R_1' et R_2' est un groupe ou un atome pouvant être converti en l'un des substituants de la classe définie dans la revendication 1 pour l'autre des R_1 et R_2 , on convertit le groupe ou l'atome en l'un des substituants de la classe définie plus haut; et, dans le cas où R_5 est un atome d'hydrogène dans le composé de formule (I), on alcoyle ou acyle éventuellement avec un agent alcoylant ayant 1 à 3 atomes de carbone ou un agent acylant ayant 1 à 8 atomes de carbone.


19. Procédé pour la préparation d'un composé de formule (I), tel que défini dans l'une quelconque des revendications 1 à 11, caractérisé en ce qu'on oxyde un composé de formule (X), ou, lorsque R_5 est un atome d'hydrogène, un sel métallique de celui-ci:



(X)

dans laquelle R_1' et R_2' sont tels que définis dans la revendication 18 et R_3 à R_5 et n sont tels que définis dans la revendication 1, et où le cycle contenant de l'azote est en position trans par rapport au groupe OR_5 ; dans le cas où l'un des R_1' et R_2' est un groupe ou un atome pouvant être converti en un atome d'hydrogène, on convertit le groupe ou l'atome en un atome d'hydrogène; dans le cas où l'autre des R_1' et R_2' est un groupe ou un atome pouvant être converti en l'un des substituants de la classe définie dans la revendication 1 pour l'autre des R_1 et R_2 , on convertit le groupe ou l'atome en l'un des substituants de la classe définie plus haut; et, dans le cas où R_5 est un atome d'hydrogène dans le composé de formule (I), on alcoyle ou acyle éventuellement avec un agent alcoylant ayant 1 à 3 atomes de carbone ou un agent acylant ayant 1 à 8 atomes de carbone.


20. Procédé pour la préparation d'un composé de formule (I), tel que défini dans l'une quelconque des revendications 1 à 11, caractérisé en ce qu'on fait réagir un composé de formule (IV):



(IV)




dans laquelle R_1' et R_2' sont tels que définis dans la revendication 18 et R_3 et R_4 sont tels que définis dans la revendication 1, avec un anion de formule (XIII):



(XIII)

dans laquelle n est tel que défini dans la revendication 1; dans le cas où l'un des R_1' et R_2' est un groupe ou un atome pouvant être converti en un atome d'hydrogène, on convertit le groupe ou l'atome en un atome d'hydrogène; dans le cas où l'autre des R_1' et R_2' est un groupe ou un atome pouvant être converti en l'un des substituants de la classe définie pour l'autre des R_1 et R_2 , on convertit le groupe ou l'atome en l'un des substituants de la classe définie plus haut; et, dans le cas où R_5 est un atome d'hydrogène dans le composé de formule (I), on alcoyle ou acyle éventuellement avec un agent alcoylant ayant 1 à 3 atomes de carbone ou un agent acylant ayant 1 à 8 atomes de carbone.

21. Procédé pour la préparation d'un composé de formule (I), tel que défini dans l'une quelconque des revendications 1 à 11, caractérisé en ce qu'on cyclise un composé de formule (XIV):



(XIV)

dans laquelle R_1' et R_2' sont tels que définis dans la revendication 18 et R_3 à R_5 et n sont tels que définis dans la revendication 1 et L_3 est un groupe mobile, et où le groupe amino substitué est en position trans par rapport au groupe $-OR_5$; dans le cas où l'un des R_1' et R_2' est un groupe ou un atome pouvant être converti en un atome d'hydrogène, on convertit le groupe ou l'atome en un atome d'hydrogène; dans le cas où l'autre des R_1' et R_2' est un groupe ou un atome pouvant être converti en l'un des substituants de la classe définie dans la revendication 1 pour l'autre des R_1 et R_2 , on convertit le groupe ou l'atome en l'un des substituants de la classe définie plus haut; et, dans le cas où R_5 est un atome d'hydrogène dans le composé de formule (I), on alcoyle ou acyle éventuellement avec un agent alcoylant ayant 1 à 3 atomes de carbone ou un agent acylant ayant 1 à 8 atomes de carbone.

22. Procédé suivant l'une quelconque des revendications 18 à 21, caractérisé en ce que R_1' et R_2' sont R_1 et R_2 respectivement.

23. Procédé suivant l'une quelconque des revendications 18 à 21, caractérisé en ce que le composé de formule (I) est isolé sous une forme acceptable du point de vue pharmaceutique.

24. Composition pharmaceutique caractérisée par un composé de formule (I) tel que défini dans l'une quelconque des revendications 1 à 17 et un support acceptable du point de vue pharmaceutique.


25. Composé de formule (I), tel que défini dans l'une quelconque des revendications 1 à 17, utile en tant que substance thérapeutique active.

26. Composé de formule (I) tel que défini dans l'une quelconque des revendications 1 à 17 ou composition telle que définie dans la revendication 24, utile dans le traitement de l'hypertension.

27. Composé de formule (XIV), tel que défini dans la revendication 21.

28. Composé de formule (III) tel que défini dans la revendication 18.

29. Composé de formule (IX):




(IX)



dans laquelle R_1' , R_2' , R_3 et R_4 sont tels que définis dans les revendications 1 et 18, le groupe azide étant en position trans par rapport au groupe hydroxy.

30. Composé de formule (XI):



(XI)

dans laquelle R_6 est un groupe méthoxy ou éthoxy, R_1' , R_2' , R_3 à R_5 et n sont tels que définis dans les revendications 1 à 18, le groupe amino substitué étant en position trans par rapport au groupe OR_5 .

31. 6 - Cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - amino - 2H - benzo[b]pyran - 3 - ol.

32. 6 - Cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (1 - céto - 4 - chlorobutylamino) - 2H - benzo[b]pyran - 3 - ol.

33. 6 - Cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (4,4 - diéthoxybutylamino) - 2H - benzo[b]pyran - 3 - ol.

34. 6 - Nitro - 3,4 - dihydro - 2 - méthyl - 3,4 - époxy - 2H - benzo[b]pyranne.


35. 6 - Nitro - 3,4 - dihydro - 2 - méthyl - trans - 4 - (éthoxycarbonylpropylamino) - 2H - benzo[b]pyran - 3 - ol.

36. 7 - Nitro - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (3 - carbéthoxypropylamino) - 2H - benzo[b]pyran - 3 - ol.

37. 6 - Chloro - 3,4 - dihydro - 2,2 - diméthyl - trans - 3 - bromo - 4 - hydroxy - 2H - benzo[b]pyranne.

38. 6 - Chloro - 3,4 - dihydro - 2,2 - diméthyl - 3,4 - époxy - 2H - benzo[b]pyranne.

39. Composé de formule (X):




(X)

dans laquelle R_1' , R_2' , R_3 à R_5 et n sont tels que définis dans les revendications 1 et 18, le cycle contenant l'azote étant en position trans par rapport au groupe OR_5 .

40. 4 - Cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - hydroxy - 1 - pyrrolidiny) - 2H - benzo[b]pyran - 3 - ol.



1. Composé de formule (I):



(I)

dans laquelle:

l'un des R_1 et R_2 est un atome d'hydrogène et l'autre est choisi dans la classe des groupes alcoylcarbonyle, alcoxycarbonyle, alcoylcarbonyle, alcoylhydroxyméthyle, nitro, cyano, chloro, trifluorométhyle, alcoylsulfinyle, alcoylsulfonyle, alcoxysulfinyle, alcoxysulfonyle, alcoylcarbonylamino, alcoxycarbonylamino ou aminosulfinyle, aminosulfonyle ou aminocarbonyle, la partie amino étant éventuellement substituée par un ou deux groupes alcoyles, ou alcoylsulfinylamino, alcoylsulfonylamino, alcoxysulfinylamino ou alcoxysulfonylamino, éthylényle substitué en position terminale par un alcoylcarbonyle, nitro ou cyano, ou $-C(\text{alcoyl})NOH$ ou $-C(\text{alcoyl})NNH_2$, les groupes alcoyles ou les parties alcoyles des radicaux contenant le groupe alcoyle ayant 1 à 6 atomes de carbone;

l'une des R_3 et R_4 est un atome d'hydrogène ou un groupe alcoyle ayant 1 à 4 atomes de carbone et l'autre est le groupe alcoyle ayant 1 à 4 atomes de carbone, ou R_3 et R_4 forment avec l'atome de carbone auquel ils sont attachés, un groupe spiroalcoyle de 3 à 6 atomes de carbone;

R_5 est un atome d'hydrogène, un groupe alcoyle ayant 1 à 3 atomes de carbone ou acyle ayant 1 à 8 atomes de carbone; et

n est 1 ou 2; le groupe lactame étant en position trans par rapport au groupe OR_5 .


2. Composé suivant la revendication 1, caractérisé en ce que l'un des R_1 et R_2 est un atome d'hydrogène et l'autre est choisi dans la classe des groupes alcoylcarbonyle, alcoxycarbonyle, alcoylcarbonyloxy, alcoylhydroxyméthyle, nitro, chloro ou cyano.

3. Composé suivant la revendication 1 ou 2, caractérisé en ce que les groupes alcoyles ou les parties alcoyles des radicaux contenant un groupe alcoyle concernant l'autre des R_1 et R_2 , sont des groupes méthyle ou éthyle.

4. Composé suivant l'une quelconque des revendications précédentes caractérisé en ce que R_3 et R_4 sont tous deux des groupes alcoyles ayant 1 à 4 atomes de carbone.

5. Composé suivant l'une quelconque des revendications précédentes caractérisé en ce que R_5 est un atome d'hydrogène.

6. Composé de formule (II):



(II)



caractérisé en ce que l'une des R_1 et R_2 est un atome d'hydrogène et l'autre est choisi dans la classe des groupes alcoylcarbonyle, alcoxycarbonyle, alcoylcarbonyloxy, alcoylhydroxyméthyle, nitro ou cyano, les groupes alcoyles ou les parties alcoyles des radicaux contenant un groupe alcoyle étant un groupe méthyle ou éthyle, et n est 1 ou 2; le groupe lactame étant trans par rapport au groupe OH.

7. Composé suivant l'une quelconque des revendications précédentes, caractérisé en ce que l'un des R_1 et R_2 est un atome d'hydrogène et l'autre est un groupe nitro ou cyano.

8. Composé suivant l'une quelconque des revendications précédentes, caractérisé en ce que R_2 est un atome d'hydrogène.

9. Composé suivant la revendication 8, caractérisé en ce que R_1 est un groupe cyano et R_3 et R_4 sont tous deux un groupe méthyle.

10. 6 - Cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pipéridinyl) - 2H - benzo[b]pyran - 3 - ol; 7 - nitro - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; acétate de 6 - cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - yle; 6 - nitro - 3,4 - dihydro - 2 - méthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - carbométhoxy - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - carbométhoxy - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pipéridinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - chloro - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; 6 - acétyl - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; ou 6 - acétyl - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pipéridinyl) - 2H - benzo[b]pyran - 3 - ol.

11. 6 - Cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol.

12. Composé suivant la revendication 11, sous la forme d'un solide.

13. Composé suivant la revendication 12, sous la forme d'un solide cristallin.


14. Composé suivant la revendication 12 ou la revendication 13, ayant une gamme de points de fusion entre 226° et 231°C inclus.

15. Composé suivant la revendication 14, ayant un point de fusion de 230 à 231°C.

16. Composé suivant l'une quelconque des revendications précédentes, caractérisé en ce qu'il est sous une forme acceptable du point de vue pharmaceutique.

17. Composé suivant l'une quelconque des revendications précédentes, caractérisé en ce qu'il est préparé par synthèse.

18. Procédé pour la préparation d'un composé de formule (I), tel que défini dans l'une quelconque des revendications 1 à 11, caractérisé en ce qu'on cyclise un composé de formule (III), ou lorsque R_5 est un atome d'hydrogène, un sel métallique de celui-ci:




(III)

où l'un des R_1 et R_2 est un atome d'hydrogène ou un groupe ou un atome pouvant être converti en un atome d'hydrogène et l'autre est l'un des substituants de la classe définie dans la revendication 1 pour l'autre des R_1 et R_2 , ou un groupe ou un atome pouvant être converti en ceux-ci, R_3 à R_5 et n sont tels que définis dans la revendication 1, L_1 est un groupe mobile, et où le groupe amino substitué est trans par rapport au groupe OR_5 ; dans le cas où l'un des R_1 et R_2 est un groupe ou un atome pouvant être converti en un atome d'hydrogène, on convertit le groupe ou l'atome en un atome d'hydrogène; dans le cas où l'autre des R_1 et R_2 est un groupe ou un atome pouvant être converti en l'un des substituants de la classe définie dans la revendication 1 pour l'autre des R_1 et R_2 , on convertit le groupe ou l'atome en l'un des substituants de la classe définie plus haut; et, dans le cas où R_5 est un atome d'hydrogène dans le composé de formule (I), on alcoyle ou acyle éventuellement avec un agent alcoylant ayant 1 à 3 atomes de carbone ou un agent acylant ayant 1 à 8 atomes de carbone.




19. Procédé pour la préparation d'un composé de formule (I), tel que défini dans l'une quelconque des revendications 1 à 11, caractérisé en ce qu'on oxyde un composé de formule (X), ou, lorsque R_5 est un atome d'hydrogène, un sel métallique de celui-ci:



(X)


dans laquelle R_1' et R_2' sont tels que définis dans la revendication 18 et R_3 à R_5 et n sont tels que définis dans la revendication 1, et où le cycle contenant de l'azote est en position trans par rapport au groupe OR_5 ; dans le cas où l'un des R_1' et R_2' est un groupe ou un atome pouvant être converti en un atome d'hydrogène, on convertit le groupe ou l'atome en un atome d'hydrogène; dans le cas où l'autre des R_1' et R_2' est un groupe ou un atome pouvant être converti en l'un des substituants de la classe définie dans la revendication 1 pour l'autre des R_1 et R_2 ; on convertit le groupe ou l'atome en l'un des substituants de la classe définie plus haut; et, dans le cas où R_5 est un atome d'hydrogène dans le composé de formule (I), on alcoyle ou acyle éventuellement avec un agent alcoylant ayant 1 à 3 atomes de carbone ou un agent acylant ayant 1 à 8 atomes de carbone.

20. Procédé pour la préparation d'un composé de formule (I), tel que défini dans l'une quelconque des revendications 1 à 11, caractérisé en ce qu'on fait réagir un composé de formule (IV):



(IV)

dans laquelle R_1' et R_2' sont tels que définis dans la revendication 18 et R_3 et R_4 sont tels que définis dans la revendication 1, avec un anion de formule (XIII):




(XIII)

dans laquelle n est tel que défini dans la revendication 1; dans le cas où l'un des R_1' et R_2' est un groupe ou un atome pouvant être converti en un atome d'hydrogène, on convertit le groupe ou l'atome en un atome d'hydrogène; dans le cas où l'autre des R_1' et R_2' est un groupe ou un atome pouvant être converti en l'un des substituants de la classe définie pour l'autre des R_1 et R_2 , on convertit le groupe ou l'atome en l'un des substituants de la classe définie plus haut; et, dans le cas où R_5 est un atome d'hydrogène dans le composé de formule (I), on alcoyle ou acyle éventuellement avec un agent alcoylant ayant 1 à 3 atomes de carbone ou un agent acylant ayant 1 à 8 atomes de carbone.



21. Procédé pour la préparation d'un composé de formule (II), tel que défini dans l'une quelconque des revendications 1 à 11, caractérisé en ce qu'on cyclise un composé de formule (XIV):



(XIV)

dans laquelle R_1' et R_2' sont tels que définis dans la revendication 18 et R_3 à R_5 et n sont tels que définis dans la revendication 1 et L_3 est un groupe mobile, et où le groupe amino substitué est en position trans par rapport au groupe $-OR_5$; dans le cas où l'un des R_1' et R_2' est un groupe ou un atome pouvant être converti en un atome d'hydrogène, on convertit le groupe ou l'atome en un atome d'hydrogène; dans le cas où l'autre des R_1' et R_2' est un groupe ou un atome pouvant être converti en l'un des substituants de la classe définie dans la revendication 1 pour l'autre des R_1 et R_2 , on convertit le groupe ou l'atome en l'un des substituants de la classe définie plus haut; et, dans le cas où R_5 est un atome d'hydrogène dans le composé de formule (II), on alcoyle ou acyle éventuellement avec un agent alcoylant ayant 1 à 3 atomes de carbone ou un agent acylant ayant 1 à 8 atomes de carbone.

22. Procédé suivant l'une quelconque des revendications 18 à 21, caractérisé en ce que R_1' et R_2' sont R_1 et R_2 respectivement.

23. Procédé suivant l'une quelconque des revendications 18 à 21, caractérisé en ce que le composé de formule (I) est isolé sous une forme acceptable du point de vue pharmaceutique.

24. Composition pharmaceutique caractérisée par un composé de formule (I) tel que défini dans l'une quelconque des revendications 1 à 17 et un support acceptable du point de vue pharmaceutique.


25. Composé de formule (I), tel que défini dans l'une quelconque des revendications 1 à 17, utile en tant que substance thérapeutique active.

26. Composé de formule (I) tel que défini dans l'une quelconque des revendications 1 à 17 ou composition telle que définie dans la revendication 24, utile dans le traitement de l'hypertension.

27. Composé de formule (XIV), tel que défini dans la revendication 21.

28. Composé de formule (III) tel que défini dans la revendication 18.

29. Composé de formule (IX):



(IX)

dans laquelle R_1' , R_2' , R_3 et R_4 sont tels que définis dans les revendications 1 et 18, le groupe azide étant en position trans par rapport au groupe hydroxy.

30. 6 - Cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - amino - 2H - benzo[b]pyran - 3 - ol.

31. 6 - Cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (1 - céto - 4 - chlorobutylamino) - 2H - benzo[b]pyran - 3 - ol.

32. 6 - Nitro - 3,4 - dihydro - 2 - méthyl - 3,4 - époxy - 2H - benzo[b]pyranne.

33. 6 - Nitro - 3,4 - dihydro - 2 - méthyl - trans - 4 - (éthoxycarbonylpropylamino) - 2H - benzo[b]pyran - 3 - ol.

34. 7 - Nitro - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (3 - carbéthoxypropylamino) - 2H - benzo[b]pyran - 3 - ol.


35. 6 - Chloro - 3,4 - dihydro - 2,2 - diméthyl - trans - 3 - bromo - 4 - hydroxy - 2H - benzo[b]pyranne.

36. 6 - Chloro - 3,4 - dihydro - 2,2 - diméthyl - 3,4 - époxy - 2H - benzo[b]pyranne.



Revendications pour les Etat contractant: AT

1. Procédé pour la préparation d'un composé de formule (I):



(I)


dans laquelle:

l'un des R_1 et R_2 est un atome d'hydrogène et l'autre est choisi dans la classe des groupes alcoylcarbonyle, alcoycarbonyle, alcoylcarbonyloxy, alcoylhydroxyméthyle, nitro, cyano, chloro, trifluorométhyle, alcoylsulfinyle, alcoylsulfonyle, alcoylsulfinyle, alcoylsulfonyle, alcoylcarbonylamino, alcoylcarbonylamino, ou aminosulfinyle, aminosulfonyle ou aminocarbonyle, la partie amino étant éventuellement substituée par un ou deux groupes alcoyles, ou alcoylsulfinylamino, alcoylsulfonylamino, alcoylcarbonyle, nitro ou cyano, ou $-C(\text{alcoyl})NOH$ ou $-C(\text{alcoyl})NNH_2$, les groupes alcoyles ou les parties

alcoyles des radicaux contenant le groupe alcoyle ayant 1 à 6 atomes de carbone; l'une des R_3 et R_4 est un atome d'hydrogène ou un groupe alcoyle ayant 1 à 4 atomes de carbone et l'autre est le groupe alcoyle ayant 1 à 4 atomes de carbone, ou R_3 et R_4 forment, avec l'atome de carbone auquel ils sont attachés, un groupe spiroalcoyle de 3 à 6 atomes de carbone;

R_5 est un atome d'hydrogène, un groupe alcoyle ayant 1 à 3 atomes de carbone ou acyle ayant 1 à 8

atomes de carbone; et n est 1 ou 2; le groupe lactame étant en position trans par rapport au groupe OR_5 , caractérisé en ce qu'on cyclise un composé de formule (III), ou lorsque R_5 est un atome d'hydrogène, un sel métallique de celui-ci:




(III)

où l'un des R_1' et R_2' est un atome d'hydrogène ou un groupe ou un atome pouvant être converti en un atome d'hydrogène et l'autre est l'un des substituants de la classe définie dans la revendication 1 pour l'autre des R_1 et R_2 , ou un groupe ou un atome pouvant être converti en ceux-ci, R_3 à R_5 et n sont tels que définis dans la revendication 1, L_1 est un groupe mobile, et où le groupe amino substitué est trans par rapport au groupe OR_5 ; dans le cas où l'un des R_1' et R_2' est un groupe ou un atome pouvant être converti en un atome d'hydrogène, on convertit le groupe ou l'atome en un atome d'hydrogène; dans le cas où l'autre des R_1' et R_2' est un groupe ou un atome pouvant être converti en l'un des substituants de la classe définie dans la revendication 1 pour l'autre des R_1 et R_2 , on convertit le groupe ou l'atome en l'un des substituants de la classe définie plus haut; et, dans le cas où R_5 est un atome d'hydrogène dans le composé de formule (I), on alcoyle ou acyle éventuellement avec un agent alcoylant ayant 1 à 3 atomes de carbone ou un agent acylant ayant 1 à 8 atomes de carbone.




2. Procédé pour la préparation d'un composé de formule (I), tel que défini dans la revendication 1, caractérisé en ce qu'on oxyde un composé de formule (X), ou, lorsque R_5 est un atome d'hydrogène, un sel métallique de celui-ci:



(X)


dans laquelle R_1' et R_2' ainsi que R_3 à R_5 et n sont tels que définis dans la revendication 1, et où le cycle contenant de l'azote est en position trans par rapport au groupe OR_5 ; dans le cas où l'un des R_1' et R_2' est un groupe ou un atome pouvant être converti en un atome d'hydrogène, on convertit le groupe ou l'atome en un atome d'hydrogène; dans le cas où l'autre des R_1' et R_2' est un groupe ou un atome pouvant être converti en l'un des substituants de la classe définie dans la revendication 1 pour l'autre des R_1 et R_2 , on convertit le groupe ou l'atome en l'un des substituants de la classe définie plus haut; et, dans le cas où R_5 est un atome d'hydrogène dans le composé de formule (I), on alcoyle ou acyle éventuellement avec un agent alcoylant ayant 1 à 3 atomes de carbone ou un agent acylant ayant 1 à 8 atomes de carbone.

3. Procédé pour la préparation d'un composé de formule (I), tel que défini dans la revendication 1, caractérisé en ce qu'on fait réagir un composé de formule (IV):



(IV)

dans laquelle R_1' et R_2' ainsi que R_3 et R_4 sont tels que définis dans la revendication 1, avec un anion de formule (XIII):




(XIII)

dans laquelle n est tel que défini dans la revendication 1; dans le cas où l'un des R_1' et R_2' est un groupe ou un atome pouvant être converti en un atome d'hydrogène, on convertit le groupe ou l'atome en un atome d'hydrogène; dans le cas où l'autre des R_1' et R_2' est un groupe ou un atome pouvant être converti en l'un des substituants de la classe définie pour l'autre des R_1 et R_2 , on convertit le groupe ou l'atome en l'un des substituants de la classe définie plus haut; et, dans le cas où R_5 est un atome d'hydrogène dans le composé de formule (I), on alcoyle ou acyle éventuellement avec un agent alcoylant ayant 1 à 3 atomes de carbone ou un agent acylant ayant 1 à 8 atomes de carbone.



4. Procédé pour la préparation d'un composé de formule (I), tel que défini dans la revendication 1, caractérisé en ce qu'on cyclise un composé de formule (XIV):



(XIV)

dans laquelle R_1' et R_2' ainsi que R_3 à R_5 et n sont tels que définis dans la revendication 1 et L_3 est un groupe mobile, et où le groupe amino substitué est en position trans par rapport au groupe $-OR_5$; dans le cas où l'un des R_1' et R_2' est un groupe ou un atome pouvant être converti en un atome d'hydrogène, on convertit le groupe ou l'atome en un atome d'hydrogène; dans le cas où l'autre des R_1' et R_2' est un groupe ou un atome pouvant être converti en l'un des substituants de la classe définie dans la revendication 1 pour l'autre des R_1 et R_2 , on convertit le groupe ou l'atome en l'un des substituants de la classe définie plus haut; et, dans le cas où R_5 est un atome d'hydrogène dans le composé de formule (I), on alcoyle ou acyle éventuellement avec un agent alcoylant ayant 1 à 3 atomes de carbone ou un agent acylant ayant 1 à 8 atomes de carbone.


5. Procédé suivant l'une quelconque des revendications 1 à 4, caractérisé en ce que l'une des R_1 et R_2 est un atome d'hydrogène et l'autre est choisi dans la classe des groupes alcoylcarbonyle, alcoxycarbonyle, alcoylcarbonyloxy, alcoylhydroxyméthyle, nitro, chloro ou cyano.

6. Procédé suivant l'une quelconque des revendications 1 à 5, caractérisé en ce que les groupes alcoyles ou les parties alcoyles des radicaux contenant des groupes alcoyles concernant l'autre des R_1 et R_2 sont des groupes méthyle ou éthyle.

7. Procédé suivant l'une quelconque des revendications précédentes, caractérisé en ce que R_3 et R_4 sont tous deux des groupes alcoyles ayant 1 à 4 atomes de carbone.

8. Procédé suivant l'une quelconque des revendications précédentes, caractérisé en ce que R_5 est un atome d'hydrogène.

9. Procédé suivant l'une quelconque des revendications 1 à 4, pour la préparation d'un composé de formule (II):



(II)

caractérisé en ce que l'un des R_1 et R_2 est un atome d'hydrogène et l'autre est choisi dans la classe des groupes alcoylcarbonyle, alcoxycarbonyle, alcoylcarbonyloxy, alcoylhydroxyméthyle, nitro ou cyano, les groupes alcoyles ou les parties alcoyles des radicaux contenant des groupes alcoyles étant des groupes méthyle ou éthyle, et n est 1 ou 2; le groupe lactame étant en position trans par rapport au groupe OH.

10. Procédé suivant l'une quelconque des revendications précédentes, caractérisé en ce que l'un des R_1 et R_2 est un atome d'hydrogène et l'autre est un groupe nitro ou cyano.

11. Procédé suivant l'une quelconque des revendications précédentes, caractérisé en ce que R_2 est un atome d'hydrogène.

12. Procédé suivant la revendication 11, dans lequel R_1 est un groupe cyano et R_3 et R_4 sont tous deux un groupe méthyle.



13. Procédé suivant l'une quelconque des revendications 1 à 4, pour la préparation du 6 - cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pipéridinyl) - 2H - benzo[b]pyran - 3 - ol; du 7 - nitro - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; de l'acétate de 6 - cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - yle; du 6 - nitro - 3,4 - dihydro - 2 - méthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; du 6 - carbométhoxy - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; du 6 - carbométhoxy - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pipéridinyl) - 2H - benzo[b]pyran - 3 - ol; du 6 - chloro - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; du 6 - acétyl - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol; ou du 6 - acétyl - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pipéridinyl) - 2H - benzo[b]pyran - 3 - ol.
14. Procédé suivant l'une quelconque des revendications 1 à 4, pour la préparation du 6 - cyano - 3,4 - dihydro - 2,2 - diméthyl - trans - 4 - (2 - oxo - 1 - pyrrolidinyl) - 2H - benzo[b]pyran - 3 - ol.
15. Procédé suivant la revendication 14, dans lequel le composé résultant est sous la forme d'un solide.
16. Procédé suivant la revendication 15, dans lequel le composé résultant est sous la forme d'un solide cristallin.
17. Procédé suivant la revendication 15 ou 16, dans lequel le composé résultant a une gamme de point de fusion entre 226 et 231°C inclus.
18. Procédé suivant la revendication 17, dans lequel le produit résultant a un point de fusion de 230 à 231°C.
19. Procédé suivant l'une quelconque des revendications 1 à 18, caractérisé en ce que R_1' et R_2' sont R_1 et R_2 respectivement.
20. Procédé suivant l'une quelconque des revendications 1 à 19, caractérisé en ce que le composé de formule (I) est isolé sous une forme acceptable du point de vue pharmaceutique.

