

PYRIMIDINE COMPOUNDS CONTAINING ACIDIC GROUPS

Cross Reference to Related Applications

[0001] This application claims the benefit of U.S. Provisional Application No. 62/430,183, filed December 5, 2016, and U.S. Provisional Application No. 62/532,230, filed July 13, 2017, the contents of which are incorporated herein by reference in their entireties.

Field of the Invention

[0002] The present disclosure relates to a class of pyrimidine derivatives having immunomodulating properties that act via TLR7 which are useful in the treatment of viral or allergic diseases and cancers.

Background of the Invention

[0003] The present disclosure relates to pyrimidine derivatives, processes for their preparation, pharmaceutical compositions containing them and their use in therapy.

[0004] The immune system is comprised of innate and acquired immunity, both of which work cooperatively to protect the host from microbial infections. It has been shown that innate immunity can recognize conserved pathogen-associated molecular patterns through toll-like receptors (TLRs) expressed on the cell surface of immune cells. Recognition of invading pathogens then triggers cytokine production (including interferon alpha(IFN α)) and upregulation of co-stimulatory molecules on phagocytes, leading to modulation of T cell function. Thus, innate immunity is closely linked to acquired immunity and can influence the development and regulation of an acquired response.


[0005] TLRs are a family of type I transmembrane receptors characterized by an NH₂-terminal extracellular leucine-rich repeat domain (LRR) and a COOH-terminal intracellular tail containing a conserved region called the Toll/IL-1 receptor (TIR) homology domain. The extracellular domain contains a varying number of LRR, which are thought to be involved in ligand binding. Eleven TLRs have been described to date in humans and mice. They differ from each other in ligand specificities, expression patterns, and in the target genes they can induce.

[0006] Ligands which act via TLRs (also known as immune response modifiers (IRMS)) have been developed, for example, the imidazoquinoline derivatives described in US Patent No. 4689338 which include the product Imiquimod for treating genital warts, and the adenine derivatives described in WO 98/01448 and WO 99/28321.

[0007] Compounds with liver targeting property are desirable. Certain moieties that aid in liver targeting that have been disclosed in references include acidic moieties. (Tu et al., Current Topics in Medicinal Chemistry, 2013, 13, 857-866; Oballa et al., J. Med. Chem., 2011, 54, 5082-5096; Pfefferkorn et al., J. Med. Chem., 2012, 55, 1318-1333, the contents of which are incorporated herein by reference in their entireties)


Summary of the Invention

[0008] The present disclosure provides a compound having the structure of Formula (1), and pharmaceutically acceptable salts thereof,




wherein

R^{1a} is selected from the group consisting of H, C₁-C₄ alkyl, -NH₂, -NHAc,

-COOH, -SO₂CH₃, -SCH₃, -OCH₃, ; and A,

wherein the alkyl is optionally substituted with -OH, -NH₂, -NHAc, -COOH, -SO₂CH₃,



-SCH₃, -OCH₃, , or A;

R^{1b} is C₂-C₅ alkyl;

X is selected from the group consisting of H and C₁-C₄ alkyl, wherein the alkyl is optionally substituted with A, -OH, or -C(CH₃)₂OH;

L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , -O-, -S-, -SO₂-, -NH-, and -CH₂CH₂-;

Y is selected from the group consisting of C₁-C₃ alkyl, aryl, and heteroaryl, wherein the alkyl, aryl, and heteroaryl are optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy;




m is an integer from zero to four; and

n is an integer from one to four; and


wherein the compound is substituted with at least one A.

[0009] The present disclosure provides a compound having the structure of Formula (1), and pharmaceutically acceptable salts thereof,




wherein

R^{1a} is selected from the group consisting of H, C_1-C_4 alkyl, $-\text{NH}_2$, $-\text{NHAc}$,



wherein the alkyl is optionally substituted with $-\text{OH}$, $-\text{NH}_2$, $-\text{NHAc}$, $-\text{COOH}$, $-\text{SO}_2\text{CH}_3$,










R^{1b} is C_2-C_5 alkyl;

X is selected from the group consisting of H and C_1-C_4 alkyl, wherein the alkyl is optionally substituted with A, $-\text{OH}$, or $-\text{C}(\text{CH}_3)_2\text{OH}$;

L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, , , $-\text{O}-$, $-\text{S}-$, $-\text{SO}_2-$, $-\text{NH}-$, and $-\text{CH}_2\text{CH}_2-$;

Y is selected from the group consisting of C_1 - C_3 alkyl, aryl, and heteroaryl, wherein the alkyl, aryl, and heteroaryl are optionally substituted with 1-5 substituents that are independently selected from A , C_1 - C_3 alkyl, and C_1 - C_3 alkoxy;

A is selected from the group consisting of , , , , , , and ,


L^2 is selected from the group consisting of a bond, $-(\text{CH}_2)_n-$, $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, , , $-(\text{CH}_2)_m-$, $-(\text{CH}_2)_n-$, $-\text{O}(\text{CH}_2\text{CH}_2)_n-$, $-\text{O}(\text{C}_1\text{-C}_4 \text{ alkylene})-$, $-\text{O}(\text{CH}_2\text{CH}_2)_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$, $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{O}(\text{CH}_2\text{CH}_2)_m-$, and $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{O}(\text{CH}_2\text{CH}_2)_m-\text{OCH}_2\text{CH}_2\text{CF}_2-$;

m is an integer from zero to four; and

n is an integer from one to four; and

wherein the compound is substituted with at least one A .

[0010] The present disclosure provides a compound having the structure of Formula (1), and pharmaceutically acceptable salts thereof,



wherein







R^{1a} is selected from the group consisting of H, C_1 - C_4 alkyl, $-\text{NH}_2$, $-\text{COOH}$, and $-\text{SO}_2\text{CH}_3$, wherein the alkyl is optionally substituted with $-\text{OH}$, $-\text{NH}_2$, $-\text{COOH}$, or $-\text{SO}_2\text{CH}_3$;


R^{1b} is C_2 - C_5 alkyl;

X is selected from the group consisting of H and C₁-C₄ alkyl, wherein the alkyl is optionally substituted with A;

L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , -O-, -S-, -SO₂-, -NH-, and -CH₂CH₂-;

Y is selected from the group consisting of C₁-C₃ alkyl, aryl, and heteroaryl, wherein the alkyl, aryl, and heteroaryl are optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy;

A is selected from the group consisting of , , , , , and ;

L² is selected from the group consisting of a bond, -(CH₂)_n-, -C(O)NH(CH₂)_n-, , , , -[O(CH₂CH₂)]_n-, -[O(C₁-C₄ alkylene)]-, and -[O(CH₂CH₂)]_n-OCH₂CH₂CF₂-;

m is an integer from zero to four; and

n is an integer from one to four; and

wherein the compound is substituted with at least one A.

[0011] In certain embodiments, the compound having the structure of Formula (1) can have any one or more of the following features,

- when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and L² is -CH₂-; then A is not -L²-COOH;
- when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and L² is -CH₂-; then A is not -L²-COOH, except when R^{1a} comprises -COOH;
- when L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then X is not -CH₃;
- when L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then X is not -CH₃, except when R^{1a} comprises -COOH;
- when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; then L² is not -CH₂-;

- (f) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; then L² is not -CH₂-, except when R^{1a} comprises -COOH;
- (g) when X is -CH₃; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then L¹ is not -CH₂-;
- (h) when X is -CH₃; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then L¹ is not -CH₂-, except when R^{1a} comprises -COOH;
- (i) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then R^{1a} comprises COOH;
- (j) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted A and -OCH₃; and L² is -CH₂-; then A is not -L²-COOH;
- (k) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted A and -OCH₃; and L² is -CH₂-; then A is not -L²-COOH, except when R^{1a} comprises -COOH;
- (l) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted A and -OCH₃; and L² is -CH₂-; and R^{1a} does not comprise COOH; then A is not -L²-COOH.


[0012] In certain embodiments, the compound having the structure of Formula (1) can have any one or more of the following features,

- (m) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and L² is -CH₂-; then A is not -L²-COOH;
- (n) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and L² is -CH₂-; then A is not -L²-COOH, except when R^{1a} comprises -COOH or -SO₂CH₃;
- (o) when L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then X is not -CH₃;
- (p) when L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then X is not -CH₃, except when R^{1a} comprises -COOH or -SO₂CH₃;
- (q) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; then L² is not -CH₂-;
- (r) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; then L² is not -CH₂-, except when R^{1a} comprises -COOH or -SO₂CH₃;
- (s) when X is -CH₃; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then L¹ is not -CH₂-;
- (t) when X is -CH₃; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then L¹ is not -CH₂-, except when R^{1a} comprises -COOH or -SO₂CH₃;
- (u) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then R^{1a} comprises COOH or -SO₂CH₃;


- (v) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted A and -OCH₃; and L² is -CH₂-; then A is not -L²-COOH;
- (w) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted A and -OCH₃; and L² is -CH₂-; then A is not -L²-COOH, except when R^{1a} comprises -COOH or -SO₂CH₃;
- (x) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted A and -OCH₃; and L² is -CH₂-; and R^{1a} does not comprise COOH or -SO₂CH₃; then A is not -L²-COOH;
- (y) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then R^{1a} is not H or alkyl substituted with -OH.

[0013] In certain embodiments, the compound having the structure of Formula (1) can have any one or more of the following features:


- (aa) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; L² is -CH₂-, -O-

(CH₂)₂-O(CH₂)₂-, or -O-(CH₂)₂-O(CH₂)₂(CF₂)₂-; and A is $\frac{s}{2}$ -L²-, then A and L¹ are not in a para position with respect to each other;


- (bb) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; L² is -CH₂-; and A

is $\frac{s}{2}$ -L²-, then A and L¹ are not in a para position with respect to each other;


- (cc) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A and -OCH₃; L² is

-CH₂-; and A is $\frac{s}{2}$ -L²-, then A and L¹ are not in a para position with respect to each other;


- (dd) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; L² is -O-(CH₂)₂-

O(CH₂)₂-; and A is $\frac{s}{2}$ -L²-, then A and L¹ are not in a para position with respect to each other;


- (ee) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; L² is -O-(CH₂)₂-

O(CH₂)₂(CF₂)₂-; and A is $\frac{s}{2}$ -L²-, then A and L¹ are not in a para position with respect to each other;


- (ff) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A; and L² is -CH₂-, -O-(CH₂)₂-O(CH₂)₂-, or -O-(CH₂)₂-O(CH₂)₂(CF₂)-, then




- (gg) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted




- (hh) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted




- (ii) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted




- (jj) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted




- (kk) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A; L² is -CH₂-, -O-(CH₂)₂-O(CH₂)₂-, or -O-(CH₂)₂-O(CH₂)₂(CF₂)-; and A




- (ll) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A; L² is -CH₂-; and A and L¹ are para position with respect to each other;




- (mm) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A and -OCH₃; L² is -CH₂-; and A and L¹ are para position with respect to




- (nn) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A; L² is -O-(CH₂)₂-O(CH₂)₂-; and A and L¹ are para position with respect to

each other; then A is not ;

- (oo) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A; L² is -O-(CH₂)₂-O(CH₂)₂(CF₂)₂-; and A and L¹ are para position with

respect to each other; then A is not .








[0014] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1a),



wherein

X is H or CH₃;


L¹ is selected from the group consisting of a bond, -CH₂-, -O-, -S-, -CF₂-, , , and -CH₂CH₂-;

A is selected from the group consisting of , , , , , , and .

L² is selected from the group consisting of a bond, -CH₂-, -CH₂CH₂-, , , -C(O)NH(CH₂)_n-, -[O(CH₂CH₂)]_n-, -[O(C₁-C₄ alkylene)]-, and -[O(CH₂CH₂)]_n-OCH₂CH₂CF₂-; -C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-; and C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-OCH₂CH₂CF₂-; and

R^3 is H, C_1 - C_3 alkyl, or C_1 - C_3 alkoxy.

[0015] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1a),




wherein

X is H or CH_3 ;

L^1 is selected from the group consisting of a bond, $-CH_2-$, $-CF_2-$, , , and $-CH_2CH_2-$;


A is selected from the group consisting of $\frac{1}{2}L^2-COOH$, $\frac{1}{2}L^2-PO_3H_2$, $\frac{1}{2}L^2-SO_3H$,



L^2 is selected from the group consisting of a bond, $-CH_2-$, $-CH_2CH_2-$, , , $-C(O)NH(CH_2)_n-$, $-[O(CH_2CH_2)]_n-$, $-[O(C_1-C_4 \text{ alkylene})]_n-$, and $-[O(CH_2CH_2)]_n-OCH_2CH_2CF_2-$; and


R^3 is H, C_1 - C_3 alkyl, or C_1 - C_3 alkoxy.


[0016] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1a),





wherein


X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$;


A^{1a} is selected from the group consisting of ,



L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, , and $-\text{CH}_2\text{CH}_2-$;


A is selected from the group consisting of ,



L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1-\text{C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$; $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-$; and $\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and


R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.

[0017] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1a),




wherein

X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$;




L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , and -CH₂CH₂-;



L² is selected from the group consisting of a bond, -CH₂-, -CH₂CH₂-, , , -C(O)NH(CH₂)_n-, -[O(CH₂CH₂)]_n-, -[O(C₁-C₄ alkylene)]-, and -[O(CH₂CH₂)]_n-OCH₂CH₂CF₂-; and

R³ is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy.


[0018] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1b),



wherein

X is H or CH₃;


L¹ is selected from the group consisting of a bond, -CH₂-, -O-, -S-, -CF₂-, , , and -CH₂CH₂-;



L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{O}(\text{CH}_2\text{CH}_2)_n-$, $-\text{O}(\text{C}_1-\text{C}_4 \text{ alkylene})-$, and $-\text{O}(\text{CH}_2\text{CH}_2)_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$; $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{O}(\text{CH}_2\text{CH}_2)_m-$; and $\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{O}(\text{CH}_2\text{CH}_2)_m-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and

R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.


[0019] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1b),



wherein

X is H or CH_3 ;


L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, and $-\text{CH}_2\text{CH}_2-$;



L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1\text{-C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and

R^3 is H, $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy.


[0020] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1b),






wherein


X is $-\text{CH}_2\text{-A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{-A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{-A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2\text{-A}^{1a}$,

A^{1a} is selected from the group consisting of , , , ,



L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, , , and $-\text{CH}_2\text{CH}_2-$;

A is selected from the group consisting of , , ,



L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1\text{-C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$


OCH₂CH₂CF₂-;

-C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-; and C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-OCH₂CH₂CF₂-;

and




R³ is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy.


[0021] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1b),






wherein


X is -CH₂-A^{1a}, -CH₂CH₂-A^{1a}, -CH₂CH₂CH₂-A^{1a}, or -CH₂C(CH₃)₂-A^{1a};

A^{1a} is selected from the group consisting of , , , ,



L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , and -CH₂CH₂-;


A is selected from the group consisting of , , ,



L² is selected from the group consisting of a bond, -CH₂-, -CH₂CH₂-, , , -C(O)NH(CH₂)_n-, -[O(CH₂CH₂)]_n-, -[O(C₁-C₄ alkylene)]-, and -[O(CH₂CH₂)]_n-OCH₂CH₂CF₂-; and


R³ is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy.


[0022] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1c),




wherein

X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$,

A^{1a} is selected from the group consisting of ,




L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, ,



R^3 is H, C_1 - C_3 alkyl, or C_1 - C_3 alkoxy; and


R^4 is H or C_1 - C_3 alkoxy.

[0023] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1c),



wherein

X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$,




L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , and -CH₂CH₂-;


R³ is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy; and

R⁴ is H or C₁-C₃ alkoxy.

[0024] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1d),




X is -CH₂-A^{1a}, -CH₂CH₂-A^{1a}, -CH₂CH₂CH₂-A^{1a}, or -CH₂C(CH₃)₂-A^{1a},



L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , -O-, -CH₂CH₂-, and -S-; and


Y is H or C₁-C₃ alkyl.

[0025] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1d),



X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$,


A^{1a} is selected from the group consisting of , , , ,



L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, , , $-\text{O}-$, and $-\text{CH}_2\text{CH}_2-$; and

Y is H or C_1-C_3 alkyl.


[0026] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1e),




wherein

R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.

[0027] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1f),




[0028] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1g),



wherein

R³ is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy.


[0029] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1h),



wherein

R⁴ is H or C₁-C₃ alkoxy.

[0030] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1i),




wherein

L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, , , $-\text{O}-$, $-\text{CH}_2\text{CH}_2-$, and $-\text{S}-$; and

Y is H or C_1 - C_3 alkyl.

[0031] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1i),




wherein


L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, , , $-\text{O}-$, and $-\text{CH}_2\text{CH}_2-$; and

Y is H or C_1 - C_3 alkyl.

[0032] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1j),



[0033] The present disclosure provides the compound of Formula (1) that is a compound of Formula (1k),



[0034] The present disclosure provides pharmaceutical compositions comprising a compound of the present disclosure, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

[0035] The present disclosure provides a method of treating a condition associated with TLR7 modulation in a subject in need thereof, comprising administering to the subject an effective amount of a compound of the present disclosure, or a pharmaceutically acceptable salt thereof. In certain embodiments, the condition is viral infection or cancer.

[0036] The present disclosure provides a compound of the present disclosure, or a pharmaceutically acceptable salt thereof, for use in treating a condition associated with TLR7 modulation. In certain embodiments, the condition is viral infection or cancer.

[0037] The present disclosure provides use of a compound of the present disclosure, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating a condition associated with TLR7 modulation. In certain embodiments, the condition is viral infection or cancer.

Detailed Description

[0038] Although specific embodiments of the present disclosure are herein illustrated and described in detail, the invention is not limited thereto. The detailed descriptions are provided as exemplary of the present invention and should not be construed as constituting any limitation of the invention. Modifications will be obvious to those skilled in the art, and all modifications that do not depart from the spirit of the invention are intended to be included with the scope of the appended claims.

Definitions

[0039] The following definitions are meant to clarify, but not limit, the terms defined. If a particular term used herein is not specifically defined, such term should not be considered indefinite. Rather, terms are used within their accepted meanings.

[0040] The term “alkyl” as used herein refers to a straight or branched saturated hydrocarbon. For example, an alkyl group can have 1 to 8 carbon atoms (i.e., (C₁-C₈)alkyl) or 1 to 6 carbon atoms (i.e., (C₁-C₆ alkyl) or 1 to 4 carbon atoms.

[0041] The term “alkylene” as used herein refers to a straight or branched saturated hydrocarbon radical having two monovalent radical centers derived by the removal of two hydrogen atoms from the same or two different carbon atoms of a parent alkane. For example, an alkylene group can have 1 to 8 carbon atoms, 1 to 6 carbon atoms, or 1 to 4 carbon atoms.

[0042] The term “alkoxy” as used herein refers to an alkyl group attached to the parent molecular moiety through an oxygen atom.

[0043] The term “halo” or “halogen” as used herein refers to fluoro, chloro, bromo and iodo.

[0044] The term “aryl” as used herein refers to a single all carbon aromatic ring or a multiple condensed all carbon ring system wherein at least one of the rings is aromatic. For example, an aryl group can have 6 to 20 carbon atoms, 6 to 14 carbon atoms, or 6 to 12 carbon atoms. Aryl includes a phenyl radical. Aryl also includes multiple condensed ring systems (e.g., ring systems comprising 2, 3 or 4 rings) having about 9 to 20 carbon atoms in which at least one ring is aromatic and wherein the other rings may be aromatic or not aromatic (i.e., carbocycle). Such multiple condensed ring systems may be optionally substituted with one or more (e.g., 1, 2 or 3) oxo groups on any carbocycle portion of the multiple condensed ring system. The rings of the multiple condensed ring system can be connected to each other via fused, spiro and bridged bonds when allowed by valency requirements. It is to be understood that the point of attachment of a multiple condensed ring system, as defined above, can be at any position of the ring system including an aromatic or a carbocycle portion of the ring.

[0045] The term “heteroaryl” as used herein refers to a single aromatic ring that has at least one atom other than carbon in the ring, wherein the atom is selected from the group consisting of oxygen, nitrogen and sulfur; the term also includes multiple condensed ring systems that have at least one such aromatic ring, which multiple condensed ring systems are further described below. Thus, the term includes single aromatic rings of from about 1 to 6 carbon atoms and about 1-4 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur in the rings. The sulfur and nitrogen atoms may also be present in an oxidized form provided the ring is aromatic. The term also includes multiple condensed ring systems (e.g., ring systems comprising 2, 3 or 4 rings) wherein a heteroaryl group, as defined above, can be condensed with one or more rings selected from heteroaryls (to form for example a naphthyridinyl such as 1,8-naphthyridinyl), heterocycles, (to form for example a 1, 2, 3, 4-tetrahydronaphthyridinyl such as 1, 2, 3, 4-tetrahydro-1,8-naphthyridinyl), carbocycles

(to form for example 5,6,7, 8-tetrahydroquinolyl) and aryls (to form for example indazolyl) to form the multiple condensed ring system. Thus, a heteroaryl (a single aromatic ring or multiple condensed ring system) has about 1-20 carbon atoms and about 1-6 heteroatoms within the heteroaryl ring. Such multiple condensed ring systems may be optionally substituted with one or more (e.g., 1, 2, 3 or 4) oxo groups on the carbocycle or heterocycle portions of the condensed ring. The rings of the multiple condensed ring system can be connected to each other via fused, spiro and bridged bonds when allowed by valency requirements. It is to be understood that the individual rings of the multiple condensed ring system may be connected in any order relative to one another. It is also to be understood that the point of attachment of a multiple condensed ring system (as defined above for a heteroaryl) can be at any position of the multiple condensed ring system including a heteroaryl, heterocycle, aryl or carbocycle portion of the multiple condensed ring system and at any suitable atom of the multiple condensed ring system including a carbon atom and heteroatom (e.g., a nitrogen).

[0046] As used herein, the term “pharmaceutically acceptable” refers to carrier(s), diluent(s), excipient(s) or salt forms that are compatible with the other ingredients of the formulation and not deleterious to the recipient of the pharmaceutical composition.

[0047] As used herein, the term “pharmaceutical composition” refers to a compound of the present disclosure optionally admixed with one or more pharmaceutically acceptable carriers, diluents, excipients, or adjuvants. Pharmaceutical compositions preferably exhibit a degree of stability to environmental conditions so as to make them suitable for manufacturing and commercialization purposes.


[0048] As used herein, the terms “effective amount,” “therapeutic amount,” or “effective dose” refer to an amount of active ingredient sufficient to elicit the desired pharmacological or therapeutic effects, thus resulting in effective prevention or treatment of a disorder. Prevention of a disorder may be manifested by delaying or preventing the progression of the disorder, as well as delaying or preventing the onset of the symptoms associated with the disorder. Treatment of the disorder may be manifested by a decrease or elimination of symptoms, inhibition or reversal of the progression of the disorder, as well as any other contribution to the well-being of the patient.

[0049] The effective dose can vary, depending upon factors such as the condition of the patient, the severity of the symptoms of the disorder, and the manner in which the pharmaceutical composition is administered. Typically, to be administered in an effective dose, compounds are required to be administered in an amount of less than 30 mg/kg of

patient weight. Often, the compounds may be administered in an amount from less than about 1 mg/kg patient weight to less than about 100 µg/kg of patient weight, and occasionally between about 10 µg/kg to less than 100 µg/kg of patient weight. The foregoing effective doses typically represent that amount administered as a single dose, or as one or more doses administered over a 24 hours period. For human patients, the effective dose of the compounds may require administering the compound in an amount of at least about 1mg/24 hr/patient, but not more than about 2400 mg/24 hr/patient, and often not more than about 500 mg/ 24 hr/ patient.


Compounds

[0050] The present disclosure provides a compound having the structure of Formula (1), and pharmaceutically acceptable salts thereof,




wherein

R^{1a} is selected from the group consisting of H, C₁-C₄ alkyl, -NH₂, -NHAc,

-COOH, -SO₂CH₃, -SCH₃, -OCH₃, ; and A,

wherein the alkyl is optionally substituted with -OH, -NH₂, -NHAc, -COOH, -SO₂CH₃,



-SCH₃, -OCH₃, , or A;

R^{1b} is C₂-C₅ alkyl;

X is selected from the group consisting of H and C₁-C₄ alkyl, wherein the alkyl is optionally substituted with A, -OH, or -C(CH₃)₂OH;

L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , -O-, -S-, -SO₂-, -NH-, and -CH₂CH₂-;

Y is selected from the group consisting of C₁-C₃ alkyl, aryl, and heteroaryl, wherein the alkyl, aryl, and heteroaryl are optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy;




m is an integer from zero to four; and

n is an integer from one to four; and


wherein the compound is substituted with at least one A.

[0051] The present disclosure provides a compound having the structure of Formula (1), and pharmaceutically acceptable salts thereof,




wherein

R^{1a} is selected from the group consisting of H, $\text{C}_1\text{-C}_4$ alkyl, $-\text{NH}_2$, $-\text{NHAc}$,



wherein the alkyl is optionally substituted with $-\text{OH}$, $-\text{NH}_2$, $-\text{NHAc}$, $-\text{COOH}$, $-\text{SO}_2\text{CH}_3$,










R^{1b} is $\text{C}_2\text{-C}_5$ alkyl;

X is selected from the group consisting of H and $\text{C}_1\text{-C}_4$ alkyl, wherein the alkyl is optionally substituted with A, $-\text{OH}$, or $-\text{C}(\text{CH}_3)_2\text{OH}$;

L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, , , $-\text{O}-$, $-\text{S}-$, $-\text{SO}_2-$, $-\text{NH}-$, and $-\text{CH}_2\text{CH}_2-$;

Y is selected from the group consisting of C_1 - C_3 alkyl, aryl, and heteroaryl, wherein the alkyl, aryl, and heteroaryl are optionally substituted with 1-5 substituents that are independently selected from A , C_1 - C_3 alkyl, and C_1 - C_3 alkoxy;

A is selected from the group consisting of , , , , , , and ,


L^2 is selected from the group consisting of a bond, $-(\text{CH}_2)_n-$, $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, , , $-\text{C}(\text{CH}_3)_2-(\text{CH}_2)_m-\text{C}(\text{CH}_3)_2-(\text{CH}_2)_n-$, $-\text{O}(\text{CH}_2\text{CH}_2)_n-$, $-\text{O}(\text{C}_1\text{-C}_4 \text{ alkylene})-$, $-\text{O}(\text{CH}_2\text{CH}_2)_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$, $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{O}(\text{CH}_2\text{CH}_2)_m-$, and $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{O}(\text{CH}_2\text{CH}_2)_m-\text{OCH}_2\text{CH}_2\text{CF}_2-$;


m is an integer from zero to four; and


n is an integer from one to four; and


wherein the compound is substituted with at least one A .

[0052] As disclosed above, R^{1a} is selected from the group consisting of H , C_1 - C_4 alkyl,


$-\text{NH}_2$, $-\text{NHAc}$, $-\text{COOH}$, $-\text{SO}_2\text{CH}_3$, $-\text{SCH}_3$, $-\text{OCH}_3$, and , wherein the alkyl is optionally substituted with $-\text{OH}$, $-\text{NH}_2$, $-\text{NHAc}$, $-\text{COOH}$, $-\text{SO}_2\text{CH}_3$, $-\text{SCH}_3$, $-\text{OCH}_3$, or

. In certain embodiments, R^{1a} is $-\text{COOH}$, which is a selection among the

substituents of A . In certain embodiments, R^{1a} is , which is a selection among the substituents of A . In certain embodiments, R^{1a} is C_1 - C_4 alkyl, wherein the alkyl is substituted with $-\text{COOH}$, which is a selection among the substituents of A . In certain embodiments, R^{1a}

is C₁-C₄ alkyl, wherein the alkyl is substituted with , which is a selection among the substituents of A.

[0053] The present disclosure provides a compound having the structure of Formula (1), and pharmaceutically acceptable salts thereof,



wherein






R^{1a} is selected from the group consisting of H, C₁-C₄ alkyl, -NH₂, -COOH, and -SO₂CH₃, wherein the alkyl is optionally substituted with -OH, -NH₂, -COOH, or -SO₂CH₃;


R^{1b} is C₂-C₅ alkyl;

X is selected from the group consisting of H and C₁-C₄ alkyl, wherein the alkyl is optionally substituted with A;

L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , -O-, -S-, -SO₂-, -NH-, and -CH₂CH₂-;

Y is selected from the group consisting of C₁-C₃ alkyl, aryl, and heteroaryl, wherein the alkyl, aryl, and heteroaryl are optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy;

A is selected from the group consisting of , , , , , and ;

L² is selected from the group consisting of a bond, -(CH₂)_n-, -C(O)NH(CH₂)_n-, , , , -[O(CH₂CH₂)]_n-, -[O(C₁-C₄ alkylene)]-, and -[O(CH₂CH₂)]_n-OCH₂CH₂CF₂-;


m is an integer from zero to four; and

n is an integer from one to four; and

wherein the compound is substituted with at least one A.


[0054] In certain embodiments, R^{1b} is C_3 alkyl, C_4 alkyl, or C_5 alkyl. In certain embodiments, R^{1b} is $-(CH_2)_2CH_3$. In certain embodiments, R^{1b} is $-(CH_2)_3CH_3$.

[0055] In certain embodiments, R^{1a} is H. In certain embodiments, R^{1a} is C_1 - C_4 alkyl, optionally substituted with -OH. In certain embodiments, R^{1a} is C_1 - C_3 alkyl, optionally substituted with -OH. In certain embodiments, R^{1a} is $-CH_2C(CH_3)_2OH$. In certain




[0056] In certain embodiments, the stereocenter of the carbon bearing R^{1A} and R^{1B} is an (S)-configuration. In certain embodiments, the stereocenter of the carbon bearing R^{1A} and R^{1B} is an (R)-configuration.

[0057] In certain embodiments, R^{1a} is C_1 - C_4 alkyl, optionally substituted with -COOH. In certain embodiments, R^{1a} is C_1 - C_3 alkyl, optionally substituted with -COOH. In



[0058] In certain embodiments, R^{1a} is C_1 - C_4 alkyl, wherein the alkyl is optionally substituted with -OH, -NH₂, -COOH, or -SO₂CH₃. In certain embodiments, R^{1a} is -NH₂, -COOH, or -SO₂CH₃. In certain embodiments, R^{1a} is C_1 - C_4 alkyl, optionally substituted with -OH, -OCH₃, -SCH₃, or -SO₂CH₃.

[0059] In certain embodiments, X is C_1 - C_4 alkyl, wherein the alkyl is substituted with

A. In certain embodiments, X is C_1 - C_4 alkyl, wherein the alkyl is substituted with , wherein L² is a bond. In certain embodiments, X is CH₃.





[0060] In certain embodiments, X is H. In certain embodiments, X is C_1 - C_4 alkyl.

[0061] In certain embodiments, L¹ is -CH₂-, -CH₂CH₂-, or -O-. In certain embodiments, L¹ is -CH₂-. In certain embodiments, L¹ is -CH₂-, -CH₂CH₂-, -O-, or -S-.

[0062] In certain embodiments, L¹ is a bond, -CF₂-, , , -S-, -SO₂-, or -NH-.

[0063] In certain embodiments, Y is C₁-C₃ alkyl or aryl. In certain embodiments, Y is aryl, wherein the aryl is substituted with C₁-C₃ alkoxy. In certain embodiments, Y is aryl, wherein the aryl is substituted with A. In certain embodiments, Y is aryl, wherein the aryl is substituted with C₁-C₃ alkoxy and A.

[0064] In certain embodiments, Y is heteroaryl, wherein the heteroaryl is optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy. In certain embodiments, Y is C₁-C₃ alkyl, wherein the alkyl is optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy. In certain embodiments, Y is aryl, wherein the aryl is optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy.

[0065] In certain embodiments, A is . In certain embodiments, A is . In certain embodiments, A is . In certain embodiments, A is . In certain embodiments, A is . In certain embodiments, A is

[0066] In certain embodiments, L^2 is $-(CH_2)_n-$. In certain embodiments, L^2 is .


In certain embodiments, L^2 is . In certain embodiments, L^2 is $-C(O)NH(CH_2)_n-$.

[0067] In certain embodiments, L² is a bond, $\text{---}\frac{\zeta}{\zeta}\text{---}(\text{CH}_2)_m\text{---}\text{C}(\text{CH}_3)_2\text{---}(\text{CH}_2)_n\text{---}\frac{\zeta}{\zeta}\text{---}$, $\text{---}[\text{O}(\text{CH}_2\text{CH}_2)]_n\text{---}$, $\text{---}[\text{O}(\text{C}_1\text{---C}_4 \text{ alkylene})]_n\text{---}$, or $\text{---}[\text{O}(\text{CH}_2\text{CH}_2)]_n\text{---OCH}_2\text{CH}_2\text{CF}_2\text{---}$.

[0068] In certain embodiments, n is one or two. In certain embodiments, n is one. In certain embodiments, n is two. In certain embodiments, n is three. In certain embodiments, n is four.








[0069] In certain embodiments, m is zero. In certain embodiments, m is one. In certain embodiments, m is two. In certain embodiments, m is three. In certain embodiments, m is four.

[0070] The present disclosure provides a compound of Formula (1), having the structure of Formula (1a), and pharmaceutically acceptable salts thereof,



[0071] In certain embodiment of Formula (1a),
X is H or CH₃;

L¹ is selected from the group consisting of a bond, -CH₂-, -O-, -S-, -CF₂-, , , and -CH₂CH₂-;

A is selected from the group consisting of , , , , , , and ;

L² is selected from the group consisting of a bond, -CH₂-, -CH₂CH₂-, , , -C(O)NH(CH₂)_n-, -[O(CH₂CH₂)]_n-, -[O(C₁-C₄ alkylene)]-, and -[O(CH₂CH₂)]_n-OCH₂CH₂CF₂-;
-C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-; and C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-OCH₂CH₂CF₂-;
and


R³ is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy.

[0072] In another embodiment of Formula (1a),

X is H or CH₃;

L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , and -CH₂CH₂-;


A is selected from the group consisting of , , ,



L² is selected from the group consisting of a bond, -CH₂-, -CH₂CH₂-, , , -C(O)NH(CH₂)_n-, -[O(CH₂CH₂)]_n-, -[O(C₁-C₄ alkylene)]-, and -[O(CH₂CH₂)]_n-OCH₂CH₂CF₂-; and

R³ is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy.


[0073] As disclosed above, in certain embodiments, X may be -CH₂-A^{1a}, -CH₂CH₂-A^{1a}, -CH₂CH₂CH₂-A^{1a}, or -CH₂C(CH₃)₂-A^{1a}; wherein A^{1a} is selected from the group



selection among the substituents of A.

[0074] Accordingly, in certain embodiments, X may be -CH₂-A^{1a}, -CH₂CH₂-A^{1a}, -CH₂CH₂CH₂-A^{1a}, or -CH₂C(CH₃)₂-A^{1a}, wherein A^{1a} is A;


wherein A is selected from the group consisting of $\text{--L}^2\text{--COOH}$, $\text{--L}^2\text{--P(O)(OH)}_2$,



L^2 is a bond.

[0075] Accordingly, in certain embodiments, X may be $\text{--CH}_2\text{--A}^{1a}$, $\text{--CH}_2\text{CH}_2\text{--A}^{1a}$, $\text{--CH}_2\text{CH}_2\text{CH}_2\text{--A}^{1a}$, or $\text{--CH}_2\text{C(CH}_3)_2\text{--A}^{1a}$, wherein A^{1a} is A;

wherein A is selected from the group consisting of $\text{--L}^2\text{--COOH}$, $\text{--L}^2\text{--P(O)(OH)}_2$,




L^2 is a bond.


[0076] In another embodiment of Formula (1a),

X is $\text{--CH}_2\text{--A}^{1a}$, $\text{--CH}_2\text{CH}_2\text{--A}^{1a}$, $\text{--CH}_2\text{CH}_2\text{CH}_2\text{--A}^{1a}$, or $\text{--CH}_2\text{C(CH}_3)_2\text{--A}^{1a}$,


A^{1a} is selected from the group consisting of --COOH , --P(O)(OH)_2 , $\text{--SO}_3\text{H}$, --N=N--NH ,



L^1 is selected from the group consisting of a bond, $\text{--CH}_2\text{--}$, --O-- , --S-- , $\text{--CF}_2\text{--}$, $\text{--C(CH}_3)_2\text{--}$,



A is selected from the group consisting of $\frac{1}{2}$ -L²-COOH, $\frac{1}{2}$ -L²-P(OH)₂, $\frac{1}{2}$ -L²-SO₃H,




L² is selected from the group consisting of a bond, -CH₂-, -CH₂CH₂-, , , -C(O)NH(CH₂)_n-, -[O(CH₂CH₂)]_n-, -[O(C₁-C₄ alkylene)]-, and -[O(CH₂CH₂)]_n-OCH₂CH₂CF₂-; -C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-; and C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-OCH₂CH₂CF₂-; and

R³ is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy.

[0077] In another embodiment of Formula (1a),


X is -CH₂-A^{1a}, -CH₂CH₂-A^{1a}, -CH₂CH₂CH₂-A^{1a}, or -CH₂C(CH₃)₂-A^{1a};

A^{1a} is selected from the group consisting of $\frac{1}{2}$ -COOH, $\frac{1}{2}$ -P(OH)₂, $\frac{1}{2}$ -SO₃H, $\frac{1}{2}$ -N=N-NH, $\frac{1}{2}$ -N=N-N-CH₂-COOH, $\frac{1}{2}$ -N=N-N-CH₂-COOH, and $\frac{1}{2}$ -N=N-N-CH₂-COOH;



L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , and -CH₂CH₂-;


A is selected from the group consisting of $\frac{1}{2}$ -L²-COOH, $\frac{1}{2}$ -L²-P(OH)₂, $\frac{1}{2}$ -L²-SO₃H,



L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1-\text{C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and

R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.




[0078] The present disclosure provides a compound of Formula (1), having the structure of Formula (1b), and pharmaceutically acceptable salts thereof,



[0079] In certain embodiment of Formula (1b),

X is H or CH_3 ;

L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, , , and $-\text{CH}_2\text{CH}_2-$;

A is selected from the group consisting of $\text{--}\frac{1}{2}\text{--}L^2\text{--COOH}$, $\text{--}\frac{1}{2}\text{--}L^2\text{--P}(\text{OH})_2$, $\text{--}\frac{1}{2}\text{--}L^2\text{--SO}_3\text{H}$, , , and , and $\text{--}\frac{1}{2}\text{--}L^2\text{--B}(\text{OH})_2$;




L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1-\text{C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-$; and $\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and

R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.

[0080] In another embodiment of Formula (1b),

X is H or CH_3 ;



L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, , , and $-\text{CH}_2\text{CH}_2-$;




A is selected from the group consisting of $\text{--}\frac{1}{2}\text{--}L^2\text{--}\text{C}(=\text{O})\text{OH}$, $\text{--}\frac{1}{2}\text{--}L^2\text{--}\text{P}(\text{OH})_2$, $\text{--}\frac{1}{2}\text{--}L^2\text{--}\text{SO}_3\text{H}$, , , and ;

L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1\text{--C}_4\text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n\text{OCH}_2\text{CH}_2\text{CF}_2-$; and

R^3 is H, $\text{C}_1\text{--C}_3$ alkyl, or $\text{C}_1\text{--C}_3$ alkoxy.


[0081] As disclosed above, in certain embodiments, X may be $-\text{CH}_2\text{--}A^{1a}$, $-\text{CH}_2\text{CH}_2\text{--}A^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{--}A^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2\text{--}A^{1a}$; wherein A^{1a} is selected from the group

consisting of , , , , , , and  or A^{1a} is selected from the group consisting of , , ,

, , and . In the above, X comprises A^{1a} , which is a selection among the substituents of A.

[0082] Accordingly, in certain embodiments, X may be $-\text{CH}_2\text{--}A^{1a}$, $-\text{CH}_2\text{CH}_2\text{--}A^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{--}A^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2\text{--}A^{1a}$; wherein A^{1a} is A;


wherein A is selected from the group consisting of $\text{--L}^2\text{--COOH}$, $\text{--L}^2\text{--P(O)(OH)}_2$,



L^2 is a bond.

[0083] Accordingly, in certain embodiments, X may be $\text{--CH}_2\text{--A}^{1a}$, $\text{--CH}_2\text{CH}_2\text{--A}^{1a}$, $\text{--CH}_2\text{CH}_2\text{CH}_2\text{--A}^{1a}$, or $\text{--CH}_2\text{C(CH}_3)_2\text{--A}^{1a}$, wherein A^{1a} is A;

wherein A is selected from the group consisting of $\text{--L}^2\text{--COOH}$, $\text{--L}^2\text{--P(O)(OH)}_2$,




L^2 is a bond.


[0084] In another embodiment of Formula (1b),

X is $\text{--CH}_2\text{--A}^{1a}$, $\text{--CH}_2\text{CH}_2\text{--A}^{1a}$, $\text{--CH}_2\text{CH}_2\text{CH}_2\text{--A}^{1a}$, or $\text{--CH}_2\text{C(CH}_3)_2\text{--A}^{1a}$,


A^{1a} is selected from the group consisting of --COOH , --P(O)(OH)_2 , $\text{--SO}_3\text{H}$, --N=N--NH ,



L^1 is selected from the group consisting of a bond, $\text{--CH}_2\text{--}$, --O-- , --S-- , $\text{--CF}_2\text{--}$, $\text{--C(CH}_3)_2\text{--}$,



A is selected from the group consisting of $\text{--L}^2\text{--COOH}$, $\text{--L}^2\text{--P(OH)}_2$, $\text{--L}^2\text{--SO}_3\text{H}$,




L^2 is selected from the group consisting of a bond, $\text{--CH}_2\text{--}$, $\text{--CH}_2\text{CH}_2\text{--}$, $\text{--C(O)NH(CH}_2)_n\text{--}$, $\text{--[O(CH}_2\text{CH}_2)]_n\text{--}$, $\text{--[O(C}_1\text{--C}_4\text{ alkylene)]--}$, and $\text{--[O(CH}_2\text{CH}_2)]_n\text{--OCH}_2\text{CH}_2\text{CF}_2\text{--}$; $\text{--C(O)NHCH}_2\text{CH}_2\text{--[O(CH}_2\text{CH}_2)]_m\text{--}$; and $\text{C(O)NHCH}_2\text{CH}_2\text{--[O(CH}_2\text{CH}_2)]_m\text{--OCH}_2\text{CH}_2\text{CF}_2\text{--}$; and

R^3 is H, $\text{C}_1\text{--C}_3$ alkyl, or $\text{C}_1\text{--C}_3$ alkoxy.

[0085] In another embodiment of Formula (1b),


X is $\text{--CH}_2\text{--A}^{1a}$, $\text{--CH}_2\text{CH}_2\text{--A}^{1a}$, $\text{--CH}_2\text{CH}_2\text{CH}_2\text{--A}^{1a}$, or $\text{--CH}_2\text{C(CH}_3)_2\text{--A}^{1a}$;

A^{1a} is selected from the group consisting of --COOH , --P(OH)_2 , $\text{--SO}_3\text{H}$, --N=N--NH ,



L^1 is selected from the group consisting of a bond, $\text{--CH}_2\text{--}$, $\text{--CF}_2\text{--}$, and $\text{--CH}_2\text{CH}_2\text{--}$;


A is selected from the group consisting of $\text{--L}^2\text{--COOH}$, $\text{--L}^2\text{--P(OH)}_2$, $\text{--L}^2\text{--SO}_3\text{H}$,




L^2 is selected from the group consisting of a bond, $\text{--CH}_2\text{--}$, $\text{--CH}_2\text{CH}_2\text{--}$, $\text{--C(O)NH(CH}_2)_n\text{--}$, $\text{--[O(CH}_2\text{CH}_2)]_n\text{--}$, $\text{--[O(C}_1\text{--C}_4\text{ alkylene)]--}$, and $\text{--[O(CH}_2\text{CH}_2)]_n\text{--OCH}_2\text{CH}_2\text{CF}_2\text{--}$; and

R^3 is H, C_1 - C_3 alkyl, or C_1 - C_3 alkoxy.


[0086] The present disclosure provides a compound of Formula (1), having the structure of Formula (1c), and pharmaceutically acceptable salts thereof,



[0087] As disclosed above, in certain embodiments, X may be $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$, wherein A^{1a} is selected from the group




[0088] Accordingly, in certain embodiments, X may be $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$, wherein A^{1a} is A;



[0089] Accordingly, in certain embodiments, X may be $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$, wherein A^{1a} is A;

wherein A is selected from the group consisting of $\frac{1}{2}-L^2-\text{COOH}$, $\frac{1}{2}-L^2-\text{P}(\text{OH})_2$,




L^2 is a bond.


[0090] In certain embodiment of Formula (1c),

X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$,

A^{1a} is selected from the group consisting of $\frac{1}{2}-\text{COOH}$, $\frac{1}{2}-\text{P}(\text{OH})_2$, $\frac{1}{2}-\text{SO}_3\text{H}$, $\frac{1}{2}-\text{N}=\text{N}-\text{NH}$,



L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, $\frac{1}{2}-\text{C}(\text{CH}_3)_2-\frac{1}{2}$,




R^3 is H, C_1 - C_3 alkyl, or C_1 - C_3 alkoxy; and

R^4 is H or C_1 - C_3 alkoxy.

[0091] In another embodiment of Formula (1c),

X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$,

A^{1a} is selected from the group consisting of $\frac{1}{2}-\text{COOH}$, $\frac{1}{2}-\text{P}(\text{OH})_2$, $\frac{1}{2}-\text{SO}_3\text{H}$, $\frac{1}{2}-\text{N}=\text{N}-\text{NH}$,




L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, $\frac{1}{2}-\text{C}(\text{CH}_3)_2-\frac{1}{2}$, $\frac{1}{2}-\text{C}(\text{CH}_3)_2-\frac{1}{2}$, and $-\text{CH}_2\text{CH}_2-$;


R^3 is H, C_1 - C_3 alkyl, or C_1 - C_3 alkoxy; and

R⁴ is H or C₁-C₃ alkoxy.


[0092] The present disclosure provides a compound of Formula (1), having the structure of Formula (1d), and pharmaceutically acceptable salts thereof,



[0093] As disclosed above, in certain embodiments, X may be -CH₂-A^{1a}, -CH₂CH₂-A^{1a}, -CH₂CH₂CH₂-A^{1a}, or -CH₂C(CH₃)₂-A^{1a}; wherein A^{1a} is selected from the group




[0094] Accordingly, in certain embodiments, X may be -CH₂-A^{1a}, -CH₂CH₂-A^{1a}, -CH₂CH₂CH₂-A^{1a}, or -CH₂C(CH₃)₂-A^{1a}; wherein A^{1a} is A;



[0095] Accordingly, in certain embodiments, X may be -CH₂-A^{1a}, -CH₂CH₂-A^{1a}, -CH₂CH₂CH₂-A^{1a}, or -CH₂C(CH₃)₂-A^{1a}; wherein A^{1a} is A;

wherein A is selected from the group consisting of $\frac{1}{2}\text{-L}^2\text{-COOH}$, $\frac{1}{2}\text{-L}^2\text{-P(O)(OH)}_2$,




L^2 is a bond.

[0096] In certain embodiment of Formula (1d),

X is $-\text{CH}_2\text{-A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{-A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{-A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2\text{-A}^{1a}$,

A^{1a} is selected from the group consisting of $\frac{1}{2}\text{-COOH}$, $\frac{1}{2}\text{-P(O)(OH)}_2$, $\frac{1}{2}\text{-SO}_3\text{H}$, $\frac{1}{2}\text{-N=N-NH}$,




L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, $\frac{1}{2}\text{-C(CH}_3)_2$, $\frac{1}{2}\text{-C(CH}_3)_2$, $-\text{O-}$, $-\text{CH}_2\text{CH}_2$, and $-\text{S-}$; and

Y is H or $\text{C}_1\text{-C}_3$ alkyl.

[0097] In another embodiment of Formula (1d),

X is $-\text{CH}_2\text{-A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{-A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2\text{-A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2\text{-A}^{1a}$,


A^{1a} is selected from the group consisting of $\frac{1}{2}\text{-COOH}$, $\frac{1}{2}\text{-P(O)(OH)}_2$, $\frac{1}{2}\text{-SO}_3\text{H}$, $\frac{1}{2}\text{-N=N-NH}$,



L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, $\frac{1}{2}\text{-C(CH}_3)_2$, $\frac{1}{2}\text{-C(CH}_3)_2$, $-\text{O-}$, and $-\text{CH}_2\text{CH}_2-$; and


Y is H or $\text{C}_1\text{-C}_3$ alkyl.

[0098] The present disclosure provides a compound of Formula (1), having the structure of Formula (1e), and pharmaceutically acceptable salts thereof,




wherein R^3 is H, $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy.

[0099] The present disclosure provides a compound of Formula (1), having the structure of Formula (1f), and pharmaceutically acceptable salts thereof,




[00100] The present disclosure provides a compound of Formula (1), having the structure of Formula (1g), and pharmaceutically acceptable salts thereof,




wherein R^3 is H, $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy.

[00101] The present disclosure provides a compound of Formula (1), having the structure of Formula (1h), and pharmaceutically acceptable salts thereof,



wherein R^4 is H or $\text{C}_1\text{-C}_3$ alkoxy.

[00102] The present disclosure provides a compound of Formula (1), having the structure of Formula (1i), and pharmaceutically acceptable salts thereof,



wherein

L^1 is selected from the group consisting of a bond, $-CH_2-$, $-CF_2-$, , , $-O-$, $-CH_2CH_2-$, and $-S-$; and


Y is H or C_1-C_3 alkyl.

[00103] In another embodiment of Formula (1i),

L^1 is selected from the group consisting of a bond, $-CH_2-$, $-CF_2-$, , , $-O-$, and $-CH_2CH_2-$; and


Y is H or C_1-C_3 alkyl.

[00104] The present disclosure provides a compound of Formula (1), having the structure of Formula (1j), and pharmaceutically acceptable salts thereof,




[00105] In certain embodiment of Formula (1j), X is H or CH_3 .

[00106] In certain embodiment of Formula (1j), R^{1a} is selected from the group consisting of H and C_1-C_4 alkyl, wherein the alkyl is optionally substituted with $-OH$, $-NH_2$, -

$NHAc$, $-COOH$, $-SO_2CH_3$, $-SCH_3$, $-OCH_3$, or .

[00107] In certain embodiment of Formula (1j), R^{1a} is selected from the group consisting of H and C_1-C_4 alkyl, wherein the alkyl is optionally substituted with $-COOH$.

[00108] The present disclosure provides a compound of Formula (1), having the structure of Formula (1k), and pharmaceutically acceptable salts thereof,




[00109] In certain embodiment of Formula (1k), X is H or CH₃.

[00110] In certain embodiment of Formula (1k), R^{1b} is C₄ alkyl.


[00111] In certain embodiment of Formula (1k),

Y is aryl or heteroaryl, wherein the aryl and heteroaryl are optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy;

A is selected from the group consisting of $\frac{1}{2}$ -L²-COOH, $\frac{1}{2}$ -L²-P(O)(OH)₂, $\frac{1}{2}$ -L²-SO₃H,




L² is selected from the group consisting of a bond, -(CH₂)_n-, -C(O)NH(CH₂)_n-,



[00112] In certain embodiment of Formula (1k),


Y is aryl, wherein the aryl is optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy;



L^2 is selected from the group consisting of a bond, $-(CH_2)_n-$, $-C(O)NH(CH_2)_n-$, , , , $-\{-(CH_2)_m-\text{C}(\text{CH}_3)_2-(CH_2)_n-\}$, $-[O(CH_2CH_2)]_n-$, $-[O(C_1\text{-}C_4 \text{ alkylene})]_n-$, and $-[O(CH_2CH_2)]_n-OCH_2CH_2CF_2$.


[00113] In certain embodiment of Formula (1k),

Y is aryl, wherein the aryl is optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy;



L^2 is selected from the group consisting of a bond, $-(CH_2)_n$, , , and 


[00114] In one variation of Formula (1), R^{1b} is $-(CH_2)_2CH_3$ and R^{1a} is H. In one variation of Formula (1), R^{1b} is $-(CH_2)_2CH_3$ and R^{1a} is C₁-C₄ alkyl, optionally substituted with -OH. In one variation of Formula (1), R^{1b} is $-(CH_2)_2CH_3$ and R^{1a} is $-CH_2C(CH_3)_2OH$.



[00115] In one variation of Formula (1), Y is aryl, wherein the aryl is substituted with C₁-C₃ alkoxy and A. In one variation of Formula (1), Y is aryl, wherein the aryl is substituted with -OCH₃ and A.

[00116] In one variation of Formula (1), R^{1b} is $-(CH_2)_3CH_3$ and R^{1a} is C₁-C₄ alkyl, optionally substituted with -COOH. In one variation of Formula (1), R^{1b} is $-(CH_2)_2CH_3$ and R^{1a} is $-CH_2COOH$.

[00117] The present disclosure provides a compound of formula (1),




having one, two, three, or more of the following features:

- a) R^{1b} is $-(CH_2)_2CH_3$;

- b) R^{1a} is H;
- c) X is $-\text{CH}_3$;
- d) L^1 is $-\text{CH}_2-$; and
- e) Y is aryl, optionally substituted with 1-5 substituents that are independently selected from A, C_1 - C_3 alkyl, and C_1 - C_3 alkoxy.


[00118] The present disclosure provides a compound of formula (1),



having one, two, or three of the following features:

- a) R^{1b} is $-(\text{CH}_2)_2\text{CH}_3$ or $-(\text{CH}_2)_3\text{CH}_3$;
- b) R^{1a} is H; and
- c) X is C_1 - C_4 alkyl, wherein the alkyl is substituted with A.


[00119] The present disclosure provides a compound of formula (1),



having one, two, or three of the following features:

- a) R^{1b} is $-(\text{CH}_2)_2\text{CH}_3$ or $-(\text{CH}_2)_3\text{CH}_3$;
- b) R^{1a} is C_1 - C_4 alkyl, substituted with $-\text{OH}$, $-\text{NH}_2$, $-\text{COOH}$, or $-\text{SO}_2\text{CH}_3$; and
- c) X is C_1 - C_4 alkyl, wherein the alkyl is substituted with A.

[00120] The present disclosure provides a compound of formula (1),




having one, two, three, or more of the following features:

- a) R^{1b} is $-(\text{CH}_2)_2\text{CH}_3$ or $-(\text{CH}_2)_3\text{CH}_3$;
- b) R^{1a} is C_1 - C_4 alkyl, optionally substituted with $-\text{COOH}$;
- c) X is $-\text{CH}_3$;
- d) L^1 is $-\text{CH}_2-$; and


- e) Y is aryl, optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy.

[00121] The present disclosure provides a compound of formula (1),




having one, two, three, or more of the following features:

- R^{1b} is $-(CH_2)_2CH_3$ or $-(CH_2)_3CH_3$;
- R^{1a} is H or C₁-C₄ alkyl, optionally substituted with $-COOH$;
- X is $-CH_3$;
- L¹ is $-CH_2-$; and
- Y is aryl, optionally substituted with 1-5 substituents that are independently selected



[00122] The present disclosure provides a compound having the structure of Formula (1), and pharmaceutically acceptable salts thereof,



wherein



R^{1a} is selected from the group consisting of H, C₁-C₄ alkyl, $-NH_2$, $-COOH$, and $-SO_2CH_3$, wherein the alkyl is optionally substituted with $-OH$, $-NH_2$, $-COOH$, or $-SO_2CH_3$;

R^{1b} is C₂-C₅ alkyl;

X is selected from the group consisting of H and C₁-C₄ alkyl, wherein the alkyl is optionally substituted with A or halogen;

L¹ is selected from the group consisting of a bond, $-CH_2-$, $-CF_2-$, , , $-O-$, $-S-$, $-SO_2-$, $-NH-$, and $-CH_2CH_2-$;

Y is selected from the group consisting of C₁-C₃ alkyl, aryl, and heteroaryl, wherein the alkyl, aryl, and heteroaryl are optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy;



m is an integer from zero to four; and

n is an integer from one to four; and

wherein the compound is substituted with at least one A,

with any one or more of the following features,

- when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and L² is -CH₂-; then A is not -L²-COOH;
- when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and L² is -CH₂-; then A is not -L²-COOH, except when R^{1a} comprises -COOH;
- when L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then X is not -CH₃;
- when L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then X is not -CH₃, except when R^{1a} comprises -COOH;
- when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; then L² is not -CH₂-;
- when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; then L² is not -CH₂-, except when R^{1a} comprises -COOH;
- when X is -CH₃; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then L¹ is not -CH₂-;

- (h) when X is -CH₃; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then L¹ is not -CH₂-, except when R^{1a} comprises -COOH;
- (i) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then R^{1a} comprises COOH;
- (j) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted A and -OCH₃; and L² is -CH₂-; then A is not -L²-COOH;
- (k) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted A and -OCH₃; and L² is -CH₂-; then A is not -L²-COOH, except when R^{1a} comprises -COOH;
- (l) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted A and -OCH₃; and L² is -CH₂-; and R^{1a} does not comprise COOH; then A is not -L²-COOH.


[00123] In certain embodiments, the compound having the structure of Formula (1) can have any one or more of the following features,

- (m) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and L² is -CH₂-; then A is not -L²-COOH;
- (n) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and L² is -CH₂-; then A is not -L²-COOH, except when R^{1a} comprises -COOH or -SO₂CH₃;
- (o) when L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then X is not -CH₃;
- (p) when L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then X is not -CH₃, except when R^{1a} comprises -COOH or -SO₂CH₃;
- (q) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; then L² is not -CH₂-;
- (r) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; then L² is not -CH₂-, except when R^{1a} comprises -COOH or -SO₂CH₃;
- (s) when X is -CH₃; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then L¹ is not -CH₂-;
- (t) when X is -CH₃; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then L¹ is not -CH₂-, except when R^{1a} comprises -COOH or -SO₂CH₃;
- (u) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then R^{1a} comprises COOH or -SO₂CH₃;
- (v) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted A and -OCH₃; and L² is -CH₂-; then A is not -L²-COOH;
- (w) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted A and -OCH₃; and L² is -CH₂-; then A is not -L²-COOH, except when R^{1a} comprises -COOH or -SO₂CH₃;


- (x) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted A and -OCH₃; and L² is -CH₂-; and R^{1a} does not comprise COOH or -SO₂CH₃; then A is not -L²-COOH;
- (y) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and A is -L²-COOH; and L² is -CH₂-; then R^{1a} is not H or alkyl substituted with -OH.

[00124] In certain embodiments, the compound having the structure of Formula (1) can have any one or more of the following features:


- (aa) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; L² is -CH₂-, -O-(CH₂)₂-

O(CH₂)₂-, or -O-(CH₂)₂-O(CH₂)₂(CF₂)₂-, and A is ; then A and L¹ are not in a para position with respect to each other;


- (bb) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; L² is -CH₂-; and A is

; then A and L¹ are not in a para position with respect to each other;


- (cc) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A and -OCH₃; L² is

-CH₂-; and A is ; then A and L¹ are not in a para position with respect to each other;


- (dd) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; L² is -O-(CH₂)₂-

O(CH₂)₂-, and A is ; then A and L¹ are not in a para position with respect to each other;

- (ee) when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; L² is -O-(CH₂)₂-

O(CH₂)₂(CF₂)₂-, and A is ; then A and L¹ are not in a para position with respect to each other;

- (ff) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A; and L² is -CH₂-, -O-(CH₂)₂-O(CH₂)₂-, or -O-(CH₂)₂-O(CH₂)₂(CF₂)₂-, then A is not

.

- (gg) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A;

and L² is -CH₂-; then A is not $\frac{1}{2}\text{-L}^2\text{-}\overset{\text{O}}{\underset{\text{OH}}{\text{P}}}\text{-OH}$;

- (hh) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A

and -OCH₃; and L² is -CH₂-; then A is not $\frac{1}{2}\text{-L}^2\text{-}\overset{\text{O}}{\underset{\text{OH}}{\text{P}}}\text{-OH}$;

- (ii) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A;

and L² is -O-(CH₂)₂-O(CH₂)₂-; then A is not $\frac{1}{2}\text{-L}^2\text{-}\overset{\text{O}}{\underset{\text{OH}}{\text{P}}}\text{-OH}$;

- (jj) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A;

and L² is -O-(CH₂)₂-O(CH₂)₂(CF₂)₂-; then A is not $\frac{1}{2}\text{-L}^2\text{-}\overset{\text{O}}{\underset{\text{OH}}{\text{P}}}\text{-OH}$;

- (kk) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A; L² is -CH₂-, -O-(CH₂)₂-O(CH₂)₂-, or -O-(CH₂)₂-O(CH₂)₂(CF₂)₂-; and A and L¹ are

para position with respect to each other; then A is not $\frac{1}{2}\text{-L}^2\text{-}\overset{\text{O}}{\underset{\text{OH}}{\text{P}}}\text{-OH}$;

- (ll) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A; L² is -CH₂-; and A and L¹ are para position with respect to each other; then A is

not $\frac{1}{2}\text{-L}^2\text{-}\overset{\text{O}}{\underset{\text{OH}}{\text{P}}}\text{-OH}$;


- (mm) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A and -OCH₃; L² is -CH₂-; and A and L¹ are para position with respect to each

other; then A is not $\frac{1}{2}\text{-L}^2\text{-}\overset{\text{O}}{\underset{\text{OH}}{\text{P}}}\text{-OH}$;

- (nn) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A; L² is -O-(CH₂)₂-O(CH₂)₂-; and A and L¹ are para position with respect to each


other; then A is not $\frac{1}{2}\text{-L}^2\text{-}\overset{\text{O}}{\underset{\text{OH}}{\text{P}}}\text{-OH}$;

- (oo) when X is -CH₃; R^{1a} is H; R^{1b} is C₄alkyl; L¹ is -CH₂-; Y is aryl substituted with A; L² is -O-(CH₂)₂-O(CH₂)₂(CF₂)₂-; and A and L¹ are para position with respect to


each other; then A is not 



















[00125] In certain embodiments, the present disclosure provides a compound of formula (1) that is not disclosed in WO 2009/067081 (PCT/SE2008/051334). In certain embodiments, the present disclosure provides a compound of formula (1) that is not disclosed in WO 2012/031140 (PCT/US2011/050231).

[00126] In certain embodiments, the present disclosure provides a compound of



[00127] The present disclosure provides for the following compounds and pharmaceutically acceptable salts thereof.




Compound	-X	-L ¹ -Y	R ^{1a} 	Compound
1	-CH ₃		-(CH ₂) ₃ CH ₃	
2	-CH ₃		-(CH ₂) ₃ CH ₃	
3	-CH ₃		-(CH ₂) ₃ CH ₃	
4	-CH ₃		-(CH ₂) ₃ CH ₃	
5	-CH ₃		-(CH ₂) ₃ CH ₃	
6	-CH ₃		-(CH ₂) ₃ CH ₃	
7	-CH ₃		-(CH ₂) ₃ CH ₃	
8	-CH ₃			
9	-CH ₃		-(CH ₂) ₃ CH ₃	
























Compound	-X	-L ¹ -Y	$\begin{array}{c} R^{1a} \\ \\ \text{---} R^{1b} \end{array}$	Compound
10	-CH ₃		-(CH ₂) ₃ CH ₃	
11	-CH ₃		-(CH ₂) ₃ CH ₃	
12	-CH ₃		-(CH ₂) ₃ CH ₃	
13	-CH ₃		-(CH ₂) ₃ CH ₃	
14	-CH ₃		-(CH ₂) ₃ CH ₃	
15			-(CH ₂) ₃ CH ₃	
16	-CH ₃		-(CH ₂) ₃ CH ₃	
17	-CH ₃		-(CH ₂) ₃ CH ₃	























Compound	-X	-L ¹ -Y	R ^{1a} R ^{1b}	Compound
18	-CH ₃			
19			-(CH ₂) ₃ CH ₃	
20	-CH ₃			
20C	-CH ₃			
21		-OCH ₃		



























[00128] The present disclosure provides for the following compounds and pharmaceutically acceptable salts thereof.



Compound	-X	-L ¹ -Y	R ^{1a} R ^{1b}	Compound
22	-CH ₃			
23	-CH ₃			
24	-CH ₃		-(CH ₂) ₃ CH ₃	
25	-CH ₃		-(CH ₂) ₃ CH ₃	
26	-CH ₃		-(CH ₂) ₄ CH ₃	
27				
28	-CH ₃		-(CH ₂) ₃ CH ₃	

29				
30	-CH ₃		-(CH ₂) ₃ CH ₃	
31	-CH ₃		-(CH ₂) ₃ CH ₃	
32	-CH ₃			
33	-CH ₃			
34	-CH ₃			
35	-CH ₃			
36	-CH ₃			

37			$-(\text{CH}_2)_3\text{CH}_3$	
38				
39	$-\text{CH}_3$			
40			$-(\text{CH}_2)_3\text{CH}_3$	
41	$-\text{CH}_3$			
42	$-\text{CH}_3$			
43			$-(\text{CH}_2)_3\text{CH}_3$	
44	$-\text{CH}_3$			

45				
46				
47			$-(CH_2)_3CH_3$	
48				
49				
50				
51	$-CH_3$			

[00129] Unless otherwise stated, structures depicted herein are also meant to include compounds which differ only in the presence of one or more isotopically enriched atoms. For example, compounds having the present structure except for the replacement of a hydrogen atom by deuterium or tritium, or the replacement of a carbon atom by ^{13}C or ^{14}C , or the

replacement of a nitrogen atom by ^{15}N , or the replacement of an oxygen atom with ^{17}O or ^{18}O are within the scope of the present disclosure. Such isotopically labeled compounds are useful as research or diagnostic tools.


General Synthetic Methods

[00130] As noted herein, the present disclosure includes specific representative compounds, which are identified herein with particularity. The compounds of the present disclosure may be made by a variety of methods, including well-known standard synthetic methods. Illustrative general synthetic methods are set out below and then specific compounds of the present disclosure are prepared in the working Examples.

[00131] In all of the examples described below, protecting groups for sensitive or reactive groups are employed where necessary in accordance with general principles of synthetic chemistry. Protecting groups are manipulated according to standard methods of organic synthesis (T. W. Green and P. G. M. Wuts, *Protecting Groups in Organic Synthesis*, 3rd Edition, John Wiley & Sons, New York (1999)). These groups are removed at a convenient stage of the compound synthesis using methods that are readily apparent to those skilled in the art. The selection of processes as well as the reaction conditions and order of their execution shall be consistent with the preparation of compounds of the present disclosure.

[00132] A representative synthesis for subject compounds is shown in Scheme 1.

Scheme 1



[00133] In Scheme 1, compound of formula (G) is an embodiment wherein L^1 is $-\text{CH}_2-$ and Y is aryl, which is appropriately substituted. Also in Scheme 1, LG is a leaving group; and R is H or alkyl. Compounds of formula (A) and (B) are commercially available starting materials. Alternatively, compounds of formula (A) and (B) can be synthesized via a variety of different synthetic routes using commercially available starting materials and/or starting materials prepared by conventional synthetic methods.

[00134] With continued reference to Scheme 1, compounds of formula (C) may be prepared by reacting a compound of formula (A) with a base, such as sodium hydride, in a suitable solvent such as tetrahydrofuran or N,N-dimethylformamide at a temperature, for example, from 0°C to room temperature (20°C), followed by addition of a compound of formula (B). The reaction is then preferably heated at a temperature, for example, from 50°C to 100°C , optionally in the presence of an additive such as potassium iodide.

[00135] Compounds of formula (D) may be prepared by reacting a compound of formula (C) with guanidine or guanidine carbonate in a suitable solvent such as methanol or ethanol at a temperature, for example, in the range from 50°C to 150°C .


[00136] Compounds of formula (E) may be prepared by reacting a compound of formula (D) with phosphorous oxychloride, at a temperature, for example, from 50°C to 110°C .

[00137] Compounds of formula (F) may be prepared by reacting a compound of formula (E) with excess of an amine of formula $\text{R}^{1a}\text{R}^{1b}\text{NH}$, in a suitable solvent such as NMP, butanol or 1,2-dioxane at a temperature, for example, from 50°C to 150°C . Alternatively, the reaction can be performed in a microwave at a temperature, for example, from 50°C to 200°C .

[00138] Compounds of formula (G) may be prepared by reacting a compound of formula (F) with a reducing agent, such as lithium aluminum hydride, in a suitable solvent such as tetrahydrofuran at a temperature, for example, from 0°C to 60°C .

[00139] A representative synthesis for subject compounds is shown in Scheme 2.

Scheme 2



[00140] In Scheme 2, compound of formula (G) is an embodiment wherein L^1 is $-\text{CH}_2-$ and Y is aryl, which is appropriately substituted. Also in Scheme 2, R is H or alkyl.


[00141] With continued reference to Scheme 2, compounds of formula (H) may be prepared by reacting a compound of formula (G) with an oxidizing agent, such as manganese oxide, in a suitable solvent such as tetrahydrofuran or N,N-dimethylformamide at a temperature, for example, from 40°C to 100°C .

[00142] Compounds of formula (I) may be prepared by reacting a compound of formula (H) via a Wittig reaction with $\text{RO}-\text{C}(\text{O})-\text{CH}=\text{PPh}_3$. The reaction may be carried out in a suitable solvent, such as tetrahydrofuran, at a temperature, for example, from 50°C to 150°C .

[00143] Compounds of formula (J) may be prepared by the reduction of a compound of formula (I) under hydrogenation conditions. The reaction may be carried out with a catalyst such as palladium on carbon under a hydrogen atmosphere in a suitable solvent such as ethyl acetate at a temperature, for example, from 20°C to 100°C .

[00144] A representative synthesis for subject compounds is shown in Scheme 3.

Scheme 3



[00145] In Scheme 3, compound of formula (G) is an embodiment wherein L^1 is $-\text{CH}_2-$ and Y is aryl, which is appropriately substituted.


[00146] With continued reference to Scheme 3, compounds of formula (K) may be prepared by reacting a compound of formula (G) with a chlorinating reagent, such as thionyl chloride in a suitable solvent such as methylene chloride at a temperature, for example, from room temperature to 50°C .

[00147] Compounds of formula (L) may be prepared by reacting a compound of formula (K) with a cyanide salt, such as potassium cyanide, in a suitable solvent such as dimethylsulfoxide or N,N-dimethylformamide (or a mixture of both solvents) at a temperature, for example, from room temperature to 50°C .

[00148] Compounds of formula (M) may be prepared by reacting a compound of formula (L) with an azido reagent, such as trimethylsilyl azide, in an azide-nitrile cycloaddition. The reaction can be run in a suitable solvent such as NMP or dioxane at a temperature, for example, from 50°C to 150°C . The reaction may be done in the presence of catalyst, such as dibutyltin oxide.

[00149] Compounds of formula (N) may be prepared by reacting a compound of formula (M) with an alkylating agent, such as 2-bromoacetate, in a suitable solvent such as acetone at a temperature, for example, from 0°C to 60°C .

Scheme 4




[00150] In Scheme 4, compound of formula (G) is an embodiment wherein L^1 is $-\text{CH}_2-$ and Y is aryl, which is appropriately substituted. Preparation of compounds of formula (K) and (L) are described above.

[00151] With continued reference to Scheme 4, compounds of formula (O) may be prepared by hydrolyzing a compound of formula (L), such as with use of a base, such as potassium hydroxide, in a suitable solvent such as ethane-1,2-diol and water (or mixtures thereof) at a temperature, for example, from 50°C to 200°C .

[00152] A representative synthesis for subject compounds is shown in Scheme 5.

Scheme 5




[00153] In Scheme 5, compound of formula (G) is an embodiment wherein L^1 is $-\text{CH}_2-$ and Y is aryl, which is appropriately substituted. Preparation of compound of formula (K) is described above.

[00154] With continued reference to Scheme 5, compounds of formula (P) may be prepared by reacting a compound of formula (K) with triethylphosphite, in a suitable solvent or neat at a temperature, for example, from 50°C to 150°C.

[00155] Compounds of formula (Q) may be prepared by reacting a compound of formula (P) with reagents to remove the ethyl groups, such as bromotrimethylsilane, in a suitable solvent such as methylene chloride at a temperature, for example, from room temperature to 60°C.

[00156] A representative synthesis for subject compounds is shown in Scheme 6.

Scheme 6



[00157] In Scheme 6, compound of formula (F') is an embodiment wherein L¹ is –CH₂– and Y is aryl, which is appropriately substituted. Also in Scheme 6, LG¹ and LG² are leaving groups. Compounds of formula (A') and (B') are commercially available starting materials. Alternatively, compounds of formula (A') and (B') can be synthesized via a variety of different synthetic routes using commercially available starting materials and/or starting materials prepared by conventional synthetic methods.

[00158] With continued reference to Scheme 6, compounds of formula (C') may be prepared by reacting a compound of formula (A') with a base, such as sodium hydride, in a suitable solvent such as tetrahydrofuran or N,N-dimethylformamide at a temperature, for example, from 0°C to room temperature (20°C), followed by addition of a compound of formula (B). The reaction is then preferably heated at a temperature, for example, from 50°C to 100°C, optionally in the presence of an additive such as potassium iodide.


[00159] Compounds of formula (D') may be prepared by reacting a compound of formula (C') with guanidine or guanidine carbonate in a suitable solvent such as methanol or ethanol at a temperature, for example, in the range from 50°C to 150°C.

[00160] Compounds of formula (E') may be prepared by reacting a compound of formula (D') with phosphorous oxychloride, at a temperature, for example, from 50°C to 110°C.

[00161] Compounds of formula (F') may be prepared by reacting a compound of formula (E') with excess of an amine of formula $R^{1a}R^{1b}NH$, in a suitable solvent such as NMP, butanol or 1,2-dioxane at a temperature, for example, from 50°C to 150°C. Alternatively, the reaction can be performed in a microwave at a temperature, for example, from 50°C to 200°C.

[00162] A representative synthesis for subject compounds is shown in Scheme 7.

Scheme 7




[00163] In Scheme 7, compound of formula (F') is an embodiment wherein L^1 is $-CH_2-$ and Y is aryl, which is appropriately substituted. Also in Scheme 7, LG^2 is a leaving group.

[00164] With continued reference to Scheme 7, compounds of formula (G') may be prepared by reacting a compound of formula (F') with an azido reagent, such as trimethylsilyl azide, in an azide-nitrile cycloaddition. The reaction can be run in a suitable solvent such as NMP or dioxane at a temperature, for example, from 50°C to 150°C. The reaction may be done in the presence of catalyst, such as dibutyltin oxide.

[00165] A representative synthesis for subject compounds is shown in Scheme 8.

Scheme 8



[00166] In Scheme 8, compound of formula (F') is an embodiment wherein L^1 is $-CH_2-$ and Y is aryl, which is appropriately substituted. Also in Scheme 8, LG^2 is a leaving group.

[00167] With continued reference to Scheme 8, compounds of formula (I') may be prepared by reacting a compound of formula (F') with triethylphosphite, in a suitable solvent or neat at a temperature, for example, from 50°C to 150°C.

[00168] Compounds of formula (J) may be prepared by reacting a compound of formula (I') with reagents to remove the ethyl groups, such as bromotrimethylsilane, in a suitable solvent such as methylene chloride at a temperature, for example, from room temperature to 60°C.

Method of Treatment

[00169] The compounds of formula (1) and their pharmaceutically acceptable salts have activity as pharmaceuticals, in particular as modulators of toll-like receptor (especially TLR7) activity, and thus may be used in the treatment of:

1. respiratory tract: obstructive diseases of the airways including: asthma, including bronchial, allergic, intrinsic, extrinsic, exercise-induced, drug-induced (including aspirin and NSAID-induced) and dust- induced asthma, both intermittent and persistent and of all severities, and other causes of airway hyper-responsiveness; chronic obstructive pulmonary disease (COPD); bronchitis, including infectious and eosinophilic bronchitis; emphysema; bronchiectasis; cystic fibrosis; sarcoidosis; farmer's lung and related diseases; hypersensitivity pneumonitis; lung fibrosis, including cryptogenic fibrosing alveolitis, idiopathic interstitial pneumonias, fibrosis complicating anti-neoplastic therapy and chronic infection, including tuberculosis and aspergillosis and other fungal infections; complications of lung transplantation; vasculitic and thrombotic disorders of the lung vasculature, and pulmonary hypertension; antitussive activity including treatment of chronic cough associated with inflammatory and secretory conditions of the airways, and iatrogenic cough; acute and chronic rhinitis including rhinitis medicamentosa, and vasomotor rhinitis; perennial and seasonal allergic rhinitis including rhinitis nervosa (hay fever); nasal polyposis; acute viral infection including the common cold, and infection due to respiratory syncytial virus, influenza, coronavirus (including SARS) and adenovirus;
2. skin: psoriasis, atopic dermatitis, contact dermatitis or other eczematous dermatoses, and delayed-type hypersensitivity reactions; phyto- and photodermatitis; seborrhoeic dermatitis, dermatitis herpetiformis, lichen planus, lichen sclerosus et atrophica, pyoderma gangrenosum, skin sarcoid, discoid lupus erythematosus, pemphigus, pemphigoid, epidermolysis bullosa, urticaria, angioedema, vasculitides, toxic erythemas, cutaneous eosinophilias, alopecia areata, male-pattern baldness,

Sweet's syndrome, Weber-Christian syndrome, erythema multiforme; cellulitis, both infective and non-infective; panniculitis; cutaneous lymphomas, non-melanoma skin cancer and other dysplastic lesions, non-malignant skin cancer, basal cell carcinoma; actinic keratosis; drug-induced disorders including fixed drug eruptions;

3. eyes: blepharitis; conjunctivitis, including perennial and vernal allergic conjunctivitis; iritis; anterior and posterior uveitis; choroiditis; autoimmune, degenerative or inflammatory disorders affecting the retina; ophthalmitis including sympathetic ophthalmitis; sarcoidosis; infections including viral, fungal, and bacterial;

4. genitourinary: nephritis including interstitial and glomerulonephritis; nephrotic syndrome; cystitis including acute and chronic (interstitial) cystitis and Hunner's ulcer; acute and chronic urethritis, prostatitis, epididymitis, oophoritis and salpingitis; vulvovaginitis; Peyronie's disease; erectile dysfunction (both male and female);

5. allograft rejection: acute and chronic following, for example, transplantation of kidney, heart, liver, lung, bone marrow, skin or cornea or following blood transfusion; or chronic graft versus host disease;

6. other auto-immune and allergic disorders including rheumatoid arthritis, irritable bowel syndrome, inflammatory bowel syndrome, colitis, Crohn's disease, systemic lupus erythematosus, multiple sclerosis, Hashimoto's thyroiditis, Graves' disease, Addison's disease, diabetes mellitus, idiopathic thrombocytopaenic purpura, eosinophilic fasciitis, hyper-IgE syndrome, antiphospholipid syndrome and Sazary syndrome;

7. oncology: treatment of common cancers including liver, lung, bladder, gastrointestinal (including gastric, colorectal, esophageal, and rectal), head and neck, prostate, breast, lung, ovarian, pancreatic, bowel and colon, stomach, skin, and brain tumors and malignancies affecting the bone marrow (including the leukaemias) and lymphoproliferative systems, such as Hodgkin's and non-Hodgkin's lymphoma; including the prevention and treatment of metastatic disease and tumour recurrences, and paraneoplastic syndromes; and

8. infectious diseases: virus diseases such as genital warts, common warts, plantar warts, hepatitis B, hepatitis C, herpes simplex virus, molluscum contagiosum, variola, human immunodeficiency virus (HIV), human papilloma virus (HPV), cytomegalovirus (CMV), varicella zoster virus (VZV), rhinovirus, adenovirus, coronavirus, influenza, parainfluenza; bacterial diseases such as tuberculosis and

mycobacterium avium, leprosy; other infectious diseases, such as fungal diseases, chlamydia, Candida, aspergillus, cryptococcal meningitis, Pneumocystis carinii, cryptosporidiosis, histoplasmosis, toxoplasmosis, trypanosome infection and leishmaniasis.

[00170] The present disclosure provides the use of a compound of formula (1) or a pharmaceutically acceptable salt thereof as a vaccine adjuvant, used together with one or more antigens against the following diseases: HBV, HPV, meningitis, Tdap, flu, rabies, tuberculosis, malaria, Staphylococcus aureus infection, and cancers (tumor-associated antigen or neo-antigen).

[00171] Thus, the present disclosure provides a compound of formula (1) or a pharmaceutically-acceptable salt thereof as hereinbefore defined for use in therapy.

[00172] In a further aspect, the present disclosure provides the use of a compound of formula (1) or a pharmaceutically acceptable salt thereof as hereinbefore defined in the manufacture of a medicament for use in therapy.

[00173] The present disclosure provides a method of treating a condition associated with TLR7 modulation in a subject in need thereof, comprising administering to the subject an effective amount of a compound of the present disclosure, or a pharmaceutically acceptable salt thereof. The present disclosure also provides a compound of the present disclosure, or a pharmaceutically acceptable salt thereof, for use in treating a condition associated with TLR7 modulation. The present disclosure provides use of a compound of the present disclosure, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating a condition associated with TLR7 modulation. In certain embodiments, the condition is viral infection or cancer.

[00174] In the context of the present specification, the term "therapy" also includes "prophylaxis" unless there are specific indications to the contrary. The terms "therapeutic" and "therapeutically" should be construed accordingly.

[00175] Prophylaxis is expected to be particularly relevant to the treatment of persons who have suffered a previous episode of, or are otherwise considered to be at increased risk of, the disease or condition in question. Persons at risk of developing a particular disease or condition generally include those having a family history of the disease or condition, or those who have been identified by genetic testing or screening to be particularly susceptible to developing the disease or condition.

[00176] In particular, the compounds of the present disclosure (including pharmaceutically acceptable salts) may be used in the treatment of asthma, COPD, allergic

rhinitis, allergic conjunctivitis, atopic dermatitis, cancer, hepatitis B, hepatitis C, HIV, HPV, bacterial infections and dermatosis.

[00177] The present disclosure still further provides a method of treating, or reducing the risk of, a disease or condition comprising or arising from abnormal cell growth (e.g. a cancer), which method comprises administering to a patient in need thereof a therapeutically effective amount of a compound of formula (1) or a pharmaceutically acceptable salt thereof as hereinbefore defined.

[00178] The present disclosure also provides a method of treating, or reducing the risk of, an obstructive airways disease or condition (e.g. asthma or COPD) which comprises administering to a patient in need thereof a therapeutically effective amount of a compound of formula (1) or a pharmaceutically acceptable salt thereof as hereinbefore defined.

[00179] For the above-mentioned therapeutic uses the dosage administered will, of course, vary with the compound employed, the mode of administration, the treatment desired and the disorder indicated. For example, the daily dosage of the compound of the present disclosure, if inhaled, may be in the range from about 0.05 micrograms per kilogram body weight ($\mu\text{g/kg}$) to about 100 micrograms per kilogram body weight ($\mu\text{g/kg}$). Alternatively, if the compound is administered orally, then the daily dosage of the compound of the present disclosure may be in the range from about 0.01 micrograms per kilogram body weight ($\mu\text{g/kg}$) to about 100 milligrams per kilogram body weight (mg/kg).

Pharmaceutical Compositions

[00180] The compounds of formula (1) and pharmaceutically acceptable salts thereof may be used on their own but will generally be administered in the form of a pharmaceutical composition in which the formula (1) compound/salt (active ingredient) is in association with a pharmaceutically acceptable adjuvant, diluent or carrier. Conventional procedures for the selection and preparation of suitable pharmaceutical formulations are described in, for example, "Pharmaceuticals - The Science of Dosage Form Designs", M. E. Aulton, Churchill Livingstone, 1988.

[00181] In certain embodiments, the administration can be oral, intravenous, subcutaneous, intramuscular, intratumoral, intradermal, intranasal, inhaled, intravesicle, topical, sublingual, buccal, intrarectal, intrathecal, intracranial, or other forms of local delivery.

[00182] Depending on the mode of administration, the pharmaceutical composition will comprise from about 0.05 to about 99 %w (per cent by weight), more particularly from about 0.05 to about 80 %w, still more particularly from about 0.10 to about 70 %w, and even

more particularly from about 0.10 to about 50 %w, of active ingredient, all percentages by weight being based on total composition.

[00183] The present disclosure also provides a pharmaceutical composition comprising a compound of formula (1) or a pharmaceutically acceptable salt thereof as hereinbefore defined, in association with a pharmaceutically acceptable adjuvant, diluent or carrier.

[00184] The present disclosure further provides a process for the preparation of a pharmaceutical composition of the present disclosure which comprises mixing a compound of formula (1) or a pharmaceutically acceptable salt thereof as hereinbefore defined with a pharmaceutically acceptable adjuvant, diluent or carrier.

[00185] The pharmaceutical compositions may be administered topically (e.g. to the skin or to the lung and/or airways) in the form, e.g., of creams, solutions, suspensions, heptafluoroalkane (HFA) aerosols and dry powder formulations, for example, formulations in the inhaler device known as the Turbuhaler[®]; or systemically, e.g. by oral administration in the form of tablets, capsules, syrups, powders or granules; or by parenteral administration in the form of a sterile solution, suspension or emulsion for injection (including intravenous, subcutaneous, intramuscular, intravascular or infusion); or by rectal administration in the form of suppositories.

[00186] Dry powder formulations and pressurized HFA aerosols of the compounds of the present disclosure (including pharmaceutically acceptable salts) may be administered by oral or nasal inhalation. For inhalation, the compound is desirably finely divided. The finely divided compound preferably has a mass median diameter of less than 10 micrometres (μm), and may be suspended in a propellant mixture with the assistance of a dispersant, such as a C₈-C₂₀ fatty acid or salt thereof, (for example, oleic acid), a bile salt, a phospholipid, an alkyl saccharide, a perfluorinated or polyethoxylated surfactant, or other pharmaceutically acceptable dispersant.

[00187] The compounds of the present disclosure may also be administered by means of a dry powder inhaler. The inhaler may be a single or a multi dose inhaler, and may be a breath actuated dry powder inhaler.

[00188] One possibility is to mix the finely divided compound of the present disclosure with a carrier substance, for example, a mono-, di- or polysaccharide, a sugar alcohol, or another polyol. Suitable carriers are sugars, for example, lactose, glucose, raffinose, melezitose, lactitol, maltitol, trehalose, sucrose, mannitol; and starch. Alternatively the finely divided compound may be coated by another substance. The powder mixture may also be

dispensed into hard gelatin capsules, each containing the desired dose of the active compound.

[00189] Another possibility is to process the finely divided powder into spheres which break up during the inhalation procedure. This spheronized powder may be filled into the drug reservoir of a multidose inhaler, for example, that known as the Turbuhaler® in which a dosing unit meters the desired dose which is then inhaled by the patient. With this system the active ingredient, with or without a carrier substance, is delivered to the patient.

[00190] For oral administration the compound of the present disclosure may be admixed with an adjuvant or a carrier, for example, lactose, saccharose, sorbitol, mannitol; a starch, for example, potato starch, corn starch or amylopectin; a cellulose derivative; a binder, for example, gelatin or polyvinylpyrrolidone; and/or a lubricant, for example, magnesium stearate, calcium stearate, polyethylene glycol, a wax, paraffin, and the like, and then compressed into tablets. If coated tablets are required, the cores, prepared as described above, may be coated with a concentrated sugar solution which may contain, for example, gum arabic, gelatin, talcum and titanium dioxide. Alternatively, the tablet may be coated with a suitable polymer dissolved in a readily volatile organic solvent.

[00191] For the preparation of soft gelatin capsules, the compound of the present disclosure may be admixed with, for example, a vegetable oil or polyethylene glycol. Hard gelatin capsules may contain granules of the compound using either the above-mentioned excipients for tablets. Also liquid or semisolid formulations of the compound of the present disclosure may be filled into hard gelatin capsules.

[00192] Liquid preparations for oral application may be in the form of syrups or suspensions, for example, solutions containing the compound of the present disclosure, the balance being sugar and a mixture of ethanol, water, glycerol and propylene glycol. Optionally such liquid preparations may contain colouring agents, flavouring agents, saccharine and/or carboxymethylcellulose as a thickening agent or other excipients known to those skilled in art.

Combination Therapy

[00193] The compounds of the present disclosure (that is, compounds of formula (1) and pharmaceutically acceptable salts thereof) may also be administered in conjunction with other compounds used for the treatment of the above conditions.

[00194] The present disclosure therefore further relates to combination therapies wherein a compound of the present disclosure or a pharmaceutical composition or formulation comprising a compound of the present disclosure is administered concurrently or

sequentially or as a combined preparation with another therapeutic agent or agents, for the treatment of one or more of the conditions listed.

[00195] The anti-cancer treatment defined hereinbefore may be applied as a sole therapy or may involve, in addition to the compound of the present disclosure, conventional surgery or radiotherapy or chemotherapy. Such chemotherapy may include one or more of the following categories of anti-tumour agents:

- (i) other antiproliferative/antineoplastic drugs and combinations thereof, as used in medical oncology, such as alkylating agents (for example cis-platin, oxaliplatin, carboplatin, cyclophosphamide, nitrogen mustard, melphalan, chlorambucil, busulphan, temozolamide and nitrosoureas); antimetabolites (for example gemcitabine and antifolates such as fluoropyrimidines like 5-fluorouracil and tegafur, raltitrexed, methotrexate, cytosine arabinoside, and hydroxyurea); antitumour antibiotics (for example anthracyclines like adriamycin, bleomycin, doxorubicin, daunomycin, epirubicin, idarubicin, mitomycin-C, dactinomycin and mithramycin); antimitotic agents (for example vinca alkaloids like vincristine, vinblastine, vindesine and vinorelbine and taxoids like taxol and taxotere and polokinese inhibitors); and topoisomerase inhibitors (for example epipodophyllotoxins like etoposide and teniposide, amsacrine, topotecan and camptothecin);
- (ii) cytostatic agents such as antioestrogens (for example tamoxifen, fulvestrant, toremifene, raloxifene, droloxifene and idoxifene), antiandrogens (for example bicalutamide, flutamide, nilutamide and cyproterone acetate), LHRH antagonists or LHRH agonists (for example goserelin, leuprorelin and buserelin), progestogens (for example megestrol acetate), aromatase inhibitors (for example as anastrozole, letrozole, vorazole and exemestane) and inhibitors of 5 α -reductase such as finasteride;
- (iii) anti-invasion agents (for example c-Src kinase family inhibitors like 4-(6-chloro-2,3-methylenedioxyanilino)-7-[2-(4-methylpiperazin-1-yl)ethoxy]-5-tetrahydropyran-4-yloxyquinazoline (AZD0530; International Patent Application WO 01/94341) and N-(2-chloro-6-methylphenyl)-2-[6-[4-(2-hydroxyethyl)piperazin-1-yl]-2-methylpyrimidin-4-ylamino]thiazole-5-carboxamide (dasatinib, BMS-354825; J. Med. Chem. 2004, 47, 6658-6661), and metalloproteinase inhibitors like marimastat, inhibitors of urokinase plasminogen activator receptor function or antibodies to Heparanase);
- (iv) inhibitors of growth factor function: for example such inhibitors include growth factor antibodies and growth factor receptor antibodies (for example the anti-erbB2

antibody trastuzumab [Herceptin™], the anti-EGFR antibody panitumumab, the anti-erbB1 antibody cetuximab [Erbix, C225] and any growth factor or growth factor receptor antibodies disclosed by Stern et al. Critical reviews in oncology/haematology, 2005, Vol. 54, ppl 1-29); such inhibitors also include tyrosine kinase inhibitors, for example inhibitors of the epidermal growth factor family (for example EGFR family tyrosine kinase inhibitors such as N-(3-chloro-4-fluorophenyl)-7-methoxy-6-(3-morpholinopropoxy)quinazolin-4-amine (gefitinib, ZD 1839), N-(3-ethynylphenyl)-6,7-bis(2-methoxyethoxy)quinazolin-4-amine (erlotinib, OSI-774) and 6-acrylamido-N-(3-chloro-4-fluorophenyl)-7-(3-morpholinopropoxy)-quinazolin-4-amine (CI 1033), erbB2 tyrosine kinase inhibitors such as lapatinib, inhibitors of the hepatocyte growth factor family, inhibitors of the platelet-derived growth factor family such as imatinib, inhibitors of serine/threonine kinases (for example Ras/Raf signalling inhibitors such as farnesyl transferase inhibitors, for example sorafenib (BAY 43-9006)), inhibitors of cell signalling through MEK and/or AKT kinases, inhibitors of the hepatocyte growth factor family, c-kit inhibitors, abl kinase inhibitors, IGF receptor (insulin-like growth factor) kinase inhibitors; aurora kinase inhibitors (for example AZD1 152, PH739358, VX-680, MLN8054, R763, MP235, MP529, VX-528 AND AX39459) and cyclin dependent kinase inhibitors such as CDK2 and/or CDK4 inhibitors;

(v) antiangiogenic agents such as those which inhibit the effects of vascular endothelial growth factor, [for example the anti-vascular endothelial cell growth factor antibody bevacizumab (Avastin™) and VEGF receptor tyrosine kinase inhibitors such as 4-(4-bromo-2-fluoroanilino)-6-methoxy-7-(1-methylpiperidin-4-ylmethoxy)quinazoline (ZD6474; Example 2 within WO 01/32651), 4-(4-fluoro-2-methylindol-5-yloxy)-6-methoxy-7-(3-pyrrolidin-1-ylpropoxy)quinazoline (AZD2171; Example 240 within WO 00/47212), vatalanib (PTK787; WO 98/35985) and SU1 1248 (sunitinib; WO 01/60814), compounds such as those disclosed in International Patent Applications WO97/22596, WO 97/30035, WO 97/32856 and WO 98/13354 and compounds that work by other mechanisms (for example linomide, inhibitors of integrin $\alpha v \beta 3$ function and angiostatin)];

(vi) vascular damaging agents such as Combretastatin A4 and compounds disclosed in International Patent Applications WO 99/02166, WO 00/40529, WO 00/41669, WO 01/92224, WO 02/04434 and WO 02/08213;

- (vii) antisense therapies, for example those which are directed to the targets listed above, such as ISIS 2503, an anti-ras antisense;
- (viii) gene therapy approaches, including for example approaches to replace aberrant genes such as aberrant p53 or aberrant BRCA1 or BRCA2, GDEPT (gene-directed enzyme pro-drug therapy) approaches such as those using cytosine deaminase, thymidine kinase or a bacterial nitroreductase enzyme and approaches to increase patient tolerance to chemotherapy or radiotherapy such as multi-drug resistance gene therapy;
- (ix) immunotherapy approaches, including for example ex-vivo and in-vivo approaches to increase the immunogenicity of patient tumour cells, such as transfection with cytokines such as interleukin 2, interleukin 4 or granulocyte-macrophage colony stimulating factor, approaches to decrease T-cell anergy, approaches using transfected immune cells such as cytokine -transfected dendritic cells, approaches using cytokine -transfected tumour cell lines and approaches using anti-idiotypic antibodies;
- (x) Checkpoint inhibitors, including but not limited to antibodies to PD-1/PD-L1, CTLA-4, TIM-3, LAG-3, OX-40, GITR, VISTA, 4-1BB, CD40, TIGIT, BTLA;
- (xi) Kinase inhibitors, including but not limited to small molecule or monoclonal antibody inhibitors of BRAF, EGFR, ALK, RAS, RAF, VEGF, HER, c-MET, MEK, FGFR, BCR-ABL, PI3K;
- (xii) Inhibitors of cancer/immune metabolism, including but not limited to inhibitors of IDO, TDO, GLS, IDH, arginase, adenosine receptor, CD73, CD39;
- (xiii) Epigenetic modulators, including but not limited to inhibitors of HDAC, bromodomain, methyl transferase;
- (xiv) Developmental pathway modulator, including but not limited to Smo, Wnt, YAP;
- (xv) Other anti-cancer or immune-oncology biologics including but not limited to oncolytic virus, BCG, CART, cytokines; and
- (xv) Antibodies including but not limited to PD-1 antibody and PD-L1 antibody.

[00196] Furthermore, for the treatment of the inflammatory diseases, COPD, asthma and allergic rhinitis the compounds of the present disclosure may be combined with agents such as tumour necrosis factor alpha (TNF-alpha) inhibitors such as anti-TNF monoclonal antibodies (for example Remicade, CDP-870 and adalimumab) and TNF receptor immunoglobulin molecules (such as Enbrel); non-selective cyclo-oxygenase COX-1/COX-2

inhibitors whether applied topically or systemically (such as piroxicam, diclofenac, propionic acids such as naproxen, flubiprofen, fenoprofen, ketoprofen and ibuprofen, fenamates such as mefenamic acid, indomethacin, sulindac, azapropazone, pyrazolones such as phenylbutazone, salicylates such as aspirin), COX-2 inhibitors (such as meloxicam, celecoxib, rofecoxib, valdecoxib, lumarcocoxib, parecoxib and etoricoxib); corticosteroids; glucocorticosteroids (whether administered by topical, oral, intramuscular, intravenous, or intra-articular routes); beta agonists; anti-histamines; methotrexate, lefunomide; hydroxychloroquine, d-penicillamine, auranofin or other parenteral or oral gold preparations.

[00197] For the treatment of the infectious diseases, the compounds of the present disclosure may be combined with agents such as nucleoside antivirals; non-nucleoside antivirals, including but not limited inhibitors of entry, polymerase, reverse transcriptase, protease, integrase; monoclonal antibodies against specific viruses; siRNA therapies; antibiotics; and antifungals.

[00198] For the treatment of HBV, the compounds of the present disclosure may be combined with agents such as antiviral nucleosides.

[00199] The present disclosure still further relates to other innate immune agonists targeting the following classes of receptors, including, but not limited to, TLRs (Toll-like receptor); NLRs (Nod-like receptor); CLRs (C-type lectin receptor); RLRs (RIG-I like receptor); and STING (stimulator of interferon gene).

[00200] The present disclosure still further relates to the combination of a compound of the present disclosure and a leukotriene biosynthesis inhibitor, 5 -lipoxygenase (5-LO) inhibitor or 5-lipoxygenase activating protein (FLAP) antagonist such as; zileuton; ABT-761; fenleuton; tepoxalin; Abbott-79175; Abbott-85761; an N-(5-substituted)-thiophene-2-alkylsulfonamide; 2,6-di-tert-butylphenolhydrazones; a methoxytetrahydropyrans such as Zeneca ZD-2138; the compound SB-210661; a pyridinyl-substituted 2-cyanonaphthalene compound such as L-739,010; a 2-cyanoquinoline compound such as L-746,530; or an indole or quinoline compound such as MK-591, MK-886, and BAY x 1005.

[00201] The present disclosure further relates to the combination of a compound of the present disclosure and a receptor antagonist for leukotrienes (LTB₄, LTC₄, LTD₄, and LTE₄) selected from the group consisting of the phenothiazin-3-ones such as L-651,392; amidino compounds such as CGS-25019c; benzoxalamines such as ontazolast; benzenecarboximidamides such as BIIL 284/260; and compounds such as zafirlukast, ablukast, montelukast, pranlukast, verlukast (MK-679), RG-12525, Ro-245913, iralukast (CGP 45715A), and BAY x 7195.

[00202] The present disclosure still further relates to the combination of a compound of the present disclosure and a phosphodiesterase (PDE) inhibitor such as a methylxanthanine including theophylline and aminophylline; a selective PDE isoenzyme inhibitor including a PDE4 inhibitor an inhibitor of the isoform PDE4D, or an inhibitor of PDE5.

[00203] The present disclosure further relates to the combination of a compound of the present disclosure and a histamine type 1 receptor antagonist such as cetirizine, loratadine, desloratadine, fexofenadine, acrivastine, terfenadine, astemizole, azelastine, levocabastine, chlorpheniramine, promethazine, cyclizine, or mizolastine; applied orally, topically or parenterally.

[00204] The present disclosure still further relates to the combination of a compound of the present disclosure and a gastroprotective histamine type 2 receptor antagonist.

[00205] The present disclosure further relates to the combination of a compound of the present disclosure and an antagonist of the histamine type 4 receptor.

[00206] The present disclosure still further relates to the combination of a compound of the present disclosure and an alpha-1/alpha-2 adrenoceptor agonist vasoconstrictor sympathomimetic agent, such as propylhexedrine, phenylephrine, phenylpropanolamine, ephedrine, pseudoephedrine, naphazoline hydrochloride, oxymetazoline hydrochloride, tetrahydrozoline hydrochloride, xylometazoline hydrochloride, tramazoline hydrochloride or ethylnorepinephrine hydrochloride.

[00207] The present disclosure further relates to the combination of a compound of the present disclosure and an anticholinergic agent including muscarinic receptor (M1, M2, and M3) antagonists such as atropine, hyoscyne, glycopyrrrolate, ipratropium bromide, tiotropium bromide, oxitropium bromide, pirenzepine or telenzepine.

[00208] The present disclosure still further relates to the combination of a compound of the present disclosure together with a beta-adrenoceptor agonist (including beta receptor subtypes 1-4) such as isoprenaline, salbutamol, formoterol, salmeterol, terbutaline, orciprenaline, bitolterol mesylate, and pirbuterol.

[00209] The present disclosure further relates to the combination of a compound of the present disclosure and a chromone, such as sodium cromoglycate or nedocromil sodium.

[00210] The present disclosure still further relates to the combination of a compound of the present disclosure together with an insulin-like growth factor type I (IGF-I) mimetic.

[00211] The present disclosure still further relates to the combination of a compound of the present disclosure and a glucocorticoid, such as flunisolide, triamcinolone acetonide,

beclomethasone dipropionate, budesonide, fluticasone propionate, ciclesonide or mometasone furoate.

[00212] The present disclosure still further relates to the combination of a compound of the present disclosure together with an inhibitor of matrix metalloproteases (MMPs), i.e., the stromelysins, the collagenases, and the gelatinases, as well as aggrecanase; especially collagenase-1 (MMP-I), collagenase-2 (MMP-8), collagenase-3 (MMP-13), stromelysin-1 (MMP-3), stromelysin-2 (MMP-10), and stromelysin-3 (MMP-11) and MMP-9 and MMP-12.

[00213] The present disclosure still further relates to the combination of a compound of the present disclosure together with modulators of chemokine receptor function such as antagonists of CCR1, CCR2, CCR2A, CCR2B, CCR3, CCR4, CCR5, CCR6, CCR7, CCR8, CCR9, CCR10 and CCR11 (for the C-C family); CXCR1, CXCR2, CXCR3, CXCR4 and CXCR5 (for the C-X-C family) and CX3CR1 for the C-X3-C family.

[00214] The present disclosure still further relates to the combination of a compound of the present disclosure together with a cytokine or modulator of cytokine function, including alpha-, beta-, and gamma- interferon; interleukins (IL) including IL1 to 15, and interleukin antagonists or inhibitors, including agents which act on cytokine signalling pathways.

[00215] The present disclosure still further relates to the combination of a compound of the present disclosure together with an immunoglobulin (Ig) or Ig preparation or an antagonist or antibody modulating Ig function such as anti-IgE (omalizumab).

[00216] The present disclosure further relates to the combination of a compound of the present disclosure and another systemic or topically-applied anti-inflammatory agent, such as thalidomide or a derivative thereof, a retinoid, dithranol or calcipotriol.

[00217] The present disclosure further relates to the combination of a compound of the present disclosure together with an antibacterial agent such as a penicillin derivative, a tetracycline, a macrolide, a beta-lactam, a fluoroquinolone, metronidazole, an inhaled aminoglycoside; an antiviral agent including acyclovir, famciclovir, valaciclovir, ganciclovir, cidofovir, amantadine, rimantadine, ribavirin, zanamavir and oseltamavir; a protease inhibitor such as indinavir, nelfinavir, ritonavir, and saquinavir; a nucleoside reverse transcriptase inhibitor such as didanosine, lamivudine, stavudine, zalcitabine or zidovudine; or a non-nucleoside reverse transcriptase inhibitor such as nevirapine or efavirenz.

[00218] In a further aspect the present disclosure provides a combination (for example for the treatment of COPD, asthma or allergic rhinitis) of a compound of formula (1) or a

pharmaceutically acceptable salt thereof as hereinbefore defined and one or more agents independently selected from:

- a non-steroidal glucocorticoid receptor (GR-receptor) agonist;
- a selective β_2 adrenoceptor agonist (such as metaproterenol, isoproterenol, isoprenaline, albuterol, salbutamol, formoterol, salmeterol, terbutaline, orciprenaline, bitolterol mesylate, pirbuterol or indacaterol);
- a phosphodiesterase inhibitor (such as a PDE4 inhibitor);
- a protease inhibitor (such as a neutrophil elastase or matrix metalloprotease MMP- 12 inhibitor);
- a glucocorticoid;
- an anticholinergic agent;
- a modulator of chemokine receptor function (such as a CCR1 receptor antagonist); and
- an inhibitor of kinase function (such as the kinases p38 or IKK).

[00219] The present disclosure also provides a pharmaceutical product comprising, in combination, a preparation of a first active ingredient which is a compound of formula (1) or a pharmaceutically acceptable salt thereof as hereinbefore defined, and a preparation of a second active ingredient which is

- a non-steroidal glucocorticoid receptor (GR-receptor) agonist;
- a selective β_2 adrenoceptor agonist;
- a phosphodiesterase inhibitor;
- a protease inhibitor;
- a glucocorticoid;
- an anticholinergic agent;
- a modulator of chemokine receptor function; or
- an inhibitor of kinase function;

for simultaneous, sequential or separate use in therapy.

[00220] In another aspect, the present disclosure provides a kit comprising a preparation of a first active ingredient which is a compound of formula (1) or a pharmaceutically acceptable salt thereof as hereinbefore defined, and a preparation of a second active ingredient which is


- a non-steroidal glucocorticoid receptor (GR-receptor) agonist;
- a selective β_2 adrenoceptor agonist;
- a phosphodiesterase inhibitor;
- a protease inhibitor;

- a glucocorticoid;
- an anticholinergic agent;
- a modulator of chemokine receptor function; or
- an inhibitor of kinase function;

and instructions for the simultaneous, sequential or separate administration of the preparations to a patient in need thereof.

Exemplary Embodiments

[00221] Embodiment I-1. A compound having the structure of Formula (1), or a pharmaceutically acceptable salt thereof,



wherein


R^{1a} is selected from the group consisting of H, C_1 - C_4 alkyl, $-NH_2$, $-COOH$, and $-SO_2CH_3$, wherein the alkyl is optionally substituted with $-OH$, $-NH_2$, $-COOH$, or $-SO_2CH_3$;


R^{1b} is C_2 - C_5 alkyl;

X is selected from the group consisting of H and C_1 - C_4 alkyl, wherein the alkyl is optionally substituted with A;

L^1 is selected from the group consisting of a bond, $-CH_2-$, $-CF_2-$, , , $-O-$, $-S-$, $-SO_2-$, $-NH-$, and $-CH_2CH_2-$;

Y is selected from the group consisting of C_1 - C_3 alkyl, aryl, and heteroaryl, wherein the alkyl, aryl, and heteroaryl are optionally substituted with 1-5 substituents that are independently selected from A, C_1 - C_3 alkyl, and C_1 - C_3 alkoxy;



L^2 is selected from the group consisting of a bond, $-(CH_2)_n-$, $-C(O)NH(CH_2)_n-$, , , $-(CH_2)_m-$, , $-(CH_2)_n-$, $-[O(CH_2CH_2)]_n-$, $-[O(C_1-C_4 \text{ alkylene})]_n-$, and $-[O(CH_2CH_2)]_n-OCH_2CH_2CF_2-$;

m is an integer from zero to four; and

n is an integer from one to four; and

wherein the compound is substituted with at least one A; and

when X is $-CH_3$; L^1 is $-CH_2-$; Y is aryl substituted with A; and L^2 is $-CH_2-$; then A is not $-L^2-COOH$, except when R^{1a} comprises $-COOH$.



[00222] Embodiment I-2. The compound of Embodiment I-1, or a pharmaceutically acceptable salt thereof, wherein R^{1b} is $-(CH_2)_2CH_3$.

[00223] Embodiment I-3. The compound of Embodiment I-1, or a pharmaceutically acceptable salt thereof, wherein R^{1b} is $-(CH_2)_3CH_3$.

[00224] Embodiment I-4. The compound of any one of Embodiments I-1 to I-3, or a pharmaceutically acceptable salt thereof, wherein R^{1a} is C_1-C_4 alkyl, optionally substituted with $-OH$.


[00225] Embodiment I-5. The compound of any one of Embodiments I-1 to I-4, or a pharmaceutically acceptable salt thereof, wherein R^{1a} is $-CH_2C(CH_3)_2OH$.

[00226] Embodiment I-6. The compound of any one of Embodiments I-1 to I-4, or

a pharmaceutically acceptable salt thereof, wherein  is  or .

[00227] Embodiment I-7. The compound of any one of Embodiments I-1 to I-3, or a pharmaceutically acceptable salt thereof, wherein R^{1a} is C_1-C_4 alkyl, optionally substituted with $-COOH$.

[00228] Embodiment I-8. The compound of any one of Embodiments I-1 to I-3

and I-7, or a pharmaceutically acceptable salt thereof, wherein  is .

[00229] Embodiment I-9. The compound of any one of Embodiments I-1 to I-3, or a pharmaceutically acceptable salt thereof, wherein R^{1a} is H.

[00230] Embodiment I-10. The compound of any one of Embodiments I-1 to I-9, or a pharmaceutically acceptable salt thereof, wherein X is C₁-C₄ alkyl, wherein the alkyl is substituted with A.

[00231] Embodiment I-11. The compound of any one of Embodiments I-1 to I-10, or a pharmaceutically acceptable salt thereof, wherein X is C₁-C₄ alkyl, wherein the alkyl is

substituted with $\frac{X}{\text{---}}\text{---L}^2\text{---}\overset{\text{O}}{\parallel}\text{C}\text{---OH}$, wherein L² is a bond.

[00232] Embodiment I-12. The compound of any one of Embodiments I-1 to I-9, or a pharmaceutically acceptable salt thereof, wherein X is CH₃.

[00233] Embodiment I-13. The compound of any one of Embodiments I-1 to I-9, or a pharmaceutically acceptable salt thereof, wherein X is H.

[00234] Embodiment I-14. The compound of any one of Embodiments I-1 to I-13, or a pharmaceutically acceptable salt thereof, wherein L¹ is -CH₂-, -CH₂CH₂-, or -O-.

[00235] Embodiment I-15. The compound of any one of Embodiments I-1 to I-14, or a pharmaceutically acceptable salt thereof, wherein L¹ is -CH₂-.

[00236] Embodiment I-16. The compound of any one of Embodiments I-1 to I-15, or a pharmaceutically acceptable salt thereof, wherein Y is C₁-C₃ alkyl or aryl.

[00237] Embodiment I-17. The compound of any one of Embodiments I-1 to I-16, or a pharmaceutically acceptable salt thereof, wherein Y is aryl, wherein the aryl is substituted with C₁-C₃ alkoxy.

[00238] Embodiment I-18. The compound of any one of Embodiments I-1 to I-17, or a pharmaceutically acceptable salt thereof, wherein Y is aryl, wherein the aryl is substituted with A.

[00239] Embodiment I-19. The compound of any one of Embodiments I-1 to I-18,

or a pharmaceutically acceptable salt thereof, wherein A is $\frac{X}{\text{---}}\text{---L}^2\text{---}\overset{\text{O}}{\parallel}\text{C}\text{---OH}$.


[00240] Embodiment I-20. The compound of any one of Embodiments I-1 to I-18,

or a pharmaceutically acceptable salt thereof, wherein A is $\frac{X}{\text{---}}\text{---L}^2\text{---}\overset{\text{O}}{\parallel}\text{P}(\text{OH})_2$.

[00241] Embodiment I-21. The compound of any one of Embodiments I-1 to I-18,


or a pharmaceutically acceptable salt thereof, wherein A is $\frac{X}{\text{---}}\text{---L}^2\text{---}\overset{\text{O}}{\parallel}\text{S}(\text{OH})_2$.

[00242] Embodiment I-22. The compound of any one of Embodiments I-1 to I-18,




or a pharmaceutically acceptable salt thereof, wherein A is

[00243] Embodiment I-23. The compound of any one of Embodiments I-1 to I-18,



or a pharmaceutically acceptable salt thereof, wherein A is

[00244] Embodiment I-24. The compound of any one of Embodiments I-1 to I-18,



or a pharmaceutically acceptable salt thereof, wherein A is

[00245] Embodiment I-25. The compound of any one of Embodiments I-1 to I-24,

or a pharmaceutically acceptable salt thereof, wherein L^2 is $-(CH_2)_n-$.

[00246] Embodiment I-26. The compound of any one of Embodiments I-1 to I-25,

or a pharmaceutically acceptable salt thereof, wherein n is one or two.

[00247] Embodiment I-27. The compound of any one of Embodiments I-1 to I-24,



or a pharmaceutically acceptable salt thereof, wherein L^2 is

[00248] Embodiment I-28. The compound of any one of Embodiments I-1 to I-24,




or a pharmaceutically acceptable salt thereof, wherein L^2 is 

[00249] Embodiment I-29. The compound of any one of Embodiments I-1 to I-24,

or a pharmaceutically acceptable salt thereof, wherein L^2 is $-C(O)NH(CH_2)_n-$







[00250] Embodiment I-30. A compound of Embodiment I-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1a).



wherein

X is H or CH₃


L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , and -CH₂CH₂-;

A is selected from the group consisting of , , , , , and ;

L² is selected from the group consisting of a bond, -CH₂-, -CH₂CH₂-, , , -C(O)NH(CH₂)_n-, -[O(CH₂CH₂)]_n-, -[O(C₁-C₄ alkylene)]-, and -[O(CH₂CH₂)]_n-OCH₂CH₂CF₂-; and







R³ is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy.

[00251] Embodiment I-31. A compound of Embodiment I-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1a),









wherein

X is -CH₂-A^{1a}, -CH₂CH₂-A^{1a}, -CH₂CH₂CH₂-A^{1a}, or -CH₂C(CH₃)₂-A^{1a};

A^{1a} is selected from the group consisting of , , , , , and ;


L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, , , and $-\text{CH}_2\text{CH}_2-$;

A is selected from the group consisting of , , , , , and  ;

L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1\text{-C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n\text{OCH}_2\text{CH}_2\text{CF}_2-$; and

R^3 is H, $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy.







[00252] Embodiment I-32. A compound of Embodiment I-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1b),



wherein

X is H or CH_3 ;


L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, , , and $-\text{CH}_2\text{CH}_2-$;

A is selected from the group consisting of , , , , , and  ;

L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1-\text{C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and





R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.


[00253] Embodiment I-33. A compound of Embodiment I-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1b),






wherein


X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$;

A^{1a} is selected from the group consisting of , , , ,



L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, , , and $-\text{CH}_2\text{CH}_2-$;


A is selected from the group consisting of , , ,



L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1-\text{C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and





R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.


[00254] Embodiment I-34. A compound of Embodiment I-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1c),



wherein

X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$;

A^{1a} is selected from the group consisting of , , , ,




L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, , , and $-\text{CH}_2\text{CH}_2-$;



R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy; and


R^4 is H or C_1-C_3 alkoxy.

[00255] Embodiment I-35. A compound of Embodiment I-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1d),



X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$,


A^{1a} is selected from the group consisting of , , , ,



L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, , , $-\text{O}-$, and $-\text{CH}_2\text{CH}_2-$; and

Y is H or C_1-C_3 alkyl.


[00256] Embodiment I-36. A compound of Embodiment I-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1e),




wherein

R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.

[00257] Embodiment I-37. A compound of Embodiment I-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1f),




[00258] Embodiment I-38. A compound of Embodiment I-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1g),



wherein

R^3 is H, C_1 - C_3 alkyl, or C_1 - C_3 alkoxy.


[00259] Embodiment I-39. A compound of Embodiment I-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1h),



wherein

R^4 is H or C_1 - C_3 alkoxy.

[00260] Embodiment I-40. A compound of Embodiment I-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1i),










wherein
















L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, , , $-\text{O}-$, and $-\text{CH}_2\text{CH}_2-$; and







Y is H or C_1 - C_3 alkyl.

[00261] Embodiment I-41. A compound, or a pharmaceutically acceptable salt, selected from the group consisting of:



Compound	-X	$-\text{L}^1-\text{Y}$	R^{1a} 
1	$-\text{CH}_3$		$-(\text{CH}_2)_3\text{CH}_3$
2	$-\text{CH}_3$		$-(\text{CH}_2)_3\text{CH}_3$
3	$-\text{CH}_3$		$-(\text{CH}_2)_3\text{CH}_3$
4	$-\text{CH}_3$		$-(\text{CH}_2)_3\text{CH}_3$
5	$-\text{CH}_3$		$-(\text{CH}_2)_3\text{CH}_3$
6	$-\text{CH}_3$		$-(\text{CH}_2)_3\text{CH}_3$

Compound	-X	-L ¹ -Y	R ^{1a} 
7	-CH ₃		-(CH ₂) ₃ CH ₃
8	-CH ₃		
9	-CH ₃		-(CH ₂) ₃ CH ₃
10	-CH ₃		-(CH ₂) ₃ CH ₃
11	-CH ₃		-(CH ₂) ₃ CH ₃
12	-CH ₃		-(CH ₂) ₃ CH ₃
13	-CH ₃		-(CH ₂) ₃ CH ₃
14	-CH ₃		-(CH ₂) ₃ CH ₃
15			-(CH ₂) ₃ CH ₃
16	-CH ₃		-(CH ₂) ₃ CH ₃
17	-CH ₃		-(CH ₂) ₃ CH ₃
18	-CH ₃		

Compound	-X	-L ¹ -Y	$\begin{array}{c} R^{1a} \\ \\ \text{---} R^{1b} \end{array}$
19			$-(CH_2)_3CH_3$
20	-CH ₃		
21		-OCH ₃	

[00262] Embodiment I-42. A pharmaceutical composition comprising a compound of any one of Embodiments I-1 to I-41, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

[00263] Embodiment I-43. A method of treating a condition associated with TLR7 modulation in a subject in need thereof, comprising administering to the subject an effective amount of a compound of any one of Embodiments I-1 to I-41, or a pharmaceutically acceptable salt thereof.

[00264] Embodiment I-44. The method of Embodiment I-43, wherein the condition is viral infection or cancer.


[00265] Embodiment I-45. A compound of any one of Embodiments I-1 to I-41, or a pharmaceutically acceptable salt thereof, for use in treating a condition associated with TLR7 modulation.

[00266] Embodiment I-46. The compound of Embodiment I-45, wherein the condition is viral infection or cancer.

[00267] Embodiment I-47. Use of a compound of any one of Embodiments I-1 to I-41, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating a condition associated with TLR7 modulation.


[00268] Embodiment I-48. The use of Embodiment I-47, wherein the condition is viral infection or cancer.

[00269] Embodiment II-1. A compound having the structure of Formula (1), or a pharmaceutically acceptable salt thereof,




wherein

R^{1a} is selected from the group consisting of H, C₁-C₄ alkyl, -NH₂, -NHAc,

-COOH, -SO₂CH₃, -SCH₃, -OCH₃, and ,

wherein the alkyl is optionally substituted with -OH, -NH₂, -NHAc, -COOH, -SO₂CH₃,








-SCH₃, -OCH₃, or ;


R^{1b} is C₂-C₅ alkyl;

X is selected from the group consisting of H and C₁-C₄ alkyl, wherein the alkyl is optionally substituted with A, -OH, or -C(CH₃)₂OH;

L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , -O-, -S-, -SO₂-, -NH-, and -CH₂CH₂-;

Y is selected from the group consisting of C₁-C₃ alkyl, aryl, and heteroaryl, wherein the alkyl, aryl, and heteroaryl are optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy;

A is selected from the group consisting of , , , , , , and ;

L² is selected from the group consisting of a bond, -(CH₂)_n-, -C(O)NH(CH₂)_n-, , , , -[O(CH₂CH₂)]_n-, -[O(C₁-C₄ alkylene)]-,

$-\text{[O(CH}_2\text{CH}_2\text{)]}_n\text{-OCH}_2\text{CH}_2\text{CF}_2\text{-}$; $-\text{C(O)NHCH}_2\text{CH}_2\text{-[O(CH}_2\text{CH}_2\text{)]}_m\text{-}$; and
 $-\text{C(O)NHCH}_2\text{CH}_2\text{-[O(CH}_2\text{CH}_2\text{)]}_m\text{-OCH}_2\text{CH}_2\text{CF}_2\text{-}$;

m is an integer from zero to four; and

n is an integer from one to four; and

wherein the compound is substituted with at least one A; and

when X is $-\text{CH}_3$; L^1 is $-\text{CH}_2\text{-}$; Y is aryl substituted with A; and L^2 is $-\text{CH}_2\text{-}$; then A is not $-\text{L}^2\text{-COOH}$, except when R^{1a} comprises $-\text{COOH}$.




[00270] Embodiment II-2. The compound of Embodiment II-1, or a pharmaceutically acceptable salt thereof, wherein R^{1b} is $-(\text{CH}_2)_2\text{CH}_3$.





[00271] Embodiment II-3. The compound of Embodiment II-1, or a pharmaceutically acceptable salt thereof, wherein R^{1b} is $-(\text{CH}_2)_3\text{CH}_3$.

[00272] Embodiment II-4. The compound of any one of Embodiments II-1 to II-3, or a pharmaceutically acceptable salt thereof, wherein R^{1a} is $\text{C}_1\text{-C}_4$ alkyl, optionally substituted with $-\text{OH}$, $-\text{OCH}_3$, $-\text{SCH}_3$, $-\text{SO}_2\text{CH}_3$

[00273] Embodiment II-5. The compound of any one of Embodiments II-1 to II-4, or a pharmaceutically acceptable salt thereof, wherein R^{1a} is $-\text{CH}_2\text{C}(\text{CH}_3)_2\text{OH}$.



[00274] Embodiment II-6. The compound of any one of Embodiments II-1 to II-4,

or a pharmaceutically acceptable salt thereof, wherein  is , ,

, , , or 

[00275] Embodiment II-7. The compound of any one of Embodiments II-1 to II-3, or a pharmaceutically acceptable salt thereof, wherein R^{1a} is $\text{C}_1\text{-C}_4$ alkyl, optionally substituted with $-\text{COOH}$.

[00276] Embodiment II-8. The compound of any one of Embodiments II-1 to II 3

and II-7, or a pharmaceutically acceptable salt thereof, wherein  is .

[00277] Embodiment II-9. The compound of any one of Embodiments II-1 to II-3, or a pharmaceutically acceptable salt thereof, wherein R^{1a} is H.

[00278] Embodiment II-10. The compound of any one of Embodiments II-1 to II-9, or a pharmaceutically acceptable salt thereof, wherein X is C₁-C₄ alkyl, wherein the alkyl is substituted with A.

[00279] Embodiment II-11. The compound of any one of Embodiments II-1 to II-10, or a pharmaceutically acceptable salt thereof, wherein X is C₁-C₄ alkyl, wherein the alkyl

is substituted with $\text{--}\overset{\text{O}}{\parallel}\text{C}\text{--}\text{OH}$, wherein L² is a bond.

[00280] Embodiment II-12. The compound of any one of Embodiments II-1 to II-9, or a pharmaceutically acceptable salt thereof, wherein X is CH₃.

[00281] Embodiment II-13. The compound of any one of Embodiments II-1 to II-9, or a pharmaceutically acceptable salt thereof, wherein X is H.

[00282] Embodiment II-14. The compound of any one of Embodiments II-1 to II-13, or a pharmaceutically acceptable salt thereof, wherein L¹ is -CH₂-, -CH₂CH₂-, -O-, or -S-.

[00283] Embodiment II-15. The compound of any one of Embodiments II-1 to II-14, or a pharmaceutically acceptable salt thereof, wherein L¹ is -CH₂-.

[00284] Embodiment II-16. The compound of any one of Embodiments II-1 to II-15, or a pharmaceutically acceptable salt thereof, wherein Y is C₁-C₃ alkyl or aryl.

[00285] Embodiment II-17. The compound of any one of Embodiments II-1 to II-16, or a pharmaceutically acceptable salt thereof, wherein Y is aryl, wherein the aryl is substituted with C₁-C₃ alkoxy.

[00286] Embodiment II-18. The compound of any one of Embodiments II-1 to II-17, or a pharmaceutically acceptable salt thereof, wherein Y is aryl, wherein the aryl is substituted with A.

[00287] Embodiment II-19. The compound of any one of Embodiments II-1 to II-18, or a pharmaceutically acceptable salt thereof, wherein A is $\text{--}\overset{\text{O}}{\parallel}\text{C}\text{--}\text{OH}$.


[00288] Embodiment II-20. The compound of any one of Embodiments II-1 to II-

18, or a pharmaceutically acceptable salt thereof, wherein A is $\text{--}\text{P}(\text{OH})_2\text{--}$.


[00289] Embodiment II-21. The compound of any one of Embodiments II-1 to II-

18, or a pharmaceutically acceptable salt thereof, wherein A is $\text{--}\text{S}(=\text{O})_2\text{--}\text{OH}$.


[00290] Embodiment II-22. The compound of any one of Embodiments II-1 to II-

18, or a pharmaceutically acceptable salt thereof, wherein A is .


[00291] Embodiment II-23. The compound of any one of Embodiments II-1 to II-

18, or a pharmaceutically acceptable salt thereof, wherein A is .

[00292] Embodiment II-24. The compound of any one of Embodiments II-1 to II-

18, or a pharmaceutically acceptable salt thereof, wherein A is .

[00293] Embodiment II-25. The compound of any one of Embodiments II-1 to II-

18, or a pharmaceutically acceptable salt thereof, wherein A is .

[00294] Embodiment II-26. The compound of any one of Embodiments II-1 to II-25, or a pharmaceutically acceptable salt thereof, wherein L^2 is $-(CH_2)_n-$.

[00295] Embodiment II-27. The compound of any one of Embodiments II-1 to II-26, or a pharmaceutically acceptable salt thereof, wherein n is one or two.

[00296] Embodiment II-28. The compound of any one of Embodiments II-1 to II-


25, or a pharmaceutically acceptable salt thereof, wherein L^2 is .

[00297] Embodiment II-29. The compound of any one of Embodiments II-1 to II-

25, or a pharmaceutically acceptable salt thereof, wherein L^2 is .

[00298] Embodiment II-30. The compound of any one of Embodiments II-1 to II-25, or a pharmaceutically acceptable salt thereof, wherein L^2 is $-C(O)NH(CH_2)_n-$.









[00299] Embodiment II-31. A compound of Embodiment II-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1a),



wherein

X is H or CH₃


L¹ is selected from the group consisting of a bond, -CH₂-, -O-, -S-, -CF₂-, , , and -CH₂CH₂-;

A is selected from the group consisting of , , , , , , , and ;

L² is selected from the group consisting of a bond, -CH₂-, -CH₂CH₂-, , , -C(O)NH(CH₂)_n-, -[O(CH₂CH₂)]_n-, -[O(C₁-C₄ alkylene)]-, and -[O(CH₂CH₂)]_n-OCH₂CH₂CF₂-; -C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-, C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-OCH₂CH₂CF₂-; and





R³ is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy.

[00300] Embodiment II-32. A compound of Embodiment II-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1a),




wherein

X is -CH₂-A^{1a}, -CH₂CH₂-A^{1a}, -CH₂CH₂CH₂-A^{1a}, or -CH₂C(CH₃)₂-A^{1a};



R³ is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy.







[00301] Embodiment II-33. A compound of Embodiment II-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1b),



wherein

X is H or CH₃;


L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, , , and $-\text{CH}_2\text{CH}_2-$;

A is selected from the group consisting of , , , , , and , and ;

L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1-\text{C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$; $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-$; $\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and







R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.

[00302] Embodiment II-34. A compound of Embodiment II-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1b),








wherein

X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$;

A^{1a} is selected from the group consisting of , , , , , , and ;


L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, , , and $-\text{CH}_2\text{CH}_2-$;

A is selected from the group consisting of , , , , , , and ,

L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1\text{-C}_4\text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_m-$, $\text{OCH}_2\text{CH}_2\text{CF}_2-$, $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-$, $\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-$, $\text{OCH}_2\text{CH}_2\text{CF}_2-$; and






R^3 is H, $\text{C}_1\text{-C}_3$ alkyl, or $\text{C}_1\text{-C}_3$ alkoxy.

[00303] Embodiment II-35. A compound of Embodiment II-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1c),



wherein

X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$;


A^{1a} is selected from the group consisting of , , , , , , and ,

L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, , , and $-\text{CH}_2\text{CH}_2-$;

R^3 is H, C_1 - C_3 alkyl, or C_1 - C_3 alkoxy; and


R^4 is H or C_1 - C_3 alkoxy.

[00304] Embodiment II-36. A compound of Embodiment II-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1d),



X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$;


A^{1a} is selected from the group consisting of , , , ,



L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, , , $-\text{O}-$, and $-\text{CH}_2\text{CH}_2-$, $-\text{S}-$; and


Y is H or C_1 - C_3 alkyl.

[00305] Embodiment II-37. A compound of Embodiment II-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1e),




wherein R^3 is H, C_1 - C_3 alkyl, or C_1 - C_3 alkoxy.

[00306] Embodiment II-38. A compound of Embodiment II-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1f),




[00307] Embodiment II-39. A compound of Embodiment II-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1g),




wherein R^3 is H, C_1 - C_3 alkyl, or C_1 - C_3 alkoxy.

[00308] Embodiment II-40. A compound of Embodiment II-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1h),



wherein R^4 is H or C_1 - C_3 alkoxy.

[00309] Embodiment II-41. A compound of Embodiment II-1, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1i),










wherein
















L^1 is selected from the group consisting of a bond, $-CH_2-$, $-CF_2-$, , , $-O-$, $-CH_2CH_2-$, and $-S-$; and

















Y is H or C_1-C_3 alkyl.


















[00310] Embodiment II-42. A compound, or a pharmaceutically acceptable salt, selected from the group consisting of:

































Compound	-X	$-L^1-Y$	R^{1a} 
1	$-CH_3$		$-(CH_2)_3CH_3$
2	$-CH_3$		$-(CH_2)_3CH_3$
3	$-CH_3$		$-(CH_2)_3CH_3$
4	$-CH_3$		$-(CH_2)_3CH_3$
5	$-CH_3$		$-(CH_2)_3CH_3$
6	$-CH_3$		$-(CH_2)_3CH_3$

Compound	-X	-L ¹ -Y	$ \begin{array}{c} R^{1a} \\ \\ \text{---} \text{C} \text{---} \\ \\ R^{1b} \end{array} $
7	-CH ₃		-(CH ₂) ₃ CH ₃
8	-CH ₃		
9	-CH ₃		-(CH ₂) ₃ CH ₃
10	-CH ₃		-(CH ₂) ₃ CH ₃
11	-CH ₃		-(CH ₂) ₃ CH ₃
12	-CH ₃		-(CH ₂) ₃ CH ₃
13	-CH ₃		-(CH ₂) ₃ CH ₃
14	-CH ₃		-(CH ₂) ₃ CH ₃
15			-(CH ₂) ₃ CH ₃
16	-CH ₃		-(CH ₂) ₃ CH ₃
17	-CH ₃		-(CH ₂) ₃ CH ₃
18	-CH ₃		

Compound	-X	-L ¹ -Y	$\begin{matrix} R^{1a} \\ \\ \text{---} \text{C} \text{---} \\ \\ R^{1b} \end{matrix}$
19			$-(CH_2)_3CH_3$
20	$-CH_3$		
21		$-OCH_3$	
22	$-CH_3$		
23	$-CH_3$		
24	$-CH_3$		$-(CH_2)_3CH_3$
25	$-CH_3$		$-(CH_2)_3CH_3$
26	$-CH_3$		$-(CH_2)_4CH_3$
27			

Compound	-X	-L ¹ -Y	$\begin{matrix} R^{1a} \\ \\ \text{---} R^{1b} \end{matrix}$
28	-CH ₃		-(CH ₂) ₃ CH ₃
29			
30	-CH ₃		-(CH ₂) ₃ CH ₃
31	-CH ₃		-(CH ₂) ₃ CH ₃
32	-CH ₃		
33	-CH ₃		
34	-CH ₃		
35	-CH ₃		
36	-CH ₃		
37			-(CH ₂) ₃ CH ₃

Compound	-X	-L ¹ -Y	$\begin{matrix} R^{1a} \\ \\ \text{---} R^{1b} \end{matrix}$
38			
39	-CH ₃		
40			-(CH ₂) ₃ CH ₃
41	-CH ₃		
42	-CH ₃		
43			-(CH ₂) ₃ CH ₃
44	-CH ₃		
45			
46			
47			-(CH ₂) ₃ CH ₃

Compound	-X	-L ¹ -Y	R ^{1a} R ^{1b}
48			
49			
50			
51	-CH ₃		

[00311] Embodiment II-43. A pharmaceutical composition comprising a compound of any one of Embodiments II-1 to II-42, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

[00312] Embodiment II-44. A method of treating a condition associated with TLR7 modulation in a subject in need thereof, comprising administering to the subject an effective amount of a compound of any one of Embodiments II-1 to II-42, or a pharmaceutically acceptable salt thereof.

[00313] Embodiment II-45. The method of Embodiment II-44, wherein the condition is viral infection or cancer.

[00314] Embodiment II-46. The method of Embodiment II-45, wherein the administration is oral, intravenous, subcutaneous, intramuscular, intratumoral, intradermal, intranasal, inhaled, intravesicle, topical, sublingual, buccal, intrarectal, intrathecal, intracranial, or other forms of local delivery.

[00315] Embodiment II-47. A compound of any one of Embodiments II-1 to II-42, or a pharmaceutically acceptable salt thereof, for use as a medicament.

[00316] Embodiment II-48. A compound of any one of Embodiments II-1 to II-42, or a pharmaceutically acceptable salt thereof, for use in treating a condition associated with TLR7 modulation.

[00317] Embodiment II-49. The compound of Embodiments II-48, wherein the condition is viral infection or cancer.

[00318] Embodiment II-50. Use of a compound of any one of Embodiments II-1 to II-42, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating a condition associated with TLR7 modulation.

[00319] Embodiment II-51. The use of Embodiment II-50, wherein the condition is viral infection or cancer.

[00320] Embodiment II-52. A pharmaceutical composition of Embodiment II-43, further comprising at least one or more additional therapeutic agents.


[00321] Embodiment II-53. The pharmaceutical composition of Embodiment II-52, wherein the at least one or more additional therapeutic agent is antiviral nucleoside.

[00322] Embodiment II-54. The pharmaceutical composition of Embodiment II-52, wherein the at least one or more additional therapeutic agent is PD-1 antibody or PD-L1 antibody.

[00323] Embodiment II-55. A method of treating HBV in a subject in need thereof, comprising administering a compound of any one of Embodiments II-1 to II-42, or a pharmaceutically acceptable salt thereof, in combination with an antiviral nucleoside.


[00324] Embodiment II-56. A method of treating cancer in a subject in need thereof, comprising administering a compound of any one of Embodiments II-1 to II-42, or a pharmaceutically acceptable salt thereof, in combination with a PD-1 antibody or PD-L1 antibody.

[00325] Embodiment III-1. A compound having the structure of Formula (1), or a pharmaceutically acceptable salt thereof,




wherein

R^{1a} is selected from the group consisting of H, C₁-C₄ alkyl, -NH₂, -NHAc,

-COOH, -SO₂CH₃, -SCH₃, -OCH₃, ; and A,

wherein the alkyl is optionally substituted with -OH, -NH₂, -NHAc, -COOH, -SO₂CH₃,




-SCH₃, -OCH₃, , or A;





R^{1b} is C₂-C₅ alkyl;

X is selected from the group consisting of H and C₁-C₄ alkyl, wherein the alkyl is optionally substituted with A, -OH, or -C(CH₃)₂OH;


L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , -O-, -S-, -SO₂-, -NH-, and -CH₂CH₂-;

Y is selected from the group consisting of C₁-C₃ alkyl, aryl, and heteroaryl, wherein the alkyl, aryl, and heteroaryl are optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy;

A is selected from the group consisting of , , ,

, , , and ,

L² is selected from the group consisting of a bond, -(CH₂)_n-, -C(O)NH(CH₂)_n-, ,

, , -[O(CH₂CH₂)]_n-, -[O(C₁-C₄ alkylene)]-, -[O(CH₂CH₂)]_n-OCH₂CH₂CF₂-, -C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-, and -C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-OCH₂CH₂CF₂-;


m is an integer from zero to four;

n is an integer from one to four;


wherein the compound is substituted with at least one A; and

when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and L² is -CH₂-; then A is not -L²-COOH, except when R^{1a} comprises -COOH or -SO₂CH₃; and

when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; L² is -CH₂-, -O-(CH₂)₂-


O(CH₂)₂-, or -O-(CH₂)₂-O(CH₂)₂(CF₂)₂-; and A is ; then A and L¹ are not in a para position with respect to each other.

[00326] Embodiment III-2. A compound having the structure of Formula (1), or a pharmaceutically acceptable salt thereof,




wherein

R^{1a} is selected from the group consisting of H, C₁-C₄ alkyl, -NH₂, -NHAc,

-COOH, -SO₂CH₃, -SCH₃, -OCH₃, and ,

wherein the alkyl is optionally substituted with -OH, -NH₂, -NHAc, -COOH, -SO₂CH₃,

-SCH₃, -OCH₃, or ;


R^{1b} is C₂-C₅ alkyl;


X is selected from the group consisting of H and C₁-C₄ alkyl, wherein the alkyl is optionally substituted with A, -OH, or -C(CH₃)₂OH;

L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , -O-, -S-, -SO₂-, -NH-, and -CH₂CH₂-;

Y is selected from the group consisting of C₁-C₃ alkyl, aryl, and heteroaryl, wherein the alkyl, aryl, and heteroaryl are optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy;


[00331] Embodiment III-7. The compound of any one of Embodiments III-1 to III-

6, or a pharmaceutically acceptable salt thereof, wherein $\begin{array}{c} R^{1a} \\ | \\ \text{---} \text{C} \text{---} R^{1b} \end{array}$ is ,



[00332] Embodiment III-8. The compound of any one of Embodiments III-1 to III-4, or a pharmaceutically acceptable salt thereof, wherein R^{1a} is C_1 - C_4 alkyl, optionally substituted with $-COOH$.


[00333] Embodiment III-9. The compound of any one of Embodiments III-1 to III-4

and III-8, or a pharmaceutically acceptable salt thereof, wherein $\begin{array}{c} R^{1a} \\ | \\ \text{---} \text{C} \text{---} R^{1b} \end{array}$ is .

[00334] Embodiment III-10. The compound of any one of Embodiments III-1 to III-4, or a pharmaceutically acceptable salt thereof, wherein R^{1a} is H.

[00335] Embodiment III-11. The compound of any one of Embodiments III-1 to III-10, or a pharmaceutically acceptable salt thereof, wherein X is C_1 - C_4 alkyl, wherein the alkyl is substituted with A.

[00336] Embodiment III-12. The compound of any one of Embodiments III-1 to III-11, or a pharmaceutically acceptable salt thereof, wherein X is C_1 - C_4 alkyl, wherein the alkyl

is substituted with , wherein L^2 is a bond.

[00337] Embodiment III-13. The compound of any one of Embodiments III-1 to III-10, or a pharmaceutically acceptable salt thereof, wherein X is CH_3 .

[00338] Embodiment III-14. The compound of any one of Embodiments III-1 to III-10, or a pharmaceutically acceptable salt thereof, wherein X is H.

[00339] Embodiment III-15. The compound of any one of Embodiments III-1 to III-14, or a pharmaceutically acceptable salt thereof, wherein L^1 is $-CH_2-$, $-CH_2CH_2-$, $-O-$, or $-S-$.


[00340] Embodiment III-16. The compound of any one of Embodiments III-1 to III-15, or a pharmaceutically acceptable salt thereof, wherein L^1 is $-CH_2-$.

[00341] Embodiment III-17. The compound of any one of Embodiments III-1 to III-16, or a pharmaceutically acceptable salt thereof, wherein Y is C₁-C₃ alkyl or aryl.


[00342] Embodiment III-18. The compound of any one of Embodiments III-1 to III-17, or a pharmaceutically acceptable salt thereof, wherein Y is aryl, wherein the aryl is substituted with C₁-C₃ alkoxy.

[00343] Embodiment III-19. The compound of any one of Embodiments III-1 to III-18, or a pharmaceutically acceptable salt thereof, wherein Y is aryl, wherein the aryl is substituted with A.


[00344] Embodiment III-20. The compound of any one of Embodiments III-1 to III-

19, or a pharmaceutically acceptable salt thereof, wherein A is .


[00345] Embodiment III-21. The compound of any one of Embodiments III-1 to III-

19, or a pharmaceutically acceptable salt thereof, wherein A is .


[00346] Embodiment III-22. The compound of any one of Embodiments III-1 to III-

19, or a pharmaceutically acceptable salt thereof, wherein A is .


[00347] Embodiment III-23. The compound of any one of Embodiments III-1 to III-

19, or a pharmaceutically acceptable salt thereof, wherein A is .


[00348] Embodiment III-24. The compound of any one of Embodiments III-1 to III-

19, or a pharmaceutically acceptable salt thereof, wherein A is .

[00349] Embodiment III-25. The compound of any one of Embodiments III-1 to III-

19, or a pharmaceutically acceptable salt thereof, wherein A is .

[00350] Embodiment III-26. The compound of any one of Embodiments III-1 to III-

19, or a pharmaceutically acceptable salt thereof, wherein A is .

[00351] Embodiment III-27. The compound of any one of Embodiments III-1 to III-26, or a pharmaceutically acceptable salt thereof, wherein L^2 is $-(CH_2)_n$.


[00352] Embodiment III-28. The compound of any one of Embodiments III-1 to III-27, or a pharmaceutically acceptable salt thereof, wherein n is one or two.

[00353] Embodiment III-29. The compound of any one of Embodiments III-1 to III-26, or a pharmaceutically acceptable salt thereof, wherein L^2 is .

[00354] Embodiment III-30. The compound of any one of Embodiments III-1 to III-26, or a pharmaceutically acceptable salt thereof, wherein L^2 is .

[00355] Embodiment III-31. The compound of any one of Embodiments III-1 to III-26, or a pharmaceutically acceptable salt thereof, wherein L^2 is $-C(O)NH(CH_2)_n$.









[00356] Embodiment III-32. A compound of Embodiment III-1 or III-2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1a),



wherein

X is H or CH_3


L^1 is selected from the group consisting of a bond, $-CH_2-$, $-O-$, $-S-$, $-CF_2-$, , , and $-CH_2CH_2-$;

A is selected from the group consisting of , , , , , , , and .

L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1-\text{C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$; $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-$; $\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and


R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.


[00357] Embodiment III-33. A compound of Embodiment III-1 or III-2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1a),



wherein


X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$;

A^{1a} is selected from the group consisting of , , , ,



L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, , , and $-\text{CH}_2\text{CH}_2-$;


A is selected from the group consisting of , , ,



L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1-\text{C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$; $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-$; $\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and

R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.

[00358] Embodiment III-34. A compound of Embodiment III-1 or III-2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1b),




wherein

X is H or CH_3 ;

L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, , , and $-\text{CH}_2\text{CH}_2-$;


A is selected from the group consisting of $-\text{L}^2-\text{C}(=\text{O})\text{OH}$, $-\text{L}^2-\text{P}(\text{OH})_2$, $-\text{L}^2-\text{SO}_3\text{H}$,



L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1-\text{C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$; $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-$; $\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and

R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.


[00359] Embodiment III-35. A compound of Embodiment III-1 or III-2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1b),



wherein


X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$;

A^{1a} is selected from the group consisting of ,



L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, , , and $-\text{CH}_2\text{CH}_2-$;


A is selected from the group consisting of ,



L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1-\text{C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$; $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-$; $\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and

R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.


[00360] Embodiment III-36. A compound of Embodiment III-1 or III-2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1c),




wherein

X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$;

A^{1a} is selected from the group consisting of ,




L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, ,



R^3 is H, C_1 - C_3 alkyl, or C_1 - C_3 alkoxy; and


R^4 is H or C_1 - C_3 alkoxy.

[00361] Embodiment III-37. A compound of Embodiment III-1 or III-2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1d),



X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$;


A^{1a} is selected from the group consisting of , , , ,



L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , -O-, and -CH₂CH₂-, -S-, and


Y is H or C₁-C₃ alkyl.

[00362] Embodiment III-38. A compound of Embodiment III-1 or III-2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1e),




wherein R³ is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy.

[00363] Embodiment III-39. A compound of Embodiment III-1 or III-22, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1f),




[00364] Embodiment III-40. A compound of Embodiment III-1 or III-22, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1g),




wherein R^3 is H, C_1 - C_3 alkyl, or C_1 - C_3 alkoxy.

[00365] Embodiment III-41. A compound of Embodiment III-1 or III-2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1h),



wherein R^4 is H or C_1 - C_3 alkoxy.


[00366] Embodiment III-42. A compound of Embodiment III-1 or III-2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1i),




wherein

L^1 is selected from the group consisting of a bond, $-CH_2-$, $-CF_2-$, , , $-O-$, $-CH_2CH_2-$, and $-S-$; and
Y is H or C_1 - C_3 alkyl.


[00367] Embodiment III-43. A compound of Embodiment III-1 or III-2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1j),



[00368] Embodiment III-44. A compound of Embodiment III-1 or III-2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1k),














































[00369] Embodiment III-45. A compound, or a pharmaceutically acceptable salt, selected from the group consisting of:


























Compound	-X	-L ¹ -Y	$\begin{matrix} R^{1a} \\ \\ \text{---} \text{C} \text{---} \\ \\ R^{1b} \end{matrix}$	Compound
1	-CH ₃		-(CH ₂) ₃ CH ₃	
2	-CH ₃		-(CH ₂) ₃ CH ₃	
3	-CH ₃		-(CH ₂) ₃ CH ₃	



























Compound	-X	-L ¹ -Y	$\begin{array}{c} R^{1a} \\ \\ \text{---} R^{1b} \end{array}$	Compound
4	-CH ₃		-(CH ₂) ₃ CH ₃	
5	-CH ₃		-(CH ₂) ₃ CH ₃	
6	-CH ₃		-(CH ₂) ₃ CH ₃	
7	-CH ₃		-(CH ₂) ₃ CH ₃	
8	-CH ₃			
9	-CH ₃		-(CH ₂) ₃ CH ₃	
10	-CH ₃		-(CH ₂) ₃ CH ₃	
11	-CH ₃		-(CH ₂) ₃ CH ₃	
12	-CH ₃		-(CH ₂) ₃ CH ₃	

Compound	-X	-L ¹ -Y	R ^{1a} 	Compound
13	-CH ₃		-(CH ₂) ₃ CH ₃	
14	-CH ₃		-(CH ₂) ₃ CH ₃	
15			-(CH ₂) ₃ CH ₃	
16	-CH ₃		-(CH ₂) ₃ CH ₃	
17	-CH ₃		-(CH ₂) ₃ CH ₃	
18	-CH ₃			
19			-(CH ₂) ₃ CH ₃	
20	-CH ₃			

Compound	-X	-L ¹ -Y	R ^{1a} 	Compound
20C	-CH ₃			
21		-OCH ₃		
22	-CH ₃			
23	-CH ₃			
24	-CH ₃		-(CH ₂) ₃ CH ₃	
25	-CH ₃		-(CH ₂) ₃ CH ₃	
26	-CH ₃		-(CH ₂) ₄ CH ₃	
27				

Compound	-X	-L ¹ -Y	R ^{1a} R ^{1b}	Compound
28	-CH ₃		-(CH ₂) ₃ CH ₃	
29				
30	-CH ₃		-(CH ₂) ₃ CH ₃	
31	-CH ₃		-(CH ₂) ₃ CH ₃	
32	-CH ₃			
33	-CH ₃			
34	-CH ₃			
35	-CH ₃			

Compound	-X	-L ¹ -Y	R ^{1a} 	Compound
36	-CH ₃			
37			-(CH ₂) ₃ CH ₃	
38				
39	-CH ₃			
40			-(CH ₂) ₃ CH ₃	
41	-CH ₃			
42	-CH ₃			

Compound	-X	-L ¹ -Y	R ^{1a} 	Compound
43			-(CH ₂) ₃ CH ₃	
44	-CH ₃			
45				
46				
47			-(CH ₂) ₃ CH ₃	
48				
49				

Compound	-X	-L ¹ -Y	R ^{1a} R ^{1b}	Compound
50				
51	-CH ₃			

[00370] Embodiment III-46. A pharmaceutical composition comprising a compound of any one of Embodiments III-1 to III-45, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

[00371] Embodiment III-47. A method of treating a condition associated with TLR7 modulation in a subject in need thereof, comprising administering to the subject an effective amount of a compound of any one of Embodiments III-1 to III-45, or a pharmaceutically acceptable salt thereof.

[00372] Embodiment III-48. The method of Embodiment III-47, wherein the condition is viral infection or cancer.

[00373] Embodiment III-49. The method of Embodiment III-47 or III-48, wherein the administration is oral, intravenous, subcutaneous, intramuscular, intratumoral, intradermal, intranasal, inhaled, intravesicle, topical, sublingual, buccal, intrarectal, intrathecal, intracranial, or other forms of local delivery.

[00374] Embodiment III-50. A compound of any one of Embodiments III-1 to III-45, or a pharmaceutically acceptable salt thereof, for use as a medicament.

[00375] Embodiment III-51. A compound of any one of Embodiments III-1 to III-45, or a pharmaceutically acceptable salt thereof, for use in treating a condition associated with TLR7 modulation.

[00376] Embodiment III-52. The compound of Embodiment III-51, wherein the condition is viral infection or cancer.

[00377] Embodiment III-53. Use of a compound of any one of Embodiments III-1 to III-45, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating a condition associated with TLR7 modulation.

[00378] Embodiment III-54. The use of Embodiment III-53, wherein the condition is viral infection or cancer.

[00379] Embodiment III-55. A pharmaceutical composition of Embodiment III-46, further comprising at least one or more additional therapeutic agents.

[00380] Embodiment III-56. The pharmaceutical composition of Embodiment III-55, wherein the at least one or more additional therapeutic agent is antiviral nucleoside.

[00381] Embodiment III-57. The pharmaceutical composition of Embodiment III-55, wherein the at least one or more additional therapeutic agent is PD-1 antibody or PD-L1 antibody.

[00382] Embodiment III-58. A method of treating HBV in a subject in need thereof, comprising administering a compound of any one of Embodiments III-1 to III-45, or a pharmaceutically acceptable salt thereof, in combination with an antiviral nucleoside.

[00383] Embodiment III-59. A method of treating cancer in a subject in need thereof, comprising administering a compound of any one of Embodiments III-1 to III-45, or a pharmaceutically acceptable salt thereof, in combination with a PD-1 antibody or PD-L1 antibody.

Examples


[00384] The following examples are provided to illustrate the present disclosure, and should not be construed as limiting thereof. In these examples, all parts and percentages are by weight, unless otherwise noted. Abbreviations in the examples are noted below.

Abbreviations

AIBN	azobisisobutyronitrile
aq.	aqueous
DCM	dichloromethane
DMAP	4-dimethylaminopyridine
EA	ethyl acetate
eq	equivalent
h	hour
HATU	(1-[Bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium 3-oxid hexafluorophosphate)
HPLC	high performance liquid chromatography
LC-MS	liquid chromatography mass spectrometry
min	minutes
NBS	N-bromosuccinimide

NMP	N-methylpyrrolidine
NMR	nuclear magnetic resonance
PE	petroleum ether
rt or r.t.	room temperature
sat.	saturated
TBAF	tetrabutylammonium flouride
TBDPS	tert-butyl diphenylsilyl
TEA	triethylamine
THF	tetrahydrofuran
TLC	thin layer chromatography

Example 1: 3-(4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxyphenyl)propanoic acid (Compound 1)



Step 1: methyl 4-(2-(ethoxycarbonyl)-3-oxobutyl)-3-methoxybenzoate

[00385] Sodium hydride (60% in mineral oil, 1.25 eq) was added in portions over 10 min to a solution of ethyl 3-oxobutanoate (1.2 eq) in THF (0.8M) at 0 °C. The resulting suspension was stirred at 0 °C for 10 min and a solution of methyl 4-(bromomethyl)-3-

methoxybenzoate (1.0 eq) in THF (0.5M) was added dropwise over 10 min. The mixture was warmed to 70 °C and stirred for 3 h. The mixture was allowed to cool and then poured into ice water and stirred for 30 min. The mixture was partitioned between EA/water. The organic layer dried over Na₂SO₄, filtered, concentrated and purified by flash chromatography on silica (eluent PE/EA = 100/1 to 5/1) to give the title compound.

Step 2: methyl 4-((2-amino-4-hydroxy-6-methylpyrimidin-5-yl)methyl)-3-methoxybenzoate

[00386] A mixture of methyl 4-(2-(ethoxycarbonyl)-3-oxobutyl)-3-methoxybenzoate (1.0 eq) and guanidine carbonate (1.0 eq) in MeOH (0.2M) was stirred overnight at 65 °C and then allowed to cool to rt. The precipitate was collected by filtration and suspended in water. The solid was collected by filtration and washed with MeOH and EA to give the title compound as a white solid.

Step 3: methyl 4-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-3-methoxybenzoate

[00387] Methyl 4-((2-amino-4-hydroxy-6-methylpyrimidin-5-yl)methyl)-3-methoxybenzoate (1.0 eq) in POCl₃ (0.5M) was stirred overnight at 100 °C under N₂. The mixture was cooled to rt and the solvent was removed. To the residue was added water and the pH of the mixture was adjusted to 7 with solid NaHCO₃. The resulting mixture was heated at 50 °C for 1h and then allowed to cool to rt. The solid was collected, washed with water, EA and dried under vacuum to give the title compound.

Step 4: methyl 4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxybenzoate

[00388] To a suspension of methyl 4-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-3-methoxybenzoate (1.0 eq) in NMP (0.3M) was added butan-1-amine (4.5 eq). The resulting mixture was stirred overnight at 125 °C and the mixture was cooled to rt. The mixture was partitioned between EA/water. The organic layer was dried over Na₂SO₄, concentrated and purified by flash chromatography on silica (eluent PE/EA = 20:1 to 1:5) to give the title compound.

Step 5: (4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxyphenyl)methanol

[00389] To a stirred solution of methyl 4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxybenzoate (1.0 eq) in THF (0.1M) was added LiAlH₄

(1M in THF, 2.0 eq) dropwise. The resulting mixture was stirred at 0 °C for 10 min and at rt for 1 h. The mixture was diluted with EA and quenched with 2N NaOH. The mixture was partitioned between EA/water. The organic layer was dried over Na₂SO₄, filtered and concentrated to give the title compound as white solid.

Step 6: 4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxybenzaldehyde

[00390] To a stirred solution of 4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxyphenyl)methanol (1.0 eq) in THF (0.06M) was added MnO₂ (5.0 eq). The resulting mixture was stirred overnight at 50 °C. The mixture was filtered. The filtrate was concentrated to give the title compound.

Step 7: ethyl (E)-3-(4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxyphenyl)acrylate

[00391] A mixture of 4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxybenzaldehyde (1.0 eq) and ethyl 2-(triphenyl-λ⁵-phosphanylidene)acetate (1.2eq) in THF (0.1M) was stirred at 70 °C for 5 h. The mixture was concentrated and purified by flash chromatography on silica (eluent 0-5% MeOH in DCM) to give the title compound as a white solid.

Step 8: ethyl 3-(4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxyphenyl)propanoate

[00392] To a stirred solution of ethyl (E)-3-(4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxyphenyl)acrylate (1.0 eq) in EA (0.02M) was added Pd/C (20% weight). The resulting mixture was stirred overnight at 50 °C under H₂. The mixture was filtered. The filtrate was concentrated and purified by flash chromatography on silica (eluent 0-5% MeOH in DCM) to give the title compound.


Step 9: 3-(4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxyphenyl)propanoic acid

[00393] To a stirred solution of ethyl 3-(4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxyphenyl)propanoate (1.0 eq) in MeOH (0.03M) was added 1 N NaOH (5.0 eq). The resulting mixture was stirred overnight at 45 °C. Solvent was removed. The residue was neutralized with 2 N HCl to pH 7. The suspension was filtered. The solid was collected and dissolved in HCl/dioxane. The solution was concentrated to give the title compound as a white solid (HCl salt).

[00394] LC-MS: [M+H] = 373.4

[00395] ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.32 (br s, 1H), 12.11 (br s, 1H), 7.83 (br s, 1H), 7.41 (br s, 2H), 6.90 (s, 1H), 6.70 (s, 2H), 3.83 (s, 3H), 3.64 (s, 2H), 3.36-3.30 (m, 2H), 2.82-2.77 (m, 2H), 2.54-2.50 (m, 2H), 2.11 (s, 3H), 1.50-1.43 (m, 2H), 1.23-1.15 (m, 2H), 0.85 (t, *J* = 7.2 Hz, 3H).

Example 2: 5-(4-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-N⁴-butyl-6-methylpyrimidine-2,4-diamine (Compound 2)



Step 1: N⁴-butyl-5-(4-(chloromethyl)-2-methoxybenzyl)-6-methylpyrimidine-2,4-diamine

[00396] To a stirred solution of 4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxyphenyl)methanol (1.0 eq, Example 1 – Step 5) in DCM (0.14M) was added SOCl₂ (1.8 eq) at rt under N₂. The mixture was stirred at rt for 1 h. The mixture was partitioned between DCM/sat. aq. NaHCO₃. The organic layer was dried over Na₂SO₄, filtered and concentrated to give the title compound.

Step 2: 2-(4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxyphenyl)acetonitrile

[00397] A mixture of *N*⁴-butyl-5-(4-(chloromethyl)-2-methoxybenzyl)-6-methylpyrimidine-2,4-diamine (1.0 eq) in 1:1 DMSO/DMF (0.14M) and NaCN (2.8 eq) was stirred overnight at rt. The mixture was partitioned between EA/water. The organic layer was dried over Na₂SO₄, filtered, concentrated and purified by flash chromatography on silica (eluent 0-5% MeOH in DCM) to give the title compound.


Step 3: 5-(4-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-N⁴-butyl-6-methylpyrimidine-2,4-diamine

[00398] To a stirred mixture of 2-(4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxyphenyl)acetonitrile (1.0 eq), Bu₂SnO (2.0 eq) in NMP (0.06M) was added TMSN₃ (10.0 eq). The resulting mixture was stirred overnight at 125 °C under N₂. The mixture was cooled to rt. The mixture was filtered to give the title compound as a white powder.

[00399] LC-MS: [M+H] = 383.5

[00400] ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.19 (s, 1H), 7.85 (br s, 1H), 7.34 (br s, 2H), 7.01 (s, 1H), 6.74 (s, 2H), 4.24 (s, 2H), 3.84 (s, 3H), 3.65 (s, 2H), 3.37-3.34 (m, 2H), 2.09 (s, 3H), 1.48-1.41 (m, 2H), 1.23-1.16 (m, 2H), 0.83 (t, *J* = 7.2 Hz, 3H).

Example 3A: 5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-N⁴-butyl-6-methylpyrimidine-2,4-diamine (Compound 3)



Step A: methyl 3-(bromomethyl)-4-methoxybenzoate

[00401] To a mixture of methyl 4-methoxy-3-methylbenzoate (1.0 eq) in CCl₄ (0.37M), NBS (1.1 eq) and AIBN (0.2 eq) was stirred overnight at 95 °C under nitrogen atmosphere. The resulting mixture was concentrated and partitioned between DCM and water, the organic layer was dried over Na₂SO₄, concentrated and purified by flash chromatography on silica (eluent PE/EA = 100:1 to 20:1) to give the title compound as a white solid.

Preparation of (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)methanol

[00402] Preparation of (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)methanol followed Example 1 – Step 1 to 5, but using methyl 3-(bromomethyl)-4-methoxybenzoate instead of methyl 4-(bromomethyl)-3-methoxybenzoate in Step 1.


Preparation of 5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-N⁴-butyl-6-methylpyrimidine-2,4-diamine

[00403] Preparation of 5-(5-((2*H*-tetrazol-5-yl)methyl)-2-methoxybenzyl)-N⁴-butyl-6-methylpyrimidine-2,4-diamine followed Example 2 – Steps 1 to 3, but using (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)methanol instead of 4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxyphenyl)methanol in Step 1. The title compound was purified by prep-HPLC using acetonitrile/water as the eluent to give a white solid.

[00404] LC-MS: [M+H] = 383.3

[00405] ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.03 (d, *J* = 7.6 Hz, 1H), 6.90 (d, *J* = 8.4 Hz, 1H), 6.73 (s, 1H), 6.60 (br s, 1H), 6.31 (s, 2H), 3.97 (s, 2H), 3.80 (s, 3H), 3.60 (s, 2H), 3.29-3.24 (m, 2H), 2.02 (s, 3H), 1.43-1.37 (m, 2H), 1.22-1.17 (m, 2H), 0.84 (t, *J* = 7.2 Hz, 3H).

Example 3B: 5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-N⁴-butyl-6-methylpyrimidine-2,4-diamine (Compound 3)



Step 1: methyl 3-(bromomethyl)-4-methoxybenzoate

[00406] To a mixture of methyl 4-methoxy-3-methylbenzoate (1.0 eq) in CCl₄ (0.37M), NBS (1.1 eq) and AIBN (0.2 eq) was stirred overnight at 95 °C under nitrogen atmosphere. The resulting mixture was concentrated and partitioned between DCM and water, the organic layer was dried over Na₂SO₄, concentrated and purified by flash chromatography on silica (eluent PE/EA = 100:1 to 20:1) to give the title compound as a white solid.

Step 2: methyl 3-(2-(ethoxycarbonyl)-3-oxobutyl)-4-methoxybenzoate

[00407] To a solution of ethyl 3-oxobutanoate (1.2 eq) in THF (0.35M) at 0 °C was added in portions 60% NaH (1.25 eq) under N₂. The resulting suspension was stirred at 0 °C for 10 min, then a solution of methyl 3-(bromomethyl)-4-methoxybenzoate (1.0 eq) in THF (2M) was added dropwise over 10 min. The resulting mixture was stirred overnight at 70 °C. The mixture was cooled down to rt and ice water was added. The mixture was extracted with ethyl acetate. The organic layer was separated and dried over Na₂SO₄, concentrated and purified by flash chromatography on silica (eluent PE/EA = 50:1-5:1) to give the title compound as a yellow oil.

Step 3: methyl 3-((2-amino-4-hydroxy-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate

[00408] To a solution of methyl 3-(2-(ethoxycarbonyl)-3-oxobutyl)-4-methoxybenzoate (1.0 eq) in MeOH (0.2M) was added guanidine carbonate (1.0 eq). The resulting mixture was stirred at 65 °C for 16 h and then cooled down to r.t. The precipitate was collected by filtration, and the solid was triturated with water and filtered. The filter cake was washed with MeOH and EtOAc, dried under vacuum to give the title compound as a white solid.

Step 4: methyl 3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate

[00409] To a suspension of methyl 3-((2-amino-4-hydroxy-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate (1.0 eq) in POCl₃ (0.55M) was stirred at 100 °C for 15 h. The reaction mixture was allowed to cool to rt, and POCl₃ was evaporated to dryness under reduced pressure. The residue was diluted with water and adjusted pH to 7 with saturated NaHCO₃. Then the mixture was heated at 50 °C for 1 h, cooled down to rt. The precipitate was collected by filtration, and the filter cake was washed with water and EtOAc, dried over vacuum to give the title compound.

Step 5: methyl 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate

[00410] A mixture of butan-1-amine (4.5 eq) and methyl 3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate (1.0 eq) in NMP (0.3M) was stirred overnight at 150 °C. The reaction was allowed to cool, diluted with EtOAc, washed with water and brine. The organic phase was dried and evaporated under reduced pressure. The

residue was suspended in diethyl ether, and the solid was collected by filtration to give the title compound as a solid.

Step 6: (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-methanol

[00411] To a stirred solution of methyl 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate (1.0 eq) in THF (0.3M) was added 2.5M LiAlH₄ (2.0 eq) in THF at 0 °C under N₂ atmosphere. The resulting mixture was stirred at 0 °C for 30 min and then at r.t. for 2 h. The mixture was diluted with EtOAc and quenched with 2N NaOH. The suspension was filtered, and the filtrate was partitioned between EA/water. The organic layer was dried over Na₂SO₄, filtered and concentrated to give the title compound as a white solid.

Step 7: N⁴-butyl-5-(5-(chloromethyl)-2-methoxybenzyl)-6-methylpyrimidine-2,4-diamine

[00412] To a stirred solution of (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)methanol (1.0 eq) in DCM (0.34M) was added SOCl₂ (1.3 eq) at rt under N₂. The mixture was stirred at rt for 1 h. The mixture was partitioned between DCM and saturated aqueous NaHCO₃. The organic layer was dried over Na₂SO₄, filtered and concentrated to give the title compound.

Step 8: 2-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-acetonitrile

[00413] A mixture of N⁴-butyl-5-(5-(chloromethyl)-2-methoxybenzyl)-6-methylpyrimidine-2,4-diamine (1.0 eq) in 1:1 DMSO/DMF (0.34M) and NaCN (2.8 eq) was stirred overnight at rt. The mixture was partitioned between EA/water. The organic layer was dried over Na₂SO₄, filtered, concentrated and purified by flash chromatography on silica (eluent 0-5% MeOH in DCM) to give the title compound.


Step 9: 5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-N⁴-butyl-6-methylpyrimidine-2,4-diamine

[00414] To a stirred mixture of 2-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)acetonitrile (1.0 eq) in NMP (0.06M) and Bu₂SnO (2.0 eq) was added TMSN₃ (10.0 eq). The resulting mixture was stirred overnight at 125 °C under N₂. The mixture was cooled down to rt, and the mixture was concentrated and purified by prep-HPLC to give the title compound as a solid.

[00415] LC-MS: $[M+H]^+ = 383.3$

[00416] ^1H NMR (400 MHz, DMSO- d_6) δ 7.03 (d, $J = 7.6$ Hz, 1H), 6.90 (d, $J = 8.4$ Hz, 1H), 6.73 (s, 1H), 6.60 (br s, 1H), 6.31 (s, 2H), 3.97 (s, 2H), 3.80 (s, 3H), 3.60 (s, 2H), 3.29-3.24 (m, 2H), 2.02 (s, 3H), 1.43-1.37 (m, 2H), 1.22-1.17 (m, 2H), 0.84 (t, $J = 7.2$ Hz, 3H).

Example 4: 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)propanoic acid (Compound 4)




[00417] The title compound was prepared according to the protocols described for Example 1 – Steps 1 to 9, but using methyl 3-(bromomethyl)-4-methoxybenzoate instead of methyl 4-(bromomethyl)-3-methoxybenzoate in Step 1.

[00418] LC-MS: $[M+H] = 373.2$

[00419] ^1H NMR (400 MHz, DMSO- d_6) δ 12.47 (s, 1H), 7.81 (t, $J = 4.8$ Hz, 1H), 7.46 (br s, 2H), 7.06 (d, $J = 8.4$ Hz, 1H), 6.91 (d, $J = 8.4$ Hz, 1H), 6.72 (s, 1H), 3.81 (s, 3H), 3.76 (s, 2H), 3.40-3.36 (m, 2H), 2.68 (t, $J = 7.2$ Hz, 2H), 2.41 (t, $J = 7.2$ Hz, 2H), 2.13 (s, 3H), 1.50-1.44 (m, 2H), 1.24-1.18 (m, 2H), 0.85 (t, $J = 7.2$ Hz, 3H).

Example 5: 2-(5-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzyl)-2H-tetrazol-2-yl)acetic acid (Compound 5)

Example 6: 2-(5-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzyl)-1H-tetrazol-1-yl)acetic acid (Compound 6)



[00420] To a stirred solution of 5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)- N^4 -butyl-6-methylpyrimidine-2,4-diamine (1.0 eq, Example 3) in acetone (0.05M) was added

K₂CO₃ (1.5 eq) at 0 °C under N₂. The resulting mixture was stirred at 0 °C for 30 min and ethyl 2-bromoacetate (1.5 eq) was added dropwise. The mixture was slowly warmed to rt and stirred for 5 h. The mixture was quenched with sat. NH₄Cl. The mixture was partitioned between DCM/water. The organic phase was dried over Na₂SO₄, concentrated and purified by prep-HPLC to give two regioisomers in 3:2 ratio as white powders. To a stirred solution of each isomer (1.0 eq) in MeOH (0.02M) was added 1N NaOH (5.0 eq). The resulting mixture was stirred at rt for 16 h. The reaction was monitored by TLC and LC-MS. The mixture was neutralized with 1N HCl and purified to give the title compounds.

[00421] Isomer 1: 2-(5-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzyl)-2*H*-tetrazol-2-yl)acetic acid

[00422] LC-MS: [M+H] = 441.5


[00423] ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.62 (br s, 1H), 7.82 (br s, 1H), 7.50 (br s, 2H), 7.11 (d, *J* = 7.6 Hz, 1H), 6.95 (d, *J* = 8.8 Hz, 1H), 6.79 (s, 1H), 5.31 (s, 2H), 4.16 (s, 2H), 3.82 (s, 3H), 3.66 (s, 2H), 3.39-3.30 (m, 2H), 2.10 (s, 3H), 1.45-1.38 (m, 2H), 1.3-1.2 (m, 2H), 0.85 (t, *J* = 7.2 Hz, 3H).

[00424] Isomer 2: 2-(5-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzyl)-1*H*-tetrazol-1-yl)acetic acid

[00425] LC-MS: [M+H] = 441.5

[00426] ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.69 (s, 1H), 7.85 (br s, 1H), 7.51 (br s, 2H), 7.10 (d, *J* = 8.0 Hz, 1H), 6.95 (d, *J* = 8.4 Hz, 1H), 6.82 (s, 1H), 5.58 (s, 2H), 4.11 (s, 2H), 3.81 (s, 3H), 3.67 (s, 2H), 3.38-3.35 (m, 2H), 2.11 (s, 3H), 1.49-1.42 (m, 2H), 1.24-1.18 (m, 2H), 0.85 (t, *J* = 7.2 Hz, 3H).

Example 7: (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzyl)phosphonic acid (Compound 7)



Step 1: *N*⁴-butyl-5-(5-(chloromethyl)-2-methoxybenzyl)-6-methylpyrimidine-2,4-diamine

[00427] The title compound was prepared according to the protocols described in Example 2 – Step 1, but using (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)methanol instead of (4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxyphenyl)methanol.

Step 2: diethyl (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzyl)phosphonate

[00428] *N*⁴-butyl-5-(5-(chloromethyl)-2-methoxybenzyl)-6-methylpyrimidine-2,4-diamine (1.0 eq) in neat triethyl phosphite (0.1M) was stirred at 140 °C for 2 h. The mixture was concentrated to give the title compound.

Step 3: (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzyl)phosphonic acid


[00429] To a stirred solution of diethyl (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzyl)phosphonate (1.0 eq) in DCM (0.04M) was added bromotrimethylsilane (4.0 eq). The resulting mixture was stirred for 2 h at rt and then partitioned between DCM/water. The organic layer was dried over Na_2SO_4 , concentrated. The crude product was purified by prep-HPLC (mobile phase $\text{CH}_3\text{CN}/\text{H}_2\text{O}/\text{NH}_3$) and freeze-

dried to give a white powder, which was re-dissolved in MeCN/H₂O/1N HCl (4:2:1) and freeze-dried to give the title compound as a white solid (HCl salt).

[00430] LC-MS: [M+H] = 395.1

[00431] ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.64 (br s, 2H), 7.82 (t, *J* = 5.6 Hz, 1H), 7.48 (br s, 2H), 7.08 (d, *J* = 8.0 Hz, 1H), 6.91 (d, *J* = 8.4 Hz, 1H), 6.74 (s, 1H), 3.81 (s, 3H), 3.66 (s, 2H), 3.37-3.34 (m, 2H), 2.80 (d, *J* = 20.9 Hz, 2H), 2.12 (s, 3H), 1.50-1.45 (m, 2H), 1.27-1.22 (m, 2H), 0.86 (t, *J* = 7.2 Hz, 3H).

Example 8: (S)-2-((5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-methylpyrimidin-4-yl)amino)pentan-1-ol (Compound 8)



Step A: (S)-1-((tert-butyldiphenylsilyl)oxy)pentan-2-amine

[00432] A mixture of (S)-2-aminopentan-1-ol (1.0 eq) in DCM (0.2M), TEA (5.0 eq), and DMAP (0.3 eq) at 0 °C was added TBDPSCl (1.5 eq). The resulting mixture was stirred at rt for 3 h. The mixture was diluted with water and DCM, separated, and the aqueous layer was back-extracted with DCM twice. The combined organic layers were dried over Na₂SO₄, concentrated under reduced pressure. The residue was purified by flash column chromatography (eluent: DCM/MeOH = 200:1 to 80:1) to give the title compound as a brown oil.

Preparation of methyl (S)-3-((2-amino-4-((1-((tert-butyldiphenylsilyl)oxy)pentan-2-yl)amino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate

[00433] Preparation of methyl (S)-3-((2-amino-4-((1-((tert-butyldiphenylsilyl)oxy)pentan-2-yl)amino)-6-methylpyrimidin-5-yl)methyl)-4-

methoxybenzoate followed Example 1 – Step 4, but using methyl 3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate instead of methyl 4-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-3-methoxybenzoate, and (*S*)-1-((*tert*-butyldiphenylsilyl)oxy)pentan-2-amine instead of butan-1-amine.

Preparation of (*S*)-(3-((2-amino-4-((1-((*tert*-butyldiphenylsilyl)oxy)pentan-2-yl)amino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)methanol

[00434] Preparation of (*S*)-(3-((2-amino-4-((1-((*tert*-butyldiphenylsilyl)oxy)pentan-2-yl)amino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)methanol followed Example 1 – Step 5, but using methyl (*S*)-3-((2-amino-4-((1-((*tert*-butyldiphenylsilyl)oxy)pentan-2-yl)amino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate instead of methyl 4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxybenzoate.

Preparation of (*S*)-5-(5-((2*H*-tetrazol-5-yl)methyl)-2-methoxybenzyl)-*N*⁴-(1-((*tert*-butyldiphenylsilyl)oxy)pentan-2-yl)-6-methylpyrimidine-2,4-diamine

[00435] Preparation of (*S*)-5-(5-((2*H*-tetrazol-5-yl)methyl)-2-methoxybenzyl)-*N*⁴-(1-((*tert*-butyldiphenylsilyl)oxy)pentan-2-yl)-6-methylpyrimidine-2,4-diamine followed Example 2 – Steps 1 to 3, but using (*S*)-(3-((2-amino-4-((1-((*tert*-butyldiphenylsilyl)oxy)pentan-2-yl)amino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)methanol instead of (4-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-3-methoxyphenyl)methanol.

Preparation of (*S*)-2-((5-(5-((2*H*-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-methylpyrimidin-4-yl)amino)pentan-1-ol


[00436] To a solution of (*S*)-5-(5-((2*H*-tetrazol-5-yl)methyl)-2-methoxybenzyl)-*N*⁴-(1-((*tert*-butyldiphenylsilyl)oxy)pentan-2-yl)-6-methylpyrimidine-2,4-diamine (1.0 eq) in anhydrous THF (0.05M) at 0 °C was added TBAF (10 eq). The resulting mixture was stirred at rt for 16 h. LCMS showed the reaction was completed. Then the mixture was concentrated under reduced pressure. The residue was purified by prep-HPLC (mobile phase CH₃CN/H₂O/HCOOH), freeze-dried to give the title compound as a white solid (formic acid salt)

[00437] LC-MS: [M+H] = 413.2

[00438] ¹H NMR (400 MHz, DMSO) δ 8.19 (s, 1H), 7.05 (d, *J* = 8.4 Hz, 1H), 6.91 (d, *J* = 8.4 Hz, 1H), 6.85 (s, 1H), 6.68-6.60 (m, 2H), 6.39-6.34 (m, 1H), 4.23-4.20 (m, 1H), 3.99

(s, 2H), 3.81 (s, 3H), 3.69-3.60 (s, 2H), 3.41-3.33 (m, 2H), 2.11 (s, 3H), 1.48-1.45 (m, 1H), 1.35-1.28 (m, 1H), 1.13-1.09 (m, 2H), 0.79 (t, $J = 7.2$ Hz, 3H).

Example 9: 2-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanoic acid (Compound 9)



Step 1: (4-methoxy-3-methylphenyl)methanol

[00439] To a solution of methyl 4-methoxy-3-methylbenzoate (1.0 eq) in anhydrous THF (0.3M) at 0 °C was added dropwise LiAlH_4 (2.5 M, 2.5 eq) under N_2 . The resulting mixture was stirred at r.t. for 2 h, then the mixture was quenched with 1.25 M NaOH solution at 0 °C and extracted with EtOAc for three times. The organic layer was dried over Na_2SO_4 , concentrated under reduced pressure to give the title compound, which was used directly to next step without further purification.

Step 2: 4-(chloromethyl)-1-methoxy-2-methylbenzene

[00440] To a solution of (4-methoxy-3-methylphenyl)methanol (1.0 eq) in DCM (0.3M) at 0 °C was added SOCl_2 (2.0 eq). The resulting mixture was stirred at r.t. for 2 h.

Then the mixture was quenched with NaHCO_3 solution and extracted with DCM. The organic layer was dried over Na_2SO_4 , concentrated under reduced pressure to give the title compound as a yellow oil.

Step 3: 2-(4-methoxy-3-methylphenyl)acetonitrile

[00441] To a solution of 4-(chloromethyl)-1-methoxy-2-methylbenzene (1.0 eq) in 1:1 DMSO/DMF (0.5M) was added NaCN (3.0 eq). The resulting mixture was stirred at r.t. for 16 h, then water was added and the mixture was extracted with EtOAc. The organic layer was washed with brine twice, dried over Na_2SO_4 , concentrated under reduced pressure. The residue was purified by flash column chromatography (eluent: PE/EA = 100:1 to 30:1) to give the title compound as a yellowish oil.

Step 4: 2-(4-methoxy-3-methylphenyl)-2-methylpropanenitrile

[00442] A mixture of 2-(4-methoxy-3-methylphenyl)acetonitrile (1.0 eq) in DMSO (0.4M) at 0 °C was added iodomethane (10 eq) and 50% NaOH solution (6.0 eq). The resulting mixture was stirred at r.t. for 3 h. Then the mixture was diluted with water and EtOAc, the aqueous layer was back-extracted with EtOAc twice. The organic layer was combined, dried over Na_2SO_4 , concentrated under reduced pressure. The residue was purified by flash column chromatography (eluent: PE/EA = 100:1 to 30:1) to give the title compound as a slightly yellow oil.

Step 5: 2-(3-(bromomethyl)-4-methoxyphenyl)-2-methylpropanenitrile

[00443] To a solution of 2-(4-methoxy-3-methylphenyl)-2-methylpropanenitrile (1.0 eq) in CCl_4 (0.2M) were added NBS (1.1 eq) and AIBN (0.2 eq). The resulting mixture was stirred at 95 °C for 16 h. Then the mixture was filtered and the filtration was concentrated under reduced pressure to give the crude product, which was purified by column chromatography on silica (eluent: PE/EA = 100:1 to 30:1) to give the title compound as a yellow oil.

Step 6: ethyl 2-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)-3-oxobutanoate

[00444] To a solution of ethyl 3-oxobutanoate (1.2 eq) in anhydrous THF (0.17M) at 0 °C was added portion-wise NaH (60% in mineral oil, 1.3 eq). After stirring for 10 min, a solution of 2-(3-(bromomethyl)-4-methoxyphenyl)-2-methylpropanenitrile (1.0 eq) in THF (0.3M) was added dropwise into the above mixture over 10 min. The resulting mixture was stirred at 70 °C for 16 h, quenched with water and the mixture was extracted with EtOAc, dried over Na_2SO_4 , concentrated under reduced pressure. The residue was purified by column

chromatography on silica (eluent: PE/EA = 50:1 to 10:1) to give the title compound as a yellow oil.

Step 7: 2-(3-((2-amino-4-hydroxy-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanenitrile

[00445] A mixture of ethyl 2-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)-3-oxobutanoate (1.0 eq) and guanidine carbonate (1.0 eq) in MeOH (0.3M) was stirred at 65 °C for 16 h. The precipitated solid was collected by filtration. The solid was washed with water and dried in vacuum to give the title compound as a white solid.

Step 8: 2-(3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanenitrile

[00446] To a solution of 2-(3-((2-amino-4-hydroxy-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanenitrile (1.0 eq) in POCl₃ (0.3M) was stirred at 100 °C for 2 h. The reaction mixture was allowed to cool to r.t. and POCl₃ was evaporated to dryness under reduced pressure. The residue was diluted with water and adjusted pH to 8 with solid NaHCO₃. Then the mixture was stirred at 50 °C for 1 h, cooled to r.t. and the precipitated solid was collected by filtration. The filter cake was washed with water, dried in vacuum to give the title compound as a white solid.

Step 9: 2-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanenitrile

[00447] A mixture of 2-(3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanenitrile (1.0 eq) in NMP (0.2M) was added butan-1-amine (4.5 eq) and stirred at 110 °C for 16 h. Then water and EtOAc were added into the above mixture, the organic layer was dried over Na₂SO₄, concentrated under reduced pressure. The residue was purified by flash column chromatography (eluent: DCM/MeOH = 100:1 to 10:1) to give the title compound as a white solid.

Step 10: 2-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanoic acid


[00448] A mixture of 2-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanenitrile (1.0 eq) in 1:1 ethane-1,2-diol/H₂O (0.075M) was added KOH (10 eq) and stirred at 150 °C for 2 h. Then the mixture was allowed to cool to r.t. and adjusted pH to 8 with 1 M HCl solution. The precipitated solid was filtered and washed with water for three times. The filter cake was dissolved in MeCN, H₂O and 4M HCl in

dioxane solution, then the solution was freeze-dried to give the title compound as a white solid (HCl salt).

[00449] LC-MS: $[M+H] = 387.2$

[00450] ^1H NMR (400 MHz, DMSO) δ 12.34 (s, 1H), 7.92 (s, 1H), 7.45 (br s, 2H), 7.20 (d, $J = 8.4$ Hz, 1H), 6.96 (d, $J = 8.4$ Hz, 1H), 6.85 (s, 1H), 3.82 (s, 3H), 3.70 (s, 2H), 3.5-3.4 (m, 2H), 2.13 (s, 3H), 1.51-1.45 (m, 2H), 1.36 (s, 6H), 1.27-1.18 (m, 2H), 0.86 (t, $J = 7.2$ Hz, 3H).

Example 10: 5-(5-(2-(2H-tetrazol-5-yl)propan-2-yl)-2-methoxybenzyl)-N⁴-butyl-6-methylpyrimidine-2,4-diamine (Compound 10)




[00451] A mixture of 2-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanenitrile (1.0 eq, Example 9 – Step 9) in dioxane (0.06M), TMSN₃ (10 eq) and Bu₂SnO (2.0 eq) was sealed into a tube reactor. The mixture was stirred at 125 °C for 16 h. LCMS showed the reaction was completed. Then the mixture was concentrated under reduced pressure. The residue was purified by prep-HPLC (mobile phase CH₃CN/H₂O/HCOOH), freeze-dried to give the title compound as a white solid (formic acid salt)

[00452] LC-MS: $[M+H]^+ = 411.2$

[00453] ^1H NMR (400 MHz, DMSO) δ 6.97 (d, $J = 6.4$ Hz, 1H), 6.88 (d, $J = 8.4$ Hz, 1H), 6.72 (s, 1H), 6.44 (br s, 1H), 6.16 (br s, 2H), 3.79 (s, 3H), 3.58 (s, 2H), 3.27-3.25 (m, 2H), 1.98 (s, 3H), 1.61 (s, 6H), 1.44-1.40 (m, 2H), 1.24-1.18 (m, 2H), 0.85 (t, $J = 7.2$ Hz, 3H).

Example 11: N⁴-butyl-5-(2-methoxy-5-(2H-tetrazol-5-yl)benzyl)-6-methylpyrimidine-2,4-diamine (Compound 11)



Step 1: 4-bromo-2-(bromomethyl)-1-methoxybenzene

[00454] A mixture of 4-bromo-1-methoxy-2-methylbenzene (1.0 eq) in CCl₄ (0.3M), NBS (1.1 eq), and AIBN (0.2 eq) was stirred at 95 °C for 3 h. Then the mixture was filtered and the filtration was concentrated under reduced pressure to give the crude product, which was purified by column chromatography on a silica gel (eluent: PE) to give the title compound as a white solid.

Step 2: ethyl 2-(5-bromo-2-methoxybenzyl)-3-oxobutanoate

[00455] To a stirred solution of ethyl 3-oxobutanoate (1.2 eq) in anhydrous THF (0.5M) at 0 °C was added portion-wise 60% NaH (1.3 eq). After stirring for 10 min, a solution of 4-bromo-2-(bromomethyl)-1-methoxybenzene (1.0 eq) in THF (0.8M) was added dropwise into the above mixture over 10 min. Then the resulting mixture was stirred at 70 °C for 16 h, water was added and extracted with EtOAc. The combined organic phase was dried over Na₂SO₄, concentrated under reduced pressure. The residue was purified by column chromatography on a silica gel (eluent: PE/EA = 100:1 to 10:1) to give the title compound as yellow oil.

Step 3: 2-amino-5-(5-bromo-2-methoxybenzyl)-6-methylpyrimidin-4-ol

[00456] A mixture of ethyl 2-(5-bromo-2-methoxybenzyl)-3-oxobutanoate (1.0 eq) and guanidine carbonate (1.0 eq) in MeOH (0.3M) was stirred at 65 °C for 16 h. After the reaction was completed, the mixture was cooled to r.t. The precipitated solid was collected by filtration and the filter cake was washed with water, dried in vacuum to give the title compound as a white solid.

Step 4: 5-(5-bromo-2-methoxybenzyl)-4-chloro-6-methylpyrimidin-2-amine

[00457] To a solution of 2-amino-5-(5-bromo-2-methoxybenzyl)-6-methylpyrimidin-4-ol (1.0 eq) in POCl₃ (0.5M) was stirred at 100 °C for 16 h. The reaction mixture was allowed to cool to r.t. and POCl₃ was evaporated to dryness under reduced pressure. The residue was diluted with water and adjusted pH to 8 with NaHCO₃ solid, and then the mixture was stirred at 50 °C for 1 h, cooled to r.t. The precipitated solid was collected by filtration. The filter cake was washed with water and MeOH, dried in vacuum to give the title compound as a white solid.

Step 5: 5-(5-bromo-2-methoxybenzyl)-N⁴-butyl-6-methylpyrimidine-2,4-diamine

[00458] A mixture of 5-(5-bromo-2-methoxybenzyl)-4-chloro-6-methylpyrimidin-2-amine (1.0 eq) in NMP (0.4M) and butan-1-amine (4.5 eq) was stirred at 130 °C for 16 h. The mixture was diluted with water and EtOAc, the organic layer was washed with brine and dried over Na₂SO₄, concentrated under reduced pressure. The residue was purified by flash column chromatography (eluent: DCM/MeOH = 500:1 to 50:1) to give the title compound as a white solid.

Step 6: 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzonitrile

[00459] A mixture of 5-(5-bromo-2-methoxybenzyl)-N⁴-butyl-6-methylpyrimidine-2,4-diamine (1.0 eq) in DMF (0.1M), Zn(CN)₂ (1.5 eq), Pd₂(dba)₃ (0.08 eq), and X-Phos (0.2 eq) was stirred at 110 °C for 1 h under Argon atmosphere. The resulting mixture was allowed to cool to r.t. and diluted with water and EtOAc. The organic layer was separated, dried over Na₂SO₄, concentrated under reduced pressure. The residue was purified by flash column chromatography (eluent: DCM/MeOH = 100:1 to 10:1) to give the title compound as a white solid.


Step 7: N⁴-butyl-5-(2-methoxy-5-(2H-tetrazol-5-yl)benzyl)-6-methylpyrimidine-2,4-diamine

[00460] A mixture of 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzonitrile (1.0 eq) in dioxane (0.07M), TMSN₃ (10 eq) and Bu₂SnO (2.0 eq) was stirred at 120 °C for 4 h. LCMS showed the reaction was completed. Then the mixture was concentrated under reduced pressure. The residue was purified by prep-HPLC (mobile phase CH₃CN/H₂O/HCOOH), freeze-dried to give the title compound as a white solid (formic acid salt).

[00461] LC-MS: [M+H] = 369.1

[00462] ¹H NMR (400 MHz, DMSO) δ 7.84 (d, *J* = 8.4 Hz, 1H), 7.56 (br s, 1H), 7.44 (s, 1H), 7.09 (d, *J* = 8.4 Hz, 1H), 7.05 (br s, 2H), 3.90 (s, 3H), 3.73 (s, 2H), 3.4-3.3 (m, 2H), 2.13 (s, 3H), 1.49-1.41 (m, 2H), 1.19-1.12 (m, 2H), 0.76 (t, *J* = 7.6 Hz, 3H).

Example 12: (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)phosphonic acid (Compound 12)



Preparation of N⁴-butyl-5-(5-iodo-2-methoxybenzyl)-6-methylpyrimidine-2,4-diamine

[00463] Preparation of *N*⁴-butyl-5-(5-iodo-2-methoxybenzyl)-6-methylpyrimidine-2,4-diamine followed the protocols described for Example 11 – Step 1 to 5, but using 4-iodo-1-methoxy-2-methylbenzene instead of 4-bromo-1-methoxy-2-methylbenzene in Step 1.

Step 1: diethyl (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)phosphonate

[00464] To a solution of *N*⁴-butyl-5-(5-iodo-2-methoxybenzyl)-6-methylpyrimidine-2,4-diamine (1.0 eq) in toluene (0.06M), diethylphosphite (3.5 eq) and triethylamine (6.0 eq) was added Pd(dppf)Cl₂ (0.05 eq). The mixture was stirred at 110 °C for 16 h. The reaction mixture was allowed to cool to r.t. and was evaporated to dryness under reduced pressure to afford the crude product which was purified by flash column chromatography (eluent: DCM/MeOH = 50:1 to 10:1) to give the title compound.


Step 2: (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-phosphonic acid

[00465] To a solution of diethyl (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)phosphonate (1.0 eq) in DCM (0.08M) was added dropwise TMSBr (20 eq). The mixture was stirred at rt for 2 h. The reaction mixture was diluted with water and DCM, the organic layer was washed with brine and dried over Na₂SO₄, concentrated under reduced pressure. The residue was purified by prep-HPLC (mobile phase CH₃CN/H₂O/HCOOH), freeze-dried to give the title compound as a white solid (formic acid salt).

[00466] LC-MS: [M+H] = 381.3

[00467] ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.00 (s, 1H), 7.99 (t, *J* = 4.8 Hz, 1H), 7.59-7.53 (m, 1H), 7.47 (br s, 2H), 7.10-7.06 (m, 2H), 3.89 (s, 3H), 3.72 (s, 2H), 3.40-3.35 (m, 2H), 2.10 (s, 3H), 1.53-1.45 (m, 2H), 1.25-1.19 (m, 2H), 0.87 (t, *J* = 7.2 Hz, 3H).

Example 13: 5-(5-(1-(2H-tetrazol-5-yl)cyclopropyl)-2-methoxybenzyl)-N⁴-butyl-6-methylpyrimidine-2,4-diamine (Compound 13)



Step 1: 1-(4-methoxy-3-methylphenyl)cyclopropane-1-carbonitrile:

[00468] To a flask containing 2-(4-methoxy-3-methylphenyl)acetonitrile (1.0 eq, Example 9 – Step 3), 1-bromo-2-chloroethane (1.5 eq), Et₃NBnCl (0.02 eq) was added 0.77g/mL NaOH solution (6.0 eq). The resulting mixture was stirred overnight at 50 °C. The mixture was partitioned between EA/water. The organic layer was dried over Na₂SO₄, concentrated and purified by flash chromatography on silica (eluent PE/EA = 100:1~20:1) to give the title compound.

Preparation of 1-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)cyclopropane-1-carbonitrile

[00469] Preparation of 1-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)cyclopropane-1-carbonitrile following protocols described in Example 9: Step 5 to 9, but using 1-(4-methoxy-3-methylphenyl)cyclopropane-1-carbonitrile instead of 2-(4-methoxy-3-methylphenyl)-2-methylpropanenitrile in Step 5.


Preparation of 5-(5-(1-(2H-tetrazol-5-yl)cyclopropyl)-2-methoxybenzyl)-N4-butyl-6-methylpyrimidine-2,4-diamine:

[00470] A mixture of 1-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)cyclopropane-1-carbonitrile (1.0 eq) in dioxane (0.1M), TMSN₃ (10.0 eq), Bu₂SnO (2.0 eq) was stirred overnight at 120 °C. The reaction was monitored by LC-MS. The mixture was concentrated and purified by prep-HPLC (mobile phase CH₃CN/H₂O/HCOOH), freeze-dried to give the title compound as a white powder (formic acid salt).

[00471] LC-MS: [M+H]⁺=409.2

[00472] ¹H NMR (400 MHz, DMSO-*d*₆) δ 8.19 (s, 1H), 7.16 (d, *J* = 8.8 Hz, 1H), 6.92 (d, *J* = 8.4 Hz, 1H), 6.77 (s, 1H), 6.69 (br s, 1H), 6.42 (br s, 2H), 3.82 (s, 3H), 3.61 (s, 2H), 3.30-3.25 (m, 2H), 2.03 (s, 3H), 1.43-1.35 (m, 4H), 1.21-1.11 (m, 4H), 0.83 (t, *J* = 7.2 Hz, 3H).

Example 14: 1-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)cyclopropane-1-carboxylic acid (Compound 14)




[00473] A mixture of 1-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)cyclopropane-1-carbonitrile (1.0 eq) in 1:1 ethylene glycol/H₂O (0.1M) and KOH (10 eq) was stirred for 3 h at 150 °C. The reaction was monitored by LC-MS until completion. The mixture was neutralized with 1 N HCl to pH = 7. The resulting suspension was filtered. The filter cake was washed with water. To the solid was added MeCN, H₂O and

4N HCl/dioxane. The solution was freeze-dried to give the title compound as a white solid (HCl salt).

[00474] LC-MS: $[M+H]^+ = 385.2$

[00475] ^1H NMR (400 MHz, DMSO- d_6) δ 12.21 (s, 1H), 12.16 (s, 1H), 7.90 (t, $J = 5.2$ Hz, 1H), 7.41 (br s, 2H), 7.16 (d, $J = 6.0$ Hz, 1H), 6.92 (d, $J = 8.4$ Hz, 1H), 6.81 (s, 1H), 3.81 (s, 3H), 3.68 (s, 2H), 3.4-3.3 (m, 2H), 2.12 (s, 3H), 1.51-1.43 (m, 2H), 1.38 (s, 2H), 1.26-1.17 (m, 2H), 0.99 (s, 2H), 0.86 (t, $J = 7.2$ Hz, 3H).

Example 15: 3-(2-amino-6-(butylamino)-5-(2-methoxybenzyl)pyrimidin-4-yl)propanoic acid (Compound 15)



Step 1: 1-(bromomethyl)-2-methoxybenzene

[00476] To a solution of 1-methoxy-2-methylbenzene (1 eq) in CCl_4 (0.9M) was added AIBN (0.2 eq) and NBS (1.1 eq). The reaction was heated at reflux for 2 h. After the reaction was completed, the reaction was concentrated and diluted with water. The aqueous phase was extracted with DCM. Combined organic layers were washed with water then brine, dried over Na_2SO_4 , and concentrated to give the title compound as a light yellow oil.

Step 2: dimethyl 2-(2-methoxybenzyl)malonate

[00477] To a solution of dimethyl malonate (1.2 eq) in THF (0.4M) was added NaH (1.25eq) at 0°C in portions under nitrogen. The solution was stirred at 0°C for 15 min, and 1-

(bromomethyl)-2-methoxybenzene (1.0 eq) was added. The reaction was stirred for 2 h at room temperature. After the reaction was completed, the reaction was quenched by addition of water. The aqueous phase was extracted with EA. Combined organic layers were washed with water, then brine, dried over Na₂SO₄, and concentrated. The crude product was purified by column chromatography (PE/EA=40:1) to give the title compound as light yellow oil.

Step 3: 2-amino-5-(2-methoxybenzyl)pyrimidine-4,6-diol

[00478] To a solution of dimethyl 2-(2-methoxybenzyl)malonate (1.0 eq) in MeOH (0.4M) was added guanidine carbonate (1.0 eq). The reaction was heated at 75 °C for 16 h under nitrogen. After cooling, the solid particles were filtered and collected from the solution to give the title compound as a white solid.

Step 4: 4,6-dichloro-5-(2-methoxybenzyl)pyrimidin-2-amine

[00479] To a suspension of 2-amino-5-(2-methoxybenzyl)pyrimidine-4,6-diol (1.0 eq) in POCl₃ (0.4M) was added 2 drops of DMF. The reaction was heated at 100 °C for 15 h under nitrogen. After cooling, the solution was poured into ice-water. The solid particles were filtered and collected from the solution to give the title compound as a white solid.

Step 5: ethyl (E)-3-(2-amino-6-chloro-5-(2-methoxybenzyl)pyrimidin-4-yl)acrylate

[00480] To a solution of 4,6-dichloro-5-(2-methoxybenzyl)pyrimidin-2-amine (1.0 eq) in 3:1 dioxane/H₂O (0.14M) was added ethyl (*E*)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)acrylate (1.2 eq), K₂CO₃ (2.0 eq), and Pd(dppf)Cl₂ (0.1 eq). The mixture was stirred at 100 °C for 3 h under nitrogen. The reaction was diluted with water, and the aqueous phase was extracted with EA. Combined organic layers were washed with water, then brine, dried over Na₂SO₄, and concentrated. The crude product was purified by column chromatography (PE/EA=40:3) to give the title compound as light yellow solid.

Step 6: ethyl (E)-3-(2-amino-6-(butylamino)-5-(2-methoxybenzyl)pyrimidin-4-yl)acrylate

[00481] To a solution of ethyl (*E*)-3-(2-amino-6-chloro-5-(2-methoxybenzyl)pyrimidin-4-yl)acrylate (1.0 eq) in NMP (0.2M) was added butan-1-amine (3.0 eq) and DIEA (3.0 eq), and the mixture was stirred at 150 °C for 2 h. After the reaction was completed, the reaction was diluted with water. The aqueous phase was extracted with EA. Combined organic layers were washed with water, then brine, dried over Na₂SO₄, and concentrated. The crude product was purified by column chromatography (DCM/MeOH=10:1) to give the title compound as a yellow oil.

Step 7: ethyl 3-(2-amino-6-(butylamino)-5-(2-methoxybenzyl)pyrimidin-4-yl)propanoate


[00482] To a solution of ethyl (*E*)-3-(2-amino-6-(butylamino)-5-(2-methoxybenzyl)pyrimidin-4-yl)acrylate (1.0 eq) in MeOH (0.04M) was added Pd/C (1/3 weight eq). The mixture was stirred at room temperature for 1 h under hydrogen. After the reaction was completed, the solid particles were filtered off and the filtrate was concentrated to give the title compound as a yellow oil.

Step 8: 3-(2-amino-6-(butylamino)-5-(2-methoxybenzyl)pyrimidin-4-yl)propanoic acid

[00483] To a solution of ethyl 3-(2-amino-6-(butylamino)-5-(2-methoxybenzyl)pyrimidin-4-yl)propanoate (1eq) in 1:1 MeOH/H₂O (0.04M) was added NaOH (5.0 eq). The solution was stirred at room temperature for 1 h. After the reaction was completed, the pH was adjusted to 7 using acetic acid, and the solution was concentrated and purified by prep-HPLC (mobile phase: NH₃.H₂O/MeCN/H₂O) to give the title compound as a white solid (ammonium salt).

[00484] LC-MS: [M+H] = 359.4

[00485] ¹H NMR(400 MHz, MeOD) δ 7.23 (t, *J* = 6.8 Hz, 1H), 6.99 (d, *J* = 8.0 Hz, 1H), 6.91-6.88 (m, 2H), 3.90 (s, 3H), 3.80 (s, 2H), 3.44 (t, *J* = 6.8 Hz, 2H), 2.85 (t, *J* = 6.0 Hz, 2H), 2.42 (t, *J* = 6.0 Hz 2H), 1.51-1.46 (m, 2H), 1.25-1.19 (m, 2H), 0.88 (t, *J* = 6.8 Hz, 3H).

Example 16: 2-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzamido)ethane-1-sulfonic acid (Compound 16)

[00486] Preparation of methyl 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate followed Example 1 – Steps 1 to 4, but using methyl 3-(bromo-methyl)-4-methoxybenzoate as the starting material in Step 1.

Step 1: 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoic acid

[00487] To a suspension of methyl 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate (1.0 eq) in MeOH (0.3M) was added 1N NaOH (5.0 eq). The

resulting mixture was stirred at 60 °C for 2 h. The reaction was monitored by TLC. The mixture was cooled to rt. The mixture was neutralized with 2N HCl and purified to give the title compound.


Step 2: 2-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenz-amido)ethane-1-sulfonic acid

[00488] To a solution of 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxy-benzoic acid (1.0 eq) in DMF (0.6M) was added HATU (1.0 eq), 2-aminoethane-1-sulfonic acid (1.1 eq) and DIEA (3.0 eq). The resulting mixture was stirred at rt for 3 h. The mixture was filtered and the filtrate was purified by prep-HPLC (mobile phase CH₃CN/H₂O/HCOOH), then freeze-dried to give the title compound as a white solid.

[00489] LC-MS: [M+H]⁺ = 452.2

[00490] ¹H NMR (400 MHz, DMSO) δ 11.89 (s, 1H), 8.35 (t, *J* = 3.2 Hz, 1H), 7.90 (s, 1H), 7.68 (d, *J* = 8.4 Hz, 1H), 7.40 (br s, 2H), 7.26 (s, 1H), 7.09 (d, *J* = 8.4 Hz, 1H), 3.90 (s, 3H), 3.71 (s, 2H), 3.48-3.42 (m, 2H), 3.39-3.33 (m, 2H), 2.62 (t, *J* = 6.8 Hz, 2H), 2.12 (s, 3H), 1.49-1.41 (m, 2H), 1.23-1.15 (m, 2H), 0.82 (t, *J* = 7.2 Hz, 3H).

Example 17: 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzenesulfonic acid (Compound 17)



[00491] Preparation of 5-(5-bromo-2-methoxybenzyl)-N⁴-butyl-6-methylpyrimidine-2,4-diamine followed Example 1 – Steps 1 to 4, but 4-bromo-2-(bromomethyl)-1-methoxybenzene as the starting material in Step 1.

Step 1: N⁴-butyl-5-(2-methoxybenzyl)-6-methylpyrimidine-2,4-diamine

[00492] To a stirred solution of 5-(5-bromo-2-methoxybenzyl)-N⁴-butyl-6-methylpyrimidine-2,4-diamine (1.0 eq) in MeOH (0.1M) was added Pd/C (10% wt eq). The resulting mixture was stirred at rt for 1 h under H₂ atmosphere, and then filtered. The filtrate was concentrated to give the title compound.

Step 2: 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzene-sulfonyl chloride


[00493] N⁴-butyl-5-(2-methoxybenzyl)-6-methylpyrimidine-2,4-diamine (1.0 eq) in ClSO₃H (0.7M) was stirred at rt for 2 h, and then poured into ice water. The mixture was partitioned between DCM/water. The organic layer was dried over Na₂SO₄, filtered, concentrated to give the title compound.

Step 3: 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzenesulfonic acid

[00494] To a stirred solution of 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzene-sulfonyl chloride (1.0 eq) in 1:1 MeCN/H₂O (0.025M) was added K₂CO₃ (2.0 eq). The resulting mixture was stirred overnight at rt and then concentrated. The residue was purified by prep-HPLC (mobile phase CH₃CN/H₂O/HCOOH), freeze-dried to give the title compound.

[00495] LC-MS: [M-H]⁺ = 379.2

[00496] ¹H NMR (400 MHz, DMSO-*d*₆) δ 11.92 (br s, 1H), 7.79 (br s, 1H), 7.46 (d, *J* = 6.8 Hz, 1H), 7.22 (br s, 2H), 7.04 (s, 1H), 6.95 (d, *J* = 8.4 Hz, 1H), 3.85 (s, 3H), 3.68 (s, 2H), 3.37-3.34 (m, 2H), 2.08 (s, 3H), 1.54-1.44 (m, 2H), 1.26-1.23 (m, 2H), 0.86 (t, *J* = 7.2 Hz, 3H).

Example 18: (S)-3-((5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-methylpyrimidin-4-yl)amino)heptan-1-ol (Compound 18)


[00497] The title compound is prepared following the protocols described in Example 8, but using (S)-3-aminoheptan-1-ol (from J. Med. Chem. 2016, 59, 7936–7949) instead of

(*S*)-2-aminopentane-1-ol in Step A. In the final step, the crude product was purified by prep-HPLC (mobile phase CH₃CN/H₂O/HCOOH), then freeze-dried to give the title compound as a white solid (formic acid salt).

[00498] LC-MS: [M+H]⁺ = 441.3

[00499] ¹H NMR (400 MHz, DMSO) δ 8.22 (s, 1H), 7.03 (d, *J* = 8.8 Hz, 1H), 6.90 (d, *J* = 8.4 Hz, 1H), 6.78 (s, 1H), 6.52 (s, 2H), 6.35 (d, *J* = 8.0 Hz, 1H), 4.27-4.21 (m, 1H), 3.99-3.91 (m, 2H), 3.80 (s, 3H), 3.62 (s, 2H), 3.39-3.26 (m, 3H), 2.07 (s, 3H), 1.64-1.50 (m, 2H), 1.48-1.37 (m, 2H), 1.24-1.14 (m, 2H), 1.12-1.04 (m, 2H), 0.76 (t, *J* = 7.2 Hz, 3H).

Example 19: 3-(2-amino-6-(butylamino)-5-(4-methoxyphenethyl)pyrimidin-4-yl)propanoic acid (Compound 19)



Step 1: 4,6-dichloropyrimidin-2-amine

[00500] To a solution of 2-aminopyrimidine-4,6-diol (1.0 eq) in POCl₃ (1M) was stirred at 100 °C for 4 h. The reaction mixture was allowed to cool to rt and the mixture was poured into ice-water. The precipitated solid was collected by filtration. The solid was washed with water and dried in vacuum to give the title compound as a yellow solid.

Step 2: ethyl (E)-3-(2-amino-6-chloropyrimidin-4-yl)acrylate

[00501] To a solution of 4,6-dichloropyrimidin-2-amine (1.0 eq) in 3:1 dioxane/H₂O (0.3M) was added ethyl (*E*)-3-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)acrylate (1.1 eq), K₂CO₃ (2.0 eq), and Pd(dppf)Cl₂ (0.1 eq). The mixture was stirred at 100 °C for 2 h under nitrogen. The reaction was diluted with water, and the aqueous layer was extracted with EA. The combined organic layers were washed with water, brine, dried over Na₂SO₄, and concentrated. The crude product was purified by column chromatography (PE/EA=10:1) to give the title compound as a yellow solid.

Step 3: ethyl (E)-3-(2-amino-6-(butylamino)pyrimidin-4-yl)acrylate

[00502] To a solution of ethyl (*E*)-3-(2-amino-6-chloropyrimidin-4-yl)acrylate (1.0 eq) in NMP (0.4M) was added butan-1-amine (3.0 eq) and DIEA (3.0 eq). The mixture was stirred at 150 °C for 3 h. The reaction was diluted with water, and the aqueous phase was extracted with EA. The combined organic layers were washed with water, brine, dried over Na₂SO₄, and concentrated. The crude product was purified by column chromatography (PE/EA=1:1) to give the title compound as a yellow solid.

Step 4: ethyl 3-(2-amino-6-(butylamino)pyrimidin-4-yl)propanoate

[00503] To a solution of ethyl (*E*)-3-(2-amino-6-(butylamino)pyrimidin-4-yl)acrylate (1.0 eq) in MeOH (0.2M) was added Pd/C (20% wt eq). The mixture was stirred at 25 °C for 2 h under H₂ and then filtered. The filtrate was concentrated to give the title compound as a colored solid.

Step 5: ethyl 3-(2-amino-6-(butylamino)-5-iodopyrimidin-4-yl)propanoate

[00504] To a solution of ethyl 3-(2-amino-6-(butylamino)pyrimidin-4-yl)propanoate (1.0 eq) in 2:1 DCM/H₂O (0.05M) was added NaOH (2.0 eq), followed by I₂ (1.0 eq). The mixture was stirred at 25 °C for 16 h. The reaction was diluted with water, and the aqueous phase was extracted with EA. The combined organic layers were washed with water, brine, dried over Na₂SO₄, and concentrated. The crude product was purified by column chromatography (PE/EA=4:3) to give the title compound as a yellow solid.

Step 6: ethyl 3-(2-amino-6-(butylamino)-5-((4-methoxyphenyl)ethynyl)pyrimidin-4-yl)propanoate

[00505] To a solution of ethyl 3-(2-amino-6-(butylamino)-5-iodopyrimidin-4-yl)propanoate (1.0 eq) in DMF (0.1M) was added 1-ethynyl-4-methoxybenzene (1.2 eq), TEA (1/5th reaction volume), CuI (0.2 eq), and Pd(PPh₃)₂Cl₂ (0.1 eq). The mixture was stirred

at 50 °C for 16 h under nitrogen. The reaction was diluted with water, and the aqueous phase was extracted with EA. The combined organic layers were washed with water, brine, dried over Na₂SO₄, and concentrated. The crude product was purified by column chromatography (PE/EA=1:1) to give the title compound as a brown oil.

Step 7: ethyl 3-(2-amino-6-(butylamino)-5-(4-methoxyphenethyl)pyrimidin-4-yl)propanoate

[00506] To a solution of ethyl 3-(2-amino-6-(butylamino)-5-(4-methoxyphenethyl)pyrimidin-4-yl)propanoate (1.0 eq) in MeOH (0.08M) was added Pd/C (30% wt eq). The mixture was stirred at 50 °C for 2 h under H₂ and then filtered. The filtrate was concentrated to give the title compound as a yellow oil.


Step 8: 3-(2-amino-6-(butylamino)-5-(4-methoxyphenethyl)pyrimidin-4-yl)propanoic acid

[00507] To a solution of ethyl 3-(2-amino-6-(butylamino)-5-(4-methoxyphenethyl)pyrimidin-4-yl)propanoate (1.0 eq) in 1:1 MeOH/H₂O (0.2M) was added NaOH (5.0 eq). The resulting mixture was stirred at 25 °C for 1 h. The pH was adjusted to 7 using acetic acid, and the solution was concentrated and purified by prep-HPLC (mobile phase: NH₄OH/MeCN/H₂O) to give the title compound as a white solid (ammonium salt).

[00508] LC-MS: [M+H]⁺ = 373.4

[00509] ¹H NMR (400 MHz, CD₃OD) δ 7.00 (d, *J* = 8.4 Hz, 2H), 6.81 (d, *J* = 8.8 Hz, 2H), 3.75 (s, 3H), 3.48 (t, *J* = 7.6 Hz, 2H), 2.75-2.71 (m, 4H), 2.43 (t, *J* = 6.0 Hz, 2H), 2.18 (t, *J* = 6.4 Hz, 2H), 1.60-1.54 (m, 2H), 1.40-1.35 (m, 2H), 0.97 (t, *J* = 6.8 Hz, 3H)

Example 20A: (S)-3-((5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-methylpyrimidin-4-yl)amino)heptanoic acid (Compound 20)



[00510] Methyl 3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate was prepared following Example 1 – Steps 1 to 3, but using methyl 3-(bromomethyl)-4-methoxybenzoate as the starting material instead of methyl 4-(bromomethyl)-3-methoxybenzoate.

Step 1: (3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)methanol

[00511] To a stirred solution of methyl 3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate (1.0 eq) in THF (0.2M) at 0 °C was added dropwise 2.5M LiAlH₄ (2.5 eq). The resulting mixture was stirred at rt for 1 h. Then the mixture was quenched with 1 M NaOH solution and filtered. The filtration was extracted with EtOAc. The organic layer was dried over Na₂SO₄, filtered, concentrated under reduced pressure. The residue was purified by flash column chromatography (eluent DCM/MeOH = 100:1 to 10:1) to give the title compound as a yellow solid.

Step 2: *tert*-butyl (S)-3-((2-amino-5-(5-(hydroxymethyl)-2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoate

[00512] A mixture of *tert*-butyl (S)-3-aminoheptanoate (4.0 eq; prepared following procedures reported in J. Med. Chem. 2016, 59, 7936–7949) and (3-((2-amino-4-chloro-6-

methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-methanol (1.0 eq) in NMP (0.8M) was stirred at 120 °C for 16 h. Then water and EtOAc were added into the reaction mixture. The organic layer was separated, dried over Na₂SO₄, concentrated under reduced pressure. The residue was purified by flash column chromatography (eluent: DCM/MeOH = 100:1 to 20:1) to give the title compound as a yellow oil.

Step 3 to 5: *tert*-butyl (S)-3-((5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-methylpyrimidin-4-yl)amino)heptanoate

[00513] The title compound was prepared following Example 2 – Steps 1 to 3, but using *tert*-butyl (S)-3-((2-amino-5-(5-(hydroxymethyl)-2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoate as the starting material. The crude product was purified by prep-HPLC (mobile phase CH₃CN/H₂O/HCOOH), and freeze-dried to give the title compound as a white solid (formic acid salt).

[00514] LC-MS: [M+H]⁺ = 511.4

[00515] ¹H NMR (400 MHz, DMSO) δ 8.20 (s, 1H), 7.04 (dd, *J* = 8.4, 2.0 Hz, 1H), 6.93 (d, *J* = 8.0 Hz, 1H), 6.77 (s, 1H), 6.60 (br s, 2H), 6.39 (br s, 1H), 4.55-4.52 (m, 1H), 4.02 (s, 2H), 3.83 (s, 3H), 3.67-3.57 (m, 2H), 2.37 (d, *J* = 6.8 Hz, 2H), 2.07 (s, 3H), 1.46-1.38 (m, 2H), 1.32 (s, 9H), 1.26-1.13 (m, 2H), 1.09-1.01 (m, 2H), 0.77 (t, *J* = 7.2 Hz, 3H).


Step 6: (S)-3-((5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-methylpyrimidin-4-yl)amino)heptanoic acid

[00516] *Tert*-butyl (S)-3-((5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-methylpyrimidin-4-yl)amino)heptanoate (1 eq) was added 3.0 M HCl/dioxane (20 eq) at 0 °C, and warmed to rt. After stirring at 1 h, the mixture was concentrated under reduced pressure. The residue was purified by prep-HPLC (mobile phase CH₃CN/H₂O/HCOOH), freeze-dried to give the title compound as a white solid.

[00517] LC-MS: [M+H]⁺ = 455.2

[00518] ¹H NMR (400 MHz, DMSO) δ 7.35 (br s, 1H), 7.07 (br s, 2H), 6.96 (d, *J* = 7.6 Hz, 1H), 6.88 (d, *J* = 8.4 Hz, 1H), 6.76 (s, 1H), 4.56-4.54 (m, 1H), 4.07-3.97 (m, 2H), 3.79 (s, 3H), 3.68-3.66 (m, 2H), 2.46-2.38 (m, 2H), 2.05 (s, 3H), 1.53-1.50 (m, 2H), 1.27-1.16 (m, 4H), 0.82 (t, *J* = 6.8 Hz, 3H).

Example 20B: (S)-3-((5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-methylpyrimidin-4-yl)amino)heptanoic acid (Compound 20)



Step 1: *tert*-butyl (E)-hept-2-enoate

[00519] A mixture of pentanal (1.0 eq) and *tert*-butyl 2-(triphenyl-λ⁵-phosphanylidene)acetate (1.05 eq) was stirred at 50 °C for 16 h. Then the mixture was concentrated under reduced pressure, and petroleum ether (PE) was added. The solids were filtered off, and the filtrate was evaporated to dryness, which was purified by flash column chromatography (eluent: 0-1% EA in PE) to give the title compound as a yellow oil.

Step 2: *tert*-butyl (*S*)-3-(benzyl((*S*)-1-phenylethyl)amino)heptanoate

[00520] To a solution of (*S*)-*N*-benzyl-1-phenylethan-1-amine (1.3 eq) in THF (0.9M) at -78 °C was added dropwise 2.5M *n*-BuLi (1.2 eq) over 20 min. The mixture was stirred at -78 °C for 10 min, then *tert*-butyl (*E*)-hept-2-enoate (1.0 eq) in THF (0.7M) was added dropwise into the above mixture. The resulting mixture was stirred at -78 °C for 30 min. Then the mixture was quenched with aq. NH₄Cl and extracted with EtOAc. The organic layer was dried over Na₂SO₄, filtered, concentrated under reduced pressure. The residue was purified by flash column chromatography (eluent: 0-1% EA in PE) to give the title compound as a yellow oil.

Step 3: *tert*-butyl (*S*)-3-aminoheptanoate

[00521] To a solution of *tert*-butyl (*S*)-3-(benzyl((*S*)-1-phenylethyl)amino)heptanoate (1.0 eq) in MeOH (0.5M) was added 10%wt Pd/C (1/10th weight equivalent). The resulting mixture was stirred at 50 °C for 16 h under H₂. The solid was filtered off and the filtrate was concentrated under reduced pressure to give the title compound as a brown oil, which was used directly to next step without further purification.

Step 4: (3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-methanol

[00522] To a stirred solution of methyl 3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate (1.0 eq; from Example 3B – Step 4) in THF (0.2M) at 0 °C was added dropwise 2.5M LiAlH₄ (2.5 eq). The resulting mixture was stirred at rt for 1 h. Then the mixture was quenched with 1 M NaOH solution and filtered. The filtration was extracted with EtOAc. The organic layer was dried over Na₂SO₄, filtered, concentrated under reduced pressure. The residue was purified by flash column chromatography (eluent DCM/MeOH = 100:1 to 10:1) to give the title compound as a yellow solid.

Step 5: *tert*-butyl (*S*)-3-((2-amino-5-(5-(hydroxymethyl)-2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoate

[00523] A mixture of *tert*-butyl (*S*)-3-aminoheptanoate (4.0 eq) and (3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-methanol (1.0 eq) in NMP (0.8M) was stirred at 120 °C for 16 h. Then water and EtOAc were added into the reaction mixture. The organic layer was separated, dried over Na₂SO₄, concentrated under reduced pressure. The residue was purified by flash column chromatography (eluent: DCM/MeOH = 100:1 to 20:1) to give the title compound as a yellow oil.

Step 6: *tert*-butyl (S)-3-((2-amino-5-(5-(chloromethyl)-2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoate

[00524] To a stirred solution of *tert*-butyl (S)-3-((2-amino-5-(5-(hydroxymethyl)-2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoate (1.0 eq) in DCM (0.05M) at 0 °C was added SOCl₂ (2.0 eq). The resulting mixture was stirred at 0 °C for 1 h. Then the mixture was quenched with NaHCO₃ solution and extracted with DCM. The organic layer was dried over Na₂SO₄, concentrated under reduced pressure to give the title compound as a yellow oil.

Step 7: *tert*-butyl (S)-3-((2-amino-5-(5-(cyanomethyl)-2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoate

[00525] A mixture of *tert*-butyl (S)-3-((2-amino-5-(5-(chloromethyl)-2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoate (1.0 eq) in 1:1 DMSO/DMF (0.12M) and NaCN (3.0 eq) was stirred at rt for 16 h. The resulting mixture was partitioned between EtOAc and water. The organic layer was washed with brine, dried over Na₂SO₄, concentrated under reduced pressure. The residue was purified by flash column chromatography (eluent: DCM/MeOH = 100:1 to 20:1) to give the title compound as a yellow oil.

Step 8: *tert*-butyl (S)-3-((5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-methylpyrimidin-4-yl)amino)heptanoate

[00526] A mixture of *tert*-butyl (S)-3-((2-amino-5-(5-(cyanomethyl)-2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoate (1.0 eq) in dioxane (1.2M), TMSN₃ (10 eq) and Bu₂SnO (2.0 eq) was stirred at 120 °C for 2 h. Then the mixture was concentrated under reduced pressure to give crude product as brown oil, which was used directly to next step without further purification.


Step 9: (S)-3-((5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-methylpyrimidin-4-yl)amino)heptanoic acid

[00527] *Tert*-butyl (S)-3-((5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-methylpyrimidin-4-yl)amino)heptanoate (1 eq) was added 3.0 M HCl/dioxane (20 eq) at 0 °C, and warmed to rt. After stirring at 1 h, the mixture was concentrated under reduced pressure. The residue was purified by prep-HPLC (mobile phase CH₃CN/H₂O/HCOOH), freeze-dried to give the title compound as a white solid (formic acid salt).

[00528] LC-MS: [M+H]⁺ = 455.2

[00529] ^1H NMR (400 MHz, DMSO) δ 7.35 (br s, 1H), 7.07 (br s, 2H), 6.96 (d, J = 7.6 Hz, 1H), 6.88 (d, J = 8.4 Hz, 1H), 6.76 (s, 1H), 4.56-4.54 (m, 1H), 4.07-3.97 (m, 2H), 3.79 (s, 3H), 3.68-3.66 (m, 2H), 2.46-2.38 (m, 2H), 2.05 (s, 3H), 1.53-1.50 (m, 2H), 1.27-1.16 (m, 4H), 0.82 (t, J = 6.8 Hz, 3H).


Example 20C: (R)-3-((5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-methylpyrimidin-4-yl)amino)heptanoic acid (Compound 20C)



[00530] The title compound was prepared by following the procedure described in Example 20B, but using (*R*)-*N*-benzyl-1-phenylethan-1-amine instead of (*S*)-*N*-benzyl-1-phenylethan-1-amine in Step 2. The title compound showed identical ^1H NMR and LCMS data as Example 20A/20B.

[00531] To demonstrate the chirality, samples of Example 20A/20B (e.g., (*S*)-enantiomer) and Example 20C (e.g., (*R*)-enantiomer) were injected on a Superchiral S-OZ column (0.46cm I.D. x 25 cm L) and eluted with Hexane/EtOH/MeOH/diethylamine (v/v/v/v = 80/6.6/13.4/0.05) at 0.8mL/min. The (*R*)-enantiomer showed a retention time of 12.2 min, whereas Example 20A/20B (e.g., (*S*)-enantiomer) showed a retention time of 13.2 min.


Example 21: (S)-3-(2-amino-6-((1-hydroxyheptan-3-yl)amino)-5-methoxypyrimidin-4-yl)propanoic acid (Compound 21)



[00532]

The title compound can be prepared according to the steps of Example 15.

Example 22: (S)-5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-6-methyl-N⁴-(1-(methylthio)heptan-3-yl)pyrimidine-2,4-diamine (Compound 22)



Step 1: methyl (S)-3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxybenzoate

[00533] A mixture of methyl 3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate (1.0 eq, from Example 3B – Step 4) in NMP (0.8M), (S)-1-(methylthio)heptan-3-amine (1.0 eq; prepared by following procedures reported in WO2014/128189, pg 8, compound D) and DIEA (3.0 eq) was stirred for 48 h at 120 °C under N₂ atmosphere. The reaction was allowed to cool, diluted with EtOAc, washed with water and brine. The organic layer was dried over Na₂SO₄, concentrated and purified by column chromatography on silica gel (eluent 0-5% MeOH in DCM) to give the title compound.

Step 2: (S)-3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxyphenylmethanol

[00534] To a stirred solution of methyl (S)-3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)-pyrimidin-5-yl)methyl)-4-methoxybenzoate (1.0 eq) in THF (0.18M) was added dropwise of 2.5M LiAlH₄ (2.0 eq) at 0 °C. The reaction was stirred at r.t. for 1 h before it was quenched with EA and 1N NaOH. The mixture was filtered, and the filtrate was partitioned between EA/water. The organic layer was dried over Na₂SO₄,

concentrated and purified by column chromatography on silica gel (eluent 0-5% MeOH in DCM) to give the title compound.

Step 3: (S)-5-(5-(chloromethyl)-2-methoxybenzyl)-6-methyl-N⁴-(1-(methylthio)heptan-3-yl)pyrimidine-2,4-diamine

[00535] To a stirred solution of (S)-3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)-pyrimidin-5-yl)methyl)-4-methoxyphenyl)methanol (1.0 eq) in DCM (0.1M) was added SOCl₂ (5.0 eq) at rt. The resulting mixture was stirred at rt for 2 h before it was quenched with sat. NaHCO₃. The mixture was partitioned between DCM/water. The organic layer was dried over Na₂SO₄ and concentrated to give the title compound.

Step 4: (S)-2-(3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxyphenyl)acetonitrile

[00536] To a stirred solution of (S)-5-(5-(chloromethyl)-2-methoxybenzyl)-6-methyl-N⁴-(1-(methylthio)-heptan-3-yl)pyrimidine-2,4-diamine (1.0 eq) in 1:1 DMSO/DMF (0.1M) was added NaCN (3.0 eq). The mixture was stirred at rt for 16 h before it was quenched with sat. NaHCO₃. The mixture was partitioned between EA/water. The organic layer was dried over Na₂SO₄, concentrated and purified to give the title compound.


Step 5: (S)-5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-6-methyl-N⁴-(1-(methylthio)-heptan-3-yl)pyrimidine-2,4-diamine

[00537] A mixture of (S)-2-(3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxyphenyl)acetonitrile (1.0 eq) in dioxane (0.15M) was added TMSN₃ (10.0 eq), Bu₂SnO (2.0 eq), and stirred overnight at 120 °C. The mixture was concentrated and purified by prep-HPLC (mobile phase CH₃CN/H₂O/ HCOOH), freeze-dried to give the title compound as a white powder (formic acid salt).

[00538] LCMS: [M+H]⁺=471.3

[00539] ¹H NMR (400 MHz, DMSO) δ 7.03 (d, *J* = 8.4 Hz, 1H), 6.92 (d, *J* = 8.4 Hz, 1H), 6.79 (s, 1H), 6.44 (s, 2H), 6.24 (d, *J* = 8.8 Hz, 1H), 4.25-4.22 (m, 1H), 4.00 (s, 2H), 3.82 (s, 3H), 3.64 (s, 2H), 2.29-2.27 (m, 2H), 2.08 (s, 3H), 1.95 (s, 3H), 1.68-1.64 (m, 2H), 1.44-1.34 (m, 2H), 1.19-1.16 (m, 2H), 1.12-1.02 (m, 2H), 0.78 (t, *J* = 7.2 Hz, 3H).

Example 23: (S)-5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-6-methyl-N⁴-(1-(methylsulfonyl)heptan-3-yl)pyrimidine-2,4-diamine (Compound 23)



(S)-5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-6-methyl-N⁴-(1-(methylthio)heptan-3-yl)pyrimidine-2,4-diamine


(S)-5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-6-methyl-N⁴-(1-(methylsulfonyl)heptan-3-yl)pyrimidine-2,4-diamine
Chemical Formula: C₂₃H₃₄N₈O₃S
Exact Mass: 502.25

[00540] To a stirred solution of (S)-5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-6-methyl-N⁴-(1-(methylthio)heptan-3-yl)pyrimidine-2,4-diamine (1.0 eq, Example 22) in acetone (0.02M) was added oxone (2.0 eq) at r.t. The resulting mixture was stirred at r.t. for 3 h before it was filtered and purified by prep-HPLC (mobile phase CH₃CN/H₂O/HCOOH), freeze-dried to give the title compound as a white powder (formic acid salt).

[00541] LCMS: [M+H]⁺=503.3

[00542] ¹H NMR (400 MHz, DMSO) δ 7.05 (d, *J* = 8.4 Hz, 1H), 6.93 (d, *J* = 8.4 Hz, 1H), 6.76 (s, 1H), 6.47 (s, 2H), 6.38 (d, *J* = 6.4 Hz, 1H), 4.25-4.23 (m, 1H), 4.04 (s, 2H), 3.82 (s, 3H), 3.66 (s, 2H), 3.01-2.85 (m, 5H), 2.05 (s, 3H), 1.92-1.90 (m, 1H), 1.81-1.80 (m, 1H), 1.49-1.34 (m, 2H), 1.18-1.15 (m, 2H), 1.11-0.99 (m, 2H), 0.77 (t, *J* = 7.2 Hz, 3H).

Example 24: (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)boronic acid (Compound 24)



Step 1: *N*⁴-butyl-5-(2-methoxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzyl)-6-methylpyrimidine-2,4-diamine

[00543] A mixture of 5-(5-bromo-2-methoxybenzyl)-*N*⁴-butyl-6-methylpyrimidine-2,4-diamine (1.0 eq, from Example 11 – Step 5) in dioxane (0.03M), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (1.2 eq), KOAc (3.0 eq), and Pd(dppf)Cl₂ (0.1 eq) was stirred at 90 °C for 4 h. Then the mixture was filtered and the filtrate was concentrated under reduced pressure to give the crude product, which was purified by column chromatography on silica gel (eluent: 0-5% MeOH in DCM) to give the title compound as a white solid.


Step 2: (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-boronic acid

[00544] To a mixture of *N*⁴-butyl-5-(2-methoxy-5-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzyl)-6-methylpyrimidine-2,4-diamine (1.0 eq) in MeOH (0.05M) was added ^tBuB(OH)₂ (2.0 eq) and 6N HCl (1/10th of total reaction volume). The mixture was stirred at rt for 16 h under nitrogen atmosphere. The mixture was concentrated under reduced pressure and purified by prep-HPLC (mobile phase: FA/H₂O/CH₃CN) to give the title compound as a solid (formic acid salt).

[00545] LCMS: [M+H]⁺=345.3

[00546] ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.76 (s, 2H), 7.66 (d, *J* = 7.6 Hz, 1H), 7.25 (s, 1H), 6.96 (d, *J* = 8.0 Hz, 1H), 6.76 (br s, 1H), 6.56 (br s, 2H), 3.86 (s, 3H), 3.63 (s, 2H), 3.30-3.28 (m, 2H), 2.06 (s, 3H), 1.44-1.41 (m, 2H), 1.18-1.15 (m, 2H), 0.82 (t, *J* = 7.2 Hz, 3H).

Example 25: 4-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzamido)butanoic acid (Compound 25)



Step 1: methyl 4-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxy-benzamido)butanoate

[00547] A mixture of 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoic acid (1.0 eq, from Example 16 – Step 1) in DMF (0.06M), methyl 4-aminobutanoate (1.0 eq), HATU (1.5 eq) and DIEA (4.0 eq) was stirred at 50 °C for 5 h. The mixture was cooled to rt and water was added. The aqueous layer was extracted with DCM. The combined organic layers were dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified by column chromatography on silica gel (eluent: DCM/MeOH = 100:1 to 10:1) to give the title compound as a brown solid.


Step 2: 4-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxy-benzamido)butanoic acid

[00548] To a stirred solution of methyl 4-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzamido)butanoate (1.0 eq) in MeOH (0.025M) was added 1N NaOH (6 eq). The resulting mixture was stirred at 50 °C for 2 h. The mixture was cooled to rt and adjusted to pH 7 using 1N HCl. The precipitated solids were collected by filtration and triturated with water, dried under vacuum to give the title compound as a white solid.

[00549] LCMS: [M+H]⁺ = 430.3

[00550] ¹H NMR (400 MHz, DMSO) δ 8.28 (br s, 1H), 7.70 (d, *J* = 9.2 Hz, 1H), 7.29 (s, 1H), 7.02 (d, *J* = 8.8 Hz, 1H), 5.99 (br s, 1H), 5.81-5.79 (m, 2H), 3.89 (s, 3H), 3.62 (s, 2H), 2.36-3.16 (m, 4H), 2.21 (t, *J* = 7.2 Hz, 2H), 1.98 (s, 3H), 1.72-1.67 (m, 2H), 1.42-1.38 (m, 2H), 1.20-1.14 (m, 2H), 0.81 (t, *J* = 7.6 Hz, 3H).

Example 26: 5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-6-methyl-N⁴-pentylpyrimidine-2,4-diamine (Compound 26)



5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-6-methyl-N⁴-pentylpyrimidine-2,4-diamine
Chemical Formula: C₂₀H₂₈N₈O
Exact Mass: 396.24


[00551] The title compound was prepared according to the procedures described in Example 3, but using pentan-1-amine instead of butan-1-amine in Step 5. The final product

was purified by prep-HPLC (mobile phase CH₃CN/H₂O/HCOOH), freeze-dried to give the title compound as a white solid (formic acid salt).

[00552] LC-MS: [M+H]⁺ = 397.2

[00553] ¹H NMR (400 MHz, DMSO) δ 7.03 (d, J = 8.8 Hz, 1H), 6.91 (d, J = 8.0 Hz, 1H), 6.74 (s, 1H), 6.64 (br s, 1H), 6.48 (br s, 2H), 3.97 (s, 2H), 3.81 (s, 3H), 3.61 (s, 2H), 3.30-3.24 (m, 2H), 2.04 (s, 3H), 1.46-1.40 (m, 2H), 1.29-1.21 (m, 2H), 1.20-1.39 (m, 2H), 0.83 (t, J = 6.8 Hz, 3H).

Example 27: (S)-3-(2-amino-6-((1-hydroxyheptan-3-yl)amino)-5-(2-methoxybenzyl)-pyrimidin-4-yl)propanoic acid (Compound 27)



Step 1: diethyl 3-oxohexanedioate

[00554] To a solution of 4-ethoxy-4-oxobutanoic acid (1.0 eq) in anhydrous THF (0.5M) at r.t. was added carbonyl di-imidazole (1.2 eq) under N₂. The resulting mixture was stirred at r.t. for 1 h, then MgCl₂ (1.0eq) and potassium 3-ethoxy-3-oxopropanoate (1.0 eq) were added into the above solution at r.t. and heated at 60°C for 2h. The mixture was filtered and the filtration was concentrated under reduced pressure to give the crude product, which was purified by column chromatography on silica (eluent: PE/EA = 100:1 to 3:1) to give the title compound as a yellow oil.

Step 2: diethyl 2-(2-methoxybenzyl)-3-oxohexanedioate

[00555] To a solution of diethyl 3-oxohexanedioate (1.2 eq) in anhydrous THF (1.2M) was added 60% NaH (1.3 eq) in portions at 0 °C. After stirring for 10 min, a solution of 1-(bromomethyl)-2-methoxybenzene (1.0 eq; from Example 15 – Step 1) in THF (1M) was added dropwise into above mixture. The resulting mixture was stirred at 60 °C for 16 h and quenched with water. The mixture was extracted with EA, and the combined organic layers were dried over Na₂SO₄, concentrated under reduced pressure. The residue was purified by flash chromatography on silica (eluent: PE/EA = 50:1 to 10:1) to give the title compound as a yellow oil.

Step 3: 3-(2-amino-6-hydroxy-5-(2-methoxybenzyl)pyrimidin-4-yl)propanoic acid

[00556] A mixture of diethyl 2-(2-methoxybenzyl)-3-oxohexanedioate (1.0 eq) and guanidine carbonate (1.0 eq) in MeOH (0.3M) was stirred at 90 °C for 16 h. The precipitate was collected by filtration, washed with water, and dried in vacuum to give the title compound as a white solid.

Step 4: 3-(2-amino-6-chloro-5-(2-methoxybenzyl)pyrimidin-4-yl)propanoic acid

[00557] A solution of 3-(2-amino-6-hydroxy-5-(2-methoxybenzyl)pyrimidin-4-yl)propanoic acid (1.0 eq) in POCl₃ (0.26M) was stirred at 100 °C for 2 h under nitrogen. The reaction was cooled to rt and POCl₃ was evaporated under reduced pressure. The residue was diluted with water and the pH was adjusted to 7 by adding solid NaHCO₃. The mixture was stirred at 50 °C for 1 h, then cooled to rt, and the precipitate was collected by filtration. The filter cake was washed with water, dried in vacuum to give the title compound as a white solid.


Step 5: (S)-3-(2-amino-6-((1-hydroxyheptan-3-yl)amino)-5-(2-methoxybenzyl)pyrimidin-4-yl)propanoic acid

[00558] A solution of 3-(2-amino-6-chloro-5-(2-methoxybenzyl)pyrimidin-4-yl)propanoic acid (1.0 eq) in neat (S)-3-aminoheptan-1-ol (1.6M; prepared from *J. Med. Chem.* **2016**, 59, 7936–7949) was stirred at 120 °C for 2 h. The mixture was diluted with water, and the aqueous phase was extracted with EA. The combined organic layers were washed with 1N HCl, brine, dried over Na₂SO₄, and concentrated. The crude was purified by prep-HPLC (mobile phase: NH₄OH/MeCN/H₂O) to give the title compound white solid.

[00559] LCMS: [M+H]⁺ = 417.3

[00560] ^1H NMR (400 MHz, CD_3OD) δ 7.24 (t, $J = 8.0$ Hz, 1H), 7.02-6.96 (m, 2H), 6.89 (t, $J = 7.2$ Hz, 1H), 4.42-4.40 (m, 1H), 3.92 (s, 3H), 3.82 (s, 2H), 3.46-3.38 (m, 2H), 2.95-2.90 (m, 2H), 2.48 (t, $J = 7.2$ Hz, 2H), 1.8-1.0 (m, 8H), 0.80 (t, $J = 7.2$ Hz, 3H).

Example 28: *N*-(3-(2*H*-tetrazol-5-yl)propyl)-3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzamide (Compound 28)



Step 1: 4-(1,3-dioxoisindolin-2-yl)butanenitrile

[00561] A mixture of 4-bromobutanenitrile (1.5 eq) and 1,3-dioxoisindolin-2-one (1.0 eq) in DMF (0.3M) was stirred at 100 °C for 8 h. The mixture was filtered and the filtrate was concentrated under reduced pressure. The residue was purified by flash chromatography on silica (eluent: PE/EA = 100:1 to 1:1) to give the title compound as a white solid.

Step 2: 4-aminobutanenitrile

[00562] A solution of 4-(1,3-dioxoisindolin-2-yl)butanenitrile (1.0 eq) in EtOH (0.26M) was added $\text{NH}_2\text{NH}_2 \cdot \text{H}_2\text{O}$ (2.0 eq) and stirred at rt for 16 h. The mixture was concentrated, then diluted with water. The aqueous phase was extracted with DCM. The combined organic phase was dried over Na_2SO_4 , concentrated under reduced pressure to give the title compound as a brown oil.

Step 3: 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-*N*-(3-cyanopropyl)-4-methoxybenzamide

[00563] A mixture of 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoic acid (1.0 eq; from Example 16 – Step 1) in DMF (0.09M), 4-aminobutanenitrile (2.0 eq), HATU (2.0 eq) DIEA (3.0 eq) was stirred at 50 °C for 5 h. The

mixture was cooled to rt, then water was added and the aqueous phase was extracted with DCM. The combined organic phase was dried over Na_2SO_4 , concentrated under reduced pressure. The residue was purified by flash chromatography on silica (PE/EA=50:1 to 1:2) to give the title compound as a brown solid.


Step 4: *N*-(3-(2*H*-tetrazol-5-yl)propyl)-3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzamide

[00564] To a stirred solution of 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-*N*-(3-cyanopropyl)-4-methoxybenzamide (1.0 eq) in NMP (0.08M) was added Bu_2SnO (2.0 eq) and TMSN_3 (10.0 eq). The resulting mixture was stirred overnight at 125 °C under N_2 . The mixture was cooled to rt, concentrated and purified by prep-HPLC (mobile phase: $\text{CH}_3\text{CN}/\text{H}_2\text{O}/0.1\%\text{NH}_4\text{OH}$) to give the title compound as a solid.

[00565] LC-MS: $[\text{M}+\text{H}]^+ = 454.5$

[00566] ^1H NMR (400 MHz, DMSO) δ 8.45 (s, 1H), 7.74 (d, $J = 8.0$ Hz, 1H), 7.32 (s, 1H), 7.04 (d, $J = 8.8$ Hz, 1H), 6.57 (br s, 1H), 6.30-6.20 (m, 2H), 3.90 (s, 3H), 3.65 (s, 2H), 3.30-3.26 (m, 4H), 2.82 (t, $J = 7.2$ Hz, 2H), 2.03 (s, 3H), 1.90 (t, $J = 6.8$ Hz, 2H), 1.46-1.38 (m, 2H), 1.23-1.12 (m, 2H), 0.80 (t, $J = 7.2$ Hz, 3H).

Example 29: (*S*)-3-((2-amino-6-(2-carboxyethyl)-5-(2-methoxybenzyl)pyrimidin-4-yl)amino)heptanoic acid (Compound 29)



Step 1: *tert*-butyl (*S,E*)-3-((2-amino-6-(3-ethoxy-3-oxoprop-1-en-1-yl)-5-(2-methoxybenzyl)-pyrimidin-4-yl)amino)heptanoate

[00567] A solution of ethyl (*E*)-3-(2-amino-6-chloro-5-(2-methoxybenzyl)pyrimidin-4-yl)acrylate (1.0 eq, from Example 15 – Step 5) in neat *tert*-butyl (*S*)-3-aminoheptanoate (0.6M, from Example 20B – Step 3) was stirred at 120 °C for 2 h. The reaction solution was diluted with water, and the aqueous layer was extracted with EA. Combined organic layers were washed with water, brine, dried over Na₂SO₄, and concentrated. The crude was purified by column chromatography (DCM/MeOH=10:1) to give the title compound as a yellow solid.

Step 2: *tert*-butyl (*S*)-3-((2-amino-6-(3-ethoxy-3-oxopropyl)-5-(2-methoxybenzyl)pyrimidin-4-yl)amino)heptanoate

[00568] To a solution of *tert*-butyl (*S,E*)-3-((2-amino-6-(3-ethoxy-3-oxoprop-1-en-1-yl)-5-(2-methoxy-benzyl)-pyrimidin-4-yl)amino)heptanoate (1.0 eq) in MeOH (0.04M) was added Pd/C (1/3 weight equivalent). The mixture was stirred under H₂ atmosphere at room temperature for 2 h. The Pd/C was filtered off, and the filtrate was concentrated to give the title compound as a yellow solid.


Step 3: (*S*)-3-((2-amino-6-(2-carboxyethyl)-5-(2-methoxybenzyl)pyrimidin-4-yl)amino)-heptanoic acid

[00569] To a solution of *tert*-butyl (*S*)-3-((2-amino-6-(3-ethoxy-3-oxopropyl)-5-(2-methoxybenzyl)-pyrimidin-4-yl)amino)heptanoate (1.0eq) in 4:1 EtOH/H₂O (0.06M) was added KOH (5.0 eq). The solution was stirred at 100°C for 3 h. pH was adjusted to 7 using acetic acid, and the solution was concentrated and then purified by prep-HPLC (mobile phase: NH₄OH/MeCN/ H₂O) to give the title compound as a white solid.

[00570] LC-MS: [M+H]⁺=431.4

[00571] ¹H NMR (400 MHz, CD₃OD) δ 7.23 (t, *J* = 7.2 Hz, 1H), 7.01-6.97 (m, 2H), 6.89 (t, *J* = 7.2 Hz, 1H), 4.61 (m, 1H), 3.91 (s, 3H), 3.82 (s, 2H), 2.94-2.92 (m, 2H), 2.47-2.41 (m, 2H), 1.6-0.9 (m, 8H), 0.80 (t, *J* = 7.2 Hz, 3H).

Example 30: (3-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzamido)propyl)phosphonic acid (Compound 30)



Step 1: diethyl (3-(1,3-dioxoisindolin-2-yl)propyl)phosphonate

[00572] A mixture of diethyl (3-bromopropyl)phosphonate (1.2 eq) and 1,3-dioxoisindolin-2-ide (1.0 eq) in DMF (0.2M) was stirred at 100 °C for 8 h. The reaction was filtered, and the filtrate was concentrated under reduced pressure to give the title compound as a white solid.

Step 2: diethyl (3-aminopropyl)phosphonate

[00573] A mixture of diethyl (3-(1,3-dioxoisindolin-2-yl)propyl)phosphonate (1.0 eq) in EtOH (0.25M) and $NH_2NH_2 \cdot H_2O$ (2.0 eq) was stirred at rt for 16 h. The reaction was filtered, the filtrate was poured in water, and the aqueous phase was extracted with DCM. The organic phase was dried over Na_2SO_4 , concentrated under reduced pressure to give the title compound as a brown oil.

Step 3: diethyl (3-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxy-benzamido)propyl)phosphonate

[00574] A mixture of 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoic acid (1.0 eq; from Example 16 – Step 1) in DMF (0.1M), diethyl (3-aminopropyl)phosphonate (2.0 eq), HATU (2.0 eq) and DIEA (3.0 eq) was stirred at 50 °C for 5 h under nitrogen. The mixture was cooled to rt, diluted with water, and the aqueous phase was extracted with DCM. The organic phase was dried over Na_2SO_4 , concentrated under reduced pressure to give the title compound as a brown solid.


Step 4: (3-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzamido)propyl)phosphonic acid

[00575] A solution of diethyl (3-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzamido)propyl)phosphonate (1.0 eq) in DCM (0.08M) and TMSBr (10.0 eq) was stirred overnight at rt under nitrogen. The mixture was concentrated, purified by prep-HPLC (mobile phase CH₃CN/H₂O/0.1%NH₄OH) to give the title compound as a solid.

[00576] LC-MS: [M+H]⁺ = 466.4

[00577] ¹H NMR (400 MHz, DMSO) δ 8.59 (s, 1H), 7.76 (d, *J* = 8.0 Hz, 1H), 7.27 (s, 1H), 7.20 (br s, 1H), 7.03 (d, *J* = 8.4 Hz, 1H), 3.88 (s, 3H), 3.64 (s, 2H), 3.35-3.28 (m, 2H), 3.25-3.16 (m, 2H), 2.07 (s, 3H), 1.74-1.62 (m, 2H), 1.48-1.33 (m, 4H), 1.22-1.11 (m, 2H), 0.81 (t, *J* = 7.2 Hz, 3H).

Example 31: 5-((5-((2*H*-tetrazol-5-yl)methyl)-2-methoxyphenyl)thio)-*N*⁴-butyl-6-methylpyrimidine-2,4-diamine (Compound 31)



Step 1: methyl 3-((dimethylcarbamothioyl)oxy)-4-methoxybenzoate

[00578] To a stirred solution of methyl 3-hydroxy-4-methoxybenzoate (1.0 eq) in DMF (0.55M) was added NaH (1.1 eq) at 0 °C. The mixture was warmed to rt and stirred for 30 min before treated with dimethylcarbamothioic chloride (1.1 eq). The mixture was stirred overnight and partitioned between ether/water. The resulting suspension was filtered. The solid was collected and dried under vacuum to give the title compound as a white solid.

Step 2: methyl 3-((dimethylcarbamoyl)thio)-4-methoxybenzoate

[00579] Methyl 3-((dimethylcarbamothioyl)oxy)-4-methoxybenzoate (1.0 eq) in PhOPh (1M) was stirred for 16 h at 260 °C. The mixture was cooled to rt and the mixture was purified by flash chromatography on silica (eluent PE/EA = 100:1-2:1) to give the title compound.

Step 3: methyl 3-mercapto-4-methoxybenzoate

[00580] To a stirred solution of methyl 3-((dimethylcarbamoyl)thio)-4-methoxybenzoate (1.0 eq) in THF (0.3M) was added MeONa (2.0 eq) at rt. The resulting mixture was stirred at 60 °C for 2 h. The mixture was cooled down to rt and quenched with 1N HCl. The mixture was partitioned between EA/water. The organic phase was dried over Na₂SO₄ and concentrated and purified by flash chromatography on silica (eluent PE/EA = 100:1~5:1) to give the title compound.

Step 4: methyl 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)thio)-4-methoxybenzoate

[00581] A mixture of methyl 3-mercapto-4-methoxybenzoate (1.0 eq) in dioxane (0.3M), *N*^t-butyl-5-iodo-6-methylpyrimidine-2,4-diamine (1.5 eq), Cs₂CO₃ (2.0 eq), CuI (2.0 eq) was stirred overnight at 100 °C. The mixture was partitioned between EA/NH₄OH. The organic layer was washed with brine, dried over Na₂SO₄, concentrated and purified by flash chromatography on silica (eluent PE/EA = 100:1~1:1) to give the title compound.

Step 5: (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)thio)-4-methoxyphenyl)-methanol

[00582] To a stirred solution of methyl 3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)thio)-4-methoxybenzoate (1.0 eq) in THF (0.2M) was added 1M LiAlH₄ (2.0 eq) dropwise at 0 °C. The resulting mixture was stirred at 0 °C for 10 min and at rt for 1 h. The mixture was diluted with EA and quenched with 2N NaOH. The mixture was

partitioned between EA/water. The organic layer was dried over Na₂SO₄, filtered and concentrated to give the title compound.

Step 6: *N*⁴-butyl-5-((5-(chloromethyl)-2-methoxyphenyl)thio)-6-methylpyrimidine-2,4-diamine

[00583] To a stirred solution of (3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)thio)-4-methoxyphenyl)methanol (1.0 eq) in DCM (0.2M) was added SOCl₂ (2.0 eq) at rt under N₂. The mixture was stirred at rt for 1 h. The mixture was partitioned between DCM and saturated aqueous NaHCO₃. The organic layer was dried over Na₂SO₄, filtered and concentrated to give the title compound.

Step 7: 2-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)thio)-4-methoxyphenyl)acetonitrile

[00584] A mixture of *N*⁴-butyl-5-((5-(chloromethyl)-2-methoxyphenyl)thio)-6-methylpyrimidine-2,4-diamine (1.0 eq) in 1:1 DMSO/DMF (0.1M) and NaCN (3.0 eq) was stirred overnight at rt. The mixture was partitioned between EA/water. The organic layer was dried over Na₂SO₄, filtered, concentrated and purified by flash chromatography on silica (eluent 0-5% MeOH in DCM) to give the title compound.


Step 8: 5-((5-((2*H*-tetrazol-5-yl)methyl)-2-methoxyphenyl)thio)-*N*⁴-butyl-6-methylpyrimidine-2,4-diamine

[00585] A mixture of 2-(3-((2-amino-4-(butylamino)-6-methylpyrimidin-5-yl)thio)-4-methoxyphenyl)acetonitrile (1.0 eq) in NMP (0.1M), Bu₂SnO (2.0 eq), and TMSN₃ (10.0 eq) was stirred for 3 h at 120 °C under N₂. The mixture was cooled down to rt and the mixture was concentrated and purified by prep-HPLC (mobile phase CH₃CN/H₂O/NH₄OH), freeze-dried to give the title compound as a white powder.

[00586] LCMS: [M+H]⁺ = 401.4

[00587] ¹H NMR (400 MHz, DMSO) δ 7.03-6.92 (m, 2H), 6.46-6.42 (m, 2H), 6.39 (s, 2H), 4.06 (s, 2H), 3.83 (s, 3H), 3.26-3.21 (m, 2H), 2.14 (s, 3H), 1.39-1.32 (m, 2H), 1.17-1.11 (m, 2H), 0.81 (t, *J* = 7.2 Hz, 3H).

Example 32: (S)-5-(5-((2*H*-tetrazol-5-yl)methyl)-2-methoxybenzyl)-*N*⁴-(1-methoxyheptan-3-yl)-6-methylpyrimidine-2,4-diamine (Compound 32)



Step 1: (3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)methanol

[00588] To a solution of methyl 3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate (1.0 eq, from Example 3B – Step 4) in THF (1.6M) was added LiAlH₄ (3.0 eq) at r.t. The reaction was stirred at r.t. for 2 h. The reaction solution was diluted with water. The aqueous layer was extracted with DCM. The organic layer was washed with water, brine, separated, dried over Na₂SO₄, and concentrated to give the title compound as a pale solid.

Step 2: (S)-3-((2-amino-4-((1-methoxyheptan-3-yl)amino)-6-methylpyrimidin-5-yl)methyl)-4-methoxy-phenyl)methanol

[00589] A mixture of (3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)methanol (1.0 eq) in NMP (0.4M) and (S)-1-methoxyheptan-3-amine (2.0 eq) was stirred at 120 °C for 16 h under nitrogen. The reaction solution was diluted with water. The aqueous phase was extracted with EA. The organic layer was washed with water, brine, dried over Na₂SO₄, and concentrated. The crude was purified by column chromatography (DCM/MeOH=50:1) to give the title compound as a yellow solid.

Step 3: (*S*)-5-(5-(chloromethyl)-2-methoxybenzyl)-*N*⁴-(1-methoxyheptan-3-yl)-6-methylpyrimidine-2,4-diamine

[00590] To a stirred solution of (*S*)-3-((2-amino-4-((1-methoxyheptan-3-yl)amino)-6-methylpyrimidin-5-yl)methyl)-4-methoxy-phenyl)methanol (1.0 eq) in DCM (0.3M) was added SOCl₂ (2.0 eq) dropwise. The resulting mixture was stirred at 20 °C for 1 h, then quenched with H₂O. The mixture was partitioned between DCM/water. The organic layer was separated, dried over Na₂SO₄, filtered, and concentrated to give the title compound as a brown oil.

Step 4: (*S*)-2-(3-((2-amino-4-((1-methoxyheptan-3-yl)amino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)acetonitrile

[00591] A mixture of (*S*)-5-(5-(chloromethyl)-2-methoxybenzyl)-*N*⁴-(1-methoxyheptan-3-yl)-6-methylpyrimidine-2,4-diamine (1.0 eq) in DMSO (0.4M) and KCN (3.0 eq) was stirred at 80 °C for 4 h, then cooled to r.t. Water was added, and the solution was extracted with DCM. The organic phase was dried over Na₂SO₄, concentrated, and purified by column chromatography (DCM/MeOH=40:1) to give the title compound as a white solid.


Step 5: (*S*)-5-(5-((2*H*-tetrazol-5-yl)methyl)-2-methoxybenzyl)-*N*⁴-(1-methoxyheptan-3-yl)-6-methylpyrimidine-2,4-diamine

[00592] To a stirred mixture of (*S*)-2-(3-((2-amino-4-((1-methoxyheptan-3-yl)amino)-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)acetonitrile (1.0 eq) in dioxane (0.2M) and Bu₂SnO (2.0 eq) was added TMSN₃ (10.0 eq). The resulting mixture was stirred at 110 °C for 4 h under N₂, then cooled down to r.t. The mixture was concentrated and purified by prep-HPLC (mobile phase 0.1%NH₃·H₂O/CH₃CN) to give the title compound as a solid.

[00593] LC-MS: [M+H]⁺ = 455.4

[00594] ¹H NMR (400 MHz, CDCl₃) δ 6.86 (s, 1H), 6.84 (d, *J* = 8.4 Hz, 1H), 6.50 (d, *J* = 8.4 Hz, 1H), 4.28-4.23 (m, 3H), 3.90 (s, 3H), 3.58 (s, 2H), 3.19-3.16 (m, 2H), 3.14 (s, 3H), 2.50 (s, 3H), 1.83-1.81 (m, 1H), 1.59-1.06 (m, 7H), 0.83 (t, *J* = 7.2 Hz, 3H).

Example 33: (S)-2-(3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanoic acid (Compound 33)



Step 1: (S)-2-(3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanenitrile

[00595] A mixture of 2-(3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanenitrile (1.0 eq, from Example 9 – Step 8) in NMP (0.6M), (S)-1-(methylthio)heptan-3-amine (1.5 eq; prepared by following procedures reported in WO2014/128189, pg 8, compound D) and DIEA (3.0 eq) was stirred at 120 °C for 48 h. Water was added to the mixture, and it was extracted with EtOAc. The organic layer was dried over Na₂SO₄ and concentrated under reduced pressure. The residue was purified by flash chromatography on silica (eluent: DCM/MeOH = 100:1~20:1) to give the title compound as a brown oil.


Step 2: (S)-2-(3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanoic acid

[00596] To a stirred solution of (S)-2-(3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanenitrile (1.0 eq) in 1:1 ethane-1,2-diol/H₂O (0.1M) was added KOH (10eq) and heated in a sealed tube at 150 °C for 16 h. The mixture was acidified by 1N HCl solution to pH = 2-3 and filtered. The filtrate was purified by prep-HPLC (mobile phase CH₃CN/H₂O/NH₃·H₂O), then freeze-dried to give the title compound as a white solid.

[00597] LC-MS: [M+H]⁺ = 475.3

[00598] ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.14 (dd, *J* = 8.4, 2.0 Hz, 1H), 6.91 (d, *J* = 8.8 Hz, 1H), 6.86 (s, 1H), 5.91 (br s, 2H), 5.72 (d, *J* = 9.2 Hz, 1H), 4.25-4.15 (m, 1H), 3.82 (s, 3H), 3.62 (s, 2H), 2.31-2.25 (m, 2H), 2.02 (s, 3H), 1.95 (s, 3H), 1.70-1.52 (m, 2H), 1.44-1.34 (m, 2H), 1.31 (s, 6H), 1.25-1.05 (m, 4H), 0.78 (t, *J* = 6.8 Hz, 3H).

Example 34: (S)-5-(2-methoxy-5-(2H-tetrazol-5-yl)benzyl)-6-methyl-N⁴-(1-(methylthio)heptan-3-yl)pyrimidine-2,4-diamine (Compound 34)



Step 1: 4-methoxy-3-methylbenzonitrile

[00599] A mixture of 4-bromo-1-methoxy-2-methylbenzene (1.0 eq) in DMF (1M), Zn(CN)₂ (1.5 eq), Pd₂(dba)₃ (0.02 eq) and S-phos (0.05 eq) was stirred at 80 °C for 16 h. The resulting mixture was concentrated and purified by column chromatography on silica gel (eluent PE/EA = 100:1~20:1) to give the title compound.

Step 2: 3-(bromomethyl)-4-methoxybenzonitrile

[00600] A mixture of 4-methoxy-3-methylbenzonitrile (1.0 eq) in CCl₄ (0.7M), NBS (1.1 eq) and AIBN (0.1 eq) was stirred at 95 °C for 16 h under nitrogen. The resulting mixture was concentrated and partitioned between DCM and water. The organic layer was dried over Na₂SO₄, concentrated, and purified by column chromatography on a silica gel (eluent PE/EA = 100:1~20:1) to give the title compound as a white solid.

Step 3: ethyl 2-(5-cyano-2-methoxybenzyl)-3-oxobutanoate

[00601] To a solution of 3-(bromomethyl)-4-methoxybenzonitrile (1.5 eq) in THF (0.6M) at 0 °C was added portion-wise 60% NaH (1.5 eq) under N₂. The resulting suspension was stirred at 0 °C for 10 min, then a solution of ethyl 3-oxobutanoate (1.0 eq) in THF (1M) was added drop-wise over 10 min at 0 °C. The resulting mixture was stirred at 70 °C for 15 h.

The reaction was cooled to r.t., and ice water was added. The mixture was extracted with ethyl acetate. The organic layer was dried over Na₂SO₄, concentrated and purified by column chromatography on silica gel (eluent PE/EA = 50:1~5:1) to give the title compound as yellow oil.

Step 4: 3-((2-amino-4-hydroxy-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzonitrile

[00602] To a solution of ethyl 2-(5-cyano-2-methoxybenzyl)-3-oxobutanoate (1.0 eq) in MeOH (0.7M) was added guanidine carbonate (1.0 eq). The resulting mixture was stirred at 70 °C for 16 h. The mixture was concentrated, and the residue was suspended in EtOAc and then filtered. The filter cake was washed with water and EtOAc. The resulting solid was collected and dried under vacuum to give the title compound as a white solid.

Step 5: 3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzonitrile

[00603] The suspension of 3-((2-amino-4-hydroxy-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzonitrile (1.0 eq) in POCl₃ (0.3M) was stirred at 100 °C for 16 h. The reaction mixture was cooled to r.t., and solvent was evaporated under reduced pressure. The residue was diluted with water, and pH was adjusted to 7 using solid NaHCO₃. The precipitate was collected by filtration, and the filter cake was washed with water and EtOAc, dried under vacuum to give the title compound.

Step 6: (S)-3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxybenzonitrile

[00604] A mixture of 3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzonitrile (1.0 eq) in NMP (0.3M) and (S)-1-(methylthio)heptan-3-amine (1.5 eq, prepared by following procedures reported in WO2014/128189, pg 8, compound D) was stirred at 150 °C for 4 h under nitrogen. The reaction solution was diluted with water, and the aqueous phase was extracted with EA. The organic layer was washed with water, brine, dried over Na₂SO₄, and concentrated. The crude was purified by column chromatography (DCM/MeOH=50:1) to give the title compound as a yellow solid.

Step 7: (S)-5-(2-methoxy-5-(2H-tetrazol-5-yl)benzyl)-6-methyl-N⁴-(1-(methylthio)heptan-3-yl)pyrimidine-2,4-diamine


[00605] To a stirred mixture of (S)-3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxybenzonitrile (1.0 eq) in dioxane (0.4M), Bu₂SnO (2.0 eq) was added TMSN₃ (10.0 eq). The resulting mixture was stirred at 110 °C for 4 h

under N₂. The mixture was cooled to r.t., and the mixture was concentrated and purified by prep-HPLC (mobile phase 0.1% NH₃·H₂O/CH₃CN) to give the title compound.

[00606] LC-MS: [M+H]⁺ = 457.3

[00607] ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.82 (d, *J* = 8.8 Hz, 1H), 7.50 (s, 1H), 7.07 (d, *J* = 8.4 Hz, 1H), 6.55 (br, 3H), 4.28-4.26 (m, 1H), 3.89 (s, 3H), 3.73 (s, 2H), 2.27 (t, *J* = 7.6 Hz, 2H), 2.14 (s, 3H), 1.88 (s, 3H), 1.67-1.65 (m, 2H), 1.41-1.39 (m, 2H), 1.12-1.04 (m, 4H), 0.69 (t, *J* = 6.8 Hz, 3H).

Example 35: (S)-3-((3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxybenzamido)propyl)phosphonic acid (Compound 35)



Step 1: methyl (S)-3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxybenzoate

[00608] A mixture of methyl 3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate (1.0 eq, from Example 3B – Step 4) in NMP (0.16M) and (S)-1-(methylthio)heptan-3-amine (1.5 eq, prepared by following procedures reported in WO2014/128189, pg 8, compound D) was stirred at 150 °C for 4 h under nitrogen. The mixture was diluted with water. The aqueous phase was extracted with EA. The organic layer

was washed with water, brine, dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography (DCM/MeOH=50:1) to give the title compound as a yellow solid.

Step 2: (*S*)-3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxybenzoic acid

[00609] A mixture of methyl (*S*)-3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxybenzoate (1.0 eq) in 1:1 EtOH/H₂O (0.02M) and NaOH (10.0 eq) was stirred at 70°C for 16 h. The mixture was neutralized using 1N HCl, and the resulting suspension was filtered. The filter cake was washed with water to give a solid which was dissolved in MeCN/H₂O. 4N HCl/dioxane was added, and the resulting solution was freeze-dried to give the title compound as a white solid.

Step 3: diethyl (*S*)-3-(3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxybenzamido)propyl)phosphonate

[00610] A mixture of (*S*)-3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxybenzoic acid (1.0 eq) in DMF (0.1M), diethyl (3-aminopropyl)phosphonate (1.1eq, from Example 30 – Step 2), HATU (1.5 eq) and DIEA (2.0 eq) was stirred at 30 °C for 16 h. The mixture was cooled to r.t., and water was added. The mixture was extracted with DCM. The organic phase was dried over Na₂SO₄, concentrated under reduced pressure to give the title compound as a colorless oil.


Step 4: (*S*)-3-(3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxybenzamido)propyl)phosphonic acid

[00611] A mixture of diethyl (*S*)-3-(3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxybenzamido)propyl)phosphonate (1.0 eq) in DCM (0.05M) and TSMBr (10.0 eq) was stirred at 35 °C for 5 h under N₂. The mixture was cooled and concentrated. The residue was purified by prep-HPLC (mobile phase CH₃CN/H₂O/NH₃·H₂O) to give the title compound.

[00612] LC-MS: [M+H]⁺ = 554.3

[00613] ¹H NMR (400 MHz, CDCl₃) δ 13.93 (s, 1H), 10.36 (s, 1H), 10.12 (s, 1H), 7.94 (d, *J* = 8.0 Hz, 1H), 7.80 (s, 1H), 6.96 (d, *J* = 8.8 Hz, 1H), 5.94 (d, *J* = 8.4 Hz, 1H), 4.23-4.21 (m, 1H), 3.98 (s, 3H), 3.73-3.30 (m, 4H), 2.76 (s, 3H), 2.43-2.41 (m, 1H), 2.05-1.80 (m, 7H), 1.70 (s, 3H), 1.55-1.20 (m, 6H), 0.86 (t, *J* = 4.0 Hz, 3H).

Example 36: (*S*)-N⁴-(1-(2*H*-tetrazol-5-yl)heptan-3-yl)-5-(2-methoxybenzyl)-6-methylpyrimidine-2,4-diamine (Compound 36)



Step 1: (S)-3-(benzyl((S)-1-phenylethyl)amino)heptyl methanesulfonate

[00614] To a stirred solution of (S)-3-(benzyl((S)-1-phenylethyl)amino)heptan-1-ol (1.0 eq, prepared from *J. Med. Chem.* **2016**, *59*, 7936-7949) and DIEA (1.1 eq) in DCM (0.5M) was added MsCl (1.1 eq) at 0 °C. The resulting mixture was warmed to r.t. over 3 h. Solvent was removed, and the residue was diluted with EA. The organic layer was washed with 1N HCl, 1N NaOH, brine, dried and concentrated to give the title compound.

Step 2: (S)-4-(benzyl((S)-1-phenylethyl)amino)octanenitrile

[00615] A mixture of (S)-3-(benzyl((S)-1-phenylethyl)amino)heptyl methanesulfonate (1.0 eq) in DMSO (0.3M) and KCN (7.5 eq) was stirred at 30 °C for 16 h. The reaction was diluted with water. The aqueous layer was extracted with EA. The organic layer was separated, dried and concentrated to give the title compound as a light yellow oil.

Step 3: (S)-N-benzyl-N-((S)-1-phenylethyl)-1-(2H-tetrazol-5-yl)heptan-3-amine

[00616] A mixture of (S)-4-(benzyl((S)-1-phenylethyl)amino)octanenitrile (1.0 eq) in dioxane (0.2M), TMSN₃ (2.5 eq) and Bu₂SnO (2.0 eq) was stirred at 120 °C in a seal tube for 18 h. The mixture was diluted with EA and then washed with water. The organic layer was dried and concentrated. The crude was purified by column chromatography (eluent PE: EA=10:1) to give the title compound as a light brown oil.

Step 4: (S)-1-(2H-tetrazol-5-yl)heptan-3-amine

[00617] To a solution of (S)-N-benzyl-N-((S)-1-phenylethyl)-1-(2H-tetrazol-5-yl)heptan-3-amine (1.0 eq) in MeOH (0.2M) was added 50% Pd/C (0.3 wt eq) under nitrogen.

The reaction was stirred at 40 °C under H₂ atmosphere for 16 h. Pd/C was filtered off, and the filtrate was concentrated to give the title compound as a light yellow solid.


Step 5: (S)-N⁴-(1-(2*H*-tetrazol-5-yl)heptan-3-yl)-5-(2-methoxybenzyl)-6-methylpyrimidine-2,4-diamine

[00618] A solution of 4-chloro-5-(2-methoxybenzyl)-6-methylpyrimidin-2-amine (1.0 eq, from Example 41 – Step 3) in NMP (0.4M) and (S)-1-(2*H*-tetrazol-5-yl)heptan-3-amine (2.0 eq) was stirred at 150 °C for 3 h under nitrogen. The reaction was diluted with water, and the aqueous was extracted with EA. The organic layers were combined and washed with water, brine, dried, and concentrated. The crude was purified by prep-HPLC (mobile phase 0.1% NH₃H₂O/CH₃CN) to give the title compound.

[00619] LC-MS: [M+H]⁺ = 411.4

[00620] ¹H NMR (400 MHz, MeOD) δ 7.17 (t, *J* = 7.2 Hz, 1H), 6.96 (d, *J* = 8.8 Hz, 2H), 6.85 (t, *J* = 7.2 Hz, 1H), 4.24-4.20 (m, 1H), 3.91 (s, 3H), 3.78-3.63 (m, 2H), 3.19-3.16 (m, 1H), 2.72 (t, *J* = 8.0 Hz, 2H), 2.28 (s, 3H), 2.28-0.80 (m, 8H), 0.77 (t, *J* = 7.2 Hz, 3H).

Example 37: 6-(2-(2*H*-tetrazol-5-yl)ethyl)-N⁴-butyl-5-(2-methoxybenzyl)pyrimidine-2,4-diamine (Compound 37)



Step 1: ethyl 5-cyano-3-oxopentanoate

[00621] To a solution of 3-cyanopropanoic acid (1.0 eq) in THF (0.5M) was added CDI (1.2 eq). The reaction solution was stirred at 25 °C for 1 h. MgCl₂ (1.0 eq) and potassium 3-ethoxy-3-oxopropanoate (1.0 eq) were added to the above solution and stirred at 60 °C for 2

h. The solution was cooled to room temperature and quenched by H₂O. The aqueous phase was extracted with EA. The combined organic layer was washed with water then brine. The organic layer was separated and dried over Na₂SO₄, concentrated and purified by column chromatography (PE/EA=3:1) to give the title compound as a yellow oil.

Step 2: diethyl 2-(2-methoxybenzyl)-3-oxohexanedioate

[00622] To a solution of ethyl 5-cyano-3-oxopentanoate (1.2 eq) in THF (0.3M) was added NaH (1.3 eq) at 0 °C in portions under nitrogen. The solution was stirred at 0 °C for 15 min, then 1-(bromomethyl)-2-methoxybenzene (1.0 eq) was added. The reaction was stirred at 40 °C for 16 h. The reaction mixture was quenched by water. The aqueous layer was extracted with EA. The combined organic layer was washed with water, brine, dried over Na₂SO₄, and concentrated. The crude was purified by column chromatography (PE/EA=5:1) to give the title compound as a light yellow oil.

Step 3: 3-(2-amino-6-hydroxy-5-(2-methoxybenzyl)pyrimidin-4-yl)propanenitrile

[00623] To a solution of diethyl 2-(2-methoxybenzyl)-3-oxohexanedioate (1.0 eq) in MeOH (0.6M) was added guanidine carbonate (1.0 eq). The reaction was heated at 70 °C for 16 h under nitrogen. After cooling, the reaction solution was concentrated under reduce pressure and added 1:1 H₂O/EA. The precipitates were filtered and collected to give the title compound as an off-white solid.

Step 4: 3-(2-amino-6-chloro-5-(2-methoxybenzyl)pyrimidin-4-yl)propanenitrile

[00624] A mixture of 3-(2-amino-6-hydroxy-5-(2-methoxybenzyl)pyrimidin-4-yl)propanenitrile (1.0 eq) and POCl₃ (0.3M) was stirred at 100 °C for 2 h under nitrogen. The reaction was cooled to room temperature and poured onto ice-water. The pH was adjusted to 7 using NaHCO₃ solution. The aqueous solution was extracted with EA. The organic layer was washed with water, brine, dried over Na₂SO₄, and concentrated to give the title compound as a grey solid.

Step 5: 3-(2-amino-6-(butylamino)-5-(2-methoxybenzyl)pyrimidin-4-yl)propanenitrile

[00625] To a solution of 3-(2-amino-6-chloro-5-(2-methoxybenzyl)pyrimidin-4-yl)propanenitrile (1.0 eq) in NMP (0.16M) was added butan-1-amine (3.0 eq) and DIEA (3.0 eq). The mixture was stirred at 100 °C for 16 h under nitrogen. The reaction solution was diluted with water and extracted with EA. The organic layer was washed with water, brine, dried over Na₂SO₄, and concentrated. The crude product was purified by column chromatography (DCM/MeOH=50:1) to give the title compound as a yellow solid.


Step 6: 6-(2-(2*H*-tetrazol-5-yl)ethyl)-*N*⁴-butyl-5-(2-methoxybenzyl)pyrimidine-2,4-diamine

[00626] To a solution of 3-(2-amino-6-(butylamino)-5-(2-methoxybenzyl)pyrimidin-4-yl)propanenitrile (1.0 eq) in dioxane (0.06M) was added TMSN₃ (2.0eq) and Bu₂SnO (2.0 eq). The mixture was stirred at 120 °C for 3 h. The solution was concentrated and purified by prep-HPLC (mobile phase: NH₄HCO₃/MeCN/H₂O) to give the title compound as a white solid.

[00627] LC-MS: [M+H]⁺ = 383.4

[00628] ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.18 (t, *J* = 8.4 Hz, 1H), 6.98 (d, *J* = 8.0 Hz, 1H), 6.81 (t, *J* = 7.2 Hz, 1H), 6.73 (d, *J* = 7.6 Hz, 1H), 6.59 (t, *J* = 5.6 Hz, 1H), 6.27 (br s, 2H), 3.84 (s, 3H), 3.65 (s, 2H), 3.30-3.25 (m, 2H), 3.02 (t, *J* = 7.6 Hz, 2H), 2.76 (t, *J* = 7.6 Hz, 2H), 1.49-1.37 (m, 2H), 1.23-1.16 (m, 2H), 0.83 (t, *J* = 7.6 Hz, 3H).

Example 38: (S)-3-(2-amino-5-(2-methoxybenzyl)-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-4-yl)propanoic acid (Compound 38)




Step 1: (S)-3-(2-amino-5-(2-methoxybenzyl)-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-4-yl)propanoic acid

[00629] A mixture of 3-(2-amino-6-chloro-5-(2-methoxybenzyl)pyrimidin-4-yl)propanoic acid (1.0 eq, from Example 27 – Step 4) and neat (S)-1-(methylthio)heptan-3-amine (4.0 eq, prepared by following procedures reported in WO2014/128189, pg 8, compound D) was stirred at 120 °C for 24 h. The reaction was diluted with water, and the aqueous solution was extracted with EA. The organic layers were combined and washed with 1N HCl, brine, dried over Na₂SO₄, concentrated and purified by prep-HPLC (mobile phase: 0.1% NH₃.H₂O/MeCN/H₂O) to give the title compound as a white solid.

[00630] LC-MS: [M+H]⁺ = 447.3

[00631] ^1H NMR (400 MHz, MeOD) δ 7.23 (t, J = 8.0 Hz, 1H), 7.02 (d, J = 8.0 Hz, 1H), 6.97 (d, J = 8.0 Hz, 1H), 6.92 (t, J = 8.0 Hz, 1H), 4.41-4.34 (m, 1H), 3.92 (s, 3H), 3.83 (s, 2H), 2.93 (t, J = 7.2 Hz, 2H), 2.49 (t, J = 6.8 Hz, 2H), 2.25 (t, J = 7.2 Hz, 2H), 1.96 (s, 3H), 1.76-0.90 (m, 8H), 0.81 (t, J = 7.6 Hz, 3H).

Example 39: 5-(5-((2*H*-tetrazol-5-yl)methyl)-2-methoxybenzyl)-6-methyl-*N*⁴-(pentan-2-yl)pyrimidine-2,4-diamine (Compound 39)



5-(5-((2*H*-tetrazol-5-yl)methyl)-2-methoxybenzyl)-6-methyl-*N*⁴-(pentan-2-yl)pyrimidine-2,4-diamine

Chemical Formula: $\text{C}_{20}\text{H}_{29}\text{N}_8\text{O}$


Exact Mass: 396.24

[00632] The title compound was prepared following the procedures described for Example 3, but using pentan-2-amine instead of butan-1-amine in Step 5.

[00633] LC-MS: $[\text{M}+\text{H}]^+ = 397.4$

[00634] ^1H NMR (400 MHz, CDCl_3) δ 7.26 (d, J = 8.4 Hz, 1H), 6.92 (s, 1H), 6.85 (d, J = 8.4 Hz, 1H), 6.49 (d, J = 6.8 Hz, 1H), 4.26 (s, 2H), 4.20-4.18 (m, 1H), 3.92 (s, 3H), 3.58 (s, 2H), 2.59 (s, 3H), 1.48-1.46 (m, 2H), 1.40-1.38 (m, 2H), 1.08 (d, J = 6.4 Hz, 3H), 0.85 (t, J = 7.2 Hz, 3H).

Example 40: 3-(5-(5-((2*H*-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-(butylamino)pyrimidin-4-yl)propanoic acid (Compound 40)



Step 1: (4-methoxy-3-methylphenyl)methanol

[00635] To a stirred solution of methyl 4-methoxy-3-methylbenzoate (1.0 eq) in THF (0.6M) was added LiAlH_4 (1.2 eq) in portions at 4°C under nitrogen. The resulting mixture was warmed to r.t. over 2 h. The mixture was quenched with aqueous Na_2SO_4 solution. The solution was filtered, and the filtrate was concentrated to give the title compound as a colorless oil.

Step 2: 4-(chloromethyl)-1-methoxy-2-methylbenzene

[00636] To a stirred solution of (4-methoxy-3-methylphenyl)methanol (1.0 eq) in DCM (1.4M) was added SOCl_2 (1.5 eq) at r.t. under N_2 . The mixture was stirred at r.t. for 1 h and then concentrated. The residue was partitioned between DCM and saturated aqueous NaHCO_3 solution. The organic layer was dried over Na_2SO_4 , filtered and concentrated to give the title compound.

Step 3: 2-(4-methoxy-3-methylphenyl)acetonitrile

[00637] A mixture of 4-(chloromethyl)-1-methoxy-2-methylbenzene (1.0 eq) in 1:1 DMSO/DMF (0.9M) and NaCN (2.0 eq) was stirred for 16 h at r.t. The mixture was

partitioned between EA and water. The organic layer was dried over Na₂SO₄, filtered, concentrated and purified by flash chromatography on silica (eluent PE/EA = 100:1~20:1) to give the title compound.

Step 4: 2-(3-(bromomethyl)-4-methoxyphenyl)acetonitrile

[00638] To a solution of 2-(4-methoxy-3-methylphenyl)acetonitrile (1.0 eq) in CCl₄ (0.35M) was added NBS (1.2 eq) and AIBN (2.0 g, 13 mmol, 0.1 eq). The resulting mixture was heated at reflux for 1 h. Then the mixture was filtered, and the filtration was concentrated under reduced pressure to give the crude product, which was purified by flash chromatography on silica (eluent: PE/EA = 40:1 to 4:1) to give the title compound as a yellow solid.

Step 5: diethyl 2-(5-(cyanomethyl)-2-methoxybenzyl)-3-oxohexanedioate

[00639] To a solution of 2-(3-(bromomethyl)-4-methoxyphenyl)acetonitrile (1.0 eq) in anhydrous THF (0.73ML) at 0 °C was added portion-wise of 60% NaH (1.2 eq). After stirring for 10 min, a solution of diethyl 3-oxohexanedioate (1.1 eq, from Example 27 – Step 1) in THF (2.4M) was added drop-wise into the above mixture over 10 min. The resulting mixture was warmed to r.t. and stirred for 20 h. The reaction was quenched with water. The mixture was extracted with EtOAc, dried over Na₂SO₄, concentrated under reduced pressure. The residue was purified by flash chromatography on silica (eluent: PE/EA = 50:1 to 10:1) to give the title compound as a yellow oil.

Step 6: 3-(2-amino-5-(5-(cyanomethyl)-2-methoxybenzyl)-6-hydroxypyrimidin-4-yl)propanoic acid

[00640] A mixture of diethyl 2-(5-(cyanomethyl)-2-methoxybenzyl)-3-oxohexanedioate (1.0 eq) in MeOH (0.3M) and guanidine carbonate (1.5 eq) was stirred at 70 °C for 16 h. Solvent was removed, and the residue was diluted with water and acidified by adding 1 N HCl to pH 5. The resulting precipitate was filtered and dried to give the title compound as a white solid.

Step 7: 3-(2-amino-6-chloro-5-(5-(cyanomethyl)-2-methoxybenzyl)pyrimidin-4-yl)propanoic acid

[00641] A mixture of 3-(2-amino-5-(5-(cyanomethyl)-2-methoxybenzyl)-6-hydroxypyrimidin-4-yl)propanoic acid (1.0 eq) in POCl₃ (1.9M) was stirred at 100 °C for 3 h. The reaction mixture was cooled to r.t., and POCl₃ was evaporated under reduced pressure. The residue was diluted with water. The pH was adjusted to 7 with solid NaHCO₃. The

precipitate was collected by filtration, washed with water, and dried under vacuum to give the title compound as a greenish solid.

Step 8: 3-(2-amino-6-(butylamino)-5-(5-(cyanomethyl)-2-methoxybenzyl)pyrimidin-4-yl)propanoic acid

[00642] A mixture of 3-(2-amino-6-chloro-5-(5-(cyanomethyl)-2-methoxybenzyl)pyrimidin-4-yl)propanoic acid (1.0 eq) in EtOH (0.3M), butan-1-amine (4 eq), and DIEA (5 eq) was stirred at 85 °C for 48 h. Solvent was removed, and the residue was purified by flash column chromatography (eluent: DCM/MeOH = 40:1) to give the title compound as a pale solid.


Step 9: 3-(5-(5-((2*H*-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-(butylamino)pyrimidin-4-yl)propanoic acid

[00643] A mixture of 3-(2-amino-6-(butylamino)-5-(5-(cyanomethyl)-2-methoxybenzyl)pyrimidin-4-yl)propanoic acid (1.0 eq) in dioxane (0.1M), TMSN₃ (5 eq) and Bu₂SnO (2.0 eq) was heated in a seal tube at 70 °C for 3 h. The mixture was concentrated under reduced pressure, and the crude was purified by prep-HPLC (mobile phase: 0.1% NH₃.H₂O in CH₃CN) to give the title compound as an off-white solid.

[00644] LC-MS: [M+H]⁺ = 441.2

[00645] ¹H NMR (400 MHz, CDCl₃) δ 7.46 (s, 1H), 7.24 (d, *J* = 8.4 Hz, 1H), 6.84 (d, *J* = 8.4 Hz, 1H), 5.78 (br s, 1H), 4.17 (s, 2H), 3.90 (s, 3H), 3.70 (s, 2H), 3.36-3.31 (m, 2H), 3.17 (t, *J* = 6.0 Hz, 2H), 2.77 (t, *J* = 6.0 Hz, 2H), 1.41-1.38 (m, 2H), 1.18-1.13 (m, 2H), 0.84 (t, *J* = 6.8 Hz, 3H).

Example 41: (S)-3-((2-amino-5-(2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoic acid (Compound 41)



Step 1: ethyl 2-(2-methoxybenzyl)-3-oxobutanoate

[00646] To a solution of 1-(bromomethyl)-2-methoxybenzene (1.2 eq) in THF (0.6M) at 0 °C was added 60% NaH (1.25 eq) in portions under N_2 . The resulting suspension was stirred at 0 °C for 10 min, then a solution of ethyl 3-oxobutanoate (1.0 eq) in THF (5M) was added drop-wise over 10 min. The resulting mixture was stirred for 15 h at 70 °C. The mixture was cooled to r.t. and ice water was added. The mixture was extracted with ethyl acetate. The organic layer was dried over Na_2SO_4 , concentrated, and purified by column chromatography on silica gel (eluent PE/EA = 50:1~5:1) to give the title compound as a yellow oil.

Step 2: 2-amino-5-(2-methoxybenzyl)-6-methylpyrimidin-4-ol

[00647] To a solution of ethyl 2-(2-methoxybenzyl)-3-oxobutanoate (1.0 eq) in MeOH (0.9M) was added guanidine carbonate (1.0 eq). The resulting mixture was stirred overnight at 65 °C. The mixture was concentrated to dryness. The residue was suspended in EtOAc and then filtered. The filter cake was washed with water and EtOAc, and then dried under vacuum to give the title compound as a white solid.

Step 3: 4-chloro-5-(2-methoxybenzyl)-6-methylpyrimidin-2-amine

[00648] A suspension of 2-amino-5-(2-methoxybenzyl)-6-methylpyrimidin-4-ol (1.0 eq) in POCl_3 (1M) was stirred at 100 °C for 24 h. The mixture was cooled to r.t. and POCl_3

was evaporated to dryness under reduced pressure. The residue was diluted with water, and pH was adjusted to 7 using solid NaHCO₃. The precipitate was collected by filtration, washed with water and EtOAc, and dried under vacuum to give the title compound.

Step 4: *tert*-butyl (*S*)-3-((2-amino-5-(2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoate

[00649] A mixture of 4-chloro-5-(2-methoxybenzyl)-6-methylpyrimidin-2-amine (1.0 eq) in NMP (1.3M) and *tert*-butyl (*S*)-3-aminoheptanoate (4.5 eq, from Example 20B – Step 3) was stirred at 120 °C for 3 h. The mixture was partitioned between water and EtOAc. The organic layer was washed with brine, dried over Na₂SO₄, and concentrated under reduced pressure. The residue was purified by flash column chromatography (eluent: DCM/MeOH = 100:1 to 30:1) to give the title compound as a yellow solid.


Step 5: (*S*)-3-((2-amino-5-(2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoic acid

[00650] A mixture of *tert*-butyl (*S*)-3-((2-amino-5-(2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoate (1.0 eq) in 1:1 EtOH/H₂O (0.1M) and NaOH (10.0 eq) was heated at 100 °C for 2 h. The reaction was cooled to r.t., and the solvent was removed. The residue was diluted with H₂O and acidified using 1N HCl to pH 2. The resulting suspension was extracted with DCM. The combined organic layers were dried and concentrated. The crude was purified by prep-TLC (CHCl₃: MeOH=10:1) to give an oil, which was dissolved in 1:1 CH₃CN/H₂O and added 4N HCl/dioxane. The solution was freeze-dried to give the title compound as a white solid.

[00651] LC-MS: [M+H]⁺ = 373.2

[00652] ¹H NMR (400 MHz, DMSO-*d*₆) δ 12.20 (br, 1H), 7.20-7.17 (m, 1H), 6.99 (d, *J* = 8.4 Hz, 1H), 6.82-6.79(m, 2H), 6.68 (br, 2H), 4.53-4.50 (m, 1H), 3.85 (s, 3H), 3.66 (s, 2H), 2.50-2.37 (m, 2H), 2.07 (s, 3H), 1.51-1.38 (m, 2H), 1.24-0.98 (m, 4H), 0.77 (t, *J* = 6.8 Hz, 3H).

Example 42: (*S*)-3-((2-amino-5-(5-(2-carboxypropan-2-yl)-2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoic acid (Compound 42)



Step 1: *tert*-butyl (*S*)-3-((2-amino-5-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoate

[00653] A mixture of 2-(3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanenitrile (1.0 eq, from Example 9 – Step 8) in NMP (0.9M) and *tert*-butyl (*S*)-3-aminoheptanoate (1.3 eq, from Example 20B – Step 3) was stirred at 120 °C for 16 h under nitrogen. The reaction mixture was diluted with water. The aqueous phase was extracted with EA. The combined organic layer was washed with water, then brine. The organic layer was separated and dried over Na₂SO₄, concentrated and purified by flash chromatography on silica (eluent DCM/MeOH=50:1) to give the title compound as a yellow solid.


Step 2: (*S*)-3-((2-amino-5-(5-(2-carboxypropan-2-yl)-2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoic acid

[00654] A mixture of *tert*-butyl (*S*)-3-((2-amino-5-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoate (1.0 eq) in 1:1 ethylene glycol/H₂O (0.1M) was added NaOH (10.0 eq) and stirred at 150 °C for 16 h. The mixture was neutralized by adding 1 N HCl. The resulting suspension was filtered. The filter cake was washed with water and purified by prep-HPLC (mobile phase: 0.1% HCOOH/MeCN/H₂O) to give the title compound as a white solid.

[00655] LC-MS: [M+H]⁺=459.3

[00656] ¹H NMR (400 MHz, DMSO-*d*₆) δ 7.19 (br, 1H), 7.18 (dd, *J* = 8.4, 2.4 Hz, 1H), 6.96 (d, *J* = 8.4 Hz, 1H), 6.84 (d, *J* = 2.0 Hz, 1H), 4.58-4.56 (m, 1H), 3.83 (s, 3H), 3.79 (s, 2H), 2.48-2.42 (m, 2H), 2.13 (s, 3H), 1.52-1.50 (m, 2H), 1.38 (s, 3H), 1.35 (s, 3H), 1.25-1.18 (m, 2H), 1.24-1.07 (m, 2H), 0.91 (t, *J* = 7.2 Hz, 3H).

Example 43: 2-(3-((2-amino-4-(butylamino)-6-(2-carboxyethyl)pyrimidin-5-yl)methyl)-4-methoxy-phenyl)-2-methylpropanoic acid (Compound 43)



Step 1: diethyl 2-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)-3-oxohexanedioate

[00657] To a solution of 2-(3-(bromomethyl)-4-methoxyphenyl)-2-methylpropanenitrile (1.1 eq, from Example 9 – Step 5) in THF (0.3M) was added NaH (1.2eq) at 0 °C in portions under nitrogen. The mixture was stirred at 0 °C for 10 min. A solution of diethyl 3-oxohexanedioate (1.0 eq, from Example 27 – Step 1) in THF (0.75M) was added to the above solution, and the reaction was stirred at 60 °C for 16 h. The reaction was quenched by water. The aqueous solution was extracted with EA. The combined organic layers were washed with water, brine, dried over Na_2SO_4 , and concentrated. The crude product was purified by column chromatography (PE/EA=10:1) to give the title compound as a yellow oil.

Step 2: 3-(2-amino-5-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)-6-hydroxypyrimidin-4-yl)propanoic acid

[00658] To a solution of diethyl 2-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)-3-oxohexanedioate (1.0 eq) in MeOH (0.7M) was added guanidine carbonate (1.0 eq). The reaction was heated at 65 °C for 16 h under nitrogen. After cooling, the reaction solution was concentrated under reduced pressure. The residue was purified by column chromatography (DCM/MeOH=5:1) to give the title compound as a white solid.

Step 3: 3-(2-amino-6-chloro-5-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)pyrimidin-4-yl)propanoic acid

[00659] A solution of 3-(2-amino-5-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)-6-hydroxypyrimidin-4-yl)propanoic acid (1.0 eq) in POCl_3 (0.3M) was stirred at 100 °C for 16 h under nitrogen. After cooling, the mixture was concentrated and poured into water. The pH value was adjusted to 8 by adding NaHCO_3 . The aqueous phase was extracted with EA. Combined organic layers were dried over Na_2SO_4 , concentrated, and purified by column chromatography (DCM/MeOH=10:1) to give the title compound as a brown solid.

Step 4: 3-(2-amino-6-(butylamino)-5-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)pyrimidin-4-yl)propanoic acid

[00660] A solution of 3-(2-amino-6-chloro-5-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)pyrimidin-4-yl)propanoic acid (1.0 eq) in neat butan-1-amine (0.15M) was stirred at 120 °C for 3 h under nitrogen. The reaction was purified by column chromatography (DCM/MeOH=10:1) to give the title compound as a brown oil.


Step 5: 2-(3-((2-amino-4-(butylamino)-6-(2-carboxyethyl)pyrimidin-5-yl)methyl)-4-methoxyphenyl)-2-methylpropanoic acid

[00661] To a solution of 3-(2-amino-6-(butylamino)-5-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)pyrimidin-4-yl)propanoic acid (1.0 eq) in 1:1 ethylene glycol/ H_2O (0.07M) was added KOH (20 eq). The mixture was stirred at 150 °C for 16 h. After cooling, pH was adjusted to 6 by adding 4N HCl. The mixture was extracted with EA. The organic layer was washed with brine, dried over Na_2SO_4 , concentrated. The crude product was purified by prep-HPLC (mobile phase: $\text{NH}_4\text{HCO}_3/\text{MeCN}/\text{H}_2\text{O}$) to give the title compound as a white solid.

[00662] LC-MS: $[\text{M}+\text{H}]^+ = 445.3$

[00663] ^1H NMR (400 MHz, CD_3OD) δ 7.27 (dd, $J=8.4, 2.4\text{Hz}$, 1H), 6.95 (d, $J=8.4\text{Hz}$, 1H), 6.93 (s, 1H), 3.89 (s, 3H), 3.78 (s, 2H), 3.45 (t, $J=7.0\text{Hz}$, 2H), 2.84 (t, $J=6.0\text{Hz}$, 2H), 2.42 (t, $J=6.4\text{Hz}$, 2H), 1.54-1.47 (m, 2H), 1.45 (s, 6H), 1.26-1.21 (m, 2H), 0.88 (t, $J=7.2\text{Hz}$, 3H).

Example 44: (*S*)-3-((2-amino-5-(2-methoxy-5-((3-phosphonopropyl)carbamoyl)benzyl)-6-methylpyrimidin-4-yl)amino)heptanoic acid (Compound 44)



Step 1: methyl (*S*)-3-((2-amino-4-((1-(*tert*-butoxy)-1-oxoheptan-3-yl)amino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate

[00664] A mixture of methyl 3-((2-amino-4-chloro-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate (1.0 eq, from Example 3B – Step 4) in NMP (2M) and *tert*-butyl (*S*)-3-aminoheptanoate (2.0 eq, from Example 20B – Step 3) was stirred at 120 °C for 16 h under nitrogen. The reaction solution was diluted with water and extracted with EA. The organic layer was washed with water, brine, dried over Na₂SO₄, and concentrated. The residue was purified by column chromatography (DCM/MeOH=50:1) to give the title as a yellow solid.

Step 2: (*S*)-3-((2-amino-4-((1-(*tert*-butoxy)-1-oxoheptan-3-yl)amino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoic acid

[00665] A mixture of methyl (*S*)-3-((2-amino-4-((1-(*tert*-butoxy)-1-oxoheptan-3-yl)amino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoate (1.0 eq) in 1:1 ethanol/H₂O (0.1M) and NaOH (10.0 eq) was stirred at 30 °C for 4 h. The mixture was neutralized by adding 1 N HCl to pH 7. The resulting suspension was filtered. The filter cake was washed with water to give the title compound.

Step 3: *tert*-butyl (*S*)-3-((2-amino-5-(5-((3-(diethoxyphosphoryl)propyl)carbamoyl)-2-methoxybenzyl)-6-methylpyrimidin-4-yl)amino)heptanoate

[00666] A mixture of (*S*)-3-((2-amino-4-((1-(*tert*-butoxy)-1-oxoheptan-3-yl)amino)-6-methylpyrimidin-5-yl)methyl)-4-methoxybenzoic acid (1.0 eq) in DMF (0.1M), diethyl (3-aminopropyl)phosphonate (2.0 eq, from Example 30 – Step 2), HATU (2.0 eq) and DIEA (3.0 eq) was stirred at 50 °C for 16 h. The mixture was cooled to r.t. and added water. The mixture was extracted with DCM. The combined organic phase was dried over Na₂SO₄, and concentrated under reduced pressure to give the title compound as a colorless oil.


Step 4: (*S*)-3-((2-amino-5-(2-methoxy-5-((3-phosphonopropyl)carbamoyl)benzyl)-6-methyl-pyrimidin-4-yl)amino)heptanoic acid

[00667] A solution of *tert*-butyl (*S*)-3-((2-amino-5-(5-((3-(diethoxyphosphoryl)propyl)carbamoyl)-2-methoxy-benzyl)-6-methylpyrimidin-4-yl)amino)heptanoate (1.0 eq) in DCM (0.08M) was added TSMBr (10.0 eq) and stirred at 35 °C for 16 h under N₂. The mixture was concentrated, and the residue was purified by trituration and prep-HPLC (mobile phase CH₃CN/H₂O/NH₃H₂O) to give the title compound.

[00668] LC-MS: [M+H]⁺ = 538.3

[00669] ¹H NMR (400 MHz, DMSO) δ 8.69 (br, 1H), 7.72 (d, *J* = 8.8 Hz, 1H), 7.50 (s, 1H), 7.01 (d, *J* = 8.4 Hz, 1H), 6.63 (br, 1H), 4.47-4.45 (m, 1H), 3.85 (s, 3H), 3.27-3.25 (m, 4H), 2.4 (d, *J* = 6.0 Hz, 2H), 2.13 (s, 3H), 1.72-1.68 (m, 2H), 1.46-1.26 (m, 4H), 1.12-0.96 (m, 2H), 0.96-0.94 (m, 2H), 0.70 (t, *J* = 7.2 Hz, 3H).

Example 45: (*S*)-3-((2-amino-6-(2-carboxyethyl)-5-(5-(2-carboxypropan-2-yl)-2-methoxy-benzyl)pyrimidin-4-yl)amino)heptanoic acid (Compound 45)



Step 1: (*S*)-3-(2-amino-6-((1-(*tert*-butoxy)-1-oxoheptan-3-yl)amino)-5-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)pyrimidin-4-yl)propanoic acid

[00670] To a solution of 3-(2-amino-6-chloro-5-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)pyrimidin-4-yl)propanoic acid (1.0 eq, from Example 43 – Step 3) in NMP (0.15M) was added *tert*-butyl (*S*)-3-aminoheptanoate (5.0 eq, from Example 20B – Step 3) and K₂CO₃ (3.0 eq). The mixture was stirred at 150 °C for 16 h. The reaction solution was diluted with water/EA and concentrated to give the title compound as a crude oil.


Step 2: (*S*)-3-((2-amino-6-(2-carboxyethyl)-5-(5-(2-carboxypropan-2-yl)-2-methoxybenzyl)-pyrimidin-4-yl)amino)heptanoic acid

[00671] To a solution of (*S*)-3-(2-amino-6-((1-(*tert*-butoxy)-1-oxoheptan-3-yl)amino)-5-(5-(2-cyanopropan-2-yl)-2-methoxybenzyl)pyrimidin-4-yl)propanoic acid (crude) in 1:1 ethylene glycol/H₂O was added excess KOH and stirred at 150 °C for 16 h. After cooling, the reaction mixture was diluted with water/EA. The pH was adjusted to 6 by adding 4N HCl, then extracted with 3:1 CHCl₃/IPA. The combined organic layer was washed with brine, dried over Na₂SO₄, and concentrated. The crude product was purified by prep-HPLC (mobile phase: NH₄OH/MeCN/H₂O) to give the title compound as a white solid.

[00672] LC-MS: [M+H]⁺ = 517.3

[00673] ¹H NMR (400 MHz, CD₃OD) δ 7.27 (dd, *J* = 8.0, 2.4 Hz, 1H), 7.00-6.95 (m, 2H), 4.61-4.57 (m, 1H), 3.90 (s, 3H), 3.80 (s, 2H), 2.94-2.87 (m, 2H), 2.50-2.38 (m, 4H), 1.56-1.53 (m, 1H), 1.48 (s, 3H), 1.45 (s, 3H), 1.46-1.44 (m, 1H), 1.27-1.17 (m, 2H), 1.08-1.03 (m, 2H), 0.79 (t, *J* = 7.4 Hz, 3H).

Example 46: (S)-3-((5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-(2-carboxyethyl)-pyrimidin-4-yl)amino)heptanoic acid (Compound 46)



Step 1: *tert*-butyl (S)-3-(3-(2-amino-6-(((S)-1-(*tert*-butoxy)-1-oxoheptan-3-yl)amino)-5-(5-(cyanomethyl)-2-methoxybenzyl)pyrimidin-4-yl)propanamido)heptanoate

[00674] A mixture of 3-(2-amino-6-chloro-5-(5-(cyanomethyl)-2-methoxybenzyl)pyrimidin-4-yl)propanoic acid (1.0 eq, from Example 40 – Step 7) and *tert*-butyl (S)-3-aminoheptanoate (3 eq., from Example 20B – Step 3) was stirred at 140 °C for 3 h. The crude product was used directly in next step.

Step 2: *tert*-butyl (S)-3-(3-(5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-(((S)-1-(*tert*-butoxy)-1-oxoheptan-3-yl)amino)pyrimidin-4-yl)propanamido)heptanoate

[00675] A mixture of *tert*-butyl (S)-3-(3-(2-amino-6-(((S)-1-(*tert*-butoxy)-1-oxoheptan-3-yl)amino)-5-(5-(cyanomethyl)-2-methoxybenzyl)pyrimidin-4-yl)propanamido)heptanoate (1.0 eq) in dioxane (0.2M), TMSN₃ (3 eq) and Bu₂SnO (2.0 eq) was heated at 110 °C in a seal tube for 6 h. The mixture was diluted with EA and washed with water. The organic layer was dried, concentrated, and the residue was purified by column chromatography (DCM:MeOH=100:1~10:1) to give the title compound as a light brown solid.

Step 3: (S)-3-((5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-(2-carboxyethyl)-pyrimidin-4-yl)amino)heptanoic acid


[00676] A solution of *tert*-butyl (S)-3-(3-(5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-(((S)-1-(*tert*-butoxy)-1-oxoheptan-3-yl)amino)pyrimidin-4-

yl)propanamido)heptanoate (1.0 eq) in ethane-1,2-diol (0.1M) was added 10M aqueous KOH (44 eq) and heated at 150 °C for 5 days. The solid was filtered off, and the filtrate was concentrated and purified by prep-HPLC (mobile phase: 0.1% NH₃.H₂O, CH₃CN) to give the title compound as a white solid.

[00677] LC-MS: [M+H]⁺ = 513.2

[00678] ¹H NMR (400 MHz, MeOD) δ 7.09 (dd, *J* = 8.4, 2.0 Hz, 1H), 6.95 (s, 1H), 6.89 (d, *J* = 8.4 Hz, 1H), 4.51-4.49 (m, 1H), 4.04 (s, 2H), 3.87 (s, 3H), 3.77 (s, 2H), 2.92-2.85 (m, 2H), 2.52-2.21 (m, 4H), 1.5-0.9 (m, 6H), 0.73 (t, *J* = 7.6 Hz, 3H).

Example 47: 3-(2-amino-6-(butylamino)-5-(2-methoxybenzyl)pyrimidin-4-yl)propan-1-ol (Compound 47)



Step 1: 3-(2-amino-6-chloro-5-(2-methoxybenzyl)pyrimidin-4-yl)propan-1-ol

[00679] To a solution of 3-(2-amino-6-chloro-5-(2-methoxybenzyl)pyrimidin-4-yl)propanoic acid (1.0 eq, from Example 27 – Step 4) in THF (0.16M) was added LAH (3.0 eq). The reaction solution was stirred at r.t. for 2 h, and then quenched with water. The aqueous layer was extracted with DCM. The combined organic layers were washed with water, brine, dried over Na₂SO₄, and concentrated to give the title compound as a pale solid.

Step 2: 3-(2-amino-6-(butylamino)-5-(2-methoxybenzyl)pyrimidin-4-yl)propan-1-ol


[00680] A solution of 3-(2-amino-6-chloro-5-(2-methoxybenzyl)pyrimidin-4-yl)propan-1-ol (1.0 eq) in EtOH (0.2M), butan-1-amine (4.0 eq), and DIEA (5 eq) was stirred at 85 °C for 3 days. The reaction solution was concentrated, diluted with 1N HCl, and extracted with DCM. The combined organic layers were washed with brine, dried over Na₂SO₄, concentrated, and purified by flash column chromatography (eluent: DCM/MeOH = 100:1~20:1) to give the title compound as a pale solid.

[00681] LC-MS: [M+H]⁺ = 345.4

[00682] ¹H NMR (400 MHz, DMSO) δ 7.80 (br, 1H), 7.40 (br, 2H), 7.22 (t, *J* = 8.0 Hz, 1H), 7.01 (d, *J* = 8.4 Hz, 1H), 6.84 (t, *J* = 7.6 Hz, 1H), 6.78 (d, *J* = 8.4 Hz, 1H), 3.85 (s,

3H), 3.72 (s, 2H), 3.38-3.34 (m, 2H), 2.50-2.44 (m, 2H), 1.61-1.54 (m, 2H), 1.50-1.42 (m, 2H), 1.34-1.15 (m, 4H), 0.84(t, $J = 7.6$ Hz, 3H).

Example 48: (S)-3-(5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-4-yl)propanoic acid (Compound 48)



Step 1: 3-(2-amino-5-(5-(cyanomethyl)-2-methoxybenzyl)-6-(((S)-1-(methylthio)heptan-3-yl)amino)-pyrimidin-4-yl)-N-((S)-1-(methylthio)heptan-3-yl)propanamide

[00683] A solution of 3-(2-amino-6-chloro-5-(5-(cyanomethyl)-2-methoxybenzyl)pyrimidin-4-yl)propanoic acid (1.0 eq, from Example 40 – Step 7) in EtOH (0.4M) and (S)-1-(methylthio)heptan-3-amine (2.3 eq) was stirred at 90 °C for 3 days. Solvent was removed, the residue was purified by column chromatography (eluent DCM: MeOH=20:1) to give the title compound as a light brown oil.

Step 2: 3-(5-(5-((2H-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-(((S)-1-(methylthio)heptan-3-yl)amino)pyrimidin-4-yl)-N-((S)-1-(methylthio)heptan-3-yl)propanamide

[00684] A mixture of 3-(2-amino-5-(5-(cyanomethyl)-2-methoxybenzyl)-6-(((S)-1-(methylthio)heptan-3-yl)amino)-pyrimidin-4-yl)-N-((S)-1-(methylthio)heptan-3-yl)propanamide (1.0 eq) in dioxane (0.04M), TMSN₃ (3.8 eq), and Bu₂SnO (2.5 eq) was heated at 90 °C for 6 h in a seal tube. The reaction solution was cooled down and used directly in the next step without any purification.


Step 3: (S)-3-(5-(5-((2*H*-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-4-yl)propanoic acid

[00685] The crude product from the previous step was dissolved in ethane-1, 2-diol and added aqueous KOH (excess). The reaction was heated at 150 °C for 5 days. The solid was filtered, and the filtrate was concentrated and purified by prep-HPLC (mobile phase: 0.1% TFA/CH₃CN/H₂O) to give the title compound as a white solid.

[00686] LC-MS: [M+H]⁺ = 529.3

[00687] ¹H NMR (400 MHz, MeOD) δ 7.20 (d, *J* = 7.6 Hz, 1H), 7.01 (d, *J* = 8.8 Hz, 1H), 6.93 (s, 1H), 4.43-4.40 (m, 1H), 4.19 (s, 2H), 3.91 (s, 3H), 3.83 (s, 2H), 2.92 (t, *J* = 6.8 Hz, 2H), 2.56 (t, *J* = 7.6 Hz, 2H), 2.24 (t, *J* = 7.6 Hz, 2H), 1.93 (s, 3H), 1.78-1.75 (m, 1H), 1.67-1.63 (m, 1H), 1.51-1.49 (m, 1H), 1.41-1.39 (m, 1H), 1.26-1.19 (m, 2H), 1.10-1.04 (m, 2H), 0.80 (t, *J* = 7.6 Hz, 3H).

Example 49: (S)-3-(5-(5-((2*H*-tetrazol-5-yl)methyl)-2-methoxybenzyl)-2-amino-6-((1-methoxyheptan-3-yl)amino)pyrimidin-4-yl)propanoic acid (Compound 49)




[00688] The title compound was prepared following the procedures described for Example 48, but using (S)-1-methoxyheptan-3-amine instead of (S)-1-(methylthio)heptan-3-amine in Step 1.

[00689] LC-MS: [M+H]⁺ = 513.3

[00690] ¹H NMR (400 MHz, CDCl₃) δ 7.21 (d, *J* = 8.4 Hz, 1H), 7.17 (s, 1H), 6.82 (d, *J* = 8.4 Hz, 1H), 6.14 (d, *J* = 8.0 Hz, 1H), 4.26-4.05 (m, 2H), 4.03 (d, *J* = 15.2 Hz, 1H), 3.88 (s, 3H), 3.74-3.61 (m, 2H), 3.33-3.25 (m, 2H), 3.17-3.05 (m, 2H), 3.04 (s, 3H), 2.72-2.68 (m, 1H), 2.53-2.48 (m, 1H), 1.81-1.76 (m, 1H), 1.55-1.50 (m, 1H), 1.41-1.15 (m, 4H), 1.03-0.95 (m, 2H), 0.78 (t, *J* = 7.6 Hz, 3H).

Example 50: (S)-3-(2-amino-5-(5-(carboxymethyl)-2-methoxybenzyl)-6-((1-(methylsulfonyl)heptan-3-yl)amino)pyrimidin-4-yl)propanoic acid (Compound 50)



Step 1: 3-(2-amino-5-(5-(cyanomethyl)-2-methoxybenzyl)-6-(((S)-1-(methylthio)heptan-3-yl)amino)-pyrimidin-4-yl)-N-((S)-1-(methylthio)heptan-3-yl)propanamide


[00691] The title compound was prepared following the procedures described for Example 48 – Step 1.

Step 2: (S)-3-(2-amino-5-(5-(carboxymethyl)-2-methoxybenzyl)-6-((1-(methylsulfonyl)heptan-3-yl)amino)pyrimidin-4-yl)propanoic acid

[00692] To a solution of KOH in 10% H_2O in nBuOH (10M) was added 3-(2-amino-5-(5-(cyanomethyl)-2-methoxybenzyl)-6-(((S)-1-(methylthio)heptan-3-yl)amino)-pyrimidin-4-yl)-N-((S)-1-(methylthio)heptan-3-yl)propanamide (1.0 eq). The resulting mixture was heated in a seal tube at 150 °C for 16 h. The mixture was cooled to r.t. and filtered. To the filtrate was added oxone (5.0 eq) and stirred at r.t. for 5 h. The reaction was concentrated under reduced pressure to 1/3 volume. The insoluble solid was filtered off and the filtrate was purified by prep-HPLC (mobile phase: 0.1% $HCOOH/MeCN/H_2O$) and freeze-dried to give the title compound as a white solid.

[00693] LC-MS: $[M+H]^+ = 537.4$

Example 51: (*S*)-2-(3-((2-amino-4-methyl-6-((1-(methylsulfonyl)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxyphenyl)acetic acid (Compound 51)



[00694] Step 1: (*S*)-2-(3-((2-amino-4-methyl-6-((1-(methylsulfonyl)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxyphenyl)acetonitrile

To a solution of (*S*)-2-(3-((2-amino-4-methyl-6-((1-(methylthio)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxyphenyl)acetonitrile (1 eq, from Example 22 – Step 4) in 1:1:1 THF/MeOH/H₂O (0.2M) was added oxone (1.2eq) in portions at r.t. The reaction was stirred at r.t. for 2 h, and then diluted with DCM. The organic layer was washed with water and brine, dried, and concentrated to give a light yellow solid, which was used in the next step directly.

[00695] Step 2: (*S*)-2-(3-((2-amino-4-methyl-6-((1-(methylsulfonyl)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxyphenyl)acetic acid

To a solution of (*S*)-2-(3-((2-amino-4-methyl-6-((1-(methylsulfonyl)heptan-3-yl)amino)pyrimidin-5-yl)methyl)-4-methoxyphenyl)acetonitrile (1.0 eq) in 1:1 MeOH/H₂O (0.1M) was added KOH (7.5 eq). The reaction was stirred at 120 °C for 4 h. Solvent was removed and HCl was added to achieve pH 9. The mixture was purified by prep-HPLC (0.1% NH₃·H₂O/CH₃CN) to give the title compound as a light yellow solid.

[00696] LCMS: [M+H]⁺=479.3

[00697] ¹H NMR (400 MHz, CD₃OD) δ 7.16 (dd, *J* = 8.4, 2.0 Hz, 1H), 6.98 (d, *J* = 2.0 Hz, 1H), 6.91 (d, *J* = 8.4 Hz, 1H), 4.30-4.22 (m, 1H), 3.89 (s, 3H), 3.77-3.68 (m, 2H), 3.32 (s, 2H), 2.91-2.70 (m, 5H), 2.32 (s, 3H), 2.08-1.95 (m, 1H), 1.81-1.69 (m, 1H), 1.56-1.05 (m, 6H), 0.83 (t, *J* = 7.2 Hz, 3H).

BIOLOGICAL EXAMPLES

Biological Example 1: HEK TLR7 Assay

[00698] HEK-Blue™ TLR7 cells were purchased from Invivogen (San Diego, California). The following description was taken from the product information sheet.

[00699] “HEK-Blue™ hTLR7 cells are designed for studying the stimulation of human TLR7 (hTLR7) by monitoring the activation of NF-κB. HEK-Blue™ hTLR7 cells were obtained by co-transfection of the hTLR7 gene and an optimized secreted embryonic alkaline phosphatase (SEAP) reporter gene into HEK293 cells. The SEAP reporter gene is placed under the control of the IFN-β minimal promoter fused to five NF-κB and AP-1-binding sites. Stimulation with a TLR7 ligand activates NF-κB and AP-1 which induce the production of SEAP, which is detected by the HEK-Blue™ Detection cell culture medium.”

[00700] A typical assay protocol involved the following steps:

1. Cells were cultured according to the product information sheet.
2. 10 mM compound stock in DMSO were first diluted to 3 mM and then 3-fold serially diluted using DMSO to afford a 10-pt dilution.
3. 3 µl of the diluted DMSO were added to 57 µl HEK-Blue™ Detection media for a further 20-fold dilution.
4. 10 µl of the diluted compound in assay media were added into 40 µl cell culture (in HEK-Blue™ Detection media) in 384-well plate. Final cell concentration = 8,000 cells per well.
5. The plates were incubated at 37 °C in 5% CO₂ for 16 h. SEAP was determined using a spectrophotometer at 620-655 nm.

[00701] HEK-TLR7 activity table below provides results.

Example	HEK-TLR7 EC ₅₀	Example	HEK-TLR7 EC ₅₀
1	C	31	C
2	C	32	A
3	B	33	A
4	C	34	A
5	C	35	A
6	C	36	B
7	C	37	B
8	B	38	A
9	B	39	B
10	B	40	B
11	B	41	B
12	B	42	B
13	B	43	C
14	C	44	B
15	B	45	C
16	B	46	B
17	A	47	B
18	A	48	A
19	C	49	B
20 (S-enantiomer)	B	50	B
20C (R-enantiomer)	C	51	B
21	C		
22	A		
23	A		
24	A		
25	B		
26	B		
27	B		
28	C		
29	B		
30	B		


EC₅₀ category:

A = <100 nM

B = 100-1000 nM

C = 1000-10000 nM

[00702] As discussed herein, R^{1A} and R^{1B} in formula (1) can create a chiral center. Compounds 20 and 20C differ by the stereocenter of the carbon bearing R^{1A} and R^{1B}.



As shown above, the (S)-enantiomer (e.g. Compound 20) is >10-fold more potent than the (R)-enantiomer (e.g. Compound 20C) in the HEK-TLR7 assay. Other examples can follow the same trend.

Biological Example 2: Pharmacokinetics experiments

[00703] BALB/c mice were dosed p.o. with 10 mg/kg compound formulated in 0.5% carboxy methylcellulose + 0.5% Tween-80 suspension. Mice were bled retro-orbitally at different time points post dose. Blood was processed into serum by centrifugation followed by protein precipitation, reverse-phase gradient elution and MRM detection via ESI+ mass spectrometry to determine compound concentration. Animals were sacrificed after the final blood collection (5 hour), liver removed and flash frozen in liquid nitrogen for tissue PK analyses. The liver-to-serum concentration (L:S) at 5 hour post dose is one parameter for identifying compounds with potential enrichment in liver.

[00704] Mouse liver-to-serum concentration (L:S) table below provides results.

Example	L:S at 5hr
3	A
9	A
10	B
11	A
12	B
13	B
15	B
17	B
18	A
20	A
22	A
23	A
24	B
29	A
30	A
32	A

33	A
34	A
40	A
41	B
42	A
43	A
44	B
45	A
46	B

L:S category:

A = >100 represents high liver-enrichment

B = 10-100 represents moderate liver-enrichment


C = <10 represents low liver-enrichment

EQUIVALENTS

[00705] While the present invention has been described in conjunction with the specific embodiments set forth above, many alternatives, modifications and other variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.


We claim:

1. A compound having the structure of Formula (1), or a pharmaceutically acceptable salt thereof,




wherein

R^{1a} is selected from the group consisting of H, C₁-C₄ alkyl, -NH₂, -NHAc,

-COOH, -SO₂CH₃, -SCH₃, -OCH₃, ; and A,

wherein the alkyl is optionally substituted with -OH, -NH₂, -NHAc, -COOH, -SO₂CH₃,








-SCH₃, -OCH₃, , or A;

R^{1b} is C₂-C₅ alkyl;

X is selected from the group consisting of H and C₁-C₄ alkyl, wherein the alkyl is optionally substituted with A, -OH, or -C(CH₃)₂OH;

L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , -O-, -S-, -SO₂-, -NH-, and -CH₂CH₂-;

Y is selected from the group consisting of C₁-C₃ alkyl, aryl, and heteroaryl, wherein the alkyl, aryl, and heteroaryl are optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy;

A is selected from the group consisting of , , , , , , and .

L^2 is selected from the group consisting of a bond, $-(CH_2)_n-$, $-C(O)NH(CH_2)_n-$, , , $-\frac{1}{2}-(CH_2)_m-\frac{1}{2}$, $-\frac{1}{2}-(CH_2)_m-\frac{1}{2}-(CH_2)_n-\frac{1}{2}$, $-[O(CH_2CH_2)]_n-$, $-[O(C_1-C_4 \text{ alkylene})]_n-$, $-[O(CH_2CH_2)]_n-OCH_2CH_2CF_2-$, $-C(O)NHCH_2CH_2-[O(CH_2CH_2)]_m-$; and $-C(O)NHCH_2CH_2-[O(CH_2CH_2)]_m-OCH_2CH_2CF_2-$;

m is an integer from zero to four;


n is an integer from one to four;

wherein the compound is substituted with at least one A; and


when X is $-CH_3$; L^1 is $-CH_2-$; Y is aryl substituted with A; and L^2 is $-CH_2-$; then A is not

$-L^2-COOH$, except when R^{1a} comprises $-COOH$ or $-SO_2CH_3$; and

when X is $-CH_3$; L^1 is $-CH_2-$; Y is aryl substituted with A; L^2 is $-CH_2-$, $-O-(CH_2)_2-$


$O(CH_2)_2-$, or $-O-(CH_2)_2-O(CH_2)_2(CF_2)-$; and A is ; then A and L^1 are not in a para position with respect to each other.

2. A compound having the structure of Formula (1), or a pharmaceutically acceptable salt thereof,




wherein

R^{1a} is selected from the group consisting of H, C_1-C_4 alkyl, $-NH_2$, $-NHAc$,

$-COOH$, $-SO_2CH_3$, $-SCH_3$, $-OCH_3$, and ,

wherein the alkyl is optionally substituted with $-OH$, $-NH_2$, $-NHAc$, $-COOH$, $-SO_2CH_3$,









$-SCH_3$, $-OCH_3$, or ;

R^{1b} is C_2-C_5 alkyl;

X is selected from the group consisting of H and C₁-C₄ alkyl, wherein the alkyl is optionally substituted with A, -OH, or -C(CH₃)₂OH;

L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , -O-, -S-, -SO₂-, -NH-, and -CH₂CH₂-;

Y is selected from the group consisting of C₁-C₃ alkyl, aryl, and heteroaryl, wherein the alkyl, aryl, and heteroaryl are optionally substituted with 1-5 substituents that are independently selected from A, C₁-C₃ alkyl, and C₁-C₃ alkoxy;

A is selected from the group consisting of , , , , , , , and .

L² is selected from the group consisting of a bond, -(CH₂)_n-, -C(O)NH(CH₂)_n-, , , , -(CH₂)_m-, -(CH₂)_n-, -[O(CH₂CH₂)]_n-, -[O(C₁-C₄ alkylene)]-, -[O(CH₂CH₂)]_n-OCH₂CH₂CF₂-, -C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-, and -C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-OCH₂CH₂CF₂-;


m is an integer from zero to four; and

n is an integer from one to four; and

wherein the compound is substituted with at least one A; and

when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; and L² is -CH₂-; then A is not -L²-COOH, except when R^{1a} comprises -COOH or -SO₂CH₃; and

when X is -CH₃; L¹ is -CH₂-; Y is aryl substituted with A; L² is -CH₂-, -O-(CH₂)₂-

O(CH₂)₂-, or -O-(CH₂)₂-O(CH₂)₂(CF₂)₂-; and A is ; then A and L¹ are not in a para position with respect to each other.


3. The compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein R^{1b} is -(CH₂)₂CH₃.

4. The compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein R^{1b} is $-(CH_2)_3CH_3$.

5. The compound of any one of claims 1-4, or a pharmaceutically acceptable salt thereof, wherein R^{1a} is C_1 - C_4 alkyl, optionally substituted with $-OH$, $-OCH_3$, $-SCH_3$, or $-SO_2CH_3$.


6. The compound of any one of claims 1-5, or a pharmaceutically acceptable salt thereof, wherein R^{1a} is $-CH_2C(CH_3)_2OH$.

7. The compound of any one of claims 1-6, or a pharmaceutically acceptable salt thereof, wherein




8. The compound of any one of claims 1-4, or a pharmaceutically acceptable salt thereof, wherein R^{1a} is C_1 - C_4 alkyl, optionally substituted with $-COOH$.

9. The compound of any one of claims 1-4 and 8, or a pharmaceutically



10. The compound of any one of claims 1-4, or a pharmaceutically acceptable salt thereof, wherein R^{1a} is H.

11. The compound of any one of claims 1-10, or a pharmaceutically acceptable salt thereof, wherein X is C_1 - C_4 alkyl, wherein the alkyl is substituted with A.

12. The compound of any one of claims 1-11, or a pharmaceutically acceptable salt thereof, wherein X is C₁-C₄ alkyl, wherein the alkyl is substituted with , wherein L² is a bond.

13. The compound of any one of claims 1-10, or a pharmaceutically acceptable salt thereof, wherein X is CH₃.

14. The compound of any one of claims 1-10, or a pharmaceutically acceptable salt thereof, wherein X is H.


15. The compound of any one of claims 1-14, or a pharmaceutically acceptable salt thereof, wherein L¹ is -CH₂-, -CH₂CH₂-, -O-, or -S-.


16. The compound of any one of claims 1-15, or a pharmaceutically acceptable salt thereof, wherein L¹ is -CH₂-.

17. The compound of any one of claims 1-16, or a pharmaceutically acceptable salt thereof, wherein Y is C₁-C₃ alkyl or aryl.

18. The compound of any one of claims 1-17, or a pharmaceutically acceptable salt thereof, wherein Y is aryl, wherein the aryl is substituted with C₁-C₃ alkoxy.

19. The compound of any one of claims 1-18, or a pharmaceutically acceptable salt thereof, wherein Y is aryl, wherein the aryl is substituted with A.


20. The compound of any one of claims 1-19, or a pharmaceutically acceptable salt thereof, wherein A is .

21. The compound of any one of claims 1-19, or a pharmaceutically acceptable salt thereof, wherein A is .


22. The compound of any one of claims 1-19, or a pharmaceutically acceptable




23. The compound of any one of claims 1-19, or a pharmaceutically acceptable




24. The compound of any one of claims 1-19, or a pharmaceutically acceptable



25. The compound of any one of claims 1-19, or a pharmaceutically acceptable




26. The compound of any one of claims 1-19, or a pharmaceutically acceptable



27. The compound of any one of claims 1-26, or a pharmaceutically acceptable salt thereof, wherein L^2 is $-(CH_2)_n-$.

28. The compound of any one of claims 1-27, or a pharmaceutically acceptable salt thereof, wherein n is one or two.


29. The compound of any one of claims 1-26, or a pharmaceutically acceptable



30. The compound of any one of claims 1-26, or a pharmaceutically acceptable salt thereof, wherein L^2 is .

31. The compound of any one of claims 1-26, or a pharmaceutically acceptable salt thereof, wherein L^2 is $-C(O)NH(CH_2)_n-$.








32. A compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1a),



wherein

X is H or CH₃


L^1 is selected from the group consisting of a bond, $-CH_2-$, $-O-$, $-S-$, $-CF_2-$, , , and $-CH_2CH_2-$;

A is selected from the group consisting of , , , , , , and .

L^2 is selected from the group consisting of a bond, $-CH_2-$, $-CH_2CH_2-$, , , $-C(O)NH(CH_2)_n-$, $-[O(CH_2CH_2)]_n-$, $-[O(C_1-C_4 \text{ alkylene})]_n-$, and $-[O(CH_2CH_2)]_n-OCH_2CH_2CF_2-$; $-C(O)NHCH_2CH_2-[O(CH_2CH_2)]_m-$; $C(O)NHCH_2CH_2-[O(CH_2CH_2)]_m-OCH_2CH_2CF_2-$; and


R^3 is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy.


33. A compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1a),



wherein


X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$;

A^{1a} is selected from the group consisting of , , , ,



L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, , , and $-\text{CH}_2\text{CH}_2-$;


A is selected from the group consisting of , , ,



L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, , , $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1-\text{C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$; $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-$; $\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and

R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.







34. A compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1b),



wherein

X is H or CH₃;


L¹ is selected from the group consisting of a bond, -CH₂-, -O-, -S-, -CF₂-, , , and -CH₂CH₂-;

A is selected from the group consisting of , , , , , and , and ;

L² is selected from the group consisting of a bond, -CH₂-, -CH₂CH₂-, , , -C(O)NH(CH₂)_n-, -[O(CH₂CH₂)]_n-, -[O(C₁-C₄ alkylene)]-, and -[O(CH₂CH₂)]_n-OCH₂CH₂CF₂-; -C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-, C(O)NHCH₂CH₂-[O(CH₂CH₂)]_m-OCH₂CH₂CF₂-; and

R³ is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy.


35. A compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1b),



wherein


X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$;

A^{1a} is selected from the group consisting of $-\text{C}(=\text{O})\text{OH}$, $-\text{P}(\text{OH})_3$, $-\text{SO}_3\text{H}$, $-\text{N}=\text{N}-\text{NH}$,



L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, $-\text{C}(\text{CH}_3)_2-$, $-\text{C}(\text{CH}_2\text{CH}_2)_2-$, and $-\text{CH}_2\text{CH}_2-$;


A is selected from the group consisting of $-\text{L}^2-\text{C}(=\text{O})\text{OH}$, $-\text{L}^2-\text{P}(\text{OH})_3$, $-\text{L}^2-\text{SO}_3\text{H}$,



L^2 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CH}_2\text{CH}_2-$, $-\text{C}(\text{CH}_3)_2-$, $-\text{C}(\text{CH}_2\text{CH}_2)_2-$, $-\text{C}(\text{O})\text{NH}(\text{CH}_2)_n-$, $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-$, $-\text{[O}(\text{C}_1-\text{C}_4 \text{ alkylene})]-$, and $-\text{[O}(\text{CH}_2\text{CH}_2)]_n-\text{OCH}_2\text{CH}_2\text{CF}_2-$; $-\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-$; $\text{C}(\text{O})\text{NHCH}_2\text{CH}_2-\text{[O}(\text{CH}_2\text{CH}_2)]_m-\text{OCH}_2\text{CH}_2\text{CF}_2-$; and

R^3 is H, C_1-C_3 alkyl, or C_1-C_3 alkoxy.


36. A compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1c),




wherein

X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$,

A^{1a} is selected from the group consisting of ,




L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{O}-$, $-\text{S}-$, $-\text{CF}_2-$, ,




R^3 is H, C_1 - C_3 alkyl, or C_1 - C_3 alkoxy; and

R^4 is H or C_1 - C_3 alkoxy.

37. A compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1d),




X is $-\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2-\text{A}^{1a}$, $-\text{CH}_2\text{CH}_2\text{CH}_2-\text{A}^{1a}$, or $-\text{CH}_2\text{C}(\text{CH}_3)_2-\text{A}^{1a}$;



L¹ is selected from the group consisting of a bond, -CH₂-, -CF₂-, , , -O-, and -CH₂CH₂-, -S-; and


Y is H or C₁-C₃ alkyl.

38. A compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1e),




wherein R³ is H, C₁-C₃ alkyl, or C₁-C₃ alkoxy.

39. A compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1f),




40. A compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1g),




wherein R^3 is H, C_1 - C_3 alkyl, or C_1 - C_3 alkoxy.

41. A compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1h),



wherein R^4 is H or C_1 - C_3 alkoxy.


42. A compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1i),




wherein

L^1 is selected from the group consisting of a bond, $-\text{CH}_2-$, $-\text{CF}_2-$, , , $-\text{O}-$, $-\text{CH}_2\text{CH}_2-$, and $-\text{S}-$; and
 Y is H or C_1 - C_3 alkyl.


43. A compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1j),



44. A compound of claim 1 or 2, or a pharmaceutically acceptable salt thereof, wherein the compound of Formula (1) is a compound of Formula (1k),














































45. A compound, or a pharmaceutically acceptable salt, selected from the group consisting of:
















































Compound	-X	-L ¹ -Y	$\begin{matrix} R^{1a} \\ \\ \text{---} \\ \\ R^{1b} \end{matrix}$	Compound
1	-CH ₃		-(CH ₂) ₃ CH ₃	
2	-CH ₃		-(CH ₂) ₃ CH ₃	
3	-CH ₃		-(CH ₂) ₃ CH ₃	



























Compound	-X	-L ¹ -Y	$\begin{array}{c} R^{1a} \\ \\ \text{---} R^{1b} \end{array}$	Compound
4	-CH ₃		-(CH ₂) ₃ CH ₃	
5	-CH ₃		-(CH ₂) ₃ CH ₃	
6	-CH ₃		-(CH ₂) ₃ CH ₃	
7	-CH ₃		-(CH ₂) ₃ CH ₃	
8	-CH ₃			
9	-CH ₃		-(CH ₂) ₃ CH ₃	
10	-CH ₃		-(CH ₂) ₃ CH ₃	
11	-CH ₃		-(CH ₂) ₃ CH ₃	
12	-CH ₃		-(CH ₂) ₃ CH ₃	

Compound	-X	-L ¹ -Y	R ^{1a} 	Compound
13	-CH ₃		-(CH ₂) ₃ CH ₃	
14	-CH ₃		-(CH ₂) ₃ CH ₃	
15			-(CH ₂) ₃ CH ₃	
16	-CH ₃		-(CH ₂) ₃ CH ₃	
17	-CH ₃		-(CH ₂) ₃ CH ₃	
18	-CH ₃			
19			-(CH ₂) ₃ CH ₃	
20	-CH ₃			

Compound	-X	-L ¹ -Y	R ^{1a} 	Compound
20C	-CH ₃			
21		-OCH ₃		
22	-CH ₃			
23	-CH ₃			
24	-CH ₃		-(CH ₂) ₃ CH ₃	
25	-CH ₃		-(CH ₂) ₃ CH ₃	
26	-CH ₃		-(CH ₂) ₄ CH ₃	
27				

Compound	-X	-L ¹ -Y	R ^{1a} 	Compound
28	-CH ₃		-(CH ₂) ₃ CH ₃	
29				
30	-CH ₃		-(CH ₂) ₃ CH ₃	
31	-CH ₃		-(CH ₂) ₃ CH ₃	
32	-CH ₃			
33	-CH ₃			
34	-CH ₃			
35	-CH ₃			

Compound	-X	-L ¹ -Y	R ^{1a} 	Compound
36	-CH ₃			
37			-(CH ₂) ₃ CH ₃	
38				
39	-CH ₃			
40			-(CH ₂) ₃ CH ₃	
41	-CH ₃			
42	-CH ₃			

Compound	-X	-L ¹ -Y	R ^{1a} 	Compound
43			-(CH ₂) ₃ CH ₃	
44	-CH ₃			
45				
46				
47			-(CH ₂) ₃ CH ₃	
48				
49				

Compound	-X	-L ¹ -Y	R ^{1a} R ^{1b}	Compound
50				
51	-CH ₃			

46. A pharmaceutical composition comprising a compound of any one of claims 1 to 45, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier.

47. A method of treating a condition associated with TLR7 modulation in a subject in need thereof, comprising administering to the subject an effective amount of a compound of any one of claims 1 to 45, or a pharmaceutically acceptable salt thereof.

48. The method of claim 47, wherein the condition is viral infection or cancer.

49. The method of claim 47 or 48, wherein the administration is oral, intravenous, subcutaneous, intramuscular, intratumoral, intradermal, intranasal, inhaled, intravesicle, topical, sublingual, buccal, intrarectal, intrathecal, intracranial, or other forms of local delivery.

50. A compound of any one of claims 1 to 45, or a pharmaceutically acceptable salt thereof, for use as a medicament.

51. A compound of any one of claims 1 to 45, or a pharmaceutically acceptable salt thereof, for use in treating a condition associated with TLR7 modulation.

52. The compound of claim 51, wherein the condition is viral infection or cancer.

53. Use of a compound of any one of claims 1 to 45, or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for treating a condition associated with TLR7 modulation.
54. The use of claim 53, wherein the condition is viral infection or cancer.
55. A pharmaceutical composition of claim 46, further comprising at least one or more additional therapeutic agents.
56. The pharmaceutical composition of claim 55, wherein the at least one or more additional therapeutic agent is antiviral nucleoside.
57. The pharmaceutical composition of claim 55, wherein the at least one or more additional therapeutic agent is PD-1 antibody or PD-L1 antibody.
58. A method of treating HBV in a subject in need thereof, comprising administering a compound of any one of claims 1 to 45, or a pharmaceutically acceptable salt thereof, in combination with an antiviral nucleoside.
59. A method of treating cancer in a subject in need thereof, comprising administering a compound of any one of claims 1 to 45, or a pharmaceutically acceptable salt thereof, in combination with a PD-1 antibody or PD-L1 antibody.