CLAIMS

1. A method of producing a reprogrammed cell, said method comprising exposing a population of human cells enriched for Stro-1^{bright} multipotential cells and/or progeny cells thereof to potency-determining factors under conditions sufficient to reprogram the cells, in which the potency-determining factors comprise:

(i) an Oct4 protein, a functional fragment of an Oct4 protein, or a nucleic acid encoding the Oct4 protein or the functional fragment of the Oct4 protein; and

(ii) a Sox2 protein, a functional fragment of the Sox2 protein, or a nucleic acid encoding the Sox2 protein or the functional fragment of the Sox2 protein; and in which the cell population is derived from adipose tissue or dental pulp.

2. A method according to Claim 1 additionally comprising culturing the exposed cell population to produce reprogrammed cells.

3. A method according to Claim 1 or 2 additionally comprising isolating the reprogrammed cells.

4. A method according to any preceding claim, in which the potency-determining factors further comprise one or more potency determining factors selected from the group consisting of Klf4, Nanog, Lin28, c-Myc, bFGF, SCF, TERT, SV40 large T antigen, HPV16E6, HPV16E7, Bmil, Fbx15, Eras, ECAT15-2, Tcl1, β-catenin, ECAT1, ESG1, Dnmt3L, ECAT8, Gdf3, Sox15, ECAT15-1, Fthl17, Sal14, Rex1, UTF1, Stella, Stat3 and Grb2.

5. A method according to any preceding claim, in which the potency-determining factors are selected from the group consisting of:

(i) a combination of Oct4, Sox2 and at least one of Nanog and Lin28;

(ii) a combination of Oct4, Klf4 and c-Myc;

(iii) a combination of Oct4, Sox2 and Klf4;

(iv) a combination of Oct4, Sox2, Klf4 and c-Myc;

(v) a combination of Oct4, Sox2, Nanog and Lin28; and

(vi) a combination of Oct4, Sox2, Klf4, c-Myc, Nanog and Lin28.

6. A method according to Claim 1, comprising introducing into the cell population one or more nucleic acids comprising a sequence encoding one or more of the potency-determining factors operably linked to a promoter.

7. A method according to Claim 6, comprising administering a plurality of nucleic acids each comprising a sequence encoding a distinct potency-determining factor operably linked to a promoter.

8. A method according to Claim 6 or 7, in which the nucleic acid(s) are within one or more vector(s).

9. A method according to Claim 8, in which the vector(s) is(are) a viral vector(s).

10. A method according to Claim 6, in which the nucleic acid(s) do(es) not integrate into the genome of the Stro-1^{bright} multipotential cells and/or progeny cells thereof.

11. A method according to Claim 1 or 2, in which the reprogrammed cells comprise pluripotent cells.

12. A method according to Claim 1 or 2, in which the reprogrammed cells (i) express a cell marker selected from the group consisting of Oct-4, Nanog, SSEA3, SSEA4, Tra-1-60 and Tra-1-81; (ii) exhibit morphology characteristic of pluripotent cells; and/or (iii) form teratomas when introduced into an immunocompromised animal.

13. A method according to Claim 1 or 2, in which the potency-determining factors consist of Oct4, Sox2, Klf4, and cMyc, or nucleic acids encoding the factors.

14. A method according to Claim 1 or 2, further comprising differentiating the reprogrammed cells into cells with less potency than the reprogrammed cells.