

CHECKPOINT BLOCKADE AND MICROSATELLITE INSTABILITY

- [01] This invention was made with government support under CA43460 and CA62924 awarded by the National Institutes of Health. The government has certain rights in the invention.

TECHNICAL FIELD OF THE INVENTION

- [02] This invention is related to the area of cancer. In particular, it relates to cancer therapy.

BACKGROUND OF THE INVENTION

- [03] Microsatellite instability (MSI) is the accumulation of sequencing errors in microsatellites. This occurs in tumors with deficiency in DNA mismatch repair. MSI is present in Lynch Syndrome which is an inherited cancer syndrome that predisposes patients to colon, endometrial, gastric cancer, ovarian, small intestine, liver, hepatobiliary, upper urinary tract, brain, and prostate cancer. MSI is also present in 10-20% of sporadic colorectal, gastric, prostate, lung, ampullary, and endometrial cancers. Between 0.3% and 13% of pancreatic cancers are reported to be MSI as well.
- [04] The importance of intact immune surveillance in controlling outgrowth of neoplastic transformation has been known for decades. Accumulating evidence shows a correlation between tumor-infiltrating lymphocytes (TILs) in cancer tissue and favorable prognosis in various malignancies. In particular, the presence of CD8+ T-cells and the ratio of CD8+ effector T-cells / FoxP3+ regulatory T-cells seems to correlate with improved prognosis and long-term survival in solid malignancies such as ovarian, colorectal and pancreatic cancer, hepatocellular carcinoma, malignant MEL and RCC. TILs can be expanded *ex vivo* and re-infused, inducing durable objective tumor responses in cancers such as melanoma.
- [05] The PD-1 receptor-ligand interaction is a major pathway hijacked by tumors to suppress immune control. The normal function of PD-1, expressed on the cell surface

of activated T-cells under healthy conditions, is to down-modulate unwanted or excessive immune responses, including autoimmune reactions. The ligands for PD-1 (PD-L1 and PD-L2) are constitutively expressed or can be induced in various tumors. Binding of either PD-1 ligand to PD-1 inhibits T-cell activation triggered through the T-cell receptor. PD-L1 is expressed at low levels on various non-hematopoietic tissues, most notably on vascular endothelium, whereas PD-L2 protein is only detectably expressed on antigen-presenting cells found in lymphoid tissue or chronic inflammatory environments. PD-L2 is thought to control immune T-cell activation in lymphoid organs, whereas PD-L1 serves to dampen unwarranted T-cell function in peripheral tissues. Although healthy organs express little (if any) PD-L1, a variety of cancers were demonstrated to express abundant levels of this T-cell inhibitor. High expression of PD-L1 on tumor cells (and to a lesser extent of PD-L2) has been found to correlate with poor prognosis and survival in various cancer types, including renal cell carcinoma (RCC), pancreatic carcinoma, hepatocellular carcinoma, ovarian carcinoma and non-small cell lung cancer (NSCLC). Furthermore, PD-1 has been suggested to regulate tumor-specific T cell expansion in patients with malignant MEL. The observed correlation of clinical prognosis with PD-L1 expression in multiple cancers suggests that the PD-1/PD-L1 pathway plays a critical role in tumor immune evasion and should be considered as an attractive target for therapeutic intervention.

- [06] Blockade of immune checkpoints such as cytotoxic T-lymphocyte antigen-4 (CTLA-4) and programmed death-1 (PD-1) is showing promise in patients with cancer. CTLA-4 and PD-1 are upregulated on activated T cells and provide inhibitory signals to T cells undergoing activation. Inhibitory antibodies directed at these receptors have been shown to break immune tolerance and promote anti-tumor immunity. MK-3475 is a humanized monoclonal IgG4 antibody against PD-1 and is showing activity in multiple tumor types including melanoma and non-small cell lung cancer (NSCLC). Previously, activity of a different PD-1 blocking antibody, BMS-936558, a fully humanized monoclonal IgG4 antibody, also showed activity in melanoma, NSCLC, and a complete response in a single patient with colorectal cancer.
- [07] MK-3475 (previously known as SCH 900475) is a potent and highly-selective humanized mAb of the IgG4/kappa isotype designed to directly block the interaction

between PD-1 and its ligands, PD-L1 and PD-L2. MK-3475 contains the S228P stabilizing mutation and has no antibody-dependent cell-mediated cytotoxicity (ADCC) or complement-dependent cytotoxicity (CDC) activity. MK-3475 strongly enhances T lymphocyte immune responses in cultured blood cells from healthy human donors, cancer patients, and primates. In T- cell activation assays using human donor blood cells, the EC50 was in the range of 0.1 to 0.3 nM. MK-3475 also modulates the level of interleukin-2 (IL-2), tumor necrosis factor alpha (TNF α), interferon gamma (IFN γ), and other cytokines. The antibody potentiates existing immune responses only in the presence of antigen and does not nonspecifically activate T- cells.

[08] The programmed death 1 (PD-1) pathway is a negative feedback system repressing Th1 cytotoxic immune responses that, if unregulated, could damage the host¹⁻³. It is upregulated in many tumors and their surrounding microenvironment. Blockade of this pathway with antibodies to PD-1 or its ligands has led to remarkable clinical responses in some patients with many different cancer types, including melanomas, non-small cell lung cancer, renal cell carcinoma, bladder cancer and Hodgkin's lymphoma⁴⁻¹⁰. The expression of ligands to PD-1 (PD-L1 or PD-L2) on the surface of tumor cells or immune cells is important but not a definitive predictive biomarker for response to PD-1 blockade^{4,6-8,11}.

[09] We were intrigued that, in reports of the effects of PD-1 blockade in human tumors, only one of 33 colorectal cancer (CRC) patients responded to this treatment, in contrast to substantial fractions of patients with melanomas, renal cell cancers, and lung tumors.^{10,12} What was different about this single patient? We hypothesized that this patient had MMR-deficiency, because MMR-deficiency occurs in a small fraction of advanced CRCs,^{13,14} somatic mutations found in tumors can be recognized by the patient's own immune system,¹⁵ and MMR-deficient cancers have 10- to 100-fold more somatic mutations than MMR-proficient CRC.¹⁶⁻¹⁸ Moreover, MMR-deficient cancers contain prominent lymphocyte infiltrates, consistent with an immune response¹⁹⁻²². And two of the tumor types that were most responsive to PD-1 blockade in a study by Topalian et al.¹⁰ had high numbers of somatic mutations as a result of exposure to cigarette smoke (lung cancers) or UV radiation (melanomas)^{23,24}. Our hypothesis was correct: the tumor of the single CRC patient who responded to PD-1

blockade was MMR-deficient²⁵. We therefore hypothesized that MMR-deficient tumors are more responsive to PD-1 blockade than are MMR-proficient tumors.

- [10] To test this hypothesis, we initiated a phase 2 clinical trial to evaluate immune checkpoint blockade in patients whose tumors had or did not have MMR-deficiency. Since MMR deficiency in tumors arises through two routes²⁶⁻²⁸, we recruited patients with Hereditary Non-Polyposis Colorectal Cancer (HNPCC, also known as Lynch Syndrome), which results from an inherited germline defect in one of four MMR genes followed by a second inactivating somatic change in the remaining wild-type allele. We also recruited patients with sporadic MMR-deficient tumors, where both alleles of a MMR gene are inactivated by somatic mutations or by epigenetic silencing²⁹. In either case, the neoplasms that arise harbor hundreds or thousands of mutations^{16,18}.
- [11] There is a continuing need in the art to improve cancer treatments so that the lives of patients are not curtailed and so that the quality of life is not diminished.

SUMMARY OF THE INVENTION

- [12] According to one embodiment of the invention a method of treating a cancer patient is provided. The cancer patient has a high mutational burden, such as found in microsatellite instable cancer (MSI). An immune checkpoint inhibitory antibody is administered to the cancer patient.
- [13] According to another embodiment of the invention a method of treating a cancer patient is provided. A sample from a cancer patient is tested for one or more microsatellite markers selected from the group consisting of BAT-25, BAT-26, MON0-27, NR-21, NR-24, Penta C, and Penta D, and determined to have microsatellite instability. The cancer is selected from the group consisting of: colon, gastric, endometrial, cholangiocarcinoma, pancreatic, and prostate cancers. An anti-PD-1 antibody is administered to the cancer patient.

- [14] According to another embodiment of the invention a method is provided for categorizing a tumor of a human. A sample from the human is tested to evaluate stability of one or more microsatellite markers. Microsatellite instability is determined in the sample. The tumor is identified as a good candidate for treatment with an immune checkpoint inhibitory antibody.
- [15] According to yet another embodiment of the invention a method is provided for categorizing a tumor of a human. A sample from the human is tested to evaluate stability of one or more microsatellite markers. Microsatellite stability in the sample is determined. The tumor is identified as a bad candidate for treatment with an immune checkpoint inhibitory antibody.
- [16] These and other embodiments which will be apparent to those of skill in the art upon reading the specification provide the art with methods for treating microsatellite instable cancers.

BRIEF DESCRIPTION OF THE DRAWINGS

- [17] Figs. 1A-1B. Clinical Responses to pembrolizumab. (Fig. 1A) Biochemical Responses. Serum protein biomarker levels were measured with each cycle and the values represent percent change from baseline. Patients were included if baseline tumor marker values were greater than the upper limit of normal. CA-125 was used for a patient with endometrial cancer; CA19-9 was used for one cholangiocarcinoma and one ampullary cancer; and CEA was used for all other patients. Green, red, and black lines represent patients with MMR-deficient CRCs, MMR-proficient CRCs, and MMR-deficient non-CRC, respectively. (Fig. 1B) Radiographic responses. Tumor responses were measured at regular intervals and values show the best fractional change of the sum of longest diameters (SLD) from the baseline measurements of each measurable tumor.
- [18] Figs. 2A-2D. Clinical benefit to pembrolizumab according to MMR status. Kaplan-Meier curves are shown for (Fig. 2A) progression-free survival in the colorectal cancer cohorts, (Fig. 2B) overall survival in the colorectal cancer cohorts, (Fig. 2C)

progression-free survival of patients with MMR-deficient cancers other than colorectal (median PFS=5.4 months; 95% CI, 3% to not estimable), and (Fig. 2D) overall survival of patients with MMR-deficient cancers other than colorectal. In both cohorts with MMR-deficient tumors (CRC and non-CRC), median overall survival was not reached. Patients in the cohort with MMR-proficient cancers had a median PFS of 2.2 months (95% CI 1.4 to 2.8%) and a median OS of 5.0 months (95% CI 3.0 to not estimable).

- [19] Fig. 3 (Figure S2.) Spider plot of radiographic response. Tumor responses were measured at regular intervals and values show percent change of the sum of longest diameters (SLD) from the baseline measurements of each measurable tumor. Patients were only included if baseline and on study treatment scans were available. Green and red represent patients with MMR-deficient and proficient CRCs, respectively. Blue represents patients with MMR-deficient cancers other than CRC.
- [20] Figs. 4A-4B (Figure S3). MMR-proficient and deficient CRCs have comparable time on treatment and duration of metastatic disease prior to study enrollment. Kaplan-Meier estimates of (Fig. 4A) time on therapy immediately prior to study enrollment (HR 0.81, 95% CI 0.38 to 1.752, $p=.60$) and (Fig. 4B) duration of metastatic disease prior to enrollment (HR 1.13, 95% CI 0.49 to 2.62, $p=.78$) on this pembrolizumab study were comparable between the MMR-deficient and proficient CRC cohorts. The short duration on prior therapy is expected in a treatment refractory CRC population.
- [21] Fig. 5 (Figure S4.) Waterfall plot of biochemical response. Serum protein biomarker levels were measured with each cycle and the values represent best percent change from baseline. Patients were included if baseline tumor marker values were greater than the upper limit of normal. CA-125 was used for a patient with endometrial cancer; CA19-9 was used for 1 cholangiocarcinoma and 1 ampullary cancer; and CEA was used for all other patients. Green and red represent patients with MMR-deficient and proficient CRCs, respectively. Blue represents patients with MMR-deficient cancers other than CRC.

- [22] Figs. 6A-6B (Figure S5.) Somatic mutations in MMR-deficient and proficient tumors. Total somatic mutations per tumor identified by exome sequencing of tumor and matched normal DNA (Fig. 6A) and correlation with objective responses (Fig. 6B) (non-parametric Wilcoxon test , $p=0.007$ and Jonckheere-Terpstra test for trend, $p=0.02$).
- [23] Fig. 7 (Figure S6). Immunohistochemistry of CD8 and PD-L1 Expression. The invasive front (yellow dashed line) from a MMR-deficient CRC (subject #16, top) and MMR-proficient CRC (subject #3, bottom). The yellow dashed line separates tumor (T) and normal (N) tissue. There is marked expression of PD-L1 (blue arrows) and CD8 (brown dots) in the MMR-deficient tumor (top panels) patient while there is very little expression of either marker in the MMR-proficient tumor (bottom panels). Representative images of tumor infiltrating lymphocytes (TIL) in another MMR-deficient CRC (subject #19, top) and MMR-proficient CRC (subject #3, bottom) immunolabeled with an antibody to CD8 (brown dots). Note the infiltration of CD8 cells in the MMR-deficient tumor. Invasive front original magnification 10x and TIL 20x.
- [24] Fig. 8 (Figure S7.) CD8 and PD-L1 Expression in the MMR-deficient and MMR-proficient tumor microenvironment. T cell density units are cells/mm² of tumor. Invasive front refers to the immune cells (TILs and macrophages) at the junction of the tumor and normal tissue. P-values obtained using an unpaired t-test.
- [25] Fig. 9 (Figure S8.) CD8 expression and clinical benefit to pembrolizumab. Correlation between the intratumoral CD8⁺ T cell density (cells/mm²) and objective response (Jonckheere-Terpstra test for trend, $p=0.02$).
- [26] Fig. 10 (Table S1.) Comparison of immune-related and RECIST response criteria (adapted from Wolchok et al. Clin Can Res 2009;15:7412-20.)

- [27] Fig. 11 (Table S2.) Immune-Related response to treatment
- [28] Fig. 12 (Table S4.) Correlation of total somatic mutations and mutation associated neoantigens (MANA) with clinical outcomes
- [29] Fig. 13 (Table S5.) Correlation of immune markers with clinical outcome

DETAILED DESCRIPTION OF THE INVENTION

- [30] The inventors have found that immune checkpoint inhibitors work best in tumors with high mutation burdens. Furthermore, tumors deficient in mismatch repair are particularly susceptible to a particular form of immunotherapy because this phenotype results in ongoing accumulation of mutations at a high frequency. The inventors have developed a treatment for cancer patients that display the microsatellite instability phenotype or other high mutational burden. The treatment involves an inhibitory antibody for an immune checkpoint. Such checkpoints include PD-1, IDO, CTLA-4, PD-L1, and LAG-3. Other immune checkpoints can be used as well. Antibodies can be administered by any means that is convenient, including but not limited to intravenous infusion, oral administration, subcutaneous administration, sublingual administration, ocular administration, nasal administration, etc.
- [31] Microsatellite instability (MSI) tumors are deficient in DNA mismatch repair which leads to a high rate of spontaneous mutations and the potential for the expression of neo-antigens. Furthermore, similar to melanoma, in MSI positive colon cancers, there is often prominent lymphocyte infiltration. Any tumors that are MSI or otherwise high mutational burden may be treated according to the invention. They may be tested for the attribute of MSI according to any method known in the art, including but not limited that described in example 1 below. Any of one or more MSI markers can be tested to determine an MSI phenotype. Samples may be tested for high mutational burden by identifying tumors with at least 100, at least 200, at least 300, at least 400, at least 500, at least 600, at least 700, at least 800, at least 900, at least 1000, at least

1100, at least 1200, at least 1300, at least 1400, at least 1500, or at least 1600 mutations per tumor genome. High mutational burden means a large number of somatic mutations in the tumor relative to normal tissues of the individual. An average number of somatic mutations in a non-MSI tumor is about 70 somatic mutations.

- [32] Any type of tumor that displays the MSI phenotype or a high mutational burden may be tested and/or treated according to the invention. These include without limitation cancers of the colon, gastric, endometrial, cholangiocarcinoma, pancreatic, and prostate cancer. Tumors of the ampulla, biliary, brain, including glioma, breast, lung, skin, esophagus, liver, kidney, ovaries, sarcoma, uterus, cervix, bladder, testes, oral cavity, tongue, and small and large bowel may also be tested and/or treated.
- [33] Testing of MSI can be accomplished by any means known in the art. One or more of the following markers may be tested: five nearly monomorphic mononucleotide repeat markers (BAT-25, BAT-26, MON0-27, NR-21 and NR-24) and two highly polymorphic pentanucleotide repeat markers (Penta C and Penta D). In one commercial system which can be used, fluorescently labeled primers (marker panel) are used for co-amplification of all seven of the above named markers. Fragments are detected after amplification for assignment of genotype/phenotype.
- [34] Samples that can be tested for MSI include tumor tissue as well as body fluids that contain nucleic acids shed from tumors. Testing for tumor DNA in such tissues and body fluids is well known.
- [35] Types of antibodies which can be used include any that are developed for the immune checkpoint inhibitors. These can be monoclonal or polyclonal. They may be single chain fragments or other fragments of full antibodies, including those made by enzymatic cleavage or recombinant DNA techniques. They may be of any isotype, including but not limited to IgG, IgM, IgE. The antibodies may be of any species source, including human, goat, rabbit, mouse, cow, chimpanzee. The antibodies may be humanized or chimeric. The antibodies may be conjugated or engineered to be

attached to another moiety, whether a therapeutic molecule or a tracer molecule. The therapeutic molecule may be a toxin, for example.

- [36] The data from the small phase 2 trial of pembrolizumab to treat tumors with and without deficiency of MMR supports the hypothesis that MMR-deficient tumors are more responsive to PD-1 blockade than are MMR-proficient tumors. MMR-deficiency occurs in many cancers, including those of the colorectum, uterus, stomach, biliary tract, pancreas, ovary, prostate and small intestine^{18,34-42}. Patients with MMR-deficient tumors of these types also benefit from anti-PD-1 therapy, as may patients whose tumors contain other DNA repair deficiencies, such as those with mutations in *POLD*, *POLE*, or *MYH*.^{18,43,44}
- [37] The hypothesis that MMR-deficient tumors stimulate the immune system is not a new idea⁴⁵, and has been supported by the dense immune infiltration and Th1-associated cytokine-rich environment observed in MMR-deficient tumors.^{19-22,46} A recent study refined these classic observations by showing that the MMR-deficient tumor microenvironment strongly expressed several immune checkpoint ligands including PD-1, PD-L1, CTLA-4, LAG-3 and IDO, indicating that their active immune microenvironment is counterbalanced by immune inhibitory signals that resists tumor elimination⁴⁷. That the immune infiltrate associated with MMR-deficient carcinomas was directed at neoantigens was the most likely explanation for both the old and new findings. The correlation of higher mutational load and higher response rate to anti-CTLA-4 in melanoma⁴¹ and anti-PD-1 in lung cancer⁴⁸ provide further support for the idea that MANA recognition is an important component of the endogenous anti-tumor immune response.
- [38] Based on the results of the current and previous studies, we suggest that the greatly (>20-fold) increased number of mutation-associated neoantigens resulting from MMR deficiency (Fig.. 12 (Table S4); also available on line at New England Journal of Medicine; incorporated by reference herein) is the basis for the enhanced anti-PD-1 responsiveness of this genetically defined subset of cancers. Though our estimates for the number of mutation-associated neoantigens in tumors is based only on *in silico* predictions of binding-affinity, this suggestion is consistent with the observation that

MMR-proficient tumors have far less infiltration of lymphocytes than MMR-deficient tumors (Fig. 7 (S6), Fig. 8 (S7) and Fig. 13 (Table S5); available on line at New England Journal of Medicine; incorporated by reference herein). Recent studies^{49,50} show that only a tiny proportion of predicted neo-epitopes are actually presented on the cell surface with MHC and are targets of endogenous T cell responses. It seems likely, though that the number of predicted mutation-associated neoantigens is proportionate to the number of actual mutation-associated neoantigens, and tumors with a high number of actual mutation-associated neoantigens are more likely to stimulate the immune system to react against the tumor. Alternative mechanisms underlying the difference in anti-PD-1 responsiveness between MMR-deficient and MMR-proficient tumors should also be considered. For example, different signaling pathways activated in MMR-deficient and MMR-proficient tumors may result in differences in secretion of soluble factors that could result in differential activation of the PD-1 pathway within the tumor microenvironment²⁶⁻²⁸. Genetic differences could effect epigenetic differences that alter the expression of tumor-associated self-antigens that in turn could alter the antigenicity of the tumor. Experimental analyses of antigen-specific immune responses as well as changes in immune microenvironments should help to define the relative contribution of these factors to the striking responsiveness of MMR-deficient tumors to PD-1 antibodies.

- [39] Several notable observations were made during the course of this study. First, changes in serum protein biomarkers, like CEA, corresponded with clinical benefit after a single dose of therapy. Declines in CEA levels preceded objective radiographic evidence by several months; perhaps other biomarkers such as circulating tumor DNA (ctDNA) may also be beneficial as surrogate markers of early response.^{51,52} Second, our results suggest that the evaluation of tumor genomes can help guide immunotherapy. They support the view that the number and type of alterations may prove useful for judging the potential utility of immune checkpoint inhibitors, even in MMR-proficient cancers^{41,48,53} Most importantly, our results demonstrate a new approach for the treatment of a specific class of tumors based solely on genetic status: *i.e.*, without regard to underlying tumor type.

[40] The above disclosure generally describes the present invention. All references disclosed herein are expressly incorporated by reference. A more complete understanding can be obtained by reference to the following specific examples which are provided herein for purposes of illustration only, and are not intended to limit the scope of the invention.

EXAMPLE 1

MSI Testing

[41] MSI testing is already standardized and performed in CLIA-certified laboratories without need for assay development. Archived tumor samples or newly obtained biopsies will be used for determining MSI. MSI status will be performed locally by CLIA certified immunohistochemistry (IHC) or PCR based tests for eligibility. Evaluable patients will be confirmed using the MSI Analysis System from Promega at Johns Hopkins. This test will determine MSI status through the insertion or deletion of repeating units in the five nearly monomorphic mononucleotide repeat markers (BAT-25, BAT-26, MON0-27, NR-21 and NR-24). At least 2 MSI loci are required to be evaluable in Cohorts A and C. Patients may be assigned to a new cohort and/or replaced based on the Promega test results.

EXAMPLE 2

METHODS

[42] Patients

[43] Treatment-refractory progressive metastatic cancer patients for this phase 2 study were recruited from three participating centers (Table 1). Three cohorts were evaluated: Cohort A was composed of patients with MMR-deficient colorectal adenocarcinomas; Cohort B was composed of patients with MMR-proficient colorectal adenocarcinomas; and Cohort C was composed of patients with MMR-deficient cancers of types other than colorectal.

[44] Study Oversight

- [45] The protocol, which can be found at NEJM.org, was approved by each site's institutional review boards, and the study was conducted in accordance with the Declaration of Helsinki and the International Conference on Harmonization Guidelines for Good Clinical Practice. All the patients provided written informed consent before study entry. The principal investigator (D.L.) and study sponsor (L.A.D.) were responsible for oversight of the study. Merck donated the study drug, reviewed the final drafts of the protocol and of this manuscript. The clinical study was primarily funded through philanthropic support.
- [46] Study Design
- [47] This phase 2 trial was conducted using a Green-Dahlberg two-stage design and consisted of the three parallel cohorts described above. The study agent, pembrolizumab (Merck), was administered at 10 mg/kg intravenously every 14 days. Pembrolizumab is a humanized monoclonal anti-PD-1 antibody of the IgG4/kappa isotype that blocks the interaction between PD-1 and its ligands, PD-L1 and PD-L2.
- [48] Safety assessments were performed before each treatment. Assessments of total tumor burden via measurements of serum biomarkers were performed at the start of each cycle. Radiologic assessments were made at 12 weeks and every 8 weeks thereafter. Further details concerning the clinical protocol are provided in the Example 3.
- [49] Analysis of mismatch repair status
- [50] Tumors with genetic defects in MMR pathways are known to harbor thousands of somatic mutations, especially in regions of repetitive DNA known as microsatellites. The accumulation of mutations in these regions of the genome is termed microsatellite instability (MSI)²⁶⁻²⁸. MMR-status was assessed using the MSI Analysis System from Promega in tumors, through the evaluation of selected microsatellite sequences particularly prone to copying errors when MMR is compromised²⁶⁻²⁸. See Supplementary Appendix for additional details.
- [51] Genomic & Bioinformatic Analyses

[52] Primary tumor samples and matched normal peripheral-blood specimens were obtained from a subset of subjects with MMR-deficient and others with MMR-proficient carcinomas where sufficient tumor tissue was available for exome sequencing³⁰ and HLA haplotyping. To assess the potential for mutant peptide binding, somatic exome data combined with the individual patient's MHC class I HLA haplotype was applied to the an epitope prediction algorithm^{31,32}. This algorithm provided an estimate of the total number of mutation-associated neoantigens in each tumor. Additional details are provided in the Supplementary Appendix (available on line at New England Journal of Medicine; incorporated by reference herein).

[53] Statistical Analysis

[54] The primary endpoints for Cohorts A and B were immune-related objective response rate (irORR) and immune-related progression-free survival (irPFS) rate at 20 weeks assessed using immune-related response criteria (irRC)³³. The primary endpoint for Cohort C was irPFS rate at 20 weeks. Immune-related criteria (i.e, criteria used to evaluate immune-based therapies) are based on radiographic responses, and unlike RECIST criteria, capture extent of disease after disease progression; these criteria are defined and compared to RECIST v1.1 in Fig. 10 (Table S1). Response rate and PFS rate at 20 weeks were evaluated and reported in this study using RECIST v1.1 and irRC (Fig. 10 (Table S1)). PFS and overall survival was summarized by Kaplan-Meier method. Details of the hypothesis, the decision rules to reject the null hypotheses and early-stopping rules for efficacy and futility, and statistical methods are provided in the Supplementary Appendix.

EXAMPLE 3

SUPPLEMENTARY METHODS

[55] PATIENTS

[56] To be eligible for participation in this study, patients had to be at least 18 years of age, have histologically confirmed evidence of previously-treated, progressive carcinoma. All patients underwent MMR status testing prior to enrollment. All patients had at

least one measurable lesion as defined by the Response Evaluation Criteria in Solid Tumors (RECIST), version 1.1, an Eastern Cooperative Oncology Group (ECOG) performance-status score of 0 or 1, and adequate hematologic, hepatic, and renal function. Eligible patients with CRC must have received at least 2 prior cancer therapies and patients with other cancer types must have received at least 1 prior cancer therapy. Patients with untreated brain metastases, history of HIV, hepatitis B, hepatitis C, clinically significant ascites/effusions, or autoimmune disease were excluded.

[57] STUDY OVERSIGHT

[58] Initial drafts of the manuscript were prepared by a subset of the authors and all authors contributed to the final manuscript. All the authors made the decision to submit the manuscript for publication. The principal investigator and study sponsor vouch for the accuracy and completeness of the data reported as well as adherence to the protocol.

[59] HLA TYPING

[60] HLA-A, HLA-B and HLA-C Sequence Based Typing can be divided into three distinct steps, as described below. A generic, A*02 specific, B generic, B group specific, C generic and C*07 specific PCR and sequencing mixes were made in the JHU core facility. Celera's AlleleSEQR HLA-B Sequence Based Typing kit was used for B generic SBT. The HLA-A typing scheme is composed of two PCR reactions, A generic and A*02 specific. A generic amplicon encompasses partial exon 1- partial exon 5. A*02 amplicon encompasses partial intron 1 - partial exon 5. HLA-B typing scheme is composed of two PCR reactions, B generic and B group specific. The B generic PCR is a multiplexed reaction containing two PCR amplicons encompassing exon 2 – exon 3 and exon 4 – exon 7. B group specific amplicon encompasses partial intron 1 - partial exon 5. HLA-C typing scheme is composed of two PCR reactions, C generic and C*07 specific. C generic and C*07 specific amplicons encompasses exons 1 - 7.

- [61] The specificity of the HLA-A and B PCR employed AmpliTaq Gold DNA polymerase. The GeneAmp High Fidelity enzyme is used for the HLA-C and C*07 PCR mixes. This enzyme is a mix of two polymerases: AmpliTaq DNA polymerase (non-proofreading polymerase) and a proofreading polymerase. This enzyme mix is necessary to produce efficient and robust amplification of the larger full length HLA-C amplicon.
- [62] PCR product purification was performed using Exonuclease I and Shrimp Alkaline Phosphatase. The A generic and B generic amplicons were bi-directionally sequenced for exons 2,3,4. The C generic amplicon was bi-directionally sequenced for exons 2,3 and sequenced in a single direction for exons 1,4,5,6,7. A*02 specific, B group specific and C*07 specific amplicons were sequenced in a single direction for exons 2,3. All sequencing reactions were performed with Big Dye Terminator V1.1 from Applied Biosystems and sequenced with an ABI Prism 3500XL Genetic Analyzer. Conexio Genomic's "Assign SBT" allele assignment software was used to process the data files.
- [63] MISMATCH REPAIR STATUS TESTING^{1,2}
- [64] Six slides of tumor and normal (uninvolved lymph node or margin of resection) were cut (5 microns each), deparaffinized (xylene), and one stained with hematoxylin and eosin (H+E). A tumor area containing at least 20% neoplastic cells, designated by a board-certified Anatomic Pathologist was macrodissected using the Pinpoint DNA isolation system (Zymo Research, Irvine, CA), digested in proteinase K for 8 hours and DNA was isolated using a QIAamp DNA Mini Kit (Qiagen, Valencia, CA). MSI was assessed using the MSI Analysis System (Promega, Madison, WI), composed of 5 pseudomonomorphic mononucleotide repeats (BAT-25, BAT-26, NR-21, NR-24 and MONO-27) to detect MSI and 2-pentanucleotide repeat loci (PentaC and PentaD) to confirm identity between normal and tumor samples, per manufacturer's instructions. Following amplification of 50-100 ng DNA, the fluorescent PCR products were sized on an Applied Biosystems 3130xl capillary electrophoresis instrument (Invitrogen, Calsbad, CA). Pentanucleotide loci confirmed identity in all cases. Controls included water as a negative control and a mixture of 80% germline DNA with 20% MSI

cancer DNA as a positive control. The size in bases was determined for each microsatellite locus and tumors were designated as MSI if two or more mononucleotide loci varied in length compared to the germline DNA.

[65] SEQUENCING ANALYSIS

[66] Samples

[67] Samples provided as FFPE blocks or frozen tissue underwent pathological review to determine tumor cellularity. Tumors were macrodissected to remove contaminating normal tissue, resulting in samples containing >20% neoplastic cells. Matched normal samples were provided as blood, saliva or normal tissue obtained from surgery.

[68] Sample Preparation and Next-Generation Sequencing³

[69] Sample preparation, library construction, exome capture, next generation sequencing, and bioinformatics analyses of tumor and normal samples were performed at Personal Genome Diagnostics, Inc. (Baltimore, Maryland). In brief, DNA was extracted from frozen or formalin-fixed paraffin embedded (FFPE) tissue, along with matched blood or saliva samples using the Qiagen DNA FFPE tissue kit or Qiagen DNA blood mini kit (Qiagen, CA). Genomic DNA from tumor and normal samples were fragmented and used for Illumina TruSeq library construction (Illumina, San Diego, CA) according to the manufacturer's instructions or as previously described⁴. Briefly, 50 nanograms (ng) - 3 micrograms (μg) of genomic DNA in 100 microliters (μl) of TE was fragmented in a Covaris sonicator (Covaris, Woburn, MA) to a size of 150-450bp. To remove fragments smaller than 150bp, DNA was purified using Agencourt AMPure XP beads (Beckman Coulter, IN) in a ratio of 1.0 to 0.9 of PCR product to beads twice and washed using 70% ethanol per the manufacturer's instructions. Purified, fragmented DNA was mixed with 36 μl of H₂O, 10 μl of End Repair Reaction Buffer, 5 μl of End Repair Enzyme Mix (cat# E6050, NEB, Ipswich, MA). The 100 μl end-repair mixture was incubated at 20°C for 30 min, and purified using Agencourt AMPure XP beads (Beckman Coulter, IN) in a ratio of 1.0 to 1.25 of PCR

product to beads and washed using 70% ethanol per the manufacturer's instructions. To A-tail, 42 µl of end-repaired DNA was mixed with 5 µl of 10X dA Tailing Reaction Buffer and 3 µl of Klenow (exo-)(cat# E6053, NEB, Ipswich, MA). The 50 µl mixture was incubated at 37°C for 30 min and purified using Agencourt AMPure XP beads (Beckman Coulter, IN) in a ratio of 1.0 to 1.0 of PCR product to beads and washed using 70% ethanol per the manufacturer's instructions. For adaptor ligation, 25 µl of A-tailed DNA was mixed with 6.7 µl of H₂O, 3.3 µl of PE-adaptor (Illumina), 10 µl of 5X Ligation buffer and 5 µl of Quick T4 DNA ligase (cat# E6056, NEB, Ipswich, MA). The ligation mixture was incubated at 20°C for 15 min and purified using Agencourt AMPure XP beads (Beckman Coulter, IN) in a ratio of 1.0 to 0.95 and 1.0 of PCR product to beads twice and washed using 70% ethanol per the manufacturer's instructions. To obtain an amplified library, twelve PCRs of 25 µl each were set up, each including 15.5 µl of H₂O, 5 µl of 5 x Phusion HF buffer, 0.5 µl of a dNTP mix containing 10 mM of each dNTP, 1.25 µl of DMSO, 0.25 µl of Illumina PE primer #1, 0.25 µl of Illumina PE primer #2, 0.25 µl of Hotstart Phusion polymerase, and 2 µl of the DNA. The PCR program used was: 98°C for 2 minutes; 12 cycles of 98°C for 15 seconds, 65°C for 30 seconds, 72°C for 30 seconds; and 72°C for 5 min. DNA was purified using Agencourt AMPure XP beads (Beckman Coulter, IN) in a ratio of 1.0 to 1.0 of PCR product to beads and washed using 70% ethanol per the manufacturer's instructions. Exonic or targeted regions were captured in solution using the Agilent SureSelect v.4 kit according to the manufacturer's instructions (Agilent, Santa Clara, CA). The captured library was then purified with a Qiagen MinElute column purification kit and eluted in 17 µl of 70°C EB to obtain 15 µl of captured DNA library. (5) The captured DNA library was amplified in the following way: Eight 30µL PCR reactions each containing 19 µl of H₂O, 6 µl of 5 x Phusion HF buffer, 0.6 µl of 10 mM dNTP, 1.5 µl of DMSO, 0.30 µl of Illumina PE primer #1, 0.30µl of Illumina PE primer #2, 0.30 µl of Hotstart Phusion polymerase, and 2 µl of captured exome library were set up. The PCR program used was: 98°C for 30 seconds; 14 cycles (exome) or 16 cycles (targeted) of 98°C for 10 seconds, 65°C for 30 seconds, 72°C for 30 seconds; and 72°C for 5 min. To purify PCR products, a NucleoSpin Extract II purification kit (Macherey-Nagel, PA) was used following the manufacturer's instructions. Paired-end sequencing, resulting in 100 bases from each end of the fragments for exome libraries and 150 bases from each end of the fragment

for targeted libraries, was performed using Illumina HiSeq 2000/2500 and Illumina MiSeq instrumentation (Illumina, San Diego, CA).

- [70] Primary Processing of Next-Generation Sequencing Data and Identification of Putative Somatic Mutations³
- [71] Somatic mutations were identified using VariantDx custom software (Personal Genome Diagnostics, Baltimore, Maryland) for identifying mutations in matched tumor and normal samples. Prior to mutation calling, primary processing of sequence data for both tumor and normal samples were performed using Illumina CASAVA software (v1.8), including masking of adapter sequences. Sequence reads were aligned against the human reference genome (version hg18) using ELAND with additional realignment of select regions using the Needleman-Wunsch method 5. Candidate somatic mutations, consisting of point mutations, insertions, and deletions were then identified using VariantDx across the either the whole exome or regions of interest. VariantDx examines sequence alignments of tumor samples against a matched normal while applying filters to exclude alignment and sequencing artifacts. In brief, an alignment filter was applied to exclude quality failed reads, unpaired reads, and poorly mapped reads in the tumor. A base quality filter was applied to limit inclusion of bases with reported phred quality score > 30 for the tumor and > 20 for the normal. A mutation in the tumor was identified as a candidate somatic mutation only when (i) distinct paired reads contained the mutation in the tumor; (ii) the number of distinct paired reads containing a particular mutation in the tumor was at least 10% of read pairs; (iii) the mismatched base was not present in >1% of the reads in the matched normal sample as well as not present in a custom database of common germline variants derived from dbSNP; and (iv) the position was covered in both the tumor and normal at > 150X. Mutations arising from misplaced genome alignments, including paralogous sequences, were identified and excluded by searching the reference genome.
- [72] Candidate somatic mutations were further filtered based on gene annotation to identify those occurring in protein-coding regions. Functional consequences were predicted using snpEff and a custom database of CCDS, RefSeq and Ensembl annotations using

the latest transcript versions available on hg18 from UCSC (<https://genome.ucsc.edu/>). Predictions were ordered to prefer transcripts with canonical start and stop codons and CCDS or Refseq transcripts over Ensembl when available. Finally mutations were filtered to exclude intronic and silent changes, while retaining mutations resulting in missense mutations, nonsense mutations, frameshifts, or splice site alterations. A manual visual inspection step was used to further remove artifactual changes.

[73] MUTANT PEPTIDE MHC BINDING PREDICTION

[74] Somatic frameshift, insertions, deletions, and missense mutations predicted to result in an amino acid change were analyzed for potential MHC class I binding based on the individual patient's HLA haplotype. Our initial analysis focused on HLA-A and HLA-B. Amino acid mutations were linked to their corresponding CCDS accession number and in instances where this was unavailable, either a Refseq or ensemble transcript was used to extract the protein sequence. To identify 8mer, 9mer, and 10mer epitopes, amino acid fragments surrounding each mutation were identified. These 15, 17, and 19 mutant amino acid fragments were analyzed by the epitope prediction program NetMHC 3.4.6. Epitopes with a predicted affinity of <50nm were considered to be strong potential binders and epitopes with a predicted affinity of <500nm were considered to be weak potential binders as suggested by the NetMHC group⁶.

[75] To further refine the total neoantigen burden, we repeated that same process for the complementary wild-type peptide for each mutant peptide. We then filtered for mutant peptides that were strong potential binders when the complementary wild-type peptide was predicted a weak potential binder. These mutant peptides are referred to as mutation-associated neoantigens (MANA). In the event that a patient had a (e.g., cases 1, 17 and 21) single MHC haplotype not supported by NetMHC 3.4, the individual haplotype was not included in our analysis.

[76] STATISTICAL METHODS

[77] Design of the trial⁷

[78] This trial was conducted using a parallel two-stage design to simultaneously evaluate the efficacy of MK-3475 and MSI as a treatment selection marker for anti-PD-1 therapy. It consisted of two-stage phase 2 studies in parallel in the three cohorts of patients described in the text. The study agent, MK-3475, was administered at 10 mg/kg intravenously every 14 days.

[79] For each of Cohort A and B, the co-primary endpoints were progression-free-survival (irPFS) at 20 weeks and objective response (irOR) assessed using immune related criteria. A step-down gatekeeping procedure was used to preserve the overall type I error. A two-stage Green-Dahlberg design was used to evaluate irPFS, with interim and final analysis after 15 and 25 patients, respectively. At stage 1, ≥ 1 of 15 free-of-progression at 20 weeks were required to proceed to the second stage, and ≥ 4 of 25 free-of-progression at 20 weeks were then required to proceed to test for irOR, with ≥ 4 of 25 responders (irCR or irPR) indicating promising efficacy in that cohort. Each cohort could be terminated for efficacy as soon as ≥ 4 free-of-progression at 20 weeks and ≥ 4 responses were confirmed, or be terminated for futility as soon as 0 of 15 in stage 1 were free-of-progression at 20 weeks or ≥ 22 subjects had disease progression by 20 weeks. This design achieves 90% power to detect a 20-week irPFS rate of 25% and 80% power to detect an irOR rate (irORR) of 21%, with an overall type I error of 0.05 at the null hypothesis of 20-week irPFS rate of 5% and irORR of 5%.

[80] For Cohort C, the primary endpoint was irPFS at 20 weeks. A two-stage Green-Dahlberg two-stage design was used, with an interim and final analysis after 14 and 21 patients; at stage 1, ≥ 1 of 14 free-of-progression at 20 weeks were required to proceed to the second stage, with ≥ 4 of 21 free-of-progression at 20 weeks at the end indicating adequate efficacy in Cohort C. The cohort could be terminated as soon as ≥ 4 free-of-progression at 20 weeks were confirmed. The design has 81% power to detect a 20-week irPFS rate of 25% with a 5% type I error at the null hypothesis of 20-week irPFS rate of 5%.

[81] Statistical analysis

- [82] Response and progression were evaluated using RECIST v1.1 and the immune-related response criteria (irRC) adopted from Wolchok et al.⁸, which uses the sum of the products of bidimensional tumor measurements and incorporates new lesions into the sum. Progression-free survival (PFS) rates and irPFS rate at 20-weeks was estimated as the proportion of patients who were free-of-disease progression and alive at 20 weeks after the initiation of pembrolizumab. Patients who had disease progression prior to 20 weeks or were enrolled for >20 weeks at the time the study data were collated were included in the analysis for estimating 20-week PFS (irPFS) rate. Patients who dropped out early due to toxicities or worsening disease and therefore did not have 20-week tumor assessment were considered as having progressive disease. ORR (irORR) was the proportion of patients who achieved best overall response of CR or PR (irCR or irPR). Patients who were in the study long enough to have tumor response evaluations were included in the analysis for estimating response rates. Among those who responded (CR or PR), duration of response was the time of first RECIST response to the time of disease progression, and was censored at the last evaluable tumor assessment for responders who had not progressed.
- [83] PFS and irPFS were defined as the time from the date of initial dose to the date of disease progression or the date of death due to any cause, whichever occurred first. PFS and irPFS were censored on the date of the last evaluable tumor assessment documenting absence of progressive disease for patients who were alive and progression-free. Overall survival (OS) was defined as the time from the date of initial dose to death due to any cause. For patients who were still alive at the time of analysis, the OS time was censored on the last date the patients were known to be alive. Survival times were summarized by the Kaplan-Meier method. As a post hoc analysis, log-rank tests were used to compare Cohort A and B and hazard ratios were estimated based on Cox models.
- [84] The association of percent CEA decline after 1 cycle with PFS or OS was assessed using landmark analysis based on Cox regression models. For correlative studies, non-parametric Wilcoxon test was used to compare mutational load between MMR-deficient and MMR-proficient patients. The effects of baseline mutational burden and

immune markers on response and survival times were examined using logistic regression and Cox regression, respectively.

[85] IMMUNOHISTOCHEMISTRY & IMAGE ANALYSIS

[86] The fraction of malignant cells exhibiting a membranous pattern of B7-H1 expression and the percentage at the invasive front were quantified by three pathologists (R.A.A., F.B., and J.M.T.) as previously reported^{9,10}. Image analysis was used to determine the number of CD8 diaminobenzidine (DAB)-stained cells. Using the H&E-stained slide for each case, we identified the following regions: i) tumor, ii) invasive front (the boundary between malignant and non-malignant tissue), and iii) normal tissue. The CD8-stained slides were scanned at 20x equivalent magnification (0.49 micrometers per pixel) on an Aperio ScanScope AT. Regions corresponding to tumor, invasive front and normal tissue (above, from the H&E) were annotated on separate layers using Aperio ImageScope v12.1.0.5029.

[87] CD8-positive lymphocyte density was calculated in each of the above regions using a custom algorithm implemented in PIP¹¹. Results were converted to Deepzoom images using the VIPS library¹² and visualized using the OpenSeadragon viewer (<http://openseadragon.github.io>).

[88] References for Example 3 only.

1. Bacher JW, Flanagan LA, Smalley RL, et al. Development of a fluorescent multiplex assay for detection of MSI-High tumors. *Disease markers* 2004;20:237-50.
2. Murphy KM, Zhang S, Geiger T, et al. Comparison of the microsatellite instability analysis system and the Bethesda panel for the determination of microsatellite instability in colorectal cancers. *The Journal of molecular diagnostics : JMD* 2006;8:305-11.
3. Jones S, Anagnostou V, Lytle K, et al. Personalized genomic analyses for cancer mutation discovery and interpretation. *Science translational medicine* 2015;7:283ra53.

4. Sausen M, Leary RJ, Jones S, et al. Integrated genomic analyses identify ARID1A and ARID1B alterations in the childhood cancer neuroblastoma. *Nature genetics* 2013;45:12-7.
5. Needleman SB, Wunsch CD. A general method applicable to the search for similarities in the amino acid sequence of two proteins. *Journal of molecular biology* 1970;48:443-53.
6. Lundegaard C, Lamberth K, Harndahl M, Buus S, Lund O, Nielsen M. NetMHC-3.0: accurate web accessible predictions of human, mouse and monkey MHC class I affinities for peptides of length 8-11. *Nucleic acids research* 2008;36:W509-12.
7. Buyse M, Michiels S, Sargent DJ, Grothey A, Matheson A, de Gramont A. Integrating biomarkers in clinical trials. *Expert review of molecular diagnostics* 2011;11:171-82.
8. Wolchok JD, Hoos A, O'Day S, et al. Guidelines for the evaluation of immune therapy activity in solid tumors: immune-related response criteria. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2009;15:7412-20.
9. Llosa NJ, Cruise M, Tam A, et al. The vigorous immune microenvironment of microsatellite instable colon cancer is balanced by multiple counter-inhibitory checkpoints. *Cancer Discov* 2015:43-51.
10. Taube JM, Anders RA, Young GD, et al. Colocalization of Inflammatory Response with B7-H1 Expression in Human Melanocytic Lesions Supports an Adaptive Resistance Mechanism of Immune Escape. *Science Translational Medicine* 2012;4:127ra37.
11. Cuka N, Hempel H, Sfanos K, De Marzo A, Cornish T. PIP: An Open Source Framework for Multithreaded Image Analysis of Whole Slide Images. *LABORATORY INVESTIGATION* 2014;94:398A-A.
12. Cupitt J, Martinez K. VIPS: an image processing system for large images. *Electronic Imaging: Science & Technology*; 1996: International Society for Optics and Photonics. p. 19-28.

EXAMPLE 4

[89] Patients

[90] 41 consecutive patients were enrolled and treated between September 2013 and January 2015. (Table 1). Recruitment included patients in pursuit of a clinical trial option who were known to have tumors with mismatch repair, or who had tumors of unknown status who were then tested. One patient in the MMR-deficient CRC cohort was enrolled under an IRB eligibility waiver allowing a grade 3 bilirubin level. A total of 32 CRC patients were enrolled into Cohorts A and B. All CRC patients received ≥ 2 prior chemotherapy regimens (median=4) except for one MMR-proficient patient who had received one chemotherapeutic and one (non-PD1-based) immunotherapeutic regimen.

[91] Nine subjects diagnosed with MMR-deficient solid tumors other than CRC were enrolled onto Cohort C. All Cohort C patients received ≥ 1 prior cancer treatments (median=2).

EXAMPLE 5

[92] Primary endpoint evaluation

[93] The irORR and irPFS at 20 weeks (Fig. 11 (Table S2)) for Cohort A were 40% (4 of 10 patients; 95% CI, 12 to 74%) and 78% (7 of 9 patients; 95% CI, 40 to 97%) and for Cohort C were 71% (5 of 7 patients; 95% CI, 29 to 96%) and 67% (4 of 6 patients; 95% CI, 22 to 96%). In Cohort B, comprised of patients with MMR-proficient CRCs, irORR and 20-week irPFS were 0% (95%CI, 0 to 20%) and 11% (2 of 18 patients; 95% CI, 1 to 35%). Both the MMR-deficient cohorts A and C reached their predefined early stopping rule for efficacy when four subjects were free-of-disease progression at 20 weeks and four objective responses were observed based on immune-related response criteria (Fig. 11 (Table S2); available on line at New England Journal of Medicine; incorporated by reference herein; and supplementary methods, above).

- [94] The median time of follow-up for patients was 32 weeks (range, 5-51 weeks) for patients with MMR-deficient CRC (Cohort A), 12 weeks (range, 2-56 weeks) for patients with MMR-proficient CRC (Cohort B) and 12 weeks (range, 4-42 weeks) for patients with MMR-deficient non-CRC tumors (Cohort C). All patients evaluable for 20-week irPFS were followed for at least 20 weeks.

EXAMPLE 6

- [95] Radiographic evaluation
- [96] Of the ten evaluable MMR-deficient CRC patients in Cohort A, four (40%; 95% CI, 12-74%) achieved objective responses by RECIST criteria (Table 2, Fig. 1 and Fig. 3 (S2)). Patients were considered not evaluable unless they underwent a 12-week scan. The disease control rate was defined as the fraction of patients who achieved an objective response or whose disease was stable, and was 90% in Cohort A (9 of 10 patients; 95% CI, 55-100%).
- [97] Of the seven evaluable patients with MMR-deficient cancer types other than CRC enrolled in Cohort C, five (71%; 95% CI, 29-96%) achieved objective responses (Table 2, Fig. 3 (S2) and Fig. 1) using RECIST criteria and the disease control rate was 71% (5 of 7 patients; 95% CI, 29-96%).
- [98] Patients in Cohort C responded faster than patients in Cohort A (median time to response by RECIST of 12 vs. 28 weeks, $p=0.03$). Furthermore, all six MMR-deficient tumors that were not associated with Lynch syndrome (100%) achieved an objective response, whereas only three of eleven tumors (27%) associated with Lynch Syndrome responded (Table S3; $p=0.009$; available on-line at New England Journal of Medicine; and incorporated by reference herein). No other baseline characteristics showed statistically significant association with objective responses.
- [99] Of the 18 patients with MMR-proficient CRCs in Cohort B, no objective responses were observed (Table 2, Fig. 3 (S2) and Fig. 1) using RECIST criteria and the disease control rate was 11% (2 of 18 patients; 95%CI, 1 to 35%).

- [100] All patients who achieved a response by RECIST criteria (Fig. 11 (Table 2)) also achieved a response by immune-related response criteria (Fig. 11 (Table S2)).

EXAMPLE 7

[101] Survival

- [102] In Cohort A, the patients with MMR-deficient CRC, median progression-free survival (PFS) and median overall survival (OS) were not reached (Fig. 2). In contrast, the patients with MMR-proficient cancers in Cohort B achieved a PFS of only 2.2 months (95% CI, 1.4-2.8) and a median OS of 5.0 months (95% CI, 3.0 to not estimable). In Cohort C (MMR-deficient non-CRC), the median PFS was 5.4 months (95% CI, 3 to not estimable) and the median OS was not reached.
- [103] A *post hoc* (Fig. 2) comparison of the MMR-deficient and proficient CRC cohorts showed hazard ratios (HR) for disease progression (HR=0.10; 95% CI, 0.03-0.37; $p<0.001$) and overall survival (HR = 0.22; 95% CI, 0.05-1.00; $p=0.05$), favoring patients with MMR-deficient CRC.
- [104] To evaluate whether the difference in survival might be due to prognostic differences, we measured the duration of time patients had been diagnosed with metastatic disease and the clinical performance of patients on their previous regimen prior to enrollment. We found that there was no significant difference between MMR-deficient vs. MMR-proficient CRC patients with respect to their duration of metastatic disease ($p=0.77$; Log-rank test) or median PFS ($p=0.60$, Log-rank test) on their prior regimens (Fig. 4 (S3)). We also performed an additional multivariate analysis of PFS and OS to examine the difference in outcomes between MMR-deficient CRC and MMR-proficient tumors adjusting for elapsed time since initial diagnosis. The magnitude of the hazard ratios for PFS (HR 0.04, 95% CI 0.01-0.21, $P<0.001$) and OS (HR 0.18, 95% CI 0.03-1.01, $P=0.05$), representing the different effect of pembrolizumab between MMR-deficient and MMR-proficient tumors, was maintained after adjusting for this potential difference.

EXAMPLE 8

[105] Safety Assessment

[106] Adverse events occurring in > 5% of patients are listed in Table 3. Select adverse events included rash/pruritus (24%), thyroiditis/hypothyroidism/hypophysitis (10%), and asymptomatic pancreatitis (15%). While the numbers were small, thyroid function abnormalities were limited to the MMR-deficient cohorts (Table 3).

EXAMPLE 9

[107] Tumor markers

[108] In the two CRC cohorts, baseline CEA levels were evaluable and above the upper limit of normal (3 mg/dl), in 29 of 32 patients prior to enrollment. Major CEA declines occurred in seven of the ten patients with MMR-deficient CRC and in none of the 19 patients with MMR-proficient CRC in which CEA was evaluable (Fig. 1 and Fig. 5 (S4)). In non-CRC MMR-deficient patients, tumor marker levels (CEA, CA19-9 or CA-125) were elevated above the upper limit of normal in four patients. CA19-9 or CA-125 declines of > 70% occurred in three of these four patients. Tumor marker kinetics of all 3 cohorts are shown in Figure 1. The level of CEA decline after 1 dose (between days 14 and 28) of pembrolizumab was predictive of both progression-free (p=0.01) and overall survival outcomes (p=0.02). The CEA response occurred well in advance of radiographic confirmation of disease control (range, 10 to 35 weeks). In contrast, patients who progressed showed rapid biomarker elevation within 30 days of initiating therapy. Thus, changes in CEA levels significantly preceded and correlated with ultimate radiographic changes.

EXAMPLE 10

[109] Genomic Analysis

[110] Analysis of whole-exome sequences showed an average of 1,782 somatic mutations per tumor in MMR-deficient patients (n = 9) compared with 73 mutations per tumor in MMR-proficient patients (n=6) (non-parametric Wilcoxon test, p=0.007) (Figs. 6A-

6B (S5); see also Table S3 which is available on-line at New England Journal of Medicine; incorporated by reference herein). Most (63%) of these mutations are predicted to alter amino acids.

- [111] These mutations were then assessed for their immunogenic potential in the context of each patient's individual MHC haplotype. We thereby identified an average of 578 and 21 potential mutation-associated neoantigens from the tumors of MMR-deficient and MMR-proficient patients, respectively (Table S3; which is available on-line at New England Journal of Medicine; incorporated by reference herein). The fraction of potential mutation-associated neoantigens among all somatic mutations was similar in both cohorts (averaging 32% and 29% in MMR-deficient and -proficient patients, respectively). High numbers of somatic mutations and potential mutation-associated neoantigens were associated with improved progression-free survival and with a trend in favor of objective response (Fig. 13 (S5) and Fig. 12 (Table S4); also available on line at New England Journal of Medicine; incorporated by reference herein).

EXAMPLE 11

[112] Immunohistochemistry

- [113] Expression of CD8 and PD-L1 were evaluated by immunohistochemistry within the tumor and at the invasive fronts of the tumor in the 30 cases in which tumor tissue was available (Fig. 7 (S6); also available on line at New England Journal of Medicine; incorporated by reference herein). Tumors from patients in Cohorts A and C contained a greater density of CD8-positive lymphoid cells than did tumors from Cohort B patients (Fig. 8 (S7); $p=0.10$) and CD8-labeling was associated with a trend favoring objective response and stable disease (Fig. 9 (S8) and Fig. 13 (Table S5); also available on line at New England Journal of Medicine; incorporated by reference herein). This CD8-positive lymphoid infiltrate was especially prominent at the invasive fronts of the tumors (Fig 8 (S7); $p=0.04$). Significant membranous PD-L1 expression only occurred in MMR-deficient patients and was prominent on tumor infiltrating lymphocytes (TILs) and tumor-associated macrophages located at the tumors' invasive fronts (Fig. 8 (S7); $p=0.04$). Expression of CD8 and PD-L1 were not statistically associated with PFS or OS (Fig. 13 (Table S5)).

Table 1. Demographic and Baseline Characteristics of Patients

Characteristic	MMR-deficient CRC n=11	MRC- proficient CRC n=21	P values ¹	MMR- deficient non-CRC n=9
Age – years				
median	46	61	0.02	57
range	(24-65)	(32-79)		(34-92)
Sex – no. (%)				
Female	5(45)	8(38)	0.72	4(44)
Male	6(55)	13(62)		5(56)
Race – no. (%)				
white	8(73)	17(81)	0.66	8(89)
black	1(9)	3(14)		0(0)
other	2(18)	1(5)		1(11)
ECOG Performance Status – no. (%) ²				
0	0(0)	6(29)	0.07	2(22)
1	11(100)	15(71)		7(78)
Diagnosis – no. (%)				
Colon	9(82)	18(86)	>0.99	0(0)
Rectal	2(18)	3(14)		0(0)
Ampullary/Cholangiocarcinoma	0(0)	N/A		4(44)
Endometrial	0(0)	N/A		2(22)
Small bowel	0(0)	N/A		2(22)
Gastric	0(0)	N/A		1(11)
Histology – no. (%)				
Well/moderately differentiated	7(64)	18(86)	0.20	4(44)
Poorly differentiated	4(36)	3(14)		3(33)
Other	0(0)	0(0)		2(22)
Stage IV – no. (%)	(11)100	21(100)	>0.99	9(100)
Liver metastases – no. (%)	6(55)	11(52)	>0.99	6(67)
Time since first diagnosis - months				
median	31	58	0.07	23
range ³	6-95	27-192		2-105
Prior systemic therapies – no. (%)				
0	0(0)	0(0)	0.89	1(11)
2	3(27)	4(19)		5(56)
3	3(27)	5(24)		1(11)
>4	5(45)	12(57)		2(22)
Detected germline mutation or known Lynch – no. (%)				
Yes	9(82)	0(0)	<0.001	4(44)
No	2(18)	21(100)		4(44)
Unknown	0(0)	0(0)		1(11)
BRAF wild type – no. (%)				
Yes	8(73)	11(52)	0.64	4(44)
No	0(0)	1(5)		0(0)
Unknown	3(27)	9(43)		5(56)
KRAS wild type – no. (%)				
Yes	6(55)	13(62)	0.72	4(44)
No	5(45)	8(38)		1(11)
Unknown	0(0)	0(0)		4(44)

MMR, mismatch repair; CRC, colorectal cancer

¹ MMR-deficient CRC versus MMR-proficient CRC² ECOG, Eastern Cooperative Oncology Group

Table 2. Objective RECIST responses

Type of Response – no. (%)	MMR-deficient CRC n=10	MRC-proficient CRC n=18	MMR-deficient non-CRC n=7
Complete Response	0(0)	0(0)	1(14) ¹
Partial Response	4(40)	0(0)	4(57) ²
Stable Disease (Week 12)	5(50)	2(11)	0(0)
Progressive Disease	1(10)	11(61)	2(29)
Not Evaluable ³	0(0)	5(28)	0(0)
Objective Response Rate (%)	40	0	71
95% CI	12-74	0-19	29-96
Disease Control Rate (%) ⁴	90	11	71
95% CI	55-100	1-35	29-96
Duration of Response – median weeks	Not reached	N/A ⁵	Not reached
Time to Response, median weeks (range)	28 (13-35)	N/A ⁵	11 (10-13)

¹ Originally PR at 12 weeks that was converted to CR at 20 weeks

² One PR at 12 weeks

³ Patients were considered not evaluable if they did not undergo a 12 week scan due to clinical progression.

⁴ The rate of disease control was defined as the percentage of patients who had a complete response, partial response or stable disease for 12 weeks or more.

⁵ No responses recorded for MMR-proficient CRC patients

Table 3. Drug-Related Adverse Events

Event – no (%) ¹	All Grades N=41	Grade 3 or 4 N=41
Any	40(98)	17(41)
Blood and Lymphatic		
Anemia	8(20)	7(17)
Lymphopenia	8(20)	8(20)
Cardiac		
Sinus tachycardia	4(10)	0
Dermatologic		
Dry skin	5(12)	0
Rash/pruritis	10(24)	0
Endocrine Disorders		
Thyroiditis/Hypothyroidism/Hypophysitis	4(10)	0
Gastrointestinal		
Abdominal Pain	10(24)	0
Anorexia	4(10)	0
Constipation	8(20)	0
Diarrhea	10(24)	2(5)
Dry mouth	5(12)	0
Nausea	5(12)	0
Bowel Obstruction	3(7)	3(7)
Hepatobiliary		
ALT, elevated	3(7)	2(5)
Pancreatitis ²	6(15)	0
Metabolism and Nutrition		
Hypoalbuminemia	4(10)	4(10)
Hyponatremia	3(7)	3(7)
Musculoskeletal		
Arthralgia	7(17)	0
Myalgia	6(15)	0
Nervous System		
Dizziness	4(10)	0
Headache	7(17)	0
Psychiatric		
Insomnia	3(7)	0
Respiratory ³		
Allergic Rhinitis	12(29)	0
Cough	4(10)	0
Dyspnea	6(15)	0
Upper Respiratory Infection	3(7)	0
Other		
Cold intolerance	6(15)	0
Edema	4(10)	0
Fatigue	13(32)	0
Fever	5(12)	0
Pain	14(34)	0

¹ Adverse Events occurring in greater than 5% of patients² All cases of pancreatitis were asymptomatic³ One incidence of pneumonitis (2%)

References

The disclosure of each reference cited is expressly incorporated herein.

1. Nishimura H, Okazaki T, Tanaka Y, et al. Autoimmune Dilated Cardiomyopathy in PD-1 Receptor-Deficient Mice. *Science* 2001;291:319-22.
2. Chen L. Co-inhibitory molecules of the B7-CD28 family in the control of T-cell immunity. *Nat Rev Immunol* 2004;4:336-47.
3. Nishimura H, Nose M, Hiai H, Minato N, Honjo T. Development of Lupus-like Autoimmune Diseases by Disruption of the PD-1 Gene Encoding an ITIM Motif-Carrying Immunoreceptor. *Immunity* 1999;11:141-51.
4. Ansell SM, Lesokhin AM, Borrello I, et al. PD-1 Blockade with Nivolumab in Relapsed or Refractory Hodgkin's Lymphoma. *The New England journal of medicine* 2015;372:311-9.
5. Hamid O, Robert C, Daud A, et al. Safety and tumor responses with lambrolizumab (anti-PD-1) in melanoma. *The New England journal of medicine* 2013;369:134-44.
6. Herbst RS, Soria JC, Kowanetz M, et al. Predictive correlates of response to the anti-PD-L1 antibody MPDL3280A in cancer patients. *Nature* 2014;515:563-7.
7. Powles T, Eder JP, Fine GD, et al. MPDL3280A (anti-PD-L1) treatment leads to clinical activity in metastatic bladder cancer. *Nature* 2014;515:558-62.
8. Topalian SL, Sznol M, McDermott DF, et al. Survival, durable tumor remission, and long-term safety in patients with advanced melanoma receiving nivolumab. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2014;32:1020-30.

9. Brahmer JR, Tykodi SS, Chow LQ, et al. Safety and activity of anti-PD-L1 antibody in patients with advanced cancer. *The New England journal of medicine* 2012;366:2455-65.
10. Topalian SL, Hodi FS, Brahmer JR, et al. Safety, activity, and immune correlates of anti-PD-1 antibody in cancer. *The New England journal of medicine* 2012;366:2443-54.
11. Taube JM, Klein A, Brahmer JR, et al. Association of PD-1, PD-1 ligands, and other features of the tumor immune microenvironment with response to anti-PD-1 therapy. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2014;20:5064-74.
12. Brahmer JR, Drake CG, Wollner I, et al. Phase I study of single-agent anti-programmed death-1 (MDX-1106) in refractory solid tumors: safety, clinical activity, pharmacodynamics, and immunologic correlates. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2010;28:3167-75.
13. Koopman M, Kortman GAM, Mekenkamp L, et al. Deficient mismatch repair system in patients with sporadic advanced colorectal cancer. *Br J Cancer* 0000;100:266-73.
14. Goldstein J, Tran B, Ensor J, et al. Multicenter retrospective analysis of metastatic colorectal cancer (CRC) with high-level microsatellite instability (MSI-H). *Annals of oncology : official journal of the European Society for Medical Oncology / ESMO* 2014;25:1032-8.
15. Segal NH, Parsons DW, Peggs KS, et al. Epitope landscape in breast and colorectal cancer. *Cancer research* 2008;68:889-92.

16. Timmermann B, Kerick M, Roehr C, et al. Somatic mutation profiles of MSI and MSS colorectal cancer identified by whole exome next generation sequencing and bioinformatics analysis. *PloS one* 2010;5:e15661.
17. Eshleman JR, Lang EZ, Bowerfind GK, et al. Increased mutation rate at the hprt locus accompanies microsatellite instability in colon cancer. *Oncogene* 1995;10:33-7.
18. Comprehensive molecular characterization of human colon and rectal cancer. *Nature* 2012;487:330-7.
19. Dolcetti R, Viel A, Doglioni C, et al. High prevalence of activated intraepithelial cytotoxic T lymphocytes and increased neoplastic cell apoptosis in colorectal carcinomas with microsatellite instability. *The American journal of pathology* 1999;154:1805-13.
20. Alexander J, Watanabe T, Wu TT, Rashid A, Li S, Hamilton SR. Histopathological identification of colon cancer with microsatellite instability. *The American journal of pathology* 2001;158:527-35.
21. Smyrk TC, Watson P, Kaul K, Lynch HT. Tumor-infiltrating lymphocytes are a marker for microsatellite instability in colorectal carcinoma. *Cancer* 2001;91:2417-22.
22. Young J, Simms LA, Biden KG, et al. Features of colorectal cancers with high-level microsatellite instability occurring in familial and sporadic settings: parallel pathways of tumorigenesis. *The American journal of pathology* 2001;159:2107-16.
23. Berger MF, Hodis E, Heffernan TP, et al. Melanoma genome sequencing reveals frequent PREX2 mutations. *Nature* 2012;485:502-6.
24. Lee W, Jiang Z, Liu J, et al. The mutation spectrum revealed by paired genome sequences from a lung cancer patient. *Nature* 2010;465:473-7.

25. Lipson EJ, Sharfman WH, Drake CG, et al. Durable cancer regression off-treatment and effective reinduction therapy with an anti-PD-1 antibody. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2013;19:462-8.
26. Boland CR, Goel A. Microsatellite instability in colorectal cancer. *Gastroenterology* 2010;138:2073-87 e3.
27. Lynch HT, de la Chapelle A. Hereditary colorectal cancer. *The New England journal of medicine* 2003;348:919-32.
28. Yamamoto H, Imai K, Perucho M. Gastrointestinal cancer of the microsatellite mutator phenotype pathway. *Journal of gastroenterology* 2002;37:153-63.
29. Herman JG, Umar A, Polyak K, et al. Incidence and functional consequences of hMLH1 promoter hypermethylation in colorectal carcinoma. *Proceedings of the National Academy of Sciences of the United States of America* 1998;95:6870-5.
30. Jones S, Anagnostou V, Lytle K, et al. Personalized genomic analyses for cancer mutation discovery and interpretation. *Science translational medicine* 2015;7:283ra53.
31. Lundegaard C, Lamberth K, Harndahl M, Buus S, Lund O, Nielsen M. NetMHC-3.0: accurate web accessible predictions of human, mouse and monkey MHC class I affinities for peptides of length 8-11. *Nucleic acids research* 2008;36:W509-12.
32. Lundegaard C, Lund O, Nielsen M. Accurate approximation method for prediction of class I MHC affinities for peptides of length 8, 10 and 11 using prediction tools trained on 9mers. *Bioinformatics* 2008;24:1397-8.
33. Wolchok JD, Hoos A, O'Day S, et al. Guidelines for the evaluation of immune therapy activity in solid tumors: immune-related response criteria. *Clinical cancer*

research : an official journal of the American Association for Cancer Research

2009;15:7412-20.

34. Maple JT, Smyrk TC, Boardman LA, Johnson RA, Thibodeau SN, Chari ST.

Defective DNA mismatch repair in long-term (> or =3 years) survivors with pancreatic cancer. *Pancreatology* 2005;5:220-7; discussion 7-8.

35. Meltzer SJ, Yin J, Manin B, et al. Microsatellite instability occurs frequently and in both diploid and aneuploid cell populations of Barrett's-associated esophageal adenocarcinomas. *Cancer research* 1994;54:3379-82.

36. Nakata B, Wang YQ, Yashiro M, et al. Prognostic value of microsatellite instability in resectable pancreatic cancer. *Clinical cancer research : an official journal of the American Association for Cancer Research* 2002;8:2536-40.

37. Comprehensive molecular characterization of gastric adenocarcinoma. *Nature* 2014;513:202-9.

38. Agaram NP, Shia J, Tang LH, Klimstra DS. DNA mismatch repair deficiency in ampullary carcinoma: a morphologic and immunohistochemical study of 54 cases. *American journal of clinical pathology* 2010;133:772-80.

39. Kandoth C, Schultz N, Cherniack AD, et al. Integrated genomic characterization of endometrial carcinoma. *Nature* 2013;497:67-73.

40. Garg K, Leitao MM, Jr., Kauff ND, et al. Selection of endometrial carcinomas for DNA mismatch repair protein immunohistochemistry using patient age and tumor morphology enhances detection of mismatch repair abnormalities. *The American journal of surgical pathology* 2009;33:925-33.

41. Snyder A, Makarov V, Merghoub T, et al. Genetic basis for clinical response to CTLA-4 blockade in melanoma. *The New England journal of medicine* 2014;371:2189-99.

42. Williams AS, Huang WY. The analysis of microsatellite instability in extracolonic gastrointestinal malignancy. *Pathology* 2013;45:540-52.
43. Jones S, Emmerson P, Maynard J, et al. Biallelic germline mutations in MYH predispose to multiple colorectal adenoma and somatic G:C→T:A mutations. *Human Molecular Genetics* 2002;11:2961-7.
44. Palles C, Cazier J-B, Howarth KM, et al. Germline mutations affecting the proofreading domains of POLE and POLD1 predispose to colorectal adenomas and carcinomas. *Nature genetics* 2013;45:136-44.
45. Bodmer W, Bishop T, Karran P. Genetic steps in colorectal cancer. *Nature genetics* 1994;6:217-9.
46. Kim H, Jen J, Vogelstein B, Hamilton SR. Clinical and pathological characteristics of sporadic colorectal carcinomas with DNA replication errors in microsatellite sequences. *The American journal of pathology* 1994;145:148-56.
47. Llosa NJ, Cruise M, Tam A, et al. The vigorous immune microenvironment of microsatellite instable colon cancer is balanced by multiple counter-inhibitory checkpoints. *Cancer Discov* 2015;43-51. [This disclosure was made by some of the joint inventors.]
48. Rizvi NA, Hellmann MD, Snyder A, et al. Mutational landscape determines sensitivity to PD-1 blockade in non-small cell lung cancer. *Science* 2015.
49. Gubin MM, Zhang X, Schuster H, et al. Checkpoint blockade cancer immunotherapy targets tumour-specific mutant antigens. *Nature* 2014;515:577-81.
50. Linnemann C, van Buuren MM, Bies L, et al. High-throughput epitope discovery reveals frequent recognition of neo-antigens by CD4+ T cells in human melanoma. *Nature medicine* 2015;21:81-5.

51. Lipson EJ, Velculescu VE, Pritchard TS, et al. Circulating tumor DNA analysis as a real-time method for monitoring tumor burden in melanoma patients undergoing treatment with immune checkpoint blockade. *Journal for immunotherapy of cancer* 2014;2:42.
52. Diaz LA, Jr., Bardelli A. Liquid biopsies: genotyping circulating tumor DNA. *Journal of clinical oncology : official journal of the American Society of Clinical Oncology* 2014;32:579-86.
53. Yadav M, Jhunjunwala S, Phung QT, et al. Predicting immunogenic tumour mutations by combining mass spectrometry and exome sequencing. *Nature* 2014;515:572-6.

We claim:

1. A method of treating a cancer patient comprising:
administering to a cancer patient an immune checkpoint inhibitory antibody, wherein the cancer patient has a microsatellite instable cancer (MSI).
2. The method of claim 1 wherein the cancer is selected from the group consisting of: colon, gastric, endometrial, cholangiocarcinoma, pancreatic, and prostate cancers.
3. The method of claim 1 wherein the antibody is an anti-PD-1 antibody.
4. The method of claim 1 wherein the antibody is an anti-IDO antibody.
5. The method of claim 1 wherein the antibody is an anti-CTLA-4 antibody.
6. The method of claim 1 wherein the antibody is an anti-PD-L1 antibody.
7. The method of claim 1 wherein the antibody is an anti-LAG-3 antibody.
8. The method of claim 1 wherein a sample from the MSI cancer patient is tested to evaluate stability of one or more microsatellite markers prior to the step of administering.
9. The method of claim 1 wherein the antibody is a humanized monoclonal antibody.
10. The method of claim 1 wherein the antibody is MK-3475.
11. The method of claim 1 wherein the antibody is an IgG4 antibody.
12. The method of claim 8 wherein a plurality of microsatellite markers is tested.
13. The method of claim 8 wherein the one or more microsatellite markers is selected from the group consisting of BAT-25, BAT-26, MON0-27, NR-21, NR-24, Penta C, and Penta D.
14. The method of claim 12 wherein the plurality of microsatellite markers comprises BAT-25, BAT-26, MON0-27, NR-21, NR-24, Penta C, and Penta D.
15. A method of treating a cancer patient comprising:
testing a sample from a cancer patient and determining instability of one or more microsatellite markers selected from the group consisting of BAT-25, BAT-26, MON0-27, NR-21, NR-24, Penta C, and Penta D; and
administering to the cancer patient an anti-PD-1 antibody, wherein the cancer is selected from the group consisting of: colon, gastric, endometrial, cholangiocarcinoma, pancreatic, and prostate cancers.

16. A method of categorizing a tumor of a human, comprising:
 - testing a sample from the human to evaluate stability of one or more microsatellite markers;
 - determining microsatellite instability in the sample;
 - identifying the tumor as a good candidate for treatment with an immune checkpoint inhibitory antibody.
17. The method of claim 16 further comprising the step of:
 - prescribing an immune checkpoint inhibitory antibody for the human.
18. The method of claim 16 wherein the one or more microsatellite markers is selected from the group consisting of BAT-25, BAT-26, MON0-27, NR-21, NR-24, Penta C, and Penta D.
19. The method of claim 16 wherein the one or more microsatellite markers is selected from the group consisting of BAT-25, BAT-26, MON0-27, NR-21, NR-24, Penta C, and Penta D.
20. A method of categorizing a tumor of a human, comprising:
 - testing a sample from the human to evaluate stability of one or more microsatellite markers;
 - determining microsatellite stability in the sample;
 - identifying the tumor as a bad candidate for treatment with an immune checkpoint inhibitory antibody.
21. The method of claim 20 wherein the human is being treated with an immune checkpoint inhibitory antibody, and further comprising the step of ceasing treatment with an immune checkpoint inhibitory antibody.
22. The method of claim 20 wherein the one or more microsatellite markers is selected from the group consisting of BAT-25, BAT-26, MON0-27, NR-21, NR-24, Penta C, and Penta D.
23. The method of claim 20 wherein the one or more microsatellite markers is selected from the group consisting of BAT-25, BAT-26, MON0-27, NR-21, NR-24, Penta C, and Penta D.
24. A method of treating a cancer patient comprising:
 - administering to a cancer patient an immune checkpoint inhibitory antibody, wherein the cancer patient has a mutational burden of at least 100 somatic mutations per tumor.
25. The method of claim 24 wherein the antibody is an anti-PD-1 antibody.

26. The method of claim 24 wherein the antibody is an anti-IDO antibody.
27. The method of claim 24 wherein the antibody is an anti-CTLA-4 antibody.
28. The method of claim 24 wherein the antibody is an anti-PD-L1 antibody.
29. The method of claim 24 wherein the antibody is an anti-LAG-3 antibody.
30. The method of claim 24 further comprising determining a high mutational burden in the tumor prior to the step of administering.