

PHENYLPYRROLIDINONE FORMYL PEPTIDE 2 RECEPTOR AGONISTS

CROSS REFERENCE TO RELATED APPLICATIONS

This application is entitled to priority pursuant to 35 U.S.C. §119(e) to U.S.
5 provisional patent application No. 62/638,556, filed March 5, 2018, which is incorporated
herein in its entirety.

BACKGROUND OF THE INVENTION

The present invention relates to novel pyrrolidinone compounds, which are formyl
10 peptide 2 (FPR2) receptor agonists and/or formyl peptide 1 (FPR1) receptor agonists,
compositions containing them, and methods of using them, for example, for the treatment
of atherosclerosis, heart failure, chronic obstructive pulmonary disease (COPD), and
related diseases.

Formyl peptide receptor 2 (FPR2) belongs to a small group of seven-
15 transmembrane domain, G protein-coupled receptors that are expressed in multiple
human tissues including immune cells and are known to be important in host defense and
inflammation. FPR2 shares significant sequence homology with FPR1 and FPR3 (Journal
of Autoimmunity 85, 2017, 64-77). Collectively, these receptors bind a number of
structurally diverse agonists, including N-formyl and nonformyl peptides which act as
20 chemo attractants and activate phagocytes. The endogenous peptide Annexin A1 and its
N-terminal fragments are examples of ligands that bind human FPR1 and FPR2. Fatty
acids such as eicosanoid, lipoxin A₄, which belongs to a class of small pro-resolution
mediators (SPMs), has also been identified as an agonist for FPR2 (Ye RD., et al.,
Pharmacol. Rev., 2009, 61, 119-61).

25 Endogenous FPR2 pro-resolution ligands, such as lipoxin A₄ and Annexin A1,
have been reported to trigger a wide array of cytoplasmatic cascades such as Gi coupling,
Ca²⁺ mobilization and β-arrestin recruitment. (Int J Mol Sci. 2013 April; 14(4): 7193–
7230). FPR2 regulates both innate and adaptive immune systems including neutrophils,
macrophages, T-, and B-cells. In neutrophils, FPR2 ligands modulate movement,
30 cytotoxicity and life span. In macrophages, agonism of FPR2 prevents apoptosis and
enhances efferocytosis. (Chandrasekharan JA, Sharma-Walia N., J. Inflamm. Res., 2015,
8, 181-92). The initiation of resolution of inflammation by FPR2 agonism is responsible

for enhancing anti-fibrotic wound healing and returning of the injured tissue to homeostasis (Romano M., et al., Eur. J. Pharmacol., 2015, 5, 49-63).

Chronic inflammation is part of the pathway of pathogenesis of many human diseases and stimulation of resolution pathways with FPR2 agonists may have both protective and reparative effects. Ischaemia-reperfusion (I/R) injury is a common feature of several diseases associated with high morbidity and mortality, such as myocardial infarction and stroke. Non-productive wound healing associated with cardiomyocyte death and pathological remodeling resulting from ischemia-reperfusion injury leads to scar formation, fibrosis, and progressive loss of heart function. FPR2 modulation is proposed to enhance myocardial wound healing post injury and diminish adverse myocardial remodeling (Kain V., et al., J. Mol. Cell. Cardiol., 2015, 84, 24-35). In addition, FPR2 pro-resolution agonists, in the central nervous system, may be useful therapeutics for the treatment of a variety of clinical I/R conditions, including stroke in brain (Gavins FN., Trends Pharmacol. Sci., 2010, 31, 266-76) and I/R induced spinal cord injury (Liu ZQ ., et al., Int. J. Clin. Exp. Med., 2015, 8, 12826-33).

In addition to beneficial effects of targeting the FPR2 receptor with novel pro-resolution agonists for treatment of I/R induced injury therapeutic, utility of these ligands can also be applied to other diseases. In the cardiovascular system both the FPR2 receptor and its pro-resolution agonists were found to be responsible for atherogenic-plaque stabilization and healing (Petri MH., et al., Cardiovasc. Res., 2015, 105, 65-74; and Fredman G., et al., Sci. Trans. Med., 2015, 7(275);275ra20). FPR2 agonists also have been shown to be beneficial in preclinical models of chronic inflammatory human diseases, including: infectious diseases, psoriasis, dermatitis, inflammatory bowel syndrome, Crohn's disease, ocular inflammation, sepsis, pain, metabolic/diabetes diseases, cancer, COPD, asthma and allergic diseases, cystic fibrosis, acute lung injury and fibrosis, rheumatoid arthritis and other joint diseases, Alzheimer's disease, kidney fibrosis, and organ transplantation (Romano M., et al., Eur. J. Pharmacol., 2015, 5, 49-63, Perrett, M., et al., Trends in Pharm. Sci., 2015, 36, 737-755).

SUMMARY OF THE INVENTION

The present invention provides novel pyrrolidinone, and their analogues thereof, which are useful as FPR2 agonists, including stereoisomers, tautomers, pharmaceutically acceptable salts, or solvates thereof.

The present invention also provides processes and intermediates for making the compounds of the present invention or stereoisomers, tautomers, pharmaceutically acceptable salts, or solvates thereof.

5 The present invention also provides pharmaceutical compositions comprising a pharmaceutically acceptable carrier and at least one of the compounds of the present invention or stereoisomers, tautomers, pharmaceutically acceptable salts, or solvates thereof.

The compounds of the invention may be used in therapy.

10 The compounds of the invention may be used in the treatment and/or prophylaxis of multiple diseases or disorders associated with FPR2, such as inflammatory diseases, heart diseases, chronic airway diseases, cancers, septicemia, allergic symptoms, HIV retrovirus infection, circulatory disorders, neuroinflammation, nervous disorders, pains, prion diseases, amyloidosis, and immune disorders. The heart diseases are selected from the group consisting of angina pectoris, unstable angina, myocardial infarction, acute
15 coronary disease, cardiac iatrogenic damage, and heart failure including, but not limited to, acute heart failure, chronic heart failure of ischemic and non-ischemic origin, systolic heart failure, diastolic heart failure, heart failure with reduced ejection fraction (HF_{REF}), and heart failure with preserved ejection fraction (HF_{PEF}).


20 The compounds of the invention can be used alone, in combination with other compounds of the present invention, or in combination with one or more other agent(s).

Other features and advantages of the invention will be apparent from the following detailed description and claims.

DESCRIPTION OF THE INVENTION

25 The invention encompasses compounds of Formula (I), which are formyl peptide 2 (FPR2) receptor agonists and/or formyl peptide 1 (FPR1) receptor agonists, compositions containing them, and methods of using them, for example, in the treatment of atherosclerosis, heart failure, chronic obstructive pulmonary disease (COPD), and related diseases.

30 One aspect of the invention is a compound of Formula (I):



wherein:

Ar¹ is phenyl, pyridinyl, or pyridazinyl and is substituted with 1 halo, haloalkyl or haloalkoxy substituent in the 4-position and 0-2 additional halo or haloalkyl substituents;

5 Ar² is phenyl or pyridinyl substituted with 0-2 substituents selected from cyano, fluoro, alkyl, haloalkyl, cycloalkyl, alkoxy, and haloalkoxy;

Ar³ is phenyl or pyridinyl and is substituted with 0-2 substituents selected from cyano, halo, hydroxyalkyl, alkoxyalkyl, (R¹R²N)alkyl, (alkyl)₂(O)P, (alkyl)(O)(NR¹)S, alkylSO₂, and alkylSO₂NH;

10 R¹ is hydrogen or alkyl; and

R² is hydrogen or alkyl; or (R¹)(R²)N taken together is azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, or morpholinyl and is substituted with 0-3 substituents selected from fluoro and alkyl; or a pharmaceutically acceptable salt thereof.

15 Another aspect of the invention is a compound of Formula (I) where Ar¹ is phenyl substituted with 1 halo, haloalkyl or haloalkoxy substituent in the 4-position and 0-2 additional halo or haloalkyl substituents.


Another aspect of the invention is a compound of Formula (I) where Ar² is phenyl substituted with 0 substituents, 1 alkyl or cycloalkyl substituent, or 2 fluoro substituents.

20 Another aspect of the invention is a compound of Formula (I) where Ar² is phenyl substituted with 0 substituents or 2 fluoro substituents.

Another aspect of the invention is a compound of Formula (I) where Ar³ is phenyl substituted with 1-2 substituents selected from cyano, halo, hydroxyalkyl, alkoxyalkyl, (R¹R²N)alkyl, (alkyl)₂(O)P, (alkyl)(O)(NR¹)S, alkylSO₂, and alkylSO₂NH.

25 Another aspect of the invention is a compound of Formula (I) where R¹ is hydrogen and R² is hydrogen.

Another aspect of the invention is a compound of Formula (II):



or a pharmaceutically acceptable salt thereof, wherein:

Ar¹ is aryl substituted with 1-2 R^{1a} and 1-2 R^{1b} or monocyclic heteroaryl with 1-3 heteroatoms selected from nitrogen, oxygen, and sulfur, and substituted with 1-2 R^{1a} and 1-2 R^{1b};

Ar² is aryl substituted with 1-4 R^{2a} or 6-membered heteroaryl with 1-2 nitrogen atoms, and substituted with 1-4 R^{2a};

Ar³ is aryl substituted with 1-4 R^{3a} or monocyclic heteroaryl with 1-3 heteroatoms selected from nitrogen, oxygen, and sulfur, and substituted with 1-4 R^{3a};

R^{1a} is hydrogen or halo;

R^{1b} is halo, haloalkyl, alkoxy, or haloalkoxy;

R^{2a} is hydrogen, cyano, halo, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl, alkoxy, or haloalkoxy; alternatively, two adjacent R^{2a} groups are taken together with the carbon atoms to which they are attached to form a heterocycle with 1-4 heteroatoms selected from nitrogen, oxygen, and sulfur;

R^{3a} is cyano, halo, alkyl, alkoxy, hydroxyalkyl, alkoxyalkyl, haloalkyl, (R¹R²N)alkyl, R¹R²N, alkylC(O)(R²)Nalkyl, (alkyl)₂(O)P, (alkoxy)₂(O)P, (alkoxy)(alkyl)(O)P, (alkyl)(O)(NR¹)S, alkylSO₂, or alkylSO₂NH; alternatively, two adjacent R^{3a} groups are taken together with the carbon atoms to which they are attached to form a heterocycle with 1-4 heteroatoms selected from nitrogen, oxygen, and sulfur;


R^{4a} or R^{4b} is independently hydrogen, alkyl, alkoxy, hydroxylalkyl, alkoxyalkyl, or haloalkoxy; alternatively, R^{4a} and R^{4b} together with the carbon atom they are both attached to form a C₃₋₆ cycloalkyl;

R^{5a} or R^{5b} is independently hydrogen, alkyl, hydroxylalkyl, alkoxyalkyl or haloalkoxyl;

R¹ is hydrogen or alkyl; and

R² is hydrogen or alkyl; or R¹R²N taken together is azetidiny, oxazolyl pyrrolidinyl, piperidinyl, piperazinyl, or morpholinyl, and is substituted with 0-3 substituents selected from halo, alkyl, and oxo.

5 Another aspect of the invention is a compound of Formula (III):



or a pharmaceutically acceptable salt thereof, wherein:

Ar¹ is phenyl substituted with 1-2 R^{1a} and 1-2 R^{1b} or 6-membered heteroaryl with 1-3 nitrogen atoms and substituted with 1R^{1a} and 1-2 R^{1b};

10 Ar³ is phenyl substituted with 1-3 R^{3a} or 5- to 6-membered heteroaryl with 1-3 nitrogen atoms and substituted with 1-3 R^{3a};

R^{1a} is hydrogen or halo;

R^{1b} is halo, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy;

15 R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy;

R^{3a} is cyano, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ alkoxyalkyl, C₁₋₄ haloalkyl, (R¹R²N)C₁₋₄ alkyl, R¹R²N, C₁₋₄ alkylC(O)(R²)NC₁₋₄ alkyl, (C₁₋₄ alkyl)₂(O)P, (C₁₋₄ alkoxy)₂(O)P, (C₁₋₄ alkoxy)(C₁₋₄ alkyl)(O)P, C₁₋₄ alkylSO₂, or C₁₋₄ alkylSO₂NH;

20 R¹R²N taken together is oxazolyl or pyrrolidinyl and is substituted with 0-3 substituents selected from halo, alkyl, and oxo;

R^{4a} or R^{4b} is independently hydrogen, C₁₋₄ alkyl, or C₁₋₄ hydroxylalkyl; alternatively, R^{4a} and R^{4b} together with the carbon atom they are both attached to form a C₃₋₆ cycloalkyl; and


25 R^{5a} or R^{5b} is independently hydrogen, C₁₋₄ alkyl, C₁₋₄ hydroxylalkyl, or C₁₋₄ alkoxyalkyl.

Another aspect of the invention is a compound of Formula (II) or (III), or a pharmaceutically acceptable salt thereof, wherein:

Ar¹ is phenyl substituted with 1-2 R^{1a} and 1-2 R^{1b}, pyridinyl substituted with 1 R^{1a} and 1-2 R^{1b}, or pyrazinyl substituted with 1 R^{1a} and 1-2 R^{1b}; and

5 Ar³ is phenyl substituted with 1-3 R^{3a}, pyrazolyl substituted with 1-3 R^{3a}, pyridinyl substituted with 1-3 R^{3a}, or pyrimidinyl substituted with 1-3 R^{3a}.

Another aspect of the invention is a compound of Formula (IV):



10 or a pharmaceutically acceptable salt thereof, wherein:

Ar¹ is phenyl substituted with 1-2 R^{1a} and 1-2 R^{1b}, pyridinyl substituted with 1 R^{1a} and 1-2 R^{1b}, or pyrazinyl substituted with 1 R^{1a} and 1-2 R^{1b};

R^{1a} is hydrogen or halo;

R^{1b} is halo, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy;


15 R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy;

R^{3a} is cyano, halo, C₁₋₃ alkyl, C₁₋₃ hydroxyalkyl, C₁₋₃ alkoxyalkyl, C₁₋₃ haloalkyl, R¹R²N, (C₁₋₃ alkyl)₂(O)P, (C₁₋₃ alkoxy)₂(O)P, (C₁₋₃ alkoxy)(C₁₋₃ alkyl)(O)P, C₁₋₃ alkylSO₂, or C₁₋₃ alkylSO₂NH;

20 R^{4a} or R^{4b} is independently hydrogen, C₁₋₃ alkyl, or C₁₋₃ hydroxylalkyl; alternatively, R^{4a} and R^{4b} together with the carbon atom they are both attached to form a C₃₋₆ cycloalkyl; and

R^{5a} or R^{5b} is independently hydrogen, C₁₋₃ alkyl, C₁₋₃ hydroxylalkyl, or C₁₋₃ alkoxyalkyl.

Another aspect of the invention is a compound of Formula (V):



or a pharmaceutically acceptable salt thereof, wherein:


- 5 R^{1a} is hydrogen or F;
 R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy;
 R^{2a} is hydrogen, halo, C₁₋₃ alkyl, C₁₋₃ haloalkyl, or C₃₋₆ cycloalkyl;
 R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂
 10 alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH; and
 R^{3a'} is halo.

Another aspect of the invention is a compound of Formula (V), or a pharmaceutically acceptable salt thereof, wherein:

- 15 R^{1a} is hydrogen or F;
 R^{1b} is F, Cl, or CF₃;
 R^{2a} is hydrogen, F, Cl, isopropyl, CF₃, or cyclopropyl;
 R^{3a} is (CH₃)₂(O)P, (CH₃CH₂)₂(O)P, (CH₃CH₂O)(CH₃)(O)P, CH₃SO₂, or
 CH₃SO₂NH; and
 R^{3a'} is F.

20

Another aspect of the invention is a compound of Formula (VI):



or a pharmaceutically acceptable salt thereof, wherein:


R^{1a} is hydrogen or halo;

R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy;

5 R^{2a} is hydrogen, halo, C₁₋₃ alkyl, or C₃₋₆ cycloalkyl;

R^{3a} is (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

Another aspect of the invention is a compound of Formula (VII):



or a pharmaceutically acceptable salt thereof, wherein:

Ar³ is pyrazolyl substituted with 1-3 R^{3a}, pyridinyl substituted with 1-3 R^{3a}, or pyrimidinyl substituted with 1-3 R^{3a};

R^{1a} is hydrogen or halo;

15 R^{1b} is halo, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy;

R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy;

R^{3a} is cyano, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ alkoxyalkyl, (R¹R²N)C₁₋₄ alkyl, R¹R²N-, (C₁₋₄ alkyl)₂(O)P, (C₁₋₄ alkoxy)₂(O)P, (C₁₋₄ alkoxy)(C₁₋₄ alkyl)(O)P, C₁₋₄ alkylSO₂, or C₁₋₄ alkylSO₂NH;


R¹ is hydrogen or alkyl;

R² is hydrogen or alkyl; or R¹R²N taken together is oxazolyl or pyrrolidinyl, and is substituted with 0-3 substituents selected from halo, alkyl, or oxo;

R^{4a} or R^{4b} is independently hydrogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkoxyalkyl, or C₁₋₄ haloalkoxy; R^{4a} and R^{4b} together with the carbon atom to which they are both attached form a C₃₋₆ cycloalkyl; and

R^{5a} or R^{5b} is independently hydrogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkoxyalkyl, or C₁₋₄ haloalkoxy.

Another aspect of the invention is a compound of Formula (VIII):



(VIII)

or a pharmaceutically acceptable salt thereof, wherein:

R^{1a} is hydrogen or halo;


R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy;

R^{2a} is hydrogen, halo, C₁₋₃ alkyl, or C₃₋₆ cycloalkyl;

R^{3a} is (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH; and

R^{5a} or R^{5b} is independently hydrogen or C₁₋₂ alkyl.

Another aspect of the invention is a compound of Formula (IX):




or a pharmaceutically acceptable salt thereof, wherein:

5 R¹ᵃ is hydrogen or halo;

R¹ᵇ is halo, C₁-₂ haloalkyl, or C₁-₂ alkoxy;

R²ᵃ is hydrogen, halo, C₁-₂ haloalkyl, C₁-₃ alkyl, or C₃-₆ cycloalkyl;



R³ᵃ is (C₁-₂ alkyl)₂(O)P, (C₁-₂ alkoxy)(C₁-₂ alkyl)(O)P, or C₁-₂ alkylSO₂NH;

10 R³ᵃ' is halo;

R⁴ᵃ or R⁴ᵇ is independently hydrogen, C₁-₂ alkyl, or C₁-₂ hydroxyalkyl; or R⁴ᵃ and R⁴ᵇ together with the carbon atom to which they are both attached form a C₃-₆ cycloalkyl; and

15 R⁵ᵃ or R⁵ᵇ is independently hydrogen, C₁-₂ alkyl, C₁-₂ hydroxyalkyl, or C₁-₂ alkoxyalkyl.

Another aspect of the invention is a compound of Formula (II), (III), (IV), (VII), or (IX), or a pharmaceutically acceptable salt thereof, wherein:

20 R⁴ᵃ is hydrogen;

R⁴ᵇ is hydrogen;

R⁵ᵃ is C₁-₂ alkyl, C₁-₂ hydroxyalkyl, or C₁-₂ alkoxyalkyl; and

R⁵ᵇ is hydrogen.

Another aspect of the invention is a compound of formula (II), (III), (IV), (VII), or (IX), or a pharmaceutically acceptable salt thereof, wherein:


R^{4a} is C₁₋₂ alkyl;


5 R^{4b} is C₁₋₂ alkyl; or R^{4a} and R^{4b} are taken together to form a cyclopropyl;


R^{5a} is hydrogen; and

R^{5b} is hydrogen.

Another aspect of the invention is a compound selected from the group consisting







5 For a compound of Formula (I), (II), (III), (IV), (V), (VI), (VII), (VIII), or (IX), the scope of any instance of a variable substituent, including Ar^1 , Ar^2 , Ar^3 , R^{1a} , R^{1b} , R^{2a} ,

R^{3a}, R^{4a}, R^{4b}, R^{5a}, R^{5b}, R¹, R² can be used independently with the scope of any other instance of a variable substituent. As such, the invention includes combinations of the different aspects.

In one non-limiting embodiment, for a compound of Formula (II), (III), (IV),
5 (VII), or (IX), R^{4a}, R^{4b}, R^{5a}, R^{5b} are all hydrogen; Ar¹ is phenyl, pyridinyl, or pyrazinyl, each substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is phenyl, pyrazolyl, pyridinyl, or pyrimidinyl, each substituted with 1-4 R^{3a}; R^{3a} is
10 halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV),
(VII), or (IX), R^{4a}, R^{4b}, R^{5a}, R^{5b} are all hydrogen; Ar¹ is phenyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl
15 substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrazolyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV),
20 (VII), or (IX), R^{4a}, R^{4b}, R^{5a}, R^{5b} are all hydrogen; Ar¹ is phenyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrimidinyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P,
25 (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV),
(VII), or (IX), R^{4a}, R^{4b}, R^{5a}, R^{5b} are all hydrogen; Ar¹ is phenyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄
30 haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is phenyl, pyrazolyl, pyridinyl, or pyrimidinyl, each substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl,

alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a}, R^{4b}, R^{5a}, R^{5b} are all hydrogen; Ar¹ is pyridinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is phenyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a}, R^{4b}, R^{5a}, R^{5b} are all hydrogen; Ar¹ is pyridinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrazolyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a}, R^{4b}, R^{5a}, R^{5b} are all hydrogen; Ar¹ is pyridinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyridinyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a}, R^{4b}, R^{5a}, R^{5b} are all hydrogen; Ar¹ is pyridinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrimidinyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a}, R^{4b}, R^{5a}, R^{5b} are all hydrogen; Ar¹ is pyrazinyl substituted with 1R^{1a}

and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is phenyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a}, R^{4b}, R^{5a}, R^{5b} are all hydrogen; Ar¹ is pyrazinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrazolyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a}, R^{4b}, R^{5a}, R^{5b} are all hydrogen; Ar¹ is pyrazinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyridinyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a}, R^{4b}, R^{5a}, R^{5b} are all hydrogen; Ar¹ is pyrazinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrimidinyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In one non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} is methyl, R^{4b} is methyl; or R^{4a} is hydrogen, R^{4b} is hydroxymethyl; or R^{4a} and R^{4b} are taken together to form a cyclopropyl; R^{5a} and R^{5b} are all hydrogen; Ar¹ is phenyl, pyridinyl, or pyrazinyl, each substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a}

is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is phenyl, pyrazolyl, pyridinyl, or pyrimidinyl, each substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

5 In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} is methyl, R^{4b} is methyl; or R^{4a} is hydrogen, R^{4b} is hydroxymethyl; or R^{4a} and R^{4b} are taken together to form a cyclopropyl; R^{5a} and R^{5b} are all hydrogen; Ar¹ is phenyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrazolyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

15 In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} is methyl, R^{4b} is methyl; R^{4a} is hydrogen, R^{4b} is hydroxymethyl; or R^{4a} and R^{4b} are taken together to form a cyclopropyl; R^{5a} and R^{5b} are all hydrogen; Ar¹ is phenyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrimidinyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

25 In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} is methyl, R^{4b} is methyl; R^{4a} is hydrogen, R^{4b} is hydroxymethyl; or R^{4a} and R^{4b} are taken together to form a cyclopropyl; R^{5a} and R^{5b} are all hydrogen; Ar¹ is phenyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is phenyl, pyrazolyl, pyridinyl, or pyrimidinyl, each substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} is methyl, R^{4b} is methyl; R^{4a} is hydrogen, R^{4b} is hydroxymethyl; or R^{4a} and R^{4b} are taken together to form a cyclopropyl; R^{5a} and R^{5b} are all hydrogen; Ar¹ is pyridinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is phenyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

10 In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} is methyl, R^{4b} is methyl; R^{4a} is hydrogen, R^{4b} is hydroxymethyl; or R^{4a} and R^{4b} are taken together to form a cyclopropyl; R^{5a} and R^{5b} are all hydrogen; Ar¹ is pyridinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrazolyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

20 In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} is methyl, R^{4b} is methyl; R^{4a} is hydrogen, R^{4b} is hydroxymethyl; or R^{4a} and R^{4b} are taken together to form a cyclopropyl; R^{5a} and R^{5b} are all hydrogen; Ar¹ is pyridinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyridinyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

30 In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} is methyl, R^{4b} is methyl; R^{4a} is hydrogen, R^{4b} is hydroxymethyl; or R^{4a} and R^{4b} are taken together to form a cyclopropyl; R^{5a} and R^{5b} are all hydrogen; Ar¹ is pyridinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋

4 alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrimidinyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

5 In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} is methyl, R^{4b} is methyl; R^{4a} is hydrogen, R^{4b} is hydroxymethyl; or R^{4a} and R^{4b} are taken together to form a cyclopropyl; R^{5a} and R^{5b} are all hydrogen; Ar¹ is pyrazinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is phenyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

15 In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} is methyl, R^{4b} is methyl; R^{4a} is hydrogen, R^{4b} is hydroxymethyl; or R^{4a} and R^{4b} are taken together to form a cyclopropyl; R^{5a} and R^{5b} are all hydrogen; Ar¹ is pyrazinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrazolyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

25 In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} is methyl, R^{4b} is methyl; R^{4a} is hydrogen, R^{4b} is hydroxymethyl; or R^{4a} and R^{4b} are taken together to form a cyclopropyl; R^{5a} and R^{5b} are all hydrogen; Ar¹ is pyrazinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyridinyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} is methyl, R^{4b} is methyl; R^{4a} is hydrogen, R^{4b} is hydroxymethyl; or R^{4a} and R^{4b} are taken together to form a cyclopropyl; R^{5a} and R^{5b} are all hydrogen; Ar¹ is pyrazinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrimidinyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

10

In one non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} and R^{4b} are all hydrogen; R^{5a} is hydrogen, R^{5b} is methyl or hydroxymethyl; Ar¹ is phenyl, pyridinyl, or pyrazinyl, each substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is phenyl, pyrazolyl, pyridinyl, or pyrimidinyl, each substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

20

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} and R^{4b} are all hydrogen; R^{5a} is hydrogen, R^{5b} is methyl or hydroxymethyl; Ar¹ is phenyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrazolyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

25

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV), (VII), or (IX), R^{4a} and R^{4b} are all hydrogen; R^{5a} is hydrogen, R^{5b} is methyl or hydroxymethyl; Ar¹ is phenyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy,

30

or C₁₋₄ haloalkoxy; Ar³ is pyrimidinyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV),
5 (VII), or (IX), R^{4a} and R^{4b} are all hydrogen; R^{5a} is hydrogen, R^{5b} is methyl or hydroxymehtyl; Ar¹ is phenyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is phenyl, pyrazolyl, pyridinyl, or pyrimidinyl, each substituted
10 with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV),
(VII), or (IX), R^{4a} and R^{4b} are all hydrogen; R^{5a} is hydrogen, R^{5b} is methyl or hydroxymehtyl; Ar¹ is pyridinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or
15 halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is phenyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV),
(VII), or (IX), R^{4a} and R^{4b} are all hydrogen; R^{5a} is hydrogen, R^{5b} is methyl or hydroxymehtyl; Ar¹ is pyridinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or
halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a}
25 is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrazolyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV),
(VII), or (IX), R^{4a} and R^{4b} are all hydrogen; R^{5a} is hydrogen, R^{5b} is methyl or
30 hydroxymehtyl; Ar¹ is pyridinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄

alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyridinyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV),
5 (VII), or (IX), R^{4a} and R^{4b} are all hydrogen; R^{5a} is hydrogen, R^{5b} is methyl or hydroxymehtyl; Ar¹ is pyridinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrimidinyl substituted with 1-4 R^{3a}; R^{3a} is halo,
10 hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV),
(VII), or (IX), R^{4a} and R^{4b} are all hydrogen; R^{5a} is hydrogen, R^{5b} is methyl or hydroxymehtyl; Ar¹ is pyrazinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or
15 halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is phenyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV),
(VII), or (IX), R^{4a} and R^{4b} are all hydrogen; R^{5a} is hydrogen, R^{5b} is methyl or hydroxymehtyl; Ar¹ is pyrazinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or
halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄
25 alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrazolyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV),
(VII), or (IX), R^{4a} and R^{4b} are all hydrogen; R^{5a} is hydrogen, R^{5b} is methyl or
30 hydroxymehtyl; Ar¹ is pyrazinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄

alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyridinyl substituted with 1-4 R^{3a}; R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In another non-limiting embodiment, for a compound of Formula (II), (III), (IV),
5 (VII), or (IX), R^{4a} and R^{4b} are all hydrogen; R^{5a} is hydrogen, R^{5b} is methyl or hydroxymehtyl; Ar¹ is pyrazinyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen or halo; R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy; Ar² is phenyl substituted with 1-4 R^{2a}; R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy; Ar³ is pyrimidinyl substituted with 1-4 R^{3a}; R^{3a} is halo,
10 hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH.

In one preferred embodiment, for a compound of Formula (II), (III), (IV), or (VII), or (IX), R^{4a} is methyl; R^{4b} is methyl; R^{4a} is hydrogen; R^{4b} is hydroxymethyl; or R^{4a} and R^{4b} are taken together to form a cyclopropyl; R^{5a} is hydrogen, R^{5b} is methyl or
15 hydroxymehtyl; R^{5a} is hydrogen; R^{5b} is hydrogen; or R^{4a}, R^{4b}, R^{5a}, R^{5b} are all hydrogen; Ar¹ is phenyl substituted with 1R^{1a} and 1-2 R^{1b}; R^{1a} is hydrogen, or F; R^{1b} is F, Cl, or CF₃; Ar² is phenyl substituted with 1-2 R^{2a}; R^{2a} is hydrogen, F, Cl, isopropyl, CF₃ CF₃, cyclopropyl; Ar³ is phenyl or pyridinyl, each substituted with 1-2 R^{3a}; R^{3a} is F, (CH₃)₂(O)P, (CH₃CH₂)₂(O)P, (CH₃CH₂O)(CH₃)(O)P, CH₃SO₂, or CH₃SO₂NH.

20

Unless specified otherwise, these terms have the following meanings. "Alkyl" means a straight or branched alkyl group composed of 1 to 6 carbons. "Alkenyl" means a straight or branched alkyl group composed of 2 to 6 carbons with at least one double bond. "Alkynyl" means a straight or branched alkyl group composed of 2 to 6 carbons
25 with at least one triple bond. "Cycloalkyl" means a monocyclic ring system composed of 3 to 7 carbons. Terms with a hydrocarbon moiety (e.g. alkoxy) include straight and branched isomers for the hydrocarbon portion. "Halo" includes fluoro, chloro, bromo, and iodo. "Haloalkyl" and "haloalkoxy" include all halogenated isomers from monohalo to perhalo. "Aryl" means a monocyclic or bicyclic aromatic hydrocarbon groups having 6
30 to 12 carbon atoms, or a bicyclic fused ring system wherein one or both of the rings is aromatic. Bicyclic fused ring systems consist of a phenyl group fused to a four- to seven-membered aromatic or non-aromatic carbocyclic ring. Representative examples of aryl

groups include but are not limited to phenyl, indanyl, indenyl, naphthyl, and tetrahydronaphthyl. "Heteroaryl" means a 5 to 7 membered monocyclic or 8 to 11 membered bicyclic aromatic ring system with 1-5 heteroatoms independently selected from nitrogen, oxygen, and sulfur. Where a bonding attachment location is not specified, the bonding may be attached at any appropriate location as understood by practitioners in the art. Combinations of substituents and bonding patterns are only those that result in stable compounds as understood by practitioners in the art. Parenthetical and multiparenthetical terms are intended to clarify bonding relationships to those skilled in the art. For example, a term such as ((R)alkyl) means an alkyl substituent further substituted with the substituent R.

The invention includes all pharmaceutically acceptable salt forms of the compounds. Pharmaceutically acceptable salts are those in which the counter ions do not contribute significantly to the physiological activity or toxicity of the compounds and as such function as pharmacological equivalents. These salts can be made according to common organic techniques employing commercially available reagents. Some anionic salt forms include acetate, acistrate, besylate, bromide, chloride, citrate, fumarate, glucouronate, hydrobromide, hydrochloride, hydroiodide, iodide, lactate, maleate, mesylate, nitrate, pamoate, phosphate, succinate, sulfate, tartrate, tosylate, and xinofoate. Some cationic salt forms include ammonium, aluminum, benzathine, bismuth, calcium, choline, diethylamine, diethanolamine, lithium, magnesium, meglumine, 4-phenylcyclohexylamine, piperazine, potassium, sodium, tromethamine, and zinc.

Some of the compounds of the invention exist in stereoisomeric forms including the structure below with the indicated carbon. The invention includes all stereoisomeric forms of the compounds including enantiomers and diastereomers. Methods of making and separating stereoisomers are known in the art. The invention includes all tautomeric forms of the compounds. The invention includes atropisomers and rotational isomers.

The invention is intended to include all isotopes of atoms occurring in the compounds. Isotopes include those atoms having the same atomic number but different mass numbers. By way of general example and without limitation, isotopes of hydrogen include deuterium and tritium. Isotopes of carbon include ^{13}C and ^{14}C . Isotopically-labeled compounds of the invention can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described herein,

using an appropriate isotopically-labeled reagent in place of the non-labeled reagent otherwise employed. Such compounds may have a variety of potential uses, for example as standards and reagents in determining biological activity. In the case of stable isotopes, such compounds may have the potential to favorably modify biological, pharmacological, or pharmacokinetic properties.

BIOLOGICAL METHODS

N-formyl peptide receptors (FPRs) are a family of chemo attractant receptors that facilitate leukocyte response during inflammation. FPRs belong to the seven-transmembrane G protein-coupled receptor superfamily and are linked to inhibitory G-proteins (Gi). Three family members (FPR1, FPR2 and FPR3) have been identified in humans and are predominantly found in myeloid cells with varied distribution and have also been reported in multiple organs and tissues. After agonist binding, the FPRs activate a multitude of physiological pathways, such as intra cellular signaling transduction, Ca^{2+} mobilization and transcription. The family interacts with a diverse set of ligands that includes proteins, polypeptides and fatty acid metabolites which activate both pro-inflammatory and pro-resolution downstream responses. FPR2 and FPR1 Cyclic Adenosine Monophosphate (cAMP) Assays were used to measure the activity of the compounds in this patent.

FPR2 and FPR1 Cyclic Adenosine Monophosphate (cAMP) Assays. A mixture of forskolin (5 μM final for FPR2 or 10 μM final for FPR1) and IBMX (200 μM final) were added to 384-well Proxiplates (Perkin-Elmer) pre-dotted with test compounds in DMSO (1% final) at final concentrations in the range of 0.020 nM to 100 μM . Chinese Hamster Ovary cells (CHO) overexpressing human FPR1 or human FPR2 receptors were cultured in F-12 (Ham's) medium supplemented with 10% qualified FBS, 250 $\mu\text{g}/\text{ml}$ zeocin and 300 $\mu\text{g}/\text{ml}$ hygromycin (Life Technologies). Reactions were initiated by adding 2,000 human FPR2 cells per well or 4,000 human FPR1 cells per well in Dulbecco's PBS (with calcium and magnesium) (Life Technologies) supplemented with 0.1% BSA (Perkin-Elmer). The reaction mixtures were incubated for 30 min at room temperature. The level of intracellular cAMP was determined using the HTRF HiRange cAMP assay reagent kit (Cisbio) according to manufacturer's instruction. Solutions of cryptate conjugated anti-cAMP and d2 fluorophore-labelled cAMP were made in a supplied lysis buffer

separately. Upon completion of the reaction, the cells were lysed with equal volume of the d2-cAMP solution and anti-cAMP solution. After a 1-h room temperature incubation, time-resolved fluorescence intensity was measured using the Envision (Perkin-Elmer) at 400 nm excitation and dual emission at 590 nm and 665 nm. A calibration curve was constructed with an external cAMP standard at concentrations ranging from 1 μ M to 0.1 pM by plotting the fluorescent intensity ratio from 665 nm emission to the intensity from the 590 nm emission against cAMP concentrations. The potency and activity of a compound to inhibit cAMP production was then determined by fitting to a 4-parametric logistic equation from a plot of cAMP level versus compound concentrations.

The examples disclosed below were tested in the FPR2 and FPR1 cAMP assay described above and found having FPR2 and/or FPR1 agonist activity. Table 1 below lists EC₅₀ values in the FPR2 and FPR1 cAMP assays measured for the following examples.

Table 1

Example	hFPR2 cAMP2 EC ₅₀ (μ M)	hFPR1 cAMP EC ₅₀ (μ M)
1	0.00043	0.20
2	0.0042	0.36
3	0.0011	0.22
4	0.00093	1.7
5	0.00068	0.23
6	0.0010	0.19
7	0.0021	0.14
8	0.0020	0.061
9	0.0021	0.031
10	0.0022	0.13
11	0.0026	0.29
12	0.0025	0.45
13	0.0065	2.5
14	0.0072	0.60
15	0.010	1.5
16	0.012	0.12
17	0.016	4.1
18	0.020	2.1
19	0.022	0.078
20	0.024	>10
21	0.031	0.39
22	0.033	1.5
23	0.0082	0.70

24	0.022	1.4
25	0.00053	0.26
26	0.00055	0.047
27	0.00050	0.11
28	0.0098	0.55
29	0.0070	0.19
30	0.00022	0.083
31	0.0040	0.43
32	0.00091	0.25
33	0.0011	0.47
34	0.0041	0.026
35	0.0086	0.052
36	0.0047	0.069
37	0.00140	0.17
38	0.0035	0.042
39	0.00058	0.057
40	0.0046	0.050
41	0.0036	0.50
42	0.0010	0.22
43	0.0033	0.12
44	0.043	0.45
45	0.0030	0.14
46	0.0048	0.093
47	0.0028	0.064
48	0.0037	0.11
49	0.0056	3.4
50	0.014	0.64
51	0.0082	1.3
52	0.010	1.1
53	0.0078	2.7
54	0.0055	0.90
55	0.022	>10
56	0.0075	7.7
57	0.00095	1.0
58	0.0063	0.61
59	0.0017	0.14
60	0.0037	0.12
61	0.023	0.37
62	0.031	0.72
63	0.00567	4.4
64	0.0088	2.9
65	0.0012	2.6
66	0.0049	0.58
67	0.00248	1.3
68	0.00053	0.15
69	0.022	5.0
70	0.000077	0.020

71	0.0021	1.28
72	0.00061	0.054
73	0.00089	0.17
74	0.0022	0.14
75	0.0034	1.18
76	0.0041	2.1
77	0.0012	0.079
78	0.0012	0.045
79	0.039	>10
80	0.0014	0.086
81	0.00052	0.027
82	0.0025	0.78
83	0.00041	0.00099
84	0.0060	0.28
85	0.00062	0.0051
86	0.00085	0.021
87	0.0016	0.0012
88	0.0053	0.0049
89	0.0044	0.025
90	0.0051	1.4
91	0.0057	0.20
92	0.0083	0.22
93	0.012	0.78
94	0.0018	0.012
95	0.032	0.20
96	0.0029	0.075
97	0.0072	0.11
98	0.0029	0.27
99	0.00079	0.24
100	0.0020	0.092
101	0.00066	0.0011
102	0.0017	0.57
103	0.00057	0.35
104	0.0052	0.049
105	0.0069	0.071
106	0.0031	0.0035
107	0.011	0.0058
108	0.0014	0.053
109	0.00044	0.39
110	0.0011	0.64
111	0.030	0.54
112	0.00027	0.39
113	0.00039	0.59
114	0.00086	0.27
115	0.0012	0.055
116	0.00121	0.89
117	0.013	0.084

118	0.017	0.32
119	0.014	0.19
120	0.00078	0.16
121	0.00030	0.0050
122	0.00081	0.61
123	0.0015	0.24
124	0.00079	0.76
125	0.00099	0.18
126	0.012	>10
127	0.018	0.92
128	0.0010	0.024
129	0.0024	0.019
130	0.0057	3.0
131	0.0012	0.061
132	0.00048	0.028
133	0.15	>5
134	0.0025	1500
135	0.0029	720

PHARMACEUTICAL COMPOSITIONS AND METHODS OF USE

The compounds of the present invention may be administered to mammals,
5 preferably humans, for the treatment of a variety of conditions and disorders associated
with the FPR2 receptor such as Behcet's disease, Sweet disease, systemic lupus
erythematosus (SLE), Wegener's granulomatosis, virus infection, diabetes, amputations,
cancers, bacterial infection, physical external injuries, physical disorders including
exposure to radiation, vasoconstriction, anaphylactic reactions, allergic reactions, rhinitis,
10 shocks (endotoxic, hemorrhagic, traumatic, splanchnic ischemia, and circulatory shocks),
rheumatoid arthritis, gout, psoriasis, benign prostatic hyperplasia, myocardial ischemia,
myocardial infarction, heart failure, brain injuries, pulmonary diseases, COPD, COAD,
COLD, acute lung injury, acute respiratory distress syndrome, chronic bronchitis,
pulmonary emphysema, asthma (allergic asthma and non-allergic asthma), cystic fibrosis,
15 kidney fibrosis, nephropathy, renal glomerular diseases, ulcerative colitis, IBD, Crohn's
disease, periodontitis, pains, Alzheimer's disease, AIDS, uveitic glaucoma, conjunctivitis,
Sjogren's syndrome, rhinitis, atherosclerosis, neuroinflammatory diseases including
multiple sclerosis, stroke, sepsis, and the like.

Unless otherwise specified, the following terms have the stated meanings. The
20 term "subject" refers to any human or other mammalian species that could potentially

benefit from treatment with a FPR2 and/or FPR1 agonist as understood by practitioners in this field. Some subjects include human beings of any age with risk factors for cardiovascular disease. Common risk factors include age, sex, weight, family history, sleep apnea, alcohol or tobacco use, physical inactivity arrhythmia or signs of insulin resistance such as acanthosis nigricans, hypertension, dyslipidemia, or polycystic ovary syndrome (PCOS). The term "patient" means a person suitable for therapy as determined by practitioners in the field. "Treating" or "treatment" cover the treatment of a patient or subject as understood by practitioners in this field. "Preventing" or "prevention" cover the preventive treatment (*i.e.*, prophylaxis and/or risk reduction) of a subclinical disease-state in a patient or subject aimed at reducing the probability of the occurrence of a clinical disease-state as understood by practitioners in this field. Patients are selected for preventative therapy based on factors that are known to increase risk of suffering a clinical disease state compared to the general population. "Therapeutically effective amount" means an amount of a compound that is effective as understood by practitioners in this field.

Another aspect of the invention are pharmaceutical compositions comprising a therapeutically effective amount of a compound of Formulae (I)-(IX) in combination with a pharmaceutical carrier.

Another aspect of the invention are pharmaceutical compositions comprising a therapeutically effective amount of a compound of Formulae (I)-(IX) in combination with at least one other therapeutic agent and a pharmaceutical carrier.

"Pharmaceutical composition" means a composition comprising a compound of the invention in combination with at least one additional pharmaceutically acceptable carrier. A "pharmaceutically acceptable carrier" refers to media generally accepted in the art for the delivery of biologically active agents to animals, in particular, mammals, including, *i.e.*, adjuvant, excipient or vehicle, such as diluents, preserving agents, fillers, flow regulating agents, disintegrating agents, wetting agents, emulsifying agents, suspending agents, sweetening agents, flavoring agents, perfuming agents, anti-bacterial agents, anti-fungal agents, lubricating agents and dispensing agents, depending on the nature of the mode of administration and dosage forms.

Pharmaceutically acceptable carriers are formulated according to a number of factors well within the purview of those of ordinary skill in the art. These include, without

limitation: the type and nature of the active agent being formulated; the subject to which the agent-containing composition is to be administered; the intended route of administration of the composition; and the therapeutic indication being targeted.

Pharmaceutically acceptable carriers include both aqueous and non-aqueous liquid media, as well as a variety of solid and semi-solid dosage forms. Such carriers can include a number of different ingredients and additives in addition to the active agent, such additional ingredients being included in the formulation for a variety of reasons, *e.g.*, stabilization of the active agent, binders, etc., well known to those of ordinary skill in the art. Descriptions of suitable pharmaceutically acceptable carriers, and factors involved in their selection, are found in a variety of readily available sources such as, for example, Allen, L.V., Jr. et al., *Remington: The Science and Practice of Pharmacy* (2 Volumes), 22nd Edition, Pharmaceutical Press (2012).

Particularly when provided as a single dosage unit, the potential exists for a chemical interaction between the combined active ingredients. For this reason, when the compound of the present invention and a second therapeutic agent are combined in a single dosage unit they are formulated such that although the active ingredients are combined in a single dosage unit, the physical contact between the active ingredients is minimized (that is, reduced). For example, one active ingredient may be enteric coated. By enteric coating one of the active ingredients, it is possible not only to minimize the contact between the combined active ingredients, but also, it is possible to control the release of one of these components in the gastrointestinal tract such that one of these components is not released in the stomach but rather is released in the intestines. One of the active ingredients may also be coated with a material that affects a sustained-release throughout the gastrointestinal tract and also serves to minimize physical contact between the combined active ingredients. Furthermore, the sustained-released component can be additionally enteric coated such that the release of this component occurs only in the intestine. Still another approach would involve the formulation of a combination product in which the one component is coated with a sustained and/or enteric release polymer, and the other component is also coated with a polymer such as a low viscosity grade of hydroxypropyl methylcellulose (HPMC) or other appropriate materials as known in the art, in order to further separate the active components. The polymer coating serves to form an additional barrier to interaction with the other component.

Another aspect of the invention is a method for treating heart disease comprising administering a therapeutically effective amount of a compound of Formula (I) to a patient.

Another aspect of the invention is a method for treating heart disease wherein the heart disease is selected from the group consisting of angina pectoris, unstable angina, myocardial infarction, heart failure, acute coronary disease, acute heart failure, chronic heart failure, and cardiac iatrogenic damage.

It will be understood that treatment or prophylaxis of heart failure may involve treatment or prophylaxis of a cardiovascular event as well. Treatment or prophylaxis as referred to herein may refer to treatment or prophylaxis of certain negative symptoms or conditions associated with or arising as a result of a cardiovascular event. By way of example, treatment or prophylaxis may involve reducing or preventing negative changes in fractional shortening, heart weight, lung weight, myocyte cross sectional area, pressure overload induced cardiac fibrosis, stress induced cellular senescence, and/or cardiac hypertrophy properties, or any combination thereof, associated with or arising as a result of a cardiovascular event. Treatment may be administered in preparation for or in response to a cardiovascular event to alleviate negative effects. Prevention may involve a pro-active or prophylactic type of treatment to prevent the cardiovascular event or to reduce the onset of negative effects of a cardiovascular event.

In one embodiment, the present invention provides the use of compounds of Formulae (I)-(IX) or a pharmaceutically acceptable salt thereof for the preparation of a pharmaceutical composition for the treatment or prophylaxis of heart failure, for example, heart failure results from hypertension, an ischemic heart disease, a non-ischemic heart disease, exposure to a cardiotoxic compound, myocarditis, Kawasaki's disease, Type I and Type II diabetes, thyroid disease, viral infection, gingivitis, drug abuse, alcohol abuse, pericarditis, atherosclerosis, vascular disease, hypertrophic cardiomyopathy, dilated cardiomyopathy, myocardial infarction, atrial fibrosis, left ventricular systolic dysfunction, left ventricular diastolic dysfunction, coronary bypass surgery, pacemaker implantation surgery, starvation, an eating disorder, muscular dystrophies, and a genetic defect. Preferably, the heart failure to be treated is diastolic heart failure, heart failure with reduced ejection fraction (HFrEF), heart failure with preserved ejection fraction

(HFpEF), acute heart failure, and chronic heart failure of ischemic and non-ischemic origin.

In one embodiment, the present invention provides the use of compounds of Formulae (I)-(IX) to treat systolic and/or diastolic dysfunction, wherein the compound is administered in a therapeutically effective amount to increase the ability of the cardiac muscle cells to contract and relax thereby increasing the filling and emptying of both the right and left ventricles, preferably, the left ventricle.

In another embodiment, the present invention provides the use of compounds of Formulae (I)-(IX) to treat heart failure wherein the compound is administered in a therapeutically effective amount to increase ejection fraction in the left ventricle.

In still another embodiment, the present invention provides the use of compounds of Formulae (I)-(IX) to treat heart failure wherein the compound is administered in a therapeutically effective amount to reduce fibrosis in heart tissue.

Another aspect of the invention is a method for treating heart disease wherein the treatment is post myocardial infarction.

Another aspect of the invention is a method for treating dermatological diseases including, but not limited to, rosacea, rosacea fulminans, sunburn, psoriasis, menopause-associated hot flashes, flushing and redness associated with hot flashes, erythema associated with hot flashes, hot flashes resulting from orchiectomyatopic dermatitis, treatment of redness and itch from insect bites, photoaging, seborrheic dermatitis, acne, allergic dermatitis, telangiectasia (dilations of previously existing small blood vessels) of the face, angioectasias, rhinophyma (hypertrophy of the nose with follicular dilation), acne-like skin eruptions (may ooze or crust), burning or stinging sensation, erythema of the skin, cutaneous hyperactivity with dilation of blood vessels of the skin, Lyell's syndrome, Stevens-Johnson syndrome, local itching and discomfort associated with hemorrhoids, hemorrhoids, erythema multiforme minor, erythema multiforme major, erythema nodosum, eye puffiness, urticaria, pruritis, purpura, varicose veins, contact dermatitis, atopic dermatitis, nummular dermatitis, generalized exfoliative dermatitis, stasis dermatitis, lichen simplex chronicus, perioral dermatitis, pseudofolliculitis barbae, granuloma annulare, actinic keratosis, basal cell carcinoma, squamous cell carcinoma, eczema, dermal wound healing, hypertrophic scars, keloids, burns, rosacea, atopic dermatitis, acne, psoriasis, seborrheic dermatitis, actinic keratoses, basal cell carcinoma,

squamous cell carcinoma, melanoma, viral warts, photoaging, photodamage, melasma, post-inflammatory hyperpigmentation, other disorders of pigmentation, and alopecia (scarring and non-scarring forms). The compounds below would be expected to have therapeutic effects in many different types of skin disease, but have been exemplified by demonstrating accelerated wound healing activity.

Another aspect of the invention is a method for treating heart disease comprising administering a therapeutically effective amount of a compound of Formula (I) to a patient in conjunction with other therapeutic agents.

The compounds of this invention can be administered by any suitable means, for example, orally, such as tablets, capsules (each of which includes sustained release or timed release formulations), pills, powders, granules, elixirs, tinctures, suspensions (including nanosuspensions, microsuspensions, spray-dried dispersions), syrups, and emulsions; sublingually; buccally; parenterally, such as by subcutaneous, intravenous, intramuscular, or intrasternal injection, or infusion techniques (*e.g.*, as sterile injectable aqueous or non-aqueous solutions or suspensions); nasally, including administration to the nasal membranes, such as by inhalation spray; topically, such as in the form of a cream or ointment; or rectally such as in the form of suppositories. They can be administered alone, but generally will be administered with a pharmaceutical carrier selected on the basis of the chosen route of administration and standard pharmaceutical practice.

The dosage regimen for the compounds of the present invention will, of course, vary depending upon known factors, such as the pharmacodynamic characteristics of the particular agent and its mode and route of administration; the species, age, sex, health, medical condition, and weight of the recipient; the nature and extent of the symptoms; the kind of concurrent treatment; the frequency of treatment; the route of administration, the renal and hepatic function of the patient, and the effect desired.

By way of general guidance, the daily oral dosage of each active ingredient, when used for the indicated effects, will range between about 0.01 to about 5000 mg per day, preferably between about 0.1 to about 1000 mg per day, and most preferably between about 0.1 to about 250 mg per day. Intravenously, the most preferred doses will range from about 0.01 to about 10 mg/kg/minute during a constant rate infusion. Compounds of

this invention may be administered in a single daily dose, or the total daily dosage may be administered in divided doses of two, three, or four times daily.

Dosage forms (pharmaceutical compositions) suitable for administration may contain from about 1 milligram to about 2000 milligrams of active ingredient per dosage unit. In these pharmaceutical compositions the active ingredient will ordinarily be present in an amount of about 0.1-95% by weight based on the total weight of the composition. A typical capsule for oral administration contains at least one of the compounds of the present invention (250 mg), lactose (75 mg), and magnesium stearate (15 mg). The mixture is passed through a 60 mesh sieve and packed into a No. 1 gelatin capsule. A typical injectable preparation is produced by aseptically placing at least one of the compounds of the present invention (250 mg) into a vial, aseptically freeze-drying and sealing. For use, the contents of the vial are mixed with 2 mL of physiological saline, to produce an injectable preparation.

The compounds of the present invention may be employed in combination with other suitable therapeutic agents useful in the treatment of the aforementioned diseases or disorders including: anti-atherosclerotic agents, anti-dyslipidemic agents, anti-diabetic agents, anti-hyperglycemic agents, anti-hyperinsulinemic agents, anti-thrombotic agents, anti-retinopathic agents, anti-neuropathic agents, anti-nephropathic agents, anti-ischemic agents, anti-hypertensive agents, anti-obesity agents, anti-hyperlipidemic agents, anti-hypertriglyceridemic agents, anti-hypercholesterolemic agents, anti-restenotic agents, anti-pancreatic agents, lipid lowering agents, anorectic agents, memory enhancing agents, anti-dementia agents, cognition promoting agents, appetite suppressants, agents for treating heart failure, agents for treating peripheral arterial disease, agents for treating malignant tumors, and anti-inflammatory agents.

The compounds of the invention may be used with at least one of the following heart failure agents selected from loop diuretics, Angiotensin converting enzyme (ACE) inhibitors, Angiotensin II receptor blockers (ARBs), angiotensin receptor-neprilysin inhibitors (ARNI), beta blockers, mineralocorticoid receptor antagonists, nitroxyl donors, RXFP1 agonists, APJ agonists and cardiogenic agents. These agents include, but are not limited to furosemide, bumetanide, torsemide, sacubitril-valsartan, thiazide diuretics, captopril, enalapril, lisinopril, carvedilol, metoprolol, bisoprolol, serelaxin, spironolactone,

eplerenone, ivabradine, candesartan, eprosartan, irbestarin, losartan, olmesartan, telmisartan, and valsartan.

The compounds of the present invention may be employed in combination with at least one of the following therapeutic agents in treating atherosclerosis:

5 anti-hyperlipidemic agents, plasma HDL-raising agents, anti-hypercholesterolemic agents, cholesterol biosynthesis inhibitors (such as HMG CoA reductase inhibitors), LXR agonist, probucol, raloxifene, nicotinic acid, niacinamide, cholesterol absorption inhibitors, bile acid sequestrants (such as anion exchange resins, or quaternary amines (*e.g.*, cholestyramine or colestipol)), low density lipoprotein receptor inducers, clofibrate,
10 fenofibrate, benzofibrate, cipofibrate, gemfibrizol, vitamin B₆, vitamin B₁₂, anti-oxidant vitamins, β -blockers, anti-diabetes agents, angiotensin II antagonists, angiotensin converting enzyme inhibitors, platelet aggregation inhibitors, fibrinogen receptor antagonists, aspirin and fibric acid derivatives.

The compounds of the present invention may be employed in combination at least
15 one of the following therapeutic agents in treating cholesterol biosynthesis inhibitor, particularly an HMG-CoA reductase inhibitor. Examples of suitable HMG-CoA reductase inhibitors include, but are not limited to, lovastatin, simvastatin, pravastatin, fluvastatin, atorvastatin, and rosuvastatin.

The compounds of the invention may be used in combination with at least one of
20 the following anti-diabetic agents depending on the desired target therapy. Studies indicate that diabetes and hyperlipidemia modulation can be further improved by the addition of a second agent to the therapeutic regimen. Examples of anti-diabetic agents include, but are not limited to, sulfonylureas (such as chlorpropamide, tolbutamide, acetohexamide, tolazamide, glyburide, gliclazide, glynase, glimepiride, and glipizide),
25 biguanides (such as metformin), thiazolidinediones (such as ciglitazone, pioglitazone, troglitazone, and rosiglitazone), and related insulin sensitizers, such as selective and non-selective activators of PPAR α , PPAR β and PPAR γ ; dehydroepiandrosterone (also referred to as DHEA or its conjugated sulphate ester, DHEA-SO₄); anti-glucocorticoids; TNF α inhibitors; dipeptidyl peptidase IV (DPP4) inhibitor (such as sitagliptin,
30 saxagliptin), GLP-1 agonists or analogs (such as exenatide), α -glucosidase inhibitors (such as acarbose, miglitol, and voglibose), pramlintide (a synthetic analog of the human hormone amylin), other insulin secretagogues (such as repaglinide, gliquidone, and

nateglinide), insulin, as well as the therapeutic agents discussed above for treating atherosclerosis.

The compounds of the invention may be used in combination with at least one of the following anti-obesity agents selected from phenylpropanolamine, phentermine, 5 diethylpropion, mazindol, fenfluramine, dexfenfluramine, phentiramine, β_3 -adrenoreceptor agonist agents; sibutramine, gastrointestinal lipase inhibitors (such as orlistat), and leptins. Other agents used in treating obesity or obesity-related disorders include neuropeptide Y, enterostatin, cholecystokinin, bombesin, amylin, histamine H_3 receptors, dopamine D_2 receptor modulators, melanocyte stimulating hormone, 10 corticotrophin releasing factor, galanin and gamma amino butyric acid (GABA).

The compounds of the present invention are also useful as standard or reference compounds, for example as a quality standard or control, in tests or assays involving the FPR2. Such compounds may be provided in a commercial kit, for example, for use in pharmaceutical research involving FPR2 activity. For example, a compound of the 15 present invention could be used as a reference in an assay to compare its known activity to a compound with an unknown activity. This would ensure the experimenter that the assay was being performed properly and provide a basis for comparison, especially if the test compound was a derivative of the reference compound. When developing new assays or protocols, compounds according to the present invention could be used to test 20 their effectiveness. The compounds of the present invention may also be used in diagnostic assays involving FPR2.

The present invention also encompasses an article of manufacture. As used herein, article of manufacture is intended to include, but not be limited to, kits and packages. The article of manufacture of the present invention, comprises: (a) a first container; (b) a 25 pharmaceutical composition located within the first container, wherein the composition, comprises a first therapeutic agent, comprising a compound of the present invention or a pharmaceutically acceptable salt form thereof; and, (c) a package insert stating that the pharmaceutical composition can be used for the treatment of dyslipidemias and the sequelae thereof. In another embodiment, the package insert states that the 30 pharmaceutical composition can be used in combination (as defined previously) with a second therapeutic agent for the treatment of dyslipidemias and the sequelae thereof. The article of manufacture can further comprise: (d) a second container, wherein components

(a) and (b) are located within the second container and component (c) is located within or outside of the second container. Located within the first and second containers means that the respective container holds the item within its boundaries. The first container is a receptacle used to hold a pharmaceutical composition. This container can be for manufacturing, storing, shipping, and/or individual/bulk selling. First container is intended to cover a bottle, jar, vial, flask, syringe, tube (*e.g.*, for a cream preparation), or any other container used to manufacture, hold, store, or distribute a pharmaceutical product. The second container is one used to hold the first container and, optionally, the package insert. Examples of the second container include, but are not limited to, boxes (*e.g.*, cardboard or plastic), crates, cartons, bags (*e.g.*, paper or plastic bags), pouches, and sacks. The package insert can be physically attached to the outside of the first container via tape, glue, staple, or another method of attachment, or it can rest inside the second container without any physical means of attachment to the first container. Alternatively, the package insert is located on the outside of the second container. When located on the outside of the second container, it is preferable that the package insert is physically attached via tape, glue, staple, or another method of attachment. Alternatively, it can be adjacent to or touching the outside of the second container without being physically attached. The package insert is a label, tag, marker, etc. that recites information relating to the pharmaceutical composition located within the first container. The information recited will usually be determined by the regulatory agency governing the area in which the article of manufacture is to be sold (*e.g.*, the United States Food and Drug Administration). Preferably, the package insert specifically recites the indications for which the pharmaceutical composition has been approved. The package insert may be made of any material on which a person can read information contained therein or thereon. Preferably, the package insert is a printable material (*e.g.*, paper, plastic, cardboard, foil, adhesive-backed paper or plastic, etc.) on which the desired information has been formed (*e.g.*, printed or applied).

CHEMISTRY METHODS

Abbreviations as used herein, are defined as follows: "1x" for once, "2x" for twice, "3x" for thrice, "°C" for degrees Celsius, "aq" for aqueous, "Col" for column, "eq" for equivalent or equivalents, "g" for gram or grams, "mg" for milligram or milligrams,

"L" for liter or liters, "mL" for milliliter or milliliters, "μL" for microliter or microliters, "N" for normal, "M" for molar, "nM" for nanomolar, "mol" for mole or moles, "mmol" for millimole or millimoles, "min" for minute or minutes, "h" for hour or hours, "rt" for room temperature, "RT" for retention time, "ON" for overnight, "atm" for atmosphere, "psi" for pounds per square inch, "conc." for concentrate, "aq" for "aqueous", "sat" or "sat'd" for saturated, "MW" for molecular weight, "mw" or "μwave" for microwave, "mp" for melting point, "Wt" for weight, "MS" or "Mass Spec" for mass spectrometry, "ESI" for electrospray ionization mass spectroscopy, "HR" for high resolution, "HRMS" for high resolution mass spectrometry, "LCMS" for liquid chromatography mass spectrometry, "HPLC" for high pressure liquid chromatography, "RP HPLC" for reverse phase HPLC, "TLC" or "tlc" for thin layer chromatography, "NMR" for nuclear magnetic resonance spectroscopy, "nOe" for nuclear Overhauser effect spectroscopy, "¹H" for proton, "δ" for delta, "s" for singlet, "d" for doublet, "t" for triplet, "q" for quartet, "m" for multiplet, "br" for broad, "Hz" for hertz, and "α", "β", "R", "S", "E", and "Z" are stereochemical designations familiar to one skilled in the art.

Ac	acetic
AcOH	acetic acid
Acn (or MeCN)	acetonitrile
BINAP	2,2'-bis(diphenylphosphino)-1,1'-binaphthyl
BISPIN	ais(pinacolato)diboron
Bn	benzyl
Boc	<i>tert</i> -butyl carbonyl
Boc ₂ O	di- <i>tert</i> -butyl dicarbonate
Bu	butyl
dba as in (Pd ₂ (dba) ₃)	dibenzylideneacetone
DCM	dichloromethane
DEAD	diethyl azodicarboxylate
DIAD	diisopropyl azodicarboxylate
DIEA	diisopropylethylamine
DMAP	4-dimethylaminopyridine
DME	dimethoxyethane

DMF	dimethylformamide
DMSO	dimethyl sulfoxide
dppf	1,1'-bis(diphenylphosphino)ferrocene
Et	ethyl
EtOH	ethanol
EtOAc	ethyl acetate
HATU	2-(7-Aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
HBTU	2-(1H-Benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate
<i>i</i> -Bu	isobutyl
<i>i</i> -Pr	isopropyl
LAH	lithium aluminum hydride
Me	methyl
MeOH	methanol
NBS	N-bromosuccinimide
NMM	<i>N</i> -methylmorpholine
NMP	N-Methylpyrrolidone
Pet	petroleum
Ph	phenyl
Pr	propyl
rt	room temperature
<i>t</i> -Bu	<i>tert</i> -butyl
TBDMS-Cl	<i>t</i> -butyldimethylchlorosilane
TEA	triethylamine
TFA	trifluoroacetic acid
THF	tetrahydrofuran
Ts	tosyl
Xantphos	4,5-Bis(diphenylphosphino)-9,9-dimethylxanthene


The disclosed compounds can be made by various methods known in the art including those of the following schemes and in the specific embodiments section. The structure numbering and variable numbering shown in the synthetic schemes are distinct

from and should not be confused with the structure or variable numbering in the claims or the rest of the specification. The variables in the schemes are meant only to illustrate how to make some of the compounds of this invention.

The disclosure is not limited to the foregoing illustrative examples and the
5 examples should be considered in all respects as illustrative and not restrictive, and all changes which come within the meaning and range of equivalency of the claims are therefore intended to be embraced.

A consideration in the planning of any synthetic route in this field is the choice of the protecting group used for protection of the reactive functional groups present in the
10 compounds described in this invention. An authoritative account describing the many alternatives to the trained practitioner is Greene, T.W. et al., *Protecting Groups in Organic Synthesis*, 4th Edition, Wiley (2007)).


Compounds having the general Formula (I): wherein A, B, C, R^x, R^y, and R^z are defined above as Ar¹, Ar² and Ar³, (R^{2a})₁₋₄, (R^{3a})₁₋₄, and (R^{1a})₁₋₂, (R^{1b})₁₋₂, respectively, can
15 be prepared by the following one or more of the synthetic Schemes.



1-Arylpyrrolidinone compounds of this invention wherein rings A, B and C are substituted phenyl rings can be prepared by the general route shown in Scheme 1, starting from a suitably protected 3-aminopyrrolidin-2-one **1a**, where PG is a protecting group
20 such as Boc or Cbz. **1a** can be prepared with methods known to one skilled in the art. Copper or Pd-catalyzed coupling of **1a** to a substituted iodobenzene or bromobenzene **1b** or other suitable halo aryl or heteroaryl compound in a suitable solvent such as butanol or dioxane or toluene, in the presence of a base such as potassium carbonate or cesium carbonate and a suitable ligand such as N,N'-dimethylethylenediamine, or xanthphos can
25 afford 1-phenylpyrrolidinones **1c**. Additional methods for this transformation include other variations of Ullmann, Goldberg, and Buchwald copper-catalyzed amidation or


Buchwald Pd-catalyzed amidation depending on the nature of ring B, using methods known to one skilled in the art for these types of couplings (see for example Yin & Buchwald *Organic Lett.* **2000**, 2, 1101; Klapers *et al. JACS*, **2001**, 123, 7727; Klapars *et al. JACS*, **2002**, 124, 7421; Yin & Buchwald *JACS*, **2002**, 124, 6043; Kiyomori, Madoux & Buchwald, *Tet. Lett.*, **1999**, 40, 2657). Subsequent palladium-catalyzed coupling of **1c** to a suitably substituted phenyl boronic acid **1d**, or analogous boronate or trifluoroborate reagent, can provide the biaryl compound **1e**. Removal of the Boc or Cbz protecting group from **1e**, followed by condensation of the resulting free amine with a suitably substituted phenyl isocyanate, **1g** or phenylcarbamate **1h** can provide ureas **1f**. Suitable isocyanates or 4-nitrophenylcarbamates are either commercially available or can be readily obtained from the corresponding aniline by methods known to one skilled in the art. Alternately, the ureas **1f** can be obtained by treatment of the deprotected 3-aminopyrrolidinone intermediate with 4-nitrophenylchloroformate to form the carbamate, followed by condensation with an appropriately substituted aniline **1j**. It will also be recognized by one skilled in the art that additional compounds of this invention wherein rings A, B or C are heteroaryl rings, such as pyridine, pyrimidine, thiazole, etc., can also be prepared using the methods outlined in Scheme 1 by substituting the appropriate heteroaryl iodide or bromide for **1b**, heteroarylboronic acid or boronate for **1d** and heteroaryl amine, isocyanate or p-nitrophenylcarbamate for **1e**. Racemic compounds were separated using either chiral HPLC or SFC to provide single enantiomers.

Scheme 1




Alternatively as described in Scheme 2, compounds of this invention can be prepared from intermediate **1c** by first deprotecting the amine and forming the urea linkage to ring A using the conditions described above for the conversion of **1e** to **1f** to provide compounds **2a**. Compound **2a** can then be coupled with an appropriate boronic acid or boronate under Pd-catalysis conditions as shown in Scheme 1 for the transformation of **1c** to **1e**. Racemic compounds can be separated using either chiral HPLC or SFC to provide single enantiomers.

Scheme 2




Additionally as shown in Scheme 3, compounds of this invention can be prepared from intermediate **2a** by conversion to boronate **3a** using iridium/Pd-catalyzed C-H borylation according to the methods of Suzuki and Miyaura followed by coupling of the resulting pinacolatoboron species with aryl or heteroaryl halides using palladium or copper catalyzed processes to provide compounds **1f**. Racemic compounds can be separated using either chiral HPLC or SFC to provide single enantiomers.

Scheme 3




Alternatively as described in Scheme 4, compounds of this invention can be prepared by brominating amine **4a** to give intermediate **4b**. Subsequent amide coupling with 2,4-dibromo-butyl chloride under basic conditions such as potassium phosphate in acetonitrile, followed by aqueous ammonia mediated ring closure can give lactam **1c**. Intermediate **1c** can be converted into final products using the reactions shown in Schemes 1-3. Racemic compounds can be separated using either chiral HPLC or SFC to provide single enantiomers.

Scheme 4



As described in Scheme 5, compounds of this invention with substituted lactams can be prepared from intermediate **5a**. Copper or Pd-catalyzed coupling of **5a** to a substituted iodobenzene or bromobenzene **1b** or other suitable halo aryl or heteroaryl compound in a suitable solvent such as butanol or dioxane or toluene, in the presence of a base such as potassium carbonate or cesium carbonate and a suitable ligand such as N,N'-dimethylethylenediamine, or Xanthphos can afford 1-phenylpyrrolidinones **5b**. Additional methods for this transformation include other variations of Ullmann, and Buchwald copper-catalyzed amidation or Buchwald Pd-catalyzed amidation depending on the nature of ring B, using methods known to one skilled in the art for these types of couplings. Subsequent preparation of a lithium lactam enolate and treatment with an azide such as trisyl azide can yield intermediate **5c** (see for example *J. Org. Chem.* **2003**, 68, 7219-7233). Reduction of azide to amine followed by protection of amine can give intermediate **5d**. Chemistry described in previous schemes can be used to convert this intermediate into compounds claimed in this patent. Racemic compounds can be separated using either chiral HPLC or SFC to provide single enantiomers.


Scheme 5



Compounds of the invention can also be synthesized using the route shown in Scheme 6. Palladium catalyzed borylation of intermediate **1c** gives intermediate **6a**.

Palladium mediated coupling of a suitably substituted aryl halides such as **6b** can give biaryl lactam **1e**. Chemistry described in previous schemes can be used to convert this intermediate into compounds claimed in this patent. Racemic compounds can be separated using either chiral HPLC or SFC to provide single enantiomers.

5



Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments that are given for illustration of the invention and are not intended to be limiting thereof.

The following methods were used in the exemplified Examples, except where noted otherwise. Purification of intermediates and final products was carried out via either normal or reverse phase chromatography. Normal phase chromatography was carried out using prepacked SiO₂ cartridges eluting with either gradients of hexanes and ethyl acetate or DCM and MeOH unless otherwise indicated. Reverse phase preparative HPLC was carried out using C18 columns with UV 220 nm or prep LCMS detection eluting with gradients of Solvent A (90% water, 10% MeOH, 0.1% TFA) and Solvent B (10% water, 90% MeOH, 0.1% TFA) or with gradients of Solvent A (95% water, 5% Acn, 0.1% TFA) and Solvent B (5% water, 95% ACN, 0.1% TFA) or with gradients of Solvent A (95% water, 2% Acn, 0.1% HCOOH) and Solvent B (98% Acn, 2% water, 0.1% HCOOH) or with gradients of Solvent A (95% water, 5% Acn, 10 mM NH₄OAc) and Solvent B (98% ACN, 2% water, 10 mM NH₄OAc) or with gradients of Solvent A (98% water, 2% Acn, 0.1% NH₄OH) and Solvent B (98% Acn, 2% water, 0.1% NH₄OH).

25

LC/MS Methods Employed in Characterization of Examples. Reverse phase analytical HPLC/MS was performed on a Waters Acquity system coupled with a Waters MICROMASS® ZQ Mass Spectrometer.

- 5 Method A: Linear gradient of 0 to 100% B over 3 min, with 0.75 min hold time at 100% B;

UV visualization at 220 nm

Column: Waters BEH C18 2.1 x 50 mm

Flow rate: 1.0 mL/min

- 10 Solvent A: 0.1% TFA, 95% water, 5% Acn

Solvent B: 0.1% TFA, 5% water, 95% Acn

- Method B: Linear gradient of 0 to 100% B over 3 min, with 0.75 min hold time at 100% B;

- 15 UV visualization at 220 nm

Column: Waters BEH C18 2.1 x 50 mm

Flow rate: 1.0 mL/min

Solvent A: 10 mM ammonium acetate, 95% water, 5% Acn

Solvent B: 10 mM ammonium acetate, 5% water, 95% Acn

20

Analytical HPLC: Methods Employed in Characterization of Examples

- Method C: Ascentis Express C18, 2.1 x 50 mm, 2.7- μ m particles; Solvent A: 95% water, 5% Acn, 0.05% TFA; Solvent B: 95% Acn, 5% water, 0.1% TFA; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 1-minute hold at 100% B; Flow: 1.1 mL/min.

- Method D: Ascentis Express C18, 2.1 x 50 mm, 2.7- μ m particles; Solvent A: 95% water, 5% Acn with 10 mM ammonium acetate; Solvent B: 95% Acn, 5% water with 10 mM ammonium acetate; Temperature: 50 °C; Gradient: 0-100% B over 3 minutes, then a 1-minute hold at 100% B; Flow: 1.1 mL/min.

- 30 Method E: Column-Kinetex C18, 75 x 3 mm, 2.6- μ m particles; Solvent A: 98% water, 2% Acn with 10 mM ammonium acetate; Solvent B: 98% Acn, 2% water with 10 mM ammonium acetate; Temperature: 25 °C; Gradient: 20-100% B over 4 minutes, then

a 0.6-minute hold at 100% B; Flow: 1.0 mL/min; then, Gradient: 100-20% B over 0.4 minutes; Flow: 1.5 mL/min, UV 220 nm.

SFC and chiral purity methods

5 Method I: Chiralpak AD-H, 250 x 4.6 mm, 5.0-µm particles; % CO₂: 60%, % Co solvent: 40% {0.2%DEA IN IPA:Acn (1:1)}, Total Flow: 4.0g/min, Back Pressure: 100bars, Temperature : 25°C, UV: 218 nm.

 Method II: Chiralpak OD-H, 250 x 4.6 mm, 5.0-µm particles; % CO₂: 60%, % Co solvent: 40% {0.2% DEA IN IPA:Acn (1:1)}, Total Flow: 4.0g/min, Back Pressure: 104
10 bars, Temperature : 24.9°C, UV: 287 nm.

 Method III: Chiralpak OJ-H, 250 x 4.6 mm, 5.0-µm particles; % CO₂: 60%, % Co solvent: 30%(0.3% DEA in Methanol), Total Flow: 4.0g/min, Back Pressure: 101 bars, Temperature : 23.6°C, UV: 272 nm.

 Method IV: Chiralpak AS-H, 250 x 4.6 mm, 5.0-µm particles; % CO₂: 60%, %
15 Co solvent: 40%(0.3% DEA in Methanol), Total Flow: 4.0g/min, Back Pressure: 102 bars, Temperature : 25.4°C, UV: 272 nm.

 Method V: Chiralcel OJ-H, 250 x 4.6 mm, 5.0-µm particles; % CO₂: 60%, % Co solvent: 40%(0.2% DEA in Methanol), Total Flow: 4.0g/min, Back Pressure: 102 bars, Temperature : 24.6°C, UV: 272 nm.

20 Method VI: Luxcellulose-2, 250 x 4.6 mm, 5.0-µm particles; % CO₂: 60%, % Co solvent: 35%(0.2% DEA in Methanol), Total Flow: 3.0 g/min, Back Pressure: 101 bars, Temperature : 23.6°C, UV: 260 nm.

 Method VII: Chiralcel AS-H, 250 x 4.6 mm, 5.0-µm particles; % CO₂: 60%, % Co solvent: 40%(0.2% DEA in Methanol), Total Flow: 4.0 g/min, Back Pressure: 101
25 bars, Temperature : 24.4°C, UV: 270 nm.

 Method VIII: Chiralpak IC, 250 x 4.6 mm, 5.0-µm particles; % CO₂: 60%, % Co solvent: 40%(0.2% DEA in Methanol), Total Flow: 4.0 g/min, Back Pressure: 101 bars, Temperature : 24.4°C, UV: 270 nm.

 Method IX: Column: chiralpak IF (250 X 4.6mm), 5 micron, mobile phase: 0.2%
30 DEA in ethanol, FLOW:1.0ml\min.

 Method X: Column: LUX AMYLOSE 2 (250 X 4.6mm), 5 micron, mobile phase: 0.2% DEA in n-hexane:ethanol:5:95, FLOW:1.0ml\min.

Method XI: Column: CHIRALCEL OD-H (250 X 4.6mm), 5 micron, mobile phase: 0.2% DEA in n-hexane:ethanol:70:30, FLOW:1.0ml\min.

Method XII: Column: CHIRAL PAK ID 250 X 4.6mm), 5 micron, mobile phase: 0.1% DEA in Methanol, FLOW: 1.0ml/min.


5

NMR Employed in Characterization of Examples. ¹H NMR spectra were obtained with Bruker or JEOL[®] Fourier transform spectrometers operating at frequencies as follows: ¹H NMR: 400 MHz (Bruker or JEOL[®]) or 500 MHz (Bruker or JEOL[®]). ¹³C NMR: 100 MHz (Bruker or JEOL[®]). Spectra data are reported in the format: chemical shift (multiplicity, coupling constants, and number of hydrogens). Chemical shifts are specified in ppm downfield of a tetramethylsilane internal standard (δ units, tetramethylsilane = 0 ppm) and/or referenced to solvent peaks, which in ¹H NMR spectra appear at 2.49 ppm for CD₂HSOCD₃, 3.30 ppm for CD₂HOD, 1.94 for CD₃CN, and 7.24 ppm for CHCl₃, and which in ¹³C NMR spectra appear at 39.7 ppm for CD₃SOCD₃, 49.0 ppm for CD₃OD, and 77.0 ppm for CDCl₃. All ¹³C NMR spectra were proton decoupled.

10

15

Intermediate 1: *3-Aminopyrrolidin-2-one*




To a stirred solution of hexamethyldisilazane (11 mL, 523 mmol) in CH₃CN (100 mL) at rt was added a solution of DL-2,4-diaminobutyric acid dihydrochloride (10 g, 52 mmol) in Acn (100 mL). The resulting reaction mixture was heated to reflux for 40 h. Then, the crude reaction mixture was poured into ice cold MeOH (400 mL), stirred at rt for 30 min and evaporated under reduced pressure. The resulting solid was dissolved in CH₂Cl₂ (700 mL), and the insoluble residue was removed by filtration under vacuum.

20

25


Then, the filtrate was concentrated under reduced pressure to give 3-aminopyrrolidin-2-one (4.1 g, 41 mmol, 78% yield) as a yellow solid. ¹H NMR (400MHz, DMSO-d₆): δ 7.59 (br. s., 2H), 3.23-3.10 (m, 1H), 3.09 - 3.03 (m, 2H), 2.23-2.22 (m, 1H), 1.70 - 1.57 (m, 1H).

30 Intermediate 2: *tert-Butyl (2-oxopyrrolidin-3-yl)carbamate*




To a stirred solution of 3-aminopyrrolidin-2-one (4.0 g, 40 mmol) in methanol-triethylamine (130 mL, 9:1) under argon atmosphere at rt, was added Boc-anhydride (9.6 mL, 41 mmol). The reaction mixture was stirred at rt overnight followed by heating
 5 to reflux for two hours. Then, the reaction mixture was cooled to rt and concentrated under reduced pressure. Ether (50 mL) was added to the crude residue and the solid was filtered through Buchner funnel to yield Intermediate 2 (4.0 g, 20 mmol, 50% yield) as brown solid. ¹H NMR (400MHz, DMSO-d₆): δ 7.69 (br. s., 1H), 6.99 (d, J=8.0 Hz, 1H), 4.06 - 3.96 (m, 1H), 3.19 - 3.10 (m, 2H), 2.29 - 2.19 (m, 1H), 1.89 - 1.76 (m, 1H), 1.35 (s, 9H).

Intermediate 3: *tert*-Butyl (1-(4-bromo-2,3-difluorophenyl)-2-oxopyrrolidin-3-yl)carbamate




To a stirred solution of Intermediate 2 (1.0 g, 5.0 mmol) in 1,4-dioxane (10 mL),
 15 were added 1,4-dibromo-2,3-difluorobenzene (1.6 g, 6.0 mmol), and Cs₂CO₃ (3.3 g, 10 mmol). The reaction mixture was purged with nitrogen for 5 min and charged with Xantphos (0.29 g, 0.50 mmol) and Pd₂(dba)₃ (0.23 g, 0.25 mmol). The reaction mixture was again purged with nitrogen for 3 min and heated at 120°C for 16 h. The reaction mixture was cooled, filtered through a Celite pad, and the filtrate was concentrated under
 20 reduced pressure. The crude product was purified using column chromatography (pet. ether-EtOAc) to yield Intermediate 3 (1.1 g, 2.8 mmol, 47% yield) as brown solid. MS(ESI) m/z: 391.2 (M+H)⁺. ¹H NMR (400MHz, DMSO-d₆) δ 7.61-7.59 (m, 1H), 7.35 - 7.28 (m, 2H), 4.39 - 4.28 (m, 1H), 3.82 - 3.64 (m, 2H), 2.42 - 2.31 (m, 1H), 2.10 - 1.97 (m, 1H), 1.40 (s, 9H).

Intermediate 4: *tert*-Butyl (1-(2,3-difluoro-2'-(methylsulfonamido)-[1,1'-biphenyl]-4-yl)-2-oxopyrrolidin-3-yl)carbamate



To a solution of Intermediate 3 (1.1 g, 2.1 mmol) in 1,4-dioxane-water (11 mL, 10:1) at rt, were added (2-(methanesulfonamido)phenyl)boronic acid (0.45 g, 2.1 mmol), and potassium phosphate, tribasic (0.74 g, 4.2 mmol). The reaction mixture was purged with nitrogen for 5 min and charged with $\text{PdCl}_2(\text{dppf})\cdot\text{CH}_2\text{Cl}_2$ (0.17 g, 0.21 mmol). The reaction mixture was again purged with nitrogen for 3 min and heated at 80 °C for 16 h. The reaction mixture was cooled and filtered through a Celite pad, and the filtrate was concentrated under reduced pressure. The crude product was purified via column chromatography (pet. ether-EtOAc) to give Intermediate 4 (0.44 g, 0.91 mmol, 43% yield) as brown solid. MS(ESI) m/z : 482.2 $[\text{M}+\text{H}]^+$; ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 9.13 (s, 1H), 7.53 - 7.44 (m, 2H), 7.38 - 7.29 (m, 4H), 7.24 - 7.17 (m, 1H), 4.36-4.31 (m, 1H), 3.86 - 3.69 (m, 2H), 2.90 (s, 3H), 2.41-2.37 (m, 1H), 2.10 - 2.02 (m, 1H), 1.41 (s, 9H).


Intermediate 5: *N*-(4'-(3-Amino-2-oxopyrrolidin-1-yl)-2',3'-difluoro-[1,1'-biphenyl]-2-yl)methanesulfonamide hydrochloride



To an ice cooled solution of Intermediate 4 (440 mg, 0.91 mmol) in 1,4-dioxane (1 mL), was added 4 N HCl in 1,4-dioxane (4.6 mL, 1 mmol), and the reaction mixture was stirred at rt for 2 h. The solvent was evaporated under reduced pressure. The gummy solid was triturated with diethyl ether (20 ml x 2) and dried to yield Intermediate 5 (350 mg, 0.84 mmol, 92% yield) as a brown solid.


MS(ESI) m/z: 418.2 [M+H]⁺; ¹H NMR (400MHz, DMSO-d₆) δ 9.18 (s, 1H), 8.71 (br. s., 3H), 7.55 - 7.47 (m, 2H), 7.45 - 7.24 (m, 4H), 4.33 - 4.25 (m, 1H), 3.97 - 3.85 (m, 2H), 2.89 (s, 3H), 2.64 - 2.54 (m, 1H), 2.30 - 2.17 (m, 1H).

5 Intermediate 6: *tert*-butyl (*R*)-(2-oxopyrrolidin-3-yl)carbamate



1-Propanephosphonic anhydride (50% in EtOAc, 210 mL, 340 mmol) was added to a solution of Boc-D-2,4-diaminobutyric acid (50 g, 230 mmol) and TEA (96 mL, 690 mmol) in DCM (1500 mL) at 0 °C. The reaction mixture was stirred at rt overnight under nitrogen. The reaction mixture was concentrated in vacuo, and the crude product was purified by column chromatography (MeOH/DCM). The product was recrystallized with EtOAc/pet. ether to obtain Intermediate 6 (32 g, 70 mmol, 70 % yield) as a white solid. MS(ESI) m/z: 201.2 (M+H)⁺. ¹H NMR (400 MHz, DMSO-d₆) δ = 7.69 (s, 1H), 6.99 (br. d., *J* = 9.0 Hz, 1H), 4.01 (q, *J* = 9.0 Hz, 1H), 3.17 - 3.09 (m, 2H), 2.28 - 2.19 (m, 1H), 1.90 - 1.75 (m, 1H), 1.39 (s, 9H).


Intermediate 7: *tert*-butyl (*R*)-(1-(2,3-difluoro-4-bromophenyl)-2-oxopyrrolidin-3-yl)carbamate



20 A reaction mixture of Intermediate 6 (20 g, 100 mmol), 2,3-difluoro-1,4-dibromobenzene (14 g, 110 mmol), potassium phosphate tribasic (32 g, 150 mmol), and cuprous iodide (7.6 g, 40 mmol) in 1,4-dioxane (250 mL) was purged with nitrogen for 5 min. N,N'-Dimethylethylenediamine (5.5 mL, 50 mmol) was added, and the reaction mixture was heated in a pressure tube at 65 °C for 12 h. The reaction mixture was diluted with EtOAc, filtered through Celite, and concentrated under reduced pressure. The crude product was purified by column chromatography (35% EtOAc in pet ether) and


recrystallized (EtOAc/pet ether) to obtain Intermediate 7 (18 g, 39 mmol, 39% yield) as a white solid. MS(ESI) m/z : 439.0 ($M+H$)⁺. ¹H NMR (400 MHz, CDCl₃) δ = 7.54 (ddd, J = 8.5, 6.3, 2.3 Hz, 1H), 7.09 (ddd, J = 8.5, 6.3, 2.3 Hz, 1H), 5.16 (br. s., 1H), 4.41 - 4.30 (m, 1H), 3.87 (m, 1H), 3.79 - 3.69 (m, 1H), 2.85 - 2.73 (m, 1H), 2.17 - 2.05 (m, 1H), 1.48 (m, 9H).

Intermediate 8: (2-bromophenyl)dimethylphosphine oxide



In a pressure tube, a reaction mixture of 1-bromo-2-iodobenzene (2.3 mL, 18 mmol), potassium phosphate tribasic (5.6 g, 27 mmol) and Xantphos (0.61 g, 1.1 mmol) in DMF (40 mL) was degassed with argon for 3 min. Dimethylphosphine oxide (1.7 g, 21 mmol), and PdOAc₂ (0.20 g, 0.88 mmol) were added. The mixture was degassed with argon, sealed and heated for 16 h at 110 °C. The reaction mixture was cooled to rt, filtered through Celite and concentrated under reduced pressure. The crude product was purified by column chromatography (CHCl₃/MeOH) to give Intermediate 8 (2.1 g, 9.0 mmol, 51% yield). MS(ESI) m/z : 233.0 ($M+H$)⁺. ¹H NMR (300 MHz, DMSO-d₆) δ = 7.97 (ddd, J = 11.9, 7.8, 2.0 Hz, 1H), 7.76 (ddd, J = 7.8, 3.9, 1.2 Hz, 1H), 7.65 - 7.44 (m, 2H), 1.83 (d, J = 13.5 Hz, 6H).


Intermediate 9: *tert*-butyl (*R*)-(1-(2'-(dimethylphosphoryl)-2,3-difluoro-[1,1'-biphenyl]-4-yl)-2-oxopyrrolidin-3-yl)carbamate



A reaction mixture of Intermediate 7 (15 g, 38 mmol), Intermediate 8 (11 g, 46 mmol), bispin (24 g, 96 mmol), and K₃PO₄ (24 g, 115 mmol) in 1,4-dioxane (250 mL) was purged with argon for 5 min. PdCl₂(dppf)-CH₂Cl₂ adduct (3.1 g, 3.8 mmol) was


added and the mixture was again purged with argon and then stirred in a pressure tube at 105 °C for 24 h. The reaction mixture was diluted with EtOAc, filtered through Celite, and concentrated under reduced pressure. The crude product was purified via column chromatography (5% MeOH in CHCl₃) to give Intermediate 9 (9.0 g, 19 mmol, 51%
 5 yield). MS(ESI) m/z: 465.4 (M+H)⁺. ¹H NMR (400 MHz, DMSO-d₆) δ = 7.92 (ddd, *J* = 12.8, 7.6, 1.3 Hz, 1H), 7.70 - 7.49 (m, 2H), 7.41 - 7.23 (m, 4H), 4.54 - 4.23 (m, 1H), 3.87 - 3.70 (m, 2H), 2.44 - 2.37 (m, 1H), 2.21 - 2.00 (m, 1H), 1.51 (br. d., *J* = 13.1 Hz, 6H), 1.42 (s, 9H).

10 Intermediate 10: (*R*)-3-amino-1-(2'-(dimethylphosphoryl)-2,3-difluoro-[1,1'-biphenyl]-4-yl)pyrrolidin-2-one hydrochloride




HCl (4 M in 1,4-dioxane, 200 mL, 800 mmol) was added to a stirred solution of Intermediate 9 (34 g, 73 mmol) in 1,4-dioxane (500 mL) at rt. The reaction mixture was
 15 stirred for 3 h and concentrated under reduced pressure to yield Intermediate 10 (28 g, 70 mmol, 95% yield), which was used in the next synthetic step without further purification. MS(ESI) m/z: 365.2 (M+H)⁺. ¹H NMR (400 MHz, DMSO-d₆) δ = 8.62 (br. s., 3H), 7.95 - 7.81 (m, 1H), 7.68 - 7.57 (m, 2H), 7.39 - 7.32 (m, 2H), 7.31 - 7.24 (m, 1H), 4.33 - 4.19 (m, 1H), 3.94 - 3.84 (m, 2H), 2.61 - 2.53 (m, 1H), 2.25 - 2.15 (m, 1H), 1.53 (d, *J* = 13.1
 20 Hz, 6H).

Intermediate 11: *phenyl (4-chloro-2-fluorophenyl)carbamate*




Phenyl chloroformate (20 mL, 160 mmol) was added slowly to a solution of 4-chloro-2-fluoroaniline (19 mL, 170 mmol) and pyridine (35 mL, 430 mmol) in DCM (250 mL) at 0 °C. The reaction mixture was gradually warmed to rt and stirred for 12 h. The reaction mixture was quenched with water. The mixture was extracted with DCM (3 x 200 mL). The combined organic layers were washed with 0.5 N HCl and 10% sodium bicarbonate, dried over sodium sulfate, and concentrated in vacuo. The solid was stirred in pet ether (100 mL) for 15 min, filtered and dried to get Intermediate 11 (36 g, 40 mmol, 78% yield). ¹H NMR (300MHz, CHLOROFORM-d) δ = 10.22 - 9.99 (m, 1H), 7.73 (br. t., *J* = 8.7 Hz, 1H), 7.52 (dd, *J* = 10.7, 2.1 Hz, 1H), 7.47 - 7.38 (m, 2H), 7.34 - 7.18 (m, 4H).

Intermediate 12: *phenyl (2-fluoro-4-(trifluoromethyl)phenyl)carbamate*




Phenyl chloroformate (4.5 mL, 36 mmol) was added slowly to a solution of 2-fluoro-4-(trifluoromethyl) aniline (8.0 g, 45 mmol) and pyridine (9.0 mL, 110 mmol) in DCM (80 mL) at 0 °C. The reaction mixture was gradually warmed to rt and stirred for 3 h. The reaction mixture was quenched with water and the mixture was extracted with DCM (3 x 200 mL). The combined organic layers were washed with 0.5 N HCl and 10% sodium bicarbonate, dried over sodium sulfate, and concentrated in vacuo. The solid was stirred in pet. ether (100 mL) for 15 min, filtered and dried to give Intermediate 11 (9.0 g, 30 mmol, 67% yield). MS(ESI) *m/z*: 317.2 (*M*+NH₄)⁺. ¹H NMR (300MHz, CHLOROFORM-d) δ = 10.43 (s, 1H), 8.04 (br. t., *J* = 8.1 Hz, 1H), 7.75 (br. d., *J* = 10.9 Hz, 1H), 7.60 (br. d., *J* = 8.6 Hz, 1H), 7.52 - 7.38 (m, 2H), 7.36 - 7.21 (m, 3H).

Intermediate 13: *(R)-1-(1-(4-bromo-2,3-difluorophenyl)-2-oxopyrrolidin-3-yl)-3-(2-fluoro-4-(trifluoromethyl)phenyl)urea*




HCl (4 M in dioxane, 32 mL, 130 mmol) was added dropwise to a solution of Intermediate 7 (10 g, 26 mmol) in DCE (300 mL) at 0 °C. The reaction mixture was stirred for 2 h at rt and concentrated under reduced pressure. The crude intermediate was dissolved in 1,4-dioxane (300 mL) and cooled to 0 °C. DIEA (22 mL, 130 mmol) was added dropwise over a period of 10 min, Intermediate 12 (6.9 g, 23 mmol) was added, and the mixture was stirred for 12 h at rt. The reaction mixture was filtered through Celite, and the filter plug was washed with EtOAc. The filtrate was dried over sodium sulfate and concentrated under reduced pressure. The crude product was purified by column chromatography (2% MeOH:DCM) to yield Intermediate 13 (12 g, 24 mmol, 95% yield). MS(ESI) m/z: 496.0 (M+H)⁺.

Intermediate 14: *(R)*-1-(1-(2,3-difluoro-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl)-2-oxopyrrolidin-3-yl)-3-(2-fluoro-4-(trifluoromethyl)phenyl)urea




A solution of Intermediate 13 (12 g, 24 mmol) and bispin (18 g, 73 mmol) in 1,4-dioxane (120 mL) was degassed with argon for 5 min. PdCl₂(dppf)-CH₂Cl₂ (4.0 g, 4.8 mmol) was added, and the reaction mixture was slowly heated to 110 °C and stirred for 16 h. The mixture was filtered through Celite, and the plug was washed with EtOAc. The filtrate was dried over sodium sulfate and concentrated under reduced pressure to give Intermediate 14 (13 g, 24 mmol, 99% yield). MS(ESI) m/z: 544.2 (M+H)⁺.

Intermediate 15: (3-bromopyridin-2-yl)dimethylphosphine oxide




To a stirred solution of 3-bromo-2-iodopyridine (5.0 g, 18 mmol) in DMF (5 mL),
5 were added dimethylphosphine oxide (1.7 g, 21 mmol), and K₃PO₄ (4.1 g, 19 mmol). The
reaction mixture was purged with nitrogen for 5 min and charged with Xantphos (1.0 g,
1.8 mmol) and Pd(OAc)₂ (0.20 g, 0.88 mmol). The reaction mixture was again purged
with nitrogen for 3 min and heated at 100 °C for 9 hours. The reaction mixture was
filtered through Celite and concentrated under reduced pressure. The crude material was
10 purified by column chromatography (5% MeOH-CHCl₃) to give Intermediate 15 (2.7 g,
12 mmol, 66 % yield) as a brown liquid. MS(ESI) m/z: 234.1 [M+H]⁺. ¹H NMR (400
MHz, DMSO-d₆) δ = 8.72 (d, J = 4.8 Hz, 1H), 8.23-8.20 (m, 1 H), 7.53 - 7.49 (m, 1 H),
1.66 (d, J = 13.6 Hz, 6H).

15 Intermediate 16: (2-Bromo-6-fluorophenyl)dimethylphosphine oxide




To a stirred solution of 1-bromo-3-fluoro-2-iodobenzene (2.0 g, 6.7 mmol) in 1,4-
dioxane (15 mL), were added dimethylphosphine oxide (0.62 g, 8.0 mmol), and K₃PO₄
(1.6 g, 7.3 mmol). The reaction mixture was purged with nitrogen for 5 min and charged
20 with Xantphos (0.39 g, 0.67 mmol) and Pd₂(dba)₃ (0.30 g, 0.33 mmol). The reaction
mixture was again purged with nitrogen for 3 min and heated at 100 °C for 9 h. The
reaction mixture was filtered through Celite and concentrated under reduced pressure.
The crude material was purified by column chromatography (0-100% EtOAc-Hexane,
followed by 3% MeOH-CHCl₃) to give Intermediate 16 (0.33 g, 1.3 mmol, 20% yield) as
25 a yellow solid. MS(ESI) m/z: 250.9 [M+H]⁺. ¹H NMR (400 MHz, DMSO-d₆) δ = 7.63 (d,
J = 8.0 Hz, 1H), 7.52 (td, J = 8.0, 6.0 Hz, 1H), 7.43 - 7.28 (m, 1H), 1.87 (d, J = 16.6 Hz,
3H), 1.86 (d, J = 16.6 Hz, 3H).

Intermediate 17: diethylphosphine oxide



To a stirred solution of 2M ethylmagnesium bromide (330 mL, 650 mmol) in THF at rt, was added diethyl phosphonate (30 g, 220 mmol) and the mixture was stirred for
 5 additional 2 hours. Then, a cold solution of K₂CO₃ (90 g, 650 mmol) in water (120 mL) was added and the reaction mixture was filtered through a Celite pad. The filtrate was concentrated under reduced pressure and azeotroped with toluene (20 mL x 3). The crude material was diluted with DCM (100 mL), filtered through a cotton bed and concentrated under reduced pressure to give Intermediate 17 (12 g, 110 mmol, 52% yield) as a
 10 yellowish liquid. MS(ESI) m/z: 107.2 [M+H]⁺.


Intermediate 18: (2-Bromophenyl)diethylphosphine oxide



To a stirred solution of 1-bromo-2-iodobenzene (10 g, 35.3 mmol) in DMF (50
 15 mL), were added diethylphosphine oxide (4.5 g, 42 mmol), and K₃PO₄ (15 g, 71 mmol). The reaction mixture was purged with nitrogen for 5 min and charged with Xantphos (1.0 g, 1.8 mmol) and palladium (II) acetate (0.40 g, 1.8 mmol). The reaction mixture was again purged with nitrogen for 3 min, heated to 100 °C and stirred for 16 h. The reaction mixture was cooled, filtered through a Celite pad, and the filtrate was concentrated under
 20 reduced pressure. The crude product was purified by column chromatography (MeOH-EtOAc) to yield the Intermediate 18 (3.2 g, 12 mmol, 35% yield) as a yellowish liquid. MS(ESI) m/z: 263.2 [M+H]⁺. ¹H NMR (400MHz, DMSO-d₆) δ = 8.05 - 7.94 (m, 1H), 7.79 - 7.68 (m, 2H), 7.64 - 7.48 (m, 1H), 2.26 - 2.03 (m, 4H), 0.93 (td, J = 17.4, 7.7 Hz, 6H).

25


Intermediate 19: Dimethyl (2-bromophenyl)phosphonate



To a stirred solution of 2-bromophenyl trifluoromethanesulfonate (2.0 g, 6.6 mmol) in toluene (100 mL), were added dimethyl phosphonate (1.1 g, 9.8 mmol), and

DIEA (1.7 mL, 9.8 mmol). The reaction mixture was purged with nitrogen for 5 min and charged with 1,3-bis(diphenylphosphino)propane (0.14 g, 0.33 mmol) and palladium (II) acetate (0.04 g, 0.17 mmol). The reaction mixture was again purged with nitrogen for 3 min and heated at 110 °C for 16 h. The reaction mixture was cooled and filtered through a Celite pad, and the filtrate was concentrated under reduced pressure. The product was purified by flash chromatography (5-10% MeOH in chloroform) to yield Intermediate 19 (1.2 g, 3.6 mmol, 55% yield) as a colorless liquid. MS(ESI) m/z: 267.1 [M+H]⁺. ¹H NMR (400MHz, CDCl₃) δ = 8.02 - 8.00 (m, 1 H) 7.70 - 7.69 (m, 1 H) 7.43 - 7.41 (m, 2 H) 3.83 - 3.73 (m, 6 H).


Intermediate 20: *Ethyl (2-bromophenyl)(methyl)phosphinate*



To a stirred solution of 1-bromo-2-iodobenzene (1.0 g, 3.5 mmol) in DMF (5 mL), were added diethyl methylphosphonite (0.58 g, 4.2 mmol), and K₃PO₄ (2.3 g, 11 mmol).

The reaction mixture was purged with nitrogen for 5 min and charged with Xantphos (0.21 g, 0.35 mmol) and palladium (II) acetate (0.040 g, 0.18 mmol). The reaction mixture was again purged with nitrogen for 3 min and heated at 110 °C for 16 h. The reaction mixture was cooled, filtered through Celite pad and the filtrate was concentrated under reduced pressure. The crude product was purified by flash chromatography (pet. ether-EtOAc) to yield the Intermediate 20 (0.35 g, 1.3 mmol, 38% yield) as a brown liquid. MS(ESI) m/z: 263.1 [M+H]⁺. ¹H NMR (400MHz, CDCl₃) δ = 8.18 - 8.13 (m, 1 H) 7.67 - 7.65 (m, 1 H) 7.64 - 7.38 (m, 2 H) 4.07- 4.03 (m, 1 H) 3.85 - 3.80 (m, 1 H) 1.86 (d, J = 15.0 Hz, 3H), 1.32 (t, J = 7.2 Hz, 3H).


Intermediate 21: *4-Bromo-3-cyclopropyl-2-fluoroaniline*



To an ice cooled solution of 3-cyclopropyl-2-fluoroaniline (0.85 g, 5.6 mmol) in DMF (11 mL) under argon atmosphere at rt, was added NBS (1.0 g, 5.6 mmol). The

reaction mixture was gradually warmed to rt over 3 h and quenched with aqueous $\text{Na}_2\text{S}_2\text{O}_3$ (10 mL). The biphasic mixture was extracted with EtOAc (30 mL x 2). The combined organic layer was washed with brine (15 mL x 2), dried over Na_2SO_4 , concentrated under reduced pressure and purified through flash column chromatography (25% EtOAc-Pet.ether) to yield Intermediate 21 (1.2 g, 5.0 mmol, 89% yield) as an orange liquid. ^1H NMR (400 MHz, CDCl_3) δ = 7.11 (d, J = 8.5 Hz, 1H), 6.51 (t, J = 8.8 Hz, 1H), 3.67 (br s, 2H), 1.80 (tt, J = 8.6, 5.7 Hz, 1H), 1.07 - 0.82 (m, 4H).

Intermediate 22: *tert*-Butyl (1-(4-bromo-3-cyclopropyl-2-fluorophenyl)-2-oxopyrrolidin-3-yl)carbamate




To a stirred solution of Intermediate 21 (1.1 g, 4.8 mmol) in Acn (10 mL) at 0 °C under nitrogen was added K_3PO_4 (1.0 g, 4.8 mmol). The resulting mixture was treated with 2,4-dibromobutanoyl chloride (0.60 mL, 4.5 mmol) and stirred for 2 h. Then, K_2CO_3 (2.0 g, 14 mmol) was added, and the mixture was stirred for 16 h. The solid was filtered through a Celite pad and the pad was washed with Acn (15 mL x 2). The filtrate was partially evaporated, ammonia (aq, 10 mL, 460 mmol) was added, and the mixture was stirred at 40 °C for 16 h. The mixture was extracted with EtOAc (20 mL x 3) and washed with brine (20 mL). The combined organics were dried over Na_2SO_4 and evaporated to give crude 3-amino-1-(4-bromo-3-cyclopropyl-2-fluorophenyl)pyrrolidin-2-one (1.0 g, 3.2 mmol, 67% yield), which was used in next step without further purification.

To a stirred solution of 3-amino-1-(4-bromo-3-cyclopropyl-2-fluorophenyl)pyrrolidin-2-one in DCM (15 mL) under argon atmosphere at rt, were added TEA (0.98 mL, 7.0 mmol) and $(\text{Boc})_2\text{O}$ (0.90 mL, 3.9 mmol). The reaction mixture was stirred at rt for 6 hours. Then, the reaction mixture was quenched with aqueous saturated NH_4Cl (20 mL) and extracted with EtOAc (20 mL x 3). The combined organic layers were washed with brine (20 mL), dried over Na_2SO_4 , concentrated under reduced pressure and purified via flash chromatography (20% EtOAc-pet. ether) to give


Intermediate 22 (1.0 g, 2.4 mmol, 69 % yield) as a yellowish gummy solid. MS(ESI) m/z: 413.1 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃) δ = 7.38 (dd, *J* = 8.5, 1.5 Hz, 1H), 7.14 - 7.09 (m, 1H), 5.18 (br. s., 1H), 4.37 - 4.33 (m, 1H), 3.86 - 3.74 (m, 1H), 3.69 - 3.62 (m, 1H), 2.79 - 2.75 (m, 1H), 2.15 - 2.01 (m, 1H), 1.80 (tt, *J* = 8.6, 5.7 Hz, 1H), 1.48 (s, 9H), 1.14 - 1.01 (m, 2H), 0.91 - 0.84 (m, 2H).

Intermediate 23: *tert*-Butyl(1-(2-cyclopropyl-2'-(dimethylphosphoryl)-3-fluoro-[1,1'-biphenyl]-4-yl)-2-oxopyrrolidin-3-yl)carbamate




To a solution of Intermediate 22 (0.50 g, 1.2 mmol) in 1,4-dioxane (10 mL) at rt, were added (2-bromophenyl)dimethylphosphine oxide (0.42 g, 1.8 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (0.92 g, 3.6 mmol), and K₂CO₃ (0.42 g, 3.0 mmol). The reaction mixture was purged with nitrogen for 5 min and then charged with PdCl₂(dppf).DCM (0.090 g, 0.12 mmol). The reaction mixture was again purged with nitrogen for 3 min and heated at 100 °C for 16 h. The reaction mixture was cooled and filtered through a Celite pad, and the filtrate was concentrated under reduced pressure. The crude intermediate was purified by flash chromatography (MeOH-CHCl₃) to yield Intermediate 23 (0.30 g, 0.62 mmol, 51% yield) as an orange solid. MS(ESI) m/z: 487.1 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃) δ = 8.25 - 8.10 (m, 1H), 7.62 - 7.48 (m, 2H), 7.33 - 7.21 (m, 2H), 7.07 - 6.95 (m, 1H), 5.26 - 5.10 (m, 1H), 4.48 - 4.27 (m, 1H), 3.93 - 3.79 (m, 1H), 3.79 - 3.67 (m, 1H), 2.88 - 2.74 (m, 1H), 2.20 - 2.01 (m, 1H), 1.54 - 1.37 (m, 16H), 1.04 - 0.85 (m, 2H), 0.73 - 0.68 (m, 2H).

Intermediate 24: 3-Amino-1-(2-cyclopropyl-2'-(dimethylphosphoryl)-3-fluoro-[1,1'-biphenyl]-4-yl)pyrrolidin-2-one hydrochloride



To an ice cooled solution of Intermediate 23 (0.33 g, 0.68 mmol) in 1,4-dioxane (1 mL) under argon atmosphere at rt, was added 4M HCl in 1,4-dioxane (3.0 mL, 12 mmol) and the mixture was stirred at rt for 2 h. The solvent was evaporated under reduced pressure to obtain a gummy solid. The crude product was triturated with pet. ether (10 mL x 2) and dried to give Intermediate 24 (0.26 mg, 0.67 mmol, 99% yield) as a brown gummy solid. MS(ESI) m/z: 387.1 [M+H]⁺.


Intermediate 25: *tert*-Butyl (1-(4-bromo-3-fluoro-2-(trifluoromethyl)phenyl)-2-oxopyrrolidin-3-yl)carbamate



To a stirred solution of 4-bromo-2-fluoro-3-(trifluoromethyl)aniline (0.70 g, 2.7 mmol) in Acn (10 mL) at 0 °C under nitrogen was added K₃PO₄ (0.58 g, 2.7 mmol). The resulting mixture was treated with 2,4-dibromobutanoyl chloride (0.29 mL, 2.2 mmol) and stirred for 2 h. Then, K₂CO₃ (1.1 g, 8.1 mmol) was added, and the reaction mixture was stirred for 16 h. The mixture was filtered through a Celite pad, and the pad was washed with Acn (15 mL x 2). The filtrate was partially evaporated and ammonia (aq) (10 mL, 462 mmol) was added. The mixture was stirred at 40 °C for 16 h and then extracted with EtOAc (20 mL x 3). The combined organics were washed with brine (20 mL), dried over Na₂SO₄ and evaporated to give crude 3-amino-1-(4-bromo-2-fluoro-3-(trifluoromethyl)phenyl)pyrrolidin-2-one (0.88 g, 2.6 mmol, 95 % yield), which was used in the next step without any further purification.


To a stirred solution of crude 3-amino-1-(4-bromo-3-fluoro-2-(trifluoromethyl)phenyl)pyrrolidin-2-one in DCM (15 mL) under argon atmosphere at rt, were added TEA (3.7 mL, 26 mmol) and (Boc)₂O (3.4 mL, 15 mmol). The reaction mixture was stirred at rt for 6 hours and then quenched with aqueous saturated NH₄Cl (20 mL) and extracted with EtOAc (20 mL x 3). The combined organic layers were washed with brine (20 mL), dried over Na₂SO₄, concentrated under reduced pressure and purified via column chromatography (30% EtOAc-pet. ether) to yield Intermediate 25 (4.8 g, 11 mmol, 82% yield) as a yellowish gummy solid. MS(ESI) m/z: 442.8 [M+H]⁺. ¹H NMR (300 MHz, CDCl₃) δ = 7.82 (t, *J* = 7.8 Hz, 1H), 7.02 (br. d., *J* = 8.7 Hz, 1H), 5.23 - 5.05 (m, 1H), 4.43 - 4.23 (m, 1H), 3.80 - 3.53 (m, 2H), 2.89 - 2.68 (m, 1H), 2.21 - 2.03 (m, 1H), 1.47 (s, 9H).

Intermediate 26: *tert*-Butyl (1-(2'-(dimethylphosphoryl)-2-fluoro-3-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)-2-oxopyrrolidin-3-yl)carbamate




To a solution of Intermediate 25 (0.60 g, 1.4 mmol) in 1,4-dioxane (10 mL) at rt, were added (2-bromophenyl)dimethylphosphine oxide (0.41 mg, 1.8 mmol), 4,4,4',4',5,5,5',5'-octamethyl-2,2'-bi(1,3,2-dioxaborolane) (0.52 g, 2.0 mmol), and, K₂CO₃ (0.56 mg, 4.1 mmol). The reaction mixture was purged with nitrogen for 5 min and then charged with PdCl₂(dppf).DCM (56 mg, 0.070 mmol). The reaction mixture was again purged with nitrogen for 3 min and heated at 100 °C for 16 h. The reaction mixture was cooled, filtered through a Celite pad and the filtrate was concentrated under reduced pressure. The crude product was purified by column chromatography (EtOAc/Pet. Ether) to give the Intermediate 26 (0.30 g, 0.58 mmol, 43% yield) as an orange solid. MS(ESI) m/z: 515.1

Intermediate 27: 3-Amino-1-(2'-(dimethylphosphoryl)-2-fluoro-3-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)pyrrolidin-2-one hydrochloride




To an ice cooled solution of Intermediate 26 (0.30 g, 0.58 mmol) in 1,4-dioxane (5 mL) under argon atmosphere at rt, was added 4M HCl in 1,4-dioxane (5 mL) and stirred at rt for 2 h. The solvent was evaporated under reduced pressure. The solid was triturated with pet. ether (10 ml x 2) and dried to give the Intermediate 27 (0.20 g, 0.48 mmol, 83% yield) as a brown gummy solid. MS(ESI) m/z: 415.1 [M+H]⁺.

Intermediate 28: (S)-1-(4-Bromo-2,3-difluorophenyl)-5-(hydroxymethyl)pyrrolidin-2-one




To a solution of (S)-5-(hydroxymethyl)pyrrolidin-2-one (1.0 g, 8.7 mmol) in 1,4-dioxane (15 mL) at rt, were added 1,4-dibromo-2,3-difluorobenzene (2.4 g, 8.7 mmol), K₃PO₄ (3.7 g, 17 mmol) and, N,N'-dimethylethylenediamine (0.15 g, 1.7 mmol). The reaction mixture was purged with nitrogen for 5 min and then charged with copper (I) iodide (0.17 g, 0.87 mmol). The reaction mixture was again purged with nitrogen for 3 min and heated at 95 °C for 12 h. The reaction mixture was cooled, filtered through a Celite pad and the filtrate was concentrated under reduced pressure. The crude compound was purified by column chromatography (3% MeOH-CHCl₃) to yield Intermediate 28 (0.80 g, 2.6 mmol, 30% yield) as a yellowish solid. MS(ESI) m/z: 305.9 [M+H]⁺. ¹H NMR (400 MHz, DMSO-d₆) δ = 7.59 (ddd, J = 8.8, 7.0, 2.3 Hz, 1H), 7.24 (ddd, J = 9.0, 7.0, 2.0 Hz, 1H), 4.80 (t, J = 5.0 Hz, 1H), 4.20 - 4.04 (m, 1H), 3.37 (dd, J = 5.0, 4.0 Hz, 2H), 2.56 - 2.45 (m, 1H), 2.43 - 2.35 (m, 1H), 2.25 - 2.18 (m, 1H), 2.07 - 1.95 (m, 1H).

Intermediate 29: (*S*)-1-(4-Bromo-2,3-difluorophenyl)-5-(((*tert*-butyldimethylsilyl)oxy)methyl)pyrrolidin-2-one



5 To a stirred solution of Intermediate 28 (0.80 g, 2.6 mmol) in DCM (10 mL) under argon atmosphere at rt, were added TEA (0.55 mL, 3.9 mmol) and *tert*-butyldimethylsilyl chloride (0.43 g, 2.9 mmol). The reaction mixture was stirred at rt for 16 h. The reaction mixture was quenched with aqueous saturated NH₄Cl (20 mL) and extracted with DCM (20 mL x 2). The combined organic layers were washed with brine (20 mL), dried over
10 Na₂SO₄, concentrated under reduced pressure. The crude product was purified via column chromatography (15% EtOAc-Pet. ether) to give Intermediate 29 (0.85 g, 2.0 mmol, 77% yield) as a white solid. MS(ESI) *m/z*: 420.2 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃) δ = 7.35 (m, 1H), 7.15 - 7.05 (m, 1H), 4.24 - 4.17 (m, 1H), 3.62 - 3.49 (m, 2H), 2.69 - 2.63 (m, 1H), 2.54 - 2.50 (m, 1H), 2.41 - 2.26 (m, 1H), 2.14 - 2.03 (m, 1H), 0.84 (s, 9H), -0.02
15 (s, 3H), -0.05 (s, 3H).


Intermediate 30: (*3R,5S*)-3-azido-1-(4-bromo-2,3-difluorophenyl)-5-(((*tert*-butyldimethylsilyl)oxy)methyl)pyrrolidin-2-one



20 To a stirred solution of Intermediate 29 (1.0 g, 2.4 mmol) in THF (20 mL) at -78 °C, lithium diisopropylamide (1.8 mL, 3.6 mmol) was added and stirring was continued for 30 min. Then, a solution of 2,4,6-triisopropylbenzenesulfonyl azide (1.0 g, 3.3 mmol) in THF (5 mL) was cannulated into the mixture and the mixture was stirred for 4 h. Then, the reaction mixture was quenched with saturated aqueous NaHCO₃ (2 mL) and gradually
25 warmed to rt. The reaction mixture was diluted with EtOAc (40 mL), washed with H₂O


(10 mL x 2) and brine (20 mL), dried over Na₂SO₄, and concentrated under reduced pressure. The crude residue was purified by column chromatography (15% EtOAc-Pet ether, 40 g column) to yield Intermediate 30 (0.85 g, 1.8 mmol, 77% yield) as an orange solid. MS(ESI) m/z: 461.1 [M+H]⁺. ¹H NMR (400 MHz, DMSO-d₆) δ = 7.65 (ddd, *J* = 9.0, 7.0, 2.0 Hz, 1H), 7.36 - 7.28 (m, 1H), 4.72 - 4.64 (m, 1H), 4.33 - 4.24 (m, 1H), 3.63 - 3.52 (m, 2H), 2.48 - 2.28 (m, 1H), 2.14 (dt, *J* = 13.3, 8.7 Hz, 1H), 0.91 - 0.66 (m, 9H), -0.06 (s, 3H), -0.08 (s, 3H).

Intermediate 31: *tert*-Butyl ((3*R*,5*S*)-1-(4-bromo-2,3-difluorophenyl)-5-(((*tert*-butyldimethylsilyl)oxy)methyl)-2-oxopyrrolidin-3-yl)carbamate




To a stirred solution Intermediate 30 (0.85 g, 1.8 mmol) in THF (8 mL) were added triphenylphosphine (0.73 g, 2.8 mmol) and H₂O (0.8 mL). The reaction mixture was stirred at rt for 3 h and heated at 65 °C for 6 h. The solvent was evaporated under reduced pressure and used for the next step without further purification. To a stirred solution of crude (5*S*)-3-amino-1-(4-bromo-2,3-difluorophenyl)-5-(((*tert*-butyldimethylsilyl)oxy)methyl)pyrrolidin-2-one in DCM (5 mL) under argon atmosphere at rt, were added Boc-anhydride (0.84 mL, 3.6 mmol) and TEA (0.75 mL, 5.4 mmol). The reaction mixture was stirred at rt for 2 h. Then, the reaction mixture was quenched with aqueous saturated NH₄Cl (15 mL) and extracted with CHCl₃ (20 mL x 2). The combined organic layers were washed with brine (15 mL), dried over Na₂SO₄, concentrated under reduced pressure. The crude product was purified via column chromatography (15% EtOAc-Pet. ether) to give Intermediate 31 (0.80 g, 1.5 mmol, 83% yield) as a white solid. MS(ESI) m/z: 535.2 [M+H]⁺. ¹H NMR (400 MHz, CDCl₃) δ = 7.41 - 7.32 (m, 1H), 7.15 - 7.01 (m, 1H), 5.02 - 5.22 (m, 1H), 4.73 - 4.61 (m, 1H), 4.52 - 4.43 (m, 1H), 4.21 - 4.10 (m, 1H), 3.61 - 3.52 (m, 1H), 2.63 - 2.51 (m, 1H), 2.35 - 2.31 (m, 1H), 1.43 (s, 9H), 0.84 (s, 9H), -0.06 (s, 3H), -0.08 (s, 3H).

Intermediate 32: *tert*-Butyl ((3*R*,5*S*)-5-(((*tert*-butyldimethylsilyl)oxy)methyl)-1-(2'-(dimethylphosphoryl)-2,3-difluoro-[1,1'-biphenyl]-4-yl)-2-oxopyrrolidin-3-yl)carbamate



To a stirred solution of Intermediate 31 (0.80 g, 1.49 mmol) in 1,4-dioxane (10 mL), were added (2-bromophenyl)dimethylphosphine oxide (0.52 g, 2.2 mmol), bis(pinacolato)diboron (1.1 g, 4.5 mmol), and K₂CO₃ (0.42 mg, 3.0 mmol). The reaction mixture was purged with argon for 5 min and charged with PdCl₂(dppf)-CH₂Cl₂ adduct (0.13 mg, 0.15 mmol). The reaction mixture was again purged with nitrogen for 3 min and heated at 100 °C for 12 h. The reaction mixture was cooled, filtered through a Celite pad and the filtrate was concentrated under reduced pressure. The crude product was purified by column chromatography (MeOH/CHCl₃) to yield Intermediate 32 (0.40 g, 0.66 mmol, 44% yield) as a brown solid. MS(ESI) *m/z*: 609.3 [M+H]⁺. ¹H NMR (400 MHz, DMSO-*d*₆) δ = 8.01 - 7.97 (m, 1H), 7.85 - 7.81 (m, 1H), 7.80 - 7.64 (m, 2H), 7.59 - 7.27 (m, 3H), 4.66 - 4.56 (m, 1H), 4.42 - 4.33 (m, 1H), 3.76 - 3.55 (m, 2H), 2.46 - 2.28 (m, 2H), 1.54 - 1.39 (m, 15H), 0.84 (s, 9H), 0.06 (s, 3H), -0.08 (s, 3H).


Intermediate 33: (3*R*,5*S*)-3-amino-1-(2'-(dimethylphosphoryl)-2,3-difluoro-[1,1'-biphenyl]-4-yl)-5-(hydroxymethyl)pyrrolidin-2-one hydrochloride



To an ice cooled solution of Intermediate 32 (0.40 g, 0.66 mmol) in 1,4-dioxane (1 mL) under argon atmosphere at rt, was added 4M HCl in 1,4-dioxane (2.5 mL, 10 mmol)


and stirred at rt for two h. The solvent was evaporated under reduced pressure and the gummy solid was further triturated with EtOAc (10 ml x 2) and dried to give Intermediate 33 (0.22 g, 0.56 mmol, 85% yield) as a brown solid. MS(ESI) m/z : 395.1 $[M+H]^+$. 1H NMR (400 MHz, DMSO- d_6) δ = 7.97 - 7.88 (m, 1H), 7.82 - 7.72 (m, 1H), 7.69 - 7.47 (m, 2H), 7.43 - 7.25 (m, 2H), 4.35 - 4.29 (m, 2H), 4.11 - 3.86 (m, 4H), 3.55 - 3.34 (m, 2H), 2.48 - 2.27 (m, 2H), 1.52 (br d, J = 13.1 Hz, 6H).

Intermediate 34: 1-(4-Bromo-2,3-difluorophenyl)-5-methylpyrrolidin-2-one



To a solution of racemic 5-methylpyrrolidin-2-one (1.0 g, 10 mmol) in 1,4-dioxane (10 mL) at rt, were added 1,4-dibromo-2,3-difluorobenzene (3.3 g, 12 mmol), K_3PO_4 (4.3 g, 20 mmol) and N,N' -dimethylethylenediamine (0.18 g, 2.0 mmol). The reaction mixture was purged with nitrogen for 5 min and then charged with copper (I) iodide (0.19 g, 1.0 mmol). The reaction mixture was again purged with nitrogen for 3 min and heated at 80 °C for 6 h. The reaction mixture was cooled, filtered through a Celite pad and the filtrate was concentrated under reduced pressure. The crude product was purified by column chromatography (pet. ether-EtOAc) to give Intermediate 34 (0.90 g, 3.1 mmol, 31% yield) as brown liquid. MS(ESI) m/z : 291.0 $[M+H]^+$. 1H NMR (400MHz, $CDCl_3$) δ = 7.38 - 7.01 (m, 1 H), 6.99 - 6.96 (m, 1 H), 4.24 - 4.20 (m, 1H), 2.62-2.51 (m, 2 H), 2.50-2.38 (m, 1 H), 1.81-1.76 (m, 1 H), 1.17 (d, J = 6.4 Hz, 3H).


Intermediate 35: 3-Azido-1-(4-bromo-2,3-difluorophenyl)-5-methylpyrrolidin-2-one



To a stirred solution of Intermediate 34 (0.20 g, 0.69 mmol) in THF (3 mL) at -78 °C, lithium diisopropylamide (0.69 mL, 1.4 mmol) was added and stirring was continued

for 40 min. Then, a solution of 2,4,6-triisopropylbenzenesulfonyl azide (0.26 g, 0.83 mmol) in THF (2 mL) was cannulated into the mixture and the mixture was stirred for 4 h. The mixture was then quenched with saturated aqueous NH_4Cl (10 mL), gradually warmed to rt and extracted with EtOAc (20 mL x 2). The combined organic layers were washed with brine (15 mL), dried over Na_2SO_4 , and concentrated under reduced pressure. The crude residue was purified by column chromatography (12% EtOAc-Pet ether) to yield Intermediate 35 (0.10 g, 0.21 mmol, 31% yield) as an orange solid. MS(ESI) m/z : 331.1 $[\text{M}+\text{H}]^+$. ^1H NMR (400MHz, CDCl_3) δ = 7.40 - 7.36 (m, 1 H), 7.04 - 6.98 (m, 1 H), 4.36 - 4.31 (m, 1H), 4.26 - 4.21 (m, 1H), 2.32 - 2.26 (m, 1 H), 2.14 - 2.043 (m, 1 H), 1.16 (d, J = 6.4 Hz, 3 H).

Intermediate 36: *tert*-Butyl (1-(4-bromo-2,3-difluorophenyl)-5-methyl-2-oxopyrrolidin-3-yl)carbamate




To a stirred solution 3-azido-1-(4-bromo-2,3-difluorophenyl)-5-methylpyrrolidin-2-one (1.5 g, 4.5 mmol) in THF (21 mL) were added triphenylphosphine (2.4 g, 9.1 mmol) and water (3 mL). The reaction mixture was stirred at rt for 30 min and heated to 65 °C for 6 h. The solvent was evaporated under reduced pressure and the crude product used for the next step without further purification.

To a stirred solution of crude 3-amino-1-(4-bromo-2,3-difluorophenyl)-5-methylpyrrolidin-2-one in DCM (20 mL) under argon atmosphere at rt, were added TEA (0.82 mL, 5.9 mmol) and Boc-anhydride (1.4 mL, 5.9 mmol). The reaction mixture was stirred at rt for 2 h. Then, the reaction mixture was quenched with aqueous saturated NH_4Cl (25 mL) and extracted with CHCl_3 (30 mL x 2). The combined organic layers were washed with brine (25 mL), dried over Na_2SO_4 , and concentrated under reduced pressure. The crude product was purified via column chromatography (50% EtOAc-Pet. ether) to give Intermediate 36 (1.3 g, 3.2 mmol, 82% yield) as a brown solid. MS(ESI) m/z : 405.2 $[\text{M}+\text{H}]^+$. ^1H NMR (400MHz, CDCl_3) δ = 7.39 - 7.26 (m, 1 H), 7.0 - 6.99 (m, 1


H), 5.29 - 5.07 (m, 1H), 4.58 - 4.31 (m, 1 H), 4.30 - 4.13 (m, 1 H), 2.51 - 2.04 (m, 1 H), 1.81 - 1.76 (m, 1 H), 1.45 (s, 9 H), 1.30 - 1.15 (m, 3 H). (Mixture of diastereomers)

Intermediate 37: *tert*-Butyl (1-(2'-(dimethylphosphoryl)-2,3-difluoro-[1,1'-biphenyl]-4-yl)-5-methyl-2-oxopyrrolidin-3-yl)carbamate



To a stirred solution of Intermediate 36 (1.0 g, 2.5 mmol) in 1,4-dioxane (15 mL), were added (2-bromophenyl)dimethylphosphine oxide (0.58 g, 2.5 mmol), bis(pinacolato)diboron (1.9 g, 7.4 mmol) and K₂CO₃ (1.0 g, 7.4 mmol). The reaction mixture was purged with nitrogen for 5 min and charged with PdCl₂(dppf).CH₂Cl₂ adduct (0.20 g, 0.25 mmol). The reaction mixture was again purged with nitrogen for 3 min and heated at 110 °C for 15 h. The mixture was cooled, filtered through a Celite pad and the filtrate was concentrated under reduced pressure. The crude compound was purified by flash chromatography (MeOH/CDCl₃) to give Intermediate 37 (0.70 g, 1.5 mmol, 59% yield) as a brown solid. MS(ESI) m/z: 479.2 [M+H]⁺. ¹H NMR (400MHz, CDCl₃) δ = 7.78-7.73 (m, 1H), 7.62 - 7.51 (m, 3H), 7.37 - 7.28 (m, 1H), 7.23 - 7.18 (m, 1H), 5.31-5.11 (m, 1H), 4.59 - 4.24 (m, 2H), 2.58-2.32 (m, 1H), 1.77 - 1.74 (m, 3H), 1.65 - 1.49 (m, 13H), 1.30 - 1.23 (m, 3H). (Mixture of diastereomers)


Intermediate 38: 3-Amino-1-(2'-(dimethylphosphoryl)-2,3-difluoro-[1,1'-biphenyl]-4-yl)-5-methylpyrrolidin-2-one hydrochloride



To an ice cooled solution of Intermediate 37 (0.70 g, 1.5 mmol) in 1,4-dioxane (5 mL) under argon atmosphere, was added 4M HCl (7.3 mL, 29 mmol) in 1,4-dioxane and stirred at rt for 2 h. The solvent was evaporated under reduced pressure to give

5 Intermediate 38 (0.50 g, 1.2 mmol, 82% yield) as brown liquid. MS(ESI) m/z : 379.2 $[M+H]^+$. 1H NMR (400MHz, $CDCl_3$) δ = 7.78 - 7.73 (m, 1H), 7.62 - 7.51 (m, 3H), 7.37 - 7.28 (m, 1H), 7.23 - 7.18 (m, 1H), 5.38 - 4.98 (m, 3H), 4.59 - 4.42 (m, 2H), 2.58-2.29 (m, 1H), 1.71 - 1.63 (m, 3H), 1.55 - 1.49 (m, 4H), 1.30 - 1.23 (m, 3H).


10 Example 1: (*R*)-*N*-(4'-(3-(3-(4-Chlorophenyl)ureido)-2-oxopyrrolidin-1-yl)-2',3'-difluoro-[1,1'-biphenyl]-2-yl)methanesulfonamide



To a stirred solution of Intermediate 5 (50 mg, 0.12 mmol), in DCE (2 mL) under nitrogen at rt, DIPEA (0.084 mL, 0.48 mmol), and 1-chloro-4-isocyanatobenzene (22.0 mg, 0.144 mmol) were added sequentially. The resulting reaction mixture was stirred at rt for 4 h. The reaction mixture was quenched with water and extracted with EtOAc. The organic layer was washed with ice cold water, brine, dried over Na_2SO_4 , and concentrated under reduced pressure. The crude compound was purified by reverse phase chromatography followed by chiral HPLC to Example 1 (16 mg, 0.028 mmol, 24%). RT = 1.732 min, 94.2%, (Method C); MS(ESI) m/z : 535.1 $[M+H]^+$; 1H NMR (400MHz, DMSO- d_6): δ 9.17 (s, 1H), 8.87 (s, 1H), 7.96 (s, 1H), 7.58 -7.29 (m, 10H), 4.64 - 4.48


(m, 1H), 3.96 -3.82 (m, 1H), 3.81 - 3.66 (m, 1H), 2.74 (s, 3H), 2.59 - 2.55 (m, 1H), 2.20-2.07 (m, 1H).

Example 25: *(R)*-1-(4-chloro-2-fluorophenyl)-3-(1-(2'-(dimethylphosphoryl)-2,3-difluoro-[1,1'-biphenyl]-4-yl)-2-oxopyrrolidin-3-yl)urea



DIEA (61 mL, 350 mmol) was added to a solution of Intermediate 10 (28 g, 70 mmol) in DCE (560 mL) slowly at 0 °C. After stirring for 2 min, Intermediate 11 (17 g, 63 mmol) was added, and the reaction mixture was slowly heated to 50 °C and stirred for 12 h. The reaction mixture was concentrated under reduced pressure, and the crude product was purified via column chromatography (5% MeOH in DCM). The product was purified by prep HPLC, washed with water, and dried to obtain Example 25 (29 g, 54 mmol, 78% yield) as a white solid.


Example 57: *(R)*-1-(1-(2'-(dimethylphosphoryl)-2,3-difluoro-[1,1'-biphenyl]-4-yl)-2-oxopyrrolidin-3-yl)-3-(2-fluoro-4-(trifluoromethyl)phenyl)urea



DIEA (5.5 mL, 6.2 mmol) was added to a solution of Intermediate 10 (2.5 g, 6.2 mmol) in DCE (70 mL) slowly at 0 °C. After stirring for 2 min, Intermediate 12 (1.7 g, 5.6 mmol) was added, and the reaction mixture was slowly heated to 50 °C and stirred for 8 h. The reaction mixture was concentrated under reduced pressure and the crude product

was purified via column chromatography (6.5% MeOH in DCM). The product was purified by prep HPLC, washed with water, and dried to obtain Example 57 (2.2 g, 3.9 mmol, 62% yield) as a white solid.


- 5 Example 73: *(R)*-1-(1-(4-(2-(dimethylphosphoryl)pyridin-3-yl)-2,3-difluorophenyl)-2-oxopyrrolidin-3-yl)-3-(2-fluoro-4-(trifluoromethyl)phenyl)urea



To a solution of Intermediate 14 (13 g, 24 mmol) and Intermediate 15 (5.6 g, 24 mmol) in 1,4 dioxane (260 mL) and water (26 mL), were added potassium phosphate (10 g, 48 mmol) and PdCl₂(dppf)-CH₂Cl₂ (2.0 g, 2.4 mmol). The reaction mixture was stirred for 30 min, slowly raised to 110 °C and stirred for 12 h. The mixture was filtered through Celite, and the plug was washed with EtOAc. The filtrate was dried over sodium sulfate and concentrated under reduced pressure. The crude compound was purified via column chromatography (2% MeOH in DCM) to yield Example 73 (9.1 g, 16 mmol, 66% yield).


15

Example 89: *(R)*-1-(1-(2-Cyclopropyl-2'-(dimethylphosphoryl)-3-fluoro-[1,1'-biphenyl]-4-yl)-2-oxopyrrolidin-3-yl)-3-(2-fluoro-4-(trifluoromethyl)phenyl)urea



To an ice cooled suspension of Intermediate 24 (0.13 g, 0.34 mmol) in DMF (2 mL) under argon atmosphere at rt, were added DIEA (0.18 mL, 1.0 mmol) and Intermediate 12 (0.12 g, 0.41 mmol). Then, the resulting solution was heated at 50 °C for 2 hours. The reaction mixture was concentrated under reduced pressure and the crude product was purified by reverse phase chromatography followed by chiral HPLC to give
 5 Example 89 (0.07 g, 0.12 mmol, 33% yield).


Example 91: *(R)*-1-(4-Chloro-2-fluorophenyl)-3-(1-(2'-(dimethylphosphoryl)-2-fluoro-3-(trifluoromethyl)-[1,1'-biphenyl]-4-yl)-2-oxopyrrolidin-3-yl)urea



10


To an ice cooled suspension of Intermediate 27 (0.080 g, 0.18 mmol) in DCE (2 mL) under argon atmosphere at rt, were added DIEA (0.06 mL, 0.36 mmol) and Intermediate 11 (0.060 g, 0.21 mmol). Then, the resulting solution was heated at 50 °C for 2 h. The reaction mixture was concentrated under reduced pressure, and the crude product was purified by reverse phase chromatography followed by chiral HPLC to yield
 15 Example 91 (0.060 g, 0.11 mmol, 60% yield) as a white solid.

Example 109: 1-(4-Chloro-2-fluorophenyl)-3-(1-(2'-(dimethylphosphoryl)-2,3-difluoro-[1,1'-biphenyl]-4-yl)-5-methyl-2-oxopyrrolidin-3-yl)urea








To an ice cooled suspension of Intermediate 38 (0.20 g, 0.48 mmol) in DCE (10 mL) under argon atmosphere at rt, were added DIEA (0.34 mL, 1.9 mmol) and Intermediate 11 (73 mg, 0.25 mmol). Then, the resulting solution was heated at 50 °C for 3 h. The reaction mixture was concentrated under reduced pressure and the crude product was purified by reverse phase chromatography followed by chiral HPLC to yield Example 109 (0.02 g, 0.04 mmol, 8.3% yield) as a white solid.







Example 112: 1-(4-Chloro-2-fluorophenyl)-3-((3R,5S)-1-(2'-(dimethylphosphoryl)-2,3-difluoro-[1,1'-biphenyl]-4-yl)-5-(hydroxymethyl)-2-oxopyrrolidin-3-yl)urea









To an ice cooled suspension of Intermediate 22 (0.11 g, 0.28 mmol) in DMF (2 mL) under argon atmosphere at rt, were added DIEA (0.15 mL, 0.84 mmol) and Intermediate 11 (0.11 g, 0.42 mmol). Then, the resulting solution was heated at 50 °C for 3 h. The reaction mixture was concentrated under reduced pressure and the crude product was purified by reverse phase chromatography followed by chiral HPLC to yield Example 112 (0.05 g, 0.08 mmol, 29% yield).






The following Examples in Table 1 were made by using the same procedure as shown above in Examples 1, 25, 57, 73, 89, 91, 109, and 112.






Ex	Structure	LCMS (M+H) +	HPLC Method, RT (min.) & Purity	¹ H NMR
2		490.1	Method D, RT = 1.75 min, 99.3%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.64 (d, J=2.2 Hz, 1H), 8.16 (t, J=8.8 Hz, 1H), 7.63 (d, J=7.6 Hz, 1H), 7.54 - 7.32 (m, 4H), 7.31 - 7.08 (m, 4H), 5.16 (t, J=5.5 Hz, 1H), 4.63 - 4.48 (m, 1H), 4.35 (d, J=5.4 Hz, 2H), 3.97 - 3.85 (m, 1H), 3.84 - 3.74 (m, 1H), 2.65 - 2.56 (m, 1H), 2.16 - 2.03 (m, 1H).
3		472.1	Method D, RT = 1.89 min, 98.9%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.86 (s, 1H), 7.63 (d, J=7.6 Hz, 1H), 7.54 - 7.34 (m, 5H), 7.34 - 7.14 (m, 4H), 6.72 (d, J=7.3 Hz, 1H), 5.16 (t, J=5.4 Hz, 1H), 4.55 (dt, J=10.1, 8.1 Hz, 1H), 4.35 (d, J=5.4 Hz, 2H), 3.95 - 3.83 (m, 1H), 3.83 - 3.72 (m, 1H), 2.60 - 2.54 (m, 1H), 2.19 - 2.06 (m, 1H).
4		588.2 (M+N H ₄) ⁺	Method D, RT = 1.66 min, 95.2%	¹ H NMR (400MHz, DMSO-d ₆) δ = 9.18 (s, 1H), 8.21 (s, 1H), 7.97 (s, 1H), 7.60 - 7.45 (m, 2H), 7.45 - 7.31 (m, 4H), 7.26 - 7.20 (m, 1H), 7.03 - 6.97 (m, 1H), 4.60 - 4.52 (m, 1H), 3.98 - 3.82 (m, 2H), 3.79 - 3.73 (m, 1H), 2.87 (s, 3H), 2.18 - 2.06 (m, 1H).
5		553.1	Method C, RT = 1.79 min, 96.8%	¹ H NMR (400MHz, DMSO-d ₆) δ = 9.18 (s, 1H), 8.65 (d, J=2.2 Hz, 1H), 8.16 (t, J=8.9 Hz, 1H), 7.56 - 7.29 (m, 6H), 7.29 - 7.13 (m, 3H), 4.64 - 4.48 (m, 1H), 3.96 - 3.87 (m, 1H), 3.78-3.74 (m, 1H), 2.74 (s, 3H), 2.62 - 2.56 (m, 1H), 2.13 - 2.01 (m, 1H).
6		569.1	Method C, RT = 1.86 min, 97.2%	¹ H NMR (400MHz, DMSO-d ₆) δ = 9.16 (s, 1H), 7.96 (s, 1H), 7.70 - 7.56 (m, 4H), 7.55 - 7.45 (m, 2H), 7.44 - 7.30 (m, 3H), 7.28 - 7.20 (m, 1H), 6.86 (d, J=7.6 Hz, 1H), 4.60-4.54 (m, 1H), 3.96 - 3.86 (m, 1H), 3.77-3.75 (m, 1H), 2.74 (s, 3H), 2.61 - 2.54 (m, 1H), 2.21 - 2.06 (m, 1H).






7		538.2	Method D, RT = 1.75 min, 95.9%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.65 (s, 1H), 8.23 - 8.05 (m, 2H), 7.91 - 7.68 (m, 2H), 7.51(d, J=7.3 Hz, 1H), 7.46 - 7.35 (m, 2H), 7.35 - 7.25 (m, 1H), 7.20 - 7.18 (m, 2H), 4.63 - 4.48(m, 1H), 3.94 - 3.87 (m, 1H), 3.83 - 3.69 (m, 1H), 3.05 (s, 3H), 2.63 - 2.54 (m, 1H), 2.10-2.05 (m, 1H).
8		467.2	Method D, RT = 1.85 min, 100%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.88 (s, 1H), 8.05 (d, J=8.1 Hz, 1H), 7.96 - 7.85 (m, 1H), 7.79 - 7.63 (m, 2H), 7.59 - 7.49 (m, 1H), 7.49 - 7.37 (m, 3H), 7.30 (d, J=8.8 Hz, 2H), 6.75 (d, J=7.6 Hz, 1H), 4.63 - 4.51 (m, 1H), 3.98 - 3.87 (m, 1H), 3.87 - 3.73 (m, 1H), 2.58 (br. s., 1H), 2.18 - 2.11 (m, 1H).
9		485.1	Method D, RT = 1.91 min, 96.8%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.65 (s, 1H), 8.15 (t, J=8.9 Hz, 1H), 8.04 (d, J=7.9 Hz, 1H), 7.94 - 7.83 (m, 1H), 7.78 - 7.60 (m, 2H), 7.57 - 7.49 (m, 1H), 7.48 - 7.34 (m, 2H), 7.20 (d, J=7.6Hz, 2H), 4.64 - 4.52 (m, 1H), 3.98 - 3.88 (m, 1H), 3.86 - 3.76 (m, 1H), 2.65 - 2.55 (m, 1H), 2.18 - 2.00 (m, 1H).
10		520.1	Method D, RT = 1.69 min, 99.1%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.88 (s, 1H), 8.15 (dd, J=7.7, 1.3 Hz, 1H), 7.89 - 7.71 (m, 2H), 7.52 (dd, J=7.3, 1.5 Hz, 1H), 7.49 - 7.37 (m, 3H), 7.36 - 7.22 (m, 3H), 6.74 (d, J=7.6 Hz, 1H), 4.62 - 4.49 (m, 1H), 3.97 - 3.85 (m, 1H), 3.84 - 3.73 (m, 1H), 3.06 (s, 3H), 2.60 - 2.54 (m, 1H), 2.19 - 2.07 (m, 1H).
11		554.2	Method D, RT = 1.81 min, 97.1%	¹ H NMR (400MHz, DMSO-d ₆) δ = 9.18 (s, 1H), 8.15 (dd, J=7.8, 1.5 Hz, 1H), 7.90 - 7.72 (m, 2H), 7.70 - 7.55 (m, 4H), 7.52 (dd, J=7.5, 1.3 Hz, 1H), 7.42 (t, J=7.6 Hz, 1H), 7.36 - 7.25 (m, 1H), 6.86 (d, J=7.3 Hz, 1H), 4.65 - 4.50 (m, 1H), 4.00 - 3.86 (m, 1H), 3.81-3.77 (m, 1H), 3.06 (s, 3H), 2.62 - 2.54 (m, 1H), 2.16 - 2.11 (m, 1H).
12		485.2	Method D, RT = 1.50 min, 96.9%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.94 (s, 1H), 7.62 (s, 1H), 7.49 - 7.34 (m, 7H), 7.31 - 7.23 (m, 2H), 6.81 (d, J=7.3 Hz, 1H), 4.61 - 4.50 (m, 1H), 4.16 (q, J=6.8 Hz, 1H), 3.92 - 3.84 (m, 1H), 3.79 - 3.74 (m, 1H), 3.60 - 3.20 (m, 2H), 2.61 - 2.54 (m, 1H), 2.17 - 2.07 (m, 1H), 1.35 (d, J=6.6 Hz, 3H).

13		554.1	Method D, RT = 1.67 min, 100%	¹ H NMR (400MHz, DMSO-d ₆) δ = 9.40 (s, 1H), 9.15 (s, 1H), 8.77 (d, J=7.1 Hz, 1H), 8.17 (d, J=2.2 Hz, 1H), 7.98 (dd, J=10.3, 2.2 Hz, 1H), 7.52 - 7.42 (m, 2H), 7.41 - 7.27 (m, 3H), 7.20 (t, J=7.1 Hz, 1H), 4.68 - 4.55 (m, 1H), 3.95-3.81 (m, 2H), 2.86 (s, 3H), 2.60 - 2.55 (m, 1H), 2.28 - 2.14 (m, 1H).
14		533.2	Method C, RT = 2.07 min, 99.2%	¹ H NMR (400MHz, DMSO-d ₆) δ = 9.19 (s, 1H), 8.96 (br. s., 1H), 7.75 (d, J=8.8 Hz, 2H), 7.65 - 7.56 (m, 4H), 7.45 (d, J=8.8 Hz, 2H), 7.41 - 7.27 (m, 4H), 6.81 (d, J=7.6 Hz, 1H), 4.58 - 4.49 (m, 1H), 3.89-3.79 (m, 2H), 2.73 (s, 3H), 2.51-2.50 (m, 1H), 2.10 - 2.01 (m, 1H).
15		537.2	Method D, RT = 1.54 min, 99.7%	¹ H NMR (400MHz, DMSO-d ₆) δ = 9.71 (s, 1H), 9.13 (s, 1H), 8.83 (s, 1H), 8.38 (s, 1H), 7.55 (d, J=7.6 Hz, 1H), 7.51 - 7.45 (m, 2H), 7.41 - 7.31 (m, 3H), 7.24 - 7.17 (m, 1H), 4.58 - 4.49 (m, 1H), 3.89-3.79 (m, 2H), 2.88 (s, 3H), 2.51-2.50 (m, 1H), 2.20 - 2.10 (m, 1H).
16		460.1	Method D, RT = 2.00 min, 92.1%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.88 (s, 1H), 7.62 - 7.50 (m, 2H), 7.50 - 7.42 (m, 3H), 7.42 - 7.33 (m, 3H), 7.33 - 7.22 (m, 2H), 6.75 (d, J=7.3 Hz, 1H), 4.65 - 4.49 (m, 1H), 3.97 - 3.84 (m, 1H), 3.84 - 3.70 (m, 1H), 2.61 - 2.53 (m, 1H), 2.21 - 2.04 (m, 1H).
17		536.1	Method D, RT = 1.66 min, 99.7%	¹ H NMR (400MHz, DMSO-d ₆) δ = 9.43 (s, 1H), 9.15 (br. s., 1H), 8.23 (d, J=2.7 Hz, 1H), 8.05 (d, J=7.8 Hz, 1H), 7.80 (dd, J=8.8, 2.7 Hz, 1H), 7.57 (d, J=8.8 Hz, 1H), 7.50 - 7.44 (m, 2H), 7.40 - 7.27 (m, 3H), 7.20 (t, J=7.1 Hz, 1H), 4.63 - 4.54 (m, 1H), 3.95-3.81 (m, 2H), 2.86 (s, 3H), 2.60 - 2.55 (m, 1H), 2.21 - 2.11 (m, 1H).
18		571.2	Method D, RT = 1.68 min, 100%	¹ H NMR (400MHz, DMSO-d ₆) δ = 10.13 (s, 1H), 9.12 (br. s., 1H), 9.05 (s, 1H), 8.73 (s, 1H), 7.89 (d, J=7.6 Hz, 1H), 7.52 - 7.46 (m, 2H), 7.41 - 7.31 (m, 3H), 7.25 - 7.16 (m, 1H), 4.66 - 4.60 (m, 1H), 3.95-3.81 (m, 2H), 2.60 - 2.55 (m, 1H), 2.88 (s, 3H), 2.24 - 2.13 (m, 1H).






19		461.1	Method D, RT = 1.73 min, 96.8%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.88 (s, 1H), 8.38 (d, J=4.6 Hz, 1H), 8.16 (t, J=7.9 Hz, 1H), 7.56 (t, J=6.2 Hz, 1H), 7.53 - 7.35 (m, 4H), 7.30 (d, J=9.0 Hz, 2H), 6.74 (d, J=7.3 Hz, 1H), 4.64 - 4.49 (m, 1H), 3.96 - 3.85 (m, 1H), 3.84 - 3.73 (m, 1H), 2.61 - 2.54 (m, 1H), 2.21 - 2.01 (m, 1H).
20		534.2	Method D, RT = 1.63 min, 98.8 %	¹ H NMR (400MHz, DMSO-d ₆) δ = 9.43 (s, 1H), 8.95 (s, 1H), 8.71 (d, J=2.2 Hz, 1H), 8.19 (dd, J=8.7, 1.8 Hz, 1H), 7.88 - 7.71 (m, 3H), 7.52 (d, J=8.8 Hz, 2H), 7.47 - 7.28 (m, 4H), 6.99 (d, J=7.3 Hz, 1H), 4.64 - 4.54 (m, 1H), 3.99 - 3.78 (m, 2H), 2.74 (s, 3H), 2.50-2.46 (m, 1H) 2.15 - 2.02 (m, 1H).
21		478.1	Method D, RT = 2.05 min, 100%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.65 (d, J=2.2 Hz, 1H), 8.16 (t, J=8.9 Hz, 1H), 7.61 - 7.50 (m, 2H), 7.50 - 7.32 (m, 5H), 7.27 - 7.14 (m, 2H), 4.64 - 4.50 (m, 1H), 3.98 - 3.85 (m, 1H), 3.84 - 3.75 (m, 1H), 2.64 - 2.55 (m, 1H), 2.17 - 2.01 (m, 1H).
22		485.2	Method D, RT = 1.55 min, 100%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.97 (s, 1H), 7.61 (s, 1H), 7.53 - 7.33 (m, 7H), 7.33 - 7.22 (m, 2H), 6.84 (d, J=7.3 Hz, 1H), 4.55 (m, 1H), 4.14 (q, J=6.8 Hz, 1H), 3.91 - 3.84 (m, 1H), 3.78 - 3.73 (m, 1H), 2.64 - 2.55 (m, 1H), 2.12 (m, 1H), 1.34 (d, J=6.6 Hz, 3H).
23		554.2	Method D, RT = 1.46 min, 94.8%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.83 (br. s., 1H), 8.63 (br. s., 1H), 8.16 (t, J = 8.9 Hz, 1H), 8.08 (br. s., 1H), 7.47 - 7.32 (m, 2H), 7.32 - 7.10 (m, 4H), 6.65 (br. s., 1H), 4.64 - 4.48 (m, 1H), 3.92 - 3.82 (m, 1H), 3.75 (t, J = 8.7 Hz, 1H), 2.95 (br. s., 3H), 2.63 - 2.54 (m, 1H), 2.21 - 1.96 (m, 1H).
24		521.1	Method D, RT = 1.62 min, 99.7%	¹ H NMR (400MHz, DMSO-d ₆) δ 9.47 (s, 1H), 8.25 (d, J=2.7 Hz, 1H), 8.15 (d, J=7.6 Hz, 1H), 8.01 - 7.89 (m, 1H), 7.88 - 7.72 (m, 3H), 7.65 (d, J=9.3 Hz, 1H), 7.55 - 7.46 (m, 1H), 7.43 (t, J=7.2 Hz, 1H), 7.31 (t, J=7.0 Hz, 1H), 4.67 - 4.56 (m, 1H), 3.97 - 3.87 (m, 1H), 3.85 - 3.77 (m, 1H), 3.06 (s, 3H), 2.63 - 2.55 (m, 1H), 2.20 - 2.10 (m, 1H).






25		536.2	Method D, RT = 1.476 min, 98.6%	^1H NMR (400 MHz, DMSO- d_6) δ = 8.64 (br. s., 1H), 8.16 (t, J = 9.0 Hz, 1H), 7.97 - 7.84 (m, 1H), 7.69 - 7.57 (m, 2H), 7.43 (dd, J = 11.0, 2.5 Hz, 1H), 7.40 - 7.32 (m, 2H), 7.32 - 7.23 (m, 1H), 7.23 - 7.12 (m, 2H), 4.56 (dt, J = 9.9, 8.1 Hz, 1H), 3.95 - 3.85 (m, 1H), 3.78 (br. t., J = 8.8 Hz, 1H), 2.63 - 2.54 (m, 1H), 2.14 - 1.99 (m, 1H), 1.52 (br. d., J = 13.1 Hz, 6H).
26		557.2	Method D, RT = 1.89 min, 99.7%	^1H NMR (400MHz, DMSO- d_6) δ = 8.95 (br. s., 1H), 8.63 (d, J = 2.2 Hz, 1H), 8.17 (t, J = 8.9 Hz, 1H), 7.52 - 7.24 (m, 7H), 7.23 - 7.13 (m, 2H), 7.12 (s, 1H), 4.63 - 4.47 (m, 1H), 3.82 (td, J = 9.5, 6.5 Hz, 1H), 3.71 - 3.59 (m, 1H), 2.86 (s, 3H), 2.63 - 2.56 (m, 1H), 2.17 - 2.04 (m, 1H), 2.01 - 1.95 (m, 1H), 1.04 - 0.83 (m, 2H), 0.80 - 0.58 (m, 2H).
27		559.2	Method D, RT = 1.868 min, 99.6%	^1H NMR (400MHz, DMSO- d_6) δ = 9.01 (s, 1H), 8.62 (d, J = 2.0 Hz, 1H), 8.17 (t, J = 8.9 Hz, 1H), 7.55 (d, J = 2.0 Hz, 1H), 7.48 - 7.32 (m, 6H), 7.32 - 7.26 (m, 1H), 7.25 - 7.12 (m, 2H), 4.60 - 4.50 (m, 1H), 3.82 - 3.71 (m, 1H), 3.56 (t, J = 9.3 Hz, 1H), 3.06 - 3.02 (m, 1H), 2.81 (s, 3H), 2.65 - 2.56 (m, 1H), 2.12 - 2.02 (m, 1H), 1.31 - 1.13 (m, 6H).
28		464.1	Method D, RT = 1.732 min, 98.6%	^1H NMR (400MHz, DMSO- d_6) δ = 8.62 (d, J = 2.2 Hz, 1H), 8.15 (t, J = 8.9 Hz, 1H), 7.84 (d, J = 2.4Hz, 1H), 7.80 - 7.68 (m, 1H), 7.49 - 7.28 (m, 2H), 7.24 - 7.06 (m, 2H), 6.73 - 6.61 (m, 1H), 4.63- 4.46 (m, 1H), 3.94 (s, 3H), 3.91 - 3.78 (m, 1H), 3.78 - 3.67 (m, 1H), 2.64 - 2.53 (m, 1H), 2.16 - 2.01 (m, 1H).
29		572.2	Method C, RT = 1.887 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.93 (s, 1H), 8.41 (t, J = 8.4 Hz, 1H), 8.15 (dd, J = 7.9, 1.3 Hz, 1H), 7.90 - 7.72 (m, 2H), 7.66 (dd, J = 11.5, 1.7 Hz, 1H), 7.51 (dd, J = 7.5, 1.3 Hz, 2H), 7.46 - 7.37 (m, 1H), 7.37 - 7.21 (m, 2H), 4.65 - 4.51 (m, 1H), 4.00 - 3.87 (m, 1H), 3.81 (t, J = 8.8 Hz, 1H), 3.06 (s, 3H), 2.65 - 2.56 (m, 1H), 2.20 - 2.02 (m, 1H).

30		580.1	Method C, RT = 1.632 min, 98.7%	^1H NMR (400MHz, DMSO- d_6) δ = 8.65 (s, 1H), 8.14 (t, J = 8.8 Hz, 1H), 8.01 - 7.87 (m, 1H), 7.72 - 7.57 (m, 2H), 7.48 - 7.36 (m, 2H), 7.29 (m, 2H), 7.23 - 7.16 (m, 1H), 7.10 (d, J = 7.6 Hz, 1H), 4.70 - 4.58 (m, 1H), 4.39 - 4.27 (m, 1H), 3.51 - 3.43 (m, 1H), 3.39 - 3.35 (m, 1H), 3.24 (s, 3H), 2.46 - 2.44 (m, 1H), 2.34 - 2.26 (m, 1H), 1.61 - 1.40 (m, 6H)
31		524.2	Method D, RT = 1.719min, 99.2%	^1H NMR (400MHz, DMSO- d_6) δ = 8.93 (d, J = 2.4 Hz, 1H), 8.41 (t, J = 8.2 Hz, 1H), 7.72 - 7.58 (m, 2H), 7.55 - 7.31 (m, 5H), 7.30 - 7.17 (m, 2H), 5.16 (t, J = 5.5 Hz, 1H), 4.64 - 4.53 (m, 1H), 4.35 (d, J = 5.6 Hz, 2H), 3.97 - 3.84 (m, 1H), 3.80 (t, J = 8.7 Hz, 1H), 2.61 (dd, J = 19.4, 6.7 Hz, 1H), 2.15 - 2.03 (m, 1H).
32		535.2	Method C, RT = 1.550 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 9.11 (br. s., 1H), 8.63 (d, J = 2.4 Hz, 1H), 8.16 (t, J = 8.9 Hz, 1H), 7.56 (t, J = 8.2 Hz, 1H), 7.50 - 7.28 (m, 7H), 7.24 - 7.12 (m, 2H), 4.60 - 4.50 (m, 1H), 3.92 - 3.82 (m, 1H), 3.74 (t, J = 8.4 Hz, 1H), 2.86 (s, 3H), 2.60 (dd, J = 12.0, 8.1 Hz, 1H), 2.14 - 2.03 (m, 1H).
33		587.2	Method C, RT = 1.726 min, 98.1%	^1H NMR (400MHz, DMSO- d_6) δ = 9.12 (br. s., 1H), 8.93 (d, J = 2.9 Hz, 1H), 8.41 (t, J = 8.3 Hz, 1H), 7.66 (d, J = 11.7 Hz, 1H), 7.56 - 7.43 (m, 3H), 7.43 - 7.27 (m, 4H), 7.23 (t, J = 7.1 Hz, 1H), 4.66 - 4.54 (m, 1H), 4.00 - 3.86 (m, 1H), 3.78 (t, J = 9.2 Hz, 1H), 2.91 (s, 3H), 2.66 - 2.57 (m, 1H), 2.16 - 2.04 (m, 1H).
34		565.2	Method C, RT = 1.606 min, 96.7%	^1H NMR (400MHz, DMSO- d_6) δ = 8.78 (s, 1H), 8.63 (d, J = 2.7 Hz, 1H), 8.16 (t, J = 8.9 Hz, 1H), 7.53 - 7.46 (m, 1H), 7.46 - 7.36 (m, 2H), 7.34 - 7.15 (m, 5H), 7.11 (dd, J = 8.4, 1.6 Hz, 1H), 4.60 - 4.49 (m, 1H), 3.94 - 3.80 (m, 1H), 3.74 - 3.72 (m, 1H), 3.66 (s, 3H), 2.90 (s, 3H), 2.63 - 2.54 (m, 1H), 2.12 - 2.01 (m, 1H).






35		502.1	Method D, RT = 1.634 min, 93.3%	^1H NMR (400MHz, DMSO- d_6) δ = 8.63 (d, J = 2.2 Hz, 1H), 8.21 - 8.10 (m, 1H), 7.59 (d, J = 7.1 Hz, 1H), 7.50 - 7.35 (m, 2H), 7.33 (t, J = 8.1 Hz, 1H), 7.28 - 7.12 (m, 4H), 7.06 (dd, J = 8.1, 1.7 Hz, 1H), 5.06 (t, J = 5.5 Hz, 1H), 4.59 - 4.48 (m, 1H), 4.33 - 4.29 (m, 2H), 3.94 - 3.81 (m, 1H), 3.75 (t, J = 8.8 Hz, 1H), 3.55 (s, 3H), 2.59 (dd, J = 12.2, 7.3 Hz, 1H), 2.14 - 1.98 (m, 1H).
36		565.1	Method D, RT = 1.669 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.68 (s, 1H), 8.37 (s, 1H), 8.16 (t, J = 8.9 Hz, 1H), 7.56 - 7.37 (m, 3H), 7.35 - 7.15 (m, 5H), 7.15 - 7.05 (m, 1H), 4.59 - 4.49 (m, 1H), 3.87 (s, 3H), 3.79 (dd, J = 9.7, 6.5 Hz, 1H), 3.69 (t, J = 8.6 Hz, 1H), 2.88 (s, 3H), 2.59 (dd, J = 11.7, 7.3 Hz, 1H), 2.13 - 1.96 (m, 1H).
37		593.2	Method C, RT = 1.902min, 97.5%	^1H NMR (400MHz, DMSO- d_6) δ = 8.97 (br. s., 1H), 8.49 - 8.30 (m, 2H), 7.65 (d, J = 11.7 Hz, 1H), 7.55 (d, J = 2.0 Hz, 1H), 7.52 (d, J = 9.0 Hz, 1H), 7.48 - 7.22 (m, 7H), 4.59 - 4.52 (m, 1H), 3.83 - 3.77 (m, 1H), 3.57 - 3.55 (m, 1H), 3.06 - 3.02 (m, 1H), 2.87 (s, 3H), 2.66 - 2.58 (m, 1H), 2.14 - 2.01 (m, 1H), 1.31 - 1.13 (m, 6H).
38		496.2	Method C, RT = 1.863 min, 92.2%	^1H NMR (400MHz, DMSO- d_6) δ = 8.67 (s, 1H), 8.38 (s, 1H), 8.17 (m, 1H), 7.59 (d, J = 7.3 Hz, 1H), 7.49 - 7.30 (m, 4H), 7.30 - 7.11 (m, 4H), 4.61 - 4.50 (m, 1H), 4.43 (s, 2H), 3.91 (s, 1H), 3.81 - 3.76 (m, 1H), 3.59 - 3.56 (m, 1H), 3.04 - 3.01 (m, 1H), 2.64 - 2.55 (m, 1H), 2.14 - 2.03 (m, 1H), 1.31 - 1.13 (m, 6H).
39		602.1	Method C, RT = 1.731 min, 92.6%	^1H NMR (400MHz, DMSO- d_6) δ = 8.92 (br. s., 1H), 8.41 (t, J = 8.4 Hz, 1H), 8.15 (d, J = 7.8 Hz, 1H), 7.91 - 7.76 (m, 2H), 7.66 (br. d., J = 12.0 Hz, 1H), 7.57 - 7.47 (m, 2H), 7.43 - 7.29 (m, 2H), 7.25 (br. d., J = 6.8 Hz, 1H), 5.13 (br. d., J = 1.7 Hz, 1H), 4.78 - 4.65 (m, 1H), 4.27 - 4.13 (m, 1H), 3.54 - 3.42 (m, 2H), 3.06 (s, 3H), 2.57 - 2.24 (m, 1H), 2.35 - 2.31 (m, 1H).






40		544.2	Method D, RT = 1.851 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.63 (d, J = 2.0 Hz, 1H), 8.17 (t, J = 8.9 Hz, 1H), 8.12 (dd, J = 7.9, 1.1 Hz, 1H), 7.82 - 7.75 (m, 1H), 7.73 - 7.64 (m, 1H), 7.54 - 7.36 (m, 3H), 7.34 - 7.25 (m, 2H), 7.23 - 7.09 (m, 2H), 4.57 - 4.52 (m, 1H), 3.84 - 3.74 (m, 1H), 3.59 (t, J = 9.0 Hz, 1H), 3.06 - 3.02 (m, 1H), 2.82 (s, 3H), 2.65 - 2.56 (m, 1H), 2.13 - 2.00 (m, 1H), 1.31 - 1.06 (m, 6H).
41		506.1	Method D, RT = 1.825 min, 97.2%	^1H NMR (400MHz, DMSO- d_6) δ = 9.15 (s, 1H), 7.78 - 7.50 (m, 5H), 7.50 - 7.43 (m, 1H), 7.43 - 7.31 (m, 2H), 7.30 - 7.16 (m, 2H), 6.85 (d, J = 7.3 Hz, 1H), 5.16 (t, J = 5.4 Hz, 1H), 4.65 - 4.52 (m, 1H), 4.35 (d, J = 5.4 Hz, 2H), 3.99 - 3.83 (m, 1H), 3.83 - 3.72 (m, 1H), 2.64 - 2.53 (m, 1H), 2.20 - 2.05 (m, 1H).
42		568.1	Method C, RT = 1.598 min, 98.9%	^1H NMR (400MHz, DMSO- d_6) δ = 8.61 (br. s., 1H), 8.22 - 8.07 (m, 2H), 7.88 - 7.73 (m, 2H), 7.58 - 7.49 (m, 1H), 7.45 - 7.25 (m, 3H), 7.23 - 7.15 (m, 1H), 7.09 (d, J = 7.1 Hz, 1H), 4.76 - 4.64 (m, 1H), 4.24 - 4.13 (m, 1H), 3.99 - 3.87 (m, 1H), 3.49 - 3.45 (m, 2H), 3.05 (s, 3H), 2.57 - 2.55 (m, 1H), 2.26 - 2.21 (m, 1H).
43		591.2	Method C, RT = 2.008 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 9.08 - 8.82 (m, 2H), 8.42 (t, J = 8.3 Hz, 1H), 7.71 - 7.61 (m, 1H), 7.52 (d, J = 8.3 Hz, 1H), 7.48 - 7.36 (m, 2H), 7.36 - 7.20 (m, 5H), 7.12 (s, 1H), 4.65 - 4.52 (m, 1H), 3.83 (td, J = 9.2, 6.1 Hz, 1H), 3.66 (t, J = 9.2 Hz, 1H), 2.86 (s, 3H), 2.67 - 2.58 (m, 1H), 2.19 - 2.03 (m, 1H), 2.00 - 1.90 (m, 1H), 1.04 - 0.83 (m, 2H), 0.79 - 0.62 (m, 2H).
44		545.2	Method C, RT = 1.776 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.63 (s, 1H), 8.33 (d, J = 7.8 Hz, 1H), 8.15 (t, J = 8.9 Hz, 1H), 7.51 (s, 1H), 7.49 - 7.30 (m, 6H), 7.25 - 7.15 (m, 2H), 4.98 (m, 1H), 4.61 - 4.51 (m, 1H), 3.89 (td, J = 9.5, 6.6 Hz, 1H), 3.77 (t, J = 8.7 Hz, 1H), 2.58 (dd, J = 13.6, 5.7 Hz, 1H), 2.16 - 2.01 (m, 1H), 1.85 (s, 3H), 1.38 (d, J = 7.1 Hz, 3H).






45		542.2	Method C, RT = 1.778 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.63 (s, 1H), 8.17 (t, J = 8.9 Hz, 1H), 7.98 (dd, J = 12.8, 7.7 Hz, 1H), 7.68 - 7.59 (m, 1H), 7.59 - 7.47 (m, 2H), 7.45 - 7.26 (m, 4H), 7.25 - 7.13 (m, 2H), 4.60 - 4.49 (m, 1H), 3.80 (d, J = 6.1 Hz, 1H), 3.61 - 3.52 (m, 1H), 3.07 - 2.99 (m, 1H), 2.59 - 2.55 (m, 1H), 2.28 - 1.99 (m, 1H), 1.33 (d, J = 13.4 Hz, 6H), 1.23 - 1.10 (m, 6H).
46		577.2	Method C, RT = 1.716 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.64 (d, J = 2.4 Hz, 1H), 8.16 (t, J = 8.9 Hz, 1H), 7.93 (dd, J = 12.3, 8.2 Hz, 1H), 7.61 (t, J = 7.5 Hz, 1H), 7.54 (t, J = 7.5 Hz, 1H), 7.42 (dd, J = 11.2, 2.4 Hz, 1H), 7.39 - 7.32 (m, 1H), 7.32 - 7.24 (m, 2H), 7.24 - 7.06 (m, 3H), 4.63 - 4.50 (m, 1H), 3.83 (td, J = 9.6, 6.7 Hz, 1H), 3.67 (t, J = 8.8 Hz, 1H), 2.65 - 2.56 (m, 1H), 2.16 - 2.05 (m, 1H), 2.00 - 1.90 (m, 1H), 1.34 (d, J = 13.2 Hz, 3H), 1.33 (d, J = 13.2 Hz, 3H), 0.98 - 0.84 (m, 2H), 0.82 - 0.65 (m, 2H).
47		560.2	Method C, RT = 1.760 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.80 (br. s., 1H), 8.62 (s, 1H), 8.25 - 8.05 (m, 2H), 7.68 - 7.51 (m, 2H), 7.51 - 7.35 (m, 2H), 7.31 - 7.10 (m, 3H), 6.93 - 6.91 (m, 1H), 4.59 - 4.50 (m, 1H), 3.77 (d, J = 6.4 Hz, 1H), 3.56 (d, J = 8.3 Hz, 1H), 3.10 (s, 3H), 3.04 - 2.98 (m, 1H), 2.65 - 2.62 (m, 1H), 2.18 - 1.96 (m, 1H), 1.39 - 1.15 (m, 6H).
48		558.2	Method C, RT = 1.861 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 10.5 (br. s., 1H), 8.64 (d, J = 2.7 Hz, 1H), 8.31 - 8.04 (m, 2H), 7.71 (dd, J = 7.6, 1.7 Hz, 1H), 7.47 - 7.35 (m, 2H), 7.35 - 7.27 (m, 1H), 7.26 - 7.14 (m, 3H), 7.10 (br. s., 1H), 4.64 - 4.50 (m, 1H), 3.81 (td, J = 9.6, 6.5 Hz, 1H), 3.65 (t, J = 9.0 Hz, 1H), 3.20 (s, 3H), 2.66 - 2.57 (m, 1H), 2.15 - 2.02 (m, 1H), 2.01 - 1.88 (m, 1H), 1.03 - 0.83 (m, 2H), 0.78 - 0.55 (m, 2H).
49		519.1	Method C, RT = 1.469 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 9.45 (s, 1H), 8.25 (d, J = 2.7 Hz, 1H), 8.00 - 7.87 (m, 2H), 7.82 (dd, J = 9.0, 2.7 Hz, 1H), 7.71 - 7.54 (m, 3H), 7.44 - 7.31 (m, 2H), 7.31 - 7.22 (m, 1H), 4.66 - 4.56 (m, 1H), 4.01 - 3.85 (m, 1H), 3.84 - 3.76 (m, 1H), 2.59 - 2.86 (m, 1H), 2.21 - 2.11 (m, 1H), 1.52 (d, J = 13.2 Hz, 6H).






50		503.2	Method D, RT = 1.498 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.64 (d, J = 1.7 Hz, 1H), 8.20 - 8.08 (m, 1H), 7.62 (s, 1H), 7.53 - 7.28 (m, 6H), 7.25 - 7.16 (m, 2H), 4.62 - 4.49 (m, 1H), 4.17 (q, J = 6.6 Hz, 1H), 3.94 - 3.85 (m, 1H), 3.82 - 3.72 (m, 1H), 3.34 (br. s., 2H), 2.65 - 2.55 (m, 1H), 2.13 - 2.02 (m, 1H), 1.35 (d, J = 6.6 Hz, 3H).
51		568.2	Method C, RT = 1.709 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.99 (s, 1H), 7.99 - 7.86 (m, 1H), 7.73 - 7.57 (m, 2H), 7.57 - 7.44 (m, 2H), 7.42 - 7.32 (m, 2H), 7.32 - 7.14 (m, 3H), 6.81 (d, J = 7.3 Hz, 1H), 4.61 - 4.49 (m, 1H), 3.94 - 3.84 (m, 1H), 3.83 - 3.71 (m, 1H), 2.61 - 2.53 (m, 1H), 2.18 - 2.05 (m, 1H), 1.52 (d, J = 13.2 Hz, 6H).
52		538.1	Method D, RT = 1.774 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.61 (d, J = 2.2 Hz, 1H), 8.23 - 8.04 (m, 2H), 7.87 - 7.63 (m, 2H), 7.49 (dd, J = 7.5, 1.1 Hz, 1H), 7.42 (dd, J = 11.1, 2.3 Hz, 1H), 7.35 (d, J = 9.0 Hz, 2H), 7.24 - 7.11 (m, 2H), 4.63 - 4.50 (m, 1H), 3.83 - 3.75 (m, 1H), 3.75 - 3.67 (m, 1H), 3.05 (s, 3H), 2.63-2.55 (m, 1H), 2.19 - 2.08 (m, 1H).
53		536.1	Method C, RT = 1.594 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.65 - 8.57 (m, 1H), 8.19 - 8.12 (m, 1H), 7.96 - 7.86 (m, 1H), 7.69 - 7.56 (m, 2H), 7.46 - 7.37 (m, 4H), 7.24 - 7.17 (m, 2H), 4.64 - 4.53 (m, 1H), 3.84 - 3.67 (m, 2H), 2.66 - 2.57 (m, 1H), 2.19 - 2.06 (m, 1H), 1.52 (d, J = 13.2 Hz, 6H).
54		516.2	Method C, RT = 1.485 min, 98.5%	^1H NMR (400MHz, DMSO- d_6) δ = 8.49 (s, 1H), 8.14 (dd, J = 7.8, 1.5 Hz, 1H), 7.91 - 7.71 (m, 2H), 7.51 (dd, J = 7.8, 1.5 Hz, 1H), 7.41 (t, J = 7.8 Hz, 1H), 7.38 - 7.22 (m, 3H), 6.92 - 6.74 (m, 2H), 6.56 (d, J = 7.3 Hz, 1H), 4.56 - 4.50 (m, 1H), 3.99 - 3.83 (m, 1H), 3.83 - 3.75 (m, 1H), 3.71 (s, 3H), 3.06 (s, 3H), 2.61 - 2.54 (m, 1H), 2.20 - 2.01 (m, 1H).

55		571.2	Method C, RT = 1.589 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 9.86 (s, 1H), 9.13 (d, J = 6.8 Hz, 1H), 8.49 (s, 1H), 8.22 (dd, J = 10.4, 1.8 Hz, 1H), 8.00 - 7.84 (m, 1H), 7.72 - 7.56 (m, 2H), 7.46 - 7.30 (m, 2H), 7.27 (t, J = 7.8 Hz, 1H), 4.74 - 4.61 (m, 1H), 4.00 - 3.87 (m, 1H), 3.86 - 3.78 (m, 1H), 2.66 - 2.56 (m, 1H), 2.30 - 2.17 (m, 1H), 1.52 (d, J = 13.2 Hz, 6H).
56		537.2	Method C, RT = 1.475 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 9.44 (s, 1H), 8.68 (d, J = 6.8 Hz, 1H), 8.19 (s, 1H), 8.05 (dd, J = 10.0, 2.2 Hz, 1H), 7.98 - 7.88 (m, 1H), 7.67-7.40 (m, 2H), 7.47 - 7.31 (m, 2H), 7.29 - 7.21 (m, 1H), 4.70 - 4.62 (m, 1H), 3.98 - 3.84 (m, 1H), 3.84 - 3.74 (m, 1H), 2.63 - 2.55 (m, 1H), 2.27 - 2.13 (m, 1H), 1.52 (d, J = 13.2 Hz, 6H).
57		570.2	Method C, RT = 1.728min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.93 (d, J = 2.0 Hz, 1H), 8.41 (t, J = 8.6 Hz, 1H), 7.98 - 7.85 (m, 1H), 7.72 - 7.55 (m, 3H), 7.51 (d, J = 8.3 Hz, 1H), 7.42 - 7.30 (m, 3H), 7.30 - 7.22 (m, 1H), 4.65 - 4.54 (m, 1H), 3.96 - 3.85 (m, 1H), 3.79-3.45 (m, 1H), 2.65 - 2.57 (m, 1H), 2.16 - 2.02 (m, 1H), 1.52 (d, J = 13.2 Hz, 6H).
58		537.2	Method C, RT = 1.671 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.75 (br. s., 1H), 8.19 (d, J = 8.1 Hz, 1H), 8.14 (t, J = 8.9 Hz, 1H), 7.78 - 7.67 (m, 2H), 7.46 - 7.23 (m, 5H), 7.23 - 7.16 (m, 1H), 4.61 - 4.50 (m, 1H), 4.27 (br. s., 1H), 3.96 - 3.85 (m, 1H), 3.82 - 3.75 (m, 1H), 2.93 - 2.89 (m, 3H), 2.62 - 2.54 (m, 1H), 2.15 - 2.04 (m, 1H).
59		537.2	Method C, RT = 1.672 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.76 (br. s., 1H), 8.26 - 8.06 (m, 2H), 7.81 - 7.64 (m, 2H), 7.51 - 7.29 (m, 5H), 7.20 (ddd, J = 9.0, 2.4, 1.3 Hz, 1H), 4.56 (dd, J = 17.9, 8.3 Hz, 1H), 4.28 (br. s., 1H), 3.92 - 3.90 (m, 1H), 3.79 (t, J = 8.3 Hz, 1H), 2.93 - 2.89 (m, 3H), 2.64 - 2.53 (m, 1H), 2.16 - 2.01 (m, 1H).






60		576.3	Method C, RT = 2.041 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.91 (d, J = 2.9 Hz, 1H), 8.43 (t, J = 8.6 Hz, 1H), 7.98 (dd, J = 12.7, 7.8 Hz, 1H), 7.71 - 7.60 (m, 2H), 7.60 - 7.47 (m, 3H), 7.37 (dd, J = 7.0, 3.5 Hz, 1H), 7.35 - 7.23 (m, 3H), 4.63 - 4.51 (m, 1H), 3.86 - 3.74 (m, 1H), 3.58 (t, J = 8.7 Hz, 1H), 3.09 - 2.99 (m, 1H), 2.66 - 2.56 (m, 1H), 2.17 - 2.02 (m, 1H), 1.34 (d, J = 13.2 Hz, 6H), 1.18 (d, J = 6.8 Hz, 3H), 1.20 (d, J = 6.8 Hz, 3H).
61		543.2	Method C, RT = 1.774 min, 99.7%	^1H NMR (400MHz, DMSO- d_6) δ = 9.44 (s, 1H), 8.67 (d, J = 5.9 Hz, 1H), 8.19 (d, J = 2.2 Hz, 1H), 8.04 (dd, J = 10.1, 2.2 Hz, 1H), 8.01 - 7.91 (m, 1H), 7.68 - 7.59 (m, 1H), 7.59 - 7.48 (m, 2H), 7.37 (dd, J = 6.4, 4.2 Hz, 1H), 7.31 (d, J = 1.0 Hz, 2H), 4.60 - 4.55 (m, 1H), 3.83 - 3.75 (m, 1H), 3.64 - 3.59 (m, 1H), 3.13 - 3.05 (m, 1H), 2.63 - 2.54 (m, 1H), 2.29 - 2.20 (m, 1H), 1.34 (d, J = 13.2 Hz, 6H), 1.19 (d, J = 6.8 Hz, 3H), 1.20 (d, J = 6.8 Hz, 3H).
62		524.2	Method C, RT = 1.607 min, 99.6%	^1H NMR (400MHz, DMSO- d_6) δ = 9.74 (br. s., 1H), 8.90 (d, J = 1.5 Hz, 1H), 8.41 (d, J = 1.2 Hz, 1H), 7.98 - 7.89 (m, 1H), 7.67 - 7.43 (m, 3H), 7.41 - 7.20 (m, 3H), 7.13 (d, J = 1.7 Hz, 1H), 4.64 - 4.54 (m, 1H), 3.83 (td, J = 9.5, 6.7 Hz, 1H), 3.73 - 3.65 (m, 1H), 2.65 - 2.56 (m, 1H), 2.24 - 2.13 (m, 1H), 2.03 - 1.93 (m, 1H), 1.34 (d, J = 13.2 Hz, 6H), 1.02 - 0.82 (m, 2H), 0.80 - 0.65 (m, 2H).
63		520.2	Method C, RT = 1.517min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 9.83 (br. s., 1H), 8.90 (s, 1H), 8.41 (d, J = 1.5 Hz, 1H), 7.99 - 7.85 (m, 1H), 7.73 - 7.53 (m, 3H), 7.51 - 7.30 (m, 2H), 7.30 - 7.20 (m, 1H), 4.67 - 4.55 (m, 1H), 3.95 - 3.87 (m, 1H), 3.80 - 3.77 (m, 1H), 2.64 - 2.56 (m, 1H), 2.28 - 2.00 (m, 1H), 1.52 (d, J = 13.2 Hz, 6H).
64		553.3	Method C, RT = 1.742min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 9.79 (s, 1H), 8.59 (s, 1H), 8.08 (dd, J = 8.8, 2.2 Hz, 2H), 7.96 - 7.86 (m, 1H), 7.79 (d, J = 8.8 Hz, 1H), 7.72 - 7.55 (m, 2H), 7.45 - 7.31 (m, 2H), 7.31 - 7.23 (m, 1H), 4.70 - 4.59 (m, 1H), 3.98 - 3.84 (m, 1H), 3.84 - 3.74 (m, 1H), 2.65 - 2.55 (m, 1H), 2.28 - 2.08 (m, 1H), 1.53 (d, J = 13.2 Hz, 6H).

65		514.3	Method C, RT = 1.496 min, 97.9%	^1H NMR (400MHz, DMSO- d_6) δ = 8.48 (s, 1H), 7.98 - 7.88 (m, 1H), 7.72 - 7.52 (m, 2H), 7.43 - 7.19 (m, 5H), 6.88 - 6.77 (m, 2H), 6.55 (d, J = 7.3 Hz, 1H), 4.58 - 4.49 (m, 1H), 3.93 - 3.83 (m, 1H), 3.77 - 3.74 (m, 1H), 3.71 (s, 3H), 2.60 - 2.54 (m, 1H), 2.23 - 1.99 (m, 1H), 1.52 (d, J = 13.2 Hz, 6H).
66		571.2	Method C, RT = 1.815 min, 96.2%	^1H NMR (400MHz, DMSO- d_6) δ = 8.41 (t, J = 8.2 Hz, 1H), 8.18 (br. s., 1H), 7.80 - 7.69 (m, 2H), 7.67 (d, J = 10.0 Hz, 1H), 7.52 (d, J = 8.6 Hz, 1H), 7.46 - 7.29 (m, 4H), 7.29 - 7.06 (m, 1H), 4.65 - 4.53 (m, 1H), 3.98 - 3.86 (m, 1H), 3.80 (t, J = 8.9 Hz, 1H), 2.93 - 2.89 (m, 3H), 2.64 - 2.55 (m, 1H), 2.16 - 2.04 (m, 1H).
67		565.2	Method C, RT = 1.804 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 9.45 (s, 1H), 8.69 (d, J = 6.4 Hz, 1H), 8.19 (s, 1H), 8.05 (d, J = 10.3 Hz, 1H), 7.89 (dd, J = 11.5, 7.3 Hz, 1H), 7.64 (quin, J = 7.8 Hz, 2H), 7.48 - 7.26 (m, 2H), 7.17 (t, J = 7.9 Hz, 1H), 4.70 - 4.59 (m, 1H), 3.96 - 3.85 (m, 1H), 3.85 - 3.74 (m, 1H), 2.63 - 2.54 (m, 1H), 2.29 - 2.11 (m, 1H), 1.80 - 1.60 (m, 4H), 0.93 - 0.88 (m, 6H).
68		598.2	Method C, RT = 2.038 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.91 (br. s., 1H), 8.41 (t, J = 8.6 Hz, 1H), 7.97 - 7.83 (m, 1H), 7.75 - 7.56 (m, 3H), 7.51 (d, J = 8.8 Hz, 1H), 7.42 - 7.23 (m, 3H), 7.22 - 7.11 (m, 1H), 4.63 - 4.51 (m, 1H), 3.96 - 3.87 (m, 1H), 3.81 - 3.76 (m, 1H), 2.64 - 2.56 (m, 1H), 2.17 - 2.02 (m, 1H), 1.80 - 1.60 (m, 4H), 0.93 - 0.88 (m, 6H).
69		586.2	Method C, RT = 1.903 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.70 (s, 1H), 8.22 (t, J = 9.2 Hz, 1H), 7.92 (dd, J = 12.5, 7.8 Hz, 1H), 7.73 - 7.55 (m, 2H), 7.49 - 7.30 (m, 3H), 7.30 - 7.23 (m, 1H), 7.23 - 7.09 (m, 2H), 4.63 - 4.52 (m, 1H), 4.00 - 3.84 (m, 1H), 3.80 - 3.75 (m, 1H), 2.64 - 2.56 (m, 1H), 2.18 - 2.01 (m, 1H), 1.52 (d, J = 13.2 Hz, 6H).






70		554.2	Method C, RT = 1.663 min, 99.2%	^1H NMR (400MHz, DMSO- d_6) δ = 8.63 (d, J = 5.4 Hz, 1H), 8.16 (t, J = 8.9 Hz, 1H), 7.79 - 7.68 (m, 1H), 7.52 - 7.36 (m, 2H), 7.35 - 7.24 (m, 1H), 7.24 - 7.08 (m, 4H), 4.61 - 4.50 (m, 1H), 3.94 - 3.82 (m, 1H), 3.79 - 3.67 (m, 1H), 2.59 (dd, J = 12.7, 7.6 Hz, 1H), 2.13 - 2.01 (m, 1H), 1.86 - 1.59 (m, 6H)
71		532.3	Method D, RT = 1.383 min, 98.9%	^1H NMR (400MHz, DMSO- d_6) δ = 8.24 (s, 1H), 7.92 (dd, J = 12.2, 8.1 Hz, 1H), 7.84 (t, J = 9.3 Hz, 1H), 7.74 - 7.55 (m, 2H), 7.37 (d, J = 6.8 Hz, 2H), 7.30 - 7.22 (m, 1H), 6.97 (d, J = 7.1 Hz, 1H), 6.87 (dd, J = 12.2, 2.6 Hz, 1H), 6.72 (dd, J = 8.9, 1.8 Hz, 1H), 4.61 - 4.48 (m, 1H), 3.95 - 3.84 (m, 1H), 3.82 - 3.59 (m, 1H), 3.74 (s, 3H), 2.62 - 2.55 (m, 1H), 2.15 - 2.02 (m, 1H), 1.52 (d, J = 13.2 Hz, 6H).
72		588.2	Method C, RT = 1.794 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.95 (dd, J = 8.2, 2.5 Hz, 1H), 8.41 (t, J = 8.2 Hz, 1H), 7.76 - 7.65 (m, 2H), 7.53 - 7.43 (m, 2H), 7.38 - 7.28 (m, 2H), 7.23 - 7.16 (m, 2H), 4.62 - 4.54 (m, 1H), 3.94 - 3.83 (m, 1H), 3.81 - 3.70 (m, 1H), 2.64 - 2.53 (m, 1H), 2.16 - 2.02 (m, 1H), 1.76 - 1.64 (m, 6H).
73		588.2 (M+N H_4) $^+$	Method D, RT = 1.647 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.94 (d, J = 2.9 Hz, 1H), 8.84 (d, J = 4.2 Hz, 1H), 8.42 (t, J = 8.3 Hz, 1H), 7.87-7.86 (m, 1H), 7.73 - 7.62 (m, 2H), 7.52 (d, J = 8.6 Hz, 1H), 7.43 - 7.25 (m, 3H), 4.65 - 4.52 (m, 1H), 3.98 - 3.86 (m, 1H), 3.82 - 3.70 (m, 1H), 2.64 - 2.56 (m, 1H), 2.16 - 2.04 (m, 1H), 1.62 (d, J = 13.2 Hz, 6H).
74		554.2 (M+N H_4) $^+$	Method D, RT = 1.498 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.84 (br d, J = 4.2 Hz, 1H), 8.64 (s, 1H), 8.15 (t, J = 8.9 Hz, 1H), 7.86 (dd, J = 3.5, 7.5 Hz, 1H), 7.70 - 7.61 (m, 1H), 7.42 (dd, J = 2.4, 11.0 Hz, 1H), 7.37 (br d, J = 8.1 Hz, 1H), 7.32 - 7.24 (m, 1H), 7.20 (br d, J = 7.6 Hz, 2H), 4.62 - 4.50 (m, 1H), 3.95 - 3.84 (m, 1H), 3.82 - 3.71 (m, 1H), 2.64 - 2.56 (m, 1H), 2.13 - 2.00 (m, 1H), 1.62 (d, J = 13.4 Hz, 6H).






75		564.2	Method C, RT = 1.723 min, 99.9%	$^1\text{H NMR}$ (400MHz, DMSO- d_6) δ = 8.62 (d, J = 2.4 Hz, 1H), 8.16 (t, J = 8.9 Hz, 1H), 7.87 (ddd, J = 11.6, 7.5, 1.7 Hz, 1H), 7.70 - 7.53 (m, 2H), 7.43 (dd, J = 11.0, 2.4 Hz, 1H), 7.36 (ddd, J = 7.5, 3.6, 1.6 Hz, 1H), 7.33 - 7.26 (m, 2H), 7.23 - 7.16 (m, 2H), 4.64 - 4.50 (m, 1H), 3.84 - 3.74 (m, 1H), 3.73 - 3.63 (m, 1H), 2.66 - 2.57 (m, 1H), 2.19 - 2.01 (m, 1H), 1.71 (m, 4H), 0.93 - 0.88 (m, 6H).
76		570.2	Method C, RT = 1.867 min, 98.9%	$^1\text{H NMR}$ (400MHz, DMSO- d_6) δ = 8.91 (d, J = 2.9 Hz, 1H), 8.41 (m, 1H), 7.91 (ddd, J = 13.0, 7.6, 1.2 Hz, 1H), 7.73 - 7.56 (m, 3H), 7.52 (d, J = 9.5 Hz, 1H), 7.46 - 7.38 (m, 3H), 7.35 (d, J = 7.1 Hz, 1H), 4.68 - 4.55 (m, 1H), 3.85 - 3.76 (m, 1H), 3.74 - 3.66 (m, 1H), 2.67 - 2.59 (m, 1H), 2.21 - 2.10 (m, 1H), 1.49 (d, J = 13.5 Hz, 6H).
77		586.2	Method C, RT = 1.781 min, 99.8%	$^1\text{H NMR}$ (400MHz, DMSO- d_6) δ = 8.95 (t, J = 3.2 Hz, 1H), 8.42 (t, J = 8.2 Hz, 1H), 8.01 - 7.90 (m, 1H), 7.70 - 7.58 (m, 3H), 7.55 - 7.47 (m, 2H), 7.39 - 7.21 (m, 3H), 4.63 - 4.53 (m, 1H), 3.97 - 3.85 (m, 1H), 3.82 - 3.70 (m, 1H), 2.64 - 2.55 (m, 1H), 2.17 - 2.03 (m, 1H), 1.48 (d, J = 12.8 Hz, 3H), 1.46 (d, J = 12.8 Hz, 3H).
78		568.1	Method C, RT = 1.691 min, 99.8%	$^1\text{H NMR}$ (400MHz, DMSO- d_6) δ = 8.65 (dd, J = 6.1, 2.4 Hz, 1H), 8.17 (t, J = 8.9 Hz, 1H), 7.98 - 7.88 (m, 1H), 7.70 - 7.57 (m, 2H), 7.54 - 7.39 (m, 3H), 7.29 (td, J = 3.5, 7.1 Hz, 1H), 7.24 - 7.13 (m, 2H), 4.63 - 4.49 (m, 1H), 3.87 - 3.76 (m, 1H), 3.73 - 3.64 (m, 1H), 2.66 - 2.57 (m, 1H), 2.16 - 2.01 (m, 1H), 1.52 - 1.44 (m, 6H).
79		500.1	Method C, RT = 1.763 min, 100%	$^1\text{H NMR}$ (400MHz, DMSO- d_6) δ = 8.73 (s, 1H), 8.63 (d, J = 2.2 Hz, 1H), 8.33 (s, 1H), 8.15 (t, J = 8.8 Hz, 1H), 7.90 (s, 1H), 7.70 (t, J = 7.2 Hz, 1H), 7.48 - 7.33 (m, 2H), 7.29 - 7.11 (m, 2H), 4.55 (dd, J = 18.1, 8.6 Hz, 1H), 3.92 - 3.84 (m, 1H), 3.81 - 3.72 (m, 1H), 2.61 (m, 1H), 2.12 - 2.00 (m, 1H).






80		602.1	Method C, RT = 1.821 min, 99.5%	^1H NMR (400MHz, DMSO- d_6) δ = 8.94 (dd, J = 5.9, 2.9 Hz, 1H), 8.42 (m, 1H), 7.99 - 7.89 (m, 1H), 7.72 - 7.55 (m, 3H), 7.54 - 7.44 (m, 3H), 7.35 (dd, J = 7.1, 2.9 Hz, 1H), 7.32 - 7.24 (m, 1H), 4.65 - 4.53 (m, 1H), 3.87 - 3.78 (m, 1H), 3.74 - 3.67 (m, 1H), 2.63 (dt, J = 6.4, 5.1 Hz, 1H), 2.17 - 2.03 (m, 1H), 1.50 - 1.45 (m, 6H).
81		552.2	Method D, RT = 2.358 min, 99.7%	^1H NMR (400MHz, DMSO- d_6) δ = 8.64 (br. s., 1H), 8.15 (t, J = 8.8 Hz, 1H), 7.94 (dd, J = 12.5, 1.5 Hz, 1H), 7.69 - 7.58 (m, 2H), 7.55 - 7.48 (m, 1H), 7.42 (dd, J = 11.3, 2.3 Hz, 1H), 7.32 (d, J = 7.8 Hz, 1H), 7.30 - 7.26 (m, 1H), 7.22 - 7.16 (m, 2H), 4.56 (s, 1H), 3.96 - 3.83 (m, 1H), 3.83 - 3.70 (m, 1H), 2.61 - 2.57 (m, 1H), 2.13 - 2.01 (m, 1H), 1.48 (d, J = 12.8 Hz, 3H), 1.46 (d, J = 12.8 Hz, 3H).
82		508.1	Method C, RT = 1.834 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.63 (d, J = 1.0 Hz, 1H), 8.15 (t, J = 8.8 Hz, 1H), 7.63 - 7.50 (m, 2H), 7.50 - 7.29 (m, 5H), 7.24 - 7.15 (m, 1H), 7.11 (d, J = 7.3 Hz, 1H), 5.10 (t, J = 4.9 Hz, 1H), 4.81 - 4.62 (m, 1H), 4.29 - 4.12 (m, 1H), 3.49 - 3.47 (m, 2H), 2.55 - 2.52 (m, 1H), 2.31 - 2.21 (m, 1H).
83		586.2	Method C, RT = 2.031 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.66 (d, J = 2.5 Hz, 1H), 8.15 (t, J = 8.8 Hz, 1H), 7.89 - 7.80 (m, 1H), 7.75 (t, J = 7.8 Hz, 1H), 7.63 - 7.53 (m, 2H), 7.43 (dd, J = 11.3, 2.3 Hz, 1H), 7.31 - 7.13 (m, 4H), 4.62 - 4.50 (m, 1H), 3.97 - 3.94 (m, 1H), 3.81 - 3.73 (m, 1H), 2.63 - 2.55 (m, 1H), 2.17 - 2.05 (m, 1H), 1.54 (d, J = 13.6 Hz, 3H), 1.52 (d, J = 13.6 Hz, 3H).
84		555.2 (M+N H ₄) ⁺	Method C, RT = 1.369 min, 92.6%	^1H NMR (400MHz, DMSO- d_6) δ = 9.47 (s, 1H), 8.97 (d, J = 5.6 Hz, 1H), 8.66 (d, J = 2.4 Hz, 1H), 8.15 (t, J = 8.9 Hz, 1H), 7.48 - 7.34 (m, 3H), 7.25 - 7.17 (m, 2H), 4.62 - 4.54 (m, 1H), 3.96 - 3.85 (m, 1H), 3.78 (t, J = 8.6 Hz, 1H), 2.63 - 2.54 (m, 1H), 2.15 - 2.04 (m, 1H), 1.69 (d, J = 13.7 Hz, 6H).






85		570.2	Method C, RT = 1.722 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.68 - 8.58 (m, 1H), 8.15 (t, J = 8.8 Hz, 1H), 7.76 - 7.65 (m, 1H), 7.52 - 7.35 (m, 3H), 7.25 - 7.15 (m, 3H), 7.13 - 7.06 (m, 1H), 4.59 - 4.48 (m, 1H), 3.91 - 3.81 (m, 1H), 3.78 - 3.67 (m, 1H), 2.55 - 2.52 (m, 1H), 2.13 - 2.01 (m, 1H), 1.73 - 1.61 (m, 6H).
86		604.1	Method C, RT = 1.834 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.98 - 8.91 (m, 1H), 8.45 - 8.38 (m, 1H), 7.76 - 7.62 (m, 2H), 7.52 (br. d., J = 9.3 Hz, 1H), 7.48 - 7.40 (m, 2H), 7.36 - 7.30 (m, 1H), 7.24 - 7.18 (m, 1H), 7.11 (d, J = 6.6 Hz, 1H), 4.62 - 4.52 (m, 1H), 3.92 - 3.83 (m, 1H), 3.78 - 3.70 (m, 1H), 2.64 - 2.56 (m, 1H), 2.14 - 2.06 (m, 1H), 1.72 - 1.64 (m, 6H).
87		602.2	Method C, RT = 1.795 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.65 (dd, J = 1.7, 8.8 Hz, 1H), 8.15 (t, J = 8.9 Hz, 1H), 7.88 (dd, J = 7.3, 12.7 Hz, 1H), 7.70 - 7.53 (m, 3H), 7.43 (dd, J = 2.4, 11.0 Hz, 1H), 7.38 - 7.30 (m, 2H), 7.24 - 7.10 (m, 2H), 4.61 - 4.52 (m, 1H), 3.96 - 3.85 (m, 1H), 3.81 - 3.71 (m, 1H), 2.61 - 2.54 (m, 1H), 2.17 - 2.03 (m, 1H), 1.64 - 1.48 (m, 6H).
88		636.2	Method C, RT = 1.919 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.94 (dd, J = 2.9, 8.6 Hz, 1H), 8.41 (m, 1H), 7.92 - 7.82 (m, 1H), 7.71 - 7.54 (m, 4H), 7.51 (d, J = 8.8 Hz, 1H), 7.39 - 7.31 (m, 3H), 4.63 - 4.53 (m, 1H), 3.96 - 3.87 (m, 1H), 3.81 - 3.73 (m, 1H), 2.63 - 2.57 (m, 1H), 2.18 - 2.06 (m, 1H), 1.61 - 1.50 (m, 6H).
89		592.2	Method C, RT = 1.872 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.96 - 8.89 (m, 1H), 8.41 (t, J = 8.3 Hz, 1H), 8.03 - 7.93 (m, 1H), 7.71 - 7.49 (m, 4H), 7.39 - 7.26 (m, 3H), 7.14 - 7.07 (m, 1H), 4.58 - 4.49 (m, 1H), 3.87 - 3.78 (m, 1H), 3.74 - 3.67 (m, 1H), 2.63 - 2.54 (m, 1H), 2.13 - 2.02 (m, 1H), 1.60 - 1.50 (m, 1H), 1.45 (d, J = 13.4 Hz, 3H), 1.42 (d, J = 13.4 Hz, 3H), 0.79 - 0.72 (m, 1H), 0.71 - 0.61 (m, 3H).

90		614.2	Method C, RT = 1.768 min, 99.7%	^1H NMR (400MHz, DMSO- d_6) δ = 8.94 (d, J = 2.0 Hz, 1H), 8.40 (t, J = 8.3 Hz, 1H), 8.01 - 7.89 (m, 1H), 7.72 - 7.58 (m, 3H), 7.51 (m, 1H), 7.40 (m, 1H), 7.34 - 7.18 (m, 3H), 4.71 - 4.58 (m, 1H), 4.33 (m, 1H), 3.48 (dd, J = 10.5, 3.9 Hz, 1H), 3.40 - 3.36 (m, 1H), 3.24 (s, 3H), 2.49 - 2.44 (m, 1H), 2.37 - 2.23 (m, 1H), 1.62 - 1.36 (m, 6H).
91		586.2	Method C, RT = 1.778 min, 99.8%	^1H NMR (400MHz, DMSO- d_6) δ = 8.71 - 8.59 (m, 1H), 8.22 - 8.12 (m, 1H), 7.93 - 7.90 (m, 1H), 7.83 - 7.75 (m, 1H), 7.70 - 7.58 (m, 2H), 7.48 - 7.33 (m, 3H), 7.24 - 7.14 (m, 2H), 4.62 (d, J = 1.2 Hz, 1H), 3.96 - 3.84 (m, 1H), 3.81 - 3.71 (m, 1H), 2.65 - 2.58 (m, 1H), 2.10 - 2.04 (m, 1H), 1.67 - 1.33 (m, 6H).
92		620.2	Method C, RT = 1.911 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 9.00 - 8.89 (m, 1H), 8.47 - 8.37 (m, 1H), 7.95 - 7.86 (m, 1H), 7.83 - 7.74 (m, 1H), 7.71 - 7.60 (m, 3H), 7.55 - 7.48 (m, 1H), 7.45 - 7.36 (m, 2H), 7.35 - 7.32 (m, 1H), 4.75 - 4.43 (m, 1H), 3.96 - 3.84 (m, 1H), 3.81 - 3.71 (m, 1H), 2.66 - 2.58 (m, 1H), 2.10 - 2.04 (m, 1H), 1.60 - 1.48 (m, 6H).
93		589.2 (M+N H ₄) ⁺	Method C, RT = 1.532 min, 94.1%	^1H NMR (400MHz, DMSO- d_6) δ = 9.47 (s, 1H), 9.03 - 8.85 (m, 2H), 8.42 (t, J = 8.2 Hz, 1H), 7.67 (br. d., J = 11.5 Hz, 1H), 7.53 (s, 1H), 7.48 - 7.30 (m, 3H), 4.60 (td, J = 10.2, 8.0 Hz, 1H), 4.01 - 3.87 (m, 1H), 3.79 (br. t., J = 8.8 Hz, 1H), 2.65 - 2.55 (m, 1H), 2.18 - 2.04 (m, 1H), 1.69 (d, J = 13.7 Hz, 6H).
94		620.2	Method C, RT = 1.932 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.95 (d, J = 3.2 Hz, 1H), 8.41 (t, J = 8.4 Hz, 1H), 7.90 - 7.80 (m, 1H), 7.76 (t, J = 7.9 Hz, 1H), 7.67 (dd, J = 1.8, 11.6 Hz, 1H), 7.63 - 7.55 (m, 2H), 7.52 (d, J = 9.0 Hz, 1H), 7.34 (d, J = 7.1 Hz, 1H), 7.29 - 7.19 (m, 2H), 4.64 - 4.55 (m, 1H), 3.96 - 3.87 (m, 1H), 3.83 - 3.74 (m, 1H), 2.99 - 2.87 (m, 1H), 2.64 - 2.56 (m, 1H), 2.16 - 2.06 (m, 1H), 1.60 - 1.48 (m, 6H).






95		568.2	Method C, RT = 1.608 min, 99%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.65 (br. s., 1H), 8.16 (t, <i>J</i> = 8.8 Hz, 1H), 7.98 - 7.87 (m, 1H), 7.74 - 7.57 (m, 3H), 7.43 (dd, <i>J</i> = 11.2, 2.4 Hz, 1H), 7.32 - 7.10 (m, 4H), 6.59 (t, <i>J</i> = 52.4 Hz, 1H), 4.60 - 4.50 (m, 1H), 3.95 - 3.85 (m, 1H), 3.81 - 3.71 (m, 1H), 2.62 - 2.55 (m, 1H), 2.17 - 2.03 (m, 1H), 1.50 (d, <i>J</i> = 13.2 Hz, 3H), 1.44 (d, <i>J</i> = 13.2 Hz, 3H).
96		532.2	Method C, RT = 1.813 min, 100%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.64 (d, <i>J</i> = 1.5 Hz, 1H), 8.16 (m, 1H), 8.04 - 7.92 (m, 1H), 7.69 - 7.54 (m, 2H), 7.43 (dd, <i>J</i> = 11.1, 2.3 Hz, 1H), 7.36 - 7.26 (m, 2H), 7.24 - 7.13 (m, 3H), 4.59 - 4.45 (m, 1H), 3.89 - 3.78 (m, 1H), 3.69 - 3.55 (m, 1H), 2.62 - 2.54 (m, 1H), 2.12 (m, 3H), 2.10 - 1.99 (m, 1H), 1.45 (br. d., <i>J</i> = 13.4 Hz, 3H), 1.36 (br. d., <i>J</i> = 13.4 Hz, 3H).
97		585.2	Method C, RT = 1.809 min, 100%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.95 (d, <i>J</i> = 2.7 Hz, 1H), 8.42 (t, <i>J</i> = 8.4 Hz, 1H), 8.02 - 7.84 (m, 1H), 7.81 - 7.74 (m, 1H), 7.70 - 7.62 (m, 2H), 7.57 (d, <i>J</i> = 7.6 Hz, 1H), 7.54 - 7.48 (m, 1H), 7.46 (dd, <i>J</i> = 6.8, 1.7 Hz, 1H), 7.42 (dd, <i>J</i> = 7.2, 1.7 Hz, 1H), 7.40 - 7.32 (m, 1H), 4.68 - 4.56 (m, 1H), 4.45 (s, 2H), 4.00 - 3.88 (m, 1H), 3.85 - 3.77 (m, 1H), 2.66 - 2.56 (m, 1H), 2.18 - 2.05 (m, 1H).
98		586.2	Method C, RT = 1.764 min, 99.9%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.93 (br. s., 1H), 8.42 (m, 1H), 7.94 - 7.90 (m, 1H), 7.71 - 7.57 (m, 3H), 7.56 - 7.49 (m, 1H), 7.49 - 7.41 (m, 1H), 7.41 - 7.26 (m, 3H), 4.67 - 4.53 (m, 1H), 3.88 - 3.79 (m, 1H), 3.76 - 3.66 (m, 1H), 2.67 - 2.56 (m, 1H), 2.20 - 2.05 (m, 1H), 1.57 - 1.48 (m, 6H).
99		552.2	Method C, RT = 1.595 min, 99.7%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.64 (br. s., 1H), 8.17 (t, <i>J</i> = 8.9 Hz, 1H), 7.95 - 7.90 (m, 1H), 7.71 - 7.57 (m, 2H), 7.50 - 7.30 (m, 4H), 7.25 - 7.09 (m, 2H), 4.65 - 4.49 (m, 1H), 3.82 (dt, <i>J</i> = 9.4, 6.4 Hz, 1H), 3.74 - 3.61 (m, 1H), 2.66 - 2.56 (m, 1H), 2.15 - 2.03 (m, 1H), 1.57 - 1.48 (m, 6H).






100		599.2	Method C, RT = 1.857 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.93 (d, J = 2.7 Hz, 1H), 8.85 (dd, J = 4.6, 1.2 Hz, 1H), 8.42 (m, 1H), 7.86 - 8.82 (m, 1H), 7.70 - 7.60 (m, 2H), 7.55 - 7.47 (m, 1H), 7.42 - 7.29 (m, 2H), 7.24 - 7.16 (m, 1H), 4.64 - 4.52 (m, 1H), 3.96 - 3.87 (m, 1H), 3.81 - 3.73 (m, 1H), 2.65 - 2.58 (m, 1H), 2.16 - 1.81 (m, 5H), 1.02 - 0.85 (m, 6H).
101		583.1 (M+N H_4) ⁺	Method C, RT = 1.746 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.69 - 8.56 (m, 1H), 8.15 (t, J = 8.9 Hz, 1H), 8.04 - 7.91 (m, 1H), 7.76 - 7.59 (m, 2H), 7.49 - 7.32 (m, 3H), 7.29 - 7.10 (m, 3H), 4.62 - 4.51 (m, 1H), 3.97 - 3.61 (m, 4H), 2.57 - 2.52 (m, 1H), 2.14 - 2.01 (m, 1H), 1.44 (d, J = 14.4 Hz, 3H), 1.12 (t, J = 7.1 Hz, 3H).
102		598.2	Method C, RT = 1.965 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.96 (d, J = 2.7 Hz, 1H), 8.46 (t, J = 8.3 Hz, 1H), 7.97 - 7.86 (m, 1H), 7.72 - 7.57 (m, 3H), 7.53 (d, J = 8.8 Hz, 1H), 7.42 - 7.30 (m, 2H), 7.30 - 7.22 (m, 2H), 4.65 (d, J = 9.0 Hz, 1H), 3.88 (br. d., J = 9.3 Hz, 1H), 3.45 (br. d., J = 9.3 Hz, 1H), 1.53 (d, J = 13.2 Hz, 6H), 1.22 (s, 3H), 1.06 (s, 3H).
103		581.2 (M+N H_4) ⁺	Method C, RT = 1.965 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.68 (d, J = 1.5 Hz, 1H), 8.22 (t, J = 9.0 Hz, 1H), 7.97 - 7.86 (m, 1H), 7.70 - 7.56 (m, 2H), 7.45 (dd, J = 11.2, 2.4 Hz, 1H), 7.41 - 7.29 (m, 2H), 7.29 - 7.17 (m, 2H), 7.12 (d, J = 9.0 Hz, 1H), 4.62 (d, J = 9.0 Hz, 1H), 3.86 (br. d., J = 9.3 Hz, 1H), 3.45 (br. d., J = 9.3 Hz, 1H), 1.54 (d, J = 13.2 Hz, 6H), 1.21 (s, 3H), 1.05 (s, 3H).
104		600.2	Method C, RT = 1.871 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.98 - 8.86 (m, 1H), 8.41 (t, J = 8.68 Hz, 1H), 8.02 - 7.92 (m, 1H), 7.77 - 7.60 (m, 3H), 7.56 - 7.48 (m, 1H), 7.46 - 7.30 (m, 3H), 7.28 - 7.18 (m, 1H), 4.65 - 4.51 (m, 1H), 3.98 - 3.81 (m, 2H), 3.81 - 3.73 (m, 1H), 3.73 - 3.61 (m, 1H), 2.65 - 2.54 (m, 1H), 2.17 - 2.02 (m, 1H), 1.44 (d, J = 14.67 Hz, 3H), 1.12 (t, J = 7.1 Hz, 3H).






105		600.2	Method C, RT = 1.871 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 9.02 - 8.90 (m, 1H), 8.41 (t, J = 8.3 Hz, 1H), 8.05 - 7.92 (m, 1H), 7.76 - 7.60 (m, 3H), 7.51 (dd, J = 8.7, 1.1 Hz, 1H), 7.46 - 7.31 (m, 3H), 7.28 - 7.15 (m, 1H), 4.65 - 4.53 (m, 1H), 3.97 - 3.74 (m, 3H), 3.74 - 3.60 (m, 1H), 2.65 - 2.56 (m, 1H), 2.18 - 2.02 (m, 1H), 1.45 (d, J = 14.43 Hz, 3H), 1.12 (t, J = 7.1 Hz, 3H).
106		580.1	Method C, RT = 1.763 min, 94.7%	^1H NMR (400MHz, DMSO- d_6) δ = 8.65 (d, J = 2.5 Hz, 1 H) 8.15 (t, J = 8.9 Hz, 1 H), 7.96 - 7.88 (m, 1 H), 7.67 - 7.64 (m, 2 H), 7.51 (d, J = 8.6 Hz, 1 H), 7.37 - 7.47 (m, 2 H), 7.27 (d, J = 8.8 Hz, 1 H), 7.16 - 7.23 (m, 2 H), 4.47 - 4.64 (m, 1 H), 3.90 - 4.01 (m, 1H), 3.82 - 3.90 (m, 1 H), 2.54 - 2.62 (m, 1 H), 2.01 - 2.15 (m, 1 H), 1.53 (d, J = 13.2 Hz, 6 H).
107		568.1	Method C, RT = 1.662 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.74 (d, J = 1.71 Hz, 1 H), 8.11 (t, J = 8.93 Hz, 1 H), 7.89 - 8.01 (m, 1 H), 7.57 - 7.71 (m, 3 H), 7.40 - 7.40 (m, 1 H), 7.34 - 7.48 (m, 1 H), 7.32 (d, J = 8.31 Hz, 1 H), 7.25 - 7.14 (m, 2 H), 7.04 (t, J = 52.4 Hz, 1H) 4.41 - 4.55 (m, 1 H), 3.89 - 4.01 (m, 1 H), 3.64 - 3.71 (m, 1 H), 2.56 - 2.52 (m, 1 H), 2.18 - 2.31 (m, 1 H), 1.30 - 1.63 (m, 6 H).
108		602.1	Method C, RT = 1.892 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.93 (d, J = 2.5 Hz, 1 H), 8.41 (t, J = 8.4 Hz, 1 H), 8.0 - 7.87 (m, 1 H), 7.77 - 7.71 (m, 1 H), 7.69 - 7.59 (m, 2 H), 7.55 - 7.49 (m, 1 H), 7.48 - 7.42 (m, 1 H), 7.41 - 7.27 (m, 2 H), 7.23 - 7.13 (m, 1 H), 4.65 - 4.54 (m, 1 H), 3.97 - 3.86 (m, 1 H), 3.82 - 3.71 (m, 1 H), 3.51 (s, 3H), 3.49 (s, 3H), 2.65 - 2.56 (m, 1 H), 2.18 - 2.02 (m, 1 H).
109		550.2	Method C, RT = 1.650 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.63 (d, J = 2.0 Hz, 1 H), 8.14 (t, J = 8.9 Hz, 1 H), 7.99 - 7.84 (m, 1 H), 7.72 - 7.55 (m, 2 H), 7.37 - 7.50 (m, 2 H), 7.35 - 7.25 (m, 2 H), 7.23 - 7.17 (m, 1 H), 7.14 (d, J = 7.1 Hz, 1 H), 4.70 - 4.58 (m, 1 H), 4.36 - 4.23 (m, 1 H), 2.36 - 2.25 (m, 2 H), 1.46 (d, J = 12.4 Hz, 6H), 1.19 (d, J = 6.4 Hz, 3 H).

110		599.2	Method C, RT = 1.851 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.96 (d, J = 3.18 Hz, 1 H), 8.84 (m, 1 H), 8.46 (t, J = 8.2 Hz, 1 H), 7.87 (m, 1H), 7.63 - 7.72 (m, 2 H), 7.52 (d, J = 9.5 Hz, 1 H), 7.19 - 7.39 (m, 3 H), 4.65 (d, J = 9.1 Hz, 1 H), 3.88 (d, J = 9.2 Hz, 1 H), 3.45 (d, J = 9.2 Hz, 1 H), 1.63 (d, J = 13.2 Hz, 6H), 1.23 (s, 3 H), 1.07 (s, 3 H).
111		566.2	Method C, RT = 1.404 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.63 (s, 1H), 8.17 (t, J = 8.8 Hz, 1H), 8.00 - 7.88 (m, 1H), 7.70 - 7.58 (m, 2H), 7.49 - 7.36 (m, 2H), 7.33 - 7.13 (m, 4H), 4.93 (t, J = 5.0 Hz, 1H), 4.68 - 4.56 (m, 1H), 4.25 - 4.12 (m, 1H), 3.44 (br. t., J = 4.0 Hz, 2H), 2.67 - 2.61 (m, 1H), 1.98 - 1.85 (m, 1H), 1.64 - 1.42 (m, 6H).
112		566.2	Method C, RT = 1.425 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.61 (s, 1H), 8.14 (t, J = 8.9 Hz, 1H), 7.98 - 7.87 (m, 1H), 7.71 - 7.56 (m, 2H), 7.46 - 7.36 (m, 2H), 7.36 - 7.24 (m, 2H), 7.22 - 7.15 (m, 1H), 7.09 (d, J = 7.6 Hz, 1H), 5.15 - 5.05 (m, 1H), 4.76 - 4.63 (m, 1H), 4.24 - 4.11 (m, 1H), 3.47 (br. d., J = 3.9 Hz, 2H), 2.53 (br. s., 1H), 2.29 - 2.20 (m, 1H), 1.60 - 1.41 (m, 6H).
113		565.2	Method C, RT = 1.719 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.84 (d, J = 4.9 Hz, 1 H), 8.68 (d, J = 2.5 Hz, 1 H), 8.22 (t, J = 8.9 Hz, 1 H), 7.87 (ddd, J = 7.8, 4.4, 1.5 Hz, 1 H), 7.67 (ddd, J = 7.8, 4.9, 2.5 Hz, 1 H), 7.44 (dd, J = 11.3, 2.5 Hz, 1 H), 7.26 - 7.38 (m, 2 H), 7.21 (ddd, J = 8.9, 2.5, 1.1 Hz, 1 H), 7.00 - 7.16 (m, 1 H), 4.62 (d, J = 9.1 Hz, 1 H), 3.86 (d, J = 9.1 Hz, 1 H), 3.44 (d, J = 9.1 Hz, 1 H), 1.63 (d, J = 13.7 Hz, 6 H), 1.19 - 1.28 (m, 3 H), 1.05 (s, 3 H).
114		600.2	Method D, RT = 1.577 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 9.06 - 8.51 (br. s., 1H), 8.45 - 8.36 (m, 1 H), 8.00 - 7.87 (m, 1 H), 7.71 - 7.57 (m, 3 H), 7.51 (dd, J = 9.17, 1.10 Hz, 1 H), 7.44 - 7.11 (m, 4 H), 4.78 - 4.61 (m, 1 H), 4.26 - 4.21 (m, 1 H), 3.50 - 3.40 (m, 3 H), 2.61 - 2.53 (m, 1 H), 2.33 - 2.22 (m, 1 H), 1.44 - 1.55 (m, 6 H).

115		584.2	Method D, RT = 1.805 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.97 - 8.85 (m, 1 H), 8.46 - 8.34 (m, 1 H), 7.99 - 7.89 (m, 1 H), 7.70 - 7.57 (m, 3 H), 7.52 (br. d., J = 8.6 Hz, 1 H), 7.46 - 7.17 (m, 4 H), 4.71 - 4.57 (m, 1 H), 4.34 - 4.15 (m, 1 H), 2.37 - 2.30 (m, 1 H), 1.75 - 1.62 (m, 1 H), 1.59 - 1.39 (m, 6 H), 1.19 (d, J = 6.4 Hz, 3 H).
116		584.2	Method D, RT = 1.802 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.92 (d, J = 2.7 Hz, 1 H), 8.40 (t, J = 8.3 Hz, 1 H), 7.97 - 7.89 (m, 1 H), 7.71 - 7.58 (m, 3 H), 7.51 (d, J = 8.8 Hz, 1 H), 7.40 (m, 1 H), 7.36 - 7.25 (m, 3 H), 4.66 (td, J = 9.2, 7.3 Hz, 1 H), 4.35 - 4.24 (m, 1 H), 2.38 - 2.24 (m, 2 H), 1.63 - 1.41 (m, 6 H), 1.20 (d, J = 6.4 Hz, 3 H).
117		558.2	Method C, RT = 1.711 min, 99.4%	^1H NMR (400MHz, DMSO- d_6) δ = 8.65 (br. s., 1 H), 8.18 (t, J = 8.9 Hz, 1 H), 7.96 (m, 1 H), 7.69 - 7.53 (m, 2 H), 7.43 (dd, J = 12.5, 2.5 Hz, 1 H), 7.37 - 7.26 (m, 2 H), 7.25 - 7.10 (m, 3 H), 4.57 (dt, J = 10.3, 8.1 Hz, 1 H), 3.95 - 3.65 (m, 2 H), 2.66 - 2.55 (m, 1 H), 2.14 - 2.03 (m, 1H), 1.80 - 1.69 (m, 1 H), 1.51 - 1.30 (m, 6 H), 0.98 - 0.83 (m, 2 H), 0.67 - 0.43 (m, 2 H).
118		491	Method C, RT = 1.921 min, 99.3%	^1H NMR (400MHz, DMSO- d_6) δ = 8.63 (d, J = 2.1 Hz, 1 H), 8.29 (dd, J = 5.0, 2.1 Hz, 1 H), 8.15 (t, J = 8.8 Hz, 1 H), 7.78 (dd, J = 7.3, 2.1 Hz, 1 H), 7.47 - 7.36 (m, 2 H), 7.35 - 7.28 (m, 1 H), 7.25 - 7.05 (m, 3 H), 4.62 - 4.48 (m, 1 H), 3.95 - 3.84 (m, 1 H), 3.89 (s, 3H), 3.83 - 3.71 (m, 1 H), 2.63 - 2.52 (m, 1 H), 2.14 - 2.03 (m, 1H).
119		545.1	Method C, RT = 1.737 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.64 (d, J = 2.0 Hz, 1H), 8.19 - 8.05 (m, 1H), 7.63 - 7.52 (m, 2H), 7.50 - 7.35 (m, 4H), 7.27 - 7.13 (m, 3H), 4.55 (td, J = 10.5, 8.0 Hz, 1H), 4.30 (t, J = 7.9 Hz, 2H), 3.96 - 3.81 (m, 3H), 3.81 - 3.72 (m, 1H), 2.61 - 2.54 (m, 1H), 2.13 - 2.02 (m, 1H).
120		585.0	Method E, RT = 2.500 min, 99.9%	^1H NMR (400MHz, DMSO- d_6) δ = 8.93 (d, J = 2.7 Hz, 1H), 8.85 (d, J = 4.7 Hz, 1H), 8.40 (t, J = 8.2 Hz, 1H), 7.90 (m, 1H), 7.72 - 7.63 (m, 2H), 7.51 (m, 1H),

				7.37 - 7.28 (m, 3H), 4.67 (m, 1H), 4.30 (m, 1H), 2.46 - 2.23 (m, 2H), 1.62 (d, $J = 13.2$ Hz, 6H), 1.19 (d, $J = 5.2$ Hz, 3H).
121		551.2	Method C, RT = 1.596 min, 90.1%	^1H NMR (400MHz, DMSO- d_6) δ = 8.85 (d, $J = 4.9$ Hz, 1H), 8.69 - 8.60 (m, 1H), 8.20 - 8.09 (m, 1H), 7.91 (dd, $J = 6.7, 4.9$ Hz, 1H), 7.68 (m, 1H), 7.43 (dd, $J = 11.0, 2.5$ Hz, 1H), 7.37 - 7.30 (m, 1H), 7.29 - 7.18 (m, 3H), 4.69 - 4.50 (m, 1H), 4.34 - 4.10 (m, 1H), 2.86 - 2.76 (m, 1H), 1.65 - 1.62 (m, 1H), 1.63 (d, $J = 13.4$ Hz, 3H), 1.62 (d, $J = 13.4$ Hz, 3H), 1.12 (d, $J = 6.1$ Hz, 3H).
122		551.2	Method C, RT = 1.592 min, 92.5%	^1H NMR (400MHz, DMSO- d_6) δ = 8.85 (d, $J = 4.2$ Hz, 1 H), 8.65 (s, 1 H), 8.15 (t, $J = 8.9$ Hz, 1 H), 7.90 (m, 1 H), 7.68 (m, 1 H), 7.42 (dd, $J = 11.1, 2.3$ Hz, 1 H), 7.36 - 7.31 (m, 2 H), 7.22 - 7.14 (m, 2 H), 4.70 - 4.59 (m, 1 H), 4.34 - 4.26 (m, 1 H), 2.37 - 2.27 (m, 2 H), 1.62 (d, $J = 13.5$ Hz, 6 H), 1.12 (d, $J = 6.1$ Hz, 3H).
123		618.2 (M+N H_4) $^+$	Method C, RT = 1.502 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.91 (d, $J = 2.7$ Hz, 1H), 8.85 (d, $J = 4.6$ Hz, 1H), 8.41 (t, $J = 8.6$ Hz, 1H), 7.89 (dd, $J = 7.5, 4.0$ Hz, 1H), 7.70 - 7.63 (m, 2H), 7.51 (br. d., $J = 8.6$ Hz, 1H), 7.34 (t, $J = 6.4$ Hz, 2H), 7.25 (d, $J = 7.3$ Hz, 1H), 5.12 (t, $J = 4.9$ Hz, 1H), 4.77 - 4.66 (m, 1H), 4.19 (br. d., $J = 8.3$ Hz, 1H), 3.49 - 3.43 (m, 2H), 2.62 - 2.53 (m, 1H), 2.33 - 2.22 (m, 1H), 1.63 (d, $J = 13.4$ Hz, 3H), 1.62 (d, $J = 13.4$ Hz, 3H).
124		600.2	Method C, RT = 1.577 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 9.19 (br. s., 1H), 8.45 (t, $J = 8.4$ Hz, 1H), 7.96 - 7.86 (m, 1H), 7.70 - 7.58 (m, 3H), 7.52 (d, $J = 8.4$ Hz, 1H), 7.42 - 7.33 (m, 2H), 7.31 - 7.22 (m, 2H), 4.87 (t, $J = 8.1$ Hz, 1H), 4.18 - 4.07 (m, 1H), 3.71 (br. d., $J = 8.8$ Hz, 1H), 3.64 - 3.58 (m, 1H), 3.4 - 3.2 (m, 2H), 2.85 - 2.72 (m, 1H), 1.52 (br. d., $J = 13.2$ Hz, 6H).
125		583.2 (M+N H_4) $^+$	Method D, RT = 1.442 min, 100%	^1H NMR (400MHz, DMSO- d_6) δ = 8.70 - 8.63 (m, 1H), 8.13 (t, $J = 8.6$ Hz, 1H), 7.97 - 7.86 (m, 1H), 7.71 - 7.58 (m, 2H), 7.45 - 7.35 (m, 3H), 7.29 - 7.17 (m, 3H), 4.37 (dd, $J = 10.0, 8.3$ Hz, 1H), 3.88 - 3.81 (m, 1H), 3.76 - 3.66 (m, 1H), 3.61 (dd, $J = 11.0, 6.6$ Hz, 2H), 2.66 - 2.56 (m, 1H), 1.53 (br. d., $J = 13.2$ Hz, 6H).

126		583.2 (M+N H ₄) ⁺	Method D, RT = 1.482 min, 94.5%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.90 (s, 1H), 8.19 (t, <i>J</i> = 8.9 Hz, 1H), 7.98 - 7.86 (m, 1H), 7.69 - 7.59 (m, 2H), 7.43 (dd, <i>J</i> = 11.2, 2.4 Hz, 1H), 7.40 - 7.34 (m, 2H), 7.30 - 7.24 (m, 1H), 7.23 - 7.18 (m, 1H), 7.12 (d, <i>J</i> = 7.8 Hz, 1H), 4.85 (t, <i>J</i> = 8.1 Hz, 1H), 4.17 - 4.07 (m, 1H), 3.71 (br. d., <i>J</i> = 9.5 Hz, 1H), 3.60 (br. dd., <i>J</i> = 9.5, 3.2 Hz, 2H), 2.76 - 2.70 (m, 1H), 1.52 (br. d., <i>J</i> = 13.2 Hz, 6H).
127		583.2 (M+N H ₄) ⁺	Method D, RT = 1.490 min, 96.5%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.91 (s, 1H), 8.20 (t, <i>J</i> = 9.1 Hz, 1H), 7.98 - 7.86 (m, 1H), 7.71 - 7.61 (m, 2H), 7.43 (dd, <i>J</i> = 11.0, 2.5 Hz, 1H), 7.40 - 7.34 (m, 2H), 7.30 - 7.24 (m, 1H), 7.21 - 7.16 (m, 1H), 7.13 (d, <i>J</i> = 8.0 Hz, 1H), 4.86 (t, <i>J</i> = 8.2 Hz, 1H), 4.16 - 4.08 (m, 1H), 3.72 (br. d., <i>J</i> = 9.3 Hz, 1H), 3.60 (br. dd., <i>J</i> = 9.3, 3.2 Hz, 2H), 2.76 - 2.71 (m, 1H), 1.53 (br. d., <i>J</i> = 13.2 Hz, 6H).
128		585.0	Method E, RT = 2,517 min, 99.7%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.93 (br. s., 1H), 8.84 (d, <i>J</i> = 3.9 Hz, 1H), 8.41 (t, <i>J</i> = 8.1 Hz, 1H), 7.93 - 7.88 (m, 1H), 7.74 - 7.61 (m, 2H), 7.51 (br. d., <i>J</i> = 8.3 Hz, 1H), 7.44 - 7.21 (m, 3H), 4.70 - 4.52 (m, 1H), 4.22 (br. d., <i>J</i> = 9.5 Hz, 1H), 2.85 - 2.78 (m, 1H), 2.57 - 2.52 (m, 1H), 1.63 (d, <i>J</i> = 13.4 Hz, 3H), 1.61 (d, <i>J</i> = 13.4 Hz, 3H), 1.13 (d, <i>J</i> = 6.4 Hz, 3H).
129		552.1	Method D, RT = 1.831 min, 100%	¹ H NMR (400MHz, DMSO-d ₆) δ = 8.71 - 8.55 (m, 1H), 8.21 - 8.09 (m, 2H), 7.91 - 7.73 (m, 2H), 7.61 - 7.50 (m, 1H), 7.42 (dd, <i>J</i> = 11.2, 2.0 Hz, 1H), 7.37 - 7.24 (m, 2H), 7.24 - 7.11 (m, 2H), 4.70 - 4.52 (m, 1H), 4.39 - 4.13 (m, 1H), 3.06 (s, 3H), 2.38 - 2.23 (m, 2H), 1.12 (d, <i>J</i> = 6.1 Hz, 3H).
130		548.2	Method C, RT = 1.578 min, 99.6%	¹ H NMR (400MHz, DMSO-d ₆) δ = 9.73 - 9.50 (m, 1H), 8.95 (d, <i>J</i> = 1.2 Hz, 1H), 8.44 (d, <i>J</i> = 1.5 Hz, 1H), 7.97 - 7.85 (m, 1H), 7.71 - 7.56 (m, 2H), 7.43 - 7.31 (m, 3H), 7.30 - 7.21 (m, 1H), 4.67 (d, <i>J</i> = 9.0 Hz, 1H), 3.89 - 3.82 (m, 1H), 3.47 (br. d., <i>J</i> = 9.5 Hz, 1H), 1.53 (d, <i>J</i> = 13.2 Hz, 6H), 1.23 (s, 3H), 1.07 (s, 3H).


131		669.4	Method D, RT = 1.764 min, 100%	¹ H NMR (400 MHz, DMSO-d ₆) δ = 8.93 (d, <i>J</i> = 2.0 Hz, 1H), 8.40 (t, <i>J</i> = 8.6 Hz, 1H), 7.94 (dd, <i>J</i> = 12.6, 7.5 Hz, 1H), 7.73 - 7.57 (m, 3H), 7.50 (br. d., <i>J</i> = 8.6 Hz, 1H), 7.41 - 7.17 (m, 4H), 4.78 - 4.66 (m, 1H), 4.47 - 4.28 (m, 1H), 3.32 - 3.20 (m, 2H), 3.29 - 3.07 (m, 2H), 2.56 - 2.48 (m, 2H), 2.46 - 2.17 (m, 6H), 1.51 (br. d., <i>J</i> = 13.2 Hz, 6H).
132		579.2 (M+N H ₄) ⁺	Method D, RT = 1.736 min, 100%	¹ H NMR (400 MHz, DMSO-d ₆) δ = 8.61 - 8.54 (m, 1H), 8.14 (t, <i>J</i> = 8.8 Hz, 1H), 7.96 - 7.88 (m, 1H), 7.68 - 7.59 (m, 2H), 7.44 (dd, <i>J</i> = 11.1, 2.6 Hz, 1H), 7.41 - 7.35 (m, 2H), 7.30 - 7.24 (m, 1H), 7.22 - 7.18 (m, 1H), 7.09 - 7.01 (m, 1H), 4.78 - 4.67 (m, 1H), 4.16 - 4.05 (m, 1H), 3.62 - 3.53 (m, 1H), 1.52 (br. d., <i>J</i> = 13.2 Hz, 6H), 0.90 - 0.71 (m, 4H).
133		524.1	Method E, RT = 1.518 min, 98.9%	¹ H NMR (400 MHz, DMSO-d ₆) δ = 8.99 (s, 1H), 7.92 (m, 1H), 7.52 - 7.72 (m, 2H), 7.33 - 7.41 (m, 2H), 7.21 - 7.29 (m, 1H), 7.04 (m, 2H), 6.73 (d, <i>J</i> = 7.5 Hz, 1H), 4.53 (m, 1H), 3.83 - 3.95 (m, 1H), 3.71 - 3.81 (m, 1H), 2.60 - 2.54 (m, 1H), 2.04 - 2.19 (m, 1H), 1.52 (br. d., <i>J</i> = 13.2 Hz, 6H).
134		518.1	Method D, RT = 1.511 min, 99.1%	
135		524.1	Method D, RT = 1.496 min, 95.0%	

It will be evident to one skilled in the art that the present disclosure is not limited to the foregoing illustrative examples, and that it can be embodied in other specific forms

without departing from the essential attributes thereof. It is therefore desired that the examples be considered in all respects as illustrative and not restrictive, reference being made to the appended claims, rather than to the foregoing examples, and all changes which come within the meaning and range of equivalency of the claims are therefore
5 intended to be embraced therein.

WE CLAIM:

1. A compound of Formula (II):



or a pharmaceutically acceptable salt thereof, wherein:

- 5 Ar¹ is aryl substituted with 1-2 R^{1a} and 1-2 R^{1b} or monocyclic heteroaryl with 1-3 heteroatoms selected from nitrogen, oxygen, and sulfur, and substituted with 1-2 R^{1a} and 1-2 R^{1b};
- Ar² is aryl substituted with 1-4 R^{2a} or 6-membered heteroaryl with 1-2 nitrogen atoms, and substituted with 1-4 R^{2a};
- 10 Ar³ is aryl substituted with 1-4 R^{3a} or monocyclic heteroaryl with 1-3 heteroatoms selected from nitrogen, oxygen, and sulfur, and substituted with 1-4 R^{3a};
- R^{1a} is hydrogen or halo;
- R^{1b} is halo, haloalkyl, alkoxy, or haloalkoxy;
- R^{2a} is hydrogen, cyano, halo, alkyl, hydroxyalkyl, haloalkyl, cycloalkyl,
- 15 alkoxy, or haloalkoxy; alternatively, two adjacent R^{2a} groups are taken together with the carbon atoms to which they are attached to form a heterocycle with 1-4 heteroatoms selected from nitrogen, oxygen, and sulfur;
- R^{3a} is cyano, halo, alkyl, alkoxy, hydroxyalkyl, alkoxyalkyl, haloalkyl, (R¹R²N)alkyl, R¹R²N, alkylC(O)(R²)Nalkyl, (alkyl)₂(O)P, (alkoxy)₂(O)P,
- 20 (alkoxy)(alkyl)(O)P, (alkyl)(O)(NR¹)S, alkylSO₂, or alkylSO₂NH; alternatively, two adjacent R^{3a} groups are taken together with the carbon atoms to which they are attached to form a heterocycle with 1-4 heteroatoms selected from nitrogen, oxygen, and sulfur;
- R^{4a} or R^{4b} is independently hydrogen, alkyl, alkoxy, hydroxylalkyl,
- 25 alkoxyalkyl, or haloalkoxy; alternatively, R^{4a} and R^{4b} together with the carbon atom they are both attached to form a C₃₋₆ cycloalkyl;

R^{5a} or R^{5b} is independently hydrogen, alkyl, hydroxylalkyl, alkoxyalkyl or haloalkoxyl;

R¹ is hydrogen or alkyl; and


R² is hydrogen or alkyl; or R¹R²N taken together is azetidiny, oxazolyl

5 pyrrolidiny, piperidiny, piperaziny, or morpholiny, and is substituted with 0-3 substituents selected from halo, alkyl, and oxo;

alkyl means a straight or branched alkyl group composed of 1 to 6 carbons;

cycloalkyl means a monocyclic ring system composed of 3 to 7 carbons.

10 2. The compound of claim 1, having Formula (III):



or a pharmaceutically acceptable salt thereof, wherein:

Ar¹ is phenyl substituted with 1-2 R^{1a} and 1-2 R^{1b} or 6-membered heteroaryl with 1-3 nitrogen atoms and substituted with 1R^{1a} and 1-2 R^{1b};

15 Ar³ is phenyl substituted with 1-3 R^{3a} or 5- to 6-membered heteroaryl with 1-3 nitrogen atoms and substituted with 1-3 R^{3a};

R^{1a} is hydrogen or halo;

R^{1b} is halo, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy;

20 R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy;

R^{3a} is cyano, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ alkoxyalkyl, C₁₋₄ haloalkyl, (R¹R²N)C₁₋₄ alkyl, R¹R²N, C₁₋₄ alkylC(O)(R²)NC₁₋₄ alkyl, (C₁₋₄ alkyl)₂(O)P, (C₁₋₄ alkoxy)₂(O)P, (C₁₋₄ alkoxy)(C₁₋₄ alkyl)(O)P, C₁₋₄ alkylSO₂, or C₁₋₄ alkylSO₂NH;

25 R¹R²N taken together is oxazolyl or pyrrolidiny and is substituted with 0-3 substituents selected from halo, alkyl, and oxo;

R^{4a} or R^{4b} is independently hydrogen, C₁₋₄ alkyl, or C₁₋₄ hydroxylalkyl;
alternatively, R^{4a} and R^{4b} together with the carbon atom they are both attached to
form a C₃₋₆ cycloalkyl; and


R^{5a} or R^{5b} is independently hydrogen, C₁₋₄ alkyl, C₁₋₄ hydroxylalkyl, or C₁₋₄
5 alkoxyalkyl.

3. The compound of claim 2, or a pharmaceutically acceptable salt
thereof, wherein:

Ar¹ is phenyl substituted with 1-2 R^{1a} and 1-2 R^{1b}, pyridinyl substituted with 1
10 R^{1a} and 1-2 R^{1b}, or pyrazinyl substituted with 1 R^{1a} and 1-2 R^{1b}; and

Ar³ is phenyl substituted with 1-3 R^{3a}, pyrazolyl substituted with 1-3 R^{3a},
pyridinyl substituted with 1-3 R^{3a}, or pyrimidinyl substituted with 1-3 R^{3a}.

4. The compound of claim 2 or 3, having Formula (IV):



or a pharmaceutically acceptable salt thereof, wherein:

Ar¹ is phenyl substituted with 1-2 R^{1a} and 1-2 R^{1b}, pyridinyl substituted with 1
R^{1a} and 1-2 R^{1b}, or pyrazinyl substituted with 1 R^{1a} and 1-2 R^{1b};

R^{1a} is hydrogen or halo;

20 R^{1b} is halo, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy;

R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆
cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy;


R^{3a} is cyano, halo, C₁₋₃ alkyl, C₁₋₃ hydroxyalkyl, C₁₋₃ alkoxyalkyl, C₁₋₃ haloalkyl, R¹R²N, (C₁₋₃ alkyl)₂(O)P, (C₁₋₃ alkoxy)₂(O)P, (C₁₋₃ alkoxy)(C₁₋₃ alkyl)(O)P, C₁₋₃ alkylSO₂, or C₁₋₃ alkylSO₂NH;

R^{4a} or R^{4b} is independently hydrogen, C₁₋₃ alkyl, or C₁₋₃ hydroxyalkyl;

- 5 alternatively, R^{4a} and R^{4b} together with the carbon atom they are both attached to form a C₃₋₆ cycloalkyl; and

R^{5a} or R^{5b} is independently hydrogen, C₁₋₃ alkyl, C₁₋₃ hydroxyalkyl, or C₁₋₃ alkoxyalkyl.

- 10 5. The compound of claim 4, having Formula (V):



or a pharmaceutically acceptable salt thereof, wherein:

R^{1a} is hydrogen or F;

R^{1b} is halo, C₁₋₂ haloalkyl, or C₁₋₂ alkoxy;

- 15 R^{2a} is hydrogen, halo, C₁₋₃ alkyl, C₁₋₃ haloalkyl, or C₃₋₆ cycloalkyl;

R^{3a} is halo, hydroxyalkyl, alkoxyalkyl, (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, C₁₋₂ alkylSO₂, or C₁₋₂ alkylSO₂NH; and

R^{3a'} is halo.

- 20 6. The compound of claim 5, or a pharmaceutically acceptable salt thereof, wherein:

R^{1a} is hydrogen or F;


R^{1b} is F, Cl, or CF_3 ;

R^{2a} is hydrogen, F, Cl, isopropyl, CF_3 , or cyclopropyl;

R^{3a} is $(CH_3)_2(O)P$, $(CH_3CH_2)_2(O)P$, $(CH_3CH_2O)(CH_3)(O)P$, CH_3SO_2 , or CH_3SO_2NH ; and

5 $R^{3a'}$ is F.

7. The compound of claim 4, having Formula (VI):



(VI)

or a pharmaceutically acceptable salt thereof, wherein:

10 R^{1a} is hydrogen or halo;


R^{1b} is halo, C_{1-2} haloalkyl, or C_{1-2} alkoxy;

R^{2a} is hydrogen, halo, C_{1-3} alkyl, or C_{3-6} cycloalkyl;

R^{3a} is $(C_{1-2} \text{ alkyl})_2(O)P$, $(C_{1-2} \text{ alkoxy})_2(O)P$, $(C_{1-2} \text{ alkoxy})(C_{1-2} \text{ alkyl})(O)P$, C_{1-2} alkyl SO_2 , or C_{1-2} alkyl SO_2NH .

15

8. The compound of claim 2 or 3, having Formula (VII):



or a pharmaceutically acceptable salt thereof, wherein:

Ar³ is pyrazolyl substituted with 1-3 R^{3a}, pyridinyl substituted with 1-3 R^{3a}, or pyrimidinyl substituted with 1-3 R^{3a};

5 R^{1a} is hydrogen or halo;

R^{1b} is halo, C₁₋₄ haloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy;

R^{2a} is hydrogen, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ haloalkyl, C₃₋₆ cycloalkyl, C₁₋₄ alkoxy, or C₁₋₄ haloalkoxy;

10 R^{3a} is cyano, halo, C₁₋₄ alkyl, C₁₋₄ hydroxyalkyl, C₁₋₄ alkoxyalkyl, (R¹R²N)C₁₋₄ alkyl, R¹R²N-, (C₁₋₄ alkyl)₂(O)P, (C₁₋₄ alkoxy)₂(O)P, (C₁₋₄ alkoxy)(C₁₋₄ alkyl)(O)P, C₁₋₄ alkylSO₂, or C₁₋₄ alkylSO₂NH;

R¹ is hydrogen or alkyl;


R² is hydrogen or alkyl; or R¹R²N taken together is oxazolyl or pyrrolidinyl, and is substituted with 0-3 substituents selected from halo, alkyl, or oxo;

15 R^{4a} or R^{4b} is independently hydrogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkoxyalkyl, or C₁₋₄ haloalkoxy; R^{4a} and R^{4b} together with the carbon atom they are both attached to form a C₃₋₆ cycloalkyl; and

R^{5a} or R^{5b} is independently hydrogen, C₁₋₄ alkyl, C₁₋₄ alkoxy, C₁₋₄ alkoxyalkyl, or C₁₋₄ haloalkoxy.

20

9. The compound of claim 8, having Formula (VIII):



or a pharmaceutically acceptable salt thereof, wherein:

R^{1a} is hydrogen or halo;


R^{1b} is halo, C_{1-2} haloalkyl, or C_{1-2} alkoxy;

5 R^{2a} is hydrogen, halo, C_{1-3} alkyl, or C_{3-6} cycloalkyl;

R^{3a} is $(C_{1-2} \text{ alkyl})_2(O)P$, $(C_{1-2} \text{ alkoxy})_2(O)P$, $(C_{1-2} \text{ alkoxy})(C_{1-2} \text{ alkyl})(O)P$, C_{1-2} alkylSO₂, or C_{1-2} alkylSO₂NH; and

R^{5a} or R^{5b} is independently hydrogen or C_{1-2} alkyl.

10 10. The compound of claim 1, having Formula (IX):




or a pharmaceutically acceptable salt thereof, wherein:

R^{1a} is hydrogen or halo;

R^{1b} is halo, C_{1-2} haloalkyl, or C_{1-2} alkoxy;

R^{2a} is hydrogen, halo, C₁₋₂ haloalkyl, C₁₋₃ alkyl, or C₃₋₆ cycloalkyl;



R^{3a} is (C₁₋₂ alkyl)₂(O)P, (C₁₋₂ alkoxy)(C₁₋₂ alkyl)(O)P, or C₁₋₂ alkylSO₂NH;

R^{3a'} is halo;

- 5 R^{4a} or R^{4b} is independently hydrogen, C₁₋₂ alkyl, or C₁₋₂ hydroxyalkyl; or R^{4a} and R^{4b} together with the carbon atom to which they are both attached form a C₃₋₆ cycloalkyl; and

 R^{5a} or R^{5b} is independently hydrogen, C₁₋₂ alkyl, C₁₋₂ hydroxyalkyl, or C₁₋₂ alkoxyalkyl.

10

11. The compound of claim 4, or a pharmaceutically acceptable salt thereof, wherein:

 R^{4a} is hydrogen;

 R^{4b} is hydrogen;

- 15 R^{5a} is C₁₋₂ alkyl, C₁₋₂ hydroxylalkyl, or C₁₋₂ alkoxyalkyl; and

 R^{5b} is hydrogen.

12. The compound of claim 4, or a pharmaceutically acceptable salt thereof, wherein:


- 20 R^{4a} is C₁₋₂ alkyl;

 R^{4b} is C₁₋₂ alkyl;

 R^{5a} is hydrogen; and

 R^{5b} is hydrogen.

- 25 13. A compound according to claim 1, of Formula (I):



wherein:

Ar¹ is phenyl, pyridinyl, or pyridazinyl and is substituted with 1 halo, haloalkyl or haloalkoxy substituent in the 4-position and 0-2 additional halo or haloalkyl substituents;

Ar² is phenyl or pyridinyl and is substituted with 0-2 substituents selected from cyano, fluoro, alkyl, haloalkyl, cycloalkyl, alkoxy, and haloalkoxy;

Ar³ is phenyl or pyridinyl and is substituted with 0-2 substituents selected from cyano, halo, hydroxyalkyl, alkoxyalkyl, (R¹R²N)alkyl, (alkyl)₂(O)P, (alkyl)(O)(NR¹)S, alkylSO₂, and alkylSO₂NH;

R¹ is hydrogen or alkyl; and

R² is hydrogen or alkyl;

or (R¹)(R²)N taken together is azetidiny, pyrrolidinyl, piperidinyl, piperazinyl, or morpholinyl and is substituted with 0-3 substituents selected from halo and alkyl;

or a pharmaceutically acceptable salt thereof.


14. A compound of claim 13 where Ar¹ is phenyl substituted with 1 halo, haloalkyl or haloalkoxy substituent in the 4-position and 0-2 additional halo or haloalkyl substituents.


15. A compound of claim 13 where Ar² is phenyl substituted with 0 substituents, 1 alkyl or cycloalkyl substituent, or 1-2 fluoro substituents.


16. A compound of claim 13 where Ar³ is phenyl substituted with 1-2 substituents selected from cyano, halo, hydroxyalkyl, alkoxyalkyl, (R¹R²N)alkyl, (alkyl)₂(O)P, (alkyl)(O)(NR¹)S, alkylSO₂, and alkylSO₂NH.

5 17. A compound of claim 13 where R¹ is hydrogen and R² is hydrogen.


18. A compound of claim 13 selected from the group consisting of













- 5 19. A compound according to claim 1, selected from the group consisting of










or a pharmaceutically acceptable salt thereof.


21. A compound according to claim 19, which is



5


or a pharmaceutically acceptable salt thereof.

22. A compound according to claim 19, which is




10 or a pharmaceutically acceptable salt thereof.

23. A compound according to claim 19, which is



or a pharmaceutically acceptable salt thereof.

24. A compound according to claim 1, which is



or a pharmaceutically acceptable salt thereof.

25. A composition comprising a compound of any one of claims 1-24, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, diluent, or excipient.

26. A compound according to any one of claims 1-24, or a pharmaceutically acceptable salt thereof or composition according to claim 25 for use in therapy.

27. A compound of any one of claims 1-24 or a pharmaceutically acceptable salt thereof for use in treating heart disease in a patient in need thereof.

28. The compound or a pharmaceutically acceptable salt thereof for use according to claim 27 wherein the heart disease is selected from the group consisting of angina pectoris, unstable angina, myocardial infarction, heart failure, acute coronary disease, acute heart failure, chronic heart failure, and cardiac iatrogenic damage.

29. The compound or pharmaceutically acceptable salt thereof for use according to 28 wherein the heart failure is selected from the group consisting of congestive heart failure, systolic heart failure, diastolic heart failure, heart failure with reduced ejection fraction (HFrEF), heart failure with preserved ejection fraction (HFpEF), acute heart failure, chronic heart failure of ischemic and non-ischemic origin.

30. The compound or pharmaceutically acceptable salt thereof for use according to claim 28 wherein the heart disease is heart failure.

31. Use of the compound of any one of claims 1-24 or a pharmaceutically acceptable salt thereof in the manufacture of a medicament for treating heart disease in a patient in need thereof.

32. The use of claim 31 wherein the heart disease is selected from the group consisting of angina pectoris, unstable angina, myocardial infarction, heart failure, acute coronary disease, acute heart failure, chronic heart failure, and cardiac iatrogenic damage.

33. The use of claim 32 wherein the heart failure is selected from the group consisting of congestive heart failure, systolic heart failure, diastolic heart failure, heart failure with reduced ejection fraction (HFrEF), heart failure with preserved ejection fraction (HFpEF), acute heart failure, chronic heart failure of ischemic and non-ischemic origin.

34. The use of claim 32 wherein the heart disease is heart failure.