

HMO production

Technical field

5 The present invention relates to the field of recombinant production of biological molecules in host cells. More particularly it relates to a method for recombinant production of human milk oligosaccharides (HMO) using a genetically modified cell expressing a protein of the major facilitator superfamily (MFS).

10 Background of the invention

Human milk oligosaccharides (HMOs) constitute the third largest solid component in human milk and are highly resistant to enzymatic hydrolysis. As a consequence, a substantial fraction of HMOs remains largely undigested and unabsorbed, which enables their passage through to the colon. In the colon, HMOs
15 may serve as substrates to shape the gut ecosystem by selectively stimulating the growth of specific saccharolytic bacteria. This selectivity is viewed as beneficial for both infants and adults since strains of *Bifidobacterium* species are believed to have a positive effect on gut health (Chichlowski M. *et al.*, (2012) J. Pediatr. Gastroenterol. Nutr. 5:251-258; Elison E. *et al.*, (2016) Brit J. Nutr, 116: 1356–1368).

20 Besides their prebiotic properties, HMOs have been linked to additional positive effects, which expands their field of application (Kunz C. *et al.*, (2014) Food Oligosaccharides: Production, Analysis and Bioactivity, 1st Edition, p 5-20, Eds. Moreno J. and Luz Sanz M., John Wiley & Sons, Ltd).

The obvious health benefits of HMOs have enabled their approval for use in
25 foods, such as infant formulas and foods, and for consumer health products. Biotechnological production of HMOs is a valuable cost-efficient and large-scale way of HMO manufacturing. It relies on genetically engineered bacteria constructed so as to express the glycosyltransferases needed for synthesis of the desired oligosaccharides and takes advantage of the bacteria's innate pool of nucleotide
30 sugars as HMO precursors. Recent developments in biotechnological production of HMOs have made it possible to overcome certain inherent limitations of bacterial expression systems. For example, HMO-producing bacterial cells may be genetically modified to increase the limited intracellular pool of nucleotide sugars in the bacteria (WO2012112777), to improve activity of enzymes involved in the HMO production
35 (WO2016040531), or to facilitate the secretion of synthesized HMOs into the extracellular media (WO2010142305, WO2017042382). Further, expression of

genes of interest in recombinant cells may be regulated by using particular promoters or other gene expression regulators, like e.g. what has recently been described in WO2019123324.

The approach described in WO2010142305 and WO2017042382 has an advantage in that it allows to reduce the metabolic burden inflicted on the producing cell by high levels of recombinant gene expression, e.g. using methods of WO2012112777, WO2016040531 or WO2019123324. This approach attracts growing attention in recombinant HMO-producing cell engineering, e.g. recently there have been described several new genes encoding sugar transporter proteins, and fermentation procedures that can facilitate the efflux of a recombinantly produced 2'-fucosyllactose (2'-FL), the most abundant HMO of human milk (WO2018077892, US201900323053, US201900323052). However, at present, there is no algorithm that is able to pinpoint the right transporter protein for the efflux of different recombinantly produced HMO structures among numerous bacterial proteins with predicted transporter function in multiple protein databases, e.g. UniProt, since the structures/factors defining substrate specificity of sugar transporters are still not well-studied and remain highly unpredictable.

Summary of the invention

Efficient sugar efflux transporter proteins for recombinantly produced HMOs and development of recombinant cells expressing said proteins are advantageous for high scale industrial HMO manufacturing.

This invention provides recombinant cells capable of producing a human milk oligosaccharide (HMO), wherein the cells are expressing a heterologous gene encoding a putative MFS (major facilitator superfamily) transporter protein, originating from the bacterium *Yersinia bercovieri*. More specifically, the invention relates to a genetically modified cell optimized for the production of an oligosaccharide, in particular an HMO, comprising a recombinant nucleic acid encoding a protein having at least 80 % sequence identity to the amino acid sequence of SEQ ID NO: 1 (Figure 4). The amino acid sequence identified herein as SEQ ID NO: 1 is the amino acid sequence that is 100 % identical to the amino acid sequence having the GenBank accession ID EEQ08298.1 (<https://www.ncbi.nlm.nih.gov/protein/EEQ08298.1/>). The MFS transporter protein having the amino acid sequence of SEQ ID NO: 1 is identified herein as "YberC protein" or "YberC transporter" or "YberC", interchangeably; a nucleic acid sequence

encoding YberC protein is identified herein as "YberC coding nucleic acid/DNA" or "*yberC* gene" or "*yberC*".

The present invention shows that the use of HMO producing recombinant cells that express YberC protein results in very distinct improvements of the HMO manufacturing process, both in fermentation and purification of the HMOs. The disclosed herein recombinant cells and methods for HMO production provide both higher yields of total produced HMOs, lower by-product formation or by-product-to-product ratio, lower biomass formation per fermentation and facilitated recovery of the HMOs during downstream processing of the fermentation broth.

Surprisingly, the expression of a DNA sequence encoding YberC in different HMO producing cells is found to be associated with the accumulation of some particular HMOs in the extracellular media and accumulation of some other HMOs in the interior of the producing cells, and in an increase in total HMO production. Surprisingly, an increase in the efflux of the produced HMOs is found to be characteristic for HMOs that consist of either tri or tetra units of monosaccharides, i.e. HMOs that are trisaccharides and tetrasaccharides, e.g. 3'-sialyllactose (3'-SL), 6'-sialyllactose (6'-SL), lacto-N-triose 2, (LNT-2), lacto-N-neotetraose (LNnT) and lacto-N-tetraose (LNT), but not for larger oligosaccharide structures, like pentasaccharides and hexasaccharides, which accumulate inside of the producing cells. Surprisingly, it is also found that in particular the total production of the HMOs lacto-N-triose 2 (LNT-2), lacto-N-tetraose (LNT) and lacto-N-neotetraose (LNnT), in the corresponding HMO producing cells expressing *yberC* is increased, while the by-product production, e.g. para-lacto-N-neohexaose (pLNnH) or para-lacto-neohexaose II (pLNH-II), in these cells, correspondingly, is often decreased and said by-product oligosaccharides typically accumulate inside of the production cells. Further, highly unexpectedly, expression of YberC protein in HMO producing cells leads to reduction in biomass formation during high-cell density fermentations and to healthier cell cultures, as reflected e.g. by a decrease in the number of dead cells at the end of fermentation, which makes the manufacturing process more efficient as more product is produced per biomass unit.

Accordingly, a first aspect of the invention relates to a genetically modified cell capable of producing one or more Human Milk Oligosaccharides (HMOs), selected from the list of 3'-sialyllactose (3'-SL), 6'-sialyllactose (6'-SL), Lacto-N-Triose-2 (LNT-2), Lacto-N-tetraose (LNT), Lacto-N-neotetraose (LNnT), Lacto-N-fucopentaose I (LNFP I), Lacto-N-fucopentaose II (LNFP II), Lacto-N-fucopentaose III (LNFP III), Lacto-N-fucopentaose IV (LNFP IV), and Lacto-N-fucopentaose V

(LNFP V), para-lacto-N-neohexaose (pLNnH), and para-lacto-neohexaose II (pLNH-II), wherein said cell comprises a recombinant nucleic acid encoding a protein of SEQ ID NO: 1, or a functional homologue thereof having an amino acid sequence that is at least 80% identical, preferably at least 85% identical, more preferably at least 90% identical to SEQ ID NO: 1. In particular, the invention relates to said genetically modified cell capable of producing one or more Human Milk Oligosaccharides (HMOs), wherein the oligosaccharide is selected from the list of Lacto-N-Triose-2 (LNT-2), Lacto-N-tetraose (LNT), Lacto-N-neotetraose (LNnT), Lacto-N-fucopentaose I (LNFP I), Lacto-N-fucopentaose II (LNFP II), Lacto-N-fucopentaose III (LNFP III), Lacto-N-fucopentaose IV (LNFP IV), and Lacto-N-fucopentaose V (LNFP V), para-lacto-N-neohexaose (pLNnH), and para-lacto-neohexaose II (pLNH-II).

A second aspect of the invention relates to a nucleic acid construct comprising a nucleic acid sequence(s) encoding an MFS transporter protein, wherein the nucleic acid sequence encoding the protein has at least 70 % sequence identity to SEQ ID NO: 2, such as at least 80 %, such as at least 85%, such as at least 95 %, such as at least 99%, as well as to a genetically modified cell comprising the nucleic acid construct, which is *Escherichia coli*.

In one aspect, the nucleic acid construct comprising a nucleic acid sequence(s) encoding an MFS transporter protein, wherein the nucleic acid sequence is at least 70% identical to SEQ ID NO: 2.

A third aspect of the invention relates to a method for the production of one or more oligosaccharides, the method comprising the steps of:

- (i) providing a genetically modified cell capable of producing an HMO, wherein said cell comprises a recombinant nucleic acid encoding a protein of SEQ ID NO: 1, or a functional homologue thereof having an amino acid sequence is at least 80 % identical, preferably at least 85 % identical, more preferably at least 90 % identical to SEQ ID NO: 1;
- (ii) culturing the cell according to (i) in a suitable cell culture medium to express said recombinant nucleic acid;
- (iii) harvesting one or more HMOs produced in step (ii).

The invention also relates to the use of a genetically modified cell or a nucleic acid construct comprising a heterologous nucleic acid sequence encoding a

Major facilitator superfamily (MFS) protein, said nucleic acid sequence having at least 70 % sequence identity to SEQ ID NO: 2, for the production of one or more Human Milk Oligosaccharides (HMOs) selected from the list of 3'-sialyllactose (3'-SL), 6'-sialyllactose (6'-SL), Lacto-N-Triose-2 (LNT-2), Lacto-N-tetraose (LNT),

5 Lacto-N-neotetraose (LNnT), Lacto-N-fucopentaose I (LNFP I), Lacto-N-fucopentaose II (LNFP II), Lacto-N-fucopentaose III (LNFP III), Lacto-N-fucopentaose IV (LNFP IV), and Lacto-N-fucopentaose V (LNFP V), para-lacto-N-neohexaose (pLNnH), and para-lacto-neohexaose II (pLNH-II), in particular selected

10 from the list of Lacto-N-Triose-2 (LNT-2), Lacto-N-tetraose (LNT), Lacto-N-neotetraose (LNnT), Lacto-N-fucopentaose I (LNFP I), Lacto-N-fucopentaose II (LNFP II), Lacto-N-fucopentaose III (LNFP III), Lacto-N-fucopentaose IV (LNFP IV), and Lacto-N-fucopentaose V (LNFP V), para-lacto-N-neohexaose (pLNnH), and para-lacto-neohexaose II (pLNH-II).

15 As mentioned above, during the culturing of genetically modified cells capable of producing one or more HMOs, which cells comprise a nucleic acid encoding an YberC transporter protein, it has surprisingly been found that the corresponding one or more HMOs are produced in high yields, while by-product and biomass formation is reduced. This is advantageous for the recovery of the HMOs

20 during downstream processes, as e.g. the overall recovery and purification procedure may comprise less steps and overall time of purification may be shortened, and therefore the recovery of the product is less laborious and more economically efficient.

These effects of increased product yields and facilitation of the product

25 recovery makes the present invention superior to the disclosures of the prior art.

Other aspects and advantageous features of the present invention are described in detail and illustrated by non-limiting working examples below.

Brief description of figures

30 **Figure 1** shows the percentage (%) relative LNT-2 concentrations in total, supernatant and pellet samples for strains with optimal expression of a glycosyltransferase gene differing only in the expression of the heterologous transporter gene *yberC0001_9420*. Contrary to the strain MP3941, the strain

35 MP3940 expresses the *yberC0001_9420* gene .r

Figure 2 shows the percentage (%) relative LNnT and by-product concentrations for the strains MP4436 and MP3863 in total, supernatant and pellet samples. Although both strains show optimal expression of the required glycosyltransferase genes, it is MP3863 that expresses the heterologous transporter gene *yberC0001_9420* under the control of the *Plac* element.

Figure 3 shows the percentage (%) relative LNT and by-product concentrations for the strains MP4473, MP4462 and MP4547 in total, supernatant and pellet samples. Although all three strains show optimal expression of the required genes, it is only MP4462 and MP4547 that express the heterologous transporter gene *yberC0001_9420* under the control of the *Plac* or *PglpF* element, respectively.

Figure 4 presents SEQ ID NO: 1.

Detailed description

In the following, embodiments of the invention will be described in further detail. Each specific variation of the features can be applied to other embodiments of the invention, unless specifically stated otherwise.

Generally, all terms used herein are to be interpreted according to their ordinary meaning in the technical field, and applicable to all aspects and embodiments of the invention, unless explicitly defined or stated otherwise. All references to "a/an/the [cell, sequence, gene, transporter, step, etc.]" are to be interpreted openly as referring to at least one instance of said cell, sequence, gene, transporter, step, etc., unless explicitly stated otherwise. The steps of any method disclosed herein do not have to be performed in the exact order disclosed, unless explicitly stated.

The present invention in general relates to a genetically modified cell for efficient production of oligosaccharides and use of the said genetically modified cell in a method of producing the oligosaccharides. In particular, the present invention relates to a genetically modified cell enabled to synthesize an oligosaccharide, preferably a heterologous oligosaccharide, in particular a human milk oligosaccharide (HMO). Accordingly, a cell of the invention is modified to express a set of recombinant nucleic acids that are necessary for synthesis of one or more HMOs by the cells (which enable the cell to synthesize one or more HMOs), such as genes encoding one or more enzymes with glycosyltransferase activity described

below. The oligosaccharide producing recombinant cell of the invention is further modified to comprise a heterologous recombinant nucleic acid sequence, preferably, a DNA sequence, encoding a MFS (major facilitator superfamily) protein, originating from the bacterium *Yersinia bercovieri*. More specifically, the invention relates to a

5 genetically modified cell optimized for the production of one or more particular oligosaccharides, in particular one or more particular HMOs, comprising a recombinant nucleic acid encoding a protein having at least 80 % sequence similarity, preferably at least 85 %, more preferably at least 90 %, and even more preferably at least 95 % sequence similarity to the amino acid sequence of SEQ ID

10 NO: 1 (Figure 4). The amino acid sequence identified herein as SEQ ID NO: 1 is an amino acid sequence that has 100 % identity with the amino acid sequence having the GenBank accession ID: EEQ08298.1 (<https://www.ncbi.nlm.nih.gov/protein/EEQ08298.1/>).

Accordingly, a first aspect of the invention relates to a genetically

15 modified cell capable of producing one or more HMOs, wherein said cell comprises a recombinant nucleic acid encoding a protein of SEQ ID NO: 1, or a functional homologue thereof having amino acid sequence is at least 80 % identical, preferably at least 85 % identical, more preferably at least 90 % identical to SEQ ID NO: 1.

The MFS transporter protein having the amino acid sequence of SEQ ID

20 NO: 1 is identified herein as "YberC protein" or "YberC transporter" or "YberC", interchangeably; a nucleic acid sequence encoding YberC protein is identified here in as "YberC coding nucleic acid /DNA" or "*yberC* gene" or "*yberC*".

By the term "Major Facilitator Superfamily (MFS)" is meant a large and exceptionally diverse family of the secondary active transporter class, which is

25 responsible for transporting a range of different substrates, including sugars, drugs, hydrophobic molecules, peptides, organic ions, etc. The specificity of sugar transporter proteins is highly unpredictable and the identification of novel transporter protein with specificity towards for example oligosaccharides requires unburden laboratory experimentation (for more details see review by Reddy V.S. *et al.*, (2012),

30 FEBS J. 279(11): 2022–2035). The term "MFS transporter" means in the present context a protein that facilitates transport of an oligosaccharide, preferably, an HMO, through the cell membrane, preferably transport of an HMO/oligosaccharide synthesized by the host cell from the cytosol to the cell exterior, preferably an

HMO/oligosaccharide comprising three or four sugar units, in particular, LNT-2, LNT,

35 LNT, 3'-SL or 6'-SL. Additionally, or alternatively, the MFS transporter, may also facilitate efflux of molecules which are not considered HMO or oligosaccharides

according to the present invention, such as lactose, glucose, cell metabolites or toxins.

The term "sequence identity of [a certain] %" in the context of two or more nucleic acid or amino acid sequences means that the two or more sequences have
5 nucleotides or amino acid residues in common in the given percent when compared and aligned for maximum correspondence over a comparison window or designated sequences of nucleic acids or amino acids (i.e. the sequences have at least 90 percent (%) identity). Percent identity of nucleic acid or amino acid sequences can be measured using a BLAST 2.0 sequence comparison algorithm with default
10 parameters, or by manual alignment and visual inspection (see e.g. <http://www.ncbi.nlm.nih.gov/BLAST/>). This definition also applies to the complement of a test sequence and to sequences that have deletions and/or additions, as well as those that have substitutions. An example of an algorithm that is suitable for determining percent identity, sequence similarity and for alignment is the BLAST
15 2.2.20 algorithm, which is described in Altschul *et al. Nucl. Acids Res.* **25**, 3389 (1997). BLAST 2.2.20 is used to determine percent sequence identity for the nucleic acids and proteins of the invention. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). Examples of sequence alignment algorithms are
20 CLUSTAL Omega (<http://www.ebi.ac.uk/Tools/msa/clustalo/>), EMBOSS Needle (http://www.ebi.ac.uk/Tools/psa/emboss_needle/), MAFFT (<http://mafft.cbrc.jp/alignment/server/>) or MUSCLE (<http://www.ebi.ac.uk/Tools/msa/muscle/>).

In the context of the invention, the term "oligosaccharide" means a
25 saccharide polymer containing a number of monosaccharide units. In some embodiments, preferred oligosaccharides are saccharide polymers consisting of three or four monosaccharide units, i.e. trisaccharides or tetrasaccharides. Preferable oligosaccharides of the invention are human milk oligosaccharides (HMOs).

30 The term "human milk oligosaccharide" or "HMO" in the present context means a complex carbohydrate found in human breast milk (for reference, see Urashima *et al.*: *Milk Oligosaccharides*. Nova Science Publisher (2011); or Chen, *Adv. Carbohydr. Chem. Biochem.* **72**, 113 (2015)). The HMOs have a core structure comprising a lactose unit at the reducing end that can be elongated by one or more
35 β -N-acetyl-lactosaminy and/or one or more β -lacto-N-biosyl units, and this core

structure can be substituted by an alpha-L-fucopyranosyl and/or an alpha-N-acetyl-neuraminy (sialyl) moiety. In this regard, the non-acidic (or neutral) HMOs are devoid of a sialyl residue, and the acidic HMOs have at least one sialyl residue in their structure. The non-acidic (or neutral) HMOs can be fucosylated or non-
5 fucosylated. Examples of such neutral non-fucosylated HMOs include lacto-N-triose 2 (LNT-2) lacto-N-tetraose (LNT), lacto-N-neotetraose (LNnT), lacto-N-neohexaose (LNnH), para-lacto-N-neohexaose (pLNnH), para-lacto-N-hexaose (pLNH) and lacto-N-hexaose (LNH). Examples of neutral fucosylated HMOs include 2'-fucosyllactose (2'-FL), lacto-N-fucopentaose I (LNFP-I), lacto-N-difucohexaose I (LNDFH-I), 3-
10 fucosyllactose (3-FL), difucosyllactose (DFL), lacto-N-fucopentaose II (LNFP-II), lacto-N-fucopentaose III (LNFP-III), lacto-N-difucohexaose III (LNDFH-III), fucosyl-lacto-N-hexaose II (FLNH-II), lacto-N-fucopentaose V (LNFP-V), lacto-N-difucohexaose II (LNDFH-II), fucosyl-lacto-N-hexaose I (FLNH-I), fucosyl-para-lacto-N-hexaose I (FpLNH-I), fucosyl-para-lacto-N-neohexaose II (F-pLNnH II) and
15 fucosyl-lacto-N-neohexaose (FLNnH). Examples of acidic HMOs include 3'-sialyllactose (3'-SL), 6'-sialyllactose (6'-SL), 3-fucosyl-3'-sialyllactose (FSL), 3'-O-sialyllacto-N-tetraose a (LST a), fucosyl-LST a (FLST a), 6'-O-sialyllacto-N-tetraose b (LST b), fucosyl-LST b (FLST b), 6'-O-sialyllacto-N-neotetraose (LST c), fucosyl-LST c (FLST c), 3'-O-sialyllacto-N-neotetraose (LST d), fucosyl-LST d (FLST d), sialyl-
20 lacto-N-hexaose (SLNH), sialyl-lacto-N-neohexaose I (SLNH-I), sialyl-lacto-N-neohexaose II (SLNH-II) and disialyl-lacto-N-tetraose (DSLNT). In the context of the present invention lactose is not regarded as an HMO species.

In some embodiments of the invention, tri-HMOs and tetra-HMOs may be a preferred, e.g. trisaccharides LNT-2, 3'-SL, 6'-SL, and tetrasaccharides LNT, LNnT,
25 FSL.

To be able to synthesize one or more HMOs, the recombinant cell of the invention comprises at least one recombinant nucleic acid which encodes a functional enzyme with glycosyltransferase activity. The galactosyltransferase gene may be integrated into the genome (by chromosomal integration) of the host cell, or
30 alternatively, it may be comprised in a plasmid DNA and expressed as plasmid-borne. If two or more glycosyltransferases are needed for the production of an HMO, e.g. LNT or LNnT, two or more recombinant nucleic acids encoding different enzymes with glycosyltransferase activity may be integrated in the genome and/or expressed from a plasmid, e.g. a β -1,3-N-acetylglucosaminyltransferase (a first
35 recombinant nucleic acid encoding a first glycosyltransferase) in combination with a

β -1,3-galactosyltransferase (a second recombinant nucleic acid encoding a second glycosyltransferase) for the production of LNT, where the first and second recombinant nucleic acid can independently from each other be integrated chromosomally or on a plasmid. In one preferred embodiment, both the first and second recombinant nucleic acid are stably integrated into the chromosome of the production cell; in another embodiment at least one of the first and second glycosyltransferase is plasmid-borne. A protein/enzyme with glycosyltransferase activity (glycosyltransferase) may be selected in different embodiments from enzymes having the activity of α -1,2-fucosyltransferase, α -1,3-fucosyltransferase, α -1,3/4-fucosyltransferase, α -1,4-fucosyltransferase, α -2,3-sialyltransferase, α -2,6-sialyltransferase, β -1,3-N-acetylglucosaminyltransferase, β -1,6-N-acetylglucosaminyltransferase, β -1,3-galactosyltransferase and β -1,4-galactosyltransferase. For example, the production of LNT requires that the modified cell expresses at least two glycosyltransferases, a β -1,3-N-acetylglucosaminyltransferase and a β -1,3-galactosyltransferase; the production of LNnT requires that the modified cell expresses an active β -1,3-N-acetylglucosaminyltransferase and an active β -1,4-galactosyltransferase; for the production of 6'-SL the modified cell has to express an active α -2,6-sialyltransferase enzyme. Some non-limiting embodiments of proteins having glycosyltransferase activity, which can be encoded by the recombinant genes comprised by the production cell, can be selected from non-limiting examples of Table 1.

Table 1

Gene	Protein Sequence ID (GenBank)	Description	HMO example
<i>lgtA_Nm</i>	WP_002248149.1	β -1,3-N-acetylglucosaminyltransferase	LNT, LNnT, LNFP-I, LNFP-II, LNFP-III, LNFP-V, LNFP-VI, LNDFH-I, LNDFH-II, pLNH, F-pLNH I, pLNnH
<i>lgtA_Nm_MC58</i>	AAF42258.1	β -1,3-N-acetylglucosaminyltransferase	LNT, LNnT, LNFP-I, LNFP-II, LNFP-III, LNFP-V, LNFP-VI, LNDFH-I, LNDFH-II, pLNH, F-pLNH I, pLNnH
<i>lgtA_Hd</i>	AAN05638.1	β -1,3-N-acetylglucosaminyltransferase	LNT, LNnT, LNFP-I, LNFP-II, LNFP-III, LNFP-V, LNFP-VI, LNDFH-I, LNDFH-II, pLNH, F-pLNH I, pLNnH
<i>lgtA_Ng_PID2</i>	AAK70338.1	β -1,3-N-acetylglucosaminyltransferase	LNT, LNnT, LNFP-I, LNFP-II, LNFP-III, LNFP-V, LNFP-VI, LNDFH-I, LNDFH-II, pLNH, F-pLNH I, pLNnH

<i>lgtA_Ng_NCCP11945</i>	<i>ACF31229.1</i>	β -1,3-N-acetylglucosaminyl-transferase	LNT, LNnT, LNFP-I, LNFP-II, LNFP-III, LNFP-V, LNFP-VI, LNDFH-I, LNDFH-II, pLNH, F-pLNH I, pLNnH
<i>lgtA_Past</i>	<i>AAK02595.1</i>	β -1,3-N-acetylglucosaminyl-transferase	LNT, LNnT, LNFP-I, LNFP-II, LNFP-III, LNFP-V, LNFP-VI, LNDFH-I, LNDFH-II, pLNH, F-pLNH I, pLNnH
<i>lgtA_Nc</i>	<i>EEZ72046.1</i>	β -1,3-N-acetylglucosaminyl-transferase	LNT, LNnT, LNFP-I, LNFP-II, LNFP-III, LNFP-V, LNFP-VI, LNDFH-I, LNDFH-II, pLNH, F-pLNH I, pLNnH
<i>lgtA_Nm_87255</i>	<i>ELK60643.1</i>	β -1,3-N-acetylglucosaminyl-transferase	LNT, LNnT, LNFP-I, LNFP-II, LNFP-III, LNFP-V, LNFP-VI, LNDFH-I, LNDFH-II, pLNH, F-pLNH I, pLNnH
<i>galT_Hp/HP0826</i>	<i>NP_207619.1</i>	β -1,4-galactosyltransferase	LNnT, LNFP-III, LNFP-VI, pLNH I, F-pLNH I, pLNnH
<i>galT_Nm/lgtB</i>	<i>AAF42257.1</i>	β -1,4-galactosyltransferase	LNnT, LNFP-III, LNFP-VI, pLNH I, F-pLNH I, pLNnH
<i>wbgO</i>	<i>WP_000582563.1</i>	β -1,3-galactosyltransferase	LNT, LNFP-I, LNFP-II, LNFP-V, LNDFH-I, LNDFH-II, pLNH, F-pLNH I
<i>cpslBJ</i>	<i>AB050723.1</i>	β -1,3-galactosyltransferase	LNT, LNFP-I, LNFP-II, LNFP-V, LNDFH-I, LNDFH-II, pLNH, F-pLNH I
<i>jhp0563</i>	<i>AEZ55696.1</i>	β -1,3-galactosyltransferase	LNT, LNFP-I, LNFP-II, LNFP-V, LNDFH-I, LNDFH-II, pLNH, F-pLNH I
<i>galTK</i>	<i>homologous to BD182026</i>	β -1,3-galactosyltransferase	LNT, LNFP-I, LNFP-II, LNFP-V, LNDFH-I, LNDFH-II, pLNH, F-pLNH I
<i>futC</i>	<i>WP_080473865.1</i>	α -1,2-fucosyl-transferase	2'-FL, DFL, LNFP-I, LNDFH-I
<i>FucT2_HpUA802</i>	<i>AAC99764.1</i>	α -1,2-fucosyl-transferase	2'-FL, DFL, LNFP-I, LNDFH-I
<i>FucT2_EcO126t</i>	<i>ABE98421.1</i>	α -1,2-fucosyl-transferase	2'-FL, DFL, LNFP-I, LNDFH-I
<i>FucT2_Hm12198</i>	<i>CBG40460.1</i>	α -1,2-fucosyl-transferase	2'-FL, DFL, LNFP-I, LNDFH-I
<i>FucT2_Pm9515</i>	<i>ABM71599.1</i>	α -1,2-fucosyl-transferase	2'-FL, DFL, LNFP-I, LNDFH-I
<i>FucT2_HpF57</i>	<i>BAJ59215.1</i>	α -1,2-fucosyl-transferase	2'-FL, DFL, LNFP-I, LNDFH-I
<i>FucT6_3_Bf</i>	<i>CAH09151.1</i>	α -1,3-fucosyl-transferase	2'-FL, 3-FL, DFL, LNFP-I, LNFP-III, LNFP-V, LNFP-VI, LNDFH-II, F-pLNH I
<i>FucT7_3_Bf</i>	<i>CAH09495.1</i>	α -1,3-fucosyl-transferase	2'-FL, 3-FL, DFL, LNFP-I, LNFP-III, LNFP-V, LNFP-VI, LNDFH-II, F-pLNH I
<i>FucT_3_Am</i>	<i>ACD04596.1</i>	α -1,3-fucosyl-transferase	2'-FL, 3-FL, DFL, LNFP-I, LNFP-III, LNFP-V, LNFP-VI, LNDFH-II, F-pLNH I
<i>MAMA_R764</i>	<i>AGC02224.1</i>	α -1,3-fucosyl-transferase	2'-FL, 3-FL, DFL, LNFP-I, LNFP-III, LNFP-V, LNFP-VI, LNDFH-II, F-pLNH I
<i>Mg791</i>	<i>AEQ33441.1</i>	α -1,3-fucosyl-transferase	2'-FL, 3-FL, DFL, LNFP-I, LNFP-III, LNFP-V, LNFP-VI, LNDFH-II, F-pLNH I

<i>Moumou_00703</i>	<i>YP_007354660</i>	α -1,3-fucosyl-transferase	2'-FL, 3-FL, DFL, LNFP-I, LNFP-III, LNFP-V, LNFP-VI, LNDFH-II, F-pLNH I
<i>futA</i>	<i>NP_207177.1</i>	α -1,3-fucosyl-transferase	2'-FL, 3-FL, DFL, LNFP-I, LNFP-III, LNFP-V, LNFP-VI, LNDFH-II, F-pLNH I
<i>fucT</i>	<i>AAB81031.1</i>	α -1,3-fucosyl-transferase	2'-FL, 3-FL, DFL, LNFP-I, LNFP-III, LNFP-V, LNFP-VI, LNDFH-II, F-pLNH I
<i>fucTIII</i>	<i>AY450598.1</i>	α -1,4-fucosyl-transferase	LNDFH-I, LNDFH-II
<i>fucTa</i>	<i>AF194963.1</i>	α -1,3/4-fucosyl-transferase	LNFP-II, LNDFH-I, LNDFH-II
<i>Pd2,6ST</i>	<i>BAA25316.1</i>	α -2,6-sialyltransferase	6'-SL
<i>PspST6</i>	<i>BAF92026.1</i>	α -2,6-sialyltransferase	6'-SL
<i>PiST6_145</i>	<i>BAF91416.1</i>	α -2,6-sialyltransferase	6'-SL
<i>PiST6_119</i>	<i>BAI49484.1</i>	α -2,6-sialyltransferase	6'-SL
<i>NST</i>	<i>AAC44541.1</i>	α -2,3-sialyltransferase	3'-SL

An aspect of the present invention is the provision of a nucleic acid construct comprising a heterologous nucleic acid sequence(s) encoding a protein capable of sugar transportation which is a major facilitator superfamily (MFS) protein as shown in SEQ ID NO: 1, or a functional homologue thereof which amino acid sequence is at least 80 % identical to SEQ ID NO: 1, wherein the nucleic acid sequence encoding the MFS protein has at least 70 % sequence identity to SEQ ID NO: 2.

By the term "heterologous nucleic acid sequence", "recombinant gene/nucleic acid/DNA encoding" or "coding nucleic acid sequence" is meant an artificial nucleic acid sequence (i.e. produced *in vitro* using standard laboratory methods for making nucleic acid sequences) that comprises a set of consecutive, non-overlapping triplets (codons) which is transcribed into mRNA and translated into a polypeptide when placed under the control of the appropriate control sequences, i.e. promoter. The boundaries of the coding sequence are generally determined by a ribosome binding site located just upstream of the open reading frame at the 5' end of the mRNA, a translational start codon (AUG, GUG or UUG), and a translational stop codon (UAA, UGA or UAG). A coding sequence can include, but is not limited to, genomic DNA, cDNA, synthetic, and recombinant nucleic acid sequences. The term "nucleic acid" includes RNA, DNA and cDNA molecules. It is understood that, as a result of the degeneracy of the genetic code, a multitude of nucleotide sequences

encoding a given protein may be produced. The term nucleic acid is used interchangeably with the term "polynucleotide". An "oligonucleotide" is a short chain nucleic acid molecule. "Primer" is an oligonucleotide, whether occurring naturally as in a purified restriction digest or produced synthetically, which is capable of acting as
5 a point of initiation of synthesis when placed under conditions in which synthesis of a primer extension product which is complementary to a nucleic acid strand is induced, (i.e., in the presence of nucleotides and an inducing agent such as DNA polymerase and at a suitable temperature and pH). The primer is preferably single stranded for maximum efficiency in amplification but may alternatively be double stranded. If
10 double stranded, the primer is first treated to separate its strands before being used to prepare extension products. Preferably, the primer is a deoxyribonucleotide. The primer must be sufficiently long to prime the synthesis of extension products in the presence of the inducing agent. The exact lengths of the primers will depend on many factors, including temperature, source of primer and the use of the method.

15 The recombinant nucleic sequence of the invention may be a coding DNA sequence, e.g. a gene, or non-coding DNA sequence, e.g. a regulatory DNA, such as a promoter sequence. One aspect of the invention relates to providing a recombinant cell comprising recombinant DNA sequences encoding enzymes necessary for the production of one or more HMOs and a DNA sequence encoding
20 the YberC transporter. Accordingly, in one embodiment the invention relates to a nucleic acid construct comprising a coding nucleic sequence, i.e. recombinant DNA sequence of a gene of interest, e.g. a glycosyltransferase gene or the *yberC* gene, and a non-coding DNA sequence, e.g. a promoter DNA sequence, e.g. a recombinant promoter sequence derived from the promoter of the *lac* operon or an
25 *g/p* operon, or a promoter sequence derived from another genomic promoter DNA sequence, or a synthetic promoter sequence, wherein the coding and promoter sequences are operably linked. The term "operably linked" refers to a functional relationship between two or more nucleic acid (e.g., DNA) segments. Typically, it refers to the functional relationship of a transcriptional regulatory sequence to a
30 transcribed sequence. For example, a promoter sequence is operably linked to a coding sequence if it stimulates or modulates the transcription of the coding sequence in an appropriate host cell or other expression system. Generally, promoter transcriptional regulatory sequences that are operably linked to a transcribed sequence are physically contiguous to the transcribed sequence, i.e.,
35 they are cis-acting.

In one embodiment, the nucleic acid construct of the invention may be a

part of the vector DNA, in another embodiment the construct is an expression cassette/cartridge that is integrated in the genome of a host cell. Accordingly, the term "nucleic acid construct" means an artificially constructed segment of nucleic acid, in particular a DNA segment, which is intended to be 'transplanted' into a target cell, e.g. a bacterial cell, to modify the expression of a gene or express a gene/coding DNA sequence which may be included in the construct. In the context of the invention, the nucleic acid construct contains a recombinant DNA sequence comprising two or more recombinant DNA sequences: essentially, a non-coding DNA sequence comprising a promoter DNA sequence and a coding DNA sequence encoding a protein of interest, e.g. the YberC protein, a glycosyltransferase, of another protein useful for production of an HMO in a host cell. Preferably, the construct comprises further non-coding DNA sequences that either regulate transcription or translation of the coding DNA of the construct, e.g. a DNA sequence facilitating ribosome binding to the transcript, a leading DNA sequence that stabilize the transcript, or transcription termination sequence.

Integration of the recombinant nucleic acid of interest comprised in the construct (expression cassette) into the bacterial genome can be achieved by conventional methods, e.g. by using linear cartridges that contain flanking sequences homologous to a specific site on the chromosome, as described for the attTn7-site (Waddell C.S. and Craig N.L., *Genes Dev.* (1988) Feb;2(2):137-49.); methods for genomic integration of nucleic acid sequences in which recombination is mediated by the Red recombinase function of the phage λ or the RecE/RecT recombinase function of the Rac prophage (Murphy, *J Bacteriol.* (1998);180(8):2063-7; Zhang *et al.*, *Nature Genetics* (1998) 20: 123-128 Muyrers *et al.*, *EMBO Rep.* (2000) 1(3): 239-243); methods based on Red/ET recombination (Wenzel *et al.*, *Chem Biol.* (2005), 12(3):349-56.; Vetcher *et al.*, *Appl Environ Microbiol.* (2005);71(4):1829-35); or positive clones, i.e. clones that carry the expression cassette, can be selected e.g. by means of a marker gene, or loss or gain of gene function.

A single copy of the expression cassette comprising a gene of interest may be sufficient to secure production of a desired HMO and achieve the desired effects according to the invention. Accordingly, in some preferred embodiments, the invention relates to a recombinant HMO producing cell that comprises one, two or three copies of a gene of interest integrated in the genomic DNA of the cell. In some embodiments the single copy of a gene is preferred.

In one preferred embodiment, recombinant coding nucleic acid sequence of

the nucleic acid construct of the invention is heterologous with respect to the promoter, which means that in the equivalent native coding sequence in the genome of species of origin is transcribed under control of another promoter sequence (i.e. not the promoter sequence of the construct). Still, with respect to the host cell, the coding DNA may be either heterologous (i.e. derived from another biological species or genus), such as e.g. the DNA sequence encoding the YberC protein expressed in *Escherichia coli* host cells, or homologous (i.e. derived from the host cell), such as e.g. genes of the colonic acid operon, the *GMAB* genes.

The term, a "regulatory element" or "promoter" or "promoter region" or "promoter element" is a nucleic acid sequence that is recognized and bound by a DNA dependent RNA polymerase during initiation of transcription. The promoter, together with other transcriptional and translational regulatory nucleic acid sequences (also termed "control sequences") is necessary to express a given gene or group of genes (an operon). In general, the transcriptional and translational regulatory sequences include, but are not limited to, promoter sequences, ribosomal binding sites, transcriptional start and stop sequences, translational start and stop sequences, and transcription enhancer or activator sequences. The "transcription start site" means the first nucleotide to be transcribed and is designated +1. Nucleotides downstream of the start site are numbered +2, +3, +4 etc., and nucleotides in the 5' opposite (upstream) direction are numbered -1, -2, -3 etc. The promoter DNA sequence of the construct can derive from a promoter region of any gene of the genome of a selected species, preferably, a promoter region of the genomic DNA of *E. coli*. Accordingly, any promoter DNA sequence that is able to bind to an RNA polymerase and initiate transcription is suitable for practicing the invention. In principle, any promoter DNA sequence can be used to control transcription of the recombinant nucleic acid of interest of the construct, different or same promoter sequences may be used to drive transcription of different genes of interest integrated in the genome of the host cell or in expression vector DNA. To have an optimal expression of the recombinant genes included in the construct, the construct may comprise further regulatory sequences, e.g. a leading DNA sequence, such as a DNA sequence derived from 5'-untranslated region (5'UTR) of a *glp* gene of *E. coli*, a sequence for ribosomal binding. Examples of the later sequences are described in WO2019123324 (incorporated herein by reference) and illustrated in non-limiting working examples herein.

In some preferred embodiments, the regulatory element for the regulation of the expression of a recombinant gene included in the construct of the invention is

glpFKX operon promoter, *PglpF*, in other preferred embodiments, the promoter is *lac* operon promoter, *Plac*.

In a further aspect, the regulatory element for the regulation of the expression of a recombinant gene included in the construct of the invention is the *mgIBAC*;
5 galactose/methyl-galactosidase transporter promoter *PmgIB* or variants thereof such as but not limited to *PmgIB_70UTR* of SEQ ID NO: 15, or *PmgIB_70UTR_SD4* of SEQ ID NO: 16.

In a further aspect, the regulatory element for the regulation of the expression of a recombinant gene included in the construct of the invention is the
10 *gatYZABCD*; tagatose-1,6-bisP aldolase promoter *PgatY* or variants thereof such as but not limited to *PgatY_U70UTR* of SEQ ID NO: 17.

The preferred regulatory element present in a genetically modified cell and/or in a nucleic acid construct of the present invention is selected from the group consisting of *PgatY_70UTR*, *PglpF*, *PglpF_SD1*, *PglpF_SD10*, *PglpF_SD2*,
15 *PglpF_SD3*, *PglpF_SD4*, *PglpF_SD5*, *PglpF_SD6*, *PglpF_SD7*, *PglpF_SD8*, *PglpF_SD9*, *Plac_16UTR*, *Plac*, *PmgIB_70UTR* and *PmgIB_70UTR_SD4*.

An especially preferred regulatory element present in a genetically modified cell and/or in a nucleic acid construct of the present invention is selected from the group consisting of *PglpF* and *Plac*.

20 In some preferred embodiments, the regulatory element for the regulation of the expression of a recombinant gene included in the construct of the invention is *glpFKX* operon promoter, *PglpF*, in other preferred embodiments, the promoter is *lac* operon promoter, *Plac*. However, any promoter enabling transcription and/or regulation of the level of transcription of one or more recombinant nucleic acids that
25 encode one or more proteins (or one or more regulatory nucleic acids) that are either necessary or beneficial to achieve an optimal level of biosynthetic production of one or more HMOs in the host cell, e.g. glucosyltransferases, proteins involved in the transmembrane transport of an HMO or HMO precursor, gene expression regulatory proteins, etc, and allowing to achieve the desired effects according to the invention is
30 suitable for practicing the invention.

Preferably, the construct of the invention comprising a gene related to the biosynthetic production of an HMO, a promoter DNA sequence, and other regulatory sequences, such as a ribosomal binding site sequence (e.g. Shine-Dalgarno sequence), expressed in the host cell enables the production of the HMO at the level
35 of at least 0,03 g/OD (optical density) of 1 liter of the fermentation media comprising

a suspension of host cells, e.g., at the level of around 0.05 g/l/OD, around 0,1 g/l/OD, or more. For the purposes of the invention, the latter level of HMO production is regarded as "sufficient" or "optimal" and the host cell capable of producing this level of a desired HMO is regarded as "suitable host cell", i.e. the cell can be further
 5 modified to express the HMO transporter protein, e.g. YberC, to achieve at least one effect described herein that is advantageous for the HMO production.

The genetically modified cell (host cell or recombinant cell) may be e.g. a bacterial or yeast cell. In one preferred embodiment, the genetically modified cell is
 10 bacterial cell. Regarding the bacterial host cells, there are, in principle, no limitations; they may be eubacteria (gram-positive or gram-negative) or archaeobacteria, as long as they allow genetic manipulation for insertion of a gene of interest and can be cultivated on a manufacturing scale. Preferably, the host cell has the property to allow cultivation to high cell densities. Non-limiting examples of bacterial host cells
 15 that are suitable for recombinant industrial production of an HMO(s) according to the invention could be *Erwinia herbicola* (*Pantoea agglomerans*), *Citrobacter freundii*, *Pantoea citrea*, *Pectobacterium carotovorum*, or *Xanthomonas campestris*. Bacteria of the genus *Bacillus* may also be used, including *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus coagulans*, *Bacillus thermophilus*, *Bacillus laterosporus*,
 20 *Bacillus megaterium*, *Bacillus mycoides*, *Bacillus pumilus*, *Bacillus lentus*, *Bacillus cereus*, and *Bacillus circulans*. Similarly, bacteria of the genera *Lactobacillus* and *Lactococcus* may be modified using the methods of this invention, including but not limited to *Lactobacillus acidophilus*, *Lactobacillus salivarius*, *Lactobacillus plantarum*, *Lactobacillus helveticus*, *Lactobacillus delbrueckii*, *Lactobacillus rhamnosus*,
 25 *Lactobacillus bulgaricus*, *Lactobacillus crispatus*, *Lactobacillus gasseri*, *Lactobacillus casei*, *Lactobacillus reuteri*, *Lactobacillus jensenii*, and *Lactococcus lactis*. *Streptococcus thermophiles* and *Propionibacterium freudenreichii* are also suitable bacterial species for the invention described herein. Also included as part of this invention are strains, modified as described here, from the genera *Enterococcus*
 30 (e.g., *Enterococcus faecium* and *Enterococcus thermophiles*), *Bifidobacterium* (e.g., *Bifidobacterium longum*, *Bifidobacterium infantis*, and *Bifidobacterium bifidum*), *Sporolactobacillus* spp., *Micromonospora* spp., *Micrococcus* spp., *Rhodococcus* spp., and *Pseudomonas* (e.g., *Pseudomonas fluorescens* and *Pseudomonas aeruginosa*). Bacteria comprising the characteristics described herein are cultured in
 35 the presence of lactose, and an oligosaccharide, such as an HMO, produced by the cell is retrieved, either from the bacterium itself or from a culture supernatant of the

bacterium. In one preferred embodiment, the genetically modified cell of the invention is an *Escherichia coli* cell.

In another preferred embodiment, the host cell is a yeast cell e.g. *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Pichia pastoris*,

5 *Kluyveromyces lactis*, *Kluyveromyces marxianus*, etc.

The HMOs produced by recombinant cells of the invention may be purified using a suitable procedure available in the art (e.g. such as described in WO2015188834, WO2017182965 or WO2017152918).

Genetically modified cells of the invention can be provided using standard
10 methods of the art e.g. those described in the manuals by Sambrook *et al.*, Wilson & Walker, "Maniatis *et al.*", and Ausubel *et al.*

A host cell suitable for the HMO production, e.g. *E. coli*, may comprise an endogenous β -galactosidase gene or an exogenous β -galactosidase gene, e.g. *E. coli* comprises an endogenous *lacZ* gene (e.g., GenBank Accession Number V00296
15 (GI:41901)). For the purposes of the invention, an HMO-producing cell is genetically manipulated to either comprise any β -galactosidase gene or to comprise an inactivated version of the gene. The gene may be inactivated by a complete or partial deletion of the corresponding nucleic acid sequence from the bacterial genome, or the gene sequence is mutated in the way that it is transcribed, or, if
20 transcribed, the transcript is not translated or if translated to a protein (i.e. β -galactosidase), the protein does not have the corresponding enzymatic activity. In this way the HMO-producing cell accumulates an increased intracellular lactose pool which is beneficial for the production of HMOs.

In some embodiments, the engendered cell contains a deficient sialic acid
25 catabolic pathway. By "sialic acid catabolic pathway" is meant a sequence of reactions, usually controlled and catalyzed by enzymes, which results in the degradation of sialic acid. An exemplary sialic acid catabolic pathway described herein is the *E. coli* pathway. In this pathway, sialic acid (Neu5Ac; N-acetylneuraminic acid) is degraded by the enzymes NanA (N-acetylneuraminic acid
30 lyase) and NanK (N-acetylmannosamine kinase) and NanE (N-acetylmannosamine-6-phosphate epimerase), all encoded from the *nanATEK-yhcH* operon, and repressed by NanR (<http://ecocyc.org/ECOLI>). A deficient sialic acid catabolic pathway is rendered in the *E. coli* host by introducing a mutation in the endogenous *nanA* (N-acetylneuraminic acid lyase) (e.g., GenBank Accession Number
35 D00067.1(GL216588)) and/or *nanK* (N-acetylmannosamine kinase) genes (e.g., GenBank Accession Number (amino acid) BAE77265.1 (GL85676015)), and/or *nanE*

(N-acetylmannosamine-6-phosphate epimerase, GI: 947745, incorporated herein by reference). Optionally, the *nanT* (N-acetylneuraminate transporter) gene is also inactivated or mutated. Other intermediates of sialic acid metabolism include: (ManNAc-6-P) N-acetylmannosamine-6-phosphate; (GlcNAc-6-P) N-

5 acetylglucosamine-6-phosphate; (GlcN-6-P) Glucosamine-6-phosphate, and (Fruc-6-P) Fructose-6-phosphate. In some preferred embodiments, *nanA* is mutated. In other preferred embodiments, *nanA* and *nanK* are mutated, while *nanE* remains functional. In another preferred embodiment, *nanA* and *nanE* are mutated, while *nanK* has not been mutated, inactivated or deleted. A mutation is one or more changes in the

10 nucleic acid sequence coding the gene product of *nanA*, *nanK*, *nanE*, and/or *nanT*. For example, the mutation may be 1, 2, up to 5, up to 10, up to 25, up to 50 or up to 100 changes in the nucleic acid sequence. For example, the *nanA*, *nanK*, *nanE*, and/or *nanT* genes are mutated by a null mutation. Null mutations as described herein encompass amino acid substitutions, additions, deletions, or insertions, which

15 either cause a loss of function of the enzyme (i.e. reduced or no activity) or loss of the enzyme (i.e. no gene product). By "deleted" is meant that the coding region is removed completely or in part such that no (functional) gene product is produced. By inactivated is meant that the coding sequence has been altered such that the resulting gene product is functionally inactive or encodes for a gene product with less

20 than 100 %, e.g. 90 %, 80 %, 70 %, 60 %, 50 %, 40 %, 30 % or 20 % of the activity of the native, naturally occurring, endogenous gene product. A "not mutated" gene or protein does not differ from a native, naturally-occurring, or endogenous coding sequence by 1, 2, up to 5, up to 10, up to 20, up to 50, up to 100, up to 200 or up to 500 or more codons, or to the corresponding encoded amino acid sequence.

25 Furthermore, the host cell (e.g., *E. coli*) also comprises a sialic acid synthetic capability. For example, the bacterium comprises a sialic acid synthetic capability through provision of an exogenous UDP-GlcNAc 2-epimerase (e.g., *neuC* of *Campylobacter jejuni* (GenBank AAK91727.1; GI:15193223) or equivalent (e.g. *neuC* of *E.coli* S88 (GenBank YP_002392936.1; GI: 218560023), a Neu5Ac

30 synthase (e.g., *neuB* of *C. jejuni* (GenBank AAK91726.1; GI:15193222) or equivalent, (e.g. *Flavobacterium limnosediminis* sialic acid synthase, GenBank GL559220424), and/or a CMP-Neu5Ac synthetase (e.g., *neuA* of *C. jejuni* (GenBank AAK91728.1; GI:15193224) or equivalent, (e.g. *Vibrio brasiliensis* CMP-sialic acid synthase, GenBank GI: 493937153).

Production of neutral N-acetylglucosamine-containing HMOs in engineered bacteria and yeasts is known in the art (see e.g. Gebus C *et al.* (2012) Carbohydrate Research 363 83-90; US10,519,475).

For the production of N-acetylglucosamine-containing HMOs, such as

5 Lacto-N-triose 2 (LNT-2), Lacto-N-tetraose (LNT), Lacto-N-neotetraose (LNnT), Lacto-N-fucopentaose I (LNFP-I), Lacto-N-fucopentaose II (LNFP-II), Lacto-N-fucopentaose III (LNFP-III), Lacto-N-fucopentaose V (LNFP-V), Lacto-N-difucohexaose I (LDFH-I), Lacto-N-difucohexaose II (LDFH-II), and Lacto-N-neodifucohexaose II (LNDFH-III), the bacterium comprises a functional *lacY* and a

10 dysfunctional *lacZ* gene, as described above, and it is engineered to comprise an exogenous UDP-GlcNAc:Gal α / β -R β -3-N-acetylglucosaminyltransferase gene, or a functional variant or fragment thereof. This exogenous UDP-GlcNAc:Gal α / β -R β -3-N-acetylglucosaminyltransferase gene may be obtained from any one of a number of sources, (see Table 1), e.g., the *lgtA* gene described from *N. meningitidis* (Genbank

15 protein Accession AAF42258.1) or *N. gonorrhoeae* (Genbank protein Accession ACF31229.1). Optionally, an additional exogenous glycosyltransferase gene may be co-expressed in the bacterium comprising an exogenous UDP-GlcNAc:Gal α / β -R β -3-N-acetylglucosaminyltransferase. For example, a β -1,4-galactosyltransferase ((Gal4T) gene is co-expressed with the UDP-GlcNAc:Gal α / β -R β -3-N-

20 acetylglucosaminyltransferase gene. This exogenous β -1,4-galactosyltransferase gene can be obtained from any one of a number of sources, e.g., the one described from *N. meningitidis*, the *lgtB* gene (Genbank protein Accession AAF42257.1), or from *H. pylori*, the *HP0826/galT* gene (Genbank protein Accession NP_207619.1). Optionally, the additional exogenous glycosyltransferase gene co-expressed in the

25 bacterium comprising an exogenous UDP-GlcNAc:Gal α / β -R β -3-N-acetylglucosaminyltransferase (Gal3T) gene is a P-1,3-galactosyltransferase gene, e.g., that described from *E. coli* 055:H7, the *wbgO* gene (Genbank protein Accession WP_000582563.1), or from *H. pylori*, the *jhp0563* gene (Genbank protein Accession AEZ55696.1), or from *Streptococcus agalactiae* type *Ib* O12 the *cps/BJ* gene

30 Genbank protein Accession AB050723). Functional variants and fragments of any of the enzymes described above are also encompassed by the disclosed invention.

A N-acetylglucosaminyltransferase gene and/or a galactosyltransferase gene, can also be operably linked to a *Pglp* and be expressed from the

corresponding genome-integrated cassette. In one embodiment, the gene that is

35 genome integrated is a gene encoding for a galactosyltransferase, e.g. *HP0826* gene encoding for the GalT enzyme from *H. pylori* (Genbank protein Accession

NP_207619.1); in another embodiment, the gene that is genome integrated is a gene encoding a β -1,3-N-acetylglucosaminyltransferase, e.g. *lgtA* gene from *N.*

meningitidis (Genbank protein Accession AAF42258.1). In these embodiments, the second gene, i.e. a gene encoding a β -1,3-N-acetylglucosaminyltransferase or

- 5 galactosyltransferase, correspondingly, may either be expressed from a genome-integrated or plasmid-borne cassette. The second gene may optionally be expressed either under the control of a *glp* promoter or under the control of any other promoter suitable for the expression system, e.g. *Plac*.

- Unless otherwise specified, all technical and scientific terms used herein
- 10 have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Singleton *et al.* (1994) Dictionary of Microbiology and Molecular Biology, second edition, John Wiley and Sons (New York) provides one of skill with a general dictionary of many of the terms used in this invention. Although any methods and materials similar or equivalent to those described herein can be
- 15 used in the practice or testing of the present invention, the preferred methods and materials are described. Most of the nomenclature and general laboratory procedures required in this application can be found in Sambrook *et al.*, Molecular Cloning: A Laboratory Manual, Vol. 1-3, Cold Spring Harbor Laboratory, Cold Spring Harbor, New York (2012); Wilson K. and Walker J., Principles and Techniques of
- 20 Biochemistry and Molecular Biology (2010), Cambridge University Press; or in Maniatis *et al.*, Molecular Cloning A laboratory Manual, Cold Spring Harbor Laboratory (2012); or in Ausubel *et al.*, Current protocols in molecular biology, John Wiley and Sohns (2010). The manuals are hereinafter referred to as "Sambrook *et al.*", "*Wilson & Walker*", "*Maniatis et al.*", "*Ausubel et al.*", correspondingly.

25

A second aspect of the invention related to a method for the production of one or more HMOs, the method comprising the steps of:

- (i) providing a genetically modified cell capable of producing an HMO, wherein said cell comprises a recombinant
- 30 nucleic acid encoding a protein of SEQ ID NO: 1, or a functional homologue thereof having amino acid sequence that is at least 80 % identical, preferably at least 85 % identical, more preferably at least 90 % identical to SEQ ID NO: 1;
- (ii) culturing the cell of (i) in a suitable cell culture medium
- 35 to express said recombinant nucleic acid;
- (iii) harvesting the one or more HMO(s) produced in step

(ii).

According to the invention, the term “culturing” (or “cultivating” or “cultivation”, also termed “fermentation”) relates to the propagation of bacterial expression cells in a controlled bioreactor according to methods known in the industry.

To produce one or more HMOs, the HMO-producing bacteria as described herein are cultivated according to the procedures known in the art in the presence of a suitable carbon source, e.g. glucose, glycerol, lactose, etc., and the produced HMO is harvested from the cultivation media and the microbial biomass formed during the cultivation process. Thereafter, the HMOs are purified according to the procedures known in the art, e.g. such as described in WO2015188834, WO2017182965 or WO2017152918, and the purified HMOs are used as nutraceuticals, pharmaceuticals, or for any other purpose, e.g. for research. Manufacturing of HMOs is typically accomplished by performing cultivation in larger volumes. The term “manufacturing” and “manufacturing scale” in the meaning of the invention defines a fermentation with a minimum volume of 5 L culture broth. Usually, a “manufacturing scale” process is defined by being capable of processing large volumes of a preparation containing the HMO or HMOs of interest and yielding amounts of the protein of interest that meet, e.g. in the case of a therapeutic compound or composition, the demands for clinical trials as well as for market supply. In addition to the large volume, a manufacturing scale method, as opposed to simple lab scale methods like shake flask cultivation, is characterized by the use of the technical system of a bioreactor (fermenter) which is equipped with devices for agitation, aeration, nutrient feeding, monitoring and control of process parameters (pH, temperature, dissolved oxygen tension, back pressure, etc.). To a large extent, the behavior of an expression system in a lab scale method, such as shake flasks, benchtop bioreactors or the deep well format described in the examples of the disclosure, does allow to predict the behavior of that system in the complex environment of a bioreactor.

With regard to the suitable cultivation medium used in the fermentation process, there are no limitations. The culture medium may be semi-defined, i.e. containing complex media compounds (e.g. yeast extract, soy peptone, casamino acids, etc.), or it may be chemically defined, without any complex compounds.

By the term “one or more HMOs” is meant that an HMO production cell may

be able to produce a single HMO structure (a first HMO) or multiple HMO structures (a second, a third, etc. HMO). In some embodiments, it may be preferred a host cell that produces a single HMO, in other preferred embodiments, a host cell producing multiple HMO structures may be preferred. Non-limiting examples for host cells

5 producing single HMO structures are 3'-SL, 6'-SL or LNT-2 producing cells. Non-limiting examples of host cells capable of producing multiple HMO structures can be FSL, LNT, LNnT, LNFP I, LNFP II, LNFP III, LNFP IV, LNFP V, pLNnH, pLNH-II producing cells.

The term "harvesting" in the context in the invention relates to collecting the

10 produced HMO(s) following the termination of fermentation. In different embodiments it may include collecting the HMO(s) included in both the biomass (i.e. the host cells) and cultivation media, i.e. before/without separation of the fermentation broth from the biomass. In other embodiments the produced HMOs may be collected separately from the biomass and fermentation broth, i.e. after/following the separation of

15 biomass from cultivation media (i.e. fermentation broth). The separation of cells from the medium can be carried out with any of the methods well known to the skilled person in the art, such as any suitable type of centrifugation or filtration. The separation of cells from the medium can follow immediately after harvesting the fermentation broth or be carried out at a later stage after storing the fermentation

20 broth at appropriate conditions. Recovery of the produced HMO(s) from the remaining biomass (or total fermentation) include extraction thereof from the biomass (i.e the production cells). It can be done by any suitable methods of the art, e.g. by sonication, boiling, homogenization, enzymatic lysis using lysozyme, or freezing and grinding.

25 After recovery from fermentation, HMO(s) are available for further processing and purification.

Purification of HMOs produced by fermentation can be done using a suitable procedure described in WO2016095924, WO2015188834, WO2017152918, WO2017182965, US20190119314 (all incorporated by reference).

30 In some embodiments of the invention, a host cell may produce several HMOs, wherein one HMO is the "product" HMO and some/all the other HMOs are "by-product" HMOs. Typically, by-product HMOs are either the major HMO precursors or products of further modification of the major HMO. In some embodiments, it may be desired to produce the product HMO in abundant amounts

35 and by-product HMOs in minor amounts. Cells and methods for HMO production described herein allow for controlled production of an HMO product with a defined

HMO profile, e.g. in one embodiment, the produced HMO mixture wherein the product HMO is a dominating HMO compared to the other HMOs (i.e. by-product HMOs) of the mixture, i.e. the product HMO is produced in higher amounts than other by-product HMOs; in other embodiments, the cell producing the same HMO-mixture may be tuned to produce one or more by-product HMOs in higher amount than product HMO. For example, during the production of LNnT, the product HMO, often a significant amount of pLNnH, the by-product HMO, is produced. With the genetically modified cells of the present invention the level of pLNnH in the LNnT product can be significantly reduced.

Advantageously, the invention provides both a decreased ratio of by-product to product and an increased overall yield of the product (and/or HMOs in total). This, less by-product formation in relation to product formation facilitates an elevated product production and increases efficiency of both the production and product recovery process, providing superior manufacturing procedure of HMOs.

In different preferred embodiments, different host cells producing either/both 3'-SL, 6'-SL, LNT-2, , FSL, LNT, LNnT, DFL, FSL, LNT, LNnT, LNFP I, LNFP II, LNFP III, LNFP IV, LNFP V, pLNnH, pLNH-II, as the product or by-product HMO, may be selected. In one preferred embodiment, the product is LNT-2 and by-product is pLNnH. In another preferred embodiment, the product is LNnT and by-product is pLNnH. In another preferred embodiment, the product is LNT and by-product is pLNH-II. In another preferred embodiment, the product is LNT2 and by-product is pLNH-II.

The invention also relates to the use of a genetically modified cell and/or a nucleic acid construct according to the invention, for the production of one or more oligosaccharides, preferably one or more human milk oligosaccharide(s).

In one embodiment, the genetically modified cell and/or the nucleic acid construct according to the invention is used in the production of a specific HMO selected from the group consisting of 3'-sialyllactose (3'-SL), 6'-sialyllactose (6'-SL), Lacto-N-Triose-2 (LNT-2), Lacto-N-tetraose (LNT), Lacto-N-neotetraose (LNnT), Lacto-N-fucopentaose I (LNFP I), Lacto-N-fucopentaose II (LNFP II), Lacto-N-fucopentaose III (LNFP III), Lacto-N-fucopentaose IV (LNFP IV), and Lacto-N-fucopentaose V (LNFP V), para-lacto-N-neohexaose (pLNnH), and para-lacto-neohexaose II (pLNH-II).

In a preferred embodiment the genetically modified cell and/or the nucleic acid construct according to the invention is used in the production of a specific HMO selected from the group consisting of Lacto-N-Triose-2 (LNT-2), Lacto-N-tetraose

(LNT), Lacto-N-neotetraose (LNnT), Lacto-N-fucopentaose I (LNFP I), Lacto-N-fucopentaose II (LNFP II), Lacto-N-fucopentaose III (LNFP III), Lacto-N-fucopentaose IV (LNFP IV), and Lacto-N-fucopentaose V (LNFP V), para-lacto-N-neohexaose (pLNnH), and para-lacto-neohexaose II (pLNH-II).

5 In a further preferred embodiment, the genetically modified cell and/or the nucleic acid construct according to the invention is used in the production of an HMO selected from the group consisting of Lacto-N-Triose-2 (LNT-2), Lacto-N-tetraose (LNT), Lacto-N-neotetraose (LNnT) and para-lacto-neohexaose II (pLNH-II).

10 In an especially preferred embodiment the genetically modified cell or the nucleic acid construct according to the invention is used in the production of a specific HMO selected from the group consisting of Lacto-N-Triose-2 (LNT-2), Lacto-N-tetraose (LNT) and Lacto-N-neotetraose (LNnT).

The invention is further illustrated by non-limiting examples and
15 embodiments below.

EXAMPLES

Materials and Methods

20 Unless otherwise noted, standard techniques, vectors, control sequence elements, and other expression system elements known in the field of molecular biology are used for nucleic acid manipulation, transformation, and expression. Such standard techniques, vectors, and elements can be found, for example, in: Ausubel *et al.* (eds.), *Current Protocols in Molecular Biology* (1995) (John Wiley & Sons);
25 Sambrook, Fritsch, & Maniatis (eds.), *Molecular Cloning* (1989) (Cold Spring Harbor Laboratory Press, NY); Berger & Kimmel, *Methods in Enzymology* 152: Guide to Molecular Cloning Techniques (1987) (Academic Press); Bukhari *et al.* (eds.), *DNA Insertion Elements, Plasmids and Episomes* (1977) (Cold Spring Harbor Laboratory Press, NY); Miller, J.H. *Experiments in molecular genetics* (1972.) (Cold spring
30 Harbor Laboratory Press, NY)

The embodiments described below are selected to illustrate the invention and are not limiting the invention in any way.

Strains

The bacterial strain used, MDO, was constructed from *Escherichia coli* K12 DH1. The *E. coli* K12 DH1 genotype is: F^- , λ^- , *gyrA96*, *recA1*, *relA1*, *endA1*, *thi-1*, *hsdR17*, *supE44*. Strains utilized in the present Examples are described in Table 2.

5 **Table 2**

Strain IDs	Product	Relevant Genotype
DH1	-	<i>F^-</i> λ^- <i>endA1</i> <i>recA1</i> <i>relA1</i> <i>gyrA96</i> <i>thi-1</i> <i>glnV44</i> <i>hsdR17</i> (<i>r_K m_K</i>)
MDO	-	<i>E. coli</i> DH1 Δ <i>lacZ</i> , Δ <i>lacA</i> , Δ <i>nanKETA</i> , Δ <i>melA</i> , Δ <i>wcaJ</i> , Δ <i>mdoH</i>
MP3941	LNT-2	<i>MDO PglpF-GlcNAcT</i>
MP3940		<i>MDO PglpF-GlcNAcT PglpF-yberC0001_9420</i>
MP4436	LNnT	<i>MDO PglpF-GlcNAcT PglpF-Gal4T</i>
MP3863		<i>MDO PglpF-GlcNAcT PglpF-Gal4T Plac-yberC0001_9420</i>
MP4473	LNT	<i>MDO PglpF-GlcNAcT PglpF-Gal3T</i>
MP4462		<i>MDO PglpF-GlcNAcT PglpF-Gal3T Plac-yberC0001_9420</i>

Media

The Luria Broth (LB) medium was made using LB Broth Powder, Millers
 10 (Fisher Scientific) and LB agar plates were made using LB Agar Powder, Millers
 (Fisher Scientific). When appropriated ampicillin ((100 μ g/mL) or any appropriate
 antibiotic), and/or chloramphenicol (20 μ g/ml) was added.

Basal Minimal medium had the following composition: NaOH (1 g/L), KOH
 (2.5 g/L), KH_2PO_4 (7 g/L), $\text{NH}_4\text{H}_2\text{PO}_4$ (7 g/L), Citric acid (0.5 g/l), Trace mineral
 15 solution (5 mL/L). The trace mineral stock solution contained: $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$ 0.82 g/L,
 Citric acid 20 g/L, $\text{MnSO}_4 \cdot \text{H}_2\text{O}$ 0.98 g/L, $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$ 3.925 g/L, $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$ 0.2
 g/L. The pH of the Basal Minimal Medium was adjusted to 7.0 with 5 N NaOH and
 autoclaved. Before inoculation the Basal Minimal medium was supplied with 1 mM
 MgSO_4 , 4 μ g/ml thiamin, 0.5 % of a given carbon source (glycerol (Carbosynth)), and
 20 when appropriate Isopropyl- β -D-Thiogalactoside (IPTG) (0.2 mM) was added.
 Thiamin, antibiotics, and IPTG were sterilized by filtration. All percentage
 concentrations for glycerol are expressed as v/v and for glucose as w/v.

M9 plates containing 2-deoxy-galactose had the following composition: 15
 g/L agar (Fisher Scientific), 2.26 g/L 5x M9 Minimal Salt (Sigma-Aldrich), 2 mM
 25 MgSO_4 , 4 μ g/mL thiamine, 0.2 % glycerol, and 0.2 % 2-deoxy-D-galactose
 (Carbosynth).

MacConkey indicator plates had the following composition: 40 g/L
 MacConkey agar Base (BD Difco™) and a carbon source at a final concentration of 1
 %.

Cultivation

Unless otherwise noted, *E. coli* strains were propagated in Luria-Bertani (LB) medium containing 0.2 % glucose at 37°C with agitation. Agar plates were
5 incubated at 37°C overnight.

Chemical competent cells and transformations

E. coli was inoculated from LB plates in 5 mL LB containing 0.2 % glucose at 37°C with shaking until OD₆₀₀ ~0.4. 2 mL culture was harvested by centrifugation
10 for 25 seconds at 13000 g. The supernatant was removed, and the cell pellet resuspended in 600 µL cold TB solutions (10 mM PIPES, 15 mM CaCl₂, 250 mM KCl). The cells were incubated on ice for 20 minutes followed by pelleting for 15 seconds at 13000 g. The supernatant was removed, and the cell pellet resuspended in 100 µL cold TB solution. Transformation of plasmids were done using 100 µL
15 competent cells and 1 to 10 ng plasmid DNA. Cells and DNA were incubated on ice for 20 minutes before heat shocking at 42°C for 45 seconds. After 2 min incubation on ice 400 µL SOC (20 g/L tryptone, 5 g/L Yeast extract, 0.5 g/L NaCl, 0.186 g/L KCl, 10 mM MgCl₂, 10 mM MgSO₄ and 20 mM glucose) was added and the cell culture was incubated at 37°C with shaking for 1 hour before plating on selective
20 plates.

Plasmid were transformed into TOP10 chemical competent cells at conditions recommended by the supplier (ThermoFisher Scientific).

DNA techniques

25 Plasmid DNA from *E. coli* was isolated using the QIAprep Spin Miniprep kit (Qiagen). Chromosomal DNA from *E. coli* was isolated using the QIAamp DNA Mini Kit (Qiagen). PCR products were purified using the QIAquick PCR Purification Kit (Qiagen). DreamTaq PCR Master Mix (ThermoFisher), Phusion U hot start PCR master mix (ThermoFisher), USER Enzyme (New England Biolab) were used as
30 recommended by the supplier. Primers were supplied by Eurofins Genomics, Germany. PCR fragments and plasmids were sequenced by Eurofins Genomics. Colony PCR was done using DreamTaq PCR Master Mix in a T100™ Thermal Cycler (Bio-Rad).

35 **Table 3:** Oligos used for amplification of plasmid backbones, promoter elements, and

genes of interest:

Name	SEQ ID NO	Oligonucleotide Sequence 5'-3'	Description
O40	5	ATTAACCCUCCAGGCATCAAATAAAACGAAAGGC	Backbone.for
O68	6	ATGCGCAAUTGTGAGTTAGCTCACTCATTAG	Plac.for
O79	7	ATTTGCGCAUCACCAATCAAATTCACGCGGCC	Backbone.rev
O113	8	AGCTGTTUCCTCCTTAGGTACCCAGCTTTTGTTC C	Plac.rev
O261	9	ATGCGCAAUAGCGGCACGCCTTGCAGATTACG	PgIpF.for
O262	10	AGCTGTTUCCTCCTTGGTTAATGTTTGTGTATGCG	PgIpF.rev
KABY721	11	AAACAGCUATGAAGAGCGCGCTGACCTTTAGC	yberC0001_9420.for
KABY722	12	AGGGTTAAUTTACGCCTCACGCACACGCG	yberC0001_9420.rev
O48	13	CCCAGCGAGACCTGACCGCAGAAC	galK.for
O49	14	CCCCAGTCCATCAGCGTGACTACC	galK.rev

Table 4: The heterologous transporter gene tested for its ability to facilitate microbial production of HMOs

Gene	Origin of Genes	Accession Number	Function
yberC0001_9420	<i>Yersinia bercovieri</i>	EEQ08298	MFS transporter

5

Table 5: The synthetic DNA elements utilized for expression of *YberC*

Sequence name	SEQ ID NO	Sequence (5'-3')	Description
<i>PgIpF</i>	3	GCGGCACGCCTTGCAGATTACGTTTGCCACACTTTT CATCCTTCTCCTGGTGACATAATCCACATCAATCGAAA ATGTTAATAAATTTGTTGCGCGAATGATCTAACAACAT GCATCATGTACAATCAGATGGAATAAATGGCGCGATAA CGCTCATTTTATGACGAGGCACACACATTTTAAGTTTCG ATATTCTCGTTTTTGCTCGTTAACGATAAGTTTACAGC ATGCCTACAAGCATCGTGGAGGTCCGTGACTTTCACG CATAACAACAAACATTAACCAAGGAGGAAACAGCT	300-nucleotide DNA expression element
<i>Plac</i>	4	ATGCGCAAATTTGTGAGTTAGCTCACTCATTAGGCACCC CAGGCTTTACACTTTATGCTTCCGGCTCGTATGTTGTG TGGAATTGTGAGCGGATAACAATTTACACAGGAAACA GCTATGACCATGATTACGCCAAGCGCGCAATTAACCCT CACTAAAGGGAACAAAAGCTGGGTACCTAAGGAGGAA ACAGCT	195-nucleotide DNA expression element
<i>yberC0001_9420</i>	2	ATGAAGAGCGCGCTGACCTTTAGCCGTCGTATCAATC CGGTGTTTCTGGCGTTTTTCGTGGTGGCGTTCCTGAG CGGTATTGCGGGTGCCTGTCAGGCGCCGACCCTGAG CCTGTTTCTGAGCACCAGGTTAAAGTGCGTCCGCTG TGGGTTGGCCTGTTTTACACCGTGAACGCGATTGCGG GTATCACCGTTAGCTTCATTCTGGCGAAGCGTAGCGA CAGCCGTGGCGATCGTCGTAAACTGATCATGGTTTGC TACCTGATGGCGGTGGGTAAGTGCCTGCTGTTTCGCGT TTAACCGTGAATATCTGACCCTGATTACCGCGGGCGTT CTGCTGGCGAGCGTGGCGAACACCGCGATGCCGAG ATTTTCGCGCTGGCGCGTGAGTATGCGGATAGCAGCG CGCGTGAAGTGTTATGTTTACGACGATTATGCGTGC GCAACTGAGCCTGGCGTGGGTGATTGGTCCGCCGCT	MFS transporter

		GAGCTTCATGCTGGCGCTGAACTATGGCTTCACCGTG ATGTTTAGCATTGCGGCGGGTATCTTTGTTCTGAGCGC GCTGGTTGTGTGGTTCATTCTGCCGAGCGTGCCGCGT GCGGAACCGGTGGTTGACGCGCCGGTGGTTGTGCAG GGCAGCCTGTTTGCGGATAAAAACGTTCTGCTGCTGT CATTGCGAGCATGCTGATGTGGACCTGCAACACCATG TACATCATTGACATGCCGCTGTATATCACCGCGAGCCT GGGTCTGCCGGAGCGTCTGGCGGGTCTGCTGATGGG CACCGCGGCGGGTCTGAAATCCCGATTATGCTGCTG GCGGGCTACAGCGTGCGTTATTCGGCAAGCGTAAAA TCATGCTGTTTGCGGTTCTGGCGGGCGTGCTGTTCTA CACCGTCTGGTTCTGTTCAAGTTTAAACCGCGCTGA TGCTGCTGCAGATTTTCAACGCGATCTTTATTGGCAT GTGGCGGGTATCGGCATGCTGTATTTCAAGACCTGA TGCCGGGTCGTGCGGGTGCGGCGACCAACCTGTTCA CCAACAGCATTAGCACCGGCGTTATCCTGGCGGGTGT GCTGCAAGGTGGCCTGACCGAAACCTGGGGTCACGAT AGCGTTTATGTGATGGCGATGGTTCTGAGCATTCTGG CGCTGATTATCTGCGCGCGTGTCGTGAGGCGTAA	
--	--	---	--

Construction of Plasmids

- Plasmid backbones containing two *I-SceI* endonuclease sites, separated by two DNA fragments appropriated for homologous recombination into the *E. coli* genome and a T1 transcriptional terminator sequence were synthesized. For
- 5 example in one plasmid backbone the *gal* operon (required for homologous recombination in *galK*), and a T1 transcriptional terminator sequence (pUC57::*gal*) was synthesized (GeneScript). The DNA sequences used for homologous recombination in the *gal* operon covered base pairs 3.628.621-3.628.720 and
- 10 3.627.572-3.627.671 in sequence *Escherichia coli* K-12 MG155 complete genome GenBank: ID: CP014225.1. Insertion by homologous recombination would result in a deletion of 949 base pairs of *galK* and a *galK*- phenotype. In similar ways backbones based on pUC57 (GeneScript) or an any other appropriated vector containing two *I-SceI* endonuclease sites, separated by two DNA fragments appropriated for
- 15 homologous recombination into the *E. coli* genome and a T1 transcriptional terminator sequence could be synthesized. Standard techniques well-known in the field of molecular biology were used for designing of primers and amplification of specific DNA sequences of the *Escherichia coli* K-12 DH1 chromosomal DNA. Such standard techniques, vectors, and elements can be found, for example, in: Ausubel
- 20 *et al.* (eds.), *Current Protocols in Molecular Biology* (1995) (John Wiley & Sons); Sambrook, Fritsch, & Maniatis (eds.), *Molecular Cloning* (1989) (Cold Spring Harbor Laboratory Press, NY); Berger & Kimmel, *Methods in Enzymology* 152: Guide to Molecular Cloning Techniques (1987) (Academic Press); Bukhari *et al.* (eds.).

Chromosomal DNA obtained from *E. coli* DH1 was used to amplify a 300 bp DNA fragment containing the promoter PglpF using oligos O261 and O262, and a 195 bp DNA fragment containing Plac using oligos O68 and O113 (Table 3).

A 1.182 bp DNA fragment containing a codon optimized version of the yberC0001_9420 gene originating from *Yersinia bercovieri* was synthesized by GeneScript (Table 5). The yberC0001_9420 gene was amplified by PCR using oligos KABY721 and KABY722.

All PCR fragments (plasmid backbones, promoter containing elements and the YberC gene) were purified, and plasmid backbones, promoter elements (PglpF, or Plac), and a YberC containing DNA fragment were assembled. The plasmids were cloned by standard USER cloning. Cloning in any appropriated plasmid could be done using any standard DNA cloning techniques. The plasmids were transformed into TOP10 cells and selected on LB plates containing 100 µg/mL ampicillin (or any appropriated antibiotic) and 0.2 % glucose. The constructed plasmids were purified and the promoter sequence and the 5' end of the YberC gene was verified by DNA sequencing (MWG Eurofins Genomics). In this way, a genetic cassette containing any promoter of interest linked to the YberC gene was constructed.

Table 6: Examples of helper and donor plasmids used for strain construction

Plasmid	Relevant genotype	Marker gene
pACBSR	<i>Para-I-SceI-λ Red, p15A ori, cam*</i>	cam
pUC57	<i>pMB1, bla</i>	bla
pUC57::gal	<i>pUC57::galTK' / T1-galKM'</i>	bla

DNA sequences of heterologous genes coding for transporters or glycosyltransferases of interest were codon optimized and synthesized by GenScript. The gene of interest encoding for the transporter protein YberC (Table 4) was amplified by PCR using appropriated primers covering the start codon, ATG, and the stop codon, TAA, of the gene (Table 3). To construct donor plasmids with any heterologous gene of interest, standard USER cloning was employed to combine the purified PCR fragments of the relevant plasmid backbone, promoter element and gene of interest. Cloning in an appropriated plasmid could be done using any standard DNA cloning technique. Following cloning the DNA was transformed into TOP10 cells and selected on LB plates containing 100 µg/mL ampicillin (or 50 mg/mL kanamycin depending on the backbone applied) and 0.2 % glucose. The

constructed plasmids were purified, and the promoter sequence and the 5' end of the gene of interest were verified by DNA sequencing (MWG Eurofins Genomics).

Construction of strains

5 The bacterial strain used, MDO, was constructed from *Escherichia coli* K-12 DH1. The *E. coli* K-12 DH1 genotype is: *F⁻, λ , gyrA96, recA1, relA1, endA1, thi-1, hsdR17, supE44*. In addition to the *E. coli* K-12 DH1 genotype MDO has the following modifications: *lacZ*: deletion of 1.5 kbp, *lacA*: deletion of 0.5 kbp, *nanKETA*: deletion of 3.3 kbp, *melA*: deletion of 0.9 kbp, *wcaJ*: deletion of 0.5 kbp, *mdoH*: deletion of 0.5 kbp, and insertion of *Plac* promoter upstream of the *gmd* gene.

10 Insertion of an expression cassette containing a promoter linked to the *yberC0001_9420* gene and to a T1 transcriptional terminator sequence was performed by Gene Gorging essentially as described by Herring *et al.* (Herring, C.D., Glasner, J.D. and Blattner, F.R. (2003) *Gene* (311). 153-163), and example of helper and donor plasmids used to construct strains presented in the present application are provided in Table 6. Briefly, the donor plasmid and the helper plasmid were co-transformed into MDO and selected on LB plates containing 0.2 % glucose, ampicillin (100 µg/mL) or kanamycin (50 mg/mL) and chloramphenicol (20 µg/mL). A single colony was inoculated in 1 mL LB containing chloramphenicol (20 µg/mL) and 10 µL of 20 % L-arabinose and incubated at 37°C with shaking for 7 to 8 hours. For integration in the *galK* loci of *E. coli* cells were then plated on M9-DOG plates and incubated at 37°C for 48 hours. Single colonies formed on MM-DOG plates were re-streaked on LB plates containing 0.2 % glucose and incubated for 24 hours at 37°C. Colonies that appeared white on MacConkey-galactose agar plates and were sensitive for both ampicillin and chloramphenicol were expected to have lost the donor and the helper plasmid and contain an insertion in the *galK* loci. Insertions in the *galK* site was identified by colony PCR using primers O48 (SEQ ID NO: 13) and O49 (SEQ ID NO: 14) and the inserted DNA was verified by sequencing (Eurofins Genomics, Germany).

25 Insertion of genetic cassettes at other loci in the *E. coli* chromosomal DNA was done in a similar way using different selection marker genes.

Deep Well Assay (DWA)

35 DWA was performed as originally described to Lv *et al* (Bioprocess Biosyst Eng (2016) 39:1737–1747) and optimized for the purposes of the current invention.

More specifically, the strains disclosed in the examples were screened in 24 deep well plates using a 4-day protocol. During the first 24 hours, cells were grown to high densities while in the next 72 hours cells were transferred to a medium that allowed induction of gene expression and product formation. Specifically, during day 1 fresh
5 inoculums were prepared using a basal minimal medium supplemented with magnesium sulphate, thiamine and glucose. After 24 hours of incubation of the prepared cultures at 34 °C with a 700 rpm shaking, cells were transferred to a new basal minimal medium (2 ml) supplemented with magnesium sulphate and thiamine to which an initial bolus of 20 % glucose solution (1 µl) and 10 % lactose solution (0.1 ml)
10 were added, then 50 % sucrose solution as carbon source was provided to the cells. After inoculation of the new medium, cells were shaken at 700 rpm at 28 °C for 72 hours. After denaturation and subsequent centrifugation, the supernatants were analyzed by HPLC.

For the analysis of total samples, the cell lysate prepared by boiling was
15 pelleted by centrifugation for 10 minutes at 4.700 rpm. The HMO concentration in the supernatant was determined by HPLC or HPAC methods.

Results

20 **Example 1. Engineering of *Escherichia coli* for LNT-2 production using the *yberC0001_9420* gene**

The *Escherichia coli* K-12 (DH1) MDO strain can be manipulated to express heterologous genes of interest. For instance, the strain MP3941 is a LNT-2 production strain optimally overexpressing a β -1,3-N-acetylglucosaminyltransferase
25 genes selected from non-limiting examples of Table 1, e.g. *lgtA*.

Analysis of total samples (Figure 3) showed that LNT-2 titer can be markedly improved when an expression cassette containing a promoter element (*PglpF*) linked to a *yberC0001_9420* gene is integrated into the chromosomal DNA of strain MP3941 in a single copy to generate the strain MP3940 (Table 2).
30 Interestingly, the extracellular fraction of LNT-2 is also much larger for strain MP3940 than for strain MP3941, a fact that is also reflected in the lower trisaccharide levels that are detected in the pellet fraction of the *yberC0001_9420*-expressing strain MP3940 compared to the corresponding levels in the non-transporter expressing strain MP3941 (Figure 1).

35

Example 2. Engineering of *Escherichia coli* for LNnT production using the *yberC0001_9420* gene

The *Escherichia coli* K-12 (DH1) MDO strain can be manipulated to express heterologous genes of interest. For instance, the strain MP4436 is a LNnT production strain overexpressing a β -1,3-N-acetylglucosaminyltransferase gene and a β -1,3-galactosyltransferase gene, selected from non-limiting examples of Table 1, e.g. *lgtA* and *galT*, at optimal levels.

As shown from the HPLC analysis of total samples, insertion of an expression cassette containing the promoter element *P_{lac}* linked to a *yberC0001_9420* gene into the chromosomal DNA of strain MP4436 in a single copy to generate the strain MP3863 (Table 2), resulted in i) lower total LNnT titers, ii) slightly higher total LNT-2 concentrations, and iii) much lower total pLNnH formation (Figure 2).

Although the total LNnT titer in the *yberC0001_9420*-expressing strain MP3863 is not improved compared to the reference strain MP4436, it is obvious from the supernatant sample analysis that the extracellular LNnT fraction is markedly increased. A similar observation is true for the LNT-2 concentration in the medium of MP3863 cultures, where the LNT-2 fraction is at least 2-fold increased compared to the corresponding fraction for the strain MP4436 (Figure 2). The latter observation could easily explain the lower LNnT titer achieved with the strain MP3863 compared to the strain MP4436, since the precursor sugar molecule LNT-2 is more efficiently transported out of the cell in the presence of the YberC0001_9420 transporter. Accordingly, LNnT and LNT-2 concentrations in the pellet sample of the *yberC0001_9420*-expressing strain MP3863 are markedly reduced. As mentioned above, pLNnH formation is largely decreased in the strain MP3863 compared to the strain MP4436. Interestingly, pLNnH is seemingly found only in the cell pellet of the strain MP3863, and, without being bound by any scientific theory, it thus does not seem to be a substrate for the YberC0001_9420 transporter (Figure 2).

Example 3. Engineering of *Escherichia coli* for LNT production using the *yberC0001_9420* gene

The *Escherichia coli* K-12 (DH1) MDO strains can be manipulated to express heterologous genes of interest. For instance, the strain MP4473 is a LNT production strain expressing a β -1,3-N-acetylglucosaminyltransferase gene and a β -1,3-galactosyltransferase gene selected from non-limiting examples of Table 1, e.g. *lgtA* and *galT*, at optimal levels.

Insertion of an expression cassette containing the promoter element *Plac* linked to a *yberC0001_9420* gene into the chromosomal DNA of strain MP4473 in a single copy to generate the strain MP4462 (Table 2), had a strong impact on total HMO titers and HMO distribution between the supernatant and pellet fractions of cell cultures (Figure 3).

As shown from the analysis of total samples, *Plac*-driven expression of the *yberC0001_9420* gene had a drastically positive effect on total LNT titer, which was accompanied by a more than 2-fold increase in total LNT-2 titer and marked reduction in total pLNH-II formation.

The significant impact of introducing *yberC0001_9420* gene under the control of an expression element of a moderate strength, e.g. *Plac*, in the genetic background of a LNT production strain with optimal glycosyltransferase expression, is highlighted further by looking at the analysis of supernatant samples. The LNT concentration in the supernatant fraction of cultures of the strain MP4462 is largely increased compared to the one measured in the medium of MP4473 cultures (Figure 3). In agreement with the higher total LNT-2 titers observed in the transporter-expressing strain MP4462 compared to the reference strain MP4473, LNT-2 concentration in the supernatant samples of former strain is at least 2-fold higher than in the latter strain. Accordingly, the LNT, but not LNT-2, concentration in the pellet of the *yberC0001_9420*-expressing strain MP4462 is markedly decreased. As mentioned above, pLNH-II formation is largely reduced in strain MP4462. Interestingly, pLNH-II is seemingly found only in the cell pellet of the strain MP4462, and, without being bound by any scientific theory, it thus does not seem to be a substrate for the *YberC0001_9420* transporter (Figure 3).