DIFLUORO PENTAPEPTIDE DERIVATIVE ANTI-INFLAMMATORY AGENTS

TECHNICAL FIELD

The subject invention relates to novel peptide derivatives which are useful as anti-inflammatory agents.

BACKGROUND OF THE INVENTION

Certain polypeptide derivatives having various biological activities are known. The following references disclose such polypeptide derivatives: U.S. Patent No. 4,242,329 issued to Claeson, Simonsson & Arielly on December 30, 1980; U.S. Patent No. 4,316,889 issued to Bajusz, Hasenohrl, Barabas & Bagdy on February 23, 1982; U.S. Patent No. 4,399,065 issued to Bajusz, Hasenohrl, Barabas & Bagdy on August 16, 1983; U.S. Patent No. 4,401,594 issued to Umezawa, Takeuchi, Aoyagi, Ishii, Saino & Someno on August 30, 1983; U.S. Patent No. 4,478,745 issued to Bajusz, Hasenohrl, Barabas & Bagdy on October 23, 1984; U.S. Patent No. 4,528,133 issued to Kasafirek, Fric, Slaby & Robalova on July 9, 1985; U.S. Patent No. 4,596,789 issued to Dutta, Stein, Trainor & Wildonger on June 24, 1986; U.S. Patent No. 4,623,639 issued to Hassall, Johnson & Roberts on November 18, 1986; U.S. Patent No. 4,703,036 issued to Bajusz, Hasenohrl, Bagdy, Barabas, Dioszegi, Fittler, Jozsa, Horvath & Jozst on October 27, 1987; U.S. Patent No. 4,880,780 issued to Trainor & Stein on November 14, 1989; U.S. Patent No. 4,883,863 issued to Abe, Yaginuma, Nagasawa & Kuroiwa on November 28, 1989; U.S. Patent No. 4,902,781 issued to Mizoue, Okazaki, Hanada, Omura & Amamoto on February 20, 1990; U.S. Patent No. 4,923,890 issued to Trainor & Stein on May 8, 1990; U.S. Patent No. 5,066,643 issued to Abeles & Gelb on November 19, 1991; PCT Patent Application No. 84/00365 of Galpin & Wilby, published February 2, 1984; PCT Patent Application No. 87/02675 of Hoover, published May 7, 1987; Japanese Patent Application No. 47-30618 of Toray Inds. Inc., published November 9, 1972; Japanese Patent Application No. 57-145846 of Mitsubishi Chem. Ind. KK, published July 19, 1974; Japanese Patent Application No. 58-164563 of Daiichi Kagaku Yaku, published March 25, 1982; European Patent Application No. 0275101 of Merrell Dow Pharmaceuticals, Inc., published July 20, 1988; Aoyagi, Miyata, Nanbo, Kojima, Matsuzaki, Ishizuka, Takeuchi & Umezawa, "Biological Activities of Leupeptins," The Journal of Antibiotics, Vol.

This Sheet Has Been Replaced

XXII, No. 11 (Nov. 1969), pp. 558-568; Bajusz, Barabas, Tolnay, Szell & Bagdy, "Inhibition of Thrombin and Trypsin by Tripeptide Aldehydes," Int. J. Peptide Protein Res., Vol. 12 (1978), pp. 217-221; Gaal, Bacsy & Rappay, "Cationic Ferritin Uptake by Cultured Anterior Pituitary Cells Treated with the Proteinase Inhibitor, BOC-DPhe-Phe-Lys-H," Histochemistry, Vol. 88 (1988), pp. 401-406; Makara, Rappay, Garamvolgyi, Nagy, Danko & Bajusz, "The Tripeptide Aldehyde, Boc-DPhe-Phe-Lysinal, is a Novel $Ca2^+$ Channel Inhibitor in Pituitary Cells," European Journal of Pharmacology, Vol. 151 (1988), pp. 147-149; Nagy, Makara, Horvath, Rappay, Kurcz & Bajusz, "Tripeptide Aldehyde Protease Inhibitors May Depress in Vitro Prolactin and Growth Hormone Release," Endocrinology, Vol. 116, No. 4 (1895), pp. 1426-1432; Rappay, Makara, Bajusz & Nagy, "Various Proteinase Inhibitors Decrease Prolactin and Growth Hormone Release by Anterior Pituitary Cells," Life Sciences, Vol. 36 (1985), pp. 549-555.

It is an object of the subject invention to provide novel difluoro pentapeptide derivatives which are useful as anti-inflammatory agents.


It is a further object of the subject invention to provide novel difluoro pentapeptide derivatives which reduce bone resorption and/or heterotopic bone formation associated with rheumatoid arthritis.

It is also an object of the subject invention to provide novel pharmaceutical compositions comprising certain difluoro pentapeptide derivatives.

It is also an object of the subject invention to provide methods for treating diseases characterized by inflammation using the subject compounds.

SUMMARY OF THE INVENTION

The subject invention involves anti-inflammatory compounds having the following structure:



wherein:

(a) -X is selected from the group consisting of cyclic alkyl having from 4 to about 15 carbon atoms; branched alkyl, with at least two branches, having from 6 to about 15 carbon atoms; and arylalkyl having from 6 to about 15 carbon atoms;

(b) n is an integer from 0 to about 2;

This Sheet Has Been Replaced

(c) -V- is selected from the group consisting of -OC(O)-, -N(Q)C(O)-, -N(Q)C(S)-, -C(O)-, -SO_2- and -P(O)(OH)-;

(d) -Q is hydrogen; or straight or branched chain alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; or -Q and -X are covalently linked forming a cyclic moiety which includes the nitrogen to which -Q is bonded and from about 5 to about 20 carbon atoms;

(e) Z is -O- or -NH-; when V is -OC(O)-, -Z- is -NH-;

(f) $-R^1$ is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^1$ is in either D or L configuration;

(g) $-R^2$ is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^2$ is in L configuration;

(h) $-R^3$ is $-(CH_2)_m$-A-NH_2 or $-(CH_2)_m$-A-B-$C(NH_2)$=NH wherein m is an integer from 1 to about 6, -A- is a covalent bond or p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-; and the carbon atom bonded to $-R^3$ is in L configuration;


(i) $-R^4$ is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^4$ is in L configuration;

(j) $-R^5$ is $-(CH_2)_m$-A-NH_2 or $-(CH_2)_m$-A-B-$C(NH_2)$=NH wherein m is an integer from 1 to about 6, -A- is a covalent bond or p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-; and the carbon atom bonded to $-R^5$ is in D or L configuration; and

(k) -Y is hydrogen or methyl.

The subject invention also involves pharmaceutical compositions comprising the above compounds, and methods for treating inflammation or pain using such compounds and compositions.

This Sheet Has Been Replaced



DETAILED DESCRIPTION OF THE INVENTION

The term "alkyl," as used herein, unless otherwise indicated, means carbon-containing chains which may be straight, branched or cyclic; which may be saturated or unsaturated; and which may be unsubstituted or substituted. As used herein, "cyclic alkyl" may have all or only a portion of the total number of carbon atoms indicated as being in the alkyl group in the cyclic ring itself. Cyclic alkyl includes monocycloalkyl, bicycloalkyl and/or tricycloalkyl.

Preferred alkyl are as noted in this paragraph unless provided otherwise in particular instances. Preferred alkyl are saturated. Preferred alkyl are unsubstituted. Preferred alkyl substituents are selected from halo, amino, hydroxy, alkoxy, cyano, nitro, aryl and trifluoromethyl. As used herein, "alkoxy" is -O-alkyl. It is preferred that substituted alkyl be mono-, di- or trisubstituted, especially monosubstituted.

The term "aryl," as used herein, unless otherwise specified, means aryl rings which may be unsubstituted or substituted. Preferred aryl is as noted in this paragraph, unless provided otherwise in particular instances. Preferred aryl are phenyl and naphthyl, especially phenyl. Preferred aryl are mono-, di-, tri- or unsubstituted; more preferred aryl are monosubstituted or unsubstituted, especially unsubstituted. Preferred aryl substituents include alkyl, halo, amino, hydroxy, alkoxy, cyano, nitro and trifluoromethyl.


The term "arylalkyl," as used herein, means alkyl substituted with aryl. A preferred arylalkyl is arylmethyl.

The term "aliphatic" as used herein, means open-chain carbon-containing portions of arylalkyls.

The following abbreviations are used herein. Abbreviations for amino acids may refer to the amino acid itself, or more often to an amino acid moiety that is part of a peptide or peptide derivative structure.

Arg	Arginyl
Orn	Ornithyl
Leu	Leucyl
Gly	Glycyl
t-Bug	t-Butylglycyl
Phe	Phenylalanyl
Adoc	Adamantyloxycarbonyl
Z	Benzyloxycarbonyl
Boc	t-Butyloxycarbonyl
DECP	Diethylcyanophosphonate

This Sheet Has Been Replaced




Me	Methyl
Et	Ethyl
TEA	Triethylamine
DMF	Dimethylformamide
TFA	Trifluoroacetate
ipa	Isopropanol
Bn	Benzyl
Lys	Lysyl
Ac	Acetyl
TLC	Thin Layer Chromatography
TSA	Toluenesulfonic acid
THF	Tetrahydrofuran
HOBt	Hydroxybenzotriazole
DCC	Dicyclohexylcarbodiimide
DIPEA	N,N-Diisopropylethylamine
Chg	Cyclohexylglycyl
Cha	Cyclohexylalanyl
Nal	Naphthylalanyl
Trp	Tryptophyl
Adg	Adamantylglycyl
pGphe	p-Guanidinophenylalanyl
3,5-Dmadoc	3,5-Dimethyladamantyloxycarbonyl
eBroc	endo-Bornyloxycarbonyl
Noc	Naphthyloxycarbonyl
Moc	Menthyloxycarbonyl
Ad	Adamantoyl
Fmoc	Fluorenylmethoxycarbonyl
Ada	Adamantaneacetyl
Adac	Adamantylaminocarbonyl
Mnoc	Morpholinyl
Norad	Noradamantoyl
Hadoc	Homoadamantyloxycarbonyl
Foc	Fenchyloxycarbonyl
Imoc	Isomenthyloxycarbonyl
Ipoc	Isopinocamphanyloxycarbonyl
AC	acetyl
DMF	dimethylformamide

This Sheet Has Been Replaced



HPLC high pressure liquid chromatography

The novel anti-inflammatory compounds of the subject invention are those having the following chemical structure:



In Structure (1), n is an integer of from 0 to about 2; n is preferably 0 or 1.

In Structure (1), $-R^1$ is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl, the aliphatic portion being saturated and having 1 or 2 carbon atoms. Preferred $-R^1$ is saturated alkyl. Preferred alkyl are unsubstituted, or are arylalkyl, especially benzyl and naphthal. Preferred cyclic alkyl are cyclic C_3-C_8 (more preferably C_5-C_6) methyl or adamantylmethyl. Preferred $-R^1$ is hydrophobic, preferably with the hydrophobicity concentrated close to the carbon atom to which $-R^1$ is bonded. Specific examples of preferred $-R^1$ include t-butyl, 1,1-dimethylpropyl, i-propyl, i-butyl, s-butyl, neo-pentyl, cyclohexyl, cyclohexylmethyl, adamantyl, adamantylmethyl, benzyl, naphthal; most preferred $-R^1$ is t-butyl.

In Structure (1), the carbon to which $-R^1$ is bonded is in either D or L, preferably D, configuration. The structure $-Z-CH(R^1)-C(O)-$ is an amino acid moiety (hereinafter -AA1-) when -Z- is -NH; preferred amino acid moieties for -AA1- include t-Bug, Val, Ile, Leu, Chg, Cha, Phe, Nal, Trp and Adg; more preferred are t-Bug, Val and Ile; most preferred -AA1- is t-Bug.

In Structure (1), $-R^2$ is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl, the aliphatic portion being saturated and having 1 or 2 carbon atoms. Preferred branched or cyclic alkyl are saturated. Preferred branched or cyclic alkyl are unsubstituted. Preferred branched alkyl have from 3 to 5 carbon atoms; most preferred branched alkyl is i-butyl. Preferred cyclic alkyl have a C_3-C_7, more preferably C_5-C_6, cyclic ring bonded to a methylene, ethylene or n-propylene (preferably methylene or ethylene, more preferably methylene) which is bonded to the carbon atom in Structure (1) to which $-R^2$ is bonded. Preferred arylalkyl

This Sheet Has Been Replaced



are benzyl, p-hydroxybenzyl and naphthal. More preferred -R^2 is selected from i-propyl, i-butyl, s-butyl, cyclohexylmethyl, benzyl and naphthal; most preferred is benzyl.

In Structure (1), the carbon atom to which -R^2 is bonded is in D or L configuration. The structure -NH-CH(R^2)-C(O)- is an amino acid moiety (hereinafter -AA^2-); preferred amino acid moieties for -AA^2- include Phe, Nal, Cha, Leu and Ile; most preferred -AA^2- is Phe.


In Structure (1), -R^3 is -$(CH_2)_m$-A-NH_2 or -$(CH_2)_m$-A-B-C(NH_2)=NH, wherein m is an integer of from 1 to about 6, -A- is a covalent bond, p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-. When -A- is a covalent bond, m is preferably 2-5, more preferably 2-4, more preferably still 3 or 4. When -A- is p-phenyl or p-cyclohexyl, m is preferably 1-4, more preferably 1-3, more preferably still 1. -A- is preferably a covalent bond. -B- is preferably -NH-. More preferred -R^3 is 3-guanidino-n-propyl or 4-amino-n-butyl.

In Structure (1), the carbon to which -R^3 is bonded is in D or L configuration, preferably in L configuration. The structure -NH-(CH(R^3)-C(O)- is an amino acid moiety (hereinafter -AA^3-); preferred amino acid moieties for -AA^3- include Arg, Lys, homo-Arg, p-amidino-phe, p-amino-phe and p-Gphe; most preferred -AA^3- is Lys.

In Structure (1), -R^4 is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl, the aliphatic portion being saturated and having 1 or 2 carbon atoms. Preferred branched or cyclic alkyl are saturated and unsubstituted. Preferred branched alkyl have from 3 to 5 carbon atoms; most preferred branched alkyl is i-butyl. Preferred cyclic alkyl have a C_3-C_7, more preferably C_5-C_6 cyclic ring bonded to a methylene, ethylene or n-propylene (preferably methylene or ethylene, more preferably methylene) which is bonded to the carbon atom in Structure (1) to which -R^4 is bonded. Preferred arylalkyl are benzyl, p-hydroxybenzyl and naphthal. More preferred -R^4 is selected from i-propyl, i-butyl, s-butyl, cyclohexylmethyl, benzyl and naphthal; most preferred is benzyl.

In Structure (1), the carbon atom to which -R^4 is bonded is in D or L configuration. The structure -NH-CH(R^4)-C(O)- is an amino acid moiety (hereinafter -AA^4-); preferred amino acid moieties for -AA^4- include Phe, Nal, Cha, Leu and Ile; most preferred -AA^4- is Leu.

This Sheet Has Been Replaced



In Structure (1), $-R^5$ is $-(CH_2)_m$-A-NH_2 or $-(CH_2)_m$-A-B-$C(NH_2)$=NH, wherein m is an integer from 1 to about 6, -A- is a covalent bond, p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-. When -A- is a covalent bond, m is preferably 2-5, more preferably 2-4, more preferably still 3 or 4. When -A- is p-phenyl or p-cyclohexyl, m is preferably 1-4, more preferably 1-3, more preferably still 1. -A- is preferably a covalent bond. -B- is preferably -NH-. More preferred $-R^5$ is 3-guanidino-n-propyl or 4-amino-n-butyl.

In Structure (1), the carbon to which $-R^5$ is bonded is in D or L configuration, preferably in L configuration. The structure -NH-CH(R^5)-C(O)- is an amino acid moiety (hereinafter $-AA^5$); preferred amino acid moieties for $-AA^5$ include Arg, homo-Arg, Lys, p-amidino-Phe, p-amino-Phe and p-Gphe; most preferred $-AA^5$- is Arg.

In Structure (1), -Y is hydrogen or methyl.


In Structure (1), -Z- is -O- or -NH-. Preferred -Z- is -NH-.

In Structure (1), -V- is selected from -OC(O)-, -N(Q)C(O)-, -N(Q)C(S)-, -C(O)-, $-SO^2$- and -P(O)(OH)-. Preferred -V- is selected from -OC(O)-, -N(Q)C(O)-, -N(Q)C(S)-, and -C(O)-. More preferred -V- is -OC(O)- or -N(Q)C(O)-. Most preferred -V- is -OC(O)-. When -V- is -OC(O)-, -Z- is -NH-.

In Structure (1), -X is selected from cyclic alkyl, having from 4 to about 15 carbon atoms; branched alkyl having from 6 to about 15 carbon atoms and at least 2 branches; and aryl, having from 6 to about 15 carbon atoms. Preferred -X has from 6 to 12 carbon atoms; more preferred -X has from 8 to 10 carbon atoms. Preferred alkyl portions of -X are saturated. Preferred -X is unsubstituted, or substituted with unsubstituted alkyl or aryl. Preferred cyclic alkyl are monocycloalkyl, bicycloalkyl, and tricycloalkyl, more preferred are bicycloalkyl and tricycloalkyl, especially tricycloalkyl. Preferred cycloalkyl have 5 or 6 carbon atoms, more preferably 6 carbon atoms, in each cyclic ring. A highly preferred -X is adamantyl. Preferred aryl -X are fluorenyl, naphthyl and phenyl, unsubstituted or substituted with alkyl. Particularly preferred aryl -X include naphthyl and fluorenyl. When -X is aryl or cyclic alkyl, n is preferably 1.

In Structure (1) when -V- is -N(Q)C(O)- or -N(Q)C(S)-, -Q is selected from hydrogen; straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; or -Q and -X are covalently linked forming a cyclic moiety which includes the nitrogen to which -Q is bonded and from about 5 to about 20 carbon atoms. Preferred -Q-X- has from 6 to 15 carbon atoms; more preferred -Q-X- has from 8 to 12 carbon atoms.

This Sheet Has Been Replaced



Preferred -Q-X- is unsubstituted or substituted with unsubstituted alkyl or aryl. Preferred cyclic

- N - X is monocyclic, bicyclic or tricyclic; more preferred is monocyclic. Most preferred -Q is hydrogen.

In Structure (1), preferred X-$(CH^2)_n$-V- (hereinafter X'-) include Adoc. Ad, Fmoc, Ada, Adac, Mnoc, Norad, Hadoc, Foc, Imoc, Ipoc, 3,5-Dmadoc, eBroc, Noc and Moc; more preferred X' is Adoc, Ipoc and eBroc; most preferred -X' is Adoc.

Preferred anti-inflammatory compounds of the subject invention include pharmaceutically-acceptable salts of the above compounds. Particularly preferred salts include salts of addition formed between a strong acid and the above compounds. Particularly preferred are such salts of addition where either $-R^3$ or $-R^5$, or both, are protonated, resulting in a positive charge on the $-R^3$ and R^5 moieties. Preferred compounds of the subject invention are often associated with one or more molecules of water in the form of hydrates.

Preferred anti-inflammatory compounds of the subject invention include the following compounds:

Adoc-D-tBug-Phe-Lys-C(O)CF_2C(O)-Leu-Arg-OMe
Adoc-D-tBug-Phe-Lys-C(O)CF_2C(O)-Gly-Arg-OMe
Adoc-D-tBug-Phe-Lys-C(O)CF_2C(O)-D-Phe-Arg-OMe
Adoc-D-tBug-Phe-Lys-C(O)CF_2C(O)-Leu-D-Arg-OMe
Mnoc-D-Phe-Phe-Arg-C(O)-CF_2C(O)-Leu-Lys-OMe
eBroc-D-t-Bug-Phe-pGphe-C(O)CF_2C(O)-Ala-Arg-OMe
Ipoc-D-Nal-Phe-Arg-C(O)CF_2C(O)-Gly-Arg-OMe
Adoc-D-Phe-Phe-Lys-C(O)CF_2C(O)-Leu-Arg-OH
Noc-D-t-Bug-Phe-Lys-C(O)CF_2C(O)-Ala-Lys-OMe

Pharmaceutical Compositions

Pharmaceutical compositions of the subject invention comprise a safe and effective amount of a difluoro pentapeptide derivative as defined hereinabove and a pharmaceutically-acceptable carrier. Such compositions typically comprise from about 0.1% to about 95% by weight of the difluoro pentapeptide derivative, preferably from about 1% to about 90% and more preferably from about 5% to about 75%.

The term "pharmaceutically-acceptable carrier," as used herein, means one or more compatible solid or liquid filler diluents or encapsulating

This Sheet Has Been Replaced



substances which are suitable for administration to a human or lower animal. The term "compatible," as used herein, means that the components of the pharmaceutical compositions are capable of being commingled with the difluoro pentapeptide derivative of the subject invention, and with each other, in a manner such that there is no interaction which would substantially reduce the pharmaceutical efficacy of the composition under ordinary use situations. Pharmaceutically-acceptable carriers must, of course, be of sufficiently high purity and sufficiently low toxicity to render them suitable for administration to the human or lower animal being treated.

Some examples of substances which can serve as pharmaceutically-acceptable carriers are sugars, such as lactose, glucose and sucrose; starches, such as cornstarch and potato starch; cellulose and its derivatives, such as sodium carboxymethylcellulose, ethylcellulose and cellulose acetate; powdered tragacanth; malt; gelatins; talc; stearic acid; magnesium stearate; calcium sulfate; vegetable oils, such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil and oil of theobroma; polyols, such as propylene glycol, glycerin, sorbitol, mannitol, and polyethylene glycol; alginic acid; pyrogen-free water; isotonic saline; phosphate buffer solutions; cocoa butter (suppository base); emulsifiers, such as TweensR; wetting agents and lubricants, such as sodium lauryl sulfate; coloring agents; flavoring agents; stabilizers; antioxidants; preservatives; as well as other non-toxic compatible substances used in pharmaceutical formulations. Other compatible pharmaceutical additives and actives may be included in the pharmaceutically-acceptable carrier for use in the compositions of the subject invention.

The pharmaceutically-acceptable carrier employed in conjunction with the difluoro pentapeptide derivative of the subject invention is used at a concentration sufficient to provide a practical size to dosage relationship. The pharmaceutically-acceptable carrier, in total, typically comprises from about 5% to about 99.9% by weight of the pharmaceutical compositions of the subject invention, preferably from about 10% to about 99%, and more preferably from about 25% to about 95%.

Total single dosages of the difluoro pentapeptide derivatives of the subject invention in pharmaceutical compositions are generally from about 1 mg to about 1000 mg, preferably from about 50 mg to about 800 mg, more preferably from about 100 mg to about 500 mg.

The choice of a pharmaceutically-acceptable carrier to be used in conjunction with the difluoro pentapeptide derivatives of the subject invention is

This Sheet Has Been Replaced



determined by the way the active is to be administered. Preferred modes of administering the actives of the subject invention are by injection, ingestion, inhalation and topically.

Methods and materials for manufacturing injectables can be found in Remington's Pharmaceutical Sciences, 17th ed., 1985, Chapter 85, p. 1518, the disclosures of which are incorporated herein by reference in their entirety. Generally, three types of injectable dosage forms are preferred: (1) aqueous solutions, (2) non-aqueous solutions; and (3) emulsions. Injectable dosage forms typically contain from about 0.001 mg/ml to about 10 mg/ml, preferably from about 0.4 mg/ml to about 0.6 mg/ml, of a compound of the subject invention. Preferred injectable dosage compositions of the subject invention include sterile aqueous solutions or emulsions of pH from about 3 to about 8 (more preferably of pH from about 4 to about 6).

Preferred pharmaceutical carriers in which the difluoro pentapeptide derivatives of the subject invention can be administered by ingestion are proteinoid microcapsules disclosed in the following patents: U.S. Patent No. 4,895,725 issued to Kantor, Steiner & Pack on January 23, 1990; U.S. Patent No. 4,925,673 issued to Steiner & Rosen on May 15, 1990; U.S. Patent No. 4,976,968 issued to Steiner on December 11, 1990; and U.S. Patent No. 4,983,402 issued to Steiner on January 8, 1991; all four patents are hereby incorporated in their entirety by reference.

Methods for Producing Anti-inflammatory Activity and Analgesia

The subject invention also encompasses methods of producing anti-inflammatory activity and/or analgesia in humans or lower animals through administering, to the human or lower animal in need of such treatment, a safe and effective amount of a difluoro pentapeptide derivative of the subject invention. The amount can be given in a single dose or multiple doses repeatedly over the course of the treatment. While dosages higher than those described herein are effective to reduce inflammation and produce analgesia, care must be taken in some individuals to prevent adverse side effects. The difluoro pentapeptide derivatives and compositions of the subject invention can be used to reduce inflammation in various disorders at the deeper structures, muscles, tendons, bursa and joints associated with disease and trauma, to treat and prevent pain.

The preferred modes of administering the difluoro pentapeptide derivatives of the subject invention are by injection, ingestion, inhalation and topically. Specific modes of administration include, without limitation, oral

This Sheet Has Been Replaced



ingestion; injection, such as intramuscular, intravenous, intraperitoneal, intradermal and subcutaneous; inhalation; and topically, such as transdermally, orally, mucosally and sublingually.

The term "safe and effective amount," as used herein, means an amount of a difluoro pentapeptide derivative or composition high enough to significantly positively modify the condition to be treated, but low enough to avoid serious side effects (at a reasonable benefit/risk ratio), within the scope of sound medical judgement. A safe and effective amount of a difluoro pentapeptide derivative or composition will vary with the particular condition being treated, the age and physical condition of the patient being treated, the severity of the condition, the duration of the treatment, the nature of concurrent therapy, the specific active employed, the particular pharmaceutically-acceptable carrier utilized, and like factors within the knowledge and expertise of the attending physician.

Preferred daily dosages of the difluoro pentapeptide derivatives of the subject invention range from about 0.1 mg/kg of body weight to about 500 mg/kg of body weight, more preferably from about 1 mg/kg to about 100 mg/kg, still more preferably from about 2 mg/kg to about 50 mg/kg. From 1 to about 4 single dosages per day may be administered, more preferably from about 2 to about 3 single dosages per day. Preferred amounts of the difluoro pentapeptide derivatives administered by injection are from about 0.1 mg/kg/day to about 50 mg/kg/day, more preferably from about 1 mg/kg/day to about 10 mg/kg/day. Preferred amounts of the difluoro pentapeptide derivatives administered by oral ingestion are from about 1 mg/kg/day to about 500 mg/kg/day, more preferably from about 5 mg/kg/day to about 100 mg/kg/day. Preferred amounts of the difluoro pentapeptide derivatives administered by inhalation are from about 0.1 mg/kg/day to about 500 mg/kg/day, more preferably from about 5 mg/kg/day to about 100 mg/kg/day. Preferred amounts of the difluoro pentapeptide derivatives administered topically are from about 1 mg/kg/day to about 500 mg/kg/day, more preferably from about 50 mg/kg/day to about 250 mg/kg/day.

Methods for Synthesizing the Compounds

The following non-limiting schemes and examples demonstrates methods of synthesizing difluoro pentapeptide derivatives of the subject invention.

I. Synthesis of Adoc-D-t-Bug-Phe-Lys-C(0)-CF_2-C(0)-Leu-Arg-OMe

Adoc-D-t-Bug-OH - To a solution of 48.8 g (0.370 M) of D-t-butylglycine in 371 ml of 1N NaOH and 159 ml of dioxane at 0 C was added 33.9 g (0.40 M) of $NaHCO_3$ followed by the dropwise addition of a solution of 80.0 g (0.40 M) of

This Sheet Has Been Replaced




adamantyl fluoroformate in 371 ml of dioxane over 3 h. The mixture is stirred at this temperature for 0.5 h and the cooling bath removed. The solution is stirred for 3 h and the pH adjusted to 10 with 1N NaOH. This solution is extracted with EtOAc and the remaining aqueous phase acidified to pH = 3. This acidic solution is extracted with EtOAc, dried and the solvent removed to afford 118 g of product that is homogeneous by TLC (94/5/1 - CH_2Cl_2/ipa/AcOH), Rf = 0.90.

Adoc-D-t-Bug-Phe-OBn - To a solution of 115 g (0.37 M) of Adoc-D-t-Bug-OH and L-Phe-OBn.pTSA (0.36 M) in 1L CH_2Cl_2 is added 74.6 g (0.74 M) of TEA followed by the addition of 60.8 g (0.37 M) of DECP and the resulting solution is stirred overnight. The solvent is removed and the product redissolved in EtOAc and washed successively with brine, 0.4N HCl, and saturated $NaHCO_3$. The solution is dried and the solvent removed. The residue is chromatographed on silica to afford 186 g of product. TLC (85/15 - hexane/ETOAc) - Rf = 0.40.

Adoc-D-tBug-Phe-OH (1) - To a dry flask under N_2 is added 34.0 g 5% Pd on carbon. The flask is cooled to -78oC and 200 ml of dry degassed MeOH is added. To this slurry is added 8.8 ml of formic acid (88%) and this is followed by the addition of a mixture of 36.0 g of Adoc-D-t-Bug-Phe-OBn (0.066 M) in 100 ml of MeOH and 4.4 ml of formic acid. The cooling bath is removed and the slurry stirred for 2 h. At this time an additional 200 ml of 4.4% formic acid in MeOH is added and the slurry stirred overnight. The mixture is filtered through glass fiber filter paper under N_2. The filtrate is refiltered through a bed of Celite to remove all traces of catalyst. Removal of the solvent yields 17.3 g of product.

α-Boc, Σ-Z-Lys-C(OH)-CF_2-C(O)-OEt (3) - To a solution of 79.2 g (1.21 M) of activated zinc in 0.600 L THF is added 2.36 g of ethyl bromodifluoroacetate and the solution brought to reflux. As soon as reflux begins a mixture of 236 g (1.16 M) of ethyl bromodifluoroacetate and 176 g (0.485 M) of (2) in 1.80 L of THF is added dropwise. For the synthesis of (2), see Burkhart, J., N. Peet and P. Bey, "A Novel Method for the Preparation of Peptidyl a-Keto Esters," Tetrahedron Letters, Vol. 31, No. 10, pp. 1385 -1388 (1990). The solution is refluxed for 1 h, cooled and quenched with saturated $KHSO_4$. The quenched product is diluted with EtOAc and washed successively with saturated $KHSO_4$, saturated $NaHCO_3$ and brine. This sequence is followed by drying with $MgSO_4$ and removal of solvent. The residue is chromatographed on silica to afford 68.0 g of pure product. Rf = 0.20 75/25 hexane/EtOAc.

This Sheet Has Been Replaced



α-Boc, Σ-Z-Lys-C(OH)-CF_2-C(O)-ONa (4) - To a solution of 1.00 g (0.002 M) of (3) in 10 ml of MeOH is added 2 ml of 1N NaOH and the solution stirred at room temperature for 5 h and the solvent is removed. The residue is dissolved in a small amount of water and the milky solution extracted with ether. The water is removed from the aqueous phase and the residue thoroughly dried by placing on the vacuum pump for 24 h. Recovered 0.069 g of product is recovered.

α-Boc, Σ-Z-Lys-C(OH)-CF_2-C(O)-Leu-Arg(NO_2)-OMe (5) - To a solution of 0.500 g (1.03 mmol) of (4) and 0.470 g (1.26 mmol) of Leu-Arg(NO_2)-OMe.HCl in 30 ml of DMF is added 0.019 g (1.26 mmol) of HOBt in one portion. This addition is followed by the dropwise addition of 0.260 g (1.26 mmol) of DCC in 3 ml DMF. Finally, 0.003 ml of DIPEA is added and the solution stirred for 48 h. The DMF is removed in vacuo and the residue dissolved in EtOAc. The solution is transferred to a separatory funnel and washed with 0.1 N HCl, saturated $NaHCO_3$, and brine. The ethyl acetate layer is dried with $MgSO_4$, filtered and the solvent removed. The residue is chromatographed on silica to afford 0.420 g of product. Rf = 0.68 - 90/10 CH_2Cl_2/MeOH.

Σ-Z-Lys-C(OH)-CF_2-C(O)-Leu-Arg(NO_2)-OMe.HCl - To a solution of (5) (17.3 g, 0.022 M) in 200 ml of dioxane is added 66.8 ml (0.220 m) of 3.3 M HCl in dioxane and the mixture stirred for 1h. At this time an additional 5 ml of 3.3 M HCl in dioxane is added and the solution stirred an additional 0.25 h. The solvent is removed and the residue placed on the vacuum pump. The solid residue is triturated with ether under N2 and filtered. Recovered 16.0 g of white solid.

Adoc-D-t-Buq-Phe-Lys(Z)-C(OH)-CF_2-C(O)-Leu-Arg(NO_2)-OMe (6) - To a solution of 16.0 g of the residue from the previous reaction (0.022 M) and 11.1 g of (1) (0.024 M) in 300 ml of CH_2Cl_2 is added 6.69 ml (0.048 M) of TEA dropwise followed by the dropwise addition of 3.64 ml (0.024 M) of DECP. The solution is stirred overnight and the solvent removed. The residue is dissolved in EtOAc and washed successively with 0.1 N HCl, saturated $NaHCO_3$, and brine. The solution is dried and the solvent removed to afford 26.6 g that is chromatographed on silica to afford 12.7 g of pure product. Rf = 0.85, 90/10 - CH_2Cl_2/MeOH.

Adoc-D-t-Buq-Phe-Lys(Z)-C(O)CF_2C(O)-Leu-Arg(NO_2)-OMe - To a slurry of 14.3 g (0.034 M) of Dess-Martin periodinane in 1.15 L of CH_2Cl_2 is added 0.910 ml of dry pyridine, followed by the dropwise addition of 12.7 g

This Sheet Has Been Replaced




(0.011 M) of (6) in 2.3 L of CH_2Cl_2. For the preparation of Dess-Martin periodidinane, see Dess, D., and J. Martin, Journal of Organic Chemistry, Vol. 48 pp. 4155 -4156 (1983). The slurry is stirred for 2 h and 58.7 g (0.24 M) of $Na_2S_2O_4$ in saturated $NaHCO_3$ added and the resulting solution stirred for 0.5 h. The phases are separated and the aqueous phase extracted with ethyl acetate. The combined organic phases are dried and the solvent removed to give 13.9 g of product that is used in the next reaction without purification. (Rf = 0.30 - 95/5 CH_2Cl_2/MeOH).


Adoc-D-t-Bug-Phe-Lys-C(O)CF_2C(O)-Leu-Arg-OMe (7) - To a solution of 13.4 g (0.012 M) of the residue from the previous reaction in 200 ml of dry DMF is added 6 g of palladium black and the resulting slurry degassed for 0.5 h. To this slurry is added 8 ml of 3.3 M HCl in dioxane. The reaction is placed on the Paar shaker and hydrogenated for 6 h. The slurry is filtered through Celite and the solvent removed. The residue is chromatographed on reverse phase HPLC (CH_3CN/H_2O) to afford 6.5 g product.

This Sheet Has Been Replaced





This Sheet Has Been Replaced



This Sheet Has Been Replaced

II. Synthesis of Adoc-D-t-Bug-Phe-Arg-C(O)-CF_2-C(O)-Ala-Arg-OMe

Adoc-D-t-Bug-OH - To a solution of 48.8 g (0.370 M) of D-t-butylglycine in 371 ml of 1N NaOH and 159 ml of dioxane at 0oC is added 33.9 g (0.40 M) of adamantyl fluoroformate in 371 ml of dioxane over 3 hours. The mixture is stirred at this temperature for 0.5 hours and the cooling bath is removed. The solution is stirred for 3 hours and the pH adjusted to 10 with 1N NaOH. This solution is extracted with EtOAc and the remaining aqueous phase acidified to pH = 3. This acidic solution is extracted with EtOAc, dried and the solvent is removed to afford 118 g of product that is homogeneous by TLC (94/5/1 - CH_2Cl_2/i-pa/AcOH), Rf = 0.90.

Adoc-D-t-Bug-Phe-OBn - To a solution of 115 g (0.37 M) of Adoc-D-t-But-OH and L-Phe-OBn.pTSA (0.36 M) in 1L CH_2Cl_2 is added 74.6 g (0.74 M) of TEA followed by the addition of 60.8 g (0.37 M) of DECP and the resulting solution is stirred overnight. The solvent is removed and the product is redissolved in EtOAc and washed successively with brine, 0.4N NCl, and saturated $NaHCO_3$. The solution is dried and the solvent is removed. The residue is chromatographed on silica to afford 186 g of product. TLC (85/15 - hexane/EtOAc) - Rf = 0.40.

Adoc-D-t-Bug-Phe-OH (1) - To a dry flask under N_2 is added 34.0 g 5% Pd on carbon. The flask is cooled to -78oC and 200 ml of dry degassed MeOH is added. To this slurry is added 8.8 ml of formic acid (88%) and this is followed by the addition of a mixture of 36.0 g of Adoc-D-t-Bug-Phe-OBn (0.066 M) in 100 ml of MeOH and 4.4 ml of formic acid. The cooling bath is removed and the slurry is stirred for 2 hours. At this time an additional 200 ml of 4.4% formic acid in MeOH is added and the slurry is stirred overnight. The mixture is filtered through glass fiber filter paper under N_2. The filtrate is refiltered through a bed of Celite to remove all traces of catalyst. Removal of the solvent yields 17.3 g of product.

α-Boc, Σ-Z-Lys-C(OH)-CF_2-C(O)-OEt (3) - To a solution of 79.2 g (1.21 M) of activated zinc in 0.600 L THF is added 2.36 g of ethyl bromodifluoroacetate and the solution is brought to reflux. As soon as reflux begins, a mixture of 236 g (1.16 M) of ethyl bromodifluoroacetate and 176 g (0.485 M) of 2 in 1.80 L of THF is added dropwise. The solution is refluxed for 1 hour, cooled and quenched with saturated $KHSO_4$. The quenched product is diluted with EtOAc and washed successively with saturated $KHSO_4$, saturated $NaHCO_3$ and brine. This sequence is followed by drying with $MgSO_4$ and

This Sheet Has Been Replaced



removal of solvent. The residue is chromatographed on silica to afford 68.0 g of pure product. Rf = 0.20 75/25 hexane/EtOAc.

α-Boc, Σ -Z-Lys-C(OH)-CF_2-C(O)-ONa (4) - To a solution of 1.00 g (0.002 M) of 3 in 10 ml of MeOH is added 2 ml of 1N NaOH. The solution is stirred at room temperature for 5 hours and the solvent is removed. The residue is dissolved in a small amount of water and the milky solution is extracted with ether. The water is removed from the aqueous phase and the residue thoroughly dried by placing on the vacuum pump for 24 hours. 0.069 g of product is recovered.

α-Boc, Σ -Z-Lys-C(OH)-CH_2-C(O)-Leu-Arg(NO_2)-OMe (5) - To a solution of 0.500 g (1.03 mmol) and 4 and 0.470 g (1.26 mmol) of Leu-Arg(NO_2)-OMe·HCl in 30 ml of DMB is added 0.019 g (1.26 mmol) of HOBt in one portion. This addition is followed by the dropwise addition of 0.260 g (1.26 mmol) of DCC in 3 ml DMF. Finally, 0.003 ml of DIPEA is added and the solution stirred for 48 hours. The DMF is removed in vacuo and the residue dissolved in EtOAc. The solution is transferred to a separatory funnel and washed with 0.1 N HCl, saturated $NaHCO_3$, and brine. The ethyl acetate layer is dried with $MgSO_4$, filtered and the solvent is removed. The residue is chromatographed on silica gel to afford 0.420 g of product. Rf = 0.68-90/10 CH_2Cl_2/MeOH.

Σ-Z-Lys-C(OH)-CH_2-C(O)-Leu-Arg(NO_2)-OMe.HCl - To a solution of 5 (17.3 g, 0.022 M) in 200 ml of dioxane is added 66.8 ml (0.220 m) of 3.3 M HCl in dioxane. The mixture is stirred for 1 hour. At this time an additional 5 ml of 3.3 M HCl in dioxane is added and the solution is stirred an additional 0.25 hour. The solvent is removed and the residue placed on the vacuum pump. The solid residue is triturated with ether under N_2 and filtered. 16.0 g of while solid is recovered.

Adoc-D-t-Buα-Phe-Lys(Z)-C(OH)-CH_2-C(O)-Leu-Arg(NO_2)-OMe - (6) To a solution of 16.0 g of the crude product from the previous reaction (0.022 M) and 11.1 g of 1 (0.024 M) in 300 ml of CH_2Cl_2 is added 6.69 ml (0.048 M) of TEA dropwise followed by the dropwise addition of 3.64 ml (0.024 M) of DECP. The solution is stirred overnight and the solvent is removed. The residue is dissolved in EtOAc and washed successively with 0.1 N HCl, saturated $NaHCO_3$, and brine. The solution is dried and the solvent is removed to afford 26.6 g that is chromatographed on silica to afford 12.7 g of pure product. Rf = 0.85, 90/10 - CH_2Cl_2/MeOH.

This Sheet Has Been Replaced



Adoc-D-t-Buq-Phe-Lys(Z)-C(O)CF_2C(O)-Leu-Arq(NO_2)-OMe - To a slurry of 14.3 g (0.034 M) of Dess-Martin periodinane in 1.15 L of CH_2Cl_2 is added 0.910 ml of dry pyridine, followed by the dropwise addition of 12.7 g (0.011 M) of the product from the previous reaction in 2.3 L of CH_2Cl_2. The slurry is stirred for 2 hours and 58.7 g (0.24 M) of $Na_2S_2O_4$ in saturated $NaHCO_3$ is added. The resulting solution is stirred for 0.5 hour. The phases are separated and the aqueous phase is extracted with ethyl acetate. The combined organic phases are dried and the solvent is removed to give 13.9 g of product that is used in the next reaction without purification (Rf = 0.30 - 95/5 CH_2Cl_2/MeOH).

Adoc-D-t-Buq-Phe-Lys-C(O)CF_2C(O)-Leu-Arq-OMe.TFA (7) - To a solution of 13.4 g (0.012 M) of the crude product from the previous reaction in 200 ml of dry DMF is added 6 g of palladium black. The resulting slurry is degassed for 0.5 hour. To this slurry is added 8 ml of 3.3 M Hcl in dioxane. The reaction is placed on the Paar shaker and hydrogenated for 6 hours. The slurry is filtered through Celite and the solvent is removed. The residue is chromatographed on reverse phase HPLC (CH_3CN/H_2O/TFA) to afford 6.5 g product.

α-Boc, γ-Bn, γ-Z-Orn-C(OH)-CF_2-C(O)OEt (9) - To a solution of 5.67 g (0.087 M) of activated zinc in dry THF is added 1.0 g of ethyl bromodifluoroacetate and the temperature is raised to reflux. At this time, 15.2 g (0.037 M) of -Bn, -Z- -Boc-Ornithinal 8 and 16.9 g (0.083 M) of ethyl bromodifluoroacetate in THF is added, and the slurry allowed to reflux for 0.5 hour. The solution is cooled and quenched with saturated $KHSO_4$. The solution is diluted with EtOAc and washed with saturated $KHSO_4$, saturated $NaHCO_3$ and brine. The solution is dried and the solvent removed. The residue is chromatographed on silica to give 6.58 g of product. Rf = 0.22 99/1 - CH_2Cl_2/ipa.

α-Boc, γ-Bn, γ-Z-Orn-C(OH)-CF_2-C(O)ONa - The ester 9 (0.965 g, 1.71 mmol) is dissolved in 15 ml MeOH and 1.71 ml of 1N NaOH is added. The solution is stirred for 3 hours and the solvent is removed. The residue is placed on the vacuum pump overnight and used in the next reaction as is (0.870).

α-Boc, γ-Bn, γ-Z-Orn-C(OH)-CF_2-C(O)-Ala-Arq(NO_2)-OMe 10 - To a solution of 0.870 g (1.56 mmol) of the crude product from the previous reaction and 0.579 g (1.90 mmol) of Ala-Arg(NO_2)-OMe·HCl in 80 ml of dry DMF is added 0.283 g (2.09 mmol) of HOBt, followed by the successive addition of 0.392 g (1.90 mmol) of DCC in 20 ml of DMF and 0.068 ml (0.39 mmol) of

This Sheet Has Been Replaced



DIPEA. The mixture is stirred for 18 hours at room temperature and the solvent is removed. The residue is dissolved in EtOAc and washed with 0.1N HCl, saturated $NaHCO_3$ and brine. The solution is dried and the solvent is removed. Chromatography on silica gel yields 0.520 g of product (Rf = 0.20 and 0.22).

α-Boc-Orn-C(OH)-CF_2-C(O)-Ala-Arg(NO_2)-OMe - The tripeptide 10 (0.338 g, 0.43 mmol) is dissolved in 10 ml of DMF and 0.15 g of palladium black is added. The slurry is degassed and 0.20 ml of 4.0 M HCl in dioxane is added. The mixture is hydrogenated on the Paar apparatus overnight and filtered through Celite. The residue is redissolved in MeOH and 0.20 g of $Pd(OH)_2$ is added. The slurry is hydrogenated on the Paar for 24 hours at 50 psi. The slurry is filtered through Celite and the solvent is removed. The residue is used in the next reaction without purification.

α-Boc-Arg(Z)$_2$-C(OH)-CF_2-C(O)-Ala-Arg(NO_2)-OMe (11) - The crude amine (0.220 g, 0.39 mmol) is dissolved in THF and 0.325 g (0.91 mmol) of N,N-Bis-Z-S-methyl-isothiourea is added. To this solution is added 0.150 g $Hg(OAc)_2$ (0.47 mmol) and the mixture is stirred overnight. The solution is diluted with EtOAc and washed successively with 0.1 HCl, saturated $NaHCO_3$ and brine. The organic phase is dried and the solvent is removed. The residue is chromatographed on silica to afford 0.170 g of product as a mixture of diastereomers. Rf = 0.60, 0.62 - 5% MeOH/CH_2Cl_2.

Adoc-D-t-Bug-Phe-Arg(Z)$_2$-C(OH)-CH_2-C(O)-Ala-Arg(NO_2)-OMe (12) - To a solution of 0.228 g (0.262 mmol) of 11 in 5 ml of dioxane is added 0.656 ml of HCl in dioxane. The solution is stirred for 1 hour and an additional 0.656 ml of HCl in dioxane is added. After stirring for an additional 1 hour, the solvent is removed and the residue is placed on the vacuum pump overnight. The hydrolysis product is dissolved in 5 ml of CH_2Cl_2 and combined with 0.120 g (0.262 mmol) of Adoc-D-t-Bug-Phe-OH. To this solution is added 0.073 ml (0.525 mmol) of TEA followed by the addition of 0.043 ml of DECP. The resulting mixture is stirred overnight. The solvent is removed and the residue chromatographed on silica to afford 0.086 g of product as a mixture of diastereomers. Rf = 0.65, 0.68 - 4% ipa/CH_2Cl_2.

Adoc-D-t-Bug-Phe-Arg(Z)$_2$-C(O)-CF_2-C(O)-Ala-Arg(NO_2)-OMe - To a slurry of 0.094 g (0.16 mmol) of Dess Martin periodinane in 5 ml CH_2Cl_2 is added 0.091 g (0.052 mmol) of 12 dissolved in 0.5 ml via syringe. The mixture is stirred for 0.5 hour and 0.385 g (1.55 mmol) of $Na_2S_2O_3$ in saturated $NaHCO_3$ is added. After 0.5 hour, the solution is extracted with EtOAc and the

This Sheet Has Been Replaced




organic phase is separated and dried with $MgSO_4$. Removal of the solvent yields 0.94 g crude product. Rf = 0.68, 0.71 - 4% ipa/CH_2Cl_2.

Adoc-D-t-Buq-Phe-Arq-C(O)-CF_2-C(O)-Ala-Arq-OMe·TFA (13) - The crude alcohol is dissolved in 2 ml DMF and 50 mg palladium black is added. The slurry is degassed and 0.040 ml of 4.0 M HCl in dioxane is added. The mixture is hydrogenated on the Paar apparatus for 20 hours at 50 psi. The slurry is filtered through Celite and the solvent is removed. The residue is chromatographed on a reverse phase column (CH_3CN/H_2O/TFA) to afford 0.044 g of pure product.


N,N'-bis-carbobenzyloxy-S-methyl-isothiourea (14) - To a solution of S-methyl-isothiourea sulfate dimer (5.00 g, 18.0 mmol) in cold (0oC) 1N NaOH (36.0 ml, 36.0 mmol) is added 11.3 ml (79.0 mmol) benzyl chloroformate and 79.0 ml (79.0 mmol) of 1N NaOH dropwise from separate addition funnels. The slurry is stirred for 0.5 hour at 0oC and 1.5 hour at room temperature. The solution is extracted with ethyl acetate, dried with magnesium sulfate and the filtrate evaporated. Chromatography on silica gel yields 2.62 g pure product. Rf = 0.40 (15% EtOAc/pet ether.

This Sheet Has Been Replaced






8




10




14

11



12

This Sheet Has Been Replaced



13

This Sheet Has Been Replaced

III. Synthesis of N-Terminating Linking Elements

Adac-D-t-Bug-Phe-OBn - To a solution of 5 ml of a 15% solution of phosgene in toluene under argon is added 0.140 ml TEA (1.00 mmol). This is followed by the addition of 0.112 g (0.750 mmol) of 1-adamantanamine in 1 ml toluene dropwise via syringe. The resulting slurry is stirred for 0.3 hour and the excess phosgene is removed by purging with argon. The solid is filtered off by suction through a sintered glass funnel and the solvent is removed. The residue is redissolved in dichloromethane and cooled to 0oC under argon. To this solution is added 0.140 ml TEA (1.00 mmol) followed by 0.100 g (0.250 mmol) d-t-Bug-Phe-OBn·HCl in one portion. The resulting solution is removed. Chromatography on silica yields 0.350 g product.

Hydrogenation of this compound followed by subsequent coupling with hydrolysis product of 11 yields compounds that are successfully elaborated to an alternative N-terminating linking element.

Methods for Testing Activity of the Compounds

The following non-limiting procedures are methods for testing the anti-inflammatory and/or analgesic activity of the difluoro pentapeptide derivatives of the subject invention.

Several enzyme inhibition assays are known to be predictive of anti-inflammatory activity for compounds. Such enzyme assays are useful for measuring the activity of compounds of the subject invention. Such enzyme assays include the following: porcine pancreatic kallikrein (PPK) - see references A, E and F; human urinary kallikrein (HUK) - see references E and F; human plasma kallikrein (HPK) - see references B and E; human plasmin (HP) - see references B and C; and urokinase (UK) - see reference D. The indicated references, which are hereby incorporated herein by reference, are the following:

(A) Lottenberg, R., U. Christensen, C.M. Jackson & P.L. Coleman, "Assay of Coagulation Proteases Using Peptide Chromogenic and Fluorogenic Substrates," Methods in Enzymology, Vol. 80, Academic Press, New York, NY (1981), pp. 341-361; (B) Geiger, R. "Kallikrein," Methods of Enzymatic Analysis, Vol. V, 3rd Edition, Bergmeyer, ed., Academic Press, New Nork, NY (1984), pp. 129-143; (C) Morris, J.P., S. Blatt, J.R. Powell, D.K. Strickland & F.J. Castellino, "Role of Lysine Binding Regions in the Kinetic Properties of Human Plasmin," Biochemistry, Vol. 20, No. 17 (August 18, 1981) p. 4811; (D) Wohl, R.C., L. Summaria & K.C. Robbins, "Kinetics of Activation of Human Plasminogen by Different Activator Species at pH 7.4 and 37oC" The Journal Biological

This Sheet Has Been Replaced



Chemistry, Vol. 255, No. 5 (March 10, 1980), pp. 2005-2013; (E) Okunishi, H., J. Burton & J. Spragg, "Specificity of Substrate Analogue Inhibitors of Human Urinary Kallikrein," Hypertension, Vol. 7, No. 3, Suppl. 1 (May-June, 1985), pp. I-72-I75; (F) Amundsen, E., J. Putter, P. Friberger, M. Knos, M. Larsbraten & G. Claeson, "Methods for the Determination of Glandular Kallikrein by Means of a Chromogenic Tripeptide Substrate," In Kinins-II part A, Fuji, S., et al., eds., Plenum Press, New York, NY (1979) pp. 83-95.

Another useful assay of activity is based on a method for determination of slow-binding enzyme inhibition disclosed in Imperiali, B. & R.H. Abeles, "Inhibition of Serine Proteases by Peptidyl Fluoromethyl Ketones," Biochemistry, Vol. 25 (1986) pp. 3760-3767. The method is modified as described below in order to study the slow binding inhibition of human plasmin, pig pancreatic kallikrein and human plasma kallikrein.

Reactions: The reaction mixtures contain 78 mM tris-HCl buffer, pH 7.4, 78 mM NaCl, 0.2 mg/ml bovine serum albumin, 0.2 mM S-2251 (D-Val-Leu-Arg-p-nitroanilide), 0.5 U/ml plasmin (1 mM), and variable concentrations of the test compound to be studied, in a total volume of 1 ml. The stock solution of plasmin is 1 U/ml in 50% glycerol. The absorbance change due to release of p-nitroaniline on enzymatic cleavage of S-2251 is monitored using an HP-8450 spectrophotometer system, set to measure $A^{400-410} - A^{470-490}$. The temperature is 30oC.

Calculations: K_{obsd} and v_s (steady state inhibited rate) are determined by fitting the progressive curve (first 20 min) to the integrated rate equation (i) using Labtech NotebookR software. Estimated k_1 is calculated for each run from v_s and the uninhibited rate v (equation ii), with (S) = 0.2 mM and K_m = 0.77 mM.

(i) $A = v_s t - ((v_s - v_o)/k_{obsd})(1 - \exp(-k_{obsd} t)) + A_o$

(ii) $k_1 = (I)/((v/v_s - 1)(1 + (s)/k_m))$

A plot of k_{obsd} vs test compound concentration for the different runs is then fit to a line, $y = mx + b$ (see equation iii); $k_{on} = m(1 + (S)/k_m)$ and $k_{off} = b$ are then calculated. Finally k_1 is calculated from equation iv.

(iii) $k_{obsd} = (I)/((v/v_s - 1)(1 + (S)/k_m))$

(iv) $k_1 = k_{off}/k_{on}$

Several in vivo assays are known to be predictive of the anti-inflammatory activity of compounds. Such in vivo assays are useful for measuring the activity of compounds of the subject invention. Such in vivo

This Sheet Has Been Replaced



assays are disclosed in the following references which are hereby incorporated herein by reference:

Winter, C.A., E.A. Risley, G.V. Nuss, "Carrageenin-Induced Edema in Hind Paw of the Rat as an Assay for Antiinflammatory Drugs," Proc. Soc. Exp. Biol., N.Y., Vol. 111 (1962), pp. 544-547; Vander Wende, C. & S. Margolin, "Analgesic Tests Based Upon Experimentally Induced Acute Abdominal Pain in Rats," Federal Proceedings., Vol. 15 (1956), p. 494; Blackham, A., J.W. Burns, J.B. Farmer, H. Radziwonik, J. Westwick, "An X-Ray Analysis of Adjuvant Arthritis in the Rat. The Effect of Prednisolone and Indomethacin," Agents and Actions, Vol. 7/1 (1977), p. 145-151; Winter, C.A. & G.W. Nuss, "Treatment of Adjuvant Arthritis in Rats with Anti-Inflammatory Drugs" Arthritis and Rheumatism, Vol. 9, No. 3 (June, 1966), pp. 394-404; and Francis, M.D., K. Hovancik & R.W. Boyce, "NE-58095: A Diphosphonate Which Prevents Bone Erosion and Preserves Joint Architecture in Experimental Arthritis," Int. J. Tiss. Reac., Vol. XI, No. 5 (1989), pp. 239-252.

Compositions and Methods of Using the Compounds

The following non-limiting examples exemplify contemplated compositions and uses for the subject invention.

Example I

Tablets are made by conventional procedures, each having the following composition:

Component	Quantity (mg)
Noc-D-t-Bug-Phe-Lys-C(O)CF_2C(O)-Leu-Arg-OMe	400
Microcrystalline cellulose	200
Pregelatinized starch	200
Povidone K-30	40
Magnesium stearate	20


One tablet is administered orally four times daily to a patient to alleviate inflammation in joints due to arthritis.

Example II

A lotion is made by conventional procedures, the lotion having the following composition:

Component	Quantity (%)
eBroc-D-tBug-Phe-Arg-C(O)CF_2C(O)-Gly-Arg-OMe	2.5
Glycerin	4.0
Methyl paraben	0.2
Propyl paraben	0.1

This Sheet Has Been Replaced



Carbopol 934	0.15
NaOH	0.46
Cetyl stearyl palmitate	1.0
Stearic acid	0.5
Lanolin fatty acids	0.5
Cetyl alcohol	3.0
Zantham gum	0.3
Sodium stearoyl-2-lactolate	0.75
Isopropyl myristate	2.0
Water	q.s.

One gram of the lotion is administered topically to the skin in the area of a burn twice daily to reduce inflammation and pain.

Example III

A solution is made by conventional means, each 2 ml of solution having the following composition:

Component	Quantity (mg)
Adoc-D-Phe-Phe-Lys-C(O)CF_2C(O)-D-Phe-Arg-OH	80
Benzalkonium chloride	40
Sterile aqueous saline solution	q.s.

A 2 ml dose of the solution is injected intramuscularly to a patient with arthritis to reduce inflammation and pain.

Example IV


A solution is made by conventional means, the solution having the following composition:

Component	Quantity (%)
Adoc-D-t-Bug-Phe-Arg-C(O)CF_2C(O)-Ala-Arg-OH	5.0
Benzalkonium chloride	0.02
Sodium carboxymethyl cellulose	0.01
Aqueous saline solution	q.s.

A 0.2 ml dose of the solution is administered by inhalation to a patient as needed to alleviate upper respiratory distress due to asthma.


While particular embodiments of the subject invention have been described, it will be obvious to those skilled in the art that various changes and modifications of the subject invention can be made without departing from the spirit and the scope of the invention. It is intended to cover, in the appended claims, all such modifications that are within the scope of this invention.

This Sheet Has Been Replaced



Claims

1. A compound, and pharmaceutically-acceptable salts and hydrates thereof, having the structure:



wherein:

(a) -X is selected from the group consisting of cyclic alkyl having from 4 to 15 carbon atoms; branched alkyl having at least two branches having from 6 to 15 carbon atoms; and aryl having from 6 to 15 carbon atoms;

(b) n is an integer from 0 to 2;

(c) -V- is selected from -OC(O)-, -N(Q)C(O)-, -N(Q)C(S)-, -C(O)-, $-SO_2-$ and -P(O)(OH)-;

d) -Q is hydrogen; or straight or branched chain alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to 6 carbon atoms; or -Q and -X are covalently linked forming a cyclic moiety which includes the nitrogen to which -Q is bonded and from 5 to 20 carbon atoms;

(e) -Z- is -O- or -NH-; when V is -OC(O)-, -Z- is -NH-;

(f) $-R^1$ is selected from straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^1$ is in either D or L configuration;

(g) $-R^2$ is selected from straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to 6 carbon atoms; cyclic alkyl,

This Sheet Has Been Replaced

* C L M S *

saturated or unsaturated with 1 or 2 double bonds, having from 3 to 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^2$ is in either D or L configuration;

(h) $-R^3$ is $-(CH_2)_m$-A-NH_2 or $-(CH_2)_m$-A-B-$C(NH_2)$=NH wherein m is an integer from 1 to 6, -A- is a covalent bond or p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-; and the carbon atom bonded to $-R^3$ is in L configuration;

(i) $-R^4$ is selected from straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^4$ is in either D or L configuration;

(j) $-R^5$ is $-(CH_2)_m$-A-NH_2 or $-(CH_2)_m$-A-B-$C(NH_2)$=NH wherein m is an integer from 1 to 6, -A- is a covalent bond or p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-; and the carbon atom bonded to $-R^5$ is in either D or L configuration; and

(k) -Y is hydrogen or methyl.

2. The compound of Claim 1 wherein:

(a) -V- is -OC(O)- or -NHC(O)-, and -Z- is -NH-;

(b) n is 0 or 1;

(c) m is an integer from 1 to 5;

(d) -X is saturated and unsubstituted bicyclo or tricyclo alkyl having from 8 to 12 carbon atoms, or unsubstituted aryl having from 8 to 12 carbon atoms; and

This Sheet Has Been Replaced

(e) $-R^1$, $-R^2$ and $-R^4$ are selected from straight chain or branched alkyl having from 1 to 5 carbon atoms, cyclic alkyl having from 3 to 6 carbon atoms, and benzyl.

3. The compound of Claim 1 or 2, wherein:

(a) $-R^3$ is $-(CH_2)_m$ -A- NH_2, wherein -A- is a covalent bond, and m is from 3 to 5;

(b) $-R^5$ is $(CH_2)_m$ -A-B-C (NH_2) = NH, wherein -A- is a covalent bond, -B- is -NH-, and m is from 2 to 4;

(c) -Y is methyl; and


(d) -V- is -OC(O)-.

4. The compound of Claim 1, 2 or 3 wherein:

(a) -X is tricyclo alkyl; and

(b) $-R^1$ and $-R^4$ are saturated and unsubstituted straight chain or branched alkyl, having from 1 to 4 carbon atoms, and $-R^2$ is benzyl.

5. A compound, and the pharmaceutically-acceptable salts and hydrates thereof, having the structure:



wherein:

(a) $-AA^1-$ is selected from t-butylglycyl, valyl, isoleucyl, leucyl, cyclohexylglycyl, cyclohexylalanyl, phenylalanyl, naphthylalanyl, tryptophyl and adamantylglycyl;

(b) $-AA^2-$ is selected from phenylalanyl, naphthylalanyl, cyclohexylalanyl, leucyl and isoleucyl;

This Sheet Has Been Replaced

* C L M S *

(c) -AA^3- is selected from arginyl, homoarginyl, lysyl, p-guanidinophenylalanyl, p-amidinophenylalanyl and p-aminophenylalanyl;

(d) -AA^4- is selected from phenylalanyl, naphthylalanyl, cyclohexylalanyl, leucyl and isoleucyl;

(e) -AA^5- is selected from arginyl, homoarginyl, lysyl, p-guanidinophenylalanyl, p-amidinophenylalanyl and p-aminophenylalanyl;

(f) -X^1 is selected from adamantyloxycarbonyl, adamantoyl, fluorenyl-methoxycarbonyl, adamantaneacetyl, adamantylaminocarbonyl, noradamantoyl, morpholinyl, homoadamantyloxycarbonyl, fenchyloxycarbonyl, isomenthyloxy-carbonyl, isopinocamphanyloxycarbonyl, 3,5-dimethyladamantyloxycarbonyl, endo-bornyloxycarbonyl, naphthyloxycarbonyl, and menthyloxycarbonyl; and

(g) -Y is hydrogen or methyl.

6. The compound of Claim 5 wherein:

(a) -AA^5- is arginyl;

(b) -AA^3- is lysyl; and

(c) -X^1 is selected from adamantyloxycarbonyl, endo-bornyloxycarbonyl and isopinocamphanyloxycarbonyl.

7. The compound of Claim 5 or 6 wherein:

(a) -AA^1- is selected from t-butylglycyl, valyl, and isoleucyl;

(b) -AA^2 is phenylalanyl; and

(c) -AA^4- is leucyl.

8. The compound of Claim 5, 6 or 7 wherein:

(a) -AA^1- is t-butylglycyl;

(b) -X^1 is adamantyloxycarbonyl; and

This Sheet Has Been Replaced



* C L M S *

(c) -Y is methyl.

9. A pharmaceutical composition comprising a compound of any of Claims 1-8 and a pharmaceutically-acceptable carrier.

10. Use of a compound of any of Claims 1-8 for manufacture of a medicament for treating inflammation or pain.

This Sheet Has Been Replaced

9608179-9 IPOS

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : C07K 7/06, A61K 37/02

A1

(11) International Publication Number: WO 94/14842

(43) International Publication Date: 7 July 1994 (07.07.94)

(21) International Application Number: PCT/US93/12349

(22) International Filing Date: 16 December 1993 (16.12.93)

(30) Priority Data:
07/995,217 22 December 1992 (22.12.92) US

(71) Applicant: THE PROCTER & GAMBLE COMPANY [US/US]; One Procter & Gamble Plaza, Cincinnati, OH 45202 (US).

(72) Inventor: MCIVER, John, McMillan; 832 Cedarhill Drive, Cincinnati, OH 45240 (US).

(74) Agents: REED, T., David et al.; The Procter & Gamble Company, 5299 Spring Grove Avenue, Cincinnati, OH 45217.(US).


(81) Designated States: AU, BB, BG, BR, BY, CA, CZ, FI, HU, JP, KP, KR, KZ, LK, LV, MG, MN, MW, NO, NZ, PL, RO, RU, SD, SK, UA, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

Published
With international search report.
Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: DIFLUORO-PENTAPEPTIDE DERIVATIVE ANTI-INFLAMMATORY AGENTS

(57) Abstract

The subject invention involves anti-inflammatory compounds having structure (I), wherein: (a) -X is cyclic alkyl, having from 4 to 15 carbon atoms; branched alkyl, having from 6 to 15 carbon atoms; or aryl having from 6 to 15 carbon atoms; (b) n is an integer from 0 to 2; (c) -V- is -OC(O)-, -N(Q)C(O)-, -N(Q)C(S)-, -C(O)-, -SO2- or -P(O)(OH)-; (d) -Q is hydrogen; or straight or branched chain alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to 6 carbon atoms; or -Q and -X are covalently linked forming a cyclic moiety which includes the nitrogen to which -Q is bonded and from 5 to 20 carbon atoms; (e) Z is -O- or -NH-; when V is -OC(O)-, -Z- is -NH-; (f) -R1 is a hydrophobic moiety; (g) -R2 is a hydrophobic moiety; (h) -R3 is -(CH2)m-A-NH2 or -(CH2)m-A-B-C(NH2)=NH wherein m is an integer from 1 to about 6, -A- is a covalent bond or p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-; (i) -R4 is a hydrophobic moiety; (j) -R5 is -(CH2)m-A-NH2 or -(CH2)m-A-B-C(NH2)=NH wherein m is an integer from 1 to about 6, -A- is a covalent bond or p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-; and (k) -Y is hydrogen or methyl. The subject invention also involves pharmaceutical compositions comprising the above compounds, and methods for treating inflammation or pain using such compounds and compositions.



(I)

REPLACED SHEETS	SG
☑ Description	: paged 1 to paged 28
☑ Claim(s)	: paged 29 to paged 33
☐ Drawing(s)	: paged ..... to paged .....
☑ Abstract	: paged 34 to paged 35
DATE OF RECEIPT	.....


This Sheet Has Been Replaced

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo			SE	Sweden
CH	Switzerland	KR	Republic of Korea	SI	Slovenia
CI	Côte d'Ivoire	KZ	Kazakhstan	SK	Slovakia
CM	Cameroon	LI	Liechtenstein	SN	Senegal
CN	China	LK	Sri Lanka	TD	Chad
CS	Czechoslovakia	LU	Luxembourg	TG	Togo
CZ	Czech Republic	LV	Latvia	TJ	Tajikistan
DE	Germany	MC	Monaco	TT	Trinidad and Tobago
DK	Denmark	MD	Republic of Moldova	UA	Ukraine
ES	Spain	MG	Madagascar	US	United States of America
FI	Finland	ML	Mali	UZ	Uzbekistan
FR	France	MN	Mongolia	VN	Viet Nam
GA	Gabon				

This Sheet Has Been Replaced





960879-9

REPLACEMENT SHEETS SG
☑ Description : paged 1 to paged 28
☑ Claim(s) : paged 29 to paged 33
☐ Drawing(s) paged ..... to paged .....
☑ Abstract : paged 34 to paged .....

DATE OF RECEIPT 26/2/97

- 1 -

REPLACED SHEETS SG
☐ Description : paged ..... to paged .....
☑ Claim(s) : paged 29 to paged 33
☐ Drawing(s) : paged ..... to paged .....
☐ Abstract : paged ..... to paged .....
DATE OF RECEIPT .....

DIFLUORO PENTAPEPTIDE DERIVATIVE ANTI-INFLAMMATORY AGENTS

TECHNICAL FIELD

The subject invention relates to novel peptide derivatives which are useful as anti-inflammatory agents.

BACKGROUND OF THE INVENTION

Certain polypeptide derivatives having various biological activities are known. The following references disclose such polypeptide derivatives: U.S. Patent No. 4,242,329 issued to Claeson, Simonsson & Arielly on December 30, 1980; U.S. Patent No. 4,316,889 issued to Bajusz, Hasenohrl, Barabas & Bagdy on February 23, 1982; U.S. Patent No. 4,399,065 issued to Bajusz, Hasenohrl, Barabas & Bagdy on August 16, 1983; U.S. Patent No. 4,401,594 issued to Umezawa, Takeuchi, Aoyagi, Ishii, Saino & Someno on August 30, 1983; U.S. Patent No. 4,478,745 issued to Bajusz, Hasenohrl, Barabas & Bagdy on October 23, 1984; U.S. Patent No. 4,528,133 issued to Kasafirek, Fric, Slaby & Robalova on July 9, 1985; U.S. Patent No. 4,596,789 issued to Dutta, Stein, Trainor & Wildonger on June 24, 1986; U.S. Patent No. 4,623,639 issued to Hassall, Johnson & Roberts on November 18, 1986; U.S. Patent No. 4,703,036 issued to Bajusz, Hasenohrl, Bagdy, Barabas, Dioszegi, Fittler, Jozsa, Horvath & Jozst on October 27, 1987; U.S. Patent No. 4,880,780 issued to Trainor & Stein on November 14, 1989; U.S. Patent No. 4,883,863 issued to Abe, Yaginuma, Nagasawa & Kuroiwa on November 28, 1989; U.S. Patent No. 4,902,781 issued to Mizoue, Okazaki, Hanada, Omura & Amamoto on February 20, 1990; U.S. Patent No. 4,923,890 issued to Trainor & Stein on May 8, 1990; U.S. Patent No. 5,066,643 issued to Abeles & Gelb on November 19, 1991; PCT Patent Application No. 84/00365 of Galpin & Wilby, published February 2, 1984; PCT Patent Application No. 87/02675 of Hoover, published May 7, 1987; Japanese Patent Application No. 47-30618 of Toray Inds. Inc., published November 9, 1972; Japanese Patent Application No. 57-145846 of Mitsubishi Chem. Ind. KK, published July 19, 1974; Japanese Patent Application No. 58-164563 of Daiichi Kagaku Yaku, published March 25, 1982; European Patent Application No. 0275101 of Merrell Dow Pharmaceuticals, Inc., published July 20, 1988; Aoyagi, Miyata, Nanbo, Kojima, Matsuzaki, Ishizuka, Takeuchi & Umezawa, "Biological Activities of Leupeptins," The Journal of Antibiotics, Vol.

REPLACEMENT SHEET
26 FEB 1997

XXII, No. 11 (Nov. 1969), pp. 558-568; Bajusz, Barabas, Tolnay, Szell & Bagdy, "Inhibition of Thrombin and Trypsin by Tripeptide Aldehydes," Int. J. Peptide Protein Res., Vol. 12 (1978), pp. 217-221; Gaal, Bacsy & Rappay, "Cationic Ferritin Uptake by Cultured Anterior Pituitary Cells Treated with the Proteinase Inhibitor, BOC-DPhe-Phe-Lys-H," Histochemistry, Vol. 88 (1988), pp. 401-406; Makara, Rappay, Garamvolgyi, Nagy, Danko & Bajusz, "The Tripeptide Aldehyde, Boc-DPhe-Phe-Lysinal, is a Novel $Ca2^+$ Channel Inhibitor in Pituitary Cells," European Journal of Pharmacology, Vol. 151 (1988), pp. 147-149; Nagy, Makara, Horvath, Rappay, Kurcz & Bajusz, "Tripeptide Aldehyde Protease Inhibitors May Depress in Vitro Prolactin and Growth Hormone Release," Endocrinology, Vol. 116, No. 4 (1895), pp. 1426-1432; Rappay, Makara, Bajusz & Nagy, "Various Proteinase Inhibitors Decrease Prolactin and Growth Hormone Release by Anterior Pituitary Cells," Life Sciences, Vol. 36 (1985), pp. 549-555.

It is an object of the subject invention to provide novel difluoro pentapeptide derivatives which are useful as anti-inflammatory agents.


It is a further object of the subject invention to provide novel difluoro pentapeptide derivatives which reduce bone resorption and/or heterotopic bone formation associated with rheumatoid arthritis.

It is also an object of the subject invention to provide novel pharmaceutical compositions comprising certain difluoro pentapeptide derivatives.

It is also an object of the subject invention to provide methods for treating diseases characterized by inflammation using the subject compounds.

SUMMARY OF THE INVENTION

The subject invention involves anti-inflammatory compounds having the following structure:



wherein:

(a) -X is selected from the group consisting of cyclic alkyl having from 4 to about 15 carbon atoms; branched alkyl, with at least two branches, having from 6 to about 15 carbon atoms; and arylalkyl having from 6 to about 15 carbon atoms;

(b) n is an integer from 0 to about 2;

REPLACEMENT SHEET
26 FEB 1997

(c) -V- is selected from the group consisting of -OC(O)-, -N(Q)C(O)-, -N(Q)C(S)-, -C(O)-, -SO_2- and -P(O)(OH)-;

(d) -Q is hydrogen; or straight or branched chain alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; or -Q and -X are covalently linked forming a cyclic moiety which includes the nitrogen to which -Q is bonded and from about 5 to about 20 carbon atoms;

(e) Z is -O- or -NH-; when V is -OC(O)-, -Z- is -NH-;

(f) $-R^1$ is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^1$ is in either D or L configuration;

(g) $-R^2$ is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^2$ is in L configuration;

(h) $-R^3$ is $-(CH_2)_m$-A-NH_2 or $-(CH_2)_m$-A-B-$C(NH_2)$=NH wherein m is an integer from 1 to about 6, -A- is a covalent bond or p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-; and the carbon atom bonded to $-R^3$ is in L configuration;

(i) $-R^4$ is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^4$ is in L configuration;

(j) $-R^5$ is $-(CH_2)_m$-A-NH_2 or $-(CH_2)_m$-A-B-$C(NH_2)$=NH wherein m is an integer from 1 to about 6, -A- is a covalent bond or p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-; and the carbon atom bonded to $-R^5$ is in D or L configuration; and

(k) -Y is hydrogen or methyl.

The subject invention also involves pharmaceutical compositions comprising the above compounds, and methods for treating inflammation or pain using such compounds and compositions.

REPLACEMENT SHEET
26 FEB 1997

DETAILED DESCRIPTION OF THE INVENTION

The term "alkyl," as used herein, unless otherwise indicated, means carbon-containing chains which may be straight, branched or cyclic; which may be saturated or unsaturated; and which may be unsubstituted or substituted. As used herein, "cyclic alkyl" may have all or only a portion of the total number of carbon atoms indicated as being in the alkyl group in the cyclic ring itself. Cyclic alkyl includes monocycloalkyl, bicycloalkyl and/or tricycloalkyl.

Preferred alkyl are as noted in this paragraph unless provided otherwise in particular instances. Preferred alkyl are saturated. Preferred alkyl are unsubstituted. Preferred alkyl substituents are selected from halo, amino, hydroxy, alkoxy, cyano, nitro, aryl and trifluoromethyl. As used herein, "alkoxy" is -O-alkyl. It is preferred that substituted alkyl be mono-, di- or trisubstituted, especially monosubstituted.

The term "aryl," as used herein, unless otherwise specified, means aryl rings which may be unsubstituted or substituted. Preferred aryl is as noted in this paragraph, unless provided otherwise in particular instances. Preferred aryl are phenyl and naphthyl, especially phenyl. Preferred aryl are mono-, di-, tri- or unsubstituted; more preferred aryl are monosubstituted or unsubstituted, especially unsubstituted. Preferred aryl substituents include alkyl, halo, amino, hydroxy, alkoxy, cyano, nitro and trifluoromethyl.

The term "arylalkyl," as used herein, means alkyl substituted with aryl. A preferred arylalkyl is arylmethyl.

The term "aliphatic" as used herein, means open-chain carbon-containing portions of arylalkyls.

The following abbreviations are used herein. Abbreviations for amino acids may refer to the amino acid itself, or more often to an amino acid moiety that is part of a peptide or peptide derivative structure.

Arg	Arginyl
Orn	Ornithyl
Leu	Leucyl
Gly	Glycyl
t-Bug	t-Butylglycyl
Phe	Phenylalanyl
Adoc	Adamantyloxycarbonyl
Z	Benzyloxycarbonyl
Boc	t-Butyloxycarbonyl
DECP	Diethylcyanophosphonate


REPLACEMENT SHEET
26 FEB 1997

Me	Methyl
Et	Ethyl
TEA	Triethylamine
DMF	Dimethylformamide
TFA	Trifluoroacetate
ipa	Isopropanol
Bn	Benzyl
Lys	Lysyl
Ac	Acetyl
TLC	Thin Layer Chromatography
TSA	Toluenesulfonic acid
THF	Tetrahydrofuran
HOBt	Hydroxybenzotriazole
DCC	Dicyclohexylcarbodiimide
DIPEA	N,N-Diisopropylethylamine
Chg	Cyclohexylglycyl
Cha	Cyclohexylalanyl
Nal	Naphthylalanyl
Trp	Tryptophyl
Adg	Adamantylglycyl
pGphe	p-Guanidinophenylalanyl
3,5-Dmadoc	3,5-Dimethyladamantyloxycarbonyl
eBroc	endo-Bornyloxycarbonyl
Noc	Naphthyloxycarbonyl
Moc	Menthyloxycarbonyl
Ad	Adamantoyl
Fmoc	Fluorenylmethoxycarbonyl
Ada	Adamantaneacetyl
Adac	Adamantylaminocarbonyl
Mnoc	Morpholinyl
Norad	Noradamantoyl
Hadoc	Homoadamantyloxycarbonyl
Foc	Fenchyloxycarbonyl
Imoc	Isomenthyloxycarbonyl
Ipoc	Isopinocamphanyloxycarbonyl
AC	acetyl
DMF	dimethylformamide

REPLACEMENT SHEET
26 FEB 1997

HPLC high pressure liquid chromatography

The novel anti-inflammatory compounds of the subject invention are those having the following chemical structure:



In Structure (1), n is an integer of from 0 to about 2; n is preferably 0 or 1.

In Structure (1), $-R^1$ is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl, the aliphatic portion being saturated and having 1 or 2 carbon atoms. Preferred $-R^1$ is saturated alkyl. Preferred alkyl are unsubstituted, or are arylalkyl, especially benzyl and naphthal. Preferred cyclic alkyl are cyclic C_3-C_8 (more preferably C_5-C_6) methyl or adamantylmethyl. Preferred $-R^1$ is hydrophobic, preferably with the hydrophobicity concentrated close to the carbon atom to which $-R^1$ is bonded. Specific examples of preferred $-R^1$ include t-butyl, 1,1-dimethylpropyl, i-propyl, i-butyl, s-butyl, neo-pentyl, cyclohexyl, cyclohexylmethyl, adamantyl, adamantylmethyl, benzyl, naphthal; most preferred $-R^1$ is t-butyl.

In Structure (1), the carbon to which $-R^1$ is bonded is in either D or L, preferably D, configuration. The structure $-Z-CH(R^1)-C(O)-$ is an amino acid moiety (hereinafter -AA1-) when -Z- is -NH; preferred amino acid moieties for -AA1- include t-Bug, Val, Ile, Leu, Chg, Cha, Phe, Nal, Trp and Adg; more preferred are t-Bug, Val and Ile; most preferred -AA1- is t-Bug.

In Structure (1), $-R^2$ is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl, the aliphatic portion being saturated and having 1 or 2 carbon atoms. Preferred branched or cyclic alkyl are saturated. Preferred branched or cyclic alkyl are unsubstituted. Preferred branched alkyl have from 3 to 5 carbon atoms; most preferred branched alkyl is i-butyl. Preferred cyclic alkyl have a C_3-C_7, more preferably C_5-C_6, cyclic ring bonded to a methylene, ethylene or n-propylene (preferably methylene or ethylene, more preferably methylene) which is bonded to the carbon atom in Structure (1) to which $-R^2$ is bonded. Preferred arylalkyl

REPLACEMENT SHEET
20 FEB 1997

are benzyl, p-hydroxybenzyl and naphthal. More preferred -R^2 is selected from i-propyl, i-butyl, s-butyl, cyclohexylmethyl, benzyl and naphthal; most preferred is benzyl.

In Structure (1), the carbon atom to which -R^2 is bonded is in D or L configuration. The structure -NH-CH(R^2)-C(O)- is an amino acid moiety (hereinafter -AA^2-); preferred amino acid moieties for -AA^2- include Phe, Nal, Cha, Leu and Ile; most preferred -AA^2- is Phe.

In Structure (1), -R^3 is -$(CH_2)_m$-A-NH_2 or -$(CH_2)_m$-A-B-C(NH_2)=NH, wherein m is an integer of from 1 to about 6, -A- is a covalent bond, p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-. When -A- is a covalent bond, m is preferably 2-5, more preferably 2-4, more preferably still 3 or 4. When -A- is p-phenyl or p-cyclohexyl, m is preferably 1-4, more preferably 1-3, more preferably still 1. -A- is preferably a covalent bond. -B- is preferably -NH-. More preferred -R^3 is 3-guanidino-n-propyl or 4-amino-n-butyl.

In Structure (1), the carbon to which -R^3 is bonded is in D or L configuration, preferably in L configuration. The structure -NH-(CH(R^3)-C(O)- is an amino acid moiety (hereinafter -AA^3-); preferred amino acid moieties for -AA^3- include Arg, Lys, homo-Arg, p-amidino-phe, p-amino-phe and p-Gphe; most preferred -AA^3- is Lys.

In Structure (1), -R^4 is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl, the aliphatic portion being saturated and having 1 or 2 carbon atoms. Preferred branched or cyclic alkyl are saturated and unsubstituted. Preferred branched alkyl have from 3 to 5 carbon atoms; most preferred branched alkyl is i-butyl. Preferred cyclic alkyl have a C_3-C_7, more preferably C_5-C_6 cyclic ring bonded to a methylene, ethylene or n-propylene (preferably methylene or ethylene, more preferably methylene) which is bonded to the carbon atom in Structure (1) to which -R^4 is bonded. Preferred arylalkyl are benzyl, p-hydroxybenzyl and naphthal. More preferred -R^4 is selected from i-propyl, i-butyl, s-butyl, cyclohexylmethyl, benzyl and naphthal; most preferred is benzyl.

In Structure (1), the carbon atom to which -R^4 is bonded is in D or L configuration. The structure -NH-CH(R^4)-C(O)- is an amino acid moiety (hereinafter -AA^4-); preferred amino acid moieties for -AA^4- include Phe, Nal, Cha, Leu and Ile; most preferred -AA^4- is Leu.

REPLACEMENT SHEET
26 FEB 1997

In Structure (1), $-R^5$ is $-(CH_2)_m$-A-NH_2 or $-(CH_2)_m$-A-B-$C(NH_2)$=NH, wherein m is an integer from 1 to about 6, -A- is a covalent bond, p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-. When -A- is a covalent bond, m is preferably 2-5, more preferably 2-4, more preferably still 3 or 4. When -A- is p-phenyl or p-cyclohexyl, m is preferably 1-4, more preferably 1-3, more preferably still 1. -A- is preferably a covalent bond. -B- is preferably -NH-. More preferred $-R^5$ is 3-guanidino-n-propyl or 4-amino-n-butyl.

In Structure (1), the carbon to which $-R^5$ is bonded is in D or L configuration, preferably in L configuration. The structure -NH-CH(R^5)-C(O)- is an amino acid moiety (hereinafter $-AA^5$); preferred amino acid moieties for $-AA^5$ include Arg, homo-Arg, Lys, p-amidino-Phe, p-amino-Phe and p-Gphe; most preferred $-AA^5$- is Arg.

In Structure (1), -Y is hydrogen or methyl.

In Structure (1), -Z- is -O- or -NH-. Preferred -Z- is -NH-.

In Structure (1), -V- is selected from -OC(O)-, -N(Q)C(O)-, -N(Q)C(S)-, -C(O)-, $-SO^2-$ and -P(O)(OH)-. Preferred -V- is selected from -OC(O)-, -N(Q)C(O)-, -N(Q)C(S)-, and -C(O)-. More preferred -V- is -OC(O)- or -N(Q)C(O)-. Most preferred -V- is -OC(O)-. When -V- is -OC(O)-, -Z- is -NH-.

In Structure (1), -X is selected from cyclic alkyl, having from 4 to about 15 carbon atoms; branched alkyl having from 6 to about 15 carbon atoms and at least 2 branches; and aryl, having from 6 to about 15 carbon atoms. Preferred -X has from 6 to 12 carbon atoms; more preferred -X has from 8 to 10 carbon atoms. Preferred alkyl portions of -X are saturated. Preferred -X is unsubstituted, or substituted with unsubstituted alkyl or aryl. Preferred cyclic alkyl are monocycloalkyl, bicycloalkyl, and tricycloalkyl, more preferred are bicycloalkyl and tricycloalkyl, especially tricycloalkyl. Preferred cycloalkyl have 5 or 6 carbon atoms, more preferably 6 carbon atoms, in each cyclic ring. A highly preferred -X is adamantyl. Preferred aryl -X are fluorenyl, naphthyl and phenyl, unsubstituted or substituted with alkyl. Particularly preferred aryl -X include naphthyl and fluorenyl. When -X is aryl or cyclic alkyl, n is preferably 1.

In Structure (1) when -V- is -N(Q)C(O)- or -N(Q)C(S)-, -Q is selected from hydrogen; straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; or -Q and -X are covalently linked forming a cyclic moiety which includes the nitrogen to which -Q is bonded and from about 5 to about 20 carbon atoms. Preferred -Q-X- has from 6 to 15 carbon atoms; more preferred -Q-X- has from 8 to 12 carbon atoms.

REPLACEMENT SHEET
26 FEB 1997

Preferred -Q-X- is unsubstituted or substituted with unsubstituted alkyl or aryl. Preferred cyclic



- N - X is monocyclic, bicyclic or tricyclic; more preferred is monocyclic. Most preferred -Q is hydrogen.

In Structure (1), preferred X-$(CH^2)_n$-V- (hereinafter X'-) include Adoc. Ad, Fmoc, Ada, Adac, Mnoc, Norad, Hadoc, Foc, Imoc, Ipoc, 3,5-Dmadoc, eBroc, Noc and Moc; more preferred X' is Adoc, Ipoc and eBroc; most preferred -X' is Adoc.

Preferred anti-inflammatory compounds of the subject invention include pharmaceutically-acceptable salts of the above compounds. Particularly preferred salts include salts of addition formed between a strong acid and the above compounds. Particularly preferred are such salts of addition where either $-R^3$ or $-R^5$, or both, are protonated, resulting in a positive charge on the $-R^3$ and R^5 moieties. Preferred compounds of the subject invention are often associated with one or more molecules of water in the form of hydrates.

Preferred anti-inflammatory compounds of the subject invention include the following compounds:

Adoc-D-tBug-Phe-Lys-C(O)CF_2C(O)-Leu-Arg-OMe
Adoc-D-tBug-Phe-Lys-C(O)CF_2C(O)-Gly-Arg-OMe
Adoc-D-tBug-Phe-Lys-C(O)CF_2C(O)-D-Phe-Arg-OMe
Adoc-D-tBug-Phe-Lys-C(O)CF_2C(O)-Leu-D-Arg-OMe
Mnoc-D-Phe-Phe-Arg-C(O)-CF_2C(O)-Leu-Lys-OMe
eBroc-D-t-Bug-Phe-pGphe-C(O)CF_2C(O)-Ala-Arg-OMe
Ipoc-D-Nal-Phe-Arg-C(O)CF_2C(O)-Gly-Arg-OMe
Adoc-D-Phe-Phe-Lys-C(O)CF_2C(O)-Leu-Arg-OH
Noc-D-t-Bug-Phe-Lys-C(O)CF_2C(O)-Ala-Lys-OMe

Pharmaceutical Compositions

Pharmaceutical compositions of the subject invention comprise a safe and effective amount of a difluoro pentapeptide derivative as defined hereinabove and a pharmaceutically-acceptable carrier. Such compositions typically comprise from about 0.1% to about 95% by weight of the difluoro pentapeptide derivative, preferably from about 1% to about 90% and more preferably from about 5% to about 75%.

The term "pharmaceutically-acceptable carrier," as used herein, means one or more compatible solid or liquid filler diluents or encapsulating

REPLACEMENT SHEET
26 FEB 1997

substances which are suitable for administration to a human or lower animal. The term "compatible," as used herein, means that the components of the pharmaceutical compositions are capable of being commingled with the difluoro pentapeptide derivative of the subject invention, and with each other, in a manner such that there is no interaction which would substantially reduce the pharmaceutical efficacy of the composition under ordinary use situations. Pharmaceutically-acceptable carriers must, of course, be of sufficiently high purity and sufficiently low toxicity to render them suitable for administration to the human or lower animal being treated.

Some examples of substances which can serve as pharmaceutically-acceptable carriers are sugars, such as lactose, glucose and sucrose; starches, such as cornstarch and potato starch; cellulose and its derivatives, such as sodium carboxymethylcellulose, ethylcellulose and cellulose acetate; powdered tragacanth; malt; gelatins; talc; stearic acid; magnesium stearate; calcium sulfate; vegetable oils, such as peanut oil, cottonseed oil, sesame oil, olive oil, corn oil and oil of theobroma; polyols, such as propylene glycol, glycerin, sorbitol, mannitol, and polyethylene glycol; alginic acid; pyrogen-free water; isotonic saline; phosphate buffer solutions; cocoa butter (suppository base); emulsifiers, such as TweensR; wetting agents and lubricants, such as sodium lauryl sulfate; coloring agents; flavoring agents; stabilizers; antioxidants; preservatives; as well as other non-toxic compatible substances used in pharmaceutical formulations. Other compatible pharmaceutical additives and actives may be included in the pharmaceutically-acceptable carrier for use in the compositions of the subject invention.

The pharmaceutically-acceptable carrier employed in conjunction with the difluoro pentapeptide derivative of the subject invention is used at a concentration sufficient to provide a practical size to dosage relationship. The pharmaceutically-acceptable carrier, in total, typically comprises from about 5% to about 99.9% by weight of the pharmaceutical compositions of the subject invention, preferably from about 10% to about 99%, and more preferably from about 25% to about 95%.

Total single dosages of the difluoro pentapeptide derivatives of the subject invention in pharmaceutical compositions are generally from about 1 mg to about 1000 mg, preferably from about 50 mg to about 800 mg, more preferably from about 100 mg to about 500 mg.

The choice of a pharmaceutically-acceptable carrier to be used in conjunction with the difluoro pentapeptide derivatives of the subject invention is

REPLACEMENT SHEET
26 FEB 1997

determined by the way the active is to be administered. Preferred modes of administering the actives of the subject invention are by injection, ingestion, inhalation and topically.

Methods and materials for manufacturing injectables can be found in Remington's Pharmaceutical Sciences, 17th ed., 1985, Chapter 85, p. 1518, the disclosures of which are incorporated herein by reference in their entirety. Generally, three types of injectable dosage forms are preferred: (1) aqueous solutions, (2) non-aqueous solutions; and (3) emulsions. Injectable dosage forms typically contain from about 0.001 mg/ml to about 10 mg/ml, preferably from about 0.4 mg/ml to about 0.6 mg/ml, of a compound of the subject invention. Preferred injectable dosage compositions of the subject invention include sterile aqueous solutions or emulsions of pH from about 3 to about 8 (more preferably of pH from about 4 to about 6).

Preferred pharmaceutical carriers in which the difluoro pentapeptide derivatives of the subject invention can be administered by ingestion are proteinoid microcapsules disclosed in the following patents: U.S. Patent No. 4,895,725 issued to Kantor, Steiner & Pack on January 23, 1990; U.S. Patent No. 4,925,673 issued to Steiner & Rosen on May 15, 1990; U.S. Patent No. 4,976,968 issued to Steiner on December 11, 1990; and U.S. Patent No. 4,983,402 issued to Steiner on January 8, 1991; all four patents are hereby incorporated in their entirety by reference.

Methods for Producing Anti-inflammatory Activity and Analgesia

The subject invention also encompasses methods of producing anti-inflammatory activity and/or analgesia in humans or lower animals through administering, to the human or lower animal in need of such treatment, a safe and effective amount of a difluoro pentapeptide derivative of the subject invention. The amount can be given in a single dose or multiple doses repeatedly over the course of the treatment. While dosages higher than those described herein are effective to reduce inflammation and produce analgesia, care must be taken in some individuals to prevent adverse side effects. The difluoro pentapeptide derivatives and compositions of the subject invention can be used to reduce inflammation in various disorders at the deeper structures, muscles, tendons, bursa and joints associated with disease and trauma, to treat and prevent pain.

The preferred modes of administering the difluoro pentapeptide derivatives of the subject invention are by injection, ingestion, inhalation and topically. Specific modes of administration include, without limitation, oral

REPLACEMENT SHEET
26 FEB 1997

ingestion; injection, such as intramuscular, intravenous, intraperitoneal, intradermal and subcutaneous; inhalation; and topically, such as transdermally, orally, mucosally and sublingually.

The term "safe and effective amount," as used herein, means an amount of a difluoro pentapeptide derivative or composition high enough to significantly positively modify the condition to be treated, but low enough to avoid serious side effects (at a reasonable benefit/risk ratio), within the scope of sound medical judgement. A safe and effective amount of a difluoro pentapeptide derivative or composition will vary with the particular condition being treated, the age and physical condition of the patient being treated, the severity of the condition, the duration of the treatment, the nature of concurrent therapy, the specific active employed, the particular pharmaceutically-acceptable carrier utilized, and like factors within the knowledge and expertise of the attending physician.

Preferred daily dosages of the difluoro pentapeptide derivatives of the subject invention range from about 0.1 mg/kg of body weight to about 500 mg/kg of body weight, more preferably from about 1 mg/kg to about 100 mg/kg, still more preferably from about 2 mg/kg to about 50 mg/kg. From 1 to about 4 single dosages per day may be administered, more preferably from about 2 to about 3 single dosages per day. Preferred amounts of the difluoro pentapeptide derivatives administered by injection are from about 0.1 mg/kg/day to about 50 mg/kg/day, more preferably from about 1 mg/kg/day to about 10 mg/kg/day. Preferred amounts of the difluoro pentapeptide derivatives administered by oral ingestion are from about 1 mg/kg/day to about 500 mg/kg/day, more preferably from about 5 mg/kg/day to about 100 mg/kg/day. Preferred amounts of the difluoro pentapeptide derivatives administered by inhalation are from about 0.1 mg/kg/day to about 500 mg/kg/day, more preferably from about 5 mg/kg/day to about 100 mg/kg/day. Preferred amounts of the difluoro pentapeptide derivatives administered topically are from about 1 mg/kg/day to about 500 mg/kg/day, more preferably from about 50 mg/kg/day to about 250 mg/kg/day.

Methods for Synthesizing the Compounds

The following non-limiting schemes and examples demonstrates methods of synthesizing difluoro pentapeptide derivatives of the subject invention.

I. Synthesis of Adoc-D-t-Bug-Phe-Lys-C(0)-CF_2-C(0)-Leu-Arg-OMe

Adoc-D-t-Bug-OH - To a solution of 48.8 g (0.370 M) of D-t-butylglycine in 371 ml of 1N NaOH and 159 ml of dioxane at 0 C was added 33.9 g (0.40 M) of $NaHCO_3$ followed by the dropwise addition of a solution of 80.0 g (0.40 M) of

REPLACEMENT SHEET
26 FEB 1997

adamantyl fluoroformate in 371 ml of dioxane over 3 h. The mixture is stirred at this temperature for 0.5 h and the cooling bath removed. The solution is stirred for 3 h and the pH adjusted to 10 with 1N NaOH. This solution is extracted with EtOAc and the remaining aqueous phase acidified to pH = 3. This acidic solution is extracted with EtOAc, dried and the solvent removed to afford 118 g of product that is homogeneous by TLC (94/5/1 - CH_2Cl_2/ipa/AcOH), Rf = 0.90.

Adoc-D-t-Bug-Phe-OBn - To a solution of 115 g (0.37 M) of Adoc-D-t-Bug-OH and L-Phe-OBn.pTSA (0.36 M) in 1L CH_2Cl_2 is added 74.6 g (0.74 M) of TEA followed by the addition of 60.8 g (0.37 M) of DECP and the resulting solution is stirred overnight. The solvent is removed and the product redissolved in EtOAc and washed successively with brine, 0.4N HCl, and saturated $NaHCO_3$. The solution is dried and the solvent removed. The residue is chromatographed on silica to afford 186 g of product. TLC (85/15 - hexane/ETOAc) - Rf = 0.40.

Adoc-D-tBug-Phe-OH (1) - To a dry flask under N_2 is added 34.0 g 5% Pd on carbon. The flask is cooled to -78oC and 200 ml of dry degassed MeOH is added. To this slurry is added 8.8 ml of formic acid (88%) and this is followed by the addition of a mixture of 36.0 g of Adoc-D-t-Bug-Phe-OBn (0.066 M) in 100 ml of MeOH and 4.4 ml of formic acid. The cooling bath is removed and the slurry stirred for 2 h. At this time an additional 200 ml of 4.4% formic acid in MeOH is added and the slurry stirred overnight. The mixture is filtered through glass fiber filter paper under N_2. The filtrate is refiltered through a bed of Celite to remove all traces of catalyst. Removal of the solvent yields 17.3 g of product.

α-Boc, Σ-Z-Lys-C(OH)-CF_2-C(O)-OEt (3) - To a solution of 79.2 g (1.21 M) of activated zinc in 0.600 L THF is added 2.36 g of ethyl bromodifluoroacetate and the solution brought to reflux. As soon as reflux begins a mixture of 236 g (1.16 M) of ethyl bromodifluoroacetate and 176 g (0.485 M) of (2) in 1.80 L of THF is added dropwise. For the synthesis of (2), see Burkhart, J., N. Peet and P. Bey, "A Novel Method for the Preparation of Peptidyl a-Keto Esters," Tetrahedron Letters, Vol. 31, No. 10, pp. 1385 -1388 (1990). The solution is refluxed for 1 h, cooled and quenched with saturated $KHSO_4$. The quenched product is diluted with EtOAc and washed successively with saturated $KHSO_4$, saturated $NaHCO_3$ and brine. This sequence is followed by drying with $MgSO_4$ and removal of solvent. The residue is chromatographed on silica to afford 68.0 g of pure product. Rf = 0.20 75/25 hexane/EtOAc.

REPLACEMENT SHEET

26 FEB 1997

α-Boc, Σ-Z-Lys-C(OH)-CF_2-C(O)-ONa (4) - To a solution of 1.00 g (0.002 M) of (3) in 10 ml of MeOH is added 2 ml of 1N NaOH and the solution stirred at room temperature for 5 h and the solvent is removed. The residue is dissolved in a small amount of water and the milky solution extracted with ether. The water is removed from the aqueous phase and the residue thoroughly dried by placing on the vacuum pump for 24 h. Recovered 0.069 g of product is recovered.

α-Boc, Σ-Z-Lys-C(OH)-CF_2-C(O)-Leu-Arg(NO_2)-OMe (5) - To a solution of 0.500 g (1.03 mmol) of (4) and 0.470 g (1.26 mmol) of Leu-Arg(NO_2)-OMe.HCl in 30 ml of DMF is added 0.019 g (1.26 mmol) of HOBt in one portion. This addition is followed by the dropwise addition of 0.260 g (1.26 mmol) of DCC in 3 ml DMF. Finally, 0.003 ml of DIPEA is added and the solution stirred for 48 h. The DMF is removed in vacuo and the residue dissolved in EtOAc. The solution is transferred to a separatory funnel and washed with 0.1 N HCl, saturated $NaHCO_3$, and brine. The ethyl acetate layer is dried with $MgSO_4$, filtered and the solvent removed. The residue is chromatographed on silica to afford 0.420 g of product. Rf = 0.68 - 90/10 CH_2Cl_2/MeOH.

Σ-Z-Lys-C(OH)-CF_2-C(O)-Leu-Arg(NO_2)-OMe.HCl - To a solution of (5) (17.3 g, 0.022 M) in 200 ml of dioxane is added 66.8 ml (0.220 m) of 3.3 M HCl in dioxane and the mixture stirred for 1h. At this time an additional 5 ml of 3.3 M HCl in dioxane is added and the solution stirred an additional 0.25 h. The solvent is removed and the residue placed on the vacuum pump. The solid residue is triturated with ether under N2 and filtered. Recovered 16.0 g of white solid.

Adoc-D-t-Bug-Phe-Lys(Z)-C(OH)-CF_2-C(O)-Leu-Arg(NO_2)-OMe (6) - To a solution of 16.0 g of the residue from the previous reaction (0.022 M) and 11.1 g of (1) (0.024 M) in 300 ml of CH_2Cl_2 is added 6.69 ml (0.048 M) of TEA dropwise followed by the dropwise addition of 3.64 ml (0.024 M) of DECP. The solution is stirred overnight and the solvent removed. The residue is dissolved in EtOAc and washed successively with 0.1 N HCl, saturated $NaHCO_3$, and brine. The solution is dried and the solvent removed to afford 26.6 g that is chromatographed on silica to afford 12.7 g of pure product. Rf = 0.85, 90/10 - CH_2Cl_2/MeOH.


Adoc-D-t-Bug-Phe-Lys(Z)-C(O)CF_2C(O)-Leu-Arg(NO_2)-OMe - To a slurry of 14.3 g (0.034 M) of Dess-Martin periodinane in 1.15 L of CH_2Cl_2 is added 0.910 ml of dry pyridine, followed by the dropwise addition of 12.7 g

REPLACEMENT SHEET
26 FEB 1997


(0.011 M) of (6) in 2.3 L of CH_2Cl_2. For the preparation of Dess-Martin periodidinane, see Dess, D., and J. Martin, Journal of Organic Chemistry, Vol. 48 pp. 4155 -4156 (1983). The slurry is stirred for 2 h and 58.7 g (0.24 M) of $Na_2S_2O_4$ in saturated $NaHCO_3$ added and the resulting solution stirred for 0.5 h. The phases are separated and the aqueous phase extracted with ethyl acetate. The combined organic phases are dried and the solvent removed to give 13.9 g of product that is used in the next reaction without purification. (Rf = 0.30 - 95/5 CH_2Cl_2/MeOH).

Adoc-D-t-Bug-Phe-Lys-C(O)CF_2C(O)-Leu-Arg-OMe (7) - To a solution of 13.4 g (0.012 M) of the residue from the previous reaction in 200 ml of dry DMF is added 6 g of palladium black and the resulting slurry degassed for 0.5 h. To this slurry is added 8 ml of 3.3 M HCl in dioxane. The reaction is placed on the Paar shaker and hydrogenated for 6 h. The slurry is filtered through Celite and the solvent removed. The residue is chromatographed on reverse phase HPLC (CH_3CN/H_2O) to afford 6.5 g product.

REPLACEMENT SHEET
26 FEB 1997



REPLACEMENT SHEET
26 FEB 1997



REPLACEMENT SHEET
26 FEB 1997

II. Synthesis of Adoc-D-t-Bug-Phe-Arg-C(O)-CF_2-C(O)-Ala-Arg-OMe

Adoc-D-t-Bug-OH - To a solution of 48.8 g (0.370 M) of D-t-butylglycine in 371 ml of 1N NaOH and 159 ml of dioxane at 0oC is added 33.9 g (0.40 M) of adamantyl fluoroformate in 371 ml of dioxane over 3 hours. The mixture is stirred at this temperature for 0.5 hours and the cooling bath is removed. The solution is stirred for 3 hours and the pH adjusted to 10 with 1N NaOH. This solution is extracted with EtOAc and the remaining aqueous phase acidified to pH = 3. This acidic solution is extracted with EtOAc, dried and the solvent is removed to afford 118 g of product that is homogeneous by TLC (94/5/1 - CH_2Cl_2/i-pa/AcOH), Rf = 0.90.

Adoc-D-t-Bug-Phe-OBn - To a solution of 115 g (0.37 M) of Adoc-D-t-But-OH and L-Phe-OBn.pTSA (0.36 M) in 1L CH_2Cl_2 is added 74.6 g (0.74 M) of TEA followed by the addition of 60.8 g (0.37 M) of DECP and the resulting solution is stirred overnight. The solvent is removed and the product is redissolved in EtOAc and washed successively with brine, 0.4N NCl, and saturated $NaHCO_3$. The solution is dried and the solvent is removed. The residue is chromatographed on silica to afford 186 g of product. TLC (85/15 - hexane/EtOAc) - Rf = 0.40.

Adoc-D-t-Bug-Phe-OH (1) - To a dry flask under N_2 is added 34.0 g 5% Pd on carbon. The flask is cooled to -78oC and 200 ml of dry degassed MeOH is added. To this slurry is added 8.8 ml of formic acid (88%) and this is followed by the addition of a mixture of 36.0 g of Adoc-D-t-Bug-Phe-OBn (0.066 M) in 100 ml of MeOH and 4.4 ml of formic acid. The cooling bath is removed and the slurry is stirred for 2 hours. At this time an additional 200 ml of 4.4% formic acid in MeOH is added and the slurry is stirred overnight. The mixture is filtered through glass fiber filter paper under N_2. The filtrate is refiltered through a bed of Celite to remove all traces of catalyst. Removal of the solvent yields 17.3 g of product.

α-Boc, Σ-Z-Lys-C(OH)-CF_2-C(O)-OEt (3) - To a solution of 79.2 g (1.21 M) of activated zinc in 0.600 L THF is added 2.36 g of ethyl bromodifluoroacetate and the solution is brought to reflux. As soon as reflux begins, a mixture of 236 g (1.16 M) of ethyl bromodifluoroacetate and 176 g (0.485 M) of 2 in 1.80 L of THF is added dropwise. The solution is refluxed for 1 hour, cooled and quenched with saturated $KHSO_4$. The quenched product is diluted with EtOAc and washed successively with saturated $KHSO_4$, saturated $NaHCO_3$ and brine. This sequence is followed by drying with $MgSO_4$ and

REPLACEMENT SHEET
26 FEB 1997

removal of solvent. The residue is chromatographed on silica to afford 68.0 g of pure product. Rf = 0.20 75/25 hexane/EtOAc.

α-Boc, Σ -Z-Lys-C(OH)-CF_2-C(O)-ONa (4) - To a solution of 1.00 g (0.002 M) of 3 in 10 ml of MeOH is added 2 ml of 1N NaOH. The solution is stirred at room temperature for 5 hours and the solvent is removed. The residue is dissolved in a small amount of water and the milky solution is extracted with ether. The water is removed from the aqueous phase and the residue thoroughly dried by placing on the vacuum pump for 24 hours. 0.069 g of product is recovered.

α-Boc, Σ -Z-Lys-C(OH)-CH_2-C(O)-Leu-Arg(NO_2)-OMe (5) - To a solution of 0.500 g (1.03 mmol) and 4 and 0.470 g (1.26 mmol) of Leu-Arg(NO_2)-OMe·HCl in 30 ml of DMB is added 0.019 g (1.26 mmol) of HOBt in one portion. This addition is followed by the dropwise addition of 0.260 g (1.26 mmol) of DCC in 3 ml DMF. Finally, 0.003 ml of DIPEA is added and the solution stirred for 48 hours. The DMF is removed in vacuo and the residue dissolved in EtOAc. The solution is transferred to a separatory funnel and washed with 0.1 N HCl, saturated $NaHCO_3$, and brine. The ethyl acetate layer is dried with $MgSO_4$, filtered and the solvent is removed. The residue is chromatographed on silica gel to afford 0.420 g of product. Rf = 0.68-90/10 CH_2Cl_2/MeOH.

Σ-Z-Lys-C(OH)-CH_2-C(O)-Leu-Arg(NO_2)-OMe.HCl - To a solution of 5 (17.3 g, 0.022 M) in 200 ml of dioxane is added 66.8 ml (0.220 m) of 3.3 M HCl in dioxane. The mixture is stirred for 1 hour. At this time an additional 5 ml of 3.3 M HCl in dioxane is added and the solution is stirred an additional 0.25 hour. The solvent is removed and the residue placed on the vacuum pump. The solid residue is triturated with ether under N_2 and filtered. 16.0 g of while solid is recovered.

Adoc-D-t-Bug-Phe-Lys(Z)-C(OH)-CH_2-C(O)-Leu-Arg(NO_2)-OMe - (6) To a solution of 16.0 g of the crude product from the previous reaction (0.022 M) and 11.1 g of 1 (0.024 M) in 300 ml of CH_2Cl_2 is added 6.69 ml (0.048 M) of TEA dropwise followed by the dropwise addition of 3.64 ml (0.024 M) of DECP. The solution is stirred overnight and the solvent is removed. The residue is dissolved in EtOAc and washed successively with 0.1 N HCl, saturated $NaHCO_3$, and brine. The solution is dried and the solvent is removed to afford 26.6 g that is chromatographed on silica to afford 12.7 g of pure product. Rf = 0.85, 90/10 - CH_2Cl_2/MeOH.

REPLACEMENT SHEET
26 FEB 1997

Adoc-D-t-Bug-Phe-Lys(Z)-C(O)CF_2C(O)-Leu-Arg(NO_2)-OMe - To a slurry of 14.3 g (0.034 M) of Dess-Martin periodinane in 1.15 L of CH_2Cl_2 is added 0.910 ml of dry pyridine, followed by the dropwise addition of 12.7 g (0.011 M) of the product from the previous reaction in 2.3 L of CH_2Cl_2. The slurry is stirred for 2 hours and 58.7 g (0.24 M) of $Na_2S_2O_4$ in saturated $NaHCO_3$ is added. The resulting solution is stirred for 0.5 hour. The phases are separated and the aqueous phase is extracted with ethyl acetate. The combined organic phases are dried and the solvent is removed to give 13.9 g of product that is used in the next reaction without purification (Rf = 0.30 - 95/5 CH_2Cl_2/MeOH).

Adoc-D-t-Bug-Phe-Lys-C(O)CF_2C(O)-Leu-Arg-OMe.TFA (7) - To a solution of 13.4 g (0.012 M) of the crude product from the previous reaction in 200 ml of dry DMF is added 6 g of palladium black. The resulting slurry is degassed for 0.5 hour. To this slurry is added 8 ml of 3.3 M Hcl in dioxane. The reaction is placed on the Paar shaker and hydrogenated for 6 hours. The slurry is filtered through Celite and the solvent is removed. The residue is chromatographed on reverse phase HPLC (CH_3CN/H_2O/TFA) to afford 6.5 g product.

α-Boc, γ-Bn, γ-Z-Orn-C(OH)-CF_2-C(O)OEt (9) - To a solution of 5.67 g (0.087 M) of activated zinc in dry THF is added 1.0 g of ethyl bromodifluoroacetate and the temperature is raised to reflux. At this time, 15.2 g (0.037 M) of -Bn, -Z- -Boc-Ornithinal 8 and 16.9 g (0.083 M) of ethyl bromodifluoroacetate in THF is added, and the slurry allowed to reflux for 0.5 hour. The solution is cooled and quenched with saturated $KHSO_4$. The solution is diluted with EtOAc and washed with saturated $KHSO_4$, saturated $NaHCO_3$ and brine. The solution is dried and the solvent removed. The residue is chromatographed on silica to give 6.58 g of product. Rf = 0.22 99/1 - CH_2Cl_2/ipa.

α-Boc, γ-Bn, γ-Z-Orn-C(OH)-CF_2-C(O)ONa - The ester 9 (0.965 g, 1.71 mmol) is dissolved in 15 ml MeOH and 1.71 ml of 1N NaOH is added. The solution is stirred for 3 hours and the solvent is removed. The residue is placed on the vacuum pump overnight and used in the next reaction as is (0.870).

α-Boc, γ-Bn, γ-Z-Orn-C(OH)-CF_2-C(O)-Ala-Arg(NO_2)-OMe 10 - To a solution of 0.870 g (1.56 mmol) of the crude product from the previous reaction and 0.579 g (1.90 mmol) of Ala-Arg(NO_2)-OMe·HCl in 80 ml of dry DMF is added 0.283 g (2.09 mmol) of HOBt, followed by the successive addition of 0.392 g (1.90 mmol) of DCC in 20 ml of DMF and 0.068 ml (0.39 mmol) of

REPLACEMENT SHEET
26 FEB 1997

DIPEA. The mixture is stirred for 18 hours at room temperature and the solvent is removed. The residue is dissolved in EtOAc and washed with 0.1N HCl, saturated $NaHCO_3$ and brine. The solution is dried and the solvent is removed. Chromatography on silica gel yields 0.520 g of product (Rf = 0.20 and 0.22).

α-Boc-Orn-C(OH)-CF_2-C(O)-Ala-Arg(NO_2)-OMe - The tripeptide 10 (0.338 g, 0.43 mmol) is dissolved in 10 ml of DMF and 0.15 g of palladium black is added. The slurry is degassed and 0.20 ml of 4.0 M HCl in dioxane is added. The mixture is hydrogenated on the Paar apparatus overnight and filtered through Celite. The residue is redissolved in MeOH and 0.20 g of $Pd(OH)_2$ is added. The slurry is hydrogenated on the Paar for 24 hours at 50 psi. The slurry is filtered through Celite and the solvent is removed. The residue is used in the next reaction without purification.

α-Boc-Arg(Z)$_2$-C(OH)-CF_2-C(O)-Ala-Arg(NO_2)-OMe (11) - The crude amine (0.220 g, 0.39 mmol) is dissolved in THF and 0.325 g (0.91 mmol) of N,N-Bis-Z-S-methyl-isothiourea is added. To this solution is added 0.150 g $Hg(OAc)_2$ (0.47 mmol) and the mixture is stirred overnight. The solution is diluted with EtOAc and washed successively with 0.1 HCl, saturated $NaHCO_3$ and brine. The organic phase is dried and the solvent is removed. The residue is chromatographed on silica to afford 0.170 g of product as a mixture of diastereomers. Rf = 0.60, 0.62 - 5% MeOH/CH_2Cl_2.

Adoc-D-t-Bug-Phe-Arg(Z)$_2$-C(OH)-CH_2-C(O)-Ala-Arg(NO_2)-OMe (12) - To a solution of 0.228 g (0.262 mmol) of 11 in 5 ml of dioxane is added 0.656 ml of HCl in dioxane. The solution is stirred for 1 hour and an additional 0.656 ml of HCl in dioxane is added. After stirring for an additional 1 hour, the solvent is removed and the residue is placed on the vacuum pump overnight. The hydrolysis product is dissolved in 5 ml of CH_2Cl_2 and combined with 0.120 g (0.262 mmol) of Adoc-D-t-Bug-Phe-OH. To this solution is added 0.073 ml (0.525 mmol) of TEA followed by the addition of 0.043 ml of DECP. The resulting mixture is stirred overnight. The solvent is removed and the residue chromatographed on silica to afford 0.086 g of product as a mixture of diastereomers. Rf = 0.65, 0.68 - 4% ipa/CH_2Cl_2.

Adoc-D-t-Bug-Phe-Arg(Z)$_2$-C(O)-CF_2-C(O)-Ala-Arg(NO_2)-OMe - To a slurry of 0.094 g (0.16 mmol) of Dess Martin periodinane in 5 ml CH_2Cl_2 is added 0.091 g (0.052 mmol) of 12 dissolved in 0.5 ml via syringe. The mixture is stirred for 0.5 hour and 0.385 g (1.55 mmol) of $Na_2S_2O_3$ in saturated $NaHCO_3$ is added. After 0.5 hour, the solution is extracted with EtOAc and the


REPLACEMENT SHEET
26 FEB 1997

organic phase is separated and dried with $MgSO_4$. Removal of the solvent yields 0.94 g crude product. Rf = 0.68, 0.71 - 4% ipa/CH_2Cl_2.

Adoc-D-t-Bug-Phe-Arg-C(O)-CF_2-C(O)-Ala-Arg-OMe·TFA (13) - The crude alcohol is dissolved in 2 ml DMF and 50 mg palladium black is added. The slurry is degassed and 0.040 ml of 4.0 M HCl in dioxane is added. The mixture is hydrogenated on the Paar apparatus for 20 hours at 50 psi. The slurry is filtered through Celite and the solvent is removed. The residue is chromatographed on a reverse phase column (CH_3CN/H_2O/TFA) to afford 0.044 g of pure product.

N,N'-bis-carbobenzyloxy-S-methyl-isothiourea (14) - To a solution of S-methyl-isothiourea sulfate dimer (5.00 g, 18.0 mmol) in cold (0oC) 1N NaOH (36.0 ml, 36.0 mmol) is added 11.3 ml (79.0 mmol) benzyl chloroformate and 79.0 ml (79.0 mmol) of 1N NaOH dropwise from separate addition funnels. The slurry is stirred for 0.5 hour at 0oC and 1.5 hour at room temperature. The solution is extracted with ethyl acetate, dried with magnesium sulfate and the filtrate evaporated. Chromatography on silica gel yields 2.62 g pure product. Rf = 0.40 (15% EtOAc/pet ether.


REPLACEMENT SHEET
26 FEB 1997



8

1) NaOH

2) Ala-Leu-OMe-HCl
DCC
HOBt




10

1)H_2

2)N,N'-bis-CBz-S-methyl-isothiourea


14



11

1) HCl

2) Adoc-D-tBug-Phe-OH
DECP
TEA




12

1) Dess-Martin

2) H_2Pd

REPLACEMENT SHEET
26 FEB 1997



13

REPLACEMENT SHEET
26 FEB 1997

III. Synthesis of N-Terminating Linking Elements

Adac-D-t-Bug-Phe-OBn - To a solution of 5 ml of a 15% solution of phosgene in toluene under argon is added 0.140 ml TEA (1.00 mmol). This is followed by the addition of 0.112 g (0.750 mmol) of 1-adamantanamine in 1 ml toluene dropwise via syringe. The resulting slurry is stirred for 0.3 hour and the excess phosgene is removed by purging with argon. The solid is filtered off by suction through a sintered glass funnel and the solvent is removed. The residue is redissolved in dichloromethane and cooled to 0oC under argon. To this solution is added 0.140 ml TEA (1.00 mmol) followed by 0.100 g (0.250 mmol) d-t-Bug-Phe-OBn·HCl in one portion. The resulting solution is removed. Chromatography on silica yields 0.350 g product.

Hydrogenation of this compound followed by subsequent coupling with hydrolysis product of 11 yields compounds that are successfully elaborated to an alternative N-terminating linking element.

Methods for Testing Activity of the Compounds

The following non-limiting procedures are methods for testing the anti-inflammatory and/or analgesic activity of the difluoro pentapeptide derivatives of the subject invention.

Several enzyme inhibition assays are known to be predictive of anti-inflammatory activity for compounds. Such enzyme assays are useful for measuring the activity of compounds of the subject invention. Such enzyme assays include the following: porcine pancreatic kallikrein (PPK) - see references A, E and F; human urinary kallikrein (HUK) - see references E and F; human plasma kallikrein (HPK) - see references B and E; human plasmin (HP) - see references B and C; and urokinase (UK) - see reference D. The indicated references, which are hereby incorporated herein by reference, are the following:

(A) Lottenberg, R., U. Christensen, C.M. Jackson & P.L. Coleman, "Assay of Coagulation Proteases Using Peptide Chromogenic and Fluorogenic Substrates," Methods in Enzymology, Vol. 80, Academic Press, New York, NY (1981), pp. 341-361; (B) Geiger, R. "Kallikrein," Methods of Enzymatic Analysis, Vol. V, 3rd Edition, Bergmeyer, ed., Academic Press, New Nork, NY (1984), pp. 129-143; (C) Morris, J.P., S. Blatt, J.R. Powell, D.K. Strickland & F.J. Castellino, "Role of Lysine Binding Regions in the Kinetic Properties of Human Plasmin," Biochemistry, Vol. 20, No. 17 (August 18, 1981) p. 4811; (D) Wohl, R.C., L. Summaria & K.C. Robbins, "Kinetics of Activation of Human Plasminogen by Different Activator Species at pH 7.4 and 37oC" The Journal Biological

REPLACEMENT SHEET
26 FEB 1997

Chemistry, Vol. 255, No. 5 (March 10, 1980), pp. 2005-2013; (E) Okunishi, H., J. Burton & J. Spragg, "Specificity of Substrate Analogue Inhibitors of Human Urinary Kallikrein," Hypertension, Vol. 7, No. 3, Suppl. 1 (May-June, 1985), pp. I-72-I75; (F) Amundsen, E., J. Putter, P. Friberger, M. Knos, M. Larsbraten & G. Claeson, "Methods for the Determination of Glandular Kallikrein by Means of a Chromogenic Tripeptide Substrate," In Kinins-II part A, Fuji, S., et al., eds., Plenum Press, New York, NY (1979) pp. 83-95.

Another useful assay of activity is based on a method for determination of slow-binding enzyme inhibition disclosed in Imperiali, B. & R.H. Abeles, "Inhibition of Serine Proteases by Peptidyl Fluoromethyl Ketones," Biochemistry, Vol. 25 (1986) pp. 3760-3767. The method is modified as described below in order to study the slow binding inhibition of human plasmin, pig pancreatic kallikrein and human plasma kallikrein.

Reactions: The reaction mixtures contain 78 mM tris-HCl buffer, pH 7.4, 78 mM NaCl, 0.2 mg/ml bovine serum albumin, 0.2 mM S-2251 (D-Val-Leu-Arg-p-nitroanilide), 0.5 U/ml plasmin (1 mM), and variable concentrations of the test compound to be studied, in a total volume of 1 ml. The stock solution of plasmin is 1 U/ml in 50% glycerol. The absorbance change due to release of p-nitroaniline on enzymatic cleavage of S-2251 is monitored using an HP-8450 spectrophotometer system, set to measure $A^{400-410} - A^{470-490}$. The temperature is 30oC.

Calculations: K_{obsd} and v_s (steady state inhibited rate) are determined by fitting the progressive curve (first 20 min) to the integrated rate equation (i) using Labtech NotebookR software. Estimated k_1 is calculated for each run from v_s and the uninhibited rate v (equation ii), with (S) = 0.2 mM and K_m = 0.77 mM.

(i) $A = v_s t - ((v_s - v_o)/k_{obsd})(1 - \exp(-k_{obsd} t)) + A_o$

(ii) $k_1 = (I)/((v/v_s - 1)(1 + (s)/k_m))$

A plot of k_{obsd} vs test compound concentration for the different runs is then fit to a line, $y = mx + b$ (see equation iii); $k_{on} = m(1 + (S)/k_m)$ and $k_{off} = b$ are then calculated. Finally k_1 is calculated from equation iv.

(iii) $k_{obsd} = (I)/((v/v_s - 1)(1 + (S)/k_m))$

(iv) $k_1 = k_{off}/k_{on}$

Several in vivo assays are known to be predictive of the anti-inflammatory activity of compounds. Such in vivo assays are useful for measuring the activity of compounds of the subject invention. Such in vivo

REPLACEMENT SHEET
26 FEB 1997

assays are disclosed in the following references which are hereby incorporated herein by reference:

Winter, C.A., E.A. Risley, G.V. Nuss, "Carrageenin-Induced Edema in Hind Paw of the Rat as an Assay for Antiinflammatory Drugs," Proc. Soc. Exp. Biol., N.Y., Vol. 111 (1962), pp. 544-547; Vander Wende, C. & S. Margolin, "Analgesic Tests Based Upon Experimentally Induced Acute Abdominal Pain in Rats," Federal Proceedings., Vol. 15 (1956), p. 494; Blackham, A., J.W. Burns, J.B. Farmer, H. Radziwonik, J. Westwick, "An X-Ray Analysis of Adjuvant Arthritis in the Rat. The Effect of Prednisolone and Indomethacin," Agents and Actions, Vol. 7/1 (1977), p. 145-151; Winter, C.A. & G.W. Nuss, "Treatment of Adjuvant Arthritis in Rats with Anti-Inflammatory Drugs" Arthritis and Rheumatism, Vol. 9, No. 3 (June, 1966), pp. 394-404; and Francis, M.D., K. Hovancik & R.W. Boyce, "NE-58095: A Diphosphonate Which Prevents Bone Erosion and Preserves Joint Architecture in Experimental Arthritis," Int. J. Tiss. Reac., Vol. XI, No. 5 (1989), pp. 239-252.

Compositions and Methods of Using the Compounds

The following non-limiting examples exemplify contemplated compositions and uses for the subject invention.

Example I

Tablets are made by conventional procedures, each having the following composition:

Component	Quantity (mg)
Noc-D-t-Bug-Phe-Lys-C(O)CF_2C(O)-Leu-Arg-OMe	400
Microcrystalline cellulose	200
Pregelatinized starch	200
Povidone K-30	40
Magnesium stearate	20

One tablet is administered orally four times daily to a patient to alleviate inflammation in joints due to arthritis.

Example II

A lotion is made by conventional procedures, the lotion having the following composition:

Component	Quantity (%)
eBroc-D-tBug-Phe-Arg-C(O)CF_2C(O)-Gly-Arg-OMe	2.5
Glycerin	4.0
Methyl paraben	0.2
Propyl paraben	0.1

REPLACEMENT SHEET
26 FEB 1997

Carbopol 934	0.15
NaOH	0.46
Cetyl stearyl palmitate	1.0
Stearic acid	0.5
Lanolin fatty acids	0.5
Cetyl alcohol	3.0
Zantham gum	0.3
Sodium stearoyl-2-lactolate	0.75
Isopropyl myristate	2.0
Water	q.s.

One gram of the lotion is administered topically to the skin in the area of a burn twice daily to reduce inflammation and pain.

Example III

A solution is made by conventional means, each 2 ml of solution having the following composition:

Component	Quantity (mg)
Adoc-D-Phe-Phe-Lys-C(O)CF_2C(O)-D-Phe-Arg-OH	80
Benzalkonium chloride	40
Sterile aqueous saline solution	q.s.

A 2 ml dose of the solution is injected intramuscularly to a patient with arthritis to reduce inflammation and pain.

Example IV

A solution is made by conventional means, the solution having the following composition:

Component	Quantity (%)
Adoc-D-t-Bug-Phe-Arg-C(O)CF_2C(O)-Ala-Arg-OH	5.0
Benzalkonium chloride	0.02
Sodium carboxymethyl cellulose	0.01
Aqueous saline solution	q.s.


A 0.2 ml dose of the solution is administered by inhalation to a patient as needed to alleviate upper respiratory distress due to asthma.

While particular embodiments of the subject invention have been described, it will be obvious to those skilled in the art that various changes and modifications of the subject invention can be made without departing from the spirit and the scope of the invention. It is intended to cover, in the appended claims, all such modifications that are within the scope of this invention.

REPLACEMENT SHEET
26 FEB 1997

Claims

1. A compound, and pharmaceutically-acceptable salts and hydrates thereof, having the structure:



wherein:

(a) -X is selected from the group consisting of cyclic alkyl having from 4 to 15 carbon atoms; branched alkyl having at least two branches having from 6 to 15 carbon atoms; and aryl having from 6 to 15 carbon atoms;

(b) n is an integer from 0 to 2;

(c) -V- is selected from -OC(O)-, -N(Q)C(O)-, -N(Q)C(S)-, -C(O)-, $-SO_2-$ and -P(O)(OH)-;

d) -Q is hydrogen; or straight or branched chain alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to 6 carbon atoms; or -Q and -X are covalently linked forming a cyclic moiety which includes the nitrogen to which -Q is bonded and from 5 to 20 carbon atoms;

(e) -Z- is -O- or -NH-; when V is -OC(O)-, -Z- is -NH-;

(f) $-R^1$ is selected from straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^1$ is in either D or L configuration;

(g) $-R^2$ is selected from straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to 6 carbon atoms; cyclic alkyl,

This Sheet Has Been Replaced

REPLACEMENT SHEET
26 FEB 1997

saturated or unsaturated with 1 or 2 double bonds, having from 3 to 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^2$ is in either D or L configuration;

(h) $-R^3$ is $-(CH_2)_m$-A-NH_2 or $-(CH_2)_m$-A-B-$C(NH_2)$=NH wherein m is an integer from 1 to 6, -A- is a covalent bond or p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-; and the carbon atom bonded to $-R^3$ is in L configuration;

(i) $-R^4$ is selected from straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^4$ is in either D or L configuration;

(j) $-R^5$ is $-(CH_2)_m$-A-NH_2 or $-(CH_2)_m$-A-B-$C(NH_2)$=NH wherein m is an integer from 1 to 6, -A- is a covalent bond or p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-; and the carbon atom bonded to $-R^5$ is in either D or L configuration; and

(k) -Y is hydrogen or methyl.

2. The compound of Claim 1 wherein:

(a) -V- is -OC(O)- or -NHC(O)-, and -Z- is -NH-;

(b) n is 0 or 1;

(c) m is an integer from 1 to 5;

(d) -X is saturated and unsubstituted bicyclo or tricyclo alkyl having from 8 to 12 carbon atoms, or unsubstituted aryl having from 8 to 12 carbon atoms; and

REPLACEMENT SHEET

26 FEB 1997

This Sheet Has Been Replaced

(e) $-R^1$, $-R^2$ and $-R^4$ are selected from straight chain or branched alkyl having from 1 to 5 carbon atoms, cyclic alkyl having from 3 to 6 carbon atoms, and benzyl.

3. The compound of Claim 1 or 2, wherein:

(a) $-R^3$ is $-(CH_2)_m$ -A- NH_2, wherein -A- is a covalent bond, and m is from 3 to 5;

(b) $-R^5$ is $(CH_2)_m$ -A-B-C (NH_2) = NH, wherein -A- is a covalent bond, -B- is -NH-, and m is from 2 to 4;

(c) -Y is methyl; and


(d) -V- is -OC(O)-.

4. The compound of Claim 1, 2 or 3 wherein:

(a) -X is tricyclo alkyl; and

(b) $-R^1$ and $-R^4$ are saturated and unsubstituted straight chain or branched alkyl, having from 1 to 4 carbon atoms, and $-R^2$ is benzyl.

5. A compound, and the pharmaceutically-acceptable salts and hydrates thereof, having the structure:



wherein:

(a) $-AA^1-$ is selected from t-butylglycyl, valyl, isoleucyl, leucyl, cyclohexylglycyl, cyclohexylalanyl, phenylalanyl, naphthylalanyl, tryptophyl and adamantylglycyl;

(b) $-AA^2-$ is selected from phenylalanyl, naphthylalanyl, cyclohexylalanyl, leucyl and isoleucyl;

This Sheet Has Been Replaced

REPLACEMENT SHEET
26 FEB 1997

(c) -AA^3- is selected from arginyl, homoarginyl, lysyl, p-guanidinophenylalanyl, p-amidinophenylalanyl and p-aminophenylalanyl;

(d) -AA^4- is selected from phenylalanyl, naphthylalanyl, cyclohexylalanyl, leucyl and isoleucyl;

(e) -AA^5- is selected from arginyl, homoarginyl, lysyl, p-guanidinophenylalanyl, p-amidinophenylalanyl and p-aminophenylalanyl;

(f) -X^1 is selected from adamantyloxycarbonyl, adamantoyl, fluorenyl-methoxycarbonyl, adamantaneacetyl, adamantylaminocarbonyl, noradamantoyl, morpholinyl, homoadamantyloxycarbonyl, fenchyloxycarbonyl, isomenthyloxy-carbonyl, isopinocamphanyloxycarbonyl, 3,5-dimethyladamantyloxycarbonyl, endo-bornyloxycarbonyl, naphthyloxycarbonyl, and menthyloxycarbonyl; and

(g) -Y is hydrogen or methyl.

6. The compound of Claim 5 wherein:

(a) -AA^5- is arginyl;

(b) -AA^3- is lysyl; and

(c) -X^1 is selected from adamantyloxycarbonyl, endo-bornyloxycarbonyl and isopinocamphanyloxycarbonyl.

7. The compound of Claim 5 or 6 wherein:

(a) -AA^1- is selected from t-butylglycyl, valyl, and isoleucyl;

(b) -AA^2 is phenylalanyl; and

(c) -AA^4- is leucyl.

This Sheet Has Been Replaced



8. The compound of Claim 5, 6 or 7 wherein:

(a) -AA^1- is t-butylglycyl;

(b) -X^1 is adamantyloxycarbonyl; and

REPLACEMENT SHEET

26 FEB 1997

(c) -Y is methyl.

9. A pharmaceutical composition comprising a compound of any of Claims 1-8 and a pharmaceutically-acceptable carrier.

10. Use of a compound of any of Claims 1-8 for manufacture of a medicament for treating inflammation or pain.

This Sheet Has Been Replaced

REPLACEMENT SHEET
26 FEB 1997

ABSTRACT

DIFLUORO-PENTAPEPTIDE DERIVATIVE ANTI-INFLAMMATORY AGENTS


The subject invention involves anti-inflammatory compounds having structure (I), wherein: (a) -X is cyclic alkyl, having from 4 to 15 carbon atoms; branched alkyl, having from 6 to 15 carbon atoms; or arly having from 6 to 15 carbon atoms; (b) n is an integer from 0 to 2; (c) -V- is -OC(O)-, -N(Q)C(O)-, -N(Q)C(S)-, -C(O)-, -SO2- or -P(O)(OH)-; (d) -Q is hydrogen; or straight or branched chain alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to 6 carbon atoms; or -Q and -X are covalently linked forming a cyclic moiety which includes the nitrogen to which -Q is bonded and from 5 to 20 carbon atoms; (e) Z is -O- or -NH-; when V is -OC(O)-, -Z- is -NH-; (f) -R1 is a hydrophobic moiety; (g) -R2 is a hydrophobic moiety; (h) -R3 is -(CH2)m-A-NH2 or -(CH2)m-A-B-C(NH2)=NH wherein m is an integer from 1 to about 6, -A- is a covalent bond or p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-; (i) -R4 is a hydrophobic moiety; (j) -R5 is -(CH2)m-A-NH2 or -(CH2)m-A-B-C(NH2)=NH wherein m is an integer from 1 to about 6, -A- is a covalent bond or p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-; and (k) -Y is hydrogen or methyl. The subject invention also involves pharmaceutical compositions comprising the above compounds, and methods for treating inflammation or pain using such compounds and compositions.

REPLACEMENT SHEET
26 FEB 1997

PROPOSED AMENDMENT

Claims

1. A compound, ~~and~~ or pharmaceutically-acceptable salts ~~and~~ or hydrates thereof, having the structure:



wherein:

(a) -X is selected from the group consisting of cyclic alkyl having from 4 to about 15 carbon atoms; branched alkyl having at least two branches having from 6 to about 15 carbon atoms; and aryl having from 6 to about 15 carbon atoms;

(b) n is an integer from 0 to 2;

(c) -V- is selected from the group consisting of -OC(O)-, -N(Q)C(O)-, -N(Q)C(S)-, -C(O)-, $-SO_2-$ and -P(O)(OH)-;

d) -Q is hydrogen; or straight or branched chain alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to 6 carbon atoms; or -Q and -X are covalently linked forming a cyclic moiety which includes the nitrogen to which -Q is bonded and from about 5 to about 20 carbon atoms;

(e) -Z- is -O- or -NH-; when V is -OC(O)-, -Z- is -NH-;

(f) $-R^1$ is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^1$ is in either D or L configuration;

(g) $-R^2$ is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl,

saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^2$ is in either D or L configuration;

(h) $-R^3$ is $-(CH_2)_m$-A-NH_2 or $-(CH_2)_m$-A-B-$C(NH_2)$=NH wherein m is an integer from 1 to about 6, -A- is a covalent bond or p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-; and the carbon atom bonded to $-R^3$ is in L configuration;

(i) $-R^4$ is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to $-R^4$ is in either D or L configuration;

(j) $-R^5$ is $-(CH_2)_m$-A-NH_2 or $-(CH_2)_m$-A-B-$C(NH_2)$=NH wherein m is an integer from 1 to about 6, -A- is a covalent bond or p-phenyl or p-cyclohexyl, and -B- is a covalent bond or -NH-; and the carbon atom bonded to $-R^5$ is in either D or L configuration; and

(k) -Y is hydrogen or methyl.

2. The compound of Claim 1 wherein

~~(a)~~ -V- is -OC(O)- or -NHC(O)-, and -Z- is -NH-;

3. The compound of claim 2 wherein:

a(b) n is 0 or 1;

b(c) m is an integer from 1 to 5; and

(d) -X is saturated and unsubstituted bicyclo or tricyclo alkyl ~~having from 8~~ to 12 carbon atoms, or ~~unsubstituted aryl~~ having from 8 to 12 carbon atoms; and

(c) $-R^1$, $-R^2$ and $-R^4$ are selected from the group consisting of straight chain or branched alkyl having from 1 to about 5 carbon atoms, cyclic alkyl having from 3 to about 6 carbon atoms, and benzyl.

{insert new claim 4}

5. The compound of Claim ~~1 or 2~~ 4 wherein:

(a) $-R^3$ is $-(CH_2)_m$ -A- NH_2, wherein -A- is a covalent bond, and m is from 3 to 5;

(b) $-R^5$ is $(CH_2)_m$ -A-B-C (NH_2) = NH, wherein -A- is a covalent bond, -B- is -NH-, and m is from 2 to about 4;

~~(c) -Y is methyl; and~~

~~(d) -V- is -OC(O)-.~~


~~4. The compound of Claim 1, 2 or 3 wherein:~~

~~(a) -X is tricyclo-alkyl; and~~

~~(b) $-R^1$ and $-R^4$ are saturated and unsubstituted straight chain or branched alkyl, having from 1 to 4 carbon atoms, and $-R^2$ is benzyl.~~

{insert new claims 6 and 7}

6. A compound, ~~and~~ or the pharmaceutically-acceptable salts ~~and~~ or hydrates thereof, having the structure:



wherein:

(a) $-AA^1-$ is selected from the group consisting of t-butylglycyl, valyl, isoleucyl, leucyl, cyclohexylglycyl, cyclohexylalanyl, phenylalanyl, naphthylalanyl, tryptophyl and adamantylglycyl;

(b) $-AA^2-$ is selected from the group consisting of phenylalanyl, naphthylalanyl, cyclohexylalanyl, leucyl and isoleucyl;

(c) -AA^3- is selected from the group consisting of arginyl, homoarginyl, lysyl, p-guanidinophenylalanyl, p-amidinophenylalanyl and p-aminophenylalanyl;

(d) -AA^4- is selected from the group consisting of phenylalanyl, naphthylalanyl, cyclohexylalanyl, leucyl and isoleucyl;

(e) -AA^5- is selected from the group consisting of arginyl, homoarginyl, lysyl, p-guanidinophenylalanyl, p-amidinophenylalanyl and p-aminophenylalanyl;

(f) -X^1 is selected from the group consisting of adamantyloxycarbonyl, adamantoyl, fluorenyl-methoxycarbonyl, adamantaneacetyl, adamantylaminocarbonyl, noradamantoyl, morpholinyl, homoadamantyloxycarbonyl, fenchyloxycarbonyl, isomenthyloxy-carbonyl, isopinocamphanyloxycarbonyl, 3,5-dimethyladamantyloxycarbonyl, endo-bornyloxycarbonyl, naphthyloxycarbonyl, and menthyloxycarbonyl; and

(g) -Y is hydrogen or methyl.

{insert new claims 9 to 11}

~~6.~~ The compound of Claim 5 wherein:

(a) -AA^5- is arginyl;

(b) -AA^3- ~~is~~ lysyl; and

(c) -X^1 is selected from adamantyloxycarbonyl, endo-bornyloxycarbonyl and isopinocamphanyloxycarbonyl.

12 7. The compound of Claim 10 ~~5 or 6~~ wherein

(a) -AA^1- is selected from t-butylglycyl, valyl, and isoleucyl;

(b) -AA^2 is phenylalanyl; and

(c) -AA^4- is leucyl.

{insert new claim 13}

14 8. The compound of Claim 13 ~~5, 6 or 7~~ wherein

~~(a)~~ -AA^1- is t-butylglycyl;

~~(b)~~ -X^1 is adamantyloxycarbonyl; and

~~(c)~~ -Y is methyl.

{insert new claim 15}


16 9. A pharmaceutical composition comprising a compound of any of Claims ~~1-8~~ 1, 5, 7, 8, 10, 14 or 15; and (a) pharmaceutically-acceptable carrier.

17 10. Use of a compound of any of Claims ~~1-8~~ 1, 5, 8, 10 or 14 for manufacture of a medicament for treating inflammation or pain.

REPLACEMENT SHEETS SG
☐ Description : paged ..... to paged .....
☑ Claim(s) : paged 29 to paged 33.
☐ Drawing(s) : paged ..... to paged .....
☐ Abstract : paged ..... to paged .....
DATE OF RECEIPT 23/8/99

What is claimed is:

1. A compound, or pharmaceutically-acceptable salts or hydrates thereof, having the structure:



wherein:

(a) —X is selected from the group consisting of cyclic alkyl having from 4 to about 15 carbon atoms; branched alkyl having at least two branches having from 6 to about 15 carbon atoms; and aryl having from 6 to about 15 carbon atoms;

(b) n is an integer from 0 to 2;

(c) —V— is selected from the group consisting of —OC(O)—, —N(Q)C(O)—, —N(Q)C(S)—, —C(O)—, —SO_2— and —P(O)(OH)—;

(d) —Q is hydrogen; or straight or branched chain alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to 6 carbon atoms; or —Q and —X are covalently linked forming a cyclic moiety which includes the nitrogen to which —Q is bonded and from about 5 to about 20 carbon atoms;

(e) Z is —O— or —NH—; when V is —OC(O)—, —Z— is —NH—;

(f) —R^1 is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to —R^1 is in either D or L configuration;

(g) —R^2 is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from

REPLACEMENT SHEET
28 AUG 1999

1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to —R^2 is in either D or L configuration;

(h) —R^3 is —$(CH_2)_m$ —A—NH_2 or —$(CH_2)_m$ —A—B—$C(NH_2)$=NH wherein m is an integer from 1 to about 6, —A— is a covalent bond or p-phenyl or p-cyclohexyl, and —B— is a covalent bond or —NH— ; and the carbon atom bonded to —R^3 is in L configuration;

(i) —R^4 is selected from the group consisting of straight or branched alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 1 to about 6 carbon atoms; cyclic alkyl, saturated or unsaturated with 1 or 2 double bonds, having from 3 to about 10 carbon atoms; and arylalkyl wherein the aliphatic portion is saturated and has 1 or 2 carbon atoms; and the carbon atom bonded to —R^4 is in either D or L configuration;

(j) —R^5 is —$(CH_2)_m$ —A—NH_2 or —$(CH_2)_m$ —A—B—$C(NH_2)$=NH wherein m is an integer from 1 to about 6, —A— is a covalent bond or p-phenyl or p-cyclohexyl, and —B— is a covalent bond or —NH— ; and the carbon atom bonded to —R^5 is in either D or L configuration; and

(k) —Y is hydrogen or methyl.

2. The compound of claim 1 wherein

—V— is —OC(O)— or —NHC(O)—, and —Z— is —NH—.

3. The compound of claim 2, wherein:

(a) n is 0 or 1;

(b) m is an integer from 1 to 5; and

(c) —R^1, —R^2 and —R^4 are selected from the group consisting of straight chain or branched alkyl having from 1 to about 5 carbon atoms, cyclic alkyl having from 3 to about 6 carbon atoms, and benzyl.

REPLACEMENT SHEET
28 AUG 1999

4. The compound of claim 3 wherein —X is saturated and unsubstituted bicyclo or tricyclo alkyl having from 8 to about 12 carbon atoms, or unsubstituted aryl having from about 8 to about 12 carbon atoms.

5. The compound of claim 4 wherein:
(a) $-R^3$ is $-(CH_2)_m-A-NH_2$, wherein —A— is a covalent bond, and m is from 3 to 5; and
(b) $-R^5$ is $-(CH_2)_m-A-B-C(NH_2)=NH$, wherein —A— is a covalent bond, —B— is —NH—, and m is from 2 to about 4.

6. The compound of claim 5 wherein —Y is methyl, —V— is —OC(O)—, and X is tricycloalkyl.

7. The compound of claim 6 wherein $-R^1$ and $-R^4$ are saturated and unsubstituted straight chain or branched alkyl, having from 1 to about 4 carbon atoms, and $-R^2$ is benzyl.

8. A compound, or the pharmaceutically-acceptable salts or hydrates thereof, having the structure:

$$X'-AA^1-AA^2-AA^3-CF_2-\overset{\overset{O}{\|}}{C}-AA^4-AA^5-OY$$

wherein:

(a) $-AA^1-$ is selected from the group consisting of t-butylglycyl, valyl, isoleucyl, leucyl, cyclohexylglycyl, cyclohexylalanyl, phenylalanyl, naphthylalanyl, tryptophyl and adamantylglycyl;

(b) $-AA^2-$ is selected from the group consisting of phenylalanyl, naphthylalanyl, cyclohexylalanyl, leucyl and isoleucyl;

(c) $-AA^3-$ is selected from the group consisting of arginyl, homoarginyl, lysyl, p-guanidinophenylalanyl, p-amidinophenylalanyl and p-aminophenylalanyl;

REPLACEMENT SHEET
28 AUG 1999

* C L M S *

(d) —AA^4 — is selected from the group consisting of phenylalanyl, naphthylalanyl, cyclohexylalanyl, leucyl and isoleucyl;

(e) —AA^5— is selected from the group consisting of arginyl, homoarginyl, lysyl, p-guanidinophenylalanyl, p-amidinophenylalanyl and p-aminophenylalanyl;

(f) —X' is selected from the group consisting of adamantyloxycarbonyl, adamantoyl, fluorenylmethoxycarbonyl, adamantaneacetyl, adamantylaminocarbonyl, noradamantoyl, morpholinyl, homoadamantyloxycarbonyl, fenchyloxycarbonyl, isomenthyloxy-carbonyl, isopinocamphanyloxycarbonyl, 3,5-dimethyladamantyloxycarbonyl, endo-bornyloxycarbonyl, naphthyloxycarbonyl and menthyloxycarbonyl; and

(g) —Y is hydrogen or methyl.

9. The compound of claim 8 wherein —AA^5 — is arginyl.

10. The compound of claim 9 wherein —AA^3 — is lysyl.

11. The compound of claim 10 wherein —X' is adamantyloxycarbonyl.

12. The compound of claim 10 wherein
(a) —AA^1 — is selected from t-butylglycyl, valyl and isoleucyl;
(b) —AA^2 — is phenylalanyl; and
(c) —AA^4 — is leucyl.

13. The compound of claim 12 wherein X' is selected from adamantyloxycarbonyl, endo-bornyloxycarbonyl and isopinocamphanyloxycarbonyl.

14. The compound of claim 13 wherein —AA^1— is t-butylglycyl, —X' is adamantyloxycarbonyl and —Y is methyl.

REPLACEMENT SHEET
28 AUG 1999

15. The compound of claim 10 wherein —Y is methyl.

16. A pharmaceutical composition comprising:
(1) a compound of any of claims 1, 5, 7, 8, 10, 14 or 15; and
(2) a pharmaceutically-acceptable carrier.

17. Use of a compound of any of claims 1, 5, 8, 10 or 14 for manufacture of a medicament for treating inflammation or pain.

REPLACEMENT SHEET
28 AUG 1999