

SUBSTITUTED IMIDAZOLES AS N-TYPE CALCIUM CHANNEL BLOCKERSCROSS-REFERENCE TO RELATED APPLICATIONS

5 Not applicable.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR
DEVELOPMENT

The research and development of the invention described below was not federally
10 sponsored.

BACKGROUND OF THE INVENTION

Calcium ions play a fundamental role in the physiology and biochemistry of organisms and of cells. The entry of calcium into cells through ion channels mediates a variety of cellular and physiological responses, including gene expression, signal
15 transduction, neurotransmitter release, muscle contraction and hormone secretion. Ion channels are classified by gating, or what opens and closes the channel to the flux of ions. Voltage-gated ion channels open or close depending on the voltage gradient across the plasma membrane, whereas ligand-gated ion channels open or close depending on the binding of ligands to the channel. The classification of voltage-gated calcium channels
20 divides them into three groups: (i) high voltage-activated channels, which include L-, N-, P- and Q-type channels; (ii) intermediate voltage-activated R-type channels; and (iii) low voltage-activated T-type channels.

The N-type calcium channel is distributed mainly in central and peripheral neurons,
25 being localized primarily to presynaptic nerve terminals. This channel regulates the calcium flux required for depolarization-evoked release of neurotransmitters from synaptic endings. The transmission of pain signals from the periphery to the central nervous system (CNS) is mediated, inter alia, by N-type calcium channels located in the spinal cord. Inhibition of the N-type calcium channel in the superficial dorsal horn leads to a decrease

in membrane excitability and neurotransmitter release, resulting in pain relief. In addition, knock-out mice lacking the N-type calcium channel exhibit reduced nociceptive behaviors in animal models of pain.


5 N-type calcium channels have been shown to mediate the development and maintenance of the neuronal sensitization processes associated with neuropathic pain and therefore provide attractive targets for the development of analgesic drugs. Three N-type calcium channel modulators are currently approved for the treatment of pain: ω -conotoxin MVIIA (ziconotide), marketed as Prialt®, potently and selectively blocks the N-type
10 calcium channel and is indicated for the management of severe chronic pain; gabapentin, marketed as Neurontin®, and pregabalin, marketed as Lyrica®, bind with high affinity to the $\alpha 2\delta$ subunit of the N-type calcium channel and are indicated for the treatment of fibromyalgia, diabetic nerve pain and/or post-herpetic neuralgia pain.

15 It is an object of the present invention to provide N-Type calcium channel blockers. It is also an object of the invention to provide a method of treating, ameliorating or preventing pain by the administration of a compound of Formula (I). And, it is an object of the invention to provide a pharmaceutical composition comprising a compound of Formula (I), useful for treating, ameliorating or preventing pain.

20

SUMMARY OF THE INVENTION

The present invention is directed to a compound of Formula (I)



25

Formula (I)

wherein

R¹ is

- i) phenyl optionally independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, difluoromethoxy, and C₁₋₄alkoxy; provided that when phenyl of group (i) is substituted with a single substituent, that substituent is at the 4-position;
- 5 ii) a heteroaryl selected from the group consisting of pyridinyl, pyrimidinyl, pyridazinyl, and pyrazinyl; wherein said heteroaryl is optionally independently substituted with one or two substituents that are chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, or C₁₋₄alkoxy;
- iii) pyrimidin-5-ylmethyl;
- 10 iv) phenylmethyl, wherein the phenyl portion of phenylmethyl is optionally independently substituted with one or two substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, and C₁₋₄alkoxy; provided that when phenylmethyl of group (iv) is substituted with a single substituent, that substituent is at the 4-position;
- 15 v) phenylsulfonyl, wherein the phenyl portion of phenylsulfonyl is optionally independently substituted with one or two substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, and C₁₋₄alkoxy; provided that when phenylsulfonyl of group (v) is substituted with a single substituent, that substituent is at the 4-position;
- 20 vi) C₁₋₄alkylsulfonyl;
- vii) C₃₋₇cycloalkylsulfonyl; or
- viii) trifluoromethylsulfonyl;

R² is


- 25 i) phenyl optionally substituted with a substituent that is selected from the group consisting of C₁₋₄alkoxy and trifluoromethoxy;
- ii) a heteroaryl selected from the group consisting of pyridinyl, pyrimidinyl, thiazolyl, triazolyl, and pyrazinyl; wherein said heteroaryl is optionally substituted with a substituent that is C₁₋₄alkyl, C₁₋₄alkoxy, trifluoromethoxy, or hydroxy;

- iii) C₃₋₇cycloalkyl; or
- iv) C₃₋₇cycloalkyl-(C₁₋₂)alkyl;

R³ is selected from the group consisting of hydrogen, chloro, and methyl;

5

G is G1, G2, or G3,



and enantiomers, diastereomers, solvates and pharmaceutically acceptable salts thereof.

10 The present invention also provides, *inter alia*, a pharmaceutical composition comprising, consisting of and/or consisting essentially of a pharmaceutically acceptable carrier, a pharmaceutically acceptable excipient, and/or a pharmaceutically acceptable diluent, and a compound of Formula (I), or a pharmaceutically acceptable salt form thereof.

15 Also provided are processes for making a pharmaceutical composition comprising, consisting of, and/or consisting essentially of admixing a compound of Formula (I) and a pharmaceutically acceptable carrier, a pharmaceutically acceptable excipient, and/or a pharmaceutically acceptable diluent.

20 The present invention further provides, *inter alia*, methods for treating or ameliorating a N-Type calcium channel-modulated disorder in a subject, including a human or other mammal in which the disease, syndrome, or condition is affected by the modulation of the N-Type calcium channel, such as pain and the diseases that lead to such pain, using a compound of Formula (I).

25 The present invention also provides, *inter alia*, methods for producing the instant compounds and pharmaceutical compositions and medicaments thereof.

DETAILED DESCRIPTION OF THE INVENTION

With reference to substituents, the term “independently” refers to the situation where when more than one substituent is possible, the substituents may be the same or different from each other.

5 The term “alkyl” whether used alone or as part of a substituent group, refers to straight and branched carbon chains having 1 to 8 carbon atoms. Therefore, designated numbers of carbon atoms (e.g. C₁₋₈) refer independently to the number of carbon atoms in an alkyl moiety or to the alkyl portion of a larger alkyl-containing substituent. In substituent groups with multiple alkyl groups such as (C₁₋₆alkyl)₂amino-, the C₁₋₆alkyl
10 groups of the dialkylamino may be the same or different.

 The term “alkoxy” refers to an -O-alkyl group, wherein the term “alkyl” is as defined above.

 The terms “alkenyl” and “alkynyl” refer to straight and branched carbon chains having 2 or more carbon atoms, wherein an alkenyl chain contains at least one double bond
15 and an alkynyl chain contains at least one triple bond.

 The term “cycloalkyl” refers to saturated or partially saturated, monocyclic or polycyclic hydrocarbon rings of 3 to 14 carbon atoms. Examples of such rings include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and adamantyl.

 The term “benzo-fused cycloalkyl” refers to a 5- to 8- membered monocyclic
20 cycloalkyl ring fused to a benzene ring. The carbon atom ring members that form the cycloalkyl ring may be fully saturated or partially saturated.

 The term “heterocyclyl” refers to a nonaromatic monocyclic or bicyclic ring system having 3 to 10 ring members and which contains carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O, and S. Included
25 within the term heterocyclyl is a nonaromatic cyclic ring of 5 to 7 members in which 1 to 2 members are nitrogen, or a nonaromatic cyclic ring of 5 to 7 members in which 0, 1 or 2 members are nitrogen and up to 2 members are oxygen or sulfur and at least one member must be either nitrogen, oxygen or sulfur; wherein, optionally, the ring contains zero to one unsaturated bonds, and, optionally, when the ring is of 6 or 7 members, it contains up to 2

unsaturated bonds. The carbon atom ring members that form a heterocycle ring may be fully saturated or partially saturated. The term "heterocyclyl" also includes two 5 membered monocyclic heterocycloalkyl groups bridged to form a bicyclic ring. Such groups are not considered to be fully aromatic and are not referred to as heteroaryl groups.

5 When a heterocycle is bicyclic, both rings of the heterocycle are non-aromatic and at least one of the rings contains a heteroatom ring member. Examples of heterocycle groups include, and are not limited to, pyrrolinyl (including 2*H*-pyrrole, 2-pyrrolinyl or 3-pyrrolinyl), pyrrolidinyl, imidazoliny, imidazolidinyl, pyrazoliny, pyrazolidinyl, piperidinyl, morpholinyl, thiomorpholinyl, and piperazinyl. Unless otherwise noted, the
10 heterocycle is attached to its pendant group at any heteroatom or carbon atom that results in a stable structure.

 The term "benzo-fused heterocyclyl" refers to a 5 to 7 membered monocyclic heterocycle ring fused to a benzene ring. The heterocycle ring contains carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O, and S.
15 The carbon atom ring members that form the heterocycle ring may be fully saturated or partially saturated. Unless otherwise noted, benzo-fused heterocycle ring is attached to its pendant group at a carbon atom of the benzene ring.

 The term "aryl" refers to an unsaturated, aromatic monocyclic or bicyclic ring of 6 to 10 carbon members. Examples of aryl rings include phenyl and naphthalenyl.

20 The term "heteroaryl" refers to an aromatic monocyclic or bicyclic aromatic ring system having 5 to 10 ring members and which contains carbon atoms and from 1 to 4 heteroatoms independently selected from the group consisting of N, O, and S. Included within the term heteroaryl are aromatic rings of 5 or 6 members wherein the ring consists of carbon atoms and has at least one heteroatom member. Suitable heteroatoms include
25 nitrogen, oxygen, and sulfur. In the case of 5 membered rings, the heteroaryl ring preferably contains one member of nitrogen, oxygen or sulfur and, in addition, up to 3 additional nitrogens. In the case of 6 membered rings, the heteroaryl ring preferably contains from 1 to 3 nitrogen atoms. For the case wherein the 6 membered ring has 3 nitrogens, at most 2 nitrogen atoms are adjacent. Examples of heteroaryl groups include

furyl, thienyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, oxadiazolyl, triazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, indolyl, isoindolyl, benzofuryl, benzothienyl, indazolyl, benzimidazolyl, benzothiazolyl, benzoxazolyl, benzisoxazolyl, benzothiadiazolyl, benzotriazolyl, quinolinyl, isoquinolinyl
 5 and quinazolinyl. Unless otherwise noted, the heteroaryl is attached to its pendant group at any heteroatom or carbon atom that results in a stable structure.


The term “halogen” or “halo” refers to fluorine, chlorine, bromine and iodine.

The term “formyl” refers to the group —C(=O)H .

The term “oxo” refers to the group $(=\text{O})$.

10 Whenever the term “alkyl” or “aryl” or either of their prefix roots appear in a name of a substituent (*e.g.*, arylalkyl, alkylamino) the name is to be interpreted as including those limitations given above for “alkyl” and “aryl.” Designated numbers of carbon atoms (*e.g.*, $\text{C}_1\text{--C}_6$) refer independently to the number of carbon atoms in an alkyl moiety, an aryl moiety, or in the alkyl portion of a larger substituent in which alkyl appears as its prefix
 15 root. For alkyl and alkoxy substituents, the designated number of carbon atoms includes all of the independent members included within a given range specified. For example C_{1-6} alkyl would include methyl, ethyl, propyl, butyl, pentyl and hexyl individually as well as sub-combinations thereof (*e.g.*, C_{1-2} , C_{1-3} , C_{1-4} , C_{1-5} , C_{2-6} , C_{3-6} , C_{4-6} , C_{5-6} , C_{2-5} , etc.).

In general, under standard nomenclature rules used throughout this disclosure, the
 20 terminal portion of the designated side chain is described first followed by the adjacent functionality toward the point of attachment. Thus, for example, a “ $\text{C}_1\text{--C}_6$ alkylcarbonyl” substituent refers to a group of the formula:



25 The term “R” at a stereocenter designates that the stereocenter is purely of the *R*-configuration as defined in the art; likewise, the term “S” means that the stereocenter is purely of the *S*-configuration. As used herein, the terms “*R” or “*S” at a stereocenter are used to designate that the stereocenter is of pure but unknown configuration. As used

herein, the term “RS” refers to a stereocenter that exists as a mixture of the *R*- and *S*-configurations. Similarly, the terms “*RS” or “*SR” refer to a stereocenter that exists as a mixture of the *R*- and *S*-configurations and is of unknown configuration relative to another stereocenter within the molecule.

5 Compounds containing one stereocenter drawn without a stereo bond designation are a mixture of two enantiomers. Compounds containing two stereocenters both drawn without stereo bond designations are a mixture of 4 diastereomers. Compounds with 2 stereocenters both labeled “RS” and drawn with stereo bond designations are a 2-component mixture with relative stereochemistry as drawn. Compounds with 2
10 stereocenters both labeled “*RS” and drawn with stereo bond designations are a 2-component mixture with relative stereochemistry unknown. Unlabeled stereocenters drawn without stereo bond designations are a mixture of the *R*- and *S*-configurations. For unlabeled stereocenters drawn with stereo bond designations, the absolute stereochemistry is as depicted.

15 Unless otherwise noted, it is intended that the definition of any substituent or variable at a particular location in a molecule be independent of its definitions elsewhere in that molecule. It is understood that substituents and substitution patterns on the compounds of Formula (I) can be selected by one of ordinary skill in the art to provide compounds that are chemically stable and that can be readily synthesized by techniques
20 known in the art as well as those methods set forth herein.

The term “subject” refers to an animal, preferably a mammal, most preferably a human, who has been the object of treatment, observation or experiment.

 The term “therapeutically effective amount” refers to an amount of an active compound or pharmaceutical agent, including a compound of the present invention, which
25 elicits the biological or medicinal response in a tissue system, animal or human that is being sought by a researcher, veterinarian, medical doctor or other clinician, which includes alleviation or partial alleviation of the symptoms of the disease, syndrome, condition, or disorder being treated.

The term “composition” refers to a product that includes the specified ingredients

in therapeutically effective amounts, as well as any product that results, directly, or indirectly, from combinations of the specified ingredients in the specified amounts.

The term "N-Type calcium channel blocker" is intended to encompass a compound that interacts with the N-Type calcium channel to substantially reduce or eliminate its functional activity, thereby decreasing the flow of calcium ions through the channel and the rise of intracellular calcium concentrations.

The term "N-Type calcium channel-modulated" is used to refer to the condition of being affected by the modulation of the N-Type calcium channel, including the condition of being affected by the inhibition of the N-Type calcium channel, such as, for example, pain, the diseases that lead to such pain and treatments that lead to the reduction of such pain.

As used herein, unless otherwise noted, the term "affect" or "affected" (when referring to a disease, syndrome, condition or disorder that is affected by the inhibition of N-Type calcium channel) shall include a reduction in the frequency and / or severity of one or more symptoms or manifestations of said disease, syndrome, condition or disorder and / or include the prevention of the development of one or more symptoms or manifestations of said disease, syndrome, condition or disorder or the development of the disease, condition, syndrome or disorder.

The compounds of Formula (I) are useful in methods for treating, ameliorating and / or preventing a disease, a syndrome, a condition or a disorder that is affected by the inhibition of N-Type calcium channel. Such methods comprise, consist of and / or consist essentially of administering to a subject, including an animal, a mammal, and a human in need of such treatment, amelioration and / or prevention, a therapeutically effective amount of a compound of Formula (I), or an enantiomer, diastereomer, solvate or pharmaceutically acceptable salt form thereof. In particular, the compounds of Formula (I) are useful for treating, ameliorating and / or preventing pain as well as diseases, syndromes, conditions or disorders causing such pain. More particularly, the compounds of Formula (I) are useful for treating, ameliorating and / or preventing acute pain, inflammatory pain and / or

neuropathic pain, comprising administering to a subject in need thereof a therapeutically effective amount of a compound of Formula (I), as herein defined.


Acute pain, as used herein, refers to pain that comes on quickly, can be of varying severity but is self-limiting and of relatively short duration. Examples of acute pain
5 include, but are not limited to, post-operative pain, post-surgical pain, toothache, burn, sunburn, insect/animal bites and stings, headache and/or any pain associated with acute trauma or injury.

Inflammatory pain refers to pain arising from an inflammatory disease, condition, syndrome or disorder, including but not limited to inflammatory bowel disease, irritable
10 bowel syndrome, visceral pain, migraine, post-operative pain, osteoarthritis, rheumatoid arthritis, back pain, low back pain, joint pain, abdominal pain, chest pain, labor pain, musculoskeletal diseases, skin diseases, toothache, pyresis, burn, sunburn, snake bite, venomous snake bite, spider bite, insect sting, neurogenic or overactive bladder, interstitial cystitis, urinary tract infection, rhinitis, contact dermatitis/hypersensitivity, itch, eczema,
15 pharyngitis, mucositis, enteritis, irritable bowel syndrome, cholecystitis, pancreatitis, post-mastectomy pain syndrome, menstrual pain, endometriosis, pain due to physical trauma, headache, sinus headache, tension headache or arachnoiditis.

A further embodiment of the present invention is directed to a method for treating, ameliorating and / or preventing neuropathic pain. Neuropathic pain refers to a disease,
20 syndrome, condition and/or disorder involving damage to the peripheral or central nervous system, including cancer pain, neurological disorders, spine and peripheral nerve surgery, brain tumor, traumatic brain injury (TBI), chemotherapy-induced pain, pain chronification, radicular pain, HIV pain, spinal cord trauma, chronic pain syndrome, fibromyalgia, chronic fatigue syndrome, lupus, sarcoidosis, peripheral neuropathy, bilateral peripheral
25 neuropathy, diabetic neuropathy, central pain, neuropathies associated with spinal cord injury, stroke, amyotrophic lateral sclerosis (ALS), Parkinson's disease, multiple sclerosis, sciatic neuritis, mandibular joint neuralgia, peripheral neuritis, polyneuritis, stump pain, phantom limb pain, bony fractures, oral neuropathic pain, Charcot's pain, complex regional pain syndrome I and II (CRPS I/II), radiculopathy, Guillain-Barre syndrome, meralgia

paresthetica, burning-mouth syndrome, optic neuritis, postfebrile neuritis, migrating neuritis, segmental neuritis, Gombault's neuritis, neuronitis, cervicobrachial neuralgia, cranial neuralgia, geniculate neuralgia, glossopharyngeal neuralgia, migrainous neuralgia, idiopathic neuralgia, intercostals neuralgia, mammary neuralgia, Morton's neuralgia,
 5 nasociliary neuralgia, occipital neuralgia, post-herpetic neuralgia, causalgia, red neuralgia, Sluder's neuralgia, splenopalatine neuralgia, supraorbital neuralgia, trigeminal neuralgia, vulvodynia, or vidian neuralgia.

Embodiments of the present invention include a compound of Formula (I)



Formula (I)


wherein

a) R¹ is


- i) phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, difluoromethoxy, and C₁₋₄alkoxy; provided that when phenyl of group (i) is substituted with a single substituent, that substituent is at the 4-position;
- ii) a heteroaryl that is pyridinyl; wherein said pyridinyl is optionally independently substituted with one or two substituents that are chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, or C₁₋₄alkoxy;
- iii) phenylmethyl, wherein the phenyl portion of phenylmethyl is optionally independently substituted with one or two substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, and C₁₋₄alkoxy; provided that when phenylmethyl of group (iii) is substituted with a single substituent, that substituent is at the 4-position;

- 5 iv) phenylsulfonyl, wherein the phenyl portion of phenylsulfonyl is optionally independently substituted with one or two substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, and C₁₋₄alkoxy; provided that when phenylsulfonyl of group (iv) is substituted with a single substituent, that substituent is at the 4-position;
- v) C₁₋₄alkylsulfonyl;
 or
 vi) trifluoromethylsulfonyl;
- 10 b) R¹ is
- i) phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, methyl, difluoromethoxy, and C₁₋₂alkoxy; provided that when phenyl of group (i) is substituted with a single substituent, that
- 15 substituent is at the 4-position; or
- ii) a heteroaryl that is pyridinyl; wherein said pyridinyl is optionally independently substituted with one or two substituents that are chloro, fluoro, bromo, cyano, trifluoromethyl, or C₁₋₄alkoxy;
- 20 c) R¹ is phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, cyano, trifluoromethyl, difluoromethoxy, and methyl; provided that when phenyl is substituted with a single substituent, that substituent is at the 4-position;
- 25 d) R² is
- i) phenyl substituted with a substituent that is selected from the group consisting of C₁₋₄alkoxy and trifluoromethoxy;

- ii) a heteroaryl selected from the group consisting of pyridinyl, pyrimidinyl, and pyrazinyl; wherein said heteroaryl is optionally substituted with a substituent that is C₁₋₄alkoxy or trifluoromethoxy;
- iii) C₃₋₇cycloalkyl; or
- 5 iv) C₃₋₇cycloalkyl-(C₁₋₂)alkyl;
- e) R² is
- i) phenyl substituted with C₁₋₄alkoxy;
- ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl;
- 10 wherein said heteroaryl is optionally substituted with C₁₋₄alkoxy; or
- iii) C₃₋₇cycloalkyl-(C₁₋₂)alkyl;
- f) R² is
- i) phenyl substituted with C₁₋₄alkoxy; or
- 15 ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl; wherein said heteroaryl is optionally substituted with C₁₋₄alkoxy;
- g) R² is
- i) phenyl substituted with methoxy; or
- 20 ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl; wherein said heteroaryl is optionally substituted with C₁₋₂alkoxy;
- h) R³ is hydrogen;
- 25 i) G is G1 or G2;




j) G is G1;



G1 ;

k) G is G2;




G2

5

and any combination of embodiments a) through k) above, provided that it is understood that combinations in which different embodiments of the same substituent would be combined are excluded;

10 and enantiomers, diastereomers, solvates and pharmaceutically acceptable salts thereof.

An embodiment of the present invention is directed to a compound of Formula (I)



Formula (I)

15

wherein

R¹ is

i) phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, difluoromethoxy, and C₁₋₄alkoxy; provided that when phenyl of group (i) is substituted with a single substituent, that substituent is at the 4-position;

20

- 5 ii) a heteroaryl that is pyridinyl; wherein said pyridinyl is optionally independently substituted with one or two substituents that are chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, or C₁₋₄alkoxy;

 iii) phenylmethyl, wherein the phenyl portion of phenylmethyl is optionally independently substituted with one or two substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, and C₁₋₄alkoxy; provided that when phenylmethyl of group (iii) is substituted with a single substituent, that substituent is at the 4-position;

10 iv) phenylsulfonyl, wherein the phenyl portion of phenylsulfonyl is optionally independently substituted with one or two substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, and C₁₋₄alkoxy; provided that when phenylsulfonyl of group (iv) is substituted with a single substituent, that substituent is at the 4-position;

 v) C₁₋₄alkylsulfonyl; or

15 vi) trifluoromethylsulfonyl;

R² is

- i) phenyl substituted with a substituent that is selected from the group consisting of C₁₋₄alkoxy and trifluoromethoxy;

20 ii) a heteroaryl selected from the group consisting of pyridinyl, pyrimidinyl, and pyrazinyl; wherein said heteroaryl is optionally substituted with a substituent that is C₁₋₄alkoxy or trifluoromethoxy;


 iii) C₃₋₇cycloalkyl; or

 iv) C₃₋₇cycloalkyl-(C₁₋₂)alkyl;

25


R³ is hydrogen, chloro, or methyl;

G is G1 or G2;



and enantiomers, diastereomers, solvates, and pharmaceutically acceptable salt forms thereof.

5 An embodiment of the present invention is directed to a compound of Formula (I)



Formula (I)

wherein

10 R¹ is


- i) phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, methyl, difluoromethoxy, and C₁₋₂alkoxy; provided that when phenyl of group (i) is substituted with a single substituent, that
- 15 substituent is at the 4-position; or
- ii) a heteroaryl that is pyridinyl; wherein said pyridinyl is optionally independently substituted with one or two substituents that are chloro, fluoro, bromo, cyano, trifluoromethyl, or C₁₋₄alkoxy;

20 R² is

- i) phenyl substituted with C₁₋₄alkoxy;
- ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl; wherein said heteroaryl is optionally substituted with C₁₋₄alkoxy; or
- iii) C₃₋₇cycloalkyl-(C₁₋₂)alkyl;


R^3 is hydrogen;

G is G1 or G2;



and enantiomers, diastereomers, solvates, and pharmaceutically acceptable salt forms thereof.

An embodiment of the present invention is directed to a compound of Formula (I)



Formula (I)

wherein


R^1 is phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, cyano, trifluoromethyl, difluoromethoxy, and methyl; provided that when phenyl is substituted with a single
15 substituent, that substituent is at the 4-position;

R^2 is

- 20 i) phenyl substituted with C_{1-4} alkoxy; or
 ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl;
 wherein said heteroaryl is optionally substituted with C_{1-4} alkoxy;

R^3 is hydrogen;


G is G1 or G2;



and enantiomers, diastereomers, solvates, and pharmaceutically acceptable salt forms thereof.

5

An embodiment of the present invention is directed to a compound of Formula (I)



Formula (I)

wherein


10 R¹ is phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, cyano, trifluoromethyl, difluoromethoxy, and methyl; provided that when phenyl is substituted with a single substituent, that substituent is at the 4-position;

15 R² is

- i) phenyl substituted with methoxy; or
- ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl; wherein said heteroaryl is optionally substituted with C₁₋₂alkoxy;

20 R³ is hydrogen;


G is G1 or G2;



and enantiomers, diastereomers, solvates, and pharmaceutically acceptable salt forms thereof.

5


An embodiment of the present invention is directed to a compound of Formula (I)



Formula (I)

wherein


- 10 R^1 is phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, cyano, trifluoromethyl, difluoromethoxy, and methyl; provided that when phenyl is substituted with a single substituent, that substituent is at the 4-position;
- 15 R^2 is
- i) phenyl substituted with methoxy; or
 - ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl; wherein said heteroaryl is optionally substituted with C_{1-2} alkoxy;
- 20 R^3 is hydrogen;
- G is G1;



G1 ;

and enantiomers, diastereomers, solvates, and pharmaceutically acceptable salt forms thereof.

5 An embodiment of the present invention is directed to a compound of Formula (I)



Formula (I)

wherein

10 R¹ is phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, cyano, trifluoromethyl, difluoromethoxy, and methyl; provided that when phenyl is substituted with a single substituent, that substituent is at the 4-position;


R² is

- 15 i) phenyl substituted with methoxy; or
 ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl; wherein said heteroaryl is optionally substituted with C₁₋₂alkoxy;

R³ is hydrogen;

20

G is G2;




G2 ;

and enantiomers, diastereomers, solvates, and pharmaceutically acceptable salt forms thereof.

Further embodiments of the present invention are directed to a compound of

5 Formula (I)



Formula (I)

selected from the group consisting of

- Cpd 1**, 4-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- 10 **Cpd 2**, 4-[1-(4-Fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyridine;
- Cpd 3**, 2-Ethoxy-5-[2-(3-methoxypyridin-4-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]pyridine;
- 15 **Cpd 4**, 1-(4-Fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 5**, 5-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-methylpyridine;
- Cpd 6**, 2-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-5-methylpyridine;
- 20 **Cpd 7**, 2-(2-Methoxyphenyl)-1-(4-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 8**, 5-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-methylpyrimidine;
- 25 **Cpd 9**, 3-(1-(3-chlorobenzyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine;

- Cpd 10**, 3-[1-(4-Fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 11**, 3-[1-(4-Chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- 5 **Cpd 12**, 4-[2-(3-Methoxypyridin-2-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 13**, 4-[1-(4-Chloro-3-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyridine;
- Cpd 14**, 2-Chloro-5-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- 10 **Cpd 15**, 4-[2-(2-Methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-methylbenzonitrile;
- Cpd 16**, 2-methoxy-3-(4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1-(2,3,4-trifluorophenyl)-1H-imidazol-2-yl)pyridine;
- 15 **Cpd 17**, 2-[1-(3-Chloro-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyridine;
- Cpd 18**, 4-[1-(3-Chloro-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyridine;
- Cpd 19**, 2-[1-(4-Chloro-3-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyridine;
- 20 **Cpd 20**, 3-[1-(3-Chloro-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-4-methoxypyridine;
- Cpd 21**, 4-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-methylbenzonitrile;
- 25 **Cpd 22**, 5-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-methylbenzonitrile;
- Cpd 23**, 3-[1-(3-Fluoro-4-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;

- Cpd 24**, 3-[1-(4-Fluoro-3-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 25**, 5-[2-(2-Methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-methylbenzonitrile;
- 5 **Cpd 26**, 3-[1-(4-Chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-4-methyl-4H-1,2,4-triazole;
- Cpd 27**, 3-[1-(2-Chlorobenzyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 28**, 3-[1-(4-Chlorobenzyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- 10 **Cpd 29**, 4-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-(trifluoromethyl)benzonitrile;
- Cpd 30**, 3-[1-(2,4-Difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- 15 **Cpd 31**, 3-[1-(2,3-difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 32**, 3-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 33**, 4-[1-(3,4-Difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyridine;
- 20 **Cpd 34**, 2-Methoxy-4-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 35**, 2-[1-(3,4-Difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyridine;
- 25 **Cpd 36**, 3-Fluoro-4-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 37**, 2-Methoxy-5-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;

- Cpd 38**, 1-(4-Chlorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 39**, 5-Chloro-2-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]pyridine;
- 5 **Cpd 40**, 3-[1-(3,4-Difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 41**, 1-(4-Chloro-3-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 42**, 1-(3,4-Difluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- 10 **Cpd 43**, 4-[2-(2-Methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 44**, 3-[1-(4-Chloro-3-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- 15 **Cpd 45**, 1-(3-Chloro-4-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 46**, 3-[1-(3-Chloro-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 47**, 2-[1-(4-Fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyrazine;
- 20 **Cpd 48**, 3-[2-(2-Methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 49**, 5-[1-(4-Chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-4-methoxypyrimidine;
- 25 **Cpd 50**, 2-[1-(4-Chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyrazine;
- Cpd 51**, 3-[1-(4-Chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-ethoxypyridine;

- Cpd 52**, 3-[1-(3,4-Difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-ethoxypyridine;
- Cpd 53**, 5-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-(trifluoromethyl)pyridine;
- 5 **Cpd 54**, 2-Methoxy-3-{4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1-[6-(trifluoromethyl)pyridin-3-yl]-1H-imidazol-2-yl}pyridine;
- Cpd 55**, 1-(4-Chlorophenyl)-2-(cyclopropylmethyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 56**, 1-(4-Chlorophenyl)-2-cyclopropyl-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- 10 **Cpd 57**, 4-[2-Cyclopropyl-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 58**, 4-[2-(Cyclopropylmethyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- 15 **Cpd 59**, 5-[2-(Cyclopropylmethyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-(trifluoromethyl)pyridine;
- Cpd 60**, 2-[1-(4-Chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-ethoxypyrazine;
- Cpd 61**, 1-(4-(difluoromethoxy)phenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- 20 **Cpd 62**, 1-(4-Bromo-3-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 63**, 1-(3-Bromo-4-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- 25 **Cpd 64**, 3-[1-(3-Bromo-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 65**, 3-[1-(4-Bromo-3-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;

- Cpd 66**, 4-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine;
- Cpd 67**, 2-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine;
- 5 **Cpd 68**, 3-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-4-methoxypyridine;
- Cpd 69**, 3-(1-(4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-4-methoxypyridine;
- Cpd 70**, 2-(1-(4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine;
- 10 **Cpd 71**, 3-(1-(2,5-difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine;
- Cpd 72**, 3-(1-(3,5-difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine;
- 15 **Cpd 73**, 3-fluoro-5-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile;
- Cpd 74**, 2-Fluoro-5-[2-(2-methoxy-phenyl)-4-(2,2,6,6-tetramethyl-tetrahydro-pyran-4-yl)-imidazol-1-yl]-benzonitrile;
- Cpd 75**, 3-[1-(6-Ethoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- 20 **Cpd 76**, 3-Fluoro-4-[2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 77**, 2-Methoxy-4-[2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- 25 **Cpd 78**, 2-Chloro-4-[2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 79**, 2-Fluoro-5-[2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;

- Cpd 80**, 2-Fluoro-4-[2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 81**, 2-Fluoro-4-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- 5 **Cpd 82**, 2,6-difluoro-4-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile;
- Cpd 83**, 3,5-difluoro-4-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile;
- Cpd 84**, 2-Chloro-5-[2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- 10 **Cpd 85**, 2-Methoxy-5-[2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 86**, 4-[2-(4-Methoxypyrimidin-5-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- 15 **Cpd 87**, 4-[2-(3-Methoxypyrazin-2-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 88**, 2-fluoro-4-(2-(3-methoxypyrazin-2-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile;
- Cpd 89**, 4-(2-(2-ethoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-fluorobenzonitrile;
- 20 **Cpd 90**, 1-(4-chlorophenyl)-2-(2-methoxyphenyl)-5-methyl-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 91**, 4-[5-Chloro-2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- 25 **Cpd 92**, 3-[5-Chloro-1-(4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 93**, 3-[5-Chloro-1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;

- Cpd 94**, 2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1-[(trifluoromethyl)sulfonyl]-1H-imidazole;
- Cpd 95**, 1-(Cyclopropylsulfonyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- 5 **Cpd 96**, 2-(2-Methoxyphenyl)-1-[(2-methylpropyl)sulfonyl]-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 97**, 1-[(4-Chlorophenyl)sulfonyl]-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 98**, 1-[(4-Fluorophenyl)sulfonyl]-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- 10 **Cpd 99**, 4-{[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]methyl} benzonitrile;
- Cpd 100**, 5-{[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]methyl} pyrimidine;
- 15 **Cpd 101**, 3-{[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]methyl} benzonitrile;
- Cpd 102**, 4-[1-(4-Chlorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 103**, 4-[1-(4-Chlorophenyl)-2-(4-methoxypyridin-3-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- 20 **Cpd 104**, 4-[1-(4-Chlorophenyl)-2-(3-methoxypyridin-2-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 105**, 4-[1-(4-Chlorophenyl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- 25 **Cpd 106**, 4-[1-(3-Bromo-4-fluorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 107**, 4-[1-(3,4-Difluorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;

- Cpd 108**, 4-[1-(3,4-Difluorophenyl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 109**, 4-{2-(2-Methoxyphenyl)-1-[6-(trifluoromethyl)pyridin-3-yl]-1H-imidazol-4-yl}-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- 5 **Cpd 110**, 4-[1-(3-Chloro-4-fluorophenyl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 111**, 4-[1-(3-Chloro-4-fluorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 112**, 4-[1-(4-Chloro-3-fluorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- 10 **Cpd 113**, 4-[1-(4-Chloro-3-fluorophenyl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 114**, 4-[1-(4-Chlorophenyl)-2-(2-ethoxypyridin-3-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- 15 **Cpd 115**, 4-[4-(4-Hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 116**, 2-Fluoro-5-[4-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 117**, 2-Fluoro-5-[4-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-2-(2-methoxyphenyl)-1H-imidazol-1-yl]benzonitrile;
- 20 **Cpd 118**, 2-Fluoro-4-[4-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-2-(2-methoxyphenyl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 119**, 2-Fluoro-4-[4-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-1-yl]benzonitrile;
- 25 **Cpd 120**, 4-[1-(4-Chlorophenyl)-2-(3-methoxypyridin-4-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 121**, 4-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-(trifluoromethyl)benzonitrile;

- Cpd 122**, 3-[1-(4-Chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]pyridin-2-ol;
- Cpd 123**, 2-Chloro-4-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- 5 **Cpd 124**, 1-(4-Chlorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyl-3,6-dihydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 125**, 1-(4-Bromo-3-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyl-3,6-dihydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 126**, 2-Fluoro-4-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyl-3,6-dihydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- 10 and pharmaceutically acceptable salt forms thereof.

For use in medicine, salts of compounds of Formula (I) refer to non-toxic

15 “pharmaceutically acceptable salts.” Other salts may, however, be useful in the preparation of compounds of Formula (I) or of their pharmaceutically acceptable salts thereof. Suitable pharmaceutically acceptable salts of compounds of Formula (I) include acid addition salts which can, for example, be formed by mixing a solution of the compound with a solution of a pharmaceutically acceptable acid such as hydrochloric acid,

20 sulfuric acid, fumaric acid, maleic acid, succinic acid, acetic acid, benzoic acid, citric acid, tartaric acid, carbonic acid or phosphoric acid. Furthermore, where the compounds of Formula (I) carry an acidic moiety, suitable pharmaceutically acceptable salts thereof may include alkali metal salts, such as sodium or potassium salts; alkaline earth metal salts, such as calcium or magnesium salts; and salts formed with suitable organic ligands, such

25 as quaternary ammonium salts. Thus, representative pharmaceutically acceptable salts include acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydrobromide,

hydrochloride, hydroxynaphthoate, iodide, isothionate, lactate, lactobionate, laurate, malate, maleate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, mucate, napsylate, nitrate, N-methylglucamine ammonium salt, oleate, pamoate (embonate), palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, salicylate, 5 stearate, sulfate, subacetate, succinate, tannate, tartrate, teoclate, tosylate, triethiodide and valerate.

Representative acids and bases that may be used in the preparation of pharmaceutically acceptable salts include acids including acetic acid, 2,2-dichloroacetic acid, acylated amino acids, adipic acid, alginic acid, ascorbic acid, L-aspartic acid, 10 benzenesulfonic acid, benzoic acid, 4-acetamidobenzoic acid, (+)-camphoric acid, camphorsulfonic acid, (+)-(1S)-camphor-10-sulfonic acid, capric acid, caproic acid, caprylic acid, cinnamic acid, citric acid, cyclamic acid, dodecylsulfuric acid, ethane-1,2-disulfonic acid, ethanesulfonic acid, 2-hydroxy-ethanesulfonic acid, formic acid, fumaric acid, galactaric acid, gentisic acid, glucoheptonic acid, D-gluconic acid, D-glucuronic acid, 15 L-glutamic acid, α -oxo-glutaric acid, glycolic acid, hippuric acid, hydrobromic acid, hydrochloric acid, (+)-L-lactic acid, ($\pm$)-DL-lactic acid, lactobionic acid, maleic acid, (-)-L-malic acid, malonic acid, ($\pm$)-DL-mandelic acid, methanesulfonic acid, naphthalene-2-sulfonic acid, naphthalene-1,5-disulfonic acid, 1-hydroxy-2-naphthoic acid, nicotinic acid, nitric acid, oleic acid, orotic acid, oxalic acid, palmitic acid, pamoic acid, phosphoric acid, 20 L-pyroglutamic acid, salicylic acid, 4-amino-salicylic acid, sebaic acid, stearic acid, succinic acid, sulfuric acid, tannic acid, (+)-L-tartaric acid, thiocyanic acid, p-toluenesulfonic acid and undecylenic acid; and bases including ammonia, L-arginine, benethamine, benzathine, calcium hydroxide, choline, deanol, diethanolamine, diethylamine, 2-(diethylamino)-ethanol, ethanolamine, ethylenediamine, N-methyl- 25 glucamine, hydrabamine, 1H-imidazole, L-lysine, magnesium hydroxide, 4-(2-hydroxyethyl)-morpholine, piperazine, potassium hydroxide, 1-(2-hydroxyethyl)-pyrrolidine, sodium hydroxide, triethanolamine, tromethamine and zinc hydroxide.

Embodiments of the present invention include prodrugs of compounds of Formula (I). In general, such prodrugs will be functional derivatives of the compounds that are

readily convertible *in vivo* into the required compound. Thus, in the methods of treating or preventing embodiments of the present invention, the term “administering” encompasses the treatment or prevention of the various diseases, conditions, syndromes and disorders described with the compound specifically disclosed or with a compound that may not be specifically disclosed, but which converts to the specified compound *in vivo* after administration to a patient. Conventional procedures for the selection and preparation of suitable prodrug derivatives are described, for example, in “Design of Prodrugs”, ed. H. Bundgaard, Elsevier, 1985.

Where the compounds according to embodiments of this invention have at least one chiral center, they may accordingly exist as enantiomers. Where the compounds possess two or more chiral centers, they may additionally exist as diastereomers. It is to be understood that all such isomers and mixtures thereof are encompassed within the scope of the present invention. Furthermore, some of the crystalline forms for the compounds may exist as polymorphs and as such are intended to be included in the present invention. In addition, some of the compounds may form solvates with water (i.e., hydrates) or common organic solvents, and such solvates are also intended to be encompassed within the scope of this invention. The skilled artisan will understand that the term compound as used herein, is meant to include solvated compounds of Formula (I).

Where the processes for the preparation of the compounds according to certain embodiments of the invention give rise to mixture of stereoisomers, these isomers may be separated by conventional techniques such as preparative chromatography. The compounds may be prepared in racemic form, or individual enantiomers may be prepared either by enantiospecific synthesis or by resolution. The compounds may, for example, be resolved into their component enantiomers by standard techniques, such as the formation of diastereomeric pairs by salt formation with an optically active acid, such as (-)-di-p-toluoyl-d-tartaric acid and/or (+)-di-p-toluoyl-l-tartaric acid followed by fractional crystallization and regeneration of the free base. The compounds may also be resolved by formation of diastereomeric esters or amides, followed by chromatographic separation and

removal of the chiral auxiliary. Alternatively, the compounds may be resolved using a chiral HPLC column.

One embodiment of the present invention is directed to a composition, including a pharmaceutical composition, comprising, consisting of, and/or consisting essentially of the (+)-enantiomer of a compound of Formula (I) wherein said composition is substantially free from the (-)-isomer of said compound. In the present context, substantially free means less than about 25 %, preferably less than about 10 %, more preferably less than about 5 %, even more preferably less than about 2 % and even more preferably less than about 1 % of the (-)-isomer calculated as.

$$\% (+) - \text{enantiomer} = \frac{(\text{mass}(+) - \text{enantiomer})}{(\text{mass}(+) - \text{enantiomer}) + (\text{mass}(-) - \text{enantiomer})} \times 100$$

Another embodiment of the present invention is a composition, including a pharmaceutical composition, comprising, consisting of, and consisting essentially of the (-)-enantiomer of a compound of Formula (I) wherein said composition is substantially free from the (+)-isomer of said compound. In the present context, substantially free from means less than about 25 %, preferably less than about 10 %, more preferably less than about 5 %, even more preferably less than about 2 % and even more preferably less than about 1 % of the (+)-isomer calculated as

$$\% (-) - \text{enantiomer} = \frac{(\text{mass}(-) - \text{enantiomer})}{(\text{mass}(+) - \text{enantiomer}) + (\text{mass}(-) - \text{enantiomer})} \times 100$$

During any of the processes for preparation of the compounds of the various embodiments of the present invention, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in *Protective Groups in Organic Chemistry, Second Edition*, J.F.W. McOmie, Plenum Press, 1973; T.W. Greene &

P.G.M. Wuts, *Protective Groups in Organic Synthesis*, John Wiley & Sons, 1991; and T.W. Greene & P.G.M. Wuts, *Protective Groups in Organic Synthesis, Third Edition*, John Wiley & Sons, 1999. The protecting groups may be removed at a convenient subsequent stage using methods known from the art.

5 Even though the compounds of embodiments of the present invention (including their pharmaceutically acceptable salts and pharmaceutically acceptable solvates) can be administered alone, they will generally be administered in admixture with a pharmaceutically acceptable carrier, a pharmaceutically acceptable excipient and/or a pharmaceutically acceptable diluent selected with regard to the intended route of
10 administration and standard pharmaceutical or veterinary practice. Thus, particular embodiments of the present invention are directed to pharmaceutical and veterinary compositions comprising compounds of Formula (I) and at least one pharmaceutically acceptable carrier, pharmaceutically acceptable excipient, and/or pharmaceutically acceptable diluent.

15 By way of example, in the pharmaceutical compositions of embodiments of the present invention, the compounds of Formula (I) may be admixed with any suitable binder(s), lubricant(s), suspending agent(s), coating agent(s), solubilizing agent(s), and combinations thereof.
Solid oral dosage forms, such as tablets or capsules, containing the compounds of the
20 present invention may be administered in at least one dosage form at a time, as appropriate. It is also possible to administer the compounds in sustained release formulations.

Additional oral forms in which the present inventive compounds may be administered include elixirs, solutions, syrups, and suspensions; each optionally containing flavoring agents and coloring agents.

25 Alternatively, compounds of Formula (I) can be administered by inhalation (intratracheal or intranasal) or in the form of a suppository or pessary, or they may be applied topically in the form of a lotion, solution, cream, ointment or dusting powder. For example, they can be incorporated into a cream comprising, consisting of, and/or consisting essentially of an aqueous emulsion of polyethylene glycols or liquid paraffin.

They can also be incorporated, at a concentration of between about 1 % and about 10 % by weight of the cream, into an ointment comprising, consisting of, and/or consisting essentially of a white wax or white soft paraffin base together with any stabilizers and preservatives as may be required. An alternative means of administration includes
5 transdermal administration by using a skin or transdermal patch.

The pharmaceutical compositions of the present invention (as well as the compounds of the present invention alone) can also be injected parenterally, for example intracavernosally, intravenously, intramuscularly, subcutaneously, intradermally or intrathecally. In this case, the compositions will also include at least one of a suitable
10 carrier, a suitable excipient, and a suitable diluent.

For parenteral administration, the pharmaceutical compositions of the present invention are best used in the form of a sterile aqueous solution that may contain other substances, for example, enough salts and monosaccharides to make the solution isotonic with blood.

15 For buccal or sublingual administration, the pharmaceutical compositions of the present invention may be administered in the form of tablets or lozenges, which can be formulated in a conventional manner.

By way of further example, pharmaceutical compositions containing at least one of the compounds of Formula (I) as the active ingredient can be prepared by mixing the
20 compound(s) with a pharmaceutically acceptable carrier, a pharmaceutically acceptable diluent, and/or a pharmaceutically acceptable excipient according to conventional pharmaceutical compounding techniques. The carrier, excipient, and diluent may take a wide variety of forms depending upon the desired route of administration (e.g., oral, parenteral, etc.). Thus for liquid oral preparations, such as suspensions, syrups, elixirs and
25 solutions, suitable carriers, excipients and diluents include water, glycols, oils, alcohols, flavoring agents, preservatives, stabilizers, coloring agents and the like; for solid oral preparations, such as powders, capsules and tablets, suitable carriers, excipients and diluents include starches, sugars, diluents, granulating agents, lubricants, binders, disintegrating agents and the like. Solid oral preparations also may be optionally coated

with substances, such as, sugars, or be enterically-coated so as to modulate the major site of absorption and disintegration. For parenteral administration, the carrier, excipient and diluent will usually include sterile water, and other ingredients may be added to increase solubility and preservation of the composition. Injectable suspensions or solutions may
5 also be prepared utilizing aqueous carriers along with appropriate additives, such as solubilizers and preservatives.

A therapeutically effective amount of a compound of Formula (I) or a pharmaceutical composition thereof includes a dose range from about 0.1 mg to about 3000 mg, or any particular amount or range therein, in particular from about 1 mg to about
10 1000 mg, or any particular amount or range therein; or, more particularly, from about 10 mg to about 500 mg, or any particular amount or range therein, of active ingredient in a regimen of about 1 to about 4 times per day for an average (70 kg) human; although, it is apparent to one skilled in the art that the therapeutically effective amount for a compound of Formula (I) will vary as will the diseases, syndromes, conditions, and disorders being
15 treated.

For oral administration, a pharmaceutical composition is preferably provided in the form of tablets containing about 0.01, about 10, about 50, about 100, about 150, about 200, about 250, and about 500 milligrams of a compound of Formula (I).

Advantageously, a compound of Formula (I) may be administered in a single daily
20 dose, or the total daily dosage may be administered in divided doses of two, three and four times daily.

Optimal dosages of a compound of Formula (I) to be administered may be readily determined and will vary with the particular compound used, the mode of administration, the strength of the preparation and the advancement of the disease, syndrome, condition or
25 disorder. In addition, factors associated with the particular subject being treated, including subject gender, age, weight, diet and time of administration, will result in the need to adjust the dose to achieve an appropriate therapeutic level and desired therapeutic effect. The above dosages are thus exemplary of the average case. There can be, of course, individual instances wherein higher or lower dosage ranges are merited, and such are within the scope

of this invention.

Compounds of Formula (I) may be administered in any of the foregoing compositions and dosage regimens or by means of those compositions and dosage regimens established in the art whenever use of a compound of Formula (I) is required for
5 a subject in need thereof.

As N-Type calcium channel blockers, the compounds of Formula (I) are useful in methods for treating and/or preventing a disease, a syndrome, a condition or a disorder in a subject, including an animal, a mammal and a human in which the disease, the syndrome, the condition or the disorder is affected by the modulation of the N-Type calcium channel.
10 Such methods comprise, consist of and/or consist essentially of administering to a subject, including an animal, a mammal, and a human in need of such treatment or prevention a therapeutically effective amount of a compound, salt or solvate of Formula (I). In particular, the compounds of Formula (I) are useful for preventing or treating pain, such as inflammatory pain or neuropathic pain, or diseases, syndromes, conditions or disorders
15 causing such pain.

GENERAL SYNTHETIC METHODS

Representative compounds of the present invention can be synthesized in accordance with the general synthetic methods described below and illustrated in the schemes and examples that follow. Since the schemes are an illustration, the invention
20 should not be construed as being limited by the chemical reactions and conditions described in the schemes. The various starting materials used in the schemes and examples are commercially available or may be prepared by methods well within the skill of persons versed in the art. The variables are as defined herein.
25


The following solvents, reagents or scientific terminology may be referred to by their abbreviations:

TLC Thin Layer Chromatography

	DCM	Dichloromethane
	DCE	1,2-Dichloroethane
	THF	Tetrahydrofuran
	MeOH	Methanol
5	EtOH	Ethanol
	IPA	Isopropyl alcohol
	n-BuOH	n-Butanol
	EtOAc	Ethyl acetate
	Et ₂ O	Diethyl ether
10	DMA	N,N-Dimethylacetamide
	DMF	N,N-Dimethylformamide
	Et ₃ N	Triethylamine
	DMSO	Dimethylsulfoxide
	DIPEA	Diisopropylethylamine (Hunig's base)
15	HEK	Human embryonic kidney
	MeI	Methyliodide
	NBS	N-Bromosuccinimide
	TFA	Trifluoroacetic acid
	PTSA	p-Toluenesulfonic acid
20	AcOH	Acetic acid
	Boc	tert-butoxycarbonyl
	Cat	Catalytic
	mL	milliliters
	mol	moles
25	mmol	millimoles
	h	hour or hours
	min	minute or minutes
	g	grams
	mg	milligrams

	μL	Microliters
	eq	Equivalents
	rt or RT	Room temperature, ambient, about 27 °C
	MS	Mass spectrometry
5	NA	Not available
	NE	No Effect
	tmhd	Dipivaloylmethanato

Scheme A illustrates a route for the synthesis of certain compounds of the present invention wherein R^2 is an optionally substituted phenyl or an optionally substituted heteroaryl as defined herein.



A compound of formula **a1** is either commercially available or may be prepared by methods known in the scientific literature. A compound of formula **a1** may be reacted with a compound of formula **a2** in the presence of an appropriate Lewis acid such as trimethylaluminum or the like, in an aprotic organic solvent such as toluene, at a


temperature from about 0 °C to about 70 °C, to afford a compound of formula **a3**. A compound of formula **a3** may be treated with a compound of formula **a4** in the presence of an appropriate inorganic base such as sodium bicarbonate, at about 100 °C, to afford a compound of formula **(I)-A**. A compound of formula **(I)-A** may be treated with NCS to afford compound of formula **(I)-A1**.

Alternatively, an appropriately substituted carboxylic acid compound of formula **a6** (commercially available or prepared by methods known in the scientific literature) may be converted to its corresponding acid chloride by the action of an appropriate chlorinating agent such as oxalyl chloride, thionyl chloride, or the like, in an organic solvent such as dichloromethane, at a temperature of about 0 °C to room temperature, to afford a compound of formula **a7**. A compound of formula **a7** may be converted to an amide of formula **a8** via treatment with an appropriately substituted amine of formula **a2**, in the presence of a non-nucleophilic tertiary amine base such as triethylamine, in an organic solvent such as dichloromethane, at a temperature of about 0 °C to room temperature.

Conversion of a compound of formula **a8** to a compound of formula **a3** may be accomplished by treatment with a chlorinating reagent such as phosphorus pentachloride, in an organic solvent such as chloroform, at about room temperature, followed by reaction with ammonia gas at a temperature of about 0 °C.

Scheme B illustrates a route for the synthesis of certain compounds of the present invention wherein R^1 is an optionally substituted phenyl as defined herein.

Scheme B




A compound of formula **b1** may be converted to its corresponding phenyl cyanide in the presence of zinc cyanide, a transition metal catalyst, dimethylformamide, and suitable ligands, to afford compounds of Formula (I)-B wherein R¹ is a cyano-substituted phenyl ring.

5

Scheme C illustrates a route for the synthesis of certain compounds of the present invention wherein R¹ is an optionally substituted phenylsulfonyl or alkylsulfonyl as defined herein.

Scheme C




10

A compound of formula **a1** may be reacted with ammonium chloride, in the presence of an appropriate Lewis acid such as trimethylaluminum or the like, in an aprotic organic solvent such as toluene, at a temperature from about 0 °C to about 70 °C, to afford a compound of formula **c1**. A compound of formula **c1** may be reacted with a compound of formula **a4** in the presence of an appropriate inorganic base such as sodium bicarbonate, at about 100 °C, to afford a compound of formula **c2**. A compound of formula **c2** may be treated with an appropriately substituted sulfonyl chloride in the presence of a non-nucleophilic tertiary amine base such as triethylamine, in an organic solvent such as dichloromethane, at a temperature of about 0 °C to room temperature, to afford a compound of formula (I)-C.

20

Scheme D illustrates a route for the synthesis of certain compounds of the present invention wherein R^{1d} is an optionally substituted phenylmethyl or heteroaryl-methyl group as defined by the scope of the invention.


Scheme D

A compound of formula **c2** may be reacted with an inorganic base such as NaH, K_2CO_3 or CS_2CO_3 , in the presence of an appropriate phenylmethyl-halide or heteroarylmethyl-halide
 5 to afford a compound of formula **(I)-D**.

Scheme E illustrates a route for the synthesis of certain compounds of the present invention wherein G is G2.

Scheme E

10




Commercially available ethoxy ethylene may be treated with a strong organic base such as an alkyllithium base, organomagnesium bromide, or the like, at 0 °C, followed by addition
 15 of compound **e1**, to afford compound **e2**. Compound **e2** may be treated with a brominating reagent such as hydrogen bromide in acetic acid, NBS, or the like, at 0 °C to room

temperature, to afford compound **e3**. A compound of formula **a3** may be treated with a compound of formula **e3** in the presence of an appropriate inorganic base such as sodium bicarbonate, at about 100 °C, to afford a compound of formula (I)-E.

- 5 Scheme F illustrates a route for the conversion of certain compounds of Formula (I)-F1, wherein G is G2, to compounds of Formulae (I)-F2 and (I)-F3.

Scheme F



- A compound of formula (I)-F1, prepared according to the methods described in Scheme E, may be converted to a compound of formula (I)-F2 (wherein G is G3) in the presence of zinc cyanide, a transition metal catalyst, dimethylformamide, and suitable ligands.
- Reaction of a compound of formula (I)-F2 may occur by the action of a hydride source such as triphenylsilane, or the like, in the presence of Mn(tmhd)₃, at a temperature of about 0 °C to room temperature in an oxygen atmosphere, to afford a compound of formula (I)-F3.
- 15 F3.

Specific Examples

- Yields reported herein refer to purified products (unless specified) and are not optimized. Analytical TLC was performed on Merck silica gel 60 F254 aluminium-backed plates. Compounds were visualized by UV light and/or stained either with iodine, potassium permanganate or ninhydrin solution. Flash column chromatography was performed on silica gel (100-200 M) or flash chromatography. ¹H-NMR spectra were recorded on a Bruker Avance-400 MHz spectrometer with a BBO (Broad Band Observe) and BBFO (Broad Band Fluorine Observe) probe. Chemical shifts (δ) are expressed in
- 20


parts per million (ppm) downfield by reference to tetramethylsilane (TMS) as the internal standard. Splitting patterns are designated as s (singlet), d (doublet), t (triplet), q (quartet), m (multiplet) and br s (broad singlet). Coupling constants (J) are given in hertz (Hz). LC-MS analyses were performed using the Electrospray Ionization (ESI) technique.

5

A. Preparation of Chemical Intermediates

Example 1

Preparation of intermediate A1; N-(4-cyanophenyl)-2-methoxybenzimidamide



10













To a stirred solution of 2-methoxybenzonitrile (10 g, 0.075 mol) in toluene (200 ml) at 0 °C was added trimethylaluminum (90 mL, 0.09 mol) drop-wise over a period of 10 min. The reaction mixture was then stirred at room temperature for 3 h followed by addition of 4-aminobenzonitrile (8.6g, 0.074mol) in toluene 100 mL. The reaction mixture was heated to 70 °C for 16 h. After confirming the completion by LCMS the reaction mixture was quenched with ice-cold water; the aqueous layer was filtered through a pad of diatomaceous earth, and the resultant filtrate was washed with ethyl acetate. The aqueous layer was extracted with ethyl acetate (2 x 100 mL). The combined organic extracts were washed with brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as a white solid. (12.8 g, 68%).

15

20

By using analogous protocols as described in the foregoing example the intermediates described in Table 1 have been prepared using appropriate starting materials






Table 1

Intermediate No.	Structure	Intermediate No.	Structure
A2		A33	
A3		A34	
A4		A35	
A5		A36	
A6		A37	
A7		A38	

Intermediate No.	Structure	Intermediate No.	Structure
A8		A39	
A9		A40	
A10		A41	
A11		A42	
A12		A43	
A13		A44	
A14		A45	


Intermediate No.	Structure	Intermediate No.	Structure
A15		A46	
A16		A47	
A17		A48	
A18		A49	
A19		A50	
A20		A51	
A21		A52	

Intermediate No.	Structure	Intermediate No.	Structure
A22		A53	
A23		A54	
A24		A55	
A25		A56	
A26		A57	
A27		A58	
A28		A59	
A29		A60	

Intermediate No.	Structure	Intermediate No.	Structure
A30		A61	
A31		A62	
A32			

Example 2

Preparation of intermediate A-63 ; 2-methoxy-N-(6-(trifluoromethyl)pyridin-3-yl)nicotinimidamide



5

Step 1. 2-Methoxy-nicotinoyl chloride: To a stirred solution of 2-methoxynicotinic acid (2 g, 0.013 mol) in DCM (20 mL) under a N₂ atmosphere was added oxalyl chloride (2 mL, 0.026 mol) drop-wise at 0 °C followed by the addition of a catalytic amount of DMF (2 drops). The reaction mixture was stirred at room temperature for 2 h. Upon completion of the reaction, the solvent was removed under reduced pressure in an inert atmosphere to provide title compound as a thick liquid (2.1 g, 98%).

Step 2. 2-Methoxy-N-(6-trifluoromethyl-pyridin-3-yl)-nicotinamide To a stirred solution of 6-(trifluoromethyl)pyridin-3-amine (1.8 g, 0.0011 mol) in DCM (20 mL) under a N₂ atmosphere was added triethylamine (4.9 mL, 0.035 mol) at 0 °C followed by the drop-wise addition of 2-methoxy-nicotinoyl chloride (2 g, 0.0011 mol) in DCM (5 mL). The reaction mixture was stirred at room temperature for 2 h. Upon completion of the

reaction, the solvent was evaporated and resultant residue was partitioned between DCM and water. The organic layer was separated and washed with brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated to afford the title compound as an off-white solid (2.02 g, 61.3%). LCMS: 298.1 $[\text{M}+\text{H}]^+$.

- 5 **Step 3. 2-methoxy-N-(6-(trifluoromethyl)pyridin-3-yl)nicotinimidamide:** To a stirred solution of PCl_5 (0.17 g, 0.0084 mol) in chloroform (5 mL) at room temperature under a N_2 atmosphere was added 2-methoxy-N-(6-trifluoromethyl-pyridin-3-yl)-nicotinamide (0.5 g, 0.00168 mol). The reaction mixture was stirred at 61 °C for 30 min. The reaction mixture was cooled to 0 °C followed by bubbling of ammonia gas for 1 h. The reaction
- 10 was quenched with NaHCO_3 , partitioned between DCM and water. The organic layer was separated and washed with brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated to afford title compound as an off-white solid (0.25 g, 50.7%). LCMS- 297.1 $[\text{M}+\text{H}]^+$. By using analogous protocols as described in the foregoing example the compounds described in **figure 1** have been prepared using appropriate starting materials





Figure 1

Example 3

Preparation of intermediate A71; 2-bromo-1-(2,2,6,6-tetramethyltetrahydro-2H-pyran-yl)ethanone




- 5 **Step 1. 2,2,6,6-tetramethyldihydro-2H-pyran-4(3H)-one:** To a stirred solution of 2,6-dimethylhepta-2,5-dien-4-one (100 g, 0.724 mol) was added 6N HCl (600 mL), then the reaction mixture was heated to 45 °C for 7 days. Upon completion, the reaction was quenched with ice cold water and extracted with ethyl acetate (4 x 150 mL). The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and
- 10 concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as a yellow liquid. (yield 28 g, 25%)

- Step 2. 2,2,6,6-tetramethyltetrahydro-2H-pyran-4-carbonitrile:** To a stirred solution of 2,2,6,6-tetramethyldihydro-2H-pyran-4(3H)-one (30 g, 0.192 mol) in dimethoxyethane (400 mL) was added tosylmethyl isocyanide (48.7 g, 0.249 eq) followed by the addition of tert-butyl alcohol (24.1 g, 0.326) at room temperature. The reaction mixture was cooled to 0 °C followed by portion-wise addition of potassium tert-butoxide (53.8 g, 0.48 mol). It was stirred at room temperature for 12 h. The reaction mixture was filtered after dilution with diethyl ether at 0 °C and the residue was further washed with diethyl ether. The resultant filtrate was concentrated to provide the title compound as a yellow semi-solid (22 g, 68%)
- Step 3. 2,2,6,6-tetramethyltetrahydro-2H-pyran-4-carboxylic acid:** To a stirred solution of 2,2,6,6-tetramethyltetrahydro-2H-pyran-4-carbonitrile (22 g, 0.131 mol) in water (400 mL) was added KOH (45 g, 0.815 mol) and the reaction mixture was allowed to reflux for 8 h. The completion of the reaction was confirmed by TLC. The reaction mixture was diluted with water and extracted with dichloromethane (3x 100 mL). The aqueous layer was acidified with 1N HCl, extracted with dichloromethane. The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as a pale yellow solid (13 g, 52%).
- Step 4. 2,2,6,6-tetramethyltetrahydro-2H-pyran-4-carbonyl chloride:** To a stirred solution of 2,2,6,6-tetramethyltetrahydro-2H-pyran-4-carboxylic acid (2.7 g, 0.0144 mol) in DCM (30 mL) was added oxalyl chloride (3.6 g, 0.0289 mol) at 0 °C. The reaction mixture was allowed stirred for 3 h at room temperature and then concentrated to provide the title compound as a black thick liquid (2.6 g, crude). It was used in the next step without further purification.
- Step 5. 2-diazo-1-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)ethanone:** To a stirred solution of 2,2,6,6-tetramethyltetrahydro-2H-pyran-4-carbonyl chloride (2.6 g, 0.012 mol) in dichloromethane (20 mL) was added trimethylsilyldiazomethane (2.9 g, 0.025 mol) at 0 °C after which it was stirred for 12 h at rt. The completion of the reaction was confirmed

by TLC. The reaction mixture was concentrated under reduced pressure to provide a crude residue of the title compound as a black thick liquid which was used in the next step without further purification. (2.5 g, crude)

Step 6. 2-bromo-1-(2,2,6,6-tetramethyltetrahydro-2H-pyran-yl)ethanone:




A71

5 To a stirred solution of 2-diazo-1-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)ethanone. (2.5 g, 0.011 mol) in diethyl ether (20 mL) was added aqueous hydrobromic acid (8 mL) drop wise at 0 °C. The stirring was continued for 3 h at 0 °C. The completion of the reaction was confirmed by TLC. The reaction mixture was quenched with 10% NaHCO₃ aqueous solution (50 mL). The combined organic layers were dried over Na₂SO₄, filtered, and concentrated under reduced pressure to provide a crude residue of the title compound as a yellow liquid which was used in the next step without further purification. (2.4 g, crude).

15 **Step 7 . 1-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)ethanone:** To a stirred solution of 2,2,6,6-tetramethyltetrahydro-2H-pyran-4-carbonitrile (2 g, 0.012 mol) in THF (25 mL) was added MeMgBr (3M in Et₂O) (4.7 mL, 0.0036 mol) at -78 °C and the reaction was slowly allowed to reach room temperature and stirred for overnight. After completion of the reaction, the reaction was quenched with saturated NH₄Cl solution at 0 °C and extracted with EtOAc (2 x 75 mL). The combined organic extracts were washed with 20 brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as thick liquid. (1.54 g, 70 %).

Step 8. 2-bromo-1-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)ethanone:


**A71**

To a stirred solution of 1-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)ethanone (0.15 g, 0.0008 mol) in EtOH (3 mL) was added Br₂ solution (0.06 mL, 0.0012 mol) at 0 °C and the reaction was stirred for overnight. After completion of the reaction, the reaction was
 5 diluted with ice water and extracted with DCM (2 x 50 mL). The combined organic extracts were washed with NaHCO₃, brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide the crude title compound which was used in the subsequent stage without purification. Yield: 0.35 g (Crude).

10

Example 4

Preparation of intermediate A72; 2-bromo-1-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)propan-1-one




15 **Step 1.** 1-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)propan-1-one: To a stirred solution of 2,2,6,6-tetramethyltetrahydro-2H-pyran-4-carbonitrile (0.5 g, 0.003 mol) in THF (5 mL) was added EtMgBr (3M in Et₂O) (1.19 mL, 0.0036 mol) at -78 °C and the reaction was slowly allowed to reach room temperature and stirred for overnight. After completion of the reaction, the reaction was quenched with saturated NH₄Cl solution at 0 °C and

extracted with EtOAc (2 x 10 mL). The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as colorless solid. (0.25 g, 42.3%)

Step 2. 2-bromo-1-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)propan-1-one: To a stirred solution of 1-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)propan-1-one (0.2 g, 0.001 mol) in EtOH (5 mL) was added Br₂ solution (0.078 mL, 0.0015 mol) at 0 °C and the reaction was stirred for overnight. After completion of the reaction, the reaction was diluted with ice water and extracted with DCM (2 x 50 mL). The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide the crude title compound which was used in the subsequent stage without purification. Yield: 0.35 g (Crude).

Example 5

Preparation of intermediate A74: 2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole




Step 1. 2-methoxybenzimidamide: To a stirred solution of ammonium chloride (6 g, 0.113 mol) in toluene (80 mL) at 0 °C was added trimethylaluminium (2.97 g, 0.041 mol)

drop wise for a period of 10 min. The reaction mixture was stirred for 3 h at room temperature followed by addition of 2-methoxybenzonitrile (5 g, 0.037 mol) in toluene (10 mL). The reaction was then heated to 70 °C for 16 h. After completion, the reaction was quenched with ice cold water, filtered through diatomaceous earth, and washed with ethyl acetate. The aqueous layer was extracted with ethyl acetate (2 x). The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as a solid (4.5 g, 80%).

Step 2. 2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole: To a stirred solution of 2-methoxybenzimidamide **A73** (0.6 g, 0.004 mol) in 1,4- dioxane (20 mL) was added 2-bromo-1-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)ethanone **A71** (1.04 g, 0.004 mol), followed by the addition of NaHCO₃ (1 g, 0.012 mol). The reaction mixture was refluxed for 5 h. The reaction was quenched with ice cold water after completion. The aqueous layer was extracted with ethyl acetate (3 x 25 mL). The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as a white solid (0.5 g crude).

Example 6

Preparation of intermediate A75: 2-bromo-1-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)ethanone




5

Step 1. 1-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)ethanone: To a stirred solution of ethoxyethene (1.84 g, 25.5 mmol) in THF (40 mL) was added ^tBuLi (16 mL, 25.6 mmol) at -78 °C. The reaction mixture was slowly allowed to warm to 10 °C followed by addition of 2,2,6,6-tetramethyldihydro-2H-pyran-4(3H)-one (2 g, 12.8 mmol). The mixture was stirred for 16 h at room temperature. The reaction was quenched by the addition of HCl (3 mL) in aqueous methanol (20 mL, MeOH : H₂O = 1:1). The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide the crude title compound as an off-white solid (1.2 g) which was used in the next step without further purification.

15 **Step 2. (See below)**

Step 3. 2,2,6,6-Tetramethyl-4-trimethylsilanylethynyl-tetrahydro-pyran-4-ol. To a solution of ethynyl-trimethyl-silane (5.5 mL, 38.4 mmol) in dry THF (25 mL) was added

- n-BuLi (32 mL, 38.4 mmol) at -78 °C and the mixture was stirred at that temperature for 45 min followed by addition of 2,2,6,6-tetramethyl-tetrahydro-pyran-4-one (5.0 g, 32 mmol) in dry THF (25 mL) at -78 °C. The mixture was stirred for 1 h and then quenched with saturated NH₄Cl solution and extracted with ethyl acetate (3 × 100 mL). The
- 5 combined organic extract was washed with water and brine solution, dried over Na₂SO₄, filtered, and concentrated under reduced pressure to afford the product as sticky white solid. Crude product was forwarded for next stage without purification. Yield: 8.0 g, crude.
- Step 4. 4-Ethynyl-2,2,6,6-tetramethyl-tetrahydro-pyran-4-ol.** To a stirred solution of 2,2,6,6-tetramethyl-4-trimethylsilanylethynyl-tetrahydro-pyran-4-ol (8.0 g, 0.031 mol) in
- 10 MeOH (120 mL) was added potassium carbonate (10.86 g, 0.078 mol) and the mixture was stirred at rt for 12 h. MeOH was evaporated to obtain a residue, to which water was added. The mixture was extracted with ethyl acetate (3 × 100 mL). The organic portion was washed with brine solution, dried over Na₂SO₄, filtered, and concentrated under reduced pressure to afford the crude product, which was triturated with n-pentane to afford the
- 15 purified product as an off-white solid. (Yield: 3.5g, 61%).
- Step 5. 1-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)ethanone.** To a stirred solution of HgO (0.185 g, 0.085 mmol) in acetone/H₂O (30 mL/ 4 mL) was added H₂SO₄ (0.03 mL) and the mixture was heated to 60 °C, followed by addition of 4-ethynyl-2,2,6,6-tetramethyl-tetrahydro-pyran-4-ol (**3**) (2.6 g, 1.42 mmol) in acetone (10 mL) drop
- 20 wise at that temperature. After consumption of starting material (TLC), the reaction mixture was concentrated, diluted with cold water (50 mL) and extracted with EtOAc (3 x 50 mL). The combined organic extracts were washed with water and brine, dried over Na₂SO₄, filtered, and concentrated under reduced pressure to afford the crude product. Yield: 2.5 g (87%).
- 25 **Step 2. 2-bromo-1-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)ethanone.**




To a stirred solution of 1-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)ethanone (1.1 g, 0.0054 mol) in CHCl_3 (30 mL) was added two drops of HBr/AcOH at 0°C , followed by addition of Br_2 (0.3 mL, 0.006 mol) in CHCl_3 (2 mL). The temperature was gradually increased to room temperature and the reaction mixture was stirred for 4 h. After completion of the reaction, the reaction mixture was diluted with DCM (100 mL). The DCM solution was washed with NaHCO_3 , brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure to provide the crude title compound as a thick liquid which is used directly for the next stage without further purification. Yield: 1.2 g (Crude).

B. Preparation of Final Compounds

Example 7

Compound 1: 4-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile







Reaction conditions: (a) NaHCO_3 , dioxane, 100°C



- To a stirred solution of N-(4-cyanophenyl)-2-methoxybenzimidamide **A1** (2.5 g, 0.0099 mol) and 2-bromo-1-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)ethanone **A71** (3.24 g, 0.0123 mol) in dioxane (50 mL) was added NaHCO₃ (1.67 g, 0.0199 mol) and the reaction mixture was refluxed at 100 °C for 5 h. After completion of the reaction the mixture was concentrated under reduced pressure and quenched with ice cold water. The aqueous layer was extracted with ethyl acetate (2 x 150 mL). The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide a residue. The residue was purified by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane to afford the title compound **1** as a white solid. ¹H NMR (400 MHz, CDCl₃) δ 7.61 – 7.57 (m, 3H), 7.38 – 7.34 (m, 1H), 7.21 (d, *J* = 8.4 Hz, 2H), 7.05 (t, *J* = 6.8 Hz, 1H), 6.96 (s, 1H), 6.71 (d, *J* = 8.4 Hz, 1H), 3.27 (s, 3H), 3.24 - 3.22 (m, 1H), 2.06 – 2.02 (m, 2H), 1.50 – 1.44 (m, 2H), 1.36 (s, 6H), 1.25 (s, 6H); LCMS: 416.3 (M+H)⁺.
- Using analogous protocols to those described in Example 7, the compounds described in Table 2 have been prepared using an appropriately substituted amidine intermediate.



Table 2



Cpd No.	Structure	Compound Name	Analytical data
---------	-----------	---------------	-----------------



Cpd No.	Structure	Compound Name	Analytical data
2		4-(1-(4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) 8.29-8.26 (m, 2H), 7.50 (d, <i>J</i> = 4.4 Hz, 1H), 7.33 (s, 1H), 7.28 - 7.16 (m, 4H), 3.32 (s, 3H), 3.15-3.08 (m, 1H), 1.90 (d, <i>J</i> = 2.8 Hz, 2H), 1.40-1.39 (m, 2H), 1.28 (s, 6H), 1.15 (s, 6H). LCMS: 410.4 [M+H] ⁺
3		4-(1-(4-ethoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) δ = 8.28 - 8.29 (m, 2H), 7.94-7.93 (m, 1H), 7.53-7.45 (m, 2H), 7.33 (s, 1H), 6.80 (d, <i>J</i> = 8.8 Hz, 1H), 4.29-4.24 (q, 2H), 3.41 (s, 3H), 3.14 - 3.08 (m, 1H), 1.93-1.89 (m, 2H), 1.36 - 1.24 (m, 3H), 1.21-1.12 (m, 6H), 1.11-1.09 (m, 2H). LCMS: 436.2 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
4		1-(4-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole	^1H NMR (400 MHz, DMSO- d_6) δ = 7.45 (d, J = 6.8 Hz, 1H), 7.36 (t, J = 6.8 Hz, 1H), 7.13 - 7.12 (m, 5H), 7.01 (J = 7.6 Hz, 1H), 6.87 (d, J = 8 Hz, 1H), 3.23 (s, 3H), 3.10 (d J = 11.6 Hz, 1H), 1.91 (d, J = 11.6 Hz, 2H), 1.36 (t, J = 12.8 Hz, 2H), 1.28(s, 6H), 1.15 (s, 6H). LCMS: 409.2 $[\text{M}+\text{H}]^+$
5		5-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-methylpyridine	^1H NMR (400 MHz, DMSO- d_6) δ = 8.18 (s, 1 H), 7.35-7.5 (m, 3 H), 7.28 (s, 1 H), 7.22 (d, J = 8.4 Hz, 1 H), 7.02 (t, J = 7.6 Hz, 1 H), 6.88 (d, J = 8.0 Hz 1 H), 3.23 (s, 3 H), 3.03-3.14 (m, 1 H), 2.43 (s, 3 H), 1.93 (t, J = 2.4 Hz, 2 H), 1.36 (t, J = 12.8 Hz, 2 H), 1.28 (s, 6 H), 1.15 (s, 6 H). LCMS: 406.5 $[\text{M}+\text{H}]^+$.



Cpd No.	Structure	Compound Name	Analytical data
6		2-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-5-methylpyridine	¹ H NMR (400 MHz, DMSO-d ₆) δ = 8.20 (s, 1H), 7.60 (d, <i>J</i> = 8 Hz, 1H), 7.50 (d, <i>J</i> = 1.2 Hz, 1H), 7.35 - 7.39 (m, 2H), 7.03 (t, <i>J</i> = 7.6 Hz, 1H), 6.87 - 6.93 (m, 2H), 3.16 (s, 3H), 3.07-3.13 (m, 1H), 2.26 (s, 3H), 1.93 (d, <i>J</i> = 2.8 Hz, 2H), 1.39-1.33 (m, 2H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS: 406.3 [M+H] ⁺
7		2-(2-methoxyphenyl)-1-(4-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.40 (d, <i>J</i> = 7.6 Hz, 1H), 7.31 (t, <i>J</i> = 8 Hz, 1H), 7.12 (s, 1H), 6.99 (q, <i>J</i> = 8.8 Hz, <i>J</i> = 8.8 Hz, 1H), 6.87 (d, <i>J</i> = 8 Hz, 1H), 3.71 (s, 3H), 3.26 (s, 3H), 3.08 (t, <i>J</i> = 12.4 Hz, 1H), 1.90 (dd, <i>J</i> = 2.8 Hz, <i>J</i> = 12.8 Hz, 2H), 1.36 (t, <i>J</i> = 12.8 Hz, 2H), 1.28 (s, 6H), 1.15 (s, 6H). LCMS: 421.2 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
8		5-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-methylpyrimidine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.47 (s, 2 H), 7.51 (d, <i>J</i> = 1.6 Hz, 1 H), 7.43 (d, <i>J</i> = 1.2 Hz, 1 H), 7.39 (s, 1 H), 7.08-7.04 (m, 1 H), 6.90 (d, <i>J</i> = 8.4 Hz 1 H), 3.27 (s, 3 H), 3.14-3.03 (m, 1 H), 2.60 (s, 3 H), 1.94 (d, <i>J</i> = 2.8 Hz, 2 H), 1.42-1.36 (m, 2 H), 1.28 (s, 6 H), 1.15 (s, 6 H) LCMS: 407.5 [M+H] ⁺
9		3-(1-(3-chlorobenzyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) δ = 8.25 (d, <i>J</i> = 4 Hz, 1H), 7.67 (d, <i>J</i> = 8 Hz, 1H), 7.30 (t, <i>J</i> = 5.2 Hz, 2H), 7.05 (t, <i>J</i> = 11.2 Hz, 3H), 6.90 (s, 1H), 4.99 (s, 2H), 3.78 (s, 3H), 3.08-3.01 (m, 1H), 1.86 (q, <i>J</i> = 2.8 & 2.9 Hz, 2H), 1.32-1.40 (m, 2H), 1.26 (s, 6H), 1.12 (s, 6H). LCMS: 440.2 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
10		3-(1-(4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	^1H NMR (400 MHz, DMSO- d_6) δ = 8.17 (d, J = 4.8 Hz, 1H), 7.90 (d, J = 7.6 Hz, 1H), 7.27 (s, 1H), 7.1-7.26 (m, 4H), 7.07 (d, J = 5.2 Hz, 1H), 3.1 (s, 1H), 1.89 (t, J = 15.6 Hz, 2H), 1.36 (t, J = 12.8 Hz, 2H), 1.28 (s, 6H), 1.14 (s, 6H). LCMS: 410.5 $[\text{M}+\text{H}]^+$
11		3-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	^1H NMR: (400 MHz, CDCl_3) δ 8.16 (d, J = 3.2 Hz, 1H), 7.88 (d, J = 6.4 Hz, 1H), 7.30 – 7.27 (m, 2H), 7.04 (d, J = 8.4 Hz, 2H), 6.97 – 6.93 (m, 2H), 3.48 (s, 3H), 3.25 – 3.19 (m, 1H), 2.03 (d, J = 10.8 Hz, 2H), 1.51 – 1.44 (m, 2H), 1.36 (s, 6H), 1.25 (s, 2H). LCMS 426.2 $(\text{M}+\text{H})^+$



Cpd No.	Structure	Compound Name	Analytical data
12		4-(2-(3-methoxypyridin-2-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) δ = 8.19 (s, 1H), 7.75 - 7.85 (m, 2H), 7.40 - 7.50 (m, 3H), 7.26 (d, <i>J</i> = 8 Hz, 2H), 3.45 (s, 3H), 3.05 - 3.20 (m, 1H), 1.92 (d, <i>J</i> = 11.6 Hz, 2H), 1.30 - 1.40 (m, 2H), 1.29 (s, 6H), 1.16 (s, 6H). LCMS: 417.5 [M+H] ⁺
13		4-(1-(4-chloro-3-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.35 (d, <i>J</i> =4.8 Hz, 1H), 8.18 (s, 1H), 7.53 (d, <i>J</i> =4.8 Hz, 1H), 7.36-7.32 (m, 1H), 6.99-6.96 (m, 2H), 6.82 (d, <i>J</i> =8.4 Hz, 1H), 3.47 (s, 3H), 3.25-3.19 (m, 1H), 2.02-1.99 (m, 2H), 1.50-1.43(m, 2H), 1.35 (s, 6H), 1.28 (s, 6H). LCMS: 444.4 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
14		2-chloro-5-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	^1H NMR (CDCl_3 , 400 MHz) δ 7.57-7.55 (m, 1H), 7.47-7.46 (m, 1H), 7.42-7.35 (m, 1H), 7.26 (s, 1H), 7.06-7.03 (m, 1H), 6.92 (s, 1H), 6.74 (d, $J=8.4$ Hz, 1H), 3.37 (s, 3H), 3.26-3.19 (m, 1H), 2.04-2.01 (m, 2H), 1.50-1.43 (m, 2H), 1.35 (s, 6H), 1.29 (s, 6H). LCMS: 450.5 $[\text{M}+\text{H}]^+$
15		4-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-methylbenzonitrile	^1H NMR (CDCl_3 , 400 MHz) δ 8.20-8.18 (m, 1H), 7.94-7.91 (m, 1H), 7.55-7.53 (m, 1H), 7.11 (s, 1H), 7.01-6.97 (m, 3H), 3.44 (s, 3H), 3.26-3.19 (m, 1H), 2.50 (s, 3H), 2.04-2.00 (m, 2H), 1.50-1.42 (m, 2H), 1.35 (s, 6H), 1.26 (s, 6H). LCMS: 431.5 $[\text{M}+\text{H}]^+$



Cpd No.	Structure	Compound Name	Analytical data
16		2-methoxy-3-(4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1-(2,3,4-trifluorophenyl)-1H-imidazol-2-yl)pyridine	¹ H NMR (400 MHz, DMSO) : 8.20-8.18 (m, 1H), 7.94-7.91 (m, 1H), 7.29-7.35 (m, 2H), 7.10-7.07 (m, 2H), 3.45 (s, 3H), 3.16-3.10 (m, 1H), 1.93-1.89 (m, 2H), 1.39-1.33 (m, 2H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS: 446.4 [M+H] ⁺
17		2-(1-(3-chloro-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) 8.18-8.17 (m, 1H), 7.47-7.35 (m, 5H), 7.02-7.00 (m, 1H), 3.51 (s, 3H), 3.13-3.07 (m, 1H), 1.92 (d, J=11.6 Hz, 2H), 1.37 (d, J=13.2 Hz, 2H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS: 444.4 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
18		4-(1-(3-chloro-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) 8.31-8.28 (m, 2H), 7.55-7.51 (m, 2H), 7.44-7.39 (m, 2H), 7.12-7.08 (m, 1H), 3.37 (s, 3H), 3.14-3.07 (m, 1H), 1.93-1.89 (m, 2H), 1.39-1.32 (m, 2H), 1.28 (s, 6H), 1.15 (s, 6H). LCMS: 444.4 [M+H] ⁺
19		2-(1-(4-chloro-3-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) 8.19-8.17 (s, 1H), 7.55-7.51 (m, 1H), 7.48-7.42 (m, 2H), 7.37 (s, 1H), 7.29-7.26 (m, 1H), 6.87-6.85 (m, 1H), 3.51 (s, 3H), 3.14-3.07 (m, 1H), 1.93-1.89 (m, 2H), 1.39-1.33 (m, 2H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS: 444.4 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
20		3-(1-(3-chloro-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-4-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) 8.51 (s, 1H), 8.48-8.47 (m, 1H), 7.54-7.52 (m, 1H), 7.43-7.38 (m, 2H), 7.15-7.11 (m, 1H), 6.98-6.97 (m, 1H) 3.40 (s, 3H), 3.05-3.15 (m, 1H), 1.94-1.90 (m, 2H), 1.39-1.32 (m, 2H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS: 444.4 [M+H] ⁺
21		4-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-methylbenzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) 7.70 (d, <i>J</i> =8 Hz, 1H), 7.49 (d, <i>J</i> =7.6 Hz, 1H), 7.41 (t, 1H), 7.34-7.33 (m, 2H), 7.05 (t, 1H), 6.98-6.96 (m, 1H), 6.89 (d, <i>J</i> =8 Hz, 1H), 3.18 (s, 3H), 3.13-3.06 (m, 1H), 2.42 (s, 3H), 1.94-1.90 (m, 2H), 1.39-1.32 (m, 2H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS: 430.5 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
22		5-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-methylbenzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) 7.58 (s, 1H), 7.47 (d, J=6.8 Hz, 1H), 7.38 (d, J=7.6 Hz, 2H), 7.31 (s, 1H), 7.25 (d, J=7.2 Hz, 1H), 7.04 (t, 1H), 6.89 (d, J=8.4 Hz, 1H), 3.22 (s, 3H), 3.12-2.98 (m, 1H), 2.44 (s, 3H), 1.92 (d, J=12.8Hz, 2H), 1.44-1.35 (m, 2H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS: 430.3 [M+H] ⁺
23		3-(1-(3-fluoro-4-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) 8.20-8.18 (m, 1H), 7.89-7.87 (m, 1H), 7.25 (s, 1H), 7.16-7.06 (m, 3H), 6.85 (d, J=7.6 Hz, 1H), 3.81 (s, 3H), 3.41 (s, 3H), 3.12-3.05 (m, 1H), 1.93-1.89 (m, 2H), 1.39-1.32 (m, 2H), 1.28 (s, 6H), 1.15 (s, 6H). LCMS: 440.5 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
24		3-(1-(4-fluoro-3-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) 8.20-8.18 (m, 1H), 7.91-7.88 (m, 1H), 7.32 (s, 1H), 7.20-7.15 (m, 1H), 7.10-7.07 (m, 1H), 7.00-6.98 (m, 1H), 6.63-6.60 (m, 1H), 3.70 (s, 3H), 3.39 (s, 3H), 3.13-3.07 (m, 1H) 1.94-1.90 (m, 2H), 1.40-1.34 (m, 2H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS: 440.5 [M+H] ⁺
25		5-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-methylbenzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) 8.20 (d, J=2.4 Hz, 1H), 7.92 (d, J= 3.6 Hz, 1H), 7.66 (s, 1H), 7.42 (d, J= 8.4 Hz, 1H), 7.37 (s, 1H), 7.29 (d, J=2 Hz, 1H), 7.11 (d, J=4.8 Hz, 1H), 3.32 (s, 3H), 3.13-3.07 (m, 1H), 2.46 (s, 3H), 1.92 (d, J= 13.2 Hz, 2H), 1.37 (d, J=12.8 Hz, 2H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS: 431.5 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
26		3-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-4-methyl-4H-1,2,4-triazole	^1H NMR: ^1H NMR (400 MHz, DMSO- d_6) δ 8.41 (bs, 1H), 7.46 (d, J = 6.8 Hz, 2H), 7.29 (t, J = 14 Hz, 3H), 3.83 (s, 3H), 1.88 (d, J = 11.6 Hz, 2H), 1.38 (d, J = 12.4 Hz, 2H), 1.28 (s, 6H), 1.15 (s, 6H). LCMS 400.46 $[\text{M}+\text{H}]^+$
27		3-(1-(2-chlorobenzyl)-4-(2-methoxypyridin-2-yl)-2-methoxypyridine-2-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	^1H NMR (CDCl_3 , 400 MHz) δ 8.27 (s, 1 H), 7.7-7.84 (m, 1 H), 7.35 (d J = 7.2 Hz, 1 H), 7.18-7.28 (m, 2 H), 7.08 (s, 1 H), 6.9-7.0 (m, 1 H), 6.68 (s, 1 H), 5.13 (s, 1 H), 5.05 (s, 1 H), 3.79 (s, 3 H), 2.68-2.80 (m, 1 H), 2.0-2.10 (m, 1 H), 1.75-1.86 (m, 1 H), 1.4-1.5 (m, 3 H), 1.16 (s, 6 H), 1.03 (s, 3 H), 0.89 (s, 3 H). LCMS: 440.4 $[\text{M}+\text{H}]^+$



Cpd No.	Structure	Compound Name	Analytical data
28		3-(1-(4-chlorobenzyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.25 (d, <i>J</i> = 3.2 Hz, 1 H), 7.6-7.7 (m, 1 H), 7.33 (d, <i>J</i> = 8.4 Hz, 2 H), 7.06-7.03 (m, 1 H), 6.98 (d, <i>J</i> = 6.8 Hz, 3 H), 4.97 (s, 2 H), 3.78 (s, 3 H), 3.05-2.99 (m, 1 H), 1.86-1.82 (m, 2 H), 1.35-1.30 (m, 1 H), 1.25 (s, 7 H), 1.12 (s, 6 H). LCMS: 440.2 [M+H] ⁺
29		4-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-(trifluoromethyl)benzonitrile	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.2-8.3 (m, 1 H), 7.99-7.97 (m, 1 H), 7.78 (d, <i>J</i> = 8 Hz, 1 H), 7.62 (s, 1 H), 7.3-7.4 (m, 1 H), 7.05-7.01 (m, 2 H), 3.4 (s, 3 H), 3.26-3.20 (m, 1 H), 2.04-2.00 (m, 2 H), 1.51-1.45 (m, 2 H), 1.35 (s, 6 H), 1.26 (s, 6 H). LCMS: 485.5 [M+H] ⁺

Cpd No.	Structure	Compound Name	Analytical data
30		3-(1-(2,4-difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.21 (s, 1 H), 7.92 (d, <i>J</i> = 7.2 Hz, 1 H), 7.46-7.1 (m, 1 H), 7.3-7.44 (m, 2 H), 7.09 (m, 2 H), 3.6 (s, 3 H), 3.1-3.2 (m, 1 H), 1.92 (d, <i>J</i> = 12.8 Hz, 2 H), 1.38 (d, <i>J</i> = 13.2 Hz, 2 H), 1.29 (s, 6 H), 1.15 (s, 6 H). LCMS: 428.5 [M+H] ⁺
31		3-(1-(2,3-difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.28 (t, <i>J</i> = 3.6 Hz, 1 H), 8.01 (t, <i>J</i> = 5.6 Hz, 1 H), 7.68 (s, 1 H), 7.55-7.63 (m, 1 H), 7.22-7.27 (m, 1 H), 7.11-7.19 (m, 2 H), 3.48 (s, 3 H), 3.29-3.23 (m, 1 H), 1.98-1.93 (m, 2 H), 1.44-1.38 (m, 2 H), 1.30 (s, 6 H), 1.17 (s, 6 H). LCMS: 428.5 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
32		3-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.75 (d, J=7.2 Hz, 1H), 7.67 (s, 1H), 7.53-7.50 (m, 2H), 7.42-7.37 (m, 3H), 7.05 (t, J=7.6 Hz, 1H), 6.68 (d, J=8 Hz, 1H), 3.18 (s, 3H), 3.10 (m, 1H), 1.93 (d, J=14.4 Hz, 2H), 1.38 (d, J=12.8 Hz, 2H), 1.32 (s, 6H), 1.15 (s, 6H). LCMS: 416.5 [M+H] ⁺
33		4-(1-(3,4-difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.35 (d, J=4.8 Hz, 1H), 8.1 (s, 1H), 7.52 (d, J=4.8 Hz, 1H), 7.15-7.10 (m, 1H), 7.03-6.99 (m, 1H), 6.95 (s, 1H), 6.84 (d, J=9.6 Hz, 1H), 3.49 (s, 3H), 3.25-3.19 (m, 1H), 2.03 (d, J=10 Hz, 2H), 1.50-1.43 (m, 2H), 1.36 (s, 6H), 1.26 (s, 6H). LCMS: 428.2 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
34		2-methoxy-4-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (CDCl ₃ , 400 MHz) δ 7.60-7.59(d, <i>J</i> =1.2 Hz, 1H), 7.52-7.50 (d, <i>J</i> =8.4Hz, 1H), 7.39 - 7.35 (m, 1H), 7.07 (t, <i>J</i> =7.6Hz, 1H), 6.98 (s, 1H), 6.85(d, <i>J</i> =2.0Hz, 1H), 6.75 (d, <i>J</i> =10.4Hz, 1H), 6.60(s, 1H), 3.60 (s, 3H), 3.26 (s, 3H), 3.26-3.20 (m, 1H), 2.06-2.02 (m, 2H), 1.51-1.45 (m, 2H), 1.36 (s, 6H), 1.26 (s, 6H). LCMS: 446.5 [M+H] ⁺
35		2-(1-(3,4-difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) δ = 8.18-8.16 (m, 1H), 7.49-7.38 (m, 3H), 7.37-7.28 (m, 2H), 6.87-6.85 (m, 1H), 3.51 (s, 3H), 3.14-3.17 (m, 1H), 1.92 (dd, <i>J</i> = 2.8 Hz, <i>J</i> = 12.8 Hz, 2H), 1.39-1.33 (m, 2H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS: 428.23 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
36		3-fluoro-4-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) δ = 8.07 (d, <i>J</i> = 10.4 Hz, 1H), 7.66 (d, <i>J</i> = 9.2 Hz, 1H), 7.51 (d, <i>J</i> = 7.2 Hz, 1H), 7.39-7.35 (m, 1H), 7.34-7.27 (m, 2H), 7.06-7.00 (m, 1H), 6.86 (d, <i>J</i> = 8.4 Hz, 1H), 3.22 (s, 3H), 3.16-3.12 (m, 1H), 1.94-1.88 (m, 2H), 1.41-1.33 (m, 2H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS: 434.48 [M+H] ⁺
37		2-methoxy-5-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) δ = 7.57 (d, <i>J</i> = 2.4 Hz, 1H), 7.46 (dd, <i>J</i> = 2 Hz, <i>J</i> = 7.6 Hz, 1H), 7.41-7.36 (m, 1H), 7.34-7.31 (m, 1H), 7.26 (s, 1H), 7.18-7.16 (m, 1H), 7.04-6.98 (m, 1H), 6.90-6.88 (m, 1H), 3.88 (s, 3H), 3.28 (s, 3H), 3.11-3.05 (m, 1H), 1.91 (dd, <i>J</i> = 2.8 Hz, <i>J</i> = 12.8 Hz, 2H), 1.38-1.32 (m, 2H), 1.28 (s, 6H), 1.15 (s, 6H). LCMS: 446.30 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
38		1-(4-chlorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole	¹ H NMR (CDCl ₃ , 400 MHz) δ 7.55 (d, <i>J</i> =6.4Hz, 1H), 7.32 (t, <i>J</i> = 7.2Hz, 1H), 7.55 – 7.23 (m, 2H), 7.05 – 6.98 (m, 3H), 6.91 (s, 1H), 6.71 (d, <i>J</i> =8.4Hz, 1H), 3.23 (s, 3H), 3.26 – 3.20 (m, 1H), 2.06 – 2.03 (m, 2H), 1.51 – 1.44 (m, 2H), 1.35 (s, 6H), 1.25 (s, 6H). LCMS: 425.3 [M+H] ⁺
39		5-chloro-2-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)pyridine	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.41 (d, <i>J</i> = 2.4Hz, 1H), 7.63 – 7.60 (m, 1H), 7.50 – 7.47 (m, 1H), 7.39 – 7.33 (m, 2H), 7.05 (t, 7.6Hz, 1H), 6.81 - 6.75 (m, 2H), 3.35 (s, 3H), 3.25 - 3.19 (m, 1H), 2.06 – 2.07 (m, 2H), 1.49 – 1.46 (m, 2H), 1.35 (s, 6H), 1.25 (s, 6H). LCMS: 426.2 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
40		3-(1-(3,4-difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.19(dd, <i>J</i> = 6.4Hz, 1.6Hz, 1H), 7.90(dd, <i>J</i> = 8.8Hz, 1.6Hz, 1H), 7.45-7.36 (m, 2H), 7.31(s, 1H), 7.10-7.07(m, 1H), 6.93(d, 8Hz, 1H), 3.388(s, 3H), 3.086 (m, 1H), 1.92-1.881(m, 2H), 1.38-1.318 (m, 2H), 1.139(s, 6H), 1.127(s, 6H); LCMS: 428.3 [M+H] ⁺


Cpd No.	Structure	Compound Name	Analytical data
41		1-(4-chloro-3-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole	¹ H NMR (CDCl ₃ , 400 MHz) δ 7.56 - 7.53 (dd, <i>J</i> = 7.2Hz, 1.2Hz, 1H), 7.37-7.31 (m, 1H), 7.27 (t, <i>J</i> = 8Hz, 1H), 7.02 (t, <i>J</i> = 7.6 Hz, 1H), 6.97 - 6.94 (dd, <i>J</i> = 9.6Hz, 2.4Hz, 1H), 6.91 (s, 1H), 6.85 (d, <i>J</i> = 8.4 Hz, 1H), 6.73 (d, <i>J</i> = 8.4Hz, 1H), 3.37 (s, 3H), 3.23 - 3.22 (m, 1H), 2.06 - 2.02 (dd, <i>J</i> = 12.8Hz, 2.8 Hz, 2H), 1.47 (t, <i>J</i> = 12.8Hz, 2H), 1.35 (s, 6H), 1.25 (s, 6H). LCMS: 443.2 [M+H] ⁺
42		1-(3,4-difluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.45(dd, <i>J</i> = 1.6Hz, 6.0Hz, 1H), 7.39(m, 2H), 7.30 (m, 2H), 7.01(t, <i>J</i> = 2.8Hz, 1H), 3.07(m, 1H), 1.90 (dd, <i>J</i> = 3.2Hz, 10.0Hz, 2H), 1.34(t, <i>J</i> = 12.8Hz, 1H), 1.27 (s, 6H), 1.13 (s, 6H); LCMS: 427.3 [M+H] ⁺


Cpd No.	Structure	Compound Name	Analytical data
43		4-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.21 – 8.19 (m, 1H), 7.96 – 7.94 (m, 1H), 7.85 (d, <i>J</i> = 8.8Hz, 2H), 7.39 (s, 1H), 7.32 (d, <i>J</i> = 8.4Hz, 2H), 7.12 – 7.09 (m, 1H), 3.12 – 3.08 (m, 1H), 1.92 – 1.88 (m, 2H), 1.39 – 1.33 (m, 2H), 1.27 (s, 6H), 1.14 (s, 6H). LCMS: 417.3 [M+H] ⁺
44		3-(1-(4-chloro-3-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.20 (m, 1H), 7.91 (dd, <i>J</i> = 8Hz, 8Hz, 1H), 7.56 (t, <i>J</i> = 8Hz, 1H), 7.35 (m, 1H), 7.1(m, 1H), 6.94 (dd, <i>J</i> = 12Hz, 8Hz, 1H), 3.30 (s, 3H), 3.0 (m, 1H), 1.91 (dd, <i>J</i> = 4Hz, 12Hz, 2H), 1.36 (t, <i>J</i> = 16Hz, 2H), 1.2(s, 6H), 1.1(s, 6H), LCMS: 444.2 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
45		1-(3-chloro-4-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.53 (dd, <i>J</i> = 8Hz, 8Hz, 1H), 7.33 (m, 1H), 7.23 (t, <i>J</i> = 7.6Hz, 1H), 7.01 (m, 2H), 6.94 (m, 1H), 6.8 (s, 1H), 6.72 (d, <i>J</i> = 12Hz, 1H), 3.30 (s, 3H), 3.22 (m, 1H), 2.04 (d, <i>J</i> = 7.6Hz, 2H), 1.46 (t, <i>J</i> = 12Hz, 2H), 1.35 (s, 6H), 1.25 (s, 6H). LCMS: 443.2 [M+H] ⁺
46		3-(1-(3-chloro-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.17 (m, 1H), 7.88 (m, 1H), 7.27 (m, 1H), 7.06 (m, 1H), 6.95 (m, 2H), 6.92 (s, 2H), 3.54 (s, <i>J</i> = 12.8Hz, 3H), 3.219 (m, 1H), 2.02 (m, 2H), 1.46 (m, 2H), 1.35 (s, 6H), 1.25 (s, 6H); LCMS: 444.2 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
47		2-(1-(4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyrazine	^1H NMR (CDCl_3 , 400 MHz) δ 8.20 (d, J = 2.4 Hz, 1H), 8.07 (d, J = 4 Hz, 1H), 7.11 (t, J = 8 Hz, 2H), 7.0 (t, J = 8 Hz, 2H), 6.96 (s, 1H), 3.6 (s, 3H), 2.05 (d, J = 4 Hz, 2H), 1.52 (d, J = 16 Hz, 2H), 1.35 (s, 6H), 1.25 (s, 6H). LCMS: 411.2 $[\text{M}+\text{H}]^+$
48		3-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	^1H NMR (CDCl_3 , 400 MHz) δ 8.18 (t, J = 2 Hz, 1H), 7.94 (d, J = 7.6 Hz, 1H), 7.58 (d, J = 8 Hz, 1H), 7.44 (t, J = 10.4 Hz, 2H), 7.33 (d, J = 8 Hz, 1H), 7.01 - 6.96 (m, 2H), 3.44 (s, 3H), 3.23 (t, J = 12.4 Hz, 1H), 2.04 - 2.01 (dd, J = 12.8 Hz, 2.4 Hz, 2H), 1.48 (t, J = 12.8 Hz, 2H), 1.36 (s, 6H), 1.26 (s, 6H). LCMS: 417.3 $[\text{M}+\text{H}]^+$



Cpd No.	Structure	Compound Name	Analytical data
49		5-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-4-methoxypyrimidine	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.75 (s, 1H), 8.71 (s, 1H), 7.32 (d, J = 8Hz, 2H), 7.06 (d, J = 8.80Hz, 2H), 6.97 (s, 1H), 3.6 (s, 3H), 3.20 - 3.26 (m, 1H), 2.0 - 2.04 (dd, J = 12.8Hz, 3.2Hz, 2H), 1.48 (t, J = 12.8Hz, 2H), 1.36 (s, 6H), 1.26 (s, 6H). LCMS: 427.4 [M+H] ⁺
50		2-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyrazine	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.21(d, J = 2.8Hz, 1H), 8.09(d, J = 2Hz, 1H), 7.28(t, J = 8.4Hz, 2H), 7.06(d, J = 8.4Hz, 2H), 6.96 (s, 1H), 3.62 (s, 3H), 3.28 (m, 1H), 2.05(dd, J = 12.8Hz, 12.8Hz, 2H), 1.49 (t, J = 13.2Hz, 2H), 1.35(s, 6H), 1.25(s, 6H). LCMS: 427.2 [M+H] ⁺


Cpd No.	Structure	Compound Name	Analytical data
51		3-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-ethoxypyridine	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.15 – 8.14 (m, 1H), 7.92 – 7.91 (m, 1H), 7.29 – 7.27 (m, 2H), 7.05 (d, <i>J</i> = 8.4Hz, 2H), 6.96 – 6.93 (m, 2H), 3.97 (dd, <i>J</i> = 14, 6.8Hz, 2H), 3.29 – 3.21 (m, 1H), 2.04 – 2.01 (m, 2H), 1.51 – 1.44 (m, 2H), 1.36 (s, 6H), 1.26 (s, 6H), 0.94 (t, <i>J</i> = 6.8Hz, 3H). LCMS: 440.3 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
52		3-(1-(3,4-difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-ethoxypyridine	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.17 (t, <i>J</i> = 3.2 Hz, 1H), 7.94 - 7.92 (dd, <i>J</i> = 6.8 Hz, 1.2 Hz, 1H), 7.46 - 7.35 (m, 2H), 7.32 (s, 1H), 7.09 - 7.06 (m, 1H), 6.94 (d, <i>J</i> = 8.4 Hz, 1H), 3.90 (q, <i>J</i> = 7.2 Hz, 2H), 3.08 (t, <i>J</i> = 12.4 Hz, 1H), 1.92 - 1.89 (dd, <i>J</i> = 12.8 Hz, 2 Hz, 2H), 1.35 (t, <i>J</i> = 12.8 Hz, 2H), 1.27 (s, 6H), 1.14 (s, 6H), 0.84 (t, <i>J</i> = 7.2 Hz, 3H). LCMS: 442.3 [M+H] ⁺


Cpd No.	Structure	Compound Name	Analytical data
53		5-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-(trifluoromethyl)pyridine	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.592(s, 1H), 7.61(m, 2H), 7.55(m, 1H) 7.393-7.354(m, 1H), 7.260(s, 1H), 7.07-7.04(m, 1H), 6.989(s, 1H), 6.7195(d, 8.4Hz, 1H), 3.28 (s, 3H), 3.23(m, 1H) 2.05 (m, 2H), 1.51(m, 2H) 1.36 (s, 6H), 1.26(s, 6H); LCMS: 460.3 [M+H] ⁺
54		2-methoxy-3-(4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1-(6-(trifluoromethyl)pyridin-3-yl)-1H-imidazol-2-yl)pyridine	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.595(d, 2.4Hz, 1H), 8.20(dd <i>J</i> = 6.8Hz, 1.6Hz, 1H), 7.99 (dd, <i>J</i> = 9.2Hz, 2Hz 1H), 7.66(m, 1H), 7.55(m, 1H), 7.01(m, 2H), 3.435(s, 3H), 3.24 (m, 1H), 2.03(m, 2H), 1.5(m, 2H), 1.369(s, 6H), 1.270(s, 6H); LCMS: 461.2 [M+H] ⁺


Cpd No.	Structure	Compound Name	Analytical data
55		1-(4-chlorophenyl)-2-(cyclopropylmethyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole	^1H NMR (400 MHz, DMSO- d_6) 7.75 (d, J = 8.4Hz, 2H), 7.42 (d, J = 8.4Hz, 2H), 6.93 (s, 1H), 2.99 (s, 1H), 2.51 (s, 2H), 1.86 - 1.82 (dd, J = 12.8Hz, 2.8Hz, 2H), 1.31 - 1.25 (m, 8H), 1.11 (s, 6H), 0.85 (m, 1H), 0.32 (d, J = 6.4Hz, 2H), - 0.047 (d, J = 4.4Hz, 2H). LCMS: 373.3 $[\text{M}+\text{H}]^+$
56		1-(4-chlorophenyl)-2-cyclopropyl-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole	^1H NMR (CDCl_3 , 400 MHz) δ 7.44(d, J = 8.8Hz, 2H), 7.34(d, J = 8.8Hz, 2H), 6.67(s, 1H), 5.29 (s, 1H), 3.0 (m, 1H), 1.93(dd, J = 12.8Hz, 13.2Hz, 2H), 1.34(m, 1H), 1.70(m, 1H), 1.25(m, 1H), 1.33(s, 6H), 1.22(s, 6H), 1.05(m, 2H), 0.87(m, 2H). LCMS: 359.3 $[\text{M}+\text{H}]^+$



Cpd No.	Structure	Compound Name	Analytical data
57		4-(2-cyclopropyl-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.95 (d, J=8 Hz, 2H), 7.71 (d, J=7.6 Hz, 2H), 7.08 (s, 1H), 2.94 (t, J=12.8 Hz, 1H), 1.80 (t, J=7.2 Hz, 3H), 1.27-1.21 (s, 8H), 1.10 (s, 6H), 0.88 - 0.85(m, 4H). LCMS: 350.3 [M+H] ⁺
58		4-(2-(cyclopropylmethyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) δ =7.99-7.99(2H,d,J=0.06Hz) , 7.64-7.62 (2H,d,J=1.44Hz), 7.08(1H,s), 3.0(3H,s), 2.59-2.57 (2H,d,J=0.96Hz), 1.87-1.83 (2H,dd,J=2.4Hz),1.31-1.25(2H,d,J=3.48Hz),1.25(6H,s),1.12(6H,s). LCMS: 364 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
59		5-(2-(cyclopropylmethyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-(trifluoromethyl)pyridine	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.87 (s, 1H), 8.19 (d, <i>J</i> = 8.4Hz, 1H), 8.06 (d, <i>J</i> = 8.4Hz, 1H), 7.15 (s, 1H), 2.61 (t, <i>J</i> = 6.4Hz, 1H), 2.06 (d, <i>J</i> = 6.4Hz, 2H), 1.88 – 1.84 (dd, <i>J</i> = 12.8Hz, 2.8Hz, 2H), 1.31 (d, <i>J</i> = 12.8Hz, 2H), 1.26 (s, 6H), 1.26 (s, 6H), 0.89 (m, 1H), 0.35 – 0.32 (dd, <i>J</i> = 7.6Hz, 4.4Hz, 2H), -0.04 - -0.04 (dd, <i>J</i> = 10Hz, 5.2Hz, 2H). LCMS: 408.5 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
60		2-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-ethoxypyrazine	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.23 (d, <i>J</i> = 2.4Hz, 1H), 8.08 (d, <i>J</i> = 2.8Hz, 1H), 7.29 (d, <i>J</i> = 8.8Hz, 2H), 7.06 (d, <i>J</i> = 8.8Hz, 2H), 6.96 (s, 1H), 4.08 - 4.02 (q, <i>J</i> = 7.2Hz, 2H), 3.31 - 3.24 (m, 1H), 2.07 - 2.03 (dd, <i>J</i> = 13.2Hz, 3.2Hz, 2H), 1.50 (t, <i>J</i> = 13.6Hz, 2H), 1.35 (s, 6H), 1.25 (s, 6H), 1.02 (t, <i>J</i> = 6.8Hz, 3H). LCMS: 441.3 [M+H] ⁺
61		1-(4-(difluoromethoxy)phenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.45 - 7.43 (dd, <i>J</i> = 7.6 Hz, 1.6 Hz, 1H), 7.38 - 7.33 (m, 1H), 7.20 (s, 1H), 7.13 (s, 1H), 7.00 (t, <i>J</i> = 5.2 Hz, 1H), 6.86 (d, <i>J</i> = 8.4 Hz, 1H), 3.21 (s, 3H) 3.08 (t, <i>J</i> = 12.8 Hz, 1H), 1.92 - 1.88 (dd, <i>J</i> = 13.2 Hz, 3.2 Hz, 2H), 1.35 (t, <i>J</i> = 12.8Hz, 2H), 1.27 (s, 6h), 1.14 (s, 6H). LCMS: 457.6 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
62		1-(4-bromo-3-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole	¹ H NMR (CDCl ₃ , 400 MHz) δ 7.56 – 7.53 (dd, <i>J</i> = 7.6Hz, 2Hz, 1H), 7.44 (t, <i>J</i> = 8.0Hz, 1H), 7.37 – 7.33 (m, 1H), 7.02 (t, <i>J</i> = 7.6Hz, 1H), 6.94 – 6.91 (m, 2H), 6.08 – 6.78 (m, 1H), 6.74 (d, <i>J</i> = 8.4Hz, 1H), 3.36 (s, 3H), 3.22 (t, <i>J</i> = 12.8Hz, 1H), 2.02 (d, <i>J</i> = 2.8Hz, 2H), 1.47 (t, <i>J</i> = 12.8Hz, 2H), 1.35 (s, 6H), 1.25 (s, 6H). LCMS: 487.2 [M+H] ⁺


Cpd No.	Structure	Compound Name	Analytical data
63		1-(3-bromo-4-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.49 - 7.51 (dd, <i>J</i> = 5.6Hz, 2.4Hz, 1H), 7.45 (d, <i>J</i> = 7.6Hz, 1H), 7.37 (t, <i>J</i> = 6.8Hz, 1H), 7.32 (t, <i>J</i> = 8.8Hz, 1H), 7.27 (s, 1H), 7.06-7.10 (m, 1H), 7.01 (t, <i>J</i> = 7.2Hz, 1H), 6.88 (d, <i>J</i> = 8.4 Hz, 1H), 3.26 (s, 3H), 3.04-3.10 (m, 1H), 1.88-1.92 (dd, <i>J</i> = 13.2Hz, 3.2Hz, 2H), 1.34 (t, <i>J</i> = 12.8Hz, 2H), 1.27 (s, 6H), 1.14 (s, 6H). LCMS: 487.2 [M+H] ⁺


Cpd No.	Structure	Compound Name	Analytical data
64		3-(1-(3-bromo-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.18 - 8.16 (dd, <i>J</i> = 4.8Hz, 2Hz, 1H), 7.89 - 7.87 (dd, <i>J</i> = 7.2Hz, 1.6Hz, 1H), 7.44 - 7.42 (dd, <i>J</i> = 6.4Hz, 2.8Hz, 1H), 7.06 (m, 3H), 6.92 (s, 1H), 3.53 (s, 3H), 3.25 - 3.18 (m, 1H), 2.04 - 2.00 (dd, <i>J</i> = 12.8Hz, 3.2Hz, 2H), 1.47 (t, <i>J</i> = 12.8Hz, 2H), 1.35 (s, 6H), 1.25 (s, 6H). LCMS: 488.2 [M+H] ⁺
65		3-(1-(4-bromo-3-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.19 - 8.17 (dd, <i>J</i> = 7.2Hz, 2Hz, 1H), 7.90 - 7.88 (dd, <i>J</i> = 7.6Hz, 1.6Hz, 1H), 7.51 - 7.46 (m, 1H), 6.99 - 6.93 (m, 3H), 6.79 (d, <i>J</i> = 8.4 Hz, 1H), 3.52 (s, 3H), 3.28 - 3.22 (m, 1H), 2.04 - 2.00 (dd, <i>J</i> = 12.8Hz, 2.8Hz, 2H), 1.54 - 1.44 (m, 2H), 1.35 (s, 6H), 1.26 (s, 6H). LCMS: 488.2 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
66		4-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.30 (d, <i>J</i> = 4.8 Hz, 1H), 8.28 (s, 1H), 7.51 (d, <i>J</i> = 4.8 Hz, 1H), 7.44 (d, <i>J</i> = 8.44 Hz, 2H), 7.36 (s, 1H), 7.16 (d, <i>J</i> = 8.8 Hz, 2H), 3.33 (s, 3H), 3.11 (t, <i>J</i> = 13.2 Hz, 1H), 1.93-1.89 (m, 2H), 1.36 (t, <i>J</i> = 12.4 Hz, 2H), 1.28 (s, 6H), 1.15 (s, 6H). LCMS 426.36 [M+H] ⁺
67		2-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.16 (d, <i>J</i> = 3.2 Hz, 1H), 7.45-7.37 (m, 4H), 7.30 (s, 1H), 7.09 (d, <i>J</i> = 8.4 Hz, 2H), 3.47 (s, 3H), 1.93-1.89 (m, 2H), 1.40-1.33 (m, 2H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS 426.36 [M+H] ⁺



Cpd No.	Structure	Compound Name	Analytical data
68		3-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-4-methoxypyridine	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.50 (s, 1H), 8.66 (d, <i>J</i> = 5.6 Hz, 1H), 7.43 (d, <i>J</i> = 8.8 Hz, 2H), 7.33 (s, 1H), 7.17 (d, <i>J</i> = 6.0 Hz, 1H), 3.36 (s, 3H), 3.14-3.08 (m, 1H), 1.94-1.89 (m, 2H), 1.4-1.33 (m, 2H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS 426.36 [M+H] ⁺
69		3-(1-(4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-4-methoxypyridine	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.49 (s, 1H), 8.53 (d, <i>J</i> = 5.6 Hz, 1H), 7.3 (s, 1H), 7.20 (d, <i>J</i> = 6.0 Hz, 4H), 6.95 (d, <i>J</i> = 5.2 Hz, 1H), 3.37 (s, 3H), 3.11-3.08 (m, 1H), 1.93-1.85 (m, 2H), 1.40-1.37 (m, 2H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS 410.23 [M+H] ⁺


Cpd No.	Structure	Compound Name	Analytical data
70		2-(1-(4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.12-8.16 (m, 1H), 7.44-7.38 (m, 1H), 7.26 (s, 1H), 7.19-7.09 (m, 4H), 3.47 (s, 3H), 3.14-3.07 (m, 1H), 1.94-1.91 (m, 2H), 1.40-1.37 (m, 2H), 1.34 (s, 6H), 1.15 (s, 6H). LCMS 410.47 [M+H] ⁺
71		3-(1-(2,5-difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.18 (d, <i>J</i> = 3.2 Hz, 1H), 7.91 (d, <i>J</i> = 5.6 Hz, 1H), 7.43-7.40 (m, 1H), 7.29 (br s, 2H), 7.26-7.22 (m, 1H), 7.10-7.07 (m, 1H), 3.40 (s, 3H), 3.12 (m, 1H), 1.91 (d, <i>J</i> = 12.8, 1H), 1.36 (dd, <i>J</i> = 12.8 & 12.8 Hz, 1H), 1.29 (s, 6H), 1.15 (s, 6H). LCMS 428.62 [M+H] ⁺

Cpd No.	Structure	Compound Name	Analytical data
72		3-(1-(3,5-difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.20 (dd, <i>J</i> = 1.6 & 5.2 Hz, 1H), 7.91 (dd, <i>J</i> = 1.6 & 5.6 Hz, 1H), 6.99 (dd, <i>J</i> = 5.2 & 7.2 Hz, 1H), 6.95 (s, 1H), 6.75 (dd, <i>J</i> = 8.4 & 8.4 Hz, 1H), 6.68 (d, <i>J</i> = 5.6 Hz, 2H), 3.50 (s, 3H), 3.21 (m, 1H), 2.02-2.00 (m, 2H), 1.50-1.43 (m, 2H), 1.35 (s, 6H), 1.20 (s, 6H). LCMS 428.51 [M+H] ⁺

Cpd No.	Structure	Compound Name	Analytical data
73		3-fluoro-5-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.22 (dd, <i>J</i> = 1.6 & 5.2 Hz, 1H), 7.96 (dd, <i>J</i> = 1.6 & 5.6 Hz, 1H), 7.31 (d, <i>J</i> = 7.2 Hz, 1H), 7.23 (d, <i>J</i> = 4.4 Hz, 1H), 7.12 (d, <i>J</i> = 9.2 Hz, 1H), 7.02 (dd, <i>J</i> = 5.2 & 7.2 Hz, 1H), 6.96 (s, 1H), 3.50 (s, 3H), 3.22 (m, 1H), 2.02 (dd, <i>J</i> = 2.8 & 13.2 Hz, 2H), 1.50-1.43 (m, 2H), 1.36 (s, 6H), 1.20 (s, 6H). LCMS 435.36 [M+H] ⁺

Cpd No.	Structure	Compound Name	Analytical data
75		3-(1-(6-ethoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.18 - 8.19(m, 1H), 7.94 (d, <i>J</i> = 2.8 Hz, 1H), 7.88 - 7.91 (m, 1H), 7.51 - 7.54 (m, 1H), 7.27 (s, 1H), 7.07 - 7.10 (m, 1H), 6.79 (d, <i>J</i> = 9.2 Hz, 1H), 4.23 - 4.28 (q, 2H), 3.42 (s, 3H), 3.10 (t, <i>J</i> = 13.2 Hz, 1H), 1.89 - 1.93 (m, 2H), 1.36 - 1.39 (m, 2H), 1.28 - 1.33 (m, 9H), 1.15 (m, 6H) LCMS 437.5[M+H] ⁺
121		4-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-(trifluoromethyl)benzonitrile	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.13 (d, <i>J</i> = 8.4Hz, 1H), 7.69 (s, 1H), 7.53-7.58 (m, 3H), 7.42 (t, <i>J</i> = 7.6 Hz, 1H), 7.07 (t, <i>J</i> = 7.6Hz, 1H), 6.88 (d, <i>J</i> = 8.4Hz, 1H), 3.16 (s, 3H), 3.09 - 3.13 (m, 1H), 1.90 - 1.94 (m, 2H), 1.32 - 1.39 (m, 2H), 1.28 (s, 6H), 1.14 (s, 6H). LCMS 453.2 [M+H] ⁺

Cpd No.	Structure	Compound Name	Analytical data
123		2-chloro-4-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (DMSO-d ₆ , 400 MHz) δ 7.59 (d, <i>J</i> = 7.6 Hz, 1H), 7.55 (d, <i>J</i> = 7.6 Hz, 1H), 7.39 (t, <i>J</i> = 6.4 Hz, 1H), 7.34 (d, <i>J</i> = 2.0 Hz, 1H), 7.08-7.05 (m, 2H), 6.96 (s, 1H), 6.75 (d, <i>J</i> = 8.4 Hz, 1H), 3.34 (s, 3H), 3.22 (m, 1H), 2.03 (dd, <i>J</i> = 2.4 & 12.8 Hz, 2H), 1.50-1.40 (m, 2H), 1.35 (s, 6H), 1.26 (s, 6H). LCMS 450.50 [M+H] ⁺
124*		1-(4-chlorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyl-3,6-dihydro-2H-pyran-4-yl)-1H-imidazole	¹ H NMR (400 MHz, CDCl ₃) δ 7.62 – 7.60 (dd, <i>J</i> = 7.6, 1.6 Hz, 1H), 7.36 – 7.32 (m, 1H), 7.26 – 7.24 (m, 2H), 7.10 (s, 1H), 7.06 – 7.00 (m, 3H), 6.71 (d, <i>J</i> = 8.4 Hz, 1H), 6.53 (s, 1H), 3.33 (s, 3H), 2.31 (s, 2H), 1.33 (d, <i>J</i> = 6.0 Hz, 12H). LCMS 423.2 [M+H] ⁺


Cpd No.	Structure	Compound Name	Analytical data
125*		1-(4-bromo-3-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyl-3,6-dihydro-2H-pyran-4-yl)-1H-imidazole	¹ H NMR:(400 MHz, DMSO-d ₆) δ 7.66 (t, <i>J</i> = 8.4Hz, 1H), 7.58 (s, 1H), 7.51-7.49 (m, 1H), 7.43 - 7.39 (m, 1H), 7.26 - 7.23 (dd, <i>J</i> = 10Hz, 2Hz, 1H), 7.04 (t, <i>J</i> = 7.6Hz, 1H), 6.93 - 6.87 (m, 2H), 6.33 (s, 1H), 3.28 (d, <i>J</i> = 10 Hz, 3H), 2.24 (s, 2H), 1.22 (d, <i>J</i> = 6.8Hz, 12H) LCMS 485.2 [M+H] ⁺

*Compounds **124** and **125** were side products obtained while synthesizing compound **38** and **62** respectively

5

Example 8

Compound 74: 2-Fluoro-5-[2-(2-methoxy-phenyl)-4-(2,2,6,6-tetramethyl-tetrahydro-pyran-4-yl)-imidazol-1-yl]-benzonitrile







To a solution of (3-bromo-4-fluoro-phenyl)-2-(2-methoxy-phenyl)-4-(2,2,6,6-tetramethyl-tetrahydro-pyran-4-yl)-1H-imidazole **63** (0.17 g, 0.00034 mol) in DMF (10 mL) was added Zn(CN)₂ (0.08 g, 0.00068 mol) under N₂. The reaction mixture was further purged with N₂



for 30 min and then stirred for 1 h at 160 °C. The reaction was then quenched with ice cold water and extracted with ethyl acetate (50 mL x 3). The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica
5 gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound **74** as a pale yellow solid. Yield: 0.11 g (72.8%). ¹H NMR (CDCl₃, 400 MHz) δ 7.54-7.57 (dd, *J* = 5.2Hz, 2.8Hz, 1H), 7.41-7.43 (dd, *J* = 5.2Hz, 2.8Hz, 1H), 7.31 - 7.38 (m, 2H), 7.13 (t, *J* = 8.4Hz, 1H), 7.04 (t, *J* = 7.6Hz, 1H), 6.90 (s, 1H), 6.73 (d, *J* = 8.4 Hz, 1H), 3.38 (s, 3H), 3.23 (m, 1H), 2.01-2.05 (dd, *J* = 12.8Hz, 3.2Hz, 2H), 1.43 - 1.50 (m, 2H), 1.36 (s, 6H),
10 1.26 (s, 6H). LCMS: 487.2 [M+H]⁺.



Using analogous protocols to those described in Example 8, the compounds described in Table 3 have been prepared using and appropriately substituted bromide intermediate.



Table 3



Cpd No.	Structure	Compound name	Analytical data
76		3-fluoro-4-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.16 (dd, <i>J</i> = 4.8Hz, 4.8Hz, 2H), 7.96(dd, <i>J</i> = 1.6Hz, 7.6Hz, 1H), 7.49(d, <i>J</i> = 4Hz, 1H), 7.38(d, <i>J</i> = 8Hz, 1H), 7.17 (t, <i>J</i> = 8Hz, 1H), 6.9 (m, 2H), 3.4 (s, 3H), 3.2 (m, 1H), 2.02(dd, <i>J</i> = 12Hz, 9.6Hz, 2H), 1.48 (t, <i>J</i> = 12Hz, 2H), 1.36(s, 6H), 1.26(s, 6H). LCMS: 435.3 [M+H] ⁺
77		2-methoxy-4-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.2(d, <i>J</i> = 1.6Hz, 1H), 7.92(d, <i>J</i> = 6.8Hz, 1H), 7.52(d, <i>J</i> = 8.4Hz, 1H), 6.99 (s, 2H), 6.81(d, <i>J</i> = 1.6Hz, 1H), 6.65 (s, 1H), 3.70 (m, 2H), 3.4 (s, 3H), 3.23(m, 1H), 2.02(d, <i>J</i> = 10.4Hz, 2H), 1.48 (t, <i>J</i> = 12.8Hz, 2H), 1.36(s, 6H), 1.26(s, 6H). LCMS: 447.3 [M+H] ⁺



Cpd No.	Structure	Compound name	Analytical data
78		2-chloro-4-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) 8.23 – 8.22 (dd, <i>J</i> = 4.8Hz, 1.6Hz, 1H), 7.98 – 7.94 (m, <i>J</i> = 2.6Hz, 1H), 7.48 (s, 1H), 7.24 – 7.12 (dd, <i>J</i> = 8.4Hz, 2.4Hz, 1H), 7.14 – 7.11 (dd, <i>J</i> = 6.8Hz, 4.8Hz, 1H), 3.34 (s, 3H) 3.08 (m, 1H), 1.93 – 1.89 (dd, <i>J</i> = 12.8Hz, 2.8Hz, 2H), 1.35 (t, <i>J</i> = 12.8Hz, 2H), 1.27 (s, 6H). LCMS: 451.2 [M+H] ⁺
79		2-fluoro-5-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.19 (d, <i>J</i> = 4.8Hz, 1H), 7.91 (t, <i>J</i> = 6Hz, 1H), 7.45- 7.43 (m, 1H), 7.36 - 7.32 (m, 1H), 7.18 (t, <i>J</i> = 8.4Hz, 1H), 7.01 - 6.98 (m, 1H), 6.92 (s, 1H), 3.52 (s, 3H), 3.22 (bs, 1H), 2.03 - 1.99 (m, 2H), 1.47 (t, <i>J</i> = 12.8Hz, 3H), 1.36 (s, 6H), 1.26 (s, 6H). LCMS: 435.3 [M+H] ⁺

Cpd No.	Structure	Compound name	Analytical data
80		2-fluoro-4-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.22 - 8.21 (dd, <i>J</i> = 4.8Hz, 1.6Hz, 1H), 7.96 - 7.94 (dd, <i>J</i> = 7.6Hz, 2Hz, 1H), 7.58 (t, <i>J</i> = 8Hz, 1H), 7.04 - 6.97 (m, 4H), 3.49 (s, 3H), 3.25 - 3.18 (m, 1H), 2.03 - 1.99 (dd, <i>J</i> = 13.2Hz, 3.2Hz, 2H), 1.47 (t, <i>J</i> = 12.8Hz, 2H), 1.36 (s, 6H), 1.26 (s, 7H). LCMS: 435 [M+H] ⁺
81		2-fluoro-4-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) 7.06 - 7.58 (dd, <i>J</i> = 7.2Hz, 1.6Hz, 1H), 7.53 (t, <i>J</i> = 8Hz, 1H), 7.41 - 7.37 (m, 1H), 7.067 (t, <i>J</i> = 7.6Hz, 1H), 7.00 (d, <i>J</i> = 8.4Hz, 2H), 6.95 (s, 1H), 6.75 (d, <i>J</i> = 8.4Hz, 1H), 3.34 (s, 3H), 3.26 - 3.19 (m, 1H), 2.05 - 2.01 (dd, <i>J</i> = 13.6Hz, 3.2Hz, 2H), 1.47 (t, <i>J</i> = 12.8Hz, 2H), 1.35 (s, 6H), 1.25 (s, 6H). LCMS: 434.3 [M+H] ⁺

Cpd No.	Structure	Compound name	Analytical data
82		2,6-difluoro-4-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.25 (d, <i>J</i> = 3.6Hz, 1H), 7.96 (d, <i>J</i> = 5.6Hz, 1H), 7.05 – 7.03(m, 1H), 6.96 (s, 1H), 6.84 (d, <i>J</i> = 8.0Hz, 2H), 3.56 (s, 3H), 3.24 – 3.18 (m, 1H), 2.02 – 1.981 (m, 2H), 1.46 (t, <i>J</i> = 12.8Hz, 2H), 1.35 (s, 6H), 1.26 (s, 6H). LCMS: 453.3 [M+H] ⁺
83		3,5-difluoro-4-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) δ ppm 7.93(d, <i>J</i> = 7.6 Hz, 2H), 7.45 (d, <i>J</i> = 7.2 Hz, 1H), 7.35 (t, <i>J</i> = 7.2 Hz, 1H), 7.26 (s, 1H), 7.00 (t, <i>J</i> = 7.6 Hz, 1H), 6.86 (d, <i>J</i> = 8.4 Hz, 1H), 3.31 (s, 3H), 1.91 (d, <i>J</i> = 12.8 Hz, 2H), 1.36 (d, <i>J</i> = 12.8 Hz, 2H), 1.28 (s, 6H), 1.14 (s, 6H); LCMS: 452.3 [M+H] ⁺


Cpd No.	Structure	Compound name	Analytical data
84		2-chloro-5-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.21(dd, <i>J</i> = 1.2Hz, 5.2Hz, 1H), 7.94(dd, <i>J</i> = 8.8Hz, 9.6Hz, 2H), 7.71(d, <i>J</i> = 8.8Hz, 1H), 7.41(m, 2H), 7.12(m, 1H), 3.35(s, 3H), 3.12(m, 1H), 1.89(d, <i>J</i> = 2.8Hz, 2H), 1.34(m, 2H), 1.27(s, 6H), 1.14(s, 6H). LCMS: 451.2 [M+H] ⁺
85		2-methoxy-5-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.19 - 8.17 (dd, <i>J</i> = 4.8 Hz, <i>J</i> ₂ = 1.6 Hz, 1H), 7.93 - 7.88 (m, 1H), 7.63 (d, <i>J</i> = 2.4 Hz, 1H), 7.37 - 7.34 (dd, <i>J</i> = 8.8 Hz, 2.8 Hz, 1H), 7.29 (s, 1H), 7.18 (d, <i>J</i> = 9.2 Hz, 1H), 7.09 - 7.06 (m, 1H), 3.88 (s, 3H), 3.39 (s, 3H), 3.08 (t, <i>J</i> = 12.8 Hz, 1H), 1.92 - 1.88 (dd, <i>J</i> = 12.8 Hz, 2.8 Hz, 2H), 1.34 (t, <i>J</i> = 12.8 Hz, 2H), 1.27 (s, 6H), 1.14 (s, 6H). LCMS: 447.3 [M+H] ⁺

Cpd No.	Structure	Compound name	Analytical data
86		4-(2-(4-methoxypyrimidin-5-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	^1H NMR (400 MHz, DMSO- d_6) δ 8.81 (s, 1H), 8.74 (s, 1H), 7.87 (d, J = 8.4 Hz, 2H), 7.48 (s, 1H), 7.41 (d, J = 8.4 Hz, 2H), 3.40 (s, 3H), 3.09 - 3.15 (m, 1H), 1.89-1.93 (dd, J = 12.8 Hz, 2.4 Hz, 2H), 1.36 (t, J = 12.8 Hz, 2H), 1.27 (s, 6H), 1.14 (s, 6H). LCMS: 418.3 $[\text{M}+\text{H}]^+$
87		4-(2-(3-methoxypyrazin-2-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	^1H NMR (400 MHz, DMSO- d_6) δ 8.27 - 8.22 (dd, J = 18 Hz, 2.8 Hz, 2H), 7.84 (d, J = 8.4 Hz, 2H), 7.48 (s, 1H), 7.30 (d, J = 8.8 Hz, 2H), 3.59 (s, 3H), 3.13 (t, J = 12.4 Hz, 1H), 1.93 - 1.89 (dd, J = 13.6 Hz, 3.2 Hz, 2H), 1.40 - 1.31 (m, 2H), 1.28 (s, 6H), 1.23 (t, J = 4 Hz, 1H), 1.14 (s, 6H). LCMS: 418.3 $[\text{M}+\text{H}]^+$

Cpd No.	Structure	Compound name	Analytical data
88		2-fluoro-4-(2-(3-methoxypyrazin-2-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.20 (d, <i>J</i> = 2.4 Hz, 1H), 8.13 (d, <i>J</i> = 2.0 Hz, 1H), 7.59 (t, <i>J</i> = 7.2 Hz, 1H), 7.03 (q, <i>J</i> = 10 Hz, 3H), 3.72 (s, 3H), 3.28 (t, <i>J</i> = 12.8 Hz, 1H), 2.05 - 2.02 (m, 2H), 1.54 - 1.45 (m, 2H), 1.35 (s, 6H), 1.25 (s, 6H). LCMS: 436.2 [M+H] ⁺
89		4-(2-(2-ethoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-fluorobenzonitrile	¹ H NMR (CDCl ₃ , 400 MHz) δ 8.18 (d, <i>J</i> = 4.8 Hz, 1H), 7.94 (d, <i>J</i> = 6.8 Hz, 1H), 7.45 (d, <i>J</i> = 3.2 Hz, 1H), 7.36 - 7.33 (m, 1H), 7.17 (t, <i>J</i> = 8.4 Hz, 1H), 6.99 (t, <i>J</i> = 5.2 Hz, 1H), 6.93 (s, 1H), 4.01 (q, <i>J</i> = 6.8 Hz, 2H), 3.22 (t, <i>J</i> = 12.4 Hz, 1H), 2.02 (d, <i>J</i> = 13.2 Hz, 2H), 1.51 - 1.42 (m, 2H), 1.36 (s, 6H), 1.26 (s, 6H), 0.98 (t, <i>J</i> = 6.8 Hz, 3H). LCMS: 449.3 [M+H] ⁺

Example 9

Compound 90: 1-(4-chlorophenyl)-2-(2-methoxyphenyl)-5-methyl-4-(2,2,6,6-tetramethyl tetrahydro-2H-pyran-4-yl)-1H-imidazole




5

To a stirred solution of N-(4-chlorophenyl)-2-methoxybenzimidamide **A 38**, (0.2 g, 0.07 mmol) and 2-bromo-1-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)propan-1-one **A72** (0.3 g, 0.108 mmol) in dioxane (10 mL) was added NaHCO₃ (0.13 g, 0.15 mmol). The reaction mixture was refluxed at 100 °C for 12 h. After completion of the reaction it was
 10 filtered through a pad of diatomaceous earth. The resultant filtrate was concentrated under reduced pressure to provide a crude mixture. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as an off-white solid. Yield: 0.055 g, 17%. ¹H NMR (400 MHz, DMSO-d₆) 7.45 (d, *J* = 7.2Hz, 1H), 7.28 – 7.23 (m, 3H), 7.00 (d, *J* = 8.4Hz, 2H), 6.93 (t, *J* = 8.0Hz, 1H), 6.64 (d, *J* =
 15 7.6Hz, 1H), 3.37 (s, 3H), 3.16 – 3.18 (m, 1H), 2.11 (s, 3H), 1.87 (t, *J* = 12.8Hz, 2H), 1.75 – 1.72 (m, 2H), 1.36 (s, 6H), 1.26 (s, 6H); LCMS: 439.3 [M+H]⁺.

Example 10

Compound 91: 4-(5-chloro-2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile


20



To a solution of 4-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile, Cpd **43**, (0.055 g, 0.012 mmol) in acetonitrile (3 mL) was added NCS (0.017 g, 0.013 mmol) at 0 °C. The reaction mixture was then heated at 80 °C for 5 h. After completion of the reaction, it was quenched with water and extracted with DCM (2 x 50 mL). The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as a white solid. Yield: 0.02 g, 34%. ¹H NMR (CDCl₃, 400 MHz) δ 8.16 – 8.14 (m, 1H), 7.89 – 7.86 (m, 1H), 7.67 (d, *J* = 8.4 Hz, 2H), 7.27 – 7.25 (m, 1H), 6.97 – 6.94 (m, 1H), 3.45 (s, 3H), 3.34 – 3.26 (m, 1H), 1.82 – 1.76 (m, 4H), 1.37 (s, 6H), 1.27 (s, 6H); LCMS: 451.2 [M+H]⁺.

Example 11

Compound 92: 3-(5-chloro-1-(4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine




Compound **92** was prepared using the procedure described Example 10 for the synthesis of Compound **91**. ¹H NMR (DMSO, 400 MHz) 8.17 (dd, *J* = 1.6 & 4.8 Hz, 1H), 7.88 (dd, *J* =

1.6 & 3.6 Hz, 1H), 7.28-7.24 (m, 4H), 7.07-7.04 (m, 1H), 3.46 (s, 3H), 3.18 (m, 1H), 1.75-1.70 (m, 2H), 1.64-1.57 (m, 2H), 1.30 (s, 6H), 1.16 (s, 6H); LCMS 444.46 [M+H]⁺.

Example 12

5 **Compound 93: 3-(5-chloro-1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine**




Compound **93** was prepared using the procedure described Example 10 for the synthesis
 10 of Compound **91**. ¹H NMR (DMSO, 400 MHz): 8.12 (d, *J* = 4.8 Hz, 1H), 7.81 (dd, *J* = 1.6 & 5.2 Hz, 1H), 7.33 (d, *J* = 8.4 Hz, 2H), 7.07 (d, *J* = 8.4 Hz, 2H), 6.93 (dd, *J* = 5.2 & 7.2 Hz, 1H), 3.51 (s, 3H), 3.32-3.20 (m, 1H), 1.90-1.80 (m, 4H), 1.37 (s, 6H), 1.26 (s, 6H); LCMS 460.44 [M+H]⁺

15

Example 13

Compound 94: 2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1((trifluoromethyl)sulfonyl)-1H-imidazole




To a stirred solution of 2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole (0.2 g, 0.0006 mol) in DCM (10 mL) at 0 °C was added triethylamine (0.129 g, 0.0012 mol) dropwise over a period of 1 min. The reaction mixture was stirred at 0 °C followed by the addition of trifluoromethanesulfonic anhydride (0.36 g, 0.0012 mol) and stirred to room temperature for 12 h. After completion, the reaction was quenched with ice cold water, then extracted with ethyl acetate (3 x 25 mL). The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as a white solid. Yield: 37 mg, 24 %. ¹H NMR (400 MHz, DMSO-d₆) 7.67 (s, 1H), 7.50 - 7.55 (m, 1H), 7.31 (d, *J* = 7.6 Hz, 1H), 7.12 (d, *J* = 8.4 Hz, 1H), 7.03 (t, *J* = 7.2 Hz, 1H), 3.74 (s, 3H), 3.16 (s, 1H), 1.87 (dd, *J* = 2.8 Hz, *J* = 12.8 Hz, 2H), 1.34 (t, *J* = 12.4 Hz, 2H), 1.26 (s, 6H), 1.14 (s, 6H); LCMS: 447.4 [M+H]⁺.

15

Example 14

Compound 95: 1-(cyclopropylsulfonyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole



To a stirred solution of 2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole (0.1 g 0.00031 mol), in THF (3 mL) was added NaH (0.031 g, 0.00079 mol) portion-wise at 0 °C under a N₂ atmosphere. The reaction mixture was stirred for 15 min followed by the addition of cyclopropylsulfonyl chloride (0.053 g 0.00038 mol) and then the reaction was stirred at rt for 16 h. The reaction mixture was cooled and quenched with water, extracted with ethyl acetate (2 x 20mL). The combined organic extracts were


washed with brine, dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as a white solid.



Yield: 0.012 g, 10 %. ^1H NMR (400 MHz, $\text{DMSO}-d_6$) δ 7.42 (t, $J = 7.6\text{Hz}$, 1H), 7.20 (d, $J = 7.2\text{Hz}$, 1H), 7.03 (d, 2H), 6.95 (t, $J = 7.6\text{Hz}$, 1H), 3.70 (s, 3H), 3.27 (s, 1H), 1.89 (s, 2H), 1.25 - 1.15 (m, 19H); LCMS 419.2 $[\text{M}+\text{H}]^+$.

By using analogous protocols to those described in the foregoing example, the compounds described in Table 4 have been prepared, using appropriately substituted aryl or alkyl sulfonyl chlorides or of sulfonic anhydrides.

10


Table 4

Cpd No.	Structure	Compound name	Analytical data
96		1-(isobutylsulfonyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole	^1H NMR (CDCl_3 , 400 MHz) δ 7.46 - 7.42 (m, 1H), 7.35 - 7.33 (m, 1H), 7.14 (s, 1H), 7.02 (t, $J = 7.6\text{Hz}$, 1H), 6.94 (d, $J = 8.4\text{Hz}$, 1H), 3.80 (s, 3H), 3.19 - 3.12 (m, 3H), 2.20 - 2.14 (m, 1H), 2.01 - 1.97 (m, 2H), 1.45 - 1.39 (m, 2H), 1.33 (s, 6H), 1.25 (s, 6H), 1.02 (d, $J = 6.4\text{Hz}$, 6H). LCMS: 435.3 $[\text{M}+\text{H}]^+$

Cpd No.	Structure	Compound name	Analytical data
97		1-((4-chlorophenyl)sulfonyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.6 (d, <i>J</i> = 8.8Hz, 2H), 7.50 (s, 1H), 6.59 (s, 1H), 7.40(t, <i>J</i> = 7.2Hz, 1H), 7.05 - 6.94 (m, 3H), 3.45(s, 3H), 1.03(t, <i>J</i> = 12.0Hz, 1H), 1.84(dd, <i>J</i> = 2.8Hz, 2.4Hz, 2H), 1.29 (d, <i>J</i> = 12Hz, 2H) 1.23(s, 6H), 1.12 (s, 6H): LCMS: 489.2 [M+H] ⁺
98		1-((4-fluorophenyl)sulfonyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole	¹ H NMR (CDCl ₃ , 400 MHz) δ 7.69 -7.66 (m, 2H), 7.50 (s, 1H), 7.49 - 7.42 (m, 3H), 7.04 (m, 3H), 7.05 - 6.94 (m, 3H), 3.47 (s, 3H), 3.02 (s, 1H), 1.84 (dd, <i>J</i> = 2.8Hz, 2.4Hz, 2H), 1.30 - 1.23(m, , 9H), 1.10 (s 6H): LCMS: 473.3 [M+H] ⁺



Example 15

Compound 99: 4-((2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)methyl)benzonitrile




- To a stirred solution of 2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole (0.1 g, 0.00031 mol), in THF was added NaH (0.031 g, 0.000791 mol) portion- wise at 0 °C under a N₂ atmosphere. The reaction mixture was stirred for 15 min
- 5 followed by the addition of 4-cyanobenzyl bromide (0.074 g, 0.00034mol). The reaction was then stirred at rt for 16 h. The reaction mixture was cooled and quenched with water, and extracted with ethyl acetate (2 x 20 mL). The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced
- 10 pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as white solid. (0.081 g, 59.5%). ¹H NMR (CDCl₃, 400 MHz) δ 7.59 (d, *J* = 8.4Hz, 2H), 7.41 - 7.36 (m, 2H), 7.14 (d, *J* = 8.0Hz, 2H), 7.01(t, *J* = 7.6Hz, 1H), 6.85 (d, *J* = 8.4Hz, 1H), 6.60 (s, 1H), 4.90(s, 2H), 3.60(s, 3H), 3.10(t, *J* = 12.8Hz, 1H), 2.0 (dd, *J* = 2.4Hz, 2.0Hz, 2H), 1.23(s, 6H), 1.13(s, 6H); LCMS: 430.5 [M+H]⁺.
- 15 By using analogous protocols to those described in the foregoing example the compounds described in Table 5 have been prepared using appropriately substituted bromide compounds.

Table 5

Cpd No.	Structure	Compound name	Analytical data
100		5-((2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)methyl)pyrimidine	¹ H NMR (CDCl ₃ , 400 MHz) δ 9.13 (s, 1H), 8.4(s 2H), 7.51 - 7.39 (m, 2H), 7.03 (t, <i>J</i> = 15.2Hz, 1H), 6.93 (d, <i>J</i> = 8.4Hz, 1H), 6.65 (s, 1H), 3.6 (s, 3H), 3.10(t, <i>J</i> = 12.8Hz, 1H) 1.98 (dd, <i>J</i> = 12.8Hz, 2H) 1.40 (t, 2H), 1.30(s, 6H), 1.23(s, 6H): LCMS: 407.2 [M+H] ⁺
101		3-((2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)methyl)benzonitrile	¹ H NMR (CDCl ₃ , 400 MHz) δ 7.56 (d, <i>J</i> = 8.0Hz, 1H), 7.43 - 7.25 (m, 5H), 7.01 (t, <i>J</i> = 7.6Hz, 1H), 6.92(d, <i>J</i> = 8.8Hz, 1H), 6.60 (s, 1H), 6.60 (s, 1H), 4.90 (s, 2H), 3.60 (s, 3H), 3.1(t, <i>J</i> = 12.8Hz, 1H), 2.0 (dd, <i>J</i> = 2.4Hz, 2.0Hz, 2H), 1.53 - 1.37 (m, 2H), 1.3(s, 6H), 1.23 (s, 6H): LCMS: 430.5 [M+H] ⁺



Example 16



Compound 102: 4-(1-(4-chlorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl)-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol







- To a stirred solution of N-(4-chlorophenyl)-2-methoxybenzimidamide **A 39** (0.37 g, 0.0014 mol) in dioxane (15 mL) was added 2-bromo-1-(4-hydroxy-2,2,6,6-tetramethyl-tetrahydro-pyran-4-yl)-ethanone **A 75** (0.4 g, 0.0014 mol) and NaHCO₃ (0.24 g, 0.00286 mol) and the mixture was stirred for 5 h at 85 °C. The solvent was evaporated and the resultant crude residue was partitioned between ethyl acetate and water. The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as a pale yellow solid. Yield: 0.12 g, 20%. ¹H NMR (400 MHz, DMSO-d₆) 7.47 -7.45(m, 1H), 7.40 – 7.35 (m, 3H), 7.26(s, 1H), 7.09 (d, *J*=8.8Hz, 2H), 7.01 (t, *J*=7.6Hz, 1H), 6.87 (d, *J*=8.4Hz, 1H), 4.82 (s, 1H), 3.21 (s, 3H), 1.87 – 1.76 (m, 4H), 1.44 (s, H), 1.10 (s, 6H); LCMS: 441.2 [M+H]⁺.
- By using analogous protocols to those described in the foregoing example the compounds described in Table 6 have been prepared using an appropriately substituted amidine intermediate.



Table 6



Cpd No.	Structure	Compound name	Analytical data
103		4-(1-(4-chlorophenyl)-2-(4-methoxypyridin-3-yl)-1H-imidazol-4-yl)-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol	¹ H NMR (400 MHz, DMSO-d ₆) δ = 8.51 (s, 1H), 8.47 (d, <i>J</i> = 5.6 Hz, 1H), 7.44 (d, <i>J</i> = 8.4 Hz, 2H), 7.36 (s, 1H), 7.17 (d, <i>J</i> = 8.4 Hz, 2H), 6.96 (d, <i>J</i> = 5.6 Hz, 1H), 4.92 (s, 1H), 3.35 (s, 3H), 1.86 (d, <i>J</i> = 13.2 Hz, 2H), 1.80 (d, <i>J</i> = 13.2 Hz, 2H), 1.46 (s, 6H), 1.12 (s, 6H). LCMS: 442.44 [M+H] ⁺
104		4-(1-(4-chlorophenyl)-2-(3-methoxypyridin-2-yl)-1H-imidazol-4-yl)-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol	¹ H NMR (400 MHz, DMSO-d ₆) δ = 8.19 (s, 1H), 7.38 - 7.44 (m, 4H), 7.32 (s, 1H), 7.08 (d, <i>J</i> = 7.6 Hz, 2H), 4.94 (s, 1H), 3.44 (s, 3H), 1.77 - 1.87 (m, 4H), 1.46 (s, 6H), 1.11 (s, 6H). LCMS: 442.48 [M+H] ⁺


Cpd No.	Structure	Compound name	Analytical data
105		4-(1-(4-chlorophenyl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-4-yl)-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ = 8.20 (d, <i>J</i> = 8 Hz, 1H), 7.93 (d, <i>J</i> = 7.2 Hz, 1H), 7.45 (d, <i>J</i> = 8.4 Hz, 2H), 7.34 (s, 1H), 7.15 (d, <i>J</i> = 8.4 Hz, 2H), 7.10 (t, <i>J</i> = 6.8 Hz, 1H), 4.93 (s, 1H), 3.34 (s, 3H), 1.77 - 1.87 (m, 4H), 1.46 (s, 6H), 1.11 (s, 6H). LCMS: 442.44 [M+H] ⁺
106		4-(1-(3-bromo-4-fluorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl)-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.46 (d, <i>J</i> = 8.0Hz, 2H), 7.34 - 7.40 (m, 2H), 7.31 (s, 1H), 7.08 - 7.12 (m, 1H), 7.02 (t, <i>J</i> = 7.2Hz, 1H), 6.80 (d, <i>J</i> = 8.4Hz, 1H), 4.80 (s, 1H), 3.25 (s, 3H), 1.80 - 1.82 (m, 4H), 1.44 (s, 6H), 1.22 (s, 6H). LCMS: 453.2 [M+H] ⁺

Cpd No.	Structure	Compound name	Analytical data
107		4-(1-(3,4-difluorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl)-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.47 (m, 1H), 7.48 (m, 2H), 7.26 (m, 2H), 7.02(t, <i>J</i> =8Hz, 1H), 6.89 (m, 2H), 4.8 (s, 1H), 3.2 (s, 3H), 1.81 (m, 4H), 1.44(s, 6H), 1.10(s, 6H). LCMS: 443.3[M+H] ⁺
108		4-(1-(3,4-difluorophenyl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-4-yl)-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol	¹ H NMR (400 MHz, DMSO-d ₆) δ 7.48(d, <i>J</i> = 7.6Hz, 1H), 7.39 (t, <i>J</i> = 8Hz, 1H), 7.34 (s, 1H), 7.2 (dd, <i>J</i> = 8Hz, 10.4Hz, 1H), 7.04 (t, <i>J</i> = 7.6Hz, 1H), 6.92 (t, <i>J</i> = 12Hz, 1H), 4.8 (s, 1H), 3.2 (s, 3H), 1.81 (m, 4H), 1.1(s, 6H), 1.4(s, 6H). LCMS: 444.2 [M+H] ⁺

Cpd No.	Structure	Compound name	Analytical data
109		4-(2-(2-methoxyphenyl)-1-(6-(trifluoromethyl)pyridin-3-yl)-1H-imidazol-4-yl)-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.574(s, 1H), 7.93(d, <i>J</i> = 8.4Hz, 1H), 7.804 (d, <i>J</i> = 10.4Hz, 1H), 7.57(d, <i>J</i> = 6Hz, 1H), 7.516 (s, 1H), 7.430 (m, 1H), 7.101(m, 1H), 6.898(d, <i>J</i> = 8.4Hz, 1H), 4.937(s, 1H), 3.16(s, 3H), 1.85(m, 4H), 1.47(s, 6H), 1.13 (s, 6H); LCMS: 476.3 [M+H] ⁺
110		4-(1-(3-chloro-4-fluorophenyl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-4-yl)-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 8.19(t, <i>J</i> = 7.2Hz, 2H), 7.91(m, 1H), 7.46 (m, 1H), 7.42 (d, <i>J</i> = 8.0Hz, 1H), 7.36 (s, 1H), 7.11 (m, 2H), 4.85(s, 1H), 3.37(s, 3H), 1.81(dd, <i>J</i> = 13.6Hz, 5.2Hz, 4H), 1.44(s, 6H), 1.10(s, 6H); LCMS: 460.2 [M+H] ⁺

Cpd No.	Structure	Compound name	Analytical data
111		4-(1-(3-chloro-4-fluorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl)-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.5 (d, <i>J</i> = 6Hz, 1H), 7.40 - 7.39 (m, 3H), 6.20 (s, 1H), 7.09 - 7.01 (m, 2H), 6.80 (d, <i>J</i> = 8.4Hz, 1H), 4.80 (s, 1H), 3.25 (s, 3H), 1.86 - 1.77 (m, 4H), 1.44 (s, 6H), 1.10 (s, 6H); LCMS: 459.2 [M+H] ⁺
112		4-(1-(4-chloro-3-fluorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl)-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol	¹ H NMR (400 MHz, DMSO- <i>d</i> ₆) δ 7.53 (t, <i>J</i> = 12Hz, 1H), 7.48 (d, <i>J</i> = 7.6Hz, 1H), 7.40 (t, <i>J</i> = 8Hz, 1H), 7.34 (s, 1H), 7.2 (dd, <i>J</i> = 10Hz, 10.4Hz, 1H), 7.01 (t, <i>J</i> = 8Hz, 1H), 6.90 (t, <i>J</i> = 8Hz, 2H), 4.8 (s, 1H), 3.2 (s, 3H), 1.81 (m, 4H), 1.44 (s, 6H), 1.10 (s, 6H). LCMS: 459.2 [M+H] ⁺


Cpd No.	Structure	Compound name	Analytical data
113		4-(1-(4-chloro-3-fluorophenyl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-4-yl)-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol	¹ H NMR (400 MHz, DMSO-d ₆) δ 8.20 (t, <i>J</i> = 4Hz, 1H), 7.93 (dd, <i>J</i> = 8Hz, 8Hz, 1H), 7.58 (t, <i>J</i> = 8Hz, 1H), 7.38 (s, 1H), 7.31 (d, <i>J</i> = 1.6Hz, 1H), 7.10 (m, 1H), 6.96 (d, <i>J</i> = 8Hz, 1H), 4.8 (s, 1H), 3.3 (s, 3H), 1.81 (m, 4H), 1.44 (s, 6H), 1.10 (s, 6H). LCMS: 460.2 [M+H] ⁺
114		4-(1-(4-chlorophenyl)-2-(2-ethoxypyridin-3-yl)-1H-imidazol-4-yl)-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol	¹ H NMR (400 MHz, DMSO-d ₆): 8.17 (d, <i>J</i> = 3.2Hz, 1H), 7.93 (d, <i>J</i> = 5.6Hz, 1H), 7.43 (d, <i>J</i> = 8.4Hz, 2H), 7.33 (s, 1H), 7.15 (d, <i>J</i> = 8.8Hz, 2H), 7.07 (m, 1H), 4.86 (s, 1H), 3.86 (q, <i>J</i> = 2.8Hz, 2H), 1.82 (q, <i>J</i> = 12.8Hz, 4H), 1.44 (s, 6H), 1.26 (s, 2H), 1.15 (s, 6H), 0.80 (q, <i>J</i> = 5.6Hz, 3H). LCMS: 456.3 [M+H] ⁺

Cpd No.	Structure	Compound name	Analytical data
115		4-(4-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR: (400 MHz, DMSO-d ₆): 8.21 – 8.19 (m, 1H), 7.98 – 7.95 (m, 1H), 7.86 (d, <i>J</i> = 8.4Hz, 2H), 7.42 (s, 1H), 7.32 (d, <i>J</i> = 8.8Hz, 2H), 7.13 – 7.10 (m, 1H), 4.90 (s, 1H), 1.89 – 1.77 (m, 4H), 1.45 (s, 6H), 1.10 (s, 6H). LCMS: 433.3 [M+H] ⁺

Example 17

Compound 116: 2-Fluoro-5-[4-(4-hydroxy-2,2,6,6-tetramethyl-tetrahydro-pyran-4-yl)-2-(2-methoxy-phenyl)-imidazol-1-yl]-benzonitrile

5



Step 1. Compound 126. 2-Fluoro-5-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyl-3,6-dihydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile. To a stirred solution of 4-[1-(3-bromo-4-fluoro-phenyl)-2-(2-methoxy-phenyl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyl-tetrahydro-pyran-4-ol, (0.16 g, 0.000635 mol) in DMF (5 mL) under an N₂ atmosphere was added Zn(CN)₂ (0.075 g, 0.00212 mol) and tetrakis-triphenylphosphine palladium (0.184 g, 0.00015 mol), purging with nitrogen for 15 min before each addition. The reaction


10



mixture was further purged for 5 min and stirred for 1.5 h at 160 °C. The reaction mixture was quenched with ice cold water and extracted with ethyl acetate (50 mL x 3). The combined organic extracts were washed with brine, dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as a white solid. (0.09 g, 63.3%), LCMS: 432.3 [M+H]⁺.


Step 2. Compound 116: 2-Fluoro-5-[4-(4-hydroxy-2,2,6,6-tetramethyl-tetrahydropyran-4-yl)-2-(2-methoxy-phenyl)-imidazol-1-yl]-benzonitrile. To a stirred solution of 2-fluoro-5-[2-(2-methoxy-phenyl)-4-(2,2,6,6-tetramethyl-3,6-dihydro-2H-pyran-4-yl)-imidazol-1-yl]-benzonitrile (0.09 g, 0.000208 mol), and Mn(tmhd)₃ (0.025 g, 0.0004172 mol) in i-PrOH (7 mL) and DCM (1 mL) at 0 °C, phenylsilane (0.045 g, 0.000417 mol) was added and the reaction mixture was stirred for 3 h from 0 °C to rt under an oxygen atmosphere. Saturated Na₂S₂O₃ solution (2 mL) was added and the mixture was stirred for 2 h. The reaction mixture was further diluted with brine (10 mL) and extracted with ethyl acetate (10 mL x 3). The combined organic extracts were dried over anhydrous Na₂SO₄, filtered, and concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using ethyl acetate in hexane afforded the title compound as an off white solid (0.05 g, 56 %). ¹H NMR (CDCl₃, 400 MHz) δ 7.54-7.56 (dd, *J* = 7.6 Hz, 1.2 Hz, 1H), 7.38 - 7.43 (m, 2H), 7.31 - 7.36 (m, 2H), 7.14 (t, *J* = 8.8Hz, 1H), 7.05 (t, *J* = 8.0Hz, 1H), 7.02 (s, 1H), 6.74 (d, *J* = 8.4Hz, 1H), 3.38 (s, 3H), 2.06 (d, *J* = 13.2Hz, 2H), 1.90 (d, *J* = 13.6Hz, 2H), 1.56 (s, 6H), 1.26 (s, 6H). LCMS: 450.3.2 [M+H]⁺

By using analogous protocols to those described in the foregoing example the compounds described in Table 7 have been prepared using appropriately substituted bromide intermediate.

Table 7

Cpd No.	Structure	Compound name	Analytical data
117		2-fluoro-5-(4-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-2-(2-methoxyphenyl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR (CDCl ₃ , 400 MHz) δ 7.54-7.56 (dd, <i>J</i> = 7.6 Hz, 1.2 Hz, 1H), 7.38 - 7.43 (m, 2H), 7.31 - 7.36 (m, 2H), 7.14 (t, <i>J</i> = 8.8Hz, 1H), 7.05 (t, <i>J</i> = 8.0Hz, 1H), 7.02 (s, 1H), 6.74 (d, <i>J</i> = 8.4Hz, 1H), 3.38 (s, 3H), 2.06 (d, <i>J</i> = 13.2Hz, 2H), 1.90 (d, <i>J</i> = 13.6Hz, 2H), 1.56 (s, 6H), 1.26 (s, 6H). LCMS: 450.3 [M+H] ⁺


Cpd No.	Structure	Compound name	Analytical data
118		2-fluoro-4-(4-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-2-(2-methoxyphenyl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR:(400 MHz, DMSO-d ₆):δ 7.89 (t, <i>J</i> = 8.0Hz, 1H), 7.52 (d, <i>J</i> = 7.2Hz, 1H), 7.43 - 7.40 (m, 2H), 7.35 (d, <i>J</i> = 10Hz, 1H), 7.07 (t, <i>J</i> = 7.6Hz, 2H), 6.90 (d, <i>J</i> = 8.4Hz, 1H), 4.88 (s, 1H), 3.22 (s, 3H), 1.85 - 1.77 (m, 4H), 1.44 (s, 6H), 1.10 (s, 6H). LCMS: 450.3 [M+H] ⁺
119		2-fluoro-4-(4-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-1-yl)benzonitrile	¹ H NMR:(400 MHz, DMSO-d ₆):δ 8.24 - 8.22 (dd, <i>J</i> = 4.8Hz, 1.6Hz, 1H), 7.99 - 7.97 (dd, <i>J</i> = 7.2Hz, 1.6Hz, 1H), 7.93 (t, <i>J</i> = 8Hz, 1H), 7.48 (s, 1H), 7.44 (d, <i>J</i> = 1.6Hz, 1H), 7.34 (bs, 1H), 7.15 - 7.12 (m, 2H), 4.92 (s, 1H), 3.33 (s, 3H), 1.85 - 1.77 (m, 4H), 1.45 (s, 6H), 1.10 (s, 6H). LCMS: 451.3 [M+H] ⁺

Cpd No.	Structure	Compound name	Analytical data
120		4-(1-(4-chlorophenyl)-2-(3-methoxypyridin-4-yl)-1H-imidazol-4-yl)-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol	¹ H NMR (DMSO-d ₆ , 400 MHz): 8.31-8.27 (m, 2H), 7.52 (d, <i>J</i> = 4.4 Hz, 1H), 7.44 (d, <i>J</i> = 18.4 Hz, 2H), 7.38 (s, 1H), 7.16 (d, <i>J</i> = 8.8 Hz, 2H), 3.32 (s, 3H) 4.95 (s, 1H), 1.87-1.84 (m, 4H), 1.46 (s, 6H), 1.11 (s, 6H). MS: ESI-MS, <i>m/z</i> 442.19 (M+1).

Example 18

Compound 122: 3-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)pyridin-2(1H)-one

5



122

To a stirred solution of 3-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine (320 mg) was added aqueous HBr (47%) 2 mL at 0-5 °C, and the mixture was stirred at rt for 16 h. The reaction was quenched with 10% NaHCO₃ solution(30ml), extracted with 10% MeOH in DCM (2 x 50 mL). The




10





combined organic extracts were dried over anhydrous Na_2SO_4 , filtered, and concentrated under reduced pressure to provide a residue. Purification by column chromatography on silica gel (100-200 mesh) using a mixture of MeOH and DCM afforded the title compound as an off white solid. Yield: 0.015 g (48.5%); LCMS 412.2 $[\text{M}+\text{H}]^+$





5





The compounds of Table 8, exemplified hereinbelow, were prepared according to the schemes and specific examples described herein.





Table 8. Compounds of Formula (I)





Cpd No.	Structure
1	
2	
3	





Cpd No.	Structure
4	
5	
6	
7	





Cpd No.	Structure
8	
9	
10	
11	





Cpd No.	Structure
12	
13	
14	
15	





Cpd No.	Structure
16	
17	
18	
19	





Cpd No.	Structure
20	
21	
22	
23	





Cpd No.	Structure
24	
25	
26	
27	





Cpd No.	Structure
28	
29	
30	
31	





Cpd No.	Structure
32	
33	
34	
35	





Cpd No.	Structure
36	
37	
38	
39	





Cpd No.	Structure
40	
41	
42	
43	





Cpd No.	Structure
44	
45	
46	
47	





Cpd No.	Structure
48	
49	
50	
51	





Cpd No.	Structure
52	
53	
54	
55	





Cpd No.	Structure
56	
57	
58	
59	





Cpd No.	Structure
60	
61	
62	
63	





Cpd No.	Structure
64	
65	
66	
67	





Cpd No.	Structure
68	
69	
70	
71	





Cpd No.	Structure
72	 <chem>COc1ccc(cc1)-c2nc(C3CC(C)(C)OC3C)c4cc(F)cc(F)n4</chem>
73	 <chem>COc1ccc(cc1)-c2nc(C3CC(C)(C)OC3C)c4cc(F)cc(F)n4</chem>
74	 <chem>COc1ccc(cc1)-c2nc(C3CC(C)(C)OC3C)c4cc(F)cc(F)n4</chem>
75	 <chem>COc1ccc(cc1)-c2nc(C3CC(C)(C)OC3C)c4cc(F)cc(F)n4</chem>





Cpd No.	Structure
76	
77	
78	
79	





Cpd No.	Structure
80	
81	
82	
83	





Cpd No.	Structure
84	
85	
86	
87	





Cpd No.	Structure
88	
89	
90	
91	





Cpd No.	Structure
92	
93	
94	
95	





Cpd No.	Structure
96	
97	
98	
99	





Cpd No.	Structure
100	
101	
102	
103	




Cpd No.	Structure
104	
105	
106	
107	

Cpd No.	Structure
108	
109	
110	
111	

Cpd No.	Structure
112	
113	
114	
115	

Cpd No.	Structure
116	
117	
118	
119	

Cpd No.	Structure
120	
121	
122	
123	

Cpd No.	Structure
124	
125	
126	

Biological Examples

5

In Vitro Assays

Example 1

Functional Assay: Calcium Influx Assay

10 A Ca flux assay using Functional Drug Screening System (FDSS, Hamamatsu) was utilized to identify novel N-type Ca channel antagonists (Dai et al 2008, Beladreddetti et al 2009). All cell culture reagents were procured from HyClone and other reagents were

from Fisher. Recombinant HEK 293 cells expressing human N-type Ca channel (Millipore) were maintained in DMEM F12 containing 10% FBS and selection antibiotics at 37 °C in 5% CO₂.

When CaV2.2 cells were 75% - 85% confluent in their culture flask, they were removed from the flask, counted, and plated at 13,000-15,000 cells (50 µL) per well in black-wall clear-bottom 384-well assay plates. Cells were incubated at 30 °C in 5% CO₂ overnight. The medium was removed and cells were loaded with Fluo8 dye (AAT Bioquest) at 37 °C and 30 °C, 30 min each in darkness. Cells were washed four times with wash buffer (16 mM HEPES pH 7.2, 2 mM CaCl₂, 1 mM MgCl₂, 5 mM glucose, 140 mM choline chloride and 2 mM KCl) leaving 15 µL buffer after the last wash. Compounds of Formula (I), serially diluted, were added to the cell plate. The plates were incubated in darkness for 15 min. Cells were depolarized with 100 mM KCl and Ca influx was measured. Data was plotted as percent inhibition vs concentration of the compound and IC₅₀ values were generated using Graphpad Prism non-linear regression analysis. Resultant data are shown in Table 9.

Example 2

Measurement of Electrophysiological Response

The recombinant cell line described above was used in the measurement of electrophysiological properties with QPatch (Sophion). Cells were maintained in culture as mentioned above. Prior to the experiment, the flasks were moved to 30 °C and incubated for 48-72 hrs. On the day of assaying, cells were detached and harvested. The cell pellet was re-suspended in SMF4HEK (HyClone)/25mM HEPES at 2-5 million/ mL and placed on the QStirrer of the QPatch for 30-60 min prior to the start of the assay. The Intracellular Buffer contained 97.9 mM CsCl, 27 mM CsF, 8.2 mM EGTA, 10 mM HEPES, 2 mM NaCl, 0.3 mM GTP, 3mM Mg-ATP pH ~7.3 adjusted with CsOH and ~280mOsm. Extracellular buffer contained 132 mM NaCl, 5.4 mM KCl, 1.8 mM CaCl₂, 0.8 mM MgCl₂, 10 mM HEPES, 10 mM glucose pH ~7.4 adjusted with NaOH/ HCl and

~300mOsm. Microtiter-Plate/Reference Buffer contained 140 mM TEA-Cl, 10mM BaCl₂, 0.8 mM MgCl₂, 10 mM HEPES pH ~7.4 adjusted with TEA-OH/HCl, and ~300mOsm. Cells were transferred to a QPatch 48 plate and gigaseal was formed. The plate was perfused with intracellular and extracellular buffer. Compounds diluted in reference buffer were applied to the extracellular site and tested following the voltage protocol as described by Finley et al (2010). In brief, cells were depolarized from resting potential of -80 mV to +20 mV and current was recorded. Data was plotted as percent inhibition vs concentration of the compound and IC₅₀ values were generated using Graphpad Prism non-linear regression analysis. Resultant data is reported in Table 9. Concentration ranges for each compound are included with its Q patch value.

Table 9
FDSS and Q patch Assay Data

Cpd No.	N Type IC ₅₀ Data (μM)		
	FDSS	Qpatch Pulse=1	Qpatch Pulse=15
1	0.041	0.365 (0.01- 1 μM)	>1 (0.01- 1 μM)
2	0.128	NE (10 μM)	6.2 (0.3-10 μM)
3	0.376	0.830 (0.01- 1 μM)	NE (0.01- 1 μM)
4	0.017	10 (0.3-10 μM)	1.06 (0.3-10 μM)
5	0.191	5.9 (0.3-10 μM)	10 (0.3-10 μM)
6	0.739	10 (0.3-10 μM)	10 (0.3-10 μM)
7	0.113	1.17 (0.3-10 μM)	0.68 (0.3-10 μM)
8	>10	0.508 (0.01- 1 μM)	> 1 (0.01- 1 μM)
9	0.793	46 % (0.01- 1 μM)	37 % (0.01- 1 μM)
10	0.401	5.92 (0.3-10 μM)	1.47 (0.3-10 μM)
11	0.082	63 % (10 μM)	21 % (10 μM)
12	0.63	28% (10 μM)	NE (10 μM)

	N Type IC ₅₀ Data (μM)		
Cpd No.	FDSS	Qpatch Pulse=1	Qpatch Pulse=15
13	0.2	>1 (0.01- 1 μM)	NE (0.01- 1 μM)
14	2.7	0.266 (0.01- 1 μM)	>1 (0.01- 1 μM)
15	0.256	poor fit(0.01-0.3 μM)	poor fit(0.01-0.3 μM)
16	0.359	poor fit(0.01-0.3 μM)	poor fit(0.01-0.3 μM)
17	0.591	4.4 (0.3-10 μM)	5.2 (0.3-10 μM)
18	0.108	0.694 (0.01- 1 μM)	0.376 (0.01- 1 μM)
19	0.263	0.966 (0.01- 1 μM)	28 % (1 μM)
20	1.06	>1 (0.01- 1 μM)	NE (0.01- 1 μM)
21	0.119	0.639 (0.01- 1 μM)	39 % (0.01- 1 μM)
22	0.075	0.590 (0.01- 1 μM)	0.847 (0.01- 1 μM)
23	0.31	21 % (0.01- 1 μM)	NE (0.01- 1 μM)
24	7.15	29 % (0.01- 1 μM)	NE (0.01- 1 μM)
25	0.818	>1 (0.01- 1 μM)	NE (0.01- 1 μM)
26	5.76	>1 (0.01- 1 μM)	>1 (0.01- 1 μM)
27	6.4	poor fit(0.01-0.3 μM)	1.9(0.01-0.3 μM)
28	1.64	poor fit(0.01-0.3 μM)	poor fit(0.01-0.3 μM)
29	0.492	poor fit(0.01-0.3 μM)	poor fit(0.01-0.3 μM)
30	0.321	67 % (10 μM)	35 % (10 μM)
31	0.769	1.58(0.01-0.3 μM)	poor fit(0.01-0.3 μM)
32	0.278	>1 (0.01- 1 μM)	NE (0.01- 1 μM)
33	0.459	>1 (0.01- 1 μM)	NE (0.01- 1 μM)

	N Type IC ₅₀ Data (μM)		
Cpd No.	FDSS	Qpatch Pulse=1	Qpatch Pulse=15
34	1.62	38 % (0.01- 1 μM)	NE (0.01- 1 μM)
35	0.838	NA	NA
36	0.04	0.376 (0.01- 1 μM)	NE (0.01- 1 μM)
37	0.424	NE (0.01- 1 μM)	NE (0.01- 1 μM)
38	0.372	2.4 (0.3-10 μM)	0.769 (0.3-10 μM)
39	0.163	1.7 (0.3-10 μM)	1.9 (0.3-10 μM)
40	0.822	52 % (10 μM)	1.94 (0.3-10 μM)
41	0.131	71 % (10 μM)	0.602 (0.3-10 μM)
42	0.194	2.6 (0.3-10 μM)	1.08 (0.3-10 μM)
43	0.291	34 % (10 μM)	7.5 (0.3-10 μM)
44	0.142	2.9 (0.3-10 μM)	3.08 (0.3-10 μM)
45	0.149	2.2 (0.3-10 μM)	0.671 (0.3-10 μM)
46	0.255	10 (0.3-10 μM)	2.5 (0.3-10 μM)
47	1.54	>1 (0.01- 1 μM)	NE (0.01- 1 μM)
48	5.41	0.222(0.01- 1 μM)	0.843 (0.01- 1 μM)
49	0.123	>1 (0.01- 1 μM)	NE (0.01- 1 μM)
50	0.215	>1 (0.01- 1 μM)	0.971 (0.01- 1 μM)
51	0.029	0.179 (0.01- 1 μM)	0.701 (0.01- 1 μM)
52	0.115	0.721 (0.01- 1 μM)	NE (0.01- 1 μM)
53	0.096	6.5 (0.3-10 μM)	2.01 (0.3-10 μM)
54	1.96	0.302 (0.01- 1 μM) Poor fit(0.01- 1 μM)	0.237 (0.01- 1 μM) Poor fit(0.01- 1 μM)
55	0.384	48 % (0.01- 1 μM)	0.780 (0.01- 1 μM)
56	1.67	NE (0.01- 1 μM)	NE (0.01- 1 μM)
57	10	NE (0.01- 1 μM)	NE (0.01- 1 μM)
58	4.81	NE (0.01- 1 μM)	NE (0.01- 1 μM)

	N Type IC ₅₀ Data (μM)		
Cpd No.	FDSS	Qpatch Pulse=1	Qpatch Pulse=15
59	4.4	poor fit(0.01-0.3 μM)	poor fit(0.01-0.3 μM)
60	0.321	0.247(0.01-0.3 μM)	poor fit(0.01-0.3 μM)
61	0.037	0.751(0.01-0.3 μM)	1.05 (0.01-0.3 μM)
62	0.177	81 % (10 μM)	0.413 (0.3-10 μM)
63	0.213	53 % (10 μM)	3.5 (0.3-10 μM)
64	0.36	4.06 (0.3-10 μM)	1.72 (0.3-10 μM)
65	0.225	58 % (10 μM)	8.6 (0.3-10 μM)
66	0.049	10 (0.3-10 μM)	1.3 (0.3-10 μM)
67	0.166	NE (10 μM)	NE (10 μM)
68	0.109	10 (0.3-10 μM)	4.6 (0.3-10 μM)
69	1.43	53 % (10 μM)	29 % (10 μM)
70	0.871	41 % (10 μM)	NE (10 μM)
71	0.422	5.83 (0.01- 1 μM)	poor fit(0.01- 1 μM)
72	2.4	poor fit(0.01- 1 μM)	poor fit(0.01- 1 μM)
73	8.16	poor fit(0.01- 1 μM)	poor fit(0.01- 1 μM)
74	0.488	36 % (10 μM)	44 % (10 μM)
75	2.060	9.8 (0.3-10 μM)	4.2 (0.3-10 μM)
76	0.151	>1 (0.01- 1 μM)	NE (0.01- 1 μM)
77	10	>1 (0.01- 1 μM)	NE (0.01- 1 μM)
78	0.234	58 % (0.01- 1 μM)	NE (0.01- 1 μM)
79	1.33	29 % (10 μM)	3.6 (0.3-10 μM)
80	0.333	NE (10 μM)	28 % (10 μM)
81	0.201	6.7 (0.3-10 μM)	2.8 (0.3-10 μM)
82	3.26	poor fit(0.01-0.3 μM)	poor fit(0.01-0.3 μM)

	N Type IC ₅₀ Data (μM)		
Cpd No.	FDSS	Qpatch Pulse=1	Qpatch Pulse=15
83	0.049	poor fit(0.01-0.3 μM)	poor fit(0.01-0.3 μM)
84	0.362	39 % (0.01- 1 μM)	NE (0.01- 1 μM)
85	6.48	45 % (0.01- 1 μM)	NE (0.01- 1 μM)
86	1.3	28 % (0.01- 1 μM)	NE (0.01- 1 μM)
87	6.2	poor fit(0.01-0.3 μM)	poor fit(0.01-0.3 μM)
88	2.17	0.349 (0.01-0.3 μM)	poor fit(0.01-0.3 μM)
89	0.205	0.517 (0.01-0.3 μM)	poor fit(0.01-0.3 μM)
90	0.12	poor fit(0.01-0.3 μM)	poor fit(0.01-0.3 μM)
91	1.43	58 % (10 μM)	62 % (10 μM)
92	0.041	0.253 (0.01- 1 μM)	0.269 (0.01- 1 μM)
93	0.15	6.5 (0.3-10 μM)	1.8 (0.3-10 μM)
94	0.551	NE (0.01- 1 μM)	NE (0.01- 1 μM)
95	10	NE (0.01- 1 μM)	NE (0.01- 1 μM)
96	0.768	10 (0.3-10 μM)	5.4 (0.3-10 μM)
97	0.515	NE (10 μM)	3.08 (0.3-10 μM)
98	0.692	9.02 (0.3-10 μM)	3.6 (0.3-10 μM)
99	10.9	NE (0.01- 1 μM)	NE (0.01- 1 μM)
100	10	10 (0.3-10 μM)	10 (0.3-10 μM)
101	3.58	>1 (0.01- 1 μM)	NE (0.01- 1 μM)
102	0.037	48 % (10 μM)	1.74 (0.3-10 μM)
103	0.367	4.3 (0.3-10 μM)	31 % (10 μM)
104	0.464	NE (10 μM)	NE (10 μM)

	N Type IC ₅₀ Data (μM)		
Cpd No.	FDSS	Qpatch Pulse=1	Qpatch Pulse=15
105	0.3	8.3 (0.3-10 μM) 0.964(0.01- 1 μM)	3.82 (0.3-10 μM) NE (0.01- 1 μM)
106	0.148	71 % (10 μM)	5.03 (0.3-10 μM)
107	0.43	3.53(0.3-10 μM)	2.42 (0.3-10 μM)
108	0.394	5.8 (0.3-10 μM)	7.4 (0.3-10 μM)
109	0.322	4.5 (0.3-10 μM)	7.9 (0.3-10 μM)
110	0.385	44 % (10 μM)	30 % (10 μM)
111	0.257	62 % (10 μM)	2.5 (0.3-10 μM)
112	0.041	2.5 (0.3-10 μM)	3 (0.3-10 μM)
113	0.083	4.4 (0.3-10 μM)	7.8 (0.3-10 μM)
114	0.607	0.668 (0.01- 1 μM)	>1 (0.01- 1 μM)
115	0.731	NE (10 μM)	NE (10 μM)
116	1.37	>1 (0.01- 1 μM)	>1 (0.01- 1 μM)
117	1.44	40 % (10 μM)	25 % (10 μM)
118	0.15	8.1 (0.3-10 μM)	10 (0.3-10 μM)
119	1.29	28% (10 μM)	NE (10 μM)
120	0.108	1.6 (0.3-10 μM)	9.1 (0.3-10 μM)
121	0.233	6.6 (0.3-10 μM)	3.3 (0.3-10 μM)
122	3.010	NE (0.01- 1 μM)	NE (0.01- 1 μM)
123	0.129	0.468 (0.01- 1 μM)	>1 (0.01- 1 μM)
124	0.103	2.5 (0.3-10 μM)	1.6 (0.3-10 μM)
125	0.313	2.1 (0.3-10 μM)	2.3 (0.3-10 μM)
126	0.234	29 % (10 μM)	1.9 (0.3-10 μM)

In Vivo Assays

Example 3

Complete Freud's Adjuvant (CFA) Induced Mechanical Allodynia

Animals arrived and were acclimatized for 1-2 weeks in a temperature-controlled room with a 12 h light/dark cycle and allowed free access to standard laboratory chow and water.

CFA (Sigma) was injected intra-plantar (75 µg/150 µL) from a concentration of 1:1 (diluted 1 mg/ mL of CFA in 1 mL of PBS). CFA-induced mechanical allodynia was quantified by a Von frey test on day 2 (48 h post injection of CFA) and animals were randomized based on iPWT response (Dixon 1980). Animals exhibiting a PWT of <5.0 g were selected for testing. Animals were administered a compound of Formula (I) through appropriate route based on their pharmacokinetic properties. Responses were measured with a Von-Frey filament. The maximum possible effect (% MPE) was determined as 100% if sensitivity in ipsilateral paw neared that of the contralateral paw. Values from the vehicle-treated animals were considered 0%. The compound effect was determined based on these values. Resultant data are reported in Table 10, hereinbelow.

Example 4

Sciatic Nerve Ligation Model of Neuropathic Pain

Left L5 and L6 spinal nerves are isolated adjacent to the vertebral column and were ligated with 5-0 silk suture distal to the dorsal root ganglion, as described by Kim and Chung (1992). The incision was closed with the help of GLUture topical tissue adhesive. At 14 days post-surgery (one day prior to test compound administration), mechanical allodynia was quantified using eight von Frey filaments, calibrated in the range 0.4–15.1 g (Nielsen et al 2005). Rats were placed into individual plastic containers on top of a suspended wire mesh grid and acclimated to the test chambers for at least 15 min. Filaments were applied perpendicular to the mid-plantar paw surface, with enough force to

cause slight buckling, and held in place for 6-8 sec until a response was noted as a sharp paw withdrawal, flinching, licking and/or biting immediately upon removal of the filament (Nielsen et al 2005). Rats exhibiting increased mechanical sensitivity (ipsilateral paw withdrawal test cut-off of < 4.0 g) were selected and randomized. Test compounds of Formula (I) were administered and a response was measured at Tmax. The maximum possible effect (% MPE) was determined to be 100% if the sensitivity in the ipsilateral paw neared that of the contralateral paw. Values from vehicle-treated animals were considered 0 %. The compound effect was determined based on these values. Resultant data are reported in Table 10, hereinbelow.

10

References

Dai, G, Haedo RJ, Warren VA, Ratliff KS, Bugianesi RM, Rush A, *et al*: A high-throughput assay for evaluating state-dependence and subtype selectivity of CaV2 calcium channel inhibitors. *Assay Drug Dev Technol* , **2008**, 6, 195-212 .

15 Beladredetti F, Tringham E, Eduljee C, Jiang X, Dong H, Hendricson A, *et al*. “ A fluorescence-based high throughput screening assay for the identification of T-type calcium channel blockers”. *Assay and Drug Development Technologies*, **2009**, 7, 266-280.

Dixon,WJ, “Efficient analysis of experimental observations”. *Ann. Rev. Pharmacol. Toxicol.*, **1980**, 20, 441-462.

20 Finley FA, Lubin ML, Neeper MP, Beck E, Liu Y, Flores CM, Qin N. “An integrated multiassay approach to the discovery of small-molecule N-type voltage-gated calcium channel antagonists”. *Assay and Drug Development Technologies*, **2010**; 8(6), 685-694.

Kim, SH and Chung, JM., “An Experimental Model For Peripheral Neuropathy Produced By Segmental Spinal Nerve Ligation In The Rat”. *Pain*, **1992**, 50, 355-363.

25 Nielsen CK, Lewis RJ, Alewood D, Drinkwater R, Palant E, Patterson M, Yaksh TL, McCumber D and Smith MT. “Anti-allodynic efficacy of the c-conopeptide, Xen2174, in rats with neuropathic pain”. *Pain*, **2005**, 1–13.


Table 10*CFA and SNL In Vivo Data*

Dose (mg/kg)	CFA (% Reversal)		SNL(%MPE)			
	Cpd 1	Cpd 11	Cpd 1	Cpd 11	Cpd 10	Cpd 38
3	NA	20.97±7.88	NA	28.29± 10.84	12.17±11.24	NA
10	NA	41.12±5.14	NA	49.01±9.93	40.53±13.64	NA
12.5	19.73±8.25	NA	22.21±7.61	NA	NA	NA
25	37.62±7.17	NA	40.9±20.29	NA	NA	NA
30	NA	55.8±11.32	NA	70.69±10.3	54.59±13.37	NA
50	58.8±10.22	NA	64.07±12.3 1	NA	NA	87.01± 9.25
60	NA	76.4±3.78	NA	96.18±3.22	NA	NA
100	75.26±8.96	NA	75±13.01	NA	85.45±8.44	NA

While the foregoing specification teaches the principles of the present invention,
 5 with examples provided for the purpose of illustration, it will be understood that the
 practice of the invention encompasses all of the usual variations, adaptations and/or
 modifications as come within the scope of the following claims and their equivalents.

We claim:

1. A compound of Formula (I)



Formula (I)

wherein

R¹ is

- i) phenyl optionally independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, difluoromethoxy, and C₁₋₄alkoxy; provided that when phenyl of group (i) is substituted with a single substituent, that substituent is at the 4-position;
- ii) a heteroaryl selected from the group consisting of pyridinyl, pyrimidinyl, pyridazinyl, and pyrazinyl; wherein said heteroaryl is optionally independently substituted with one or two substituents that are chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, or C₁₋₄alkoxy;
- iii) pyrimidin-5-ylmethyl;
- iv) phenylmethyl, wherein the phenyl portion of phenylmethyl is optionally independently substituted with one or two substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, and C₁₋₄alkoxy; provided that when phenylmethyl of group (iv) is substituted with a single substituent, that substituent is at the 4-position;
- v) phenylsulfonyl, wherein the phenyl portion of phenylsulfonyl is optionally independently substituted with one or two substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋

alkyl, and C₁₋₄alkoxy; provided that when phenylsulfonyl of group (v) is substituted with a single substituent, that substituent is at the 4-position;


- vi) C₁₋₄alkylsulfonyl;
- vii) C₃₋₇cycloalkylsulfonyl; or
- viii) trifluoromethylsulfonyl;

R² is

- i) phenyl optionally substituted with a substituent that is selected from the group consisting of C₁₋₄alkoxy and trifluoromethoxy;
- ii) a heteroaryl selected from the group consisting of pyridinyl, pyrimidinyl, thiazolyl, triazolyl, and pyrazinyl; wherein said heteroaryl is optionally substituted with a substituent that is C₁₋₄alkyl, C₁₋₄alkoxy, trifluoromethoxy, or hydroxy;
- iii) C₃₋₇cycloalkyl; or
- iv) C₃₋₇cycloalkyl-(C₁₋₂)alkyl;

R³ is selected from the group consisting of hydrogen, chloro, or methyl;

G is G1, G2, or G3,



and enantiomers, diastereomers, and pharmaceutically acceptable salts thereof.

2. The compound of claim 1 wherein R¹ is

- i) phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, difluoromethoxy, and C₁₋₄alkoxy; provided that

when phenyl of group (i) is substituted with a single substituent, that substituent is at the 4-position;

- ii) a heteroaryl that is pyridinyl; wherein said pyridinyl is optionally independently substituted with one or two substituents that are chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, or C₁₋₄alkoxy;
- iii) phenylmethyl, wherein the phenyl portion of phenylmethyl is optionally independently substituted with one or two substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, and C₁₋₄alkoxy; provided that when phenylmethyl of group (iii) is substituted with a single substituent, that substituent is at the 4-position;
- iv) phenylsulfonyl, wherein the phenyl portion of phenylsulfonyl is optionally independently substituted with one or two substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, and C₁₋₄alkoxy; provided that when phenylsulfonyl of group (iv) is substituted with a single substituent, that substituent is at the 4-position;
- v) C₁₋₄alkylsulfonyl; or
- vi) trifluoromethylsulfonyl.


3. The compound of claim 2 wherein R¹ is

- i) phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, methyl, difluoromethoxy, and C₁₋₂alkoxy; provided that when phenyl of group (i) is substituted with a single substituent, that substituent is at the 4-position; or
- ii) a heteroaryl that is pyridinyl; wherein said pyridinyl is optionally independently substituted with one or two substituents that are chloro, fluoro, bromo, cyano, trifluoromethyl, or C₁₋₄alkoxy.


4. The compound of claim 3 wherein R^1 is phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, cyano, trifluoromethyl, difluoromethoxy, and methyl; provided that when phenyl is substituted with a single substituent, that substituent is at the 4-position.
5. The compound of claim 1 wherein R^2 is
 - i) phenyl substituted with a substituent that is selected from the group consisting of C_{1-4} alkoxy and trifluoromethoxy;
 - ii) a heteroaryl selected from the group consisting of pyridinyl, pyrimidinyl, and pyrazinyl; wherein said heteroaryl is optionally substituted with a substituent that is C_{1-4} alkoxy or trifluoromethoxy;
 - iii) C_{3-7} cycloalkyl; or
 - iv) C_{3-7} cycloalkyl- (C_{1-2}) alkyl.
6. The compound of claim 5 wherein R^2 is
 - i) phenyl substituted with C_{1-4} alkoxy;
 - ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl; wherein said heteroaryl is optionally substituted with C_{1-4} alkoxy; or
 - iii) C_{3-7} cycloalkyl- (C_{1-2}) alkyl.
7. The compound of claim 6 wherein R^2 is
 - i) phenyl substituted with C_{1-4} alkoxy; or
 - ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl; wherein said heteroaryl is optionally substituted with C_{1-4} alkoxy.
8. The compound of claim 7 wherein R^2 is
 - i) phenyl substituted with methoxy; or
 - ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl; wherein said heteroaryl is optionally substituted with C_{1-2} alkoxy.

9. The compound of claim 1 wherein R^3 is hydrogen.


10. The compound of claim 1 wherein G is G1 or G2;




11. The compound of claim 10 wherein G is G1;



12. The compound of claim 11 wherein G is G2;



13. A compound of Formula (I)



Formula (I)

wherein

R^1 is

- i) phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano,

trifluoromethyl, C₁₋₄alkyl, difluoromethoxy, and C₁₋₄alkoxy; provided that when phenyl of group (i) is substituted with a single substituent, that substituent is at the 4-position;


- ii) a heteroaryl that is pyridinyl; wherein said pyridinyl is optionally independently substituted with one or two substituents that are chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, or C₁₋₄alkoxy;
- iii) phenylmethyl, wherein the phenyl portion of phenylmethyl is optionally independently substituted with one or two substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, and C₁₋₄alkoxy; provided that when phenylmethyl of group (iii) is substituted with a single substituent, that substituent is at the 4-position;
- iv) phenylsulfonyl, wherein the phenyl portion of phenylsulfonyl is optionally independently substituted with one or two substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, C₁₋₄alkyl, and C₁₋₄alkoxy; provided that when phenylsulfonyl of group (iv) is substituted with a single substituent, that substituent is at the 4-position;
- v) C₁₋₄alkylsulfonyl; or
- vi) trifluoromethylsulfonyl;

R² is

- i) phenyl substituted with a substituent that is selected from the group consisting of C₁₋₄alkoxy and trifluoromethoxy;
- ii) a heteroaryl selected from the group consisting of pyridinyl, pyrimidinyl, and pyrazinyl; wherein said heteroaryl is optionally substituted with a substituent that is C₁₋₄alkoxy or trifluoromethoxy;
- iii) C₃₋₇cycloalkyl; or
- iv) C₃₋₇cycloalkyl-(C₁₋₂)alkyl;


R³ is selected from the group consisting of hydrogen, chloro, or methyl;

G is G1 or G2;



and enantiomers, diastereomers, and pharmaceutically acceptable salt forms thereof.

14. A compound of Formula (I)



Formula (I)

wherein

R^1 is

- i) phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, bromo, cyano, trifluoromethyl, methyl, difluoromethoxy, and C_{1-2} alkoxy; provided that when phenyl of group (i) is substituted with a single substituent, that substituent is at the 4-position; or
- ii) a heteroaryl that is pyridinyl; wherein said pyridinyl is optionally independently substituted with one or two substituents that are chloro, fluoro, bromo, cyano, trifluoromethyl, or C_{1-4} alkoxy;


R^2 is

- i) phenyl substituted with C_{1-4} alkoxy;

- ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl; wherein said heteroaryl is optionally substituted with C₁₋₄alkoxy; or
- iii) C₃₋₇cycloalkyl-(C₁₋₂)alkyl;


R³ is hydrogen;

G is G1 or G2;



and enantiomers, diastereomers, and pharmaceutically acceptable salt forms thereof.

15. A compound of Formula (I)



Formula (I)

wherein


R¹ is phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, cyano, trifluoromethyl, difluoromethoxy, and methyl; provided that when phenyl is substituted with a single substituent, that substituent is at the 4-position;

R² is

- i) phenyl substituted with C₁₋₄alkoxy; or
- ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl; wherein said heteroaryl is optionally substituted with C₁₋₄alkoxy;


R^3 is hydrogen;

G is G1 or G2;



and enantiomers, diastereomers, and pharmaceutically acceptable salt forms thereof.

16. A compound of Formula (I)



Formula (I)

wherein


R^1 is phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, cyano, trifluoromethyl, difluoromethoxy, and methyl; provided that when phenyl is substituted with a single substituent, that substituent is at the 4-position;

R^2 is

- i) phenyl substituted with methoxy; or
- ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl; wherein said heteroaryl is optionally substituted with C_{1-2} alkoxy;


R^3 is hydrogen;

G is G1 or G2;



and enantiomers, diastereomers, and pharmaceutically acceptable salt forms thereof.

17. A compound of Formula (I)



Formula (I)

wherein


R^1 is phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, cyano, trifluoromethyl, difluoromethoxy, and methyl; provided that when phenyl is substituted with a single substituent, that substituent is at the 4-position;

R^2 is

- i) phenyl substituted with methoxy; or
- ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl; wherein said heteroaryl is optionally substituted with C_{1-2} alkoxy;


R^3 is hydrogen;

G is G1;



and enantiomers, diastereomers, and pharmaceutically acceptable salt forms thereof.

18. A compound of Formula (I)



Formula (I)

wherein


R¹ is phenyl independently substituted with one to three substituents that are selected from the group consisting of chloro, fluoro, cyano, trifluoromethyl, difluoromethoxy, and methyl; provided that when phenyl is substituted with a single substituent, that substituent is at the 4-position;

R² is

- i) phenyl substituted with methoxy; or
- ii) a heteroaryl selected from the group consisting of pyridinyl and pyrazinyl; wherein said heteroaryl is optionally substituted with C₁₋₂alkoxy;

R³ is hydrogen;


G is G2;



G2 ;

and enantiomers, diastereomers, and pharmaceutically acceptable salt forms thereof.

19. A compound of Formula (I)



Formula (I)

selected from the group consisting of

- Cpd 1**, 4-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 2**, 4-[1-(4-Fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyridine;
- Cpd 3**, 2-Ethoxy-5-[2-(3-methoxypyridin-4-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]pyridine;
- Cpd 4**, 1-(4-Fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 5**, 5-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-methylpyridine;
- Cpd 6**, 2-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-5-methylpyridine;
- Cpd 7**, 2-(2-Methoxyphenyl)-1-(4-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 8**, 5-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-methylpyrimidine;
- Cpd 9**, 3-(1-(3-chlorobenzyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine;
- Cpd 10**, 3-[1-(4-Fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 11**, 3-[1-(4-Chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;

- Cpd 12**, 4-[2-(3-Methoxypyridin-2-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 13**, 4-[1-(4-Chloro-3-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyridine;
- Cpd 14**, 2-Chloro-5-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 15**, 4-[2-(2-Methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-methylbenzonitrile;
- Cpd 16**, 2-methoxy-3-(4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1-(2,3,4-trifluorophenyl)-1H-imidazol-2-yl)pyridine;
- Cpd 17**, 2-[1-(3-Chloro-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyridine;
- Cpd 18**, 4-[1-(3-Chloro-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyridine;
- Cpd 19**, 2-[1-(4-Chloro-3-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyridine;
- Cpd 20**, 3-[1-(3-Chloro-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-4-methoxypyridine;
- Cpd 21**, 4-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-methylbenzonitrile;
- Cpd 22**, 5-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-methylbenzonitrile;
- Cpd 23**, 3-[1-(3-Fluoro-4-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 24**, 3-[1-(4-Fluoro-3-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 25**, 5-[2-(2-Methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-methylbenzonitrile;

- Cpd 26**, 3-[1-(4-Chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-4-methyl-4H-1,2,4-triazole;
- Cpd 27**, 3-[1-(2-Chlorobenzyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 28**, 3-[1-(4-Chlorobenzyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 29**, 4-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-(trifluoromethyl)benzonitrile;
- Cpd 30**, 3-[1-(2,4-Difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 31**, 3-(1-(2,3-difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine;
- Cpd 32**, 3-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 33**, 4-[1-(3,4-Difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyridine;
- Cpd 34**, 2-Methoxy-4-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 35**, 2-[1-(3,4-Difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyridine;
- Cpd 36**, 3-Fluoro-4-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 37**, 2-Methoxy-5-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 38**, 1-(4-Chlorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 39**, 5-Chloro-2-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]pyridine;

- Cpd 40**, 3-[1-(3,4-Difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 41**, 1-(4-Chloro-3-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 42**, 1-(3,4-Difluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 43**, 4-[2-(2-Methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 44**, 3-[1-(4-Chloro-3-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 45**, 1-(3-Chloro-4-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 46**, 3-[1-(3-Chloro-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 47**, 2-[1-(4-Fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyrazine;
- Cpd 48**, 3-[2-(2-Methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 49**, 5-[1-(4-Chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-4-methoxypyrimidine;
- Cpd 50**, 2-[1-(4-Chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-methoxypyrazine;
- Cpd 51**, 3-[1-(4-Chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-ethoxypyridine;
- Cpd 52**, 3-[1-(3,4-Difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-ethoxypyridine;
- Cpd 53**, 5-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-(trifluoromethyl)pyridine;

- Cpd 54**, 2-Methoxy-3-{4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1-[6-(trifluoromethyl)pyridin-3-yl]-1H-imidazol-2-yl}pyridine;
- Cpd 55**, 1-(4-Chlorophenyl)-2-(cyclopropylmethyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 56**, 1-(4-Chlorophenyl)-2-cyclopropyl-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 57**, 4-[2-Cyclopropyl-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 58**, 4-[2-(Cyclopropylmethyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 59**, 5-[2-(Cyclopropylmethyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-(trifluoromethyl)pyridine;
- Cpd 60**, 2-[1-(4-Chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-3-ethoxypyrazine;
- Cpd 61**, 1-(4-(difluoromethoxy)phenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 62**, 1-(4-Bromo-3-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 63**, 1-(3-Bromo-4-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 64**, 3-[1-(3-Bromo-4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 65**, 3-[1-(4-Bromo-3-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 66**, 4-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine;
- Cpd 67**, 2-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine;

- Cpd 68**, 3-(1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-4-methoxypyridine;
- Cpd 69**, 3-(1-(4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-4-methoxypyridine;
- Cpd 70**, 2-(1-(4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-3-methoxypyridine;
- Cpd 71**, 3-(1-(2,5-difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine;
- Cpd 72**, 3-(1-(3,5-difluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl)-2-methoxypyridine;
- Cpd 73**, 3-fluoro-5-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile;
- Cpd 74**, 2-Fluoro-5-[2-(2-methoxy-phenyl)-4-(2,2,6,6-tetramethyl-tetrahydro-pyran-4-yl)-imidazol-1-yl]-benzonitrile;
- Cpd 75**, 3-[1-(6-Ethoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 76**, 3-Fluoro-4-[2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 77**, 2-Methoxy-4-[2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 78**, 2-Chloro-4-[2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 79**, 2-Fluoro-5-[2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 80**, 2-Fluoro-4-[2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 81**, 2-Fluoro-4-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;

- Cpd 82**, 2,6-difluoro-4-(2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile;
- Cpd 83**, 3,5-difluoro-4-(2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile;
- Cpd 84**, 2-Chloro-5-[2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 85**, 2-Methoxy-5-[2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 86**, 4-[2-(4-Methoxypyrimidin-5-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 87**, 4-[2-(3-Methoxypyrazin-2-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 88**, 2-fluoro-4-(2-(3-methoxypyrazin-2-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)benzonitrile;
- Cpd 89**, 4-(2-(2-ethoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl)-2-fluorobenzonitrile;
- Cpd 90**, 1-(4-chlorophenyl)-2-(2-methoxyphenyl)-5-methyl-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 91**, 4-[5-Chloro-2-(2-methoxypyridin-3-yl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 92**, 3-[5-Chloro-1-(4-fluorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 93**, 3-[5-Chloro-1-(4-chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]-2-methoxypyridine;
- Cpd 94**, 2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1-[(trifluoromethyl)sulfonyl]-1H-imidazole;
- Cpd 95**, 1-(Cyclopropylsulfonyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;

- Cpd 96**, 2-(2-Methoxyphenyl)-1-[(2-methylpropyl)sulfonyl]-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 97**, 1-[(4-Chlorophenyl)sulfonyl]-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 98**, 1-[(4-Fluorophenyl)sulfonyl]-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazole;
- Cpd 99**, 4-{[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]methyl} benzonitrile;
- Cpd 100**, 5-{[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]methyl} pyrimidine;
- Cpd 101**, 3-{[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]methyl} benzonitrile;
- Cpd 102**, 4-[1-(4-Chlorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 103**, 4-[1-(4-Chlorophenyl)-2-(4-methoxypyridin-3-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 104**, 4-[1-(4-Chlorophenyl)-2-(3-methoxypyridin-2-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 105**, 4-[1-(4-Chlorophenyl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 106**, 4-[1-(3-Bromo-4-fluorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 107**, 4-[1-(3,4-Difluorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 108**, 4-[1-(3,4-Difluorophenyl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 109**, 4-{2-(2-Methoxyphenyl)-1-[6-(trifluoromethyl)pyridin-3-yl]-1H-imidazol-4-yl}-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;

- Cpd 110**, 4-[1-(3-Chloro-4-fluorophenyl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 111**, 4-[1-(3-Chloro-4-fluorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 112**, 4-[1-(4-Chloro-3-fluorophenyl)-2-(2-methoxyphenyl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 113**, 4-[1-(4-Chloro-3-fluorophenyl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 114**, 4-[1-(4-Chlorophenyl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 115**, 4-[4-(4-Hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 116**, 2-Fluoro-5-[4-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 117**, 2-Fluoro-5-[4-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-2-(2-methoxyphenyl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 118**, 2-Fluoro-4-[4-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-2-(2-methoxyphenyl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 119**, 2-Fluoro-4-[4-(4-hydroxy-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-2-(2-methoxypyridin-3-yl)-1H-imidazol-1-yl]benzonitrile;
- Cpd 120**, 4-[1-(4-Chlorophenyl)-2-(3-methoxypyridin-4-yl)-1H-imidazol-4-yl]-2,2,6,6-tetramethyltetrahydro-2H-pyran-4-ol;
- Cpd 121**, 4-[2-(2-Methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]-2-(trifluoromethyl)benzonitrile;
- Cpd 122**, 3-[1-(4-Chlorophenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-2-yl]pyridin-2-ol;
- Cpd 123**, 2-Chloro-4-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyltetrahydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;

Cpd 124, 1-(4-Chlorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyl-3,6-dihydro-2H-pyran-4-yl)-1H-imidazole;

Cpd 125, 1-(4-Bromo-3-fluorophenyl)-2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyl-3,6-dihydro-2H-pyran-4-yl)-1H-imidazole;

Cpd 126, 2-Fluoro-4-[2-(2-methoxyphenyl)-4-(2,2,6,6-tetramethyl-3,6-dihydro-2H-pyran-4-yl)-1H-imidazol-1-yl]benzonitrile;

and pharmaceutically acceptable salt forms thereof.

20. A pharmaceutical composition comprising a compound of claim 1 or 19 and at least one of a pharmaceutically acceptable carrier, a pharmaceutically acceptable excipient, and a pharmaceutically acceptable diluent.
21. The pharmaceutical composition of claim 20, wherein the composition is a solid oral dosage form.
22. The pharmaceutical composition of claim 20, wherein the composition is a syrup, an elixir or a suspension.
23. A method for treating inflammatory pain in a subject in need thereof comprising administering to the subject a therapeutically effective amount of a compound of claim 1 or 19.
24. The method of claim 23 wherein the inflammatory pain is due to inflammatory bowel disease, irritable bowel syndrome, visceral pain, migraine, post-operative pain, osteoarthritis, rheumatoid arthritis, back pain, lower back pain, joint pain, abdominal pain, chest pain, labor pain, musculoskeletal diseases, skin diseases, toothache, pyresis, burn, sunburn, snake bite, venomous snake bite, spider bite, insect sting, neurogenic/overactive bladder, interstitial cystitis, urinary tract infection, rhinitis, contact dermatitis/hypersensitivity, itch, eczema, pharyngitis,

mucositis, enteritis, irritable bowel syndrome, cholecystitis, pancreatitis, postmastectomy pain syndrome, menstrual pain, endometriosis, pain, pain due to physical trauma, headache, sinus headache, tension headache or arachnoiditis.

25. A method for treating neuropathic pain in a subject in need thereof comprising administering to the subject a therapeutically effective amount of a compound of claim 1 or 19.
26. The method of claim 25 wherein the neuropathic pain is cancer pain, neurological disorders, spine and peripheral nerve surgery, brain tumor, traumatic brain injury (TBI), chemotherapy-induced pain, pain chronification, radicular pain, HIV pain, spinal cord trauma, chronic pain syndrome, fibromyalgia, chronic fatigue syndrome, lupus, sarcoidosis, peripheral neuropathy, bilateral peripheral neuropathy, diabetic neuropathy, central pain, neuropathies associated with spinal cord injury, stroke, amyotrophic lateral sclerosis (ALS), Parkinson's disease, multiple sclerosis, sciatic neuritis, mandibular joint neuralgia, peripheral neuritis, polyneuritis, stump pain, phantom limb pain, bony fractures, oral neuropathic pain, Charcot's pain, complex regional pain syndrome I and II (CRPS I/II), radiculopathy, Guillain-Barre syndrome, meralgia paresthetica, burning-mouth syndrome, optic neuritis, postfebrile neuritis, migrating neuritis, segmental neuritis, Gombault's neuritis, neuronitis, cervicobrachial neuralgia, cranial neuralgia, geniculate neuralgia, glossopharyngeal neuralgia, migrainous neuralgia, idiopathic neuralgia, intercostals neuralgia, mammary neuralgia, Morton's neuralgia, nasociliary neuralgia, occipital neuralgia, postherpetic neuralgia, causalgia, red neuralgia, Sluder's neuralgia, splenopalatine neuralgia, supraorbital neuralgia, trigeminal neuralgia, vulvodynia, or vidian neuralgia.