

DESCRIPTION

METHOD FOR ANALYZING OLIGOMERIC PROANTHOCYANIDIN (OPC)

TECHNICAL FIELD

This invention relates to a method for determining
5 oligomeric proanthocyanidin (generic name for a mixture of
n-polymers of flavan-3-ol: $n \geq 2$) contained in analytes or
samples to be analyzed, such as naturally occurring
substances, foods and drinks, pharmaceuticals and/or
cosmetics.

10 BACKGROUND ART

Proanthocyanidin (OPC) is said to be one of the
efficacious components of the French Paradox, since it is
contained in wine as well (1995, Clin. Chim. Acta. 235,
207-219). An antioxidant action, a peripheral circulation
15 improving action, a blood flow improving effect, a hepatic
function improving effect (2004, Japan Food Science, 403,
January Issue, 40-45), and a platelet aggregation
suppressing effect (Officially Published Patent Gazette
2003-527418) are known as the medicinal benefits of
20 proanthocyanidin. Development of a method for convenient
qualitative and quantitative evaluation of OPC, which is
such an active ingredient, is therefore desired.

Known methods for analyzing proanthocyanidin include
reversed phase HPLC by high-performance liquid
25 chromatography-mass spectrometry (LC-MS) (2003, Biosci.
Biotechnol. Biochem., 67, (5), 1140-1142), and normal phase
HPLC involving gradient elution by LC-MS (2003, J. Agric.
Food Chem., 51, 7513-7521). However, both of these methods

require MS detectors, and none of them are said to be convenient. Furthermore, proanthocyanidin is present as very many stereoisomers, owing to the stereoisomerism of flavan-3-ols which are the constituents of proanthocyanidin.

5 There are limits on the compounds available as standard substances. Thus, its quantitative analysis has been impossible, except for some known compounds. Besides, proanthocyanidin exists in the natural world in forms ranging from the monomer flavan-3-ol to a dimer, a trimer, 10 and further to n-mers of a higher polymerization degree. The analysis by reversed phase HPLC has shown that peaks of flavan-3-ol (monomer), the dimer and the trimer overlap.

As described above, no convenient method has been existent for the qualitative and quantitative evaluation of 15 OPC.

DISCLOSURE OF THE INVENTION

Under these circumstances, the inventors have focused attention on the desire for the development of a novel method of analysis which can determine the abundance 20 ratios and contents of n-mers contained in natural substances, foods and beverages, and pharmaceutical products, without interference from flavan-3-ol which is the monomer. It is an object of the present invention, therefore, to provide a novel method for assaying OPC which 25 is contained in analytes, such as natural substances, foods and beverages.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a view showing the results of analysis of

catechin, and a dimer and a trimer of flavan-3-ol by normal phase HPLC.

FIG. 2 is a view showing the results of analysis of analytes, which have catechin and proanthocyanidin mixed
5 therein, by normal phase HPLC.

EFFECTS OF THE INVENTION

The inventors conducted various studies in an attempt to solve the aforementioned problems. As a result, they have elucidated, from the aspects of experimentation
10 and molecular weight, that (a) the number of anthocyanidins produced by hydrolysis of oligomeric proanthocyanidin (OPC) is always $n/2$, regardless of the magnitude of the polymerization degree, n , of flavan-3-ol contained in OPC. This has led them to work out a convenient method of OPC
15 assay. The inventors have further found the unexpected fact that (b) the separation of flavan-3-ol polymers of different polymerization degrees n , which have hitherto been difficult to separate by chromatography, can be easily achieved by high performance liquid chromatography using a
20 normal phase column.

The method of the present invention uses the above fact (a) and/or the fact (b) to analyze the amount of oligomeric proanthocyanidin (OPC) in natural substances, foods and beverages, pharmaceuticals and/or cosmetics, and
25 also analyze the proportions and/or contents of n -polymers in OPC in a convenient manner.

According to the method of the present invention, oligomeric proanthocyanidin (OPC) can be assayed without

interference from the flavan-3-ol monomer often contained in oligomeric proanthocyanidin.


Also, the method of the present invention can assay flavan-3-ol polymers for respective polymers of oligomeric proanthocyanidin in analytes, without requiring a mass spectrometer. Thus, this method is very suitable for the analysis of oligomeric proanthocyanidin in natural substances, foods and beverages, pharmaceuticals and/or cosmetics.

10 MODES FOR CARRYING OUT THE INVENTION

The analytes targeted by the method of the present invention are arbitrary samples, which are expected to contain a mixture of n-polymers ($n \geq 2$) of flavan-3-ol (hereinafter referred to as oligomeric proanthocyanidin or OPC), such as natural substances (grape seeds, tamarind, apple, bark, pine bark-derived polyphenol, tea leaves, cocoa, etc., and/or their treatment products (extract, etc.)), foods and beverages, pharmaceuticals and/or cosmetics.

Oligomeric proanthocyanidin, typically, includes at least one of compounds represented by the following general formulas:


[Chemical formulas 1]




The present invention targets, particularly, the above oligomeric proanthocyanidin which is

proanthocyanidins including at least one of procyanidins B1, B2, B3 and B4 of the structural formula 1 where $n = 0$, procyanidins B5, B6, B7 and B8 of the structural formula 2 where $n_1 = 1$ and $n_2 = 0$, and procyanidins C1, C2 and C4 of the structural formula 1 where $n = 1$. The above B1, B2 ... C1, C2 ... are stereoisomers of the respective compounds. The structures of the main procyanidins, OPC and catechin are illustrated below.


[Chemical formulas 1a]




Catechin




PB7(cate-4 α ,6-cate)




PB1 (epi-4 β ,8-cate)



PB2 (epi-4 β ,8-epi)



PB3 (cate-4 α ,8-cate)



PB5 (epi-4 β ,6-epi)

(1) Determination of total amount of OPC

According to the present invention, there is no need to consider the types and polymerization degrees of flavan-3-ol polymers contained in oligomeric proanthocyanidin

(OPC) in analytes and, when anthocyanidin occurring upon hydrolysis of OPC is assayed, the total amount of OPC can be determined from its value. The method of the present invention can be performed in the following sequence:

5 Hydrolysis of OPC

 The hydrolysis of oligomeric proanthocyanidin can be performed by thermal decomposition under acidic conditions. The preferred conditions include, for example, the use of an acid/lower alcohol mixture as acidic conditions, the
10 acid being preferably hydrochloric acid, sulfuric acid or nitric acid, and the lower alcohol being preferably propanol, butanol, pentanol and/or isopentanol. The thermal decomposition can be performed, with the temperature being 50 to 100°C, preferably 80 to 100°C, more
15 preferably 85 to 95°C, and the reaction time being 30 minutes or more, preferably 1 hour or more. The concentration of the acid can be chosen from the range of 0.1N to 2N, preferably 0.4N to 1N.

 In performing hydrolysis, if the analyte is a liquid
20 sample containing oligomeric proanthocyanidin in a low concentration, the sample may be concentrated by a suitable method, for example, lyophilization or drying or solidification to dryness under reduced pressure.

 The analyte (including the concentrate) is dissolved
25 in the above-mentioned acid/lower alcohol mixture at a concentration of 0.01 to 1%, preferably, 0.05 to 0.2% (0.5 to 2 mg/ml), whereby it can be hydrolyzed.

 A concrete example of the hydrolysis is as follows:

A sample (0.5 mg) containing proanthocyanidin is dissolved in 1 ml of 0.6N HCl/butanol in a glass test tube, and the solution is allowed to stand for 2 hours in a water bath at 90°C. After completion of the reaction, the absorption spectrum at 700 to 400 nm is measured with UV-265 (Shimadzu Corp.). Measurement of the absorbance at 551 nm confirms whether the hydrolysis reaction has been fully carried out to produce anthocyanidin.

Measurement of amount of anthocyanidin

10 The measurement of the amount of anthocyanidin produced by hydrolysis may be made by any publicly known method and, for example, can be performed easily by high performance liquid chromatography or the absorbance method. The absorbance method can be performed using the hydrolyzate unchanged and, preferably, measures the absorbance at 550 to 552 nm at which anthocyanidin shows a maximum absorption in the visible absorption spectrum. At this wavelength, the influence of components other than anthocyanidin produced by the hydrolysis of oligomeric proanthocyanidin (e.g., catechin: maximum absorption 270 nm) can be disregarded.

 An example of the HPLC process is concretely described as follows: Anthocyanidin is assayed using an HPCL column YMC-ODS-A312 (6 mm x 150 mm, YMC), 1 ml/min of a mixture acetic acid:methanol:water = 15:17.5:67.5, and a column temperature of 40°C, with an area value at A520 nm being detected. Cyanidin chloride, delphinidin chloride, and pelargonidin chloride (all available from Funakoshi)

can be used as standard substances. The retention times of these standard substances are all different, and a mixture of them or any one of them selected can be used as a standard substance. As the HPLC column, not only the above
5 C18-based resin, but also the C8-, C30-, or polymer-based C18 resin can make analysis similarly, as long as it is a reversed phase resin.

Calculation of total amount of OPC

To determine the amount of oligomeric
10 proanthocyanidin in the analyte based on the amount of anthocyanidin assayed above, the following procedure is effected: a) The amount of anthocyanidin produced from one n-polymer by hydrolysis is $n/2$, regardless of the magnitude of the polymerization degree n of the flavan-3-ol polymer
15 contained in OPC. Based on this fact, the amount of anthocyanidin produced by hydrolysis is multiplied by a factor 2, whereby the total amount of OPC in a predetermined amount of the analyte can be found, for example, in mg. Alternatively, b) the amount of
20 anthocyanidin produced by hydrolysis of OPC is in proportion to the total amount of OPC, regardless of the magnitude of the polymerization degree n of the flavan-3-ol polymer contained in OPC. Based on this fact, the amount of anthocyanidin produced by hydrolysis is compared with
25 the amount of anthocyanidin produced by hydrolysis of a known amount of an OPC standard substance, and the total amount (e.g., in mg) of OPC in the analyte and/or the proportion (a%) of OPC in the analyte can be determined by

a calibration curve or the following equation:

Total amount of OPC (mg) = (measured value of sample/measured value of standard) x amount (mg) of standard (Equation 1)

5 Proportion (a%) of OPC in analyte = (total amount (mg) of OPC/amount of analyte) x 100 (Equation 2)

The standard substance used can be selected, as appropriate, from procyanidin B1 and procyanidin B2 (both available from Funakoshi).

10 (2) Assay of each of n-polymers constituting OPC

In particularly preferred embodiments of the present invention, in addition to or separately from the determination of the total weight of OPC in the predetermined weight of the analyte in (1) above, the proportions of various flavan-3-ol polymers in OPC are clarified by high performance liquid chromatography (HPLC). By so doing, there is also provided a novel method of OPC assay which determines the weights of respective polymers of different polymerization degrees, n, contained in the predetermined weight of the analyte.

15

20

The column used in the high performance liquid chromatography is preferably a normal phase column, and particularly preferably a normal phase column packed with a silica gel-based resin. Studies by the inventors have shown that proanthocyanidin having different polymerization degrees (in this case, n = 1 is also included) can be separated into constituents very satisfactorily by chromatography on a normal phase column. Judging from the

25

fact that flavan-3-ols have high polarity, it has not been considered so far that they can be separated by a normal phase column. Furthermore, the method of the present invention enables measurement with the use of an
5 ultraviolet detector, and does not require a mass spectrometric detector.

The conditions for high performance liquid chromatography may be determined as desired. If they are concretely illustrated, Inertsil SIL (4.6 mm ϕ x 150 mm, GL
10 Sciences Inc.), for example, is used as the column, the eluant is, for example, a mixture of hexane, methanol, tetrahydrofuran, and formic acid. Preferably, isocratic elution (about 1 ml/min) with hexane:MeOH:THF:HCOOH = 45:40:14:1 is performed. Analysis can be made even at a
15 flow velocity of 0.3 to 1.5 ml/min with the use of hexane:MeOH:THF:HCOOH = 40-60:30-50:10-20:0.1-5. The column temperature is 10 to 60°C, preferably 40°C, and the detector is preferably a photodiode array detector, which is used in the collection of spectrum data at 240 to 400 nm.
20 This is because OPC has a maximum absorption at 280 nm, but in the case of a sample incorporating other polyphenols in mixed form, different peaks at wavelengths other than 280 nm constitute the maximum absorption, thus making it possible to distinguish the other polyphenols from OPC and
25 exclude them. However, in an environment where only a detector of a single wavelength can be used, analysis can be made only with the absorption at A280 nm.

The total peak area of all polymers separated by

chromatography (all polymers including a dimer or polymers of higher degrees of polymerization) is the sum of the area values at A280 nm of peaks having the maximum absorption at 280 nm. The column may be Shimpack PREP-SIL(H) (4.6 mmφ x 300 mm, Shimadzu Corp.) or Supersher Si60 (4.5 mmφ x 100 mm, Merck & Co.) as well as Inertsil SIL (4.6 mmφ x 150 mm, GL Sciences Inc.).

The proportion of each n-polymer constituting OPC (b%) can be calculated from the following equation based on the peak area obtained upon high performance liquid chromatography:

$$b\% = \{\text{peak area (each n-polymer)}\} / \{\text{total peak area (all polymers)}\} \times 100 \dots (\text{Equation 3})$$

Based on the so obtained proportion of each n-polymer and the weight of OPC obtained in the aforementioned (1), the weight of each n-polymer contained in the predetermined weight of the analyte can be determined, for example, in mg by the following equation:

$$\text{Amount of n-polymer (mg)} = \text{total amount of OPC (mg)} \times \text{proportion of n-polymer (b\%)} / 100 \dots (\text{Equation 4})$$

Next, the present invention will be described more concretely by Examples. However, the present invention is not limited to these Examples.

Example 1 Investigation of acid hydrolysis conditions

The decomposition of flavangenol by acid was observed over time. Flavangenol (1 mg) was dissolved in 1 ml of 0.6N HCl/BuOH, and the solution was heated in a hot water bath at 90°C. After a lapse of 20 minutes until

140 minutes later, sampling was done at intervals of 20 minutes. The sample taken was diluted 1:10 with butanol, and measured for the visible absorption spectrum at 400 to 700 nm. The maximum absorption was present at 551 nm for all samples. The results of the measurements are shown in Table 1.

Table 1 Changes over time in acid-decomposed flavangenol

Time	Absorbance (551 nm)
20 min	0.732
40 min	0.796
60 min	0.819
80 min	0.830
100 min	0.867
120 min	0.877
140 min	0.823

The outcome was that the absorbance increased slowly with the passage of time during acid decomposition, and reached its peak in 120 minutes. Based on this outcome, the period of hydrolysis was set at 120 min (2 hours) in subsequent experiments.

Example 2 Production and assay of anthocyanidin upon acid hydrolysis

A sample (0.5 mg) containing proanthocyanidin was dissolved in 1 ml of 0.6N HCl/butanol in a glass test tube, and the solution was allowed to stand for 2 hours in a water bath at 90°C. After completion of the reaction, the absorption spectrum at 700 to 400 nm was measured with UV-265 (Shimadzu Corp.), and the absorbance at 551 nm was

determined. The solution after completion of the reaction was subjected to HPLC under the following conditions to assay anthocyanidin:

Column: YMC-ODS-A312, 6 mm ϕ x 150 mm

5 Mobile phase: CH₃COOH:MeOH:H₂O = 15:17.5:67.5

Detection: A520 nm (measured at 400 to 600 nm by PDA)

As standard substances for assay, delphinidin, cyanidin and pelargonidin were purchased from Funakoshi.

10 The standard substance, delphinidin, was eluted in 4.2 minutes, with $\lambda_{\max}$ of 535 nm, cyanidin was eluted in 5.5 minutes, with $\lambda_{\max}$ of 525 nm, and pelargonidin was eluted in 8.0 minutes, with $\lambda_{\max}$ of 515 nm. The components from the acid hydrolyzate of the sample, corresponding to these
15 conditions, were assayed as delphinidin, cyanidin and pelargonidin.

The samples analyzed were flavangenol, grape seed polyphenol, tea polyphenol, apple polyphenol, and tamarind, and procyanidin B1 (Funakoshi) was used as the standard
20 substance for OPC.

The results are shown in Tables 2 and 3.

Table 2 OPC content by absorbance method

	A550 nm	OPC content
Procyanidin B1	1.6905	100.0%
Flavangenol	0.8897	52.6%
Apple	0.3804	22.5%
Grape seeds	0.7523	44.5%
Tamarind	0.7641	45.2%

Table 3 OPC content by HPLC method

Sample	Cyanidin	OPC content
	µg/ml	
Procyanidin B1	66.19	100.0%
Flavangenol	29.45	44.5%
Tamarind	28.64	43.3%
Apple	13.48	20.4%
Grape seeds	27.81	42.0%
Green tea	1.84+1.98*	5.8%

*: cyanidin 1.84 µg/ml + delphinidin 1.98 µg/ml

Example 3 Analysis of catechin, dimer and trimer by 5 normal phase HPLC

The standard substances for catechin and proanthocyanidin were analyzed by normal phase HPLC under the following conditions:

Samples (0.1 mg each) of (+)-catechin (Nacalai
10 Tesque), (-)-epicatechin (Wako Pure Chemical Industries), and procyanidin B1 (Funakoshi) were each dissolved in 1 ml of a mobile phase, and the solution was filtered through a 0.45 µm filter, and then subjected to HPLC under the conditions shown below.

15 The trimer was synthesized by the method described in Example 5 to be offered below.

Column: Inertsil SIL, 4.6 mmφ x 150 mm

Mobile phase: hexane:MeOH:THF:HCOOH = 45:40:14:1

Detection: A280 nm (measured at 240 to 400 nm by

20 PDA)

Under these conditions, the monomers ((+)-catechin and (-)-epicatechin) were eluted in 2.9 minutes, the dimer in 3.6 minutes, and the trimer in 4.3 minutes.

Their chromatograms are shown in FIG. 1.

5 Example 4 Analysis by normal phase HPLC

Samples containing catechin and proanthocyanidin in mixed form were analyzed by normal phase HPLC under the conditions shown below.

The sample (1 to 2 mg) containing proanthocyanidin
10 was dissolved in 1 ml of a mobile phase, and the solution was filtered through a 0.45 μ m filter, and then subjected to HPLC under the following conditions.

The samples used were apple polyphenol, grape seed polyphenol, flavangenol, and tamarind.

15 Column: Inertsil SIL, 4.6 mm ϕ x 150 mm

Mobile phase: hexane:MeOH:THF:HCOOH = 45:40:14:1

Detection: A280 nm (measured at 240 to 400 nm by PDA)

The chromatograms as analytic patterns are shown in
20 FIG. 2. In these patterns, the symbol $\times$ represents polyphenols having no maximum absorption at 280 nm and different from OPC.

The concentrations of the dimer and the trimer in each sample were determined by the equation $c\% = a\% \times b\%$.

25 The results are shown in Table 4.

Table 4 Contents of dimer and trimer

	OPC purity		A280 nm ratio	Abundance ratio
	a%		b%	c%
Flavangenol	44.5%	Dimer	55.4%	24.7%
	44.5%	Trimer	32.0%	14.2%
Apple polyphenol	20.4%	Dimer	31.6%	6.4%
	20.4%	Trimer	23.5%	4.8%
Grape seed polyphenol	42.0%	Dimer	12.5%	5.3%
	42.0%	Trimer	5.8%	2.4%
Tamarind	43.3%	Dimer	22.8%	9.9%
	43.3%	Trimer	23.2%	10.0%

Example 5 Synthesis of OPC

The synthesis of OPC was performed in the following manner in accordance with a paper (1981, J.C.S. Perkin I. 1235-1245).

(+)-Taxifolin (500 mg) was dissolved in 50 ml of ethanol, and 200 mg of NaBH₄ was added. Then, 1 g of (+)-catechin was added and dissolved. Then, HCl was added, and the mixture was stirred for 1 hour. The reaction product was purified by reversed phase HPLC to obtain a trimer.

CLAIMS

1. A method for determining a total weight of oligomeric proanthocyanidin (OPC) contained in a predetermined weight of an analyte,

said method comprising measuring a weight of anthocyanidin produced by hydrolysis of OPC in the analyte, separately measuring an amount of anthocyanidin produced by hydrolysis of a known weight of an OPC standard substance, and

determining the total weight of OPC in the analyte and/or a weight ratio of OPC (a%) in the analyte by a calibration curve or the following equation

based on a fact that an amount of anthocyanidin produced by hydrolysis of OPC is in proportion to the total amount of OPC, regardless of an amount ratio of n-polymers contained in OPC and different in a magnitude of a polymerization degree n:

Total weight of OPC = (measured value of sample/measured value of standard) x weight of standard substance (Equation 1)

Weight ratio of OPC (a%) in analyte = (total weight of OPC/weight of analyte) x 100 (Equation 2)

2. The method according to Claim 1, wherein the standard substance is procyanidin B1.

3. The method according to any one of Claims 1 to 2, wherein the hydrolysis is performed by thermal decomposition under acidic conditions.

4. The method according to Claim 3, wherein the acidic conditions are a mixture of an acid and a lower alcohol.
5. The method according to Claim 3 or 4, wherein the acid is hydrochloric acid, sulfuric acid, or nitric acid.
6. The method according to Claim 4 or 5, wherein the lower alcohol is propanol, butanol, pentanol and/or isopentanol.
7. The method according to any one of Claims 3 to 6, wherein the thermal decomposition is performed at a temperature of 50 to 100°C, preferably 80 to 100°C, more preferably 85 to 95°C, for a reaction time of 30 minutes or more, preferably 1 hour or more.
8. The method according to any one of Claims 1 to 7, wherein the measurement of the amount of anthocyanidin is made by high performance liquid chromatography and/or an absorbance method.
9. The method according to Claim 8, wherein the absorbance method is performed by measuring absorbance at 550 to 552 nm at which anthocyanidin shows a maximum absorption in a visible absorption spectrum.
10. A method for assaying oligomeric proanthocyanidin (OPC), contained in an analyte, for n-polymers of flavan-3-ol different in a polymerization degree n, and comprising the steps of:

a) determining a total weight of OPC, contained in a predetermined weight of the analyte, by the method of any one of Claims 1 to 9;

b) separately from the step a), analyzing the analyte by high performance liquid chromatography (HPLC) to separate respective n-polymers of different polymerization degrees, n, contained in OPC, and determining quantitative proportions of the respective n-polymers to an amount of all n-polymers, namely, OPC; and

c) determining a weight of each n-polymer in the analyte by results of a) and b).

11. The method according to Claim 10, wherein a column used in the high performance liquid chromatography is a normal phase column.

12. The method according to Claim 11, wherein a resin packed in the normal phase column is a silica gel-based resin.

13. The method according to Claim 11 or 12, wherein, in the high performance liquid chromatography, an eluant is a mixture of hexane, methanol, tetrahydrofuran, and formic acid, a mixing ratio of the mixture is hexane/methanol/tetrahydrofuran/formic acid = 40-60:30-50:10-20:0.1-5, preferably, a ratio of 45:40:14:1 in isocratic elution, a column temperature is 10 to 60°C, preferably 40°C, and an absorbance at 280 nm is measured.

14. The method according to any one of Claims 10 to 13, wherein the quantitative proportion of each n-polymer (b%) is calculated from the following equation based on a peak area obtained upon the high performance liquid chromatography:

$$b\% = \{\text{peak area (each n-polymer)}\} / \{\text{total peak area (all polymers)}\} \times 100 \dots (\text{Equation 3})$$


15. The claim according to Claim 10, wherein the weight of each n-polymer in the step (c) is determined by the following equation based on the weight of OPC determined by (a) and the proportion of each n-polymer determined by (b):

$$\text{Amount of n-polymer(mg)} = \text{total weight of OPC(mg)} \times \text{proportion of n-polymer (b\%)/100} \dots (\text{Equation 4})$$


16. The method according to any one of Claims 1 to 15, wherein the oligomeric proanthocyanidin to be assayed contains at least one of compounds represented by the following general formulas:

[Chemical formulas 1]

Structural formula 2




Structural formula 4




n = integer of 0 to 18,
 $n1$ = integer of 1 to 18,
 $n2$ = integer of 0 to 18, and
 R = H or OH.

Structural formula 1



Structural formula 3



17. The method according to Claim 16, herein the oligomeric proanthocyanidin is proanthocyanidins containing

at least one of procyanidins B1, B2, B3 and B4 of Structural Formula 1 where $n = 0$, procyanidins B5, B6, B7 and B8 of Structural Formula 2 where $n_1 = 1$ and $n_2 = 0$, and procyanidins C1, C2 and C4 of Structural Formula 1 where $n = 1$.

18. The method according to any one of Claims 1 to 17, wherein the analyte is a natural substance (grape seeds, tamarind, apple, bark, pine bark-derived polyphenol, tea leaves, cocoa, or the like, and/or a treatment product thereof (extract or the like)), a food or a beverage, a pharmaceutical and/or a cosmetic.

19. A method for determining proportions of n -polymers of different polymerization degrees of oligomeric proanthocyanidin (OPC), contained in an analyte, by high performance liquid chromatography (HPLC).

20. The method according to Claim 19, wherein a column used in the high performance liquid chromatography is a normal phase column.

21. The method according to Claim 19 or 20, wherein a resin packed in the normal phase column is a silica gel-based resin.

22. The method according to any one of Claims 19 to 21, wherein, in the high performance liquid chromatography, an eluant is a mixture of hexane, methanol, tetrahydrofuran, and formic acid, a mixing ratio of the mixture is hexane/methanol/tetrahydrofuran/formic acid = 40-60:30-50:10-20:0.1-5, preferably, a ratio of 45:40:14:1 in isocratic elution, a column temperature is 10 to 60°C, preferably 40°C, and an absorbance at 280 nm is measured.

23. The method according to any one of Claims 19 to 22, wherein the proportion of each n-polymer (b%) is calculated from the following equation based on a peak area obtained upon the high performance liquid chromatography:

$$b\% = \{\text{peak area (each n-polymer)}\} / \{\text{total peak area (all polymers)}\} \times 100 \dots \text{(Equation 3)}$$

24. The method according to any one of Claims 1 to 23, further comprising a step of lyophilizing the analyte, or solidifying the analyte to dryness under reduced pressure, followed by powdering, if the analyte is a liquid including a soft drink or a refreshing drink, a tea beverage, or an alcoholic beverage.