

PHARMACEUTICAL COMPOSITIONS COMPRISING 2,3,5-TRIMETHYL-6-NONYLCYCLOHEXA-2,5-DIENE-1,4-DIONE

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 63/219,784, filed July 8, 2021, the contents of which is herein incorporated by reference in its entirety.

FIELD

[0002] Provided herein is pharmaceutical composition of 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione (**Compound 1**), and/or the hydroquinone form thereof, wherein **Compound 1** is optionally provided as a hydrate, and/or solvate thereof; and methods useful for treating or suppressing a disease or disorder such as an α -synucleinopathy, a tauopathy, an autistic spectrum disorder, a pervasive developmental disorder, a liver disease, liver damage, dementia, and reperfusion injury in a subject using such pharmaceutical compositions.

BACKGROUND

[0003] U.S. Publication No. 2007/0072943 describes certain quinone compounds, pharmaceutical compositions, and methods of treating certain mitochondrial disorders. U.S. Publication No. 2010/0063161 describes the compound 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione, pharmaceutical compositions, and methods for treating pervasive developmental disorders and Attention Deficit Hyperactivity Disorder (ADHD). US Publication No. US Publication No. 2020/0121618 describes the compound 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione, compositions and methods useful for treating or protecting biological systems against damage caused by inflammation and/or oxidative stress. US Patent No. 11,174,212 B2 describes a polymorphic form of the compound 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione, compositions and methods useful for treating or suppressing a disorder selected from the group consisting of α -synucleinopathies, tauopathies, ALS, traumatic brain injury, and reperfusion injury.

[0004] Self-emulsifying drug delivery systems (SEDDS), are lipid-based formulations mixtures of oils, surfactants, co-surfactants, and a solubilized compound. SEDDS are an important way to improve the oral absorption of highly lipophilic compounds with poor aqueous solubility. Following oral administration, SEDDS spontaneously emulsify and rapidly

disperse in gastrointestinal fluids, yielding micro- or nanoemulsions containing the solubilized compound.


[0005] What are needed are pharmaceutical compositions comprising 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione, for example, a SEDDS formulation. Pharmaceutical compositions provided herein show good mitigation of observed positive food effect, good bioavailability, no birefringence when stored, little or no degradation of 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione during storage, and/or little or no reduction in potency.

SUMMARY

[0006] In a first aspect provided herein is a pharmaceutical composition comprising:

a) about 48 wt/wt% to about 70 wt/wt% of a pharmaceutically acceptable oil, about 12 wt/wt% to about 25 wt/wt% of a propylene glycol laurates composition, and about 8 wt/wt% to about 20 wt/wt% of a polysorbate 80;

b) about 1 wt/wt% to about 15 wt/wt% of **Compound 1**:



and/or the hydroquinone form thereof; wherein **Compound 1**, and/or the hydroquinone thereof is optionally a hydrate thereof, and/or solvate thereof; and wherein when **Compound 1**, and/or hydroquinone thereof, is in the form of a hydrate and/or solvate, then the about 1 wt/wt% to about 15 wt/wt% of **Compound 1**, and/or its hydroquinone, does not include the weight of the water in the hydrate or the weight of the solvent in the solvate; and

c) 0 wt/wt% to about 2% wt/wt% of an optional flavorant;

wherein the wt/wt% of **Compound 1**, the pharmaceutically acceptable oil, the propylene glycol laurates composition, the polysorbate 80, and optional flavorant total 100%.

[0007] In a second aspect provided herein is a method of preparing a pharmaceutical formulation, comprising:

step a) mixing about 48 wt/wt% to about 70 wt/wt% of the pharmaceutically acceptable oil, about 12 wt/wt% to about 25 wt/wt% of the propylene glycol laurates composition, and about 8 wt/wt% to about 20 wt/wt% of the polysorbate 80 composition, and 0 wt/wt% to about 2% wt/wt% of an optional flavorant;

step b) adding about 1 wt/wt% to about 15%, of **Compound 1** and/or the hydroquinone form thereof, wherein **Compound 1** and/or the hydroquinone form thereof is optionally a hydrate and/or solvate thereof; to the mixture from step a) and mixing,


and wherein when the **Compound 1**, and/or hydroquinone thereof, is in the form of a hydrate and/or solvate, then the about 1 wt/wt% to about 15 wt/wt% of **Compound 1** and/or its hydroquinone, does not include the weight of any the water in the hydrate or the weight of the solvent in the solvate; and

wherein the wt/wt% of **Compound 1**, the pharmaceutically acceptable oil, the propylene glycol laurates composition, the polysorbate 80, and optional flavorant total 100%.

[0008] In a third aspect, provided herein are methods of using the pharmaceutical composition or pharmaceutical compositions described herein. In some embodiments, the methods are useful for treating or suppressing a disease or disorder such as an α -synucleinopathy, a tauopathy, an autistic spectrum disorder, a pervasive developmental disorder, a liver disease, liver damage, dementia, and reperfusion injury in a subject using such pharmaceutical compositions.

[0009] In a fourth aspect provided herein is a pharmaceutical composition comprising:

a) about 85 wt/wt% to about 99 wt/wt% of a pharmaceutical acceptable oil;



b) about 1 wt/wt% to about 15% of **Compound 1**:

and/or the hydroquinone form thereof; wherein **Compound 1**, and/or the hydroquinone thereof is optionally a hydrate thereof, and/or solvate thereof; and wherein when **Compound 1**, and/or hydroquinone thereof, is in the form of a hydrate and/or solvate, then the about 1 wt/wt% to about 15 wt/wt% of **Compound 1**, and/or its hydroquinone, does not include the weight of the water in the hydrate or the weight of the solvent in the solvate; and


c) 0 wt/wt% to about 2% wt/wt% of an optional flavorant;

wherein the wt/wt% of **Compound 1**, the pharmaceutically acceptable oil, and optional flavorant total 100%.

[0010] In a fifth aspect, provided herein is a pharmaceutical composition comprising:

a) about 48 wt/wt% to about 70 wt/wt% of a pharmaceutically acceptable oil, about 12 wt/wt% to about 25 wt/wt% of a co-surfactant, and about 8 wt/wt% to about 20 wt/wt% of a surfactant;

b) about 1 wt/wt% to about 15 wt/wt% of **Compound 1**:



and/or the hydroquinone form thereof; wherein **Compound 1**, and/or the hydroquinone thereof is optionally a hydrate thereof, and/or solvate thereof; and wherein when **Compound 1**, and/or hydroquinone thereof, is in the form of a hydrate and/or solvate, then the about 1 wt/wt% to about 15 wt/wt% of **Compound 1**, and/or its hydroquinone, does not include the weight of the water in the hydrate or the weight of the solvent in the solvate; and


c) 0 wt/wt% to about 2% wt/wt% of an optional flavorant;

wherein the wt/wt% of **Compound 1**, the pharmaceutically acceptable oil, the surfactant, the co-surfactant, and the optional flavorant total 100%.

[0011] In a sixth aspect, provided herein is a pharmaceutical composition comprising:

a) about 65 wt/wt% to about 80 wt/wt% of a Gelucire-like oil, about 8 wt/wt% to about 10 wt/wt% of an excipient like TPGS, and about 8 wt/wt% to about 10 wt/wt% of a polysorbate 80;

b) about 1 wt/wt% to about 17 wt/wt% or about 1 wt/wt% to about 13 wt/wt% or about



1 wt/wt% to about 10 wt/wt% of **Compound 1**: and/or the hydroquinone form thereof; wherein **Compound 1**, and/or the hydroquinone thereof is optionally a hydrate thereof, and/or solvate thereof; and wherein when **Compound 1**, and/or hydroquinone thereof, is in the form of a hydrate and/or solvate, then the about 1 wt/wt% to about 15 wt/wt% of **Compound 1**, and/or its hydroquinone, does not include the weight of the water in the hydrate or the weight of the solvent in the solvate; and

c) 0 wt/wt% to about 2% wt/wt% of an optional flavorant;

wherein the wt/wt% of **Compound 1**, the pharmaceutically acceptable oil, the surfactant, the co-surfactant, and the optional flavorant total 100%.

BRIEF DESCRIPTION OF THE DRAWINGS

[0012] Fig. 1 shows the physical stability under polarized microscope for Gelucire 44/14-based formulation of **Compound 1** as described in Example 4.

[0013] Fig. 2 shows the physical stability under polarized microscope for Labrafac WL 1349-based formulation of **Compound 1** as described in Example 4.

[0014] Fig. 3 shows monkey PK profiles for formulations in monkeys as described in Example 10.

[0015] Fig. 4a monkey PK profiles for SEDDS Formulations 1-6 (all formulations first 8 hrs, Linear scale) as described in Example 10.

[0016] Fig. 4b shows monkey PK profiles for SEDDS Formulations 1-6 (all formulations, Linear scale) as described in Example 10.

[0017] Fig. 4c shows monkey PK profiles for SEDDS Formulations 1-6 as described in Example 10.

[0018] Fig. 5a shows monkey PK parameter comparison of the SEDDS Formulations 1-6 - $T_{1/2}$ as described in Example 10.

[0019] Fig. 5b shows monkey PK parameter comparison of the SEDDS Formulations 1-6 - C_{Max} as described in Example 10.

[0020] Fig. 5c shows monkey PK parameter comparison of the SEDDS Formulations 1-6 - AUC_{Last} as described in Example 10.

[0021] Fig. 6 shows moisture abstractability of the liquid formulation or its components in hard gelatin capsule as described in Example 11.

DETAILED DESCRIPTION

[0022] The present disclosure provides a pharmaceutical composition of **Compound 1**, and/or the hydroquinone form thereof, and methods useful for treating or suppressing a disease or disorder such as an α -synucleinopathy, a tauopathy, an autistic spectrum disorder, a pervasive developmental disorder, a liver disease, liver damage, dementia, and reperfusion injury in a subject using such pharmaceutical compositions.

Definitions

[0023] When referring to the compounds provided herein, the following terms have the following meanings unless indicated otherwise. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art. In the event that there is a plurality of definitions for a term herein, those in this section prevail unless stated otherwise.

[0024] Use of the singular herein includes the plural and vice versa unless expressly stated to be otherwise, or obvious from the context that such is not intended. That is, “a” and “the” refer to one or more of whatever the word modifies. For example, “a therapeutic agent” includes one therapeutic agent, two therapeutic agent s, etc. Likewise, “a capsule” may refer to one, two or more capsules, and “the disease” may mean one disease or a plurality of diseases. By the same token, words such as, without limitation, “capsules” and “therapeutic agents” would refer to one capsule or therapeutic agent as well as to a plurality of capsules or therapeutic agents, unless, again, it is expressly stated or obvious from the context that such is not intended.


[0025] As used herein, unless specifically defined otherwise, any words of approximation such as without limitation, “about,” “essentially,” “substantially,” when used in connection with various terms such as temperatures, doses, amounts, or weight percent of ingredients of a composition or a dosage form, mean e.g. a temperature, dose, amount, or weight percent that is recognized by those of ordinary skill in the art to provide an effect equivalent to that obtained from the specified temperature dose, amount, or weight percent. Specifically, the terms “about” and “approximately,” when used in this context, contemplate a temperature, dose, amount, or weight percent, etc. within 15%, within 10%, within 5%, within 4%, within 3%, within 2%, within 1%, or within 0.5% of the specified temperature, dose, amount, or weight percent, etc. In addition, any of the embodiments herein, where ranges or numbers are expressed with “about,” i.e. “about 48 wt/wt% to about 70 wt/wt%,” can be replaced with a range or number that does not recite “about,” i.e. “48 wt/wt% to 70 wt/wt%.”

[0026] As used herein, any ranges presented are inclusive of the end-points. For example, “a temperature between 10 °C and 30 °C” or “a temperature from 10 °C to 30 °C” includes 10 °C and 30 °C, as well as any specific temperature in between. Similarly, a temperature of 20 °C ± 10 °C would cover the same range as “a temperature between 10 °C and 30 °C.”

[0027] “Alpha-synuclein” and “α-synuclein” are used interchangeably herein.

[0028] As used herein “birefringent crystals” refers to non-cubic crystals.

[0029] “Hydroquinone form” indicates the form of the compound when a two electron reduction of the quinone ring is effected, providing a net conversion of the two oxo groups to two hydroxy groups. For example, the hydroquinone form of the quinone compound:



[0030] As used herein, “initial potency” is measured on the day the pharmaceutical composition is prepared and before it is stored on the same day as it is prepared.

[0031] As used herein, “initial total impurity level” is measured on the day the pharmaceutical composition is prepared and before it is stored, on the same day as it is prepared.

[0032] As used herein, “initial degradation level” is measured on the day the pharmaceutical composition is prepared and before it is stored on the same day as it is prepared.

[0033] As used herein, “Gelucire-like oil” refers to a composition comprising a small fraction of mono, di- and triglycerides and mainly PEG-32 (MW 1500) mono and diesters of lauric acid (C_{12}). In some or any embodiments, the Gelucire-like oil refers to a composition consisting of a small fraction of mono, di- and triglycerides and mainly PEG-32 (MW 1500) mono and diesters of lauric acid (C_{12}). In some or any embodiments, the Gelucire-like oil is Gelucire 44/14.

[0034] As used herein “Labrafac-like oil” refers to an oil of medium chain triglyceride of fractionated vegetable C_8 and C_{10} fatty acids (mainly fractionated coconut oil or palm kernel oil) with an HLB of 1. In some embodiments, Labrafac brand oil can be replaced with another oil comprising medium-chain triglycerides. In some or any embodiments, Labrafac-like oil means Labrafac WL 1349 (alternatively named Labrafac Lipophile WL 1349) or Labrafac PG. In some or any embodiments, Labrafac-like oil means Labrafac WL 1349. In some or any embodiments, Labrafac-like oil means Labrafac PG.

[0035] As used herein “propylene glycol laurates composition” refers to a composition comprising propylene glycol mono- and di-esters of lauric acid (C_{12}), mainly composed of monoesters and a small fraction of di-esters with HLB value 3 and can be used as a nonionic water-insoluble co-surfactant. In some embodiments, the propylene glycol laurates

composition consists of propylene glycol mono-and di-esters of lauric acid (C₁₂), mainly composed of monoesters and a small fraction of di-esters. In some embodiments, the propylene glycol laurates composition is Lauroglycol 90. In other embodiments, the skilled person would recognize that Lauroglycol 90 can be replaced with another propylene glycol laurates composition, including, for example Capmul PG-12.

[0036] As used herein “polysorbate 80,” refers to polyoxyethylene sorbitan monooleate containing 20 units of oxyethylene, which are hydrophilic, hygroscopic, nonionic surfactants that can be used as an emulsifying agent. In some embodiments, the polysorbate is Tween 80. In other embodiments, the skilled person would recognize that Tween 80 can be replaced with another polysorbate 80.

[0037] As used herein “Low-fat food for Testing” refers, in some embodiments, to food having 11.4% fat (by calories); in some embodiments, 15% fat (by calories). One example is provided in Example 10. A preparation of a “Low-fat food for Testing” is provided in Example 13a. “Low-fat food for Testing” may be provided as a single food item or may be in the form a meal.

[0038] As used herein “Medium-fat food for Testing” refers, in some embodiments, to food having 37.9% fat (by calories); in some embodiments, 38% fat (by calories). One example is provided in Example 10. A preparation of a “Medium-fat food for Testing” is provided in Example 13b. “Medium-fat food for Testing” may be provided as a single food item or may be in the form a meal.

[0039] As used herein “Low-fat food” refers to food having 25% fat or less (by calories), about 100-125 kcal or less (based on a 400-500 kcal total meal), in some embodiments 11.4% fat (by calories); in some embodiments, 15% fat (by calories). “Low-fat food” may be provided as a single food item or may be in the form a meal.

[0040] As used herein “Medium-fat food” refers to food having 25-50% fat (by calories), in some embodiments, 37.9% fat (by calories); in some embodiments, 38% fat (by calories). One example is provided in paragraph [00213] which immediately follows Table 22. Another example is provided in Example 13b. “Medium-fat food” may be provided as a single food item or may be in the form a meal.

[0041] As used herein “High-fat food” refers to food having 50% fat or more (by calories), about 500-600 kcal or more (based on a 800-1000 kcal total meal). “High-fat food” may be provided as a single food item or may be in the form a meal.

[0042] The term “pharmaceutically acceptable oil” as used herein is a GMP grade oil that can be consumed orally by humans. In some embodiments, the pharmaceutically acceptable oil comprises medium-chain triglycerides. In some embodiments, the pharmaceutically acceptable oil is one that solubilizes **Compound 1** at room temperature. In some embodiments, the pharmaceutically acceptable oil is selected from one or more of the group consisting of: a Labrafac-like oil (including Labrafac WL 1349 and Labrafac PG), sesame oil, cottonseed oil, soybean oil, olive oil, and corn oil. In some embodiments, the pharmaceutically acceptable oil is selected from one or more of the group consisting of: a Labrafac-like oil (including Labrafac WL 1349 and Labrafac PG), cottonseed oil, soybean oil, olive oil, and corn oil. In some embodiments, the pharmaceutically acceptable oil is selected from one or more of the group consisting of: a Labrafac-like oil (including Labrafac WL 1349 and Labrafac PG), sesame oil, cottonseed oil, soybean oil, and corn oil. In some embodiments, the pharmaceutically acceptable oil is selected from one or more of the group consisting of: a Labrafac-like oil (including Labrafac WL 1349 and Labrafac PG), cottonseed oil, soybean oil, and corn oil. In some or any embodiments, one pharmaceutically acceptable oil is selected.

[0043] As used herein, potency = $(100\% - \text{total impurities by HPLC}) \times (100\% - \text{water content}\% - \text{total residual solvent}\% - \text{Residue on ignition}\%)$. Potency may be calculated as follows $(\% \text{ area purity by HPLC}/100) \times (100 - \% \text{wt/wt water content (KF)} - \% \text{wt/wt residual solvents} - \% \text{wt/wt residue on ignition (ROI)})$. In various embodiments, the potency of the 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione is at least about 95%, at least about 96%, at least about 97%, at least about 98%, at least about 99%, or at least about 99.5%.

[0044] As used herein “solvate” refers to a compound described herein that further includes a stoichiometric or non-stoichiometric amount of solvent bound by non-covalent intermolecular forces. Where the solvent is water, the solvate is a hydrate.

[0045] As used herein, the term “formulate” refers to the preparation of a drug, *e.g.*, **Compound 1**, in a form suitable for oral or transdermal administration to a mammalian patient, in some embodiments, a human. Thus, “formulation” can include the addition of pharmaceutically acceptable excipients, diluents, or carriers.

[0046] A pharmaceutical composition can comprise a unit dose formulation, where the unit dose is a dose sufficient to have a therapeutic effect.

[0047] By “stable pharmaceutical composition” is meant any pharmaceutical composition having sufficient stability to have utility as a pharmaceutical product. In some embodiments, a

stable pharmaceutical composition has sufficient stability to allow storage at a convenient temperature, in some embodiments, between -20 °C and 40 °C, in some embodiments, about 2 °C to about 30 °C, for a reasonable period of time, *e.g.*, the shelf-life of the product which can be as short as one month but is typically six months or longer, in some embodiments, one year or longer, in some embodiments, twenty-four months or longer, and in some embodiments, thirty-six months or longer. The shelf-life or expiration can be that amount of time where the active ingredient degrades to a point below 90% purity. For purposes of the present description stable pharmaceutical composition includes reference to pharmaceutical compositions with specific ranges of impurities as described herein. In some embodiments,, a stable pharmaceutical composition is one which has minimal degradation of the active ingredient, *e.g.*, it retains at least about 85% of un-degraded active, in some embodiments, at least about 90%, and in some embodiments, at least about 95%, after storage at 2-30 °C for a 2-3 year period of time.

[0048] The term “pharmaceutically acceptable” as used herein means that the thing that is pharmaceutically acceptable, *e.g.*, components, including containers, of a pharmaceutical composition, does not cause unacceptable loss of pharmacological activity or unacceptable adverse side effects. Examples of pharmaceutically acceptable components are provided in The United States Pharmacopeia (USP), The National Formulary (NF), adopted at the United States Pharmacopeial Convention, held in Rockville, Md. in 1990 and FDA Inactive Ingredient Guide 1990, 1996 issued by the U.S. Food and Drug Administration (both are hereby incorporated by reference herein, including any drawings). Other grades of solutions or components that meet necessary limits and/or specifications that are outside of the USP/NF can also be used.

[0049] The term “pharmaceutical composition” as used herein shall mean a composition that is made under conditions such that it is suitable for administration to humans, *e.g.*, it is made under good manufacturing practice (GMP) conditions and contains pharmaceutically acceptable excipients, *e.g.*, without limitation, stabilizers, bulking agents, buffers, carriers, diluents, vehicles, solubilizers, and binders. As used herein “pharmaceutical composition” includes, but is not limited to, a pre-drying solution(s) or dispersion(s) as well as a liquid form ready for injection or infusion after reconstitution of a dry-powder preparation.

[0050] A “pharmaceutical dosage form” as used herein means the pharmaceutical compositions disclosed herein being in a form of a capsule (usually referred to as single oral solid dosage form) or in a container and in an amount suitable for reconstitution and administration of one or more doses, typically about 1-2, 1-3, 1-4, 1-5, 1-6, 1-10, or about 1-

20 doses. The ultimate dosage form can be sterile, fluid and stable under the conditions of manufacture and storage. The prevention of the growth of microorganisms can be accomplished by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, sorbic acid, thimerosal, and the like.

[0051] The single or multiple dosages which can be used include an amount of **Compound 1** independently selected from about 0.1 mg/kg to about 600 mg/kg body weight, or about 1.0 mg/kg to about 500 mg/kg body weight, or about 1.0 mg/kg to about 400 mg/kg body weight, or about 1.0 mg/kg to about 300 mg/kg body weight, or about 1.0 mg/kg to about 200 mg/kg body weight, or about 1.0 mg/kg to about 100 mg/kg body weight, or about 1.0 mg/kg to about 50 mg/kg body weight, or about 1.0 mg/kg to about 30 mg/kg body weight, or about 1.0 mg/kg to about 10 mg/kg body weight, or about 10 mg/kg to about 600 mg/kg body weight, or about 10 mg/kg to about 500 mg/kg body weight, or about 10 mg/kg to about 400 mg/kg body weight, or about 10 mg/kg to about 300 mg/kg body weight, or about 10 mg/kg to about 200 mg/kg body weight, or about 10 mg/kg to about 100 mg/kg body weight, or about 50 mg/kg to about 150 mg/kg body weight, or about 100 mg/kg to about 200 mg/kg body weight, or about 150 mg/kg to about 250 mg/kg body weight, or about 200 mg/kg to about 300 mg/kg body weight, or about 250 mg/kg to about 350 mg/kg body weight, or about 200 mg/kg to about 400 mg/kg body weight, or about 300 mg/kg to about 400 mg/kg body weight, or about 250 mg/kg to about 300 mg/kg body weight, or about 300 mg/kg body weight. In some embodiments, the amount of **Compound 1** is about 90 mg, about 100 mg, about 110 mg, or about 120 mg per dose. In some embodiments, the amount of **Compound 1** is about 100 mg per dose. In some embodiments, the maximum clinical dose is 1000 mg/day. Compounds of the present disclosure may be administered in a single daily dose, or the total daily dosage may be administered in divided dosage of two, three or four times daily.

[0052] Single or multiple doses can be administered. In some embodiments, the dose is administered once, twice, three times, four times, five times, or six times. In some embodiments, the dose is administered once per day, twice per day, three times per day, or four times per day. In some embodiments, the dose is administered every hour, every two hours, every three hours, every four hours, every 6 hours, every 12 hours, or every 24 hours.

[0053] Also provided are articles of manufacture and kits comprising a pharmaceutical composition which comprises **Compound 1** and/or the hydroquinone form thereof, for use in any of the methods described herein.

[0054] As used herein, the term “excipient” means the substances used to formulate active pharmaceutical ingredients (API) into pharmaceutical formulations; in an embodiment, an excipient does not lower or interfere with the primary therapeutic effect of the API. In some embodiments, an excipient is therapeutically inert. The term “excipient” encompasses carriers, diluents, vehicles, solubilizers, stabilizers, bulking agents, and binders. Excipients can also be those substances present in a pharmaceutical formulation as an indirect or unintended result of the manufacturing process. In some embodiments, excipients are approved for or considered to be safe for human and animal administration, *e.g.*, generally regarded as safe (GRAS) substances. GRAS substances are listed by the Food and Drug administration in the Code of Federal Regulations (C.F.R.) at 21 C.F.R. §182 and 21 C.F.R. § 184, incorporated herein by reference. In some embodiments, excipients include, but are not limited to, hexitols, including mannitol and the like.

[0055] As used herein, the terms “subject” and “patient” are used interchangeably herein. The terms “subject” and “subjects” refer to an animal, such as a mammal including a non-primate (*e.g.*, a cow, pig, horse, cat, dog, rat, and mouse) and a primate (*e.g.*, a monkey such as a cynomolgous monkey, a chimpanzee and a human), and for example, a human. In certain embodiments, the subject is a human.

[0056] As used herein, the terms “therapeutic agent” and “therapeutic agents” refer to any agent(s) which can be used in the treatment or prevention of a disease or one or more symptoms thereof. In certain embodiments, the term “therapeutic agent” includes a compound as described herein. In certain embodiments, a therapeutic agent is an agent which is known to be useful for, or has been or is currently being used for the treatment or prevention of a disease or one or more symptoms thereof.

[0057] “Therapeutic use” of the compounds discussed herein is defined as using one or more of the compounds discussed herein to treat or suppress a disorder, as defined herein. A

[0058] As used herein “therapeutically effective amount” of a compound is an amount of the compound, which, when administered to a subject, is sufficient to reduce or eliminate either a disorder or one or more symptoms of a disorder, or to retard the progression of a disorder or of one or more symptoms of a disorder, or to reduce the severity of a disorder or of one or more symptoms of a disorder, or to suppress the clinical manifestation of a disorder, or to suppress the manifestation of adverse symptoms of a disorder. A therapeutically effective amount can be given in one or more administrations.

[0059] As used herein “Treating” or “treatment of” a disorder with the pharmaceutical compositions and methods discussed herein is defined as administering a pharmaceutical composition discussed herein, with or without additional therapeutic agents, in order to reduce or eliminate either the disorder or one or more symptoms of the disorder, or to retard the progression of the disorder or of one or more symptoms of the disorder, or to reduce the severity of the disorder or of one or more symptoms of the disorder. “Suppression” of a disorder with the pharmaceutical compositions and methods discussed herein is defined as administering a pharmaceutical composition discussed herein, with or without additional therapeutic agents, in order to suppress the clinical manifestation of the disorder, or to suppress the manifestation of adverse symptoms of the disorder. The distinction between treatment and suppression is that treatment occurs after adverse symptoms of the disorder are manifest in a subject, while suppression occurs before adverse symptoms of the disorder are manifest in a subject. Suppression may be partial, substantially total, or total. In some embodiments, genetic screening can be used to identify patients at risk of the disorder. The pharmaceutical compositions and methods disclosed herein can then be administered to asymptomatic patients at risk of developing the clinical symptoms of the disorder, in order to suppress the appearance of any adverse symptoms.

[0060] In certain embodiments, the pharmaceutical composition is stable when stored at about 15 °C to about 25 °C and about 60% relative humidity for a duration of time of at least one month, at least two months, at least 3 months, or at least six months. In certain embodiments, the pharmaceutical composition is stable when stored at about 25 °C and about 60% relative humidity for a duration of time of at least one month, at least two months, or at least 3 months. In certain embodiments, the pharmaceutical composition is stable when stored at about 40 °C and about 75% relative humidity for a duration of time of at least one month, at least two months, at least three months, or at least six months. In certain embodiments, the pharmaceutical composition is stable when stored at about 40 °C and about 75% relative humidity for a duration of time of at least one month, at least two months, or at least three months.

[0061] In certain embodiments, the pharmaceutical composition is an oral pharmaceutical composition. In certain embodiments, the pharmaceutical composition is a transdermal pharmaceutical composition.

[0062] In certain embodiments, the pharmaceutical composition is an oral pharmaceutical composition and has bioavailability when administered with a low-fat food which differs from

bioavailability when administered with a medium-fat food by a percentage difference of about 90% or less, about 85% or less, about 80% or less, about 75% or less, about 70% or less, or about 65% or less.

[0063] In certain embodiments, the oral pharmaceutical composition bioavailability for Medium-fat Food for Testing is about 45 to about 55%, in some embodiments, or about 40% to about 50% in some embodiments, for a pharmaceutical formulation made from a vehicle which comprises a): about 60 wt/wt% a pharmaceutically acceptable oil (e.g. Labrafac WL 1349, sesame oil, cottonseed oil, soybean oil, Labrafac PG, olive oil, or corn oil), about 20 wt/wt% propylene glycol laurates composition (e.g. Lauroglycol 90), and about 20 wt/wt% polysorbate 80 (e.g. Tween 80); wherein the % weight of the components in a) total 100%. In certain embodiments, the oral pharmaceutical composition bioavailability for Medium-fat Food for Testing is about 49% to about 50%, in some embodiments, for a pharmaceutical formulation made from a vehicle which comprises in a): about 60 wt/wt% a pharmaceutically acceptable oil (e.g. Labrafac WL 1349, sesame oil, cottonseed oil, soybean oil, Labrafac PG, olive oil, or corn oil), about 20 wt/wt% propylene glycol laurates composition (e.g. Lauroglycol 90), and about 20 wt/wt% polysorbate 80 (e.g. Tween 80); wherein the % weight of the components in a) total 100%. In certain embodiments, the oral pharmaceutical composition bioavailability for Low-fat Food for Testing is about 10 to about 15%, in some embodiments, for a pharmaceutical formulation made from a vehicle which comprises in a): about 60 wt/wt% a pharmaceutically acceptable oil (e.g. Labrafac WL 1349, sesame oil, cottonseed oil, soybean oil, Labrafac PG, olive oil, or corn oil), about 20 wt/wt% propylene glycol laurates composition (e.g. Lauroglycol 90), and about 20 wt/wt% polysorbate 80 (e.g. Tween 80); wherein the % weight of the components in a) total 100%. In certain embodiments, the oral pharmaceutical composition bioavailability for Low-fat Food for Testing is about 13%, in some embodiments, for a pharmaceutical formulation made from a vehicle which comprises in a): about 60 wt/wt% a pharmaceutically acceptable oil (e.g. Labrafac WL 1349, sesame oil, cottonseed oil, soybean oil, Labrafac PG, olive oil, or corn oil), about 20 wt/wt% propylene glycol laurates composition (e.g. Lauroglycol 90), and about 20 wt/wt% polysorbate 80 (e.g. Tween 80); wherein the % weight of the components in a) total 100%.

[0064] In certain embodiments, the oral pharmaceutical composition bioavailability for Medium-fat Food for Testing is about 15% to about 40%, in some embodiments, about 25% to about 35%, in some embodiments, for a pharmaceutical formulation made from a vehicle which comprises in a): about 65 wt/wt% a pharmaceutically acceptable oil (e.g. Labrafac WL 1349,

sesame oil, cottonseed oil, soybean oil, Labrafac PG, olive oil, or corn oil), about 20 wt/wt% propylene glycol laurates composition (e.g. Lauroglycol 90), and about 15 wt/wt% polysorbate 80 (e.g. Tween 80); wherein the % weight of the components in a) total 100%. In certain embodiments, the oral pharmaceutical composition bioavailability for Medium-fat Food for Testing is about 29% for a pharmaceutical formulation made from a vehicle which comprises in a): about 65 wt/wt% a pharmaceutically acceptable oil (e.g. Labrafac WL 1349, sesame oil, cottonseed oil, soybean oil, Labrafac PG, olive oil, or corn oil), about 20 wt/wt% propylene glycol laurates composition (e.g. Lauroglycol 90), and about 15 wt/wt% polysorbate 80 (e.g. Tween 80); wherein the % weight of the components in a) total 100%. In certain embodiments, the oral pharmaceutical composition bioavailability for Low-fat Food for Testing is about 10% to about 15% for a pharmaceutical formulation made from a vehicle which comprises in a): about 65 wt/wt% a pharmaceutically acceptable oil (e.g. Labrafac WL 1349, sesame oil, cottonseed oil, soybean oil, Labrafac PG, olive oil, or corn oil), about 20 wt/wt% propylene glycol laurates composition (e.g. Lauroglycol 90), and about 15 wt/wt% polysorbate 80 (e.g. Tween 80); wherein the % weight of the components in a) total 100%. In certain embodiments, the oral pharmaceutical composition bioavailability for Medium-fat Food for Testing is about 14% for a pharmaceutical formulation made from a vehicle which comprises in a): about 65 wt/wt% a pharmaceutically acceptable oil (e.g. Labrafac WL 1349, sesame oil, cottonseed oil, soybean oil, Labrafac PG, olive oil, or corn oil), about 20 wt/wt% propylene glycol laurates composition (e.g. Lauroglycol 90), and about 15 wt/wt% polysorbate 80 (e.g. Tween 80); wherein the % weight of the components in a) total 100%.

[0065] In certain embodiments, the pharmaceutical composition shows no birefringent crystals when stored at about 15 °C to about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least about one month. In certain embodiments, the pharmaceutical composition shows no birefringent crystals when stored at about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least about one month. In certain embodiments, the pharmaceutical composition showed no birefringent when stored at about 15 °C to about 25 °C and about 60% relative humidity for a duration of time of at least about one month. In certain embodiments, the pharmaceutical composition showed no birefringent when stored at about 25 °C and about 60% relative humidity for a duration of time of at least about one month. In certain embodiments, the pharmaceutical composition showed no birefringent when stored at about 40 °C and about 75% relative humidity for a duration of time of at least about one month, at least

three months, or at least six months. In certain embodiments, the pharmaceutical composition showed no birefringent when stored at about 40 °C and about 75% relative humidity for a duration of time of at least about one month.

[0066] In certain embodiments, the pharmaceutical composition's total impurity level increases by 25% or less from an initial total impurity level to a subsequent total impurity level. In some embodiments, the pharmaceutical composition is stored at about 40 °C and about 75% relative humidity for at least about one month, at least about two months, or at least about three months. In certain embodiments, optionally the initial total impurity level is measured on the day the pharmaceutical composition is prepared and before it is stored (on the same day as it is prepared) at 40 °C and about 75% relative humidity. In some or any embodiments, the subsequent total impurity level is tested after about one month, about two months, or about 3 months in storage. In certain embodiments, the initial total impurity level is about 2% or less, about 1% or less, about 0.5% or less, about 0.4% or less, or about 0.3% or less. In some embodiments, the pharmaceutical composition's total impurity level increases from an initial total impurity level to a subsequent impurity level by 25% or less, 20% or less, 15% or less.

[0067] In certain embodiments, the vehicle a) in the pharmaceutical composition comprises about 60% w/w pharmaceutically acceptable oil (for example, a Labrafac-like oil (Labrafac WL 1349 or Labrafac PG), sesame oil, cottonseed oil, soybean oil, olive oil, or corn oil); about 20 w/w % propylene glycol laurates composition (Lauroglycol 90); and about 20 w/w % polysorbate 80 (Tween 80); wherein the % weight of the components in a) total 100%. In certain embodiments, the vehicle a) in the pharmaceutical composition comprises about 65% w/w pharmaceutically acceptable oil (for example, a Labrafac-like oil (Labrafac WL 1349 or Labrafac PG), sesame oil, cottonseed oil, soybean oil, olive oil, or corn oil); about 20 w/w % propylene glycol laurates composition (Lauroglycol 90); and about 15 w/w % polysorbate 80 (Tween 80); wherein the % weight of the components in a) total 100%.

[0068] In certain embodiments, the pharmaceutical composition comprises about 100 to about 120 mg of **Compound 1**, per g of a)+b)+c). In certain embodiments, the pharmaceutical composition comprises about 100 mg of **Compound 1**, to g of a). In certain embodiments, the pharmaceutical composition comprises about 120 mg of **Compound 1**, to g of a) +b)+c).

[0069] In certain embodiments, **Compound 1** is provided in a capsule and the weight of **Compound 1** in the capsule is selected from the group consisting of about 50 mg and about 60 mg. In certain embodiments, **Compound 1** is provided in a capsule and the weight of

Compound 1 in the capsule is about 50 mg. In certain embodiments, the weight of **Compound 1** in the capsule is about 60 mg.

[0070] In certain embodiments, the pharmaceutical composition shows about 2% or less change in potency from an initial potency after storing at about 15 °C to about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least one month. In certain embodiments, the pharmaceutical composition shows about 2% or less change in potency from an initial potency after storing at about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least one month. In certain embodiments, the pharmaceutical composition shows about 2% or less change in potency from an initial potency after storing at about 15 °C to about 25 °C and about 60% relative humidity for a duration of time of at least one month. In certain embodiments, the pharmaceutical composition shows about 2% or less change in potency from an initial potency after storing at about 25 °C and about 60% relative humidity for a duration of time of at least one month. In certain embodiments, the pharmaceutical composition shows about 2% or less change in potency from an initial potency after storing stored at about 40 °C and about 75% relative humidity for a duration of time of at least one month. In certain embodiments, optionally the initial potency is measured on the day the pharmaceutical composition is prepared and before it is stored on the same day as it is prepared.

[0071] In certain embodiments, the pharmaceutical composition shows about 1% or less degradation from an initial degradation level after storing at about 15 °C to about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least one month. In certain embodiments, the pharmaceutical composition shows about 1% or less degradation from an initial degradation level after storing at about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least one month. In certain embodiments, the pharmaceutical composition shows about 1% or less degradation from an initial degradation level after storing at about 15 °C to about 25 °C and about 60% relative humidity for a duration of time of at least one month. In certain embodiments, the pharmaceutical composition shows about 1% or less degradation from an initial degradation level after storing at about 25 °C and about 60% relative humidity for a duration of time of at least one month. In certain embodiments, the pharmaceutical composition shows about 1% or less degradation from an initial degradation level after storing at about 40 °C and about 75% relative humidity for a duration of time of at least one month. In

certain embodiments, optionally the initial degradation level is measured on the day the pharmaceutical composition is prepared and before it is stored on the same day as it is prepared.

[0072] In certain embodiments, the pharmaceutical composition is in a capsule. In certain embodiments, the pharmaceutical composition is a hard gelatin capsule. In certain embodiments, any water content in the gelatin of the hard gelatin capsule is about 20% or less, about 10% to about 20%, or about 11% to about 16%, when stored for about one month, about one or more months, about three months, or about three or more months.

[0073] In certain embodiments, the oral pharmaceutical composition is a solution. In certain embodiments, the oral pharmaceutical composition comprises about 50 mg to about 120 mg of **Compound 1** per g of a) +b)+c). In certain embodiments, the oral pharmaceutical composition's strength is selected from the group consisting of about 60 mg/mL and about 100 mg/mL of **Compound 1**.

[0074] In certain embodiments, the oral pharmaceutical composition is stable when stored at about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least about one month, at least about three months, or at least about six months. In certain embodiments, the oral pharmaceutical composition is stable when stored at about 5 °C for a period of at least about 4 weeks. In certain embodiments, stability studies are performed following ICH stability guidelines.

[0075] In certain embodiments, the oral pharmaceutical composition comprises an optional flavorant. In certain embodiments, the amount of the optional flavorant is about 0 to about 2% wt/wt%. In certain embodiments, the amount of the optional flavorant is about 0 to about 1 wt/wt%. In certain embodiments, the optional flavorant is less than 1% wt/wt%. In certain embodiments, the flavorant is an oil-soluble flavorant.

[0076] In certain embodiments, the flavorant is selected from the group consisting of natural flavors, natural fruit flavors, artificial flavors, artificial fruit flavors, flavor enhancers and mixtures thereof. In certain embodiments, the flavorant is selected from one or more of natural flavors. In certain embodiments, the flavorant is selected from one or more of natural fruit flavors. In certain embodiments, the flavorant is selected from one or more of artificial flavors. In certain embodiments, the flavorant is selected from one or more of artificial fruit flavors. In certain embodiments, the flavorant is selected from one or more of flavor enhancers.

[0077] In certain embodiments, the flavorant has a flavor selected from the group consisting of raspberries, punch, cherry, strawberries, and blueberries. In certain embodiments,

the flavorant has the flavor of raspberries. In certain embodiments, the flavorant has the flavor of punch. In certain embodiments, the flavorant has the flavor of cherry. In certain embodiments, the flavorant has the flavor of strawberries. In certain embodiments, the flavorant has the flavor of blueberries.


[0078] In certain embodiments, the flavorant has an odor selected from the group consisting of raspberries, punch, cherry, strawberries, and blueberries. In certain embodiments, the flavorant has the odor of raspberries. In certain embodiments, the flavorant has the odor of punch. In certain embodiments, the flavorant has the odor of cherry. In certain embodiments, the flavorant has the odor of strawberries. In certain embodiments, the flavorant has the odor of blueberries.

[0079] In certain embodiments, the pharmaceutically acceptable oil is selected from one or more (in some embodiments, 1, 2, or 3; in some embodiments, 1 or 2; in some embodiments, 1) of the group consisting of: a Labrafac-like oil (including Labrafac WL 1349 and Labrafac PG), sesame oil, cottonseed oil, soybean oil, olive oil, and corn oil. In some embodiments, the pharmaceutically acceptable oil is selected from one or more (in some embodiments, 1, 2, or 3; in some embodiments, 1 or 2; in some embodiments, 1) of the group consisting of: a Labrafac-like oil (including Labrafac WL 1349 and Labrafac PG), cottonseed oil, soybean oil, olive oil, and corn oil. In some embodiments, the pharmaceutically acceptable oil is selected from one or more (in some embodiments, 1, 2, or 3; in some embodiments, 1 or 2; in some embodiments, 1) of the group consisting of: a Labrafac-like oil (including Labrafac WL 1349 and Labrafac PG), sesame oil, cottonseed oil, soybean oil, and corn oil. In some embodiments, the pharmaceutically acceptable oil is selected from one or more (in some embodiments, 1, 2, or 3; in some embodiments, 1 or 2; in some embodiments, 1) of the group consisting of: a Labrafac-like oil (including Labrafac WL 1349 and Labrafac PG), cottonseed oil, soybean oil, and corn oil.

[0080] In some or any embodiments, the pharmaceutically acceptable oil is not a Gelucire-like oil.

[0081] In some or any embodiments of the sixth aspect, the Gelucire-like oil is Gelucire 44/14. In some or any embodiments of the sixth aspect, the ratio of the Gelucire-like oil to a surfactant (e.g. TP GS) to the polysorbate 80 is about 80:10:10.


[0082] In certain embodiments, the pharmaceutical composition comprises b)



Compound 1

, or one or more members selected from the group consisting of hydrates and solvates thereof.


[0083] In certain embodiments, the pharmaceutical composition comprises b)



Compound 1

(not as a hydrate or solvate thereof).


[0084] In certain embodiments, the pharmaceutical composition comprises b), the



Compound 1

hydroquinone form of , or one or more members selected from the group consisting of hydrates and solvates thereof.

[0085] In certain embodiments, the pharmaceutical composition comprises b), the



Compound 1

hydroquinone form of (not as a hydrate or solvate thereof).

[0086] Description of the polymorph, which can be used herein to prepare the pharmaceutical compositions disclosed herein, are provided in PCT Application Publication No. WO 2020/081879, which is hereby incorporated by reference in its entirety.

[0087] Particle size distribution can be determined by laser diffraction, using a Malvern 3000 Mastersizer. Settings are shown in Table A.

Table A. Malvern Instrument Settings

<i>Particle Type</i>	
<i>Non-spherical particle mode</i>	<i>Yes</i>
<i>Is Fraunhofer type</i>	<i>No</i>

<i>Material Properties</i>	
<i>Refractive index</i>	<i>1.480</i>
<i>Absorption index</i>	<i>0.001</i>
<i>Particle density</i>	<i>1.00 g/cm³</i>
<i>Different optical properties in blue light</i>	<i>Yes</i>
<i>Refractive index (in blue light)</i>	<i>1.480</i>
<i>Absorption index (in blue light)</i>	<i>0.001</i>

<i>Dispersant properties</i>	
<i>Dispersant name</i>	<i>Water</i>
<i>Refractive index</i>	<i>1.330</i>
<i>Level sensor threshold</i>	<i>100.000</i>

<i>Measurement Duration</i>	
<i>Background measurement duration (red)</i>	<i>15.00 s</i>
<i>Sample measurement duration (red)</i>	<i>15.00 s</i>
<i>Perform blue light measurement?</i>	<i>Yes</i>
<i>Background measurement duration (blue)</i>	<i>15.00 s</i>
<i>Sample measurement duration (blue)</i>	<i>15.00 s</i>
<i>Assess light background stability</i>	<i>No</i>

<i>Measurement sequence</i>	
<i>Aliquots</i>	<i>1</i>
<i>Automatic number of measurements</i>	<i>No</i>
<i>Pre-alignment delay</i>	<i>0.00 s</i>
<i>Number of measurements</i>	<i>3</i>
<i>Delay between measurements</i>	<i>0.00 s</i>

<i>Pre-measurement delay</i>	<i>0.00 s</i>
<i>Close measurement window after measurement</i>	<i>No</i>
<i>Measurement obscuration settings</i>	
<i>Auto start measurement</i>	<i>No</i>
<i>Obscuration low limit</i>	<i>1.00 %</i>
<i>Obscuration high limit</i>	<i>10.00 %</i>
<i>Enable obscuration filtering</i>	<i>No</i>
<i>Measurement alarms</i>	
<i>Use Background Check</i>	<i>No</i>
<i>Background Check Limits</i>	<i>[1,200],[20,60]</i>

[0088] Particles are generally polydisperse, i.e., not all the same size. One measure of polydispersity is the ratio D90/D10. D10 represents the particle diameter corresponding to 10% cumulative (from 0 to 100%) undersize particle size distribution (i.e. the percentage of particles smaller than D10 is 10%). D90 represents the particle diameter corresponding to 90% cumulative (from 0 to 100%) undersize particle size distribution (i.e. the percentage of particles smaller than D90 is 90%). D90 and D10 are determined by laser diffraction, discussed above, unless expressly stated otherwise.

[0089] In some embodiments, the polymorph, which can be used herein to prepare the pharmaceutical compositions disclosed herein, is according to any one of Embodiments 1-10.

[0090] Embodiment 1A. In certain embodiments, provided is a method, according to the second aspect in the Summary, of preparing a pharmaceutical composition described herein, comprising adding about 1 wt/wt% to about 12 wt/wt%, about 4 wt/wt% to about 12 wt/wt%, about 5 wt/wt% to about 10 wt/wt%, 5 wt/wt%, about 6 wt/wt%, about 7 wt/wt%, about 8 wt/wt%, about 9 wt/wt%, or about 10 wt/wt% **Compound 1** and/or the hydroquinone form thereof per g of a) +b)+c), wherein **Compound 1** and/or the hydroquinone form thereof is optionally a hydrate thereof, and/or solvate thereof; to the mixture from step a)+b)+c).

[0091] Embodiment 1B. In certain embodiments, provided is a method, according to the second aspect in the Summary, of preparing a pharmaceutical composition described herein, comprising adding about 1 wt/wt% to about 12 wt/wt%, about 4 wt/wt% to about 12 wt/wt%,

about 5 wt/wt% to about 10 wt/wt%, 5 wt/wt%, about 6 wt/wt%, about 7 wt/wt%, about 8 wt/wt%, about 9 wt/wt%, or about 10 wt/wt% **Compound 1** and/or the hydroquinone form thereof per g of a) +b)+c), wherein **Compound 1** and/or the hydroquinone form thereof is optionally a hydrate thereof, and/or solvate thereof; to the mixture from step a)+b)+c). In certain embodiments, provided is a method, according to the second aspect in the Summary, of preparing a pharmaceutical composition described herein, wherein step b) occurs at room temperature.

[0092] Embodiment 2. In embodiments of the first aspect in the Summary, the second aspect in the Summary, and Embodiments 1A and 1B, **Compound 1** is a polymorph of an anhydrate of 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione, wherein a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.10, 12.12, and 16.14, and wherein the characteristic peaks are measured using a Cu K α 1 source, and a wavelength of 1.540598 Å. In some embodiments, the characteristic peaks are measured at room temperature, in some embodiments at a temperature of 23-25 °C. In some embodiments, **Compound 1** in step b) is a polymorph of an anhydrate of 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione, wherein a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.10, 12.12, and 16.14, and wherein the characteristic peaks are measured using a Cu K α 1 source, and a wavelength of 1.540598 Å. In some embodiments, the characteristic peaks are measured at room temperature, in some embodiments at a temperature of 23-25 °C.

[0093] Embodiment 2A. The polymorph of Embodiment 2, comprising characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.10, 11.77, 12.12, and 16.14.

[0094] Embodiment 3. The polymorph of Embodiment 2, comprising characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.10, 11.77, 12.12, 16.14, and 22.41.

[0095] Embodiment 4. The polymorph of any one of Embodiments 2-3, wherein the angular positions may vary by ± 0.1 .

[0096] Embodiment 5. The polymorph of any one of Embodiments 2-3, wherein the angular positions may vary by ± 0.05 .

[0097] Embodiment 6. The polymorph of any one of Embodiments 2-5, wherein the polymorph has a powder x-ray diffraction pattern substantially as shown in any one of Figures 5, 11, 14, and 16 as provided in WO 2020/081879.

[0098] Embodiment 7. The polymorph of any one of Embodiments 2-6, having a differential scanning calorimetry (DSC) thermogram substantially as shown in Figure 7 as provided in WO 2020/081879.

[0099] Embodiment 8. The polymorph of any one of Embodiments 2-7, wherein a DSC thermogram has a single endothermic peak at about 47 to about 53°C.

[00100] Embodiment 9. The polymorph of any one of Embodiments 2-8, having a thermogravimetric analysis (TGA) thermogram substantially as shown in Figure 8 as provided in WO 2020/081879.

[00101] Embodiment 10. The polymorph of any one of Embodiments 2-9, having a ¹H NMR spectrum substantially as shown in Figure 6 as provided in WO 2020/081879.

[00102] Embodiment 11. A composition comprising the polymorph of any one of Embodiments 2-10, wherein at least about 95% by mole of the 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione is the polymorph, exclusive of any solvents, carriers or excipients.

[00103] Embodiment 12. A composition comprising the polymorph of any one of Embodiments 2-10, wherein at least about 99% by mole of the 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione is the polymorph, exclusive of any solvents, carriers or excipients.

[00104] Embodiment 13. A composition comprising the polymorph of any one of Embodiments 2-10, or the composition of Embodiment 11 or 12, wherein at least about 95% a/a as measured by HPLC of the composition is the 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione, exclusive of any solvents, carriers or excipients.

[00105] Embodiment 14. A composition comprising the polymorph of any one of Embodiments 2-10, or the composition of Embodiment 11 or 12, wherein at least about 99% a/a as measured by HPLC of the composition is the 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione, exclusive of any solvents, carriers or excipients.

[00106] Embodiment 15. A composition comprising the polymorph of any one of Embodiments 2-10, or the composition of any one of Embodiments 11-14, wherein the potency of the 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione is at least about 95%.

[00107] Embodiment 16. A composition comprising the polymorph of any one of Embodiments 2-10, or the composition of any one of Embodiments 11-14, wherein the potency of the 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione is at least about 99%.

[00108] Embodiment 17. A composition comprising the polymorph of any one of Embodiments 2-10, or the composition of any one of Embodiments 11-16, wherein the polymorph is present as a plurality of particles, wherein the particles have a ratio of D90:D10 less than about 11:1.

[00109] Embodiment 18. A composition comprising the polymorph of any one of Embodiments 2-10, or the composition of any one of Embodiments 11-16, wherein the polymorph is present as a plurality of particles, wherein the particles have a ratio of D90:D10 less than about 7:1.

[00110] Embodiment 19. A composition comprising the polymorph of any one of Embodiments 2-10, or the composition of any one of Embodiments 11-18, wherein the polymorph was recrystallized by a solvent comprising about 75-85% IPA/water.


[00111] Embodiment 20. A composition comprising the polymorph of any one of Embodiments 2-10, or the composition of any one of Embodiments 11-19, wherein the polymorph was recrystallized by a solvent comprising about 80-85% IPA/water.

[00112] Embodiment 21. A composition comprising the polymorph of any one of Embodiments 2-10, or the composition of any one of Embodiments 11-19, wherein the polymorph was recrystallized by a solvent comprising about 85% IPA/water.

Additional Embodiments

[00113] Embodiment 21A. Provided herein is a pharmaceutical composition comprising:

a) 55-75 wt/wt% of Labrafac, 15-25 wt/wt% of Lauroglycol 90, and 10-20 wt/wt% of Tween 80, wherein the wt/wt% of Labrafac, Lauroglycol 90, and Tween 80 total 100%; and



b) about 90 to about 120 mg of **Compound 1**: and/or the hydroquinone form thereof per g of a); wherein the **Compound 1**, and/or the hydroquinone

thereof is optionally a hydrate thereof, and/or solvate thereof; and wherein when **Compound 1**, and/or hydroquinone thereof, is in the form of a hydrate and/or solvate, then the about 90 to about 120 mg weight of **Compound 1**, and/or its hydroquinone does not include the weight of the water in the hydrate or the weight of the solvent in the solvate.

[00114] Embodiment 22. The pharmaceutical composition of embodiment 21A, wherein the pharmaceutical composition is stable when stored at about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least about one month.

[00115] Embodiment 23. The pharmaceutical composition of any one of embodiments 21A-22, wherein the pharmaceutical composition shows no birefringent crystals when stored at about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least about one month.

[00116] Embodiment 24. The pharmaceutical composition of any one of embodiments 21A-23, wherein the pharmaceutical composition shows no birefringent crystals when stored at about 25 °C and about 60% relative humidity for a duration of at least about one month.

[00117] Embodiment 25. The pharmaceutical composition of any one of embodiments 21A-23, wherein the pharmaceutical composition shows no birefringent crystals when stored at about 40 °C and about 75% relative humidity for a duration of at least about one month.

[00118] Embodiment 26. The pharmaceutical composition of any one of embodiments 21A-25, wherein the total impurity level increases by 25% or less from an initial total impurity level to a subsequent total impurity level, when stored at about 40 °C and about 75% relative humidity for at least about three months, optionally wherein the initial total impurity level is measured on the day the pharmaceutical composition is prepared and before it is stored, on the same day as it is prepared, at 40 °C and about 75% relative humidity; and wherein the subsequent total impurity level is tested after about 3 months in storage.

[00119] Embodiment 27. The pharmaceutical composition of embodiment 26, wherein the subsequent total impurity level is about 1% or less, about 0.5% or less, or about 0.4% or less.

[00120] Embodiment 28. The pharmaceutical composition of any one of embodiments 21A-27, wherein a) comprises about 60% w/w Labrafac; about 20% w/w Lauroglycol 90; and about 20% w/w Tween 80.

[00121] Embodiment 29. The pharmaceutical composition of any one of embodiments 21A-27, wherein a) comprises about 65% w/w Labrafac; about 20% w/w Lauroglycol 90; and about 15% w/w Tween 80.

[00122] Embodiment 30. The pharmaceutical composition of any one of embodiments 21A-29, wherein the pharmaceutical composition comprises about 100 mg of **Compound 1** to g of a).

[00123] Embodiment 31. The pharmaceutical composition of any one of embodiments 21A-30, wherein the total weight of the pharmaceutical composition is selected from the group consisting of about 50 mg to about 60 mg.


[00124] Embodiment 32. The pharmaceutical composition of any one of embodiments 21A-31, wherein the total weight of the pharmaceutical composition is about 50 mg.

[00125] Embodiment 33. The pharmaceutical composition of any one of embodiments 21A-31, wherein the pharmaceutical composition is about 60 mg.

[00126] Embodiment 34. The pharmaceutical composition of embodiment 21A-32 or 21A-33, wherein the pharmaceutical composition shows about 2% or less change in potency from an initial potency after storing at about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least one month; optionally wherein the initial potency is measured on the day the pharmaceutical composition is prepared and before it is stored on the same day as it is prepared.

[00127] Embodiment 35. The pharmaceutical composition of embodiment 21A-32 or 21A-33, wherein the pharmaceutical composition shows about 1% or less degradation from an initial degradation level after storing at about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least one month; optionally wherein the initial degradation level is measured on the day the pharmaceutical composition is prepared and before it is stored on the same day as it is prepared.


[00128] Embodiment 36. The pharmaceutical composition of any one of embodiments 1-35, wherein b) is




Compound 1

, or one or more members selected from the group consisting of hydrates thereof and solvates thereof.

[00129] Embodiment 37. The pharmaceutical composition of any one of embodiments 1-36, wherein b) is


**Compound 1**

[00130] Embodiment 38. The pharmaceutical composition of any one of embodiments 1-35, wherein b) is the hydroquinone form of

**Compound 1**

, or one or more members selected from the group consisting of hydrates thereof, and solvates thereof.

[00131] Embodiment 39. The pharmaceutical composition of any one of embodiments 1-35 and 38, wherein b) is the hydroquinone form of

**Compound 1**

[00132] Embodiment 40. The pharmaceutical composition of any one of embodiments 21A-39, wherein the pharmaceutical composition is an oral pharmaceutical composition.

[00133] Embodiment 41. The pharmaceutical composition of embodiment 40, wherein the oral pharmaceutical composition is in a capsule.

[00134] Embodiment 42. The pharmaceutical composition of embodiment 41, wherein the capsule is a hard gelatin capsule.

[00135] Embodiment 43. The pharmaceutical composition of embodiment 42, wherein any water content in the gelatin of the hard gelatin capsule is about 20% or less, about 10% to about 20%, or about 11% to about 16%, when stored for about one month, about one or more months, about three months, or about three or more months.

[00136] Embodiment 44. The pharmaceutical composition of any one of embodiments 40-43, wherein the pharmaceutical composition has bioavailability when administered with a low-fat food which differs from bioavailability when administered with a medium-fat food by a percentage difference of about 90% or less, about 85% or less, about 80% or less, about 75% or less, about 70% or less, or about 65% or less.

[00137] Embodiment 45. The pharmaceutical composition of any one of embodiments 40-43, wherein oral bioavailability for medium-fat food is about 40 to about 50%, in some embodiments, about 49%.

[00138] Embodiment 46. The pharmaceutical composition of embodiments 40-43 and 45, wherein oral bioavailability for low-fat food is about 10% to about 15%, in some embodiments, about 13%.

[00139] Embodiment 47. The pharmaceutical composition of any one of embodiments 20-23, wherein oral bioavailability for medium-fat food is about 25% to about 35%, in some embodiments, about 29%.

[00140] Embodiment 48. The pharmaceutical composition of embodiments 40-43 and 47, wherein oral bioavailability for low-fat food is about 10% to about 15%, in some embodiments, about 14%.

[00141] Embodiment 49. The pharmaceutical composition of any one of embodiments 21A-48, wherein the oral pharmaceutical composition comprises about 100 mg to about 120 mg of **Compound 1** per g of a).

[00142] Embodiment 50. The pharmaceutical composition of any one of embodiments 21A-42, wherein the pharmaceutical composition is a transdermal pharmaceutical composition.

[00143] Embodiment 51. A method of preparing a pharmaceutical composition of any one of embodiments 21A-50, comprising:

step a) mixing 55-75 wt/wt% of Labrafac, 15-25 wt/wt% of Lauroglycol 90, and 10-20 wt/wt% of Tween 80, wherein the wt/wt% of Labrafac, Lauroglycol 90, and Tween 80 total 100%;

step b) adding about 90 mg to about 120 mg of **Compound 1** and/or the hydroquinone form thereof per g of a), wherein **Compound 1** and/or the hydroquinone form thereof is optionally a hydrate and/or solvate thereof; to the mixture from step a) and mixing;

and wherein when the **Compound 1**, and/or hydroquinone thereof, is in the form of a hydrate and/or solvate, then the about 90 to about 120 mg weight of **Compound 1** and/or its hydroquinone, does not include the weight of any the water in the hydrate or the weight of the solvent in the solvate.

[00144] Embodiment 52. The method of embodiment 51, wherein step b) occurs at room temperature.

[00145] Embodiment 53. The method of embodiment 51 or 52, wherein **Compound 1** in step b) is a polymorph of an anhydrate of 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-

dione, wherein a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.10, 12.12, and 16.14, and wherein data are obtained with a Cu K α 1 source, a wavelength of 1.540598 Å, and at a temperature of 23-25 °C.

[00146] Embodiment 54. The method of any one of claims 51-53, wherein **Compound 1** is added and mixed under light protection or yellow light.

[00147] Embodiment 55. A pharmaceutical composition prepared by the method of any one of embodiments 51-54.

[00148] Embodiment 56. A method for treating or suppressing a disease or disorder selected from the group consisting of an α -synucleinopathy, a tauopathy, an autistic spectrum disorder, a pervasive developmental disorder, a liver disease, liver damage, dementia, and reperfusion injury, comprising administering a pharmaceutical composition of any one of embodiments 1-49 or a pharmaceutical composition of embodiment 55.

[00149] Embodiment 57. The method of embodiment 56, wherein the α -synucleinopathy is selected from the group consisting of: Parkinson's Disease (idiopathic and genetic), Parkinson's Disease with dementia (PDD), multisystem atrophy (MSA), Frontotemporal Dementia, Dementia with Lewy Bodies (DLB), Gaucher's disease (GD), Neurodegeneration with Brain Iron Accumulation (NBIA), and neuroaxonal dystrophies (PLA2G6-associated neurodegeneration).

[00150] Embodiment 58. The method of embodiment 57, wherein the Parkinson's Disease is that wherein the patient has a mutation in one or more of the following genes: *MAPT* (Microtubule-associated protein tau), *PRKN* (parkin), *PINK1* (PINK1), LRRK2 (leucine-rich repeat kinase 2), *GBA* (glucocerebrosidase), *SNCA* (alpha synuclein), *PARK7* (DJ-1), and/or *UCHL1* (ubiquitin carboxyl-terminal esterase L1).

[00151] Embodiment 59. The method of embodiment 56, wherein the tauopathy is selected from the group consisting of: Alzheimer's disease, dementia pugilistica, Guam Amyotrophic lateral sclerosis-Parkinsonism-Dementia (Guam ALS/PD), Pick Disease, Argyrophilic grain dementia, Nieman-Pick type C, Subacute sclerosing panencephalitis (SSPE), Progressive supranuclear palsy (PSP), multisystem atrophy (MSA), Corticobasoganglionic degeneration, Frontotemporal dementia with parkinsonism-17 (FTDP-17), Postencephalitic Parkinsonism (PEP), and Autosomal recessive Parkinsonism.

[00152] Embodiment 60. The method of embodiment 56, wherein the liver disease is selected from the group consisting of NASH/NAFL, pediatric NAFLD, alcoholic hepatitis, cholestatic liver disease, viral hepatitis, drug-induced liver toxicity, hemachromatosis,

Wilson's disease, liver transplant reperfusion injury, hepatic insufficiency where the hepatic insufficiency is due to injury, SIRS, sepsis, or severe illness; and drug-induced liver toxicity, such as cisplatin-induced liver toxicity and acetaminophen-induced liver toxicity.

[00153] Embodiment 61. The method of embodiment 56, wherein the autistic spectrum disorder or pervasive developmental disorder is selected from the group consisting of autistic disorder, Asperger's syndrome, childhood disintegrative disorder (CDD), Rett's disorder, PDD-not otherwise specified (PDD-NOS), and attention deficit/hyperactivity disorder (ADHD).

[00154] Embodiment 62. The method of any one of embodiments 56-61, wherein the pharmaceutical composition is administered orally.

[00155] In certain embodiments, provided herein are methods of treating or suppressing a disease or disorder selected from the group consisting of an α -synucleinopathy, a tauopathy, an autistic spectrum disorder, a pervasive developmental disorder, a liver disease, liver damage, dementia, and reperfusion injury, comprising administering an pharmaceutical composition, as provided herein. In certain embodiments, provided herein are methods of treating or suppressing a disease or disorder selected from the group consisting of an α -synucleinopathy, a tauopathy, an autistic spectrum disorder, a pervasive developmental disorder, a liver disease, liver damage, dementia, and reperfusion injury, comprising administering an transdermal pharmaceutical composition, as provided herein.

[00156] In certain embodiments, provided herein are methods of using the pharmaceutical composition for treating or suppressing α -synucleinopathy. In certain embodiments, the α -synucleinopathy is selected from the group consisting of: Parkinson's Disease (idiopathic and genetic), Parkinson's Disease with dementia (PDD), multisystem atrophy (MSA), Frontotemporal Dementia, Dementia with Lewy Bodies (DLB), Gaucher's disease (GD), Neurodegeneration with Brain Iron Accumulation (NBIA), and neuroaxonal dystrophies (PLA2G6-associated neurodegeneration).

[00157] In certain embodiments, the Parkinson's Disease is that wherein the patient has a mutation in one or more of the following genes: *MAPT* (Microtubule-associated protein tau), *PRKN* (parkin), *PINK1* (PINK1), *LRRK2* (leucine-rich repeat kinase 2), *GBA* (glucocerebrosidase), *SNCA* (alpha synuclein), *PARK7* (DJ-1), and/or *UCHL1* (ubiquitin carboxyl-terminal esterase L1).

[00158] In certain embodiments, provided herein are methods of using the pharmaceutical composition for treating or suppressing tauopathy. In certain embodiments, the tauopathy is selected from the group consisting of: Alzheimer's disease, dementia pugilistica, Guam

Amyotrophic lateral sclerosis-Parkinsonism-Dementia (Guam ALS/PD), Pick Disease, Argyrophilic grain dementia, Nieman-Pick type C, Subacute sclerosing panencephalitis (SSPE), Progressive supranuclear palsy (PSP), multisystem atrophy (MSA), Corticobasoganglionic degeneration, Frontotemporal dementia with parkinsonism-17 (FTDP-17), Postencephalitic Parkinsonism (PEP), and Autosomal recessive Parkinsonism.

[00159] In certain embodiments, provided herein are methods of using the pharmaceutical composition for treating or suppressing a liver disease or liver damage. In certain embodiments, the liver disease or liver damage is selected from the group consisting of NASH/NAFL, pediatric NAFLD, alcoholic hepatitis, cholestatic liver disease, viral hepatitis, drug-induced liver toxicity, hemochromatosis, Wilson's disease, liver transplant reperfusion injury, hepatic insufficiency where the hepatic insufficiency is due to injury, SIRS, sepsis, or severe illness; and drug-induced liver toxicity, such as cisplatin-induced liver toxicity and acetaminophen-induced liver toxicity.

[00160] In certain embodiments, provided herein are methods of using the pharmaceutical composition for treating or suppressing autistic spectrum disorder or pervasive developmental disorder. In certain embodiments, the autistic spectrum disorder or pervasive developmental disorder is selected from the group consisting of autistic disorder, Asperger's syndrome, childhood disintegrative disorder (CDD), Rett's disorder, PDD-not otherwise specified (PDD-NOS), and attention deficit/hyperactivity disorder (ADHD).

[00161] In certain embodiments, provided herein are methods for treating or suppressing dementia. In certain embodiments, the dementia is selected from the group consisting of vascular dementia, dementia associated with Huntington's disease, dementia associated with Creutzfeldt-Jakob disease, dementia associated with normal pressure hydrocephalus, dementia associated with Wernicke-Korsakoff syndrome, and dementia associated with posterior cortical atrophy (PCA).

[00162] In certain embodiments, provided herein are methods for treating or suppressing a disease or disorder selected from the group consisting of an α -synucleinopathy, a tauopathy, an autistic spectrum disorder, a pervasive developmental disorder, a liver disease, liver damage, dementia, and reperfusion injury, comprising orally administering a pharmaceutical composition provided herein.

Preparation of Compound 1

[00163] Compound 1 and its polymorph can be manufactured by methods known in the art, for example, Examples 1A, 1B, 2, 3A, and 3B as described in International Patent Application No. PCT/US2019/056836 (WO 2020/081879) which is incorporated by reference herein in its entirety, including any figures, for all purposes.

Manufacture of Pharmaceutical Compositions

[00164] Typical dosage forms are prepared according to the second aspect. In certain embodiments, step b) occurs at room temperature. In certain embodiments, step a) is mixing about 65:20:15 by weight of a pharmaceutically acceptable oil:a propylene glycol laurates composition:a polysorbate 80. In certain embodiments, step a) is mixing about 65:20:15 by weight of a Labrafac-like oil (Labrafac WL 1349):Lauroglycol 90:Tween 80. In certain embodiments, step a) is mixing about 65:20:20 by weight of a pharmaceutically acceptable oil:a propylene glycol laurates composition:a polysorbate 80. In certain embodiments, step a) is mixing about 65:20:20 by weight a Labrafac-like oil (Labrafac WL 1349):Lauroglycol 90:Tween 80. In certain embodiments, **Compound 1** in step b) is a polymorph of an anhydrate of 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione. In some embodiments, **Compound 1** is measured under light protection or yellow light. In some embodiments, the formulation is added to a capsule. In some embodiments, the formulation solution is stored before encapsulation under light protection or yellow light. In some embodiments, the formulation is added to a capsule under light protection or yellow light.

Dosage and Unit Dosage Forms

[00165] In human therapeutics, the doctor will determine the posology which is considered most appropriate according to a treatment and according to the age, weight, stage of the infection and other factors specific to the subject to be treated. In certain embodiments, the pharmaceutical composition of the disclosure can be administered at the recommended maximum clinical dosage or at lower doses. Dosage levels of the active compounds in the compositions of the disclosure may be varied so as to obtain a desired response depending on the severity of the disorder and the response of the individual.

[00166] In certain embodiments, the pharmaceutical composition is administered for at least 24 weeks. In certain embodiments, the pharmaceutical composition is administered for at least 28 weeks. In certain embodiments, the pharmaceutical composition is administered at a dose selected from 150 mg BID and 250 mg BID. In certain embodiments, the pharmaceutical

composition is administered at a dose of 150 mg BID. In certain embodiments, the pharmaceutical composition is administered at a dose of 250 mg BID. In certain embodiments, the pharmaceutical composition is administered at a dose selected from 100 mg BID and 500 mg BID. In certain embodiments, the pharmaceutical composition is administered at a dose of 100 mg BID. In certain embodiments, the pharmaceutical composition is administered at a dose of 500 mg BID. In certain embodiments, the pharmaceutical composition is administered at a dose of 10 capsules total for each day, wherein the weight of the pharmaceutical composition in each capsule is 50 mg for each day. In certain embodiments, the pharmaceutical composition is administered at a dose of 6 capsules total for each day, wherein the weight of the pharmaceutical composition in each capsule is 50 mg. In certain embodiments, the pharmaceutical composition is administered at dose ranging from about 100 mg to about 1000 mg daily dosage for a period of at least 14 days. The weights of **Compound 1** provided in this paragraph do not include the weight of any water in a hydrate of **Compound 1** or the weight of any solvent in a solvate of **Compound 1**.

[00167] In certain embodiments, the pharmaceutical composition administered dosages show no treatment emergent serious adverse events (TEAEs). In certain embodiments, the pharmaceutical composition administered dosages show no TEAEs leading to discontinuation of the pharmaceutical composition. In certain embodiments, the pharmaceutical composition administered dosages show no TEAEs leading to death.

Methods of Administration

[00168] The pharmaceutical compositions disclosed herein may be administered enterally, orally, sublingually, rectally, or topically in dosage unit formulations. In some embodiments, suitable modes of administration include oral, transdermal, transmucosal, iontophoretic, intraperitoneal, rectal, and gastrointestinal.

[00169] Oral administration is a preferred route of administration, and formulations suitable for oral administration are preferred formulations.

[00170] Oral administration may include administration in Low-fat Food. Oral administration may include administration in Medium-fat Food. Oral administration may include administration in food such as soft foods or semi-solid foods. Soft foods and semi-solid foods include applesauce, pudding (including chocolate), and jam or spread (including strawberry-flavored).

[00171] SEDDS formulations may be more amenable to administration via a feeding tube, as a feeding tube may be difficult to rinse with water. An advantage of the SEDDS formulations provided and claimed herein is that a feeding tube can be rinsed effectively with water. In certain embodiments, administration may be by feeding tube, at a dose of about 0.25 mL to about 10 mL, about 0.25 mL to about 7.5 mL, or about 2.5 mL, of a 100 mg/mL solution. In certain embodiments, administration may be by feeding tube, with a solution at a concentration of about 36 mg/mL to about 120 mg/mL. In certain embodiments, the dose is administered in a vehicle as provided in Formulations 1-6. In some or any embodiments, the ratio of oil to propylene glycol laurates composition to polysorbate 80 is about 65:20:15.

[00172] Topical administration is another preferred route of administration, and formulations suitable for topical administration are preferred formulations.

Formulations for topical administration may include lotions, tinctures, creams, emulsions, ointments, sprays, gels, and the like, and may further be formulated in other suitable formulations such as sunscreens, moisturizing lotions and creams, facial gels and creams, etc. Topical administration may also involve the use of transdermal administration such as transdermal patches or iontophoresis devices.

EXAMPLES

[00173] In the following examples, the abbreviations have the meanings provided in the table below.

ADI	Allowable Daily Intake
DSE	Drug Safety Evaluation
FaSSIF	fasted State Simulated Intestinal Fluid
HLB	hydrophilic-lipophilic balance
o/w	oil/water or oil-in-water
RH	relative humidity
RRT	relative retention time
RT	room temperature
SEDDS	Self-Emulsifying Drug Delivery System
API	Active Pharmaceutical Ingredient
IIG	FDA Inactive ingredient Guide
TPGS	D- α -tocopheryl polyethylene glycol succinate
PO	By mouth
ICH	The International Council for Harmonisation
CRC	Child-resistant Cap
BE	Bioequivalent
Vehicle C	60:20:20 Labrafac WL 1349:Lauroglycol 90:Tween 80
Vehicle D	65:20:15 Labrafac WL 1349:Lauroglycol 90:Tween 80

[00174] Certain densities are provided below where the vehicle for each is 65:20:15 Labrafac WL 1349, Lauroglycol 90, Tween 80:

<i>Solution</i>	<i>Density</i>
<i>60 mg/mL solution</i>	<i>961 mg/mL</i>
<i>100 mg/mL solution</i>	<i>962.9 mg/mL</i>
<i>vehicle (without Compound 1)</i>	<i>959.9 mg/mL</i>

[00175] Densities were measured using procedures known to those of ordinary skill in the art.

Example 1

Screening of solubilizers

[00176] Different solubilizers were screened for the formulation of **Compound 1**. Table 1 showed screening of excipients for the formulation of **Compound 1**. Compounds of Entries 6, 9, 11, 15, and 16 at both concentrations were soluble. Compounds of Entries 7 (200 mg/mL dilution), 8 (200 mg/mL dilution), and 17 (400 mg/mL dilution) were initially soluble but crashed out later (after overnight). Compounds of entries 1-5, 10, and 12-14, were not soluble. Gelucire 44/14 and Labrafac Lipo WL1349 seemed promising as solubilizers for **Compound 1**, and thus further experimentation were conducted with them (Entries 16 and 11 respectively). Although, caproyl also looked promising, it was found to be toxic in monkeys and thus, not pursued.

Table 1: Initial Screening of Potential Solubilizers for **Compound 1**

Entry	Excipient	Initial Mass	400 mg/mL dilution	add	200 mg/mL dilution
1	PEG 400	500	1.25	1.25	2.50
2	tween 80	500	1.25	1.25	2.50
3	corn oil	500	1.25	1.25	2.50
4	cottonseed oil	500	1.25	1.25	2.50
5	span 80	500	1.25	1.25	2.50
6	capryol PGMC	500	1.25	1.25	2.50
7	capryol 90	500	1.25	1.25	2.50
8	lauroglycol 90	500	1.25	1.25	2.50
9	lauroglycol FCC	500	1.25	1.25	2.50
10	plurol oleique CC 497	500	1.25	1.25	2.50
11	labrafac lipo WL 1349	500	1.25	1.25	2.50
12	labrasol	500	1.25	1.25	2.50
13	labrafil M1944CS	500	1.25	1.25	2.50
14	triethyl citrate	500	1.25	1.25	2.50

Entry	Excipient	Initial Mass	400 mg/mL dilution	add	200 mg/mL dilution
15	vitamin E TPGS	500	1.25	1.25	2.50
16	gelucire 44/14	500	1.25	1.25	2.50
17	kolliphor HS 15	500	1.25	1.25	2.50

[00177] Based on the initial screening as shown in Table 1 and based on limits of excipients allowed by FDA, the solubility of **Compound 1** in the oils in Table 1A were measured. Based on the solubility data, Labrafac WL 1349 was further pursued.

Table 1A: Solubility of **Compound 1**

	Solubility, mg/mL	Temp, °C
<i>Gelucire 44/14</i>	<i>269</i>	<i>45</i>
<i>TPGS</i>	<i>195</i>	<i>45</i>
<i>PEG400</i>	<i>11</i>	<i>37</i>
<i>Solutol HS15</i>	<i>94</i>	<i>37</i>
<i>Sesame oil</i>	<i>139</i>	<i>RT</i>
<i>Olive oil</i>	<i>135</i>	<i>RT</i>
<i>Labrafac WL 1349</i>	<i>Between 100 and 150 mg/g</i>	<i>RT</i>
<i>Soybean oil</i>	<i>< 150 mg/g</i>	<i>RT</i>

Example 2

Screening of Gelucire-Based Formulation

[00178] Certain combinations of Gelucire 44/14 with Solutol HS15, TPGS and Tween 80 were screened as shown in Table 2 at 200 mg/g concentration of **Compound 1**. Physical stability was observed microscopically at week one at 25 °C/60%RH and at 5 °C. Bioperformance was assessed visual observation after 10X dilution in FaSSIF. Based on the crash resistance test performed with ten times dilution with FaSSIF, the most promising formulation combination for Gelucire 44/14 appeared to be 80% w/w Gelucire 44/14, 10% w/w TPGS and 10% w/w Tween 80.

Table 2: Screening of Different Potential Gelucire-Based Formulations for **Compound 1** at 200 mg/g Concentration (i.e. 200 mg of **Compound 1** in 1 g of **Compound 1** + vehicle)

Component	%w/w	%w/w	%w/w	%w/w	%w/w	%w/w
Gelucire 44/14	85	80	90	90	80	70

Component	%w/w	%w/w	%w/w	%w/w	%w/w	%w/w
Solutol HS15	5	--	--	--	--	--
TPGS	5	10	10	--	--	10
Tween 80	5	10		10	20	20
Physical stability	Crystallization	Crystallization	Crystallization	Crystallization	Crystallization	Crystallization
Bioperformance	A lot of precipitation	Clear with minute crystals	A lot of precipitation	precipitates	precipitates	precipitates

[00179] In the above table and in this paragraph, the %w/w are based on the total weight of the vehicle without compound. The 80% w/w Gelucire 44/14, 10% w/w TPGS and 10% w/w Tween 80 formulation vehicles were further tested at lower **Compound 1** concentrations of 150 mg/g (i.e. 150 mg of **Compound 1** in 1 g of **Compound 1** + vehicle) and 100 mg/g (i.e. 100 mg of **Compound 1** in 1 g of **Compound 1** + vehicle) to minimize the crystallization risk upon storage. Physical stability was observed microscopically at week one at 25 °C/60%RH and at 5 °C. Bioperformance was assessed visual observation after 10X dilution in FaSSIF. As shown in Table 3, crystallization is observed after 1 h and 1.5 hour for **Compound 1** concentrations of 150 mg/g (i.e. 150 mg of **Compound 1** in 1 g of **Compound 1** + vehicle) and 100 mg/g (i.e. 100 mg of **Compound 1** in 1 g of **Compound 1** + vehicle), respectively.

Table 3: Feasibility of Selected Gelucire-Based Formulation at Different Concentrations of **Compound 1**

Component	Feasibility @ 150 mg/g	Feasibility @ 100 mg/g
	%w/w	%w/w
Gelucire 44/14	80	80
Solutol HS15	--	--
TPGS	10	10
Tween 80	10	10
Physical stability	Crystallization observed under microscope when compared to vehicle	Not clear from Microscope when compared to vehicle
Bioperformance risk	A lot of precipitation at RT within 1 h	Clear up to 1 h, Started seeing precipitation at RT after 1.5 h

[00180] In the above table, the %w/w are based on the total weight of the vehicle without compound.

*Example 3**Screening of Labrafac-Based Formulation*

[00181] To develop a SEDDS formulation of **Compound 1**, the combinations of Labrafac WL 1349 with co-surfactant Lauroglycol 90 (low HLB value) and surfactant Tween 80 (High HLB value) were tested to form a SEDDS formulation (Table 4).

Table 4: Labrafac-Based SEDDS considered for **Compound 1**

Component	Formulation A	Formulation B
	%w/w	%w/w
Labrafac WL 1349	60	80
LauroGlycol	20	10
Tween 80	20	10

[00182] In the above table, the %w/w are based on the total weight of the vehicle without compound. Formulation A formed a clear single phase solution vehicle whereas Formulation B did not form a single phase with Tween 80 forming a separate phase.

[00183] Formulation A was further evaluated as potential SEDDS formulation of **Compound 1**.

*Example 4**Comparison of Gelucire-Based vs. Labrafac-Based Formulations*

[00184] The Gelucire 44/14 formulation in Table 5 was prepared by melting at 45 °C and mixing the required quantities of TPGS (melted at 37 °C) and Tween 80 at RT to form a clear solution vehicle at 45 °C. **Compound 1** was then solubilized by mixing at 45 °C to form a clear solution.

[00185] The Labrafac-based formulation in Table 5 was prepared by mixing Labrafac WL 1349 with Lauroglycol 90 and Tween 80 at RT to form a clear solution vehicle. **Compound 1** was then solubilized by mixing advantageously at RT to form a clear solution. In the table below, the %w/w are based on the total weight of the vehicle without compound.

[00186] Two formulations selected for Gelucire 44/14-based and Labrafac-WL-1349-based formulation vehicles with **Compound 1** at 100 mg/g (i.e. 100 mg of **Compound 1** in 1 g of **Compound 1** + vehicle) are shown and compared in Table 5. The Gelucire 44/14-based formulation is a semi-solid at RT, while the Labrafac-WL-1349-based formulation is advantageously a liquid at RT. The Gelucire 44/14-based formulation, when diluted to 10 times with FaSSIF, forms a clear solution. However, **Compound 1** precipitation is observed after

about 1.5 hour. The Labrafac-WL-1349-based formulation forms an oil-in-water emulsion when diluted to 10 times with water, and **Compound 1** is crash-resistant for at least about 24 hours, being predominantly solubilized in the oil phase.

Table 5: Comparison of Gelucire-Based and Labrafac-Based Formulations for **Compound 1** (100 mg/g) (i.e. 100 mg of **Compound 1** in 1 g of **Compound 1** + vehicle)

Gelucire Vehicle 1		Vehicle C	
Component	% w/w	Component	% w/w
Gelucire 44/14	80	Labrafac WL 1349	60
TPGS	10	LauroGlycol	20
Tween 80	10	Tween 80	20
Water soluble components and thus forms a clear solution upon dilution		Oil based formulation and thus forms an o/w emulsion upon dilution in water	
Upon 10X dilution in FaSSIF, the API is crash resistant up to 1.5 h		Upon 10 X dilution in water, the API is crash resistant up to 24 h with API predominantly solubilized in the oil phase	
Semi-solid formulation at RT		Liquid formulation at RT	
Physically unstable at RT within 1 month		Physically stable at RT up to 1 month	
Chemically stable at 25°C/60%RH and 40°C/75%RH for 1 month		Chemically stable at 25°C/60%RH and 40°C/75%RH for 1 month	

Example 4A

[00187] Physical stability evaluation of both Gelucire 44/14-based and Labrafac-WL-1349-based formulation vehicles was done under polarized microscope. The Gelucire 44/14-based formulation as shown in Figure 1 clearly crystallized out at 25 °C/60%RH at one month with liquid phase separation observed visually. Hence, the Gelucire 44/14 -based formulation was physically unstable upon storage for one month. The Labrafac-WL-1349-based formulation as shown in Figure 2 was still a clear solution with no birefringent crystals observed both at 25 °C/60%RH and 40 °C/75%RH for at one month. Therefore, Labrafac-WL-1349-based formulation was physically stable upon storage.

Example 4B

[00188] The chemical stability evaluation were also performed. Both Gelucire 44/14-based and Labrafac-WL-1349-based formulations are chemically stable at 25 °C/60%RH and 40 °C/75%RH at one month with less than 0.05 percentage points of degradation observed in either formulations as shown in Table 6 and 7 respectively.

[00189] Based on the solubility, and physical stability data, the Labrafac-WL-1349-based formulation was further pursued, and discussed herein.

Table 6: Chemical Stability of Selected Gelucire-Based Formulation of **Compound 1** at 100 mg/g Concentration (i.e. 100 mg of **Compound 1** in 1 g of **Compound 1** + vehicle)

Sample Id	% Assay (% LC)	% Impurity (% Area)						Total % Impurity
		RRT 0.66	RRT 0.78	RRT 0.80	RRT 0.94	Compound 1	RRT 1.44	
Initial	100.8	0.06	0.05	0.13	0.08	99.68		0.32
1M_RT	101.1	0.06	0.04	0.18	0.08	99.60	0.03	0.39
1M_40°C/75%RH	102.2	0.06	0.04	0.14	0.09	99.62	0.04	0.37

[00190] In the above table, 1M means 1 month.

Table 7: Chemical Stability of Selected Labrafac-WL-1349-Based Formulation of **Compound 1** at 100 mg/g Concentration (i.e. 100 mg of **Compound 1** in 1 g of **Compound 1** + vehicle)

Sample Id	% Assay (% LC)	% Impurity (% Area)						Total % Impurity
		RRT 0.66	RRT 0.78	RRT 0.80	RRT 0.94	Compound 1	RRT 1.44	
Initial	101.3	0.06	0.05	0.12	0.08	99.69		0.31
1M_RT	100.6	0.06	0.03	0.08	0.08	99.71	0.03	0.28
1M_40°C/75%RH	101.5	0.07	0.04	0.05	0.08	99.73	0.04	0.27

[00191] In the above table, 1M means 1 month.

Example 5

Stability of Labrafac-WL-1349-Based Formulation Compared to Sesame Oil-based Formulation

[00192] A sesame oil formulation was prepared by mixing 100 mg **Compound 1** per g of **Compound 1** and sesame oil.

[00193] The physical and chemical stability of Vehicle C Labrafac-WL-1349-based formulation of **Compound 1** at 100 mg/g and sesame oil-based formulation were performed.

[00194] Both formulations were physically stable for up to 3 months at 25 °C/60%RH and 40 °C/75%RH conditions with no birefringent crystals observed under polarized microscope (data not shown).

[00195] As shown in Table 8, the formulation was relatively stable with total impurity levels increasing only from 0.31% at initial to 0.38% at three months at 40 °C/75%RH. Table 10 showed the formulation vehicle used.

Table 8: Chemical Stability of Labrafac-WL-1349-Based Formulation of **Compound 1** at 100 mg/g Concentration (i.e. 100 mg of **Compound 1** in 1 g of **Compound 1** + vehicle) up to 6 months and of Sesame Oil-based Formulation of **Compound 1** at 100 mg/g Concentration (i.e. 100 mg of **Compound 1** in 1 g of **Compound 1** + vehicle) up to 3 months

Sample Id	% Assay (% LC)	% Impurity (% Area)									
		RRT 0.19	RRT 0.66	RRT 0.78	RR T 0.80	RR T 0.85	RRT 0.94	Cmpd 1	RRT 1.44	RRT 1.46	Total % Impurity
Initial	101.3	ND	0.06	0.05	0.12	ND	0.08	99.69	ND	ND	0.31
1M 25°C/ 60%RT	100.6	ND	0.06	0.03	0.08	ND	0.08	99.71	0.03	0.03	0.28
3M 25°C/ 60%RT	103.4	ND	0.07	ND	0.04	ND	0.08	99.72	0.06	0.03	0.28
6M 25°C/ 60%RT	101.7	0.02	0.06	0.02	0.04	0.03	0.08	99.64	0.07	0.04	0.36
1M 40°C/ 75%RH	101.5	ND	0.07	0.03	0.05	ND	0.08	99.73	0.04	0.04	0.27
3M 40°C/ 75%RH	102.5	0.04	0.06	ND	0.04	0.04	0.07	99.61	0.08	0.05	0.38
6M 40°C/ 75%RH	100.8	0.05	0.05	ND	0.04	0.04	0.07	99.58	0.09	0.07	0.41

[00196] In the above table, 1M, 3M, and 6M means 1 month, 3 months and 6 months respectively.

Table 8a: Chemical Stability of Sesame Oil-based Formulation of **Compound 1** at 100 mg/g Concentration (i.e. 100 mg of **Compound 1** in 1 g of **Compound 1** + vehicle) up to 3 months

Sample Id	% Assay (% LC)	% Impurity (% Area)										
		RRT 0.66	RRT 0.68	RRT 0.76	RRT 0.78	RRT 0.80	RRT 0.94	Cmpd 1	RRT 1.42	RRT 1.44	RRT 1.46	Total % Impurity
Initial	103.6	0.06	ND	0.03	0.04	0.14	0.09	99.62	ND	0.03	ND	0.39
1M 25°C/ 60%RT	100.7	0.06	ND	0.02	0.04	0.18	0.08	99.58	ND	0.03	ND	0.41
3M 25°C/ 60%RT	101.6	0.06	0.06	ND	0.04	0.17	0.08	99.43	0.05	0.06	0.04	0.56
1M 40°C/ 75%RH	99.9	0.06	ND	0.03	0.04	0.17	0.08	99.57	ND	0.04	ND	0.42
3M 40°C/ 75%RH	101.8	0.06	0.06	0.03	0.04	0.16	0.09	99.37	0.03	0.09	0.06	0.62

[00197] In the above table, 1M and 3M means 1 month and 3 months respectively.

Example 6

ADI Limits for Labrafac-WL-1349-Based Formulation Components

[00198] The Allowable Daily Intake (ADI) values for each of the excipients or components of the Labrafac-WL-1349-based Formulation C-1 is shown in Table 9. At the maximum possible clinical dose of 1000 mg/day of **Compound 1**, the ADI limit for Vehicle C is exceeded for Tween 80 (1800 mg vs. 1500 mg ADI limit). Labrafac WL 1349 and Lauroglycol 90 levels are well within the limit at this highest dose level. The Tween 80 level was reduced from 20% w/w to 15% w/w in the vehicle (Vehicle D) to keep within the ADI limits. The composition of an optimized Vehicle D is Labrafac WL 1349 65%w/w, Lauroglycol 90 20%w/w and Tween 80 15% w/w. This vehicle still forms a clear single-phase solution at RT.

Table 9: ADI Limits for Excipients in Labrafac-WL-1349-Based Formulation C-1 Based on Input From DSE

Ingredient					Formulation	250	500	1000
	IIG	ADI (mg/kg/day)		ADI (mg/day)	Formulation C-1	mg per day	mg per day	mg per day
Cmpd 1					100 mg / g	NA	NA	NA
Labrafac WL 1349 (Gattefosse)/medium-chain triglycerides	3390/5mL	90 ^(a)	Calculated from 47 week rat data.	5400	540	1350	2700	5400
Lauroglycol 90 (Gattefosse)/Propylene glycol monolaurate	235 mg	500 ^(b)	EU limit for food additive	30000	180	450	900	1800
Tween 80 (Croda)/Polysorbate 80	418 mg	25	EU, FDA limit for food additive	1500	180	450	900	1800
Total					1000	NA	NA	NA

(a) The ADI is calculated based on NOAEL of 9 g/kg from 47 weeks study in Wistar rats using procedures disclosed in *Food Chem Toxicol.* **2000**, 38(1), 79-98.

(b) The ADI is 500 mg/kg per day for adult patients and children > 5 years and 50 mg/kg for children < 5 years and > 1 month.

[00199] The IIG values in Table 10 are the limits of the excipients on the FDA website.

[00200] The actual compositions of **Compound 1** (at 100 mg/g concentration, i.e. 100 mg of **Compound 1** in 1 g of **Compound 1** + vehicle) with 18% Tween 80 (60:20:20 Labrafac-WL-1349:Lauroglycol 90:Tween 80) (Formulation C-1) and 13.5% Tween 80 (65:20:15

Labrafac-WL-1349:Lauroglycol 90:Tween 80) (Formulation D-1) are shown in Table 10 and Table 11, respectively.

Table 10: Composition of **Compound 1** in the Vehicle C

Formulation C-1		
Components	%w/w	%w/w
Compound 1	10	--
Labrafac WL 1349	54	60%
Lauroglycol 90	18	20%
Tween 80	18	20%

Table 11: Composition of **Compound 1** in the Vehicle D

Formulation D-1		
Components	%w/w	%w/w
Compound 1	10	--
Labrafac WL 1349	58.5	65%
Lauroglycol 90	18	20%
Tween 80	13.5	15%

[00201] The actual compositions of **Compound 1** (at 60 mg/mL concentration) with 14.07% Tween 80 (65:20:15 Labrafac-WL-1349:Lauroglycol 90:Tween 80) (Formulation E-1) and **Compound 1** (at 100 mg/mL concentration) with 13.44% Tween 80 (65:20:15 Labrafac-WL-1349:Lauroglycol 90:Tween 80) (Formulation F-1) are shown in Table 12 and Table 13, respectively.

Table 12: Composition of **Compound 1** in the Vehicle E

Formulation E-1		
Components	%w/w	%w/w without Compound 1
Compound 1	6.243	---
Labrafac WL 1349	60.94	65%
Lauroglycol 90	18.75	20%
Tween 80	14.07	15%

Table 13: Composition of **Compound 1** in the Vehicle F

Formulation F-1		
Components	%w/w	%w/w without Compound 1
Compound 1	10.39	---
Labrafac WL 1349	58.25	65%
Lauroglycol 90	17.92	20%

Formulation F-1		
Components	%w/w	%w/w without Compound 1
Tween 80	13.44	15%

Example 7

Stability of 50 mg and 60 mg Strength Capsules Prepared from 100 mg/g (Formulation D-1) and 120 mg/g (Formulation D-2) Concentrations, Respectively

[00202] The physical and chemical stability of 50 mg strength capsules of Formulation D-1 (100 mg/g concentration in Vehicle D, i.e. 100 mg of **Compound 1** in 1 g of **Compound 1** + Vehicle D) was assessed at accelerated conditions of temperature and humidity. The physical and chemical stability of 60 mg strength capsules of **Compound 1** Formulation D-2 (120 mg/g concentration in Vehicle D, i.e. 100 mg of **Compound 1** in 1 g of **Compound 1** + Vehicle D) was also assessed at accelerated conditions of temperature and humidity. Size 00 Hard Gelatin Capsules of Swedish orange color were used for the study.

[00203] Both strengths are physically stable for up to one month at 5 °C, 25 °C/60%RH and 40 °C/75%RH conditions with no birefringent crystals observed under polarized microscope (data not shown).

Example 8

Chemical Stability of formulation in Capsules

[00204] The chemical stability of 50 mg and 60 mg strength capsules of **Compound 1** Labrafac-based Formulations D-1 and D-2, respectively, analyzed by HPLC, is shown in Table 14 and Table 15, respectively. About 2% or less change in potency from initial to a duration of time of at least one month at 5 °C, 25 °C/60%RH and 40 °C/75%RH for both strengths were observed. Also, about 1 percentage point or less degradation was observed from initial to all conditions tested for both strengths.

Table 14: Chemical Stability of Formulation D-1 (50 mg Strength Capsules of **Compound 1** in Vehicle D (15% Tween 80 based on weight of vehicle) at 100 mg/g Concentration) up to three months

Sample Id	% Assay (% LC)	% Impurity (% Area)								Total % Impurity
		RRT 0.66	RRT 0.76	RRT 0.78	RRT 0.79	RRT 0.80	RRT 0.94	Compound 1	RRT 1.46	
Initial	98.4	0.06	0.04	0.03	0.02	0.18	0.08	99.56	0.03	0.44
1M_5C	100.9	0.06	0.04	0.02	0.03	0.17	0.08	99.58	0.03	0.43

Sample Id	% Assay (% LC)	% Impurity (% Area)								
		RRT 0.66	RRT 0.76	RRT 0.78	RRT 0.79	RRT 0.80	RRT 0.94	Compound 1	RRT 1.46	Total % Impurity
3M_5C	101.8	0.06	0.02	0.04	0.03	0.18	0.08	99.54	0.03	0.44
1M_25C/ 60%RH	100.2	0.06	0.04	ND	0.03	0.18	0.09	99.57	0.03	0.43
3M_25C/ 60%RH	100.1	0.06	0.02	0.05	0.03	0.17	0.08	99.55	0.03	0.44
1M_40C/ 75%RH	99.7	0.06	0.04	ND	0.03	0.17	0.08	99.57	0.04	0.42
3M_40C/ 75%RH	100.1	0.07	0.02	0.04	0.03	0.12	0.09	99.57	0.05	0.42

Table 15: Chemical Stability of Formulations D-2 (60 mg Strength Capsules of **Compound 1** in Vehicle D (15% Tween 80 based on weight of vehicle) at 120 mg/g Concentration) up to three months

Sample Id	% Assay (% LC)	% Impurity (%Area)										Total % Impurity
		RRT ~ 0.19	RRT ~ 0.66	RRT ~ 0.76	RRT~ 0.78	RRT ~ 0.79	RRT~ 0.80	RRT~ 0.94	Compound 1	RRT ~ 1.44	RRT ~ 1.46	
Initial	102.7	ND	0.06	0.04	0.02	0.02	0.18	0.08	99.57	ND	0.03	0.43
1M_5C	100.7	ND	0.06	0.02	0.04	0.03	0.20	0.07	99.55	ND	0.03	0.45
3M_5C	101.2	ND	0.06	0.02	0.04	0.02	0.20	0.08	99.48	0.04	0.05	0.51
1M_25C/ 60%RH	100.8	ND	0.06	0.03	0.04	0.03	0.18	0.08	99.55	ND	0.03	0.45
3M_25C/ 60%RH	99.0	ND	0.06	0.02	0.04	0.03	0.18	0.08	99.49	0.04	0.05	0.50
1M_40C/ 75%RH	101.2	ND	0.06	0.02	0.04	0.03	0.17	0.08	99.56	ND	0.04	0.44
3M_40C/ 75%RH	100.3	0.03	0.06	0.02	0.04	0.03	0.14	0.08	99.51	0.04	0.05	0.49

Example 9

Chemical Stability of formulation solutions

[00205] The chemical stability of 60 mg/mL and 100 mg/mL strengths of **Compound 1** Labrafac-WL-1349-based Formulations E-1 and F-1, respectively, analyzed by HPLC, is shown in Table 16 and Table 17, respectively. No change in potency was detected from initial measurement to a measurement at two weeks or at one month, for each of the two test conditions (25 °C/60%RH and 40 °C/75%RH. In addition, no degradation was observed from initial to all conditions tested for both solutions.

Table 16: Chemical Stability of Formulation E-1 (at 60 mg/mL Concentration of **Compound 1** - Labrafac-WL-1349-Based Formulation) up to three months

Compound 1 Oral Solution 60 mg/mL						
Sample Id	Appearance	% Assay	% Impurity Individual unspecified	% Impurity Total	Water Content (%)	Water Activity (aw)
Initial	Clear yellow liquid	101.0	ND	ND	0.04	0.1685
2W_25 °C /60%RH	Clear yellow liquid	101.5	ND	ND	0.18	-
1M_25 °C /60%RH	Clear yellow liquid	101.5	ND	ND	0.07	-
2W_40 °C /75%RH	Clear yellow liquid	100.8	ND	ND	0.13	-
1M_40 °C /75%RH	Clear yellow liquid	101.9	ND	ND	0.09	-

Table 17: Chemical Stability of Formulation F-1 (at 100 mg/mL Concentration of **Compound 1** – Labrafac-WL-139-Based Formulation) up to three months

Compound 1 Oral Solution 100 mg/mL						
Sample Id	Appearance	% Assay	% Impurity Individual unspecified	% Impurity Total	Water Content (%)	Water Activity (aw)
Initial	Clear yellow liquid	100.5	ND	ND	0.05	0.1710
2W_25C/60%RH	Clear yellow liquid	100.1	ND	ND	0.16	-
1M_25C/60%RH	Clear yellow liquid	100.9	ND	ND	0.08	-
2W_40C/75%RH	Clear yellow liquid	99.8	ND	ND	0.17	-
1M_40C/75%RH	Clear yellow liquid	100.6	ND	ND	0.07	-

Example 10

Monkey PK Study

[00206] A monkey PK study was conducted with a sesame oil formulation of 100 mg/g of **Compound 1**, a Gelucire 44/14-based formulation of 100 mg/g **Compound 1** (Gelucire Form 1), and two different Labrafac-WL-1349-based formulations of 100 mg/g **Compound 1**. In this paragraph, 100 mg/g is 100 mg of **Compound 1** in 1 g of **Compound 1** + vehicle. The formulations were administered with Low-fat Food for Testing and Medium-fat Food for Testing and Labrafac-WL-1349-based formulation of **Compound 1** administered with Low-fat Food for Testing at dose of 125 mg, PO (Table 18). Each of the formulations in Table 18 were administered as 4 capsules/dose. In the table below, the %w/w are based on the total weight of the vehicle without compound. Four monkeys per cohort were used.

Table 18: Monkey PK Study Design for **Compound 1** Formulations

Formulations:	Meal
P1G1: Sesame oil solution	Medium Fat Food for Testing
P2G1: Gelucire Vehicle 1 (80 w/w% Gelucire 44/14, 10 w/w% TPGS and 10 w/w% Tween 80)	Medium Fat Food for Testing
PO1: Labrafac Vehicle C (60 wt/wt% Labrafac WL 1349, 20 wt/wt% Lauroglycol 90, 20 wt/wt% Tween 80)	Low Fat Food for Testing
P1G2: Vehicle C (60 wt/wt% Labrafac WL 1349, 20 wt/wt% Lauroglycol 90, 20 wt/wt% Tween 80)	Medium Fat Food for Testing
PO2: Vehicle D (65 wt/wt% Labrafac WL 1349, 20 wt/wt% Lauroglycol 90, 15 wt/wt% Tween 80)	Low Fat Food for Testing
PO3: Vehicle D (65 wt/wt% Labrafac WL 1349, 20 wt/wt% Lauroglycol 90, 15 wt/wt% Tween 80)	Medium Fat Food for Testing
Low Fat: 11.4% fat (by calories), about 34 kcal	
Medium Fat: 37.9% fat (by calories), about 62 kcal	
Dose: 125 mg PO Single Dose	

[00207] As shown in Table 19, bioavailability appears to be superior for Gelucire 44/14-based formulation (60.0%). Bioavailability of Labrafac-based formulation P1G2 (49.1%) is comparable to sesame oil-based formulation P1G1 (~40.8%) with Labrafac formulation P1G2 having slightly higher exposure at 49.1%. Another observation is the absence of a second peak in the PK profiles of Gelucire 44/14-based and Labrafac-based formulations of **Compound 1** compared to sesame oil-based formulation. This second peak in the sesame oil formulation indicates an unknown absorption phenomenon.

[00208] Also as shown in Table 19, better mitigation of observed positive food effect (*i.e.* a smaller difference in bioavailability between an administration with Low-fat Food for Testing

and an administration with Medium-fat Food for Testing) is best with Labrafac-WL-1349-based formulation with Vehicle D (where the vehicle alone contains 15 wt/wt% Tween 80) compared to Labrafac-WL-1349-based formulation with Vehicle C (where the vehicle alone contains 20 wt/wt% Tween 80). With a low-fat meal, the Labrafac-WL-1349-based formulation with Vehicle D (where the vehicle alone contains 15 wt/wt% Tween 80) has a comparable bioavailability to Labrafac-WL-1349-based formulation with Vehicle C (where the vehicle alone contains 20 wt/wt% Tween 80). However, with Medium Fat Food for Testing, a formulation using Vehicle D has a lower bioavailability of 29.4% compared to 49.1% bioavailability obtained with a formulation using Vehicle C. Thus, the lesser difference in bioavailability between Low Fat Food for Testing and Medium Fat Food for Testing for Vehicle D (15% Tween 80) when compared to Vehicle C (20% Tween 80) is surprising. This means that a formulation of Compound 1 in Vehicle D could be administered advantageously with or without food. In the table below, the wt/wt% refers to the amount of Tween in the vehicle (without compound).

Table 19: Monkey PK Study Results (PK Parameters) for Various Formulations of **Compound 1** (administered PO, 125mg)

Formulation	Food (Fat%)	T _{1/2} (h)	T _{max} (h)	C _{max} (h)	AUC _{last} (h*ng/mL)	AUC _{0-24h} (h*ng/mL)	C _{avg, 0-24h} (ng/mL)	C _{24h} (ng/mL)	F** (%)
Sesame Solution* (P1G1)	37.9%	6.94	5.3	2080	10800	10200	416	77.8	40.8
Gelucire 44/14 (P2G1)		8.25	4.0	4050	15800	15400	641	43.4	60.0
20 wt/wt% Tween (PO1)	11.4%	5.60	4.0	742	3220	3220	134	10.5	13.4
20 wt/wt% Tween (P1G2)	37.9%	8.87	4.5	2690	12900	12600	524	37.7	49.1
15 wt/wt% Tween (PO2)	11.4%	7.25	4.0	689	3040	3010	125	13.6	14.0
15 wt/wt% Tween (PO3)	37.9%	5.15	4.0	1110	5660	5660	236	17.2	29.4

* One animal was excluded from the calculation; ** F calculated relative to IV dose of 0.5 mg/kg (1.3 mg) in Study II (doses were adjusted for B.Wt).

a: T_{last} 16.5 hr; b: T_{last} 14.0 hr.

Example 10a

Solubility Screening of Additional Pharmaceutically Acceptable Oils ("oils"), Surfactants and Co-surfactants

[00209] A total of 36 different excipients including pharmaceutically acceptable oils, surfactants and co-surfactants were evaluated for **Compound 1**. Solubility is as shown in Table 20. Higher solubility of **Compound 1** in oils is desirable to increase drug loading.

Table 20: SEDDS Excipient Solubility Screening Studies

	Component	Solubility mg/mL	Temperature	
Glycerides	Sesame Oil	139	RT	Oils
	Olive Oil	135	RT	
	Soybean Oil	110.59	RT	
	Corn Oil	115.34	RT	
	Cottonseed Oil	111.76	RT	
	Maisine CC	90.34	RT	
	Pe Ceol	IIG of 0.8 mg		
	Labrafac Lipo WL1349	140.92	RT	
	Labrafac PG	221.91	RT	
	Labrafac MC60	Semi-solid		
	Castor Oil	Laxative		
	Peanut Oil	Non-GMP		
Polyglycerol esters	Plurol oleique CC 497	50-100 (range)		Co-surfactants
Propylene glycol esters	Lauroglycol FC C	159.7	RT	
	Capryol P GMC	183.74	RT	
	Capryol 90	124.54	RT	
	Lauroglycol 90	133.48	RT	
Polyoxyglycerides PEG esters	Labrafil M1944CS	100.75	RT	Water Dispersible Surfactant
	Labrafil M2125CS	118.61	RT	
	Labrafil M2130CS	Solid ²		
	Labrasol	35.73	RT	
	TPGS	195	45	
	Gelucire 44/14	269	45	
	Gelucire 48/16	Solid		
	Gelucire 50/13	Solid		
Additional Surfactants	Tween 80	25-50 (range)		Other Surfactant
	Tween 20	Not Filterable		
	Tween 40	Non-GMP Non-GMP		

	Component	Solubility mg/mL	Temperature	
	Tween 60	50-100 (range)		
	Span 80	Solid		
	Span 20	Non-GMP		
	Span 60	Non-GMP		
	Triethyl citrate	30.06	RT	
	Poloxamer 124	Non-GMP		
	Poloxamer 331	Non-GMP		

SEDDS Formulation Screening: Phase Clarity and Solubility Studies

[00210] As shown in Table 20, the SEDDS formulation can be achieved with a range of pharmaceutically acceptable oils. However, the choice of surfactant and co-surfactant may be limited due to several reasons: (a) some commercially available surfactants are non-GMP in nature and hence cannot be used for humans, (b) semi-solid physical state of some of the excipients makes them inappropriate for oral solution formulation, and/or (c) phase miscibility issues with some surfactants causing physical instability including drug precipitation. In order to achieve an optimized SEDDS formulation of a combination of oil, surfactant, and co-surfactant. Several formulation design related aspects are considered, for example: safe amount for daily intake for excipients, drug solubility in SEDDS formulations, crash resistance/drug precipitation, physical and chemical stability, bioperformance, and pharmacokinetics. Based on these aforementioned criteria, two co-surfactants (Lauroglycol 90 and Capryol 90) and a surfactant (Tween 80) were selected to prepare additional SEDDS formulations. Sesame oil only formulation exhibited a second peak as shown in Fig. 3.

[00211] A total of 15 SEDDS formulations were screened for the phase clarity and solubility studies as shown in Table 21. In the table below, the %w/w are based on the total weight of the vehicle without compound. Most of the SEDDS formulations (e.g., olive-oil based, Labrafac-WL-1349-based etc.) were clear and provided high drug solubility. Highest solubility of **Compound 1** was observed with Labrafac PG and Labrafac WL 1349 based formulations. The SEDDS formulations that exhibited hazy or had significantly lower drug solubility were omitted from further screening. Hence, a total of 6 formulations (Table 21, Prototype # 4, 6, 10, 12, 14, and 15) were selected for a monkey PK study.

Table 21: Compositions of SEDDS Formulations Evaluated in Monkey PK Study

Prototype	Excipient Components			Observations	
	Oil Solubilizer (65% wt/wt)	Co-surfactant (20% wt/wt)	Surfactant (15% wt/wt)	Visual Clarity	Compound 1 Solubility (mg/mL at RT)
1	Maisine CC	Caproyl 90	Tween 80	Clear	108.54
2	Maisine CC	Lauroglycol 90	Tween 80	Clear	103.85
3	Olive Oil	Caproyl 90	Tween 80	Clear	130.2
4	Olive Oil	Lauroglycol 90	Tween 80	Clear	135.32
5	Corn Oil	Caproyl 90	Tween 80	Hazy	NA
6	Corn Oil	Lauroglycol 90	Tween 80	Clear	131.45
7	Sesame Oil	Caproyl 90	Tween 80	Hazy	NA
8	Sesame Oil	Lauroglycol 90	Tween 80	Clear	136.13
9	Labrafac WL 1349	Caproyl 90	Tween 80	Clear	129.08
10	Labrafac WL 1349	Lauroglycol 90	Tween 80	Clear	176.2
11	Soybean Oil	Caproyl 90	Tween 80	Hazy	NA
12	Soybean Oil	Lauroglycol 90	Tween 80	Clear	128.17
13	Cottonseed Oil	Caproyl 90	Tween 80	Hazy	NA
14	Cottonseed Oil	Lauroglycol 90	Tween 80	Clear	132.47
15	Labrafac PG	Lauroglycol 90	Tween 80	Clear	233.10

[00212] The six formulations from Table 21 that exhibited phase clarity and good to excellent solubility for **Compound 1** were evaluated in monkey PK studies as shown in Table 22. In the table below, the %w/w are based on the total weight of the vehicle and compound. All six formulations were crash resistant. Maisine CC based formulations from Table 21 were not further evaluated due to relatively lower solubility of **Compound 1** in these formulations. Caproyl based co-surfactant formulations were eliminated due to toxicity issues found with the excipient.

Table 22: Compositions of SEDDS Formulations Evaluated in Monkey PK Study

Formulation #	Cmpd 1 (% wt/wt)	Solubilizer (% wt/wt)	Surfactant (% wt/wt)	Co-surfactant (% wt/wt)	Crash Resistant
1	10.39	Cottonseed oil (58.25)	Tween 80 (17.92)	Lauroglycol 90 (13.44)	Yes

Formulation #	Cmpd 1 (% wt/wt)	Solubilizer (% wt/wt)	Surfactant (% wt/wt)	Co-surfactant (% wt/wt)	Crash Resistant
2	10.39	Soybean oil (58.25)	Tween 80 (17.92)	Lauroglycol 90 (13.44)	Yes
3	10.39	Labrafac PG (58.25)	Tween 80 (17.92)	Lauroglycol 90 (13.44)	Yes
4	10.39	Olive oil (58.25)	Tween 80 (17.92)	Lauroglycol 90 (13.44)	Yes
5	10.39	Corn oil (58.25)	Tween 80 (17.92)	Lauroglycol 90 (13.44)	Yes
6	10.39	Labrafac WL 1349 (58.25)	Tween 80 (17.92)	Lauroglycol 90 (13.44)	Yes

[00213] Three monkeys per group were evaluated and dosed with 50 mg/kg of **Compound 1**, in the respective formulations. The monkeys were fed 30 minutes prior to dosing with a medium fat liquid diet, 5 mL/kg containing 0.23 g/mL of total fat. Monkey PK study results are shown in Fig. 4a-4c. All of the SEDDS formulations were found to be orally bioavailable in monkeys and showed similar T_{max} (Table 23). Most of the SEDDS formulations, including the Labrafac WL 1349-based formulation (Formulation 6), exhibited desirable PK profile, including the unexpected absence of a second peak. A second peak was observed with the sesame oil formulation and Formulation 4. In the first four hours, soybean, Labrafac PG, and Olive oil-based formulations showed a similar PK profile compared to the Labrafac WL 1349-based formulation (Fig. 5a-5c). Cottonseed oil-based formulation 1 showed the highest C_{max} and AUC (Fig. 5b and 5c).

[00214] The Labrafac WL 1349-based formulation showed the lowest PK variability and the desired plasma levels with minimal peak-to-trough ratio (difference between C_{max} and C_{trough}), Table 23. The peak (bump) in the elimination phase observed with the sesame oil-only formulation and Formulation 4 was not observed with Formulations 1-3 and, 5, and 6.

Table 23: Monkey PK Study Results of SEDDS Formulations

Group	Animal	$T_{1/2}$ hr	T_{max} hr	C_{max}	AUC_{last}
Formulation 1 (Cottonseed oil based)	1	8.8	4.0	3030	14593
	2	10.0	4.0	2310	14177
	3	3.9	4.0	2070	6532
	Mean	7.6	4.0	2470	11767
	SD	3.3	0.0	500	4539
	%RSD	43.1	0.0	20.2	38.6
Formulation 2 (Soybean oil based)	4	6.8	4.0	1080	5014
	5	9.9	4.0	1530	8430
	6	9.9	4.0	408	2147

Group	Animal	T _{1/2} hr	T _{max} hr	C _{max}	AUC _{last}
	Mean	8.9	4.0	1006	5197
	SD	1.8	0.0	565	3146
	%RSD	20.5	0.0	56.1	60.5
Formulation 3 (Labrafac PG based)	7	3.4	4.0	575	3402
	8	5.8	4.0	2070	7641
	9	4.9	4.0	1730	10991
	Mean	4.7	4.0	1458	7345
	SD	1.2	0.0	784	3803
	%RSD	25.1	0.0	53.7	51.8
Formulation 4 (Olive oil based)	10	6.0	6.0	1480	10354
	11	4.0	2.0	1340	7180
	12	7.6	2.0	1230	8027
	Mean	5.9	3.3	1350	8520
	SD	1.8	2.3	125	1643
	%RSD	30.7	69.3	9.3	19.3
Formulation 5 (Corn oil based)	13	9.6	4.0	2510	12584
	14	4.0	4.0	1410	6986
	15	4.8	6.0	1160	6665
	Mean	6.2	4.7	1693	8745
	SD	3.0	1.2	718	3329
	%RSD	49.0	24.7	42.4	38.1
Formulation 6	16	18.8	6.0	989	6079
	17	7.3	4.0	772	4807
	18	10.7	4.0	795	5967
	Mean	12.3	4.7	852	5618
	SD	5.9	1.2	119	704
	%RSD	48.4	24.7	14.0	12.5

Table 23a: Monkey PK Study Results of SEDDS Formulations - Bioavailability

Group	Animal	AUC _{last} /D	AUC _{inf}	AUC _{inf} /D	AUC%extrap	F%
Formulation 1 (Cottonseed oil based)	1	292	14660	293	0.46	47
	2	284	14284	286	0.75	45
	3	131	6570	131	0.58	21
	Mean	235	11838	237	0.60	38
	SD	90.8	4566	91.3	0.15	14
	%RSD	38.6	38.6	38.6	24.8	38.6
Formulation 2 (Soybean oil based)	4	100	5066	101	1.03	16
	5	169	8598	172	1.95	27
	6	42.9	2224	44.5	3.45	7
	Mean	104	5296	106	2.14	17
	SD	62.9	3193	63.9	1.22	10
	%RSD	60.5	60.3	60.3	56.9	60.3

Group	Animal	AUClast/D	AUCinf	AUCinf/D	AUC%extrap	F%
Formulation 3 (Labrafac PG based)	7	68.0	3429	68.6	0.77	11
	8	153	7685	154	0.57	24
	9	220	11030	221	0.36	35
	Mean	147	7381	148	0.57	23
	SD	76.1	3810	76.2	0.21	12
	%RSD	51.8	51.6	51.6	36.8	51.6
Formulation 4 (Olive oil based)	10	207	10465	209	1.06	33
	11	144	7278	146	1.35	23
	12	161	8105	162	0.95	26
	Mean	170	8616	172	1.12	27
	SD	32.9	1654	33.1	0.21	5
	%RSD	19.3	19.2	19.2	18.3	19.2
Formulation 5 (Corn oil based)	13	252	12661	253	0.60	40
	14	140	7055	141	0.98	22
	15	133	6701	134	0.54	21
	Mean	175	8806	176	0.71	28
	SD	66.6	3343	66.9	0.24	11
	%RSD	38.1	38.0	38.0	34.0	38.0
Formulation 6	16	122	6309	126	3.65	20
	17	96.1	4863	97.3	1.15	15
	18	119	6117	122	2.45	19
	Mean	112	5763	115	2.42	18
	SD	14.1	785	15.7	1.25	2
	%RSD	12.5	13.6	13.6	51.6	13.6

[00215] Certain formulation provided herein were tested for stability, as provided herein, and demonstrated good stability under various conditions. Other formulations, such as Formulations 1-5, provided herein, have not been tested for stability but would be expected to demonstrate comparable stability to those that were tested.

[00216] In certain embodiments, 20 wt/wt% Lauroglycol (wt/wt% based on total weight of vehicle) may advantageously assist in maintaining a one phase solution.

[00217] One advantage for Formulation 6 compared to a formulation of **Compound 1** in sesame oil only is that Formula 6, with 10%w/w and 12% w/w **Compound 1**, are stable at RT, where the maximum advisable loading of **Compound 1** in sesame oil only is 10% w/w. In addition, while the formulation of **Compound 1** in sesame oil only demonstrated good chemical stability at RT, Formulation 6 demonstrated somewhat better chemical stability at accelerated conditions and for a longer term. Still another advantage of Formulation 6 compared to the formulation of **Compound 1** in sesame oil only relates to micellization and

solubilization in gastrointestinal fluids; Formulation 6 is designed to self-emulsify where sesame oil only formulation must rely on intestinal surfactants. Finally, the sesame oil formulation shows a 2.7-fold food effect (low fat to medium fat) where Formulation 6 shows a 1.96-fold food effect (low fat to medium fat).

Example 11

*Manufacturability of **Compound 1** Labrafac-WL-1349-Based Formulation Capsules*

(i). Viscosity of Labrafac-Based Formulation Vehicle

[00218] A Brookfield viscometer with CP-51 spindle (Medium – High viscosity spindle) was calibrated. With highest 200 rpm (CP-51), Formulation D (Labrafac-WL-1349-based formulation vehicle with 15% Tween 80) has a viscosity of 23.56 cP. The acceptable viscosity for the encapsulation of the DP-vehicle mixing blend should be in between 0.222 and 3000 Cp.

(ii). Moisture Abstractability of Labrafac-WL-1349-Based Formulation Vehicle Components

[00219] Moisture abstractability of the liquid formulation components in Hard Gelatin Capsule is shown in Figure 6. The water equilibrium may be describe by the curves in Figure 3 depending on the moisture/drying history. As long as the gelatin water content remains in the range of 11 to 16%, no significant effect is expected on capsules brittleness potential. Table 25 shows the analysis of individual components of Labrafac-WL-1349-based formulation on moisture abstractability.

[00220] Both Labrafac-WL-1349 and Lauroglycol 90 have a very low potential to absorb moisture. While Tween 80 is hygroscopic, it is only 15% w/w in the vehicle and is not expected to influence the integrity of the capsule shell upon storage.

Table 25: Analysis of Individual Components of Labrafac-WL-1349-Based Formulation C-1 and D-1 on Moisture Abstractability

Component	Formulation C-1 (20% Tween 80)		Formulation D-1 (15% Tween 80)		Description / Criteria	Risk level
	% w/w	mg / unit	% w/w	mg / unit		
Cmpd 1	10	50	10	50		NA
Labrafac Lipophile WL 1349	54	270	58.5	292.5	Medium chain triglyceride of fractionated vegetable C8 and C10 fatty acids (mainly fractionated coconut oil or palm kernel oil) with an HLB of 1; non rancidable; no to very low hygroscopicity	Low

Component	Formulation C-1 (20% Tween 80)		Formulation D-1 (15% Tween 80)		Description / Criteria	Risk level
	% w/w	mg / unit	% w/w	mg / unit		
Lauroglycol 90	18	90	18	90	Non-ionic water insoluble surfactant used as co-surfactant, consists of propylene glycol mono- and di-stearates of lauric acid (C12), with HLB value 3	Low
Tween 80	18	90	13.5	67.5	Polysorbates containing 20 units of oxyethylene are hydrophilic, hygroscopic, nonionic surfactants that are used as emulsifying agents	Med - High

Example 12

Manufacturing Process for Oral Solution of **Compound 1**

[00221] Each of the oil, propylene glycol laurates composition (Lauroglycol 90), and polysorbate 80 (Tween 80) were weighed and added to a container, and were mixed until a uniform solution was formed. To this solution was added **Compound 1**. The resulting mixture was stirred until a yellow solution was obtained. The solution was then test visually and under microscope for the absence of **Compound 1** crystals. The solution was filled in an amber glass bottle, closed and sealed with CRC cap. The optional flavorant can be added to either the mixture without **Compound 1** or to the mixture containing **Compound 1**.

Example 13

Manufacturing Process for Semi-Meal Procedure

Example 13a: Low fat meat (containing 15% calories from fat):

[00222] The homogenized meal was prepared as follow:

30 g daily food (approximately 6% fat by weight) was weighed in a container. 90 mL water was added into the container and contents homogenized.

[00223] For low diet groups, animals were fasted overnight and given homogenized normal meal by gavage at 37.5 mL/animal (approximately 15 mL/kg) about 30 mins prior to dosing. The rest diet, approximately 90 g, was given to animals 4 hours post-dose.

Example 13b: Medium fat meat (containing 38% calories from fat):

[00224] The homogenized meal was prepared as follow:

80 g of food weighed in a container. 15 mL eatable corn oil and some water were added into the container and placed it to soften them for about 30 min. Homogenized 20 min to mix well and made up the volume to 250 mL with water.

[00225] For medium diet groups, animals were fasted overnight and given homogenized normal meal by gavage at 37.5 mL/animal (approximately 15 mL/kg) about 30 mins prior to dosing. The rest diet (about 103 g), was given to animals 4 hours post-dose.

[00226] The disclosures of all publications, patents, patent applications and published patent applications referred to herein by an identifying citation are hereby incorporated herein by reference in their entirety.


[00227] Although the foregoing invention has been described in some detail by way of illustration and example for purposes of clarity of understanding, it is apparent to those skilled in the art that certain minor changes and modifications will be practiced. Therefore, the description and examples should not be construed as limiting the scope of the invention.

What is claimed is:

1. A pharmaceutical composition comprising:

a) about 48 wt/wt% to about 70 wt/wt% of a pharmaceutically acceptable oil, about 12 wt/wt% to about 25 wt/wt% of a propylene glycol laurates composition, and about 8 wt/wt% to about 20 wt/wt% of a polysorbate 80;

b) about 1 wt/wt% to about 15 wt/wt% of **Compound 1**:



and/or the hydroquinone form thereof; wherein **Compound 1**, and/or the hydroquinone thereof is optionally a hydrate thereof, and/or solvate thereof; and wherein when **Compound 1**, and/or hydroquinone thereof, is in the form of a hydrate and/or solvate, then the about 1 wt/wt% to about 15 wt/wt% of **Compound 1**, and/or its hydroquinone, does not include the weight of the water in the hydrate or the weight of the solvent in the solvate; and


c) 0 wt/wt% to about 2% wt/wt% of an optional flavorant;

wherein the wt/wt% of **Compound 1**, the pharmaceutically acceptable oil, the propylene glycol laurates composition, the polysorbate 80, and optional flavorant total 100%.

2. The pharmaceutical composition of claim 1, comprising:

a) about 52 wt/wt% to about 65 wt/wt% of a pharmaceutically acceptable oil, about 17 wt/wt% to about 20 wt/wt% of a propylene glycol laurates composition, and about 13 wt/wt% to about 20 wt/wt% of a polysorbate 80;

b) about 1 wt/wt% to about 12 wt/wt% of **Compound 1**:



and/or the hydroquinone form thereof; wherein **Compound 1**, and/or the hydroquinone thereof is optionally a hydrate thereof, and/or solvate thereof; and wherein when **Compound 1**, and/or hydroquinone thereof, is in the form of a hydrate and/or solvate, then the about 1 wt/wt% to about 12 wt/wt% of **Compound 1**, and/or its hydroquinone, does not include the weight of the water in the hydrate or the weight of the solvent in the solvate; and

c) 0 wt/wt% to about 2% wt/wt% of an optional flavorant;

wherein the wt/wt% of **Compound 1**, the pharmaceutically acceptable oil, the propylene glycol laurates composition, the polysorbate 80, and optional flavorant total 100%.

3. The pharmaceutical composition of claim 1 or 2, wherein the ratio of the pharmaceutically acceptable oil to propylene glycol laurates composition to polysorbate 80 is
about 55-75 parts by weight of the pharmaceutically acceptable oil,
about 15-25 parts by weight of the propylene glycol laurates composition,
about 10-20 parts by weight of the polysorbate 80, or
wherein the ratio is 55-75:15-25:10-20 by weight; and
wherein the parts in the ratio of the pharmaceutically acceptable oil to propylene glycol laurates composition to polysorbate 80 total 100.

4. The pharmaceutical composition of any one of claims 1-3, wherein the ratio of the pharmaceutically acceptable oil to propylene glycol laurates composition to polysorbate 80 is
about 60-65 parts by weight of the pharmaceutically acceptable oil,
about 20-25 parts by weight of the propylene glycol laurates composition,
about 15-20 parts by weight of the polysorbate 80, or
wherein the ratio is 60-65:20-25:15-20; and
wherein the parts in the ratio of the pharmaceutically acceptable oil to propylene glycol laurates composition to polysorbate 80 total 100.

5. The pharmaceutical composition of any one of claims 1-4, wherein the ratio of the pharmaceutically acceptable oil to propylene glycol laurates composition to polysorbate 80 is
about 60 parts by weight of the pharmaceutically acceptable oil,
about 20 parts by weight of the propylene glycol laurates composition,
about 20 parts by weight of the polysorbate 80, or
wherein the ratio is 60:20: 20.

6. The pharmaceutical composition of any one of claims 1-4, wherein the ratio of the pharmaceutically acceptable oil to propylene glycol laurates composition to polysorbate 80 is
about 65 parts by weight of the pharmaceutically acceptable oil,
about 20 parts by weight of the propylene glycol laurates composition,
about 15 parts by weight of the polysorbate 80, or
wherein the ratio is 60:20:15.

7. The pharmaceutical composition of any one of claims 1-6, wherein the pharmaceutically acceptable oil is
selected from the group consisting of one or more of: a Labrafac-like oil (Labrafac WL 1349 or Labrafac PG), sesame oil, cottonseed oil, soybean oil, olive oil, and corn oil;
or
selected from the group consisting of one or more of: a Labrafac-like oil (Labrafac WL 1349 or Labrafac PG), cottonseed oil, soybean oil, and corn oil; and
optionally wherein one pharmaceutically acceptable oil is selected.
8. The pharmaceutical composition of any one of claims 1-7, wherein the pharmaceutically acceptable oil is a Labrafac-like oil (Labrafac WL 1349 or Labrafac PG),.
9. The pharmaceutical composition of any one of claims 1-7, wherein the pharmaceutically acceptable oil is sesame oil.
10. The pharmaceutical composition of any one of claims 1-7, wherein the pharmaceutically acceptable oil is soybean oil.
11. The pharmaceutical composition of any one of claims 1-7, wherein the pharmaceutically acceptable oil is Labrafac WL 1349.
12. The pharmaceutical composition of any one of claims 1-7, wherein the pharmaceutically acceptable oil is olive oil.
13. The pharmaceutical composition of any one of claims 1-7, wherein the pharmaceutically acceptable oil is corn oil.
14. The pharmaceutical composition of any one of claims 1-7, wherein the pharmaceutically acceptable oil is cottonseed oil.
15. The pharmaceutical composition of any one of claims 1-14, wherein the propylene glycol laurates composition is Lauroglycol 90.
16. The pharmaceutical composition of any one of claims 1-15, wherein the polysorbate 80 is Tween 80.
17. The pharmaceutical composition of any one of claims 1-16, wherein the pharmaceutical composition comprises about 1 wt/wt% to about 12 wt/wt%, about 4 wt/wt% to about 12 wt/wt%, or about 5 wt/wt% to about 10 wt/wt% of **Compound 1**.

18. The pharmaceutical composition of any one of claims 1-17, wherein the pharmaceutical composition comprises about 5 wt/wt%, about 6 wt/wt%, about 7 wt/wt%, about 8 wt/wt%, about 9 wt/wt%, or about 10 wt/wt% of **Compound 1**.
19. The pharmaceutical composition of any one of claims 1-18, wherein the pharmaceutical composition comprises about 5 wt/wt% of **Compound 1**.
20. The pharmaceutical composition of any one of claims 1-18, wherein the pharmaceutical composition comprises about 6 wt/wt% of **Compound 1**.
21. The pharmaceutical composition of any one of claims 1-18, wherein the pharmaceutical composition comprises about 10 wt/wt% of **Compound 1**.
22. The pharmaceutical composition of any one of claims 1-21, wherein the pharmaceutical composition contains about 50 mg to about 120 mg of **Compound 1** in the composition, about 50 mg to about 100 mg of **Compound 1** in the composition, about 60 mg to about 100 mg of **Compound 1** in the composition, about 100 mg to about 120 mg of **Compound 1** in the composition, about 50 mg of **Compound 1** in the composition, about 60 mg of **Compound 1** in the composition, or about 100 mg of **Compound 1** in the composition.
23. The pharmaceutical composition of claim 1-22, wherein the pharmaceutical composition is stable when stored at about 15 °C to about 25 °C or at about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least about one month, at least about three months, or at least about six months.
24. The pharmaceutical composition of any one of claims 1-22, wherein the pharmaceutical composition shows no birefringent crystals when stored at about 15 °C to about 25 °C or at about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least about one month.
25. The pharmaceutical composition of any one of claims 1-22, wherein the pharmaceutical composition shows no birefringent crystals when stored at about 15 °C to about 25 °C or at about 25 °C and about 60% relative humidity for a duration of at least about one month or at least about six months.
26. The pharmaceutical composition of any one of claims 1-22, wherein the pharmaceutical composition shows no birefringent crystals when stored at about 40 °C and about 75% relative humidity for a duration of at least about one month.

27. The pharmaceutical composition of any one of claims 1-26, wherein the total impurity level increases by 25% or less from an initial total impurity level to a subsequent total impurity level, when stored at about 40 °C and about 75% relative humidity for at least about three months, optionally wherein the initial total impurity level is measured on the day the pharmaceutical composition is prepared and before it is stored on the same day as it is prepared; and wherein the subsequent total impurity level is tested after about 3 months in storage.


28. The pharmaceutical composition of any one of claims 1- 27, wherein the subsequent total impurity level is about 1% or less, about 0.5% or less, or about 0.4% or less.

29. The pharmaceutical composition of claim 27 or 28, wherein the pharmaceutical composition shows about 2% or less change in potency from an initial potency after storing at about 15 °C to 25 °C or at about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least one month; optionally wherein the initial potency is measured on the day the pharmaceutical composition is prepared and before it is stored on the same day as it is prepared.

30. The pharmaceutical composition of claim 27 or 28, wherein the pharmaceutical composition shows about 1% or less degradation from an initial degradation level after storing at about 15 °C to 25 °C or at about 25 °C and about 60% relative humidity or at about 40 °C and about 75% relative humidity for a duration of time of at least one month; optionally wherein the initial degradation level is measured on the day the pharmaceutical composition is prepared and before it is stored on the same day as it is prepared.

31. The pharmaceutical composition of any one of claims 1-22, wherein the pharmaceutical composition is stable when stored at about 5 °C for a period of at least about 4 weeks.


32. The pharmaceutical composition of any one of claims 1-31, wherein b) is



Compound 1


, or one or more members selected from the group consisting of hydrates thereof and solvates thereof.

33. The pharmaceutical composition of any one of claims 1-32, wherein b) is



Compound 1


34. The pharmaceutical composition of any one of claims 1-31, wherein b) is the hydroquinone form of



Compound 1

, or one or more members selected from the group consisting of hydrates thereof, and solvates thereof.

35. The pharmaceutical composition of any one of claims 1-31, and 34, wherein b) is the hydroquinone form of



Compound 1

36. The pharmaceutical composition of any one of claims 1-35, wherein the pharmaceutical composition is an oral pharmaceutical composition.

37. The pharmaceutical composition of claim 36, wherein the oral pharmaceutical composition is in a capsule.

38. The pharmaceutical composition of claim 37, wherein the capsule is a hard gelatin capsule.

39. The pharmaceutical composition of claim 40, wherein any water content in the gelatin of the hard gelatin capsule is about 20% or less, about 10% to about 20%, or about 11% to about 16%, when stored for about one month, about one or more months, about three months, or about three or more months.

40. The pharmaceutical composition of claims 1-36, wherein the oral pharmaceutical composition is a solution.

41. The pharmaceutical composition of any one of claims 36-40, wherein the pharmaceutical composition has bioavailability when administered with a Low-fat Food for Testing which differs from bioavailability when administered with a Medium-fat Food for Testing by a percentage difference of about 90% or less, about 85% or less, about 80% or less, about 75% or less, about 70% or less, or about 65% or less; optionally wherein the pharmaceutically acceptable oil is a Labrafac-like oil (preferably, Labrafac WL 1349), the propylene glycol laurates composition is Lauroglycol 90, and the polysorbate 80 is Tween 80; additionally optionally wherein the ratio by weight of the pharmaceutically acceptable oil to the propylene glycol laurates composition to the polysorbate 80 is about 65:20:15.

42. The pharmaceutical composition of any one of claims 36-40, wherein oral bioavailability for Medium-fat Food for Testing is about 45% to about 55%, about 40 to about 50%, about 49%; optionally wherein the pharmaceutically acceptable oil is a Labrafac-like oil (preferably, Labrafac WL 1349), the propylene glycol laurates composition is Lauroglycol 90, and the polysorbate 80 is Tween 80; additionally optionally wherein the ratio by weight of the Labrafac-like oil (preferably, Labrafac WL 1349) to the Lauroglycol 90 to the polysorbate 80 is about 60:20:20.

43. The pharmaceutical composition of claims 36-40 and 42, wherein oral bioavailability for Low-fat Food for Testing is about 10% to about 15%, in some embodiments, about 13%; optionally wherein the pharmaceutically acceptable oil is a Labrafac-like oil (preferably, Labrafac WL 1349), the propylene glycol laurates composition is Lauroglycol 90, and the polysorbate 80 is Tween 80; additionally optionally wherein the ratio by weight of the pharmaceutically acceptable oil to the propylene glycol laurates composition to the polysorbate 80 is about 60:20:20.

44. The pharmaceutical composition of any one of claims 36-40, wherein oral bioavailability for Medium-fat Food for Testing is about 15% to about 40%, 25% to about 35%, about 29%; optionally wherein the ratio by weight of the pharmaceutically acceptable oil to the propylene glycol laurates composition to the polysorbate 80 is about 65:20:15; and additionally optionally wherein the pharmaceutically acceptable oil is a Labrafac-like oil (Labrafac WL 1349 or Labrafac PG), cottonseed oil, soybean oil, olive oil, or corn oil; the propylene glycol laurates composition is Lauroglycol 90; and the polysorbate 80 is Tween 80.

45. The pharmaceutical composition of claims 36-40 and 44, wherein oral bioavailability for Low-fat Food for Testing is about 10% to about 15%, in some embodiments, about 14%; optionally wherein the pharmaceutically acceptable oil is a Labrafac-like oil (preferably, Labrafac WL 1349), the propylene glycol laurates composition is Lauroglycol 90, and the polysorbate 80 is Tween 80; additionally optionally wherein the ratio by weight of the pharmaceutically acceptable oil to the propylene glycol laurates composition to the polysorbate 80 is about 65:20:15.

46. The pharmaceutical composition of any one of claims 1-45, wherein the pharmaceutical composition comprises a flavorant.

47. The pharmaceutical composition of claims 1-46, wherein the amount of the flavorant is less than 1% wt/wt%.

48. The pharmaceutical composition of claims 46 and 47, wherein the flavorant is selected from the group consisting of natural flavors, natural fruit flavors, artificial flavors, artificial fruit flavors, flavor enhancers and mixtures thereof.

49. The pharmaceutical composition of any one of claims 46-48, wherein the flavorant is oil soluble.

50. The pharmaceutical composition of any one of claims 46-49, wherein the flavorant has flavor selected from the group consisting of raspberries, punch, cherry, strawberries, and blueberries.

51. The pharmaceutical composition of any one of claims 46-50, wherein the flavorant has odor selected from the group consisting of raspberries, punch, cherry, strawberries, and blueberries.

52. The pharmaceutical composition of any one of claims 1-45, wherein the pharmaceutical composition does not comprise a flavorant.

53. The pharmaceutical composition of any one of claims 1-35, wherein the pharmaceutical composition is a transdermal pharmaceutical composition.

54. A method of preparing a pharmaceutical composition of any one of claims 1-53, comprising:

step a) mixing about 48 wt/wt% to about 70 wt/wt% of a pharmaceutically acceptable oil, about 12 wt/wt% to about 25 wt/wt% of a propylene glycol laurates composition, about 8 wt/wt% to

about 20 wt/wt% of a polysorbate 80, and 0 wt/wt% to about 2% wt/wt% of an optional flavorant;

step b) adding about 1 wt/wt% to about 15%, of **Compound 1** and/or the hydroquinone form thereof, wherein **Compound 1** and/or the hydroquinone form thereof is optionally a hydrate and/or solvate thereof; to the mixture from step a) and mixing,

and wherein when the **Compound 1**, and/or hydroquinone thereof, is in the form of a hydrate and/or solvate, then the about 1 wt/wt% to about 15 wt/wt% of **Compound 1** and/or its hydroquinone, does not include the weight of any the water in the hydrate or the weight of the solvent in the solvate; and

wherein the wt/wt% of **Compound 1**, the pharmaceutically acceptable oil, the propylene glycol laurates composition, the polysorbate 80, and optional flavorant total 100%.

55. The method of claim 54, wherein step b) occurs at room temperature.

56. The method of claim 54 or 55, wherein **Compound 1** in step b) is a polymorph of an anhydrate of 2,3,5-trimethyl-6-nonylcyclohexa-2,5-diene-1,4-dione, wherein a powder X-ray diffraction pattern for the polymorph comprises characteristic peaks at least at the following angular positions, wherein the angular positions may vary by ± 0.2 : 4.10, 12.12, and 16.14, and wherein the characteristic peaks are measured with a Cu K α 1 source, a wavelength of 1.540598 Å, and at room temperature or at a temperature of 23-25 °C.

57. The method of any one of claims 54-56, wherein the mixing occurs until a clear formulation solution forms.

58. The method of claim 57, wherein the solution is assessed visually and under microscope for the absence of crystals of **Compound 1**.

59. The method of any one of claims 54-58, wherein **Compound 1** is added and mixed under light protection or yellow light.

60. A pharmaceutical composition prepared by the method of any one of claims 54-58.

61. A method for treating or suppressing a disease or disorder selected from the group consisting of an α -synucleinopathy, a tauopathy, an autistic spectrum disorder, a pervasive developmental disorder, a liver disease, liver damage, dementia, and reperfusion injury, comprising administering a pharmaceutical composition of any one of claims 1-53 or a pharmaceutical composition of claim 60 to a patient in need thereof.

62. The method of claim 61, wherein the α -synucleinopathy is selected from the group consisting of: Parkinson's Disease (idiopathic and genetic), Parkinson's Disease with dementia (PDD), multisystem atrophy (MSA), Frontotemporal Dementia, Dementia with Lewy Bodies (DLB), Gaucher's disease (GD), Neurodegeneration with Brain Iron Accumulation (NBIA), and neuroaxonal dystrophies (PLA2G6-associated neurodegeneration).

63. The method of claim 62, wherein the Parkinson's Disease is that wherein the patient has a mutation in one or more of the following genes: *MAPT* (Microtubule-associated protein tau), *PRKN* (parkin), *PINK1* (PINK1), *LRRK2* (leucine-rich repeat kinase 2), *GBA* (glucocerebrosidase), *SNCA* (alpha synuclein), *PARK7* (DJ-1), and/or *UCHL1* (ubiquitin carboxyl-terminal esterase L1).

64. The method of claim 61, wherein the tauopathy is selected from the group consisting of: Alzheimer's disease, dementia pugilistica, Guam Amyotrophic lateral sclerosis-Parkinsonism-Dementia (Guam ALS/PD), Pick Disease, Argyrophilic grain dementia, Nieman-Pick type C, Subacute sclerosing panencephalitis (SSPE), Progressive supranuclear palsy (PSP), multisystem atrophy (MSA), Corticobasoganglionic degeneration, Frontotemporal dementia with parkinsonism-17 (FTDP-17), Postencephalitic Parkinsonism (PEP), and Autosomal recessive Parkinsonism.

65. The method of claim 61, wherein the liver disease is selected from the group consisting of NASH/NAFL, pediatric NAFLD, alcoholic hepatitis, cholestatic liver disease, viral hepatitis, drug-induced liver toxicity, hemochromatosis, Wilson's disease, liver transplant reperfusion injury, hepatic insufficiency where the hepatic insufficiency is due to injury, SIRS, sepsis, or severe illness; and drug-induced liver toxicity, such as cisplatin-induced liver toxicity and acetaminophen-induced liver toxicity.

66. The method of claim 61, wherein the autistic spectrum disorder or pervasive developmental disorder is selected from the group consisting of autistic disorder, Asperger's syndrome, childhood disintegrative disorder (CDD), Rett's disorder, Rett syndrome (RTT), PDD-not otherwise specified (PDD-NOS), and attention deficit/hyperactivity disorder (ADHD).

67. The method of claim 61, wherein the reperfusion injury is selected from the group consisting of cardiac reperfusion injury, renal reperfusion injury, and stroke.

68. The method of any one of claims 61-67, wherein the pharmaceutical composition is administered orally.

69. The method of any one of claims 61-68, wherein the pharmaceutical composition is administered for at least 24 weeks or for at least 28 weeks.
70. The method of claim 69, wherein the pharmaceutical composition is administered at a dose selected from 150 mg BID and 250 mg BID.
71. The method of claim 70, wherein the pharmaceutical composition is administered at a dose of 10 capsules total per day, wherein the weight of **Compound 1** in each capsule is 50 mg.
72. The method of claim 70, wherein the pharmaceutical composition is administered at a dose of 6 capsules total per day, wherein the weight of **Compound 1** in each capsule is 50 mg.
73. The method of any one of claims 61-72, wherein the patient does not experience any emergent serious adverse events (TEAEs), no TEAEs leading to discontinuation of the pharmaceutical composition, and/or no TEAEs leading to death.
74. The method of any one of claims 61-68, wherein the pharmaceutical composition is administered at a dose ranging from about 100 mg to about 1000 mg daily dose for a period of at least 14 days.
75. The method of any one of claims 61-74, wherein the pharmaceutical composition does not comprise the optional flavorant and the pharmaceutical composition is administered in or with food.
76. The method of claim 75, wherein the food is a Low-fat Food or a Medium fat Food.
77. The method of claim 76, wherein the food is a Low-fat Food.
78. The method of claim 76, wherein the food is a Medium fat Food.