


Quinoline derivatives

The present invention is concerned with novel quinoline derivatives, their manufacture, pharmaceutical compositions containing them and their use as medicaments. The active compounds of the present invention are useful in treating obesity and other disorders.

5 In particular, the present invention relates to compounds of the general formula




wherein

- R^1 is selected from the group consisting of hydrogen,
lower alkyl, lower alkenyl,
10 cycloalkyl or lower cycloalkylalkyl, wherein the cycloalkyl ring may be
unsubstituted or substituted by one or two groups selected from the group
consisting of lower alkyl, lower hydroxyalkyl and lower alkoxyalkyl,
lower hydroxyalkyl,
lower alkoxyalkyl, and
15 lower heterocyclalkyl, wherein the heterocyclalkyl ring may be unsubstituted or
substituted by one or two groups selected from lower alkyl and halogen;
- R^2 is selected from the group consisting of hydrogen,
lower alkyl, lower alkenyl,
cycloalkyl or lower cycloalkylalkyl, wherein the cycloalkyl ring may be
20 unsubstituted or substituted by one or two groups selected from the group
consisting of lower alkyl, lower hydroxyalkyl and lower alkoxyalkyl,
lower hydroxyalkyl,
lower alkoxyalkyl, and


lower heterocyclalkyl, wherein the heterocyclalkyl ring may be unsubstituted or substituted by one or two groups selected from lower alkyl and halogen; or

- 5 R^1 and R^2 together with the nitrogen atom to which they are attached form a 4-, 5-, 6- or 7-membered saturated or partly unsaturated heterocyclic ring optionally containing a further heteroatom selected from nitrogen, oxygen or sulfur, said saturated or partly unsaturated heterocyclic ring being unsubstituted or substituted by one, two or three groups independently selected from the group consisting of lower alkyl, halogen, halogenalkyl, cyano, hydroxy, hydroxyalkyl, lower alkoxy, oxo, phenyl, benzyl, pyridyl and carbamoyl;


- 10 A is selected from



A 1



A 2



A 3

wherein

- m is 0, 1 or 2;
 15 R^3 is lower alkyl;

n is 0, 1 or 2;

R^7 is lower alkyl;

p is 0, 1 or 2;

q is 0, 1 or 2;

- 20 R^5 is hydrogen or lower alkyl;

and pharmaceutically acceptable salts thereof.

The compounds of formula I are antagonists and/or inverse agonists at the histamine 3 receptor (H3 receptor).

Histamine (2-(4-imidazolyl) ethylamine) is one of the aminergic neurotransmitters which is widely distributed throughout the body, e. g. the gastrointestinal tract (Burks 1994 in Johnson L.R. ed., Physiology of the Gastrointestinal Tract, Raven Press, NY, pp. 211 – 242). Histamine regulates a variety of digestive pathophysiological events like
5 gastric acid secretion, intestinal motility (Leurs et al., Br J. Pharmacol. 1991, 102, pp 179-185), vasomotor responses, intestinal inflammatory responses and allergic reactions (Raithel et al., Int. Arch. Allergy Immunol. 1995, 108, 127-133). In the mammalian brain, histamine is synthesized in histaminergic cell bodies which are found centrally in the tuberomammillary nucleus of the posterior basal hypothalamus. From there, the cell
10 bodies project to various brain regions (Panula et al., Proc. Natl. Acad. Sci. USA 1984, 81, 2572-2576; Inagaki et al., J. Comp. Neurol 1988, 273, 283 – 300).

According to current knowledge, histamine mediates all its actions in both the central nervous system (CNS) and in the periphery through four distinct histamine receptors, the histamine H1, H2, H3 and H4 receptors.

15 H3 receptors are predominantly localized in the CNS. As an autoreceptor, H3 receptors constitutively inhibit the synthesis and secretion of histamine from histaminergic neurons (Arrang et al., Nature 1983, 302, 832-837; Arrang et al., Neuroscience 1987, 23, 149-157). As heteroreceptors, H3 receptors also modulate the release of other neurotransmitters such as acetylcholine, dopamine, serotonin and
20 norepinephrine among others in both the central nervous system and in peripheral organs, such as lungs, cardiovascular system and gastrointestinal tract (Clapham & Kilpatrick, Br. J. Pharmacol. 1982, 107, 919- 923; Blandina et al. in The Histamine H3 Receptor (Leurs RL and Timmermann H eds, 1998, pp 27-40, Elsevier, Amsterdam, The Netherlands). H3 receptors are constitutively active, meaning that even without
25 exogenous histamine, the receptor is tonically activated. In the case of an inhibitory receptor such as the H3 receptor, this inherent activity causes tonic inhibition of neurotransmitter release. Therefore it may be important that a H3R antagonist would also have inverse agonist activity to both block exogenous histamine effects and to shift the receptor from its constitutively active (inhibitory) form to a neutral state.


30 The wide distribution of H3 receptors in the mammalian CNS indicates the physiological role of this receptor. Therefore the therapeutic potential as a novel drug development target in various indications has been proposed.

The administration of H3R ligands - as antagonists, inverse agonists, agonists or partial agonists - may influence the histamine levels or the secretion of neurotransmitters
35 in the brain and the periphery and thus may be useful in the treatment of several disorders. Such disorders include obesity, (Masaki et al; Endocrinol. 2003, 144, 2741-

2748; Hancock et al., European J. of Pharmacol. 2004, 487, 183-197), cardiovascular disorders such as acute myocardial infarction, dementia and cognitive disorders such as attention deficit hyperactivity disorder (ADHD) and Alzheimer's disease, neurological disorders such as schizophrenia, depression, epilepsy, Parkinson's disease, and seizures
5 or convulsions, sleep disorders, narcolepsy, pain, gastrointestinal disorders, vestibular dysfunction such as Morbus Meniere, drug abuse and motion sickness (Timmermann, J. Med. Chem. 1990, 33, 4-11).

It is therefore an object of the present invention to provide selective, directly acting H3 receptor antagonists respectively inverse agonists. Such antagonists / inverse agonists
10 are useful as therapeutically active substances, particularly in the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors.

In a first aspect, there is provided compounds of the general formula



wherein


20 R^1 is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, cycloalkyl or lower cycloalkylalkyl, wherein the cycloalkyl ring may be unsubstituted or substituted by one or two groups selected from the group consisting of lower alkyl, lower hydroxyalkyl and lower alkoxyalkyl, lower hydroxyalkyl,
25 lower alkoxyalkyl, and lower heterocyclalkyl, wherein the heterocycl ring may be unsubstituted or substituted by one or two groups selected from lower alkyl and halogen;

R^2 is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, cycloalkyl or lower cycloalkylalkyl, wherein the cycloalkyl ring may be
30 unsubstituted or substituted by one or two groups selected from the group consisting of lower alkyl, lower hydroxyalkyl and lower alkoxyalkyl, lower hydroxyalkyl, lower alkoxyalkyl, and lower heterocyclalkyl, wherein the heterocycl ring may be unsubstituted or


substituted by one or two groups selected from lower alkyl and halogen; or

R¹ and R² together with the nitrogen atom to which they are attached form a 4-, 5-, 6- or 7-membered saturated or partly unsaturated heterocyclic ring optionally containing a further heteroatom selected from nitrogen, oxygen or sulfur, said saturated or partly unsaturated heterocyclic ring being unsubstituted or substituted by one, two or three groups independently selected from the group consisting of lower alkyl, halogen, halogenalkyl, cyano, hydroxy, hydroxyalkyl, lower alkoxy, oxo, phenyl, benzyl, pyridyl and carbamoyl;


A is selected from



A 1



A 2



A 3

wherein

m is 0, 1 or 2;

R³ is lower alkyl;

n is 0, 1 or 2;

R⁷ is lower alkyl;

p is 0, 1 or 2;


q is 0, 1 or 2;

R⁵ is hydrogen or lower alkyl;

and pharmaceutically acceptable salts thereof.

In a second aspect, there is provided a process for the manufacture of compounds as defined above, which process comprises


a) reacting a compound of the formula II




wherein R is lower alkyl,
with an alcohol of the formula III



wherein A is as defined in claim 1,
in the presence of a trialkylphosphine or triphenylphosphine and of a diazo compound
to obtain a compound of the formula IV




and converting the ester of formula IV into the acid of formula V



under acidic or basic conditions,
and coupling the compound of formula V with an amine of the formula VI



wherein R^1 and R^2 are as defined in claim 1, with the help of an coupling agent under basic conditions to obtain a compound of the formula I



I

wherein A, R^1 and R^2 are as defined in claim 1, and if desired, converting the compound obtained into a pharmaceutically acceptable salt.

In a third aspect, there is provided compounds as defined above when manufactured by a process according to the second aspect.

In a fourth aspect, there is provided pharmaceutical compositions comprising a compound as defined above and a pharmaceutically acceptable carrier and/or adjuvant.

In a fifth aspect, there is provided pharmaceutical compositions according to the fourth aspect for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors.

In a sixth aspect, there is provided compounds as defined above for use as therapeutically active substances.

In a seventh aspect, there is provided compounds as defined above for use as therapeutically active substances for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors.

In an eighth aspect, there is provided the use of compounds as defined above for the preparation of medicaments for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors.

In a ninth aspect, there is provided the use according to the eighth aspect for the treatment and/or prevention of obesity.

In the present description the term "alkyl", alone or in combination with other groups, refers to a branched or straight-chain monovalent saturated aliphatic hydrocarbon radical of one to twenty carbon atoms, preferably one to sixteen carbon atoms, more preferably one to ten carbon atoms.

The term "lower alkyl" or " C_1-C_8 -alkyl", alone or in combination, signifies a straight-chain or branched-chain alkyl group with 1 to 8 carbon atoms, preferably a straight or branched-chain alkyl group with 1 to 6 carbon atoms and particularly preferred a straight or branched-chain alkyl group with 1 to 4 carbon atoms. Examples of straight-chain and branched C_1-C_8 alkyl groups are methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tert.-butyl, the isomeric pentyls, the isomeric hexyls, the isomeric heptyls and the isomeric octyls, preferably methyl and ethyl and most preferred methyl.

The term "lower alkenyl" or " C_2-C_8 -alkenyl", alone or in combination, signifies a straight-chain or branched alkyl group comprising an olefinic bond and up to 8, preferably up to 6, particularly preferred up to 4 carbon atoms. Examples of alkenyl groups are ethenyl, 1-propenyl, 2-propenyl, isopropenyl, 1-butenyl, 2-butenyl, 3-butenyl and isobutenyl. A preferred example is 2-propenyl.

The term "alkoxy" refers to the group $R'-O-$, wherein R' is alkyl. The term "lower alkoxy" refers to the group $R'-O-$, wherein R' is lower alkyl and the term "lower alkyl" has the previously given significance. Examples of lower alkoxy groups are e.g. methoxy, ethoxy, n-propoxy, isopropoxy, n-butoxy, isobutoxy, sec. butoxy and tert.butoxy, preferably methoxy and ethoxy and most preferred methoxy.

The term "lower alkoxyalkyl" or "alkoxy- C_1-C_8 -alkyl" refers to lower alkyl groups as defined above wherein at least one of the hydrogen atoms of the lower alkyl group is

replaced by an alkoxy group as defined above. Among the preferred lower alkoxyalkyl groups are methoxymethyl, methoxyethyl and ethoxymethyl, with methoxymethyl being especially preferred.

The term "halogen" refers to fluorine, chlorine, bromine and iodine, with fluorine,
5 chlorine and bromine being preferred.

The term "lower halogenalkyl" or "halogen-C₁-C₈-alkyl" refers to lower alkyl groups as defined above wherein at least one of the hydrogen atoms of the lower alkyl group is replaced by a halogen atom, preferably fluoro or chloro, most preferably fluoro. Among the preferred halogenated lower alkyl groups are trifluoromethyl,
10 difluoromethyl, fluoromethyl and chloromethyl, with trifluoromethyl being especially preferred.

The term "lower halogenalkoxy" or "halogen-C₁-C₈-alkoxy" refers to lower alkoxy groups as defined above wherein at least one of the hydrogen atoms of the lower alkoxy group is replaced by a halogen atom, preferably fluoro or chloro, most preferably fluoro.
15 Among the preferred halogenated lower alkyl groups are trifluoromethoxy, difluoromethoxy, fluormethoxy and chloromethoxy, with trifluoromethoxy being especially preferred.

The term "lower hydroxyalkyl" or "hydroxy-C₁-C₈-alkyl" refers to lower alkyl groups as defined above wherein at least one of the hydrogen atoms of the lower alkyl
20 group is replaced by a hydroxy group. Examples of lower hydroxyalkyl groups are hydroxymethyl or hydroxyethyl.

The term "cycloalkyl" or "C₃-C₇-cycloalkyl" means a cycloalkyl ring containing 3 to 7 carbon atoms, such as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl or cycloheptyl. The cycloalkyl ring may be optionally substituted as defined herein.
25 Especially preferred is cyclopropyl.

The term "lower cycloalkylalkyl" or "C₃-C₇-cycloalkyl-C₁-C₈-alkyl" refers to lower alkyl groups as defined above wherein at least one of the hydrogen atoms of the lower alkyl group is replaced by a cycloalkyl group as defined above. Examples of preferred lower cycloalkylalkyl groups are cyclopropylmethyl or cyclopropylmethyl wherein the
30 cyclopropyl group is substituted by lower alkyl, lower hydroxyalkyl or lower alkoxyalkyl, preferably methoxymethyl.

The term "heterocyclyl" means a monovalent saturated or partly unsaturated ring incorporating one, two, or three heteroatoms chosen from nitrogen, oxygen or sulfur. The heterocyclyl ring may be optionally substituted as defined herein. Examples of

heterocyclyl moieties include azetidiny, oxetanyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, dihydropyridyl, piperidyl, piperazinyl, morpholinyl, or thiomorpholinyl, azepinyl, dihydropyrrolyl, pyrrolidinyl, pyrazolidinyl, imidazolinyl, dihydrofuryl, tetrahydrofuryl, dihydropyranyl or tetrahydropyranyl. Especially preferred are oxetanyl, piperidyl and morpholinyl.

The term "lower heterocyclalkyl" or "heterocycl-C₁-C₈-alkyl" refers to lower alkyl groups as defined above wherein at least one of the hydrogen atoms of the lower alkyl group is replaced by a heterocycl group as defined above. An examples of a preferred lower heterocyclalkyl group is 3-fluorooxetan-3-yl.

The term "a 4-, 5-, 6- or 7-membered saturated or partly unsaturated heterocyclic ring optionally containing a further heteroatom selected from nitrogen, oxygen or sulfur" refers to a saturated or partly unsaturated N-heterocyclic ring, which may optionally contain a further nitrogen, oxygen or sulfur atom, such as 2,5-dihydropyrrolidinyl, pyrrolidinyl, imidazolidinyl, pyrazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, piperidyl, piperazinyl, morpholinyl, thiomorpholinyl, or 3,6-dihydro-2H-pyridinyl. The heterocyclic ring may be unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, lower alkoxy and halogen.

The term "carbamoyl" refers to the group -CO-NH₂.


The term "pharmaceutically acceptable salts" refers to those salts which retain the biological effectiveness and properties of the free bases or free acids, which are not biologically or otherwise undesirable. The salts are formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, preferably hydrochloric acid, and organic acids such as acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, salicylic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, N-acetylcystein and the like. In addition these salts may be prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from an inorganic base include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium salts and the like. Salts derived from organic bases include, but are not limited to salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, lysine, arginine, N-ethylpiperidine, piperidine, polymine resins and the like. The compound of formula I can also be present in the form of zwitterions.

Particularly preferred pharmaceutically acceptable salts of compounds of formula I are the hydrochloride salts.

The compounds of formula I can also be solvated, e.g. hydrated. The solvation can be effected in the course of the manufacturing process or can take place e.g. as a
5 consequence of hygroscopic properties of an initially anhydrous compound of formula I (hydration). The term pharmaceutically acceptable salts also includes physiologically acceptable solvates.

“Isomers” are compounds that have identical molecular formulae but that differ in the nature or the sequence of bonding of their atoms or in the arrangement of their
10 atoms in space. Isomers that differ in the arrangement of their atoms in space are termed “stereoisomers”. Stereoisomers that are not mirror images of one another are termed “diastereoisomers”, and stereoisomers that are non-superimposable mirror images are termed “enantiomers”, or sometimes optical isomers. A carbon atom bonded to four nonidentical substituents is termed a “chiral center”.

15 In detail, the present invention relates to compounds of the general formula




wherein

- R¹ is selected from the group consisting of hydrogen,
lower alkyl, lower alkenyl,
20 cycloalkyl or lower cycloalkylalkyl, wherein the cycloalkyl ring may be
unsubstituted or substituted by one or two groups selected from the group
consisting of lower alkyl, lower hydroxyalkyl and lower alkoxyalkyl,
lower hydroxyalkyl,
lower alkoxyalkyl, and
25 lower heterocyclalkyl, wherein the heterocyclalkyl ring may be unsubstituted or
substituted by one or two groups selected from lower alkyl and halogen;
- R² is selected from the group consisting of hydrogen,
lower alkyl, lower alkenyl,
cycloalkyl or lower cycloalkylalkyl, wherein the cycloalkyl ring may be
30 unsubstituted or substituted by one or two groups selected from the group
consisting of lower alkyl, lower hydroxyalkyl and lower alkoxyalkyl,

lower hydroxyalkyl,
 lower alkoxyalkyl, and
 lower heterocyclalkyl, wherein the heterocyclalkyl ring may be unsubstituted or substituted by one or two groups selected from lower alkyl and halogen; or

- 5 R^1 and R^2 together with the nitrogen atom to which they are attached form a 4-, 5-, 6- or 7-membered saturated or partly unsaturated heterocyclic ring optionally containing a further heteroatom selected from nitrogen, oxygen or sulfur, said saturated or partly unsaturated heterocyclic ring being unsubstituted or substituted by one, two or three groups independently selected from the group
 10 consisting of lower alkyl, halogen, halogenalkyl, cyano, hydroxy, hydroxyalkyl, lower alkoxy, oxo, phenyl, benzyl, pyridyl and carbamoyl;

A is selected from



15 wherein

m is 0, 1 or 2;

R^3 is lower alkyl;

n is 0, 1 or 2;

R^7 is lower alkyl;

20 p is 0, 1 or 2;

q is 0, 1 or 2;

R^5 is hydrogen or lower alkyl;

and pharmaceutically acceptable salts thereof.

Preferred compounds of formula I according to the present invention are those,
 25 wherein R^1 is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl,

cycloalkyl, lower cycloalkylalkyl, wherein the cycloalkyl ring may be unsubstituted or substituted by one or two groups selected from lower alkyl, lower hydroxyalkyl or lower alkoxyalkyl, lower hydroxyalkyl, lower alkoxyalkyl, and lower heterocyclalkyl, wherein the heterocycl ring may be unsubstituted or substituted with one or two groups
5 selected from lower alkyl or halogen, and
R² is hydrogen or lower alkyl.

More preferred are compounds of formula I according to present invention, wherein R¹ is selected from the group consisting of lower alkyl, cycloalkyl, lower cycloalkylalkyl, wherein the cycloalkyl ring may be unsubstituted or substituted by lower
10 alkoxyalkyl,
lower alkoxyalkyl, and lower heterocyclalkyl wherein the heterocycl ring may be unsubstituted or substituted with one or two groups selected from lower alkyl or halogen,
and R² is hydrogen or lower alkyl.

15 Especially preferred are those compounds of formula I, wherein R¹ and R² are lower alkyl.

Another group of preferred compounds of formula I according to present invention are those, wherein R¹ and R² together with the nitrogen atom to which they are attached form a 4-, 5-, 6- or 7-membered saturated or partly unsaturated heterocyclic
20 ring optionally containing a further heteroatom selected from nitrogen, oxygen or sulfur, said heterocyclic ring being unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, halogen, halogenalkyl, cyano, hydroxy, hydroxyalkyl, lower alkoxy, oxo, phenyl, benzyl, pyridyl and carbamoyl.


Preferred compounds of formula I are those, wherein R¹ and R² together with the
25 nitrogen atom to which they are attached form a 4-, 5-, 6- or 7-membered saturated or partly unsaturated heterocyclic ring optionally containing a further heteroatom selected from nitrogen, oxygen or sulfur, said heterocyclic ring being unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, halogen, halogenalkyl, cyano, hydroxy, lower alkoxy, oxo, phenyl, benzyl, pyridyl and carbamoyl.

30 More preferred are those compounds of formula I, wherein R¹ and R² together with the nitrogen atom to which they are attached form a 4-, 5-, or 6- or 7-membered saturated or partly unsaturated heterocyclic ring optionally containing a further heteroatom selected from nitrogen, oxygen or sulfur, said heterocyclic ring being unsubstituted or substituted by one, two or three groups independently selected from
35 lower alkyl, halogen, halogenalkyl, cyano, hydroxy, lower alkoxy, and oxo.

Even more preferred are compounds of formula I, wherein R^1 and R^2 together with the nitrogen atom to which they are attached form a heterocyclic ring selected from the group consisting of morpholine, piperidine, 2,5-dihydropyrrole, pyrrolidine, azepane, piperazine, azetidine, thiomorpholine and 3,6-dihydro-2H-pyridine, said heterocyclic
 5 ring being unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, halogen, halogenalkyl, cyano, hydroxy, lower alkoxy, and oxo.

Especially preferred are compounds of formula I, wherein R^1 and R^2 together with the nitrogen atom to which they are attached form a heterocyclic ring selected from morpholinyl, thiomorpholinyl, 1,1-dioxothiomorpholinyl, 3,6-dihydro-2H-pyridinyl,
 10 piperidinyl, 2-methylpiperidinyl, 3-methylpiperidinyl, 4-hydroxypiperidinyl, 4,4-difluoropiperidinyl, 2,5-dihydropyrrolyl, 4-methylpiperidinyl, pyrrolidinyl, 2-methylpyrrolidinyl, 2-trifluoromethylpyrrolidinyl, 2-cyanopyrrolidinyl, 3-hydroxypyrrolidinyl and azetidiny.

Furthermore, compounds of formula I according to the present invention, wherein
 15 A signifies




A 1

and wherein m is 0, 1 or 2, and R^3 is lower alkyl, are preferred.

Within this group, those compounds of formula I are preferred, wherein m is 0, thus meaning pyrrolidine groups are preferred.

20 A further preferred group includes those compounds of formula I, wherein m is 1, thus meaning piperidine groups are also preferred.


Also preferred are compounds of formula I according to the present invention, wherein A signifies



A 2

and wherein n is 0, 1 or 2; and R⁴ is lower alkyl, , with those compounds, wherein n is 0, thus meaning pyrrolidine derivatives, being more preferred.

Further preferred compounds of formula I according to the present invention are those, wherein A signifies



5

A 3

wherein p is 0, 1 or 2, q is 0, 1 or 2, and R⁵ is hydrogen or lower alkyl.

Within this group, those compounds of formula I are preferred, wherein p is 1, thus meaning piperidine groups are preferred. Especially preferred are those compounds of formula I, wherein p is 1 and q is 1.

10 Furthermore, compounds wherein R⁵ is hydrogen, are preferred.

Examples of preferred compounds of formula I are the following:

[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(4-methoxy-piperidin-1-yl)-methanone,

15 (2,5-dihydro-pyrrol-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone,

6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid ethyl-methyl-amide,

[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(2-methyl-pyrrolidin-1-yl)-methanone,

20 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid tert-butylamide,

6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid cyclopropylmethyl-propyl-amide,

6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(2-methyl-piperidin-1-yl)-methanone,

(4-hydroxy-piperidin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone,

[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(3-methyl-piperidin-1-yl)-methanone,

5 (3,4-dihydro-1H-isoquinolin-2-yl)-[6-(3-piperidin-1-yl-propoxy)-naphthalen-2-yl]-methanone 1:1 hydrochloride,

[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-thiomorpholin-4-yl-methanone,

10 [6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(2-trifluoromethyl-pyrrolidin-1-yl)-methanone,

6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid ethyl-(2-methoxyethyl)-amide,

azetidin-1-yl-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone,

15 (3,6-dihydro-2H-pyridin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone,

6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid (3-fluoro-oxetan-3-ylmethyl)-amide,

6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid (1-methoxymethyl-cyclopropylmethyl)-amide,

20 {6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid [2-(tetrahydropyran-4-yl)-ethyl]-amide,

[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-morpholin-4-yl-methanone,

6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid cyclohexyl-methyl-amide,

25 (4,4-difluoro-piperidin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone,

(3-hydroxy-pyrrolidin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone,

[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-pyrrolidin-1-yl-methanone,

(R)-1-[6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile,

5 (1,1-dioxo-thiomorpholin-4-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone,

6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid dimethylamide

(2,5-dihydro-pyrrol-1-yl)-[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-methanone,

10 ([6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-(4-methoxy-piperidin-1-yl)-methanone,

6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid ethyl-methylamide,

([6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-(4-methyl-piperidin-1-yl)-methanone,

15 [6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-(2-methyl-pyrrolidin-1-yl)-methanone,

6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid tert-butylamide,

6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid cyclopropylmethyl-propyl-amide,

20 {[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-(2-methyl-piperidin-1-yl)-methanone,

[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-(3-methyl-piperidin-1-yl)-methanone,

25 6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid isopropyl-methylamide,

[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-thiomorpholin-4-yl-methanone,

[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-(2-trifluoromethyl-pyrrolidin-1-yl)-methanone,

6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid ethyl-(2-methoxyethyl)-amide,

5 azetidin-1-yl-[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-methanone,

(3,6-dihydro-2H-pyridin-1-yl)-[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-methanone,

6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid (3-fluoro-oxetan-3-ylmethyl)-amide,

10 6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid (1-methoxymethyl-cyclopropylmethyl)-amide,

6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid [2-(tetrahydropyran-4-yl)-ethyl]-amide,

15 (4,4-difluoro-piperidin-1-yl)-[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-methanone,

[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-morpholin-4-yl-methanone,

(4-methoxy-piperidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,

20 (4-methyl-piperidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone

morpholin-4-yl-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,

(2-methyl-pyrrolidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,

25 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid cyclopropylmethyl-propyl-amide,

[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-(2-trifluoromethyl-pyrrolidin-1-yl)-methanone,

- (2,5-dihydro-pyrrol-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
- 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid ethyl-(2-methoxy-ethyl)-amide,
- 5 azetidin-1-yl-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
- (3,6-dihydro-2H-pyridin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
- (4,4-difluoro-piperidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
- 10 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid cyclohexyl-methylamide,
- 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid (1-methoxymethyl-cyclopropylmethyl)-amide,
- 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid (3-fluoro-oxetan-3-ylmethyl)-amide,
- 15 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid [2-(tetrahydro-pyran-4-yl)-ethyl]-amide,
- 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid (tetrahydro-pyran-4-yl)-amide,
- 20 (2-methyl-piperidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
- [6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-pyrrolidin-1-yl-methanone,
- (R)-1-[6-(3-piperidin-1-yl-propoxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile,
- 25 (1,1-dioxo--thiomorpholin-4-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
- (4-methyl-piperidin-1-yl)-{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-methanone,

{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-pyrrolidin-1-yl-methanone,

(R)-1-{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carbonyl}-pyrrolidine-2-carbonitrile,

5 (1,1-dioxo-thiomorpholin-4-yl)-{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-methanone,

(4-methoxy-piperidin-1-yl)-{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-methanone,

10 {6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-morpholin-4-yl-methanone,

azetidin-1-yl-{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-methanone,

6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid [2-(tetrahydro-pyran-4-yl)-ethyl]-amide,

15 6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid (3-fluoro-oxetan-3-ylmethyl)-amide,

(2-methyl-pyrrolidin-1-yl)-{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-methanone,

(S)-1-[6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile,

20 [6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(4-methyl-piperidin-1-yl)-methanone,

(4-hydroxymethyl-piperidin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone,

6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid isobutyl-amide,

25 6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid cyclohexyl-methyl-amide,

and pharmaceutically acceptable salts thereof.

Particularly preferred compounds of formula I of the present invention are the following:

[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(4-methoxy-piperidin-1-yl)-methanone,

5 [6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(2-methyl-pyrrolidin-1-yl)-methanone,

[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(2-trifluoromethyl-pyrrolidin-1-yl)-methanone,

azetidin-1-yl-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone,

10 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid (3-fluoro-oxetan-3-ylmethyl)-amide,

6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid (1-methoxymethyl-cyclopropylmethyl)-amide,

[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-pyrrolidin-1-yl-methanone,

15 (R)-1-[6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile,

azetidin-1-yl-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,

(R)-1-[6-(3-piperidin-1-yl-propoxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile,

20 (S)-1-[6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile,

and pharmaceutically acceptable salts thereof.

Furthermore, the pharmaceutically acceptable salts of the compounds of formula I and the pharmaceutically acceptable esters of the compounds of formula I individually
25 constitute preferred embodiments of the present invention.

Compounds of formula I may form acid addition salts with acids, such as conventional pharmaceutically acceptable acids, for example hydrochloride, hydrobromide, phosphate, acetate, fumarate, maleate, salicylate, sulphate, pyruvate, citrate, lactate, mandelate, tartrate, and methanesulphonate. Preferred are the


hydrochloride salts. Also solvates and hydrates of compounds of formula I and their salts form part of the present invention.

Compounds of formula I can have one or more asymmetric carbon atoms and can exist in the form of optically pure enantiomers, mixtures of enantiomers such as, for example, racemates, optically pure diastereoisomers, mixtures of diastereoisomers, diastereoisomeric racemates or mixtures of diastereoisomeric racemates. The optically active forms can be obtained for example by resolution of the racemates, by asymmetric synthesis or asymmetric chromatography (chromatography with a chiral adsorbent or eluant). The invention embraces all of these forms.

It will be appreciated, that the compounds of general formula I in this invention may be derivatised at functional groups to provide derivatives which are capable of conversion back to the parent compound in vivo. Physiologically acceptable and metabolically labile derivatives, which are capable of producing the parent compounds of general formula I in vivo are also within the scope of this invention.

A further aspect of the present invention is the process for the manufacture of compounds of formula I as defined above, which process comprises

reacting a compound of the formula II



II


wherein R is lower alkyl,

with an alcohol of the formula III




III

wherein A is as defined herein before,
in the presence of a trialkylphosphine or triphenylphosphine and of a diazo compound to obtain a compound of the formula IV



IV


and converting the ester of formula IV into the acid of formula V




V

under acidic or basic conditions,

- 5 and coupling the compound of formula V with an amine of the formula VI



wherein R^1 and R^2 are as defined herein before, with the help of an coupling agent under basic conditions to obtain a compound of the formula I



I


- 10 wherein A, R^1 and R^2 are as defined herein before,
and if desired,
converting the compound obtained into a pharmaceutically acceptable salt.

- Coupling agents for the reaction of compounds of formula V with amines of formula VI are for example N,N'-carbonyldiimidazole (CDI), N,N'-
15 dicyclohexylcarbodiimide (DCC), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI), 1-[bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium-3-oxide hexafluorophosphate (HATU), 1-hydroxy-1,2,3-benzotriazole (HOBT), or O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU).

In more detail, the compounds of formula I can be manufactured by the methods given below, by the methods given in the examples or by analogous methods.

Appropriate reaction conditions for the individual reaction steps are known to a person skilled in the art. Starting materials are either commercially available or can be prepared
 5 by methods analogous to the methods given below, by methods described in references cited in the text or in the examples, or by methods known in the art.

Scheme 1




Compounds of formula II can be manufactured starting from commercially
 10 available 6-methoxy-quinoline (1) that can be converted to the N-oxide 2 by reaction with hydrogen peroxide and a solvent like acetic acid under refluxing conditions. The 6-methoxy-quinoline-1-oxide is reacted with silver cyanide and benzoyl chloride to obtain the 6-methoxy-quinoline-2-carbonitrile 3 via a modification of the Reissert'sche reaction (Ber., 38, 1610 (1905)). Hydrolysis of the cyano group can be affected by employing an
 15 acidic or basic medium. We find it convenient to use a base like sodium hydroxide and after acidic treatment of the mixture we obtained the corresponding 6-methoxy-quinoline-2-carboxylic acid 4. Removal of the methyl group with an acid like hydrobromic acid 48% in water gives 6-hydroxy-quinoline-2-carboxylic acid (5). The acid can be esterified with an alcohol like ethanol and an acid like sulfuric acid to yield
 20 the 6-hydroxy-quinoline-2-carboxylic acid ester II, for example 6-hydroxy-quinoline-2-carboxylic acid ethyl ester (IIa).

The preparation of compounds of formula I of the present invention may be carried out in sequential or convergent synthetic routes. Syntheses of the invention are shown in the following scheme. The skills required for carrying out the reaction and

purification of the resulting products are known to those in the art. The substituents and indices used in the following description of the processes have the significance given above unless indicated to the contrary.

Scheme 2



5

Compounds of general formula I can be prepared according to scheme 2 as follows:

- a) The syntheses of ethers are widely described in literature and the procedures are known to those in the art (For reaction conditions described in literature affecting such reactions see for example: Comprehensive Organic Transformations: A Guide to Functional Group Preparations, 2nd Edition, Richard C. Larock. John Wiley & Sons, New York, 1999). The transformation can be affected by employing reaction conditions which are commonly utilised in the so called "*Mitsunobu reaction*" which is known to those in the art and widely described (Hughes, David L. The Mitsunobu reaction. Organic Reactions (John Wiley & Sons, New York, 1992, 42, 335-656) We find it convenient to couple an ester of formula II with alcohols of formula III (either commercially available or accessible by methods described in references or by methods known in the art; as appropriate) under conditions employing a phosphine like a trialkylphosphine such as tributylphosphine ((n-Bu)₃P), triphenylphosphine (Ph₃P) and the like and a diazo-compound like diethyl-azodicarboxylate (DEAD), diisopropyl-azodicarboxylate (DIAD) (optionally polymer bound), tetramethyl azodicarboxamide and the like in a solvent commonly used in such transformations like tetrahydrofuran (THF), toluene, dichloromethane and the like. There is no

20

particular restriction on the nature of the solvent to be employed, provided that it has no adverse effect on the reaction or the reagents involved and that it can dissolve the reagents, at least to some extent. The reaction can take place over a wide range of temperatures, and the precise reaction temperature is not critical to the invention.

5 We find it convenient to carry out the reaction with heating from ambient temperature to reflux. The time required for the reaction may also vary widely, depending on many factors, notably the reaction temperature and the nature of the reagents. However, a period of from 0.5 h to several days will usually suffice to yield the compounds of formula IV.

10 b) The hydrolysis of esters are widely described in literature and the procedures are known to those in the art (For reaction conditions described in literature affecting such reactions see for example:). The transformation can be affected by employing acidic or basic medium. We find it convenient to use acidic conditions employing an acid like HCl and a solvent like dioxane, THF and the like. There is no particular
15 restriction on the nature of the solvent to be employed, provided that it has no adverse effect on the reaction or the reagents involved and that it can dissolve the reagents, at least to some extent. The reaction can take place over a wide range of temperatures, and the precise reaction temperature is not critical to the invention. We find it convenient to carry out the reaction with heating from ambient
20 temperature to reflux. The time required for the reaction may also vary widely, depending on many factors, notably the reaction temperature and the nature of the reagents. However, a period of from 0.5 h to several days will usually suffice to yield the title compounds VI.

25 c) The coupling of carboxylic acids with amines is widely described in literature and the procedures are known to those in the art (For reaction conditions described in literature affecting such reactions see for example: Comprehensive Organic Transformations: A Guide to Functional Group Preparations, 2nd Edition, Richard C. Larock. John Wiley & Sons, New York, 1999). 6-Alkoxy-quinoline-2-carboxylic acids of formula IV can conveniently be transformed to the respective amide through
30 coupling with an amine V (either commercially available or accessible by methods described in references or by methods known in the art; as appropriate) by employing the usage of coupling agents. For example coupling agents like N,N'-carbonyldiimidazole (CDI), N,N'-dicyclohexylcarbodiimide (DCC), 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride (EDCI), 1-
35 [bis(dimethylamino)methylene]-1H-1,2,3-triazolo[4,5-b]pyridinium-3-oxide hexafluorophosphate (HATU), 1-hydroxy-1,2,3-benzotriazole (HOBT), O-benzotriazol-1-yl-N,N,N',N'-tetramethyluronium tetrafluoroborate (TBTU) and the

like can equally well be employed to affect such transformation. We find it convenient to carry out the reaction in a solvent like dimethylformamide (DMF) and in the presence of a base. There is no particular restriction on the nature of the solvent to be employed, provided that it has no adverse effect on the reaction or the reagents involved and that it can dissolve the reagents, at least to some extent. Examples for suitable solvents include: DMF, dichloromethane (DCM), dioxane, THF, and the like. There is no particular restriction on the nature of the base used in this stage, and any base commonly used in this type of reaction may equally be employed here. Examples of such bases include triethylamine and diisopropylethylamine, and the like. The reaction can take place over a wide range of temperatures, and the precise reaction temperature is not critical to the invention. We find it convenient to carry out the reaction with heating from ambient temperature to reflux. The time required for the reaction may also vary widely, depending on many factors, notably the reaction temperature and the nature of the reagents. However, a period of from 0.5 h to several days will usually suffice to yield amide derivatives of formula I.

As described above, the compounds of formula I of the present invention can be used as medicaments for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors. Examples of such diseases are obesity, metabolic syndrome (syndrome X), neurological diseases including Alzheimer's disease, dementia, age-related memory dysfunction, mild cognitive impairment, cognitive deficit, attention deficit hyperactivity disorder, epilepsy, neuropathic pain, inflammatory pain, migraine, Parkinson's disease, multiple sclerosis, stroke, dizziness, schizophrenia, depression, addiction, motion sickness and sleep disorders including narcolepsy, and other diseases including asthma, allergy, allergy-induced airway responses, congestion, chronic obstructive pulmonary disease and gastro-intestinal disorders. The use as medicament for the treatment and/or prevention of obesity is preferred.

The invention therefore also relates to pharmaceutical compositions comprising a compound as defined above and a pharmaceutically acceptable carrier and/or adjuvant.

Further, the invention relates to compounds as defined above for use as therapeutically active substances, particularly as therapeutic active substances for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors. Examples of such diseases are obesity, metabolic syndrome (syndrome X), neurological diseases including Alzheimer's disease, dementia, age-related memory dysfunction, mild cognitive impairment, cognitive deficit, attention deficit hyperactivity disorder, epilepsy, neuropathic pain, inflammatory pain, migraine, Parkinson's disease,

multiple sclerosis, stroke, dizziness, schizophrenia, depression, addiction, motion sickness and sleep disorders including narcolepsy, and other diseases including asthma, allergy, allergy-induced airway responses, congestion, chronic obstructive pulmonary disease and gastro-intestinal disorders.

5 In another embodiment, the invention relates to a method for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors. Examples of such diseases are obesity, metabolic syndrome (syndrome X), neurological diseases including Alzheimer's disease, dementia, age-related memory dysfunction, mild cognitive impairment, cognitive deficit, attention deficit hyperactivity disorder, epilepsy,
10 neuropathic pain, inflammatory pain, migraine, Parkinson's disease, multiple sclerosis, stroke, dizziness, schizophrenia, depression, addiction, motion sickness and sleep disorders including narcolepsy, and other diseases including asthma, allergy, allergy-induced airway responses, congestion, chronic obstructive pulmonary disease and gastro-intestinal disorders. A method for the treatment and/or prevention of obesity is
15 preferred.

The invention further relates to the use of compounds of formula I as defined above for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors. Examples of such diseases are obesity, metabolic syndrome (syndrome X), neurological diseases including Alzheimer's disease, dementia, age-related
20 memory dysfunction, mild cognitive impairment, cognitive deficit, attention deficit hyperactivity disorder, epilepsy, neuropathic pain, inflammatory pain, migraine, Parkinson's disease, multiple sclerosis, stroke, dizziness, schizophrenia, depression, addiction, motion sickness and sleep disorders including narcolepsy, and other diseases including asthma, allergy, allergy-induced airway responses, congestion, chronic
25 obstructive pulmonary disease and gastro-intestinal disorders. The use of compounds of formula I as defined above for the treatment and/or prevention of obesity is preferred.

In addition, the invention relates to the use of compounds of formula I as defined above for the preparation of medicaments for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors. Examples of such
30 diseases are obesity, metabolic syndrome (syndrome X), neurological diseases including Alzheimer's disease, dementia, age-related memory dysfunction, mild cognitive impairment, cognitive deficit, attention deficit hyperactivity disorder, epilepsy, neuropathic pain, inflammatory pain, migraine, Parkinson's disease, multiple sclerosis, stroke, dizziness, schizophrenia, depression, addiction, motion sickness and sleep
35 disorders including narcolepsy, and other diseases including asthma, allergy, allergy-induced airway responses, congestion, chronic obstructive pulmonary disease and

gastro-intestinal disorders. The use of compounds of formula I as defined above for the preparation of medicaments for the treatment and/or prevention of obesity is preferred.

The compounds of formula I and their pharmaceutically acceptable salts possess valuable pharmacological properties. Specifically, it has been found that the compounds
5 of the present invention are good histamine 3 receptor (H3R) antagonists and/or inverse agonists.

The following test was carried out in order to determine the activity of the compounds of formula (I).

Binding assay with ^3H -(R) α -methylhistamine

10 Saturation binding experiments were performed using HR3-CHO membranes prepared as described in Takahashi, K, Tokita, S., Kotani, H. (2003) J. Pharmacol. Exp. Therapeutics 307, 213-218.

An appropriate amount of membrane (60 to 80 μg protein/well) was incubated with increasing concentrations of ^3H -(R) α -Methylhistamine dihydrochloride (0.10 to 10
15 nM). Non specific binding was determined using a 200 fold excess of cold (R) α -Methylhistamine dihydrobromide (500 nM final concentration). The incubation was carried out at room temperature (in deep-well plates shaking for three hours). The final volume in each well was 250 μl . The incubation was followed by rapid filtration on GF/B filters (pre-soaked with 100 μl of 0.5% PEI in Tris 50 mM shaking at 200 rpm for two
20 hours). The filtration was made using a cell-harvester and the filter plates were then washed five times with ice cold washing buffer containing 0.5 M NaCl. After harvesting, the plates were dried at 55 $^{\circ}\text{C}$ for 60min, then we added scintillation fluid (Microscint 40, 40 microl in each well) and the amount of radioactivity on the filter was determined in Packard top-counter after shaking the plates for two hours at 200 rpm at room
25 temperature.

Binding Buffer: 50 mM Tris-HCl pH 7.4 and 5 mM $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ pH 7.4. Washing Buffer: 50 mM Tris-HCl pH 7.4 and 5 mM $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ and 0.5 M NaCl pH 7.4.

Indirect measurement of affinity of H3R inverse agonists: twelve increasing concentrations (ranging from 10 μM to 0.3 nM) of the selected compounds were always
30 tested in competition binding experiments using membrane of the human HR3-CHO cell line. An appropriate amount of protein, e.g. approximately 500cpm binding of RAMH at K_d , were incubated for 1 hour at room temperature in 250 μl final volume in 96-well plates in presence of ^3H -(R) α -Methylhistamine (1 nM final concentration = K_d).

Non-specific binding was determined using a 200 fold excess of cold (R) α -Methylhistamine dihydrobromide.

All compounds were tested at a single concentration in duplicates. Compounds that showed an inhibition of [^3H]-RAMH by more than 50% were tested again to determine
5 IC_{50} in a serial dilution experiment. K_i 's were calculated from IC_{50} based on Cheng-Prusoff equation (Cheng, Y, Prusoff, WH (1973) Biochem Pharmacol 22, 3099-3108).

The compounds of the present invention exhibit K_i values within the range of about 1 nM to about 1000 nM, preferably of about 1 nM to about 100 nM, and more preferably of about 1 nM to about 30 nM. The following table shows measured values for
10 some selected compounds of the present invention.

The following table shows measured values for some selected compounds of the present invention.

	K_i (nM)
Example 2	200
Example 14	78
Example 64	450

The compounds of formula (I) and their pharmaceutically acceptable salts and
15 esters can be used as medicaments, e.g. in the form of pharmaceutical preparations for enteral, parenteral or topical administration. They can be administered, for example, perorally, e.g. in the form of tablets, coated tablets, dragées, hard and soft gelatine capsules, solutions, emulsions or suspensions, rectally, e.g. in the form of suppositories,
parenterally, e.g. in the form of injection solutions or infusion solutions, or topically, e.g.
20 in the form of ointments, creams or oils.

The production of the pharmaceutical preparations can be effected in a manner which will be familiar to any person skilled in the art by bringing the described compounds of formula (I) and their pharmaceutically acceptable, into a galenical administration form together with suitable, non-toxic, inert, therapeutically compatible
25 solid or liquid carrier materials and, if desired, usual pharmaceutical adjuvants.

Suitable carrier materials are not only inorganic carrier materials, but also organic carrier materials. Thus, for example, lactose, corn starch or derivatives thereof, talc,

stearic acid or its salts can be used as carrier materials for tablets, coated tablets, dragées and hard gelatine capsules. Suitable carrier materials for soft gelatine capsules are, for example, vegetable oils, waxes, fats and semi-solid and liquid polyols (depending on the nature of the active ingredient no carriers are, however, required in the case of soft gelatine capsules). Suitable carrier materials for the production of solutions and syrups are, for example, water, polyols, sucrose, invert sugar and the like. Suitable carrier materials for injection solutions are, for example, water, alcohols, polyols, glycerol and vegetable oils. Suitable carrier materials for suppositories are, for example, natural or hardened oils, waxes, fats and semi-liquid or liquid polyols. Suitable carrier materials for topical preparations are glycerides, semi-synthetic and synthetic glycerides, hydrogenated oils, liquid waxes, liquid paraffins, liquid fatty alcohols, sterols, polyethylene glycols and cellulose derivatives.

Usual stabilizers, preservatives, wetting and emulsifying agents, consistency-improving agents, flavour-improving agents, salts for varying the osmotic pressure, buffer substances, solubilizers, colorants and masking agents and antioxidants come into consideration as pharmaceutical adjuvants.

The dosage of the compounds of formula (I) can vary within wide limits depending on the disease to be controlled, the age and the individual condition of the patient and the mode of administration, and will, of course, be fitted to the individual requirements in each particular case. For adult patients a daily dosage of about 1 mg to about 1000 mg, especially about 1 mg to about 100 mg, comes into consideration. Depending on the dosage it is convenient to administer the daily dosage in several dosage units.

The pharmaceutical preparations conveniently contain about 0.1-500 mg, preferably 0.5-100 mg, of a compound of formula (I).


The following examples serve to illustrate the present invention in more detail. They are, however, not intended to limit its scope in any manner.

Examples

Intermediate 1


6-Hydroxy-quinoline-2-carboxylic acid ethyl ester

a) 6-Methoxy-quinoline 1-oxide




6-Methoxy quinoline (15 g, 0.094 mol) was dissolved in acetic acid (97 ml) and treated with hydrogen peroxide (37 ml). The mixture was stirred to 100 °C for 2 hours. After vaporation to dryness 100 ml of water was added to the residue until a precipitate appears. Filtration and washing with water gives a yellow precipitate that is dried under vacuum to yield 13.5 g of the title compound as a light yellow solid (82 %). MS (m/e): 176.3 (M+ H) ⁺.

b) 6-Methoxy-quinoline-2-carbonitrile




6-Methoxy-quinoline-1-oxide (13.48 g, 0.076mol) in 30 ml chloroform was treated with 3.87 g (0.028 mol) of benzoyl chloride and 3.65 g (0.027 mol) of silver cyanide. The mixture was stirred for 4 hours at room temperature and stirred to reflux for additional 14 h. After evaporation of the solvent to half volume the mixture was cooled to 0 °C and a precipitate of silver salt appeared. The mixture was filtered and the solution was concentrated and the solid residue was washed in diethyl ether and dried under vacuum to yield 11 g (82 %) of the title compound as light brown solid. MS (m/e)= 185.3 (M+ H) ⁺.

c) 6-Methoxy-quinoline-2-carboxylic acid




6-Methoxy-quinoline-2-carbonitrile (9.3 g, 0.050 mol) in 96 ml methanol was treated with 240 ml of 20% NaOH and the mixture was heated to 120 °C in a sealed tube overnight. After cooling to 0 °C a precipitate appeared. The mixture was filtered to get a sodium salt that was suspended in water. HCl 25% was added until pH 3-4 to get the acid as a precipitate that was filtered and dried under vacuum. The mother liquid of the first filtration was acidified with HCl 25% until pH 3-4 until a precipitate appeared. The precipitate was filtered and the solid was washed in water and dried under vacuum to yield 9.6 g (86 %) of the title compound as light brown solid. MS (m/e)= 204 (M+ H) ⁺.

d) 6-Hydroxy-quinoline-2-carboxylic acid

6-Methoxy-quinoline-2-carboxylic acid (4 g, 0.019 mol) was suspended in hydrobromic acid 48% in water (80 ml) and the mixture was heated at 125 °C overnight.

- 5 After cooling to 0 °C ammonium hydroxide was added until the pH was 6-7 followed by addition of HCl until the pH was 3-4 and the compound precipitated. The solid was filtrated, washed with water and dried under vacuum to yield 3.5 g (0.0185 mol, 97% of theory) of the title compound as a yellow solid MS (m/e)= 190.1 (M+H)⁺.


e) 6-Hydroxy-quinoline-2-carboxylic acid ethyl ester

10

6-Hydroxy-quinoline-2-carboxylic acid (2.3 g, 0.012 mol) was dissolved in absolute ethanol (150 ml). Sulfuric acid (0.550ml, 0.0096 mol) was added and the mixture was refluxed for 16 hours. After cooling to room temperature the ethanol was evaporated and 60 ml of ethyl acetate, 50 ml of cooled water were added. The pH was
15 adjusted to 7 with solid NaHCO₃. Extraction and concentration of the organic phase gave the title product as a light yellow solid that was used for the next steps without purification (2.1 g, 81% of theory). MS (m/e)= 218.4 (M+ H)⁺.


Intermediate 2(6-(1-Isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride

a) 6-(1-Isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid ethyl ester




A mixture of 1 g (0.0046 mol) of 6-hydroxy-quinoline-2-carboxylic acid ethyl ester, 2.4 g (0.0092 mol) of triphenylphosphine (Fluka), 2.4 g (0.0092 mol) of 1-isopropyl-piperidin-4-ol and 1.6 ml (0.0092 mol) of di-tert.-butyl azadicarboxylate 40% in toluene in 100 ml THF was stirred for a prolonged period of time at 35 °C. The mixture was filtered through a pad of silica and washed with 30 ml THF. The mixture was evaporated to dryness and purified on silica eluting with a gradient of DCM/ MeOH 98/2. The product fractions were evaporated and the residue was triturated with diethyl ether to yield after drying under vacuum 1.3 mg (83 %) of the title compound as white solid. MS (m/e): 343.3 (M+ H)⁺.

b) 6-(1-Isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride




6-(1-Isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid ethyl ester (1.1 g, 0.0033 mol) was dissolved in dioxane (25 ml). HCl 37% (2.06 ml, 0.066 mol) was added and the mixture was stirred at 85 °C for 16 hours. Dioxane was evaporated and toluene (3 x 15 ml) was used to remove the remaining water by azeotropic distillation. After evaporation of the toluene the title compound was obtained as a yellow solid after drying under vacuum (1.3 g, 100% of theory). MS (m/e)= 315.2 (M+ H)⁺.


Intermediate 36-(1-Isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloridea) 6-(1-Isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid ethyl ester

- 5 A mixture of 0.960 g (0.0044 mol) of 6-hydroxy-quinoline-2-carboxylic acid ethyl ester, 2.32 g (0.0088 mmol) of triphenylphosphine (Fluka), 0.739g (0.0057 mmol) of 1-isopropyl-3-pyrrolidinol and 1.61 ml (0.0088 mmol) of di-tert.-butyl azadicarboxylate 40% in toluene in 100 ml THF was stirred for a prolonged period of time at 35 °C. The mixture was filtered through a pad of silica and washed with 30 ml of THF. The mixture
10 was evaporated to dryness and purified on silica eluting with a gradient of DCM/ MeOH 98/2. The product fractions were evaporated and the residue was triturated with diethyl ether to yield after drying under vacuum 1.35 g (94 %) of the title compound as white solid. MS (m/e): 329.3 (M+ H)⁺.


15 b) 6-(1-Isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride



- 6-(1-Isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid ethyl ester (0.5 g, 0.002 mol) was dissolved in dioxane (20 ml). HCl 37% (0.94 ml, 0.03 mol) was added and the mixture was stirred at 85 °C for 16 hours. Dioxane was evaporated and toluene (3 x 15 ml) was used to remove the remaining water by azeotropic distillation. After
20 evaporation of the toluene the title compound was obtained as a yellow solid after drying under vacuum (0.570 g, 100 % of theory). MS (m/e)= 301.2 (M+ H)⁺.


Intermediate 46-(3-Piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloridea) 6-(3-Piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid ethyl ester

- 5 A mixture of 0.400 g (0.002 mol) of 6-hydroxy-quinoline-2-carboxylic acid ethyl ester, 0.966 g (0.004 mmol) of triphenylphosphine (Fluka), 0.396 g (0.003 mmol) of 3-piperidin-1-yl-propan-1-ol and 0.68 ml (0.004 mmol) of di-tert.-butyl azadicarboxylate 40% in toluene in 40 ml THF was stirred for a prolonged period of time at 35 °C. The mixture was filtered through a pad of silica and washed with 30 ml THF. The mixture
- 10 was evaporated to dryness and purified on silica eluting with a gradient of DCM/ MeOH/NH₄OH from 97/3/0.5 to 90/10/0.5. The product fractions were evaporated and the residue was triturated with diethyl ether to yield after drying under vacuum 0.600 g (93 %) of the title compound as white solid. MS (m/e): 343.3 (M+ H)⁺.


b) 6-(3-Piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride

15

- 6-(3-Piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid ethyl ester (0.6 g, 0.002 mol) was dissolved in dioxane (20 ml). HCl 37% (1.08 ml, 0.035 mol) was added and the mixture was stirred at 85 °C for 16 hours. Dioxane was evaporated and toluene (3 x 10 ml) was used to remove the remaining water by azeotropic distillation. After evaporation
- 20 of the toluene the title compound was obtained as a yellow solid after drying under vacuum (0.592 g, 96% of theory). MS (m/e)= 315.3(M+ H)⁺.

Intermediate 56-[2-(1-Methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid 1:1 hydrochloridea) 6-[2-(1-Methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid ethyl ester

- 5 A mixture of 0.400 g (0.002 mol) of 6-hydroxy-quinoline-2-carboxylic acid ethyl ester, 0.966 g (0.004 mmol) of triphenylphosphine (Fluka), 0.396 g (0.003 mmol) of 1-methyl-2-pyrrolidineethanol and 0.68 ml (0.004 mmol) di-tert.-butyl azodicarboxylate 40% in toluene in 40 ml THF was stirred for a prolonged period of time at 35 °C. The mixture was filtered through a pad of silica and washed with 30 ml THF. The mixture was
- 10 evaporated to dryness and purified on silica eluting with a gradient of DCM/MeOH/NH₄OH from 97/3/0.5 to 90/10/0.5. The product fractions were evaporated and the residue was triturated with diethyl ether to yield after drying under vacuum 0.317 g (53 %) of the title compound as white solid. MS (m/e): 329.2 (M+ H)⁺.

b) 6-[2-(1-Methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid 1:115 hydrochloride

- 6-[2-(1-Methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid ethyl ester (0.310 g, 0.001 mol) was dissolved in dioxane (10 ml). HCl 37% (0.580 ml, 0.019 mol) was added and the mixture was stirred at 85 °C for 16 hours. Dioxane was evaporated and
- 20 toluene (3 x 10 ml) was used to remove the remaining water by azeotropic distillation. After evaporation of the toluene the title compound was obtained as a yellow solid after drying under vacuum (0.395g, 96% of theory). MS (m/e)= 301.3 (M+ H)⁺.

Example 1(2,5-Dihydro-pyrrol-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone

6-(1-Isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride
 5 (30 mg, 0.086 mmol, see intermediate 2) was dissolved in DMF (0.300 ml). 1,1'-
 Carbonyl-diimidazole (17 mg, 0.10 mmol) was added and the mixture was stirred for
 half an hour. 4-Methoxy-piperidine was added (6 mg, 0.014 mmol) and the mixture was
 stirred overnight. The mixture was diluted with 0.4 ml methanol and subjected to
 preparative HPLC purification on reversed phase material eluting with a gradient of
 10 acetonitrile/water/ triethylamine. The product fractions were evaporated to dryness to
 yield 4.7 mg (14 %) of the title compound as light brown solid. MS (m/e): 412.4 (MH⁺,
 100%).

According to the procedure described for the synthesis of example 1 further
 derivatives have been synthesised from 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-
 15 carboxylic acid 1:1 hydrochloride and the respective amines . The results are shown in
 table 1 and comprise examples 2 to 26.

Table 1

No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
2	(2,5-dihydro-pyrrol-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone	365.4	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 5-dihydropyrrole (commercially available)	366.4
3	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid ethyl-methyl-amide	355.48	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and ethyl-methyl-amide (commercially available)	356.4
4	[6-(1-isopropyl-	381.52	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1	382.3

No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
	piperidin-4-yloxy)-quinolin-2-yl]-(2-methyl-pyrrolidin-1-yl)-methanone		hydrochloride and 2-methyl-pyrrolidine (commercially available)	
5	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid tert-butylamide	369.51	6-(1-Isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and tert-butylamide (commercially available)	370.3
6	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid cyclopropylmethyl-propyl-amide	409.57	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and cyclopropylmethyl-propyl-amide (commercially available)	410.5
7	6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(2-methyl-piperidin-1-yl)-methanone	395.55	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 2-methyl-piperidine (commercially available)	396.3
8	(4-hydroxy-piperidin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone	397.52	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 4-hydroxy-piperidine (commercially available)	398.4
9	[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(3-methyl-piperidin-1-yl)-methanone	395.55	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 3-methyl-piperidine (commercially available)	396.3

No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
10	(3,4-dihydro-1H-isoquinolin-2-yl)-[6-(3-piperidin-1-yl-propoxy)-naphthalen-2-yl]-methanone 1:1 hydrochloride	369.51	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and isopropyl-methyl-amine (commercially available)	370.4
11	[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-thiomorpholin-4-yl-methanone	399.56	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and thiomorpholine (commercially available)	400.4
12	[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(2-trifluoromethyl-pyrrolidin-1-yl)-methanone	435.49	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 2-trifluoromethyl-pyrrolidine (commercially available)	436.4
13	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid ethyl-(2-methoxy-ethyl)-amide	399.53	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and ethyl-(2-methoxy-ethyl)-amine (commercially available)	400.5
14	azetidin-1-yl-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone	353.47	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and azetidine (commercially available)	354.4
15	(3,6-dihydro-2H-pyridin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-	379.5	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and	380.5

No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
	quinolin-2-yl]-methanone		1,2,3,6-tetrahydro-pyridine (commercially available)	
16	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid (3-fluoro-oxetan-3-ylmethyl)-amide	411.55	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and (3-fluoro-oxetan-3-yl)-methylamine (commercially available)	412.4
17	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid (1-methoxymethyl-cyclopropylmethyl)-amide	411.55	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and (1-methoxymethyl-cyclopropyl)-methylamine (commercially available)	412.4
18	{6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid [2-(tetrahydro-pyran-4-yl)-ethyl]-amide	425.57	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 2-(tetrahydro-pyran-4-yl)-ethylamine (commercially available)	426.4
19	[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-morpholin-4-yl-methanone	383.4	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and morpholine (commercially available)	384.3
20	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid cyclohexyl-methyl-amide	409.57	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and cyclohexyl-methyl-amide	410.6

No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
			(commercially available)	
21	(4,4-difluoro-piperidin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone	417.497	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 4,4-difluoro-piperidine (commercially available)	418.3
22	(3-hydroxy-pyrrolidin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone	383.4	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 3-hydroxy-pyrrolidine (commercially available)	384.3
23	[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-pyrrolidin-1-yl-methanone	367.49	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and pyrrolidine (commercially available)	368.2
24	(R)-1-[6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile	392.5	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and ((R)- pyrrolidine-2-carbonitrile (commercially available)	393.3
25	(1,1-dioxo--thiomorpholin-4-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone	431.5	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and thiomorpholine 1,1-dioxide	432.4
26	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid	341.5	6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and	342.2

No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
	dimethylamide		dimethylamide (commercially available)	

Example 27(2,5-Dihydro-pyrrol-1-yl)-[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-methanone

- 5 According to the procedure described for the synthesis of example 1, (2,5-dihydro-pyrrol-1-yl)-[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-methanone was synthesized from 6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride (intermediate 3) and 2,5-dihydropyrrole (commercially available). The title compound was yielded in 18 % (5.7mg) as off-white solid. MS (m/e): 352.5 (MH⁺,
 10 100%).

Accordingly, further derivatives have been synthesised from 6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and the respective amines. The results are shown in table 2 and comprise examples 28 to 46.

Table 2

Ex. No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
28	([6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-(4-methoxy-piperidin-1-yl)-methanone	397.52	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 4-methoxy-piperidine	398.4
29	6-(1-isopropyl-pyrrolidin-3-yloxy)-	341.45	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and	342.2

Ex. No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
	quinoline-2-carboxylic acid ethyl-methyl-amide		ethyl-methyl-amide (commercially available)	
30	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl)-(4-methyl-piperidin-1-yl)-methanone	381.52	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and 4-methyl-piperidine (commercially available)	382.3
31	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl)-(2-methyl-pyrrolidin-1-yl)-methanone	367.49	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and 2-methyl-pyrrolidine (commercially available)	368.3
32	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid tert-butylamide	355.48	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and tert-butylamide (commercially available)	356.4
33	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid cyclopropylmethyl-propyl-amide	395.55	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and cyclopropylmethyl-propyl-amide (commercially available)	396.3
34	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl)-(2-methyl-piperidin-1-yl)-methanone	381.52	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and 2-methyl-piperidine (commercially available)	382.3
35	6-(1-isopropyl-	381.52	6-(1-isopropyl-pyrrolidin-3-yloxy)-	382.3

Ex. No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
	pyrrolidin-3-yloxy)-quinolin-2-yl]-(3-methyl-piperidin-1-yl)-methanone		quinoline-2-carboxylic acid acid 1:1 hydrochloride and 3-methyl-piperidine (commercially available)	
36	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid isopropyl-methyl-amide	355.48	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and isopropyl-methyl-amine (commercially available)	356.4
37	[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-thiomorpholin-4-yl-methanone	385.53	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and thiomorpholine (commercially available)	386.3
38	[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-(2-trifluoromethyl-pyrrolidin-1-yl)-methanone	421.46	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and 2-trifluoromethyl-pyrrolidine (commercially available)	422.3
39	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid ethyl-(2-methoxy-ethyl)-amide	385.51	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and ethyl-(2-methoxy-ethyl)-amine (commercially available)	386.4
40	azetidin-1-yl-[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-methanone	339.44	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and azetidine (commercially available)	340.3

Ex. No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
41	(3,6-dihydro-2H!-pyridin-1-yl)-[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-methanone	365.48	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and 1,2,3,6-tetrahydro-pyridine (commercially available)	366.2
42	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid (3-fluoro-oxetan-3-ylmethyl)-amide	387.45	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and (3-fluoro-oxetan-3-yl)-methylamine	388.3
43	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid (1-methoxymethyl-cyclopropylmethyl)-amide	397.52	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and (1-methoxymethyl-cyclopropyl)-methylamine	398.4
44	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid [2-(tetrahydro-pyran-4-yl)-ethyl]-amide	411.55	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and 2-(tetrahydro-pyran-4-yl)-ethylamine	412.4
45	(4,4-difluoro-piperidin-1-yl)-[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-methanone	403.5	6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid acid 1:1 hydrochloride and 4,4-difluoro-piperidine (commercially available)	404.4

Ex. No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
46	[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-morpholin-4-yl-methanone	369.4	6-(1-Isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and morpholine (commercially available)	370.3

Example 47(4-Methoxy-piperidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone

According to the procedure described for the synthesis of example 1, (4-methoxy-piperidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone was synthesized from 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride (intermediate 4) and 4-methoxy-piperidine (commercially available). The title compound was yielded in 22% (7.8mg) as off-white solid. MS (m/e): 412.4 (MH⁺, 100%).

Accordingly, further derivatives have been synthesised from 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and the respective amines. The results are shown in table 3 and comprise examples 48 to 66.

Table 3

Ex. No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
48	(4-methyl-piperidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone	395.55	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 4-methyl-piperidine (commercially available)	396.3

Ex. No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
49	morpholin-4-yl-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone	383.49	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and morpholine (commercially available)	384.3
50	(2-methyl-pyrrolidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone	381.52	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 2-methyl-pyrrolidine (commercially available)	382.4
51	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid cyclopropylmethyl-propyl-amide	409.57	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and cyclopropylmethyl-propyl-amide (commercially available)	410.4
52	[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-(2-trifluoromethyl-pyrrolidin-1-yl)-methanone	435.49	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 2-trifluoromethyl-pyrrolidine (commercially available)	436.3
53	(2,5-dihydro-pyrrol-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone	365.48	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 2,5-dihydropyrrole (commercially available)	366.3

Ex. No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
54	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid ethyl-(2-methoxy-ethyl)-amide	399.53	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and ethyl-(2-methoxy-ethyl)-amine (commercially available)	400.3
55	azetidin-1-yl-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone	353.47	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and azetidine (commercially available)	354.3
56	(3,6-dihydro-2H-pyridin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone	379.5	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 1,2,3,6-tetrahydro-pyridine (commercially available)	380.4
57	(4,4-difluoro-piperidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone	417.5	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 4,4-difluoro-piperidine (commercially available)	418.4
58	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid cyclohexyl-methyl-amide	409.57	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and cyclohexyl-methyl-amide (commercially available)	410.4

Ex. No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
59	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid (1-methoxymethyl-cyclopropylmethyl)-amide	411.55	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and (1-methoxymethyl-cyclopropyl)-methylamine (commercially available)	412.4
60	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid (3-fluoro-oxetan-3-ylmethyl)-amide	401.48	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and (3-fluoro-oxetan-3-yl)-methylamine (commercially available)	402.4
61	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid [2-(tetrahydro-pyran-4-yl)-ethyl]-amide	425.57	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 2-(tetrahydro-pyran-4-yl)-ethylamine (commercially available)	426.3
62	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid (tetrahydro-pyran-4-yl)-amide	397.52	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 4-amino-tetrahydropyran (commercially available)	398.3
63	(2-methyl-piperidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone	395.55	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and 2-methyl-piperidine (commercially available)	396.3

Ex. No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
64	[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-pyrrolidin-1-yl-methanone	367.49	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and pyrrolidine (commercially available)	368.2
65	(R)-1-[6-(3-piperidin-1-yl-propoxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile	392.5	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and ((R)-pyrrolidine-2-carbonitrile (commercially available)	393.3
66	(1,1-dioxo--thiomorpholin-4-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone	431.55	6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid 1:1 hydrochloride and thiomorpholine 1,1-dioxide	432.4

Example 67(4-Methyl-piperidin-1-yl)-[6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl]-methanone

5 According to the procedure described for the synthesis of example 1, (4-methyl-piperidin-1-yl)-[6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl]-methanone was synthesized from 6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid 1:1 hydrochloride (intermediate 5) and 4-methyl-piperidine (commercially available). The title compound was yielded in 13% (4.2mg) as white solid. MS (m/e):

10 382.4 (MH⁺, 100%).

According to the procedure described for the synthesis of example 67 further derivatives have been synthesised from 6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-

quinoline-2-carboxylic acid 1:1 hydrochloride and the respective amines. The results are shown in table 4 and comprise examples 68 to 76.

Table 4

Ex. No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
68	{6-[2-(1-Methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-pyrrolidin-1-yl-methanone	353.47	6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid 1:1 hydrochloride and pyrrolidine (commercially available)	354.3
69	(R)-1-[6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile	378.48	6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid 1:1 hydrochloride and ((R)-pyrrolidine-2-carbonitrile (commercially available)	379.4
70	(1,1-dioxo-thiomorpholin-4-yl)-{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-methanone	417.53	6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid 1:1 hydrochloride and thiomorpholine 1,1-dioxide	418.4
71	(4-methoxy-piperidin-1-yl)-{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-methanone	397.52	6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid 1:1 hydrochloride and 4-methoxy-piperidine (commercially available)	398.3
72	{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-morpholin-4-yl-methanone	369.46	6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid 1:1 hydrochloride and morpholine (commercially available)	370.3

Ex. No	Systematic name	MW	Starting materials	MW found (M+H) ⁺
73	azetidin-1-yl-{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-methanone	339.44	6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid 1:1 hydrochloride and azetidine (commercially available)	340.3
74	6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid [2-(tetrahydro-pyran-4-yl)-ethyl]-amide	411.55	6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid 1:1 hydrochloride and 2-(tetrahydro-pyran-4-yl)-ethylamine (commercially available)	412.4
75	6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid (3-fluoro-oxetan-3-ylmethyl)-amide	387.45	6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid 1:1 hydrochloride and (3-fluoro-oxetan-3-yl)-methylamine (commercially available)	388.3
76	(2-methyl-pyrrolidin-1-yl)-{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-methanone	367.49	6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid 1:1 hydrochloride and 2-methyl-pyrrolidine (commercially available)	368.2

Example 77(S)-1-[6-(1-Isopropyl-piperidin-4-yloxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile

- 5 According to the procedure described for the synthesis of example 1, (S)-1-[6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile was synthesized from 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride, (S)-cyanopyrrolidine (commercially available) and O-(benzotriazol-1-

yl)-N,N,N',N'-tetramethyluronium tetrafluoroborate instead of 1,1'-carbonyl-diimidazole. The title compound was yielded in 45% (50 mg) as colorless foam. MS (m/e): 392.9 (M).

Example 78

5 [6-(1-Isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(4-methyl-piperidin-1-yl)-methanone

According to the procedure described for the synthesis of example 1, [6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(4-methyl-piperidin-1-yl)-methanone was synthesized from 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1
10 hydrochloride, 4-methylpiperidine (commercially available) and 1,1'-carbonyl-diimidazole. MS (m/e): 396.6 (M+H).

Example 79

(4-Hydroxymethyl-piperidin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone

15 According to the procedure described for the synthesis of example 1, (4-hydroxymethyl-piperidin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone was synthesized from 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride, 4-hydroxymethyl piperidine (commercially available) and 1,1'-carbonyl-diimidazole. MS (m/e): 412.5 (M+H).

20 Example 80

6-(1-Isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid isobutyl-amide

According to the procedure described for the synthesis of example 1, (6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid isobutyl-amide was synthesized from 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid 1:1
25 hydrochloride, isobutylamine (commercially available) and 1,1'-carbonyl-diimidazole. MS (m/e): 370.6 (M+H).

Example 816-(1-Isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid cyclohexyl-methyl-amide

5 According to the procedure described for the synthesis of example 27, 6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid cyclohexyl-methyl-amide was synthesized from 6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid 1:1 hydrochloride (intermediate 3), N-methylcyclohexylamine (commercially available) and 1,1'-carbonyl-diimidazole. MS (m/e): 396.4 (M+H).

Example A

Film coated tablets containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	<u>Per tablet</u>	
Kernel:		
Compound of formula (I)	10.0 mg	200.0 mg
Microcrystalline cellulose	23.5 mg	43.5 mg
Lactose hydrous	60.0 mg	70.0 mg
Povidone K30	12.5 mg	15.0 mg
Sodium starch glycolate	12.5 mg	17.0 mg
Magnesium stearate	1.5 mg	4.5 mg
(Kernel Weight)	120.0 mg	350.0 mg
Film Coat:		
Hydroxypropyl methyl cellulose	3.5 mg	7.0 mg
Polyethylene glycol 6000	0.8 mg	1.6 mg
Talc	1.3 mg	2.6 mg
Iron oxyde (yellow)	0.8 mg	1.6 mg
Titanium dioxide	0.8 mg	1.6 mg

- 5 The active ingredient is sieved and mixed with microcrystalline cellulose and the mixture is granulated with a solution of polyvinylpyrrolidone in water. The granulate is mixed with sodium starch glycolate and magnesiumstearate and compressed to yield kernels of 120 or 350 mg respectively. The kernels are lacquered with an aqueous solution / suspension of the above mentioned film coat.

Example B

Capsules containing the following ingredients can be manufactured in a conventional manner:

<u>Ingredients</u>	<u>Per capsule</u>
Compound of formula (I)	25.0 mg
Lactose	150.0 mg
Maize starch	20.0 mg
Talc	5.0 mg

5 The components are sieved and mixed and filled into capsules of size 2.

Example C

Injection solutions can have the following composition:

Compound of formula (I)	3.0 mg
Gelatine	150.0 mg
Phenol	4.7 mg
Sodium carbonate	to obtain a final pH of 7
Water for injection solutions	ad 1.0 ml

Example D

Soft gelatin capsules containing the following ingredients can be manufactured in a conventional manner:

Capsule contents

Compound of formula (I)	5.0 mg
Yellow wax	8.0 mg
Hydrogenated Soya bean oil	8.0 mg
Partially hydrogenated plant oils	34.0 mg
Soya bean oil	110.0 mg
Weight of capsule contents	165.0 mg

Gelatin capsule

Gelatin	75.0 mg
Glycerol 85 %	32.0 mg
Karion 83	8.0 mg (dry matter)
Titanium dioxide	0.4 mg
Iron oxide yellow	1.1 mg

- 5 The active ingredient is dissolved in a warm melting of the other ingredients and the mixture is filled into soft gelatin capsules of appropriate size. The filled soft gelatin capsules are treated according to the usual procedures.

Example E


Sachets containing the following ingredients can be manufactured in a conventional manner:

Compound of formula (I)	50.0 mg
Lactose, fine powder	1015.0 mg
Microcrystalline cellulose (AVICEL PH 102)	1400.0 mg
Sodium carboxymethyl cellulose	14.0 mg
Polyvinylpyrrolidone K 30	10.0 mg
Magnesiumstearate	10.0 mg
Flavoring additives	1.0 mg

- 5 The active ingredient is mixed with lactose, microcrystalline cellulose and sodium carboxymethyl cellulose and granulated with a mixture of polyvinylpyrrolidone in water. The granulate is mixed with magnesiumstearate and the flavouring additives and filled into sachets.


Claims

1. Compounds of the general formula




wherein


- 5 R^1 is selected from the group consisting of hydrogen,
lower alkyl, lower alkenyl,
cycloalkyl or lower cycloalkylalkyl, wherein the cycloalkyl ring may be
unsubstituted or substituted by one or two groups selected from the group
consisting of lower alkyl, lower hydroxyalkyl and lower alkoxyalkyl,
10 lower hydroxyalkyl,
lower alkoxyalkyl, and
lower heterocyclalkyl, wherein the heterocyclalkyl ring may be unsubstituted or
substituted by one or two groups selected from lower alkyl and halogen;
- 15 R^2 is selected from the group consisting of hydrogen,
lower alkyl, lower alkenyl,
cycloalkyl or lower cycloalkylalkyl, wherein the cycloalkyl ring may be
unsubstituted or substituted by one or two groups selected from the group
consisting of lower alkyl, lower hydroxyalkyl and lower alkoxyalkyl,
lower hydroxyalkyl,
20 lower alkoxyalkyl, and
lower heterocyclalkyl, wherein the heterocyclalkyl ring may be unsubstituted or
substituted by one or two groups selected from lower alkyl and halogen; or
- 25 R^1 and R^2 together with the nitrogen atom to which they are attached form a 4-, 5-, 6- or
7-membered saturated or partly unsaturated heterocyclic ring optionally
containing a further heteroatom selected from nitrogen, oxygen or sulfur, said
saturated or partly unsaturated heterocyclic ring being unsubstituted or
substituted by one, two or three groups independently selected from the group
consisting of lower alkyl, halogen, halogenalkyl, cyano, hydroxy, hydroxyalkyl,
lower alkoxy, oxo, phenyl, benzyl, pyridyl and carbamoyl;
- 30 A is selected from



A 1



A 2



A 3

wherein

m is 0, 1 or 2;

5 R³ is lower alkyl;

n is 0, 1 or 2;

R⁷ is lower alkyl;

p is 0, 1 or 2;

q is 0, 1 or 2;

10 R⁵ is hydrogen or lower alkyl;

and pharmaceutically acceptable salts thereof.

2. Compounds of formula I according to claim 1, wherein R¹ is selected from the group consisting of hydrogen, lower alkyl, lower alkenyl, cycloalkyl or lower cycloalkylalkyl, wherein the cycloalkyl ring may be unsubstituted or substituted by one or two groups selected from lower alkyl, lower hydroxyalkyl or lower alkoxyalkyl, lower hydroxyalkyl, lower alkoxyalkyl, and lower heterocyclalkyl, wherein the heterocyclalkyl ring may be unsubstituted or substituted with one or two groups selected from lower alkyl or halogen, and

15 R² is hydrogen or lower alkyl.

3. Compounds of formula I according to claims 1 or 2, wherein R¹ is selected from the group consisting of lower alkyl, cycloalkyl or lower cycloalkylalkyl, wherein the cycloalkyl ring may be unsubstituted or substituted by lower alkoxyalkyl, lower alkoxyalkyl, and lower heterocyclalkyl wherein the heterocyclalkyl ring may be unsubstituted or substituted

25

with one or two groups selected from lower alkyl or halogen,
and R² is hydrogen or lower alkyl.

4. Compounds of formula I according to claim 3, wherein R¹ and R² are lower alkyl.


5 5. Compounds of formula I according to any one of claims 1 to 4, wherein R¹ and R² together with the nitrogen atom to which they are attached form a 4-, 5-, 6- or 7-membered saturated or partly unsaturated heterocyclic ring optionally containing a further heteroatom selected from nitrogen, oxygen or sulfur, said heterocyclic ring being unsubstituted or substituted by one, two or three groups independently selected from
10 lower alkyl, halogen, halogenalkyl, cyano, hydroxy, lower alkoxy, oxo, phenyl, benzyl, pyridyl and carbamoyl.

6. Compounds of formula I according to any one of claims 1 to 5, wherein R¹ and R² together with the nitrogen atom to which they are attached form a 4-, 5-, or 6- or 7-membered saturated or partly unsaturated heterocyclic ring optionally containing a
15 further heteroatom selected from nitrogen, oxygen or sulfur, said heterocyclic ring being unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, halogen, halogenalkyl, cyano, hydroxy, lower alkoxy, and oxo.

7. Compounds of formula I according to any one of claims 1 to 6, wherein R¹ and R² together with the nitrogen atom to which they are attached form a heterocyclic ring
20 selected from the group consisting of morpholine, piperidine, 2,5-dihydropyrrole, pyrrolidine, azepane, piperazine, azetidine, thiomorpholine and 3,6-dihydro-2H-pyridine, said heterocyclic ring being unsubstituted or substituted by one, two or three groups independently selected from lower alkyl, halogen, halogenalkyl, cyano, hydroxy, lower alkoxy, and oxo.

25 8. Compounds of formula I according to any one of claims 1 to 7, wherein R¹ and R² together with the nitrogen atom to which they are attached form a heterocyclic ring selected from morpholinyl, thiomorpholinyl, 1,1-dioxothiomorpholinyl, 3,6-dihydro-2H-pyridinyl, piperidinyl, 2-methylpiperidinyl, 3-methylpiperidinyl, 4-hydroxypiperidinyl, 4,4-difluoropiperidinyl, 2,5-dihydropyrrolyl, 4-methylpiperidinyl,
30 pyrrolidinyl, 2-methylpyrrolidinyl, 2-trifluoromethylpyrrolidinyl, 2-cyanopyrrolidinyl, 3-hydroxypyrrolidinyl and azetidiny.

9. Compounds of formula I according to any one of claims 1 to 8, wherein A signifies




A 1

wherein m is 0, 1 or 2, and R³ is lower alkyl.

10. Compounds of formula I according to claim 9, wherein m is 0.

11. Compounds of formula I according to claim 9, wherein m is 1.

12. Compounds of formula I according to any one of claims 1 to 8, wherein A signifies




A 2

wherein n is 0, 1 or 2; and R⁴ is lower alkyl.

13. Compounds of formula I according to claim 12, wherein n is 0.

14. Compounds of formula I according to any one of claims 1 to 8, wherein A signifies



A 3

wherein p is 0, 1 or 2, q is 0, 1 or 2, and R⁵ is hydrogen or lower alkyl.

15. Compounds of formula I according to claim 14, wherein p is 1.

16. Compounds of formula I according to claim 14 or 15, wherein R⁵ is hydrogen.

17. Compounds of formula I according to claim 1, selected from the group consisting of

- [6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(4-methoxy-piperidin-1-yl)-
- 5 methanone,
- (2,5-dihydro-pyrrol-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-
- methanone,
- 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid ethyl-methyl-amide,
- [6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(2-methyl-pyrrolidin-1-yl)-
- 10 methanone,
- 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid tert-butylamide,
- 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid cyclopropylmethyl-
- propyl-amide,
- 6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(2-methyl-piperidin-1-yl)-methanone,
- 15 (4-hydroxy-piperidin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-
- methanone,
- [6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(3-methyl-piperidin-1-yl)-
- methanone,
- (3,4-dihydro-1H-isoquinolin-2-yl)-[6-(3-piperidin-1-yl-propoxy)-naphthalen-2-yl]-
- 20 methanone 1:1 hydrochloride,
- [6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-thiomorpholin-4-yl-methanone,
- [6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-(2-trifluoromethyl-pyrrolidin-1-yl)-
- methanone,
- 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid ethyl-(2-methoxy-ethyl)-
- 25 amide,
- azetidin-1-yl-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone,
- (3,6-dihydro-2H-pyridin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-
- methanone,
- 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid (3-fluoro-oxetan-3-
- 30 ylmethyl)-amide,
- 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid (1-methoxymethyl-
- cyclopropylmethyl)-amide,
- {6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid [2-(tetrahydro-pyran-4-
- yl)-ethyl]-amide,
- 35 [6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-morpholin-4-yl-methanone,
- 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid cyclohexyl-methyl-
- amide,
- (4,4-difluoro-piperidin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-

- methanone,
(3-hydroxy-pyrrolidin-1-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-
methanone,
[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-pyrrolidin-1-yl-methanone,
5 (R)-1-[6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carbonyl]-pyrrolidine-2-
carbonitrile,
(1,1-dioxo--thiomorpholin-4-yl)-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-
methanone,
6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid dimethylamide,
10 (2,5-dihydro-pyrrol-1-yl)-[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-
methanone,
{[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-(4-methoxy-piperidin-1-yl)-
methanone,
6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid ethyl-methyl-amide,
15 {[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-(4-methyl-piperidin-1-yl)-
methanone,
[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-(2-methyl-pyrrolidin-1-yl)-
methanone,
6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid tert-butylamide,
20 6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid cyclopropylmethyl-
propyl-amide,
{[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-(2-methyl-piperidin-1-yl)-
methanone,
[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-(3-methyl-piperidin-1-yl)-
25 methanone,
6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid isopropyl-methyl-
amide,
[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-thiomorpholin-4-yl-methanone,
[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-(2-trifluoromethyl-pyrrolidin-1-yl)-
30 methanone,
6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid ethyl-(2-methoxy-
ethyl)-amide,
azetidin-1-yl-[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-methanone,
(3,6-dihydro-2H!-pyridin-1-yl)-[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-
35 methanone,
6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid (3-fluoro-oxetan-3-
ylmethyl)-amide,
6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid (1-methoxymethyl-


- cyclopropylmethyl)-amide,
 6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid [2-(tetrahydro-pyran-4-yl)-ethyl]-amide,
 (4,4-difluoro-piperidin-1-yl)-[6-(1-isopropyl-pyrrolidin-3-yloxy)-quinolin-2-yl]-
 5 methanone,
 [6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-morpholin-4-yl-methanone,
 (4-methoxy-piperidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
 (4-methyl-piperidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
 morpholin-4-yl-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
 10 (2-methyl-pyrrolidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid cyclopropylmethyl-propyl-
 amide,
 [6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-(2-trifluoromethyl-pyrrolidin-1-yl)-
 methanone,
 15 (2,5-dihydro-pyrrol-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid ethyl-(2-methoxy-ethyl)-
 amide,
 azetidin-1-yl-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
 (3,6-dihydro-2H-pyridin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-
 20 methanone,
 (4,4-difluoro-piperidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid cyclohexyl-methyl-amide,
 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid (1-methoxymethyl-
 cyclopropylmethyl)-amide,
 25 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid (3-fluoro-oxetan-3-
 ylmethyl)-amide,
 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid [2-(tetrahydro-pyran-4-yl)-
 ethyl]-amide,
 6-(3-piperidin-1-yl-propoxy)-quinoline-2-carboxylic acid (tetrahydro-pyran-4-yl)-
 30 amide,
 (2-methyl-piperidin-1-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
 [6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-pyrrolidin-1-yl-methanone,
 (R)-1-[6-(3-piperidin-1-yl-propoxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile,
 (1,1-dioxo--thiomorpholin-4-yl)-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-
 35 methanone,
 (4-methyl-piperidin-1-yl)-[6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl]-
 methanone,
 [6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl]-pyrrolidin-1-yl-methanone,

- (R)-1-[6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile,
 (1,1-dioxo-thiomorpholin-4-yl)-{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-methanone,
 5 (4-methoxy-piperidin-1-yl)-{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-methanone,
 {6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-morpholin-4-yl-methanone,
 azetidin-1-yl-{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-methanone,
 6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid [2-(tetrahydro-
 10 pyran-4-yl)-ethyl]-amide,
 6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinoline-2-carboxylic acid (3-fluoro-oxetan-3-ylmethyl)-amide,
 (2-methyl-pyrrolidin-1-yl)-{6-[2-(1-methyl-pyrrolidin-2-yl)-ethoxy]-quinolin-2-yl}-methanone,
 15 (S)-1-[6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile,
 {6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl}-(4-methyl-piperidin-1-yl)-methanone,
 (4-hydroxymethyl-piperidin-1-yl)-{6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl}-
 20 methanone,
 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid isobutyl-amide,
 6-(1-isopropyl-pyrrolidin-3-yloxy)-quinoline-2-carboxylic acid cyclohexyl-methyl-amide,
 and pharmaceutically acceptable salts thereof.
- 25 18. Compounds of formula I according to claim 1, selected from the group consisting of
 {6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl}-(4-methoxy-piperidin-1-yl)-methanone,
 {6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl}-(2-methyl-pyrrolidin-1-yl)-
 30 methanone,
 {6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl}-(2-trifluoromethyl-pyrrolidin-1-yl)-methanone,
 azetidin-1-yl-[6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl]-methanone,
 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid (3-fluoro-oxetan-3-
 35 ylmethyl)-amide,
 6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carboxylic acid (1-methoxymethyl-cyclopropylmethyl)-amide,
 {6-(1-isopropyl-piperidin-4-yloxy)-quinolin-2-yl}-pyrrolidin-1-yl-methanone,

- (R)-1-[6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile,
 azetidin-1-yl-[6-(3-piperidin-1-yl-propoxy)-quinolin-2-yl]-methanone,
 (R)-1-[6-(3-piperidin-1-yl-propoxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile,
 5 (S)-1-[6-(1-isopropyl-piperidin-4-yloxy)-quinoline-2-carbonyl]-pyrrolidine-2-carbonitrile,
 and pharmaceutically acceptable salts thereof.

19. A process for the manufacture of compounds according to any one of claims 1 to 18, which process comprises

- 10 a) reacting a compound of the formula II




II

wherein R is lower alkyl,
 with an alcohol of the formula III




- 15 wherein A is as defined in claim 1,
 in the presence of a trialkylphosphine or triphenylphosphine and of a diazo compound
 to obtain a compound of the formula IV



IV

and converting the ester of formula IV into the acid of formula V


- 65 -




V

under acidic or basic conditions,

and coupling the compound of formula V with an amine of the formula VI



- 5 wherein R^1 and R^2 are as defined in claim 1, with the help of an coupling agent
under basic conditions to obtain a compound of the formula I



I

wherein A, R^1 and R^2 are as defined in claim 1,
and if desired,

- 10 converting the compound obtained into a pharmaceutically acceptable salt.

20. Compounds according to any one of claims 1 to 18 when manufactured by a
process according to claim 19.

21. Pharmaceutical compositions comprising a compound according to any one of
claims 1 to 18 and a pharmaceutically acceptable carrier and/or adjuvant.

- 15 22. Pharmaceutical compositions according to claim 21 for the treatment and/or
prevention of diseases which are associated with the modulation of H3 receptors.

23. Compounds according to any one of claims 1 to 18 for use as therapeutically
active substances.

- 20 24. Compounds according to any one of claims 1 to 18 for use as therapeutically
active substances for the treatment and/or prevention of diseases which are associated
with the modulation of H3 receptors.

25. The use of compounds according to any one of claims 1 to 18 for the preparation of medicaments for the treatment and/or prevention of diseases which are associated with the modulation of H3 receptors.

5

26. The use according to claim 25 for the treatment and/or prevention of obesity.