


ORGANIC COMPOUNDS

This invention relates to organic compounds, their preparation and use as pharmaceuticals.

An aspect of the present invention provides compounds of formula (I) or stereoisomers or pharmaceutically acceptable salts thereof,



wherein

R^1 is selected from $\text{NH-C}_1\text{-C}_8\text{-alkyl}$, $\text{NHC(O)C}_1\text{-C}_8$ hydroxyalkyl, $\text{NHC(O)C}_1\text{-C}_8$ aminoalkyl, $\text{NHCO}_2\text{C}_1\text{-C}_8\text{-alkyl}$, $\text{NHCO}_2\text{C}_2\text{-C}_8$ hydroxyalkyl, NHC(O)-3-to 12- membered heterocyclic group containing from 1 to 4 heteroatoms selected from the group consisting of oxygen, nitrogen and sulfur, that group being optionally substituted, by $\text{C}_1\text{-C}_8\text{-alkyl}$, $\text{NHC(O)-C}_6\text{-C}_{10}\text{-aryl}$ optionally substituted by $\text{C}_1\text{-C}_8\text{-alkyl}$ or $\text{O-C}_1\text{-C}_8\text{-alkyl}$, $\text{NH-C}_1\text{-C}_8\text{-alkoxycarbonyl}$ optionally substituted by $\text{C}_6\text{-C}_{10}\text{-aryl}$, and $\text{NHC(O)-C}_1\text{-C}_8\text{-alkyl}$ optionally substituted by halo, OH, $\text{C}_1\text{-C}_8\text{-alkyl}$, COOH or $\text{C}_1\text{-C}_8\text{-alkoxycarbonyl}$;

R^2 is $\text{C}_1\text{-C}_8\text{-alkyl}$ substituted by OH, halogen $\text{C}_6\text{-C}_{10}\text{-aryl}$ optionally substituted by OH, SO_2R^{10} , $\text{SC}_1\text{-C}_8\text{-alkyl}$, CN, halogen, $\text{O-C}_7\text{-C}_{14}\text{-aralkyl}$, or $\text{O-C}_1\text{-C}_8\text{-alkyl}$, a $\text{C}_3\text{-C}_{15}\text{-carbocyclic}$ group optionally substituted by $\text{O-C}_7\text{-C}_{14}$ aralkyl, $\text{C}_3\text{-C}_{15}\text{-carbocyclic}$ group, $\text{O-C}_1\text{-C}_8\text{-alkyl}$, $\text{C}_2\text{-C}_8\text{-alkenyl}$, $\text{C}_2\text{-C}_8\text{-alkynyl}$ or $\text{C}_1\text{-C}_8\text{-alkyl}$, $\text{O-C}_1\text{-C}_8\text{-alkyl}$, $\text{-SO}_2\text{-C}_1\text{-C}_8\text{-alkyl}$, a 3- to 12-membered heterocyclic group containing from 1 to 4 ring nitrogen atoms and optionally containing from 1 to 4 other heteroatoms selected from the group consisting of oxygen and sulfur, that group being optionally

substituted by 3- to 12-membered heterocyclic group containing from 1 to 4 ring nitrogen atoms and optionally containing from 1 to 4 other heteroatoms selected from the group consisting of oxygen and sulfur, C₇-C₁₄ aralkyl, or C₆-C₁₄-aryl optionally substituted by O-C₇-C₁₄ aralkyl, or

R² is a C₃-C₁₅-carbocyclic group optionally substituted by O-C₇-C₁₄ aralkyl, C₃-C₁₅-carbocyclic group, O-C₁-C₈-alkyl, or C₁-C₈-alkyl, or

R² is a 3- to 12-membered heterocyclic group containing from 1 to 4 ring nitrogen atoms and optionally containing from 1 to 4 other heteroatoms selected from the group consisting of oxygen and sulfur, that group being optionally substituted by 3- to 12-membered heterocyclic group containing from 1 to 4 ring nitrogen atoms and optionally containing from 1 to 4 other heteroatoms selected from the group consisting of oxygen and sulfur, C₇-C₁₄ aralkyl, or C₆-C₁₄-aryl optionally substituted by O-C₇-C₁₄ aralkyl;

R³ is hydrogen, halo, C₂-C₈-alkenyl, C₂-C₈-alkynyl or C₁-C₈-alkoxycarbonyl, or

R³ is amino optionally substituted by C₃-C₈-cycloalkyl optionally substituted by amino, hydroxy, C₇-C₁₄-aralkyloxy, -SO₂-C₆-C₁₀-aryl or -NH-C(=O)-NH-R^{3c}, or

R³ is amino substituted by R^{3a}, -R^{3a}-C₇-C₁₄-aralkyl or a C₅-C₁₅-carbocyclic group optionally substituted by OH, C₁-C₈-alkyl or C₁-C₈-alkoxycarbonyl, or

R³ is aminocarbonyl optionally substituted by R^{3b}, or

R³ is C₁-C₈-alkylamino optionally substituted by OH, R^{3b}, amino, di(C₁-C₈-alkyl)amino, -NH-C(=O)-C₁-C₈-alkyl, -NH-SO₂-C₁-C₈-alkyl, -NH-C(=O)-NH-R^{3c}, -NH-C(=O)-NH-C₁-C₈-alkyl-R^{3b}, a C₅-C₁₅-carbocyclic group or by C₆-C₁₀-aryl optionally substituted by C₆-C₁₀-aryloxy, or

R^3 is C_1 - C_8 -alkylaminocarbonyl or C_3 - C_8 -cycloalkylamino-carbonyl optionally substituted by amino, C_1 - C_8 -alkylamino, di(C_1 - C_8 -alkyl)amino or $-NH-C(=O)-NH-R^{3d}$, or

R^3 is a 3- to 12-membered heterocyclic group containing from 1 to 4 ring nitrogen atoms and optionally containing from 1 to 4 other heteroatoms selected from the group consisting of oxygen and sulfur, that group being optionally substituted by 0-3 R^4 ;

R^{3a} and R^{3b} are each independently a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur; optionally substituted by halo, cyano, oxo, OH, carboxy, nitro, C_1 - C_8 -alkyl, C_1 - C_8 -alkylcarbonyl, OH- C_1 - C_8 -alkyl, C_1 - C_8 -haloalkyl, amino- C_1 - C_8 -alkyl, amino(OH) C_1 - C_8 -alkyl and C_1 - C_8 -alkoxy optionally substituted by aminocarbonyl;

R^{3c} is a 5- or 6-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, which is optionally substituted by a 5- or 6-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur;

R^{3d} are independently a 5- or 6-membered heterocyclic ring containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, said 5- or 6-membered heterocyclic ring being optionally substituted by halo, cyano, oxo, OH, carboxy, amino, nitro, C_1 - C_8 -alkyl, C_1 - C_8 -alkylsulfonyl, aminocarbonyl, C_1 - C_8 -alkylcarbonyl, C_1 - C_8 -alkoxy optionally substituted by aminocarbonyl, or a 5- or 6-membered heterocyclic ring containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, said ring also being optionally substituted by halo, cyano, oxo, OH, carboxy, amino, nitro, C_1 - C_8 -alkyl, C_1 - C_8 -alkylsulfonyl, aminocarbonyl, C_1 - C_8 -alkylcarbonyl, C_1 - C_8 -alkoxy optionally substituted by aminocarbonyl;

R^4 is selected from OH, C_1 - C_8 -alkyl optionally substituted by OH, C_1 - C_8 -alkoxy, C_7 - C_{14} -aralkyl optionally substituted with OH, O- C_1 - C_8 -alkyl, halogen C_6 - C_{10} -aryl, or O- C_6 - C_{10} -aryl, C_1 - C_8 -alkoxy, C_6 - C_{10} -aryl optionally substituted by OH, C_1 - C_8 -alkyl, O- C_1 - C_8 -alkyl or -halogen, O- C_6 - C_{10} -aryl optionally substituted by OH, C_1 - C_8 -alkyl, O- C_1 - C_8 -alkyl or -halogen, $NR^{4a}R^{4b}$, $NHC(O)R^{4c}$, $NHS(O)_2R^{4d}$, $NHS(O)_2R^{4e}$, $NR^{4f}C(O)NR^{4e}R^{4h}$, $NR^{4f}C(O)NR^{4g}R^{4h}$, $NR^{4i}C(O)OR^{4j}$, C_1 - C_8 -alkylcarbonyl, C_1 - C_8 -alkoxycarbonyl, di(C_1 - C_8 -alkyl)aminocarbonyl, $COOR^{4k}$, $C(O)R^{4l}$, $NHC(O)R^{4q}$, $NHC(=NR^{4m})N(R^{4n})R^{4o}$, and a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur optionally substituted by $COOR^{4p}$;

R^{4a} , R^{4c} , R^{4f} , R^{4h} and R^{4i} are, independently, H, or C_1 - C_8 -alkyl;

R^{4b} is H, C_1 - C_8 -alkyl a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by $0-3R^5$ or C_6 - C_{10} -aryl;

R^{4d} , R^{4e} , and R^{4j} are, independently, C_1 - C_8 -alkyl or a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by $0-3R^5$;

R^{4g} is C_1 - C_8 -alkyl optionally substituted by a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted with SO_2R^{10} , CN, or $0-3R^5$, or

R^{4g} is a C_6 - C_{10} -aryl optionally substituted by OH, C_1 - C_8 -alkyl, O- C_1 - C_8 -alkyl, SO_2R^{10} or -halogen, or

R^{4g} is a C_7 - C_{14} -aralkyl optionally substituted by OH, O- C_1 - C_8 -alkyl, halogen, C_6 - C_{10} -aryl, SO_2R^{10} , CN, $-C(=NH)NH_2$, or O- C_6 - C_{10} -aryl, or

R^{4g} is a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by 0-3R⁵;

R^{4k} is H, C₁-C₈-alkyl, C₆-C₁₀-aryl or a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur;

R^{4l} is C₁-C₈-alkyl, C₆-C₁₀-aryl, NHR⁶ or a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur;

R^{4m} is H or CN;

R⁴ⁿ is H or C₁-C₈ alkyl;

R^{4o} is H, C₁-C₈-alkyl optionally substituted by OH or by a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted with SO₂R¹⁰, CN, or 0-3R⁵, C₁-C₈-alkoxy, C₇-C₁₄-aralkyl optionally substituted with OH, O-C₁-C₈-alkyl, halogen C₆-C₁₀-aryl, or O-C₆-C₁₀-aryl, C₁-C₈-alkoxy, C₆-C₁₀-aryl optionally substituted by OH, C₁-C₈-alkyl, O-C₁-C₈-alkyl SO₂R¹⁰ or -halogen;

R^{4p} is H, C₁-C₈-alkyl or C₇-C₁₄-aralkyl;

R^{4q} is C₆-C₁₀-aryl optionally substituted by OH, C(=NH)NH₂, or SO₂NH₂, or a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur optionally substituted by 0-3R⁵ or a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by 0-3R⁵;

R^5 is selected from OH, C_1 - C_8 -alkyl optionally substituted by OH, CN, SO_2R^{10} or halogen, C_7 - C_{14} -aralkyl optionally substituted with OH, O- C_1 - C_8 -alkyl, C_6 - C_{10} -aryl, or O- C_6 - C_{10} -aryl, C_1 - C_8 -alkoxy, C_6 - C_{10} -aryl optionally substituted by OH, C_1 - C_8 -alkyl, O- C_1 - C_8 -alkyl or -halogen, O- C_6 - C_{10} -aryl optionally substituted by OH, C_1 - C_8 -alkyl, O- C_1 - C_8 -alkyl optionally substituted by halogen, $NR^{5a}R^{5b}$, $NHC(O)R^{5c}$, $NHS(O)_2R^{5d}$, $NHS(O)_2R^{5e}$, $NR^{5f}C(O)NR^{5g}R^{5h}$, $NR^{5i}C(O)OR^{5j}$, C_1 - C_8 -alkylcarbonyl, C_1 - C_8 -alkoxycarbonyl, di(C_1 - C_8 -alkyl)aminocarbonyl, $COOR^{5k}$, $C(O)R^{5l}$, a $C(O)$ - C_6 - C_{10} -aryl optionally substituted by OH, -COOH, C_1 - C_8 -alkyl, O- C_1 - C_8 -alkyl, -halogen, or SO_2R^{10} , $C(O)NHR^{5m}$ or a 3-12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by $O-3R^7$;

R^{5a} , R^{5b} , R^{5c} , R^{5f} , R^{5h} and R^{5i} are, independently, H, C_1 - C_8 -alkyl or C_6 - C_{10} -aryl;

R^{5d} , R^{5e} , R^{5g} , R^{5j} and R^{5m} are, independently, C_1 - C_8 -alkyl or a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by $COOR^8$;

R^{5k} is H, C_1 - C_8 -alkyl, C_6 - C_{10} -aryl or a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur;

R^{5l} is C_1 - C_8 -alkyl, C_6 - C_{10} -aryl or a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by $COOR^9$;

R^6 is $COOR^{6a}$ or a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by $COOR^{6b}$;

R^{6a} , R^{6b} , R^7 , R^8 and R^9 are selected from H, C_1 - C_8 -alkyl and C_7 - C_{14} -aralkyl; and

R^{10} is C_1 - C_8 -alkyl optionally substituted by halogen, C_6 - C_{10} -aryl optionally substituted by OH, C_1 - C_8 -alkyl, O- C_1 - C_8 -alkyl or -halogen, or $NR^{4a}R^{4b}$.

According to formula (I) R^1 is suitably $NHC(O)C_1$ - C_8 hydroxyalkyl. R^1 is preferably $NHC(O)C_1$ - C_2 hydroxyalkyl (e.g. a 2-hydroxy-acetamide group, a 2-hydroxy-propionamide group, or a 3-hydroxy-propionamide group).

According to formula (I), R^1 is also suitably $NHCO_2C_1$ - C_8 -alkyl, R^1 is preferably $NHCO_2CH_3$.

According to formula (I), R^2 is suitably selected from C_1 - C_8 -alkyl optionally substituted by OH, halogen or C_6 - C_{10} -aryl optionally substituted by OH or O- C_1 - C_8 alkyl, preferably C_6 - C_{10} -aryl is phenyl substituted by one OCH_3 or one OH.

According to formula (I), R^3 is suitably a N-bonded 3-to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur. This heterocyclic group is preferably a pyrrolidine, or pyrazole. The heterocyclic group is optionally substituted by $NR^{4f}C(O)NR^{4g}R^{4h}$, $NR^{4a}R^{4b}$, $NHC(O)R^{4q}$ and $NHC(=NR^{4m})N(R^{4n})R^{4o}$, where R^{4a} is preferably H or C_1 - C_8 alkyl (e.g. methyl) and R^{4f} and R^{4h} are preferably H. R^{4b} is selected from H and 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by 0-3 R^5 . R^{4b} is preferably H or a 4,5 dihydro-1H imidazole.

According to formula (I), R^{4g} is suitably C_1 - C_8 -alkyl optionally substituted by a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted with SO_2R^{10} , CN, or 0-3 R^5 . R^{4g} is preferably a methylene substituted by a pyridine where the pyridine is optionally substituted by one CN.

According to formula (I), R^{4g} is also suitably C_6 - C_{10} -aryl optionally substituted by OH, C_1 - C_8 -alkyl, O- C_1 - C_8 -alkyl, SO_2R^{10} , or -halogen, R^{4g} is preferably a phenyl that is optionally substituted by one OH or one SO_2NH_2 .

According to formula (I), R^{4g} is also suitably C_7 - C_{14} -aralkyl optionally substituted by OH, O- C_1 - C_8 -alkyl, halogen, C_6 - C_{10} -aryl, SO_2R^{10} , CN, $-C(=NH)NH_2$, or O- C_6 - C_{10} -aryl. R^{4g} is preferably a benzyl group optionally substituted by one OH or one $-C(=NH)NH_2$.

According to formula (I), R^{4g} is also suitably 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by 0-3 R^5 ; R^{4g} is preferably a pyridine optionally substituted by CN or pyrrolidine substituted by a C(O)- C_6 - C_{10} -aryl optionally substituted by OH, -COOH, C_1 - C_8 -alkyl, O- C_1 - C_8 -alkyl, -halogen, or SO_2R^{10} . Preferably the C(O)- C_6 - C_{10} -aryl is C(O)-benzoic acid.

According to formula (I), R^{4m} is CN.


According to formula (I) R^{4n} is H or C_1 - C_8 alkyl. Preferably, R^{4n} is H.

According to formula (I), R^{4o} is H, C_1 - C_8 -alkyl optionally substituted by OH or by a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted with SO_2R^{10} , CN, or 0-3 R^5 , C_1 - C_8 -alkoxy, C_7 - C_{14} -aralkyl optionally substituted with OH, O- C_1 - C_8 -alkyl, halogen C_6 - C_{10} -aryl, or O- C_6 - C_{10} -aryl, C_1 - C_8 -alkoxy, C_6 - C_{10} -aryl optionally substituted by OH, C_1 - C_8 -alkyl, O- C_1 - C_8 -alkyl SO_2R^{10} or -halogen. R^{4o} is preferably a methylene substituted by an unsubstituted pyridine or phenyl optionally substituted by SO_2NH_2 .

According to formula (I), R^{4q} is suitably phenyl substituted by OH, $C(=NH)NH_2$, or SO_2NH_2 .

According to formula (I), R^{4q} is also suitably 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur optionally substituted by a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by 0-3 R^5 . Preferably R^{4q} is a 6 membered heterocyclic group (e.g. pyridine) substituted by a 6 membered heterocyclic group (e.g. morpholine).

Another aspect of the invention is compounds of formula Ia



where

R^1 is $\text{NHCO}_2\text{C}_1\text{-C}_8\text{-alkyl}$ or $\text{NH C(O) C}_1\text{-C}_8\text{ hydroxylalkyl}$;

R^4 is $\text{NR}^{4f}\text{C(O)NR}^{4g}\text{R}^{4h}$, $\text{NR}^{4a}\text{R}^{4b}$, NHC(O)R^{4q} and $\text{NHC(=NR}^{4m})\text{N(R}^{4n})\text{R}^{4o}$;

R^{4a} is selected from H and $\text{C}_1\text{-C}_8\text{ alkyl}$;

R^{4b} is selected from H, $\text{C}_1\text{-C}_8\text{ alkyl}$ and 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by 0-3 R^5 ;

R^{4f} and R^{4h} are H;

R^{4g} is C_1 - C_8 -alkyl optionally substituted by a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted with CN, or

R^{4g} is a C_6 - C_{10} -aryl optionally substituted by OH or SO_2R^{10} , or

R^{4g} is a C_7 - C_{14} -aralkyl optionally substituted by OH - $C(=NH)NH_2$, or

R^{4g} is a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by $O-3R^5$;

R^{4m} is CN;

R^{4n} is H;

R^{4o} is H, C_1 - C_8 -alkyl optionally substituted by OH or by a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted with SO_2R^{10} , a C_7 - C_{14} -aralkyl optionally substituted with OH, $O-C_1$ - C_8 -alkyl, halogen, C_6 - C_{10} -aryl, or $O-C_6$ - C_{10} -aryl;

R^{4q} is C_6 - C_{10} -aryl optionally substituted by OH, $C(=NH)NH_2$, or SO_2NH_2 , or a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur optionally substituted by a 3- to 12-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur;


R^5 is $C(O)-C_6-C_{10}$ -aryl optionally substituted by OH, $-COOH$, C_1-C_8 -alkyl, $O-C_1-C_8$ -alkyl, -halogen, or SO_2R^{10} ;

R^{10} is NH_2 ;

R^{11} and R^{12} are, independently, selected from H, OH, halogen and $O-C_1-C_8$ -alkyl; and

R^{13} is selected from H or OH.

An aspect of the present invention provides compounds of formula (II) or stereoisomers or pharmaceutically acceptable salts thereof,



wherein

W is CH_2 or O, with the proviso that when W is O, then R^{11a} is not a N-bonded substituent;

R^{11a} is $-NH_2$, $-NH-C_1-C_8$ -alkylcarbonyl, $-NH-C_3-C_8$ -cycloalkylcarbonyl, $-NHSO_2-C_1-C_8$ -alkyl, $-NH-C_7-C_{14}$ -aralkylcarbonyl or $-NHC(=O)-C(=O)-NH-C_1-C_8$ -alkyl optionally substituted by R^{11b} , or

R^{11a} is selected from CH_2OH , $CH_2-O-C_1-C_8$ -alkyl, $C(O)-O-C_1-C_8$ -alkyl, $C(O)NH_2$, and $C(O)-NH-C_1-C_8$ -alkyl;

R^{11b} is a 3- to 12-membered heterocyclic ring containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, said 3- to 12-membered heterocyclic ring being optionally substituted by halo, cyano, oxo, hydroxy, carboxy, amino, nitro, C₁-C₈-alkyl, C₁-C₈-alkylsulfonyl, aminocarbonyl, C₁-C₈-alkylcarbonyl or C₁-C₈-alkoxy optionally substituted by aminocarbonyl;

R^{12a} is C₁-C₈-alkyl substituted by OH, halogen C₆-C₁₀-aryl optionally substituted by OH, SO₂R¹⁰, SC₁-C₈-alkyl, CN, halogen, O-C₇-C₁₄-aralkyl, or O-C₁-C₈-alkyl, a C₃-C₁₅-carbocyclic group optionally substituted by O-C₇-C₁₄ aralkyl, C₃-C₁₅-carbocyclic group, O-C₁-C₈-alkyl, C₂-C₈-alkenyl, C₂-C₈-alkynyl or C₁-C₈-alkyl, O-C₁-C₈-alkyl, -SO₂-C₁-C₈-alkyl, a 3- to 12-membered heterocyclic group containing from 1 to 4 ring nitrogen atoms and optionally containing from 1 to 4 other heteroatoms selected from the group consisting of oxygen and sulfur, that group being optionally substituted by 3- to 12-membered heterocyclic group containing from 1 to 4 ring nitrogen atoms and optionally containing from 1 to 4 other heteroatoms selected from the group consisting of oxygen and sulfur, C₇-C₁₄ aralkyl, or C₆-C₁₄-aryl optionally substituted by O-C₇-C₁₄ aralkyl, or

R^{12a} is a C₃-C₁₅-carbocyclic group optionally substituted by O-C₇-C₁₄ aralkyl, C₃-C₁₅-carbocyclic group, O-C₁-C₈-alkyl, or C₁-C₈-alkyl, or

R^{12a} is a 3- to 12-membered heterocyclic group containing from 1 to 4 ring nitrogen atoms and optionally containing from 1 to 4 other heteroatoms selected from the group consisting of oxygen and sulfur, that group being optionally substituted by 3- to 12-membered heterocyclic group containing from 1 to 4 ring nitrogen atoms and optionally containing from 1 to 4 other heteroatoms selected from the group consisting of oxygen and sulfur, C₇-C₁₄ aralkyl, or C₆-C₁₄-aryl optionally substituted by O-C₇-C₁₄ aralkyl;

R^{14} is selected from $NR^{14a}R^{14b}$, $NR^{14f}C(O)NR^{14g}R^{14h}$, $NHC(O)R^{14q}$, and $NHC(=NR^{14m})N(R^{14n})R^{14o}$;

R^{14a} , R^{14c} , R^{14f} , R^{14h} and R^{14i} are, independently, H, C_1 - C_8 -alkyl or C_6 - C_{10} -aryl;

R^{14b} is H, C_1 - C_8 -alkyl a 3- to 10-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by $O-3R^{15}$ or C_6 - C_{10} -aryl;

R^{14g} is C_1 - C_8 -alkyl optionally substituted by a 3- to 10-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted with SO_2R^{16} , CN, or $O-3R^{15}$, or

R^{14g} is a C_6 - C_{10} -aryl optionally substituted by OH, C_1 - C_8 -alkyl, $O-C_1$ - C_8 -alkyl, SO_2R^{16} or -halogen, or

R^{14g} is a C_7 - C_{14} -aralkyl optionally substituted by OH, $O-C_1$ - C_8 -alkyl, halogen, C_6 - C_{10} -aryl, SO_2R^{16} , CN, $-C(=NH)NH_2$, or $O-C_6$ - C_{10} -aryl, or

R^{14g} is a 3- to 10-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by $O-3R^{15}$;

R^{14m} is CN;

R^{14n} is H or C_1 - C_8 alkyl;

R^{14o} is H, C_1 - C_8 -alkyl optionally substituted by OH or by a 3- to 10-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted with SO_2R^{16} , CN, or $O-3R^{15}$, C_1 - C_8 -alkoxy, C_7 - C_{14} -aralkyl optionally substituted with OH, $O-C_1$ - C_8 -

alkyl, halogen C₆-C₁₀-aryl, or O-C₆-C₁₀-aryl, C₁-C₈-alkoxy, C₆-C₁₀-aryl optionally substituted by OH, C₁-C₈-alkyl, O-C₁-C₈-alkyl SO₂R¹⁶ or -halogen;

R^{14p} is H, C₁-C₈-alkyl or C₇-C₁₄-aralkyl;

R^{14q} is a 3- to 10-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur optionally substituted by 0-3R¹⁵ or a 3- to 10-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by 0-3R¹⁵;

R¹⁵ is selected from CN, or halogen, O-C₁-C₈-alkyl optionally substituted by halogen, a C(O)-C₆-C₁₀-aryl optionally substituted by OH, -COOH, C₁-C₈-alkyl, O-C₁-C₈-alkyl, -halogen, or SO₂R¹⁶; and

R¹⁶ is C₁-C₈-alkyl optionally substituted by halogen, C₆-C₁₀-aryl optionally substituted by OH, C₁-C₈-alkyl, O-C₁-C₈-alkyl or -halogen, or NR^{14a}R^{14b}.

According to formula (II), R^{11a} is suitably a 3- to 10-membered heterocyclic group containing from 1 to 4 ring nitrogen atoms and optionally containing from 1 to 4 other heteroatoms selected from the group consisting of oxygen and sulfur, that group being optionally substituted by oxo, C₁-C₈-alkoxy, C₆-C₁₀-aryl, R^{11b} or by C₁-C₈-alkyl optionally substituted by hydroxyl. R^{11a} is preferably tetrazole substituted by an ethyl group or a triazole substituted by an ethanol group.

According to formula (II), R^{11a} is also suitably -NH-C₁-C₈-alkylcarbonyl or -NH-C₃-C₈-cycloalkylcarbonyl. The -NH-C₁-C₈-alkylcarbonyl group is preferably a acetamide group or a propionamide group. The -NH-C₃-C₈-cycloalkylcarbonyl is preferably a cyclobutane carboxylic acid amide group.

According to formula (II), R^{11a} is also suitably $C(O)-NH-C_1-C_8$ -alkyl, preferably $C(O)-NH$ -ethyl.

According to formula (II), R^{12a} is suitably selected from C_1-C_8 -alkyl optionally substituted by OH, halogen or C_6-C_{10} -aryl optionally substituted by OH or $O-C_1-C_8$ alkyl, preferably C_6-C_{10} -aryl is phenyl substituted by OCH_3 or OH.

According to formula (II), R^{14} is suitably $NR^{14f}C(O)NR^{14g}R^{14h}$, $NR^{14a}NR^{14b}$, $NHC(O)R^{14q}$ and $NHC(=NR^{14m})N(R^{14n})R^{14o}$, where R^{14a} is preferably H or C_1-C_8 alkyl (e.g. methyl) and R^{14f} and R^{14h} are preferably H. R^{14b} is selected from H and 3- to 10-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by 0-3 R^{15} . R^{14b} is preferably H or a 4,5 dihydro-1H imidazole.

According to formula (II), R^{14g} is C_1-C_8 -alkyl optionally substituted by a 3- to 10-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted with SO_2R^{16} , CN, or 0-3 R^{15} . R^{14g} is preferably a methylene substituted by a pyridine where the pyridine is optionally substituted by CN.

According to formula (II), R^{14g} is also suitably C_6-C_{10} -aryl optionally substituted by OH, C_1-C_8 -alkyl, $O-C_1-C_8$ -alkyl, SO_2R^{16} , or -halogen, R^{14g} is preferably a phenyl that is optionally substituted by OH or SO_2NH_2 .

According to formula (II), R^{14g} is also suitably C_7-C_{14} -aralkyl optionally substituted by OH, $O-C_1-C_8$ -alkyl, halogen, C_6-C_{10} -aryl, SO_2R^{16} , CN, $-C(=NH)NH_2$, or $O-C_6-C_{10}$ -aryl. R^{14g} is preferably a benzyl group optionally substituted by OH or $-C(=NH)NH_2$.

According to formula (II), R^{14g} is also suitably 3- to 10-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by 0-3 R^{15} ; R^{14g} is preferably a pyridine

optionally substituted by CN or pyrrolidine substituted by a C(O)-C₆-C₁₀-aryl optionally substituted by OH, -COOH, C₁-C₈-alkyl, O-C₁-C₈-alkyl, -halogen, or SO₂R¹⁶. Preferably the C(O)-C₆-C₁₀-aryl is C(O)-benzoic acid.

According to formula (II), R^{14m} is CN.

According to formula (II) R¹⁴ⁿ is H or C₁-C₈ alkyl. Preferably, R¹⁴ⁿ is H.

According to formula (II), R^{14o} is H, C₁-C₈-alkyl optionally substituted by OH or by a 3- to 10-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted with SO₂R¹⁶, CN, or 0-3R¹⁵, C₁-C₈-alkoxy, C₇-C₁₄-aralkyl optionally substituted with OH, O-C₁-C₈-alkyl, halogen C₆-C₁₀-aryl, or O-C₆-C₁₀-aryl, C₁-C₈-alkoxy, C₆-C₁₀-aryl optionally substituted by OH, C₁-C₈-alkyl, O-C₁-C₈-alkyl SO₂R¹⁶ or -halogen. R^{14o} is preferably a methylene substituted by an unsubstituted pyridine or phenyl optionally substituted by SO₂NH₂.

According to formula (II) R^{14q} is a 3- to 10-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur optionally substituted by a 3- to 10-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur, optionally substituted by 0-3R¹⁵. Preferably R^{14q} is a 6 membered heterocyclic group (e.g. pyridine) substituted by a 6 membered heterocyclic group (e.g. morpholine).

DEFINITIONS

Terms used in the specification have the following meanings:

"Optionally substituted" means the group referred to can be substituted at one or more positions by any one or any combination of the radicals listed thereafter.

"Halo" or "halogen", as used herein, may be fluorine, chlorine, bromine or iodine. Preferably halo is chlorine.

"C₁-C₈-alkyl", as used herein, denotes straight chain or branched alkyl having 1-8 carbon atoms. Preferably C₁-C₈-alkyl is C₁-C₄-alkyl.

"C₁-C₈-alkoxy", or as used herein, denotes straight chain or branched alkoxy having 1-8 carbon atoms. Preferably, C₁-C₈-alkoxy is C₁-C₄-alkoxy.

"C₃-C₈-cycloalkyl", as used herein, denotes cycloalkyl having 3-8 ring carbon atoms, e.g., a monocyclic group, such as a cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl or cyclooctyl, any of which can be substituted by one or more, usually one or two, C₁-C₄-alkyl groups; or a bicyclic group, such as bicycloheptyl or bicyclooctyl.

"C₁-C₈-alkylamino" and "di(C₁-C₈-alkyl)amino", as used herein, denote amino substituted respectively by one or two C₁-C₈-alkyl groups as hereinbefore defined, which may be the same or different.

"C₁-C₈-alkylcarbonyl" and "C₁-C₈-alkoxycarbonyl", as used herein, denote C₁-C₈-alkyl or C₁-C₈-alkoxy, respectively, as hereinbefore defined attached by a carbon atom to a carbonyl group.

"C₆-C₁₀-aryl", as used herein, denotes a monovalent carbocyclic aromatic group that contains 6-10 carbon atoms and which may be, e.g., a monocyclic group, such as phenyl; or a bicyclic group, such as naphthyl.

"C₇-C₁₄-aralkyl", as used herein, denotes alkyl, e.g., C₁-C₄-alkyl, as hereinbefore defined, substituted by C₆-C₁₀-aryl as hereinbefore defined. Preferably, C₇-C₁₄-aralkyl is C₇-C₁₀-aralkyl, such as phenyl-C₁-C₄-alkyl.

"C₁-C₈-alkylaminocarbonyl" and "C₃-C₈-cycloalkylaminocarbonyl" as used herein denote C₁-C₈-alkylamino and C₃-C₈-cycloalkylamino respectively as hereinbefore defined attached by a carbon atom to a carbonyl group. Preferably C₁-C₈-alkylaminocarbonyl and C₃-C₈-cycloalkyl-aminocarbonyl are C₁-C₄-alkylaminocarbonyl and C₃-C₈-cycloalkylaminocarbonyl respectively.

"C₅-C₁₅-carbocyclic group" as used herein denotes a carbocyclic group having 5 to 15 ring carbon atoms, for example a monocyclic group, either aromatic or non-aromatic, such as a cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl or phenyl, or a bicyclic group such as bicyclooctyl, bicyclononyl, bicyclodecyl, indanyl or indenyl, again any of which can be substituted by one or more, usually one or two, C₁-C₄-alkyl groups. Preferably the C₅-C₁₅-carbocyclic group is a C₅-C₁₀-carbocyclic group, especially phenyl, cyclohexyl or indanyl. The C₅-C₁₅-carbocyclic group can unsubstituted or substituted. Preferred substituents on the heterocyclic ring include halo, cyano, OH, carboxy, amino, aminocarbonyl, nitro, C₁-C₁₀-alkyl, C₁-C₁₀-alkoxy and C₃-C₁₀-cycloalkyl, especially OH or amino.

"C₃-C₁₅-carbocyclic group", as used herein, denotes a carbocyclic group having 3-15 ring carbon atoms, e.g., a monocyclic group, either aromatic or non-aromatic, such as a cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl or phenyl; or a bicyclic group, such as bicyclooctyl, bicyclononyl, bicyclodecyl, indanyl or indenyl, again any of which can be substituted by one or more, usually one or two, C₁-C₄-alkyl groups. Preferably the C₃-C₁₅-carbocyclic group is a C₅-C₁₀-carbocyclic group, especially phenyl, cyclohexyl or indanyl. The C₅-C₁₅-carbocyclic group can unsubstituted or substituted. Substituents on the heterocyclic ring include halo, cyano, OH, carboxy, amino, aminocarbonyl, nitro, C₁-C₁₀-alkyl, C₁-C₁₀-alkoxy and C₃-C₁₀-cycloalkyl.

"3- to 12-membered heterocyclic ring containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur", as used herein, may be, e.g., furan, pyrrole, pyrrolidine, pyrazole, imidazole, triazole, isotriazole, tetrazole, thiadiazole, isothiazole, oxadiazole, pyridine, piperidine, pyrazine, oxazole, isoxazole,

pyrazine, pyridazine, pyrimidine, piperazine, pyrrolidine, morpholino, triazine, oxazine or thiazole. Preferred heterocyclic rings include piperazine, pyrrolidine, morpholino, imidazole, isotriazole, pyrazole, tetrazole, thiazole, triazole, thiadiazole, pyridine, piperidine, pyrazine, furan, oxazole, isoxazole, oxadiazole and azetidine. The 3- to-12-membered heterocyclic ring can be unsubstituted or substituted.

"5- or 6-membered heterocyclic group containing at least one ring heteroatom selected from the group consisting of nitrogen, oxygen and sulfur" as used herein may be, for example, a saturated or unsaturated heterocyclic group such as furanyl, pyrrolyl, pyrrolidinyl, pyrazolyl, imidazolyl, triazolyl, isotriazolyl, tetrazolyl, thiadiazolyl, isothiazolyl, oxadiazolyl, pyridinyl, piperidinyl, pyrazinyl, oxazolyl, isoxazolyl, pyrazinyl, pyridazinyl, pyrimidinyl, piperazinyl, pyrrolidinyl, morpholinyl, triazinyl, oxazinyl or thiazolyl. Preferred 5- or 6-membered heterocyclic groups include pyrazolyl, imidazolyl, pyrrolidinyl, pyridinyl and piperidinyl. The 5- or 6-membered heterocyclic group can be unsubstituted or substituted. Preferred substituents include halo, cyano, oxo, OH, carboxy, amino, nitro, C₁-C₈-alkyl (optionally substituted by hydroxy), C₁-C₈-alkylsulfonyl, aminocarbonyl, C₁-C₈-alkylcarbonyl, C₁-C₈-alkoxycarbonyl, and C₁-C₈-alkoxy optionally substituted by aminocarbonyl. Especially preferred substituents include chloro, cyano, carboxy, amino, C₁-C₈-alkoxycarbonyl, C₁-C₄-alkoxy and C₁-C₄-alkyl optionally substituted by OH.

Throughout this specification and in the claims that follow, unless the context requires otherwise, the word "comprise", or variations, such as "comprises" or "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integers or steps.

Throughout this specification and in the claims that follow, unless the context requires otherwise, the word "comprise", or variations, such as "comprises" or "comprising", will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of

integers or steps. As understood by one skilled in the art only combinations of substituents that are chemically possible are embodiments of the invention.

Especially preferred specific compounds of formula (I) and formula (II) are those described hereinafter in the Examples.

Stereoisomers are those compounds where there is an asymmetric carbon atom. The compounds exist in individual optically active isomeric forms or as mixtures thereof, e.g., as diastereomeric mixtures. The present invention embraces both individual optically active *R* and *S* isomers, as well as mixtures thereof. Individual isomers can be separated by methods well known to those skilled in the art, e.g. chiral high performance liquid chromatography (HPLC).

Tautomers are one of two or more structural isomers that exist in equilibrium and are readily converted from one isomeric form to another.


The compounds of the invention may exist in both unsolvated and solvated forms. The term 'solvate' is used herein to describe a molecular complex comprising the compound of the invention and one or more pharmaceutically acceptable solvent molecules, for example, ethanol. The term 'hydrate' is employed when said solvent is water.

SYNTHESIS

Another embodiment of the present invention provides a process for the preparation of compounds of formula (I) in free or pharmaceutically acceptable salt form, which comprises the steps of:

21

(i) reacting a compound of formula (Ib)



wherein

 R^1 , and R^2 are as defined in Claim 1;

Z is H or a protecting group; and

X is a leaving group,

with a compound of formula (Ic)




wherein

 R^3 is as defined in Claim 1; and

removing any protecting groups and recovering the resultant compound of formula (I), in free or pharmaceutically acceptable salt form.

The compound of formula (Ic) may be prepared by reacting a compound of formula (Id)




wherein

 R^1 and Z are as defined in Claim 1; and

L represents a leaving group or a protected derivative thereof with a 2,6-dihalopurine, e.g., 2,6-dichloropurine,

to provide a compound of formula (Ie)



wherein


R^1 and Z are defined in Claim 1; and

X and X^2 are halogen.

Compound of formula (Ie) can be reacted with R^2NH_2 under conventional conditions to provide compound of formula (Ib).

Another embodiment of the present invention provides a process for the preparation of compounds of formula (II) in free or pharmaceutically acceptable salt form, which comprises the steps of:

- (i) reacting a compound of formula (IIa)



wherein


R^{11a} , W and R^{12a} are as defined in Claim 4;

Z is H or a protecting group; and

X is a leaving group,

with a compound of formula (IIb)

23




(IIb)

wherein

R¹⁴ is as defined in Claim 1; and

removing any protecting groups and recovering the resultant compound of formula (I), in free or pharmaceutically acceptable salt form.

The compound of formula (IIb) may be prepared by reacting a compound of formula (IIc)




(IIc)

wherein

R^{11a} and Z are as defined hereinbefore; and

L represents a leaving group or a protected derivative thereof with a 2,6-dihalopurine, e.g., 2,6-dichloropurine,

to provide a compound of formula (IIId)



(IIId)

wherein

R^{11a} and Z are defined hereinbefore; and

X and X² are halogen.

Compound of formula (IIId) can be reacted with R²NH₂ under conventional conditions to provide compound of formula (IIa).

The compounds of formula (I) and formula (II) can be prepared, for example, using the reactions and techniques described below and in the Examples. The compounds of formula (I) and formula (II) can be prepared analogously to the preparations described in Applicant's patent applications PCT/EP2005/011344, GB 0500785.1, and GB 0505219.6. The reactions may be performed in a solvent appropriate to the reagents and materials employed and suitable for the transformations being effected. It will be understood by those skilled in the art of organic synthesis that the functionality present on the molecule should be consistent with the transformations proposed. This will sometimes require a judgment to modify the order of the synthetic steps or to select one particular process scheme over another in order to obtain a desired compound of the invention.

The various substituents on the synthetic intermediates and final products shown in the following reaction schemes can be present in their fully elaborated forms, with suitable protecting groups where required as understood by one skilled in the art, or in precursor forms which can later be elaborated into their final forms by methods familiar to one skilled in the art. The substituents can also be added at various stages throughout the synthetic sequence or after completion of the synthetic sequence. In many cases, commonly used functional group manipulations can be used to transform one intermediate into another intermediate, or one compound of formula (I) into another compound of formula (I) or one compound of formula (II) into another compound of formula (II). Examples of such manipulations are conversion of an ester or a ketone to an alcohol; conversion of an ester to a ketone; interconversions of esters, acids and amides; alkylation, acylation and sulfonylation of alcohols and amines; and many others. Substituents can also be added using common reactions, such as alkylation, acylation, halogenation or oxidation. Such manipulations are well-known in the art, and many reference works summarize procedures and methods for such manipulations. Some reference works which give examples and references to the primary literature of organic synthesis for many functional group manipulations, as well as other transformations

commonly used in the art of organic synthesis are *March's Organic Chemistry*, 5th Edition, Wiley and Chichester, Eds. (2001); *Comprehensive Organic Transformations*, Larock, Ed., VCH (1989); *Comprehensive Organic Functional Group Transformations*, Katritzky et al. (series editors), Pergamon (1995); and *Comprehensive Organic Synthesis*, Trost and Fleming (series editors), Pergamon (1991). It will also be recognized that another major consideration in the planning of any synthetic route in this field is the judicious choice of the protecting group used for protection of the reactive functional groups present in the compounds described in this invention. Multiple protecting groups within the same molecule can be chosen such that each of these protecting groups can either be removed without removal of other protecting groups in the same molecule, or several protecting groups can be removed using the same reaction step, depending upon the outcome desired. An authoritative account describing many alternatives to the trained practitioner is T. W. Greene and P. G. M. Wuts, *Protective Groups In Organic Synthesis*, Wiley and Sons, 1999. It is understood by those skilled in the art that only combinations of substituents that are chemically possible are embodiments of the present invention.

Compounds of formula (I) and formula (II) in free form may be converted into salt form, and vice versa, in a conventional manner. The compounds in free or salt form can be obtained in the form of hydrates or solvates containing a solvent used for crystallisation. Compounds of formula I can be recovered from reaction mixtures and purified in a conventional manner. Isomers, such as stereoisomers, may be obtained in a conventional manner, e.g. by fractional crystallisation or asymmetric synthesis from correspondingly asymmetrically substituted, e.g. optically active, starting materials.

Compounds of formula (I) and formula (II) and their pharmaceutically acceptable salts are useful as pharmaceuticals. In particular, they activate the adenosine A_{2A} receptor, i.e. they act as A_{2A} receptor agonists. Their properties as A_{2A} agonists may be demonstrated using the method described by L. J. Murphree *et al* in *Molecular Pharmacology* 61, 455-462 (2002).

Compounds of the Examples hereinbelow have K_i values below 1.0 μM in the above assay. For example, the compounds of Examples 1, 14, 20, 36, 69, and 178 have K_i values of 0.0083, 0.0025, 0.0016, 0.0030, 0.0043, and 0.0080 μM respectively.

Having regard to their activation of the adenosine A_{2A} receptor, compounds of formula I in free or pharmaceutically acceptable salt form, hereinafter alternately referred to as "agents of the invention", are useful in the treatment of conditions which respond to the activation of the adenosine A_{2A} receptor, particularly inflammatory or allergic conditions. Treatment in accordance with the invention may be symptomatic or prophylactic. Accordingly, agents of the invention are useful in the treatment of inflammatory or obstructive airways diseases, resulting, for example, in reduction of tissue damage, airways inflammation, bronchial hyperreactivity, remodelling or disease progression. Inflammatory or obstructive airways diseases and conditions to which the present invention is applicable include acute lung injury (ALI), adult/acute respiratory distress syndrome (ARDS), chronic obstructive pulmonary, airways or lung disease (COPD, COAD or COLD), including chronic bronchitis or dyspnea associated therewith, emphysema, as well as exacerbation of airways hyperreactivity consequent to other drug therapy, in particular other inhaled drug therapy. The invention is also applicable to the treatment of bronchitis of whatever type or genesis including, e.g., acute, arachidic, catarrhal, croupus, chronic or phthinoïd bronchitis. Further inflammatory or obstructive airways diseases to which the present invention is applicable include bronchiectasis, pneumoconiosis (an inflammatory, commonly occupational, disease of the lungs, frequently accompanied by airways obstruction, whether chronic or acute, and occasioned by repeated inhalation of dusts) of whatever type or genesis, including, for example, aluminosis, anthracosis, asbestosis, chalicosis, ptilosis, siderosis, silicosis, tabacosis and byssinosis.

Other inflammatory or obstructive airways diseases to which the present invention is applicable include asthma of whatever type or genesis including both intrinsic (non-allergic) asthma and extrinsic (allergic) asthma, mild asthma, moderate asthma, severe asthma, bronchitic asthma, exercise-induced asthma, occupational asthma and asthma

induced following bacterial infection. Treatment of asthma is also to be understood as embracing treatment of subjects, e.g. of less than 4 or 5 years of age, exhibiting wheezing symptoms and diagnosed or diagnosable as “wheezy infants”, an established patient category of major medical concern and now often identified as incipient or early-phase asthmatics. (For convenience this particular asthmatic condition is referred to as “wheezy-infant syndrome”).

Prophylactic efficacy in the treatment of asthma will be evidenced by reduced frequency or severity of symptomatic attack, e.g. of acute asthmatic or bronchoconstrictor attack, improvement in lung function or improved airways hyperreactivity. It may further be evidenced by reduced requirement for other, symptomatic therapy, i.e. therapy for or intended to restrict or abort symptomatic attack when it occurs, for example anti-inflammatory (e.g. cortico-steroid) or bronchodilatory. Prophylactic benefit in asthma may in particular be apparent in subjects prone to “morning dipping”. “Morning dipping” is a recognised asthmatic syndrome, common to a substantial percentage of asthmatics and characterised by asthma attack, e.g. between the hours of about 4 to 6 am, i.e. at a time normally substantially distant from any previously administered symptomatic asthma therapy.

Having regard to their anti-inflammatory activity, in particular in relation to inhibition of eosinophil activation, agents of the invention are also useful in the treatment of eosinophil related disorders, e.g. eosinophilia, in particular eosinophil related disorders of the airways (e.g. involving morbid eosinophilic infiltration of pulmonary tissues) including hyper-eosinophilia as it effects the airways and/or lungs as well as, for example, eosinophil-related disorders of the airways consequential or concomitant to Löffler’s syndrome, eosinophilic pneumonia, parasitic (in particular metazoan) infestation (including tropical eosinophilia), bronchopulmonary aspergillosis, polyarteritis nodosa (including Churg-Strauss syndrome), eosinophilic granuloma and eosinophil-related disorders affecting the airways occasioned by drug-reaction.

Agents of the invention are also useful in the treatment of inflammatory or allergic conditions of the skin, for example psoriasis, contact dermatitis, atopic dermatitis, alopecia areata, erythema multiforma, dermatitis herpetiformis, scleroderma, vitiligo, hypersensitivity angiitis, urticaria, bullous pemphigoid, lupus erythematosus, pemphigus, epidermolysis bullosa acquisita, and other inflammatory or allergic conditions of the skin.

Agents of the invention may also be used for the treatment of other diseases or conditions, in particular diseases or conditions having an inflammatory component, for example, treatment of diseases and conditions of the eye such as conjunctivitis, keratoconjunctivitis sicca, and vernal conjunctivitis, diseases affecting the nose including allergic rhinitis, and inflammatory disease in which autoimmune reactions are implicated or having an autoimmune component or aetiology, including autoimmune haematological disorders (e.g. haemolytic anaemia, aplastic anaemia, pure red cell anaemia and idiopathic thrombocytopenia), systemic lupus erythematosus, polychondritis, scleroderma, Wegener granulomatosis, dermatomyositis, chronic active hepatitis, myasthenia gravis, Steven-Johnson syndrome, idiopathic sprue, autoimmune inflammatory bowel disease (e.g. ulcerative colitis and Crohn's disease), endocrine ophthalmopathy, Grave's disease, sarcoidosis, alveolitis, chronic hypersensitivity pneumonitis, multiple sclerosis, primary biliary cirrhosis, uveitis (anterior and posterior), keratoconjunctivitis sicca and vernal keratoconjunctivitis, interstitial lung fibrosis, psoriatic arthritis and glomerulonephritis (with and without nephrotic syndrome, e.g. including idiopathic nephrotic syndrome or minimal change nephropathy).

Further, agents of the invention may also be used for the treatment of cystic fibrosis, pulmonary hypertension, pulmonary fibrosis, inflammatory bowel syndrome, wound healing, diabetic nephropathy as described in WO 05/107463, reduction of inflammation in transplanted tissue as described in US 2005/182018, inflammatory diseases caused by pathogenic organisms as described in WO 03/086408, and cardiovascular conditions as described in WO 03/029264.

Also, the agents of the invention may be used to assess the severity of coronary artery stenosis as described in WO 00/078774 and useful in conjunction with radioactive imaging agents to image coronary activity and useful in adjunctive therapy with angioplasty as described in WO 00/78779.

Agents of the invention are also useful in combination with a protease inhibitor for prevention of organ ischaemia and reperfusion injury as described in WO 05/003150, and in combination with an integrin antagonist for treating platelet aggregation as described in WO 03/090733.

Agents of the invention are also useful in promoting wound healing in bronchial epithelial cells as described in *AJP-Lung* 290: 849-855.

Other diseases or conditions which may be treated with agents of the invention include diabetes, e.g. diabetes mellitus type I (juvenile diabetes) and diabetes mellitus type II, diarrheal diseases, ischemia/reperfusion injuries, retinopathy, such as diabetic retinopathy or hyperbaric oxygen-induced retinopathy, conditions characterised by elevated intraocular pressure or secretion of ocular aqueous humor, such as glaucoma, ischemic tissue/organ damage from reperfusion, bedsores, as agents for promoting sleep, as agents for treating demyelinating diseases, eg multiple sclerosis and as neuroprotective agents for eg, cerebral haemorrhagic injury and spinal cord ischaemi-reperfusion injury.


The effectiveness of an agent of the invention in inhibiting inflammatory conditions, for example in inflammatory airways diseases, may be demonstrated in an animal model, e.g. a mouse or rat model, of airways inflammation or other inflammatory conditions, for example as described by Szarka et al, *J. Immunol. Methods* (1997) 202:49-57; Renzi et al, *Am. Rev. Respir. Dis.* (1993) 148:932-939; Tsuyuki et al., *J. Clin. Invest.* (1995) 96:2924-2931; Cernadas et al (1999) *Am. J. Respir. Cell Mol. Biol.* 20:1-8; and Fozard et al (2002) *European Journal of Pharmacological* 438, 183-188.

The agents of the invention are also useful as co-therapeutic agents for use in combination with other drug substances such as anti-inflammatory, bronchodilatory, antihistamine or anti-tussive drug substances, particularly in the treatment of obstructive or inflammatory airways diseases such as those mentioned hereinbefore, for example as potentiators of therapeutic activity of such drugs or as a means of reducing required dosaging or potential side effects of such drugs. An agent of the invention may be mixed with the other drug substance in a fixed pharmaceutical composition or it may be administered separately, before, simultaneously with or after the other drug substance.

Accordingly the invention includes a combination of an agent of the invention as hereinbefore described with an anti-inflammatory, bronchodilatory, antihistamine or anti-tussive drug substance, said agent of the invention and said drug substance being in the same or different pharmaceutical composition.

Suitable anti-inflammatory drugs include steroids, in particular glucocorticosteroids such as budesonide, beclamethasone dipropionate, fluticasone propionate, ciclesonide or mometasone furoate, or steroids described in WO 02/88167, WO 02/12266, WO 02/100879, WO 02/00679 (especially those of Examples 3, 11, 14, 17, 19, 26, 34, 37, 39, 51, 60, 67, 72, 73, 90, 99 and 101), WO 03/35668, WO 03/48181, WO 03/62259, WO 03/64445, WO 03/72592, WO 04/39827 and WO 04/66920; non-steroidal glucocorticoid receptor agonists, such as those described in DE 10261874, WO 00/00531, WO 02/10143, WO 03/82280, WO 03/82787, WO 03/86294, WO 03/104195, WO 03/101932, WO 04/05229, WO 04/18429, WO 04/19935 and WO 04/26248; LTB₄ antagonists such as BIIL 284, CP-195543, DPC11870, LTB₄ ethanolamide, LY 293111, LY 255283, CGS025019C, CP-195543, ONO-4057, SB 209247, SC-53228 and those described in US 5451700; LTD₄ antagonists such include montelukast, pranlukast, zafirlukast, accolate, SR2640, Wy-48,252, ICI 198615, MK-571, LY-171883, Ro 24-5913 and L-648051; PDE4 inhibitors such cilomilast (Ariflo® GlaxoSmithKline), Roflumilast (Byk Gulden), V-11294A (Napp), BAY19-8004 (Bayer), SCH-351591 (Schering-Plough), Arofylline (Almirall Prodesfarma), PD189659 / PD168787 (Parke-Davis), AWD-12-281 (Asta Medica), CDC-801 (Celgene), SelCID(TM) CC-10004 (Celgene), VM554/UM565

(Vernalis), T-440 (Tanabe), KW-4490 (Kyowa Hakko Kogyo), and those disclosed in WO 92/19594, WO 93/19749, WO 93/19750, WO 93/19751, WO 98/18796, WO 99/16766, WO 01/13953, WO 03/104204, WO 03/104205, WO 03/39544, WO 04/000814, WO 04/000839, WO 04/005258, WO 04/018450, WO 04/018451, WO 04/018457, WO 04/018465, WO 04/018431, WO 04/018449, WO 04/018450, WO 04/018451, WO 04/018457, WO 04/018465, WO 04/019944, WO 04/019945, WO 04/045607 and WO 04/037805; adenosine A_{2B} receptor antagonists such as those described in WO 02/42298; and beta-2 adrenoceptor agonists such as albuterol (salbutamol), metaproterenol, terbutaline, salmeterol fenoterol, procaterol, and especially, formoterol, carmoterol and pharmaceutically acceptable salts thereof, and compounds (in free or salt or solvate form) of formula I of WO 0075114, which document is incorporated herein by reference, preferably compounds of the Examples thereof, especially a compound of formula



and pharmaceutically acceptable salts thereof, as well as compounds (in free or salt or solvate form) of formula I of WO 04/16601, and also compounds of EP 1440966, JP 05025045, WO 93/18007, WO 99/64035, US 2002/0055651, US 2005/0133417, US 2005/5159448, WO 01/42193, WO 01/83462, WO 02/66422, WO 02/ 70490, WO 02/76933, WO 03/24439, WO 03/42160, WO 03/42164, WO 03/72539, WO 03/91204, WO 03/99764, WO 04/16578, WO 04/22547, WO 04/32921, WO 04/33412, WO 04/37768, WO 04/37773, WO 04/37807, WO 04/39762, WO 04/39766, WO 04/45618 WO 04/46083 , WO 04/80964, EP1460064, WO 04/087142, WO 04/089892, EP 01477167, US 2004/0242622, US 2004/0229904, WO 04/108675, WO 04/108676, WO 05/033121, WO 05/040103, WO 05/044787, WO 05/058867, WO 05/065650, WO 05/066140 and WO 05/07908.

Suitable bronchodilatory drugs include anticholinergic or antimuscarinic agents, in particular ipratropium bromide, oxitropium bromide, tiotropium salts and CHF 4226 (Chiesi), and glycopyrrolate, but also those described in EP 424021, US 3714357, US 5171744, US 2005/171147, US 2005/182091, WO 01/04118, WO 02/00652, WO 02/51841, WO 02/53564, WO 03/00840, WO 03/33495, WO 03/53966, WO 03/87094, WO 04/018422, WO 04/05285 and WO 05/077361.

Suitable dual anti-inflammatory and bronchodilatory drugs include dual beta-2 adrenoceptor agonist / muscarinic antagonists such as those disclosed in US 2004/0167167, US 2004/0242622, US 2005/182092, WO 04/74246 and WO 04/74812.

Suitable antihistamine drug substances include cetirizine hydrochloride, acetaminophen, clemastine fumarate, promethazine, loratidine, desloratidine, diphenhydramine and fexofenadine hydrochloride, activastine, astemizole, azelastine, ebastine, epinastine, mizolastine and tefenadine as well as those disclosed in JP 2004107299, WO 03/099807 and WO 04/026841.

Other useful combinations of agents of the invention with anti-inflammatory drugs are those with antagonists of chemokine receptors, e.g. CCR-1, CCR-2, CCR-3, CCR-4, CCR-5, CCR-6, CCR-7, CCR-8, CCR-9 and CCR10, CXCR1, CXCR2, CXCR3, CXCR4, CXCR5, particularly CCR-5 antagonists such as Schering-Plough antagonists SC-351125, SCH-55700 and SCH-D, Takeda antagonists such as N-[[4-[[[6,7-dihydro-2-(4-methylphenyl)-5H-benzo-cyclohepten-8-yl]carbonyl]amino]phenyl]-methyl]tetrahydro-N,N-dimethyl-2H-pyran-4-amin-ium chloride (TAK-770), and CCR-5 antagonists described in US 6166037 (particularly claims 18 and 19), WO 00/66558 (particularly claim 8), WO 00/66559 (particularly claim 9), WO 04/018425 and WO 04/026873.

In accordance with the foregoing, the invention also provides a method for the treatment of a condition responsive to activation of the adenosine A_{2A} receptor, for example an

inflammatory or allergic condition, particularly an inflammatory or obstructive airways disease, which comprises administering to a subject, particularly a human subject, in need thereof a compound of formula (I) or formula (II) in free form or in the form of a pharmaceutically acceptable salt. In another aspect the invention provides a compound of formula (I) or formula (II), in free form or in the form of a pharmaceutically acceptable salt, for use in the manufacture of a medicament for the treatment of a condition responsive to activation of the adenosine A_{2A} receptor, particularly an inflammatory or obstructive airways disease.

The agents of the invention may be administered by any appropriate route, e.g. orally, for example in the form of a tablet or capsule; parenterally, for example intravenously; by inhalation, for example in the treatment of inflammatory or obstructive airways disease; intranasally, for example in the treatment of allergic rhinitis; topically to the skin, for example in the treatment of atopic dermatitis; or rectally, for example in the treatment of inflammatory bowel disease.

In a further aspect, the invention also provides a pharmaceutical composition comprising a compounds of formula (I) and formula (II) in free form or in the form of a pharmaceutically acceptable salt, optionally together with a pharmaceutically acceptable diluent or carrier therefor. The composition may contain a co-therapeutic agent such as an anti-inflammatory, broncho-dilatory, antihistamine or anti-tussive drug as hereinbefore described. Such compositions may be prepared using conventional diluents or excipients and techniques known in the galenic art. Thus oral dosage forms may include tablets and capsules. Formulations for topical administration may take the form of creams, ointments, gels or transdermal delivery systems, e.g. patches. Compositions for inhalation may comprise aerosol or other atomizable formulations or dry powder formulations.

When the composition comprises an aerosol formulation, it preferably contains, for example, a hydro-fluoro-alkane (HFA) propellant such as HFA134a or HFA227 or a mixture of these, and may contain one or more co-solvents known in the art such as ethanol (up to 20% by weight), and/or one or more surfactants such as oleic acid or

sorbitan trioleate, and/or one or more bulking agents such as lactose. When the composition comprises a dry powder formulation, it preferably contains, for example, the compounds of formula (I) or formula (II) having a particle diameter up to 10 microns, optionally together with a diluent or carrier, such as lactose, of the desired particle size distribution and a compound that helps to protect against product performance deterioration due to moisture e.g. magnesium stearate. When the composition comprises a nebulised formulation, it preferably contains, for example, the compound of formula (I) or formula (II) either dissolved, or suspended, in a vehicle containing water, a co-solvent such as ethanol or propylene glycol and a stabiliser, which may be a surfactant.

The invention includes (A) a compounds of formula (I) or formula (II) in inhalable form, e.g. in an aerosol or other atomisable composition or in inhalable particulate, e.g. micronised, form, (B) an inhalable medicament comprising a compounds of formula (I) or formula (II) in inhalable form; (C) a pharmaceutical product comprising a compounds of formula (I) or formula (II) in inhalable form in association with an inhalation device; and (D) an inhalation device containing a compounds of formula (I) or formula (II) in inhalable form.


Dosages of compounds of formula (I) or formula (II) employed in practising the present invention will of course vary depending, for example, on the particular condition to be treated, the effect desired and the mode of administration. In general, suitable daily dosages for administration by inhalation are of the order of 0.005 to 10 mg, while for oral administration suitable daily doses are of the order of 0.05 to 100 mg.

The invention is illustrated by the following Examples.

EXAMPLES

Preferred compounds of formula I
















35







































are shown in Table 1 below.



















TABLE 1

















Ex.	R ¹	R ²	R ³
1			
2			
3			
4			



















5			
6			
7			
8			
9			




















10			
11			
12			
13			
14			
15			
















16			
17			
18			
19			
20			
21			



















22			
23			
24			
25			
26			
27			






















28			
29	H-		
30			
31			
32			
33			






















34			
35			
36			
37			
38			
39			






















40			
41			
42			
43			
44		-H	
45			
46			



















47			
48			
49			
50			
51			
















52			
53			
54			
55			
56			
57			

58			
59			
60			
61			
62			
63			
64			
















65			
66			
67			
68			
69			
70			
71			



















72			
73			
74			
75			
76			
77			
78			



















79			
80			
81			
82			
83			
84			



















85			
86			
87			
88			
89			

50






















90	 <chem>CCC(=O)C</chem>	 <chem>CC(c1ccccc1)(c2ccccc2)CC</chem>	 <chem>CN1CCC[C@H]1NC(=O)c2ccncc2</chem>
91	 <chem>CCC(=O)C</chem>	 <chem>CC(C)CC(C)C</chem>	 <chem>CN1CCC[C@H]1NC(=O)c2ccncc2</chem>
92	 <chem>CCC(=O)C</chem>	 <chem>CC(C)(C)C(C)(C)CC</chem>	 <chem>CN1CCC[C@H]1NC(=O)c2ccncc2</chem>
93	 <chem>CCC(=O)C</chem>	 <chem>CC(C)CC(C)C</chem>	 <chem>CN1CCC[C@H]1NC(=O)c2cc3ncoc3c2C</chem>
94	 <chem>C1CCC1C(=O)N</chem>	 <chem>CC(c1ccccc1)(c2ccccc2)CC</chem>	 <chem>CN1CCC[C@H]1NC(=O)Cc2ccccc2</chem>






















95			
96			
97			
98			
99			
100			


















101			
102			
103			
104			
105			
106			






















107			
108			
109			
110			
111			
112			















113			
114			
115			
116			
117			
118			

119			
120			
121			
122			
123			
124			
125			













126			
127			
128			
129			
130			
131			
132			

133			
134			
135			
136			
137			
138			













139			
140			
141			
142			
143			
144			
145			



















146			
147			
148			
149			
150			



















60















151			
152			
153			
154			













61

155			
156			
157			
158			













159			
160			
161			
162			
163			
164			













165			
166			
167			
168		-H	
169			-Cl
170		-H	
171			













172			
173			
174		-H	
175			
176			













177			
178			
179			
180			













Ex.	R ¹	R ²	R ³
181			
182			
183			
184			













185	 <chem>CC(=O)CCO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)CC</chem>	 <chem>CN1CCCC1NC(=O)NC2=CC=C(O)C=C2</chem>
186	 <chem>CC(=O)CCO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)CC</chem>	 <chem>CN1CCCC1NC(=O)NC2=CC=C(S(=O)(=O)N)C=C2</chem>
187	 <chem>CC(=O)CCO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)CC</chem>	 <chem>CN1CCCC1NC(=O)NC2=CC=C(S(=O)(=O)N)C=C2</chem>
188	 <chem>CC(=O)CCO</chem>	 <chem>CC(C1=CC=C(O)C=C1)(C2=CC=C(O)C=C2)CC</chem>	 <chem>CN1CCCC1NC(=O)NC2=CC=C(S(=O)(=O)N)C=C2</chem>













189			
190			
191			
192			

193	 <chem>CC(=O)CO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)O</chem>	 <chem>CN1CCCC1C(=O)NCc2ccc(O)cc2</chem>
194	 <chem>CC(=O)CO</chem>	 <chem>CC(O)Cc1ccc(O)cc1</chem>	 <chem>CN1CCCC1C(=O)NCc2ccncc2</chem>
195	 <chem>CC(=O)CO</chem>	 <chem>CC(O)Cc1ccc(O)cc1</chem>	 <chem>CN1CCCC1C(=O)NCc2ccncc2</chem>
196	 <chem>CC(=O)CO</chem>	 <chem>CC(O)Cc1ccc(O)cc1</chem>	 <chem>CN1CCCC1C(=O)NCc2ccc(O)cc2</chem>













197	 <chem>CC(=O)CO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)CC</chem>	 <chem>Cn1cc(C(=O)Nc2cccnc2)cn1</chem>
198	 <chem>CC(=O)CO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)CC</chem>	 <chem>Cn1cc(C(=O)NCCc2ccncc2)cn1</chem>
199	 <chem>CC(=O)CO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)CC</chem>	 <chem>Cn1cc(C(=O)NCCc2ccc(O)cc2)cn1</chem>
200	 <chem>CC(=O)CO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)CC</chem>	 <chem>Cn1cc(C(=O)Nc2ccc(S(=O)(=O)N)cc2)cn1</chem>

201	 <chem>CC(=O)CO</chem>	 <chem>CC(c1ccccc1)c2ccccc2</chem>	 <chem>Cn1cc(C(=O)Nc2ccc(S(=O)(=O)N)cc2)nn1</chem>
202	 <chem>CC(=O)CO</chem>	 <chem>CC(c1ccc(O)cc1)c2ccc(O)cc2</chem>	 <chem>Cn1cc(C(=O)Nc2ccncc2)nn1</chem>
203	 <chem>CC(=O)CO</chem>	 <chem>CC(c1ccc(O)cc1)c2ccc(O)cc2</chem>	 <chem>Cn1cc(C(=O)NCCc2ccncc2)nn1</chem>
204	 <chem>CC(=O)CO</chem>	 <chem>CC(c1ccc(O)cc1)c2ccc(O)cc2</chem>	 <chem>Cn1cc(C(=O)NCCc2ccc(O)cc2)nn1</chem>













205			
206			
207			
208			













209			
210			
211			
212			













213	 <chem>CC(=O)O</chem>	 <chem>CC(c1ccccc1)(c2ccccc2)c3ccccc3</chem>	 <chem>CC1=CN=C(C(=O)Nc2ccc(S(=O)(=O)N)cc2)N1</chem>
214	 <chem>CC(=O)O</chem>	 <chem>CC(c1ccccc1)(c2ccccc2)c3ccccc3</chem>	 <chem>CC1=CN=C(C(=O)Nc2ccc(S(=O)(=O)N)cc2)N1</chem>
215	 <chem>CC(=O)O</chem>	 <chem>CC(C)(O)Cc1ccccc1</chem>	 <chem>CC1=CN=C(C(=O)Nc2ccc(O)cc2)N1</chem>
216	 <chem>CC(=O)O</chem>	 <chem>CC(C)(O)Cc1ccccc1</chem>	 <chem>CC1=CN=C(C(=O)Nc2ccc(S(=O)(=O)N)cc2)N1</chem>
















217			
218			
219			
220			
















221	 <chem>CC(=O)O</chem>	 <chem>CC(C1=CC=C(O)C=C1)-C2=CC=C(O)C=C2</chem>	 <chem>Cn1cc(C(=O)Nc2cccnc2)nn1</chem>
222	 <chem>CC(=O)O</chem>	 <chem>CC(C1=CC=C(O)C=C1)-C2=CC=C(O)C=C2</chem>	 <chem>Cn1cc(C(=O)NCC2=CC=CC=N2)nn1</chem>
223	 <chem>CC(=O)O</chem>	 <chem>CC(C1=CC=C(O)C=C1)-C2=CC=C(O)C=C2</chem>	 <chem>Cn1cc(C(=O)NCC2=CC=C(O)C=C2)nn1</chem>
224	 <chem>CC(=O)O</chem>	 <chem>CC(C1=CC=C(O)C=C1)-C2=CC=C(O)C=C2</chem>	 <chem>Cn1cc(C(=O)Nc2ccc(S(=O)(=O)N)cc2)nn1</chem>
















225			
226			
227			
228			
















229			
230			
231			
232			



















233			
234			
235			
236			







Example	R1	R2	R3
237			
238			
239			
240			
241			

242			
243			
244			
245			
246			

247	 <chem>CC(=O)O[C@H](C)C(=O)OC</chem>	 <chem>CCC(c1ccccc1)c2ccccc2</chem>	 <chem>CN1CC[C@H](NC(=O)Nc2ccncc2)C1</chem>
248	 <chem>CC(=O)CC(=O)O</chem>	 <chem>CCC(c1ccccc1)c2ccccc2</chem>	 <chem>CN1CC[C@H](NC(=O)Nc2ccncc2)C1</chem>
249	 <chem>CC(=O)OC(C)(C)C(=O)OC</chem>	 <chem>CCC(c1ccccc1)c2ccccc2</chem>	 <chem>CN1CC[C@H](NC(=O)Nc2ccncc2)C1</chem>
250	 <chem>CC(=O)OC[C@H](C)C(=O)OC</chem>	 <chem>CCC(c1ccccc1)c2ccccc2</chem>	 <chem>CN1CC[C@H](NC(=O)Nc2ccncc2)C1</chem>
251	 <chem>CC(=O)OC(=O)CC</chem>	 <chem>CCC(c1ccccc1)c2ccccc2</chem>	 <chem>CN1CC[C@H](NC(=O)Nc2ccncc2)C1</chem>

252			
253			
254			
255			
256			

257			
258			
259			
260			
261			
262			

263			
264			

Preparation of intermediate compounds

Abbreviations used are as follows: CDI is 1,1'-carbonyldiimidazole, DCM is dichloromethane, DIPEA is diisopropylethylamine, DMAP is 4-dimethylaminopyridine, DMF is dimethyl-formamide, DMSO is dimethylsulfoxide, LCMS is liquid chromatographic mass spectroscopy, TEA is triethylamine, TFA is trifluoroacetic acid, THF is tetrahydrofuran, EtOH is ethanol, IPA is *iso*-propylalcohol and TLC is thin-layer chromatography.

Intermediate A

1-(R)-Pyrrolidin-3-yl-3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-urea hydrochloride:

A1: Imidazole-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide:

A suspension comprising CDI (2.29 g, 14 mmol) and triethylamine (3.8 ml, 27 mmol) in dry DCM (20 ml) is treated portionwise over 5 minutes with 3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-ylamine dihydrochloride (prepared using the procedure described in

international patent application WO 01/94368) (2.88 g, 13 mmol). The reaction mixture is stirred at room temperature for 4.5 hours to yield the title compound as a 0.43 M solution in DCM.

A2: (R)-3-[3-(3,4,5,6-Tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-pyrrolidine-1-carboxylic acid tert-butyl ester trifluoroacetate

To a solution of imidazole -1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide (18 ml of a 10 mg/ml solution in DCM)(A1) is added (R)-3-amino-1-N-Boc-pyrrolidine (136 mg, 0.74 mmol) in iso-propanol (3 ml). The reaction mixture is stirred at room temperature overnight and then diluted with DCM (25 ml). This mixture is washed with 0.1 M HCl, water, brine, dried (MgSO₄) and concentrated *in vacuo*. Purification of the crude residue by C-18 reverse phase column chromatography eluting with acetonitrile : water : TFA (0.1%) (gradient of 0 to 100% acetonitrile) yields the title compound.

A3: 1-(R)-Pyrrolidin-3-yl-3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-urea hydrochloride:

A solution of (R)-3-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-pyrrolidine-1-carboxylic acid tert-butyl ester trifluoroacetate (0.2 g, 0.51 mmol) in 1.25 M HCl in MeOH (10 ml) is stirred at room temperature over night. The solvent was removed in *vacuo* to yield the title compound.

Intermediate B

1,3-Di(R)-pyrrolidin-3-yl-urea

B1: 1,3-Bis-((R)-1-benzyl-pyrrolidin-3-yl)-urea

A solution comprising (R)-1-benzyl-pyrrolidin-3-ylamine (5.0 g, 28.4 mmol) in DCM (10 ml) is treated with CDI (2.3 g, 14.2 mmol) and the reaction mixture is stirred at room temperature for 48 hours. The solvent is removed *in vacuo* and the resulting residue is dissolved in ethyl acetate. This portion is washed with water followed by brine, dried (MgSO₄) and concentrated *in vacuo* to yield the title compound as pale orange solid.

B2: 1,3-Di(R)-pyrrolidin-3-yl-urea

To a solution of 1,3-bis-((R)-1-benzyl-pyrrolidin-3-yl)-urea (5.34 g, 14.1 mmol) in ethanol (80 ml) under an inert atmosphere of Argon is added palladium hydroxide on carbon (1.07 g). The reaction mixture is purged with Argon and placed under an atmosphere of hydrogen for two days after which time, the mixture is filtered and the catalyst washed with ethanol. The organic portions are combined and concentrated *in vacuo* to yield the title compound as a white solid.

Intermediate CImidazole-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide

A stirred solution of CDI (1.1 g, 6.77 mmol) in DCM (100 ml) is treated with 3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-ylamine (WO 9965895 EP 21973) (1 g, 5.64 mmol in 50 ml of DCM) added dropwise over 30 minutes. The reaction mixture is stirred at room temperature for 15 minutes to yield the title compound as a 10 mg/ml solution in DCM. The compound is used in solution in subsequent reactions. This solution consists of the imidazole-urea (Intermediate C) together with variable amounts of the corresponding isocyanate and imidazole. This solution is used in the subsequent steps since the imidazole-urea intermediate and isocyanate intermediate are equally suitable as precursors to ureas.

Intermediate D**(2S,3S,4R,5R)-5-[2-Amino-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide**

Step D1: (3aS,4S,6R,6aR)-6-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxole-4-carboxylic acid ethylamide

The title compound is prepared by the procedure of Preparation of aminopurine-b-D-ribofuranuronamide derivatives as antiinflammatories. Di Ayres, Barry Edward; Gregson, Michael; Ewan, George Blanch; Keeling, Suzanne Elaine; Bell, Richard. (Glaxo Group Limited, UK). PCT Int. Appl. (1996), 49 pp. WO 9602553.

Step D2: (2S,3S,4R,5R)-5-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

A solution of (3aS,4S,6R,6aR)-6-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,2-dimethyl-tetrahydro-furo[3,4-d][1,3]dioxole-4-carboxylic acid ethylamide (Step D1) in TFA/water (2:1) is stirred at RT overnight. The reaction mixture is concentrated *in vacuo* to afford the titled compound.

Intermediate E

(2S,3S,4R,5R)-5-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide trifluoroacetate

Step E1: (2S,3S,4R,5R)-5-[6-(2,2-Diphenyl-ethylamino)-2-((R)-3-BOC-amino-pyrrolidin-1-yl)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide (2S,3S,4R,5R)-5-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide (Intermediate D) (1g, 1.91mmol), (3R)-3-(BOC-amino)pyrrolidine (1.068g, 5.74mmol) and sodium iodide (287mg, 1.91mmol) is dissolved in acetonitrile (10ml) and NMP (0.5ml). The reaction mixture is heated using microwave radiation at 160°C for 30 minutes in the Personal Chemistry EmrysTM Optimizer microwave reactor. The reaction mixture is concentrated *in vacuo* and purified by C-18 reverse phase column chromatography eluting with acetonitrile:water (0.1% TFA) (gradient 0-100% acetonitrile) to afford the titled compound.

Step E2: (2S,3S,4R,5R)-5-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide trifluoroacetate

(2S,3S,4R,5R)-5-[6-(2,2-Diphenyl-ethylamino)-2-((R)-3-BOC-amino-pyrrolidin-1-yl)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide (step E1) is dissolved in DCM and TFA and stirred at RT overnight. The reaction mixture is concentrated *in vacuo* to afford the titled compound.

Intermediate F

[4-(2-Amino-ethyl)-imidazo[1-yl]-acetic acid cyclohexyl ester:

F1: [4-(2-Amino-ethyl)-imidazo[1-yl]-acetic acid methyl ester sulphate:

A solution comprising [4-(2-amino-ethyl)-imidazo[1-yl]-acetic acid (prepared according to the procedure of Jain, Rahul; Cohen, Louis A. Regiospecific alkylation of histidine and histamine at N-1. Tetrahedron (1996), 52(15), 5363-70) (7.6 g, 44.8 mmol) in methanol (100 ml) is treated with concentrated sulphuric acid (3 drops) and heated to reflux for 18 hours. Molecular sieves are added to the reaction mixture which is refluxed for a further 3 days. The mixture is filtered and concentrated *in vacuo*. The solid is taken up in water and basified to pH10 using sodium hydroxide. The solution is extracted with DCM using a continuous liquid-liquid extraction system to yield the title product.

F2: [4-(2-Amino-ethyl)-imidazo[1-yl]-acetic acid cyclohexyl ester sulphate:

[4-(2-Amino-ethyl)-imidazo[1-yl]-acetic acid methyl ester sulphate (1 g, 3.6 mmol) is suspended in cyclohexanol (50 ml) and treated with concentrated sulphuric acid (5 drops). The reaction mixture is heated to 110 °C for 4 hours and concentrated *in vacuo*. The residue is dissolved in saturated sodium bicarbonate solution and the mixture is concentrated *in vacuo*. The resulting solid is triturated with methanol, filtered and the filtrate is reduced *in vacuo*. The crude solid is dissolved in water, washed with DCM and the aqueous portion is concentrated *in vacuo* to yield the title product.

Intermediate G

[(1S,2R,3S,4R)-4-(2,6-Dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionyl-carbamic acid tert-butyl ester:

G1: (1S,4R)-4-(2,6-Dichloro-purin-9-yl)-cyclopent-2-enol

2,6-Dichloropurine (10 g, 52.90 mmol), (1S,4R)-cis 4-acetoxy-2-cyclopenten-1-ol (10 g, 70.40 mmol), tris(dibenzylideneacetone)dipalladium(0) (3.20 g, 3.50 mmol) and polymer supported triphenylphosphine (3 mmol/g, 11.60 g, 35.00 mmol) are placed in an oven-dried flask under an atmosphere of argon. Dry deoxygenated THF (80 ml) is added and the reaction mixture is stirred gently for 5 minutes. Triethylamine (20 ml) is added and the reaction mixture is stirred at 50°C. The reaction is shown to be complete by LCMS after 1 hour. The reaction mixture is allowed to cool, filtered and the solvent is removed *in vacuo*. The title compound is obtained after purification by flash column chromatography (silica, dichloromethane / methanol 25:1). ¹H nmr (CDCl₃, 400 MHz);

8.30(s, 1H), 6.40(m, 1H), 5.90(m, 1H), 5.50(m, 1H), 4.95(m, 1H), 3.05(m, 1H), 2.10(m, 1H), MS (ES+) m/e 271 (MH^+).

G2: Carbonic acid (1S,4R)-4-(2,6-dichloro-purin-9-yl)-cyclopent-2-enyl ester ethyl ester
(1S,4R)-4-(2,6-Dichloro-purin-9-yl)-cyclopent-2-enol (9.5 g, 35.05 mmol) is placed in an oven-dried flask under an atmosphere of argon. Dry THF (200mL) is added followed by dry pyridine (5.54 g, 70.1 mmol). Ethyl chloroformate (15.21 g, 140.2 mmol) is added slowly so that the temperature does not rise above 40°C and the reaction mixture is stirred at room temperature. The reaction is shown to be complete by LCMS after 1 hour. The solvent is removed *in vacuo* and the residue is partitioned between dichloromethane (200mL) and water (200mL). The organic layer is washed with water (150 ml) and brine (150 ml), dried over $MgSO_4$, filtered and the solvent is removed *in vacuo*. The title compound is obtained after crystallisation from methanol. 1H nmr ($CDCl_3$, 400 MHz); 8.20(s, 1H), 6.45(m, 1H), 6.25(m, 1H), 5.75(m, 1H), 5.70(m, 1H), 4.25(q, 2H), 3.20(m, 1H), 2.05(m, 1H), 1.35(t, 3H), MS (ES+) m/e 343 (MH^+).

G3: [(1S,4R)-4-(2,6-Dichloro-purin-9-yl)-cyclopent-2-enyl]-propionyl-carbamic acid tert-butyl ester

Carbonic acid (1S,4R)-4-(2,6-dichloro-purin-9-yl)-cyclopent-2-enyl ester ethyl ester (1.00 g, 2.92 mmol), propionyl-carbamic acid tert-butyl ester (Intermediate W) (0.55 g, 3.21 mmol) and triphenylphosphine (0.115 g, 0.44 mmol) are placed under an inert atmosphere of Argon. THF (10 ml) is added followed by tris(dibenzylideneacetone)dipalladium(0) (0.13 g, 0.15 mmol). The reaction mixture is stirred at 50° for 1 hour. The solvent is removed *in vacuo* and purification by chromatography on silica eluting with EtOAc/hexane (1:4) affords the title product. 1H nmr ($CDCl_3$, 400 MHz); 8.70(s, 1H), 6.15(m, 1H), 5.85(m, 1H), 5.80(m, 1H), 5.60(m, 1H), 3.15(m, 1H), 2.75(q, 2H), 2.10(m, 1H), 1.55(s, 9H), 1.15(t, 3H), MS (ES+) m/e 426 (MH^+).

G4: [(1S,2R,3S,4R)-4-(2,6-Dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionyl-carbamic acid tert-butyl ester:

[(1S,4R)-4-(2,6-dichloro-purin-9-yl)-cyclopent-2-enyl]-propionyl-carbamic acid tert-butyl ester (11.37 g, 26.7 mmol), methanesulfonamide (2.54 g, 26.7 mmol) and AD-mixa (55 g) are placed in a flask of water (100 ml) and *t*-butanol (100 ml). Osmium tetroxide (4% in water) is added and the reaction mixture is stirred vigorously at room temperature overnight. Sodium sulfite (40 g) is added and the mixture is stirred at room temperature for a further hour and then partitioned between EtOAc and water. The organic portion is separated, dried (MgSO₄) and concentrated *in vacuo*. The crude product is purified by chromatography on silica eluting with DCM:MeOH (25:1 increasing to 10:1) to afford the title compound.

Intermediate H

4-[(Imidazole-1-carbonyl)-amino]-piperidine-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide:

H1: [1-(3,4,5,6-Tetrahydro-2H-[1,2']bipyridinyl-4-ylcarbamoyl)-piperidin-4-yl]-carbamic acid tert-butyl ester:

To a suspension of 4-N-Boc-amino-piperidine (0.396 g, 1.85 mmol) in iso-propanol (5 ml) is added imidazole-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide (50 ml of a 10 mg/ml solution in DCM, 1.85 mmol) and the reaction mixture is stirred at room temperature overnight. The solvent is removed *in vacuo* and recrystallisation of the solid from methanol yields the title product.


H2: 4-Amino-piperidine-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide dihydrochloride

[1-(3,4,5,6-Tetrahydro-2H-[1,2']bipyridinyl-4-ylcarbamoyl)-piperidin-4-yl]-carbamic acid tert-butyl ester (0.45 g, 1.12 mmol) is treated with 4 M HCl (in dioxane)(2.5 ml) and methanol (1 ml, co-solvent) and the reaction mixture is allowed to stir at room temperature for 1 hour. The solvent is removed *in vacuo* and the resulting solid is dried in a vacuum oven to yield the title product.

H3: 4-[(Imidazole-1-carbonyl)-amino]-piperidine-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide:

The title product is prepared analogously to imidazole-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide (Intermediate A1) by replacing 3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-ylamine dihydrochloride with 4-amino-piperidine-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide dihydrochloride (Intermediate H2).

Intermediate I



This compound is prepared analogously to 4-[(imidazole-1-carbonyl)-amino]-piperidine-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide (Intermediate H) by replacing imidazole-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide with 4-[(imidazole-1-carbonyl)-amino]-piperidine-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide.

Intermediate J

N-{(1S,2R,3S,4R)-4-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide

J1: (1S,4R)-4-(2,6-Dichloro-purin-9-yl)-cyclopent-2-enol

2,6-Dichloropurine (10 g, 52.90 mmol), (1S,4R)-cis 4-acetoxy-2-cyclopenten-1-ol (10 g, 70.40 mmol), tris(dibenzylideneacetone)dipalladium(0) (3.20 g, 3.50 mmol) and polymer supported triphenylphosphine (3 mmol/g, 11.60 g, 35.00 mmol) are placed in an oven-dried flask under an atmosphere of argon. Dry deoxygenated THF (80 ml) is added and the reaction mixture is stirred gently for 5 minutes. Triethylamine (20 ml) is added and the reaction mixture is stirred at 50°C. The reaction is shown to be complete by LCMS after 1 hour. The reaction mixture is allowed to cool, filtered and the solvent is removed *in vacuo*. The title compound is obtained after purification by flash column chromatography (silica, dichloromethane / methanol 25:1). ¹H nmr (CDCl₃, 400 MHz); 8.30(s, 1H), 6.40(m, 1H), 5.90(m, 1H), 5.50(m, 1H), 4.95(m, 1H), 3.05(m, 1H), 2.10(m, 1H), MS (ES+) *m/e* 271 (MH⁺).

J2: Carbonic acid (1S,4R)-4-(2,6-dichloro-purin-9-yl)-cyclopent-2-enyl ester ethyl ester
(1S,4R)-4-(2,6-Dichloro-purin-9-yl)-cyclopent-2-enol (9.5 g, 35.05 mmol) is placed in an oven-dried flask under an atmosphere of argon. Dry THF (200mL) is added followed by dry pyridine (5.54 g, 70.1 mmol). Ethyl chloroformate (15.21 g, 140.2 mmol) is added slowly so that the temperature does not rise above 40°C and the reaction mixture is stirred at room temperature. The reaction is shown to be complete by LCMS after 1 hour. The solvent is removed *in vacuo* and the residue is partitioned between dichloromethane (200mL) and water (200mL). The organic layer is washed with water (150 ml) and brine (150 ml), dried over MgSO₄, filtered and the solvent is removed *in vacuo*. The title compound is obtained after crystallisation from methanol. ¹H nmr (CDCl₃, 400 MHz); 8.20(s, 1H), 6.45(m, 1H), 6.25(m, 1H), 5.75(m, 1H), 5.70(m, 1H), 4.25(q, 2H), 3.20(m, 1H), 2.05(m, 1H), 1.35(t, 3H), MS (ES+) *m/e* 343 (MH⁺).

J3: Di-Boc-[(1S,4R)-4-(2,6-dichloro-purin-9-yl)-cyclopent-2-enyl]-amine
Carbonic acid (1S,4R)-4-(2,6-dichloro-purin-9-yl)-cyclopent-2-enyl ester ethyl ester (2.5 g, 7.29 mmol), di-*t*-butyl iminodicarboxylate (1.74 g, 8.02 mmol), tris(dibenzylideneacetone)dipalladium(0) (0.33 g, 0.36 mmol) and triphenylphosphine (0.29 g, 1.09 mmol) are placed in an oven-dried flask under an atmosphere of argon. Dry deoxygenated THF (30ml) is added and the reaction mixture is stirred at room temperature. The reaction is shown to be complete by LCMS after 3 hours. The solvent is removed *in vacuo* and the title compound is obtained after purification by flash column chromatography (silica, ethyl acetate / isohexane 4:1) ¹H nmr (CDCl₃, 400 MHz); 8.70(s, 1H), 6.20(m, 1H), 5.85(m, 1H), 5.80(m, 1H), 5.40(m, 1H), 3.20(m, 1H), 2.15(m, 1H), 1.55(s, 18H), MS (ES+) *m/e* 470 (MH⁺).

J4: (1S,2R,3S,5R)-3-(Di-Boc-amino)-5-(2,6-dichloro-purin-9-yl)-cyclopentane-1,2-diol
The title compound is prepared from di-Boc-[(1S,4R)-4-(2,6-dichloro-purin-9-yl)-cyclopent-2-enyl]-amine using a procedure analogous to that use to prepare (1R,2S,3R,5S)-3-(6-{[bis-(4-methoxy-phenyl)-methyl]-amino}-2-chloro-purin-9-yl)-5-(di-Boc-amino)-cyclopentane-1,2-diol (Intermediate in the preparation of Intermediate

ZA). ^1H nmr (CDCl_3 , 400 MHz); 8.35(s, 1H), 4.80(m, 1H), 4.70(m, 1H), 4.50(m, 1H), 3.85(m, 1H), 3.75(m, 1H), 3.10(m, 1H), 2.75(m, 1H), 2.55(m, 1H), 1.55(s, 18H), MS (ES+) m/e 504 (MH^+).

J5: (1S,2R,3S,5R)-3-Amino-5-(2,6-dichloro-purin-9-yl)-cyclopentane-1,2-diol trifluoroacetate

A solution of (1S,2R,3S,5R)-3-(Di-Boc-amino)-5-(2,6-dichloro-purin-9-yl)-cyclopentane-1,2-diol (0.550 g, 1.09 mmol) in DCM (4 ml) is treated with TFA (2 ml) and stirred at room temperature for 2 hours. The solvent is removed in vacuo and afford the title product which is used in the next step without further purification.

MS (ES+) m/e 304 (MH^+).

J6: N-[(1S,2R,3S,4R)-4-(2,6-Dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide

A solution of 1S,2R,3S,5R)-3-amino-5-(2,6-dichloro-purin-9-yl)-cyclopentane-1,2-diol trifluoroacetate (0.304 g, 1.00 mmol) in THF (10 ml) is treated with DIPEA (0.387 g, 3.00 mmol) followed by propionyl chloride (0.093 g, 1.00 mmol). The reaction mixture is stirred at room temperature for 2 hours. The solvent is removed *in vacuo* and the title compound is obtained after purification by reverse phase column chromatography (IsoluteTM C18, 0-100% acetonitrile in water – 0.1% TFA). MS (ES+) m/e 360 (MH^+).

J7: N-[(1S,2R,3S,4R)-4-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide

N-[(1S,2R,3S,4R)-4-(2,6-Dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide (160 mg, 0.44 mmol) is dissolved in THF (5 ml) under an atmosphere of argon.

Diisopropylamine (69 mg, 0.53 mmol) is added followed by 2,2-diphenylethylamine (96 mg, 0.49 mmol) and the reaction mixture is stirred at 50° C. The reaction is shown to be complete by LCMS after 2 hours. The solvent is removed *in vacuo* and the title compound is obtained after purification by reverse phase column chromatography (IsoluteTM C18, 0-100% acetonitrile in water – 0.1% TFA). ^1H nmr (MeOD, 400 MHz); 8.00(s, 1H), 7.40-7.15(m, 10H), 4.75(m, 1H), 4.60(m, 1H), 4.50(m, 1H), 4.20(m, 3H),

3.95(m, 1H), 2.85(m, 1H), 2.40(q, 2H), 2.10(m, 1H), 1.20 (t, 3H), MS (ES+) m/e 521 (MH^+).

The final compound of Intermediate J may also be prepared using the following process:

JJ1: {2-Chloro-9-[(1R,4S)-4-(di-Boc-amino)-cyclopent-2-enyl]-9H-puri

n-6-yl}-(2,2-diphenyl-ethyl)-amine

(1S,2R,3S,5R)-3-(Di-Boc-amino)-5-(2,6-dichloro-purin-9-yl)-cyclopentane-1,2-diol

(13.0g, 27.66 mmol) is dissolved in THF (250 ml) under an atmosphere of argon.

Diisopropylamine (4.28 g, 33.19 mmol) is added followed by 2,2-diphenylethylamine(6.0 g, 30.43 mmol) and the reaction mixture is stirred at 50°C. The reaction is shown to be complete by LCMS after 18 hours. The solvent is removed *in vacuo* and the reaction mixture is partitioned between dichloromethane (250 ml) and 0.1M HCl (250 ml). The organic layer is washed with water (200 ml) and brine (200 ml), dried over $MgSO_4$, filtered and the solvent is removed *in vacuo* to give the title compound. 1H nmr ($CDCl_3$, 400 MHz); 8.05(s, 1H), 7.30-7.10(m, 10H), 6.00(m, 1H), 5.70(m, 2H), 5.60(m, 1H), 5.20(m, 1H), 4.30(m, 1H), 4.20(m, 1H), 3.65(m, 1H), 3.05(m, 1H), 2.00(m, 1H), 1.70(m, 1H), 1.40(s, 18H), MS (ES+) m/e 631 (MH^+).

JJ2: (1R,2S,3R,5S)-3-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-5-(di-Boc-amino)-cyclopentane-1,2-diol

A solution of {2-Chloro-9-[(1R,4S)-4-(di-Boc-amino)-cyclopent-2-enyl]-9H-purin-6-yl}-(2,2-diphenyl-ethyl)-amine (2.9 g, 4.6 mmol) in THF (60 ml) is treated with 4-methylmorpholine N-oxide (1.1g, 9.3 mmol) and osmium tetroxide (4% solution in water) (6 ml) and the mixture is stirred at room temperature for 48 hours. The solvent is removed under reduced pressure and the residue is purified by column chromatography on silica gel eluting with a gradient system of methanol : dichloromethane (0:100 by volume) gradually changing to methanol : dichloromethane (4:96 by volume) to afford the title compound. LCMS (electrospray): m/z [MH^+] 665.34

JJ3: (1S,2R,3S,5R)-3-Amino-5-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-cyclopentane-1,2-diol trifluoroacetate
(1R,2S,3R,5S)-3-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-5-(di-Boc-amino)-cyclopentane-1,2-diol (10.3 g, 15.50 mmol) is dissolved in dichloromethane (50 ml). TFA (25ml) is added and the reaction mixture is stirred at room temperature. The reaction is shown to be complete by LCMS after 2 hours. The solvent is removed *in vacuo* to give the title compound. ¹H nmr (MeOD, 400 MHz); 7.90(s, 1H), 7.30-7.10(m, 10H), 4.65(m, 1H), 4.50(m, 1H), 4.40(m, 1H), 4.20(m, 1H), 4.10(m, 2H), 3.50(m, 1H), 2.75(m, 1H), 2.15(m, 1H), MS (ES+) *m/e* 465 (MH⁺).

JJ4: N-{(1S,2R,3S,4R)-4-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide
(1S,2R,3S,5R)-3-Amino-5-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-cyclopentane-1,2-diol trifluoroacetate (9.50 g, 16.42 mmol) and diisopropylethylamine (6.36 g, 49.27 mmol) are placed in a flask with dry THF (150 ml). Propionyl chloride (1.52 g, 16.42mmol) is added dropwise and the reaction mixture is stirred at room temperature. The reaction is shown to be complete by LCMS after 1 hour. The solvent is removed *in vacuo* and the residue is partitioned between dichloromethane (250 ml) and water (250 ml). The organic layer is washed with water (200 ml) and brine (200 ml), dried over MgSO₄, filtered and the solvent is removed *in vacuo*. The solid is recrystallised from 1,2-dichloroethane to give the title compound. ¹H nmr (MeOD, 400 MHz); 8.00(s, 1H), 7.40-7.15(m, 10H), 4.75(m, 1H), 4.60(m, 1H), 4.50(m, 1H), 4.20(m, 3H), 3.95(m, 1H), 2.85(m, 1H), 2.40(q, 2H), 2.10(m, 1H), 1.20 (t, 3H), MS (ES+) *m/e* 521 (MH⁺).

Intermediate K

Cyclobutanecarboxylic acid {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -amide:


A solution of (1S,2R,3S,5R)-3-amino-5-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-cyclopentane-1,2-diol hydrochloride (Intermediate JJ3) (100 mg, 0.2 mmol) in dry THF (1 ml) is treated with diisopropylethylamine (0.17 ml, 1mmol) and

cyclobutanecarboxylic acid chloride (0.023 ml, 0.2 mmol) and the mixture is stirred at room temperature for 48 hours. The solvent is removed under reduced pressure. The residue is purified by reverse-phase chromatography eluting with a gradient system of acetonitrile (0.1% TFA) : water (0.1% TFA) (0:100 by volume) gradually changing to acetonitrile (0.1% TFA) : water (0.1% TFA) (100:0 by volume) to afford the title compound (51mg). LCMS (electrospray): m/z $[MH^+]$ 547.26. 1H nmr (MeOD, 400 MHz); 8.00(s, 1H), 7.40-7.25(m, 8H), 7.20-7.15 (m, 2H), 4.70(m, 1H), 4.50(m, 2H), 4.20(m, 2H), 3.95(m, 1H), 2.85(m, 1H), 2.30(m, 2H), 2.20(m, 2H), 2.05(m, 2H), 1.90(m, 1H)

Intermediate L


{(1S,2R,3S,4R)-4-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-carbamic acid benzyl ester

L1: Preparation of intermediate L1




A cooled (0 °C) solution of benzyl carbamate (4.0 g, 27 mmol) in THF (100 ml) under an inert atmosphere of Argon is treated with potassium iodide (3.2 g of a 35 %w/w dispersion in oil, 28 mmol) portionwise over 10 minutes. The reaction mixture is allowed to warm to room temperature over 30 minutes after which time benzyl chloroformate (5.0 g, 29 mmol) is added. After stirring at room temperature for 2 hours, the reaction is quenched with water (20 ml). The THF is removed *in vacuo* and the resulting mixture is partitioned between EtOAc and 2M HCl. The organic portion is separated and washed with brine, dried (MgSO₄) and concentrated *in vacuo*. The resulting oil is purified by chromatography on silica eluting with 1:3 EtOAc/*iso*-hexane to yield a product which is recrystallised from DCM/*iso*-hexane to afford the title product.

L2: Preparation of intermediate L2:




A solution comprising carbonic acid (1S,4R)-4-(2,6-dichloro-purin-9-yl)-cyclopent-2-enyl ester ethyl ester (Intermediate J2) (2.0 g, 5.83 mmol), Intermediate L1 (2.2 g, 7.58 mmol) and triphenyl phosphine (229 mg, 0.9 mmol) in THF (20 ml) is stirred at room temperature for 30 minutes. Tris(dibenzylideneacetone)dipalladium (0) (238 mg, 0.3 mmol) is added and the resulting mixture is stirred at room temperature for 1.5 hours. The solvent is removed *in vacuo* and the crude product is purified by chromatography on silica eluting with MeOH/DCM (gradient of 0 to 1 % MeOH) to yield the title compound.

L3: Preparation of intermediate L3:



This compound is prepared analogously to 2-chloro-9-[(1R,4S)-4-(di-Boc-amino)-cyclopent-2-enyl]-9H-purin-6-yl}-(2,2-diphenyl-ethyl)-amine (Intermediate JJ1) by replacing (1S,2R,3S,5R)-3-(Di-Boc-amino)-5-(2,6-dichloro-purin-9-yl)-cyclopentane-1,2-diol (Intermediate J4) with Intermediate L2.

L4: Preparation of intermediate L4:



This compound is prepared analogously to (1R,2S,3R,5S)-3-[2-chloro-6-(2,2-diphenylethylamino)-purin-9-yl]-5-(di-Boc-amino)-cyclopentane-1,2-diol (Intermediate JJ2) by replacing {2-Chloro-9-[(1R,4S)-4-(di-Boc-amino)-cyclopent-2-enyl]-9H-purin-6-yl}-(2,2-diphenylethyl)-amine with Intermediate L3.

L5: {(R)-1-[9-((1R,2S,3R,4S)-4-Benzyloxycarbonylamino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenylethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester

A suspension of Intermediate L4 (1.03 g, 1.4 mmol) and (3R)-(+)-3-(Boc-amino)pyrrolidine (1.03 g, 5.5 mmol) in acetonitrile (2 ml) is treated with sodium iodide (ca. 2 mg) and then heated using microwave radiation in a Personal Chemistry Emrys™ Optimizer microwave reactor at 160°C. After 1 hour, the solvent is removed *in vacuo* and the crude residue is partitioned between DCM and 0.2 M HCl. The organic layer is separated and the aqueous portion is extracted with DCM. The combined organic extracts are washed with saturated sodium bicarbonate solution, water, brine, dried (MgSO₄) and concentrated *in vacuo* to afford the title compound as a brown oil. MS (ES⁺) *m/e* 745 (MH⁺).

L6: {(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenylethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-carbamic acid benzyl ester:

A solution of {(R)-1-[9-((1R,2S,3R,4S)-4-benzyloxycarbonylamino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenylethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester (Intermediate L5) (1.24 g, 1.7 mmol) in MeOH (3 ml) is treated with 4M HCl in dioxane (5 ml) and stirred at room temperature for 2 hours. The solvent is removed *in*

vacuo and purification is carried out by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% HCl). The fractions are collected and the MeCN is removed *in vacuo*. The remaining aqueous portion is basified with saturated sodium bicarbonate solution and extracted with DCM. The combined organic extracted are dried (MgSO₄) and concentrated *in vacuo* to afford the title product. MS (ES+) *m/e* 649 (MH⁺).

Intermediate M

N-{(3aR,4S,6R,6aS)-6-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,2-dimethyl-tetrahydro-cyclopenta[1,3]dioxol-4-yl}-propionamide:

M1: {(R)-1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid benzyl ester:

A solution of (R)-pyrrolidin-3-yl-carbamic acid benzyl ester hydrochloride (0.88 g, 3.45 mmol) in DCM is free-based using sodium hydrogen carbonate solution to yield (R)-pyrrolidin-3-yl-carbamic acid benzyl ester (0.487 g, 2.22 mmol). This amine is added to N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Intermediate J) (0.5 g, 0.96 mmol) and TEA (0.224 g, 2.22 mmol) and then dissolved in NMP (7 ml). The reaction mixture is heated using microwave radiation in a Personal Chemistry Emrys™ Optimizer microwave reactor at 190 °C for 1 hour. The resulting mixture is purified by chromatography on silica eluting with 5% MeOH in DCM to yield the title compound.

M2: {(R)-1-[9-((3aS,4R,6S,6aR)-2,2-Dimethyl-6-propionylamino-tetrahydro-cyclopenta[1,3]dioxol-4-yl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid benzyl ester:

A solution of {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid benzyl ester (0.63 g, 0.89 mmol) in acetone (10 ml) and 2,2-dimethyloxopropane (5 ml) is treated with toluenesulfonic acid (ca.60 mg) and then stirred at room temperature overnight. The mixture is basified using ammonium hydroxide and the solvent is removed *in vacuo*. The crude product is partitioned between DCM and water and the organic portion is washed

with brine, dried over MgSO_4 , filtered and the solvent is removed *in vacuo* to give the title compound. $[\text{MH}^+ 745]$.

M3: N-{{(3aR,4S,6R,6aS)-6-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,2-dimethyl-tetrahydro-cyclopenta[1,3]dioxol-4-yl}-propionamide:

To a solution of {(R)-1-[9-((3aS,4R,6S,6aR)-2,2-dimethyl-6-propionylamino-tetrahydro-cyclopenta[1,3]dioxol-4-yl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid benzyl ester (0.598 g, 0.79 mmol) in ethanol (7.5 ml) under an inert atmosphere of Argon is added palladium hydroxide on carbon (10 mg). The reaction mixture is purged with Argon and placed under an atmosphere of hydrogen overnight. The mixture is filtered and purified by chromatography on silica eluting with 5 % MeOH in DCM to yield the title compound. $[\text{MH}^+ 611]$.

Intermediate N

(R)-[1,3']Bipyrrolidinyl

N1: (R)-1'-Benzyl-[1,3']bipyrrolidinyl:

An ice-cooled solution solution of 2,5-dimethoxytetrahydrofuran (19.11 ml, 0.147 mol) and 6M sulphuric acid (37.2 ml) in THF (200 ml) is treated dropwise with (R)-(1)-benzyl-3-aminopyrrolidine (10 g, 0.057 mol) 6M sulphuric acid (37.2 ml) in THF (150 ml) and sodium borohydride pellets (8.62 g, 0.227 mol) simultaneously, ensuring the temperature remains below 10°C. The reaction mixture is allowed to warm to room temprature and water (10 ml) is added to aid dissolution of the sodium hydroxide pellets. After stirring at room temperature for 12 days, the mixture is cooled with the use on an ice-bath and water is added (500 ml). The solution is basified by addition of sodium hydroxide pellets ($\text{pH} < 10$) and then filtered under vacuum. The filtrate is extracted with diethyl ether and DCM and the organic portions are combined and concentrated *in vacuo*. The crude residue is sonicated in diethyl ether and filtered under vacuum. The filtrate is reduced *in vacuo* again and the resulting crude is dissolved in MeCN (8 ml) and purified by reverse phase column chromatography (Isolute™ C18, 0-100% MeCN in water – 0.1% TFA) to yield the title product.

N2: (R)-[1,3']Bipyrrolidinyl:

A solution of (R)-1'-benzyl-[1,3']bipyrrolidinyl (0.517 g, 2.24 mmol) in methanol (25 ml) under an atmosphere of Argon is treated with palladium hydroxide on carbon (0.1 g). The reaction mixture is placed under an atmosphere of hydrogen and stirred at room temperature overnight and then filtered through celite™. The filtrate is concentrated *in vacuo* to yield the title product as a dark orange oil.

Intermediate O(R)-N-Pyrrolidin-3-yl-isonicotinamide:O1: (R)-3-[(Pyridine-4-carbonyl)-amino]-pyrrolidine-1-carboxylic acid tert-butyl ester:

A cooled (0°C) stirred solution of (R)-3-amino-pyrrolidine-1-carboxylic acid tert-butyl ester (1.0 g, 5.36 mmol) and TEA (1.5 ml, 11.0 mmol) in THF (10 ml) is treated dropwise over 1 minute with pyridine-4 carbonyl chloride hydrochloride (0.935 g, 5.25 mmol). After 5 minutes, the reaction mixture is allowed to warm to room temperature and stirred overnight. The resulting mixture is diluted with EtOAc and washed twice with saturated sodium bicarbonate solution followed by brine. The organic portion is dried (MgSO₄) and concentrated *in vacuo*. The crude product is purified by recrystallisation from EtOAc/*iso*-hexane to afford the title product. [MH⁺ 292].

O2: (R)-N-Pyrrolidin-3-yl-isonicotinamide:

A solution of (R)-3-[(pyridine-4-carbonyl)-amino]-pyrrolidine-1-carboxylic acid tert-butyl ester (1.38 g, 4.74 mmol) in MeOH (6 ml) is treated with 2M HCl (5 ml) and left to stand at room temperature overnight. The resulting mixture is diluted with MeOH and added to 12 ml of Dowex resin (50Wx2-200). After 30 minutes, the resin is washed with water until neutral and then further washed off with MeOH and 2% ammonia. The solvent is removed *in vacuo* to afford the title compound as a crystalline solid. [MH⁺ 192].

Intermediate P5-Methyl-isoxazole-3-carboxylic acid (R)-pyrrolidin-3-ylamide:

TEA (0.42 ml, 3.0 mmol) is added to a cooled (-10°C) solution of 5-methyl isoxazole-3-carbonyl chloride (0.44 g, 2.95 mmol) in THF (5 ml). To this turbid mixture is added dropwise, (R)-3-amino-1-N-pyrrolidine (0.5 g, 2.68 mmol) in THF (2 ml) and the reaction mixture is allowed to warm to room temperature over 30 minutes. After standing at room temperature overnight, the reaction mixture is diluted with EtOAc (30 ml) and washed with water (2 x 5 ml), brine, dried (MgSO₄) and concentrated *in vacuo*. The resulting oil is dissolved in MeOH (5 ml) is treated dropwise with 6M HCl (1.15 ml). After standing at room temperature for 4 days, the reaction mixture is concentrated *in vacuo* and co-evaporated with MeOH/EtOAc. The crude residue is triturated with EtOAc to afford the title compound. [MH⁺ 196].

Intermediate Q

N-{(1S,2R,3S,4R)-4-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide

Q1: Acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl} - methyl ester

A suspension of (1S,2R,3S,5R)-3-amino-5-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-cyclopentane-1,2-diol dihydrochloride (Intermediate JJ3) (250 mg, 0.46 mmol) in dry THF (10 ml) is treated with TEA (0.188 g, 1.86 mmol) followed by acetoxyacetylchloride (0.064 g, 0.46 mmol) and then stirred at room temperature for 30 minutes. The solvent is removed *in vacuo* and the solvent is partitioned between DCM and 0.1M HCl. The organic portion is separated and washed with brine, dried (MgSO₄) and concentrated *in vacuo* to afford the title product.

Q2: N-{(1S,2R,3S,4R)-4-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide:

To a suspension of acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl} - methyl ester (0.2 g, 0.35 mmol) in MeOH (10 ml) is added potassium carbonate (0.098 g, 0.7 mmol) and the reaction mixture is stirred at room temperature for 1 hour. The solvent is removed *in vacuo* and

the solvent is partitioned between DCM and water. The organic portion is separated, dried (MgSO₄) and concentrated *in vacuo* to afford the title product.

Intermediate R

3-Isocyanato-benzenesulfonamide:

To a vigorously stirred solution of 3-aminobenzenesulphonamide (1 g, 5.8 mmol) in dry dioxane (25 ml) is added trichloromethyl chloroformate (1.72 g, 8.7 mmol) and the reaction mixture is heated to reflux for 3 hours. The solvent is removed *in vacuo* to yield the title product which is used without further purification.

Intermediate S

4-Isocyanato-benzenesulfonamide:

This compound is prepared analogously to Intermediate R by replacing 3-aminobenzenesulphonamide with 4-aminobenzenesulphonamide.

Intermediate T

{(1S,2R,3S,4R)-4-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-carbamic acid methyl ester:

This compound is prepared analogously to cyclobutanecarboxylic acid {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-amide (Intermediate K) by replacing cyclobutanecarboxylic acid chloride with methylchloroformate.

Intermediate UA

(3-Hydroxy-benzyl)-carbamic acid phenyl ester

3-Hydroxybenzylamine (200mg, 1.62mmol) and Sodium hydrogen carbonate (273mg, 3.25mmol) suspended in water/DCM (4ml, 1:1) is treated with phenyl chloroformate (0.204ml, 1.62mmol). After stirring at RT overnight, the reaction mixture is diluted with

more DCM/water and the organic phase is separated. The organic portion is concentrated *in vacuo* to afford the titled compound. (MH+ 244)

Intermediate UB

Pyridin-3-yl-carbamic acid phenyl ester

Phenyl chloroformate (0.733ml, 5.84mmol) is suspended in pyridine/DCM (3ml, 2:1). The solution is stirred at 0°C, and 3-aminopyridine (500mg, 5.31mmol) dissolved in DCM (1ml) is added drop-wise. The reaction mixture is at 0°C for 1 hour. The solvent is removed *in vacuo* and the residue is dissolved in ethyl acetate. This organic portion is washed with 0.1M HCl and then concentrated *in vacuo* to afford the titled compound. (MH+ 215).

Intermediates UC-UE

These compounds namely,

- (3-Sulfamoyl-phenyl)-carbamic acid phenyl ester (Intermediate **UC**)
- (4-Sulfamoyl-phenyl)-carbamic acid phenyl ester (Intermediate **UD**)
- Pyridin-2-ylmethyl-carbamic acid phenyl ester (Intermediate **UE**)

are prepared analogously to Intermediate UB by replacing 3-aminopyridine with the appropriate amine.

Intermediate VA

3-((R)-3-Pyrrolidin-3-ylureido)-benzenesulfonamide

VA1: 3-[3-((R)-1-Benzyl-pyrrolidin-3-yl)-ureido]-benzenesulfonamide

A solution of (R)-N-benzyl-3-aminopyrrolidine (14.9 g, 0.084 mol) in methanol (100 mL) is added to a suspension of (3-sulfamoyl-phenyl)-carbamic acid phenyl ester (25 g, 0.084 mol). The resulting pale orange solution is stirred at mild reflux (DrySyn @ 80° C) for two hours, then allowed to cool to room temperature, before the volatile components are removed under reduced pressure. The orange syrup is purified by flash column chromatography (silica; DCM/methanol 10:1) to give a beige foamed solid .

VA2: 3-((R)-3-Pyrrolidin-3-ylureido)-benzenesulfonamide

A solution of 3-[3-((R)-1-benzyl-pyrrolidin-3-yl)-ureido]-benzenesulfonamide (25 g, 0.067 mol) in ethanol (250 mL) is purged with nitrogen, and palladium hydroxide (2.5 g, 20% w/w) is added. The suspension is purged with hydrogen and stirred under a positive pressure of hydrogen for 24 hours. Filtration through Celite® (filter material) and removal of solvent under reduced pressure gives the product as a colourless waxy solid.

Intermediate VB1-Pyridin-3-yl-3-(R)-pyrrolidin-3-yl-ureaVB1: 1-((R)-1-Benzyl-pyrrolidin-3-yl)-3-pyridin-3-yl-urea

A solution of pyridin-3-yl-carbamic acid phenyl ester (1.6 g) in dry THF (20 ml) is treated with (R)-1-benzyl-pyrrolidin-3-ylamine (1.9 g, 1.05 eq) and then heated using microwave radiation at 110° C for 1000 s. The solvent is removed *in vacuo* and purification of the crude product by chromatography on silica eluting with DCM followed by EtOAc and EtOH affords the title compound as an oil. (MH⁺ 297).

VB2: 1-Pyridin-3-yl-3-(R)-pyrrolidin-3-yl-urea

The titled compound is prepared from 1-((R)-1-benzyl-pyrrolidin-3-yl)-3-pyridin-3-yl-urea analogously to 3-((R)-3-pyrrolidin-3-ylureido)-benzenesulfonamide (Step VA1).

Intermediate VC1-Pyridin-3-yl-3-(R)-pyrrolidin-3-yl-urea

This compound is prepared from (4-sulfamoyl-phenyl)-carbamic acid phenyl ester (Intermediate UD) analogously to Intermediate VA.

Intermediate VD

N-((1S,2R,3S,4R)-4-[6-((S)-1-Benzyl-2-hydroxy-ethylamino)-2-chloro-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide:

This compound is prepared analogously to (N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide trifluoroacetate (Example 7 step 3) by replacing 4,4'-(2-aminoethylidene)bis-phenol (Example 7 step1) with (S)-2-amino-3-phenyl-propan-1-ol.

Intermediate W

Propionyl-carbamic acid tert-butyl ester

The title compound is prepared from propyl-carbamic acid tert-butyl ester using the procedure described by Ken-ichi Takana et al in *Chem. Pharm. Bull.* 1988, 36, 3125. ¹H nmr (CDCl₃, 400 MHz); 7.25(br s, 1H), 2.75(q, 2H), 1.50(s, 9H), 1.15(t, 3H).

Intermediate X

Bis-(4-methoxy-phenyl)-methanone oxime

4,4'-Dimethoxybenzophenone (25 g, 103 mmol) is suspended in ethanol (150 ml) and pyridine (30 ml). Hydroxylamine hydrochloride (21.50 g, 310 mmol) is added and the reaction mixture is refluxed. The reaction is shown to be complete by TLC after 3 hours. The reaction mixture is allowed to cool and the solvent is removed *in vacuo*. The residue is partitioned between ethyl acetate (500 ml) and water (500 ml). The organic layer dried is over MgSO₄, filtered and the solvent removed *in vacuo*. The title compound is obtained following crystallisation from ethylacetate/ cyclohexane. ¹H nmr (CDCl₃, 400 MHz); 7.70(s, 1H), 7.40 (d of d, 4H), 6.95(d, 2H), 6.85(d, 2H), 3.85(s, 3H), 3.80(s, 3H).

Intermediate Y

C,C-Bis-(4-methoxy-phenyl)-methylamine

Bis-(4-methoxy-phenyl)-methanone oxime (20 g, 77.82 mmol) is suspended in ammonia .880 (450 ml) and ethanol (90 ml). Ammonium acetate (3.00 g, 38.91 mmol) is added followed by the portionwise addition of zinc dust (25.29 g, 389.10 mmol). Once the addition is complete the reaction mixture is slowly heated to 50°C. When the

effervescence has ceased the reaction mixture is refluxed. The reaction is shown to be complete by TLC after 4 hours. The reaction mixture is allowed to cool and ethyl acetate is added (250 ml). The reaction mixture is filtered through Celite™ and the phases are separated. The organic layer dried is over MgSO₄, filtered and the solvent removed *in vacuo* to give the title compound. ¹H nmr (CDCl₃, 400 MHz); 7.25 (d, 4H), 6.80 (d, 4H), 5.10(s, 1H), 3.75(s, 6H).

Intermediate Z

Biphenyl-2-yl-carbamic acid 1-[2-((R)-3-amino-pyrrolidin-1-yl)-2-oxo-ethyl]-piperidin-4-yl ester

Z1: {(R)-1-[2-(4-Hydroxy-piperidin-1-yl)-acetyl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester:

To a solution of (R)-pyrrolidin-3-yl-carbamic acid tert-butyl ester (1.5 g, 8.1 mmol) in THF (150 ml) is added TEA (2.3 ml, 16.1 mmol) followed by dropwise addition of chloroacetyl chloride (0.67 ml, 8.5 mmol). The reaction mixture is allowed to stir at room temperature for 2 hours and then treated TEA (2.3 ml, 16.1 mmol) followed by 4-piperidinol (4.07 g, 40.3 mmol). After stirring at 50 °C for 18 hours, the solvent is removed in vacuo and purification of the crude residue by reverse phase column chromatography (Isolute™ C18, 0-100% MeOH in water – 0.1% TFA) to yield the title product. [MH+ 328.19].

Z2: ((R)-1-{2-[4-(Biphenyl-2-ylcarbamoxyloxy)-piperidin-1-yl]-acetyl}-pyrrolidin-3-yl)-carbamic acid tert-butyl ester:

{(R)-1-[2-(4-Hydroxy-piperidin-1-yl)-acetyl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester (Intermediate Z1) (520 mg, 1.6 mmol) and 2-biphenylisocyanate (930 mg, 2.65 mmol) are dissolved in NMP (2 ml) and heated to 70 °C overnight. Purification is carried out by reverse phase column chromatography (Isolute™ C18, 0-100% MeOH in water – 0.1% TFA). The fractions containing the product are concentrated *in vacuo* to remove the acetonitrile and the aqueous is treated with saturated sodium bicarbonate solution. The product is extracted with DCM and the combined organic portions are concentrated in vacuo to afford the title product. [MH+ 523.24].

Z3: Biphenyl-2-yl-carbamic acid 1-[2-((R)-3-amino-pyrrolidin-1-yl)-2-oxo-ethyl]-piperidin-4-yl ester:

A solution of ((R)-1-[2-[4-(biphenyl-2-ylcarbamoxyloxy)-piperidin-1-yl]-acetyl]-pyrrolidin-3-yl)-carbamic acid tert-butyl ester (Intermediate Z2) (1.17 g, 2.24 mmol) in DCM (10 ml) is treated with TFA (5 ml) and stirred at room temperature for 2 hours. The solution is basified by addition of saturated sodium bicarbonate solution and then extracted with DCM. The combined organic portions are washed with water, brine, dried (MgSO₄) and concentrated *in vacuo* to yield the title product. [MH⁺ 423.20].

Intermediate ZA

N-[(1S,2R,3S,4R)-4-(6-Amino-2-chloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate

Bis-(4-methoxy-phenyl)-methyl-(2-chloro-9H-purin-6-yl)-amine

2,6-Dichloropurine (9.50 g, 50.29 mmol) is dissolved in THF (200 ml) under an atmosphere of argon. Diisopropylamine (7.14 g, 55.32 mmol) is added followed by C,C-bis-(4-methoxy-phenyl)-methylamine (see preparation of intermediates) (12.22 g, 50.29 mmol) and the reaction mixture is stirred at 50°C. The reaction is shown to be complete by LCMS after 5 days. The solvent is removed *in vacuo* and replaced with MeOH (250 mL). The resulting precipitate is filtered off and dried to give the title compound. ¹H nmr (d₆-DMSO, 400 MHz); 8.20(br s, 1H), 7.25(d, 4H), 6.90(d, 4H), 3.75(s, 6H), 3.15(m, 1H), MS (ES⁺) *m/e* 396 (MH⁺).

(1S,4R)-4-(6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-2-chloro-purin-9-yl)-cyclopent-2-enol

Bis-(4-methoxy-phenyl)-methyl-(2-chloro-9H-purin-6-yl)-amine (13 g, 32.87 mmol) is placed in an oven-dried flask under an atmosphere of argon. Dry deoxygenated THF (100 ml) and dry DMSO (2 ml) are added and the suspension is cooled on an ice-bath. Sodium hydride 95% (0.79 g, 32.87 mmol) is then slowly added and the solution is stirred at room temperature for 30 minutes. (1S,4R)-cis 4-Acetoxy-2-cyclopenten-1-ol (4.9 g, 34.5 mmol) and triphenylphosphine (1.36 g, 5.17 mmol) are placed in an oven-dried flask

under an atmosphere of argon. Dry deoxygenated THF (50 ml) is added. This solution is added to the anion solution via syringe. Tetrakis(triphenylphosphine)palladium(0) (2 g, 1.73 mmol) is then added and the mixture is stirred at 50°C. The reaction is shown to be complete by LCMS after 2 hours. The reaction mixture is allowed to cool and the solvent is removed *in vacuo*. The residue is taken up in methanol (50 ml) and the resulting precipitate is filtered off and dried to give the title compound. ¹H nmr (CDCl₃, 400 MHz); 9.10(m, 1H), 8.10(m, 1H), 7.30(d, 4H), 6.90(d, 4H), 6.55(d, 1H), 6.20(m, 1H), 5.95(m, 1H), 5.40(m, 1H), 5.30(d, 1H), 4.70(m, 1H), 3.70(s, 6H), 2.90(m, 1H), 1.70(m, 1H), MS (ES+) *m/e* 478 (MH⁺).

Carbonic acid (1S,4R)-4-(6-{[bis-(4-methoxy-phenyl)-methyl]-amino}-2-chloro-purin-9-yl)-cyclopent-2-enyl ester ethyl ester

(1S,4R)-4-(6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-2-chloro-purin-9-yl)-cyclopent-2-enol (8.00 g, 16.75 mmol) is placed in an oven-dried flask under an atmosphere of argon. Dry pyridine (80 ml) is added followed by diisopropylamine (16 ml). A catalytic amount of DMAP is added followed by 3-oxy-benzotriazole-1-carboxylic acid ethyl ester (6.94 g, 33.50 mmol, see preparation of intermediates). The reaction mixture is stirred at room temperature. The reaction is shown to be complete by TLC after 18 hours. The solvent is removed *in vacuo* and the residue is partitioned between ethyl acetate (500 ml) and 2M HCl (200 ml). The organic layer is washed with water (150 ml) and brine (150 ml), dried over MgSO₄, filtered and the solvent is removed *in vacuo*. The title compound is obtained after purification by flash column chromatography (silica, dichloromethane / methanol 50:1). ¹H nmr (CDCl₃, 400 MHz); 7.80(s, 1H), 7.25(d of d, 4H), 6.85(d of d, 4H), 6.65(m, 1H), 6.50(m, 1H), 6.35(m, 1H), 6.15(m, 1H), 5.65(m, 2H), 4.25(q, 2H), 3.80(s, 6H), 3.10(m, 1H), 1.95(m, 1H), 1.35(t, 3H).

[Bis-(4-methoxy-phenyl)-methyl]-{2-chloro-9-[(1R,4S)-4-(di-Boc-amino)-cyclopent-2-enyl]-9H-purin-6-yl}-amine

Carbonic acid (1S,4R)-4-(6-{[bis-(4-methoxy-phenyl)-methyl]-amino}-2-chloro-purin-9-yl)-cyclopent-2-enyl ester ethyl ester (2.00 g, 3.64 mmol), di-*t*-butyl iminodicarboxylate (0.87 g, 4.00 mmol) and triphenylphosphine (0.14 g, 0.55 mmol) are placed in an oven-

dried flask under an atmosphere of argon. Dry deoxygenated THF (20 ml) is added followed by tetrakis(triphenylphosphine)palladium(0) (0.21 g, 0.18 mmol) and the mixture is stirred at room temperature. The reaction is shown to be complete by LCMS after 3 hours. The solvent is removed *in vacuo* and the title compound is obtained after purification by flash column chromatography (silica, iso-hexane / ethyl acetate 4:1). ¹H nmr (CDCl₃, 400 MHz); 8.20(s, 1H), 7.25(d, 4H), 6.85(d, 4H), 6.60(m, 1H), 6.35(m, 1H), 6.10(m, 1H), 5.80(m, 1H), 5.65(m, 1H), 5.35(m, 1H), 3.80(s, 6H), 3.15(m, 1H), 2.10(m, 1H), 1.55(s, 18H).

(1R,2S,3R,5S)-3-(6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-2-chloro-purin-9-yl)-5-(di-Boc-amino)-cyclopentane-1,2-diol

[Bis-(4-methoxy-phenyl)-methyl]-{2-chloro-9-[(1R,4S)-4-(di-Boc-amino)-cyclopent-2-enyl]-9H-purin-6-yl}-amine (0.75 g, 1.11 mmol) is dissolved in THF (15 ml). N-Methylmorpholine N-oxide (0.26 g, 2.22 mmol) is added followed by osmium tetroxide (1.5 ml, 4% in water). The reaction mixture is stirred at room temperature. The reaction is shown to be complete by LCMS after 18 hours. The solvent is removed *in vacuo* and the title compound is obtained after purification by flash column chromatography (silica, dichloromethane / methanol 50:1). ¹H nmr (CDCl₃, 400 MHz); 7.75(s, 1H), 7.25(m, 4H), 6.85(m, 4H), 6.60(m, 2H), 5.70(m, 1H), 4.70(m, 2H), 4.60(m, 1H), 4.45(m, 1H), 3.80(s, 6H), 3.70(m, 1H), 3.40(m, 1H), 3.25(m, 1H), 2.65(m, 1H), 2.50(m, 1H), 1.55(s, 18H).

(1S,2R,3S,5R)-3-Amino-5-(6-amino-2-chloro-purin-9-yl)-cyclopentane-1,2-diol trifluoroacetate

(1R,2S,3R,5S)-3-(6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-2-chloro-purin-9-yl)-5-(di-Boc-amino)-cyclopentane-1,2-diol (600 mg, 0.84 mmol) is dissolved in dichloromethane (4 ml). TFA (2 ml) is added and the reaction mixture is stirred at room temperature. The reaction is shown to be complete by LCMS after 18 hours. The solvent is removed *in vacuo* and the title compound is obtained after purification by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% TFA). ¹H nmr (MeOD, 400 MHz); 8.10(s, 1H), 4.80(m, 1H), 4.60(m, 1H), 4.30(m, 1H), 3.60(m, 1H), 2.85(m, 1H), 2.30(m, 1H). MS (ES+) *m/e* 285 (MH⁺).

N-[(1S,2R,3S,4R)-4-(6-Amino-2-chloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate

(1S,2R,3S,5R)-3-Amino-5-(6-amino-2-chloro-purin-9-yl)-cyclopentane-1,2-diol trifluoroacetate (intermediate for preparing Example 1) (20 mg, 39 μ mol) and diisopropylethylamine (25 mg, 190 μ mol) are placed in a flask with dry THF (1ml). Propionyl chloride (3.6 mg, 39 μ mol) is added and the reaction mixture is stirred at room temperature. The reaction is shown to be complete by LCMS after 3 hours. The solvent is removed *in vacuo* and the title compound is obtained, which can be purified by reverse phase column chromatography (IsoluteTM C18, 0-100% acetonitrile in water – 0.1% TFA). ¹H nmr (MeOD, 400 MHz); 8.10(s, 1H), 4.75(m, 1H), 4.60(m, 1H), 4.20(m, 1H), 4.00(m, 1H), 3.75(m, 1H), 3.25(m, 1H), 2.85(m, 1H), 2.40(q, 2H), 2.10(m, 1H), 1.20(t, 3H), MS (ES+) *m/e* 341 (MH⁺).

Intermediate ZB

Pyridin-3-yl-carbamic acid phenyl ester:

A solution of pyridine (2 ml) in DCM (10 ml) is treated with phenylchloroformate (1.83 g, 11.7 mmol). To this solution is added 3-aminopyridine (1.0 g, 10.6 mmol) in DCM (8 ml) which results in an exotherm of 20 °C. The reaction mixture is stirred at room temperature for 2 hours and then concentrated *in vacuo*. The residue is partitioned between EtOAc and water and the organic portion is separated. This organic portion is washed with water, saturated sodium bicarbonate solution, dried (MgSO₄) and concentrated *in vacuo* to afford the title compound as a white solid. (MH⁺ 215.13)

Intermediate ZC

Pyridin-2-ylmethyl-carbamic acid phenyl ester

The title compound is prepared analogously to pyridin-3-yl-carbamic acid phenyl ester, by substituting C-pyridin-2-yl-methylamine for 3-aminopyridine.

Intermediate ZD

(3-Hydroxy-benzyl)-carbamic acid phenyl ester

The title compound is prepared analogously to pyridin-3-yl-carbamic acid phenyl ester, by substituting 3-aminomethyl-phenol for 3-aminopyridine.

Intermediate ZE

(4-Sulfamoyl-phenyl)-carbamic acid phenyl ester

The title compound is prepared analogously to pyridin-3-yl-carbamic acid phenyl ester, by substituting 4-amino-benzenesulfonamide for 3-aminopyridine.

Intermediate ZF

(3-Sulfamoyl-phenyl)-carbamic acid phenyl ester

The title compound is prepared analogously to pyridin-3-yl-carbamic acid phenyl ester, by substituting 3-amino-benzenesulfonamide for 3-aminopyridine.

Intermediate ZG

N-{(1S,2R,3S,4R)-4-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-3-hydroxy-propionamide

ZG1: 3-tert-Butoxy-N-[(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide

The title compound is prepared as described for (R)-2-benzyloxy-N-[(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide (Example 181, Step 1), by replacing (R)-2-benzyloxy-propionic acid with 3-tert-butoxypropionic acid.

ZG2: 3-tert-Butoxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide

The title compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Intermediate J7), by substituting 3-tert-butoxy-N-[(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide (Intermediate ZG1) for N-[(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide.

ZG3: {(R)-1-[9-[(1R,2S,3R,4S)-4-(3-tert-Butoxy-propionylamino)-2,3-dihydroxy-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester

The title compound is prepared analogously to {(R)-1-[9-[(1R,2S,3R,4S)-4-((R)-2-benzyloxy-propionylamino)-2,3-dihydroxy-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester (Example 181, step 3), by substituting 3-tert-butoxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Intermediate ZG2) for (R)-2-benzyloxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Example 181, step 2).

ZG4: N-{(1S,2R,3S,4R)-4-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-3-hydroxy-propionamide

The title compound is prepared analogously to (R)-N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-benzyloxy-propionamide (Example 181, step 4), by substituting {(R)-1-[9-[(1R,2S,3R,4S)-4-(3-tert-butoxy-propionylamino)-2,3-dihydroxy-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester (Intermediate ZG3) for {(R)-1-[9-[(1R,2S,3R,4S)-4-((R)-2-Benzyloxy-propionylamino)-2,3-dihydroxy-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester (Example 181, step 3).

Intermediate ZH

N-((1S,2R,3S,4R)-4-{2-((R)-3-Amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide

The title compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-3-hydroxy-propionamide (Intermediate ZG), by substituting 4,4'-(2-aminoethylidene)bis-phenol (prepared as described by Schelkun, R.M. et al. Bioorg. Med. Chem. Lett. (1999), 9(16), pp 2447-2452) for 2,2-diphenyl-ethylamine.

Intermediate ZI**2-Amino-1,1-bis-(4-chloro-phenyl)-ethanol**

The title compound is prepared by combining 4,4'-dichlorobenzophenone (5g, 20mmol) and zinc iodide (480mg, 1.49mmol) in DCM (100mL). Trimethylsilyl cyanide (2.17g, 21.9mmol) is added, and the reaction is stirred at room temperature for 18 hours. The reaction is washed with water (100mL) and dried over magnesium sulfate, before filtering and removal of the solvent under reduced pressure. The residue is redissolved in dry THF (40mL), 1.0M borane in THF (40mL) is added, and the reaction is stirred at reflux for 24 hours. After cooling, the volatile components are removed under reduced pressure, the residue is taken up in methanol (100mL). Concentrated hydrochloric acid is added, and the reaction is refluxed for a further 2 hours, before once more removing the volatile components under reduced pressure, to give the title compound as a hydrochloride salt.

Intermediate ZJ

Acetic acid [(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl-carbamoyl]-methyl ester

The title compound is prepared analogously to Intermediate J6, from Intermediate J5, replacing propionyl chloride with acetoxyacetylchloride.

Intermediate ZK**Acetic acid ((1S,2R,3S,4R)-4-{6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-2,3-dihydroxy-cyclopentylcarbamoyl)-methyl ester**

Acetic acid [(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl-carbamoyl]-methyl ester (Intermediate ZJ; 1 equivalent) and 4,4'-(2-aminoethylidene)bis-phenol (1.1 equivalents; prepared as described by Schelkun, R.M. *et al. Bioorg. Med. Chem. Lett.* (1999), **9(16)**, pp 2447-2452) are combined in dry THF and treated with DIPEA (1.2 equivalents) and stirred at 50°C overnight. The reaction is diluted with ethyl acetate and washed consecutively with water (x2) and brine, before drying over magnesium sulfate, filtering and removal of the volatile components under reduced pressure to afford the title compound.

Intermediate ZL

Acetic acid ((1S,2R,3S,4R)-4-{6-[2,2-bis-(4-chloro-phenyl)-2-hydroxy-ethylamino]-2-chloro-purin-9-yl}-2,3-dihydroxy-cyclopentylcarbamoyl)-methyl ester

The title compound is prepared analogously to acetic acid ((1S,2R,3S,4R)-4-{6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-2,3-dihydroxy-cyclopentylcarbamoyl)-methyl ester (Intermediate ZK), by substituting 2-amino-1,1-bis-(4-chloro-phenyl)-ethanol (Intermediate ZI) for 4,4'-(2-aminoethylidene)bis-phenol.

Intermediate ZM

Acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-((S)-1-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl}-methyl ester

The title compound is prepared by combining acetic acid [(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl-carbamoyl]-methyl ester (1 equivalent) and (S)-2-amino-3-phenyl-propan-1-ol 1 equivalent) in dichloromethane with triethylamine (1.1 equivalents) and stirring overnight. The reaction is diluted with dichloromethane, and washed consecutively with 0.1M hydrochloric acid, water and brine, before drying over magnesium sulfate. Filtration and removal of the volatile components under reduced pressure, affords the title compound.

Intermediate ZN

Acetic acid ((1S,2R,3S,4R)-4-{2-chloro-6-[(S)-1-hydroxymethyl-2-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentylcarbamoyl)-methyl ester

The title compound is prepared analogously to acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-((S)-1-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl}-methyl ester (intermediate ZM), by substituting 4-((S)-2-amino-3-hydroxy-propyl)-phenol for (S)-2-amino-3-phenyl-propan-1-ol.

Intermediate ZO

N-((1S,2R,3S,4R)-4-{2-((R)-3-Amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-chloro-phenyl)-2-hydroxy-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

ZO1: Acetic acid {(1S,2R,3S,4R)-4-[6-[2, 2-bis-(4-chloro-phenyl)-2-hydroxy-ethylamino]-2-((R)-3-tert-butoxycarbonylamino-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl} -methyl ester

A suspension of acetic acid ((1S,2R,3S,4R)-4-{6-[2, 2-bis-(4-chloro-phenyl)-2-hydroxy-ethylamino]-2-chloro-purin-9-yl}-2,3-dihydroxy-cyclopentylcarbamoyl)-methyl ester (intermediate ZL; 1 equivalent) and (3R)-(+)-3-(Boc-amino)pyrrolidine (4 equivalents) in acetonitrile is treated with a catalytic amount of sodium iodide and then heated using microwave radiation in a Personal Chemistry Emrys™ Optimizer microwave reactor at 160°C. After 1 hour, the solvent is removed *in vacuo*; purification by column chromatography / crystallisation affords the title compound.

ZO2: N-((1S,2R,3S,4R)-4-{2-((R)-3-Amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-chloro-phenyl)-2-hydroxy-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

A solution of acetic acid {(1S,2R,3S,4R)-4-[6-[2, 2-bis-(4-chloro-phenyl)-2-hydroxy-ethylamino]-2-((R)-3-tert-butoxycarbonylamino-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl} -methyl ester (Intermediate ZO1) in MeOH (~0.5M) is treated with an equal volume of 4M HCl in dioxane and stirred at room temperature for 2 hours. The solvent is removed *in vacuo* and purification is carried out by column chromatography / crystallisation to afford the title compound.

Intermediate ZP

N-{(1S,2R,3S,4R)-4-[2-(4-Amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide.

ZP1: N-{(1S,2R,3S,4R)-4-[6-(2,2-Diphenyl-ethylamino)-2-hydrazino-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide

The title compound was prepared analogously to N-[(1S,2R,3S,4R)-4-(6-{[bis-(4-methoxy-phenyl)-methyl]-amino}-2-hydrazino-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide (Example 168, step 2), by substituting acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl} -methyl ester (Intermediate Q1) for N-[(1S,2R,3S,4R)-4-(6-{[Bis-(4-methoxy-phenyl)-

methyl]-amino}-2-chloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide
(example 168, step1).

ZP2: N- {(1S,2R,3S,4R)-4-[6-(2,2-Diphenyl-ethylamino)-2-(4-nitro-pyrazol-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide

The title compound is prepared analogously to 1-[6-{[bis-(4-methoxy-phenyl)-methyl]-amino}-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid ethyl ester (example 168, step 3), by reaction of N- {(1S,2R,3S,4R)-4-[6-(2,2-diphenyl-ethylamino)-2-hydrazino-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide (Intermediate ZP1) and sodium nitromalondehyde (prepared as described by Fanta P.E. Org. Syntheses, Coll. Vol. 4 (1963), pp 844-845).

ZP3: N- {(1S,2R,3S,4R)-4-[2-(4-Amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide

The title compound is prepared by dissolving N- {(1S,2R,3S,4R)-4-[6-(2,2-Diphenyl-ethylamino)-2-(4-nitro-pyrazol-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide (Intermediate ZP2) in methanol (~1.0M), adding a 2:1 mixture by weight of activated carbon and ferric chloride and an excess of hydrazine monohydrate, and stirring the resulting mixture at 65°C for four hours. Filtration through Celite™, removal of the volatile components under reduced pressure, and purification by column chromatography / crystallisation, affords the title compound.

Intermediate ZQ

N-((1S,2R,3S,4R)-4-{2-(4-Amino-pyrazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

The title compound is prepared analogously to N- {(1S,2R,3S,4R)-4-[2-(4-amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide (Intermediate ZP), by substituting acetic acid ((1S,2R,3S,4R)-4-{6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-2,3-dihydroxy-cyclopentylcarbamoyl)-methyl ester (Intermediate ZK) for acetic acid {(1S,2R,3S,4R)-4-

[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl} - methyl ester (Intermediate Q1).

Intermediate ZR

N-{(1S,2R,3S,4R)-4-[2-(4-Amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide

The title compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-(4-amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide (Intermediate ZP), by substituting N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide (Intermediate J) for acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl} - methyl ester (Intermediate Q1).

Intermediate ZS

1-[9-[(1R,2S,3R,4S)-2,3-Dihydroxy-4-(2-hydroxy-acetyl-amino)-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid ethyl ester

The title compound is prepared analogously to 1-[6-{[bis-(4-methoxy-phenyl)-methyl]-amino}-9-[(1R,2S,3R,4S)-2,3-dihydroxy-4-propionyl-amino-cyclopentyl]-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid ethyl ester (example 168, step3), substituting N-{(1S,2R,3S,4R)-4-[6-(2,2-diphenyl-ethylamino)-2-hydrazino-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide (Intermediate ZP1) for N-[(1S,2R,3S,4R)-4-(6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-2-hydrazino-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide (Example 168, step 2).

Intermediate ZT

N-{(1S,2R,3S,4R)-4-[2-(4-Amino-pyrazol-1-yl)-6-((S)-1-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide

The title compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-(4-amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide (Intermediate ZP), by substituting acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-((S)-1-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-

cyclopentylcarbamoyl} -methyl ester (Intermediate ZM) for acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl} -methyl ester (Intermediate Q1).

Intermediate ZU

N-{(1S,2R,3S,4R)-4-[2-(4-Amino-imidazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide

ZU1: N-{(1S,2R,3S,4R)-4-[6-(2,2-Diphenyl-ethylamino)-2-(4-nitro-imidazol-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide

The title compound is prepared by dissolving acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl} -methyl ester (Intermediate Q1) in N-methyl-2-pyrrolidinone, followed by potassium carbonate (5 equivalents) and 4-nitro-1H-imidazole (10 equivalents). The mixture is heated by microwave irradiation to 150°C for two hours, then diluted with ethyl acetate and washed consecutively with water (x2) and brine, before drying over magnesium sulfate.

Filtration, removal of the volatile components under reduced pressure and purification by flash column chromatography / crystallisation affords the title compound.

ZU2: N-{(1S,2R,3S,4R)-4-[2-(4-Amino-imidazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide

The title compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-(4-amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide (Intermediate ZP3), by substituting N-{(1S,2R,3S,4R)-4-[6-(2,2-Diphenyl-ethylamino)-2-(4-nitro-imidazol-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide (Intermediate ZU1) for N-{(1S,2R,3S,4R)-4-[6-(2,2-Diphenyl-ethylamino)-2-(4-nitro-pyrazol-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide (Intermediate ZP2).

Intermediate ZV

N-((1S,2R,3S,4R)-4-{2-(4-Amino-imidazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

The title compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-(4-amino-imidazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide (Intermediate ZU) by substituting acetic acid ((1S,2R,3S,4R)-4-{6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-2,3-dihydroxy-cyclopentylcarbamoyl)-methyl ester (Intermediate ZK) for acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl}-methyl ester (Intermediate Q1) in step ZU1.

Intermediate ZW

N-{(1S,2R,3S,4R)-4-[2-(3-Amino-[1,2,4]triazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide

The title compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-(4-amino-imidazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide (Intermediate ZU) from {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl}-methyl ester (Intermediate Q1) by substituting 3-nitro-1H-[1,2,4]triazole for 4-nitro-1H-imidazole in step ZU1.

Intermediate ZX

N-((1S,2R,3S,4R)-4-{2-(3-Amino-[1,2,4]triazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

The title compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-(4-amino-imidazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide (Intermediate ZU) by substituting acetic acid ((1S,2R,3S,4R)-4-{6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-2,3-dihydroxy-cyclopentylcarbamoyl)-methyl ester (Intermediate ZK) for acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl}-methyl ester (Intermediate Q1) and 3-nitro-1H-[1,2,4]triazole for 4-nitro-1H-imidazole in step ZU1.

Intermediate ZY

(2S,3S,4R,5R)-5-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

The title compound is prepared as described in WO 9602553.

Preparation of Specific Examples:

Example 1

N-{(1S,2R,3S,4R)-4-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide trifluoroacetate

Step 1: 2-Benzoyloxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-acetamide

The title compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Intermediate J) by replacing propionyl chloride with benzyloxy-acetyl chloride.

Step 2: N-{(1S,2R,3S,4R)-4-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-benzyloxy-acetamide trifluoroacetate

A solution of 2-benzyloxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-acetamide (80 mg, 0.13 mmol) in NMP: MeCN (1 ml of a 1:1 mixture) is treated with sodium iodide (6 mg, 0.04 mmol) followed by (3R)-3-aminopyrrolidine (34 mg, 0.4 mmol). The reaction mixture is heated using microwave radiation in a Personal Chemistry Emrys™ Optimizer microwave reactor at 200 °C. The reaction is shown to be complete by LCMS after 30 minutes. The title compound is obtained after purification by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% TFA).

Step 3: N-{(1S,2R,3S,4R)-4-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide trifluoroacetate

A solution of N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-benzyloxy-acetamide trifluoroacetate (0.022 g, 0.03 mmol) in ethanol (2 ml) under an atmosphere of Argon is treated with palladium hydroxide on carbon (0.05 g, 20%w/w carbon). The reaction mixture is placed under an atmosphere of hydrogen and stirred at room temperature for 30 hours and then filtered through celite™. The filtrate is concentrated *in vacuo* and purification of the crude by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% TFA) yields the title product. (MH+ 573.4)

Example 2

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide To a stirred solution of N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide trifluoroacetate (Example 1) (1.1 g, 2 mmol) in NMP (2 ml) and MeOH (10 ml) is added dropwise phenyl chloroformate (0.47 g, 3 mmol). The reaction mixture is allowed to stir at room temperature for 3 hours and then treated with 3-hydroxybenzylamine (0.55 g, 4.45 mmol) and stirred for a further 2 hours at 80 °C. Purification of the product by C-18 reverse phase column chromatography eluting with acetonitrile : water : TFA (0.1%) (gradient of 0 to 100% acetonitrile) yields the title compound. (MH+ 722.31)

Example 3

((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid benzyl ester

This compound is prepared analogously to Example 2 by replacing N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide trifluoroacetate with {(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-carbamic acid benzyl ester (Intermediate L). (MH+ 783.3)

Example 4**N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide hydrochloride**

A solution comprising N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide trifluoroacetate (Example 1) (0.5 g, 0.87 mmol) and pyridin-3-yl-carbamic acid phenyl ester (Intermediate ZB) (0.198 g, 0.87 mmol) in NMP (1ml) is stirred at 100°C for 1 hour. The solvent is removed *in vacuo* and the title compound is obtained after purification by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% HCl). (MH+ 707.65)

Example 5**N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(3-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide hydrochloride**

This compound is prepared analogously to Example 4 by replacing pyridin-3-yl-carbamic acid phenyl ester (Intermediate ZB) with (3-sulfamoyl-phenyl)-carbamic acid phenyl ester (EP 365484). (MH+ 707.65)


Example 6**N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(4-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide hydrochloride**

This compound is prepared analogously to Example 4 by replacing pyridin-3-yl-carbamic acid phenyl ester (Intermediate ZB) with (4-sulfamoyl-phenyl)-carbamic acid phenyl ester (JP 2002283758). (MH+ 771.60)

Example 7

N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

Step 1:



A reaction mixture comprising (1S,2R,3S,5R)-3-(Di-Boc-amino)-5-(2,6-dichloro-purin-9-yl)-cyclopentane-1,2-diol (Intermediate J4) (3.00g, 5.95 mmol) and 4,4'-(2-aminoethylidene)bis-phenol (prepared according to the preparation of R.M.Schelkun et al. Bioorg. Med. Chem. Lett. 9 (1999) 2447-2452.) (1.5 g, 6.54 mmol) in dry THF (20 ml) is treated with DIPEA (1.2 ml, 7.14 mmol) and stirred at 50°C overnight. The solvent is removed *in vacuo* and the residue is partitioned between DCM and 2M HCl. The organic portion is separated, dried (Mg SO₄) and concentrated *in vacuo* to afford the title compound.

Step 2: (1S,2R,3S,5R)-3-Amino-5-{6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-cyclopentane-1,2-diol

A solution of Example 7 step 1 (3.45 g, 5.80 mmol) in methanol (7 ml) is treated with 4M HCl in dioxane (3 ml) and the reaction mixture is stirred at room temperature overnight. The solvent is removed *in vacuo* and the residue is partitioned between saturated sodium bicarbonate solution and DCM. The organic portion is separated, dried (MgSO₄) and concentrated *in vacuo* to afford the title compound.

Step 3: N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide trifluoroacetate:

A mixture comprising (1S,2R,3S,5R)-3-amino-5-{6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-cyclopentane-1,2-diol (1.03 g, 1.93 mmol) and

potassium carbonate (1.34 g, 9.65 mmol) in DMF (5 ml) is treated with acetoxyacetyl chloride (0.499 g, 4.63 mmol) and TEA (0.78 g, 7.72 mmol) and stirred at room temperature overnight. The solvent is removed *in vacuo* and the residue is dissolved in methanol and 5M KOH. The resulting mixture is filtered to remove the undissolved potassium carbonate and then concentrated *in vacuo*. Purification of the residue by preparative HPLC eluting with acetonitrile : water : TFA (0.1%) (gradient of 5 to 100% acetonitrile) yields the title compound. (MH⁺ 555.42).

Step 4: ((R)-1-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-9-[(1R,2S,3R,4S)-2,3-dihydroxy-4-(2-hydroxy-acetylamino)-cyclopentyl]-9H-purin-2-yl}-pyrrolidin-3-yl)-carbamic acid tert-butyl ester hydrochloride

A mixture comprising N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide trifluoroacetate (475 mg, 0.71 mmol), (R)-pyrrolidin-3-yl-carbamic acid tert-butyl ester (0.529 g, 2.84 mmol) and sodium iodide (106 mg, 0.71 mmol) in acetonitrile (0.5 ml) is heated using microwave radiation in a Personal Chemistry Emrys™ Optimizer microwave reactor at 160 °C. The reaction is shown to be complete by LCMS after 30 minutes. The title compound is obtained after purification by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% HCl).

Step 5: N-((1S,2R,3S,4R)-4-{2-((R)-3-Amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide hydrochloride

This compound is prepared analogously to (1S,2R,3S,5R)-3-amino-5-{6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-cyclopentane-1,2-diol (step 2)

Step 6: N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-{(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

A solution comprising N-((1S,2R,3S,4R)-4-{2-((R)-3-amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide hydrochloride (15 mg, 0.02 mmol) and (3-hydroxy-benzyl)-carbamic acid phenyl ester (prepared according to the method of McDonnell M.E. et al *Bioorganic and Medicinal Chemistry Letters* 14 (2004) 531-534) (6 mg, 0.026 mmol) in MeOH (0.5 ml) is treated with TEA (33 μ l, 0.23 mmol) and then heated at 100°C for 30 minutes. Purification of the crude product by preparative HPLC eluting with acetonitrile : water : TFA (0.1%) (gradient of 10 to 100% acetonitrile) yields the title compound. (MH+ 803.46)

Examples 8-10

These compounds namely,

N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-[3-(3-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide trifluoroacetate (MH+ 803.46) (Example 8),

N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-[3-(4-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide trifluoroacetate (MH+ 803.45) (Example 9) and

N-[(1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide trifluoroacetate (MH+ 739.36) (Example 10) are prepared analogously to Example 6 by replacing (2-hydroxy-benzyl)-carbamic acid phenyl ester with (2-sulfamoyl-benzyl)-carbamic acid phenyl ester, (3-sulfamoyl-benzyl)-carbamic acid phenyl ester and pyridin-2-ylmethyl-carbamic acid phenyl ester respectively. These carbamic acid phenyl esters are prepared according to the method of McDonnell M.E. et al. *Bioorganic and Medicinal Chemistry Letters* 14 (2004) 531-534.

Example 11

((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid methyl ester trifluoroacetate

Step 1: ((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid benzyl ester

A solution of ((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-carbamic acid benzyl ester (Intermediate L) (0.2 g, 0.31 mmol) in NMP/THF (3 ml of a 1:2 mixture) is treated with TEA (0.05 g, 0.46 mmol) followed by chloroformic acid phenyl ester (0.053 g, 0.34 mmol). The reaction mixture is stirred at room temperature for 2 hours and then treated with 2-amino methylpyridine (0.076 g, 0.62 mmol). The mixture is heated to 50°C overnight and then the solvent is removed in vacuo. Purification by C-18 reverse phase column chromatography eluting with acetonitrile : water : HCl (0.1%) (gradient of 0 to 100% acetonitrile) affords the product in acetonitrile which is subsequently washed with saturated sodium bicarbonate solution and extracted with DCM (3 times). The combined organic portions are dried (MgSO₄) and concentrated *in vacuo* to afford the title compound. (MH⁺ 783.3)

Step 2: 1-[(R)-1-[9-((1R,2S,3R,4S)-4-Amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl]-3-pyridin-2-ylmethyl-urea

A solution of ((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid benzyl ester (0.13 g, 166 µmol) in ethanol (9 ml) under an inert atmosphere of Argon is treated with Palladium (10% on charcoal) (44 mg). The reaction mixture is placed under an atmosphere of hydrogen and stirred at room temperature overnight and then filtered through celite™. The filtrate is concentrated *in vacuo* to afford the title compound. (MH⁺ 649.89)

Step 3: ((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid methyl ester trifluoroacetate

A solution of 1-[(R)-1-[9-((1R,2S,3R,4S)-4-amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl]-3-pyridin-2-ylmethyl-urea (17 mg,

26 μmol) in THF (1 ml) is treated with TEA (5 mg, 50 μmol) followed by methyl chloroformate (2.7 mg, 29 μmol). The reaction mixture is stirred at room temperature overnight and purification of the crude mixture by preparative LC-MS affords the title compound. (MH⁺ 707.49)

Example 12

[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{2-[3-(3,4,5,6-tetrahydro-2H-[1,2'-bipyridinyl-4-yl]-ureido]-ethylcarbamoyl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-carbamic acid tert-butyl ester

Step 1: 9-[(1R,4S)-4-(tert-Butoxycarbonyl-propionyl-amino)-cyclopent-2-enyl]-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester:

This compound is prepared analogously to Di-Boc-[(1S,4R)-4-(2,6-dichloro-purin-9-yl)-cyclopent-2-enyl]-amine (Intermediate J3) by replacing di-t-butyl iminodicarboxylate with propionyl-carbamic acid tert-butyl ester.

Step 2: 9-[(1R,2S,3R,4S)-4-(tert-butoxycarbonyl-propionyl-amino)-2,3-dihydroxy-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester

To a stirred suspension comprising 9-[(1R,4S)-4-(tert-butoxycarbonyl-propionyl-amino)-cyclopent-2-enyl]-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester (6.6 g, 10.82 mmol), methane sulphonamide (1.03 g, 10.82 mmol) and AD-mix-a (16.23 g) in t-butanol (40 ml) and water (40 ml) is added osmium tetroxide (3 ml of a 4% solution in water). The reaction mixture is stirred vigorously for 36 hours. The reaction mixture is partitioned between ethyl acetate and water and the organic portion is dried (MgSO₄) and concentrated *in vacuo*. The title product is precipitated from methanol. Further product is derived from the mother liquor by chromatography on silica eluting with DCM : methanol (25:1).

Step 3: {(1S,2R,3S,4R)-4-[2-(2-Amino-ethylcarbamoyl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-carbamic acid tert-butyl ester

A solution comprising 9-[(1R,2S,3R,4S)-4-(tert-butoxycarbonyl-propionyl-amino)-2,3-dihydroxy-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl

ester (3.00 g, 4.66 mmol) in ethylenediamine (5 ml, 75 mmol) is heated at 90°C for 1 hour. After cooling to room temperature, the ethylenediamine is removed *in vacuo* and purification by C-18 reverse phase column chromatography eluting with water (100%) followed by MeOH (100%) yields the title compound.

Step 4: [(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{2-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-ethylcarbamoyl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-carbamic acid tert-butyl ester

To a suspension of {(1S,2R,3S,4R)-4-[2-(2-amino-ethylcarbamoyl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-carbamic acid tert-butyl ester (2.5 g, 4.05 mmol) in IPA (10 ml) is added imidazole-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide (Intermediate C) (151 ml of a 10 mg/ml solution in DCM, 5.56 mmol) and the reaction mixture is stirred at room temperature overnight. The solvent is removed *in vacuo* and the crude product is purified by chromatography on silica eluting with DCM/MeOH (25:1 increasing to 15:1). The resulting solid is further purified by dissolving in DCM and washing the solution with water, drying (MgSO₄) and concentrating *in vacuo* to afford the title product. (MH⁺ 820.7)

Example 13

[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{2-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-ethylcarbamoyl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-carbamic acid methyl ester trifluoroacetate

Step 1: 9-((1R,2S,3R,4S)-4-Amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid {2-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-ethyl}-amide dihydrochloride

The title compound is prepared analogously to (1S,2R,3S,5R)-3-Amino-5-{6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-cyclopentane-1,2-diol (example 7 step 2) by replacing N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide hydrochloride with (1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{2-[3-(3,4,5,6-tetrahydro-2H-

[1,2']bipyridinyl-4-yl)-ureido]-ethylcarbamoyl} -purin-9-yl)-2,3-dihydroxy-cyclopentyl]-carbamic acid tert-butyl ester (Example 12).

Step 2: [(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{2-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-ethylcarbamoyl} -purin-9-yl)-2,3-dihydroxy-cyclopentyl]-carbamic acid methyl ester trifluoroacetate

A mixture comprising 9-((1R,2S,3R,4S)-4-amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid {2-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-ethyl} -amide dihydrochloride (20 mg, 25 μ mol), TEA (12.7 mg, 125 μ mol) in THF (2 ml) is treated with methyl chloroformate (6.0 mg, 63 μ mol) and left to stir at room temperature overnight. The solvent is removed *in vacuo* and purification of the crude product by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% TFA) yields the title product. (MH⁺ 778.6)

Example 14

9-[(1R,2S,3R,4S)-2,3-Dihydroxy-4-(2-hydroxy-acetyl-amino)-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid {2-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-ethyl}-amide trifluoroacetate

Step 1: Acetic acid [(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{2-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-ethylcarbamoyl} -purin-9-yl)-2,3-dihydroxy-cyclopentylcarbamoyl]-methyl ester trifluoroacetate

This compound is prepared analogously to Example 13 by replacing methyl chloroformate with acetoxyacetyl chloride.

Step 2: 9-[(1R,2S,3R,4S)-2,3-Dihydroxy-4-(2-hydroxy-acetyl-amino)-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid {2-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-ethyl} -amide trifluoroacetate

A solution of acetic acid [(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{2-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-ethylcarbamoyl} -purin-9-yl)-2,3-dihydroxy-cyclopentylcarbamoyl]-methyl ester trifluoroacetate (0.015 g, 16 μ mol) in methanol (1 ml) is treated with potassium carbonate (0.01 g, 10 μ mol) and the reaction mixture is allowed to stir at room temperature for 1 hour. The solvent is removed *in*

vacuo and purification of the crude by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% TFA) yields the title product. (MH+ 778.6)

Example 15

N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-((R)-3-pyrrolidin-3-ylureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

N-((1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (Intermediate Q) (60 mg, 0.11 mmol) and 1,3-di(R)-pyrrolidin-3-yl-urea (Intermediate B) (91 mg, 0.46 mmol) in dry DMSO (0.2 ml) is stirred at 100°C for 2 hours. Purification of the resulting mixture by C-18 reverse phase column chromatography eluting with acetonitrile : water : TFA (0.1%) (gradient of 20 to 70% acetonitrile) yields the product which is treated with saturated sodium bicarbonate solution and passed through the C-18 reverse phase column again. The column is washed first with water followed by MeOH (with 1% ammonia) to elute the product which is concentrated *in vacuo* to afford the title compound. (MH+ 685.2)

Example 16

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-((R)-3-pyrrolidin-3-ylureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (Example 15) by replacing 1,3-di(R)-pyrrolidin-3-yl-urea (Intermediate B) with 1-(R)-pyrrolidin-3-yl-3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-urea hydrochloride (Intermediate A). (MH+ 388.8)

Example 17

4-[(R)-3-(3-[(R)-1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl)-ureido)-pyrrolidine-1-carbonyl]-benzoic acid trifluoroacetate

Step 1: N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-((R)-3-pyrrolidin-3-ylureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate

The title compound is prepared analogously to N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-benzyloxy-acetamide trifluoroacetate Example 1 (Step 2) by replacing 2-benzyloxy-N-((1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-acetamide with N-((1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-propionamide (Intermediate J) and by replacing (3R)-3-aminopyrrolidine with 1,3-di(R)-pyrrolidin-3-yl-urea (Intermediate B).

Step 2: 4-[(R)-3-(3-((R)-1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl)-ureido)-pyrrolidine-1-carbonyl]-benzoic acid trifluoroacetate

A solution of N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-((R)-3-pyrrolidin-3-ylureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate (step 1) (0.02 g, 29 μ mol) DIPEA (0.0075 g, 58 μ mol) in NMP (0.2 ml) is treated with a solution of terephthaloyl chloride (0.006 g, 14.5 μ mol) in NMP (0.1 ml). After stirring at room temperature for 1 hour the reaction mixture is purified by C-18 reverse phase column chromatography eluting with acetonitrile : water : TFA (0.1%) (gradient of 0 to 100% acetonitrile) to yield the title compound. (MH⁺ 831.6)

Example 18

N-((R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl)-6-morpholin-4-yl-nicotinamide trifluoroacetate

Step 1: ((R)-1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl)-carbamic acid tert-butyl ester trifluoroacetate:

A reaction mixture comprising N-((1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-propionamide (Intermediate J) (2.5

g, 4.80 mmol) and (3R)-(+)-(3-Boc-amino)pyrrolidine (2.5 g, 13.6 mmol) in DMSO (8 ml) is heated at 100 °C overnight. The resulting mixture is purified by reverse phase column chromatography (Isolute™ C18, 0-100% MeOH in water – 0.1% TFA) to yield the title product.

Step 2: N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide {(R)-1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester trifluoroacetate (3.22 g, 4.80 mmol) is dissolved in 1.25 M HCl in MeOH (60 ml, 75 mmol) and left to stir at room temperature overnight. The solvent is removed *in vacuo* and the crude product is dissolved in a minimal volume of EtOH/saturated sodium carbonate solution and purified by reverse phase column chromatography (Isolute™ C18, 0-100% MeOH in water) to yield the title product.

Step 3: N-{(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-6-morpholin-4-yl-nicotinamide trifluoroacetate

A solution of N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (22.8 mg, 0.04 mmol) in THF (1 ml) is treated with TEA (7.3 mg, 0.072 mmol) and then added to 6-morpholinonicotinoyl chloride (8.2 mg, 0.036 mmol). The reaction mixture is shaken and then allowed to stand at room temperature overnight. The solvent is removed *in vacuo* and purification by C-18 reverse phase column chromatography eluting with acetonitrile : water : TFA (0.1%) (gradient of 0 to 100% acetonitrile) yields the title compound. (MH⁺ 761.4)

Example 19

N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate

A suspension of N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Example 18 step 2) (0.12 mg, 230 μ mol) and sodium hydrogencarbonate (27 mg, 253 μ mol) in DMSO (300 μ l) is treated with phenyl chlorocarbonate (36 mg, 230 μ mol) and then stirred at room temperature for 3 hours. This reaction mixture is added to 2-picolyamine (4.1 mg, 38 μ mol) and stirred at 80 °C for 5 hours. Purification of the crude product by C-18 reverse phase column chromatography eluting with acetonitrile : water : TFA (0.1%) (gradient of 0 to 100% acetonitrile) to yield the title compound. (MH⁺ 705.4)

Example 20

N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-4-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate

Step 1: Imidazole-1-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide:

A mixture comprising N-{(3aR,4S,6R,6aS)-6-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,2-dimethyl-tetrahydro-cyclopenta[1,3]dioxol-4-yl}-propionamide (Intermediate M) (0.24 g, 394 μ mol) and CDI (0.275 g, 1.7 mmol) in dry DCM (6 ml) is stirred at room temperature for 3 hours. The resulting solution is purified by chromatography on silica eluting with 100% DCM changing to 5% MeOH in DCM to afford the title compound as a yellow oil. The oil consists of the imidazole-urea intermediate together with variable amounts of the corresponding isocyanate and imidazole which are equally suitable as precursors to ureas.

Step 2: N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-4-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate

Pyridylmethylamine (4.3 mg, 40 μ mol) is treated with a solution of imidazole-1-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide (25 mg, 40 μ mol) in DCM (1 ml) and the reaction mixture is stirred at room temperature overnight. The

solvent is removed *in vacuo* and the residue is treated with TFA (0.5 ml) and water (0.5 ml). After stirring at room temperature for 3 hours, the reaction mixture is concentrated *in vacuo* and the resulting crude is purified by mass directed preparative LC-MS eluting with acetonitrile: water: trifluoroacetic acid to afford the title compound. (MH+ 705.4)

Example 21

N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-[(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate

This compound is prepared analogously to Example 20 by replacing 4-pyridylmethylamine with 3-hydroxybenzylamine. (MH+720.4)

Example 22

N-[(R)-1-[6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-pyrrolidin-3-yl]-6-morpholin-4-yl-nicotinamide hydrochloride

Step 1: N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide hydrochloride

A reaction mixture comprising [(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionyl-carbamic acid tert-butyl ester (Intermediate G) (2.00g, 4.35 mmol) and 4,4'-(2-aminoethylidene)bis-phenol (prepared by the procedure of R.M.Schelkun et al. Bioorg. Med. Chem. Lett. 9 (1999) 2447-2452.) (1.19 g, 5.20 mmol) in dry THF (40 ml) is treated with DIPEA (0.67 g, 5.20 mmol) and stirred at room temperature for 2 days. The solvent is removed *in vacuo* and the title compound is obtained after purification by reverse phase column chromatography (Isolute™ C18, 20-70% acetonitrile in water – 0.1% HCl).

Step 2: 2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide

This compound is prepared analogously to (1S,2R,3S,5R)-3-mino-5-{6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-cyclopentane-1,2-diol (Example 7 step 2).

Step 3: {(R)-1-[6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester trifluoroacetate

This compound is prepared analogously to ((R)-1-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-9-[(1R,2S,3R,4S)-2,3-dihydroxy-4-(2-hydroxy-acetylamino)-cyclopentyl]-9H-purin-2-yl}-pyrrolidin-3-yl)-carbamic acid tert-butyl ester hydrochloride (Example 7 step 4).

Step 4: N-((1S,2R,3S,4R)-4-{2-((R)-3-Amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide

A solution of {(R)-1-[6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester trifluoroacetate (0.4 g, 0.57 mmol) in DCM (5 ml) and TFA (2.5 ml) is stirred at room temperature for 2 hours. The solvent is removed in vacuo to afford the title compound.

Step 5: N-{(R)-1-[6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-pyrrolidin-3-yl}-6-morpholin-4-yl-nicotinamide hydrochloride

A suspension of N-((1S,2R,3S,4R)-4-{2-((R)-3-amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (20 mg, 33 μ mol) in dry THF is treated with NMP (0.5 ml) followed by TEA (13.4 mg, 0.13 mmol) and 6-morpholinonicotinoyl chloride (8.3 mg, 37 μ mol). The reaction mixture is stirred at room temperature for 30 minutes and then the solvent is removed *in vacuo*.

Purification by C-18 reverse phase column chromatography eluting with acetonitrile : water : HCl (0.1%) (gradient of 0 to 100% acetonitrile) yields the title compound. (MH⁺ 793.4)

Example 23**N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide hydrochloride**

Step 1: {(R)-1-[6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid phenyl ester

A mixture comprising N-((1S,2R,3S,4R)-4-{2-((R)-3-amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 22 step 4) (50 mg, 83 μ mol) and potassium carbonate (46 mg, 332 μ mol) in NMP (1 ml) is treated with phenylchloroformate and stirred at room temperature for 90 minutes. This mixture was used in the next step without purification.

Step 2: N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide hydrochloride

A mixture comprising {(R)-1-[6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid phenyl ester (12 mg, 16 μ mol) and 2-(aminomethyl)pyridine (5.4 mg, 50 μ mol) in NMP (0.5 ml) is heated to 100°C for 2 hours. After cooling to room temperature, purification by C-18 reverse phase column chromatography eluting with acetonitrile : water : HCl (0.1%) (gradient of 0 to 100% acetonitrile) yields the title compound. (MH⁺ 737.5)

Example 24**N-((1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-{(R)-3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl)-propionamide hydrochloride**

This compound is prepared analogously to N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-

yl}-2,3-dihydroxy-cyclopentyl)-propionamide hydrochloride (Example 23) by replacing 2-(aminomethyl)pyridine with the appropriate amine. (MH+ 752.5)

Example 25

N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide:

A solution of N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide trifluoroacetate (0.2 g, 0.28 mmol) in dry THF (10 ml) is treated with TEA (113 mg, 1.12 mmol) followed by 3-isocyanato-pyridine (38 mg, 0.31 mmol). The reaction mixture is heated at 50 °C overnight. The solvent is removed *in vacuo* and purification by reverse phase column chromatography (Isolute™ C18, 10-50% acetonitrile in water – 0.1% TFA) affords the product. The product is further purified by dissolving in MeOH and treating with saturated sodium bicarbonate solution. The mixture is passed through a pre-washed (400 ml MeOH followed by 400 ml water) eluting with 0.5% ammonia 880: water (100 ml) followed by water (400 ml) and finally MeOH to afford the title compound. (MH+ 693.4)

Example 26

((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid methyl ester hydrochloride

Step 1: ((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid benzyl ester trifluoroacetate

A solution comprising ((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-carbamic acid benzyl ester (Intermediate L) (0.1 g, 0.15 mmol), pyridine-3-isocyanate (0.02 g, 0.17 mmol) and TEA (0.017 g, 0.17 mmol) in THF (2 ml) is stirred at room temperature overnight. The solvent is removed *in vacuo* and purification is carried out by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% TFA). The

fractions are collected and the MeCN is removed *in vacuo*. The remaining aqueous portion is basified with saturated sodium bicarbonate solution and extracted with DCM. The combined organic extracted are dried (MgSO₄) and concentrated *in vacuo* to afford the title product. MS (ES+) *m/e* 769 (MH⁺).

Step 2: 1-{(R)-1-[9-((1R,2S,3R,4S)-4-Amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-3-pyridin-3-yl-urea

To a solution of ((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid benzyl ester trifluoroacetate (step 1) (35 mg, 46 μmol) in ethanol (1 ml) under an inert atmosphere of argon is added 10% palladium on carbon (10 mg). The reaction mixture is purged with argon and placed under a positive atmosphere of hydrogen overnight after which time, the mixture is filtered through celite and the catalyst washed with ethanol. The organic portions are combined and concentrated *in vacuo* to yield the title compound. MS (ES+) *m/e* 635 (MH⁺).

Step 3: ((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid methyl ester hydrochloride

This compound is prepared analogously to Example 11 by replacing 1-{(R)-1-[9-((1R,2S,3R,4S)-4-amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-3-pyridin-2-ylmethyl-urea with 1-{(R)-1-[9-((1R,2S,3R,4S)-4-Amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-3-pyridin-3-yl-urea. (MH⁺ 693.5)

Example 27

N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-[(R)-3-[3-(3-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide hydrochloride

A suspension of N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide (Example 18 step 2)

(20 mg, 0.035 mmol) and 3-isocyanato-benzenesulfonamide (Intermediate S) (17.7 mg, 0.095 mmol) in THF (1 ml) and DMF (1 ml) is stirred at room temperature overnight. The solvent is removed in vacuo and purification by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% HCl) affords the title compound. (MH+ 769.5)

Example 28

N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-[(R)-3-[3-(4-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide hydrochloride

This compound is prepared analogously to Example 27 with the appropriate isocyanate. (MH+ 769.5)

Example 29

1-[(R)-1-[9-((1R,2S,3R,4S)-4-Amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl]-3-pyridin-2-ylmethyl-urea

Step 1: ((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid benzyl ester.

A solution of {(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-carbamic acid benzyl ester (Intermediate L) (0.2 g, 0.31 mmol) in NMP/THF (3 ml of a 1:2 mixture) is treated with TEA (0.05 g, 0.46 mmol) followed by chloroformic acid phenyl ester (0.053 g, 0.34 mmol). The reaction mixture is stirred at room temperature for 2 hours and then treated with 2-amino methylpyridine (0.076 g, 0.62 mmol). The mixture is heated to 50°C overnight and then the solvent is removed in vacuo. Purification by C-18 reverse phase column chromatography eluting with acetonitrile : water : HCl (0.1%) (gradient of 0 to 100% acetonitrile) affords the product in acetonitrile which is subsequently washed with saturated sodium bicarbonate solution and extracted with DCM (3 times). The combined organic portions are dried (MgSO₄) and concentrated *in vacuo* to afford the title compound. (MH+ 783.3)

Step 2: 1-{(R)-1-[9-((1R,2S,3R,4S)-4-Amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-3-pyridin-2-ylmethyl-urea

A solution of ((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid benzyl ester (0.13 g, 166 μ mol) in ethanol (9 ml) under an inert atmosphere of Argon is treated with Palladium (10% on charcoal) (44 mg). The reaction mixture is placed under an atmosphere of hydrogen and stirred at room temperature overnight and then filtered through celite™. The filtrate is concentrated *in vacuo* to afford the title compound. (MH+ 649.89)

Example 30-32

These compounds namely,

N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-acetamide trifluoroacetate (MH+ 691.5) (Example 30),

Cyclopropanecarboxylic acid ((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-amide trifluoroacetate (MH+ 717.5) (Example 31) and

cyclobutanecarboxylic acid ((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-amide trifluoroacetate (MH+ 731.5) (Example 32)

are prepared analogously (Example 11) by replacing methyl chloroformate with the appropriate acid chloride.

Example 33

N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(N'-cyano-N''-pyridin-3-ylmethyl-guanidino)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide

Step 1: (1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-cyano-2-phenyl-isoureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentanecarboxylic acid ethylamide

A solution of N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Example 18 step 2) (50 mg) and diphenyl cyanocarbodiimide (21 mg) in dry DCM (2 ml) is treated with TEA (13 μ l) and then stirred at room temperature for 5 hours. The solvent is removed *in vacuo* and the resulting residue is partitioned between EtOAc and water. The organic portion is separated, dried (Na_2SO_4) and concentrated in *vacuo* to afford the crude product which is purified by preparative HPLC to afford the title compound. (MH+ 715).

Step 2: N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(N'-cyano-N"-pyridin-3-ylmethyl-guanidino)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide

A mixture comprising (1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-cyano-2-phenyl-isoureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentanecarboxylic acid ethylamide (80 mg, 0.11 mmol) and 2-(aminomethyl)pyridine (35 μ l, 0.33 mmol) in absolute ethanol (1 ml) is using microwave radiation in a Personal Chemistry EmrysTM Optimizer microwave reactor at 100 °C for 2500 s followed by 120 °C for 1 hour. The solvent is removed *in vacuo* and purification is carried out by preparative HPLC. The fractions are combined, treated with saturated sodium bicarbonate solution and then extracted with ethyl acetate (3x). The organic portions are dried (Na_2SO_4) and concentrated in *vacuo* to afford the title compound. (MH+ 729)

Example 34

((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-((R)-3-pyrrolidin-3-ylureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid methyl ester

This compound is prepared analogously to {(R)-1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester trifluoroacetate (Example 18 step 1) by replacing N-

{{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide (Intermediate J) with {{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -carbamic acid methyl ester (Intermediate T) and by replacing (3R)-(+)-(3-Boc-amino)pyrrolidine with 1,3-di(R)-pyrrolidin-3-yl-urea (Intermediate B). (MH+ 685.2)

Example 35

N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-phenyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide hydrochloride

This compound is prepared analogously to Example 27 by replacing 3-isocyanato-benzenesulfonamide (Intermediate S) with phenyl isocyanate. (MH+ 690.9)

Example 36

4-[(R)-3-(3-{(R)-1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-ureido)-pyrrolidine-1-carbonyl]-benzoic acid methyl ester

This compound is prepared analogously to Example 17 by replacing terephthaloyl chloride with methyl-4-chlorocarbonyl benzoate. Purification is carried out by C-18 reverse phase column chromatography eluting with acetonitrile : water : HCl (0.1%) (gradient of 20 to 100% acetonitrile) to yield the title compound. (MH+ 845.58)

Example 37

N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate

A suspension of N-{{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide (Example 18 step 2) (20 mg, 38 μ mol) and sodium hydrogencarbonate (4.5 mg, 42 μ mol) in DMSO (300 μ l) is treated with phenyl chloroformate (36 mg, 230 μ mol) and then stirred at room temperature for 3 hours. This reaction mixture is added to 3-picolyamine (4.1 mg, 38

μmol) and stirred at 80 °C for 5 hours. Purification of the crude product by C-18 reverse phase column chromatography eluting with acetonitrile : water : TFA (0.1%) (gradient of 0 to 100% acetonitrile) to yield the title compound. (MH+ 705.4)

Example 38

((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid benzyl ester trifluoroacetate

A solution comprising {(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -carbamic acid benzyl ester (Intermediate L) (0.1 g, 0.15 mmol), pyridine-3-isocyanate (0.02 g, 0.17 mmol) and TEA (0.017 g, 0.17 mmol) in THF (2 ml) is stirred at room temperature overnight. The solvent is removed *in vacuo* and purification is carried out by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% TFA). The fractions are collected and the MeCN is removed *in vacuo*. The remaining aqueous portion is basified with saturated sodium bicarbonate solution and extracted with DCM. The combined organic extracted are dried (MgSO₄) and concentrated *in vacuo* to afford the title product. (MH+ 769.5)

Example 39

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(3-hydroxy-phenyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate

Step 1: Imidazole-1-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-p-ropionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl} -amide:

A mixture comprising N-{(3aR,4S,6R,6aS)-6-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,2-dimethyl-tetrahydro-cyclopenta[1,3]dioxol-4-yl} -propionamide (Intermediate M) (0.24 g, 394 μmol) and CDI (0.275 g, 1.7 mmol) in dry DCM (6 ml) is stirred at room temperature for 3 hours. The resulting solution is purified by chromatography on silica eluting with 100% DCM changing to 5% MeOH in DCM to

afford the title compound as a yellow oil. The oil consists of the imidazole-urea intermediate together with variable amounts of the corresponding isocyanate and imidazole which are equally suitable as precursors to ureas.

Step 2: N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(3-hydroxy-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate:

3-Aminophenol (4.3 mg, 40 μ mol) is treated with a solution of imidazole-1-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide (25 mg, 40 μ mol) in DCM (1 ml) and the reaction mixture is stirred at room temperature overnight. The solvent is removed *in vacuo* and the residue is treated with TFA (0.5 ml) and water (0.5 ml). After stirring at room temperature for 3 hours, the reaction mixture is concentrated *in vacuo* and the resulting crude is purified by mass directed preparative LC-MS eluting with acetonitrile: water: trifluoroacetic acid to afford the title compound. (MH⁺ 706.4)

Example 40

N-[(1S,2R,3S,4R)-4-[2-[(R)-3-(4-Acetylamino-benzenesulfonylamino)-pyrrolidin-1-yl]-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate

N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide (Example 18 step 2) (23 mg, 40 μ mol) is treated with a solution of 4-acetamidobenzenesulfonyl chloride (9.5 mg, 0.039 mmol) in NMP (0.5 ml) and stirred at room temperature overnight. Purification by preparative LCMS affords the title compound. (MH⁺ 768.50)

Example 41

N-[(1S,2R,3S,4R)-4-[2-[(R)-3-(4,5-Dihydro-1H-imidazol-2-ylamino)-pyrrolidin-1-yl]-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide

A mixture comprising N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide (Example

18 step 2) (30 mg, 0.053 mmol) and 4,5-dihydro-1H-imidazole-2-thiol hydroiodide (24 mg, 0.106 mmol) in absolute ethanol (2 ml) is treated with DMAP (catalytic amount) and TEA (29 μ l, 0.212 mmol). The resulting mixture is heated using microwave radiation in a Personal Chemistry Emrys™ Optimizer microwave reactor at 150 °C for 4000 s. The solvent is removed in vacuo and purification of the crude product by preparative HPLC ((30-95% acetonitrile in water – 0.1% TFA) affords the title compound. (MH+ 639.63)

Example 42

N-{(1S,2R,3S,4R)-4-[2-{N'-[1-Cyclohexyl-methylidene]-hydrazino}-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide

Step 1: N-{(1S,2R,3S,4R)-4-[6-(2,2-Diphenyl-ethylamino)-2-hydrazino-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide:

N-{(1S,2R,3S,4R)-4-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Intermediate J) (1.0 g, 1.91 mmol) and hydrazine monohydrate (12 ml) is stirred for 72 h and then isopropyl alcohol (10 ml) is added. The solvent is removed *in vacuo* and the residue is water (10 ml) and stirred for 12 h. The fine solid obtained is filtered, washed with water and dried *in vacuo* to afford the product. LC-MS (0.1% formic acid, acetonitrile) (MH+ 517)

Step 2: N-{(1S,2R,3S,4R)-4-[2-{N'-[1-Cyclohexyl-methylidene]-hydrazino}-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide

To a solution of N-{4-[6-(2,2-diphenyl-ethylamino)-2-hydrazino-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (step 1) (0.1 g, 0.19 mmol) in dry methanol (5 ml) is added cyclohexane carboxaldehyde (0.026 g, 0.23 mmol). The reaction mixture is heated at reflux for 12 h. The reaction mixture is concentrated *in vacuo* purification by preparative TLC affords the title compound. LC-MS (0.1% formic acid, acetonitrile): (MH+ 611.45)

Example 43

1-[(R)-1-[9-[(1R,2S,3R,4S)-2,3-Dihydroxy-4-propylamino-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl]-3-pyridin-3-yl-urea hydrochloride

A solution of 1-[(R)-1-[9-[(1R,2S,3R,4S)-4-amino-2,3-dihydroxy-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl]-3-pyridin-3-yl-urea (Example 26 step 2) (17 mg, 26 μmol) in DCE (1 ml) is treated with propanaldehyde (1.5 mg, 26 μmol) and placed under an atmosphere of argon. Sodium triacetoxyborohydride (10 mg, 51 μmol) is added and the reaction mixture is stirred at room temperature overnight. The mixture is quenched with 2M NaOH (5 drops) and the solvent is removed *in vacuo*. The residue is purified by C-18 reverse phase column chromatography eluting with acetonitrile : water : HCl (0.1%) (gradient of 20 to 100% acetonitrile) to yield the title compound. (MH⁺ 677.14)

Example 44

N-[(1S,2R,3S,4R)-4-(6-Amino-2-{N'-[1-cyclohexylmethylidene]hydrazino}-purin-9-yl)-2,3-dihydroxycyclopentyl]propionamide

This compound is prepared analogously to Example 42 by replacing N-[(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide (Intermediate J) with N-[(1S,2R,3S,4R)-4-(6-amino-2-chloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (Intermediate ZA). (MH⁺ 431.33)

Example 45

(R)-3-(3-[(R)-1-[9-[(1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl]-ureido)-pyrrolidine-1-carboxylic acid pyridin-3-ylamide hydrochloride

This compound is prepared analogously to Example 23 by replacing {(R)-1-[6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-9-[(1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl]-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid phenyl ester with N-[(1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-[(R)-3-pyrrolidin-3-ylureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate

(Example 17 step 1) and by replacing (aminomethyl)pyridine with pyridin-3-yl-carbamic acid phenyl ester (Intermediate ZB). (MH+ 803.8)

Examples 46-47

These compounds namely,

9-[(1R,2S,3R,4S)-4-(3,3-dimethyl-ureido)-2,3-dihydroxy-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid {2-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-ethyl}-amide trifluoroacetate (MH+ 791.6) (Example 46) and acetic acid [(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{2-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-ethylcarbamoyl}-purin-9-yl)-2,3-dihydroxy-cyclopentylcarbamoyl]-methyl ester trifluoroacetate (Example 47) are prepared analogously to Example 13 by replacing methyl chloroformate with either dimethylcarbamyl chloride or acetoxyacetyl chloride respectively.

Example 48

9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-methanesulfonylamino-ethyl)-amide

Step 1: 6-(2,2-Diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester

6-(2,2-Diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester hydrochloride (prepared using the procedure described in international patent application WO 2001/94368) (35 g, 85.3 mmol) is placed in a flask under an atmosphere of argon. Dry CHCl_3 (300 ml) and N,O-bis(trimethylsilyl)acetamide (61 ml) are added and the reaction mixture is refluxed for 1 hour. The reaction mixture is allowed to cool and any volatiles removed *in vacuo*. To the resulting oil is added MeOH (300 ml). The resulting white solid is filtered and washed with MeOH (2 x 200 ml) and then dried in a vacuum oven to give the title compound. ^1H NMR (DMSO, 400 MHz).

Step 2: 6-(2,2-Diphenyl-ethylamino)-9-((1R,4S)-4-hydroxy-cyclopent-2-enyl)-9H-purine-2-carboxylic acid methyl ester

To 6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester (5g 13.4mmol) under an atmosphere of argon is added dry deoxygenated tetrahydrofuran (100 ml) and

dry dimethyl sulfoxide (2 ml). Sodium hydride 95% (0.32 g, 13.4 mmol) is then added and the solution is stirred at 40°C. Separately to (1S,4R)-cis 4-Acetoxy-2-cyclopenten-1-ol (1.89 g, 13.4 mmol), triphenylphosphine (0.53 g, 2.0 mmol) and tris(dibenzylideneacetone)dipalladium(0) (0.69 g, 0.67 mmol) is added dry deoxygenated tetrahydrofuran (20 ml) and the mixture stirred at room temperature for 10 minutes. This solution is added to the anion solution via syringe and the resulting mixture is then stirred at 80°C. The reaction is shown to be complete by LCMS after 2 hours. The reaction mixture is allowed to cool, methanol is added and a solid is filtered. The filtrate is concentrated *in vacuo* and the title compound is obtained by precipitation from dichloromethane/hexane. ¹H NMR (MeOD, 400 MHz); 8.15(s, 1H), 7.40-7.15(m, 10H), 6.20(m, 1H), 5.95(m, 1H), 5.50(m, 2H), 4.75(m, 2H), 4.55(m, 1H), 4.10(m 2H), 3.90(s, 2H), 3.80(s, 1H), 2.9(m, 1H), 1.75(m, 1H).

Step 3: 6-(2,2-Diphenyl-ethylamino)-9-((1R,4S)-4-ethoxycarbonyloxy-cyclopent-2-enyl)-9H-purine-2-carboxylic acid methyl ester

6-(2,2-Diphenyl-ethylamino)-9-((1R,4S)-4-hydroxy-cyclopent-2-enyl)-9H-purine-2-carboxylic acid methyl ester (2.80 g, 6.14 mmol) is placed in an oven-dried flask under an atmosphere of argon. Dry tetrahydrofuran (30 ml) is added followed by dry pyridine (0.97 g, 12.3 mmol). Ethyl chloroformate (2.66 g, 24.6 mmol) is added slowly and the reaction mixture is stirred at room temperature. The reaction is shown to be complete by LCMS after 3 hours. The solvent is removed *in vacuo* and the residue is partitioned between dichloromethane (200 ml) and 1M HCl (2x 200 ml). The organic layer is washed with water (2 x 100 ml) and brine (2 x 100 ml), dried over MgSO₄, filtered and the solvent is removed *in vacuo*. The title compound is obtained after purification by flash column chromatography (silica, 4% MeOH in dichloromethane). MS (ES+) *m/e* 528.3 (MH⁺).

Step 4: 9-((1R,4S)-4-Di-tert-butoxycarbonylamino-cyclopent-2-enyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester

6-(2,2-Diphenyl-ethylamino)-9-((1R,4S)-4-ethoxycarbonyloxy-cyclopent-2-enyl)-9H-purine-2-carboxylic acid methyl ester (2.2 g, 4.2 mmol) is dissolved in deoxygenated

tetrahydrofuran. The resultant solution is stirred under an atmosphere of argon at room temperature. Di-*t*-butyl iminodicarboxylate (0.9 g, 4.2 mmol), triphenylphosphine (0.16 g, 0.63 mmol) and triethylamine (0.42 g, 4.2 mmol) are added followed by tris(dibenzylideneacetone)dipalladium(0) (0.22 g, 0.21 mmol). The reaction mixture is then stirred at 45°C for 4 hours, allowed to cool to room temperature, methanol is added and the reaction mixture filtered. The filtrate is concentrated *in vacuo*. The resultant oil is purified by column chromatography (silica, 80% ether in hexane) to yield the title compound, MS (ES+) *m/e* 536.4 (MH⁺).

Step 5: 9-((1R,2S,3R,4S)-4-Di-*tert*-butoxycarbonylamino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester

The title compound is prepared from 9-((1R,4S)-4-di-*tert*-butoxycarbonylamino-cyclopent-2-enyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester using a procedure analogous to that of (1R,2S,3R,5S)-3-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-5-(di-Boc-amino)-cyclopentane-1,2-diol (Intermediate JJ2). MS (ES+) *m/e* 689.4 (MH⁺).

Step 6: 9-((1R,2S,3R,4S)-4-Amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester

9-((1R,2S,3R,4S)-4-Di-*tert*-butoxycarbonylamino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester (0.5 g, 0.73 mmol) is dissolved in dioxane and stirred under an atmosphere of argon. 4M HCl in dioxane (3.68 ml, 14.5 mmol) is added and the resultant solution is stirred for 20 hours then concentrated *in vacuo*. The title compound is obtained by flash column chromatography (Isolute™ C18, 0-100% acetonitrile in water). MS (ES+) *m/e* 489.3 (MH⁺).

Step 7: 9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester

9-((1R,2S,3R,4S)-4-Amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester hydrochloride (200 mg, 0.36 mmol) is dissolved in tetrahydrofuran (5 ml). Diisopropylethylamine (0.16 ml, 0.9 mmol) is added

and the solution is stirred for 10 minutes. Propionyl chloride (33 mg, 0.36 mmol) is added and the reaction mixture is stirred at room temperature for 1 hour. The reaction is quenched with methanol and the title compound is obtained by flash column chromatography (Isolute™ C18, 0-100% acetonitrile in water). MS (ES+) *m/e* 545.3 (MH⁺).

Step 8: 9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-amino-ethyl)-amide

9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester (62 mg, 1.0 mmol) is dissolved in ethylene diamine (3.4 ml, 51 mmol) and the solution is stirred at 105°C. The reaction is shown to be complete by LCMS after 45 minutes. The reaction mixture is concentrated *in vacuo* and the title compound is obtained after purification by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water). MS (ES+) *m/e* 573.4 (MH⁺).

Step 9: 9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-methanesulfonylamino-ethyl)-amide

A solution of 9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-amino-ethyl)-amide (0.01 g, 0.017 mmol) in chloroform (1 ml) is treated with mesyl chloride (1 ml of a 3 mg/ml solution in chloroform) and TEA (0.003 ml). After stirring at 5°C for 1 hour, the reaction mixture is diluted with DCM and extracted with 1 M HCl. The organic portion is isolated and concentrated *in vacuo* to yield the title product. (MH⁺ 651.5)

Example 49-55

These compounds,

9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid {2-[3-(3-methoxy-phenyl)-ureido]-ethyl}-amide trifluoroacetate (MH⁺ 722.4) (Example 49),

9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid {2-[3-(3-cyano-phenyl)-ureido]-ethyl} -amide trifluoroacetate (MH+ 717.5) (Example 50),

[3-(2-{[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carbonyl]-amino}-ethyl)-ureido]-acetic acid ethyl ester trifluoroacetate (MH+ 702.5) (Example 51),

9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid [2-(3-ethyl-ureido)-ethyl]-amide trifluoroacetate (MH+ 644.5) (Example 52),

9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid {2-[3-(4-methoxy-phenyl)-ureido]-ethyl} -amide trifluoroacetate (MH+ 722.5) (Example 53),

9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid {2-[3-(4-cyano-phenyl)-ureido]-ethyl} -amide trifluoroacetate (MH+ 717.5) (Example 54) and

4-[3-(2-{[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carbonyl]-amino}-ethyl)-ureido]-benzoic acid methyl ester trifluoroacetate (MH+ 750.4) (Example 55)

are prepared analogously to Example 48 by replacing mesyl chloride (step 9) with the appropriate isocyanate

Example 56

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(4-phenoxy-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate

N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide (Example 18 step 2) (0.023 g, 40 μ mol) is treated with a solution of 4-phenoxyphenyl isocyanate (0.0078 g, 39 μ mol) in NMP (0.5 ml). After stirring at room temperature overnight purification is carried out using mass directed preparative LC-MS eluting with acetonitrile: water: trifluoroacetic acid to afford the title compound. (MH+ 782.5)

Examples 57-90

These compounds namely,

N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-[(R)-3-[3-(2-phenoxy-phenyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (MH+ 782.5) (Example 57),

N-[(1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(4-trifluoromethyl-benzenesulfonylamino)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate (MH+ 779.4) (Example 58),

N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-[(R)-3-[3-(3,4,5-trimethoxy-phenyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (MH+ 780.5) (Example 59),

N-[(1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-((E)-2-phenyl-ethenesulfonylamino)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate (MH+ 737.5) (Example 60),

(3-[(R)-1-[9-[(1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl]-ureido)-acetic acid ethyl ester trifluoroacetate (MH+ 700.5) (Example 61),

Cyclopropanecarboxylic acid [(R)-1-[9-[(1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl]-amide trifluoroacetate (MH+ 639.5) (Example 62),

N-{(R)-1-[9-[(1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-propionamide trifluoroacetate (MH+ 627.5) (Example 63),

N-{(1S,2R,3S,4R)-4-[2-[(R)-3-(3,3-dimethyl-ureido)-pyrrolidin-1-yl]-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide trifluoroacetate (MH+ 642.5) (Example 64),

N-{(R)-1-[9-[(1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-benzamide trifluoroacetate (MH+ 689.5) (Example 65),

N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-isopropyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate (MH+ 656.5) (Example 66),

N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-ethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate (MH+ 642.5) (Example 67),

{(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid benzyl ester trifluoroacetate (MH+ 705.3) (Example 68)

and N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(4-hydroxy-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (MH+ 745.6) (Example 69),

N-[(1S,2R,3S,4R)-4-[2-{(R)-3-[3-(2,3-dihydro-benzo[1,4]dioxin-6-yl)-ureido]-pyrrolidin-1-yl}-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (MH+ 748.5) (Example 70),

benzo[1,2,5]thiadiazole-5-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (MH+ 733.4) (Example 71),

quinoxaline-6-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (MH+ 727.5) (Example 72),

N-[(1S,2R,3S,4R)-4-[2-{(R)-3-[3-(3,4-dihydro-2H-benzo[b][1,4]dioxepin-7-yl)-ureido]-pyrrolidin-1-yl}-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (MH+ 762.5) (Example 73),

N-[(1S,2R,3S,4R)-4-[2-{(R)-3-[3-(4-Cyano-phenyl)-ureido]-pyrrolidin-1-yl}-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (MH+ 715.5) (Example 74),

4-(4-chloro-benzenesulfonyl)-3-methyl-thiophene-2-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (MH+ 870.5) (Example 75),

3-chloro-4-(propane-2-sulfonyl)-thiophene-2-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (MH+ 821.4) (Example 76),

N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(4-trifluoromethoxy-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (MH+ 774.4) (Example 77),

N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(3-methylsulfanyl-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (MH+ 736.4) (Example 78),

N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(4-methylsulfanyl-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (MH+ 736.4) (Example 79),

quinoxaline-2-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (MH+ 727.5) (Example 80),

benzo[1,2,5]thiadiazole-4-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (MH+ 733.4) (Example 81),

5-(3-{(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-ureido)-isophthalic acid dimethyl ester trifluoroacetate (MH+ 806.5) (Example 82),

N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[2-(1H-indol-3-yl)-2-oxo-acetylamin]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (MH+ 742.5) (Example 83),

pyridine-2-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (MH+ 676.5) (Example 84),

1-(4-methoxy-phenyl)-5-methyl-1H-pyrazole-4-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (MH+ 785.5) (Example 85),

3-{(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-ylcarbamoyl}-piperidine-1-carboxylic acid benzylester trifluoroacetate (MH+816.6) (Example 86),

N-{(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-4-dipropylsulfamoyl-benzamide trifluoroacetate (MH+ 838.4) (Example 87),

1-(4-trifluoromethyl-pyrimidin-2-yl)-piperidine-4-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (MH+ 828.4) (Example 88),

N-{(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-nicotinamide trifluoroacetate (MH+ 676.4) (Example 89),

N-{(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-isonicotinamide trifluoroacetate (MH+ 676.3) (Example 90),

are prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(4-phenoxy-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (Example 56) by replacing 4-phenoxyphenyl isocyanate with the appropriate isocyanate or acid chloride. Reactions using acid chlorides also have excess triethylamine added.

Example 91-92

These compounds namely,

N-{(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(1-ethyl-propylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-isonicotinamide hydrochloride (MH 566.4+) (Example 91) and

N-{(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(3,3-dimethyl-butylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-isonicotinamide hydrochloride (MH+ 580.5) (Example 92),

are prepared analogously to N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-benzyloxy-acetamide trifluoroacetate (Example 1 step 2) by replacing (3R)-3-aminopyrrolidine with (R)-N-pyrrolidin-3-yl-isonicotinamide (Intermediate O) and by replacing 2-benzyloxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-acetamide with the appropriate starting compound. The preparation of the starting compounds are either described herein or can be prepared from [(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionyl-carbamic acid tert-butyl ester (Intermediate G) and the appropriate amine using a procedure analogous to 2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 22 steps 1 and 2)

Example 93

5-Methyl-isoxazole-3-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(1-ethyl-propylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide hydrochloride

This compound is prepared analogously to N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-((R)-3-pyrrolidin-3-ylureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (Example 15) by replacing N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide (Intermediate Q) with N-{(1S,2R,3S,4R)-4-[2-Chloro-6-(1-ethyl-propylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide [prepared from [(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionyl-carbamic acid tert-butyl ester (Intermediate G) and the appropriate amine using a procedure analogous to 2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 22 steps 1 and 2)] and by replacing 1,3-di(R)-pyrrolidin-3-yl-urea (Intermediate B) with 5-methyl-isoxazole-3-carboxylic acid (R)-pyrrolidin-3-ylamide (Intermediate P). (MH⁺ 570.4)

Example 94

Cyclobutanecarboxylic acid {(1S,2R,3S,4R)-4-[2-((R)-1-benzyl-pyrrolidin-3-ylamino)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-amide trifluoroacetate

This compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-benzyloxy-acetamide trifluoroacetate Example 1 (Step 2) by replacing 2-benzyloxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-acetamide with cyclobutanecarboxylic acid {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-amide (Intermediate K) and by replacing (3R)-3-aminopyrrolidine with (R)-1-benzyl-pyrrolidin-3-ylamine. (MH⁺ 687.5)

Example 95

9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid {2-[3-(2-diisopropylamino-ethyl)-ureido]-ethyl}-amide trifluoroacetate

To a solution comprising 9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-amino-ethyl)-amide (25 mg, 0.04 mmol) (Example 48 step 8) in toluene (0.8 ml) and IPA (0.4 ml) is added imidazole-1-carboxylic acid (2-diisopropylamino-ethyl)-amide (10 mg, 0.04 mmol) in DCM (0.44 ml). The reaction mixture is stirred at room temperature under an inert atmosphere of Argon overnight and then the solvent is removed *in vacuo*. Purification of the resulting crude product by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% TFA) yields the title product. (MH⁺ 372.3)

Example 96-98

These compounds namely,

N-{(1S,2R,3S,4R)-4-[2-((R)-1-benzyl-pyrrolidin-3-ylamino)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide trifluoroacetate (MH⁺ 661.6) (Example 96),

{(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester trifluoroacetate (MH⁺ 661.6) (Example 97) and
 (R)-3-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-ylamino]-pyrrolidine-1-carboxylic acid tert-butyl ester trifluoroacetate (MH⁺ 671.5) (Example 98),
 are prepared analogously to N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-benzyloxy-acetamide trifluoroacetate Example 1 (Step 2) by replacing 2-benzyloxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-acetamide with N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Intermediate J) and by replacing (3R)-3-aminopyrrolidine with the appropriate amine.

Example 99

9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-propionylamino-ethyl)-amide

Step 1: 9-((1R,2S,3R,4S)-4-Di-tert-butoxycarbonylamino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-amino-ethyl)-amide:

The title compound is prepared analogously to 9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-amino-ethyl)-amide (Example 48, step 8) by replacing 9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester with 9-((1R,2S,3R,4S)-4-di-tert-butoxycarbonylamino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester (Example 48 step 5).

Step 2: 9-((1R,2S,3R,4S)-4-Amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-amino-ethyl)-amide hydrochloride

The title compound is prepared from 9-((1R,2S,3R,4S)-4-amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-amino-ethyl)-

amide hydrochloride analogously to (1S,2R,3S,5R)-3-amino-5-{6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-cyclopentane-1,2-diol (Example 7 step 2).

Step 3: 9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-propionylamino-ethyl)-amide

The title compound is prepared analogously to 9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester (Example 48 step 7) by replacing 9-((1R,2S,3R,4S)-4-amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester hydrochloride with 9-((1R,2S,3R,4S)-4-amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-amino-ethyl)-amide hydrochloride. (MH+ 629.5)

Example 100

N-((1S,2R,3S,4R)-4-[2-(2-Amino-ethylamino)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate

The title compound is prepared analogously to N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-benzyloxy-acetamide trifluoroacetate (Example 1 step 2) by replacing 2-benzyloxy-N-((1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-acetamide with N-((1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-propionamide (Intermediate J) and by replacing (3R)-3-aminopyrrolidine with ethylene diamine. (MH+545.4)

Example 101

9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-{3-[1-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-ylcarbamoyl)-piperidin-4-yl]-ureido}-ethyl) amide trifluoroacetate

Step 1: {(1S,2R,3S,4R)-4-[6-(2,2-Diphenyl-ethylamino)-2-(2-{3-[1-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-ylcarbamoyl)-piperidin-4-yl]-ureido}-ethylcarbamoyl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-carbamamic acid tert-butyl ester

The title compound is prepared analogously to [(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{2-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-ethylcarbamoyl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-carbamamic acid tert-butyl ester (Example 12) by replacing imidazole-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide (Intermediate C) with 4-[(imidazole-1-carbonyl)-amino]-piperidine-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide (Intermediate H).

Step 2: 9-((1R,2S,3R,4S)-4-Amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-{3-[1-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-ylcarbamoyl)-piperidin-4-yl]-ureido}-ethyl)-amide dihydrochloride

A solution of {(1S,2R,3S,4R)-4-[6-(2,2-diphenyl-ethylamino)-2-(2-{3-[1-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-ylcarbamoyl)-piperidin-4-yl]-ureido}-ethylcarbamoyl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-carbamamic acid tert-butyl ester (76.7 mg, 81 μ mol) in MeOH (0.5 ml) is treated with 4M HCl in dioxane (0.5 ml). After stirring at room temperature for 1 hour, the solvent is removed *in vacuo* to afford the title compound which is used crude in the next step.

Step 3: 9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-{3-[1-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-ylcarbamoyl)-piperidin-4-yl]-ureido}-ethyl) amide trifluoroacetate

The title compound is prepared analogously to 9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester (Example 48 step 7) by replacing 9-((1R,2S,3R,4S)-4-amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid methyl ester hydrochloride with 9-((1R,2S,3R,4S)-4-amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-{3-[1-(3,4,5,6-tetrahydro-2H-

[1,2']bipyridinyl-4-ylcarbamoyl)-piperidin-4-yl]-ureido}-ethyl)-amide dihydrochloride.
(MH+ 451.8)

Example 102-104

These compounds namely,

{1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-piperidin-4-yl}-carbamic acid tert-butyl ester trifluoroacetate (MH+ 685.6) (Example 102),

N-{(1S,2R,3S,4R)-4-[2-(1-benzyl-piperidin-4-ylamino)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide trifluoroacetate (MH+ 675.5) (Example 103) and N-{(1S,2R,3S,4R)-4-[2-[1,4]diazepan-1-yl-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide trifluoroacetate (MH+585.5) (Example 104)

are prepared analogously to N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-benzyloxy-acetamide trifluoroacetate Example 1 (Step 2) by replacing 2-benzyloxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-acetamide with N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Intermediate J) and by replacing (3R)-3-aminopyrrolidine with the appropriate amine.

Example 105 and 106

These compounds namely,

{(R)-1-[9-[(1R,2S,3R,4S)-4-(cyclobutanecarbonyl-amino)-2,3-dihydroxy-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester trifluoroacetate (MH+ 697.3) (Example 105) and

(S)-3-[9-[(1R,2S,3R,4S)-4-(cyclobutanecarbonyl-amino)-2,3-dihydroxy-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-ylamino]-pyrrolidine-1-carboxylic acid tert-butyl ester trifluoroacetate (MH+697.3) (Example 106)

are prepared analogously to N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-benzyloxy-

acetamide trifluoroacetate Example 1 (Step 2) by replacing 2-benzyloxy-N-
 {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-
 cyclopentyl} -acetamide with cyclobutanecarboxylic acid {(1S,2R,3S,4R)-4-[2-chloro-6-
 (2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -amide (Intermediate
 K) and by replacing (3R)-3-aminopyrrolidine with the appropriate amine.

Example 107

**N-{(1S,2R,3S,4R)-4-[6-(2,2-diphenyl-ethylamino)-2-(3,4,5,6-tetrahydro-2H-
 [1,2']bipyridinyl-4-ylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide
 trifluoroacetate**

This compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-
 pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-
 benzyloxy-acetamide trifluoroacetate Example 1 (Step 2) by replacing 2-benzyloxy-N-
 {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-
 cyclopentyl} -acetamide with N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-
 ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide (Intermediate J) and
 by replacing (3R)-3-aminopyrrolidine with 3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-
 ylamine. (MH⁺ 331.7)

Example 108

**4-[3-(2-{[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-
 -6-(2,2-diphenyl-ethylamino)-9H-purine-2-carbonyl]-amino}-ethyl)-ureido]-
 piperidine-1-carboxylic acid benzyl ester**

To a solution of 9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-
 diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-amino-ethyl)-amide (Example 48
 step 8) (0.1 g, 174 mmol) in chloroform (5 ml) is added 4-isocyanato-Z-piperidine (0.045
 g, 0.174 mmol) in chloroform (5 ml). The reaction mixture is allowed to stir at room
 temperature overnight and then methanol is added to quench any residual isocyanate.
 The solvent is removed *in vacuo* to yield the title compound. (MH⁺833.5)

Example 109

9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid {2-[3-(5-methyl-3-phenyl-isoxazol-4-yl)-ureido]-ethyl}-amide trifluoroacetate

This compound is prepared analogously to 4-[3-(2-{[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carbonyl]-amino}-ethyl)-ureido]-piperidine-1-carboxylic acid benzyl ester (Example 108) by replacing 4-isocyanato-Z-piperidine with 4-isocyanato-5-methyl-3-phenyl-isoxazole. (MH+773.5)

Examples 110-112

These compounds namely,

N-{(1S,2R,3S,4R)-4-[2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide trifluoroacetate (MH+ 599.5) (Example 110),

N-{(1S,2R,3S,4R)-4-[6-(2,2-diphenyl-ethylamino)-2-((R)-3-methylamino-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide trifluoroacetate (MH+ 585.4) (Example 111) and

(4-{2-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-ylamino]-ethyl}-imidazol-1-yl)-acetic acid trifluoroacetate (MH+ 654.3) (Example 112)

are prepared analogously to N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-benzyloxy-acetamide trifluoroacetate Example 1 (Step 2) by replacing 2-benzyloxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-acetamide with N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Intermediate J) and by replacing (3R)-3-aminopyrrolidine with the appropriate amine.

Example 113

9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-{3-[1-(4-methoxy-phenylcarbamoyl)-piperidin-4-yl]-ureido}-ethyl)-amide trifluoroacetate

Step 1: 9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid [2-(3-piperidin-4-yl-ureido)-ethyl]-amide

A solution of 4-[3-(2-{[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carbonyl]-amino}-ethyl)-ureido]-piperidine-1-carboxylic acid benzyl ester (Example 108) (0.145 g, 0.174 mmol) in methanol (1 ml) under an atmosphere of Argon is treated with palladium hydroxide on carbon (0.054 g, 20%w/w carbon). The reaction mixture is placed under an atmosphere of hydrogen and stirred at room temperature for 72 hours and then filtered. The filtrate is concentrated *in vacuo* to yield the title compound as a green oil.

Step 2: 9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-{3-[1-(4-methoxy-phenylcarbamoyl)-piperidin-4-yl]-ureido}-ethyl)-amide trifluoroacetate

This compound is prepared analogously to 4-[3-(2-{[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carbonyl]-amino}-ethyl)-ureido]-piperidine-1-carboxylic acid benzyl ester (Example 108) by replacing 9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-amino-ethyl)-amide with 9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid [2-(3-piperidin-4-yl-ureido)-ethyl]-amide and by replacing 4-isocyanato-Z-piperidine with 1-isocyanato-4-methoxy-benzene. (MH+848.6)

Example 114-115

These compounds namely,

9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-{3-[1-(4-cyano-phenylcarbamoyl)-piperidin-4-yl]-ureido}-ethyl)-amide trifluoroacetate (MH+843.6) (Example 114) and ((4-[3-(2-{[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carbonyl]-amino}-ethyl)-ureido]-piperidine-1-carbonyl]-amino)-acetic acid ethyl ester trifluoroacetate (MH+ 828.6) (Example 115) are prepared analogously to Example 113 by replacing 1-isocyanato-4-methoxy-benzene with the appropriate isocyanate.

Example 116

9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid [2-(3-{1-[1-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-ylcarbamoyl)-piperidin-4-yl]-ureido)-ethyl]-amide trifluoroacetate

The title compound is prepared analogously to 9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-{3-[1-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-ylcarbamoyl)-piperidin-4-yl]-ureido}-ethyl) amide trifluoroacetate (Example 101) by replacing 4-[(imidazole-1-carbonyl)-amino]-piperidine-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amidewith Intermediate I. (MH+514.9)

Examples 117- 125

These compounds namely,

N-((1S,2R,3S,4R)-4-[6-[2,2-bis-(4-methoxy-phenyl)-ethylamino]-2-((R)-3-dimethylamino-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide trifluoroacetate (MH+ 659.4) (Example 117),

N-((1S,2R,3S,4R)-4-{2-((R)-3-Dimethylamino-pyrrolidin-1-yl)-6-[2-(4-fluoro-phenyl)-2-phenyl-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate (MH+ 617.4)(Example 118),

N-{(1S,2R,3S,4R)-4-[2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-(1-ethyl-propylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide trifluoroacetate (MH+ 489.3)

(Example 119),

N-((1S,2R,3S,4R)-4-{2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-[(9H-fluoren-9-ylmethyl)-amino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate (MH+ 597.4) (Example 120),

N-{(1S,2R,3S,4R)-4-[2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-((S)-1-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide trifluoroacetate (MH+ 553.4) (Example 121),

N-((1S,2R,3S,4R)-4-{2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-[(naphthalen-1-ylmethyl)-amino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate (MH+ 559.4) (Example 122),

N-{(1S,2R,3S,4R)-4-[6-[(2'-cyano-biphenyl-4-ylmethyl)-amino]-2-((R)-3-dimethylamino-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide trifluoroacetate (MH+ 610.3) (Example 123),

N-{(1S,2R,3S,4R)-4-[2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-(3,3-dimethyl-butylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide trifluoroacetate (MH+ 503.4) (Example 124) and

N-((1S,2R,3S,4R)-4-{2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-[2-(4-sulfamoyl-phenyl)-ethylamino]-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate (MH+ 602.3) (Example 125)

are prepared analogously to N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-benzyloxy-acetamide trifluoroacetate (Example 1 step 2) by replacing (3R)-3-aminopyrrolidine with dimethyl-(R)-pyrrolidin-3-yl-amine and by replacing 2-benzyloxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-acetamide with the appropriate starting compounds. The preparation of the starting compounds are either described herein or can be prepared from [(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionyl-carbamic acid tert-butyl ester (Intermediate G) and the appropriate amine using a procedure analogous to 2R,3S,4R)-4-

{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl} -2,3-dihydroxy-cyclopentyl)-propionamide (Example 22 steps 1 and 2)

Example 126

N-[(1S,2R,3S,4R)-4-[2-[(R)-3-(2-acetylamino-4-methyl-thiazole-5-sulfonylamino)-pyrrolidin-1-yl]-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate

N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide (Example 18 step 2) (0.023 g, 40 μ mol) is treated with a solution of 2-acetamido-4-methyl-5-thiazolesulfonyl chloride (0.0104 g, 39 μ mol) in NMP (0.5 ml). After stirring at room temperature overnight purification is carried out using mass directed preparative LC-MS eluting with acetonitrile: water: trifluoroacetic acid to afford the title compound. (MH⁺ 789.4)

Examples 127-132

These compounds namely,

N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-[(R)-3-[3-(3-phenyl-isoxazol-4-yl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (MH⁺ 771.5) (Example 127),

5-pyridin-2-yl-thiophene-2-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (MH⁺ 758.4) (Example 128),

4-[(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-ylcarbamoyl]-piperidine-1-carboxylic acid benzylester trifluoroacetate (MH⁺ 816.6) (Example 129),

N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-[(R)-3-[3-(3-methyl-5-phenyl-isoxazol-4-yl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (MH⁺ 771.5) (Example 130),

1,3-dimethyl-1H-thieno[2,3-c]pyrazole-5-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (MH⁺ 749.4) (Example 131),
 1-methyl-1H-benzotriazole-5-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (MH⁺ 730.5) (Example 132),
 are prepared analogously to Example 126 by replacing 2-acetamido-4-methyl-5-thiazolesulfonyl chloride with the appropriate isocyanate or acid chloride. Reactions using acid chlorides also have triethylamine added.

Example 133

N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-formamide hydrochloride

Step 1: ((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid benzyl ester trifluoroacetate

A solution comprising {(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-carbamic acid benzyl ester (Intermediate L) (0.1 g, 0.15 mmol), pyridine-3-isocyanate (0.02 g, 0.17 mmol) and TEA (0.017 g, 0.17 mmol) in THF (2 ml) is stirred at room temperature overnight. The solvent is removed *in vacuo* and purification is carried out by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% TFA). The fractions are collected and the MeCN is removed *in vacuo*. The remaining aqueous portion is basified with saturated sodium bicarbonate solution and extracted with DCM. The combined organic extracted are dried (MgSO₄) and concentrated *in vacuo* to afford the title product. MS (ES⁺) *m/e* 769 (MH⁺).

Step 2: 1-[(R)-1-[9-((1R,2S,3R,4S)-4-Amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl]-3-pyridin-3-yl-urea

To a solution of ((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid benzyl

ester trifluoroacetate (35 mg, 46 μ mol) in ethanol (1 ml) under an inert atmosphere of Argon is added 10% palladium on carbon (10 mg). The reaction mixture is purged with Argon and placed under a positive atmosphere of hydrogen overnight after which time, the mixture is filtered through celite and the catalyst washed with ethanol. The organic portions are combined and concentrated *in vacuo* to yield the title compound. MS (ES+) *m/e* 635 (MH^+).

Step 3: N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-formamide hydrochloride

Acetic anhydride (1.9 mg, 19 mmol) and formic acid (1.4 mg, 30 mmol) are stirred together for 30 minutes and then added to a solution of 1-{(R)-1-[9-((1R,2S,3R,4S)-4-amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-3-pyridin-3-yl-urea (11 mg, 17 μ mol) in THF (0.5 ml). The reaction mixture is stirred at room temperature overnight and purification by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% HCl) affords the title compound. (MH^+ 663.5)

Example 134

(R)-3-Dimethylamino-pyrrolidine-1-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate Step 1: Imidazole-1-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide:

A mixture comprising N-{(3aR,4S,6R,6aS)-6-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,2-dimethyl-tetrahydro-cyclopenta[1,3]dioxol-4-yl}-propionamide (Intermediate M) (0.24 g, 394 μ mol) and CDI (0.275 g, 1.7 mmol) in dry DCM (6 ml) is stirred at room temperature for 3 hours. The resulting solution is purified by chromatography on silica eluting with 100% DCM changing to 5% MeOH in DCM to afford the title compound as a yellow oil. The oil consists of the imidazole-urea

intermediate together with variable amounts of the corresponding isocyanate and imidazole which are equally suitable as precursors to ureas.

Step 2: (R)-3-Dimethylamino-pyrrolidine-1-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate:

Dimethyl-(R)-pyrrolidin-3-yl-amine (4.6 mg, 40 μ mol) is treated with a solution of imidazole-1-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide (25 mg, 40 μ mol) in DCM (1 ml) and the reaction mixture is stirred at room temperature overnight. The solvent is removed *in vacuo* and the residue is treated with TFA (0.5 ml) and water (0.5 ml). After stirring at room temperature for 3 hours, the reaction mixture is concentrated *in vacuo* and the resulting crude is purified by mass directed preparative LC-MS eluting with acetonitrile: water: trifluoroacetic acid to afford the title compound. (MH⁺ 711.5)

Example 135

4-(2-cyano-ethyl)-piperazine-1-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate

This compound is prepared analogously to (R)-3-dimethylamino-pyrrolidine-1-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (Example 134) by replacing dimethyl-(R)-pyrrolidin-3-yl-amine with the appropriate amine. (MH⁺ 697.4)

Example 136 -155

These compounds namely,

N-[(1S,2R,3S,4R)-4-(2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-{[(1R,3S)-2,2-dimethyl-3-(2-methyl-propenyl)-cyclopropylmethyl]-amino}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate (MH⁺ 555.6) (Example 136),

N- {(1S,2R,3S,4R)-4-[6-((1R,2R)-2-benzyloxy-cyclopentylamino)-2-((R)-3-dimethylamino-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 593.5) (Example 137),

N- {(1S,2R,3S,4R)-4-[6-((1S,2S)-2-benzyloxy-cyclopentylamino)-2-((R)-3-dimethylamino-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 593.5) (Example 138),

N- {(1S,2R,3S,4R)-4-[6-((1S,2S)-bicyclopentyl-2-ylamino)-2-((R)-3-dimethylamino-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 555.6) (Example 139),

N- {(1S,2R,3S,4R)-4-[6-((R)-1-benzyl-pyrrolidin-3-ylamino)-2-((R)-3-dimethylamino-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 578.5) (Example 140),

N- {(1S,2R,3S,4R)-4-[6-((S)-1-benzyl-pyrrolidin-3-ylamino)-2-((R)-3-dimethylamino-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 578.5) (Example 141),

N- {(1S,2R,3S,4R)-4-[2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-(2-piperidin-1-yl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 530.5) (Example 142),

N- {(1S,2R,3S,4R)-4-[6-[2-(4-benzyl-piperidin-1-yl)-ethylamino]-2-((R)-3-dimethylamino-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 620.5) (Example 143),

N- {(1S,2R,3S,4R)-4-[2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-((S)-2-phenyl-1-pyrrolidin-1-ylmethyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 606.5) (Example 144),

N- {(1S,2R,3S,4R)-4-[2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-ylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 579.5) (Example 145),

N- {(1S,2R,3S,4R)-4-[2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-((S)-1-hydroxymethyl-3-methyl-butylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 519.5) (Example 146),

N- {(1S,2R,3S,4R)-4-[2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-((S)-1-hydroxymethyl-3-methylsulfanyl-propylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 537.5) (Example 147),

N- {(1S,2R,3S,4R)-4-[6-((R)-1-benzyl-2-hydroxy-ethylamino)-2-((R)-3-dimethylamino-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 553.5) (Example 148),

N- {(1S,2R,3S,4R)-4-[2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-((1S,2S)-2-hydroxy-1-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 569.5) (Example 149),

N-((1S,2R,3S,4R)-4- {2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-[(S)-2-hydroxy-1-(4-hydroxy-benzyl)-ethylamino]-purin-9-yl} -2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate (MH+ 569.5) (Example 150),

N-((1S,2R,3S,4R)-4- {2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-[(S)-2-hydroxy-1-(1H-imidazol-4-ylmethyl)-ethylamino]-purin-9-yl} -2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate (MH+ 543.5) (Example 151),

N-((1S,2R,3S,4R)-4- {2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-[(S)-2-hydroxy-1-(1H-indol-3-ylmethyl)-ethylamino]-purin-9-yl} -2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate (MH+ 592.4) (Example 152),

N- {(1S,2R,3S,4R)-4-[6-((S)-1-benzyl-2-methoxy-ethylamino)-2-((R)-3-dimethylamino-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 567.4) (Example 153),

N- {(1S,2R,3S,4R)-4-[6-[(S)-1-(4-benzyloxy-benzyl)-2-hydroxy-ethylamino]-2-((R)-3-dimethylamino-pyrrolidin-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide trifluoroacetate (MH+ 659.4) (Example 154) and

N-((1S,2R,3S,4R)-4- {2-((R)-3-dimethylamino-pyrrolidin-1-yl)-6-[(1S,2S)-2-hydroxy-1-hydroxymethyl-2-(4-methylsulfanyl-phenyl)-ethylamino]-purin-9-yl} -2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate (MH+ 615) (Example 155)

are prepared analogously to N- {(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-benzyloxy-acetamide trifluoroacetate (Example 1 step 2) by replacing (3R)-3-aminopyrrolidine with dimethyl-(R)-pyrrolidin-3-yl-amine and by replacing 2-benzyloxy-N- {(1S,2R,3S,4R)-4-

[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} - acetamide with the appropriate starting compounds. The preparation of the starting compounds are either described herein or can be prepared from [(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionyl-carbamic acid tert-butyl ester (Intermediate G) and the appropriate amine using a procedure analogous to 2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl} -2,3-dihydroxy-cyclopentyl)-propionamide (Example 22 steps 1 and 2).

Example 156

N-{(R)-1-[6-[2,2-Bis-(4-methoxy-phenyl)-ethylamino]-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-pyrrolidin-3-yl}-isonicotinamide hydrochloride

This compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-benzyloxy-acetamide trifluoroacetate (Example 1 step 2) by replacing (3R)-3-aminopyrrolidine with (R)-N-pyrrolidin-3-yl-isonicotinamide (Intermediate O) and by replacing 2-benzyloxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -acetamide with N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-methoxy-phenyl)-ethylamino]-2-chloro-purin-9-yl} -2,3-dihydroxy-cyclopentyl)-propionamide {prepared from [(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionyl-carbamic acid tert-butyl ester (Intermediate G) and the appropriate amine using a procedure analogous to 2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl} -2,3-dihydroxy-cyclopentyl)-propionamide (Example 22 steps 1 and 2)}. (MH⁺ 736.5)

Example 157

5-Methyl-isoxazole-3-carboxylic acid {(R)-1-[6-[2,2-bis-(4-methoxy-phenyl)-ethylamino]-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide hydrochloride

are prepared analogously to Example 93 by replacing N-{(1S,2R,3S,4R)-4-[2-chloro-6-(1-ethyl-propylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide with N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-methoxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide [prepared from [(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionyl-carbamic acid tert-butyl ester (Intermediate G) and the appropriate amine using a procedure analogous to 2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 22 steps 1 and 2)]. (MH⁺ 740.5)

Example 158

N-{(R)-1-[6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-pyrrolidin-3-yl}-isonicotinamide hydrochloride

A mixture comprising N-((1S,2R,3S,4R)-4-{2-((R)-3-amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 22 step 4) (20 mg, 33 μ mol) in THF (0.5 ml) and NMP (0.5 ml) is treated with TEA (13 mg, 0.13 mmol) followed by isonicotinoyl chloride hydrochloride (16 mg, 83 μ mol). After stirring at room temperature for 2 hours, the solvent is removed *in vacuo* and purification by C-18 reverse phase column chromatography eluting with acetonitrile : water : HCl (0.1%) (gradient of 0 to 100% acetonitrile) yields the title compound. (MH⁺ 708.4)

Example 159

1-Methyl-1H-benzotriazole-5-carboxylic acid {(R)-1-[6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide hydrochloride

This compound is prepared analogously to N-{(R)-1-[6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-pyrrolidin-3-yl}-isonicotinamide hydrochloride (Example 158) by replacing

isonicotinoyl chloride hydrochloride with 1-methyl-1H-1,2,3-benzotriazole-5-carbonyl chloride. (MH+ 762.4)

Example 160-162

These compounds namely,

N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-3-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide hydrochloride (MH+ 737.5) (Example 160),

N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-4-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide hydrochloride (MH+ 737.2) (Example 161),

N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-4-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide hydrochloride (MH+ 740.5) (Example 162),

are prepared analogously to N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide hydrochloride (Example 23) by replacing 2-(aminomethyl)pyridine with the appropriate amine.

Example 163

2-Amino-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-acetamide hydrochloride

A solution of 1-{(R)-1-[9-((1R,2S,3R,4S)-4-amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-3-pyridin-3-yl-urea (Example 26 step 2) (17 mg, 26 μ mol) in THF (1 ml) is treated with tert-butoxycarbonylamino-acetic acid 2,5-dioxo-pyrrolidin-1-yl ester (9 mg, 29 μ mol) and stirred at room temperature overnight. The resulting solution is treated with 1.25 M HCl in EtOH (1 ml) and stirred at room temperature for 2 days. Purification by C-18 reverse phase column chromatography eluting with acetonitrile : water : HCl (0.1%) (gradient of 0 to 100% acetonitrile) yields the title compound. (MH+ 691.99)

Example 164**N-{(1S,2R,3S,4R)-4-[(R)-2-[1,3']Bipyrrolidinyl-1'-yl-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide hydrochloride**

The title compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-benzyloxy-acetamide trifluoroacetate (Example 1 step 2) by replacing 2-benzyloxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-acetamide with N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Intermediate J) and by replacing (3R)-3-aminopyrrolidine with (R)-[1,3']bipyrrolidinyl (Intermediate N). (MH⁺ 625.4)

Example 165**N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{2-[3-(3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-ureido]-ethylamino}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate**

A solution of N-{(1S,2R,3S,4R)-4-[2-(2-amino-ethylamino)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide trifluoroacetate (Example 100) (27 mg, 37 µmol) in IPA (0.5 ml) is treated with imidazole-1-carboxylic acid (3,4,5,6-tetrahydro-2H-[1,2']bipyridinyl-4-yl)-amide (Intermediate C) (1.1 ml of a 10 mg/ml solution in DCM, 40 µmol). After stirring at room temperature for 7 days, the solvent is removed *in vacuo* and purification of the residue by C-18 reverse phase column chromatography eluting with acetonitrile : water : TFA (0.1%) (gradient of 0 to 100% acetonitrile) yields the title compound. (MH⁺ 748.6)

Example 166**Biphenyl-2-yl-carbamic acid 1-{2-[(R)-3-(3-{(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-**

pyrrolidin-3-yl}-ureido)-pyrrolidin-1-yl]-2-oxo-ethyl}- piperidin-4-yl ester trifluoroacetate

This compound is prepared analogously to (R)-3-dimethylamino-pyrrolidine-1-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (Example 134) by replacing dimethyl-(R)-pyrrolidin-3-yl-amine with Intermediate Z (MH+/2 510.62)

Example 167

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(2-methyl-5-phenyl-furan-3-yl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide

This compound is prepared analogously to (Example 126) by replacing 2-acetamido-4-methyl-5-thiazolesulfonyl chloride with 3-isocyanato-2-methyl-5-phenyl-furan and by changing the solvent to THF. (MH+ 770.48)

Example 168

1-[6-Amino-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid ethyl ester

Step 1: N-[(1S,2R,3S,4R)-4-(6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-2-chloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide

To a solution of N-[(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide (Intermediate J7) (2.6 g, 7.22 mmol) in dry THF (26 ml) is added C,C-Bis-(4-methoxy-phenyl)-methylamine (Intermediate Y) (3.5 g, 14.44 mmol). The mixture is heated at 50°C for 12 hours and then concentrated *in vacuo*. The residue is dissolved in chloroform and washed sequentially with 1.5N HCl, water and saturated aqueous brine solution. The organic phase is dried over anhydrous sodium sulphate and concentrated *in vacuo*. The crude product is purified by chromatography on silica (60 – 120 mesh) eluting with 2% methanol in chloroform to afford the title product.

LC-MS (0.1% formic acid, acetonitrile): (MH+ 567)

Step 2: N-[(1S,2R,3S,4R)-4-(6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-2-hydrazino-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide

A mixture comprising N-[(1S,2R,3S,4R)-4-(6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-2-chloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide (1.6 g, 2.82 mmol), and hydrazine monohydrate (14 ml) is stirred at room temperature for 72 h. Then isopropyl alcohol (10 ml) is added and the solvent was decanted off to afford a gummy mixture. It is dissolved in water (10 ml) and stirred for 12 h. The fine solid obtained is filtered, washed with water and dried *in vacuo* to afford the title product which is used in the next step without further purification. LC-MS (0.1% formic acid, acetonitrile): (MH+ 563).

Step 3: 1-[6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid ethyl ester:

To a solution of N-[(1S,2R,3S,4R)-4-(6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-2-hydrazino-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide (0.1 g, 0.177 mmol) in dry ethyl alcohol (5 ml) is added 2-formyl-3-oxo-propionic acid ethyl ester (synthesised from ethyl-3,3-diethoxy-propionate, as described in: Bertz S.H., Dabbagh G. and Cotte P.; J. Org. Chem. (1982) 47, pp 2216-2217) (0.033 g, 0.231 mmol). The reaction mixture is heated at reflux for 8 hours then concentrated *in vacuo*. The crude residue is purified by chromatography on silica (60 – 120 mesh) eluting with 3% methanol in chloroform to afford the title compound. LC-MS (0.1% formic acid, acetonitrile): (MH+ 671).

Step 4: 1-[6-Amino-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid ethyl ester:

A cooled (0°C) solution of 1-[6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid ethyl ester (0.1 g, 0.149 mmol) in dry dichloromethane (4 ml) is treated dropwise with TFA (2 ml). The mixture is stirred at room temperature for 12 h and then concentrated *in vacuo*. The residue is co-evaporated with chloroform three

times to remove excess trifluoro acetic acid and purification of the residue by preparative HPLC affords the title compound.

LC-MS (0.1% formic acid, acetonitrile): (MH⁺ 445.3).

Example 169

Isoxazole-5-carboxylic acid {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-amide

This compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Intermediate JJ4) by replacing propionyl chloride with isoxazole-5-carbonyl chloride. (MH⁺ 560.28)

Example 170

N-{(1S,2R,3S,4R)-4-[6-Amino-2-(4-quinolin-4-yl-pyrazol-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide

This compound is prepared analogously to Example 168 by replacing 2-formyl-3-oxopropionic acid ethyl ester with 2-(4-quinolyl)-malaldehyde. (MH⁺ 500.3)

Example 171

N-{(1S,2R,3S,4R)-4-[6-(2,2-Diphenyl-ethylamino)-2-(4-pyridin-2-yl-pyrazol-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide

This compound is prepared analogously to Example 42 by replacing cyclohexane carboxaldehyde with 2-pyridinyl-propanedial. The reaction is carried out in ethanol. (MH⁺ 630.40)

Example 172

N-{(1S,2R,3S,4R)-4-[6-(2,2-Diphenyl-ethylamino)-2-(4-pyridin-4-yl-pyrazol-1-yl)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide

This compound is prepared analogously to Example 42 by replacing cyclohexane carboxaldehyde with 4-pyridinyl-propanedial. The reaction is carried out in ethanol. (MH⁺ 630.41)

Example 173**1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid methylamide**

Step 1: 1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid ethyl ester

This compound is prepared analogously to 1-[6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid ethyl ester (Example 168 step 3) by replacing N-[(1S,2R,3S,4R)-4-(6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-2-hydrazino-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide with N-{(1S,2R,3S,4R)-4-[6-(2,2-diphenyl-ethylamino)-2-hydrazino-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide.

Step 2: 1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid methylamide

A mixture of 1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid ethyl ester (0.07 g, 0.112 mmol) and 40 % aqueous methyl amine solution (3 ml) is heated to 65 °C for 12h. The reaction mixture is concentrated *in vacuo* and purification of the crude residue by chromatography on silica eluting with 4% methanol in chloroform yields the title compound.

LC-MS (0.1% formic acid, acetonitrile): (MH⁺ 610.41)

Example 174**1-[6-Amino-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid methylamide**

Step 1: 1-[6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid methylamide:

A mixture of 1-[6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid ethyl ester (Example 168, step 3) (0.2 g, 0.298 mmol) and 40 % aqueous methyl amine solution (5 ml) is heated to 65 °C for 12h. The reaction mixture concentrated *in vacuo* and purification of the crude residue by chromatography on silica eluting with 3% methanol in chloroform yields the title compound. LC-MS (0.1% formic acid, acetonitrile): 656 (MH⁺).

Step 2: 1-[6-Amino-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid methylamide

A cooled (0°C) solution of 1-[6-{[Bis-(4-methoxy-phenyl)-methyl]-amino}-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid methylamide (0.08 g, 0.122 mmol) in dry dichloromethane (4 ml) is treated slowly with trifluoroacetic acid (2 ml). The reaction mixture is stirred at room temperature for 48 h and then concentrated *in vacuo*. The residue was co-evaporated with chloroform three times to remove excess trifluoro acetic acid and purification of the crude product by preparative HPLC affords the title compound. LC-MS (0.1% formic acid, acetonitrile): (MH⁺ 430.28)

Example 175

N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide trifluoroacetate:

This compound is prepared analogously to Example 23 by replacing {(R)-1-[6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid phenyl ester with N-((1S,2R,3S,4R)-4-{2-((R)-3-amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide hydrochloride (Example 7 step 5) and by replacing 2-(aminomethyl)pyridine with pyridin-3-yl-carbamic acid phenyl ester (Intermediate ZB). (MH⁺ 725.32)

Example 176**4-[(R)-3-(3-[(R)-1-[9-[(1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl)-ureido)-pyrrolidine-1-carbonyl]-amino}-piperidine-1-carboxylic acid benzyl ester trifluoroacetate**

The title compound is prepared analogously to 4-[3-(2-{[9-[(1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carbonyl]-amino}-ethyl)-ureido]-piperidine-1-carboxylic acid benzyl ester (Example 108) by replacing 9-[(1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purine-2-carboxylic acid (2-amino-ethyl)-amide with N-[(1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-[(R)-3-pyrrolidin-3-ylureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate (Example 17 step 1). (MH⁺ 943.4)

Example 177**N-[(1S,2R,3S,4R)-4-{2-[(S)-1-benzyl-pyrrolidin-3-yl)-methyl-amino]-6-[2-(4-fluorophenyl)-2-phenyl-ethylamino]-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate**

This compound is prepared analogously to N-[(1S,2R,3S,4R)-4-[2-[(R)-3-amino-pyrrolidin-1-yl]-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-2-benzyloxy-acetamide trifluoroacetate (Example 1 step 2) by replacing (3R)-3-(Boc-amino)pyrrolidine with ((S)-1-benzyl-pyrrolidin-3-yl)-methyl-amine and by replacing 2-benzyloxy-N-[(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-acetamide with N-[(1S,2R,3S,4R)-4-{2-chloro-6-[2-(4-fluorophenyl)-2-phenyl-ethylamino]-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-propionamide trifluoroacetate [prepared from [(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionyl-carbamic acid tert-butyl ester (Intermediate G) and the appropriate amine using a procedure analogous to 2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-chloro-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-propionamide (Example 22 steps 1 and 2)]. (MH⁺ 693.5)

Example 178**N-[(1S,2R,3S,4R)-4-[2-[(R)-3-[3-(4-Benzyloxy-phenyl)-ureido]-pyrrolidin-1-yl]-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide trifluoroacetate**

This compound is prepared analogously to Example 126 by replacing 2-acetamido-4-methyl-5-thiazolesulfonyl chloride with 1-benzyloxy-4-isocyanato-benzene. (MH+ 796.49)

Example 179-180

These compounds namely,

(R)-3-amino-pyrrolidine-1-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (MH+ 677.5) (Example 179) and

(S)-3-amino-pyrrolidine-1-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (MH+ 683.4) (Example 180)

are prepared analogously to (R)-3-dimethylamino-pyrrolidine-1-carboxylic acid {(R)-1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-amide trifluoroacetate (Example 134) by replacing dimethyl-(R)-pyrrolidin-3-yl-amine with the appropriate amine.

Example 181**(R)-N-[(1S,2R,3S,4R)-4-[6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl)-ureido]-pyrrolidin-1-yl]-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-2-hydroxy-propionamide**

Step 1: (R)-2-Benzyloxy-N-[(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide

The title compound is prepared by dissolving (R)-2-benzyloxy-propionic acid (1 equivalent) in dichloromethane with 1,3-dicyclohexylcarbodiimide (1 equivalent) and a catalytic amount of 4-dimethylaminopyridine, stirring for five minutes, then adding (1S,2R,3S,5R)-3-amino-5-(2,6-dichloro-purin-9-yl)-cyclopentane-1,2-diol (Intermediate

J5; 1-equivalent) in dichloromethane. The reaction is stirred at room temperature until determined to be complete, the solvent is removed under reduced pressure, and the title compound purified by column chromatography / crystallisation.

Step 2: (R)-2-Benzoyloxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide

The title compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Intermediate J7), by substituting acetic acid (R)-2-benzoyloxy-N-[(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide for N-[(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide.

Step 3: {(R)-1-[9-[(1R,2S,3R,4S)-4-((R)-2-Benzoyloxy-propionylamino)-2,3-dihydroxy-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester

A suspension of (R)-2-benzoyloxy-N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (1 equivalent) and (3R)-(+)-3-(Boc-amino)pyrrolidine (4 equivalents) in acetonitrile is treated with a catalytic amount of sodium iodide and then heated using microwave radiation in a Personal Chemistry Emrys™ Optimizer microwave reactor at 160°C. After 1 hour, the solvent is removed *in vacuo*; purification by column chromatography / crystallisation affords the title compound.

Step 4: (R)-N-{(1S,2R,3S,4R)-4-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-benzoyloxy-propionamide

A solution of {(R)-1-[9-[(1R,2S,3R,4S)-4-((R)-2-Benzoyloxy-propionylamino)-2,3-dihydroxy-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-carbamic acid tert-butyl ester in MeOH (~0.5M) is treated with an equal volume of 4M HCl in dioxane and stirred at room temperature for 2 hours. The solvent is removed *in vacuo* and purification is carried out by column chromatography / crystallisation to afford the title compound.

Step 5: (R)-2-Benzoyloxy-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide.

A solution comprising (R)-N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-benzoyloxy-propionamide (1 equivalent) and pyridin-3-yl-carbamic acid phenyl ester (Intermediate ZB) (1 equivalent) in NMP is stirred at 100°C for 1 hour. The solvent is removed *in vacuo* and the title compound is obtained after purification by column chromatography / crystallisation.

Step 6: (R)-N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-propionamide

To a stirred solution of (R)-2-Benzoyloxy-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide in ethanol is added 10 equivalents of ammonium formate and 20mol% of 10% palladium on carbon. The mixture is stirred at 80°C for five hours, allowed to cool and filtered through Celite™. Removal of the solvent under reduced pressure affords the title compound.

Example182

(S)-N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-propionamide

Step 1: Acetic acid (S)-1-[(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentylcarbamoyl]-ethyl ester

The title compound is prepared analogously to Intermediate J6, from Intermediate J5, replacing propionyl chloride with acetic acid (S)-1-chlorocarbonyl-ethyl ester.

Step 2: Acetic acid (S)-1-{{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl} -ethyl ester

The title compound is prepared analogously to N-{{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide (Intermediate J7), by substituting acetic acid (S)-1-[(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentylcarbamoyl]-ethyl ester (Example 182, step 1) for N-[(1S,2R,3S,4R)-4-(2,6-dichloro-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide.

Step 3: Acetic acid (S)-1-{{(1S,2R,3S,4R)-4-[2-((R)-3-tert-butoxycarbonylamino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl} -ethyl ester.

The title compound is prepared analogously to {(R)-1-[9-[(1R,2S,3R,4S)-4-((R)-2-benzyloxy-propionylamino)-2,3-dihydroxy-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl} -carbamic acid tert-butyl ester (Example 181, step 3), by substituting acetic acid (S)-1-{{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl} -ethyl ester (Example 182, step 2) for (R)-2-benzyloxy-N-{{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -propionamide (Example 181, step 2).

Step 4: (S)-N-{{(1S,2R,3S,4R)-4-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-propionamide

The title compound is prepared analogously to (R)-N-{{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-benzyloxy-propionamide (Example 181, step 4), by substituting acetic acid (S)-1-{{(1S,2R,3S,4R)-4-[2-((R)-3-tert-butoxycarbonylamino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl} -ethyl ester (Example 182, step 3) for {(R)-1-[9-[(1R,2S,3R,4S)-4-((R)-2-benzyloxy-propionylamino)-2,3-dihydroxy-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl} -carbamic acid tert-butyl ester (Example 181, step 3).

Step 5: (S)-N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-propionamide.

The title compound was prepared analogously to (R)-2-benzyloxy-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 181, step 5), by substituting (S)-N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-hydroxy-propionamide (Example 182, step 4) for (R)-N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-benzyloxy-propionamide (Example 181, step 4).

Example 183

N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide

The title compound is prepared analogously to (R)-2-benzyloxy-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 181, step 5), by substituting N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Intermediate ZG) for (R)-N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-benzyloxy-propionamide (Example 181, step 4).

Example 184

N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide

The title compound was prepared analogously to (R)-2-benzyloxy-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 181, step 5), by substituting N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-

yl]-2,3-dihydroxy-cyclopentyl} -3-hydroxy-propionamide (Intermediate ZG) for (R)-N-
{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-
9-yl]-2,3-dihydroxy-cyclopentyl} -2-benzyloxy-propionamide (Example 181, step 4) and
pyridin-2-ylmethyl-carbamic acid (Intermediate ZC) for pyridin-3-yl-carbamic acid
phenyl ester (Intermediate ZB).

Example 185

**N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-[3-(3-hydroxy-benzyl)-
ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-
propionamide**

The title compound was prepared analogously to (R)-2-benzyloxy-N-((1S,2R,3S,4R)-4-
{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-
yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 181, step 5), by substituting N-
{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-
yl]-2,3-dihydroxy-cyclopentyl} -3-hydroxy-propionamide (Intermediate ZG) for (R)-N-
{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-
9-yl]-2,3-dihydroxy-cyclopentyl} -2-benzyloxy-propionamide (Example 181, step 4) and
(3-Hydroxy-benzyl)-carbamic acid phenyl ester (Intermediate ZD) for pyridin-3-yl-
carbamic acid phenyl ester (Intermediate ZB).

Example 186

**N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-[3-(4-sulfamoyl-phenyl)-
ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-
propionamide**

The title compound was prepared analogously to (R)-2-benzyloxy-N-((1S,2R,3S,4R)-4-
{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-
yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 181, step 5), by substituting N-
{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-
yl]-2,3-dihydroxy-cyclopentyl} -3-hydroxy-propionamide (Intermediate ZG) for (R)-N-
{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-
9-yl]-2,3-dihydroxy-cyclopentyl} -2-benzyloxy-propionamide (Example 181, step 4) and

(4-Sulfamoyl-phenyl)-carbamic acid phenyl ester (Intermediate ZE) for pyridin-3-yl-carbamic acid phenyl ester (Intermediate ZB).

Example 187

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-[3-(3-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide

The title compound was prepared analogously to (R)-2-benzyloxy-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 181, step 5), by substituting N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Intermediate ZG) for (R)-N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-2-benzyloxy-propionamide (Example 181, step 4) and (3-Sulfamoyl-phenyl)-carbamic acid phenyl ester (Intermediate ZF) for pyridin-3-yl-carbamic acid phenyl ester (Intermediate ZB).

Example 188

N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-[3-(4-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-[3-(4-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 186), by substituting N-((1S,2R,3S,4R)-4-{2-((R)-3-amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Intermediate ZH) for N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Intermediate ZG).

Example 189

N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide

The title compound is prepared analogously to N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Example 184), by substituting N-((1S,2R,3S,4R)-4-{2-((R)-3-amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Intermediate ZH) for N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Intermediate ZG).

Example 190

N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-{(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 185), by substituting N-((1S,2R,3S,4R)-4-{2-((R)-3-amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Intermediate ZH) for N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Intermediate ZG).

Example 191

N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-{(R)-3-[3-(3-sulfamoyl-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(3-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-

dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 187), by substituting N-((1S,2R,3S,4R)-4-{2-((R)-3-amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Intermediate ZH) for N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Intermediate ZG).

Example 192

N-((1S,2R,3S,4R)-2,3-Dihydroxy-4-{6-(2-hydroxy-2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-cyclopentyl)-2-hydroxy-acetamide

Step 1: N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-chloro-phenyl)-2-hydroxy-ethylamino]-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

A solution comprising: N-((1S,2R,3S,4R)-4-{2-((R)-3-Amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-chloro-phenyl)-2-hydroxy-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (Intermediate ZO; 1 equivalent) and pyridin-3-yl-carbamic acid phenyl ester (Intermediate ZB; 1 equivalent) in NMP is stirred at 100°C for 1 hour. The solvent is removed *in vacuo* and the title compound is obtained after purification by column chromatography / crystallisation.

Step 2: N-((1S,2R,3S,4R)-2,3-Dihydroxy-4-{6-(2-hydroxy-2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-cyclopentyl)-2-hydroxy-acetamide

To a stirred solution of N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-chloro-phenyl)-2-hydroxy-ethylamino]-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide in ethanol is added 10 equivalents of ammonium formate and 20mol% of 10% palladium on carbon. The mixture is stirred at 80°C for five hours, allowed to cool and filtered through Celite™. Removal of the solvent under reduced pressure affords the title compound.

Example 193**N-((1S,2R,3S,4R)-2,3-Dihydroxy-4-[2-((R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl)-6-(2-hydroxy-2,2-diphenyl-ethylamino)-purin-9-yl]-cyclopentyl)-2-hydroxy-acetamide**

The title compound is prepared analogously to N-((1S,2R,3S,4R)-2,3-dihydroxy-4-{6-(2-hydroxy-2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-cyclopentyl)-2-hydroxy-acetamide (Example 192), by substituting (3-hydroxy-benzyl)-carbamic acid phenyl ester (Intermediate ZD) for pyridin-3-yl-carbamic acid phenyl ester (Intermediate ZB).

Example 194**N-((1S,2R,3S,4R)-2,3-Dihydroxy-4-{6-((S)-1-hydroxymethyl-2-phenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-cyclopentyl)-2-hydroxy-acetamide**

The title compound is prepared analogously to (R)-2-benzyloxy-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 181, step 5), by substituting Acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-((S)-1-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbonyl}-methyl ester (intermediate ZM) for (R)-N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-benzyloxy-propionamide (Example 181, step 4).

Example 195**N-((1S,2R,3S,4R)-2,3-Dihydroxy-4-{6-[(S)-1-hydroxymethyl-2-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-cyclopentyl)-2-hydroxy-acetamide**

The title compound is prepared analogously to (R)-2-benzyloxy-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 181, step 5), by substituting Acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-((S)-1-hydroxymethyl-2-(4-hydroxy-phenyl)-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbonyl}-methyl ester

(Intermediate ZN) for (R)-N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-benzyloxy-propionamide (Example 181, step 4).

Example 196

N-((1S,2R,3S,4R)-2,3-Dihydroxy-4-{2-[(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl]-6-[(S)-1-hydroxymethyl-2-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-cyclopentyl)-2-hydroxy-acetamide

The title compound is prepared analogously to N-((1S,2R,3S,4R)-2,3-Dihydroxy-4-{6-[(S)-1-hydroxymethyl-2-(4-hydroxy-phenyl)-ethylamino]-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-cyclopentyl)-2-hydroxy-acetamide (Example 195), by substituting (3-hydroxy-benzyl)-carbamic acid phenyl ester (Intermediate ZD) for pyridin-3-yl-carbamic acid phenyl ester (Intermediate ZB).

Example 197

N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[4-(3-pyridin-3-yl-ureido)-pyrazol-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

The title compound is prepared analogously to (R)-2-benzyloxy-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 181, step 5), by substituting N-{(1S,2R,3S,4R)-4-[2-(4-amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide (Intermediate ZP) for (R)-N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-benzyloxy-propionamide (Example 181, step 4).

Example 198

N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[4-(3-pyridin-2-ylmethyl-ureido)-pyrazol-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

The title compound is prepared analogously to N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Example 184), by substituting N-

{(1S,2R,3S,4R)-4-[2-(4-amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide (Intermediate ZP) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-3-hydroxy-propionamide (Intermediate ZG).

Example 199

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{4-[3-(3-hydroxy-benzyl)-ureido]-pyrazol-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 185), by substituting N-{(1S,2R,3S,4R)-4-[2-(4-amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide (Intermediate ZP) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-3-hydroxy-propionamide (Intermediate ZG).

Example 200

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{4-[3-(3-sulfamoyl-phenyl)-ureido]-pyrazol-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(3-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 187), by substituting N-{(1S,2R,3S,4R)-4-[2-(4-amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide (Intermediate ZP) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-3-hydroxy-propionamide (Intermediate ZG).

Example 201

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{4-[3-(4-sulfamoyl-phenyl)-ureido]-pyrazol-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-[3-(4-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 186), by substituting N-[(1S,2R,3S,4R)-4-[2-(4-amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide (Intermediate ZP) for N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Intermediate ZG).

Example 202

N-[(1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[4-(3-pyridin-3-yl-ureido)-pyrazol-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

The title compound is prepared analogously to (R)-2-benzyloxy-N-[(1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 181, step 5), by substituting N-[(1S,2R,3S,4R)-4-{2-(4-amino-pyrazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (Intermediate ZQ) for (R)-N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-2-benzyloxy-propionamide (Example 181, step 4).

Example 203

N-[(1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[4-(3-pyridin-2-ylmethyl-ureido)-pyrazol-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Example 184), by substituting N-[(1S,2R,3S,4R)-4-{2-(4-amino-pyrazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (Intermediate ZQ) for N-

{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -3-hydroxy-propionamide (Intermediate ZG).

Example 204

N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-{4-[3-(3-hydroxy-benzyl)-ureido]-pyrazol-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 185), by substituting N-((1S,2R,3S,4R)-4-{2-(4-amino-pyrazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (Intermediate ZQ) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -3-hydroxy-propionamide (Intermediate ZG).

Example 205

N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-{4-[3-(3-sulfamoyl-phenyl)-ureido]-pyrazol-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(3-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 187), by substituting N-((1S,2R,3S,4R)-4-{2-(4-amino-pyrazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (Intermediate ZQ) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -3-hydroxy-propionamide (Intermediate ZG).

Example 206

N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-{4-[3-(4-sulfamoyl-phenyl)-ureido]-pyrazol-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-[3-(4-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 186), by substituting N-((1S,2R,3S,4R)-4-{2-(4-amino-pyrazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (Intermediate ZQ) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-3-hydroxy-propionamide (Intermediate ZG).

Example 207

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-[4-[3-(3-hydroxy-benzyl)-ureido]-pyrazol-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-[(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 185), by substituting N-{(1S,2R,3S,4R)-4-[2-(4-amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Intermediate ZR) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-3-hydroxy-propionamide (Intermediate ZG).

Example 208

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-[4-(3-pyridin-2-ylmethyl-ureido)-pyrazol-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide

The title compound is prepared analogously to N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Example 184), by substituting N-{(1S,2R,3S,4R)-4-[2-(4-amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Intermediate ZR) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-3-hydroxy-propionamide (Intermediate ZG).

Example 209

1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenylethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid 3-hydroxy-benzylamide

The title compound is prepared analogously to 1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenylethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid methylamide (Example 173), by substituting 3-hydroxybenzylamine for methylamine.

Example 210

1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenylethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid (pyridin-2-ylmethyl)-amide

The title compound is prepared analogously to 1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenylethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid methylamide (Example 173), by substituting C-pyridin-2-yl-methylamine for methylamine.

Example 211

1-[9-[(1R,2S,3R,4S)-2,3-Dihydroxy-4-(2-hydroxy-acetyl-amino)-cyclopentyl]-6-(2,2-diphenylethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid (pyridin-2-ylmethyl)-amide

The title compound is prepared analogously to 1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenylethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid methylamide (Example 173), by substituting 1-[9-[(1R,2S,3R,4S)-2,3-dihydroxy-4-(2-hydroxy-acetyl-amino)-cyclopentyl]-6-(2,2-diphenylethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid ethyl ester (Intermediate ZS) for 1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenylethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid ethyl ester (Example 173, step 1) and C-pyridin-2-yl-methylamine for methylamine.

Example 212

1-[9-[(1R,2S,3R,4S)-2,3-Dihydroxy-4-(2-hydroxy-acetylamino)-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid 3-hydroxy-benzylamide

The title compound is prepared analogously to 1-[9-[(1R,2S,3R,4S)-2,3-Dihydroxy-4-(2-hydroxy-acetylamino)-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid (pyridin-2-ylmethyl)-amide (Example 211), by substituting 3-hydroxybenzylamine for C-pyridin-2-yl-methylamine.

Example 213

1-[9-[(1R,2S,3R,4S)-2,3-Dihydroxy-4-(2-hydroxy-acetylamino)-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid (3-sulfamoyl-phenyl)-amide

The title compound is prepared analogously to 1-[9-[(1R,2S,3R,4S)-2,3-Dihydroxy-4-(2-hydroxy-acetylamino)-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid (pyridin-2-ylmethyl)-amide (Example 211), by substituting 3-amino-benzenesulfonamide for C-pyridin-2-yl-methylamine.

Example 214

1-[9-[(1R,2S,3R,4S)-2,3-Dihydroxy-4-(2-hydroxy-acetylamino)-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid (4-sulfamoyl-phenyl)-amide

The title compound is prepared analogously to 1-[9-[(1R,2S,3R,4S)-2,3-Dihydroxy-4-(2-hydroxy-acetylamino)-cyclopentyl]-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid (pyridin-2-ylmethyl)-amide (Example 211), by substituting 4-amino-benzenesulfonamide for C-pyridin-2-yl-methylamine.

Example 215

N-{(1S,2R,3S,4R)-2,3-Dihydroxy-4-[2-{4-[3-(3-hydroxy-benzyl)-ureido]-pyrazol-1-yl}-6-((S)-1-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-cyclopentyl}-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 185), by substituting N-[(1S,2R,3S,4R)-4-[2-(4-amino-pyrazol-1-yl)-6-((S)-1-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide (intermediate ZT) for N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Intermediate ZG).

Example 216

N-[(1S,2R,3S,4R)-2,3-Dihydroxy-4-(6-((S)-1-hydroxymethyl-2-phenyl-ethylamino)-2-{4-[3-(4-sulfamoyl-phenyl)-ureido]-pyrazol-1-yl}-purin-9-yl)-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(4-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 186), by substituting N-[(1S,2R,3S,4R)-4-[2-(4-amino-pyrazol-1-yl)-6-((S)-1-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide (intermediate ZT) for N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Intermediate ZG).

Example 217

N-[(1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[4-(3-pyridin-3-yl-ureido)-imidazol-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to (R)-2-benzyloxy-N-[(1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 181, step 5), by substituting N-[(1S,2R,3S,4R)-4-[2-(4-amino-imidazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide (Intermediate ZU) for (R)-N-

{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-benzyloxy-propionamide (Example 181, step 4).

Example 218

N-[(1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[4-(3-pyridin-2-ylmethyl-ureido)-imidazol-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 184), by substituting N-{(1S,2R,3S,4R)-4-[2-(4-amino-imidazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide (Intermediate ZU) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -3-hydroxy-propionamide (Intermediate ZG).

Example 219

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{4-[3-(3-hydroxy-benzyl)-ureido]-imidazol-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 185), by substituting N-{(1S,2R,3S,4R)-4-[2-(4-amino-imidazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide (Intermediate ZU) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -3-hydroxy-propionamide (Intermediate ZG).

Example 220

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{4-[3-(3-sulfamoyl-phenyl)-ureido]-imidazol-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(3-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 187), by substituting N-

{(1S,2R,3S,4R)-4-[2-(4-amino-imidazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide (Intermediate ZU) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-3-hydroxy-propionamide (Intermediate ZG).

Example 221

N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[4-(3-pyridin-3-yl-ureido)-imidazol-1-yl]-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

The title compound is prepared analogously to (R)-2-benzyloxy-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide (Example 181, step 5), by substituting N-((1S,2R,3S,4R)-4-{2-(4-Amino-imidazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (intermediate ZV) for (R)-N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-benzyloxy-propionamide (Example 181, step 4).

Example 222

N-((1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[4-(3-pyridin-2-ylmethyl-ureido)-imidazol-1-yl]-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

The title compound is prepared analogously to N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Example 184), by substituting N-((1S,2R,3S,4R)-4-{2-(4-amino-imidazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (intermediate ZV) for N-((1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Intermediate ZG).

Example 223**N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-{4-[3-(3-hydroxy-benzyl)-ureido]-imidazol-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide**

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 185), by substituting N-((1S,2R,3S,4R)-4-{2-(4-Amino-imidazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (intermediate ZV) for N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Intermediate ZG).

Example 224**N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-{4-[3-(3-sulfamoyl-phenyl)-ureido]-imidazol-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide**

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(3-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 187), by substituting N-((1S,2R,3S,4R)-4-{2-(4-amino-imidazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (intermediate ZV) for N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Intermediate ZG).

Example 225**N-[(1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[3-(3-pyridin-3-yl-ureido)-[1,2,4]triazol-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide**

The title compound is prepared analogously to (R)-2-benzyloxy-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-

2,3-dihydroxy-cyclopentyl)-propionamide (Example 181, step 5), by substituting N-
{(1S,2R,3S,4R)-4-[2-(3-Amino-[1,2,4]triazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide (Intermediate ZW) for (R)-N-
{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-benzyloxy-propionamide (Example 181, step 4).

Example 226

N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[3-(3-pyridin-2-ylmethyl-ureido)-[1,2,4]triazol-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

The title compound is prepared analogously to N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Example 184), by substituting N-
{(1S,2R,3S,4R)-4-[2-(3-Amino-[1,2,4]triazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide (Intermediate ZW) for N-
{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -3-hydroxy-propionamide (Intermediate ZG).

Example 227

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{3-[3-(3-hydroxy-benzyl)-ureido]-[1,2,4]triazol-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 185), by substituting N-
{(1S,2R,3S,4R)-4-[2-(3-Amino-[1,2,4]triazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -2-hydroxy-acetamide (Intermediate ZW) for N-
{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl} -3-hydroxy-propionamide (Intermediate ZG).

Example 228

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{3-[3-(3-sulfamoyl-phenyl)-ureido]-[1,2,4]triazol-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(3-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 187), by substituting N-{(1S,2R,3S,4R)-4-[2-(3-Amino-[1,2,4]triazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide (Intermediate ZW) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-3-hydroxy-propionamide (Intermediate ZG).

Example 229

N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{3-[3-(4-sulfamoyl-phenyl)-ureido]-[1,2,4]triazol-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(4-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 186), by substituting N-{(1S,2R,3S,4R)-4-[2-(3-Amino-[1,2,4]triazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide (Intermediate ZW) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-3-hydroxy-propionamide (Intermediate ZG).

Example 230

N-[(1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[3-(3-pyridin-3-yl-ureido)-[1,2,4]triazol-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to (R)-2-benzyloxy-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl)-propionamide (Example 181, step 5), by substituting N-((1S,2R,3S,4R)-4-{2-(3-Amino-[1,2,4]triazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-

ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (intermediate ZX) for (R)-N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenylethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-2-benzyloxy-propionamide (Example 181, step 4).

Example 231

N-[(1S,2R,3S,4R)-4-{6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-[3-(3-pyridin-2-ylmethyl-ureido)-[1,2,4]triazol-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-{6-(2,2-Diphenylethylamino)-2-[(R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide (Example 184), by substituting N-[(1S,2R,3S,4R)-4-{2-(3-Amino-[1,2,4]triazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (intermediate ZX) for N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenylethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Intermediate ZG).

Example 232

N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-{3-[3-(3-hydroxy-benzyl)-ureido]-[1,2,4]triazol-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenylethylamino)-2-{(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 185), by substituting N-[(1S,2R,3S,4R)-4-{2-(3-Amino-[1,2,4]triazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (Intermediate ZX) for N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenylethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Intermediate ZG).

Example 233**N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-{3-[3-(3-sulfamoyl-phenyl)-ureido]-[1,2,4]triazol-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide**

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-((R)-3-[3-(3-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl)-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 187), by substituting N-((1S,2R,3S,4R)-4-{2-(3-amino-[1,2,4]triazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (Intermediate ZX) for N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Intermediate ZG).

Example 234**N-[(1S,2R,3S,4R)-4-(6-[2,2-Bis-(4-hydroxy-phenyl)-ethylamino]-2-{3-[3-(4-sulfamoyl-phenyl)-ureido]-[1,2,4]triazol-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide**

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-((R)-3-[3-(4-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl)-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 186), by substituting N-((1S,2R,3S,4R)-4-{2-(3-amino-[1,2,4]triazol-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (Intermediate ZX) for N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Intermediate ZG).

Example 235**N-[(1S,2R,3S,4R)-4-[2-[2-(4-Chloro-phenyl)-ethoxy]-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-propionamide**

Step 1: N-[(3aR,4S,6R,6aS)-6-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,2-dimethyl-tetrahydro-cyclopenta[1,3]dioxol-4-yl]-propionamide

The title compound is prepared by dissolving N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (Intermediate J) in a 2:1 mixture of acetone and 2,2-dimethoxypropane with a catalytic amount of toluene-4-sulfonic acid, and stirring at room temperature overnight. The volatile components are removed under reduced pressure to afford the title compound.

Step 2: N-{(3aR,4S,6R,6aS)-6-[2-[2-(4-Chloro-phenyl)-ethoxy]-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,2-dimethyl-tetrahydro-cyclopenta[1,3]dioxol-4-yl}-propionamide

The title compound is prepared by adding N-{(3aR,4S,6R,6aS)-6-[2-Chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,2-dimethyl-tetrahydro-cyclopenta[1,3]dioxol-4-yl}-propionamide (example 235, step 1) to a premixed solution of sodium hydride (60% in oil) and 2-(4-chloro-phenyl)-ethanol (1 equivalent) in dry THF. The reaction is stirred at 50°C for 48 hours, before quenching residual sodium hydride with excess aqueous ammonium chloride. The reaction mixture is then partitioned between ethyl acetate and water; the organic phase is washed consecutively with water and brine before drying over magnesium sulfate. Filtration and removal of the volatile components under reduced pressure yields the crude product; purification by column chromatography / crystallisation affords the title compound.

Step 3: N-{(1S,2R,3S,4R)-4-[2-[2-(4-Chloro-phenyl)-ethoxy]-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide

The title compound is prepared by dissolving N-{(3aR,4S,6R,6aS)-6-[2-[2-(4-Chloro-phenyl)-ethoxy]-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,2-dimethyl-tetrahydro-cyclopenta[1,3]dioxol-4-yl}-propionamide (Example 235, Step 2) in THF, adding an equal volume of 1.0M hydrochloric acid, and stirring at room temperature for 48 hours, before diluting with water, and extracting into ethyl acetate. The organic phase is dried over magnesium sulfate, filtered and the volatile components removed under reduced pressure, to yield the title compound.

Example 236

4-{3-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-prop-2-ynyl}-cyclohexanecarboxylic acid methyl ester

The title compound is prepared by combining N-{(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-propionamide (intermediate J), 4-prop-2-ynyl-cyclohexanecarboxylic acid methyl ester (prepared as described by Rieger J.M., Brown M.L., Sullivan G.W., Linden J. and Macdonald T.L.; *J. Med. Chem.*(2001), **44**, 531-539), copper (I) iodide, triphenylphosphine and dichlorobis(triphenylphosphine)palladium(II) in a 2:1 mixture of triethylamine / DMF, and heating by microwave irradiation for 3600 seconds at 120°C. Purification by column chromatography affords the title compound.

Example 237

(2S,3S,4R,5R)-5-(6-(2,2-Diphenyl-ethylamino)-2-{4-[3-(3-hydroxy-benzyl)-ureido]-pyrazol-1-yl}-purin-9-yl)-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

Step 1: (2S,3S,4R,5R)-5-[2-(4-Amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

The title compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-(4-amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide (Intermediate ZP), by substituting (2S,3S,4R,5R)-5-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide (Intermediate ZY) for acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl}-methyl ester (Intermediate Q1) in step ZP1.

Step 2: (2S,3S,4R,5R)-5-(6-(2,2-Diphenyl-ethylamino)-2-{4-[3-(3-hydroxy-benzyl)-ureido]-pyrazol-1-yl}-purin-9-yl)-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-

dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 185), by substituting (2S,3S,4R,5R)-5-[2-(4-Amino-pyrazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide (Example 237, step 1) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-3-hydroxy-propionamide (Intermediate ZG).

Example 238

(2S,3S,4R,5R)-5-(6-(2,2-Diphenyl-ethylamino)-2-{4-[3-(3-hydroxy-benzyl)-ureido]-imidazol-1-yl}-purin-9-yl)-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

Step 1: (2S,3S,4R,5R)-5-[2-(4-Amino-imidazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

The title compound is prepared analogously to N-{(1S,2R,3S,4R)-4-[2-(4-amino-imidazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-2-hydroxy-acetamide (Intermediate ZU), by substituting (2S,3S,4R,5R)-5-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide (Intermediate ZY) for acetic acid {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl}-methyl ester (Intermediate Q1) in step ZU1.

Step 2: (2S,3S,4R,5R)-5-(6-(2,2-Diphenyl-ethylamino)-2-{4-[3-(3-hydroxy-benzyl)-ureido]-imidazol-1-yl}-purin-9-yl)-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 185), by substituting (2S,3S,4R,5R)-5-[2-(4-amino-imidazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide (Example 238, step 1) for N-{(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-3-hydroxy-propionamide (Intermediate ZG).

Example 239**(2S,3S,4R,5R)-5-(6-(2,2-Diphenyl-ethylamino)-2-{3-[3-(3-hydroxy-benzyl)-ureido]-[1,2,4]triazol-1-yl}-purin-9-yl)-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide**

Step 1: (2S,3S,4R,5R)-5-[2-(3-Amino-[1,2,4]triazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-[2-(3-Amino-[1,2,4]triazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide (intermediate ZW), by substituting (2S,3S,4R,5R)-5-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide (Intermediate ZY) for acetic acid [(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentylcarbamoyl]-methyl ester (Intermediate Q1).

Step 2: (2S,3S,4R,5R)-5-(6-(2,2-Diphenyl-ethylamino)-2-{3-[3-(3-hydroxy-benzyl)-ureido]-[1,2,4]triazol-1-yl}-purin-9-yl)-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

The title compound is prepared analogously to N-[(1S,2R,3S,4R)-4-(6-(2,2-diphenyl-ethylamino)-2-{(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Example 185), by substituting (2S,3S,4R,5R)-5-[2-(3-amino-[1,2,4]triazol-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide (Example 239, step 1) for N-[(1S,2R,3S,4R)-4-[2-((R)-3-amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl]-3-hydroxy-propionamide (Intermediate ZG).

Example 240**(2S,3S,4R,5R)-5-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(3-hydroxy-benzyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide**

(2S,3S,4R,5R)-5-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide trifluoroacetate

(Intermediate E) and (3-Hydroxy-benzyl)-carbamic acid phenyl ester (Intermediate UA) are dissolved in methanol and TEA. The reaction mixture is heated using microwave radiation at 100°C for 30 minutes in the Personal Chemistry EmrysTM Optimizer microwave reactor. The reaction mixture is concentrated *in vacuo* and purified by C-18 reverse phase column chromatography eluting with acetonitrile:water (0.1% TFA) (gradient 0-100% acetonitrile) to afford the titled compound.

Example 241

(2S,3S,4R,5R)-5-(6-(2,2-Diphenyl-ethylamino)-2-((R)-3-[3-(4-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl)-purin-9-yl)-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

This compound is prepared analogously to Example 240 by replacing (3-Hydroxy-benzyl)-carbamic acid phenyl ester (Intermediate UA) with (4-Sulfamoyl-phenyl)-carbamic acid phenyl ester (Intermediate UD).

Example 242

(2S,3S,4R,5R)-5-(6-(2,2-Diphenyl-ethylamino)-2-((R)-3-[3-(3-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl)-purin-9-yl)-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

This compound is prepared analogously to Example 240 by replacing (3-Hydroxy-benzyl)-carbamic acid phenyl ester (Intermediate UA) with (3-Sulfamoyl-phenyl)-carbamic acid phenyl ester (Intermediate UC).

Example 243

(2S,3S,4R,5R)-5-(6-(2,2-Diphenyl-ethylamino)-2-((R)-3-(3-pyridin-2-ylmethyl-ureido)-pyrrolidin-1-yl)-purin-9-yl)-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

This compound is prepared analogously to Example 240 by replacing (3-Hydroxy-benzyl)-carbamic acid phenyl ester (Intermediate UA) with Pyridin-2-ylmethyl-carbamic acid phenyl ester (Intermediate UE).

Example 244**(2S,3S,4R,5R)-5-[2-[(R)-3-[3,4-Dioxo-2-(pyridin-3-ylamino)-cyclobut-1-enylamino]-pyrrolidin-1-yl]-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide**

Step 244a: (2S,3S,4R,5R)-5-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(2-methoxy-3,4-dioxo-cyclobut-1-enylamino)-pyrrolidin-1-yl]-purin-9-yl}-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

A mixture comprising (2S,3S,4R,5R)-5-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide trifluoroacetate (Intermediate E) and 3,4-dimethoxy-3-cyclobutene-1,2-dione in absolute EtOH and cat.DMAP is heated using microwave radiation in a Personal Chemistry EmrysTM Optimizer microwave reactor at 120°C for 1 hour. The solvent is removed *in vacuo* and the resulting crude product partitioned between ethyl acetate and water. The organic portion is separated, dried (Na₂SO₄) and concentrated *in vacuo*. Purification by column chromatography on silica gel eluting with EtOAc/*iso*-hexane (30-100% EtOAc) affords the titled compound.

Step 244b: (2S,3S,4R,5R)-5-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(2-methoxy-3,4-dioxo-cyclobut-1-enylamino)-pyrrolidin-1-yl]-purin-9-yl}-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

A mixture comprising (2S,3S,4R,5R)-5-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(2-methoxy-3,4-dioxo-cyclobut-1-enylamino)-pyrrolidin-1-yl]-purin-9-yl}-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide (step244a) in absolute EtOH and catalytic amount of TsOH is heated using microwave radiation in a Personal Chemistry EmrysTM Optimizer microwave reactor at 150 °C for 4000 seconds. The solvent is removed *in vacuo* and the resulting crude product partitioned between ethyl acetate and water. The organic portion is separated, dried (Na₂SO₄) and concentrated *in vacuo*. Purification by C-18 reverse phase column chromatography eluting with acetonitrile:water (0.1% TFA) (gradient 0-100% acetonitrile) to afford the titled compound.

Example 245

(2S,3S,4R,5R)-5-(6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-[2-(3-hydroxy-benzylamino)-3,4-dioxo-cyclobut-1-enylamino]-pyrrolidin-1-yl]-purin-9-yl)-3,4-dihydroxy-tetrahydro-furan-2-carboxylic acid ethylamide

This compound is prepared analogously to Example 244 by replacing 3-aminopyridine with 3-hydroxybenzylamine.

Example 246 – 253

These compounds are prepared from the product of Example 26 step 2 following an analogous procedure to Example 26 step 3, replacing methyl chloroformate with the appropriate acid chloride or anhydride.

Example 254

(R)-N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-propionamide

Step 1: (R)-N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-phenoxy-propionamide.

HATU (61 mg, 0.16 mmol) and R-(+)-2-benzyloxypropionic acid (32 mg, 0.16 mmol) are dissolved in DMF (5 ml) and after stirring for 5 minutes, the solution is treated with 1-{(R)-1-[9-((1R,2S,3R,4S)-4-Amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-3-pyridin-3-yl-urea (Example 26 step 2) (0.1 g, 0.16 mmol) in DMF (0.5 ml). DIPEA (56 µl, 0.32 mmol) is added and the resulting solution is stirred for 2 hours. The mixture is then treated with sat. Na₂CO₃ and 1 ml MeOH and then partitioned between EtOAc and water. The organic portion is separated and concentrated *in vacuo*. The resulting crude product is purified by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% HCl) to afford the title product.

Step 2: (R)-N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-propionamide

A solution of (R)-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-phenoxy-propionamide (83 mg, 0.104 mmol) in EtOH (20 ml) and THF (5 ml) under an inert atmosphere is treated with palladium hydroxide (20%w/w on carbon, 32 mg) followed by acetic acid (2 ml). The reaction mixture is placed under an atmosphere of hydrogen for two weeks after which time, the mixture is filtered and concentrated *in vacuo*. The resulting crude product is washed with EtOH (3 times), filtered and then concentrated *in vacuo*. The residue is dissolved in MeOH (2 ml) and purification by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.1% NH₃) affords the title compound as a white solid. (MH⁺ 707.4)

Example 255

(S)-N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-propionamide

A mixture comprising 1-{(R)-1-[9-((1R,2S,3R,4S)-4-amino-2,3-dihydroxy-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-pyrrolidin-3-yl}-3-pyridin-3-yl-urea (Example 26 step 2) (0.1 g, 0.16 mmol) and TEA (24 µl, 0.18 mmol) in THF (7 ml) at room temperature is treated with (S)-(-)-2-acetoxy-propionyl chloride (24 mg, 0.16 mmol) in MeCN (1 ml) at (0 ° C) over 1 minute. The mixture is allowed to stir at room temperature for 18 h and then treated with sat. Na₂CO₃ (1 ml) and MeOH (1 ml) and then stirred for a further 2 days. The solvent is removed *in vacuo* and purification by reverse phase column chromatography (Isolute™ C18, 0-100% acetonitrile in water – 0.3% NH₃) affords the title compound as a white solid. (MH⁺ 707.7)

Example 256

N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-2-methyl-propionamide

This compound is prepared analogously to Example 255 by replacing (S)-(-)-2-acetoxy-propionyl chloride with 2-acetoxy isobutyryl chloride. The ester hydrolysis is carried out in the presence of 1M NaOH in MeOH. (MH⁺ 721.5)

Example 257

N-((1S,2R,3S,4R)-4-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-((S)-1-hydroxymethyl-2-phenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide

This compound is prepared from N-((1S,2R,3S,4R)-4-[6-((S)-1-benzyl-2-hydroxy-ethylamino)-2-chloro-purin-9-yl]-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (Intermediate VD) analogously to N-((1S,2R,3S,4R)-4-{2-((R)-3-amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide hydrochloride (Example 7 step 5). (MH⁺ 527.26)

Example 258

{(1S,2R,3S,4R)-4-[2-((R)-3-Amino-pyrrolidin-1-yl)-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-carbamic acid methyl ester trifluoroacetate

This compound is prepared from {(1S,2R,3S,4R)-4-[2-chloro-6-(2,2-diphenyl-ethylamino)-purin-9-yl]-2,3-dihydroxy-cyclopentyl}-carbamic acid methyl ester (Intermediate T) analogously to N-((1S,2R,3S,4R)-4-{2-((R)-3-amino-pyrrolidin-1-yl)-6-[2,2-bis-(4-hydroxy-phenyl)-ethylamino]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide hydrochloride (Example 7 step 5). (MH⁺ 572.21)

Examples 259-261

These compounds namely,

N-((1S,2R,3S,4R)-4-{6-((S)-1-Benzyl-2-hydroxy-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-hydroxy-acetamide (MH+ 647.01) (Example 259),

N-[(1S,2R,3S,4R)-4-(6-((S)-1-Benzyl-2-hydroxy-ethylamino)-2-{(R)-3-[3-(4-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-2-hydroxy-acetamide (MH+ 724.99) (Example 260) and

[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-{(R)-3-[3-(3-sulfamoyl-phenyl)-ureido]-pyrrolidin-1-yl}-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-carbamic acid methyl ester (MH+ 771.34) (Example 261)

are prepared from the appropriate starting compounds and corresponding pyrrolidinyl-urea (preparations described herein) analogously to Example 15.

Example 262

1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid (pyridin-2-ylmethyl)-amide

Step 1: 1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid QBA289

A solution comprising 1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid ethyl ester (Example 173 step 1) (0.4 g, 0.64 mmol) in water (3 ml) is treated with 1M KOH in MeOH (6 ml) and stirred at room temperature for 48 h. The solvent is removed *in vacuo* and the resulting crude product is dissolved in water (5 ml) and acidified to pH 3-4 with 1.5N HCl. The solution is extracted with EtOAc and the organic portion is dried over sodium sulfate and concentrated *in vacuo*. Purification by chromatography on silica eluting with 5% MeOH in chloroform affords the title compound.

Step 2: 1-[9-((1R,2S,3R,4S)-2,3-Dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2

-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid (pyridin-2-ylmethyl)-amide

A mixture comprising 1-[9-((1R,2S,3R,4S)-2,3-dihydroxy-4-propionylamino-cyclopentyl)-6-(2,2-diphenyl-ethylamino)-9H-purin-2-yl]-1H-pyrazole-4-carboxylic acid (0.06 g, 0.01 mmol) in dry DCM (10 ml) is treated with 2-aminomethyl pyridine (0.021 g, 0.2 mmol) followed by HOBt (0.027 g, 0.2 mmol), 1-[Bis(dimethylamino)methylene]-1H-benzotriazoliumhexafluorophosphate-3-oxide HBTU (0.076 g, 0.2 mmol), N-methyl morpholine (0.02 g, 0.2 mmol) and DMAP (1 ml). The resulting mixture is stirred at room temperature for 24 h and then concentrated *in vacuo*. The resulting residue purified by preparative TLC eluting with 10% MeOH in chloroform to afford the title compound. (MH⁺ 687.1)

Example 263

N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-3-hydroxy-propionamide hydrochloride

This compound is prepared analogously to (R)-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-2-phenoxy-propionamide (Example 254 step 1) by replacing R-(+)-2-benzyloxypionic acid with 3-tert-butoxypionic acid. (MH⁺ 707.4)

Example 264


(R)-2-Benzyloxy-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide hydrochloride

This compound is prepared analogously to (R)-N-((1S,2R,3S,4R)-4-{6-(2,2-diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-

cyclopentyl)-2-phenoxy-propionamide (Example 254 step 1) by replacing R-(+)-2-benzyloxypropionic acid with (R)-2-benzyloxy-propionic acid. (MH⁺ 797.7)

Claims:
















1. A compound of formula (I), or stereoisomers or pharmaceutically acceptable salts thereof,







































(I)



















wherein R¹, R² and R³ are


















Ex.	R ¹	R ²	R ³
1			
2			
3			
4			



















5			
6			
7			
8			
9			




















10			
11			
12			
13			
14			
15			
















16			
17			
18			
19			
20			
21			



















22			
23			
24			
25			
26			
27			






















28			
29	H-		
30			
31			
32			
33			






















34			
35			
36			
37			
38			
39			






















40			
41			
42			
43			
44		-H	
45			
46			



















47			
48			
49			
50			
51			
















52			
53			
54			
55			
56			
57			
















58			
59			
60			
61			
62			
63			
64			



















65			
66			
67			
68			
69			
70			
71			



















72			
73			
74			
75			
76			
77			
78			






















79			
80			
81			
82			
83			
84			



















85			
86			
87			
88			
89			






















90	 <chem>CCC(=O)C</chem>	 <chem>CC(c1ccccc1)(c2ccccc2)CC</chem>	 <chem>CN1CC[C@H](NC(=O)c2ccncc2)C1</chem>
91	 <chem>CCC(=O)C</chem>	 <chem>CC(C)CC(C)C</chem>	 <chem>CN1CC[C@H](NC(=O)c2ccncc2)C1</chem>
92	 <chem>CCC(=O)C</chem>	 <chem>CC(C)(C)CC(C)C</chem>	 <chem>CN1CC[C@H](NC(=O)c2ccncc2)C1</chem>
93	 <chem>CCC(=O)C</chem>	 <chem>CC(C)CC(C)C</chem>	 <chem>CN1CC[C@H](NC(=O)c2cc3ncoc3c2C)C1</chem>
94	 <chem>CC(=O)C1CCC1</chem>	 <chem>CC(c1ccccc1)(c2ccccc2)CC</chem>	 <chem>CN1CC[C@H](NC(=O)Cc2ccccc2)C1</chem>



















95			
96			
97			
98			
99			
100			


















101			
102			
103			
104			
105			
106			






















107			
108			
109			
110			
111			
112			
113			















114			
115			
116			
117			
118			
119			













120			
121			
122			
123			
124			
125			
126			













127			
128			
129			
130			
131			
132			



















133			
134			
135			
136			
137			
138			



















139			
140			
141			
142			
143			
144			
145			















146			
147			
148			
149			
150			













151			
152			
153			
154			













155			
156			
157			
158			













159			
160			
161			
162			
163			
164			













165			
166			
167			
168		-H	
169			-Cl
170		-H	
171			













172			
173			
174		-H	
175			
176			













177			
178			
179			
180			













Ex.	R ¹	R ²	R ³
181			
182			
183			
184			













185			
186			
187			
188			













189			
190			
191			
192			













193	 <chem>CC(=O)CO</chem>	 <chem>CC1(Cc2ccccc2)C(O)CCCC1</chem>	 <chem>CN1CCCC1C(=O)NCc2ccc(O)cc2</chem>
194	 <chem>CC(=O)CO</chem>	 <chem>CC(C)C(O)Cc1ccccc1</chem>	 <chem>CN1CCCC1C(=O)Nc1ccncc1</chem>
195	 <chem>CC(=O)CO</chem>	 <chem>CC(C)C(O)Cc1ccc(O)cc1</chem>	 <chem>CN1CCCC1C(=O)Nc1ccncc1</chem>
196	 <chem>CC(=O)CO</chem>	 <chem>CC(C)C(O)Cc1ccc(O)cc1</chem>	 <chem>CN1CCCC1C(=O)NCc2ccc(O)cc2</chem>













197	 <chem>CC(=O)CO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)C</chem>	 <chem>CN1C=NC=C1NC(=O)NC2=CC=CC=N2</chem>
198	 <chem>CC(=O)CO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)C</chem>	 <chem>CN1C=NC=C1NC(=O)NCC2=CC=CC=N2</chem>
199	 <chem>CC(=O)CO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)C</chem>	 <chem>CN1C=NC=C1NC(=O)NCC3=CC=C(O)C=C3</chem>
200	 <chem>CC(=O)CO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)C</chem>	 <chem>CN1C=NC=C1NC(=O)NCC3=CC=C(S(=O)(=O)N)C=C3</chem>













201			
202			
203			
204			













205			
206			
207			
208			













209			
210			
211			
212			













213			
214			
215			
216			
















217			
218			
219			
220			
















221	 <chem>CC(=O)CO</chem>	 <chem>CC(c1ccc(O)cc1)c2ccc(O)cc2</chem>	 <chem>Cn1cc[nH]1NC(=O)Nc2ccncc2</chem>
222	 <chem>CC(=O)CO</chem>	 <chem>CC(c1ccc(O)cc1)c2ccc(O)cc2</chem>	 <chem>Cn1cc[nH]1NC(=O)NCc2ccncc2</chem>
223	 <chem>CC(=O)CO</chem>	 <chem>CC(c1ccc(O)cc1)c2ccc(O)cc2</chem>	 <chem>Cn1cc[nH]1NC(=O)NCc2ccc(O)cc2</chem>
224	 <chem>CC(=O)CO</chem>	 <chem>CC(c1ccc(O)cc1)c2ccc(O)cc2</chem>	 <chem>Cn1cc[nH]1NC(=O)Nc2ccc(S(=O)(=O)N)cc2</chem>
















225	 <chem>CC(=O)CO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)CC</chem>	 <chem>CC1=CN=C(NC(=O)Nc2ccc(cc2N)nc3ccncc3)N1</chem>
226	 <chem>CC(=O)CO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)CC</chem>	 <chem>CC1=CN=C(NC(=O)NCCc2ccncc2)N1</chem>
227	 <chem>CC(=O)CO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)CC</chem>	 <chem>CC1=CN=C(NC(=O)NCCc2ccc(cc2)O)N1</chem>
228	 <chem>CC(=O)CO</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)CC</chem>	 <chem>CC1=CN=C(NC(=O)Nc2ccc(cc2)S(=O)(=O)N)N1</chem>
















229			
230			
231			
232			



















233	 <chem>CC(=O)CO</chem>	 <chem>CC(c1ccc(O)cc1)-c2ccc(O)cc2</chem>	 <chem>Cn1nc(NC(=O)Nc2ccc(S(=O)(=O)N)cc2)nn1</chem>
234	 <chem>CC(=O)CO</chem>	 <chem>CC(c1ccc(O)cc1)-c2ccc(O)cc2</chem>	 <chem>Cn1nc(NC(=O)Nc2ccc(S(=O)(=O)N)cc2)nn1</chem>
235	 <chem>CCC(=O)C</chem>	 <chem>CC(c1ccccc1)-c2ccccc2</chem>	 <chem>COCCc1ccc(Cl)cc1</chem>
236	 <chem>CCC(=O)C</chem>	 <chem>CC(c1ccccc1)-c2ccccc2</chem>	 <chem>COC(=O)[C@H]1CCCC[C@@H]1CC#C</chem>







Example	R1	R2	R3
237			
238			
239			
240			
241			

242	 <chem>CC(=O)c1ccoc1</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)C3=CC=CC=C3</chem>	 <chem>CN1CC[C@H](NC(=O)NC2=CC=CC=N2)C1</chem>
243	 <chem>CC(=O)c1c(C)n(C)c(C)c1</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)C3=CC=CC=C3</chem>	 <chem>CN1CC[C@H](NC(=O)NC2=CC=CC=N2)C1</chem>
244	 <chem>CC(=O)c1c(C)nc(C)o1</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)C3=CC=CC=C3</chem>	 <chem>CN1CC[C@H](NC(=O)NC2=CC=CC=N2)C1</chem>
245	 <chem>CC(=O)c1ccncc1</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)C3=CC=CC=C3</chem>	 <chem>CN1CC[C@H](NC(=O)NC2=CC=CC=N2)C1</chem>
246	 <chem>CC(=O)c1ccccc1OC</chem>	 <chem>CC(C1=CC=CC=C1)(C2=CC=CC=C2)C3=CC=CC=C3</chem>	 <chem>CN1CC[C@H](NC(=O)NC2=CC=CC=N2)C1</chem>

247			
248			
249			
250			
251			

252			
253			
254			
255			
256			

257			
258			
259			
260			
261			
262			

263			
264			

2. The compound according to claim 1 or stereoisomers or pharmaceutically acceptable salts thereof, represented by ((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-pyridin-3-yl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid methyl ester, or a pharmaceutically acceptable salt thereof.

3. The compound according to claim 1 or stereoisomers or pharmaceutically acceptable salts thereof, represented by ((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-((R)-3-pyrrolidin-3-ylureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-carbamic acid methyl ester, or a pharmaceutically acceptable salt thereof.

4. The compound according to claim 1 or stereoisomers or pharmaceutically acceptable salts thereof, represented by N-((1S,2R,3S,4R)-4-{6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-(3-phenyl-ureido)-pyrrolidin-1-yl]-purin-9-yl}-2,3-dihydroxy-cyclopentyl)-propionamide, or a pharmaceutically acceptable salt thereof.

5. The compound according to claim 1 or stereoisomers or pharmaceutically acceptable salts thereof, represented by N-[(1S,2R,3S,4R)-4-(6-(2,2-Diphenyl-ethylamino)-2-[(R)-3-[3-(3-hydroxy-phenyl)-ureido]-pyrrolidin-1-yl]-purin-9-yl)-2,3-dihydroxy-cyclopentyl]-propionamide, or a pharmaceutically acceptable salt thereof.

6. A compound according to any one of claims 1 to 5 or stereoisomers or pharmaceutically acceptable salts thereof for use as a pharmaceutical.
7. A compound according to any one of claims 1 to 5 or stereoisomers or pharmaceutically acceptable salts thereof in combination with an anti-inflammatory, bronchodilatory, anti-histamine or anti-tussive drug substance, said compound and said drug substance being in the same or different pharmaceutical composition.
8. A pharmaceutical composition comprising as active ingredient a compound according to any one of claims 1 to 5 or stereoisomers or pharmaceutically acceptable salts thereof, together with a pharmaceutically acceptable diluent or carrier.
9. A compound according to any one of claims 1 to 5 or stereoisomers or pharmaceutically acceptable salts thereof for use in the treatment of an inflammatory or obstructive airways disease.