(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number
WO 01/01140 A1

(51) International Patent Classification[7]: G01N 33/543, 33/50, A61K 9/127

(21) International Application Number: PCT/GB00/02465

(22) International Filing Date: 27 June 2000 (27.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
9915074.0 28 June 1999 (28.06.1999) GB

(71) Applicant *(for all designated States except US)*: **PROVALIS UK LIMITED** [GB/GB]; Newtech Square, Deeside Industrial Park, Deeside, Flintshire CH5 2NT (GB).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: **NEW, Roger** [GB/GB]; Flat 10, Leinster Mansions, No. 1 Langland Gardens, London NW3 6QB (GB). **TOTH, Istvan** [GB/AU]; 5 Zelita Road, Moggill, QSL 4070 (AU).


(74) Agents: **DANIELS, Jeffrey, Nicholas** et al.; Page White & Farrer, 54 Doughty Street, London WC1N 2LS (GB).

(81) Designated States *(national)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States *(regional)*: ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: EPITOPES FORMED BY NON-COVALENT ASSOCIATION OF CONJUGATES



(57) Abstract: A composition for interacting with a ligand, which composition comprises a non-covalent association of a plurality of distinct conjugates, each conjugate comprising a head group and a tail group, wherein the tail groups of the conjugates form a hydrophobic aggregation and the conjugates are movable within the association so that, in the presence of a ligand, at least two of the head groups are appropriately positioned to form an epitope capable of interacting with the ligand more strongly than each of head groups individually.

WO 01/01140 A1

Published:
— *With international search report.*
— *Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.